

Description

This invention relates to novel aminosulfonylalkyl nitrosoureas having an antitumor activity and to the preparation thereof. The invention is also concerned with pharmaceutical compositions containing said compounds.

Background of the invention

Specific 1 - (2 - chloroethyl) - 1 - nitrosoureas, e.g. carmustine (BCNU; 1,2 - bis - (2 - chloroethyl) - nitrosourea), lomustine (CCNU; 1 - (2 - chloroethyl) - 3 - cyclohexyl - 1 - nitrosourea) and semustine (Methyl-CCNU; 1 - (2 - chloroethyl) - 3 - (4 - methyl - cyclohexyl) - 1 - nitrosourea) are used as chemotherapeutic agents for the treatment of a number of experimental and clinical tumours (1).

It is generally known, however, that these compounds have several serious toxic effects, e.g. damaging effects on liver, kidney, lung organs, and also bone marrow depression, neurotoxicity and gastrointestinal toxicity (1-4).

One new compound of the same type, chlorozotocin or 2 - [3 - (2 - chloroethyl) - 3 - nitrosourido] - D - glucopyranose, which is under clinical evaluation, has been shown to possess decreased bone marrow toxicity (5), but the other types of toxicities are still evident, e.g., liver and lung toxicity (6, 7).

It is also generally known that after some time tumours tend to develop resistance to the anticancer therapy employed, thus rendering the therapy without curative effect.

Thus, the need exists for new and improved chemotherapeutic agents of this type, either or both with increased antitumour activity and decreased toxicity.

General description of the invention

It has now, surprisingly, been found that the novel aminosulfonyl nitrosoureas of the present invention corresponding to the general formula (I) as defined below, possess improved chemotherapeutic properties in comparison with several known and clinically used nitrosourea antitumour agents. Thus, the compounds of the invention show improved therapeutic indexes and good oral cytostatic activity against the Walker 256 carcinosarcoma and the L1210 lymphatic leukemia. The new compounds also possess a greater ameliorative or curative potential as shown by the L1210 experimental results.

The compounds of the invention can be employed in disorders responsive to treatment with anticancer agents both in single therapy and in combination with other anticancer agents or immunosuppressive agents. The compounds can be used as such or combined with either solid or liquid carriers or diluents and made available in varying amounts in such pharmaceutical forms as e.g., tablets, pills, capsules, pellets, powders, ointments, suppositories and aqueous or non-aqueous suspensions and solutions.

Objects of the invention

Accordingly, one object of the invention is to provide new compounds of the general formula (I), having the aforesaid activity, preferably also with a relatively low degree of toxicity.

A second object is to provide such type of compounds, which can be employed in disorders responsive to treatment with anticancer agents and with immunosuppressive agents for the amelioration or palliation thereof.

Another object of the invention is to provide processes for preparing the new compounds having the general formula (I).


A further object of the invention is to provide a method of treating a living animal body suffering from a disorder responsive to treatment with anticancer agents and with immunosuppressive agents, which comprises the step of administering to said living animal body an effective amount of a compound having the general formula (I), said compound being administered in an amount sufficient to at least mitigate said disorder.

Yet another object of the invention is to provide compositions containing as an active ingredient one or more of the compounds, having the general formula (I), preferably together with a pharmaceutically acceptable carrier and, if desired, other pharmacologically active agents.

Other objects of the invention will be apparent to one skilled in the art, and still other objects will become apparent hereinafter.

Summary of the invention

According to the invention there are provided novel compounds having the general formula:



wherein A is lower alkyl or halolower alkyl; B is hydrogen, lower alkenyl, lower alkynyl, 3-6 C cycloalkyl, $-\text{RSO}_2\text{NR}^3\text{R}^4$, or lower alkyl optionally monosubstituted with lower alkoxy, or 3-6 C cycloalkyl;

R is straight or branched 2—5 C alkylene, optionally monosubstituted with $\text{SO}_2\text{NR}^3\text{R}^4$ or CONR^3R^4 , said alkylene always containing at least two carbon atoms separating the N of the urea from any $\text{SO}_2\text{NR}^1\text{R}^2$ or $\text{SO}_2\text{NR}^3\text{R}^4$;

R^1 and R^2 are the same or different and selected from hydrogen, 5—6 C cycloalkyl, lower alkoxy, phenyl, benzyl and straight or branched 1—6 C alkyl, optionally monosubstituted with hydroxy, lower alkoxy, $\text{SO}_2\text{NR}^3\text{R}^4$, or CONR^3R^4 ;

R^1 and R^2 may also together form a 4—5 C alkylene;

R^3 and R^4 are the same or different and selected from hydrogen or lower alkyl, R^3 and R^4 may also together form a 4—5 C alkylene.

In this disclosure the expression "lower" means that the group referred to contains 1—4 C (C=carbon) atoms, inclusive. Thus, lower alkyl, lower alkenyl, lower alkynyl and lower alkoxy include for instance: methyl, ethyl, propyl, isopropyl, butyl, secondary butyl, isobutyl, tertiary butyl, allyl, propargyl, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, secondary butoxy and tertiary butoxy.

Furthermore the expression "halo" means fluorine, chlorine, bromine or iodine.

As regards the substituent A, it is preferred that this substituent is 2-chloroethyl or 2-fluoroethyl, especially 2-chloroethyl.

Also, as regards the substituent B, it is preferred that this substituent is hydrogen or lower alkyl, especially hydrogen.

Furthermore, such compounds are preferred wherein R is unsubstituted straight 2—5 C alkylene or wherein R is ethylene, especially unsubstituted ethylene.

Especially preferred compounds are such wherein one of the substituents R^1 and R^2 is methyl and the other is lower alkyl or 2-hydroxyethyl, or wherein R^1 and R^2 are the same and are hydrogen, methyl, ethyl, or propyl, or wherein R^1 and R^2 together are 5 C alkylene.

Most preferred compounds are such wherein R^1 and R^2 are hydrogen and/or methyl.

The following compounds are preferred:

- a) 1 - (2 - chloroethyl) - 3 - [2 - (dimethylaminosulfonyl)ethyl] - 1 - nitrosourea
- b) 1 - (2 - chloroethyl) - 3 - [2 - (diethylaminosulfonyl)ethyl] - 1 - nitrosourea
- c) 3 - [2 - (aminosulfonyl)ethyl] - 1 - (2 - chloroethyl) - 1 - nitrosourea
- d) 1 - (2 - chloroethyl) - 1 - nitroso - 3 - [2 - (1 - piperidinosulfonyl)ethyl] - urea
- e) 1 - (2 - chloroethyl) - 3 - [2 - (methylaminosulfonyl)ethyl] - 1 - nitrosourea
- f) 1 - (2 - chloroethyl) - 3 - [2 - (dipropylaminosulfonyl)ethyl] - 1 - nitrosourea
- g) 1 - (2 - chloroethyl) - 3 - [2 - (2 - hydroxyethylaminosulfonyl)ethyl] - 1 - nitrosourea
- h) 1 - (2 - chloroethyl) - 3 - [2 - (N - 2 - hydroxyethyl - N - methyl - aminosulfonyl) - ethyl] - 1 - nitrosourea
- i) 1 - (2 - chloroethyl) - 3 - (2 - [2 - methoxyethylaminosulfonyl)ethyl] - 1 - nitrosourea
- j) 1 - (2 - chloroethyl) - 3 - [2 - (dimethylaminosulfonyl)ethyl] - 3 - methyl - 1 - nitrosourea
- k) 1 - (2 - chloroethyl) - 3 - (5 - [dimethylaminosulfonyl]pentyl) - 1 - nitrosourea
- l) 1 - (2 - chloroethyl) - 3 - (3 - [dimethylaminosulfonyl]propyl) - 1 - nitrosourea
- m) 1 - (2 - chloroethyl) - 3 - (4 - [dimethylaminosulfonyl]butyl) - 1 - nitrosourea
- n) 1 - (2 - chloroethyl) - 3 - (2 - [N - ethyl - N - propylaminosulfonyl)ethyl] - 1 - nitrosourea


Methods of preparation

The compounds having the general formula (I) may be prepared by conventional methods.

A general process (Method 1 below) for preparing compounds having the general formula (I) is as follows:

Method 1

Compounds of the general formula (I) are prepared by N-nitrosation of a corresponding urea having the general formula



wherein R^1 , R^2 , R, B and A are as defined above.


Among other methods for preparing compounds having the general formula (I) the following may be mentioned:

Method 2

An amine having the general formula



wherein R^1 , R^2 , R and B are as defined above, is reacted with an N - alkyl - N - nitroso - carbamoyl derivative of the general formula



wherein A is as defined before, and wherein L is a suitable leaving group.

Methods 1 and 2 are illustrated by the following processes:

A. A process according to Method 1, characterized by reacting a urea (II) with a nitrosating agent. Several different such agents are known and the following may be mentioned: Metal nitrites, such as alkali metal nitrites, e.g., sodium, potassium and lithium nitrite, in combination with an acid, e.g. hydrochloric acid, acetic acid, formic acid and methane sulfonic acid; nitrosyl chloride (NOCl); dinitrogen trioxide (N_2O_3); dinitrogen tetroxide (N_2O_4); mixtures of various nitrogen oxides; nitrosyl sulfuric acid ($\text{HO}_3\text{S}-\text{ONO}$); nitrosonium tetrafluoroborate (NOBF_4); alkyl nitrites, e.g. propyl nitrite and isoamyl nitrite; and nitric acid in combination with a suitable reducing agent, e.g., copper dust.

Although the process may be performed at ambient temperatures, normally subambient temperatures such as around 0°C and sometimes even substantially lower, e.g., around -60°C , are employed, thus to prevent thermal degradation of the compounds of the general formula (I). Normally, temperatures of -10° to $+25^\circ\text{C}$, inclusive, are suitable. Also protection from light is advantageous to prevent the corresponding photolytic degradation.

The process may be carried out in an inert solvent medium which, depending on the choice of nitrosating agent, may be a chlorinated hydrocarbon, e.g., carbon tetrachloride, methylene chloride, chloroform; dimethyl formamide, acetic anhydride, pyridine, an alcohol, e.g., ethanol, water; and in nitrosation with a metal nitrite the acid used in this conjunction may also function as the solvent medium, either alone or in a mixture with the above inert solvent medium. Mixtures of more than one solvent may also be employed.

Some nitrosating agents, e.g., nitrosyl chloride, dinitrogen tetroxide and nitrosylsulphuric acid, release a strong acid in the reaction with a urea (II), and in such cases a suitable base, e.g., potassium acetate, may be used in the process to neutralize such formed strong acid. This may similarly also be accomplished by using e.g., pyridine as both solvent and base.

Nitrosation of a urea (II) with an alkyl nitrite is performed in the presence of a suitable nucleophilic anionic catalyst, e.g., chloride or isothiocyanat anions.

Some of the foregoing methods for preparation of compounds of the general formula (I) are summarized in reference (10).

Method 1 above is further illustrated in Examples 1—3, inclusive.

B. A process according to Method 2, characterized by reacting an amine (III) with an N - alkyl - N - nitroso carbamoyl derivative (IV), wherein the group L of (IV) is a suitable leaving group.

Among such leaving groups, L, the following may be mentioned: Azido (N_3-); substituted phenoxy, i.e. ortho- or para-nitrophenoxy, ortho- or para-cyanophenoxy, polyhalophenoxy such as 2,4,5-trichloro-, penta-chloro-, or pentafluorophenoxy; and 1-(2,5-pyrrolidinedione)oxy-.

Although the process may be performed at ambient temperatures, subambient temperatures, such as around 0°C , may be employed and may also be advantageous to prevent thermal degradation of the compounds of the general formulas (I) and (IV). Normally, temperatures of -10° to $+25^\circ\text{C}$, inclusive, are suitable. Also, protection from light is advantageous to prevent the corresponding photolytic degradation.

Suitable inert solvents which may be employed in the process are, e.g., dimethyl formamide, chlorinated hydrocarbons such as methylene chloride, alcohols such as methanol, ethanol, n- and iso-propanol, and pyridine.

Some of the foregoing procedures for preparation of compounds of the general formula (I) are given in reference (11) and references given therein.

Method 2 above is further illustrated in Examples 4—8, inclusive.

Methods of preparation of intermediates

a) The compounds of the general formula (II), which are useful in Method 1 above as starting materials

for the preparation of the compounds of the general formula (I) are prepared by known methods, for example:

Methods a-1 and a-2 below:

5 a-1) A compound of the general formula (III) is reacted with an alkyl isocyanate to give a compound of the general formula (II), e.g. according to reference (12). This method is further illustrated in Example 9.

a-2) A compound of the general formula (III) is reacted with phosgene to give the corresponding isocyanate or carbamoyl chloride (e.g. according to reference [18]), followed by reaction with a primary amine as in the above method to give a compound of the general formula (II).

10 This two step procedure is further illustrated in Example 10.

b) The compounds of the general formula (III), which are useful as starting materials in Method 2 above for the preparation of the compounds of the general formula (I), are prepared by known methods, for example methods b-1, b-2, b-3 and b-4 below:

b-1) An aminoalkylsulfonic acid having the general formula



wherein R and B are as defined above, is converted to its N-protected derivative, e.g. the N-phtaloyl derivative (B=hydrogen), the N-benzyloxycarbonyl derivative, or the N - [2,2,2 - trichloroethoxy-carbonyl]derivative, converted with phosphorous pentachloride to the corresponding chlorosulfonyl-derivative, reacted with an amine of the general formula



wherein R¹ and R² are as defined above, and the N-protecting group is removed to give a compound of the general formula (III), e.g. according to reference (19) and (20).

This method is further illustrated in Example 11.

35 b-2) A bis(aminoalkyl)disulfide having the general formula



wherein R and B are as defined above, is converted to its N-protected derivative, e.g. the N-phtaloyl derivative (B=hydrogen), the N-benzyloxycarbonyl derivative, or the N - [2,2,2 - trichloroethoxycarbonyl]derivative, oxidized e.g. with chlorine to the corresponding chlorosulfonyl derivative, reacted with an amine of the general formula




wherein R¹ and R² are as defined above, and the N-protecting group is removed to give a compound of the general formula (III), e.g. according to reference (21).

b-3) A dihaloalkene, e.g. a dibromoalkene having the general formula



wherein R has the same meaning as defined above, is reacted with phtalimide to a bromoalkylphtalimide, then reacted with thiourea to a S-phtalimidoalkyl thiuronium bromide, oxidized with e.g. chlorine to a phtalimidoalkylsulfonyl chloride, reacted with an amine of the general formula



wherein R^1 and R^2 are as defined above, and the phthalimido group removed to give a compound of the general formula (III), e.g. according to reference (22).

This method is further illustrated in Example 12.

b-4) Other routes for the preparation of compounds of the general formula (III) are also known, as described e.g. in reference (9).

c) The compounds of the general formula (IV), which are useful in Method 2 above as starting materials for the preparation of the compounds of the general formula (I) are prepared by known methods, e.g. according to references (10), (11), (12) and references given therein.

In synthesizing the compounds having the general formulas (II), (III) and (IV), each group of compounds involved must be compatible with the process in question or, if necessary, protected during one or more reaction steps and then converted to the desired group. Pertinent examples of groups that may be protected are hydroxy, carboxyl, sulfonic acid, primary or secondary amino groups. Examples of such protecting groups are found in, e.g., references (13, 14, 15).

The compounds of the invention are generally characterized by the pharmacological activity hereinbefore stated, making them useful in counteracting certain physiological abnormalities in a living animal body. Effective quantities of the pharmacologically active compounds of the invention may be administered to a living animal body in any one of various ways, for example, orally as in capsules or tablets, parenterally in the form of sterile solutions, suspensions, and by pellet implantation. Among routes of parenteral administration are intravenously, subcutaneously, intramuscularly, intraperitoneally, intraarterially, and intradermally. Other modes of administration are vaginally, rectally, and topically, as e.g., in the form of ointments, suppositories, and powders.

As representative of living animal bodies, which may be treated with the compounds and compositions of the invention, and according to the method of treatment of the invention, for alleviation of the same and/or similar conditions as those described, in addition to homo sapiens the following may be mentioned: domestic animals such as dogs and cats, and farm animals such as horses, cows, sheep, and goats.

Pharmaceutical formulations are usually prepared from a predetermined quantity of one or more of the compounds of the invention. Such formulations may take the form of powders, syrups, suppositories, ointments, solutions, pills, capsules, pellets or tablets, suspensions, emulsions, oil solutions, etc, with or without, but preferably with, any one of a large variety of pharmaceutically acceptable vehicles or carriers. When in a mixture with a pharmaceutical vehicle or carrier, the active ingredient usually comprises from about 0.01 to about 75 per cent, normally from about 0.05 to about 15 per cent, by weight of the composition. Carriers such as cellulose, sugar, talc, commonly used synthetic and natural gums, water, and the like, may be used in such formulations. Binders such as polyvinylpyrrolidone and lubricants such as sodium stearate, may be used to form tablets. Disintegrating agents such as starch may also be included in tablets.

Although relatively small quantities of the active materials of the invention, even as low as 5.0 milligrams, may be used in cases of administration to subjects having a relatively low body weight, unit dosages are preferably five milligrams or above and preferably ten, twenty-five, or fifty, or even higher, depending of course upon the subject treated and the particular result desired, as will be apparent to one skilled in the art. Broader ranges appear to be 1 to 500 milligrams per unit dose.

The active agents of the invention may be combined for administration with other pharmaceutically active agents such as analgesics, steroids or hormones, or the like, or with buffers, antacids, or the like, and the proportion of the active agent or agents in the compositions may be varied widely. It is only necessary that the active ingredient of the invention constitutes an effective amount, i.e., such that a suitable effective dosage will be obtained consistent with the dosage form employed. Obviously, several unit dosage forms may be administered at about the same time. The exact individual dosage as well as daily dosages in a particular case will of course be determined according to well established medical and/or veterinary principles under the supervision of the physician or veterinarian in charge. As a rule, however, when used therapeutically, the present compounds may be administered in a quantity of 1 to 500 milligrams, preferred ranges being 5—100 milligrams per day per subject or patient, divided into 1 to 4 or more doses, over a suitable period and depending upon the subject and the type of subject being treated.

Detailed description of the invention

The following examples are intended to illustrate but not to limit the scope of the invention, although the compounds named are of particular interest for our intended purposes. These compounds have been designated by underlined numbers in the examples where their systematic names are given. The compounds are later referred to by a number code, a:b, where a means the number of the example wherein the preparation of the compound in question is described, and b refers to the order of the compounds prepared according to that example. Thus, compound 1:2 means the second compound prepared according to Example 1.

The structures of the compounds found in Examples 1—12 are confirmed the NMR and elementary analysis. The NMR data are obtained using a 60 MHz instrument (Perkin Elmer R 12).

Example 1

1 - (2 - chloroethyl) - 3 - [2 - (dimethylaminosulfonyl)ethyl]urea (54.2 g, 0.02 mol) is dissolved in a mixture of glacial acetic acid (20 ml) and acetic anhydride (100 ml). The solution is cooled to 0°C and sodium nitrite (27.6 g, 0.4 mol) is added while stirring over a period of 2 hours. The reaction mixture is maintained at 0°C for 10 hours and then added to a mixture of ice and water. The reaction product is extracted with ether, which is then washed with water, 5% aqueous sodium carbonate solution, and water. After drying over anhydrous sodium sulphate, the ether is evaporated *in vacuo*.

The product, 1 - (2 - chloroethyl) - 3 - [2 - (dimethylaminosulfonyl)ethyl] - 1 - nitroso-urea (1), is isolated and purified by preparative HPLC, is pure by TLC and has a melting point of 65—67.5°C.

In essentially the same manner the following compounds are obtained from the corresponding starting materials.

2. 1-(2-chloroethyl)-3-(2-[diethylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 68°C.
3. 3-(2-[aminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea, m.p. 115—118°C.
- 15 4. 1-(2-chloroethyl)-3-(2-[1-piperidinylaminosulfonyl]ethyl)-urea, m.p. 83°C.
5. 1-(2-chloroethyl)-3-(2-[methylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 84—86°C.
6. 1-(2-chloroethyl)-3-(2-[dipropylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 69°C.
7. 3-(2-[benzylaminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea, m.p. 76—78°C.
8. 1-(2-chloroethyl)-3-(2-[2-methoxyethylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. about 15°C.
- 20 9. 3-(2-[butylaminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea, m.p. 55—56°C.
10. 3-(2-[t-butylaminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea
11. 1-(2-chloroethyl)-3-(2-[cyclohexylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 85—87°C.
12. 3-(2-[bis(2-hydroxyethyl)aminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea
13. 1-(2-chloroethyl)-3-(2-[1-methylpentylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 24—26°C.
- 25 14. 1-(2-chloroethyl)-3-(2-[2-hydroxyethylaminosulfonyl]ethyl)-1-nitroso-urea
15. 1-(2-chloroethyl)-3-(2-[N-2-hydroxyethyl-N-methyl-aminosulfonyl]ethyl)-1-nitroso-urea, m.p. about 15°C.
16. 1-(2-chloroethyl)-3-(2-[1,1-dimethyl-2-hydroxyethylaminosulfonyl]ethyl)-1-nitroso-urea
17. 3-(2-[N-benzyl-N-methylaminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea, m.p. 74—75°C.
- 30 18. 1-(2-chloroethyl)-3-(2-[N-methyl-N-phenylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 83°C.
19. 1-(2-chloroethyl)-3-(5-[dimethylaminosulfonyl]pentyl)-1-nitroso-urea, m.p. 59—60°C.
20. 1-(2-chloroethyl)-3-(2-[N-methoxy-N-methylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 55°C.
21. 1-(2-chloroethyl)-3-(3-[dimethylaminosulfonyl]propyl)-1-nitroso-urea, m.p. 99°C.
22. 1-(2-chloroethyl)-3-(2-[4-methylbenzylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 70°C.
- 35 23. 1-(2-chloroethyl)-3-(2-[2-(dimethylaminosulfonyl)ethylaminosulfonyl]ethyl)-1-nitroso-urea, m.p. 70—72°C.
24. 1-(2-chloroethyl)-3-(4-[dimethylaminosulfonyl]butyl)-1-nitroso-urea, m.p. 83°C.
25. 3-(2-[dimethylaminosulfonyl]ethyl)-1-(2-fluoroethyl)-1-nitroso-urea
26. 3-(2-[dimethylaminosulfonyl]ethyl)-1-methyl-1-nitroso-urea
- 40 27. 1-(2-bromoethyl)-3-(2-[dimethylaminosulfonyl]ethyl)-1-nitroso-urea
28. 1-(2-chloroethyl)-3-(2-[N-cyclohexyl-N-methylaminosulfonyl]ethyl)-1-nitroso-urea
29. 1-(2-chloroethyl)-3-(2-[cyclopentylaminosulfonyl]ethyl)-1-nitroso-urea
30. 3-(2-[2-(aminosulfonyl)ethylaminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea
31. 1-(2-chloroethyl)-3-(2-[2-(methylaminosulfonyl)ethylaminosulfonyl]ethyl)-1-nitroso-urea
- 45 32. 3-(2-[2-(aminocarbonyl)ethylaminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea
33. 1-(2-chloroethyl)-3-(2-[2-(methylaminocarbonyl)ethylaminosulfonyl]ethyl)-1-nitroso-urea
34. 1-(2-chloroethyl)-3-(2-[2-(dimethylaminocarbonyl)ethylaminosulfonyl]ethyl)-1-nitroso-urea
35. 1-(2-chloroethyl)-3-(1-[dimethylaminocarbonyl]-2-[dimethylaminosulfonyl]ethyl)-1-nitroso-urea
36. 1-(2-chloroethyl)-3-(2-[dimethylaminosulfonyl]propyl)-1-nitroso-urea
- 50 37. 3-(bis-2,3-[dimethylaminosulfonyl]propyl)-1-(2-chloroethyl)-1-nitroso-urea
38. 3-(bis-1,3-[dimethylaminosulfonyl]propan-2-yl)-1-(2-chloroethyl)-1-nitroso-urea
39. 1-(2-chloroethyl)-3-(1-[dimethylaminocarbonyl]-3-[dimethylaminosulfonyl]propan-2-yl)-1-nitroso-urea
40. 1-(2-chloroethyl)-3-(1-[dimethylaminosulfonyl]-2-methylpropan-2-yl)-1-nitroso-urea
- 55 41. 1-(2-chloroethyl)-3-(1-[dimethylaminosulfonyl]propan-2-yl)-1-nitroso-urea
42. 1-(2-chloroethyl)-3-(1-[dimethylaminosulfonyl]butan-2-yl)-1-nitroso-urea
43. 3-(bis-1,3-[aminosulfonyl]propan-2-yl)-1-(2-chloroethyl)-1-nitroso-urea
44. 3-(1-[aminocarbonyl]-2-[aminosulfonyl]ethyl)-1-(2-chloroethyl)-1-nitroso-urea
45. 1-(2-chloroethyl)-3-(2-[dimethylaminosulfonyl]ethyl)-3-methyl-1-nitroso-urea
- 60 46. 3-allyl-1-(2-chloroethyl)-3-(2-[dimethylaminosulfonyl]ethyl)-1-nitroso-urea
47. 1-(2-chloroethyl)-3-propargyl-3-(2-[dimethylaminosulfonyl]ethyl)-1-nitroso-urea
48. 1-(2-chloroethyl)-3-(2-methoxyethyl)-3-(2-[dimethylaminosulfonyl]ethyl)-1-nitroso-urea
49. 1-(2-chloroethyl)-3-cyclohexyl-3-(2-[dimethylaminosulfonyl]ethyl)-1-nitroso-urea
50. 1-(2-chloroethyl)-3-cyclopropylmethyl-3-(2-[dimethylaminosulfonyl]ethyl)-1-nitroso-urea
- 65 51. 3,3-bis-[2-(dimethylaminosulfonyl)ethyl]-1-nitroso-urea

52. 1-(2-chloroethyl)-3-(2-[N-ethyl-N-methylaminosulfonyl]ethyl)-1-nitrosoourea, m.p. ~25°C

53. 1-(2-chloroethyl)-3-(2-[N-ethyl-N-propylaminosulfonyl]ethyl)-1-nitrosoourea, m.p. ~20°C

54. 1-(2-chloroethyl)-3-(2-[dimethylaminosulfonyl]ethyl)-3-ethyl-1-nitrosoourea.

5 Example 2

To a cooled (-60°C) solution of dinitrogen tetroxide (4.14 g, 0.045 mol) in glacial acetic acid (50 ml) anhydrous sodium acetate (7.38 g, 0.09 mol) is added. The temperature is raised to 0°C, and 1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl)urea (7.73 g, 0.03 mol) is added while stirring. After continued stirring at 0°C for 30 minutes the mixture is added to a mix of ice and water. The reaction product
10 is extracted with ether, which is then washed with water, 5% aqueous sodium carbonate solution, and water. After drying over anhydrous sodium sulphate the ether is evaporated *in vacuo*. The product, 1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl) - 1 - nitrosoourea (the same compound as 1:1), is isolated and purified by preparative HPLC, is pure by TLC and has a melting point of 65—67.5°C.

15 Example 3

1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl)urea (5.16 g, 0.02 mol) is dissolved in a mixture of glacial acetic acid (100 ml) and acetic anhydride (40 ml), and freshly fused potassium acetate (16 g) and phosphorus pentoxide (4 g) is added. The resulting mixture is cooled to +5°C, and a solution of nitrosyl chloride (5 g) in acetic anhydride (25 ml) is added dropwise while stirring. After the addition,
20 stirring is continued for 2 hours at +10°C. The reaction mixture is added to a mix of ice and water, and the reaction product is extracted with diethyl ether which is then washed with water, 5% aqueous sodium carbonate solution and water. After drying over anhydrous sodium sulphate the diethyl ether is evaporated *in vacuo*. The product, 1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl) - 1 - nitrosoourea (the same compound as 1:1), is isolated and purified by preparative HPLC, is pure by TLC, and has a melting
25 point of 65—67.5°C.

Example 4

A solution of 2 - amino - N,N - dimethyl - ethanesulfonamide hydrochloride (4.72, 0.025 mol) in abs. ethanol (30 ml) is cooled to 0°C. Triethyl amine (3.5 ml) is added and the mixture is added dropwise to a
30 solution of N - (2 - chloroethyl) - N - nitroso carbamoyl azide (5.4 g, 0.03 mol). The reaction mixture is stirred for 2 hours at 0°C and is then evaporated to dryness *in vacuo*. Diethyl ether is added to the residue, and the precipitated triethyl amine hydrochloride is filtered off. The diethyl ether solution is evaporated *in vacuo* to dryness. The product, 1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl) - 1 - nitrosoourea (the same compound as 1:1), is isolated and purified by preparative HPLC, is pure by TLC and
35 has a melting point of 65—67.5°C.

In essentially the same manner compounds 1:2 to 1:44, inclusive, and 1:54 are obtained from the corresponding starting materials.

Example 5

40 To a solution of 2 - amino - N,N - dimethyl - ethanesulfonamide hydrochloride (5 g, 0.026 mol) in methanol (60 ml) is added triethyl amine (2.68 g). The mixture is stirred for 10 minutes, and p-nitrophenyl N - (2 - chloroethyl) - N - nitroso carbamate (7.25 g, 0.026 mol) is added. After stirring at room temperature for 2 hours the mixture is evaporated *in vacuo* to dryness, and methylene chloride (50 ml) and triethyl amine (10 ml) is added. The resulting solution is washed twice with water (50 ml), twice with 2 M
45 hydrochloric acid (40 ml), and with water. After drying over anhydrous sodium sulphate the solution is evaporated to dryness *in vacuo*. The product, 1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl) - 1 - nitrosoourea (the same compound as 1:1), is isolated and purified by preparative HPLC, is pure by TLC and has a melting point of 65—76.5°C.

50 In essentially the same manner compounds 1:2 to 1:44, inclusive, and 1:54 are obtained from the corresponding starting materials.

1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl) - 1 - nitrosoourea (the same compound as 1:1) is also prepared essentially in the same manner from the same above mentioned amine and o-nitrophenyl N - (2 - chloroethyl) - N - nitroso carbamate or o-cyanophenyl N - (2 - chloroethyl) - N -
55 nitroso carbamate.

Example 6

To a solution of 2-aminoethanesulfonamide hydrochloride (5 g, 0.030 mol) in methanol (70 ml) triethylamine (3.13 g) is added. The mixture is stirred for 10 minutes and p-nitrophenyl N - (2 - chloroethyl) - N - nitroso carbamate (8.75 g, 0.032 mol) is added. After stirring at room temperature for 2
60 hours the mixture is evaporated *in vacuo* to dryness, and the residue is extracted with methylene chloride (30 ml). The undissolved crystalline material is filtered off and washed with methylene chloride. The product obtained, 3 - (2 - [aminosulfonyl]ethyl) - 1 - (2 - chloroethyl) - 1 - nitrosoourea (the same compound as 1:3), is pure in TLC without further chromatographic purification and is dried *in vacuo* over
65 phosphorus pentoxide. The melting point of the product obtained is 115—118°C.

Example 7

To a solution of 2 - amino - N,N - dimethyl - ethanesulfonamide hydrochloride (5 g, 0.026 mol) in methanol (60 ml) triethyl amine (2.68 g) is added. The mixture is stirred for 10 minutes and 1 - [N - (2 - chloroethyl)N - nitrosoaminocarbonyloxy] - 2,5 - pyrrolidine - dione (6.94 g, 0.028 mol) is added. After stirring at room temperature for 2 hours the reaction mixture is evaporated *in vacuo* to dryness and the residue is dissolved in methylene chloride. The solution is washed with water (4×50 ml) and dried over sodium sulphate. After filtration the dry solution is evaporated *in vacuo* to dryness. The product, 1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl) - 1 - nitrosoarea (the same compound as 1:1), is isolated and purified by preparative HPLC, is pure in TLC and has a melting point of 65—67.5°C.

Example 8

To a solution of 2 - aminoethanesulfonamide hydrochloride (5 g, 0.030 mol) in methanol (70 ml) triethyl amine (3.13 g) is added. The mixture is stirred for 10 minutes and p - cyanophenyl - N - (2 - chloroethyl) - N - nitroso carbamate (8.11 g, 0.032 mol) is added. After stirring at room temperature for 3 hours the reaction mixture is evaporated *in vacuo* to dryness and the residue is extracted with cold methylene chloride. The undissolved crystalline material is filtered off and washed with cold methylene chloride. The product obtained, 3 - (2 - [aminosulfonyl]ethyl) - 1 - (2 - chloroethyl) - 1 - nitrosoarea (the same compound as 1:3), is pure in TLC without further chromatographic purification, and is dried *in vacuo* over phosphorus pentoxide. The melting point of the product obtained is 115—118°C.

Preparation of intermediates (Examples 9—12)

Example 9

2 - amino - N,N - dimethylethanesulfonamide hydrochloride (5.64 g, 0.03 mol) is dissolved in chloroform (50 ml) while adding triethylamine (4.7 ml). The solution is cooled to 0°C and a mixture of 2-chloroethyl isocyanate (2.83 ml, 0.04 mol) and chloroform (25 ml) is added dropwise. The reaction mixture is stirred for 4 hours at room temperature and washed with water (3×50 ml). The organic phase is dried over anhydrous sodium sulphate, filtered and evaporated *in vacuo* to dryness. The product obtained is 1 - (2 - chloroethyl) - 3 - (2 - [dimethylaminosulfonyl]ethyl) - urea, m.p. 126°C.

In essentially the same manner other compounds of the general formula (II) above are obtained from the corresponding compounds of the general formula (III) by reaction with the corresponding isocyanate.

Example 10

2 - amino - N,N - dimethylethanesulfonamide (4.72 g, 0.025 mol) is dissolved in chloroform (50 ml) while adding triethyl amine (3.8 ml, 0.027 mol). The solution is cooled to 0°C and triethyl amine (3.8 ml, 0.027 mol) is added. A solution of phosgene (3 g, 0.03 mol) in chloroform (25 ml) is added dropwise while stirring. The reaction mixture is stirred for about 1 hour. A solution of 2-fluoroethyl amine hydrochloride (3.0 g, 0.03 mol), triethyl amine (4 ml, 0.03 mol) in chloroform (50 ml) is added dropwise while stirring and cooling. The reaction mixture is stirred at room temperature for 1 hour and then washed with 0.1 M hydrochloric acid (50 ml) and water (2×50 ml). The organic phase is dried and evaporated *in vacuo* to dryness. The product obtained is 3 - (2 - [dimethylaminosulfonyl]ethyl) - 1 - (2 - fluoroethyl) - urea.

In essentially the same manner the following compound is obtained from the corresponding starting materials.

2. 1-(2-bromoethyl)-3-(2-[dimethylaminosulfonyl]ethyl)-urea.

Example 11

A mixture of taurine (100 g, 0.8 mol), potassium acetate (83.8 g, 1.4 mol) and glacial acetic acid (283 ml) is boiled under stirring and reflux for 10 minutes.

To the boiling mixture is added phthalic acid anhydride (126.2 g, 0.85 mol) and the reaction mixture is boiled for another 2.5 hrs.

After cooling to room temperature the mixture is filtered and the solid is washed successively with glacial acetic acid and ethanol.

The product, the potassium salt of 2-phthalimidoethanesulfonic acid, is dried *in vacuo*. It is characterized by NMR, and elementary analysis.

The above product (132 g, 0.45 mol) is suspended in 660 ml of benzene. In order to remove traces of moisture 150 ml of the benzene is distilled off. The mixture is cooled to room temperature, whereupon phosphorus pentachloride (67.5 g, 0.32 mol) is added while stirring. The mixture is then refluxed for 1 hr and a further amount (67.5 g, 0.32 mol) of phosphorus pentachloride is added. The stirring and heating is continued for another 1.5 hrs, whereupon the solvent is removed *in vacuo*. Another 200 ml of benzene are added and the mixture is again evaporated to dryness. The residue is mixed with 900 g of crushed ice and left overnight. The solid, 2-phthalimidoethanesulfonyl chloride, is filtered off, washed with water and dried *in vacuo*. The product is characterized by NMR, and has a melting point of 159—162°C.

The above product (58.7 g, 0.215 mol) is dissolved in 400 ml of toluene. A solution of dimethylamine (32.3 ml, 0.43 mol) in 80 ml of toluene is added with cooling and stirring.

The reaction mixture is stirred at room temperature for 2 hrs, whereupon 250 ml of water are added

slowly. The precipitated compound, 2-phthalimidoethylsulfonyl dimethylamide, is washed with water and dried *in vacuo*. The product is pure in TLC, is characterized by NMR, and has a melting point of 154—156°C.

The above product (55 g, 0.195 mol) is suspended in 360 ml of 95% ethanol whereupon hydrazine hydrate (9.8 g, 0.195 mol) is added. The reaction mixture is boiled under reflux for 3 hrs.

5 The precipitated product is filtered off and the filtrate is evaporated to dryness. The residue is combined with the filtered product and the mixture is slurried in 180 ml of hot water, pH is adjusted to about 3 and the mixture left overnight in the refrigerator.

The mixture is filtered and the filtrate is evaporated to dryness. The residue, 2 - aminoethylsulfonyl dimethylamide hydrochloride, is recrystallised from 240 ml of isopropanol.

10 The product is pure in TLC, is characterized by NMR and quantitative analysis, and has a melting point of 144—146°C.

In essentially the same manner other compounds of the general formula (III) above are obtained from the corresponding aminoalkylsulfonic acid.

15 Example 12

A mixture of phthalimide (100 g, 0.68 mol), potassium carbonate (50 g, 0.36 mol) and trimethylenedibromide (350 g, 1.73 mol.) is gradually heated until reaction begins. The reaction is carried out in a vessel equipped with an efficient double mantled condenser.

20 When a clear reaction mixture is obtained, the heating is maintained for 2 hrs whereupon excess of trimethylenedibromide is distilled off by means of steam distillation. The residue is recrystallised from 100 ml of ethanol.

The crystallised product is extracted with petroleum ether. The extract is evaporated to dryness, leaving the 3-bromopropylphthalimide (m.p. 72—73°C) as a white crystalline solid.

25 A mixture of the above product (103 g, 0.38 mol.), thiourea (29.3 g, 0.38 mol.) and 200 ml of 95% aqueous ethanol is boiled under reflux for 8 hrs.

On cooling, the S-3-phthalimidopropyl thiuronium bromide precipitates and is filtered off and washed with ethanol. Recrystallisation from water gives a pure product (m.p. 225—227°C).

To a hot aqueous solution of the above product (50 g, 0.15 mol.) is added a solution of potassium acetate (56 g., 0.57 mol.) in 50 ml of hot water. The mixture is stirred and kept at 80°C for 1 hr.

30 After cooling, the crystallised thiuronium acetate is filtered off and then suspended in a mixture of 340 ml of water and 4.5 g of concentrated hydrochloric acid. On cooling (0—5°C) chlorine is passed into the solution while stirring. After about 3 hrs. the 3-phthalimidopropane-1-sulfonyl chloride is filtered off and washed with water. The product is purified by dissolving it in hot benzene, filtering the solution and precipitating the product by addition of light petroleum.

35 The product has a m.p. of 77—78°C (when dried *in vacuo* over H₂SO₄).

The above product (61.9 g, 0.215 mol.) is dissolved in 400 ml of toluene. A solution of dimethylamine (32.3 ml, 0.43 mol.) in 80 ml of toluene is added with cooling and stirring.

40 The reaction mixture is stirred at room temperature for 2 hrs, whereupon 250 ml of water are added slowly. The precipitated compound, 3 - phthalimidopropane - 1 - sulfonyldimethylamide is washed with water and dried *in vacuo*. The product is pure in TLC and is characterized by NMR.

The above product (55 g, 0.195 mol.) is suspended in 360 ml of 95% ethanol whereupon hydrazine hydrate (9.75 g, 0.195 mol.) is added. The reaction mixture is boiled under reflux for 3 hrs.

45 The precipitated product is filtered off and the filtrate is evaporated to dryness. The residue is combined with the filtered product and the mixture is slurried in 180 ml of hot water, pH is adjusted to about 3 and the mixture left overnight in the refrigerator.

The mixture is filtered and the filtrate is evaporated to dryness. The residue, crude 3 - aminopropyl - 1 - sulfonyldimethylamide hydrochloride is recrystallised from 240 ml of isopropanol and dried *in vacuo*.

The product is pure in TLC and is characterized by NMR and quantitative analysis.

50 Example 13

This example illustrates the effect of compounds of the present invention in inhibiting the growth of several tumours. From the results obtained it is obvious that the compounds of the present invention have improved toxicological and therapeutic properties, resulting in increased therapeutic indexes, in comparison with commercially available antitumour agents.

55 A therapeutic index is an indication of the therapeutic usefulness of a compound, and such an index may be defined in various ways. One type of therapeutic index commonly used is the ratio LD50/ED50, wherein LD50 is the dose that causes a 50 per cent lethality, and ED50 is the dose that causes a 50 per cent reduction of tumour size. Another type of therapeutic index, useful in evaluation of increasing survival time, is the ratio between maximum and minimum effective dose. Both these types of therapeutic indexes

60 are used and given in the experimental results below.

The experimental design and interpretation of the results are in accordance with the standards set by the CCNSC (see reference (16)) and by the DR & D (see reference (17)).

The manner of administration of the compounds is by the oral (p. o.) route in each case.

Some of the results obtained are given in Table 1—2.3 below.

65 This example shows that the new compounds are useful to prevent the growth of tumours and can

therefore be employed in treating a living animal body suffering from disorders responsive to treatment with anti-cancer agents and immunosuppressive agents.

Walker carcinosarcoma 256

- 5 Experimental animals:
Female Sprague Dawley rats

Tumour implants:

- 10 Tumour pieces with 2—3 mm diameter, implanted subcutaneously

Therapy:

The compound is administered p.o. once on the day after the implantation

Termination:

- 15 The animals are killed on the 9th day after implantation

Evaluation:

- Weight of tumour is determined in test animals and compared with those of control animals. For each compound the LD50 and ED50 dose (as defined above) are determined. The therapeutic index, T.I., (LD50/ED50) is calculated.

TABLE 1
Antitumour results on walker carcinosarcoma 256

Compound	LD50 mg/kg	ED50 mg/kg	T.I. LD50/ED50
BCNU	50	1	50
CCNU	90	1	90
Cyclophosphamide	180	6	30
1:1	50	0.25	200
1:2	60	0.4	150
1:3	40	0.2	200
1:4	125	1	125
1:5	40	0.25	160
1:6	250	2	125
1:7	250	2	125
1:14	90	0.6	150
1:15	90	0.6	150
1:19	250	1	250
1:24	250	2	125
1:28	>250	2.5	>100
1:53	250	1.7	147

In preliminary experiments the following additional compounds are found to exhibit significant activity in the foregoing test in a dose of 8 mg/kg: 1:8 to 1:13, inclusive; 1:16 to 1:18, inclusive; 1:20 to 1:23, inclusive; 1:25 to 1:45, inclusive; 1:48, 1:52 and 1:54.

Lymphatic leukemia L1210

Experimental animals:
Female BDF₁ mice

- 5 Tumour implants:
10⁵ tumour cells implanted intraperitoneally (i.p.)

Therapy:

The compounds were administered p.o. once on the day following the implantation.

10

Termination:

The animals were killed on the 60th day after the implantation. Living animals on the 60th day are termed long term survivors.

15 Evaluation:

a) Evaluation of therapeutic dose range for survival time increase: Survival times at various dose levels are determined, and the survival time of the test animals (t) is expressed as the percentage of that of the control animals (c):

20

$$t \cdot 100 / c (\%)$$

The maximum effective dose and the minimum effective dose required to achieve $t \cdot 100 / c \geq 125\%$ are determined, and a therapeutic index as defined below is calculated:

25

$$T.I._s = \text{max. effect. dose} / \text{min. effect. dose.}$$

Some of the results obtained are shown in Table 2:1.

b) Evaluation of therapeutic dose range for long term survival: Long time survival is determined at various dose levels and is expressed as surviving test animals on day 60 (T) as the percentage of that of the total number of test animals (T₀):

30

$$T \cdot 100 / T_0 (\%)$$

The maximum effective dose and the minimum effective dose to achieve long term survival $T \cdot 100 / T_0 > 20\%$ are determined and a therapeutic index, T.I._{1s} is calculated:

35

$$T.I._{1s} = \text{max. effect. dose} / \text{min. effect. dose.}$$

Some of the results obtained are shown in Table 2:2.

c) Evaluation of cure potential: The normal survival time for test animals receiving no therapeutic treatment is 9—11 days. Surviving test animals on the 60th day after implantation showing no sign of tumour presence are regarded as cured animals. Long term survival is determined as under b) above, and the effective dose or dose range to achieve a long term survival $T \cdot 100 / T_0 \geq 90\%$ is determined as an evaluation of the cure potential. Some of the results obtained are shown in Table 2:3.

45

50

55

60

65

TABLE 2:1
Evaluation of therapeutic range for survival time increase

Compound	Maximum effective dose ^{x)} mg/kg	Minimum effective dose ^{x)} mg/kg	T.I.s max. eff. dose min. eff. dose
BCNU	63	16	4
Methyl-CCNU	31	8	4
1:1	63	8	8
1:2	125	16	8
1:4	125	8	16
1:5	63	8	8
1:6	250	31	8
1:9	125	16	8
1:11	250	31	8
1:14	125	16	8
1:15	125	16	8
1:17	500	63	8
1:20	125	16	8
1:21	63	8	8
1:24	250	31	8
1:28	250	31	8
1:53	500	31	16

^{x)} Median survival time per cent of control $\geq 125\%$

The following additional compounds are found to exhibit significant activity (i.e. median survival time per cent of control $\geq 125\%$) in a dose of 125 mg/kg in the foregoing test: 1:10; 1:13; 1:18; 1:25—1:27, inclusive; 1:29—1:45, inclusive; and 1:54.

TABLE 2:2
Evaluation of therapeutic range for long term survival

Compound	Maximum effective dose ^{x)} mg/kg	Minimum effective dose ^{x)} mg/kg	T.I.s max. eff. dose min. eff. dose
BCNU	63	63	1
Methyl CCNU	none	none	none
Chlorozotocin	"	"	"
1:1	63	31	2
1:2	125	63	2
1:3	31	16	2
1:4	125	31	4
1:7	250	125	2
1:8	63	31	2
1:9	125	63	2
1:11	125	63	2
1:14	125	31	4
1:15	125	31	4
1:17	250	125	2
1:19	125	63	2
1:20	125	63	2
1:21	63	31	2
1:23	125	63	2
1:24	250	63	4
1:28	250	125	2
1:52	63	31	2
1:53	250	125	2

^{x)} Dose levels to achieve >20% long term survivors, i.e. test animal survivors on day 60.

TABLE 2:3
Evaluation of cure potential

5

10

15

20

25

30

35

40

45

50

55

60

65

Compound	Dose or dose range for cure ^{*)} mg/kg
BCNU	none
Methyl CCNU	"
Chlorozotocin	"
1:1	31—63
1:4	125
1:5	31
1:6	125
1:8	31—63
1:9	63
1:11	125
1:12	125
1:14	63
1:15	63
1:16	250
1:19	125
1:20	63—125
1:21	31—63
1:23	125
1:24	125
1:28	125
1:53	125—150

^{*)} Cured defined as test with $\geq 90\%$ long term survivors, i.e. test animal survivors on day 60.

Example 14

Manufacturing process for tablets à 20 mg

Model batch of 1000 tablets

5	I	{ Solid active compound, mesh ^{*)} 70	20 g
		{ Lactosum, ph. Nord	210 g
		{ Amylum maidis, ph. Nord.	75 g
		{ Kollidon 25, B.A.S.F.	3.5 g
10	II	Aqua purificata	q.s.
	III	{ Talcum, ph. Nord.	15 g
		{ Magnesii stearas, ph. Nord.	1.5 g
15		Weight of 1000 tablets	325 g

Weight of 1 tablet: 325 mg

^{*)} The mesh standard is according to the international system of code DIN 4189/1968.

20 Punch:

10.5 mm round, flat, scored, bevel-edged

Mix the screened substances I thoroughly and then moisten with II, whereupon the mixture is granulated through a stainless sieve No. 10 (mesh 25). Dry the granulate in an oven at a maximum temperature of 25°C, then repeat sieving through sieve No. 10. Add the substances under III and mix thoroughly. Punch tablets with a gross weight of about 325 mg.

25

Example 15

Suspension for injection 20 mg/ml

30	Active compound, mesh 100	20 mg
	Sodium chloride	8 mg
	Carboxy methylcellulose	1 mg
35	Benzyl alcohol	1 mg
	Distilled water to make	1 ml

40

Example 16

Oral suspension 5 mg/ml

	Active compound, mesh 100	5 mg
45	Sorbitol	600 mg
	Flavouring compound	q.s.
	Colour	q.s.
50	Water to make	1 ml

55

Example 17

Suppositories à 25 mg

	Active compound	25 mg
60	Cocoa butter	q.s.

65

Example 18
Ointment 2%

5	Active compound	2 g
	Triethanolamine	1 g
	Glycerol	7 g
10	Cetanol	2.5 g
	Lanoline	2.5 g
15	Stearic acid	20 g
	Sorbitan monooleate	0.5 g
	Sodium hydroxide	0.2 g
20	Methyl paraben	0.3 g
	Propyl paraben	0.1 g
25	Ethanol	0.9 g
	Water to make	100 g

Example 19
Capsules à 10 mg

30	Active compound	10 mg
	Magnesium stearate	2 mg
35	Talcum	188 mg

The substances are mixed and filled into capsules

Example 20

40	10 mg Sterile powder to be dissolved in water for injection	
	Water-soluble active compound	10 mg
	Sodium chloride	4 mg
45	Methyl paraben	0.7 mg
	Propyl paraben	0.3 mg

50 The substances are dissolved in distilled water
The solution is dispensed in vials and freeze-dried.

In the foregoing Examples 14—20 relating to compositions the Active Compounds are those covered by the general formula (I) above. Those Active Compounds which are disclosed in the foregoing Examples 55 1—8 are preferred as Active Compounds.

Also, it is to be noted that two or more Active Compounds of the invention may be used in combination in the compositions illustrated, and also if desired in combination with other pharmacologically active agents.


60 Various modifications and equivalents will be apparent to one skilled in the art and may be used in the compounds, compositions and methods of the present invention without departing from the spirit or scope thereof, and it is therefore to be understood that the invention is not to be limited to the specific examples and embodiments disclosed herein.

Literature references

1. Carter, Bakowski, Hellman—Chemotherapy of Cancer, Wiley, 1977, p. 70—72
2. Wasserman—Cancer 36 (1975), 1258
3. Schacht—Cancer 48 (1981), 1328
4. Weiss—Cancer Treatment Reviews 8 (1981), 111
5. Schein—Cancer and Chemotherapy, vol. III, Academic Press, 1981, p. 37
6. Belt—Cancer Treatment Reports 64 (1980):12, 1235
7. Ahlgren—Cancer Treatment Reports 65 (1981), 223
8. Sandler, Karo—Functional Group Preparations, vol. 2 (1971), Academic Press, chapter 17
9. Houben-Weyl, vol. 9 (1955), 343
10. Sandler, Karo—Functional Group Preparations, vol. 2 (1971), Academic Press, chapters 10, 11 and 17
11. Martinez—J. Med. Chem. 25 (1982), 178
12. Sandler, Karo—Functional Group Preparations, vol. 2 (1971), Academic Press, chapter 6
13. McOmie—Protective Groups in Organic Chemistry, Plenum Press, 1973
14. Greene—Protective Groups in Organic Synthesis, Wiley, 1981
15. Hoyben-Weyl, vol. 15/1 (1974)
16. Cancer Chemotherapy Reports, December, 1962
17. Cancer Chemotherapy Reports, September, 1972, vol. 3 No. 2
18. Sandler, Karo—Functional Group Preparations, vol. 1 (1968), Academic Press, chapter 12
19. Mead, Koepfli—J. Org. Chem. 12 (1947), 295
20. McIlwain—J. Chem. Soc. 1941, 75
21. Brynes—J. Med. Chem. 21 (1981), 45
22. Griffin, Hey—J. Chem. Soc. 1952, 3334

Claims

1. Novel compounds having the general formula:



wherein A is lower alkyl or halolower alkyl;

B is hydrogen; lower alkenyl; lower alkynyl; 3—6 C cycloalkyl; $-\text{RSO}_2\text{NR}^3\text{R}^4$, or lower alkyl optionally monosubstituted with lower alkoxy or 3—6 C cycloalkyl;

R is straight or branched 2—5 C alkylene, optionally monosubstituted with $\text{SO}_2\text{NR}^3\text{R}^4$ or CONR^3R^4 , said alkylene always containing at least two carbon atoms separating the nitrogen atom of the urea from any $\text{SO}_2\text{NR}^1\text{R}^2$ or $\text{SO}_2\text{NR}^3\text{R}^4$;

R^1 and R^2 are the same or different and selected from hydrogen, 5—6 C cycloalkyl, lower alkoxy, phenyl, benzyl, and straight or branched 1—6 C alkyl, optionally monosubstituted with hydroxy, lower alkoxy, $\text{SO}_2\text{NR}^3\text{R}^4$, or CONR^3R^4 ; R^1 and R^2 may also together form a 4—5 C alkylene;

R^3 and R^4 are the same or different and selected from hydrogen or lower alkyl or R^3 and R^4 together may form a 4—5 C alkylene.

2. Compounds according to claim 1, characterized in that A is chloroethyl or fluoroethyl, preferably chloroethyl.

3. Compounds according to claim 1—2, characterized in that R is unsubstituted 2—5 C alkylene, preferably ethylene.

4. Compounds according to any of the claims 1—3, characterized in that B is hydrogen or lower alkyl, preferably hydrogen.

5. Compounds according to any of the claims 1—4, characterized in that one of the substituents R^1 and R^2 is methyl and the other of the substituents R^1 and R^2 is lower alkyl or 2-hydroxyethyl.

6. Compounds according to any of the claims 1—4, characterized in that R^1 and R^2 are the same and selected from the group hydrogen, methyl, ethyl and propyl, or wherein R^1 and R^2 together are 5 C alkylene.

7. Compounds according to any of the claims 1—4, characterized in that R^1 and R^2 are hydrogen and/or methyl.

8. Compounds according to claim 1 selected from the following group:


- a) 1-(2-chloroethyl)-3-[2-(dimethylaminosulfonyl)ethyl]-1-nitrosourea
- b) 1-(2-chloroethyl)-3-[2-(diethylaminosulfonyl)ethyl]-1-nitrosourea
- c) 3-[2-(aminosulfonyl)ethyl]-1-(2-chloroethyl)-1-nitrosourea
- d) 1-(2-chloroethyl)-1-nitroso-3-[2-(1-piperidinosulfonyl)ethyl]urea
- e) 1-(2-chloroethyl)-3-[2-(methylaminosulfonyl)ethyl]-1-nitrosourea

Literature references

1. Carter, Bakowski, Hellman—Chemotherapy of Cancer, Wiley, 1977, p. 70—72
2. Wasserman—Cancer 36 (1975), 1258
3. Schacht—Cancer 48 (1981), 1328
- 5 4. Weiss—Cancer Treatment Reviews 8 (1981), 111
5. Schein—Cancer and Chemotherapy, vol. III, Academic Press, 1981, p. 37
6. Belt—Cancer Treatment Reports 64 (1980):12, 1235
7. Ahlgren—Cancer Treatment Reports 65 (1981), 223
8. Sandler, Karo—Functional Group Preparations, vol. 2 (1971), Academic Press, chapter 17
- 10 9. Houben-Weyl, vol. 9 (1955), 343
10. Sandler, Karo—Functional Group Preparations, vol. 2 (1971), Academic Press, chapters 10, 11 and 17
11. Martinez—J. Med. Chem. 25 (1982), 178
12. Sandler, Karo—Functional Group Preparations, vol. 2 (1971), Academic Press, chapter 6
- 15 13. McOmie—Protective Groups in Organic Chemistry, Plenum Press, 1973
14. Greene—Protective Groups in Organic Synthesis, Wiley, 1981
15. Houben-Weyl, vol. 15/1 (1974)
16. Cancer Chemotherapy Reports, December, 1962
17. Cancer Chemotherapy Reports, September, 1972, vol. 3 No. 2
- 20 18. Sandler, Karo—Functional Group Preparations, vol. 1 (1968), Academic Press, chapter 12
19. Mead, Koepfli—J. Org. Chem. 12 (1947), 295
20. McIlwain—J. Chem. Soc. 1941, 75
21. Brynes—J. Med. Chem. 21 (1981), 45
22. Griffin, Hey—J. Chem. Soc. 1952, 3334

Claims

1. Novel compounds having the general formula:



wherein A is lower alkyl or halolower alkyl;

B is hydrogen; lower alkenyl; lower alkynyl; 3—6 C cycloalkyl; $-RSO_2NR^3R^4$, or lower alkyl optionally monosubstituted with lower alkoxy or 3—6 C cycloalkyl;

R is straight or branched 2—5 C alkylene, optionally monosubstituted with $SO_2NR^3R^4$ or $CONR^3R^4$, said alkylene always containing at least two carbon atoms separating the nitrogen atom of the urea from any $SO_2NR^1R^2$ or $SO_2NR^3R^4$;

R^1 and R^2 are the same or different and selected from hydrogen, 5—6 C cycloalkyl, lower alkoxy, phenyl, benzyl, and straight or branched 1—6 C alkyl, optionally monosubstituted with hydroxy, lower alkoxy, $SO_2NR^3R^4$, or $CONR^3R^4$; R^1 and R^2 may also together form a 4—5 C alkylene;

R^3 and R^4 are the same or different and selected from hydrogen or lower alkyl or R^3 and R^4 together may form a 4—5 C alkylene.

2. Compounds according to claim 1, characterized in that A is chloroethyl or fluoroethyl, preferably chloroethyl.

3. Compounds according to claim 1—2, characterized in that R is unsubstituted 2—5 C alkylene, preferably ethylene.

4. Compounds according to any of the claims 1—3, characterized in that B is hydrogen or lower alkyl, preferably hydrogen.

5. Compounds according to any of the claims 1—4, characterized in that one of the substituents R^1 and R^2 is methyl and the other of the substituents R^1 and R^2 is lower alkyl or 2-hydroxyethyl.

6. Compounds according to any of the claims 1—4, characterized in that R^1 and R^2 are the same and selected from the group hydrogen, methyl, ethyl and propyl, or wherein R^1 and R^2 together are 5 C alkylene.

7. Compounds according to any of the claims 1—4, characterized in that R^1 and R^2 are hydrogen and/or methyl.


8. Compounds according to claim 1 selected from the following group:

- a) 1-(2-chloroethyl)-3-[2-(dimethylaminosulfonyl)ethyl]-1-nitrosourea
- b) 1-(2-chloroethyl)-3-[2-(diethylaminosulfonyl)ethyl]-1-nitrosourea
- c) 3-[2-(aminosulfonyl)ethyl]-1-(2-chloroethyl)-1-nitrosourea
- d) 1-(2-chloroethyl)-1-nitroso-3-[2-(1-piperidinosulfonyl)ethyl]urea
- 65 e) 1-(2-chloroethyl)-3-[2-(methylaminosulfonyl)ethyl]-1-nitrosourea



- f) 1-(2-chloroethyl)-3-[2-(dipropylaminosulfonyl)ethyl]-1-nitrosourea
 g) 1-(2-chloroethyl)-3-[2-(2-hydroxyethylaminosulfonyl)ethyl]-1-nitrosourea
 h) 1-(2-chloroethyl)-3-[2-(N-2-hydroxyethyl-N-methyl-aminosulfonyl)ethyl]-1-nitrosourea
 i) 1-(2-chloroethyl)-3-[2-(2-methoxyethylaminosulfonyl)ethyl]-1-nitrosourea
 j) 1-(2-chloroethyl)-3-[2-(dimethylaminosulfonyl)ethyl]-3-methyl-1-nitrosourea
 k) 1-(2-chloroethyl)-3-[5-(dimethylaminosulfonyl)pentyl]-1-nitrosourea
 l) 1-(2-chloroethyl)-3-[3-(dimethylaminosulfonyl)propyl]-1-nitrosourea
 m) 1-(2-chloroethyl)-3-[4-(dimethylaminosulfonyl)butyl]-1-nitrosourea
 n) 1-(2-chloroethyl)-3-[2-(N-ethyl-N-propylaminosulfonyl)ethyl]-1-nitrosourea

9. A method for the preparation of a compound of the formula



wherein A is lower alkyl or halolower alkyl;


B is hydrogen; lower alkenyl; lower alkynyl; 3-6 C cycloalkyl; $-\text{RSO}_2\text{NR}^3\text{R}^4$, or lower alkyl optionally monosubstituted with lower alkoxy or 3-6 C cycloalkyl;

R is straight or branched 2-5 C alkylene, optionally monosubstituted with $\text{SO}_2\text{NR}^3\text{R}^4$ or CONR^3R^4 , said alkylene always containing at least two carbon atoms separating the nitrogen atom of the urea from any $\text{SO}_2\text{NR}^1\text{R}^2$ or $\text{SO}_2\text{NR}^3\text{R}^4$;

R¹ and R² are the same or different and selected from hydrogen, 5-6 C cycloalkyl, lower alkoxy, phenyl, benzyl, and straight or branched 1-6 C alkyl, optionally monosubstituted with hydroxy, lower alkoxy, $\text{SO}_2\text{NR}^3\text{R}^4$, or CONR^3R^4 ; R¹ and R² may also together form a 4-5 C alkylene;

R³ and R⁴ are the same or different and selected from hydrogen or lower alkyl or R³ and R⁴ together may form a 4-5 C alkylene; characterized by

a) N-nitrosation of a corresponding urea of the formula



wherein R¹, R², R, B and A are as defined in claim 1 or

b) reacting an amine of the general formula



wherein R¹, R², R and B are as defined above;


with an N-alkyl-N-nitroso-carbamoyl derivative of the general formula



wherein A is as defined before and wherein L is a suitable leaving group.

10. A process according to claim 9 characterized in that a compound according to any of claims 1-8 is prepared.

11. A pharmaceutical preparation comprising as an active agent an effective amount of a compound of the formula



wherein A is lower alkyl or halolower alkyl;



B is hydrogen; lower alkenyl; lower alkynyl; 3—6 C cycloalkyl; $-\text{RSO}_2\text{NR}^3\text{R}^4$, or lower alkyl optionally monosubstituted with lower alkoxy or 3—6 C cycloalkyl;

R is straight or branched 2—5 C alkylene, optionally monosubstituted with $\text{SO}_2\text{NR}^3\text{R}^4$ or CONR^3R^4 , said alkylene always containing at least two carbon atoms separating the nitrogen atom of the urea from any

5 $\text{SO}_2\text{NR}^1\text{R}^2$ or $\text{SO}_2\text{NR}^3\text{R}^4$;

R^1 and R^2 are the same or different and selected from hydrogen, 5—6 C cycloalkyl, lower alkoxy, phenyl, benzyl, and straight or branched 1—6 C alkyl, optionally monosubstituted with hydroxy, lower alkoxy, $\text{SO}_2\text{NR}^3\text{R}^4$, or CONR^3R^4 ; R^1 and R^2 may also together form a 4—5 C alkylene;


10 R^3 and R^4 are the same or different and selected from hydrogen or lower alkyl or R^3 and R^4 together may form a 4—5 C alkylene;

in association with a pharmaceutically acceptable carrier for use as anticancer agents.

12. A pharmaceutical preparation according to claim 11 comprising as an active ingredient a compound according to any of the claims 1—8.

15 Patentansprüche

1. Neue Verbindungen der allgemeinen Formel



25 worin A niederes Alkyl oder niederes Halogenalkyl, B Wasserstoff, niederes Alkenyl, niederes Alkynyl, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, $-\text{RSO}_2\text{NR}^3\text{R}^4$ oder gegebenenfalls mit niederem Alkoxy oder Cycloalkyl mit 3 bis 6 Kohlenstoffatomen monosubstituiertes niederes Alkyl ist,

R gerad- oder verzweigt-kettiges Alkyl mit 2 bis 5 Kohlenstoffatomen, gegebenenfalls monosubstituiert mit $\text{SO}_2\text{NR}^3\text{R}^4$ oder CONR^3R^4 , bedeutet, worin dieses Alkyl immer wenigstens zwei das Stickstoffatom des Harnstoffs von jeglichem $\text{SO}_2\text{NR}^1\text{R}^2$ oder $\text{SO}_2\text{NR}^3\text{R}^4$ trennende Kohlenstoffatome

30 enthält, R^1 und R^2 gleich oder verschieden und aus Wasserstoff, Cycloalkyl mit 5 bis 6 Kohlenstoffatomen, niederem Alkoxy, Phenyl, Benzyl und gerad- oder verzweigt-kettigem, gegebenenfalls mit Hydroxy, niederem Alkoxy, $\text{SO}_2\text{NR}^3\text{R}^4$ oder CONR^3R^4 monosubstituiertem Alkyl mit 1 bis 6 Kohlenstoffatomen ausgewählt sind, R^1 und R^2 ebenso zusammen ein Alkyl mit 4 bis 5 Kohlenstoffatomen bilden können,

35 R^3 und R^4 gleich oder verschieden sind und aus Wasserstoff oder niederem Alkyl ausgewählt sind oder R^3 und R^4 zusammen ein Alkyl mit 4 bis 5 Kohlenstoffatomen bilden können.

2. Verbindungen nach Anspruch 1, dadurch gekennzeichnet, daß A Chlorethyl oder Fluorethyl, vorzugsweise Chlorethyl, ist.

3. Verbindungen nach Anspruch 1 und 2, dadurch gekennzeichnet, daß R unsubstituiertes Alkyl mit 2

40 bis 5 Kohlenstoffatomen, vorzugsweise Ethyl, ist.

4. Verbindungen nach einem der Ansprüche 1 bis 3, dadurch gekennzeichnet, daß B Wasserstoff oder niederes Alkyl, vorzugsweise Wasserstoff, ist.

5. Verbindungen nach einem der Ansprüche 1 bis 4, dadurch gekennzeichnet, daß einer der Substituenten R^1 und R^2 Methyl und der andere der Substituenten R^1 und R^2 niederes Alkyl oder

45 2-Hydroxyethyl ist.

6. Verbindungen nach einem der Ansprüche 1 bis 4, dadurch gekennzeichnet, daß R^1 und R^2 gleich sind und aus der Gruppe aus Wasserstoff, Methyl, Ethyl und Propyl ausgewählt sind oder worin R^1 und R^2 zusammen Alkyl mit 5 Kohlenstoffatomen sind.

7. Verbindungen nach einem der Ansprüche 1 bis 4, dadurch gekennzeichnet, daß R^1 und R^2

50 Wasserstoff und/oder Methyl sind.

8. Verbindungen nach Anspruch 1, ausgewählt aus der folgenden Gruppe

(a) 1-(2-Chlorethyl)-3-[2-dimethylaminosulfonyl]ethyl-1-nitrosoharnstoff

(b) 1-(2-Chlorethyl)-3-[2-(diethylaminosulfonyl)ethyl]-1-nitrosoharnstoff

(c) 3-[2-(Aminosulfonyl)ethyl]-1-(2-chloroethyl)-1-nitrosoharnstoff

55 (d) 1-(2-Chlorethyl)-1-nitroso-3-[2-(1-piperidinsulfonyl)ethyl]-harnstoff

(e) 1-(2-Chlorethyl)-3-[2-(methylaminosulfonyl)ethyl]-1-nitrosoharnstoff

(f) 1-(2-Chlorethyl)-3-[2-(dipropylaminosulfonyl)ethyl]-1-nitrosoharnstoff

(g) 1-(2-Chlorethyl)-3-[2-(2-hydroxyethylaminosulfonyl)ethyl]-1-nitrosoharnstoff

(h) 1-(2-Chlorethyl)-3-[2-(N-2-hydroxyethyl-N-methyl-aminosulfonyl)ethyl]-1-nitrosoharnstoff

60 (i) 1-(2-Chlorethyl)-3-[2-[2-methoxyethylaminosulfonyl]ethyl]-1-nitrosoharnstoff

(j) 1-(2-Chlorethyl)-3-[2-(dimethylaminosulfonyl)ethyl]-3-methyl-1-nitrosoharnstoff

(k) 1-(2-Chlorethyl)-3-[5-[dimethylaminosulfonyl]pentyl]-1-nitrosoharnstoff


(l) 1-(2-Chlorethyl)-3-[3-[dimethylaminosulfonyl]propyl]-1-nitrosoharnstoff

(m) 1-(2-Chlorethyl)-3-[4-[dimethylaminosulfonyl]butyl]-1-nitrosoharnstoff

65 (n) 1-(2-chlorethyl)-3-[2-[N-ethyl-N-propylaminosulfonyl]ethyl]-1-nitrosoharnstoff



9. Verfahren zur Herstellung einer Verbindung der Formel



worin A niederes Alkyl oder niederes Halogenalkyl ist,

10 B Wasserstoff, niederes Alkenyl, niederes Alkynyl, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, —RSO₂NR³R⁴ oder gegebenenfalls mit niederem Alkoxy oder Cycloalkyl mit 3 bis 6 Kohlenstoffatomen monosubstituiertes niederes Alkyl ist,


R gegebenenfalls mit SO₂NR³R⁴ oder CONR³R⁴ monosubstituiertes gerad- oder verzweigt-kettiges Alkyl mit 2 bis 5 Kohlenstoffatomen, wobei dieses Alkyl immer wenigstens zwei das Stickstoffatom des Harnstoffs von jeglichem SO₂NR¹R² oder SO₂NR³R⁴ trennende Kohlenstoffatome enthält,

15 R¹ und R² gleich oder verschieden sind und aus Wasserstoff, Cycloalkyl mit 5 bis 6 Kohlenstoffatomen niederes Alkoxy, Phenyl, Benzyl und gegebenenfalls mit Hydroxy, niederem Alkoxy, SO₂NR³R⁴ oder CONR³R⁴ monosubstituiertes, gerad- oder verzweigt-kettiges Alkyl mit 1 bis 6 Kohlenstoffatomen ausgewählt sind,


20 R¹ und R² ebenso zusammen ein Alkyl mit 4 bis 5 Kohlenstoffatomen bilden können,

R³ und R⁴ gleich oder verschieden sind und aus Wasserstoff oder niederem Alkyl ausgewählt sind und R³ und R⁴ zusammen Alkyl mit 4 bis 5 Kohlenstoffatomen bilden können, gekennzeichnet durch


a) N-Nitrosierung eines entsprechenden Harnstoffes der Formel



30 worin R¹, R², R, B und A die in Anspruch 1 angegebene Bedeutung haben oder
b) Umsetzung eines Amins der allgemeinen Formel




40 worin R¹, R², R und B die obenangegebene Bedeutung haben,
mit einem N-Alkyl-N-nitroso-carbamoylderivat der allgemeinen Formel



worin A die obenangegebene Bedeutung hat und worin L eine geeignete Abgangsgruppe darstellt.

50 10. Verfahren nach Anspruch 9, dadurch gekennzeichnet, daß eine Verbindung nach einem der Ansprüche 1 bis 8 hergestellt wird.

11. Pharmazeutische Zubereitung, welche als aktiven Bestandteil eine wirksame Menge einer Verbindung der Formel



worin A niederes Alkyl oder niederes Halogenalkyl ist,

B Wasserstoff, niederes Alkenyl, niederes Alkynyl, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, —RSO₂NR³R⁴ oder gegebenenfalls mit niederem Alkoxy oder Cycloalkyl mit 3 bis 6 Kohlenstoffatomen monosubstituiertes niederes Alkyl ist,

65 R gegebenenfalls mit SO₂NR³R⁴ oder CONR³R⁴ monosubstituiertes gerad- oder verzweigt-kettiges



Alkylen mit 2 bis 5 Kohlenstoffatomen ist, wobei dieses Alkyl immer wenigstens zwei das Stickstoffatom des Harnstoffs von jeglichem $\text{SO}_2\text{NR}^1\text{R}^2$ oder $\text{SO}_2\text{NR}^3\text{R}^4$ trennende Kohlenstoffatome enthält,

R^1 und R^2 gleich oder verschieden sind und aus Wasserstoff, Cycloalkyl mit 5 bis 6 Kohlenstoffatomen, niederem Alkoxy, Phenyl, Benzyl und gegebenenfalls mit Hydroxy, niederem Alkoxy, $\text{SO}_2\text{NR}^3\text{R}^4$ oder CONR^3R^4 monosubstituiertem gerad- oder verzweigt-kettigem Alkyl mit 1 bis 6 Kohlenstoffatomen ausgewählt sind,

R^1 und R^2 ebenso zusammen ein Alkyl mit 4 bis 5 Kohlenstoffatomen bilden können,


R^3 und R^4 gleich oder verschieden sind und aus Wasserstoff oder niederem Alkyl ausgewählt sind oder R^3 und R^4 zusammen Alkyl mit 4 bis 6 Kohlenstoffatomen bilden können,

in Verbindung mit einem pharmazeutisch annehmbaren Träger zur Anwendung als Mittel gegen Krebs.

12. Pharmazeutische Zubereitung nach Anspruch 11, welches als aktiver Bestandteil eine Verbindung nach einem der Ansprüche 1 bis 8 enthält.

15 Revendications

1. Nouveaux composés ayant la formule générale:



dans laquelle A est alcoyle inférieur ou halogénoalcoyle inférieur;

B est hydrogène; alcényle inférieur; alcynyle inférieur; cycloalcoyle à 3—6 C;

— $\text{RSO}_2\text{NR}^3\text{R}^4$ ou alcoyle inférieur éventuellement monosubstitué par alcoxy inférieur ou cycloalcoyle à 3—6 C;

R est alcoylène à 2—5 C, rectiligne ou ramifié, éventuellement monosubstitué par $\text{SO}_2\text{NR}^3\text{R}^4$ ou CONR^3R^4 , ledit alcoylène contenant toujours au moins deux atomes de carbone séparant l'atome d'azote de l'urée de tout $\text{SO}_2\text{NR}^1\text{R}^2$ ou $\text{SO}_2\text{NR}^3\text{R}^4$;

R^1 et R^2 sont identiques ou différents et choisis parmi hydrogène, cycloalcoyle à 5—6 C, alcoxy inférieur, phényle, benzyle, et alcoyle à 1—6 C, rectiligne ou ramifié, éventuellement monosubstitué par hydroxy, alcoxy inférieur, $\text{SO}_2\text{NR}^3\text{R}^4$ ou CONR^3R^4 ; R^1 et R^2 peuvent également former ensemble un alcoylène à 4—5 C;

R^3 et R^4 sont identiques ou différents et choisis parmi hydrogène ou alcoyle inférieur ou R^3 et R^4 peuvent former ensemble un alcoylène à 4—5 C.

2. Composés selon la revendication 1, caractérisés en ce que A est chloroéthyle ou fluoroéthyle, de préférence chloroéthyle.

3. Composés selon la revendication 1—2, caractérisés en ce que R est alcoylène à 2—5 C non substitué, de préférence éthylène.

4. Composés selon l'une quelconque des revendications 1—3, caractérisés en ce que B est hydrogène ou alcoyle inférieur, de préférence hydrogène.

5. Composés selon l'une quelconque des revendications 1—4, caractérisés en ce que l'un des substituants R^1 et R^2 est méthyle et l'autre des substituants R^1 et R^2 est alcoyle inférieur ou 2-hydroxyéthyle.

6. Composés selon l'une quelconque des revendications 1—4, caractérisés en ce que R^1 et R^2 sont identiques et choisis dans le groupe hydrogène, méthyle, éthyle et propyle, ou dans lesquels R^1 et R^2 sont ensemble alcoylène à 5 C.


7. Composés selon l'une quelconque des revendications 1—4, caractérisés en ce que R^1 et R^2 sont hydrogène et/ou méthyle.

8. Composés selon la revendication 1 choisis dans le groupe suivant:

- a) 1-(2-chloroéthyl)-3-[2-(diméthylaminosulfonyl)éthyl]-1-nitrosourée
- b) 1-(2-chloroéthyl)-3-[2-(diéthylaminosulfonyl)éthyl]-1-nitrosourée
- c) 3-[2-(aminosulfonyl)éthyl]-1-(2-chloroéthyl)-1-nitrosourée
- d) 1-(2-chloroéthyl)-1-nitroso-3-[2-(1-pipéridinosulfonyl)éthyl]-urée
- e) 1-(2-chloroéthyl)-3-[2-(méthylaminosulfonyl)éthyl]-1-nitrosourée
- f) 1-(2-chloroéthyl)-3-[2-(dipropylaminosulfonyl)éthyl]-1-nitrosourée
- g) 1-(2-chloroéthyl)-3-[2-(2-hydroxyéthylaminosulfonyl)éthyl]-1-nitrosourée
- h) 1-(2-chloroéthyl)-3-[2-(N-2-hydroxyéthyl-N-méthyl-aminosulfonyl)éthyl]-1-nitrosourée
- i) 1-(2-chloroéthyl)-3-[2-(2-méthoxyéthylaminosulfonyl)éthyl]-1-nitrosourée
- j) 1-(2-chloroéthyl)-3-[2-(diméthylaminosulfonyl)éthyl]-3-méthyl-1-nitrosourée
- k) 1-(2-chloroéthyl)-3-(5-[diméthylaminosulfonyl]pentyl)-1-nitrosourée
- l) 1-(2-chloroéthyl)-3-(3-[diméthylaminosulfonyl]propyl)-1-nitrosourée
- m) 1-(2-chloroéthyl)-3-(4-[diméthylaminosulfonyl]butyl)-1-nitrosourée
- n) 1-(2-chloroéthyl)-3-[2-(N-éthyl-N-propylaminosulfonyl)éthyl]-1-nitrosourée



9. Procédé pour la préparation d'un composé de formule



dans laquelle A est alcoyle inférieur ou halogénoalcoyle inférieur;


B est hydrogène; alcényle inférieur; alcynyle inférieur; cycloalcoyle à 3—6 C; $-\text{RSO}_2\text{NR}^3\text{R}^4$, ou alcoyle inférieur éventuellement monosubstitué par alcoxy inférieur ou cycloalcoyle à 3—6 C;

R est alcoylène à 2—5 C, rectiligne ou ramifié, éventuellement monosubstitué par $\text{SO}_2\text{NR}^3\text{R}^4$ ou CONR^3R^4 , ledit alcoylène contenant toujours au moins deux atomes de carbone séparant l'atome d'azote de l'urée de tout $\text{SO}_2\text{NR}^1\text{R}^2$ ou $\text{SO}_2\text{NR}^3\text{R}^4$;

R¹ et R² sont identiques ou différents et choisis parmi hydrogène, cycloalcoyle à 5—6 C, alcoxy inférieur, phényle, benzyle, et alcoyle à 1—6 C, rectiligne ou ramifié, éventuellement monosubstitué par hydroxy, alcoxy inférieur, $\text{SO}_2\text{NR}^3\text{R}^4$ ou CONR^3R^4 ; R¹ et R² peuvent également former ensemble un alcoylène à 4—5 C;

R³ et R⁴ sont identiques ou différents et choisis parmi hydrogène ou alcoyle inférieur ou R³ et R⁴ peuvent former ensemble un alcoylène à 4—5 C; caractérisé par:

a) N-nitrosation d'une urée correspondante de formule



dans laquelle R¹, R², R, B et A sont tels que définis dans la revendication 1 ou

b) réaction d'une amine de formule générale



dans laquelle R¹, R², R et B sont tels que définis plus haut;


avec un dérivé de N-alcoyl-N-nitroso-carbamoyl de formule générale



dans laquelle A est tel que défini précédemment et dans laquelle L est un groupe partant convenable.

10. Procédé selon la revendication 9, caractérisé en ce que l'on prépare un composé selon l'une quelconque des revendications 1—8.

11. Préparation pharmaceutique comprenant, en tant qu'agent actif, une quantité efficace d'un composé de formule



dans laquelle A est alcoyle inférieur ou halogénoalcoyle inférieur;

B est hydrogène; alcényle inférieur; alcynyle inférieur; cycloalcoyle à 3—6 C; $-\text{RSO}_2\text{NR}^3\text{R}^4$, ou alcoyle inférieur éventuellement monosubstitué par alcoxy inférieur ou cycloalcoyle à 3—6 C;

R est alcoylène à 2—5 C, rectiligne ou ramifié, éventuellement monosubstitué par $\text{SO}_2\text{NR}^3\text{R}^4$ ou CONR^3R^4 , ledit alcoylène contenant toujours au moins deux atomes de carbone séparant l'atome d'azote de l'urée de tout $\text{SO}_2\text{NR}^1\text{R}^2$ ou $\text{SO}_2\text{NR}^3\text{R}^4$;

R¹ et R² sont identiques ou différents et choisis parmi hydrogène, cycloalcoyle à 5—6 C, alcoxy



inférieur, phényle, benzyle, et alcoyle à 1—6 C, rectiligne ou ramifié, éventuellement monosubstitué par hydroxy, alcoxy inférieur, $\text{SO}_2\text{NR}^3\text{R}^4$ ou CONR^3R^4 ; R^1 et R^2 peuvent également former ensemble un alcoylène à 4—5 C;

R^3 et R^4 sont identiques ou différents et choisis parmi hydrogène ou alcoyle inférieur ou R^3 et R^4 peuvent former ensemble un alcoylène à 4—5 C;
5 en association avec un support pharmaceutiquement acceptable pour l'utilisation en tant qu'agent anti-cancéreux.

12. Préparation pharmaceutique selon la revendication 11 comprenant en tant qu'ingrédient actif un composé selon l'une quelconque des revendications 1—8.

10

15

20

25

30

35

40

45

50

55

60

65

