


MODULATORS OF CHEMOKINE RECEPTOR ACTIVITY, CRYSTALLINE FORMS AND PROCESS

[0001] This application claims priority from U.S. Provisional Application Nos.
5 60/834,235, and 60/896,026 filed July 28, 2006 and March 21, 2007, respectively, the
disclosures of both of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention provides N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-
10 2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-
yl)cyclohexyl)acetamide, or a pharmaceutically acceptable salt, solvate or prodrug,
thereof, having an unexpected combination of desirable pharmacological
characteristics. Crystalline forms of the present invention are also provided.
Pharmaceutical compositions containing the same and methods of using the same as
15 agents for the treatment of inflammatory, allergic, autoimmune, metabolic, cancer
and/or cardiovascular diseases is also an objective of this invention. The present
invention also provides a process for preparing compounds of Formula (I), including
N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-
ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide:



20

wherein R¹, R⁸, R⁹, R¹⁰, and  are as described herein. Compounds that are
useful intermediates of the process are also provided herein.

BACKGROUND OF THE INVENTION

[0003] Chemokines are chemotactic cytokines, of molecular weight 6-15 kDa, that are released by a wide variety of cells to attract and activate, among other cell types, macrophages, T and B lymphocytes, eosinophils, basophils and neutrophils (reviewed in: Charo and Rasonhoff, *New Eng. J. Med.* **2006**, *354*, 610-621; Luster, *New Eng. J. Med.* **1998**, *338*, 436-445; and Rollins, *Blood* **1997**, *90*, 909-928). There are two major classes of chemokines, CXC and CC, depending on whether the first two cysteines in the amino acid sequence are separated by a single amino acid (CXC) or are adjacent (CC). The CXC chemokines, such as interleukin-8 (IL-8), neutrophil-activating protein-2 (NAP-2) and melanoma growth stimulatory activity protein (MGSA) are chemotactic primarily for neutrophils and T lymphocytes, whereas the CC chemokines, such as RANTES, MIP-1 α , MIP-1 β , the monocyte chemotactic proteins (MCP-1, MCP-2, MCP-3, MCP-4, and MCP-5) and the eotaxins (-1 and -2) are chemotactic for, among other cell types, macrophages, T lymphocytes, eosinophils, dendritic cells, and basophils. There also exist the chemokines lymphotactin-1, lymphotactin-2 (both C chemokines), and fractalkine (a CX₃C chemokine) that do not fall into either of the major chemokine subfamilies.

[0004] The chemokines bind to specific cell-surface receptors belonging to the family of G-protein-coupled seven-transmembrane-domain proteins (reviewed in: Horuk, *Trends Pharm. Sci.* **1994**, *15*, 159-165) which are termed "chemokine receptors." On binding their cognate ligands, chemokine receptors transduce an intracellular signal through the associated trimeric G proteins, resulting in, among other responses, a rapid increase in intracellular calcium concentration, changes in cell shape, increased expression of cellular adhesion molecules, degranulation, and promotion of cell migration. There are at least ten human chemokine receptors that bind or respond to CC chemokines with the following characteristic patterns (reviewed in Zlotnik and Oshie *Immunity* **2000**, *12*, 121): CCR-1 (or "CKR-1" or "CC-CKR-1") [MIP-1 α , MCP-3, MCP-4, RANTES] (Ben-Barruch, et al., *Cell* **1993**, *72*, 415-425, and Luster, *New Eng. J. Med.* **1998**, *338*, 436-445); CCR-2A and CCR-2B (or "CKR-2A"/"CKR-2B" or "CC-CKR-2A"/"CC-CKR-2B") [MCP-1, MCP-2, MCP-3, MCP-4, MCP-5] (Charo, et al., *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 2752-2756, and Luster, *New Eng. J. Med.* **1998**, *338*, 436-445); CCR-3 (or "CKR-3" or "CC-CKR-3") [eotaxin-1, eotaxin-2, RANTES, MCP-3, MCP-4] (Combadiere, et

al., *J. Biol. Chem.* **1995**, *270*, 16491-16494, and Luster, *New Eng. J. Med.* **1998**, *338*, 436-445); CCR-4 (or "CKR-4" or "CC-CKR-4") [TARC, MDC] (Power, et al., *J. Biol. Chem.* **1995**, *270*, 19495-19500, and Luster, *New Eng. J. Med.* **1998**, *338*, 436-445); CCR-5 (or "CKR-5" OR "CC-CKR-5") [MIP-1 α , RANTES, MIP-1 β] (Sanson, et al., *Biochemistry* **1996**, *35*, 3362-3367); CCR-6 (or "CKR-6" or "CC-CKR-6") [LARC] (Baba, et al., *J. Biol. Chem.* **1997**, *272*, 14893-14898); CCR-7 (or "CKR-7" or "CC-CKR-7") [ELC] (Yoshie et al., *J. Leukoc. Biol.* **1997**, *62*, 634-644); CCR-8 (or "CKR-8" or "CC-CKR-8") [I-309] (Napolitano et al., *J. Immunol.*, **1996**, *157*, 2759-2763); CCR-10 (or "CKR-10" or "CC-CKR-10") [MCP-1, MCP-3] (Bonini, et al., *DNA and Cell Biol.* **1997**, *16*, 1249-1256); and CCR-11 [MCP-1, MCP-2, and MCP-4] (Schweickert, et al., *J. Biol. Chem.* **2000**, *275*, 90550).

[0005] In addition to the mammalian chemokine receptors, mammalian cytomegaloviruses, herpesviruses and poxviruses have been shown to express, in infected cells, proteins with the binding properties of chemokine receptors (reviewed in: Wells and Schwartz, *Curr. Opin. Biotech.* **1997**, *8*, 741-748). Human CC chemokines, such as RANTES and MCP-3, can cause rapid mobilization of calcium via these virally encoded receptors. Receptor expression may be permissive for infection by allowing for the subversion of normal immune system surveillance and response to infection. Additionally, human chemokine receptors, such as CXCR4, CCR2, CCR3, CCR5 and CCR8, can act as co-receptors for the infection of mammalian cells by microbes as with, for example, the human immunodeficiency viruses (HIV).

[0006] The chemokines and their cognate receptors have been implicated as being important mediators of inflammatory, infectious, and immunoregulatory disorders and diseases, including asthma and allergic diseases; as well as autoimmune pathologies, such as rheumatoid arthritis and multiple sclerosis; and metabolic diseases, such as atherosclerosis and diabetes (reviewed in: Charo and Rasonhoff, *New Eng. J. Med.* **2006**, *354*, 610-621; Z. Gao and W. A. Metz, *Chem. Rev.* **2003**, *103*, 3733; P. H. Carter, *Current Opinion in Chemical Biology* **2002**, *6*, 510; Trivedi, et al, *Ann. Reports Med. Chem.* **2000**, *35*, 191; Saunders and Tarby, *Drug Disc. Today* **1999**, *4*, 80; Premack and Schall, *Nature Medicine* **1996**, *2*, 1174). For example, the chemokine monocyte chemoattractant-1 (MCP-1) and its receptor CC Chemokine

Receptor 2 (CCR-2) play a pivotal role in attracting leukocytes to sites of inflammation and in subsequently activating these cells. When the chemokine MCP-1 binds to CCR-2, it induces a rapid increase in intracellular calcium concentration, increased expression of cellular adhesion molecules, and the promotion of leukocyte migration. Demonstration of the importance of the MCP-1/CCR-2 interaction has been provided by experiments with genetically modified mice. MCP-1^{-/-} mice were unable to recruit monocytes into sites of inflammation after several different types of immune challenge (Bao Lu, et al., *J. Exp. Med.* **1998**, *187*, 601). Likewise, CCR-2^{-/-} mice were unable to recruit monocytes or produce interferon- γ when challenged with various exogenous agents; moreover, the leukocytes of CCR-2 null mice did not migrate in response to MCP-1 (Landin Boring, et al., *J. Clin. Invest.* **1997**, *100*, 2552), thereby demonstrating the specificity of the MCP-1/CCR-2 interaction. Two other groups have independently reported equivalent results with different strains of CCR-2^{-/-} mice (William A. Kuziel, et al., *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 12053, and Takao Kurihara, et al., *J. Exp. Med.* **1997**, *186*, 1757). The viability and generally normal health of the MCP-1^{-/-} and CCR-2^{-/-} animals is noteworthy, in that disruption of the MCP-1/CCR-2 interaction does not induce physiological crisis. Taken together, these data lead one to the conclusion that molecules that block the actions of MCP-1/CCR2 would be useful in treating a number of inflammatory and autoimmune disorders (reviewed in: M. Feria and F. Díaz-González, *Exp. Opin. Ther. Patents* **2006**, *16*, 49; and J. Dawson, W. Miltz, and C. Wiessner, *C. Exp. Opin. Ther. Targets* **2003**, *7*, 35). This hypothesis has now been validated in a number of different animal disease models, as described below.

[0007] It is known that MCP-1 is upregulated in patients with rheumatoid arthritis (Alisa Koch, et al., *J. Clin. Invest.* **1992**, *90*, 772 – 779). Moreover, several preclinical studies have demonstrated the potential therapeutic value of antagonism of the MCP-1/CCR2 interaction in treating rheumatoid arthritis. A DNA vaccine encoding MCP-1 was shown recently to ameliorate chronic polyadjuvant-induced arthritis in rats (Sawsan Youssef, et al., *J. Clin. Invest.* **2000**, *106*, 361). Likewise, the disease symptoms could be controlled via direct administration of antibodies for MCP-1 to rats with collagen-induced arthritis (Hiroomi Ogata, et al., *J. Pathol.* **1997**, *182*, 106), or streptococcal cell wall-induced arthritis (Ralph C. Schimmer, et al., *J.*

Immunol. **1998**, *160*, 1466). Perhaps most significantly, a peptide antagonist of MCP-1, MCP-1(9-76), was shown both to prevent disease onset and to reduce disease symptoms (depending on the time of administration) in the MRL-lpr mouse model of arthritis (Jiang-Hong Gong, et al., *J. Exp. Med.* **1997**, *186*, 131). Moreover, it has
5 been demonstrated the administration of small molecule CCR2 antagonists reduced clinical score in rodent models of arthritis (C. M. Brodmerkel, et al, *J. Immunol.* **2005**, *175*, 5370; and M. Xia, et al. US Patent Application 0069123, 2006). Administration of an anti-CCR2 antibody had varying effects on murine CIA, depending on the time of administration (H. Bruhl, et al. *J. Immunol.* **2004**, *172*, 890).
10 Recent studies with CCR2^{-/-} mice have suggested that deletion of CCR2 can exacerbate rodent arthritis models in specific experimental circumstances (M. P. Quinones, et al. *J. Clin. Invest.* **2004**, *113*, 856; M. P. Quinones, et al. *J. Mol. Med.* **2006**, *84*, 503).
[0008] It is known that MCP-1 is upregulated in atherosclerotic lesions, and it has
15 been shown that circulating levels of MCP-1 are reduced through treatment with therapeutic agents (Abdolreza Rezaie-Majd, et al, *Arterioscler. Thromb. Vasc. Biol.* **2002**, *22*, 1194 – 1199). Several key studies have demonstrated the potential therapeutic value of antagonism of the MCP-1/CCR2 interaction in treating atherosclerosis. For example, when MCP-1^{-/-} mice are crossed with LDL receptor-
20 deficient mice, an 83% reduction in aortic lipid deposition was observed (Long Gu, et al., *Mol. Cell* **1998**, *2*, 275). Similarly, when MCP-1 was genetically ablated from mice which already overexpressed human apolipoprotein B, the resulting mice were protected from atherosclerotic lesion formation relative to the MCP-1^{+/+} apoB control mice (Jennifa Gosling, et al., *J. Clin. Invest.* **1999**, *103*, 773). Likewise, when
25 CCR-2^{-/-} mice are crossed with apolipoprotein E^{-/-} mice, a significant decrease in the incidence of atherosclerotic lesions was observed (Landin Boring, et al, *Nature* **1998**, *394*, 894; T. C. Dawson, et al. *Atherosclerosis* **1999**, *143*, 205). Finally, when apolipoprotein E^{-/-} mice are administered a gene encoding a peptide antagonist of CCR2, then lesion size is decreased and plaque stability is increased (W. Ni, et al.
30 *Circulation* **2001**, *103*, 2096 – 2101). Transplantation of bone marrow from CCR2^{-/-} mice into ApoE3-Leiden mice inhibited early atherogenesis (J. Guo, et al.

Arterioscler. Thromb. Vasc. Biol. **2003**, *23*, 447), but had minimal effects on advanced lesions (J. Guo, et al. *Arterioscler. Thromb. Vasc. Biol.* **2005**, *25*, 1014).

[0009] Patients with type 2 diabetes mellitus typically exhibit insulin resistance as one of the hallmark features of the disease. Insulin resistance is also associated with the grouping of abnormalities known as the "metabolic syndrome" or "syndrome X," which includes obesity, atherosclerosis, hypertension, and dyslipidemia (reviewed in: Eckel, et al. *Lancet* **2005**, *365*, 1415). It is well-recognized that inflammation plays a role in exacerbating the disease process in type 2 diabetes and the "syndrome X" pathologies (reviewed in: Chen, *Pharmacological Research* **2006**, *53*, 469; Neels and Olefsky, *J. Clin. Invest.* **2006**, *116*, 33; Danadona and Aljada, *Am J Cardiol.* **2002** *90*, 27G-33G; Pickup and Crook, *Diabetologia* **1998**, *41*, 1241). MCP-1 is recognized as playing a role in obesity-induced insulin resistance. In culture, human preadipocytes constitutively expressed MCP-1 (Gerhardt, *Mol. Cell. Endocrinology* **2001**, *175*, 81). CCR2 is expressed on adipocytes; Addition of MCP-1 to differentiated adipocytes in vitro decreases insulin-stimulated glucose uptake and the expression of several adipogenic genes (LpL, adipsin, GLU-4), α 2, β 3-adrenergic receptor, and PPAR γ) (P. Sartipy and D. Loskutoff, *Proc. Natl. Acad. Sci USA* **1999**, *96*, 6902). Patients with type 2 diabetes had greater levels of circulating MCP-1 than non-diabetic controls (S. Nomura, et al. *Clin. Exp. Immunol.* **2000**, *121*, 437), and release of MCP-1 from adipose tissue could be reduced by treatment with anti-diabetic therapies such as metformin or thiazolidinediones (J. M. Bruun, et al. *J. Clin. Endocrinol. Metab.* **2005**, *90*, 2282). Likewise, MCP-1 was also overexpressed in murine experimental models of obesity, and was primarily produced by adipose tissue (Sartipy and Loskutoff, *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 7265). In obese mice, the expression of MCP-1 both preceded and occurred concurrently with the onset of insulin resistance (H. Xu, et al. *J. Clin. Invest.* **2003**, *112*, 1821). Another study showed that the expression of MCP-1 positively correlated with body mass in the perigonadal adipose tissue of mice (Weisberg, et al. *J. Clin. Invest.* **2003**, *112*, 1796). Consistent with these data, the development of insulin resistance in *db/db* mice was ameliorated either via genetic deletion of MCP-1 or by gene-induced expression of a dominant negative peptide (H. Kanda, et al. *J. Clin. Invest.* **2006**, *116*, 1494). The logical converse could also be demonstrated: overexpression of MCP-1 in adipose

tissue promoted insulin resistance (N. Kamei, et al. *J. Biol. Chem.* **2006**, *281*, 26602). One conflicting result showing that genetic deletion of MCP-1 does not effect insulin resistance in the *db/db* mouse has also appeared (F. Y. Chow, et al. *Diabetologia* **2007**, *50*, 471). Consistent with the data on MCP-1, direct studies with CCR2 (the MCP-1 receptor) have showed that it plays a role in the formation of obesity and obesity-induced insulin resistance. Maintenance of a high fat diet increased the numbers of circulating CCR2⁺ inflammatory monocytes in both wild-type (C. L. Tsou, et al. *J. Clin. Invest.* **2007**, *117*, 902) and ApoE^{-/-} mice (F. Tacke, et al. *J. Clin. Invest.* **2007**, *117*, 185). Genetic deletion of CCR2 reduced numbers of activated macrophages in murine adipose tissue (C. N. Lumeng, et al. *Diabetes* **2007**, *56*, 16), but did not affect a population of M2 adipose macrophages thought to maintain the “lean” state (C. N. Lumeng, et al. *J. Clin. Invest.* **2007**, *117*, 175). Genetic deletion of CCR2 reduced diet-induced obesity and improved insulin sensitivity in diet-induced obesity model (S. P. Weisberg, et al. *J. Clin. Invest.* **2006**, *116*, 115; P Cornelius, RP Gladue, RS Sebastian, WO patent 2006/013427 A2), 2006), depending on experimental conditions (A. Chen, et al. *Obes. Res.* **2005**, *13*, 1311). Administration of a small molecule CCR2 antagonist also improved insulin sensitivity in this same model (S. P. Weisberg, et al. *J. Clin. Invest.* **2006**, *116*, 115).

[0010] Two studies described the important role of CCR2 in hypertension-induced vascular inflammation, remodeling, and hypertrophy (E Bush et al., *Hypertension* **2000**, *36*, 360; M Ishibashi, et al. *Circ. Res.* **2004**, *94*, 1203).

[0011] It is known that MCP-1 is upregulated in human multiple sclerosis, and it has been shown that effective therapy with interferon β -1b reduces MCP-1 expression in peripheral blood mononuclear cells, suggesting that MCP-1 plays a role in disease progression (Carla Iarlori, et al., *J. Neuroimmunol.* **2002**, *123*, 170 – 179). Other studies have demonstrated the potential therapeutic value of antagonism of the MCP-1/CCR-2 interaction in treating multiple sclerosis; all of these studies have been demonstrated in experimental autoimmune encephalomyelitis (EAE), the conventional animal model for multiple sclerosis. Administration of antibodies for MCP-1 to animals with EAE significantly diminished disease relapse (K. J. Kennedy, et al., *J. Neuroimmunol.* **1998**, *92*, 98). Furthermore, two reports have shown that CCR-2 ^{-/-} mice are resistant to EAE (B. T. Fife, et al., *J. Exp. Med.* **2000**, *192*, 899;

L. Izikson, et al., *J. Exp. Med.* **2000**, *192*, 1075). A subsequent report extended these initial observations by examining the effects of CCR2 deletion in mice from different strains (S. Gaupp, et al. *Am. J. Pathol.* **2003**, *162*, 139). Notably, administration of a small molecule CCR2 antagonist also blunted disease progression in C57BL/6 mice (C. M. Brodmerkel, et al. *J. Immunol.* **2005**, *175*, 5370).

[0012] It is known that MCP-1 is upregulated in patients who develop bronchiolitis obliterans syndrome after lung transplantation (Martine Reynaud-Gaubert, et al., *J. of Heart and Lung Transplant.*, **2002**, *21*, 721 – 730; John Belperio, et al., *J. Clin. Invest.* **2001**, *108*, 547 – 556). In a murine model of bronchiolitis obliterans syndrome, administration of an antibody to MCP-1 led to attenuation of airway obliteration; likewise, CCR2 $-/-$ mice were resistant to airway obliteration in this same model (John Belperio, et al., *J. Clin. Invest.* **2001**, *108*, 547 – 556). These data suggest that antagonism of MCP-1/CCR2 may be beneficial in treating rejection of organs following transplantation. In addition, studies have shown that disruption of MCP-1/CCR2 axis was able to prolong the survival of islet transplant (I Lee et al. *J Immunol* **2003**, *171*, 6929; R Abdi et al., *J Immunol* **2004**, *172*, 767). In rat graft models, CCR2 and MCP-1 was shown to be upregulated in grafts that develop graft vasculopathy (K Horiguchi et al., *J Heart Lung Transplant.* **2002**, *21*, 1090). In another study, anti-MCP-1 gene therapy attenuated graft vasculopathy (A Saiura et al., *Artheroscler Thromb Vasc Biol* **2004**, *24*, 1886). One study described inhibition of experimental vein graft neointimal formation by blockage of MCP-1 (H Tatewaki et al., *J Vasc Surg.* **2007**, *45*, 1236).

[0013] Other studies have demonstrated the potential therapeutic value of antagonism of the MCP-1/CCR2 interaction in treating asthma. Sequestration of MCP-1 with a neutralizing antibody in ovalbumin-challenged mice resulted in marked decrease in bronchial hyperresponsiveness and inflammation (Jose-Angel Gonzalo, et al., *J. Exp. Med.* **1998**, *188*, 157). It proved possible to reduce allergic airway inflammation in *Schistosoma mansoni* egg-challenged mice through the administration of antibodies for MCP-1 (Nicholas W. Lukacs, et al., *J. Immunol.* **1997**, *158*, 4398). Consistent with this, MCP-1 $-/-$ mice displayed a reduced response to challenge with *Schistosoma mansoni* egg (Bao Lu, et al., *J. Exp. Med.* **1998**, *187*, 601).

- [0014] Other studies have demonstrated the potential therapeutic value of antagonism of the MCP-1/CCR2 interaction in treating kidney disease. Administration of antibodies for MCP-1 in a murine model of glomerulonephritis resulted in a marked decrease in glomerular crescent formation and deposition of type I collagen (Clare M. Lloyd, et al., *J. Exp. Med.* **1997**, *185*, 1371). In addition, MCP-1
5 -/- mice with induced nephrotoxic serum nephritis showed significantly less tubular damage than their MCP-1 +/+ counterparts (Gregory H. Tesch, et al., *J. Clin. Invest.* **1999**, *103*, 73).
- [0015] Several studies have demonstrated the potential therapeutic value of antagonism of the MCP-1/CCR2 interaction in treating systemic lupus erythematosus.
10 CCR2^{-/-} mice exhibited prolonged survival and reduced renal disease relative to their WT counterparts in a murine model of systemic lupus erythematosus (G. Perez de Lema, et al. *J. Am. Soc. Neph.* **2005**, *16*, 3592). These data are consistent with the disease-modifying activity found in recent studies on genetic deletion of MCP-1 (S.
15 Shimizu, et al. *Rheumatology (Oxford)* **2004**, *43*, 1121; Gregory H. Tesch, et al., *J. Exp. Med.* **1999**, *190*, 1813) or administration of a peptide antagonist of CCR2 (H. Hasegawa, et al. *Arthritis & Rheumatism* **2003**, *48*, 2555) in rodent models of lupus.
- [0016] A remarkable 30-fold increase in CCR2⁺ lamina propria lymphocytes was observed in the small bowels from Crohn's patients relative to non-diseased ileum (S.
20 J. Connor, et al. *Gut* **2004**, *53*, 1287). Also of note, there was an expansion in the subset of circulating CCR2⁺/CD14⁺/CD56⁺ monocytes in patients with active Crohn's disease relative to controls. Several rodent studies have demonstrated the potential therapeutic value of antagonism of the MCP-1/CCR2 interaction in treating Crohn's disease/colitis. CCR-2^{-/-} mice were protected from the effects of dextran sodium
25 sulfate-induced colitis (Pietro G. Andres, et al., *J. Immunol.* **2000**, *164*, 6303). Administration of a small molecule antagonist of CCR2, CCR5, and CXCR3 (murine binding affinities = 24, 236, and 369 nM, respectively) also protected against dextran sodium sulfate-induced colitis (H. Tokuyama, et al. *Int. Immunol.* **2005**, *17*, 1023). Finally, MCP-1^{-/-} mice showed substantially reduced colonic damage (both
30 macroscopic and histological) in a hapten-induced model of colitis (W. I. Khan, et al. *Am. J. Physiol. Gastrointest. Liver Physiol.* **2006**, *291*, G803).

- [0017] Two reports described the overexpression of MCP-1 in the intestinal epithelial cells and bowel mucosa of patients with inflammatory bowel disease (H. C. Reinecker, et al., *Gastroenterology* **1995**, *108*, 40, and Michael C. Grimm, et al., *J. Leukoc. Biol.* **1996**, *59*, 804).
- 5 [0018] One study described the association of promoter polymorphism in the MCP-1 gene with scleroderma (systemic sclerosis) (S Karrer et al., *J Invest Dermatol.* **2005**, *124*, 92). In related models of tissue fibrosis, inhibition of CCR2/MCP-1 axis reduced fibrosis in skin (T Yamamoto and K Nishioka, *J Invest Dermatol* **2003**, *121*, 510; AM Ferreira et al., *J Invest Dermatol.* **2006**, *126*, 1900), lung (T Okuma et al., *J*
- 10 *Pathol.* **2004**, *204*, 594; M Gharaee-Kermani et al., *Cytokine* **2003**, *24*, 266), kidney (K Kitagawa et al., *Am J Pathol.* **2004**, *165*, 237; T Wada et al., *J Am Soc Nephrol* **2004**, *15*, 940), heart (S Hayashidani et al., *Circulation* **2003**, *108*, 2134), and liver (S Tsuruta et al., *Int J Mol Med.* **2004**, *14*, 837).
- [0019] One study has demonstrated the potential therapeutic value of antagonism
- 15 of the MCP-1/CCR2 interaction in treating alveolitis. When rats with IgA immune complex lung injury were treated intravenously with antibodies raised against rat MCP-1 (JE), the symptoms of alveolitis were partially alleviated (Michael L. Jones, et al., *J. Immunol.* **1992**, *149*, 2147).
- [0020] Several studies have shown the potential therapeutic value of antagonism
- 20 of the MCP-1/CCR2 interaction in treating cancer (reviewed in: M. J. Craig and R. D. Loberg, *Cancer Metastasis Rev.* **2006**, *25*, 611; I. Conti and B. Rollins, *Seminars in Cancer Biology* **2004**, *14*, 149; R. Giles and R. D. Loberg, *Curr. Cancer Drug Targets* **2006**, *6*, 659). When immunodeficient mice bearing human breast carcinoma cells were treated with an anti-MCP-1 antibody, inhibition of lung micrometastases
- 25 and increases in survival were observed (Rosalba Salcedo, et al., *Blood* **2000**, *96*, 34 – 40). Using human clinical tumor specimens, CCR2 expression was associated with prostate cancer progression (Y. Lu, et al. *J. Cell. Biochem.* **2007**, *101*, 676). *In vitro*, MCP-1 expression has been shown to mediate prostate cancer cell growth and invasion (Y. Lu, et al. *Prostate* **2006**, *66*, 1311); furthermore, MCP-1 expressed by
- 0 prostate cancer cells induced human bone marrow progenitors for bone resorption (Y. Lu, et al., *Cancer Res.* **2007**, *67*, 3646).

- [0021] Multiple studies have described the potential therapeutic value of antagonism of the MCP-1/CCR2 interaction in treating restenosis. In humans, MCP-1 levels correlate directly with risk for restenosis (F. Cipollone, et al. *Arterioscler. Thromb. Vasc. Biol.* **2001**, *21*, 327). Mice deficient in CCR2 or in MCP-1 showed reductions in the intimal area and in the intima/media ratio (relative to wildtype littermates) after arterial injury (Merce Roque, et al. *Arterioscler. Thromb. Vasc. Biol.* **2002**, *22*, 554; A. Schober, et al. *Circ. Res.* **2004**, *95*, 1125; W. J. Kim, et al. *Biochem Biophys Res Commun.* **2003**, *310*, 936). In mice, transfection of a dominant negative inhibitor of MCP-1 in the skeletal muscle (K. Egashira, et al. *Circ. Res.* **2002**, *90*, 1167) also reduced intimal hyperplasia after arterial injury. Blockade of CCR2 using a neutralizing antibody reduced neointimal hyperplasia after stenting in primates (C. Horvath, et al. *Circ. Res.* **2002**, *90*, 488).
- [0022] Two reports describe the overexpression of MCP-1 rats with induced brain trauma (J. S. King, et al., *J. Neuroimmunol.* **1994**, *56*, 127, and Joan W. Berman, et al., *J. Immunol.* **1996**, *156*, 3017). In addition, studies have shown that both CCR2^{-/-} (O. B. Dimitrijevic, et al. *Stroke* **2007**, *38*, 1345) and MCP-1^{-/-} mice (P. M. Hughes, et al. *J. Cereb. Blood Flow Metab.* **2002**, *22*, 308) are partially protected from ischemia/reperfusion injury.
- [0023] It is known that monocytes/macrophages play an important role in the development of neuropathic pain (Liu T, van Rooijen N, Tracey DJ, *Pain* **2000**, *86*, 25). Consistent with this notion, a potential role for CCR2 in the treatment of both inflammatory and neuropathic pain has been described recently. CCR2^{-/-} mice showed altered responses to inflammatory pain relative to their WT counterparts, including reduced pain behavior after intraplantar formalin injection and slightly reduced mechanical allodynia after intraplantar CFA injection (C. Abbadie, et al. *Proc. Natl. Acad. Sci., USA* **2003**, *100*, 7947). In addition, CCR2^{-/-} mice did not display significant mechanical allodynia after sciatic nerve injury. Likewise, a small molecule CCR2 antagonist reduced mechanical allodynia to ~80% of pre-injury levels after oral administration (C. Abbadie, J. A. Lindia, and H. Wang, WO PCT 110376, 2004).
- [0024] One study described the critical role of MCP-1 in ischemic cardiomyopathy (N. G. Frangogiannis, et al., *Circulation* **2007**, *115*, 584). Another

study described the attenuation of experimental heart failure following inhibition of MCP-1 (S Hayashidani et al., *Circulation* **2003**, *108*, 2134).

[0025] Other studies have provided evidence that MCP-1 is overexpressed in various disease states not mentioned above. These reports provide correlative evidence that MCP-1 antagonists could be useful therapeutics for such diseases. Another study has demonstrated the overexpression of MCP-1 in rodent cardiac allografts, suggesting a role for MCP-1 in the pathogenesis of transplant arteriosclerosis (Mary E. Russell, et al. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 6086). The overexpression of MCP-1 has been noted in the lung endothelial cells of patients with idiopathic pulmonary fibrosis (Harry N. Antoniades, et al., *Proc. Natl. Acad. Sci. USA* **1992**, *89*, 5371). Similarly, the overexpression of MCP-1 has been noted in the skin from patients with psoriasis (M. Deleuran, et al., *J. Dermatol. Sci.* **1996**, *13*, 228, and R. Gillitzer, et al., *J. Invest. Dermatol.* **1993**, *101*, 127); correlative findings with predominance of CCR2+ cells have also been reported (C. Vestergaard, et al. *Acta Derm. Venerol.* **2004**, *84*, 353). Finally, a recent report has shown that MCP-1 is overexpressed in the brains and cerebrospinal fluid of patients with HIV-1-associated dementia (Alfredo Garzino-Demo, WO 99/46991).

[0026] In addition, CCR2 polymorphism has been shown to be associated with sarcoidosis at least in one subset of patients (P. Spagnolo, et al. *Am J Respir Crit Care Med.* **2003**, *168*, 1162).

[0027] It should also be noted that CCR-2 has been implicated as a co-receptor for some strains of HIV (B. J. Doranz, et al., *Cell* **1996**, *85*, 1149). It has also been determined that the use of CCR-2 as an HIV co-receptor can be correlated with disease progression (Ruth I. Connor, et al., *J. Exp. Med.* **1997**, *185*, 621). This finding is consistent with the recent finding that the presence of a CCR-2 mutant, CCR2-64I, is positively correlated with delayed onset of HIV in the human population (Michael W. Smith, et al., *Science* **1997**, *277*, 959). Although MCP-1 has not been implicated in these processes, it may be that MCP-1 antagonists that act via binding to CCR-2 may have beneficial therapeutic effects in delaying the disease progression to AIDS in HIV-infected patients.

[0028] It should be noted that CCR2 is also the receptor for the human chemokines MCP-2, MCP-3, and MCP-4 (Luster, *New Eng. J. Med.* **1998**, *338*, 436-

445). Since the new compounds of formula (I) described herein antagonize MCP-1 by binding to the CCR-2 receptor, it may be that these compounds of formula (I) are also effective antagonists of the actions of MCP-2, MCP-3, and MCP-4 that are mediated by CCR-2. Accordingly, when reference is made herein to “antagonism of MCP-1,” it is to be assumed that this is equivalent to “antagonism of chemokine stimulation of CCR-2.”

[0029] Accordingly, compounds that modulate chemokine activity could demonstrate a wide range of utilities in treating inflammatory, allergic, autoimmune, metabolic, cancer and/or cardiovascular diseases. Patent publication WO2005021500 A1 (incorporated herein by reference and assigned to present applicant) discloses compounds that modulate MCP-1, MCP-2, MCP-3 and MCP-4 activity via CCR2. The reference also discloses various processes to prepare these compounds including multistep syntheses that include the introduction and subsequent removal of protecting groups.

[0030] It is desirable to find new compounds with improved pharmacological characteristics compared with known chemokine modulators. For example, it is desirable to find new compounds with improved CCR-2 inhibitory activity and selectivity for CCR-2 versus other G protein-coupled receptors (i.e. 5HT2A receptor). It is also desirable to find compounds with advantageous and improved characteristics in one or more of the following categories:


- (a) pharmaceutical properties (i.e. solubility, permeability, amenability to sustained release formulations);
- (b) dosage requirements (e.g., lower dosages and/or once-daily dosing);
- (c) factors which decrease blood concentration peak-to-trough characteristics (i.e. clearance and/or volume of distribution);
- (d) factors that increase the concentration of active drug at the receptor (i.e. protein binding, volume of distribution);
- (e) factors that decrease the liability for clinical drug-drug interactions (cytochrome P450 enzyme inhibition or induction, such as CYP 2D6 inhibition, see G.K. Dresser, J.D. Spence, D.G. Bailey, *Clin. Pharmacokinet.* **2000**, 38, 41-57, which is hereby incorporated by reference);

(f) factors that decrease the potential for adverse side-effects (e.g. pharmacological selectivity beyond G protein-coupled receptors, potential chemical or metabolic reactivity, limited CNS penetration, and/or ion-channel selectivity). It is especially desirable to find compounds having a desirable combination of the
5 aforementioned pharmacological characteristics.

[0031] It is also desirable in the art to provide new and/or improved processes to prepare such compounds. These processes may be characterized, without limitation, by a) facile adaptation to larger scale production, such as pilot plant or manufacturing scales; b) process steps and/or techniques enabling improvements in the purity
10 (including chiral purity), stability and/or ease of handling of intermediates and/or final compounds; and/or c) fewer process steps.

SUMMARY OF THE INVENTION

[0032] The present invention provides a novel antagonist or partial
15 agonist/antagonist of MCP-1 receptor activity: N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, or a pharmaceutically acceptable salt, solvate or prodrug, thereof, having an unexpected combination of desirable pharmacological
20 characteristics. Crystalline forms of the present invention are also provided. Pharmaceutical compositions containing the same and methods of using the same as agents for the treatment of inflammatory diseases, allergic, autoimmune, metabolic, cancer and/or cardiovascular diseases is also an objective of this invention. The present disclosure also provides a process for preparing compounds of Formula (I), including N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-
25 (trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide:



wherein R¹, R⁸, R⁹, R¹⁰, and  are as described herein. Compounds that are useful intermediates of the process are also provided herein.

[0033] The present disclosure also provides the use of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, or a pharmaceutically acceptable salt, solvate or prodrug, for the manufacture of a medicament for the treatment of inflammatory diseases, allergic, autoimmune, metabolic, cancer and/or cardiovascular diseases.

BRIEF DESCRIPTION OF THE DRAWINGS

10 [0034] Figure 1. Experimental and simulated powder patterns of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, hemi-hydrate, Form H0.5-4.

[0035] Figure 2. Experimental and simulated powder patterns of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, Form N-2.

15 [0036] Figure 3. Experimental and simulated powder patterns of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, 1.75 moles H₂O, Form H1.75-5.

[0037] Figure 4. Experimental and simulated powder patterns of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, mono-ethanol solvate, Form E-1.

20 [0038] Figure 5. Experimental and simulated powder patterns of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, mono-acetic acid solvate, Form HAC-1.

25 [0039] Figure 6. Experimental and simulated powder patterns of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, mono-*R*-propylene glycol solvate, Form RPG-3.


- [0040] Figure 7. DSC of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, hemihydrate, Form H0.5-4.
- [0041] Figure 8. TGA of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, hemihydrate, Form H0.5-4.
- [0042] Figure 9. DSC of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, Form N-2.
- [0043] Figure 10. TGA of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, Form N-2.
- [0044] Figure 11. Vapor Sorption Isotherm of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, Form N-2 at 25 °C.
- [0045] Figure 12. DSC of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, 1.75 moles H₂O, Form H1.75-5.
- [0046] Figure 13. TGA of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, 1.75 moles H₂O, Form H1.75-5.
- [0047] Figure 14. DSC of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, monoacetic acid solvate, Form HAC-1.
- [0048] Figure 15. TGA of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, monoacetic acid solvate, Form HAC-1.
- [0049] Figure 16. DSC of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, monoethanol solvate, Form E-1.

- [0050] Figure 17. TGA of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, mono-ethanol solvate, Form E-1.
- [0051] Figure 18. DSC of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, mono-
5 R-propylene glycol solvate, Form RPG-3.
- [0052] Figure 19. TGA of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, mono-R-propylene glycol solvate, Form RPG-3.
- 10 [0053] Figure 20. Intradermal challenge model in cynomolgus monkey: Example 1 inhibited mononuclear cell recruitment to skin.
- [0054] Figure 21. 48-hour TG peritonitis in hCCR2 KI mouse: Example 1 inhibition of monocyte/macrophage infiltration into peritoneal cavity.
- [0055] Figure 22. hCCR2 KI mouse EAE (experimental autoimmune
15 encephalomyelitis): Example 1 treatment reduced clinical score.
- [0056] Figure 23. Proton NMR spectra of Example 1, 2nd Alternative Preparation, Step 3 – Compound 7.
- [0057] Figure 24. Proton NMR spectra of Example 1, 2nd Alternative Preparation, Step 4 – Compound 8.
- 20 [0058] Figure 25. Proton NMR spectra of Example 1, 2nd Alternative Preparation, Step 4 – Compound 9.
- [0059] Figure 26. Proton NMR spectra of Example 1, 2nd Alternative Preparation, Step 4 – Compound 10
- [0060] Figure 27. Proton NMR spectra of Example 1, 2nd Alternative
25 Preparation, Step 5 – Compound 11.

DETAILED DESCRIPTION

- [0061] The present invention provides a novel antagonist or partial agonist/antagonist of MCP-1 receptor activity: N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-
30 yl)cyclohexyl)acetamide, or a pharmaceutically acceptable salt, solvate or prodrug, thereof, having an unexpected combination of desirable pharmacological characteristics. Crystalline forms of the present invention are also provided.

Pharmaceutical compositions containing the same and methods of using the same as agents for the treatment of inflammatory, allergic, autoimmune, metabolic, cancer and/or cardiovascular diseases is also an objective of this invention. The present invention also provides a process for preparing compounds of Formula (I), including
 5 N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide:



wherein R^1 , R^8 , R^9 , R^{10} , and HET are as described herein. Compounds that are useful intermediates of the process are also provided herein.

10 **[0062]** [N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, unexpectedly demonstrates a desirable combination of pharmacological characteristics including a surprisingly high degree of oral bioavailability in combination with indications that it is highly efficacious and has excellent safety
 15 criteria.

[0063] Known modulators of CCR2 receptors, such as those disclosed in patent publication WO2005021500 published March 10, 2005 (US Patent No. 7,163,937, issued January 16, 2007, assigned to present Applicant) that demonstrate an adequate degree of membrane permeability (a critical factor of oral bioavailability), are not
 20 sufficiently efficacious, as measured by their CCR2-binding ability (a measure of efficacy), and/or they lack appropriate criteria for safety as indicated by ion channel selectivity as measured by hERG and Na⁺ ion channel studies.

[0064] In contrast, as illustrated by the data presented herein in the section titled "Comparative Pharmacological Characteristics", *infra*, the relatively polar molecule,
 25 N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-

ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide demonstrates a surprisingly high degree of membrane permeability, and yet maintains potent CCR2 binding ability along with excellent ion channel selectivity.

- [0065] Accordingly, the present invention provides a new chemokine modulator
5 having improved pharmacological characteristics that is expected to be useful in treating inflammatory, allergic, autoimmune, metabolic, cancer and/or cardiovascular diseases.

EMBODIMENTS

- [0066] In one embodiment, the disclosure is directed to N-((1R,2S,5R)-5-(*tert*-
10 butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, and pharmaceutically acceptable salts, thereof.

[0067] Another embodiment is a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base.

- [0068] Another embodiment is a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, wherein the crystalline form comprising the N-2
15 Form.

- [0069] Another embodiment is the N-2 Form characterized by (or having) unit
20 cell parameters substantially equal to the following:

Cell dimensions:

$$a = 18.7240(4)$$

$$b = 8.0171(2)$$

$$c = 19.6568(5)$$

25 $\alpha = 90$

$$\beta = 114.935(2)$$

$$\gamma = 90$$

$$V(\text{\AA}^3) = 2675.7(1)$$

Space group $P2_12_12_1$

30 Molecules/unit cell 2

wherein said crystal is at a temperature of about +22°C (RT).

- [0070] Another embodiment in the N-2 Form characterized by (or having) a powder x-ray diffraction pattern comprising three or more of 2θ values ($\text{CuK}\alpha$ $\lambda=1.5418\text{ \AA}$) selected from 5.5, 9.1, 12.1, 14.0 and 19.2, at a temperature of about 22°C .
- 5 [0071] Another embodiment is the N-2 Form characterized by (or having) a powder x-ray diffraction pattern further comprising four or more of 2θ values ($\text{CuK}\alpha$ $\lambda=1.5418\text{ \AA}$) selected from 5.5, 9.1, 12.1, 14.0 and 19.2 at a temperature of about 22°C .
- [0072] Another embodiment is the N-2 Form characterized by (or having)
- 10 fractional atomic coordinates substantially as listed in Table 3.
- [0073] Another embodiment is the N-2 Form characterized by (or having) a power x ray diffraction pattern substantially according to Figure 2.
- [0074] Another embodiment is a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, comprising Form H1.75-5 (1.75 moles of water),
- 15 characterized by the unit cell parameters found in Table 1; 3 or 4 or more 2θ values ($\text{CuK}\alpha$ $\lambda=1.5418\text{ \AA}$) selected from Table 9; fractional atomic coordinate substantially as listed in Table 4 and/or a powder x-ray diffraction pattern substantially according to Figure 3.
- 20 [0075] Another embodiment is a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, comprising Form H0.5-4 (hemi-hydrate) characterized by the unit cell parameters found in Table 1; 3 or 4 or more 2θ values ($\text{CuK}\alpha$ $\lambda=1.5418\text{ \AA}$) selected from Table 9; fractional atomic coordinate substantially
- 25 as listed in Table 2 and/or a powder x-ray diffraction pattern substantially according to Figure 1.
- [0076] Another embodiment is a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, comprising Form E-1 (mono-ethanol solvate)
- 30 characterized by the unit cell parameters found in Table 1; 3 or 4 or more 2θ values ($\text{CuK}\alpha$ $\lambda=1.5418\text{ \AA}$) selected from Table 9; fractional atomic coordinate substantially

as listed in Table 5 and/or a powder x-ray diffraction pattern substantially according to Figure 4.

[0077] Another embodiment is a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, comprising Form HAC-1 (mono-acetic acid solvate) characterized by the unit cell parameters found in Table 1; 3 or 4 or more 2 θ values (CuK α λ =1.5418Å) selected from Table 9; fractional atomic coordinate substantially as listed in Table 6 and/or a powder x-ray diffraction pattern substantially according to Figure 5.

10 [0078] Another embodiment is a crystalline form patterns of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, comprising Form IPA-1 (mono-isopropanol solvate) characterized by the unit cell parameters found in Table 1; 3 or 4 or more 2 θ values (CuK α λ =1.5418Å) selected from Table 9; and/or
15 fractional atomic coordinates substantially as listed in Table 7.

[0079] Another embodiment is a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base comprising Form RPG-3 (mono-*R*-propylene glycol solvate) characterized by the unit cell parameters found in Table 1; 3 or 4 or
20 more 2 θ values (CuK α λ =1.5418Å) selected from Table 9; fractional atomic coordinate substantially as listed in Table 8 and/or a powder x-ray diffraction pattern substantially according to Figure 6.

[0080] Another embodiment is a pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a compound of the Examples.

25 [0081] Another embodiment is a method for modulation of chemokine or chemokine receptor activity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0082] Another embodiment is a method for modulation of CCR-2 receptor activity comprising administering to a patient in need thereof a therapeutically
30 effective amount of a compound of the Examples.

[0083] Another embodiment is a method for modulation of MCP-1, MCP-2, MCP-3 and MCP-4, and MCP-5 activity that is mediated by the CCR2 receptor

comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0084] Another embodiment is a method for modulation of MCP-1 activity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0085] Another embodiment is a method for inhibiting CCR2 and CCR5 activity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0086] Another embodiment is a method for treating disorders, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples said disorders being selected from diabetes, obesity, metabolic syndrome, stroke, neuropathic pain, ischemic cardiomyopathy, psoriasis, hypertension, scleroderma, osteoarthritis, aneurism, fever, cardiovascular disease, Crohn's disease, congestive heart failure, autoimmune diseases, HIV-infection, HIV-associated dementia, psoriasis, idiopathic pulmonary fibrosis, transplant arteriosclerosis, physically- or chemically-induced brain trauma, inflammatory bowel disease, alveolitis, colitis, systemic lupus erythematosus, nephrotoxic serum nephritis, glomerulonephritis, asthma, multiple sclerosis, atherosclerosis, vasculitis, vulnerable plaques, rheumatoid arthritis, restenosis, venous neointimal hyperplasia, dialysis-graft neointimal hyperplasia, arterio-venous shunt intimal hyperplasia, organ transplantation, chronic allograft nephropathy, and cancer.

[0087] Another embodiment is a method for treating disorders, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples, wherein said disorders are selected from diabetes, obesity, Crohn's disease, psoriasis, idiopathic pulmonary fibrosis, transplant arteriosclerosis, physically- or chemically-induced brain trauma, inflammatory bowel disease, alveolitis, colitis, systemic lupus erythematosus, nephrotoxic serum nephritis, glomerulonephritis, asthma, multiple sclerosis, atherosclerosis, and rheumatoid arthritis, restenosis, organ transplantation, and cancer.

[0088] Another embodiment is a method for treating disorders, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples, wherein said disorders are selected from diabetes,

obesity, Crohn's disease, systemic lupus erythematosus, glomerulonephritis, multiple sclerosis, atherosclerosis, restenosis, and organ transplantation.

[0089] Another embodiment is a method for treating disorders, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples, wherein said disorders are selected from multiple sclerosis, atherosclerosis, Crohn's disease, and diabetes.

[0090] Another embodiment is a method for treating disorders, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples, wherein said disorders are selected from restenosis, organ transplantation, and cancer.

[0091] Another embodiment is a method for treating diabetes, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0092] Another embodiment is a method for treating Crohn's disease, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0093] Another embodiment is a method for treating multiple sclerosis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0094] Another embodiment is a method for treating atherosclerosis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0095] Another embodiment is a method for treating restenosis, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

[0096] Another embodiment is a method for treating organ transplantation, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.


[0097] Another embodiment is a method for treating cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.

- [0098] Another embodiment is a method for treating cancer, wherein the cancer is selected from breast cancer, liver cancer, prostate cancer and melanoma.
- [0099] Another embodiment is a method for treating inflammatory, allergic, autoimmune, metabolic, cancer and/or cardiovascular diseases comprising
5 administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.
- [00100] Another embodiment is a method for treating inflammatory, allergic, autoimmune, metabolic, cancer and/or cardiovascular diseases which are at least partially mediated by CCR-2, comprising administering to a patient in need thereof a
10 therapeutically effective amount of a compound of the Examples.
- [00101] Another embodiment is a method for modulation of CCR2 activity comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.
- [00102] Another embodiment is a method for modulation of MIP-1 β and RANTES
15 activity that is mediated by the CCR5 receptor comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the Examples.
- [00103] Another embodiment is a compound of Examples in the preparation of a medicament for the treatment of diabetes, obesity, metabolic syndrome, stroke, neuropathic pain, ischemic cardiomyopathy, psoriasis, hypertension, scleroderma,
20 osteoarthritis, aneurism, fever, cardiovascular disease, Crohn's disease, congestive heart failure, autoimmune diseases, HIV-infection, HIV-associated dementia, psoriasis, idiopathic pulmonary fibrosis, transplant arteriosclerosis, physically- or chemically-induced brain trauma, inflammatory bowel disease, alveolitis, colitis, systemic lupus erythematosus, nephrotoxic serum nephritis, glomerulonephritis,
25 asthma, multiple sclerosis, atherosclerosis, vasculitis, vulnerable plaques, rheumatoid arthritis, restenosis, venous neointimal hyperplasia, dialysis-graft neointimal hyperplasia, arterio-venous shunt intimal hyperplasia, organ transplantation, chronic allograft nephropathy, and cancer.
- [00104] Another embodiment is a compound of the Examples for use in therapy.
- 30 [00105] The invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. This invention also encompasses all combinations of alternative aspects and embodiments of the invention noted herein.

It is understood that any and all embodiments may be taken in conjunction with any other embodiment to describe additional embodiments of the present invention. Furthermore, any elements of an embodiment (including preferred aspects) are meant to be combined with any and all other elements from any of the embodiments to describe additional embodiments.

PROCESS EMBODIMENTS


[00106] The present disclosure also provides a novel process for making compounds of formula I:



including N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, or a pharmaceutically acceptable salt thereof.

[00107] In a 1st embodiment, the present disclosure provides a novel process for preparing a compound of formula IV, the process comprising:

coupling an amino acid derivative of structure III, or a salt thereof, with a cyclohexanone of formula II, or a salt thereof (see preparation in WO2005021500), to afford a compound of structure IV, or a salt thereof having a substituted amide side chain



II


III

IV

wherein:

R_a and R_b are independently C_{1-6} alkoxy;

or R_a and R_b together with the carbon to which they are both attached combine
5 to form a carbonyl, a thiocarbonyl, a cyclic acetal or cyclic thioacetal, wherein the
cyclic acetal or cyclic thioacetal is selected from -O-Z-O- and -S-Z-S-, Z is -



$(CT_1T_2)_2$, $-(CT_1T_2)_3$, or $(T_3)_{0-4}$, and T_1 , T_2 and T_3 at each occurrence is
independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, halogen, hydroxy,
cyano, nitro, CF_3 , OC_{1-4} alkyl, OCF_3 , and $C(=O)C_{1-4}$ alkyl (preferably T_1 , T_2 and T_3 are
10 each hydrogen);

R_1 , R_2 and R_3 are independently hydrogen or an amine-protecting group;

R_4 is lower C_{1-6} alkyl or optionally substituted benzyl;

Y is halogen, SMe, $S(Me)^+R_{12}$, or OSO_2R_{13} ;

V is OH, halogen or OSO_2R_{13} ;

15 R_{12} is hydrogen, C_{1-6} alkyl, $-(CH_2)C(O)O C_{1-6}$ alkyl, or $-(CH_2)C(O)O C_{1-6}$ alkyl; and

R_{13} at each occurrence is C_{1-6} alkyl.

[00108] Preferred amine-protecting groups are groups that can be removed by
hydrolysis or hydrogenolysis under standard conditions. Such groups include without
20 limitation, a carbobenzyloxy (Cbz) group, a tert-butyloxycarbonyl (BOC), a
fluorenylmethyloxycarbonyl (Fmoc) group, a benzyl (Bn) group or a p-
methoxybenzyl (PMB) group. More preferred are Cbz, BOC, or Bn groups
(especially Cbz and Bn).

[00109] In a 2nd embodiment, the present disclosure provides a novel process
25 wherein:

R_a and R_b together with the carbon atoms to which they are both attached
combine to form a carbonyl or a 1, 3-dioxolane group (especially a 1,3-dioxolane
group);

R_1 is hydrogen;

30 R_2 is Cbz;

R₃ is hydrogen;

R₄ is C₁₋₆alkyl;

Y is S(Me); and

V is OH.

5 [00110] In a 3rd embodiment, where Y is -SMe, the disclosure provides a process further comprising alkylating the compound of formula IV with a group R₁₂X, where X is a halogen, to form a sulfonium salt thereof. The alkylating agent is preferably methyl iodide.

[00111] In a 4th embodiment, the disclosure provides a process wherein
10 cyclohexanone of formula IV is a toluenesulfonate salt.

[00112] In a 5th embodiment, the disclosure provides a process wherein the compound of formula IV is a sulfonium salt.


[00113] In a 6th embodiment, the disclosure provides a process wherein the coupling is conducted under an inert atmosphere, such as nitrogen or argon
15 (preferably nitrogen) in an aprotic solvent such as propionitrile, isopropyl acetate, n-butyl acetate, tert-butyl acetate or acetonitrile (especially acetonitrile and/or ethyl acetate).

[00114] In a 7th embodiment, the disclosure provides a process wherein the coupling can be achieved by the contacting a compound of formula II with a diimide
20 coupling reagent in the presence of an activator, and a tertiary amine base. The diimide coupling reagent includes reagents, for example such as EDAC. Examples of activators includes HOBt ((said term includes hydrates thereof) and N',N'-4-dimethylamino-pyridine. A tertiary amine base, includes for example, triethylamine, N,N-disopropyl-N-ethyl amine and tri-n-propylamine.

25 [00115] In a 8th embodiment, the disclosure provides a process wherein the diimide coupling reagent is EDAC, the activator, is HOBt, (said term includes hydrates thereof) and the tertiary amine base is triethylamine.

[00116] In a 9th embodiment, disclosure provides a process wherein the mole ratios of a compound of formula II to the diimide coupling reagent to the activator to the
30 tertiary amine about one to about 0.090 – 1.50 to about 0.95-1.50 to about 2.00 to 3.00, respectively. Said mole ratios are preferably one to about 0.095 – 1.05 to about 0.95-1.10 and to about 2.10 to 2.20, respectively.

[00117] In a 10th embodiment, the disclosure provides a novel process for preparing a compound of formula V having an ester moiety:



the process, comprising:


- 5 cyclizing the amino acid derivative side chain of a compound of formula IV, or a salt thereof, to afford a compound of formula V having an ester moiety.

[00118] In an 11th embodiment where R_a and R_b are independently C_{1-6} alkoxy, or R_a and R_b together with the carbon to which they are both attached combine to form a thiocarbonyl, a cyclic acetal or cyclic thioacetal, the disclosure provides a process
10 which optionally further comprises the step of hydrolyzing the R_a and R_b groups so that the combination of the R_a and R_b together with the carbon to which they are both attached form a carbonyl. Hydrolyzing can be conducted in a solvent such as acetone, butanone, acetonitrile and isopropanol, or aqueous solutions thereof, and is preferably conducted in aqueous acetone. Preferably, the hydrolysis step follows the
15 cyclization step.

[00119] In a 12th embodiment, the disclosure provides a process wherein the cyclization is conducted by combining a compound of formula IV, or a salt thereof, with a base in the presence of a solvent. Such bases may be, for example without limitation, cesium carbonate, cesium bicarbonate, potassium carbonate, sodium tert-butylate, sodium hexamethyldisilazide, and preferably cesium carbonate.
20

[00120] In a 13th embodiment, the disclosure provides a process wherein the cyclization is conducted under an inert atmosphere, such as nitrogen or argon (preferably nitrogen) in a solvent including, for example without limit, DMSO, DMF, DMA, N-methylpyrrolidone, sulfolane (especially DMSO and/or DMF).


25 [00121] In a 14th embodiment, the disclosure provides process for preparing a compound of formula VI:



the process comprising:


hydrolyzing the ester moiety of the compound of formula V with a hydrolyzing agent to form the acid of compound VI at temperatures from about -5 to about 5 °C. Ester hydrolyzing agents are well known to those of skill in the art and include alkali metal hydroxides, MOH, where M is Li, Na or K, preferably the hydrolyzing agent is aqueous NaOH. Preferably the hydrolyzing step is performed under biphasic conditions with an organic solvent that is partially miscible in water. Preferred organic solvents are acyclic or cyclic ethers including THF, 2-methyl THF, 1,2-dimethoxyethane, 1,4-dioxane, especially THF.

[00122] Alternatively, in a 15th embodiment, where R_a and R_b together with the atom to which they are both attached combine to form a carbonyl, the disclosure provides a process for preparing a compound of formula VII:




the process comprising:

reacting a compound of formula VIa, HO-Z-OH, with a compound of formula IV (optionally *in situ*) in the presence of an acid catalyst to give a compound of formula VII,

wherein Z is $-(CT_1T_2)_2-$, $-(CT_1T_2)_3-$, or ; and
 T_1 , T_2 and T_3 at each occurrence is independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, halogen, hydroxy, cyano, nitro, CF_3 , OC_{1-4} alkyl, OCF_3 , and $C(=O)C_{1-4}$ alkyl (preferably T_1 , T_2 and T_3 are each hydrogen).

- 5 Preferably the compound of formula VIa is ethylene glycol and the acid catalyst is p-toluenesulfonic acid, or a hydrate thereof.

[00123] In a 16th embodiment, the disclosure provides a process for preparing a compound of formula VIII:




- 10 the process comprising:

transforming the ketal of formula VII to an intermediate isocyanate of formula VIII. The transformation is preferably conducted via a Curtius rearrangement.

[00124] In a 17th embodiment, the disclosure provides a process wherein the transforming step is conducted via a Curtius rearrangement comprising the steps of:


- 15 a) mixing a substantially anhydrous solution of formula VII with a base (the base is for example, without limitation, an alkylamine, especially a tertiary amine, preferably triethylamine);
- b) adding a haloformate (for example, a chloroformate, preferably $i\text{-BuO}_2\text{CCl}$) to the solution at a temperature of from about $-10\text{ }^\circ\text{C}$ to about $0\text{ }^\circ\text{C}$ to form a mixed anhydride of the acid of formula VII;
- 20 c) treating the mixed anhydride with an azide reagent (preferably NaN_3) in the presence of a phase transfer catalyst (preferably a tetralkylammonium salt such as tetrabutylammonium bromide at about 5 mol%) at a temperature of about $-10\text{ }^\circ\text{C}$ to about $0\text{ }^\circ\text{C}$ to form the acid azide of formula VIIa:



VIIa ; and

d) heating a substantially anhydrous solution of the acyl azide of formula VIIa to form the corresponding isocyanate of formula VIII. Preferably the substantially anhydrous solution of acyl azide is dried over molecular sieves.

- 5 [00125] In a 18th embodiment, the disclosure provides a process for preparing a compound of formula IX having a ketal moiety:




the process comprising:

- 10 contacting the isocyanate of formula VIII (optionally *in situ*) with a compound of formula R₁₀COW (e.g. acetic acid) in the presence of a corresponding acid anhydride (i.e. (R₁₀CO)₂O) to form the amide of formula IX wherein:

R₁₀ is C₁₋₆alkyl (R₁₀ is preferably methyl); and

W is OH or OC₁₋₆alkyl.


- 15 [00126] In a 19th embodiment the disclosure provides a process for preparing a compound of formula X:



the process comprising:

hydrolyzing the ketal moiety of the amide of formula IX to form a compound of formula X. Ketal hydrolyzing conditions and reagents are well known to those of skill in the art. Preferably hydrolysis is conducted by heating a solution of
 5 compound IX having a ketal moiety in an organic solvent (for example acetone) and hydrochloric acid (about 1 N) at about 45 °C to about 55 °C for about 2-4 hours.

[00127] In a 20th embodiment, the disclosure provides a process for preparing a compound of formula XI:




10

the process comprising:

reductively aminating the compound of formula X with an amine of formula HNR_8R_9 in the presence of a Lewis Acid followed by a reducing agent to form a compound of formula XI having a pyrrolidonyl amine moiety.

5 [00128] In a 21st embodiment, the disclosure provides a process wherein reductively aminating comprises the steps of:

a) adding a Lewis Acid (preferably a titanium reagent including, without limit, $\text{TiCl}_2(\text{O-iso-propyl})_2$) to a solution of compound X and the amine having the formula HNR_8R_9 in an aprotic solvent to form an imine-enamine of formula XA:



Xa


; and

b) treating the imine-enamine of formula Xa with a reducing agent (preferably borane dimethyl sulfide) to afford the compound of formula XI having a pyrrolindonyl amine moiety.

- 5 In the foregoing steps, the aprotic solvent can be for example, without limitation, dichloromethane, acetonitrile, DMSO, DMF, and N-methyl-pyrrolidinone (preferably dichloromethane).

[00129] In a 22nd embodiment, the disclosure provides a process wherein the amine of formula HN(R₈)(R₉) is preferably *tert*-butyl amine.

- 10 [00130] In a 23rd embodiment, the disclosure provides process for preparing a compound of formula XII:



XII

the process comprising:

- 15 deprotecting the pyrrolidonyl amine of formula XI to form a compound of formula XII.

[00131] In a 24th embodiment, the disclosure provides a process wherein the deprotecting step is carried out by hydrogenating a solution of the compound of formula XII in the presence of a catalyst such as palladium. Preferably the

hydrogenation is carried out at about 20 to about 40 psig in a solvent, including, without limitation, methanol, over 5% Pd/C catalyst at about 25 °C for about two to about six hours.

[00132] In a 25th embodiment, the disclosure provides a process for preparing a compound of formula I comprising coupling the compound of formula XII with a


compound of formula  to afford a compound of formula I wherein:

HET is a 3-14 membered heteroaryl ring having one to four heteroatoms selected from N, O or S (preferably one to three heteroatoms, especially one to two nitrogen atoms) in at least one of the rings (HET is preferably a 6-substituted quinazolin-4-yl, more preferably 6-trifluoromethyl-quinazolin-4-yl); and

LG is a leaving group selected from halogen or OSO₂R₁₆, wherein R₁₆ is phenyl, a 5- to 7-membered heteroaryl having one or more atoms selected from N, S, or O, C₁₋₆alkyl, or a 3- to 7-membered cycloalkyl, all of which are optionally substituted by one to three groups selected from halogen, CF₃ and C₁₋₆alkyl (preferably LG is a halogen, especially chlorine).

[00133] A leaving group as used herein includes, without limitation, groups such as halogens, mesylate, nonaflates, sulfonates, tosylates and triflates. A preferred leaving group is halogen or OSO₂R₁₆, wherein R₁₆ is phenyl, a 5- to 7-membered heteroaryl having one or more atoms selected from N, S or O, C₁₋₆alkyl, or a 3- to 7-membered cycloalkyl, all of which are optionally substituted by one to three groups selected from halogen, CF₃, and C₁₋₆alkyl. In the most preferred embodiment LG is a halogen, especially chlorine.


[00134] In a 26th embodiment, the disclosure provides a process for preparing a compound of formula X:




XI


comprising the steps of:

hydrolyzing the ester moiety of the compound of formula V with a hydrolyzing agent to form the acid of compound VI at temperatures from about -5 to about 5 °C:




reacting a compound of formula VIa, HO-Z-OH, with a compound of formula IV (optionally *in situ*) in the presence of an acid catalyst to give a compound of formula VII having a carboxylic acid moiety:






transforming the carboxylic acid moiety of the ketal of formula VII to a corresponding intermediate isocyanate of formula VIII:




VIII;

contacting the isocyanate of formula VIII (optionally *in situ*), with a compound of formula $R_{10}COW$ (e.g. acetic acid) in the presence of a corresponding acid anhydride (i.e. $(R_{10}CO)_2O$) to form the amide of formula IX having a ketal moiety:



IX; and

hydrolyzing the ketal moiety of the amide of formula IX to form the compound of formula X:



X;

wherein:

R₁, and R₂ are independently hydrogen or an amine-protecting group;

5 R₄ and R₁₀ are independently C₁₋₆alkyl or optionally substituted benzyl;

R₈ and R₉ are independently hydrogen or C₁₋₆alkyl;

W is OH or OC₁₋₆alkyl;



Z is -(CT₁T₂)₂-, -(CT₁T₂)₃-, or (T₃)₀₋₄; and

T₁, T₂ and T₃ at each occurrence are independently selected from hydrogen,

10 C₁₋₄alkyl, C₂₋₄alkenyl, halogen, hydroxy, cyano, nitro, CF₃, OC₁₋₄alkyl, OCF₃, and C(=O)C₁₋₄alkyl.

[00135] In a 27th embodiment, the disclosure provides a process for preparing a compound of formula XI wherein:

the compound of formula VII is (7*R*,8*S*)-8-((3*S*)-3-

15 (((benzyloxy)carbonyl)amino)-2-oxo-1-pyrrolidinyl)-1,4-dioxaspiro[4.5]decane-7-carboxylic acid, or a salt thereof;

the compound of formula VIIa is benzyl ((3*S*)-1-((7*R*,8*S*)-7-(azidocarbonyl)-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;


the compound of formula VIII is benzyl ((3*S*)-1-((7*R*,8*S*)-7-isocyanato-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;

20 the compound of formula IX is benzyl ((3*S*)-1-((7*R*,8*S*)-7-acetamido-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;

the compound of formula X is benzyl ((3*S*)-1-((1*S*,2*R*)-2-acetamido-4-oxocyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof; and

25 the compound of formula XI is benzyl ((3*S*)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.

[00136] In a 28th embodiment, the disclosure provides a process for preparing a compound of formula I:



5

the process comprising:

hydrolyzing the ester moiety of the compound of formula V with a hydrolyzing agent (preferably an alkali hydroxide, especially sodium hydroxide) to form a compound of formula VI;

reacting a compound of formula Va, HO-Z-OH, with the compound of formula VI, optionally *in situ*, in the presence of acid catalyst (preferably p-toluenesulfonic acid, or a hydrate thereof) to afford a compound of formula VII having a carboxylic acid moiety;

transforming the carboxylic acid moiety of the ketal of formula VII to form a corresponding intermediate isocyanate of formula VIII;

contacting the isocyanate of formula VIII, optionally *in situ*, with a compound of formula $R_{10}COW$ (preferably acetic acid) in the presence of a corresponding acid anhydride, $(R_{10}CO)_2O$ (preferably acetic anhydride), to form the amide of formula IX having a ketal moiety;

hydrolyzing the ketal moiety of the amide of formula IX to form the compound of formula X; and

reductively aminating the compound of formula X with an amine of formula HNR_8R_9 (preferably *tert*-butyl amine) in the presence of a Lewis acid (preferably a titanium reagent such as $TiCl_2(O\text{-}iso\text{-}propyl)_2$) to form a compound of formula XI having a pyrrolidinyl amine moiety;

deprotecting the pyrrolidinyl amine of formula XI to form a compound of formula XII; and


coupling the compound of formula XII with a compound of formula



to afford a compound of formula I;

wherein:

R_1 , and R_2 are independently hydrogen or an amine-protecting group;
 R_4 and R_{10} are independently C_{1-6} alkyl or optionally substituted benzyl;
 R_8 and R_9 are independently hydrogen or C_{1-6} alkyl;
 W is OH or OC_{1-6} alkyl;


Z is $-(CT_1T_2)_{2-}$, $-(CT_1T_2)_{3-}$, or  $(T_3)_{0-4}$;

T₁, T₂ and T₃ at each occurrence is independently selected from hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, halogen, hydroxy, cyano, nitro, CF₃, OC₁₋₄alkyl, OCF₃, and C(=O)C₁₋₄alkyl (preferably Z is -(CH₂)₂-);

HET is a 3-14 membered heteroaryl ring having one to four heteroatoms selected from N, O or S (preferably one to three heteroatoms, especially one to two nitrogen atoms) in at least one of the rings (HET is preferably a 6-substituted quinazolin-4-yl, more preferably 6-trifluoromethyl-quinazolin-4-yl); and

LG is a leaving group selected from halogen or OSO₂R₁₆, wherein R₁₆ is phenyl, a 5- to 7-membered heteroaryl having one or more atoms selected from N, S, or O, C₁₋₆alkyl, or a 3- to 7-membered cycloalkyl, all of which are optionally substituted by one to three groups selected from halogen, CF₃ and C₁₋₆alkyl (preferably LG is a halogen, especially chlorine).


[00137] In a 29th embodiment, the disclosure provides a compound of formula VI, or a salt thereof:



VI

wherein:

R₁ and R₂ are independently selected from hydrogen and an amine-protecting group selected from BOC, Cbz, and benzyl (preferably R₁ is hydrogen and R₂ is Cbz);




Z is -(CT₁T₂)₂-, -(CT₁T₂)₃-, or (T₃)₀₋₄; and

T₁, T₂ and T₃ at each occurrence are independently selected from hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, halogen, hydroxy, cyano, nitro, CF₃, OC₁₋₄alkyl, OCF₃, and C(=O)C₁₋₄alkyl (preferably Z is -(CH₂)₂-).

A preferred compound of formula VI is (1R,2S)-2-((S)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-5-oxocyclohexanecarboxylic acid or a salt thereof.

[00138] In a 30th embodiment, the disclosure provides a compound of formula VI that is (1*R*,2*S*)-2-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-5-oxocyclohexanecarboxylic acid or a salt thereof.


5 [00139] In a 31st embodiment, the disclosure provide a novel compound of formula VII, or a salt thereof:



VII

wherein:

10 R_1 and R_2 are independently selected from hydrogen and an amine-protecting group selected from BOC, Cbz, and benzyl (preferably R_1 is hydrogen and R_2 is Cbz);


Z is $-(CT_1T_2)_2-$, $-(CT_1T_2)_3-$, or  $(T_3)_{0-4}$; and

T_1 , T_2 and T_3 at each occurrence are independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, halogen, hydroxy, cyano, nitro, CF_3 , OC_{1-4} alkyl, OCF_3 , and
15 $C(=O)C_{1-4}$ alkyl (preferably Z is $-(CH_2)_2-$).

A preferred compound of formula VII is (7*R*,8*S*)-8-((3*S*)-3-(((benzyloxy)carbonyl)amino)-2-oxo-1-pyrrolidinyl)-1,4-dioxaspiro[4.5]decane-7-carboxylic acid.

[00140] In a 32nd embodiment, the disclosure provides a compound of formula VII that is (7*R*,8*S*)-8-((3*S*)-3-(((benzyloxy)carbonyl)amino)-2-oxo-1-pyrrolidinyl)-1,4-dioxaspiro[4.5]decane-7-carboxylic acid, or a salt thereof.


[00141] In a 33rd embodiment, the disclosure provides novel compounds of formula VIIa, or a salt thereof:



VIIa

wherein:

- R₁ and R₂ are independently selected from hydrogen and an amine-protecting group selected from BOC, Cbz, and benzyl (preferably R₁ is hydrogen and R₂ is Cbz);




T₁, T₂ and T₃ at each occurrence are independently selected from hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, halogen, hydroxy, cyano, nitro, CF₃, OC₁₋₄alkyl, OCF₃, and C(=O)C₁₋₄alkyl (preferably Z is $-(CH_2)_2-$).

- A preferred compound of formula VIIa is benzyl ((3*S*)-1-((7*R*,8*S*)-7-(azidocarbonyl)-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate.


[00142] In a 34th embodiment, the disclosure provides a compound of formula VIIa that is benzyl ((3*S*)-1-((7*R*,8*S*)-7-(azidocarbonyl)-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.

- [00143] In a 35th embodiment, the disclosure provides a compound of formula VIII, or a salt thereof:



wherein:

R₁ and R₂ are independently selected from hydrogen and an amine-protecting group selected from BOC, Cbz, or benzyl (preferably R₁ is hydrogen and R₂ is Cbz);




T₁, T₂ and T₃ at each occurrence are independently selected from hydrogen,
 5 C₁₋₄alkyl, C₂₋₄alkenyl, halogen, hydroxy, cyano, nitro, CF₃, OC₁₋₄alkyl, OCF₃, and
 C(=O)C₁₋₄alkyl (preferably Z is $-(CH_2)_2-$).

A preferred compound of formula VIII is benzyl ((3*S*)-1-((7*R*,8*S*)-7-isocyanato-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidiny)carbamate.

[00144] In a 36th embodiment, the disclosure provides a compound of formula VIII
 10 that is benzyl ((3*S*)-1-((7*R*,8*S*)-7-isocyanato-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidiny)carbamate, or a salt thereof.


[00145] In a 37th embodiment, the disclosure provides a compound of formula IX,
 or a salt thereof:



15 IX

wherein:

R₁ and R₂ are independently selected from hydrogen and an amine-protecting group selected from BOC, Cbz, or benzyl (preferably R₁ is hydrogen and R₂ is Cbz);




20 T₁, T₂ and T₃ at each occurrence are independently selected from hydrogen,
 C₁₋₄alkyl, C₂₋₄alkenyl, halogen, hydroxy, cyano, nitro, CF₃, OC₁₋₄alkyl, OCF₃, and
 C(=O)C₁₋₄alkyl (preferably Z is $-(CH_2)_2-$); and

R₁₀ is C₁₋₆alkyl (preferably methyl).

A preferred compound of formula IX is benzyl ((3*S*)-1-((7*R*,8*S*)-7-acetamido-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate.

[00146] In a 38th embodiment, the disclosure provides a compound of formula IX that is benzyl ((3*S*)-1-((7*R*,8*S*)-7-acetamido-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.

[00147] In a 39th embodiment, the disclosure provides a compound of formula X, or a salt thereof:



X

10 wherein:


R₁ and R₂ are independently hydrogen or an amine-protecting group selected from BOC, Cbz, and benzyl; and

R₁₀ is C₁₋₆alkyl.

15 Preferably R₁ is hydrogen, R₂ is Cbz and R₁₀ is methyl. A preferred compound of formula X is benzyl ((3*S*)-1-((1*S*,2*R*)-2-acetamido-4-oxocyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate.

[00148] In a 40th embodiment, the disclosure provides a compound of formula X that is benzyl ((3*S*)-1-((1*S*,2*R*)-2-acetamido-4-oxocyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.

20 [00149] In a 40th embodiment, the disclosure provides a compound of formula XI, or a salt thereof:



XI

wherein:

R₁ and R₂ are independently hydrogen or an amine-protecting group selected from BOC, Cbz, and benzyl;


5 R₈ and R₉ are independently hydrogen or C₁₋₆alkyl; and

R₁₀ is C₁₋₆alkyl.

Preferably R₁ is hydrogen, R₂ is Cbz, R₈ is hydrogen, R₉ is *tert*-butyl, and R₁₀ is methyl. A preferred compound of formula XI is benzyl ((3*S*)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate.

10 **[00150]** In a 41st embodiment, the disclosure provides a compound of formula XI that is benzyl ((3*S*)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.

[00151] In a 42nd embodiment, the disclosure provides a compound of formula XII, or a salt thereof:



XII

15

wherein:

R₁ is hydrogen or an amine-protecting group selected from BOC, Cbz, and benzyl;

R₈ and R₉ are independently hydrogen or C₁₋₆alkyl; and

20 R₁₀ is C₁₋₆alkyl.

Preferably R₁ is hydrogen, R₈ is hydrogen, R₉ is tert-butyl, and R₁₀ is methyl.
 A preferred compound of formula XII is *N*-((1*R*,2*S*,5*R*)-2-((3*S*)-3-amino-2-oxo-1-pyrrolidinyl)-5-(*tert*-butylamino)cyclohexyl)acetamide.

[00152] In a 43rd embodiment, the disclosure provides a compound of formula XII
 5 that is *N*-((1*R*,2*S*,5*R*)-2-((3*S*)-3-amino-2-oxo-1-pyrrolidinyl)-5-(*tert*-butylamino)cyclohexyl)acetamide, or a salt thereof.

[00153] In a 44th embodiment, the disclosure provides a compound selected from:
 (7*R*,8*S*)-8-((3*S*)-3-(((benzyloxy)carbonyl)amino)-2-oxo-1-pyrrolidinyl)-1,4-dioxaspiro[4.5]decane-7-carboxylic acid, or a salt thereof;
 10 benzyl ((3*S*)-1-((7*R*,8*S*)-7-(azidocarbonyl)-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;
 benzyl ((3*S*)-1-((7*R*,8*S*)-7-isocyanato-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;
 benzyl ((3*S*)-1-((7*R*,8*S*)-7-acetamido-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;
 15 benzyl ((3*S*)-1-((1*S*,2*R*)-2-acetamido-4-oxocyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof; and
 benzyl ((3*S*)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.

[00154] In a 45th embodiment, the disclosure provides a process wherein:
 a compound of formula VII is (7*R*,8*S*)-8-((3*S*)-3-(((benzyloxy)carbonyl)amino)-2-oxo-1-pyrrolidinyl)-1,4-dioxaspiro[4.5]decane-7-carboxylic acid, or a salt thereof;
 a compound of formula VIIa is benzyl ((3*S*)-1-((7*R*,8*S*)-7-(azidocarbonyl)-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;
 25 a compound of formula VIII is benzyl ((3*S*)-1-((7*R*,8*S*)-7-isocyanato-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;
 a compound of formula IX is benzyl ((3*S*)-1-((7*R*,8*S*)-7-acetamido-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;
 30 the compound of formula X is benzyl ((3*S*)-1-((1*S*,2*R*)-2-acetamido-4-oxocyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof; and

a compound of formula XI is benzyl ((3*S*)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.

[00155] In a 46th embodiment, the disclosure provides a process wherein a compound of formula I is N-((1*R*,2*S*,5*R*)-5-(*tert*-butylamino)-2-((*S*)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide or a salt thereof.

[00156] The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. Thus, the above embodiments should not be considered limiting. Any and all embodiments of the present invention may be taken in conjunction with any other embodiment or embodiments to describe additional embodiments. Each individual element (including preferred aspects) of the embodiments is its own independent embodiment. Furthermore, any element of an embodiment is meant to be combined with any and all other elements from any embodiment to describe an additional embodiment. In addition, the present invention encompasses combinations of different embodiment, parts of embodiments, definitions, descriptions, and examples of the invention noted herein.

DEFINITIONS

[00157] The following are definitions of terms used in this specification and appended claims. The initial definition provided for a group or term herein applies to that group or term throughout the specification and claims, individually or as part of another group, unless otherwise indicated.

[00158] The term "alkyl" refers to straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 6 carbon atoms. When numbers appear in a subscript after the symbol "C", the subscript defines with more specificity the number of carbon atoms that a particular group may contain. For example, "C₁₋₆alkyl" refers to straight and branched chain alkyl groups with one to six carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, and so forth. The subscript "0" refers to a bond. Thus, the term hydroxy(C₀₋₂)alkyl or (C₀₋₂)hydroxyalkyl includes hydroxy, hydroxymethyl and hydroxyethyl. Alkyl groups may be substituted with one to three groups selected from (C₁₋₆)alkyl, (C₂₋₆)alkenyl, hydroxy, halogen, cyano, nitro, CF₃, O(C₁₋₆alkyl), OCF₃, C(=O)H, C(=O)(C₁₋₆alkyl),

- CO₂H, CO₂(C₁₋₆alkyl), NHCO₂(C₁₋₆alkyl), -S(C₁₋₆alkyl), NH₂, NH(C₁₋₆alkyl), N(C₁₋₆alkyl)₂, N(CH₃)₃⁺, SO₂(C₁₋₆alkyl), C(=O)(C₁₋₄alkylene)NH₂, C(=O)(C₁₋₄alkylene)NH(alkyl), C(=O)(C₁₋₄alkylene)N(C₁₋₄alkyl)₂, C₃₋₇cycloalkyl, phenyl, benzyl, phenylethyl, phenyloxy, benzyloxy, naphthyl, a four- to seven-membered heterocyclo, and/ or a five- to six- membered heteroaryl. When a substituted alkyl is substituted with an aryl, heterocyclo, cycloalkyl, or heteroaryl group, said ringed systems are as defined below and thus may have zero, one, two, or three substituents, also as defined below.
- [00159]** When the term “alkyl” is used together with another group, such as in “arylalkyl”, this conjunction defines with more specificity at least one of the substituents that the substituted alkyl will contain. For example, “arylalkyl” refers to a substituted alkyl group as defined above where at least one of the substituents is an aryl, such as benzyl. Thus, the term aryl(C₀₋₄)alkyl includes a substituted lower alkyl having at least one aryl substituent and also includes an aryl directly bonded to another group, *i.e.*, aryl(C₀)alkyl.
- [00160]** The term “alkenyl” refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms and at least one double bond. Alkenyl groups of 2 to 6 carbon atoms and having one double bond are most preferred. Alkenyl groups may be substituted as described above for alkyl groups.
- [00161]** The term “alkynyl” refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms and at least one triple bond. Alkynyl groups of 2 to 6 carbon atoms and having one triple bond are most preferred. Alkynyl groups may be substituted as described above for alkyl groups.
- [00162]** The term “alkylene” refers to bivalent straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms, *e.g.*, {-CH₂-}_n, wherein *n* is 1 to 12, preferably 1-8. Lower alkylene groups, that is, alkylene groups of 1 to 2 carbon atoms, are most preferred. The terms “alkenylene” and “alkynylene” refer to bivalent radicals of alkenyl and alkynyl groups, respectively, as defined above. Alkenylene groups may be substituted as described above for alkyl groups.
- [00163]** The term “alkoxy” refers to an oxygen atom substituted by alkyl, as defined herein. For example, the term “alkoxy” or includes the group -O-C₁₋₆alkyl.

- [00164] When a subscript is used with reference to an alkoxy, thioalkyl or aminoalkyl, the subscript refers to the number of carbon atoms that the group may contain in addition to heteroatoms.
- [00165] It should be understood that the selections for all groups, including for
 5 examples, alkoxy, thioalkyl, and aminoalkyl, will be made by one skilled in the field to provide stable compounds.
- [00166] The term “carbonyl” refers to a bivalent carbonyl group -C(=O)- .
- [00167] The term “acyl” refers to a carbonyl group linked to an organic radical, more particularly, the group C(=O)R_e , as well as the bivalent group $\text{-C(=O)R}_e\text{-}$,
 10 which are linked to organic radicals. The group R_e can be selected from alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, or heteroaryl as defined herein, or when appropriate, the corresponding bivalent group, *e.g.*, alkylene.
- [00168] The term “cycloalkyl” refers to fully saturated and partially unsaturated hydrocarbon rings (and therefore includes “cycloalkenyl rings”) of 3 to 9, preferably
 15 3 to 7 carbon atoms. The term “cycloalkyl” includes such rings having zero, one, two, or three substituents selected from $(\text{C}_{1-4})\text{alkyl}$, $(\text{C}_{2-4})\text{alkenyl}$, halogen, hydroxy, cyano, nitro, CF_3 , $\text{O}(\text{C}_{1-4}\text{alkyl})$, OCF_3 , C(=O)H , $\text{C(=O)(C}_{1-4}\text{alkyl)}$, CO_2H , $\text{CO}_2(\text{C}_{1-4}\text{alkyl})$, $\text{NHCO}_2(\text{C}_{1-4}\text{alkyl})$, $\text{S}(\text{C}_{1-4}\text{alkyl})$, NH_2 , $\text{NH}(\text{C}_{1-4}\text{alkyl})$, $\text{N}(\text{C}_{1-4}\text{alkyl})_2$, $\text{N}(\text{C}_{1-4}\text{alkyl})_3^+$, $\text{SO}_2(\text{C}_{1-4}\text{alkyl})$, $\text{C(=O)(C}_{1-4}\text{alkylene)NH}_2$, $\text{C(=O)(C}_{1-4}\text{alkylene)NH(alkyl)}$,
 20 and/or $\text{C(=O)(C}_{1-4}\text{alkylene)N(C}_{1-4}\text{alkyl)}_2$. The term “cycloalkyl” also includes such rings having a second ring fused thereto (*e.g.*, including benzo, heterocyclo, or heteroaryl rings) or having a carbon-carbon bridge of 3 to 4 carbon atoms.
- [00169] The term “halo” or “halogen” refers to chloro, bromo, fluoro and iodo.
- [00170] The term “haloalkyl” means a substituted alkyl having one or more halo
 25 substituents. For example, “haloalkyl” includes mono, bi, and trifluoromethyl.
- [00171] The term “haloalkoxy” means an alkoxy group having one or more halo substituents. For example, “haloalkoxy” includes OCF_3 .
- [00172] The term “heteroatoms” shall include oxygen, sulfur and nitrogen.
- [00173] The term “aryl” refers to phenyl, biphenyl, fluorenyl, 1-naphthyl and 2-
 30 naphthyl. The term “aryl” includes such rings having zero, one, two or three substituents selected from $(\text{C}_{1-4})\text{alkyl}$, $(\text{C}_{2-4})\text{alkenyl}$, halogen, hydroxy, cyano, nitro, CF_3 , $\text{O}(\text{C}_{1-4}\text{alkyl})$, OCF_3 , C(=O)H , $\text{C(=O)(C}_{1-4}\text{alkyl)}$, CO_2H , $\text{CO}_2(\text{C}_{1-4}\text{alkyl})$,

NHCO₂(C₁₋₄alkyl), S(C₁₋₄alkyl), NH₂, NH(C₁₋₄alkyl), N(C₁₋₄alkyl)₂, N(C₁₋₄alkyl)₃⁺, SO₂(C₁₋₄alkyl), C(=O)(C₁₋₄alkylene)NH₂, C(=O)(C₁₋₄alkylene)NH(alkyl), and/or C(=O)(C₁₋₄alkylene)N(C₁₋₄alkyl)₂.

[00174] The terms “heterocyclo” or “heterocyclic” refers to substituted and unsubstituted non-aromatic (which may be partially or fully saturated) 3- to 15-membered rings having one to four heteroatoms. Such rings can be 3-to 7-membered monocyclic groups, 7-to 11-membered bicyclic groups, and 10-to 15-membered tricyclic groups. Each ring of the heterocyclo group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less, and further provided that the ring contains at least one carbon atom. The fused rings completing bicyclic and tricyclic groups may contain only carbon atoms and may be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms may optionally be oxidized and the nitrogen atoms may optionally be quaternized. The heterocyclo group may be attached at any available nitrogen or carbon atom. The heterocyclo ring may contain zero, one, two or three substituents selected from (C₁₋₄)alkyl, (C₂₋₄)alkenyl, halogen, hydroxy, cyano, nitro, CF₃, O(C₁₋₄alkyl), OCF₃, C(=O)H, C(=O)(C₁₋₄alkyl), CO₂H, CO₂(C₁₋₄alkyl), NHCO₂(C₁₋₄alkyl), S(C₁₋₄alkyl), NH₂, NH(C₁₋₄alkyl), N(C₁₋₄alkyl)₂, N(C₁₋₄alkyl)₃⁺, SO₂(C₁₋₄alkyl), C(=O)(C₁₋₄alkylene)NH₂, C(=O)(C₁₋₄alkylene)NH(alkyl), and/or C(=O)(C₁₋₄alkylene)N(C₁₋₄alkyl)₂. Exemplary heterocyclic groups include azetidiny, pyrrolidiny, oxetanyl, imidazoliny, oxazolidiny, isoxazoliny, thiazolidiny, isothiazolidiny, tetrahydrofuranyl, piperidyl, piperaziny, 2-oxopiperaziny, 2-oxopiperidyl, 2-oxopyrrolodiny, 2-oxoazepiny, azepiny, 4-piperidonyl, tetrahydropyranyl, morpholiny, thiamorpholiny, thiamorpholiny sulfoxide, thiamorpholiny sulfone, 1,3-dioxolane, quinuclidiny, and tetrahydro-1,1-dioxothienyl and the like.

[00175] The term “heteroaryl” refers to substituted and unsubstituted aromatic 3- to 14-membered rings having one to four heteroatoms selected from O, S, or N in at least one of the rings. Said rings can be 5- or 6-membered monocyclic groups, 9- or 10-membered bicyclic groups, and 11- to 14-membered tricyclic groups. Each ring of the heteroaryl group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of

heteroatoms in each ring is four or less and each ring has at least one carbon atom. The fused rings completing the bicyclic and tricyclic groups may contain only carbon atoms and may be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms may optionally be oxidized and the nitrogen atoms may optionally be

5 quaternized. Heteroaryl groups which are bicyclic or tricyclic must include at least one fully aromatic ring but the other fused ring or rings may be aromatic or non-aromatic. The heteroaryl group may be attached at any available nitrogen or carbon atom of any ring. The heteroaryl ring system may contain zero, one, two or three substituents selected from (C₁₋₄)alkyl, (C₂₋₄)alkenyl, halogen, hydroxy, cyano, nitro,

10 CF₃, O(C₁₋₄alkyl), OCF₃, C(=O)H, C(=O)(C₁₋₄alkyl), CO₂H, CO₂(C₁₋₄alkyl), NHCO₂(C₁₋₄alkyl), S(C₁₋₄alkyl), NH₂, NH(C₁₋₄alkyl), N(C₁₋₄alkyl)₂, N(C₁₋₄alkyl)₃⁺, SO₂(C₁₋₄alkyl), C(=O)(C₁₋₄alkylene)NH₂, C(=O)(C₁₋₄alkylene)NH(alkyl), and/or C(=O)(C₁₋₄alkylene)N(C₁₋₄alkyl)₂.

[00176] Exemplary heteroaryl groups include pyrrolyl, pyrazolyl, pyrazolinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, furanyl, thienyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, indolyl, benzothiazolyl, benzodioxolyl, benzoxazolyl, benzothienyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, indoliziny, benzofuranyl, chromonyl, coumarinyl, benzopyranyl, cinnolinyl, quinoxalinyl,

20 indazolyl, pyrrolopyridyl, furopyridyl, dihydroisoindolyl, tetrahydroquinolinyl and the like. Particular heteroaryl groups include, for example, 6-substituted quinazolin-4-yl and 6-trifluoromethyl-quinazolin-4-yl.

[00177] Where a group is referred to as "optionally substituted", the term is defined herein to include both a substituted and unsubstituted group.

25 [00178] The compounds herein described may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins,

30 C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and

may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

5 [00179] One enantiomer of compounds disclosed herein may display superior activity compared with the other. Thus, all of the stereochemistries are considered to be a part of the present invention. When required, separation of the racemic material can be achieved by HPLC using a chiral column or by a resolution using a resolving agent such as camphonic chloride as in Steven D. Young, et al, *Antimicrobial Agents*
10 *and Chemotherapy*, 1995, 2602-2605.

[00180] The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or
15 other problem or complication, commensurate with a reasonable benefit/risk ratio.

[00181] As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or
20 organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic,
25 sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

0 [00182] The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the

free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's*

5 *Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

[00183] Since prodrugs are known to enhance numerous desirable qualities of pharmaceuticals (e.g., solubility, bioavailability, manufacturing, etc.) the compounds of the present invention may be delivered in prodrug form. Thus, the present
10 invention is intended to cover prodrugs of the presently claimed compounds, methods of delivering the same and compositions containing the same. "Prodrugs" are intended to include any covalently bonded carriers which release an active parent drug of the present invention *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs the present invention are prepared by modifying
15 functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of the present invention wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug of the present invention is administered to a mammalian subject, it cleaves to form a free hydroxyl, free amino,
20 or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of the present invention.

[00184] "Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity
25 from a reaction mixture, and formulation into an efficacious therapeutic agent. The present invention is intended to embody stable compounds.

[00185] "Therapeutically effective amount" is intended to include an amount of a compound of the present invention alone or an amount of the combination of compounds claimed or an amount of a compound of the present invention in
30 combination with other active ingredients effective to inhibit MCP-1 or effective to treat or prevent disorders.

[00186] As used herein, “treating” or “treatment” cover the treatment of a disease-state in a mammal, particularly in a human, and include: (a) preventing the disease-state from occurring in a mammal, in particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it; (b) inhibiting the disease-state, i.e., arresting its development; and/or (c) relieving the disease-state, i.e., causing regression of the disease state.

[00187] The names used herein to characterize a specific form, e.g., “N-2”, should not be considered limiting with respect to any other substance possessing similar or identical physical and chemical characteristics, but rather it should be understood that these designations are mere identifiers that should be interpreted according to the characterization information also presented herein.

[00188] The present invention provides, at least in part, crystalline forms of the free base of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, as a novel material, in particular in a pharmaceutically acceptable form. In certain preferred embodiments, crystalline forms of the free base are in substantially pure form. Preferred embodiments of crystalline forms of the free base are disclosed in Example 2 as the E-1, HAC-1, IPA-1, N-2, RPG-3, H0.5-4, and H1.75-5 Forms.

[00189] As used herein “polymorph” refers to crystalline forms having the same chemical composition but different spatial arrangements of the molecules, atoms, and/or ions forming the crystal.

[00190] As used herein “solvate” refers to a crystalline form of a molecule, atom, and/or ions that further contains molecules of a solvent or solvents incorporated into the crystalline structure. The solvent molecules in the solvate may be present in a regular arrangement and/or a non-ordered arrangement. The solvate may comprise either a stoichiometric or nonstoichiometric amount of the solvent molecules. For example, a solvate with a nonstoichiometric amount of solvent molecules may result from partial loss of solvent from the solvate.

[00191] As used herein “amorphous” refers to a solid form of a molecule, atom, and/or ions that is not crystalline. An amorphous solid does not display a definitive X-ray diffraction pattern.

[00192] As used herein, "substantially pure," when used in reference to a crystalline form, means a compound having a purity greater than 90 weight %, including greater than 90, 91, 92, 93, 94, 95, 96, 97, 98 and 99 weight %, and also including equal to about 100 weight % of the compound, based on the weight of the compound. The remaining material comprises other form(s) of the compound, and/or reaction impurities and/or processing impurities arising from its preparation. For example, a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide may be deemed substantially pure in that it has a purity greater than 90 weight %, as measured by means that are at this time known and generally accepted in the art, where the remaining less than 10 weight % of material comprises other form(s) of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide and/or reaction impurities and/or processing impurities.

[00193] Samples of the crystalline forms may be provided with substantially pure phase homogeneity, indicating the presence of a dominant amount of a single crystalline form and optionally minor amounts of one or more other crystalline forms. The presence of more than one crystalline form in a sample may be determined by techniques such as powder X-ray diffraction (PXRD) or solid state nuclear magnetic resonance spectroscopy (SSNMR). For example, the presence of extra peaks in the comparison of an experimentally measured PXRD pattern with a simulated PXRD pattern may indicate more than one crystalline form in the sample. The simulated PXRD may be calculated from single crystal X-ray data. See Smith, D.K., "A FORTRAN Program for Calculating X-Ray Powder Diffraction Patterns," Lawrence Radiation Laboratory, Livermore, California, UCRL-7196 (April 1963).

[00194] Preferably, the crystalline form has substantially pure phase homogeneity as indicated by less than 10%, preferably less than 5%, and more preferably less than 2% of the total peak area in the experimentally measured PXRD pattern arising from the extra peaks that are absent from the simulated PXRD pattern. Most preferred is a crystalline form having substantially pure phase homogeneity with less than 1% of the total peak area in the experimentally measured PXRD pattern arising from the extra peaks that are absent from the simulated PXRD pattern.

[00195] Procedures for the preparation of crystalline forms are known in the art. The crystalline forms may be prepared by a variety of methods, including for example, crystallization or recrystallization from a suitable solvent, sublimation, growth from a melt, solid state transformation from another phase, crystallization from a supercritical fluid, and jet spraying. Techniques for crystallization or recrystallization of crystalline forms from a solvent mixture include, for example, evaporation of the solvent, decreasing the temperature of the solvent mixture, crystal seeding a supersaturated solvent mixture of the molecule and/or salt, freeze drying the solvent mixture, and addition of antisolvents (countersolvents) to the solvent mixture.

10 [00196] The forms may be characterized and distinguished using single crystal X-ray diffraction, which is based on unit cell and intensity measurements of a single crystal of a form at a fixed analytical temperature. A detailed description of unit cell and intensity analysis is provided in Stout & Jensen, X-Ray Structure Determination: A Practical Guide, Macmillan Co., New York (1968), Chapter 3, which is herein

15 incorporated by reference. Alternatively, the unique arrangement of atoms in spatial relation within the crystalline lattice may be characterized according to the observed fractional atomic coordinates. See Stout & Jensen reference for experimental determination of fractional coordinates for structural analysis. Another means of characterizing the crystalline structure is by powder X-ray diffraction analysis in

20 which the experimental or observed diffraction profile is compared to a simulated profile representing pure powder material, both at the same analytical temperature, and measurements for the subject form characterized as a series of 2θ values and intensities.

[00197] The term "negligible weight loss," as employed herein, as characterized

25 by TGA indicates the presence of a neat (non-solvated) crystal form.

[00198] The term "negligible % water uptake," as employed herein, as characterized by moisture-sorption isotherm indicates that the form tested is non-hygroscopic.

[00199] In one embodiment of the invention, a crystalline form of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide is provided in substantially pure form. This crystalline form may be employed in pharmaceutical compositions which may

optionally include one or more other components selected, for example, from the group consisting of excipients, carriers, and one of other active pharmaceutical ingredients or active chemical entities of different molecular structures.

5 [00200] Preferably, the crystalline form has substantially pure phase homogeneity as indicated by less than 10%, preferably less than 5%, and more preferably less than 2% of the total peak area in the experimentally measured PXRD pattern arising from the extra peaks that are absent from the simulated PXRD pattern. Most preferred is a crystalline form having substantially pure phase homogeneity with less than 1% of the total peak area in the experimentally measured PXRD pattern arising from the
10 extra peaks that are absent from the simulated PXRD pattern.

[00201] In another embodiment, a composition is provided consisting essentially of the crystalline forms of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide. The composition of this embodiment may comprise at least 90 weight % of the form,
15 based on its weight in the composition.

[00202] The presence of reaction impurities and/or processing impurities may be determined by analytical techniques known in the art, such as, for example, chromatography, nuclear magnetic resonance spectroscopy, mass spectrometry or infrared spectroscopy.

20 [00203] Crystalline forms may be prepared by a variety of methods, including for example, crystallization or recrystallization from a suitable solvent, sublimation, growth from a melt, solid state transformation from another phase, crystallization from a supercritical fluid, and jet spraying. Techniques for crystallization or recrystallization of crystalline forms from a solvent mixture include, for example,
25 evaporation of the solvent, decreasing the temperature of the solvent mixture, crystal seeding a supersaturated solvent mixture of the molecule and/or salt, freeze drying the solvent mixture, and addition of antisolvents (countersolvents) to the solvent mixture. High throughput crystallization techniques may be employed to prepare crystalline forms including polymorphs.

30 [00204] Crystals of drugs, including polymorphs, methods of preparation, and characterization of drug crystals are discussed in Solid-State Chemistry of Drugs,

S.R. Byrn, R.R. Pfeiffer, and J.G. Stowell, 2nd Edition, SSCI, West Lafayette, Indiana (1999).

[00205] For crystallization techniques that employ solvent, the choice of solvent or solvents is typically dependent upon one or more factors, such as solubility of the compound, crystallization technique, and vapor pressure of the solvent. Combinations of solvents may be employed; for example, the compound may be solubilized into a first solvent to afford a solution, followed by the addition of an antisolvent to decrease the solubility of the compound in the solution and to afford the formation of crystals. An "antisolvent" is a solvent in which the compound has low solubility. Suitable solvents for preparing crystals include polar and nonpolar solvents.

[00206] In one method to prepare crystals, N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide is suspended and/or stirred in a suitable solvent to afford a slurry, which may be heated to promote dissolution. The term "slurry," as used herein, means a saturated solution of the free base, which may also contain an additional amount of the compound to afford a heterogeneous mixture of the compound and a solvent at a given temperature. Suitable solvents in this regard include, for example, polar aprotic solvents and polar protic solvents, and mixtures of two or more of these, as disclosed herein.

[00207] Seed crystals may be added to any crystallization mixture to promote crystallization. As will be clear to the skilled artisan, seeding is used as a means of controlling growth of a particular crystalline form or as a means of controlling the particle size distribution of the crystalline product. Accordingly, calculation of the amount of seeds needed depends on the size of the seed available and the desired size of an average product particle as described, for example, in "Programmed cooling of batch crystallizers," J.W. Mullin and J. Nyvlt, *Chemical Engineering Science* **1971**, 26, 369-377. In general, seeds of small size are needed to effectively control the growth of crystals in the batch. Seeds of small size may be generated by sieving, milling, or micronizing of larger crystals, or by micro-crystallization of solutions. Care should be taken that milling or micronizing of crystals does not result in any

change in crystallinity from the desired crystal form (i.e., change to amorphous or to another polymorph).

[00208] A cooled mixture may be filtered under vacuum, and the isolated solids may be washed with a suitable solvent, such as cold recrystallization solvent, and
5 dried under a nitrogen purge to afford the desired crystalline form. The isolated solids may be analyzed by a suitable spectroscopic or analytical technique, such as SSNMR, DSC, PXRD, or the like, to assure formation of the preferred crystalline form of the product. The resulting crystalline form is typically produced in an amount of greater than about 70 weight % isolated yield, but preferably greater than
10 90 weight % based on the weight of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide originally employed in the crystallization procedure. The product may be co-milled or passed through a mesh screen to de-lump the product, if necessary.

[00209] Crystalline forms may be prepared directly from the reaction medium of
15 the final process step for preparing N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide. This may be achieved, for example, by employing in the final process step a solvent or mixture of solvents from which the free base may be crystallized. Alternatively, crystalline forms may be obtained by distillation or solvent addition techniques.
20 Suitable solvents for this purpose include any of those solvents described herein, including protic polar solvents, such as alcohols, and aprotic polar solvents, such as ketones.

[00210] By way of general guidance, the reaction mixture may be filtered to remove any undesired impurities, inorganic salts, and the like, followed by washing
25 with reaction or crystallization solvent. The resulting solution may be concentrated to remove excess solvent or gaseous constituents. If distillation is employed, the ultimate amount of distillate collected may vary, depending on process factors including, for example, vessel size, stirring capability, and the like. By way of general guidance, the reaction solution may be distilled to about 1/10 the original
30 volume before solvent replacement is carried out. The reaction may be sampled and assayed to determine the extent of the reaction and the wt % product in accordance with standard process techniques. If desired, additional reaction solvent may be

added or removed to optimize reaction concentration. Preferably, the final concentration is adjusted to about 50 wt % at which point a slurry typically results.

[00211] It may be preferable to add solvents directly to the reaction vessel without distilling the reaction mixture. Preferred solvents for this purpose are those which may ultimately participate in the crystalline lattice, as discussed above in connection with solvent exchange. Although the final concentration may vary depending on desired purity, recovery and the like, the final concentration of the free base in solution is preferably about 4% to about 7%. The reaction mixture may be stirred following solvent addition and simultaneously warmed. By way of illustration, the reaction mixture may be stirred for about 1 hour while warming to about 70°C. The reaction is preferably filtered hot and washed with either the reaction solvent, the solvent added or a combination thereof. Seed crystals may be added to any crystallization solution to initiate crystallization.

[00212] The various forms described herein may be distinguishable from one another through the use of various analytical techniques known to one of ordinary skill in the art. Such techniques include, but are not limited to, X-ray powder diffraction (PXRD) and/or thermogravimetric analysis (TGA). Specifically, the forms may be characterized and distinguished using single crystal x-ray diffraction, which is based on unit cell measurements of a single crystal of a given form at a fixed analytical temperature. A detailed description of unit cells is provided in Stout & Jensen, X-Ray Structure Determination: A Practical Guide, Macmillan Co., New York (1968), Chapter 3, which is herein incorporated by reference. Alternatively, the unique arrangement of atoms in spatial relation within the crystalline lattice may be characterized according to the observed fractional atomic coordinates. Another means of characterizing the crystalline structure is by powder x-ray diffraction analysis in which the diffraction profile is compared to a simulated profile representing pure powder material, both run at the same analytical temperature, and measurements for the subject form characterized as a series of 2θ values (usually four or more).

[00213] Other means of characterizing the form may be used, such as solid state nuclear magnetic resonance (SSNMR) spectroscopy, differential scanning calorimetry (DSC), thermography and gross examination of the crystalline or amorphous


morphology. These parameters may also be used in combination to characterize the subject form.

[00214] One of ordinary skill in the art will appreciate that an X-ray diffraction pattern may be obtained with a measurement error that is dependent upon the measurement conditions employed. In particular, it is generally known that intensities in a X-ray diffraction pattern may fluctuate depending upon measurement conditions employed and the shape or morphology of the crystal. It should be further understood that relative intensities may also vary depending upon experimental conditions and, accordingly, the exact order of intensity should not be taken into account. Additionally, a measurement error of diffraction angle for a conventional X-ray diffraction pattern is typically about 0.2° or less, preferably about 0.1° (as discussed hereinafter), and such degree of measurement error should be taken into account as pertaining to the aforementioned diffraction angles. Consequently, it is to be understood that the crystal forms of the instant invention are not limited to the crystal forms that provide X-ray diffraction patterns completely identical to the X-ray diffraction patterns depicted in the accompanying Figures disclosed herein. Any crystal forms that provide X-ray diffraction patterns substantially identical to those disclosed in the accompanying Figures fall within the scope of the present invention. The ability to ascertain substantial identities of X-ray diffraction patterns is within the purview of one of ordinary skill in the art.

SYNTHESIS

Scheme 1


Hydrolysis of ketoester V to ketoacid VI



[00215] Ketoester V is hydrolyzed to its corresponding ketoacid VI by suspending V in an organic solvent partially miscible with water, such as acyclic or cyclic ethers including THF, 2-methyl THF, 1,2-dimethoxyethane, 1,4-dioxane, THF being preferred, and adding aqueous base such as aqueous solutions of alkali metal hydroxides MOH, where M is Li, Na or K, 1N NaOH being the preferred base, at -5°C to +5°C. The biphasic mixture is then agitated at $\leq 5^\circ\text{C}$ for at least one hour. Low temperature for the base addition and reaction is important to minimize the epimerization at the carbon adjacent to the ester group. Water-nonmiscible solvent, preferably methyl *tert*-butyl ether is then added and the layers are separated. The product is then transferred from the aqueous back to the organic solvent, preferably dichloromethane, by adjusting the pH with acid, preferably 3N HCl, and VI is used in solution for the next step.

Scheme 2


Ketalization of ketoacid VI to ketal acid VII



[00216] The solution of VI, preferably in dichloromethane, is exchanged via distillation into a non hygroscopic, higher boiling solvent, such as toluene, trifluorotoluene, xylenes, higher boiling esters such as *n*-butyl or isobutyl acetate, preferably toluene. A glycol of formula HO-Z-OH, (wherein Z is as defined *supra*) is then added, preferably ethylene glycol (1.2 eq), followed by a catalytic amount (0.5-2 M %) of an acid, preferably *p*-toluenesulfonic acid, and the mixture is distilled at atmospheric pressure until formation of compound VII is complete. The product VII crystallizes upon addition of ethyl acetate after cooling to approximately 70 °C. After further cooling to room temperature, VII is isolated by filtration and subsequent drying in about 70 % yield (for HO(CH₂)₂OH, R₁=H, R₂=CBz).

Scheme 3

Transformation of ketal acid VII to nonisolated ketal isocyanate VIII and then to ketal amide IX via acid activation/azidation, Curtius rearrangement and acylation




- [00217] The ketoacid **VII** is first activated by transforming it to its mixed anhydride using tertiary amines, preferably triethylamine, and haloformates, preferably isobutyl chloroformate in dry solvents, such as toluene, trifluorotoluene, 1,2-dichloroethane, 1-chlorobutane, xylenes, preferably dry toluene, by addition of a haloformate to a precooled solution of **VII** and trialkylamine. The preferred temperature for mixed anhydride formation is -10°C to 0°C . After approximately 30 min, an aqueous solution of alkali metal azide, preferably ~30 wt % sodium azide and a phase transfer catalyst, such as tetralkylammonium salts, preferably tetrabutylammonium bromide (5 mol %) is added and the biphasic mixture is vigorously agitated for about 1h at -10°C to 0°C . The organic phase is then separated and the resulting acyl azide solution is dried, 4 Å molecular sieves being preferred drying agent. The Curtius rearrangement and the concomitant trapping of the isocyanate **VIII** *in situ* with a carboxylic acid to form a ketal amide **IX** is accomplished by first adding a carboxylic acid, preferably acetic acid, and its corresponding anhydride, preferably acetic anhydride, to a dry solution of the acyl azide, and then heating the mixture $80-90^{\circ}\text{C}$ for 1-4 hours. The use of an anhydride in conjunction with carboxylic acid is critical to minimize impurity formation. After partial removal of the solvent and carboxylic acid by distillation, the product

crystallizes upon cooling to room temperature. IX is isolated by filtration and drying in 65-78 % yield (for $Z = -(CH_2)_2-$, $R_1=H$, $R_2=CBz$, $R_{10}=Me$).

Scheme 4


Hydrolysis of ketal amide IX to ketoamide X



[00218] The ketal hydrolysis of compound IX to the ketoamide X is accomplished by heating a solution of IX in organic, water miscible solvent, preferably acetone and an aqueous solution of strong acid, preferably 1N HCl, for 2-4 hrs. The preferred temperature for hydrolysis is 45-55°C. After removal of acetone, the product is extracted to dichloromethane, which is exchanged to ethyl acetate by distillation. The product X crystallizes from ethyl acetate upon cooling to room temperature, and is isolated by filtration and drying in 85-90 % yield (for $HO(CH_2)_2OH$, $R_1=H$, $R_2=CBz$, $R_{10}=Me$).

Scheme 5

Reductive amination of ketoamide X to the aminoamide XI




[00219] To a dry solution of X, preferably in dichloromethane, a primary or secondary amine is added, preferably *tert*-butylamine (5 eq), followed by a Lewis

acid, preferably $\text{TiCl}_2(\text{OPr-i})_2$ (1.2 eq), at -20°C to 0°C . The resulting imine mixture is warmed to 10 - 20°C and borane (1.1-1.2 eq) is added as a complex with dimethyl sulfide or THF, preferably dimethyl sulfide. The reaction mixture is agitated for 4-6 h and ethyl acetate saturated with water is then added. The titanium salts are removed by filtration and the product **XI** is extracted from the organic filtrate to water as its salt with an aqueous acid, preferably 1N HCl. Dichloromethane is then added and aqueous base, preferably concentrated ammonium hydroxide is added to the agitated biphasic mixture until pH is adjusted to 8.0-8.5. The product-rich dichloromethane phase is then separated and washed twice with aq ammonium chloride solution to remove the undesired trans isomer of **XI**, and finally with water. Dichloromethane is exchanged into ethyl acetate by distillation and **XI** crystallizes from ethyl acetate upon cooling and heptane addition. **XI** is isolated by filtration and drying in 65-70 % yield (for $\text{R}_1=\text{H}$, $\text{R}_2=\text{CBz}$, $\text{R}_8=\text{H}$, $\text{R}_9=\text{tert-Bu}$, $\text{R}_{10}=\text{Me}$).

Scheme 6


Deprotection of pyrrolidonyl amine XI



[00220] Removal of the amine protecting group R₂, where R₂ is CO₂CH₂Ph or CH₂Ph, is accomplished by hydrogenating a solution of **XI** in an alcohol, preferably methanol, in the presence of Pd catalyst, preferably 5wt % Pd/C, for several hours. The catalyst is then removed by filtration, and methanol is exchanged into ethyl acetate by distillation. The product **XII**, which crystallizes from ethyl acetate upon cooling and heptane addition, is isolated by filtration and drying in 90-95 % yield (for R₁=H, R₂=CBz, R₈=H, R₉=*tert*-Bu, R₁₀=Me).

Scheme 7

Coupling of amine XII with heterocycle bearing a leaving group HET-LG



- [00221] Synthesis of compound I is accomplished by coupling the
- 5 pyrrolidonylamine XII and a heterocycle bearing a leaving group in the presence of a tertiary amine, preferably triethylamine, in a compatible solvent, such as dichloromethane, isopropanol or acetonitrile, dichloromethane being preferred. All components are therefore combined and the solution is reacted for 24-48 hrs at room temperature. A solution of previously prepared crude heterocyclic component may
- 10 also be employed. After reaction completion, dichloromethane is washed with dilute acid, preferably aqueous 5 wt % acetic acid, the aqueous phase is separated and dichloromethane is then exchanged into ethyl acetate by distillation. The product I, which crystallizes from ethyl acetate upon cooling and heptane addition, is isolated by filtration and drying in 75-80 % yield (for $R_1=H$, $R_8=H$, $R_9=tert\text{-}Bu$, $R_{10}=Me$,
- 15 $HET=6\text{-(trifluoromethyl)quinazolin-4-yl}$).

- [00222] For the process of this invention, starting materials are commercially available or can be readily prepared by one of ordinary skill in the art. Solvents, temperatures, pressures, starting materials having the desired groups, and other
- 20 reaction conditions, may be readily selected as appropriate by one of ordinary skill in the art. The process can be scaled up in order to prepare larger quantities of the compound of formula I, such as in a commercial production facility.

EXAMPLES

[00223] The following Examples illustrate embodiments of the inventive compounds and starting materials, and are not intended to limit the scope of the claims.

[00224] As appropriate, reactions were conducted under an atmosphere of dry
5 nitrogen (or argon). For anhydrous reactions, Dri-Solv solvents from EM were employed. For other reactions, reagent grade or HPLC grade solvents were utilized. Unless otherwise stated, all commercially obtained reagents were used as received.


[00225] LC/MS measurements were obtained using a Shimadzu HPLC/Waters ZQ single quadrupole mass spectrometer hybrid system. Data for the peak of interest are
10 reported from positive-mode electrospray ionization. NMR (nuclear magnetic resonance) spectra were typically obtained on Bruker or JEOL 400 MHz and 500 MHz instruments in the indicated solvents. All chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance as the internal standard. ¹H-NMR spectral data are typically reported as follows: chemical shift, multiplicity (s = singlet,
15 br s = broad singlet, d = doublet, dd = doublet of doublets, t = triplet, q = quartet, sep = septet, m = multiplet, app = apparent), coupling constants (Hz), and integration.

[00226] One of skill in the art will recognize the standard abbreviations utilized herein, throughout the specification. For ease of reference, the abbreviations include, but are not necessarily limited to: sat. = saturated, HPLC = high-performance liquid
20 chromatography, AP = area percent, KF = Karl-Fischer, RT = room temperature, mmol = millimoles, HRMS = high-resolution mass spectroscopy, TBTU = O-benzotriazol-2-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate, MTBE = TBME = tert-butyl methyl ether, EDAC = N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride, EDC = N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide, TEA =
25 triethylamine, DPPA = diphenyl phosphoryl azide, IPA = isopropyl alcohol, TFA = trifluoroacetic acid, DCM = dichloromethane, THF = tetrahydrofuran, DMF = N,N-dimethylformamide, BOP = (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate, EtOAc = Ethyl acetate, DMSO = dimethylsulfoxide, °C =
degrees Celsius, eq = equivalent or equivalents, g = gram or grams, mg = milligram
30 or milligrams, mL (or ml) = milliliter or milliliters, h = hour or hours, M = molar, N = normal, min = minute or minutes, MHz = megahertz, tlc = thin layer chromatography, v/v = volume to volume ratio, and ca. = about.

[00227] "α", "β", "R" and "S" are stereochemical designations familiar to those skilled in the art.

EXAMPLE 1

N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide



[00228] **Example 1, Step 1:** (1R, 2S, 5R)-*tert*-Butyl 2-benzyloxycarbonylamino-7-oxo-6-aza-bicyclo[3.2.1]octane-6-carboxylate (89.6 g, 0.24 mol, see: P. H. Carter, et al. PCT application WO 2005/021500) was dissolved in ethyl acetate (1.5 L) and the resulting solution was washed with sat. NaHCO₃ (2 x 0.45 L) and sat. NaCl (1 x 0.45 L). The solution was dried (Na₂SO₄) and then filtered directly into a 3-necked 3 L round-bottom flask. The solution was purged with direct nitrogen injection before being charged with 10% Pd/C (13.65 g) under nitrogen atmosphere. The flask was evacuated and back-filled with hydrogen; this was repeated twice more. Hydrogen was bubbled through the solution for 30 min and then the reaction was stirred under 1 atm H₂ for 18 h. The flask was evacuated, back-filled with nitrogen, and charged with fresh catalyst (6 g of 10% Pd/C). Hydrogen was bubbled through the solution for 30 min and then the reaction was stirred under 1 atm H₂ for 18 h. The flask was evacuated and back-filled with nitrogen. The mixture was filtered through Celite; the filter pad was then washed with ethyl acetate. The filtrate (~1.6 L EtOAc volume) was diluted with acetonitrile (0.3 L) and charged sequentially with *L*-N-Cbz-methionine (68 g, 0.24 mol), TBTU (77 g, 0.24 mol), and N,N-diisopropylethylamine (42 mL, 0.24 mol). The reaction was stirred at room temperature for 4 h, during which time it changed from a suspension to a clear solution. The reaction was quenched with the addition of sat. NH₄Cl (0.75 L) and water (0.15 L); the mixture was diluted further with EtOAc (0.75 L). The phases were mixed and separated and the organic phase was washed with sat. Na₂CO₃ (2 x 0.9 L) and sat. NaCl (1 x 0.75 L). The solution was dried (Na₂SO₄), filtered, and concentrated *in vacuo* to give

(1*R*,2*S*,5*R*)- *tert*-butyl 2-((*S*)-2-(benzyloxycarbonylamino)-4-(methylthio)butanamido)-7-oxo-6-aza-bicyclo[3.2.1]octane-6-carboxylate as an oil, which was taken into the next step without further purification. LC/MS for primary peak: [M-Boc+H]⁺ = 406.3; [M+Na]⁺ = 528.3. ¹H-NMR (400 MHz, d₄-MeOH): δ 7.36 (m, 5H), 5.11 (s, 2H), 4.32 (m, 1H), 4.2 (m, 1H), 4.0 (m, 1H), 2.5 – 2.7 (m, 3H), 2.25 (m, 1H), 2.11 (s, 3H), 2.05 (m, 4H), 1.9 (m, 1H), 1.7 (m, 2H), 1.54 (s, 9H). Also present are EtOAc [1.26 (t), 2.03 (s), 4.12 (q)] and *N,N,N,N*-tetramethylurea [2.83 (s)].

[00229] **Example 1, Step 2:** A sample of (1*R*,2*S*,5*R*)- *tert*-butyl 2-((*S*)-2-(benzyloxycarbonylamino)-4-(methylthio)butanamido)-7-oxo-6-aza-bicyclo[3.2.1]octane-6-carboxylate (0.24 mol assumed; see previous procedure) was dissolved in iodomethane (1,250 g) and stirred for 48 h at room temperature. The reaction was concentrated *in vacuo*. The residue was dissolved in dichloromethane and concentrated *in vacuo*. This was repeated twice more. The resultant sludge was dissolved in dichloromethane (0.4 L) and poured into a rapidly stirring solution of MTBE (4.0 L). The resultant yellow solids were collected via suction filtration and dried under high vacuum to afford the sulfonium salt (179 g). This material was taken into the next step without further purification. LC/MS for primary peak: [M-Me₂S+H]⁺ = 458.4; [M]⁺ = 520.4. ¹H-NMR (400 MHz, d₄-MeOH): δ 7.35 (m, 5H), 5.09 (s, 2H), 4.33 (m, 1H), 4.28 (m, 1H), 3.98 (m, 1H), 3.3 – 3.45 (m, 2H), 2.97 (s, 3H), 2.94 (s, 3H), 2.78 (m, 1H), 2.0 – 2.3 (m, 4H), 1.7 (m, 2H), 1.52 (s, 9H). Also present are MTBE [1.18 (s), 3.2 (s)] and traces of *N,N,N,N*-tetramethylurea [2.81 (s)].

[00230] **Example 1, Step 3:** All of the sulfonium salt from the previous step (0.24 mol assumed) was dissolved in DMSO (2.0 L). The resultant solution was stirred under nitrogen at room temperature and charged with cesium carbonate (216 g) portionwise. The suspension was stirred at room temperature for 3 h and then filtered to remove the solids. The solution was divided into ~0.22 L portions and worked up as follows: the reaction mixture (~0.22 L) was diluted with ethyl acetate (1.5 L) and washed successively with water (3 x 0.5 L) and brine (1 x 0.3 L). The organic phase was dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The desired (1*R*,2*S*,5*R*)- *tert*-butyl 2-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-7-oxo-6-azabicyclo[3.2.1]octane-6-carboxylate (90.8 g, 83%) was obtained as a

microcrystalline foam, free from tetramethyl urea impurity. LC/MS for primary peak: $[M-Boc+H]^+ = 358.4$; $[M+Na]^+ = 480.4$. 1H -NMR (400 MHz, d_4 -MeOH): δ 7.35 (m, 5H), 5.12 (s, 2H), 4.35 (m, 2H), 4.2 (m, 1H), 3.6 (m, 1H), 3.3 (m, 1H), 2.64 (m, 1H), 2.28 – 2.42 (m, 2H), 2.15 (m, 1H), 1.7 – 2.0 (m, 5H), 1.55 (s, 9H). If desired, this material can be isolated as a solid by dissolving in MTBE (1 volume), adding to heptane (3.3 volumes), and collecting the resultant precipitate.

[00231] Example 1, Step 4: A stirring solution of (1*R*,2*S*,5*R*)-*tert*-butyl 2-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-7-oxo-6-azabicyclo[3.2.1]octane-6-carboxylate (108 g, 0.236 mol) in THF (1 L) was charged with lithium hydroxide monohydrate (21.74 g, 0.519 mol). Water (0.3 L) was added slowly, such that the temperature did not exceed 20 °C. The reaction was stirred at room temperature overnight and the volatiles were removed *in vacuo*. The pH was adjusted to ~4 through the addition of 1N HCl (450 mL) and NaH_2PO_4 . The resultant white precipitates were collected by filtration and washed with water (2 x 1 L). The solid was dissolved in dichloromethane (1.5 L) and water (~ 1 L). The organic layer was dried (Na_2SO_4), filtered, and concentrated *in vacuo*. The residue was dissolved in EtOAc (0.7 L) and the resultant solution was heated at reflux for 1 h. Solids separated after cooling to RT, and were collected *via* filtration. These solids were purified by recrystallization in isopropanol to afford the desired (1*R*,2*S*,5*R*)-2-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-5-(*tert*-butoxycarbonylamino)cyclohexanecarboxylic acid as a white solid (104.5 g, 93% yield). LC/MS for primary peak: $[M-tBu+H]^+ = 420.2$; $[M-Boc+H]^+ = 376.2$; $[M+H]^+ = 476.2$. 1H -NMR (400 MHz, d_4 -MeOH): δ 7.35 (m, 5H), 5.11 (s, 2H); 4.35 (m, 2H), 3.71 (m, 1H), 3.45 – 3.6 (m, 2H), 2.99 (m, 1H), 2.41 (m, 1H), 2.15 (m, 1H), 2.0 (m, 2H), 1.6 – 1.9 (m, 4H), 1.46 (s, 9H).

[00232] Example 1, Step 5: A 3 L round bottom flask was charged with (1*R*,2*S*,5*R*)-2-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-5-(*tert*-butoxycarbonylamino)cyclohexanecarboxylic acid (75.5 g, 0.158 mol), EDC•HCl (33.5 g, 0.175 mol), 1-hydroxybenzotriazole (23.6 g, 0.175 mol), and dichloromethane (1 L). The reaction was stirred at room temperature for 2 h, during which time it changed from a white suspension to a clear solution. Ammonia (gas) was bubbled into the solution until the pH was strongly basic (paper) and the reaction

was stirred for 10 min; this ammonia addition was repeated and the reaction was stirred for an additional 10 min. Water was added. The organic phase was washed with sat. NaHCO₃, NaH₂PO₄, and brine before being concentrated *in vacuo*. The residue was slurried with acetonitrile (0.5 L) and then concentrated in to give
5 (1*R*,2*S*,5*R*)-2-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-5-(*tert*-butoxycarbonylamino)cyclohexanecarboxamide as a white solid (75.9 g, ~100%), which was used in the next step without further purification. LC/MS for primary peak: [M-Boc+H]⁺ = 375.3; [M+H]⁺ = 475.4; [M-*t*Bu+H]⁺ = 419.3. ¹H-NMR (400 MHz, d₄-MeOH): δ 7.35 (m, 5H), 5.11 (s, 2H), 4.25 (m, 2H), 3.70 (m, 1H), 3.6 (m,
10 1H), 3.45 (m, 1H), 2.91 (m, 1H), 2.38 (m, 1H), 2.12 (m, 1H), 1.9 – 2.05 (m, 2H), 1.65 – 1.9 (m, 4H), 1.46 (s, 9H).

[00233] **Example 1, Step 6:** The reaction was run in three equal portions and combined for aqueous workup. A 5 L, 3-necked round bottom flask was charged with (1*R*,2*S*,5*R*)-2-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-5-(*tert*-
15 butoxycarbonylamino)cyclohexanecarboxamide (25.3 g, 53 mmol), acetonitrile (1.9 L), and 2.6 L of water/ice. The mixture was stirred and cooled to 0 °C. Iodobenzene diacetate (25.77 g, 80 mmol) was added and the reaction was stirred for 2 h; another 0.5 eq of iodobenzene diacetate was added. The reaction was stirred for 9 h (reaction temp < 10 °C). The mixture was charged with 8 eq N,N-diisopropylethylamine and 2
20 eq acetic anhydride. Over the next thirty minutes, 4 eq N,N-diisopropylethylamine and 2 eq acetic anhydride were added every ten minutes, until the reaction had proceeded to completion (HPLC). The acetonitrile was removed *in vacuo*; some solid separated from the residue, and this was collected by filtration. The remaining residue was extracted with dichloromethane (3 L, then 1 L). The organic phase was
25 washed sequentially with water, sat. NaHCO₃, and brine. The collected solids were added to the organic phase, along with activated carbon (15 g). The mixture was stirred for 30 minutes at 40 °C before being filtered and concentrated *in vacuo*. The residue was dissolved in EtOAc (1 L), and the resultant solution was stirred at 75 °C for 1 h before being allowed to cool to room temperature. A solid separated and was
30 collected by filtration. This solid was purified further by recrystallization: it was first dissolved in 0.5 L CH₂Cl₂, then concentrated *in vacuo*, then re-crystallized from 1 L EtOAc; this was repeated three times. The solids obtained from the mother liquors of

the above were recrystallized three times using the same method. The combined solids were recrystallized twice more from acetonitrile (0.7 L) to provide 66 g (84%) of *tert*-butyl (1*R*,3*R*,4*S*)-3-acetamido-4-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)cyclohexylcarbamate (purity >99.5% by HPLC). LC/MS for primary peak: $[M+H]^+ = 489.4$; $[M-tBu+H]^+ = 433.3$. $^1\text{H-NMR}$ (400 MHz, d_4 -MeOH): δ 7.3 – 7.4 (m, 5H), 5.11 (s, 2H), 4.35 (m, 1H), 4.15 (m, 1H), 4.04 (m, 1H), 3.8 (m, 1H), 3.6 (m, 2H), 2.44 (m, 1H), 2.12 (m, 1H), 1.87 – 2.05 (m, 4H), 1.87 (s, 3H), 1.55 – 1.7 (m, 2H), 1.46 (s, 9H). The stereochemical fidelity of the Hofmann rearrangement was confirmed through X-ray crystal structure analysis of this compound, as shown in Figure 1.

[00234] Example 1, Step 7: To a solution of *tert*-butyl (1*R*,3*R*,4*S*)-3-acetamido-4-((*S*)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)cyclohexylcarbamate (100 g, 0.205 mol) in dichloromethane (400 ml) was added TFA (400 ml) at -20 °C. The reaction solution was stirred at room temperature for 2 h. The solvent and most of TFA were removed under reduced pressure, and the residue was diluted with dichloromethane (2 L) and aqueous K_2CO_3 solution (2 L). The pH was adjusted to 10 with 1N HCl. The aqueous layer was extracted with dichloromethane (3 x 1L). The combined organic layer was dried over Na_2SO_4 , and concentrated to give benzyl (S)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-aminocyclohexyl)-2-oxopyrrolidin-3-ylcarbamate as an oil (81 g, 100% yield). This amine was used directly in the next step without further purification.

[00235] Example 1, Step 8: A solution of benzyl (S)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-aminocyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (13.3 g, 34 mmol) and 3, 5-di-*tert*-butylcyclohexa-3,5-diene-1,2-dione (7.54 g, 34 mmol) in methanol (160 ml) was stirred at room temperature for 2 h. The solution was concentrated and diluted with acetone (132 ml) and water (33 ml), followed by addition of Dowex-50WX8-200 (33 g). The reaction was stirred at room temperature for 2 h. Dowex-50WX8-200 was removed by filtration and washed with dichloromethane (300 ml). The filtrate was concentrated under vacuum to remove most of acetone. The residue was diluted with dichloromethane (200 ml) and washed with aqueous NaHCO_3 solution (200 ml) and brine (200 ml). The combined aqueous layers were extracted with dichloromethane (2 x 100 ml). The combined organic extracts were dried over Na_2SO_4 and concentrated.

The product benzyl (S)-1-((1S,2R)-2-acetamido-4-oxocyclohexyl)-2-oxopyrrolidin-3-ylcarbamate was obtained as a solid (12 g, 90% yield) by crystallization in EtOAc (100 ml) and Hexane (200 ml). ¹H-NMR (500 MHz, DMSO-*d*₆) δ ppm 7.99 (d, *J*=9.35 Hz, 1 H), 7.44 (d, *J*=8.80 Hz, 1 H), 7.28-7.39 (m, 5 H), 5.03 (s, 2 H), 4.50 (s, 1 H), 4.31 (d, *J*=12.10 Hz, 1 H), 4.18 (q, *J*=8.98 Hz, 1 H), 3.27 (m, 2 H), 2.82 (dd, *J*=15.12, 5.22 Hz, 1 H), 2.52-2.65 (m, 1 H), 2.40 (dd, *J*=12.92, 4.67 Hz, 1 H), 2.15-2.31 (m, 2 H), 2.09 (d, *J*=15.40 Hz, 1 H), 1.90 (m, 1 H), 1.81 (s, 3 H), 1.68 (m, 1 H). *m/z*: 388.46 [M+H].

[00236] **Example 1, Step 9:** To a solution of TiCl₄ (1M in dichloromethane, 36 ml, 36 mmol) in dichloromethane (30 ml) at 0° C was added Ti(OiPr)₄ (10.8 ml, 36 mmol). The mixture was then stirred at room temperature for 10 min. To a solution of benzyl (S)-1-((1S,2R)-2-acetamido-4-oxocyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (23.25 g, 60 mmol) in dichloromethane (600 ml) was added *tert*-butylamine (30 ml, 300 mmol) at room temperature, followed by the addition of the TiCl₄/Ti(OiPr)₄ solution at -50 °C. The reaction was allowed to warm slowly to room temperature. The reaction was finished after 2 h (The reaction was monitored on HPLC by quenching an HPLC sample with NaBH₄ in methanol). The solution was cooled to 10 °C and BH₃•SMe₂ (1M in dichloromethane, 66 ml, 66 mmol) was added. The mixture was stirred at room temperature for 5 h then quenched with Na₂CO₃ aqueous solution (300 ml). The precipitate was filtered off. The two layers were separated and the aqueous layer was extracted with dichloromethane (600 ml). The combined dichloromethane layers were extracted with 1N HCl twice (150 ml and 15 ml). (The product and the undesired *trans* isomer were both in the acidic aqueous phase.) The combined acidic aqueous layers were neutralized with 12 M aqueous solution of NH₄OH (12 ml) to pH~8 and extracted with dichloromethane twice (600 ml, 450 ml). (The product was in organic phase, while the *trans* isomer was still in aqueous layer.) The combined organic layers were washed with NH₄Cl aqueous solution 3 times (3 x 200 ml) until there was no *trans* isomer left in organic layer. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by crystallization in EtOAc/Hexane(200 ml / 800 ml) to give the desired benzyl (S)-1-((1S,2R,4R)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (20.80 g, 78% yield) as a white solid with 99.5% purity. ¹H-NMR (500

MHz, *DMSO-d*₆) δ ppm 8.76 (s, 1H), 7.27 - 7.46 (m, 6 H), 5.03 (m, 2 H), 4.14 (m, 1 H), 4.07 (q, *J*=8.80 Hz, 1 H), 3.83(m, 1H), 3.36 (m, 2H), 2.91 (s, 1H), 2.18 (m, 1H), 2.04 (m, 1H), 1.78 (s, 3H), 1.41 - 1.74 (m, 7H), 1.04 (s, 9H). *m/z*: 445.54 [M+H].

[00237] **Example 1, Step 10:** To a solution of benzyl (*S*)-1-((1*S*,2*R*,4*R*)-2-

5 acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (43.3 g, 98 mmol) in methanol (400 ml), 10% wet Pd/C (4.34 g) was added. The mixture was evacuated and back-filled with hydrogen with a hydrogen balloon. The mixture was stirred at room temperature for 5 h. The mixture was filtered and washed with methanol (500 ml) and concentrated under vacuum to dryness. The crude product
10 obtained was distilled with IPA (2 x 100 ml) under reduced pressure to give product N-((1*R*,2*S*,5*R*)-2-((*S*)-3-amino-2-oxopyrrolidin-1-yl)-5-(*tert*-butylamino)cyclohexyl)acetamide as an oil (30 g, 98% yield). This amine was used in the next step without further purification.

[00238] **Example 1, Step 11:** To a solution of N-((1*R*,2*S*,5*R*)-2-((*S*)-3-amino-2-oxopyrrolidin-1-yl)-5-(*tert*-butylamino)cyclohexyl)acetamide (30 g, 97 mmol) in IPA
15 (400 ml) was added TEA (27 ml, 195 mmol) and 4-chloro-6-(trifluoromethyl)quinazoline (25 g, 107 mmol; see P. H. Carter, et al. PCT application WO 2005/021500). The mixture was stirred at room temperature overnight and then stirred at 70 °C for 1h. The resulted solution was concentrated under reduced pressure
20 to dryness. The residue was dissolved in dichloromethane (1 L) and extracted with acetic acid solution I (prepared by combining 700 mL of water and 22.6 mL of glacial acetic acid) twice (500 ml, 200 ml). The acidic aqueous layer (pH 4-5) was extracted with dichloromethane (2 x 300 ml). The dichloromethane layer was extracted with acetic acid solution II (300 ml; prepared by combining 300 mL water with 4 mL of
25 glacial acetic acid). The combined acetic acid layers were basified with 1 M NaOH to pH > 12 and extracted with dichloromethane (3 x 700 ml). The combined organic layers were dried and concentrated to give the crude product as a solid (45.6 g, 93% yield). The crude product was purified by recrystallization from EtOAc (400 ml)/Hexane (900 ml) to give 42.86 g (88%) N-((1*R*,2*S*,5*R*)-5-(*tert*-butylamino)-2-
30 ((*S*)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide with 99.7% purity. ¹H-NMR (500 MHz, *DMSO-d*₆) δ ppm 9.71 (1 H, br. s.), 9.02 (1 H, s), 8.71 (1 H, d, *J*=7.97 Hz), 8.59 (1 H, s), 8.04 (1 H, dd,

$J=8.66, 1.79$ Hz), 7.88 (1 H, d, $J=8.52$ Hz), 4.91 - 5.13 (1 H, m), 4.30 - 4.57 (1 H, m),
 3.86 (1 H, dt, $J=11.89, 3.71, 3.64$ Hz), 3.43 - 3.57 (1 H, m), 3.35 - 3.45 (1 H, m), 3.04
 (1 H, t, $J=3.85$ Hz), 2.23 - 2.40 (1 H, m), 2.05 - 2.22 (1 H, m), 1.90 - 1.98 (1 H, m),
 1.86 - 1.93 (3 H, m), 1.50 - 1.78 (5 H, m), 0.98 - 1.15 (9 H, m). ^{13}C -NMR (126 MHz,
 5 *DMSO-d*₆) δ ppm 171.23, 169.35, 159.54, 156.87, 151.17, 128.97, 128.20, 125.76 (1
 C, q, $J=30.52$ Hz), 121.55 (1 C, br. s.), 124.04 (1 C, q, $J=272.11$ Hz), 114.31, 53.26,
 52.39, 50.81, 47.56, 45.70, 42.77, 34.52, 32.17, 29.14 (3 C, s), 26.49, 23.29, 20.30.
 ^{19}F -NMR (471 MHz, *DMSO-d*₆) δ ppm -60.34 (s). m/z : 507.0 [M+H]. Anal. Calcd
 for C₂₅H₃₃N₆O₂F₃: C, 59.27; H, 6.56; N, 16.59; F, 11.25 Found: C, 59.44; H, 6.64; N,
 10 16.74; F, 10.99.

ALTERNATIVE PREPARATION OF EXAMPLE 1

[00239] Example 1, Alternative Preparation, Step 1: An oven-dried 3-neck
 15 round-bottom flask was equipped with a dried stir bar, a dried reflux condenser, and
 two septa. After cooling under N₂, the flask was charged sequentially with
 (1R,2S,5R)-2-((S)-3-(benzyloxycarbonylamino)-2-oxopyrrolidin-1-yl)-5-(*tert*-
 butoxycarbonylamino)cyclohexanecarboxylic acid (60 g, 126 mmol; see Example 1,
 Step 4), acetonitrile (800 mL), N-methylmorpholine (27.7 mL, 252 mmol), and
 20 diphenylphosphoryl azide (29.9 mL, 139 mmol). The reaction was stirred at RT for 1
 h 40 min, at which time 2-trimethylsilylethanol (90 mL, 631 mmol) was added. The
 reaction was set to heat, and reached reflux 30 min later. It was allowed to reflux for
 1 h, at which time it was allowed to cool to 50 °C gradually and then cooled to 15 °C
 with external cooling. The reaction was quenched with the addition of acetic acid
 25 (1.734 mL, 30.3 mmol). The reaction was concentrated *in vacuo* and then dissolved
 in EtOAc (1.2 L). It was washed sequentially with water (1 x 0.3 L), sat. NaHCO₃ (2
 x 0.3 L), 1N HCl (1 x 0.3 L), and brine (2 x 0.3 L). The organic phase was dried
 (Na₂SO₄), filtered, and concentrated *in vacuo*. A solid appeared very early on in the
 concentration process. After the volatiles were removed, 800 mL 10%
 30 EtOAc/Hexanes was added, and the mixture was stirred overnight. The solid was
 collected and dried to yield *tert*-butyl (1R,3R,4S)-4-((S)-3-benzyloxycarbonylamino-
 2-oxopyrrolidin-1-yl)-3-((2-

trimethylsilyl)ethoxycarbonylamino)cyclohexylcarbamate (60.5 g, 102 mmol, 81 % yield). HPLC showed that the material was 72% pure, with two 12% impurities.

This material was taken into the next step without purification. The filtrate was later concentrated to yield another 4.38 g of product. Total yield = 64.9 g (87%).

- 5 **[00240] Example 1, Alternative Preparation, Step 2:** A dry 500 mL round-bottom flask was equipped with a stir bar and charged sequentially with *tert*-butyl (1R,3R,4S)-4-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)-3-((2-trimethylsilyl)ethoxycarbonylamino)cyclohexylcarbamate (60.5 g), CH₂Cl₂ (180 mL), and a solution of para-toluenesulfonic acid monohydrate (19.48 g, 102 mmol) in
- 10 CH₂Cl₂ (120 mL) and methanol (30 mL). The mixture was placed on a rotary evaporator and the bulk of the CH₂Cl₂ was removed (bath temp ca. 20 °C). When the mixture began to foam, the vacuum was released, and the bath temperature increased to 46 °C (the temperature varied between 44 and 51 °C; it was controlled with the addition of external ice). The mixture was rotated at this temperature for exactly one
- 15 hour (gas evolution was visible throughout) and then diluted with EtOAc (1 L). The organic phase was washed with 0.5 N NH₄OH (2 x 250 mL). The aqueous washes were combined and set aside. The organic phase was washed with sat. NH₄Cl (1 x 250 mL) and sat. NaCl (1 x 250 mL); these aqueous washes were discarded. The initial combined NH₄OH washes were back-extracted with EtOAc (1 x 250 mL), and
- 20 that organic extract was washed with sat. NH₄Cl (1 x 60 mL) and sat. NaCl (1 x 60 mL). All of the organic extracts were combined, dried (Na₂SO₄), filtered, and concentrated. The residue was purified by elution through a SiO₂ plug (13 cm wide x 7.5 cm tall). The first eluant was pure EtOAc (ca. 4 L). The second eluant was 1:9 (10% NH₄OH in MeOH)/CH₂Cl₂ (ca. 5 L). The fractions containing the desired
- 25 product were pooled together and evaporated to afford the desired 2-(trimethylsilyl)ethyl (1R,2S,5R)-5-amino-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)cyclohexylcarbamate (31.6 g, 64.4 mmol, 63 % yield).

- [00241] Example 1, Alternative Preparation, Step 3:** A stirring solution of 2-(trimethylsilyl)ethyl (1R,2S,5R)-5-amino-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)cyclohexylcarbamate (400 mg, 0.82 mmol) in acetonitrile (3 mL)
- 30 was charged sequentially with diisopropylethylamine (315.8 mg, 3 eq) and bromoacetonitrile (109.5 mg, 1.1 eq). The mixture was stirred at 40 °C for 30 h. The

solvent was removed under reduce pressure. The residue was purified by silica gel column chromatography, using 1.5% of methanol in dichloromethane as the eluant. The desired 2-(trimethylsilyl)ethyl(1R,2S,5R)-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)-5-(cyanomethylamino)cyclohexylcarbamate was obtained as a white solid (400 mg, 93%). LC/MS found $[M + H]^+ = 530$.

[00242] Example 1, Alternative Preparation, Step 4: A stirring solution of 2-(trimethylsilyl)ethyl(1R,2S,5R)-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)-5-(cyanomethylamino)cyclohexylcarbamate (400 mg, 0.76 mmol) in dichloromethane (5 mL), was cooled to 0 °C and charged with m-CPBA (372.6 mg, 2.2 eq) in portions. The mixture was stirred at room temperature for 1.5 h. Saturated $\text{Na}_2\text{S}_2\text{O}_3$ solution (3 mL) and saturated NaHCO_3 solution (3 mL) were added and the mixture was stirred at room temperature for 0.5h. The mixture was diluted with dichloromethane (80 mL), washed with saturated NaHCO_3 (20 mL) and brine (20 mL). The solution was dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. The residue obtained was dissolved in methanol (5 mL) and the solution was charged with $\text{NH}_2\text{OH}\cdot\text{HCl}$ (262.7 mg, 5 eq). The mixture was stirred at 60 °C for 2.5 h. After cooling to room temperature, the mixture was diluted with dichloromethane (80 mL) and filtered through a pad of celite. The filtrate was washed with saturated NaHCO_3 (2 x 20 mL). The aqueous washes were extracted with dichloromethane (30 mL). The dichloromethane layers were combined and washed with brine (30 mL). The solution was dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo* to give 2-(trimethylsilyl)ethyl (1R,2S,5R)-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)-5-(hydroxyamino)cyclohexylcarbamate (350 mg, 91%). LC/MS found $[M + H]^+ = 507$.

[00243] Example 1, Alternative Preparation, Step 5: A solution of 2-(trimethylsilyl)ethyl (1R,2S,5R)-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)-5-(hydroxyamino)cyclohexylcarbamate (350 mg, 0.69 mmol) in acetone (5 mL) was stirred at room temperature for 16 h. The mixture was concentrated *in vacuo*. The residue was dissolved in anhydrous THF (7 mL) and cooled to 0 °C. A solution of MeMgBr (1.1 mL, 3M in diethyl ether, 5 eq) was added dropwise. The mixture was stirred at room temperature for 1.5h. The reaction was quenched with water (5 mL) at 0 °C. The mixture was diluted with ethyl acetate (100 mL) and filtered through a

pad of celite. The filtrate was washed with brine (30 mL). The solution was dried over anhydrous MgSO_4 , filtered, and concentrated *in vacuo*. The residue was dissolved in 2 mL of acetonitrile and 1 mL of CS_2 was added. The mixture was stirred at room temperature for 1 hour. The solvent was removed and the crude product was

5 purified by silica gel column chromatography, using 1.5% of methanol in dichloromethane as the eluant, to provide the desired 2-(trimethylsilyl)ethyl (1R,2S,5R)-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)-5-(*tert*-butylamino)cyclohexylcarbamate (160 mg, 42%). LC/MS found $[\text{M} + \text{H}]^+ = 547$.

[00244] Example 1, Alternative Preparation, Step 6: A stirring solution of 2-

10 (trimethylsilyl)ethyl (1R,2S,5R)-2-((S)-3-benzyloxycarbonylamino-2-oxopyrrolidin-1-yl)-5-(*tert*-butylamino)cyclohexylcarbamate (100 mg, 0.183 mmol) in dichloromethane (3 mL) was charged with trifluoroacetic acid (2 mL). The reaction was stirred for 2 h at room temperature and concentrated *in vacuo*. The residue was dissolved in dichloromethane (50 mL) and washed with saturated NaHCO_3 (20 mL).

15 The aqueous layer was extracted with dichloromethane (3 x 30 mL). The dichloromethane layers were combined and dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo* to give benzyl (S)-1-((1S,2R,4R)-2-amino-4-(*tert*-butylamino)cyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (66 mg, 90%). LC/MS found $[\text{M} + \text{H}]^+ = 403$.

20 **[00245] Example 1, Alternative Preparation, Step 7:** A solution of benzyl (S)-1-((1S,2R,4R)-2-amino-4-(*tert*-butylamino)cyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (22 mg, 0.055 mmol) in dichloromethane (2 mL) was charged sequentially with triethylamine (11.1 mg, 2 eq) and acetic anhydride (6.1 mg, 1.1 eq). The reaction was stirred for 1.5 h at room temperature, diluted with dichloromethane

25 (50 mL) and washed with saturated NaHCO_3 (20 mL). The organic phase was dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo* to give benzyl (S)-1-((1S,2R,4R)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (22 mg, 90%). LC/MS found $[\text{M} + \text{H}]^+ = 445$.


[00246] Example 1, Alternative Preparation, Step 8: To a solution of benzyl

30 (S)-1-((1S,2R,4R)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxopyrrolidin-3-ylcarbamate (22 mg, 0.05 mmol) in methanol (2 mL) was added $\text{Pd}(\text{OH})_2$ (20 mg of 50% wet catalyst). The flask was evacuated and back-filled with hydrogen from a

hydrogen balloon. The mixture was stirred at room temperature for 1 h and the catalyst was removed by filtration. The filtrate was concentrated *in vacuo* to provide N-((1R,2S,5R)-2-((S)-3-amino-2-oxopyrrolidin-1-yl)-5-(*tert*-butylamino)cyclohexyl)acetamide (13 mg, 85%). LC/MS found $[M + H]^+ = 311$.


- 5 [00247] **Example 1, Alternative Preparation, Step 9:** To a solution of N-((1R,2S,5R)-2-((S)-3-amino-2-oxopyrrolidin-1-yl)-5-(*tert*-butylamino)cyclohexyl)acetamide (70 mg, 0.225 mmol) in isopropanol (3 mL) was added 4-chloro-6-(trifluoromethyl)quinazoline (63 mg, 1.2 eq) and triethylamine (56.9 mg, 2.5 eq). The mixture was stirred at room temperature for 1.5 h. The solvent
10 was removed under reduced pressure. The residue was purified with preparative HPLC to provide the title compound as its bis-TFA salt (110 mg, 67%). LC/MS found $[M + H]^+ = 507$.

2ND ALTERNATIVE PREPARATION OF EXAMPLE 1




- 15
20
25
- [00248] **Example 1, 2nd Alternative Preparation, Step 1a:** To a hydrogenator were charged ethyl (7R,8S)-8-((S)-1-phenyl-ethylamino)-1,4-dioxaspiro[4.5]decane-7-carboxylate 4-toluenesulfonate salt **1A** (1417 g, 2.8 moles, c.f.: WO2004098516, prepared analogous to US Pat.6,835,841), ethanol (200 proof, 11.4 L), and 10% Pd/C catalyst (50% wet, 284 g). The mixture was inerted with nitrogen, then pressurized with hydrogen gas (45 psig) and agitated vigorously at approx. 40 °C until starting material was consumed (HPLC). The suspension was cooled, purged with nitrogen gas and the catalyst was removed by filtration while inerted. The spent catalyst was washed with ethanol (4.3 L). The filtrate and washings were combined and concentrated under vacuum to a volume of 2–3 L while maintaining the batch between 40°–60 °C. Isopropyl acetate (5 L) was charged and the mixture was concentrated to a volume of ~2 L until most ethanol was removed (<0.5%) and residual moisture content was <1,000 ppm. Batch volume was adjusted to ~7.5 L by

the addition of isopropyl acetate. The mixture was heated to 80 °C until clear, then cooled 65°–70 °C. Seed crystals of **1** (5 g) were added and the batch was cooled to 50 °C over 2 hours, then further cooled to 20 °C over 4 hours and held for ~10 hours. The resulting slurry was filtered and the cake was washed with isopropyl acetate (2 L). The product was dried under vacuum at ~35 °C until volatiles were reduced below ~1% (LOD). Ethyl (7*R*,8*S*)-8-amino-1,4-dioxa-spiro[4.5]decane-7-carboxylate 4-toluenesulfonate salt **1** was obtained as a white, crystalline solid (936 g, 83% yield; HPLC purity: 99.8%). ¹H-NMR: (300MHz, CDCl₃) 8.14-7.89 (brs, 3H), 7.75 (d, J 9.0Hz, 2H), 7.15 (d, J 8.0Hz, 2H), 4.22-4.04 (m, 2H), 4.01-3.77 (m, 4H), 3.55-3.43 (m, 1H), 3.20-3.13 (m, 1H), 2.40-2.27 (m, 4H), 2.21-1.94 (m, 2H), 1.81-1.51 (m, 3H), 1.23 (t, J 7.0Hz, 3H); HPLC: Waters Xterra MS C18 4.6 mm x 150 mm i.d., 3.5µm particle size, 0.05% NH₄OH (5% ACN, 95% H₂O, solvent A), to 0.05% NH₄OH (95% ACN, 5% H₂O, solvent B), 5% B to 20% B in 10 minutes, changed to 95% B in 25 minutes, and then changed to 5% B in 1 minute; 11.1 minutes (aminoester **1**).



[00249] **Example 1, 2nd Alternative Preparation, Step 1b:** Ethyl (7*R*,8*S*)-8-amino-1,4-dioxaspiro[4.5]decane-7-carboxylate 4-toluenesulfonate salt **1** (450.1g; the product of reductive deprotection of a known compound – see R. J. Cherney, WO 2004/098516 and G. V. Delucca & S. S. Ko, WO 2004/110993), was combined with 1-ethyl-3-(3-dimethyl-amino-propyl)carbo-diimide hydrochloride (236.3g), 1-hydroxy benzotriazole hydrate (171.9g), *N*-carbobenzyloxy-*L*-methionine (333.4g) and acetonitrile (3.1 L). To the stirred mixture was added triethylamine (249.5g) below 30 °C. Upon reaction completion (HPLC), the mixture was diluted with ethyl acetate (8.2 L) and washed with aqueous 25% potassium bicarbonate solution (2×4.5


L) followed by water (4.5 L). The organic phase was separated and concentrated under reduced pressure to obtain a solution of ethyl (7*R*,8*S*)-8-((*S*)-2-benzyloxycarbonylamino-4-methylsulfanyl-butylamino)-1,4-dioxaspiro[4.5]decane-7-carboxylate **2** (1.4 L). Methyl iodide (2.39 kg) was added, the vessel was shielded from light and the mixture was held under slow agitation for approx. 24 h. To the thick yellow precipitate was added methyl *tert*-butyl ether (2.7 L) and the mixture was held for approx. 1 h. The product was isolated by filtration and the cake was washed with methyl *tert*-butyl ether (2x1.4 L), then dried under vacuum, yielding [(*S*)-3-benzyloxy-carbonylamino-3-((7*R*,8*S*)-7-ethoxycarbonyl-1,4-dioxaspiro[4.5]dec-8-ylcarbonyl)-propyl]-dimethylsulfonium iodide **3** (671.4 g, ~94% yield) as an off-white solid (HPLC purity 99.9%).



[00250] Example 1, 2nd Alternative Preparation, Step 2: Sulfonium salt 3
 (619.4 g), and cesium carbonate (416.8 g) and anhydrous dimethyl sulfoxide (6.2 L) were combined in a reactor equipped with a scrubber to neutralize volatile sulfides. Vigorous agitation was maintained until complete conversion was obtained (HPLC). Ethyl acetate (12.4 L) was added, followed by 20 % brine (3 L). The organic phase was separated, washed twice with brine (2x3 L) and evaporated to obtain a solution of ethyl (7*R*,8*S*)-8-((*S*)-3-benzyloxycarbonylamino-2-oxo-pyrrolidin-1-yl)-1,4-dioxaspiro[4.5]decane-7-carboxylate **4** in ethyl acetate (~0.8 L). Acetone (2.55 L) was added, followed by aqueous 0.5 M hydrochloric acid solution (2.3 L). With good mixing, the solution was heated to 50 to 60 °C until conversion of **4** to ethyl (1*R*,2*S*)-2-((*S*)-3-benzyloxycarbonylamino-2-oxo-pyrrolidin-1-yl)-5-oxo-cyclohexanecarboxylate **5** was complete (HPLC). The mixture was concentrated under reduced pressure while below 40 °C, cooled to ~30 °C, and water (4.1 L) was

added. The resulting slurry was cooled to 5 to 10 °C and agitated for ~1 hour. The product was filtered and the cake was washed with water (2x2.5 L). Upon deliquoring, the cake was dried to a constant weight below 40 °C in a vacuum oven. Cyclohexanone **5** (272g, 70% yield) was obtained (HPLC purity 98.7%).

5



[00251] Example 1, 2nd Alternative Preparation, Step 3: Cyclohexanone 5

(100g) was suspended in THF (500 mL) and cooled to 0 °C. 1N NaOH (271g) was added at 0-5°C and the biphasic mixture was agitated at ≤ 5 °C for at least one hour.

10 MTBE (500 mL) was added and the layers were separated. The bottom aqueous layer was washed again with MTBE (500 mL) and the layers were separated.


Dichloromethane (500 mL) was charged to the product-rich aqueous layer, and the mixture was cooled to 0 °C. 3N HCl (156 g) was charged maintaining ≤ 5 °C. After stirring for at least 10 min, the mixture was warmed to 20-25 °C, and the layers were

5 separated. The aqueous layer was extracted with dichloromethane (100 mL). The organic layers were combined and solvent was exchanged into toluene via distillation.


The toluene solution volume was adjusted to approximately 1L, and p-toluenesulfonic acid monohydrate (0.24 g) was added. Ethylene glycol (16.22 g) was added, and the mixture was distilled at atmospheric pressure until formation of compound 7 was

10 complete and the pot volume was approximately 500-700 mL. The solution was cooled to approximately 70 °C, and ethyl acetate (500 mL) was added maintaining approximately 70 °C. The mixture was cooled and filtered to give 73 g (70 % yield) of compound 7, (7R,8S)-8-((3S)-3-(((benzyloxy)carbonyl)amino)-2-oxo-1-pyrrolidiny)-1,4-dioxaspiro[4.5]decane-7-carboxylic acid.


5



[00252] **Example 1, 2nd Alternative Preparation, Step 4:** To a slurry of compound **7** (147 g) in dry toluene (370 mL) was charged triethylamine (32.7 g) at 15-25 °C. After the slurry became a solution after 10-15 min stirring at 25°C, the flask was cooled to -10°C and isobutyl chloroformate (44.1g) was charged at -10°-0°C. The mixture was agitated at -10°- 0°C for about 30 min. A solution of sodium azide (42 g) and tetrabutylammonium bromide (5.2 g) in water (130 mL) was added at -10°- 0°C. The biphasic slurry was vigorously agitated for at least one hour and toluene (1750 mL) followed by water (300 mL) were added. The two phases were stirred for at least 10 min and the top organic layer was separated and dried with 4 Å molecular sieves. Acetic anhydride (76 mL) and acetic acid (28 mL) were added and the solution was heated to 80-90 °C for 1-4 hours, until <2 AP of intermediate **9** was detected by HPLC. The solvent was partially distilled off to approximately two thirds of the initial volume under atmospheric pressure, the solution was cooled to ambient temperature and the resultant slurry was stirred for 16 hours. Heptane (350 mL) was slowly added and the slurry was stirred for 1h. The solids were filtered, washed with toluene/heptane 4:1 (300 mL), and dried to give 109 g (78% yield) of compound **10**, benzyl ((3*S*)-1-((7*R*,8*S*)-7-acetamido-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidiny)carbamate.




[00253] **Example 1, 2nd Alternative Preparation, Step 5:** To a solution of compound **10** (109 g) in acetone (760 mL) was charged 1N HCl (760 mL). The mixture was heated to 50 °C for 2.5 hours. The acetone was distilled off under reduced pressure and the product was extracted with dichloromethane twice 1 x 1 L and 1 x 0.5 L. Dichloromethane layers were combined and dichloromethane was exchanged into ethyl acetate by distillation until the b.p. on the kettle reached 78°C and the final volume was approximately 10 mL/g compound **10** input. The ethyl acetate slurry was cooled to ambient temperature, agitated for 16 hrs and the solids were filtered and washed with ethyl acetate (400 mL). The solid was dried to give 84 g (87% yield) of compound **11**, benzyl ((3*S*)-1-((1*S*,2*R*)-2-acetamido-4-oxocyclohexyl)-2-oxo-3-pyrrolidiny)carbamate.




[00254] **Example 1, 2nd Alternative Preparation, Step 6:** The $\text{TiCl}_2(\text{OPr}^i)_2$ reagent was pre-formed by adding $\text{Ti}(\text{OPr}^i)_4$ (11.5 mL) to a solution of 1M TiCl_4 in dichloromethane (39 mL) at 5-10°C and subsequent stirring at ambient temperature for 15 min. Compound **11** (25 g) was dissolved in dichloromethane (500 mL) and *t*-butylamine (34 mL) was added at room temperature. After 10 min the solution was cooled to -25 to -20 °C and the preformed titanium reagent solution was added at a temperature below -20°C. The mixture was allowed to warm to ambient temperature and it was agitated for 1h. A sample was taken to confirm the complete imine formation by miniquench with sodium borohydride in methanol (absence of alcohols indicated the complete consumption of the starting ketone **11**). Borane dimethyl sulfide (7.0 mL) was then added at 0-5°C and the reaction mixture was warmed to ambient temperature and stirred for at least 5 hours. Dichloromethane was partially (about half) evaporated under reduced pressure and wet ethyl acetate (300 mL, preformed by agitating ethyl acetate with water) was added within 30-60 min. The

resulting slurry was agitated for at least 4 h, the solids were filtered and washed several times with dichloromethane until no more than 5 M % of the product was retained in the cake. The filtrate and washes were combined, 1N HCl (200 mL) was added and the biphasic mixture was agitated for at least 30 min (gas evolution ceased after ~20 min). The product rich aqueous layer (upper phase) was separated and dichloromethane (500 mL) was added. Concentrated ammonium hydroxide was added to the agitated biphasic mixture until pH was adjusted to 8-8.5 (~15 mL). The organic phase was separated and washed 2 x 100 mL with 14 wt % ammonium chloride, to remove the undesired trans isomer of compound 12, and finally with water (25 mL). Dichloromethane was exchanged into ethyl acetate by distillation under normal pressure until the b.p. on the kettle reached 78°C and the final volume was approximately 5 mL/g compound 11 input (~140 mL). The solution was cooled to ambient temperature. Heptane (250 mL) was added slowly at 40-50°C and compound 12 started to crystallize. The slurry was agitated at room temperature for 3 hrs and the solids were filtered, washed with heptane (100 mL) and dried to give 20.1 g (70 % yield) of compound 12, benzyl ((3*S*)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, as white, fluffy crystals.




[00255] Example 1, 2nd Alternative Preparation, Step 7: Compound 12 (20 g) was dissolved in methanol (400 mL) and 5 % Pd/C catalyst (1.8 g, 9 wt %) was added. The mixture was hydrogenated at 25 °C and 25 psig for 3h. The catalyst was removed by filtration, and the methanol was exchanged into ethyl acetate by continuous distillation. The product crystallized from ethyl acetate (160 mL) upon cooling. Heptane (160 mL) was added at 25 °C, the slurry was agitated for 1h and the product was filtered, washed with heptane and dried to give 12.8 g (94 % yield) of

compound 13, *N*-((1*R*,2*S*,5*R*)-2-((3*S*)-3-amino-2-oxo-1-pyrrolidinyl)-5-(*tert*-butylamino)cyclohexyl)acetamide.



- 5 [00256] **Example 1, 2nd Alternative Preparation, Step 8:** 6-(trifluoromethyl)-4-quinazolinol, 14 (5 g; see P. H. Carter, et al. PCT application WO 2005/021500) was suspended in dichloromethane (100 mL). *N,N*-diisopropylethylamine (4.2 mL, 1.05 eq) and DMF (0.4 mL, 0.2 eq) were added. Oxalyl chloride (3.0 mL, 1.5 eq) was then added to the agitated slurry at 20-25 °C under cooling (exothermic addition). The
- 10 orange slurry was agitated at 30-35 °C for 2h. Steady gas evolution was observed for ~1.5 hrs at which point the slurry became an orange solution. After cooling to 20 °C, the reaction solution was added dropwise to 20 wt % aq K₂HPO₄ (50 mL) under vigorous agitation and gas evolution. The lower organic phase was separated and washed one more time with 20 wt % aq K₂HPO₄ (50 mL). The organic solution was
- 15 used as is for the next step within 16h.



- [00257] **Example 1, 2nd Alternative Preparation, Step 8:** To a solution of 15 in dichloromethane (22 mL, 5.5 mmol), solid 13 (1.55 g, 5 mmol) was added and the mixture was agitated at room temperature until the solids dissolved. Triethylamine (1.4 mL, 11 mmol) was added and the mixture was agitated for 24 hrs at ambient temperature. Water (10 mL) was added, the two phases were agitated for 10 min and the organic phase was separated. Dichloromethane was exchanged into ethyl acetate
- 10

by continuous distillation until the b.p. on the kettle reached 78°C and the final volume was approximately 10 mL/g compound **13** input (~15 mL). The slurry was cooled to ambient temperature, heptane (15 mL) was slowly added and the slurry was agitated for 16 h. The solids were filtered, washed with heptane/ethyl acetate 1:1 (5 mL) and dried to give 1.83 g (72 % yield) of beige crystals (form N-2 by XRD) of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, Example 1.

EXAMPLE 2

Crystal Forms of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide
[00258] Various crystal forms of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base were prepared and characterized as described below.

PROCEDURES FOR CHARACTERIZING THE FORMS

Single Crystal Data

[00259] Data were collected on a Bruker-Nonius (BRUKER AXS, Inc., 5465 East Cheryl Parkway Madison, WI 53711 USA) CAD4 serial diffractometer. Unit cell parameters were obtained through least-squares analysis of the experimental diffractometer settings of 25 high-angle reflections. Intensities were measured using Cu K α radiation ($\lambda = 1.5418 \text{ \AA}$) at a constant temperature with the θ -2 θ variable scan technique and were corrected only for Lorentz-polarization factors. Background counts were collected at the extremes of the scan for half of the time of the scan. Alternately, single crystal data were collected on a Bruker-Nonius Kappa CCD 2000 system using Cu K α radiation ($\lambda = 1.5418 \text{ \AA}$). Indexing and processing of the measured intensity data were carried out with the HKL2000 software package (Otwinowski, Z. & Minor, W. (1997) in *Macromolecular Crystallography*, eds. Carter, W.C. Jr & Sweet, R.M. (Academic, NY), Vol. 276, pp. 307-326) in the Collect program suite. (Collect Data collection and processing user interface: Collect: Data collection software, R. Hooff, Nonius B.V., 1998.) Alternately, single crystal data were collected on a Bruker-AXS APEX2 CCD system using Cu K α radiation (λ

= 1.5418 Å). Indexing and processing of the measured intensity data were carried out with the APEX2 software package/program suite (APEX2 Data collection and processing user interface: APEX2 User Manual, v1.27; **BRUKER AXS, Inc.**, 5465 East Cheryl Parkway Madison, WI 53711 USA).

5 [00260] When indicated, crystals were cooled in the cold stream of an Oxford cryo system (Oxford Cryosystems Cryostream cooler: J. Cosier and A.M. Glazer, *J. Appl. Cryst.*, 1986, *19*, 105) during data collection.

[00261] The structures were solved by direct methods and refined on the basis of observed reflections using either the SDP (SDP, Structure Determination Package, Enraf-Nonius, Bohemia NY 11716. Scattering factors, including f' and f'' , in the SDP software were taken from the "International Tables for Crystallography", Kynoch Press, Birmingham, England, 1974; Vol. IV, Tables 2.2A and 2.3.1) software package with minor local modifications or the crystallographic packages MAXUS (maXus solution and refinement software suite: S. Mackay, C.J. Gilmore, C. Edwards, M. Tremayne, N. Stewart, K. Shankland. maXus: a computer program for the solution and refinement of crystal structures from diffraction data or SHELXTL⁴. The derived atomic parameters (coordinates and temperature factors) were refined through full matrix least-squares. The function minimized in the refinements was $\sum_w (|F_o| - |F_c|)^2$. R is defined as $\sum ||F_o| - |F_c|| / \sum |F_o|$ while $R_w = [\sum_w (|F_o| - |F_c|)^2 / \sum_w |F_o|^2]^{1/2}$ where w is an appropriate weighting function based on errors in the observed intensities. Difference maps were examined at all stages of refinement. Hydrogens were introduced in idealized positions with isotropic temperature factors, but no hydrogen parameters were varied.

X-ray Powder Diffraction Data (PXRD)

25 [00262] PXRD data were obtained using a Bruker C2 GADDS . The radiation was Cu K α (40 KV, 50mA). The sample-detector distance was 15 cm. Powder samples were placed in sealed glass capillaries of 1mm or less in diameter; the capillary was rotated during data collection. Data were collected for $3 \leq 2\theta \leq 35^\circ$ with a sample exposure time of at least 2000 seconds. The resulting two-dimensional diffraction arcs were integrated to create a traditional 1-dimensional PXRD pattern with a step
30 size of 0.02 degrees 2θ in the range of 3 to 35 degrees 2θ .

Differential Scanning Calorimetry (DSC)

[00263] DSC experiments were performed in a TA Instruments™ model Q1000 or 2920. The sample (about 2-6 mg) was weighed in an aluminum pan and recorded accurately recorded to a hundredth of a milligram, and transferred to the DSC. The instrument was purged with nitrogen gas at 50mL/min. Data were collected between room temperature and 300°C at 10°C/min heating rate. The plot was made with the endothermic peaks pointing down.

10 Thermal Gravimetric Analysis (TGA)

[00264] TGA experiments were performed in a TA Instruments™ model Q500 or 2950. The sample (about 10-30 mg) was placed in a platinum pan previously tared. The weight of the sample was measured accurately and recorded to a thousandth of a milligram by the instrument. The furnace was purged with nitrogen gas at 100 mL/min. Data were collected between room temperature and 300°C at 10°C/min heating rate.

PREPARATION AND ANALYSIS OF THE FORMS

[00265] The unit cell data and other properties for these examples are presented in Table 1. The unit cell parameters were obtained from single crystal X-ray crystallographic analysis. A detailed account of unit cells can be found in Chapter 3 of Stout & Jensen, *X-Ray Structure Determination: a Practical Guide*, (MacMillan, 1968).

[00266] Fractional atomic coordinates for Examples 2a, b, c, d, e, f, and g, and the conditions at which they were measured are presented in Tables 2-9.

[00267] Additionally, characteristic powder x-ray diffraction peak positions (degrees $2\theta \pm 0.1$) @ RT for Examples 2a, b, c, d, e, and f are presented in Table 9, all of which are based on high quality patterns collected with a diffractometer ($\text{CuK}\alpha$) with a spinning capillary with 2θ calibrated with a NIST other suitable standard.

[00268] Finally, Figures 1, 2, 3, 4, 5 and 6 present XRPD patterns for Examples 2a, b, c, e, d and f, Figures 8, 10, 13, 15, 17 and 19 disclose the TGA of Examples 2a, b, c, d, e and f, respectively. Figures 7, 9, 12, 14, 16 and 18 disclose the DSC of

Examples 2a, b, c, d, e and f, respectively. Figure 11 discloses the Vapor Sorption Isotherm of Example 2b.

Form Preparation, DSC and TGA Characterization

- 5 [00269] **Example 2a, Form H0.5-4:** 150 mg of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base was dissolved in warm n-butyl acetate saturated with water. Heptane was added until a persistent cloud was observed. The slurry was allowed to cool to RT. Form H0.5-4 is characterized by 0.5 mole water per mole N-
- 10 ((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base. Form H0.5-4 was characterized by a DSC thermogram having a broad endotherm onset typically in the range between ca. RT and ca. 67 °C in agreement with the TGA curve; at higher temperatures other events may ensue. Form H0.5-4 was characterized by a TGA
- 15 thermal curve having a weight loss typically from 0.6% to about 1.4% up to ca. 100 °C. The theoretical weight loss for Form H0.5-4 is 1.7%, however, it is not unusual for unstable hydrates to become partially dehydrated upon drying.
- [00270] **Example 2b, Form N-2:** 1 g of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-
- 20 yl)cyclohexyl)acetamide, free base was dissolved in 10 mL of water-free EtOAc at 77°C. The solution was cooled to 70°C. 10 mg of seeds of N-2 were added. To the slurry, 18 mL of n-heptane was added over 1 hour with a syringe pump. The slurry was cooled from 70°C to 20°C over 1 hour, and agitated at 20°C overnight. The solid was isolated by filtration, washed with 3 mL of n-heptane, dried at 50°C in a vacuum
- 25 oven overnight. Form N-2 is N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base, a neat form (without additional molecules of water or solvent). Form N-2 was characterized by a DSC thermogram having an endothermic onset typically between ca. 230 °C and ca. 232 °C as a single melt with no other transformations. Form N-2
- 30 was characterized by a TGA curve having negligible weight loss at up to ca. 200 °C and in agreement with the single-crystal structure.

[00271] **Example 2c, Form H1.75-5:** A slurry of 50 mg of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base was vigorously agitated in 1 mL water for more than 16 hours. Form H1.75-5 is characterized by 1.75 moles of water per one mole of
5 N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base. Form H1.75-5 was characterized by a DSC thermogram having an endotherm onset typically between ca. RT and ca. 70 °C in agreement with the TGA curve; at higher temperatures other events may ensue. Form H1.75-5 was characterized by TGA curve having a weight
10 loss of ca. 4.3% to ca. 5.3% at temperatures up to ca 100 °C. Theoretical weight loss is ca. 5.9%, however, it is not unusual for unstable hydrates to become partially dehydrated upon drying.

[00272] **Example 2d, HAC-1:** 100 mg of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base was dissolved in 0.1 mL of HOAc at 80°C. To
15 this, 0.2 mL of t-BuOAc was added and the solution was cooled to 20°C. The solution was evaporated to dryness. The resulting solid was agitated in heptane at 50 °C for 15 hours, followed by cooling to 20°C. HAC-1 was filtered and dried at 25°C under vacuum overnight. Form HAC-1 is characterized by 1 mole of acetic acid per one
20 mole of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base. Form HAC-1 was characterized by a DSC thermogram having an endothermic onset typically at ca. 100 °C in agreement with the TGA curve; at higher temperatures other events may ensue. Form HAC-1 was also characterized by a TGA curve having
25 ca. 15.3% weight loss up to ca. 200 °C. Theoretical weight loss is ca. 10.5%, however it is not unusual for minor amount of high boiling point solvents to remain associated with the solid. In cases like this, PXRD is diagnostic of form but not sensitive to small amounts of adventitious solvent.

[00273] **Example 2e, Form E-1:** 50 mg of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base was dissolved in <1mL of boiling ethanol. The
30 solution was cooled to RT and allowed to slowly evaporate. Form E-1 is

characterized by 1 mole of ethanol per one mole of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base. Form E-1 was characterized by a DSC thermogram having an endothermic onset typically at ca. 100 °C in agreement with the TGA curves; at higher temperatures other events may ensue. Form E-1 was characterized by a TGA thermal curve having a weight loss of ca. 7.1% to ca. 7.6% up to ca. 150 °C. Theoretical weight loss is ca. 8.3%, however, it is not unusual for unstable solvates to become partially desolvated upon drying.

[00274] **Example 2f, RPG-3:** 30 to 40 mg of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base was dissolved in 2 mL of racemic propylene glycol. Water was added until a cloud was observed. The solvent was allowed to slowly evaporate to dryness. Form RPG-3 is characterized by 1 molecule of *R*-propylene glycol per one molecule of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base. Form RPG-3 was characterized by a DSC thermogram having an endothermic onset at ca. 70 °C in agreement with the TGA curve, at higher temperatures other events may ensue. Form RPG-3 was characterized by a TGA curve having a weight loss of ca. 16.4% up to ca. 110 °C. Theoretical weight loss is ca. 13.1%, however it is not unusual for minor amount of high boiling-point solvents to remain associated with the solid. In cases like this, PXRD is diagnostic of form but not sensitive to small amounts of adventitious solvent.

[00275] **Example 2g, Form IPA-1:** 40 mg of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base was slurried in <1mL of isopropyl alcohol. The slurry was gently heated to dissolve the remaining solid. The solution was cooled to RT and allowed to slowly evaporate until crystals were observed. Form IPA-1 is characterized by 1 mole of isopropyl alcohol per one mole of N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide, free base.

TABLE 1
Unit Cell Parameters

Compound	Form	T	a(Å)	b(Å)	c(Å)	α°	β°	γ°	V(Å ³)
Exp 2a	H0.5-4	-50	17.7845(7)	7.6215(3)	20.9510(9)	90	109.062(3)	90	2684.1(2)
Exp 2b	N-2	RT	18.7240(4)	8.0171(2)	19.6568(5)	90	114.935(2)	90	2675.7(1)
Exp 2c	H1.75-5	-70	12.7648(2)	34.3194(7)	12.9659(2)	90	99.053(1)	90	5609.4(2)
Exp 2d	HAC-1	RT	7.9766(7)	11.058(2)	33.348(4)	90	90	90	2941.4(6)
Exp 2e	E-1	-50	7.9866(3)	11.2594(6)	32.680(2)	90	90	90	2938.7(2)
Exp 2f	RPG-3	-50	10.3004(3)	10.5475(4)	15.4784(6)	90.045(3)	102.476(2)	109.083(2)	1547.0(1)
Exp 2g	IPA-1	RT	8.4487(2)	11.6615(3)	31.3800(9)	90	90	90	3091.7(1)

TABLE 1 (continued)
Unit Cell Parameters

Compound	Z'	Vm	sg	dcalc
Exp 2a	2	671	P2 ₁	1.235
Exp 2b	2	669	P2 ₁	1.258
Exp 2c	4	701	P2 ₁	1.274
Exp 2d	1	735	P2 ₁ 2 ₁ 2 ₁	1.279
Exp 2e	1	735	P2 ₁ 2 ₁ 2 ₁	1.249
Exp 2f	2	774	P1	1.257
Exp 2g	1	773	P2 ₁ 2 ₁ 2 ₁	1.217

5 [00276] The variables used in Table 1 are defined below:

T = temperature in Centigrade for the crystallographic data (RT is room temperature which is about +22°C)

V = volume of unit cell

Z' = number of drug molecules per asymmetric unit

10 Vm = V(unit cell) / (Z drug molecules per cell)

sg = space group

dcalc = calculated crystal density

TABLE 2

15 **Atomic Coordinates for Example 2a, Form H0.5-4**

Atom	X	Y	Z	Atom	X	Y	Z
F1	0.1718	0.5612	0.2388	F138	0.9730	0.9700	0.6210
F2	0.1576	0.4167	0.1518	F139	1.0220	0.7300	0.6430
F3	0.1277	0.6827	0.1463	F140	0.9470	0.7710	0.5532
N7	0.3729	0.0177	-0.0187	O137	0.5289	1.1320	0.2702
N9	0.3967	0.3613	0.0716	H10	0.3464	0.3422	0.0601
N11	0.5034	0.5324	0.1275	H16	0.1889	-0.1347	-0.0192
N12	0.1513	-0.3723	-0.0803	H85	0.2780	0.4332	0.1193
N15	0.2070	-0.0340	-0.0258	H88	0.4895	0.3182	0.0357

Atom	X	Y	Z	Atom	X	Y	Z
N18	0.4944	0.7660	0.2016	H90	0.2688	-0.4938	-0.1162
O23	0.3677	0.2704	-0.0786	H91	0.2791	-0.3427	-0.1635
O25	0.2371	0.2420	0.0154	H93	0.2277	0.1128	-0.0967
C82	0.2630	0.6176	0.1822	H95	0.3290	-0.0521	-0.1140
C83	0.4281	0.4888	0.1165	H99	0.3760	-0.3135	-0.0574
C84	0.3027	0.5269	0.1466	H100	0.3100	-0.3067	-0.0228
C86	0.3810	0.5754	0.1513	H102	0.5234	0.0450	0.0773
C87	0.4447	0.2515	0.0409	H103	0.4804	0.1033	0.1290
C89	0.2647	-0.3681	-0.1237	H105	0.1184	-0.0766	-0.1500
C92	0.2304	-0.0122	-0.0852	H106	0.1831	-0.0821	-0.1859
C94	0.3163	-0.0776	-0.0729	H110	0.3980	0.9063	0.2542
C96	0.4172	0.7161	0.1933	H112	0.4292	-0.1628	0.0516
C97	0.1821	0.5681	0.1800	H113	0.3692	-0.0449	0.0742
C98	0.3221	-0.2755	-0.0632	H115	0.2715	0.8230	0.2452
C101	0.4735	0.0847	0.0816	H117	0.0540	-0.3914	-0.0071
C104	0.1724	-0.1130	-0.1448	H118	0.1047	-0.5613	-0.0042
C107	0.3915	0.1858	-0.0261	H119	0.0122	-0.5664	-0.0392
C108	0.2116	0.0919	0.0195	H121	0.0157	-0.2070	-0.1065
C109	0.3748	0.8103	0.2276	H122	-0.0438	-0.3568	-0.1418
C111	0.4081	-0.0451	0.0507	H123	0.0156	-0.2818	-0.1762
C114	0.2994	0.7605	0.2220	H126	0.1446	-0.3645	-0.1779
C116	0.0592	-0.4960	-0.0309	H128	0.5835	0.7006	0.1755
C120	0.0087	-0.3106	-0.1338	H130	0.2266	0.0448	0.1189
C124	0.0696	-0.4465	-0.0977	H131	0.1603	-0.0732	0.0701
C125	0.1783	-0.3101	-0.1359	H132	0.1430	0.1249	0.0800
C127	0.5309	0.6696	0.1690	H134	0.0091	-0.6633	-0.1497
C129	0.1827	0.0426	0.0774	H135	0.1005	-0.6984	-0.1175
C133	0.0611	-0.6143	-0.1412	H136	0.0686	-0.5851	-0.1833
F4	0.9541	0.9380	0.5714	H141	0.1830	-0.4440	-0.0580
F5	0.9826	0.6800	0.5759	H14	0.6945	1.0312	0.5339
F6	1.0259	0.8300	0.6655	H21	0.7741	1.5073	0.4379
N8	0.6040	1.3694	0.4758	H28	0.7049	1.4602	0.3064
N13	0.6631	1.0203	0.5573	H29	0.6130	1.4814	0.2902
N17	0.7581	1.7462	0.3780	H32	0.5464	1.0615	0.5430
N19	0.6266	0.8793	0.6383	H34	0.6724	1.6789	0.4661
N20	0.7501	1.4082	0.4350	H35	0.5807	1.7062	0.4509
N22	0.7080	0.6517	0.7080	H38	0.8152	0.9372	0.5749
O24	0.7581	1.1336	0.4774	H41	0.5545	1.4575	0.3823
O26	0.5442	1.1244	0.4164	H44	0.5540	1.3350	0.5873
C27	0.6643	1.5023	0.3242	H45	0.6406	1.2709	0.6290
C30	0.6817	0.9034	0.6081	H48	0.6272	1.8887	0.3818
C31	0.5921	1.1293	0.5402	H49	0.5627	1.7587	0.3377
C33	0.6205	1.6624	0.4324	H52	0.6566	1.2721	0.3752
C36	0.5758	1.2019	0.4701	H55	0.6234	1.5433	0.5534
C37	0.8201	0.8496	0.6069	H56	0.6965	1.4170	0.5619

Atom	X	Y	Z	Atom	X	Y	Z
C39	0.7568	0.8126	0.6288	H58	0.8887	1.9136	0.4290
C40	0.6067	1.4694	0.4172	H59	0.9292	1.8759	0.3743
C42	0.8902	0.7591	0.6317	H60	0.9085	1.7203	0.4146
C43	0.6042	1.2934	0.5840	H62	0.6632	1.7578	0.2922
C46	0.7651	0.6822	0.6794	H64	0.8824	1.3038	0.5620
C47	0.6162	1.7663	0.3697	H65	0.8870	1.4370	0.5064
C50	0.8099	1.8166	0.3410	H66	0.9097	1.2396	0.5019
C51	0.6689	1.3952	0.3877	H69	0.7263	1.9822	0.2803
C53	0.7895	1.2775	0.4740	H70	0.8132	2.0339	0.2846
C54	0.6389	1.4243	0.5469	H71	0.7823	2.0761	0.3450
C57	0.8917	1.8331	0.3947	H73	0.8374	1.5813	0.3078
C61	0.6751	1.6998	0.3362	H74	0.8486	1.7393	0.2638
C63	0.8750	1.3182	0.5148	H75	0.7629	1.6701	0.2558
C67	0.9615	0.8066	0.6143	H77	0.8443	0.5011	0.7349
C68	0.7802	1.9934	0.3099	H79	0.9456	0.5638	0.6960
C72	0.8152	1.6904	0.2872	H81	0.6060	0.7320	0.7067
C76	0.8381	0.5893	0.7029	H142	0.7500	1.8390	0.4090
C78	0.8982	0.6264	0.6799	H143	0.5333	1.1298	0.3123
C80	0.6445	0.7516	0.6862	H144	0.5194	1.0278	0.2538

TABLE 3
Atomic Coordinates for Example 2b, Base Form N-2

Atom	X	Y	Z	Atom	X	Y	Z
N1	0.7824	1.3195	0.3238	C71	0.9289	1.4591	0.5167
N2	0.1685	0.3056	-0.0178	H72	0.9378	1.4237	0.5668
H3	0.1633	0.1989	-0.0196	H73	0.9764	1.4359	0.5102
N4	0.6223	0.8215	0.1887	C74	0.8719	1.7574	0.6080
N5	0.2627	0.4840	-0.0849	C75	0.6719	1.2862	0.4846
N6	0.6868	0.9765	0.2943	H76	0.6246	1.2933	0.4390
H7	0.7030	0.9877	0.3420	H77	0.6814	1.3914	0.5103
N8	0.4185	0.9718	0.0742	H78	0.6660	1.2009	0.5162
N9	0.0989	0.0258	-0.1003	C79	0.9173	1.6199	0.6639
N10	0.7913	1.3680	0.4754	H80	0.8851	1.5217	0.6541
H11	0.7826	1.4618	0.4918	H81	0.9305	1.6578	0.7142
N12	0.5331	0.6005	0.1813	H82	0.9647	1.5943	0.6584
N13	0.3353	0.7474	0.0475	C83	0.9176	1.9198	0.6225
H14	0.3028	0.6896	0.0583	H84	0.9680	1.8999	0.6220
N15	0.8501	1.6946	0.5311	H85	0.9252	1.9631	0.6707
N16	0.4567	1.1267	0.1887	H86	0.8887	1.9992	0.5842
O17	0.7470	1.1042	0.4435	C87	0.7960	1.7901	0.6131
O18	0.2114	0.7470	-0.1165	H88	0.7660	1.8705	0.5758
O19	0.2229	0.5239	0.0583	H89	0.8064	1.8325	0.6620
O20	0.8588	1.0890	0.3360	H90	0.7667	1.6881	0.6048

Atom	X	Y	Z	Atom	X	Y	Z
F21	0.5881	0.9034	0.4961	C91	0.2069	0.3753	0.0494
F22	0.4828	0.7754	0.4659	C92	0.1886	0.3981	-0.1294
F23	0.5874	0.6500	0.5199	H93	0.1617	0.4641	-0.1751
C24	0.1049	-0.1653	-0.0049	C94	0.1345	0.3960	-0.0889
H25	0.1473	-0.2167	-0.0125	H95	0.1249	0.5115	-0.0787
H26	0.0773	-0.2487	0.0097	C96	0.3636	0.8982	0.1612
H27	0.1257	-0.0826	0.0338	C97	0.0560	0.3169	-0.1396
C28	0.0052	-0.2107	-0.1371	H98	0.0300	0.3865	-0.1836
H29	-0.0334	-0.1565	-0.1806	H99	0.0226	0.3123	-0.1131
H30	-0.0204	-0.2889	-0.1178	C100	0.0655	0.1409	-0.1647
H31	0.0423	-0.2685	-0.1506	H101	0.0138	0.0994	-0.1999
C32	-0.0102	0.0236	-0.0618	C102	0.2651	0.6526	-0.0801
H33	0.0176	0.1111	-0.0276	C103	0.3727	0.8727	0.0930
H34	-0.0371	-0.0447	-0.0399	C104	0.3129	0.8065	0.1831
H35	-0.0478	0.0713	-0.1077	H105	0.2826	0.7213	0.1525
C36	0.6347	0.8573	0.2590	C106	0.4090	1.0275	0.2081
C37	0.7956	1.1614	0.3094	C107	0.1998	0.2249	-0.1544
C38	0.5429	0.6399	0.2526	H108	0.2268	0.1548	-0.1108
C39	0.5969	0.8058	0.3661	H109	0.2321	0.2314	-0.1820
H40	0.6303	0.8901	0.3943	C110	0.0479	-0.0823	-0.0781
C41	0.5928	0.7673	0.2949	C111	0.3943	0.5437	-0.0102
C42	0.7174	1.0886	0.2543	H112	0.4319	0.5306	0.0416
H43	0.7243	1.0288	0.2140	H113	0.4226	0.5469	-0.0417
C44	0.8437	1.4188	0.3806	C114	0.1202	0.1477	-0.2041
H45	0.8919	1.4020	0.3733	H115	0.0955	0.2128	-0.2496
C46	0.8266	1.6034	0.3730	H116	0.1283	0.0355	-0.2179
H47	0.7791	1.6255	0.3797	C117	0.3076	0.8411	0.2485
H48	0.8177	1.6396	0.3230	C118	0.2292	0.2619	0.1158
C49	0.5524	0.7201	0.3942	H119	0.2846	0.2720	0.1469
C50	0.6668	1.2414	0.2237	H120	0.2172	0.1487	0.0989
H51	0.6690	1.2785	0.1776	H121	0.2001	0.2924	0.1441
H52	0.6125	1.2182	0.2137	C122	0.3349	0.4024	-0.0337
C53	0.8609	1.3569	0.4595	H123	0.3502	0.3144	-0.0588
H54	0.8773	1.2398	0.4636	H124	0.3290	0.3556	0.0092
C55	0.4991	0.5496	0.2836	C125	0.3458	0.7009	-0.0200
H56	0.4668	0.4624	0.2569	H126	0.3682	0.7937	-0.0371
C57	0.7010	1.3718	0.2841	C127	0.4561	1.0941	0.1233
H58	0.6975	1.4822	0.2628	H128	0.4859	1.1659	0.1085
H59	0.6744	1.3722	0.3171	C129	0.4030	1.0567	0.2762
C60	0.9137	1.6472	0.5099	H130	0.4339	1.1390	0.3084
H61	0.9621	1.7039	0.5438	C131	0.2548	0.7395	0.2713
C62	0.7404	1.2441	0.4666	C132	0.3536	0.9679	0.2955
C63	0.5039	0.5898	0.3526	H133	0.3500	0.9907	0.3403
H64	0.4747	0.5300	0.3726	F134	0.1847	0.7359	0.2156
C65	0.5722	0.6959	0.1556	F135	0.2364	0.8060	0.3220

Atom	X	Y	Z	Atom	X	Y	Z
H66	0.5642	0.6734	0.1065	F136	0.2859	0.5920	0.2960
C67	0.8949	1.7014	0.4308	F137	0.1977	0.8329	0.2710
H68	0.9412	1.6866	0.4211	F138	0.2168	0.6170	0.2301
H69	0.8818	1.8193	0.4256	F139	0.2934	0.6790	0.3395
C70	0.5532	0.7632	0.4678	H140	0.8280	1.7679	0.5060
				H141	0.1330	-0.0400	-0.1062

TABLE 4
Atomic Coordinates for Example 2c, Base Form H1.75-5

Atom	X	Y	Z	Atom	X	Y	Z
C1	0.3942	0.1176	0.2183	C51	0.6094	0.0443	-0.0547
C2	0.2287	0.1174	0.0897	C52	0.6750	0.0382	0.0420
N3	0.1192	0.1293	0.0594	C53	0.9963	0.1572	0.4974
N4	0.1798	0.0501	0.1322	C54	0.7176	-0.0005	0.0684
N5	0.3753	0.0509	0.2822	C55	1.1985	0.0503	0.7049
N6	-0.0927	0.1070	-0.1055	C56	1.1539	0.1133	0.4965
N7	-0.3691	0.1154	-0.3165	C57	1.0229	0.1608	0.6162
N8	-0.2260	0.1440	-0.1970	C58	1.0367	0.1185	0.4572
O9	0.1323	0.1298	-0.1137	C59	0.7341	0.1669	0.3280
O10	0.0301	0.0456	0.0130	C60	1.1829	0.1168	0.6162
C11	0.0785	0.1332	-0.0419	C61	1.1435	0.1563	0.6496
C12	0.2431	0.0735	0.0697	C62	0.5636	0.1744	0.0306
C13	-0.2272	0.0747	-0.2281	C63	0.6973	0.1581	0.4320
C14	0.0812	0.0382	0.0995	C64	0.7972	0.1517	0.5115
C15	0.4135	0.0738	0.2008	C65	1.3015	0.0605	0.7775
C16	0.2774	0.1283	0.2010	C66	1.2257	0.0279	0.6122
C17	0.3592	0.0623	0.0929	C67	1.1287	0.0258	0.7652
C18	-0.3242	0.0799	-0.2978	C68	0.9017	0.0664	0.4282
C19	-0.1859	-0.0329	-0.2601	C69	0.8592	0.0308	0.4749
C20	-0.3192	0.1441	-0.2671	F70	0.7706	-0.0040	0.1659
C21	-0.1804	0.1091	-0.1748	F71	0.7890	-0.0119	0.0101
C22	-0.1842	0.0372	-0.2149	F72	0.6477	-0.0283	0.0583
C23	0.0394	0.1336	0.1281	N73	0.7366	0.2979	0.1640
C24	-0.3330	0.0112	-0.3340	N74	0.9286	0.2769	0.3415
C25	-0.2347	0.0067	-0.2683	N75	0.9490	0.3548	0.4399
C26	0.0308	0.0146	0.1792	N76	0.6679	0.2686	0.0076
C27	-0.3753	0.0465	-0.3473	N77	0.5468	0.3015	-0.1239
C28	-0.0381	0.1410	-0.0545	N78	1.1302	0.3587	0.6021
C29	0.4391	0.0161	0.3216	O79	0.7689	0.2552	0.3756
C30	0.4331	-0.0152	0.2365	O80	0.8000	0.3561	0.3182
C31	-0.0565	0.1490	0.0571	C81	0.6779	0.3005	0.0685
C32	0.5574	0.0267	0.3640	C82	0.8021	0.2636	0.1953
C33	0.3907	0.0006	0.4127	C83	0.6253	0.3363	0.0330

Atom	X	Y	Z	Atom	X	Y	Z
F34	-0.1107	-0.0374	-0.1835	C84	0.8293	0.2641	0.3145
F35	-0.1514	-0.0419	-0.3469	C85	0.9704	0.2829	0.4523
F36	-0.2560	-0.0605	-0.2519	C86	0.9612	0.3232	0.6101
N37	0.9782	0.0854	0.4895	C87	1.0815	0.3238	0.6395
N38	0.8832	0.1624	0.4564	C88	0.6395	0.3717	0.0871
N39	0.6198	0.1729	0.1273	C89	0.9221	0.3194	0.4932
N40	0.7167	0.1350	0.2533	C90	0.5107	0.3694	-0.1081
N41	0.5429	0.1472	-0.0401	C91	0.5605	0.3350	-0.0649
N42	1.1331	0.0855	0.6708	C92	0.6021	0.2715	-0.0843
O43	0.9140	0.1821	0.2960	C93	0.5266	0.4037	-0.0538
O44	0.8654	0.0771	0.3382	C94	0.9119	0.2659	0.1620
C45	0.8542	0.1720	0.3563	C95	0.5911	0.4051	0.0443
C46	0.5876	0.1117	-0.0112	C96	1.1535	0.3924	0.6738
C47	0.6629	0.1387	0.1561	C97	0.6068	0.4429	0.0984
C48	0.6954	0.0688	0.1114	C98	0.8832	0.3710	0.3595
C49	0.6512	0.1060	0.0865	C99	1.0912	0.2839	0.4757
C50	0.5679	0.0801	-0.0813	C100	1.1264	0.2872	0.5933
C101	0.9810	0.2861	0.2520	O151	0.4292	0.1756	0.7593
C102	0.9190	0.4094	0.3235	O152	0.3473	0.1636	0.8339
C103	1.0516	0.4113	0.6921	H153	0.4310	0.1254	0.2979
C104	1.2130	0.3807	0.7821	H154	0.4349	0.1341	0.1645
C105	1.2167	0.4208	0.6206	H155	0.2779	0.1337	0.0447
F106	0.6836	0.4427	0.1806	H156	0.2164	0.0417	0.2100
F107	0.5186	0.4543	0.1375	H157	0.3714	0.0703	0.3474
F108	0.6281	0.4715	0.0364	H158	-0.0576	0.0785	-0.0868
N109	0.3031	0.2523	0.0895	H159	0.2173	0.0675	-0.0123
N110	0.3202	0.3294	0.1970	H160	0.4992	0.0681	0.2061
N111	0.5022	0.3356	0.3522	H161	0.2372	0.1127	0.2569
N112	0.0946	0.2741	-0.0674	H162	0.2676	0.1594	0.2134
N113	-0.0336	0.2377	-0.1697	H163	0.3639	0.0302	0.0854
N114	-0.2016	0.2650	-0.2433	H164	0.4006	0.0751	0.0350
O115	0.1682	0.3314	0.0829	H165	-0.3534	0.1727	-0.2828
O116	0.1499	0.2237	0.1243	H166	-0.1099	0.0321	-0.1626
C117	0.2052	0.2373	0.0644	H167	0.0215	0.1059	0.1611
C118	0.2992	0.2930	0.2475	H168	0.0659	0.1539	0.1907
C119	-0.0515	0.3401	-0.0764	H169	-0.3750	-0.0142	-0.3730
C120	0.3416	0.2960	0.3644	H170	-0.0484	0.0061	0.1467
C121	-0.1222	0.3701	-0.0923	H171	0.0783	-0.0115	0.2000
C122	0.4698	0.2615	0.2174	H172	0.0297	0.0317	0.2489
C123	-0.1737	0.2997	-0.1931	H173	-0.4526	0.0496	-0.3983
C124	-0.0044	0.2713	-0.1206	H174	-0.0573	0.1673	-0.1021
C125	-0.2438	0.3311	-0.2090	H175	0.4770	-0.0409	0.2650
C126	-0.0754	0.3040	-0.1278	H176	0.3484	-0.0242	0.2146
C127	0.1701	0.2420	-0.0535	H177	0.4596	-0.0045	0.1679
C128	0.3488	0.2578	0.2007	H178	-0.1282	0.1348	0.0707

Atom	X	Y	Z	Atom	X	Y	Z
C129	0.5097	0.2640	0.3350	H179	-0.0650	0.1804	0.0686
C130	0.4624	0.2986	0.3883	H180	0.5963	-0.0009	0.3924
C131	-0.0968	0.4077	-0.0362	H181	0.5930	0.0381	0.3032
C132	0.3460	0.2683	0.0015	H182	0.5596	0.0461	0.4284
C133	-0.1317	0.2372	-0.2271	H183	0.4332	-0.0251	0.4433
C134	0.5253	0.3677	0.4289	H184	0.3902	0.0220	0.4712
C135	-0.2190	0.3662	-0.1602	H185	0.3084	-0.0081	0.3820
C136	0.2745	0.2511	-0.0941	H186	0.9987	0.0754	0.5691
C137	0.5618	0.4023	0.3689	H187	0.7487	0.1066	0.2786
C138	0.6152	0.3567	0.5186	H188	1.1096	0.0983	0.7400
C139	0.4264	0.3788	0.4754	H189	0.7448	0.0641	0.1858
C140	0.2540	0.3461	0.1186	H190	0.5192	0.0845	-0.1565
C141	0.2900	0.3831	0.0758	H191	0.5930	0.0200	-0.1086
F142	0.0020	0.4176	-0.0257	H192	1.0402	0.1792	0.4633
F143	-0.1497	0.4372	-0.0806	H193	1.1791	0.0849	0.4744
F144	-0.1239	0.4079	0.0550	H194	1.1975	0.1355	0.4619
O145	0.4346	0.1106	0.5123	H195	0.9975	0.1890	0.6409
O146	0.1371	0.2037	0.3256	H196	0.9825	0.1382	0.6530
O147	0.6629	0.2454	0.5702	H197	1.0259	0.1190	0.3726
O148	0.4523	0.2291	0.5970	H198	0.6972	0.1932	0.2915
O149	0.3986	0.3042	0.6776	H199	1.2687	0.1146	0.6366
O150	0.3459	0.1840	0.4326	H200	1.1831	0.1795	0.6123
H201	1.1631	0.1593	0.7328	H251	1.1684	0.4299	0.5453
H202	0.5306	0.2029	0.0082	H252	0.3950	0.3437	0.2241
H203	0.6519	0.1824	0.4545	H253	0.5749	0.3293	0.3230
H204	0.6483	0.1323	0.4245	H254	0.1193	0.3022	-0.0335
H205	0.8029	0.1216	0.5375	H255	0.2147	0.2880	0.2355
H206	0.7957	0.1702	0.5801	H256	0.0233	0.3437	-0.0249
H207	1.3466	0.0340	0.8006	H257	0.3083	0.3215	0.3958
H208	1.3500	0.0795	0.7383	H258	0.3160	0.2703	0.4028
H209	1.2843	0.0747	0.8476	H259	0.5050	0.2366	0.1841
H210	1.2710	0.0024	0.6388	H260	0.4927	0.2878	0.1790
H211	1.1517	0.0185	0.5635	H261	-0.3192	0.3277	-0.2606
H212	1.2687	0.0462	0.5671	H262	0.1323	0.2153	-0.0854
H213	1.1703	-0.0007	0.7926	H263	0.3327	0.2320	0.2439
H214	1.1105	0.0421	0.8313	H264	0.4885	0.2372	0.3700
H215	1.0556	0.0182	0.7149	H265	0.5951	0.2670	0.3470
H216	0.7976	0.0176	0.4203	H266	0.4874	0.2957	0.4712
H217	0.9226	0.0100	0.4973	H267	0.3419	0.2998	0.0019
H218	0.8269	0.0390	0.5453	H268	0.4278	0.2594	0.0040
H219	0.7349	0.3217	0.2190	H269	-0.1537	0.2100	-0.2663
H220	1.0259	0.3684	0.4660	H270	-0.2738	0.3905	-0.1734
H221	1.2052	0.3497	0.5802	H271	0.3087	0.2249	-0.1215
H222	0.7595	0.2370	0.1705	H272	0.2609	0.2719	-0.1576
H223	0.9510	0.2576	0.4964	H273	0.5805	0.4266	0.4203

Atom	X	Y	Z	Atom	X	Y	Z
H224	0.9305	0.2994	0.6498	H274	0.6328	0.3938	0.3368
H225	0.9306	0.3505	0.6365	H275	0.5006	0.4101	0.3051
H226	1.1023	0.3226	0.7234	H276	0.6311	0.3804	0.5741
H227	0.6892	0.3732	0.1634	H277	0.5931	0.3310	0.5600
H228	0.8361	0.3164	0.4802	H278	0.6878	0.3500	0.4880
H229	0.4622	0.3687	-0.1842	H279	0.4455	0.4020	0.5307
H230	0.5955	0.2449	-0.1310	H280	0.3644	0.3877	0.4139
H231	0.4887	0.4301	-0.0865	H281	0.4001	0.3535	0.5147
H232	0.9411	0.2373	0.1483	H282	0.2315	0.3935	0.0139
H233	0.9065	0.2830	0.0906	H283	0.3050	0.4046	0.1373
H234	1.1232	0.2576	0.4474	H284	0.3642	0.3778	0.0457
H235	1.1194	0.3088	0.4371	H297	0.5031	0.1166	0.5720
H236	1.2129	0.2887	0.6081	H298	0.4013	0.1383	0.4827
H237	1.1010	0.2618	0.6307	H291	0.1415	0.2116	0.2456
H238	0.9841	0.3173	0.2399	H292	0.0561	0.1944	0.3301
H239	1.0614	0.2747	0.2630	H285	0.7189	0.2533	0.6386
H240	0.8608	0.4208	0.2603	H286	0.7007	0.2488	0.5014
H241	0.9250	0.4304	0.3884	H287	0.4531	0.2087	0.6605
H242	0.9943	0.4069	0.2987	H288	0.5334	0.2354	0.5869
H243	1.0681	0.4363	0.7411	H289	0.4187	0.2763	0.6478
H244	1.0091	0.4217	0.6147	H290	0.4459	0.3091	0.7532
H245	1.0017	0.3911	0.7227	H293	0.3695	0.2054	0.4921
H246	1.2276	0.4062	0.8315	H294	0.2663	0.1913	0.3921
H247	1.1658	0.3601	0.8194	H295	0.4701	0.1653	0.8329
H248	1.2882	0.3671	0.7752	H296	0.4645	0.1639	0.6962
H249	1.2341	0.4465	0.6673	H299	0.2731	0.1523	0.8513
H250	1.2881	0.4078	0.6030	H300	0.4117	0.1542	0.8909

TABLE 5

Atomic Coordinates for Example 2e, Base Form E-1

Atom	X	Y	Z	Atom	X	Y	Z
F18	0.2662	0.4002	0.0274	F23	0.0301	0.3722	0.0039
F19	0.0473	0.4191	0.0033	C18	0.1648	0.3165	0.0297
F20	0.1401	0.2386	0.0059	H31	0.2375	0.5477	0.2387
O2	0.3035	0.7582	0.1985	H41	0.5018	0.4801	0.2585
O28	0.4277	0.5819	0.1087	H42	0.4761	0.3850	0.2161
N1	0.5559	0.6627	0.1978	H51	0.6480	0.5043	0.1733
N6	0.2412	0.4989	0.1766	H52	0.7376	0.5463	0.2204
N8	0.0422	0.3993	0.2139	H61	0.2881	0.5282	0.1472
N10	-0.1606	0.2714	0.1828	H91	-0.1519	0.3157	0.2435
N27	0.6744	0.6813	0.1115	H111	-0.2586	0.1707	0.1140
N29	0.9824	0.8036	0.1064	H121	-0.1207	0.1964	0.0469
C2	0.3877	0.6687	0.2028	H141	0.2343	0.4431	0.0976

Atom	X	Y	Z	Atom	X	Y	Z
C3	0.3235	0.5438	0.2136	H211	0.5921	0.8399	0.1984
C4	0.4822	0.4775	0.2252	H221	0.4827	0.7942	0.1303
C5	0.6231	0.5426	0.2028	H231	0.6846	0.8993	0.0877
C7	0.1130	0.4242	0.1780	H232	0.6444	0.9699	0.1348
C9	-0.0900	0.3261	0.2139	H241	0.9421	0.9813	0.1221
C11	-0.1488	0.2295	0.1107	H251	1.0517	0.8711	0.1826
C12	-0.0761	0.2450	0.0735	H252	0.8644	0.9484	0.1930
C13	0.0600	0.3235	0.0687	H261	0.8551	0.7433	0.2231
C14	0.1243	0.3844	0.1017	H262	0.8872	0.6821	0.1741
C15	-0.0857	0.2914	0.1453	H271	0.8051	0.6804	0.1028
C16	0.0498	0.3680	0.1409	H291	1.0794	0.7695	0.1261
C17	0.1207	0.3503	0.0270	H294	0.7893	0.5312	0.0656
C21	0.6470	0.7650	0.1817	H292	0.6583	0.4189	0.0854
C22	0.6168	0.7817	0.1361	H293	0.5951	0.5011	0.0421
C23	0.7030	0.8953	0.1212	H311	0.9885	0.9169	0.0098
C24	0.8922	0.8952	0.1304	H312	0.8322	0.8277	0.0324
C25	0.9185	0.8727	0.1765	H313	0.8750	0.9701	0.0523
C26	0.8353	0.7590	0.1910	H321	1.2632	0.9515	0.0466
C28	0.5782	0.5936	0.0987	H322	1.1556	1.0063	0.0897
C29	0.6596	0.5024	0.0711	H323	1.2983	0.8878	0.0951
C30	1.0633	0.8393	0.0676	H331	1.1943	0.7459	0.0197
C31	0.9326	0.8931	0.0389	H332	1.2251	0.6857	0.0690
C32	1.2044	0.9276	0.0745	H333	1.0345	0.6616	0.0430
C33	1.1314	0.7245	0.0492	H971	-0.8355	0.1305	0.1615
O99	-0.4402	0.1232	0.1894	H972	-0.7164	0.0043	0.1747
C97	-0.7313	0.0985	0.1797	H973	-0.7551	0.1157	0.2115
C98	-0.5812	0.1608	0.1676	H981	-0.5951	0.2544	0.1716
F21	0.3052	0.3620	0.0283	H982	-0.5565	0.1430	0.1349
F22	0.1907	0.2403	0.0143	H991	-0.3383	0.1855	0.1847

TABLE 6
Atomic Coordinates for Example 2d, Base Form HAC-1

Atom	X	Y	Z	Atom	X	Y	Z
C1	0.0418	0.3612	0.1413	H32	-0.1504	0.2133	0.0512
N2	-0.1614	0.2566	0.1827	H21	0.2024	0.4361	0.1001
N3	0.5574	0.6476	0.1969	H19	-0.2633	0.1786	0.1141
O4	0.4002	0.5938	0.1104	H15	-0.1260	0.2877	0.2400
N5	0.0480	0.3793	0.2136	H6	0.2703	0.5248	0.1514
N6	0.2349	0.4911	0.1766	H14	0.2507	0.5355	0.2354
N7	0.6581	0.6819	0.1120	H25A	0.5017	0.4595	0.2502
O9	0.3141	0.7518	0.2032	H25B	0.4643	0.3765	0.2131
C10	0.6548	0.7519	0.1823	H22A	0.6302	0.4915	0.1724
C11	0.1091	0.4116	0.1779	H22B	0.7255	0.5203	0.2124

Atom	X	Y	Z	Atom	X	Y	Z
C12	0.6143	0.7805	0.1389	H10	0.6208	0.8184	0.1990
C13	-0.0977	0.2822	0.1453	H12	0.4961	0.7944	0.1361
C14	0.3251	0.5321	0.2128	H31A	0.7369	0.5221	0.0663
C15	-0.0849	0.3037	0.2135	H31B	0.5607	0.5108	0.0459
C16	0.5496	0.5972	0.0993	H31C	0.6086	0.4250	0.0815
C17	0.8422	0.7397	0.1888	H17A	0.8651	0.7215	0.2164
N18	0.9734	0.7976	0.1050	H17B	0.8826	0.6751	0.1722
C19	-0.1674	0.2302	0.1113	H23A	0.6885	0.9061	0.0978
C20	0.0380	0.3268	0.0698	H23B	0.6640	0.9626	0.1405
C21	0.1085	0.3827	0.1029	H28A	0.8869	0.9177	0.1920
C22	0.6188	0.5229	0.1991	H28B	1.0495	0.8473	0.1796
C23	0.7063	0.8942	0.1260	H27	0.9448	0.9635	0.1263
C24	0.3943	0.6572	0.2042	H18	1.0626	0.7570	0.1189
C25	0.4808	0.4583	0.2218	H39A	1.2479	0.9507	0.0498
C27	0.8951	0.8864	0.1321	H39B	1.2853	0.8792	0.0894
C28	0.9305	0.8535	0.1757	H39C	1.1647	0.9911	0.0903
F29	0.2449	0.4047	0.0277	H38A	0.9706	0.9352	0.0175
C30	1.0501	0.8412	0.0669	H38B	0.8802	0.9740	0.0573
C31	0.6205	0.5052	0.0708	H38C	0.8310	0.8527	0.0360
C32	-0.1018	0.2511	0.0743	H40A	1.1575	0.7492	0.0195
C33	0.1058	0.3467	0.0300	H40B	1.0092	0.6805	0.0403
F36	0.1168	0.2494	0.0085	H40C	1.1861	0.6845	0.0608
F37	0.0041	0.4132	0.0068	H7	0.7708	0.6743	0.1022
C38	0.9217	0.9066	0.0421	H35A	-0.8134	0.1022	0.1656
C39	1.2014	0.9229	0.0747	H35B	-0.7586	0.1045	0.2107
C40	1.1073	0.7290	0.0448	H35C	-0.7035	-0.0029	0.1830
O8	-0.4446	0.1296	0.1963	H8	-0.3548	0.1817	0.1876
C26	-0.5724	0.1491	0.1735				
O34	-0.5673	0.2213	0.1458				
C35	-0.7251	0.0824	0.1840				

TABLE 7
Atomic Coordinates for Example 2g, Base Form IPA-1

Atom	X	Y	Z	Atom	X	Y	Z
F1	0.1519	0.3786	0.0243	H4B	0.8724	0.7912	0.1232
F3	-0.0140	0.2540	-0.0043	H3	0.6171	0.6809	0.1026
F2	0.0260	0.4241	0.0085	H1	0.1162	0.5505	0.1460
N4	0.8062	0.8075	0.1016	H10	0.3362	0.7935	0.1099
N3	0.5159	0.6834	0.1054	H9	0.3991	0.8799	0.1736
N2	0.3481	0.7197	0.1890	H6A	0.4782	0.9618	0.1051
N1	0.0675	0.5440	0.1700	H6B	0.5369	0.8805	0.0686
N5	-0.1311	0.4643	0.2095	H22	0.0665	0.4473	0.0929
N6	-0.2941	0.3121	0.1828	H5	0.7491	0.9735	0.1042

Atom	X	Y	Z	Atom	X	Y	Z
O1	0.1140	0.8031	0.1697	H21	-0.3360	0.1865	0.1167
O2	0.2925	0.5779	0.1031	H12A	0.5507	0.4357	0.0989
C10	0.4463	0.7911	0.1197	H12B	0.6252	0.5211	0.0660
C9	0.4471	0.8051	0.1677	H12C	0.4678	0.4565	0.0549
C4	0.9030	0.8194	0.0629	H8A	0.6698	0.7392	0.1786
C24	-0.0632	0.3232	0.0669	H8B	0.6137	0.8216	0.2153
C19	-0.0909	0.3946	0.1378	H16	0.0302	0.6334	0.2243
C6	0.5348	0.8912	0.0992	H13A	0.4882	0.6277	0.2283
C20	-0.2139	0.3158	0.1454	H13B	0.4421	0.5607	0.1863
C11	0.4342	0.5879	0.0964	H3B	0.9771	0.9859	0.0712
C22	-0.0159	0.3962	0.0980	H3C	1.0944	0.9144	0.0431
C5	0.7048	0.9021	0.1156	H3D	1.0915	0.8973	0.0926
C17	-0.0508	0.4684	0.1726	H23	-0.2112	0.1922	0.0537
C21	-0.2559	0.2399	0.1123	H18	-0.3052	0.3848	0.2368
C14	0.1869	0.7250	0.1863	H2A	0.7158	0.7853	0.0234
C12	0.5279	0.4917	0.0774	H2B	0.8604	0.8380	-0.0010
C8	0.6155	0.8107	0.1847	H2C	0.7518	0.9169	0.0270
C16	0.1187	0.6161	0.2052	H7A	0.6551	0.9825	0.1718
C13	0.4041	0.6134	0.2080	H7B	0.8129	0.9118	0.1736
C3	1.0281	0.9129	0.0679	H1A	1.0319	0.6803	0.0846
C23	-0.1824	0.2437	0.0749	H1B	1.0662	0.7077	0.0366
C18	-0.2485	0.3861	0.2114	H1C	0.9087	0.6458	0.0495
C2	0.7981	0.8420	0.0246	H15A	0.2542	0.5916	0.2593
C7	0.7045	0.9108	0.1635	H15B	0.2564	0.4841	0.2288
C1	0.9852	0.7023	0.0579	H3A	-0.4990	0.1985	0.1863
C15	0.2577	0.5672	0.2298	H27	-0.6926	0.2734	0.1759
C25	0.0153	0.3263	0.0246	H26A	-0.7745	0.0861	0.1320
F4	0.1640	0.2980	0.0315	H26B	-0.6492	0.1771	0.1173
F5	-0.0739	0.3770	-0.0036	H26C	-0.8259	0.2136	0.1243
F6	0.0440	0.2214	0.0112	H28A	-0.8028	0.1216	0.2237
O3	-0.5666	0.1486	0.1890	H28B	-0.9203	0.1527	0.1868
C27	-0.7093	0.1903	0.1763	H28C	-0.8515	0.2497	0.2160
C26	-0.7420	0.1648	0.1343				
C28	-0.8290	0.1780	0.2024				
C29	-0.8460	0.1070	0.1744				
C30	-0.6990	0.2550	0.1410				

TABLE 8

Atomic Coordinates for Example 2f, Base Form RPG-3

Atom	X	Y	Z	Atom	X	Y	Z
F1	1.4700	0.5720	0.6614	N2	0.6350	0.3813	0.3765
F7	1.4119	0.5575	0.7000	N6	0.2687	0.8628	0.3992
F2	1.4315	0.5398	0.7741	N4	0.3398	0.7349	0.3149

Atom	X	Y	Z	Atom	X	Y	Z
F8	1.5600	0.5336	0.8131	N1	0.8898	0.3354	0.4598
F3	1.6320	0.5630	0.7860	N5	0.0527	0.9107	0.3489
F9	1.6111	0.5864	0.6960	C12	0.4202	0.2078	0.3911
N9	0.7997	-0.0295	0.8348	O1	0.6129	0.7507	0.2839
N8	0.7989	0.1232	0.9974	C10	0.7273	0.4981	0.3447
N11	1.1628	-0.0946	0.6829	C6	0.8597	0.4735	0.3332
N10	1.0937	0.0679	0.7424	C9	0.7661	0.6239	0.4071
N7	0.5331	0.0253	1.0441	C20	0.1173	0.7695	0.2548
N12	1.3802	-0.0414	0.6342	C19	0.0264	0.8385	0.2716
O3	0.8190	0.0931	0.7142	C5	0.9561	0.4627	0.4193
O4	1.0177	0.2159	0.9737	C17	0.2435	0.7914	0.3234
C34	0.7187	0.1296	0.9094	C8	0.8573	0.6109	0.4957
C45	1.3119	0.1383	0.6926	O2	0.4273	0.3887	0.2980
C37	0.6769	0.0002	0.8509	C24	-0.0460	0.6680	0.1174
C30	0.4799	0.0412	0.9508	C16	0.4606	0.7449	0.3864
C42	1.1869	0.0349	0.7058	C22	0.0802	0.6875	0.1776
C31	0.5850	0.1596	0.9161	C14	0.5786	0.7190	0.3544
C49	1.4737	0.3643	0.7029	C7	0.9892	0.5850	0.4828
C33	0.5750	-0.1179	0.8869	C23	-0.1359	0.7367	0.1342
C47	1.3487	0.2751	0.7153	C18	0.1707	0.9200	0.4061
C40	0.8584	0.0198	0.7679	C25	-0.0848	0.5788	0.0372
C32	0.4447	-0.0846	0.8933	C13	0.5736	0.6259	0.4906
C41	0.9773	-0.0327	0.7663	C21	-0.1010	0.8179	0.2096
C39	1.0115	-0.0750	0.8618	C4	0.9438	0.1985	0.3554
C35	0.9405	0.1730	1.0250	C3	0.9469	0.2256	0.4509
C38	0.8681	-0.1166	0.8863	C15	0.4317	0.6466	0.4565
C28	0.4760	0.0765	1.1130	C2	1.0956	0.2572	0.5066
C43	1.2625	-0.1234	0.6483	C11	0.4953	0.3347	0.3508
C44	1.4064	0.0935	0.6556	C1	0.8466	0.1026	0.4843
C46	1.5338	0.1870	0.6430	F4	-0.1040	0.4554	0.0598
C48	1.5649	0.3202	0.6666	F5	0.0211	0.6100	-0.0030
C36	1.0000	0.1703	1.1211	F6	-0.1869	0.5900	-0.0238
C29	0.5552	0.0482	1.2020	F10	-0.2221	0.5070	0.0190
C27	0.3192	0.0052	1.1006	F11	-0.0370	0.4791	0.0400
C26	0.5156	0.2276	1.1108	F12	-0.0360	0.6334	-0.0302
C50	1.5103	0.5068	0.7292	O7	0.3015	0.5242	0.1820
N3	0.6420	0.6560	0.4169	O8	0.5067	0.7155	0.1016
C55	0.4573	0.5719	0.0792	H12C	0.3410	0.1518	0.3473
C56	0.3131	0.5089	0.0936	H10A	0.6770	0.5138	0.2867
C54	0.4592	0.5517	-0.0152	H6A	0.8326	0.3912	0.2958
O5	1.1135	0.3136	0.8309	H6B	0.9113	0.5467	0.3027
O6	0.9224	0.3638	0.6775	H9	0.8236	0.6994	0.3801
C52	0.9658	0.4383	0.7607	H95	1.0446	0.4603	0.4067
C53	1.0998	0.4403	0.8150	H8B	0.8035	0.5371	0.5249
C51	0.9706	0.5801	0.7445	H8C	0.8845	0.6928	0.5336

Atom	X	Y	Z	Atom	X	Y	Z
H8	0.7521	0.0843	1.0354	H16	0.4967	0.8363	0.4153
H10	1.1043	0.1515	0.7516	H22	0.1413	0.6448	0.1660
H34	0.7767	0.2034	0.8813	H7A	1.0479	0.6636	0.4601
H37	0.6246	0.0147	0.7931	H77B	1.0425	0.5720	0.5399
H30	0.3926	0.0620	0.9462	H23	-0.2199	0.7261	0.0931
H31A	0.5395	0.1776	0.8580	H18	0.1897	0.9731	0.4584
H31B	0.6110	0.2396	0.9557	H13A	0.6288	0.6865	0.5425
H33A	0.5480	-0.1985	0.8476	H13B	0.5604	0.5340	0.5057
H33B	0.6213	-0.1349	0.9450	H21	-0.1626	0.8605	0.2204
H47	1.2884	0.3066	0.7391	H4A	0.8479	0.1679	0.3217
H32A	0.3953	-0.0742	0.8343	H4B	0.9856	0.1305	0.3502
H32B	0.3818	-0.1587	0.9172	H4C	0.9960	0.2797	0.3330
H41	0.9434	-0.1120	0.7240	H15A	0.3587	0.5626	0.4312
H39A	1.0471	-0.1496	0.8633	H15B	0.4034	0.6838	0.5038
H39B	1.0801	-0.0006	0.9013	H2A	1.1552	0.3410	0.4913
H38A	0.8152	-0.2110	0.8689	H2B	1.1310	0.1867	0.4955
H38B	0.8786	-0.0998	0.9495	H2C	1.0948	0.2640	0.5683
H43	1.2434	-0.2140	0.6327	H1A	0.8496	0.1213	0.5456
H46	1.5955	0.1582	0.6190	H1B	0.8758	0.0261	0.4785
H48	1.6483	0.3822	0.6582	H1C	0.7522	0.0839	0.4497
H36A	1.0650	0.1213	1.1285	H99	0.9009	0.3500	0.5070
H36B	0.9249	0.1270	1.1498	H7	0.3628	0.5029	0.2157
H36C	1.0484	0.2607	1.1471	H8A	0.5335	0.7307	0.1555
H29A	0.5101	-0.0427	1.2144	H55	0.5214	0.5322	0.1164
H29B	0.5546	0.1090	1.2479	H56A	0.2502	0.5475	0.0555
H29C	0.6509	0.0606	1.1995	H56B	0.2814	0.4136	0.0755
H27A	0.2727	0.0063	1.0399	H54A	0.4034	0.5980	-0.0510
H27B	0.2829	0.0501	1.1386	H54B	0.4208	0.4574	-0.0335
H27C	0.3022	-0.0863	1.1155	H54C	0.5546	0.5867	-0.0221
H26A	0.6150	0.2666	1.1142	H5	1.0835	0.2863	0.8748
H26B	0.4925	0.2636	1.1602	H6	0.9213	0.2865	0.6847
H26C	0.4641	0.2486	1.0564	H52	0.8933	0.3997	0.7941
H100	0.5160	-0.0550	1.0510	H53A	1.1740	0.4907	0.7866
H2	0.6741	0.3386	0.4153	H53B	1.1148	0.4884	0.8717
H4	0.3295	0.6914	0.2656	H51A	1.0458	0.6226	0.7158
H12A	0.3879	0.2319	0.4404	H51B	0.9866	0.6294	0.8002
H12B	0.4843	0.1597	0.4113	H51C	0.8826	0.5788	0.7073

TABLE 9

[00277] Characteristic powder x-ray diffraction peak positions (degrees $2\theta \pm 0.1$) @ RT for Examples 2a, b, d, c, d, e, f, and g based on a high quality pattern collected

with a diffractometer (CuK α) with a spinning capillary with 2 θ calibrated with a NIST other suitable standard

Exp 2a	Exp 2b	Exp 2c	Exp 2d	Exp 2e	Exp 2f	Exp 2g
5.2	5.5	5.1	5.3	5.4	6.3	6.9
7.9	9.1	6.9	9.6	9.4	9.0	8.7
17.1	12.1	7.4	13.7	11.2	11.7	9.8
17.6	14.0	10.2	14.7	13.7	15.0	10.3
19.6	19.2	18.0	19.5	19.1	17.6	11.8

COMPARATIVE PHARMACOLOGICAL CHARACTERISTICS

- 5 [00278] Assays and data comparing the pharmacological characteristics of Example 1 and compounds found in WO2005021500 (corresponding to US Pat No.7,163,937, assigned to present Applicant) are presented below.

Human peripheral blood mononuclear cell binding ("CCR2 Binding")

- [00279] See e.g. Yoshimura et al., *J. Immunol.* **1990**, 145, 292. The human CCR2
10 binding assay was established with human peripheral blood mononuclear cells (hPBMCs) using 125 I-human MCP-1 as the tracer ligand. hPBMCs were isolated from human leukopak (Biological Specialty Inc.) using a standard protocol with Ficoll-Hypaque (Mediatech Cellgro). Isolated hPBMCs were washed and diluted to 1×10^7 /ml in binding buffer (RPMI-1640, 0.1% BSA, 20 mM Hepes, pH 7.4). 125 I-MCP-1 (NEN/Perk Elmer) was diluted to 0.45 nM in binding buffer. The compound
15 was diluted in binding buffer at 3-fold the final concentrations used in the binding assay. The binding assay was performed using a 96-well filter plate (Millipore). Total 125 I-MCP-1 binding was assessed as follows: to each reaction of a total volume of 150 μ l were added 5×10^5 cells, 0.15 nM 125 I-MCP-1, and compound such that the
20 final concentration ranged from 0 to 100 nM. The plate was incubated at room temperature for 30 minutes followed by three washes with RPMI-1640, 0.1% BSA, 0.4 M NaCl, 20 mM Hepes, pH 7.4 using a vacuum manifold filtration (Millipore). After washing, the plate was air-dried for 60 minutes at room temperature. This was followed by adding 25 μ l of Microscint 20 into each well. The plate was sealed and
25 counted on the Trilux for 1 minute. Non-specific binding was determined in the presence of 300 nM cold MCP-1 (PeproTech Inc.). Specific 125 I-MCP-1 was calculated as the difference between total and non-specific binding. All conditions

were tested in duplicate. The IC₅₀ is defined as the concentration of competing compound required to reduce specific binding by 50%.

hERG Flux

[00280] HEK293 cells stably-expressing hERG channels were grown (37 °C, 5% CO₂) in Dulbecco's Modified Eagle's Media supplemented with 10% Sigma fetal bovine serum, non-essential amino acids, 2mM L-glutamine and 500 µg/ml G418, at incubator. Cell dissociation buffer was used to extract the cells from flasks, which were then plated into 384-well Corning poly-D-lysine coated black/clear plates at a density of 2 x 10⁴ cells per well (20 µl) in 10% serum media, and incubated for 15-24 hours at 37 °C in a 5% CO₂ incubator until a confluent monolayer of cells was obtained.

[00281] A 2 mM stock of BTC-AM dye (Molecular Probes, Eugene, OR) was prepared in 100% DMSO and then added 1:1 to 10% (w/v) pluronic acid in DMSO on the day of assay. The dye was then diluted in hERG external EP buffer (140 mM NaCl, 4.0 mM KCl, 1.8 mM CaCl₂, 1.0 mM MgCl₂, 10 mM HEPES, pH 7.3 and 10 mM glucose; all buffer components obtained from Sigma Chemical). This BTC dye mixture (30 µl) was added to the cells and produced a final loading concentration of 2.5 µM. Cells are incubated at 21 °C for 45 minutes.

[00282] Test compounds were diluted to 10 mM DMSO in 60 µl. These compounds were then serially-diluted at a 1:2 ratio in DMSO in columns 1-10 and 11-20 of a 384-well plate. Assay-ready plates were generated by stamping 2.5 µl from the DMSO serially diluted plate, which was prepared on the Velocity 11 BioCel. Aqueous plates were created by adding 48 µl of EP buffer and then were diluted 30 - 45 minutes before the assay was read on the FLIPR. After dye loading, aqueous-diluted compounds were added to the cells of the three replicate plates (10 µl) yielding a ten point concentration range of 80 µM to 0.156 nM. Final DMSO concentration in the assay is 1%. Assay-ready aqueous plates were prepared and diluted on a Cybio liquid handler.

[00283] Cells loaded with dye were read on the FLIPR384 (Molecular Devices, Sunnyvale, CA), which excites the dye using the 488 nm line of an argon laser. Emission was filtered using a 540 ± 30 nm bandpass filter. hERG channels are stimulated to open by the addition of 20 µl/well EP buffer containing 66 mM K₂SO₄

and 1.3 mM Ti_2SO_4 (Sigma/Aldrich). For each plate, data were collected every second for a period of 12 seconds, at which time the Ti^+ -containing stimulus buffer was added. Data collection proceeded every second for 48 seconds, and then continued every three seconds for an additional 2 minutes.

- 5 [00284] The dynamic range of the assay was determined from *blanks* and *totals* wells. The *totals* wells (columns 21 and 22) define maximal hERG activation for the plate (no test compound present), and the *blanks* wells (columns 23 and 24) define 100% hERG inhibition. The *blanks* wells contain 400 nM of either of the standard hERG inhibitors dofetilide (Ficker *et al.*, 1998) or E-4031. Raw data points in each
10 sample well were first corrected for cell/signal variation, negative control (blanks) background, and normalized to the positive controls (totals) using the online FLIPR software. Test compound concentration response curves for the hERG Ti^+ flux data were then fit using Excel Fit (ID Business Solutions Limited, Surrey, UK) with a single-site logistic equation, $Y = A + ((B-A)/1 + ((C/X)^D))$ where A= maximal
15 inhibition. Data were analyzed by fitting maximum amplitudes of change in fluorescence for Ti^+ flux for a given condition of test compound. Potencies (IC_{50} values) of compounds were calculated from the average of triplicate wells.

Sodium channel, site 2 binding assay

- [00285] See also: W. A. Catterall, et al. *J. Biol. Chem.* **1981**, 256, 8922. The
20 standard binding buffer contained 50 mM HEPES, 50 mM Tris-HCl, pH 7.4, 130 mM Choline Chloride, 5.4 mM KCl, 0.8 mM MgCl_2 , 5.5 mM glucose, 40 $\mu\text{g/mL}$ LqT. Binding reactions were initiated by adding synaptosomes (prepared from Wistar rat brain) to the reaction mixture containing 5 nM [^3H]-Batrachotoxin in a standard binding buffer and the compound to be tested at the desirable concentration. Samples
25 were then mixed and incubated at 37 °C for 60 minutes. The reactions were stopped by adding ice-cold washing buffer containing 50 mM HEPES, 50 mM Tris-HCl, pH 7.4, 1.8 mM CaCl_2 , 0.8mM MgCl_2 and 1 mg/mL bovine serum albumin. The synaptosomes were immediately collected onto glass fiber filters and washed 3 times with washing buffers. The radioactivity of [^3H]-Batrachotoxin remaining on the
0 filters was counted using liquid scintillation spectrometers.

Parallel Artificial Membrane Permeability Assay (PAMPA)

[00286] The Parallel Artificial Membrane Permeability Assay (PAMPA) consists of a specially formulated lecithin-based lipid combination referred to as the gastrointestinal tract (GIT) lipid. The GIT lipid is used to form a membrane in a sandwich plate assembly similar to that used in the Caco-2 assays. The GIT lipid
5 closely resembles *in vivo* membrane composition and performance as measured by standard compounds that are known to be passively absorbed in humans. PAMPA is widely used as an *in vitro* model for permeability screening of discovery compounds. The rate of passage of compounds through the PAMPA membrane is used to determine a permeability coefficient (P_c), which can be related to the *in vivo* passive
10 permeability of the compound.

[00287] The permeability coefficient (P_c) of a particular compound is examined in a pH-dependent setting with apical and basolateral pH of 7.4. All experiments are conducted in triplicate determinations.

[00288] Compounds (10 mM stocks in 100% DMSO) were diluted 1:100 in pH 7.4
15 donor well buffer (pION CAT # 110151), providing a 100 μ M assay solution in 1% DMSO. Compound diluted in donor well buffer was transferred to a Whatman Unifilter plate and filtered prior to dispensing 200 μ l into the donor well of the assay plate (pION CAT #110163). The PAMPA membrane was formed by pipetting 4 μ l of the lipid solution (pION CAT #110169) onto the filter plate (VWR CAT #13503).
20 The membrane was then covered with 200 μ l of acceptor well buffer at pH 7.4 (pION CAT #110139). The PAMPA assay plate (donor side and acceptor side) was combined and allowed to incubate at room temperature for 4 hours. The plate was then disassembled and spectrophotometer plates (VWR CAT #655801) were filled (150 μ l/well). The donor, acceptor, reference, and blank plates were read in the
25 SpectraMax UV plate reader. Data was captured by the pION software, which analyzes the spectra and generates P_c values.

CCR2 Chemotaxis

[00289] The human CCR2 chemotaxis assay was conducted with the human monocytic cell line, THP-1. THP-1 cells were first labeled with the fluorescent dye
30 Calcein-AM in phenol red-free, BSA-free RPMI-1640 (pH 7.4) at 37 °C for 30 minutes with gentle mixing every 15 minutes. The labeled cells were then washed and re-suspended at 1×10^5 /ml in chemotaxis buffer (phenol red-free RPMI-1640, 0.1%

BSA, pH 7.4). The test compound was diluted in chemotaxis buffer such that the final assay concentration ranged from 0.01 nM to 1 μ M. The ligand MCP-1 (PeproTech Inc.) was diluted to 20 nM in chemotaxis buffer. To perform the assay, an equal volume of test compound dilutions was mixed with an equal volume of labeled THP-1 cells (Mixture 1), and an equal volume of test compound dilutions was mixed with an equal volume of diluted MCP-1 ligand (Mixture 2). Both mixtures were incubated independently at 37 °C for 10 minutes followed by gentle mixing. MCP-1-induced chemotaxis was then measured in a chemotaxis plate (Becton Dickinson) by placing 50 μ l of Mixture 1 in the top chamber and 225 μ l of Mixture 2 in the bottom chamber. The plate was covered with a lid and incubated at 37°C for 30 minutes. 30 minutes later, the plate was read on a Cytofluor. All conditions were tested in duplicate. For signal to noise determination, 50 μ l of labeled THP-1 cells alone (5×10^4 /well) were placed into the top chamber and 225 μ l of ligand MCP-1 alone was placed in the bottom chamber (final concentration of 10 nM). The inhibition achieved by graded concentrations of test compound was calculated as a percentage of the compound-free MCP-1 control. The IC₅₀ is defined as the concentration of test compound required to reach 50% inhibition of cellular chemotaxis.

hERG Patch Clamp

[00290] Whole-cell patch-clamp was used to directly measure hERG currents in HEK-293 cells stably expressing the cloned hERG potassium channel α subunit. The compound was tested in an aqueous buffer with pH 7.4 at room temperature. Repetitive test pulses (0.05 Hz) were applied from a holding potential of -80 mV to +20 mV for 2 seconds and tail currents were elicited following the test pulses by stepping the voltage to -65 mV. The effects from the compound were calculated by measuring inhibition of peak tail current

Sodium channel Patch Clamp

[00291] Whole-cell patch-clamp was used to directly measure inward sodium currents in HEK-293 cells expressing the human cardiac sodium channel, SCN5A. The compound was tested at a protein-free aqueous buffer. For determining steady state inhibition, sodium currents were elicited every 5 seconds using the following

voltage protocol: cells were held at a potential of -90 mV and stepped to -20 mV for 60 ms. Effects were calculated by measuring inhibition of peak current during the test pulse to -20 mV. Rate-dependence of inhibition was assessed by stimulation at frequencies of 1 Hz and 4 Hz.

5 **Single-dose Pharmacokinetics in Rats**

[00292] Male Sprague-Dawley rats (250-300 g) were used for the pharmacokinetic studies. Rats were fasted overnight prior to PO dosing and fed 4 h post dose. Blood samples (~0.3 mL) were collected from the jugular vein into K₂EDTA-containing tubes and then centrifuged at 4°C (1500-2000xg) to obtain plasma. In an oral
10 bioavailability study, 2 groups of animals (N=2-3 per group) received the test compound either as an intravenous (IV) infusion (over 10 min) via the jugular vein or by oral gavage. Serial blood samples were obtained at 0.17 (for IV only), 0.25, 0.5, 0.75, 1, 2, 4, 6, 8, and 24 h post dose. Plasma samples, obtained by centrifugation at 4°C (1500-2000xg), were stored at -20°C until analysis by LC/MS/MS.

15 **Single-dose Pharmacokinetics in Monkeys**

[00293] The pharmacokinetics of various test compounds were evaluated in male Cynomolgus monkeys in a crossover-design. Monkeys were fasted overnight prior to PO dosing and fed 4 h post dose. A group of 1-3 animals (3 to 5 kg) received the compound by IV infusion (over 10 min) via a femoral vein and by oral gavage, with a
20 1-week washout between treatments. Serial blood samples (~0.3 mL) were collected from a femoral artery at 0.17 (IV only), 0.25, 0.5, 0.75, 1, 2, 4, 6, 8, and 24 h post dose, and centrifuged at 4°C (1500-2000xg) to obtain plasma. Samples were stored at -20°C until analysis by LC/MS/MS.

Data analysis for pharmacokinetic assays

25 [00294] The pharmacokinetic parameters were obtained by non-compartmental analysis of plasma concentration vs. time data (KINETICA™ software, Version 4.2, InnaPhase Corporation, Philadelphia, PA). The peak concentration (C_{max}) and time for C_{max} were recorded directly from experimental observations. The area under the curve from time zero to the last sampling time (AUC(0-T)) was calculated using a
30 combination of linear and log trapezoidal summations. The total plasma clearance (CL_{TP}), steady-state volume of distribution (V_{ss}), apparent elimination half-life (T_{1/2}) and mean residence time (MRT) were estimated after IV administration.

Estimations of T1/2 was made using a minimum of 3 time points with quantifiable concentrations. The absolute oral bioavailability (F) was estimated as the ratio of dose-normalized AUC values following oral and IV doses.

[00295] Find below data for each compound as measured in the assays described above.

Table 10. Comparative In Vitro Data

Compound	CCR2 Binding IC ₅₀ (nM)	hERG FLUX IC ₅₀ (nM)	Na ⁺ channel binding (% inhibition)	PAMPA permeability (nm/sec)
Example 12as, WO2005021500	0.27 (1)	2,800	Not available	Not available
Example 12aj WO2005021500	0.43 ± 0.06 (2)	770	Not available	Not available
Example 2k WO2005021500	0.88 ± 0.60 (23)	51,000	97%, 10,000 nM	529 ± 157 (9)
Example 12bd WO2005021500	1.15 ± 0.07 (2)	>80,000	54%, 10,000 nM	392
Example 8a WO2005021500	1.83 ± 0.80 (12)	>80,000	3%, 10,000 nM 33%, 30,000 nM	94 ± 58 (10)
Example 8e, WO2005021500	2.20 ± 0.03 (2)	>80,000	6%, 10,000 nM	2 ± 2 (2)
Example 9c, WO2005021500	0.96 ± 0.26 (19)	>80,000	48%, 10,000 nM 75%, 30,000 nM	145 ± 71 (8)
Example 1 Present Invention	2.74 ± 1.34 (15)	>80,000	13%, 10,000 nM 32%, 30,000 nM	560 ± 86 (5)

Table 11a. Additional Comparative In Vitro Data

Compound	CCR2 Chemotaxis IC ₅₀ (nM)	hERG patch clamp (% Inhib.)	Na ⁺ channel patch clamp (% Inhib.)
Example 2k WO2005021500	0.24 ± 0.16 (12)	83%, 10,000 nM	52%, 10,000 nM 90%, 30,000 nM
Example 8a WO2005021500	2.63 ± 1.24 (4)	4%, 10,000 nM	22%, 10,000 nM 49%, 30,000 nM
Example 9c, WO2005021500	0.21	4%, 10,000 nM	19%, 10,000 nM 39%, 30,000 nM
Example 1, Present Invention	0.75 ± 0.42 (16)	12%, 10,000 nM 19%, 30,000 nM	29%, 30,000 nM

Table 11b. Comparative In Vivo Pharmacokinetic Data in the Rat

Compound	Dose IV/PO (mg/kg)	Cl (mL/min/kg)	F%	Oral AUC (nM·h)
Example 2k WO2005021500	2.5 / 25	40	68	9294
Example 8a WO2005021500	6 / 72	42	1.4	690
Example 9c,	4 / 43	54	14	1855

WO2005021500				
Example 1, Present Invention	2 / 10	25	79	10169

Table 11c. Comparative In Vivo Pharmacokinetic Data in the Monkey

Compound	Dose IV/PO (mg/kg)	Cl (mL/min/kg)	F%	Oral AUC (nM*h)
Example 2k WO2005021500	1 / 1.4	25	46	862
Example 8a WO2005021500	1 / 11	14	9.4	1896
Example 9c, WO2005021500	1 / 10	12	26	6763
Example 1, Present Invention	1 / 1	12	95	2352

UTILITY

- 5 [00296] Representative compounds of the examples are shown to be modulators of chemokine receptor activity using assays known by those skilled in the art. In this section, we describe such assays and give their literature reference. More assays are described herein in the section titled "Comparative Pharmacological Characteristics", *supra*. By displaying activity in these assays of MCP-1 antagonism,
- 10 compounds of the examples are expected to be useful in the treatment of human diseases associated with chemokines and their cognate receptors. The definition of activity in these assays is a compound demonstrating an IC₅₀ of 30 μ M or lower in concentration when measured in a particular assay.
- 15 Antagonism of MCP-1 Binding to Human PBMC
(Yoshimura et al., *J. Immunol.* 1990, 145, 292)
- [00297] At least one compound described in the examples has activity in the antagonism of MCP-1 binding to human PBMC (human peripheral blood mononuclear cells) described here.
- 20 [00298] Millipore filter plates (#MABVN1250) are treated with 100 μ l of binding buffer (0.5% bovine serum albumin, 20 mM HEPES buffer and 5 mM magnesium chloride in RPMI 1640 media) for thirty minutes at room temperature. To measure binding, 50 μ l of binding buffer, with or without a known concentration compound, is combined with 50 μ l of ¹²⁵I labeled human MCP-1 (to give a final concentration of

150 pM radioligand) and 50 µl of binding buffer containing 5×10^5 cells. Cells used for such binding assays can include human peripheral blood mononuclear cells isolated by Ficoll-Hypaque gradient centrifugation, human monocytes (Weiner et al., *J. Immunol. Methods*. 1980, 36, 89), or the THP-1 cell line which expresses the endogenous receptor. The mixture of compound, cells and radioligand are incubated at room temperature for thirty minutes. Plates are placed onto a vacuum manifold, vacuum applied, and the plates washed three times with binding buffer containing 0.5M NaCl. The plastic skirt is removed from the plate, the plate allowed to air dry, the wells punched out and counted. The percent inhibition of binding is calculated using the total counts obtained in the absence of any competing compound and the background binding determined by addition of 100 nM MCP-1 in place of the test compound.

Antagonism of MCP-1-induced Calcium Influx

(Sullivan, et al. *Methods Mol. Biol.*, 114, 125-133 (1999))

[00299] At least one compounds described in the examples have activity in the antagonism of MCP-1-induced calcium influx assay described here.

[00300] Calcium mobilization is measured using the fluorescent Ca^{2+} indicator dye, Fluo-3. Cells are incubated at 8×10^5 cells/ml in phosphate-buffered saline containing 0.1% bovine serum albumin, 20 mM HEPES buffer, 5 mM glucose, 1% fetal bovine serum, 4 µM Fluo-3 AM and 2.5 mM probenecid for 60 minutes at 37°C. Cells used for such calcium assays can include human monocytes isolated as described by Weiner et al., *J. Immunol. Methods*, 36, 89-97 (1980) or cell lines which expresses the endogenous CCR2 receptor such as THP-1 and MonoMac-6. The cells are then washed three times in phosphate-buffered saline containing 0.1% bovine serum albumin, 20 mM HEPES, 5 mM glucose and 2.5 mM probenecid. The cells are resuspended in phosphate-buffered saline containing 0.5% bovine serum albumin, 20 mM HEPES and 2.5 mM probenecid at a final concentration of $2-4 \times 10^6$ cells/ml. Cells are plated into 96-well, black-wall microplates (100 µl/well) and the plates centrifuged at 200 x g for 5 minutes. Various concentrations of compound are added to the wells (50 µl/well) and after 5 minutes, 50 µl/well of MCP-1 is added to give a final concentration of 10 nM. Calcium mobilization is detected by using a

fluorescent-imaging plate reader. The cell monolayer is excited with an argon laser (488 nm) and cell-associated fluorescence measured for 3 minutes, (every second for the first 90 seconds and every 10 seconds for the next 90 seconds). Data are generated as arbitrary fluorescence units and the change in fluorescence for each well determined as the maximum-minimum differential. Compound-dependent inhibition is calculated relative to the response of MCP-1 alone.

Antagonism of MCP-1-induced Human PBMC Chemotaxis

(Bacon et al., *Brit. J. Pharmacol.* 1988, 95, 966)

- 10 [00301] At least one compounds described in the examples have activity in the antagonism of MCP-1-induced human PBMC chemotaxis assay described here.
- [00302] Neuroprobe MBA96-96-well chemotaxis chamber, Polyfiltronics MPC 96 well plate, and Neuroprobe polyvinylpyrrolidone-free polycarbonate PFD5 8-micron filters are warmed in a 37 °C incubator. Human Peripheral Blood Mononuclear Cells (PBMCs) (Boyum et al., *Scand. J. Clin. Lab Invest. Suppl.* 1968, 97, 31), freshly isolated via the standard ficoll density separation method, are suspended in DMEM at 1 x 10⁷ c/ml and warmed at 37°C. A 60nM solution of human MCP-1 is also warmed at 37°C. Dilutions of test compounds are made up at 2x the concentration needed in DMEM. The PBMC suspension and the 60nm MCP-1 solution are mixed 1:1 in polypropylene tubes with prewarmed DMEM with or without a dilution of the test compounds. These mixtures are warmed in a 37°C tube warmer. To start the assay, add the MCP-1/compound mixture into the wells of the Polyfiltronics MPC 96 well plate that has been placed into the bottom part of the Neuroprobe chemotaxis chamber. The approximate volume is 400µl to each well and there should be a positive meniscus after dispensing. The 8 micron filter is placed gently on top of the 96 well plate, a rubber gasket is attached to the bottom of the upper chamber, and the chamber is assembled. A 200µl volume of the cell suspension/compound mixture is added to the appropriate wells of the upper chamber. The upper chamber is covered with a plate sealer, and the assembled unit is placed in a 37°C incubator for 45 minutes. After incubation, the plate sealer is removed and all the remaining cell suspension is aspirated off. The chamber is disassembled and the filter gently removed. While holding the filter at a 90 degree angle, unmigrated cells are washed

away using a gentle stream of phosphate buffered saline and the top of the filter
wiped with the tip of a rubber squeegee. Repeat this wash twice more. The filter is
air dried and then immersed completely in Wright Geimsa stain for 45 seconds. The
filter is then washed by soaking in distilled water for 7 minutes, and then a 15 second
5 additional wash in fresh distilled water. The filter is again air dried. Migrated cells
on the filter are quantified by visual microscopy.

[00303] Mammalian chemokine receptors provide a target for interfering with or
promoting immune cell function in a mammal, such as a human. Compounds that
inhibit or promote chemokine receptor function are particularly useful for modulating
10 immune cell function for therapeutic purposes. Accordingly, the present invention is
directed to compounds which are useful in the prevention and/or treatment of a wide
variety of inflammatory, infectious, and immunoregulatory disorders and diseases,
including asthma and allergic diseases, infection by pathogenic microbes (which, by
definition, includes viruses), as well as autoimmune pathologies such as the
15 rheumatoid arthritis and atherosclerosis.

[00304] For example, an instant compound which inhibits one or more functions of
a mammalian chemokine receptor (e.g., a human chemokine receptor) may be
administered to inhibit (i.e., reduce or prevent) inflammation or infectious disease.
As a result, one or more inflammatory process, such as leukocyte emigration,
20 adhesion, chemotaxis, exocytosis (e.g., of enzymes, histamine) or inflammatory
mediator release, is inhibited.

[00305] Similarly, an instant compound which promotes one or more functions of
the mammalian chemokine receptor (e.g., a human chemokine) as administered to
stimulate (induce or enhance) an immune or inflammatory response, such as
15 leukocyte emigration, adhesion, chemotaxis, exocytosis (e.g., of enzymes, histamine)
or inflammatory mediator release, resulting in the beneficial stimulation of
inflammatory processes. For example, eosinophils can be recruited to combat
parasitic infections. In addition, treatment of the aforementioned inflammatory,
allergic and autoimmune diseases can also be contemplated for an instant compound
0 which promotes one or more functions of the mammalian chemokine receptor if one
contemplates the delivery of sufficient compound to cause the loss of receptor
expression on cells through the induction of chemokine receptor internalization or the

delivery of compound in a manner that results in the misdirection of the migration of cells.

[00306] In addition to primates, such as humans, a variety of other mammals can be treated according to the method of the present invention. For instance, mammals, including but not limited to, cows, sheep, goats, horses, dogs, cats, guinea pigs, rats or other bovine, ovine, equine, canine, feline, rodent or murine species can be treated. However, the method can also be practiced in other species, such as avian species. The subject treated in the methods above is a mammal, male or female, in whom modulation of chemokine receptor activity is desired. "Modulation" as used herein is intended to encompass antagonism, agonism, partial antagonism and/or partial agonism.

CCR5 Binding and Functional Assays

[00307] Cell derivation and cell culture: A pool of HT1080 cells stably expressing endogenous CC chemokine receptor 5 (CCR5) were developed using the methods outlined by Harrington, Sherf, and Rundlett (see United States patents US 6,361,972 and US 6,410,266). The highest-expressing clones were isolated using repetitive flow cytometry, followed by sub-cloning. These cells were then cultured in 6-well dishes at 3×10^5 cells/well and transfected with a DNA vector containing the chimeric HA-tagged G protein Gqi5 (Molecular Devices; 5 micrograms of linearized vector DNA in 15 microL of Ex-Gen from Fermentes was used for the transfection). Two days after transfection, the wells were combined and plated into P100 plates. Seven days after plating, colonies were picked, expanded, and analyzed for Gqi5 content by Western blot. A clone (designated as 3559.1.6) having high expression of Gqi5 (from transfection) and of CCR5 (endogenous) was selected and used for the experiments described below. The HT1080 cells (clone 3559.1.6) were cultured with alpha-MEM supplemented with 10% dialyzed fetal bovine serum, 2% penicillin/streptomycin/glutamine, and 500 microgram/mL hygromycin B (final concentration) at 37° C with 5% CO₂ in a humidified atmosphere.

[00308] Membrane Preparation: A cell pellet containing 1×10^8 HT1080 cells (clone 3559.1.6) was resuspended in 5 mL of ice-cold Membrane Prep Buffer (50 mM HEPES, 5 mM MgCl₂, 1 mM CaCl₂) and homogenized at high-speed on a Polytron homogenizer for 20 sec on ice. The homogenate was diluted with another

25 mL of Membrane Prep Buffer and centrifuged for 12 min (48,000 x g at 4°C). The cell pellet was resuspended in 5 mL of Membrane Prep Buffer before being rehomogenized as described previously. The homogenate was diluted with 5 mL of Membrane Prep Buffer and assayed for CCR5 protein concentration.

- 5 **[00309]** Binding assay: The freshly-prepared homogenate from the Membrane Preparation described above was diluted in Binding buffer (50 mM HEPES, 5 mM MgCl₂, 1 mM CaCl₂, 0.1% BSA; one complete protease inhibitor tablet was added before assay) to achieve a final protein concentration of 10 micrograms/well (solid white 96-well plates from Corning, Inc.). This membrane preparation was mixed
- 10 with WGA-SPA beads (Amerhsam; pre-soaked in Binding buffer) to give a concentration of 200 micrograms/well. The membrane/SPA bead mix (100 microliters/well) was then added to a plate that had been pre-dotted with 2 microliters DMSO containing various concentrations of test articles (pure DMSO for negative control; various concentrations of examples of this invention for test articles; 500 nM
- 15 MIP-1 beta as a positive control). The binding assay was initiated through the addition of 50 microliters of [¹²⁵I]-MIP-1 beta (Perkin Elmer; material was diluted in Binding buffer such that the addition of 50 microliters/well gives a final concentration of 0.1 nM [¹²⁵I]-MIP-1 beta). The plate was sealed and allowed to stand at room temperature for 4 – 6 h before being counted on a Packard TopCount. The percentage
- 20 bound for the test article was calculated, using negative and positive controls to define the window for each experiment.

- [00310]** Fluorometric Imaging Plate Reader (FLIPR)-based Functional assay: HT1080 cells (clone 3559.1.6) were plated at 10,000 cells/well (30 microliters) in 384-well plates (black/clear bottom Biocoat PDL, Beckton Dickinson) and charged
- 25 with 30 microliters/well of Fluro-4 AM fluorescent dye (prepared by dissolving 1 mg Fluro-4 AM in 440 microliters DMSO and diluting with 100 microliters of pluronic solution before diluting further with 10 mL of Hanks buffer). The cells were incubated at 37° C with 5% CO₂ for 30 min before being washed three times and suspended in Assay Buffer (20 mM HEPES, 1.2 mM CaCl₂, 5 mM MgCl₂, 2.5 mM
- 30 Probenecid, 0.5% BSA, 1x Hanks). The test article was serially diluted in DMSO and then diluted 1:10 with Assay Buffer before being added to the cells (10 microliters/well). Using FLIPR, the plates were read (10 – 70 sec) for induction of

flux (i.e. agonist activity). The cells were then further charged with Agonist Solution (30 microliters/well; prepared by diluting 30 microliters of 100 microMolar MIP-1 beta in 100 mL of Assay Buffer; this protocol delivers a final concentration of 5 nM MIP-1 beta in the assay) and the plates were read using FLIPR for one minute.

- 5 Antagonist activity of the test article was determined relative to 0.4% DMSO/Buffer negative control.

[00311] At least one compound of the disclosure is an inhibitor of both CCR2 and CCR5 and may be used to treat diseases associated with either chemokine. The compounds of the present invention are considered dual antagonists.

- 10 [00312] Diseases or conditions of human or other species which can be treated with inhibitors of chemokine receptor function, include, but are not limited to: inflammatory or allergic diseases and conditions, including respiratory allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, hypersensitivity pneumonitis, eosinophilic cellulitis (e.g., Well's syndrome),
15 eosinophilic pneumonias (e.g., Loeffler's syndrome, chronic eosinophilic pneumonia), eosinophilic fasciitis (e.g., Shulman's syndrome), delayed-type hypersensitivity, interstitial lung diseases (ILD) (e.g., idiopathic pulmonary fibrosis, or ILD associated with rheumatoid arthritis, systemic lupus erythematosus, ankylosing spondylitis, systemic sclerosis, Sjogren's syndrome, polymyositis or
20 dermatomyositis); systemic anaphylaxis or hypersensitivity responses, drug allergies (e.g., to penicillin, cephalosporins), eosinophilia-myalgia syndrome due to the ingestion of contaminated tryptophan, insect sting allergies; autoimmune diseases, such as rheumatoid arthritis, psoriatic arthritis, multiple sclerosis, systemic lupus erythematosus, myasthenia gravis, juvenile onset diabetes; glomerulonephritis,
25 autoimmune thyroiditis, Behcet's disease; graft rejection (e.g., in transplantation), including allograft rejection or graft-versus-host disease; inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis; spondyloarthropathies; scleroderma; psoriasis (including T-cell mediated psoriasis) and inflammatory dermatoses such as an dermatitis, eczema, atopic dermatitis, allergic contact
30 dermatitis, urticaria; vasculitis (e.g., necrotizing, cutaneous, and hypersensitivity vasculitis); eosinophilic myositis, eosinophilic fasciitis; cancers with leukocyte infiltration of the skin or organs. Other diseases or conditions in which undesirable

inflammatory responses are to be inhibited can be treated, including, but not limited to, vasculitis, vulnerable plaques, venous neointimal hyperplasia reperfusion injury, dialysis-graft neointimal hyperplasia, artio-venous shunt intimal hyperplasia, atherosclerosis, certain hematologic malignancies, cytokine-induced toxicity (e.g.,
5 septic shock, endotoxic shock), polymyositis, dermatomyositis. Infectious diseases or conditions of human or other species which can be treated with inhibitors of chemokine receptor function, include, but are not limited to, HIV.

[00313] Diseases or conditions of humans or other species which can be treated with promoters of chemokine receptor function, include, but are not limited to:
10 immunosuppression, such as that in individuals with immunodeficiency syndromes such as AIDS or other viral infections, individuals undergoing radiation therapy, chemotherapy, therapy for autoimmune disease or drug therapy (e.g., corticosteroid therapy), which causes immunosuppression; immunosuppression due to congenital deficiency in receptor function or other causes; and infections diseases, such as
15 parasitic diseases, including, but not limited to helminth infections, such as nematodes (round worms); (Trichuriasis, Enterobiasis, Ascariasis, Hookworm, Strongyloidiasis, Trichinosis, filariasis); trematodes (flukes) (Schistosomiasis, Clonorchiasis), cestodes (tape worms) (Echinococcosis, Taeniasis saginata, Cysticercosis); visceral worms, visceral larva migraines (e.g., Toxocara), eosinophilic
20 gastroenteritis (e.g., Anisaki sp., Phocanema sp.), cutaneous larva migraines (Ancylostoma braziliense, Ancylostoma caninum). The compounds of the present invention are accordingly useful in the prevention and treatment of a wide variety of inflammatory, infectious and immunoregulatory disorders and diseases.

[00314] In addition, treatment of the aforementioned inflammatory, allergic and
5 autoimmune diseases can also be contemplated for promoters of chemokine receptor function if one contemplates the delivery of sufficient compound to cause the loss of receptor expression on cells through the induction of chemokine receptor internalization or delivery of compound in a manner that results in the misdirection of the migration of cells.

0 [00315] In another aspect, the instant invention may be used to evaluate the putative specific agonists or antagonists of a G protein coupled receptor. The present invention is directed to the use of these compounds in the preparation and execution

of screening assays for compounds that modulate the activity of chemokine receptors. Furthermore, the compounds of this invention are useful in establishing or determining the binding site of other compounds to chemokine receptors, e.g., by competitive inhibition or as a reference in an assay to compare its known activity to a compound with an unknown activity. When developing new assays or protocols, compounds according to the present invention could be used to test their effectiveness. Specifically, such compounds may be provided in a commercial kit, for example, for use in pharmaceutical research involving the aforementioned diseases. The compounds of the instant invention are also useful for the evaluation of putative specific modulators of the chemokine receptors. In addition, one could utilize compounds of this invention to examine the specificity of G protein coupled receptors that are not thought to be chemokine receptors, either by serving as examples of compounds which do not bind or as structural variants of compounds active on these receptors which may help define specific sites of interaction.

15 **[00316]** Compounds disclosed herein are useful to treat or prevent disorders selected from rheumatoid arthritis, osteoarthritis, septic shock, atherosclerosis, aneurism, fever, cardiovascular effects, haemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, malaria, Crohn's disease, inflammatory bowel diseases, mycobacterial infection, meningitis, psoriasis, congestive heart failure, fibrotic

20 diseases, cachexia, graft rejection, autoimmune diseases, skin inflammatory diseases, multiple sclerosis, radiation damage, hyperoxic alveolar injury, HIV, HIV dementia, non-insulin dependent diabetes mellitus, asthma, allergic rhinitis, atopic dermatitis, idiopathic pulmonary fibrosis, bullous pemphigoid, helminthic parasitic infections, allergic colitis, eczema, conjunctivitis, transplantation, familial eosinophilia,

25 eosinophilic cellulitis, eosinophilic pneumonias, eosinophilic fasciitis, eosinophilic gastroenteritis, drug induced eosinophilia, cystic fibrosis, Churg-Strauss syndrome, lymphoma, Hodgkin's disease, colonic carcinoma, Felty's syndrome, sarcoidosis, uveitis, Alzheimer, Glomerulonephritis, and systemic lupus erythematosus, esophageal squamous cell carcinoma, neuropathic pain, and obesity.

30 **[00317]** In another aspect, the compounds are useful to treat or prevent inflammatory disorders selected from rheumatoid arthritis, osteoarthritis, atherosclerosis, aneurism, fever, cardiovascular effects, Crohn's disease,

inflammatory bowel diseases, psoriasis, congestive heart failure, multiple sclerosis, autoimmune diseases, skin inflammatory diseases.

[00318] In another aspect, the compounds are used to treat or prevent inflammatory disorders selected from rheumatoid arthritis, osteoarthritis,

5 atherosclerosis, Crohn's disease, inflammatory bowel diseases, and multiple sclerosis.

[00319] In another aspect, examples disclosed herein may be useful in for the treatment of a variety of cancers, including, but not limited to, the following:

carcinoma including that of the bladder (including accelerated and metastatic bladder cancer), breast, colon (including colorectal cancer), kidney, liver, lung

10 (including small and non-small cell lung cancer and lung adenocarcinoma), ovary, prostate, testes, genitourinary tract, lymphatic system, rectum, larynx, pancreas (including exocrine pancreatic carcinoma), esophagus, stomach, gall bladder, cervix, thyroid, and skin (including squamous cell carcinoma);

hematopoietic tumors of lymphoid lineage including leukemia, acute
15 lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma, histiocytic lymphoma, and Burkett's lymphoma;

hematopoietic tumors of myeloid lineage including acute and chronic myelogenous leukemias, myelodysplastic syndrome, myeloid leukemia, and
20 promyelocytic leukemia;

tumors of the central and peripheral nervous system including astrocytoma, neuroblastoma, glioma, and schwannomas;

tumors of mesenchymal origin including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and

25 other tumors including melanoma, xenoderma pigmentosum, keratoactanthoma, seminoma, thyroid follicular cancer, and teratocarcinoma.

[00320] In another embodiment, disclosed herein are methods of treating cancer, wherein the cancer is selected from breast cancer, liver cancer, prostate cancer, and melanoma. Additionally, compounds disclosed herein may be useful in the treatment
30 of ovarian cancer, and multiple myeloma.

[00321] The present invention provides methods for the treatment of a variety of non-cancerous proliferative diseases.

[00322] Combined therapy to prevent and treat inflammatory, infectious and immunoregulatory disorders and diseases, including asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis and atherosclerosis, and those pathologies noted above is illustrated by the combination of the compounds of this invention and other compounds which are known for such utilities. For example, in the treatment or prevention of inflammation, the present compounds may be used in conjunction with an anti-inflammatory or analgesic agent such as an opiate agonist, a lipoxygenase inhibitor, a cyclooxygenase-2 inhibitor, an interleukin inhibitor, such as an interleukin-1 inhibitor, a tumor necrosis factor inhibitor, an NMDA antagonist, an inhibitor of nitric oxide or an inhibitor of the synthesis of nitric oxide, a non-steroidal anti-inflammatory agent, a phosphodiesterase inhibitor, or a cytokine-suppressing anti-inflammatory agent, for example with a compound such as acetaminophen, aspirin, codeine, fentanyl, ibuprofen, indomethacin, ketorolac, morphine, naproxen, phenacetin, piroxicam, a steroidal analgesic, sufentanyl, sunlindac, interferon alpha and the like. Similarly, the instant compounds may be administered with a pain reliever; a potentiator such as caffeine, an H2-antagonist, simethicone, aluminum or magnesium hydroxide; a decongestant such as phenylephrine, phenylpropanolamine, pseudophedrine, oxymetazoline, ephinephrine, naphazoline, xylometazoline, propylhexedrine, or levodesoxy-ephedrine; and antitussive such as codeine, hydrocodone, caramiphen, carbetapentane, or dextromethorphan; a diuretic; and a sedating or non-sedating antihistamine. Likewise, compounds disclosed herein may be used in combination with other drugs that are used in the treatment/prevention/suppression or amelioration of the diseases or conditions for which compound of the present invention are useful. Such other drugs may be administered, by a route and in an amount commonly used therefore, contemporaneously or sequentially with a compound of the present invention. When a compound is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound of the present invention may be used. Accordingly, the pharmaceutical compositions include those that also contain one or more other active ingredients, in addition to a compound of the present disclosure.

[00323] Examples of other active ingredients that may be combined with a compound of the present invention, either administered separately or in the same pharmaceutical compositions, include, but are not limited to: (a) integrin antagonists such as those for selectins, ICAMs and VLA-4; (b) steroids such as beclomethasone, methylprednisolone, betamethasone, prednisone, dexamethasone, and hydrocortisone; (c) immunosuppressants such as cyclosporin, tacrolimus, rapamycin and other FK-506 type immunosuppressants; (d) antihistamines (H1-histamine antagonists) such as bromopheniramine, chlorpheniramine, dexchlorpheniramine, triprolidine, clemastine, diphenhydramine, diphenylpyraline, tripeleminamine, hydroxyzine, methdilazine, promethazine, trimetopazine, azatadine, cyproheptadine, antazoline, pheniramine pyrilamine, astemizole, terfenadine, loratadine, cetirizine, fexofenadine, descarboethoxyloratadine, and the like; (e) non-steroidal anti-asthmatics such as β_2 -agonists (terbutaline, metaproterenol, fenoterol, isoetharine, albuterol, bitolterol, and pirbuterol), theophylline, cromolyn sodium, atropine, ipratropium bromide, leukotriene antagonists (zafirlukast, montelukast, pranlukast, iralukast, pobelukast, SKB-102,203), leukotriene biosynthesis inhibitors (zileuton, BAY-1005); (f) non-steroidal antiinflammatory agents (NSAIDs) such as propionic acid derivatives (alminoprofen, benxaprofen, bucloxic acid, carprofen, fenbufen, fenoprofen, fluprofen, flurbiprofen, ibuprofen, indoprofen, ketoprofen, miroprofen, naproxen, oxaprozin, piroprofen, pranoprofen, suprofen, tiaprofenic acid, and tioxaprofen), acetic acid derivatives (indomethacin, acemetacin, alclofenac, clidanac, diclofenac, fenclofenac, fenclozic acid, fentiazac, furofenac, ibufenac, isoxepac, oxpinac, sulindac, tiopinac, tolmetin, zidometacin, and zomepirac), fenamic acid derivatives (flufenamic acid, meclofenamic acid, mefenamic acid, niflumic acid and tolfenamic acid), biphenylcarboxylic acid derivatives (diflunisal and flufenisal), oxicams (isoxicam, piroxicam, sudoxicam and tenoxicam), salicylates (acetyl salicylic acid, sulfasalazine) and the pyrazolones (apazone, bezpiperylon, feprazone, mofebutazone, oxyphenbutazone, phenylbutazone); (g) cyclooxygenase-2 (COX-2) inhibitors; (h) inhibitors of phosphodiesterase type IV (PDE-IV); (i) other antagonists of the chemokine receptors; (j) cholesterol lowering agents such as HMG-CoA reductase inhibitors (lovastatin, simvastatin and pravastatin, fluvastatin, atorvastatin, and other statins), sequestrants (cholestyramine and colestipol), nicotonic acid, fenofibric acid

derivatives (gemfibrozil, clofibrat, fenofibrate and benzaifibrate), and probucol; (k) anti-diabetic agents such as insulin, sulfonylureas, biguanides (metformin), α -glucosidase inhibitors (acarbose) and glitazones (troglitazone and pioglitazone); (l) preparations of interferons (interferon α -2a, interferon-2B, interferon α -N3, 5 interferon beta-1a, interferon beta-1b, interferon gamma-1b); (m) antiviral compounds such as efavirenz, nevirapine, indinavir, ganciclovir, lamivudine, famciclovir, and zalcitabine; (o) other compound such as 5-aminosalicylic acid and prodrugs thereof, antimetabolites such as azathioprine and 6-mercaptopurine, and cytotoxic cancer chemotherapeutic agents. The weight ratio of the compound of the 10 present invention to the second active ingredient may be varied and will depend upon the effective doses of each ingredient.

[00324] Generally, an effective dose of each will be used. Thus, for example, when a compound is combined with an NSAID the weight ratio of the compound of the present invention to the NSAID will generally range from about 1000:1 to about 15 1:1000, or alternatively from about 200:1 to about 1:200. Combinations of a compound of the present invention and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

[00325] In treating cancer, a combination of chemotherapeutic agents and/or other 20 treatments (e.g., radiation therapy) is often advantageous. The second (or third) agent may have the same or different mechanism of action than the primary therapeutic agent. It may be especially useful to employ cytotoxic drug combinations wherein the two or more drugs being administered act in different manners or in different phases of the cell cycle, and/or where the two or more drugs have overlapping 25 toxicities or side effects, and/or where the drugs being combined each has a demonstrated efficacy in treating the particular disease state manifested by the patient.

[00326] Accordingly, compounds disclosed herein (or other formulae disclosed herein) may be administered in combination with other anti-cancer and cytotoxic 30 agents and treatments useful in the treatment of cancer or other proliferative diseases. The invention herein further comprises use of the compounds herein (or other formulae disclosed herein), in preparing medicaments for the treatment of cancer,

and/or it comprises the packaging of the compounds of herein together with instructions that the compounds be used in combination with other anti-cancer or cytotoxic agents and treatments for the treatment of cancer. The present invention further comprises combinations of the compounds of and one or more additional
5 agents in kit form, e.g., where they are packaged together or placed in separate packages to be sold together as a kit, or where they are packaged to be formulated together.

[00327] The second (or more) anti-cancer agents may be selected from any one or more of the following:

10 alkylating agents (including nitrogen mustards, alkyl sulfonates, nitrosoureas, ethylenimine derivatives, and triazenes); anti-angiogenics (including matrix metalloproteinase inhibitors); antimetabolites (including adenosine deaminase inhibitors, folic acid antagonists, purine analogues, and pyrimidine analogues); antibiotics or antibodies (including monoclonal antibodies, CTLA-4 antibodies,
15 anthracyclines); aromatase inhibitors;

cell-cycle response modifiers; enzymes; farnesyl-protein transferase inhibitors;

hormonal and antihormonal agents and steroids (including synthetic analogs, glucocorticoids, estrogens/anti-estrogens [e.g., SERMs], androgens/anti-androgens,
20 progestins, progesterone receptor agonists, and luteinizing hormone-releasing [LHRH] agonists and antagonists); insulin-like growth factor (IGF)/insulin-like growth factor receptor (IGFR) system modulators (including IGFR1 inhibitors); integrin-signaling inhibitors; kinase inhibitors (including multi-kinase inhibitors and/or inhibitors of Src kinase or Src/abl, cyclin dependent kinase [CDK] inhibitors,
25 panHer, Her-1 and Her-2 antibodies, VEGF inhibitors, including anti-VEGF antibodies, EGFR inhibitors, mitogen-activated protein [MAP] inhibitors, MEK inhibitors, Aurora kinase inhibitors, PDGF inhibitors, and other tyrosine kinase inhibitors or serine /threonine kinase inhibitors;

microtubule-disruptor agents, such as ecteinascidins or their analogs and
0 derivatives; microtubule-stabilizing agents such as taxanes, and the naturally-occurring epothilones and their synthetic and semi-synthetic analogs;

microtubule-binding, destabilizing agents (including vinca alkaloids); and

topoisomerase inhibitors; prenyl-protein transferase inhibitors; platinum coordination complexes; signal transduction inhibitors; and other agents used as anti-cancer and cytotoxic agents such as biological response modifiers, growth factors, and immune modulators.

5

[00328] Additionally, the compounds of the present invention can be formulated or co-administered with other therapeutic agents that are selected for their particular usefulness in addressing side effects associated with the aforementioned conditions. For example, compounds of the invention may be formulated with agents to prevent
10 nausea, hypersensitivity and gastric irritation, such as antiemetics, and H₁ and H₂ antihistaminics.

The above other therapeutic agents, when employed in combination with the compounds of the present invention, can be used, for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as otherwise determined by one
15 of ordinary skill in the art.

[00329] The compounds are administered to a mammal in a therapeutically effective amount. By "therapeutically effective amount" it is meant an amount of a compound of the present disclosure that, when administered alone or in combination with an additional therapeutic agent to a mammal, is effective to prevent or
20 ameliorate the disease condition or the progression of the disease.

DOSAGE AND FORMULATION

[00330] The compounds of this disclosure can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release
25 formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. They may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the
30 basis of the chosen route of administration and standard pharmaceutical practice.

[00331] The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic

characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. A physician or veterinarian can determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the disorder.

[00332] By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to 1000 mg/kg of body weight, or between about 0.01 to 100 mg/kg of body weight per day, or alternatively, between about 1.0 to 20 mg/kg/day. Intravenously, the doses will range from about 1 to about 10 mg/kg/minute during a constant rate infusion. Compounds of this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily. In one embodiment, the daily oral dosage of the active ingredient is between 3 and 600 mg either administered once daily or in divided doses administered twice daily. Alternatively, the active ingredient may be administered in doses of 10-20 mg administered twice daily or 40 to 100 mg administered once daily. Alternatively, the active ingredient may be administered a dose of 12.5 mg twice a day or 75 mg once a day. Alternatively, the active ingredient may be administered in doses of 3, 10, 30, 100, 300, and 600 mg administered either once or twice a day.

[00333] Compounds of this invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal skin patches. When administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

[00334] The compounds are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as pharmaceutical carriers) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

[00335] For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

[00336] The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

[00337] Compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic acid and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

[00338] Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 100 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will

ordinarily be present in an amount of about 0.5-95% by weight based on the total weight of the composition.

[00339] Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

[00340] Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

[00341] In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration may contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

[00342] Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, a standard reference text in this field.

[00343] Representative useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as follows:

Capsules

[00344] A large number of unit capsules can be prepared by filling standard two-piece hard gelatin capsules each with 100 milligrams of powdered active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.

Soft Gelatin Capsules

[00345] A mixture of active ingredient in a digestable oil such as soybean oil, cottonseed oil or olive oil may be prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100
5 milligrams of the active ingredient. The capsules should be washed and dried.

Tablets

[00346] Tablets may be prepared by conventional procedures so that the dosage unit is 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide,
10 5 milligrams of magnesium stearate, 275 milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

Injectable

15 [00347] A parenteral composition suitable for administration by injection may be prepared by stirring 1.5% by weight of active ingredient in 10% by volume propylene glycol and water. The solution should be made isotonic with sodium chloride and sterilized.

20 Suspension

[00348] An aqueous suspension can be prepared for oral administration so that each 5 mL contain 100 mg of finely divided active ingredient, 200 mg of sodium carboxymethyl cellulose, 5 mg of sodium benzoate, 1.0 g of sorbitol solution, U.S.P., and 0.025 mL of vanillin.

25

[00349] Where the compounds of this invention are combined with other anticoagulant agents, for example, a daily dosage may be about 0.1 to 100 milligrams of the compound of Formula I and about 1 to 7.5 milligrams of the second anticoagulant, per kilogram of patient body weight. For a tablet dosage form, the
30 compounds of this invention generally may be present in an amount of about 5 to 10 milligrams per dosage unit, and the second anti-coagulant in an amount of about 1 to 5 milligrams per dosage unit.

[00350] Where two or more of the foregoing second therapeutic agents are administered with the compound of the examples, generally the amount of each component in a typical daily dosage and typical dosage form may be reduced relative to the usual dosage of the agent when administered alone, in view of the additive or synergistic effect of the therapeutic agents when administered in combination.

[00351] Particularly when provided as a single dosage unit, the potential exists for a chemical interaction between the combined active ingredients. For this reason, when the compound of the examples and a second therapeutic agent are combined in a single dosage unit they are formulated such that although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized (that is, reduced). For example, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. One of the active ingredients may also be coated with a material which effects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product in which the one component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a low viscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component.

[00352] These as well as other ways of minimizing contact between the components of combination products of the present invention, whether administered in a single dosage form or administered in separate forms but at the same time by the same manner, will be readily apparent to those skilled in the art, once armed with the present disclosure.

5 [00353] Additionally, certain compounds disclosed herein may be useful as metabolites of other compounds. Therefore, in one embodiment, compounds may be useful either as a substantially pure compound, which may also then be incorporated into a pharmaceutical composition, or may be useful as metabolite which is generated after administration of the prodrug of that compound. In one embodiment, a compound may be useful as a metabolite by being useful for treating disorders as described herein.

10 [00354] "Substantially pure" as used herein is intended to include a compound having a purity greater than about 90 weight percent, including about 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, and 100 percent.

15 [00355] As one example, a compound disclosed herein may be substantially pure in having a purity greater than about 90 percent (by weight), where the remaining less than about 10 percent of material comprises other metabolite of the compound, a prodrug of the compound, and/or reaction and/or processing impurities arising from its preparation.

[00356] Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise that as specifically described herein.

20

IN VIVO ASSAYS AND EFFICACY

[00357] N-((1R,2S,5R)-5-(*tert*-butylamino)-2-((S)-2-oxo-3-(6-(trifluoromethyl)quinazolin-4-ylamino)pyrrolidin-1-yl)cyclohexyl)acetamide (also referred to as "Example 1") was evaluated in the following in vivo assays as described below.

25

Section 1. Example 1 blocked mononuclear cell recruitment to the skin following intradermal (ID) MCP-1 challenge in cynomolgus monkey.

30 [00358] Intradermal injection of MCP-1 results in the infiltration of mononuclear cells to the injection site. This model was initially developed to assess the inhibitory effect of CCR2 antagonists on the infiltration of mononuclear cells to the skin tissue

injected with human MCP-1. The cellular infiltrate can be measured semi-quantitatively by histological scoring.

Methods

5 [00359] Each monkey was dosed with Example 1 or its vehicle control (0.05 N HCl) once daily for three days. Example 1 was orally administered at doses of 0, 5, 10, or 20 mg/kg to groups of 4 cynomolgus monkeys (2 per sex per group). Immediately after dosing on Day 3, all animals received 2 intradermal injections of 10 µg (50 µL/injection) of human MCP-1 (R & D Systems) and 2 intradermal
10 injections of its DPBS control (50 µl/injection) at separate sites on the dorsal thorax. Dermal biopsies of all sites were obtained at approximately 18 hours following MCP-1 (or DPBS) challenge. Biopsies were processed for semi-quantitative histological evaluation. Representative sections of skin samples were examined by light microscopy; microscopic lesions and cellular infiltration were noted and their
15 incidences were tabulated.

[00360] In addition to biopsy analysis, blood was collected and evaluated for complete blood counts and cell differentials. Also evaluated were plasma samples for compound (and metabolite) concentrations, and serum samples for systemic MCP-1 levels.

20

Results

[00361] The recruitment of mononuclear cells to the skin of vehicle-treated control animals in response to MCP-1 challenge was significant (mean histologic score of 2.0 with a range of 1-3, Table 10). Example 1 at 5, 10, and 20 mg/kg inhibited this
25 dermal mononuclear cell infiltration by 75%, 95% and 95%, respectively (Table 10 and Figure 20). The compound also blocked the infiltration of other cell types such as eosinophils and neutrophils (Table 12). The plasma concentrations of Example 1 at 18 hours and their relationship to levels of inhibition and Cyno chemotaxis IC90 values are summarized in Table 12. Based on the IC50 value of 7.1 ± 2.7 nM for
30 Example 1 in cyno chemotaxis assay, the 5, 10 and 20 mg/kg doses resulted in free plasma concentrations of 0.8-, 2.1-, and 4.3-fold the chemotaxis IC90 at 18 hours post dosing (Table 12).

TABLE 12

Summary of effects of Example 1 on infiltration of mononuclear cells and other cell types in response to MCP-1 challenge in cynomolgus monkeys ^{a,b}

Doses mg/kg	Free plasma concentration (nM)	Fold Chemotaxis IC90	Mononuclear cell score (range) (inhibition%)	PMN ^c score (range)	Eos ^d score (range)	Total cell score (range)
0	0	0	2.0 (1 - 3) (0%)	0.3 (0 - 1)	1.5 (0.5 - 3)	4.0 (2 - 6 ^e)
5	99	0.8	0.5 (0 - 1) (75%)	0.1 (0 - 0.5)	1.4 (1 - 2)	2.0 (1 - 3)
10	248	2.1	0.1 (0 - 0.5) (95%)	0.1 (0 - 0.5)	0.8 (0.5 - 1)	1.0 (0.5 - 2)
20	512	4.3	0.1 (0 - 0.5) (95%)	0 (0 - 0)	0.8 (0.5 - 1)	0.9 (0.5 - 1)

5 ^a An arbitrary scaling system from 0 to 4 was utilized with each number representing a particular designation of inflammatory infiltrate as follows: 0, unremarkable number of inflammatory cells; 0.5, trace; 1, minimal; 2, mild; 3, moderate; 4, marked infiltration.

^b Mean values are an average of 8 MCP-1 site biopsies, representing 2 separate
10 biopsies from 4 monkeys per group. Ranges represent the spread of average histological scores for 2 biopsies per animal.

^c PMN stands for polymorphonuclear cell (neutrophil).

^d Eos is an abbreviation for eosinophils.

^e Total score is a mathematical sum of mean values for each cell type to demonstrate
15 the dose response.

[00362] Evaluation of changes in serum inflammatory mediators showed an increase (approximately 3-4 fold) in MCP-1 level in Example 1-treated groups relative to vehicle control. In addition, complete blood count (CBC) analysis showed

an increase (~2-fold) in neutrophils in Example 1-treated groups, relative to vehicle control, at 18 hours on Day 4 following three days of dosing.

[00363] To refine the dose (concentration) response of Example 1 in a more readily quantifiable system, we used hCCR2 KI mice to evaluate the effect of

- 5 Example 1 monocyte/macrophage infiltration in thioglycollate (TG)-induced peritonitis model with flow cytometry-based methodology.

Section 2. Example 1 inhibited monocyte/macrophage infiltration in 48-hour TG peritonitis model in hCCR2 KI mouse

- 10 [00364] TG-induced peritonitis model has been used as a model of recruitment of monocytes/macrophages to inflammation site. Both in-house and published studies have demonstrated that monocyte/macrophage recruitment in this model is CCR2-dependent. See Boring L. et al., *Impaired monocyte migration and reduced type 1 (Th1) cytokine responses in C-C chemokine receptor 2 knockout mice*. J Clin Invest., 15 100(10):2552-61. (1997); and Kuziel, W.A. et al., *Severe reduction in leukocyte adhesion and monocyte extravasation in mice deficient in CC chemokine receptor 2*. Proc Natl Acad Sci U S A. 94(22):12053-8 (1997).

Methods

- 20 [00365] For the 48-hour TG peritonitis study, Example 1 was dosed twice a day with the first dose given one hour prior to TG injection. Total peritoneal cell counts were obtained on isolated cells by a cell counter. Blood was also collected in heparin from the retro-orbital sinus at the end of each study for flow cytometry and in EDTA for determination of drug concentration.
- 25 [00366] For flow cytometric analysis, peritoneal exudate cells (1×10^6) were washed once with FACS buffer (PBS/0.5% BSA) and resuspended in FACS buffer. Cells were incubated with an Fc-blocking antibody (BD Pharmingen) on ice for 15 min followed by addition of the following antibodies (BD Pharmingen): PE conjugated anti-F4/80, FITC conjugated anti-Ly6C, and Alexa 647 conjugated anti-
- 30 hCCR2. After 45 min on ice, cells were fixed by BD Cytofix for 15 min on ice, washed twice with FACS buffer, and resuspended in 200 μ l FACS buffer. Cellular events (40,000) were acquired for each sample and data were analyzed using FloJo

software (TreeStar). A FSC/SSC gate was set to include all monocytes (low SSC, higher FSC) while excluding granulocytes from the analysis. This gated population was then analyzed for Ly6C (FITC), F4/80 (PE) expression. Peritoneal monocytes/macrophage numbers were determined by multiplying total peritoneal cell counts obtained by the cell counter and the percentage of monocytes/macrophages identified by F4/80⁺ cells from flow cytometry. Statistical significance of differences between means was analyzed using the paired two-tailed t test with significance set at p values below 0.05.

10 Results

[00367] Example 1 was evaluated in the hCCR2 KI mouse TG peritonitis model to determine its EC50 in inhibiting monocyte/macrophage infiltration. Mice were administered thioglycollate, and dosed orally with Example 1 at 1, 25, or 100 mg/kg BID. Forty eight hours post TG treatment, peritoneal lavage was obtained for cellular infiltrate analysis by flow cytometry.

[00368] To distinguish between the recruited monocyte/macrophages versus resident macrophages and granulocytes, staining of both F4/80 and Ly6C monocyte/macrophage surface markers was used to define the recruited monocyte/macrophages. A dose-dependent inhibition in monocyte/macrophage infiltration was observed (Figure 21). Doses of 1, 25, and 100 mg/kg gave an inhibition of 24%, 74% and 78%, respectively. In three separate studies with multiple doses, the average EC50 for inhibition of monocyte/macrophage infiltration by this analysis was estimated to be 3.9 nM.

[00369] To assess the in vivo level of receptor occupancy by Example 1 in the 48-hour thioglycollate peritonitis model in the hCCR2 KI mouse, plasma levels of both Example 1 and mouse MCP-1 were measured. The caveat for this estimation is that only CCR2 and its major ligand MCP-1, were taken into consideration. The receptor occupancy of a ligand in the presence of a competitive inhibitor is defined by the Gaddum equation:

$$\frac{[RL]}{[R]} = \frac{1}{1 + (K_d / [L]) (1 + [I] / K_i)}$$

[00370] Since Example 1 is a competitive inhibitor of MCP-1 binding to CCR2, the amounts of both mouse MCP-1/CCR2 receptor complex and Example 1/CCR2 receptor complex can be determined using the serum levels of both mouse MCP-1 and protein-unbound Example 1 in plasma. The K_d for mouse MCP-1 binding to hCCR2 is 0.91 +/- 0.08 nM (n=8) which was determined in cold competition ligand binding experiments using 125 I-human MCP-1. The average K_i for Example 1 binding to hCCR2 is 1.3 nM. The fraction of mouse MCP-1/CCR2 receptor complexes is determined using the form of the equation described above. To determine the fraction of Example 1/CCR2 complexes the equation is re-defined as:

$$\frac{[RI]}{[R]} = \frac{1}{1 + (K_i / [I]) (1 + [L] / K_d)}$$

Finally, the amount of CCR2 free is determined from:

$$[CCR2]_{total} = [CCR2]_{free} + [mouse\ MCP-1/CCR2] + [Example\ 1/CCR2]$$

[00371] As shown in Table 13, the percent inhibition of monocyte/macrophage infiltration into the peritoneum at 48 hour reflects the percentage of Example 1/CCR2 receptor complex.

TABLE 13

Determination of in vivo receptor occupancy of Example 1 in blood of hCCR2 KI mice in the 48-hour TG peritonitis model

Dose (mg/kg)	Concentration of Mouse MCP-1 in plasma (nM)	Concentration of free Example 1 in plasma (nM) (fold IC90 CCR2 binding)	% mouse MCP-1 bound CCR2	% of Example 1-bound CCR2	% free CCR2	% inhibition of monocyte/macrophage infiltration ^a
100	0.015	53 (1.8)	0.04	97.6	2.4	78
25	0.017	14 (0.5)	0.16	91.2	8.6	74
1	0.005	1.4 (0.05)	0.26	51.4	48.3	24
0 (vehicle)	0	0	0	0	100	0

Section 3. Chronic Efficacy Studies

Experimental autoimmune encephalomyelitis (EAE)

Methods

[00372] To assess the effect of Example 1 on chronic models of disease, we used the EAE model of multiple sclerosis in hCCR2 KI mice. To study the effect of Example 1 on EAE model, 10 mice per group were used. On day 0, hCCR2 KI mice were immunized subcutaneously with a total of 200 μ l of 300 μ g myelin oligodendrocyte glycoprotein (MOG) 35-55 (Genemed Synthesis) mixed 1:1 with 300 μ g Mycobacterium tuberculosis (H37Ra) (Becton-Dickinson) in incomplete Freund's adjuvant (IFA) (Sigma-Aldrich). On day 0 (two hours post-immunization) and day 2, mice were injected intraperitoneally with 100 μ l of 400 ng pertussis toxin. Clinical scoring began on day 10, continued three times per week throughout the study, and was based on a scale of 0-5: 0, no signs of disease; 0.5, partial tail weakness; 1, limp tail or waddling gait with tail tonic; 1.5, waddling gait with partial tail weakness; 2, waddling gait with limp tail (ataxia); 2.5 (ataxia with partial limb paralysis; 3, full paralysis of one limb; 3.5, full paralysis of one limb with partial paralysis of a second limb; 4, full paralysis of two limbs; 4.5, moribund; 5, death. Oral dosing of Example 1 at 25 mg/kg and 55 mg/kg (BID) was initiated on day 1.

Results

[00373] Example 1 at both doses reduced the area under curve (AUC) of the clinical score by 49% ($p < 0.05$) (Figure 22). The IC₅₀ is 3.7 nM for Example 1 in ¹²⁵I-mouse MCP-1 binding to hCCR2-expressing cells, hPBMCs (mimicking hCCR2 KI setting). Based on this IC₅₀ value, the 25 and 55 mg/kg doses resulted in a free plasma trough concentration of 1- and 3-fold the binding IC₉₀. Histological evaluation of the spinal cord on Day 22 did not demonstrate a significant difference in total inflammatory cellular infiltrate between mice treated with Example 1 versus vehicle. A marked neutrophil infiltrate was observed in mice treated with compound.

30

What is claimed is:

1. A compound selected from:

(7*R*,8*S*)-8-((3*S*)-3-(((benzyloxy)carbonyl)amino)-2-oxo-1-pyrrolidinyl)-1,4-dioxaspiro[4.5]decane-7-carboxylic acid, or a salt thereof;

benzyl ((3*S*)-1-((7*R*,8*S*)-7-(azidocarbonyl)-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;

benzyl ((3*S*)-1-((7*R*,8*S*)-7-isocyanato-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;

benzyl ((3*S*)-1-((7*R*,8*S*)-7-acetamido-1,4-dioxaspiro[4.5]dec-8-yl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof;

benzyl ((3*S*)-1-((1*S*,2*R*)-2-acetamido-4-oxocyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof; and

benzyl ((3*S*)-1-((1*S*,2*R*,4*R*)-2-acetamido-4-(*tert*-butylamino)cyclohexyl)-2-oxo-3-pyrrolidinyl)carbamate, or a salt thereof.