THERMOSTABLE CAS9 NUCLEASES

FIELD OF THE INVENTION

The present invention relates to the field of genetic engineering and more particularly to nucleic acid editing and genome modification. The present invention concerns genetic engineering tools in the form of nucleases which can be configured for sequence-directed site-specific binding, nicking, cutting and modification of genetic material; also ribonucleoproteins which exert activity, particularly nuclease activity, on sequence specific sites of genetic material, and modified nucleases and ribonucleoproteins for use as markers. The invention therefore also concerns associated expression constructs for delivery and expression of nucleases and guide RNAs within non-human cells. Further, the invention concerns the sequence-specific editing of nucleic acids *in vitro* or *in vivo* and methods used to achieve that. A particular area to which the invention relates is the genetic manipulation of thermophilic organisms, particularly microorganisms.

BACKGROUND TO THE INVENTION

It was first demonstrated in 2007 that CRISPR-Cas is an adaptive immune system in many bacteria and most archaea (Barrangou *et al.*, 2007, Science 315: 1709-1712), Brouns *et al.*, 2008, Science 321: 960-964). Based on functional and structural criteria, two classes of CRISPR-Cas systems that each comprise three types have so far been characterized, most of which use small RNA molecules as guide to target complementary DNA sequences (Makarova *et al.*, 2015, Nat Rev Microbiol 13: 722-736; Mahanraju *et al.*, 2016, Science 353: aad5147).

In a recent study by the Doudna/Charpentier labs, a thorough characterization of the effector enzyme of the class 2/type II CRISPR-Cas system (Cas9) was performed, including demonstration that the introduction of designed CRISPR RNA guides (with specific spacer sequences) targets complementary sequences (protospacers) on a plasmid, causing double strand breaks of this plasmid (Jinek *et al.*, 2012, Science 337: 816-821). Following Jinek *et al.*, 2012, Cas9 is used as a tool for genome editing.

Cas9 has been used to engineer the genomes of a range of eukaryotic cells (*e.g.* fish, plant, man) (Charpentier and Doudna, 2013, Nature 495: 50-51).

In addition, Cas9 has been used to improve yields of homologous recombination in bacteria by selecting for dedicated recombination events (Jiang *et al.*, 2013, Nature Biotechnol 31: 233-239). To achieve this, a toxic fragment (Targeting construct) is co-transfected with a rescuing fragment carrying the desired alteration (Editing construct, carrying point mutation or deletions). The Targeting construct consists of Cas9 in combination with a design CRISPR and an antibiotic resistance marker, defining the site of the desired recombination on the host chromosome; in the presence of the corresponding antibiotic, integration of the Targeting construct in the host chromosome is selected for. Only when the additional recombination occurs of the Editing construct with the CRISPR target site on the host chromosome, the host can escape from the auto-immunity problem. Hence, in the presence of the antibiotic, only the desired (marker-free) mutants are able to survive and grow. A related strategy to select for subsequent removal of the integrated Targeting construct from the chromosome is presented as well, generating a genuine marker free mutant.

It has been established in recent years that CRISPR-Cas mediated genome editing constitutes a useful tool for genetic engineering. It has been established that the prokaryotic CRISPR systems serve their hosts as adaptive immune systems (Jinek *et al.*, 2012, Science 337: 816-821) and can be used for quick and effective genetic engineering (Mali *et al.*, 2013, Nat Methods 10:957-963, for example), requiring only modification of the guide sequence in order to target sequences of interest.

However, there is a continuing need for the development of agents with improved sequence-specific nucleic acid detection, cleavage and manipulation under a variety of experimental conditions for application in the area of genetic research and genome editing. In particular, currently available sequence-specific genome editing tools, including Cas9, are not applicable for use in all conditions or organisms, for example, sequence-specific nucleases are relatively thermo-sensitive and therefore not applicable for use in strictly thermophilic microorganisms (which are capable of growth between 41°C and 122°C and grow optimally in the ranges of temperatures from >45°C to 80°C with hyperthermophiles capable of optimal growth above 80°C), for

example, microorganisms that are used in industrial fermentations or for *in vitro* laboratory processes conducted at elevated temperatures.

To date there is no experimental evidence for active Cas9 proteins in thermophiles. Based on a comparative genome screening by Chylinski *et al*. (2014; Nucleic Acids Research 42: 6091-61-05) on the presence of Cas9 in bacteria it was found that the Type II-C CRISPR-Cas system is only present in approximately 3.3% of all bacterial genomes. Among thermophilic bacteria, the Type II system is underrepresented based on statistical analysis (P=0.0019). In addition, no Type II system has been found in archaea however, this could possibly be due to the absence of the RNase III protein (involved in the Type II system) in archaea. Chylinski, *et al*., (2014; Nucleic Acids Research 42: 6091-6105) did describe the classification and evolution of type II CRISPR-Cas systems, in particular, two species are identified which exhibit these systems, however these species grow maximally at 55°C and do not exhibit strictly thermophilic growth with optimum growth temperature 60-80°C, with hyperthermophiles capable of growing optimally above 80°C.

Despite the rarity of the CRISPR-Cas system in bacterial genomes and in particular the fact that Cas9 has been found only in bacteria (not archaea) with optimal growth temperatures below 45°C, the inventors have surprisingly discovered several thermostable Cas9 variants which enable genome editing to be carried out at elevated temperatures. The inventors have also discovered optimised protospacer adjacent motif (PAM) sequences that work with the thermostable Cas9 variants to enable genome editing to be carried out over a wide range of temperatures, including at the elevated temperatures. These Cas9 nucleases, and RNA molecules that are designed with knowledge of the associated PAM sequences, provide novel tools for genetic engineering at elevated temperatures and are of particular value in the genetic manipulation of thermophilic organisms; particularly microorganisms.

A phylogenetic re-assessment of the thermophilic genus *Geobacillus* has recently been performed, which resulted in the creation of a new genus, *Parageobacillus*. Consequently, some species previously of the genus *Geobacillus* have been systematically re-assigned to *Parageobacillus* and re-named accordingly (Aliyu *et al*., (2016) Systematic and Applied Microbiology 39:527-533).

Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and the CRISPR-associated (Cas) proteins provide adaptive and heritable immunity in prokaryotes against invading genetic elements (Brouns *et al*. Science 321, (2008); Barrangou et al. CRISPR provides acquired resistance against viruses in prokaryotes. Science 315, (2007); Wright *et al*. Cell 164, 29–44 (2016); Mohanraju *et al*. Science 353, aad5147 (2016)). CRISPR-Cas systems are subdivided into two classes (1 and 2) and six types (I-VI), depending on their complexity and signature protein (Makarova *et al. Nat. Rev. Microbiol.* 13, 722–736 (2015)). Class 2 systems, including type-II CRISPR-Cas9 and type V CRISPR-Cas12a (previously called CRISPR-Cpf1) have recently been exploited as genome engineering tools for both eukaryotes (Komor *et al*. Cell 168, 20–36 (2017); Puchta, Curr. Opin. Plant Biol. 36, 1–8 (2017); Xu *et al*. J. Genet. Genomics 42, 141–149 (2015); Tang *et al*. Nat. Plants 3, 17018 (2017); Zetsche *et al*. Nat. Biotechnol. 35, 31–34 (2016)) and prokaryotes (Mougiakos, *et al*. Trends Biotechnol. 34, 575–587 (2016)). These systems are among the simplest CRISPR-Cas systems known as they introduce targeted double stranded DNA breaks (DSBs) based on a ribonucleoprotein (RNP) complex formed by a single Cas endonuclease and an RNA guide.

To date, *Streptococcus pyogenes* Cas9 (SpCas9) is the best characterized and most widely employed Cas9 for genome engineering. Although a few other type-II systems have been characterized, none of them is derived from a thermophilic organism (Nakade, *et al*. Bioengineered 1–9 (2017). doi:10.1080/21655979.2017.1282018). Characterization of such CRISPR-Cas systems would be interesting to gain fundamental insights as well as to develop novel applications.

Although basic genetic tools are available for a number of thermophiles (Taylor *et al*. Microb. Biotechnol. 4, 438–448 (2011); Olson, et al. Curr. Opin. Biotechnol. 33, 130–141 (2015); Zeldes, *et al*. Front. Microbiol. 6, 1209 (2015)), the efficiency of these tools is still too low to enable full exploration and exploitation of this interesting group of organisms. Based on our finding that SpCas9 is not active *in vivo* ≥42°C, we have previously developed an SpCas9-based engineering tool for facultative thermophiles, combining homologous recombination at elevated temperatures and SpCas9-based counter-selection at moderate temperatures (Mougiakos *et al. ACS Synth. Biol.* 6, 849–861 (2017)). However, a Cas9-based editing and silencing tool for obligate

thermophiles is not yet available as SpCas9 is not active at or above 42°C (Mougiakos *et al. ACS Synth. Biol.* 6, 849–861 (2017)) and to date no thermophilic Cas9 has been characterized.

SUMMARY

The inventors have discovered and characterised ThermoCas9: an RNA-guided DNA-endonuclease from the CRISPR-Cas type-IIC system of the thermophilic bacterium *Geobacillus thermodenitrificans* T12. The inventors have surprisingly shown its *in vitro* activity over a broad temperature range, demonstrated the importance of the sgRNA-structure for thermostability and applied ThermoCas9 for *in vivo* genome editing across a wide temperature range.

In one aspect, the present disclosure refers to a nucleoprotein complex comprising a Cas protein, at least one targeting RNA molecule which recognises a target nucleic acid sequence in a double stranded target polynucleotide, and the target polynucleotide, wherein

the Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith;

the double stranded target polynucleotide comprises a target nucleic acid strand, comprising said target nucleic acid sequence, and a non-target nucleic acid strand, comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence and a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3'; wherein the nucleoprotein complex is not in a human cell.

In another aspect, the present disclosure refers to a transformed non-human cell, having a double stranded target polynucleotide comprising a target nucleic acid sequence, wherein the double stranded target polynucleotide comprises a target nucleic acid strand, comprising said target nucleic acid sequence, and a non-target nucleic acid strand, comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence, said cell comprising:

a clustered regularly interspaced short palindromic repeat (CRISPR)-associated (Cas) protein having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith;

at least one targeting RNA molecule which recognizes the target nucleic acid sequence in the target nucleic acid strand, wherein the non-target strand further comprises a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3'; and

an expression vector comprising a nucleic acid encoding at least one of said Cas protein and said targeting RNA molecule.

In another aspect, the present disclosure refers to an *in vitro* method of binding, cleaving, marking or modifying a double stranded target polynucleotide, wherein the double stranded target polynucleotide comprises a target nucleic acid strand comprising a target nucleic acid sequence, and a non-target nucleic acid strand comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence, said method comprising:

a. designing at least one targeting RNA molecule, wherein the targeting RNA molecule recognizes the target sequence in the target strand, and the non-target strand further comprises a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3';

b. forming a ribonucleoprotein complex comprising the targeting RNA molecule and a Cas protein, wherein the isolated Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith; and

c. the ribonucleoprotein complex binding, cleaving, marking or modifying the target polynucleotide; and wherein said method is not used in human cells.

In yet another aspect, the present disclosure refers to the *in vitro* use of at least one targeting RNA molecule and a Cas protein to form a nucleoprotein complex for binding, cleaving, marking or modifying a double stranded target polynucleotide comprising a target nucleic acid sequence, wherein:

the double stranded target polynucleotide comprises a target nucleic acid strand, comprising said target nucleic acid sequence, and a non-target nucleic acid strand, comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence;

the Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith;

the at least one targeting RNA molecule recognizes the target sequence;

the non-target nucleic acid strand further comprises a protospacer adjacent motif (PAM) sequence directly adjacent to the 3' end of the protospacer nucleic acid sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3'; and wherein said use is not in a human cell.

Accordingly, the present disclosure provides an isolated clustered regularly interspaced short palindromic repeat (CRISPR)-associated (Cas) protein or polypeptide comprising;

a. the amino acid motif EKDGKYYC [SEQ ID NO: 2]; and/or
b. the amino acid motif $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and/or
c. the amino acid motif X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and/or
d. the amino acid motif X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and/or
e. the amino acid motif $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

For the avoidance of doubt proteins, polypeptides or nucleic acids coding for Cas proteins of the present disclosure may also be referred to as "GtCas9" or

“ThermoCas9”. “GtCas9” and “ThermoCas9” are used interchangeably throughout the specification and have the same meaning.

A polypeptide in the context of this disclosure may be viewed as a fragment of the full length Cas protein. Such fragments may be inactive and used in ways and for purposes not associated directly with binding, editing and/or cutting of genetic material, for example for standards in assays or raising antibodies or the like.

In preferred examples however, the Cas protein or polypeptide is functional and capable of cleavage, binding, marking or modifying at a temperature in the range 20°C and 100°C, inclusive, when associated with at least one targeting RNA molecule, and a polynucleotide comprising a target nucleic acid sequence recognised by the targeting RNA molecule. Preferably the Cas protein or polypeptide is functional and capable of said cleavage, binding, marking or modifying at a temperature in the range 50°C and 70°C, for example 55°C or 60°C.

In particular examples, the disclosure may provide a Cas protein or polypeptide comprising the amino acid motif EKDGKYYC [SEQ ID NO: 2]. In other examples, the Cas proteins or polypeptides may further comprise the amino acid motif $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine.

In other examples the Cas proteins or polypeptides defined herein may additionally further comprise the amino acid motif X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine.

In other examples, the Cas proteins or polypeptides defined herein may additionally further comprise the amino acid motif X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine.

In other examples, the Cas proteins or polypeptides defined herein may additionally further comprise the amino acid motif $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID

NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

In accordance with the present disclosure, it may be appreciated that a Cas protein or polypeptide of the disclosure may comprise any of the motifs of SEQ ID NOs 2 to 6, either alone or in combination. The following summarises each of the combinations of motifs which may characterize Cas proteins or polypeptides of the disclosure:

EKDGKYYC [SEQ ID NO: 2].

EKDGKYYC [SEQ ID NO: 2]; and $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine.

EKDGKYYC [SEQ ID NO: 2]; and $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine.

EKDGKYYC [SEQ ID NO: 2]; and $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine.

EKDGKYYC [SEQ ID NO: 2]; and $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected

from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

EKDGKYYC [SEQ ID NO: 2]; and $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

EKDGKYYC [SEQ ID NO: 2]; and $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

EKDGKYYC [SEQ ID NO: 2]; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

EKDGKYYC [SEQ ID NO: 2]; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine.

EKDGKYYC [SEQ ID NO: 2]; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine.

EKDGKYYC [SEQ ID NO: 2]; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

EKDGKYYC [SEQ ID NO: 2]; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine.

EKDGKYYC [SEQ ID NO: 2]; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

EKDGKYYC [SEQ ID NO: 2]; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

$X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is

independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine.

$X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine.

$X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

$X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

$X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of

Alanine, Glutamate or Arginine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine.

$X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine.

X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine.

In another example, the present disclosure provides an isolated Cas protein or polypeptide fragment thereof having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 77% identity therewith, wherein the Cas protein or polypeptide fragment thereof comprises any of the following motifs or amino acids, either alone or in combination:

An RuvC-I domain comprising IGLDIGITSIG [SEQ ID NO: 23], preferably IGLDIGITSIGWAVINLD [SEQ ID NO: 24].

A Bridge domain comprising RSARR [SEQ ID NO: 25], preferably PRRLARSARRRLRRRKHRLERIRRL [SEQ ID NO: 26]; and/or

An α-helical/recognition lobe domain comprising WQLR [SEQ ID NO: 27]; and/or
An α-helical/recognition lobe domain comprising HLAKRRG [SEQ ID NO: 28], preferably LARILLHLAKRRG [SEQ ID NO: 29]; and/or
An α-helical/recognition lobe domain comprising IFAKQ [SEQ ID NO: 30], preferably EIKLIFAKQ [SEQ ID NO: 31]; and/or
An α-helical/recognition lobe domain comprising IWASQR [SEQ ID NO: 32]; and/or KVGFCTFEPKEKRAPK [SEQ ID NO: 33]; and/or
FTVWEHINKLRL [SEQ ID NO: 34]; and/or

An RuvC-II domain comprising the motif IANPVVMRALTQ [SEQ ID NO: 35], preferably IANPVVMRALTQARKVVNAIIKKYG [SEQ ID NO: 36]; and/or
An RuvC-II domain comprising the motif ELAR [SEQ ID NO: 37], preferably IHIELARE [SEQ ID NO: 38]; and/or

An HNH domain comprising the motif QNGKCAY [SEQ ID NO: 39], preferably IVKFKLWSEQNGKCAY [SEQ ID NO: 40]; and/or
An HNH domain comprising the motif VDHVIP [SEQ ID NO: 41], preferably VDHVIPYSRSLDDSYTNKVL [SEQ ID NO: 42]; and/or

An RuvC-III domain comprising the motif DTRYISRFLAN [SEQ ID NO: 43]; and/or
An RuvC-III domain comprising the motif VYTVNGRITAHLRSRW [SEQ ID NO: 44]; and/or

An RuvC-III domain comprising the motif HHAVDA [SEQ ID NO: 45], preferably HHAVDAAIVA [SEQ ID NO: 46]; and/or

Preferably, the present disclosure provides an isolated Cas protein or polypeptide fragment thereof having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 77% identity therewith, wherein the Cas protein or polypeptide fragment thereof comprises each of the amino acid motifs [SEQ ID NO: 23] to [SEQ ID NO: 46] in combination.

In another example, the present disclosure provides an isolated Cas protein or polypeptide fragment thereof having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 77% identity therewith. Preferably the Cas protein or polypeptide is capable of binding, cleavage, marking or modifying at a temperature in the range 20°C and 100°C inclusive. Preferably the Cas protein or polypeptide is capable of said cleavage, binding, marking or modifying at a temperature in the range between 20°C and 70°C, for example 25°C, 55°C, 60°C or 65°C. Preferably the Cas protein or polypeptide is capable of said cleavage, binding, marking or modifying at a temperature in the range between 50°C and 70°C, for example 55°C or 60°C. Preferably the Cas protein or polypeptide is capable of said cleavage, binding, marking or modifying at a temperature in the range between 30°C and 80°C, at a temperature between 37°C and 78°C, preferably at a temperature above 55°C; more preferably at a temperature between 55°C and 80°C; even more preferably at a temperature between 55°C and 65°C or 60°C and 65°C.

The present disclosure also provides uses of a targeting RNA molecule and a Cas protein or polypeptide provided herein, for binding, cleaving, marking or modifying a target polynucleotide comprising a target nucleic acid sequence. The targeting RNA molecule recognizes the target nucleic acid sequence on a target nucleic acid strand of the polynucleotide.

The target polynucleotide that comprises the target nucleic acid sequence may be double stranded and so comprise a target nucleic acid strand, comprising said target nucleic acid sequence, and a non-target nucleic acid strand, comprising a protospacer nucleic acid sequence. The protospacer nucleic acid sequence is substantially

complementary to the target nucleic acid sequence and pairs with it in the double stranded target polynucleotide. The non-target nucleic acid strand may further comprise a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence. The PAM sequence may be at least 6, 7, or 8 nucleic acids in length. Preferably, the PAM sequence has a cytosine in the fifth position. Preferably the PAM sequence comprises the sequence 5'-NNNNC-3', so that from the 5'-end the PAM sequence begins 5'-NNNNC-3'. Additionally or alternatively, the PAM sequence may have an adenine in the eighth position, so that the PAM sequence comprises the sequence 5'-NNNNNNNA-3', and from the 5'-end the PAM sequence begins 5'-NNNNNNNA-3'. Additionally or alternatively, the PAM sequence may have a cytosine in one or more of the first, second, third, fourth, and sixth positions, such that from the 5'-end the PAM sequence begins 5'-CNNNN-3', 5'-NCNNN-3', 5'-NNCNN-3', 5'-NNNCN-3', and/or 5'-NNNNNC-3'. Optionally the PAM sequence comprises, so that from the 5'-end the PAM sequence begins, 5'-CCCCCCNA-3' [SEQ ID NO: 10], and further preferably the PAM sequence comprises, so that from the 5'-end the PAM sequence begins, 5'-CCCCCCAA-3' [SEQ ID NO: 11]. Other preferred PAM sequences include 5'-ATCCCCAA-3' [SEQ ID NO: 21] and 5'-ACGGCCAA-3' [SEQ ID NO: 22].

Preferably, the Cas protein or polypeptide is capable of the binding, cleaving, marking or modifying at a temperature in the range 40°C to 80°C inclusive, preferably in the range 45°C to 80°C inclusive, and further preferably in the range 50°C to 80°C inclusive. For example, the binding, cleaving, marking or modifying occurs at a temperature of 45°C, 46°C, 47°C, 48°C, 49°C, 50°C, 51°C, 52°C, 53°C, 54°C, 55°C, 56°C, 57°C, 58°C, 59°C, 60°C, 61°C, 62°C, 63°C, 64°C, 65°C, 66°C, 67°C, 68°C, 69°C, 70°C, 71°C, 72°C, 73°C, 74°C, 75°C, 76°C, 77°C, 78°C, 79°C or 80°C. More preferably the Cas protein or polypeptide is capable of the binding, cleaving, marking or modifying at a temperature in the range 55 to 65°C. In preferred examples, a Cas protein or polypeptide fragment of the disclosure may comprises an amino acid sequence of at least 75% identity; preferably at least 85%; more preferably at least 90%; even more preferably at least 95% identity to SEQ ID NO: 1.

The Cas protein or polypeptide may be used in combination with a targeting RNA molecule that recognizes a target nucleic acid sequence on the target nucleic acid

strand, where the non-target nucleic acid sequence has a PAM sequence directly adjacent the 3' end of the protospacer sequence on the non-target strand, as disclosed herein. Thus, the PAM sequence may comprise the sequence 5'-NNNNC-3', and the Cas protein may bind, cleave, mark or modify the target strand at a temperature in the range 20°C and 100°C inclusive, preferably in the range 30°C and 90°C inclusive, in the range 37°C and 78°C inclusive, in the range 40°C and 80°C inclusive, in the range 50°C and 70°C inclusive, or in the range 55°C and 65°C, inclusive. Preferably from the 5'-end the PAM sequence begins 5'-NNNNC-3' and the Cas protein may bind, cleave, mark or modify the target strand at a temperature in the range 20°C and 100°C inclusive, preferably in the range 30°C and 90°C inclusive, in the range 37°C and 78°C inclusive, in the range 40°C and 80°C inclusive, in the range 50°C and 70°C inclusive, or in the range 55°C and 65°C, inclusive. Preferably from the 5'-end the PAM sequence begins 5'-NNNNNNNA-3' and the Cas protein may bind, cleave, mark or modify the target strand at a temperature in the range 20°C and 100°C inclusive, preferably in the range 30°C and 90°C inclusive, in the range 37°C and 78°C inclusive, in the range 40°C and 80°C inclusive, the range 50°C and 70°C inclusive, or in the range 55°C and 65°C, inclusive. Further preferably the 5'-end of the PAM sequence begins 5'-NNNNCNNA-3' [SEQ ID NO: 47] and the Cas protein may bind, cleave, mark or modify the target strand at a temperature in the range 20°C and 100°C inclusive, preferably in the range 30°C and 90°C inclusive, in the range 37°C and 78°C inclusive, in the range 40°C and 80°C inclusive, in the range 50°C and 70°C inclusive, or in the range 55°C and 65°C, inclusive.

More particularly, a Cas protein or polypeptide of the disclosure may comprise an amino acid sequence with a percentage identity with SEQ ID NO:1 as follows: at least 60%, at least 61%, at least 62%, at least 63%, at least 64%, at least 65%, at least 66%, at least 67%, at least 68%, at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, at least 99.5% or at least 99.8%. The percentage identity may be at least 89%. The percentage identity may be at least 90%. Preferably the percentage identity will be at least 95%, for example 98%.

The percentage amino acid sequence identity with SEQ ID NO: 1 is determinable as a function of the number of identical positions shared by the sequences in a selected comparison window, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

A Cas protein or polypeptide fragment of the disclosure may be characterised in terms of both the reference sequence SEQ ID NO: 1 and any aforementioned percentage variant thereof as defined by percentage sequence identity, alone or in combination with any of the aforementioned amino acid motifs (i.e. SEQ ID NOS 2 and/or 3 and/or 4 and/or 5 and/or 6) as essential features.

The disclosure provides a use of a targeting RNA molecule as provided herein and a Cas protein or polypeptide of the disclosure for binding, cleaving, marking or modifying a target nucleic acid strand comprising a target nucleic acid sequence. Preferably said binding, cleaving, marking or modifying occurs at a temperature disclosed herein, for example at a temperature of between 20 and 100°C. The disclosure also provides a method of binding, cleaving, marking or modifying a target nucleic acid sequence in a target nucleic acid strand comprising designing a targeting RNA molecule as provided herein and forming a ribonucleoprotein complex comprising the targeting RNA molecule and a Cas protein or polypeptide of the disclosure. Preferably the ribonucleoprotein complex binding, cleaving, marking or modifying the target nucleic acid sequence at a temperature disclosed herein, for example at a temperature of between 37 and 100°C.

The uses and methods of the disclosure may be carried out, and the nucleoproteins of the disclosure formed and used, *in vivo*, for example in bacterial cells. The uses and methods of the disclosure may be carried out, and the nucleoproteins of the disclosure formed and used, *in vivo*, except in human cells. Alternatively the uses and methods of the disclosure may be carried out, and the nucleoproteins of the disclosure formed and used, *in vitro*. The Cas protein of the disclosure may be provided in isolated form, for example when used *in vitro* or when added to cells by transfection, the Cas protein may be heterologously expressed, for example following transient or stable transformation of the cell by nucleic acid encoding the Cas protein, the targeting RNA

molecule may be transcribed from an expression vector following transient or stable transformation of the cell by nucleic acid encoding the RNA molecule, and/or the RNA molecule may be provided in isolated form, for example when used *in vitro* or when added to cells by transfection. In preferred examples, the Cas protein or polypeptide is expressed from the genome of a host cell, following stable intergration of a nucleic acid encoding the Cas protein or polypeptide in the genome of the host cell. Thus the Cas protein and/or RNA molecule may be added to the *in vivo* or *in vitro* environment using any artificial or contrived method for adding a protein or nucleic acid molecule to a cell in which it is not otherwise present.

The polynucleotide comprising the target nucleic acid sequence may be cleaved by the Cas protein, and optionally the cleavage may be DNA cleavage. The target nucleic acid strand comprising the target sequence may be double stranded DNA and the method or use may result in a double stranded break in the polynucleotide comprising the target nucleic acid sequence. The polynucleotide comprising the target nucleic acid sequence may be double stranded DNA, the Cas protein may lack the ability to cut the double stranded DNA and the use or method may result in gene silencing of the polynucleotide.

The Cas protein or polypeptide may be provided for the methods, uses and nucleoproteins of the disclosure at a concentration of 250 nM or less, for example at a concentration of 200 nM or less, 150 nM or less, 100 nM or less, 50 nM or less, 25nM or less, 10 nM or less, 5nM or less, 1nM or less or 0.5 nM or less. Alternatively, the Cas protein or polypeptide may be provided at a concentration of at least 0.5 nM, at least 1nM, at least 5nM, at least 10nM, at least 25nM, at least 50 nM, at least 100 nM, at least 150 nM, at least 200 nM, or at least 250 nM. The PAM sequence of the disclosure may have an adenine in the eighth position, so that the PAM sequence comprises the sequence 5'-NNNNNNNA-3', and the concentration of Cas protein or polypeptide may be 100 nM or less, 50 nM or less, 25nM or less, 10 nM or less, 5nM or less, 1nM or less or 0.5 nM or less. The PAM sequence may comprise the sequence 5'-NNNNCNNA-3' [SEQ ID NO: 47], and the concentration of Cas protein or polypeptide may be 100 nM or less, 50 nM or less, 25nM or less, 10 nM or less, 5nM or less, 1nM or less or 0.5 nM or less. The PAM sequence may comprise the sequence 5'-CCCCCCNA-3' [SEQ ID NO: 10], and the concentration of Cas protein

or polypeptide may be 100 nM or less, 50 nM or less, 25nM or less, 10 nM or less, 5nM or less, 1nM or less or 0.5 nM or less.

Also, the disclosure provides nucleic acids encoding any of the aforementioned proteins or polypeptides of the disclosure. The nucleic acids may be isolated or in the form of expression constructs.

In all aforementioned examples of the present disclosure, amino acid residues may be substituted conservatively or non-conservatively. Conservative amino acid substitutions refer to those where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not alter the functional properties of the resulting polypeptide.

Similarly it will be appreciated by a person of average skill in the art that nucleic acid sequences may be substituted conservatively or non-conservatively without affecting the function of the polypeptide. Conservatively modified nucleic acids are those substituted for nucleic acids which encode identical or functionally identical variants of the amino acid sequences. It will be appreciated by the skilled reader that each codon in a nucleic acid (except AUG and UGG; typically the only codons for methionine or tryptophan, respectively) can be modified to yield a functionally identical molecule. Accordingly, each silent variation (i.e. synonymous codon) of a polynucleotide or polypeptide, which encodes a polypeptide of the present disclosure, is implicit in each described polypeptide sequence.

The disclosure provides a transformed cell, having a target nucleic acid sequence in a double stranded target polynucleotide, said cell comprising a Cas protein or polypeptide as provided herein and at least one targeting RNA molecule as provided herein, and an expression vector comprising a nucleic acid encoding at least one of said Cas protein and said targeting RNA molecule. The Cas protein and targeting RNA molecule may enable or permit binding, cleaving, marking or modifying of the target sequence to occur in the transformed cell at a raised temperature, or at a range of temperatures, for example between 37 and 100°C, as disclosed herein. The disclosure further provides a method of binding, cleaving, marking or modifying a target nucleic acid in a cell comprising either 1) transforming, transfecting or transducing the cell with

an expression vector comprising a nucleotide sequence encoding a Cas protein or polypeptide of the disclosure and a nucleotide sequence encoding a targeting RNA molecule of the disclosure; or 2) transforming, transfecting or transducing the cell with an expression vector comprising a nucleotide sequence encoding a Cas protein or polypeptide of the disclosure and a further expression vector comprising a nucleotide sequence encoding a targeting RNA molecule of the disclosure; or 3) transforming, transfecting or transducing the cell with an expression vector comprising a nucleotide sequence encoding a Cas protein or polypeptide of the disclosure, and delivering a targeting RNA molecule as provided herein to, or into the cell. The Cas protein or polypeptide may be expressed from the genome of the transformed cell, for example following stable integration into the genome of a nucleotide sequence encoding the Cas protein or polypeptide.

The disclosure also provides kits comprising one or more of the reagents for carrying out the uses and methods of the disclosure, or for generating the transformed cells or nucleoprotein complex of the disclosure, said kits including: a Cas protein or polypeptide of the disclosure or an expression vector comprising a nucleic acid sequence encoding a Cas protein or polypeptide of the disclosure; and/or a targeting RNA molecule of the disclosure or an expression vector comprising a nucleic acid sequence encoding a targeting RNA molecule of the disclosure. The kits may further include instructions for carrying out the disclosure, for example instructions for how to design a targeting RNA molecule in accordance with the disclosure.

RNA Guides and Target Sequences

Cas proteins of the invention allow for sequence-specific binding, cleavage, tagging, marking or modification of target nucleic acids at elevated temperatures. Target nucleic acids may be DNA (single-stranded or double-stranded), RNA or synthetic nucleic acids. A particularly useful application of the present invention is the sequence-specific targeting and modification of genomic DNA by one or more Cas proteins of the invention in complex with one or more guide RNAs (gRNAs) that complementarily bind to a targeted sequence of the genomic DNA. Consequently, the target nucleic acid is preferably double-stranded DNA. Such targeting may be performed *in vitro* or *in vivo*. Preferably such targeting is performed *in vivo*. In this

way, Cas proteins of the invention may be used to target and modify specific DNA sequences located in the genomic DNA of a cell. It is envisaged that the Cas system may be used to modify genomes in a variety of cell types of and/or in different organisms.

The gRNAs, also called targeting RNA molecules, recognize the target nucleic acid sequence on the polynucleotide target strand. The RNA molecules may be designed to recognize a target sequence in a double stranded target polynucleotide, wherein the non-target strand comprises a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence. Disclosed herein are PAM sequences that work in an optimal manner with the Cas proteins and polypeptides of the invention. With knowledge of these PAM sequences, gRNAs may be designed for use with the Cas proteins and polypeptides of the invention across the temperature ranges and increased temperatures of the invention.

Accordingly, the present invention provides a ribonucleoprotein complex comprising a Cas protein or a polypeptide of the invention as hereinbefore described, and further comprising at least one RNA molecule which has a targeting function in that it recognizes a particular nucleotide sequence in a target polynucleotide. The present invention also provides use of at least one targeting RNA molecule and a Cas protein or polypeptide for binding, cleaving, marking or modifying a target nucleic acid strand, and a method of binding, cleaving, marking or modifying a target nucleic acid sequence in a target nucleic acid strand using a ribonucleoprotein or nucleoprotein of the invention, as well as transformed non-human cells having the Cas protein or polypeptide and targeting RNA molecule. The target polynucleotide may further comprise a defined PAM sequence directly adjacent the 3' end of a protospacer sequence, in accordance with a PAM sequence provided herein. The PAM sequence may be 6, 7, or 8 nucleic acids in length, or longer, preferably 8 nucleic acids in length. Preferably, the RNA molecule is a single-stranded RNA molecule, e.g. a CRISPR RNA (crRNA) and is associated, e.g. by hybridization with a tracrRNA. The targeting RNA may be a chimera of a crRNA and tracrRNA. The aforementioned RNA molecules may have a ribonucleotide sequence of at least 90% identity, or complementarity to a target nucleotide sequence. Optionally, the RNA molecule has a ribonucleotide sequence of at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at

least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% identity or complementarity to a target nucleotide sequence. The preferred target nucleotide sequence is a DNA.

In a preferred aspect, the present invention provides a ribonucleoprotein complex as hereinbefore described, wherein the at least one targeting RNA molecule is substantially complementary along its length to a target DNA sequence.

The targeting RNA molecule may be bound to or associated with the target sequence within the nucleoprotein complex, so that the target polynucleotide, comprising the target sequence and PAM sequence on the non-target strand, may be associated with and so form part of a nucleoprotein complex of the invention.

Alteration of the sequence of the RNA guide which associates with the Cas protein of the invention therefore allows the Cas protein to be programmed to mark or cut double-stranded DNA at sites complementary to the guide RNA.

Preferably, the length of the at least one targeting RNA molecule in a ribonucleoprotein complex of the invention is in the range 35 to 135 residues, optionally in the range 35 to 134 residues, 35 to 133 residues, 35 to 132 residues, 35 to 131 residues, 35 to 130 residues, 35 to 129 residues, 35 to 128 residues, 35 to 127 residues, 35 to 126 residues, 35 to 125 residues, 35 to 124 residues, 35 to 123 residues, 35 to 122 residues, 35 to 121 residues, 35 to 120 residues, 35 to 119 residues, 35 to 118 residues, 35 to 117 residues, 35 to 116 residues, 35 to 115 residues, 35 to 114 residues, 35 to 113 residues, 35 to 112 residues, 35 to 111 residues, 35 to 100 residues, 35 to 109 residues, 35 to 108 residues, 35 to 107 residues, 35 to 106 residues, 35 to 105 residues, 35 to 104 residues, 35 to 103 residues, 35 to 102 residues, 35 to 101 residues, 35 to 100 residues, 35 to 99 residues, 35 to 98 residues, 35 to 97 residues, 35 to 96 residues, 35 to 95 residues, 35 to 94 residues, 35 to 93 residues, 35 to 92 residues, 35 to 91 residues, 35 to 90 residues, 35 to 89 residues, 35 to 88 residues, 35 to 87 residues, 35 to 86 residues, 35 to 85 residues, 35 to 84 residues, 35 to 83 residues, 35 to 82 residues, 35 to 81 residues, 35 to 80 residues, 35 to 79 residues, 35 to 78 residues, 35 to 77 residues, 35 to 76 residues, 35 to 75 residues, 35 to 74 residues, 35 to 73 residues, 35 to 72 residues, 35 to 71 residues,

35 to 70 residues, 35 to 69 residues, 35 to 68 residues, 35 to 67 residues, 35 to 66 residues, 35 to 65 residues, 35 to 64 residues, 35 to 63 residues, 35 to 62 residues, 35 to 61 residues, 35 to 60 residues, 35 to 59 residues, 35 to 58 residues, 35 to 57 residues, 35 to 56 residues, 35 to 55 residues, 35 to 54 residues, 35 to 53 residues, 35 to 52 residues, 35 to 51 residues, 35 to 50 residues, 35 to 49 residues, 35 to 48 residues, 35 to 47 residues, 35 to 46 residues, 35 to 45 residues, 35 to 44 residues, 35 to 43 residues, 35 to 42 residues, 35 to 41 residues, 35 to 40 residues, 35 to 39 residues, 35 to 38 residues, 35 to 37 residues, 35 to 36 residues or 35 residues. Preferably, the length of the at least one RNA molecule is in the range 36 to 174 residues, 37 to 173 residues, 38 to 172 residues, 39 to 171 residues, 40 to 170 residues, 41 to 169 residues, 42 to 168 residues, 43 to 167 residues, 44 to 166 residues, 45 to 165 residues, 46 to 164 residues, 47 to 163 residues, 48 to 162 residues, 49 to 161 residues, 50 to 160 residues, 51 to 159 residues, 52 to 158 residues, 53 to 157 residues, 54 to 156 residues, 36 to 74 residues, 37 to 73 residues, 38 to 72 residues, 39 to 71 residues, 40 to 70 residues, 41 to 69 residues, 42 to 68 residues, 43 to 67 residues, 44 to 66 residues, 45 to 65 residues, 46 to 64 residues, 47 to 63 residues, 48 to 62 residues, 49 to 61 residues, 50 to 60 residues, 51 to 59 residues, 52 to 58 residues, 53 to 57 residues, 54 to 56 residues.

In preferred aspects, the present invention provides a ribonucleoprotein complex, wherein the complementary portion of the at least one RNA molecule is at least 30 residues long. Alternatively, the complementary portion of the at least one RNA molecule may be 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74 or 75 residues long.

The targeting RNA molecule will preferably require a high specificity and affinity for the target nucleic acid sequence. A dissociation constant (K_d) in the range 1 μM to 1 pM, preferably 1 nM to 1pM; more preferably 1 – 100 pM is desirable as may be determined by native gel electrophoresis, or alternatively isothermal titration calorimetry, surface plasmon resonance, or fluorescence based titration methods. Affinity may be determined using an electrophoretic mobility shift assay (EMSA), also called gel retardation assay (see Semenova *et al*. (2011) PNAS 108: 10098–10103).

The targeting RNA molecule is preferably modeled on what are known from nature in prokaryotes as CRISPR RNA (crRNA) molecules. The structure of crRNA molecules is already established and explained in more detail in Jore *et al.*, 2011, Nature Structural & Molecular Biology 18: 529–537. In brief, a mature crRNA of type I-E is often 61 nucleotides long and consists of a 5' "handle" region of 8 nucleotides, the "spacer" sequence of 32 nucleotides, and a 3' sequence of 21 nucleotides which form a hairpin with a tetranucleotide loop (Fig 5). Type I systems differ from type II (Cas9) and details of different systems are described in Van der Oost 2014 Nat Rev Micr 12: 479–492. In type II (Cas9) systems there is a different processing mechanism, making use of a second RNA (tracrRNA) and two ribonucleases. Rather than a hairpin, the mature crRNA in type II remains attached to a fragment of the tracrRNA (Fig. 5). However, the RNA used in the invention does not have to be designed strictly to the design of naturally occurring crRNA, whether in length, regions or specific RNA sequences. What is clear though, is that RNA molecules for use in the invention may be designed based on gene sequence information in the public databases or newly discovered, and then made artificially, e.g. by chemical synthesis in whole or in part. The RNA molecules of the invention may also be designed and produced by way of expression in genetically modified cells or cell free expression systems and this option may include synthesis of some or all of the RNA sequence.

The structure and requirements of crRNA in type II (Cas9) has also been described in Jinek *et al.*, 2012 *ibid*. In type I, there is a so-called "SEED" portion forming the 5' end of the spacer sequence and which is flanked 5' thereto by the 5' handle of 8 nucleotides. Semenova *et al.* (2011, PNAS 108: 10098-10103), have found that all residues of the SEED sequence should be complementary to the target sequence, although for the residue at position 6, a mismatch may be tolerated (Fig. 5). In type II, there is a SEED of 10-12 nucleotides that is located at the 3' end of the spacer (Fig. 5) (reviewed by Van der Oost 2014 *ibid.*). Similarly, when designing and making an RNA component of a ribonucleoprotein complex of the invention directed at a target locus (i.e. sequence), the necessary match and mismatch rules for the type II SEED sequence can be applied.

The invention therefore includes a method of detecting and/or locating a single base change in a target nucleic acid molecule comprising contacting a nucleic acid sample

with a ribonucleoprotein complex of the invention as hereinbefore described, or with a Cas protein or polypeptide and separate targeting RNA component of the invention as hereinbefore described, and wherein the sequence of the targeting RNA (including when in the ribonucleoprotein complex) is such that it discriminates between a normal allele and a mutant allele by virtue of a single base change at, for example, position 6 of a contiguous sequence of 8 nucleotide residues.

Without wishing to be bound by a particular theory, a design rule which may be used in preparing a targeting RNA component of ribonucleoprotein complexes of the invention involves the so-called "PAM" (protospacer adjacent motif) sequence in a double stranded target polynucleotide. In the type I-E system of *E.coli*, the PAM sequence may be a conserved triplet of nucleotide residues, such as 5'-CTT-3', 5'-CAT-3', 5'-CCT-3', 5'-CAC-3', 5'-TTT-3', 5'-ATT-3', and 5'-AWG-3', wherein W is A, T or U. In Type I, a PAM sequence located in the targeted strand is usually at a position corresponding to 5' of the SEED. In Type II, however, the PAM is located at the other end, on the displaced, or non-target, strand close to the 3' end of the crRNA spacer, at a position corresponding to 3' of the seed (Fig. 5) (Jinek *et al.*, 2012, *op. cit.*). For *Streptococcus pyogenes* Cas9, the PAM sequence has a conserved pair of nucleotide residues, 5'-NGG-3'. Recently, different Cas9 variants (Type IIA and Type IIC) (Ran *et al.*, 2015 Nature 520:186-191) – Fig 1A) have been characterized, and PAMs have been revealed (see Ran *et al.*, 2015, *ibid.* – Fig 1C). Currently established Cas9 PAMs include: Type IIA 5'-NGGNNNN-3' (*Streptococcus pyogenes*), 5'-NNGTNNN-3' (*Streptococcus pasteurianus*), 5'-NNGGAAN-3' (*Streptococcus thermophilus*), 5'-NNGGGNN-3' (*Staphylococcus aureus*), and Type IIC 5'-NGGNNNN-3' (*Corynebacterium difteriae*), 5'-NNGGGTN-3' (*Campylobacter lari*), 5'-NNNCATN-3' (*Parvobaculum lavamentivorans*), 5'-NNNNGTA-3' (*Neiseria cinerea*). Cas9 of *Geobacillus thermodenitrificans* T12 (this invention) belongs to Type IIC (Ran *et al.*, 2015, *ibid.*). The inventors have surprisingly found that the choice of PAM sequences for use with the invention can influence the temperature(s) at which the Cas proteins and polypeptides of the invention will interact with a target sequence. In particular, the inventors have found a preference for an 8-mer PAM sequence to confer activity across a broad temperature range, with a cytosine in the 5th position after the 3' end of the target sequence, and/or an adenine in the 8th position. There is also a

preference for cytosine in the 1st, 2nd, 3rd, 4th and/or 6th position of the PAM sequence after the 3' end of the protospacer sequence.

In particular aspects, interaction with a target sequence within a broad temperature range of, for example 20°C to 100°C, 20°C to 80°C, 30 to 80°C, 20°C to 70°C or 25°C to 65 °C may be achieved by utilizing a PAM sequence of 5'-NNNNCVAA-3' [SEQ ID NO: 48]. There is no specific preference for the for the first 4 PAM positions. The first 4 nucleotides can therefore conveniently be any nucleotide (NNNN). Preferably, interaction with a target sequence within such a broad temperature range may be achieved by utilizing a PAM sequence of 5'-NNNNCSAA-3' [SEQ ID NO: 49]. Optimally, the PAM may be of the sequence 5'-NNNNCGAA-3' [SEQ ID NO: 50] or 5'-NNNNCCAA-3' [SEQ ID NO: 51].

Where interaction with a target sequence is required at ≥30°C, e.g. 30°C to 100°C, preferably in the range 30°C to 70°C, 30°C to 65°C, or 45°C to 65°C the PAM sequence may optimally be of the sequence 5'-NNNNCNAA-3' [SEQ ID NO: 52] or 5'-NNNNCMCA-3' [SEQ ID NO: 53]. There is no specific preference for the for the first 4 PAM positions. The first 4 nucleotides can therefore conveniently be any nucleotide (NNNN). Optionally, for example, the PAM sequence may be 5'-CCCCCNAA-3'or 5'-CCCCCMCA-3'. Optionally, for example, the PAM sequence may be selected from 5'-CCCCCAAA-3', 5'-CCCCCATA-3', 5'-CCCCCAGA-3', 5'-CCCCCACA-3', 5'-CCCCCTAA-3', 5'-CCCCCTTA-3', 5'-CCCCCTGA-3', 5'-CCCCCTCA-3', 5'-CCCCCGAA-3', 5'-CCCCCGTA-3', 5'-CCCCCGGA-3', 5'-CCCCCGCA-3', 5'-CCCCCCAA-3' [SEQ ID NO: 11], 5'-CCCCCCTA-3', 5'-CCCCCCGA-3', or 5'-CCCCCCCA-3'.

In embodiments of the invention, a targeting RNA molecule may have a length in the range of 35 – 200 residues. In preferred embodiments, the portion of the RNA which is complementary to and used for targeting a desired nucleic acid sequence is from 15 to 32 residues long. In the context of a naturally-occurring crRNA, this would correspond to the spacer portion as shown for example in figure 1 of Semenova *et al.* (2011 *ibid.*).

A ribonucleoprotein complex of the invention may have a targeting component comprising 8 residues derived from the CRISPR repeat 5' to the RNA sequence which has substantial complementarity to the DNA target sequence. The RNA sequence having complementarity to the DNA target sequence would be understood to correspond in the context of a crRNA as being the spacer sequence. The 5' flanking sequence of the RNA would be considered to correspond to the 5' handle of a crRNA; as shown for example in figure 1 of Semenova *et al.* (2011 *ibid.*).

A ribonucleoprotein complex of the invention may have a hairpin and tetranucleotide loop forming sequence 3' to the targeting RNA sequence which has complementarity to a DNA target sequence, i.e. 3' to what would correspond to the 3' handle flanking the spacer sequence in a crRNA; for example as shown in figure 1 of Semenova *et al.* (2011 *ibid.*).

Without wishing to be bound by a particular theory, in a preferred ribonucleoprotein complex and double stranded target polynucleotide, the non-target nucleic acid strand, which does not pair with the targeting RNA of the ribonucleoprotein complex, may comprise a directly 3' adjacent PAM sequence selected from one or more of 5'-NNNNCNNA-3' [SEQ ID NO: 47], 5'-CNNNCNN-3', 5'-NNNCCNN-3', 5'-NNCNCNN-3', 5'-NNNNCCN-3', and 5'-NCNNCNN-3'. Optionally, for example the PAM sequence may be selected from 5'-NNNNC-3', 5'-NNNNCNNA-3' [SEQ ID NO: 47], 5'-CNNNC-3', 5'-CNNNCNNA-3', 5'-NCNNC-3', 5'-NCNNCNNA-3', 5'-NNCNC-3', 5'-NNCNCNNA-3', 5'-NNNCC-3', 5'-NNNCCNNA-3', 5'-NNNNCC-3', 5'-NNNNCCNA-3', 5'-CCNNC-3', 5'-CCNNCNNA-3', 5'-CNCNC-3', 5'-CNCNCNNA-3', 5'-CNNCCN-3', 5'-CNNCCNNA-3', 5'-CNNNCC-3', 5'-CNNNCCNA-3', 5'-CCCNCN-3', 5'-CCCNCNNA-3', 5'-CCNCCN-3', 5'-CCNCCNNA-3', 5'-CCNNCC-3', 5'-CCNNCCNA-3', 5'-CCCCC-3' [SEQ ID NO: 12], 5'-CCCCCNNA-3' [SEQ ID NO: 13], 5'-CCCCCC-3' [SEQ ID NO: 14], 5'-CCCCCCNA-3' [SEQ ID NO: 10], 5'-NCCNC-3', 5'-NCCNCNNA-3', 5'-NCCCC-3', 5'-NCCCCNNA-3', 5'-NCCCCC-3' [SEQ ID NO: 15], 5'-NCCCCCNA-3' [SEQ ID NO: 16], 5'-NNCCC-3', 5'-NNCCCNNA-3', 5'-NNCCCC-3', 5'-NNCCCCNA-3', 5'-NNNCCC-3', and 5'-NNNCCCNA-3'. The PAM sequence may be 5'-CNCCCCAC-3' [SEQ ID NO: 17], 5'-CCCCCCAG-3' [SEQ ID NO: 18], 5'-CCCCCCAA-3' [SEQ ID NO: 11], 5'-CCCCCCAT-3' [SEQ ID NO: 19], 5'-CCCCCCAC-3' [SEQ ID NO: 20], 5'-ATCCCCAA-3' [SEQ ID NO: 21], or 5'-ACGGCCAA-3' [SEQ

ID NO: 22]. Preferably the PAM sequence will be of the sequence 5'-NNNNCNNA-3' [SEQ ID NO: 47]. However, it will be appreciated that other combinations of nucleotides may be used depending on the desired application and/or concentration of Cas protein or polypeptide. In particular, there is no specific preference for the for the first 4 PAM positions. The first 4 nucleotides can therefore conveniently be any nucleotide (NNNN). These sequences correspond to what is termed "protospacer adjacent motif" or "PAM" in the context of naturally occurring crRNAs. In type IIC CRISPR/Cas systems these PAM sequences facilitate stable interaction with the Cascade/crRNA complex with its dsDNA target, in order to ensure high degree of specificity of the crRNA – both in the natural system targets and therefore preferably also of the RNAs according to the present invention – for the target sequence. Preferably the sequence directly adjacent the protospacer will not be 5'-NNNCATN-3'.

Additionally, the PAM sequence may be of the sequence 5'-NNNNCNNA-3' [SEQ ID NO: 47], for example 5'-NNNNCNAA-3' [SEQ ID NO: 52], or 5'-NNNNCMCA-3' [SEQ ID NO: 53].

One of the limitations of the mesophilic SpCas9 is that it only displays activity between 25 and 44°C; above these temperatures SpCas9 activity rapidly decreases to undetectable levels (Mougiakos et al., 2017, ACS Synth Biol. 6: 849-861). In contrast to the 25–44°C range of its mesophilic orthologue SpCas9, ThermoCas9 of the present invention is active *in vitro* in a much broader temperature range of 20-70°C. The extended activity and stability of ThermoCas9 allows for its application in molecular biology techniques that require DNA manipulation at temperatures of 20-70°C, as well as its exploitation in harsh environments that require robust enzymatic activity. ThermoCas9 may also therefore be used as a genome editing tool for both thermophilic and mesophilic organisms.

In addition to having a broad functional temperature activity range, that is, to be functional at both low and high temperatures, for example at both 20°C and 70°C ,or 20°C and 65°C or 25°C and 65°C, the ability to manipulate the range of temperatures at which ThermoCas9 is capable of targeted cleavage or binding or at which targeted cleavage or binding takes place efficiently, by modifying structural features of

ThermoCas9 or associated elements (such as, for example, the sgRNA or tracRNA), would enable a greater level of control to be exerted over nucleic acid sequence manipulation. However, until now little was known about the molecular determinants of Cas9 activity at particular temperatures.

The inventors have identified several factors that are important for conferring the thermostability of ThermoCas9, one of which is the PAM preferences of ThermoCas9. The PAM preferences of ThermoCas9 are very strict for activity in the lower part of the temperature range (≤ 30°C), whereas more variety in the PAM is allowed for activity at the moderate to optimal temperatures (37°C to 60°C). As such, the PAM sequence may be altered to obtain the most efficient binding, cleavage, marking or modification of the target at a given temperature. This provides a great deal of flexibility in application of the ThermoCas9, depending on the particular application. For instance in some applications a very broad temperature range of target binding, cleavage, marking or modification may be desirable, for example 20°C to 70°C, preferably 20°C to 65°C or 25°C to 65°C. Binding, cleavage, marking or modification of a target sequence within such a broad temperature range may be achieved by utilizing a PAM sequence of 5'-NNNNCVAA-3' [SEQ ID NO: 48]. Preferably, binding, cleavage, marking or modification of a target sequence within such a broad temperature range may be achieved by utilizing a PAM sequence of 5'-NNNNCSAA-3' [SEQ ID NO: 49], for example 5'-NNNNCGAA-3' [SEQ ID NO: 50] or 5'-NNNNCCAA-3' [SEQ ID NO: 51]. There is no specific preference for the for the first 4 PAM positions. The first 4 nucleotides can therefore conveniently be any nucleotide (NNNN). Optionally, for example 5'-CCCCCGAA-3' or 5'-CCCCCCAA-3' [SEQ ID NO: 11].

Where binding, cleavage, marking or modification of the target is required at ≥30°C, e.g. 30°C to 100°C, preferably in the range 30°C to 70°C, 30°C to 65°C, or 45°C to 65°C the PAM sequence may optimally be of the sequence 5'-NNNNCNAA-3' [SEQ ID NO: 52] or 5'-NNNNCMCA-3' [SEQ ID NO: 53]. There is no specific preference for the for the first 4 PAM positions. The first 4 nucleotides can therefore conveniently be any nucleotide (NNNN). Optionally, for example the PAM sequence may be 5'-CCCCCNAA-3' or 5'-CCCCCMCA-3'. Optionally, for example the PAM sequence may be selected from 5'-CCCCCAAA-3', 5'-CCCCCATA-3', 5'-CCCCCAGA-3', 5'-CCCCCACA-3', 5'-CCCCCTAA-3', 5'-CCCCCTTA-3', 5'-CCCCCTGA-3', 5'-

CCCCCTCA-3', 5'-CCCCCGAA-3', 5'-CCCCCGTA-3', 5'-CCCCCGGA-3', 5'-CCCCCGCA-3', 5'-CCCCCCAA-3' [SEQ ID NO: 11], 5'-CCCCCCTA-3', 5'-CCCCCCGA-3', or 5'-CCCCCCCA-3'.

The PAM sequences of the invention provided herein comprise the sequences disclosed herein, for example as 6-mer, 7-mer or 8-mer sequences. The 6-mer, 7-mer or 8-mer sequences may begin directly 3' of the protospacer sequence on the non-target strand, with no additional nucleic acids interspaced between the protospacer sequence, complimentary to that bound by the target RNA, and the 5' end of the PAM sequence. However, it will be appreciated that there may be additional nucleic acids forming part of the PAM sequence at the 3' end of the 6-mer, 7-mer or 8-mer sequences. Additionally or alternatively, the non-target strand may comprise additional nucleic acids 3' of the PAM sequence.

A nucleoprotein complex of the invention may comprise a ribonucleoprotein complex of the invention and the target nucleic acid strand of nucleic acid, with which the ribonucleoprotein is associated.

Binding, Cleavage, Marking and Modifying Temperatures

The temperature range, including optimal temperature range of the activity, for example nuclease activity, of the Cas proteins of the present invention is significantly higher than that of known Cas9 proteins. Also, the upper extent of the range in which it retains activity is much higher than that of known Cas9 proteins. A higher optimal temperature and functional range provides a significant advantage in genetic engineering at high temperatures and therefore, for example, in the editing of the genomes of thermophilic organisms, many of which have utility in a range of industrial, agricultural and pharmaceutical processes conducted at elevated temperatures. Thus the methods, uses, nucleoproteins and transformed cells of the invention may be useful in industrial processes, for example providing genome editing for metabolic engineering purposes. The presence of the PAM sequences of the invention, directly adjacent to the protospacer sequence in the non-target strand, improve the specificity of the Cas proteins and polypeptides for the target sequences, and support the use of

the Cas proteins and polypeptides at higher temperatures and across larger functional temperature ranges.

In accordance with a significantly greater thermostability, Cas proteins of the present invention retain function, for example nuclease activity, across a much greater temperature range of than that of known Cas9 proteins. Also, the upper extent of the range in which it retains activity is much higher than that of known Cas9 proteins. A higher optimal temperature and functional range provides a significant advantage in genetic engineering at high temperatures and therefore, for example, in the editing of the genomes of thermophilic and mesophilic organisms, many of which have utility in a range of industrial, agricultural and pharmaceutical processes conducted at elevated temperatures. The extended activity and stability of ThermoCas9 allows for its application in molecular biology techniques that require DNA manipulation within a broad range of temperatures, for example 20-70°C, as well as its exploitation in harsh environments that require robust enzymatic activity. ThermoCas9 may also therefore be used as a genome editing tool for both thermophilic and mesophilic organisms.

Advantageously, the inventors have also shown that Cas proteins of the invention can also be used to direct transcriptional control of target sequences, for example silencing transcription by sequence-specific binding to target sequences. ThermoCas9 may also therefore be used as a transcriptional control tool in both thermophilic and mesophilic organisms, for example in silencing or activating transcription of target genes. ThermoCas9 may also therefore be used as a gene-silencing tool in both thermophilic and mesophilic organisms.

Advantageously, Cas proteins or polypeptides of the invention are capable of nucleic acid binding, cleavage, marking or modifying at a temperature from 20°C to 100°C but are particularly useful at elevated temperatures, for example at a temperature between 41°C and 122°C, preferably at a temperature between 50°C and 100°C. Cas proteins and polypeptides of the invention are capable of binding, cleaving, marking or modifying DNA, RNA and synthetic nucleic acids. Cas proteins or polypeptides of the invention may also provide operability for nuclease activity, gene editing and nucleic acid marking applications at temperatures in the range 20 to 50 °C, for example.

Where a temperature range is included herein, it is intended that the endpoints are included in the disclosed temperature range, i.e. that the range is "inclusive". For example, where it is stated that there is activity at a temperature in the range between 20°C and 100°C, the temperatures of 20°C and 100°C are included in said range.

Preferably, Cas proteins or polypeptides of the invention, when associated with suitable gRNA (guide RNA, also called targeting RNA molecule) which recognizes a target sequence in the polynucleotide molecule(s) to be bound, cleaved, marked or modified, does so at temperatures in the range 20°C to 100°C, optionally in the range 20°C to 70°C, 20°C to 65°C, 25°C to 70°C, 25°C to 65°C, 55°C to 100°C, 50°C to 70°C, 55°C to 70°C, or 55°C to 65°C.

Preferably, Cas proteins or polypeptides of the invention, when associated with suitable gRNA (guide RNA, also called targeting RNA molecule) which recognizes a target sequence in the polynucleotide molecule(s) to be bound, cleaved, marked or modified, does so at temperatures in the range 50°C to 100°C, optionally in the range 55°C to 100°C, 60°C to 100°C, 65°C to 100°C, 70°C to 100°C, 75°C to 100°C, 80°C to 100°C, 85°C to 100°C, 90°C to 100°C, 95°C to 100°C. More preferably, Cas proteins of the invention cleave, mark or modify nucleic acids at temperatures in the range 51°C to 99°C, 52°C to 98°C, 53°C to 97°C, 54°C to 96°C, 55°C to 95°C, 56°C to 94°C, 57°C to 93°C, 58°C to 92°C, 59°C to 91°C, 60°C to 90°C, 61°C to 89°C, 62°C to 88°C, 63°C to 87°C, 64°C to 86°C, 65°C to 85°C, 66°C to 84°C, 67°C to 83°C, 68°C to 82°C, 69°C to 81°C, 70°C to 80°C, 71°C to 79°C, 72°C to 78°C, 73°C to 77°C, 74°C to 76°C, or at a temperature of 75°C. Preferably, Cas proteins of the invention bind, cleave, mark or modify nucleic acids at temperatures in the range 60°C to 80°C, 61°C to 79°C, 62°C to 78°C, 63°C to 77°C, 64°C to 76°C, 60°C to 75°C, 60°C to 70°C. Optimally Cas proteins of the invention bind, cleave, mark or modify nucleic acids at temperatures in the range 60°C to 65°C, preferably at 65°C.

Target RNA molecules may be designed for use with the Cas proteins and polypeptides of the invention, wherein the target RNA molecules bind to the target sequence in a target strand, and the non-target strand further comprises a PAM sequence provided herein immediately 3' of the protospacer sequence. The PAM sequence may comprise 5'-NNNNNNNA-3', preferably 5'-NNNNCNNA-3' [SEQ ID

NO: 47], optionally, for example 5'-CCCCCCNA-3' [SEQ ID NO: 10] or 5'-CCCCCCAA-3' [SEQ ID NO: 11], and the uses, methods, transformed cells, and nucleoproteins of the invention may provide binding, cleaving, marking and/or modifying of the target strand across the temperature range of from 55 °C to 65 °C, preferably across the temperature range of from 50 °C to 70 °C , from 40 °C to 65 °C, from 45°C to 75 °C, from 37°C to 78°C and/or from 20°C to 80°C.

The PAM sequence may be altered to obtain the most efficient cleavage of the target at a given temperature. This provides a great deal of flexibility in application of Cas proteins of the present invention, depending on the particular application. Where binding, cleavage, marking or modifying activity, for example cleavage activity is required within a temperature range of 20°C to 100°C, preferably 20°C to 70°C, or 20°C to 65°C or 25°C to 65°C, then activity may be achieved by utilizing a PAM sequence of 5'-NNNNCVAA-3' [SEQ ID NO: 48], preferably, activity within such a temperature range may be achieved by utilizing a PAM sequence of 5'-NNNNCSAA-3' [SEQ ID NO: 49], for example 5'-NNNNCGAA-3' [SEQ ID NO: 50] or 5'-NNNNCCAA-3' [SEQ ID NO: 51]. Optionally, for example 5'-CCCCCGAA-3' [SEQ ID NO: 52] or 5'-CCCCCCAA-3' [SEQ ID NO: 11].

The inventors have found that thermostability of ThermoCas9 increases along with association of a guide (sgRNA) to form a ribonucleoprotein complex. The guide (sgRNA) may suitably comprise a tracrRNA and a crRNA. In such an arrangement, the guide may suitably comprise a crRNA which comprises a nucleotide spacer-fragment and repeat-fragment. The crRNA may suitably be 17-20 nt in length. Optionally, the crRNA may be 17 nt in length. Alternatively, the crRNA may be 18 nt in length, 19 nt in length or 20 nt in length. The guide may also comprise a tracrRNA (anti-repeat fragment (that base pairs with repeat fragment of crRNA)). The tracrRNA and crRNA can be separated by a synthetic linker. The following guide represents a preferred arrangement: 5'-[crRNA (17-20 nucleotide spacer-fragment & repeat-fragment) – (optional: synthetic loop to link the two RNAs) – tracrRNA (anti-repeat fragment (that base pairs with repeat fragment of crRNA) & some variable stem-loop structures (as to which see below), that may be truncated to some extent in some systems)]-3'.

Usually, the tracrRNA will be provided as part of a chimeric single-guide RNA (sgRNA), for example comprising a crRNA and a tracrRNA. The tracrRNA may consist of an anti-repeat region followed by one or more hairpin structures, preferably two or more hairpin structures or more preferably three or more hairpin structures. The presence of the full-length repeat/anti-repeat hairpin (formed by the 3'-end of the crRNA part (repeat) of the and the 5'-end of the complementary tracrRNA part (anti-repeat) in a synthetic sgRNA chimera fused by a 4-nucleotide linker, e.g. 5'-GAAA-3')) at the spacer distal end functions as an anchor to the nuclease, but is not essential to target selection and cleavage activity. For example deletions at the spacer distal end of up to 50-nt long deletion of the tracrRNA part can be tolerated with little to no effect on the DNA cleavage efficiency. Accordingly, for example a deletion of the spacer distal end of the full-length repeat-anti-repeat hairpin may be made up to 50 nt, up to 45 nt, up to 40nt, up to 35 nt, up to 30 nt, up to 25 nt, up to 20nt, up to 15 nt, up to 10 nt, or up to 5 nt, without compromise in terms of target DNA cleavage efficiency.

Surprisingly, the inventors have also found that the structure of the tracrRNA influences thermostability of the ThermoCas9 and efficiency of activity, in particular cleavage activity. Specifically, the number of hairpin (or stem-loop) structures in the tracrRNA or the sgRNA can be modified in order to obtain the most efficient binding, cleavage, marking or modifying of the target at a given temperature. This provides a great deal of flexibility in application of Cas proteins of the present invention, depending on the particular application. Optionally, the tracrRNA or the sgRNA may be provided with a nucleic acid sequence which is capable of forming one or more stem-loop structures, two or more stem-loop structures or three or more stem-loop structures. Optionally, the tracrRNA or the sgRNA may be provided with a nucleic acid sequence which is arranged to form one or more stem-loop structures, two or more stem-loop structures or three or more stem-loop structures. Preferably, the sgRNA will be provided with a nucleic acid sequence which is capable of forming at least three stem-loop structures.

Optionally where binding, cleavage, marking or modifying activity, for example cleavage activity is required within a temperature range of 20°C to 60°C, preferably 37°C to 60°C, or 37°C, 40°C, 45°C, 50°C, 55°C or 60°C then activity may be achieved

by utilizing a sgRNA sequence which is capable of forming one or more stem-loop structures.

Optionally where binding, cleavage, marking or modifying activity, for example cleavage activity is required within a temperature range of 20°C to 65°C, preferably 37°C to 65°C, more preferably 45°C to 55°C or 37°C, 40°C, 45°C, 50°C, 55°C or 60°C then activity may be achieved by utilizing a sgRNA sequence which is capable of forming two or more stem-loop structures.

Optionally where binding, cleavage, marking or modifying activity, for example cleavage activity is required within a temperature range of 20°C to 100°C, preferably 20°C to 70°C, more preferably 37°C to 65°C or 37°C, 40°C, 45°C, 50°C, 55°C, 60°C or 65°C then activity may be achieved by utilizing a sgRNA sequence which is capable of forming three or more stem-loop structures.

Preferably, the portion of the sgRNA corresponding to the tracrRNA will comprise the sequence; AAGGGCUUUCUGCCUAUAGGCAGACUGCCC [SEQ ID NO: 54] which exemplifies the 5' hairpin. Preferably, the portion of the sgRNA corresponding to the tracrRNA will further comprise the sequence; GUGGCGUUGGGGAUCGCCUAUCGCC [SEQ ID NO: 55], which exemplifies the 'middle' hairpin. Preferably, the portion of the sgRNA corresponding to the tracrRNA will further comprise the sequence; CGCUUUCUUCGGGCAUUCCCCACUCUUAGGCGUUUU [SEQ ID NO: 56], which exemplifies the 3' hairpin.

Preferably, the portion of the sgRNA corresponding to the tracrRNA will comprise the sequence;
AAGGGCUUUCUGCCUAUAGGCAGACUGCCCGUGGCGUUGGGGAUCGCCUAU CGCC [SEQ ID NO: 57] i.e. including the 5' hairpin and the middle hairpin.

Preferably, the portion of the sgRNA corresponding to the tracrRNA may comprise the sequence;
AAGGGCUUUCUGCCUAUAGGCAGACUGCCCGUGGCGUUGGGGAUCGCCUAU CGCCCGCUUUCUUCGGGCAUUCCCCACUCUUAGGCGUUUU [SEQ ID NO: 58] i.e. including the 5' hairpin, the middle hairpin and the 3' hairpin.

The inventors have discovered that the number of predicted stem-loops of the tracrRNA scaffold plays a crucial role in DNA cleavage, in particular at elevated temperatures. They have determined that, although the presence of three stem-loops of the tracrRNA scaffold is not essential for cleavage activity, when all three loops are present, the cleavage is most efficient at all temperatures in the range indicating that a full length tracrRNA is required for optimal ThermoCas9-based DNA cleavage at elevated temperatures. In contrast, removal of the 3' hairpin results in a decrease in efficiency of cleavage. Moreover, the inventors found that removal of both the middle and the 3' hairpins, results in a drastically decline in the cleavage efficiency of ThermoCas9, particularly at the upper and lower extremes of the functional temperature ranges. Preferably, where binding, cleavage, marking or modification of a target sequence is required at elevated temperatures, for example 45°C to 100°C, 50°C to 100°C, 50°C to 70°C, 50°C to 65°C, 55°C to 65°C or within a broad temperature range such as 20°C to 100°C, 20°C to 70°C, 20°C to 65°C. Preferably, ThermoCas9 associated with a sgRNA with three stem-loop structures will remain stable and capable of binding, cleavage, marking or modification of a target sequence for at least 1 min, at least 2 min, at least 3 min, at least 4 min or at least 5 min, preferably 5min at a selected temperature in the range 20°C to 100°C, 20°C to 70°C, 20°C to 65°C, 45°C to 100°C, 50°C to 100°C, 50°C to 70°C, 50°C to 65°C or 55°C to 65°C.

Additionally, the inventors have also discovered that the length of the spacer sequence of the sgRNA can be varied in order to manipulate the efficiency of ThermoCas9 activity, for example binding, cleavage, marking or modifying activity. Typically, the spacer sequence will be in the range 18nt to 25nt in length. Optionally the spacer sequence will be 18nt, 19nt, 20nt, 21nt, 22nt, 23nt, 24nt or 25nt in length. Preferably, spacer lengths of 19nt, 20nt, 21nt or 23nt will be used, since Cas9 proteins of the invention cleave target sequences with the highest efficiency when associated with sgRNAs with these spacer lengths. The cleavage efficiency drops significantly when a spacer of 18nt is used. Preferably, the length of the spacer will be 23nt.

In all aspects of the invention, Cas proteins or polypeptides may be obtained or derived from bacteria, archaea or viruses; or alternatively may be synthesised *de novo*. In

preferred embodiments, a Cas protein or polypeptide of the invention is derived from a thermophilic prokaryotic organism, which may be classified as an archaea or bacterium, but is preferably a bacterium. More preferably a Cas protein or polypeptide of the invention will be derived from a thermophilic bacterium. Herein, the term thermophilic is to be understood as meaning capable of survival and growth at relatively high temperatures, for example in the context of the invention, capable of nucleic acid cleavage, binding or modification at a temperature between 41 and 122 °C (106 and 252 °F). Preferably a Cas protein or polypeptide of the invention may be isolated from one or more thermophilic bacteria and will function above 60°C. Preferably a Cas protein or polypeptide of the invention may be isolated from one or more thermophilic bacteria and will function in the range 60°C to 80°C and optimally between 60°C and 65°C. In preferred embodiments, a Cas protein or polypeptide of the invention is derived from *Geobacillus* sp. More preferably, a Cas protein of the invention is derived from *Geobacillus thermodenitrificans*. Even more preferably, a Cas protein of the invention is derived from *Geobacillus thermodenitrificans* T12. A Cas protein or polypeptide of the invention may be derived from a virus.

Functional Moieties

Advantageously, the ability of Cas proteins, polypeptides and ribonucleoprotein complexes of the invention to target any polynucleotide sequence in a sequence-specific manner may be exploited in order to modify the target nucleic acid in some way, for example by cleaving it and/or marking it and/or modifying it. It will therefore be appreciated that additional proteins may be provided along with the Cas protein or polypeptide to achieve this. Accordingly, the Cas proteins or polypeptides of the invention may further comprise at least one functional moiety and/or the Cas proteins, polypeptides or ribonucleoprotein complexes of the present invention may be provided as part of a protein complex comprising at least one further protein. In a preferred aspect the present invention provides a Cas protein, polypeptide or a ribonucleoprotein complex wherein the Cas protein or at least one further protein further comprises at least one functional moiety. The at least one functional moiety may be fused or linked to the Cas protein. Preferably, the at least one functional moiety may be translationally fused to the Cas protein through expression in natural or artificial protein expression systems. Alternatively, the at least one functional moiety

may be covalently linked by a chemical synthesis step to the Cas protein. Preferably, the at least one functional moiety is fused or linked to the N-terminus and/or the C-terminus of the Cas protein; preferably the C-terminus.

Desirably, the at least one functional moiety will be a protein. It may be a heterologous protein or alternatively may be native to the bacterial species from which the Cas protein was derived. The at least one functional moiety may be a protein; optionally selected from a helicase, a nuclease, a helicase-nuclease, a DNA methylase, a histone methylase, an acetylase, a phosphatase, a kinase, a transcription (co-)activator, a transcription repressor, a DNA binding protein, a DNA structuring protein, a marker protein, a reporter protein, a fluorescent protein, a ligand binding protein, a signal peptide, a subcellular localisation sequence, an antibody epitope or an affinity purification tag.

In a particularly preferred aspect, the present invention provides a Cas protein, polypeptide, or a ribonucleoprotein complex, wherein the at least one functional moiety is a marker protein, for example GFP.

Nuclease Activity

A Cas ribonucleoprotein of the invention has nucleic acid binding, cleavage, marking or modification activity at a temperature, preferably an elevated temperature, disclosed herein, for example at a temperature between 50°C and 100°C. The ribonucleoproteins of the invention may be capable of binding, cleaving, marking or modifying DNA, RNA or synthetic nucleic acids. In preferred aspects Cas ribonucleoproteins of the invention are capable of cleaving DNA in a sequence-specific manner, in particular double-stranded DNA.

Cas proteins, polypeptides or ribonucleoproteins of the invention may have more than one nuclease domain. Site-specific nucleases can permit the generation of double strand breaks (DSBs) at selected positions along a strand of DNA. In a target host cell, this enables DSBs to be made at specific pre-selected positions in the genome. The creation of such breaks by site-specific nucleases prompts the endogenous

cellular repair machinery to be repurposed in order to insert, delete or modify DNA at desired positions in the genome of interest.

One or more nuclease activity sites of the protein or polypeptide molecule may be inactivated, e.g. so as to allow the activity of another functional moiety linked or fused to the protein or polypeptide, e.g. a nuclease domain such as Fok1 nuclease.

Therefore notwithstanding the fact that the Cas proteins, polypeptides and ribonucleoproteins of the invention may have endogenous nuclease activity, for certain applications it may be desirable to inactivate the native nuclease activity of the Cas protein and provide a Cas protein or a ribonucleoprotein complex wherein the native Cas9 nuclease activity is inactivated and the Cas protein is linked to at least one functional moiety. Reducing the incidence of mis-targeting events by complementation of the native Cas9 nuclease activity is one such application. This may desirably be achieved by inactivation of the native Cas9 nuclease activity of the Cas protein or ribonucleoprotein complex and provision of a heterologous nuclease, preferably fused to the Cas protein. Accordingly, the present invention provides a Cas protein or a ribonucleoprotein complex, wherein the at least one functional moiety is a nuclease domain, preferably a FokI nuclease domain. In a particularly preferred aspect, the Cas protein or ribonucleoprotein complex of the invention fused to a FokI nuclease domain is provided as part of a protein complex, preferably comprising another Cas protein or ribonucleoprotein complex of the invention fused to a FokI nuclease domain and wherein the two complexes target opposite strands of the target genomic DNA.

For some applications it may be desirable to completely attenuate the nuclease activity of the Cas protein, polypeptide or ribonucleoprotein, for example in applications where the Cas protein or ribonucleoprotein complex is utilised to recognise and modify a specific target sequence in a nucleic acid, for instance to mark it as part of a diagnostic test. In such applications, the nuclease activity of the Cas protein may be inactivated and the functional moiety fused to the Cas protein may be a protein; optionally selected from a helicase, a nuclease, a helicase-nuclease, a DNA methylase, a histone methylase, an acetylase, a phosphatase, a kinase, a transcription (co-)activator, a transcription repressor, a DNA binding protein, a DNA structuring protein, a marker

protein, a reporter protein, a fluorescent protein, a ligand binding protein, a signal peptide, a subcellular localisation sequence, an antibody epitope or an affinity purification tag.

In a preferred aspect, a catalytically inactive, or "dead" Cas protein or polypeptide (dCas) lacking nuclease activity may be bound to a target nucleic acid sequence and thereby sterically repress activity of that sequence. For example, a target RNA may be designed that is complementary to a promoter or exonic sequence of a gene, so that binding of the dCas and target RNA to the gene sterically represses transcriptional initiation or elongation of the gene sequence, thereby repressing expression of the gene. Alternatively, the methods and uses described herein can use modified nuclease variants of gtCas9 that are nickases. A nickase can be created via a mutation in either one of the HNH or the RuvC catalytic domains of the gtCas9 nuclease. This has been shown for *S. pyogenes* Cas9 (spCas) with spCas9-mutants D10A and H840A, which have an inactive RuvC or HNH nuclease domain, respectively. The combination of these two mutations leads to a catalytically dead Cas9 variant (Standage-Beier, K. et al., 2015, ACS Synth. Biol. 4, 1217–1225; Jinek, M. et al., 2012, Science 337, 816– 821; Xu, T. et al., 2015, Appl. Environ. Microbiol. 81, 4423–4431). Based on sequence homology (Figure 3), these residues can be D8 (D17 in Figure 3) and D581 or H582 (Figure 3) in gtCas9.
Preferably, the mutations D8A and H582A in gtCas9 (ThermoCas9) can be used to create a catalytically inactive, or "dead" Cas protein or polypeptide variant of ThermoCas9 (dCas) which lacks nuclease activity. Such a dCas may usefully find application as, for example, an efficient thermoactive transcriptional silencing CRISPRi tool, being able to steadily and specifically bind to DNA elements without introducing dsDNA breaks. Advantageously, such a system could, amongst other things, greatly facilitate metabolic studies of thermophiles.

In a particularly preferred aspect, the present invention provides a Cas protein or a ribonucleoprotein complex, wherein the nuclease activity of the Cas protein is inactivated and the at least one functional moiety is a marker protein, for example GFP. In this way it may be possible to specifically target a nucleic acid sequence of interest and to visualize it using a marker which generates an optical signal. Suitable markers may include for example, a fluorescent reporter protein, e.g. Green

Fluorescent Protein (GFP), Yellow Fluorescent Protein (YFP), Red Fluorescent Protein (RFP), Cyan Fluorescent Protein (CFP) or mCherry. Such a fluorescent reporter gene provides a suitable marker for visualisation of protein expression since its expression can be simply and directly assayed by fluorescence measurement. Alternatively, the reporter nucleic acid may encode a luminescent protein, such as a luciferase (e.g. firefly luciferase). Alternatively, the reporter gene may be a chromogenic enzyme which can be used to generate an optical signal, e.g. a chromogenic enzyme (such as beta-galactosidase (LacZ) or beta-glucuronidase (Gus)). Reporters used for measurement of expression may also be antigen peptide tags. Other reporters or markers are known in the art, and they may be used as appropriate.

Because the marker may be visualized, in certain embodiments where the target nucleic acid is RNA, specifically mRNA, it may be possible to quantify the transcriptional activity of a gene by detection and quantification of the optical signal provided by the marker, particularly where the optical signal generated by the marker is directly proportionate to the quantity of the expression product. Therefore in preferred embodiments of the invention, Cas proteins or ribonucleoproteins of the invention may be used to assay expression products of a gene of interest.

In one aspect, the gtCas9 described herein may be used in a homologous recombination (HR) mediated genome modification method in microbial cells. Such methods involve HR and site-directed gtCas9 activity, whereby counter selection occurs by the gtCas9 activity removing microbes which do not have a desired modification introduced by HR.

Thus the methods and uses provided herein allow the process of homologous recombination to be favoured during a first step such that the microbial genome can be modified with the desired mutation and a second step in which unmodified cells can be targeted by the gtCas9 ribonuclease complex to introduce a DSDB into the genomes of the unmodified cells. Due to an absence of an efficient non-homologous end joining (NHEJ) repair mechanism in the majority of microbes, DSDB typically leads to cell death. Thus, these methods and uses increase overall the population of microbial cells with the desired mutation whilst eliminating any unmodified microbial

cells. Preferably, such methods and uses are used in microbes that have substantially no endogenous NHEJ repair mechanism. Alternatively, the methods and uses may be applied to microbes that have an endogenous NHEJ repair mechanism. The methods and uses described herein may be applied to microbes that have an endogenous NHEJ repair mechanism but wherein the NHEJ repair mechanism is either conditionally reduced or the NHEJ activity is knocked out.

The methods and uses provided herein may utilise a sequence of the homologous recombination polynucleotide that has at least one mis-match with the guide RNA, such that the guide RNA is no longer able to recognise the modified genome. This means that the gtCas9 ribonuclease complex will not recognise the modified genome. Therefore, no DSDB can be introduced by the gtCas9 ribonuclease complex and so the modified cells will survive. However, the cells with unmodified genomes will still have substantial complementarity to the guide RNA and consequently can be cleaved site-specifically by the gtCas9 ribonuclease complex.

In another aspect of the methods and uses of the invention, the way in which the gtCas9 ribonucleoase complex is prevented from acting to cleave the microbial genome is not so much to modify or eliminate the sequence targeted by the guide, but rather the PAM required by the gtCas9 ribonuclease complex. The PAM is either modified or eliminated in order to blind the gtCas9 ribonuclease complex to the specific cutting site. Therefore, methods and uses of the invention may include those using a sequence of the homologous recombination polynucleotide that does not include a PAM sequence recognised by the gtCas9 ribonuclease complex. Therefore, no DSDB can be introduced by the gtCas9 ribonuclease complex and so the HR modified cells will survive. However, the unmodified cells will still be recognised by the gtCas9 ribonuclease complex and its guide and so consequently are cleaved site-specifically.

Thus methods and uses are provided herein that rely on HR to modify the genome of the microbe. Preferably, the upstream flank and downstream flanks are 0.5 kilobases (kb) to 1.0 kb each in length. However, recombination using larger or shorter fragments is possible as well. The homologous recombination polynucleotide may further comprise a polynucleotide sequence between the upstream and downstream

flanking regions. This polynucleotide sequence could for example contain a modification that is to be introduced into the microbial genome.

Whilst homologous recombination relies upon the upstream and downstream flanks having substantial complementarity to the target regions, mismatches can be accommodated as well. Therefore, in some embodiments, homologous recombination is known to occur between DNA segments with extensive homology to the upstream and downstream flanks. In alternative embodiments, the upstream and downstream flanks have complete complementarity to the target regions. The upstream and downstream flanks need not be identical in size. However, in some instances the upstream and downstream flanks are identical in size. The efficiency of homologous recombination will vary depending on the likelihood of homologous recombination of the smallest fragment length of the flank. However, even if the homologous recombination process is inefficient, advantageously the method described herein will select for any microbial cell that has the desired modification over the unmodified microbial cell. Homologous recombination also allows large deletions (e.g. 50 kb or more) to be made encompassing complete gene clusters. Homologous recombination is also used for recombineering, which is a well-known method to allow for recombination over smaller fragments (45-100 nt). The methods and uses described herein can optionally further comprise at least another homologous recombination polynucleotide or a polynucleotide comprising a sequence encoding a homologous recombination polynucleotide having a sequence substantially complementary to a second target region containing the target in the microbial genome.

In preferred embodiments, the methods and uses described herein utilise a homologous recombination polynucleotide that is DNA. In some embodiments the DNA is single stranded. In other embodiments, the DNA is double stranded. In further embodiments, the DNA is double stranded and plasmid borne.

HR in the methods and uses provided herein may be used to remove a polynucleotide sequence from the microbial genome. Alternatively, HR in the methods and uses provided herein may be used to insert one or more gene(s), or fragment(s) thereof, in to the microbial genome. As a further alternative, HR in the methods and uses

provided herein may be used to modify or replace at least one nucleotide in the microbial genome. Consequently, the methods and uses provided herein may be used for any desired kind of genome modification.

Alternatively, the gtCas9 described herein may be used in a HR mediated genome modification method in microbial cells, whereby the gtCas9 activity introduces DSDB and can induce cellular HR in microbial cells, as has been shown for spCas9 (Jiang *et al.* (2013) Nature Biotech, 31, 233-239; Xu *et al.* (2015) Appl Environ Microbiol, 81, 4423-4431; Huang *et al*. (2015) Acta Biochimica et Biophysica Sinica, 47, 231-243). .

Alternatively, homologous recombination may be facilitated through recombineering, *e.g.*, by introducing an oligonucleotide into a microbial cell expressing a gene coding for RecT or beta protein as reviewed by Mougiakos *et al*. ((2016), Trends Biotechnol. 34: 575-587). In a further embodiment, the Cas9 can be combined with Multiplex Automated Genome Engineering (MAGE) as exemplified by Ronda *et al*. ((2016), Sci. Rep. 6: 19452.)

Throughout, the reference sequences of the Cas proteins of the invention may be defined as a nucleotide sequence encoding the amino acid sequence. For example the amino acid sequence of the motifs defined in SEQ ID's 2 to 6 also includes all nucleic acid sequences which encode that amino acid sequence.

Accordingly, the present invention also provides an isolated nucleic acid molecule encoding a Cas protein comprising;

a. the amino acid motif EKDGKYYC [SEQ ID NO: 2]; and/or
b. the amino acid motif $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and/or
c. the amino acid motif X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and/or

d. the amino acid motif X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and/or
e. the amino acid motif $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine;

wherein the Cas protein is capable of DNA binding, cleavage, marking or modification between 50°C and 100°C when associated with at least one targeting RNA molecule, and a polynucleotide comprising a target nucleic acid sequence recognised by the targeting RNA molecule.

In another aspect the present invention also provides an isolated nucleic acid molecule encoding a clustered regularly interspaced short palindromic repeat (CRISPR)-associated (Cas) protein having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 77% identity therewith.

In another aspect the present invention also provides an isolated nucleic acid molecule, further comprising at least one nucleic acid sequence encoding a peptide which upon translation is fused to the Cas protein.

In another aspect the present invention also provides an isolated nucleic acid molecule, wherein the at least one nucleic acid sequence fused to the nucleic acid molecule encoding the Cas protein encodes a protein selected from a helicase, a nuclease, a helicase-nuclease, a DNA methylase, a histone methylase, an acetylase, a phosphatase, a kinase, a transcription (co-)-activator, a transcription repressor, a DNA binding protein, a DNA structuring protein, a marker protein, a reporter protein, a fluorescent protein, a ligand binding protein, a signal peptide, a subcellular localisation sequence, an antibody epitope or an affinity purification tag.

ThermoCas9 Nuclease Activity: Divalent Cations

Previously characterized, mesophilic Cas9 endonucleases employ divalent cations to catalyze the generation of DSBs in target DNA. The inventors have shown that ThermoCas9 can mediate dsDNA cleavage in the presence of any of the following divalent cations: Mg^{2+}, Ca^{2+}, Mn^{2+}, Co^{2+}, Ni^{2+}, and Cu^{2+}.

ThermoCas9 Nuclease Activity: Substrates

The inventors have also surprisingly shown that despite reports that certain type-IIC systems were efficient single stranded DNA cutters ((Ma, *et al.*, *Mol. Cell* 60, 398–407 (2015); Zhang, *et al.*, *Mol. Cell* 60, 242–255 (2015)), ThermoCas9 cannot direct cleavage of ssDNA. The nuclease activity of ThermoCas9 is limited to dsDNA substrates.

Expression Vectors

Nucleic acids of the present invention may be isolated. However, in order that expression of the nucleic acid sensing construct may be carried out in a chosen cell, the polynucleotide sequence encoding the Cas protein or ribonucleoprotein will preferably be provided in an expression construct. In some embodiments, the polynucleotide encoding the Cas protein or ribonucleoprotein will be provided as part of a suitable expression vector. In certain embodiments an expression vector of the present invention (with or without nucleotide sequence encoding amino acid residues which on expression will be fused to a Cas protein) may further comprise a nucleotide sequence encoding a targeting RNA molecule as hereinbefore defined. Consequently, such expression vectors can be used in an appropriate host to generate a ribonucleoprotein complex of the invention which can target a desired nucleotide sequence. Alternatively, nucleotide sequences encoding a targeting RNA molecule as hereinbefore defined may be provided in a separate expression vector or alternatively may be delivered to a target cell by other means.

Suitable expression vectors will vary according to the recipient cell and suitably may incorporate regulatory elements which enable expression in the target cell and preferably which facilitate high-levels of expression. Such regulatory sequences may be capable of influencing transcription or translation of a gene or gene product, for example in terms of initiation, accuracy, rate, stability, downstream processing and mobility.

Such elements may include, for example, strong and/or constitutive promoters, 5' and 3' UTR's, transcriptional and/or translational enhancers, transcription factor or protein binding sequences, start sites and termination sequences, ribosome binding sites, recombination sites, polyadenylation sequences, sense or antisense sequences, sequences ensuring correct initiation of transcription and optionally poly-A signals ensuring termination of transcription and transcript stabilisation in the host cell. The regulatory sequences may be plant-, animal-. bacteria-, fungal- or virus derived, and preferably may be derived from the same organism as the host cell. Clearly, appropriate regulatory elements will vary according to the host cell of interest. For example, regulatory elements which facilitate high-level expression in prokaryotic host cells such as in *E. coli* may include the pLac, T7, P(Bla), P(Cat), P(Kat), trp or tac promoters. Regulatory elements which facilitate high-level expression in eukaryotic host cells might include the AOX1 or GAL1 promoter in yeast or the CMV- or SV40-promoters, CMV-enhancer, SV40-enhancer, *Herpes simplex virus* VIP16 transcriptional activator or inclusion of a globin intron in animal cells. In plants, constitutive high-level expression may be obtained using, for example, the *Zea mays* ubiquitin 1 promoter or 35S and 19S promoters of cauliflower mosaic virus.

Suitable regulatory elements may be constitutive, whereby they direct expression under most environmental conditions or developmental stages, developmental stage specific or inducible. Preferably, the promoter is inducible, to direct expression in response to environmental, chemical or developmental cues, such as temperature, light, chemicals, drought, and other stimuli. Suitably, promoters may be chosen which allow expression of the protein of interest at particular developmental stages or in response to extra- or intra-cellular conditions, signals or externally applied stimuli. For example, a range of promoters exist for use in *E. coli* which give high-level expression at particular stages of growth (e.g. osmY stationary phase promoter) or in response to particular stimuli (e.g. HtpG Heat Shock Promoter).

Suitable expression vectors may comprise additional sequences encoding selectable markers which allow for the selection of said vector in a suitable host cell and/or under particular conditions.

The invention also includes a method of modifying a target nucleic acid in a cell, comprising transfecting, transforming or transducing the cell with any of the expression vectors as hereinbefore described. The methods of transfection, transformation or transduction are of the types well known to a person of skill in the art. Where there is one expression vector used to generate expression of a ribonucleoprotein complex of the invention and when the targeting RNA is added directly to the cell then the same or a different method of transfection, transformation or transduction may be used. Similarly, when there is one expression vector being used to generate expression of a ribonucleoprotein complex of the invention and when another expression vector is being used to generate the targeting RNA *in situ* via expression, then the same or a different method of transfection, transformation or transduction may be used.

In other embodiments, mRNA encoding the Cas protein or polypeptide is introduced into a cell so that the Cascade complex is expressed in the cell. The targeting RNA which guides the Cas protein complex to the desired target sequence is also introduced into the cell, whether simultaneously, separately or sequentially from the mRNA, such that the necessary ribonucleoprotein complex is formed in the cell.

Accordingly, the invention also provides a method of modifying, i.e. cleaving, tagging, modifying, marking or binding, a target nucleic acid comprising contacting the nucleic acid with a ribonucleoprotein complex as hereinbefore defined.

In addition, the invention also includes a method of modifying a target nucleic acid comprising contacting the nucleic acid with a Cas protein or polypeptide as hereinbefore defined, in addition to a targeting RNA molecule as hereinbefore defined.

In accordance with the above methods, modification of target nucleic acid may therefore be carried out *in vitro* and in a cell-free environment. In a cell-free environment, addition of each of the target nucleic acid, the Cas protein and the targeting RNA molecule may be simultaneous, sequential (in any order as desired), or separately. Thus it is possible for the target nucleic acid and targeting RNA to be added simultaneously to a reaction mix and then the Cas protein or polypeptide of the invention to be added separately at a later stage.

Equally, the modification of the target nucleic acid may be made *in vivo*, that is *in situ* in a cell, whether an isolated cell or as part of a multicellular tissue, organ or organism. In the context of whole tissue and organs, and in the context of an organism, the method may desirably be carried out *in vivo* or alternatively may be carried out by isolating a cell from the whole tissue, organ or organism, treating the cell with ribonucleoprotein complex in accordance with the method and subsequently returning the cell treated with ribonucleoprotein complex to its former location, or a different location, whether within the same or a different organism.

In these embodiments, the ribonucleoprotein complex or the Cas protein or polypeptide requires an appropriate form of delivery into the cell. Such suitable delivery systems and methods are well known to persons skilled in the art, and include but are not limited to cytoplasmic or nuclear microinjection. In preferred modes of delivery, an Adeno-associated virus (AAV) is used; this delivery system is not disease causing in humans and has been approved for clinical use in Europe.

Accordingly the present invention provides a method of modifying a target nucleic acid comprising contacting the nucleic acid with:

a. a ribonucleoprotein complex as hereinbefore defined; or

b. a protein or protein complex as hereinbefore defined and an RNA molecule as hereinbefore defined.

In a further aspect the present invention provides a method of modifying a target nucleic acid in a cell, comprising transforming, transfecting or transducing the cell with an expression vector comprising nucleotide sequences encoding a ribonucleoprotein complex as hereinbefore defined; or alternatively transforming, transfecting or transducing the cell with an expression vector comprising nucleotide sequences encoding a protein or protein complex as hereinbefore defined and a further expression vector comprising a nucleotide sequence encoding a targeting RNA molecule as hereinbefore defined.

In a further aspect, the present invention provides a method of modifying a target nucleic acid in a cell comprising transforming, transfecting or transducing the cell with an expression vector comprising nucleotide sequences encoding a protein or protein

complex as hereinbefore defined, and then delivering a targeting RNA molecule as hereinbefore defined into the cell.

In embodiments where the guide (i.e. targeting) RNA (gRNA) molecule and the Cas protein or polypeptide are provided separately rather than as part of a ribonucleoprotein complex, the gRNA molecule requires an appropriate form of delivery into a cell, whether simultaneously, separately or sequentially with the Cas protein or protein complex. Such forms of introducing RNA into cells are well known to a person of skill in the art and may include in vitro or ex vivo delivery via conventional transfection methods. Physical methods, such as microinjection and electroporation, as well as calcium co-precipitation, and commercially available cationic polymers and lipids, and cell-penetrating peptides, cell-penetrating (biolistic) particles may each be used. For example, viruses, particularly preferred is AAV, may be used as delivery vehicles, whether to the cytoplasm and/or nucleus, for example via the (reversible) fusion of Cas protein complex of the invention or a ribonucleoprotein complex of the invention to the viral particle.

In another aspect the present invention provides a method of modifying a target nucleic acid, wherein the at least one functional moiety is a marker protein or reporter protein and the marker protein or reporter protein associates with the target nucleic acid; preferably wherein the marker is a fluorescent protein, for example a green fluorescent protein (GFP).

In the aforementioned methods of modifying a target nucleic acid, the functional moiety may be a marker and the marker associates with the target nucleic acid; preferably wherein the marker is a protein; optionally a fluorescent protein, e.g. green fluorescent protein (GFP), yellow fluorescent protein (YFP), red fluorescent protein (RFP) or mCherry. Whether *in vitro*, *ex vivo* or *in vivo*, then methods of the invention can be used to directly visualise a target locus in a nucleic acid molecule, preferably in the form of a higher order structure such as a supercoiled plasmid or chromosome, or a single stranded target nucleic acid such as mRNA. Direct visualisation of a target locus may use electron micrography, or fluorescence microscopy. However, it will be appreciated that in the context of methods of the invention, other kinds of label may

be used as the marker including organic dye molecules, radiolabels and spin labels which may be small molecules.

In methods of the invention for modifying a target nucleic acid wherein the target nucleic acid is dsDNA, the functional moiety may be a nuclease or a helicase-nuclease, and the modification is preferably a single stranded or a double stranded break at a desired locus. In this way unique sequence specific cutting of DNA can be engineered by using a suitable functional moiety fused to a ribonucleoprotein complex. The chosen sequence of the RNA component of the final ribonucleoprotein complex provides the desired sequence specificity for the action of the functional moiety.

Therefore, the invention also provides a method of non-homologous end joining of a dsDNA molecule in a cell at a desired locus to remove at least a part of a nucleotide sequence from the dsDNA molecule; optionally to knockout the function of a gene or genes, wherein the method comprises making double stranded breaks using any of the methods of modifying a target nucleic acid as hereinbefore described.

The invention further provides a method of homologous recombination of a nucleic acid into a dsDNA molecule in a cell at a desired locus in order to modify an existing nucleotide sequence or insert a desired nucleotide sequence, wherein the method comprises making a double stranded break at the desired locus using any of the methods of modifying a target nucleic acid as hereinbefore described.

The invention therefore also provides a method of modifying gene expression in an organism comprising modifying a target nucleic acid sequence according to any of the methods hereinbefore described, wherein the nucleic acid is dsDNA and the functional moiety is selected from a DNA modifying enzyme (e.g. a methylase or acetylase), a transcription activator or a transcription repressor.

The invention additionally provides a method of modifying gene expression in an organism comprising modifying a target nucleic acid sequence according to any of the methods hereinbefore described, wherein the nucleic acid is an mRNA and the functional moiety is a ribonuclease; optionally selected from an endonuclease, a 3' exonuclease or a 5' exonuclease.

The target nucleic acid may be DNA, RNA or synthetic nucleic acid. Preferably the target nucleic acid is DNA; preferably dsDNA.

However, the target nucleic acid can be RNA; preferably mRNA. Alternatively therefore, the present invention also provides methods of modifying a target nucleic acid, wherein the target nucleic acid is RNA.

In another aspect the present invention provides a method of modifying a target nucleic acid, wherein the nucleic acid is dsDNA, the at least one functional moiety is a nuclease or a helicase-nuclease, and the modification is a single- stranded or a double- stranded break at a desired locus.

In another aspect the present invention provides a method of modifying a target nucleic acid in a cell, wherein modification results in a silencing of gene expression at a desired locus; and wherein the method includes the steps of;
a. making double-stranded breaks in the dsDNA molecule; and
b. repair of the dsDNA molecule in the cell by non-homologous end joining (NHEJ).

In another aspect the present invention provides a method of modifying a target nucleic acid in a cell; wherein the existing nucleotide sequence is modified or deleted and/or a desired nucleotide sequence is inserted at a desired location wherein the method includes the steps of;
a. making a double stranded break at the desired locus; and
b. repair of the dsDNA molecule in the cell by homologous recombination.

In another aspect the present invention provides a method of modifying gene expression in a cell comprising modifying a target nucleic acid sequence as hereinbefore described; wherein the nucleic acid is dsDNA and the functional moiety is selected from a DNA modifying enzyme (*e.g.* a methylase or acetylase), a transcription activator or a transcription repressor.

In another aspect the present invention provides a method of modifying gene expression in a cell comprising modifying a target nucleic acid sequence as

hereinbefore described, wherein the nucleic acid is an mRNA and the functional moiety is a ribonuclease; optionally selected from an endonuclease, a 3' exonuclease or a 5' exonuclease.

In another aspect the present invention provides a method of modifying a target nucleic acid as hereinbefore described, wherein the method is carried out at a temperature between 45°C and 100°C. Preferably, the method is carried out at a temperature at or above 50°C. More preferably, the method is carried out at a temperature between 55°C and 80°C. Optimally, the method is carried out at a temperature between 60°C and 65°C. Alternatively, the method may be carried out at a temperature between 20°C and 45°C. More preferably, at a temperature between 30°C and 45°C. Even more preferably at a temperature between 37°C and 45°C.

In any of the methods of modifying a target nucleic acid hereinbefore described, the cell may be a prokaryotic cell or alternatively, may be a eukaryotic cell.

Host Cells

Advantageously, the present invention is of broad applicability and host cells of the present invention may be derived from any genetically tractable organism which can be cultured. Accordingly, the present invention provides a host cell transformed by a method as hereinbefore described. The invention provides a transformed cell, having a target nucleic acid sequence in a double stranded target polynucleotide, said cell comprising a Cas protein or polypeptide as provided herein and at least one targeting RNA molecule as provided herein, and an expression vector comprising a nucleic acid encoding at least one of said Cas protein and said targeting RNA molecule.

Appropriate host cells may be prokaryotic or eukaryotic. In particular, commonly used host cells may be selected for use in accordance with the present invention including prokaryotic or eukaryotic cells which are genetically accessible and which can be cultured, for example prokaryotic cells, fungal cells, plant cells and animal cells . Preferably, host cells will be selected from a prokaryotic cell, a fungal cell, a plant cell, a protist cell or an animal cell. Preferably, host cells will be selected from a prokaryotic cell, a fungal cell, a plant cell, a protist cell or an animal cell except a human cell.

Preferably, host cells will not include human cells, including embryonic stem cells. Preferred host cells for use in accordance with the present invention are commonly derived from species which typically exhibit high growth rates, are easily cultured and/or transformed, display short generation times, species which have established genetic resources associated with them or species which have been selected, modified or synthesized for optimal expression of heterologous protein under specific conditions. In preferred embodiments of the invention where the protein of interest is eventually to be used in specific industrial, agricultural, chemical or therapeutic contexts, an appropriate host cell may be selected based on the desired specific conditions or cellular context in which the protein of interest is to be deployed. Preferably the host cell will be a prokaryotic cell. In preferred embodiments the host cell is a bacterial cell. The host cell may for instance be an *Escherichia coli* (*E. coli*) cell. Preferably the host cell will be a cell of a thermophilic bacterium.

Methods and uses of the invention described herein may be used to modify genomes of bacterial cells. In particular embodiments, the bacteria are thermophilic bacteria, preferably the bacteria are selected from: *Acidithiobacillus* species including ***Acidithiobacillus caldus***; *Aeribacillus* species including *Aeribacillus pallidus; Alicyclobacillus* species including *Alicyclobacillus acidocaldarius*, *Alicyclobacillus acidoterrestris*, *Alicyclobacillus cycloheptanicusl, Alicyclobacillus hesperidum; Anoxybacillus* species including *Anoxybacillus caldiproteolyticus*, *Anoxybacillus flavithermus*, *Anoxybacillus rupiensis*, *Anoxybacillus tepidamans*; *Bacillus* species including ***Bacillus caldolyticus***, ***Bacillus caldotenax***, ***Bacillus caldovelox***, ***Bacillus coagulans***, ***Bacillus clausii***, ***Bacillus hisashii***, ***Bacillus licheniformis***, ***Bacillus methanolicus***, ***Bacillus smithii including Bacillus smithii ET138***, ***Bacillus subtilis***, *Bacillus thermocopriae*, *Bacillus thermolactis*, ***Bacillus thermoamylovorans***, *Bacillus thermoleovorans*; *Caldibacillus* species including *Caldibacillus debilis*; *Caldicellulosiruptor* species including ***Caldicellulosiruptor bescii***, ***Caldicellulosiruptor hydrothermalis***, *Caldicellulosiruptor kristjanssonii*, *Caldicellulosiruptor kronotskyensis*, *Caldicellulosiruptor lactoaceticus*, *Caldicellulosiruptor obsidiansis*, *Caldicellulosiruptor owensensis*, *Caldicellulosiruptor saccharolyticus*; *Clostridium* species including *Clostridium clariflavum*, *Clostridium straminisolvens*, *Clostridium tepidiprofundi*, *Clostridium thermobutyricum*, ***Clostridium thermocellum***, ***Clostridium thermosuccinogenes***, *Clostridium*

thermopalmarium; *Deinococcus* species including ***Deinococcus cellulosilyticus***, ***Deinococcus deserti***, ***Deinococcus geothermalis***, ***Deinococcus murrayi***, ***Deinococcus radiodurans***; *Defluviitalea* species including *Defluviitalea phaphyphila*, *Desulfotomaculum* species including *Desulfotomaculum carboxydivorans*, *Desulfotomaculum nigrificans*, *Desulfotomaculum salinum*, *Desulfotomaculum solfataricum*; *Desulfurella* species including *Desulfurella acetivorans*; *Desulfurobacterium* species including *Desulfurobacterium thermolithotrophum*; *Geobacillus* species including *Geobacillus icigianus*, *Geobacillus caldoxylosilyticus*, *Geobacillus jurassicus*, *Geobacillus galactosidasius*, ***Geobacillus kaustophilus***, *Geobacillus lituanicus*, ***Geobacillus stearothermophilus***, *Geobacillus subterraneus*, *Geobacillus thermantarcticus*, *Geobacillus thermocatenulatus*, ***Geobacillus thermodenitrificans***, ***Geobacillus thermoglucosidans***, ***Geobacillus thermoleovorans***, *Geobacillus toebii*, ***Geobacillus uzenensis***, *Geobacillus vulcanii*, *Geobacillus zalihae*; *Hydrogenobacter* species including *Hydrogenobacter thermophiles*; *Hydrogenobaculum* species including *Hydrogenobaculum acidophilum*; *Ignavibacterium* species including *Ignavibacterium album*; *Lactobacillus* species including ***Lactobacillus bulgaricus***, ***Lactobacillus delbrueckii***, *Lactobacillus ingluviei*, ***Lactobacillus thermotolerans***; *Marinithermus* species including *Marinithermus hydrothermalis; Moorella* species including ***Moorella thermoacetica***; *Oceanithermus* species including *Oceanithermus desulfurans*, *Oceanithermus profundus*; *Paenibacillus* species including *Paenibacillus* sp. J2, *Paenibacillus marinum*, *Paenibacillus thermoaerophilus; Persephonella* species including *Persephonella guaymasensis*, *Persephonella hydrogeniphila*, *Persephonella marina*; *Rhodothermus* species including ***Rhodothermus marinus***, *Rhodothermus obamensis*, *Rhodothermus profundi*; *Sulfobacillus* species including *Sulfobacillus acidophilus*; *Sulfurihydrogenibium* species including *Sulfurihydrogenibium azorense*, *Sulfurihydrogenibium kristjanssonii*, *Sulfurihydrogenibium rodmanii*, *Sulfurihydrogenibium yellowstonense*; *Symbiobacterium* species including *Symbiobacterium thermophilum*, *Symbiobacterium toebii*; *Thermoanaerobacter* species including ***Thermoanaerobacter brockii***, ***Thermoanaerobacter ethanolicus***, *Thermoanaerobacter italicus*, *Thermoanaerobacter kivui*, *Thermoanaerobacter marianensis*, *Thermoanaerobacter mathranii*, *Thermoanaerobacter pseudoethanolicus*, *Thermoanaerobacter wiegelii*; *Thermoanaerobacterium* species including *Thermoanaerobacterium aciditolerans*,

Thermoanaerobacterium aotearoense, *Thermoanaerobacterium ethanolicus*, ***Thermoanaerobacterium pseudoethanolicus***, ***Thermoanaerobacterium saccharolyticum***, ***Thermoanaerobacterium thermosaccharolyticum***, ***Thermoanaerobacterium xylanolyticum***; *Thermobacillus* species including *Thermobacillus composti*, *Thermobacillus xylanilyticus*; *Thermocrinis* species including *Thermocrinis albus*, *Thermocrinis ruber; Thermodulfatator* species including *Thermodesulfatator atlanticus*, *Thermodesulfatator autotrophicus*, *Thermodesulfatator indicus*; *Thermodesulfobacterium* species including *Thermodesulfobacterium commune*, *Thermodesulfobacterium hydrogeniphilum*; *Thermodesulfobium* species including *Thermodesulfobium narugense*; *Thermodesulfovibrio* species including *Thermodesulfovibrio aggregans*, *Thermodesulfovibrio thiophilus*, *Thermodesulfovibrio yellowstonii*; *Thermosipho* species including *Thermosipho africanus*, *Thermosipho atlanticus*, *Thermosipho melanesiensis*; *Thermotoga* species including ***Thermotoga maritima***, *Thermotoga neopolitana*, ***Thermotoga* sp. RQ7**; *Thermovibrio* species including *Thermovibrio ammonificans*, *Thermovibrio ruber*; *Thermovirga* species including *Thermovirga lienii* and *Thermus* species including ***Thermus aquaticus***, ***Thermus caldophilus***, ***Thermus flavus***, ***Thermus scotoductus***, ***Thermus thermophilus***; *Thiobacillus neapolitanus.*

In another aspect, a method or use described herein can be used to modify bacteria that are mesophilic. In preferred embodiments, the bacteria are selected from: *Acidithiobacillus* species including ***Acidithiobacillus caldus***; *Actinobacillus* species including ***Actinobacillus succinogenes;*** *Anaerobiospirillum* species including *Anaerobiospirillum succiniciproducens; Bacillus* species including ***Bacillus alcaliphilus***, ***Bacillus amyloliquefaciens***, *Bacillus circulans*, ***Bacillus cereus***, ***Bacillus clausii***, *Bacillus firmus*, *Bacillus halodurans*, *Bacillus hisashii, Bacillus lautus*, *Bacillus lentus*, ***Bacillus licheniformis***, ***Bacillus megaterium***, *Bacillus pumilus*, ***Bacillus smithii***, ***Bacillus subtilis***, ***Bacillus thuringiensis***; *Basfia* species including ***Basfia succiniciproducens***; *Brevibacillus* species including ***Brevibacillus brevis***; ***Brevibacillus laterosporus***; *Clostridium* species including ***Clostridium acetobutylicum***, ***Clostridium autoethanogenum***, ***Clostridium beijerinkii***, *Clostridium carboxidivorans*, ***Clostridium cellulolyticum***, ***Clostridium ljungdahlii***, ***Clostridium pasteurianum***, ***Clostridum perfringens***, *Clostridium ragsdalei*,

Clostridium saccharobutylicum, *Clostridium saccharoperbutylacetonium*; *Corynebacterium* species including ***Corynebacterium glutamicum***; *Desulfitobacterium* species including ***Desulfitobacterium dehalogenans***, *Desulfitobacterium hafniense*; *Desulfotomaculum* species including *Desulfotomaculum acetoxidans*, *Desulfotomaculum gibsoniae*, *Desulfotomaculum reducens, Desulfotomaculum ruminis*; *Enterobacter* species including *Enterobacter asburiae*; *Enterococcus* species including ***Enterococcus faecalis***; *Escherichia* species including ***Escherichia coli***; *Lactobacillus* species including ***Lactobacillus acidophilus***, ***Lactobacillus amylophilus***, ***Lactobacillus amylovorus***, *Lactobacillus animalis*, *Lactobacillus arizonensis*, ***Lactobacillus bavaricus***, ***Lactobacillus brevis***, ***Lactobacillus buchneri***, ***Lactobacillus bulgaricus***, ***Lactobacillus casei***, ***Lactobacillus corynoformis***, ***Lactobacillus crispatus***, ***Lactobacillus curvatus***, ***Lactobacillus delbrueckii***, ***Lactobacillus fermentum***, ***Lactobacillus gasseri***, ***Lactobacillus helveticus***, ***Lactobacillus johnsonii***, ***Lactobacillus pentosus***, ***Lactobacillus plantarum***, ***Lactobacillus reuteri***, ***Lactobacillus rhamnosus***, ***Lactobacillus sakei***, ***Lactobacillus salivarius***, ***Lactobacillus sanfriscensis***; *Lactococcus* species, including ***Lactococcus lactis***; *Mannheimia* species including ***Mannheimia succiniciproducens***; *Paenibacillus* species including ***Paenibacillus alvei***, *Paenibacillus beijingensis*, *Paenibacillus borealis*, *Paenibacillus dauci*, *Paenibacillus durus*, *Paenibacillus graminis*, ***Paenibacillus larvae***, *Paenibacillus lentimorbus*, *Paenibacillus macerans*, *Paenibacillus mucilaginosus*, *Paenibacillus odorifer*, ***Paenibacillus polymyxa***, *Paenibacillus stellifer*, *Paenibacillus terrae*, *Paenibacillus wulumuqiensis*; *Pediococcus* species including ***Pediococcus acidilactici***, *Pediococcus claussenii*, *Pediococcus ethanolidurans*, ***Pediococcus pentosaceus***; *Propionibacterium* species, including ***P. acidipropionici***, ***P. freudenreichii***, ***P. jensenii***; *Salmonella typhimurium; Sporolactobacillus* species including *Sporolactobacillus inulinus*, *Sporolactobacillus laevolacticus*; *Staphylococcus aureus; Streptococcus* species including ***Streptococcus agalactiae***, ***Streptococcus bovis***, *Streptococcus equisimilis*, *Streptococcus feacalis*, *Streptococcus mutans*, *Streptococcus oralis*, *Streptococcus pneumonia*, *Streptococcus pyogenes*, *Streptococcus salivarius*, ***Streptococcus thermophilus***, *Streptococcus sobrinus*, *Streptococcus uberis*; *Streptomyces* species including *Streptomyces achromogenes*, *Streptomyces avermitilis*, ***Streptomyces coelicolor***, *Streptomyces griseus*, ***Streptomyces lividans***, ***Streptomyces parvulus***,

Streptomyces venezuelae, Streptomyces vinaceus; *Tetragenococcus* species including ***Tetragenococcus halophilus*** and *Zymomonas* species including ***Zymomonas mobilis***. *Pseudomonas* species including ***Pseudomonas putida***, *Pseudomonas aeruginosa, Pseudomonas alcaligenes, Pseudomonas anguilliseptica, Pseudomonas argentinensis, Pseudomonas borbori, Pseudomonas citronellolis, Pseudomonas flavescens, Pseudomonas mendocina, Pseudomonas nitroreducens, Pseudomonas oleovorans, Pseudomonas pseudoalcaligenes, Pseudomonas resinovorans, Pseudomonas straminea, Pseudomonas asplenii, Pseudomonas aurantiaca, Pseudomonas aureofaciens, Pseudomonas chlororaphis, Pseudomonas corrugate, Pseudomonas fragi, Pseudomonas lundensis, Pseudomonas taetrolens, Pseudomonas antarctica, Pseudomonas azotoformans, 'Pseudomonas blatchfordae', Pseudomonas brassicacearum, Pseudomonas brenneri, Pseudomonas cedrina, Pseudomonas corrugate, Pseudomonas fluorescens, Pseudomonas gessardii, Pseudomonas libanensis, Pseudomonas mandelii, Pseudomonas marginalis, Pseudomonas mediterranea, Pseudomonas meridiana, Pseudomonas migulae, Pseudomonas mucidolens, Pseudomonas orientalis, Pseudomonas panacis, Pseudomonas protegens, Pseudomonas proteolytica, Pseudomonas rhodesiae, Pseudomonas synxantha, Pseudomonas thivervalensis, Pseudomonas tolaasii, Pseudomonas veronii, Pseudomonas denitrificans, Pseudomonas pertucinogena, Pseudomonas cremoricolorata, Pseudomonas entomophila, Pseudomonas fulva, Pseudomonas monteilii, Pseudomonas mosselii, Pseudomonas oryzihabitans, Pseudomonas parafulva, Pseudomonas plecoglossicida, Pseudomonas putida, Pseudomonas balearica, Pseudomonas luteola, Pseudomonas stutzeri, Pseudomonas amygdali, Pseudomonas avellanae, Pseudomonas caricapapayae, Pseudomonas cichorii, Pseudomonas coronafaciens, Pseudomonas ficuserectae, 'Pseudomonas helianthi', Pseudomonas meliae, Pseudomonas savastanoi, Pseudomonas syringae, 'Pseudomonas tomato', Pseudomonas viridiflava, Pseudomonas abietaniphila, Pseudomonas acidophila, Pseudomonas agarici, Pseudomonas alcaliphila, Pseudomonas alkanolytica, Pseudomonas amyloderamosa, Pseudomonas asplenii, Pseudomonas azotifigens, Pseudomonas cannabina, Pseudomonas coenobios, Pseudomonas congelans, Pseudomonas costantinii, Pseudomonas cruciviae, Pseudomonas delhiensis, Pseudomonas excibis, Pseudomonas extremorientalis, Pseudomonas frederiksbergensis, Pseudomonas fuscovaginae, Pseudomonas gelidicola, Pseudomonas grimontii, Pseudomonas*

indica, Pseudomonas jessenii, Pseudomonas jinjuensis, Pseudomonas kilonensis, Pseudomonas knackmussii, Pseudomonas koreensis, Pseudomonas lini, Pseudomonas lutea, Pseudomonas moraviensis, Pseudomonas otitidis, Pseudomonas pachastrellae, Pseudomonas palleroniana, Pseudomonas papaveris, Pseudomonas peli, Pseudomonas perolens, Pseudomonas poae, Pseudomonas pohangensis, Pseudomonas protegens, Pseudomonas psychrophila, Pseudomonas psychrotolerans, Pseudomonas rathonis, Pseudomonas reptilivora, Pseudomonas resiniphila, Pseudomonas rhizosphaerae, Pseudomonas rubescens, Pseudomonas salomonii, Pseudomonas segitis, Pseudomonas septica, Pseudomonas simiae, Pseudomonas suis, Pseudomonas thermotolerans, Pseudomonas toyotomiensis, Pseudomonas tremae, Pseudomonas trivialis, Pseudomonas turbinellae, Pseudomonas tuticorinensis, Pseudomonas umsongensis, Pseudomonas vancouverensis, Pseudomonas vranovensis, Pseudomonas xanthomarina. Preferably the mesophilic bacterium is Pseudomonas putida.

In a further aspect, a method or use defined herein could be used to modify the genome of yeast or fungi. In particular embodiments, the fungal species are mesophilic, preferably the fungi is selected from: an *Aspergillus* species including, but not limited to, ***Aspergillus nidulans***, ***Aspergillus niger***, *Aspergillus oryzae* and *Aspergillus terreus,* more preferably the *Aspergillus* species is *Aspergillus nidulans* or *Aspergillus niger*. Alternatively, the mesophilic fungal species could be a *Candida* species. A method or use defined herein could be used to modify the genome of yeast species including, but not limited to, *Saccharomyces* species including ***Saccharomyces cerevisiae***, *Schizosaccharomyces* species including ***Schizosaccharomyces pombe***, *Pichia* species including, but not limited to ***Pichia pastoris***, ***Pichia stipitis***. A method or use defined herein could be used to modify the genome of fungal species including, but not limited to, *Hansenula* species including ***Hansenula polymorpha***, *Penicillium* species including, but not limited to ***P. brasilianum***, ***P. chrysogenum***, *Yarrowia* species including ***Yarrowia lipolytica***.

The invention further relates to use of a method as defined herein to modify a yeast or fungal species that are thermophilic, preferably the fungi or yeast is selected from: *Aspergillus* species including ***Aspergillus fumigatus***, ***Aspergillus nidulans***, *Aspergillus terreus*, *Aspergillus versicolor*; *Canariomyces* species including

Canariomyces thermophile; Chaetomium species including *Chaetomium mesopotamicum*, ***Chaetomium thermophilum***; *Candida* species including *Candida bovina*, *Candida sloofii*, *Candida thermophila*, ***Candida tropicalis***, ***Candida krusei*** (=*Issatchenkia orientalis*); *Cercophora* species including *Cercophora coronate*, *Cercophora septentrionalis*; *Coonemeria* species including *Coonemeria aegyptiaca*; *Corynascus* species including *Corynascus thermophiles*; *Geotrichum* species including *Geotrichum candidum*; *Kluyveromyces* species including ***Kluyveromyces fragilis***, ***Kluyveromyces marxianus***; *Malbranchea* species including *Malbranchea cinnamomea*, *Malbranchea sulfurea*; *Melanocarpus* species including *Melanocarpus albomyces*;*Myceliophtora* species including *Myceliophthora fergusii*, *Myceliophthora thermophila*; *Mycothermus* species including *Mycothermus thermophiles* (=*Scytalidium thermophilum/Torula thermophila*); *Myriococcum* species including *Myriococcum thermophilum*; *Paecilomyces* species including *Paecilomyces thermophila*; *Remersonia* species including *Remersonia thermophila*; *Rhizomucor* species including ***Rhizomucor pusillus***, *Rhizomucor tauricus*; ***Saccharomyces*** species including ***Saccharomyces cerevisiae***, ***Schizosaccharomyces*** species including ***Schizosaccharomyces pombe***, Scytalidium species including *Scytalidium thermophilum*; *Sordaris* species including *Sordaria thermophila*; *Thermoascus* species including *Thermoascus aurantiacus*, *Thermoascus thermophiles*; *Thermomucor* species including *Thermomucor indicae-seudaticae* and *Thermomyces* species including *Thermomyces ibadanensis*, ***Thermomyces lanuginosus***.

In the aforementioned lists, microbes identified in **bold** typeface have been found to be particularly suitable/applicable in use for the present invention.

Some preferred embodiments of the present invention include one or more thermophilic microbes selected from: Thermophilic *bacilli*, including *Aeribacillus, Alicyclobacillus, Anoxybacillus, Bacillus, Geobacillus*; *Paenibacillus* species; Thermophilic *clostridia*, including *Anaerobacter, Anaerobacterium, Caldicellulosiruptor, Clostridium, Moorella, Thermoanaerobacter, Thermoanaerobacterium, Thermobrachium, Thermohalobacter* species or one or more thermophilic *Lactobacillus* species and mesophilic bacteria selected from *Bacillus* species, *Escherichia coli*, *Lactobacillus* species *Lactococcus* species, *Propionibacterium* species and *Pseudomonas* species.

BRIEF DESCRIPTION OF THE FIGURES

The invention will now be described in detail with reference to a specific embodiment and with reference to the accompanying drawings, in which:

Figure 1 shows a Neighbour-Joining tree of Cas9 protein sequences. All sequences having a sequence similarity above 40% with strain T12 based on pBLAST or PSI-BLAST were included, as well as currently well-characterized sequences (*S. pyogenes, S. thermophiles and A. naeslundii*), as well as all currently identified thermophilic sequences also when these were below 40% identity. For all thermophilic sequences, the percentage identity to T12 is indicated after the strain name. Gene identifier (gi) numbers are indicated before the species name. Legend: Closed circles: thermophilic (optimum above 60°C) Cas9 sequences, closed squares: thermotolerant (optimum <50°C) Cas9 sequences, open triangle: Cas9 sequence currently most used for genome editing purposes from mesophilic origin; no sign: mesophilic Cas9. Values at the nodes represent 1000-replicate bootstrap values; scale bar represents estimated amino acid substitutions per site.

Figure 2 shows a Neighbour-Joining tree of Cas9 gene sequences. Identity at the gene level was extremely poor; sequences from the same organisms as those used for the protein alignment were used for the gene alignment. Gene identifier (gi) numbers are indicated before the species name. Legend: Closed circles: thermophilic (optimum above 60°C) Cas9 sequences, closed squares: thermotolerant (optimum <50°C) Cas9 sequences, open triangle: Cas9 sequence currently most used for genome editing purposes from mesophilic origin; no sign: mesophilic Cas9. Values at the nodes represent 1000-replicate bootstrap values.

Figure 3 shows a protein sequence alignment for gtCas9 (SEQ ID NO: 1) (Type II-C) with well-characterized Type II-C (*A. naeslundii*/'ana'; SEQ ID NO: 8) and Type II-A (*S. pyogenes*/'pyo'; SEQ ID NO: 9 and *S. thermophilus*) Cas9 sequences. Important active site residues are well conserved and indicated with black arrows. Protein domains as described for Ana-Cas9 and Pyo-Cas9 (Jinek, *et al.*, 2014, Science 343:

1247997) are indicated with shaded boxes and similarly coloured letters. The PAM recognition domain has been determined for the *S. pyogenes* Type II-A system but not for any Type II-C system and is therefore only indicated in the *S. pyogenes* sequence.

Figure 4 shows protein architecture of *A. naeslundii* Cas9 (Cas9-Ana) (Jinek *et al.*, 2014). gtCas9 belongs to the same Type II-C CRISPR system and active site residues could be identified.

Figure 5 shows a comparison of crRNA-guided targeting of complementary dsDNA. Base pairing is indicated with dashed lines. RNA is depicted in black, DNA in grey. Base pairing between crRNA spacer and target protospacer is indicated with thick black dashed line, base pairing between DNA strands and between RNA strands is indicated with thick grey dashed lines. The 5' end of the crRNA is indicated. Note that PAM (small white box) in Type I resides downstream of target strand (protospacer), whereas in Type II it resides at the other end on the displaced strand. Likewise, the seed (the predicted sequence of the guide where base pairing with target DNA strand starts, and where no mismatches are allowed) is located close to the PAM, and as such differs in types I and II (Van der Oost, 2014 *ibid.*). Panel A shows a schematic of a Type I Cascade system of *E. coli.* crRNA has internal spacer (grey box, 31-32 nt that allows for target recognition), flanked bt a 8 nt 5' handle and a 29 nt 3' handle that consists of a stem-loop structure (hairpin) (Jore 2011 *ibid.*). Panel B shows a schematic of a Type II Cas9 system of *S. pyogenes*. crRNA basepairs with tracrRNA, that allows for processing by RNaseIII (opposite black triangles). Additionally, the 5' end of the crRNA is trimmed by an RNase (black triangle), typically resulting in a 20 nt spacer. Note that a synthetic loop may be introduced to link the crRNA and tracrRNA, resulting in a single guide RNA (sgRNA) (Jinek *et al.*, 2012 *ibid.*).

Figure 6 shows an alignment of sequences of the *G.thermodenitrificans* T12 type IIc CRISPR system.

Figure 7 shows six single hits obtained to provide an *in silico* PAM prediction for gtCas9.

Figure 8 shows a weblogo combining the results of the alignments illustrated in Figure 7. The weblogo was generated using weblogo.berkeley.edu.

Figure 9 shows the results of an *in vitro* cleavage assay at 60°C targeting plasmids with purified gtCas9. The plasmids included specific 8 nucleotide-long sequence variants of the PAM sequences.

Figure 10 shows the results of *in vitro* assays to investigate the effect of gtCas9 concentration, using a targeted plasmid with the CCCCCCAA [SEQ ID NO: 11] PAM sequence.

Figure 11 shows the results of *in vitro* assays using a targeted plasmid with the CCCCCCAA [SEQ ID NO: 11] PAM sequence over a range of temperatures.

Figure 12 shows the results of *in vivo* genome editing of *Bacillus smithii* ET138 cells using gtCas9 and 8nt PAM sequences, by the growth or absence of colonies of the *Bacillus smithii* ET138 cells on selection plates, as explained in Example 9. Colonies are indicated with arrows in Figure 12.

Figure 13 shows the results of a PCR screen for colonies in which the *pyrF* gene was deleted. The colonies were generated following transformation of *Bacillus smithii* ET138 cells with construct 3 (negative control). 15 colonies were screened but none showed the deletion genotype -2.1 kb band size and instead all showed the wild type -2.9kb band size, as explained in Example 9.

Figure 14 shows the results of a PCR screen for colonies in which the *pyrF* gene was deleted. The colonies were generated following transformation of *Bacillus smithii* ET138 cells with construct 1 (PAM sequence ATCCCCAA [SEQ ID NO: 21]). 20 colonies were screened and one showed the deletion genotype -2.1 kb band size whilst the rest showed both the wild type -2.9kb band size and the deletion genotype -2.1 kb band size, as explained in Example 9. No wild type only genotypes were observed.

Figure 15 shows the *Geobacillus thermodenitrificans* T12 type-IIC CRISPR-Cas locus encodes a thermostable Cas9 homolog, ThermoCas9.

(A) Schematic representation of the genomic locus encoding ThermoCas9. The domain architecture of ThermoCas9 based on sequence comparison, with predicted active sites residues highlighted in red. A homology model of ThermoCas9 generated using Phyre 2 (Kelley *et al. Nat. Protoc.* 10, 845–858 (2015)) is shown, with different colours for the domains.

(B) Phylogenetic tree of Cas9 orthologues that are highly identical to ThermoCas9. Evolutionary analysis was conducted in MEGA7 (Kumar *et al. Mol. Biol. Evol.* 33, 1870–1874 (2016)).

(C) SDS-PAGE of ThermoCas9 after purification by metal-affinity chromatography and gel filtration. The migration of the obtained single band is consistent with the theoretical molecular weight of 126 kD of the apo-ThermoCas9.

Figure 16 shows ThermoCas9 PAM analysis.

(A) Schematic illustrating the *in vitro* cleavage assay for discovering the position and identity (5′-NNNNNNN-3') of the protospacer adjacent motif (PAM). Black triangles indicate the cleavage position.

(B) Sequence logo of the consensus 7nt long PAM of ThermoCas9, obtained by comparative analysis of the ThermoCas9-based cleavage of target libraries. Letter height at each position is measured by information content.

(C) Extension of the PAM identity to the 8th position by *in vitro* cleavage assay. Four linearized plasmid targets, each containing a distinct 5′-CCCCCCAN-3' PAM, were incubated with ThermoCas9 and sgRNA at 55˚C for 1 hour, then analysed by agarose gel electrophoresis.

(D) *In vitro* cleavage assays for DNA targets with different PAMs at 30˚C and 55˚C. Sixteen linearized plasmid targets, each containing one distinct 5'-CCCCCNNA-3' [SEQ ID NO: 13] PAM, were incubated with ThermoCas9 and sgRNA, then analysed for cleavage efficiency by agarose gel electrophoresis. See also Figure 21.

Figure 17 shows ThermoCas9 is active at a wide temperature range and its thermostability increases when bound to sgRNA.

(A) Schematic representation of the sgRNA and a matching target DNA. Target DNA is shown as a rectangular with black outline, and the PAM is shown as a dark grey, horizontal ellipse with back outline. The crRNA is shown as a dark grey rectangular with black outline and the site where the 3'-end of the crRNA is linked with 5'-end of the tracrRNA is shown as a black, vertical ellipse. The black box with the white letters and the light grey box with the black letters indicate the predicted three and two loops at the 3'-side of the tracrRNA, respectively. The 41-nt truncation of the repeat/anti-repeat region –formed by the complementary 3'-end of the crRNA and the 5'-end of the tracrRNA- is indicated with a long, light grey, vertical, dotted line. The predicted 3' position of the first tracrRNA loop is marked with a black triangle and a black dotted line. The predicted 3' position of the second tracrRNA loop is marked with a white triangle and a black dotted line. The predicted 3' position of the third tracrRNA loop is marked with a white triangle and a white dotted line.

(B) The importance of the predicted three stem-loops of the tracrRNA scaffold was tested by transcribing truncated variants of the sgRNA and evaluating their ability to guide ThermoCas9 to cleave target DNA at various temperatures. Average values of at least two biological replicates are shown, with error bars representing S.D.

(C) To identify the maximum temperature, endonuclease activity of ThermoCas9:sgRNA RNP complex was assayed after incubation at 60˚C, 65˚C and 70˚C for 5 or 10 min. The pre-heated DNA substrate was added and the reaction was incubated for 1 hour at the corresponding temperature.

(D) Comparison of active temperature range of ThermoCas9 and SpCas9 by activity assays conducted after 5 min of incubation at the indicated temperature. The pre-heated DNA substrate was added and the reaction was incubated for 1 hour at the same temperature.

Figure 18 shows ThermoCas9-based genome engineering in thermophiles.

(A) Schematic overview of the basic pThermoCas9_Δgene-of-interest (goi) construct. The *thermocas9* gene was introduced either to the pNW33n (*B. smithii*) or to the pEMG (*P. putida*) vector. Homologous recombination flanks were introduced upstream *thermocas9* and encompassed the 1kb (*B.smithii*) or 0.5kb (*P. putida*) upstream and 1kb or 0.5 kb downstream region of the gene of

interest (goi) in the targeted genome. A sgRNA-expressing module was introduced downstream the *thermocas9* gene. As the origin of replication (ori), replication protein (rep), antibiotic resistance marker (AB) and possible accesory elements (AE) are backbone specific, they are represented with dotted outline.

(B) Agarose gel electrophoresis showing the resulting products from genome-specific PCR on ten colonies from the ThermoCas9-based *pyrF* deletion process from the genome of *B. smithii* ET 138. All ten colonies contained the Δ*pyrF* genotype and one colony was a clean Δ*pyrF* mutant, lacking the wild type product.

(C) Schematic overview of the basic pThermoCas9i_goi construct. Aiming for the expression of a catalytically inactive ThermoCas9 (Thermo-dCas9:D8A, H582A mutant), the corresponding mutations were introduced to create the *thermo-dcas9* gene. The *thermo-dcas9* gene was introduced to the pNW33n vector. An sgRNA-expressing module was introduced downstream the *thermo-dcas9.*

(D) Graphical representation of the production, growth and RT-qPCR results from the *ldhL* silencing experiment using Thermo-dCas9. The graphs represent the lactate production, optical density at 600nm and percentage of *ldhL* transcription in the repressed cultures compared to the control cultures. Average values from at least two biological replicates are shown, with error bars representing S.D.

Figure 19 shows a multiple sequence alignment of Type II-A, B and C Cas9 orthologues. Cas9 protein sequences of *Streptococcus pyogenes* (Sp), *Streptococcus thermophilus* (St), *Wolinella succinogenes* (Ws), *Neisseria meningitides* (Nm), *Actinomyces naeslundii* (An), and *Geobacillus thermodenitrificans* (*Thermo*) were aligned using ClustalW1 in MEGA7 2 with default settings; ESPript3 was used to generate the visualization. Strictly conserved residues are shown in white text on grey background; similar residues are shown in black text in white vertical rectangules with black outline. Pyramids indicate the two conserved nuclease domains in all sequences. Horizontal black arrows and curls indicate β-strands and α-helices, respectively, in the SpCas9 secondary structure (protein database nr 4CMP4). Structural domains are indicated for SpCas9 and ThermoCas9 using the same colour scheme as in Figure 15A.

Figure 20 shows *in silico* PAM determination results. Panel (A) shows the two hits obtained with phage genomes using CRISPRtarget6. Panel (B) shows sequence logo of the consensus 7nt long PAM of ThermoCas9, obtained by *in silico* PAM analysis. Letter height at each position is measured by information content.

Figure 21 shows ThermoCas9 PAM discovery. *In vitro* cleavage assays for DNA targets with different PAMs at 20˚C, 37˚C, 45˚C and 60˚C. Seven (20˚C) or sixteen (37˚C, 45˚C, 60˚C) linearized plasmid targets, each containing a distinct 5'-CCCCCNNA-3' [SEQ ID NO: 13] PAM, were incubated with ThermoCas9 and sgRNA, then analysed by agarose gel electrophoresis.

Figure 22 shows activity of ThermoCas9 at a wide temperature range using sgRNA containing one loop. The importance of the predicted three stem loops of the tracrRNA scaffold was tested by transcribing truncated variations of the sgRNA and evaluating their ability to guide ThermoCas9 to cleave target DNA at various temperatures. Shown above is the effect of one loop on the activity of ThermoCas9 at various temperatures. Average values from at least two biological replicates are shown, with error bars representing S.D.

Figure 23 shows ThermoCas9 mediates dsDNA targeting using divalent cations as catalysts and does not cleave ssDNA. Panel (A) shows *in vitro* plasmid DNA cleavage by ThermoCas9 with EDTA and various metal ions. M = 1 kb DNA ladder. Panel (B) shows activity of ThermoCas9 on ssDNA substrates. M= 10 bp DNA ladder.

Figure 24 shows spacer selection for the *ldhL* silencing experiment. Schematic representation of the spacer (sgRNA)-protospacer annealing during the *ldhL* silencing process; the selected protospacer resides on the non-template strand and 39nt downstream the start codon of the *ldhL* gene.

Figure 25 shows a map of plasmid pThermoCas9_ppΔpyrF consisting of the pEMG backbone, the *Pseudomonas putida pyrF* flanking region and the *thermocas9* gene and a *Pseudomonas putida pyrF* targeting sgRNA.

Figure 26 shows the results of capillary gel electrophoresis showing the resulting products from genomespecific PCR on the obtained colonies from the ThermoCas9-based *pyrF* deletion process from the genome of *Pseudomonas putida*. The 1854 bp band and the 1112 bp band corresponds to the *pyrF* and Δ*pyrF* genotype, respectively.

Below are polynucleotide and amino acid sequences of Cas proteins used in accordance with the invention.

[SEQ ID NO: 1] *Geobacillus thermodenitrificans* T12 Cas9 protein AA sequence

MKYKIGLDIGITSIGWAVINLDIPRIEDLGVRIFDRAENPKTGESLALPRRLARSARRR
LRRRKHRLERIRRLFVREGILTKEELNKLFEKKHEIDVWQLRVEALDRKLNNDELARI
LLHLAKRRGFRSNRKSERTNKENSTMLKHIEENQSILSSYRTVAEMVVKDPKFSLH
KRNKEDNYTNTVARDDLEREIKLIFAKQREYGNIVCTEAFEHEYISIWASQRPFASK
DDIEKKVGFCTFEPKEKRAPKATYTFQSFTVWEHINKLRLVSPGGIRALTDDERRLIY
KQAFHKNKITFHDVRTLLNLPDDTRFKGLLYDRNTTLKENEKVRFLELGAYHKIRKAI
DSVYGKGAAKSFRPIDFDTFGYALTMFKDDTDIRSYLRNEYEQNGKRMENLADKVY
DEELIEELLNLSFSKFGHLSLKALRNILPYMEQGEVYSTACERAGYTFTGPKKKQKT
VLLPNIPPIANPVVMRALTQARKVVNAIIKKYGSPVSIHIELARELSQSFDERRKMQK
EQEGNRKKNETAIRQLVEYGLTLNPTGLDIVKFKLWSEQNGKCAYSLQPIEIERLLE
PGYTEVDHVIPYSRSLDDSYTNKVLVLTKENREKGNRTPAEYLGLGSERWQQFETF
VLTNKQFSKKKRDRLLRLHYDENEENEFKNRNLNDTRYISRFLANFIREHLKFADSD
DKQKVYTVNGRITAHLRSRWNFNKNREESNLHHAVDAAIVACTTPSDIARVTAFYQ
RREQNKELSKKTDPQFPQPWPHFADELQARLSKNPKESIKALNLGNYDNEKLESL
QPVFVSRMPKRSITGAAHQETLRRYIGIDERSGKIQTVVKKKLSEIQLDKTGHFPMY
GKESDPRTYEAIRQRLLEHNNDPKKAFQEPLYKPKKNGELGPIIRTIKIIDTTNQVIPL
NDGKTVAYNSNIVRVDVFEKDGKYYCVPIYTIDMMKGILPNKAIEPNKPYSEWKEMT
EDYTFRFSLYPNDLIRIEFPREKTIKTAVGEEIKIKDLFAYYQTIDSSNGGLSLVSHDN
NFSLRSIGSRTLKRFEKYQVDVLGNIYKVRGEKRVGVASSSHSKAGETIRPL*

[SEQ ID NO: 7] *Geobacillus thermodenitrificans* T12 Cas9 DNA Sequence

ATGAAGTATAAAATCGGTCTTGATATCGGCATTACGTCTATCGGTTGGGCTGTC
ATTAATTTGGACATTCCTCGCATCGAAGATTTAGGTGTCCGCATTTTTGACAGAG
CGGAAAACCCGAAAACCGGGGAGTCACTAGCTCTTCCACGTCGCCTCGCCCGC
TCCGCCCGACGTCGTCTGCGGCGTCGCAAACATCGACTGGAGCGCATTCGCC

GCCTGTTCGTCCGCGAAGGAATTTTAACGAAGGAAGAGCTGAACAAGCTGTTT
GAAAAAAAGCACGAAATCGACGTCTGGCAGCTTCGTGTTGAAGCACTGGATCG
AAAACTAAATAACGATGAATTAGCCCGCATCCTTCTTCATCTGGCTAAACGGCG
TGGATTTAGATCCAACCGCAAGAGTGAGCGCACCAACAAAGAAAACAGTACGAT
GCTCAAACATATTGAAGAAAACCAATCCATTCTTTCAAGTTACCGAACGGTTGCA
GAAATGGTTGTCAAGGATCCGAAATTTTCCCTGCACAAGCGTAATAAAGAGGAT
AATTACACCAACACTGTTGCCCGCGACGATCTTGAACGGGAAATCAAACTGATT
TTCGCCAAACAGCGCGAATATGGGAACATCGTTTGCACAGAAGCATTTGAACAC
GAGTATATTTCCATTTGGGCATCGCAACGCCCTTTTGCTTCTAAGGATGATATC
GAGAAAAAAGTCGGTTTCTGTACGTTTGAGCCTAAAGAAAAACGCGCGCCAAAA
GCAACATACACATTCCAGTCCTTCACCGTCTGGGAACATATTAACAAACTTCGT
CTTGTCTCCCCGGGAGGCATCCGGGCACTAACCGATGATGAACGTCGTCTTAT
ATACAAGCAAGCATTTCATAAAAATAAAATCACCTTCCATGATGTTCGAACATTG
CTTAACTTGCCTGACGACACCCGTTTTAAAGGTCTTTTATATGACCGAAACACCA
CGCTGAAGGAAAATGAGAAAGTTCGCTTCCTTGAACTCGGCGCCTATCATAAAA
TACGGAAAGCGATCGACAGCGTCTATGGCAAAGGAGCAGCAAAATCATTTCGT
CCGATTGATTTTGATACATTTGGCTACGCATTAACGATGTTTAAAGACGACACCG
ACATTCGCAGTTACTTGCGAAACGAATACGAACAAAATGGAAAACGAATGGAAA
ATCTAGCGGATAAAGTCTATGATGAAGAATTGATTGAAGAACTTTTAAACTTATC
GTTTTCTAAGTTTGGTCATCTATCCCTTAAAGCGCTTCGCAACATCCTTCCATAT
ATGGAACAAGGCGAAGTCTACTCAACCGCTTGTGAACGAGCAGGATATACATTT
ACAGGGCCAAAGAAAAAACAGAAAACGGTATTGCTGCCGAACATTCCGCCGAT
CGCCAATCCGGTCGTCATGCGCGCACTGACACAGGCACGCAAAGTGGTCAATG
CCATTATCAAAAAGTACGGCTCACCGGTCTCCATCCATATCGAACTGGCCCGG
GAACTATCACAATCCTTTGATGAACGACGTAAAATGCAGAAAGAACAGGAAGGA
AACCGAAAGAAAAACGAAACTGCCATTCGCCAACTTGTTGAATATGGGCTGACG
CTCAATCCAACTGGGCTTGACATTGTGAAATTCAAACTATGGAGCGAACAAAAC
GGAAAATGTGCCTATTCACTCCAACCGATCGAAATCGAGCGGTTGCTCGAACCA
GGCTATACAGAAGTCGACCATGTGATTCCATACAGCCGAAGCTTGGACGATAG
CTATACCAATAAAGTTCTTGTGTTGACAAAGGAGAACCGTGAAAAAGGAAACCG
CACCCCAGCTGAATATTTAGGATTAGGCTCAGAACGTTGGCAACAGTTCGAGAC
GTTTGTCTTGACAAATAAGCAGTTTTCGAAAAAGAAGCGGGATCGACTCCTTCG
GCTTCATTACGATGAAAACGAAGAAAATGAGTTTAAAAATCGTAATCTAAATGAT
ACCCGTTATATCTCACGCTTCTTGGCTAACTTTATTCGCGAACATCTCAAATTCG

CCGACAGCGATGACAAACAAAAAGTATACACGGTCAACGGCCGTATTACCGCC
CATTTACGCAGCCGTTGGAATTTTAACAAAAACCGGGAAGAATCGAATTTGCAT
CATGCCGTCGATGCTGCCATCGTCGCCTGCACAACGCCGAGCGATATCGCCCG
AGTCACCGCCTTCTATCAACGGCGCGAACAAAACAAAGAACTGTCCAAAAAGAC
GGATCCGCAGTTTCCGCAGCCTTGGCCGCACTTTGCTGATGAACTGCAGGCGC
GTTTATCAAAAAATCCAAAGGAGAGTATAAAAGCTCTCAATCTTGGAAATTATGA
TAACGAGAAACTCGAATCGTTGCAGCCGGTTTTTGTCTCCCGAATGCCGAAGC
GGAGCATAACAGGAGCGGCTCATCAAGAAACATTGCGGCGTTATATCGGCATC
GACGAACGGAGCGGAAAAATACAGACGGTCGTCAAAAAGAAACTATCCGAGAT
CCAACTGGATAAAACAGGTCATTTCCCAATGTACGGGAAAGAAAGCGATCCAAG
GACATATGAAGCCATTCGCCAACGGTTGCTTGAACATAACAATGACCCAAAAAA
GGCGTTTCAAGAGCCTCTGTATAAACCGAAGAAGAACGGAGAACTAGGTCCTAT
CATCCGAACAATCAAAATCATCGATACGACAAATCAAGTTATTCCGCTCAACGAT
GGCAAAACAGTCGCCTACAACAGCAACATCGTGCGGGTCGACGTCTTTGAGAA
AGATGGCAAATATTATTGTGTCCCTATCTATACAATAGATATGATGAAAGGGATC
TTGCCAAACAAGGCGATCGAGCCGAACAAACCGTACTCTGAGTGGAAGGAAAT
GACGGAGGACTATACATTCCGATTCAGTCTATACCCAAATGATCTTATCCGTATC
GAATTTCCCCGAGAAAAAACAATAAAGACTGCTGTGGGGGAAGAAATCAAAATT
AAGGATCTGTTCGCCTATTATCAAACCATCGACTCCTCCAATGGAGGGTTAAGT
TTGGTTAGCCATGATAACAACTTTTCGCTCCGCAGCATCGGTTCAAGAACCCTC
AAACGATTCGAGAAATACCAAGTAGATGTGCTAGGCAACATCTACAAAGTGAGA
GGGGAAAAGAGAGTTGGGGTGGCGTCATCTTCTCATTCGAAAGCCGGGGAAAC
TATCCGTCCGTTATAA

DETAILED DESCRIPTION

Example 1: Isolation of *Geobacillus thermodenitrificans*

G. thermodenitrificans was surprisingly discovered during a search of a library of ± 500 isolates for a thermophile capable of degrading lignocellulosic substrates under anaerobic conditions. At first a library of ± 500 isolates was established which, after several selection rounds by isolation on cellulose and xylan, was trimmed down to 110 isolates. This library of 110 isolates consisted solely of *Geobacillus* isolates with *G. thermodenitrificans* representing 79% of the library.

The isolated *G. thermodenitrificans* strain has been named "T12". The Cas9 protein from *G. thermodenitrificans* T12 has been named "gtCas9".

Example 2: Defining the essential consensus sequences for Cas9 in *Geobacillus thermodenitrificans*

The following database searches and alignments were performed:
pBLAST and nBLAST were performed on the in-house BLAST server, in which either the protein or gene sequence of *G. thermodenitrificans* T12 was used as query sequence. This database was last updated May 2014 and therefore does not contain the most recently added *Geobacillus* genomes, but normal online BLAST was not used to prevent publication of the T12 sequence. Sequence identities found to be greater than 40% in the BLAST search are included in Figure 1.

To include more recent sequence data, the sequence of *Geobacillus* MAS1 (most closely related to gtCas9) was used to perform a PSI-BLAST on the NCBI website (Johnson *et al.*, 2008 Nucleic Acids Res. 36(Web Server issue): W5-9). Two consecutive rounds of PSI-BLAST were performed, in which only sequences that met the following criteria were used for the next round: minimum sequence coverage of 96% in the first round and 97% in the second and third round, minimum identity 40%, only one strain per species.

The sequences resulting from the PSI-BLAST, as well as the sequences with more than 40% identity to T12 from the internal server pBLAST that did not appear in the PSI-BLAST were aligned together with currently well-characterized mesophilic sequences and all currently identified thermophilic sequences also if these were more distantly related, from which a Neighbour-Joining tree was constructed (see Figure 1). Alignment was performed in Mega6 using ClustalW, after which a tree was constructed using the Neighbour-Joining method and bootstrap analysis was performed using 1000 replicates.

When BLASTn was performed using *Geobacillus sp.* MAS1 as the query sequence, only *Geobacillus sp.* JF8 Cas9 was identified with 88% identity, indicating very little

homology at the gene level. Figure 2 is a Neighbour-Joining tree of Clustal-aligned Cas9 gene sequences.

Protein sequences of *G. thermodenitrificans* T12, *A. naeslundii* and *S. pyogenes* were further analyzed for protein domain homology (see Figure 3) by aligning them in CloneManager using BLOSUM62 with default settings.

Example 3: Identifying core amino acid motifs which are essential for the function of CAS9 and those which confer thermostability in thermophilic Cas9 nucleases

Percentages identity of the above described aligned protein sequences are provided in Figure 1. gtCas9 belongs to Type II-C. The best-studied and recently crystalized structure of a Type II-C system is from *Actinomyces naeslundii* (Jinek *et al.*, 2014, Science 343: 1247997). This protein sequence shows only 20% identity to gtCas9 but can be used to estimate highly conserved residues. Two well-characterized Type II-A systems (*S. pyogenes* and *S. thermophilus*) were also included in the analyses (Jinek *et al.*, 2014, Science 343: 1247997; Nishimasu *et al.*, 2014, Cell 156: 935-949). Alignments of these four protein sequences are shown in Figure 3; Figure 4 shows the protein architecture as determined for *A. naeslundii* ('Ana-Cas9') (Jinek *et al.*, 2014, Science 343: 1247997). The length of Cas9 from t12 (gtCas9) and *Actinomyces naeslundii* is highly similar (*A. naeslundii* 1101 aa, gtCas9 1082 aa) and gtCas9 is expected to have similar protein architecture but this remains to be determined, as the overall sequence identity to cas9-Ana is only 20%. All active side residues described by Jinek *et al.* (Jinek *et al.*, 2014, Science 343: 1247997) in Cas9 from *A. naeslundii* and *S. pyogenes* could be identified in gtCas9 (see Figure 3). The PAM-binding domain has been determined for the *S. pyogenes* Type II-A system but not for any Type II-C system and is therefore only indicated in the *S. pyogenes* sequence. Moreover, the PAM-recognition site varies strongly, not only between CRISPR systems but also between species containing the same system.

Example 4: Determination of the PAM sequence of *G. thermodenitrificans* gtCas9

It has been established that the prokaryotic CRISPR systems serve their hosts as adaptive immune systems (Jinek *et al.*, 2012, Science 337: 816-821) and can be used for quick and effective genetic engineering (Mali *et al.*, 2013, Nat Methods 10: 957-963.).

Cas9 proteins function as sequence-specific nucleases for the type II CRISPR systems (Makarova *et al.*, 2011, Nat Rev Micro 9: 467-477). Small crRNA molecules, which consist of a "spacer" (target) linked to a repetition region, are the transcription and processing products of a CRISPR loci. "Spacers" naturally originate from the genome of bacteriophages and mobile genetic elements, but they can also be designed to target a specific nucleotide sequence during a genetic engineering process (Bikard *et al.*, 2013, Nucleic Acids Research 41: 7429-7437). The crRNA molecules are employed by the Cas9 as guides for the identification of their DNA targets. The spacer region is identical to the targeted for cleavage DNA region, the "protospacer" (Brouns *et al.*, 2012, Science 337: 808-809). A PAM (Protospacer Adjacent Motif), next to the protospacer, is required for the recognition of the target by the Cas9 (Jinek *et al.*, 2012, Science 337: 816-821).

In order to perform *in vitro* or *in vivo* PAM-determination studies for Type II systems, it is necessary to *in silico* predict the CRISPR array of the system, the tracrRNA-expressing module. The CRISPR array is used for the identification of the crRNA module. The tracrRNA-expressing sequence is located either within a 500 bp-window flanking Cas9 or between the Cas genes and the CRISPR locus (Chylinski, K., *et al.* (2014) Classification and evolution of type II CRISPR-Cas systems. *Nucleic Acids Res.* 42, 6091-6105). The tracrRNA should consist of a 5'-sequence with high level of complementarity to the direct repeats of the CRISPR array, followed by a predicted structure of no less than two stem-loop structures and a Rho-independent transcriptional termination signal (Ran, F.A., *et al.* (2015) In vivo genome editing using Staphylococcus aureus Cas9. Nature 520, 186-191). The crRNA and tracrRNA molecule can then be used to design a chimeric sgRNA module. The 5'-end of the sgRNA consists of a truncated 20 nt long spacer followed by the 16-20 nt long truncated repeat of the CRISPR array. The repeat is followed by the corresponding truncated anti-repeat and the stem loop of the tracrRNA module. The repeat and anti-repeat parts of the sgRNA are generally connected by a GAAA linker (Karvelis, T., *et*

al. (2015) Rapid characterization of CRISPR-Cas9 protospacer adjacent motif sequence elements. Genome Biol. 16, 253).

The *cas* genes (the cas9 followed by the cas1 and the cas2 genes) of the *G.thermodenitrificans* T12 type IIc CRISPR system are transcribed using the antisense strand of the T12 chromosome. The cas2 gene is followed by a 100bp long DNA fragment which upon transcription forms an RNA structure with multiple loops. This structure obviously acts as a transcriptional terminator.

A CRISPR array with 11 repeats and 10 spacer sequences is located upstream of the transcriptional termination sequence and the leader of the array is located at the 5' end of the array. The DNA locus which is transcribed into the tracrRNA is expected to be downstream of the *cas9* gene. The alignment of the 325bp long sequence right downstream of the *cas9* gene with the 36bp long repeat from the CRISPR array revealed that there is a 36bp long sequence in the tracrRNA locus almost identical to the repeat (shown in Figure 6). This result led us to the conclusion that the direction of the transcription of the tracrRNA locus should be opposite to the direction of the transcription of the CRISPR array. Consequently the 5'-end of the tracrRNA will be complementary to the 3'-end of the crRNA, leading to the formation of the –required by the Cas9- dual-RNA molecule.

Example 5: Target generation with randomized PAM

Two different spacers from the CRISPR II loci of the *G. thermodenitrificans* T12 strain were amplified by PCR using the *G. thermodenitrificans* T12 genomic DNA as template. Two pairs of degenerate primers were used for the amplification of each spacer:

Firstly, a pair that cause the introduction of six random nucleotides upstream of the "protospacer" fragment were used, leading to the production of a pool of protospacers with randomized PAM sequences.

Secondly, a pair that cause the introduction of six random nucleotides downstream of the "protospacer" fragment were used, leading to the production of a pool of protospacers with randomized PAM sequences.

The produced fragments were ligated to the pNW33n vector, producing 4 pools of "protospacer" constructs, with all the possible 4096 different combinations of 6-nucleotide long PAMs each. The assembled DNA was used for the transformation of *G. thermodenitrificans* T12 cells. The cells were plated on chloramphenicol selection and more than 2 x 10^6 cells from each protospacer pool will be pooled. The plasmid DNA was extracted from the pools, the target region will be PCR amplified and the products sent for deep sequencing. The PAMs with the fewest reads will be considered active and the process will be repeated only with pNW33n constructs that contain spacers with these PAMs. Reduced transformation efficiency of the *G. thermodenitrificans* T12 will confirm the activity of the PAMs.

Example 6: *In vitro* determination of PAM sequences for gtCas9

Construction of the pRham:*cas9*$_{gt}$ vector

The *cas9*$_{gt}$ gene was PCR amplified from the *G. thermodenitrificans* T12 genome, using the BG6927 and BG6928 primers, and combined with the pRham C-His Kan Vector (Lucigen) in one mixture. The mixture was used for transforming *E. cloni* thermo-competent cells according to the provided protocol. 100µl from the transformation mixture were plated on LB+50kanamycin plates for overnight growth at 37°C. Out of the formed *E. cloni*:: pRham:*cas9*$_{gt}$ single colonies 3 were randomly selected and inoculated in 10ml LB medium containing 50µg/ml kanamucin. Glycerol stocks were prepared from the cultures by adding sterile glycerol to 1ml from each culture up to a final concentration of 20% (v/v). The glycerol stocks were stored at -80°C. The remaining 9ml from each culture were used for plasmid isolation according to the "GeneJET Plasmid Miniprep Kit" (Thermoscientific) protocol. The plasmids were sent for sequence verification of the *cas9*$_{gt}$ and one of the plasmids was verified to contain the gene with the right sequence. The corresponding culture was further used for heterologous expression and purification of the gtCas9.

Heterologous Expression of gtCas9 in *E. cloni*::pRham: *cas9*$_{gt}$ vector

An *E. cloni*:: pRham:*cas9*$_{gt}$ preculture was prepared after inoculating 10ml LB+50kanamycin with the corresponding glycerol stocks. After overnight growth at 37°C and 180rpm, 2ml from the preculture were used for inoculating 200ml of LB+50kanamycin medium. The *E. cloni*::pRham: *cas9*$_{gt}$ culture was incubated at 37°C, 180rpm until an OD_{600} of 0.7. The gtCas9 expression was then induced by adding L-rhamnose to a final concentration of 0.2%w/v. The expression was allowed to proceed for 8h, after which the cultures were centrifuged for 10 minutes at 4700rpm, 4°C to harvest the cells. The medium was discarded and the pelleted cells were either stored at -20°C or used for the preparation of the cell free extract (CFE) according to the following protocol:

1. Resuspend the pellet in 20ml Sonication Buffer (20mM Sodium Phosphate buffer (pH=7.5), 100mM NaCl, 5mM MgCl2, 5%(v/v) Glycerol , 1mM DTT)
2. Disrupt 1 ml of cells by sonication (8 pulses of 30 seconds, cool for 20 seconds on ice in between)
3. Centrifuge for 15 minutes at 35000g, 4°C in order to precipitate insoluble parts
4. Remove the supernatant and store it at 4°C or on ice

Designing and construction of the PAM library targeting sgRNA module for gtCas9

After *in silico* determination of the tracrRNA expressing DNA module in the genome of *G. thermodenitrificans* T12 strain (see Example 4 above), a single guide (sg)RNA expressing DNA module that combines the crRNA and tracrRNA modules of the CRISPR/Cas9 system in a single molecule was designed. The spacer at the 5'-end of the sgRNA was designed to be complementary to the protospacer of the plasmid library and the module was set under the transcriptional control of a T7 promoter. The pT7_sgRNA DNA module was synthesized by Baseclear and received in a pUC57 vector, forming the pUC57:pT7_sgRNA vector. DH5α competent *E.coli* cells (NEB) were transformed with the vector and the transformation mixture was plated on LB-agar plates containing 100μg/ml ampicillin. The plates were incubated overnight at 37°C. Three of the formed single colonies were inoculated in 10ml LB medium containing 100μg/ml ampicillin. Glycerol stocks were prepared from the cultures by adding sterile glycerol to 1ml from each culture up to a final concentration of 20% (v/v). The glycerol stocks were stored at -80°C. The remaining 9ml from each culture were used for plasmid isolation according to the "GeneJET Plasmid Miniprep Kit" (Thermoscientific) protocol. The isolated plasmid was used as a PCR template for

amplification of the pT7_sgRNA module. The 218bp long pT7_sgRNA DNA module (of which the first 18bp correspond to the pT7) was obtained using the primers BG6574 and BG6575. The complete PCR mixture was run on a 1.5% agarose gel. The band with the desired size was excised and purified according to the "Zymoclean™ Gel DNA Recovery Kit" protocol.

In vitro transcription (IVT) was performed using the "HiScribe™ T7 High Yield RNA Synthesis Kit" (NEB). The purified pT7_sgRNA DNA module was used as template. The IVT mixture was mixed with an equal volume of RNA loading dye (NEB) and heated at 70°C for 15 minutes in order to disrupt the secondary structure. The heat treated IVT mixture was run on a denaturing Urea-PAGE and the resulting polyacrylamide gel was embaptised for 10 minutes in 100ml 0.5x TBE buffer containing 10µl of SYBR Gold (Invitrogen) for staining purposes. The band at the desired size (200nt) was excised and the sgRNA was purified according to the following RNA purification protocol:

1. Cut RNA gel fragments with a scalpel and add 1ml of RNA elution buffer, leave overnight at room temperature.
2. Divide 330µl aliquots into new 1.5ml tubes.
3. Add 3 volumes (990µl) of pre-chilled (-20°C) 100% EtOH.
4. Incubate for 60 minutes at -20°C.
5. Centrifuge for 20 minutes at 13000rpm in a microfuge at room temperature.
6. Remove EtOH, wash pellet with 1ml 70% EtOH.
7. Centrifuge for 5 minutes at 13000rpm in a microfuge at room temperature.
8. Remove 990µl of the supernatant.
9. Evaporate the rest EtOH in a thermomixer at 55°C for 15 to 20 minutes.
10. Resuspend pellet in 20µl MQ, store at -20°C.

<u>Designing and construction of a 7nt long PAM library, and linearization of the library</u>

The design and construction of the PAM library was based on the pNW33n vector. A 20bp long protospacer was introduced to the vector, flanked at its 3'side by a 7 degenerate nucleotides long sequence; the degenerate sequence serves as the PAM and when the protospacer is flanked by a right PAM then it can be recognized as a target by an sgRNA loaded Cas9 and cleaved. The PAM library was prepared according to the following protocol:

1. Prepare the SpPAM double stranded DNA insert by annealing the single stranded DNA oligos 1 (BG6494) and 2 (BG6495)
   I. 10µl 10x NEBuffer 2.1
   II. 1µl 50µM oligo 1 (~1.125µg)
   III. 1µl 50µM oligo 2 (~1.125µg)
   IV. 85 µl MQ
   V. Incubate the mixture at 94°C for 5min and cool down to 37°C at a rate of 0.03°C/sec
2. Add 1 µl Klenow 3'->5' exo$^-$ polymerase (NEB) to each annealed oligos mixture and then add 2.5 µl of 10µM dNTPs. Incubate at 37°C for 1h and then at 75°C for 20min.
3. Add 2µl of the HF-BamHI and 2µl of the BspHI restriction enzymes to 46µl of the annealing mixture. Incubate at 37°C for 1h. This process will lead to the SpPAMbb insert with sticky ends. Use the Zymo DNA cleaning and concentrator kit (Zymo Research) to clean the created insert.
4. Digest pNW33n with the HF-BamHI and BspHI (NEB) and purify the 3.400bp long linear pNW33nbb fragment with sticky ends, using the Zymo DNA cleaning and concentrator kit (Zymo Research).
5. Ligate 50ng of pNW33nBB with 11ng of the SPPAMbb insert using the NEB T4 ligase according to the provided protocol. Purify the ligation mixture using the Zymo DNA cleaning and concentrator kit (Zymo Research).
6. Transform DH10b electro-competent cells (200µl of cells with 500ng of DNA). Recover the cells in SOC medium (200µl cells in 800µl SOC) for an hour and then inoculate 50ml of LB+12.5 µg/ml chloramphenicol with the recovered cells. Incubate overnight the culture at 37°C and 180rpm.
7. Isolate plasmid DNA from the culture using the JetStar 2.0 maxiprep kit (GENOMED).
8. Use the SapI (NEB) restriction according to the provided protocol for linearizing the isolated plasmids.

Designing and execution of the PAM determination reactions

The following cleavage reaction was set up for gtCas9-induced introduction of dsDNA breaks to the PAM library members that contain the right PAM downstream of the 3' end of the targeted protospacer:

1. 2.5µg of *E. cloni*::pRham:*cas9*$_{gt}$ CFE per reaction
2. sgRNA to 30nM final concentration
3. 200ng of linearized PAM library per reaction
4. 2µl of cleavage buffer (100mM Sodium Phosphate buffer (pH=7.5), 500mM NaCl, 25mM MgCl2, 25%(v/v) Glycerol, 5mM DTT)
5. MQ water up to 20µl final volume

The reaction was incubated for 1h at 60°C and stopped after adding 4µl of 6x gel loading dye (NEB). The reaction mixture was then loaded to a 1% agarose gel. The gel was subjected to an 1h and 15min long electrophoresis at 100V and then it was incubated for 30min in 100ml 0.5xTAE buffer containing 10µl of SYBR Gold dye (ThermoFisher). After visualizing the DNA bands with blue light, the band that corresponded to the successfully cleaved and PAM containing DNA fragments was cut-off the gel and gel purified using the "Zymoclean™ Gel DNA Recovery Kit" according to the provided protocol.

Tagging of the PAM-containing gtCAs9 cleaved DNA fragments for sequencing

The Cas9-induced DNA breaks are usually introduced between the 3rd and the 4th nucleotide of a protospacer, proximally to the PAM sequence. As a result, it is not possible to design a pair of primers that can PCR amplify the PAM-containing part of the cleaved DNA fragments, in order to further on sequence and determine the PAM sequence. For this purpose a 5-step process was employed:

Step 1: A-Tailing with Taq Polymerase

A-Tailing is a process to add a non-templated adenine to the 3' end of a blunt, double-stranded DNA molecule using Taq polymerase

Reaction components:

- gtCas9-cleaved and PAM-containing DNA fragments – 200ng
- 10X ThermoPol® Buffer (NEB) – 5µl
- 1 mM dATP – 10 µl
- Taq DNA Polymerase (NEB) – 0.2µl
- H_2O – up to 50µl final reaction volume
- Incubation time – 20 min
- Incubation temperature – 72°C

Step 2: Construction of the sequencing adaptors

Two complementary short ssDNA oligonucleotides were phosphorylated and annealed to form the sequencing adaptor for the PAM-proximal site of the DNA fragments from step 1. One of the oligonucleotides had an additional thymine at its 3' end, in order to facilitate the ligation of the adaptor to the A-tailed fragments.

Adaptor Oligonucleotides phosphorylation (Separate phosphorylation reactions for each oligo)

- 100 µM oligonucleotide stock – 2 µL
- 10X T4 DNA ligase buffer (NEB) – 2 µL
- Sterile MQ water – 15 µL
- T4 Polynucleotide Kinase (NEB) – 1 µL
- Incubation time – 60 min
- Incubation temperature – 37°C
- T4 PNK inactivation – 65°C for 20 min

Annealing of the phosphorylated oligonucleotides

- Oligonucleotide 1 – 5 µL from the corresponding phosphorylation mixture
- Oligonucleotide 1 – 5 µL from the corresponding phosphorylation mixture
- Sterile MQ water – 90 µL
- Incubate the phosphorylated oligos at 95°C for 3 minutes. Cool the reaction slowly at room temperature for ~ 30 min to 1 hr

Step 3: Ligation of the gtCas9-cleaved, A-tailed fragments with the sequencing adaptors

The products of step 1 and 2 were ligated according to the following protocol:

- 10X T4 DNA Ligase Buffer – 2 µl
- Product step 1 – 50 ng
- Product step 2 – 4 ng
- T4 DNA Ligase – 1 µl
- Sterile MQ water – to 20 µl
- Incubation time – 10 min
- Incubation temperature – 20-25°C
- Heat inactivation at 65°C for 10 min

Step 4: PCR amplification of a 150-nucleotides long PAM-containing fragment

5µl from the ligation mixture of step 4 were used as template for PCR amplification using Q5 DNA polymerase (NEB). The oligonucleotide with the thymine extension from step 2 was employed as the forward primer and the reverse primer was designed to anneal 150 nucleotides downstream of the PAM sequence.

The same sequence was amplified using non-gtCas9 treated PAM-library DNA as template. Both PCR products were gel purified and sent for Illumina HiSeq 2500 paired-end sequencing (Baseclear).

Analysis of the sequencing results and determination of the candidate PAM sequences

After analysing the sequencing results the following frequency matrices were constructed. The matrices depict the relative abundance of each nucleotide at every PAM position of the gtCas9 digested and non-digested libraries:

Non-digested	**pos1**	**pos2**	**pos3**	**pos4**	**pos5**	**pos6**	**pos7**
A	19.22	20.83	19.12	24.43	24.59	21.75	18.22
C	34.75	30	31.9	30.54	25.96	27.9	27.17
T	19.16	22.19	25.34	21.28	26.09	26	21.56
G	26.87	26.98	23.64	23.75	23.36	24.35	33.05

Digested	**pos1**	**pos2**	**pos3**	**pos4**	**pos5**	**pos6**	**pos7**
A	10.63	18.65	14.6	14.49	3.36	8.66	27.54
C	**66.22**	**49.59**	**56.82**	**60.35**	**92.4**	**62.26**	34.94
T	8.09	11.21	19.12	12.15	2.35	14.66	5.58
G	15.05	20.54	9.45	13.01	1.89	14.43	31.94

These results indicate a clear preference for targets with cytosine at the 5th PAM position and preference for targets with cytosines at the first 4 PAM positions.

Example 7: *In silico* PAM prediction for gtCas9

In silico predictions of PAMs are possible if enough protospacer sequences are available in genome databases. The *in silico* prediction of gtCas9 PAM started with identification of hits of spacers from the CRISPR array in the genome of *G. thermodenitrificans* T12 strain by comparison to sequences in genome databases such as GenBank. The "CRISPR finder" (http://crispr.u-psud.fr/Server/) tool was used to identify candidate CRISPR loci in T12. The identified CRISPR loci output was then

loaded into "CRISPR target" (http://bioanalysis.otago.ac.nz/CRISPRTarget/crispr_analysis.html) tool, which searches selected databases and provides an output with matching protospacers. These protospacer sequences were then screened for unique hits and for complementarity to spacers – for example, mismatches in the seed sequence were considered to be likely false positive hits and were excluded from further analysis. Hits with identity to prophage sequences and (integrated) plasmids demonstrated that the obtained hits were true positives. Overall, this process yielded 6 single hits (Figure 7). Subsequently, the flanking regions (3' for Type II gtCas nuclease) of the remaining, unique protospacer hits were aligned and compared for consensus sequences using a WebLogo (http://weblogo.berkeley.edu/logo.cgi) (Crooks_GE, Hon_G, Chandonia_JM, Brenner_SE WebLogo: A sequence logo generator, *Genome Research*, 14:1188-1190, (2004)) tool (Figure 8).

The *in silico* results were comparable to the *in vitro* PAM identification experimental results (see Example 6) in which there was a bias for the identity of the 5th residue of the PAM sequence to be a cytosine.

Example 8: Determination of 8 nucleotide long PAM sequences for gtCas9

The *in silico* data from Example 8 suggested that gtCas9 had some preference for adenosine at the 8th position, therefore further PAM determination experiments were carried out where the 8th position of the PAM sequence was also tested. This is consistent with the characterisation of mesophilic *Brevibacillus laterosporus* SSP360D4 (Karvelis *et al.*, 2015) Cas9 PAM sequence which was found to extend between the 5th and the 8th positions at the 3' end of a protospacer.

Specific 8 nucleotide-long sequence variants of the PAMs were trialed with gtCas9:

1) CNCCCCAC [SEQ ID NO: 17],
2) CCCCCCAG [SEQ ID NO: 18],
3) CCCCCCAA [SEQ ID NO: 11],
4) CCCCCCAT [SEQ ID NO: 19],
5) CCCCCCAC [SEQ ID NO: 20],
6) NNNNTNNC (negative control PAM)

After performing an *in vitro* cleavage assay at 60°C targeting these (non-linearized) plasmids with purified gtCas9 and the same sgRNA as before (see Example 6) an increased gtCas9 cleavage activity when the CCCCCCAA [SEQ ID NO: 11] sequence was employed as PAM was observed (Figure 9). However, cleavage activity was clearly detectable for all the tested PAM sequences, even for the negative control PAM sequence a faint cleavage band was observed. Without wishing to be bound to a particular theory, it is possible that use of high gtCas9 concentration contributed to the cleavage observed with the negative control. It has been generally observed that high Cas9 concentrations in *in vitro* assays lead to Cas9-induced DNA cleavage without stringent PAM requirement.

Cas9 concentration in general is known to influence the efficiency of the Cas9 induced DNA cleavage (higher Cas9 concentration results in higher Cas9 activity). This was also observed when performing *in vitro* assays using the targeted plasmid with the CCCCCCAA [SEQ ID NO: 11] PAM sequence and different gtCas9 concentrations (Figure 10)

The targeted plasmid with the CCCCCCAA [SEQ ID NO: 11] PAM sequence for *in vitro* assays as described above was conducted over a wide temperature range between 38 and 78°C (Figure 11). Surprisingly, gtCas9 was active at all the temperatures showing the highest activity between 40.1 and 64.9°C.

Thus the optimal temperature range of Cas9 from *Geobacillus* species is much higher than that of Cas9 proteins which have been characterised to date. Similarly the upper extent of the range in which it retains nuclease activity is much higher than that of known Cas9 proteins. A higher optimal temperature and functional range provides a significant advantage in genetic engineering at high temperatures and therefore in editing the genomes of thermophilic organisms, which have utility in a range of industrial, agricultural and pharmaceutical processes conducted at elevated temperatures.

Example 9: *In vivo* genome editing of *Bacillus smithii* ET138 with gtCas9 and 8 nucleotide length PAM sequences

To confirm that the 8 nucleotide PAMs were also recognised by gtCas9 *in vivo*, an experiment was designed to delete the *pyrF* gene in the genome of *Bacillus smithii* ET138 at 55°C.

This method relies upon providing a homologous recombination template construct in which regions complimentary to the upstream and downstream of the target (*pyrF)* gene are provided to *B. smithii* ET 138 cells. Introduction of the template allows for the process of homologous recombination to be used to introduce the homologous recombination template (with no *pyrF gene)* into the genome such that it also replaces the WT *pyrF* gene in the genome of a cell.

Inclusion of a gtCas9 and a sgRNA in the homologous recombination construct can be used to introduce double stranded DNA breaks (DSDBs) into bacterial genomes that contain WT *pyrF.* DSDBs in a bacterial genome typically results in cell death. Therefore, a sgRNA that recognises a sequence in the WT *pyrF* could result in DSDB and death of cells containing the WT *pyrF* only. Introduction of DSDB is also dependent on a suitable PAM sequence being located downstream at the 3' end of the protospacer that is recognised by gtCas9.

The pNW33n plasmid was used as a backbone to clone:

i) the $cas9_{gt}$ gene under the control of an in-house developed glucose repressible promoter; and

ii) the 1kb upstream and 1kb downstream regions of the *pyrF* gene in the genome of *B. smithii* ET138 as a template for homologous recombination that would result in deletion of the *pyrF* gene from the genome of *B. smithii* ET138; and

iii) single guide RNA (sgRNA) expressing module under the transcriptional control of a constitutive promoter.

Three separate constructs were generated in which the sequence of the single guide RNAs differed at the first 20 nucleotides, which correspond to the sequence that guides the gtCas9 to its specific DNA target in the genome (also known as the spacer). The three different spacer sequences were designed to target three different candidate

protospacers all in the *pyrF* gene of *B. smithii* ET138. The constructs are herein referred to as constructs 1, 2 and 3 respectively.

The three different targeted protospacers had at their 3'-end the following candidate PAM sequences:

1. TCCATTCC (negative control according to the results of the *in vitro* assays; 3'-end of the protospacer targeted by the sgRNA encoded on construct number 3)
2. ATCCCCAA (3'-end of the protospacer targeted by the sgRNA encoded on construct number 1; [SEQ ID NO: 21])
3. ACGGCCAA (3'-end of the protospacer targeted by the sgRNA encoded on construct number 2, [SEQ ID NO: 22])

After transforming *B. smithii* ET 138 cells with one of the three constructs and plating on selection plates, the following results were obtained:

1. When the cells were transformed with the construct targeting the protospacer that had the negative control TCCATTCC PAM sequence at the 3' end (construct number 3) the transformation efficiency was not affected (Figure 12 A). The number of colonies was in the same range as the number of colonies after transformation with the pNW33n positive control construct (Figure 12 B). Of the 15 colonies that were subjected to colony PCR to screen for colonies in which the *pyrF* gene was deleted, none showed the deletion genotype -2.1kb expected band size-, all were wild-type -2.9kb expected band size- (Figure 13). This indicates that the tested PAM was indeed not recognised by the gtCas9 *in vivo*.
2. When the cells were transformed with construct number 1 only a few colonies were obtained (Figure 12 C) when compared to the positive control (cells transformed with pNW33n). 20 colonies were subjected to colony PCR to screen for colonies in which the *pyrF* gene was deleted. The majority (19) of the colonies contained both the wild type and *pyrF* deletion genotype whilst one colonies had a *pyrF* deletion genotype (Figure 14). This result indicated that the PAM sequence ATCCCCAA [SEQ ID NO: 21] is recognised *in vivo* by gtCas9 because no WT only genotypes were observed. The reduced transformation efficiency is also indicative that a proportion of the cell population has been reduced, which could be attributable to cell death caused of WT only genotype cells by DSDB due to successful targeting by gtCas9.

3. When the cells were transformed with construct number 2 no colonies were obtained (Figure 12 D). The lack of colonies is indicative that all of the cell population had been successfully targeted by the gtCas9, which led to cell death caused by DSDB. This suggests that ACGGCCAA [SEQ ID NO: 22] PAM sequence is recognised by gtCas9.

These results indicate that gtCas9 is active at 55°C *in vivo* with the above mentioned PAM sequences, a result that comes in agreement with the *in vitro* PAM determination results. Moreover it can be used as a genome editing tool at the same temperature in combination with a plasmid borne homologous recombination template.

Example 10: ThermoCas9 identification and purification

We recently isolated and sequenced *Geobacillus thermodenitrificans* strain T12, a Gram positive, moderately thermophilic bacterium with an optimal growth temperature at 65°C (Daas *et al. Biotechnol. Biofuels* 9, 210 (2016)). Contrary to previous claims that type II CRISPR-Cas systems are not present in thermophilic bacteria (Li *et al. Nucleic Acids Res.* 44, e34–e34 (2016)), the sequencing results revealed the existence of a type-IIC CRISPR-Cas system in the genome of *G. thermodenitrificans* T12 (Figure 15A). The Cas9 endonuclease of this system (ThermoCas9) was predicted to be relatively small (1082 amino acids) compared to other Cas9 orthologues, such as SpCas9 (1368 amino acids). The size difference is mostly due to a truncated REC lobe, as has been demonstrated for other small Cas9 orthologues (Figure 19)(Ran *et al. Nature* 520, 186–191 (2015)). Furthermore, ThermoCas9 was expected to be active at least around the temperature optimum of *G. thermodenitrificans* T12 (Daas *et al. Biotechnol. Biofuels* 9, 210 (2016)). Using the ThermoCas9 sequence as query, we performed BLAST-P searches in the NCBI/non-redundant protein sequences dataset, and found a number of highly identical Cas9 orthologues (87-99% identity at protein level, Table 1), mostly within the *Geobacillus* genus, supporting the idea that ThermoCas9 is part of a highly conserved defense system of thermophilic bacteria (Figure 15B). These characteristics suggested it may be a potential candidate for exploitation as a genome editing and silencing tool for thermophilic microorganisms, and for conditions at which enhanced protein robustness is required.

We initially performed *in silico* prediction of the crRNA and tracrRNA modules of the *G. thermodenitrificans* T12 CRISPR-Cas system using a previously described approach (Mougiakos *et al. Trends Biotechnol.* 34, 575–587 (2016); Ran *et al. Nature* 520, 186–191 (2015)). Based on this prediction, a 190 nt sgRNA chimera was designed by linking the predicted full size crRNA (30 nt long spacer followed by 36 nt long repeat) and tracrRNA (36 nt long anti-repeat followed by a 88 nt sequence with three predicted hairpin structures). ThermoCas9 was heterologously expressed in *E. coli* and purified to homogeneity. Hypothesizing that the loading of the sgRNA to the ThermoCas9 would stabilize the protein, we incubated purified apo-ThermoCas9 and ThermoCas9 loaded with *in vitro* transcribed sgRNA at 60°C and 65°C, for 15 and 30 min. SDS-PAGE analysis showed that the purified ThermoCas9 denatures at 65°C but not at 60°C, while the denaturation temperature of ThermoCas9-sgRNA complex is above 65°C (Figure 15C). The demonstrated thermostability of ThermoCas9 implied its potential as a thermo-tolerant CRISPR-Cas9 genome editing tool, and encouraged us to analyze some relevant molecular features in more detail.

Table 1. pBLAST results of Cas9 protein sequences from Figure 1B compared to ThermoCas9.

Species	**% identity[a]**
Geobacillus 47C-IIb	99
Geobacillus 46C-IIa	89
Geobacillus LC300	89
Geobacillus jurassicus	89
Geobacillus MAS1	88
Geobacillus stearothermophilus	88
Geobacillus stearothermophilus ATCC 12980	88
Geobacillus Sah69	88
Geobacillus stearothermophilus	88
Geobacillus kaustophilus	88
Geobacillus stearothermophilus	88
Geobacillus genomosp. 3	87

Geobacillus genomosp. 3	87
Geobacillus subterraneus	87
Effusibacillus pohliae	86

Example 11: ThermoCas9 PAM determination

The first step towards the characterization of ThermoCas9 was the *in silico* prediction of its PAM preferences for successful cleavage of a DNA target. We used the 10 spacers of the *G. thermodenitrificans* T12 CRISPR locus to search for potential protospacers in viral and plasmid sequences using CRISPRtarget (Biswas *et al*. *RNA Biol.* 10, 817–827 (2013)). As only two hits were obtained with phage genomes (Figure 20A), it was decided to proceed with an *in vitro* PAM determination approach. We *in vitro* transcribed the predicted sgRNA sequence that contained a spacer for ThermoCas9-based targeting linear dsDNA substrates with a matching protospacer. The protospacer was flanked at its 3'-end by randomized 7-base pair (bp) sequences. After performing ThermoCas9-based cleavage assays at 55°C, the cleaved members of the library (together with a non-targeted library sample as control) were deep-sequenced and compared in order to identify the ThermoCas9 PAM preference (Figure 16A). The sequencing results revealed that ThermoCas9 introduces double stranded DNA breaks that, in analogy to mesophilic Cas9 variants, are located mostly between the 3rd and the 4th PAM proximal nucleotides. Moreover, the cleaved sequences revealed that ThermoCas9 recognizes a 5'-NNNNCNR-3' PAM, with subtle preference for cytosine at the 1st, 3rd, 4th and 6th PAM positions (Figure 16B). Recent studies have revealed the importance of the 8th PAM position for target recognition of certain Type IIC Cas9 orthologues (Karvelis *et al. Genome Biol.* 16, 253 (2015); Kim *et al*. *Genome Res.* 24, 1012–9 (2014)). For this purpose, and taking into account the results from the *in silico* ThermoCas9 PAM prediction, we performed additional PAM determination assays. This revealed optimal targeting efficiency in the presence of an adenine at the 8th PAM position (Figure 16C). Interestingly, despite the limited number of hits, the aforementioned *in silico* PAM prediction (Figure 20B) also suggested the significance of a cytosine at the 5th and an adenine at the 8th PAM positions.

To further clarify the ambiguity of the PAM at the 6th and 7th PAM positions, we generated a set of 16 different target DNA fragments in which the matching

protospacer was flanked by 5'-CCCCCNNA-3' [SEQ ID NO: 13] PAMs. Cleavage assays of these fragments (each with a unique combination of the 6th and 7th nucleotide) were performed in which the different components (ThermoCas9, sgRNA guide, dsDNA target) were pre-heated separately at different temperatures (20, 30, 37, 45, 55 and 60°C) for 10 min before combining and incubating them for 1 hour at the corresponding assay temperature. When the assays were performed at temperatures between 37°C and 60°C, all the different DNA substrates were cleaved (Figure 16D, Figure 21). However, the most digested target fragments consisted of PAM sequences (5th to 8th PAM positions) 5'-CNAA-3' and 5'-CMCA-3', whereas the least digested targets contained a 5'-CAKA-3' PAM. At 30°C, only cleavage of the DNA substrates with the optimal PAM sequences (5th to 8th PAM positions) 5'-CNAA-3' and 5'-CMCA-3' was observed (Figure 2D). Lastly, at 20°C only the DNA substrates with (5th to 8th PAM positions) 5'-CVAA-3' and 5'-CCCA PAM sequences were targeted (Figure 21), making these sequences the most preferred PAMs. These findings demonstrate that at its lower temperature limit, ThermoCas9 only cleaves fragments with a preferred PAM. This characteristic could be exploited during *in vivo* editing processes, for e.g. to avoid off-target effects.

Example 12: Thermostability and truncations

The predicted tracrRNA consists of the anti-repeat region followed by three hairpin structures (Figure 17A). Using the tracrRNA along with the crRNA to form a sgRNA chimera resulted in successful guided cleavage of the DNA substrate. It was observed that a 41-nt long deletion of the spacer distal end of the full-length repeat-anti-repeat hairpin (Figure 17A), most likely better resembling the dual guide's native state, had little to no effect on the DNA cleavage efficiency. The effect of further truncation of the predicted hairpins (Figure 17A) on the cleavage efficiency of ThermoCas9 was evaluated by performing a cleavage time-series in which all the components (sgRNA, ThermoCas9, substrate DNA) were pre-heated separately at different temperatures (37-65°C) for 1, 2 and 5 min before combining and incubating them for 1 hour at various assay temperatures (37-65°C). The number of predicted stem-loops of the tracrRNA scaffold seemed to play a crucial role in DNA cleavage; when all three loops were present, the cleavage efficiency was the highest at all tested temperatures, whereas the efficiency decreased upon removal of the 3' hairpin (Figure 17B). Moreover, the cleavage efficiency drastically dropped upon removal of both the middle

and the 3' hairpins (Figure 22). Whereas pre-heating ThermoCas9 at 65°C for 1 or 2 min resulted in detectable cleavage, the cleavage activity was abolished after 5 min incubation. The thermostability assay showed that sgRNA variants without the 3' stem-loop result in decreased stability of the ThermoCas9 protein at 65°C, indicating that a full length tracrRNA is required for optimal ThermoCas9-based DNA cleavage at elevated temperatures. Additionally, we also varied the lengths of the spacer sequence (from 25 to 18 nt) and found that spacer lengths of 23, 21, 20 and 19 cleaved the targets with the highest efficiency. The cleavage efficiency drops significantly when a spacer of 18 nt is used.

In vivo, the ThermoCas9:sgRNA RNP complex is probably formed within minutes. Together with the above findings, this motivated us to evaluate the activity and thermostability of the RNP. Pre-assembled RNP complex was heated at 60, 65 and 70°C for 5 and 10 min before adding pre-heated DNA and subsequent incubation for 1 hour at 60, 65 and 70°C. Strikingly, we observed that the ThermoCas9 RNP was active up to 70°C, in spite of its pre-heating for 5 min at 70°C (Figure 17C). This finding confirmed our assumption that the ThermoCas9 stability strongly correlates with the association of an appropriate sgRNA guide (Ma et al., *Mol. Cell* 60, 398–407 (2015)).

It would be advantageous in some applications for the ThermoCas9 to have a broad temperature activity range, that is, to be functional at both low and high temperatures. Also, in some circumstances, it would be advantageous if the activity of the ThermoCas9 could be restricted to narrower temperature ranges, for example, active at only low or only high temperatures. Consequently, the ability to manipulate the range of temperatures at which ThermoCas9 is capable of targeted cleavage or binding or at which targeted cleavage or binding takes place efficiently, by modifying structural features of ThermoCas9 or associated elements (such as the sgRNA), would enable a greater level of control to be exerted over nucleic acid sequence manipulation. Hence, we set out to compare the ThermoCas9 temperature range to that of the *Streptococcus pyogenes* Cas9 (SpCas9). Both Cas9 homologues were subjected to *in vitro* activity assays between 20 and 65°C. Both proteins were incubated for 5 min at the corresponding assay temperature prior to the addition of the sgRNA and the target DNA molecules. In agreement with previous analysis[26], the mesophilic SpCas9 was active only between 25 and 44°C (Figure 17D); above these temperature SpCas9 activity rapidly decreased to undetectable levels. In contrast,

ThermoCas9 cleavage activity could be detected between 25 and 65°C (Figure 17D). This indicates the potential to use ThermoCas9 as a genome editing tool for both thermophilic and mesophilic organisms.

Previously characterized, mesophilic Cas9 endonucleases employ divalent cations to catalyze the generation of DSBs in target DNA (Jinek *et al. Science* 337, 816–821 (2012); Chen et al. *J. Biol. Chem.* 289, 13284–13294 (2014)). To evaluate which cations contribute to DNA cleavage by ThermoCas9, plasmid cleavage assays were performed in the presence of one of the following divalent cations: Mg^{2+}, Ca^{2+}, Mn^{2+}, Co^{2+}, Ni^{2+}, and Cu^{2+}; an assay with the cation-chelating agent EDTA was included as negative control. As expected, target dsDNA was cleaved in the presence of divalent cations and remained intact in the presence of EDTA (Figure 23A). Based on reports that certain type-IIC systems were efficient single stranded DNA cutters (Ma *et al. Mol. Cell* 60, 398–407 (2015); Zhang *et al. Mol. Cell* 60, 242–255 (2015)), we tested the activity of ThermoCas9 on ssDNA substrates. However, no cleavage was observed, indicating that ThermoCas9 is a dsDNA nuclease (Figure 23B).

Example 13: ThermoCas9-based gene deletion in the thermophile *B. smithii*

We set out to develop a ThermoCas9-based genome editing tool for thermophilic bacteria. Here, we show a proof of principle using *Bacillus smithii* ET 138 cultured at 55°C. In order to use a minimum of genetic parts, we followed a single plasmid approach. We constructed a set of pNW33n-based pThermoCas9 plasmids containing the *thermocas9* gene under the control of the native *xylL* promoter (P_{xylL}), a homologous recombination template for repairing Cas9-induced double stranded DNA breaks within a gene of interest, and a sgRNA expressing module under control of the constitutive *pta* promoter (P_{pta}) from *Bacillus coagulans* (Figure 4A).

The first goal was the deletion of the full length *pyrF* gene from the genome of *B. smithii* ET 138. The pNW33n-derived plasmids pThermoCas9_bsΔpyrF1 and pThermoCas9_bsΔpyrF2 were used for expression of different ThermoCas9 guides with spacers targeting different sites of the *pyrF* gene, while a third plasmid (pThermoCas9_ctrl) contained a random non-targeting spacer in the sgRNA expressing module. Transformation of *B. smithii* ET 138 competent cells at 55°C with the control plasmids pNW33n (no guide) and pThermoCas9_ ctrl resulted in the

formation of ~200 colonies each. Out of 10 screened pThermoCas9_ ctrl colonies, none contained the Δ*pyrF* genotype, confirming findings from previous studies that homologous recombination in *B. smithii* ET 138 is not sufficient to obtain clean mutants (Mougiakos *et al. ACS Synth. Biol.* 6, 849–861 (2017); Bosma *et al. Microb. Cell Fact.* 14, 99 (2015)). In contrast, transformation with the pThermoCas9_bsΔpyrF1 and pThermoCas9_bsΔpyrF2 plasmids resulted in 20 and 0 colonies respectively, confirming the *in vivo* activity of ThermoCas9 at 55°C and verifying the above described broad *in vitro* temperature range of the protein. Out of ten pThermoCas9_ΔpyrF1 colonies screened, one was a clean Δ*pyrF* mutant whereas the rest had a mixed wild type/Δ*pyrF* genotype (Figure 4B), proving the applicability of the system, as the designed homology directed repair of the targeted pyrF gene was successful. Nonetheless, in the tightly controlled SpCas9-based counter-selection system we previously developed the *pyrF* deletion efficiency was higher (Olson *et al.*, *Curr. Opin. Biotechnol.* 33, 130–141 (2015)). The low number of obtained transformants and clean mutants in the ThermoCas9-based tool can be explained by the low homologous recombination efficiency in *B. smithii* (Olson *et al.*, *Curr. Opin. Biotechnol.* 33, 130–141 (2015)) combined with the constitutive expression of highly active ThermoCas9. It is anticipated that the use of a tightly controllable promoter will increase efficiencies.

Example 14: ThermoCas9-based gene deletion in the mesophile *Pseudomonas putida*

To broaden the applicability of the ThermoCas9-based genome editing tool, and to evaluate whether *in vitro* results could be confirmed *in vivo*, its activity in the mesophilic Gram-negative bacterium *P. putida* KT2440 was evaluated by combining homologous recombination and ThermoCas9-based counter-selection. For this organism, a Cas9-based tool has not been reported to date. Once more, we followed a single plasmid approach. We constructed the pEMG-based pThermoCas9_ppΔpyrF plasmid containing the *thermocas9* gene under the control of the 3-methylbenzoate-inducible Pm-promoter, a homologous recombination template for deletion of the *pyrF* gene and a sgRNA expressing module under the control of the constitutive P3 promoter. After transformation of *P. putida* KT2440 cells and PCR confirmation of plasmid integration, a colony was inoculated in selective liquid medium for overnight culturing at 37°C. The overnight culture was used for inoculation of selective medium and ThermoCas9

expression was induced with 3-methylbenzoate. Subsequently, dilutions were plated on non-selective medium, supplemented with 3-methylbenzoate. For comparison, a parallel experiment without inducing ThermoCas9 expression with 3-methylbenzoate was performed. The process resulted in 76 colonies for the induced culture and 52 colonies for the non-induced control culture. For the induced culture, 38 colonies (50%) had a clean deletion genotype and 6 colonies had mixed wild-type/deletion genotype. On the contrary, only 1 colony (2%) of the non-induced culture had the deletion genotype and there were no colonies with mixed wild-type/deletion genotype retrieved (Figure 24). These results show that ThermoCas9 can be used as an efficient counter-selection tool in the mesophile *P. putida* KT2440 when grown at 37°C.

Example 15: ThermoCas9-based gene silencing

An efficient thermoactive transcriptional silencing CRISPRi tool is currently not available. Such a system could be useful in a number of applications. For example, such a system would greatly facilitate metabolic studies of thermophiles. A catalytically dead variant of ThermoCas9 could serve this purpose by steadily binding to DNA elements without introducing dsDNA breaks. To this end, we identified the RuvC and HNH catalytic domains of ThermoCas9 and introduced the corresponding D8A and H582A mutations for creating a dead (d)ThermoCas9. After confirmation of the designed sequence, Thermo-dCas9 was heterologously produced, purified and used for an *in vitro* cleavage assay with the same DNA target as used in the aforementioned ThermoCas9 assays; no cleavage was observed confirming the catalytic inactivation of the nuclease.

Towards the development of a Thermo-dCas9-based CRISPRi tool, we aimed for the transcriptional silencing of the highly expressed *ldhL* gene from the genome of *B. smithii* ET138. We constructed the pNW33n-based vectors pThermoCas9i_*ldh*L and pThermoCas9i_ctrl. Both vectors contained the *thermo-dCas9* gene under the control of P_{xylL} promoter and a sgRNA expressing module under the control of the constitutive P_{pta} promoter (Figure 4C). The pThermoCas9i_*ldh*L plasmid contained a spacer for targeting the non-template DNA strand at the 5' end of the 138 *ldhL* gene in *B. smithii* ET 138 (Figure S7). The position and targeted strand selection were based on previous studies (Bikard *et al. Nucleic Acids Res.* 41, 7429–7437 (2013); Larson *et al. Nat. Protoc.* 8, 2180–2196 (2013)), aiming for the efficient down-regulation of the *ldhL*

gene. The pThermoCas9i_ctrl plasmid contained a random non-targeting spacer in the sgRNA-expressing module. The constructs were used to transform *B. smithii* ET 138 competent cells at 55°C followed by plating on LB2 agar plates, resulting in equal amounts of colonies. Two out of the approximately 700 colonies per construct were selected for culturing under microaerobic lactate-producing conditions for 24 hours, as described previously (Bosma *et al. Appl. Environ. Microbiol.* 81, 1874–1883 (2015)). The growth of the pThermoCas9i_*ldh*L cultures was 50% less than the growth of the pThermoCas9i_ctrl cultures (Figure 4E). We have previously shown that deletion of the *ldhL* gene leads to severe growth retardation in *B. smithii* ET 138 due to a lack of Ldh-based NAD^+-regenerating capacity under micro-aerobic conditions (Bosma *et al. Microb. Cell Fact.* 14, 99 (2015)). Thus, the observed decrease in growth is likely caused by the transcriptional inhibition of the *ldhL* gene and subsequent redox imbalance due to loss of NAD^+-regenerating capacity. Indeed, HPLC analysis revealed 40% reduction in lactate production of the *ldhL* silenced cultures, and RT-qPCR analysis showed that the transcription levels of the *ldhL* gene were significantly reduced in the pThermoCas9i_*ldh*L cultures compared to the pThermoCas9i_ctrl cultures (Figure 4E).

Example 16: Summary

Most CRISPR-Cas applications are based on RNA-guided DNA interference by Class 2 CRISPR-Cas proteins, such as Cas9 and Cas12a (Komor *et al., Cell* 168, 20–36 (2017); Puchta, *Curr. Opin. Plant Biol.* 36, 1–8 (2017); Xu *et al. J. Genet. Genomics* 42, 141–149 (2015); Tang *et al. Nat. Plants* 3, 17018 (2017); Zetsche *et al. Nat. Biotechnol.* 35, 31–34 (2016); Mougiakos *et al., Trends Biotechnol.* 34, 575–587 (2016)). Prior to this work, no Class 2 CRISPR-Cas immune systems were identified and characterized in thermophilic microorganisms, in contrast to the highly abundant Class 1 CRISPR-Cas systems present in thermophilic bacteria and archaea (Makarova *et al., Nat. Rev. Microbiol.* 13, 722–736 (2015); Weinberger et al., *MBio* 3, e00456-12 (2012)), a few of which have been used for genome editing of thermophiles (Li *et al. Nucleic Acids Res.* 44, e34–e34 (2016)). As a result, the application of CRISPR-Cas technologies was mainly restricted to temperatures below 42°C, due to the mesophilic nature of the employed Cas-endonucleases. Hence, this has excluded application of these technologies in obligate thermophiles and in experimental approaches that require elevated temperatures and/or improved protein stability.

The inventors have characterized ThermoCas9, a Cas9 orthologue from the thermophilic bacterium *G. thermodenitrificans* T12, a strain that we previously isolated from compost (Daas et al., *Biotechnol. Biofuels* 9, 210 (2016)). Data mining revealed additional Cas9 orthologues in the genomes of other thermophiles, which were nearly identical to ThermoCas9, for the first time showing that CRISPR-Cas type II systems do exist in thermophiles, at least in some branches of the *Bacillus* and *Geobacillus* genera. The inventors have shown that ThermoCas9 is active *in vitro* in a wide temperature range of 20-70°C, which is much broader than the 25–44°C range of its mesophilic orthologue SpCas9. The extended activity and stability of ThermoCas9 allows for its application in molecular biology techniques that require DNA manipulation at temperatures of 20-70°C, as well as its exploitation in harsh environments that require robust enzymatic activity. Furthermore, the inventors have identified several factors that are important for conferring the thermostability of ThermoCas9. Firstly, the inventors have demonstrated that the PAM preferences of ThermoCas9 are very strict for activity in the lower part of the temperature range (≤ 30°C), whereas more variety in the PAM is allowed for activity at the moderate to optimal temperatures (37-60°C). Secondly, the inventors have demonstrated that ThermoCas9 activity and thermostability strongly depends on the association with an appropriate sgRNA guide. Without wishing to be bound by any particular theory, the inventors hypothesise that this stabilization of the multi-domain Cas9 protein is most likely the result of a major conformational change from an open/flexible state to a rather compact state, as described for SpCas9 upon guide binding (Jinek *et al. Science* 343, 1247997–1247997 (2014)).

Based on the here described characterization of the novel ThermoCas9, the inventors have successfully developed genome engineering tools for strictly thermophilic prokaryotes. We showed that ThermoCas9 is active *in vivo* at 55°C and 37°C and we adapted the current Cas9-based engineering technologies for the thermophile *B. smithii* ET 138 and the mesophile *P. putida* KT2440. Due to the wide temperature range of ThermoCas9, it is anticipated that the simple, effective and single plasmid-based ThermoCas9 approach will be suitable for a wide range of thermophilic and mesophilic microorganisms that can grow at temperatures from 37°C up to 70°C. This complements the existing mesophilic technologies, allowing their use for a large group of organisms for which these efficient tools were thus far unavailable.

Screening natural resources for novel enzymes with desired traits is unquestionably valuable. Previous studies have suggested that the adaptation of a mesophilic Cas9 orthologue to higher temperatures, with directed evolution and protein engineering, would be the best approach towards the construction of a thermophilic Cas9 protein[29]. Instead, we identified a clade of Cas9 in some thermophilic bacteria, and transformed one of these thermostable ThermoCas9 variants into a powerful genome engineering tool for both thermophilic and mesophilic organisms. With this study, we further stretched the potential of the Cas9-based genome editing technologies and open new possibilities for using Cas9 technologies in novel applications under harsh conditions or requiring activity over a wide temperature range.

Example 17: Materials and Methods

a. Bacterial strains and growth conditions

The moderate thermophile *B. smithii* ET 138 Δ*sigF* Δ*hsdR* (Mougiakos, *et al.*, (2017) ACS Synth. Biol. 6, 849–861) was used for the gene editing and silencing experiments using ThermoCas9. It was grown in LB2 medium (Bosma, *et al. Microb. Cell Fact.* 14, 99 (2015)) at 55°C. For plates, 30 g of agar (Difco) per liter of medium was used in all experiments. If needed chloramphenicol was added at the concentration of 7 μg/mL. For protein expression, *E. coli* Rosetta (DE3) was grown in LB medium in flasks at 37°C in a shaker incubator at 120 rpm until an $OD_{600\ nm}$ of 0.5 was reached after the temperature was switched to 16°C. After 30 min, expression was induced by addition of isopropyl-1-thio-β-d-gal-actopyranoside (IPTG) to a final concentration of 0.5 mM, after which incubation was continued at 16°C. For cloning PAM constructs for 6th and 7th, and 8th positions, DH5-alpha competent *E. coli* (NEB) was transformed according to the manual provided by the manufacturer and grown overnight on LB agar plates at 37°C. For cloning degenerate 7-nt long PAM library, electro-competent DH10B *E. coli* cells were transformed according to standard procedures (Sambrook, Fritsch & Maniatis, T. *Molecular cloning : a laboratory manual*. (Cold Spring Harbor Laboratory, 1989) and grown on LB agar plates at 37°C overnight. *E. coli* DH5α *λpir* (Invitrogen) was used for *P. putida* plasmid construction using the transformation procedure described by Ausubel *et al.* (*Current Protocols in Molecular Biology*. (John Wiley & Sons, Inc., 2001). doi:10.1002/0471142727). For all *E. coli* strains, if required chloramphenicol was used in concentrations of 25 mg/L and kanamycin in 50 mg/L.

Pseudomonas putida KT2440 (DSM 6125) strains were cultured at 37°C in LB medium unless stated otherwise. If required, kanamycin was added in concentrations of 50 mg/L and 3-methylbenzoate in a concentration of 3 mM.

b. ThermoCas9 expression and purification

ThermoCas9 was PCR-amplified from the genome of *G. thermodenitrificans* T12, then cloned and heterologously expressed in *E. coli* Rosetta (DE3) and purified using FPLC by a combination of Ni^{2+}-affinity, anion exchange and gel filtration chromatographic steps. The gene sequence was inserted into plasmid pML-1B (obtained from the UC Berkeley MacroLab, Addgene #29653) by ligation-independent cloning using oligonucleotides (Table 2) to generate a protein expression construct encoding the ThermoCas9 polypeptide sequence (residues 1-1082) fused with an N-terminal tag comprising a hexahistidine sequence and a Tobacco Etch Virus (TEV) protease cleavage site. To express the catalytically inactive ThermoCas9 protein (Thermo-dCas9), the D8A and H582A point mutations were inserted using PCR and verified by DNA sequencing.

Table 2 | Oligonucleotides used in this study.

	Oligo	Sequence	Description	SEQ ID
PAM Library construction	BG6494	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**NNNNNNNCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with 7-nt long random PAM sequence	59
	BG6495	TATGCC***GGATCC***TCAGACCAAGTTTACTCATATATACTTTAGATTGATTTAAAACTTCATTTTTAATTTAAAAGGATCTAG	RV for construction of *in vitro* target DNA sequences	60
	BG7356	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-T-	Adaptor when annealed with BG7357, ligates to A-tailed ThermoCas9 cleaved fragments	61
	BG7357	CTGTCTCTTATACACATCTGACGCTGCCGACGA	Adaptor when annealed with BG7356, ligates to A-tailed ThermoCas9 cleaved fragments	62
	BG7358	TCGTCGGCAGCGTCAG	FW sequencing adaptor for PCR amplification of the ThermoCas9 cleaved fragments	63
	BG7359	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACCATGATTACGCCAAGC	RV sequencing adapter for PCR amplification of the ThermoCas9 cleaved fragments	64
	BG7616	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGGTCATGAGATTATCAAAAAGGATCTTC	RV sequencing adaptor for PCR amplification of the control fragments	65
	BG8157	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCCAGCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM "CCCCCCAG"	66
	BG8158	TATGCC**TCATGAGATTATCAAAAAGGATCTTCA**CCCCCCCAACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM "CCCCCCAA"	67
	BG8159	**TATGCC*TCATGA*GATTATCAAAAAGGATCTTCA**CCCCCCCATCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM "CCCCCCAT"	68
	BG8160	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM "CCCCCCAC"	69
	BG8161	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**NNNNTNNCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM "NNNNTNN"	70

WO 2018/108338 PCT/EP2017/070796

Oligo	Sequence	Description	SEQ ID
BG8363	ACGGTTATCCACAGAATCAG	FW for PCR linearization of PAM identification libraries	71
BG8364	CGGGATTGACTTTTAAAAAAGG	RV for PCR linearization of PAM identification libraries	72
BG8763	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCAAA**CTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "AA"	73
BG8764	**TATGCC*TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCATA**CTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "AT"	74
BG8765	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCAGA**CTAGATCCTTTTAAATTAAAAATG AAGTTTTAAATCAATC	FW for construction of in *vitro* target DNA with PAM position 6&7 "AG"	75
BG8766	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCACA**CTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "AC"	76
BG8767	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCTAA**CTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "TA"	77
BG8768	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCTTA**CTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "TT"	78
BG8769	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCTGA**CTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "TG"	79
BG8770	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCTCA**CTAGATCCTTTTAAATTAAAAATGA AGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "TC"	80
BG8771	TATGCC***TCATGA*****GATTATCAAAAAGGATCTTCA** **CCCCCCGAA**CTAGATCCTTTTAAATTAAAAATG AAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "GA"	81

	Oligo	Sequence	Description	SEQ ID
	BG8772	TATGCC***TCATG***AGATTATCAAAAAGGATCTTCA**C**CCCCCCGTACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "GT"	82
	BG8773	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCGGACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "GG"	83
	BG8774	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCGCACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "GC"	84
	BG8775	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCCAACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "CA"	85
	BG8776	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCCTACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "CT"	86
	BG8777	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCCGACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "CG"	87
	BG8778	TATGCC***TCATGA***GATTATCAAAAAGGATCTTCA**C**CCCCCCCCACTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATC	FW for construction of *in vitro* target DNA with PAM position 6&7 "CC"	88
sgRNA module for *in vitro* transcription	BG6574	AAGCTTGAAATAATACGACTCACTATAGG	FW for PCR amplification of the sgRNA template for the first PAM identification process (30nt long spacer)	89
	BG6576	AAAAAAGACCTTGACGTTTTCC	FW for PCR amplification of the sgRNA template for the first PAM identification process	90
	BG9307	AAGCTTGAAATAATACGACTCACTATAGGTGAGATTATCAAAAAGGATCTTCACGTC	RV for PCR amplification of the sgRNA template for all the PAM identification processes except the first one (25nt long spacer)	91
	BG9309	AAAACGCCTAAGAGTGGGGAATG	RV for PCR amplification of the 3-hairpins long sgRNA template for all the PAM identification processes except the first one	92
	BG9310	AAAAGGCGATAGGCGATCC	RV for PCR amplification of the 2-hairpins long sgRNA template for all the PAM identification processes except the first one	93
	BG9311	AAAACGGGTCAGTCTGCCTATAG	RV for PCR amplification of the 1-hairpin long sgRNA template for all the PAM identification processes except the first one	94

	Oligo	Sequence	Description	SEQ ID
	BG9308	AAGCTTGAAATAATACGACTCACTATAGGTGAGATTATCAAAAAGGATCTTCACGTC	pT7 and 25nt spacer sgRNA Fw	95
	BG10118	AAGCTTGAAATAATACGACTCACTATAGGAGATTATCAAAAAGGATCTTCACGTCA	pT7 and 24nt spacer sgRNA Fw	96
	BG10119	AAGCTTGAAATAATACGACTCACTATAGGAAGATTATCAAAAAGGATCTTCACGTCATAG	pT7 and 23nt spacer sgRNA Fw	97
	BG10120	AAGCTTGAAATAATACGACTCACTATAGGATTATCAAAAAGGATCTTCACGTCATAGT	pT7 and 22nt spacer sgRNA Fw	98
	BG10121	AAGCTTGAAATAATACGACTCACTATAGGAATTATCAAAAAGGATCTTCACGTCATAGTT	pT7 and 21nt spacer sgRNA Fw	99
	BG10122	AAGCTTGAAATAATACGACTCACTATAGGTTATCAAAAAGGATCTTCACGTCATAGTT	pT7 and 20nt spacer sgRNA Fw	100
	BG10123	AAGCTTGAAATAATACGACTCACTATAGGTATCAAAAAGGATCTTCACGTCATAGTTC	pT7 and 19nt spacer sgRNA Fw	101
	BG10124	AAGCTTGAAATAATACGACTCACTATAGGATCAAAAAGGATCTTCACGTCATAGTTC	pT7 and 18nt spacer sgRNA Fw	102
	BG9312	AAAACGCCTAAGAGTGGGGAATGCCCGAAGAAAGCGGGCGATAGGCGATCC	3 loops sgRNA OH Rv	103
Editing and	BG8191	AAGCTTGGCGTAATCATGGTC	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	104
	BG8192	TCATGAGTTCCCATGTTGTG	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	105
	BG8194	tatggcgaatcacaacatgggaactcatgaGAACATCCTCTTTCTTAG	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	106

Oligo	Sequence	Description	SEQ ID
BG8195	gccgatatcaagaccgattttatacttcatTTAAGTTACCTCCTCGATTG	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	107
BG8196	ATGAAGTATAAAATCGGTCTTG	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	108
BG8197	TAACGGACGGATAGTTTC	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	109
BG8198	gaaagccggggaaactatccgtccgttataAATCAGACAAAATGGCCTGCTTATG	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	110
BG8263	gaactatgacactttattttcagaatggacGTATAACGGTATCCATTTTAAGAATAATCC	For the construction of the pThermoCas9_ctrl plasmid	111
BG8268	accgttatacgtccattctgaaaataaagtGTCATAGTTCCCCTGAGAT	For the construction of the pThermoCas9_ctrl plasmid	112
BG8210	aacagctatgaccatgattacgccaagcttCCCTCCCATGCACAATAG	For the construction of the pThermoCas9_ctrl plasmid & pThermoCas9_bsΔpyrF1/2	113
BG8261	gaactatgacatcatggagttttaaatccaGTATAACGGTATCCATTTTAAGAATAATCC	For the construction of the pThermoCas9_bsΔpyrF1	114
BG8266	accgttatactggatttaaaactccatgatGTCATAGTTCCCCTGAGAT	For the construction of the pThermoCas9_bsΔpyrF2	115
BG8317	gaactatgaccacccagcttacatcaacaaGTATAACGGTATCCATTTTTAAGAATAATCC	For the construction of the pThermoCas9_ΔbspyrF2	116
BG8320	accgttatacttgttgatgtaagctgggtgGTCATAGTTCCCCTGAGAT	For the construction of the pThermoCas9_bsΔpyrF2	117
BG9075	CTATCGGCATTACGTCTATC	For the construction of the pThermoCas9i_ctrl	118
BG9076	GCGTCGACTTCTGTATAGC	For the construction of the pThermoCas9i_ctrl	119
BG9091	TGAAGTATAAAATCGGTCTTGCTATCGGCATTACGTCTATC	For the construction of the pThermoCas9i_ctrl	120
BG9092	CAAGCTTCGGCTGTATGGAATCACAGCGTCGACTTCTGTATAGC	For the construction of the pThermoCas9i_ctrl	121
BG9077	GCTGTGATTCCATACAG	For the construction of the pThermoCas9i_ctrl	122

WO 2018/108338 PCT/EP2017/070796

Oligo	Sequence	Description	SEQ ID
BG9267	GGTGCAGTAGGTTGCAGCTATGCTTGTATAACGGTATCCAT	For the construction of the pThermoCas9i_ctrl	123
BG9263	AAGCATAGCTGCAACCTACTGCACCGTCATAGTTCCCCTGAGATTATCG	For the construction of the pThermoCas9i_ctrl	124
BG9088	TCATGACCAAAATCCCTTAACG	For the construction of the pThermoCas9i_ctrl	125
BG9089	TTAAGGGATTTTGGTCATGAGAACATCCTCTTTCTTAG	For the construction of the pThermoCas9i_ctrl	126
BG9090	GCAAGACCGATTTTATACTTCATTTAAG	For the construction of the pThermoCas9i_ctrl	127
BG9548	GGATCCCATGACGCTAGTATCCAGCTGGGTCATAGTTCCCCTGAGATTATCG	For the construction of the pThermoCas9i_ldhL	128
BG9601	TTCAATATTTTTTTTGAATAAAAAATACGATACAATAAAAATGTCTAGAAAAAGATAAAAATG	For the construction of the pThermoCas9i_ldhL	129
BG9600	TTTTTTATTCAAAAAAAATATTGAATTTTAAAAATGATGGTGCTAGTATGAAG	For the construction of the pThermoCas9i_ldhL	130
BG9549	CCAGCTGGATACTAGCGTCATGGGATCCGTATAACGGTATCCATTTTAAGAATAATCC	For the construction of the pThermoCas9i_ldhL	131
BG8552	TCGGGGGTTCGTTTCCCTTG	FW to check genomic *pyrF* deletion KO check	132
BG8553	CTTACACAGCCAGTGACGGAAC	RV to check genomic *pyrF* deletion KO check	133
BG2365	GCCGGCGTCCCGGAAAACGA	For the construction of the pThermoCas9_ppΔpyrF	134
BG2366	GCAGGTCGGGTTCCTCGCATCCATGCCCCCGAACT	For the construction of the pThermoCas9_ppΔpyrF	135
BG2367	ggcttcggaatcgttttccgggacgccggcACGGCATTGGCAAGGCCAAG	For the construction of the pThermoCas9_ppΔpyrF	136
BG2368	gacacaggcatcggtGCAGGGTCTCTTGGCAAGTC	For the construction of the pThermoCas9_ppΔpyrF	137
BG2369	gccaagagaccctgCACCGATGCCTGTGTCGAACC	For the construction of the pThermoCas9_ppΔpyrF	138

	Oligo	Sequence	Description	SEQ ID
	BG2370	cttggcggaaaacgtcaaggtctttttacACGCGCATCAACTTCAAGGC	For the construction of the pThermoCas9_ppΔpyrF	139
	BG2371	atgac**gagctgttcaccagcagcgc**TATTATTGAAGCATTTATCAGGG	For the construction of the pThermoCas9_ppΔpyrF	140
	BG2372	GTAAAAAAGACCTTGACGTTTTC	For the construction of the pThermoCas9_ppΔpyrF	141
	BG2373	tatgaagcgggccatTTGAAGACGAAAGGGCCTC	For the construction of the pThermoCas9_ppΔpyrF	142
	BG2374	taata**gcgctgctggtgaacagctc**GTCATAGTTCCCCTGAGATTATCG	For the construction of the pThermoCas9_ppΔpyrF	143
	BG2375	tggagtcatgaacatATGAAGTATAAAATCGGTCTTG	For the construction of the pThermoCas9_ppΔpyrF	144
	BG2376	ccctttcgtcttcAAATGGCCCGCTTCATAAGCAG	For the construction of the pThermoCas9_ppΔpyrF	145
	BG2377	gattttatacTTCATATGTTCATGACTCCATTATTATTG	For the construction of the pThermoCas9_ppΔpyrF	146
	BG2378	gggggcatggatgCGAGGAACCCGACCTGCATTGG	For the construction of the pThermoCas9_ppΔpyrF	147
	BG2381	ACACGGCGGATGCACTTACC	FW for confirmation of plasmid integration and *pyrF* deletion in *P. putida*	148
	BG2382	TGGACGTGTACTTCGACAAC	RV for confirmation of *pyrF* deletion in *P. putida*	149
	BG2135	ACACGGCGGATGCACTTACC	RV for confirmation of plasmid integration in *P. putida*	150
Sequencing primers	BG8196	TGGACGTGTACTTCGACAAC	*thermocas9* seq. 1	151
	BG8197	TAACGGACGGATAGTTTC	*thermocas9* seq. 2	152
	BG6850	GCCTCATGAATGCAGCGATGGTCCGGTGTTC	*pyr*F US	153
	BG6849	GCCTCATGAGTTCCCATGTTGTGATTC	*pyrF* DS	154

	Oligo	Sequence	Description	SEQ ID
	BG6769	CAATCCAACTGGGCTTGAC	*thermocas9* seq. 3	155
	BG6841	CAAGAACTTTATTGGTATAG	*thermocas9* seq. 4	156
	BG6840	TTGCAGAAATGGTTGTCAAG	*thermocas9* seq. 5	157
	BG9215	GAGATAATGCCGACTGTAC	pNW33n backbone seq. 1	158
	BG9216	AGGGCTCGCCTTTGGGAAG	pNW33n backbone seq. 2	159
	BG9505	GTTGCCAACGTTCTGAG	*thermocas9* seq. 6	160
	BG9506	AATCCACGCCGTTTAG	*thermocas9* seq. 7	161
Cleavage assay	BG8363	ACGGTTATCCACAGAATCAG	FW for PCR linearization of DNA target	162

	Oligo	Sequence	Description	SEQ ID
	BG8364	CGGGATTGACTTTTAAAAAAGG	RV for PCR linearization of DNA target	163
	BG9302	AAACTTCATTTTTAATTTAAAAGGATCTAGAACCCCCCGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAA	Non-template strand oligonucleotide for ssDNA cleavage assays	164
	BG9303	TTTCTACGGGGTCTGACGCTCAGTGGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCCCCCCAACTAGATCCTTTTAAATTAAAAATGAAGTTT	Template strand oligonucleotide for ssDNA cleavage assays	165
	BG9304	TTTCTACGGGGTCTGACGCTCAGTGGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACGGGGGGTTCTAGATCCTTTTAAATTAAAAATGAAGTTT	Template strand oligonucleotide for ssDNA cleavage assays	166
ThermoCas9 expression and RT-qPCR	BG7886	TACTTCCAATCCAATGCAAAGTATAAAATCGGTCTTGATATCG	FW LIC_*thermocas9*	167
	BG7887	TTATCCACTTCCAATGTTATTATAACGGACGGATAGTTTCCCCGGCTTTC	RV LIC_*thermocas9*	168
ThermoCas9 expression	BG9665	ATGACGAAAGGAGTTTCTTATTATG	RV qPCR check *ldhl*	169
	BG9666	AACGGTATTCCGTGATTAAG	FW qPCR check *ldhl*	170

Restriction sites are shown in italics. The PAMs are colored red. Spacer regions are shown in bold. Nucleotides in lowercase letters correspond to primer overhangs for HiFi DNA Assembly. LIC: Ligase Independent cloning; FW: Forward primer; RV: Reverse primer.

The proteins were expressed in *E. coli* Rosetta 2 (DE3) strain. Cultures were grown to an OD_{600nm} of 0.5-0.6. Expression was induced by the addition of IPTG to a final concentration of 0.5 mM and incubation was continued at 16°C overnight. Cells were harvested by centrifugation and the cell pellet was resuspended in 20 mL of Lysis Buffer (50 mM sodium phosphate pH 8, 500 mM NaCl, 1mM DTT, 10 mM imidazole) supplemented with protease inhibitors (Roche cOmplete, EDTA-free) and lysozyme. Once homogenized, cells were lysed by sonication (Sonoplus, Bandelin) using a using an ultrasonic MS72 microtip probe (Bandelin), for 5-8 minutes consisting of 2s pulse and 2.5s pause at 30% amplitude and then centrifuged at 16,000×*g* for 1 hour at 4°C to remove insoluble material. The clarified lysate was filtered through 0.22 micron filters (Mdi membrane technologies) and applied to a nickel column (Histrap HP, GE Lifesciences), washed and then eluted with 250 mM imidazole. Fractions containing ThermoCas9 were pooled and dialyzed overnight into the dialysis buffer (250 mM KCl, 20 mM HEPES/KOH, and 1 mM DTT, pH 7.5). After dialysis, sample was diluted 1:1 in 10 mM HEPES/KOH pH 8, and loaded on a heparin FF column pre-equilibrated in IEX-A buffer (150 mM KCl, 20 mM HEPES/KOH pH 8). Column was washed with IEX-A and then eluted with a gradient of IEX-C (2M KCl, 20 mM HEPES/KOH pH 8). The sample was concentrated to 700 µL prior to loading on a gel filtration column (HiLoad 16/600 Superdex 200) via FPLC (AKTA Pure). Fractions from gel filtration were analysed by SDS-PAGE; fractions containing ThermoCas9 were pooled and concentrated to 200 µL (50 mM sodium phosphate pH 8, 2 mM DTT, 5% glycerol, 500 mM NaCl) and either used directly for biochemical assays or frozen at -80°C for storage.

c. *In vitro* synthesis of sgRNA

The sgRNA module was designed by fusing the predicted crRNA and tracrRNA sequences with a 5'-GAAA-3' linker. The sgRNA-expressing DNA sequence was put under the transcriptional control of the T7 promoter. It was synthesized (Baseclear, Leiden, The Netherlands) and provided in the pUC57 backbone. All sgRNAs used in the biochemical reactions were synthesized using the HiScribe™ T7 High Yield RNA Synthesis Kit (NEB). PCR fragments coding for sgRNAs, with the T7 sequence on the 5' end, were utilized as templates for in vitro transcription reaction. T7 transcription

was performed for 4 hours. The sgRNAs were run and excised from urea-PAGE gels and purified using ethanol precipitation.

d. *In vitro* cleavage assay

In vitro cleavage assays were performed with purified recombinant ThermoCas9. ThermoCas9 protein, the *in vitro* transcribed sgRNA and the DNA substrates (generated using PCR amplification using primers described in Table 2) were incubated separately (unless otherwise indicated) at the stated temperature for 10 min, followed by combining the components together and incubating them at the various assay temperatures in a cleavage buffer (100 mM sodium phosphate buffer (pH=7), 500 mM NaCl, 25 mM $MgCl_2$, 25 (V/V%) glycerol, 5 mM dithiothreitol (DTT)) for 1 hour. Each cleavage reaction contained 160 nM of ThermoCas9 protein, 4 nM of substrate DNA, and 150 nM of synthetized sgRNA. Reactions were stopped by adding 6x loading dye (NEB) and run on 1.5% agarose gels. Gels were stained with SYBR safe DNA stain (Life Technologies) and imaged with a Gel DocTM EZ gel imaging system (Bio-rad).

e. Library construction for *in vitro* PAM screen

For the construction of the PAM library, a 122-bp long DNA fragment, containing the protospacer and a 7-bp long degenerate sequence at its 3'-end, was constructed by primer annealing and Klenow fragment (exo-) (NEB) based extension. The PAM-library fragment and the pNW33n vector were digested by BspHI and BamHI (NEB) and then ligated (T4 ligase, NEB). The ligation mixture was transformed into electro-competent *E. coli* DH10B cells and plasmids were isolated from liquid cultures. For the 7nt-long PAM determination process, the plasmid library was linearized by SapI (NEB) and used as the target. For the rest of the assays the DNA substrates were linearized by PCR amplification.

f. PAM screening assay

The PAM screening of thermoCas9 was performed using *in vitro* cleavage assays, which consisted of (per reaction): 160 nM of ThermoCas9, 150 nM in vitro transcribed sgRNA, 4 nM of DNA target, 4 µl of cleavage buffer (100 mM sodium phosphate buffer pH 7.5, 500 mM NaCl, 5 mM DTT, 25% glycerol) and MQ water up to 20 µl final reaction volume. The PAM containing cleavage fragments from the 55°C reactions

were gel purified, ligated with Illumina sequencing adaptors and sent for Illumina HiSeq 2500 sequencing (Baseclear). Equimolar amount of non-thermoCas9 treated PAM library was subjected to the same process and sent for Illumina HiSeq 2500 sequencing as a reference. HiSeq reads with perfect sequence match to the reference sequence were selected for further analysis. From the selected reads, those present more than 1000 times in the ThermoCas9 treated library and at least 10 times more in the ThermoCas9 treated library compared to the control library were employed for WebLogo analysis (Crooks *et al.*, *Genome Res.* 14, 1188–1190 (2004)).

g. Editing and silencing constructs for *B. smithii* and *P. putida*

All the primers and plasmids used for plasmid construction were designed with appropriate overhangs for performing NEBuilder HiFi DNA assembly (NEB), and they are listed in Table 2 and 3 respectively. The fragments for assembling the plasmids were obtained through PCR with Q5 Polymerase (NEB) or Phusion Flash High-Fidelity PCR Master Mix (ThermoFisher Scientific), the PCR products were subjected to 1% agarose gel electrophoresis and they were purified using Zymogen gel DNA recovery kit (Zymo Research). The assembled plasmids were transformed to chemically competent *E. coli* DH5α cells (NEB), or to *E. coli* DH5α λpir (Invitrogen) in the case of *P. putida* constructs, the latter to facilitate direct vector integration. Single colonies were inoculated in LB medium, plasmid material was isolated using the GeneJet plasmid miniprep kit (ThermoFisher Scientific) and sequence verified (GATC-biotech) and 1µg of each construct transformed of *B. smithii* ET 138 electro-competent cells, which were prepared according to a previously described protocol (Bosma, *et al. Microb. Cell Fact.* 14, 99 (2015)). The MasterPure™ Gram Positive DNA Purification Kit (Epicentre) was used for genomic DNA isolation from *B. smithii* and *P. putida* liquid cultures.

For the construction of the pThermoCas9_ctrl, pThermoCas9_bsΔpyrF1 and pThermoCas9_bsΔpyrF2 vectors, the pNW33n backbone together with the Δ*pyrF* homologous recombination flanks were PCR amplified from the pWUR_Cas9sp1_hr vector (Mougiakos, *et al. ACS Synth. Biol.* 6, 849–861 (2017)) (BG8191and BG8192). The native P_{xylA} promoter was PCR amplified from the genome of *B. smithii* ET 138 (BG8194 and BG8195). The *thermocas9* gene was PCR amplified from the genome of *G. thermodenitrificans* T12 (BG8196 and BG8197). The P_{pta} promoter was PCR

amplified from the pWUR_Cas9sp1_hr vector (Mougiakos, *et al. ACS Synth. Biol.* 6, 849–861 (2017)) (BG8198 and BG8261_2/BG8263_nc2/ BG8317_3). The spacers followed by the sgRNA scaffold were PCR amplified from the pUC57_T7t12sgRNA vector (BG8266_2/BG8268_nc2/8320_3 and BG8210).

A four-fragment assembly was designed and executed for the construction of the pThermoCas9i_ldhL vectors. Initially, targeted point mutations were introduced to the codons of the *thermocas9* catalytic residues (mutations D8A and H582A), through a two-step PCR approach using pThermoCas9_ctrl as template. During the first PCR step (BG9075, BG9076), the desired mutations were introduced at the ends of the produced PCR fragment and during the second step (BG9091, BG9092) the produced fragment was employed as PCR template for the introduction of appropriate assembly-overhangs. The part of the *thermocas9* downstream the second mutation along with the *ldhL* silencing spacer was PCR amplified using pThermoCas9_ctrl as template (BG9077 and BG9267). The sgRNA scaffold together with the pNW33n backbone was PCR amplified using pThermoCas9_ctrl as template (BG9263 and BG9088). The promoter together with the part of the *thermocas9* upstream the first mutation was PCR amplified using pThermoCas9_ctrl as template (BG9089, BG9090)

A two-fragment assembly was designed and executed for the construction of pThermoCas9i_ctrl vector. The spacer sequence in the pThermoCas9i_ldhL vector was replaced with a random sequence containing BaeI restriction sites at both ends. The sgRNA scaffold together with the pNW33n backbone was PCR amplified using pThermoCas9_ctrl as template (BG9548, BG9601). The other half of the construct consisted of *Thermo-dCas9* and promoter was amplified using pThermoCas9i_ldhL as template (BG9600, BG9549).

A five-fragment assembly was designed and executed for the construction of the *P. putida* KT2440 vector pThermoCas9_ppΔpyrF. The replicon from the suicide vector pEMG was PCR amplified (BG2365, BG2366). The flanking regions of *pyrF* were amplified from KT2440 genomic DNA (BG2367, BG2368 for the 576-bp upstream flank, and BG2369, BG2370 for the 540-bp downstream flank). The flanks were fused in an overlap extension PCR using primers BG2367 and BG2370 making use of the overlaps of primers BG2368 and BG2369. The sgRNA was amplified from the pThermoCas9_ctrl plasmid (BG2371, BG2372). The constitutive P3 promoter was

amplified from pSW_I-SceI (BG2373, BG2374). This promoter fragment was fused to the sgRNA fragment in an overlap extension PCR using primers BG2372 and BG2373 making use of the overlaps of primers BG2371 and BG2374. ThermoCas9 was amplified from the pThermoCas9_ctrl plasmid (BG2375, BG2376). The inducible Pm-XylS system, to be used for 3-methylbenzoate induction of ThermoCas9 was amplified from pSW_I-SceI (BG2377, BG2378).

Table 3. Plasmids used in this study

Plasmid	Description	Restriction sites used	Primers	Source
pNW33n	*E. coli-Bacillus* shuttle vector, cloning vector, Cam^R	-	-	BGSC
pUC57_T7sgRNAfull	pUC57 vector containing DNA encoding the sgRNA under the control of T7 promoter; serves as a template for *in vitro* transcription of full length Repeat/Antirepeat sgRNAs			Baseclear
pMA2_T7sgRNAtruncated R/AR	Vector containing DNA encoding the truncated Repeat/Antirepeat part of the sgRNA under the control of T7 promoter; serves as a template for *in vitro* transcription of truncated Repeat/Antirepeat sgRNAs	-	-	Gen9
pRARE	T7 RNA polymerase based expression vector, Kan^R	-	-	EMD Millipore
pML-1B	*E. coli* Rosetta™ (DE3) plasmid, encodes rare tRNAs, Cam^R	-	-	Macrolab, Addgene
pEMG	*P. putida* suicide vector, used as template for replicon and Kan^R		See Table 2	1
pSW_I-SceI	*P. putida* vector containing *I-SceI*, used as template for *xylS* and P_{Pm}		See Table 2	1
pWUR_Cas9sp1_hr	pNW33n with spCas9-module containing spacer targeting the *pyrF* gene. This plasmid was used as a template for constructing the ThermoCas9 based constructs	-	-	[2]
pThermo_Cas9	*thermocas9* with N-term. His-tag and TEV cleavage site in pML-1B. Expression vector for ThermoCas9	SspI and Ligase Independent Cloning	BG7886 and BG7887	This study
pThermo_dCas9	cas9*dthermo*cas9 with N-term. His-tag and TEV cleavage site in pML-1B. Expression vector for catalytically inactive (dead) dThermoCas9	SspI and Ligase Independent Cloning	BG7886 and BG7888	This study

Plasmid	Description	Restriction sites used	Primers	Source
pNW-PAM7nt	Target sequence in pNW33n vector containing a 7-nt degenerate PAM for *in vitro* PAM determination assay	BamHI and BspHI	See Table 2	This study
pNW63-pNW78	Target sequence in pNW33n vector containing distinct nucleotides at the 6th and 7th positions of the PAM (CCCCCNNA)	BamHI and BspHI	See Table 2	This study
pThermoCas9_ctrl	pNW33n with ThermoCas9-module[1] containing a non-targeting spacer. Used as a negative control	-	See Table 2	This study
pThermoCas9_bsΔpyrF1	pNW33n with ThermoCas9-module[1] containing spacer 1 targeting the *pyrF* gene and the fused us+ds *pyrF*-flanks	-	See Table 2	This study
pThermoCas9_bsΔpyrF2	pNW33n with ThermoCas9-module[1] containing spacer 2 targeting the *pyrF* gene and the fused us+ds *pyrF*-flanks	-	See Table 2	This study
pThermoCas9i_ctrl	pNW33n with Thermo-dCas9-module[2] containing a non-targeting spacer. Used as a wild-type control	-	See Table 2	This study
pThermoCas9i_ldhL	pNW33n with Thermo-dCas9-module[2] containing spacer 2 targeting the *ldhL* gene	-	See Table 2	This study
pThermoCas9_ppΔpyrF	pEMG with ThermoCas9-module[3] for *Pseudomonas putida* containing a spacer targeting the a spacer targeting the *pyrF* gene and the fused us+ds *pyrF*-flanks	-	See Table 2	This study

[1] The ThermoCas9 module contains *thermocas9* under the native P_{xylL} promoter followed by the sgRNA under the *B. coagulans* P_{pta} promoter (Figure 4).

[2] Like the ThermoCas9 module, but with the *thermo-dCas9* instead of *thermocas9* (Figure 4).

[3] The ThermoCas9 module for *Pseudomonas putida* contains *thermocas9* under the transcriptional control of the inducible Pm-XylS system followed by the sgRNA under the constitutive P3 promoter.

h. Editing protocol for *P. putida*

Transformation of the plasmid to *P. putida* was performed according to Choi *et al.* (Choi et al., *J. Microbiol. Methods* 64, 391–397 (2006)). After transformation and selection of integrants, overnight cultures were inoculated. 10µl of overnight culture was used for inoculation of 3ml fresh selective medium and after 2 hours of growth at 37°C ThermoCas9 was induced with 3-methylbenzoate. After an additional 6h, dilutions of the culture were plated on non-selective medium supplemented with 3-methylbenzoate. For the control culture the addition of 3-methylbenzoate was omitted in all the steps. Confirmation of plasmid integration in the *P. putida* chromosome was done by colony PCR with primers BG2381 and BG2135. Confirmation of *pyrF* deletion was done by colony PCR with primers BG2381 and BG2382.

i. RNA isolation

RNA isolation was performed by the phenol extraction based on a previously described protocol (van Hijum *et al. BMC Genomics* 6, 77 (2005)). Overnight 10 mL cultures were centrifuged at 4°C and 4816×*g* for 15 min and immediately used for RNA isolation. After removal of the medium, cells were suspended in 0.5 mL of ice-cold TE buffer (pH 8.0) and kept on ice. All samples were divided into two 2mL screw-capped tubes containing 0.5 g of zirconium beads, 30 µL of 10% SDS, 30 µL of 3 M sodium acetate (pH 5.2), and 500 µL of Roti-Phenol (pH 4.5−5.0, Carl Roth GmbH). Cells were disrupted using a FastPrep-24 apparatus (MP Biomedicals) at 5500 rpm for 45 s and centrifuged at 4°C and 10 000 rpm for 5 min. 400 µL of the water phase from each tube was transferred to a new tube, to which 400 µL of chloroform−isoamyl alcohol (Carl Roth GmbH) was added, after which samples were centrifuged at 4 °C and 18 400×*g* for 3 min. 300 µL of the aqueous phase was transferred to a new tube and mixed with 300 µL of the lysis buffer from the high pure RNA isolation kit (Roche). Subsequently, the rest of the procedure from this kit was performed according to the manufacturer's protocol, except for the DNase incubation step, which was performed for 45 min. The concentration and integrity of cDNA was determined using Nanodrop-1000 Integrity and concentration of the isolated RNA was checked on a NanoDrop 1000.

j. Quantification of mRNA by RT-qPCR

First-strand cDNA synthesis was performed for the isolated RNA using SuperScriptTM III Reverse Transcriptase (Invitrogen) according to manufacturer's protocol. qPCR was performed using the PerfeCTa SYBR Green Supermix for iQ from Quanta Biosciences. 40 ng of each cDNA library was used as the template for qPCR. Two sets of primers were used; BG9665:BG9666 amplifying a 150-nt long region of the *ldhL* gene and BG9889:BG9890 amplifying a 150-nt long sequence of the *rpoD* (RNA polymerase sigma factor) gene which was used as the control for the qPCR. The qPCR was run on a Bio-Rad C1000 Thermal Cycler.

k. HPLC

A high-pressure liquid chromatography (HPLC) system ICS-5000 was used for lactate quantification. The system was operated with Aminex HPX 87H column from Bio-Rad Laboratories and equipped with a UV1000 detector operating on 210 nm and a RI-150 40°C refractive index detector. The mobile phase consisted of 0.16 N H_2SO_4 and the column was operated at 0.8 mL/min. All samples were diluted 4:1 with 10 mM DMSO in 0.01 N H_2SO_4.

The following section of the description consists of numbered paragraphs simply providing statements of the invention already described herein. The numbered paragraphs in this section are not claims. The claims are set forth below in the later section headed “claims”.

1. An isolated clustered regularly interspaced short palindromic repeat (CRISPR)-associated (Cas) protein or polypeptide comprising;

a. the amino acid motif EKDGKYYC [SEQ ID NO: 2]; and/or

b. the amino acid motif $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and/or

c. the amino acid motif X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and/or

d. the amino acid motif X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and/or

e. the amino acid motif $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine;

wherein the Cas protein is capable of nucleic acid cleavage between 50°C and 100°C when associated with at least one targeting RNA molecule, and a polynucleotide comprising a target nucleic acid sequence recognised by the targeting RNA molecule.

2. An isolated Cas protein or polypeptide fragment having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 77% identity therewith, wherein the Cas protein is capable of binding, cleaving, modifying or marking a polynucleotide comprising a target nucleic acid sequence at a temperature between

50°C and 100°C when associated with at least one RNA molecule which recognizes the target sequence.

3. A Cas protein or polypeptide fragment as in numbered paragraph 1 or 2, wherein the Cas protein or fragment is capable of nucleic acid binding, cleavage, marking or modification at a temperature between 50°C and 75°C, preferably at a temperature above 60°C; more preferably at a temperature between 60°C and 80°C; more preferably at a temperature between 60°C and 65°C.

4. A Cas protein or polypeptide fragment as in any of numbered paragraphs 1 to 3, wherein the nucleic acid binding, cleavage, marking or modification is DNA cleavage.

5. A Cas protein or polypeptide fragment as in any preceding numbered paragraph, wherein the amino acid sequence comprises an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 77% identity therewith.

6. A Cas protein or polypeptide fragment as in any preceding numbered paragraph, wherein the Cas protein is obtainable from a bacterium, archaeon or virus.

7. A Cas protein or polypeptide fragment as in any preceding numbered paragraph, wherein the Cas protein is obtainable from *Geobacillus* sp., preferably from *Geobacillus thermodenitrificans*.

8. A ribonucleoprotein complex comprising a Cas protein as in any preceding numbered paragraph, and comprising at least one targeting RNA molecule which recognises a sequence in a target polynucleotide.

9. A ribonucleoprotein complex as in numbered paragraph 8, wherein the targeting RNA molecule comprises a crRNA and optionally a tracrRNA.

10. A ribonucleoprotein complex as in any of numbered paragraphs 7 to 9, wherein the length of the at least one RNA molecule is in the range 35 – 135 nucleotide residues.

11. A ribonucleoprotein complex as in numbered paragraph 8 or 9, wherein the target sequence is 31 or 32 nucleotide residues in length.

12. A Cas protein or polypeptide as in any of numbered paragraphs 1 to 7 or a ribonucleoprotein complex as in any of 8 to 11, wherein the protein or polypeptide is provided as part of a protein complex comprising at least one further functional or non-functional protein.

13. A Cas protein, polypeptide, or ribonucleoprotein complex as in numbered paragraph 12, wherein the Cas protein or polypeptide, and/or the at least one further protein further comprise at least one functional moiety.

14. A Cas protein or polypeptide, or ribonucleoprotein complex as in numbered paragraph 13, wherein the at least one functional moiety is fused or linked to the N-terminus and/or the C-terminus of the Cas protein, polypeptide or ribonucleoprotein complex; preferably the N-terminus.

15. A Cas protein or polypeptide, or a ribonucleoprotein complex as in numbered paragraph 13 or 14, wherein the at least one functional moiety is a protein; optionally

selected from a helicase, a nuclease, a helicase-nuclease, a DNA methylase, a histone methylase, an acetylase, a phosphatase, a kinase, a transcription (co-)activator, a transcription repressor, a DNA binding protein, a DNA structuring protein, a marker protein, a reporter protein, a fluorescent protein, a ligand binding protein, a signal peptide, a subcellular localisation sequence, an antibody epitope or an affinity purification tag.

16. A Cas protein or polypeptide, or a ribonucleoprotein complex as in numbered paragraph 15, wherein the native activity of the Cas9 nuclease activity is inactivated and the Cas protein is linked to at least one functional moiety.

17. A Cas protein or polypeptide, or a ribonucleoprotein complex as in numbered paragraph 15 or 16, wherein the at least one functional moiety is a nuclease domain; preferably a FokI nuclease domain.

18. A Cas protein or polypeptide, or a ribonucleoprotein complex as in any of numbered paragraphs 15 to 17, wherein the at least one functional moiety is a marker protein, for example GFP.

19. An isolated nucleic acid molecule encoding a Cas protein or polypeptide, comprising;

a. the amino acid motif EKDGKYYC [SEQ ID NO: 2]; and/or

b. the amino acid motif $X_1X_2CTX_3X_4$ [SEQ ID NO: 3] wherein X_1 is independently selected from Isoleucine, Methionine or Proline, X_2 is independently selected from Valine, Serine, Asparagine or Isoleucine, X_3 is independently selected from Glutamate or Lysine and X_4 is one of Alanine, Glutamate or Arginine; and/or

c. the amino acid motif X_5LKX_6IE [SEQ ID NO: 4] wherein X_5 is independently selected from Methionine or Phenylalanine and X_6 is independently selected from Histidine or Asparagine; and/or

d. the amino acid motif X_7VYSX_8K [SEQ ID NO: 5] wherein X_7 is Glutamate or Isoleucine and X_8 is one of Tryptophan, Serine or Lysine; and/or

e. the amino acid motif $X_9FYX_{10}X_{11}REQX_{12}KEX_{13}$ [SEQ ID NO: 6] wherein X_9 is Alanine or Glutamate, X_{10} is Glutamine or Lysine, X_{11} is Arginine or Alanine, X_{12} is Asparagine or Alanine and X_{13} is Lysine or Serine;

wherein the Cas protein or polypeptide is capable of DNA binding, cleavage, marking or modification between 50°C and 100°C when associated with at least one targeting RNA molecule, and a polynucleotide comprising a target nucleic acid sequence recognised by the targeting RNA molecule.

20. An isolated nucleic acid molecule encoding a clustered regularly interspaced short palindromic repeat (CRISPR)-associated (Cas) protein having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 77% identity therewith; or a polypeptide fragment thereof.

21. An isolated nucleic acid molecule as in numbered paragraph 19 or 20, further comprising at least one nucleic acid sequence encoding an amino acid sequence which upon translation is fused with the Cas protein or polypeptide.

22. An isolated nucleic acid molecule as in numbered paragraph 21, wherein the at least one nucleic acid sequence fused to the nucleic acid molecule encoding the Cas protein or polypeptide, encodes a protein selected from a protein selected from a helicase, a nuclease, a helicase-nuclease, a DNA methylase, a histone methylase, an acetylase, a phosphatase, a kinase, a transcription (co-)activator, a transcription repressor, a DNA binding protein, a DNA structuring protein, a marker protein, a

reporter protein, a fluorescent protein, a ligand binding protein, a signal peptide, a subcellular localisation sequence, an antibody epitope or an affinity purification tag.

23. An expression vector comprising a nucleic acid molecule as in any of numbered paragraphs 19 to 22.

24. An expression vector as in numbered paragraph 23, further comprising a nucleotide sequence encoding at least one targeting RNA molecule.

25. A method of modifying a target nucleic acid comprising contacting the nucleic acid with:

a. a ribonucleoprotein complex of any of numbered paragraphs 6 to 11; or

b. a protein or protein complex of any of numbered paragraphs 12 to 18 and at least one targeting RNA molecule as defined in any of numbered paragraphs 6 to 11; and wherein said method is not used in human cells.

26. A method of modifying a target nucleic acid in a non-human cell, comprising transforming, transfecting or transducing the cell with an expression vector of numbered paragraph 24; or alternatively transforming, transfecting or transducing the cell with an expression vector of numbered paragraph 23 and a further expression vector comprising a nucleotide sequence encoding a targeting RNA molecule as defined in any of numbered paragraphs 6 to 11

27. A method of modifying a target nucleic acid in a non-human cell comprising transforming, transfecting or transducing the cell with an expression vector of numbered paragraph 23, and then delivering a targeting RNA molecule as defined in any of numbered paragraphs 6 to 11 to or into the cell.

28. A method of modifying a target nucleic acid as in any of numbered paragraphs 25 to 28, wherein the at least one functional moiety is a marker protein or reporter protein and the marker protein or reporter protein associates with the target nucleic acid; preferably wherein the marker is a fluorescent protein, for example a green fluorescent protein (GFP).

29. A method as in any of numbered paragraphs 25 to 28, wherein the target nucleic acid is DNA; preferably dsDNA.

30. A method as in any of numbered paragraphs 25 to 28, wherein the target nucleic acid is RNA.

31. A method of modifying a target nucleic acid as in numbered paragraph 29, wherein the nucleic acid is dsDNA, the at least one functional moiety is a nuclease or a helicase-nuclease, and the modification is a single-stranded or a double-stranded break at a desired locus.

32. A method of silencing gene expression at a desired locus according to any of the methods in any of numbered paragraphs 26, 27, 29 or 31.

33. A method of modifying or deleting and/or inserting a desired nucleotide sequence at a desired location according to any of the methods as in any of numbered paragraphs 26, 27, 29 or 31.

34. A method of modifying gene expression in a non-human cell comprising modifying a target nucleic acid sequence as in a method of any of numbered

paragraphs 25 to 29; wherein the nucleic acid is dsDNA and the functional moiety is selected from a DNA modifying enzyme (e.g. a methylase or acetylase), a transcription activator or a transcription repressor.

35. A method of modifying gene expression in a non-human cell comprising modifying a target nucleic acid sequence as in a method of numbered paragraph 30, wherein the nucleic acid is an mRNA and the functional moiety is a ribonuclease; optionally selected from an endonuclease, a 3' exonuclease or a 5' exonuclease.

36. A method of modifying a target nucleic acid as in any of numbered paragraphs 25 to 35, wherein the method is carried out at a temperature between 50°C and 100°C.

37. A method of modifying a target nucleic acid as in numbered paragraph 36, wherein the method is carried out at a temperature at or above 60°C, preferably between 60°C and 80°C, more preferably between 60°C and 65°C.

38. A method as in any of numbered paragraphs 25 to 37 wherein the cell is a prokaryotic cell.

39. A method as in any of numbered paragraphs 25 to 38 wherein the cell is a eukaryotic cell.

40. A host cell transformed by a method as in any of numbered paragraphs 22 to 36; wherein the cell is not a human cell.

CLAIMS

1. A nucleoprotein complex comprising a Cas protein, at least one targeting RNA molecule which recognises a target nucleic acid sequence in a double stranded target polynucleotide, and the target polynucleotide, wherein

the Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith;

the double stranded target polynucleotide comprises a target nucleic acid strand, comprising said target nucleic acid sequence, and a non-target nucleic acid strand, comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence and a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3'; wherein the nucleoprotein complex is not in a human cell.

2. A transformed non-human cell, having a double stranded target polynucleotide comprising a target nucleic acid sequence, wherein the double stranded target polynucleotide comprises a target nucleic acid strand, comprising said target nucleic acid sequence, and a non-target nucleic acid strand, comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence, said cell comprising:

a clustered regularly interspaced short palindromic repeat (CRISPR)-associated (Cas) protein having an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith;

at least one targeting RNA molecule which recognizes the target nucleic acid sequence in the target nucleic acid strand, wherein the non-target strand further comprises a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3'; and

an expression vector comprising a nucleic acid encoding at least one of said Cas protein and said targeting RNA molecule.

3. A transformed cell as claimed in claim 2, wherein the Cas protein is expressed from an expression vector.

4. A nucleoprotein complex as claimed in claim 1, or a transformed cell as claimed in claim 2 or claim 3, wherein the Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 95% identity therewith.

5. A nucleoprotein complex as claimed in claim 1 or claim 4, or transformed cell as claimed in any of claims 2 to 4, wherein the nucleoprotein complex occurs at a temperature between 20°C and 100°C, at a temperature between 30°C and 80°C, or at a temperature between 37°C and 78°C, preferably at a temperature above 55°C; more preferably at a temperature between 55°C and 80°C; even more preferably at a temperature between 55°C and 65°C or 60°C and 65°C.

6. A nucleoprotein complex or a transformed cell as claimed in any preceding claim, wherein the double stranded target polynucleotide comprising the target nucleic acid sequence is cleaved by the Cas protein; preferably wherein said cleavage is DNA cleavage.

7. A transformed cell as claimed in any of claims 2 to 5, wherein (i) binding, cleaving, marking or modifying of the double stranded target polynucleotide is at a temperature between 20°C and 100°C, at a temperature between 30°C and 80°C, or at a temperature between 37°C and 78°C; preferably at a temperature above 55°C; more preferably at a temperature between 55°C and 80°C; even more preferably at a temperature between 55°C and 65°C or 60°C and 65°C; and/or (ii) the target polynucleotide comprising the target sequence is double stranded DNA and said cleaving results in a double stranded break in the target polynucleotide.

8. A transformed cell or a nucleoprotein complex as claimed in any of claims to 5, wherein, the Cas protein lacks nuclease activity to cut the double stranded DNA yet there is gene silencing of the target polynucleotide.

9. A transformed cell or a nucleoprotein complex as claimed in any preceding claim, wherein the PAM sequence comprises 5'-NNNNCNNA-3' [SEQ ID NO: 47]; preferably wherein the PAM sequence comprises 5'-NNNNCSAA-3' [SEQ ID NO: 48]; more preferably wherein the PAM sequence comprises 5'-NNNNCCAA-3' [SEQ ID NO: 50].

10. A nucleoprotein complex as claimed in claim 1 or claim 4, or a transformed cell as claimed in any of claims 2 to 4, wherein the nucleoprotein complex forms at a temperature between 20°C and 70°C; preferably at a temperature between 25°C and 65°C.

11. A transformed cell as claimed in any of claims 3 to 5, wherein binding, cleaving, marking or modifying of the double stranded target polynucleotide occurs at a temperature between 20°C and 70°C; preferably at a temperature between 25°C and 65°C.

12. A transformed cell or a nucleoprotein complex as claimed in any preceding claim wherein the cell is a prokaryotic cell or wherein the nucleoprotein complex is in a prokaryotic cell.

13. A transformed cell or a nucleoprotein complex as claimed in any preceding claim, wherein the targeting RNA molecule comprises a crRNA and a tracrRNA.

14. A transformed cell or a nucleoprotein complex as claimed in any preceding claim, wherein the length of the at least one targeting RNA molecule is in the range 35 – 200 nucleotide residues; and/or wherein the target nucleic acid sequence is from 15 to 32 nucleotide residues in length.

15. A transformed cell or a nucleoprotein complex as claimed in any preceding claim, wherein the nuclease activity of the Cas protein is optionally inactive and the Cas protein further comprises at least one functional moiety; preferably a protein; optionally selected from a helicase, a nuclease, a helicase-nuclease, a DNA methylase, a histone methylase, an acetylase, a phosphatase, a kinase, a transcription (co-)activator, a transcription repressor, a DNA binding protein, a DNA structuring protein, a marker protein, a reporter protein, a fluorescent protein, a ligand binding protein, a signal peptide, a subcellular localisation sequence, an antibody epitope or an affinity purification tag, for example a green fluorescent protein (GFP).

16. An *in vitro* method of binding, cleaving, marking or modifying a double stranded target polynucleotide, wherein the double stranded target polynucleotide comprises a target nucleic acid strand comprising a target nucleic acid sequence,

and a non-target nucleic acid strand comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence, said method comprising:

a. designing at least one targeting RNA molecule, wherein the targeting RNA molecule recognizes the target sequence in the target strand, and the non-target strand further comprises a protospacer adjacent motif (PAM) sequence directly adjacent the 3' end of the protospacer sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3';

b. forming a ribonucleoprotein complex comprising the targeting RNA molecule and a Cas protein, wherein the isolated Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith; and

c. the ribonucleoprotein complex binding, cleaving, marking or modifying the target polynucleotide; and wherein said method is not used in human cells.

17. *In vitro* use of at least one targeting RNA molecule and a Cas protein to form a nucleoprotein complex for binding, cleaving, marking or modifying a double stranded target polynucleotide comprising a target nucleic acid sequence, wherein:

the double stranded target polynucleotide comprises a target nucleic acid strand, comprising said target nucleic acid sequence, and a non-target nucleic acid strand, comprising a protospacer nucleic acid sequence complementary to the target nucleic acid sequence;

the Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 89% identity therewith;

the at least one targeting RNA molecule recognizes the target sequence;

the non-target nucleic acid strand further comprises a protospacer adjacent motif (PAM) sequence directly adjacent to the 3' end of the protospacer nucleic acid sequence, wherein the PAM sequence comprises 5'-NNNNCNN-3'; and wherein said use is not in a human cell.

18. An *in vitro* method as claimed in claim 16 or an *in vitro* use as claimed in claim 17, wherein the Cas protein has an amino acid sequence of SEQ ID NO: 1 or a sequence of at least 95% identity therewith.

19. An *in vitro* method as claimed in claim 16 or claim 18, or an *in vitro* use as claimed in claim 17 or claim 18, wherein the nucleoprotein complex forms at a temperature between 20°C and 100°C, at a temperature between 30°C and 80°C, or at a temperature between 37°C and 78°C, preferably at a temperature above 55°C; more preferably at a temperature between 55°C and 80°C; even more preferably at a temperature between 55°C and 65°C or 60°C and 65°C.

20. An *in vitro* method as claimed in any of claims 16, 18 or 19, or an *in vitro* use as claimed in any of claims 17 to 19, wherein the PAM sequence comprises 5'-NNNNCNNA-3', 5'-CNNNCNN-3', 5'-NNNCCNN-3', 5'-NNCNCNN-3', 5'-NNNNCCN-3', and/or 5'-NCNNCNN-3'; preferably wherein the PAM sequence comprises 5'-CCCCCCNA-3' [SEQ ID NO: 10], more preferably wherein the PAM sequence comprises 5'-CCCCCCAA-3' [SEQ ID NO: 11].

21. An *in vitro* method as claimed in any of claims 16, 18, 19 or 20, or an *in vitro* use as claimed in any of claims 17 to 20, wherein the Cas protein further comprises at least one functional moiety; preferably a protein; optionally selected from a helicase, a nuclease, a helicase-nuclease, a DNA methylase, a histone methylase, an acetylase, a phosphatase, a kinase, a transcription (co-)activator, a transcription repressor, a DNA binding protein, a DNA structuring protein, a marker protein, a reporter protein, a fluorescent protein, a ligand binding protein, a signal peptide, a subcellular localisation sequence, an antibody epitope or an affinity purification tag, for example a green fluorescent protein (GFP).