
PYRAZOLYL QUINOXALINE KINASE INHIBITORS

FIELD OF THE INVENTION

The invention relates to new quinoxaline derivative compounds, to pharmaceutical compositions comprising said compounds, to processes for the preparation of said compounds and to the use of said compounds in the treatment of diseases, e.g. cancer.

SUMMARY

According to the disclosure there is provided compounds of formula (I):



(I)

including any tautomeric or stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2, 3 or 4;

R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, cyanoC_{1-4}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with **-**NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-haloC_{1-6}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-S(=O)$_2$-$NR^{14}R^{15}$, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –Si(CH_3)$_3$, C_{1-6}alkyl substituted with -P(=O)(OH)$_2$ or C_{1-6}alkyl substituted with -P(=O)(OC_{1-6}alkyl)$_2$;

each R^{1a} is independently selected from hydrogen, C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -NH(C_{3-8}cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl substituted with one or more fluoro atoms;

each R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{2-4}alkynyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkyl, haloC_{1-4}alkoxy, hydroxyhaloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, haloC_{1-4}alkoxyC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, hydroxyhaloC_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkyl substituted with R^{13}, C_{1-4}alkyl substituted with -C(=O)-R^{13}, C_{1-4}alkoxy substituted with R^{13}, C_{1-4}alkoxy substituted with -C(=O)-R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with -NR^7R^8, C_{1-4}alkyl substituted with –C(=O)-NR^7R^8, C_{1-4}alkoxy substituted with -NR^7R^8, C_{1-4}alkoxy substituted with –C(=O)-NR^7R^8, -NR^7R^8 and -C(=O)-NR^7R^8; or when two R^2 groups are attached to adjacent carbon atoms they may be taken together to form a radical of formula:

-O-(C(R^{17})$_2$)$_p$-O-;

-X-CH=CH-; or

-X-CH=N-; wherein R^{17} represents hydrogen or fluorine, p represents 1 or 2 and X represents O or S;

R^3 represents hydroxyl, C_{1-6}alkoxy, hydroxyC_{1-6}alkoxy, C_{1-6}alkoxy substituted with -$NR^{10}R^{11}$, C_{1-6}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, haloC_{1-6}alkyl optionally substituted with –O-C(=O)-C_{1-6}alkyl, hydroxyC_{1-6}alkyl optionally substituted with –O-C(=O)-C_{1-6}alkyl, hydroxyC_{2-6}alkenyl, hydroxyC_{2-6}alkynyl, hydroxyhaloC_{1-6}alkyl, cyanoC_{1-6}alkyl, C_{1-6}alkyl substituted with carboxyl, C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-O-C(=O)–, C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–, C_{1-6}alkyl substituted with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups or with –O-C(=O)-C_{1-6}alkyl, C_{2-6}alkenyl substituted with C_{1-6}alkoxy, C_{2-6}alkynyl substituted with C_{1-6}alkoxy, C_{1-6}alkyl substituted with R^9 and optionally substituted with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-R^9, C_{1-6}alkyl substituted with hydroxyl and R^9, C_{2-6}alkenyl substituted with R^9, C_{2-6}alkynyl substituted with R^9, C_{1-6}alkyl substituted with -$NR^{10}R^{11}$, C_{2-6}alkenyl substituted with -$NR^{10}R^{11}$, C_{2-6}alkynyl substituted with -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with one or two halogens and -$NR^{10}R^{11}$, -C_{1-6}alkyl-C(R^{12})=N-O-R^{12}, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-haloC_{1-6}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -S(=O)$_2$-haloC_{1-}

$_6$alkyl, C_{1-6}alkyl substituted with $-S(=O)_2-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-NR^{12}-S(=O)_2-C_{1-6}$alkyl, C_{1-6}alkyl substituted with $-NH-S(=O)_2-$haloC_{1-6}alkyl, C_{1-6}alkyl substituted with $-NR^{12}-S(=O)_2-NR^{14}R^{15}$, R^{13}, C_{1-6}alkyl substituted with $-P(=O)(OH)_2$ or C_{1-6}alkyl substituted with $-P(=O)(OC_{1-6}alkyl)_2$;

R^4 and R^5 each independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, $-S(=O)_2-C_{1-6}$alkyl, $-S(=O)_2-$haloC_{1-6}alkyl, $-S(=O)_2-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-S(=O)_2-C_{1-6}$alkyl, C_{1-6}alkyl substituted with $-S(=O)_2-$haloC_{1-6}alkyl, C_{1-6}alkyl substituted with $-S(=O)_2-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-NH-S(=O)_2-C_{1-6}$alkyl, C_{1-6}alkyl substituted with $-NH-S(=O)_2-$haloC_{1-6}alkyl, C_{1-6}alkyl substituted with $-NH-S(=O)_2-NR^{14}R^{15}$, R^{13} or C_{1-6}alkyl substituted with R^{13};

R^6 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C_{1-6}alkyl, cyanoC_{1-6}alkyl, hydroxyl, carboxyl, hydroxyC_{1-6}alkyl, halogen, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}alkoxyC_{1-6}alkyl, C_{1-6}alkyl-O-C(=O)-, $-NR^{14}R^{15}$, $-C(=O)-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-C(=O)-NR^{14}R^{15}$, $-S(=O)_2-C_{1-6}$alkyl, $-S(=O)_2-$haloC_{1-6}alkyl, $-S(=O)_2-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-S(=O)_2-C_{1-6}$alkyl, C_{1-6}alkyl substituted with $-S(=O)_2-$haloC_{1-6}alkyl, C_{1-6}alkyl substituted with $-S(=O)_2-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-NH-S(=O)_2-C_{1-6}$alkyl, C_{1-6}alkyl substituted with $-NH-S(=O)_2-$haloC_{1-6}alkyl or C_{1-6}alkyl substituted with $-NH-S(=O)_2-NR^{14}R^{15}$;

R^7 and R^8 each independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl or C_{1-6}alkoxyC_{1-6}alkyl;

R^9 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl each optionally and each independently being substituted with 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from =O, C_{1-4}alkyl, hydroxyl, carboxyl, hydroxyC_{1-4}alkyl, cyano, cyanoC_{1-4}alkyl, C_{1-4}alkyl-O-C(=O)-, C_{1-4}alkyl substituted with C_{1-}

$_4$alkyl-O-C(=O)-, C_{1-4}alkyl-C(=O)-, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, halogen, haloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkyl, -$NR^{14}R^{15}$, -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkoxy, –S(=O)$_2$-C_{1-4}alkyl, –S(=O)$_2$-haloC_{1-4}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –NH-S(=O)$_2$-C_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-S(=O)$_2$-haloC_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-S(=O)$_2$-$NR^{14}R^{15}$, R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with R^{13}, phenyl optionally substituted with R^{16}, phenylC_{1-6}alkyl wherein the phenyl is optionally substituted with R^{16}, a 5 or 6-membered aromatic monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S wherein said heterocyclyl is optionally substituted with R^{16};
or when two of the substituents of R^9 are attached to the same atom, they may be taken together to form a 4 to 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S;

R^{10} and R^{11} each independently represent hydrogen, carboxyl, C_{1-6}alkyl, cyanoC_{1-6}alkyl, C_{1-6}alkyl substituted with –$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –C(=O)-$NR^{14}R^{15}$, haloC_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, R^6, C_{1-6}alkyl substituted with R^6, -C(=O)-R^6, -C(=O)-C_{1-6}alkyl, –C(=O)-hydroxyC_{1-6}alkyl, -C(=O)-haloC_{1-6}alkyl,-C(=O)-hydroxyhaloC_{1-6}alkyl, C_{1-6}alkyl substituted with –Si$(CH_3)_3$, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-haloC_{1-6}alkyl, -S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-S(=O)$_2$-$NR^{14}R^{15}$;

R^{12} represents hydrogen or C_{1-4}alkyl optionally substituted with C_{1-4}alkoxy;


R^{13} represents C_{3-8}cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C_{3-8}cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C_{1-6}alkyl, -C(=O)-C_{1-6}alkyl, C_{1-6}alkoxy, or -$NR^{14}R^{15}$;

R^{14} and R^{15} each independently represent hydrogen, or haloC_{1-4}alkyl, or C_{1-4}alkyl optionally substituted with a substituent selected from hydroxyl, C_{1-4}alkoxy, amino or mono-or di(C_{1-4}alkyl)amino;

R^{16} represents hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{1-4}alkoxy, -$NR^{14}R^{15}$ or –C(=O)$NR^{14}R^{15}$;

the *N*-oxides thereof, the pharmaceutically acceptable salts thereof or the solvates thereof.

In one example there is provided compounds of formula (I^0):



(I)

including any stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2, 3 or 4;

R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with -NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-$NR^{14}R^{15}$, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –$Si(CH_3)_3$, C_{1-6}alkyl substituted with -P(=O)$(OH)_2$ or C_{1-6}alkyl substituted with -P(=O)$(OC_{1-6}alkyl)_2$;

each R^2 is independently selected from halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{2-4}alkynyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkyl, haloC_{1-4}alkoxy, hydroxyhaloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, haloC_{1-4}alkoxyC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, hydroxyhaloC_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkyl substituted with R^{13}, C_{1-4}alkyl substituted with -C(=O)-R^{13}, C_{1-4}alkoxy substituted with R^{13}, C_{1-4}alkoxy substituted with -C(=O)-R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with -NR^7R^8, C_{1-4}alkyl substituted with –C(=O)-NR^7R^8, C_{1-4}alkoxy substituted with -NR^7R^8, C_{1-4}alkoxy substituted with –C(=O)-NR^7R^8, -NR^7R^8 or -C(=O)-NR^7R^8;

R^3 represents hydroxyl, C_{1-6}alkoxy, hydroxyC_{1-6}alkoxy, C_{1-6}alkoxy substituted with -$NR^{10}R^{11}$, C_{1-6}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, haloC_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyC_{2-6}alkenyl, hydroxyC_{2-6}alkynyl, hydroxyhaloC_{1-6}alkyl, cyanoC_{1-6}alkyl, C_{1-6}alkyl substituted with carboxyl, C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-O-C(=O)–, C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–, C_{1-6}alkyl substituted with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{2-6}alkenyl substituted with C_{1-6}alkoxy, C_{2-6}alkynyl substituted with C_{1-6}alkoxy, C_{1-6}alkyl substituted with R^9, C_{1-6}alkyl substituted with –C(=O)-R^9, C_{1-6}alkyl substituted with hydroxyl and R^9, C_{2-6}alkenyl substituted with R^9, C_{2-6}alkynyl substituted with R^9, C_{1-6}alkyl substituted with -$NR^{10}R^{11}$, C_{2-6}alkenyl substituted with -$NR^{10}R^{11}$, C_{2-6}alkynyl substituted with -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with one or two halogens and -$NR^{10}R^{11}$, -C_{1-6}alkyl-$C(R^{12})$=N-O-R^{12}, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-$NR^{14}R^{15}$, R^{13}, C_{1-6}alkyl substituted with -P(=O)$(OH)_2$ or C_{1-6}alkyl substituted with -P(=O)$(OC_{1-6}alkyl)_2$;

R^4 and R^5 independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NH- $S(=O)_2$-$NR^{14}R^{15}$, R^{13} or C_{1-6}alkyl substituted with R^{13};

R^6 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C_{1-6}alkyl, cyanoC_{1-}

$_{6}$alkyl, hydroxyl, carboxyl, hydroxyC_{1-6}alkyl, halogen, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}alkoxyC_{1-6}alkyl, C_{1-6}alkyl-O-C(=O)-, -$NR^{14}R^{15}$, -C(=O)-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$;

R^7 and R^8 independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl or C_{1-6}alkoxyC_{1-6}alkyl;

R^9 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl each optionally and each independently being substituted with 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from =O, C_{1-4}alkyl, hydroxyl, carboxyl, hydroxyC_{1-4}alkyl, cyano, cyanoC_{1-4}alkyl, C_{1-4}alkyl-O-C(=O)-, C_{1-4}alkyl substituted with C_{1-4}alkyl-O-C(=O)-, C_{1-4}alkyl-C(=O)-, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, halogen, haloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkyl, -$NR^{14}R^{15}$, -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkoxy, –$S(=O)_2$-C_{1-4}alkyl, –$S(=O)_2$-haloC_{1-4}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-C_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$, R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with R^{13}, phenyl optionally substituted with R^{16}, phenylC_{1-6}alkyl wherein the phenyl is optionally substituted with R^{16}, a 5 or 6-membered aromatic monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S wherein said heterocyclyl is optionally substituted with R^{16};
or when two of the substituents of R^9 are attached to the same atom, they may be taken together to form a 4 to 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S;

R^{10} and R^{11} each independently represent hydrogen, C_{1-6}alkyl, cyanoC_{1-6}alkyl, C_{1-6}alkyl substituted with –$NR^{14}R^{15}$, haloC_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or

two hydroxyl groups, R^6, C_{1-6}alkyl substituted with R^6, -C(=O)-R^6, -C(=O)-C_{1-6}alkyl, –C(=O)-hydroxyC_{1-6}alkyl, -C(=O)-haloC_{1-6}alkyl,-C(=O)-hydroxyhaloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$Si(CH_3)_3$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, -$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$;

R^{12} represents hydrogen or C_{1-4}alkyl optionally substituted with C_{1-4}alkoxy;

R^{13} represents C_{3-8}cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C_{3-8}cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C_{1-6}alkyl, -C(=O)-C_{1-6}alkyl, C_{1-6}alkoxy, or -$NR^{14}R^{15}$;


R^{14} and R^{15} each independently represent hydrogen, or haloC_{1-4}alkyl, or C_{1-4}alkyl optionally substituted with a substituent selected from hydroxyl, C_{1-4}alkoxy, amino or mono-or di(C_{1-4}alkyl)amino;

R^{16} represents hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{1-4}alkoxy, -$NR^{14}R^{15}$ or –C(=O)$NR^{14}R^{15}$;

the *N*-oxides thereof, the pharmaceutically acceptable salts thereof or the solvates thereof.

WO2006/092430, WO2008/003702, WO01/68047, WO2005/007099, WO2004/098494, WO2009/141386, WO 2004/030635, WO 2008/141065, WO 2011/026579, WO 2011/028947 and WO 00/42026 which each disclose a series of heterocyclyl derivatives.

According to one aspect of the disclosure, there is provided a compound of formula (VI):



(VI)

including any tautomeric or stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2, 3 or 4;

R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, cyanoC_{1-4}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with -NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-haloC_{1-6}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-S(=O)$_2$-$NR^{14}R^{15}$, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –Si(CH_3)$_3$, C_{1-6}alkyl substituted with -P(=O)(OH)$_2$ or C_{1-6}alkyl substituted with -P(=O)(OC_{1-6}alkyl)$_2$;

each R^{1a} is independently selected from hydrogen, C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -NH(C_{3-8}cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl substituted with one or more fluoro atoms;

each R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{2-4}alkynyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkyl, haloC_{1-4}alkoxy, hydroxyhaloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, haloC_{1-4}alkoxyC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, hydroxyhaloC_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkyl substituted with R^{13}, C_{1-4}alkyl substituted with -C(=O)-R^{13}, C_{1-4}alkoxy substituted with R^{13}, C_{1-4}alkoxy substituted with -C(=O)-R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with -NR^7R^8, C_{1-4}alkyl substituted with –C(=O)-NR^7R^8, C_{1-4}alkoxy substituted with -NR^7R^8, C_{1-4}alkoxy substituted with –C(=O)-NR^7R^8, -NR^7R^8 and -C(=O)-NR^7R^8; or when two R^2 groups are attached to adjacent carbon atoms they may be taken together to form a radical of formula:

-O-(C(R^{17})$_2$)$_p$-O-;

-X-CH=CH-; or

-X-CH=N-; wherein R^{17} represents hydrogen or fluorine, p represents 1 or 2 and X represents O or S;

R^4 and R^5 each independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may

optionally be substituted with one or two hydroxyl groups, –S(=O)$_2$-C$_{1-6}$alkyl, –S(=O)$_2$-haloC$_{1-6}$alkyl, –S(=O)$_2$-NR^{14}R^{15}, C$_{1-6}$alkyl substituted with -S(=O)$_2$-C$_{1-6}$alkyl, C$_{1-6}$alkyl substituted with -S(=O)$_2$-haloC$_{1-6}$alkyl, C$_{1-6}$alkyl substituted with –S(=O)$_2$-NR^{14}R^{15}, C$_{1-6}$alkyl substituted with –NH-S(=O)$_2$-C$_{1-6}$alkyl, C$_{1-6}$alkyl substituted with –NH-S(=O)$_2$-haloC$_{1-6}$alkyl, C$_{1-6}$alkyl substituted with –NH- S(=O)$_2$-NR^{14}R^{15}, R^{13} or C$_{1-6}$alkyl substituted with R^{13};

R^6 represents C$_{3-8}$cycloalkyl, C$_{3-8}$cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; said C$_{3-8}$cycloalkyl, C$_{3-8}$cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C$_{1-6}$alkyl, cyanoC$_{1-6}$alkyl, hydroxyl, carboxyl, hydroxyC$_{1-6}$alkyl, halogen, haloC$_{1-6}$alkyl, hydroxyhaloC$_{1-6}$alkyl, C$_{1-6}$alkoxy, C$_{1-6}$alkoxyC$_{1-6}$alkyl, C$_{1-6}$alkyl-O-C(=O)-, -NR^{14}R^{15}, -C(=O)-NR^{14}R^{15}, C$_{1-6}$alkyl substituted with -NR^{14}R^{15}, C$_{1-6}$alkyl substituted with -C(=O)-NR^{14}R^{15}, –S(=O)$_2$-C$_{1-6}$alkyl, –S(=O)$_2$-haloC$_{1-6}$alkyl, –S(=O)$_2$-NR^{14}R^{15}, C$_{1-6}$alkyl substituted with –S(=O)$_2$-C$_{1-6}$alkyl, C$_{1-6}$alkyl substituted with –S(=O)$_2$-haloC$_{1-6}$alkyl, C$_{1-6}$alkyl substituted with –S(=O)$_2$-NR^{14}R^{15}, C$_{1-6}$alkyl substituted with –NH-S(=O)$_2$-C$_{1-6}$alkyl, C$_{1-6}$alkyl substituted with –NH-S(=O)$_2$-haloC$_{1-6}$alkyl or C$_{1-6}$alkyl substituted with –NH-S(=O)$_2$-NR^{14}R^{15};

R^7 and R^8 each independently represent hydrogen, C$_{1-6}$alkyl, hydroxyC$_{1-6}$alkyl, haloC$_{1-6}$alkyl, hydroxyhaloC$_{1-6}$alkyl or C$_{1-6}$alkoxyC$_{1-6}$alkyl;

R^{13} represents C$_{3-8}$cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C$_{3-8}$cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C$_{1-6}$alkyl, -C(=O)-C$_{1-6}$alkyl, C$_{1-6}$alkoxy, or -NR^{14}R^{15}; and

R^{14} and R^{15} each independently represent hydrogen, or haloC$_{1-4}$alkyl, or C$_{1-4}$alkyl optionally substituted with a substituent selected from hydroxyl, C$_{1-4}$alkoxy, amino or mono-or di(C$_{1-4}$alkyl)amino.

DETAILED DESCRIPTION OF THE INVENTION

Unless the context indicates otherwise, references to formula (I) in all sections of this document (including the uses, methods and other aspects of the invention) include references to all other sub-formula (e.g. I', I'', I''', I^0, $I^{0'}$, $I^{0''}$, $I^{0'''}$), sub-groups, preferences, embodiments and examples as defined herein.

The prefix "C_{x-y}" (where x and y are integers) as used herein refers to the number of carbon atoms in a given group. Thus, a C_{1-6}alkyl group contains from 1 to 6 carbon

atoms, a C_{3-6}cycloalkyl group contains from 3 to 6 carbon atoms, a C_{1-4}alkoxy group contains from 1 to 4 carbon atoms, and so on.

The term 'halo' or 'halogen' as used herein refers to a fluorine, chlorine, bromine or iodine atom.

The term 'C_{1-4}alkyl', or 'C_{1-6}alkyl' as used herein as a group or part of a group refers to a linear or branched saturated hydrocarbon group containing from 1 to 4 or 1 to 6 carbon atoms. Examples of such groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl or hexyl and the like.

The term 'C_{2-4}alkenyl' or 'C_{2-6}alkenyl' as used herein as a group or part of a group refers to a linear or branched hydrocarbon group containing from 2 to 4 or 2 to 6 carbon atoms and containing a carbon carbon double bond.

The term 'C_{2-4}alkynyl' or 'C_{2-6}alkynyl' as used herein as a group or part of a group refers to a linear or branched hydrocarbon group having from 2 to 4 or 2 to 6 carbon atoms and containing a carbon carbon triple bond.

The term 'C_{1-4}alkoxy' or 'C_{1-6}alkoxy' as used herein as a group or part of a group refers to an –O-C_{1-4}alkyl group or an –O-C_{1-6}alkyl group wherein C_{1-4}alkyl and C_{1-6}alkyl are as defined herein. Examples of such groups include methoxy, ethoxy, propoxy, butoxy, and the like.

The term 'C_{1-4}alkoxyC_{1-4}alkyl' or 'C_{1-6}alkoxyC_{1-6}alkyl' as used herein as a group or part of a group refers to a C_{1-4}alkyl–O-C_{1-4}alkyl group or a C_{1-6}alkyl–O-C_{1-6}alkyl group wherein C_{1-4}alkyland C_{1-6}alkyl are as defined herein. Examples of such groups include methoxyethyl, ethoxyethyl, propoxymethyl, butoxypropyl, and the like.

The term 'C_{3-8}cycloalkyl' as used herein refers to a saturated monocyclic hydrocarbon ring of 3 to 8 carbon atoms. Examples of such groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl and the like.

The term 'C_{3-8}cycloalkenyl' as used herein refers to a monocyclic hydrocarbon ring of 3 to 8 carbon atoms having a carbon carbon double bond.

The term 'hydroxyC_{1-4}alkyl' or 'hydroxyC_{1-6}alkyl' as used herein as a group or part of a group refers to a C_{1-4}alkyl or C_{1-6}alkyl group as defined herein wherein one or more than one hydrogen atom is replaced with a hydroxyl group. The terms 'hydroxyC_{1-4}alkyl' or 'hydroxyC_{1-6}alkyl' therefore include monohydroxyC_{1-4}alkyl, monohydroxyC_{1-6}alkyl and also polyhydroxyC_{1-4}alkyl and polyhydroxyC_{1-6}alkyl. There may be one, two, three or more hydrogen atoms replaced with a hydroxyl group, so the hydroxyC_{1-4}alkyl or hydroxyC_{1-6}alkyl may have one, two, three or more hydroxyl groups. Examples of such groups include hydroxymethyl, hydroxyethyl, hydroxypropyl and the like.

The term 'haloC_{1-4}alkyl' or 'haloC_{1-6}alkyl' as used herein as a group or part of a group refers to a C_{1-4}alkyl or C_{1-6}alkyl group as defined herein wherein one or more than one hydrogen atom is replaced with a halogen. The term 'haloC_{1-4}alkyl' or 'haloC_{1-6}alkyl' therefore include monohaloC_{1-4}alkyl, monohaloC_{1-6}alkyl and also polyhaloC_{1-4}alkyl and polyhaloC_{1-6}alkyl. There may be one, two, three or more hydrogen atoms replaced with a halogen, so the haloC_{1-4}alkyl or haloC_{1-6}alkyl may have one, two, three or more halogens. Examples of such groups include fluoroethyl, fluoromethyl, trifluoromethyl or trifluoroethyl and the like.

The term 'hydroxyhaloC_{1-4}alkyl' or 'hydroxyhaloC_{1-6}alkyl' as used herein as a group or part of a group refers to a C_{1-4}alkyl or C_{1-6}alkyl group as defined herein wherein one or more than one hydrogen atom is replaced with a hydroxyl group and one or more than one hydrogen atom is replaced with a halogen. The term 'hydroxyhaloC_{1-4}alkyl' or 'hydroxyhaloC_{1-6}alkyl' therefore refers to a C_{1-4}alkyl or C_{1-6}alkyl group wherein one, two, three or more hydrogen atoms are replaced with a hydroxyl group and one, two, three or more hydrogen atoms are replaced with a halogen.

The term 'hydroxyC_{1-4}alkoxy' or 'hydroxyC_{1-6}alkoxy' as used herein as a group or part of a group refers to an –O-C_{1-4}alkyl group or an –O-C_{1-6}alkyl group wherein the C_{1-4}alkyl and C_{1-6}alkyl group is as defined above and one or more than one hydrogen atom of the C_{1-4}alkyl or C_{1-6}alkyl group is replaced with a hydroxyl group. The term 'hydroxyC_{1-4}alkoxy' or 'hydroxyC_{1-6}alkoxy' therefore include monohydroxyC_{1-4}alkoxy,

monohydroxyC_{1-6}alkoxy and also polyhydroxyC_{1-4}alkoxy and polyhydroxyC_{1-6}alkoxy. There may be one, two, three or more hydrogen atoms replaced with a hydroxyl group so the hydroxyC_{1-4}alkoxy or hydroxyC_{1-6}alkoxy may have one, two, three or more hydroxyl groups. Examples of such groups include hydroxymethoxy, hydroxyethoxy, hydroxypropoxy and the like.

The term 'haloC_{1-4}alkoxy' or 'haloC_{1-6}alkoxy' as used herein as a group or part of a group refers to a –O-C_{1-4}alkyl group or a –O-C_{1-6} alkyl group as defined herein wherein one or more than one hydrogen atom is replaced with a halogen. The terms 'haloC_{1-4}alkoxy' or 'haloC_{1-6}alkoxy' therefore include monohaloC_{1-4}alkoxy, monohaloC_{1-6}alkoxy and also polyhaloC_{1-4}alkoxy and polyhaloC_{1-6}alkoxy. There may be one, two, three or more hydrogen atoms replaced with a halogen, so the haloC_{1-4}alkoxy or haloC_{1-6}alkoxy may have one, two, three or more halogens. Examples of such groups include fluoroethyloxy, difluoromethoxy or trifluoromethoxy and the like.

The term 'hydroxyhaloC_{1-4}alkoxy' as used herein as a group or part of a group refers to an –O-C_{1-4}alkyl group wherein the C_{1-4}alkyl group is as defined herein and wherein one or more than one hydrogen atom is replaced with a hydroxyl group and one or more than one hydrogen atom is replaced with a halogen. The term 'hydroxyhaloC_{1-4}alkoxy' therefore refers to a –O-C_{1-4}alkyl group wherein one, two, three or more hydrogen atoms are replaced with a hydroxyl group and one, two, three or more hydrogen atoms are replaced with a halogen.

The term 'haloC_{1-4}alkoxyC_{1-4}alkyl' as used herein as a group or part of a group refers to a C_{1-4}alkyl–O-C_{1-4}alkyl group wherein C_{1-4}alkyl is as defined herein and wherein in one or both of the C_{1-4}alkyl groups one or more than one hydrogen atom is replaced with a halogen. The term 'haloC_{1-4} alkoxyC_{1-4}alkyl' therefore refers to a C_{1-4}alkyl–O-C_{1-4}alkyl group wherein in one or both of the C_{1-4}alkyl groups one, two, three or more hydrogen atoms are replaced with a halogen and wherein C_{1-4} alkyl is as defined herein. Preferably, in one of the C_{1-4}alkyl groups one or more than one hydrogen atom is replaced with a halogen. Preferably, haloC_{1-4}alkoxyC_{1-4}alkyl means C_{1-4}alkyl substituted with haloC_{1-4}alkoxy.

The term 'hydroxyhaloC_{1-4}alkoxyC_{1-4}alkyl' as used herein refers to a C_{1-4}alkyl–O-C_{1-4}alkyl group wherein C_{1-4}alkyl is as defined herein and wherein in one or both of the C_{1-4}alkyl groups one or more than one hydrogen atom is replaced with a hydroxyl group and one or more than one hydrogen atom is replaced with a halogen. The terms 'hydroxyhaloC_{1-4}alkoxyC_{1-4}alkyl' therefore refers to a C_{1-4}alkyl–O-C_{1-4}alkyl group wherein in one or both of the C_{1-4}alkyl groups one, two, three or more hydrogen atoms are replaced with a hydroxyl group and one, two, three or more hydrogen atoms are replaced with a halogen and wherein C_{1-4}alkyl is as defined herein.

The term 'hydroxyC_{2-6}alkenyl' as used herein refers to a C_{2-6}alkenyl group wherein one or more than one hydrogen atom is replaced with a hydroxyl group and wherein C_{2-6}alkenyl is as defined herein.

The term 'hydroxyC_{2-6}alkynyl' as used herein refers to a C_{2-6}alkynyl group wherein one or more than one hydrogen atom is replaced with a hydroxyl group and wherein C_{2-6}alkynyl is as defined herein.

The term phenylC_{1-6}alkyl as used herein refers to a C_{1-6}alkyl group as defined herein which is substituted with one phenyl group.

The term cyanoC_{1-4}alkyl or cyanoC_{1-6}alkyl as used herein refers to a C_{1-4}alkyl or C_{1-6}alkyl group as defined herein which is substituted with one cyano group.

The term "heterocyclyl" as used herein shall, unless the context indicates otherwise, include both aromatic and non-aromatic ring systems. Thus, for example, the term "heterocyclyl group" includes within its scope aromatic, non-aromatic, unsaturated, partially saturated and fully saturated heterocyclyl ring systems. In general, unless the context indicates otherwise, such groups may be monocyclic or bicyclic and may contain, for example, 3 to 12 ring members, more usually 5 to 10 ring members. Reference to 4 to 7 ring members include 4, 5, 6 or 7 atoms in the ring and reference to 4 to 6 ring members include 4, 5, or 6 atoms in the ring. Examples of monocyclic groups are groups containing 3, 4, 5, 6, 7 and 8 ring members, more usually 3 to 7, and preferably 5, 6 or 7 ring members, more preferably 5 or 6 ring members. Examples of bicyclic groups are those containing 8, 9, 10, 11 and 12 ring members, and more usually

9 or 10 ring members. Where reference is made herein to heterocyclyl groups, the heterocyclyl ring can, unless the context indicates otherwise, be optionally substituted (i.e. unsubstituted or substituted) by one or more substituents as discussed herein.

The heterocyclyl groups can be heteroaryl groups having from 5 to 12 ring members, more usually from 5 to 10 ring members. The term "heteroaryl" is used herein to denote a heterocyclyl group having aromatic character. The term "heteroaryl" embraces polycyclic (e.g. bicyclic) ring systems wherein one or more rings are non-aromatic, provided that at least one ring is aromatic. In such polycyclic systems, the group may be attached by the aromatic ring, or by a non-aromatic ring.

Examples of heteroaryl groups are monocyclic and bicyclic groups containing from five to twelve ring members, and more usually from five to ten ring members. The heteroaryl group can be, for example, a five membered or six membered monocyclic ring or a bicyclic structure formed from fused five and six membered rings or two fused six membered rings, or two fused five membered rings. Each ring may contain up to about five heteroatoms typically selected from nitrogen, sulphur and oxygen. Typically the heteroaryl ring will contain up to 4 heteroatoms, more typically up to 3 heteroatoms, more usually up to 2, for example a single heteroatom. In one embodiment, the heteroaryl ring contains at least one ring nitrogen atom. The nitrogen atoms in the heteroaryl rings can be basic, as in the case of an imidazole or pyridine, or essentially non-basic as in the case of an indole or pyrrole nitrogen. In general the number of basic nitrogen atoms present in the heteroaryl group, including any amino group substituents of the ring, will be less than five.

Examples of five membered heteroaryl groups include but are not limited to pyrrole, furan, thiophene, imidazole, furazan, oxazole, oxadiazole, oxatriazole, isoxazole, thiazole, thiadiazole, isothiazole, pyrazole, triazole and tetrazole groups.

Examples of six membered heteroaryl groups include but are not limited to pyridine, pyrazine, pyridazine, pyrimidine and triazine.

A bicyclic heteroaryl group may be, for example, a group selected from:

a) a benzene ring fused to a 5- or 6-membered ring containing 1, 2 or 3 ring heteroatoms;
b) a pyridine ring fused to a 5- or 6-membered ring containing 0, 1, 2 or 3 ring heteroatoms;
c) a pyrimidine ring fused to a 5- or 6-membered ring containing 0, 1 or 2 ring heteroatoms;
d) a pyrrole ring fused to a 5- or 6-membered ring containing 0, 1, 2 or 3 ring heteroatoms;
e) a pyrazole ring fused to a 5- or 6-membered ring containing 0, 1 or 2 ring heteroatoms;
f) an imidazole ring fused to a 5- or 6-membered ring containing 0, 1 or 2 ring heteroatoms;
g) an oxazole ring fused to a 5- or 6-membered ring containing 0, 1 or 2 ring heteroatoms;
h) an isoxazole ring fused to a 5- or 6-membered ring containing 0, 1 or 2 ring heteroatoms;
i) a thiazole ring fused to a 5- or 6-membered ring containing 0, 1 or 2 ring heteroatoms;
j) an isothiazole ring fused to a 5- or 6-membered ring containing 0, 1 or 2 ring heteroatoms;
k) a thiophene ring fused to a 5- or 6-membered ring containing 0, 1, 2 or 3 ring heteroatoms;
l) a furan ring fused to a 5- or 6-membered ring containing 0, 1, 2 or 3 ring heteroatoms;
m) a cyclohexyl ring fused to a 5- or 6-membered ring containing 1, 2 or 3 ring heteroatoms; and
n) a cyclopentyl ring fused to a 5- or 6-membered ring containing 1, 2 or 3 ring heteroatoms.

Particular examples of bicyclic heteroaryl groups containing a five membered ring fused to another five membered ring include but are not limited to imidazothiazole (e.g. imidazo[2,1-b]thiazole) and imidazoimidazole (e.g. imidazo[1,2-a]imidazole).

Particular examples of bicyclic heteroaryl groups containing a six membered ring fused to a five membered ring include but are not limited to benzofuran, benzothiophene, benzimidazole, benzoxazole, isobenzoxazole, benzisoxazole, benzthiazole, benzisothiazole, isobenzofuran, indole, isoindole, indolizine, indoline, isoindoline, purine (e.g., adenine, guanine), indazole, pyrazolopyrimidine (e.g. pyrazolo[1,5-a]pyrimidine), triazolopyrimidine (e.g. [1,2,4]triazolo[1,5-a]pyrimidine), benzodioxole, imidazopyridine and pyrazolopyridine (e.g. pyrazolo[1,5-a]pyridine) groups.

Particular examples of bicyclic heteroaryl groups containing two fused six membered rings include but are not limited to quinoline, isoquinoline, chroman, thiochroman, chromene, isochromene, chroman, isochroman, benzodioxan, quinolizine, benzoxazine, benzodiazine, pyridopyridine, quinoxaline, quinazoline, cinnoline, phthalazine, naphthyridine and pteridine groups.

Examples of polycyclic heteroaryl groups containing an aromatic ring and a non-aromatic ring include, tetrahydroisoquinoline, tetrahydroquinoline, dihydrobenzthiene, dihydrobenzfuran, 2,3-dihydro-benzo[1,4]dioxine, benzo[1,3]dioxole, 4,5,6,7-tetrahydrobenzofuran, tetrahydrotriazolopyrazine (e.g. 5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazine), indoline and indane groups.

A nitrogen-containing heteroaryl ring must contain at least one ring nitrogen atom. Each ring may, in addition, contain up to about four other heteroatoms typically selected from nitrogen, sulphur and oxygen. Typically the heteroaryl ring will contain up to 3 heteroatoms, for example 1, 2 or 3, more usually up to 2 nitrogens, for example a single nitrogen. The nitrogen atoms in the heteroaryl rings can be basic, as in the case of an imidazole or pyridine, or essentially non-basic as in the case of an indole or pyrrole nitrogen. In general the number of basic nitrogen atoms present in the heteroaryl group, including any amino group substituents of the ring, will be less than five.

Examples of nitrogen-containing heteroaryl groups include, but are not limited to, pyridyl, pyrrolyl, imidazolyl, oxazolyl, oxadiazolyl, thiadiazolyl, oxatriazolyl, isoxazolyl, thiazolyl, isothiazolyl, furazanyl, pyrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, triazolyl (e.g., 1,2,3-triazolyl, 1,2,4-triazolyl), tetrazolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzoxazolyl, benzisoxazole, benzthiazolyl and benzisothiazole, indolyl, 3H-indolyl,

isoindolyl, indolizinyl, isoindolinyl, purinyl (e.g., adenine [6-aminopurine], guanine [2-amino-6-hydroxypurine]), indazolyl, quinolizinyl, benzoxazinyl, benzodiazinyl, pyridopyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, phthalazinyl, naphthyridinyl and pteridinyl.

Examples of nitrogen-containing polycyclic heteroaryl groups containing an aromatic ring and a non-aromatic ring include tetrahydroisoquinolinyl, tetrahydroquinolinyl, and indolinyl.

The term "non-aromatic group" embraces, unless the context indicates otherwise, unsaturated ring systems without aromatic character, partially saturated and fully saturated heterocyclyl ring systems. The terms "unsaturated" and "partially saturated" refer to rings wherein the ring structure(s) contains atoms sharing more than one valence bond i.e. the ring contains at least one multiple bond e.g. a C=C, C≡C or N=C bond. The term "fully saturated" refers to rings where there are no multiple bonds between ring atoms. Saturated heterocyclyl groups include piperidine, morpholine, thiomorpholine, piperazine. Partially saturated heterocyclyl groups include pyrazolines, for example 2-pyrazoline and 3-pyrazoline.

Examples of non-aromatic heterocyclyl groups are groups having from 3 to 12 ring members, more usually 5 to 10 ring members. Such groups can be monocyclic or bicyclic, for example, and typically have from 1 to 5 heteroatom ring members (more usually 1, 2, 3 or 4 heteroatom ring members), usually selected from nitrogen, oxygen and sulphur. The heterocyclyl groups can contain, for example, cyclic ether moieties (e.g. as in tetrahydrofuran and dioxane), cyclic thioether moieties (e.g. as in tetrahydrothiophene and dithiane), cyclic amine moieties (e.g. as in pyrrolidine), cyclic amide moieties (e.g. as in pyrrolidone), cyclic thioamides, cyclic thioesters, cyclic ureas (e.g. as in imidazolidin-2-one) cyclic ester moieties (e.g. as in butyrolactone), cyclic sulphones (e.g. as in sulpholane and sulpholene), cyclic sulphoxides, cyclic sulphonamides and combinations thereof (e.g. thiomorpholine).

Particular examples include morpholine, piperidine (e.g. 1-piperidinyl, 2-piperidinyl, 3-piperidinyl and 4-piperidinyl), piperidone, pyrrolidine (e.g. 1-pyrrolidinyl, 2-pyrrolidinyl and 3-pyrrolidinyl), pyrrolidone, azetidine, pyran (2H-pyran or 4H-pyran),

dihydrothiophene, dihydropyran, dihydrofuran, dihydrothiazole, tetrahydrofuran, tetrahydrothiophene, dioxane, tetrahydropyran (e.g. 4-tetrahydro pyranyl), imidazoline, imidazolidinone, oxazoline, thiazoline, 2-pyrazoline, pyrazolidine, piperazone, piperazine, and N-alkyl piperazines such as N-methyl piperazine. In general, preferred non-aromatic heterocyclyl groups include saturated groups such as piperidine, pyrrolidine, azetidine, morpholine, piperazine and N-alkyl piperazines.

In a nitrogen-containing non-aromatic heterocyclyl ring the ring must contain at least one ring nitrogen atom. The heterocylic groups can contain, for example cyclic amine moieties (e.g. as in pyrrolidine), cyclic amides (such as a pyrrolidinone, piperidone or caprolactam), cyclic sulphonamides (such as an isothiazolidine 1,1-dioxide, [1,2]thiazinane 1,1-dioxide or [1,2]thiazepane 1,1-dioxide) and combinations thereof. Particular examples of nitrogen-containing non-aromatic heterocyclyl groups include aziridine, morpholine, thiomorpholine, piperidine (e.g. 1-piperidinyl, 2-piperidinyl, 3-piperidinyl and 4-piperidinyl), pyrrolidine (e.g. 1-pyrrolidinyl, 2-pyrrolidinyl and 3-pyrrolidinyl), pyrrolidone, dihydrothiazole, imidazoline, imidazolidinone, oxazoline, thiazoline, 6H-1,2,5-thiadiazine, 2-pyrazoline, 3-pyrazoline, pyrazolidine, piperazine, and N-alkyl piperazines such as N-methyl piperazine.

The heterocyclyl groups can be polycyclic fused ring systems or bridged ring systems such as the oxa- and aza analogues of bicycloalkanes, tricycloalkanes (e.g. adamantane and oxa-adamantane). For an explanation of the distinction between fused and bridged ring systems, see *Advanced Organic Chemistry*, by Jerry March, 4th Edition, Wiley Interscience, pages 131-133, 1992.

The heterocyclyl groups can each be unsubstituted or substituted by one or more substituent groups. For example, heterocyclyl groups can be unsubstituted or substituted by 1, 2, 3 or 4 substituents. Where the heterocyclyl group is monocyclic or bicyclic, typically it is unsubstituted or has 1, 2 or 3 substituents.

The term ‘aryl’ as used herein refers to carbocyclyl aromatic groups including phenyl, naphthyl, indenyl, and tetrahydronaphthyl groups.

In one embodiment R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, cyanoC_{1-4}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with -NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-$NR^{14}R^{15}$, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –$Si(CH_3)_3$, C_{1-6}alkyl substituted with -$P(=O)(OH)_2$ or C_{1-6}alkyl substituted with -$P(=O)(OC_{1-6}alkyl)_2$.

In one embodiment R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with -NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, or C_{1-6}alkyl substituted with –$Si(CH_3)_3$.

In one embodiment R^1 represents hydrogen.

In one embodiment R^1 represents C_{1-6}alkyl. R^1 may represent -CH_3, -CD_3, -CH_2CH_3, -$CH_2CH_2CH_3$, -$CH_2CH(CH_3)_2$, -$CH(CH_3)_2$, -$CH_2CH(CH_3)_2$. In one embodiment R^1 represents -CH_3. In another embodiment R^1 represents -CD_3.

In one embodiment R^1 represents C_{2-4}alkenyl. R^1 may represent –CH_2-CH=CH_2.

In one embodiment R^1 represents hydroxyC_{1-6}alkyl. R^1 may represent -CH_2CH_2OH, -$CH_2C(CH_3)_2OH$ or $CH_2CHOHCH_2OH$.

In one embodiment R^1 represents haloC_{1-6}alkyl. R^1 may represent -CH_2CH_2F, $CH_2CH_2CH_2Cl$ or CH_2CH_2Br.

In one embodiment R^1 represents C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups. R^1 may represent -$CH_2CH_2OCH_3$.

In one embodiment R^1 represents C_{1-6}alkyl substituted with -NR^4R^5.

In one embodiment when R^1 represents C_{1-6}alkyl substituted with -NR^4R^5, R^4 and R^5 each represent hydrogen. R^1 may represent -$CH_2CH_2NH_2$ or -$CH_2CH_2CH_2NH_2$.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with -NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3. R^1 may represent -$CH_2CH_2NHCH_3$.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with -NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents –$S(=O)_2$-$NR^{14}R^{15}$ where R^{14} and R^{15} each represent C_{1-4}alkyl optionally substituted with hydroxyl, for example –CH_3. R^1 may represent -$CH_2CH_2NHS(=O)_2N(CH_3)_2$.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with -NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents –$S(=O)_2$-C_{1-6}alkyl. R^1 may represent -$CH_2CH_2NHS(=O)_2CH_3$.

In one embodiment R^1 represents C_{1-6}alkyl substituted with –C(=O)-NR^4R^5.

In one embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, R^4 and R^5 each represent C_{1-6}alkyl, for example –CH_3. R^1 may represent -$CH_2C(=O)N(CH_3)_2$.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3. R^1 may represent -$CH_2C(=O)NHCH_3$ or -$C(CH_3)_2C(=O)NHCH_3$.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents hydroxyC_{1-6}alkyl, for example –CH_2CH_2OH. R^1 may represent -$C(CH_3)_2C(=O)NHCH_2CH_2OH$ or -$CH_2C(=O)NHCH_2CH_2OH$.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, for example –$CH_2CH_2OCH_3$. R^1 may represent -$CH_2C(=O)NHCH_2CH_2OCH_3$ or -$C(CH_3)_2C(=O)NHCH_2CH_2OCH_3$.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents C_{1-6}alkyl substituted with R^{13}. R^{13} may represent a saturated 5 membered monocyclic heterocyclyl containing at least one nitrogen heteroatom, for example pyrrolidine. R^1 may represent –CH_2-C(=O)-NH-CH_2-CH_2-(pyrrolidin-1-yl).

In another embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, one of R^4 and R^5 represents hydrogen and the other represents C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl. R^1 may represent -$CH_2CH_2CH_2NHCH_2CH_2$-$S(=O)_2$-CH_3.

In one embodiment R^1 represents –$S(=O)_2$-C_{1-6}alkyl. R^1 may represent –$S(=O)_2$-CH_3.

In one embodiment R^1 represents –$S(=O)_2$-$NR^{14}R^{15}$. R^{14} and R^{15} may each represent C_{1-4}alkyl optionally substituted with hydroxyl, for example R^{14} and R^{15} may both represent –CH_3. R^1 may represent –$S(=O)_2$-$N(CH_3)_2$.

In one embodiment R^1 represents C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl. R^1 may represent -$CH_2CH_2S(=O)_2$-CH_3.


In one embodiment R^1 represents C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl. R^1 may represent -$CH_2CH_2NHS(=O)_2$-CH_3.

In one embodiment R^1 represents R^6. R^6 may represent a saturated 4, 5 or 6 membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, which may optionally be substituted.

In one embodiment when R^1 represents R^6, R^6 represents piperidinyl, for example 4-piperidinyl .

In one embodiment when R^1 represents R^6, R^6 represents tetrahydropyranyl, for example 2-tetrahydropyranyl or 4-tetrahydropyranyl.

In one embodiment when R^1 represents R^6, R^6 represents tetrahydrofuranyl, for example 3-tetrahydrofuranyl.

In another embodiment when R^1 represents R^6, R^6 represents azetidinyl substituted by one hydroxyC_{1-6}alkyl group. The hydroxyC_{1-6}alkyl group may be $-CH_2CH_2OH$. R^6 may represent 

In another embodiment when R^1 represents R^6, R^6 represents piperidinyl substituted by one C_{1-6}alkyl-O-C(=O)- group. The C_{1-6}alkyl-O-C(=O)- group may be $(CH_3)_3C$-O-C(=O)-. R^6 may represent 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3C$-O-C(=O)-. In another embodiment when R^1 represents R^6, R^6 represents piperidinyl substituted by one $-S(=O)_2$-C_{1-6}alkyl group. The $-S(=O)_2$-C_{1-6}alkyl group may be $-S(=O)_2CH_3$. R^6 may represent 4-piperidinyl substituted on the nitrogen atom with $-S(=O)_2CH_3$. In another embodiment when R^1 represents R^6, R^6 represents piperidinyl substituted by one C_{1-6}alkyl group. The C_{1-6}alkyl group may be $-CH_3$. R^6 may represent 4-piperidinyl substituted on the nitrogen atom with $-CH_3$.

In one embodiment R^1 represents C_{1-6}alkyl substituted with R^6. R^6 may represent a saturated 4, 5 or 6 membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, which may optionally be substituted. R^6 may represent pyrrolidinyl, thiophenyl, piperidinyl, morpholinyl, piperazinyl, tetrahydropyranyl. R^1 may represent methyl or ethyl each substituted with 4-piperidinyl, 4-piperazinyl, 1-pyrrolidinyl or 4-tetrahydropyranyl. R^1 may represent propyl substituted with morpholinyl where the morpholinyl is linked to the propyl through the N heteroatom. In another embodiment the heterocyclyl may be substituted by one substituent selected from halogen, C_{1-6}alkyl, hydroxyl, hydroxyC_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}alkyl-O-C(=O)-. The substituent may be –Cl, $-CH_3$, -OH, $-CH_2CH_2OH$, $-CH_2CH_2CH_2OH$, $-OCH_3$, $(CH_3)_3C$-O-C(=O)-.

R^1 may represent methyl, ethyl or propyl each substituted with 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3C$-O-C(=O)-, 4-piperidinyl substituted on the nitrogen atom with $-CH_3$, 4-piperazinyl substituted on the nitrogen atom (N1) with $(CH_3)_3C$-O-C(=O)-, 4-piperazinyl substituted on the nitrogen atom (N1) with $-CH_2CH_2OH$, 4-piperazinyl substituted on the nitrogen atom (N1) with $-CH_2CH_2CH_2OH$, 4-piperidinyl substituted in the 4 position by –OH, or 4-piperidinyl substituted in the 4 position by $-O-CH_3$. R^1 may represent methyl substituted with 2-thiophenyl substituted in the 5 position

with chlorine. In another embodiment the heterocyclyl may be substituted by two substituents selected from hydroxyl, C_{1-6}alkoxy, C_{1-6}alkyl-O-C(=O)-. The substituent may be -OH, -OCH_3, $(CH_3)_3$C-O-C(=O)-. R^1 may represent methyl substituted with 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)- and in the 4 position by –OH.

In one embodiment R^1 represents C_{1-6}alkyl substituted with –C(=O)-R^6. R^6 may represent a saturated 4, 5 or 6 membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, which may optionally be substituted. R^6 may represent piperazinyl or pyrrolidinyl.

In one embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-R^6, R^6 represents piperazinyl. R^1 may represent –$C(CH_3)_2$-C(=O)-(piperazin-4-yl).

In another embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-R^6, R^6 represents piperazinyl substituted by one C_{1-6}alkyl-O-C(=O)- group, for example $C(CH_3)_3$-O-C(=O)-. R^1 may represent –$C(CH_3)_2$-C(=O)-(piperazin-4-yl) substituted on the nitrogen atom in the 1 position by $C(CH_3)_3$-O-C(=O)-.

In another embodiment when R^1 represents C_{1-6}alkyl substituted with –C(=O)-R^6, R^6 represents pyrrolidinyl substituted by one hydroxyl group. R^1 may represent –CH_2-C(=O)-(pyrrolidin-1-yl) substituted in the 3 position by –OH.

In one embodiment R^1 represents hydroxyC_{1-6}alkyl substituted with R^6, R^6 may represent a saturated 4, 5 or 6 membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, which may optionally be substituted. R^6 may represent piperidinyl, for example 1-piperidinyl. R^1 may represent –$CH_2CHOHCH_2$-piperidin-1-yl.

In one embodiment R^1 represents C_{1-6}alkyl substituted with –$Si(CH_3)_3$. R^1 may represent -$CH_2Si(CH_3)_3$.

In one embodiment R^1 represents cyanoC_{1-4}alkyl. R^1 may represent -CH_2CH_2CN.

In one embodiment each R^{1a} is independently selected from hydrogen, C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -

NH(C_{3-8}cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl substituted with one or more fluoro atoms.

In one embodiment each R^{1a} is independently selected from hydrogen, C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with di(C_{1-4}alkyl)amino, and C_{1-4}alkyl substituted with one or more fluoro atoms.
In one embodiment one or two R^{1a} represents hydrogen. In one embodiment each R^{1a} represents hydrogen.

In one embodiment one or two R^{1a} represents C_{1-4}alkyl, for example $-CH_3$, $-CH_2CH_3$. In one embodiment each R^{1a} represents C_{1-4}alkyl, for example $-CH_3$.

In one embodiment one or two R^{1a} represents hydroxyC_{1-4}alkyl, for example $-CH_2OH$, $-CH_2CH_2OH$, $-CH_2CH_2CH_2OH$.

In one embodiment one or two R^{1a} represents C_{1-4}alkyl substituted with di(C_{1-4}alkyl)amino, for example $-CH_2N(CH_3)_2$. In one embodiment one or two R^{1a} represents C_{1-4}alkyl substituted with one or more fluoro atoms, for example $-CF_3$.

In one embodiment:
(i) one R^{1a} represents hydrogen and the other R^{1a} represents C_{1-4}alkyl, for example $-CH_3$, $-CH_2CH_3$;
(ii) one R^{1a} represents hydrogen and the other R^{1a} represents hydroxyC_{1-4}alkyl, for example $-CH_2OH$, $-CH_2CH_2OH$, $-CH_2CH_2CH_2OH$;
(iii) one R^{1a} represents hydrogen and the other R^{1a} represents C_{1-4}alkyl substituted with one or more fluoro atoms, for example $-CF_3$; or
(iv) each R^{1a} independently represents C_{1-4}alkyl, for example each R^{1a} represents $-CH_3$.

In one embodiment, R^1 is methyl and R^{1a} is hydrogen or methyl.

In one embodiment each R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkyl, haloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkoxy substituted with R^{13}, $-C(=O)-R^{13}$, C_{1-4}alkyl substituted with NR^7R^8, C_{1-4}alkoxy substituted with NR^7R^8, $-NR^7R^8$ and $-C(=O)-$

NR^7R^8; or when two R^2 groups are attached to adjacent carbon atoms they may be taken together to form a radical of formula -O-(C(R^{17})$_2$)$_p$-O- wherein R^{17} represents hydrogen or fluorine and p represents 1 or 2.

In one embodiment each R^2 is independently selected from halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkoxy substituted with R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with NR^7R^8, C_{1-4}alkoxy substituted with NR^7R^8, -NR^7R^8 or -C(=O)-NR^7R^8;

In one embodiment one or more R^2 represents hydroxyl.

In one embodiment one or more R^2 represents halogen, for example fluorine, chlorine or bromine.

In one embodiment one or more R^2 represents cyano.

In one embodiment one or more R^2 represents C_{1-4}alkyl, for example –CH_3.

In one embodiment one or more R^2 represents C_{2-4}alkenyl, for example -CH=CH_2.

In one embodiment one or more R^2 represents C_{1-4}alkoxy, for example CH_3O-, $(CH_3)_2CHO$-, CH_3CH_2O-, or CD_3O-.

In one embodiment one or more R^2 represents hydroxyC_{1-4}alkyl, for example -CH_2OH.

In one embodiment one or more R^2 represents hydroxyC_{1-4}alkoxy, for example –OCH_2CH_2OH.

In one embodiment one or more R^2 represents haloC_{1-4}alkyl, for example –CF_3.

In one embodiment one or more R^2 represents haloC_{1-4}alkoxy, for example -OCH_2CH_2F or –O-CHF_2-. In one embodiment one or more R^2 represents -OCH_2CH_2F or –O-CHF_2 or –OCF_3.

In one embodiment one or more R^2 represents C_{1-4}alkoxyC_{1-4}alkyl, for example -$CH_2CH_2OCH_3$.

In one embodiment one or more R^2 represents R^{13}. R^{13} may represent a saturated 5 membered monocyclic heterocyclyl containing two oxygen heteroatoms, for example dioxolanyl, particularly 2-dioxolanyl.

In one embodiment one or more R^2 represents C_{1-4}alkoxy substituted with R^{13}. R^{13} may represent C_{3-8}cycloalkyl, for example cyclopropyl. One or more R^2 may represent –$OCH_2C_3H_5$.

In one embodiment one or more R^2 represents -C(=O)-R^{13}. R^{13} may represent a saturated 5 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyrrolidinyl. R^2 may represent –C(=O)-(1-pyrrolidinyl).

In one embodiment one or more R^2 represents C_{1-4}alkyl substituted with -NR^7R^8. In one embodiment R^7 and R^8 each represent hydrogen. One or more R^2 may represent -CH_2NH_2. In another embodiment R^7 and R^8 may each independently represent C_{1-6}alkyl, for example –CH_2CH_3 or –CH_3. One or more R^2 may represent -$CH_2N(CH_2CH_3)_2$, -$CH_2N(CH_3)_2$ or -$CH_2N(CH_2CH_3)(CH_3)$.

In one embodiment one or more R^2 represents C_{1-4}alkoxy substituted with -NR^7R^8. In one embodiment one of R^7 and R^8 represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3. One or more R^2 may represent –$OCH_2CH_2NHCH_3$. In one embodiment R^7 and R^8 each represent hydrogen. One or more R^2 may represent –$OCH_2CH_2NH_2$.

In one embodiment one or more R^2 represents -NR^7R^8. In one embodiment one of R^7 and R^8 represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3. In one embodiment each of R^7 and R^8 represents C_{1-6}alkyl, for example –CH_3.

In one embodiment one or more R^2 represents -C(=O)-NR^7R^8. In one embodiment one of R^7 and R^8 represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3.

In one embodiment when two R^2 groups are attached to adjacent carbon atoms they may be taken together to form a radical of formula -O-(C(R^{17})$_2$)$_p$-O- wherein R^{17} represents hydrogen and p represents 1.

In one embodiment n is equal to 0. In one embodiment n is equal to 1. In one embodiment n is equal to 2. In one embodiment n is equal to 3. In one embodiment n is equal to 4.

In one embodiment n is equal to 1. R^2 may be at the 3-position. R^2 may represent
(i)haloC_{1-4}alkoxy, for example –O-CHF_2;
(ii)C_{1-4}alkoxy, for example CH_3O- or $(CH_3)_2CHO$- ;
(iii)cyano; ; or
(iv) -NR^7R^8, for example –$NHCH_3$.

In one embodiment n is equal to 1. R^2 may be at the 3-position. R^2 may represent haloC_{1-4}alkoxy, for example –OCF_3.
In one embodiment n is equal to 1. R^2 may be at the 3-position. R^2 may represent C_{1-4}alkoxy, for example CH_3O-. In one embodiment n is equal to 1. R^2 may be at the 3-position. R^2 may represent -NR^7R^8 where R^7 and R^8 each independently represent C_{1-6}alkyl, for example –$N(CH_3)_2$.
In one embodiment n is equal to 2. One R^2 may be at the 3-position and the other may be at the 5-position:
(i) each R^2 may represent C_{1-4}alkoxy, for example each R^2 may be CH_3O-, or the R^2 at the 3-position may be $(CH_3)_2CHO$- and the R^2 at the 5-position may be CH_3O-, or the R^2 at the 3-position may be CH_3O- and the R^2 at the 5-position may be CD_3O-;
(ii) the R^2 at the 3-position may represent halogen, for example fluorine, chlorine or bromine, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O-, CD_3O- or CH_3CH_2O- ;
(iii) the R^2 at the 3-position may represent C_{1-4}alkyl, for example –CH_3, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O- ;
(iv) the R^2 at the 3-position may represent cyano, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O-;

(v) the R^2 at the 3-position may represent C_{1-4}alkyl substituted with NR^7R^8, for example -CH_2NH_2 or –$CH_2N(CH_3)_2$ or -$CH_2N(CH_2CH_3)_2$ or -$CH_2N(CH_2CH_3)(CH_3)$, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O- ;

(vi) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, and the R^2 at the 5-position may represent -C(=O)-NR^7R^8, for example –C(=O)$NHCH_3$ or –C(=O)NH_2 ;

(vii) the R^2 at the 3-position may represent hydroxyC_{1-4}alkoxy, for example –OCH_2CH_2OH, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O- ;

(viii) the R^2 at the 3-position may represent -C(=O)-R^{13}, for example –C(=O)-(pyrrolidin-1-yl), and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O-;

(ix) the R^2 at the 3-position may represent C_{1-4}alkoxy substituted with R^{13}, for example –$OCH_2C_3H_5$, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O- ;

(x) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, and the R^2 at the 5-position may represent C_{1-4}alkoxy substituted with NR^7R^8, for example –$OCH_2CH_2NHCH_3$ or – $OCH_2CH_2NH_2$;

(xi) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, and the R^2 at the 5-position may represent C_{2-4}alkenyl, for example -$CH=CH_2$;

(xii) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, and the R^2 at the 5-position may represent C_{1-4}alkoxyC_{1-4}alkyl, for example -$CH_2CH_2OCH_3$;or the R^2 at the 3-position may be CH_3O- and the R^2 at the 5-position may be CH_3OCH_2-;

(xiii) the R^2 at the 3-position may represent R^{13}, for example 2-dioxolanyl, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O-;

(xiv) the R^2 at the 3-position may represent hydroxyC_{1-4}alkoxy, for example –OCH_2CH_2OH, and the R^2 at the 5-position may represent halogen, for example fluorine;

(xv) the R^2 at the 3-position may represent haloC_{1-4}alkoxy, for example -OCH_2CH_2F, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O- ;

(xvi) the R^2 at the 3-position may represent halogen, for example fluorine, and the R^2 at the 5-position may represent -C(=O)-NR^7R^8, for example –C(=O)$NHCH_3$;

(xvii) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, and the R^2 at the 5-position may represent halogen, for example fluorine ; or

(xviii) the R^2 at the 3-position may represent represents hydroxyC_{1-6}alkyl, for example -CH_2OH, and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O- .

In one embodiment n is equal to 2. One R^2 may be at the 3-position and the other may be at the 5-position:

(i) the R^2 at the 3-position may represent hydroxyl and the R^2 at the 5-position may represent C_{1-4}alkoxy, for example CH_3O-;

(ii) each R^2 may represent halogen, for example chlorine;

(iii) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O- and the R^2 at the 5-position may represent C_{1-4}alkyl substituted with -NR^7R^8 where R^7 and R^8 may each independently represent C_{1-6}alkyl, for example -$CH_2N(CH_2CH_3)_2$;

(iv) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, and the R^2 at the 5-position may represent haloC_{1-4}alkoxy, for example -$OCHF_2$;

(v) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, and the R^2 at the 5-position may represent haloC_{1-4}alkyl, for example -CHF_2; or

(vi) each R^2 may represent hydroxyl.

In one embodiment n is equal to 2. One R^2 may be at the 3-position and the other may be at the 5-position. Each R^2 may represent C_{1-4}alkoxy, for example each R^2 may be CH_3O-, $(CH_3)_2CHO$-, CH_3CH_2O-, CD_3O-. In one embodiment both R^2 are for example CH_3O-, or CD_3O-. In one embodiment both R^2 are CH_3O-.

In one embodiment n is equal to 2. One R^2 may be at the 4-position and the other may be at the 5-position. Each R^2 may represent C_{1-4}alkoxy, for example each R^2 may be CH_3O-.

In one embodiment n is equal to 2. One R^2 may be at the 5-position and the other may be at the 6-position. Each R^2 may represent C_{1-4}alkoxy, for example each R^2 may be CH_3O-.

In one embodiment n is equal to 2. One R^2 may be at the 2-position and the other may be at the 5-position:

(i) each R^2 may represent C_{1-4}alkoxy, for example each R^2 may be CH_3O-; or

(ii) the R^2 at the 2-position may be halogen, for example chlorine, and the R^2 at the 5 position may represent C_{1-4}alkoxy, for example CH_3O-.

In one embodiment n is equal to 3. One R^2 may be at the 2-position, one may be at the 3-position and one may be at the 5-position:
(i) the R^2 at the 2-position may represent halogen, for example chlorine, the R^2 at the 3-position and the 5-position may each represent C_{1-4}alkoxy, for example each of these R^2 may be CH_3O- ; or
(ii) the R^2 at the 2-position may represent C_{1-4}alkyl, for example $-CH_3$, the R^2 at the 3-position and the 5-position may each represent C_{1-4}alkoxy, for example each of these R^2 may be CH_3O- .

In one embodiment n is equal to 3. One R^2 may be at the 3-position, one may be at the 4-position and one may be at the 5-position:
(i) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, the R^2 at the 4-position and the 5-position may each represent halogen, for example fluorine; or ;
(ii) the R^2 at the 3-position may represent C_{1-4}alkoxy, for example CH_3O-, the R^2 at the 4-position and the 5-position may be taken together to form a radical of formula -O-$(C(R^{17})_2)_p$-O- wherein R^{17} represents hydrogen and p represents 1.

In one embodiment n is equal to 3. One R^2 may be at the 2-position, one may be at the 3-position and one may be at the 5-position: (i) the R^2 at the 2-position may represent halogen, for example fluorine, the R^2 at the 3-position and the 5-position may each represent C_{1-4}alkoxy, for example CH_3O-.

In one embodiment n is equal to 4. One R^2 may be at the 2-position, one may be at the 3-position, one may be at the 5-position and one may be at the 6-position, the R^2 at the 2-position and the 6-position may each represent halogen, for example chlorine or fluorine, the R^2 at the 3-position and the 5-position may each represent C_{1-4}alkoxy, for example CH_3O-.

R^3 may represent C_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, hydroxyC_{2-6}alkynyl, haloC_{1-6}alkyl, haloC_{1-6}alkyl optionally substituted (e.g. substituted) with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups or with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with R^9, C_{1-6}alkyl substituted with -$NR^{10}R^{11}$,

C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with one or two halogens and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with carboxyl, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with R^9 and optionally substituted with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with hydroxyl and R^9, -C_{1-6}alkyl-$C(R^{12})$=N-O-R^{12}, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-R^9, C_{2-6}alkenyl substituted with R^9, C_{2-6}alkynyl substituted with R^9, hydroxyC_{1-6}alkoxy, C_{2-6}alkenyl, C_{2-6}alkynyl, R^{13}, C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)– or C_{1-6}alkyl substituted with -P(=O)(OC_{1-6}alkyl)$_2$.

R^3 may represent C_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with R^9, C_{1-6}alkyl substituted with -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with one or two halogens and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with carboxyl, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with hydroxyl and R^9, -C_{1-6}alkyl-$C(R^{12})$=N-O-R^{12}, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-R^9, C_{2-6}alkynyl substituted with R^9, hydroxyC_{1-6}alkoxy, C_{2-6}alkenyl, C_{2-6}alkynyl, R^{13} or C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–.

In one embodiment R^3 represents C_{1-6}alkyl. R^3 may represent –CH_3, -CH_2CH_3, -$CH_2CH_2CH_3$ or –$CH_2CH(CH_3)_2$.

In one embodiment R^3 represents hydroxyC_{1-6}alkyl. R^3 may represent –CH_2CH_2OH, –$CH_2CH_2CH_2OH$, –$CH_2CHOHCH_3$, –$CH_2CHOHCH_2CH_3$, –$CH_2CHOHCH(CH_3)_2$, -$CH_2CH_2C(OH)(CH_3)_2$, -$CH_2CHOHCH_2OH$ or –$CH_2C(CH_3)_2OH$. R^3 may represent –CD_2CD_2OH or –$CD_2CD_2CD_2OH$. R^3 may represent –$CH(CH_3)CH_2OH$.

In one embodiment R^3 represents haloC_{1-6}alkyl. R^3 may represent $-CH_2CH_2CH_2Cl$ or $-CH_2CH_2CH_2CH_2Cl$. R^3 may represent $-CH_2CH_2F$ or $-CH_2CH_2I$.

In one embodiment R^3 represents haloC_{1-6}alkyl optionally substituted with $-O-C(=O)-C_{1-6}$alkyl. R^3 may represent $-CH_2CH(CF_3)-O-C(=O)CH_3$.

In one embodiment R^3 represents hydroxyhaloC_{1-6}alkyl, for example R^3 may represent $-CH_2CHOHCF_3$.

In one embodiment R^3 represents hydroxyC_{2-6}alkynyl, for example R^3 may represent $-CH_2-C\equiv C-CH_2OH$ or $-CH_2-C\equiv C-C(CH_3)_2OH$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with $-C(=O)-C_{1-6}$alkyl, for example R^3 may represent $CH_3-C(=O)-CH_2-$, $(CH_3)_2CH-C(=O)-CH_2-$

In one embodiment R^3 represents C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups. R^3 may represent $-CH_2CH_2OCH_3$, $-CH_2CH_2OCH_2CH_3$ or $-CH_2CHOHCH_2OCH_3$.

In one embodiment R^3 represents C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups or with $-O-C(=O)-C_{1-6}$alkyl. R^3 may represent $-CH_2CH(-O-C(=O)CH_3)CH_2OCH_3$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with R^9.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents optionally substituted C_{3-8}cycloalkyl, for example cyclopropyl or cyclopentyl. R^3 may represent $-CH_2-C_3H_5$ or $-CH_2C_5H_9$.

In one embodiment where the C_{3-8}cycloalkyl is cyclopropyl it is substituted by one hydroxyC_{1-4}alkyl, for example $-CH_2OH$.

In one embodiment where the C_{3-8}cycloalkyl is cyclopropyl it is substituted by one 6-membered aromatic monocyclic heterocyclyl containing one nitrogen heteroatom, for example 4-pyridinyl.

In another embodiment where the C_{3-8}cycloalkyl is cyclopropyl it is substituted by one C_{1-6}alkyl-O-C(=O)-, for example CH_3CH_2-O-C(=O)-.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing a nitrogen and an oxygen heteroatom, for example isoxazolyl. In one embodiment the heterocyclyl is substituted with one or two C_{1-4}alkyl groups, for example $-CH_3$ groups. R^3 may represent methyl substituted with 5-isoxazoyl substituted in the 3 position with $-CH_3$ or methyl substituted with 3-isoxazoyl substituted in the 5 position with $-CH_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted saturated 6 membered monocyclic heterocyclyl containing a nitrogen and an oxygen heteroatom, for example morpholinyl. R^3 may represent ethyl or propyl substituted by 4-morpholinyl. R^3 may represent methyl substituted by 3-morpholinyl. R^3 may represent methyl substituted by 6-morpholinyl.
In one embodiment the heterocyclyl is substituted with one or two C_{1-4}alkyl groups, for example $-CH_3$ groups. R^3 may represent ethyl or propyl substituted by 4-morpholinyl substituted in the 2 and 6 positions by $-CH_3$. R^3 may represent methyl substituted by 3-morpholinyl substituted in the 5 position by two $-CH_3$. R^3 may represent methyl substituted by 6-morpholinyl substituted in the 4 position by $-CH(CH_3)_2$.In one embodiment the heterocyclyl is substituted with one C_{1-4}alkyl group, for example $-CH(CH_3)_2$, and one =O. R^3 may represent methyl substituted by 6-morpholinyl substituted in the 3 position by =O and 4 position by $-CH(CH_3)_2$.
In another embodiment the heterocyclyl is substituted with phenylC_{1-6}alkyl, wherein the phenyl is optionally substituted with R^{16}, for example $-CH_2-C_6H_5$. R^3 may represent methyl substituted by 2-morpholinyl substituted in the 4 position by $-CH_2-C_6H_5$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated or an aromatic 3, 4, 5 or 6 membered monocyclic heterocyclyl containing one or two oxygen heteroatoms, for example ethylene oxide (oxiranyl), trimethylene oxide (oxetanyl), tetrahydrofuranyl, dioxolanyl, tetrahydropyranyl or furanyl. R^3 may be methyl substituted with 2-tetrahydrofuranyl, 2-dioxolane, ethylene oxide, 2-furanyl or 4-tetrahydropyranyl,

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted 4 membered heterocyclyl containing one oxygen heteroatom, for example oxetanyl, and the heterocyclyl is substituted with one C_{1-4}alkyl group, for example $-CH_3$. R^3 may be methyl substituted with 3-oxetanyl substituted in the 3 position by $-CH_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted 4 membered heterocyclyl containing one oxygen heteroatom, for example oxetanyl, and the heterocyclyl is substituted with one C_{1-4}alkyl substituted with $-NR^{14}R^{15}$ group where one of R^{14} and R^{15} is hydrogen and the other is C_{1-4}alkyl, for example $-CH(CH_3)_2$. R^3 may be methyl substituted with 3-oxetanyl substituted in the 3 position by $-CH_2NHCH(CH_3)_2$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 6 membered monocyclic heterocycle containing one or two nitrogen heteroatoms, for example pyridinyl or pyrazinyl. R^3 may represent methyl substituted with 3-pyridinyl or 2-pyrazinyl. R^3 may represent propyl substituted with 4-pyridinyl.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 6 membered monocyclic heterocycle containing two nitrogen heteroatoms, for example pyrimidinyl. R^3 may represent methyl or propyl substituted with 2-pyrimidinyl .

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 6 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyridinyl, substituted with one halogen, for example chlorine or bromine. R^3 may represent methyl substituted with 3-pyridinyl substituted in the 6 position by chlorine or 2-pyridinyl substituted in the 6 position by bromine.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 6 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyridinyl, substituted with:

(i) one C_{1-4}alkyl, for example $-CH_3$. R^3 may represent propyl substituted with 6-pyridinyl substituted in the 4 position by $-CH_3$; or

(ii) one C_{1-4}alkoxy, for example $-OCH_3$. R^3 may represent propyl substituted with 2-pyridinyl substituted in the 3 position by $-OCH_3$. R^3 may represent methyl substituted with 2-pyridinyl substituted in the 6 position by $-OCH_3$;

(iii) one C_{1-4}alkyl substituted by $-NR^{14}R^{15}$. In one embodiment R^{14} and R^{15} each represent hydrogen. R^3 may represent methyl substituted with 6-pyridinyl substituted in the 2 position by $-CH_2NH_2$; or

(iv) one $-NR^{14}R^{15}$. In one embodiment one of R^{14} and R^{15} represents hydrogen and the other represents C_{1-4}alkyl, for example $-CH_3$. R^3 may represent methyl substituted with 6-pyridinyl substituted in the 2 position by $-NHCH_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 6 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example pyrimidinyl, substituted with:

(i) one or two C_{1-4}alkoxy groups, for example $-OCH_3$. R^3 may represent propyl substituted with 2-pyrimidinyl substituted in the 4 position by $-OCH_3$. R^3 may represent methyl substituted with 2-pyrimidinyl substituted in the 4 and 6 positions by $-OCH_3$;

(ii) one hydroxyl group, for example –OH. R^3 may represent propyl substituted with 2-pyrimidinyl substituted in the 4 position by –OH.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted saturated 6 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example piperazinyl. R^3 may represent methyl substituted with 3-piperazinyl.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted saturated 6 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example piperazinyl substituted with R^{13}, for example said R^{13} representing piperidinyl being substituted with one C_{1-4}alkyl-C(=O)-, for example $-C(=O)-CH_3$. R^3 may represent ethyl substituted with 1-piperazinyl substituted in the 4 position with 4-piperidinyl substituted in the 1 position with $-C(=O)-CH_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted saturated 6 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example piperazinyl substituted with C_{1-4}alkyl substituted with $-C(=O)-NR^{14}R^{15}$. R^3 may represent ethyl substituted with 1-piperazinyl substituted in the 4 position with $-CH_2C(=O)NHCH(CH_3)_2$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a partially saturated 6 membered monocyclic heterocyclyl containing one nitrogen heteroatom which may optionally be substituted. R^3 may represent ethyl or propyl substituted with 1,2,3,6-tetrahydropyridine.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted saturated 4 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example azetidinyl.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 4 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example azetidinyl, and the heterocyclyl is substituted with one or two halogens, for example fluorine. R^3 may represent propyl substituted by 1-azetidinyl substituted in the 3 position by two fluorines.In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 4 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example azetidinyl, and the heterocyclyl is substituted with one hydroxyl group. R^3 may represent propyl substituted by 1-azetidinyl substituted in the 3 position by one -OH.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 5 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyrrolidinyl. R^3 may represent ethyl or propyl substituted with 1-pyrrolidinyl or 2-pyrrolidinyl.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 5 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyrrolidinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with:

a) one or two halogens, for example fluorine. R^3 may represent propyl substituted with 1-pyrrolidinyl substituted in the 3 position by two fluorines or with 1-pyrrolidinyl substituted in the 3 position by one fluorine;

b) one haloC_{1-4}alkyl, for example $-CH_2Cl$. R^3 may represent propyl substituted with 1-pyrrolidinyl substituted in the 2 position by $-CH_2Cl$;

c) one hydroxyl group. R^3 may represent ethyl or propyl substituted with 1-pyrrolidinyl substituted in the 3 position by –OH;

d) one =O group. R^3 may represent ethyl or propyl substituted with 1-pyrrolidinyl substituted in the 2 position by =O;

e) one $-S(=O)_2-C_{1-4}$alkyl group and the C_{1-4}alkyl may be $-CH_3$. R^3 may represent propyl substituted with 1-pyrrolidinyl substituted in the 3 position by $-S(=O)_2-CH_3$;

f) one $-NR^{14}R^{15}$ group. In one embodiment R^{14} and R^{15} each represent hydrogen. R^3 may represent ethyl or propyl substituted with 1-pyrrolidinyl substituted in the 3 position with $-NH_2$. In another embodiment R^{14} and R^{15} each independently represent C_{1-4}alkyl optionally substituted with hydroxyl, for example $-CH_3$. R^3 may represent ethyl substituted with 1-pyrrolidinyl substituted in the 3 position with $-N(CH_3)_2$. In another embodiment one of R^{14} and R^{15} is hydrogen and the other is C_{1-4}alkyl optionally substituted with hydroxyl, for example $-CH_3$. R^3 may represent propyl substituted with 1-pyrrolidinyl substituted in the 3 position with $-NHCH_3$;

g) one or two C_{1-4}alkyl groups, for example $-CH_3$ or $-CH(CH_3)_2$. R^3 may represent ethyl or propyl substituted with 1-pyrrolidinyl substituted in the 2 position with $-CH_3$, 1-pyrrolidinyl substituted in the 2 and the 5 position with $-CH_3$ or 1-pyrrolidinyl substituted in the 2 position with two $-CH_3$;

h) one carboxyl group. R^3 may represent ethyl substituted with 1-pyrrolidinyl substituted in the 2 position with $-C(=O)OH$;

i) one hydroxyC_{1-4}alkyl, for example $-CH_2OH$, $-C(CH_3)_2OH$ or $-CH_2CH_2OH$. R^3 may represent ethyl or propyl substituted with 1-pyrrolidinyl substituted in the 2 position with $-CH_2OH$;

j) R^{13}. In one embodiment R^{13} represents a saturated 6-membered monocyclic heterocyclyl containing one nitrogen heteroatom. In another embodiment R^{13} represents a saturated 6-membered monocyclic heterocyclyl containing one nitrogen and one oxygen heteroatom. In a further embodiment R^{13} represents a saturated 6-membered monocyclic heterocyclyl containing one nitrogen and one oxygen heteroatom, and the heterocyclyl is substituted, for example substituted with two C_{1-6}alkyl groups, for example two $-CH_3$ groups. R^3 may represent propyl substituted with 1-pyrrolidinyl substituted in the 3 position by 1-piperidinyl, or propyl substituted with 1-pyrrolidinyl substituted in the 3 position by 4-morpholinyl substituted in positions 2 and 6 by $-CH_3$;

k) one cyano group. R^3 may represent ethyl or propyl substituted with 1-pyrrolidinyl substituted in the 3 position by $-CN$;

l) one cyanoC_{1-4}alkyl, for example $-CH_2CN$. R^3 may represent propyl substituted with 1-pyrrolidinyl substituted in the 2 position by $-CH_2CN$.R^3 may represent ethyl substituted with 1-pyrrolidinyl substituted in the 2 position by $-CH_2CN$;

m) one C_{1-4}alkyl substituted with $-NH-S(=O)_2$-haloC_{1-4}alkyl, for example $-CH_2NH-S(=O)_2-CF_3$. R^3 may represent propyl substituted with 1-pyrrolidinyl substituted in the 2 position by $-CH_2NH-S(=O)_2-CF_3$; or

n) one C_{1-6}alkyl-O-C(=O)-, for example $(CH_3)_3C-O-C(=O)-$ or $CH_3-O-C(=O)-$. R^3 may represent methyl or ethyl substituted by 2-pyrrolidinyl substituted in the 1 position by $(CH_3)_3C-O-C(=O)-$ or substituted by 1-pyrrolidinyl substituted in the 2 position by $CH_3-O-C(=O)-$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 5 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyrrolidinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with a 6-membered aromatic monocyclic heterocyclyl containing one or two nitrogen heteroatoms, for example pyridinyl or pyrimidinyl, and optionally substituted with R^{16}. In one embodiment R^{16} represents C_{1-4}alkoxy, for example $-OCH_3$. R^3 may represent methyl substituted by 3-pyrrolidinyl substituted in the 1-position by 2-pyridinyl substituted in the 3-position by $-OCH_3$. R^3 may represent methyl substituted by 3-pyrrolidinyl substituted in the 1-position by 2-pyrimidinyl substituted in the 4-position by $-OCH_3$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example piperidinyl. R^3 may represent methyl, ethyl or propyl substituted by 4-piperidinyl or 1-piperidinyl.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example piperidinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with :

a) one or two halogens, for example fluorine. R^3 may represent ethyl substituted by 1-piperidinyl substituted at the 4 position by two fluorines;

b) one hydroxyl group. R^3 may represent methyl or ethyl substituted by 1-piperidinyl substituted at the 4 position by one –OH or 4-piperidinyl substituted at the 4 position by one –OH;

c) one $-NR^{14}R^{15}$ group. In one embodiment R^{14} and R^{15} each represent hydrogen. R^3 may represent ethyl substituted by 1-piperidinyl substituted at the 3 position or the 4 position by $-NH_2$. In another embodiment R^{14} and R^{15} each independently represent C_{1-4}alkyl optionally substituted with hydroxyl, for example $-CH_3$. R^3 may represent ethyl substituted by 1-piperidinyl substituted at the 4 position by $-N(CH_3)_2$;

d) one or two C_{1-4}alkyl groups, for example $-CH_3$ or $-CH(CH_3)_2$. R^3 may represent methyl, ethyl or propyl substituted by 1-piperidinyl substituted at the 2 position by $-CH_3$, 1-piperidinyl substituted at the 2 and the 6 position by $-CH_3$, 4-piperidinyl substituted at the 1 position by $-CH(CH_3)_2$, 4-piperidinyl substituted at the 1 position by $-CH_3$, 1-piperidinyl substituted at the 3 and the 5 position by $-CH_3$;

e) one hydroxyC_{1-4}alkyl, for example $-CH_2OH$, $-C(CH_3)_2OH$ or $-CH_2CH_2OH$. R^3 may represent ethyl substituted by 1-piperidinyl substituted in the 4 position by $-C(CH_3)_2OH$, 1-piperidinyl substituted in the 4 position by $-CH_2CH_2OH$; 1-piperidinyl substituted in the 4 position by $-CH_2OH$;

f) one cyano group. R^3 may represent ethyl or propyl substituted with 1-piperidinyl substituted at the 3 position with $-CN$;

g) one C_{1-6}alkyl-O-C(=O)-, for example CH_3CH_2-O-C(=O)-, $(CH_3)_3$C-O-C(=O)- or CH_3-O-C(=O)-. R^3 may represent methyl or ethyl substituted with 1-piperidinyl substituted in the 4 position by CH_3CH_2-O-C(=O)-, 4-piperidinyl substituted in the 1 position by $(CH_3)_3$C-O-C(=O)-;

h) one C_{1-6}alkyl-O-C(=O)-, for example $(CH_3)_3$C-O-C(=O)-, and one hydroxyl group. R^3 may represent methyl substituted with 4-piperidinyl substituted in the 4 position by $-OH$ and in the 1 position by $(CH_3)_3$C-O-C(=O)-;

i) one C_{1-6}alkyl-O-C(=O)-, for example $(CH_3)_3$C-O-C(=O)-, and one C_{1-4}alkoxy group, for example $-OCH_3$. R^3 may represent methyl substituted with 4-piperidinyl substituted in the 4 position by $-OCH_3$ and in the 1 position by $(CH_3)_3$C-O-C(=O)- ;

j) one C_{1-4}alkoxy group, for example $-OCH_3$. R^3 may represent methyl or ethyl substituted with 1-piperidinyl substituted in the 4 position by $-OCH_3$ or 4-piperidinyl substituted in the 4 position by $-OCH_3$;

k) one haloC_{1-4}alkyl group, for example $-CF_3$. R^3 may represent propyl substituted with 1-piperidinyl substituted in the 4 position by $-CF_3$; or

l) one -C(=O)-$NR^{14}R^{15}$ where R^{14} and R^{15} both represent hydrogen. R^3 may represent ethyl substituted with 1-piperidinyl substituted in the 3 position by -C(=O)-NH_2. R^3 may

represent ethyl or propyl substituted with 1-piperidinyl substituted in the 2 position by -C(=O)-NH_2.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example piperidinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with :

a) one =O. R^3 may represent ethyl substituted by 1-piperidinyl substituted at the 4 position by =O, or propyl substituted by 1-piperidinyl substituted at the 2 position by =O;

b) one C_{1-6}alkyl substituted with -$NR^{14}R^{15}$ where R^{14} and R^{15} both represent hydrogen. R^3 may represent ethyl substituted with 1-piperidinyl substituted in the 4 position by –CH_2NH_2.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example piperidinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with a 6-membered aromatic monocyclic heterocyclyl containing two nitrogen heteroatoms, for example pyrimidinyl, and optionally substituted with R^{16}. In one embodiment R^{16} represents C_{1-4}alkoxy, for example –OCH_3. R^3 may represent methyl substituted by 4-piperidinyl substituted in the 1-position by 2-pyrimidinyl substituted in the 4-position by –OCH_3.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a bicyclic heterocyclyl containing a benzene ring fused to a 5- or 6-membered ring containing 1, 2 or 3 ring heteroatoms. In one embodiment the bicyclic heterocyclyl contains a benzene ring fused to a 5-membered ring containing 1 ring heteroatom. In one embodiment the ring heteroatom is a nitrogen heteroatom. In one embodiment the bicyclic heterocyclyl is substituted with two =O groups on the 5-membered ring containing one ring heteroatom. R^3 may represent ethyl, propyl or butyl substituted with isoindolyl-1,3,-dione (e.g. isoindol-2-yl-1,3-dione, also known as phtalimidyl).R^3 may represent –$CH(CH_3)CH_2$- substituted with isoindolyl-1,3,-dione.

In one embodiment when R^3 represents C_{1-6}alkyl (for example ethyl or propyl) substituted with R^9, R^9 represents an optionally substituted monocyclic heterocycl

containing at least one heteroatom selected from N, O or S. In one embodiment R^9 represents a 4, 5 or 6 membered monocyclic saturated heterocycle substituted with two substituents which are attached to the same atom and which are taken together to form a 4 to 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; and R^3 represents C_{1-6}alkyl (for example ethyl or propyl) substituted with a 4, 5 or 6 membered monocyclic saturated heterocycle substituted with two substituents which are attached to the same atom and which are taken together to form a 4 to 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S. For example R^3 may represent ethyl substituted with 2-oxa-6-aza-spiro[3.3]heptane or R^3 may represent ethyl substituted with 1-piperidinyl substituted on the 4 position by 1,4-dioxolane e.g. to form 1, 4-dioxa-8-aza-spiro[4.5]decane.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing one sulphur heteroatom, for example thiophene. R^3 may represent methyl substituted with 2-thiophenyl. In one embodiment the aromatic 5 membered monocyclic heterocyclyl containing one sulphur heteroatom is substituted with one chlorine. R^3 may represent methyl substituted with 2-thiophenyl substituted at the 5 position by chlorine.
In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing one sulphur and one nitrogen heteroatom, for example thiazole. The 5-membered heterocyclyl may be substituted with for example one C_{1-4}alkyl, for example $-CH_3$. R^3 may represent methyl substituted with 4-thiazolyl substituted in the 2 position by $-CH_3$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example piperazinyl. R^3 may represent ethyl or propyl substituted with 1-piperazinyl. In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example piperazinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with :

a) one C_{1-4}alkyl-C(=O)-, for example CH_3-C(=O)-. R^3 may represent ethyl substituted with 1-piperazinyl substituted in the 4 position by CH_3-C(=O)-;

b) one hydroxyC_{1-4}alkyl, for example $-CH_2CH_2OH$. R^3 may represent ethyl substituted with 1-piperazinyl substituted in the 4 position by $-CH_2CH_2OH$;
c) one or two C_{1-4}alkyl, for example $-CH_3$. R^3 may represent ethyl or propyl substituted with 1-piperazinyl substituted at the 3 and 5 positions by $-CH_3$ or 1-piperazinyl substituted at the 4 position by $-CH_3$;
d) one =O. R^3 may represent ethyl substituted with 1-piperazinyl substituted in the 3 position by =O; or
e) one -C(=O)-R^{13}. R^{13} may be C_{3-8}cycloalkyl, for example cyclopropyl. R^3 may represent ethyl substituted with 1-piperazinyl substituted in the 4 position by -C(=O)-C_3H_5.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example piperazinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with two phenylC_{1-6}alkyl groups wherein the phenyl is substituted with R^{16}. R^{16} may represent C_{1-4}alkoxy, for example CH_3O-. R^3 may represent methyl substituted with 2-piperazinyl substituted in the 1 and 4 position by methylphenyl wherein the phenyl is substituted in the 4 position by CH_3O-.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an aromatic 5 membered monocyclic heterocyclyl containing four nitrogen heteroatoms, for example tetrazolyl. R^3 may represent ethyl substituted with 5-tetrazolyl.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an aromatic 5 membered monocyclic heterocyclyl containing one oxygen and two nitrogen heteroatoms, for example 1, 3, 4-oxadiazolyl. The heterocyclyl may be substituted. For example the heterocyclyl may be substituted with one $-NR^{14}R^{15}$ group, where each of R^{14} and R^{15} is hydrogen. Alternatively one of R^{14} and R^{15} may be hydrogen and the other may represent C_{1-4}alkyl optionally substituted with hydroxyl, for example $-CH_2CH_2OH$. R^3 may represent methyl substituted with 2-(1, 3, 4-oxadiazolyl) substituted at the 5 position by $-NH_2$ or 2-(1, 3, 4-oxadiazolyl) substituted at the 5 position by -NH$-CH_2CH_2OH$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing two

nitrogen heteroatoms, for example pyrazolyl or imidazolyl. R^3 may represent methyl, ethyl or propyl substituted with 1-pyrazolyl or 2-imidazoyl. R^3 may represent methyl substituted with 3-pyrazolyl or 5-pyrazolyl.The heterocyclyl may be substituted. For example the heterocyclyl may be substituted with one or two C_{1-4}alkyl, for example $-CH_3$ or $-CH_2CH_3$. R^3 may represent methyl, ethyl or propyl substituted with 1-imidazolyl substituted at the 2 position by $-CH_3$, 3-pyrazolyl substituted at the 1 and 5 positions by $-CH_3$, 1-imidazolyl substituted at the 2 and 5 positions by $-CH_3$, 1-imidazolyl substituted at the 2 and 4 positions by $-CH_3$, 2-imidazolyl substituted at the 1 position by $-CH_3$ or 2-imidazolyl substituted at the 1 position by $-CH_2CH_3$. R^3 may represent methyl substituted with 2-imidazolyl substituted at the 5 position by $-CH_3$. R^3 may represent ethyl substituted with 1-pyrazolyl substituted at the 3 position by $-CH_3$. R^3 may represent methyl substituted with 4-pyrazolyl substituted at the 1 position by $-CH_3$.In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example imidazolyl. The heterocyclyl may be substituted. For example the heterocyclyl is substituted with one C_{1-4}alkyl, for example $-CH_3$, and with one $-S(=O)_2-NR^{14}R^{15}$. R^{14} and R^{15} may each represent C_{1-4}alkyl, for example $-CH_3$. R^3 may represent methyl substituted with 2-imidazolyl substituted in the 3 position by $-S(=O)_2-N(CH_3)_2$ and in the 5 position by $-CH_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example pyrazolyl. The heterocyclyl may be substituted. For example the heterocyclyl is substituted with R^{13}. R^{13} may represent a saturated 6 membered monocyclic heterocyclyl containing one oxygen heteroatom. R^3 may represent methyl substituted with 5-pyrazolyl substituted in the 2 position by 2-tetrahydropyran. R^3 may represent methyl substituted with 3-pyrazolyl substituted in the 1 position by 2-tetrahydropyran.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example imidazolyl. The heterocyclyl may be substituted. For example the heterocyclyl is substituted with $-S(=O)_2-NR^{14}R^{15}$. R^{14} and R^{15} may each represent C_{1-4}alkyl optionally substituted with a substituent selected from hydroxyl, C_{1-}

$_4$alkoxy, amino or mono-or di(C_{1-4}alkyl)amino, for example $-CH_3$. R^3 may represent methyl substituted with 2-imidazoyl substituted in the 1 position by $-S(=O)_2-N(CH_3)_2$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing three nitrogen heteroatoms, for example triazolyl. R^3 may represent methyl substituted with 4-(1, 2, 3-triazolyl. The heterocyclyl may be substituted. For example the heterocyclyl is substituted with

a) one hydroxyC_{1-4}alkyl group, for example $-CH_2CH_2OH$. R^3 may represent methyl substituted with 4-(1, 2, 3-triazolyl) substituted in the 1 position by $-CH_2CH_2OH$ or 4-(1,2,3-triazolyl) substituted in the 2 position by $-CH_2OH$; or

b) one C_{1-4}alkyl substituted with C_{1-6}alkyl-O-C(=O)- group, for example $-CH_2-C(=O)-OCH_2CH_3$. R^3 may represent methyl substituted with 4-(1, 2, 3-triazolyl) substituted in the 1 position by $-CH_2-C(=O)-OCH_2CH_3$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted aromatic 5 membered monocyclic heterocyclyl containing three nitrogen heteroatoms, for example triazolyl. R^3 may represent ethyl substituted with 1-(1, 2, 4-triazolyl. The heterocyclyl may be substituted. For example the heterocyclyl is substituted with one C_{1-4}alkyl group, for example $-CH_3$. R^3 may represent ethyl or propyl substituted with 1-(1, 2, 4-triazolyl) substituted in the 3 position by $-CH_3$. R^3 may represent ethyl or propyl substituted with 2-(1, 2, 4-triazolyl) substituted in the 3 position by $-CH_3$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 5 membered monocyclic heterocyclyl containing one nitrogen and one oxygen heteroatom, for example oxazolidinyl. The heterocyclyl may be substituted, for example substituted with one =O. R^3 may represent ethyl or propyl substituted with 3-oxazolidinyl substituted in the 2 position by =O. R^3 may represent methyl substituted with 5-oxazolidinyl substituted in the 2 position by =O. The heterocyclyl may be substituted, for example substituted with one =O and one C_{1-6}alkyl. R^3 may represent methyl substituted with 5-oxazolidinyl substituted in the 2 position by =O and in the 3 position by $-CH(CH_3)_2$.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 6 membered monocyclic heterocyclyl containing one nitrogen and one sulphur heteroatom, for example thiomorpholinyl. The heterocyclyl may be substituted, for example substituted with two =O groups on the sulphur heteroatom. R^3 may represent propyl substituted with 4-thiomorpholinyl substituted in the 1 position by two =O groups.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 7 membered monocyclic heterocyclyl containing two nitrogen heteroatoms, for example homopiperazinyl. R^3 may represent ethyl substituted with 1-homopiperazinyl.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents a saturated 7 membered monocyclic heterocyclyl containing one nitrogen and one oxygen heteroatom, for example homomorpholinyl. R^3 may represent ethyl substituted with homomorpholinyl.

In another embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents phenyl or naphthyl, in particular phenyl. R^3 may represent $-CH_2-C_6H_5$. When R^9 represents phenyl or naphthyl, in particular phenyl, the phenyl or naphthyl group may be substituted, for example by one chlorine. R^3 may represent methyl substituted with phenyl substituted in the 2, 3 or 4 position by chlorine.

In one embodiment R^3 represents cyanoC_{1-6}alkyl, for example $-CH_2CH_2CN$ or $-CH_2CH_2CH_2CN$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with hydroxyl, halo or $-NR^{10}R^{11}$. In a further embodiment R^3 represents C_{1-6}alkyl substituted with hydroxyl or $-NR^{10}R^{11}$. In a yet further embodiment R^3 represents C_{1-6}alkyl substituted with $-NR^{10}R^{11}$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with hydroxyl, halo or $-NR^{10}R^{11}$, wherein the C_{1-6}alkyl group is a straight chain alkyl group e.g. 2-ethyl, n-propyl, n-butyl. In one embodiment R^3 represents C_{1-4}alkyl substituted with $-NR^{10}R^{11}$. In one embodiment R^3 represents C_{1-4}alkyl substituted $-NR^{10}R^{11}$, wherein the C_{1-4}alkyl group is

a straight chain alkyl group e.g. 2-ethyl, n-propyl, n-butyl. In one embodiment R^3 represents C_{1-4}alkyl substituted with -$NR^{10}R^{11}$, wherein the C_{1-4}alkyl group is an ethyl group (-CH_2CH_2-).

In one embodiment when R^3 represents C_{1-6}alkyl substituted with -$NR^{10}R^{11}$, R^{10} and R^{11} have the following meanings :

a) each of R^{10} and R^{11} represent hydrogen. R^3 may represent –$CH_2CH_2NH_2$, –$CH_2CH_2CH_2NH_2$ or –$CH_2CH_2CH_2CH_2NH_2$. R^3 may represent –$CH_2CH(CH_3)NH_2$, -$CH(CH_3)CH_2NH_2$, ;

b) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3, –CH_2CH_3 or –$CH(CH_3)_2$. R^3 may represent –$CH_2CH_2NHCH_3$, –$CH_2CH_2CH_2NHCH_3$, -$CH_2CH_2NHCH_2CH_3$, -$CH_2CH_2NHCH(CH_3)_2$, -CD_2-CD_2-$NHCH(CH_3)_2$ or –$CH_2CH_2CH_2NHCH(CH_3)_2$. R^3 may represent -$CH(CH_3)CH_2NHCH(CH_3)_2$;

c) each of R^{10} and R^{11} independently represent C_{1-6}alkyl, for example –CH_2CH_3 or –$CH(CH_3)_2$. R^3 may represent –$CH_2CH_2N(CH_2CH_3)_2$, –$CH_2CH_2N(CH_2CH_3)(CH(CH_3)_2)$. Each of R^{10} and R^{11} may independently represent C_{1-6}alkyl, for example –CH_3. R^3 may represent –$CH_2CH_2N(CH_3)_2$ or –$CH_2CH_2N(CH_3)CH(CH_3)_2$;

d) one of R^{10} and R^{11} represents hydrogen and the other represents haloC_{1-6}alkyl, for example –CH_2CF_3, –CH_2CHF_2 or –CH_2CH_2F. R^3 may represent –$CH_2CH_2CH_2NHCH_2CF_3$, -$CH_2CH_2NHCH_2CHF_2$ or –$CH_2CH_2NHCH_2CH_2F$. HaloC_{1-6}alkyl may be –$C(CH_3)_2CH_2F$. R^3 may represent -$CH(CH_3)CH_2NHCH_2CF_3$, -$CH_2CH(CH_3)NHCH_2CF_3$, -$CH_2CH_2NHCH_2CF_3$, -$CH_2CH_2CH_2NHCH_2CHF_2$ -$CH_2CH_2NHCH_2CH_2CF_3$, - $CH_2CH_2CH_2NHCH_2CHF_2$, -$CH_2CH_2CH_2NHC(CH_3)_2CH_2F$, -$CD_2$-$CD_2$-$NHCH_2CF_3$;

e) one of R^{10} and R^{11} represents hydrogen and the other represents -C(=O)-C_{1-6}alkyl, for example -C(=O)-Me. R^3 may represent –CH_2CH_2NH-C(=O)-CH_3 ;

f) one of R^{10} and R^{11} represents hydrogen and the other represents –$S(=O)_2$-C_{1-6}alkyl, for example –$S(=O)_2$-CH_3, –$S(=O)_2$-CH_2CH_3 or –$S(=O)_2$-$CH(CH_3)_2$. R^3 may represent -CH_2CH_2NH–$S(=O)_2$-CH_3, -$CH_2CH_2CH_2NH$–$S(=O)_2$-CH_3, -CH_2CH_2NH–$S(=O)_2$-CH_2CH_3 or -CH_2CH_2NH–$S(=O)_2$-$CH(CH_3)_2$;

g) one of R^{10} and R^{11} represents hydrogen and the other represents –$S(=O)_2$-$NR^{14}R^{15}$, where R^{14} and R^{15} each represent C_{1-4}alkyl optionally substituted with hydroxyl, for

example -CH_3. R^3 may represent -CH_2CH_2NH–$S(=O)_2$-$N(CH_3)_2$ or -$CH_2CH_2CH_2NH$–$S(=O)_2$-$N(CH_3)_2$;

h) one of R^{10} and R^{11} represents hydrogen and the other represents hydroxyC_{1-6}alkyl, for example –CH_2CH_2OH. R^3 may represent –$CH_2CH_2NHCH_2CH_2OH$;

i) one of R^{10} and R^{11} represents hydrogen and the other represents -C(=O)-hydroxyhaloC_{1-6}alkyl, for example -C(=O)-$C(OH)(CH_3)CF_3$. R^3 may represent –$CH_2CH_2CH_2NH$-C(=O)-$C(OH)(CH_3)CF_3$ or –CH_2CH_2NH-C(=O)-$C(OH)(CH_3)CF_3$;

j) one of R^{10} and R^{11} represents hydrogen and the other represents -C(=O)-R^6. R^6 may represent C_{3-8}cycloalkyl, for example cyclopropyl. R^3 may represent –CH_2CH_2NH-C(=O)-C_3H_5. Alternatively, R^6 may represent a saturated 6-membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example piperidinyl. The heterocyclyl may be substituted, for example substituted by one C_{1-6}alkyl group, for example –CH_3 to form N-methyl piperidinyl. R^3 may represent –CH_2CH_2NH-C(=O)-(piperidin-3-yl) where the piperidinyl is substituted at the 1 position by –CH_3;

k) one of R^{10} and R^{11} represents hydrogen and the other represents cyanoC_{1-6}alkyl, for example –CH_2CH_2CN. R^3 may represent –$CH_2CH_2NHCH_2CH_2CN$ R^3 may represent –$CH_2CH_2CH_2NHCH_2CH_2CN$;

l) one of R^{10} and R^{11} represents hydrogen and the other represents R^6. R^6 may represent C_{3-8}cycloalkyl, for example cyclopropyl or cyclopentyl, or R^6 may represent a saturated 6-membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example piperidinyl. The heterocyclyl may be substituted, for example substituted with four C_{1-6}alkyl groups, for example –CH_3 to form for example 2,2,6,6-tetramethyl-piperidinyl. R^3 may represent –$CH_2CH_2NHC_3H_5$, –$CH_2CH_2NHC_5H_9$ or –CH_2CH_2NH-(2,2,6,6-tetramethyl-piperidin-4-yl). For example, the heterocyclyl may be substituted by one -$S(=O)_2NR^{14}R^{15}$, for example -$S(=O)_2NH_2$. R^3 may represent -CH_2CH_2NH-(piperidin-4-yl) where the piperidnyl is substituted in the 1 position by -$S(=O)_2NH_2$;

m) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl substituted with R^6. R^6 may represent C_{3-8}cycloalkyl, for example cyclopropyl. R^3 may represent –$CH_2CH_2NHCH_2C_3H_5$. Alternatively R^6 may represent a saturated, 5-membered monocyclic heterocyclyl containing one oxygen heteroatom. R^3 may represent –$CH_2CH_2NHCH_2$-(tetrahydrofuran-2-yl). Alternatively R^6 may represent an aromatic, 6-membered monocyclic heterocyclyl containing one nitrogen heteroatom. R^3 may represent –$CH_2CH_2NHCH_2$-(pyridin-6-yl);

n) one of R^{10} and R^{11} represents hydrogen and the other represents -C(=O)-haloC_{1-6}alkyl, for example –C(=O)-CF_3. R^3 may represent –$CH_2CH_2NHC(=O)$-CF_3 or -$CH_2CH_2CH_2NHC(=O)$-CF_3;

o) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl substituted with –$Si(CH_3)_3$. R^3 may represent –$CH_2CH_2NHCH_2Si(CH_3)_3$; or

p) one of R^{10} and R^{11} represents C_{1-6}alkyl and the other represents C_{1-6}alkyl substituted with R^6. R^6 may represent phenyl. R^6 may represent phenyl substituted with –$NR^{14}R^{15}$ where R^{14} and R^{15} each represent hydrogen. In one embodiment one of R^{10} and R^{11} represents –CH_3 and the other represents –CH_2-C_6H_5. R^3 may represent –$CH_2CH_2N(CH_3)CH_2$-C_6H_5. In one embodiment one of R^{10} and R^{11} represents –$CH(CH_3)_2$ and the other represents –CH_2-C_6H_5 wherein the phenyl is substituted in the 4-position by –NH_2.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with -$NR^{10}R^{11}$, R^{10} and R^{11} have the following meanings:

a) one of R^{10} and R^{11} represents C_{1-6}alkyl, for example –$CH(CH_3)_2$ and the other represents C_{1-6}alkyl substituted with –$NR^{14}R^{15}$ where R^{14} and R^{15} each represent hydrogen. R^3 may represent -$CH_2CH_2N(CH(CH_3)_2)CH_2CH_2CH_2NH_2$;

b) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl substituted with –C(=O)-$NR^{14}R^{15}$ where R^{14} and R^{15} each represent hydrogen. R^3 may represent –$CH_2CH_2CH_2NHCH_2C(=O)NH_2$ or –$CH_2CH_2NHCH_2C(=O)NH_2$;

c) one of R^{10} and R^{11} represents C_{1-6}alkyl, for example –CH_3 and the other represents C_{1-6}alkoxy, for example –OCH_3. R^3 may represent -$CH_2CH_2CH_2N(CH_3)$-OCH_3.

d) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkoxy, for example –OCH_3. R^3 may represent -CH_2CH_2NH-OCH_3; or

e) one of R^{10} and R^{11} represents hydrogen and the other represents hydroxyhaloC_{1-6}alkyl, for example –$CH_2CHOHCF_3$. R^3 may represent -$CH_2CH_2NHCH_2CHOHCF_3$.

f) one of R^{10} and R^{11} represents hydrogen and the other represents carboxyl (i.e. –C(=O)-OH); R^3 may represent -$CH_2CH_2CH_2NHCOOH$.

In one embodiment R^{10} represents hydrogen or C_{1-6}alkyl, for example hydrogen, –CH_3, -CH_2CH_3 or -$CH(CH_3)_2$. In one embodiment R^{10} is hydrogen.

In one embodiment R^{11} represents hydrogen, C_{1-6}alkyl, haloC_{1-6}alkyl, -C(=O)-C_{1-6}alkyl, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, hydroxyC_{1-6}alkyl, -C(=O)-hydroxyhaloC_{1-6}alkyl, -C(=O)-R^6, cyanoC_{1-6}alkyl, R^6, -C(=O)-R^6, C_{1-6}alkyl substituted with R^6, -C(=O)-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –Si$(CH_3)_3$.

In one embodiment R^{11} represents hydrogen, –CH_3, –CH_2CH_3 or –$CH(CH_3)_2$, -CH_2CF_3, –CH_2CHF_2 or –CH_2CH_2F, -C(=O)-CH_3, –S(=O)$_2$-CH_3, –S(=O)$_2$-CH_2CH_3 , –S(=O)$_2$-$CH(CH_3)_2$, –S(=O)$_2$-$N(CH_3)_2$, –CH_2CH_2OH, -C(=O)-$C(OH)(CH_3)CF_3$, -C(=O)- cyclopropyl, –CH_2CH_2CN, cyclopropyl, cyclopentyl, 2,2,6,6-tetramethyl-piperidinyl, -$CH_2C_3H_5$, -CH_2-tetrahydrofuranyl, -C(=O)-(1-methyl-piperidin-3-yl), –C(=O)-CF_3, –$CH_2Si(CH_3)_3$, –CH_2-C_6H_5.

In one embodiment R^3 represents –$CH_2CH_2NH_2$, –$CH_2CH_2CH_2NH_2$, –$CH_2CH_2CH_2CH_2NH_2$, –$CH_2CH_2NHCH_3$, –$CH_2CH_2CH_2NHCH_3$, -$CH_2CH_2NHCH_2CH_3$, -$CH_2CH_2NHCH(CH_3)_2$, –$CH_2CH_2CH_2NHCH(CH_3)_2$, –$CH_2CH_2N(CH_2CH_3)_2$, –$CH_2CH_2N(CH_2CH_3)(CH(CH_3)_2)$, –$CH_2CH_2CH_2NHCH_2CF_3$, -$CH_2CH_2NHCH_2CHF_2$ or –$CH_2CH_2NHCH_2CH_2F$, –CH_2CH_2NH-C(=O)-CH_3, -CH_2CH_2NH–S(=O)$_2$-CH_3, -$CH_2CH_2CH_2NH$–S(=O)$_2$-CH_3, -CH_2CH_2NH–S(=O)$_2$-CH_2CH_3, -CH_2CH_2NH–S(=O)$_2$-$CH(CH_3)_2$, -CH_2CH_2NH–S(=O)$_2$-$N(CH_3)_2$, -$CH_2CH_2CH_2NH$–S(=O)$_2$-$N(CH_3)_2$, –$CH_2CH_2NHCH_2CH_2OH$, –$CH_2CH_2CH_2NH$-C(=O)-$C(OH)(CH_3)CF_3$, –CH_2CH_2NH-C(=O)-$C(OH)(CH_3)CF_3$, –CH_2CH_2NH-C(=O)-C_3H_5, –$CH_2CH_2NHCH_2CH_2CN$, $CH_2CH_2NHC_3H_5$, –$CH_2CH_2NHC_5H_9$, -CH_2CH_2-NHCO-(piperidin-3-yl) where the piperidin-3-yl is substituted in the 1 position by _-CH_3, –$CH_2CH_2NHCH_2C_3H_5$, –$CH_2CH_2NHCH_2$(tetrahydrofuran-2-yl), –CH_2CH_2NHC(=O)-CF_3, -$CH_2CH_2CH_2NHC$(=O)-CF_3 , –CH_2CH_2NH-(2,2,6,6-tetramethyl-piperidin-4-yl), –$CH_2CH_2NHCH_2Si(CH_3)_3$, –$CH_2CH_2N(CH_3)CH_2$-C_6H_5.

In one embodiment R^3 represents C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$.
In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, each of R^{10} and R^{11} represents hydrogen. R^3 may represent –$CH_2CHOHCH_2NH_2$.
In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3, –$CH(CH_3)_2$. R^3 may represent –$CH_2CHOHCH_2NHCH_3$ or –$CH_2CHOHCH_2NHCH(CH_3)_2$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, one of R^{10} and R^{11} represents hydrogen and the other represents haloC_{1-6}alkyl, for example $–CH_2CF_3$. R^3 may represent $–CH_2CHOHCH_2NHCH_2CF_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, one of R^{10} and R^{11} represents C_{1-6}alkyl, for example $–CH(CH_3)_2$, and the other represents –C(=O)-haloC_{1-6}alkyl, for example $-C(=O)-CH_2Cl$. R^3 may represent $–CH_2CHOHCH_2N(CH(CH_3)_2)-C(=O)CH_2Cl$.

In one embodiment R^3 represents hydroxyC_{1-6}alkyl, wherein hydroxyC_{1-6}alkyl includes $–CD_2CD_2OH$, $–CH_2CH_2CH_2OH$, $–CD_2CD_2CD_2OH$, $–CH_2CHOHCH_3$, $–CH_2CHOHCH_2CH_3$, $–CH_2CHOHCH(CH_3)_2$, $-CH_2CH_2C(OH)(CH_3)_2$, $-CH_2CHOHCH_2OH$ or $–CH_2C(CH_3)_2OH$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with one or two halo atoms and -$NR^{10}R^{11}$. In one embodiment each of R^{10} and R^{11} represents hydrogen. R^3 may represent $–CH_2CHFCH_2NH_2$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl. R^3 may represent $–CH_2C(=O)-O-CH_2CH_3$ or $-CH_2CH_2-C(=O)-O-CH_2CH_3$R^3 may represent $–CH(CH_3)C(=O)-O-CH_2CH_3$.

In one embodiment R^3 represents C_{1-6}alkyl (for example methyl) substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–. R^3 may represents$–CH_2-C(=O)-CH_2OCH_3$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with -C(=O)-$NR^{10}R^{11}$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, the C_{1-6}alkyl group is a straight chain alkyl group e.g. n-ethyl, n-propyl, n-butyl. In one embodiment R^3 represents C_{1-4}alkyl substituted with -C(=O)-$NR^{14}R^{15}$. In one embodiment when R^3 represents C_{1-4}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, the C_{1-4}alkyl group is a straight chain alkyl group e.g. n-ethyl, n-propyl, n-butyl. In one embodiment when R^3 represents C_{1-6}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, the C_{1-6}alkyl group is an ethyl group ($-CH_2CH_2-$).

In one embodiment when R^3 represents C_{1-6}alkyl substituted with -C(=O)-$NR^{10}R^{11}$, R^{10} and R^{11} have the following meanings :

a) R^{10} and R^{11} each represent hydrogen. R^3 may represent $-CH_2C(=O)NH_2$;

b) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl, e.g. $-CH_3$. R^3 may represent $-CH_2C(=O)NHCH_3$; C_{1-6}alkyl may be $-CH(CH_3)_2$. R^3 may represent $-CH_2C(=O)NHCH(CH_3)_2$ or $-CH_2CH_2C(=O)NHCH(CH_3)_2$;

c) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, for example $-CH_2CH_2OCH_3$. R^3 may represent $-CH_2C(=O)$-$NHCH_2CH_2OCH_3$;

d) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl substituted with R^6. R^6 may be a saturated 5-membered monocyclic heterocycle containing one nitrogen heteroatom, for example pyrrolidinyl. Alternatively R^6 may be an aromatic 5-membered monocyclic heterocycle containing two nitrogen heteroatoms, for example imidazolyl. R^3 may represent $-CH_2C(=O)$-NH-CH_2CH_2-(pyrrolidin-1-yl) or $-CH_2C(=O)$-NH-CH_2CH_2-(imidazol-2-yl);

e) one of R^{10} and R^{11} represents hydrogen and the other represents hydroxyC_{1-6}alkyl, for example $-CH_2CH_2OH$. R^3 may represent $-CH_2C(=O)$-$NHCH_2CH_2OH$; or

f) one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl substituted with $-NR^{14}R^{15}$ where R^{14} and R^{15} are both hydrogen. R^3 may represents -$CH_2C(=O)$-$NHCH_2CH_2NH_2$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with -C(=O)-$NR^{10}R^{11}$, R^{10} and R^{11} have the following meanings:

a) one of R^{10} and R^{11} represents hydrogen and the other represents haloC_{1-6}alkyl, for example $-CH_2CF_3$. R^3 may represent -$CH_2CH_2C(=O)$-$NHCH_2CF_3$;

b) one of R^{10} and R^{11} represents C_{1-6}alkyl, for example $-CH_3$ and the other represents C_{1-6}alkoxy, for example $-OCH_3$. R^3 may represent -$CH_2CH_2C(=O)N(CH_3)$-OCH_3.

c) one of R^{10} and R^{11} represents hydrogen and the other represents R^6. R^6 may be a six membered monocyclic heterocyclyl containing one or two nitrogen atoms and optionally substituted with one C_{1-6}alkyl or C_{1-6}alkoxy. R^3 may represent -$CH_2C(=O)$NH-(pyridin-2-yl) wherein the pyridin-2-yl is substituted in the 3-position by $-OCH_3$, -$CH_2C(=O)$NH-(pyridin-6-yl) wherein the pyridin-6-yl is substituted in the 4-position by $-CH_3$ or -$CH_2C(=O)$NH-(pyrimidin-2-yl) wherein the pyrimidin-2-yl is substituted in the 4-position

by –OCH_3. R^3 may represent -$CH_2C(=O)NH$-(pyridin-3-yl), -$CH_2C(=O)NH$-(pyridin-6-yl) or -$CH_2C(=O)NH$-(pyridin-4-yl).

In one embodiment R^3 represents C_{1-6}alkyl substituted with carboxyl. R^3 may represent –$CH_2C(=O)OH$ or – $CH_2CH_2C(=O)OH$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with –O-$C(=O)$-$NR^{10}R^{11}$. In one embodiment one of R^{10} and R^{11} represents hydrogen and the other represents C_{1-6}alkyl, for example –CH_3. R^3 may represent -CH_2CH_2-O-$C(=O)$-$NHCH_3$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-C_{1-6}alkyl. In one embodiment R^{12} represents hydrogen. R^3 may represent –CH_2CH_2NH-$S(=O)_2$-CH_3, –$CH_2CH_2CH_2NH$-$S(=O)_2$-CH_3, -CH_2CH_2NH-$S(=O)_2$-$CH(CH_3)_2$ or –CH_2CH_2NH-$S(=O)_2$-CH_2CH_3.

In one embodiment R^3 represents C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-$NR^{14}R^{15}$. In one embodiment R^{12} represents hydrogen and R^{14} and R^{15} each represent –CH_3. R^3 may represent –CH_2CH_2NH-$S(=O)_2$-$N(CH_3)_2$ or – $CH_2CH_2CH_2NH$-$S(=O)_2$-$N(CH_3)_2$.
In one embodiment R^3 represents C_{1-6}alkyl substituted with R^9 and optionally substituted with –O-$C(=O)$-C_{1-6}alkyl.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9 and optionally substituted with –O-$C(=O)$-C_{1-6}alkyl, R^9 represents 5-membered unsaturated ring fused to a 6-membered unsaturated ring, for example a furan ring fused to a pyridine ring, or a pyrrole ring fused to a pyridine ring, wherein the pyrrole ring is optionally substituted with one C_{1-4}alkyl, for example –CH_3. In one embodiment R^9 represents 1H-pyrrolo[3,2-b]pyridinyl, 1-methyl-1H-pyrrolo[3,2-b]pyridinyl or furo[3,2-b]pyridinyl.

In one embodiment R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9, R^9 represents a saturated 5 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyrrolidinyl. R^3 may represent propyl substituted with –OH and 1-pyrrolidinyl.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9, R^9 represents a saturated 5 membered monocyclic heterocyclyl containing one nitrogen heteroatom, for example pyrrolidinyl, and the heterocyclyl is substituted. For example the heterocyclyl is substituted with

a) two halo's, for example two fluorines. R^3 may represent propyl substituted with –OH and 1-pyrrolidinyl where the 1-pyrrolidinyl is substituted at the 3 position by two fluorines; or

b) a cyano group. R^3 may represent propyl substituted with –OH and 1-pyrrolidinyl where the 1-pyrrolidinyl is substituted at the 3 position by a cyano group.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9, R^9 represents a saturated 6 membered monocyclic heterocycle containing one nitrogen and one oxygen heteroatom, for example morpholinyl. R^3 may represent propyl substituted with –OH and 4-morpholinyl.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9, R^9 represents a saturated 6 membered monocyclic heterocycle containing one nitrogen heteroatom, for example piperidinyl. R^3 may represent propyl substituted with –OH and 1-piperidinyl.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9, R^9 represents an aromatic 5 membered monocyclic heterocycle containing three nitrogen heteroatoms, for example 1, 2, 4-triazolyl. The heterocycle may be substituted by one C_{1-4}alkyl, for example $–CH_3$. R^3 may represent propyl substituted with –OH and 2-(1, 2, 4-triazolyl) substituted in the 3 position by $–CH_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9, R^9 represents an aromatic 5 membered monocyclic heterocycle containing two nitrogen heteroatoms, for example imidazolyl. The heterocycle may be substituted by one C_{1-4}alkyl, for example $–CH_3$. R^3 may represent propyl substituted with –OH and 1-imidazolyl substituted in the 2 position by $–CH_3$.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with hydroxyl and R^9, R^9 represents an optionally substituted bicyclic heterocyclyl containing one nitrogen

heteroatom, said bicyclic heterocyclyl may be substituted for example with two =O groups. R^3 may represent propyl substituted with hydroxyl and isoindole-1,3-dione.

In one embodiment R^3 represents -C_{1-6}alkyl-$C(R^{12})$=N-O-R^{12}. R^{12} may independently be chosen from hydrogen and C_{1-4}alkyl optionally substituted with C_{1-4}alkyloxy, for example $-CH_3$ or $-CH(CH_3)_2$. R^3 may represent $-CH_2C(CH_3)$=N-O-H, $-CH_2C(CH_2OCH_3)$=N-O-H or $-CH_2C(CH(CH_3)_2)$=N-O-H.

In one embodiment R^3 represents $-S(=O)_2-NR^{14}R^{15}$, where R^{14} and R^{15} may each be C_{1-4}alkyl. R^3 may be $-S(=O)_2-N(CH_3)_2$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl. R^3 may be $-CH_2CH_2-S(=O)_2-CH_3$.

In one embodiment R^3 represents C_{1-6}alkyl substituted with $-C(=O)-R^9$. R^9 may represent a saturated 5-membered monocyclic heterocycle containing one nitrogen heteroatom, for example pyrrolidinyl. R^3 may represent $-CH_2-C(=O)-R^9$ and R^9 is 1-pyrrolidinyl.

In one embodiment R^3 represents C_{2-6}alkenyl substituted with R^9. R^9 may represent an optionally substituted aromatic 6-membered monocyclic heterocycle containing one or two nitrogen heteroatoms, for example pyridinyl or pyrimidinyl. The heterocyclyl may be substituted, for example with one C_{1-4}alkyl or one C_{1-4}alkoxy substituent, for example $-CH_3$ or $-OCH_3$. R^3 may represent $-CH_2CH$=CH-(2-pyrimidinyl), $-CH_2CH$=CH-(2-pyrimidinyl) wherein the 2-pyrimidinyl is substituted in the 4-position by $-OCH_3$, $-CH_2CH$=CH-(2-pyridinyl) wherein the 2-pyridinyl is substituted in the 4-position by $-CH_3$ or $-CH_2CH$=CH-(2-pyridinyl) wherein the 2-pyridinyl is substituted in the 3-position by $-OCH_3$.

In one embodiment R^3 represents C_{2-6}alkynyl substituted with R^9. R^9 may represent an optionally substituted aromatic 5-membered monocyclic heterocycle containing two nitrogen heteroatoms, for example imidazolyl. The heterocyclyl may be substituted, for example substituted with one C_{1-4}alkyl substituent, for example $-CH_3$. R^3 may represent -CH_2-C≡C—(2-imidazolyl) wherein the 2-imidazolyl is substituted in the 1 position by –

CH_3 or $-CH_2-C\equiv C-$(5-imidazolyl) wherein the 5-imidazolyl is substituted in the 1 position by $-CH_3$.

In one embodiment R^3 represents C_{2-6}alkynyl substituted with R^9.

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted aromatic 6-membered monocyclic heterocycle containing one or two nitrogen heteroatoms, for example pyridinyl, pyrimidinyl or pyrazinyl. R^3 may represent $-CH_2-C\equiv C-$ (4-pyridinyl), $-CH_2-C\equiv C-$ (3-pyridinyl), $-CH_2-C\equiv C-$ (2-pyridinyl), $-CH_2-C\equiv C-$ (2-pyrimidinyl), $-CH_2-C\equiv C-$ (6-pyrazinyl).

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted aromatic 6-membered monocyclic heterocycle containing one or two nitrogen heteroatoms, for example pyridinyl, pyrimidinyl or pyrazinyl and the heterocyclyl may be substituted, for example substituted with:

a) one hydroxyC_{1-4}alkyl. R^3 may represent $-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2 or 4-position with $-CH_2OH$;

b) one C_{1-4}alkoxy, for example $-OCH_3$, $-OCH_2CH_3$. R^3 may represent $-CH_2-C\equiv C-$ (4-pyridinyl) substituted in the 6-position with $-OCH_3$,$-CH_2-C\equiv C-$ (2-pyridinyl) substituted in the 3 or 5-position with $-OCH_3$,$-CH_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 4 or 6-position with $-OCH_3$,
$-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2, 4 or 5-position with $-OCH_3$,-$CH_2-C\equiv C-$ (6-pyrimidinyl) substituted in the 4 -position with $-OCH_3$,
$-CH_2-C\equiv C-$ (5-pyrazinyl) substituted in the 6 -position with $-OCH_3$,
$-CH_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 6-position with $-OCH_2CH_3$,
$-C(CH_3)_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 4-position with $-OCH_3$-$CH_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 4-position with $-OCH(CH_3)_2$;

c) one cyano. R^3 may represent $-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2 or the 4-position with cyano, $-CH_2-C\equiv C-$ (4-pyridinyl) substituted in the 5 or 6-position with cyano;

d) one – $NR^{14}R^{15}$. R^3 may represent $-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2 or 4-position with $-NH_2$, $-CH_2-C\equiv C-$ (6-pyrimidinyl) substituted in the 2-position with $-NH_2$, $-CH_2-C\equiv C-$ (2-pyridinyl) substituted in the 3-position with $-NH_2$, -$CH_2-C\equiv C-$ (3-pyrazinyl) substituted in the 6-position with $-NH_2$, $-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 5-position with $-NHCH_3$,

e) one C_{1-4}alkyl, for example $-CH_3$ or $-CH_2CH_3$. R^3 may represent -CH_2 —C≡C— (6-pyridinyl) substituted in the 3 or 4-position with $-CH_3$, -CH_2 —C≡C— (2-pyridinyl) substituted in the 3-position with $-CH_3$, -CH_2 —C≡C— (2-pyrimidinyl) substituted in the 4-position with $-CH_3$, -CH_2 —C≡C— (2-pyrimidinyl) substituted in the 6-position with $-CH_2CH_3$,

f) one C_{1-4}alkyl, for example $-CH_3$ and one $-NR^{14}R^{15}$, for example $-NH_2$. R^3 may represent -CH_2 —C≡C— (6-pyrimidinyl) substituted in the 2-position with $-CH_3$ and in the 4-position with $-NH_2$;

g) one halogen, for example –Cl and one – $NR^{14}R^{15}$, for example $-NH_2$. R^3 may represent -CH_2 —C≡C— (6-pyrimidinyl) substituted in the 2-position with $-NH_2$ and in the 4-position with –Cl,

h) one halogen, for example –Br, –Cl or –F. R^3 may represent -CH_2 —C≡C— (2-pyrazinyl) substituted in the 3-position with –Cl, -CH_2 —C≡C— (3-pyrazinyl) substituted in the 5-position with –Cl, -CH_2 —C≡C— (2-pyridinyl) substituted in the 3-position with –F, -CH_2 —C≡C— (5-pyridinyl) substituted in the 6-position with –Br;

i) one $-C(=O)-NR^{14}R^{15}$. R^3 may represent -CH_2 —C≡C— (6-pyridinyl) substituted in the 4-position with $-C(=O)-NH_2$;

j) one C_{1-4}alkyl-O-C(=O)-. R^3 may represent -CH_2 —C≡C— (6-pyridinyl) substituted in the 5-position with CH_3-O-C(=O)-, -CH_2 —C≡C— (2-pyrimidinyl) substituted in the 6-position with CH_3-O-C(=O)-;

k) one haloC_{1-4}alkyl. R^3 may represent -CH_2 —C≡C— (2-pyridinyl) substituted in the 3-position with – CF_3.

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted aromatic 5-membered monocyclic heterocyclyl containing one nitrogen and one sulphur heteroatom, for example thiazolyl. R^3 may represent -CH_2 —C≡C— (5-thiazolyl).

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted phenyl. R^3 may be -CH_2 —C≡C— (phenyl). The phenyl may be substituted, for example with one C_{1-4}alkoxy. R^3 may represent -CH_2 —C≡C— (phenyl) where the phenyl is substituted in the 5-position by $-OCH_3$.

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted saturated 4-membered monocyclic heterocycle containing one nitrogen heteroatom, for example azetidinyl. The heterocyclyl may be substituted, for example with:

a) one hydroxyl and one C_{1-4}alkyl-O-C(=O)-. R^3 may represent -CH_2 —C≡C— (3-azetidinyl) substituted in the 1-position by $(CH_3)_3$C-O-C(=O)- and in the 3-position by –OH;

b) one hydroxyl. R^3 may represent -CH_2 —C≡C— (3-azetidinyl) substituted in the 3-position by –OH.

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted saturated 5-membered monocyclic heterocycle containing one nitrogen heteroatom, for example pyrrolidinyl. The heterocyclyl may be substituted, for example with:

a) one hydroxyl and one C_{1-4}alkyl-O-C(=O)-. R^3 may represent -CH_2 —C≡C— (3-pyrrolidinyl) substituted in the 1-position by $(CH_3)_3$C-O-C(=O)- and in the 3-position by –OH;

b) one hydroxyl. R^3 may represent -CH_2 —C≡C— (3-pyrrolidinyl) substituted in the 3-position by –OH.

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted saturated 6-membered monocyclic heterocycle containing one nitrogen heteroatom, for example piperidinyl. R^3 may represent -CH_2 —C≡C— (4-piperidinyl). The heterocyclyl may be substituted, for example with:

a) one hydroxyl. R^3 may represent -CH_2 —C≡C— (4-piperidinyl) substituted in the 4-position by –OH;

b) one C_{1-4}alkyl-O-C(=O)-. R^3 may represent -CH_2 —C≡C— (4-piperidinyl) substituted in the 1-position by $(CH_3)_3$C-O-C(=O)- .

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted saturated 5-membered monocyclic heterocycle containing one oxygen heteroatom, for example tetrahydrofuranyl. The heterocyclyl may be substituted, for example with one hydroxyl. R^3 may represent -CH_2 —C≡C— (4-tetrahydrofuranyl) substituted in the 3-position by –OH.

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents an optionally substituted saturated 6-membered monocyclic heterocycle containing one oxygen heteroatom, for example tetrahydropyranyl. The heterocyclyl may be substituted, for example with one hydroxyl. R^3 may represent -CH_2—C≡C— (4-tetrahydropyranyl) substituted in the 4-position by –OH.

In one embodiment when R^3 represents C_{2-6}alkynyl substituted with R^9, R^9 represents a C_{3-8}cycloalkyl, for example cyclohexyl.
R^3 may represent -CH_2—C≡C— (cyclohexyl).

In one embodiment R^3 represents C_{2-6}alkynyl (e.g. –CH_2—C≡C—) substituted with R^9, wherein R^9 represents C_{3-8}cycloalkyl or 3 to 12 membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, said C_{3-8}cycloalkyl or 3 to 12 membered monocyclic or bicyclic heterocyclyl each optionally and each independently being substituted with 1, 2, 3, 4 or 5 substituents as defined herein.

In one embodiment R^3 represents C_{2-6}alkynyl (e.g. –CH_2—C≡C—) substituted with R^9, wherein R^9 represents an optionally substituted 4 to 8-membered monocyclic or bridged heterocyclyl, for example R^9 represents an optionally substituted azetidinyl, pyrrolidinyl, imidazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, piperidinyl, tetrahydrofuranyl, tetrahydropyranyl, or 2,5-diaza-bicyclo[2.2.1]heptanyl.

In one embodiment R^3 represents C_{2-6}alkynyl (e.g. –CH_2—C≡C—) substituted with R^9, wherein R^9 represents

- an optionally substituted aromatic 5- or 6-membered monocyclic heterocyclyl, for example imidazolyl, thiazolyl, pyridinyl, pyrimidinyl or pyrazinyl.
- an optionally substituted saturated 4-, 5-,or 6-membered monocyclic heterocyclyl, for example azetidinyl, pyrrolidinyl, piperidinyl, tetrahydrofuranyl, tetrahydropyranyl.
- an optionally substituted 6 to 8 membered bridged heterocyclyl group, for example 2,5-diaza-bicyclo[2.2.1]heptanyl.
- a C_{3-8}cycloalkyl, for example cyclohexyl.

In one embodiment R^3 represents C_{2-6}alkynyl (e.g. $-CH_2-C\equiv C-$) substituted with R^9, wherein R^9 represents

- an optionally substituted aromatic 5-membered monocyclic heterocycle containing two nitrogen heteroatoms, for example imidazolyl,
- an optionally substituted aromatic 6 membered monocyclic heterocycle containing one nitrogen heteroatom, for example pyridinyl,
- an optionally substituted aromatic 6 membered monocyclic heterocycle containing one or two nitrogen heteroatoms, for example pyridinyl, pyrimidinyl or pyrazinyl
- an optionally substituted aromatic 5-membered monocyclic heterocyclyl containing one nitrogen and one sulphur heteroatom, for example thiazolyl,
- an optionally substituted saturated 4-membered monocyclic heterocycle containing one nitrogen heteroatom, for example azetidinyl,
- an optionally substituted saturated 5-membered monocyclic heterocycle containing one nitrogen heteroatom, for example pyrrolidinyl,
- an optionally substituted saturated 5-membered monocyclic heterocycle containing one oxygen heteroatom, for example tetrahydrofuranyl,
- an optionally substituted saturated 6-membered monocyclic heterocycle containing one oxygen heteroatom, for example tetrahydropyranyl,
- a C_{3-8}cycloalkyl, for example cyclohexyl or
- a 6 to 8 membered bridged heterocyclyl group, for example 2,5-diaza-bicyclo[2.2.1]heptanyl.

In one embodiment when R^3 represents C_{1-6}alkyl substituted with R^9, R^9 represents an optionally substituted 6 to 8 membered bridged heterocyclyl group, for example 2,5-diaza-bicyclo[2.2.1]heptanyl optionally substituted by $-C(=O)-O-C_4$alkyl.

In one embodiment R^3 represents C_{1-6}alkyloxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups. R^3 may represent $-CH_2CHOHCH_2OCH_3$.

In one embodiment R^3 represents C_{2-6}alkenyl. R^3 may represent $-CH_2-CH=CH_2$.

In one embodiment R^3 represents C_{2-6}alkynyl. R^3 may represent $^{-CH_2-C\equiv C-H}$. R^3 may represent $-C(CH_3)_2-C\equiv C-H$.

In one embodiment R^3 represents R^{13}.

In one embodiment when R^3 represents R^{13}, R^{13} represents a saturated 4-membered monocyclic heterocycle containing one oxygen heteroatom. R^3 may represent 3–oxetanyl.

In another embodiment when R^3 represents R^{13}, R^{13} represents an optionally substituted C_{3-8}cycloalkyl. For example the C_{3-8}cycloalkyl may be substituted with one $NR^{14}R^{15}$ where one of R^{14} and R^{15} represents hydrogen and the other represents C_{1-4}alkyl optionally substituted with hydroxyl, for example $–CH(CH_3)_2$. R^3 may represent cyclohexanyl substituted in the 4 position by $-NH–CH(CH_3)_2$.

In one embodiment of the invention R^3 represents C_{1-6}alkyl substituted by R^9, wherein R^9 is a saturated heterocyclyl substituted by R^{13}, wherein R^{13} is a saturated heterocyclyl which is optionally substituted, for example substituted by $–C(=O)-C_{1-6}$alkyl. In one embodiment R^9 is piperazinyl substituted by R^{13}, wherein R^{13} is piperidinyl substituted by $–C(=O)-C_{1-6}$alkyl.

In one embodiment of the invention R^3 represents C_{1-6}alkyl substituted with $-P(=O)(OC_{1-6}alkyl)_2$. R^3 may represent $–CH_2CH_2P(=O)(OCH_2CH_3)_2$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $–CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example $–CH_2CH_2NHCH(CH_3)_2$.

In a further embodiment of the invention R^1 represents C_{1-6}alkyl, for example $–CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, R^3 represents C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example $–CH_2CH_2-CH_2-NHCH_2CF_3$.

In a further embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, R^3 represents C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example $-CH_2CH_2NH_2$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{2-6}alkynyl substituted with $-R^9$, for example -CH_2 —C≡C— (2-pyridinyl).

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{2-6}alkynyl substituted with $-R^9$, for example -CH_2 —C≡C— (2-pyridinyl) substituted in the 3-position by $-OCH_3$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CD_3O-, and R^3 represents C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example R^3 may represent -CD_2-CD_2-$NHCH(CH_3)_2$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{2-6}alkynyl substituted with $-R^9$, for example -CH_2 —C≡C— (6-pyridinyl) substituted in the 2-position by $-NH_2$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{2-6}alkynyl substituted with $-R^9$, for example -CH_2 —C≡C— (2-pyrimidinyl) substituted in the 4-position by $-OCH_3$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH(CH_3)_2$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CD_3O-, and R^3 represents C_{2-6}alkynyl substituted with $-R^9$, for example -CH_2 —C≡C— (4-pyridinyl).

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH(CH_3)_2$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups or with –O-C(=O)-C_{1-6}alkyl, for example $-CH_2CHOHCH_2OCH_3$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{2-6}alkynyl substituted with $-R^9$, for example -CH_2 —C≡C— (6-pyridinyl) substituted in the 4-position by $-CH_3$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl substituted with $-NR^4R^5$, for example $-CH_2CH_2CH_2NH_2$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents hydroxyhaloC_{1-6}alkyl, for example $-CH_2CHOHCF_3$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 4 and two R^2 represent C_{1-4}alkoxy, for example CH_3O-, and two R^2 represent halogen, for example F, and R^3 represents C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example $-CH_2CH_2NH(CH(CH_3)_2)$.

In one embodiment of the invention R^1 represents C_{1-6}alkyl, for example $-CH_3$, each R^{1a} represents hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy, for example CH_3O-, and R^3 represents C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example – $CH_2CH_2CH_2NH_2$.

In a further embodiment the compound of formula (I) as defined herein is selected from the following compounds or is one of the following compounds:

N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine (compound 84)

3-{4-[3-(4-{7-[(Cyclopropylmethyl)(3,5-dimethoxyphenyl)amino]quinoxalin-2-yl}-1H-pyrazol-1-yl)propyl]piperazin-1-yl}propan-1-ol or HCl salt thereof (compound 130)

N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine (compound 4)

2-[4-(7-{(Cyclopropylmethyl)[3-(2-hydroxyethoxy)-5-methoxyphenyl]amino}quinoxalin-2-yl)-1H-pyrazol-1-yl]ethanol or HCl salt thereof (compound 131)

N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-imidazol-2-yl)prop-2-yn-1-yl]-3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-amine (compound 300)

1-(3-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}propyl)pyrrolidin-2-one (compound 132)

(3S)-1-(2-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}ethyl)pyrrolidine-3-carbonitrile (compound 133)

N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]-N'-(2,2,2-trifluoroethyl)propane-1,3-diamine (compound 5)

2-(4-{7-[(3,5-Dimethoxyphenyl){2-[(1-methylethyl)amino]ethyl}amino]quinoxalin-2-yl}-1H-pyrazol-1-yl)-N-methylacetamide or HCl salt thereof (compound 134)

N-(3,5-Dimethoxyphenyl)-N-[3-(1-ethyl-1H-pyrazol-4-yl)quinoxalin-6-yl]-N'-(1-methylethyl)ethane-1,2-diamine or HCl salt thereof (compound 135)

N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-{3-[1-(tetrahydro-2H-pyran-4-ylmethyl)-1H-pyrazol-4-yl]quinoxalin-6-yl}ethane-1,2-diamine or HCl salt thereof (compound 136)

(2S)-3-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}propane-1,2-diol (compound 98)

N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-[3-(1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine or HCl salt thereof (compound 137)

N-(3,5-Dimethoxyphenyl)-N-(1H-imidazol-2-ylmethyl)-3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-amine (compound 99)

3-{(Cyclopropylmethyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}-5-fluoro-N-methylbenzamide (compound 138)

1-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}-3-[(2,2,2-trifluoroethyl)amino]propan-2-ol (compound 139)

3-[(2-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}ethyl)amino]propanenitrile (compound 140)

4-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}-2-methylbutan-2-ol (compound 141)

(2S)-1-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}-3-[(2,2,2-trifluoroethyl)amino]propan-2-ol (compound 142)

N-[2-(4-Acetylpiperazin-1-yl)ethyl]-N-(3,5-dimethoxyphenyl)-3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-amine (compound 143)

4-(2-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}ethyl)piperazin-2-one (compound 144)

(2S)-1-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}-3-[(1-methylethyl)amino]propan-2-ol or HCl salt thereof (compound 145)

N-(3,5-Dimethoxyphenyl)-3-(1-methyl-1H-pyrazol-4-yl)-N-(pyrazin-2-ylmethyl)quinoxalin-6-amine (compound 146)

N-(3,5-Dimethoxyphenyl)-N-{3-[1-(1-methylethyl)-1H-pyrazol-4-yl]quinoxalin-6-yl}-N'-(2,2,2-trifluoroethyl)propane-1,3-diamine or HCl salt thereof (compound 147)

(2R*)-3-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}-1,1,1-trifluoropropan-2-ol (relative stereochemistry)(Compound 148)

(2S*)-3-{(3,5-Dimethoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}-1,1,1-trifluoropropan-2-ol (relative stereochemistry) (compound 149) ;

a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof.

In a further embodiment the compound of formula (I) as defined herein is selected from the following compounds or is one of the following compounds:

N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]-N'-(2,2,2-trifluoroethyl)propane-1,3-diamine (compound 5)

N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine (compound 4)

N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine (compound 84)

N-(3,5-Dimethoxyphenyl)-N-{3-[1-(1-methylethyl)-1H-pyrazol-4-yl]quinoxalin-6-yl}-N'-(2,2,2-trifluoroethyl)propane-1,3-diamine or HCl salt thereof (compound 147)

N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-[3-(1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine or HCl salt thereof (compound 137)

N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-{3-[1-(tetrahydro-2H-pyran-4-ylmethyl)-1H-pyrazol-4-yl]quinoxalin-6-yl}ethane-1,2-diamine or HCl salt thereof (compound 136)

N-(3,5-Dimethoxyphenyl)-N-[3-(1-ethyl-1H-pyrazol-4-yl)quinoxalin-6-yl]-N'-(1-methylethyl)ethane-1,2-diamine or HCl salt thereof (compound 135)

2-(4-{7-[(3,5-Dimethoxyphenyl){2-[(1-methylethyl)amino]ethyl}amino]quinoxalin-2-yl}-1H-pyrazol-1-yl)-N-methylacetamide or HCl salt thereof (compound 134)

N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-imidazol-2-yl)prop-2-yn-1-yl]-3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-amine (compound 300) ;
a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof.

In a further embodiment the compound of formula (I) as defined herein is selected from the following compounds or is one of the following compounds:
N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]-N'-(2,2,2-trifluoroethyl)propane-1,3-diamine (compound 5)
N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine (compound 4)
N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine (compound 84) ;
a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof.

In a further embodiment the compound of formula (I) as defined herein is selected from the following compounds or is one of the following compounds:
N-(3,5-Dimethoxyphenyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]propane-1,3-diamine; (compound 93)
2-(4-{7-[(3,5-Dimethoxyphenyl){2-[(1-methylethyl)amino]ethyl}amino]quinoxalin-2-yl}-1H-pyrazol-1-yl)ethanol; (compound 691)
N-(3,5-Dimethoxyphenyl)-N'-(1-methylethyl)-N-(3-{1-[2-(methylsulfonyl)ethyl]-1H-pyrazol-4-yl}quinoxalin-6-yl)ethane-1,2-diamine; (compound 678)
N-(3,5-Dimethoxyphenyl)-3-(1-methyl-1H-pyrazol-4-yl)-N-(3-pyridin-2-ylprop-2-yn-1-yl)quinoxalin-6-amine; (compound 691)
N-(3,5-Dimethoxyphenyl)-N-[3-(3-methoxypyridin-2-yl)prop-2-yn-1-yl]-3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-amine; (compound 652)
N-{3,5-Bis[($^2H_3$)methyloxy]phenyl}-N'-(1-methylethyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]($^2H_4$)ethane-1,2-diamine; (compound 618)
N-[3-(6-Aminopyridin-2-yl)prop-2-yn-1-yl]-N-(3,5-dimethoxyphenyl)-3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-amine; (compound 689)
N-(3,5-Dimethoxyphenyl)-N-[3-(4-methoxypyrimidin-2-yl)prop-2-yn-1-yl]-3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-amine; (compound 688)
N-{3,5-Bis[($^2H_3$)methyloxy]phenyl}-3-[1-(1-methylethyl)-1H-pyrazol-4-yl]-N-(3-pyridin-4-ylprop-2-yn-1-yl)quinoxalin-6-amine; (compound 653)

1-[(3,5-Dimethoxyphenyl){3-[1-(1-methylethyl)-1H-pyrazol-4-yl]quinoxalin-6-yl}amino]-3-methoxypropan-2-ol; or its hydrochloric acid salt; (compound 657)


N-(3,5-Dimethoxyphenyl)-3-(1-methyl-1H-pyrazol-4-yl)-N-[3-(4-methylpyridin-2-yl)prop-2-yn-1-yl]quinoxalin-6-amine; (compound 634)

3-[{3-[1-(3-Aminopropyl)-1H-pyrazol-4-yl]quinoxalin-6-yl}(3,5-dimethoxyphenyl)amino]-1,1,1-trifluoropropan-2-ol; or an enantiomer thereof;(compound 660 and 661)

N-(2,6-Difluoro-3,5-dimethoxyphenyl)-N'-(1-methylethyl)-N-[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]ethane-1,2-diamine (compound 687);

a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof.

According to an aspect of the invention there is provided compounds of formula (I):



(I)

including any tautomeric or stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2, 3 or 4;

R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, cyanoC_{1-4}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with $-NR^4R^5$, C_{1-6}alkyl substituted with $-C(=O)-NR^4R^5$, $-S(=O)_2-C_{1-6}$alkyl, $-S(=O)_2-NR^{14}R^{15}$, C_{1-6}alkyl substituted with $-S(=O)_2-C_{1-6}$alkyl, C_{1-6}alkyl substituted with $-NH-S(=O)_2-C_{1-6}$alkyl, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with $-C(=O)-R^6$, hydroxyC_{1-6}alkyl substituted with R^6, or C_{1-6}alkyl substituted with $-Si(CH_3)_3$;

each R^{1a} is independently selected from hydrogen, C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with di(C_{1-4}alkyl)amino, and C_{1-4}alkyl substituted with one or more fluoro atoms;

each R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkyl, haloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkoxy substituted with R^{13}, $-C(=O)-R^{13}$, C_{1-4}alkyl substituted with NR^7R^8, C_{1-4}alkoxy substituted with NR^7R^8, $-NR^7R^8$ and $-C(=O)-NR^7R^8$; or when two R^2 groups are attached to adjacent carbon atoms they may be

taken together to form a radical of formula -O-(C(R^{17})$_2$)$_p$-O- wherein R^{17} represents hydrogen or fluorine and p represents 1 or 2;

R^3 represents C_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, hydroxyC_{2-6}alkynyl, haloC_{1-6}alkyl, haloC_{1-6}alkyl optionally substituted with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups or with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with R^9, C_{1-6}alkyl substituted with -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with one or two halogens and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with carboxyl, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –NR^{12}-S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with and R^9 and optionally substituted with –O-C(=O)-C_{1-6}alkyl, C_{1-6}alkyl substituted with hydroxyl and R^9, -C_{1-6}alkyl-C(R^{12})=N-O-R^{12}, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-R^9, C_{2-6}alkenyl substituted with R^9, C_{2-6}alkynyl substituted with R^9, hydroxyC_{1-6}alkoxy, C_{2-6}alkenyl, C_{2-6}alkynyl, R^{13} or C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)– or C_{1-6}alkyl substituted with -P(=O)(OC_{1-6}alkyl)$_2$;

R^4 and R^5 each independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-haloC_{1-6}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-$NR^{14}R^{15}$, R^{13} or C_{1-6}alkyl substituted with R^{13};

R^6 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C_{1-6}alkyl, cyanoC_{1-6}alkyl, hydroxyl, carboxyl, hydroxyC_{1-6}alkyl, halogen, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}

$_6$alkyl, C_{1-6}alkoxy, C_{1-6}alkoxyC_{1-6}alkyl, C_{1-6}alkyl-O-C(=O)-, -$NR^{14}R^{15}$, -C(=O)-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$;

R^7 and R^8 each independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl or C_{1-6}alkoxyC_{1-6}alkyl;

R^9 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, naphthyl, or 3 to 12 membered monocyclic or bicyclic heterocyclyl each optionally and each independently being substituted with 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from =O, C_{1-4}alkyl, hydroxyl, carboxyl, hydroxyC_{1-4}alkyl, cyano, cyanoC_{1-4}alkyl, C_{1-4}alkyl-O-C(=O)-, C_{1-4}alkyl substituted with C_{1-4}alkyl-O-C(=O)-, C_{1-4}alkyl-C(=O)-, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, halogen, haloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkyl, -$NR^{14}R^{15}$, -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkoxy, –$S(=O)_2$-C_{1-4}alkyl, –$S(=O)_2$-haloC_{1-4}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-C_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$, R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with R^{13}, phenyl optionally substituted with R^{16}, phenylC_{1-6}alkyl wherein the phenyl is optionally substituted with R^{16}, a 5 or 6-membered aromatic monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S wherein said heterocyclyl is optionally substituted with R^{16};
or when two of the substituents of R^9 are attached to the same atom, they may be taken together to form a 4 to 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S;

R^{10} and R^{11} each independently represent hydrogen, C_{1-6}alkyl, cyanoC_{1-6}alkyl, C_{1-6}alkyl substituted with –$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –C(=O)-$NR^{14}R^{15}$, haloC_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, R^6,

C_{1-6}alkyl substituted with R^6, -C(=O)-R^6, -C(=O)-C_{1-6}alkyl, –C(=O)-hydroxyC_{1-6}alkyl, -C(=O)-haloC_{1-6}alkyl,-C(=O)-hydroxyhaloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$Si(CH_3)_3$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, -$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$;

R^{12} represents hydrogen or C_{1-4}alkyl optionally substituted with C_{1-4}alkoxy;


R^{13} represents C_{3-8}cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C_{3-8}cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C_{1-6}alkyl, -C(=O)-C_{1-6}alkyl, C_{1-6}alkoxy, or -$NR^{14}R^{15}$;

R^{14} and R^{15} each independently represent hydrogen, or haloC_{1-4}alkyl, or C_{1-4}alkyl optionally substituted with a substituent selected from hydroxyl, C_{1-4}alkoxy, amino or mono- or di(C_{1-4}alkyl)amino;

R^{16} represents hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{1-4}alkoxy, -$NR^{14}R^{15}$ or –C(=O)$NR^{14}R^{15}$;

the *N*-oxides thereof, the pharmaceutically acceptable salts thereof or the solvates thereof.

In one embodiment there is provided a compound of formula (I^0):



(I^0)

including any stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2, 3 or 4;

R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with -NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-}

$_6$alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, or C_{1-6}alkyl substituted with –Si$(CH_3)_3$;

each R^2 is independently selected from halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkoxy substituted with R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with NR^7R^8, C_{1-4}alkoxy substituted with NR^7R^8, -NR^7R^8 or -C(=O)-NR^7R^8;

R^3 represents C_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxy groups, C_{1-6}alkyl substituted with R^9, C_{1-6}alkyl substituted with -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with hydroxyl and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with one or two halogens and -$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with carboxyl, C_{1-6}alkyl substituted with –O-C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –NR^{12}-S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with hydroxyl and R^9, -C_{1-6}alkyl-C(R^{12})=N-O-R^{12}, C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, C_{1-6}alkyl substituted with –C(=O)-R^9, C_{2-6}alkynyl substituted with R^9, hydroxyC_{1-6}alkoxy, C_{2-6}alkenyl, C_{2-6}alkynyl, R^{13}, or C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–;

R^4 and R^5 independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-haloC_{1-6}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-$NR^{14}R^{15}$, R^{13} or C_{1-6}alkyl substituted with R^{13};

R^6 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4, 5, 6 or 7-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4, 5, 6 or 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C_{1-6}alkyl, cyanoC_{1-6}alkyl, hydroxyl, carboxyl, hydroxyC_{1-6}alkyl, halogen, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}alkoxyC_{1-6}alkyl, C_{1-6}alkyl-O-C(=O)-, -$NR^{14}R^{15}$,

-C(=O)-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$;

R^7 and R^8 independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl or C_{1-6}alkoxyC_{1-6}alkyl;

R^9 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl or a 3 to 12 membered monocyclic or bicyclic heterocyclyl containing at least one heteroatom selected from N, O or S, said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, aryl or a 3 to 12 membered monocyclic or bicyclic heterocyclyl each optionally and each independently being substituted with 1 to 5 substituents, each substituent independently being selected from =O, C_{1-4}alkyl, hydroxyl, carboxyl, hydroxyC_{1-4}alkyl, cyano, cyanoC_{1-4}alkyl, C_{1-4}alkyl-O-C(=O)-, C_{1-4}alkyl substituted with C_{1-6}alkyl-O-C(=O)-, C_{1-4}alkyl-C(=O)-, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, halogen, haloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkyl, -$NR^{14}R^{15}$, -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -$NR^{14}R^{15}$, C_{1-4}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, C_{1-4}alkoxy, –$S(=O)_2$-C_{1-4}alkyl, –$S(=O)_2$-haloC_{1-4}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-C_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-4}alkyl, C_{1-4}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$, R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with R^{13}, phenyl optionally substituted with R^{16}, phenylC_{1-6}alkyl wherein the phenyl is optionally substituted with R^{16}, a 5 or 6-membered aromatic monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S wherein said heterocyclyl is optionally substituted with R^{16};

or when two of the substituents of R^9 are attached to the same atom, they may be taken together to form a 4, 5, 6 or 7-membered saturated monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S ;

R^{10} and R^{11} each independently represent hydrogen, C_{1-6}alkyl, cyanoC_{1-6}alkyl, C_{1-6}alkyl substituted with –$NR^{14}R^{15}$, haloC_{1-6}alkyl, hydroxyC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, R^6, C_{1-6}alkyl substituted with R^6, -C(=O)-R^6, -C(=O)-C_{1-6}alkyl, –C(=O)-hydroxyC_{1-6}alkyl, -C(=O)-haloC_{1-6}alkyl,-C(=O)-hydroxyhaloC_{1-6}alkyl, C_{1-}

$_6$alkyl substituted with –$Si(CH_3)_3$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, -$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$;

R^{12} represents hydrogen or C_{1-4}alkyl optionally substituted with C_{1-4}alkoxy;


R^{13} represents C_{3-8}cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C_{3-8}cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C_{1-6}alkyl, -(C=O)-C_{1-6}alkyl, C_{1-6}alkoxy, or -$NR^{14}R^{15}$;

R^{14} and R^{15} each independently represent hydrogen, or haloC_{1-4}alkyl, or C_{1-4}alkyl optionally substituted with a substituent selected from hydroxyl, C_{1-4}alkoxy, amino or mono- or di(C_{1-4}alkyl)amino;

R^{16} represents hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{1-4}alkoxy, -$NR^{14}R^{15}$ or –C(=O)$NR^{14}R^{15}$;

and a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof.

In one embodiment there is provided a compound of formula (I^0):



(I^0)

including any stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2, 3 or 4;

R^1 represents hydrogen,

C_{1-6}alkyl, for example -CH_3, -CD_3, -CH_2CH_3, -$CH_2CH_2CH_3$, -$CH_2CH(CH_3)_2$, -$CH(CH_3)_2$, -$CH_2CH(CH_3)_2$,

C_{2-4}alkenyl, for example –CH_2-CH=CH_2,

hydroxyC_{1-6}alkyl, for example -CH_2CH_2OH, -$CH_2C(CH_3)_2OH$ or $CH_2CHOHCH_2OH$,

haloC_{1-6}alkyl, for example -CH_2CH_2F, $CH_2CH_2CH_2Cl$ or CH_2CH_2Br,

C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, for example -$CH_2CH_2OCH_3$,

C_{1-6}alkyl substituted with -NR^4R^5, for example -$CH_2CH_2NH_2$ or -$CH_2CH_2CH_2NH_2$, -$CH_2CH_2NHCH_3$, -$CH_2CH_2NHS(=O)_2N(CH_3)_2$, -$CH_2CH_2NHS(=O)_2CH_3$,

C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, for example -$CH_2C(=O)N(CH_3)_2$, -$CH_2C(=O)NHCH_3$ or -$C(CH_3)_2C(=O)NHCH_3$, -$C(CH_3)_2C(=O)NHCH_2CH_2OH$ or -$CH_2C(=O)NHCH_2CH_2OH$, -$CH_2C(=O)NHCH_2CH_2OCH_3$ or -$C(CH_3)_2C(=O)NHCH_2CH_2OCH_3$, –$CH_2$-C(=O)-NH-$CH_2$-$CH_2$-(pyrrolidin-1-yl), -$CH_2CH_2CH_2NHCH_2CH_2$-S(=O)$_2$-$CH_3$,

–S(=O)$_2$-C_{1-6}alkyl, for example –S(=O)$_2$-CH_3,

–S(=O)$_2$-$NR^{14}R^{15}$, for example –S(=O)$_2$-$N(CH_3)_2$,

C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, for example -CH_2CH_2S(=O)$_2$-CH_3,

C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, for example -$CH_2CH_2NHS(=O)_2$-CH_3,

R^6, for example 4-piperidinyl ,2-tetrahydropyranyl or 4-tetrahydropyranyl, 4-tetrahydrofuranyl , 3-azetidinyl substituted in the 1 position by –CH_2CH_2OH, 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)-, 4-piperidinyl substituted on the nitrogen atom with –S(=O)$_2$$CH_3$, 4-piperidinyl substituted on the nitrogen atom with –CH_3 ,

C_{1-6}alkyl substituted with R^6, for example methyl or ethyl each substituted with 4-piperidinyl, 4-piperazinyl, 1-pyrrolidinyl or 4-tetrahydropyranyl; propyl substituted with morpholinyl where the morpholinyl is linked to the propyl through the N heteroatom; methyl, ethyl or propyl each substituted with 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)-, 4-piperidinyl substituted on the nitrogen atom with –CH_3, 4-piperazinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)-, 4-piperazinyl substituted on the nitrogen atom with -CH_2CH_2OH, 4-piperazinyl substituted on the nitrogen atom with -$CH_2CH_2CH_2OH$, 1-piperidinyl substituted in the 1 position by –OH, 1-piperidinyl substituted in the 1 position by –O-CH_3; methyl substituted with 2-thiophenyl substituted in the 5 position with chlorine; methyl substituted with 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)- and in the 4 position by –OH,

C_{1-6}alkyl substituted with –C(=O)-R^6, for example –C$(CH_3)_2$-C(=O)-(piperazin-4-yl), –C$(CH_3)_2$-C(=O)-(piperazin-4-yl) substituted on the nitrogen atom in the 1 position by $(CH_3)_3$C-O-C(=O)-, –CH_2-C(=O)-(pyrrolidin-1-yl) substituted in the 3 position by –OH,

hydroxyC_{1-6}alkyl substituted with R^6, for example $-CH_2CHOHCH_2-$ substituted with 1-piperidinyl,

C_{1-6}alkyl substituted with $-Si(CH_3)_3$, for example $-CH_2Si(CH_3)_3$, or

cyanoC_{1-4}alkyl, for example $-CH_2CH_2CN$;

each R^2 is independently selected from

hydroxyl,

halogen, for example fluorine, chlorine or bromine,

cyano,

C_{1-4}alkyl, for example $-CH_3$,

C_{2-4}alkenyl, for example $-CH=CH_2$,

C_{1-4}alkoxy, for example CH_3O-, $(CH_3)_2CHO-$, CH_3CH_2O-, CD_3O-,

hydroxyC_{1-4}alkyl, for example $-CH_2OH$,

hydroxyC_{1-4}alkoxy, for example $-OCH_2CH_2OH$,

haloC_{1-4}alkyl, for example $-CF_3$,

haloC_{1-4}alkoxy, for example $-OCH_2CH_2F$, CHF_2O- or $-OCF_3$,

C_{1-4}alkoxyC_{1-4}alkyl, for example $-CH_2CH_2OCH_3$,

R^{13}, for example 2-dioxolanyl,

C_{1-4}alkoxy substituted with R^{13}, for example $-OCH_2C_3H_5$,

$-C(=O)-R^{13}$, for example $-C(=O)$-(1-pyrrolidinyl),

C_{1-4}alkyl substituted with NR^7R^8, for example $-CH_2N(CH_2CH_3)_2$, $-CH_2N(CH_3)_2$ or $-CH_2N(CH_2CH_3)(CH_3)$,

C_{1-4}alkoxy substituted with NR^7R^8, for example $-OCH_2CH_2NH_2$,

$-NR^7R^8$, for example $-NHCH_3$ or $-N(CH_3)_2$,

$-C(=O)-NR^7R^8$; for example $-C(=O)-NHCH_3$, or

two R^2 groups are attached to adjacent carbon atoms and together to form a radical of formula $-O-(C(R^{17})_2)_p-O-$ wherein R^{17} represents hydrogen and p represents 1;

R^3 represents

C_{1-6}alkyl, for example $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$ or $-CH_2CH(CH_3)_2$,

hydroxyC_{1-6}alkyl, for example $-CH_2CH_2OH$, $-CH_2CH_2CH_2OH$, $-CH_2CHOHCH_3$, $-CH_2CHOHCH_2CH_3$, $-CH_2CHOHCH(CH_3)_2$, $-CH_2CH_2C(OH)(CH_3)_2$, $-CH_2CHOHCH_2OH$, $-CH_2C(CH_3)_2OH$, $-CD_2CD_2OH$, $-CD_2CD_2CD_2OH$, or $-CH(CH_3)CH_2OH$,

hydroxyhaloC_{1-6}alkyl, for example $-CH_2CHOHCF_3$,

haloC_{1-6}alkyl, for example $-CH_2CH_2CH_2Cl$, $-CH_2CH_2CH_2CH_2Cl$, $-CH_2CH_2F$ or $-CH_2CH_2I$,

haloC_{1-6}alkyl optionally substituted with –O-C(=O)-C_{1-6}alkyl, for example $-CH_2CH(CF_3)$-O-C(=O)CH_3, hydroxyC_{2-6}alkynyl, for example–CH_2 —C≡C— CH_2OH or -CH_2 —C≡C— $C(CH_3)_2OH$,

C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, for example CH_3-C(=O)–CH_2-, $(CH_3)_2CH$-C(=O)-CH_2-,

C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxy groups, for example $-CH_2CH_2OCH_3$, $-CH_2CH_2OCH_2CH_3$ or $-CH_2CHOHCH_2OCH_3$,

C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups or with –O-C(=O)-C_{1-6}alkyl, for example $-CH_2CH(-O-C(=O)CH_3)CH_2OCH_3$,

C_{1-6}alkyl substituted with R^9, for example

$-CH_2$-C_3H_5 or $-CH_2C_5H_9$,

C_{1-6}alkyl substituted with cyclopropyl substituted by $-CH_2OH$, CH_3CH_2-O-C(=O)-4-pyridinyl,

methyl substituted with 5-isoxazolyl which is substituted in the 3 position with $-CH_3$. or substituted with 3-isoxazolyl which is substituted in the 5 position by $-CH_3$,

ethyl or propyl substituted by 4-morpholinyl, methyl substituted by 3-morpholinyl, methyl substituted by 6-morpholinyl,

ethyl or propyl substituted by 4-morpholinyl which is substituted in the 2 and 6 positions by $-CH_3$,

methyl substituted by 2-morpholinyl which is substituted in the 4 position by -CH_2-C_6H_5 methyl substituted by 3-morpholinyl substituted in the 5 position by two $-CH_3$, methyl substituted by 6-morpholinyl substituted in the 4 position by $-CH(CH_3)_2$,methyl substituted by 6-morpholinyl substituted in the 3 position by =O and 4 position by $-CH(CH_3)_2$, methyl substituted by 2-morpholinyl substituted in the 4 positon by -CH_2-C_6H_5, methyl substituted with 2-tetrahydrofuranyl, 2-dioxolane, ethylene oxide, 2-furanyl, or 4-tetrahydropyranyl,

methyl substituted with 3-oxetanyl which is substituted in the 3 position by $-CH_3$, methyl substituted with 3-oxetanyl substituted in the 3 position by $-CH_2NHCH(CH_3)_2$,

methyl substituted with 3-pyridinyl or 2-pyrazinyl or propyl substituted with 4-pyridinyl

methyl or propyl substituted with 2-pyrimidinyl,

methyl substituted with 3-pyridinyl which is substituted in the 6 position by chlorine or methyl substituted with 2-pyridinyl which is substituted in the 6 position by bromine,

propyl substituted with 6-pyridinyl substituted in the 4 position by $-CH_3$, propyl substituted with 6-pyridinyl substituted in the 3 position by $-OCH_3$, methyl substituted with 2-pyridinyl substituted in the 6 position by $-OCH_3$, methyl substituted with 6-pyridinyl substituted in the 2 position by $-CH_2NH_2$, methyl substituted with 6-pyridinyl substituted in the 2 position by $-NHCH_3$,

propyl substituted with 2-pyrimidinyl substituted in the 4 position by $-OCH_3$, methyl substituted with 2-pyrimidinyl substituted in the 4 and 6 positions by $-OCH_3$, propyl substituted with 2-pyrimidinyl substituted in the 4 position by –OH,

methyl substituted with 3-piperazinyl,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by 4-piperidinyl being substituted in the 1 position by $-C(=O)-CH_3$, ethyl substituted with 1-piperazinyl substituted in the 4 position with $-CH_2C(=O)NHCH(CH_3)_2$,

ethyl or propyl substituted with 1, 2, 3, 6-tetrahydropyridine,

C_{1-6}alkyl substituted with azetidinyl,

propyl substituted by 1-azetidinyl which is substituted in the 3 position by two fluorines,

propyl substituted by 1-azetidinyl which is substituted in the 3 position by one -OH,

ethyl or propyl substituted with 1-pyrrolidinyl or 2-pyrrolidinyl,

propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by two fluorines or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by one fluorine,

propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by $-CH_2Cl$,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by –OH,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by =O,

propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by $-S(=O)_2-CH_3$,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position with $-NH_2$,

ethyl substituted with 1-pyrrolidinyl which is substituted in the 3 position with –$N(CH_3)_2$, propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position with –$NHCH_3$,

ethyl or propyl substituted with a) 1-pyrrolidinyl which is substituted in the 2 position with –CH_3; b) 1-pyrrolidinyl which is substituted in the 2 and the 5 position with –CH_3; or c) 1-pyrrolidinyl which is substituted in the 2 position with two –CH_3,

ethyl substituted with 1-pyrrolidinyl which is substituted in the 2 position with –C(=O)OH,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by –CH_2OH or with pyrrolidinyl which is substituted with –$C(CH_3)_2OH$ or –CH_2CH_2OH,

propyl substituted with a) 1-pyrrolidinyl which is substituted in the 3 position by 1-piperidinyl, or b) 1-pyrrolidinyl which is substituted in the 3 position by 4-morpholinyl being substituted in positions 2 and 6 by –CH_3,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by –CN,

propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by –CH_2CN,or ethyl substituted with 1-pyrrolidinyl substituted in the 2 position by –CH_2CN,

propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by –CH_2NH-$S(=O)_2$-CF_3,

methyl or ethyl substituted by a) 2-pyrrolidinyl which is substituted in the 1 position by $(CH_3)_3$C-O-C(=O)- or b) 1-pyrrolidinyl which is substituted in the 2 position by CH_3-O-C(=O)-, methyl substituted by 3-pyrrolidinyl substituted in the 1-position by 2-pyridinyl substituted in the 3-position by –OCH_3 or methyl substituted by 3-pyrrolidinyl substituted in the 1-position by 2-pyrimidinyl substituted in the 4-position by –OCH_3,

methyl, ethyl or propyl substituted by 4-piperidinyl or 1-piperidinyl,

ethyl substituted by 1-piperidinyl which is substituted at the 4 position by two fluorines,

methyl or ethyl substituted by a) 1-piperidinyl which is substituted at the 4 position by one –OH or b) 4-piperidinyl which is substituted at the 4 position by one –OH,

ethyl substituted by 1-piperidinyl which is substituted at the 3 position or the 4 position by –NH_2,

ethyl substituted by 1-piperidinyl which is substituted at the 4 position by –$N(CH_3)_2$,

methyl, ethyl or propyl substituted by a) 1-piperidinyl which is substituted at the 2 position by –CH_3, b) 1-piperidinyl which is substituted at the 2 and the 6 position by –

CH_3, c) 4-piperidinyl which is substituted at the 1 position by $-CH(CH_3)_2$, d) 4-piperidinyl which is substituted at the 1 position by $-CH_3$, e) 1-piperidinyl which is substituted at the 3 and the 5 position by $-CH_3$,

ethyl substituted by a) 1-piperidinyl which is substituted in the 4 position by $-C(CH_3)_2OH$, b) 1-piperidinyl which is substituted in the 4 position by $-CH_2CH_2OH$, c) 1-piperidinyl which is substituted in the 4 position by $-CH_2OH$,

ethyl or propyl substituted with 1-piperidinyl which is substituted at the 3 position with –CN,

methyl or ethyl substituted with a) 1-piperidinyl which is substituted in the 4 position by CH_3CH_2-O-C(=O)-, or b) 4-piperidinyl which is substituted in the 1 position by $(CH_3)_3$C-O-C(=O)-,

methyl substituted with 4-piperidinyl which is substituted in the 4 position by –OH and in the 1 position by $(CH_3)_3$C-O-C(=O)-,

methyl substituted with 4-piperidinyl which is substituted in the 4 position by $-OCH_3$ and in the 1 position by $(CH_3)_3$C-O-C(=O)- ,

methyl or ethyl substituted with a) 1-piperidinyl which is substituted in the 4 position by $-OCH_3$ or b) 4-piperidinyl which is substituted in the 4 position by $-OCH_3$,

propyl substituted with 1-piperidinyl which is substituted in the 4 position by $-CF_3$,

ethyl substituted with1-piperidinyl which is substituted in the 3 position by -C(=O)-NH_2, ethyl or propyl substituted with 1-piperidinyl substituted in the 2 position by -C(=O)-NH_2,ethyl substituted by 1-piperidinyl substituted at the 4 position by =O, or propyl substituted by 1-piperidinyl substituted at the 2 position by =O,

ethyl substituted with 1-piperidinyl substituted in the 4 position by $-CH_2NH_2$,

methyl substituted by 4-piperidinyl substituted in the 1-position by 2-pyrimidinyl substituted in the 4-position by $-OCH_3$,

ethyl, propyl or butyl substituted with isoindole-1, 3-dione, $-CH(CH_3)CH_2$-substituted with isoindolyl-1,3,-dione,

ethyl substituted with 2-oxa-6-aza-spiro[3.3.]heptane,

ethyl substituted with 1, 4-dioxa-8-aza-spiro[4.5]decane,

methyl substituted with 2-thiophenyl,

methyl substituted with 2-thiophenyl which is substituted at the 5 position by chlorine,

methyl substituted with 4-thiazolyl which is substituted in the 2 position by $-CH_3$,

ethyl or propyl substituted with 1-piperazinyl,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by CH_3-C(=O)-,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by –CH_2CH_2OH,

ethyl or propyl substituted with a) 1-piperazinyl which is substituted at the 3 and 5 positions by –CH_3 or b) 1-piperazinyl which is substituted at the 4 position by –CH_3,

ethyl substituted with 1-piperazinyl which is substituted in the 3 position by =O,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by -C(=O)-C_3H_5,

methyl substituted with 2-piperazinyl substituted in the 1 and 4 position by methylphenyl wherein the phenyl is substituted in the 4 position by CH_3O-, ethyl substituted with 5-tetrazolyl,

methyl substituted with a) 2-(1, 3, 4-oxadiazoyl) which is substituted at the 5 position by –NH_2 or b) 2-(1, 3, 4-oxadiazolyl) which is substituted at the 5 position by -NH–CH_2CH_2OH,

methyl, ethyl or propyl substituted with 1-pyrazolyl or 2-imidazolyl,methyl substituted with 3-pyrazolyl or 5-pyrazolyl,

methyl, ethyl or propyl substituted with a) 1-imidazolyl which is substituted at the 2 position by –CH_3, b) 3-pyrazolyl which is substituted at the 1 and 5 positions by –CH_3, c) 1-imidazolyl which is substituted at the 2 and 5 positions by –CH_3, d) 1-imidazolyl which is substituted at the 2 and 4 positions by –CH_3, e) 2-imidazolyl which is substituted at the 1 position by –CH_3 or f) 2-imidazolyl which is substituted at the 1 position by –CH_2CH_3,methyl substituted with 2-imidazolyl substituted at the 5 position by –CH_3 ethyl substituted with 1-pyrazolyl substituted at the 3 position by –CH_3

methyl substituted with 4-pyrazolyl substituted at the 1 position by –CH_3methyl substituted with 2-imidazolyl substituted in the 3 position by –$S(=O)_2$-$N(CH_3)_2$ and in the 5 position by –CH_3, methyl substituted with 5-pyrazolyl substituted in the 2 position by 2-tetrahydropyran, or methyl substituted with 3-pyrazolyl substituted in the 1 position by 2-tetrahydropyran methyl substituted with 2-imidazolyl which is substituted in the 1 position by –$S(=O)_2$-$N(CH_3)_2$,

methyl substituted with 4-(1, 2, 3-triazolyl),

methyl substituted with a) 4-(1, 2, 3-triazolyl) which is substituted in the 1 position by –CH_2CH_2OH or b) 4-(1, 2, 3-triazolyl) which is substituted in the 2 position by –CH_2OH,

methyl substituted with 4-(1, 2, 3-triazolyl) which is substituted in the 1 position by $-CH_2C(=O)-OCH_2CH_3$,

ethyl substituted with 1-(1, 2, 4-triazolyl,

ethyl or propyl substituted with 1-(1, 2, 4-triazolyl) substituted in the 3 position by $-CH_3$,

ethyl or propyl substituted with 2-(1, 2, 4-triazolyl) substituted in the 3 position by $-CH_3$,

ethyl or propyl substituted with 3-oxazolidinyl which is substituted in the 2 position by =O,methyl substituted with 5-oxazolidinyl substituted in the 2 position by =O,methyl substituted with 5-oxazolidinyl substituted in the 2 position by =O and in the 3 position by $-CH(CH_3)_2$,

propyl substituted with 4-thiomorpholinyl which is substituted in the 1 position by two =O groups,

ethyl substituted with 1- homopiperazinyl,

ethyl substituted with homomorpholinyl,

$-CH_2-C_6H_5$,

methyl substituted with phenyl which is substituted in the 2, 3 or 4 position by chlorine,cyanoC_{1-6}alkyl, for example $-CH_2CH_2CN$ or $-CH_2CH_2CH_2CN$,

C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example $-CH_2CH_2NH_2$, $-CH_2CH_2CH_2NH_2$, $-CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH(CH_3)NH_2$, $-CH(CH_3)CH_2NH_2$, $-CH_2CH_2NHCH_3$, $-CH_2CH_2CH_2NHCH_3$, $-CH_2CH_2NHCH_2CH_3$, $-CH_2CH_2NHCH(CH_3)_2$, $-CD_2-CD_2-NHCH(CH_3)_2$, $-CH_2CH_2CH_2NHCH(CH_3)_2$, $-CH(CH_3)CH_2NHCH(CH_3)_2$, $-CH_2CH_2N(CH_2CH_3)_2$, $-CH_2CH_2N(CH_2CH_3)(CH(CH_3)_2)$, $-CH_2CH_2N(CH_3)_2$ or $-CH_2CH_2N(CH_3)CH(CH_3)_2$, $-CH_2CH_2CH_2NHCH_2CF_3$, $-CH_2CH_2NHCH_2CHF_2$ or $-CH_2CH_2NHCH_2CH_2F$, $-CH(CH_3)CH_2NHCH_2CF_3$, $-CH_2CH(CH_3)NHCH_2CF_3$, $-CH_2CH_2NHCH_2CF_3$, $-CH_2CH_2CH_2NHCH_2CHF_2$ $-CH_2CH_2NHCH_2CH_2CF_3$, $-CH_2CH_2CH_2NHCH_2CHF_2$, $-CH_2CH_2CH_2NHC(CH_3)_2CH_2F$, $-CD_2-CD_2-CD_2-NHCH_2CF_3$, $-CH_2CH_2NH-C(=O)-CH_3$, $-CH_2CH_2NH-S(=O)_2-CH_3$, $-CH_2CH_2CH_2NH-S(=O)_2-CH_3$, $-CH_2CH_2NH-S(=O)_2-CH_2CH_3$ or $-CH_2CH_2NH-S(=O)_2-CH(CH_3)_2$, $-CH_2CH_2NH-S(=O)_2-N(CH_3)_2$ or $-CH_2CH_2CH_2NH-S(=O)_2-N(CH_3)_2$, $-CH_2CH_2NHCH_2CH_2OH$, $-CH_2CH_2CH_2NH-C(=O)-C(OH)(CH_3)CF_3$ or $-CH_2CH_2NH-C(=O)-C(OH)(CH_3)CF_3$,

$-CH_2CH_2NH-C(=O)-C_3H_5$,

$-CH_2CH_2NH-C(=O)$-(piperidin-3-yl) where the piperidinyl is substituted at the 1 position by $-CH_3$,

$-CH_2CH_2NHCH_2CH_2CN$, $-CH_2CH_2CH_2NHCH_2CH_2CN$,

$-CH_2CH_2NHC_3H_5$, $-CH_2CH_2NHC_5H_9$ or $-CH_2CH_2NH$-(2,2,6,6-tetramethyl-piperidin-4-yl)

$-CH_2CH_2NH$-(piperidin-4-yl) where the piperidinyl is substituted in the 1 position by $-S(=O)_2NH_2$,

$-CH_2CH_2NHCH_2C_3H_5$, $-CH_2CH_2NHCH_2$-(tetrahydrofuran-2-yl), $-CH_2CH_2NHCH_2$-(pyridin-6-yl),

$-CH_2CH_2NHC(=O)-CF_3$ or $-CH_2CH_2CH_2NHC(=O)-CF_3$;

$-CH_2CH_2NHCH_2Si(CH_3)_3$,

$-CH_2CH_2N(CH_3)CH_2-C_6H_5$.,

one of R^{10} and R^{11} represents $-CH(CH_3)_2$ and the other represents $-CH_2-C_6H_5$ wherein the phenyl is substituted in the 4-position by $-NH_2$,

$-CH_2CH_2N(CH(CH_3)_2)CH_2CH_2CH_2NH_2$,

$-CH_2CH_2CH_2NHCH_2C(=O)NH_2$ or $-CH_2CH_2NHCH_2C(=O)NH_2$,

$-CH_2CH_2CH_2N(CH_3)-OCH_3$,

$-CH_2CH_2NH-OCH_3$, or

$-CH_2CH_2NHCH_2CHOHCF_3$;

$-CH_2CH_2CH_2NHCOOH$.

C_{1-6}alkyl substituted with hydroxyl and $-NR^{10}R^{11}$, for example $-CH_2CHOHCH_2NH_2$, $-CH_2CHOHCH_2NHCH_3$ or $-CH_2CHOHCH_2NHCH(CH_3)_2$, $-CH_2CHOHCH_2NHCH_2CF_3$, $-CH_2CHOHCH_2N(CH(CH_3)_2)-C(=O)CH_2Cl$,

C_{1-6}alkyl substituted with one or two halogens and $-NR^{10}R^{11}$, for example $-CH_2CHFCH_2NH_2$,

C_{1-6}alkyl substituted with $-C(=O)-O-C_{1-6}$alkyl, for example $-CH_2-C(=O)-O-CH_2CH_3$ or $-CH_2CH_2-C(=O)-O-CH_2CH_3$, $-CH(CH_3)C(=O)-O-CH_2CH_3$,

C_{1-6}alkyl (for example methyl) substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–, for example $-CH_2-C(=O)-CH_2OCH_3$,

C_{1-6}alkyl substituted with $-O-C(=O)-NR^{10}R^{11}$, for example $-CH_2-C(=O)NH_2$, $-CH_2-C(=O)NHCH_3$, $-CH_2C(=O)NHCH(CH_3)_2$ or $-CH_2CH_2C(=O)NHCH(CH_3)_2$, $-CH_2-C(=O)-NHCH_2CH_2OCH_3$, $-CH_2-C(=O)-NH-CH_2CH_2$-(pyrrolidin-1-yl) or $-CH_2-C(=O)-NH-CH_2CH_2$-(imidazol-2-yl), $-CH_2-C(=O)-NHCH_2CH_2OH$, $-CH_2-C(=O)-NHCH_2CH_2NH_2$, - $CH_2CH_2C(=O)-NHCH_2CF_3$ $-CH_2CH_2C(=O)N(CH_3)-OCH_3$, -

$CH_2C(=O)NH$-(pyridin-2-yl) wherein the pyridin-2-yl is substituted in the 3-position by $-OCH_3$, -$CH_2C(=O)NH$-(pyridin-6-yl) wherein the pyridin-6-yl is substituted in the 4-position by $-CH_3$ or -$CH_2C(=O)NH$-(pyrimidin-2-yl) wherein the pyrimidin-2-yl is substituted in the 4-position by $-OCH_3$, -$CH_2C(=O)NH$-(pyridin-3-yl), -$CH_2C(=O)NH$-(pyridin-6-yl) or -$CH_2C(=O)NH$-(pyridin-4-yl),

C_{1-6}alkyl substituted with carboxyl, for example $-CH_2C(=O)OH$ or $-CH_2CH_2C(=O)OH$,

C_{1-6}alkyl substituted with $-O-C(=O)-NR^{10}R^{11}$, for example -$CH_2CH_2-O-C(=O)-NHCH_3$,

C_{1-6}alkyl substituted with $-NR^{12}-S(=O)_2-C_{1-6}$alkyl, for example $-CH_2CH_2NH-S(=O)_2-CH_3$, $-CH_2CH_2CH_2NH-S(=O)_2-CH_3$, -$CH_2CH_2NH-S(=O)_2-CH(CH_3)_2$ or $-CH_2CH_2NH-S(=O)_2-CH_2CH_3$,

C_{1-6}alkyl substituted with $-NR^{12}-S(=O)_2-NR^{14}R^{15}$, for example $-CH_2CH_2NH-S(=O)_2-N(CH_3)_2$ or $-CH_2CH_2CH_2NH-S(=O)_2-N(CH_3)_2$,

C_{1-6}alkyl substituted with R^9 and optionally substituted with $-O-C(=O)-C_{1-6}$alkyl, R^9 represents 1H-pyrrolo[3,2-b]pyridinyl, 1-methyl-1H-pyrrolo[3,2-b]pyridinyl or furo[3,2-b]pyridinyl,

C_{1-6}alkyl substituted with hydroxyl and R^9, for example

propyl substituted with –OH and 1-pyrrolidinyl,

propyl substituted with –OH and 1-pyrrolidinyl where the 1-pyrrolidinyl is substituted at the 3 position by two fluorines,

propyl substituted with –OH and 1-pyrrolidinyl where the 1-pyrrolidinyl is substituted at the 3 position by a cyano group,

propyl substituted with –OH and 4-morpholinyl,

propyl substituted with –OH and 1-piperidinyl,

propyl substituted with –OH and 2-(1, 2, 4-triazolyl) substituted in the 3 position by $-CH_3$,

propyl substituted with –OH and 1-imidazolyl substituted in the 2 position by $-CH_3$,

propyl substituted with –OH and isoindole-1, 3-dione,

-C_{1-6}alkyl-$C(R^{12})=N-O-R^{12}$, for example $-CH_2C(CH_3)=N-O-H$, $-CH_2C(CH_2OCH_3)=N-O-H$ or $-CH_2C(CH(CH_3)_2)=N-O-H-S(=O)_2-NR^{14}R^{15}$, for example $-S(=O)_2-N(CH_3)_2$,C_{1-6}alkyl substituted with -$S(=O)_2-C_{1-6}$alkyl, for example $-CH_2CH_2-S(=O)_2-CH_3$,

C_{1-6}alkyl substituted with $-C(=O)-NR^{10}R^{11}$, for example

$-CH_2C(=O)NH_2$,

$-CH_2C(=O)NHCH_3$,

$-CH_2C(=O)-NHCH_2CH_2OCH_3$,

$-CH_2C(=O)$-NH-CH_2CH_2-(pyrrolidin-1-yl) or $-CH_2C(=O)$-NH-CH_2CH_2-(imidazol-2-yl),$-CH_2C(=O)-NHCH_2CH_2OH$,$-CH_2C(=O)-NHCH_2CH_2NH_2$,

C_{1-6}alkyl substituted with $-C(=O)-R^9$, for example $-CH_2C(=O)-R^9$ and R^9 is 1-pyrrolidinyl,

C_{2-6}alkenyl substituted with R^9, for example $-CH_2CH=CH$-(2-pyrimidinyl), $-CH_2CH=CH$-(2-pyrimidinyl) wherein the 2-pyrimidinyl is substituted in the 4-position by $-OCH_3$, $-CH_2CH=CH$-(2-pyridinyl) wherein the 2-pyridinyl is substituted in the 4-position by $-CH_3$ or $-CH_2CH=CH$-(2-pyridinyl) wherein the 2-pyridinyl is substituted in the 3-position by $-OCH_3$,

C_{2-6}alkynyl substituted with R^9, for example

$-CH_2-C\equiv C-$(2-imidazolyl) wherein the 2-imidazolyl is substituted in the 1 position by $-CH_3$ or $-CH_2-C\equiv C-$(5-imidazolyl) wherein the 5-imidazolyl is substituted in the 1 position by $-CH_3$,$-CH_2-C\equiv C-$ (4-pyridinyl), $-CH_2-C\equiv C-$ (3-pyridinyl), $-CH_2-C\equiv C-$ (2-pyridinyl), $-CH_2-C\equiv C-$ (2-pyrimidinyl), $-CH_2-C\equiv C-$ (6-pyrazinyl),-$CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2 or 4-position with $-CH_2OH$,

$-CH_2-C\equiv C-$ (4-pyridinyl) substituted in the 6-position with $-OCH_3$,$-CH_2-C\equiv C-$ (2-pyridinyl) substituted in the 3 or 5-position with $-OCH_3$,

$-CH_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 4 or 6-position with $-OCH_3$,

$-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2, 4 or 5-position with $-OCH_3$,

$-CH_2-C\equiv C-$ (6-pyrimidinyl) substituted in the 4 -position with $-OCH_3$,

$-CH_2-C\equiv C-$ (5-pyrazinyl) substituted in the 6 -position with $-OCH_3$,

$-CH_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 6-position with $-OCH_2CH_3$,

$-C(CH_3)_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 4-position with $-OCH_3$,

$-CH_2-C\equiv C-$ (2-pyrimidinyl) substituted in the 4-position with $-OCH(CH_3)_2$;

$-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2 or the 4-position with cyano, -$CH_2-C\equiv C-$ (4-pyridinyl) substituted in the 5 or 6-position with cyano;

$-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 2 or 4-position with $-NH_2$, -$CH_2-C\equiv C-$ (6-pyrimidinyl) substituted in the 2-position with $-NH_2$, $-CH_2-C\equiv C-$ (2-pyridinyl) substituted in the 3-position with $-NH_2$, $-CH_2-C\equiv C-$ (3-pyrazinyl) substituted in the 6-position with $-NH_2$, $-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 5-position with $-NHCH_3$,

$-CH_2-C\equiv C-$ (6-pyridinyl) substituted in the 3 or 4-position with $-CH_3$, -$CH_2-C\equiv C-$ (2-pyridinyl) substituted in the 3-position with $-CH_3$, $-CH_2-C\equiv C-$ (2-

pyrimidinyl) substituted in the 4-position with –CH_3, -CH_2 —C≡C— (2-pyrimidinyl) substituted in the 6-position with –CH_2CH_3,

-CH_2 —C≡C— (6-pyrimidinyl) substituted in the 2-position with –CH_3 and in the 4-position with –NH_2,

-CH_2 —C≡C— (6-pyrimidinyl) substituted in the 2-position with –NH_2 and in the 4-position with –Cl,

-CH_2 —C≡C— (2-pyrazinyl) substituted in the 3-position with –Cl, -CH_2 —C≡C— (3-pyrazinyl) substituted in the 5-position with –Cl, -CH_2 —C≡C— (2-pyridinyl) substituted in the 3-position with –F, -CH_2 —C≡C— (5-pyridinyl) substituted in the 6-position with –Br;

-CH_2 —C≡C— (6-pyridinyl) substituted in the 4-position with –C(=O)-NH_2;

-CH_2 —C≡C— (6-pyridinyl) substituted in the 5-position with CH_3-O-C(=O)-, -CH_2 —C≡C— (2-pyrimidinyl) substituted in the 6-position with CH_3-O-C(=O)-;

-CH_2 —C≡C— (2-pyridinyl) substituted in the 3-position with – CF_3,

-CH_2 —C≡C— (5-thiazolyl),

-CH_2 —C≡C— (phenyl),

-CH_2 —C≡C— (phenyl) where the phenyl is substituted in the 5-position by -OCH_3,

-CH_2 —C≡C— (3-azetidinyl) substituted in the 1-position by $C(CH_3)_3$-O-C(=O)- and in the 3-position by –OH,

-CH_2 —C≡C— (3-azetidinyl) substituted in the 3-position by –OH,

-CH_2 —C≡C— (3-pyrrolidinyl) substituted in the 1-position by $C(CH_3)_3$-O-C(=O)- and in the 3-position by –OH,

-CH_2 —C≡C— (3-pyrrolidinyl) substituted in the 3-position by –OH,

-CH_2 —C≡C— (4-piperidinyl),

-CH_2 —C≡C— (4-piperidinyl) substituted in the 4-position by –OH,

-CH_2 —C≡C— (4-piperidinyl) substituted in the 1-position by $C(CH_3)_3$-O-C(=O)-,

-CH_2 —C≡C— (4-tetrahydrofuranyl) substituted in the 3-position by –OH,

-CH_2 —C≡C— (4-tetrahydropyranyl) substituted in the 4-position by –OH,

-CH_2 —C≡C— (cyclohexyl),

C_{1-6}alkyl substituted with R^9, R^9 represents a 6 to 8 membered bridged heterocyclyl group, for example 2,5-diaza-bicyclo[2.2.1]heptanyl optionally substituted by –C(=O)-O-C_4alkyl, C_{1-6}alkyloxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, for example –$CH_2CHOHCH_2OCH_3$,

C_{2-6}alkenyl, for example $-CH_2-CH=CH_2$,

C_{2-6}alkynyl, for example $-CH_2-C\equiv C-H$ or $-C(CH_3)_2-C\equiv C-H$.,


C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–, for example $-CH_2-C(=O)-CH_2OCH_3$, or

R^{13}, for example 3–oxetanyl, , cyclohexanyl substituted in the 4 position by -NH–$CH(CH_3)_2$,

C_{1-6}alkyl substituted by R^9, wherein R^9 is a saturated heterocyclyl substituted by R^{13}, wherein R^{13} is a saturated heterocyclyl which is optionally substituted, for example substituted by $-C(=O)-C_{1-6}$alkyl. In one embodiment R^9 is piperazinyl substituted by R^{13}, wherein R^{13} is piperidinyl substituted by $-C(=O)-C_{1-6}$alkyl,

C_{1-6}alkyl substituted by R^9, wherein R^9 is a saturated heterocyclyl substituted by R^{13}, wherein R^{13} is a saturated heterocyclyl which is optionally substituted, for example substituted by $-C(=O)-C_{1-6}$alkyl. In one embodiment R^9 is piperazinyl substituted by R^{13}, wherein R^{13} is piperidinyl substituted by $-C(=O)-C_{1-6}$alkyl.

In one embodiment there is provided a compound of formula (I^0):



(I^0)

including any stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2,or 3;

R^1 represents hydrogen,

C_{1-6}alkyl, for example $-CH_3$, $-CD_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)_2$, $-CH_2CH(CH_3)_2$,

C_{2-4}alkenyl, for example $-CH_2-CH=CH_2$,

hydroxyC_{1-6}alkyl, for example $-CH_2CH_2OH$, $-CH_2C(CH_3)_2OH$ or $CH_2CHOHCH_2OH$,

haloC_{1-6}alkyl, for example $-CH_2CH_2F$, $CH_2CH_2CH_2Cl$ or CH_2CH_2Br,

C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, for example $-CH_2CH_2OCH_3$,

C_{1-6}alkyl substituted with -NR^4R^5, for example -$CH_2CH_2NH_2$ or -$CH_2CH_2CH_2NH_2$, -$CH_2CH_2NHCH_3$, -$CH_2CH_2NHS(=O)_2N(CH_3)_2$, -$CH_2CH_2NHS(=O)_2N(CH_3)_2$,

C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, for example -$CH_2C(=O)N(CH_3)_2$, -$CH_2C(=O)NHCH_3$ or -$C(CH_3)_2C(=O)NHCH_3$ -$C(CH_3)_2C(=O)NHCH_2CH_2OH$ or -$CH_2C(=O)NHCH_2CH_2OH$, -$CH_2C(=O)NHCH_2CH_2OCH_3$ or -$C(CH_3)_2C(=O)NHCH_2CH_2OCH_3$, –$CH_2$-C(=O)-NH-$CH_2$-$CH_2$-(pyrrolidin-1-yl), -$CH_2CH_2CH_2NHCH_2CH_2$-$S(=O)_2$-$CH_3$,

–$S(=O)_2$-C_{1-6}alkyl, for example –$S(=O)_2$-CH_3,

–$S(=O)_2$-$NR^{14}R^{15}$, for example –$S(=O)_2$-$N(CH_3)_2$,

C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, for example -$CH_2CH_2S(=O)_2$-CH_3,

C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, for example -$CH_2CH_2NHS(=O)_2$-CH_3,

R^6, for example 2-tetrahydropyranyl, 3-azetidinyl substituted in the 1 position by –CH_2CH_2OH, 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)-, 4-piperidinyl substituted on the nitrogen atom with –$S(=O)_2CH_3$,

C_{1-6}alkyl substituted with R^6, for example methyl or ethyl each substituted with 4-piperidinyl, 4-piperazinyl, 1-pyrrolidinyl or 4-tetrahydropyranyl; propyl substituted with morpholinyl where the morpholinyl is linked to the propyl through the N heteroatom; methyl, ethyl or propyl each substituted with 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)-, 4-piperidinyl substituted on the nitrogen atom with –CH_3, 4-piperazinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)-, 4-piperazinyl substituted on the nitrogen atom with -CH_2CH_2OH, 4-piperazinyl substituted on the nitrogen atom with -$CH_2CH_2CH_2OH$, 1-piperidinyl substituted in the 1 position by –OH, 1-piperidinyl substituted in the 1 position by –O-CH_3; methyl substituted with 2-thiophenyl substituted in the 5 position with chlorine; methyl substituted with 4-piperidinyl substituted on the nitrogen atom with $(CH_3)_3$C-O-C(=O)- and in the 4 position by –OH,

C_{1-6}alkyl substituted with –C(=O)-R^6, for example –$C(CH_3)_2$-C(=O)-(piperazin-4-yl), –$C(CH_3)_2$-C(=O)-(piperazin-4-yl) substituted on the nitrogen atom in the 1 position by $C(CH_3)_3$-O-C(=O)-, –CH_2-C(=O)-(pyrrolidin-1-yl) substituted in the 3 position by –OH,

hydroxyC_{1-6}alkyl substituted with R^6, for example –$CH_2CHOHCH_2$- substituted with 1-piperidinyl; or

C_{1-6}alkyl substituted with –$Si(CH_3)_3$, for example -$CH_2Si(CH_3)_3$;

each R^2 is independently selected from

halogen, for example fluorine, chlorine or bromine,

cyano,

C_{1-4}alkyl, for example $-CH_3$,

C_{2-4}alkenyl, for example -CH=CH_2,

C_{1-4}alkoxy, for example CH_3O-, $(CH_3)_2CHO$-, CH_3CH_2O-, CD_3O-,

hydroxyC_{1-4}alkyl, for example -CH_2OH,

hydroxyC_{1-4}alkoxy, for example $-OCH_2CH_2OH$,

haloC_{1-4}alkoxy, for example -OCH_2CH_2F or CHF_2O-,

C_{1-4}alkoxyC_{1-4}alkyl, for example -$CH_2CH_2OCH_3$,

R^{13}, for example 2-dioxolanyl,

C_{1-4}alkoxy substituted with R^{13}, for example $-OCH_2C_3H_5$,

-C(=O)-R^{13}, for example –C(=O)-(1-pyrrolidinyl),

C_{1-4}alkyl substituted with NR^7R^8, for example -$CH_2N(CH_2CH_3)_2$, -$CH_2N(CH_3)_2$ or -$CH_2N(CH_2CH_3)(CH_3)$,

C_{1-4}alkoxy substituted with NR^7R^8, for example $-OCH_2CH_2NH_2$,

-NR^7R^8, for example $-NHCH_3$, or

-C(=O)-NR^7R^8; for example –C(=O)-$NHCH_3$;

R^3 represents

C_{1-6}alkyl, for example $-CH_3$, -CH_2CH_3, -$CH_2CH_2CH_3$ or $-CH_2CH(CH_3)_2$,

hydroxyC_{1-6}alkyl, for example $-CH_2CH_2OH$, $-CH_2CH_2CH_2OH$, $-CH_2CHOHCH_3$, $-CH_2CHOHCH_2CH_3$, $-CH_2CHOHCH(CH_3)_2$, -$CH_2CH_2C(OH)(CH_3)_2$, -$CH_2CHOHCH_2OH$ or $-CH_2C(CH_3)_2OH$,

hydroxyhaloC_{1-6}alkyl, for example $-CH_2CHOHCF_3$,

haloC_{1-6}alkyl, for example $-CH_2CH_2CH_2Cl$ or -$CH_2CH_2CH_2CH_2Cl$,

C_{1-6}alkyl substituted with –C(=O)-C_{1-6}alkyl, for example CH_3-C(=O)–CH_2-, $(CH_3)_2CH$-C(=O)-CH_2-,

C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxy groups, for example $-CH_2CH_2OCH_3$, $-CH_2CH_2OCH_2CH_3$ or $-CH_2CHOHCH_2OCH_3$,

C_{1-6}alkyl substituted with R^9, for example

$-CH_2$-C_3H_5 or $-CH_2C_5H_9$,

C_{1-6}alkyl substituted with cyclopropyl substituted by $-CH_2OH$ or CH_3CH_2-O-C(=O)-

methyl substituted with 5-isoxazoyl which is substituted in the 3 position with $-CH_3$. or substituted with 3-isoxazoyl which is substituted in the 5 position by $-CH_3$, 3

ethyl or propyl substituted by 4-morpholinyl

ethyl or propyl substituted by 4-morpholinyl which is substituted in the 2 and 6 positions by $-CH_3$

methyl substituted by 2-morpholinyl which is substituted in the 4 positon by $-CH_2-C_6H_5$

methyl substituted with 2-tetrahydrofuranyl, 2-dioxolane, ethylene oxide, 2-furanyl, or 4-tetrahydropyranyl,

methyl substituted with 3-oxetanyl which is substituted in the 3 position by $-CH_3$.

methyl substituted with 3-pyridinyl or 2-pyrazinyl.

methyl substituted with 3-pyridinyl which is substituted in the 6 position by chlorine or methyl substituted with 2-pyridinyl which is substituted in the 6 position by bromine,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by 4-piperidinyl being substituted in the 1 position by $-C(=O)-CH_3$,

ethyl or propyl substituted with 1, 2, 3, 6-tetrahydropyridine,

C_{1-6}alkyl substituted with azetidinyl,

propyl substituted by 1-azetidinyl which is substituted in the 3 position by two fluorines,propyl substituted by 1-azetidinyl which is substituted in the 3 position by one -OH,

ethyl or propyl substituted with 1-pyrrolidinyl or 2-pyrrolidinyl,

propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by two fluorines or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by one fluorine,

propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by $-CH_2Cl$,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by $-OH$,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by $=O$,

propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by $-S(=O)_2-CH_3$,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position with $-NH_2$,ethyl substituted with 1-pyrrolidinyl which is substituted in the 3 position with –

$N(CH_3)_2$, propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position with –$NHCH_3$,

ethyl or propyl substituted with a) 1-pyrrolidinyl which is substituted in the 2 position with –CH_3; b) 1-pyrrolidinyl which is substituted in the 2 and the 5 position with –CH_3; or c) 1-pyrrolidinyl which is substituted in the 2 position with two –CH_3,

ethyl substituted with 1-pyrrolidinyl which is substituted in the 2 position with –C(=O)OH,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by –CH_2OH or with pyrrolidinyl which is substituted with –$C(CH_3)_2OH$ or –CH_2CH_2OH,

propyl substituted with a) 1-pyrrolidinyl which is substituted in the 3 position by 1-piperidinyl, or b) 1-pyrrolidinyl which is substituted in the 3 position by 4-morpholinyl being substituted in positions 2 and 6 by –CH_3,

ethyl or propyl substituted with 1-pyrrolidinyl which is substituted in the 3 position by –CN, propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by –CH_2CN,

propyl substituted with 1-pyrrolidinyl which is substituted in the 2 position by –CH_2NH-$S(=O)_2$-CF_3,

methyl or ethyl substituted by a) 2-pyrrolidinyl which is substituted in the 1 position by $(CH_3)_3$C-O-C(=O)- or b) 1-pyrrolidinyl which is substituted in the 2 position by CH_3-O-C(=O)-,

methyl, ethyl or propyl substituted by 4-piperidinyl or 1-piperidinyl,

ethyl substituted by 1-piperidinyl which is substituted at the 4 position by two fluorines,

methyl or ethyl substituted by a) 1-piperidinyl which is substituted at the 4 position by one –OH or b) 4-piperidinyl which is substituted at the 4 position by one –OH,

ethyl substituted by 1-piperidinyl which is substituted at the 3 position or the 4 position by –NH_2,

ethyl substituted by 1-piperidinyl which is substituted at the 4 position by –$N(CH_3)_2$,

methyl, ethyl or propyl substituted by a) 1-piperidinyl which is substituted at the 2 position by –CH_3, b) 1-piperidinyl which is substituted at the 2 and the 6 position by –CH_3, c) 4-piperidinyl which is substituted at the 1 position by –$CH(CH_3)_2$, d) 4-piperidinyl which is substituted at the 1 position by –CH_3, e) 1-piperidinyl which is substituted at the 3 and the 5 position by –CH_3,

ethyl substituted by a) 1-piperidinyl which is substituted in the 4 position by –$C(CH_3)_2OH$, b) 1-piperidinyl which is substituted in the 4 position by –CH_2CH_2OH, c) 1-piperidinyl which is substituted in the 4 position by –CH_2OH,

ethyl or propyl substituted with 1-piperidinyl which is substituted at the 3 position with –CN,

methyl or ethyl substituted with a) 1-piperidinyl which is substituted in the 4 position by CH_3CH_2-O-C(=O)-, or b) 4-piperidinyl which is substituted in the 1 position by $(CH_3)_3$C-O-C(=O)-, ,

methyl substituted with 4-piperidinyl which is substituted in the 4 position by –OH and in the 1 position by $(CH_3)_3$C-O-C(=O)-,

methyl substituted with 4-piperidinyl which is substituted in the 4 position by –OCH_3 and in the 1 position by $(CH_3)_3$C-O-C(=O)- ,

methyl or ethyl substituted with a) 1-piperidinyl which is substituted in the 4 position by –OCH_3 or b) 4-piperidinyl which is substituted in the 4 position by –OCH_3,

propyl substituted with 1-piperidinyl which is substituted in the 4 position by –CF_3,

ethyl substituted with1-piperidinyl which is substituted in the 3 position by -C(=O)-NH_2,

ethyl, propyl or butyl substituted with isoindole-1, 3-dione,

ethyl substituted with 2-oxa-6-aza-spiro[3.3.]heptane,

ethyl substituted with 1, 4-dioxa-8-aza-spiro[4.5]decane,

methyl substituted with 2-thiophenyl,

methyl substituted with 2-thiophenyl which is substituted at the 5 position by chlorine,methyl substituted with 4-thiazolyl which is substituted in the 2 position by –CH_3,

ethyl or propyl substituted with 1-piperazinyl,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by CH_3-C(=O)-,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by –CH_2CH_2OH,

ethyl or propyl substituted with a) 1-piperazinyl which is substituted at the 3 and 5 positions by –CH_3 or b) 1-piperazinyl which is substituted at the 4 position by –CH_3,

ethyl substituted with 1-piperazinyl which is substituted in the 3 position by =O,

ethyl substituted with 1-piperazinyl which is substituted in the 4 position by -C(=O)-C_3H_5,

ethyl substituted with 5-tetrazolyl,

methyl substituted with a) 2-(1, 3, 4-oxadiazoyl) which is substituted at the 5 position by $-NH_2$ or b) 2-(1, 3, 4-oxadiazoyl) which is substituted at the 5 position by -NH–CH_2CH_2OH,

methyl, ethyl or propyl substituted with 1-pyrazoyl or 2-imidazoyl,

methyl, ethyl or propyl substituted with a) 1-imidazoyl which is substituted at the 2 position by $-CH_3$, b) 3-pyrazolyl which is substituted at the 1 and 5 positions by $-CH_3$, c) 1-imidazolyl which is substituted at the 2 and 5 positions by $-CH_3$, d) 1-imidazolyl which is substituted at the 2 and 4 positions by $-CH_3$, e) 2-imidazolyl which is substituted at the 1 position by $-CH_3$ or f) 2-imidazolyl which is substituted at the 1 position by $-CH_2CH_3$,methyl substituted with 2-imidazolyl which is substituted in the 1 position by $-S(=O)_2-N(CH_3)_2$,

methyl substituted with 4-(1, 2, 3-triazolyl),

methyl substituted with a) 4-(1, 2, 3-triazolyl) which is substituted in the 1 position by $-CH_2CH_2OH$ or b) 4-(1, 2, 3-triazolyl) which is substituted in the 2 position by $-CH_2OH$,

methyl substituted with 4-(1, 2, 3-triazolyl) which is substituted in the 1 position by $-CH_2C(=O)-OCH_2CH_3$,

ethyl or propyl substituted with 3-oxazolidinyl which is substituted in the 2 position by =O,

propyl substituted with 4-thiomorpholinyl which is substituted in the 1 position by two =O groups,

ethyl substituted with 1- homopiperazinyl,

$-CH_2-C_6H_5$,

methyl substituted with phenyl which is substituted in the 2, 3 or 4 position by chlorine,

C_{1-6}alkyl substituted with $-NR^{10}R^{11}$, for example $-CH_2CH_2NH_2$, $-CH_2CH_2CH_2NH_2$ or $-CH_2CH_2CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, $-CH_2CH_2CH_2NHCH_3$, $-CH_2CH_2NHCH_2CH_3$, $-CH_2CH_2NHCH(CH_3)_2$ or $-CH_2CH_2CH_2NHCH(CH_3)_2$, $-CH_2CH_2N(CH_2CH_3)_2$, $-CH_2CH_2N(CH_2CH_3)(CH(CH_3)_2)$, $-CH_2CH_2CH_2NHCH_2CF_3$, $-CH_2CH_2NHCH_2CHF_2$ or $-CH_2CH_2NHCH_2CH_2F$, $-CH_2CH_2NH-C(=O)-CH_3$, $-CH_2CH_2NH-S(=O)_2-CH_3$, $-CH_2CH_2CH_2NH-S(=O)_2-CH_3$, $-CH_2CH_2NH-S(=O)_2-CH_2CH_3$ or $-CH_2CH_2NH-S(=O)_2-CH(CH_3)_2$ $-CH_2CH_2NH-S(=O)_2-N(CH_3)_2$ or $-CH_2CH_2CH_2NH-S(=O)_2-N(CH_3)_2$ $-CH_2CH_2NHCH_2CH_2OH$, $-CH_2CH_2CH_2NH-C(=O)-C(OH)(CH_3)CF_3$ or $-CH_2CH_2NH-C(=O)-C(OH)(CH_3)CF_3$ $-CH_2CH_2NH-C(=O)-C_3H_5$ $-CH_2CH_2NH-C(=O)-$(piperidin-3-yl)

where the piperidinyl is substituted at the 1 position by $-CH_3$; $-CH_2CH_2NHCH_2CH_2CN$ $-CH_2CH_2NHC_3H_5$, $-CH_2CH_2NHC_5H_9$ or $-CH_2CH_2NH$-(2,2,6,6-tetramethyl-piperidin-4-yl), $-CH_2CH_2NHCH_2C_3H_5$, $-CH_2CH_2NHCH_2$-(tetrahydrofuran-2-yl), $-CH_2CH_2NHC(=O)-CF_3$ or $-CH_2CH_2CH_2NHC(=O)-CF_3$, $-CH_2CH_2NHCH_2Si(CH_3)_3$,$-CH_2CH_2N(CH_3)CH_2-C_6H_5$, $-CH_2CH_2NH$-(piperidin-4-yl) where the piperidinyl is substituted in the 1 position by $-S(=O)_2NH_2$,

C_{1-6}alkyl substituted with hydroxyl and $-NR^{10}R^{11}$, for example $-CH_2CHOHCH_2NH_2$, $-CH_2CHOHCH_2NHCH_3$ or $-CH_2CHOHCH_2NHCH(CH_3)_2$, $-CH_2CHOHCH_2NHCH_2CF_3$,

C_{1-6}alkyl substituted with one or two halogens and $-NR^{10}R^{11}$, for example $-CH_2CHFCH_2NH_2$,

C_{1-6}alkyl substituted with $-C(=O)-O-C_{1-6}$alkyl, for example $CH_2-C(=O)-O-CH_2CH_3$ or $-CH_2CH_2-C(=O)-O-CH_2CH_3$,

C_{1-6}alkyl substituted with $-O-C(=O)-NR^{10}R^{11}$, for example $-CH_2-C(=O)NH_2$, $-CH_2-C(=O)NHCH_3$, $-CH_2-C(=O)-NHCH_2CH_2OCH_3$, $-CH_2-C(=O)-NH-CH_2CH_2$-(pyrrolidin-1-yl) or $-CH_2-C(=O)-NH-CH_2CH_2$-(imidazol-2-yl), $-CH_2-C(=O)-NHCH_2CH_2OH$, $-CH_2-C(=O)-NHCH_2CH_2NH_2$,

C_{1-6}alkyl substituted with carboxyl, for example $-CH_2C(=O)OH$ or $-CH_2CH_2C(=O)OH$,

C_{1-6}alkyl substituted with $-O-C(=O)-NR^{10}R^{11}$, for example $-CH_2CH_2-O-C(=O)-NHCH_3$,

C_{1-6}alkyl substituted with $-NR^{12}-S(=O)_2-C_{1-6}$alkyl, for example $-CH_2CH_2NH-S(=O)_2-CH_3$, $-CH_2CH_2CH_2NH-S(=O)_2-CH_3$, $-CH_2CH_2NH-S(=O)_2-CH(CH_3)_2$ or $-CH_2CH_2NH-S(=O)_2-CH_2CH_3$,

C_{1-6}alkyl substituted with $-NR^{12}-S(=O)_2-NR^{14}R^{15}$, for example $-CH_2CH_2NH-S(=O)_2-N(CH_3)_2$ or $-CH_2CH_2CH_2NH-S(=O)_2-N(CH_3)_2$,

C_{1-6}alkyl substituted with hydroxyl and R^9, for example

propyl substituted with –OH and 1-pyrrolidinyl,

propyl substituted with –OH and 1-pyrrolidinyl where the 1-pyrrolidinyl is substituted at the 3 position by two fluorines,

propyl substituted with –OH and 1-pyrrolidinyl where the 1-pyrrolidinyl is substituted at the 3 position by a cyano group,

propyl substituted with –OH and 4-morpholinyl,

propyl substituted with –OH and 1-piperidinyl,

propyl substituted with –OH and isoindole-1, 3-dione,

-C_{1-6}alkyl-C(R^{12})=N-O-R^{12}, for example –$CH_2C(CH_3)$=N-O-H, –$CH_2C(CH_2OCH_3)$=N-O-H or –$CH_2C(CH(CH_3)_2)$=N-O-H

C_{1-6}alkyl substituted with –C(=O)-$NR^{10}R^{11}$, for example

–$CH_2C(=O)NH_2$,

–$CH_2C(=O)NHCH_3$,

–$CH_2C(=O)$-$NHCH_2CH_2OCH_3$,

–$CH_2C(=O)$-NH-CH_2CH_2-(pyrrolidin-1-yl) or –$CH_2C(=O)$-NH-CH_2CH_2-(imidazol-2-yl),–$CH_2C(=O)$-$NHCH_2CH_2OH$,-$CH_2C(=O)$-$NHCH_2CH_2NH_2$,

C_{1-6}alkyl substituted with –C(=O)-R^9, for example –$CH_2C(=O)$-R^9 and R^9 is 1-pyrrolidinyl,

C_{2-6}alkynyl substituted with R^9, for example -CH_2-C≡C—(2-imidazolyl) wherein the 2-imidazolyl is substituted in the 1 position by –CH_3 or -CH_2-C≡C—(5-imidazolyl) wherein the 5-imidazolyl is substituted in the 1 position by –CH_3,


C_{2-6}alkenyl, for example –CH_2-CH=CH_2,

C_{2-6}alkynyl, for example -CH_2-C≡C—H,

C_{1-6}alkyl substituted with C_{1-6}alkoxyC_{1-6}alkyl-C(=O)–, for example -CH_2–C(=O)-CH_2OCH_3, or

R^{13}.

In one embodiment the compound of formula (I) or formula (I^0) is a compound of formula ($I^{0'}$):




($I^{0'}$)

including any stereochemically isomeric form thereof;

and a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof, wherein n, R^2 and R^3 are as defined herein.


In one embodiment the compound of formula (I) or formula (I^0) is a compound of formula ($I^{0''}$)



($I^{0''}$)

including any stereochemically isomeric form thereof;
and a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof, wherein R^2 and R^3 are as defined herein.

In one embodiment the compound of formula (I) or formula (I^0) is a compound of formula ($I^{0'''}$)



($I^{0'''}$)

including any stereochemically isomeric form thereof;
and a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof, wherein R^3 is as defined herein.

In one embodiment there is provided a compound of formula ($I^{0'''}$) wherein R^3 is as defined in any of the embodiments above, in particular as defined at pages 86 line 20 to page 92 line 17.

In one embodiment the compound of formula (I) is a compound wherein one R^{1a} is selected from hydrogen, C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -NH(C_{3-8}cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl substituted with one or more fluoro atoms; and the other R^{1a} is selected from C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -NH(C_{3-8}cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl

substituted with one or more fluoro atoms; and wherein n, R^1, R^2 and R^3 are as defined herein.


In one embodiment the compound of formula (I) is a compound wherein each R^{1a} is independently selected from C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -NH(C_{3-8}cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl substituted with one or more fluoro atoms; and the other R^{1a} is selected from C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -NH(C_{3-8}cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl substituted with one or more fluoro atoms; and wherein n, R^1, R^2 and R^3 are as defined herein.

In one embodiment the compound of formula (I) is a compound wherein each R^{1a} is hydrogen; and wherein n, R^1, R^2 and R^3 are as defined herein.

In one embodiment every alkyl group within the R^3 definition is a C_{1-4}alkyl group.

In one embodiment every alkyl group within the R^3 definition is a linear C_{1-6}alkyl group, in particular a linear C_{1-4}alkyl group.

In one embodiment the compound of formula (I) is a compound of formula (I'):




(I')

including any stereochemically isomeric form thereof;

and a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof, wherein n, R^{1a}, R^2 and R^3 are as defined herein.


In one embodiment the compound of formula (I) is a compound of formula (I'')



(I'')

including any stereochemically isomeric form thereof;
and a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof, wherein R^{1a}, R^2 and R^3 are as defined herein.

In one embodiment the compound of formula (I) is a compound of formula (I''')



(I''')

including any stereochemically isomeric form thereof;
and a *N*-oxide thereof, a pharmaceutically acceptable salt thereof or a solvate thereof, wherein R^{1a} and R^3 is as defined herein.

In one embodiment there is provided a compound of formula (I), (I'), (I''), (I'''), (I^0), ($I^{0'}$), ($I^{0''}$) or ($I^{0'''}$) wherein every alkyl group within the R^3 definition is a linear C_{1-6}alkyl group. In one embodiment there is provided a compound of formula (I), (I'), (I''), (I'''), (I^0), ($I^{0'}$), ($I^{0''}$) or ($I^{0'''}$) wherein every alkyl group within the R^3 definition is a C_{1-4}alkyl group. In one embodiment there is provided a compound of formula (I), (I'), (I''), (I'''), (I^0), ($I^{0'}$), ($I^{0''}$) or ($I^{0'''}$) wherein every alkyl group within the R^3 definition is a linear C_{1-4}alkyl group.

For the avoidance of doubt, it is to be understood that each general and specific preference, embodiment and example for one substituent may be combined with each


general and specific preference, embodiment and example for one or more, preferably, all other substituents as defined herein and that all such embodiments are embraced by this application.

Methods for the Preparation of Compounds of Formula (I)

In this section, as in all other sections of this application unless the context indicates otherwise, references to formula (I) also include all other sub-groups and examples thereof as defined herein.

In general, compounds of formula (I) can be prepared according to the following reaction Scheme 1.

Scheme 1



In scheme 1, an intermediate of formula (IV) is prepared by reacting an intermediate of formula (II) wherein W_1 and W_2, each independently represent a suitable leaving group, such as for example halo, e.g. chloro or bromo and the like, with an intermediate of formula (III) in the presence of a suitable catalyst, such as for example

tetrakis(triphenylphosphine)palladium (0) or palladium (II) acetate , a suitable base, such as for example sodium carbonate, a suitable ligand, such as for example triphenylphosphine, and a suitable solvent or solvent mixture, such as for example ethylene glycol dimethylether and water. An intermediate of formula (II) wherein W_1 is chloro and W_2 is bromo can be prepared by reacting 7-bromo-2(1H)-quinoxalinone with phosphorus oxychloride, or alternatively with thionyl chloride and N,N-dimethylformamide in a suitable solvent, such as, for example toluene. An intermediate of formula (IV) can also be prepared by reacting 7-bromo-2-(1H-pyrazol-4-yl)quinoxaline with an intermediate W_{10}-R^1 wherein W_{10} represents a suitable leaving group, such as for example halo, e.g. bromo and the like. An intermediate of formula (IV) wherein the R^1 substituent carries a suitable protective group can be prepared according to the same protocol but wherein 7-bromo-2-(1H-pyrazol-4-yl)quinoxaline is reacted with an intermediate W_{10}-R^1-P wherein P represents a suitable protective group, such as for example –C(=O)-O-C$(CH_3)_3$. The intermediate of formula (IV) is then further reacted in a next step with an intermediate of formula (V) in the presence of a suitable catalyst, such as for example palladium (II) acetate, a suitable base, such as sodium tert-butoxide or Cs_2CO_3, a suitable ligand, such as for example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], and a suitable solvent or solvent mixture, such as for example dioxane or ethylene glycol dimethylether and water, resulting in an intermediate of formula (VI). Said intermediate of formula (VI) can then be reacted with an intermediate of formula (VII) wherein W_3 represents a suitable leaving group, such as for example halo, e.g. bromo and wherein R^x and R^y represent C_{1-4}alkyl, and R^z represent C_{1-4}alkyl or phenyl, for instance R^x and R^y represent CH_3 and R^z represents $C(CH_3)_3$ or phenyl, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide or N,N-dimethylacetamide, resulting in an intermediate of formula (VIII). Intermediates of formula (VIII) or intermediates of formula (VIII) wherein the R^1 substituent carries a suitable protective group can also be prepared by reacting an intermediate of formula (IV) or an intermediate of formula (IV) wherein the R^1 substituent carries a suitable protective group with an intermediate of formula (XXIII') wherein $R^{3a'}$ represent –C_{1-6}alkyl-O-Si$(R^x)(R^y)(R^z)$ in the presence of a suitable catalyst, such as for example palladium (II) acetate, a suitable ligand, such as for example racemic -2,2'-bis(diphenylphosphino)-1,1'-binaphtyl, a suitable base, such as for example Cs_2CO_3, and a suitable solvent, such as for example 1,2-dimethoxyethane. Intermediates of

formula (VIII) can be converted into a compound of formula (I) wherein R^3 represents –C_{1-6}alkyl-OH, said compounds being represented by formula (I-a) or compounds of formula (I-a) wherein the R^1 substituent carries a suitable protective group,, by reaction with tetrabutylammonium fluoride in the presence of a suitable solvent, such as for example tetrahydrofuran. This type of reaction can also be performed in the presence of a suitable acid, such as for example acetic acid or HCl, and a suitable solvent, such as for example tetrahydrofurane or dioxane. Alternatively, an intermediate of formula (VI) can react with an intermediate of formula (VII') wherein W_3 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide or N,N-dimethylacetamide, resulting in an intermediate of formula (XXV) which can then be deprotected in the presence of a suitable acid, such as for example HCl, and a suitable solvent, such as for example an alcohol, e.g. methanol or isopropanol, to give a compound of formula (I-a).The compounds of formula (I-a) or compounds of formula (I-a) wherein the R^1 substituent carries a suitable protective group can be reacted with methanesulfonyl chloride in the presence of a suitable base, such as for example triethylamine, diisopropylethanamine or N,N-dimethyl-4-aminopyridine, and a suitable solvent, such as for example dichloromethane or tetrahydrofuran,, to result in an intermediate of formula (IX) (mesylate derivative) or an intermediate of formula (IX') (chloride derivative) or intermediates of formula (IX) or (IX') wherein the R^1 substituent carries a suitable protective group. Intermediates of formula (IX) or (IX') can then be reacted with an intermediate of formula (X) to obtain a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with $NR^{10}R^{11}$, said compounds being represented by formula (I-b) or compounds of formula (I-b) wherein the R^1 substituent carries a suitable protective group. This reaction may optionally be performed in the presence of a suitable base, such as for example triethylamine, K_2CO_3, Na_2CO_3 or sodium hydride and optionally a suitable solvent, such as for example acetonitrile, tetrahydrofuran, dioxane, N,N-dimethylformamide, 1-methyl-pyrrolidinone, a suitable alcohol, e.g. 1-butanol and the like. This type of reaction can also be performed with a suitable salt of the intermediate of formula (X), e.g. HCl salt of intermediate of formula (X), or may be performed in the presence of potassium iodide. In this way compounds wherein R^3 represents iodoC_{1-6}alkyl can be obtained. Compounds of formula (I-b) wherein the R^1 substituent carries a suitable protective group can be converted in a compound of formula (I-b) by reaction with a suitable acid, such as for

example trifluoroacetic acid, in the presence of a suitable solvent, such as for example dichloromethane.


Intermediates of formula (IX) can also react with a suitable nitrogen containing ring within the definition of R^9, said ring being represented by formula (XXI) or a suitable salt of an intermediate of formula (XXI), in the presence of a suitable solvent, such as for example acetonitrile, 1-methyl-2-pyrrolidinone, or an alcohol, e.g. 1-butanol, optionally in the presence of potassium iodide or a suitable base, such as for example Na_2CO_3, K_2CO_3 or triethylamine, resulting in a compound of formula (I-d). Intermediates of formula (IX) can also react with an intermediate of formula (X-a) wherein P represents a suitable protective group, such as for example $-C(=O)-O-C(CH_3)_3$, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example dimethylacetamide, resulting in an intermediate of formula (XXX) which can be deprotected to a compound of formula (I-b-1) in the presence of a suitable acid, such as for example HCl or trifluoroacetic acid, and a suitable solvent, such as for example dichloromethane or an alcohol, e.g. methanol. Intermediates of formula (XXX) can also be prepared by reacting an intermediate of formula (VI) with an intermediate of formula W_6-C_{1-6}alkyl-$NR^{10}P$ wherein W_6 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, or $-O-S(=O)_2-CH_3$, and P is as defined above, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, e.g. N,N-dimethylformamide or N,N-dimethylacetamide. Alternatively compounds of formula (1-d) or (1-b-1) can also be prepared by reacting respectively an intermediate of formula (VI) with an intermediate of formula W_6-C_{1-6}alkyl-Ncycle or W_6-C_{1-6}alkyl-NHR^{10} wherein W_6 is as defined above.

Intermediates of formula (VI) can react with W_6-R^{3a} wherein W_6 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, or $-O-S(=O)_2-CH_3$, and R^{3a} represents optionally substituted C_{1-6}alkyl, such as for example $-CH_2-C_3H_5$, in the presence of a suitable base, such as for example sodium hydride or Cs_2CO_3, and a suitable solvent, such as for example N,N-dimethylformamide, N,N-dimethylacetamide or acetonitrile, resulting in a compound of formula (I-c). In this way, compounds of formula (I-c) wherein R^3 represents $-S(=O)_2-N(CH_3)_2$ can also be prepared by reacting an intermediate of formula (VI) with dimethylsulfamoyl chloride, in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,N-dimethylformamide.

Compounds of formula (I-c) wherein R^{3a} represents $-CH_2$-C(OH)(R')(R'') wherein R' represents optionally substituted C_{1-4}alkyl and R'' represents hydrogen or optionally substituted C_{1-4}alkyl, said compounds being represented by formula (I-c-1), can be prepared by reacting the intermediate of formula (VI) with an intermediate of formula (XXII) in the presence of a suitable base, such as for example sodium hydride, Cs_2CO_3, or potassium hydroxide, and a suitable solvent, such as for example N,N-dimethylformamide, N,N-dimethylacetamide, acetonitrile or water.
Intermediates of formula (IV) can also react with an intermediate of formula (XXIII) in the presence of a suitable catalyst, such as for example palladium (II) acetate or tris(dibenzylideneacetone)dipalladium (0), a suitable base, such as for example sodium tert-butoxide, a suitable ligand, such as for example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] or 2-dicyclohexylphosphino-2'-(N,N-dimethylamino)biphenyl, and a suitable solvent, such as for example dioxane, resulting in a compound of formula (I-c).

Compounds of formula (I-b) wherein R^{11} is C_{1-6}alkyl substituted with amino, said compounds being represented by formula (I-b-2), can also be prepared according to the following reaction Scheme 1A.

Scheme 1A




In Scheme 1A, a compound of formula (I-b-1) is reacted with N-(3-bromopropyl)phtalimide in the presence of a suitable base, such as for example potassium carbonate, and a suitable solvent, such as for example acetonitrile, resulting in an intermediate of formula (XXXVI) which can be converted into a compound of

formula (I-b-2) by reaction with hydrazine in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol.

Compounds of formula (I-b) wherein R^1 is hydrogen, said compounds being represented by formula (I-b-3) can be prepared according to reaction Scheme 1A1.

Scheme 1A1




In Scheme 1A1, an intermediate of formula (I-a-1) is reacted with methanesulfonyl chloride in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example dichloromethane resulting in an intermediate of formula (IX-1) wherein R^u represents $–O-S(=O)_2-CH_3$, which is converted into a compound of formula (I-b-3) by reaction with an intermediate of formula (X) in the presence of a suitable solvent, such as for example acetonitrile.

It is considered to be within the knowledge of the person skilled in the art to recognize in which condition and for which definitions of R^{1a} in the reactions of Scheme 1a and Scheme 1a1 a protective group may be appropriate for the reactions to be carried out. For instance, a hydroxyl group within the definition of R^{1a} may be protected with a tert. butyldimethylsilyl moiety; a NH group within the definition of R^{1a} may be protected with a $–C(=O)-O-C(CH_3)_3$ group.

It is also considered to be within the knowledge of the person skilled in the art to recognize appropriate deprotection reactions.

Compounds of formula (I) wherein R^3 represents optionally substituted C_{2-6}alkynyl, said compounds being represented by formula (I-k), can be prepared according to reaction Scheme 1B.

Scheme 1B




(VI) (I-k)

In Scheme 1B, an intermediate of formula (VI) is reacted with an intermediate of formula W_{11}-R^{3b} wherein R^{3b} represents optionally substituted C_{2-6}alkynyl and W_{11} represents a suitable leaving group such as for example halo, e.g. chloro, or –O-S(=O)$_2$-CH_3, in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,N-dimethylformamide. The intermediate W_{11}-R^{3b} wherein W_{11} represents –O-S(=O)$_2$-CH_3, can be prepared by reacting the corresponding alcohol derivative with methanesulfonyl chloride in the presence of a suitable base, such as for example triethylamine or 4-dimethylaminopyridine, and a suitable solvent, such as for example dichloromethane.

Compounds of formula (I-k), wherein R^{3b} represents C_{2-6}alkynyl substituted with hydroxyl, said compounds being represented by formula (I-k-1), can be prepared according to the following reaction Scheme 1C.


Scheme 1C



In Scheme 1C, an intermediate of formula (VI) is reacted with an intermediate of formula (XXXVIII) in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,n-dimethylformamide, resulting in an intermediate of formula (VIII'), which is converted into a compound of formula (I-k-1) by reaction with a suitable acid, such as for example trifluoroacetic acid, in the presence of a suitable solvent, such as for example tetrahydrofuran.

Compounds of formula (I-k), wherein R^{3b} represents C_{2-6}alkynyl, said compounds being represented by formula (I-k-2), can be prepared according to the following reaction Scheme 1D.


Scheme 1D



In Scheme 1D, a compound of formula (I-k-2) is prepared by deprotecting an intermediate of formula (XXXXII) in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example an alcohol, e.g. methanol and the like. Said intermediate of formula (XXXXII) can be prepared by reacting an intermediate of formula (VI) with W_{13}-C_{2-6}alkynyl-$Si(CH_3)_3$ in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,N-dimethylformamide.

Compounds of formula (I), wherein R^3 represents ethyl substituted with $-P(=O)(OC_{1-6}alkyl)_2$, said compounds being represented by formula (I-l), can be prepared according to the following reaction Scheme 1E.


Scheme 1E



In scheme 1E, an intermediate of formula (VI) is reacted with di(C_{1-6}alkyl)vinylphosphonate in the presence of a suitable catalyst, such as for example tri-N-butylphosphine, and a suitable solvent, such as for example acetonitrile resulting in a compound of formula (I-l).

Intermediates of formula (VI) can also be prepared according to the following reaction Scheme 2.

Scheme 2




In Scheme 2, an intermediate of formula (XII) is prepared by reacting an intermediate of formula (XI) wherein W_1 represents a suitable leaving group, such as for example halo, e.g. chloro and the like, with an intermediate of formula (III) in the presence of a suitable catalyst, such as for example tetrakis(triphenylphosphine)palladium (0), a suitable base, such as for example Na_2CO_3, and a suitable solvent or solvent mixture, such as for example ethylene glycol dimethylether and water. The intermediate of formula (XII) is hydrogenated in a next step to an intermediate of formula (XIII) in the presence of a suitable catalyst, such as for example Nickel, and a suitable solvent, such as for example an alcohol, e.g. methanol, or tetrahydrofuran, or mixtures thereof.

Intermediates of formula (XIII) can also be prepared by reacting an intermediate of formula (IV) with NH_4OH in the presence of Cu_2O. In a next step, the intermediate of formula (XIII) is reacted with an intermediate of formula (XIV) wherein W_5 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, in the presence of a suitable catalyst, such as for example palladium (II) acetate, a suitable base, such as for example sodium tert-butoxide, a suitable ligand, such as for example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], and a suitable solvent, such as for example ethylene glycol dimethyl ether or dioxane, resulting in an intermediate of formula (VI). This reaction may also be performed in the presence of $Pd_2(dba)_3$ as catalyst, Xphos as ligand, a suitable base, such as for example Cs_2CO_3, and a suitable solvent, such as for example an alcohol, e.g. butanol.
Intermediates of formula (IV) wherein R^1 is hydrogen can be converted into an intermediate of formula (IV) wherein R^1 is other than hydrogen, said R^1 being represented by $R^{1'}$, by reaction with W_{14}-$R^{1'}$ wherein W_{14} is a suitable leaving group, such as for example halo, e.g. bromo, in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,N-dimethylformamide.


Intermediates of formula (VI) can alternatively also be prepared according to the following reaction Scheme 3.

Scheme 3




In Scheme 3, an intermediate of formula (XV) is reacted with an intermediate of formula (V) in the presence of a suitable catalyst, such as for example palladium (II) acetate, a suitable base, such as for example sodium tert-butoxide, a suitable ligand, such as for

example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], and a suitable solvent, such as for example ethylene glycol dimethyl ether, resulting in an intermediate of formula (XVI). In a next step, the intermediate of formula (XVI) is reacted with $P(=O)Cl_3$ or chlorosuccinimide, optionally in the presence of a solvent, such as for example acetonitrile, resulting in an intermediate of formula (XVII) which is converted into an intermediate of formula (VI) by reaction with an intermediate of formula (III) in the presence of a suitable catalyst, such as for example tetrakis(triphenylphosphine)palladium (0) or tris(dibenzylideneacetone)dipalladium (0), a suitable base, such as for example Na_2CO_3 or K_3PO_4, optionally in the presence of a suitable ligand, such as for example 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl, and a suitable solvent, such as for example ethylene glycol dimethylether.

In the above reaction, an intermediate of formula (III) can react in its protected form, such as for example  . The resulting protected intermediate of formula (VI) can be converted into the deprotected $-C_{1-6}$alkyl-OH intermediate by reaction with tetrabutylammonium fluoride, in the presence of a suitable solvent, such as for example tetrahydrofuran. Said $-C_{1-6}$alkyl-OH can be converted into $-C_{1-6}$alkyl-NH_2 by first reacting the $-C_{1-6}$alkyl-OH with methanesulfonyl chloride in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example dichloromethane, followed by reacting the obtained intermediate with di-tert-butyl-iminocarboxylate in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,N-dimethylformamide, followed by reaction with a suitable acid, such as for example trifluoroacetic acid, in a suitable solvent, such as for example dichloromethane.

Intermediates of formula (VIII) can alternatively also be prepared according to the following reaction Scheme 4.


Scheme 4



In Scheme 4, an intermediate of formula (XVII) is reacted with an intermediate of formula (VII) in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide, resulting in an intermediate of formula (XVIII). The intermediate of formula (XVIII) can then be reacted with an intermediate of formula (III) in the presence of a suitable catalyst, such as for example $Pd_2(dba)_3$, a suitable base, such as for example K_3PO_4, a suitable ligand, such as for example 2-dicyclohexylphosphino-2',6'-dimethoxy-biphenyl or S-Phos, and a suitable solvent, such as for example dioxane or water or mixtures thereof.

Intermediates of formula (VIII') can be prepared according to the following reaction Scheme 4A.

Scheme 4A




(XVIII) (XXXVII) (VIII')

In Scheme 4A, an intermediate of formula (XVIII) is reacted with an intermediate of formula (XXXVII) in the presence of a suitable catalyst, such as for example tetrakis(triphenylphisphine)palladium (0), and a suitable solvent, such as for example toluene.

Intermediates of formula (VIII') can be further reacted according to the following reaction Scheme 4B.

Scheme 4B




In Scheme 4B, intermediates of formula (VIII') wherein R^1 represents hydrogen, said intermediates being represented by formula (VIII'-a), can be converted into an intermediate of formula (VIII') wherein R^1 represents haloC_{1-6}alkyl, said intermediates being represented by formula (VIII'-b) by reaction with W_{12}-C_{1-6}alkyl-halo wherein W_{12} represents a suitable leaving group, such as for example halo, e.g. chloro, in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,N-dimethylformamide. Said intermediates of formula (VIII'-b) can be converted into an intermediate of formula (VIII'-c) wherein R^1 represents an optionally substituted R^6, by reaction with optionally substituted R^6 in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example acetonitrile. When in an intermediate of formula (VIII'-c) the R^6 carries a hydroxyl group as in an intermediate of formula (VIII'-c-1), then said hydroxyl group can be protected by a suitable protective group P, such as for example –O-C(=O)-C_{1-6}alkyl, by reaction with

C_{1-6}alkyl-C(=O)-W_{12}, in the presence of a suitable base, such as for example triethylamine, 4-dimethylaminopyridine, and a suitable solvent, such as for example dichloromethane, resulting in an intermediate of formula (VIII'-c-2) which can be converted into an intermediate of formula (XXXIX) by reaction with tetrabutylammonium fluoride in the presence of a suitable solvent, such as for example tetrahydrofuran. Said intermediate of formula (XXXIX) can be converted into an intermediate of formula (XXXX) by reaction with methansulfonyl chloride in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example dichloromethane, which can be converted into an intermediate of formula (XXXXI) by reaction with an intermediate of formula (X) in a suitable solvent, such as for example acetonitrile. Said intermediate of formula (XXXXI) can then be deprotected into a compound of formula (I-b-4) in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example an alcohol, e.g. methanol and the like.

Intermediates of formula (VIII') can also be reacted to prepare compounds of the present invention according to the reaction schemes as presented in Scheme 1. It is considered to be within the knowledge of the person skilled in the art to recognize in which condition and for which definitions of R^{1a} a protective group may be appropriate for the reactions to be carried out. For instance, a hydroxyl group within the definition of R^{1a} may be protected with a tert. butyldimethylsilyl moiety; a NH group within the definition of R^{1a} may be protected with a –C(=O)-O-$C(CH_3)_3$ group.
It is also considered to be within the knowledge of the person skilled in the art to recognize appropriate deprotection reactions.

Compounds of formula (I) wherein R^3 represents optionally substituted C_{1-6}alkyl, said compounds being represented by formula (I-c), can also be prepared according to the below reaction Scheme 5.


Scheme 5



In Scheme 5, an intermediate of formula (XVII) is reacted with W_6-R^{3a} wherein W_6 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, and R^{3a} represents optionally substituted C_{1-6}alkyl, such as for example $-CH_2-C_3H_5$, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide, resulting in an intermediate of formula (XIX). In a next step, the intermediate of formula (XIX) is reacted with an intermediate of formula (III) in the presence of a suitable catalyst, such as for example tetrakis(triphenyl)phosphine palladium or $Pd_2(dba)_3$ (tris(dibenzylideneacetone) dipalladium (0)), a suitable ligand, such as 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl, a suitable base, such as for example Na_2CO_3 or K_3PO_4, and a suitable solvent, such as for example ethylene glycol dimethylether or dioxane or water.

Compounds of formula (I-c) can alternatively also be prepared according to the below reaction Scheme 6.


Scheme 6



In Scheme 6, an intermediate of formula (IV) is reacted with R^{3a}-NH_2 in the presence of a suitable catalyst, such as for example palladium (II) acetate, a suitable base, such as for example sodium tert-butoxide, and a suitable ligand, such as for example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], resulting in an intermediate of formula (XX) which is reacted in a next step with an intermediate of formula (XIV) in the presence of a suitable catalyst, such as for example palladium (II) acetate or $Pd_2(dba)_3$ (tris(dibenzylidene acetone) dipalladium (0)), a suitable ligand such as for example 2-dicyclohexylphosphino-tris-isopropyl-biphenyl or 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], a suitable base, such as for example sodium tert-butoxide, and a suitable solvent, such as for example ethylene glycol dimethylether.

Compounds of formula (I) wherein R^3 is C_{1-6}alkyl substituted with 5-amino-1,3,4-oxadiazolyl can be prepared according to the below reaction Scheme 7.


Scheme 7



In Scheme 7, a compound of formula (I-h) is reacted with NH_2-NH_2 in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol resulting in an intermediate of formula (XXXI) which is then reacted in a next step with W_8-CN, wherein W_8 represents a suitable leaving group, such as for example halo, e.g. bromo, in the presence of a suitable base, such as for example $NaHCO_3$, and a suitable solvent, such as for example water or dioxane.

Compounds of formula (I) wherein R^3 is C_{1-6}alkyl substituted with 3,3-dimethyl-morpholine can be prepared according to the below reaction Scheme 7.A

Scheme 7A




In Scheme 7A, a compound of formula (I-j'') is reacted with 2-amino-2-methyl-1-propanol in the presence of a suitable base, such as for example NaH and in the presence of a suitable solvent, such as for example N,N-dimethylformamide resulting in an intermediate of formula (XXXII) of which the NH2 moiety is protected by a suitable protecting group P, such as for example –C(=O)-O-C(CH3)3, by reaction with for


instance di-tert-butyl dicarbonate in the presence of a suitable solvent, such as for example dioxane, and a suitable base, such as for example NaHCO3, resulting in an intermediate of formula (XXXIII). In a next step, said intermediate is reacted with methanesulfonyl chloride in the presence of a suitable solvent, such as for example dichloromethane, and a suitable base, such as for example triethylamine resulting in an intermediate of formula (XXXIV) which is converted into an intermediate of formula (XXXV) by reaction with a suitable acid, such as for example trifluoroacetic acid, in the presence of a suitable solvent, such as for example dichloromethane. The intermediate of formula (XXXV) is converted into a compound of formula (I-j') by reaction with a suitable base, such as for example *N,N*-diisopropylethylamine and triethylamine in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.

As already shown above, compounds of formula (I) or some of the above-described intermediates can be prepared by deprotecting the corresponding protected compounds. Other protection-deprotection reactions are shown in the following reaction Scheme 8.

Scheme 8



In Scheme 8, compounds of formula (I) wherein R^1 represents hydroxyC_{1-6}alkyl, said compounds being represented by formula (I-e), can be prepared by deprotecting an


intermediate of formula (XXVI) in the presence of a suitable acid, such as for example HCl or trifluoroacetic acid, or a suitable de-silylating agent, such as for example tetrabutyl ammonium fluoride, and a suitable solvent, such as an alcohol, e.g. methanol, or tetrahydrofuran. Intermediates of formula (XXVI) can be prepared by reacting a compound of formula (I) wherein R^1 is hydrogen, said compounds being represented by formula (I-f), with an intermediate of formula (XXIV) wherein W9 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, and P represents a suitable protective group, such as for example –O-Si(CH_3)$_2$($C(CH_3)_3$) or , in the presence of a suitable base, such as for example sodium hydride or K_2CO_3, and a suitable solvent, such as for example N,N-dimethylformamide or acetonitrile.

Compounds of formula (I) wherein R^1 represents C_{1-6}alkyl substituted with –C(=O)-R^6 wherein R^6 is an appropriate nitrogen containing ring linked to the C(=O) moiety via the nitrogen atom, said compounds being represented by formula (I-g), can be prepared by reacting an intermediate of formula (XXIX) with an intermediate of formula (XXI) in the presence of suitable peptide coupling reagents such as, 1-hydroxy-benzotriazole and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide HCl. Intermediates of formula (XXIX) can be prepared by reacting an intermediate of formula (XXVIII) with LiOH in the presence of a suitable solvent, such as for example tetrahydrofuran or water. Intermediates of formula (XXVIII) can be prepared by reacting a compound of formula (I-f) with an intermediate of formula (XXVII) wherein W_9 is as defined above, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide.

Compounds of formula (I-i) can be prepared starting from an intermediate of formula (XXIX) by reaction with NHR^4R^5 in the presence of suitable peptide coupling reagents such as 1-hydroxy-benzotriazole and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide HCl and a suitable base, such as triethylamine, and a suitable solvent, such as for example dichloromethane.

Further protection-deprotection reactions can also be used as outlined in the following reaction Scheme 9.

Scheme 9



In Scheme 9, the following reaction conditions apply:

A ; in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide.

B : in the presence of a suitable catalyst, such as for example palladium (II)acetate, a suitable base, such as for example sodium tert-butoxide, a suitable ligand, such as for example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], and a suitable solvent, such as for example dioxane or ethylene glycol dimethylether.

C : in the presence of a suitable catalyst, such as for example palladium (II)acetate, a suitable base, such as for example sodium tert-butoxide, a suitable ligand, such as for example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], and a suitable solvent, such as for example dioxane or ethylene glycol dimethylether.


D : in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example dichloromethane.

E : in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example 1-methyl-2-pyrrolidinone.

F : in the presence of hydrazine monohydrate, and a suitable solvent, such as for example an alcohol, e.g. ethanol.


G : in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example tetrahydrofuran.

It is considered to be within the knowledge of the person skilled in the art to recognize in which condition and on which part of the molecule a protective group may be appropriate. For instance, protective group on the R^1 substituent or on the pyrrazole moiety, or protective group on the R^3 substituent or on the R^2 substituent or combinations thereof. The skilled person is also considered to be able to recognize the most feasible protective group, such as for example –C(=O)-O-C_{1-4}alkyl or




or O-Si$(CH_3)_2(C(CH_3)_3)$ or -CH_2-O-CH_2CH_2-O-CH_3.

The present invention also comprises deuterated compounds. These deuterated compounds may be prepared by using the appropriate deuterated intermediates during the synthesis process. For instance an intermediate of formula (IV-a)



can be converted into an intermediate of formula (IV-b)



by reaction with iodomethane-D3 in the presence of a suitable base, such as for example cesium carbonate, and a suitable solvent, such as for example acetonitrile.

The compounds of formula (I) may also be converted into each other via art-known reactions or functional group transformations.

For instance, compounds of formula (I) wherein R^1 represents tetrahydropyranyl can be converted into a compound of formula (I) wherein R^1 represents hydrogen, by reaction with a suitable acid, such as for example HCl or trifluoroacetic acid, in the presence of a suitable solvent, such as for example dichloromethane, dioxane, or an alcohol, e.g. methanol, isopropanol and the like.

Compounds of formula (I) wherein R^1 or R^3 represent monohaloalkyl, can be converted into a compound of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl substituted with a ring moiety as defined hereinabove by the intermediate of formula (XXI) and linked to the C_{1-6}alkyl moiety by the nitrogen atom, by reaction with an intermediate of formula (XXI) optionally in the presence of a suitable base, such as for example triethylamine or K_2CO_3 or sodium hydride, and optionally in the presence of a suitable solvent, such as for example acetonitrile, N,N-dimethylformamide or 1-methyl-2-pyrrolidinone.

Compounds of formula (I) wherein R^1 or R^3 represents C_{1-6}alkyl-OH, can be converted into a compound of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl-F by reaction with diethylaminosulfur trifluoride in the presence of a suitable solvent, such as for example dichloromethane and in the presence of catalytic amounts of an alcohol, such as for example ethanol. Likewise, a compound of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl substituted with R^6 or R^9 wherein said R^6 or R^9 is substituted with OH, can be converted into a compound of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl substituted with R^6 or R^9 wherein said R^6 or R^9 is substituted with F, by reaction with diethylaminosulfur trifluoride in the presence of a suitable solvent, such as for example dichloromethane.

Compounds of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl substituted with R^6 or R^9 wherein said R^6 or R^9 is substituted with $-C(=O)-O-C_{1-6}$alkyl, can be converted into a

compound of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl substituted with R^6 or R^9 wherein said R^6 or R^9 is substituted with $-CH_2$-OH, by reaction with $LiAlH_4$ in the presence of a suitable solvent, such as for example tetrahydrofuran.

Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with 1,3-dioxo-2H-isoindol-2-yl, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with amino, by reaction with hydrazine monohydrate in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol.

Compounds of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl substituted with amino, can be converted into a compound of formula (I) wherein R^1 or R^3 represents C_{1-6}alkyl substituted with $-NH$-$S(=O)_2$-C_{1-6}alkyl, by reaction with Cl-$S(=O)_2$-C_{1-6}alkyl in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example dichloromethane.




Compounds of formula (I) wherein R^1 or R^3 represents C_{1-6}alkyl substituted with halo, can be converted into a compound of formula (I) wherein R^1 or R^3 represent C_{1-6}alkyl substituted with NR^4R^5 or $NR^{10}R^{11}$, by reaction with NHR^4R^5 or $NHR^{10}R^{11}$, either using such amino in large excess or in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example acetonitrile, N,N-dimethylacetamide or 1-methyl-pyrrolidinone.

Compounds of formula (I) wherein R^1 represents hydrogen, can be converted into a compound of formula (I) wherein R^1 represents polyhaloC_{1-6}alkyl or polyhydroxyC_{1-6}alkyl or C_{1-6}alkyl or $-S(=O)_2$-$NR^{14}R^{15}$ or $-S(=O)_2$-C_{1-6}alkyl, by reaction with polyhaloC_{1-6}alkyl-W or polyhydroxyC_{1-6}alkyl-W or C_{1-6}alkyl-W or W-$S(=O)_2$-$NR^{14}R^{15}$ or W-$S(=O)_2$-C_{1-6}alkyl, wherein W represents a suitable leaving group, such as for example halo, e.g. bromo and the like, in the presence of a suitable base, such as for example sodium hydride or K_2CO_3 or triethylamine or 4-dimethylamino-pyridine or diisopropylamine, and a suitable solvent, such as for example N,N-dimethylformamide or acetonitrile or dichloromethane.

Compounds of formula (I) wherein R^1 represents hydrogen can also be converted into a compound of formula (I) wherein R^1 represents C_{1-6}alkyl-OH, by reaction with W-C_{1-6}alkyl-O-$Si(CH_3)_2(C(CH_3)_3)$ in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide.

Compounds of formula (I) wherein R^1 represents hydrogen, can also be converted into compound of formula (I) wherein R^1 represents ethyl substituted with $-S(=O)_2$-C_{1-6}alkyl, by reaction with C_{1-6}alkyl-vinylsulfone, in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example an alcohol, e.g.

methanol or by reaction with C_{1-6}alkyl-2-bromoethylsulfone in the presence of a suitable deprotonating agent, such as for example NaH, and a suitable solvent, such as for example dimethyformamide.

Compounds of formula (I) wherein R^1 represents hydrogen can also be converted into a compound of formula (I) wherein R^1 represents $-CH_2$-CHOH-CH_2-, by reaction with  in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide, wherein  represents a suitable nitrogen containing ring within the definition of R^6.

Compounds of formula (I) wherein R^1 represents C_{1-6}alkyl substituted with R^6 wherein said R^6 is substituted with $-C(=O)$-O-C_{1-6}alkyl or $-S(=O)_2$-$NR^{14}R^{15}$ or wherein R^3 represents C_{1-6}alkyl substituted with R^9 wherein said R^9 is substituted with $-C(=O)$-O-C_{1-6}alkyl or $-S(=O)_2$-$NR^{14}R^{15}$, can be converted into a compound of formula (I) wherein the R^6 or R^9 is unsubstituted, by reaction with a suitable acid, such as for example HCl and a suitable solvent, such as for example dioxane, acetonitrile or an alcohol, e.g. isopropylalcohol. Compounds of formula (I) wherein R^1 represents C_{1-6}alkyl substituted with R^6 wherein said R^6 is a ring moiety comprising a nitrogen atom which is substituted with $-CH_2$-OH or wherein R^3 represents C_{1-6}alkyl substituted with R^9 wherein said R^9 is a ring moiety comprising a nitrogen atom which is substituted with $-CH_2$-OH, can be converted into a compound of formula (I) wherein the R^6 or R^9 is unsubstituted, by reaction with sodium hydroxide, in the presence of a suitable solvent, such as for example tetrahydrofuran.

Compounds of formula (I) wherein R^1 represents C_{1-6}alkyl substituted with R^6 or R^3 represents C_{1-6}alkyl substituted with R^9, wherein said R^6 or said R^9 is unsubstituted, can be converted into a compound of formula (I) wherein said R^6 or said R^9 is substituted with C_{1-6}alkyl, by reaction with W-C_{1-6}alkyl wherein W is as defined above, in the presence of a suitable base. Such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide.

Compounds of formula (I) wherein R^1 or R^3 represent hydroxyC_{1-6}alkyl, can be converted into the corresponding carbonyl compound, by reaction with dess-Martin-

periodinane, in the presence of a suitable solvent, such as for example dichloromethane.

Compounds of formula (I) wherein R^1 represents C_{1-6}alkyl substituted with R^6 or R^3 represents C_{1-6}alkyl substituted with R^9, wherein said R^6 or said R^9 is substituted with C_{1-6}alkyl-halo, can be converted into a compound of formula (I) wherein said R^6 or said R^9 is substituted with C_{1-6}alkyl-CN, by reaction with sodium cyanide, in the presence of a suitable solvent, such as for example water or an alcohol, e.g. ethanol.

Compounds of formula (I) wherein R^1 represents C_{1-6}alkyl substituted with R^6 wherein said R^6 is unsubstituted or wherein R^3 represents C_{1-6}alkyl substituted with R^9 wherein said R^9 is unsubstituted, can be converted into a compound of formula (I) wherein R^6 or R^9 is substituted with $-CH_3$ or $-CH(CH_3)_2$, by reaction with formaldehyde or acetone and $NaBH_3CN$, in the presence of a suitable solvent, such as for example tetrahydrofuran or an alcohol, e.g. methanol.

Compounds of formula (I) wherein R^1 contains a R^6 substituent substituted with OH or wherein R^3 contains a R^9 substituent substituted with OH, can be converted into a compound of formula (I) wherein the R^6 or R^9 substituent is substituted with C_{1-6}alkyloxy, by reaction with W-C_{1-6}alkyl, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example N,N-dimethylformamide.

Compounds of formula (I) wherein R^1 contains a R^6 substituent substituted with C_{1-6}alkyloxy or wherein R^3 contains a R^9 substituent substituted with C_{1-6}alkyloxy, can be converted into a compound of formula (I) wherein the R^6 or R^9 substituent is substituted with –OH by reaction with a suitable acid, such as for example hydrochloric acid.

Compounds of formula (I) wherein R^1 contains a R^6 substituent substituted with halo or wherein R^3 contains a R^9 substituent substituted with halo can be converted into a compound of formula (I) wherein the R^6 or R^9 substituent is substituted with $-NR^{14}R^{15}$ by reaction with $NHR^{14}R^{15}$ in a suitable sovent, such as for example 1-methyl-pyrrolidinone.

Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with COOH, by reaction with LiOH in the presence of a suitable solvent, such as for example tetrahydrofuran. Said compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with COOH, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with $-C(=O)-NH_2$ or –C(=O)-$NHCH_3$ or $-C(=O)NR^{10}R^{11}$, by reaction with $NH(Si(CH_3)_3)_2$ or $MeNH_3^+Cl^-$ or $NHR^{10}R^{11}$ in the presence of suitable peptide coupling reagents such as for example 1-(3-

dimethylaminopropyl)-3-ethylcarbodiimide HCl and 1-hydroxybenzotriazole, a suitable base, such as for example triethylamine and a suitable solvent such as for example dichloromethane or N,N-dimethylformamide. Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, can also be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with 2-imidazolyl, by reaction under N_2 with ethylenediamine and trimethylaluminium in the presence of a suitable solvent, such as for example toluene and heptane. This compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with 2-imidazolyl, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –C(=O)-NH-$(CH_2)_2$-NH_2 by reaction with sodium hydroxide. Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with COOH, can also be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –C(=O)-N(CH_3)(OCH_3) by reaction with dimethylhydroxylamine, in the presence of carbonyldiimidazole and a suitable solvent, such as for example dichloromethane.

Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with , can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with 2 OH's, by reaction with a suitable acid, such as for example trifluoroacetic acid, and a suitable solvent, such as for example dioxane or water. These compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with , can also be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with OH and $NR^{10}R^{11}$, by reaction with $NH_2R^{10}R^{11}$ optionally in salt form, such as for example $NHR^{10}R^{11+}Cl^-$, optionally in the presence of a suitable base, such as for example sodium hydride or Na_2CO_3 or triethylamine or KI, and in the presence of a suitable solvent, such as for example N,N-dimethylformamide or an alcohol, e.g. 1-butanol or ethanol.

Compounds of formula (I) wherein R^3 represents C_{1-3}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, can be converted into a compound of formula (I) wherein R^3 represents C_{1-3}alkyl substituted with –C(CH3)$_2$-OH, by reaction with iodomethane and Mg powder, in the presence of a suitable solvent, such as for example diethylether or tetrahydrofuran.

Compounds of formula (I) wherein R^3 represents C_{1-5}alkyl substituted with –C(=O)-O-C_{1-6}alkyl, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –OH, by reaction with $LiAlH_4$ in a suitable solvent, such as for example tetrahydrofuran.

Compounds of formula (I) wherein R^3 represents C_{1-5}alkyl substituted with –OH, can be converted into a compound of formula (I) wherein R^3 represents C_{1-5}alkyl substituted with –O-C(=O)-C_{1-6}alkyl by reaction with Cl-C(=O)-C1-6alkyl in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example tetrahydrofuran.


Compounds of formula (I) wherein R^3 represents $–CH_2-CH=CH_2$, can be converted into a compound of formula (I) wherein R^3 represents $–CH_2-CHOH-CH_2-OH$, by reaction with potassium permanganate, and a suitable solvent, such as for example acetone or water.


Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –C(=O)-C_{1-4}alkyl, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –C(C_{1-4}alkyl)=N-OH, by reaction with hydroxylamine, in the presence of a suitable base, such as for example pyridine, and a suitable solvent, such as for example an alcohol, e.g. ethanol.

Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with NH_2, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with -NH-C(=O)-R^6 or with -NH-C(=O)-C_{1-6}alkyl or with -NH-C(=O)-polyhydroxyC_{1-6}alkyl or with -NH-C(=O)-polyhaloC_{1-6}alkyl or with -NH-C(=O)-polyhydroxypolyhaloCvalkyl, by reaction with the corresponding COOH analogue, e.g. R^6-COOH or CF_3-C(CH_3)(OH)-COOH and the like, in the presence of suitable peptide coupling reagents such as 1-hydroxy-benzotriazole and 1-(3-dimethylamino)propyl)carbodiimide optionally in the presence of a suitable base, such as for example triethylamine. Said compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with NH_2, can also be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with NH-C(=O)-CF_3, by reaction with trifluoroacetic anhydride, in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example tetrahydrofuran. Said compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with NH_2, can also be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with –NH-polyhaloC_{1-6}alkyl, e.g. $–NH-CH_2-CH_2-F$, by reaction with polyhaloC_{1-6}alkyl-W, with W as defined above, e.g. iodo-2-fluoroethane, in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example N,N-dimethylformamide or dioxane.

Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with cyano, can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted

with tetrazolyl by reaction with sodium azide, and $NH_4^+Cl^-$ in the presence of a suitable solvent, such as for example N,N-dimethylformamide.

Compounds of formula (I) wherein R^3 represents -CH2-C≡CH, can be converted into a compound of formula (I) wherein R^3 represents , by reaction with ethyl azidoacetate in the presence of CuI and a suitable base, such as for example diisopropylamine, and a suitable solvent, such as for example tetraydrofuran.

Compounds of formula (I) wherein R^3 represents -CH2-C≡CH, can be converted into a compound of formula (I) wherein R^3 represents  by reaction with sodium azide and formaldehyde, in the presence of a suitable catalyst, such as for example $CuSO_4$ and sodium L ascorbate, a suitable acid, such as for example acetic acid, and a suitable solvent, such as for example dioxane.

Compounds of formula (I) wherein R^3 represent C_{2-6}alkynyl, can be converted into a compound of formula (I) wherein R^3 represents C_{2-6}alkynyl substituted with R^9, by reaction with W-R^9 wherein W is as defined above, in the presence of a suitable catalyst, such as for example dichlorobis(triphenylphosphine)palladium, a suitable co-catalyst such as CuI, a suitable base, such as for example triethylamine, and a suitable solvent, such as for example dimethylsulfoxide.

Compounds of formula (I) wherein R^3 comprises R^9 substituted with halo, can be converted into a compound of formula (I) wherein R^3 comprises R^9 substituted with -$NR^{14}R^{15}$ by reaction with $NHR^{14}R^{15}$ in the presence of a suitable solvent, such as for example 1-methyl-2-pyrrolidinone.

Compounds of formula (I) wherein R^3 comprises C_{2-6}alkynyl, can be hydrogenated into a compound of formula (I) wherein R^3 comprises C_{2-6}alkyl in the presence of a suitable catalyst, such as for example palladium on charcoal, and a suitable solvent, such as for example ethylacetate.

Compounds of formula (I) wherein R^3 comprises C_{2-6}alkynyl, can be hydrogenated into a compound of formula (I) wherein R^3 comprises C_{2-6}alkenyl in the presence of a suitable catalyst, such as for example Lindlar catalyst, and a suitable solvent, such as for example ethylacetate.

Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with -P(=O)(OC_{1-6}alkyl)$_2$ can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl

substituted with -P(=O)(OH)$_2$ by reaction with bromotrimethylsilane in the presence of a suitable solvent, such as for example dichloromethane.

Compounds of formula (I) wherein the R^9 substituent is substituted with =O, can be converted into the corresponding reduced R^9 substituent by reaction with a suitable reducing agent, such as for example $LiAlH_4$ in a suitable solvent, such as for example tetrahydrofuran.

Compounds of formula (I) wherein R^3 comprises -NHR^{10} can be converted into a compound of formula (I) wherein R^3 comprises -NR^{10}-(C=O)-optionally substituted C_{1-6}alkyl, by reaction with the corresponding W-(C=O)-optionally substituted C_{1-6}alkyl wherein W represents a suitable leaving group, such as for example halo, e.g. chloro and the like, in the presence of a suitable base, such as for example triethylamine, and a suitable solvent, such as for example acetonitrile.

Compounds of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with NR^{10}(benzyl) can be converted into a compound of formula (I) wherein R^3 represents C_{1-6}alkyl substituted with NHR^{10}, by reaction with 1-chloroethylchloroformate in the presence of a suitable solvent, such as for example dichloromethane

Compounds of formula (I) wherein R^1 represents unsubstituted piperidine, can be converted into a compound of formula (I) wherein R^1 represents 1-methyl-piperidine, by reaction with iodomethane in the presence of a suitable base, such as for example potassium carbonate, and a suitable solvent, such as for example acetonitrile.

Compounds of formula (I) wherein R^1 represents hydrogen can be converted into a compound of formula (I) wherein R^1 represents optionally substituted C_{1-6}alkyl, by reaction with optionally substituted C_{1-6}alkyl-W wherein W represents a suitable leaving group, such as for example halo, e.g. bromo and the like, in the presence of a suitable base, such as for example potassium carbonate, and a suitable solvent, such as for example acetonitrile.


Compounds of formula (I) wherein R^2 represents halo, e.g. bromo, can be converted into a compound of formula (I) wherein R^2 represents cyano, by reaction with zinc cyanide, in the presence of a suitable catalyst, such as for example $Pd_2(dba)_3$ and a suitable ligand, such as for example 1,1-bis(diphenylphosphino)ferrocene, in the presence of a suitable solvent, such as for example N,N-dimethylformamide.

Said R^2 substituent being cyano can be converted into –CH_2-NH_2 by hydrogenation in the presence of NH_3 and Nickel.

Compounds of formula (I) wherein R^2 represents $-OCH_3$ can be converted into a compounds of formula (I) wherein R^2 represents –OH by reaction with boron tribromide in the presence of a suitable solvent, such as for example dichloromethane.
Compounds of formula (I) wherein R^2 represents –OH can be converted into a compounds of formula (I) wherein R^2 represents $-OCH_3$ by reaction with methyl iodine in the presence of a suitable base, such as for example potassium carbonate, and a suitable solvent, such as for example N,N-dimethylformamide.
Compounds of formula (I) wherein R^2 represents hydrogen, can be converted into a compound of formula (I) wherein R^2 represents $-CHOH-CF_3$ by reaction with trifluoroacetaldehyde methyl hemiketal.


A further aspect of the invention is a process for the preparation of a compound of formula (I) as defined herein, which process comprises:

(i) deprotecting a compound of formula (XXX) wherein P represents a suitable protective group, such as for example a butyloxycarbonyl-group ($-CO_2C(CH_3)_3$) in the presence of a suitable acid, such as for example HCl or trifluoroacetic acid;



(XXX) or

(ii) the reaction of a compound of the formula (IX) or (IX'):




(IX) : R^u is $-O-(S=O)_2-CH_3$

(IX') : R^u is Cl

or a protected form thereof, with an appropriately substituted amine or a reactive derivative thereof, such as for example $NHR^{10}R^{11}$ (X), $NHR^{10}P$ (X-a) or  (XXI),

for example in a sealed vessel, in the presence of a suitable base, such as for example sodium hydride and/or in the presence or absence of a solvent such as acetonitrile, N,N-dimethylformamide or N,N-dimethylacetamide; or


(iii) the reaction of a compound of the formula (VI):



(VI)

or a protected form thereof, with a compound of formula W_6-C_{1-6}alkyl-$NR^{10}P$ wherein P represents a suitable protective group and W_6 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, or –O-S(=O)$_2$-CH_3, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, e.g. N,N-dimethylformamide or N,N-dimethylacetamide, followed by removing P and optionally removing any further protecting group present; or


(iv) the reaction of a compound of the formula (VI):



(VI)

or a protected thereof, with a compound of formula W_6-C_{1-6}alkyl-NHR^{10} wherein W_6 represents a suitable leaving group, such as for example halo, e.g. bromo and the like, or –O-S(=O)$_2$-CH_3, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, e.g. N,N-dimethylformamide or N,N-dimethylacetamide;


(v) the reaction of a compound of formula (XXXVI)



(XXXVI)

with hydrazine in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol;


(vi) the reaction of a compound of formula (IX-1) wherein R^u represents $–O-S(=O)_2-CH_3$,



(IX-1) (X)

with an intermediate of formula (X) in the presence of a suitable solvent, such as for example acetonitrile;


(vii) the reaction of a compound of formula (VI)



(VI)

with an intermediate of formula $W_{11}-R^{3b}$ wherein R^{3b} represents optionally substituted C_{2-6}alkynyl and W_{11} represents a suitable leaving group such as for example halo, e.g. chloro, or $–O-S(=O)_2-CH_3$, in the presence of a suitable base, such as for example NaH, and a suitable solvent, such as for example N,N-dimethylformamide;


(viii) the reaction of a compound of formula (VIII') wherein R^x and R^y represent C_{1-4}alkyl, and R^z represent C_{1-4}alkyl or phenyl,



(VIII')

with a suitable acid, such as for example trifluoroacetic acid, in the presence of a suitable solvent, such as for example tetrahydrofuran;


(viii) deprotecting a compound of formula (XXXXII)



(XXXXII)

in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example an alcohol, e.g. methanol and the like;


(ix) the reaction of a compound of formula (VI)



(VI)

with di(C_{1-6}alkyl)vinylphosphonate in the presence of a suitable catalyst, such as for example tri-N-butylphosphine, and a suitable solvent, such as for example acetonitrile;


(x) deprotecting a compound of formula (XXXXI)



(XXXXI)

in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example an alcohol, e.g. methanol and the like;

(xi) the reaction of a compound of formula (XIX) with a compound of formula (III)




(XIX)


(III)

in the presence of a suitable catalyst, such as for example tetrakis(triphenyl)phosphine palladium or $Pd_2(dba)_3$ (tris(dibenzylideneacetone) dipalladium (0)), a suitable ligand, such as 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl, a suitable base, such as for example Na_2CO_3 or K_3PO_4, and a suitable solvent, such as for example ethylene glycol dimethylether or dioxane or water;

(xii) the reaction of a compound of formula (XX) wherein R^{3a} represents optionally substituted C_{1-6}alkyl, with a compound of formula (XIV)



(XX)




(XIV)

in the presence of a suitable catalyst, such as for example palladium (II) acetate or $Pd_2(dba)_3$ (tris(dibenzylidene acetone) dipalladium (0)), a suitable ligand such as for example 2-dicyclohexylphosphino-tris-isopropyl-biphenyl or 1,1'-[1,1'-binaphthalene]-

2,2'-diylbis[1,1-diphenylphosphine], a suitable base, such as for example sodium tert-butoxide, and a suitable solvent, such as for example ethylene glycol dimethylether;


(xiii) the reaction of a compound of formula (XXXI)



(XXXI)


with W_8-CN, wherein W_8 represents a suitable leaving group, such as for example halo, e.g. bromo, in the presence of a suitable base, such as for example $NaHCO_3$, and a suitable solvent, such as for example water or dioxane;


(xiv) the reaction of a compound of formula (XXXV)



(XXXV)

with a suitable base, such as for example *N,N*-diisopropylethylamine and triethylamine, in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol;


(xv) deprotecting a compound of formula (XXVI) wherein P represents a suitable protective group such as for example –O-Si$(CH_3)_2(C(CH_3)_3)$ or ,



(XXVI)

in the presence of a suitable acid, such as for example HCl or trifluoroacetic acid, or a suitable de-silylating agent, such as for example tetrabutyl ammonium fluoride, and a suitable solvent, such as an alcohol, e.g. methanol, or tetrahydrofuran;


(xvi) the reaction of a compound of formula (XXIX) with a compound of formula (XXI)



(XXIX)

in the presence of suitable peptide coupling reagents such as, 1-hydroxy-benzotriazole and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide HCl;


(xvii) the reaction of a compound of formula (XXIX)



(XXIX)

with NHR^4R^5 in the presence of suitable peptide coupling reagents such as 1-hydroxy-benzotriazole and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide HCl and a suitable base, such as triethylamine, and a suitable solvent, such as for example dichloromethane;


(xviii) reacting the below compound



:

with NHR^7R^8 in the presence of a suitable base, such as for example K_2CO_3, and a suitable solvent, such as for example tetrahydrofuran;


(xviii) deprotecting the below compound



in the presence of hydrazine monohydrate, and a suitable solvent, such as for example an alcohol, e.g. ethanol;

wherein R^1, R^{1a}, R^2, R^{10}, and n are as defined herein; and optionally thereafter converting one compound of the formula (I) into another compound of the formula (I).

A further embodiment is a process for synthesis of a compound of formula (VI) wherein:



1) a compound of formula (II) is reacted with an intermediate of formula (III) in the presence of a suitable catalyst, such as for example tetrakis(triphenylphosphine)palladium (0) or palladium (II) acetate , a suitable base, such as for example sodium carbonate, a suitable ligand, such as for example triphenylphosphine, and a suitable solvent or solvent mixture, such as for example ethylene glycol dimethylether and water; wherein W_1 and W_2, each independently represent a suitable leaving group, such as for example halo, e.g. chloro or bromo;

and then

2) a compound of formula (IV) is reacted with an intermediate of formula (V) in the presence of a suitable catalyst, such as for example palladium (II) acetate, a suitable base, such as sodium tert-butoxide or Cs_2CO_3, a suitable ligand, such as for example 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine], and a suitable solvent or solvent mixture, such as for example dioxane or ethylene glycol dimethylether and water;

wherein optionally the intermediate of formula (II) wherein W_1 is chloro and W_2 is bromo is prepared by reacting 7-bromo-2(1H)-quinoxalinone with phosphorus oxychloride, or alternatively with thionyl chloride and N,N-dimethylformamide in a suitable solvent, such as, for example toluene;

or vice versa, wherein a compound of formula (II) is reacted with an intermediate of formula (V) first and then reacted with an intermediate of formula (III) using the methods described above.

In a further embodiment the invention provides a novel intermediate. In one embodiment the invention provides a novel intermediate of formula (II)-(XXXI). In another embodiment the invention provides a novel intermediate of formula (VI) or formula (IX). In another embodiment the invention provides a compound of formula (I-a)-(I-i).

Pharmaceutically Acceptable Salts, Solvates or Derivatives thereof

In this section, as in all other sections of this application, unless the context indicates otherwise, references to formula (I) include references to all other sub-groups, preferences, embodiments and examples thereof as defined herein.

Unless otherwise specified, a reference to a particular compound also includes ionic forms, salts, solvates, isomers, tautomers, N-oxides, esters, prodrugs, isotopes and protected forms thereof, for example, as discussed below; preferably, the ionic forms, or salts or tautomers or isomers or N-oxides or solvates thereof; and more preferably, the ionic forms, or salts or tautomers or solvates or protected forms thereof, even more preferably the salts or tautomers or solvates thereof. Many compounds of the formula (I) can exist in the form of salts, for example acid addition salts or, in certain cases salts of organic and inorganic bases such as carboxylate, sulphonate and phosphate salts. All such salts are within the scope of this invention, and references to compounds of the formula (I) include the salt forms of the compounds. It will be appreciated that references to “derivatives” include references to ionic forms, salts, solvates, isomers, tautomers, N-oxides, esters, prodrugs, isotopes and protected forms thereof.

According to one aspect of the invention there is provided a compound as defined herein or a salt, tautomer, N-oxide or solvate thereof. According to a further aspect of the invention there is provided a compound as defined herein or a salt or solvate thereof. References to compounds of the formula (I) and sub-groups thereof as defined herein include within their scope the salts or solvates or tautomers or N-oxides of the compounds.

The salt forms of the compounds of the invention are typically pharmaceutically acceptable salts, and examples of pharmaceutically acceptable salts are discussed in Berge *et al*. (1977) "Pharmaceutically Acceptable Salts," *J. Pharm. Sci.,* Vol. 66, pp. 1-19. However, salts that are not pharmaceutically acceptable may also be prepared as intermediate forms which may then be converted into pharmaceutically acceptable salts. Such non-pharmaceutically acceptable salts forms, which may be useful, for example, in the purification or separation of the compounds of the invention, also form part of the invention.

The salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods such as methods described in *Pharmaceutical Salts: Properties, Selection, and Use*, P. Heinrich Stahl (Editor), Camille G. Wermuth (Editor), ISBN: 3-90639-026-8, Hardcover, 388 pages, August 2002. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media such as ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are used. The compounds of the invention may exist as mono- or di-salts depending upon the pKa of the acid from which the salt is formed.

Acid addition salts may be formed with a wide variety of acids, both inorganic and organic. Examples of acid addition salts include salts formed with an acid selected from the group consisting of acetic, 2,2-dichloroacetic, adipic, alginic, ascorbic (e.g. L-ascorbic), L-aspartic, benzenesulphonic, benzoic, 4-acetamidobenzoic, butanoic, (+) camphoric, camphor-sulphonic, (+)-(1*S*)-camphor-10-sulphonic, capric, caproic, caprylic, cinnamic, citric, cyclamic, dodecylsulphuric, ethane-1,2-disulphonic, ethanesulphonic, 2-hydroxyethanesulphonic, formic, fumaric, galactaric, gentisic, glucoheptonic, D-gluconic, glucuronic (e.g. D-glucuronic), glutamic (e.g. L-glutamic), α-oxoglutaric, glycolic, hippuric, hydrobromic, hydrochloric, hydriodic, isethionic, lactic (e.g. (+)-L-lactic, (±)-DL-lactic), lactobionic, maleic, malic, (-)-L-malic, malonic, (±)-DL-mandelic, methanesulphonic, naphthalenesulphonic (e.g.naphthalene-2-sulphonic), naphthalene-1,5-disulphonic, 1-hydroxy-2-naphthoic, nicotinic, nitric, oleic, orotic, oxalic, palmitic, pamoic, phosphoric, propionic, L-pyroglutamic, pyruvic, salicylic, 4-amino-salicylic, sebacic, stearic, succinic, sulphuric, tannic, (+)-L-tartaric, thiocyanic, toluenesulphonic (e.g. *p*-toluenesulphonic), undecylenic and valeric acids, as well as acylated amino acids and cation exchange resins.

One particular group of salts consists of salts formed from acetic, hydrochloric, hydriodic, phosphoric, nitric, sulphuric, citric, lactic, succinic, maleic, malic, isethionic, fumaric, benzenesulphonic, toluenesulphonic, methanesulphonic (mesylate), ethanesulphonic, naphthalenesulphonic, valeric, acetic, propanoic, butanoic, malonic, glucuronic and lactobionic acids. Another group of acid addition salts includes salts formed from acetic, adipic, ascorbic, aspartic, citric, DL-Lactic, fumaric, gluconic,

glucuronic, hippuric, hydrochloric, glutamic, DL-malic, methanesulphonic, sebacic, stearic, succinic and tartaric acids.

If the compound is anionic, or has a functional group which may be anionic (e.g., -COOH may be $-COO^-$), then a salt may be formed with a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as Na^+ and K^+, alkaline earth metal cations such as Ca^{2+} and Mg^{2+}, and other cations such as Al^{3+}. Examples of suitable organic cations include, but are not limited to, ammonium ion (i.e., NH_4^+) and substituted ammonium ions (e.g., NH_3R^+, $NH_2R_2^+$, NHR_3^+, NR_4^+).

Examples of some suitable substituted ammonium ions are those derived from: ethylamine, diethylamine, dicyclohexylamine, triethylamine, butylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino acids, such as lysine and arginine. An example of a common quaternary ammonium ion is $N(CH_3)_4^+$.


Where the compounds of the formula (I) contain an amine function, these may form quaternary ammonium salts, for example by reaction with an alkylating agent according to methods well known to the skilled person. Such quaternary ammonium compounds are within the scope of formula (I). Compounds of the formula (I) containing an amine function may also form N-oxides. A reference herein to a compound of the formula (I) that contains an amine function also includes the N-oxide. Where a compound contains several amine functions, one or more than one nitrogen atom may be oxidised to form an N-oxide. Particular examples of N-oxides are the N-oxides of a tertiary amine or a nitrogen atom of a nitrogen-containing heterocycle. N-Oxides can be formed by treatment of the corresponding amine with an oxidizing agent such as hydrogen peroxide or a per-acid (e.g. a peroxycarboxylic acid), see for example *Advanced Organic Chemistry*, by Jerry March, 4th Edition, Wiley Interscience, pages. More particularly, N-oxides can be made by the procedure of L. W. Deady (*Syn. Comm.* (1977), **7**, 509-514) in which the amine compound is reacted with *m*-chloroperoxybenzoic acid (MCPBA), for example, in an inert solvent such as dichloromethane.

The compounds of the invention may form solvates, for example with water (i.e., hydrates) or common organic solvents. As used herein, the term "solvate" means a physical association of the compounds of the present invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. The term "solvate" is intended to encompass both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include compounds of the invention in combination with water, isopropanol, ethanol, methanol, DMSO, ethyl acetate, acetic acid or ethanolamine and the like. The compounds of the invention may exert their biological effects whilst they are in solution.

Solvates are well known in pharmaceutical chemistry. They can be important to the processes for the preparation of a substance (e.g. in relation to their purification, the storage of the substance (e.g. its stability) and the ease of handling of the substance and are often formed as part of the isolation or purification stages of a chemical synthesis. A person skilled in the art can determine by means of standard and long used techniques whether a hydrate or other solvate has formed by the isolation conditions or purification conditions used to prepare a given compound. Examples of such techniques include thermogravimetric analysis (TGA), differential scanning calorimetry (DSC), X-ray crystallography (e.g. single crystal X-ray crystallography or X-ray powder diffraction) and Solid State NMR (SS-NMR, also known as Magic Angle Spinning NMR or MAS-NMR). Such techniques are as much a part of the standard analytical toolkit of the skilled chemist as NMR, IR, HPLC and MS. Alternatively the skilled person can deliberately form a solvate using crystallisation conditions that include an amount of the solvent required for the particular solvate. Thereafter the standard methods described above, can be used to establish whether solvates had formed. Also encompassed by formula (I) are any complexes (e.g. inclusion complexes or clathrates with compounds such as cyclodextrins, or complexes with metals) of the compounds.

Furthermore, the compounds of the present invention may have one or more polymorph (crystalline) or amorphous forms and as such are intended to be included in the scope of the invention.

Compounds of the formula (I) may exist in a number of different geometric isomeric, and tautomeric forms and references to compounds of the formula (I) include all such forms. For the avoidance of doubt, where a compound can exist in one of several geometric isomeric or tautomeric forms and only one is specifically described or shown, all others are nevertheless embraced by formula (I). Other examples of tautomeric forms include, for example, keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/enediamines, nitroso/oxime, thioketone/enethiol, and nitro/aci-nitro.



keto enol enolate

Where compounds of the formula (I) contain one or more chiral centres, and can exist in the form of two or more optical isomers, references to compounds of the formula (I) include all optical isomeric forms thereof (e.g. enantiomers, epimers and diastereoisomers), either as individual optical isomers, or mixtures (e.g. racemic mixtures) of two or more optical isomers, unless the context requires otherwise. The optical isomers may be characterised and identified by their optical activity (i.e. as + and – isomers, or *d* and *l* isomers) or they may be characterised in terms of their absolute stereochemistry using the "R and S" nomenclature developed by Cahn, Ingold and Prelog, see *Advanced Organic Chemistry* by Jerry March, 4th Edition, John Wiley & Sons, New York, 1992, pages 109-114, and see also Cahn, Ingold & Prelog (1966) *Angew. Chem. Int. Ed. Engl.*, **5**, 385-415. Optical isomers can be separated by a number of techniques including chiral chromatography (chromatography on a chiral support) and such techniques are well known to the person skilled in the art. As an alternative to chiral chromatography, optical isomers can be separated by forming diastereoisomeric salts with chiral acids such as (+)-tartaric acid, (-)-pyroglutamic acid, (-)-di-toluoyl-L-tartaric acid, (+)-mandelic acid, (-)-malic acid, and (-)-camphorsulphonic, separating the diastereoisomers by preferential crystallisation, and then dissociating the salts to give the individual enantiomer of the free base.

Where compounds of the formula (I) exist as two or more optical isomeric forms, one enantiomer in a pair of enantiomers may exhibit advantages over the other enantiomer, for example, in terms of biological activity. Thus, in certain circumstances, it may be desirable to use as a therapeutic agent only one of a pair of enantiomers, or only one of

a plurality of diastereoisomers. Accordingly, the invention provides compositions containing a compound of the formula (I) having one or more chiral centres, wherein at least 55% (e.g. at least 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95%) of the compound of the formula (I) is present as a single optical isomer (e.g. enantiomer or diastereoisomer). In one general embodiment, 99% or more (e.g. substantially all) of the total amount of the compound of the formula (I) may be present as a single optical isomer (e.g. enantiomer or diastereoisomer).

The compounds of the invention include compounds with one or more isotopic substitutions, and a reference to a particular element includes within its scope all isotopes of the element. For example, a reference to hydrogen includes within its scope ^{1}H, ^{2}H (D), and ^{3}H (T). Similarly, references to carbon and oxygen include within their scope respectively ^{12}C, ^{13}C and ^{14}C and ^{16}O and ^{18}O. The isotopes may be radioactive or non-radioactive. In one embodiment of the invention, the compounds contain no radioactive isotopes. Such compounds are preferred for therapeutic use. In another embodiment, however, the compound may contain one or more radioisotopes. Compounds containing such radioisotopes may be useful in a diagnostic context.

Esters such as carboxylic acid esters and acyloxy esters of the compounds of formula (I) bearing a carboxylic acid group or a hydroxyl group are also embraced by formula (I). In one embodiment of the invention, formula (I) includes within its scope esters of compounds of the formula (I) bearing a carboxylic acid group or a hydroxyl group. In another embodiment of the invention, formula (I) does not include within its scope esters of compounds of the formula (I) bearing a carboxylic acid group or a hydroxyl group. Examples of esters are compounds containing the group -C(=O)OR, wherein R is an ester substituent, for example, a C_{1-6} alkyl group, a heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-6} alkyl group. Particular examples of ester groups include, but are not limited to, $-C(=O)OCH_3$, $-C(=O)OCH_2CH_3$, $-C(=O)OC(CH_3)_3$, and -C(=O)OPh. Examples of acyloxy (reverse ester) groups are represented by -OC(=O)R, wherein R is an acyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Particular examples of acyloxy groups include, but are not limited to, $-OC(=O)CH_3$ (acetoxy), $-OC(=O)CH_2CH_3$, $-OC(=O)C(CH_3)_3$, -OC(=O)Ph, and $-OC(=O)CH_2Ph$.

For example, some prodrugs are esters of the active compound (e.g., a physiologically acceptable metabolically labile ester). By "prodrugs" is meant for example any compound that is converted *in vivo* into a biologically active compound of the formula (I). During metabolism, the ester group (-C(=O)OR) is cleaved to yield the active drug. Such esters may be formed by esterification, for example, of any of the carboxylic acid groups (-C(=O)OH) in the parent compound, with, where appropriate, prior protection of any other reactive groups present in the parent compound, followed by deprotection if required.

Examples of such metabolically labile esters include those of the formula -C(=O)OR wherein R is: C_{1-6}alkyl (e.g., -Me, -Et, -nPr, -iPr, -nBu, -sBu, -iBu, -tBu); C_{1-6}aminoalkyl [e.g., aminoethyl; 2-(N,N-diethylamino)ethyl; 2-(4-morpholino)ethyl); and acyloxy-C_{1-7}alkyl [e.g., acyloxymethyl; acyloxyethyl; pivaloyloxymethyl; acetoxymethyl; 1-acetoxyethyl; 1-(1-methoxy-1-methyl)ethyl-carbonyloxyethyl; 1-(benzoyloxy)ethyl; isopropoxy-carbonyloxymethyl; 1-isopropoxy-carbonyloxyethyl; cyclohexyl-carbonyloxymethyl; 1-cyclohexyl-carbonyloxyethyl; cyclohexyloxy-carbonyloxymethyl; 1-cyclohexyloxy-carbonyloxyethyl; (4-tetrahydropyranyloxy) carbonyloxymethyl; 1-(4-tetrahydropyranyloxy)carbonyloxyethyl; (4-tetrahydropyranyl)carbonyloxymethyl; and 1-(4-tetrahydropyranyl)carbonyloxyethyl]. Also, some prodrugs are activated enzymatically to yield the active compound, or a compound which, upon further chemical reaction, yields the active compound (for example, as in antigen-directed enzyme pro-drug therapy (ADEPT), gene-directed enzyme pro-drug therapy (GDEPT) and ligand-directed enzyme pro-drug therapy (LIDEPT) etc.). For example, the prodrug may be a sugar derivative or other glycoside conjugate, or may be an amino acid ester derivative.

Protein Tyrosine Kinases (PTK)

The compounds of the invention described herein inhibit or modulate the activity of certain tyrosine kinases, and thus the compounds will be useful in the treatment or prophylaxis, in particular the treatment, of disease states or conditions mediated by those tyrosine kinases, in particular FGFR.

FGFR

The fibroblast growth factor (FGF) family of protein tyrosine kinase (PTK) receptors regulates a diverse array of physiologic functions including mitogenesis, wound healing, cell differentiation and angiogenesis, and development. Both normal and malignant cell growth as well as proliferation are affected by changes in local concentration of FGFs, extracellular signalling molecules which act as autocrine as well as paracrine factors. Autocrine FGF signalling may be particularly important in the progression of steroid hormone-dependent cancers to a hormone independent state. FGFs and their receptors are expressed at increased levels in several tissues and cell lines and overexpression is believed to contribute to the malignant phenotype. Furthermore, a number of oncogenes are homologues of genes encoding growth factor receptors, and there is a potential for aberrant activation of FGF-dependent signalling in human pancreatic cancer (Knights et al., Pharmacology and Therapeutics 2010 125:1 (105-117); Korc M. et al Current Cancer Drug Targets 2009 9:5 (639-651)).

The two prototypic members are acidic fibroblast growth factor (aFGF or FGF1) and basic fibroblast growth factor (bFGF or FGF2), and to date, at least twenty distinct FGF family members have been identified. The cellular response to FGFs is transmitted via four types of high affinity transmembrane protein tyrosine-kinase fibroblast growth factor receptors (FGFR) numbered 1 to 4 (FGFR1 to FGFR4).

Disruption of the FGFR1 pathway should affect tumor cell proliferation since this kinase is activated in many tumor types in addition to proliferating endothelial cells. The over-expression and activation of FGFR1 in tumor- associated vasculature has suggested a role for these molecules in tumor angiogenesis.

A recent study has shown a link between FGFR1 expression and tumorigenicity in Classic Lobular Carcinomas (CLC). CLCs account for 10-15% of all breast cancers and, in general, lack p53 and Her2 expression whilst retaining expression of the oestrogen receptor. A gene amplification of 8p12-p11.2 was demonstrated in ~50% of CLC cases and this was shown to be linked with an increased expression of FGFR1. Preliminary studies with siRNA directed against FGFR1, or a small molecule inhibitor of the receptor, showed cell lines harbouring this amplification to be particularly sensitive to inhibition of this signalling pathway. Rhabdomyosarcoma (RMS) is the most common pediatric soft tissue sarcoma likely results from abnormal proliferation and differentiation during skeletal myogenesis. FGFR1 is over-expressed in primary rhabdomyosarcoma

tumors and is associated with hypomethylation of a 5' CpG island and abnormal expression of the AKT1, NOG, and BMP4 genes.

Fibroblast growth factor receptor 2 has high affinity for the acidic and/or basic fibroblast growth factors, as well as the keratinocyte growth factor ligands. Fibroblast growth factor receptor 2 also propagates the potent osteogenic effects of FGFs during osteoblast growth and differentiation. Mutations in fibroblast growth factor receptor 2, leading to complex functional alterations, were shown to induce abnormal ossification of cranial sutures (craniosynostosis), implying a major role of FGFR signalling in intramembranous bone formation. For example, in Apert (AP) syndrome, characterized by premature cranial suture ossification, most cases are associated with point mutations engendering gain-of-function in fibroblast growth factor receptor 2. In addition, mutation screening in patients with syndromic craniosynostoses indicates that a number of recurrent *FGFR2* mutations accounts for severe forms of Pfeiffer syndrome. Particular mutations of FGFR2 include W290C, D321A, Y340C, C342R, C342S, C342W, N549H, K641R in FGFR2.

Several severe abnormalities in human skeletal development, including Apert, Crouzon, Jackson-Weiss, Beare-Stevenson cutis gyrata, and Pfeiffer syndromes are associated with the occurrence of mutations in fibroblast growth factor receptor 2. Most, if not all, cases of Pfeiffer Syndrome (PS) are also caused by de novo mutation of the fibroblast growth factor receptor 2 gene, and it was recently shown that mutations in fibroblast growth factor receptor 2 break one of the cardinal rules governing ligand specificity. Namely, two mutant splice forms of fibroblast growth factor receptor, FGFR2c and FGFR2b, have acquired the ability to bind to and be activated by atypical FGF ligands. This loss of ligand specificity leads to aberrant signalling and suggests that the severe phenotypes of these disease syndromes result from ectopic ligand-dependent activation of fibroblast growth factor receptor 2.

Genetic aberrations of the FGFR3 receptor tyrosine kinase such as chromosomal translocations or point mutations result in ectopically expressed or deregulated, constitutively active, FGFR3 receptors. Such abnormalities are linked to a subset of multiple myelomas and in bladder, hepatocellular, oral squamous cell carcinoma and cervical carcinomas. Accordingly, FGFR3 inhibitors would be useful in the treatment of

multiple myeloma, bladder and cervical carcinomas. FGFR3 is also over-expressed in bladder cancer, in particular invasive bladder cancer. FGFR3 is frequently activated by mutation in urothelial carcinoma (UC). Increased expression was associated with mutation (85% of mutant tumors showed high-level expression) but also 42% of tumors with no detectable mutation showed over-expression, including many muscle-invasive tumors.

Over expression of FGFR4 has been linked to poor prognosis in both prostate and thyroid carcinomas. In addition a germline polymorphism (Gly388Arg) is associated with increased incidence of lung, breast, colon, liver (HCC) and prostate cancers. In addition, a truncated form of FGFR4 (including the kinase domain) has also been found to be present in 40% of pituitary tumours but not present in normal tissue. FGFR4 overexpression has been observed in liver, colon and lung tumours. FGFR4 has been implicated in colorectal and liver cancer where expression of its ligand FGF19 is frequently elevated.

Fibrotic conditions are a major medical problem resulting from abnormal or excessive deposition of fibrous tissue. This occurs in many diseases, including liver cirrhosis, glomerulonephritis, pulmonary fibrosis, systemic fibrosis, rheumatoid arthritis, as well as the natural process of wound healing. The mechanisms of pathological fibrosis are not fully understood but are thought to result from the actions of various cytokines (including tumor necrosis factor (TNF), fibroblast growth factors (FGF's), platelet derived growth factor (PDGF) and transforming growth factor beta. (TGFβ) involved in the proliferation of fibroblasts and the deposition of extracellular matrix proteins (including collagen and fibronectin). This results in alteration of tissue structure and function and subsequent pathology.

A number of preclinical studies have demonstrated the up-regulation of fibroblast growth factors in preclinical models of lung fibrosis. TGFβ1 and PDGF have been reported to be involved in the fibrogenic process and further published work suggests the elevation of FGF's and consequent increase in fibroblast proliferation, may be in response to elevated TGFβ1. The potential therapeutic benefit of targeting the fibrotic mechanism in conditions such as idiopathic pulmonary fibrosis (IPF) is suggested by the reported clinical effect of the anti-fibrotic agent pirfenidone . Idiopathic pulmonary fibrosis (also

referred to as Cryptogenic fibrosing alveolitis) is a progressive condition involving scarring of the lung. Gradually, the air sacs of the lungs become replaced by fibrotic tissue, which becomes thicker, causing an irreversible loss of the tissue's ability to transfer oxygen into the bloodstream. The symptoms of the condition include shortness of breath, chronic dry coughing, fatigue, chest pain and loss of appetite resulting in rapid weight loss. The condition is extremely serious with approximately 50% mortality after 5 years.

As such, the compounds which inhibit FGFR will be useful in providing a means of preventing the growth or inducing apoptosis in tumours, particularly by inhibiting angiogenesis. It is therefore anticipated that the compounds will prove useful in treating or preventing proliferative disorders such as cancers. In particular tumours with activating mutants of receptor tyrosine kinases or upregulation of receptor tyrosine kinases may be particularly sensitive to the inhibitors. Patients with activating mutants of any of the isoforms of the specific RTKs discussed herein may also find treatment with RTK inhibitors particularly beneficial.

Vascular Endothelial Growth Factor (VEGFR)

Chronic proliferative diseases are often accompanied by profound angiogenesis, which can contribute to or maintain an inflammatory and/or proliferative state, or which leads to tissue destruction through the invasive proliferation of blood vessels. .

Angiogenesis is generally used to describe the development of new or replacement blood vessels, or neovascularisation. It is a necessary and physiological normal process by which vasculature is established in the embryo. Angiogenesis does not occur, in general, in most normal adult tissues, exceptions being sites of ovulation, menses and wound healing. Many diseases, however, are characterized by persistent and unregulated angiogenesis. For instance, in arthritis, new capillary blood vessels invade the joint and destroy cartilage. In diabetes (and in many different eye diseases), new vessels invade the macula or retina or other ocular structures, and may cause blindness. The process of atherosclerosis has been linked to angiogenesis. Tumor growth and metastasis have been found to be angiogenesis-dependent.

The recognition of the involvement of angiogenesis in major diseases has been

accompanied by research to identify and develop inhibitors of angiogenesis. These inhibitors are generally classified in response to discrete targets in the angiogenesis cascade, such as activation of endothelial cells by an angiogenic signal; synthesis and release of degradative enzymes; endothelial cell migration; proliferation of endothelial cells; and formation of capillary tubules. Therefore, angiogenesis occurs in many stages and attempts are underway to discover and develop compounds that work to block angiogenesis at these various stages.

There are publications that teach that inhibitors of angiogenesis, working by diverse mechanisms, are beneficial in diseases such as cancer and metastasis, ocular diseases, arthritis and hemangioma.

Vascular endothelial growth factor (VEGF), a polypeptide, is mitogenic for endothelial cells *in vitro* and stimulates angiogenic responses *in vivo*. VEGF has also been linked to inappropriate angiogenesis. VEGFR(s) are protein tyrosine kinases (PTKs). PTKs catalyze the phosphorylation of specific tyrosine residues in proteins involved in cell function thus regulating cell growth, survival and differentiation.

Three PTK receptors for VEGF have been identified: VEGFR-1 (Flt-1) ; VEGFR-2 (Flk-1 or KDR) and VEGFR-3 (Flt-4). These receptors are involved in angiogenesis and participate in signal transduction. Of particular interest is VEGFR-2, which is a transmembrane receptor PTK expressed primarily in endothelial cells. Activation of VEGFR-2 by VEGF is a critical step in the signal transduction pathway that initiates tumour angiogenesis. VEGF expression may be constitutive to tumour cells and can also be upregulated in response to certain stimuli. One such stimuli is hypoxia, where VEGF expression is upregulated in both tumour and associated host tissues. The VEGF ligand activates VEGFR-2 by binding with its extracellular VEGF binding site. This leads to receptor dimerization of VEGFRs and autophosphorylation of tyrosine residues at the intracellular kinase domain of VEGFR- 2. The kinase domain operates to transfer a phosphate from ATP to the tyrosine residues, thus providing binding sites for signalling proteins downstream of VEGFR-2 leading ultimately to initiation of angiogenesis.

Inhibition at the kinase domain binding site of VEGFR-2 would block phosphorylation of tyrosine residues and serve to disrupt initiation of angiogenesis.

Angiogenesis is a physiologic process of new blood vessel formation mediated by various cytokines called angiogenic factors. Although its potential pathophysiologic role in solid tumors has been extensively studied for more than 3 decades, enhancement of angiogenesis in chronic lymphocytic leukemia (CLL) and other malignant hematological disorders has been recognized more recently. An increased level of angiogenesis has been documented by various experimental methods both in bone marrow and lymph nodes of patients with CLL. Although the role of angiogenesis in the pathophysiology of this disease remains to be fully elucidated, experimental data suggest that several angiogenic factors play a role in the disease progression. Biologic markers of angiogenesis were also shown to be of prognostic relevance in CLL. This indicates that VEGFR inhibitors may also be of benefit for patients with leukemia's such as CLL.

In order for a tumour mass to get beyond a critical size, it must develop an associated vasculature. It has been proposed that targeting a tumor vasculature would limit tumor expansion and could be a useful cancer therapy. Observations of tumor growth have indicated that small tumour masses can persist in a tissue without any tumour-specific vasculature. The growth arrest of nonvascularized tumors has been attributed to the effects of hypoxia at the center of the tumor. More recently, a variety of proangiogenic and antiangiogenic factors have been identified and have led to the concept of the "angiogenic switch," a process in which disruption of the normal ratio of angiogenic stimuli and inhibitors in a tumor mass allows for autonomous vascularization. The angiogenic switch appears to be governed by the same genetic alterations that drive malignant conversion: the activation of oncogenes and the loss of tumour suppressor genes. Several growth factors act as positive regulators of angiogenesis. Foremost among these are vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), and angiogenin. Proteins such as thrombospondin (Tsp-1), angiostatin, and endostatin function as negative regulators of angiogenesis.

Inhibition of VEGFR2 but not VEGFR1 markedly disrupts angiogenic switching, persistent angiogenesis, and initial tumor growth in a mouse model. In late-stage tumors, phenotypic resistance to VEGFR2 blockade emerged, as tumors regrew during treatment after an initial period of growth suppression. This resistance to VEGF blockade involves reactivation of tumour angiogenesis, independent of VEGF and

associated with hypoxia-mediated induction of other proangiogenic factors, including members of the FGF family. These other proangiogenic signals are functionally implicated in the revascularization and regrowth of tumours in the evasion phase, as FGF blockade impairs progression in the face of VEGF inhibition.

There is evidence for normalization of glioblastoma blood vessels in patients treated with a pan-VEGF receptor tyrosine kinase inhibitor, AZD2171, in a phase 2 study. MRI determination of vessel normalization in combination with circulating biomarkers provides for an effective means to assess response to antiangiogenic agents.

PDGFR

A malignant tumour is the product of uncontrolled cell proliferation. Cell growth is controlled by a delicate balance between growth-promoting and growth-inhibiting factors. In normal tissue the production and activity of these factors results in differentiated cells growing in a controlled and regulated manner that maintains the normal integrity and functioning of the organ. The malignant cell has evaded this control; the natural balance is disturbed (via a variety of mechanisms) and unregulated, aberrant cell growth occurs. A growth factor of importance in tumour development is the platelet-derived growth factor (PDGF) that comprises a family of peptide growth factors that signal through cell surface tyrosine kinase receptors (PDGFR) and stimulate various cellular functions including growth, proliferation, and differentiation.

Advantages of a selective inhibitor

Development of FGFR kinase inhibitors with a differentiated selectivity profile provides a new opportunity to use these targeted agents in patient sub-groups whose disease is driven by FGFR deregulation. Compounds that exhibit reduced inhibitory action on additional kinases, particularly VEGFR2 and PDGFR-beta, offer the opportunity to have a differentiated side-effect or toxicity profile and as such allow for a more effective treatment of these indications. Inhibitors of VEGFR2 and PDGFR-beta are associated with toxicities such as hypertension or oedema respectively. In the case of VEGFR2 inhibitors this hypertensive effect is often dose limiting, may be contraindicated in certain patient populations and requires clinical management.

Biological Activity and Therapeutic Uses

The compounds of the invention, and subgroups thereof, have fibroblast growth factor receptor (FGFR) inhibiting or modulating activity and/or vascular endothelial growth factor receptor (VEGFR) inhibiting or modulating activity, and/or platelet derived growth factor receptor (PDGFR) inhibiting or modulating activity, and which will be useful in preventing or treating disease states or conditions described herein. In addition the compounds of the invention, and subgroups thereof, will be useful in preventing or treating diseases or condition mediated by the kinases. References to the preventing or prophylaxis or treatment of a disease state or condition such as cancer include within their scope alleviating or reducing the incidence of cancer.

As used herein, the term "modulation", as applied to the activity of a kinase, is intended to define a change in the level of biological activity of the protein kinase. Thus, modulation encompasses physiological changes which effect an increase or decrease in the relevant protein kinase activity. In the latter case, the modulation may be described as "inhibition". The modulation may arise directly or indirectly, and may be mediated by any mechanism and at any physiological level, including for example at the level of gene expression (including for example transcription, translation and/or post-translational modification), at the level of expression of genes encoding regulatory elements which act directly or indirectly on the levels of kinase activity. Thus, modulation may imply elevated/suppressed expression or over- or under-expression of a kinase, including gene amplification (i.e. multiple gene copies) and/or increased or decreased expression by a transcriptional effect, as well as hyper- (or hypo-)activity and (de)activation of the protein kinase(s) (including (de)activation) by mutation(s). The terms “modulated”, “modulating” and “modulate” are to be interpreted accordingly.

As used herein, the term “mediated”, as used e.g. in conjunction with a kinase as described herein (and applied for example to various physiological processes, diseases, states, conditions, therapies, treatments or interventions) is intended to operate limitatively so that the various processes, diseases, states, conditions, treatments and interventions to which the term is applied are those in which the kinase plays a biological role. In cases where the term is applied to a disease, state or condition, the biological role played by a kinase may be direct or indirect and may be necessary and/or sufficient for the manifestation of the symptoms of the disease, state or condition (or its aetiology

or progression). Thus, kinase activity (and in particular aberrant levels of kinase activity, e.g. kinase over-expression) need not necessarily be the proximal cause of the disease, state or condition: rather, it is contemplated that the kinase mediated diseases, states or conditions include those having multifactorial aetiologies and complex progressions in which the kinase in question is only partially involved. In cases where the term is applied to treatment, prophylaxis or intervention, the role played by the kinase may be direct or indirect and may be necessary and/or sufficient for the operation of the treatment, prophylaxis or outcome of the intervention. Thus, a disease state or condition mediated by a kinase includes the development of resistance to any particular cancer drug or treatment.

Thus, for example, the compounds of the invention may be useful in alleviating or reducing the incidence of cancer.

More particularly, the compounds of the formulae (I) and sub-groups thereof are inhibitors of FGFRs. For example, compounds of the invention have activity against FGFR1, FGFR2, FGFR3, and/or FGFR4, and in particular FGFRs selected from FGFR1, FGFR2 and FGFR3; or in particular the compounds of formula (I) and sub-groups thereof are inhibitors of FGFR4.

Preferred compounds are compounds that inhibit one or more FGFR selected from FGFR1, FGFR2, FGFR3, and FGFR4. Preferred compounds of the invention are those having IC_{50} values of less than 0.1 μM.

Compounds of the invention also have activity against VEGFR.

In addition many of the compounds of the invention exhibit selectivity for the FGFR 1, 2, and/or 3, and/or 4 compared to VEGFR (in particular VEGFR2) and/or PDGFR and such compounds represent one preferred embodiment of the invention. In particular, the compounds exhibit selectivity over VEGFR2. For example, many compounds of the invention have IC_{50} values against FGFR1, 2 and/or 3 and/or 4 that are between a tenth and a hundredth of the IC_{50} against VEGFR (in particular VEGFR2) and/or PDGFR B. In particular preferred compounds of the invention have at least 10 times greater activity against or inhibition of FGFR in particular FGFR1, FGFR2, FGFR3 and/or FGFR4 than

VEGFR2. More preferably the compounds of the invention have at least 100 times greater activity against or inhibition of FGFR in particular FGFR1, FGFR2, FGFR3 and/or FGFR4 than VEGFR2. This can be determined using the methods described herein.

As a consequence of their activity in modulating or inhibiting FGFR, and/or VEGFR kinases, the compounds will be useful in providing a means of preventing the growth or inducing apoptosis of neoplasias, particularly by inhibiting angiogenesis. It is therefore anticipated that the compounds will prove useful in treating or preventing proliferative disorders such as cancers. In addition, the compounds of the invention could be useful in the treatment of diseases in which there is a disorder of proliferation, apoptosis or differentiation.

In particular tumours with activating mutants of VEGFR or upregulation of VEGFR and patients with elevated levels of serum lactate dehydrogenase may be particularly sensitive to the compounds of the invention. Patients with activating mutants of any of the isoforms of the specific RTKs discussed herein may also find treatment with the compounds of the invention particularly beneficial. For example, VEGFR overexpression in acute leukemia cells where the clonal progenitor may express VEGFR. Also, particular tumours with activating mutants or upregulation or overexpression of any of the isoforms of FGFR such as FGFR1, FGFR2 or FGFR3 or FGFR4 may be particularly sensitive to the compounds of the invention and thus patients as discussed herein with such particular tumours may also find treatment with the compounds of the invention particularly beneficial. It may be preferred that the treatment is related to or directed at a mutated form of one of the receptor tyrosine kinases, such as discussed herein. Diagnosis of tumours with such mutations could be performed using techniques known to a person skilled in the art and as described herein such as RTPCR and FISH.

Examples of cancers which may be treated (or inhibited) include, but are not limited to, a carcinoma, for example a carcinoma of the bladder, breast, colon (e.g. colorectal carcinomas such as colon adenocarcinoma and colon adenoma), kidney, epidermis, liver, lung (for example adenocarcinoma, small cell lung cancer and non-small cell lung carcinomas), oesophagus, head and neck, gall bladder, ovary, pancreas (e.g. exocrine

pancreatic carcinoma), stomach, gastrointestinal (also known as gastric) cancer (e.g. gastrointestinal stromal tumours), cervix, endometrium, thyroid, prostate, or skin (for example squamous cell carcinoma or dermatofibrosarcoma protuberans); a hematopoietic tumour of lymphoid lineage, for example leukemia, acute lymphocytic leukemia, chronic lymphocytic leukemia, B-cell lymphoma (e.g. diffuse large B-cell lymphoma), T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, hairy cell lymphoma, or Burkett's lymphoma; a hematopoietic tumour of myeloid lineage, for example leukemias, acute and chronic myelogenous leukemias, chronic myelomonocytic leukemia (CMML), myeloproliferative disorder, myeloproliferative syndrome, myelodysplastic syndrome, or promyelocytic leukemia; multiple myeloma; thyroid follicular cancer; a tumour of mesenchymal origin (e.g. Ewing's sarcoma), for example fibrosarcoma or rhabdomyosarcoma; a tumour of the central or peripheral nervous system, for example astrocytoma, neuroblastoma, glioma (such as glioblastoma multiforme) or schwannoma; melanoma; seminoma; teratocarcinoma; osteosarcoma; xeroderma pigmentosum; keratoctanthoma; thyroid follicular cancer; or Kaposi's sarcoma.

Certain cancers are resistant to treatment with particular drugs. This can be due to the type of the tumour or can arise due to treatment with the compound. In this regard, references to multiple myeloma includes bortezomib sensitive multiple myeloma or refractory multiple myeloma. Similarly, references to chronic myelogenous leukemia includes imitanib sensitive chronic myelogenous leukemia and refractory chronic myelogenous leukemia. Chronic myelogenous leukemia is also known as chronic myeloid leukemia, chronic granulocytic leukemia or CML. Likewise, acute myelogenous leukemia, is also called acute myeloblastic leukemia, acute granulocytic leukemia, acute nonlymphocytic leukaemia or AML.

The compounds of the invention can also be used in the treatment of hematopoetic diseases of abnormal cell proliferation whether pre-malignant or stable such as myeloproliferative diseases. Myeloproliferative diseases ("MPD"s) are a group of diseases of the bone marrow in which excess cells are produced. They are related to, and may evolve into, myelodysplastic syndrome. Myeloproliferative diseases include polycythemia vera, essential thrombocythemia and primary myelofibrosis. A further

haematological disorder is hypereosinophilic syndrome. T-cell lymphoproliferative diseases include those derived from natural Killer cells.

In addition the compounds of the invention can be used to gastrointestinal (also known as gastric) cancer e.g. gastrointestinal stromal tumours. Gastrointestinal cancer refers to malignant conditions of the gastrointestinal tract, including the esophagus, stomach, liver, biliary system, pancreas, bowels, and anus.

Thus, in the pharmaceutical compositions, uses or methods of this invention for treating a disease or condition comprising abnormal cell growth, the disease or condition comprising abnormal cell growth in one embodiment is a cancer.

Particular subsets of cancers include multiple myeloma, bladder, cervical, prostate and thyroid carcinomas, lung, breast, and colon cancers.

A further subset of cancers includes multiple myeloma, bladder, hepatocellular, oral squamous cell carcinoma and cervical carcinomas.

The compound of the invention, having FGFR such as FGFR1 inhibitory activity, may be particularly useful in the treatment or prevention of breast cancer in particular Classic Lobular Carcinomas (CLC).

As the compounds of the invention have FGFR4 activity they will also be useful in the treatment of prostate or pituitary cancers, or they will be useful in the treatment of breast cancer, lung cancer, prostate cancer, liver cancer (HCC) or lung cancer.

In particular the compounds of the invention as FGFR inhibitors, are useful in the treatment of multiple myeloma, myeloproliferatoive disorders, endometrial cancer, prostate cancer, bladder cancer, lung cancer, ovarian cancer, breast cancer, gastric cancer, colorectal cancer, and oral squamous cell carcinoma.

Further subsets of cancer are multiple myeloma, endometrial cancer, bladder cancer, cervical cancer, prostate cancer, lung cancer, breast cancer, colorectal cancer and thyroid carcinomas.

In particular the compounds of the invention are useful in the treatment of multiple myeloma (in particular multiple myeloma with t(4;14) translocation or overexpressing FGFR3), prostate cancer (hormone refractory prostrate carcinomas), endometrial cancer (in particular endometrial tumours with activating mutations in FGFR2) and breast cancer (in particular lobular breast cancer).

In particular the compounds are useful in the treatment of lobular carcinomas such as CLC (Classic lobular carcinoma).

As the compounds have activity against FGFR3 they will be useful in the treatment of multiple myeloma and bladder cancer.

In particular the compounds are useful for the treatment of t(4;14) translocation positive multiple myeloma.

In one embodiment the compounds may be useful for the treatment of sarcoma. In one embodiment the compounds may be useful for the treatment of lung cancer, e.g. squamous cell carcinoma.

As the compounds have activity against FGFR2 they will be useful in the treatment of endometrial, ovarian, gastric and colorectal cancers. FGFR2 is also overexpressed in epithelial ovarian cancer, therefore the compounds of the invention may be specifically useful in treating ovarian cancer such as epithelial ovarian cancer.

In one embodiment, the compounds may be useful for the treatment of lung cancer, in particular NSCLC, squamous cell carcinoma, liver cancer, kidney cancer, breast cancer, colon cancer, colorectal cancer, prostate cancer.

Compounds of the invention may also be useful in the treatment of tumours pre-treated with VEGFR2 inhibitor or VEGFR2 antibody (e.g. Avastin).

In particular the compounds of the invention may be useful in the treatment of VEGFR2-resistant tumours. VEGFR2 inhibitors and antibodies are used in the treatment of

thyroid and renal cell carcinomas, therefore the compounds of the invention may be useful in the treatment of VEGFR2-resistant thyroid and renal cell carcinomas.

The cancers may be cancers which are sensitive to inhibition of any one or more FGFRs selected from FGFR1, FGFR2, FGFR3, FGFR4, for example, one or more FGFRs selected from FGFR1, FGFR2 or FGFR3.

Whether or not a particular cancer is one which is sensitive to inhibition of FGFR or VEGFR signalling may be determined by means of a cell growth assay as set out below or by a method as set out in the section headed “Methods of Diagnosis”.

The compounds of the invention, and in particular those compounds having FGFR, or VEGFR inhibitory activity, may be particularly useful in the treatment or prevention of cancers of a type associated with or characterised by the presence of elevated levels of FGFR, or VEGFR, for example the cancers referred to in this context in the introductory section of this application.

The compounds of the present invention may be useful for the treatment of the adult population. The compounds of the present invention may be useful for the treatment of the pediatric population.

It has been discovered that some FGFR inhibitors can be used in combination with other anticancer agents. For example, it may be beneficial to combine an inhibitor that induces apoptosis with another agent which acts via a different mechanism to regulate cell growth thus treating two of the characteristic features of cancer development. Examples of such combinations are set out below.

The compounds of the invention may be useful in treating other conditions which result from disorders in proliferation such as type II or non-insulin dependent diabetes mellitus, autoimmune diseases, head trauma, stroke, epilepsy, neurodegenerative diseases such as Alzheimer’s, motor neurone disease, progressive supranuclear palsy, corticobasal degeneration and Pick’s disease for example autoimmune diseases and neurodegenerative diseases.

One sub-group of disease states and conditions that the compounds of the invention may be useful consists of inflammatory diseases, cardiovascular diseases and wound healing.

FGFR, and VEGFR are also known to play a role in apoptosis, angiogenesis, proliferation, differentiation and transcription and therefore the compounds of the invention could also be useful in the treatment of the following diseases other than cancer; chronic inflammatory diseases, for example systemic lupus erythematosus, autoimmune mediated glomerulonephritis, rheumatoid arthritis, psoriasis, inflammatory bowel disease, autoimmune diabetes mellitus, Eczema hypersensitivity reactions, asthma, COPD, rhinitis, and upper respiratory tract disease; cardiovascular diseases for example cardiac hypertrophy, restenosis, atherosclerosis; neurodegenerative disorders, for example Alzheimer's disease, AIDS-related dementia, Parkinson's disease, amyotropic lateral sclerosis, retinitis pigmentosa, spinal muscular atropy and cerebellar degeneration; glomerulonephritis; myelodysplastic syndromes, ischemic injury associated myocardial infarctions, stroke and reperfusion injury, arrhythmia, atherosclerosis, toxin-induced or alcohol related liver diseases, haematological diseases, for example, chronic anemia and aplastic anemia; degenerative diseases of the musculoskeletal system, for example, osteoporosis and arthritis, aspirin-sensitive rhinosinusitis, cystic fibrosis, multiple sclerosis, kidney diseases and cancer pain.

In addition, mutations of FGFR2 are associated with several severe abnormalities in human skeletal development and thus the compounds of invention could be useful in the treatment of abnormalities in human skeletal development, including abnormal ossification of cranial sutures (craniosynostosis), Apert (AP) syndrome, Crouzon syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrate syndrome, and Pfeiffer syndrome.

The compound of the invention, having FGFR such as FGFR2 or FGFR3 inhibitory activity, may be particularly useful in the treatment or prevention of the skeletal diseases. Particular skeletal diseases are achondroplasia or thanatophoric dwarfism (also known as thanatophoric dysplasia).

The compound of the invention, having FGFR such as FGFR1, FGFR2 or FGFR3 inhibitory activity, may be particularly useful in the treatment or prevention in pathologies in which progressive fibrosis is a symptom. Fibrotic conditions in which the compounds of the inventions may be useful in the treatment of include diseases exhibiting abnormal or excessive deposition of fibrous tissue for example in liver cirrhosis, glomerulonephritis, pulmonary fibrosis, systemic fibrosis, rheumatoid arthritis, as well as the natural process of wound healing. In particular the compounds of the inventions may also be useful in the treatment of lung fibrosis in particular in idiopathic pulmonary fibrosis.

The over-expression and activation of FGFR and VEGFR in tumor- associated vasculature has also suggested a role for compounds of the invention in preventing and disrupting initiation of tumor angiogenesis. In particular the compounds of the invention may be useful in the treatment of cancer, metastasis, leukemia's such as CLL, ocular diseases such as age-related macular degeneration in particular wet form of age-related macular degeneration, ischemic proliferative retinopathies such as retinopathy of prematurity (ROP) and diabetic retinopathy, rheumatoid arthritis and hemangioma.

The activity of the compounds of the invention as inhibitors of FGFR1-4, VEGFR and/or PDGFR A/B can be measured using the assays set forth in the examples below and the level of activity exhibited by a given compound can be defined in terms of the IC_{50} value. Preferred compounds of the present invention are compounds having an IC_{50} value of less than 1μM, more preferably less than 0.1 μM.

The invention provides compounds that have FGFR inhibiting or modulating activity, and which may be useful in preventing or treating disease states or conditions mediated by FGFR kinases.

In one embodiment, there is provided a compound as defined herein for use in therapy, for use as a medicine. In a further embodiment, there is provided a compound as defined herein for use in the prophylaxis or treatment, in particular in the treatment, of a disease state or condition mediated by a FGFR kinase.

Thus, for example, the compounds of the invention may be useful in alleviating or reducing the incidence of cancer. Therefore, in a further embodiment, there is provided a compound as defined herein for use in the prophylaxis or treatment, in particular the treatment, of cancer. In one embodiment, the compound as defined herein is for use in the prophylaxis or treatment of FGFR-dependent cancer. In one embodiment, the compound as defined herein is for use in the prophylaxis or treatment of cancer mediated by FGFR kinases.

Accordingly, the invention provides *inter alia*:

- A method for the prophylaxis or treatment of a disease state or condition mediated by a FGFR kinase, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.
- A method for the prophylaxis or treatment of a disease state or condition as described herein, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.
- A method for the prophylaxis or treatment of cancer, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.
- A method for alleviating or reducing the incidence of a disease state or condition mediated by a FGFR kinase, which method comprises administering to a subject in need thereof a compound of the formula (I) as defined herein.
- A method of inhibiting a FGFR kinase, which method comprises contacting the kinase with a kinase-inhibiting compound of the formula (I) as defined herein.
- A method of modulating a cellular process (for example cell division) by inhibiting the activity of a FGFR kinase using a compound of the formula (I) as defined herein.
- A compound of formula (I) as defined herein for use as a modulator of a cellular process (for example cell division) by inhibiting the activity of a FGFR kinase.
- A compound of formula (I) as defined herein for use in the prophylaxis or treatment of cancer, in particular the treatment of cancer.
- A compound of formula (I) as defined herein for use as a modulator (e.g. inhibitor) of FGFR.
- The use of a compound of formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of a disease state or condition

mediated by a FGFR kinase, the compound having the formula (I) as defined herein.

- The use of a compound of formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of a disease state or condition as described herein.
- The use of a compound of formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment, in particular the treatment, of cancer.
- The use of a compound of formula (I) as defined herein for the manufacture of a medicament for modulating (e.g. inhibiting) the activity of FGFR.
- Use of a compound of formula (I) as defined herein in the manufacture of a medicament for modulating a cellular process (for example cell division) by inhibiting the activity of a FGFR kinase.
- The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for prophylaxis or treatment of a disease or condition characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4).
- The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of a cancer, the cancer being one which is characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4).
- The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of cancer in a patient selected from a sub-population possessing a genetic aberrations of FGFR3 kinase.
- The use of a compound of the formula (I) as defined herein for the manufacture of a medicament for the prophylaxis or treatment of cancer in a patient who has been diagnosed as forming part of a sub-population possessing a genetic aberrations of FGFR3 kinase.
- A method for the prophylaxis or treatment of a disease or condition characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4), the method comprising administering a compound of the formula (I) as defined herein.
- A method for alleviating or reducing the incidence of a disease or condition characterised by up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or

FGFR3 or FGFR4), the method comprising administering a compound of the formula (I) as defined herein.

- A method for the prophylaxis or treatment of (or alleviating or reducing the incidence of) cancer in a patient suffering from or suspected of suffering from cancer; which method comprises (i) subjecting a patient to a diagnostic test to determine whether the patient possesses a genetic aberrations of FGFR3 gene; and (ii) where the patient does possess the said variant, thereafter administering to the patient a compound of the formula (I) as defined herein having FGFR3 kinase inhibiting activity.
- A method for the prophylaxis or treatment of (or alleviating or reducing the incidence of) a disease state or condition characterised by up-regulation of an FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4); which method comprises (i) subjecting a patient to a diagnostic test to detect a marker characteristic of up-regulation of a FGFR kinase (e.g. FGFR1 or FGFR2 or FGFR3 or FGFR4) and (ii) where the diagnostic test is indicative of up-regulation of a FGFR kinase, thereafter administering to the patient a compound of the formula (I) as defined herein having FGFR kinase inhibiting activity.

In one embodiment, the disease mediated by FGFR kinases is a oncology related disease (e.g. cancer). In one embodiment, the disease mediated by FGFR kinases is a non-oncology related disease (e.g. any disease disclosed herein excluding cancer). In one embodiment the disease mediated by FGFR kinases is a condition described herein. In one embodiment the disease mediated by FGFR kinases is a skeletal condition described herein. Particular abnormalities in human skeletal development, include abnormal ossification of cranial sutures (craniosynostosis), Apert (AP) syndrome, Crouzon syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrate syndrome, Pfeiffer syndrome, achondroplasia and thanatophoric dwarfism (also known as thanatophoric dysplasia).

Mutated Kinases

Drug resistant kinase mutations can arise in patient populations treated with kinase inhibitors. These occur, in part, in the regions of the protein that bind to or interact with the particular inhibitor used in therapy. Such mutations reduce or increase the capacity of the inhibitor to bind to and inhibit the kinase in question. This can occur at any of the amino acid residues which interact with the inhibitor or are important for supporting the

binding of said inhibitor to the target. An inhibitor that binds to a target kinase without requiring the interaction with the mutated amino acid residue will likely be unaffected by the mutation and will remain an effective inhibitor of the enzyme.

A study in gastric cancer patient samples showed the presence of two mutations in FGFR2, Ser167Pro in exon IIIa and a splice site mutation 940-2A-G in exon IIIc. These mutations are identical to the germline activating mutations that cause craniosynotosis syndromes and were observed in 13% of primary gastric cancer tissues studied. In addition activating mutations in FGFR3 were observed in 5% of the patient samples tested and overexpression of FGFRs has been correlated with a poor prognosis in this patient group.

In addition there are chromosomal translocations or point mutations that have been observed in FGFR which give rise to gain-of-function, over-expressed, or constitutively active biological states.

The compounds of the invention would therefore find particular application in relation to cancers which express a mutated molecular target such as FGFR. Diagnosis of tumours with such mutations could be performed using techniques known to a person skilled in the art and as described herein such as RTPCR and FISH.

It has been suggested that mutations of a conserved threonine residue at the ATP binding site of FGFR would result in inhibitor resistance. The amino acid valine 561 has been mutated to a methionine in FGFR1 which corresponds to previously reported mutations found in Abl (T315) and EGFR (T766) that have been shown to confer resistance to selective inhibitors. Assay data for FGFR1 V561M showed that this mutation conferred resistance to a tyrosine kinase inhibitor compared to that of the wild type.

Methods of Diagnosis

Prior to administration of a compound of the formula (I), a patient may be screened to determine whether a disease or condition from which the patient is or may be suffering is one which would be susceptible to treatment with a compound having activity against FGFR, and/or VEGFR.

For example, a biological sample taken from a patient may be analysed to determine whether a condition or disease, such as cancer, that the patient is or may be suffering from is one which is characterised by a genetic abnormality or abnormal protein expression which leads to up-regulation of the levels or activity of FGFR, and/or VEGFR or to sensitisation of a pathway to normal FGFR, and/or VEGFR activity, or to upregulation of these growth factor signalling pathways such as growth factor ligand levels or growth factor ligand activity or to upregulation of a biochemical pathway downstream of FGFR, and/or VEGFR activation.

Examples of such abnormalities that result in activation or sensitisation of the FGFR, and/or VEGFR signal include loss of, or inhibition of apoptotic pathways, up-regulation of the receptors or ligands, or presence of mutant variants of the receptors or ligands e.g PTK variants. Tumours with mutants of FGFR1, FGFR2 or FGFR3 or FGFR4 or up-regulation, in particular over-expression of FGFR1, or gain-of-function mutants of FGFR2 or FGFR3 may be particularly sensitive to FGFR inhibitors.

For example, point mutations engendering gain-of-function in FGFR2 have been identified in a number of conditions. In particular activating mutations in FGFR2 have been identified in 10% of endometrial tumours.

In addition, genetic aberrations of the FGFR3 receptor tyrosine kinase such as chromosomal translocations or point mutations resulting in ectopically expressed or deregulated, constitutively active, FGFR3 receptors have been identified and are linked to a subset of multiple myelomas, bladder and cervical carcinomas. A particular mutation T674I of the PDGF receptor has been identified in imatinib-treated patients. In addition, a gene amplification of 8p12-p11.2 was demonstrated in ~50% of lobular breast cancer (CLC) cases and this was shown to be linked with an increased expression of FGFR1. Preliminary studies with siRNA directed against FGFR1, or a small molecule inhibitor of the receptor, showed cell lines harbouring this amplification to be particularly sensitive to inhibition of this signalling pathway.

Alternatively, a biological sample taken from a patient may be analysed for loss of a negative regulator or suppressor of FGFR or VEGFR. In the present context, the term “loss” embraces the deletion of a gene encoding the regulator or suppressor, the

truncation of the gene (for example by mutation), the truncation of the transcribed product of the gene, or the inactivation of the transcribed product (e.g. by point mutation) or sequestration by another gene product.

The term up-regulation includes elevated expression or over-expression, including gene amplification (i.e. multiple gene copies) and increased expression by a transcriptional effect, and hyperactivity and activation, including activation by mutations. Thus, the patient may be subjected to a diagnostic test to detect a marker characteristic of up-regulation of FGFR, and/or VEGFR. The term diagnosis includes screening. By marker we include genetic markers including, for example, the measurement of DNA composition to identify mutations of FGFR, and/or VEGFR. The term marker also includes markers which are characteristic of up regulation of FGFR and/or VEGFR, including enzyme activity, enzyme levels, enzyme state (e.g. phosphorylated or not) and mRNA levels of the aforementioned proteins.

The diagnostic tests and screens are typically conducted on a biological sample selected from tumour biopsy samples, blood samples (isolation and enrichment of shed tumour cells), stool biopsies, sputum, chromosome analysis, pleural fluid, peritoneal fluid, buccal spears, biopsy or urine.

Methods of identification and analysis of mutations and up-regulation of proteins are known to a person skilled in the art. Screening methods could include, but are not limited to, standard methods such as reverse-transcriptase polymerase chain reaction (RT-PCR) or in-situ hybridization such as fluorescence in situ hybridization (FISH).

Identification of an individual carrying a mutation in FGFR, and /or VEGFR may mean that the patient would be particularly suitable for treatment with a FGFR, and /or VEGFR inhibitor. Tumours may preferentially be screened for presence of a FGFR, and /or VEGFR variant prior to treatment. The screening process will typically involve direct sequencing, oligonucleotide microarray analysis, or a mutant specific antibody. In addition, diagnosis of tumours with such mutations could be performed using techniques known to a person skilled in the art and as described herein such as RT-PCR and FISH.

In addition, mutant forms of, for example FGFR or VEGFR2, can be identified by direct sequencing of, for example, tumour biopsies using PCR and methods to sequence PCR

products directly as hereinbefore described. The skilled artisan will recognize that all such well-known techniques for detection of the over expression, activation or mutations of the aforementioned proteins could be applicable in the present case.

In screening by RT-PCR, the level of mRNA in the tumour is assessed by creating a cDNA copy of the mRNA followed by amplification of the cDNA by PCR. Methods of PCR amplification, the selection of primers, and conditions for amplification, are known to a person skilled in the art. Nucleic acid manipulations and PCR are carried out by standard methods, as described for example in Ausubel, F.M. *et al.*, eds. (2004) Current Protocols in Molecular Biology, John Wiley & Sons Inc., or Innis, M.A. *et al.*, eds. (1990) PCR Protocols: a guide to methods and applications, Academic Press, San Diego. Reactions and manipulations involving nucleic acid techniques are also described in Sambrook *et al.*, (2001), 3rd Ed, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press. Alternatively a commercially available kit for RT-PCR (for example Roche Molecular Biochemicals) may be used, or methodology as set forth in United States patents 4,666,828; 4,683,202; 4,801,531; 5,192,659, 5,272,057, 5,882,864, and 6,218,529 and incorporated herein by reference. An example of an in-situ hybridisation technique for assessing mRNA expression would be fluorescence in-situ hybridisation (FISH) (see Angerer (1987) Meth. Enzymol., **152**: 649).

Generally, in situ hybridization comprises the following major steps: (1) fixation of tissue to be analyzed; (2) prehybridization treatment of the sample to increase accessibility of target nucleic acid, and to reduce nonspecific binding; (3) hybridization of the mixture of nucleic acids to the nucleic acid in the biological structure or tissue; (4) post-hybridization washes to remove nucleic acid fragments not bound in the hybridization, and (5) detection of the hybridized nucleic acid fragments. The probes used in such applications are typically labelled, for example, with radioisotopes or fluorescent reporters. Preferred probes are sufficiently long, for example, from about 50, 100, or 200 nucleotides to about 1000 or more nucleotides, to enable specific hybridization with the target nucleic acid(s) under stringent conditions. Standard methods for carrying out FISH are described in Ausubel, F.M. *et al.*, eds. (2004) Current Protocols in Molecular Biology, John Wiley & Sons Inc and Fluorescence In Situ Hybridization: Technical Overview by John M. S. Bartlett in Molecular Diagnosis of Cancer, Methods and

Protocols, 2nd ed.; ISBN: 1-59259-760-2; March 2004, pps. 077-088; Series: Methods in Molecular Medicine.

Methods for gene expression profiling are described by (DePrimo *et al*. (2003), *BMC Cancer*, **3**:3). Briefly, the protocol is as follows: double-stranded cDNA is synthesized from total RNA Using a (dT)24 oligomer for priming first-strand cDNA synthesis, followed by second strand cDNA synthesis with random hexamer primers. The double-stranded cDNA is used as a template for *in vitro* transcription of cRNA using biotinylated ribonucleotides. cRNA is chemically fragmented according to protocols described by Affymetrix (Santa Clara, CA, USA), and then hybridized overnight on Human Genome Arrays.

Alternatively, the protein products expressed from the mRNAs may be assayed by immunohistochemistry of tumour samples, solid phase immunoassay with microtitre plates, Western blotting, 2-dimensional SDS-polyacrylamide gel electrophoresis, ELISA, flow cytometry and other methods known in the art for detection of specific proteins. Detection methods would include the use of site specific antibodies. The skilled person will recognize that all such well-known techniques for detection of upregulation of FGFR, and/or VEGFR, or detection of FGFR, and/or VEGFR variants or mutants could be applicable in the present case.

Abnormal levels of proteins such as FGFR or VEGFR can be measured using standard enzyme assays, for example, those assays described herein. Activation or overexpression could also be detected in a tissue sample, for example, a tumour tissue. By measuring the tyrosine kinase activity with an assay such as that from Chemicon International. The tyrosine kinase of interest would be immunoprecipitated from the sample lysate and its activity measured.

Alternative methods for the measurement of the over expression or activation of FGFR or VEGFR including the isoforms thereof, include the measurement of microvessel density. This can for example be measured using methods described by Orre and Rogers (Int J Cancer (1999), **84(2)** 101-8). Assay methods also include the use of markers, for example, in the case of VEGFR these include CD31, CD34 and CD105.

Therefore all of these techniques could also be used to identify tumours particularly suitable for treatment with the compounds of the invention.

The compounds of the invention are particular useful in treatment of a patient having a mutated FGFR. The G697C mutation in FGFR3 is observed in 62% of oral squamous cell carcmonas and causes constitutive activation of the kinase activity. Activating mutations of FGFR3 have also been identified in bladder carcinoma cases. These mutations were of 6 kinds with varying degrees of prevelence: R248C, S249C, G372C, S373C, Y375C, K652Q. In addition, a Gly388Arg polymorphism in FGFR4 has been found to be associated with increased incidence and aggressiveness of prostate, colon, lung, liver (HCC) and breast cancer.

Therefore in a further aspect the invention includes use of a compound according to the invention for the manufacture of a medicament for the treatment or prophylaxis of a disease state or condition in a patient who has been screened and has been determined as suffering from, or being at risk of suffering from, a disease or condition which would be susceptible to treatment with a compound having activity against FGFR.

Particular mutations a patient is screened for include G697C, R248C, S249C, G372C, S373C, Y375C, K652Q mutations in FGFR3 and Gly388Arg polymorphism in FGFR4.

In another aspect the invention includes a compound of the invention for use in the prophylaxis or treatment of cancer in a patient selected from a sub-population possessing a variant of the FGFR gene (for example G697C mutation in FGFR3 and Gly388Arg polymorphism in FGFR4).

MRI determination of vessel normalization (e.g. using MRI gradient echo, spin echo, and contrast enhancement to measure blood volume, relative vessel size, and vascular permeability) in combination with circulating biomarkers (circulating progenitor cells (CPCs), CECs, SDF1, and FGF2) may also be used to identify VEGFR2-resistant tumours for treatment with a compound of the invention.

Pharmaceutical Compositions and Combinations

In view of their useful pharmacological properties, the subject compounds may be

formulated into various pharmaceutical forms for administration purposes.

In one embodiment the pharmaceutical composition (e.g. formulation) comprises at least one active compound of the invention together with one or more pharmaceutically acceptable carriers, adjuvants, excipients, diluents, fillers, buffers, stabilisers, preservatives, lubricants, or other materials well known to those skilled in the art and optionally other therapeutic or prophylactic agents.

To prepare the pharmaceutical compositions of this invention, an effective amount of a compound of the present invention, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. The pharmaceutical compositions can be in any form suitable for oral, parenteral, topical, intranasal, ophthalmic, otic, rectal, intra-vaginal, or transdermal administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets.

Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, to aid solubility for example, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause a significant deleterious effect

to the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment. It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient, calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

The compound of the invention is administered in an amount sufficient to exert its anti-tumour activity.

Those skilled in the art could easily determine the effective amount from the test results presented hereinafter. In general it is contemplated that a therapeutically effective amount would be from 0.005 mg/kg to 100 mg/kg body weight, and in particular from 0.005 mg/kg to 10 mg/kg body weight. It may be appropriate to administer the required dose as single, two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 0.5 to 500 mg, in particular 1 mg to 500 mg, more in particular 10 mg to 500 mg of active ingredient per unit dosage form.

Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 % by weight, more preferably from 0.1 to 70 % by weight, even more preferably from 0.1 to 50 % by weight of the compound of the present invention, and, from 1 to 99.95 % by weight, more preferably from 30 to 99.9 % by weight, even more preferably from 50 to 99.9 % by weight of a pharmaceutically acceptable carrier, all percentages being based on the total weight of the composition.

As another aspect of the present invention, a combination of a compound of the present invention with another anticancer agent is envisaged, especially for use as a medicine, more specifically for use in the treatment of cancer or related diseases.

For the treatment of the above conditions, the compounds of the invention may be advantageously employed in combination with one or more other medicinal agents, more particularly, with other anti-cancer agents or adjuvants in cancer therapy. Examples of anti-cancer agents or adjuvants (supporting agents in the therapy) include but are not limited to:

- platinum coordination compounds for example cisplatin optionally combined with amifostine, carboplatin or oxaliplatin;
- taxane compounds for example paclitaxel, paclitaxel protein bound particles (Abraxane^{TM}) or docetaxel;
- topoisomerase I inhibitors such as camptothecin compounds for example irinotecan, SN-38, topotecan, topotecan hcl;
- topoisomerase II inhibitors such as anti-tumour epipodophyllotoxins or podophyllotoxin derivatives for example etoposide, etoposide phosphate or teniposide;
- anti-tumour vinca alkaloids for example vinblastine, vincristine or vinorelbine;
- anti-tumour nucleoside derivatives for example 5-fluorouracil, leucovorin, gemcitabine, gemcitabine hcl, capecitabine, cladribine, fludarabine, nelarabine;
- alkylating agents such as nitrogen mustard or nitrosourea for example cyclophosphamide, chlorambucil, carmustine, thiotepa, mephalan (melphalan), lomustine, altretamine, busulfan, dacarbazine, estramustine, ifosfamide optionally in combination with mesna, pipobroman, procarbazine, streptozocin, telozolomide, uracil;

- anti-tumour anthracycline derivatives for example daunorubicin, doxorubicin optionally in combination with dexrazoxane, doxil, idarubicin, mitoxantrone, epirubicin, epirubicin hcl, valrubicin;
- molecules that target the IGF-1 receptor for example picropodophilin;
- tetracarcin derivatives for example tetrocarcin A;
- glucocorticoïden for example prednisone;
- antibodies for example trastuzumab (HER2 antibody), rituximab (CD20 antibody), gemtuzumab, gemtuzumab ozogamicin, cetuximab, pertuzumab, bevacizumab, alemtuzumab, eculizumab, ibritumomab tiuxetan, nofetumomab, panitumumab, tositumomab, CNTO 328;
- estrogen receptor antagonists or selective estrogen receptor modulators or inhibitors of estrogen synthesis for example tamoxifen, fulvestrant, toremifene, droloxifene, faslodex, raloxifene or letrozole;
- aromatase inhibitors such as exemestane, anastrozole, letrazole, testolactone and vorozole;
- differentiating agents such as retinoids, vitamin D or retinoic acid and retinoic acid metabolism blocking agents (RAMBA) for example accutane;
- DNA methyl transferase inhibitors for example azacytidine or decitabine;
- antifolates for example premetrexed disodium;
- antibiotics for example antinomycin D, bleomycin, mitomycin C, dactinomycin, carminomycin, daunomycin, levamisole, plicamycin, mithramycin;
- antimetabolites for example clofarabine, aminopterin, cytosine arabinoside or methotrexate, azacitidine, cytarabine, floxuridine, pentostatin, thioguanine;
- apoptosis inducing agents and antiangiogenic agents such as Bcl-2 inhibitors for example YC 137, BH 312, ABT 737, gossypol, HA 14-1, TW 37 or decanoic acid;
- tubuline-binding agents for example combrestatin, colchicines or nocodazole;
- kinase inhibitors (e.g. EGFR (epithelial growth factor receptor) inhibitors, MTKI (multi target kinase inhibitors), mTOR inhibitors) for example flavoperidol, imatinib mesylate, erlotinib, gefitinib, dasatinib, lapatinib, lapatinib ditosylate, sorafenib, sunitinib, sunitinib maleate, temsirolimus;
- farnesyltransferase inhibitors for example tipifarnib;
- histone deacetylase (HDAC) inhibitors for example sodium butyrate, suberoylanilide hydroxamide acid (SAHA), depsipeptide (FR 901228), NVP-LAQ824, R306465, JNJ-26481585, trichostatin A, vorinostat;

- Inhibitors of the ubiquitin-proteasome pathway for example PS-341, MLN .41 or bortezomib;
- Yondelis;
- Telomerase inhibitors for example telomestatin;
- Matrix metalloproteinase inhibitors for example batimastat, marimastat, prinostat or metastat.
- Recombinant interleukins for example aldesleukin, denileukin diftitox, interferon alfa 2a, interferon alfa 2b, peginterferon alfa 2b
- MAPK inhibitors
- Retinoids for example alitretinoin, bexarotene, tretinoin
- Arsenic trioxide
- Asparaginase
- Steroids for example dromostanolone propionate, megestrol acetate, nandrolone (decanoate, phenpropionate), dexamethasone
- Gonadotropin releasing hormone agonists or antagonists for example abarelix, goserelin acetate, histrelin acetate, leuprolide acetate
- Thalidomide, lenalidomide
- Mercaptopurine, mitotane, pamidronate, pegademase, pegaspargase, rasburicase
- BH3 mimetics for example ABT-737
- MEK inhibitors for example PD98059, AZD6244, CI-1040
- colony-stimulating factor analogs for example filgrastim, pegfilgrastim, sargramostim; erythropoietin or analogues thereof (e.g. darbepoetin alfa); interleukin 11; oprelvekin; zoledronate, zoledronic acid; fentanyl; bisphosphonate; palifermin.
- a steroidal cytochrome P450 17alpha-hydroxylase-17,20-lyase inhibitor (CYP17), e.g. abiraterone, abiraterone acetate.

The compounds of the present invention also have therapeutic applications in sensitising tumour cells for radiotherapy and chemotherapy.

Hence the compounds of the present invention can be used as "radiosensitizer" and/or “chemosensitizer” or can be given in combination with another "radiosensitizer" and/or “chemosensitizer”.

The term "radiosensitizer", as used herein, is defined as a molecule, preferably a low molecular weight molecule, administered to animals in therapeutically effective amounts to increase the sensitivity of the cells to ionizing radiation and/or to promote the treatment of diseases which are treatable with ionizing radiation.

The term “chemosensitizer”, as used herein, is defined as a molecule, preferably a low molecular weight molecule, administered to animals in therapeutically effective amounts to increase the sensitivity of cells to chemotherapy and/or promote the treatment of diseases which are treatable with chemotherapeutics.

Several mechanisms for the mode of action of radiosensitizers have been suggested in the literature including: hypoxic cell radiosensitizers (e.g., 2- nitroimidazole compounds, and benzotriazine dioxide compounds) mimicking oxygen or alternatively behave like bioreductive agents under hypoxia; non-hypoxic cell radiosensitizers (e.g., halogenated pyrimidines) can be analogoues of DNA bases and preferentially incorporate into the DNA of cancer cells and thereby promote the radiation-induced breaking of DNA molecules and/or prevent the normal DNA repair mechanisms; and various other potential mechanisms of action have been hypothesized for radiosensitizers in the treatment of disease.

Many cancer treatment protocols currently employ radiosensitizers in conjunction with radiation of x-rays. Examples of x-ray activated radiosensitizers include, but are not limited to, the following: metronidazole, misonidazole, desmethylmisonidazole, pimonidazole, etanidazole, nimorazole, mitomycin C, RSU 1069, SR 4233, EO9, RB 6145, nicotinamide, 5-bromodeoxyuridine (BUdR), 5- iododeoxyuridine (IUdR), bromodeoxycytidine, fluorodeoxyuridine (FudR), hydroxyurea, cisplatin, and therapeutically effective analogs and derivatives of the same.

Photodynamic therapy (PDT) of cancers employs visible light as the radiation activator of the sensitizing agent. Examples of photodynamic radiosensitizers include the following, but are not limited to: hematoporphyrin derivatives, Photofrin, benzoporphyrin derivatives, tin etioporphyrin, pheoborbide-a, bacteriochlorophyll-a, naphthalocyanines, phthalocyanines, zinc phthalocyanine, and therapeutically effective analogs and derivatives of the same.

Radiosensitizers may be administered in conjunction with a therapeutically effective amount of one or more other compounds, including but not limited to: compounds which promote the incorporation of radiosensitizers to the target cells; compounds which control the flow of therapeutics, nutrients, and/or oxygen to the target cells; chemotherapeutic agents which act on the tumour with or without additional radiation; or other therapeutically effective compounds for treating cancer or other diseases.

Chemosensitizers may be administered in conjunction with a therapeutically effective amount of one or more other compounds, including but not limited to: compounds which promote the incorporation of chemosensitizers to the target cells; compounds which control the flow of therapeutics, nutrients, and/or oxygen to the target cells; chemotherapeutic agents which act on the tumour or other therapeutically effective compounds for treating cancer or other disease. Calcium antagonists, for example verapamil, are found useful in combination with antineoplastic agents to establish chemosensitivity in tumor cells resistant to accepted chemotherapeutic agents and to potentiate the efficacy of such compounds in drug-sensitive malignancies.

In view of their useful pharmacological properties, the components of the combinations according to the invention, i.e. the one or more other medicinal agent and the compound according to the present invention may be formulated into various pharmaceutical forms for administration purposes. The components may be formulated separately in individual pharmaceutical compositions or in a unitary pharmaceutical composition containing all components.

The present invention therefore also relates to a pharmaceutical composition comprising the one or more other medicinal agent and the compound according to the present invention together with a pharmaceutical carrier.

The present invention further relates to the use of a combination according to the invention in the manufacture of a pharmaceutical composition for inhibiting the growth of tumour cells.

The present invention further relates to a product containing as first active ingredient a compound according to the invention and as further active ingredient one or more

anticancer agent, as a combined preparation for simultaneous, separate or sequential use in the treatment of patients suffering from cancer.

The one or more other medicinal agents and the compound according to the present invention may be administered simultaneously (e.g. in separate or unitary compositions) or sequentially in either order. In the latter case, the two or more compounds will be administered within a period and in an amount and manner that is sufficient to ensure that an advantageous or synergistic effect is achieved. It will be appreciated that the preferred method and order of administration and the respective dosage amounts and regimes for each component of the combination will depend on the particular other medicinal agent and compound of the present invention being administered, their route of administration, the particular tumour being treated and the particular host being treated. The optimum method and order of administration and the dosage amounts and regime can be readily determined by those skilled in the art using conventional methods and in view of the information set out herein.

The weight ratio of the compound according to the present invention and the one or more other anticancer agent(s) when given as a combination may be determined by the person skilled in the art. Said ratio and the exact dosage and frequency of administration depends on the particular compound according to the invention and the other anticancer agent(s) used, the particular condition being treated, the severity of the condition being treated, the age, weight, gender, diet, time of administration and general physical condition of the particular patient, the mode of administration as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that the effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. A particular weight ratio for the present compound of formula (I) and another anticancer agent may range from 1/10 to 10/1, more in particular from 1/5 to 5/1, even more in particular from 1/3 to 3/1.

The platinum coordination compound is advantageously administered in a dosage of 1 to 500mg per square meter (mg/m^2) of body surface area, for example 50 to 400 mg/m^2, particularly for cisplatin in a dosage of about 75 mg/m^2 and for carboplatin in about 300mg/m^2 per course of treatment.

The taxane compound is advantageously administered in a dosage of 50 to 400 mg per square meter (mg/m^2) of body surface area, for example 75 to 250 mg/m^2, particularly for paclitaxel in a dosage of about 175 to 250 mg/m^2 and for docetaxel in about 75 to 150 mg/m^2 per course of treatment.

The camptothecin compound is advantageously administered in a dosage of 0.1 to 400 mg per square meter (mg/m^2) of body surface area, for example 1 to 300 mg/m^2, particularly for irinotecan in a dosage of about 100 to 350 mg/m^2 and for topotecan in about 1 to 2 mg/m^2 per course of treatment.

The anti-tumour podophyllotoxin derivative is advantageously administered in a dosage of 30 to 300 mg per square meter (mg/m^2) of body surface area, for example 50 to 250mg/m^2, particularly for etoposide in a dosage of about 35 to 100 mg/m^2 and for teniposide in about 50 to 250 mg/m^2 per course of treatment.

The anti-tumour vinca alkaloid is advantageously administered in a dosage of 2 to 30 mg per square meter (mg/m^2) of body surface area, particularly for vinblastine in a dosage of about 3 to 12 mg/m^2 , for vincristine in a dosage of about 1 to 2 mg/m^2 , and for vinorelbine in dosage of about 10 to 30 mg/m^2 per course of treatment.

The anti-tumour nucleoside derivative is advantageously administered in a dosage of 200 to 2500 mg per square meter (mg/m^2) of body surface area, for example 700 to 1500 mg/m^2, particularly for 5-FU in a dosage of 200 to 500mg/m^2, for gemcitabine in a dosage of about 800 to 1200 mg/m^2 and for capecitabine in about 1000 to 2500 mg/m^2 per course of treatment.

The alkylating agents such as nitrogen mustard or nitrosourea is advantageously administered in a dosage of 100 to 500 mg per square meter (mg/m^2) of body surface area, for example 120 to 200 mg/m^2, particularly for cyclophosphamide in a dosage of about 100 to 500 mg/m^2 , for chlorambucil in a dosage of about 0.1 to 0.2 mg/kg, for carmustine in a dosage of about 150 to 200 mg/m^2 , and for lomustine in a dosage of about 100 to 150 mg/m^2 per course of treatment.

The anti-tumour anthracycline derivative is advantageously administered in a dosage of 10 to 75 mg per square meter (mg/m^2) of body surface area, for example 15 to 60 mg/m^2, particularly for doxorubicin in a dosage of about 40 to 75 mg/m^2, for daunorubicin in a dosage of about 25 to 45mg/m^2 , and for idarubicin in a dosage of about 10 to 15 mg/m^2 per course of treatment.

The antiestrogen agent is advantageously administered in a dosage of about 1 to 100 mg daily depending on the particular agent and the condition being treated. Tamoxifen is advantageously administered orally in a dosage of 5 to 50 mg, preferably 10 to 20 mg twice a day, continuing the therapy for sufficient time to achieve and maintain a therapeutic effect. Toremifene is advantageously administered orally in a dosage of about 60mg once a day, continuing the therapy for sufficient time to achieve and maintain a therapeutic effect. Anastrozole is advantageously administered orally in a dosage of about 1mg once a day. Droloxifene is advantageously administered orally in a dosage of about 20-100mg once a day. Raloxifene is advantageously administered orally in a dosage of about 60mg once a day. Exemestane is advantageously administered orally in a dosage of about 25mg once a day.

Antibodies are advantageously administered in a dosage of about 1 to 5 mg per square meter (mg/m^2) of body surface area, or as known in the art, if different. Trastuzumab is advantageously administered in a dosage of 1 to 5 mg per square meter (mg/m^2) of body surface area, particularly 2 to 4mg/m^2 per course of treatment.
These dosages may be administered for example once, twice or more per course of treatment, which may be repeated for example every 7, 14, 21 or 28 days.

The compounds of formula (I), the pharmaceutically acceptable addition salts, in particular pharmaceutically acceptable acid addition salts, and stereoisomeric forms thereof can have valuable diagnostic properties in that they can be used for detecting or identifying the formation of a complex between a labelled compound and other molecules, peptides, proteins, enzymes or receptors.

The detecting or identifying methods can use compounds that are labelled with labelling agents such as radioisotopes, enzymes, fluorescent substances, luminous substances, etc. Examples of the radioisotopes include ^{125}I, ^{131}I, ^{3}H and ^{14}C. Enzymes are usually

made detectable by conjugation of an appropriate substrate which, in turn catalyses a detectable reaction. Examples thereof include, for example, beta-galactosidase, beta-glucosidase, alkaline phosphatase, peroxidase and malate dehydrogenase, preferably horseradish peroxidase. The luminous substances include, for example, luminol, luminol derivatives, luciferin, aequorin and luciferase.

Biological samples can be defined as body tissue or body fluids. Examples of body fluids are cerebrospinal fluid, blood, plasma, serum, urine, sputum, saliva and the like.

General Synthetic Routes


The following examples illustrate the present invention but are examples only and are not intended to limit the scope of the claims in any way.

Experimental Part

Hereinafter, the term 'CH_3CN' means acetonitrile, 'DCM' means dichloromethane, 'TBAF' means tetrabutylammonium fluoride, 'K_2CO_3' means potassium carbonate, '$MgSO_4$' means magnesium sulphate, 'MeOH' means methanol, 'EtOH' means ethanol, 'EtOAc' means ethyl acetate, 'Et_3N' means triethylamine, 'HOBt' means 1-hydroxy-1H-benzotriazole, 'DPPP' means 1,3-propanediylbis[diphenylphosphine, 'DIPE' means diisopropyl ether, 'THF' means tetrahydrofuran, 'NH_4Cl' means ammonium chloride, '$Pd(PPh_3)_4$' means tetrakis(triphenylphosphine)palladium, 'DIPEA' means *N*-ethyl-*N*-(1-methylethyl)- 2-propylamine, 'DMF' means *N,N*-dimethylformamide, 'NaH' means sodium hydride, '$Pd_2(dba)_3$' means tris(dibenzylideneacetone) dipalladium (0), 'HOAc' means acetic acid, 'PPh_3' means triphenylphosphine, 'NH_4OH' means ammonium hydroxide, 'TBDMSCl' means *tert*-butyldimethylsilyl chloride, 'S-Phos' means dicyclohexyl(2',6'-dimethoxy[1,1'-biphenyl]-2-yl)-phosphine, 'X-Phos' means dicyclohexyl[2',4',6'-tris(1-methylethyl)[1,1'-biphenyl]-2-yl]-phosphine, 'Na_2SO_4' means sodium sulfate, 'i-PrOH' means 2-propanol, 't-BuOH' means 2-methyl-2-propanol, 'K_3PO_4' means potassium phosphate, MP means melting point.


A. Preparation of the intermediates

Example A1

a-1) Preparation of intermediate 1 

7-bromo-2(1H)-quinoxalinone (47.2g; 210mmol) was added to phosphorus oxychloride (470mL).The reaction mixture was stirred at 100°C for 2 hours, cooled down to room temperature and evaporated to dryness. The crude product was taken up into DCM and poured onto ice, water and K_2CO_3 powder. The mixture was filtered over celite. The celite was washed twice with DCM. The organic layer was decanted, dried over $MgSO_4$, filtered and evaporated to dryness to give 49g (96%) of intermediate 1 (grey solid). MP=146°C.



Intermediate 1 was alternatively also prepared using the following procedure: Thionyl chloride (407.5 mL; 5.59 mol), then N,N-dimethylformamide (34.6mL; 0.45mol) were added dropwise to a mixture of 7-bromo-2(1H)-quinoxalinone (500g; 2.24mol) in toluene (7.61L). The reaction mixture was stirred at 80°C for 17 hours then cooled to 35°C and poured cautiously onto water. The bi-phasic mixture was stirred for 30 minutes and then decanted. The organic layer was evaporated to dryness and the residue crystallized in methyl-tert-butyl ether, filtered and the precipitate washed with methyl-tert-butyl ether and dried to give 407g (74.7%) of intermediate 1. Filtrate was evaporated and re-crystallized in methyl-tert-butyl ether to provide a second fraction of 72g (13.2%) of intermediate 1.


b-1) Preparation of intermediate 2 

Under N_2, intermediate 1 (20g; 82.1mmol), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (17.1g; 82.1mmol), 2M sodium carbonate aqueous solution (41.1mL; 82.1mmol) in ethylene glycol dimethyl ether (200mL) were degassed by bubbling nitrogen through for 15 minutes. Tetrakis(triphenylphosphine)palladium (0) (0.95g; 0.82mmol) was added and heated at reflux for 15 hours. The reaction mixture was poured into water and extracted with EtOAc. The organic layer was dried over

$MgSO_4$, filtered and evaporated to dryness to give 29.9g. The crude compound was purified by chromatography over silica gel (Irregular SiOH,20-45µm, 1000g MATREX; mobile phase 0.1% NH_4OH, 98% DCM, 2% CH_3OH). The pure fractions were collected and concentrated till dryness to give 19.5g (82%) of intermediate 2. MP =172°C.

Intermediate 2 was alternatively also prepared using the following procedure: Intermediate 1 (502g; 2.06mol), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (450.42g; 2.16mol), triphenylphosphine (10.82g; 0.041mol) and palladium(II)acetate were added to a mixture of sodium carbonate (240.37g; 2.267mol), 1,2-dimethoxyethane (5.48L) and water (1.13L). The reaction mixture was stirred at reflux for 20 hours, then 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (42.9g; 0.206 mol) was added and the reaction mixture refluxed until complete conversion (4 hours). The reaction mixture was poured out in water, stirred for 2 hours at room temperature, filtered and the precipitate was washed with water. The precipitate was then triturated in methanol and filtered. The precipitate was washed with methanol and dried to give 532.2g (89%) of intermediate 2 (off-white powder).

Intermediate prepared according to the above protocol	starting from
	











c-1) Preparation of intermediate 3

A mixture of intermediate 2 (20g; 69.2mmol), 3,5-dimethoxyaniline (10.6g; 69.2mmol), sodium tert-butoxide (20g; 0.21mol) and 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine (2.2g; 3.5mmol) in dioxane (500mL) was degassed at room temperature under N_2 flow. After 10 minutes, palladium(II) acetate (0.78g; 3.5mmol) was added portionwise at room temperature under N_2 flow. The reaction mixture was heated at 90°C overnight. The reaction mixture was cooled to room temperature and

partitioned between water and EtOAc. The organic layers were combined, dried over $MgSO_4$, filtered and concentrated to give 40g of crude compound. This residue was taken up into DCM/ Et_2O (3/7) and the mixture was stirred for 30 minutes. The precipitate was filtered off and dried to give 20g of intermediate 3 (brown solid). The filtrate was evaporated to dryness to give 40g of a crude compound which was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 450 g MATREX; Mobile phase 0.1% NH_4OH, 98% DCM, 2% CH_3OH). The pure fractions were concentrated to give 4.2g of intermediate 3 (brown solid). MP=199°C (DSC).

Overall yield = 96.8%.


Intermediate prepared according to the above protocol	starting from
 intermediate 49	
	
	

 D=deuterium	

Intermediate 3 was alternatively also prepared using the following procedure.



A mixture of intermediate 2 (80g; 277mmol), 3,5-dimethoxyaniline (47.6g; 304mmol) and cesium carbonate (108.2g; 332mmol) in 1,2-dimethoxyethane (1.1L) was stirred at 80°C under N_2 flow and then cooled to room temperature (solution A). In another flask under N_2, a mixture of palladium(II)acetate (0.62g; 2.8mmol) and racemic -2,2'-bis(diphenylphosphino)-1,1'-binaphtyl (1.76g; 2.8mmol) was stirred at 40°C for 15 minutes and then added to solution A at 35°C. The new reaction mixture was stirred at 80°C for 20 hours, cooled to 50°C and water was added (1.11L). The reaction mixture was seeded with crystals of intermediate 3 and extra water (0.55L) was added before cooling to room temperature. The precipitate was filtered off and washed with water, then recrystallized in isopropylalcohol (with seeding). The prepcipitate was filtered off, washed with diisopropylether and dried to provide 79.2g (79.2%) of intermediate 3.


Intermediate 3 was alternatively also prepared using the following procedure.

a-2) Preparation of intermediate 4 

2-Chloro-7-nitroquinoxaline (27.8g, 133mmol), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (30.4g, 146mmol), 2M Na_2CO_3 aqueous solution (66.3mL, 133mmol) in ethylene glycol dimethyl ether (330mL) were degassed with N_2 for 15 minutes. Tetrakis(triphenylphosphine)palladium (0) (1.5g, 1.33mmol) was added and the reaction mixture was heated at 100°C for 7 hours. The reaction was poured into water. The precipitate was filtered off, taken-up with EtOAc, then filtered and dried under vacuum to give 31.4g (93%) of intermediate 4 (yellow solid). MP=231°C (DSC).



Intermediate prepared according to the	starting from

above protocol	
	



b-2) Preparation of intermediate 5


A mixture of intermediate 4 (15.7g, 61.5mmol) and Raney nickel (16g) in CH_3OH (380mL) and THF (60mL) was hydrogenated under a 3 bars pressure overnight. The reaction mixture was filtered on a celite pad which was washed 3 times with CH_3OH/ DCM (50/50), then several times with a mixture of MeOH/acetone. The combined filtrates were evaporated till dryness to give 13.1g (95%) of intermediate 5 (brown solid). MP=240°C (DSC).

Intermediate prepared according to the above protocol	starting from
	

Intermediate 5 was alternatively also prepared using the following procedure.

A 200mL stainless steel autoclave was charged under N_2 atmosphere with intermediate 2 (5g, 17.3mmol), NH_4OH (100mL) and Cu_2O (0.1g) The autoclave was closed and the reaction was carried out for 16 hours at a temperature of 150°C. The reaction mixture was extracted with DCM, the organic layer was washed with water, dried ($MgSO_4$) and filtered. The filtrate was evaporated till dryness and the residue was purified by chromatography over silica gel (kromasil C18 100A 5µm, Eka nobel; mobile phase, from









90% of a 0.25% solution of ammonium bicarbonate in water, 10% MeOH to 100% MeOH). The pure fractions were collected to give 2.4g (61.6%) of intermediate 5.




c-2) Preparation of intermediate 3

The experiment has been performed 3 times on the following amount.

A mixture of intermediate 5 (2.12g, 9.4mmol), 1-bromo-3,5-dimethoxybenzene (2.25g, 10.4mmol), sodium tert-butoxide (2.71g, 28.3mmol) and 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] (0.29g, 0.47mmol) in ethylene glycol dimethyl ether (40mL) was degassed with N_2 for 10 minutes. Palladium(II) acetate (0.21g, 0.94mmol) was added and the mixture was heated at 135°C for 60 minutes under microwave irradiation The mixture was cooled to room temperature, poured into H_2O and EtOAc. The 3 experiments were combined for the work up. The mixture was filtered over celite. The filtrate was extracted with EtOAc. The combined organic layers were dried over $MgSO_4$, filtered and evaporated to dryness to give 11.3g of crude compound. The residue was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, (450g) MATREX; mobile phase 0.1% NH_4OH, 95% DCM, 5% iPrOH). The pure fractions were collected and the solvent was evaporated, yielding 7.6g (74%) of intermediate 3 (brown solid).


Intermediate prepared according to the above protocol	starting from
	 and 
	 and 
	



a-4) Preparation of intermediate 6


Tert-butyldimethylsilyl chloride (2.096g, 13.9mmol) was added to 3-chloro-5-methoxybenzenemethanol (2g, 11.6mmol) in DCM (40mL) at 0°C, followed by imidazole (2.5g, 36.85mmol). The reaction mixture was slowly allowed to warm to room temperature and stirred overnight. The reaction mixture was partitioned between EtOAc and water. The 2 phases were separated, the organic phase was dried ($MgSO_4$), filtered

and concentrated to give an oil which solidified on standing. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40 µm, 90g; mobile phase 30% EtOAc, 70% pentane). The fractions were collected and the solvent was evaporated, yielding 2.56g (77%) of intermediate 6.



b-4) Preparation of intermediate 7

Intermediate 6 (1.39g, 3.9mmol), intermediate 5 (0.7g, 3.1mmol), Cs_2CO_3 (3g, 0.3mmol), tris(dibenzilideneacetone)dipalladium (0.28g, 0.3mmol) and X-Phos (0.33g, 0.68mmol) in t-BuOH (20mL) were stirred at 100°C under microwave irradiation for 3 hours. The reaction mixture was filtered through celite and the filtrate was concentrated to ~1/3 of the initial volume. H_2O and EtOAc were added and the organic phase was separated, dried ($MgSO_4$), filtered and concentrated. The residue was purified by chromatography over silica gel (Hyperprep C18 HS BDS 100A 8mu (Shandon); mobile phase gradient from 70% of a 0.25% solution of ammonium bicarbonate in water / 30% CH_3CN to 10% of a 0.25% solution of ammoniumbicarbonate in water /90% CH_3CN). The pure fractions were collected and the solvent was evaporated, yielding 418mg of intermediate 7.




a-5) Preparation of intermediate 8

A mixture of intermediate 13 (see hereinafter) (9.45g, 29.9mmol), 1-(1-methylethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (8.48g, 35.9mmol), potassium phosphate (15.88g, 74.8mmol) and dicyclohexyl(2',6'-dimethoxy[1,1'-biphenyl]-2-yl)phosphine (1.23g, 3.0mmol) in dioxane (125mL) and H_2O (25mL) was degassed at room temperature under N_2 flow. After 10 minutes, $Pd(PPh_3)_4$ (1.73g, 1.5mmol) was added portionwise. The reaction mixture was then heated at 80°C overnight, then cooled to room temperature and poured out into ice water. EtOAc was added and the organic layer was washed with water, then with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (20.2g) was purified by chromatography over silica gel

(Irregular SiOH, 20-45µm, 1000g MATREX; mobile phase 95% DCM, 5% MeOH, 0.1% NH_4OH). The product fractions were collected and the solvent was evaporated, yielding 10 g (85%). of intermediate 8.


Example A2



Preparation of intermediate 9

NaH (1.77g; 44.27mmol) was added portionwise to a solution of intermediate 3 (8g; 22.13 mmol) in *N,N*-dimethylformamide (160mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour. Then, (2-bromoethoxy)-*tert*-butyldimethylsilane (9.5mL; 44.27mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 1 hour at 5°C then, allowed to warm to room temperature and stirred overnight. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 17g of a residue which was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 200g; mobile phase gradient from 100% DCM to 96% DCM, 4% MeOH).
The pure fractions were collected and concentrated yielding 11g (95%) of intermediate 9.


Intermediate 9 was alternatively also prepared using the following procedure.



a) Preparation of intermediate 40


A mixture of 3,5-dimethoxyphenylamine (250g; 1.63mol), cesium carbonate (319g; 0.98mol) and water (0.33L) in 1,2-dimethoxyethane (2L) was heated to 60°C. Then carbonochloridic acid, 2-chloroethyl ester (250g; 1.75mol) was added dropwise at this

temperature over 1 hour. A solution of potassium hydroxide (458g; 8.2mol) in water (1.3L) was added in one portion. The reaction mixture was stirred at 60°C for 30 minutes, then heated at 100°C to distill off 1,2-dimethoxyethane using a Dean-Starck trap. The residue was cooled to 50°C and extracted with methyl-tert-butyl ether (1.14L). The organic layer was washed with water, dried ($MgSO_4$), filtered and the filtrate was evaporated till dryness. The residue was crystallized in a mixture of methyl-tert-butyl ether and heptane. The precipitate was filtered off and dried to provide 241.8g (75%) of intermediate 40.



b) Preparation of intermediate 41

TBDMSCl (262.7g; 1.74mol) was added portionwise over 10 minutes, under a N_2 atmosphere, to a solution of intermediate 40 (327.4g; 1.66mol) and 1H-imidazole (124.3g; 1.825mol) in DCM (3.3L) at room temperature. Upon completion of the reaction, water (3.3L) was added and the organic layer was decanted, washed with water (3.3L), dried ($MgSO_4$), filtered and the filtrate was filtered on silica gel and concentrated to give 496g (95.9%) of intermediate 41, used crude for the next step.




c) Preparation of intermediate 9

Under an inert atmosphere, a solution of palladium(II) acetate (1.16g; 5.2mmol), racemic 2,2'-bis(diphenylphosphino)-1,1'-binaphtyl (4.4g; 6.9mmol) in 1,2-dimethoxyethane (52mL) was added at room temperature to a solution of intermediate 2 (100g; 346mmol), intermediate 41 (118.5g; 380.5mmol) and cesium carbonate (135g; 415mmol) in 1,2-dimethoxyethane (1.4L). The reaction mixture was heated at 80°C over 1 hour, stirred at this temperature for 2 hours and refluxed overnight. Water (0.5L) and DCM (1.5L) were then added at room temperature and the organic layer was separated, washed with


water and evaporated till dryness to provide crude intermediate 9 (211g) which can directly be used into the next step.

Example A3




Preparation of intermediate 10

Methanesulfonyl chloride (3.8mL; 49.33mmol) was added dropwise to a solution of compound 1 (10g; 24.66mmol) and Et_3N (8.58mL; 61.67mmol) in DCM (250mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then 1 hour at room temperature. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness (30°C). The residue was precipitated by addition of DIPE. The solid was filtered yielding, after drying, 10.09g (94%) of intermediate 10 (red solid). MP = 161°C (kofler).

Intermediate prepared according to the above protocol	**starting from**
Intermediate 17a 	compound 3

Example A4





a-1) Preparation of intermediate 11


NaH (1.1g; 27.67mmol) was added portionwise to a solution of intermediate 3 (5g; 13.83mmol) in *N*,*N*-dimethylformamide (80mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour then (3-bromopropoxy) (1,1-dimethylethyl)dimethylsilane (6.41mL, 27.67mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred 1 hour at 5°C then warmed to room temperature and stirred overnight. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give a crude residue (9.1g). Purification by chromatography over silica gel (Irregular SiOH, 15-40µm; mobile phase gradient from 100% DCM to 98% DCM, 2% MeOH) afforded after concentration of the pure fractions 7g (94%) of intermediate 11.

Intermediate 11 was alternatively also prepared using the following procedure.

NaH (31.65g, 60%w/w in oil; 0.79 mol) was added portionwise over 15 minutes to a cooled (-2°C) solution of intermediate 3 (130g; 0.36 mol) in N,N-dimethylacetamide. The reaction mixture was stirred at -2°C for 30 minutes before addition of (3-bromopropoxy) (1,1-dimethylethyl)dimethylsilane (100.2g; 0.4 mol). The reaction mixture was further stirred at -2°C for 1.5 hours and overnight at room temperature after removal of the cooling system. The reaction mixture was then poured out in water (2.5L), DCM (1L) was added and the pH was adjusted to 6 with acetic acid. The layers were separated, the organic layer was washed with water, dried ($MgSO_4$), filtered and concentrated till dryness to provide 167.2g (87%) of intermediate 11.


Intermediate prepared according to the above protocol	starting from
 Intermediate 62	 intermediate 3

Intermediate 11 was alternatively also prepared using the following procedure.



a-2) Preparation of intermediate 12

7-Bromo-2(1H)-quinoxalinone (25g; 0.11mol), 3,5-dimethoxyaniline (20.42g; 0.133mol), sodium tert-butoxide (32g; 0.333mol), 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] (6.9g; 0.011mol) in ethylene glycol dimethyl ether (400mL) were degassed with N_2 for 10 minutes. Palladium(II) acetate (2.5g; 0.011mol) was added and the mixture refluxed for 5 hours. The reaction mixture was cooled to room temperature and the solvent was concentrated under vacuum to150mL. The residue was poured onto ice water (1.5L) under stirring and EtOAc was added (100mL). The suspension was stirred at room temperature overnight and the precipitate was filtered off, washed with water, then CH_3CN and dried yielding 33g of intermediate 12.




b-2-a) Preparation of intermediate 13

Intermediate 12 (30g; 0.1mol) was added portionwise at room temperature to phosphorus oxychloride (415mL). Then the reaction mixture was heated at 80°C and stirred at this temperature for 40 minutes. The mixture was cooled to room temperature and phosphorous oxychloride was removed under vacuum. The residue was carefully poured onto an aqueous solution of K_2CO_3. The aqueous layer was extracted with DCM.

The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 450g; mobile phase, gradient from 100% DCM to 98%DCM, 2%MeOH). The product fractions were collected and the solvent was evaporated, to give 22.6g (70%) of intermediate 13. MP=137°C (Kofler).



Intermediate 13 was alternatively also prepared using the following procedure.


b-2-b) N-Chlorosuccinimide (11.23g; 84.08mmol) was added portionwise at room temperature to a suspension of PPh_3 (22.05g, 84.08mmol) in dioxane (500mL). The reaction mixture was stirred for 30 minutes. Intermediate 12 (5g; 16.8mmol) was added and the reaction mixture was refluxed for 5 hours, then cooled to room temperature and basified with Et_3N (10mL) under stirring. The suspension was stirred overnight and the insoluble material was removed by filtration. The filtrate was concentrated and the residue (35g) was purified by chromatography over silica gel (Irregular SiOH, 15-40 µm, 400g; mobile phase 100% DCM). The pure fractions were collected and evaporated to dryness, yielding 2g (37%) of intermediate 13.



c-2) Preparation of intermediate 14



NaH (1.48g; 37.1mmol) was added portionwise to a solution of intermediate 13 (9g; 28.50mmol) in DMF (100mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then, (3-bromopropoxy)(1,1-dimethylethyl)dimethylsilane (8.58mL; 37.1mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 1 hour at 5°C then allowed to warm to room temperature and stirred for 4 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (17.5g) was purified by chromatography over silica gel (Irregular SiOH, 20-45 µm, 1000g, MATREX; mobile phase 98% DCM, 2% Cyclohexane). The pure fractions were collected and the solvent was evaporated, yielding 13.3g (95%) of intermediate14.

Intermediate prepared according to the above protocol	starting from
	




d-2) Preparation of intermediate 11

A mixture of intermediate 14 (15.5g; 31.8mmol), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (9.9g; 47.6mmol), potassium phosphate (13.5g; 63.5mmol) and dicyclohexyl(2',6'-dimethoxy[1,1'-biphenyl]-2-yl)phosphine (1.3g; 3.2mmol) in dioxane (380mL) and H_2O (150mL) was stirred at room temperature under N_2 flow. After 10 minutes, $Pd_2(dba)_3$ (1.45g; 1.6mmol) was added portionwise at room temperature under N_2 flow. The reaction mixture was heated at 80°C overnight. The reaction mixture was cooled to room temperature and poured out into ice water. The mixture was filtered over celite. Celite was washed with DCM. The organic layer was washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated, yielding 21g (99%) of intermediate 11.


Intermediate prepared according to the above protocol	starting from
 Intermediate 56	 (intermediate 14=A4 c-2)

Example A5



a) Preparation of intermediate 15



Methanesulfonyl chloride (3.53mL, 45.77mmol) was added dropwise to a solution of compound 3 (9.6g, 22.88mmol) and triethylamine (7.96mL, 57.21mmol) in DCM (250mL) at 5°C under a N_2 flow. The reaction mixture was stirred for 1 hour allowing the temperature to rise to room temperature. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The crude residue was taken up into DIPE. The precipitated was filtered yielding, after drying, 10.5g (92%) of intermediate 15.




b) Preparation of intermediate 16

Methanesulfonyl chloride (0.97mL, 12.52mmol) was added dropwise to a suspension of compound 2 (0.98g, 2.50mmol) and Et_3N (2.09mL, 15.02mmol) in DCM (50mL) at 5°C under N_2. The mixture was stirred at room temperature for 3 hours. The solution was

evaporated at room temperature yielding 1.38g of intermediate 16. The residue was used without purification for the next step.


Intermediate prepared according to the above protocol	starting from
 Intermediate 48	 Compound 49



c) Preparation of intermediate 143

Methanesulfonyl chloride (519µL, 6.7 mmol) was added dropwise to a solution of compound 389 (1.5 g, 3.35 mmol), triethylamine (1.2 mL, 8.4 mmol), 4-dimethylaminopyridine (40.95 mg, 0.335 mmol) in DCM (50 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then at room temperature for 36 hours. Water and DCM were added and the organic layer was washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was crystallised from acetonitrile and Et_2O. The resulting solid was filtered and dried to give 622 mg (35%) of a yellow solid intermediate 143.

Example A6





a-1) Preparation of intermediate 17


NaH (16.88g; 0.42mol) in suspension in heptane was slowly added to a solution of intermediate 17a (100g; 0.201mol) and 1,1-dimethylethyl ester N-(2,2,2-trifluoroethyl)-carbamic acid (48.03g; 0.241mol) in *N,N*-dimethylacetamide (1 L) at 0°C. The reaction mixture was stirred for 1 hour at 0°C, allowed to warm to room tempature in 1 hour and stirred at room temperature for 5 hours.The reaction mixture was carefully quenched with water (1 L) and the solution was extracted twice with DCM. The combined organic layers were washed with water, decanted and evaporated to dryness. The residue was dissolved in toluene, the organic layer was washed with water and evaporated to dryness to provide 147g of intermediate 17 which was used without further purification in the next step.

Intermediate 17 was alternatively also prepared using the following procedure

a-2) NaH (1g; 24.94mmol) was added portionwise to a solution of intermediate 3 (4.5g; 12.47mmol) and intermediate 69 (5.02g; 14.96mmol) in DMF (47mL) at 5°C. The reaction mixture was heated at 60°C for 1 hour, then cooled to room temperature, poured onto iced water and extracted with EtOAc. The organic layer was decanted, washed with water then brine, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was combined with an analogously prepared product fraction (using 1.4g of intermediate 3)and then purified by chromatography over silica gel (Irregular SiOH, 15/40µm) mobile phase gradient from 99%DCM/1%CH_3OH to 97%DCM/3%MeOH). The pure fractions were collected and evaporated to dryness yielding 5.8g (77%) of intermediate 17. MP=113°C.









Intermediate prepared according to the above protocol	starting from
 intermediate 50	 Intermediate 48 (prepared according to A5b)









Example A7




a) Preparation of intermediate 18

NaH (830mg; 20.75mmol) was added portionwise to a solution of intermediate 3 (5g; 13.84mmol) in *N,N*-dimethylformamide (150mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour then a solution of 2-(3-bromopropoxy)tetrahydro-2H-pyran (3.5mL; 20.75mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 1 hour at 5°C, then allowed to warm to room temperature. The reaction was stirred at room temperature for 4 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to give 8.46 g of intermediate 18.

Intermediate prepared according to the above protocol	starting from
	
 Intermediate 59	
a mixture of  ,  and  intermediate 60	


 Intermediate 51	
 Intermediate 55	
 Intermediate 57	
 Intermediate 58	

Example A8





a) Preparation of intermediate 19

NaH (882mg; 22.04mmol) was added portionwise to a solution of intermediate 13 (5.8g; 18.4mmol) in DMF (100mL) under N_2 at 5°C. The reaction mixture was stirred for 20 minutes and (bromomethyl)cyclopropane (2.2mL; 22.04mmol) was added dropwise. The mixture was stirred for another 20 minutes at 5°C, then at room temperature for 1.5 hour. The reaction mixture was poured into H_2O and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness, yielding 6.7g (98%) of intermediate 19.




b) Preparation of intermediate 20

A mixture of intermediate 19 (3g; 8.1mmol), 1-Boc-pyrazole-4-boronic acid pinacol ester (2.86g; 9.7mmol), potassium phosphate (3.44g; 16.2mmol), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.33g; 0.811mmol) in dioxane (60mL) and H_2O (6mL) was stirred at room temperature under N_2 flow. After 10 minutes, tris(dibenzylideneacetone)dipalladium (0.3g; 0.41mmol) was added portionwise at room temperature and the mixture was heated at 80°C overnight .The reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the mixture was filtered through a layer of celite. The celite was washed with EtOAc, then the filtrate was extracted with EtOAc, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase 0.05% NH_4OH, 99% DCM, 1% iPrOH). The pure fractions were collected and evaporated to dryness, yielding 1.48g (36%) of intermediate 20.

Intermediate prepared according to the above protocol	starting from
 intermediate 47	

Example A9




a-1) Preparation of intermediate 21

7-Bromo-2-chloroquinoxaline (10g, 41.1mmol), 1-(tetrahydro-2H-pyran-2-yl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (11.42g, 41.1mmol), sodium carbonate 2M (20.5mL, 41.1mmol) in ethylene glycol dimethyl ether (100mL) were degassed with N_2 for 15 minutes, $Pd(PPh_3)_4$ (1.4g, 1.2mmol) was added and heated at reflux for 20 hours. The reaction mixture was cooled to room temperature, poured into H_2O and EtOAc. The precipitate was filtered and dried under vacuum to give 12 g (84%) of intermediate 21.


Intermediate 21 was alternatively also prepared using the following procedure.

a-2) Trifluoroacetic acid (5.55µl; 0.075mmol) was added dropwise to a solution of 7-bromo-2-(1H-pyrazol-4-yl)quinoxaline (410mg; 1.5mmol) and 3,4-dihydro-2H-pyran (0.16mL; 1.8mmol) in toluene (4mL) and the reaction mixture was heated to 60°C for 2 days, then cooled to room temperature and evaporated till dryness, yielding 550mg of intermediate 21.



b) Preparation of intermediate 22


A mixture of intermediate 21 (1.5g; 4.2mmol), aniline (0.58mL; 6.23mmol), sodium tert-butoxide (1.2g; 12.5mmol) and 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] (260mg; 0.42mmol) in ethylene glycol dimethyl ether (45mL) was degassed with N_2 for 30 minutes, then palladium(II) acetate (93.7mg; 0.42mmol) was added. The reaction mixture was refluxed for 4 hours. H_2O/ice was added and the product was extracted with EtOAc. The organic layer was washed with H_2O, a saturated aqueous solution of NaCl, dried ($MgSO_4$), filtered and the solvent was evaporated till dryness. The crude product was purified by chromatography over silica gel (Irregular SiOH 15-40 µm, 90g; mobile phase gradient from 99%DCM/1%MeOH to 97%DCM/3%MeOH/0.1%NH_4OH). The pure fractions were collected and the solvent was evaporated till dryness yielding 1.1g (70%) of intermediate 22 A fraction (0.7g) was re-purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm;mobile phase Gradient from 100% DCM to 0.4% NH_4OH, 96% DCM, 4% CH_3OH). The pure fractions were collected and the solvent was evaporated till dryness, yielding 0.071g (4.5%) of intermediate 22.



c) Preparation of compound 123


NaH (116.3mg; 2.9mmol) was added portionwise to a solution of intermediate 22 (0.9g; 2.4mmol) in DMF (14mL) at 5°C. The reaction mixture was stirred for 30 minutes. (Bromomethyl)cyclopropane (0.28 mL; 2.9mmol) was added dropwise and the reaction mixture was stirred for 1 hour at 5°C, then at room temperature overnight. The reaction mixture was poured into H_2O and extracted twice with EtOAc. The organic layer was washed with a saturated aqueous solution of NaCl, dried ($MgSO_4$), filtered and the

solvent was evaporated till dryness. The crude product was purified by chromatography over silica gel (Irregular SiOH, 30g, 15-40µm; mobile phase 98% DCM/ 2% CH_3OH). The pure fractions were collected and the solvent was evaporated till dryness to give 0.5g (48%) of compound. A fraction (0.4g) was re-purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g, PharmPrep MERCK; mobile phase 99% DCM, 1% MeOH). The pure fractions were collected and the solvent was evaporated, yielding 85mg (8%)of compound 123.










d) Preparation of compound 54


At 5°C, HCl/i-PrOH (80µl 5/6N; 0.4mmol) was added to a solution of compound 123 (85mg; 0.2mmol) in CH_3OH (5mL). The reaction mixture was stirred at 5°C for 4 hours. Diethyl ether (8 mL) was added and the mixture was stirred for 30 minutes , then the precipitate was filtered and dried under vacuum, yielding 58mg (71%) of compound 54 MP=138°C (Kofler).



e) Preparation of intermediate 23

The reaction was done under a nitrogen atmosphere. NaH (0.058g; 1.46mmol) was added portionwise to a solution of compound 54 (0.25g; 0.73mmol) in DMF (5mL) at 5°C. The reaction mixture was stirred for 30 minutes, then 2-(2-bromoethoxy)tetrahydro-2H-pyran (0.23mL; 1.46mmol) was added dropwise and the reaction mixture was further stirred overnight at room temperature. The reaction mixture was poured into an aqueous solution of potassium carbonate and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The crude product was purified by chromatography over silica gel (Irregular SiOH, 15-40µm 30g; mobile phase 0.1% NH_4OH, 99% DCM, 1% CH_3OH). The pure fractions were collected and the solvent was evaporated till dryness, yielding 250mg (72%) of intermediate 23.

Intermediate/compound prepared according to the above protocol	starting from
 Intermediate 54	
 Intermediate 52	 Compound 17
Intermediate 65 	 Compound 17
	7-bromo-2-(1H-pyrazol-4-yl)quinoxaline




Intermediate 691 used to prepare compound 691 according to B14A, was prepared in an analogous way:

The experiment has been performed 4 times on the following amounts.



A mixture of compound 137 (HCl salt) (2 g; 4.6 mmol), 2-bromoethoxy-t-butyl dimethylsilane (1.3 mL; 7.4 mmol) and K_2CO_3 (1.3 g; 9.3 mmol) in CH_3CN (80 mL) was stirred at 80°C for 24 hours. The reaction was poured out into ice water and EtOAc was added. The organic layers were combined, separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (12.3 g) was purified by chromatography over silica gel (SiOH 15-40µm, 450; mobile phase gradient from 0.5% NH_4OH, 97% DCM, 3% MeOH to 0.5% NH_4OH, 90% DCM, 10% MeOH). The pure fractions were collected and concentrated to give 6 g of intermediate 691

Example A10




Preparation of intermediate 24

To a solution of intermediate 65 (1.1g; 2.25mmol) in THF (15mL) and H_2O (15mL) was added lithium hydroxide monohydrate (0.34g; 4.5mmol). The reaction mixture was stirred overnight at room temperature. THF was evaporated and H_2O and HCl were added. The precipitate was filtered and dried, yielding 976mg (94%) of intermediate 24.

Intermediate prepared according to the above protocol	starting from
 Intermediate 53	


Example A11



Preparation of intermediate 25


A solution of intermediate 2 (1g; 0.35mmol), cyclopropanemethylamine (0.51g, 6.9mmol) 1,1'-[1,1'-Binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] (0.215g, 0.35mmol) and sodium tert-butoxide (1.0g, 10.4mmol) in ethyleneglycol-dimethylether (15mL) was degassed with N_2 for 10 minutes. Then palladium(II) acetate (47% Pd) (77.6mg, 0.35mmol) was added and the reaction was heated under microwave irradiation to 135°C for 30 minutes. The reaction mixture was cooled to room temperature, then poured into an aqueous solution of K_2CO_3 and extracted with EtOAc. The organic layers were combined and dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm; mobile phase, gradient from 10% DCM to 95%DCM/5%MeOH/0.1%NH_4OH). The pure fractions were collected and evaporated, yielding 710mg (74%) of intermediate 25. MP=149°C (kofler).

Example A12



a) Preparation of intermediate 26


Methanesulfonyl chloride (61µL, 0.78mmol) was added dropwise to a solution of compound 24 (0.13g, 0.26mmol), Et_3N (0.18mL, 1.3mmol) in DCM (10mL) at 5°C under N_2. The solution was stirred for 1.5 hours at 10°C. The solution was poured out into ice water, the organic layer was extracted, dried ($MgSO_4$) and evaporated to dryness at room temperature, yielding 137 mg of intermediate 26.



b) Preparation of intermediate 27

A solution of intermediate 26 (0.31g; 0.0006mol), phthalimide (0.17g, 0.0012mol) and K_2CO_3 (0.21 g; 0.0015mol) in 1-methyl-2-pyrrolidinone (10mL) were heated at 150°C for 15 hours. The mixture was cooled to room temperature and evaporated to dryness. The residue was taken up with DCM, then an aqueous K_2CO_3 solution (10%) were added. The organic layer was separated, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase 0.1%NH_4OH/ 99% DCM/ 1% MeOH). The product fractions were collected and the solvent was evaporated, yielding 212mg (63%) of intermediate 27.


Example A13



a) Preparation of intermediate 28

Hydrazine monohydrate (2.57mL, 0.083mol) was added to a solution of compound 65 (3.71g, 8.29mmol) in EtOH (35mL). The mixture was stirred overnight at reflux. Hydrazine monohydrate (2.57mL, 0.083mol) was added again and the mixture was refluxed for 15 hours. After cooling down to room temperature, the precipitate was filtered off, washed with EtOH and dried to give 2.6g (72%) of intermediate 28.


Example A14



a) Preparation of intermediate 29


NaH (0.077g; 2mmol) was added portionwise to a solution of compound 107(0.63g; 1.2mmol) in DMF (10mL). The mixture was stirred at 10°C for 60 minutes, then ethyl bromoacetate (0.16mL, 1.45mmol) was added. The resulting mixture was stirred at room temperature for 2 hours. The mixture was poured into water and the product was extracted with EtOAc. The organic layer was washed with water, brine, dried ($MgSO_4$), filtered and evaporated until dryness. The residue (1g) was crystallized from diethyl ether. The precipitate was filtered and dried, yielding 0.55g (75%) of intermediate 29.

Example A15



a) Preparation of intermediate 30


To a mixture of intermediate 2 (700mg; 2.4mmol), intermediate 39 (781mg; 2.66mmol), sodium tert-butoxide (698mg; 7.3mmol), 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] (151mg; 0.24mmol) in dioxane (12mL) was degassed at room temperature under N_2 flow. After 10 minutes, palladium (II) acetate (109mg; 0.48mmol) was added at room temperature under N_2 flow. The reaction was performed under microwave irradiation at 130°C for 1 hour. The reaction mixture was poured out onto ice water and filtered over celite. Celite was washed with DCM. The organic layer was decanted, dried ($MgSO_4$), filtered and evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase 0.1%NH4OH, 97% DCM, 3%iPrOH). The product fractions were collected and the solvent was evaporated, yielding 320mg (26%) of intermediate 30.




b) Preparation of intermediate 31

Intermediate 30 (300mg, 0.598mmol) in HCl (3N) (10.96mL, 33mmol) and THF (10mL) was stirred at 65°C for 2 hours, then for 6 hours at 70°C,and poured out onto ice. The solution was made basic with K_2CO_3 powder and extracted with DCM. The organic layer was dried ($MgSO_4$), filtered and evaporated, yielding 270mg (98%) of intermediate 31.

Example A16


a) Preparation of intermediate 32 

Acetic anhydride (3.24mL) was added portionwise over ten minutes to a stirred suspension of 3,5-dimethoxyaniline (5g, 32.64mmol) in toluene (25mL). After stirring at room temperature for 17 hours, petroleum ether was added and the precipitate collected by suction filtration and dried under vacuum. The crude product (6.1g, 96%) was used in the next step without further purification.

b) Preparation of intermediate 33 


N-(3,5-dimethoxy-phenyl)-acetamide (intermediate 32) (15g, 76.8mmol) was dissolved in AcOH (50mL) . The solution was cooled to 0°C and 32% aqueous hydrochloric acid solution (41mL, 461mmol) was added. A solution of sodium chlorate (3.5g, 33mmol) in water (4mL) was added. The mixture was stirred for 30 minutes at 0°C. The reaction mixture was poured out onto ice and water and made basic with K_2CO_3 powder. The precipitate was filtered off and washed with water.

The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase 80%DCM, 20% EtOAc) to give 8.8g (50%) of intermediate 33.

c) Preparation of intermediate 34 

Potassium hydroxide (10.7g, 192mmol) was added to a solution of *N*-(2-Chloro-3,5-dimethoxy-phenyl)-acetamide (intermediate 33) (8.8g, 38.3mmol) in EtOH (500mL) and water (50mL) and the reaction mixture heated to reflux for 18 hours. Upon cooling, water was added (ca 30mL) and the EtOH removed in vacuum. The residue was then partitioned between water and diethyl ether. The organic layer was separated, dried ($MgSO_4$), filtered and concentrated to afford 7g (97%) of intermediate 34 (white solid).


Example A17



Preparation of intermediate 35


A mixture of 2,4-dimethoxy-6-nitrotoluene (2g, 10.1mmol) and nickel (2g) in MeOH (30mL) was hydrogenated under a 3 bars pressure for 6 hours. The product was filtered over a celite pad which was washed 3 times with a solution of MeOH/DCM (50/50). The combined filtrates were evaporated till dryness to give 1.68g (99%) of intermediate 35.

Example A18



a) Preparation of intermediate 36

A mixture of 3-amino-5-methoxy-benzoic acid (300mg, 1.8mmol), 1-hydroxybenzotriazole (292mg, 2.1 mmol), *N*-Ethyl-*N*'-β-dimethylaminocarbodiimide hydrochloride (413mg, 2.1mmol), and ethyl amine (2.7mL, 5.4mmol , 2M in MeOH) in dimethylformamide (6mL) was stirred at room temperature overnight. The solvent was evaporated under vacuum and the residue partitioned between DCM and water. The organic layer was separated and the aqueous layer was extracted with further DCM. The combined organic layers were dried (Na_2SO_4) and concentrated. The residue was purified by column chromatography over silica gel eluting with 2% MeOH/DCM. The desired product fractions were collected and the solvent was evaporated, yielding 150mg (43%) of intermediate 36 (colourless oil).




b) Preparation of intermediate 135

A mixture of 3-amino-5-fluorobenzoic acid (10 g; 64.5 mmol), methylamine in THF (96.7 mL; 193.4 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (14.8 g; 77.4 mmol), 1-hydroxybenzotriazole (10.5 g; 77.4 mmol), in *N,N*-dimethylformamide (150 mL) was stirred at room temperature for 18 hours. The reaction mixture was poured out into a solution of sodium hydroxide 1 N and DCM was added. The organic

layer was separated, washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated.

The aqueous layer was neutralized with concentrated HCl and extracted with EtOAc. The organic layer was separated, dried ($MgSO_4$), filtered and evaporated till dryness to give 5g of 3-amino-5-methoxy-N-methyl-benzamide (intermediate 135).


Example A19



Preparation of intermediate 37

A solution of deoxofluor in toluene (0.478mmol; 0.176mL) was added dropwise to a solution of compound 124 (0.159mmol; 90mg) in DCM (8mL) at 5°C under N_2 flow. After 5 minutes, EtOH (a drop) was added. The mixture was stirred at 5°C for 1 hour, then overnight at room temperature. The reaction mixture was poured out into ice water and DCM was added. The mixture was basified with K_2CO_3 10% and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The obtained residue (0.090g) was purified by column chromatography over silica gel (Irregular SiOH,15/40µm, 30g; mobile phase gradient from 100%DCMto 97%DCM /3%MeOH). The product fractions were collected and the solvent was evaporated. The residue (0.070g, 77%) was crystallized from diethyl ether/CH_3CN, filtered and dried under vacuum, yielding 0.055g (60%) of intermediate 37.


Example A20



Preparation of intermediate 38

In a round bottom flask, 3,5-dimethoxybenzenamine (500mg, 3.26mmol), 3-oxetanone (588mg, 8.16mmol) and acetic acid (374µL, 6.53mmol) were diluted in MeOH (21mL). The reaction mixture was stirred at room temperature for 1 hour. Then, sodium cyanoborohydride (410mg, 6.53mmol) in MeOH (5mL) was added and the reaction mixture was stirred overnight at room temperature. Then, NaOH 3N (15mL) was added and the mixture was stirred for 1 hour at room temperature. The reaction mixture was partitioned between water and DCM. The organic layer was dried ($MgSO_4$), filtered and concentrated. The residue (1g) was purified by chromatography over silica gel (Irregular SiOH, 15 – 40 µm; mobile phase 100% DCM). The desired fractions were collected and the solvent was evaporated, yielding 377 mg (55%) of intermediate 38 (colorless oil).


Example A21



Preparation of intermediate 39

Sodium cyanoborohydride (4.55g, 72.5mmol) was added to a solution of 3,5-dimethoxyaniline (3.7g, 24.15mmol), 1,4-cyclohexanedione mono-ethylene ketal (15g, 96.6mmol) and acetic acid (5.5mL, 96mmol) in CH_3CN (50mL) at room temperature (exothermicity observed). The reaction mixture was stirred overnight. Aqueous $NaHCO_3$ solution was added and the mixture was extracted twice with EtOAc. The combined organic layers were washed with brine, dried ($MgSO_4$), filtered and dried. The residue (21g) was purified by column chromatography over silica gel (Irregular SiO_2, 15-40µm, 90g; mobile phase gradient from 100% DCM to 7%CH_3OH / 93% DCM). The pure fractions were collected and evaporated to dryness to give 4.2g (59%) of intermediate 39.


Example A22



Preparation of intermediate 42

A solution of 3-bromo-5-methoxy phenol (3.12g; 15.4 mmol), 2-(2-bromoethoxy)tetrahydro-2H-pyran (2.66 mL; 16.9 mmol) and K_2CO_3 (1.63g; 11.8 mmol) was heated at 80°C in CH_3CN (40 mL) overnight. The solution was cooled and the mixture was poured into cooled water, the product was extracted with EtOAc, the organic layer was washed with H_2O and dried ($MgSO_4$), filtered and evaporated to dryness (5.5g). The residue was purified by chromatography over silica gel (irregular SiOH, 15-40 µm, 200 g; mobile phase 80 % cyclohexane, 20 EtOAc). The product fractions were collected and the solvent was evaporated, yielding 3.7g (73%) of intermediate 42.


Example A23



Preparation of intermediate 43

Sodium hydride (1.03g, 25.86 mmol) was added portion wise to a solution of 3-bromo-5-methoxy phenol (3.5g, 17.24 mmol) in DMF (20 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 0.5 hour, then a solution of deuteriated-iodomethane (1.29mL, 20.69 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 1 hour at 5°C, then allowed to warm up to room temperature and stirred for 2 hours. The reaction was poured out into ice water, and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to give 4g of intermediate 43, used without further purification for the next step.


Example A24



Preparation of intermediate 44

A solution of 3-bromo-5-methoxyphenol (2g, 9.85 mmol), 1-bromo-2-fluoroethane (1.56g, 0.012mol) and K_2CO_3 (1.4g, 10 mmol) was heated at 80°C in CH_3CN (30 mL) overnight. The solution was cooled and the mixture was poured into cooled water, the product was extracted with Et_2O. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness to give 2.27 g of intermediate 44 used without further purification for the next step.


Example A25



Preparation of intermediate 45

Under N_2 at 10°C, Hunig's base (9.64mL; 55.16 mmol) was added to a solution of 3-bromo-5-methoxy phenol (5.6g, 27.58 mmol) in THF (100mL). 2-methoxyethoxymethylchloride (CAS 3970-21-6) (6.3 mL, 55.16 mmol) was added and the solution was stirred at room temperature overnight. The solution was poured into cooled water, and the product was extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness to give 8g (99.6%) of intermediate 45 used without further purification for the next step.


Example A26



Preparation of intermediate 46

A solution of 3-bromo-5-methoxyphenol (0.3g, 1.5 mmol), 2-iodopropane (0.21 mL, 1.6 mmol) and K_2CO_3 (1.63g, 12 mmol) was heated at 80°C in CH_3CN (20mL) for 24 hours. The solution was cooled and the mixture was poured into cooled water, the product was extracted with EtOAc. The organic layer was washed with H_2O and dried ($MgSO_4$), filtered and evaporated to dryness to give 350 mg (97%) of intermediate 46 used without further purification for the next step.


Example A26A



Preparation of intermediate 136


NaH (0.74 g; 18.4 mmol) was added portionwise to a solution of (3-chloro-5-methoxyphenyl) methanol (2.9 g; 16.7 mmol) in *N,N*-dimethylformamide (30 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour. Then ethyl iodide (0.96 mL; 12.0 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was allowed to warm to room temperature and stirred for 18 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.8 g (25%) of intermediate 136.

Example A27




a) Synthesis of intermediate 66

A mixture of (3-bromopropoxy)-tert-butyldimethylsilane (20g; 79mmol) and 2,2,2-trifluoroethylamine (31mL; 395mmol) in DMSO (140mL) was heated at 80°C for 18 hours. The reaction mixture was cooled to room temperature, water was added and the mixture was extracted with Et_2O. The organic layer was dried ($MgSO_4$), filtered and evaporated till dryness to provide 19.5g (91%) of intermediate 66.




b) Synthesis of intermediate 67

Di-tert-butyl-dicarbonate (7.96; 36.5mmol), triethylamine (6mL; 43.11mmol) and N,N-dimethyl-4-aminopyridine (202mg; 1.7mmol) were added to a solution of intermediate 66 (9g; 33.16mmol) in DCM (90mL). The reaction mixture was stirred at room temperature for 2 hours, diluted with DCM and water. The organic layer was decanted, washed successively with water, a solution of HCl (0.5N) and an aqueous solution of K_2CO_3 (10%). The organic layer was dried ($MgSO_4$), filtered and evaporated till dryness to provide 11.3g (92%) of intermediate 67.



c) Synthesis of intermediate 68


A mixture of intermediate 67 (10.8g; 29.1mmol) and tetrabutylammonium fluoride (34.9 mL of a 1M solution in THF; 34.9mmol) in THF (80mL) was stirred at room temperature overnight. Water was added and the reaction mixture extracted with DCM. The organic layer was dried ($MgSO_4$), filtered and evaporated till dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µM, 80g; mobile phase, gradient from 99%DCM, 1%MeOH to 96%DCM, 4% MeOH). The pure fractions were collected and evaporated till dryness to provide 3.65g (49%) of intermediate 68.



d) Synthesis of intermediate 69

Methane sulfonyl chloride (431µL; 5.8mmol) was added dropwise to a solution of intermediate 68 (1g; 3.9mmol) and triethylamine (811µL; 5.8mmol) in DCM (15mL) at 5°C under N_2 flow. The reaction mixture was stirred for 30 minutes at room temperature. The reaction mixture was evaporated till dryness and the resulting intermediate 69 was used without further purification for the next step.


Example A28



a) Preparation of intermediate 70

The experiment has been performed 5 times on the following amount.


NaH (0.25 g; 5.4 mmol) was added portionwise to a solution of 2-amino-2-methyl-1-propanol (1.54 mL; 16.1 mmol) in *N, N*-dimethylformamide (12 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 15 minutes. Then, compound 76 (1.4 g; 3.35 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 10.5 g of a residue which was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g; mobile phase 1% NH_4OH, 90% DCM, 10% MeOH). The pure fractions were collected and concentrated yielding 3.6 g (42%) of intermediate 70.



b) Preparation of intermediate 71


Di-tert-butyl dicarbonate (0.24 g; 1.1 mmol) was added to a solution of intermediate 70 (0.62 g; 1.1 mmol) and $NaHCO_3$(0.19 g; 2.3 mmol) in dioxane (15 mL) and water (15 mL). The mixture was stirred at room temperature for 18 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (0.8 g) was purified by chromatography over silica gel (SiOH, 5 µm 150*30 mm; mobile phase

0.2% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and the solvent was evaporated to give 0.59 g (85%) of intermediate 71.



c) Preparation of intermediate 72


Methanesulfonyl chloride (0.96 mL; 12.4 mmol) was added dropwise to a solution of intermediate 71 (2.7 g, 4.45 mmol) and triethylamine (1.86 mL; 13.35 mmol) in DCM (25 mL) at 5°C under a N_2 flow. The reaction mixture was stirred for 18 hours allowing the temperature to rise to room temperature. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (4.1 g) was purified by chromatography over silica gel (Irregular SiOH, 20-45 µm, 450 g ; mobile phase 0.2% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and the solvent was evaporated to give 3 g (100%) of intermediate 72.



d) Preparation of intermediate 73

Trifluoroacetic acid (0.97 mL; 13.1 mmol) was added to a solution of intermediate 72 (0.6 g; 0.87 mmol) in DCM (12.5 mL) at 0°C. The reaction was stirred at room temperature for 1 hour. The mixture was poured out into ice water and DCM was added. The mixture was basified with a solution of $NaHCO_3$ and the organic layer was separated, dried over $MgSO_4$, filtered and the solvent was evaporated to give 597 mg of intermediate 73 used without further purification for the next step.


Example A29



Preparation of intermediate 74


Methanesulfonyl chloride (3.32 mL; 42.9 mmol) was added dropwise to a solution of compound 606 (6 g; 14.3 mmol) and triethylamine (10 mL; 71.5 mmol) in DCM (240 mL) at 5°C under a N_2 flow. The reaction mixture was stirred for 1 hour at 5°C and allowed to rise to room temperature for 1 hour. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 9.6 g of intermediate 74 used without further purification for the next step

Example A30



a) Preparation of intermediate 75


NaH (11.4 g; 82.5 mmol) was added portionwise to a solution of 4,4,5,5-tetramethyl-2-(1H-pyrazol-4-yl)-1,3,2-dioxaborolane (4 g; 20.6 mmol) in acetone (60 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 15 minutes. Then, acetamide, 2-bromo-*N*-methyl (6.3 g; 41.3 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at 65°C for 24 hours. The reaction mixture was cooled to room temperature. The precipitate was filtered off and washed with DCM. The filtrate was evaporated till dryness, taken up into DIPE/diethyl ether and stirred at room temperature for 15 minutes. The precipitate was filtered off and washed with DCM. The filtrate was evaporated till dryness to afford 9 g of intermediate 75 used without further purification for the next step.



b) Preparation of intermediate 76

A mixture of intermediate 14 (5.7 g; 11.7 mmol), intermediate 75 (*N*-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-acetamide) (6.2 g; 23.5 mmol), potassium phosphate (7.5g; 35.2mmol) and dicyclohexyl(2',6'-dimethoxy[1,1'-biphenyl]-2-yl)phosphine (0.482 g; 1.2 mmol) in dioxane (140 mL) and H_2O (60 mL) was stirred at room temperature under N_2 flow. After 10 minutes, $Pd_2(dba)_3$ (1 g; 1.2 mmol) was added portionwise at room temperature under N_2 flow. The reaction mixture was heated at 80°C for 4 hours. The reaction mixture was cooled to room temperature and poured out into ice water. The mixture was filtered over a pad of celite®, washed with DCM. The organic layer was washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (8.3 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 450g; mobile phase 0.1% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and concentrated to give 3.5 g (51%) of intermediate 76.


Example A31



Preparation of intermediate 77


Methanesulfonyl chloride (0.73 mL; 9.4 mmol) was added dropwise to a solution of compound 614(1.5 g; 3.15 mmol) and triethylamine (2.2 mL; 15.7 mmol) in DCM (40 mL) at 5°C under a N_2 flow. The reaction mixture was stirred for 1 hour at 5°C and allowed the temperature to rise to room temperature for 1 hour. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 2.5 g of intermediate 77 used without further purification for the next step.

Example A32




a) Preparation of intermediate 78

NaH (0.44 g; 10.9 mmol) was added portionwise to a solution of 7-bromo-2-(1H-pyrazol-4-yl)quinoxaline (1.5 g; 5.45 mmol) in *N, N*-dimethylformamide (40 mL) at 0°C under N_2 flow. The reaction mixture was stirred at 5°C for 15 minutes. Then, carbamic acid, N-(3-bromopropyl)-1,1-dimethylethyl ester (2.6 g; 10.9 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was poured out into ice water, EtOAc was added. The organic layer was separated, washed with water, diethyl ether, dried ($MgSO_4$), filtered and evaporated to afford 1.3 g of intermediate 78 used without further purification for the next step.




b) Preparation of intermediate 79

Under an inert atmosphere, a solution of palladium acetate (0.11 g; 0.48 mmol), racemic 2,2'-bis (diphenylphosphino)-1,1'-binaphtyl (0.3 g; 0.48 mmol)) was added to room temperature to a solution of intermediate 41 (3.3 g; 10.6 mmol), intermediate 78 (4.2 g; 9.63 mmol) and cesium carbonate (3.8 g; 11.6 mmol) in dimethoxyethane (50 mL). The reaction mixture was stirred at 85°C for 3 days. The reaction mixture was cooled to room temperature and poured out into ice water, K_2CO_3 10% and EtOAc was added. The mixture was filtered over a pad of celite®. The organic layer was washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (8.5 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 450g ; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated to give 3.3 g (52%) of intermediate 79.




c) Preparation of intermediate 80

A 1M solution of tetrabutylammonium fluoride in THF (5.5 mL; 5.5 mmol) was added dropwise to a solution of intermediate 79 (3.3 g; 5 mmol) in THF (60 mL) at room temperature. The reaction mixture was stirred at room temperature for 3 hours. The mixture was poured out into ice water and EtOAc was added. The mixture was basified with K_2CO_3 10% and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue was crystallized from diethyl ether /CH_3CN. The precipitate was filtered off, dried in vacuum to provide 2 g (73%) of intermediate 80.



d) Preparation of intermediate 81


Methanesulfonyl chloride (0.85 mL; 10.9 mmol) was added dropwise to a solution of intermediate 80 (2 g; 3.65 mmol) and triethylamine (2.54 mL; 18.2 mmol) in DCM (50 mL) at 5°C under a N_2 flow. The reaction mixture was stirred for 1 hour at 5°C and allowed the temperature to rise to room temperature for 2 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 2.5 g of intermediate 81 used without further purification for the next step.



e) Preparation of intermediate 82


A mixture of intermediate 81 (2.5 g; 4 mmol) and isopropylamine (5.2 mL; 59.9 mmol) in acetonitrile (25 mL) was heated at 100°C in a sealed vessel for 18 hours. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, EtOAc was added. The organic layer was separated, washed with a solution of $NaHCO_3$, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (3 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 450g; mobile phase from 0.1% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and concentrated to give 1.1 g (47%) of intermediate 82.

Example A33



a) Preparation of intermediate 83


A mixture of 5-bromo-benzene-1,3-diol (7.3 g; 38.6 mmol), cesium carbonate (37.75 g; 115.9 mmol) and iodomethane-D3 (4.8 mL; 77.25 mmol) in CH_3CN (150 mL) was stirred at 80°C for 18 hours. The reaction mixture was cooled to room temperature and poured out into ice water and EtOAc was added. The organic layer was washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to afford 5.3 g of intermediate 83 used without further purification for the next step.



b) Preparation of intermediate 84

Under an inert atmosphere, a solution of palladium acetate (0.21 g; 0.9 mmol), racemic 2,2'-bis (diphenylphosphino)-1,1'-binaphtyl (0.57 g; 0.9 mmol) was added to room temperature to a solution of intermediate 5 (2 g; 10.6 mmol), intermediate 83 (2.45 g; 11 mmol) and sodium tert butoxide (2.64 g; 27.4 mmol) in dioxane (150 mL). The reaction mixture was stirred at 100°C for 4 days. The reaction mixture was cooled to room temperature and poured out into ice water and EtOAc was added. The mixture was filtered over a pad of celite®. The organic layer was washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (6 g) was purified by


chromatography over silica gel (Irregular SiOH, 15-40µm, 300g; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated. The residue (4 g) was crystallized from diethyl ether. The precipitate was filtered off, dried in vacuum to provide 3.6 g (90%) of intermediate 84. MP: 198°C (DSC)



c) Preparation of intermediate 85

NaH (0.107 g; 2.69 mmol) was added portionwise to intermediate 84 (0.49 g; 1.35 mmol) in *N,N*-dimethylformamide (10 mL). The reaction mixture was stirred at 5°C for 1 hour. Then, a solution of deuterated (2-bromoethoxy)(1,1-dimethylethyl)dimethyl-silane (deuterated version of CAS 86864-60-0; prepared by art-known deuteration method) (0.65 g; 2.7 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture allowed to warm to room temperature and stirred for 4 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to afford 0.88 g of intermediate 85 used without further purification for the next step.


Example A34



Preparation of intermediate 86

Methanesulfonyl chloride (0.17 mL, 2.1 mmol) was added dropwise to a solution of compound 617 (0.294 g, 0.7 mmol) and triethylamine (0.49 mL, 3.5 mmol) in DCM (5 mL) at 5°C under a N_2 flow. The reaction mixture was stirred for 1 hour at 5°C and allowed the temperature to rise to room temperature for 1 hour. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.45 g of intermediate 86 used without further purification for the next step.


Example A35




Preparation of intermediate 87

A mixture of compound 4 (1.3 g; 2.9 mmol), N-(3-bromopropyl) phtalimide (1.56 g; 5.8 mmol) and K_2CO_3 (0.805 g; 5.8 mmol) in CH_3CN (100 mL) was stirred at 80°C for 48 hours. The reaction mixture was cooled to room temperature, poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated until dryness. The residue (0.566 g) was purified by chromatography over silica gel (SiOH, 15-40µm, 50g; mobile phase 0.1% NH_4OH, 96% DCM, 4% MeOH). The product fractions were collected and the solvent was evaporated to give 1.26 g (34%) of intermediate 87.

Example A36A




Preparation of intermediate 88



A mixture of intermediate 88b (see A4c-2) (0.53 g; 1.1 mmol), 1,3,5-trimethyl-4-(tributylstannyl)- 1H-Pyrazole (Synthesis, (13), 1949-1958; 2001) (1.33 g; 3.33 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.064 g; 0.055


mmol) in toluene (3 mL) was stirred at 160°C for 40 minutes using one single mode microwave (Biotage).The reaction mixture was cooled to room temperature and evaporated till dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 40µm ; mobile phase gradient from 90% DCM, 10% Heptane to 100% DCM, then 99% DCM 1% MeOH). The pure fractions were collected and concentrated to give 0.41 g (68%) of intermediate 88.



Intermediate 88b solution of tetrabutylammonium fluoride (3.016mmol;3.016ml) dropwise to a solution of intermediate 88b (2.742mmol;1.30g) in THF (25ml). The reaction mixture was stirred at room temperature overnight. The reaction mixture was poured out into ice water,EtOAc was added and the organic layer was separated,washed with brine, dried ($MgSO_4$),filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (SiO_2=30g-15/40µm) Eluent: CH_2Cl_2 100 to CH_2Cl_2 98 / MeOH 2, yielding intermediate 88a.

Alternative pyrazole derivatives which can be used in the above protocol can be prepared as follows :


A)




a) Preparation of intermediate 125

N-butyllithium 1.6M in hexane (33.5 mL; 53.6 mmol) was added dropwise to a solution of 1-methylpyrazole (4 g; 48.8 mmol) in THF (66 mL) at -78°C under N_2 flow. The reaction mixture was stirred at 0°C, then (2-bromoethoxy)-tert-butyldimethylsilane (12.5 mL; 58.5 mmol) was added to the solution at -78°C and was stirred for 1 hour. The temperature of the reaction mixture was allowed to rise to room temperature and stirred for 18 hours.


The reaction mixture was poured out into ice water and EtOAc was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (16 g) was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 1000 g; mobile phase 65% Heptane, 35% EtOAc). The pure fractions were collected and concentrated to give 3 g (25%) of intermediate 125 .

b) Preparation of intermediate 126 

Pyridinium bromide perbromide 95% (3.5 g; 10.8 mmol) was added to a solution of intermediate 125 (2.6 g; 10.8 mmol) in MeOH (130 mL). The reaction mixture was stirred at 0°C for 1 hour and room temperature for 18 hours. The solvent was evaporated and the residue was poured out into water and K_2CO_3 10%. DCM was added and the organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (2.5 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 300 g; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated to give 2 g (92%) of intermediate 126 .

c) Preparation of intermediate 127 


Tert-butyldimethylsilyl chloride (1.9 g; 12.7 mmol), imidazole (1.6 g; 23.4 mmol) were successively added to a solution of intermediate 126 (2 g; 9.75 mmol) in *N, N*-dimethylformamide (7mL).The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was quenched with water and extracted with Et_2O. The organic layer was decanted, washed with water, then brine, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (SiOH, 10-40µm, 90 g; mobile phase from 100% DCM to 97% DCM, 3% MeOH). The pure fractions were collected and concentrated to give 2.8 g (90%) of intermediate 127.



d) Preparation of intermediate 128


N-butyllithium 1.6M in hexane (0.22 mL; 0.35 mmol) was added dropwise to a solution of intermediate 127 (0.102 g; 0.32 mmol) in Et_2O (1.5 mL) at -78°C under N_2 flow. The reaction mixture was stirred for 30 minutes, then tributyltin chloride (0.095 mL; 0.35 mmol) was added to the solution and was stirred at room temperature for 18 hours. The reaction mixture was poured out into ice water and Et_2O was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (0.160 g) was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60 g; mobile phase 80% Heptane , 20% EtOAc). The pure fractions were collected and concentrated to give 0.055 g (32%) of intermediate 128 .

B)



a) Preparation of intermediate 129


N-butyllithium 1.6M in hexane (25 mL; 40.2 mmol) was added dropwise to a solution of 1-methylpyrazole (3 mL; 35.5 mmol) in THF (50 mL) at -78°C under N_2 flow. The reaction mixture was stirred at 0°C, then Eschenmoser's salt (8.1 g; 43.85 mmol) was added to the solution at -78°C and was stirred for 1 hour. The the temperature of the reaction mixture was allowed to rise to room temperature and stirred for 18 hours. The reaction mixture was poured out into ice water and EtOAc was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to afford 3.1 g of intermediate 129.



b) Preparation of intermediate 130

Pyridinium bromide perbromide 95% (6.9 g; 21.6 mmol) was added to a solution of intermediate 130 (3 g; 21.6 mmol) in MeOH (200 mL). The reaction mixture was stirred


at 0°C for 1 hour and room temperature for 18 hours. The solvent was evaporated and the residue was poured out into water and K_2CO_3 10%. DCM was added and the organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (3.1 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 450 g; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated to give 1.35 g (29%) of intermediate 130 .



c) Preparation of intermediate 131

N-butyllithium 1.6M in hexane (0.8 mL; 1.26 mmol) was added dropwise to a solution of intermediate 130 (0.25 g; 1.15 mmol) in Et_2O/THF (1/2) (3 mL) at -78°C under N_2 flow. The reaction mixture was stirred for 30 minutes. Then tributyltin chloride (1.58 mL; 5.8 mmol) was added to the solution and was stirred at room temperature for 18 hours. The reaction mixture was poured out into ice water and Et_2O was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.52 g of intermediate 131 used without further purification for the next step.


Example A36B



Preparation of intermediate 89

Methanesulfonyl chloride (0.066 mL; 0.85 mmol) was added dropwise to a solution of compound 622 (0.185 g; 0.43 mmol), triethylamine (0.14 mL; 0.98 mmol) and 4-dimethylaminopyridine (0.005 g; 0.043 mmol) in THF (5 mL) at 5°C under a N_2 flow. The temperature of the reaction mixture was allowed to rise to room temperature for 2 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.26 g (yellow oil) of intermediate 89 used without further purification for the next step.


Example A37



Preparation of intermediate 91

Methanesulfonyl chloride (1 mL; 12.8 mmol) was added dropwise to a solution of 1-piperidinecarboxylic acid, 4-(3-hydroxy-1-propyn-1-yl)-, 1,1-dimethylethyl ester (2 g; 8.5 mmol), triethylamine (1.8 mL; 12.8 mmol) and 4-dimethylaminopyridine (10.4 g; 85 mmol) in DCM (20 mL) at 5°C under a N_2 flow. The temperature pof the reaction mixture was allowed to rise to room temperature for 18 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 1.41g of intermediate 91 used without further purification for the next step.


Example A38



Preparation of intermediate 92

NaH (0.24 g; 6.0 mmol) was added portionwise to intermediate 3 (1 g; 3.0 mmol) in *N, N*-dimethylformamide (30 mL). The reaction mixture was stirred at 10°C for 1 hour. Then 2-butyn-1-ol, 4-[[(1,1-dimethylethyl)dimethylsilyl]oxy]-, 1-methanesulfonate (4.2 g; 15.0 mmol) was added dropwise under N_2 flow. The reaction mixture was stirred at room temperature for 18 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (4.2 g) was purified by chromatography over silica gel (Irregular SiOH 15-40µm, 300g; mobile phase 60% Heptane, 4% MeOH, 36% EtOAc). The pure fractions were collected and concentrated to give 0.185 g (11%) of intermediate 92.


Example A39




Preparation of intermediate 93

Methanesulfonyl chloride (9.9 mL; 127.7 mmol) was added dropwise to a solution of compound 2 (10 g; 25.55 mmol), triethylamine (24.9 mL; 178.8 mmol) in DCM (400 mL) at 5°C under a N_2 flow. The reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 17.6 g of intermediate 93 used without further purification for the next step.

Example A40




Preparation of intermediate 94



(see A4c-2)(9.5 g; 20 mmol), 4,4,5,5-tetramethyl-2-(1H-pyrazol-4-yl)-1,3,2-dioxaborolane (4.3 g; 22 mmol), potassium phosphate (8.5 g; 40 mmol) in dioxane (1 L) and water (120 mL) were degassed with N_2 for 15 minutes, then S-Phos (0.83 g; 2 mmol) and $Pd_2(dba)_3$ (7.6 g; 6.6 mmol) was added. The reaction mixture was heated at 80°C for 15 hours. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, EtOAc was added and was filtered off on a pad of Celite®. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (18.5 g) was purified by chromatography over silica gel (Irregular SiOH 20-45µm, 1000g ; mobile phase


96%DCM, 4%MeOH). The pure fractions were collected and concentrated to give 5.1 g (51%) of intermediate 94.

Example A41




a) Preparation of intermediate 95

NaH (0.2 g; 4.75 mmol) was added portionwise to intermediate 94 (2 g; 4 mmol) in *N, N*-dimethylformamide (30 mL). The reaction mixture was stirred at 10°C for 1 hour. Then 1-bromo-3-chloropropane (0.5 mL; 4.75 mmol) was added dropwise under N_2 flow. The reaction mixture was stirred at room temperature for 1 hour. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to afford 2.5 g of intermediate 95 used without further purification for the next step.



b) Preparation of intermediate 96


A mixture of intermediate 95 (1.1 g; 1.48 mmol), 1-(2-hydroxylethyl)piperazine (0.407g; 2.95 mmol), K_2CO_3 (1.92 g; 14.74 mmol) in CH_3CN (10 mL) was stirred at 90°C for 12 hours. The reaction mixture was poured out into ice water and EtOAc was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.9 g of intermediate 96 used without further purification for the next step.



c) Preparation of intermediate 97


A mixture of intermediate 96 (0.56g; 0.83 mmol), acetyl chloride (0.12 mL; 1.66 mmol), triethylamine (0.27 mL; 1.9 mmol) and 4-dimethylaminopyridine (0.01 g; 0.083 mmol)

was stirred in DCM (10 mL) at 5°C under a N_2 flow. The reaction mixture was stirred to room temperature for 18 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.85g of intermediate 97 used without further purification for the next step.



d) Preparation of intermediate 98


A 1M solution of tetrabutylammonium fluoride in THF (2.5 mL, 2.5 mmol) was added dropwise to a solution of intermediate 97, (0.75 g, 0.84 mmol) in THF (5 mL) at room temperature. The reaction mixture was stirred at room temperature for 3 hours. The mixture was poured out into ice water, basified with K_2CO_3 10% and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. DCM and a few MeOH were added, then the insoluble fraction was filtered off and the filtrate was evaporated. The residue and the precipitate were combined and dissolved in DCM. The organic layer was washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (0.5 g) was purified by chromatography over silica gel (SiOH, 15-40µm, 90g ; mobile phase from 0.3% NH_4OH, 97% DCM, 3%MeOH to 1% NH_4OH, 90% DCM, 10%MeOH). The pure fractions were collected and concentrated to give 0.238 g (47%) of intermediate 98.



e) Preparation of intermediate 99

Methanesulfonyl chloride (0.1 mL; 1.3 mmol) was added dropwise to a solution of intermediate 98 (0.19 g; 0.26 mmol) and triethylamine (0.11 mL; 0.78 mmol) in DCM (5 mL) at 5°C under a N_2 flow. The reaction mixture was stirred at 10°C for 2 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer


was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.51 g of intermediate 99-AAA used without further purification for the next step.



f) Preparation of intermediate 100


A mixture of intermediate 99 (0.51 g; 0.26 mmol) and isopropylamine (5.9 mL; 68.9 mmol) in acetonitrile (1 mL) was heated at 100°C in a sealed vessel for 12 hours. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, DCM was added. The organic layer was separated, washed, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (0.59 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g ; mobile phase 0.7% NH_4OH, 93% DCM, 7%MeOH). The pure fractions were collected and concentrated to give 0.09 g (54%) of intermediate 100.

Example A42




a) Preparation of intermediate 101

A mixture of intermediate 5 (3 g; 13.3 mmol) , intermediate 45 (3.9 g; 13.3 mmol), sodium tert-butoxide (3.9 g; 40 mmol) and 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine (0.83 g; 1.33 mmol) in ethylene glycol dimethyl ether (100mL) was degassed with N_2 for 10 minutes. Palladium(II) acetate (0.3 g; 1.33 mmol) was added and the mixture was stirred at 90°C for 2 hours. The mixture was cooled down to room temperature, poured into H_2O and DCM. The mixture was filtered off on a pad of celite®. The filtrate was extracted with DCM. The combined organic layers were dried ($MgSO_4$), filtered and evaporated to dryness to give 5 g of crude compound. The residue was purified by chromatography over silica gel (SiOH, 20-45µm, 40g; Mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and the solvent was evaporated, yielding 3.6g (62%) of intermediate 101.




b) Preparation of intermediate 102

NaH (0.37 g; 9.2 mmol) was added portionwise to a solution of intermediate 101 (2 g; 4.6 mmol) in *N,N*-dimethylformamide (20 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 30 minutes. Then (2-bromoethoxy)-*tert*-butyldimethylsilane (1.3 mL; 6.0 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 15 hours at room temperature. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 3 g of intermediate 102.



c) Preparation of intermediate 103


A 1M solution of tetrabutylammonium fluoride in THF (5 mL; 5 mmol) was added dropwise to a solution of intermediate 102 (3 g; 5 mmol) in THF (50 mL) at room temperature. The reaction mixture was stirred at room temperature for 15 hours. The mixture was poured out into ice water, basified with K_2CO_3 10% and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$) and the solvent was evaporated to dryness. The residue (3g) was purified by chromatography over silica gel (SiOH, 15-40µm,40g; mobile phase 0.1% NH_4OH, 95% DCM, 5%MeOH). The pure fractions were collected and concentrated to give 2.2 g (61%) of intermediate 103.



d) Preparation of intermediate 104

Methanesulfonyl chloride (0.7 mL; 9.2 mmol) was added dropwise to a solution of intermediate 103 (2.2 g; 4.6 mmol), triethylamine (1.6 mL; 11.5 mmol) in DCM (30 mL) at 5°C under a N_2 flow. The reaction mixture was stirred at 10°C for 2 hours. The


reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 2.8 g of intermediate 104 used without further purification for the next step.



e) Preparation of intermediate 105


A mixture of intermediate 104 (2 g; 3.6 mmol) and 2-propanamine (1.6 mL; 17.9 mmol) in acetonitrile (15 mL) was heated at 100°C in a sealed vessel for 18 hours. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, EtOAc was added. The organic layer was separated, washed with bine, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (2.2 g) was purified by chromatography over silica gel (SiOH, 15-40µm, 40g; mobile phase 0.1% NH_4OH, 95% DCM, 5%MeOH). The pure fractions were collected and concentrated to give 0.8 g (43%) of intermediate 105.

Example A43



Preparation of intermediate 107

Methanesulfonyl chloride (0.19 mL; 2.4 mmol) was added dropwise to a solution of compound 625 (0.69 g; 1.2 mmol) (prepared according to the procedure described in B39 starting from




which is prepared according to the

procedure described in A2c) starting from intermediate 41 and intermediate 106), triethylamine (0.4 mL; 3 mmol) in DCM (10 mL) at 5°C under a N_2 flow. The reaction mixture was stirred at room temperature for 3 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 0.8 g of intermediate 107 as an orange oil used without further purification for the next step.

Intermediate 107 was converted into compound 650 according to the procedure described in B3 (first alternative protocol).


Example A43A



Preparation of intermediate 106


NaH (0.3 g; 7.2 mmol) was added portionwise to a solution of 7-bromo-2-(1H-pyrazol-4-yl)quinoxaline (1.6 g; 6 mmol) in *N,N*-dimethylformamide (100 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour. Then 4-methylsulfonyloxy-1-piperidinecarboxylate CAS [141699-59-4] (3.5 g; 12.6 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 18 hours at 100°C. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (8.4 g) was purified by chromatography over silica gel (Irregular, SiOH, 20-40µm; 450g ; mobile phase 0.1% NH_4OH, 98% DCM, 2%MeOH). The pure fractions were collected and concentrated to give 3.7 g (67%) of intermediate 106 (yellow oil).

Example A44




Preparation of intermediate 109

NaH (0.29 g; 7.4 mmol) was added portionwise to intermediate 108



(1.5 g; 3.7 mmol) (prepared according to the procedure described in A33b) in *N,N*-dimethylformamide (25 mL). The reaction mixture was stirred at 0°C for 30 minutes. Then 3-bromo-(1-trimethylsilyl)-1-propyne (1.6 mL; 10.2 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (2 g) was purified by chromatography over silica gel (SiOH 15-40µm, 80g mobile phase 98% DCM, 2% MeOH)to give 1.4 g of intermediate 109.

Example A45




Preparation of intermediate 110

A mixture of compound 4 (0.5 g; 1.2 mmol), 4-nitrobenzyl bromide (0.29 g; 1.35 mmol) and K_2CO_3 (0.24 g; 51.8 mmol) in CH_3CN (20 mL) was stirred at room temperature for 48 hours. The reaction mixture was cooled to room temperature, poured out into ice


water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated until dryness. The residue (0.8 g) was purified by chromatography over silica gel (Stability SiOH, 5µm, 150*30mm; mobile phase gradient from 71%Heptane,1%MeOH, 28%EtOAc to 20%MeOH, 80%EtOAc). The product fractions were collected and the solvent was evaporated to give 0.34 g (52%) of intermediate 110.

Example A46



a) Preparation of intermediate 113

NaH (0.52 g; 13 mmol) was added portionwise to 7-bromo-2-(1H-pyrazolyl-4-yl)quinoxaline (3 g; 11 mmol) in *N, N*-dimethylformamide (30 mL). The reaction mixture was stirred at 5°C for 1 hour. Then 4-bromomethyltetrahydropyran (2.4 mL; 13 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then was allowed to room temperature and stirred for 18 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was crystallized from DIPE and CH_3CN. The precipitate was filtered and dried to give 2.6 g (64%) of intermediate 113.




b) Preparation of intermediate 112

Under an inert atmosphere, a solution of palladium acetate (0.08 g; 0.35 mmol), racemic 2,2'-bis (diphenylphosphino)-1,1'-binaphtyl (0.22 g; 0.35 mmol) was added to room temperature to a solution of intermediate 113 (2.6 g; 7.0 mmol), 3,5-dimethoxyaniline (1 g; 7.0 mmol) and sodium tert butoxide (2 g; 21 mmol) in dioxane (40 mL). The reaction mixture was stirred at 90°C for 18 hours. The reaction mixture was cooled to room temperature and poured out into ice water and DCM was added. The mixture was filtered over a pad of celite®. The organic layer was washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (3.5 g) was purified by


chromatography over silica gel (SiOH, 15-40µm, mobile phase 98% DCM, 2% MeOH). The pure fractions were collected and concentrated to give 1.6 g (63%) of intermediate 112.

Example A47



a) Preparation of intermediate 114


Intermediate 13 (9 g; 28.5 mmol), intermediate 132 (20.9 g; 57 mmol), potassium phosphate (12.1 g; 57 mmol) in dioxane (200 mL) and water (80 mL) were degassed with N_2 for 15 minutes, then S-Phos (1.2 g; 2.9 mmol) and $Pd_2(dba)_3$ (1.3 g, 1.4 mmol) weres added. The reaction mixture was heated at 80°C for 6 hours. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, EtOAc was added and was filtered off on a pad of Celite®. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (28 g) was purified by chromatography over silica gel (Irregular SiOH 20-45µm, 1000g ; mobile phase 99%DCM, 1%MeOH). The pure fractions were collected and concentrated to give 13.6 g (92%) of intermediate 114.




Intermediate 132 was prepared as follows :

NaH (77.3 mmol ; 3 g) was added to a solution of 4,4,5,5 tetramethyl-2-(1H-pyrazol-4-yl)-1,3,2-dioxaborolane (10g; 51.5 mmol) in *N, N*-dimethylformamide (150 mL) at room temperature under N_2 flow.The reaction was stirred at room temperature for 1 hour. Then a solution of (3-bromopropoxy)-tert-butyldimethylsilane (18.5 mL; 77.3 mmol) was


added dropwise at room temperature under N_2 flow.The reaction mixture was stirred at room temperature overnight. The reaction mixture was poured out into ice water and EtOAc was added.The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to give 23.8 g (70%) of intermediate 132 used without further purification.

Intermediate  was prepared according to the above protocol for intermediate 114.



b) Preparation of intermediate 115


A 1M solution of tetrabutylammonium fluoride in THF (24 mL; 24 mmol) was added dropwise to a solution of intermediate 114 (12.5 g; 24 mmol) in THF (250 mL) at room temperature. The reaction mixture was stirred at room temperature for 18 hours. The mixture was poured out into ice water, basified with K_2CO_3 10% and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried to afford 8.8 g (90%) of intermediate 115. MP: 118°C (Kofler).



c) Preparation of intermediate 116


Methanesulfonyl chloride (1.9 mL; 24.7 mmol) was added dropwise to a solution of intermediate 115 (2 g; 5.0 mmol), triethylamine (4.9 mL; 34.5 mmol) in DCM (80 mL) at 5°C under a N_2 flow. The reaction mixture was stirred at room temperature for 3 hours.

The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 3.4 g of intermediate 116 used without further purification for the next step.



d) Preparation of intermediate 117


NaH (0.42 g; 10.4 mmol) was added portionwise to di-tert-butyl-iminocarboxylate (2.3 g; 10.4 mmol) in *N,N*-dimethylformamide (40 mL). The reaction mixture was stirred at 10°C for 30 minutes. Then intermediate 116 (2.5 g; 5.2 mmol) was added dropwise under N_2 flow. The reaction mixture was stirred at room temperature for 18 hours, then stirred at 50°C for 4 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (4 g) was purified by chromatography over silica gel (15-40µm; 80 g; mobile phase 98%DCM, 20% MeOH). The pure fractions were collected and concentrated to give 1.7 g (54%) of intermediate 117.



e) preparation of intermediate 118


Trifluoroacetic acid (3 mL; 39.7 mmol) was added to a solution of intermediate 117 (1.5 g; 2.5 mmol) in DCM (20 mL). The reaction was stirred at room temperature for 5 hours. The reaction mixture was poured out into ice water, basified with K_2CO_3 10%, EtOAc was added. The layers were separated then the aqueous layer was evaporated till dryness. The residue was dissolved in MeOH. The precipitate was filtered off and the filtrate was evaporated till dryness. The residue was dissolved in DCM. The precipitate was filtered off and the filtrate was evaporated till dryness to afford 0.45 g (45%) of intermediate 118 . MP: 96°C (Kofler).

Example A48




a) Preparation of intermediate 119

Thionyl chloride (26 mL; 359 mmol) was added dropwise to a solution of 3-amino-5-methoxybenzoic acid (10g; 59.82 mmol) in MeOH (150 mL) at 0°C. The reaction mixture was stirred for 2 hours at room temperature. The precipitate was filtered off, washed with DIPE and dried under vacuo at 50°C to give 8.6 g (79%) of intermediate 119 (a white solid) .




b) Preparation of intermediate 120

A solution of lithium 2.4M in THF (35.8 mL; 85.9 mmol) was added dropwise to a solution of intermediate 119 (8.62 g; 39.6 mmol) in dry THF (150 mL) at 0°C under a N_2 flow. The reaction mixture was stirred for 2 hours at room temperature. The reaction mixture was treated with NH_4Cl and stirred for 10 minutes at 0°C. The precipitate was filtered off and washed with EtOAc.The filtrate was separated and the organic layer was washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (6 g) was purified by chromatography over silica gel (200g; mobile phase gradient from 100% DCM to 7%MeOH in DCM). The pure fractions were collected and concentrated to give 3.26 g of intermediate 120.




c) Preparation of intermediate 121

Under an inert atmosphere, a solution of tris(dibenzylacetone)palladium (0) (1.12 g; 1.2 mmol), 2-dicyclohexylphospino-2',4',6'-tri-isopropyl-1,1'-biphenyl (1.28 g; 2.7 mmol) was added to room temperature to a solution of intermediate 2 (3.52 g; 12.2 mmol), intermediate 120 (3.3 g; 17.45 mmol) and cesium carbonate (11.9 g; 36.49 mmol) in *t*-BuOH (80 mL). The reaction mixture was stirred at 105°C for 1 hour using one single mode microwave. The reaction mixture was cooled to room temperature, poured out into ice water (400 mL) and was stirred for 15 minutes. The precipitate was filtered off and washed with water. The precipitate was dissolved in DCM/MeOH (95/5) and the insoluble product was filtered off and dried to give 4.7 g of intermediate 121 used without further purification for the next step.



d) Preparation of intermediate 122


MnO_2 (5.65 g; 65 mmol) was added to a solution of intermediate 121 (4.7 g; 13 mmol) in THF (270 mL). The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was filtered over a pad of celite®. The filtrate was evaporated to give 1.5 g (32%) of intermediate 122 used without further purification for the next step.



e) Preparation of intermediate 123

Intermediate 122 (0.3 g; 0.64 mmol) and diethylamine (0.14 g; 1.9 mmol) were added to Pd/C 10% (0.05 g) and 0.2 mL of a 4% solution of thiophene in DIPE in MeOH/THF (100 mL) under N_2 flow. The reaction mixture was stirred at 50°C under 75 atm H_2 atmosphere until 1 eq hydrogen was absorbed. The reaction mixture was filtered over a pad of celite®. The filtrate was evaporated to give 0.354 g of the intermediate 123 .


Example A49



Preparation of intermediate 124


A mixture of 3-bromo-5-methoxyphenol (2 g; 9.8 mmol), cesium carbonate (6.4 g; 19.7 mmol) in *N,N*-dimethylformamide (20 mL) and water (4 mL) was degassed under N_2 flow for 1 hour, then acetic acid-2-chloro-2,2-difluoro-sodium salt (5.3 g; 34.5 mmol) was added. The reaction mixture was stirred at 120°C for 2 days. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (2.5 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g ; mobile phase gradient from 95%Heptane, 5%EtOAc to 90%Heptane, 10%EtOAc). The pure fractions were collected and concentrated to give 0.56 g (23%) of intermediate 124.

Example A50



a) Preparation of intermediate 133

A mixture of 2-chloro-4-methoxypyrimidine (1.24 g; 8.5 mmol), 4-piperidinemethanol (1.2 g; 10.25 mmol) and K_2CO_3 (2.4g; 17.0 mmol) in CH_3CN (15 mL) was stirred at 80°C for 18 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (1.8 g) was purified by chromatography over silica gel (SiOH 15-40µm, 40g; mobile phase 0.1%NH_4OH, 99%DCM, 1%MeOH). The pure fractions were collected and concentrated to give 1.6 g (83%) of intermediate 133 .




b) Preparation of intermediate 134

Methanesulfonyl chloride (0.94 mL; 12.1 mmol) was added dropwise to a solution of intermediate 133 (0.54 g; 2.42 mmol), triethylamine (2.4 mL; 16.9 mmol) in DCM (15 mL) at 5°C under a N_2 flow. The reaction mixture was stirred at 10°C for 1 hour. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (1.1 g) was purified by chromatography over silica gel (SiOH 15-40µm, 40g; mobile phase from 99%DCM, 1%MeOH). The pure fractions were collected and concentrated to give 0.5 g (69%) of intermediate 134.

This intermediate was used in the preparation of compound 839.


Example A51



a) Preparation of intermediate 137


4-Methyl-1-dimethylsulfamoylimidazole (2.9 g, 15.6 mmol) was diluted in THF (105 mL). The resulting solution was cooled down to -78°C and *N* butyl lithium 2M in cyclohexane (11.7 mL, 18.7 mmol) was added dropwise. The reaction mixture was stirred for 30

minutes at -78°C, *N N*-dimethylformamide (7.6 mL, 98.0 mmol) was added and the mixture was stirred for 1 hour at -78°C, then allowed to raise to room temperature in 1 hour.The reaction mixture was neutralized with an aqueous solution of NH_4Cl and then poured out into water and EtOAc. The organic layer was dried ($MgSO_4$), filtered and concentrated to afford 3.7 g of intermediate 137.



b) Preparation of intermediate 138


A mixture of intermediate 137 (3.7g; 17 mmol) was dissolved in MeOH (32 mL). Then the reaction mixture was cooled down to 0°C and sodium borohydride (0.6 g; 17 mmol) was added. The mixture was stirred for 1 hour at 0°C. The reaction mixture was then concentrated, poured out into water and EtOAc. The organic layer was dried ($MgSO_4$), filtered and concentrated to afford 2.9 g (78%) of intermediate 138. It was directly used in the next step without any further purification.



c) Preparation of intermediate 139

Intermediate 138 (3.2 g; 14.3 mmol) was dissolved in THF (26 mL) and *N N*-dimethylformamide (13 mL). Then the solution was cooled down to 0°C and triethylamine (4.1 mL; 28.6 mmol) followed by methanesulfonyl chloride (1.3 mL; 17.2 mmol) and lithium chloride (1.8 g; 43 mmol) were successively added. The mixture was stirred at room temperature over 2 hours. The reaction mixture was poured out into EtOAc and water. The organic layer was washed once with brine, dried ($MgSO_4$), filtered and concentrated. The residue (3.5 g) was purified by chromatography over silica gel (mobile phase gradient from 100% DCM to 0.1% NH_4OH, 99% DCM, 1% MeOH). The pure fractions were collected, the solvent was evaporated to afford 2.2 g (70%) of intermediate 139 used to prepare compound 695.


Example A52



Preparation of

A mixture of 3,5-dimethoxyboronic acid (18.5 g; 101.5 mmol), 1-(2,2,2-trifluoroethyl)-4-piperidinemethanamine (16.6g; 61.7 mmol), copper (II) acetate (18.5 g; 101.5 mmol) and triethylamine (59.8 mL;425 mmol) in DCM (350 mL) was stirred at room temperature for 18 hours. The mixture was filtered and the filtrate was evaporated till dryness. The residue was purified by chromatography over silica gel (mobile phase gradient from 89% petroleum ether/11% ethyl acetate to 45% petroleum ether/55% ethyl acetate). The pure fractions were collected and the solvent was evaporated to give 3.8 g (19%) of compound.

Example A53




Preparation of intermediate 142

A mixture of intermediate 15 (1.8 g; 3.6 mmol) and glycine tert butyl ester (2.5 g; 18 mmol) in *N,N*-dimethylformamide (25 mL) was stirred at 80°C for 6 hours in a sealed tube. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (2.52 g) was purified by chromatography over silica gel (SiOH 20-45µm, 450 g; mobile phase 0.1% NH_4OH, 96% DCM, 4% MeOH). The pure fractions were collected and concentrated to afford 0.96 g (50%) of intermediate 142_used whitout further purification for the next step.

B. Preparation of the compounds


Example B1



Preparation of compound 1

Tetrabutylammonium fluoride (38.5mL, 38.5mmol;) was added dropwise to a solution of intermediate 9 (20g, 38.5mmol) in THF (350mL) at room temperature. The reaction mixture was stirred at room temperature for 5 hours. The mixture was poured out into ice water and EtOAc was added. The mixture was basified with K_2CO_3 10% and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was triturated from diethyl ether, filtered and dried under vacuum, yielding 11.7g (75%) of compound 1. MP = 153°C (DSC).

Compound 1 was alternatively prepared using the following procedure.
525g (1.01mol) of intermediate 9 was dissolved in a mixture of THF (0.89L), acetic acid (2.68L) and water (0.89L) and the reaction mixture was stirred at 50°C upon complete conversion to the alcohol. The reaction mixture was evaporated till dryness. The residue was taken up in DCM (3.68L) and water (3.68L) and the pH of the mixture was adjusted to 7 using ammonia. The layers were separated. The aqueous layer was extracted with DCM (0.5L) and the organic layers combined, dried ($MgSO_4$), filtered and evaporated till dryness. The residue was crystallized from toluene. The precipitate was filtered off, washed with toluene and dried to provide 204g (49.8% yield) of compound 1.




a) Preparation of compound 2

A mixture of intermediate 47 (1.50g; 2.476mmol), HCl 3N (2mL) in dioxane (25mL) was heated at 70°C overnight. The reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the mixture was basified with an aqueous solution of K_2CO_3 (10%). The organic layer was separated, washed with brine,

dried ($MgSO_4$), filtered and the solvent was evaporated. The compound was triturated from diethyl ether, filtered and dried under vacuum, yielding 0.790g (81%) of compound 2. MP=169°C (DSC).

Example B2



a) Preparation of compound 3


Tetrabutylammonium fluoride (14.6 mL, 14.6 mmol) was added dropwise to a solution of intermediate 11 (6.5g, 12.2mmol) in THF (100mL) at room temperature. The reaction mixture was stirred at room temperature overnight. The mixture was poured out into ice water and EtOAc was added. The mixture was basified with an aqueous solution of K_2CO_3 (10%) and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness to give 7.8g of crude compound, which was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 450g MATREX; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and evaporated to yield 4.9g (96%) of compound 3. The compound was crystallized from Et_2O/CH_3CN, the precipitate was filtered and dried to give 4.37g (85%) of compound 3. MP = 168°C (Kofler).

Compound 3 was alternatively also prepared using the following procedure. Intermediate 11 (167.2g; 313mmol) was added to a mixture of acetic acid (846mL), THF (282mL) and water (282mL) and the mixture was stirred at 50°C for 18 hours and evaporated till dryness. The crude compound 3 was used without further purification to prepare intermediate 17a.

Compound 3 was alternatively also prepared using the following procedure B2b. b)HCl/i-PrOH (11.3mL; 56.5mmol) was added dropwise to a solution of intermediate 18 (8.5g; 16.87mmol) in CH_3OH (100mL) at 10°C, and the mixture was stirred for 1 hour at room temperature. Ice water was added to the solution which was basified with NH_4OH. The product was extracted with DCM. The organic layer was dried ($MgSO_4$) and evaporated to dryness. The residue was purified by chromatography over silica gel

(Irregular SiOH, 15-40µm, 200g; mobile phase, 97%DCM, 3%CH_3OH, 0.1%NH_4OH). The pure fractions were collected and evaporated to dryness to afford 3.7g (52%) of compound 3 and 1.2g of an impure fraction This impure fraction was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase 0.5% NH_4OH, 97% DCM, 3% CH_3OH). The pure fractions were collected and the solvent was evaporated, yielding 700 mg (10%) of compound 3.

Example B3



Preparation of compound 4

A mixture of intermediate 10 (8.7g; 17.99mmol) and isopropylamine (61.3mL, 719.68mmol) was heated at 90°C for 3 hours in a sealed vessel. The reaction mixture was cooled to room temperature and the mixture was evaporated till dryness. DCM and water were added and the organic layer was separated, washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (8g) was crystallized from Et_2O/CH_3CN, filtered and dried under vacuum at 60°C, yielding 6.68g (83%) of compound 4.MP = 142°C (DSC).

Compound 4 was alternatively prepared using the following procedure.
A mixture of intermediate 10 (322g; 666mmol) and 2-propanamine (196.8g; 3.3mol) in acetonitrile (2.66L) was heated at 100°C in a sealed vessel for 18 hours. The reaction mixture was cooled to room temperature and concentrated to ~30% of its initial volume. Water (1.5L), 2-methyltetrahydrofurane (2.5L) and $NaHCO_3$ (50g) were added. The layers were separated, the organic layer was washed with a solution made of 50g of $NaHCO_3$ in water (1L), dried ($MgSO_4$), filtered over silica gel and evaporated till dryness. The residue was crystallized from 2-propanol. The precipitate was filtered off, dried in vacuum to provide 257.2g (86.5%) of compound 4.


Compound 4 was alternatively prepared using the following procedure.

Intermediate 3 (20.0g; 55.3mmol), then tetra-N-butylammonium bromide (9.06g; 27.7mmol) were added at 2°C under inert atmosphere to a solution of potassium hydroxide (46.6g; 830mmol) in THF (387mL) and water (6mL). The reaction was stirred at room temperature for 2 hours before portionwise addition of N-(2-chloroethyl)-2-propanamine HCl (CAS[6306-61-2]), and then at 50°C upon complete conversion. Water was added, layers were separated and the organic layer concentrated, taken up in DCM/water, neutralized with HCl to neutral pH. Organic layer was washed with water, dried ($MgSO_4$), filtered and evaporated till dryness to give 26.6g of compound 4.

Compound 4 as a HCl salt (.1HCl) was prepared using the following procedure.
To a stirred mixture of 2-methyltetrahydrofuran (1.5L) and KOH (140g, 250mmol) was added water (30mL). Then intermediate 3 (60g, 166mmol)) and tetrabutylammoniumbromide (13.4g, 41mmol) were added and the mixture was heated at 50°C for 1hour while stirring. Then N-(2-chloroethyl)-2-propanamine HCl (CAS[6306-61-2]) (48g, 299mmol) was added in 1 portion. The mixture was stirred for 18 hours at 50°C. When the conversion was complete, water (600mL) was added to the reaction mixture. The layers were separated and the organic layer was concentrated. The residue was dissolved in 2-propanol (120mL) and HCl in 2-propanol was added at 60°C. After cooling, the HCl-salt was isolated via filtration. After drying at 50°C in a vacuum drying oven the HCl-salt was obtained in 83% yield (compound 4a).

To 51.69g (107mmol) of the HCl salt from the previous step was added water (258mL) and DCM (258mL). The pH of the reaction mixture was adjusted using ammonium hydroxide (17.25mL) until pH = 9.5. The layers were separated and the organic layer was concentrated. The residue was crystallized from 2-propanol (258mL). After drying at 50°C under vacuum compound 4 was obtained in 91% yield (43.4g).


Example B3A



Preparation of compound 6

Intermediate 48 (7.2g; 12.7mmol), 3,3-difluoropyrrolidine hydrochloride (7.3g; 50.7mmol), sodium carbonate (6.72g; 63.42mmol), potassium iodide (2.1g; 12.7mmol) in 1-butanol (220mL) were heated to 90°C for 15 hours. The mixture was cooled to room temperature and poured into H_2O/K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 35-40µm, Grace Resolv; mobile phase gradient from 100% DCM to 95% DCM, 5% MeOH, 0.1% NH_4OH). The desired product fraction was collected and the solvent was evaporated, yielding 3.2g (44%) of compound 6.

Example B3B





Preparation of compound 580

A mixture of intermediate 10 (2.8 g; 5.8 mmol) and 1,4-dioxa-8-azaspiro[4-5]decane (1.5 g; 18 mmol) in 1-methyl-2-pyrrolidinone (10 mL) in a sealed tube was heated at 140°C using one single mode microwave (Biotage Initiator EXP 60) for 1 hour. The reaction mixture was evaporated till dryness. The crude product (6 g) was purified by chromatography over silica gel (15-40 µm 300 g; mobile phase, 0.2% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and the solvent was evaporated till dryness to give 1.9 g (61%) of compound 580.

Example B3C


Preparation of compound 666 and 665





A mixture of intermediate 10 (0.3 g; 0.6 mmol), 2-piperidine-2-carboxamide (0.32 g; 2.5 mmol), potassium iodide (0.1 g; 0.6 mmol) and carbonate sodium (0.41 g; 4.4 mmol) in 1-butanol (12 mL) was stirred at 85°C for 4 days. The reaction was poured into ice water and EtOAc was added. The organic layer was washed with brine, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (0.33 g) was purified by preparative LC on (irregular, SiOH 15-40 µM, 30 g, mobile phase gradient from 0.1%NH_4OH, 98%DCM, 2%MeOH to 0.1%NH_4OH, 96%DCM, 4%MeOH). The pure fractions were collected and the solvent was evaporated. The first product (0.1 g) was crystallized from diethyl ether. The precipitate was filtered and dried to give 0.081 g (25%) of compound 665. MP: 206°C (Kofler). The second product (0.1 g) was crystallized from diethyl ether. The precipitate was filtered and dried to give 0.082 g (25%) of compound 666. MP: 163°C (Kofler).

Example B3D





Preparation of compound 677

Intermediate 10 (1.3 g; 2.7 mmol), methoxylamine hydrochloride (2.3 g; 26.9 mmol) in triethylamine (15 mL; 107.5 mmol) were heated at 90°C for 5 hours in a sealed tube. The reaction was poured out into ice water. The organic layer was separated and washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (2 g) was purified by chromatography over silica gel (SiOH, 15-40µm, 300 g; mobile phase, 96% DCM, 4% *i*-PrOH). The pure fractions were collected and concentrated. The

residue (0.38 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.32 g (27%) of compound 677 MP: 177°C (DSC).

Example B3E



Preparation of compound 923 (free base) and compound 886 (HCl salt)


and

as a HCl salt

A mixture of intermediate 10 (1.0g; 2.07 mmol) and 3-pyrroline (628µL, 8.3 mmol) in aetonitrile (4 mL) was heated at 90°C for 90 minutes in a microwave biotage device. The reaction mixture was cooled to room temperature and the mixture was evaporated until dryness. DCM and water were added and the organic layer was separated, washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated.The residue was chromatograohied over silica gel (5µm, mobile phase: gradient from NH_4OH 0.2%, DCM 98%, MeOH 2% to NH_4OH 0.8%, DCM 92%, MeOH 8%). The eluted fractions were evaporated and the residue was dissolved in DCM, and stirred at room temperature under air bubbling.for 24 hours. The solvent was evaporated to give a yellow foam which was chromatographied over silica gel (SiOH 10µm 60g, mobile phase 0.1% NH_4OH, 98% DCM, 2% MeOH). The desired product fractions were evaporated to provide 100mg (11%) of compound 923. This compound was converted into the HCl salt in MeOH. The precipitate was filtered off, washed with MeOH and dried to give 41 mg (4%) of compound 886.


Example B3F

Preparation of compound 891 and 894



.HCl and

as a HCl salt

compound 891 compound 894

and preparation of compound 924 and 925




and



compound 924 compound 925

A mixture of intermediate 143 (622 mg, 1.2 mmol) in isopropylamine (8.06 mL, 94.6 mmol) was heated at 120°C in a sealed vessel during 48 hours..The reaction mixture was cooled to room temperature and DCM was added. The organic layer was washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated to give a yellow oil. This residue was purified by chromatography over silica gel (5µm, mobile phase: gradient from 100% DCM to 0.7% NH_4OH, 93% DCM, 7% MeOH). The desired product fractions were collected and evaporated yielding 33 mg (6%) of compound 924 and 40 mg (7%) of compound 925. Compound 924 was converted into HCl salt in MeOH. The precipitate was filtered off, washed with Et_2O and dried to give 25mg (4%) of compound 891. Compound 925 was converted into HCl salt in MeOH. The solvent was evaporated, the residue was triturated into Et_2O, filtered off, washed with Et_2O and dried to give 51mg (7%) of residue. This fraction was taken up from MeOH, and stirred 10 minutes at room temperature. The solvent was evaporated to dryness. The product was triturated, and dried, to give 24mg (3%) of compound 894.

Example B4



a) Preparation of compound 5

A solution of intermediate 17a (0.2g; 0.402mmol) in 2,2,2-trifluoroethylamine (2mL; 25mmol) was heated at 90°C in a sealed tube for 12 hours. The reaction mixture was

cooled to room temperature and poured out into ice water. EtOAc was added and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, 98% DCM, 2% CH_3OH). The pure fractions were collected and the solvent was evaporated. The residue (0.14g, 69%) was crystallized from DIPE/diethyl ether/pentane (1/1/1). The precipitate was filtered and dried under vacuum, yielding 0.134 g (67%) of compound 5, MP=126°C (DSC).


Compound 5 was alternatively also prepared using the following procedure B4b.

b) 3M HCl (60mL) was added to a solution of intermediate 17 (9.49mmol; 5.7g) in CH_3OH (120mL) at room temperature. The reaction mixture was heated at 60°C overnight. The reaction mixture was cooled to room temperature, diluted with DCM and poured onto an iced solution of K_2CO_3 (10%). The mixture was stirred for 30 minutes and the organic layer was decanted, washed with water, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by HPLC. The residue (5.3g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase 0.1% NH_4OH, 98% DCM, 2% CH_3OH). The pure fractions were collected and evaporated to dryness. The oily residue (3.93g, 83%) was crystallized from DiPE/diethyl ether/CH_3CN. The precipitate was filtered off and dried, yielding 3.7g (78%) of compound 5.


Compound 5 was alternatively also prepared using the following procedure.

A mixture of intermediate 17 (268.5g; 447mol) and trifluoroacetic acid (0.5L) in DCM (2.24L) was stirred at room temperature for 18 hours and then at 50°C for 1 hour. The reaction mixture was evaporated till dryness, taken up in toluene (0.3L) and evaporated again. The residue was dissolved in DCM (3L) and water (2L) and the pH adjusted to neutral with ammonia. The layers were separated, the aqueous layer was extracted with DCM (0.3L) and the organic layers combined and evaporated till dryness. The residue was dissolved in EtOAc (1.5L) and stirred for 1 hour with a mixture of silica gel (275g).The silica gel was filtered off, washed with EtOAc and the filtrate evaporated till dryness to give 226g of compound 5. It was crystallized from 2-propanol, filtered and dried to provide 180.8g (80%) of compound 5.

Example B4A




Preparation of compound 7



and compound 8

A methylamine solution in absolute ethyl alcohol (5.15 mL, 33% w/w, 41.4mmol) was added dropwise to a suspension of intermediate 10 (2g, 4.1mmol), K_2CO_3 (2.86g, 20.7mmol) in dry CH_3CN (40mL) at room temperature. The mixture was heated at 80°C overnight in a sealed vessel. The reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (1.85g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase gradient from 0.1% NH_4OH, 95% DCM, 5% MeOH to 0.1% NH_4OH, 90% DCM, 10% MeOH). The desired fractions were collected and the solvents were evaporated, yielding 0.30g of Fraction I (15%) and 1.25g of Fraction II (72%). Fraction I was crystallized from diethyl ether, filtered and dried under vacuum, yielding 0.240g (12%) of compound 7. MP=160-162°C. Fraction II was taken up with DCM and an aqueous solution of K_2CO_3 (10%). The mixture was stirred for 1 hour, then the organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The product was crystallized from diethyl ether/CH_3CN, filtered and dried under vacuum at 60°C, yielding 1.05g (59%) of compound 8. MP=180-182°C (Kofler).


Example B4B



Preparation of compound 679


A mixture of intermediate 3-{(3-Fluoro-5-methoxyphenyl)[3-(1-methyl-1H-pyrazol-4-yl)quinoxalin-6-yl]amino}propyl methanesulfonate prepared according to A3 0.35 g; 0.72 mmol), (S)-(+)-2-pyrrolidine methanol (0.1 mL; 1 mmol) and triethylamine (0.4 mL; 2.9 mmol) in 1-methyl-2-pyrrolidinone (1 mL) was heated at 140°C for days in a sealed tube. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.12 g) was purified by chromatography over silica gel (SiOH 5µm; mobile phase gradient from 0.52% NH_4OH, 98% DCM, 2% MeOH to 0.8% NH_4OH, 92% DCM, 8% MeOH). The pure fractions were collected and concentrated to give 0.031 g (9 %) of compound 679

Example B4C




Preparation of compound 694 as a HCl salt


NaH (0.24 g; 5.9 mmol) was added portionwise to 2-pyrrolidinone (0.46 mL; 5.9 mmol) in *N,N*-dimethylformamide (30 mL) at 5°C under N_2 flow. The reaction mixture was stirred

at 5°C for 1 hour, then intermediate  prepared according to A5 (1 g; 2 mmol) was added at 5°C under N_2 flow. The reaction mixture was allowed to warm to room temperature and stirred for 18 hours. The reaction was poured out into ice water. A precipitate was filtered, washed with water. The organic layer was separated and washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.8 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300 g ; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated. The residue was dissolved in isopropyl alcohol and stirred at 0°C, then 0.5 mL of HCl *i*-PrOH 5N was added dropwise. Diethyl ether was added and the solution was stirred at 0°C for 1 hour and the precipitate was filtered and dried to afford 0.33 g (26%) of compound 694. MP: 197°C (DSC).

Example B5


Preparation of compound 9 

NaH (0.556g; 13.9mmol) was added portionwise to a solution of intermediate 49 (3g; 6.95mmol) in DMF (85mL) at 5°C under N_2.The reaction mixture was stirred for 30 minutes. 1-bromo-3-chloropropane (2mL; 20.9mmol) was added dropwise and the mixture was stirred for 15 hours at room temperature, then poured into H_2O/K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The obtained residue was purified by Chromatography over silica gel (Irregular SiOH 15-40µm, 90g MERCK; mobile phase gradient from 100% DCM to 97% DCM, 3% MeOH, 0.1 % NH4OH). The desired product fractions were collected and the solvent was evaporated, yielding 2.94g (86%) of compound 9.




a) Preparation of compound 10

NaH (925mg, 23.1mmol) was added portionwise to a solution of intermediate 3 (4.18g, 11.6mmol) in DMF (52mL) at 5°C. The mixture was stirred at 5°C for 30 minutes, then a solution of 4-(phenylmethyl)-2-morpholinemethanol 2-methanesulfonate (4.95g, 17.3mmol) in DMF (13.5mL) was added. The reaction mixture was heated at 60°C for 18 hours. The mixture was poured into water and the product was extracted with EtOAc. The organic layer was washed with water and brine, dried ($MgSO_4$), filtered and evaporated. The obtained residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The desired product fraction was collected and the solvent was evaporated, yielding 2.74g (43%, purity 90%) of a yellow foam. A sample (440mg) was purified by achiral super critical fluid chromatography (AMINO 6µm 150x21.2mm; mobile phase, 0.3% 2-propylamine, 20% MeOH, 80% CO_2). The desired product fraction was collected and the solvent was evaporated, yielding 356mg of a residue which was crystallized with DCM/Acetone/diethyl ether. The precipitate was filtered off and dried to give 188mg of compound 10. MP=134°C (Kofler).




b-1) Preparation of compound 11

To a solution of intermediate 3 (67mg, 0.18mmol) in tetrahydrofuran (4mL) was added NaH (12mg, 0.28mmol). The suspension was stirred at room temperature until no bubbles were observed, cooled to 0°C and methyl iodide (0.08mL, 1.3mmol) was added drop wise. The reaction mixture was stirred at room temperature overnight, diluted with EtOAc and washed with brine. The organic layer was separated, dried (Na_2SO_4) and concentrated. The crude residue was purified by chromatography over silica gel to afford 38mg (54%) of compound 11 (yellow powder).



b-2) Preparation of compound 12


To a solution of intermediate 3 (100mg, 0.277mmol) in tetrahydrofuran (3mL) was added potassium hexamethyldisilazide (0.5M in toluene, 12mg, 0.831mmol). The reaction mixture was stirred at room temperature for 30 minutes and propyl bromide (0.30mL) was added drop wise. The reaction mixture was stirred at room temperature for a further 3 hours and diluted with DCM and water. The solid residue was removed by filtration, dissolved in MeOH and combined with the other organic extracts, dried ($MgSO_4$) and then concentrated. The crude residue was purified by chromatography over silica gel to afford 10mg (9%) of compound 12 (yellow powder).




b-3) Preparation of compound 13

A mixture of intermediate 3 (50mg, 0.139mmol), cesium carbonate (226mg, 0.693mmol) and 1-bromo-2-methyl propane (95mg, 0.693mmol) in CH_3CN (1 mL) was heated. The reaction mixture was heated in a CEM Discovery microwave at 100 °C for 1 hour. Upon cooling, the reaction mixture was partitioned between DCM and water. The organic layer was separated and the aqueous layer was extracted with further DCM. The combined organic layers were dried ($MgSO_4$) and concentrated. The crude residue was purified by chromatography over silica gel to afford 5 mg (9%) of compound 13 (yellow powder).


Example B6

Preparation of compound 14  and

compound 14a  as a HCl salt

NaH (513.5mg, 12.8mmol) was added portionwise to a solution of intermediate 8 (2.5g, 6.4mmol) in DMF (25mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then glycidyl methyl ether (1.1mL, 12.8mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 1 hour at 5°C, then allowed to warm up to room temperature. The reaction was stirred at 80°C for 5 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase 0.1% NH_4OH, 97.5% DCM, 2.5% MeOH). The desired product fractions were collected and the solvent was evaporated yielding 0.66g (21.5%) of compound 14 which was converted into its HCl salt with HCl/2-propanol (5-6N) in MeOH. The mixture was evaporated, and the resulting solid was triturated into diethyl ether, filtered and dried to give 0.488g (15%) of compound 14a (0.95eq HCl) (mp = 110°C, kofler).


Example B7

Preparation of compound 15 

HCl 3N (13.5mL) was added to a solution of intermediate 50 (2g, 2.98mmol) in CH_3OH (65mL) at 5°C. The reaction mixture was stirred at room temperature for 2.5 hours then

heated at 60°C overnight. The solution was poured into ice water and basified with an aqueous solution of K_2CO_3 (10%). The product was extracted with DCM. The organic layer was washed with water, dried ($MgSO_4$), filtered and evaporated till dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 450g MATREX; mobile phase 0.1% NH_4OH, 96% DCM, 4% MeOH). The desired fractions were collected and the solvent was evaporated. The residue was purified by achiral super critical fluid chromatography on (DIETHYLAMINOPROPYL 5µm 150x21.2mm; mobile phase 0.3% 2-propylamine, 80% CO_2, 20% MeOH). The desired fractions were collected and the solvent was evaporated. The residue was crystallized from CH_3CN/DIPE, the precipitate was filtered off and dried, yielding 760mg (53%) of compound 15. MP= 121°C (DSC).


Example B8



Preparation of compound 16

HCl/i-PrOH (0.33mL, 0.0017mol) was added dropwise to a solution of intermediate 51 (0.25g, 0.0004mol) in CH_3OH (6mL) at 10°C. Then the mixture was stirred for 3 hours. The solution was concentrated, taken up with iced water, basified with NH_4OH and the product was extracted with DCM. The organic layer was dried ($MgSO_4$) and evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm 30g; mobile phase 1% NH_4OH, 92% DCM, 8% MeOH). The desired product fraction was collected and the solvent was evaporated, yielding 138mg (78%) of compound 16, MP=80°C (Kofler).

Example B9



a) Preparation of compound 17

HCl 3N (4mL) was added dropwise to a solution of intermediate 20 (1.5g, 3.0mmol) in dioxane (20mL) at room temperature. The reaction mixture was heated at 70°C overnight. The reaction was cooled to room temperature and poured out into ice water. EtOAc was added and the mixture was basified with an aqueous solution of K_2CO_3 (10%). The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The compound was crystallized from diethyl ether, filtered and dried under vacuum at 60°C, yielding 1g (83%) of compound 17. MP=158-160°C (Kofler).

Compound 17 was alternatively also prepared using the following procedure B9b.


b) Under N_2, intermediate 19 (3.0g; 8.1mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (1.9g; 9.7mmol), sodium carbonate 2M (6.1mL; 12.2mmol) in ethylene glycol dimethyl ether (30mL) were degassed by bubbling nitrogen through for 10 minutes. $Pd(PPh_3)_4$ (0.75g; 0.65mmol) was added and the mixture was heated at reflux for 15 hours. The residue was poured into ice water and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 90g; mobile phase gradient from 100%DCM to 95%DCM, 5%MeOH, 0.1%NH_4OH)15-40µm, 90g). The pure fractions were collected and evaporated to dryness. The obtained residue was crystallized in DIPE, filtered and dried, yielding 1.66g (51%) of compound 17.

Compound 17 was alternatively also prepared using the following procedure B9c.

c) A mixture of intermediate 19 (3.3g, 8.9mmol), 1,1-dimethylethyl ester 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole-1-carboxylic acid (3.15g, 10.7mmol), potassium phosphate (3.79g, 17.8mmol), dicyclohexyl(2',6'-dimethoxy[1,1'-biphenyl]-2-yl)phosphine (0.37g, 0.9mmol) in dioxane (60mL) and H_2O (6mL) was stirred at room temperature under N_2 flow. After 10 minutes, $Pd_2(dba)_3$ (0.408g, 0.446mmol) was added portionwise at room temperature and the mixture was heated at 80°C overnight. The

reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the mixture was filtered through a layer of celite. The celite was washed with EtOAc, then the filtrate was extracted with EtOAc, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15/40µm 30g MERCK; mobile phase, gradient 100% DCM to 97% DCM, 3% MeOH). The desired product fraction was collected and the solvent was evaporated, yielding 3.30g (73%) of compound 17.


Example B10



Preparation of compound 18

At 5°C, HCl/i-PrOH 5/6N (213 µl; 1.06mmol) was added to a solution of intermediate 23 (250mg; 0.53mmol) in CH_3OH (5 mL). The reaction mixture was stirred at 5°C for 3 hours. H_2O and ice were added. An aqueous solution of K_2CO_3 (10%) was added until pH became basic and the product was extracted with DCM. The organic layer was washed with H_2O, brine dried ($MgSO_4$), filtered and the solvent was evaporated. The crude product was taken up in diethyl ether, filtered and dried under vacuum, yielding: 64 mg (31%) of compound18. MP=132°C (Kofler).

Example B11




Preparation of compound 19

A mixture of intermediate 52 (0.99g, 1.8mmol) in HCl 3N (3mL) and dioxane (17mL) was heated at 70°C overnight. The reaction mixture was cooled to room temperature and poured out into ice water and EtOAc was added. The solution was basified with an aqueous solution of K_2CO_3 (10%) and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm 30g MERCK; mobile phase,


gradient 100% DCM to 98% DCM, 2% MeOH). The pure fractions were collected and evaporated to dryness, yielding 782mg (97%) of compound 19. MP=130°C (Kolfer).

Example B12



Preparation of compound 20

*N*3-(ethylcarbonimidoyl)-*N*1,*N*1-dimethyl-1,3-propanediamine hydrochloride (1:1) (0.12g, 0.76mmol) was added portionwise to a solution of intermediate 24 (0.23g, 0.505mmol), 3-pyrrolidinol (0.061g, 0.76mmol), 1-hydroxybenzotriazole (0.1g, 0.76mmol), Et_3N (0.105mL, 0.76mmol) in DCM (10mL) at room temperature. The reaction mixture was stirred for 15 hours. The mixture was poured into H_2O and extracted with DCM. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was crystallized in DIPE, filtered and dried. The product fraction was purified by chromatography over silica gel (Spherical SiOH, 10µm 60g, PharmPrep MERCK; mobile phase 0.5% NH_4OH, 94% DCM, 6% MeOH). The pure fractions were collected and evaporated to dryness. The residue was crystallized with DIPE, filtered and dried, yielding 186 mg (70%) of compound 20. MP= 203.4°C (DSC).




a) Preparation of compound 21

*N*3-(ethylcarbonimidoyl)-*N*1,*N*1-dimethyl-1,3-propanediamine hydrochloride (1:1) (227mg; 1.46mmol) was added to a mixture of intermediate 53 (550mg; 0.98mmol), methylamine hydrochloride (329mg; 4.88mmol), Et_3N (0.95mL; 6.83mmol), 1-hydroxybenzotriazole (198mg; 1.46mmol) in DCM (40mL) at room temperature. The reaction mixture was stirred for 20 hours, then stirred for 2 days, poured into H_2O and extracted with DCM. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g Merck; mobile phase, gradient from 100% DCM to 90% DCM, 10% MeOH,

0.1% NH_4OH). The desired product fraction was collected and the solvent was evaporated, yielding 76 mg (17%) of compound 21 which was crystallized in diethyl ether to give 59 mg (13%) of compound 21. MP=204.5°C (DSC).

Compound 21 can also be prepared from the corresponding $-O-Si(CH_3)_2-C(CH_3)_3$ intermediate according to procedures described above, such as for example in Example B2.


Example B13



Preparation of compound 22

A mixture of intermediate 25 (0.4g, 1.43mmol), 1-bromo-3-isopropoxybenzene (0.46mL, 2.86mmol), sodium tert-butoxide (0.032g, 0.14mmol) and 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] (0.413g, 4.30mmol) in ethylene glycol dimethyl ether (3mL) was degassed with N_2 for 10 minutes. Palladium(II) acetate (47% Pd) (0.032g, 0.14mmol) was added and the mixture was heated at 135°C under microwave irradiation for 60 minutes. The mixture was cooled to room temperature, poured into H_2O/K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm 30g; mobile phase, 0.1% NH_4OH, 99% DCM, 1% MeOH). The pure fractions were collected and the solvent was evaporated. The residue was purified by chromatography over silica gel (X-Bridge-C18 5µm 30*150mm; mobile phase, gradient from 40% of a 0.5% solution of NH_4HCO_3 in water, 60% CH_3CN to 100% CH_3CN). The pure fractions were collected and the solvent was evaporated. The residue (0.187 g) was crystallized from DIPE/pentane (80/20), then the precipitate was filtered and dried under vacuum, yielding 0.128 g (22%) of compound 22. MP=109°C (DSC).


Example B14



Preparation of compound 23

A solution of intermediate 54 (0.4g, 0.666mmol) and tetrabutylammonium fluoride (0.73mL, 0.73mmol) in THF (10 mL) was stirred at 0°C for 2 hours. Water and EtOAc were added, the organic layer was separated, washed with water, then brine, dried ($MgSO_4$), filtered and the solvent was evaporated till dryness. The residue (0.4 g) was first purified by chromatography over silica gel (Irregular SiOH 15-40µm 300g MERCK; mobile phase, gradient from 98% DCM, 2% MeOH to 95% DCM, 5% MeOH). The pure fractions were collected and the solvent was evaporated. The residue was then purified by achiral super critical fluid chromatography on (AMINO 6µm 150x21.2mm; mobile phase, 0.3% 2-propylamine, 80% CO_2, 20% EtOH). The pure fractions were collected and the solvent was evaporated. The residue (0.165 g, 51%) was crystallized from DIPE, the precipitate was filtered and dried under vacuum, yielding 0.150 g (46%) of compound 23. MP=134°C (Kofler).

Example B14A




Preparation of compound 691

A 1M solution of tetrabutylammonium fluoride in THF (12.7 mL; 12.7 mmol) was added dropwise to a solution of intermediate 691 (5g; 8.5 mmol) in THF (50 mL) at room temperature. The reaction mixture was stirred at room temperature for 3 hours. The mixture was poured out into ice water and EtOAc was added. The mixture was basified with K_2CO_3 10% and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (3.5 g) was

crystallized from diethyl ether. The precipitate was filtered off, dried in vacuum to provide 3.2 g (80%) of compound 691 MP: 99°C (DSC).


Example B15



Preparation of compound 24


HCl/i-PrOH (276µL, 1.38mmol) was added dropwise to a solution of intermediate 55 (183mg, 0.35mmol) in CH_3OH (2mL) at 10°C and then the mixture was stirred for 3 hours. Diethyl ether was added and the precipitate was filtered and dried, yielding 126mg (76%) of compound 24. MP=80°C.

Example B16

Preparation of compound 25  as a HCl salt

A mixture of intermediate 16 (1.37g, 2.5mmol) in pyrrolidine (30mL) was heated at 80°C for 3 hours. The mixture was cooled to room temperature and was evaporated until dryness. The residue was taken up in DCM and H_2O. The organic layer was extracted with DCM, dried ($MgSO_4$), filtered and evaporated to dryness. The residue (3 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 90g MERCK; mobile phase, gradient from 98% DCM, 2% MeOH to 95% DCM, 5%MeOH). The pure fractions were collected and the solvent was evaporated, yielding the free base. The residue was dissolved in i-PrOH, then 1.04 mL of HCl 5N/I-PrOH (4 eq.) was added dropwise at 5°C. The salt was filtered, washed with DIPE and dried under vacuum at 60°C, yielding 0.53g (40%) of compound 25. MP = 259°C (DSC).


Example B17



Preparation of compound 26

HCl 3N (2mL) was added dropwise to a solution of intermediate 56 (0.3g, 0.484mmol) in dioxane (8mL). The solution was heated at 70°C for 3 hours. The reaction was cooled to room temperature and poured out into ice water. EtOAc was added and the mixture was basified with an aqueous solution of K_2CO_3 (10%). The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.19g) was crystallized from DIPE/CH_3CN. The precipitate was filtered and dried under vacuum, yielding 0.112g (56%) of compound 26. MP = 202°C (DSC).


Example B18



Preparation of compound 27

Intermediate 57 (0.425g; 0.88mmol), HCl 3N (3mL) and dioxane (8mL) were heated to 60°C overnight. The mixture was cooled to room temperature, poured into H_2O and basified with K_2CO_3. The product was extracted with EtOAc, dried ($MgSO_4$), filtered and evaporated to dryness, yielding 322mg (95%) of compound 27. MP=178°C (DSC).


Example B19



Preparation of compound 28

A mixture of intermediate 58 (0.3g, 0.486mmol) and amberlyst 15 ion exchange resin (0.03g) in MeOH (8 mL) was stirred at 45°C for 3 hours. The resin was filtered. The filtrate was poured out into water and was basified with an aqueous solution of K_2CO_3 (10%). EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.2g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g MERCK; mobile phase, gradient from 98% DCM, 2% MeOH to 95% DCM, 5% MeOH). The pure fractions were collected and the solvent was evaporated. The residue (0.18g) was crystallized from DIPE/CH_3CN (80/20). The precipitate was filtered off and dried under vacuum, yielding 0.114 g (52%) of compound 28. MP=142°C (DSC).

Example B20




Preparation of compound 29

A 2M solution of methylamine in THF(4.8mL, 0.0097mol) was added to a solution of intermediate 26 (0.14g, 0.0003mol) and K_2CO_3 (0.1g, 0.0007mol) in THF (5mL). The solution was heated to 100°C in a sealed tube for 24 hours, then cooled to room temperature and poured into H_2O/NaCl. The mixture was extracted with DCM. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography (Irregular SiOH,15-40µm, 30g; mobile phase, 95%DCM,-

5%MeOH, 0.1%NH_4OH). The product fractions were collected and the solvent was evaporated, yielding 103mg (86%) of compound 29. MP= 80°C (Kofler).


Example B21



Preparation of compound 30

HCl/i-PrOH (0.4mL, 0.002mol) was added dropwise to a solution of intermediate 59 (0.31g, 0.0005mol) in CH_3OH (5mL) at 10°C and the mixture was stirred for 2 hours. The solution was evaporated to dryness, then the residue was taken up with ice water, basified with NH_4OH and the product was extracted with DCM. The organic layer was dried ($MgSO_4$) and evaporated to dryness. The residue was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm).;mobile phase, gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 1.1% NH_4OH, 89% DCM, 11% MeOH). The product fractions were collected and the solvent was evaporated, yielding 106mg (47%) of compound 30. MP=80°C (Kofler).

Example B22




Preparation of compound 31

To intermediate 28 (0.4g, 0.92mmol) in dioxane (8mL) was added cyanogenbromide (0.099g, 0.93mmol) at room temperature. Then sodium hydrogenocarbonate (0.0775g, 0.92mmol) in H_2O (distilled, 4.8mL) was added. The mixture was stirred at room temperature for 5 hours. The mixture was extracted with EtOAc and dried ($MgSO_4$),

filtered and evaporated till dryness. The residue was taken up by diethyl ether, filtered and dried to give 0.42g (99%) of compound 31. MP = 254°C (Kofler).


Example B23



Preparation of compound 32

Hydrazine monohydrate (81µL, 2.58mmol) was added to a solution of intermediate 27 (0.21g, 0.37mmol) in EtOH (10mL). The mixture was heated at 80°C for 5 hours. The mixture was cooled to room temperature, evaporated and the residue was poured into water. The aqueous layer was extracted with DCM, washed with brine, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography (Irregular SiOH, 15-40µm, 10g; mobile phase, 95%DCM – 5%MeOH - 0.5% NH_4OH). The pure fractions were collected and the solvent was evaporated, yielding 97mg (59%) of compound 32, MP= 80°C (Kofler).

Example B24




Preparation of compound 33

Intermediate 31 (270mg, 0.59mmol), sodium triacetatohydroborate (312mg, 1.475mmol) and isopropylamine (100µl, 1.2mmol) in CH_3CN (6mL) were stirred at room temperature for 24 hours. Isopropylamine (500µl, 5.8mmol) was added and the reaction mixture was stirred at room temperature for 12 hours, then sodium triacetatohydroborate (312mg, 1.5mmol) was added and the reaction mixture was stirred for 24 hours. 10% K_2CO_3 aqueous solution was added. The reaction mixture was extracted twice with DCM, dried ($MgSO_4$), filtered and evaporated. The residue (437mg) was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm mobile phase, gradient from 0.2%

NH_4OH, 98% DCM, 2% MeOH to 0.8% NH_4OH, 92% DCM, 8% MeOH). The desired product fraction was collected and the solvent was evaporated, to give 113 mg of compound 33 (cis).


Example B25



Preparation of compound 34

Intermediate 29 (0.5g, 0.8mmol) and methylamine 40% in H_2O (28mL, 0.33mol) were heated in dioxane (20mL) at 120°C in a sealed tube for 5 hours. The solution was cooled and evaporated to dryness. The residue was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 1.4% NH_4OH, 86% DCM, 14% MeOH). The pure fractions were collected and the solvent was evaporated. The product was crystallized with diethyl ether. The precipitate was filtered and dried, yielding 118mg (31%) of compound 34, MP= 174°C (DSC).

Example B26




Preparation of compound 35

To a mixture of intermediate 60 (268mg, 0.51mmol) was added THF (20mL), followed by tetrabutylammonium fluoride (2.53mL, solution 1M in THF; 2.53mmol). The reaction mixture was stirred at room temperature overnight. The reaction mixture was concentrated. EtOAc and H_2O were added and the two phases were separated. The organic phase was dried ($MgSO_4$), filtered and concentrated. The residue was purified by chromatography over silica gel (Hyperprep C18 HS BDS 100A 8mu (Shandon);

mobile phase, gradient from 90% of a 0.25% solution of ammoniumbicarbonate in water, 10% CH_3CN to 100% CH_3CN). The product fractions were collected and the solvent was evaporated. The residue was dissolved in CH_3CN/H_2O and lyophilized, yielding 55mg of compound 35.


Example B27




Preparation of compound 36

A mixture of 7-bromo-2-(1-methyl-1H-pyrazol-4-yl)quinoxaline (521mg, 1.8mmol), intermediate 38 (377mg, 1.8mmol), sodium tert-butoxide (520mg, 5.4mol) in dioxane (10mL) was degassed at room temperature under N_2 flow. After 10 minutes, palladium(II) acetate (47% Pd) (20mg, 0.09mmol) and 1,1'-[1,1'-binaphthalene]-2,2'-diylbis[1,1-diphenylphosphine] (56mg, 0.09 mmol) were added portionwise at room temperature under N_2 flow. The reaction mixture was heated at 90°C overnight, then cooled to room temperature and partitioned between water and DCM. The organic layer was dried ($MgSO_4$), filtered and concentrated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g mobile phase, 98% DCM, 2% MeOH). The product fractions were collected and the solvent was evaporated. The residue was again purified by achiral super critical fluid chromatography on (2 ETHYLPYRIDINE 6µm 150x21.2mm; mobile phase, 0.3% 2-propylamine, 15% MeOH, 85% CO_2). The product fractions were collected and the solvent was evaporated. The residue was taken up in diethyl ether, filtered and dried, yielding 0.209g (27%) of compound 36. MP= 164°C (kofler).

Example B27A




Preparation of compound 920



A mixture of intermediate (see A52) (0.5g; 1.5mmol), intermediate 2 (0.36g; 1.3mmol) and sodium tert-butoxide (0.36g; 1.3mmol) in dry dioxane (40mL) was degassed at room temperature under N_2 flow. After 10 minutes, 2-dicyclohexylphosphino-2'-(N,N-dimethylamino)biphenyl (50mg; 0.13mmol) and tris(dibenzylideneacetone)dipalladium(0) (115mg; 0.13mmol) were added and the reaction mixture was heated at 100°C for 4 hours. The reaction mixture was cooled to room temperature , poured into a mixture of water and brine, filtered through a pad of Celite®, extracted with EtOAc, washed with water, dried ($MgSO_4$), filtered and evaporated to dryness to give 1.1g of residue. The residue was purified by chromatography over silica gel (5µm, mobile phase: gradient from 71% Heptane, 1% MeOH, 28% AcOEt to 20% MeOH, 80% AcOEt). The desired fractions were collected and evaporated to give 240 mg of residue. The residue was taken-up in Et_2O, filtered and dried to give 144mg of compound 920, mp = 123°C (DSC).

Example B28




Preparation of compound 37

At 0°C, potassium permanganate (0.117g, 0.738mmol) was added to a solution of compound 51 (0.326g, 0.738mmol) in acetone (10mL) and H_2O (2.5mL). The solution was stirred at room temperature overnight and was then poured on ice water. DCM was

added and the mixture was filtered through a celite layer. The organic layer was extracted, dried ($MgSO_4$) and evaporated to dryness.The residue (0.23g) was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, 95% DCM, 5% MeOH, 0.1% NH_4OH). The desired product fraction was collected and the solvent was evaporated, yielding 0.150g of compound 37, which was crystallized in DIPE, filtered and dried, yielding 0.139g (40%) of compound 37. MP = 154°C (DSC).



Example B29



Preparation of compound 38


A mixture of intermediate 62 (3.9g, 8.3mmol) and K_2CO_3 (1.15g, 8.3mmol) in MeOH (150mL) was stirred at room temperature for 18 hours. The reaction mixture was quenched with water and extracted with DCM. The organic layer was decanted, washed with brine, dried ($MgSO_4$), filtered and evaporated to dryness. The solid residue was taken up with diethyl ether and the precipitate was filtered off and dried, yielding 2.84g (85%) of compound 38. MP = 168°C, (Kofler).

Example B30

Preparation of compound 39  and compound 40 as a HCl salt 

Intermediate 10 (365mg, 0.75mmol) in 3,5-dimethylpiperidine (5mL) was heated to 80°C overnight. Then 5mL of 3,5-dimethylpiperidine was added to the solution and heated at 80°C for 5 hours. The solution was evaporated to dryness, then the residue was poured into H_2O and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated. The residue (853mg) was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0.1% NH_4OH, 99% DCM, 1% MeOH to 0.8% NH_4OH, 92% DCM, 8% MeOH). The desired product fractions were collected and the solvents were evaporated to give 41.8mg (11%) of Fraction I and 115.7 mg (31%) of compound 39. MP= 80°C (Kofler) (gummed). Fraction I was dissolved in isopropyl alcohol. The mixture was stirred at 0°C, then 67µL (4 eq) of HCl in isopropyl alcohol 5N was added dropwise to the mixture. Diethyl ether was added to the solution and was stirred at 0°C for 1 hour. The precipitate was filtered and dried to give 38.3mg (10%) of compound 40 MP=80°C (Kofler) (gummed).


Example B31




Preparation of compound 41

A mixture of intermediate 37 (0.22g, 0.39mmol), hydrazine monohydrate (0.085 mL, 2.72mmol) in EtOH (5mL) was heated at 80°C overnight. The reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated till dryness. The residue (0.250g) was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase 0.1% NH_4OH, 96% DCM, 4% MeOH). The product fractions were collected and the solvent was evaporated. The residue (0.120g, 70%) was crystallized from diethyl ether/CH_3CN, filtered and dried under vacuum at 60°C, yielding 0.110g (65%) of compound 41. MP168°C(Kofler); 169°C (DSC).

Example B32




Preparation of compound 33 and compound 43



Intermediate 31 (270mg, 0.59mmol), sodium triacetoxyborohydride (312mg, 1.48mmol) and isopropylamine (100µl, 1.2mmol) in CH_3CN (6mL) were stirred at room temperature for 24 hours. Isopropylamine (500µl, 5.8mmol) was added and the reaction mixture was

stirred at room temperature for 12 hours, then sodium triacetoxyborohydride (312mg, 1.5mmol) was added and the mixture was stirred for 24 hours. 10% K_2CO_3 aqueous solution was added. The reaction mixture was extracted twice with DCM, dried ($MgSO_4$), filtered and evaporated. The residue (437mg) was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 0.8% NH_4OH, 92% DCM, 8% MeOH). The product fractions were collected and the solvent was evaporated to give 113mg (38%) of compound 33 and 42mg (14%) of compound 43


Example B33



Preparation of compound 604

N, N-diisopropylethylamine (0.86 mL; 5.2 mmol) and triethylamine (0.73 mL; 5.2 mmol) were added to a solution of intermediate 73 (0.6 g; 0.87 mmol) in methanol (7.5 mL). The reaction was stirred at 80°C for 15 hours, cooled down to room temperature and diluted with DCM and water. The organic layer was separated, washed with water, dried ($MgSO_4$), filtered and evaporated till dryness. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60 g; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected, the solvent was evaporated. The residue (0.25 g, 59%) was crystallized from diethyl ether / CH_3CN. The precipitate was filtered off and dried under vaccum, yielding 215 mg (51%) of compound 604 . MP: 157°C (DSC)

Example B34




Preparation of compound 605

NaH (1.1 g; 27.7 mmol) was added portionwise to *N, N*-dimethylformamide (100 mL), after few minutes intermediate 3 (5 g; 13.8 mmol) was added portionwise at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 30 minutes. Then, a solution of ethyl-2-bromopropionate (3.6 mL; 27.7 mmol) in *N, N*-dimethylformamide (7 mL) was added dropwise at 5°C under N_2 flow. The reaction mixture was allowed to warm to room temperature and stirred for 3 hours. The reaction was poured out into ice water. The precipitate was filtered, washed with water. The organic layer was separated and washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (7.51 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 450g ; mobile phase 0.1% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and concentrated yielding 5.3 g (84%) of compound 605.


Example B35

Preparation of a compound 607 (mixture of enantiomers)



A solution of intermediate 74 (8 g; 16.08 mmol) and potassium phtalimide (6 g; 32.16 mmol) in CH_3CN (110 mL) was heated at 120°C for 2 hours using one single mode microwave. The reaction mixture was cooled to room temperature and poured out into ice water. The precipitate was filtered, washed with water and DCM. The organic layer was separated and washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated to give 7.4 g of compound 607 used without further purification for the next step.


Example B36



Preparation of compound 313

A 1M solution of tetrabutylammonium fluoride in THF (7.7 mL; 7.7 mmol) was added dropwise to a solution of intermediate 76 (3.5g; 5.9 mmol) in THF (75 mL) at room temperature. The reaction mixture was stirred at room temperature for 3 hours. The mixture was poured out into ice water and EtOAc was added. The mixture was basified with K_2CO_3 10% and the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (4.4 g) was crystallized from diethyl ether. The precipitate was filtered off, dried in vacuum to provide 2.62 g (93%) of compound 313 MP: 176°C (DSC).


Example B37



Preparation of compound 615

A mixture of intermediate 77 (2 g; 3.6 mmol) and isopropylamine (1.55 g; 18 mmol) in acetonitrile (30 mL) was heated at 100°C in a sealed vessel for 18 hours. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, EtOAc was added. The organic layer was separated, washed with a solution of $NaHCO_3$, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (2.5 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 450g; mobile phase 0.5% NH_4OH, 90% DCM, 10% MeOH). The pure fractions were collected and concentrated. The residue (0.85 g) was crystallized from diethyl ether, the precipitate was filtered off, dried in vacuum to provide 0.76 g (41%) of compound 615.MP: 134°C (DSC)


Example B38



Preparation of compound 616

Trifluoroacetic acid (6.5 mL; 84.8 mmol) was added to a solution of intermediate 82 in DCM (50mL) at 10°C. The reaction mixture was stirred at room temperature for 18 hours. The reaction mixture was concentrated, the residue was taken-up with DCM, washed with K_2CO_3 10%. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was crystallized from diethyl ether. The precipitate was filtered off, dried in vacuum to provide 0.65 g (65%) of compound 616.MP: 170°C (Kofler)

Example B39




Preparation of compound 617

A 1M solution of tetrabutylammonium fluoride in THF (1.82 mL, 1.8 mmol) was added dropwise to a solution of intermediate 85 (0.88 g, 1.65 mmol) in THF (20 mL) at room temperature. The reaction mixture was stirred at room temperature for 18 hours. The mixture was poured out into ice water and EtOAc was added. The mixture was basified with K_2CO_3 10%, the organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness.

The residue (0.68 g) was purified by chromatography over silica gel (Spherical SiOH, 10 µm, 60g; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated. The residue (0.54 g) was crystallized from diethyl ether. The precipitate was filtered off, dried in vacuum to provide 0.444 g (65%) of compound 617. MP: 149°C (DSC)


Example B40



Preparation of compound 618

A mixture of intermediate 86 (0.446 g; 0.91 mmol) and isopropylamine (6.2 mL; 72.3 mmol) in acetonitrile (14 mL) was heated at 140°C in a sealed vessel for 1 hour using one single mode microwave. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, EtOAc was added. The organic layer was separated, washed with a solution of $NaHCO_3$, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (0.423 g) was purified by chromatography over silica gel (Spherical SiOH, 10 µm, 60g ; mobile phase 0.5% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and concentrated.The residue (0.3 g) was crystallized from diethyl ether, the precipitate was filtered off, dried in vacuum to provide 0.21 g (52%) of compound 618.MP: 139°C (DSC).

Example B41




Preparation of compound 619 as a HCl salt

A mixture of intermediate 87 (1.26 g; 0.99 mmol), hydrazine monohydrate (0.22 mL; 7.0 mmol) in EtOH (20 mL) was stirred at 80°C for 3 hours. The reaction mixture was cooled to room temperature and poured out into ice water. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated until dryness. The residue (0.566 g) was purified by chromatography over silica gel (SiOH, 5µm, 150*30 mm; mobile phase gradient 0.2% NH_4OH, 98% DCM, 2% MeOH to 1.2% NH_4OH, 88% DCM, 12% MeOH). The product fractions were collected and the solvent was evaporated. The residue (0.385g, 77%) was dissolved in isopropyl alcohol. The reaction mixture was stirred at 0°C then 0.6 mL of HCl in isopropyl alcohol 5N was

added dropwise to the solution. Diethyl ether was added to the solution and was stirred at 0°C for 1 hour.

The precipitate was filtered off, dried in vacuum to provide 0.42 g (69%) of compound 619.MP: 210°C (Kofler)


Example B42



a) Preparation of compound 620

A mixture of intermediate10 (1.4 g; 2.9 mmol), tert butyl (1S, 4S)-(-)-2,5-diazabicyclo [2.2.1]heptane -2-carboxylate (0.69 g; 3.5 mmol) and K_2CO_3 (0.8 g; 5.8 mmol) in CH_3CN (20 mL) was stirred at 80°C for 48 hours.

The reaction mixture was cooled to room temperature, poured out into ice water EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated until dryness. The residue (1.6 g) was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 450g ; mobile phase 0.1% NH_4OH, 94% DCM, 6% MeOH). The product fractions were collected and the solvent was evaporated. The residue (0.74 g) was purified with super critical fluid chromatography (Amino 6µm, 150*21.1 mm; mobile phase 90% CO_2, 10% MeOH). The product fractions were collected and the solvent was evaporated. The residue (0.6 g, 36%) was crystallized from diethyl ether. The precipitate was filtered off, dried in vacuum to provide 0.444 g (26%) of compound 620 . MP: 114°C (Kofler)




b) Preparation of compound 621 as a HCl salt

A 5N solution of HCl in i-PrOH (0.48 mL, 2.4 mmol) was added dropwise to a solution of compound 620 (0.35 g, 0.6 mmol) in CH_3OH (10 mL) at 5°C and the mixture was then stirred 3 days at room temperature. Diethyl ether was added and the precipitate was

filtered off, was dried under vacuum to afford 0.33 g (94%) of the compound 621.MP: >260°C (Kofler)


Example B43



Preparation of compound 622

A 1M solution of tetrabutylammonium fluoride in THF (1.1 mL; 1.1 mmol) was added dropwise to a solution of intermediate 88(0.43 g; 0.79 mmol) in THF (6 mL) at room temperature. The reaction mixture was stirred at room temperature for 18 hours. The mixture was poured out into ice water. The precipitate was filtered off, washed with water and CH_3CN and dried to afford 0.13 g (40%) of compound 622. MP: 190°C (Kofler)


Example B44



Preparation of compound 623 as a HCl salt

A mixture of intermediate 89 (0.26 g; 0.43 mmol) and isopropylamine (5 mL) in acetonitrile (2 mL) was heated at 90°C in a sealed vessel for 18 hours. The reaction mixture was cooled to room temperature. The reaction mixture was poured out into ice water, DCM was added. The organic layer was separated, washed, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (0.28 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g; mobile phase 0.4% NH_4OH, 96% DCM, 4% MeOH). The pure fractions were collected and concentrated. The residue (0.156g, 77%) was dissolved in CH_3CN. HCl in isopropyl alcohol 5N was added dropwise to the solution. The solvent was evaporated, dried in vacuum to give 0.162 g (70%) of compound 623. MP: 133°C (Kofler)


Example B45



Preparation of compound 630

NaH (0.54 g; 13.3 mmol) was added portionwise to intermediate 3 (2.4 g; 6.66 mmol) in *N N*-dimethylformamide (36 mL). The reaction mixture was stirred at 0°C for 30 minutes. Then intermediate 91 (2.2 mL; 10 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at room temperature for 18 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (4.2 g) was purified by chromatography over silica gel (Irregular SiOH 15-40µm, 300g; mobile phase 0.1% NH_4OH, 98.5% DCM, 1.5% MeOH). The pure fractions were collected and concentrated to give 0.793 g (21%) of compound 630. MP: 67°C (Kofler).


Example B46



Preparation of compound 632

Trifluoroacetic acid (0.073 mL; 0.25 mmol) was added to a solution of intermediate 92 (0.135 g; 0. 25 mmol) in THF (5 mL). The reaction was stirred at room temperature for 24 hours. The reaction mixture was poured out into ice water; EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.151 g) was purified by chromatography over silica gel (SiOHSiOH, 5µm, 150*30 mm; mobile phase gradient from 70% Heptane, 2% MeOH, 28% EtOAc to 20% MeOH, 80% EtOAc). The pure fractions were collected and concentrated. The residue (0.04 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.033 g (31%) of compound 632. MP: 156°C (DSC)


Example B47



Preparation of compound 638

NaH (0.65 g; 16.6 mmol) was added portionwise to intermediate 3 (3 g; 8.3 mmol) in *N, N*-dimethylformamide (25 mL). The reaction mixture was stirred at 10°C for 30 minutes. Then 3-chloro-3-methyl-1-butyne (1.2 g; 10.8 mmol) was added dropwise under N_2 flow. The reaction mixture was stirred at room temperature for 48 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (4 g) was purified by chromatography over silica gel (mobile phase gradient from 71% Heptane, 1% MeOH, 28% EtOAc to 20% MeOH, 80% EtOAc). The pure fractions were collected and concentrated to give 0.152 g (4%) of compound 638_used without further purification for the next step.

Example B47A




Preparation of compound 919

Sodium hydride (0.24g; 6.1mmol) was added portionwise to a solution of intermediate 3 (1.1g; 3mmol) in DMF (10mL) at 5°C under N_2 flow.The reaction mixture was stirred at 5°C for 30 minutes. Then a solution of (4-chloro-2-butyn-1-yl)-benzene (1g; 6.1mmo)) was added dropwise at 5°C under N_2 flow.The reaction mixture was stirred for 1 hour at 5°C, then stirred at room temperature overnight. The reaction was poured out into ice water and EtOAc was added.The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue was purified by chromatography over silica gel (SiOH 15-40µm 300g, mobile phase: 0.1% NH_4OH, 98% DCM, 2% MeOH). The desired fractions were collected and evaporated to give 0.66g of residue which was then purified by achiral Supercritical Fluid Chromatography on (5µm, mobile phase, 60% CO_2, 40% mixture of MeOH/iPrOH 50/50 v/v). The desired

fractions were collected and evaporated to give 282mg (19%) of compound 919. This fraction was crystallized from Et_2O to give 143mg of compound 919 (mp= 130°C).


Example B48



Preparation of compound 641

A mixture of intermediate 17a (0.3 g; 0.6 mmol), glycinamide hydrochloride (0.2 g; 1.8 mmol), potassium iodide (0.1g; 0.6 mmol), sodium carbonate (0.32 g; 3.0 mmol) in 1-BuOH (12 mL) was stirred at 85°C for 18 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.28 g) was purified by chromatography over silica gel (SiOH 5µm, 150*30 mm; mobile phase gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 0.9% NH_4OH, 91% DCM, 9% MeOH). The pure fractions were collected and concentrated. The residue (0.100 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.081 g (28%) of compound 641. MP: 155°C (DSC).

Example B49




Preparation of compound 137

A mixture of intermediate 93 (12.8 g; 23.4 mmol) and isopropylamine (61 mL; 500 mmol) in acetonitrile (500 mL) was heated at 100°C in a sealed vessel for 18 hours. The reaction mixture was cooled down to room temperature. The reaction was poured out into ice water and EtOAc was added .The organic layer was washed with brine, dried ($MgSO_4$), filtered over silica gel and evaporated till dryness. The residue (13 g) was purified by chromatography over silica gel (Irregular SiOH 20-40µm, 1000g; mobile phase 0.5% NH_4OH, 95%DCM, 10%MeOH). The pure fractions were collected and

concentrated to give 8 g (55%) of the free base which was converted into its HCl salt as compound 137.


Example B50



Preparation of compound 2

A 1M solution of tetrabutylammonium fluoride in THF (30.3 mL; 30.3 mmol) was added dropwise to a solution of intermediate 94 (10.2 g; 20.2 mmol) in THF (70 mL) at room temperature. The reaction mixture was stirred at room temperature for 3 hours. The mixture was poured out into ice water, basified with K_2CO_3 10% and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried to afford 5.9 g (75%) of compound 2 . MP: 169°C (DSC).


Example B51



Preparation of compound 644 as a HCl salt

A mixture of intermediate 100 (0.09 g; 0.14 mmol) and K_2CO_3 (0.058g; 0.42mmol) in MeOH (1.1 mL) was stirred at room temperature for 1 hour. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (0.2 g) was purified by chromatography over silica gel (Irregular SiOH 15-40µm, 30g ; mobile phase 0.5%NH_4OH, 93%DCM, 7%MeOH). The pure fractions were collected and concentrated. The residue was dissolved in EtOH/CH_3CN and acidified with HCl/i-PrOH 5N. The precipitate was filtered off and dried to give 0.053 g (46%) of compound 644as a chlorhydrate.


Example B52



Alternative preparation of compound 93

Intermediate 3 (10 g; 27.7 mmol) was added to a solution of potassium hydroxide (27.4 g; 415 mmol), tetrabutylammonium bromide (1.34 g; 4.0 mmol) in (280 mL) and water (3 mL). The reaction mixture was stirred at 50°C for 30 minutes, then 3-bromopropylamine hydrochloride (9.7 g; 44.3 mmol) was added portionwise and stirred at 50°C for 2 hours. The reaction mixture was cooled to room temperature. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (20 g) was purified by chromatography (Irregular SiOH 20-45µm, 1000g; mobile phase 1%NH_4OH, 90%DCM, 10%MeOH). The pure fractions were collected and concentrated. The residue was crystallized from diethyl ether. The precipitate was filtered and dried to give 10.5 g (90%) of compound 93. MP : 178°C (DSC)


Example B53



Preparation of compound 645


A 5N solution of HCl in i-PrOH (2.5 mL; 12.5 mmol) was added dropwise to a solution of intermediate 105 (0.8 g, 1.54 mmol) in CH_3OH (25 mL) at 10°C and then the mixture was stirred for 18 hours at room temperature. The red precipitate was filtered, rinsed with diethyl ether and dried. The precipitate was taken up with DCM and washed with a solution of NaOH 1M. The organic layer was washed with water, dried ($MgSO_4$), filtered and evaporated till dryness. The residue was crystallized from diethyl ether. The precipitate was filtered and dried under vaccum to give 0.22 g (34%) of compound 645. MP : 188°C (DSC)

Example B54




Preparation of compound 646

A 5N solution of HCl in i-PrOH (1.1 mL; 5.7 mmol) was added dropwise to a solution of




intermediate (0.8 g; 1.4 mmol) (prepared according to the procedure described for intermediate 103 in A42a-c) in CH_3OH (20 mL) at 10°C and then the mixture was stirred for 18 hours. The reaction mixture was taken up with DCM and washed with a solution of hydroxide de sodium 1M, the organic layer was washed with water, dried over $MgSO_4$, filtered and evaporated till dryness. The residue (1.3 g) was purified by chromatography over silica gel (Irregular, SiOH, 15-40µm, 300g ; mobile phase from 0.5% NH_4OH, 93% DCM, 7%MeOH to 1% NH_4OH, 90% DCM, 10%MeOH). The pure fractions were collected and concentrated. The residue was crystallized from diethyl ether, the precipitate was filtered and dried under vaccum to give 0.11 g (19%) of compound 646. MP: 125°C (Kofler)

Example B55



Preparation of compound 647 as a HCl salt


A 5N solution of HCl in i-PrOH (0.7 mL; 3.4 mmol) was added dropwise to a solution of



intermediate (0.5 g; 0.8 mmol) (prepared according to

the procedure described for intermediate 105 in A42e) in CH_3OH (20 mL) at 10°C and then the mixture was stirred 18 hours. The reaction mixture was evaporated till dryness and the residue was taken up with DCM and basified with a solution of sodium hydroxide 1N. The organic layer was washed with water, dried ($MgSO_4$), filtered and evaporated till dryness. The residue was crystallized from diethyl ether and 1mL of HCl 3N. The precipitate was filtered and dried to afford 0.2 g (49%) of compound 647. MP: 133°C (Kofler)


Example B56



Preparation of compound 655

K_2CO_3 (0.38 g; 2.7 mmol) was added to intermediate 109 (1.4 g; 2.7 mmol) in MeOH (40 mL). The solution was stirred at room temperature for 3 hours. The reaction mixture was poured out into water and EtOAc was added. The organic layer was dried ($MgSO_4$), filtered and evaporated till dryness to give 1.2 g of compound 655 .

Example B57




Preparation of compound 658

Intermediate 110 (0.4 g; 0.67 mmol) was hydrogenated at room temperature in MeOH (20 mL) with Nickel de Raney (0.4 g; 6.88 mmol) as a catalyst in pressure vessel (3 bars). After 5 hours the catalyst was filtered off on a pad of Celite® and the filtrate was concentrated in vacuo until dryness. The resiude (0.32 g) was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g ; mobile phase, 0.1% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and evaporated to


dryness. The residue (0.19 g) was crystallized from diethyl ether. The precipitate was filtered and dried to give 0.16 g (42%) of compound 658. MP: 152°C (DSC).

Example B58



Preparation of compound 659


K_2CO_3 (0.17 g, 1.19 mmol) was added to intermediate 111




(0.66 g, 1.19 mmol) (prepared according to the procedure described in A44 starting from intermediate 112) in MeOH (20 mL). The solution was stirred at room temperature for 3 hours.The reaction mixture was poured out into water and EtOAc was added. The organic layer was dried over $MgSO_4$, filtered and evaporated till dryness. The residue was crystallized from CH_3CN and diethyl ether, the precipitate was filtered and dried to give 0.25 g (44%) of compound 659. MP: 106°C (DSC).

Example B59

Preparation of compounds 660 and 661



compound 660




and compound 661

NaH (0.19 g; 4.7 mmol) was added portionwise to intermediate 118 (0.95 g; 2.4 mmol) in *N,N*-dimethylformamide (10 mL). The reaction mixture was stirred at 5°C for 1 hour.


Then 1,2-epoxy-3,3,3-trifluoropropane (0.4 mL; 4.7 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then was allowed to rise to room temperature, stirred for 18 hours and 3 hours at 60°C. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (1.44 g) was purified by chromatography over silica gel (Irregular, SiOH 20-45µm, 450g, , mobile phase gradient from 0.5%NH_4OH, 98% DCM, 2% MeOH to 1%NH_4OH, 88% DCM, 12% MeOH). The pure fractions were collected and evaporated to provide 1.44g of residue. The enantiomers were separated by chiral super critical fluid chromatography (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 0.3% 2-propylamine, 55% CO_2, 45%MeOH). The desired product fractions were collected and the solvent was evaporated. The first eluted enantiomer (0.15 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.11 g (9%) of compound 660 (R*, MP=154°C (DSC)). The second eluted enantiomer (0.15 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.116 g (10%) of compound 661 (S*, MP=151°C (DSC)).

Example B59A

Preparation of compounds 926 (free base) and 892 (HCl salt)



and



as a HCl salt


Compound 926 compound 892

The following reaction was performed two times:

Sodium hydride (2.0 g, 49.8 mmol) was added portionwise to a solution of intermediate 3 (9 g, 24.9 mmol) in DMF (140 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then 1.2-epoxy-3-methylbutane (5.3 mL, 49.8 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred 1 hour at 5°C, then stirred at 80°C for 3 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to give a brown oil. The residue was purified by

chromatography over silica gel (SiOH 20-45µm, mobile phase (0.1% NH_4OH, 97.5% DCM, 2.5% MeOH). The desired product fraction were collected and evaporated to provide 1.2g (11%) of compound 389 and 3.36g (25%) of compound 926. This later fraction was repurified by chromatography over silica gel (SiOH 20-45µm 450g, mobile phase 0.1% NH_4OH, 98% DCM, 2% MeOH). The product fraction was collected and evaporated to provide 1.1 g (8%) of compound 926. A fraction (300 mg) was converted into the HCl salt in MeOH. The solid was filtered, washed with Et_2O and dried to give 159 mg of a red powder compound 892.

Example B60





Preparation of compound 664

NaH (0.11 g; 2.8 mmol) was added portionwise to intermediate 3 (0.5 g; 1.4 mmol) in *N, N*-dimethylformamide (3 mL). The reaction mixture was stirred at 5°C for 1 hour. Then dimethylsulfamoyl chloride (0.3 mL; 2.8 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then was allowed to rise to room temperature and stirred for 6 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.8 g) was purified by chromatography over silica gel (SiOH 5µm, 150*30mm, mobile phase gradient from 100% DCM to 0.4%NH_4OH, 96% DCM, 4% MeOH). The pure fractions were collected and evaporated. The residue (0.05 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.046 g (7%) of compound 664. MP: 80°C (Kofler).

Example B61

Preparation of compound 667 and 668






compound 667 and compound 668

A solution of intermediate 10 (1 g; 2 mmol), 3-methyl-1H-1,2,4-triazole (0.35 g, 4.2 mmol) and K_2CO_3 (0.72 g; 5.2 mmol) in 1-methyl-2-pyrrolidinone (35 mL) was stirred at 135°C for 18 hours. The reaction mixture was cooled down to room temperature and diluted with EtOAc and water. The organic layer was separated, washed with water, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (1.8 g) was purified by chromatography over silica gel (SiOH 20-45µm, 450 g; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and the solvent was evaporated. The residue (0.72 g) was separated by chiral super critical fluid chromatography (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 0.3% 2-propylamine, 50% CO_2, 50% isopropanol). The desired product fractions were collected and the solvent was evaporated. The first product was crystallized from diethyl ether. The precipitate was filtered and dried to give 0.25 g (26%) of compound 667 . MP: 181°C (DSC).
The second product was crystallized from diethyl ether. The precipitate was filtered and dried to give 0.27 g (28%) of compound 668. MP: 137°C (DSC).

Example B62




Preparation of compound 669

The experiment has been performed 6 times on the following amounts.
A mixture of intermediate 3 (0.5 g; 1.4 mmol), diethyl(vinyl)phosphonate (0.5 mL; 3 mmol) and tri-N-butylphosphine (0.035 mL; 0.1 mmol) in CH_3CN (2 mL) was stirred at 140°C for 15 hours in a sealed tube. The reaction mixture was cooled down to room temperature and diluted with DCM and water. The organic layers were separated,


combined, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (7 g) was purified by chromatography over silica gel (SiOH 20-45µm, 450 g; mobile phase 0.1% NH_4OH, 95% DCM, 5% iPrOH). The pure fractions were collected, the solvent was evaporated. The residue (3.1 g) was crystallized from CH_3CN and diethyl ether, the precipitate was filtered off and dried to give 0.88 g (21%) of compound 669 . MP: 122°C (DSC).

Example B63



Preparation of compound 693 as a HCl salt


Trifluoroacetic acid (7 mL; 94.7 mmol) was added to a solution of



d according to a protocol as described in B3B (1.2 g; 2 mmol) in DCM (60 mL). Then the solution was stirred at room temperature for 2 hours. The reaction mixture was poured out into ice water and basified with NH_4OH. The product was extracted with DCM. The organic layer was dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was crystallized from diethyl ether and the precipitate was filtered off.

The precipitate was dissolved in isopropyl alcohol and stirred at 0°C, then 0.8 mL of HCl *i*-PrOH 5N was added dropwise. Diethyl ether was added and the solution was stirred at 0°C for 1 hour. The precipitate was filtered and dried to afford 0.48 g (35%) of compound 693 MP: 151°C (DSC).


Example B64



Preparation of compound 846

Lithium hydroxide monohydrate (0.085 g; 2.0 mmol) was added portionwise to a solution of intermediate 142 (0.72 g; 1.4 mmol) in THF (20 mL) and H_2O (6 mL) at room temperature. The reaction mixture was stirred at 70°C for 24 hours. The reaction mixture was evaporated until dryness. The residue was taken up with diethyl ether. The precipitate was filtered off and dried under vacuum, yielding 0.577 g (88%) of compound 846. MP: 170°C (Kofler)

Example B65





Preparation of compound 763

Reaction performed in a microwave device (biotage) in a sealed tube. Intermediate 88a (198.6 mg, 0.552 mmol), intermediate 131 (520 mg, 1.21 mmol) and tetrakis(triphenylphosphine)palladium (0) (31.89 mg, 0.028 mmol) in toluene (2.6 ml) were stirred at 160 °C for 40 minutes. Toluene (2.6 ml) was added and the reaction mixture was stirred at 160 °C for 40 minutes. Water was added and the reaction mixture was extracted with AcOEt. The organic layer was dried (MgSO4), filtered and dried to give a yellow oil. This oil was crystallized from CH_3CN. The crystals were dried (room temperature) to give compound 763 as a yellow powder. MP : 176 °C.


C. Conversion reactions

Conversion 1

Preparation of compound 44a  and compound 44.  HCl

HCl (5.53mL; 27.65mmol) was added to a solution of compound 6 (3.2g; 5.53mmol) in CH_3OH (70mL) and heated to 60°C for 8 hours. The reaction mixture was cooled to room temperature, poured into H_2O, basified with K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm 30g MERCK; mobile phase, gradient 100% DCM to 90% DCM, 10% MeOH, 0.1% NH_4OH). The desired fraction was collected and the solvent was evaporated. The compound 44a 1.95g (71%) was dissolved in diisopropyl alcohol and HCl (5 to 6N in alcohol) (3mL), stirred for 30 minutes and evaporated to dryness. The residue was crystallized in diethyl ether, yielding 1.537g (47%) of compound 44. MP = 215.29°C (DSC).


Conversion 2



Preparation of compound 45

Compound 9 (3.02g; 5.95mmol) in pyrrolidine (50mL) was heated at 70°C for 2 hours. The reaction mixture was cooled to room temperature and evaporated to dryness. The residue was poured into H_2O and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue (4.04g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 90g; mobile phase, gradient from 100% DCM to 90% DCM/10% MeOH/0.1% NH_4OH). The desired fractions were collected and the solvent was evaporated, yielding 1.83g (57%) of compound 45.

Conversion 2A




Preparation of compound 344 as a HCl salt


A solution of compound 310 (0.93 g; 2.1 mmol), pyrrolidine (0.52 mL; 6.4 mL), K_2CO_3 (0.3 g; 2.2 mmol) in CH_3CN (50 mL) was stirred at 80°C for 24 hours. The reaction mixture was cooled down to room temperature, poured out into ice water and extracted with EtOAc. The organic layer was separated and washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.9 g) was purified by chromatography over silica gel (SiOH, 5µm; mobile phase gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 1.3% NH_4OH, 87% DCM, 13% MeOH). The pure fractions were collected and concentrated. The residue (0.52 g) was dissolved in MeOH and converted into the hydrochloric acid salt with HCl/2-propanol. Et_2O was added and the precipitate was stirred for 30 minutes, filtered off and dried to afford 0.55 g (47%) of compound 344. MP: 162°C (DSC)

Conversion 2B

Preparation of compound 692 and 563




as a HCl salt



as a HCl salt


NaH (0.13 g; 3.3 mmol) was added portionwise to 2,4-dimethylimidazole (0.3 g; 3 mmol) in *N,N*-dimethylformamide (25 mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 30 minutes, then compound 236 (1 g; 2.4 mmol) was added at 5°C under N_2 flow. The reaction mixture was allowed to warm to room temperature and stirred for 18 hours. The reaction was poured out into ice water. The organic layer was separated and washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (1.8 g) was purified by chromatography over silica gel (SiOH, 15-40µm, 300 g; mobile phase 0.5% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and concentrated. The residue (1 g) was purified by achiral super critical fluid chromatography (Amino 6µm; mobile phase 0.3% isopropylamine, 15% MeOH, 85% CO_2). The pure fractions were collected and the solvent was evaporated till dryness. The first fraction (0.44 g) was further purified by chromatography over silica gel (SiOH, 5µm; mobile phase gradient from 0.4% NH_4OH, 96% DCM, 4% MeOH to 1.5% NH_4OH, 85% DCM, 15% MeOH). The pure fractions were collected and concentrated. The residue (0.38 g) was dissolved in acetone, then HCl 4N in dioxane was added dropwise. Diethyl ether was added and the precipitate was filtered and dried to afford 0.39 g (27%) of compound 692 . MP: 157°C (DSC).

The second fraction was dissolved in CH_3CN, then HCl 4N in dioxane was added dropwise. The precipitate was filtered and dried to afford 0.11 g (8%) of compound 563 . MP: 201°C (DSC).



can be prepared according to the above protocol.


Conversion 3



Preparation of compound 46

Hydrazine monohydrate (0.15mL; 4.8mmol) was added to a solution of compound 47 (0.420g; 0.7mmol) in EtOH (20 mL). The mixture was heated at 80°C for 24 hours. The mixture was cooled to room temperature, evaporated and the residue was poured into water. The organic layer was extracted with DCM, washed with brine, dried ($MgSO_4$), filtered and evaporated to dryness. The crude product was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0% NH_4OH, 100% DCM, 0% MeOH to 0.8% NH_4OH, 92% DCM, 8% MeOH). The pure fractions were collected and the solvent was evaporated till dryness, yielding 56mg (71%) of compound 46.

Conversion 4




Preparation of compound 48

Methanesulfonyl chloride (0.093mL, 1.2mmol) was added to a solution of compound 93 (250mg, 0.6mmol) and Et_3N (0.25mL, 1.8mmol) in DCM (10mL) at 5°C. The mixture was stirred at room temperature for 24 hours. The reaction was poured out into ice water and


DCM was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The crude product was crystallized from diethyl ether. The precipitate was filtered and dried under vacuum to give 118mg (40%) of compound 48. MP=189°C (DSC).

Conversion 5



a) Preparation of compound 50


NaH (44.8 mg, 1.12mmol) was added portionwise to a solution of compound 17 (0.3g, 0.75mmol) in DMF (5mL) at 5°C under N_2.The reaction mixture was stirred for 30 minutes, then 1,2-dibromoethane (0.194mL, 2.24mmol) was added dropwise. The reaction mixture was stirred at room temperature for 5 hours, then poured into H_2O/K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm MERCK; mobile phase, gradient 100% DCM to 97% DCM, 3% MeOH, 0.1% NH_4OH). The pure fractions were collected and evaporated to dryness, yielding 0.236g (63%) of compound 50.



b) Preparation of compound 52


Compound 17 (214mg; 0.53mmol), 1-chloro-2-methyl-2-propanol (0.13mL; 1.28mmol), K_2CO_3 (147mg; 1.1mmol) in DMF (9mL) were heated to 120°C for 72 hours. The reaction mixture was cooled to room temperature, poured into H_2O/K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue (277mg) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g; mobile phase, gradient from 100%DCM to 90%DCM, 10%MeOH, 0.1%NH_4OH) The pure fractions were collected and evaporated to dryness. The residue

(226mg) was crystallized in diethyl ether, yielding 178mg (90%) of compound 52. MP=159°C(DSC).




c) Preparation of compound 53

A mixture of compound 54(130mg; 0.38 mmol), iodomethane (23.7 µl; 0.38mmol) and K_2CO_3 (105.3mg; 0.76mmol) in CH_3CN (10 mL) was refluxed overnight. More iodomethane (23.7 µl; 0.38mmol) and K_2CO_3 (105.3mg; 0.76mmol) were added and the reaction mixture was refluxed 8 more hours. The reaction mixture was poured onto water and the product was extracted with EtOAc. The organic layer was washed with brine, dried ($MgSO_4$), filtered off and the solvent was evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g; mobile phase, 0.1% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and the solvent was evaporated till dryness. The residue was crystallized in diethyl ether, filtered and dried, yielding 29mg (21%) of compound 53.




d) Preparation of compound 55

NaH (0.59g, 1.495mmol) was added portionwise to a solution of compound 17 (0.3g, 0.75mmol) in DMF (6mL). The mixture was stirred at 0°C for 1 hour, then 1-(2-oxiranylmethyl)-piperidine (0.316mg, 2.24mmol) was added. The resulting mixture was stirred at 5°C for 1 hour and at 90°C overnight. The mixture was poured out into water and extracted with DCM. The organic layer was dried, filtered and concentrated till dryness. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, 0.7% NH_4OH, 93% DCM, 7% MeOH). The pure fractions were collected and the solvent was evaporated). The pure fractions were collected and the solvent was evaporated, yielding 0.045 g (11%) of compound 55.



e) Preparation of compound 56


NaH (179.3mg, 4.5mmol) was added portionwise to a solution of compound 17 (1.5g, 3.7mmol) in DMF (20mL). The mixture was stirred at 0°C for 1 hour, then (2-bromoethoxy)-tert-butyldimethylsilane (0.96mL, 04.5mmol) was added. The resulting mixture was stirred at room temperature for 4 hours. The mixture was poured out into water and extracted with DCM. The organic layer was dried, filtered and concentrated till dryness to give 2.1g of a crude residue. Tetrabutylammonium fluoride (3.75mL, 1M solution in THF, 3.75mmol) was added dropwise to a solution of the above residue in THF (25mL) at room temperature and stirred at room temperature for 5 hours. The reaction mixture was poured out into ice water, basified with K_2CO_3 and extracted by EtOAc. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and concentrated till dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm 50g ;. mobile phase, gradient 100% DCM to 97% DCM, 3% MeOH, 0.1% NH_4OH).The desired product fractions were collected and the solvent was evaporated, yielding 1.3g (77%) of compound 56 which was triturated in Et_2O, filtered and dried under vacuum at 60°C to give 1.22g (73%) of compound 56. MP=147.5°C (DSC).



f) Preparation of compound 57


Compound 16 (0.02g, 0.046mmol), methyl vinyl sulfone (33µL, 0.4mmol), Et_3N (15.5mL, 0.11mmol) in CH_3OH (2 mL) were heated to 120°C under microwave irradiation for 30 minutes. The mixture was evaporated to dryness and purified by chromatography over silica gel (Irregular SiOH, 15-40 µm, 30g; mobile phase 95% DCM , 5% MeOH , 0.5%

NH_4OH). The desired fractions were collected and the solvent was evaporated, yielding 22.3 mg (90%) of compound 57. MP= 80°C (Kofler).




g) Preparation of compound 58

Dimethylsulfamoylchloride (0.06 mL, 0.56mmol) was added dropwise to a solution of compound 17 (0.15g, 0.37mmol), 4-methylaminopyridine (0.0045g, 0.037mmol), Et_3N (0.104mL, 0.75mmol) in DCM (5mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then overnight at room temperature. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15/40µm, 30g MERCK; mobile phase, gradient 100% DCM to 97% DCM, 3% MeOH). The desired fractions were collected and the solvent was evaporated. The compound was crystallized from diethyl ether, filtered and dried under vacuum at 60°C, yielding 0.065g (34%) of compound 58. MP= 163°C (DSC).



h) Preparation of compound 59


Compound 60 (prepared according to conversion 7 reaction from compound 127) (0.073g, 0.15mmol) was dissolved in DCM (5mL), *N,N*-diisopropylethylamine (0.037mL, 0.23mmol) was added. To this solution, methanesulfonylchloride (0.035mL, 0.23mmol) was added dropwise at 0°C and the mixture was stirred overnight. Water and DCM were added. The organic layer was extracted with DCM. The organic layer was dried, filtered and concentrated. The residue (0.1 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g MERCK; mobile phase, 98%DCM, 2% MeOH). The pure fractions were collected and the solvent was evaporated. The residue (0.089g) was crystallized from DIPE. The precipitate was filtered, dried under vacuum, yielding 0.04 g (47%) of compound 59. MP = 200°C (Kofler).



i) Preparation of compound 51

NaH (0.25mmol) was added portionwise to a solution of compound 17 (0.125 mmol) in DMF (4mL). The mixture was stirred at 5°C for 30 minutes, then allyl bromide (0.19 mmol) was added. The resulting mixture was stirred at room temperature for 2 hours. The mixture was poured into water and the product was extracted with EtOAc. The organic layer was washed with water, brine, dried ($MgSO_4$), filtered and evaporated till dryness, yielding 60 mg (100%) of compound 51.


Conversion 6



Preparation of compound 61

Compound 50 (0.319g, 0.63mmol), K_2CO_3 (0.347g, 2.51mmol), methylamine in 2M THF (0.94 mL, 1.88mmol) in CH_3CN (25mL) were heated at 80°C for 15 hours. The mixture was cooled to room temperature, poured into H_2O/K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, gradient from 0.2% NH_4OH, 95% DCM, 5% MeOH to 0.2% NH_4OH, 90% DCM, 10% MeOH). The pure fractions were collected and evaporated to dryness. The product was crystallized with DIPE and pentane, filtered and dried, yielding 157mg (55%) of compound 61. MP= 103°C (DSC).


Conversion 7



Preparation of compound 62

Compound 63 (0.280g, 0.46mmol), 3N HCl (4mL) and dioxane (4mL) were heated to 60°C for 5 hours. The mixture was cooled to room temperature, poured into H_2O and basified with K_2CO_3. The product was extracted with EtOAc, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was crystallized with DIPE and diethyl ether, filtered and dried, yielding 100mg (43%) of compound 62. MP= 221°C (DSC).


Conversion 8



Preparation of compound 64

Lithium hydroxide monohydrate(43mg; 1.0mmol) was added portionwise to a solution of compound 65 (230 mg; 0.5mmol) in THF (5mL) and H_2O (2mL) at room temperature. The reaction mixture was stirred at room temperature overnight. The reaction mixture was evaporated till dryness. The residue was taken up with water and the mixture was acidified with HCl 3N. After stirring, the precipitate was filtered, washed with water and dried under vacuum, yielding 0.206g (88%) of compound 64.


Conversion 9



Preparation of compound 66

A mixture of compound 67 (0.245g, 0.53mmol), zinc cyanide (0.093g, 0.79mmol), $Pd_2(dba)_3$ (0.024g, 0.026mmol), Zinc (0.017g, 0.26mmol) and 1,1'-bis(diphenylphosphino)ferrocene (0.036g, 0.066mmol) in *N,N*-dimethylacetamide (2mL) was heated at 140°C for 1 hour under microwave irradiation. The reaction was poured out into ice water and DCM was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.27g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g MERCK; mobile phase, gradient from 98% DCM, 2% MeOH to 94% DCM, 6 % MeOH).The pure fractions were collected and the solvent was evaporated. The residue (0.2g, 92%) was crystallized from DIPE. The precipitate was filtered, dried under vacuum, yielding 0.046g (21%) of compound 66. MP=143°C.


Conversion 10



Preparation of compound 68 as a HCl salt

A solution of compound 66 (0.1g, 0.24mmol) and Nickel (0.1g, 1.70mmol) in ammonia and MeOH (4mL of a 7N solution) was hydrogenated under 2 atmospheres of H_2 for 3 hours at room temperature, using Nickel as the catalyst. The catalyst was removed by filtration through celite, washed with DCM and the filtrate was concentrated. The residue (0.1g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g MERCK; mobile phase gradient from 98% DCM, 2% MeOH to 94% DCM, 6% MeOH). The pure fractions were collected and the solvent was evaporated. The residue (0.075g, 74%) was dissolved in iPrOH, 0.11 mL of HCl 5N/ iPrOH was added dropwise at 5°C. The salt was filtered, washed with DIPE and dried under vacumn at 60°C, yielding 0.032g (29%) of compound 68.


Conversion 11



Preparation of compound 69

A mixture of compound 64 (Li-salt) (500mg, 1.18mmol), 1,1,1-trimethyl-*N*-(trimethylsilyl)silanamine (0.5mL, 2.35mmol), *N*3-(ethylcarbonimidoyl)-*N*1,*N*1-dimethyl-1,3-propanediamine hydrochloride (1:1) (365mg, 2.35mmol), HOBt (318mg, 2.35mmol), Et_3N (0.33mL, 2.35mmol) in DMF (80mL) was stirred at room temperature overnight. The mixture was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (167mg) was triturated from diethyl ether, filtered and dried under vacuum to give 141mg (29%) of compound 69. MP = 264°C (DSC).


Conversion 12



Preparation of compound 70 as a HCl salt

HCl (0.496mL; 2.5mmol) was added dropwise to a solution of compound 71 (277mg; 0.50mmol) in isopropyl alcohol (20mL). The reaction mixture was heated at 50°C for 4 hours, then 70°C for 4 hours. The mixture was poured into H_2O and basified with K_2CO_3, then extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, MERCK; mobile phase, gradient 100% DCM to 80% DCM, 20% MeOH, 0.1% NH_4OH). The product fractions were collected and the solvent was evaporated. The residue (110mg) was dissolved in diisopropyl alcohol and HCl (0.2mL of a 5 to 6N in isopropyl alcohol) was added. The mixture was stirred for 30 minutes and evaporated to dryness. Then the residue was crystallized in diethyl ether, yielding 110mg (39%) of compound 70. M.P=163°C (DSC).


Conversion 13



Preparation of compound 72

Formaldehyde (0.045mL, 0.60mmol) was added to a solution of compound 73 (prepared according to conversion 7 reaction from compound 128) (0.15g, 0.30mmol) in MeOH (2mL) and THF (2mL) at room temperature. Then sodium cyanoborohydride (0.028g, 0.45mmol) was added and the mixture was stirred at room temperature for 1 hour. The mixture was poured out into ice. The organic layer was extracted with DCM, dried ($MgSO_4$), filtered off and evaporated till dryness. The residue (0.1g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g MERCK; mobile phase, gradientfrom 95% DCM, 5% MeOH to 80% DCM, 20% MeOH).. The pure fractions were collected and the solvent was evaporated. The residue (60mg, 39%) was crystallized from DIPE/ diethyl ether. The precipitate was filtered, dried under vacuum, yielding 0.046g (30%) of compound 72. MP=120°C (Kofler).

Conversion 14




Preparation of compound 74

A mixture of compound 64 (0.14g, 0.33mmol), methylamine hydrochloride (0.052g, 1.67mmol), *N*3-(ethylcarbonimidoyl)-*N*1,*N*1-dimethyl-1,3-propanediamine hydrochloride (1:1) (0.077g, 0.50mmol), 1-hydroxybenzotriazole (0.068g, 0.50mmol), triethylamine (0.325mL, 2.34mmol) in DCM (14mL) was stirred at room temperature overnight. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Stability Silica 5µm 150x30.0mm; mobile phase, gradient from 0% NH_4OH, 100% DCM, 0% MeOH to

0.7% NH_4OH, 93% DCM, 7% MeOH). The product fraction was collected and the solvent was evaporated. The residue was triturated from diethyl ether, filtered and dried under vacuum at 60°C, yielding 0.078g (54%) of compound 74. MP= 252-254°C (Kofler).


Conversion 15



Preparation of compound 75

Trifluoroacetic acid (1.07mL; 14.37mmol) was added to a solution of compound 76 (2g; 4.79mmol) in H_2O (19.5mL) and dioxane (80mL). The reaction mixture was heated to reflux for 5 hours, poured into H_2O and basified with K_2CO_3, extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 35-40µm, 80g Grace Resolv; mobile phase, gradient 100% DCM to 90% DCM, 10% MeOH, 0.1% NH_4OH). The product fraction was collected and the solvent was evaporated. The residue (2.1g) was crystallized in Et_2O and CH_3CN, yielding 1.61g (77%) of compound 75. MP =187°C (DSC).

Conversion 16




Preparation of compound 75

At 0°C, potassium permanganate (0.11g, 0.7mol) was added to a solution of compound 121 (0.28g, 0.0007mol) in acetone (8mL)/H_2O (2.5mL). The solution was stirred at room temperature for 4 hours and then poured into ice water. DCM was added and the mixture was filtered through a celite layer. The organic layer was extracted, dried ($MgSO_4$) and evaporated to dryness. The residue (200mg) was purified by chromatography over silica gel (Stability Silica 5µm 150x30.0mm;mobile phase gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 1.1% NH_4OH, 89% DCM, 11% MeOH). The

pure fractions were collected and the solvent was evaporated. The residue (100mg, 33%) was crystallized from CH_3CN/diethyl ether, yielding 77mg (25%) of compound 75. MP= 186°C (DSC).


Conversion 17




Preparation of compound 78

Iodomethane (0.5mL, 8.0mmol) was added very slowly to a suspension of Mg (0.196g, 8.0 mmol) in diethyl ether (2mL) at room temperature under N_2. When the Grignard reagent was started, diethyl ether (10mL) was added and the reaction was stirred for 30 minutes. This mixture was added dropwise to a solution of compound 65 (0.240g, 0.54mmol) in THF (12mL) at room temperature under N_2. The reaction mixture was refluxed for 2 hours, then cooled to room temperature. The mixture was poured into H_2O/NH_4Cl and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness to afford a crude residue (0.248g) which was purified by super critical fluid chromatography (CYANO 6µm 150x21.1 mm; mobile phase, 0.3% isopropylamine, 7% MeOH, 93% CO_2). The pure fractions were evaporated yielding 90 mg of compound 78 which was crystallized in Et_2O to afford 57mg (24%) of compound 78. MP = 162°C(DSC).

Conversion 18


Preparation of compound 79a  and compound 79




as a HCl salt

A mixture of compound 76 (0.505gg; 1.21mmol) and methylamine in 2M THF (6.05mLmL, 12.1mmol) in DMF (8mL) was heated at 100°C for 15 hours in a sealed vessel, cooled to room temperature and poured into $H_2O$and K_2CO_3, extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm 30g MERCK; mobile phase, gradient from 100% DCM to 90% DCM, 10% MeOH, 0.1% NH_4OH). The pure fractions were evaporated yielding 0.406g (75%) of compound 79a which was dissolved in diisopropyl alcohol. HCl (5 to 6N) was added. The mixture was stirred for 30 minutes, evaporated to dryness. Then the residue was crystallized in Et_2O, yielding 0.4g (62 %) of compound 79. MP = 224°C (DSC).

Conversion 19


Preparation of compound 80  (E-isomer)

and compound 81  (Z-isomer)

Hydroxylamine hydrochloride (0.043g, 0.62mmol) was added to a solution of compound 82 (0.13g, 0.31mmol) and pyridine (0.13mL) in EtOH (4mL) at room temperature. The mixture was stirred at room temperature overnight. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, 0.1% NH_4OH, 98% DCM, 2% MeOH). Two different residues were collected and the solvent was evaporated for each of them. The first residue was crystallized from DIPE/CH_3CN (90/10). The precipitate was filtered and dried under vacuum, yielding 0.087g (64%) of compound 80 (E-isomer). MP=144°C (Kofler).

The second residue (0.068g) was crystallized from DIPE/CH_3CN (90/10). The precipitate was filtered and dried under vacuum, yielding 0.051g (38%) of compound 81 (Z-isomer). MP=199°C (Kofler).


Conversion 20



Preparation of compound 83

*N*3-(ethylcarbonimidoyl)-*N*1,*N*1-dimethyl-1,3-propanediamine hydrochloride (1:1) (129mg; 0.83mmol) was added to a solution of compound 84 (223mg; 0.55mmol), 1-methyl-3-piperidinecarboxylic acid hydrochloride (1:1) (148.8mg; 0.82mmol), 1-hydroxybenzotriazole (112mg; 0.615mmol), 4-methylporpholine (182µl; 1.66mmol) in DMF (8mL) at room temperature. The reaction mixture was stirred for 24 hours, then poured into H_2O/K_2CO_3 and extracted with EtOAc. The organic layer was dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, 0.5% NH_4OH, 97% DCM, 3% MeOH). The product fractions were collected and the solvent was evaporated. The residue was crystallized in diethyl ether, yielding 122mg (42%) of compound 83. MP142°C (DSC).

Conversion 21




Preparation of compound 85

At 0°C, under N_2, diethylaminosulfur trifluoride (0.224 mL, 1.68mmol) in DCM (2mL) was added dropwise to a solution of compound 56 (0.250g, 0.56mmol) in DCM (4mL). The mixture was stirred overnight at room temperature. An aqueous solution of K_2CO_3 (10%) was added. The mixture was extracted with DCM. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (246mg) was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0% NH_4OH, 100% DCM, 0% MeOH to 0.3% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and evaporated to dryness. The residue

(58mg) was crystallized with DIPE, filtered and dried, yielding 36mg (14%) of compound 85.


Conversion 22



Preparation of compound 86

A mixture of compound 122 (0.5g, 1.21mmol), sodium azide (0.235g, 3.62mmol), ammonium chloride (194mg; 3.62mmol) in *N,N*-dimethylformamide (10mL) was heated at 140°C for 72 hours. The reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the organic layer was separated. The aqueous layer was acidified with HCl 3N. EtOAc was added and the mixture was stirred. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.42g) was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, 93% DCM, 7% MeOH). The product fractions were collected and the solvent was evaporated. The residue (0.110g, 20%) was crystallized from diethyl ether/CH_3CN, filtered and dried under vacuum at 60°C, yielding 0.070g (12%) of compound 86. MP= 196°C (DSC).

Conversion 23




Preparation of compound 87

Diethylaminosulfur trifluoride (276µl; 2.25mmol) was added dropwise to a solution of compound 88 (550mg; 1.12mmol) in DCM (14mL) at 0°C.The reaction mixture was stirred at room temperature for 2 hours and was then poured into H_2O/K_2CO_3. The organic layer was extracted, dried ($MgSO_4$), filtered and evaporated to dryness. The residue (629mg) was purified by chromatography over silica gel (Spherical SiOH, 10µm,

60g PharmPrep MERCK; mobile phase, 0.5% NH_4OH, 97% DCM, 3% MeOH). The product fractions were collected and the solvent was evaporated. The residue (100mg) was purified by achiral super critical fluid chromatography on (2 ETHYLPYRIDINE 6µm 150x21.2mm; mobile phase, 0.3% 2-propylamine, 87% CO_2, 13% MeOH). The product fractions were collected and the solvent was evaporated. The residue (0.08g) was crystallized in Et_2O yielding 72mg (15%) of compound 87.


Conversion 24



Preparation of compound 89

$LiAlH_4$ (0.031g, 0.82mmol) was added portionwise to a mixture of compound 90 (0.2g, 0.41mmol) in THF (10mL) at 5°C under N_2. The mixture was stirred at 5°C for 3 hours. EtOAc followed by H_2O was added dropwise to the mixture at -5°C. The suspension was passed through a short pad of celite. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0% NH_4OH, 100% DCM, 0% MeOH to 0.8% NH_4OH, 92% DCM, 8% MeOH). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from DIPE. The precipitate was filtered and dried under vacuum, yielding 73mg (40%) of compound 89. MP = 126°C (DSC).

Conversion 25




Preparation of compound 91

Copper(I) iodide (52.697mg, 0.28mmol), then *N,N*-diisopropylethylamine (0.829mL, 4.75mmol) were added at 5°C to a solution of compound 38 (1.105g, 2.78mmol) and

ethyl azidoacetate (1.38mL, 5.53mmol) in THF (35mL). The reaction mixture was stirred at room temperature for 18 hours. The mixture was quenched with water and extracted with EtOAc. The organic layer was decanted, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase, 0.1% NH_4OH, 97% DCM, 3% MeOH). The fractions were collected yielding 430mg of residue which was furtherpurified by achiral super critical fluid chromatography on (AMINO 6µm 150x21.2mm; mobile phase, 0.3% 2-propylamine, 90% CO_2, 10% MeOH). The pure fractions were collected and evaporated to dryness yielding two fractions.

The first fraction (90mg) was crystallized from CH_3CN/DiPE. The precipitate was filtered off and dried yielding 74mg (5%) of compound 91, MP=88°C (DSC). The second fraction yielded 360mg (25%) of compound 91.


Conversion 26



Preparation of compound 92

Compound 93 (740mg, 1.77mmol), Et_3N (0.54mL, 3.89mmol) and trifluoroacetic anhydride (0.37mL, 2.65mmol) in THF (25mL) were stirred at room temperature overnight. The reaction mixture was poured into water and extracted with DCM. The organic layer was washed with an aqueous solution of K_2CO_3 (10%), then with water, then dried ($MgSO_4$), filtered and evaporated to dryness. The residue (800mg) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30g; mobile phase, 0.2% NH_4OH, 98% DCM, 2% MeOH). The product fractions were collected and the solvent was evaporated. The residue (730mg) was crystallized from diethyl ether/DIPE to give 465mg (51%) of compound 92. MP=139°C (DSC).


Conversion 27



Preparation of compound 300

A suspension of compound 38 (1.38g; 3.46mmol), 2-iodo-1-methyl-1H-imidazole (0.45g; 2.16mmol) and Et_3N (3.0mL; 21.6mmol) in DMSO (25mL) was degassed under N_2. Dichlorobis(triphenylphosphine)-palladium (304mg; 0.43mmol) and copper(I) iodide (41mg; 0.22mmol) were added and the reaction mixture was stirred at 90°C for 1.5 hours. The reaction mixture was cooled to room temperature, poured onto waterand extracted with EtOAc. The organic layer was decanted, washed with brine, dried ($MgSO_4$), filtered and evaporated to dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g MERCK; mobile phase, 0.4% NH_4OH, 96% DCM, 4% MeOH). The pure fractions were collected and evaporated to dryness. The residue (780mg) was then purified by achiral super critical fluid chromatography on (2 AMINO 6µm 150x21.2mm; mobile phase, 0.3% 2-propylamine, 80% CO_2, 20% MeOH). The pure fractions were collected and evaporated to dryness, yielding 430mg (41%) of compound 300. This fraction was taken up with CH_3CN. The precipitate was filtered off and dried yielding 377mg (36%) of compound 300. MP=192°C (DSC).

Conversion 28




Preparation of compound 94

A mixture of compound 109 (2.5g, 5.29mmol) in NaOH 3M (7mL) and THF (40mL) was stirred at room temperature for 18 hours. The reaction mixture was quenched with a 10% solution of NH_4Cl and EtOAc was added. The pH was adjusted to 4.5 by adding HCl 3N. The organic layer was decanted, washed with NH_4Cl saturated, dried ($MgSO_4$),


filtered and evaporated to dryness. The residue was crystallized from EtOH. The precipitate was filtered off, washed with EtOH, then diethyl ether and dried, yielding 2.02g (86%) of compound 94. MP=101 °C (DSC).

Conversion 29

Preparation of compound 95a




and compound 95 as a HCl salt



A mixture of compound 93 (0.15g, 0.36mmol), 3,3,3-trifluoro-2-hydroxy-2-methylpropanoic acid (0.085g, 0.54mmol), *N*3-(ethylcarbonimidoyl)-*N*1,*N*1-dimethyl-1,3-propanediamine hydrochloride (1:1) (0.083g, 0.54mmol), 1-hydroxybenzotriazole (0.073g, 0.54mmol), Et_3N(0.075mL, 0.54mmol) in DCM (4mL) was stirred at room temperature overnight. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.250g) was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g PharmPrep MERCK; mobile phase, 0.1% NH_4OH, 98% DCM, 2% MeOH). The product fractions were collected and the solvent was evaporated. The compound 95a was dissolved in CH_3CN, cooled to 5°C and a solution of HCl 5N/iPrOH (0.3mL) was added dropwise. The mixture was evaporated till dryness at room temperature. The mixture was taken up with diethyl ether, then the precipitate was filtered off and dried under vacuum at 60°C, yielding 0.172g (80%) of compound 95.


Conversion 30




Preparation of compound 96

Acetone (0.322 mL, 4.361mmol) was added to a solution of compound 70 (0.2g, 0.436mmol) in MeOH (5mL) and THF (5mL) at room temperature. Then sodium cyanoborohydride (0.055g, 0.872mmol) was added and the mixture was stirred at room temperature overnight. Acetone (0.129mL, 1.745mmol) and sodium cyanoborohydride (0.055g, 0.872mmol) were added and the mixture was stirred at room temperature for the weekend. The mixture was poured out into ice, then the organic layer was extracted with DCM, dried ($MgSO_4$), filtered off and evaporated till dryness. The residue (254mg) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g Merck; mobile phase, gradient from 100% DCM to 90% DCM, 10% CH3OH, 0.1% NH4OH). The pure fractions were collected and evaporated to dryness. The product (236mg) was crystallized with DIPE, filtered and dried, yielding 186mg (85%) of compound 96. MP= 168°C (DSC).

Conversion 31


Preparation of compound 97  and

compound 98 

* means relative stereochemistry


The enantiomers of compound 75 (5.4g) were separated by chiral super critical fluid chromatography (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 0.3% 2-propylamine, 40% CO2, 60% MeOH). The desired product fractions were collected and the solvent was evaporated. The first eluted enantiomer (2.1g) was crystallized in diethyl ether, yielding 1.965g (36%) of compound 97 (R*, MP=188°C (DSC)). The second enantiomer (2.1g) was crystallized in diethyl ether, yielding 2g (37%) of compound 98 (S*, MP=186°C (DSC).

Conversion 32

Preparation of compound 99 

A mixture of compound 100 (0.5g, 0.91mmol) in HCl 4M in dioxane (2mL) and CH_3CN (10mL) was heated at 50°C overnight. The mixture was poured out into ice, basified with K_2CO_3 and extracted with DCM. The organic layer was dried ($MgSO_4$), filtered and evaporated till dryness to give 0.4g (99%) of compound 99.


Conversion 33



Preparation of compound 101

Sodium hydride (0.054g, 1.36 mmol) was added portionwise to a solution of compound 99 (0.4g, 0.9 mmol) in DMF (4mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then iodomethane (68 µL, 1.09 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred for 1 hour at 5°C, then at room temperature overnight. The reaction was poured out into ice and extracted with EtOAc. The organic layer was washed with brine, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (0.71g) was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 0.8% NH_4OH, 92% DCM, 8% MeOH). The pure fractions were evaporated till dryness. The residue was crystallized from diethyl ether and dried, yielding 0.172g (42%) of compound 101. MP = 186°C, (Kofler).


Conversion 34



Preparation of compound 102


3,3-Bis(bromomethyl)oxetane (1.592g, 6.52mmol) was added to compound 84 (2.2g, 5.44mmol) and sodium carbonate (0.961g, 9.1mmol) in 1,4-dioxane (80mL) at room temperature. The reaction mixture was stirred at reflux for 7 days, then cooled to room temperature. The mixture was filtered. The filtrate was concentrated under reduced pressure. The residue was purified by chromatography over silicagel (mobile phase, gradient from99%DCM, 1%of a solution of NH_3 in MeOH to 97.5%DCM, 2.5% of a solution of NH_3 in MeOH). The pure fractions were collected and concentrated under reduced pressure, yielding 880mg (33%) of compound 102.


Conversion 35




Preparation of compound 103 as a HCl salt

Sodium cyanide (0.094g, 1.92mmol) was added portionwise to a solution of compound 104 (0.5g, 0.96mmol) in EtOH (10mL) and H_2O (3mL) at room temperature. The reaction mixture was heated at 60°C overnight. The reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the solution was basified with an aqueous solution of K_2CO_3 (10%).The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.63g) was purified by chromatography over silica gel (Spherical SiOH, 10µm 60g PharmPrep MERCK; mobile phase 0.1% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and the solvent was evaporated, yielding 0.37g of compound (75%). This compound was further purified by chiral super critical fluid chromatography on (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 0.3% 2-propylamine, 60% EtOH, 40% CO_2). The desired product fractions were collected and the solvent was evaporated. The residue (0.240g, 49%) was dissolved in CH_3CN and cooled at 5°C. A solution of HCl 5N/i-PrOH (0.28mL) was added dropwise at 5°C. The solution was evaporated till dryness. The residue was triturated with diethyl ether, filtered and dried under vacuum at 60°C, yielding 0.250g (42%) of compound 103.

Preparation of compound 105  as a HCl salt

a-1) Preparation of intermediate 63 


and compound 126 

Methanesulfonyl chloride (0.18mL; 2.31mmol) was added dropwise to a solution of compound 108 (580mg; 1.15mmol), Et_3N (0.4mL; 2.88mmol) in DCM (10mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 2 hours. The reaction mixture was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated, yielding 0.65g (97%) of intermediate 63 and compound 126.

a-2) Sodium cyanide (0.110g, 2.24mmol) was added portionwise to a solution of intermediate 63 (0.65g, 1.12mmol) in EtOH (10mL) and H_2O (3mL) at room temperature. The reaction mixture was heated at 60°C overnight. The reaction mixture was cooled to room temperature and poured out into ice water. EtOAc was added and the solution was basified with an aqueous solution of K_2CO_3 (10%).The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue

(was purified by chromatography over silica gel (Sunfire Silica 5µm 150x30.0mm; mobile phase, gradient from 0% NH_4OH, 100% DCMto 0.5% NH_4OH, 95% DCM, 5% MeOH). The fractions were collected and the solvent was evaporated. The residue was further purified by chiral super critical fluid chromatography (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 0.3% 2-propylamine, 40% EtOH, 60% CO_2). The product fractions were collected and the solvent was evaporated. The residue (0.220g, 38%) was dissolved from CH_3CN and cooled at 5°C. A solution of HCl 5N/iPrOH (0.258mL) was added dropwise at 5°C and the mixture was evaporated till dryness. The residue was triturated from diethyl ether, filtered and dried under vacuum at 60°C, yielding 0.215g (32%) of compound 105.


Conversion 36



Preparation of compound 109

Sodium azide (84.1mg, 1.29mmol) was added at 5°C to a solution of formaldehyde (0.65mL, 8.62mmol) and HOAc (74µl, 1.29mmol) in dioxane (1.5mL). The reaction mixture was stirred for 15 minutes and a solution of compound 38 (310mg, 0.78mmol) in dioxane (1.5mL) was added. The reaction mixture was stirred at 5°C for 10 minutes, then sodium L-ascorbate (34mg, 0.17mmol) was added, followed by a solution of copper sulfate in water (0.53mL, 0.043mmol). The reaction mixture was allowed to warm to room temperature and stirred for 3 hours. Water was added and the reaction mixture was extracted with EtOAc. The organic layer was decanted, washed with brine, dried ($MgSO_4$), filtered and evaporated to dryness, yielding 367mg (100%) of compound 109.


Conversion 37



Preparation of compound 110

To a solution of compound 111 (prepared according to conversion 5a reaction from compound 129) (170mg, 0.29mmol) in DCM (20mL) was added 1-chloroethyl chloroformate (37µl, 0.34mmol) and the reaction was stirred at room temperature for 90 minutes. The solvent was removed under reduced pressure. To the residue was added MeOH (20mL) and the solution was heated to 40°C for 1 hour. The reaction mixture was cooled to room temperature and evaporated under reduced pressure to yield a red solid. The residue (170mg) was purified by chromatography over silica gel (Hyperprep C18 HS BDS 100A 8mu (Shandon); mobile phase, gradient from 80% of a 0.5% solution of ammonium carbonate in water, 20% MeOH to 20% of a 0.5% solution of ammonium carbonate in water ,80% MeOH). The product fractions were collected and the solvent was evaporated, yielding 64mg (44%) of compound 110.


Conversion 38



Preparation of compound 82

Dess-martin periodinane (5.16mL, 1.55mmol) was added dropwise at 0°C to compound 113 (0.59g, 1.41mmol) in DCM (10mL) under N_2 flow. The mixture was stirred at room temperature for 2 hours, poured out into ice and basified with an aqueous solution of K_2CO_3 (10%). The organic layer was extracted with DCM, dried ($MgSO_4$), filtered off and evaporated till dryness. The residue was purified by chromatography over silica gel (Irregular SiOH, 15-40µm; mobile phase, gradient from 98%DCM/ 2% MeOH to 95% DCM/ 5% MeOH). The pure fractions were collected and the solvent was evaporated yielding 0.47g (65%) of compound 82.


Conversion 39



Preparation of compound 114

Sodium hydride (104mg; 2.61mmol) was added portionwise to a solution of compound 115 (500mg; 0.87mmol) in *N,N*-dimethylformamide (8mL) at 5°C under N_2 flow. The reaction mixture was stirred at 5°C for 1 hour, then a solution of iodomethane (0.16mL; 2.61mmol) was added dropwise at 5°C. The reaction mixture was stirred overnight at room temperature.The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The obtained residue (0.55g) was purified by chromatography over silica gel (Irregular SiOH,15/40µm, 30g; mobile phase, gradient from 100%DCM to 96%DCM, 4%MeOH. The product fractions were collected and the solvent was evaporated, yielding 0.39g (76%) of compound 114.

Conversion 40




Preparation of compound 116

Sodium hydride (0.066g, 1.66mmol) was added portionwise to a solution of compound 117 (0.51g, 0.83mmol) in *N,N*-dimethylformamide (10mL) at 5°C under N_2 flow. The mixture was stirred for 1 hour at 5°C, then iodomethane (0.103mL, 1.66mmol) was added portionwise at 5°C .The reaction mixture was stirred for 1 hour at 5°C, then warmed to room temperature. The mixture was stirred at room temperature overnight. The mixture was poured out into ice water and EtOAc was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue was purified by chromatography over silica gel (Irregular SiOH, 15/40µm, 30g MERCK; mobile phase, gradient from 100% DCM to 98% DCM, 2% MeOH). The product


fractions were collected and the solvent was evaporated, yielding 0.400g (76%) of compound 116.

Conversion 41



a) Preparation of compound 118


To a stirred mixture of ethylenediamine (0.226mL, 3.38mmol) and dry toluene (15mL) cooled on an ice bath and under nitrogen was added dropwise trimethylaluminium in heptane (1M, 4mL, 4mmol). The mixture was stirred at room temperature for 60 minutes and then compound 65 (300mg, 0.670mmol) in dry toluene (7mL) was added. The mixture was heated to reflux for 3 hours and was then allowed to cool to room temperature. MeOH (50mL) was added cautiously. The mixture was stirred at room temperature for 5 minutes and then filtered through Celite. The organic layers were concentrated and purified over chromatography over silica gel. The desired fractions were collected and the solvent evaporated, yielding compound 118 ((4,5-dihydro-1H-imidazol-2-ylmethyl)-(3,5-dimethoxy-phenyl)-[3-(1-methyl-1H-pyrazol-4-yl)-quinoxalin-6-yl]-amine) (100 mg).



b) Preparation of compound 119

Compound 118 was heated to 100°C in aqueous sodium hydroxide (2N, 5mL) overnight to promote the ring opening reaction. 1,4-dioxane was added (5mL) and the reaction was continued for a further 10 hours at 100°C. The reaction was allowed to cool and was extracted with EtOAc (2 x). The organic layers were dried ($MgSO_4$) and concentrated. Hydrochloric acid in MeOH was added and the product was precipitated with diethyl ether. The bright red solid was isolated by filtration and was dried in a vacuum oven to give compound 119 (N-(2-Amino-ethyl)-2-{(3,5-dimethoxy-phenyl)-[3-(1-methyl-1H-pyrazol-4-yl)-quinoxalin-6-yl]-amino}-acetamide) (80mg).


Conversion 42



Preparation of compound 120

To a solution of compound 84 (36mg, 0.89mol, 1 equiv.) in dioxane (3mL) and DMF (1.5mL) was added 1-iodo-2-fluoroethane (16mg, 0.89mmol) and K_2CO_3 (25mg, 1.78mmol). The reaction mixture was heated to 90°C for 5.5 hours and a further amount of DMF was added (1.5 mL) and the reaction mixture was heated to 100°C for 1.5 hour. The solvents were removed under reduced pressure and the reaction mixture was partitioned between EtOAc and water. The layers were separated, dried ($MgSO_4$) and concentrated under reduced pressure. The crude mixture was purified by HPLC to give compound 120 (17mg).


Conversion 43



Preparation of compound 124

A solution of compound 76 (0.254g; 0.608 mmol), potassium phtalimide (0.68g, 3.65mmol) in N-methyl-pyrrolidone (5 mL) was heated under microwave irradiation for 1.5 hour at 150°C. The solution was cooled and the mixture was poured into cooled water. The product was extracted with EtOAc. The organic layer was washed with H_2O, dried ($MgSO_4$), filtered and evaporated to dryness to give compound 124 used without further purification for the next step.


Conversion 44



Preparation of compound 125

Compound 124 was heated in EtOH (20mL) with hydrazine monohydrate (0.57 mL; 18.25mmol) at 80°C for 5 hours. The mixture was cooled to room temperature, evaporated and the residue was poured into water. The organic layer was extracted with DCM, washed with brine, dried ($MgSO_4$), filtered and evaporated to dryness to give 400 mg of crude product. The residue was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60g, PharmPrep MERCK; mobile phase 0.5% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and evaporated to give 140 mg (53%) of compound 125. MP = 99°C (DSC).

Conversion 45




Preparation of compound 606

A solution of compound 605 (5.3 g; 11.55 mmol) in THF dry (105 mL) was added dropwise to a solution of lithium aluminium hydride (0.789 g; 20.79 mmol) in THF dry (105 mL) at 0°C under a N_2 flow. The reaction mixture was stirred for 2 hours at 0°C. EtOAc was added dropwise to the reaction mixture, then water was added dropwise. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The residue (5 g) was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 1000g ; mobile phase gradient from 0.1% NH_4OH, 97% DCM, 3% MeOH to 0.1% NH_4OH, 94% DCM, 6% MeOH). The pure fractions were collected and concentrated. The residue (4 g, 75%) was crystallized from DIPE. The precipitate was filtered off and dried under vaccum, yielding 3.5 g (65%) of the compound 606. MP: 97°C (DSC)

Conversion 46

Preparation of compounds 608, 609, 610, 611




*means relative stereochemistry

Compound 607 (7.4g; 6.74 mmol), hydrazine monohydrate (2.52 mL; 80.94 mmol) in EtOH (240 mL) was heated at 80°C for 2 hours. The reaction mixture was cooled to room temperature and poured out into ice water. DCM was added and the organic layer was separated, washed with water, dried ($MgSO_4$), filtered and the solvent was evaporated until dryness. The residue (5.1 g) was purified by chromatography over silica gel (Irregular; SiOH 20-45µm; 450 g ; mobile phase 0.5% NH_4OH, 93% DCM, 7% MeOH). The product fractions were collected and the solvent was evaporated to give 1.1 g of fraction I = compound 896 (enantiomeric mixture) and 1.1 g of fraction II = compound 897 (enantiomeric mixture).

The enantiomers of fraction I and II were separated by chiral super critical fluid chromatography (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 0.3% 2-propylamine, 60% CO_2, 40% isopropanol). The desired product fractions were collected and the solvent was evaporated. The first eluted enantiomer of fraction I (0.52 g) was crystallized in CH_3CN, yielding 0.325 g (12%) of compound 608 (R*, MP=159°C(DSC)). The second enantiomer of fraction I (0.53 g) was crystallized in CH_3CN, yielding 0.284 g (10%) of compound 609 (S*, MP=155°C (DSC)).


The first eluted enantiomer of fraction II (0.47 g) was crystallized in CH_3CN/ diethyl ether, yielding 0.327 g (12%) of compound 610 (R*, MP=150°C (DSC)). The second enantiomer of fraction II (0.475 g) was crystallized in CH_3CN, yielding 0.258 g (9%) of compound 611 (S*, MP=148°C (DSC)).

Conversion 47



a) Preparation of compound 612


Boron tribromide (11.55 mL; 11.55 mmol) was added dropwise to a solution of compound 202 in DCM (10 mL) at 0°C. The solution was allowed to rise slowly to room temperature and stirred for 3 days. The reaction was quenched with MeOH at 0°C. Then, a solution of saturated NH_3 was added to neutralize the reaction mixture. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (1 g) was purified by chromatography over silica gel (C18, 10µm, 250 g, 5 cm; mobile phase 0.25% $(NH_4)_2CO_3$ solution in water, CH_3CN). The pure fractions were collected and concentrated to afford 0.160 g (22%) of compound 612.



b) Preparation of compound 613

Potassium carbonate (0.057 g; 0.41 mmol) was added to a solution of compound 612(0.080 g; 0.21 mmol) in *N,N*-dimethylformamide (15 mL). The reaction mixture was stirred at room temperature for 1 hour. Then methyl iodine (0.013 mL; 0.21 mmol) was added to the reaction mixture and stirred at room temperature for 4 days. The reaction mixture was concentrated under reduced pressure to approximately1/3 of its initial volume. The residue was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.05 g) was purified by chromatography over silica gel (RP Vydac Denali C18, 10µm, 250 g, 5 cm; mobile phase 0.25% $(NH_4)_2CO_3$ solution in water, CH_3CN). The pure fractions were collected and concentrated. The residue (0.025 g) was separated by chiral super critical fluid chromatography (CHIRALPAK Diacel OJ-H 20x25 0mm; mobile phase, CO_2, MeOH with 0.2% 2-propylamine). The desired product fractions were collected and the solvent was evaporated to give 0.007 g (9%) of compound 613.


Conversion 48



Preparation of compound 625


Iodomethane (0.096 mL; 1.54 mmol) was added to a solution of compound 624 (0.73 g; 1.54 mmol) and K_2CO_3 (0.213 g; 1.54 mmol) in CH_3CN (20 mL). The reaction mixture was stirred at 60°C for 5 hours. The reaction mixture was poured out into ice water and EtOAc was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.666 g) was purified by chromatography over silica gel (Spherical SiOH, 10µm, 60 g ; mobile phase gradient from 0.5% NH_4OH, 95% DCM, 5% MeOH to 1% NH_4OH, 90% DCM, 10% MeOH). The pure fractions were collected and concentrated to give 0.3 g (38%) of compound 625. MP: 156°C (DSC).

Conversion 49



a) Preparation of compound 626

Compound 38 (2g; 5.0 mmol) was dissolved in THF (80 mL), then the solution was cooled at -78°C and n-butyllithium 1.6M in hexane (3.1 mL; 5.0 mmol) was added. The reaction mixture was allowed to slowly rise to -30°C and stirred for 45 minutes. 1-Boc-azetidinone (0.715 g; 4.17 mmol) in THF (8 mL) was added to the reaction mixture at -78°C and stirred for 1 hour, then the reaction mixture was allowed to rise to room temperature for 1 hour. The solution was poured out into ice water and NH_4Cl, EtOAc were added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (2.86 g) was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 450 g; mobile phase 0.1% NH_4OH, 96% DCM, 4% MeOH). The pure fractions were collected and concentrated to give 0.343 g (15%) of compound 626 .




b) Preparation of compound 627

Trifluoroacetic acid (1.4 mL; 17.9 mmol) was added to a solution of compound 626 (0.17 g; 0.3 mmol). The reaction was stirred at 70°C for 2 hours. The reaction mixture was cooled to room temperature and evaporated. The reaction mixture was poured out into ice water, DCM was added and basified with NH_4OH. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.35 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30 g; mobile phase 0.5% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and concentrated to give 0.048 g (34%) of compound 627.

Conversion 50


Preparation of compounds 628 and 629



NaH (0.22 g; 5.56 mmol) was added portionwise to compound 14 (0.5 g; 1.1 mmol) in THF (30 mL). The reaction mixture was stirred at 0°C for 1 hour. Then, acetyl chloride (0.8 mL; 11.1 mmol) was added dropwise at 5°C under N_2 flow. The reaction mixture was stirred at 50°C for 18 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.51 g) was purified by chromatography over silica gel (SiOH 5µm, 250*30 mm; mobile phase gradient from 100% DCM to 0.5% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and concentrated to give 220 mg of product. The enantiomers were separated by chiral super critical fluid chromatography (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 60% CO_2, 40% isopropanol). The desired product fractions were collected and the solvent was evaporated. The first eluted enantiomer (0.105 g) was crystallized in diethyl ether, yielding 0.050 g (9%) of compound 628 (S*, MP=122°C). The second enantiomer

(0.096 g) was crystallized in diethyl ether, yielding 0.051 g (9%) of compound 629 (R*, MP=124°C).


Conversion 51



Preparation of compound 631 as a HCl salt

Trifluoroacetic acid (0.52 mL; 6.9 mmol) was added to a solution of compound 630 (0.4 g; 0.7 mmol) in DCM (7 mL). The reaction was stirred at room temperature for 24 hours. The reaction mixture was poured out into ice water, DCM was added and basified with K_2CO_3. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.5 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 30 g; mobile phase 0.5% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated. HCl in isopropyl alcohol 5N was added dropwise to the residue (0.41 g). Acetone and diethyl ether were added.The precipitate was filtered off and dried in vacuum to give 0.383 g (98%) of compound 631. MP: 189°C (Kofler)

Conversion 52




Preparation of compound 633

1,1' carbonyldiimidazole (1.1 g, 6.6 mmol) was added portion wise to a solution of compound 297 (2.4 g; 5.6 mmol) in DCM (60 mL). Then the recation mixture was stirred at room temperature for 1 hour. N,O-dimethylhydroxylamine hydrochloride (0.65 g; 6.6 mmol) was added and the reaction mixture was stirred at room temperature for 72 hours. The reaction mixture was quenched with water and extracted with DCM. The organic layer was decanted, dried ($MgSO_4$), filtered and evaporated to dryness. The residue (2.6 g) was purified by chromatography over silica gel (Irregular SiOH 15-40µm, 300g; mobile phase 0.3% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and concentrated to give 0.185 g (11%). The residue (0.5 g) was crystallized

from diethyl ether. The precipitate was filtered and dried to afford 0.28 g (11%) of compound 633. MP: 130°C (DSC)

Conversion 53




Preparation of compound 635

Compound 634 (0.26 g; 0.53 mmol) was hydrogenated at room temperature in EtOAc (10 mL) with Pd/C (0.05 g) as a catalyst under atmospheric pressure. After 18 hours the catalyst was filtered off on a pad of Celite® and the filtrate was concentrated in vacuo until dryness. The residue (0.256 g) was purified by chromatography over silica gel (SiOHSiOH, 5µm, 150*30 mm; mobile phase gradient from 100% DCM to 0.8% NH_4OH, 92% DCM, 8% MeOH). The pure fractions were collected and concentrated. The residue (0.085 g) was crystallized from CH_3CN/DIPE. The precipitate was filtered and dried to afford 0.075 g (29%) of compound 635. MP: 110°C (DSC)

Conversion 54


Preparation of compounds 637 and 636



Compound 634 (0.38 g; 0.78 mmol) was hydrogenated at room temperature in EtOAc (40 mL) with Lindlar catalyst (0.075 g) as a catalyst under atmospheric pressure. After 9 hours the catalyst was filtered off on a pad of Celite ®, washed with DCM/MeOH and the filtrate was concentrated in vacuo until dryness. The residue (0.474 g) was purified by chromatography over silica gel (SiOHSiOH, 5µm, 150*30 mm; mobile phase gradient from 100% DCM to 0.8% NH_4OH, 92% DCM, 8% MeOH). The pure fractions were collected and concentrated to give two fractions. The first fraction (0.135 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.099 g (26%) of compound 636 (Z) . MP: >260°C (Kofler). The second fraction was crystallized

from diethyl ether. The precipitate was filtered and dried to afford 0.048 g (13%) of compound 637 (E) . MP: 80°C (Kofler).


Conversion 55



Preparation of compound 640

Potassium tert-butoxide (0.054 g; 0.48 mmol) was added to a solution of compound 809 (0.24 g; 0.48 mmol) in THF (15 mL) and the reaction mixture was stirred at 10°C for 2 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.44 g) was purified by chromatography over silica gel (SiOH 5µm, 150*30 mm; mobile phase gradient from 70% Heptane, 2% MeOH, 28% EtOAc to 20% MeOH, 80% EtOAc). The pure fractions were collected and concentrated. The residue (0.132 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.087 g (37%) of compound 640. MP: 241°C (DSC).


Conversion 56



Preparation of compound 642

A mixture of compound 137 (0.51 g; 1.1 mmol), 3-bromopropionitrile (0.11 mL; 1.4 mmol) and K_2CO_3 (0.8g; 5.6 mmol) in CH_3CN (15 mL) was stirred at 80°C for 6 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.5 g) was purified by chromatography over silica gel (SiOH 5µm, 150*30 mm; mobile phase gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 0.9% NH_4OH, 91% DCM, 9% MeOH). The pure fractions were collected and concentrated. The residue (0.35 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.257 g (47%) of compound 642. MP: 127°C (DSC).


Conversion 57



Preparation of compound 643 as a HCl salt

3-hydroxytetrahydrofuran (0.19 mL; 2.3 mmol) and triphenylphosphine (0.61; 2.3 mmol) was added to a solution of compound 137 (0.5 g; 1.16 mmol) in THF (14 mL) under N_2 flow. The reaction mixture was stirred at room temperature for 10 minutes, then diisopropyl azodicarboxylate (0.46 mL; 2.3 mmol) was added and the reaction mixture was stirred for 24 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was separated and washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (2 g) was purified by chromatography over silica gel (SiOH5µm, 150*30 mm; mobile phase 0.5% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and concentrated. The residue (0.19 g) was dissolved in MeOH, 2.3 mL of HCl *i*-PrOH was added, then the chlorhydrate was crystallized from diethyl ether. The precipitate was filtered off and dried to afford 0.178 g (25%) of compound 643.MP: 160°C (Kofler)


Conversion 58



Preparation of compound 648

A mixture of compound 297 (1.65 g, 3.8 mmol), 2,2,2-trifluoroethylamine (1.4 mL, 9.4 mmol), 1-hydroxybenzotriazole (3.6g, 9.4 mmol), triethylamine (1 mL, 7.5 mmol) in *N,N*-dimethylformamide (50 mL) was stirred at room temperature for 18 hours. The reaction mixture was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (3.2 g) was purified by chromatography over silica gel (Irregular Silica 20x40; mobile phase 0.1% NH_4OH, 97% DCM, 3% MeOH). The product fraction was collected and the solvent was evaporated. The residue was triturated from diethyl ether, filtered and dried under vacuum at 60°C, yielding 1.15 g (65%) of compound 648, MP= 196°C (DSC).


Conversion 59



Preparation of compound 651

Trifluroacetic acid (1 mL; 14.3 mmol) was added to a solution of compound 650 (0.44g; 0.7 mmol) in DCM (5.2 mL) at 0-5°C. The reaction mixture was stirred at room temperature for 2 hours, then was quenched with K_2CO_3 10%. The organic layer was washed with water, dried ($MgSO_4$), filtered and evaporated till dryness. The residue (0.45 g) was purified by chromatography over silica gel (Irregular Silica 15x40; 30g , mobile phase 1% NH_4OH, 90% DCM, 10% MeOH). The product fraction was collected and the solvent was evaporated. The residue was crystallized from diethyl ether/CH_3CN, filtered and dried under vacuum at 60°C, yielding 0.26 g (72%) of compound 651, MP= 122°C (Kofler).

Conversion 60




Preparation of compound 652

A suspension of compound 38 (1g; 3.5 mmol), 2-bromo-3-methoxypyridine (0.25 g; 0.35 mmol) and Et_3N (3.0 mL; 21.5 mmol) in DMSO (20 mL) was degassed under N_2 flow. Dichlorobis(triphenylphosphine)-palladium (0.25 g; 0.36 mmol) and copper(I) iodide (0.034 g; 0.18 mmol) were added and the reaction mixture was stirred at 90°C for 40 minutes. The reaction mixture was cooled to room temperature, poured out into water and EtOAc was added. The mixture was filtered off on a pad of celite®. The organic layer was decanted, washed with brine, dried ($MgSO_4$), filtered and evaporated to dryness. The residue (1.4 g) was purified by chromatography over silica gel (Irregular SiOH, 20-40µm, 450g ; mobile phase, 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and evaporated to dryness. The residue was crystallized from diethyl


ether. The precipitate was filtered and dried to give 0.6 g (65%) of compound 652. MP: 144°C (DSC)

Conversion 61

Preparation of compounds 656 and 657



as a HCl salt




as a HCl salt

Compound 14a (3.4 g) was purified by chromatography over silica gel (Irregular SiOH, 15-40µm, 300g; mobile phase, 0.1% NH_4OH, 98% DCM, 2% MeOH). The pure fractions were collected and evaporated to dryness.The residue (1 g) was separated by chiral super critical fluid chromatography (CHIRALPAK AD-H 5µm 250x20mm; mobile phase, 40% 2-propylamine, 60% CO_2). The desired product fractions were collected and the solvent was evaporated. The first eluted enantiomer (0.5 g) was dissolved in diethyl ether, 5 equivalents of HCl in *i*-PrOH were added and stirred at room temperature for 18 hours. The precipitate was filtered and dried to afford 0.29 g (8%) of compound 656(R*, MP=95°C (Kofler)). The second enantiomer (0.55 g) was purified by achiral SFC on (Amino 6µm 150*21.2 mm, mobile phase, 90% CO_2, 10%MeOH). The desired product fractions were collected and the solvent was evaporated. The residue (0.47 g) was dissolved in diethyl ether, 5 equivalents of HCl in *i*-PrOH were added and stirred at room temperature for 18 hours. The precipitate was filtered and dried to afford 0.36 g (11%) of compound 657 (S*, MP=110°C (Kofler)).


Conversion 62

Preparation of compound 663



Compound 662 (0.25 g; 0.49 mmol) in HCl (1M in H_2O) (12.2 mL; 12.2 mmol) was stirred at 60°C for 24 hours. The reaction mixture was cooled down to room temperature, evaporated till dryness. Then the residue was taken up DCM and washed with K_2CO_3 10%. The organic layer was separted and dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (0.2 g) was purified by chromatography over silica gel (SiOH 5µm, 150*30mm, mobile phase, gradient from 0.2%NH_4OH, 98% DCM, 2% MeOH to 1.1%NH_4OH, 89% DCM, 11% MeOH). The pure fractions were collected and evaporated. The residue was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.1 g (41%) of compound 663. MP: 200°C (DSC).


Conversion 63



Preparation of compound 670 1,1'carbonyldiimidazole (0.5 g, 3 mmol) was added to a solution of compound 125 (1.2 g, 2.8 mmol) in THF (20.5 mL) at 0°C under N_2 flow. The reaction mixture was stirred at room temperature for 2 hours. The reaction was poured out into ice water and EtOAc was added. The organic layer was washed with brine, dried ($MgSO_4$), filtered and evaporated. The residue (1.3 g) was purified by chromatography over silica gel (Irregular SiOH, 20-45µm, 300 g; mobile phase 0.2% NH_4OH, 96% DCM, 4% iPrOH). The pure fractions were collected, the solvent was evaporated. The residue (0.98 g) was crystallized from CH_3CN and diethyl ether. The precipitate was filtered off and dried to give 0.8 g (64%) of compound 670 ._MP: 157°C (DSC).


Conversion 64

Preparation of compound 671 and 672



A mixture of compound 76 (1.5 g; 3.6 mmol) and 3-methyl-1H-1,2,4-triazole (3.7 mL; 28.9 mmol) in 1-methyl-2-pyrrolidinone (4 mL) in a sealed tube was heated at 140°C using one single mode microwave (Biotage Initiator EXP 60) for 40 minutes. The reaction mixture was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The crude product (2.1 g) was purified by chromatography over silica gel (15-40µm 300 g; mobile phase, 0.5% NH_4OH, 93% DCM, 7% MeOH). The pure fractions were collected and the solvent was evaporated till dryness. The residue (1 g) was purified by chromatography over silica gel (Cyano 6µm 150*21 mm; mobile phase, 90 %CO_2, 10% EtOH). The desired product fractions were collected and the solvent was evaporated. The first isomer (0.3 g) was crystallized in CH_3CN / diethyl ether, yielding 0.26 g (15%) of compound 671 MP=144°C (DSC). The second isomer (0.34 g) was crystallized in CH_3CN / diethyl ether, yielding 0.26 g (15%) of compound 672 MP=194°C (DSC).


Conversion 65



Preparation of compound 673


A mixture of compound 584 (0.64 g; 1.2 mmol) and methylamine in 2M THF (3 mL; 6 mmol) in 1-methyl-2-pyrrolidinone (5 mL) was heated at 140°C for 24 hours in a sealed tube. The reaction mixture was poured out into ice water and EtOAc was added. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The crude product (1 g) was purified by chromatography over silica gel (5µm; mobile phase gradient from 100% DCM to 0.6% NH_4OH, 94% DCM, 6% MeOH). The pure fractions were collected and the solvent was evaporated till dryness. The residue was crystallized from acetone and diethyl ether. The precipitate was filtered and dried to give 0.34 g (58%) of compound 673. MP: 180°C (Kofler)

Conversion 66




a) Preparation of compound 674

Chloroacetyl chloride (0.23mL; 2.9 mmol) was was added dropwise to a solution of compound 409 (1.3 g , 2.7 mmol) and triethylamine (1.14 ml, 8.2 mmol) in acetonitrile (40 ml) at 0°C under nitrogen . The raction mixture was stirred for 2 hours at room temperature, then at 110°C overnight. Water was added and the reaction mixture was extracted with DCM, dried ($MgSO_4$), filtered and dried to provide 1.5g of compound 674 used without further purification in the next step.



b) Preparation of compound 675

Potassium tert butoxide was added portionwise to a solution of compound 674 (2.6 g; 4.7 mmol) in isopropanol (58 mL) and THF (58 mL). The reaction mixture was stirred at room temperature for 2 hours, then was poured out into ice water and DCM was added. The organic layer was separated, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The crude product (2.1 g) was purified by chromatography over silica gel (20-45µm 450 g; mobile phase gradient from 0.2% NH_4OH, 96.5% DCM, 3.5% MeOH to 1% NH_4OH, 89% DCM, 10% MeOH). The pure fractions were collected and the solvent was evaporated till dryness. The residue (0.61 g) was crystallized from diethyl ether and CH_3CN. The precipitate was filtered and dried to give 0.49 g (21%) of compound 675.MP: 187°C (Kofler)




c) Preparation of compound 676

Lithium aluminium hydride (0.028 g; 0.73 mmol) was added to a solution of compound 674 (0.25 g; 0.48 mmol) in THF (20 mL) under N_2 flow between 0-5°C.The reaction mixture was stirred between 0-5°C for 1 hour. EtOAc was added dropwise to the reaction mixture, then water was added dropwise. The organic layer was separated, washed with brine, dried ($MgSO_4$), filtered and the solvent was evaporated to dryness. The crude product (1 g) was purified by chromatography over silica gel (5µm; mobile phase, gradient from 100% DCM to 0.6% NH_4OH, 94% DCM, 6% MeOH). The pure fractions were collected and the solvent was evaporated till dryness. The residue (0.155 g) was purified by achiral Super critical fluid chromatography (2-ethylpyridine 6µm; mobile phase 0.3% isopropylamine, 20% MeOH, 80% CO_2). The pure fractions were collected and the solvent was evaporated till dryness.

The residue (0.053 g) was purified by chromatography over silica gel (15-40µm 10 g; mobile phase gradient from 100% DCM to 0.6% NH_4OH, 94% DCM, 6% MeOH). The pure fractions were collected and the solvent was evaporated till dryness to afford 0.043 g (18%) of compound 677 MP: 88°C (Kofler)

Conversion 67




a) Preparation of compound 678

The experiment has been performed 3 times on the following amount.

A mixture of compound 137 (HCl salt) (1 g; 2.3 mmol), 2-bromoethyl-methylsulfone (0.5 mL; 2.8 mmol) and K_2CO_3 (0.6 g; 4.6 mmol) in CH_3CN (33 mL) was stirred at 80°C for 2 hours. The reaction was poured out into ice water and EtOAc was added. The organic layers were separated and washed with brine, combined, dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (5.2 g) was purified by chromatography over silica gel (SiOH 15-40µm, 450; mobile phase gradient from 0.5% NH_4OH, 96% DCM, 4% MeOH to 0.5% NH_4OH, 90% DCM, 10% MeOH). The pure fractions were collected and concentrated. The residue (3.2 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 2.2 g (78%) of compound 678. MP: 148°C (DSC).


Conversion 68



a) Preparation of compound 680

Compound 681 (0.97 g; 1.4 mmol) in trifluoroacetic acid (28.5mL) was heated at 100°C for 24 hours in a sealed tube. The reaction mixture was evaporated till dryness. The crude product was diluted in DCM and basified with $NaHCO_3$. The organic layer was separated and dried ($MgSO_4$), filtered and the solvent was evaporated. The residue (1.2 g) was purified by chromatography over silica gel (SiOH 15-40µm, 300 g; mobile phase gradient from 0.5% NH_4OH, 92% DCM, 8% MeOH to 0.5% NH_4OH, 90% DCM, 10% MeOH). The pure fractions were collected and concentrated. The residue (0.4 g) was purified by chromatography over silica gel (SiOH 10µm, 60 g; mobile phase 0.5% NH_4OH, 93% DCM, 7% MeOH). The pure fractions were collected and concentrated. The residue was crystallized from DIPE/CH_3CN. The precipitate was filtered and dried to give 0.29 g (45%) of compound 680. MP: 167°C (DSC).


Conversion 69



a) Preparation of compound 682

Palladium 10% on carbon (0.65 g; 6 mmol) was added to a solution of compound 10 (1.5 g; 2.7 mmol) in MeOH (30 mL). The reaction mixture was stirred at room temperature under 3 bars. After 24 hours the catalyst was filtered off on a pad of Celite® and the filtrate was concentrated. The residue (1.2 g) was purified by chromatography over silica gel (SiOH 20-40µm, 450 g; mobile phase gradient from 0.2% NH_4OH, 96% DCM, 4% MeOH to 0.2% NH_4OH, 95% DCM, 5% MeOH). The pure fractions were collected and concentrated. . The residue (0.25 g) was purified by achiral super critical fluid chromatography (2-ethylpyridine 6µm; mobile phase 0.3% isopropylamine, 20% MeOH, 80% CO_2). The pure fractions were collected and the solvent was evaporated till dryness. The residue was crystallized from diethyl ether and CH_3CN. The precipitate was filtered off and dried to afford 0.15 g (12%) of compound 682.MP: 149°C (Kofler).


Conversion 70



a) Preparation of compound 683

A mixture of compound 84 (1 g; 2.5 mmol) and 1.2-epoxy-3.3.3-trifluoropropane (0.4 mL; 4.9 mmol) in MeOH (15 mL) was heated at 60°C for 2 hours. The reaction mixture was cooled down to room temperature and evaporated till dryness. The residue (1.6 g) was purified by chromatography over silica gel (SiOH, 15-40µm, 300 g; mobile phase gradient from, 0.1% NH_4OH, 98% DCM, 2% MeOH to 0.1% NH_4OH, 97% DCM, 3% MeOH). The pure fractions were collected and concentrated. The residue (0.56 g) was crystallized from diethyl ether. The precipitate was filtered and dried to afford 0.2 g (16%) of compound 683. MP: 123°C (DSC).


Conversion 71



Preparation of compound 685 as a HCl salt

At 5°C, A 5N solution of HCl in *i*-PrOH 5/6N (2.4 mL; 12 mmol) was added to a solution of compound 686 (0.9 g; 1.7 mmol) in CH_3OH (3 mL). The reaction mixture was stirred at 5°C for 2 hours, then for 15 hours at room temperature. The precipitate was filtered off and dried under vaccum to afford 0.425 g (52%) of compound 685. MP=203°C (Kofler).


Conversion 72



Preparation of compound 696 as a HCl salt

Hydrogen chloride (4M in dioxane) (6.8 mL; 27.2 mmol) was added to a solution of compound 695 (1.9 g; 3.4 mmol) in CH_3CN (37 mL) and was stirred at 50°C for 18 hours. The reaction mixture was poured out into ice water. The precipitate was filtered and dried to give 0.3 g (15%) of compound 696. MP: 188°C (Kofler)


Conversion 73

Preparation of compound 902 

.HBr


A mixture of compound 669 (200 mg, 0.38 mmol) and bromotrimethylsilane (3.16 ml , 23.975 mmol) in anhydrous DCM (4 ml) was stirred at room temperature for 3 hours. The solvent was evaporated, the resulting residue was diluted with MeOH-Water (1:1, 10 ml) and stirred for 20 minutes. The precipitate was filtered, washed with AcOEt and dried to give 149 mg (82%) of compound 902.

Conversion 74

Preparation of compound 906 

Compound 93 (340mg, 0.81mmol) was added at 0°C to trifluoroacetaldehyde methyl hemiketal (311µL, 3.25 mmol) and the mixture was stirred at 0°C for 4h30. The mixture was evaporated and the residue purified by chromatography over silica gel (5µm. mobile phase: gradient from 0.2% NH_4OH, 98% DCM, 2% MeOH to 1.3% NH_4OH, 87% DCM, 13% MeOH). The desired product fractions were collected and the solvent was evaporated.yielding 41 mg. The residuet was taken up in Et_2O filtered and dried.to give 29 mg of compound 906.


Conversion 75



Preparation of compound 918

A mixture of compound 584 (397 mg,; 0.75 mmol), dimethylamine (3 mL of a 2.0 M solution in tetrahydrofuran; 6 mmol) in 1-methyl-2-pyrrolidinone (11 mL) was stirred at 140°C for 24 hours in a sealed tube. The mixture was poured into ice-water and EtOAc was added. The organic layer was separated, dried ($MgSO_4$), filtered and the filtrate was evaporated until dryness to afford 607 mg of residue, which was purified by chromatography over silica gel (15-40µm, 90g, mobile phase DCM/CH_3OH/NH_4OH: 98/2/0.1) The desired fractions were collected and evaporated to dryness to give 461 mg of residue which was repurified by chromatography over silica gel (5µm, mobile phase: gradient from 100% DCM to 0.6% NH_4OH, 94% DCM, 6% MeOH) The desired fractions were collected and evaporated to dryness to give 390 mg. This residue was purified by achiral Supercritical Fluid Chromatography on (Diethylaminopropyl 5µm, mobile phase 0.3% isopropylamine, 92% CO_2, 8% MeOH). The desired fractions were collected and evaporated to dryness to give 233 mg of residue which were.crystallized from Et_2O. The precipitate was filtered and dried to give 211 mg (57%) of compound 918.

Conversion 76












Preparation of compound 757



















At 5°C, under N_2 atmosphere, NaH (447.83 mg, 11.2 mmol) was added to a mixture of compound 4 (2g, 4.48 mmol) in DMF (40mL). The reaction mixture was stirred at 10°C for 30 minutes, then iodomethane (0.335 ml, 5.375 mmol) was asdded dropwise. The reaction mixture was cooled to room temperature and stirred at room temperature for 2 hours. Poured into H_2O+NaCl and extracted with AcOEt. The organic layer was washed with H_2O, dried ($MgSO_4$), filtered and evaporated to dryness to give 2g of residue. The residue was purified by flash chromatography over silica gel (15-40µm, 40g, CH_2Cl_2/CH_3OH/NH_4OH: 96/4/0.1). The pure fractions were collected and evaporated to dryness to give 2 fractions: 1.05g of compound 757 and 0.3g of compound 757.



















The following compounds were prepared according to reaction protocols of one of the above Examples using alternative starting materials as appropriate. Those indicated with NMR* have NMR data hereinafter.
















In the table =CoX (or =BX) indicates that the preparation of this compound is described in Conversion X (or Method BX).
In the table ~CoX (or ~BX) indicates that this compound is prepared according to Conversion X (or Method BX).
















As understood by a person skilled in the art, compounds synthesised using the protocols as indicated may exist as a solvate e.g. hydrate, and/or contain residual solvent or minor impurities. Compounds isolated as a salt form, may be integer stoichiometric i.e. mono- or di-salts, or of intermediate stoichiometry.



















		 as a HCl salt
200;~B5b-1;NMR*	204;~B5b-2;NMR*	209;~B5
		
201;~B5b-1;NMR*	13;=B5 b-3;NMR*	210;~B5
		
11;=B5 b-1;NMR*	205;~B5b-2;NMR*	211;~B5
















		
202;~B5/~B5b-1;NMR*	206;~Co 5a	1;=B1
	 as a HCl salt	
12;=B5 b-2;NMR*	207;~B11	212;~B12
		
17;=B9 a/b/c	56;=Co5e/~B14	213;~B12
		
203;~B9c	208;~B5	127;~B9c
		
214;~Co5g	217;~B9a	222;~B5
		
215;~B5	63;~Co5a	223;~Co5a



















		
216;~B12	3;=B2 a/b	224;~B5
		
22;=B13	218;~B5	225;~Co5a
		 as a HCl salt
60;~Co7	219;~B3/B4a	62;=Co7
		
122;~B5	220;~B5	23;=B14
		
58;=Co5g	221;~B5	226;~B12
		 as a HCl salt
















8;=B4A ~B3a/B3/B4a	230;~Co5a	64;=Co8
		
84;~Co3;NMR*	128;~Co5a	233;~B1/B2a
		
227;~Co5a	73;~Co7	19;=B11
		 as a HCl salt
65;~B5 b-1	231;~Co5f	234;=B11
		
228;~B5	59;=Co5h	116;=Co40
 as a HCl salt		 as a HCl salt
229;~Co6	52;=Co5b	130;~Co6;NMR*
















		
117;~B9c	232;~B5	235;~Co7
	 as a HCl salt	
69;=Co11	68;=Co10	242;~B5
	 as a HCl salt	 as a HCl salt
236;~B5	239;~Co7	243;~Co6
		 as a HCl salt
237;~B5	240;~Co7	244;~Co5a
 as a HCl salt		
238;~B1/B2a	67;~B5	245;~B5
















		
66;=Co9	50;~Co5a	246;~B12
	 as a HCl salt	
2;~B1a/B17;=B50	241;~Co2/B5	24;=B15
		 as a HCl salt
113;~B6	55;=Co5d	247;=B15
		
248;~B12	252;~B5	256;~B5
		
249;~B1/B2a	253;~B12	26;=B17
		



















250;~B12	254;~B12	257;~B5
		
251;~Co7	74;=Co14	258;~B13
		
21;=B12a/~B1/B2a	255;~B1/B2a/B3/B4a	75;=Co15/Co16
		
71;~B5	70 ;=Co12	7;~ B3/B3A
	 as a HCl salt	
72;=Co13	70a;=Co12	259;~B14
	 as a HCl salt	
260;~B1/B2a	264;~B3/B4a	82;~Co38
















		
27;=B18	265;~B3/B4a	270;~B5
	 as a HCl salt	
78;=Co16/~B6	266;~B3/B4a	271;~Co5g
		
261;~Co12	76;~B5	272;~Co2
		
262;~Co4	267;~Co5g	273;~B5
 as a HCl salt		 as a HCl salt
25;=B16	268;~Co4	274;~B5
	 as a	
















	HCl salt	
263;~B5	269;~B5	121;~B5
		 as a HCl salt
275;~B3/B4a	37;=B28	79a and 79;=Co18
		 as a HCl salt
4;=B3;NMR*	278;~B3/B4a	602;~B3/B4a
		
9;~B5	279;~Co1	80;~Co19
		
45;~Co2	280;~B3a/B3/B4a	81;=Co19
		
















276;~B1/B2a	281;~B15	284;~Co1
	 as a HCl salt	 as a HCl salt
28;=B19	282;~B15	285;~Co1
	 as a HCl salt	
277;~B5	283;~B2b	20;=B12
		
286;~B3/B4a	88;~Co2/B3/B4a	295;~Co5f
	 as a HCl salt	
287;~B3/B4a	291;~Co2/B3/B4a	296;~Co18
		 as a HCl salt
29;=B20	292;~B3/B4a	54;;~Co1
















 as a HCl salt		 as a HCl salt
288;~B15	293;~Co1	297;~Co8
	 as a HCl salt	
289;~B3/B4a	294;~Co1	298;~B8
		 as a HCl salt
290;~B5	16;=B8	299;~B3/B4a
		
49;~B1	305;~Co5a	312;~B19
		
301;~Co3	61;=Co6	313;=B36
















		
93;~Co3;;NMR;=B52*	306;~Co14	314;~Co14
 as a HCl salt		
302;~B10	307;~Co14	30;=B21
 as a HCl salt	 as a HCl salt	
303;~B3/B4a	308;~B10	38;~B29
		
304;~B10	310;~B5	315;~B10
		
123;=A9c/~B5	311;~Co1	48;=Co4
		
















316;~Co4	57;=Co5f	83;=Co20
		
317;~B3/B4a	323;~B9a	327;~B10
		
318;~Co1	85;=Co21	328;~B7
		 as a HCl salt
319;~B3/B4a	324;~B3/B4a	329;~B7
		 as a HCl salt
320;~Co5f	325;~Co14	330;~B3/B4a
 as a HCl salt		
321;~B7	86;=Co22	331;~B3/B4a
















		 as a HCl salt
322;~B5	326;~B19	332;~B3/B4a
 as a HCl salt		
333;~Co8	339;~B10	343;~B3/B4a
		 as a HCl salt
334;~Co14	32;=B23	344;=Co2A
	 as a HCl salt	
335;~Co4	131;~B8;NMR*	31;=B22
		
336;~B5	90;~B5	345;~B14
















		
337;~Co1	340;~B14	346;~B9
		 as a HCl salt
109;=Co36	341;~Co5f	347;~B3/B4a
	 as a HCl salt	 as a HCl salt
338;~Co1/Co3	342;~B3/B4a	348;~B3/B4a
		
349;~B5	354;~B5	358;~B5
		
350;~B3/B4a	355;~B5	359;~B5
















		
351;~B3/B4a	356;~B5	360;~B5
		 as a HCl salt
89;=Co24	53;=Co5c	361;~Co2
		
352;~B12	91;=Co25	362;~B3/B4a
 as a HCl salt		
87;=Co23	18;=B10	363;~Co3
		
353;~B3/B4a	357;~B3/B4a	364;~B7
















		 as a HCl salt
365;~Co3	371;~B3/B4a	376;~B3/B4a
 as a HCl salt	 as a HCl salt	
366;~B3/B4a	372;~B3/B4a	377;~B3a
 as a HCl salt		
367;~B3/B4a	92;=Co26	378;~B3/B4a
		
300;=Co27	115;~B6	379;~B7
	 as a HCl salt	
368;~Co3	373;~B15	380;~Co3














	 as a HCl salt	
369;~Co27	374;~Co18	381;~B3/B4a
		
370;~B9b	375;~Co26	382;~Co2/B3/B4a
		
132;~B3/B4a;NMR*	133;~B3a;NMR*	392;~B1/B2a/B3/B4a
		 as a HCl salt
108;~B3/B4a	386;~B3/B4a	393;~B1/B2a/B3/B4a
 as a HCl salt		
383;~B3/B4a	387;~B8	394;~B1/B2a

















		
384;~B3/B4a	389;~B6	395;~B3a
	 as a HCl salt	
385;~Co5c	390;~B3a	94;=Co28
		 as a HCl salt
39;=B30	114;=Co39	396;~Co2
 as a HCl salt		
40;=B30	391;~Co12	397;~B6
	 as a HCl salt	
398;~Co12	405;~B1/B2a/B3/B4a	411;~B13
















		 as a HCl salt
399;~B1/B2a	406;~B3/B4a	412;~Co12
		 as a HCl salt
400;~B1/B2a	124;~B6/Co18	413;~Co2
	 as a HCl salt	 as a HCl salt
401;~B1/B2a	407;~B5	414;~B3/B4a
	 as a HCl salt	
402;~Co5a	408;~B5	415;~Co5a
		
403;~B1/B2a	409;~Co18	416;~Co38
















	 as a HCl salt	
404;~B1/B2a/B3/B4a	410;~Co18	417;~Co38
		
418;~B1/B2a	425;~B3/B4a	430;~B3/B4a
		
419;~B3/B4a	426;~B3/B4a	125;~Co3
	 as a HCl salt	
420;~ B3/B4a	427;~ B3/B4a	431;~B3a/B3/B4a
		 as a HCl salt
421;~Co3	428;~Co3	432;~Co1
















		
422;~B1/B2a/B3/B4a	6;~B3a	433;~B15
		
423;~B1/B2a/B3/B4a	34;=B25	434;~Co24
 as a HCl salt		 as a HCl salt
424;~B1/B2a/B3/B4a	429;~B5	435;~Co1
	 as a HCl salt	
5;=B1a/B4a/b; NMR*	440;~Co1	447;~Co4
 as a HCl salt	 as a HCl salt	 as a HCl salt
436;~Co3	441;~B3/B4a	448;~B3/B4a



















 and  as a HCl salt		
95a and 95;=Co29	442;~B3/B4a	449;~B3/B4a
		 as a HCl salt
437;~B10/Co1	443;~B3/B4a	450;~B3a
 as a HCl salt		
438;~Co1	444;~B3/B4a	451;~B3/B4a
		 as a HCl salt
439;~B3/B4a	445;~Co4	452;~B3/B4a
















 as a HCl salt		
134;~B3/B4a;NMR*	446;~B3/B4a	461;~B6
	 as a HCl salt	
454;~Co29	453;~Co1	462;~Co5c
 as a HCl salt	 and  as a HCl salt	
455;~B3/B4a	44a and 44 ;=Co1	463;~Co5a
		
456;~B3/B4a	458;~Co1	464;~B3a
	 as a	
















	HCl salt	
96;=Co30	459;~B3a	465;~Co30
		
457;~B9a	460;~B1/B2a/B3/B4a	466;~B3/B4a
	 as a HCl salt	
467;~B3/B4a	474;~B9a	479;~B1
	 as a HCl salt	
468;~B1/B2a	135;~B3/B4a;NMR*	480;~Co30
 as a HCl salt		
469;~B3/B4a	475;~Co1	97;=Co31/~Co15
		
470;~B3/B4a	476;~B1/B2a	98;=Co31/~Co15;NMR*
















 as a HCl salt		
471;~B3/B4a	477;~Co1	481;~B6
		
472;~B3/B4a	478;~Co3	482;~Co3
 as a HCl salt	 as a HCl salt	 as a HCl salt
473;~B3/B4a	136;~B3/B4a;NMR*	483;~Co3
		
484;~B5	491;~B1/B2a	497;~B3/B4a
		 as a HCl salt
485;~B3/B4a	492;~B3/B4a	119;=Co41b
















		
486;~B5	493;~B3/B4a	498;~B5b-1
		
487;~B3/B4a	494;~Co3	99;=Co32;NMR*
		
488;~Co3	495;~B5	138;~B5b-1;NMR*
	 as a HCl salt	 as a HCl salt
489;~B3a	137;~Co1;NMR*;=B49	499;~Co5
 as a HCl salt		
490;~Co18	496;~B3/B4a	500;~B1/B2a
















		 as a HCl salt
501;~Co18	506;~Co19	512;~B3/B4a
		
502;~Co18	507;~Co19	513;~B3/B4a
		
503;~Co4	508;~B5	514;~B3/B4a
		
139;~Co18;;NMR*	509;~B3a	515;~B3/B4a
		
504;~B25	510;~B3/B4a	516;~B5
		
















140;~B3/B4a;NMR*	141;~B5/B1/B2a;NMR*	35;=B26
		
505;~B1/B2a	511;~B3/B4a;	517;~B3/B4a
 as a HCl salt		
518;~B3/B4a	142;~Co18;NMR*	529;~B3a
		
519;~B3/B4a	524;~Co3	530;~B3/B4a
	 as a HCl salt	
102;=Co34	525;~Co18	10;=B5a
	 as a HCl salt	 as a HCl salt
520;~B1/B2a	526;~Co18	531;~B3/B4a

















 as a HCl salt		
521;~B3/B4a	527;~Co2	36;=B27
		
522;~B3/B4a	143;~B3/B4a;NMR*	532;~B3/B4a
		
523;~Co18	528;~B3/B4a	533;~B3a
 as a HCl salt		
534;~B3a	104;~A5	545;~B5
 as a HCl salt		
535;~B3/B4a	603;~B3/B4a	41;= B31
















		
536;~Co19	540;~Co18	33;=B32
		
537;~Co19	541;~B3/B4a	43;=B32
		 a s a HCl salt
538;~B5	542;~B5	546;~B3/B4a
		
144;~B3/B4a;NMR*	543;~B3/B4a	547;~B3/B4a
		 a s a HCl salt
539;~B5	544;~Co4	548;~B3/B4a



















		 as a HCl salt
549;~B5	47;~B3/B4a	145;~Co18; NMR*
		
550;~B6	556;~Co3	129;~Co1
		
551;~B3/B4a	46;~Co3	111;~Co5a
		
552;~B5	557;~B5a	560;~B1/B2a
 as a HCl salt		
553;~B3a	558;~Co3	561;~B3/B4a



















 as a HCl salt	 as a HCl salt	
554;~B3/B4a	103 ;=Co35	562;~B3/B4a
	 as a HCl salt	 as a HCl salt
555;~B3/B4a	559;~Co18	563;=Co2B
	 as a HCl salt	 as a HCl salt
564;~B3/B4a	110;=Co37	578;~B3/B4a
	 as a HCl salt	
565;~B1/B2a	570;~B3/B4a	579;~B1/B2a
 as a HCl salt		
566;~B3/B4a	571;~B5	580;=B3B



















		 as a HCl salt
567;~B3/B4a	574;~B1/B2a	581;~B3a
		
568;~B5	575;~B1/B2a	582;~B3/B4a
		
569;~B5	576;~B5	583;~B3/B4a
		
146;~B5;NMR*	577;~B3/B4a	584;~B5a
 .oxalate		
585;~B3/B4a	589;~Co16	100;~B5 b-3



















 as a HCl salt		
586;~B3/B4a	590;~B3	101;=Co33
 as a HCl salt		 and  as a HCl salt
105;=Co35a	591;~B3/B4a	14 and 14a ;~B6
 as a HCl salt	 as a HCl salt	
147;~B3/B4a;NMR*	592;~Co1	594;~B5
		
587;~B5b-1	593;~B6	595;~Co5
		



















588;~Co16	148;~B6;NMR*	596;~Co3
		
120;=Co42	149;~B6;NMR*	610;=Co46
		
15;~B7	118;=Co41a	611=Co46
		
597;~Co2	126;~A5	697;~B33
		
598;~B10	51;~Co5a	698;~Co46
		 as a HCl salt
599;~B5	604;=B33	699;~Co46



















		
600;~B5	608;=Co46	700;~Co27
		
601;~B5	609;=Co46	613;=Co47b
		
701;~B3	705;~B53	712;~B1
		 .trifluoroacetate
702;~Co46	706;~B53	616;=B38
		
703;~Co46	707;~B53	713;~B4
	 as a HCl salt	 as a HCl salt
645;=B53	647;B55	619;=B41



















	 as a HCl salt	
704;~B4b	708;~B55	714;~Co27
	 as a HCl salt	
615;=B37	709;~B38	690;~Co27
		
646;B54	710;~Co14	721;~B4
		
648;=Co58	711;~B39	722;~B3
		
715;~B2	620;=B42a	731;~Co27
 as a HCl salt		



















716;~B3	725;~B1	732;~Co49
		
717;~B4a	726;~B2	733;~B4b
		 as a HCl salt
718;~B4a	727;~B44	734;~B4b
		
719;~B4a	728;~B4b	624;~B11
	 as a HCl salt	
720;~B1	621;=B42b	735;~B5
		
723;~B43	729;~B3	736;~B43
	 as a trifluoroacetate salt	



















724;~B3	730;~Co1	625;~B1
		 as a HCl salt
737;=Co48	651;=Co59	744;~B3
		
652;=Co60	617;=B39	745;~Co27
		
650;~B3(1st altern. protocol)	741;~Co27	746;~B1
		
738;~Co27	742;~Co27	747;~Co27
		
739;~Co27	618;=B40	689;~Co27
		
740;~Co27	743;~B5	748;~B1



















		
749;~B43	753;~Co27	760;~Co27
		 as a HCl salt
622;=B43	754;~Co49b	623;=B44
		
688;~Co60	755;~B1	761;~B4
		
750;~Co27	756;~B3	762;~Co27
		
655;;=B56	757; =co76	763;~B65
		 as a HCl salt
751;~Co49a	758;~Co49b	764;~B44
















		
626;=Co49a	653;~Co60	765;~Co27
		
752;~Co27	759;~Co27	766;~Co27
	 as a HCl salt	
767;~Co27	657;=Co61	627;=Co49b
 as a HCl salt		
768;~B3	658;=B57	775;~B2
		
769;~B9/B29	771;~Co27	776;~B29
		
770;~B44	772;~Co27	777;~Co27


















 as a HCl salt		
644;=B51	773;~B44	778;~Co50
 as a HCl salt		
656;=Co61	774;~B4	779;~Co50
		
628;=Co50	785;~Co27	793;~Co27
		 as a HCl salt
629;=Co50	786;~Co27	631;=Co51
		
780;~Co31	787; ~Co27	794;~Co49a
		



















781;~Co31	788;~Co27	795;~Co27
 as a HCl salt		
782;~B51	789;~Co27	796;~Co27
		
634;~Co27	790;=B45	797;~A4
		
783;~B29	791;~Co27	798;~Co49a
		
784;~Co27	792;~Co27	659;=B58
	 as a HCl salt	
662;~Co53	803;~B6	801;~B5b
		 as a
















		HCl salt
799;~Co54	636;=Co54	802;~B6
		
661;=B59	637;=Co54	807;~Co54
		
660;=B59	804;~Co27	808;~Co54
		
633;=Co52	805;~Co27	809;~Co27
		
800;~Co27	806;~Co27	810;~Co53
		
635;=Co53	811;~Co53	813;~B1
		
638;=B47	812;~B5	814;~Co27















		
815;~B5	632;=B46	828;~B5b
		
816;~Co27	821;~B4a	829;~Co27
		
817~Co27	822;~B29	830;~Co14
		
640;=Co55	823;~Co14	663;=Co62
		
818;~B2	824;~Co14	664;=B60
		
819;~Co54	825;~Co55	831;~Co27
















		
687;~B3	826;~Co27	832;~Co27
		
820;~Co27	827;~Co27	833;~B5a
		
834;~Co14	840;~Co57	607;=B35
		
835;~B5a	841;~B48	606;=Co45
		
837;~B48	842~B5a	605;=B34
















		
641;=B48	665;=B3C	612;=Co47a
	 as a HCl salt	
642;=Co56	643;=Co57	630;=B45
		
838;~B48	836;~B48	666;=B3C
		
839;~B5a		844;~B3C
		 as a HCl salt
845;~Co27	851;~Co26	696;=Co 72
 . Lithium (+1)		
846;=B64	669;=B62	856;~B41
















		 as a HCl salt
847;~B2	852;~Co58	857;~A15
		
667;=B61	853;~Co58	671;=Co 64
		
848;~Co58	670;=Co 63	858;~B3
		
849;~B3	854;~B5	673;=Co 65
		
850;~B61	855;~B5	676;=Co 66c)
		












859;~B61	682;=Co 69	867;~Co61
		
860;~B61	863;~B59	868;~Co61
		 as a HCl salt
678;=Co 67	864;~B59	694=B4C
		 .oxalate
680;=Co 68	865;=Co35	869;~B3
 as a HCl salt		 a s a HCl salt
861;~Co3	866;~Co35	692;=Co2B
		
691;=B14A	679;=B4B	675;=Co 66b)

 as a HCl salt	 as a HCl salt	
862;~B3	693;=B63	870;~B5
 as a HCl salt		 as a HCl salt
685;=Co 71		875;~Co61
		
871;~B6	877;~Co64	672;=Co 64
		
872;~B4a	878;~Co63	674;=Co 66a)
	 as a HCl salt	
677;=B3D	879;~B4B	681;=B5a

		
873;~B3	880;~A2	683;=Co 70
 as a HCl salt		
874;~Co61	668;=B61	686;=B5a
		
695;~B6	882;~B5	884;~B5
 as a HCl salt	 as a HCl salt	
881;~B5	883;~B5	885;~B5a
 as a HCl salt		
886;~B3E	887;~B3	888;~B3C

		 as a HCl salt
889;~Co15	890;~B5b1	891;~B3F
 as a HCl salt		 as a HCl salt
892;~B59A	893;~B5b1	894;~B3F
		
895;~B3	896;=Co46	897;=Co46
		
898;~B5b1	899;~Co47	900;~Co11
	 .HBr	
901;~Co11	902;~Co73	903;~B4A

		
904;~B4A	905;~B5b1	906;~Co74
		
907;~Co47	908;~Co47	909;~Co47
		
910;~Co47	911;~Co47	912;~B1
	 as a oxalate salt	
913;~Co47	914;~Co49a	915;~B1
		
916;~B57	917;=Co47	918;~Co75

		
919;~B47A	920;~B27A	921;~B5a
		
922;~B5a	923;=B3E	924;=B3F
		 as a HCl salt
925;=B3F	926;=B59A	927;~B3
	 as a HCl salt	
928;~B1	4a;=B3	

Analytical Part

LC/GC/NMR

General procedure A

The HPLC measurement was performed using an Alliance HT 2790 (Waters) system comprising a quaternary pump with degasser, an autosampler, a column oven (set at 40 °C, unless otherwise indicated), a diode-array detector (DAD) and a column as specified

in the respective methods below. Flow from the column was split to a MS spectrometer. The MS detector was configured with an electrospray ionization source. Mass spectra were acquired by scanning from 100 to 1000 in 1 second using a dwell time of 0.1 second. The capillary needle voltage was 3 kV and the source temperature was maintained at 140 °C. Nitrogen was used as the nebulizer gas. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

Method 1

In addition to the general procedure A: Reversed phase HPLC was carried out on an Xterra MS C18 column (3.5 µm, 4.6 x 100 mm) with a flow rate of 1.6 ml/min. Three mobile phases (mobile phase A: 95% 25 mM ammoniumacetate + 5 % acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 1 % A, 49 % B and 50 % C in 6.5 minutes, to 1 % A and 99 % B in 1 minute and hold these conditions for 1 minute and reequilibrate with 100 % A for 1.5 minutes. An injection volume of 10 µl was used. Cone voltage was 10 V for positive ionization mode and 20 V for negative ionization mode.

Method 2

In addition to the general procedure A: Column heater was set at 45 °C. Reversed phase HPLC was carried out on an Atlantis C18 column (3.5 µm, 4.6 x 100 mm) with a flow rate of 1.6 ml/min. Two mobile phases (mobile phase A: 70 % methanol + 30 % H_2O; mobile phase B: 0.1 % formic acid in H_2O/methanol 95/5) were employed to run a gradient condition from 100 % B to 5 % B + 95 % A in 9 minutes and hold these conditions for 3 minutes. An injection volume of 10 µl was used.
Cone voltage was 10 V for positive ionization mode and 20 V for negative ionization mode.

Method 3

In addition to the general procedure A: Reversed phase HPLC was carried out on an Xterra MS C18 column (3.5 µm, 4.6 x 100 mm) with a flow rate of 1.6 ml/min. Three mobile phases (mobile phase A: 95% 25 mM ammoniumacetate + 5 % acetonitrile; mobile phase B: acetonitrile; mobile phase C: methanol) were employed to run a gradient condition from 100 % A to 50 % B and 50 % C in 6.5 minutes, to 100 % B in 1

minute, 100 % B for 1 minute and reequilibrate with 100 % A for 1.5 minutes. An injection volume of 10 µl was used.
Cone voltage was 10 V for positive ionization mode and 20 V for negative ionization mode.

Method 9

In addition to the general procedure A: Reversed phase HPLC was carried out on a Waters Xterra-RP C18 column (3.5 µm, 4.6 x 100 mm) with a flow rate of 0.8 ml/min. Two mobile phases (mobile phase A: 100 % 7 mM ammonium acetate; mobile phase B: 100 % acetonitrile) were employed to run a gradient condition from 80 % A and 20 % B (hold for 0.5 minute) to 90 % B in 4.5 minutes, 90 % B for 4 minutes and reequilibrated with initial conditions for 3 minutes. An injection volume of 5 ml was used. Cone voltage was 20 V for positive and negative ionization mode. Mass spectra were acquired by scanning from 100 to 1000 in 0.4 seconds using an interscan delay of 0.3 seconds.

Method 10

In addition to the general procedure A: Reversed phase HPLC was carried out on a Xterra-MS C18 column (3.5 µm, 4.6 x 100 mm) with a flow rate of 0.8 ml/min. Two mobile phases (mobile phase A: 100 % 7 mM ammonium acetate; mobile phase B: 100 % acetonitrile; were employed to run a gradient condition from 80 % A , 20 % B (hold for 0.5 minute) to 10 % A, 90 % B in 4.5 minutes, hold at 10 % A and 90 % B for 4 minutes and reequilibrated with initial conditions for 3 minutes. An injection volume of 10 ml was used. Cone voltage was 20 V for positive and negative ionization mode. Mass spectra were acquired by scanning from 100 to 1000 in 0.4 seconds using an interscan delay of 0.3 seconds

General procedure B

The LC measurement was performed using an Acquity UPLC (Waters) system comprising a binary pump, a sample organizer, a column heater (set at 55 °C), a diode-array detector (DAD) and a column as specified in the respective methods below. Flow from the column was split to a MS spectrometer. The MS detector was configured with an electrospray ionization source. Mass spectra were acquired by scanning from 100 to 1000 in 0.18 seconds using a dwell time of 0.02 seconds. The capillary needle voltage was 3.5 kV and the source temperature was maintained at 140 °C. Nitrogen was used as the nebulizer gas. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

Method 4

In addition to the general procedure B: Reversed phase UPLC (Ultra Performance Liquid Chromatography) was carried out on a bridged ethylsiloxane/silica hybrid (BEH) C18 column (1.7 µm, 2.1 x 50 mm; Waters Acquity) with a flow rate of 0.8 ml/min. Two mobile phases (mobile phase A: 0.1 % formic acid in H_2O/methanol 95/5; mobile phase B: methanol) were used to run a gradient condition from 95 % A and 5 % B to 5 % A and 95 % B in 1.3 minutes and hold for 0.2 minutes. An injection volume of 0.5 µl was used. Cone voltage was 10 V for positive ionization mode and 20 V for negative ionization mode.

Method 5

In addition to the general procedure B: Reversed phase UPLC (Ultra Performance Liquid Chromatography) was carried out on a bridged ethylsiloxane/silica hybrid (BEH) C18 column (1.7 µm, 2.1 x 50 mm; Waters Acquity) with a flow rate of 0.8 ml/min. Two mobile phases (25 mM ammonium acetate in H_2O/acetonitrile 95/5; mobile phase B: acetonitrile) were used to run a gradient condition from 95 % A and 5 % B to 5 % A and 95 % B in 1.3 minutes and hold for 0.3 minutes. An injection volume of 0.5 µl was used. Cone voltage was 30 V for positive ionization mode and 30 V for negative ionization mode.

General procedure C

The LC measurement was performed using a UPLC (Ultra Performance Liquid Chromatography) Acquity (Waters) system comprising a binary pump with degasser, an

autosampler, a diode-array detector (DAD) and a column as specified in the respective methods below, the column is hold at a temperature of 40°C. Flow from the column was brought to a MS detector. The MS detector was configured with an electrospray ionization source. The capillary needle voltage was 3 kV and the source temperature was maintained at 130 °C on the Quattro (triple quadrupole mass spectrometer from Waters). Nitrogen was used as the nebulizer gas. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

Method 6

In addition to the general procedure C: Reversed phase UPLC was carried out on a Waters Acquity BEH (bridged ethylsiloxane/silica hybrid) C18 column (1.7 µm, 2.1 x 100 mm) with a flow rate of 0.35 ml/min. Two mobile phases (mobile phase A: 95 % 7 mM ammonium acetate / 5 % acetonitrile; mobile phase B: 100 % acetonitrile) were employed to run a gradient condition from 90 % A and 10 % B (hold for 0.5 minutes) to 8 % A and 92 % B in 3.5 minutes, hold for 2 min and back to the initial conditions in 0.5 min, hold for 1.5 minutes. An injection volume of 2 µl was used. Cone voltage was 20 V for positive and negative ionization mode. Mass spectra were acquired by scanning from 100 to 1000 in 0.2 seconds using an interscan delay of 0.1 seconds.

Method 7

In addition to the general procedure C: Reversed phase UPLC was carried out on a Waters Acquity BEH (bridged ethylsiloxane/silica hybrid) C18 column (1.7 µm, 2.1 x 100 mm) with a flow rate of 0.343 ml/min. Two mobile phases (mobile phase A: 95 % 7 mM ammonium acetate / 5 % acetonitrile; mobile phase B: 100 % acetonitrile) were employed to run a gradient condition from 84.2 % A and 15.8 % B (hold for 0.49 minutes) to 10.5 % A and 89.5 % B in 2.18 minutes, hold for 1.94 min and back to the initial conditions in 0.73 min, hold for 0.73 minutes. An injection volume of 2 ml was used. Cone voltage was 20V for positive and negative ionization mode. Mass spectra were acquired by scanning from 100 to 1000 in 0.2 seconds using an interscan delay of 0.1 seconds.

General procedure D

The HPLC measurement was performed using an Alliance HT 2795 (Waters) system comprising a quaternary pump with degasser, an autosampler, a diode-array detector

(DAD) and a column as specified in the respective methods below, the column is hold at a temperature of 30°C. Flow from the column was split to a MS spectrometer. The MS detector was configured with an electrospray ionization source. The capillary needle voltage was 3 kV and the source temperature was maintained at 100 °C on the LCT (Time of Flight Zspray mass spectrometer from Waters). Nitrogen was used as the nebulizer gas. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

Method 8

In addition to the general procedure D: Reversed phase HPLC was carried out on a Supelco Ascentis Express C18 column (2.7 µm, 3.0 x 50 mm) with a flow rate of 0.7 ml/min. Two mobile phases (mobile phase A: 100 % 7 mM ammonium acetate; mobile phase B: 100 % acetonitrile) were employed to run a gradient condition from 80 % A and 20 % B (hold for 0.5 minute) to 5% A and 95 % B in 2.5 minutes, hold for 4.5 minutes and back to the initial conditions in 1.5 minutes and hold for 1 min. An injection volume of 5 l was used. Cone voltage was 20 V for positive and negative ionization mode. Mass spectra were acquired by scanning from 100 to 1000 in 0.4 seconds using an interscan delay of 0.3 seconds.

Table A1: physico-chemical data

If physio-chemical data were generated multiple times for a compound then all data is listed

Comp No.	Melting Point (°C)	Kofler (K) or DSC	HPLC Rt (min)	MS M+ (H^+)	LC/ GC/ MS method
202			1.39 1.25	416 416	4 5
17	159	K	3.83	402	6
203	110	K	4.66	458	6
206	101.91	DSC	4.15	460	6
207	126	K	3.41	459	6

Comp No.	Melting Point (°C)	Kofler (K) or DSC	HPLC Rt (min)	MS M+ (H^+)	LC/ GC/ MS method
538			4.04	390	6
144	225.35 222	DSC K	2.96	488	6
539			0.64	473	5
603			1.01		5
540	130.63	DSC	3.19	505	6

	126 126	K K			
56	147.56 141.67 144	DSC DSC K	3.73 3.73	446 446	6 6
208	158	K	4.57	444	6
209	166.2	DSC	4.14	422	6
210	176.79 176	DSC K	3.89	434	6
211	187.05	DSC	3.79	446	6
1	152.97 152.4 154	DSC DSC K	3.12 3.12	406 406	6 6
212	145.81	DSC	3.8	487	6
213	170.82	DSC	3.7	473	6
214	207.88 209	DSC K	4.29	480	6
215	190	K	3.8	416	6
216	162.91	DSC	3.54	556	6
22	108.61	DSC	4.52	414	6
60	90	K	3.55	485	6
122	202.94	DSC	3.42	415	6
58	165 163.09	K DSC	4.53	509	6
3	168 145	K K	3.22 3.23	420 420	6 6
218	204.29	DSC	3.59	448	6
219	215.88	DSC	3.77	535	6
220	94.96	DSC	3.67	390	6
221	123.95	DSC	4.02	416	6
222	142	K	4.23	404	6
223			3.69	501	6

541	162.74	DSC	2.99	530	6
542	147.16 155	DSC K	3.85	487	6
543	149.47	DSC	3.04	517	6
544			3.43	542	6
41	168 169.24	K DSC	2.86	437	6
33	83	K	3.02	501	6
43	107	K	3.08	501	6
546	135	K	3.13	501	6
547	217	K	3.22	473	6
548	135	K	3.13	501	6
551			1.02	515	5
552			0.87 0.63	500 500	4 5
553			3.22	489	6
554			3.48	491	6
556			2.72	396	6
557	189	K	3.42	470	6
558	94	K	3	449	6
103	140	K	3.94	512	6
559	98.59	DSC	2.79	477	6
145	94.44	DSC	2.8	477	6
129			1.1 1.13	495 495	4 5
111			1.27 1.33	593 593	5 4

224	128.96	DSC	3.6	427	6
62	220.78	DSC	3.64	513	6
23	134	K	3.33	487	6
226	164.73	DSC	4.04	501	6
8	181 185 179.61	K K DSC	2.7 2.75	419 419	6 6
84	from: 133.17 to 139.07	DSC	from 2.65 to 2.67	405	6
227	165.51	DSC	3.87	508	6
65			3.7	448	6
228	132	K	4.31	400	6
229	123.16	DSC	4.01	529	6
73			3.5	499	6
231	106.81	DSC	3.72	565	6
59	200	K	4.08	563	6
52	159.08	DSC	3.99	474	6
232			3.84	381	6
64			2.31	420	6
233	183.6	DSC	2.87	436	6
234	130	K	3.44	445	6
130	218	K DSC	3.54 2.76	586 586	6 7
235	154.8 154	DSC K	3.37	515	6
69	260.14 >260 263.87	DSC K DSC	2.8 2.84	419 419	6 6

			1.27 1.27 1.28	593 593 593	5 5 5
560			0.74	547	5
561	219	K	3.19	475	6
562			3.8	487	6
563	193 200.75	K DSC	3.2	498	6
564			2.99	567	6
565	152	K	3.34	408	6
566	183	K	3.09	484	6
567			3.1	484	6
568	164.5 167	DSC K	3.4	453	6
569	139 139.45	K DSC	3.38	453	6
146	163 158.48	K DSC	3.36	454	6
110			0.78	503	5
570			0.88	489	5
575			0.71	475	5
576			1.36	579	5
577			3.57	487	6
578			3.5	485	6
581			3.17	501	6
582	127	K	4.03	489	6
583	136.49	DSC	3	599	6
584	140	K	4.07	531	6

236			1.28	438	5
			1.28	438	5
237	80	K	4.44	420	6
238	133.66	DSC	2.68	449	6
66			3.95	411	6
2	169.26	DSC	2.94	392	6
	169.13	DSC	2.94	392	6
113	177	K	3.27	420	6
68	165	K	2.86	415	6
239	160	K	3.48	514	6
240	132	K	3.55	529	6
	132.96	DSC			
241	167.79	DSC	2.97	473	6
	149.64	DSC	2.96	473	6
55			3.9	543	6
242	196.25	DSC	3.53	429	6
243	146.13	DSC	3.52	558	6
244	155	K	4	499	6
245	70	K	3.37	487	6
246	149.27	DSC	3.78	531	6
24	80	K	3.44	446	6
249			2.63	477	6
250	104.64	DSC	4.13	545	6
251	127.97	DSC	3.7	556	6
21	204.45	DSC	2.86	463	6
	205.05	DSC	2.86	463	6
72	120	K	3.67	513	6
253	167.45	DSC	3.5	503	6
254	173.43	DSC	3.77	517	6
74	253	K	2.97	433	6
70a	163.31	DSC	2.86	459	6
256	171	K	3.69	457	6
	173.42	DSC			
26	202.6	DSC	3.04	406	6
257	140.28	DSC	4.37	492	6
75	186.76	DSC	2.87	436	6

585			2.87	488	6
586			3.82	471	6
105	135	K	3.94	512	6
147			4.25	529	6
592	194	K	2.89	421	6
593	168	K	4.01	502	6
148	149.13	DSC	3.67	474	6
149	146.61	DSC	3.67	474	6
100	73	K	3.63	549	6
			3.63	549	6
101	186	K	3.16	456	6
14 + 14a	110	K	3.62	478	6
594	214.83	DSC			
595	105	K	3.13	470	8
15	114.76	DSC	3.62	487	6
604	157	DSC	3.16	489	
608	159	DSC	2.66	419	8
609	155	DSC	2.65	419	8
610	150	DSC	2.66	419	8
611	148	DSC	2.68	419	8
697	171	DSC	2.97	461	6
698	111	DSC	3.03	461	6
699	150	DSC	2.99	461	6
700	133	DSC	2.85	477	7
613			1.26	402	4
701	68	DSC	3.18	503	6
702	129	DSC	3.89	501	6
703	80	K	3.92	501	6
645	188	DSC	2.44	433	6
704	151	DSC	3.74	487	6
615	134	DSC	2.62	519	6

	186.1	DSC	2.87	436	6
			2.87	436	6
7			3.31	463	6
259	163.47	DSC	3.14	445	6
	156	K			
260	188.74	DSC	3.03	498	6
27	176.76	DSC	3.94	385	6
78	162.1	DSC	3.38	434	6
	161.7	DSC	3.38	434	6
	165.99	DSC	3.17	434	8
261	180.29	DSC	2.79	445	6
	183	K			
262	130.64	DSC	3.21	483	6
25	245.28	DSC	2.96	445	6
	259.39	DSC	2.96	445	6
263	168.45	DSC	3.74	473	6
264			3.14	459	6
265	190	K	2.51	476	6
	186.88	DSC			
	186.88	DSC			
266	157.15	DSC	2.52	476	6
76	158.57	DSC	3.5	418	6
267	160.27	DSC	4.02	552	6
268	158.2	DSC	3.45	512	6
	177.91	DSC	3.46	512	6
269	155.79	DSC	4.46	444	6
271	181.91	DSC	3.82	523	6
272	178.46	DSC	2.77	433	6
	176.95	DSC	2.78	433	6
	180	K			
273			2.73	540	6
274	176.3	DSC	4.45	456	6
4	from		From	447	6
	134	DSC	2.95		
		or	to		
	to 142	K	2.99		
276	188.62	DSC	3.12	512	6

646	125	K	2.51	392	6
648	196	DSC	3.42	515	6
705	135	DSC	3.3	487	6
706	182	K	2.58	392	6
707	186	K	2.23	391	6
647	133	K	2.3	419	6
708	152	K	3.1	473	6
709	175	K	3.25	544	6
710	206	DSC	3.24	461	6
711	166	DSC	3.15	426	6
712	124	DSC	3.02	376	6
616	170	K	2.42	246	6
713	127	K	3.83	507	6
619	210	K	3.07	505	6
714	128	K	3.62	477	6
690	138	K	3.63	477	6
716	247	DSC	2.81	417	6
717	125	DSC	3.59	483	6
718	103	DSC	3.66	501	6
719	119	DSC	4.59	509	6
720	175	DSC			
721			0.67	528	4

28	142.35	DSC	2.98	450	6
37	154.05	DSC	3.55	476	6
278			2.66	511	6
279	193.59 199	DSC K	2.58	405	6
281			2.55	503	6
283	128.64	DSC	2.87	487	6
79a + 79			2.81	489	6
602	158.84	DSC	4.07	509	6
80	144	K	3.33	433	6
81	199	K	3.32	433	6
285	183.33	DSC	2.8 2.83	459 459	6 6
286	114.64 114	DSC K	3.24	473	6
287	156 156.77	K DSC	3.49	445	6
29	80	K	3.03	459	6
288	142	K	2.37	449	6
291			2.85	489	6
293			3.96	390	6
294	184	K	2.52 2.54	391 391	6 6
16	80	K	3.21	432	6
20	203.41	DSC	3.53	529	6
295	155.31	DSC	2.91	565	6
296	126.47	DSC	3.03	503	6
54	138	K	3.93	342	6
297	140	K	2.45	434	6
298			2.41	489	6
299	201	DSC	3.41	475	6
49	168.25	DSC	2.85	490	7
93	175.51 176.71	DSC DSC	2.72 2.62	419 419	6 6
302	222.79	DSC	2.77	503	6
303	194.82	DSC	2.8	474	6

722	138	K	3.57	455	6
724	176	DSC	3.07	433	6
620	114	K	3.89	587	6
725	227	DSC	3.11	474	6
727	109	DSC	3.01	461	6
728	109	DSC	3.77	497	6
621	>260	K	2.7	486	6
729	183	DSC	2.99	515	6
730	139	K	3.4	501	6
731	237	DSC	339	478	6
732	160	K	3.46	458	6
733	70	K			
734	178	K	3.97	507	6
624	147	K	2.65	475	6
735	143	DSC	3.56	408	6
736	68	K	2.39	450	7
625	104	K	3.06	575	7
737	156	DSC	2.28	489	7
652	142	DSC	2.84	507	7
650	66	K	2.98	616	7
738	197	DSC	2.99	502	7
739	179	DSC	2.99	502	7
740	167	DSC	2.97	483	7
651	122	K	2.05	516	7
617	149	DSC	2.44	416	7
741	85	K	3.34	506	7
742	115	K	3.2	507	7
618	139	DSC	2.3	457	7
743	135	DSC	2.83	481	7
744	200	K	2.23	530	7

304	131.93 130	DSC K	2.54	449	6
305	117.44	DSC	4.26	500	6
61	103.14	DSC	3.57	459	6
306	185 182.27	K DSC	3.07	477	6
307	132 168.8	K DSC	3.33	473	6
308	192	K	2.5	435	6
311	262.42	DSC	3.07	429	6
312	184.29	DSC	2.42	477	6
313	176.49 176.31	DSC DSC	2.95 2.96	477 477	6 6
314	201.43 201	DSC K	2.78	463	6
30	80	K	2.72	450	6
38	168	K	3.63	400	6
315	139	K	2.98	424	6
48	189.27	DSC	3.27	497	6
316	158.73	DSC	3.51	526	6
320	193.76	DSC			
321	265.41	DSC	2.8	474	6
57			3.29	538	6
323	166.37	DSC	4.19	478	6
85		DSC	4.16	448	6
324	144.1	DSC	3.6 3.6	473 473	6 6
325			2.76	516	6
86	195.95	DSC	2.52	458	6
326			3.07	438	6
83	141.79	DSC	2.89	530	6
327			3.86	434	6
329			2.87	488	6
330	139.15	DSC	2.96	502	6
332	141.98	DSC	2.92	475	6
333	150	K	2.56	503	6
334	125	K	2.75	513	6

745	110	K	3.36	476	7
746	62	K	2.44	464	7
747	68	K	2.67	507	7
689	203	DSC	2.77	492	7
749	86	K	2.64	448	7
622	190	K	2.57	434	7
688	175	DSC	6.04	508	7
750	148	DSC	2.61	493	7
655			8.11	506	10
751			2.99	585	7
752	169	K	3.01	502	7
753	155	DSC	3.01	491	7
754	167	K	2.19	499	7
755	152	K	2.72	434	7
756	118	K	2.23	461	7
757	126	K	2.64	461	7
758			2.22	485	7
653	146	DSC	311	511	7
759	180	K	2.85	502	7
760	221	K	2.51	507	7
623	133	K	2.43	475	7
761	119	K	3.2	515	7
762	137	K	3.18	507	7
763	176	K	2.27	463	7
764	126	K	2.56	475	7
765	199	DSC	2.65	492	7
766	125	DSC	2.93	507	7
767	220	K	2.9	527	7
768	158	DSC	2.79	515	7
770	130	K	2.27	477	7
644	96	K	2.06	603	7

335	122.79	DSC	3.34	497	6
336	60	K	3.9 3.9	442 442	6 6
338	198.91	DSC	2.6	405	6
339	187.62 80	DSC K	2.3	530	6
32			2.98	445	6
131	183.2	DSC	3.14	476	6
340	184.2	DSC	3.99	522	6
342	228.85	DSC	3.7	487	6
343	80 (199)	K	3.96	545	6 (6)
344	161.91	DSC	3.06	477	6 (6)
31	244 250.38 254	K DSC K	2.93 2.93	459 459	6 6
345	115.62	DSC	3.65	501	6
346	131.39	DSC	4.99	608	6
347	135	K	3.1	487	6
348	169.44	DSC	3.8	503	6
350			0.83	562	5
351	110	K	2.89	516	6
89	126.66	DSC	3.33	446	6
352			2.75	530	6
87	125	K	3.63	491	6
353	168	K	3.46	456	6
354	70	K	4.05	458	6
355	184 184.23	K DSC	4.44	486	6
356	154.94	DSC	3.71	460	6
53			4.19	356	6
91			3.44	529	6
18	132	K	3.8	386	6
357	129.47	DSC	3.53	503	6
358	114.12 110	DSC K	4.37	486	6

656	95	K	2.83	478	7
657	110	K	2.83	478	7
658	152	DSC	3.19	552	7
771	93	K	2.56	492	7
772	154	DSC	2.84	493	7
773			2.3	504	7
774	80	K	3	513	7
627			2.2	471	7
775	133	K	2.51	432	7
776			6.62	428	9
777	119	DSC	3.11	535	7
778	148	K	3.13	516	7
779	148	K	3.13	516	7
628	122	K	2.84	492	7
629	124	K	2.84	492	7
780	188	K	3.12	502	7
781	186	K	3.12	502	7
782	170	K	4.31	617	7
634	80	K	621	491	7
783	220	K	6.05	468	7
784	163	DSC	2.61	493	7
785	190	K	2.83	575	7
786	125	DSC	2.79	492	7
787	181	DSC	2.95	503	7
788	146	K	2.57	507	7
789	204	K	2.49	520	7
790	67	K	3.47	583	7
791	130	K	2.94	576	7
792	153	DSC	2.82	478	7

359	114.97 126	DSC K	4.34	486	6
360	133.89 130	DSC K	4.15	452	6
361	188	K	2.87	473	6
362	80	K	3.77	501	6
363	172.65 175.38	DSC DSC	2.69 2.68	511 511	6 6
364			2.94 2.91	488 488	6 6
365	164.92	DSC	2.83	433	6
366			3.21	501	6
367			3.17	487	6
300	193.25 191.77 191.77	DSC DSC DSC	3.4 3.4 3.4	480 480 480	6 6 6
368			2.63	497	6
369			3.4	480	6
370	94	K	3.51	459	6
372	176.98	DSC	3.3	556	6
92	148.61 138.58 148.29	DSC DSC DSC	3.67 3.6 3.6	515 515 515	6 6 6
373	165	K	2.34	435	6
374	206.48	DSC	3.65	525	6
375	188.66 229.66	DSC DSC	3.61 3.61	501 501	6 6
376	214.67	DSC	3.5	586	6
377	112.31	DSC	3.89	495	6
378			3.25 3.26	531 531	6 6
379	80 146.1	K DSC	2.8 2.83	488 488	6 6
380	70	K	2.54	449	6
381	132.63	DSC	3.98	512	6
382	118 113	K K	2.91 2.94	461 461	6 6

793	161	DSC	3	508	7
631	194	DSC	2.39	483	7
794			2.6	500	7
795	182	K	2.95	507	7
796	189	DSC	3.04	508	7
797			3.18	443	8
798			2.62	486	7
659	106	DSC	3.05	484	7
662	122	K	3.07	512	7
799	114	DSC	3.05	510	7
661	154	DSC	2.47	518	7
660	151	DSC	2.47	518	7
633	130	DSC	2.69	477	7
800	134	DSC	3.05	512	7
801	127	DSC	3.21	514	7
802	80	K	2.19	493	7
803	80	K	2.19	493	7
636	80	K	3.35	493	7
637	>260	K	3.07	493	7
804	195	DSC	3.1	522	7
805	156	DSC	2.91	535	7
806	182	K	3.19	536	7
807	96	K	3.01	509	7
808	162	DSC	3.25	509	7
809	150	DSC	2.93	506	7

	109.12 112.07 104.44	DSC DSC DSC	2.95	461	6
132	132.47 130.95 138	DSC DSC K	3.31 3.31	487 487	6 6
383	120	DSC	2.92 2.91	503 503	6 6
384	80	K	2.99	489	6
385	119.77	DSC	4.93	488	6
39			4.33	501	6
40			4.66	501	6
133	126.47 125.21	DSC DSC	3.55 3.55	484 484	6 6
386	80	K	2.98	544	6
387	162.58	DSC	2.45	466	6
389	156.45 160	DSC K	3.75	448	6
390			3.7	498	6
391	190	K	2.63	475	6
393	193.78	DSC	3.22	410	6
395	118.99	DSC	2.83	475	6
94	115.24	DSC	3.1 3.11	443 443	6 6
396	156	K	3.14	487	6
397	132.82	DSC	3.24	450	6
398			2.96	473	6
399			3.12	409	6
400			3.12	409	6
401	138	K	3.47	408	6
405	191.73	DSC	3.01	406	6
407	146	K	4.37	412	6
408	130	K	4.1	444	6
410			2.83	477	6
411			8.92 5.93	458 458	2 1
412	180	K	2.85	489	6

810	131	DSC	2.76	482	7
635	111	DSC	3.11	495	7
811	127	DSC	3.08	511	7
812	170	DSC	2.65	454	7
813	166	K	2.4	442	7
814	160	DSC	2.98	495	7
815	123	DSC	3.24	483	7
816	176	DSC	3.12	555	7
817	155	DSC	3.01	491	7
640	241	DSC	2.71	506	7
819	80	K	2.96	480	7
687	162	DSC	2.23	483	7
820	161	DSC	3.11	545	7
632	157	DSC	2.59	430	7
821	127	K	296	537	7
822	135	K	2.78	436	7
823	218	DSC	2.53	526	7
824	212	DSC	2.82	510	7
825	237	DSC	2.55	492	7
826	202	DSC	2.74	543	7
827	187	DSC	2.82	544	7
828	209	DSC	2.56	468	7
829	154	DSC	3.24	536	7
830	241	DSC	2.65	527	7
663	200	DSC	2.38	498	7
664	80	K	2.77	469	7
831	178	DSC	2.84	536	7

413	146	K	3.03	507	6
414			3.1	451	6
422	234.9	DSC	3.28	420	6
424			3.16	440	6
426			2.82	447	6
428			0.56	432	5
34	173.67	DSC	2.46	462	6
430			3.78	498	6
125	99.31	DSC	2.58	435	6
431	147.77	DSC	3.99	509	6
432	251.77	DSC	2.79	436	6
433			2.39	423	6
434			2.96	487	6
435			2.89	447	6
5	from 123.12 to 128	DSC or K	3.63 to 3.91	501	8 or 6
436			2.86	489	6
95a + 95			3.49	559	6
437			2.66	433	6
438	140.67	DSC	2.44	391	6
440			2.9	437	6
441			3.04	481	6
442	120.06	DSC	3.39	461	6
443	138.47	DSC	2.94	450	6
444	137.16	DSC	2.95	450	6
445	182.81 227.95 227.75	DSC DSC DSC	3.31	473	6
447			3.05	447	6
448	80	K	3.6	470	6
449	118.96	DSC	3.05	447	6
450	224.87	DSC	3.6	470	6
452			3.13	461	6
453			3.04	396	6

832	111	DSC	3.67	482	7
833	153	DSC	2.79	493	7
834	198	DSC	2.63	475	7
835	144	DSC	3.24	552	7
837			2.63	472	7
641	155	K	2.27	476	7
642	127	DSC	2.36	486	7
838	171	DSC	2.28	462	7
839	126	DSC	3.49	567	7
840	158	DSC	2.44	517	7
841			3.08	463	7
842	154	DSC	3.24	553	7
665	206	K	3.3	516	7
643	160	K	2.4	503	7
836			2.57	496	7
606	142	K	3.25	420	6
666	163	K	2.68	516	7
844	125	K	2.68	530	7
845	167	DSC	3.18	512	7
846	170	K	2.49	477	6
847			0.79	435	5
667	181	DSC	3.07	471	7
848	223	K	3.13	496	7
849	80	K	3.11	453	7
850	142	DSC	3.67	513	7
851	137	DSC	4	543	7
669	122	DSC	3.55	526	7
852	233	K	3.43	496	7
853	254	K	3.12	496	7
670	157	K	3.03	461	6
855			1.01 0.86	501 501	4 5

134	184 194	DSC K	2.71 2.72	504 504	6 6
454	244.06	DSC	3.25	545	8
455	155	K	3.13	586	8
456	143.58	DSC	2.9	502	8
96	168.08	DSC	2.94	501	8
457			3.2	571	6
458	171.31	DSC	3.34	470	6
44a + 44	215.29	DSC	3.84	495	6
459	99.32	DSC	4.23	593	6
460			3.14	438	6
463			3.72	568	6
464			4.39	555	6
465	141.31	DSC	2.91	473	6
467	135	K	2.89	503	6
468	130	K	2.84	503	6
469	135	K	3.05	487	6
471	140	K	3.14	489	6
473	140	K	3.14	489	6
474			3.56	590	6
135	194	DSC K	3.13	461	6
475	152.98	DSC	3.18	420	6
476	170.9	DSC	3.6	448	6
477	80	K	3.05	380	6
478	70	K	2.93	407	6
136	166	DSC	3.14	531	6
480		DSC	3.13	515	6
97	187.39 187.8	DSC DSC	2.87 2.87	436 436	6 6
98	186.98 186.27	DSC DSC	2.87 2.87	436 436	6 6
481	172.5	DSC	3.67	474	6

696	188	K	3.07	456	6
856	97	K	2.96	482	6
857	180	K	3.43	487	6
671	144	DSC	2.97	501	6
858	154	K	3.58	470	6
673	181 180	DSC K	3.76	482	6
676	88	K	3.71	503	6
859	80	K	3.22	485	6
860	151	DSC	3.21	485	6
678	148	DSC	2.27 2.87	539 539	7 6
680	167	DSC	2.58	460	6
861	156	K	3.22	447	6
691	99	DSC	2.17 2.73	477 477	7 6
862	178	K	3.44	573	6
682	149	K	2.86	461	6
863	170	K	3.77	462	6
864	154	K	3.77	462	6
865	146	DSC	3.81	498	6
866	144	DSC	3.81	498	6
679			2.99	491	6
693	151	DSC	2.79	502	6
867	178	K	2.98	424	6
868	178	K	2.98	424	6
694	197	DSC	3.49	501	6
869	138	K	3.46	489	6
692	157	K	3.19	498	6
675	187	K	3.44	517	6
870	173	K	3.34	456	6
685	203	K	3.17	442	6

482	106	K	2.82	419	6
483	134	K	3.01	433	6
484			3.03	461	6
485			0.62	474	5
487	143.13	DSC	4.82	488	1
488	96.65	DSC	2.73	393	6
			2.73	393	6
489	128.03	DSC	3.4	551	6
490	223.8	DSC	2.63	449	6
493	87.6	DSC	3.05	435	6
494	135.12	DSC	3.02	447	6
495			3.83	486	6
137			2.8	433	6
499	130	K	3.34	501	6
500			0.69	421	5
501	154.26	DSC	2.63	449	6
502	153.17	DSC	2.65	449	6
503			3.48	511	6
139			3.53	517	6
504	101.17	DSC	2.53	476	6
	104.18	DSC	2.53	476	6
140	123.19	DSC	3.25	458	6
	128.99	DSC	3.26	458	6
506			3.42	463	6
507			3.36	463	6
508	165.94	DSC	3.49	446	6
509			3.42	634	6
510	70	K	2.77	407	6
141	150	K	3.29	448	8
	171.28	DSC	3.54	448	6
	156.7	DSC			
511			0.74	502	5
512	180	K	3.41	473	6
513	160.65	DSC	3.55	469	6
514	139	K	3.3	489	6
	135.37	DSC			
515			4.78	516	3

871	154	DSC	3.05	450	6
872	89	DSC	4.08	515	6
677	177	DSC	3.44	435	6
873	183	DSC	3.07	457	6
874	107	K	3.24	450	6
875	98	K	3.24	450	6
877	213	DSC	2.94	500	6
878	143	DSC	3.52	503	6
879			0.89	531	5
880	135	DSC	3.45	501	6
668	137	DSC			
672	194	DSC			
683	123	DSC			
881	156.5	DSC	4.53	440	6
882	164	K	4.46	454	6
883	194	DSC	4.35	399	6
884	80	K	3.58	483	6
885	148	DSC	5.13	640	6
886	130	K	3.97	455	6
887	245	DSC	2.81	457	6
888			2.96	500	6
889			0.6		5
890			6.18	444	1
891	186	K	3.89	489	6
892	157	K	4.32	534	6
893			1.05	443	4
894			3.58	489	6
895	182	DSC	3.56	470	6
896	166	DSC	2.74	419	6
897	153	DSC	2.75	419	6

516			0.98 1.08	471 471	5 4
35			0.92	416	5
517	187.18	DSC			
518	187.66	DSC	3.01	479	6
519	163.93	DSC	3.08	516	6
102			0.89	487	4
521			0.91	475	5
522	134.16	DSC	2.99	503	6
523	125.77	DSC	3.53	517	6
142	126.8	DSC	3.53	517	6
524	122.12	DSC	3	433	6
525	223.34	DSC	3.32	514	6
526	234.8	DSC	3.32	514	6
527	144 145.21	K DSC	3.19	484	6
143	184.72	DSC	3.15	516	6
528	110.16	DSC	2.86	518	6
529	141.1	DSC	3.06	459	6
530	132.62 132.62	DSC DSC	2.96	488	6
10			4.28	551	6
531			3.53	489	6
36	160.14	DSC	3.38	418	6
532	164 165	K DSC	3.06	470	6
533	180.45 181.17	DSC DSC	2.71	449	6
534	205.99	DSC	3.6	475	6
535	180 179.19	K DSC	3.4	473	6
536	209.5	DSC	3.72	461	6
537	201.67	DSC	3.7	461	6

898			6.24	430	1
899			3.6	480	6
900	214	DSC	3.47	475	6
901	208	DSC	3.02	433	6
902	>260	K	2.34	470	6
903	153	DSC	3.65	513	6
904	181	DSC	3.07	471	6
905			0.98	457	5
906			2.83	517	6
907			2.97	491	7
908			3.19	501	7
909			2.99	491	7
910			3.18	501	7
911			3.13	501	7
912	102	K	2.89	474	7
913	170	K	3.1	538	7
914	166	K	3.01	599	7
915			3.22	542	7
916	70	K	3.53	552	7
917	222	DSC	2.73	517	7
918			3.36	496	7
919	130	DSC	3.5	490	7
920	123.5	DSC	3.39	541	7
921	123	DSC	3.07	493	7
922	123	DSC	3.07	493	7
927	152	K	2.52	483	7
928	130	K	5.42	442	9

NMR Data

The below NMR experiments were carried out using a Bruker Avance 500 and a Bruker Avance DRX 400 spectrometers at ambient temperature, using internal deuterium lock and equipped with reverse triple-resonance(^{1}H, ^{13}C,^{15}N TXI) probe head for the 500MHz and with reverse double-resonance (^{1}H, ^{13}C, SEI) probe head for the 400MHz. Chemical shifts (δ) are reported in parts per million (ppm).

Compound 131

^{1}H NMR (500 MHz, DMSO-d_6) δ 9.01 (s, 1H), 8.60 (s, 1H), 8.28 (s, 1H), 7.79 (d, *J* = 9.1 Hz, 1H), 7.27 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.16 (d, *J* = 2.5 Hz, 1H), 6.38 - 6.49 (m, 3H), 4.82 (br. s., 2H), 4.23 (t, *J* = 5.7 Hz, 2H), 3.96 (t, *J* = 4.9 Hz, 2H), 3.80 (t, *J* = 5.7 Hz, 2H), 3.71 - 3.75 (m, 5H), 3.69 (t, *J* = 4.9 Hz, 2H), 1.05 - 1.26 (m, 1H), 0.42 - 0.51 (m, 2H), 0.16 - 0.25 (m, 2H)

Compound 149

^{1}H NMR (500 MHz, DMSO-d_6) δ 9.00 (s, 1H), 8.58 (s, 1H), 8.22 (s, 1H), 7.80 (d, *J* = 9.1 Hz, 1H), 7.31 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.25 (d, *J* = 2.5 Hz, 1H), 6.80 (br. s., 1H), 6.49 (d, *J* = 1.9 Hz, 2H), 6.42 (br. s., 1H), 4.15 - 4.31 (m, 2H), 3.89 - 4.00 (m, 4H), 3.74 (s, 6H)

Compound 148

^{1}H NMR (500 MHz, DMSO-d_6) δ 9.00 (s, 1H), 8.58 (s, 1H), 8.22 (s, 1H), 7.80 (d, *J* = 9.1 Hz, 1H), 7.31 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.25 (d, *J* = 2.5 Hz, 1H), 6.80 (br. s., 1H), 6.49 (d, *J* = 1.9 Hz, 2H), 6.42 (br. s., 1H), 4.15 - 4.31 (m, 2H), 3.89 - 4.00 (m, 4H), 3.74 (s, 6H)

Compound 147

^{1}H NMR (400 MHz, DMSO-d_6) δ 9.76 (br. s., 1H), 9.01 (s, 1H), 8.64 (s, 1H), 8.22 (s, 1H), 7.79 (d, *J* = 9.3 Hz, 1H), 7.26 (dd, *J* = 2.8, 9.3 Hz, 1H), 7.18 (d, *J* = 2.8 Hz, 1H), 6.36 - 6.51 (m, 3H), 4.58 (spt, *J* = 6.6 Hz, 1H), 4.03 - 4.19 (m, 2H), 3.93 (t, *J* = 7.3 Hz, 2H), 3.75 (s, 6H), 3.09 - 3.20 (m, 2H), 2.08 (td, *J* = 7.3, 14.5 Hz, 2H), 1.49 (d, *J* = 6.6 Hz, 6H)

Compound 146

^{1}H NMR (400 MHz, DMSO-d_6) δ 8.99 (s, 1H), 8.71 (s, 1H), 8.65 (t, *J* = 1.5 Hz, 1H), 8.51 - 8.56 (m, 2H), 8.19 (s, 1H), 7.79 (d, *J* = 9.1 Hz, 1H), 7.39 (dd, *J* = 2.8, 9.1 Hz, 1H), 7.22 (d, *J* = 2.8 Hz, 1H), 6.55 (d, *J* = 2.1 Hz, 2H), 6.37 (t, *J* = 2.1 Hz, 1H), 5.31 (s, 2H), 3.91 (s, 3H), 3.72 (s, 6H)

Compound 145

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.98 (br. s., 2H), 8.55 (s, 1H), 8.20 (s, 1H), 7.79 (d, *J* = 9.1 Hz, 1H), 7.34 (dd, *J* = 2.6, 9.1 Hz, 1H), 7.26 (d, *J* = 2.6 Hz, 1H), 6.55 (d, *J* = 1.9 Hz, 2H), 6.43 (s, 1H), 4.17 (br. s., 1H), 3.88 - 3.99 (m, 6H), 3.75 (s, 6H), 3.30 (td, *J* = 6.3, 11.9 Hz, 1H), 3.02 - 3.16 (m, 1H), 2.96 (q, *J* = 9.6 Hz, 1H), 1.22 (d, *J* = 6.3 Hz, 6H)

Compound 144

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.96 (s, 1H), 8.55 (s, 1H), 8.21 (s, 1H), 7.77 (d, *J* = 9.5 Hz, 1H), 7.73 (br. s., 1H), 7.27 (dd, *J* = 2.7, 9.5 Hz, 1H), 7.14 (d, *J* = 2.7 Hz, 1H), 6.45 (d, *J* = 2.2 Hz, 2H), 6.38 - 6.41 (m, 1H), 3.99 (t, *J* = 6.6 Hz, 2H), 3.93 (s, 3H), 3.73 (s, 6H), 3.14 (br. s., 2H), 3.01 (s, 2H), 2.68 (t, *J* = 6.6 Hz, 2H), 2.63 (t, *J* = 5.2 Hz, 2H)

Compound 143

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.96 (s, 1H), 8.55 (s, 1H), 8.21 (s, 1H), 7.77 (d, *J* = 9.1 Hz, 1H), 7.28 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.14 (d, *J* = 2.5 Hz, 1H), 6.44 (d, *J* = 2.2 Hz, 2H), 6.32 - 6.42 (m, 1H), 3.98 (t, *J* = 6.9 Hz, 2H), 3.93 (s, 3H), 3.73 (s, 6H), 3.35 - 3.43 (m, 4H), 2.63 (t, *J* = 6.9 Hz, 2H), 2.44 (t, *J* = 4.9 Hz, 2H), 2.38 (t, *J* = 4.9 Hz, 2H), 1.97 (s, 3H)

Compound 142

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.95 (s, 1H), 8.55 (s, 1H), 8.20 (s, 1H), 7.75 (d, *J* = 9.5 Hz, 1H), 7.34 (dd, *J* = 2.7, 9.5 Hz, 1H), 7.22 (d, *J* = 2.7 Hz, 1H), 6.49 (d, *J* = 1.9 Hz, 2H), 6.38 (t, *J* = 1.9 Hz, 1H), 5.16 (d, *J* = 5.1 Hz, 1H), 3.83 - 4.00 (m, 5H), 3.69 - 3.78 (m, 7H), 3.19 - 3.31 (m, 2H), 2.68 - 2.78 (m, 1H), 2.66 (td, *J* = 6.1, 12.1 Hz, 1H), 2.30 - 2.40 (m, 1H)

Compound 141

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.95 (s, 1H), 8.55 (s, 1H), 8.21 (s, 1H), 7.76 (d, *J* = 9.1 Hz, 1H), 7.26 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.14 (d, *J* = 2.5 Hz, 1H), 6.34 - 6.44 (m, 3H), 4.49 (s, 1H), 3.84 - 3.99 (m, 5H), 3.74 (s, 6H), 1.66 - 1.86 (m, 2H), 1.16 (s, 6H)

Compound 140

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.95 (s, 1H), 8.55 (s, 1H), 8.20 (s, 1H), 7.76 (d, *J* = 9.1 Hz, 1H), 7.28 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.14 (d, *J* = 2.5 Hz, 1H), 6.46 (d, *J* = 1.9 Hz, 2H), 6.41 (t, *J* = 1.9 Hz, 1H), 3.83 - 3.96 (m, 5H), 3.74 (s, 6H), 2.82 (t, *J* = 6.7 Hz, 2H), 2.78 (t, *J* = 6.7 Hz, 2H), 2.57 (t, *J* = 6.7 Hz, 2H), 2.22 (br. s., 1H)

Compound 139

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.95 (s, 1H), 8.55 (s, 1H), 8.20 (s, 1H), 7.75 (d, *J* = 9.5 Hz, 1H), 7.34 (dd, *J* = 2.7, 9.5 Hz, 1H), 7.22 (d, *J* = 2.7 Hz, 1H), 6.49 (d, *J* = 1.9 Hz, 2H), 6.38 (t, *J* = 1.9 Hz, 1H), 5.16 (d, *J* = 5.1 Hz, 1H), 3.83 - 4.00 (m, 5H), 3.69 - 3.78 (m, 7H), 3.19 - 3.31 (m, 2H), 2.68 - 2.78 (m, 1H), 2.66 (td, *J* = 6.1, 12.1 Hz, 1H), 2.30 - 2.40 (m, 1H)

Compound 137

^{1}H NMR (400 MHz, DMSO-d_6) δ 9.10 (br.s., 3H), 8.49 (s, 2H), 7.84 (d, *J* = 9.6 Hz, 1H), 7.39 (dd, *J* = 2.7, 9.6 Hz, 1H), 7.29 (d, *J* = 2.7 Hz, 1H), 6.52 (d, *J* = 2.1 Hz, 2H), 6.46 (t, *J* = 2.1 Hz, 1H), 4.21 (t, *J* = 7.3 Hz, 2H), 3.76 (s, 6H), 3.35 (m, 1H), 3.15 (br. s., 2H), 1.25 (d, *J* = 6.1 Hz, 6H)

Compound Number 98

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.95 (s, 1H), 8.56 (s, 1H), 8.21 (s, 1H), 7.75 (d, *J* = 9.1 Hz, 1H), 7.34 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.25 (d, *J* = 2.5 Hz, 1H), 6.50 (d, *J* = 2.2 Hz, 2H), 6.37 (t, *J* = 2.2 Hz, 1H), 5.01 (d, *J* = 2.8 Hz, 1H), 4.70 - 4.79 (m, 1H), 4.03 (dd, *J* = 3.6, 14.9 Hz, 1H), 3.92 (s, 3H), 3.81 (br. s., 1H), 3.73 (s, 6H), 3.68 (dd, *J* = 8.1, 14.9 Hz, 1H), 3.36 - 3.48 (m, 2H)

Compound 136

^{1}H NMR (500 MHz, DMSO-d_6) δ 9.03 (br.s., 2H), 8.58 (s, 1H), 8.25 (s, 1H), 7.83 (d, *J* = 9.5 Hz, 1H), 7.36 (dd, *J* = 2.5, 9.5 Hz, 1H), 7.23 (d, *J* = 2.5 Hz, 1H), 6.52 (d, *J* = 1.9 Hz, 2H), 6.46 (t, *J* = 1.9 Hz, 1H), 4.19 - 4.21 (m, 2H), 4.10 (d, *J* = 6.9 Hz, 2H), 3.84 (dd, *J* =

2.8, 11.7 Hz, 2H), 3.76 (s, 6H), 3.31 - 3.38 (td, *J* = 6.1, 11.7, 1H), 3.27 (t, *J* = 11.7 Hz, 2H), 3.14 - 3.18 (m, 2H), 2.16 - 2.08 (m, 1H), 1.43 (d, *J* = 11.7 Hz, 2H), 1.18 - 1.37 (m, 8H)

Compound 135

^{1}H NMR (500 MHz, DMSO-d_6) δ 9.03 (br.s., 2H), 8.62 (s, 1H), 8.23 (s, 1H), 7.83 (d, *J* = 9.5 Hz, 1H), 7.36 (dd, *J* = 2.5, 9.5 Hz, 1H), 7.25 (d, *J* = 2.5 Hz, 1H), 6.51 (d, *J* = 2.2 Hz, 2H), 6.46 (t, *J* = 2.2 Hz, 1H), 3.76 (s, 6H), 4.12 - 4.27 (m, 4H), 3.30 - 3.43 (m, 1H), 3.07 - 3.19 (m, 2H), 1.44 (t, *J* = 7.2 Hz, 3H), 1.25 (d, *J* = 6.3 Hz, 6H)

Compound 134

^{1}H NMR (500 MHz, DMSO-d_6) δ 9.05 (s, 1H), 8.93 (m, 1H), 8.57 (s, 1H), 8.25 (s, 1H), 8.18 (q, *J* = 4.6 Hz, 1H), 7.83 (d, *J* = 9.1 Hz, 1H), 7.33 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.28 (d, *J* = 2.5 Hz, 1H), 6.52 (d, *J* = 2.2 Hz, 2H), 6.46 (t, *J* = 2.2 Hz, 1H), 4.88 (s, 2H), 4.19 (t, *J* = 7.6 Hz, 2H), 3.75 (s, 6H), 3.29 - 3.42 (m, 1H), 3.16 (br. s., 2H), 2.64 (d, *J* = 4.6 Hz, 3H), 1.25 (d, *J* = 6.6 Hz, 6H)

Compound Number 5

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.95 (s, 1H), 8.55 (s, 1H), 8.20 (s, 1H), 7.75 (d, *J* = 9.1 Hz, 1H), 7.28 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.12 (d, *J* = 2.5 Hz, 1H), 6.40 (s, 3H), 3.93 (s, 3H), 3.88 (t, *J* = 7.1 Hz, 2H), 3.74 (s, 6H), 3.11 - 3.28 (m, 2H), 2.68 - 2.72 (m,, 2H), 2.39 - 2.48 (m, 1H), 1.78 (quin, *J* = 7.1 Hz, 2H)

Compound 133

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.97 (s, 1H), 8.56 (s, 1H), 8.21 (s, 1H), 7.77 (d, *J* = 9.5 Hz, 1H), 7.27 (dd, *J* = 2.5, 9.5 Hz, 1H), 7.12 (d, *J* = 2.5 Hz, 1H), 6.29 - 6.49 (m, 3H), 3.96 (t, *J* = 6.8 Hz, 2H), 3.93 (s, 3H), 3.74 (s, 6H), 3.19 - 3.29 (m, 1H), 2.70 - 2.85 (m, 5H), 2.42 - 2.46 (m, 1H), 2.10 - 2.24 (m, 1H), 1.88 - 1.98 (m, 1H)

Compound 132

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.96 (s, 1H), 8.55 (s, 1H), 8.20 (s, 1H), 7.76 (d, *J* = 9.3 Hz, 1H), 7.23 (dd, *J* = 2.8, 9.3 Hz, 1H), 7.07 (d, *J* = 2.8 Hz, 1H), 6.41 (s, 3H), 3.93 (s, 3H), 3.81 (t, *J* = 7.4 Hz, 2H), 3.74 (s, 6H), 3.23 - 3.32 (m, 4H), 2.23 (t, *J* = 8.1 Hz, 2H), 1.93 (m, 2H), 1.84 (m,2H)

Compound Number 300

^{1}H NMR (400 MHz, DMSO-d_6) δ 9.05 (s, 1H), 8.54 - 8.63 (m, 1H), 8.24 (s, 1H), 7.84 (d, *J* = 9.1 Hz, 1H), 7.41 (d, *J* = 2.5 Hz, 1H), 7.34 (dd, *J* = 2.5, 9.1 Hz, 1H), 7.20 (s, 1H), 6.88 (s, 1H), 6.50 (s, 2H), 6.42 - 6.47 (m, 1H), 4.99 (s, 2H), 3.93 (s, 3H), 3.74 (s, 6H), 3.53 (s, 3H)

Compound Number 4

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.95 (s, 1H), 8.55 (s, 1H), 8.20 (s, 1H), 7.76 (d, *J* = 9.1 Hz, 1H), 7.27 (dd, *J* = 2.8, 9.1 Hz, 1H), 7.13 (d, *J* = 2.8 Hz, 1H), 6.46 (d, *J* = 2.2 Hz, 2H), 6.40 (t, *J* = 2.2 Hz, 1H), 3.93 (s, 3H), 3.88 (t, *J* = 6.9 Hz, 2H), 3.74 (s, 6H), 2.79 (t, *J* = 6.9 Hz, 2H), 2.70 (m, 1H), 1.69 (br. s., 1H), 0.95 (d, *J* = 6.3 Hz, 6H)

Compound Number 84

^{1}H NMR (500 MHz, DMSO-d_6) δ 8.94 (s, 1H), 8.55 (s, 1H), 8.20 (s, 1H), 7.75 (d, *J* = 9.3 Hz, 1H), 7.28 (dd, *J* = 2.5, 9.30 Hz, 1H), 7.15 (d, *J* = 2.5 Hz, 1H), 6.45 (d, *J* = 2.2 Hz, 2H), 6.38 - 6.42 (m, 1H), 3.93 (s, 3H), 3.82 (t, *J* = 7.1 Hz, 2H), 3.74 (s, 6H), 2.80 (t, *J* = 7.1 Hz, 2H), 1.55 (br. s., 2H)

Compound 130

^{1}H NMR (500 MHz, DMSO-d_6) δ 9.00 (s, 1H), 8.69 (s, 1H), 8.33 (s, 1H), 7.79 (d, *J* = 9.5 Hz, 1H), 7.28 (dd, *J* = 2.7, 9.5 Hz, 1H), 7.14 (d, *J* = 2.7 Hz, 1H), 6.38 - 6.47 (m, 3H), 5.55 (br.s., 1H), 4.34 (t, *J* = 6.6 Hz, 2H), 3.62 - 3.91 (m, 12H), 3.36 - 3.55 (m, 6H), 3.09 - 3.31 (m, 4H), 2.28 - 2.38 (m, 2H), 1.75 - 1.97 (m, 2H), 1.10 - 1.23 (m, 1H), 0.43 - 0.52 (m, 2H), 0.15 - 0.24 (m, 2H)

The below NMR experiments were carried out using a Bruker Avance AV400 spectrometer, using an internal deuterium lock and equipped with a 4-nucleus (^{1}H, ^{13}C, ^{19}F, ^{31}P) probe head. Chemical shifts (δ) are reported in parts per million (ppm) at 27°C.

Compound 138

^{1}H NMR (400 MHz, DMSO-d6): 9.07 (1H, s), 8.59 (1H, s), 8.56-8.47 (1H, m), 8.27-8.21 (1H, m), 7.87 (1H, d), 7.54-7.47 (1H, m), 7.43-7.32 (3H, m), 7.27-7.18 (1H, m), 3.98-3.89

(3H, m), 3.83 (2H, d), 2.76 (3H, d), 1.23-1.13 (1H, m), 0.50-0.41 (2H, m), 0.22-0.14 (2H, m).

Compound Number 99

^{1}H NMR (400 MHz, Me-d3-OD): 8.89 (1H, s), 8.40 (1H, s), 8.23 (1H, s), 7.79 (1H, d), 7.41 (1H, dd), 7.30 (1H, d), 7.01 (2H, s), 6.53 (2H, s), 6.47-6.40 (1H, m), 4.57 (2H, s), 4.01 (3H, s), 3.77 (7H, s).

Compound 200

^{1}H NMR (400 MHz, DMSO-d6): 8.96 (1H, s), 8.56 (1H, s), 8.21 (1H, s), 7.76 (1H, d), 7.25 (1H, dd), 7.11 (1H, d), 6.46-6.36 (3H, m), 3.99-3.82 (5H, m), 3.75 (6H, s), 1.23 (3H, t).

Compound 201

^{1}H NMR (400 MHz, DMSO-d6):: 8.92 (1H, s), 8.54 (1H, s), 8.20 (1H, s), 7.76 (1H, d), 6.99 (1H, dd), 6.81 (2H, dd), 6.64 (1H, d), 3.92 (6H, d), 3.88-3.73 (5H, m), 1.24 (3H, t).

Compound 11

^{1}H NMR (400 MHz, DMSO-d6): 8.98 (1H, s), 8.56 (1H, s), 8.22 (1H, s), 7.78 (1H, d), 7.30 (1H, dd), 7.16 (1H, d), 6.43 (2H, d), 6.40 (1H, t), 3.94 (3H, s), 3.74 (6H, s), 3.41 (3H, s).

Compound 202

^{1}H NMR (400 MHz, DMSO-d6): 8.96 (1H, s), 8.56 (1H, s), 8.21 (1H, s), 7.77 (1H, d), 7.26 (1H, dd), 7.13 (1H, d), 6.42 (3H, s), 3.93 (3H, s), 3.82-3.70 (8H, m), 1.24-1.12 (1H, m), 0.53-0.43 (2H, m), 0.26-0.16 (2H, m).

Compound 12

^{1}H NMR (400 MHz, DMSO-d6): 8.96 (1H, s), 8.56 (1H, s), 8.21 (1H, s), 7.76 (1H, d), 7.26 (1H, dd), 7.08 (1H, d), 6.41 (3H, dd), 3.93 (3H, s), 3.79 (2H, t), 3.75 (6H, s), 1.73-1.63 (2H, m), 0.96 (3H, t)

Compound 204

^{1}H NMR (400 MHz, DMSO-d6): 9.00-8.94 (1H, m), 8.59-8.53 (1H, m), 8.25-8.18 (1H, m), 7.77 (1H, d), 7.30 (1H, dd), 7.17 (1H, d), 6.44 (2H, d), 6.40 (1H, t), 4.03 (2H, t), 3.94 (3H, s), 3.74 (6H, s), 3.60 (2H, t), 3.29 (3H, s).

Compound 13

^{1}H NMR (400 MHz, DMSO-d6): 8.97 (1H, s), 8.56 (1H, s), 8.21 (1H, s), 7.77 (1H, d), 7.30 (1H, dd), 7.10 (1H, d), 6.41 (3H, s), 3.93 (3H, s), 3.74 (6H, s), 3.69 (2H, d), 2.09-1.97 (1H, m), 0.98 (6H, d).

Compound 205

^{1}H NMR (400 MHz, DMSO-d6): 9.02 (1H, s), 8.60-8.54 (1H, m), 8.22 (1H, s), 7.82 (1H, d), 7.36 (1H, dd), 7.24 (1H, d), 6.48 (2H, d), 6.40 (1H, t), 6.32 (1H, s), 5.25 (2H, s), 3.97-3.89 (3H, m), 3.78-3.69 (7H, m), 3.29 (3H, s), 2.18 (3H, s).

Pharmacological part

Biological assays A

FGFR1 (enzymatic assay)

In a final reaction volume of 30 µL, FGFR1 (h) (25 ng/ml) was incubated with 50 mM HEPES pH 7.5, 6mM $MnCl_2$, 1 mM DTT, 0,1 mM Na_3VO_4, 0,01% Triton-X-100, 500 nM Btn-Flt3 and 5 µM ATP in the presence of compound (1% DMSO final). After incubation for 60 minutes at room temperature the reaction was stopped with 2.27 nM EU-anti P-Tyr, 7 mM EDTA, 31.25 nM SA-XL-665 and 0.02% BSA which was present for 60 minutes at room temperature. Time-Resolved Fluorescence Resonance Energy Transfer (TR-FRET) signal (ex340 nm. Em 620 nm, em 655 nm) was measured afterwards and results are expressed in RFU (Relative Fluorescence Units). In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} (-logIC_{50}) value.

FGFR2 (enzymatic assay)

In a final reaction volume of 30 µL, FGFR2 (h) (150 ng/ml) was incubated with 50 mM HEPES pH 7.5, 6mM $MnCl_2$, 1 mM DTT, 0,1 mM Na_3VO_4, 0,01% Triton-X-100, 500 nM Btn-Flt3 and 0.4 µM ATP in the presence of compound (1% DMSO final). After incubation for 60 minutes at room temperature the reaction was stopped with 2.27 nM EU-anti P-Tyr, 7 mM EDTA, 31.25 nM SA-XL-665 and 0.02% BSA which was present

for 60 minutes at room temperature. Time-Resolved Fluorescence Resonance Energy Transfer (TR-FRET) signal (ex340 nm. Em 620 nm, em 655 nm) was measured afterwards and results are expressed in (Relative Fluorescence Units). In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} (-logIC_{50}) value.

FGFR3 (enzymatic assay)

In a final reaction volume of 30 µL, FGFR3 (h) (40 ng/ml) was incubated with 50 mM HEPES pH 7.5, 6mM $MnCl_2$, 1 mM DTT, 0,1 mM Na_3VO_4, 0,01% Triton-X-100, 500 nM Btn-Flt3 and 25 µM ATP in the presence of compound (1% DMSO final). After incubation for 60 minutes at room temperature the reaction was stopped with 2.27 nM EU-anti P-Tyr, 7 mM EDTA, 31.25 nM SA-XL-665 and 0.02% BSA which was present for 60 minutes at room temperature. Time-Resolved Fluorescence Resonance Energy Transfer (TR-FRET) signal (ex340 nm. Em 620 nm, em 655 nm) was measured afterwards and results are expressed in RFU (Relative Fluorescence Units). In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} (-logIC_{50}) value.

FGFR4 (enzymatic assay)

In a final reaction volume of 30 µL, FGFR4 (h) (60 ng/ml) was incubated with 50 mM HEPES pH 7.5, 6mM $MnCl_2$, 1 mM DTT, 0,1 mM Na_3VO_4, 0,01% Triton-X-100, 500 nM Btn-Flt3 and 5 µM ATP in the presence of compound (1% DMSO final). After incubation for 60 minutes at room temperature the reaction was stopped with 2.27 nM EU-anti P-Tyr, 7 mM EDTA, 31.25 nM SA-XL-665 and 0.02% BSA which was present for 60 minutes at room temperature. Time-Resolved Fluorescence Resonance Energy Transfer (TR-FRET) signal (ex340 nm. Em 620 nm, em 655 nm) was measured afterwards and results are expressed in RFU (Relative Fluorescence Units). In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} (-logIC_{50}) value.

KDR (VEGFR2) (enzymatic assay)

In a final reaction volume of 30 µL, KDR (h) (150 ng/ml) was incubated with 50 mM HEPES pH 7.5, 6mM $MnCl_2$, 1 mM DTT, 0,1 mM Na_3VO_4, 0,01% Triton-X-100, 500 nM Btn-Flt3 and 3 µM ATP in the presence of compound (1% DMSO final). After incubation

for 120 minutes at room temperature the reaction was stopped with 2.27 nM EU-anti P-Tyr, 7 mM EDTA, 31.25 nM SA-XL-665 and 0.02% BSA which was present for 60 minutes at room temperature. Time-Resolved Fluorescence Resonance Energy Transfer (TR-FRET) signal (ex340 nm. Em 620 nm, em 655 nm) was measured afterwards and results are expressed in RFU (Relative Fluorescence Units). In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} (-logIC_{50}) value.

Ba/F3-FGFR1 (minus IL3 or plus IL3) (cellular proliferation assay)

In a 384 well plate, 100 nl of compound dilution in DMSO was sprayed before adding 50 µl cell culture medium (phenol red free RPMI-1640, 10 % FBS, 2 mM L-Glutamine and 50 µg/ml Gentamycin) containing 20000 cells per well of Ba/F3-FGFR1-transfected cells. Cells were put in an incubator at 37°C and 5 % CO_2. After 24 hours, 10 µl of Alamar Blue solution (0.5 mM $K_3Fe(CN)_6$, 0.5 mM $K_4Fe(CN)_6$, 0.15 mM Resazurin and 100 mM Phosphate Buffer) was added to the wells, incubated for 4 hours at 37°C and 5% CO_2 before RFU's (Relative Fluorescence Units) (ex. 540 nm., em. 590 nm.) were measured in a flurorescence plate reader.
In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} (-logIC_{50}) value.
As a counterscreen the same experiment was performed in the presence of 10 ng/ml murine IL3.

Ba/F3-FGFR3 (minus IL3 or plus IL3) (cellular proliferation assay)

In a 384 well plate, 100 nl of compound dilution in DMSO was sprayed before adding 50 µl cell culture medium (phenol red free RPMI-1640, 10 % FBS, 2 mM L-Glutamine and 50 µg/ml Gentamycin) containing 20000 cells per well of Ba/F3-FGFR3-transfected cells. Cells were put in an incubator at 37°C and 5 % CO_2. After 24 hours, 10 µl of Alamar Blue solution (0.5 mM $K_3Fe(CN)_6$, 0.5 mM $K_4Fe(CN)_6$, 0.15 mM Resazurin and 100 mM Phosphate Buffer) was added to the wells, incubated for 4 hours at 37°C and 5% CO_2 before RFU's (Relative Fluorescence Units) (ex. 540 nm., em. 590 nm.) were measured in a flurorescence plate reader.
In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} (-logIC_{50}) value.

As a counterscreen the same experiment was performed in the presence of 10 ng/ml murine IL3.

Ba/F3-KDR (minus IL3 or plus IL3) (cellular proliferation assay)

In a 384 well plate, 100 nl of compound dilution in DMSO was sprayed before adding 50 µl cell culture medium (phenol red free RPMI-1640, 10 % FBS, 2 mM L-Glutamine and 50 µg/ml Gentamycin) containing 20000 cells per well of Ba/F3-KDR-transfected cells. Cells were put in an incubator at 37°C and 5 % CO_2. After 24 hours, 10 µl of Alamar Blue solution (0.5 mM $K_3Fe(CN)_6$, 0.5 mM $K_4Fe(CN)_6$, 0.15 mM Resazurin and 100 mM Phosphate Buffer) was added to the wells, incubated for 4 hours at 37°C and 5% CO_2 before RFU's (Relative Fluorescence Units) (ex. 540 nm., em. 590 nm.) were measured in a flurorescence plate reader.

In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} ($-\log IC_{50}$) value.

As a counterscreen the same experiment was performed in the presence of 10 ng/ml murine IL3.

Ba/F3-Flt3 (minus IL3 or plus IL3) (cellular proliferation assay)

In a 384 well plate, 100 nl of compound dilution in DMSO was sprayed before adding 50 µl cell culture medium (phenol red free RPMI-1640, 10 % FBS, 2 mM L-Glutamine and 50 µg/ml Gentamycin) containing 20000 cells per well of Ba/F3-Flt3-transfected cells. Cells were put in an incubator at 37°C and 5 % CO_2. After 24 hours, 10 µl of Alamar Blue solution (0.5 mM $K_3Fe(CN)_6$, 0.5 mM $K_4Fe(CN)_6$, 0.15 mM Resazurin and 100 mM Phosphate Buffer) was added to the wells, incubated for 4 hours at 37°C and 5% CO_2 before RFU's (Relative Fluorescence Units) (ex. 540 nm., em. 590 nm.) were measured in a flurorescence plate reader.

In this assay, the inhibitory effect of different compound concentrations (range 10 µM to 0.1 nM) was determined and used to calculate an IC_{50} (M) and pIC_{50} ($-\log IC_{50}$) value.

As a counterscreen the same experiment was performed in the presence of 10 ng/ml murine IL3.

Data for the compounds of the invention in the above assays are provided in Table A2.

Table A2

Co. No.	FGFR 1	FGFR 2	FGFR 3	FGFR 4	VEGFR 2 (KDR)	BAF3-FGFR1 (MIN IL3)	BAF3-FGFR1 (PLUS IL3)	BAF3-FGFR3 (MIN IL3)	BAF3-FGFR3 (PLUS IL3)	BAF3-KDR (MIN IL3)	BAF3-KDR (PLUS IL3)	BAF3-FLT3 (MIN IL3)	BAF3_FLT3 (PLUS IL3)
	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}	pIC_{50}
200	8.53	8.11	8.73	7.92	7.05	6.41	<5	6.61	<5	5.79	<5	5.2	<5
11	8.36	7.91	8.66	7.76	7.43	6.3	<5	6.43	<5	6.06	<5	5.71	<5
202	8.11 8.18	7.71 7.9	~8.33 8.45	7.61 7.6	6.58 6.72	5.95 6.15	<5 <5	6.1 5.93	<5 <5	5.14 5.18	<5 <5	<5 <5	<5 <5
17	7.93	7.48	8.15	7.13	6.39	5.95	<5	5.67	<5	<5	<5	<5	<5
203	7.26	7.19	7.75	6.73	6.17	5.49	<5	5.54	<5	<5	<5	<5	<5
204	8.27	7.93	8.47	7.55	6.79	6.53	<5	6.64	<5	5.35	<5	<5	<5
205	8.85	8.46	9	8.55	7.07	6.42	<5	7.09	<5	5.52	<5	5.24	<5
206	8.15	7.91	8.52	7.46	6.67	5.79	<5	6.15	<5	<5	<5	<5	<5
207	8.86	8.32	8.64	7.45	7.18	5.59	<5	~5.56	~5.24	<5	<5	<5	<5
56	8.35 8.37	8.19 8.09	8.68 8.55	7.55 7.57	6.92 6.97	6.12 6.28	<5 <5	5.72 5.9	<5 <5	<5 5.06	<5 <5	<5 <5	<5 <5
208	7.35	7.21	7.26	6.8	5.82	5.73	<5	5.66	<5	<5	<5	<5	<5

209	5.92	~6.07	6.22	5.29	6.18	<5	<5	<5	<5	<5	<5	<5	<5
210	8.37	8.03	8.45	7.75	6.75	6.7	<5	6.53	<5	5.26	<5	<5	<5
211	8.29	7.88	8.13	7.39	6.7	6.44	<5	6.01	<5	5.26	<5	<5	<5
1	8.59 8.64	8.5 8.41	9.05 8.95	8.25 8.27	7.07 7.1	7.04 6.56	<5 <5	6.89 6.86	<5 <5	6.03 5.6	<5 <5	<5 5.09	<5 <5
212	8.43	8.15	8.86	7.83	6.92	6.36	<5	5.98	<5	~5.02	<5	<5	<5
213	8.33	8.09	8.55	7.74	6.65	6.45	<5	6.06	<5	<5	<5	<5	<5
214	7.16	7	7.39	6.33	5.74	<5	<5	<5	<5	<5	<5	<5	<5
216	8.86	8.28	8.58	7.66	6.79	5.69	<5	6.05	~5.02	~5.06	~5.09	~5.26	~5.2
22	5.99	6.07	6.12	<5	5.92	<5	<5	<5	<5	<5	<5	<5	<5
60	8.99	8.42	9.01	7.84	7.37	5.8	~5.19	6.15	~5.27	~5.27	~5.26	~5.27	~5.08
122	8.77	8.49	9.1	8.21	7.12	6.36	<5	6.87	<5	5.47	<5	<5	<5
58	6.17	6.39	6.78	<6	<6	<5	<5	~5.09	<5	<5	<5	<5	<5
3	8.71 8.77	8.63 8.49	9.03 8.97	8.29 8.25	7.75 7.39	7.24	<5	7.1	<5	5.91	<5	5.22	<5
218	8.42	~8.05	8.29	7.63	6.62	6.71	<5	6.61	<5	<5	<5	<5	<5
219	6.95	6.92	7.23	6.12	<6	5.05	<5	5.23	<5	<5	<5	<5	<5
220	7.27	7.01	7.53	6.59	6.48	5.44	<5	5.34	<5	<5	<5	<5	<5
221	6.63	6.62	7.08	<6	6.97	<5	<5	~5	<5	<5	<5	<5	<5
223	8.74	8.44	9.04	8.37	6.95	6.73	<5	7	<5	5.3	<5	<5	<5
224	7.84	7.83	8.2	7.51	7.19	5.64	<5	5.6	<5	5.37	<5	5.23	<5

62	8.9	8.45	9.06	7.92	7.55	5.88	~5.25	6.06	~5.25	~5.27	~5.27	~5.27	~5.26
23	8.73	8.54	9.15	8.35	6.91	7.02	<5	6.74	<5	5.28	<5	<5	<5
226	8.16	8.07	8.69	7.52	6.46	6	<5	6.11	<5	<5	<5	<5	<5
8	8.84	8.48	8.61	7.98	7.15	5.96	<5	7.12	<5	<5	<5	<5	<5
	8.84	8.82	8.49	7.95	7.55	6.07	<5	7.07	<5	<5	<5	<5	<5
84	8.76	8.55	8.75	8.1	7.43	6.57	<5	6.92	<5	5.63	<5	<5	<5
	8.67	8.8	8.73	8.03	7.56	7.29	<5	7.16	<5	5.96	<5	<5	<5
	8.76	8.61	8.71	8.12	7.44	6.75	<5	6.82	<5	5.73	<5	<5	<5
	8.66	8.71	8.73	7.95	7.43	6.47	<5	7.06	<5	5.75	<5	<5	<5
	8.74	8.48	8.62	7.8	7.3	6.58	<5	7.15	<5	5.57	<5	<5	<5
	8.92	8.47	8.69	7.88	7.08	6.1	<5	7.03	<5	5	<5	<5	<5
	8.96	8.56	8.74	8.06	7.46	6.08	<5	7.07	<5	<5	<5	<5	<5
	8.78	8.4	8.66	7.97	7.52	5.94	<5	7.11	<5	<5	<5	<5	<5
	8.81	8.64	8.64	7.98	7.27	6.87	<5	7.29	<5	5.73	<5	<5	<5
227	8.27	8.03	8.6	7.8	6.78	5.76	<5	6.57	<5	<5	<5	<5	<5
65	8.51	8.19	8.35	7.49	6.46	6	<5	6.56	<5	<5	<5	<5	<5
229	8.53	8.14	8.95	7.64	6.68	5.75	<5	6.39	<5	<5	<5	<5	<5
73	8.73	8.53	8.64	7.46	7.31	6.09	~5.07	6	~5.25	~5.23	~5.24	~5.25	<5
231	8.16	8.29	8.45	7.38	7	5.71	<5	6.23	<5	~5.08	<5	<5	<5
59	7.76	7.95	8.07	6.99	6.59	6.34	<5	6.55	<5	<5	<5	<5	<5
52	8.04	7.82	8.21	7.28	6.91	5.7	<5	6.17	<5	<5	<5	<5	<5
232	<6	<6	6.17	<6	<6	<5	<5	<5	<5	<5	<5	<5	<5

64	7.67	7.39	7.29	7.09	<6	6.17	<5	5.75	<5	<5	<6	<5
233	8.76	8.57	8.93	8.13	7.27	6.66	<5	6.97	<5	5.59	<5	<5
19	8.09	8.01	8.17	6.93	6.86	5.2	<5	~5.68	~5.23	<5	~5.18	<5
130	8.32	8.19	8.54	7.37	7.69	6.16	~5.25	6.06	~5.33	~5.62	~5.3	~5.26
235	8.78	8.53	8.68	7.42	7.26	5.72	<5	6.15	<5	<5	<5	<5
69	7.72 7.72	7.33 7.43	7.93 7.77	6.95 7.02	6.16 6.05	5.99 5.55	<5 <5	6.29 6.1	<5 <5	<5 <5	<5 <5	<5 <5
237	6.84	6.99	7.35	6.5	6.14	~5.16	<5	~5.17	<5	<5	<5	<5
238	9.01	8.57	8.88	7.76	7.54	6.2	<5	6.89	<5	5.32	<5	<5
66	6.54	6.96	7.22	6.18	<6	<5	<5	~5.04	<5	<5	<5	<5
2	~8.64 8.56	8.48 8.33	9.02 8.79	8.03 7.97	6.86 7.34	6.51 7.1	<5 <5	7.04 7.05	<5 <5	5.41 5.49	<5 <5	<5 <5
113	8.35	8.18	8.71	7.8	6.69	6.71	<5	6.95	<5	5.45	<5	<5
68	5.86	5.99	~6.06	<5	5.91	<5	<5	~5.11	~5.15	<5	<5	5.21
239	~8.66	8.5	8.76	7.33	7.14	5.94	<5	5.91	~5.28	5.25	~5.21	~5.26
240	8.7	8.26	8.75	7.17	7.01	6.08	<5	5.71	~5.26	~5.14	~5.26	~5.26
241	9.12 9.43	8.86 8.78	9.05 8.73	8.51 8.59	7.74 7.61	8.02 7.97	<5 <5	8.03 8.01	~5.06 ~5.08	6.69 6.65	<5 <5	~5.09 ~5.14
55	8.72	8.52	8.75	7.24	7.12	6.1	~5.16	5.95	~5.27	~5.24	~5.25	~5.26
242	8.62	8.61	8.9	8.01	7.44	6.96	<5	7	<5	5.77	<5	<5
243	8.66	8.36	8.89	7.61	7.04	6.01	<5	6.01	~5.25	~5.02	~5.2	~5.21
												~5.19

244	8.27	8.29	8.49	7	6.76	5.84	<5	5.77	~5.25	~5.05	~5.08	~5.24	<5
245	8.2	8.02	8.37	6.71	6.87	6.91	<5	6.82	<5	5.94	<5	<5	<5
246	8.22	8.09	8.65	7.28	6.47	5.69	<5	6.11	<5	<5	<5	<5	<5
24	7.88	7.94	8.52	7.32	6.82	5.97	<5	5.86	<5	~5.01	<5	<5	<5
249	8.45	8.31	8.65	7.09	7.36	6.55	<5	6.56	<5	5.64	<5	<5	<5
250	8.35	8.08	8.59	7.37	6.49	5.73	<5	6.09	<5	<5	<5	<5	<5
251	7.73	7.73	7.93	6.12	6.34	5.31	<5	~5.29	~5.26	~5.12	~5.23	~5.26	~5.37
21	8.75	8.48	9.07	8.07	6.79	6.75	<5	6.8	<5	5.56	<5	<5	<5
72	8.8	~8.5	8.91	7.33	7.22	6.1	~5.05	6.03	~5.29	~5.28	~5.26	~5.27	~5.19
253	8.54	8.33	8.74	7.6	6.78	5.92	<5	6.07	<5	<5	<5	<5	<5
254	8.41	8.19	8.52	7.52	6.35	6.13	<5	6.11	<5	<5	<5	<5	<5
74	7.78	7.9	8	7.07	<6	6.14	<5	6.42	<5	<5	<5	<5	<5
70a	8.57	8.52	8.23	7.58	7.23	6.91	<5	7.13	<5	5.77	<5	<5	<5
256	9.21	8.54	8.88	8.31	7.28	7	<5	6.89	<5	<5	<5	<5	<5
26	8.76	8.47	8.9	8.11	7.14	6.85	<5	6.76	<5	5.64	<5	5.22	<5
257	7.96	7.58	7.9	7.59	6.49	6.46	<5	6.57	<5	<5	<5	<5	<5
75	8.73 8.5	8.42 8.42	8.87 8.79	8.12 8.1	7.15 6.77	7.05 6.69	<5 <5	6.9 7.11	<5 <5	5.7 5.5	<5 <5	<5 <5	<5 <5
7	8.3	8.17	8.57	7.92	6.66	6.36	<5	6.58	<5	5.26	<5	<5	<5
259	8.81	8.63	9.1	8.4	7.28	7.24	<5	7.17	<5	5.45	<5	<5	<5
260	8.79	8.51	8.9	8.18	7.47	7.2	<5	6.88	<5	5.58	<5	<5	<5

27	6.13	6.56	6.3	<6	6.4	5.67	<5	<5	6.17	<5	5.32	<5
78	8.13	7.93	8.4	7.42	6.77	6.19	<5	6.12	5.24	<5	<5	<5
	8.22	8.11	8.36	7.45	6.96	6.63	<5	6.46	~5.68	<5	<5	<5
	8.1	8.03	8.43	7.52	6.68	6.26	<5	6.26	5.38	<5	<5	<5
261	8.71	8.5	8.39	8.14	7.38	5.96	<5	7.07	<5	<5	<5	<5
262	9.29		9.35	8.91	7.86	7.74	<5	7.87	6.75	<5	<5	<5
25	9.11	8.64	8.43	7.92	7.04	7.36	<5	7.22	5.73	<5	<5	<5
	8.78	8.61	8.52	7.78	7.11	7.35	<5	7.36	5.86	<5	<5	<5
263	8.75	8.52	8.75	8.14	7.39	6.72	<5	6.75	5.54	<5	<5	<5
264	8.97	8.75	8.45	8.09	7.78	7.47	<5	7.58	6.12	<5	<5	<5
265	8.79	8.59	8.44	8.04	7.05	6.51	<5	6.65	5.24	<5	<5	<5
266	8.76	8.62	8.33	7.97	7.37	5.79	<5	6.9	<5	<5	<5	<5
76	8.46	8.19	8.43	7.65	7.05	7.18	5	6.92	6.08	5.02	5.09	<5
267	7.83	7.7	8	7.43	6.2	5.69	<5	5.76	<5	<5	<5	<5
268	9.43	8.74	9.14	9.05¶	7.96	7.74	<5	8.09	6.69	<5	<5	<5
	9.25	8.75	9.41	8.78	7.96	7.86	<5	7.89	6.69	<5	<5	<5
269	6.28	6.44	6.51	<6	<6	<5	<5	<5	<5	<5	<5	<5
271	8.14	7.87	8.58	7.61	6.75	6.14	<5	6.19	~5.06	<5	<5	<5
272	9.02	8.75	8.82	8.37	7.6	7.36	<5	7.77	6.24	<5	~5.18	<5
273	7.11	7.09	6.71	<6	6.29	5.73	<5	5.68	5.29	<5	<5	<5
274	6.1	6.27	6.65	<6	~6	<5	<5	<5	<5	<5	<5	<5

121	8.12	7.86	8.2	7.02	6.63	6.14	<5	~6.06	<5	5.55	<5	5.35	<5
4	9.01	8.35	8.42	8.21	7.32	7.23	<5	7.31	<5	5.48	<5	~5.05	<5
	8.77	8.81	8.49	8.18	7.43	7.08	<5	7.82	<5	6.05	<5	<5	<5
	8.73	8.76	8.5	8.23	7.46	7.72	<5	7.86	<5	6.11	<5	5	<5
	8.96	8.73	8.57	8.31	7.46	7.6	<5	8.05	<5	5.96	<5	<5	<5
	8.92	8.67	8.44	8.25	7.33	8.07	<5	8.07	<5	6.48	<5	<5	<5
	9.09	8.58	8.6	8.29	7.45	7.61	<5	8.13	<5	6.04	<5	<5	<5
	9.01	8.64	8.56	8.33	7.5	7.61	<5	7.98	~5.12	6	<5	<5	<5
	8.99	8.56	8.6	8.18	7.58	7.89	<5	7.8	<5	6.24	<5	<5	<5
276	8.6	8.46	8.93	8.17	7.23	6.99	<5	~7.17	<5	5.65	<5	<5	<5
28	8.62	8.47	8.83	8.09	7.35	7.15	<5	7.14	<5	5.58	<5	<5	<5
37	8.04	7.99	8.62	7.59	7.14	5.88	~5.07	5.82	<5	~5.1	<5	<5	<5
278	8.46	8.34	8.44	7.66	7.18	5.92	<5	7.09	<5	<5	<5	<5	<5
279	8.43	8.3	8.4	7.5	7.16	5.76	<5	7.04	<5	<5	<5	<5	<5
280	8.66	8.39	8.44	7.87	7.31	6.94	<5	7.49	<5	5.34	<5	<5	<5
281	8.41	8.36	8.18	7.59	7.14	7.33	<5	7.1	<5	5.94	<5	~5.15	<5
283	6.39	6.65	6.68	<6	6.31	5.23	<5	~5.21	<5	~5.18	<5	<5	<5
79/79a	8.62	8.55	8.58	7.82	6.91	7.29	<5	7.58	<5	5.8	<5	<5	<5
602	8.06	8.14	8.13	7.51	7	6.57	<5	6.6	<5	5.28	<5	<5	<5
81	8.21	8.16	8.84	7.48	7.02	6.61	<5	6.43	<5	5.24	<5	<5	<5
80	8.33	8.13	8.72	7.59	6.86	6.69	<5	6.59	<5	5.31	<5	<5	<5

284	9.07	8.8	8.68	8.4	7.39	7.69	<5	7.82	~5.08	5.72	<5	<5
	9.26	8.8	8.8	8.33	7.73	7.56	<5	7.77	~5.01	6.17	<5	~5.1	<5
286	8.75	8.49	8.56	8.14	7.68	7.24	<5	7.32	5.03	5.86	<5	<5	<5
287	8.33	8.3	8.63	7.8	7.69	6.38	<5	6.75	<5	<5	<5	<5	<5
29	~6.1	6.1	6.32	<6	~5.96	<5	<5	5.24	5.28	<5	<5	5.22	<5
288	7.65	~7.77	7.73	7.04	6.76	5.15	<5	6.58	<5	<5	<5	<5	<5
291	8.87	8.71	8.77	8.39	7.97	7.72	<5	7.79	<5	6.65	<5	<5	<5
293	7.01	6.99	7.49	6.49	6.23	<5	<5	<5	<5	<5	<5	<5	<5
294	8.67	8.4	8.64	7.69	7.28	5.78	<5	6.9	<5	<5	<5	<5	<5
	8.62	8.6	8.52	7.75	7.07	6.24	<5	6.94	<5	5.4	<5	<5	<5
16	7.81	7.78	8.34	7.39	6.53	6.17	<5	5.78	<5	<5	<5	<5	<5
20	8.25	8.31	8.84	7.85	7.16	6.07	<5	6.06	<5	~5	<5	<5	<5
295	8.86	8.61	8.75	8.14	7.6	7.97	<5	7.79	<5	6.32	<5	<5	<5
296	8.61	8.53	8.46	8	6.87	7.47	<5	7.65	<5	5.96	<5	<5	<5
54	6.16	6.07	6.11	<6	~6.04	<5	<5	<5	<5	<5	<5	<5	<5
297	8.19	8.35	8.5	7.66	7.24	5.35	<5	5.56	<5	<5	<5	<5	<5
298	8.42	8.45	8.18	7.65	6.72	6.71	<5	6.37	<5	5.21	<5	<5	<5
299	8.06	8.23	8.48	7.72	7.31	6.81	<5	6.88	<5	5.74	<5	<5	<5
49	8.93	8.34	9.19	8.33	7.53	7.13	<5	7.06	<5	5.58	<5	5.43	<5
93	8.95	8.66	8.86	8.23	7.65	6.98	<5	6.97	<5	5.81	<5	<5	<5
	8.85	8.46	8.64	8.04	7.58	6.8	<5	6.42	<5	5.47	<5	~5.01	<5

302	8.97	8.72	8.68	8.28	7.59	7.93	<5	7.91	<5	6.64	<5	<5	<5
303	8.43	8.48	8.33	7.66	7.92	6.73	<5	6.7	~5.01	6.43	<5	<5	<5
304	8.15	7.82	8.13	7.33	7.13	5.74	<5	7.08	<5	<5	<5	<5	<5
305						6.38	<5	6.67	<5	5.07	<5	<5	<5
61						5.83	~5.19	~5.67	~5.28	~5.15	~5.25	5.12	~5.12
306						5.74	<5	5.64	<5	<5	<5	<5	<5
307						5.74	<5	5.63	<5	<5	<5	<5	<5
308						6.1	<5	6.87	<5	<5	<5	6.84	<5
311						6.38	<5	6.45	<5	5.38	<5	<5	<5
312						5.76	<5	5.86	<5	<5	<5	<5	<5
313	8.67	8.53	9.01	8.27	7.08	6.98 6.56	<5 <5	7.1 7.09	<5 <5	5.65 5.56	<5 <5	<5 5.09	<5 <5
314						5.92	<5	6.05	<5	<5	<5	<5	<5
30						6.71	<5	6.32	<5	5.42	<5	5.21	<5
38	8.69	8.21	8.55	7.78	7.06	5.93	<5	6.5	<5	5.14	<5	<5	<5
315						6.6	<5	6.07	<5	5.34	<5	<5	<5
48						7.43	<5	7.53	<5	6.45	<5	<5	<5
316						7.44	<5	7.48	<5	6.35	<5	<5	<5
321	8.85	8.7	8.68	7.99	7.72	6.6	<5	6.92	<5	5.81	<5	<5	<5
57	7.97	7.84	8.42	7.35	6.54	5.92	<5	6.31	<5	<5	<5	<5	<5
323	6.58	6.78	6.84	<6	<6	<5	<5	<5	<5	<5	<5	<5	<5

85	7.78	7.84	8.18	7.04	6.59	5.56	<5	6.14	<5	<5	<5	<5	<5
324	8.81	8.65	8.5	7.9	7	7.12	<5	7.46	<5	5.63	<5	<5	<5
325	7.57	7.7	7.3	6.51	6.12	5.79	<5	6.26	<5	<5	<5	<5	<5
86	8.39	8.13	8.58	7.85	7.1	<5	<5	<5	<5	<5	<5	<5	<5
326	8.04	8.05	8.48	7.46	6.79	6.18	<5	6.39	<5	5.12	<5	<5	<5
83	8.47	8.45	7.97	7.35	6.98	6.68	<5	6.86	<5	5.63	<5	<5	<5
327	6.88	7.11	7.52	6.51	6.16	<5	<5	5	<5	<5	<5	<5	<5
328	9.14	8.71	8.78	8.29	7.7	6.94	<5	7.13	5.11	5.76	<5	~5.12	<5
330	8.85	8.65	8.51	7.75	7.47	6.49	<5	6.7	~5.08	~5.56	~5.08	~5.19	<5
332	8.82	8.67	8.52	7.85	7.33	7.27	<5	7.57	<5	6.04	<5	<5	<5
333	8.47	8.39	8.42	7.86	7.15	5.68	<5	6.13	<5	<5	<5	<5	<5
334	7.37	7.2	7.29	6.61	<6	<5	<5	<5	<5	<5	<5	<5	<5
335	9.06	8.84	9.23	8.68	7.67	7.13	<5	7.6	<5	6.09	<5	<5	<5
336	8.51	8.5	8.7	7.88	6.86	6.36	<5	6.85	<5	5.37	<5	<5	<5
338	9.21	~8.93	8.89	8.16	7.43	6.52	<5	7.12	<5	5.53	<5	<5	<5
339	8.24	8.19	8.04	7.16	7.17	5.92	<5	6.11	<5	<5	<5	<5	<5
32	6.56	6.4	6.61	<6	<6	<5	<5	5.37	<5	<5	<5	<5	<5
131	7.94	8.01	8.47	7.35	6.71	6.1	<5	6.31	<5	5.07	<5	<5	<5
340	8.05	8.05	8.39	7.3	6.6	6.54	<5	7.26	<5	5.22	<5	<5	<5
342	8.91	8.6	8.43	7.93	7.26	7.31	<5	7.59	<5	5.83	<5	<5	<5
343	8.55	8.47	8.42	7.6	7.02	6.52	<5	6.56	<5	5.28	<5	<5	<5

344	8.68	8.47	8.18	7.35	7.24	6.9	<5	7.22	<5	5.95	<5	6.12	<5
31	9.01	8.51	9.21	8.42	7.24	6.85	<5	7.41	<5	5.71	<5	5.32	<5
	8.86	8.61	9.06	8.51	7.34	6.96	<5	7.29	<5	5.87	<5	5.53	<5
345	8.38	8.41	8.73	7.63	7.06	5.82	<5	6.32	~5.02	~5.01	<5	~5.01	<5
346	~6.64	6.55	6.45	6.59	<6	5.17	<5	5.68	<5	<5	<5	<5	<5
347	8.85	8.67	8.64	8.24	7.2	7.46	<5	7.99	<5	6.09	<5	~5.12	<5
348	8.07	7.91	8.04	7.37	6.61	6.21	<5	6.61	<5	5.21	<5	<5	<5
351	8.34	8.29	7.8	6.98	7.04	5.93	<5	6.59	<5	~5.38	<5	<5	<5
89	8.43	8.35	8.52	7.6	7.01	6.33	<5	6.82	<5	5.31	<5	<5	<5
352	8.98	8.54	8.64	8.29	7.37	7.5	<5	7.89	<5	5.89	<5	<5	<5
87	9.01	8.73	8.88	8.46	7.62	7.51	<5	7.89	<5	6.39	<5	5.07	<5
353	7.68	7.73	8.07	7.06	6.33	5.64	<5	6.2	<5	<5	<5	<5	<5
354	8.51	8.08	8.43	7.7	6.57	6.2	<5.52	6.68	<5.52	<5.52	<5.52	<5.52	<5.52
355	7.49	7.21	7.19	6.21	6.23	5.19	<5	5.55	<5	<5	<5	<5	<5
356	~8.07	8.06	8.23	7.49	6.83	6.2	<5	6.56	<5	5.22	<5	<5	<5
53	6.45	6.47	6.38	~6.15	6.34	<5	<5	<5	<5	<5	<5	<5	<5
91	7.91	7.71	8.25	7.47	6.59	<5	<5	~5.27	<5	<5	<5	<5	<5
18	7.3	6.9	6.98	6.58	6.64	5.44	<5	6.07	<5	<5	<5	<5	<5
357	8.74	8.37	8.38	7.7	7.11	6.91	<5	7.38	<5	5.69	<5	<5	<5
358	6.83	7.01	6.97	6.64	<6	<5	<5	5.12	<5	<5	<5	<5	<5
359	7.65	7.57	7.66	7.12	6.14	5.69	<5	6.15	<5	<5	<5	<5	<5

360	7.87	7.67	7.89	6.9	6.16	5.5	<5	5.98	<5	<5	<5	<5	<5
361	7.47	7.42	7.27	6.15	6.1	5.67	<5	6.05	<5	5.17	<5	5.25	<5
362	8.62	8.26	8.32	7.78	7.09	6.6	<5	7.17	<5	5.54	<5	<5	<5
363	8.92	8.71	8.95	8.36	7.51	6.52	<5	6.7	<5	5.28	<5	<5	<5
	9	8.75	8.84	8.35	7.33	5.8	<5	6.57	<5	5.07	<5	<5	<5
364	8.41	8.19	8.23	7.59	7.69	5.94	<5	6.45	<5	5.8	<5	<5	<5
365	8.64	8.41	8.57	7.97	7.52	6.29	<5	6.94	<5	5.74	<5	<5	<5
366	8.94	8.6	8.62	8.18	7.31	7.23	<5	7.92	<5	5.95	<5	<5	<5
367	9.26	8.86	8.94	8.54	7.47	7.17	<5	7.63	<5	6.02	<5	<5	<5
300	8.15	7.99	8.38	7.9	7.01	>8	<5	>8	<5	5.29	<5	5.09	<5
	8.14	8.46	8.41	7.87	7.2	7.9	5.28	8.18	~5.12	5.56	5.1	5.33	5.17
	8.35	8.41	8.59	7.84	7.02	6.92	<5	7.81	<5	5.21	<5	5.11	<5
368	~8.96	8.7	8.81	8.09	7.23	5.93	<5	6.41	<5	5.14	<5	<5	<5
369	7.76	7.93	7.5	6.73	7.63	5.2	<5	5.57	<5	<5	<5	<5	<5
370	8.68	8.57	8.61	8	6.86	5.95	<5	6.51	<5	<5	<5	<5	<5
371	8.8	8.68	8.42	8.17	7.24	6.04	~5.03	6.64	~5.17	5.58	5.13	~5.22	~5.13
92	9.09	8.57	8.99	8.74	7.64	6.65	<5	7.72	<5	5.79	<5	<5	<5
	8.96	8.6	8.97	8.73	7.81	7.39	<5	7.55	<5	6.16	<5	<5	<5
	9.04	8.45	8.99	8.58	7.65	7.74	<5	7.83	<5	6.23	<5	<5	<5
373	7.96	7.97	7.97	7.26	6.62	5.63	<5	6.1	<5	<5	<5	<5	<5
374						6.55	<5	6.96	<5	5.15	<5	<5	<5

375	8.63	8.59	9.06	8.19	6.99	6.88 7.13	<5 <5	7.36 7.17	<5 <5	5.3 5.84	<5 <5	<5 <5
376						6.92	<5	7.32	<5	5.91	<5	<5
377						6.26	<5	6.75	<5	5.41	<5	<5
378	8.46	8.41	8.19	7.66	6.86	7.18 7.43	<5 <5	7.37 7.57	<5 <5	5.43 5.83	<5 <5	<5 <5
379	8.23	8.52	8.25	7.51	7.22	6.13 6.48	<5 <5	6.49 6.59	~5.19 5.1	5.14 ~5.49	<5 ~5.08	<5 ~5.16
380						6.69	<5	6.85	<5	5.32	<5	<5
381						6.81	<5	7.14	<5	5.59	<5	<5
382	8.96	8.69	8.67	8.05	6.93	7.29 7.5	<5 <5	7.63 7.69	~5.05 <5	5.56 5.79	<5 <5	<5 <5
132	8.12	8.54	8.43	7.73	7.46	6.26 6.84	<5 <5	6.75 6.65	<5 <5	5.4 5.68	<5 <5	<5 <5
108	9	8.86	8.74	8.19	7.5	7.67 7.45	<5 <5	7.68 7.59	<5 <5	6.15 5.98	<5 <5	<5 <5
384						7.21	<5	7.51	<5	5.51	<5	<5
385						5.49	<5	5.81	<5	<5	<5	<5
39						6.56	<5	6.8	<5	5.23	<5	<5
40						6.25	<5	6.47	<5	5.19	<5	<5
133	8.52	8.28	8.73	7.94	7.08	6.82 6.74	<5 <5	7.06 7.12	<5 <5	5.62 5.64	<5 <5	<5 <5

386					~5.56	<5	5.79	~5.11	<5	5.05	<5	<5
387					5.25	<5	5.57	<5	<5	<5	<5	<5
389					6.67	<5	6.98	<5	5.23	<5	<5	<5
390					7.17	<5	7.25	<5	5.62	<5	<5	<5
391					5.8	<5	6.13	<5	<5	<5	<5	<5
392	7.91	7.77	8.19	7.14	6.9	5.95	<5	6.05	<5	5.34	<5	5.37	<5
395	9.35	8.81	8.87	8.72	8.08	7.78	<5	7.88	<5	6.75	<5	<5	<5
94	8.76	8.29	9.05	8.2	7.1	6.41	<5	6.74	<5	5.31	<5	<5	<5
	8.47	8.47	8.85	8.19	7.07	6.9	<5	7.1	<5	5.62	<5	<5	<5
396	8.66	8.45	8.23	8.05	7.59	7.78	<5	8.02	~5.16	6.62	<5	5.08	<5
397	8.43	8.36	8.96	7.98	6.87	6.75	<5	~7.07	<5	5.26	<5	<5	<5
398	8.85	8.59	8.67	8	7.75	6.44	<5	6.54	~5.25	5.95	<5	~5.16	<5
399	8.58	8.37	9.05	8.24	7.08	6.59	<5.52	6.87	<5	5.65	<5	5.1	<5
400	8.55	8.46	9.01	8.15	7.14	6.55	<5	6.5	<5	5.67	<5	5.1	<5
401	7.75	7.78	8.31	7.14	6.76	5.75	<5	5.93	<5	5.33	<5	<5	<5
405	7.2	6.89	7.08	6.59	<6	5.62	<5	5.79	<5	<5	<5	<5	<5
407	7.21	6.84	7.25	6.35	6.11	<5	<5	~5.08	<5	<5	<5	<5	<5
408	6.41	6.74	6.49	<6	~6	<6	<5	6.38	<5	<5	<5	<5	<5
409	8.84	8.55	8.58	8.11	6.57	6.82	<5	6.78	<5	5.47	<5	<5	<5
411	6.62	6.75	6.75	6	6.13	<5	<5	<5	<5	<5	<5	<5	<5
412	8.14	8.36	8.1	7.42	6.93	6.35	<5	6.03	<5	5.31	<5	<5	<5

413	8.74	8.5	8.37	8.17	7.76	7.79	<5	7.94	5.09	6.4	<5	5.11	<5
414	8.54	8.43	8.27	7.64	7.17	6.65	<5	6.57	~5.15	5.89	<5	~5.98	<5
422	8.72	8.42	8.74	8.12	7.49	6.67	<5	7.29	<5	5.84	<5	<5	<5
423	8.73	8.46	8.81	8.18	7.42	7.4	<5	7.65	<5	6.07	<5	~5.07	<5
426	6.94	6.8	6.69	<6	<6	5.43	<5	5.59	<5	<5	<5	~5.05	<5
428	8.52	8.46	8.38	7.17	7.73	5.99	<5	5.8	<5	5.25	<5	<5	<5
34	8.7	8.4	8.54	7.93	7.04	5.88	<5	5.81	<5	<5	<5	<5	<5
430	8.15	8	8.32	7.7	7.26	6.13	<5	6.26	<5	5.25	<5	<5	<5
125	8.49	8.31	8.5	7.84	6.87	6.38	<5	6.38	<5	5.49	<5	<5	<5
431	7.9	7.83	8.01	7.49	6.78	5.8	<5	6.17	<5	~5.13	<5	<5	<5
432	8.91	8.55	8.52	8.22	7.26	7.04	<5	7.32	<5	5.81	<5	<5	<5
433	7.94	7.78	7.99	7.51	7.18	5.67	<5	5.63	<5	5.24	<5	5.1	<5
434	8.66	8.4	8.94	8.33	7.07	6.49	<5	7.03	<5	5.49	<5	5.09	<5
435	8.9	8.73	8.71	8.15	6.9	6.66	<5	6.63	~5.19	5.51	<5	5.28	<5
5	8.14	7.89	8.12	7.6	6.87	~6.13	<5	5.49	<5	<5	<5	<5	<5
	8.12	7.96	8.2	7.82	7.01	6.32	<5	~6.61	<5	5.06	<5	<5	<5
	8.14	7.89	8.31	7.74	7.1	6.33	<5	6.51	<5	5.11	<5	<5	<5
	8.04	8.3	8.54	7.81	6.92	6.45	<5	6.58	<5	5.43	<5	<5	<5
	8.02	8.2	8.66	7.67	6.96	6.08	<5	6.4	<5	5.32	<5	<5	<5
	8.1	8.26	8.59	7.6	7	6.22	<5	6.45	<5	5.37	<5	<5	<5
	7.91	7.94	8.08	7.58	6.66	5.87	<5	6.62	<5	5.23	<5	<5	<5
	8.22	8.26	8.45	7.81	6.86	6.24	<5	6.62	<5	5.29	<5	<5	<5

436	8.93	8.68	8.88	8.19	7.56	6.7	<5	6.72	<5	5.9	<5	<5	<5
95/95 a	8.77	8.55	8.77	8.52	7.68	6.83	<5	6.75	<5	5.97	<5	<5	<5
437	6.81	6.87	6.8	<6	<6	5.31	<5	5.35	<5	<5	~5.66	<5	<5
438	6.91	6.71	6.4	<6	<6	5.26	<5	5.41	<5	<5	<5	<5	<5
440	8.56	8.32	8.07	7.36	7.09	6.62	<5	6.57	~5.02	5.67	<5	5.93	<5
441	8.65	8.46	8.31	8.27	7.75	8.19	<5	8.39	~5.11	6.96	<5	5.1	<5
442	8.79	8.52	8.41	7.98	7.06	7.37	<5	7.37	<5	5.76	~5.61	<5	<5
443	9.06	8.64	8.54	8.32	7.33	7.53	<5	7.86	~5.13	5.89	<5	<5	<5
444	9.04	8.5	8.56	8.36	7.33	7.62	<5	7.85	~5.1	5.96	<5	<5	<5
445	8.58	8.32	8.8	8.22	6.76	6.53	<5	6.85	<5	5.46	<5	<5	<5
447	8.74	8.33	8.92	8.49	6.88	7.08	<5	7.57	<5	5.65	<5	<5	<5
448	8.8	8.41	8.64	7.96	7.31	6.38	<5	6.37	<5	5.57	<5	~5.11	<5
449	8.61	8.45	8.37	7.84	7.2	6.77	<5	6.86	<5	5.63	<5	<5	<5
450	8.69	8.54	8.53	7.97	7.26	7.07	<5	7.21	<5	5.6	<5	<5	<5
452	8.54	8.19	7.9	7.81	7.44	7.81	<5	7.94	~5.16	6.4	<5	5.06	<5
453	7.98	7.74	8	6.88	6.64	5.68	<5	5.76	<5	<5	<5	<5	<5
134	8.72 8.63	8.47 8.55	8.31 8.45	8.08 8.21	6.98 6.98	7.66 7.79	<5 <5	7.69 7.95	<5 <5	5.39 5.76	<5 <5	<5 <5	<5 <5
454	8.77	8.43	8.67	8.17	7.35	6.74	<5	7.19	<5	5.83	<5	<5	<5
455	8.54	8.34	8.28	8.1	7.21	7.1	<5	7.22	<5	5.74	<5	<5	<5

456	8.74	8.47	8.5	8.05	7.13	6.44	<5	6.6	~5.22	5.69	<5	<5	<5
96	7.6	7.7	7.51	6.74	6.32	5.77	<5	5.9	~5.21	~5.06	<5	~5	<5
457	9.08	8.58	8.85	8.07	7.3	6.19	<5	6.31	~5.25	5.46	<5	~5.11	~5.05
458	8.44	8.31	8.55	7.89	6.97	6.79	<5	6.81	<5	5.28	<5	<5	<5
44/44a	8.1	7.98	7.84	7.51	6.51	6.06	<5	6.09	<5	5.04	<5	<5	<5
459	8.24	8.06	8.18	7.54	6.76	6.27	<5	6.84	<5	5.15	<5	<5	<5
460	7.76	8.04	8.17	6.94	7.05	6.7	<5	~6.69	<5	5.53	<5	<5	<5
463			8.54	7.65		7.42	<5			6.04	<5	<5	<5
464	8.58	8.65	8.57	8.4	7.88	6.86	<5	7.34	<5	5.71	<5	<5	<5
465	8.31	8.36	8.38	7.4	7.06	6.34	<5	6.73	~5.15	5.35	<5	<5	<5
467	9.05	8.69	8.77	8.33	7.43	7.76	<5	7.95	<5	5.77	<5	<5	<5
468	8.02	7.71	7.72	7.16	6.61	5.52	<5	5.61	<5	~5.04	<5	<5	<5
469	9.23	8.76	8.77	8.52	7.4	8.12	<5	8.08	<5	5.8	<5	<5	<5
470	8.75	8.56	8.55	7.83	7.1	6.6	<5	7.03	<5	5.38	<5	<5	<5
472	8.69	8.61	8.63	8.08	7.58	7.32	<5	7.62	<5	5.57	<5	<5	<5
474	8.39	8.48	8.37	7.75	7.9	5.8	~5.56	6.04	~5.5	~5.62	~5.29	~5.37	~5.38
135	9.11	8.75	8.86	8.47	7.74	7.79	<5	7.83	~5.11	5.82	<5	<5	<5
475	~7.88	7.83	8.35	7.24	6.85	6.08	<5	6.06	<5	5.09	<5	<5	<5
476	8.32	8.6	9.15	8.37	7.57	6.85	<5	7.02	<5	5.46	<5	5.53	<5
477	8.06	7.96	8.39	7.46	6.74	5.92	<5	5.98	<5	<5	<5	<5	<5

478	7.91	7.97	8.06	7.01	7.02	5.44	<5	5.7	<5	5.04	<5	<5	<5
136	8.82	8.66	8.64	8.36	7.63	8.02	<5	8.06	~5.11	5.88	<5	~5.06	<5
480	8.9	8.64	8.68	8	7.57	6.7	~5.24	6.78	~5.36	5.89	~5.1	~5.3	~5.25
97	8.38	8.51	8.94	7.86	6.95	6.96	<5	7	<5	5.62	<5	5.06	<5
	8.13	8.19	8.75	7.9	6.94	6.38	<5	6.8	<5	5.12	<5	<5	<5
98	8.88	8.56	9.03	8.32	7.21	7.01	<5	7.25	<5	5.28	<5	<5	<5
	8.64	8.62	8.96	8.24	6.79	7.13	<5	7.59	<5	5.82	<5	<5	<5
481	8.64	8.49	9.07	8.42	6.92	6.89	<5	7.3	<5	5.27	<5	<5	<5
482	8.95	8.6	8.79	8	7.59	6.49	<5	6.45	<5	5.59	<5	<5	<5
483	8.82	8.74	8.88	8.32	7.69	6.41	<5	6.82	<5	5.68	<5	<5	<5
485	8.25	8.16	7.97	7.3	7.75	7.33	<5	7.39	<5	5.91	<5	5.13	<5
486	8.57	8.45	8.38	7.7	7.01	6.34	<5	6.71	<5	5.17	<5	<5	<5
487	8.35	8.36	8.29	7.62	7.48	6.35	<5	6.4	~5.25	~5.56	<5	~5.14	~5
488	7.99	8.12	8.02	7.26	6.73	6.11	<5	6.5	<5	5.32	<5	<5	<5
	7.92	7.88	8.01	7.41	6.65	5.7	<5	6.58	<5	5.13	<5	<5	<5
489	8.9	8.55	8.79	8.11	7.54	7.42	<5	7.9	<5	6.34	<5	~5.1	<5
490	8.62	8.65	8.5	7.88	6.9	7.02	<5	7.26	<5	5.78	<5	<5	<5
493	8.67	8.53	8.57	7.95	6.99	7.27	<5	7.35	<5	5.79	<5	5.25	<5
494	9.21	8.64	8.99	8.26	7.69	7.11	<5	7.22	<5	6	<5	~5.11	<5
495	8.17	8.35	8.24	7.58	6.93	6.7	<5	6.9	<5	5.47	<5	<5	<5
137	8.79	8.47	8.37	8.12	6.98	7.57	<5	7.74	<5	5.88	<5	<5	<5

119	7.8	7.89	7.64	6.92	6.13	5.71	<5	5.94	<5	<5	<5	<5
498	8.33	8.3	8.42	6.85	7.05	6.73	<5	7.06	<5	5.65	<5	5.01	<5
99	9.07	8.72	9.09	8.95	7.02	7.54	<5	8.12	<5	5.63	<5	<5	<5
	8.99	8.79	9.19	8.79	7.29	7.73	<5	8.11	<5	5.59	<5	<5	<5
138	8.5	8.32	8.56	7.06	7.34	6.92	<5	6.98	<5	6.16	<5	5.39	<5
	8.34	8.3	8.52	7.07	7.16	6.64	<5	7.11	<5	5.81	<5	5.11	<5
	8.53	8.5	8.6	7.13	7.49	6.95	~5.18	6.83	5.25	6.21	5.29	5.61	5.17
499	8.74	8.4	8.56	8.13	7.5	7.39	<5	7.7	<5	6.09	<5	~5.16	<5
500	8.73	8.55	8.58	7.36	7.66	7.06	<5	7.24	<5	6.29	<5	<5	<5
	8.71	8.15	8.62	7.53	7.71								
501	8.7	8.75	8.56	7.95	7.12	7.19	<5	7.08	<5	5.88	<5	<5	<5
502	8.76	8.75	8.62	8.06	7.2	7.44	<5	7.49	<5	5.96	<5	<5	<5
503	8.91	8.74	8.96	8.65	7.83	7.83	<5	7.9	<5	6.32	<5	<5	<5
139	8.18	8.28	8.39	7.87	6.86	7.02	<5	6.97	<5	5.29	<5	<5	<5
504	9.06	8.9	8.93	8.27	7.39	7.01	<5	6.62	<5	5.15	<5	<5	<5
	8.99	8.82	8.83	8.1	7.39	6.95	<5	6.71	<5	5.16	<5	<5	<5
140	8.22	8.33	8.5	8.07	7.12	6.42	<5	7.04	<5	5.5	<5	<5	<5
	8.59	8.68	8.71	8.16	7.48	6.82	<5	7.16	<5	5.53	<5	<5	<5
506	7.55	7.91	7.87	6.92	6.7	6.13	<5	6.27	<5	5.37	<5	<5	<5
507	8.1	8.22	8.22	7.32	6.66	6.51	<5	6.41	<5	5.22	<5	<5	<5
508	7.56	7.89	7.96	6.97	7.07	6.44	<5	6.19	<5	5.31	<5	<5	<5

509			8.57	8.43		>8	<5			7.05	<5	5.31	<5
510	8.25	8.34	8.1	7.34	6.89	6.95	<5	6.76	<5	5.53	<5	<5	<5
141	9.01	8.82	9.22	8.42	7.63	7.49	<5	7.58	<5	6.16	<5	<5	<5
	8.9	8.45	8.92	8.37	7.49	6.86	<5	6.99	<5	5.68	<5	<5	<5
511	8.08	8.13	8	7.55	7.46	7.71	<5	8.1	<5	6.94	<5	5.68	<5
512	8.6	8.62	8.52	7.88	7.13	7.72	<5	7.75	<5	5.73	<5	<5	<5
513	8.86	8.7	9.08	8.17	7.27	7.12	<5	7.15	<5	5.68	<5	<5	<5
514	8.31	8.56	8.45	7.81	7.51	6.91	<5	7	<5	5.94	<5	<5	<5
515	7.77	7.94	7.47	6.73	6.63	6.13	~5.24	6.34	5.29	~5.45	5.19	~5.33	~5.09
35	7.3	7.45	7.32	6.39	6.21	5.72	<5	5.64	<5	5.08	<5	5.41	<5
517	7.95	8.22	8.32	7.54	7.34	6.77	<5	7.01	<5	5.89	<5	<5	<5
518	8.37	8.55	8.2	7.43	7.2	7.02	<5	7.58	5.2	6.08	<5	5.25	<5
519	8.74	8.62	8.62	8.1	7.07	7.5	<5	7.3	<5	5.74	<5	<5	<5
102	8.53	8.47	8.52	7.78	7.24	7.07	<5	7.37	<5	5.93	<5	<5	<5
521	8.76	8.77	8.56	8.42	7.74	8.17	<5	8.19	<5	6.3	<5	5.01	<5
522	8.71	8.51	8.51	7.93	6.98	7.52	<5	7.58	<5	~6.02	<5	<5	<5
523	8.14	8.42	8.44	8.08	6.68	6.6	<5	6.84	<5	5.22	<5	<5	<5
142	8.36	8.39	8.53	7.97	6.75	6.7	<5	6.94	<5	5.48	<5	<5	<5
524	8.75	8.55	8.71	8.11	7.61	6.85	<5	7.2	<5	5.75	<5	<5	<5
525	8.03	8.43	8.49	7.72	6.64	6.73	<5	7.05	<5	5.54	<5	<5	<5
526	8.29	8.47	8.32	7.76	7.07	7.42	<5	7.48	<5	5.86	<5	<5	<5

527	8.18	8.26	8.42	7.77	7.69	6.32	<5	6.67	~5.01	5.94	~5.02	5.11	<5
143	8.01	8.21	7.95	7.3	7.03	6.66	<5	7.02	<5	5.75	<5	<5	<5
528	8.44	8.58	8.24	7.54	7.44	6.89	<5	7.18	<5	6.19	<5	<5	<5
529	8.64	8.52	8.47	8.06	7.35	7.27	<5	7.51	5.13	6.07	~5	5.12	<5
530	8.44	8.59	8.43	7.62	7.46	6.92	<5	7.34	5.21	6.34	<5	5.1	<5
10	8.21	8.12	8.21	7.96	6.43	6.97	<5	7.13	<5	<5	<5	<5	<5
531	8.57	8.67	8.65	7.97	7.34	7.36	<5	7.68	<5	6.24	<5	<5	<5
36	7.97	8.29	8.66	7.6	6.84	6.35	<5	6.59	<5	5.52	<5	5.13	<5
532	7.28	7.36	7.22	6.47	6.24	5.82	<5	6.09	<5	<5	<5	<5	<5
533	8.76	8.65	8.49	8.01	7.34	7.37	<5	7.63	<5	6.31	<5	<5	<5
534	8.63	8.66	8.46	7.92	7.1	7.41	<5	7.74	<5	6.16	<5	<5	<5
535	8.81	8.7	8.4	8.18	7.43	7.48	<5	7.99	5.15	6.35	<5	<5	<5
536	7.03	7.33	7.7	6.76	<6	5.66	<5	5.85	<5	<5	<5	<5	<5
537	8.16	8.14	7.93	7.23	6.95	6.2	<5	6.18	<5	5.33	<5	5.12	<5
538	7.25 7.25	7.52 7.52	7.8 7.8	6.99 6.99	6.38 6.38	5.39 5.39	<5 <5	5.44 5.44	<5 <5	5.18 5.18	<5 <5	<5 <5	<5 <5
144	7.92	8.31	8.41	7.52	7	6.45	<5	6.58	<5	5.34	<5	<5	<5
539	7.76	8.1	7.79	7.15	6.35	6.5	<5	6.65	<5	5.35	<5	<5	<5
603	8.53	8.56	8.22	7.6	7.14	7.52	<5	7.51	<5	5.74	<5	<5	<5
540	8.32	8.28	8.35	7.57	6.68	7.24	<5	7.37	<5	5.5	<5	<5	<5
541	8.62	8.81	8.66	8.13	7.56	7.67	<5	7.83	<5	6.38	<5	<5	<5

542	8.52	8.5	8.59	8.24	7.14	6.67	<5	6.77	<5	5.11	<5	<5	<5
543	8.7	8.54	8.39	7.73	7.18	7.52	<5	7.62	<5	5.68	<5	<5	<5
544	8.14	8.11	7.71	7.08	6.71	6.66	<5	6.79	<5	5.37	<5	<5	<5
41	8.73	8.78	8.83	8.3	7.57	7.35	<5	7.36	<5	5.93	<5	<5	<5
33	8.73	8.75	8.51	8.17	7.47	7.88	<5	7.87	<5	6.02	<5	5.07	<5
43	8.55	8.62	8.24	7.6	7.16	6.94	~5.26	7.04	~5.34	5.65	~5.3	~5.42	~5.27
546	8.69	8.6	8.45	8.15	6.9	8.31	<5	8.09	<5	5.56	<5	<5	<5
547	7.48	7.62	7.86	6.89	6.58	6.13	<5	6.3	<5	<5	<5	<5	<5
548	8.53	8.71	8.39	8.13	7.18	7.86	<5	7.96	~5.14	5.89	~5.02	5.14	<5
551	8.33	8.51	8.6	7.72	7.63	6.88	<5	7.1	<5	5.95	<5	<5	<5
552	8.14	8.4	8.01	7.11	7.43	7.35	<5	7.07	<5	6.08	<5	5.09	<5
553	8.31	8.36	8.27	7.85	7.17	7.26	<5	7.2	<5	5.88	<5	<5	<5
554	8.14	8.32	8.22	7.89	7.2	7.14	<5	7.07	<5	6	~5.04	5.15	<5
556	7.88	8.15	8.01	7.38	6.93	6.23	<5	6.21	<5	5.16	<5	<5	<5
557	7.82	8.12	8.01	6.97	6.61	6.09	<5	6.22	<5	<5	<5	<5	<5
558	8.39	8.56	8.32	7.66	6.78	7.01	<5	6.99	~5.17	5.55	5.09	5.44	<5
103	8.7	8.73	8.75	8.06	7.47	7.37	<5	7.3	<5	5.95	<5	5.07	<5
559	8.6	8.61	8.39	7.71	6.44	7.3	<5	7.19	<5	5.58	<5	<5	<5
145	8.72	8.72	8.48	7.92	6.7	7.68	<5	7.67	<5	5.52	<5	<5	<5
561	7.5	7.9	8.05	7.03	6.52	6.2	<5	6.31	<5	5.18	<5	<5	<5
562	8.57	8.49	8.3	7.86	7.39	7.49	<5	~7.7	<5	5.9	<5	<5	<5

563	8	8.35	8.11	7.32	7.97	6.23	<5	6.28	<5	5.72	<5	5.09	<5
564	8.37	8.51	8.26	7.71	7.23	7.11	<5	6.91	<5	5.77	<5	<5	<5
565	8.1	8.28	8.47	7.58	6.95	6.44	<5	6.64	<5	5.24	<5	<5	<5
566	7	7.22	7.27	6.51	<6	5.83	<5	5.88	<5	<5	<5	<5	<5
567	7.18	7.23	7.32	6.82	6.17	6.1	<5	6.24	<5	<5	<5	<5	<5
568	8.57	8.65	8.9	7.91	7.11	6.85	<5	7.06	<5	5.52	<5	<5	<5
569	7.77	7.81	7.78	6.93	6.7	5.85	<5	6.05	<5	~5.09	<5	<5	<5
146	8.62	8.67	8.78	7.79	6.96	6.98	<5	7.01	<5	5.23	<5	<5	<5
110	8.25	8.19	8.2	7.66	7.39	7.46	<5	7.81	<5	6.09	<5	<5	<5
570	8.88	8.95	8.71	8.26	7.47	7.46	<5	7.81	~5.03	5.89	<5	5.18	<5
577	8.43	8.42	8.18	7.75	7.28	7.74	<5	7.74	<5	6.26	<5	<5	<5
578	8.76	8.9	8.6	8.22	7.53	8.03	<5	8.11	~5.04	6.51	<5	5.2	<5
581	~8.76	8.72	8.28	7.94	6.95	7.82	<5	7.94	<5	5.82	<5	~5.04	<5
582	7.49	7.68	7.72	7.08	6.57	5.5	<5	5.97	<5	<5	<5	<5	<5
583			7.97	7.22		7.2	<5			5.82	<5	<5	<5
584	7.85	8.08	8.05	7.4	6.57	6.68	<5	6.72	<5	<5	<5	<5	<5
585	8.32	8.36	8.15	7.21	7.4	6.74	<5	6.6	~5.1	6.05	<5	5.25	<5
586	8.54	8.67	8.65	7.94	7.2	7.75	<5	7.62	<5	5.69	<5	<5	<5
105	8.69	8.51	8.74	8.03	7.52	7.99	<5	7.78	<5	5.97	<5	<5	<5
147	7.99	8.2	8.28	7.45	6.9	6.68	<5	6.78	<5	~5.07	<5	<5	<5
588	8.01	7.95	8.04	6.63	7.02								

120	8.71	8.34	8.42	8.13	7.37	6.77	<5	6.93	<5	5.6	<5	<5	<5
589	7.76	7.91	7.99	6.61	6.72								
590	8.64	8.5	8.4	7.95	7.96								
592	8.78	8.68	8.49	8.01	6.87	7.1	<5	7.17	<5	5.71	<5	5.21	<5
593	8.56	8.55	9	8.49	6.87	6.99	<5	7.35	<5	5.49	<5	<5	<5
148	8.93	8.74	9.21	8.6	6.86	7.3	<5	>8	<5	5.49	<5	<5	<5
149	8.36	8.4	8.63	7.93	6.56	6.88	<5	7.26	<5	5.12	<5	<5	<5
100	8.74	8.21	8.43	7.67	6.6	6.28	<5	6.43	<5	<5	<5	<5	<5
101	8.12	7.69	8.05	7.34	6.44	6.59	<5	6.44	<5	~5.01	<5	<5	<5
14/14 a	8.76	8.3	8.75	8.25	7.16	6.93	<5	7.28	<5	5.37	<5	<5	<5
594	8.71	8.29	8.8	7.23	7.48	6.75	<5	~7.11	<5	5.53	<5	5.15	<5
595	8.15	7.55	7.66	7.03	6.52	5.28	<5	5.27	<5	<5	<5	<5	<5
604	9.12	8.47	8.51	8.34	7.65	~7.59	<5	~7.54	<5	6.06	<5	5.07	<5
608	8.33	8.12	8	7.35	6.99	6.7	<5	6.88	<5	5.07	<5	<5	<5
609	8.42	8.13	8.09	7.42	6.91	6.63	<5	6.6	<5	5.22	<5	5.06	5
610	8.68	8.21	8.29	7.68	7.25	6.92	<5	~7.13	<5	5.14	<5	<5	<5
611	8.47	8.12	8.29	7.83	6.57	6.63	<5	6.93	<5	<5	<5	<5	<5
697	8.66	8.37	8.4	7.89	7.26	7.26	<5	6.7	<5	5.82	<5	<5	<5
698	8.85	8.31	8.28	7.94	7.1	7.3	<5	~8.18	<5	5.95	<5	<5	<5
699	8.82	8.36	8.42	7.78	7.19	~7.43	<5	~7.54	<5	5.57	<5	~5	<5

700	9.01	8.47	8.52	8.15	~6	7.98	5.26	8.27	<5	<5	5.25	<5
	8.57	8.09	8.32	7.99	6.5	7.79	5.25	~8.49	<5	<5	5.4	<5
613	8.39	7.91	8.42	7.5	7.08	6.35	<5	~6.61	<5	5.13	<5	<5
701	8.26	8	7.84	6.94	~6.23	6.81	<5	6.89	<5	<5	<5	<5
702	8.1	7.94	8.28	7.49	6.96	6.01	<5	6.57	<5	<5	<5	<5
703	8.32	8.03	8.61	8.25	7.07	6.98	<5	7.22	<5	5.12	<5	<5
645	8.6	8.28	8.66	7.66	7.41	7.27	<5	~6.95	<5	5.83	<5	5.85	<5
704	8.38	8.07	8.59	7.88	7.24	6.26	<5	6.94	<5	5.28	<5	5.07	<5
15	8.15	7.92	8.28	7.41	6.87	6.36	<5	6.53	<5	<5	<5	<5	<5
615	8.83	8.56	8.66	7.98	7	7.56	<5	6.89	<5	5.25	<5	<5	<5
646	8.53	8.37	8.91	7.86	7.31	6.07	<5	~6.18	<5	5.18	<5	4.99	<5
648	9.01	8.69	9.31	8.43	7.27	7.34	<5	7.52	<5	5.64	<5	<5	<5
705	8.1	7.99	8.34	7.42	7.21	6	<5	6.58	<5	5.42	<5	5.24	<5
	8.24	7.92	8.05	7.71	6.86	6.4	<5	6.81	<5	5.27	<5	5.17	<5
	8.28	8.06	8.19	7.6	7.15	6.24	<5	6.76	<5	5.35	<5	5.25	<5
706	8.6	8.41	9.08	7.77	7.32	6.49	<5	6.79	<5	5.59	<5	5.11	<5
707	8.17	7.63	8.23	6.8	7.11	6.09	<5	6.13	<5	<5	<5	<5	<5
647	9.04	8.61	8.92	7.74	7.34	6.88	<5	7.04	<5	<5	<5	5.19	<5
708	8.11	7.85	7.94	7.47	6.65	5.92	<5	6.2	<5	4.99	<5	~5	<5
709	8.45	8.24	8.42	7.7	7.42	6.22	5.04	6.31	~5.02	5.28	<5	5.32	5.14
710	7.55	7.48	7.65	6.81	6.39	5.75	<5	6.08	<5	<5	<5	<5	<5

7011	8.79	8.43	9.02	8.27	7.41	7.24	<5	~7.38	<5	5.79	<5	5.13	<5
712	7.42	7.41	7.69	6.82	6.96	5.65	<5	5.95	<5	5.24	<5	<5	<5
616	9.11	8.24	8.14	7.68	7.39	7.22	<5	~6.99	<5	5.59	<5	5.04	<5
713	7.88	8.01	8.12	7.59	7.01	7.07	<5	6.69	<5	5.33	<5	<5	<5
619	8.49	8.3	7.91	7.5	7.03	7.23	~5.18	6.81	~5.28	5.41	~5.2	~5.35	~5.33
714	7.24	7.38	7.18	7.04	6.06	5.58	<5	~5.6	<5	<5	<5	<5	<5
690	7.98 8.37	8.01 7.89	7.97 8.24	7.53 7.73	6.34 6.2	7.9 7.49	5.02 5.26	~7.97 ~7.62	~5.02 <5	<5 5.13	<5 5	5.15 5.24	<5 5.01
716	8.22	8.14	7.82	7.27	7.01	6.97	<5	~6.57	<5	5.59	<5	5.36	<5
717	8.8	8.36	8.76	8.06	7.32	7.19	<5	~6.69	<5	5.56	<5	<5	<5
718	8.38	8.09	8.19	7.9	6.98	5.61	<5	5.46	<5	<5	<5	5.04	<5
719	6.36	6.36	6.6	6.27	6.08	<5	<5	<5	<5	<5	<5	<5	<5
721	6.88	6.99	6.65	<6	<6	<5	<5	<5	<5	<5	<5	<5	<5
722	7.96	7.92	7.54	7.35	6.72	5.62	~5.24	5.53	~5.24	5.19	5.21	5.31	~5.23
724	8.81	8.2	8.49	7.77	7.05	7.3	<5	~6.99	<5	5.92	<5	<5	<5
620	7.73	7.5	7.7	6.91	7.02	6.17	<5	6.46	<5	5.11	<5	<5	<5
725	8.72	8.18	8.73	7.75	7.13	7.29	<5	6.96	<5	5.73	<5	5.07	<5
727	8.8	8.33	8.29	8.12	7.04	7.44	5.21	6.91	~5.14	5.8	<5	~5.16	<5
728	8.58	8.42	8.49	7.81	7.22	6.87	<5	6.81	<5	5.53	<5	<5	<5
621	8.57	8.22	8.25	7.66	7.49	6.87	<5	7.02	<5	5.64	<5	<5	<5
729	8.8	8.04	8.01	7.8	7.5	7.68	5.17	7.13	~5.21	6.33	<5	5.33	5.14

730	9.12	8.53	9.11	8.79	7.47	7.52	5.03	6.99	<5	6.13	<5	<5	<5
731	8.08	7.8	8.03	7.64	6.85	7.28	5.15	7.26	5.14	5.21	5.04	5.24	5.12
732	8.84	8.38	8.92	8.16	6.77	6.9	<5	7.09	<5	5.14	<5	<5	<5
733	8.93	8.47	8.7	8.25	7.18	7.8	5.12	6.98	~5.15	5.92	<5	~5.11	<5
734	6.59	6.36	6.46	6.15	6.21	<5	<5	<5	<5	<5	<5	<5	<5
624	8.73	8.2	8.78	7.64	7.4								
735	8.56	8.18	8.92	7.77	7.23	6.6	5.52	6.59	<5	<5	<5	5.81	5.2
736	8.19	7.75	8.48	7.72	6.78	6.65	<5	6.4	<5	5.4	<5	5.31	<5
625	8.51	8.12	8.85	8.11	7.33	6.54	<5	6.45	<5	<5	<5	5.22	<5
737	8.7	~8.19	8.85	7.89	7.42	7.21	~5.01	6.72	~5.05	5.71	~5.02	5.45	<5
652	7.31 7.65 7.71	7.31 7.44 7.41	7.65 7.46 7.62	6.96 7.23 7.18	<6 6.43 6.45	7.1 7 7.05	<5 <5 <5	7.37 ~7.4 7.39	<5 <5 <5	<5 <5 <5	<5 <5 <5	<5 <5 5.15	<5 <5 <5
650	9.17	8.43	8.5	8.19	7.95	7.05	5.35	7	~5.23	6.01	5.42	5.37	5.32
738	8.19 8.67	7.91 8.1	7.96 8.38	7.94 8.29	5.78 6.31	8.17 8.59	5.11 5.53	8.59 8.04	<5 <5	<5 5.06	<5 5.03	5.08 5.16	<5 <5
739	7.77	7.47	7.6	7.36	6.52	6.51	<5	6.29	<5	<5	<5	<5	<5
740	8.18 8.38	8.19 7.99	8.1 8.21	7.54 7.84	6.43 6.63	7.68 7.77	5.47 5.25	7.71 7.67	<5 <5	5.46 5.34	5.13 5.23	5.32 5.35	<5 <5
651	9.44	8.54	8.47	8.07	7.7	8.15	<5	7.65	~5.12	5.61	5.75	5.24	<5
617	8.62	8.25	8.82	7.91	7.11	7.21	5.21	7.06	<5	5.86	<5	5.47	<5

741	6.87	6.39	6.77	6.23	<6	5.55	5.08	5.35	<5	5.04	<5	5.08	<5
742	6.92	6.51	6.7	<6	<6	5.98	<5	6.14	<5	<5	<5	<5	<5
618	9.34	8.76	8.55	8.21	7.63	7.81	<5	~7.19	<5	6.04	<5	5.1	<5
743	8.57	8.28	8.64	7.85	7.58	6.58	<5	5.88	<5	5.51	<5	5.01	<5
	8.42	8.14	8.56	7.71	7.57	6.41	~5.02	6.41	<5	5.55	~5	5.11	<5
744	8.85	8.53	8.34	7.71	7.19	7.28	5.37	7.32	~5.25	6.1	5.4	5.41	5.21
745	6.58	6.4	6.54	6.42	<6	5.23	<5	5.15	<5	<5	<5	<5	<5
746	8.26	7.98	8.35	7.84	6.74	6.72	<5	6.95	<5	5.22	<5	5.65	<5
747	7.55	7.33	7.32	7.16	6.4	6.56	~5.06	6.28	<5	5.04	<5	5.06	<5
689	7.48	7.37	7.24	7.04	6.22	6.62	5.12	6.34	<5	5.01	<5	5.21	<5
749	7.31	7.35	7.46	7.11	6.17	5.57	<5	5.92	<5	<5	<5	<5	<5
622	7.26	7.11	7.42	6.92	6.2	5.64	<5	5.91	<5	5.1	<5	<5	<5
688	8.58	8.13	7.9	7.98	6.33	8.7	5.27	8.69	<5	5.1	5.06	5.13	5.04
	8.77	8.37	8.64	8.13	~6.33	8.56	5.32	8.5	~5.03	5.11	5.09	5.2	5.08
	8.96	8.28	8.46	8.18	6.22	8.54	5.2	~7.02	<5	<5	5.04	5.27	5.05
750	8.72	8.26	8.3	7.95	6.51	7.72	5.87	7.51	~5.59	5.65	5.61	5.66	~5.53
752	7.12	6.64	6.79	6.43	6.08	5.85	5.17	5.99	~5.14	5.29	5.21	5.33	5.17
753	6.65	6.49	6.6	6.16	<6	5.38	<5	5.59	<5	<5	<5	5.09	<5
754	8.23	8.15	7.99	6.68	7.01	5.73	<5	5.58	<5	<5	<5	<5	<5
755	8.04	7.77	8.07	7.49	6.75	6.2	5.01	6.29	<5	5.37	<5	5.6	5.02
756	8.87	8.54	8.37	7.89	7.47	7.67	<5	~7.75	<5	6.08	<5	<5	<5

757	8.85	8.41	8.3	7.71	7.37	7.5	<5	~7.44	<5	6.04	<5	<5	<5
758	8.06	8.14	7.75	6.83	6.68	6.11	<5	6.05	<5	5.13	<5	<5	<5
653	8.21	8.02	7.96	7.55	6.46	7.8	5.22	~8.21	~5.08	5.23	5.14	5.31	~5.09
759	8.11	8.07	7.86	7.2	7.52	6.32	5.96	6.45	5.69	5.84	5.81	6.12	5.91
760	8.22	8.35	8.25	7.75	~6.55	7.61	5.85	7.94	5.61	5.76	5.69	5.98	5.74
623	8.09	7.97	7.5	7.42	6.61	6.76	<5	7.03	<5	5.54	<5	<5	<5
761	8	7.75	7.9	7.19	6.99	5.72	<5	5.75	<5	5.09	<5	<5	<5
762	7.04	6.54	6.66	6.18	<6	5.15	<5	5.09	<5	<5	<5	<5	<5
763	7.39	7.14	7.62	6.41	<6	5.75	<5	5.77	<5	<5	<5	<5	<5
764	8.75	8.47	8.15	7.69	<6	7.16	5.17	6.99	~5.08	5.75	~5	5.19	5.02
765	7.98	8.01	8.19	7.54	7.41	7.18	<5	7.63	<5	5.08	<5	5.12	<5
766	6.87	6.64	6.71	6.27	<6	<5	<5	<5	<5	<5	<5	<5	<5
767	8.19	7.79	7.89	7.45	6.7	7.07	5.69	7.28	~5.56	<5	5.6	5.67	5.53
768	8.6	8.16	8	7.16	<6	6.46	<5	6.38	<5	5.31	<5	<5	<5
770	8.74	8.48	7.98	7.69	7.78	7.5	<5	~7.62	<5	5.65	~5	<5	<5
644	8.81	8.29	8.62	7.69	<6	7.22	<5	7.1	<5	5.62	<5	<5	<5
656	8.59	8.46	8.96	8.14	6.81	7.16	<5	~7.63	<5	5.49	<5	5.11	<5
	8.73	8.5	9.29	8.38	7.28	7.21	<5	7.46	<5	5.64	<5	5.21	<5
657	8.32	8.27	8.6	7.61	<6	7	<5	7.14	<5	5.56	<5	<5	<5
	8.58	8.38	8.87	7.87	7.32	7.15	<5	7.28	<5	5.49	<5	<5	<5
658	8.41	8.23	7.95	7.48	6.85	7.33	<5	7.31	<5	5.56	<5	<5	<5

771	8.78	8.53	8.48	8.33	6.63	7.98	6.06	8.23	5.63	5.68	5.64	5.77	5.6
772	7.84 7.91	7.89 7.99	8.11 8.33	7.76 7.58	<6 6.99	6.34 6.32	5.73 6.09	6.41 6.36	<5 <5	5.88 5.91	5.36 5.66	6.07 6.16	5.74 5.71
773	7.9	7.82	7.18	7.02	6.9	6.41	<5	6.51	<5	5.11	<5	<5	<5
774	8	8.12	8.04	7.65	7.08	6.71	<5	6.75	<5	5.37	<5	5.02	<5
627	8.4	8.43	8.21	7.65	6.86	6.75	<5	6.53	<5	5.17	<5	<5	<5
775	8.4	8.47	8.69	7.9	7.44	7.17	<5	7.21	<5	5.74	<5	5.16	<5
777	7.28	7.36	7.55	6.85	~6.54	6.98	<5	7.51	<5	<5	<5	<5	<5
778	7.52	7.44	7.26	6.5	6.17	7.58	<5	7.73	<5	5.25	<5	<5	<5
779	7.41	7.23	7.02	6.32	~6	6.6	<5	6.74	<5	<5	<5	<5	<5
628	7.43	7.32	7.46	6.75	6.69	6.57	<5	6.69	<5	5.16	<5	<5	<5
629	7.59	7.59	7.66	6.7	6.64	6.83	<5	7.04	<5	5.08	<5	<5	<5
780	8.66	8.23	8.46	8.32	7.05	7.7	<5	~7.96	<5	5.37	<5	<5	<5
781	8.23	8.11	8.34	7.78	6.86	7.08	<5	7.15	<5	5.14	<5	<5	<5
782	8.66	8.38	8.13	7.78	7.74	7.59	<5	7.44	<5	5.86	<5	<5	<5
634	8.05	7.82	7.75	7.56	<6	7.96	5.23	8.03	<5	<5	<5	5.05	<5
783	7.69	7.67	7.81	6.73	6.7	6.3	<5	6.24	<5	5.31	<5	<5	<5
784	7.08	7.15	7.21	6.82	6.42	5.4	<5	5.88	<5	<5	<5	<5	<5
785	8.18	7.91	7.84	7.45	7.11	8.58	<5	8.23	<5	5.06	<5	<5	<5
786	7.9	7.79	7.89	7.46	6.32	7.72	5.16	7.97	<5	5.07	5.05	5.13	5.02
787	7.56	7.36	7.48	7.17	6.26	7.31	5.13	7.72	<5	5.11	5.07	5.12	~5

788	8.62	8.34	8.28	7.83	6.27	8.35	5.11	8.47	<5	5.05	<5	5.13	<5
789	8.75	8.23	8.33	8.01	6.1	8.55	<5	8.61	<5	<5	<5	<5	<5
790	6.56	6.59	6.53	<6	<6	5.21	<5	5.23	<5	<5	<5	<5	<5
791	7.9	7.77	7.68	7.83	6.86	8.11	5.17	8.38	<5	5.66	5.04	5.2	<5
792	7.81	7.69	7.73	7.4	6.45	7.33	5.43	7.75	<5	5.51	~5	5.34	5.12
793	7.15	7.05	6.92	6.64	<6	6.94	5.72	7.32	5.38	5.41	5.38	5.5	5.4
631	8.15	8.24	7.53	6.89	6.47	6.51	5.2	6.49	~5.14	5.33	~5.1	5.29	~5.09
794	7.78	7.85	7.48	6.84	6.06	6.16	<5	6.11	<5	<5	<5	<5	<5
795	8.5	8.15	8.18	8.3	5.71	8.32	5.36	8.63	~5.18	5.3	5.27	5.43	~5.25
796	7.21	7.05	7.12	6.76	6.1	6.83	<5	7.23	<5	<5	<5	<5	<5
798	7.53	7.66	7.72	7.06	6.02	6.08	<5	6.33	<5	<5	<5	<5	<5
659	8.26	8.11	8.59	7.56	6.75	6.69	<5	7.12	<5	5.15	<5	5.01	<5
662	7.55	7.69	7.63	7.26	7.11	6	<5	6.21	<5	5.27	<5	<5	<5
799	7.42	7.6	7.57	7.13	6.95	6.76	<5	6.87	<5	5.52	<5	<5	<5
661	8.71	8.53	8.63	7.72	7.05	6.72	<5	6.66	<5	<5	<5	<5	<5
660	8.98	8.6	8.9	8.09	7.23	7.16	<5	7.14	<5	5.18	<5	<5	<5
633	8.29	8.37	8.53	7.81	6.81	6.76	<5	6.85	<5	5.38	<5	<5	<5
800	7.9	7.8	7.74	7.28	6.44	6.66	5.47	6.85	~5.33	5.48	5.5	5.42	5.41
801	7.58	7.55	7.65	7.19	6.49	5.74	<5	5.77	<5	<5	<5	<5	<5
802	8.68	8.44	8.85	7.77	7.25	6.71	<5	6.99	<5	5.17	<5	<5	<5
803	8.4	8.14	8.42	7.16	7.02	6.59	<5	6.67	<5	5.16	<5	<5	<5

636	7.35	7.4	7.12	6.5	6.01	5.66	<5	5.95	<5	5.11	<5	<5	<5
637	7.11	7.33	6.94	6.68	<6	5.65	<5	6.05	<5	<5	<5	<5	<5
804	8.16	8.02	7.72	7.5	<6	8.44	5.12	8.61	<5	5.15	5.09	5.1	5.06
805	7.71	7.63	7.59	7.33	<6	7.13	<5	7.63	<5	<5	<5	<5	<5
806	7.04	6.75	6.75	<6	<6	5.14	<5	5.06	<5	<5	<5	<5	<5
807	7.05	6.86	7	6.71	<6	5.58	<5	5.79	<5	<5	<5	<5	<5
808	7.32	7.03	7.14	6.6	6.19	5.96	<5	6	<5	5.22	<5	5.04	<5
809	7.89	7.49	7.73	7.32	6.02	7.47	5.23	~7.53	<5	5.09	5.11	5.13	5.01
810	7.73	7.67	7.88	7.2	6.91	6.26	5.59	6.24	<5	5.71	<5	5.28	<5
635	8.05	7.97	7.94	7.48	6.75	6.59	<5	~6.53	<5	5.25	<5	<5	<5
811	7.58	7.47	7.6	7.09	6.78	6.14	<5	~6.03	<5	5.15	<5	<5	<5
812	8.25	8.09	8.03	7.19	6.72	7.07	<5	6.8	<5	5.21	<5	<5	<5
813	8.86	8.59	8.8	8.67	7.93	8.63	<5	8.27	<5	6.87	<5	5.5	<5
814	7.35	7.47	7.56	7.18	<6	7.03	5.1	7.14	<5	5.03	<5	<5	<5
815	7.48	7.62	7.82	7.03	6.37	6.42	<5	6.3	<5	<5	<5	<5	<5
816	6.96	7.09	7.29	6.63	~6.03	6.81	<5	6.93	<5	<5	<5	<5	<5
817	7.65	7.59	7.77	7.27	6.25	7.35	5.12	~7.49	<5	<5	<5	<5	<5
640	7.25	7.64	7.72	7.41	6.61	5.82	<5	5.62	<5	<5	<5	<5	<5
819	7.74	7.78	7.93	6.95	7.12	6.28	<5	6.17	<5	5.47	<5	<5	<5
687	8.75	8.45	8.31	7.95	7.76	>8	5.22	>8	<5	7.61	<5	5.85	<5
820	6.79	7.08	7.04	6.27	<6	5.74	<5	5.81	<5	<5	<5	<5	<5

632	8.81	8.63	8.92	7.86	6.98	7.23	5.16	7.03	<5	5.69	<5	5.1	<5
821	8.86	8.53	8.68	8.56	7.85	>8	<5	>8	<5	6.61	<5	4.99	<5
822	9.03	8.79	9.1	8.45	7.65	7.82	<5	7.66	<5	6.2	<5	5.19	<5
823	7.13	6.92	7.08	6.31	<6	5.71	<5	5.7	<5	<5	<5	<5	<5
824	7.36	7.49	7.57	7.12	<6	6.04	<5	6.1	<5	<5	<5	<5	<5
825	8.08	8.24	8.47	7.86	7.04	6.64	<5	6.81	<5	5.3	<5	<5	<5
826	8.76	8.51	8.62	8.36	7.64	9.22	5.14	9.41	<5	5.97	5.09	5.32	<5
827	8.73	8.47	8.64	8.5	7.58	9.2	5.33	9.42	5.01	6.33	5.3	5.48	5.15
828	8.28	8.28	8.8	7.69	7.07	7.12	<5	7.22	<5	5.62	<5	<5	<5
829	7.82	7.52	7.54	7	<6	8.22	5.34	8.43	<5	5.11	~5	5.07	<5
830	7.54	7.6	7.62	6.87	6.15								
663	8.81	8.71	~8.95	8.32	7.39	7.55	<5	~7.61	<5	6.66	<5	<5	<5
664	7.79	~7.69	7.98	7	~6.61	5.64	<5	5.34	5.03	<5	<5	<5	<5
831	7.69	~8	7.75	7.21	6.42	6.13	5.38	6.76	5.44	5.19	5.23	5.5	5.3
832	6.45	6.61	6.43	<6	<6	5.02	<5	<5	<5	<5	<5	<5	<5
833	7.29	7.43	7.4	~6.5	6.98	5.54	<5	5.78	<5	5.11	<5	<5	<5
834	7.98	8.25	8.48	7.79	6.71	7.07	<5	7.25	<5	5.11	<5	<5	<5
837	8.73	~8.85	8.6	7.88	7.68								
641	9	~9.05	8.92	8.25	7.43	7.5	<5			6.22	<5	<5	<5
642	8.86	8.98	8.77	8.42	7.74	>8	<5			6.42	<5	<5	<5
838			8.6	7.85		7.05	<5			5.98	<5	<5	<5

839			<6	<6		<5	<5		<5	<5	<5	<5
840			8.51	8.13		>8	<5		6.57	<5	5.16	<5
881	5.1	5.2	<5	<5	5.8	<5	<5	<5	<5	<5	<5	<5
882	5.3	5.4	5.4	5.1	5.3	<5	<5	<5	<5	<5	<5	<5
883	5.7	5.8	5.7	<5	6.2	<5	<5	<5	<5	<5	<5	<5
884	6.0	6.3	6.1	5.5	6.2	<5	<5	<5	<5	<5	<5	<5
885	7.17	6.92	7.33	6.5	<6	5.33	<5		<5	<5	<5	<5
886	7.44	7.67	7.76	7.06	6.31	~5.49	<5	5.91	<5	<5	<5	<5
887	7.41	7.6	7.75	6.91	6.29	5.51	<5	5.71	<5	<5	<5	<5
888	6.8	7.07	6.74	<6	<6	5.4	<5	~5.14	<5	<5	<5	<5
889	8.38	8.38	8.39	7.14	7.26	5.27	<5	5.71	<5	<5	<5	<5
890	6.46	6.62	6.42	<6	6.21	<5	<5	<5	<5	<5	<5	<5
891	7.44	7.43	7.36	6.87	<6	5.96	<5	6.15	<5	<5	5.02	<5
892	6.69	6.89	6.65	<6	<6	<5	<5	<5	<5	<5	<5	<5
893	6.36	6.38	6.34	<6	6.09	5.04	<5	5.01	<5	<5	5.07	<5
894	6.98	6.73	6.55	<6	<6							
895	7.26	7.16	7.49	6.64	<6	5.44	<5	5.55	<5	<5	<5	<5
896	8.47	7.96	7.98	7.4	6.7	6.87	<5	6.87	<5	5.46	<5	<5
897	8.87	8	8.34	7.9	6.83	7.14	<5	7.23	<5	5.59	<5	<5
898	6.77	6.63	6.45	<6	6.16	5.15	<5	~5.12	<5	<5	<5	<5
899	7.09	6.78	6.76	6.42	<6	5.39	<5	5.22	<5	<5	<5	<5

900	7.62	7.53	7.54	6.66	<6	5.32	<5	5.33	<5	<5	<5	<5
901	7.78	7.7	8.27	7.32	<6	6.37	<5	6.56	<5	<5	<5	<5
902	8.48	8.33	8.81	8.33	7.32	<5	<5	<5	<5	<5	<5	<5
903	6.87	6.84	7.19	6.07	<6	<5	<5	<5	<5	<5	<5	<5
904	7.1	7.26	7.59	6.89	6.09	5.73	<5	6.01	<5	<5	<5	<5
905	<6	<6	<6	<6	6.14	<5	<5	<5	<5	<5	5.26	5.05
906	7.7	7.21	7.35	6.63	6.25	5.21	<5	5.23	<5	<5	<5	<5
907	6.92	6.35	6.91	6.35	<6	5.12	<5	~5.09	<5	<5	<5	<5
908	7.21	6.55	6.83	6.49	<6	7.01	<5	5.41	<5	<5	<5	<5
909	6.83	6.83	6.84	6.52	<6	5.2	<5	~5.03	<5	<5	<5	<5
910	6.66	6.18	6.33	6.04	<6	5.12	<5	<5	<5	<5	<5	<5
911	6.76	6.57	6.56	6.47	<6	5.04	<5	5.06	<5	<5	<5	<5
912	7.19	7.11	7.29	6.64	6.26	5.59	<5	5.65	<5	<5	<5	<5
913	<6	~6	<6	<6	<6	5.1	<5	5.05	<5	<5	<5	<5
914	6.8	6.72	6.21	<6	<6	5.08	<5	5.09	<5	<5	<5	<5
915	<6	<6	<6	<6	<6	<5	<5	<5	<5	<5	<5	<5
916	8	7.69	7.86	7.23	6.63	6.91	<5	6.81	<5	5.26	<5	<5
917	<6	<6	<6	<6	<6	<6	<5	6.09	<5	<5	<5	<5
918	7.04	6.99	7.21	6.42	6.08	5.7	<5	~5.63	<5	<5	<5	<5
919	6.59	6.53	6.91	<6	<6	<5	<5	~5.05	<5	<5	<5	<5
920	7.39	7.15		6.87	~6.31	5.6	<5		<5	<5	<5	<5

921	~6.63				6.47	<5	<5		<5	<5	<5	<5
922	6.92	7.21			6.75		<5		<5	<5	<5	<5
923	7.44	7.67	7.76	7.06	6.31	~5.49	<5	5.91	<5	<5	<5	<5
924	7.44	7.43	7.36	6.87	<6	5.96	<5	6.15	<5	<5	5.02	<5
926	6.69	6.89	6.65	<6	<6	<5	<5	<5	<5	<5	<5	<5
925	6.98	6.73	6.55	<6	<6							

Biological Assays B

FGFR3, VEGFR2 and PDGFR *in vitro* Kinase Inhibitory Activity Assays

Enzymes (from Upstate), prepared at 2x final concentration, were incubated with test compounds, biotinylated Flt3 substrate (biotin-VASSDNEYFYVDF) (Cell Signalling Technology Inc.) and ATP in the appropriate assay buffer (Table 1). The reaction was allowed to proceed for 3 hours (FGFR3), 1 hour (VEGFR2, PDGFR-beta) at room temperature on a plate shaker at 700 rpm before being stopped with 35 mM EDTA, pH 8 (FGFR3, VEGFR2) or 55 mM EDTA, pH 8 (PDGFR-beta). 5x detection mix (50mM HEPES pH 7.5, 0.1% BSA, 11.34 nM Eu-anti-pY (PY20) (PerkinElmer) 74 nM SA-XL665 (Cisbio) for FGFR3, 50 mM HEPES, pH 7.5, 0.1% BSA, 11.34 nM Eu-anti-pY (PY20), 187.5 nM SA-XL665 for VEGFR2 and 50 mM HEPES, pH 7.5, 0.1% BSA, 11.34 nM Eu-anti-pY (PT66) (PerkinElmer), 375 nM SA-XL665 (Cisbio) for PDGFR-beta) was then added to each well and the plate sealed and incubated at room temperature for one hour on a plate shaker at 700 rpm. The plate was then read on a Packard Fusion plate reader or a BMG Pherastar both in TRF mode.

Table 1: Final assay conditions for FGFR3, VEGFR2 and PDGFR-beta assays

Enzyme	1 x Assay Buffer	Flt3 substrate concentration	ATP concentration
FGFR3	A	0.125 µM	8 µM
VEGFR2	B	0.5 µM	0.5 µM
PDGFR-beta	C	1 µM	70 µM

Kinase Assay buffers were:

A: 50 mM HEPES pH 7.5, 6 mM $MnCl_2$, 1 mM DTT, 0.01 % TritonX-100

B: 50 mM HEPES pH 7.5, 6 mM $MnCl_2$, 1 mM DTT, 0.01 % TritonX-100, 0.1 mM Sodium orthovanadate

C: 20 mM HEPES pH 7.5, 10 mM $MnCl_2$, 0.01% Triton X-100, 1 mM DTT, 0.1 mM Sodium orthovanadate

FGFR3 and VEGFR2 Data for the compounds of the invention in the above assays are provided in Table A3.

Ba/F3-TEL-FGFR3 & Ba/F3 (WT) cell proliferation assays

Stably transfected Ba/F3-TEL-FGFR3 cells were plated out into black 96-well tissue culture plates with clear bottoms in RPMI medium containing 10% FBS and 0.25 mg/ml G418 at a density of 5 x 10^3 cells/well (200 µl per well). The parental wild-type Ba/F3 cells (DSMZ no.: ACC 300) were plated out into black 96-well tissue culture plates with clear bottoms in RPMI medium containing 10% FBS and 2 ng/ml mouse IL-3 (R&D Sysems) at a density of 2.5 x 10^3 cells/well (200 µl per well). Plates were placed in an incubator overnight before adding the compounds the following day. Dilutions of compounds were made in DMSO starting at 10 mM and were diluted into the wells to give a final DMSO concentration of 0.1% in assay. Compounds were left on the cells for 72 hours before the plates were removed from the incubator and 20 µl of Alamar Blue™ (Biosource) was added to each well. Plates were placed in the incubator for 4-6 hours before reading plates at 535 nm (excitation) / 590 nm (emission) on a Fusion plate reader (Packard). Where inhibition is high an IC_{50} can be determined.

Data for the compounds of the invention in the above assays are provided in Table A3.

Table A3


Compound number	FGFR3	VEGFR2	BAF3_TEL_FGFR3	BAF3_WT
	pIC_{50} (µM)	pIC_{50} (µM)	pIC_{50} (µM)	pIC_{50} (µM)
201	8.96	7.31	7.89	5.55
12	8.09	6.66	7.27	5.44
13	7.77	5.96	6.85	
215	6.21			
222	7.89	5.85	6.74	5.24
228	7.22	5.96		
19	8.30	6.72	6.74	5.77
24		6.77	7.43	
247		6.77	7.43	
70 & 70a	8.54	7.25	8.43	6.31
281		7.31	8.66	

282		7.31	8.66	
284	9.29	7.70	8.96	5.64
285	9.29	7.70	8.96	5.64
88 & 291	9.19	8.05	9.24	5.72
305	8.72	6.30	7.58	
61	8.29	6.80	6.46	5.72
306	7.51	5.82	6.89	
307		5.72	6.46	
308	8.82	7.57	8.57	
311	8.54	6.92	7.96	
312	8.77	7.44	6.64	
314	7.89	6.31	6.80	
30	8.52	6.89	7.58	
315	8.42	6.82	7.70	
48		7.92	8.92	
316		7.80	9.05	
328	8.85	7.80	8.06	5.96
329	8.85	7.80	8.06	5.96
331	8.74	7.48	8.60	5.13
332	8.74	7.48	8.60	5.13
371	8.87	7.44	8.20	6.21
372	8.87	7.44	8.20	6.21
374	8.62	6.68	8.21	
377	8.49	6.92	8.38	
380	9.09	7.64	8.77	5.04
381	8.82	7.17	8.42	
108	9.21	7.52		
383	9.21	7.52		
384	8.80	7.23	9.05	5.28
39	8.58	6.82	7.96	
40	8.21	6.37	7.66	
386	7.57	6.17	6.92	6.08
389	9.15	6.72	8.43	
390		7.57	9.23	
391	8.15	7	7.31	
392	8.44	7.04	7.92	5.72
393	8.44	7.04	7.92	5.72

404	7	5.82		
405	7	5.82		
409	9.05	6.70	8.89	5.60
410	9.05	6.70	8.89	5.60
423		7.68	9.17	5.72
424		7.68	9.17	5.72
426	6.82	5.77		
427	6.82	5.77		
463		7.47	9.34	6.35
466		7.47	9.41	5.46
467		7.47	9.41	5.46
470 & 471	8.74	7.09	8.96	5.80
472	9	7.60	9.04	
473	9	7.60	9.04	
484	8.70		9.10	6.23
509		8.28		
516	5.72	5.21		
516	7	6.70		

CLAIMS

1. A compound of formula (VI):



(VI)

including any tautomeric or stereochemically isomeric form thereof, wherein

n represents an integer equal to 0, 1, 2, 3 or 4;

R^1 represents hydrogen, C_{1-6}alkyl, C_{2-4}alkenyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, cyanoC_{1-4}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with -NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NR^{12}-$S(=O)_2$-$NR^{14}R^{15}$, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –$Si(CH_3)_3$, C_{1-6}alkyl substituted with -$P(=O)(OH)_2$ or C_{1-6}alkyl substituted with -$P(=O)(OC_{1-6}alkyl)_2$;

each R^{1a} is independently selected from hydrogen, C_{1-4}alkyl, hydroxyC_{1-4}alkyl, C_{1-4}alkyl substituted with amino or mono- or di(C_{1-4}alkyl)amino or -$NH(C_{3-8}$cycloalkyl), cyanoC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl, and C_{1-4}alkyl substituted with one or more fluoro atoms;

each R^2 is independently selected from hydroxyl, halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{2-4}alkynyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkyl, haloC_{1-4}alkoxy, hydroxyhaloC_{1-4}alkyl, hydroxyhaloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, haloC_{1-4}alkoxyC_{1-4}alkyl, C_{1-4}alkoxyC_{1-4}alkyl wherein each C_{1-4}alkyl may optionally be substituted with one or two hydroxyl groups, hydroxyhaloC_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkyl substituted with R^{13}, C_{1-4}alkyl substituted with -C(=O)-R^{13}, C_{1-4}alkoxy substituted with R^{13}, C_{1-4}alkoxy substituted with -C(=O)-R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with -NR^7R^8, C_{1-4}alkyl substituted with –C(=O)-NR^7R^8, C_{1-4}alkoxy

substituted with -NR^7R^8, C_{1-4}alkoxy substituted with –C(=O)-NR^7R^8, -NR^7R^8 and -C(=O)-NR^7R^8; or when two R^2 groups are attached to adjacent carbon atoms they may be taken together to form a radical of formula:

-O-$(C(R^{17})_2)_p$-O-;

-X-CH=CH-; or

-X-CH=N-; wherein R^{17} represents hydrogen or fluorine, p represents 1 or 2 and X represents O or S;

R^4 and R^5 each independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with -$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –NH- $S(=O)_2$-$NR^{14}R^{15}$, R^{13} or C_{1-6}alkyl substituted with R^{13};

R^6 represents C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S; said C_{3-8}cycloalkyl, C_{3-8}cycloalkenyl, phenyl, 4 to 7-membered monocyclic heterocyclyl, optionally and each independently being substituted by 1, 2, 3, 4 or 5 substituents, each substituent independently being selected from cyano, C_{1-6}alkyl, cyanoC_{1-6}alkyl, hydroxyl, carboxyl, hydroxyC_{1-6}alkyl, halogen, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}alkoxyC_{1-6}alkyl, C_{1-6}alkyl-O-C(=O)-, -$NR^{14}R^{15}$, -C(=O)-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -C(=O)-$NR^{14}R^{15}$, –$S(=O)_2$-C_{1-6}alkyl, –$S(=O)_2$-haloC_{1-6}alkyl, –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-haloC_{1-6}alkyl, C_{1-6}alkyl substituted with –$S(=O)_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-$S(=O)_2$-haloC_{1-6}alkyl or C_{1-6}alkyl substituted with –NH-$S(=O)_2$-$NR^{14}R^{15}$;

R^7 and R^8 each independently represent hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, hydroxyhaloC_{1-6}alkyl or C_{1-6}alkoxyC_{1-6}alkyl;

R^{13} represents C_{3-8}cycloalkyl or a saturated 4 to 6-membered monocyclic heterocyclyl containing at least one heteroatom selected from N, O or S, wherein said C_{3-8}cycloalkyl or monocyclic heterocyclyl is optionally substituted with 1, 2 or 3 substituents each independently selected from halogen, hydroxyl, C_{1-6}alkyl, -C(=O)-C_{1-6}alkyl, C_{1-6}alkoxy, or -$NR^{14}R^{15}$; and

R^{14} and R^{15} each independently represent hydrogen, or haloC_{1-4}alkyl, or C_{1-4}alkyl optionally substituted with a substituent selected from hydroxyl, C_{1-4}alkoxy, amino or mono-or di(C_{1-4}alkyl)amino.

2. A compound according to claim 1 wherein R^1 represents hydrogen, C_{1-6}alkyl, hydroxyC_{1-6}alkyl, haloC_{1-6}alkyl, C_{1-6}alkoxyC_{1-6}alkyl wherein each C_{1-6}alkyl may optionally be substituted with one or two hydroxyl groups, C_{1-6}alkyl substituted with -NR^4R^5, C_{1-6}alkyl substituted with –C(=O)-NR^4R^5, –S(=O)$_2$-C_{1-6}alkyl, –S(=O)$_2$-$NR^{14}R^{15}$, C_{1-6}alkyl substituted with -S(=O)$_2$-C_{1-6}alkyl, C_{1-6}alkyl substituted with –NH-S(=O)$_2$-C_{1-6}alkyl, R^6, C_{1-6}alkyl substituted with R^6, C_{1-6}alkyl substituted with –C(=O)-R^6, hydroxyC_{1-6}alkyl substituted with R^6, or C_{1-6}alkyl substituted with –Si(CH_3)$_3$; wherein each R^{1a} is hydrogen.

3. A compound according to claim 1 wherein each R^{1a} is hydrogen.

4. A compound according to any one of the preceding claims wherein R^1 represents C_{1-6}alkyl.

5. A compound according to any one of the preceding claims wherein R^1 represents CH_3- or CD_3-.

6. A compound according to any one of the preceding claims wherein R^2 is independently selected from halogen, cyano, C_{1-4}alkyl, C_{2-4}alkenyl, C_{1-4}alkoxy, hydroxyC_{1-4}alkyl, hydroxyC_{1-4}alkoxy, haloC_{1-4}alkoxy, C_{1-4}alkoxyC_{1-4}alkyl, R^{13}, C_{1-4}alkoxy substituted with R^{13}, -C(=O)-R^{13}, C_{1-4}alkyl substituted with NR^7R^8, C_{1-4}alkoxy substituted with NR^7R^8, -NR^7R^8 and -C(=O)-NR^7R^8.


7. A compound according to claim 6 wherein R^2 represents C_{1-4}alkoxy.

8. A compound according to claim 6 or claim 7 wherein R^2 represents CH_3O- or CD_3O-.

9. A compound according to claim 1 wherein R^1 represents C_{1-6}alkyl, each R^{1a} is hydrogen, n represents an integer equal to 2 and each R^2 represents C_{1-4}alkoxy.


10. A compound according to claim 1 wherein R^1 represents $-CH_3$, each R^{1a} is hydrogen, n represents an integer equal to 2 and each R^2 represents CH_3O-.

11. A compound according to claim 1 wherein the compound is




.

12. A compound according to claim 1 wherein the compound is




.

13. A compound according to claim 1 wherein the compound is




.


14. A compound according to claim 1 wherein the compound is selected from



,



and



.

15. A compound as defined in any one of claims 1 to 14 or a pharmaceutically acceptable salt or solvate thereof.

PYRAZOLYL QUINOXALINE KINASE INHIBITORS

ABSTRACT

The invention relates to new quinoxaline derivative compounds, to pharmaceutical compositions comprising said compounds, to processes for the preparation of said compounds and to the use of said compounds in the treatment of diseases, e.g. cancer.

[No figure]