

1 / 15

**SEQ ID NO: 2** Amino acid sequence of the wild-type alpha chain variable domain. CDRs are underlined and CDR1, 2, 3 are designated SEQ ID NO: 41, 42 and 43, respectively. The constant region is shown in italics. The FM2 and FM3 regions are in bold and are designated SEQ ID NO: 47 and 48, respectively.

AQSV AQPEDQVNVAEGNPLTVKCTYSVSGNPY**LFWYVQYPN**RG**LQFLLK**YITGDNLV**KGSY**  
**GFEAEFNKSQTSFHLKKPSALVSDS****ALYFCAVRDINS**GAGSYQLTFGKGTKLSVIPNIQNP  
*DPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDKTVLDMRSMDFKSNSAVAWS*  
*NKSDFACANAFNNSIIPEDTFFPSP*ESS

**SEQ ID NO: 3** Amino acid sequence of the wild-type beta chain variable domain. CDRs are underlined and are designated SEQ ID NO: 44, 45 and 46, respectively. The constant region is shown in italics. The FM2 and FM3 regions are in bold and are designated SEQ ID NO: 49 and 50, respectively.

SAVISQKPSRDICQRGTSLTIQCVDSQVT**MFWYRQQPGQSLTLIATANQ**SEAT**YESGF**  
**VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLC**SVGGSGGADTQYFGPGTRLTVLEDLKNVF  
*PPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWVWNGKEVHSGVSTDPQPLKEQPA*  
*LNDSRYCLSSRLRVSATFWQNPRNHFR*CQVQFYGLSENDEWTQDRAKPVTQIVSAEAWGRA  
D

FIG. 1

2 / 15

**SEQ ID NO: 4** Amino acid sequence of the soluble extracellular region of a native alpha chain of the invention. CDRs are underlined. The constant region is shown in italics and the non-native cysteine residue is shown in bold (at position 48 of constant region).

AQSVAQPEDQVNVAEGNPLTVKCTYSVSGNPYLEFWYVQYPNRGLQFLLKYITGDNLVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDINSGAGSYQLTFGKGTKLSVIPNIQNP  
DPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDK**C**VLDMRSMDFKSNSAVAWS  
NKSDFACANAFNNSIIPEDTFFPSPSS

**SEQ ID NO: 5** Amino acid sequence of the soluble extracellular region of a native beta chain of the invention. CDRs are underlined. The constant region is shown in italics and the non-native cysteine residue is shown in bold (at position 57 of constant region). Additional non-native amino acids at position 75 and position 89 of the constant region are also shown in bold.

SAVISQKPSRDICQRGTSLTIQCVDSQVTMMFWYRQQPGQSLTLIATANQGSEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCVGGSGGADTQYFGPGTRLTVLEDLKNVF  
PPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWVNGKEVHSGV**C**TDPQPLKEQPA  
LNDSTRY**A**LSSRLRVSATFWQ**D**PRNHFRQCQVQFYGLSENDEWTQDRAKPVTQIVSAEAWGRA  
D

FIG. 2

3 / 15

Amino acid sequences of mutated TCR alpha chain variable regions of the invention

**SEQ ID NO: 6** mutant alpha chain (a12) CDRs are underlined, **mutations in bold**

AQSVAQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKYITGDNLVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRD**SDQH**AGSYQLTFGKGTKLSVIP

**SEQ ID NO: 7** mutant alpha chain (a24) CDRs are underlined, **mutations in bold**

AQSVAQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKY**LGDS**ALVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDINSGAGSYQLTFGKGTKLSVIP

**SEQ ID NO: 8** mutant alpha chain (a24l) CDRs are underlined, **mutations in bold**

AQSVAQPEDLVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKY**LGDS**ALVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDINSGAGSYQLTFGKGTKLSVIP

**SEQ ID NO: 9** mutant alpha chain (a28) CDRs are underlined, **mutations in bold**

AQSVAQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKYITGDNLVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVR**SSRQ**HAGSYQLTFGKGTKLSVIP

**SEQ ID NO: 10** mutant alpha chain (a78l) CDRs are underlined, **mutations in bold**

AQSVAQPEDLVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKY**LGDS**ALVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDIDSGAGSYQLTFGKGTKLSVIP

**SEQ ID NO: 11** mutant alpha chain (a82l) CDRs are underlined, **mutations in bold**

AQSVAQPEDLVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKY**LGDS**ALVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDI**R**SGAGSYQLTFGKGTKLSVIP

**SEQ ID NO: 12** mutant alpha chain (a82) CDRs are underlined, **mutations in bold**

AQSVAQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKY**LGDS**ALVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDI**R**SGAGSYQLTFGKGTKLSVIP

**SEQ ID NO: 51** mutant alpha chain (a86) CDRs are underlined, **mutations in bold**

AQSVAQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRLQFLLKY**LTDD**NLVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDINSGAGSYQLTFGKGTKLSVIP

FIG. 3

4 / 15

Amino acid sequences of mutated TCR beta chain variable regions of the invention

**SEQ ID NO: 13** mutant beta chain (b5) CDRs are underlined, mutations in bold

SAVISQKPSRDI**K**QRGTSLT**I**QCQVDSQVTMMFWYRQQPGQSPTLIATA**AWT**GGEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 14** mutant beta chain (b12) CDRs are underlined, mutations in bold

SAVISQKPSRDI**K**QRGTSLT**I**QCQVDKRLALMFWYRQQPGQSPTLIATA**AWT**GGEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 15** mutant beta chain (b52) CDRs are underlined, mutations in bold

SAVISQKPSRDI**K**QRGTSLT**I**QCQVDKRLALMFWYRQQPGQSPTLIATA**AWT**GGEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGAADTQYFGPGTRLTVL

**SEQ ID NO: 16** mutant beta chain (b56) CDRs are underlined, mutations in bold

SAVISQKPSRDI**K**QRGTSLT**I**QCQVDKRLAMMFWYRQQPGQSPTLIATA**AWT**GGEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 17** mutant beta chain (b56l) CDRs are underlined, mutations in bold

SAVISQKPSRDI**K**QRGTSLT**I**QCQVDKRLAMMFWYRQQPGQSLTLIATA**AWT**GGEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 18** mutant beta chain (b65) CDRs are underlined, mutations in bold

SAVISQKPSRDI**K**QRGTSLT**I**QCQVDKRLALMFWYRQQPGQSPTLIATA**AWT**GSEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 19** mutant beta chain (b65l) CDRs are underlined, mutations in bold

SAVISQKPSRDI**K**QRGTSLT**I**QCQVDKRLALMFWYRQQPGQSLTLIATA**AWT**GSEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 20** mutant beta chain (b67) CDRs are underlined, mutations in bold

FIG. 4

5 / 15

SAVISQKPSRDIKQRGTSLT IQCQVDKRLAMMFWYRQQPGQSP TLIATAWTGGEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGAADTQYFGPGTRLTVL

**SEQ ID NO: 21** mutant beta chain (b67I) CDRs are underlined, mutations in bold

SAVISQKPSRDIKQRGTSLT IQCQVDKRLAMMFWYRQQPGQSLTLIATAWTGGEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGAADTQYFGPGTRLTVL

**SEQ ID NO: 22** mutant beta chain (b68I) CDRs are underlined, mutations in bold

SAVISQKPSRDIKQRGTSLT IQCQVDKRLAMMFWYRQQPGQSP TLIATAWTGSEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 23** mutant beta chain (b68I) CDRs are underlined, mutations in bold

SAVISQKPSRDIKQRGTSLT IQCQVDKRLAMMFWYRQQPGQSLTLIATAWTGSEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGGADTQYFGPGTRLTVL

**SEQ ID NO: 52** mutant beta chain (b71) CDRs are underlined, mutations in bold

SAVISQKPSRDIKQRGTSLT IQCQVDSQVAMMFWYRQQPGQSP TLIATAWQGSEATYESGF  
VIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCSVGGSGAADTQYFGPGTRLTVL

**FIG. 4** Cont'd

Amino acid sequences of alpha and beta chains of ImmTAC molecules of the invention

ImmTAC1 (S2C)

**SEQ ID NO: 32** ImmTAC alpha chain (comprising SEQ ID NO: 7(a24) and the constant domain of SEQ ID NO: 4, truncated by 8 amino acids)

AQSV AQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPN RGLQFLLKYLGDSALVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDINS GAGSYQLTFGKGTKLSVIPNIQNP  
DPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKSDVYITDKCVLDMRSMDFKSNSAVAWS  
NKSDFACANAFNNSIIPEDT

**SEQ ID NO: 33** ImmTAC beta chain (comprising an anti-CD3 scFv (**bold**) fused via a linker (*italics*) to a TCR beta chain comprising SEQ ID NO: 15(b52) and the constant domain of SEQ ID NO: 3)

AIQMTQSPSSLSASVGDRTITCRASQDIRNYLNWYQKPGKAPKLLIYYTSRLESGVPSR  
FSGSGSGTDYTLTISSSLQPEDFATYYCQQGNTLPWTFGQGTKVEIKGGGGSGGGSGGGGS  
GGGGSGGGSEVQLVESGGGLVQPGGSLRLSCAASGYSTGYTMNWVRQAPGKGLEWVALIN  
PYKGVSTYNQKFKDRFTISVDKSKNTAYLQMNSLRAEDTAVYYCARS GYYGDSDWYFDVWG  
QGTLVTVSSGGGGSSAVISQKPSRDIKQRGTSLT IQCQVDKRLALMFWYRQQPGQSPTLIA  
TAWTGGEATYESGFVIDKFPISRPNLTFSTLT VSNMSPEDSSIYLCVGGSGAADTQYFGP  
GTRLTVLEDLKNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWVNGKEVHS  
GVCTDPQPLKEQPALNDSRYALSSRLRV SATFWQDPRNHFR CQVQFYGLSENDEWTQDRAK  
PVTQIVSAEAWGRAD

FIG. 5

7 / 15

Amino acid sequences of alpha and beta chains of ImmTAC molecules of the invention

ImmTAC2 (S2E)

**SEQ ID NO: 35** ImmTAC alpha chain (comprising SEQ ID NO: 7(a24) and the constant domain of SEQ ID NO: 4, truncated by 8 amino acids)

AQSVAPQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPNRGLQFLLKYLGD~~SALVKGSY~~  
GF~~EAEFNKSQTSFHLKKPSALVSDS~~ALYFCAVRDINS~~GAGSYQLTFGKGTKLSVIPNIQNP~~  
DPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDS~~DVYITDKCVLDMRSMDFKSNSAVAWS~~  
NK~~SDFACANAFNNSII~~PEDT

**SEQ ID NO: 36** ImmTAC beta chain (comprising an anti-CD3 scFv (**bold**) fused via a linker (*italics*) to a TCR beta chain comprising SEQ ID NO: 18(b65) and the constant domain of SEQ ID NO: 5)

AIQMTQSPSSLSASVGDRVTITCRASQDIRNYLNWY**QQKPGKAPKLLIYYTSRLESGVPSR**  
**FSGSGSGTDYTLTISS**LQPEDFATYYC**QQGNTLPWTFGQGTKVEIK**GGGGSGGGSGGGGS  
GGGGSGGGSEVQLVESGGGLVQPGGSLRLS**CAASGYSTGYTMNWVRQAPGKGLEW**VALIN  
PYKGVSTYNQKFKDRFTISVDKSKNTAYLQ**MNSLRAEDTAVYYCAR**SGYYGDSDWYFDVWG  
**QGT**LV**TVSS**GGGGSSAVISQKPSRDIKQRGTS**LT**IQ**Q**QVDKRLALMFWYRQQPGQSPTLIA  
TAWTGSEATYESGFVIDKFPISRPNLTFSTLTVSNMSPEDSSIYLC**SVGGSGGADTQYFGP**  
GTRLTVLEDLKNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWVNGKEVHS  
GVCTDPQPLKEQPALNDSRYALSSRLRVSATFWQDPRNHFR**CQVQFYGLSEN**DEWTQDRAK  
PVTQIVSAEAWGRAD

FIG. 6

8 / 15

Amino acid sequences of alpha and beta chains of ImmTAC molecules of the invention

ImmTAC3 (S2G)

**SEQ ID NO: 37** ImmTAC alpha chain (comprising SEQ ID NO: 12(a82) and the constant domain of SEQ ID NO: 4, truncated by 8 amino acids)

AQSV AQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPN RGLQFLLKYLGD SALVKGSY  
GFEAEFNKSQTSFHLKKPSALVSDSALYFCAVRDIRSGAGSYQLTFGKGTKLSVIPNIQNP  
DPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDKCVLDMRSMDFKSNSAVAWS  
NKSDFACANAFNNSIIIPEDT

**SEQ ID NO: 38** ImmTAC beta chain (comprising an anti-CD3 scFv (**bold**) fused via a linker (*italics*) to a TCR beta chain comprising SEQ ID NO: 15(b52) and the constant domain of SEQ ID NO: 5)

AIQMTQSPSSLSASVGDRVITITCRASQDIRNYLNWY**QQKPGKAPKLLIYYTSRLESGVPSR**  
**FSGSGSGTDYTLTISSLPEDFATYYCQQGNTLPWTFGQGTKVEIK**GGGGSGGGSGGGGS  
GGGGSGGGSEVQLVESGGGLVQPGGSLRLS**CAASGYSTGYTMNWVRQAPGKGLEWVALIN**  
**PYKGVSTYNQKFKDRFTISVDKSKNTAYLQMNSLRAEDTAVYYCARSGYYGDSDWYFDVWG**  
**QGTTLVTVSS**GGGGSSAVISQKPSRDIKQRGTS**LTIQ**CQVDKRLALMFWYRQQPGQSPTLIA  
TAWTGGEATYESGFVIDKFPISRPNLTFSTLTVSNMSPEDSSIYLCVGGSGAADTQYFGP  
GTRLTVLEDLKNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWVNGKEVHS  
GVCTDPQPLKEQPALNDSRYALSSRLRVSATFWQDPRNHFR**CQVQFYGLSENDEWTQDRAK**  
PVTQIVSAEAWGRAD

FIG. 7

Amino acid sequences of alpha and beta chains of ImmTAC molecules of the invention

ImmTAC4 (S2I)

SEQ ID NO: 39 ImmTAC alpha chain (comprising SEQ ID NO: 12(a82) and the constant domain of SEQ ID NO: 4, truncated by 8 amino acids)

AQSV AQPEDQVNVAEGNPLTVKCTYSVSGNPYLFWYVQYPN RGLQFL LKYLGD SALVKGSY  
GF EA EFNKSQTSFHLKKPSALVSDSALYFCAVRDIRSGAGSYQLTFGKGTKLSVIPNIQNP  
DPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDKCVLDMRSMDFKSNSAVAWS  
NKSD FACANAFNNSII PEDT

SEQ ID NO: 40 ImmTAC beta chain (comprising an anti-CD3 scFv (bold) fused via a linker (italics) to a TCR beta chain comprising SEQ ID NO: 18(b65) and the constant domain of SEQ ID NO: 5)

AIQMTQSPSSLSASVGDRVTITCRASQDIRNYLNWY**QQKPGKAPKLLIYYTSRLESGVPSR**  
**FSGSGSGTDYTLTISS***LQPEDFATYYCQQGNTLPWTFGQGTKVEIKGGGGSGGGGSGGGGS*  
*GGGGSGGGSEVQLVESGGGLVQPGGSLRLS*CAASGYSTGYTMNWVR**QAPGKGLEW**VALIN  
PYKGVSTYNQKFKDRFTISVDKSKNTAYLQMN**SLRAEDTAVYYCAR**SGYYGDSDWYFDVWG  
**QGT***LVTVSSGGGGSS*AVISQKPSRDIKQRGTS**LT***IQ*CQVDKRLALMFWYRQQPGQSPTLIA  
TAWTGSEATYESGFVIDKFPISRPNLTFSTLTVSNMSPEDSSIYLC**SVGGSGGADTQYFGP**  
GTRLTVLEDLKNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYDPDHVELSWWVNGKEVHS  
GVCTDPQPLKEQPALNDSRYALSSRLRVSATFWQDPRNHFR**CQVQFYGLSENDEWTQDRAK**  
PVTQIVSAEAWGRAD

FIG. 8

10 / 15

Functional testing of ImmTAC molecules of the invention (Exp. No. PCA006)




FIG. 9

Further functional testing of ImmTAC molecules of the invention (Exp. No. PCA005)




FIG. 10

12 / 15

Further specificity testing of ImmTAC molecules of the invention (Exp. No. PCA002/3)




FIG. 11

Further specificity testing of ImmTAC molecules of the invention (Exp. No. PCA002/3)




FIG. 11 Cont'd

14 / 15

Real-time tumour cell killing by ImmTAC redirected T cells (Exp. No. PCA004)




FIG. 12

SUBSTITUTE SHEET (RULE 26)

15 / 15

Real-time tumour cell killing by ImmTAC redirected T cells (Exp. No. PCA004)




FIG. 12 Cont'd