

Description

This invention relates to novel shampoo compositions and more particularly, to shampoo compositions which have high detergent action on oily dirt and exhibit good foamability at the time of shampooing.

Known shampoo compositions have usually utilized, as a base, anionic surface active agents such as alkylsulfates and polyoxyethylene alkylsulfates, or amphoteric surface active agents such as alkylbetaines and alkylamine oxides either singly or in combination thereof. However, shampoo compositions using these types of materials do not have satisfactory detergency against oily soil such as of triglycerides which may impart objectionable oiliness to hair or skin. As a consequence, there are produced some troubles such as stickiness of hair or skin and a hard-to-groom problem of hair due to "greasiness", especially, inherent to the younger generation. Thus, there is a high demand for shampoo compositions with improved detergency.

On the other hand, it is known that for washing of wool, ethylene oxide added type nonionic surface active agents have higher detergency against oils and fats including triglycerides than the anionic surface active agents. Known shampoo compositions to which ethylene oxide added type nonionic surface active agents are incorporated have high detergency against oily soil. However, foamability which is one of the important properties of shampoo is substantially lost. This loss becomes considerable especially under highly oil-soiling conditions. Accordingly, ethylene oxide added type nonionic surface active agents are used only for hair condition controlling purposes so as to impart smoothness to hair.

Under these circumstances, the present inventors made intensive studies to overcome the above drawback and found that incorporation of a specific type of ethylene oxide-added type nonionic surface active agent and N-acyl-N-methyl-beta-alanine or its salt to known shampoo compositions comprising, as a base, anionic surface active agents (exclude N-acyl-N-methyl-beta-alanine and its salts in the present specification) or amphoteric surface active agents or mixtures thereof results in shampoo compositions which have high detergency against oily soil and exhibit good foamability. The present invention has been accomplished on the basis of the above finding.

Accordingly, one object of the invention is to provide a shampoo composition which comprises, in addition to an anionic surface active agent and/or an amphoteric surface active agent as a base, the following two ingredients (A) and (B),

(A) 0.5 to 10 wt% of an N-acyl-N-methyl-beta-alanine or its salt represented by the formula (I)



In which R represents a linear or branched, saturated or unsaturated hydrocarbon group having from 7 to 21 carbon atoms, and M represents hydrogen, an alkali metal, ammonium, alkyl-substituted ammonium with an alkyl moiety having from 1 to 3 carbon atoms, alkanolamine having a hydroxyalkyl group having from 1 to 3 carbon atoms, or basic amino acid, and (B) 0.5 to 20 wt% of one or more of ethylene oxide-added type nonionic surface active agents selected from polyoxyethylene alkyl or alkenyl ethers, polyoxyethylene alkylphenyl ethers, polyoxyethylene fatty acid esters, and polyoxyethylene hardened castor oils.

Preferable examples of ingredient (A) used in the present invention include N-acyl-N-methyl-beta-alanine or its salts represented by the formula (I) in which RCO— represents an acyl group such as lauroyl, myristoyl, palmitoyl, stearoyl, oleoyl or isostearoyl. M in the formula (I) includes alkali metals, ammonium, alkyl-substituted ammonium with an alkyl moiety having from 1 to 3 carbon atoms, alkanolamines having a hydroxyalkyl group having from 1 to 3 carbon atoms, or basic amino acids such as lysine and arginine. Of these compounds, N-lauroyl-N-methyl-beta-alanine, or sodium and thiethanolamine salts thereof are preferred. The (A) ingredient is used in the range of from 0.5 to 10 wt% (hereinafter referred to simply as %), preferably from 1 to 6%, of the total composition.

Preferable examples of (B) ingredient are mentioned below.

(1) Polyoxyethylene alkyl or alkenyl ethers which have an alkyl or alkenyl group containing from 8 to 20 carbon atoms on average and which contain 3 to 20 moles of ethylene oxide.

(2) Polyoxyethylene alkylphenyl ethers which have an alkyl group containing from 8 to 12 carbon atoms on average and which contain 3 to 20 moles of ethylene oxide.

(3) Polyoxyethylene fatty acid esters which have an alkyl or alkenyl group containing from 8 to 20 carbon atoms on average and which contain 8 to 100 moles of ethylene oxide.


(4) Polyoxyethylene hardened castor oil which contain 3 to 200 moles of ethylene oxide on average.

Of these ethylene oxide-added type nonionic surface active agents, polyoxyethylene alkyl or alkenyl ethers and polyoxyethylene hardened castor oils are preferred. These (B) ingredients are used in amounts ranging from 0.5 to 20%, preferably 0.5 to 10% and most preferably from 1 to 10%, of the total composition.

The anionic surface active agents and amphoteric surface active agents used as a base of the shampoo composition according to the present invention are as follows.


(1) Anionic surface active agents

- 1) Linear or branched alkylbenzenesulfonates having an alkyl group containing from 10 to 16 carbon atoms on average.
- 2) Alkyl or alkenyl ethersulfates which have a linear or branched alkyl or alkenyl group containing from 8 to 20 carbon atoms on average and which contain from 0.5 to 8 moles of ethylene oxide on average in one molecule.
- 3) Alkyl- or alkenylsulfates having an alkyl or alkenyl group containing from 10 to 20 carbon atoms on average.
- 4) Olefin sulfonates having from 10 to 20 carbon atoms on average in one molecule.
- 5) Alkanesulfonates having from 10 to 20 carbon atoms on average in one molecule.
- 6) Saturated or unsaturated fatty acid salts having from 10 to 20 carbon atoms on average in one molecule.
- 7) Alkyl- or alkenyl ethoxy carboxylates which have an alkyl or alkenyl group containing from 10 to 20 carbon atoms on average and which contain 0.5 to 8 moles of ethylene oxide in one molecule.
- 8) alpha-sulfo fatty acids or their esters represented by the following formula



in which Y represents an alkyl group having from 1 to 3 carbon atoms or a counter ion of the anionic surface active agent, Z represents a cation of the anionic surface active agent, and R_2 represents an alkyl or alkenyl group having from 10 to 20 carbon atoms.

9) Succinic acid derivatives of the following formula




in which R_3 represents an alkyl or alkenyl group having from 6 to 20 carbon atoms, and Y_1 and Y_2 independently represent a cation.

Examples of the cations of these anionic surface active agents include alkali metal ions such as sodium and potassium, alkaline earth metal ions such as calcium and magnesium, ammonium ion, alkanolamines having from 1 to 3 alkanol groups containing 2 or 3 carbon atoms, e.g. monoethanolamine, diethanolamine, triethanolamine and triisopropanolamine, and basic amino acids such as lysine and arginine. Preferably, sodium ammonium and triethanolamine are used as the cation.


(2) Amphoteric surface active agents

1) Alkylamine oxides represented by the following formula




in which R_6 represents an alkyl or alkenyl group having from 10 to 20 carbon atoms, and R_7 and R_8 may be the same or different and are an alkyl group having from 1 to 3 carbon atoms.

2) Alkylbetaines or sulfobetaines represented by the following formula




in which R_9 represents an alkyl or alkenyl group having from 10 to 20 carbon atoms, R_{10} and R_{11} independently represent an alkyl group having from 1 to 4 carbon atoms, p is an integer of 1 to 3, and X' represents $-COO^{\ominus}$ or $-SO_3^{\ominus}$ group.

3) Imidazoline-type amphoteric surface active agents represented by the following formula



in which R_{12} represents a fatty acid radical having from 10 to 20 carbon atoms on average, R_{13} represents hydrogen, sodium or CH_2COOMe , and Z' represents COOMe , CH_2COOMe or



in which Me represents sodium, hydrogen or an organic base, and G represents OH, an acidic salt, or an anionic surface active sulfate or sulfide.

4) Amidoamine compounds represented by the following general formula



in which R_{14} represents a saturated or unsaturated aliphatic hydrocarbon group having from 7 to 21 carbon atoms, R_{15} represents a hydrogen atom or a hydroxyalkyl group having from 1 to 4 carbon atoms, R'_{15} may be the same as R_{15} or is $-\text{CH}_2\text{CH}_2\text{COOM}'$, and M' represents a hydrogen atom, an alkali metal, an ammonium salt, or an alkanolamine salt.

Of these shampoo bases, there are most preferably used anionic surface active agents such as linear or branched alkylsulfates having from 10 to 20 carbon atoms on average, polyoxyethylene alkylsulfates whose alkyl group has from 8 to 20 carbon atoms (the average number of addition moles ranging from 0.5 to 8), and olefin sulfonates having from 10 to 20 carbon atoms on average; and alkylamine oxides, alkylbetaines and imidazoline-type amphoteric surface active agents in which the alkyl group in each compound has from 10 to 20 carbon atoms on average. The shampoo bases may be used singly or in combination and are used in an amount of from 5 to 30%, preferably 10 to 25%, of the shampoo composition.

The shampoo composition according to the present invention should be in the form of paste or liquid using water in view of the manner of application thereof. The pH of the paste or liquid is preferred to range from 5.5 to 7.5.

Aside from the above-described essential ingredients, the shampoo composition of the invention may further comprise ingredients ordinarily used in known shampoo compositions, including a foam stabilizer such as alkylolamides; solubilizers such as propylene glycol, glycerine and urea; viscosity modifiers such as ethanol, inorganic salts, higher alcohols, hydroxyethyl cellulose and hydroxypropyl cellulose; pH controlling agents such as phosphoric acid and citric acid; hair conditioners such as cationized cellulose, cationic surface active agents, glycerine ethylene glycol esters and carboxylic acid esters; and perfumes, colorants, UV absorbers, antioxidants, anti-dandruff agents, bactericides and preservatives.

The present invention is more particularly described by way of examples, which should not be construed as limiting the present invention.

Performances of shampoo compositions in examples were evaluated according to the following methods.

1) Foaming Test Method

(a) To an aqueous 1% solution of shampoo composition was added 0.5% of artificial soil, followed by agitating in a cylinder container at 40°C for 5 minutes by means of a flat propeller under conditions where the propeller was rotated at a rotation frequency of 1000 r.p.m. and reversed every 10 seconds. Thirty seconds after completion of the agitation, an amount of foams was measured for evaluation. The amount of foams is determined by foam height in cylinder container and is an indication of forming power.

(b) Organoleptic Evaluation Using Blanket

Twenty five grams of wool cloth was soaked with hot water (40°C), on which was applied 0.5 g of a shampoo composition, followed by kneading for about 2 minutes. The amount of foams was organoleptically evaluated.

Evaluation Standard of Foaming

- ⊙ Good
- Moderate

△ Rather poor
X Poor

2) Detergency Test Method

(A) Artificially Soiled Cloth

An artificial oily soil was uniformly dispersed in solvent. #2003 cloth was brought to contact with the dispersion and dried to uniformly deposit the artificial oily soil thereon. The cloth was cut into 10 cm × 10 cm test pieces used for subsequent experiments.

(B) Washing Conditions and Method

In water of 4°DH was dissolved a shampoo composition to prepare one liter of a 3% shampoo aqueous solution. Five pieces of the artificially soiled cloth and the aqueous shampoo solution were placed, as they are, in a stainless beaker for targetmeter, followed by agitating at 100 r.p.m. at 40°C for 10 minutes in the targetmeter. The pieces were each rinsed with running water, pressed with an iron and subjected to measurement of reflectivity.

(C) Calculation of Detergency Rate

The detergency rate was calculated according to the following equation.

The reflectivities of the original cloth prior to washing and before and after washing of soiled cloth at 460 mμ were measured by the use of an autographic recording colorimeter, from which the detergency rate (%) was calculated according to the following equation.

$$\text{Detergency rate (\%)} = \frac{\begin{array}{c} \text{Reflectivity} \\ \text{after} \\ \text{washing} \end{array} - \begin{array}{c} \text{Reflectivity} \\ \text{before} \\ \text{washing} \end{array}}{\begin{array}{c} \text{Reflectivity} \\ \text{of} \\ \text{original cloth} \end{array} - \begin{array}{c} \text{Reflectivity} \\ \text{before} \\ \text{washing} \end{array}} \times 100$$

The rate was indicated as an average of five cloths. The pH of the aqueous shampoo solution prior to washing was found to be 5.5.

(D) Extraction of Residual Lipid

The soiled cloth after the washing was subjected to extraction for 4 hours by the Soxhlet extraction method using an extraction solvent of chloroform/methanol (50% v/v). Thereafter, the extract was dried under reduced pressure using a rotary evaporator.

(E) Calculation of Residue of Triglyceride

To the extract obtained by drying in (D) was added a given amount of chloroform for dissolution, after which a predetermined amount of the solution was spottily applied onto a silica gel plate (20 cm × 20 cm, using Wako gel B5) for thin-layer chromatography, made by ourselves, followed by developing with a first developer solution of benzene/hexane (50/50% v/v) to about 18 cm of the plate. Subsequently, it was developed with a second developer solution of hexane/diethyl ether/acetic acid (70/30/1% v/v) to about 12 cm of the plate, followed by spraying 50% v/v sulfuric acid as usual and heating for color development. Thereafter, residual triglyceride was quantitatively determined using a densitometer (made by Shimadzu Seisakusho), from which the residual rate was calculated.

Example 1

Shampoo compositions of the following formulations were prepared to evaluate their performances by the foaming and detergency tests. The results are shown in Table 1.

Table 1

Composition/ Evaluation	Compara- tive Product 1	Compara- tive Product 2	Compara- tive Product 3	Product 1 of Invention
Sodium Polyoxyethylene (2.5) Lauryl Ether Sulfate	10	10	10	10
Polyoxyethylene (80) Hardened Castor Oil	-	5	-	5
Polyoxyethylene (4) Adduct of Lauric Alkanolamide	-	-	5	-
Sodium N-lauroyl- N-methyl-beta- alanine	-	-	5	5
Phosphoric Acid	suitable amount	suitable amount	suitable amount	suitable amount
pH (5% Aqueous Solution)	5.5	5.5	5.5	5.5
Foaming Power (mm)				
Artificial Soil Added	109	32	65	98
Foaming Power Organoleptic Evaluation Using Wool Cloth	0	x	△	0
Detergency Rate (%)	48.8	44.3	47.5	51.3
Residue of Triglyceride (%)	10.3	7.9	9.4	7.0

Example 2

Shampoo compositions of the following formulations were prepared to evaluate the foaming strength and detergency rate. The results are shown in Table 2.

EP 0 145 865 B1

Table 2

Composition/ Evaluation	Compara- tive Product 4	Compara- tive Product 5	Compara- tive Product 6	Compara- tive Product 7	Compara- tive Product 8	Product 2 of Invention
Sodium Polyoxyethylene (2.5) Lauryl Ether Sulfate	12	12	12	12	12	12
Diethanolamine Laurate	3	3	3	3	3	3
Polyoxyethylene (3) Alkyl(*) Ether	-	-	0.8	0.8	0.8	0.8
Polyoxyethylene (120) Hardened Castor Oil	-	0.8	0.8	0.8	0.8	0.8
Sodium N-lauroyl- N-methyl-beta- alanine	-	-	-	-	-	3
Sodium N- lauroylsarcosine	-	-	-	3	-	-
Sodium N- lauroylglutamate	-	-	-	-	3	-
Phosphoric Acid	suitable amount	suitable amount	suitable amount	suitable amount	suitable amount	suitable amount
pH	5.5	5.5	5.5	5.5	5.5	5.5
Foaming Power (mm) No Artificial Soil Added	230	220	210	220	220	230
Foaming Power (mm) Artificial Soil Added	120	101	72	103	92	119
Foaming Power Organoleptic Evaluation Using Wool Cloth	⊙~0	△	x	△	△	⊙~0
Detergency Rate (%)	48.3	49.2	50.0	51.0	50.4	54.0
Residue of Triglyceride (%)	10.9	7.1	6.7	6.0	6.9	3.9

(Note) * Linear and branched alkyl group having from 11 to 15 carbon atoms.

EP 0 145 865 B1

Example 3

Shampoo compositions of the following formulations were prepared for evaluation by the foaming test. The results are shown in Table 3.

Table 3

Composition/ Evaluation	A	B	C	D	E	F	G
Sodium Polyoxyethylene (2.5) Lauryl Ether Sulfate	15	15	15	15	15	15	15
Lauric ethanolamide	2.5	2.5	2.5	2.5	2.5	2.5	2.5
Polyoxyethylene (3) Alkyl(*) Ether	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Polyoxyethylene (60) Hardened Castor Oil	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Sodium N-lauroyl- N-methyl-beta- alanine	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Phosphoric acid	suit- able amount	suit- able amount	suit- able amount	suit- able amount	suit- able amount	suit- able amount	suit- able amount
pH	5.0	5.5	6.0	6.5	7.0	7.5	8.0
Foaming Power (mm)							
Artificial Soil Added	98	134	138	130	124	124	120
Foaming Power Organoleptic Evaluation Using	△	⊙	⊙	⊙	⊙~0	⊙~0	0

(Note) * Same as in Table 2.

Claims for the Contracting States: CH DE GB LI NL

1. A shampoo composition of the type which comprises an anionic surface active agent and/or amphoteric surface active agent as a base, characterized by further comprising:
(A) 0.5 to 10 wt% of an N-acyl-N-methyl-beta-alanine or its salt represented by the formula



in which R represents a linear or branched, saturated or unsaturated hydrocarbon group having from 7 to



EP 0 145 865 B1

Example 3

Shampoo compositions of the following formulations were prepared for evaluation by the foaming test. The results are shown in Table 3.

Table 3							
Composition/ Evaluation	A	B	C	D	E	F	G
Sodium Polyoxyethylene (2.5) Lauryl Ether Sulfate	15	15	15	15	15	15	15
Lauric ethanolamide	2.5	2.5	2.5	2.5	2.5	2.5	2.5
Polyoxyethylene (3) Alkyl(*) Ether	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Polyoxyethylene (60) Hardened Castor Oil	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Sodium N-lauroyl- N-methyl-beta- alanine	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Phosphoric acid	suit- able amount	suit- able amount	suit- able amount	suit- able amount	suit- able amount	suit- able amount	suit- able amount
pH	5.0	5.5	6.0	6.5	7.0	7.5	8.0
Foaming Power (mm)							
Artificial Soil Added	98	134	138	130	124	124	120
Foaming Power Organoleptic Evaluation Using	△	⊙	⊙	⊙	⊙~0	⊙~0	0

(Note) * Same as in Table 2.

Claims for the Contracting States: CH DE GB LI NL

1. A shampoo composition of the type which comprises an anionic surface active agent and/or amphoteric surface active agent as a base, characterized by further comprising:

(A) 0.5 to 10 wt% of an N-acyl-N-methyl-beta-alanine or its salt represented by the formula



in which R represents a linear or branched, saturated or unsaturated hydrocarbon group having from 7 to




EP 0 145 865 B1

- 21 carbon atoms, and M represents hydrogen, an alkali metal, ammonium, alkyl-substituted ammonium with an alkyl moiety having from 1 to 3 carbon atoms, alkanolamine having a hydroxyalkyl group having from 1 to 3 carbon atoms, or basic amino acid, and (B) 0.5 to 20 wt% of one or more of ethylene oxide-added type nonionic surface active agents selected from polyoxyethylene alkyl or alkenyl ethers, polyoxyethylene alkylphenyl ethers, polyoxyethylene fatty acid esters, and polyoxyethylene hardened castor oils.
2. The shampoo composition according to Claim 1, wherein the ingredient (a) is N-lauroyl-N-methyl-beta-alanine or its sodium salt or triethanolamine salt.
3. The shampoo composition according to Claim 1 or 2, wherein the composition has a pH ranging from 5.5 to 7.5.

Claims for the Contracting State: AT

1. A method for manufacturing a shampoo composition of the type which comprises an anionic surface active agent and/or amphoteric surface active agent as a base, characterized by adding the following two ingredients (A) and (B)
- (A) 0.5 to 10 wt% of an N-acyl-N-methyl-beta-alanine or its salt represented by the formula




(I)

- in which R represents a linear or branched, saturated or unsaturated hydrocarbon group having from 7 to 21 carbon atoms, and M represents hydrogen, an alkali metal, ammonium, alkyl-substituted ammonium with an alkyl moiety having from 1 to 3 carbon atoms, alkanolamine having a hydroxyalkyl group having from 1 to 3 carbon atoms, or basic amino acid, and (B) 0.5 to 20 wt% of one or more of ethylene oxide-added type nonionic surface active agents selected from polyoxyethylene alkyl or alkenyl ethers, polyoxyethylene alkylphenyl ethers, polyoxyethylene fatty acid esters, and polyoxyethylene hardened castor oils.
2. Method according to Claim 1, wherein the ingredient (a) is N-lauroyl-N-methyl-beta-alanine or its sodium salt or triethanolamine salt.
3. Method according to Claim 1 or 2, wherein the composition has a pH ranging from 5.5 to 7.5.

Patentansprüche für die Vertragsstaaten: CH DE GB LI NL

1. Schampoozusammensetzung des Typs, bei dem ein anionisches oberflächenaktives Mittel und/oder amphoterer oberflächenaktives Mittel als eine Basis vorliegt, dadurch gekennzeichnet, daß die Zusammensetzung ferner umfasst:
- (A) 0,5 bis 10 Gew.-% eines N-Acyl-N-methyl-beta-alanins oder dessen Salz gemäß der Formel



(I)

- in der R für eine lineare oder verzweigte, gesättigte oder ungesättigte Kohlenwasserstoffgruppe mit 7 bis 21 Kohlenstoffatomen steht und M Wasserstoff, ein Alkalimetall, Ammonium, Alkyl-substituiertes Ammonium mit einer Alkyleinheit mit 1 bis 3 Kohlenstoffatomen, Alkanolamin mit einer Hydroxyalkylgruppe mit 1 bis 3 Kohlenstoffatomen oder eine basische Aminosäure bedeutet und
- (B) 0,5 bis 20 Gew.-% von einem oder mehreren der nicht-ionischen oberflächenaktiven Mittel des Ethylenoxidaddierten Typs, ausgewählt aus Polyoxyethylenalkyl- oder -Alkenylethern, Polyoxyethylenalkylphenylethern, Polyoxyethylefettensäureestern und Polyoxyethylen-gehärtete Rhizinusöle.
2. Schampoozusammensetzung gemäß Anspruch 1, wobei der Bestandteil (A) N-Lauroyl-N-methyl-beta-alanin oder dessen Natriumsalz oder dessen Triethanolaminsalz ist.
3. Schampoozusammensetzung gemäß Anspruch 1 oder 2, wobei die Zusammensetzung einen pH im Bereich von 5,5 bis 7,5 aufweist.

Patentansprüche für den Vertragsstaat: AT

1. Verfahren zur Herstellung einer Schampoozusammensetzung des Typs, bei dem ein anionisches oberflächenaktives Mittel und/oder amphoterer oberflächenaktives Mittel als eine Basis vorliegt, dadurch



EP 0 145 865 B1

gekennzeichnet, daß man die folgenden beiden Bestandteile (A) und (B) zusetzt.

(A) 0,5 bis 10 Gew.% eines N-Acyl-N-methyl-beta-alanins oder dessen Salz gemäß der Formel



in der R für eine lineare oder verzweigte, gesättigte oder ungesättigte Kohlenwasserstoffgruppe mit 7 bis 21 Kohlenstoffatomen steht und M Wasserstoff, ein Alkalimetall, Ammonium, Alkyl-substituiertes Ammonium mit einer Alkyleinheit mit 1 bis 3 Kohlenstoffatomen, Alkanolamin mit einer Hydroxyalkylgruppe mit 1 bis 3 Kohlenstoffatomen oder eine basische Aminosäure bedeutet und

(B) 0,5 bis 20 Gew.% von einem oder mehreren der nicht-ionischen oberflächenaktiven Mittel des Ethylenoxidaddierten Typs, ausgewählt aus Polyoxyethylenalkyl- oder -Alkenylethern, Polyoxyethylenalkylphenylethern, Polyoxyethylefetsäureestern und Polyoxyethylen-gehärtete Rhizinusöle.

2. Verfahren gemäß Anspruch 1, wobei der Bestandteil (A) N-Lauroyl-N-methyl-beta-alanin oder dessen Natriumsalz oder dessen Triethanolaminsalz ist.

3. Verfahren gemäß Anspruch 1 oder 2, wobei die Zusammensetzung einen pH im Bereich von 5,5 bis 7,5 aufweist.

Revendications pour les Etats contractants: CH DE GB LI NL

1. Composition de shampoing du type qui comprend un agent tensio-actif anionique et/ou un agent tensio-actif amphotère en tant que base, caractérisée par le fait qu'elle comprend en outre:

(A) 0,5 à 10% en poids d'une N-acyl-N-méthyl-bêta-alanine, ou d'un sel de cette dernière, représentée par la formule:



dans laquelle:

R représente un groupe hydrocarboné saturé ou insaturé, linéaire ou ramifié, ayant de 7 à 21 atomes de carbone; et

M représente hydrogène, un métal alcalin, ammonium, ammonium substitué par alkyle, avec une fraction alkyle ayant de 1 à 3 atomes de carbone, alcanolamine présentant un groupe hydroxyalkyle ayant de 1 à 3 atomes de carbone, ou acide aminé basique, et

(B) 0,5 à 20% en poids d'un ou plusieurs agents tensio-actifs non-ioniques du type à oxyde d'éthylène ajouté, choisi parmi les polyoxyéthylène alkyle ou alcényl éthers, les polyoxyéthylène alkylphenyl éthers, les esters d'acides gras polyoxyéthylénés et les huiles de ricin durcies polyoxyéthylénées.

2. Composition de shampoing selon la revendication 1, dans laquelle l'ingrédient (A) est la N-lauroyl-N-méthyl-bêta-alanine, ou son sel de sodium ou son sel de triéthanolamine.

3. Composition de shampoing selon la revendication 1 ou 2, dans laquelle la composition a un pH se situant dans la plage de 5,5 à 7,5.

Revendications pour l'Etat contractant: AT

1. Procédé de fabrication d'une composition de shampoing du type qui comprend un agent tensio-actif anionique et/ou un agent tensio-actif amphotère en tant que base, caractérisé par l'addition des deux ingrédients (A) et (B) suivants:

(A) 0,5 à 10% en poids d'une N-acyl-N-méthyl-bêta-alanine, ou d'un sel de cette dernière, représentée par la formule:





EP 0 145 865 B1

dans laquelle:

R représente un groupe hydrocarboné saturé ou insaturé, linéaire ou ramifié, ayant de 7 à 21 atomes de carbone; et

M représente hydrogène, un métal alcalin, ammonium, ammonium substitué par alkyle, avec une fraction alkyle ayant de 1 à 3 atomes de carbone, alcanolamine présentant un groupe hydroxyalkyle ayant de 1 à 3 atomes de carbone, ou acide aminé basique, et

(B) 0,5 à 20% en poids d'un ou plusieurs agents tensio-actifs non-ioniques du type à oxyde d'éthylène ajouté, choisi parmi les polyoxyéthylène alkyle ou alcényl éthers, les polyoxyéthylène alkylphenyl éthers, les esters d'acides gras polyoxyéthylénés et les huiles de ricin durcies polyoxyéthylénées.

2. Procédé selon la revendication 1, dans lequel l'ingrédient (A) est la N-lauroyl-N-méthyl-bêta-alanine, ou son sel de sodium ou son sel de triéthanolamine.

3. Procédé selon la revendication 1 ou 2, dans lequel la composition a un pH se situant dans la plage de 5,5 à 7,5.