

Description**BACKGROUND OF THE INVENTION**

5 1) Field of the Invention

This invention relates to a novel cellulase, a CMCase isolated from the cellulase, and a microorganism capable of producing the cellulase.

10 2) Description of the Prior Art

Cellulases consist of a complicated enzymatic system serving as a catalyst for an enzymatic reaction in which cellulose and similar polysaccharides are decomposed into glucose, cellobiose or celloligosaccharides. Cellulases are considered to be a general name for enzymes which are called, depending upon the mechanism of activity, C₁ enzyme, C_x enzyme and beta-glucosidase, or exo-beta-glucanase, endo-beta-glucanase, cellobiase and the like. Over a long term, cellulases have been studied mainly for the purpose of effectively utilizing biomass resources. For instance, the source of cellulase supply has relied mainly on fungi belonging to the genera *Trichoderma*, *Aspergillus*, *Acremonium*, and *Humicola*. However, the cellulases deriving their origin from microorganisms including the fungi involve a diversity of working specificities and physicochemical properties of constituting enzyme groups, which have not been completely clarified yet.

Of these cellulases, those cellulases which have the high action on carboxymethyl cellulose (CMC) or C_x enzymatic action are generally called CMCases. In recent years, novel industrial utility of cellulases including the CMCases have been developed, particularly, as an additive for use in detergent compositions for clothing. However, so far as cellulases produced by microorganism in the natural field and, particularly, the above-mentioned cellulases originating from microorganism are concerned, they are, in most cases, so unstable that their activity is lost in an alkaline pH range and are so-called acidic and neutral cellulases (whose optimum working pH is in the range of 4 to 6). So-called alkaline cellulases which meet the requirement for detergent compositions for clothing, i.e. they have a maximum activity and are resistant in an alkaline range, are very small in number.

For instance, with regard to alkaline cellulases which are usable in detergent compositions for clothing, there are known only several methods of producing alkaline cellulases originating from alkalophilic microorganisms. These methods include a method in which microorganisms belonging to the genus *Bacillus* are cultivated and cellulase A is collected from the medium (Japanese Patent Publication No. 50-28515), a method in which alkalophilic bacteria belonging to the genus *Cellulomonas* are cultivated to produce alkaline cellulase 301-A (Japanese Patent Application Laid-open No. 58-224686), a method for producing CMCase by cultivation of alkalophilic *Bacillus* No. 1139 (Horikoshi et al, J. Gen. Microbiol., Vol. 131, page 3339 (1985)), and a method of producing an alkaline cellulase using a strain belonging to the genus *Streptomyces* (Japanese Patent Application Laid-open No. 61-19483).

Accordingly, there is a demand for an alkaline cellulase which has an optimum pH and a function as an enzyme for detergents for clothing in an alkaline range.

SUMMARY OF THE INVENTION

The present inventors made intensive studies to find, in the natural field, microorganism capable of producing alkali cellulases and found that a microorganism collected from a sample of soil of Haga-gun, Tochigi-ken, Japan and belonging to the genus *Bacillus* (Strain KSM-635, deposited as FERM BP-1485) was able to produce novel alkaline cellulase K effective as an additive for detergent compositions for clothing. Moreover, it was also found that when the alkaline cellulase was further purified, novel CMCases I and II were obtained as main components.

Accordingly, the invention provides a novel alkaline cellulase K enzyme as defined in claim 1 having an optimum pH at an alkaline side.

Another object of the invention is to provide novel CMCase I and II enzymes as defined in claim 3 and 4.

A further object of the invention is to provide said microorganism capable of producing these novel enzymes and a method for producing alkaline cellulase K by cultivating the microorganism and collecting the resulting alkaline cellulase K from the culture product.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph showing a growth temperature range of *Bacillus* sp KSM-635; (FERM BP-1485);
 Fig. 2 is a graph showing a growth pH range;
 Fig. 3 is a graph showing a working pH range of CMCase of alkaline cellulase K;
 Fig. 4 is a graph showing pH stability;
 Fig. 5 is a graph showing a working temperature range of CMCase of alkaline cellulase K;
 Fig. 6 is a graph showing thermal stability;
 Fig. 7 is a graph showing the relation between elution fraction and activity in gel chromatography;
 Fig. 8 is a graph showing the relation between pH at which CMCase I is reacted and relative activity;
 Fig. 9 is a graph showing the relation between pH at which CMCase I is treated and residual activity;
 Fig. 10 is a graph showing the relation between reaction temperature (pH 9.0) of CMCase I and relative activity;
 Fig. 11 is a graph showing the relation between treating temperature of CMCase I (pH 9.0) and residual activity;
 Fig. 12 is a UV absorption spectrum of CMCase I of the invention;
 Fig. 13 is a graph showing the relation between reaction pH for CMCase II and relative activity;
 Fig. 14 is a graph showing the relation between treating pH of CMCase II and residual activity;
 Fig. 15 is a graph showing the relation between reaction temperature of CMCase II (pH 9.0) and relative activity;
 Fig. 16 is a graph showing the relation between treating temperature for CMCase II (pH 9.0) and residual activity;
 Fig. 17 is a UV absorption spectrum of CMCase II; and
 Fig. 18 is a view showing the results of sodium dodecylsulfate electrophoresis of CMCases I and II of the invention.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

The microorganism used for producing the novel enzymes according to the invention has the following mycological properties. The media used for classification of strains are those media 1 to 24 and contain a suitable amount of a sterilized 1.0 wt% sodium carbonate (Na_2CO_3) solution unless otherwise indicated.

Compositions of Test Media for Classification (wt%)

Medium 1:	Bacto peptone 0.5; meat extract 0.3; Bacto agar-agar 1.5
Medium 2:	Bacto peptone 0.5; meat extract 0.3
Medium 3:	Bacto peptone 0.5; meat extract 0.3; NaCl 7.0
Medium 4:	Bacto peptone 0.5; meat extract 0.3; Bacto gelatin 20.0
Medium 5:	Bacto litmus milk 10.5
Medium 6:	Bacto peptone 0.5; meat extract 0.3; KNO_3 0.1
Medium 7:	Bacto peptone 0.7; glucose 0.5; NaCl 0.5
Medium 8:	Bacto peptone 3.0; meat extract 0.3; sodium thiosulfate 0.005; cysteine hydrochloride 0.02; iron ammonium citrate 0.05; Bacto agar-agar 0.5
Medium 9:	Bacto peptone 1.5; meat extract 0.4; lactose 1.0; sucrose 1.0; glucose 1.0; NaCl 0.5; sodium thiosulfate 0.008; sodium sulfite 0.04; ferrous sulfate 0.02; phenol red 0.002; Bacto agar-agar 1.5
Medium 10:	Bacto peptone 1.5; yeast extract 0.5; soluble starch 2.0; K_2HPO_4 0.1; Bacto agar-agar 1.5; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02
Medium 11:	ammonium phosphate 0.1; KCl 0.02; yeast extract 0.05; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02; sugars 1.0 (sterilized separately through filtration)
Medium 12:	potassium monohydrogen phosphate 0.1; ammonium dihydrogen phosphate 0.1; sodium citrate 0.2; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.03; NaCl 0.5; Bromothymol Blue 0.0024; Bacto agar-agar 1.5
Medium 13:	yeast extract 0.05; cysteine hydrochloride 0.01; sodium citrate 0.3; NaCl 0.5; sodium thiosulfate 0.008; iron ammonium citrate 0.04; glucose 0.02; potassium dihydrogen-phosphate 0.15; phenol red 0.0012; Bacto agar-agar 1.5
Medium 14:	ammonium phosphate 0.1; potassium dihydrogenphosphate 0.05; sodium citrate 0.2; Bacto agar-agar 1.5; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02
Medium 15:	yeast extract 0.05; Na_2SO_4 0.1; KH_2PO_4 0.1; glucose 1.0; inorganic nitrogen sources

- suitable amounts"
- Medium 16: yeast extract 0.05; Na_2SO_4 0.1; KH_2PO_4 0.1; glucose 1.0; inorganic nitrogen sources suitable amounts"; $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 0.05; $\text{MnSO}_4 \cdot 4-8\text{H}_2\text{O}$ 0.001; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 0.001 (sterilized separately through filtration); $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02 (sterilized separately through filtration)
- Medium 17: King A medium "Eiken" (available from Eiken Chem Co., Ltd.), indicated amounts
- Medium 18: King B medium "Eiken" (available from Eiken Chem. Co., Ltd.), indicated amounts
- Medium 19: potato dextrose agar-agar medium "Eiken" (Eiken Chem. Co., Ltd.), indicated amounts
- Medium 20: Bacto peptone 0.25; salt 0.25; yeast extract 0.25; mannitol 0.5; Bacto agar-agar 2.0
- Medium 21: urea medium "Eiken" (available from Eiken Chem. Co., Ltd.), indicated amounts
- Medium 22: Bacto peptone 0.1; NaCl 0.5; KH_2PO_4 0.2; yeast extract 0.05; glucose 0.1; urea 2.0; phenol red 0.001
- Medium 23: Bacto peptone 0.5; yeast extract 0.5; K_2HPO_4 0.1; glucose 1.0; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02
- Medium 24: yeast extract 0.5; glucose 1.0; casein (Hammerstein, Merc Inc.) 0.5; K_2HPO_4 0.1; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02; Bacto agar-agar 1.5

(Mycological Properties)

1. Results of Microscopic Observation

The *Bacillus* has a size of 0.5 to 1.2 μm x 1.5 to 4.0 μm and make an endospore (0.7 to 1.2, μm x 1.0 to 2.0 μm) at one end of the bacillus body. Periflagella Motility: positive. Gram staining: positive.

2. The state of growth in various media

- (1) Meat broth and agar-agar medium (medium 1)

The colony is circular in shape and is flat on the surface thereof. The colony is white or yellow in color and is semi-transparent and glossy.

- (2) Meat broth liquid medium (medium 2)

Growing and becoming turbid.

- (3) 7% Salt meat broth liquid medium (medium 3)

Growing and becoming turbid.

- (4) Meat broth-gelatin stab culture (medium 4)

Not growing.

- (5) Litmus milk culture (medium 5)

Coagulation and peptonization of the milk: negative. Because of the alkalinity of the medium 5, no change in color of the litmus is recognized.

3. Physiological Properties

- (1) Reduction of nitrates and denitrification reaction

Reduction of nitric acid: positive.

Denitrification: negative (medium 6).

- (2) MR test (medium 7)

Because of the alkalinity of the medium, Methyl Red does not undergo any change, thus the judgement being impossible.

- (3) VP test (medium 7)

Positive.

- (4) Formation of indole (medium 8)

Negative with respect to the reaction against an indole-producing test filter paper ("Nissan", made by Nissui Seiyaku K.K.) and also to the color change with Kovacs' indole reagent.

* Sodium nitrate was added in 0.25%, sodium nitrite was in 0.2025%, ammonium chloride was in 0.158%, and ammonium phosphate was in 0.195% (each corresponding to 0.0412 N%).

** Sodium nitrate was added in an amount of 0.25%, sodium nitrite was in an amount of 0.2025%, ammonium chloride was added in an amount of 0.158%, and ammonium phosphate was in an amount of 0.195% (each corresponding to 0.0412 N%).

(5) Formation of hydrogen sulfide (medium 9)

Negative.

(6) Hydrolysis of starch

A plate agar-agar medium, i.e. medium 10, which has been rendered acidic by the use of 4 N HCl is negative when determined by an ordinary detection method using the iodine reaction. In the liquid medium 11, formation of soluble starch is negative.

(7) Utility of citric acid

Medium 11: positive.

Medium 12 (Koser's (Simons') citric acid/agar-agar plate medium): negative. A liquid medium which is obtained by eliminating agar-agar from the medium 12 and, instead, 0.05% of yeast extract is added, is positive.

Medium 13 (Christensen's agar-agar plate medium): positive, but no change with phenol red is observed because of the alkalinity.

Medium 14: negative (not growing). Positive with respect to a liquid medium which is obtained by eliminating agar-agar from the medium 14 and adding 0.05% of yeast extract.

(8) Utilization of inorganic nitrogen sources

Medium 15: negative to pseudopositive for all of nitric acid, nitrous acid and ammonia.

Medium 16: the present medium containing very small amounts of metal salts is positive with respect to nitric acid and nitrous acid. Ammonium chloride: pseudopositive. Ammonium phosphate: positive.

(9) Formation of dyes

Not growing in the King A medium (medium 17) and not possible to judge.

Light yellow (involving no fluorescence) in the King B medium (medium 18). Growing in a potato/dextrose/agar-agar medium (medium 19) and in a mannitol/yeast extract/agar-agar medium (medium 20) but negative with respect to the formation of dyes.

(10) Urease

Medium 21: not growing. Negative when the formation ammonia with Nessler's reagent was confirmed after removal of phenol red from the medium 21 and cultivation of the present organism.

Medium 22 (Christensen's urea medium to which yeast extract is added): negative when the formation of ammonia with Nessler's reagent was confirmed after removal of phenol red from the medium and cultivation of the present organism. Negative when tested by changing the urea concentration in the medium 22 to 0.1, 0.2, 0.5, 1.0 and 2.0% on the presupposition of cytotoxin.

(11) Oxidase

"Positive" or "negative" is not clear.

(12) Catalase

Positive.

(13) Range of Growth (medium 23)

A temperature gradient incubator was used for shaking culture in an L-type test tube for 3 days. The growing temperature range was from 20 to 45 °C and an optimum growing temperature range was from 29 to 37 °C (Fig. 1).

In order to determine a pH range for growth, a test was conducted in which the concentration of Na_2CO_3 in the medium 23 was changed to change an initial pH of the medium. As a result it was found that the growing pH range was from 8 to 11 and an optimum growing pH range was from 9.5 to 10.2 (Fig. 2). On the other hand, when the pH of the medium was controlled by the use of K_2CO_3 , it was found that the growing amount was very small and the optimum growing pH was about 9.

(14) Behavior on oxygen

Aerobic.

(15) O-F test

Undergoing no change in color because of the alkalinity and growing only in an aerobic condition.

(16) Utility of Sugars (medium 11)

Utilizable carbon sources: D-ribose, L-arabinose, D-xylose, D-glucose, D-mannose, D-fructose, maltose, sucrose, trehalose, mannitol, inositol, and glycerin.

Non-utilizable carbon sources: D-galactose, lactose, sorbitol, starch, dextrin, and raffinose.

(17) Hydrolysis of casein (medium 24)

Microorganism was grown up in an agar-agar plate culture, into which 30% trichloroacetic acid was poured for judgement, revealing that any transparent zone was not formed around the bacillus colonies, from which it was judged as negative.

(18) Requirement for nutrients

As is shown in Table 1, biotin (desthiobiotin) is essential for the growth.

Table 1

Concentration of Vitamin ($\mu\text{g/ml}$)		Degree of Growth of <i>Bacillus</i> sp KSM-635 (Absorbance, 600 nm/6 days)
Biotin	Desthiobiotin	
0.001	0	4.6
0.01	0	4.6
0	0.001	4.3
0	0.01	4.4
nil		0.0

Reference is made to Bergey's Manual of Determinative Bacteriology, eighth edition, with respect to the above mycological properties, with the result that the present strain is considered as a spore-forming microorganism belonging to the genus *Bacillus*. However, since the present strain does not grow in a neutral pH but grows well only at a highly alkaline pH range, it can be provisionally determined as an alkalophilic microorganism, which has been recently reported by Horikoshi and Akiba in "Alkalophilic Microorganisms", Japan Scientific Society Press (Tokyo), 1982, and is distinguished from known bacilli growing in neutrality.

Because the mycological properties of the present strain do not coincide with those of known alkalophilic bacilli, the present strain has been determined as a novel strain and named *Bacillus* sp KSM-635, and is now deposited as FERM No. 8872 at the Fermentation Research Institute of Japan.

For the production of alkaline cellulase K by the use of the above novel microorganism, *Bacillus* sp KSM-635, the strain of *Bacillus* sp KSM-635 or its variant is cultivated in a medium to produce alkaline cellulase K. This product is subjected to an ordinary enzyme purification method to isolate and purify it.

For the production by fermentation of alkaline cellulase K, a suitable medium is sterilized such as by heating and the strain of *Bacillus* sp KSM-635 (FERM BP-1485) is inoculated in the medium, followed by shaking or aeration spinner cultivation at 22 to 40°C, preferably from 26 to 37°C, for 1 to 4 days. Good results are obtained when the pH is adjusted to 8 to 11. Since the production-by-fermentation medium is alkaline in nature, it may sometimes be expanded, which will be solved by the addition of a suitable amount of an antifoamer at suitable times.

The production of the alkaline cellulase K requires a suitable combination of nitrogen sources and carbon sources contained in a culture medium. These nutrient sources are not critical. For instance, nitrogen sources include inorganic sources such as ammonium nitrate, ammonium sulfate, ammonium chloride, ammonium phosphate, sodium nitrate and the like, and corn gluten meal, bean flour, corn steep liquor, casamino acid, yeast extract, Pharmamedia, sardine meal, meat extract, peptone, Hypro, Ajipower, corn soybean meal, coffee grounds, cotton seed oil meal, Cultivator, Amiflex, Ajipron, Zest, Ajix, and the like. Examples of the carbon sources include plant fibers such as chaff, bran, filter paper, ordinary papers, sawdust and the like, and waste molasses, invert sugar, CMC, Avicel, cellulosic cotton, xylan, pectin and the like. Further utilizable carbon sources include, for example, ribose, arabionse, xylose, glucose, mannose, fructose, maltose, sucrose, trehalose, mannitol, inositol, glycerin and the like, and organic acids such as acetic acid, citric acid and the like. Any media using combinations of these nitrogen and carbon sources may be used and the nutrient sources should not be limited to any specific ones. Aside from these ingredients, phosphoric acid and inorganic salts such as of Mg^{2+} , Ca^{2+} , Mn^{2+} , Zn^{2+} , Co^{2+} , Na^+ , K^+ and the like, and, if necessary, inorganic and organic trace nutrient sources may be suitably added to the medium.

The alkaline cellulase K may be obtained from the thus obtained culture product according to any known techniques of collection and purification of ordinary enzymes as will be particularly described in examples appearing hereinafter.

For instance, the culture product may be subjected to centrifugal separation or ultrafiltration to separate the bacillus cells. The resultant culture broth are subjected to ordinary isolation methods including, for example, salting-out, isoelectric precipitation, solvent precipitation (using methanol, ethanol, isopropanol or the like), thereby separating the protein as a precipitate. Alternatively, ultrafiltration (e.g. Diaflow Membrane YC, available from Amicon Co., Ltd.) may be used for concentration to obtain alkaline cellulase K. After precipitation in ammonium sulfate (30 to 70% saturated fraction) for the salting-out method or 75% ethanol

for the solvent precipitation method, the enzyme may be filtered, centrifugally separated or desalted to obtain a freeze-dried powder. The desalting may be effected by an ordinary procedure such as dialysis or gel filtration using Sephadex G-25 or the like. Moreover, the purification of the enzyme is possible by a suitable combination of, for example, a hydroxy apatite chromatography, an ion exchange chromatography using DEAE-Sephadex or DEAE-cellulose and a molecular sieve gel chromatography using Sephadex or Bio-gel.

The resultant alkaline cellulase K of the invention have the following physical and chemical properties.

(1) Activity

The present enzyme has a Cx enzymatic activity of acting on CMC. However, it also acts on phosphoric acid-swollen cellulose and has such action specificities as exhibiting an activity as an enzyme acting on crystalline cellulose (cellulosic cotton) or Avicel which is cellulose having high crystallinity (i.e. Avicelase), and an activity of a C₁ enzyme typical of which is a filterpaper-degrading activity (FPDase), and a beta-glucosidase activity on cellobiose and celooligosaccharide. In addition, the enzyme slightly acts on the artificial substrate of PNPC to liberate p-nitrophenol.

(2) Substrate specificity

The alkaline cellulase K has no degradation activity on xylan, amylose, dextrin, pectin, inulin and cardolan. The Avicelase and FPDase activities are about 0.3% of the CMCase activity. The degradation activity on the artificial substrate of p-nitrophenyl cellobioside (PNPC) was about 1.5 to 1.8% of the CMCase activity (Tabel 2).

Table 2

Action of Enzyme	Specfic Activity (units/g of enzyme)
β -Glucosidase	0.7
PNPCase*	5.2
CMCase	298
FPDase	1.2
Avicelase	1.1

* PNPC degrading activity

(3) Working pH and optimum working pH

The working pH of the present enzyme is from 4 to 12 and the optimum working pH is approximately from 9 to 10. At about a pH of 10.5, there appears a shoulder (Fig. 3).

(4) pH stability

Stable pH values, which are determined by allowing to stand in buffer solutions of different pHs at 40 °C for 10 minutes and 30 minutes, are, respectively, from 4.5 to 10.5 and 6.8 to 10 (Fig. 4). When allowed to stand at 5 °C, the pH is 4 to 11 and is stable over at least one month.

(5) Working temperature range and working optimum temperature

The present enzyme works in a wide temperature range of from 10 °C to 65 °C. When the reaction is effected in a glycine buffer solution (pH 9) for 20 minutes, the working optimum temperature is found to be about 40 °C (Fig. 5).

(6) Thermal stability

When thermally treated in a glycine buffer solution (pH 9) at different temperatures for 20 minutes, the present enzyme is not inactivated at all at about 40 °C, and has a residual activity of about 50% at 60 °C

and about 25% at 70 °C (Fig. 6).

(7) Measuring methods of enzymatic activities and proteins

(i) CMCase activity

0.1 ml of an enzyme solution was added to a substrate solution composed of 0.2 ml of CMC (2.5%), 0.1 ml of a 0.5M glycine buffer solution (pH 9.0) and 0.1 ml of deionized water, followed by reaction at 40 °C for 20 minutes. After completion of the reaction, reducing sugar was quantitatively determined by a 3,5-dinitrosalicylic acid (DNS) method. More specifically, 1 ml of a DNS reagent was added to 0.5 ml of the reaction solution and heated at 100 °C for 5 minutes for color development, followed by cooling and dilution with 4.5 ml of deionized water. This was subjected to colorimetry at an absorbance of 535 nm. The enzyme titer determined as one unit is an amount of the enzyme which is capable of producing reducing sugar corresponding to 1 μ mol of glucose in one minute under the above-described conditions.

(ii) Decomposition activity of PNPC

A suitable amount of CMCase was acted at 30 °C on 1.0 ml of a reaction solution containing 100 μ mol of a phosphate buffer solution (pH 7.0) and 0.1 μ mol of PNPC (Sigma Co., Ltd.), after which 0.3 ml of 1 M Na₂CO₃ and 1.7 ml of deionized water were added in this order, followed by subjecting the resultant free p-nitrophenol to colorimetry at 400 nm. The amount of the enzyme capable of liberating 1 μ mol of free p-nitrophenol in one minute under the above conditions was determined as one unit.

(iii) Avicelase and FPDase activities

2 ml of a reaction solution for measurement of the CMCase activity was provided, in which there was used, instead of the CMC substrate, 20 mg of Avicel (Merk Inc.) or a bulk piece of a filter paper having a width of 0.5 cm and a length of 5 cm (filter paper for determination of cellulase activity, Toyo No. 51-Specific), thereby determining Avicelase and FPDase activities. An amount of the enzyme capable of liberating 1 μ mol of reducing sugar, calculated as glucose, in one minute under the above conditions was determined as one unit.

(iv) The quantitative determination of proteins was effected by the use of Bio Lad Protein Assay Kit (Bio Lad Co., Ltd.) and bovine serum albumin was used for calculation as a standard protein.

(8) Influence of chelating agents

The resistance of an enzyme for detergents to chelating agents in a builder used as a reaction composition is the most important factor. Alkaline cellulase K was pretreated with EDTA (0.5 mM), EGTA (0.5 mM), NTA (0.5 mM), sodium tripolyphosphate (STPP, 50 mg/ml) and zeolite (5 mg/ml) to determine a residual activity, but any influence was not recognized.

(9) Influence of proteinases

Proteinases function to improve the detergency of detergent compositions. Accordingly, it is a matter of course to further improve the detergency by adding cellulases to proteinase-containing detergents. For this purpose, it is necessary to satisfy the requirement that cellulases for detergents are not hydrolyzed by proteinase and can stably maintain the activity. Alkaline cellulase K has a good resistance to actually employed proteinases for detergents (e.g. API-21, Maxatase, Alkalase and the like) and ordinary proteinases (e.g. pronase) (Table 3).

Table 3

Added Proteinase	Concentration (wt%)	Relative Residual Activity (%)*
API-21	0.002	98
(Showa Denko)	0.02	100
	0.2	114
Maxatase	0.002	113
(Gist)	0.02	99
	0.2	99
Alkalase	0.002	110
(Novo)	0.02	108
	0.2	99
Pronase	0.002	99
(Sigma)	0.02	98
	0.2	102

*Treated with the respective proteinases at 15°C for 12 hours. The activity of an enzyme preparation not treated was taken as 100% and the activity of the respective treated preparations was indicated as an index to the non-treated preparation.

(10) Influence of metals

Divalent metal ions (Hg^{2+} , Cu^{2+} and the like) at a suitable concentration give an effect of inhibition. A slight degree of inhibition is experienced by means of monoiodoacetic acid and p-chloromercuribenzoate.

(11) Influence of surface active agents

Alkaline cellulase K suffers little inhibition of activity by means of various surface active agents such as, for example, LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride.

(12) Molecular weight (gel chromatography using Sephadex G100)

Having a maximum peak at $180,000 \pm 10,000$ (Fig. 7).

The alkaline cellulase K of the invention as compared with known cellulases has the following properties.

The present alkaline cellulase has an optimum pH in a high pH range and is thus clearly distinguishable from cellulases, which have an optimum pH in an acidic range, of molds or germs belonging to the genera *Trichoderma*, *Penicillium*, *Aspergillus* ("Cellulases" by Kazutoshi Nishizawa, Tokyo Nankodo, 1974), *Acromonium* (Japanese Patent Publication No. 59-166081), *Humicola* (Japanese Patent Publication No. 61-16316) and the like.

When compared with the alkaline cellulase disclosed in Japanese Patent Publication No. 50-28515, the present alkaline cellulase has a molecular weight having a maximum peak at $180,000 \pm 10,000$. On the other hand, the known cellulase has a molecular weight of 15,000 to 30,000. The cellulase of *Bacillus* No. 1139 had a molecular weight of 92,000. Moreover, other physical and chemical properties are different from those of these known cellulases. Thus, the present cellulase is clearly distinguishable from these cellulases.

For obtaining CMCase I and CMCase II from the alkaline cellulase K, the alkaline cellulase K is purified by means of a preparative high performance liquid chromatography, a hydroxy apatite chromatography and a DEAE-Toyopearl chromatography and the like, thereby isolating the respective constituents. CMCase I and CMCase II have different molecular weights and electrically charging properties slightly. For instance, they can be separated from each other by elution using a linear concentration gradient of NaCl in the DEAE-Toyopearl chromatography.

The thus obtained CMCase I and CMCase II have the following properties.

CMCase I:

(1) Activity

The present enzyme has a C_x enzymatic activity of acting on CMC. The enzyme has also acts on phosphoric acid-swollen cellulose and has such activity specificities as exhibiting the activity of an enzyme (Avicelase) acting on crystalline cellulose (cellulosic cotton) and Avicel which is a high crystallinity, and activity of a C_1 enzyme such as a filter paper degrading activity (FPDase) and a beta-glucosidase activity of acting on cellobiose and cellobiosaccharides. Furthermore, it slightly acts on PNPC which is an artificial substrate to liberate p-nitrophenol.

(2) Substrate specificity

The main activity of the present enzyme is the CMCase activity and has about 0.3%, based on the main activity, of Avicelase and FPDase activities (C_1 activity). The artificial PNPC degrading activity is about 1.5 to 1.8% (Table 4). On the other hand, it has not any degrading activity on xylan, amylose, dextrin, pectin, inulin and curdlan.

Table 4

Substrate to be Reacted	Enzymatic Activity (specific activity, units/mg of protein) of CMCase I
CMC	6.56
Filter Paper	0.020
Avicel	0.019
Cellobiose	0.010
PNPC	0.099

(3) Working pH and optimum working pH

The working pH of the present enzyme is from 3 to 12.5 and the optimum working pH is from 6 to 11.5 (Fig. 8). The pH at which the working is the highest is about 9.5.

The working pH range with respect to PNPC is from 4 to 11 and the optimum pH is about 7.

(4) pH stability

Stable pH values were determined by allowing the present enzyme to stand at different pH values at 30 °C for 1 hour and measuring a residual activity. As a result, the enzyme was very stable at a pH of from 5 to 12 and was not inactivated (Fig. 9).

(5) Measuring methods of enzymatic activities and protein

The CMCase activity, PNPC degrading activity, Avicelase and FPDase activity, and the content of proteins were measured in the same method as with the alkaline cellulase K.

(6) Working temperature range and working optimum temperature

The working temperature range of CMCase I is in the range of from 10 to 60 °C. The preferable temperature range is from 22 to 53 °C. The optimum working temperature of the present enzyme is 40 °C. The present enzyme has a good resistance at low temperature and has about 50% of the activity at a low temperature of 20 °C (Fig. 10).

(7) Thermal stability

When thermally treated at 50 °C for 30 minutes, the enzyme has residual activity of about 50% (in glycine buffer solution: pH 9.0).

(8) Influences of metals

Metals and ions give influences on the physicochemical properties, particularly a viscosity, of CMC. It will be apparent that when the activity of the enzymatic components of the present invention is measured using a CMC substrate, the factors of the reaction kinetics of the alkaline cellulase K are not correctly reflected.

Accordingly, on the basis of the indication that the CMCase I has the decomposition activity on PNPC, influences of metals on the enzymatic activity were determined. As a result, it was found that Zn^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} and Hg^{2+} inhibited the activity of the CMCase I. On the other hand; Mn^{2+} and Ba^{2+} promoted to activate slightly.

(9) Influence of chelating agents

When CMC was used as a substrate, CMCase I suffered no inhibition with EDTA, EGTA, NTA, STPP and zeolite.

(10) Influence of sugars

Similar to (8), PNPC was used as a substrate for the CMCase I to determine the influences of various sugars. Cellobiose inhibited both enzymatic components and thus exhibited the mode of product inhibition, but other disaccharides, e.g. lactose and maltose, gave no influence. The activity was not inhibited with monosaccharides including glucosamine, N-acetylglucosamine, ribose, arabinose, sorbose, xylose, fructose, galactose, glucose, and their derivatives such as 3-O-methyl-beta-D-glucose, alpha-methyl-beta-D-glucose, alpha-methyl-D-glucoside, alpha-methyl-beta-mannoside and 2-deoxyglucose, and other saccharides such as rhamnose.

(11) Influence of salt concentration

A phosphate buffer solution, a bicine-Na buffer solution and a tris-HCl were used and salt (0 to 250 mM) was used as an ion-strength adjusting agent. The decomposition activity of PNPC was used as an index to determine an effect of the ion strength on the CMCase I. As a result, it was found that the ion strength had no relation to the enzymatic activity with respect to the promotion or inhibition.

(12) Influence of surface active agents

The activity was little inhibited by means of LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), dimethyldialkylammonium chloride, and taurocholic acid.

(13) Molecular weight

The molecular weight determined by gel chromatography (Toyoparl 55S, available from Toyo Soda Co., Ltd.) was $145,000 \pm 10,000$.

(14) UV absorption spectrum

The present enzyme was dissolved in a bicine-sodium buffer solution and subjected to measurement of a UV absorption spectrum, revealing that it had a maximum absorption at about 280 nm with the existence of a shoulder absorption at 290 nm being found by checking a differential absorption spectrum (Fig. 12).

(15) Detection of sugars

The purified enzyme protein was subjected to a color development test using a phenol/sulfuric acid method, with the result that a maximum absorption appeared at 480 nm. This result means that the present enzyme contains a sugar. The sugar was detected by gas chromatography using an Alditol/acetic acid method, indicating that (N-acetyl)glucosamine was contained as a sugar. The present enzyme had a content of the sugar of from 1.3 to 4.0 wt%.

(16) Resistance to proteinases

Proteinases for detergents, e.g. API (showa Denko), Maxatase (Gist Co., Ltd.) and Alkalase (Nobo Co., Ltd.) (0.0002 to 0.1 wt%) were used in coexistence with the present enzyme and the residual activity of the preparations pre-incubated at 15°C for 12 hours was measured. Inactivation was not recognized. Thus, the present enzyme was found to have a strong resistance to the proteinases (Table 5).

Table 5

Added Proteinase	Concentration (wt%)	Relative Residual Activity (%) of CMCase I
Reference (no addition)	-	100
API-21	0.1	99
(Showa Denko)	0.01	112
Maxatase	0.1	102
(Gist)	0.01	108
Alkalase	0.1	111
(Novo)	0.01	116

CMCase II:

(1) Activity

The present enzyme has a C_x enzymatic activity of acting on CMC. The enzyme has also acts on phosphoric acid-swollen cellulose and has such activity specificities as exhibiting the activity of an enzyme (Avicelase) acting on crystalline cellulose (cellulosic cotton) and Avicel which is a high crystallinity, and activity of a C₁ enzyme such as FPDase and a betaglucosidase activity on cellobiose and cellooligosaccharides. Furthermore, it slightly acts on PNPC which is an artificial substrate to liberate p-nitrophenol.

(2) Substrate specificity

The main activity of the present enzyme is the CMCase activity and has about 0.3%, based on the main activity, of Avicelase and FPDase activities (C₁ activity). The artificial substrate PNPC degrading activity is about 1.5 to 2.0% (Table 6). On the other hand, it has not any degrading activity on xylan, amylose, dextrin, pectin, inulin and curdlan.

Table 6

Substrate to be Reracted	Enzyme Activity (specific activity, units/mg of protein) of CMCase II
CMC	7.06
Filter Paper	0.022
Avicel	0.020
Cellobiose	0.010
PNPC	0.140

(3) Working pH and optimum working pH

The working pH of the present enzyme is from 3 to 12 and the optimum working pH is from 8 to 11.5 (Fig. 13). The pH at which the working is the highest is about 9.5.

The working pH range of the present enzyme with respect to PNPC is from 4 to 11 and the optimum pH is about 7.

(4) pH stability

Stable pH values were determined by allowing the present enzyme to stand at different pH values at 30 °C for 1 hour and measuring a residual activity. As a result, the enzyme was very stable at a pH of from 5 to 12 and was not deactivated (Fig. 14).

(5) Measuring methods of enzymatic activities and proteins

The CMCase activity, PNPC degrading activity, Avicelase and FPDase activity, and the content of proteins were measured in the same manner as with the alkaline cellulase K.

(6) Working temperature range and working optimum temperature

The working temperature range of CMCase II is in the range of from 5 to 58 °C. The preferable temperature range is from 14 to 45 °C. The optimum working temperature of the present enzyme is 30 to 40 °C. At 15 °C, the present enzyme has an activity of about 50% of the case of 30 to 40 °C where the maximum activity is shown (Fig. 15).

(7) Thermal stability

When thermally treated at 40 °C for 30 minutes, the enzyme has a residual activity of about 75% (in glycine buffer solution: pH 9.0) (Fig. 16).

(8) Influences of metals

Metals and ions give influences on the physicochemical properties, particularly a viscosity, of CMC. It will be apparent that when the activity of the enzymatic components of the present invention is measured using a CMC substrate, the factors of the reaction kinetics of the alkaline cellulase K are not correctly reflected. Accordingly, on the basis of the indication that the CMCase II has the decomposition activity on PNPC, influences of metals on the enzymatic activity were determined. As a result, it was found that Zn^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} , Hg^{2+} and Ca^{2+} inhibited the activity. On the other hand, Mg^{2+} , Mn^{2+} , Ca^{2+} , Ba^{2+} , Fe^{2+} , Li^{+} , K^{+} , and Na^{+} promoted to active the present enzyme to 1.1 to 2.4 times the original activity.

(9) Influence of chelating agents

When CMC was used as a substrate, the present enzyme suffered no inhibition with EDTA, EGTA, NTA, STPP and zeolite.

(10) Influence of sugars

Similar to (8), PNPC was used as a substrate for the CMCase II to determine the influences of various sugars. Cellobiose inhibited both enzymatic components and thus exhibited the mode of product inhibition, but other disaccharides, e.g. lactose and maltose, gave no influence. The activity was not inhibited by monosaccharides including glucosamine, N-acetylglucosamine, ribose, arabinose, sorbose, xylose, fructose, galactose, glucose, and their derivatives such as 3-O-methyl-beta-D-glucose, alpha-methyl-beta-D-glucose, alpha-methyl-D-glucoside, alpha-methyl-beta-mannoside and 2-deoxyglucose, and other saccharides such as rhamnose.

(11) Influence of salt concentration

A phosphate buffer solution, a bicine-Na buffer solution and a tris-HCl were used and NaCl (0 to 250 mM) was used as an ion-strength adjusting agent. The decomposition activity of PNPC was used as an index to determine an effect of the ion strength on the CMCase II. As a result, it was found that the ion strength had no relation to the enzymatic activity with respect to the promotion or inhibition.

(12) Influence of surface active agents

The activity was little inhibited by means of LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), dimethyldialkylammonium chloride, and taurocholic acid.

(13) Molecular weight

The molecular weight determined by gel chromatography (Toyopearl 55S, available from Toyo Soda Co., Ltd.) was $170,000 \pm 20,000$.

(14) UV absorption spectrum

The present enzyme was dissolved in a bicine-sodium buffer solution and subjected to measurement of a UV absorption spectrum, revealing that it had a maximum absorption at about 280 nm with the existence of a shoulder absorption at 290 nm being found by checking a differential absorption spectrum (Fig. 17).

(15) Detection of sugars

The purified enzyme protein was subjected to a color development test using a phenol/sulfuric acid method, with the result that a maximum absorption appeared at 480 nm. This result revealed that the present enzyme contains a sugar. The sugar was detected by gas chromatography using an Aldehytol/acetic acid method, indicating that (N-acetyl)glucosamine was contained as a sugar. The present enzyme had a content of the sugar of from 1.3 to 4.0 wt%.

(16) Resistance to proteinases

Proteinases for detergents, e.g. API (Showa Denko), Maxatase (Gist Co., Ltd.) and Alkalase (Novo Co., Ltd.) (0.0002 to 0.1 wt%) were used in coexistence with the present enzyme and the residual activity of the preparations pre-incubated at 15°C for 12 hours was measured. Inactivation was not recognized. Thus, the present enzyme was found to have a strong resistance to the proteinases (Table 7).

Table 7

Added Proteinase	Concentration (wt%)	Relative Residual Activity (%) of CMCase II
Reference (no addition)	-	100
API-21 (Showa Denko)	0.1 0.01	102 115
Maxatase (Gist)	0.1 0.01	100 102
Alkalase (Novo)	0.1 0.01	104 108

The comparison in properties between CMCases I and II and those of a known cellulase is as follows.

The present enzymes have an optimum pH in a high pH range and are thus clearly distinguishable from cellulases, which have an optimum pH in an acidic range, of molds or germs belonging to the genera *Trichoderma*, *Penicillium*, *Aspergillus* ("Cellulases" by Kazutoshi Nishizawa, Tokyo Nankodo, 1974), *Acromonium* (Japanese Patent Publication No. 59-166081), *Humicola* (Japanese Patent Publication No. 61-16316) and the like.

When compared with the cellulase disclosed in Japanese Patent Publication No. 50-28515 and the cellulase reported in J. Gen. Microbiol., Vol. 131, page 3339 (1985), it will be found that the CMCase I has a molecular weight of $145,000 \pm 10,000$ and the CMCase II has a molecular weight of $170,000 \pm 20,000$, whereas the alkali cellulase of Japanese Patent Publication No. 50-28515 has a molecular weight of 15,000 to 30,000 and the molecular weight of *Bacillus* No. 1139 is 92,000. Moreover, the enzymes of the present invention apparently differ from these known cellulases in various reaction kinetic properties and the existence of the sugar therein.

The alkaline cellulase K and CMCases I and II obtained according to the invention are specific enzymes which stably work in a wide range including an alkaline side.

For instance, the alkaline cellulase K has, at a pH of 11, a relative activity of about 75 to 80% based on the activity at the optimum pH. Although the optimum pH is at a strongly alkaline side, the activity is shown at a strongly acidic side of about a pH 4 or lower.

Likewise, the CMCase I has a relative activity of about 75 to 80% at a pH of 11 based on the activity at an optimum pH and exhibits an activity even at a pH of about 3.5. The CMCase II has a relative activity of about 70% at a pH of 11, based on the activity at an optimum pH and exhibits an activity at a pH of about 3.

As will be apparent from the above, the alkaline cellulase K and the CMCases I and II of the present invention are enzyme groups which shown good activity at the most strongly alkaline side among hitherto known alkaline cellulases.

Moreover, these enzymes has the features that their activity is shown even at low temperatures and that they have a strong resistance to surface active agents, chelating agents and proteinases. Accordingly, the alkaline cellulase K and the CMCases I and II of the invention can be effectively utilized not only as an additive for clothing detergents, but also as a biomass and in other fields.

The present invention is described in more detail by way of references and examples.

Example 1

A soil collected at Ichikai-machi, Haga-gun, Tochigi-ken, Japan as suspended in a sterilized saline solution and thermally treated at 80°C for 30 minutes. The thermally treated solution was suitably diluted and spread plated on a plate (composed of: 1% of a meat extract (Oxoid Co., Ltd.), 1% of Bacto peptone (Difco Co., Ltd.), 1% of NaCl, 0.1% of KH₂PO₄, 0.5% of Na₂CO₃ (separately sterilized) and 1.5% of Bacto agar-agar), followed by cultivation at 30°C for 3 days, thereby forming colonies. Transplantation on a 2% CMC-containing the above plate according to a replica method was effected to again form colonies, followed by pouring a Congo Red dye solution to obtain colonies whose periphery was transparent. The thus obtained colonies were collected from the plate and high titer CMC-ase-producing bacilli were screened.

By the above procedure, Bacillus sp KSM-635 (FERM BP-1485) of the invention was isolated.

Example 2

The strain of Bacillus sp KSM-635 (FERM BP-1485) was aerobically cultivated in a liquid medium consisting of 1.5% of meat extract, 0.5% of yeast extract, 1% of CMC, 0.1% of KH₂PO₄ and 0.75% of Na₂CO₃ at 34°C for 2 days. 3 liters of cooled ethanol (-10°C) were gradually added to 1 liter of a supernatant liquid of the culture product to cause proteins to precipitate. The precipitate was dissolved in a minimum amount of sterilized deionized water and neutralized with diluted acetic acid, followed by dialysis with running water for 15 hours and freeze-drying to obtain 8.2 g of alkaline cellulase K. The thus obtained cellulase K had the following enzymatic activities shown in Table 8.

Table 8

Kind of Enzyme	Specific Activity (units/g of enzyme powder)
β -Glucosidase	0.6
PNPCase*	7
CMCase	325
FPDase	1.2
Avicelase	1.1

* PNPC degrading activity

Example 3

A Hundred ml of a medium (pH 8.4 to 8.6) containing 1% of CMC, 2% of polypeptone, 0.1% of KH₂PO₄, 0.1% of yeast extract and 0.75% of Na₂CO₃ was placed in a 500 ml Erlenmeyer flask and sterilized by a usual manner, followed by inoculation of the strain of Bacillus sp KSM-635 (FERM BP-1485) and shaking culture at 30°C for 4 days. After completion of the culture, the bacillus cells were centrifugally removed and the resultant supernatant liquid was subjected to measurement of the CMCase activity. The activity was found to be 3,100 units/liter. This supernatant liquid was treated in the same manner as in Example 2 to obtain 9.2 g of alkaline cellulase K.

Example 4

The strain of *Bacillus* sp KSM-635 (FERM BP-1485) was inoculated in a medium of Example 2 in which 1.5 % of meat extract was added instead of Bacto peptone, followed by shaking culture at 30 °C for 3 days. After the culture, a centrifugally separated supernatant liquid was subjected to measurement of the CMCase activity. As a result, the activity was found to be 3,200 units/liter.

Example 5

One liter of the supernatant liquid obtained in Example 4 was purified according to the following procedure to obtain CMCase I and II. The purification was conducted by (1) treatment with streptomycin, (2) fractionation with ammonium sulfate (30 to 75% saturated precipitation fraction), (3) a preparative high performance liquid chromatography (e.g. SW 3000 G column (Toyo Soda Co., Ltd.)), (4) DEAE-Toyopearl (Toyo Soda Co., Ltd.) chromatography, (5) hydroxy apatite (Seikagaku Ind. Co., Ltd.) chromatography, and (6) DEAE-Toyopearl chromatography. At the sixth stage of the purification, when elution by a NaCl linear concentration gradient (from 0.25 M NaCl to 0.35 M NaCl) was effected, CMCase I and CMCase II were eluted according to the order of the elution speed to obtain 24 mg of CMCase I and 15 mg of CMCase II. The thus obtained CMCase I and II were subjected to electrophoresis according to the Davis method (Davis D.J., Ann. N.Y. Acad. Sci., Vol. 121, page 404 (1964)), followed by dyeing with Coomassie Brilliant Blue. As a result, it was found that a single band was obtained for the respective enzymes.

Example 6

The CMCase I and II obtained in Example 5 were subjected to electrophoresis using sodium dodecylsulfate according to a conventional procedure. The results are shown in Fig. 18. From the results, it was found that the CMCase I and II were association products of different molecule species having a lowest molecular weight of $30,000 \pm 2,000$ and a highest molecular weight of $120,000 \pm 15,000$. The CMCase I and II are assumed to be associated products of enzyme proteins which are strongly associated by some physical and chemical interactions and have a lowest molecular weight of $30,000 \pm 2,000$. However, how the protein is associated is not known at present. It is considered that the main protein species of the CMCase I has a molecular weight of from 55,000 to 65,000 and the CMCase II has a molecular weight of from 95,000 to 120,000. Accordingly, the products obtained in this example are merely an example of the results of the sodiumdodecylsulfate electrophoresis and should not be construed as limiting exact subunit structures of the CMCase I and II.

Claims

Claims for the following Contracting States : DE, FR, GB, IT, NL

1. Alkaline cellulase K, which is obtainable from a culture product of *Bacillus* sp KSM-635 deposited as FERM BP-1485, having the following physical and chemical properties:

- (1) Activity

Having a Cx enzymatic activity of acting on carboxymethyl cellulose along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity;

- (2) Specificity on Substrates

Acting on carboxymethyl cellulose(CMC), crystalline cellulose, Avicel, cellobiose, and p-nitrophenyl cellobioside(PNPC);

- (3) Working pH and optimum pH

Having a working pH in the range of 4 to 12 and an optimum pH in the range of 9 to 10;

- (4) Stable pH

Having stable pH values of 4.5 to 10.5 and 6.8 to 10 when allowed to stand at 40 °C for 10 minutes and 30 minutes, respectively;

- (5) Working temperature range and working optimum temperature

Working in a wide temperature range of from 10 to 65 °C with an optimum temperature being recognized at about 40 °C;

- (6) Influences of chelating agents

The activity not impeded with ethylenediamine tetraacetic acid (EDTA), ethyleneglycol-bis-(β -aminoethylether) N,N,N',N''-tetraacetic acid (EGTA), N,N-bis(carboxymethyl)glycine (nitrilotriacetic acid) (NTA), sodium tripolyphosphate (STPP) and zeolite.

Example 4

The strain of *Bacillus* sp KSM-635 (FERM BP-1485) was inoculated in a medium of Example 2 in which 1.5 % of meat extract was added instead of Bacto peptone, followed by shaking culture at 30 °C for 3 days.

After the culture, a centrifugally separated supernatant liquid was subjected to measurement of the CMCase activity. As a result, the activity was found to be 3,200 units/liter.

Example 5

One liter of the supernatant liquid obtained in Example 4 was purified according to the following procedure to obtain CMCase I and II. The purification was conducted by (1) treatment with streptomycin, (2) fractionation with ammonium sulfate (30 to 75% saturated precipitation fraction), (3) a preparative high performance liquid chromatography (e.g. SW 3000 G column (Toyo Soda Co., Ltd.)), (4) DEAE-Toyopearl (Toyo Soda Co., Ltd.) chromatography, (5) hydroxy apatite (Seikagaku Ind. Co., Ltd.) chromatography, and (6) DEAE-Toyopearl chromatography. At the sixth stage of the purification, when elution by a NaCl linear concentration gradient (from 0.25 M NaCl to 0.35 M NaCl) was effected, CMCase I and CMCase II were eluted according to the order of the elution speed to obtain 24 mg of CMCase I and 15 mg of CMCase II. The thus obtained CMCase I and II were subjected to electrophoresis according to the Davis method (Davis D.J., *Ann. N.Y. Acad. Sci.*, Vol. 121, page 404 (1964)), followed by dyeing with Coomassie Brilliant Blue. As a result, it was found that a single band was obtained for the respective enzymes.

Example 6

The CMCase I and II obtained in Example 5 were subjected to electrophoresis using sodium dodecylsulfate according to a conventional procedure. The results are shown in Fig. 18. From the results, it was found that the CMCase I and II were association products of different molecule species having a lowest molecular weight of $30,000 \pm 2,000$ and a highest molecular weight of $120,000 \pm 15,000$. The CMCase I and II are assumed to be associated products of enzyme proteins which are strongly associated by some physical and chemical interactions and have a lowest molecular weight of $30,000 \pm 2,000$. However, how the protein is associated is not known at present. It is considered that the main protein species of the CMCase I has a molecular weight of from 55,000 to 65,000 and the CMCase II has a molecular weight of from 95,000 to 120,000. Accordingly, the products obtained in this example are merely an example of the results of the sodium dodecylsulfate electrophoresis and should not be construed as limiting exact subunit structures of the CMCase I and II.

Claims

Claims for the following Contracting States : DE, FR, GB, IT, NL

1. Alkaline cellulase K, which is obtainable from a culture product of *Bacillus* sp KSM-635 deposited as FERM BP-1485, having the following physical and chemical properties:

- (1) Activity

Having a Cx enzymatic activity of acting on carboxymethyl cellulose along with a weak Ci enzymatic activity and a weak beta-glucosidase activity;

- (2) Specificity on Substrates

Acting on carboxymethyl cellulose(CMC), crystalline cellulose, Avicel, cellobiose, and p-nitrophenyl cellobioside(PNPC);

- (3) Working pH and optimum pH

Having a working pH in the range of 4 to 12 and an optimum pH in the range of 9 to 10;

- (4) Stable pH

Having stable pH values of 4.5 to 10.5 and 6.8 to 10 when allowed to stand at 40 °C for 10 minutes and 30 minutes, respectively;

- (5) Working temperature range and working optimum temperature

Working in a wide temperature range of from 10 to 65 °C with an optimum temperature being recognized at about 40 °C;

- (6) Influences of chelating agents

The activity not impeded with ethylenediamine tetraacetic acid (EDTA), ethyleneglycol-bis-(β -aminoethylether) N,N,N',N''-tetraacetic acid (EGTA), N,N-bis(carboxymethyl)glycine (nitrilotriacetic acid) (NTA), sodium tripolyphosphate (STPP) and zeolite.



(7) Influences of surface active agents

Undergoing little inhibition of activity by means of surface active agents such as sodium linear alkylbenzenesulfonates (LAS), sodium alkylsulfates (AS), sodium polyoxyethylene alkylsulfates (ES), sodium alpha-olefinsulfonates (AOS), sodium alpha-sulfonated aliphatic acid esters (alpha-SFE), sodium alkylsulfonates (SAS), polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride.

(8) Influences of proteinases

Having a strong resistance to proteinases.

(9) Molecular weight (determined by gel chromatography, using sephadex®G 100)

Having a maximum peak at $180,000 \pm 10,000$.

2. A method for producing alkaline cellulase K having the features defined in claim 1, which comprises cultivating alkaline cellulase K producing *Bacillus* sp KSM-635, deposited as FERM BP-1485, and collecting alkaline cellulase K from the resulting culture product.

3. CMCase I, which is obtainable from a culture product of *Bacillus* sp KSM-635, deposited as FERM BP-1485, having the following physical and chemical properties:

(1) Activity

Having the Cx enzymatic activity of acting on CMC along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity.

(2) Specificity on substrates

Acting on CMC, crystalline cellulose, Avicel, cellubiose, and PNPC.

(3) Working pH and optimum pH

Having a working pH in the range of 3 to 12.5 and an optimum pH in the range of 6 to 11.5.

(4) Stable pH

Not inactivated at a pH of 5 to 12 when allowed to stand at 30°C for 1 hour.

(5) Working temperature and working optimum temperature

Working in a temperature range of from 10 to 60°C with an optimum temperature being at 22 to 53°C.

(6) Influences of chelating agents

The activity not impeded with EDTA, EGTA, NTA, STPP and zeolite.

(7) Influences of surface active agents

Undergoing little inhibition of activity by means of LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), dimethyldialkylammonium chloride, and taurocholic acid.

(8) Resistance to proteinases

Having a strong resistance to proteinases.

(9) Molecular weight (determined by gel chromatography using Toyopearl®55S)

Having a molecular weight of this enzyme of $145,000 \pm 10,000$.

(10) UV absorption spectrum

This enzyme has a maximum absorption at 280 nm and a shoulder absorption at 290 nm, thus sugar components being contained.

4. CMCase II which is obtainable from a culture product of *Bacillus* sp KSM-635, deposited as FERM BP-1485, having the following physical and chemical properties:

(1) Activity

Having the Cx enzymatic activity of acting on CMC along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity.

(2) Specificity on substrates

Acting on CMC, crystalline cellulose, Avicel, cellubiose and PNPC.

(3) Working pH and optimum pH

Having working pH in the range of 3 to 12.5 and an optimum pH in the range of 6 to 11.5.

(4) pH stability

Not inactivated at a pH of 5 to 12 when allowed to stand at 30°C for 1 hour.

(5) Working temperature and working optimum temperature

Working in a temperature range of from 5 to 58°C with an optimum temperature being at 14 to 45°C.



(6) Influences of chelating agents

The activity not impeded with EDTA, EGTA, NTA, STPP and zeolite.

(7) Influences of surface active agents

Undergoing little inhibition of activity by means of LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride.

(8) Resistance to proteinases

Having a strong resistance to proteinases.

(9) Molecular weight (determined by Gel chromatography using Toyopearl®55S)

The molecular weight of CMCase II is $170,000 \pm 20,000$.

(10) UV absorption spectrum

Having a maximum absorption at 280 nm and a shoulder absorption at 290 nm.

5. Bacteria *Bacillus* sp KSM-635, deposited as FERM BP-1485, capable of producing alkaline cellulase K and having the following enzymatic properties:

(1) Activity

Having a C_x enzymatic activity of acting on CMC along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity.

(2) Specificity on Substrates

Acting on CMC, crystalline cellulose, Avicel, cellubiose, and PNPC.

(3) Working pH and optimum pH

Having a working pH in the range of 4 to 12 and an optimum pH in the range of 9 to 10 (Fig. 3).

(4) pH stability

Having stable pH values of 4.5 to 10.5 and 6.8 to 10 when allowed to stand at 40°C for 10 minutes and 30 minutes, respectively (Fig. 4).

(5) Working temperature range and working optimum temperature

Working in a wide temperature range of from 10 to 65°C with an optimum temperature being recognized at about 40°C (Fig. 5).

(6) Influences of chelating agents

The activity not impeded with EDTA, EGTA, NTA, STPP and zeolite.

(7) Influences of surface active agents

Undergoing little inhibition of activity by means of surface active agents such as LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride.

(8) Influences of proteinases

Having a strong resistance to proteinases.

(9) Molecular weight (determined by gel chromatography using sephadex®G100)

Having a maximum peak at $180,000 \pm 10,000$

40 Claims for the following Contracting State : ES

1. A method for producing alkaline cellulase K, CMCase I and CMCase II obtainable from the bacteria *Bacillus* sp KSM-635 deposited as FERM BP-1485, characterized in that it comprises cultivation of said bacteria aerobically at 22 to 40°C, at pH 8 to 11 by using a culture medium which contains a suitable combination of nitrogen sources and carbon sources, followed by isolation of the desired products by means of known techniques of collection and purification, whereby alkaline cellulase K is isolated by means of separation of the bacillus cells such as centrifugation, filtration and so on, followed by ordinary isolation methods such as salting-out, isoelectric precipitation, solvent precipitation to separate the products as a precipitate; and whereby CMCase I and CMCase II are obtained from alkaline cellulase K, by means of a preparative high performance liquid chromatography, a hydroxy apatite chromatography or a DEAE-Toyopearl® chromatography, thereby isolating the respective constituent.

2. A method according to claim 1, characterised in that the alkaline cellulase K has the following physical and chemical properties:

(1) Activity

Having a C_x enzymatic activity of acting on carboxymethyl cellulose along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity;



(2) Specificity on Substrates

Acting on carboxymethyl cellulose(CMC), crystalline cellulose, Avicel , cellobiose, and p-nitrophenyl cellobioside(PNPC);

(3) Working pH and optimum pH

Having a working pH in the range of 4 to 12 and an optimum pH in the range of 9 to 10;

(4) Stable pH

Having stable pH values of 4.5 to 10.5 and 6.8 to 10 when allowed to stand at 40 °C for 10 minutes and 30 minutes, respectively;

(5) Working temperature range and working optimum temperature

Working in a wide temperature range of from 10 to 65 °C with an optimum temperature being recognized at about 40 °C;

(6) Influences of chelating agents

The activity not impeded with ethylenediamine tetraacetic acid (EDTA), ethyleneglycol-bis-(β -aminoethylether) N,N,N',N''-tetraacetic acid (EGTA), N,N-bis(carboxymethyl)glycine (nitrilotriacetic acid) (NTA), sodium tripolyphosphate (STPP) and zeolite;

(7) Influences of surface active agents

Undergoing little inhibition of activity by means of surface active agents such as sodium linear alkylbenzenesulfonates (LAS), sodium alkylsulfates (AS), sodium polyoxyethylene alkylsulfates (ES), sodium alpha-olefinsulfonates (AOS), sodium alpha-sulfonated aliphatic acid esters (alpha-SFE), sodium alkylsulfonates (SAS), polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride;

(8) Influences of proteinases

Having a strong resistance to proteinases;

(9) Molecular weight (determined by gel chromatography, using sephadex®G 100)

Having a maximum peak at $180,000 \pm 10,000$.

3. A method according to claim 1, characterised in that the product CMCase I has the following physical and chemical properties:

(1) Activity

Having the C_x enzymatic activity of acting on CMC along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity;

(2) Specificity on substrates

Acting on CMC, crystalline cellulose, Avicel, cellubiose, and PNPC;

(3) Working pH and optimum pH

Having a working pH in the range of 3 to 12.5 and an optimum pH in the range of 6 to 11.5;

(4) Stable pH

Not inactivated at a pH of 5 to 12 when allowed to stand at 30 °C for 1 hour;

(5) Working temperature and working optimum temperature

Working in a temperature range of from 10 to 60 °C with an optimum temperature being at 22 to 53 °C;

(6) Influences of chelating agents

The activity not impeded with EDTA, EGTA, NTA, STPP and zeolite;

(7) Influences of surface active agents

Undergoing little inhibition of activity by means of LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), dimethyldialkylammonium chloride, and taurocholic acid;

(8) Resistance to proteinases

Having a strong resistance to proteinases;

(9) Molecular weight (determines by gel chromatography using Toyopearl®55S)

Having a molecular weight of this enzyme of $145,000 \pm 10,000$;

(10) UV absorption spectrum

This enzyme has a maximum absorption at 280 nm and a shoulder absorption at 290 nm, thus sugar components being contained.

4. A method according to claim 1, characterised in that the product CMCase II has the following physical and chemical properties:

(1) Activity

Having the C_x enzymatic activity of acting on CMC along with a weak C₁ enzymatic activity and



a weak beta-glucosidase activity;

(2) Specificity on substrates

Acting on CMC, crystalline cellulose, Avicel, cellubiose and PNPC;

(3) Working pH and optimum pH

Having working pH in the range of 3 to 12.5 and an optimum pH in the range of 6 to 11.5;

(4) pH stability

Not inactivated at a pH of 5 to 12 when allowed to stand at 30 °C for 1 hour;

(5) Working temperature and working optimum temperature

Working in a temperature range of from 5 to 58 °C with an optimum temperature being at 14 to 45 °C;

(6) Influences of chelating agents

The activity not impeded with EDTA, EGTA, NTA, STPP and zeolite;

(7) Influences of surface active agents

Undergoing little inhibition of activity by means of LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride;

(8) Resistance to proteinases

Having a strong resistance to proteinases;

(9) Molecular weight (determined by gel chromatography using Toyopearl®55S)

The molecular weight of CMCase II is 170,000 ± 20,000;

(10) UV absorption spectrum

Having a maximums absorption at 280 nm and a shoulder absorption at 290 nm.

5. A method according to claim 1, characterised in that the bacteria Bacillus sp KSM-635 deposited as FERM BP-1485 is capable of producing an enzyme having the following enzymatic properties:

(1) Activity

Having a C_x enzymatic activity of acting on CMC along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity;

(2) Specificity on Substrates

Acting on CMC, crystalline cellulose, Avicel, cellubiose, and PNPC;

(3) Working pH and optimum pH

Having a working pH in the range of 4 to 12 and an optimum pH in the range of 9 to 10 (Fig. 3);

(4) pH stability

Having stable pH values of 4.5 to 10.5 and 6.8 to 10 when allowed to stand at 40 °C for 10 minutes and 30 minutes, respectively (Fig. 4);

(5) Working temperature range and working optimum temperature

Working in a wide temperature range of from 10 to 65 °C with an optimum temperature being recognized at about 40 °C (Fig. 5);

(6) Influences of chelating agents

The activity not impeded with EDTA, EGTA, NTA, STPP and zeolite;

(7) Influences of surface active agents

Undergoing little inhibition of activity by means of surface active agents such as LAS, AS, ES, AOS, alpha-SFE, SAS, polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride;

(8) Influences of proteinases

Having a strong resistance to proteinases;

(9) Molecular weight (determined by gel chromatography using sephadex®G100)

Having a maximum peak at 180,000 ± 10,000.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : DE, FR, GB, IT, NL

1. Alkalische Cellulase K, erhältlich aus einem Kulturprodukt von Bacillus sp KSM-635, hinterlegt als FERM BP-1485, mit den nachstehenden physikalischen und chemischen Eigenschaften:

(1) Aktivität

C_x-enzymatische Aktivität auf Carboxymethylcellulose zusammen mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β-Glucosidaseaktivität;



(2) Spezifität auf Substrate

Wirkung auf Carboxymethylcellulose (CMC), kristalline Cellulose, Avicel, Cellobiose und p-Nitrophenylcellobiosid (PNPC);

(3) Arbeits-pH-Wert und optimaler pH-Wert

Mit einem Arbeits-pH-Wert im Bereich von 4 bis 12 und einem optimalen pH-Wert im Bereich von 9 bis 10;

(4) pH-Wert-Stabilität

Stabil bei pH-Werten von 4,5 bis 10,5 und 8,8 bis 10, wenn bei 40 °C für 10 Minuten bzw. 30 Minuten belassen;

(5) Arbeitstemperaturbereich und optimale Arbeitstemperatur

Wirkt in einem breiten Temperaturbereich von 10 bis 65 °C mit einer optimalen Temperatur bei etwa 40 °C;

(6) Einflüsse von chelatbildenden Mitteln

Die Aktivität ist nicht von Ethylendiamintetraessigsäure (EDTA), Ethylenglycol-bis-(β -aminoethylether)-N,N,N',N'-tetraessigsäure (EGTA), N,N-Bis(carboxymethyl)glycin (Nitrilotriessigsäure) (NTA), Natriumtripolyphosphat (STPP) und Zeolith beeinflusst.

(7) Einflüsse von Tensiden

Unterliegt kaum Inhibierung der Aktivität durch Tenside wie lineare Natriumalkylbenzolsulfonate (LAS), Natriumalkylsulfate (AS), Natriumpolyoxyethylenalkylsulfate (ES), Natrium- α -olefinsulfonate (AOS), Natrium- α -sulfonierte aliphatische Säureester (α -SFE), Natriumalkylsulfonate (SAS), Polyoxyethylen-sekundäre Alkylether, Fettsäuresalze (Natriumsalze) und Dimethyldialkylammoniumchlorid.

(8) Einfluß von Proteinase

Mit hoher Beständigkeit gegenüber Proteinase.

(9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Sephadex® G 100)

Mit einem maximalen Peak bei 180000 $\pm$ 10000.

2. Verfahren zur Herstellung alkalischer Cellulase K mit den in Beispiel 1 definierten Merkmalen, umfassend das Züchten von alkalische Cellulase K erzeugendem Bacillus sp KSM-635, hinterlegt als FERM BP-1485, und Gewinnen der alkalischen Cellulase K aus dem erhaltenen Kulturprodukt.

3. CMCase 1, erhältlich aus einem Kulturprodukt von Bacillus sp KSM-635, hinterlegt als FERM BP-1485, mit den nachstehenden physikalischen und chemischen Eigenschaften:

(1) Aktivität

Cx-enzymatische Aktivität auf CMC zusammen mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β -Glucosidaseaktivität.

(2) Spezifität auf Substrate

Wirkt auf CMC, kristalline Cellulose, Avicel, Cellubiose und PNPC.

(3) Arbeits-pH-Wert und optimaler pH-Wert

Mit einem Arbeits-pH-Wert in dem Bereich von 3 bis 12,5 und einem optimalen pH-Wert im Bereich von 6 bis 11,5.

(4) pH-Wert-Stabilität

Nicht inaktiviert bei einem pH-Wert von 5 bis 12, wenn bei 30 °C für 1 Stunde belassen.

(5) Arbeitstemperatur und optimale Arbeitstemperatur

Wirkt in einem Temperaturbereich von 10 bis 60 °C mit einer optimalen Temperatur von 22 bis 53 °C.

(6) Einflüsse von chelatbildenden Mitteln

Die Aktivität wird nicht von EDTA, EGTA, NTA, STPP und Zeolith beeinflusst.

(7) Einflüsse von Tensiden

Unterliegt kaum Inhibierung der Aktivität durch LAS, AS, ES, AOS, α -SFE, SAS, Polyoxyethylen-sekundäre Alkylether, Fettsäuresalze (Natriumsalze), Dimethyldialkylammoniumchlorid und Taurocholsäure.

(8) Beständigkeit gegen Proteinase

Mit hoher Beständigkeit gegen Proteinase

(9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Toyopearl® 55S)

Dieses Enzym weist ein Molekulargewicht von 145000 $\pm$ 10000 auf.

(10) UV-Absorptionsspektrum

Das Enzym weist eine maximale Absorption bei 280 nm oder eine Schulterabsorption bei 290



nm auf, somit sind Zuckerbestandteile enthalten.

4. CMCase II, erhältlich aus einem Kulturprodukt von *Bacillus* sp KSM-635, hinterlegt als FERM BP-1485, mit den nachstehenden physikalischen und chemischen Eigenschaften:

(1) Aktivität

Cx-enzymatische Aktivität auf CMC zusammen mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β -Glucosidaseaktivität.

(2) Spezifität auf Substrate

Wirkt auf CMC, kristalline Cellulose, Avicel, Cellubiose und PNPC.

(3) Arbeits-pH-Wert und optimaler pH-Wert

Mit einem Arbeits-pH-Wert in dem Bereich von 3 bis 12,5 und einem optimalen pH-Wert im Bereich von 6 bis 11,5.

(4) pH-Wert-Stabilität

Nicht aktiviert bei einem pH-Wert von 5 bis 12, wenn bei 30 °C für 1 Stunde belassen.

(5) Arbeitstemperatur und optimale Arbeitstemperatur

Wirkt in einem Temperaturbereich von 5 bis 58 °C mit einer optimalen Temperatur von 14 bis 45 °C.

(6) Einflüsse von chelatbildenden Mitteln

Die Aktivität wird nicht von EDTA, EGTA, NTA, STPP und Zeolith beeinflusst.

(7) Einflüsse von Tensiden

Unterliegt kaum Inhibierung der Aktivität durch LAS, AS, ES, AOS, α -SFE, SAS, Polyoxyethylensekundäre Alkylether, Fettsäuresalze (Natriumsalze) und Dimethyldialkylammoniumchlorid.

(8) Beständigkeit gegen Proteinase

Mit hoher Beständigkeit gegen Proteinase

(9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Toyopearl® 55S)

Das Molekulargewicht von CMCase II ist 170000 $\pm$ 20000.

(10) UV-Absorptionsspektrum

Das Enzym weist eine maximale Absorption bei 280 nm oder eine Schulterabsorption bei 290 nm auf.

5. Bakterien *Bacillus* sp KSM-635, hinterlegt als FERM BP-1485, die in der Lage sind, alkalische Cellulase K zu erzeugen, mit den nachstehenden enzymatischen Eigenschaften:

(1) Aktivität

Cx-enzymatische Aktivität auf CMC zusammen mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β -Glucosidaseaktivität.

(2) Spezifität auf Substrate

Wirkt auf CMC, kristalline Cellulose, Avicel, Cellobiose und PNPC.

(3) Arbeits-pH-Wert und optimaler pH-Wert

Mit einem Arbeits-pH-Wert in dem Bereich von 4 bis 12 und einem optimalen pH-Wert im Bereich von 9 bis 10 (Fig. 3).

(4) pH-Wert-Stabilität

Stabil bei pH-Werten von 4,5 bis 10,5 und 6,8 bis 10, wenn bei 40 °C für 10 Minuten bzw. 30 Minuten belassen (Fig. 4).

(5) Arbeitstemperatur und optimale Arbeitstemperatur

Wirkt in einem weiten Temperaturbereich von 10 bis 65 °C mit einer optimalen Temperatur von etwa 40 °C (Fig. 5).

(6) Einflüsse von chelatbildenden Mitteln

Die Aktivität wird nicht von EDTA, EGTA, NTA, STPP und Zeolith beeinflusst.

(7) Einflüsse von Tensiden

Unterliegt kaum Inhibierung der Aktivität durch Tenside, wie LAS, AS, ES, AOS, α -SFE, SAS, Polyoxyethylensekundären Alkylethern, Fettsäuresalzen (Natriumsalze) und Dimethyldialkylammoniumchlorid.

(8) Beständigkeit gegen Proteinase

Mit hoher Beständigkeit gegen Proteinase

(9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Sephadex® G 100)

Mit einem maximalen Peak bei 180000 $\pm$ 10000 (Fig. 7).



Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zur Herstellung von alkalischer Cellulase K, CMCase I und CMCase II, erhältlich aus den Bakterien *Bacillus* sp KSM-635, hinterlegt als FERM BP-1485, dadurch gekennzeichnet, daß es die Züchtung der Bakterien aerob bei 22 bis 40 °C, bei pH-Wert 8 bis 11 unter Verwendung eines Kulturmediums, bestehend aus einer geeigneten Kombination von Stickstoffquellen und Kohlenstoffquellen, gefolgt von Gewinnung der gewünschten Produkte durch übliche Verfahren zum Isolieren und Reinigen, wodurch alkalische Cellulase K durch Trennen der Bacillenzellen mit Zentrifugieren, Filtrieren usw. isoliert wird, gefolgt von üblichen Isolierungsmethoden wie Aussalzen, isoelektrische Fällung, Lösungsfällung und Trennen der Produkte als Präzipitat und wobei CMCase I und CMCase II aus alkalischer Cellulase K durch präparative Hochdruckflüssigchromatografie, Hydroxyapatitchromatografie oder DEAE-Toyopearl®-Chromatografie erhalten wird, wodurch die betreffenden Bestandteile isoliert werden.

2. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß die alkalische Cellulase K die nachstehenden physikalischen und chemischen Eigenschaften aufweist:

(1) Aktivität

Cx-enzymatische Aktivität auf Carboxymethylcellulose zusammen mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β -Glucosidaseaktivität;

(2) Spezifität auf Substrate

Wirkung auf Carboxymethylcellulose (CMC), kristalline Cellulose, Avicel, Cellobiose und p-Nitrophenylcellobiosid (PNPC);

(3) Arbeits-pH-Wert und optimaler pH-Wert

Mit einem Arbeits-pH-Wert im Bereich von 4 bis 12 und einem optimalen pH-Wert im Bereich von 9 bis 10;

(4) pH-Wert-Stabilität

Stabil bei pH-Werten von 4,5 bis 10,5 und 6,8 bis 10, wenn bei 40 °C für 10 Minuten bzw. 30 Minuten belassen;

(5) Arbeitstemperaturbereich und optimale Arbeitstemperatur

Wirkt in einem breiten Temperaturbereich von 10 bis 65 °C mit einer optimalen Temperatur bei etwa 40 °C;

(6) Einflüsse von chelatbildenden Mitteln

Die Aktivität ist nicht von Ethylendiamintetraessigsäure (EDTA), Ethylenglycol-bis-(β -aminoethylether)-N,N,N',N'-tetraessigsäure (EGTA), N,N-Bis(carboxymethyl)glycin (Nitrilotriessigsäure) (NTA), Natriumtripolyphosphat (STPP) und Zeolith beeinflusst.

(7) Einflüsse von Tensiden

Unterliegt kaum Inhibierung der Aktivität durch Tenside, wie lineare Natriumalkylbenzolsulfonate (LAS), Natriumalkylsulfate (AS), Natriumpolyoxyethylenalkylsulfate (ES), Natrium- α -olefinsulfonate (AOS), Natrium- α -sulfonierte aliphatische Säureester (α -SFE), Natriumalkylsulfonate (SAS), Polyoxyethylen-sekundäre Alkylether, Fettsäuresalze (Natriumsalze) und Dimethyldialkylammoniumchlorid.

(8) Einfluß von Proteinasen

Mit hoher Beständigkeit gegenüber Proteinasen.

(9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Sephadex® G 100)

Mit einem maximalen Peak bei 180000 $\pm$ 10000.

3. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß das Produkt CMCase I die nachstehenden physikalischen und chemischen Eigenschaften aufweist:

(1) Aktivität

Cx-enzymatische Aktivität auf CMC zusammen mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β -Glucosidaseaktivität.

(2) Spezifität auf Substrate

Wirkt auf CMC, kristalline Cellulose, Avicel, Cellubiose und PNPC.

(3) Arbeits-pH-Wert und optimaler pH-Wert

Mit einem Arbeits-pH-Wert in dem Bereich von 3 bis 12,5 und einem optimalen pH-Wert im Bereich von 6 bis 11,5.

(4) pH-Wert-Stabilität

Nicht aktiviert bei einem pH-Wert von 5 bis 12, wenn bei 30 °C für 1 Stunde belassen.



- (5) Arbeitstemperatur und optimale Arbeitstemperatur
Wirkt in einem Temperaturbereich von 10 bis 60 °C mit einer optimalen Temperatur von 22 bis 53 °C.
- (6) Einflüsse von chelatbildenden Mitteln
Die Aktivität wird nicht von EDTA, EGTA, NTA, STPP und Zeolith beeinflusst.
- (7) Einflüsse von Tensiden
Unterliegt kaum Inhibierung der Aktivität durch LAS, AS, ES, AOS, α -SFE, SAS, Polyoxyethylen- sekundäre Alkylether, Fettsäuresalze (Natriumsalze), Dimethyldialkylammoniumchlorid und Tauro- cholinsäure.
- (8) Beständigkeit gegen Proteinase
Mit hoher Beständigkeit gegen Proteinase
- (9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Toyopearl® 55S)
Dieses Enzym weist ein Molekulargewicht 145000 $\pm$ 10000 auf.
- (10) UV-Absorptionsspektrum
Das Enzym weist eine maximale Absorption bei 280 nm oder eine Schulterabsorption bei 290 nm auf, somit sind Zuckerbestandteile enthalten.
4. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß das Produkt CMCCase II die nachstehen- den physikalischen und chemischen Eigenschaften aufweist:
- (1) Aktivität
Cx-enzymatische Aktivität auf CMC zusammen mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β -Glucosidaseaktivität.
- (2) Spezifität auf Substrate
Wirkt auf CMC, kristalline Cellulose, Avicel, Cellobiose und PNPC.
- (3) Arbeits-pH-Wert und optimaler pH-Wert
Mit einem Arbeits-pH-Wert in dem Bereich von 3 bis 12,5 und einem optimalen pH-Wert im Bereich von 6 bis 11,5.
- (4) pH-Wert-Stabilität
Nicht aktiviert bei einem pH-Wert von 5 bis 12, wenn bei 30 °C für 1 Stunde belassen.
- (5) Arbeitstemperatur und optimale Arbeitstemperatur
Wirkt in einem Temperaturbereich von 5 bis 58 °C mit einer optimalen Temperatur von 14 bis 45 °C.
- (6) Einflüsse von chelatbildenden Mitteln
Die Aktivität wird nicht von EDTA, EGTA, NTA, STPP und Zeolith beeinflusst.
- (7) Einflüsse von Tensiden
Unterliegt kaum Inhibierung der Aktivität durch LAS, AS, ES, AOS, α -SFE, SAS, Polyoxyethylen- sekundäre Alkylether, Fettsäuresalze (Natriumsalze) und Dimethyldialkylammoniumchlorid.
- (8) Beständigkeit gegen Proteinase
Mit hoher Beständigkeit gegen Proteinase
- (9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Toyopearl® 55S)
Das Molekulargewicht von CMCCase II ist 170000 $\pm$ 20000.
- (10) UV-Absorptionsspektrum
Das Enzym weist eine maximale Absorption bei 280 nm oder eine Schulterabsorption bei 290 nm auf.
5. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß Bakterien *Bacillus* sp KSM-635, hinterlegt als FERM BP-1485, in der Lage sind, ein Enzym mit den nachstehenden enzymatischen Eigenschaften zu erzeugen:
- (1) Aktivität
Cx-enzymatische Aktivität auf CMC und mit einer schwachen C₁-enzymatischen Aktivität und einer schwachen β -Glucosidaseaktivität.
- (2) Spezifität auf Substrate
Wirkt auf CMC, kristalline Cellulose, Avicel, Cellubiose und PNPC.
- (3) Arbeits-pH-Wert und optimaler pH-Wert
Mit einem Arbeits-pH-Wert in dem Bereich von 4 bis 12 und einem optimalen pH-Wert im Bereich von 9 bis 10 (Fig. 3).
- (4) pH-Wert-Stabilität
Stabil bei pH-Werten von 4,5 bis 10,5 und 6,8 bis 10, wenn bei 40 °C für 10 Minuten bzw. 30



Minuten belassen (Fig.4).

(5) Arbeitstemperatur und optimale Arbeitstemperatur

Wirkt in einem weiten Temperaturbereich von 10 bis 65 °C mit einer optimalen Temperatur von etwa 40 °C (Fig. 5).

(6) Einflüsse von chelatbildenden Mitteln

Die Aktivität wird nicht von EDTA, EGTA, NTA, STPP und Zeolith beeinflusst.

(7) Einflüsse von Tensiden

Unterliegt kaum Inhibierung der Aktivität durch Tenside, wie LAS, AS, ES, AOS, α -SFE, SAS, Polyoxyethylensekundäre Alkylether, Fettsäuresalze (Natriumsalze) und Dimethyldialkylammoniumchlorid.

(8) Beständigkeit gegen Proteinase

Mit hoher Beständigkeit gegen Proteinase

(9) Molekulargewicht (bestimmt durch Gelchromatografie unter Verwendung von Sephadex® G 100)

Mit einem maximalen Peak bei 180000 $\pm$ 10000 (Fig. 7).

Revendications

Revendications pour les Etats contractants suivants : DE, FR, GB, IT, NL

1. Cellulase alcaline K, qui peut être obtenue à partir d'un produit de culture de *Bacillus* sp KSM-635 déposé sous le numéro FERM BP-1485, ayant les propriétés physiques et chimiques suivantes :

(1) Activité

Elle présente une activité enzymatique Cx d'action sur la carboxy méthyl cellulose, en même temps qu'une faible activité enzymatique C₁ et une faible activité bêta-glucosidase ;

(2) Spécificité pour les Substrats

Elle agit sur la carboxy méthyl cellulose (CMC), la cellulose cristalline, l'Avicel, la cellobiose, et le p-nitrophényl cellobioside (PNPC) ;

(3) pH d'action et pH optimal

Elle présente un pH d'action se situant dans la plage de 4 à 12 et un pH optimal se situant dans la plage de 9 à 10 ;

(4) Stabilité au pH

Elle est stable aux valeurs de pH stable de 4,5 à 10,5 et de 6,8 à 10, lorsqu'on la laisse reposer à 40 °C respectivement pendant 10 minutes et pendant 30 minutes ;

(5) Plage de température d'activité et température d'activité optimale

Elle a une activité dans une large plage de température de 10 à 65 °C, une température optimale étant reconnue à environ 40 °C ;

(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'acide éthylène diamine tétra acétique (EDTA), l'acide éthylène glycolbis-(β -aminoéthyléther) N,N,N',N''-tétra acétique (EGTA), la N,N-bis(carboxyméthyl)glycine (acide nitrilo triacétique) (NTA), le tripolyphosphate de sodium (STPP) et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par des agents tensio-actifs tels que les (alkyl linéaire) benzène sulfonates de sodium (LAS), les alkyl sulfates de sodium (AS), les alkyl sulfates de sodium polyoxyéthylénés (ES), les alpha-oléfine sulfonates de sodium (AOS), les esters d'acides aliphatiques alpha-sulfonés sodiques (alpha-SFE), les alkyl sulfonates de sodium (SAS), les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium), et le chlorure de diméthyl dialkyl ammonium ;

(8) Influences des protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel, à l'aide de Sephadex® G 100)

Elle présente un pic maximum à 180 000 $\pm$ 10 000.

2. Procédé de fabrication de cellulase alcaline K ayant les caractéristiques telles que définies à la revendication 1, qui comprend les opérations consistant à cultiver *Bacillus* sp KSM-635 produisant de la cellulase alcaline K, déposé sous le numéro FERM BP-1485, et à recueillir la cellulase alcaline K à partir du produit de culture résultant.

3. CMCCase I, qui peut être obtenue à partir d'un produit de culture de *Bacillus* sp KSM-635, déposé sous le numéro FERM BP-1485, ayant les propriétés physiques et chimiques suivantes :



(1) Activité

Elle présente l'activité enzymatique Cx d'action sur la CMC en même temps qu'une faible activité enzymatique C₁ et une faible activité bêta-glucosidase ;

(2) Spécificité pour les substrats

Elle agit sur la CMC, la cellulose cristalline, l'Avicel, la cellobiose, et le PNPC ;

(3) pH d'activité et pH optimal

Elle présente un pH d'activité se situant dans la plage de 3 à 12,5 et un pH optimal se situant dans la plage de 6 à 11,5 ;

(4) Stabilité au pH

Elle n'est pas inactivée à un pH de 5 à 12 lorsqu'on la laisse reposer à 30 °C pendant 1 heure ;

(5) Température d'activité et température d'activité optimale

Elle a une activité dans une plage de température de 10 à 60 °C, une température optimale se situant à une valeur de 22 à 53 °C ;

(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'EDTA, l'EGTA, le NTA, le STPP et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par les LAS, les AS, les ES, les AOS, les alpha-SFE, les SAS, les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium), le chlorure de diméthyl dialkyl ammonium, et l'acide taurocholique ;

(8) Résistance aux protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel à l'aide de Toyopearl® 55S)

La masse moléculaire de cette enzyme est de 145 000 ± 10 000 ;

(10) Spectre d'absorption du rayonnement UV

Cette enzyme présente une absorption maximale à 280 nm et un épaulement à 290 nm, des composants de type sucre étant de ce fait contenus.

4. CMCase II qui peut être obtenue à partir du produit de culture de *Bacillus* sp KSM-635, déposé sous le numéro FERM BP-1485, ayant les propriétés physiques et chimiques suivantes :

(1) Activité

Elle présente l'activité enzymatique Cx d'action sur la CMC en même temps qu'une faible activité enzymatique C₁ et une faible activité bêta-glucosidase ;

(2) Spécificité pour les substrats

Elle agit sur la CMC, la cellulose cristalline, l'Avicel, la cellobiose, et le PNPC ;

(3) pH d'activité et pH optimal

Elle présente un pH d'activité se situant dans la plage de 3 à 12,5 et un pH optimal se situant dans la plage de 6 à 11,5 ;

(4) Stabilité au pH

Elle n'est pas inactivée à un pH de 5 à 12 lorsqu'on la laisse reposer à 30 °C pendant 1 heure ;

(5) Température d'activité et température d'activité optimale

Elle a une activité dans une plage de température de 5 à 58 °C, une température optimale se situant à une valeur de 14 à 45 °C ;

(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'EDTA, l'EGTA, le NTA, le STPP et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par les LAS, les AS, les ES, les AOS, les alpha-SFE, les SAS, les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium) et le chlorure de diméthyl dialkyl ammonium ;

(8) Résistance aux protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel à l'aide de Toyopearl® 55S)

La masse moléculaire de la CMCase II est de 170 000 ± 20 000 ;

(10) Spectre d'absorption du rayonnement UV

Elle présente une absorption maximale à 280 nm et un épaulement à 290 nm.

5. Bactérie *Bacillus* sp KSM-635, déposée sous le numéro FERM BP-1485, capable de produire la cellulase alcaline K ayant les propriétés enzymatiques suivantes :



(1) Activité

Elle présente une activité enzymatique C_x d'action sur la CMC en même temps qu'une faible activité enzymatique C_1 et une faible activité bêta-glucosidase ;

(2) Spécificité pour les substrats

Elle agit sur la CMC, la cellulose cristalline, l'Avicel, la cellobiose, et le PNPC ;

(3) pH d'activité et pH optimal

Elle présente un pH d'activité se situant dans la plage de 4 à 12 et un pH optimal se situant dans la plage de 9 à 10 (Fig. 3) ;

(4) Stabilité au pH

Elle est stable aux valeurs de pH stable de 4,5 à 10,5 et de 6,8 à 10, lorsqu'on la laisse reposer à 40 °C respectivement pendant 10 minutes et 30 minutes (Fig. 4) ;

(5) Plage de température d'activité et température d'activité optimale

Elle a une activité dans une large plage de température de 10 à 65 °C, une température optimale étant reconnue à environ 40 °C (Fig. 5) ;

(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'EDTA, l'EGTA, le NTA, le STPP et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par des agents tensio-actifs tels que les LAS, les AS, les ES, les AOS, les alpha-SFE, les SAS, les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium), et le chlorure de diméthyl dialkyl ammonium ;

(8) Influences des protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel à l'aide de Sephadex® G 100)

Elle présente un pic maximum à 180 000 $\pm$ 10 000 (Fig.7).

Revendications pour l'Etat contractant suivant : ES

1. Procédé de fabrication de cellulase alcaline K, de CMCase I et de CMCase II pouvant être obtenues à partir de la bactérie *Bacillus* sp KSM-635 déposée sous le numéro FERM BP-1485, caractérisé par le fait qu'il comprend la culture aérobie de ladite bactérie, à une température de 22 à 40 °C, à un pH de 8 à 11, à l'aide d'un milieu de culture qui contient une combinaison appropriée de sources d'azote et de sources de carbone, en faisant suivre par l'isolement des produits désirés à l'aide de techniques connues de collecte et de purification, ce par quoi la cellulase alcaline K est isolée par séparation des cellules de bacillus, telle qu'une centrifugation, une filtration, etc., en faisant suivre par des procédés ordinaires d'isolement, tels que le relargage, la précipitation isoélectrique, la précipitation par solvant, pour séparer les produits sous la forme d'un précipité ; et ce par quoi la CMCase I et CMCase II sont obtenues à partir de la cellulase alcaline K, au moyen d'une chromatographie liquide haute performance de préparation, une chromatographie sur hydroxy apatite ou une chromatographie sur DEAE-Toyopearl®, permettant ainsi d'isoler les constituants respectifs.

2. Procédé selon la revendication 1, caractérisé par le fait que la cellulase alcaline K présente les propriétés physiques et chimiques suivantes :

(1) Activité

Elle présente une activité enzymatique C_x d'action sur la carboxy méthyl cellulose, en même temps qu'une faible activité enzymatique C_1 et une faible activité bêta-glucosidase ;

(2) Spécificité pour les substrats

Elle agit sur la carboxy méthyl cellulose (CMC), la cellulose cristalline, l'Avicel, la cellobiose, et le p-nitrophényl cellobioside (PNPC) ;

(3) pH d'activité et pH optimal

Elle a une activité à un pH se situant dans la plage de 4 à 12 et un pH optimal se situant dans la plage de 9 à 10 ;

(4) Stabilité au pH

Elle est stable aux valeurs de pH de 4,5 à 10,5 et de 6,8 à 10, lorsqu'on la laisse reposer à 40 °C respectivement pendant 10 minutes et pendant 30 minutes ;

(5) Plage de température d'activité et température d'activité optimale

Elle agit dans une large plage de température de 10 à 65 °C, une température optimale étant reconnue à environ 40 °C ;



(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'acide éthylène diamine tétra acétique (EDTA), l'acide éthylène glycol-bis-(β -aminoéthyléther) N,N,N',N''-tétra acétique (EGTA), la N,N-bis(carboxyméthyl)glycine (acide nitrilo triacétique) (NTA), le tripolyphosphate de sodium (STPP) et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par des agents tensio-actifs tels que les (alkyl linéaire) benzène sulfonates de sodium (LAS), les alkylsulfates de sodium (AS), les alkylsulfates de sodium polyoxyéthylénés (ES), les alpha-oléfine sulfonates de sodium (AOS), les esters d'acides aliphatiques alpha-sulfonés sodiques (alpha-SFE), les alkyl sulfonates de sodium (SAS), les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium), et le chlorure de diméthyl dialkyl ammonium ;

(8) Influences des protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel, à l'aide de Sephadex® G.100)

Elle présente un pic maximum à $180\,000 \pm 10\,000$.

3. Procédé selon la revendication 1, caractérisé par le fait que la CMCase I obtenue comme produit présente les propriétés physiques et chimiques suivantes :

(1) Activité

Elle présente l'activité enzymatique Cx d'action sur la CMC, en même temps qu'une faible activité enzymatique C₁ et une faible activité bêta-glucosidase ;

(2) Spécificité pour les substrats

Elle agit sur la CMC, la cellulose cristalline, l'Avicel, la cellobiose, et le PNPC ;

(3) pH d'activité et pH optimal

Elle présente un pH d'activité se situant dans la plage de 3 à 12,5 et un pH optimal se situant dans la plage de 6 à 11,5 ;

(4) Stabilité au pH

Elle n'est pas inactivée à un pH de 5 à 12 lorsqu'on la laisse reposer à 30 °C pendant 1 heure ;

(5) Température d'activité et température d'activité optimale

Elle agit dans une plage de température de 10 à 60 °C, une température optimale se situant à une valeur de 22 à 53 °C ;

(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'EDTA, l'EGTA, le NTA, le STPP et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par les LAS, les AS, les ES, les AOS, les alpha-SFE, les SAS, les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium), le chlorure de diméthyl dialkyl ammonium, et l'acide taurocholique ;

(8) Résistance aux protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel à l'aide de Toyopearl® 55S)

La masse moléculaire de cette enzyme est de $145\,000 \pm 10\,000$;

(10) Spectre d'absorption du rayonnement UV

Cette enzyme présente une absorption maximale à 280 nm et un épaulement à 290 nm, des composants de type sucre étant de ce fait contenus en son sein.

4. Procédé selon la revendication 1, caractérisé par le fait que la CMCase II obtenue comme produit présente les propriétés physiques et chimiques suivantes :

(1) Activité

Elle présente l'activité enzymatique Cx d'action sur la CMC en même temps qu'une faible activité enzymatique C₁ et une faible activité bêta-glucosidase ;

(2) Spécificité pour les substrats

Elle agit sur la CMC, la cellulose cristalline, l'Avicel, la cellobiose, et le PNPC ;

(3) pH d'activité et pH optimal

Elle présente un pH d'activité se situant dans la plage de 3 à 12,5 et un pH optimal se situant dans la plage de 6 à 11,5 ;

(4) Stabilité au pH

Elle n'est pas inactivée à un pH de 5 à 12 lorsqu'on la laisse reposer à 30 °C pendant 1 heure ;



(5) Température d'activité et température d'activité optimale

Elle agit dans une plage de température de 5 à 58 °C, une température optimale se situant à une valeur de 14 à 45 °C ;

(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'EDTA, l'EGTA, le NTA, le STPP et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par les LAS, les AS, les ES, les AOS, les alpha-SFE, SAS, les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium) et le chlorure de diméthyl dialkyl ammonium ;

(8) Résistance aux protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel à l'aide de Toyopearl® 55S)

La masse moléculaire de la CMCCase II est de 170 000 ± 20 000 ;

(10) Spectre d'absorption UV

Elle présente une absorption maximale à 280 nm et un épaulement à 290 nm.

5. Procédé selon la revendication 1, caractérisé par le fait que la bactérie *Bacillus* sp KSM-635, déposée sous le numéro FERM BP-1485, est capable de produire une enzyme ayant les propriétés enzymatiques suivantes :

(1) Activité

Elle présente une activité enzymatique C_x d'action sur CMC en même temps qu'une faible activité enzymatique C_1 et une faible activité bêta-glucosidase ;

(2) Spécificité pour les substrats

Elle agit sur la CMC, la cellulose cristalline, l'Avicel, la cellobiose, et le PNPC ;

(3) pH d'activité et pH optimal

Elle présente un pH d'activité se situant dans la plage de 4 à 12 et un pH optimal se situant dans la plage de 9 à 10 (Fig. 3) ;

(4) Stabilité de pH

Elle est stable aux valeurs de pH de 4,5 à 10,5 et de 6,8 à 10, lorsqu'on la laisse reposer à 40 °C respectivement, pendant 10 minutes et 30 minutes (Fig. 4) ;

(5) Plage de température d'activité et température d'activité optimale

Elle agit dans une large plage de température de 10 à 65 °C, une température optimale étant reconnue à environ 40 °C (Fig. 5) ;

(6) Influences des agents chélatants

L'activité n'est pas empêchée par l'EDTA, l'EGTA, le NTA, le STPP et la zéolite ;

(7) Influences des agents tensio-actifs

Elle subit une faible inhibition d'activité par des agents tensio-actifs tels que les LAS, les AS, les ES, les AOS, les alpha-SFE, les SAS, les (alkyl secondaire) éthers polyoxyéthylénés, les sels d'acides gras (sels de sodium), et le chlorure de diméthyl dialkyl ammonium ;

(8) Influences des protéinases

Elle présente une forte résistance aux protéinases ;

(9) Masse moléculaire (déterminée par chromatographie sur gel à l'aide de Sephadex® G 100)

Elle présente un pic maximum à 180 000 ± 10 000.

