


PROCESS FOR THE SYNTHESIS OF DERIVATIVES OF 3-AMINO-
TETRAHYDROFURAN-3-CARBOXYLIC ACID AND USE THEREOF AS
MEDICAMENTS

5 **Background and summary**

The present disclosure relates to a process for the manufacturing of substituted
3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) and
their precursors in high optical purity


10



to the precursors of the synthesis of substituted 3-amino-tetrahydrofuran-3-
carboxylic acid amides of general formula (I) in high optical purity, and to the
15 tautomers, enantiomers, diastereomers, mixtures and salts of substituted 3-
amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in high
optical purity, particularly the physiologically acceptable salts thereof with
inorganic or organic acids or bases, which have valuable properties.

20 Thus, the present disclosure relates to the stereoselective preparation of
compounds of the above general formula (I).

Within the meaning of the present disclosure, the high optical purity of the substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) and of the precursors of these substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) is with respect to the carbon atom in position 3 of the tetrahydrofuran ring, the position of which is depicted by the numbering "3" in the following structural formula (I)



Within the meaning of the present disclosure, "highly optical pure" means in an enantiomeric excess of more than 96%, preferably of more than 98%.

The disclosure also relates to pharmaceutical compositions containing a compound or a physiologically acceptable salt of a compound of the above general formula (I) according to the embodiments defined below and in the Examples, optionally together with one or more inert carriers and/or diluents.

The disclosure also relates to the use of a compound or a physiologically acceptable salt of a compound according to the embodiments defined below and in the Examples, for preparing a pharmaceutical composition with an inhibitory effect on Factor Xa, an inhibitory effect on related serine proteases, and/or an antithrombotic activity.

Although the pharmacologically valuable properties of the compounds in accordance with the present disclosure constitute the basic prerequisite for effective use of the compounds in a pharmaceutical composition, an active substance must in any case satisfy additional requirements in order to be
5 accepted for use as a drug. These parameters are largely connected with the physicochemical nature of the active substance. Hence, there continues to be a need for crystalline forms of active substances, which can be conveniently formulated for administration to patients and which are pure and highly crystalline in order to fulfil exact pharmaceutical requirements and
10 specifications.

Preferably, such compounds will be readily formed and have favourable bulk characteristics. Examples of favourable bulk characteristics are drying times, filterability, solubility, intrinsic dissolution rate and stability in general.

15

As the crystal modification of an active substance is important to the reproducible active substance content of a preparation, there is a need to clarify as far as possible any existing polymorphism of an active substance present in crystalline form. If there are different polymorphic modifications of an active
20 substance care must be taken to ensure that the crystalline modification of the substance does not change in the pharmaceutical preparation later produced from it. Otherwise, this could have a harmful effect on the reproducible potency of the drug. Against this background, active substances characterised by only slight polymorphism are preferred.

25

Decreased levels of organic solvents in the crystal lattice are also favourable, due in part to potential solvent toxicity to the recipient as a function of the solvent.

- 5 Another criterion which may be of exceptional importance under certain circumstances depending on the choice of formulation or the choice of manufacturing process is the solubility and dissolution rate of the active substance. If for example pharmaceutical solutions are prepared (e.g. for infusions) it is essential that the active substance should be sufficiently soluble
10 in physiologically acceptable solvents. For drugs which are to be taken orally, it is in general very important that the active substance should be sufficiently soluble over a suitable range of pH, and bioavailable.

Hence, without being restrictive, examples of the parameters which need to be
15 controlled are the stability of the starting substance under various environmental conditions, the stability during production of the pharmaceutical formulation and the stability in the final compositions of the drug.

The pharmaceutically active substance used to prepare the pharmaceutical
20 compositions should therefore have great stability which is ensured even under all kinds of environmental conditions.


The present disclosure further provides a pharmaceutically active substance which is not only characterised by high pharmacological potency but also
25 satisfies the physicochemical requirements of high purity and high crystallinity

mentioned hereinbefore, in order to fulfil as far as possible exact pharmaceutical requirements and specifications.

Hence, the present disclosure relates as well to an anhydrous crystalline form of
5 a compound which may be obtained via the manufacturing process in accordance with the present disclosure, namely the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide, to a process for the manufacture thereof, and to the use thereof in a pharmaceutical
10 composition. The structure of this compound is depicted below in the form of the free base as example 2. The characterizing data of its anhydrous crystalline form are described further below in the experimental section.


The present disclosure also provides a process for the manufacture of the
15 anhydrous crystalline form of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide, which is described further below in the experimental section.

In a first aspect, there is provided a substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I)



in an enantiomeric excess of more than 96% at the carbon atom in position 3 of the tetrahydrofuran ring, wherein

D denotes D¹ a substituted bicyclic ring system of formula (II),



wherein

K¹ and K⁴

each independently of one another denote a -CH₂, -CHR^{7a}, -CR^{7b}R^{7c} or a -C(O) group, and

R^{7a}/R^{7b}/R^{7c}

each independently of one another denote a fluorine atom, a hydroxy, C₁₋₅-alkyloxy, amino, C₁₋₅-alkylamino, di-(C₁₋₅-alkyl)-amino, C₃₋₅-cycloalkyleneimino or C₁₋₅-alkylcarbonylamino group,

a C₁₋₅-alkyl group which may be substituted by 1-3 fluorine atoms,

or

two groups R^{7b}/R^{7c} together with the cyclic carbon atom may form a 3, 4, 5-, 6-

5 or 7-membered saturated carbocyclic group

wherein the methylene groups thereof may be substituted by 1-2 C₁₋₃-alkyl or

CF₃- groups, and/or

the methylene groups thereof, if they are not bound to a heteroatom, may be

substituted by 1-2 fluorine atoms, and

10

K² and K³

each independently of one another denote a -CH₂-, -CHR^{8a}-, -CR^{8b}R^{8c}- or
a -C(O)- group, and

15 R^{8a}/R^{8b}/R^{8c}

each independently of one another denote a C₁₋₅-alkyl group which may be
substituted by 1-3 fluorine atoms,

or two groups R^{8b}/R^{8c} together with the cyclic carbon atom may form a 3, 4, 5-,

20 6- or 7-membered saturated carbocyclic group, and

in all there should be no more than four groups selected from among R^{7a}, R^{7b},
R^{7c}, R^{8a}, R^{8b} and R^{8c} in formula (II), and

25 X denotes a NR¹ group, wherein

R¹ denotes a hydrogen atom or a hydroxy, C₁₋₃-alkyloxy, amino, C₁₋₃-alkylamino, di-(C₁₋₃-alkyl)-amino, a C₁₋₅-alkyl, C₂₋₅-alkenyl-CH₂, C₂₋₅-alkynyl-CH₂ or C₃₋₆-cycloalkyl group,

5 wherein the methylene and methyl groups present in the above-mentioned groups

may additionally be substituted by a C₁₋₃-alkyl, carboxy, C₁₋₅-alkoxycarbonyl group or

by a hydroxy, C₁₋₅-alkyloxy, amino, C₁₋₅-alkylamino, C₁₋₅-dialkylamino or C₄₋₇-

10 cycloalkyleneimino group,

provided that O-C-O or O-C-N or N-C-N-bonds are excluded and/or

one to three hydrogen atoms may be replaced by fluorine atoms, provided that the methylene or methyl groups are not directly bound to a nitrogen atom,

15 and wherein

A¹ denotes either N or CR¹⁰,

A² denotes either N or CR¹¹,


20

A³ denotes either N or CR¹²,


wherein R¹⁰, R¹¹ and R¹² each independently of one another represent

a hydrogen, fluorine, chlorine, bromine or iodine atom, or a C₁₋₅-alkyl, CF₃, C₂₋₅-alkenyl, C₂₋₅-alkynyl, a cyano, carboxy, C₁₋₅-alkyloxycarbonyl, hydroxy, C₁₋₃-alkyloxy, CF₃O, CHF₂O, CH₂FO, or


5 D denotes D² a group of general formula



wherein A denotes A⁴, a group



10 or wherein A denotes A⁵, a group of general formula



15

wherein

m is the number 1 or 2,

5 X^1 denotes a carbonyl, thiocarbonyl, $C=NR^{9c}$, $C=N-OR^{9c}$, $C=N-NO_2$, $C=N-CN$ or sulphonyl group,

X^2 denotes an oxygen atom or a $-NR^{9b}$ group,

10 X^3 denotes a carbonyl, thiocarbonyl, $C=NR^{9c}$, $C=N-OR^{9c}$, $C=N-NO_2$, $C=N-CN$ or sulphonyl group,

X^4 denotes an oxygen or sulphur atom or a $-NR^{9c}$ group,

15 R^{9a} in each case independently of one another denotes a hydrogen or halogen atom or a C_{1-5} -alkyl, hydroxy, hydroxy- C_{1-5} -alkyl, C_{1-5} -alkoxy, C_{1-5} -alkoxy- C_{1-5} -alkyl, amino, C_{1-5} -alkylamino, di- $(C_{1-5}$ -alkyl)-amino, amino- C_{1-5} -alkyl, C_{1-5} -alkylamino- C_{1-5} -alkyl, di- $(C_{1-5}$ -alkyl)-amino- C_{1-5} -alkyl, aminocarbonyl, C_{1-5} -alkylaminocarbonyl, di- $(C_{1-5}$ -alkyl)-aminocarbonyl or C_{1-5} -alkylcarbonylamino group, while
20 in the previously mentioned substituted 5- to 7-membered groups A^5 the heteroatoms F, Cl, Br, I, O or N optionally introduced with R^{9a} as substituent are not separated by precisely one carbon atom from a heteroatom selected from among N, O, S,

R^{9b} each independently of one another denote a hydrogen atom or a C_{1-5} -alkyl group,


R^{9c} each independently of one another denote a hydrogen atom, a C_{1-5} -alkyl,
5 C_{1-5} -alkylcarbonyl, C_{1-5} -alkyloxycarbonyl or C_{1-5} -alkylsulphonyl group,

R^4 denotes a hydrogen or halogen atom, a C_{1-3} -alkyl or C_{1-3} -alkoxy group,
while the hydrogen atoms of the C_{1-3} -alkyl or C_{1-3} -alkoxy group may optionally
be wholly or partly replaced by fluorine atoms, a C_{2-3} -alkenyl, C_{2-3} -alkynyl,
10 nitrile, nitro or amino group,

R^5 denotes a hydrogen or halogen atom or a C_{1-3} -alkyl group,

R^3 denotes a hydrogen atom or a C_{1-3} -alkyl group, and
15

M denotes a thiophene ring according to formula (III),



20 which is bound to the carbonyl group in formula (I) via the 2-position and which
is substituted in the 5-position by R^2 and optionally additionally by R^6 , wherein

R^2 denotes

R^{2a} a hydrogen, fluorine or iodine atom, or

R^{2b} a methoxy, C₁₋₂-alkyl, formyl, NH₂CO, or

R^{2c} a chlorine, bromine or ethynyl group,

- 5 R⁶ denotes a hydrogen, fluorine, chlorine, bromine or iodine atom or a C₁₋₂-alkyl or amino group,

wherein, unless otherwise stated, by the term "halogen atom" mentioned
hereinbefore in the definitions is meant an atom selected from among fluorine,
10 chlorine, bromine and iodine,


and wherein the alkyl, alkenyl, alkynyl and alkyloxy groups contained in the
previously mentioned definitions which have more than two carbon atoms may,
unless otherwise stated, be straight-chain or branched and the alkyl groups in
15 the previously mentioned dialkylated groups, for example the dialkylamino
groups, may be identical or different,

and the hydrogen atoms of the methyl or ethyl groups contained in the
foregoing definitions, unless otherwise stated, may be wholly or partly replaced
20 by fluorine atoms,


the tautomers, enantiomers, diastereomers, mixtures and salts thereof.

In a second aspect, there is provided a compound as defined above, which is selected from the following list of compounds and mixtures and salts thereof:



(S)-3-[(5-bromo-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1R)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1S)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(S)-5-chloro-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-bromo-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-oxo-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide



(S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide




, and

(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-(3,5,5-trimethyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




In a third aspect, there is provided a method for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) as defined above in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein the enantiomers are separated via chiral column chromatography.


In a fourth aspect, there is provided a method for the preparation of substituted 3-amino tetrahydrofuran-3-carboxylic acid amides of general formula (Ia) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



comprising reacting a compound of the general formula (IVa)



with a compound of the general formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring




optionally further comprising cleaving protecting groups,


wherein K1, K2, K3, K4, X, A1, A2, A3, R2, R3 and R6 are as defined above, and

wherein Q is a hydroxy or C₁₋₄-alkyloxy group, a halogen atom or a C₁₋₅-alkyloxycarbonyloxy or acyloxy group.


In a fifth aspect, there is provided a method for the preparation of substituted 3-amino tetrahydrofuran-3-carboxylic acid amides of general formula (Ib) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



comprising reacting a compound of the general formula (IVb)



with a compound of the general formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring




optionally further comprising cleaving protecting groups

wherein A, R₄, R₅, R₂, R₃ and R₆ are as defined above and



wherein Q is a hydroxy or C₁₋₄-alkyloxy group, a halogen atom or a C₁₋₅-alkyloxycarbonyloxy or acyloxy group.

In a sixth aspect, there is provided a method for the preparation of substituted 3-amino tetrahydrofuran-3-carboxylic acid amides of general formula (Ic) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring




comprising the steps of:


a) reacting a compound of the formula (IV) with a compound of the formula (VI) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring,



and subsequently cleaving the protecting group PG to obtain a compound of the formula (VII) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring; and



b) reacting said compound (VII) of step a) with a compound of formula (VIII)




wherein

Q is a hydroxy or C₁₋₄-alkyloxy group, a halogen atom or a C₁₋₅-alkyloxycarbonyloxy or acyloxy group,

PG is a hydrogen atom or a protective group for the amino function, and

D, R₃, R₂ and R₆ are as defined above.

In a seventh aspect, there is provided a method for preparing a compound of the formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring




by enzymatic resolution of a racemic mixture of said compound of the formula (V),

wherein R² and R⁶ are as defined above, and

wherein Q is a straight or substituted C₁₋₁₂-alkyloxy group, or allyloxy or substituted allyloxy group, or C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group.

In an eighth aspect, there is provided a compound of the formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring,




wherein R² and R⁶ are as defined above, and

wherein Q is a hydroxy or C₁₋₁₂-alkyloxy group, or allyloxy or substituted allyloxy group, or a halogen atom or a C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group.


In a ninth aspect, there is provided the compound of the formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring as defined above, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of said compound of the general formula (V) has the S-configuration.

In a tenth aspect, there is provided a method for preparing a compound of the formula (VII) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



by chemical resolution of a racemic mixture of said compound of the formula (VII) with a chiral acid, wherein D and R₃ are as defined above.

In an eleventh aspect, there is provided a compound of the formula (VII) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



wherein D and R₃ are as defined above.

Brief Description of the Figures

Figure 1 shows the X-ray powder diffractogram of the anhydrous crystalline form of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide, recorded using a STOE Stadi P-diffractometer fitted with a location-

sensitive detector (OED) and a Cu anode as the x-ray source and a Germanium monochromator (CuK α radiation, λ = 1.54056 Å, 40 kV, 40 mA).


Figure 2 shows a light microscopy photograph of crystals of the anhydrous crystalline form of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide.

Figure 3 shows the thermoanalysis and determination of the melting point and loss on drying (DSC/TG) of the anhydrous crystalline form of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide, recorded using a DSC and TG and evaluated by the peak onset (heating rate: 10°C/min) for the melting point and by the weight loss step between room temperature and 180 °C for the loss on drying. The values given are determined using a DSC 821e and a TGA/STDA 851e, both made by Mettler Toledo.

Detailed Description of the Invention

A 1st embodiment of the present invention includes those compounds of general formula (I) wherein

D denotes D¹ a substituted bicyclic ring system of formula (II),



wherein

K^1 and K^4

- 5 each independently of one another denote a $-\text{CH}_2$, $-\text{CHR}^{7a}$, $-\text{CR}^{7b}\text{R}^{7c}$ or a $-\text{C}(\text{O})$ group, and

$\text{R}^{7a}/\text{R}^{7b}/\text{R}^{7c}$

- each independently of one another denote a fluorine atom, a hydroxy, C_{1-5} -alkyloxy, amino, C_{1-5} -alkylamino, di- $(\text{C}_{1-5}\text{-alkyl})$ -amino, C_{3-5} -cycloalkyleneimino or C_{1-5} -alkylcarbonylamino group,
10 a C_{1-5} -alkyl group which may be substituted by 1-3 fluorine atoms, or

- 15 two groups $\text{R}^{7b}/\text{R}^{7c}$ together with the cyclic carbon atom may form a 3, 4, 5-, 6- or 7-membered saturated carbocyclic group

wherein the methylene groups thereof may be substituted by 1-2 C_{1-3} -alkyl or CF_3 - groups, and/or

- the methylene groups thereof, if they are not bound to a heteroatom, may be
20 substituted by 1-2 fluorine atoms, and

K^2 and K^3

each independently of one another denote a $-\text{CH}_2$, $-\text{CHR}^{8a}$, $-\text{CR}^{8b}\text{R}^{8c}$ or a $-\text{C}(\text{O})$ - group, and

$R^{8a}/R^{8b}/R^{8c}$

each independently of one another denote a C₁₋₅-alkyl group which may be substituted by 1-3 fluorine atoms,

- 5 or two groups R^{8b}/R^{8c} together with the cyclic carbon atom may form a 3, 4, 5-, 6- or 7-membered saturated carbocyclic group, and

in all there should be no more than four groups selected from among R^{7a} , R^{7b} , R^{7c} , R^{8a} , R^{8b} and R^{8c} in formula (II), and

10

X denotes a NR^1 group, wherein

R^1 denotes a hydrogen atom or a hydroxy, C₁₋₃-alkyloxy, amino, C₁₋₃-alkylamino, di-(C₁₋₃-alkyl)-amino, a C₁₋₅-alkyl, C₂₋₅-alkenyl-CH₂,

- 15 C₂₋₅-alkynyl-CH₂ or C₃₋₆-cycloalkyl group,

wherein the methylene and methyl groups present in the above-mentioned groups

may additionally be substituted by a C₁₋₃-alkyl, carboxy, C₁₋₅-alkoxycarbonyl group or

- 20 by a hydroxy, C₁₋₅-alkyloxy, amino, C₁₋₅-alkylamino, C₁₋₅-dialkylamino or C₄₋₇-cycloalkyleneimino group,

provided that O-C-O or O-C-N or N-C-N-bonds are excluded and/or

one to three hydrogen atoms may be replaced by fluorine atoms, provided that the methylene or methyl groups are not directly bound to a nitrogen atom,

25

and wherein

A^1 denotes either N or CR^{10} ,

5 A^2 denotes either N or CR^{11} ,


A^3 denotes either N or CR^{12} ,

wherein R^{10} , R^{11} and R^{12} each independently of one another represent


10

a hydrogen, fluorine, chlorine, bromine or iodine atom, or a C_{1-5} -alkyl, CF_3 , C_{2-5} -alkenyl, C_{2-5} -alkynyl, a cyano, carboxy, C_{1-5} -alkyloxycarbonyl, hydroxy, C_{1-3} -alkyloxy, CF_3O , CHF_2O , CH_2FO , or


15 D denotes D^2 a group of general formula



wherein A denotes A^4 , a group



20 or wherein A denotes A^5 , a group of general formula



5

wherein

m is the number 1 or 2,

10 X^1 denotes a carbonyl, thiocarbonyl, $C=NR^{9c}$, $C=N-OR^{9c}$, $C=N-NO_2$, $C=N-CN$ or sulphonyl group,

X^2 denotes an oxygen atom or a $-NR^{9b}$ group,

15 X^3 denotes a carbonyl, thiocarbonyl, $C=NR^{9c}$, $C=N-OR^{9c}$, $C=N-NO_2$, $C=N-CN$ or sulphonyl group,

X^4 denotes an oxygen or sulphur atom or a $-NR^{9c}$ group,

20 R^{9a} in each case independently of one another denotes a hydrogen or halogen atom or a C_{1-5} -alkyl, hydroxy, hydroxy- C_{1-5} -alkyl, C_{1-5} -alkoxy, C_{1-5} -

alkoxy-C₁₋₅-alkyl, amino, C₁₋₅-alkylamino, di-(C₁₋₅-alkyl)-amino, amino-C₁₋₅-alkyl, C₁₋₅-alkylamino-C₁₋₅-alkyl, di-(C₁₋₅-alkyl)-amino-C₁₋₅-alkyl, aminocarbonyl, C₁₋₅-alkylaminocarbonyl, di-(C₁₋₅-alkyl)-aminocarbonyl or C₁₋₅-alkylcarbonylamino group, while

- 5 in the previously mentioned substituted 5- to 7-membered groups A⁵ the heteroatoms F, Cl, Br, I, O or N optionally introduced with R^{9a} as substituent are not separated by precisely one carbon atom from a heteroatom selected from among N, O, S,

- 10 R^{9b} each independently of one another denote a hydrogen atom or a C₁₋₅-alkyl group,

R^{9c} each independently of one another denote a hydrogen atom, a C₁₋₅-alkyl, C₁₋₅-alkylcarbonyl, C₁₋₅-alkyloxycarbonyl or C₁₋₅-alkylsulphonyl group,

15


R⁴ denotes a hydrogen or halogen atom, a C₁₋₃-alkyl or C₁₋₃-alkoxy group, while the hydrogen atoms of the C₁₋₃-alkyl or C₁₋₃-alkoxy group may optionally be wholly or partly replaced by fluorine atoms, a C₂₋₃-alkenyl, C₂₋₃-alkynyl, nitrile, nitro or amino group,

20

R⁵ denotes a hydrogen or halogen atom or a C₁₋₃-alkyl group,

R³ denotes a hydrogen atom or a C₁₋₃-alkyl group, and

- 25 M denotes a thiophene ring according to formula (III),



which is bound to the carbonyl group in formula (I) via the 2-position and which
 5 is substituted in the 5-position by R^2 and optionally additionally by R^6 , wherein

R^2 denotes

R^{2a} a hydrogen, fluorine or iodine atom, or

R^{2b} a methoxy, C_{1-2} -alkyl, formyl, NH_2CO , or

10 R^{2c} a chlorine, bromine atom or an ethynyl group,

R^6 denotes a hydrogen, fluorine, chlorine, bromine or iodine atom or a C_{1-2} -alkyl or amino group,

15 wherein, unless otherwise stated, by the term "halogen atom" mentioned
 hereinbefore in the definitions is meant an atom selected from among fluorine,
 chlorine, bromine and iodine,

and wherein the alkyl, alkenyl, alkynyl and alkyloxy groups contained in the
 20 previously mentioned definitions which have more than two carbon atoms may,
 unless otherwise stated, be straight-chain or branched and the alkyl groups in
 the previously mentioned dialkylated groups, for example the dialkylamino
 groups, may be identical or different,

and the hydrogen atoms of the methyl or ethyl groups contained in the foregoing definitions, unless otherwise stated, may be wholly or partly replaced by fluorine atoms,

5

the tautomers, enantiomers, diastereomers, mixtures and salts thereof.

Examples of the C₁₋₆-alkyl groups mentioned hereinbefore in the definitions are the methyl, ethyl, 1-propyl, 2-propyl, *n*-butyl, *sec*-butyl, *tert*-butyl, 1-pentyl, 2-pentyl, 3-pentyl, *neo*-pentyl, 3-methyl-2-butyl, 1-hexyl, 2-hexyl, 3-hexyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 3-methyl-3-pentyl, 2-methyl-3-pentyl, 2,2-dimethyl-3-butyl or 2,3-dimethyl-2-butyl group.

10

Examples of the C₁₋₅-alkyloxy groups mentioned hereinbefore in the definitions are the methyloxy, ethyloxy, 1-propyloxy, 2-propyloxy, *n*-butoxy, *sec*-butoxy, *tert*-butoxy, 1-pentyloxy, 2-pentyloxy, 3-pentyloxy or *neo*-pentyloxy group.

15

Examples of the C₂₋₅-alkenyl groups mentioned hereinbefore in the definitions are the ethenyl, 1-propen-1-yl, 2-propen-1-yl, 1-buten-1-yl, 2-buten-1-yl, 3-buten-1-yl, 1-penten-1-yl, 2-penten-1-yl, 3-penten-1-yl, 4-penten-1-yl, 1-hexen-1-yl, 2-hexen-1-yl, 3-hexen-1-yl, 4-hexen-1-yl, 5-hexen-1-yl, but-1-en-2-yl, but-2-en-2-yl, but-1-en-3-yl, 2-methyl-prop-2-en-1-yl, pent-1-en-2-yl, pent-2-en-2-yl, pent-3-en-2-yl, pent-4-en-2-yl, pent-1-en-3-yl, pent-2-en-3-yl, 2-methyl-but-1-en-1-yl, 2-methyl-but-2-en-1-yl, 2-methyl-but-3-en-1-yl or 2-ethyl-prop-2-en-1-yl group.

20


25

Examples of the C₂₋₅-alkynyl groups mentioned hereinbefore in the definitions are the ethynyl, 1-propynyl, 2-propynyl, 1-butyn-1-yl, 1-butyn-3-yl, 2-butyn-1-yl, 3-butyn-1-yl, 1-pentyn-1-yl, 1-pentyn-3-yl, 1-pentyn-4-yl, 2-pentyn-1-yl, 2-pentyn-3-yl, 3-pentyn-1-yl, 4-pentyn-1-yl, 2-methyl-1-butyn-4-yl, 3-methyl-1-butyn-1-yl or 3-methyl-1-butyn-3-yl group.

A 2nd embodiment of the present invention includes those compounds of general formula (I), wherein

10

D denotes a substituted bicyclic ring system of formula (II),



(II)

wherein

15

K¹ and K⁴

each independently of one another denote a -CH₂, -CHR^{7a}, or a -CR^{7b}R^{7c} group, wherein

20 R^{7a}/R^{7b}/R^{7c}

each independently of one another denote a fluorine atom, a hydroxy, methoxy or C₁₋₂-alkyl group which may be substituted by 1-3 fluorine atoms,

wherein the two groups R^{7b}/R^{7c} cannot both simultaneously be bound to the cyclic carbon atom via a heteroatom, except if $-C(R^{7b}R^{7c})-$ corresponds to a $-CF_2$ group, or

- 5 two groups R^{7b}/R^{7c} may form, together with the cyclic carbon atom, a 3-, 4- or 5-membered saturated carbocyclic group, and

K^2 and K^3

- each independently of one another represent a $-CH_2$, $-CHR^{8a}$, or
10 $-CR^{8b}R^{8c}$ group, and

$R^{8a}/R^{8b}/R^{8c}$

each independently of one another denote a C_{1-2} -alkyl group which may be substituted by 1-3 fluorine atoms,

15

or two groups R^{8b}/R^{8c} may form, together with the cyclic carbon atom, a 3-, 4-, 5-membered carbocyclic group, and

in all in formula (II) there should be no more than four groups selected from

- 20 among R^{7a} , R^{7b} , R^{7c} , R^{8a} , R^{8b} and R^{8c} , and

X denotes an NR^1 group, wherein

- 25 R^1 denotes a hydrogen atom or

a C₁₋₂-alkyl or C₃₋₄-cycloalkyl group,

wherein the methylene and methyl groups present in the above-mentioned

groups may additionally be substituted by a methyl group ,

5 and wherein

A¹ denotes CR¹⁰,

A² denotes CR¹¹,

10


A³ denotes CR¹²,

wherein R¹⁰, R¹¹ and R¹² each independently of one another represent

15 a hydrogen, fluorine, chlorine, bromine atom or a Methyl, CF₃, a cyano,


Methoxy, CF₃O, CHF₂O, CH₂FO- group, or

D denotes D² a group of general formula




20

wherein A denotes A⁴, a group



or wherein A denotes A⁵, a group of general formula



wherein

10 m is the number 1 or 2,

X^1 denotes a carbonyl or C=N-CN group,

X^3 denotes a carbonyl or C=N-CN group,

15

X^4 denotes an oxygen atom,

R^{9a} in each case independently of one another denotes a hydrogen atom or a C₁₋₂-alkyl group, while


R^4 denotes a hydrogen or fluorine, chlorine or bromine atom, a methyl or a methoxy group,

5 R^5 denotes a hydrogen, fluorine or chlorine atom or a methyl group,

R^3 denotes a hydrogen atom and

M denotes a thiophene ring according to formula (III),

10



which is bound to the carbonyl group in formula (I) via the 2-position and which is substituted in the 5-position by R^2 and optionally additionally by R^6 , wherein

15

R^2 denotes

R^{2c} a chlorine, bromine atom or an ethynyl group,


R^6 denotes a hydrogen atom,

20

the tautomers, diastereomers, mixtures and salts thereof.

A 3rd embodiment of the present invention includes all those compounds of the first and second embodiment, wherein

D denotes a substituted bicyclic ring system of formula (II),




wherein K1, K2, K3, K4, X, A1, A2 and A3 are as defined in the 1st or 2nd embodiment,

the tautomers, diastereomers, mixtures and salts thereof.

10 A 4th embodiment of the present invention includes all those compounds of the first and second embodiment, wherein

D denotes D² a group of general formula



15


wherein R4 and R5 are as defined in the 1st or 2nd embodiment,

the tautomers, diastereomers, mixtures and salts thereof.

A 5th embodiment of the present invention includes all those compounds of the

20 first and second embodiment, wherein

D denotes D² a group of general formula



5 or wherein A denotes A⁵, wherein A⁵, R⁴ and R⁵ are as defined in the 1st or 2nd embodiment,


the tautomers, diastereomers, mixtures and salts thereof.

A 6th embodiment of the present invention includes all those compounds of the
 10 previous embodiments in high optical purity at the carbon in position 3 of the tetrahydrofuran ring, wherein the amino-tetrahydrofuran carboxylic acid amide moiety has the R-configuration.

A 7th embodiment of the present invention includes all those compounds of the
 15 previous embodiments in high optical purity at the carbon in position 3 of the tetrahydrofuran ring, wherein the amino-tetrahydrofuran carboxylic acid amide moiety has the S-configuration.


The following preferred compounds of general formula (I) are mentioned by way
 20 of example, both as the tautomers, mixtures and salts thereof:

(S)-3-[(5-bromo-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



5



(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and


(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-

10 tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1R)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and


15 (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1S)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




- 5 (S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-oxo-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide



- 5 (S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide




(S)-5-ethynyl-thiophene-2-carboxylic acid-*N*-{3-[4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide




and


- (S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(pyrrolidin-1-yl-carbonyl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




- (S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(pyrrolidin-1-yl-carbonyl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




- (S)-5-ethynyl-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(2-oxo-azepan-1-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,




- 5 (S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(2-oxo-azepan-1-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide





- 5 (S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-cyanimino-morpholin-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(3-cyanimino-morpholin-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




- (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((2S)-2,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
- (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((2R)-2,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




- (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((4S)-3,4-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
- (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((2R)-3,4-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




the following compound,




5 the following compound,




(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(2,2,3-trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3,4,4-trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3,5,5-trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-9-fluor-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-8-fluor-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



as well as

(R)-3-[(5-bromo-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide,

5

(R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide,

(3R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
10 (3R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide,

(R)-5-chloro-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-cyanimin-
15 [1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

(R)-5-chloro-thiophene-2-carboxylic acid-N-{3-[4-(5-cyanimin-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

20 (R)-5-bromo-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-oxo-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

(R)-5-bromo-thiophene-2-carboxylic acid-N-{3-[4-(5-cyanimin-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

25

(R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,

5 (R)-5-ethinyl-thiophene-2-carboxylic acid-*N*-{3-[4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,

(R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(pyrrolidin-1-yl-carbonyl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

10 (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(pyrrolidin-1-yl-carbonyl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

(R)-5-ethinyl-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,

15

(R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(2-oxo-azepan-1-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,

20 (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(2-oxo-azepan-1-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide,


(R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

(R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(5-cyanimin-
[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,


(R)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-cyanimino-
5 morpholin-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

(R)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(3-cyanimino-
morpholin-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide,

10 the following compound



the following compound



15 (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-(2,2,3-trimethyl-2,3,4,5-
tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide,

(R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-(3,4,4-trimethyl-2,3,4,5-
tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide,

(R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3,5,5-trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide,


- 5 (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-9-fluor-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide,
and

- (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-8-fluor-2,3,4,5-
10 tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide.


In a preferred embodiment, the following compounds are preferred:

(S)-3-[(5-bromo-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

15




(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



20


(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

5




(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1R)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1S)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

10




(S)-5-chloro-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




15


(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-oxo-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide



- 5 (S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[4-(5-cyanimin-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide



10


(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-(3,5,5-trimethyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




- 5 The invention also relates to physiologically acceptable salts of the compounds according to the embodiments defined above and the Examples.

Description of the manufacturing process of the invention

- 10 a) The preparation of a compound of general formula (Ia)




or




wherein A, A¹ to A³, K¹ to K⁴, X and R¹ to R⁶ are defined as in embodiment 1,
 and which may optionally be protected at any amino, hydroxy, or carboxy
 groups present by the usual protective groups such as for example those
 described in T.W. Greene, P.G.M. Wuts in "Protective Groups in Organic
 5 Synthesis" and the protective groups of which may be cleaved in a manner
 known from the literature,
 is described in the exemplifying embodiments or can be done according to
 known procedures from the literature or may be carried out for example
 according to one of the following formula schemes 1a or 1b or 2:

10


Scheme 1a



Scheme 1b



Scheme 2



where

Q denotes a hydroxy or C₁₋₄-alkyloxy group, a halogen atom or a

5 C₁₋₅-alkyloxycarbonyloxy or acyloxy group and

PG denotes a hydrogen atom or a protective group for the amino function

known from the literature such as for example a tert.-butoxycarbonyl,

benzyloxycarbonyl, p-methoxybenzyloxycarbonyl, allyloxycarbonyl,

10 ethyloxycarbonyl, isopropylloxycarbonyl, 2,2,2-trichloroethyloxycarbonyl,

methyloxycarbonyl, 9-fluorenylmethyloxycarbonyl, 2-

trimethylsilylethoxycarbonyl, phenylethoxycarbonyl, acetyl or a trifluoroacetyl group.

Enantiomerically pure compounds Ia, Ib and Ic can be obtained either by chiral
5 chromatography or chemical resolution of racemic Ia, Ib or Ic or optically active intermediates V, VI or VII can be employed in the synthetic steps described in Scheme 1a, 1b and 2.

Accordingly, in a further embodiment, the present invention includes a method
10 for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ia) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring comprising reacting a compound of the general formula (IVa) with a compound of the general formula (V) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring optionally further comprising
15 cleaving protecting groups, wherein K1, K2, K3, K4, X, A1, A2, A3, R2, R3 and R6, and Q are as defined hereinabove. The amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (V) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ia) may have the R-configuration or the S-configuration.

20

In a further embodiment, the present invention also includes a method for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ib) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring comprising reacting a compound of the general formula
25 (IVb) with a compound of the general formula (V) in high optical purity at the

carbon in position 3 of the tetrahydrofuran ring, optionally further comprising cleaving protecting groups, wherein A, R₄, R₅, R₂, R₃ and R₆, and Q are as defined hereinabove. The amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (V) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ib) may have the R-configuration or the S-configuration.

In a further embodiment, the present invention also includes a method for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ic) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring comprising the steps of: a) reacting a compound of the formula (IV) with a compound of the formula (VI) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring to obtain a compound of the formula (VII) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring, optionally cleaving the amino protecting group; and b) reacting said compound (VII) of step a) with a compound of formula (VIII), wherein Q, PG, D, R₃, R₂ and R₆ are as defined hereinabove. The amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (VI) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ic) may have the R-configuration or the S-configuration.

The reaction steps i) –iii) described in Scheme 1 and 2 may for example be carried out as described in the Examples or under conditions known from the literature, for example as follows:

i) acylation of an amine (IV) or (VII) with an optionally activated carboxylic acid (V) or (VI) or (VIII)


The acylation is expediently carried out with a corresponding halide or
5 anhydride in a solvent such as methylene chloride, chloroform, carbon tetrachloride, ether, ethyl acetate, tetrahydrofuran, dioxane, benzene, toluene, acetonitrile, dimethylformamide, sodium hydroxide solution or sulpholane, optionally in the presence of an inorganic or organic base at temperatures between -20 and 200°C, but preferably at temperatures between -10 and
10 160°C.

The acylation may however also be carried out with the free acid, optionally in the presence of an acid-activating agent or a dehydrating agent, for example in the presence of isobutyl chloroformate, thionyl chloride, trimethylchlorosilane,
15 hydrogen chloride, sulphuric acid, methanesulphonic acid, *p*-toluenesulphonic acid, phosphorus trichloride, 1-propylphosphonic acid cyclic anhydride, phosphorus pentoxide, 2-ethoxy-1-ethoxycarbonyl-1.2-dihydroquinoline (EEDQ), *N,N*-dicyclohexylcarbodiimide, *N,N*-dicyclohexylcarbodiimide/camphorsulphonic acid,
20 *N,N*-dicyclohexylcarbodiimide/*N*-hydroxysuccinimide or 1-hydroxy-benzotriazole, *N,N*-carbonyldiimidazole, *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyl-uronium tetrafluoroborate/*N*-methylmorpholine, *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate/*N*-ethyl-diisopropylamine, *O*-pentafluorophenyl-*N,N,N',N'*-tetramethyluronium-hexafluorophosphate/triethylamine, *N,N*-thionyl-diimidazole or
25

triphenylphosphine/carbon tetrachloride, at temperatures between -20 and 200°C, but preferably at temperatures between -10 and 160°C.

The acylation may also be carried out with a carboxylic acid ester (V) or (VI) and the amine (IV) by activation with trimethylaluminium.

The acylation of a compound of general formula (IV) may however also be carried out with a reactive carboxylic acid derivative of general formula (R- or S- IX)



(R- or S-IX),

wherein R⁶ and R² are defined as in embodiment 1. The acylation is then conveniently carried out in a solvent such as for example toluene, tetrahydrofuran or dimethylformamide, with the addition of an acid such as acetic acid or camphorsulphonic acid or optionally in the presence of a Lewis acid such as zinc chloride or copper(II)chloride and optionally by the addition of amine bases such as for example diisopropylethylamine, triethylamine or N-methylmorpholine, at temperatures between -10 and 100°C, for example using a microwave oven or as described in P.Wipf et al., Helvetica Chimica Acta, 69, 1986, 1153.

Compounds of general formula (IX) may be prepared from compounds of general formula (V), expediently in a solvent or mixture of solvents such as dichloromethane, trichloromethane, carbon tetrachloride, benzene, chlorobenzene, toluene, xylene, hexamethyldisiloxane, ether, tetrahydrofuran, dioxane, acetonitrile, pyridine, optionally in the presence of *N,N'*-dicyclohexylcarbodiimide, *N,N'*-dicyclohexylcarbodiimide/*N*-hydroxysuccinimide or 1-hydroxy-benzotriazole, *N,N'*-carbonyldiimidazole, *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyl-uronium tetrafluoroborate/*N*-methylmorpholine, *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate/*N*-ethyl-diisopropylamine, or in acetic anhydride at temperatures between -20 and 200°C, but preferably at temperatures between -10 and 100°C.

Other methods of amide coupling are described for example in P.D. Bailey, I.D. Collier, K.M. Morgan in "Comprehensive Functional Group Interconversions", Vol. 5, page 257ff., Pergamon 1995 or in Supplementary Volume 22 to Houben-Weyl, Thieme Verlag, 2003 and literature cited therein.

ii) or iii) Cleaving a protective group

20

The optional subsequent cleaving of any protective group used is carried out hydrolytically, for example, in an aqueous solvent, e.g. in water, isopropanol/water, tetrahydrofuran/water or dioxane/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali metal base such as lithium hydroxide, sodium hydroxide or

25


potassium hydroxide or by ether cleavage, e.g. in the presence of iodotrimethylsilane, at temperatures between 0 and 100°C, preferably at temperatures between 10 and 50°C.

- 5 A benzyl, methoxybenzyl or benzyloxycarbonyl group may, however, be cleaved hydrogenolytically, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a solvent such as methanol, ethanol, ethyl acetate, dimethylformamide, dimethylformamide/acetone or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid at temperatures
10 between 0 and 50°C, but preferably at room temperature, and under a hydrogen pressure of 1 to 7 bar, but preferably 1 to 5 bar.

A protective group may however also be cleaved by the methods described in T.W. Greene, P.G.M. Wuts in "Protective Groups in Organic Synthesis".

15

- (b) The components of general formula IV (including IVa and IVb)



- 20 wherein D and R³ are defined as in embodiment 1, and which may optionally be protected at any amino, hydroxy, or carboxy groups present by the usual protective groups such as for example those described in T.W. Greene, P.G.M. Wuts in "Protective Groups in Organic Synthesis" and the protective groups of which may be cleaved in a manner


known from the literature in the course of the synthesis sequence to form compounds of formula (I),

are known from the literature, or their synthesis is described in the exemplifying embodiments, or they may be prepared for example using methods of synthesis

- 5 known from the literature or analogously to methods of synthesis known from the literature as described for example in DE4429079, US4490369, DE3515864, US5175157, DE1921861, WO85/00808 or in G. Bobowski et al., J.Heterocyclic Chem. 16, 1525, 1979 or in P.D. Johnson et al., Bioorg. Med. Chem. Letter 2003, 4197, or in WO04/46138, WO05/111014, WO05/111029 or
10 WO06/34822.

Fragments bridged in the azepine moiety as shown in formula II-1 or II-2 may for example be prepared analogously to J.W. Coe et al. J. Med. Chem., 2005, 48, 3474 or J.W. Coe et al. , US Patent application US2005/0020616,
15 WO05/111014, WO05/111029 or WO06/34822.

- For example, a compound of general formula (IV), wherein R³ denotes a hydrogen atom and A1, A2, A3, K1, K2, K3, K4 and X are defined as in embodiment 1, may be prepared by reduction of the nitro group of a compound
20 of general formula (III)




(III),

wherein A1, A2, A3, K1, K2, K3, K4 and X are defined as in embodiment 1, as follows.

The reduction of the nitro group is for example conveniently carried out in a solvent or mixture of solvents such as water, aqueous ammonium chloride solution, hydrochloric acid, sulphuric acid, phosphoric acid, formic acid, acetic acid, acetic anhydride with base metals such as iron, zinc, tin or sulphur compounds such as ammonium sulphide, sodium sulphide or sodium dithionite or by catalytic hydrogenation with hydrogen, for example under a pressure between 0.5 and 100 bar, but preferably between 1 and 50 bar, or with hydrazine as reducing agent, conveniently in the presence of a catalyst such as for example Raney nickel, palladium charcoal, platinum oxide, platinum on mineral fibres or rhodium, or with complex hydrides such as lithium aluminium hydride, sodium borohydride, sodium cyanoborohydride, diisobutylaluminium hydride, conveniently in a solvent or mixture of solvents such as water, methanol, ethanol, isopropanol, pentane, hexane, cyclohexane, heptane, benzene, toluene, xylene, ethyl acetate, methylpropionate, glycol, glycoldimethylether, diethyleneglycoldimethylether, dioxane, tetrahydrofuran, *N*-methylpyrrolidinone, or *N*-ethyl-diisopropylamine, *N*-C₁₋₅-alkylmorpholine, *N*-C₁₋₅-alkylpiperidine, *N*-C₁₋₅-alkylpyrrolidine, triethylamine, pyridine, for example at temperatures between -30 and 250°C, but preferably between 0 and 150°C.

(c) The components of general formula




wherein R^4 , R^5 , R^6 and R^2 are defined as in embodiment 1, and where

Q denotes a hydroxy or C_{1-4} -alkyloxy group, a halogen atom or a C_{1-5} -alkyloxycarbonyloxy or acyloxy group

- 5 which may optionally be protected at any amino, hydroxy, carboxy or thiol groups present by the usual protective groups such as for example those described in T.W. Greene, P.G.M. Wuts in "Protective Groups in Organic Synthesis" and the protective groups of which may be cleaved in a manner known from the literature in the course of the synthesis sequence to form
 - 10 compounds of formula (I),
- are known from the literature, or their synthesis is described in the exemplifying embodiments, or they may be prepared for example using methods of synthesis known from the literature or analogously to methods of synthesis known from the literature as described for example in WO04/46138, WO05/111014,
- 15 WO05/111029 or WO06/34822.

Scheme 3


-49-



For example they may also be prepared according to Scheme 3 by reacting a compound (VIII) with an amine (VI-1), where Q denotes a hydroxy or C_{1-4} -alkyloxy group, a halogen atom or an alkyloxycarbonyloxy or acyloxy group and Q-I denotes a hydroxy or C_{1-4} -alkyloxy group, which may optionally be converted into Q after the acylation step by saponification and activation as described above. The acylation may be carried out according to the acylation conditions described above.

- 10 The amino acid derivatives (VI-1) are known from the literature or may be prepared analogously to methods known from the literature as described in the Examples, for example, from commercially obtainable amino acid derivatives.

d) The components of general formula V or



(VII),


wherein D and R³ are defined as in embodiment 1,

can be synthesized as described in the exemplifying embodiments, or they may


be prepared for example using methods of synthesis known from the literature

5 or they may be prepared according to Scheme 4.


Scheme 4




Scheme 5



Scheme 5(a)




Scheme 5(b)



5


e) Preparation of a compound of general formula (XI)



wherein R is C₁-C₁₂-alkyl, aryl, or aryl-C₁-C₁₂-alkyl or a heterocycle, for example R is C₁-C₄-alkyl, aryl, or aryl-C₁-C₄-alkyl, alternatively R is methyl, n-butyl or iso-butyl, benzyl, phenethyl.

10

In one embodiment, the present invention relates to a process for the preparation of a compound of the formula (XI). In this process, a Strecker reaction is used in a "one-pot" process followed by in situ formation of ester avoiding the isolation of intermediate product **21** from the reaction mixture. For
5 example, the new process is shown as follow:



In one embodiment, the process uses ketone (X) as the starting material in a
10 reaction with a nitrogen source (for example ammonium acetate or ammonia),
in alcohol (for example methanol or ethanol) and a cyanide source (for example
a cyanide salt, such as sodium cyanide or potassium sodium). The reaction is
for example carried out at ambient temperature. This results in the production of
intermediate **21**. Typically, intermediate **21** is not isolated and is directly treated
15 with alcohol R-OH in the presence of acid (for example HCl) to generate the
desired ester. The process is generally operated in mild condition and readily
available starting materials are used.

Optically pure derivatives of the compounds of the general formula (XI) can be
obtained by the resolution methods described hereinbelow, including chemical
20 resolution for example with L-Mandelic acid, or enzymatic resolution, for
example with alcalase.

Accordingly, in a further embodiment, the invention relates to a compound of
the general formula (XI) in high optical purity. In one aspect the compound of

the general formula (XI) in high optical purity is in the R-configuration or in the S-configuration.

In the reactions described hereinbefore any reactive groups present such as
5 hydroxy, carboxy, amino, alkylamino or imino groups may be protected during the reaction by conventional protective groups which are cleaved again after the reaction.

For example a protecting group for a hydroxy group might be the methoxy,
10 benzyloxy, trimethylsilyl, acetyl, benzoyl, *tert.*-butyl, trityl, benzyl or tetrahydropyranyl group.

Protecting groups for a carboxyl group might be the trimethylsilyl, methyl, ethyl, *tert.*-butyl, benzyl or tetrahydropyranyl group.

15

A protecting group for an amino, alkylamino or imino group might be the acetyl, trifluoroacetyl, benzoyl, ethoxycarbonyl, *tert.*-butoxycarbonyl, benzyloxycarbonyl, benzyl, methoxybenzyl or 2,4-dimethoxybenzyl group and additionally, for the amino group, the phthalyl group.

20

For example a protecting group for an ethynyl group might be the trimethylsilyl, diphenylmethylsilyl, *tert.*-butyldimethylsilyl or a 1-hydroxy-1-methyl-ethyl group.

Other protective groups which may be used and their removal are described in T.W. Greene, P.G.M. Wuts, "Protective Groups in Organic Synthesis", Wiley, 1991 and 1999.

- 5 Any protective group used is optionally subsequently cleaved for example by hydrolysis in an aqueous solvent, e.g. in water, isopropanol/water, tetrahydrofuran/water or dioxane/water, in the presence of an acid such as trifluoroacetic acid, hydrochloric acid or sulphuric acid or in the presence of an alkali metal base such as lithium hydroxide, sodium hydroxide or potassium
- 10 hydroxide or by means of ether splitting, e.g. in the presence of iodotrimethylsilane, at temperatures between 0 and 100°C, preferably at temperatures between 10 and 50°C.

- A benzyl, methoxybenzyl or benzyloxycarbonyl group, however, is cleaved by
- 15 hydrogenolysis, for example, e.g. with hydrogen in the presence of a catalyst such as palladium/charcoal in a solvent such as methanol, ethanol, ethyl acetate, dimethylformamide, dimethylformamide/acetone or glacial acetic acid, optionally with the addition of an acid such as hydrochloric acid at temperatures between 0 and 50°C, but preferably at room temperature, and under a hydrogen
- 20 pressure of 1 to 7 bar, but preferably 1 to 5 bar.

- A methoxybenzyl group may also be cleaved in the presence of an oxidising agent such as cerium(IV)ammonium nitrate in a solvent such as methylene chloride, acetonitrile or acetonitrile/water at temperatures between 0 and 50°C,
- 25 but preferably at room temperature.

A methoxy group is conveniently cleaved in the presence of boron tribromide in a solvent such as methylene chloride at temperatures between -35 and -25°C.

- 5 A 2,4-dimethoxybenzyl group, however, is preferably cleaved in trifluoroacetic acid in the presence of anisole.

A *tert.*-butyl or *tert.*-butoxycarbonyl group is preferably cleaved by treatment with an acid such as trifluoroacetic acid or hydrochloric acid, optionally using a
10 solvent such as methylene chloride, dioxane or ether.

A phthalyl group is preferably cleaved in the presence of hydrazine or a primary amine such as methylamine, ethylamine or *n*-butylamine in a solvent such as methanol, ethanol, isopropanol, toluene/water or dioxane at temperatures
15 between 20 and 50°C.


An allyloxycarbonyl group is cleaved by treatment with a catalytic amount of tetrakis-(triphenylphosphine)-palladium(0), preferably in a solvent such as tetrahydrofuran and preferably in the presence of an excess of a base such as
20 morpholine or 1,3-dimedone at temperatures between 0 and 100°C, preferably at room temperature and under inert gas, or by treatment with a catalytic amount of tris-(triphenylphosphine)-rhodium(I)chloride in a solvent such as aqueous ethanol and optionally in the presence of a base such as 1,4-diazabicyclo[2,2,2]octane at temperatures between 20 and 70°C.

Furthermore, the compounds of formula (I) obtained may be converted into the salts thereof, particularly for pharmaceutical use into the physiologically acceptable salts with inorganic or organic acids. Acids which may be used for this purpose include for example hydrochloric acid, hydrobromic acid, sulphuric acid, methanesulphonic acid, phosphoric acid, fumaric acid, succinic acid, lactic acid, citric acid, tartaric acid or maleic acid.

Moreover, if the new compounds of formula (I) contain a carboxy group, they may subsequently, if desired, be converted into the salts thereof with inorganic or organic bases, particularly for pharmaceutical use into the physiologically acceptable salts thereof. Suitable bases for this purpose include for example sodium hydroxide, potassium hydroxide, cyclohexylamine, ethanolamine, diethanolamine, and triethanolamine.

According to the invention the optically pure derivatives of 3-Amino-tetrahydrofuran-3-carboxylic acid e.g. I, V, VI, VII or IX can be obtained for example in analogy to the following methods:

Method I: Chemical resolution with mandelic acid




Method II: Separation of the enantiomers via chiral chromatography

- 5 The separations can be performed on various DAICEL columns like AD-H, OD-H, AS-H, OJ-H, IA, IB and Kromasil DMB, TBB. Especially useful are DAICEL AD-H, OJ-H and IA columns.


Chiral chromatography can be used to

- 10 separate racemic 7 into its S- and R-enantiomer




-60-

or b) to separate racemic derivatives 11 like



or c) to separate racemic mixtures of I


5




10

Method III: Enzymatic resolution

-61-



Method IV: Alternative chemical resolution with mandelic acid



5

Accordingly, in a further embodiment, the present invention includes a method for preparing a compound of the formula (V) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring by enzymatic resolution of a racemic mixture of said compound of the formula (V), preferably with alcalase, wherein

10 R₂ and R₆ are as defined hereinabove, and wherein Q is a straight or

substituted C₁₋₁₂-alkyloxy group, allyloxy or substituted allyloxy group, a C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group, preferably Q is a straight or substituted C₁₋₄-alkyloxy.

- 5 In another embodiment, the present invention includes a compound of the formula (V) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring, wherein R₂ and R₆ are as defined hereinabove, and wherein Q is a hydroxy or substituted C₁₋₁₂-alkyloxy group, a halogen atom or a C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group, or substituted allyloxy group,
- 10 preferably Q is a substituted C₁₋₁₂-alkyloxy or substituted allyloxy group. In one aspect of the invention, the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (V) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring has the S-configuration. In another aspect, the compound of the general formula (V) in high optical purity at the carbon in
- 15 position 3 of the tetrahydrofuran ring is obtainable by the process described above.

In a further embodiment, the present invention includes a method for preparing a compound of the formula (VI) in high optical purity at the carbon in position 3

20 of the tetrahydrofuran ring by enzymatic resolution of a racemic mixture of said compound of the formula (VI), preferably with alcalase, wherein Q is straight or substituted C₁₋₁₂-alkyloxy group, or a C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group, or substituted allyloxy group, preferably Q is a substituted C₁₋₄-alkyloxy group, and PG is a hydrogen atom or a protective group for the amino function

25 as defined hereinabove.

In another embodiment, the present invention includes a compound of the formula (VI) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring, wherein Q is a hydroxy or straight or substituted

5 C₁₋₁₂-alkyloxy group, a halogen atom or a C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group, or substituted allyloxy, preferably Q is a straight or substituted

C₁₋₄-alkyloxy group, and PG is a hydrogen atom or a protective group for the amino function as defined hereinabove. In one aspect of the invention, the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the


10 general formula (VI) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring has the S-configuration. In another aspect, the compound of the general formula (VI) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring is obtainable by the process described above.

In another embodiment, the present invention includes a method for preparing a
15 compound of the formula (VII) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring by chemical resolution of a racemic mixture of said compound of the formula (VII) with a chiral acid, preferably with L-mandelic acid, wherein D and R₃ are as defined hereinabove.

In another embodiment, the present invention includes a compound of the
20 formula (VII) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring, wherein D and R₃ are as defined hereinabove. In one aspect of the invention, the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (VII) in high optical purity at the carbon in position 3 of the tetrahydrofuran ring has the S-configuration. In another
25 aspect, the compound of the general formula (VII) in high optical purity at the


carbon in position 3 of the tetrahydrofuran ring is obtainable by the process described above.

Especially useful 3-Amino-tetrahydrofuran-3-carboxylic acid precursors for the manufacturing of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) are the following compounds:



10 Determination of the absolute stereochemistry

The absolute configuration of the compounds can be determined by X-ray crystallography as is exemplified in the experimental part for (S)-3-[(5-chlorothiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid 7:



Description of the pharmaceutical useful properties of the compounds

- 5 As already mentioned hereinbefore, the compounds of general formula (I) and the tautomers, enantiomers, diastereomers and physiologically acceptable salts thereof have valuable pharmacological properties, particularly an antithrombotic activity which is preferably based on an effect on thrombin or factor Xa, for example on a thrombin-inhibiting or factor Xa-inhibiting activity, on a prolonging
- 10 effect on the aPTT time and on an inhibitory effect on related serine proteases such as e.g. urokinase, factor VIIa, factor IX, factor XI and factor XII.

The compounds listed in the Experimental Section were investigated for their effect on the inhibition of factor Xa as follows:

15

Method:

Enzyme-kinetic measurement with chromogenic substrate. The quantity of *p*-nitroaniline (pNA) released from the colourless chromogenic substrate by human factor Xa is determined photometrically at 405 nm. It is proportional to

20 the activity of the enzyme used. The inhibition of the enzyme activity by the test substance (in relation to the solvent control) is determined at various concentrations of test substance and from this the IC_{50} is calculated, as the concentration which inhibits the factor Xa used by 50 %.

Material:

Tris(hydroxymethyl)-aminomethane buffer (100 mMol) and sodium chloride (150 mMol), pH 8.0 plus 1 mg/ml Human Albumin Fraction V, protease-free

- 5 Factor Xa (Calbiochem), spec. activity: 217 IU/mg, final concentration: 7 IU/ml for each reaction mixture

Substrate S 2765 (Chromogenix), final concentration: 0.3 mM/l (1 KM) for each reaction mixture

10

Test substance: final concentration 100, 30, 10, 3, 1, 0.3, 0.1, 0.03, 0.01, 0.003, 0.001 μ Mol/l

Procedure:

- 15 10 μ l of a 23.5-times concentrated starting solution of the test substance or solvent (control), 175 μ l of TRIS/HSA buffer and 25 μ l of a 65.8 U/L Factor Xa working solution are incubated for 10 minutes at 37°C. After the addition of 25 μ l of S 2765 working solution (2.82 mMol/l) the sample is measured in a photometer (SpectraMax 250) at 405 nm for 600 seconds at 37°C.

20

Evaluation:

1. Determining the maximum increase (Δ OD/minutes) over 21 measuring points.

25

2. Determining the % inhibition based on the solvent control.
3. Plotting a dosage/activity curve (% inhibition vs substance concentration).
- 5 4. Determining the IC_{50} by interpolating the X-value (substance concentration) of the dosage/activity curve at Y = 50 % inhibition.

All the compounds tested had an IC_{50} value of less than 10 $\mu\text{mol/L}$. Surprisingly, for the compounds with S-configuration at the amino-tetrahydrofuran carboxylic acid amide moiety higher potency was observed in the FXa assay.

For example:

(S)-3-[(5-Bromo-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide showed an IC_{50} at an approximately 9-fold lower plasma concentration compared to the (R)-enantiomer,

(S)-3-[(5-Chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide showed an IC_{50} at an approximately 8-fold lower plasma concentration compared to the (R)-enantiomer,

(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

showed an IC_{50} at an approximately 10-fold lower plasma concentration compared to the corresponding (3R,5R)-diastereoisomer and (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

5 showed an IC_{50} at an approximately 7-fold lower plasma concentration compared to the corresponding (3R,5S)-diastereoisomer.

(S)-5-chloro-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-cyanimin-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide showed an

10 IC_{50} at an approximately 17-fold lower plasma concentration compared to the (R)-enantiomer,

(S)-5-bromo-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-oxo-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide showed an

15 IC_{50} at an approximately 16-fold lower plasma concentration compared to the (R)-enantiomer,

(S)-5-chloro-thiophene-2-carboxylic acid-N-{3-[4-(5-cyanimin-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide showed an IC_{50} at an

20 approximately 15-fold lower plasma concentration compared to the (R)-enantiomer and

(S)-5-bromo-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide showed an IC_{50} at an

approximately 15-fold lower plasma concentration compared to the (R)-enantiomer.

In view of their pharmacological properties the new compounds and the
5 physiologically acceptable salts thereof are suitable for the prevention and
treatment of venous and arterial thrombotic diseases, such as for example the
prevention and treatment of deep leg vein thrombosis, thrombophlebitis, for
preventing reocclusions after bypass operations or angioplasty (PT(C)A), and
occlusion in peripheral arterial diseases, and for preventing and treating
10 pulmonary embolism, disseminated intravascular coagulation and severe
sepsis, for preventing and treating DVT in patients with exacerbated COPD, for
treating ulcerative colitis, for preventing and treating coronary thrombosis, for
preventing thromboembolic events associated with atrial fibrillation, e.g., stroke
and the occlusion of shunts.

15

In addition, the compounds according to the invention are suitable for
antithrombotic support in thrombolytic treatment, such as for example with
alteplase, reteplase, tenecteplase, staphylokinase or streptokinase, for
preventing long-term restenosis after PT(C)A, for the prevention and treatment
20 of ischaemic events in patients with all forms of coronary heart disease, for
preventing metastasis and the growth of tumours and inflammatory processes,
e.g. in the treatment of pulmonary fibrosis or pulmonary arterial hypertension,,
for preventing and treating rheumatoid arthritis, for preventing and treating
fibrin-dependent tissue adhesions and/or the formation of scar tissue and for
25 promoting wound healing processes.

Compounds may also have utility as anticoagulant agents in connection with the preparation, storage fractionation or use of whole blood ; and in the coating of invasive devices such as prostheses, artificial valves and catheters in reducing
5 the risk of thrombus formation.

In view of their pharmacological properties the new compounds and the physiologically acceptable salts thereof are also suitable for the treatment of Alzheimer's and Parkinson's disease. One explanation for this arises for
10 example from the following findings, from which it can be concluded that thrombin inhibitors or factor Xa inhibitors, by inhibiting thrombin formation or thrombin activity, may be valuable drugs for treating Alzheimer's and Parkinson's disease. Clinical and experimental studies indicate that neurotoxic mechanisms, for example the inflammation which is associated with the
15 activation of proteases of the clotting cascade, are involved in the dying of neurones following brain injury. Various studies point to the involvement of thrombin in neurodegenerative processes, for example following a stroke, repeated bypass operations or traumatic brain injury. An increased thrombin activity has been demonstrated some days after peripheral nerve damage, for
20 example. It has also been shown that thrombin causes a neurite retraction, as well as glia proliferation, and apoptosis in primary cultures of neurones and neuroblastoma cells (for a summary see: *Neurobiol. Aging* 2004, 25(6), 783-793). Moreover, various *in vitro* studies on the brains of patients with Alzheimer's disease indicated that thrombin plays a role in the pathogenesis of
25 this disease (*Neurosci. Lett.* 1992, 146, 152-54). A concentration of immune-

reactive thrombin has been detected in neurite plaques in the brains of Alzheimer's patients. It has been demonstrated *in vitro* that thrombin also plays a part in the regulation and stimulation of the production of the "Amyloid Precursor Protein" (APP) as well as in the cleaving of the APP into fragments
5 which can be detected in the brains of Alzheimer's patients. Moreover, it has been demonstrated that the thrombin-induced microglial activation leads *in vivo* to the degeneration of nigral dopaminergic neurones. These findings lead one to conclude that microglial activation, triggered by endogenous substance(s) such as thrombin, for example, are involved in the neuropathological process of
10 the cell death of dopaminergic neurones of the kind which occurs in patients with Parkinson's disease (*J. Neurosci.* 2003, 23, 5877-86).

The new compounds and the physiologically acceptable salts thereof are also suitable for the prophylaxis and treatment of arterial vascular diseases as
15 combination therapy with lipid lowering agents such as HMG-CoA reductase inhibitors; and vasodilators, in particular ACE-inhibitors, angiotensin II antagonists, renin inhibitors, β -receptor antagonists, α -receptor antagonists, diuretics, Ca-channel blockers, or stimulators of soluble guanylate cyclase.

20 By increasing the antithrombotic efficacy, the new compounds and the physiologically acceptable salts thereof are also suitable in combination therapy with other anticoagulants e.g. unfractionated heparin, low molecular heparins or fondaparinux, or direct thrombin inhibitors e.g. recombinant hirudin or small molecule synthetic inhibitors.

Similarly, the compounds and the physiologically acceptable salts thereof are also suitable for the prophylaxis and treatment of arterial vascular diseases as combination therapy with platelet aggregation inhibitors. e.g., aspirin, clopidogrel or a glycoprotein-IIb/IIIa antagonist or a thrombin receptor
5 antagonist.

The dosage required to achieve such an effect is appropriately 0.001 to 3 mg/kg body weight, preferably 0.003 to 1.0 mg/kg body weight by intravenous route, and 0.003 to 30 mg/kg body weight, preferably 0.01 to 10 mg/kg body weight by
10 oral route, in each case administered 1 to 4 times a day.

For this purpose, the compounds of formula (I) prepared according to the invention may be formulated, optionally together with other active substances, with one or more inert conventional carriers and/or diluents, e.g. with corn
15 starch, lactose, glucose, microcrystalline cellulose, magnesium stearate, polyvinylpyrrolidone, citric acid, tartaric acid, water, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol, cetylstearyl alcohol, carboxymethylcellulose or fatty substances such as hard fat or suitable mixtures thereof, to produce conventional galenic preparations such
20 as plain or coated tablets, capsules, powders, suspensions or suppositories.

The new compounds and the physiologically acceptable salts thereof may be used therapeutically in conjunction with acetylsalicylic acid, with inhibitors of platelet aggregation such as fibrinogen receptor antagonists (e.g. abciximab,
25 eptifibatide, tirofiban, roxifiban), with physiological activators and inhibitors of

- the clotting system and the recombinant analogues thereof (e.g. Protein C, TFPI, antithrombin), with inhibitors of ADP-induced aggregation (e.g. clopidogrel, ticlopidine), with P₂T receptor antagonists (e.g. cangrelor) or with combined thromboxane receptor antagonists/synthetase inhibitors (e.g. terbogrel) or with a thrombin receptor antagonist (e.g. SCH-530348).

Experimental section

- The Examples that follow are intended to illustrate the invention, without restricting its scope.

As a rule, melting points and/or IR, UV, ¹H-NMR and/or mass spectra have been obtained for the compounds prepared. Unless otherwise stated, R_f values were determined using ready-made silica gel 60 F₂₅₄ TLC plates (E. Merck, Darmstadt, Item no. 1.05714) without chamber saturation. The R_f values given under the heading Alox were determined using ready-made aluminium oxide 60 F₂₅₄ TLC plates (E. Merck, Darmstadt, Item no. 1.05713) without chamber saturation. The R_f values given under the heading Reversed-phase-8 (RP-8) were determined using ready-made RP-8 F_{254s} TLC plates (E. Merck, Darmstadt, Item no. 1.15684) without chamber saturation. The ratios given for the eluants refer to units by volume of the solvents in question. For chromatographic purification silica gel made by Messrs Millipore (MATREXTM, 35-70 μm) was used. Unless more detailed information is provided as to the configuration,

it is not clear whether the products are pure stereoisomers or mixtures of enantiomers and diastereomers.


The following abbreviations are used in the test descriptions:

5	BOC	tert.-butoxycarbonyl
	CDI	1,1'-carbonyldiimidazole
	DIPEA	<i>N</i> -ethyl-diisopropylamine
	DMAP	4-dimethylzaminopyridine
	DMF	<i>N,N</i> -dimethylformamide
10	DMSO	dimethylsulfoxide
	D-DTTA	(+)- <i>O,O'</i> -di- <i>p</i> -toluoyl-D-tartaric acid
	EDC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
	EtOAc	ethylacetate
	sat.	saturated
15	h	hour(s)
	HATU	<i>O</i> -(7-azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium-hexafluorophosphate
	HOBT	1-hydroxy-benzotriazole
	IPA	isopropanol
20	LiHMDS	lithium hexamethyldisilazide
	NaHMDS	sodium hexamethyldisilazide
	i. vac.	<i>in vacuo</i>
	conc.	concentrated
	min	minute(s)
25	MsCl	methanesulfonyl chloride

	MTBE	methyl-tert.butylether
	Me-THF	2-methyl-tetrahydrofuran
	NMM	<i>N</i> -methyl-morpholine
	Pd ₂ dba ₃	bis(dibenzylideneacetone)palladium (0)
5	n-PrOH	1-propanol
	[Rh(COD)Cl] ₂	chloro(1,5-cyclooctadiene)rhodium
	R _f	retention factor
	R _t	retention time
	rt	room temperature
10	TBTU	O-(benzotriazol-1-yl)- <i>N,N,N,N</i> -tetramethyluronium tetrafluoroborate
	TEA	triethylamine
	TFA	trifluoroacetic acid
	TFFA	trifluoroacetic anhydride
15	THF	tetrahydrofuran
	TsCl	para-toluene sulfonic acid chloride
	TsOH	para-toluene sulfonic acid.
	Walphos	1-[2- (2'-diphenylphosphinophenyl)ferrocenyl]ethyldiphenylphosphine

20

The term "thiophen-2-yl" or "thien-2-yl" denotes the group shown in the box:



The HPLC-MS data were obtained under the following conditions:

Method A

The mobile phase used was:

5 A: water with 0.15% HCOOH

B: acetonitrile

	time in min	%A	%B	flow rate in ml/min
	0.00	95	5	1.00
10	2.00	95	5	1.00
	9.00	2	98	1.00

The stationary phase used was Zorbax StableBond C18 column; 8 μ m ; 50 mm
x 90 mm

15

A) Determination of the absolute stereochemistry of compound 7:

The crystal structure of compound **7** was determined by direct methods. The absolute configuration was determined by refinement of the Flack parameter (Flack H D (1983), Acta Cryst. A39, 876-881).

20

Crystal data

$C_{10}H_{10}ClNO_4S$ $V = 1100.7 (4) \text{ \AA}^3$

$M_r = 275.70$ $Z = 4$

$D_x = 1.664 \text{ Mg m}^{-3}$

25 $a = 7.8200 (16) \text{ \AA}$ $\text{Cu K}\alpha$

- $b = 8.7700(18) \text{ \AA}$ $\mu = 4.91 \text{ mm}^{-1}$
 $c = 16.050(3) \text{ \AA}$ $T = 100(2) \text{ K}$
 $\alpha = 90^\circ$ Prism, colourless
 $\beta = 90^\circ$ $0.1 \times 0.1 \times 0.2 \text{ mm}$
5 $\gamma = 90^\circ$

Data collection

Saturn 944 CCD mounted on AFC11K diffractometer

1627 independent reflections

- 10 oscillation scans 1610 reflections with $I > 2\sigma(I)$

Absorption correction: empirical (using intensity measurements)

$$R_{\text{int}} = 0.052$$

3840 measured reflections $\theta_{\text{max}} = 63.7^\circ$

15 Refinement

Refinement on F^2 $w = 1/[\sigma^2(F_o^2) + (0.0388P)^2 + 0.2397P]$

where $P = (F_o^2 + 2F_c^2)/3$

$$R[F^2 > 2\sigma(F^2)] = 0.030 \quad (\Delta/\sigma)_{\text{max}} = 0.001$$

$$wR(F^2) = 0.072 \quad \Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$$

- 20 $S = 1.08$ $\Delta\rho_{\text{min}} = -0.27 \text{ e \AA}^{-3}$

1627 reflections Extinction correction: none

158 parameters

H atoms treated by a mixture of independent and constrained refinement

Flack parameter: 0.009 (17)

Data collection: Saturn 944 CCD mounted on AFC11K/RU200 rotating anode generator; cell refinement: D*trek; data reduction: D*trek; program(s) used to solve structure: *SHELXS97* program(s) used to refine structure: *SHELXL97* molecular graphics: *XP*.

- 5 The configuration of the chiral carbon atom is S. The structure is shown in Figure 4.

B) Preparation of 3-amino-tetrahydro-furan-3-carboxylic acid derived intermediates:

10

Preparation of rac-3-amino-tetrahydro-furan-3-carboxylic acid phenethyl ester x para-toluenesulfonic acid (2):

- To a solution of 3-amino-tetrahydrofuran-3-carboxylic acid **1** (100 g, 163 mmol) in 2-phenylethanol (456 mL) and toluene (400 mL) was added TsOH.H₂O (174 g, 916 mmol) under nitrogen and the mixture was heated to reflux while the water was collected via a Dean-Strak trap. At the completion of the reaction, the reaction mixture was concentrated to 1/3 of its original volume and cooled to ambient temperature. MTBE (1000 ml) was added and the mixture was stirred for 1 h. The slurry was filtered and the wet cake was washed twice with MTBE and dried to give the desired product **2** (284 g) in 92% yield.
- 15
20

¹HNMR (DMSO-D₆, 400 MHz) δ 8.60 (br s, 3H), 7.47 (d, J = 8.0 Hz, 2 H), 7.20-7.50 (m, 5H), 7.11 (d, J = 7.9 Hz, 2 H), 4.44 (dt, J = 6.6, 1.1 Hz, 2H), 3.97 (m, 1H), 3.84 (m, 3H), 2.97 (t, J = 6.6 Hz, 2H), 2.32 (m, 1H), 2.30 (s, 3H), 2.09 (m, 1H)

25

Preparation of rac-3-amino-tetrahydro-furan-3-carboxylic acid phenethyl ester (3):

To a 1.5 L of 5% NaHCO₃ aqueous solution was added 2 (283 g) and 1.5 L of ethyl acetate and the resulting mixture was stirred at ambient temperature for 30 min. The organic phase was removed and the aqueous phase was extracted once with 1.5 L of ethyl acetate. The combined organic phases were dried and concentrated to dryness to yield 157 g of 3 in 97% yield.

Preparation of (S)-3-amino-tetrahydro-furan-3-carboxylic acid phenethyl ester x L-mandelic acid (4):

A mixture of 3 (157 g), MTBE (0.94 L), MeCN (0.94 L), water (0.094 L), and L-mandelic acid (152.2 g) was heated to 50-52 °C for 30 min. The mixture was cooled to 41-43 °C in an hour and 1.29 g of seed was added. After stirred for 30 min, the mixture was cooled to 0 °C and stirred for 3 h. The slurry was filtered and the wet cake was washed twice with MTBE (75 mL) and dried to give crude 4 (113.5 g) in 43.9% yield and 87% de.

The above salt was recrystallized from a mixture of MTBE (908 mL), CH₃CN (908 mL), and water (54.5 mL) to give 102.0 g of product in 89.9% yield and 98.2% de. One more crystallization from MTBE (816 mL), CH₃CN (816 mL), and water (48.9 mL) gave 94.3 g of 4 in 92.5% yield (35.0% overall yield) and 99.7% de. The enantiomerically purity was analyzed by chiral HPLC with chiralpak AD-H column, 250 cm x 4.6 mm; 5 µm; 2 mL/min, 220 nm; 95% heptane/5% IPA; *r*_t = 9.3 min ((S)-isomer); *r*_t = 10.1 min, ((R)-isomer).

Preparation of (S)-3-Amino-tetrahydro-furan-3-carboxylic acid phenethyl ester 5:

A suspension of 4 (94.3 g) in 750 mL of EtOAc was added 750 mL of 5% NaHCO₃ solution and the mixture was stirred for 30 min. After the organic phase was removed, the aqueous phase was extracted once with EtOAc (750 mL). The combined organic phases were dried and concentrated to dryness to give 57.6 g of 5 in >99% yield.

Preparation of (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid phenethylester (6):


A mixture of 5 (20 g, 85.1 mmol), 5-chloro-thiophene-2-carboxylic acid (14.5 g, 89.4 mmol), HOBT.H₂O (13.8 g, 102 mmol), EDC.HCl (19.6 g, 102.1 mmol), and DMF (200 mL) was cooled to 10-15 °C. Then TEA (17.8 mL) was added over 5 min and the mixture was warmed to ambient temperature and stirred for 2 h. EtOAc (200 mL) and water (200 mL) was added to the mixture and stirred for 20 min. The organic phase was removed and the aqueous phase was extracted once with EtOAc (200 mL). The combined organic phases were washed with 200 mL of 5% NaCl solution and dried and concentrated to give 6 (29 g) in 90% yield.

¹HNMR (CDCl₃, 400 MHz) δ 7.15-7.30 (m, 6H), 6.89 (br s, 1H), 4.42 (t, *J* = 6.9 Hz, 2H), 3.90-4.12 (m, 4 H), 2.96 (t, *J* = 6.9 Hz, 2H), 2.53 (m, 1H), 2.29 (m, 1H).

Preparation of (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid (7):

A solution of **6** (40 g) in MeOH (200 mL) was cooled to 10-15 °C, followed by addition of 1N NaOH (200 mL) over 5 min. The mixture was warmed to ambient temperature and stirred for 30 min. The mixture was concentrated to remove most of MeOH and MTBE (200 mL) was added and stirred the mixture for 10 min. The organic phase was removed and aqueous phase was washed once with 200 mL of MTBE. The aqueous layer was cooled to 0 °C and 3N HCl solution was added to a pH = 2-3. Then 200 mL of Me-THF and 18 g of NaCl was added and the mixture was stirred for 10 min. The aqueous phase was extracted once with 100 mL of Me-THF. The combined organic phase was concentrated and 100 mL of heptane was added. The slurry was filtered and the wet cake was washed with heptane (50 mL x 2) and dried to give **7** (23.3 g) in 99% yield.

Racemic compound **7** could also be prepared following the procedure described for example 1e:



For separation of the racemic compound into its respective enantiomers, a conventional analytical HPLC system with DAICEL AD-H 250mm x 4,6mm chiral column has been used, eluting with 0,1% Acetic Acid in Hexane(80%) / EtOH(20%) as liquid phase. At a flow rate of 1ml/min, retention times for the enantiomers are 9,2 min and 12,5 min.

Alternatively, the separation of this racemate can be achieved on HPLC with DAICEL OJ-H chiral column, eluting with 0,1% Acetic Acid in Hexane(80%) / EtOH(20%). At a flow rate of 1ml/min, retention times for the enantiomers are 6,05 min and 8,07 min respectively.

5

Preparation of (S)-2-(5-chloro-thiophene-2-yl)-3,7-dioxa-1-aza-spiro[4.4]non-1-en-4-one (8)

50 mg (0.18 mmol) (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid **7** were stirred for 2h in 45 ml acetic anhydride at 85°C.

10 The reaction mixture was evaporated down i. vac., the residue was taken up twice each in toluene and dichloromethane and completely concentrated by evaporation. The crude title compound was reacted directly without any further purification.

15 **Preparation of benzyl 3-tert.-butoxycarbonylamino-tetrahydro-furan-3-carboxylate (9):**

A mixture of 7.0 g 3-tert.-butoxycarbonylamino-tetrahydro-furan-3-carboxylic acid, 80 ml DMF and 4.6 g K₂CO₃ was stirred for 15 min, then 3.6 ml benzylbromide was added dropwise and it was stirred for 3 days. The mixture
20 was filtered and the liquid phase was concentrated i.vac.. The residue was diluted with CH₂Cl₂, and the mixture was washed with water and saturated NaCl-solution. The organic phase was dried with Na₂SO₄ and concentrated i.vac. to yield the title compound in 58% yield.

25 **Preparation of benzyl 3-amino-tetrahydro-furan-3-carboxylate (10):**

A mixture of 5.2 g benzyl 3-tert.-butoxycarbonylamino-tetrahydro-furan-3-carboxylate, 200 ml CH_2Cl_2 and 20 ml TFA was stirred for 3 h and concentrated i. vac. to yield the title compound in quantitative yield.

5 **Preparation of 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid benzylester 11 and separation of the enantiomers (S-12 and R-12):**

1.59 g (9.8 mmol) 5-chloro-thiophene-2-carboxylic acid is dissolved in 30 ml DMF and stirred with 3.61 g (10.7 mmol) benzyl 3-amino-tetrahydro-furan-3-carboxylate and 3.46 g (10.8 mmol) TBTU and 4.3 ml (39 mmol) NMM at room temperature for 20 h. Then the mixture is evaporated down and purified by chromatography on silica gel (eluant: dichloromethane/ethanol 100:0 to 94:6).

Yield: quantitative

R_f value: 0.59 (silica gel; dichloromethane/ethanol = 9:1)

15 $\text{C}_{17}\text{H}_{16}\text{ClNO}_4\text{S}$ (365.83)

Mass spectrum: $(\text{M}+\text{H})^+$ = 366/368 (chlorine isotopes)


For separation of the racemic mixture into its respective enantiomers (**S-12 and R-12**), a conventional HPLC system with DAICEL IA, 250mm x 4,6mm chiral column has been used, eluting with EtOH(2%) / CHCl_3 (20%) / Hexane (68%). At a flow rate of 1ml/min, retention times for the enantiomers are 12,3min and 20,7min.

Alternatively, the separation of this racemate can be achieved on supercritical fluid chromatography with DAICEL IA chiral column, eluting with EtOH(15%) /

CHCl_3 (10%) / supercritical CO_2 (75%). At a flow rate of 70ml/min, retention times for the enantiomers are 3,57min and 5,13 min respectively.

Preparation of (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid (7):


To a solution of (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid benzylester **S-12** (5.8 g) in ethanol (100 mL) was added 1N NaOH (63 mL). The mixture stirred for 90 min and the mixture was concentrated i.vac.. Cold 1 N HCl was added and the mixture was stirred overnight. The precipitate was filtered and dried to give the title compound in 99% yield.



Preparation of rac-3-Amino-tetrahydro-furan-3-carboxylic acid isobutyl ester hydrogen chloride (13):

To a mixture of amino acid **1** (190.5 g, 1.45 mol) in 2-methyl-1-propanol (1.9 L) at 0 °C was added dropwise thionyl chloride (211.6 mL, 2.0 equiv.) over 20 min. The mixture was heated to 89 °C, and kept at 89 °C for 0.5 h, then heated to 108 °C and stirred at this temperature for 1.5 h. The mixture was cooled to rt and concentrated to remove most of 2-methyl-1-propanol. The residue was treated with *t*-butyl methyl ether (1 L) to form a suspension and stirred at rt for 0.5 h. The mixture was filtered and the cake was washed twice with *t*-butyl


methyl ether (0.2 L x 2) to provide the desired salt **13** (287 g, 1.28 mol, 88%) as a white solid.



5


Preparation of rac-3-[(5-Chloro-thiophene-2-carbonyl)-amino]-tetrahydrofuran-3-carboxylic acid isobutyl ester (14**):**

To a mixture of salt **13** (287 g, 1.28 mol), 5-chlorothiophene-2-carboxylic acid
10 (219 g, 1.35 mol, 1.05 equiv), HOBT.H₂O (208 g, 1.54 mol, 1.2 equiv), and
EDC.HCl (295 g, 1.54 mol, 1.2 equiv) in anhydrous DMF at 0 °C was added
triethylamine over 10 min. The mixture was allowed to warm to rt and stirred for
3 h. EtOAc (2 L) and water (2 L) was added and the aqueous layer was
removed. The EtOAc layer was further washed with water (2 L) and 5% NaCl
15 solution (2 L) and concentrated. The residue was passed through a short silica
gel plug (eluent: Hexane/EtOAc 4:1 → 1:1) to give the desired amide **14** (335 g,
1.01 mol, 82%) as an oil. **14**: ¹HNMR (CDCl₃, 400 MHz) δ = 7.47 (s, 1H), 7.40 (d,
J = 4.0 Hz, 1H), 6.86 (d, *J* = 4.0 Hz, 1H), 4.26 (d, *J* = 9.6 Hz, 1H), 3.90-4.08 (m,
4H), 2.60 (m, 1H), 2.38 (m, 1H), 1.96 (m, 1H), 0.90 (d, 4.04 (m, *J* = 6.8 Hz, 6H);
20 ¹³CNMR (CDCl₃, 100 MHz) δ = 172.1, 163.5, 161.6, 136.7, 136.0, 128.3, 127.1,
76.2, 72.0, 67.9, 66.0, 60.5, 37.4, 27.7, 19.0; ESI-MS: *m/z* 332 [*M*+H]⁺, 685
[2*M*+Na]⁺.



Preparation of (S)-3-[(5-Chloro-thiophene-2-carbonyl)-amino]-tetrahydrofuran-3-carboxylic acid isobutyl ester (15):

- 5 To a 0.1 M phosphate buffer solution (pH = 6.7, NaH₂PO₄·H₂O: 100.5 g, Na₂HPO₄: 80.8 g, H₂O: 6.55 L) at rt was added Alcalase (564 mL, > 0.75U/mL, 2 U/mol) and racemic ester **14** (70 g, 0.211 mol) in acetone (3.275 L). The pH value of the resulting mixture was about 7.30. The mixture was stirred at rt until the ee of the remaining ester reached 93% by chiral HPLC (about 30 h). EtOAc
- 10 (2 L) was added and the aqueous layer was further extracted with EtOAc (1 L). The combined EtOAc layer was concentrated, passed through a silica gel plug (hexane: EtOAc = 1:1) to give the optical enriched ester **15** (28 g, 25%, 93% ee).




15

Preparation of (S)-3-[(5-Chloro-thiophene-2-carbonyl)-amino]-tetrahydrofuran-3-carboxylic acid [(S)-7]:

- The optically enriched ester **15** (43.5 g, 0.131 mol) was dissolved in MeOH (400 mL). To the solution at 0 °C was added 1 N NaOH solution (400 mL) and the
- 20


resulting mixture was stirred at rt for 0.5 h. The mixture was then concentrated to remove most of methanol. The resulting aqueous solution was washed once with *t*-butyl methyl ether (200 mL), and then neutralized with 3 N HCl at 0 °C to pH = 1-2. The mixture was saturated by adding solid NaCl and extracted with Me-THF (500 mL). The Me-THF layer was dried over Na₂SO₄ and concentrated to give crude acid (S)-7 as a white solid. The solid was then recrystallized three times from water (40 mL hot water/ g) to give desired acid in 99.5% ee (21 g, 48%). (S)-7: ¹HNMR (DMSO-D₆, 400 MHz) δ = 12.75 (s, 1H), 8.98 (s, 1H), 7.77 (d, *J* = 4.0 Hz, 1H), 7.21 (d, *J* = 4.0 Hz, 1H), 4.12 (d, *J* = 9.3 Hz, 1H), 3.92 (d, *J* = 9.3 Hz, 1H), 3.84 (t, *J* = 7.0 Hz, 2H), 2.35 (m, 2H).



Preparation of rac-3-Amino-tetrahydro-furan-3-carboxylic acid butyl ester

(16):


To a mixture of amino acid 1 (10 g, 76 mmol) and *n*-butanol (100 mL) at 0 °C was added SOCl₂ (18.2 g, 153 mmol, 2 equiv.) over 10 min. The mixture was heated to 110 °C over 10 min and stirred at the temperature for 4 h. After cooled to rt, the mixture was concentrated to remove *n*-butanol and Me-THF (100 mL) was added. To the mixture was carefully added sat. NaHCO₃ (100 mL) and the resulting mixture was stirred at rt for 30 min. The organic layer was separated and the aqueous layer was washed with Me-THF (50 mL). The combined organic phase was washed with 3% NaCl solution (50 mL) and concentrated to give the crude butyl ester 16 as an oil (14.3 g, 99.4%).



Preparation of (S)-3-Amino-tetrahydro-furan-3-carboxylic acid butyl ester

5 (S)-mandelic acid (17):


Butyl ester (16, 10 g, 0.053 mol) and 50 mL of MeCN was dissolved in a 250 mL reactor. (S)-Mandelic acid (4.88 g, 0.032 mol, 0.6 equiv) was added followed by addition of another 60 mL of MeCN. The mixture was heated to 70 °C and the clear solution was then cooled to 20 °C over 12 h and kept at 20 °C
10 for 1 h. The slurry was filtered and the mother liquid was recharged back to the reactor for wash. The cake was washed with MTBE (20 mL x 2) and dried under vacuum to give 7.8 g of salt with 90% de (43%). 7.5 g of the salt (90% de) was charged into a 250 mL reactor followed by MeCN (90 mL) and H₂O (0.9 mL). The mixture was heated to 70 °C and the clear solution was then cooled to 0 °C
15 over 12 h and kept at 0 °C for 1 h. The slurry was filtered and the wet cake was washed with cold MTBE (20 mL x 2, 0-5 °C) and dried under vacuum to give 17 as white crystal (6.5 g, 99.0% ee, 87%).



Preparation of (S)-3-tert-Butoxycarbonylamino-tetrahydro-furan-3-carboxylic acid butyl ester (18):

Salt 17 (33.9 g, 100 mmol) was dissolved in Me-THF (150 mL) and H₂O (150 mL). To the mixture at rt was added solid NaHCO₃ (12.6 g, 150 mmol) portion wise and the mixture was stirred at rt until no gas was generated. The organic phases were separated and the aqueous layer was washed with Me-THF (50 mL). The combined Me-THF was washed with 3% NaCl (100 mL) and concentrated to 100 mL. To the mixture was added Boc₂O (21.8 g, 10 mmol) in one portion. The mixture was heated to 50 °C and stirred at the temperature for 2 h. After cooled to rt, the mixture was washed with sat. NaHCO₃ (100 mL) and brine (100 mL) and concentrated to give crude 18 as an oil (27.3 g, 95%),


10



Preparation of [(S)-3-(3-Methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-ylcarbamoyl)-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester (19):

To a solution of 18 (6.1 g, 34.8 mmol) and 3-Methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-ylamine (10 g, 34.8 mmol) in THF (100 mL) at -10 °C was added dropwise LHMDS (87 mL, 1.0 M in THF, 2.5 equiv.) over 10 min. The mixture was warmed up to 0 °C over 1 h and monitored by HPLC analysis. Sat. NH₄Cl (50 mL) was added at ~0 °C to quench the reaction and the mixture was further diluted with EtOAc (100 mL). The aqueous layer was separated and extracted once with EtOAc (50 mL). The combined organic phase was washed with brine (50 mL) and concentrated. The light yellow solid was recrystallized

from IPA (85 mL) at 0 °C to give 19 as a white solid (11.5 g, 85%, 98.3 A% purity).




5

Preparation of (S)-3-Amino-tetrahydro-furan-3-carboxylic acid (3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-amide hydrogen chloride (20):


To a solution of 19 (2.00 g) in MeOH (10 mL) was added HCl /MeOH (~9N, 10 mL) at ambient temperature. The resulting clear solution was stirred at ambient temperature for 1 h and then concentrated to give a white solid. MeOH (5 mL) and MTBE (30 mL) were added and the slurry was stirred at room temp for 0.5 h. The slurry was filtered, washed with MTBE (5 mL) and dried under reduced pressure to give salt 20 as a white solid (1.8 g, 96%, >99% purity).

15



Preparation of (S)-3-amino-tetrahydrofuran-3-carboxylic acid n-butyl ester via chemical resolution with L-mandelic acid

20



- 3-Aminotetrahydro-furan-3-carboxylic acid (75 g, 0.572 mol) and 750 mL *n*-
- 5 butanol are combined into a 2 L reaction flask. The resulting suspension is cooled while stirring to about $3\text{ }^\circ\text{C}$. SOCl_2 (131.6 g, 1.144 mol) is added cautiously over 25 min to the stirred suspension (exothermic). After addition is complete, the resulting mixture is heated to $110\text{ }^\circ\text{C}$ over 1h. HCl gas is generated during the course of heating and the generation of gas is vigorous
- 10 when the temperature approaches over $100\text{ }^\circ\text{C}$. The reaction mixture is stirred $110\text{ }^\circ\text{C}$ for 4 h.
- When the reaction is complete, as indicated by LC/MS, the mixture is cooled to $\sim 70\text{ }^\circ\text{C}$. The reaction mixture is distilled to a minimum volume ($\sim 200\text{ mL}$) while controlling the temperature at $70\text{--}75\text{ }^\circ\text{C}$ at $95\text{--}100\text{ mbar}$. About 460 g
- 15 solvents/ SOCl_2 is collected. The concentrated reaction mixture is cooled to room temperature and 2-methyl-tetrahydrofuran (Me-THF) (600 mL) is added. 8% NaHCO_3 (750 mL) is added cautiously, in portions, while vigorously stirring. CO_2 is generated over the course of the addition. The resulting mixture is stirred for 15 min and then the phases are allowed to stand for 15 min. The
- 20 phases are separated and then the aqueous phase is stirred with an additional 400 mL of Me-THF 15 min. After standing for 15 min, the organic layer is separated and combined with the first organic extract. 3% NaCl (375 mL) is added to the combined organic extracts and the mixture is stirred the mixture for

15 min and then allowed to stand for 15 min. The aqueous layer (~1055g) is removed. The organic solution is distilled to a minimum volume (~ 250 mL). MeCN (1000 mL) is added in one portion and the resulting solution is distilled to a minimum volume (~ 250 mL). About 800g solvents were collected. NMR
5 assay of the concentrated product indicated 94.0 g of the desired racemic *n*-butyl ester was obtained (88%).

Chemical Resolution

MeCN (1125 mL) is added to the concentrated racemic *n*-butyl ester. A GC
10 analysis should be conducted at this stage and the content of Me-THF should be controlled to be < 5%. If the amount of Me-THF is >5%, the distillation of solvent and addition of 1125 mL of MeCN should be repeated.

L-Mandelic Acid (60.9 g, 0.4 mol) is added to the ester solution in one portion while stirring resulting in the formation of white solids. The mixture is heated to
15 70 °C and held at that temperature for 30 min, resulting in a clear solution. The solution is cooled to 20 °C over 12 h and then held at 20 °C for 1 h. The resulting slurry is filtered. The mother liquid is added back to the reaction flask to wash the flask. The filter cake is washed twice with MTBE (200 mL X 2) and the resulting solid is dried at ~50 °C under house vacuum for 3 h. (S)-3-Amino-
20 tetrahydrofuran-3-carboxylic acid *n*-butyl ester L-mandelic acid salt (82.0 g, 42%) is obtained as a white solid in 86 % de.


Enrichment Procedure

The 86% de ester (82 g) is placed in a 2 L flask, under N₂ and suspended in 980
25 mL of CH₃CN, and 19.7 mL of water. The mixture is heated to about 70 °C to

dissolve the salt, resulting in a clear solution. The solution is held at 70 °C for 30 min and then cooled 20 °C to 23 °C over 12 h. The resulting slurry is filtered and the wet filter cake is washed twice with 150 mL of MTBE. The product is dried at 50 °C under reduced pressure resulting in 66 g (81% yield) of (S)-3-amino-tetrahydrofuran-3-carboxylic acid *n*-butyl ester L-mandelic acid salt in 99% de.

Preparation of (S)-3-amino-tetrahydrofuran-3-carboxylic acid *n*-butyl ester via chemical resolution with L-mandelic acid

10



To a solution of dihydrofuran-3-one (10.0 g, 116.2 mmol) in MeOH (50 mL) was added a solution of 7 N NH_3/MeOH (33 mL) at ambient temperature. The mixture was cooled to 0 °C and 7.66 g of AcOH was added over 5 min while maintaining the temperature below 25 °C. The mixture was stirred at ambient temperature for 10 min and 5.64 g of NaCN was added in one portion. The mixture was heated to 50 °C, stirred for 2 h and then concentrated to remove the MeOH and ammonia. EtOAc (40 mL) was added to the concentrated mixture and then stirred for 15 min. The slurry was filtered and the wet cake was washed twice with 20 mL EtOAc. The combined filtrates, containing the desired 3-amino-tetrahydrofuran-3-carbonitrile, were concentrated and to the residue was added 20 mL of *n*-BuOH. The resulting mixture was cooled to 0 °C

and then 100 mL of 5.5 N HCl in *n*-BuOH was added. The resulting mixture was stirred at ambient temperature for 12 h.

The mixture was cooled to 0 °C and 20 mL water was added. The mixture was then concentrated to remove most of the *n*-BuOH. Saturated NaHCO₃ was

5 added to the residue to pH>7 and the resulting mixture was extracted twice with 200 mL of 2-methyltetrahydrofuran. The organic extracts were washed once with brine (80 mL) and then concentrated to give 18.3 g of the title compound (85% overall yield). ¹H NMR (CDCl₃, ppm): 2.02 (s, 2H), 2.09-2.15 (m, 1H), 2.43-2.50 (m, 1H), 3.75-3.78 (d, J=9.04, 1H), 3.98-4.08 (m, 3H). ¹³C NMR
10 (CDCl₃, ppm) 40.80, 54.03, 67.51, 78.72, 122.63.

Chemical Resolution

MeCN (1125 mL) is added to the concentrated racemic *n*-butyl ester. A GC analysis should be conducted at this stage and the content of Me-THF should
15 be controlled to be < 5%. If the amount of Me-THF is >5%, the distillation of solvent and addition of 1125 mL of MeCN should be repeated.

L-Mandelic Acid (60.9 g, 0.4 mol) is added to the ester solution in one portion while stirring resulting in the formation of white solids. The mixture is heated to 70 °C and held at that temperature for 30 min, resulting in a clear solution. The
20 solution is cooled to 20 °C over 12 h and then held at 20 °C for 1 h. The resulting slurry is filtered. The mother liquid is added back to the reaction flask to wash the flask. The filter cake is washed twice with MTBE (200 mL X 2) and the resulting solid is dried at ~50 °C under house vacuum for 3 h. (S)-3-Amino-tetrahydrofuran-3-carboxylic acid *n*-butyl ester L-mandelic acid salt (82.0 g,
25 42%) is obtained as a white solid in 86 % de.

Enrichment Procedure

The 86% de ester (82 g) is placed in a 2 L flask, under N₂ and suspended in 980 mL of CH₃CN, and 19.7 mL of water. The mixture is heated to about 70 °C to
 5 dissolve the salt, resulting in a clear solution. The solution is held at 70 °C for 30 min and then cooled 20 °C to 23 °C over 12 h. The resulting slurry is filtered and the wet filter cake is washed twice with 150 mL of MTBE. The product is dried at 50 °C under reduced pressure resulting in 66 g (81% yield) of (S)-3-amino-tetrahydrofuran-3-carboxylic acid *n*-butyl ester L-mandelic acid salt in
 10 99% de.


Alternatively, various alcohols (ROH) are used in as shown in the Table below.

Entry	Alcohol (ROH)	Conditions	Yield of esters
1	PhCH ₂ CH ₂ OH	H ₂ SO ₄ , 90 °C, 6h	10%
2	PhCH ₂ CH ₂ OH	H ₂ SO ₄ , CuI, 90 °C, 6h	9%
3	PhCH ₂ CH ₂ OH	TsOH , 90 °C, 6h	~39%
4	PhCH ₂ CH ₂ OH	TsOH, CuI, 90 °C, 6h	~39%
5	PhCH ₂ CH ₂ OH	Conc. HCl, 75 °C, 3h	15%
6	PhCH ₂ CH ₂ OH	2 N HCl, rt, 18h	trace
7	PhCH ₂ CH ₂ OH	6 N HCl, rt, 2h	~30%
8	MeOH	9N HCl, rt, 18h	Product on LC yield N/A
9	PhCH ₂ CH ₂ OH	~7 N HCl, rt, 18h	~33%
10	PhCH ₂ CH ₂ CH ₂ OH	~6 N HCl, rt, 18 h	~ 9%

11	nBuOH	~2 N HCl, 50 °C, 2h	~15%
12	nBuOH	~9 N HCl, rt, 18h	~70%
13	nBuOH	~9 N HCl, rt, 18h	~73%
14	Iso-BuOH	~9 N HCl, rt, 18 h	~69%

Preparation of (S)-3-[(5-Chloro-thiophene-2-carbonyl)-amino]-tetrahydro-furan-3-carboxylic acid ((S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-3-

5 benzazepin-7-yl)-amide



10 Resolution of (RS)-1 to prepare (S)-1:

To a 2 L jacketed-flask with condenser and mechanical stirrer was charged crude (RS)-1 (84 g), *n*PrOH (840 mL), and *D*-DTTA (122 g). The mixture was heated to 60 °C and then water (252 mL) was added. The mixture was then heated to reflux to become a clear solution. After stirring at reflux for ~ 0.5 h, the mixture
5 was cooled to ambient temperature over 2 h, and further cooled to 0 °C over 0.5 h and kept at 0 °C for 1h. The slurry was filtered and washed with cold solvent to give the salt (101 g) in 39% yield and 95% de.

The salt was then recrystallized from *n*PrOH/water (800mL/200mL) to give the (S)-1 (95 g) in >99% de and 37% overall yield.

10

Preparation of 4 from 2:

To a 2 L reaction was charged 2-(4-nitrophenyl)ethanol **2** (150 g, 0.882 mol), dichloromethane (1 L) and methanesulfonyl chloride (75.8 mL, 0.971 mol, 1.1 eq) at 20-25 °C. The mixture was cooled to -10 °C, and then triethylamine (107 g,
15 1.06 mol, 1.2 eq) was added slowly over 2 h while maintaining the internal temperature < 5 °C. The mixture was then warmed to 25 °C and stirred at this temperature for 1.5 h. Additional MsCl (4 mL, 0.05 eq) was added in one portion. The mixture was further stirred at 20-25 °C for 1 h. 1 N HCl (800 mL) was added and the mixture was stirred at 20-25 °C for 10 min. The organic
20 layer was separated and the aqueous layer was discarded. The organic phase was washed with 5% NaCl solution (500 mL) and concentrated to about 800 mL. To the solution was charged methanesulfonic acid (285 mL, 4.41 mol, 5 eq) in one portion, followed by addition of 1,3-dibromo-5,5-hydantoin (151 g, 0.53 mol, 0.6 eq) in several portions over 5 min. The mixture was stirred at 25-32 °C
25 for 2 h. Additional 1,3-dibromo-5,5-hydantoin (25 g, 0.09 mol, 0.1 eq) was

added and the mixture was further stirred at 25-30 °C for 4 h. The reaction mixture was cooled to 0 °C and water (800 mL) was cautiously added over 30 min while controlling the temperature < 35 °C. The organic phase was separated and washed sequentially with 10% Na₂S₂O₃ (500 mL), 3% NaHCO₃ (500 mL), 5% NaCl (500 mL), and concentrated to give crude **4** as an oil, which was directly used in the next step.

Preparation of 5 via amination:

To a solution of TEA (180 mL) and methylallylamine (96.5 g) in DMF (250 mL) at 5 °C was added dropwise a solution of crude **4** in DMF (200 mL) over 1 h while controlling the reaction temperature < 15 °C. The mixture was kept stirring at 15-18 °C for 2 h, then quenched with 3% NaHCO₃ (1 L) and EtOAc (1 L). The organic phase was separated and the aqueous phase was further extracted once with EtOAc (500 mL). The combined organic phase was washed with brine, concentrated to about 1 L, and cooled to 0-5 °C. HCl (~64.8 g) gas was bubbled through the solution to form a slurry. The slurry was filtered and the wet cake was washed with EtOAc (200 mL), dried at 20-25 °C with house vacuum over 12 h to give the desired HCL salt of **5** as a white solid (241 g, 0.75 mol, 85% overall yield from **2**). ¹HNMR (400 MHz, CDCl₃) δ 8.40 (d, *J* = 2.3 Hz, 1H), 8.10 (dd, *J* = 8.4, 2.3 Hz, 1H), 7.42 (d, *J* = 8.4 Hz, 1H), 5.83 (m, 1H), 5.17 (m, 2H), 3.09 (d, *J* = 6.4 Hz, 2H), 3.01 (m, 2H), 2.65 (m, 2H), 2.34 (s, 3H)

Synthesis of compound 6 via a Heck reaction:

To a 2 L 3-necked flask equipped with mechanical stirrer and thermometer at room temperature was charged aryl amine (85 g, 256 mmol), dioxane (640 mL)

and triethylamine (52 g, 511 mol, 2 equiv.) under argon. The mixture was degassed with argon for 15 min then Pd₂dba₃ (11.7 g) and PtBu₃.HBF₄ (7.4 g) were added under argon. The whole mixture was further degassed at room temperature for 5 min, and then heated to 100 °C and stirred at this temperature under argon for 1.5 h. The mixture was cooled to ~ 50 °C and distilled under house vacuum to remove most of dioxane, and then EtOAc (0.8 L) and 3% NaHCO₃ (0.5 L) solution were added. The mixture was filtered through a pad of diatomaceous earth to remove some precipitated palladium species. The organic phase was separated and the aqueous phase was extracted once with EtOAc. The combined organic phase was washed with brine (0.4 L) and concentrated to about 0.5 L. 5-6 M HCl in IPA (56 mL) was added at 0 °C and the resulting slurry was filtered. The wet cake was washed and dried to give the HCL salt of **6** as a yellowish solid (66 g, 84% yield, 94% purity). ¹HNMR (400 MHz, CDCl₃) δ 8.10 (d, *J* = 2.4 Hz, 1H), 8.03 (dd, *J* = 8.2, 2.4 Hz, 1H), 7.24 (d, *J* = 8.3 Hz, 1H), 5.37 (d, *J* = 1.1 Hz, 1H), 5.35 (d, *J* = 1.0 Hz, 1H), 3.34 (s, 2H), 2.98 (m, 2H), 2.81 (m, 2H), 2.43 (s, 3H).

Preparation of 9 by using CDI as coupling reagent:

To a suspension of 5-chloro-thiophenecarboxylic acid (25.7 g) in 100 mL of EtOAc was added CDI (24.85 g) in one portion at 20-25 °C over 10 min while gas was generated vigorously. The mixture was heated to reflux for 30 min and then cooled to room temperature. DMAP (1.44 g) was added followed by the amino acid butyl ester **8** (22 g) in EtOAc (150 mL) in one portion. The mixture was reflux for 20 h and cooled to 0-5 °C. 3 N HCl (150 mL) was added and the resulting mixture was stirred at room temperature for 10 min. The organic phase

was separated, washed with 5% NaHCO₃ (150 mL) and 5% NaCl solution (50 mL). The organic phase was concentrated to ~ 50 mL and MeOH (150 mL) was added. The mixture was further concentrated to ~ 100 mL.

- 5 To the above solution was added 2 N NaOH (90 mL, 1.5 equiv) and the mixture was stirred at room temperature for 2 h. The mixture was distilled to remove most of the MeOH and then cooled to 0 °C. 12 N HCl (~20 mL) was added dropwise to adjust the pH to 1~ 2 while maintaining the internal temperature <30 °C. The resulting mixture was extracted with Me-THF (200 mL). The
10 organic layer was separated, washed with 5% NaCl (150 mL), and concentrated to ~100 mL. To the residue was added heptane (100 mL) to form a slurry, which was filtered to give the desired product **9** (31.3 g, 96% yield, >98% purity) as a white solid.


15 ***Preparation of (S)-1 via asymmetric Hydrogenation of 6·HCl:***

- [Rh(COD)Cl]₂ (4.9 mg, 0.01 mmol) and Walphos (13.16 mg, 0.02 mmol) were stirred in 2 mL degassed MeOH at for 10 min before being transferred to a solution of compound **6** (127 mg, 0.5 mmol) in MeOH (2 mL). The mixture was stirred at room temperature under 100 psi H₂ for 12 h. HPLC showed complete
20 reduction of the double bond. To the mixture was added 10% Pd/C (10 mg). The mixture was further stirred at room temperature under 100 psi H₂ for 2 h, then filtered, and concentrated to give pure compound **(S)-1**. Chiral HPLC showed 79% ee.

25 ***Preparation of 10:***

Carboxylic acid **9** (3.04 g, 11.0 mmol, 1.05 eq) was dissolved in dry THF (50 mL) at room temperature followed by sequential addition of TEA (5.11 mL, 36.8 mmol, 3.5 eq) and propylphosphonic acid anhydride in EtOAc (50% w/w, , 7.0 g, 6.48 mL, 11.0 mmol, 1.05 eq). The mixture was stirred at room temperature for 5 10 min. To the mixture was added aniline (S)-1 (2.0 g, 10.5 mmol, 1.0 eq) in THF (10 mL) at room temperature. The resulting mixture was stirred at reflux for 2 h. LC showed 88 A% conversion of product along with 12% mixed anhydride. The mixture was quenched by adding 40 mL saturated NaHCO₃ solution and then was concentrated to remove most of THF. To the residue was added Me-10 THF (40 mL). The organic phase was separated and the aqueous layer was further washed with Me-THF (20 mL). The combined Me-THF extracts were washed with brined, dried over Na₂SO₄, concentrated, and purified by column chromatography (EtOAc/EtOH/Et₃N = 2:1:0.06) to give the desired product **10** (4.13 g, 9.2 mmol, 83%) as a white solid. ¹HNMR (400 MHz, CCl₃) δ 8.41 (s, 15 1H), 7.42 (s, 1H), 7.35 (d, *J* = 4.0 Hz, 1H), 7.27-7.33 (m, 2H), 7.06 (d, *J* = 8.0 Hz, 1H), 6.91 (d, *J* = 4.0 Hz, 1H), 4.34 (d, *J* = 9.4 Hz, 1H), 4.19-4.31 (m, 2H), 4.15 (d, *J* = 9.4 Hz, 1H), 3.19 (br, 1H), 3.03 (br, 1H), 2.75-2.95 (m, 3H), 2.71 (d, *J* = 12.3 Hz, 1H), 2.48 (m, 1H), 2.37 (s, 3H), 2.20-2.33 (br, 2H), 1.38 (d, *J* = 7.2 Hz, 3H); ¹³CNMR (100 MHz, CCl₃) δ 18.5, 35.4, 35.9, 47.6, 57.3, 64.3, 65.1, 20 68.1, 72.6, 117.7, 127.2, 127.7, 129.9, 135.7, 136.1, 137.4, 138.0, 146.2, 160.6, 171.7; ESI MS: 448 [M+H].

Preparation of (S)-3,5-Dimethyl-2,3,4,5-tetrahydro-1H-3-benzazepin-7-ylamine



Preparation of 11 and 12:

- 5 To a solution of **4** (120 g, 370 mmol) in dry DMF (750 mL) at ambient temperature is added allylamine (127 g, 2220 mmol). The reaction mixture is stirred at room temperature for 1 h. HPLC showed completion of the reaction. EtOAc (500 mL) and water (500 mL) were added to the reaction mixture. The organic layer was separated and the aqueous layer was extracted with EtOAc
- 10 (200 mL). The combined organic layer was washed with brine (2x250 mL) and concentrated to give the desired product **11** (103 g) in 97% yield.

- To a solution of **11** in CH_2Cl_2 (1 L) at 0°C was added TEA (2 eq) and TFAA (1.2 eq) over 0.5 h. The mixture was allowed to warm to room temperature and was
- 15 stirred for 3 h. Water (0.5 L) was added to quench the reaction and the resulting mixture was further stirred at room temperature for 10 min. The organic layer was separated and the aqueous layer was extracted with CH_2Cl_2 (200 mL). The combined CH_2Cl_2 was washed with water (0.5 L), brine (0.5 L), then concentrated to give the desired product as a brown solid (95% yield). $^1\text{HNMR}$

(400 MHz, CDCl_3) δ 8.45 (m, 1H), 8.14 (m, 1H), [7.44 (d, $J = 8.4$ Hz, major); 7.40 (d, $J = 8.4$ Hz, minor); 1H], [5.82 (m, minor), 5.68 (m, major); 1H], 5.20-5.32 (m, 2H), 3.92 (d, $J = 5.5$ Hz, 1H), 3.62 (t, $J = 7.8$ Hz, 2H), 3.18 (m, 2H).

5 **Preparation of 13:**

To a mixture of aryl bromide **12** (2.6 g, 6.82 mmol, 1.0 eq), Pd_2dba_3 (250 mg, 0.273 mmol, 0.04 eq), and *N*-methyldicyclohexylamine (2.00 g, 10.23 mmol, 1.5 eq) in anhydrous dioxane (60 mL, 0.1 M) under argon was added 10% (w/w) *t*- Bu_3P in hexane (1.62 mL, 0.55 mmol, 0.08 eq). The mixture was stirred under
10 argon at 80 °C for 2 h and then cooled to room temperature, quenched by adding water (40 mL) and EtOAc (40 mL). The organic phase was separated and the aqueous layer was washed once with EtOAc (40 mL). The combined EtOAc was washed with water (40 mL) and brine (40 mL), dried over Na_2SO_4 , and purified by column chromatography to give **13** (1.1 g, 3.77 mmol, 54%) as a
15 yellow solid along with 5-10% 8-membered ring by-product. (two atropisomers):
 ^1H NMR (400 MHz, CDCl_3) δ 8.19 (m, 1H), 8.11 (m, 1H), 7.33 (m, 1H), 5.40-5.60 (m, 2H), 4.48 (d, $J = 15.9$ Hz, 2H), 3.89 (m, 2H), 3.15 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 33.1, 35.22, 44.8, 46.2, 47.7, 50.4, 50.7, 114.9, 117.7, 118.7, 120.5, 122.3, 123.1, 123.4, 123.6, 123.7, 130.5, 130.6, 141.0, 141.4, 142.7,
20 143.2, 143.3, 143.9, 147.4.

Preparation of 14:

To a 300 mL autoclave was added Wilkinson catalyst $\text{RhCl}(\text{PPh}_3)_3$ (1.78 g, 1.92 mmol, 0.04 eq) followed by a solution of compound 18 (14.4 g, 48.0 mmol, 1.0
25 eq) in THF (100 mL). The mixture was stirred at room temperature under 30-40

psi H₂ for 12 h. LC showed complete conversion and no trace of over-reduced by-product was observed. Note: Around 10% of 8-membered ring by-product, which was generated from Heck reaction, was seen. The mixture was concentrated and purified by column chromatography (hexane →
5 hexane/EtOAc = 3/1) to give the desired hydrogenation product as a white solid. To a solution of the hydrogenation product in THF (80 mL) at 0 °C was added a solution of NaOH (1.98 g, 49.6 mmol) in water (20 mL). The mixture was stirred at room temperature for 2 h. LC showed completion of hydrolysis. The mixture was concentrated and the residue was treated with Me-THF (250 mL) and water
10 (100 mL). The Me-THF layer was separated and the aqueous layer was washed with Me-THF (100 mL). The combined Me-THF was washed with brine (100 mL), dried over Na₂SO₄, and concentrated to give the crude product (8.9 g, 43.2 mmol, 90% yield) as a yellow solid.

15 ***Preparation of 15:***

A mixture of racemic amine **14** (8.0 g, 38.8 mmol, 1 eq) and *L*-madelic acid (4.43 g, 29.1 mmol, 0.75 eq) in acetone (90 mL) and water (9 mL) was heated to reflux to become a transparent solution. The mixture was cooled to 0 °C over 6 h while stirring. The resulting slurry was filtered to give enantiomerically
20 enriched salt with 58 % ee (7.0 g, 50.4 % yield). The salt was further crystallized from ethanol-water for 5 consecutive times to give the salt (2.5 g, 18 %) with 97.0 % ee. The salt was then treated with 2 N NaOH (20 mL) and Me-THF (50 mL). The Me-THF layer was separated and the aqueous layer was extracted with Me-THF. The combined Me-THF layer was washed with brine (20 mL),


dried over Na_2SO_4 , and concentrated to give compound **15** as a yellow crystalline solid (1.44 g, 7.0 mmol, ee: 97.0%, 18% yield).

Preparation of 16 and (S)-1:

5 The intermediate **15** (1.44 g, 7.0 mmol) was dissolved in 10 mL of HCOOH . To the mixture at room temperature was added 37% HCHO (0.81 g, 37% w/w, 10.5 mmol). The mixture was stirred at 90 °C for 3 h, and concentrated to give a yellow oily product. The residue was diluted with Me-THF (50 mL) and 2 N NaOH (20 mL) and stirred at room temperature for 10 min. The aqueous layer
10 was further washed with Me-THF and the combined Me-THF was washed with brine and concentrated to give methylated compound **16**.

The above residue (1.8 g, 8.2 mmol, 98.2% ee) was dissolved in MeOH (20 mL) and 10% Pd/C (200 mg) was added. The mixture was stirred at room
15 temperature under 100 psi H_2 for 12 h, filtered to remove Pd/C, and concentrated. The residue was purified by column chromatography [$\text{EtOAc}(\text{Et}_3\text{N}$ 0.06 v/v)/ MeOH = 100/0 → 50/50 as eluent) to give the aniline compound **(S)-1** (1.57 g, 8.2 mmol, 100%, 98.2% ee, 98.5A% HPLC purity) as a brown oil.
 ^1H NMR (400 MHz, CD_3OD) δ 6.82 (d, J = 7.9 Hz, 1H), 6.61 (s, 1H), 6.49 (dd, J
20 = 7.8, 2.2 Hz, 1H), 3.07 (m, 1H), 2.93 (m, 1H), 2.60-2.85 (m, 3H), 2.30 (s, 3H), 2.18 (br, 2H), 1.32 (d, J = 7.3 Hz, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 19.2, 35.6, 47.9, 59.2, 66.0, 114.4, 114.6, 131.0, 132.3, 146.9, 147.0; ESI MS: 191 $[\text{M}+\text{H}]$.

Preparation of **(S)-3,5-Dimethyl-2,3,4,5-tetrahydro-1H-3-benzazepin-7-ylamine**
25



5 Preparation of 17:

To a solution of olefin **6** (200 mg, 0.92 mmol) in THF (2 mL) was added $RhCl(PPh_3)_3$ (34 mg, 0.037 mmol, 0.04 eq). The mixture was stirred at room temperature under 30–40 psi H_2 for 12 h, then concentrated and purified by column chromatography to give the desired amine **17** as a thick oil. **17**: 1H NMR (400 MHz, $CDCl_3$) δ 8.05 (d, $J = 2.2$ Hz, 1H), 8.00 (dd, $J = 8.2, 2.3$ Hz, 1H), 7.24 (d, $J = 8.2$ Hz, 1H), 3.29 (m, 1H), 3.22 (m, 1H), 2.80–3.10 (m, 2H), 2.89 (d, $J = 12.3$ Hz, 1H), 2.37 (s, 3H), 2.15–2.30 (m, 2H), 1.44 (d, $J = 7.2$ Hz, 3H).


Preparation of 18:

Same procedure was used for the resolution of (RS)-**1** by using DTTA.

C) Examples:

20 Example 1

(R)- and (S)-3-[(5-bromo-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



(a) 7-nitro-2,3,4,5-tetrahydro-1H-benzo[d]azepine

- 5 8.4 g (29.0 mmol) 3-trifluoroacetyl-7-nitro-2,3,4,5-tetrahydro-1H-benzo[d]azepine are suspended under a nitrogen atmosphere in 80 ml of methanol and combined with 5 ml NaOH solution (50%) and stirred for 2 h at 70°C.

The methanol is distilled off using the rotary evaporator, the residue is

- 10 combined with water and extracted with *tert.*-butylethylether. The organic phase is washed with NaOH solution (50%) and sat. sodium chloride solution, dried on sodium sulphate and evaporated to dryness i. vac..

Yield: 5.1 g (91%)

R_f value: 0.28 (aluminium oxide; dichloromethane/ethanol = 95:5)

- 15 C₁₀H₁₂N₂O₂ (192.22)

Mass spectrum: (M+H)⁺ = 193

(b) 3-methyl-7-nitro-2,3,4,5-tetrahydro-1H-benzo[d]azepine

- 20 5.0 g (26.0 mmol) 7-nitro-2,3,4,5-tetrahydro-1H-benzo[d]azepine are mixed in 9.8 ml formic acid with 15.5 ml formalin solution in water (37%) at room temperature, and stirred overnight at 70°C. The reaction mixture is made alkaline with NaOH solution (50%) while cooling with an ice bath and extracted

with *tert.*-butylmethylether. The organic phase is dried on sodium sulphate and evaporated to dryness i. vac..

Yield: 4.8 g (90%)

R_f value: 0.65 (aluminium oxide; dichloromethane/ethanol = 95:5)

5 C₁₁H₁₄N₂O₂ (206.24)

Mass spectrum: (M+H)⁺ = 207

(c) 3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-ylamine

10 4.8 g (23.2 mmol) 3-methyl-7-nitro-2,3,4,5-tetrahydro-1*H*-benzo[d]azepine are dissolved in 45 ml of methanol and combined with 400 mg Pd/C 10%. The mixture is hydrogenated in a Parr apparatus at room temperature at 3 bar hydrogen pressure for 5 hours. Then the catalyst is filtered off and the filtrate is evaporated down i. vac..

15 Yield: 3.9 g (96 %)

R_f value: 0.36 (aluminium oxide; dichloromethane/ethanol = 98:2)

C₁₁H₁₆N₂ (176.26)

Mass spectrum: (M+H)⁺ = 177

20 (d) 3-amino-tetrahydro-furan-3-carboxylic acid-hydrochloride

3.5 g (15.1 mmol) 3-*tert.*-butoxycarbonylamino-tetrahydro-furan-3-carboxylic acid are dissolved in 150 ml of 1-molar hydrochloric acid and stirred for 1 h at room temperature. Then the reaction mixture is lyophilised.

25 Yield: 2.5 g (100 %)

$C_5H_9NO_3 \cdot HCl$ (167.59)

Mass spectrum: $(M+H)^+$ = 132

(e) 3-[(5-bromo-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic
5 acid

3.1 g (14.9 mmol) 5-bromo-thiophene-2-carboxylic acid in 50 ml
dichloromethane are combined with 5.4 ml (74.6 mmol) thionyl chloride with
stirring at room temperature and stirred for 3.5 h at reflux temperature. Then the
10 reaction mixture is evaporated to dryness.

2.5 g (14.9 mmol) 3-amino-tetrahydro-furan-3-carboxylic acid-hydrochloride are
dissolved in 2.0 ml (14.9 mmol) TEA and 150 ml acetonitrile and combined with
5.9 ml (22.4 mmol) *N,O*-bis-(trimethylsilyl)-trifluoro-acetamide with stirring and
refluxed for 4 h with stirring. The reaction mixture is combined with 4.1 ml (29.8
15 mmol) TEA and the solution of the prepared acid chloride in 50 ml acetonitrile,
stirred for 15 min at reflux temperature and then cooled slowly to room
temperature. Then the mixture is evaporated to dryness i. vac., the residue is
combined with water and 2-molar sodium carbonate solution and washed with
diethyl ether. The aqueous phase is adjusted to pH 1 with 20 ml conc.
20 hydrochloric acid, the precipitate is suction filtered and dried at 50°C in the
vacuum drying cupboard.

Yield: 3.6 g (75 %)

$C_{10}H_{10}BrNO_4S$ (320.16)

Mass spectrum: $(M-H)^-$ = 318/320 (bromine isotopes)

(f) 3-[(5-bromo-thiophen-2-yl)-carbonylamino]-*N*-(3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

700.0 mg (2.19 mmol) 3-[(5-bromo-thiophen-2-yl)-carbonylamino]-tetrahydro-
5 furan-3-carboxylic acid are combined with 890.0 mg (2.34 mmol) HATU and
601.0 µl (5.47 mmol) NMM in 10 ml DMF with stirring at room temperature and
stirred for 10 min. Then 385.0 mg (2.19 mmol) 3-methyl-2,3,4,5-tetrahydro-1*H*-
benzo[*d*]azepin-7-ylamine are added and the mixture is stirred overnight at
65°C. The reaction mixture is combined with water and sat. sodium hydrogen
10 carbonate solution, the precipitate is filtered off and purified by chromatography
on aluminium oxide (eluant: dichloromethane/ethanol 100:0 to 98:2).

Yield: 850.0 mg (81 %)

R_f value: 0.62 (aluminium oxide; dichloromethane/ethanol = 95:5)


C₂₁H₂₄BrN₃O₃S (478.40)

15 Mass spectrum: (M+H)⁺ = 478/480 (bromine isotopes)

For separation of the racemic mixture into its respective enantiomers, a
conventional HPLC system with DAICEL AD-H 250mm x 4,6mm chiral column
has been used, eluting with (0,2% Diethylamine in Hexane)/ isopropanol 70/30
20 as liquid phase. At a flow rate of 1ml/min, retention times for the enantiomers
are 13.6 min and 16.4 min.

Example 2

(*R*)- and (*S*)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-(3-methyl-2,3,4,5-
25 tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



(a) benzyl 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylate

5

1.59 g (9.8 mmol) 5-chloro-thiophene-2-carboxylic acid is dissolved in 30 ml DMF and stirred with 3.61 g (10.7 mmol) benzyl 3-amino-tetrahydro-furan-3-carboxylate and 3.46 g (10.8 mmol) TBTU and 4.3 ml (39 mmol) NMM at room temperature for 20 h. Then the mixture is evaporated down and purified by chromatography on silica gel (eluant: dichloromethane/ethanol 100:0 to 94:6).

10

Yield: quantitative

R_f value: 0.59 (silica gel; dichloromethane/ethanol = 9:1)

$C_{17}H_{16}ClNO_4S$ (365.83)

Mass spectrum: $(M+H)^+$ = 366/368 (chlorine isotopes)

15

(b) 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid

20

3.6 g (9.8 mmol) benzyl 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylate are dissolved in 60 ml of ethanol and combined with 39.1 ml (39.1 mmol) 1-molar aqueous sodium hydroxide solution and stirred for 6 h at room temperature. After evaporation i. vac. the residue is combined with 1-

molar aqueous hydrochloric acid while cooling with an ice bath, the precipitate is suction filtered and dried at 60°C in the vacuum drying cupboard.

Yield: 2.5 g (91 %)

R_f value: 0.13 (silica gel; dichloromethane/ethanol 9:1)

5 C₁₀H₁₀ClNO₄S (275.71)

Mass spectrum: (M-H)⁻ = 274/276 (chlorine isotopes)

(c) 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-(3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

10

Prepared analogously to Example 2(a) from 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-tetrahydro-furan-3-carboxylic acid and 3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-ylamine with TBTU and TEA in THF at room temperature with subsequent purification by chromatography with aluminium oxide (eluant: dichloromethane/ethanol 100:0 to 97:3).

15

Yield: 67 %

R_f value: 0.63 (aluminium oxide; dichloromethane/ethanol = 95:5)

C₂₁H₂₄ClN₃O₃S (433.95)

Mass spectrum: (M+H)⁺ = 434/436 (chlorine isotopes)

20

For separation of the racemic mixture into its respective enantiomers, a conventional analytical HPLC system with DAICEL IA 250mm x 4,6mm chiral column has been used, eluting with EtOH as liquid phase. At a flow rate of 0,5ml/min, retention times for the enantiomers are 13,10 min and 16,30 min.

25

Alternatively, for separation of the racemic mixture into its respective enantiomers, a conventional HPLC system with DAICEL AD-H 250mm x 4,6mm chiral column has been used, eluting with (0,2% Cyclohexylamine in Hexane)/ isopropanol 70/30 as liquid phase. At a flow rate of 1ml/min, retention times for the enantiomers are 12.8 min and 15.2 min.

d) (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide was also prepared according to the following procedure:

10

To a mixture of 1.08 g (6.13 mmol) **7** and 1.86 g (6.75 mmol) 3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-ylamine in 50 ml THF, was added 2.98 ml TEA and 4.7 ml 1-propylphosphonic acid cyclic anhydride (50% in EtOAc). The mixture was refluxed for 2 h, the solvents were removed i.vac. and the crude mixture was purified by chromatography (Method A) to yield the title compound in 89%.

15

or to the following procedure:

20

To a solution of 5-chloro-thiophene-2-carboxylic acid in MeCN (0.1 g, 0.615 mmol) was added TsCl (0.106g, 0.554 mmol) in one portion. After the mixture was cooled to 0 °C, NMM (0.38 mL, 310 mg, 3.08 mmol, 5 equiv.) was added slowly and the mixture was warmed to room temperature and stirred for 3 h, then heated to 50 °C for 0.5 h. The reaction was monitored by HPLC for the

25

disappearance of TsCl (<1% by area). Salt 20 (0.141 g, 0.388 mmol, KF ~1%) was added and the reaction mixture was stirred at ambient temperature for 2 h and then concentrated to remove MeCN. EtOAc (50 mL) and sat. NaHCO₃ (50 mL) was added and the mixture was stirred at rt for 15 min. The organic phase
5 was separated, washed with sat. NaHCO₃ solution (50 mL) and brine (50 mL), and concentrated to give the desired example (S)-2 as a white solid (75% based on HPLC assay).

The following solubility and solid state characteristics of the compound (S)-3-
10 [(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and of its anhydrous crystalline form are relevant to the present invention.

Solubility properties of the compound (S)-3-[(5-chloro-thiophen-2-yl)-
15 carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

Solubility and dissolution rate in aqueous media

20 The table below shows the values of solubility of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide in different aqueous media.

medium	solubility [mg/ml]	pH of the saturated solution
water	0.28	8.6
0.1 N HCl	> 10	1.0
pH 2.2	> 10	2.4
pH 4.0	> 10	4.3
pH 6.0	> 10	6.3
pH 7.4	1.8	7.5
0.01 N NaOH	0.046	11.8

The table below shows the values of intrinsic dissolution rate of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide in aqueous media. The intrinsic dissolution rate was determined in aqueous media covering a range of pH 1.1 – 7.4 using the rotating disc method which maintains a constant surface area. 5 mg of drug substance was compressed to form a disc at 356.1 N for 60 s. These discs were mounted to specially designed sample holders which fit into a Sotax dissolution tester. The dissolution media (37°C) were stirred at 200 rpm. Samples were automatically withdrawn every second minute from the dissolution vessel and assayed via UV spectrophotometry. The intrinsic dissolution rate expressed in $\mu\text{g}/\text{cm}^2/\text{min}$ was calculated using the slope of the concentration versus time plot and from the linear portion of the slope of the dissolution curve, volume of dissolution medium (35ml) and area (diameter: 2 mm) of the exposed disk.

pH of aqueous medium	dissolution rate [$\mu\text{g}/\text{cm}^2/\text{min}$]
1.1	8220
2.3	4520
3.2	4000
4.1	3110
5.1	4015
6.0	4140
7.4	240

From the above results, it can be concluded that the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-

- 5 benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide has a pH dependent solubility profile in aqueous media with excellent solubility in acidic media and a reduced solubility in neutral and basic media due to the lower solubility of the free base. Furthermore, the compound exhibits up to pH 6.0 very fast dissolution rates, and even acceptable dissolution rates at pH 7.4.

10

Solid state properties of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

- 15 In the solid state, the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide appears as a white microcrystalline powder.

Manufacturing process of the anhydrous crystalline form of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide.

5

The anhydrous crystalline form of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide may be manufactured by drying a preparation of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide at a temperature above 130°C and maintaining it under dry atmosphere.

10

Solid state properties (crystallinity and polymorphism)

15 of the anhydrous crystalline form of the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

The compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide is highly crystalline. The X-ray powder diffraction diagram is shown in Figure 1.

20

The X-ray powder reflection and intensities (standardised) are shown in the following table.

25

2 θ [°]	d-value [Å]	Intensity I/I ₀ [%]
5.17	17.07	39
8.39	10.54	85
11.48	7.70	10
12.29	7.19	11
12.93	6.84	23
14.93	5.93	11
15.56	5.69	6
16.23	5.46	26
16.81	5.27	8
17.28	5.13	28
18.85	4.70	85
19.11	4.64	100
19.72	4.50	8
20.26	4.38	13
20.46	4.34	30
20.75	4.28	4
21.06	4.22	4
21.79	4.08	5
21.97	4.04	6
22.43	3.96	12
22.82	3.89	13
23.09	3.85	7
24.02	3.70	20
24.69	3.60	15
25.54	3.49	9
26.01	3.42	5
26.59	3.35	32
27.52	3.24	3
27.78	3.21	3
28.89	3.09	25
29.31	3.05	9
30.13	2.96	9
30.95	2.89	6

In the above table, the value " 2θ [$^\circ$]" denotes the angle of diffraction in degrees and the value " d_{hkl} [$\text{\AA}$]" denotes the specified distances in $\text{\AA}$ between the lattice planes.

- 5 According to the findings shown in the above table, the present invention further relates to the crystalline anhydrous form of (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide, characterised in that in the x-ray powder diagram has, inter alia, the characteristic values $d = 3.35 \text{ \AA}$, $4.34 \text{ \AA}$,
10 $4.64 \text{ \AA}$, $4.70 \text{ \AA}$, $10.54 \text{ \AA}$ and $17.07 \text{ \AA}$ (most prominent peaks in the diagram).

The material crystallizes in rod-like crystals and tends to agglomerate in larger aggregates, as shown in Figure 2.

- 15 The thermoanalysis of the crystalline anhydrous form of (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide according to the invention, as shown in Figure 3 (DSC/TG diagram), shows a $T_{fus} = 185 \pm 3 \text{ }^\circ\text{C}$ (DSC: $10 \text{ K}\cdot\text{min}^{-1}$ heating rate), in the form of a strong endothermic peak. A
20 closer look to the TG-trace (confirmed by TG-IR coupling experiments) shows a weight loss of approx. 1.0 – 2.0% up to approximately 180°C . This weight loss may indicate the presence of absorbed water on the surface of the microcrystalline material. Thermal decomposition starts above 240°C indicating a congruent melting process at 185°C .


Thus, the present invention further relates to the crystalline anhydrous form of (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide, characterised by a melting point of $T_{m.p.} = 185 \pm 3^{\circ}\text{C}$ (determined by DSC; evaluation using peak-maximum; heating rate: 10°C/min).

From the above data, it can be concluded that the compound (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide is characterized by its high solubility in acidic media and its high crystallinity. The crystalline polymorphic form is characterized as an anhydrous form, present as single stable polymorphic form.

Example 3

(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1R)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1S)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

-121-



(a) 1-methyl-2-oxo-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine

- 5 8.0 g (37 mmol) 2-chloro-*N*-(2-phenylethyl)-propanamide and 15 g (112 mmol) aluminium trichloride were carefully mixed at 90°C and heated to 150°C for 6 h. The mixture was diluted with water and methanol and extracted with EtOAc. The combined organic layers were dried with Na₂SO₄, concentrated and purified by chromatography (method A) to give the title compound,

10

(b) 1-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine

- 2.7 g (15 mmol) of 1-methyl-2-oxo-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine was added to 46 ml 1M BH₃-THF complex solution and stirred at room temperature under a nitrogen atmosphere overnight. 50 ml of methanol were carefully added, followed by 30 ml of 2M HCl. The mixture was extracted with EtOAc, the combined organic layers were dried with Na₂SO₄, concentrated and purified by chromatography (method A) to give the title compound as formic acid salt.
- 15

(c) 1,3-dimethyl-7-nitro-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine and 1,3-dimethyl-8-nitro-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine.

Following the procedure for example 1 b 1-methyl-2,3,4,5-tetrahydro-1*H*-

5 benzo[d][azepine was methylated to give 1,3-dimethyl-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine.

1.79 g (10 mmol) of 1,3-dimethyl-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine was mixed with 3.7 ml conc. H₂SO₄ and 0.71 ml 65% HNO₃ at -5°C and stirred for 1 h by -5°C to 0°C. The mixture was poured into 100 ml of ice cold water and 10 M NaOH was added. The mixture was extracted with EtOAc, the combined organic layers were dried with Na₂SO₄, concentrated and purified by chromatography on silica gel (eluant: dichloromethane:95%ethanol/5% ammonia 99:1 to 95:5) to give a mixture of the title compounds.

15 (d) 3,5-dimethyl-7-amino-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine and 3,5-dimethyl-8-amino-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine

1.4 g (6.3 mmol) of a mixture of 1,3-dimethyl-7-nitro-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine and 1,3-dimethyl-8-nitro-2,3,4,5-tetrahydro-1*H*-benzo[d][azepine, 20 ml methanol and 0.20 g 10% palladium on charcoal was stirred for 5.5 h under an atmosphere of hydrogen (50 psi). It was filtered, concentrated and the mixture was purified by chromatography on silica gel (eluant: dichloromethane:95%ethanol/5% ammonia 99:1 to 80:20) to give 0.45 g of regioisomer B: rac-7-Amino-1,3-dimethyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepine

25

R_f value: 0.75 (silica gel; dichloromethane/ethanol/ammonia = 80:20:2)

C₁₂H₁₈N₂ (190.28)

Mass spectrum: (M+H)⁺ = 191

and 0.55 g of regioisomer A: rac-7-Amino-3,5-dimethyl-2,3,4,5-tetrahydro-1H-

5 benzo[d]azepine

R_f value: 0.70 (silica gel; dichloromethane/ethanol/ammonia = 80:20:2)

C₁₂H₁₈N₂ (190.28)

Mass spectrum: (M+H)⁺ = 191.

10 e) Regioisomer A and S-configured carboxylic acid 7 were coupled according to the procedure described in example 2d to give a mixture of the 3S-diastereoisomers.

R_f value: 0.75 (silica gel; dichloromethane/ethanol/ammonia = 80:20:2)

C₂₂H₂₆ClN₃O₃S (447.979)

15 Mass spectrum: (M+H)⁺ = 448/450 chloro isotopes.

For separation of the mixture of diastereoisomers into the pure single stereoisomers (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and

20 (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide a conventional HPLC system with DAICEL AD-H 250mm x 4,6mm chiral column has been used, eluting with 0,2% Cyclohexylamine in Hexane(80%) /

EtOH(20%) as liquid phase. At a flow rate of 1ml/min, retention times for the stereoisomers are 10,75min and 16,5min.

Each of the diastereoisomers shows the following

Mass spectrum: $(M+H)^+$ = 448/450 chloro isotopes.

5

Alternatively, the separation of the mixture of diastereoisomers can be achieved by supercritical fluid chromatography with DAICEL AD-H chiral column, eluting with 0,2% Cyclohexylamine in EtOH(45%) / supercritical CO₂(65%). At a flow
10 rate of 5ml/min, retention times for the stereoisomers are 3,94min and 4,08min respectively.

f) Regioisomer B and S-configured carboxylic acid **7** can be coupled according to the procedure described above to give a diastereomeric mixture of
15 the regioisomers. (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1R)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and

(3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1S)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amid.

20 The diastereoisomers could be separated by chiral chromatography (DAICEL AS-H, 250 x 4.6 mm, eluens: Methanol and 45% Diethylamine). At a flow rate of 5ml/min, retention times for the stereoisomers are 6.5 min and 8.5 min.


Each of the diastereoisomers shows the following

Mass spectrum: $(M+H)^+$ = 448/450 chloro isotopes.

25

Example 3-A

- (3R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
- 5 (3R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



- Regioisomer A can be coupled separately with the R-configured enantiomer
- 10 of carboxylic acid 7 following the procedure for example 3 to give the 3R-diastereoisomers.

Both diastereoisomers showed the following characteristics


R_f value: 0.6 (silica gel; dichloromethane/ethanol/ammonia = 80:20:2)

C₂₂H₂₆ClN₃O₃S (447.979)

- 15 Mass spectrum: (M+H)⁺ = 448/450 chloro isotopes.

Example 4

- (R)- and (S)-5-chloro-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-cyanimin-
- 20 [1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide



500 mg (1.9 mmol) 2-(5-chloro-thiophene-2-yl)-3,7-dioxo-1-aza-spiro[4.4]non-1-en-4-one are stirred with 0.45 g (1.8 mmol) 3-methyl-4-(5-oxo-[1.4]oxazepan-4-yl)-aniline in 4.5 ml of toluene with 5.0 ml DMF and 500 μ l glacial acetic acid at 80°C for 5 h. Then the reaction mixture was concentrated poured into 100ml semisat. sodium hydrogen carbonate/100 ml EtOAc. After extraction with EtOAc, the combined organic phases were dried over magnesium sulphate and evaporated down i. vac.. The racemic mixture was separated into the enantiomers by using a chiral chromatography column (500X50mm, DAICEL AD-column, eluent: Ethanol 30 ml/min):

Enantiomer 1: R_t = 86 min


Mass spectrum: $(M+H)^+$ = 502/504 (chlorine isotopes)

Enantiomer 2 R_t = 136 min.

15 Mass spectrum: $(M+H)^+$ = 502/504 (chlorine isotopes)

Example 5

(R)- and (S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide



The (S)-Enantiomer was obtained by following the procedure for example 4 and using (S)-2-(5-chloro-thiophene-2-yl)-3,7-dioxaspiro[4.4]non-1-en-4-one **8** and 3-methyl-4-(5-oxo-[1,4]oxazepan-4-yl)-aniline.

5 $C_{22}H_{22}ClN_5O_4S$ (487.96)

Mass spectrum: $(M+H)^+$ = 488/490 (chlorine isotopes)

The (R)-Enantiomer was obtained by following the same procedure using (R)-2-(5-chloro-thiophene-2-yl)-3,7-dioxaspiro[4.4]non-1-en-4-one and 3-methyl-4-(5-oxo-[1,4]oxazepan-4-yl)-aniline.


10

$C_{22}H_{22}ClN_5O_4S$ (487.96)


Mass spectrum: $(M+H)^+$ = 488/490 (chlorine isotopes)

15 Example 6

(R)- and (S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-oxo-[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl]-amide



The racemic mixture was prepared according to the following scheme as described in WO2005111029:



For separation of the racemic mixture into its respective enantiomers, a conventional analytical HPLC system with DAICEL AD-H 250mm x 4,6mm chiral column has been used, eluting with 0,2% Cyclohexylamine in
 10 Hexane(80%) / IPA(20%) as liquid phase. At a flow rate of 1ml/min, retention times for the enantiomers are 23,70min and 28,40min.

Enantiomer 1: $R_t = 23.70$ min


Mass spectrum: $(M+H)^+ = 520/522$ (bromine isotopes)

Enantiomer 2: $R_t = 28.40$ min.

Mass spectrum: $(M+H)^+$ = 520/522 (bromine isotopes)

Example 7

(R)- and (S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(5-cyanimin-
5 [1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide



The racemic mixture was prepared from rac-3-[(5-chloro-thiophen-2-yl)-
carbonylamino]-tetrahydro-furan-3-carboxylic acid **rac-7** and 4-(5cyanimino-
10 [1,4]oxazepan-4-yl)anilin (prepared in analogy to procedures described in
WO2005/111029) in analogy to procedure 4..

$C_{22}H_{22}BrN_5O_4S$ (532.411)


Mass spectrum: $(M+H)^+$ = 532/534 (bromine isotopes)

- 15 For separation of the racemic compound into its respective enantiomers, a
conventional analytical HPLC system with DAICEL AD-H 250mm x 4,6mm
chiral column has been used, eluting with 0,2% Acetic Acid in Hexane(80%) /
EtOH(20%) as liquid phase. At a flow rate of 1ml/min, retention times for the
enantiomers are 9 min and 12,2 min.
- 20 Alternatively, the separation of this racemate can be achieved on HPLC with
DAICEL OJ-H chiral column, eluting with 0,2% Acetic Acid in Hexane(80%) /

EtOH(20%) as liquid phase. At a flow rate of 1ml/min, retention times for the enantiomers are 6,10 min and 8,10 min respectively.

Example 8

- 5 (R)- and (S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide



The racemic mixture was prepared as described in WO2005/111029.


- For separation of the racemic mixture into its respective enantiomers, a
10 conventional HPLC system with a DAICEL OD 250mm x 20 mm chiral column
has been used, eluting first with hexane and after the first peak has been
eluated with hexane/ethanol 55/45 as liquid phases.

$C_{21}H_{22}BrN_3O_5S$ (508.39)

- Mass spectrum: $(M+H)^+$ = 508/510 (bromine isotopes) for each
15 enantiomer.

Example 9

(R)- and (S)- 5-ethynyl-thiophene-2-carboxylic acid-*N*-{3-[4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide





The racemic mixture was prepared as described in WO2006/034822.

For separation of the racemic compound into its respective enantiomers, a conventional analytical HPLC system with DAICEL AD-H 250mm x 4,6mm chiral column has been used, eluting with hexane/ethanol 1/1 as liquid phase.
 5 Retention times for the enantiomers were 13.9 min and 22.2 min. For the preparative separation a DAICEL AD-H 500mm x 50mm chiral column has been used, eluting with ethanol as liquid phase.


10 The following racemic compounds can be prepared in enantiomerically pure form analogously to the methods described in the Examples above or by synthetic pathways known from the literature:

A (S)- and (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-
 15 (pyrrolidin-1-yl-carbonyl)-phenylcarbonyl]-tetrahydrofuran-3-yl}-amide







B (S)- and (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(pyrrolidin-1-yl-carbonyl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




5




C (S)- and (R)-5-ethynyl-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(3-oxo-morpholin-4-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide




10




D (S)- and (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(2-oxo-azepan-1-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide




5




E (S)- and (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[4-(2-oxo-azepan-1-yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide




10





F (S)- and (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




5




G (S)- and (R)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(5-cyanimin-[1.4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




H (S)- and (R)-5-chloro-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(3-
5 cyanimino-morpholin-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide






I (S)- and (R)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[3-chloro-4-(3-cyanimino-morpholin-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide





5



J (S)- 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-((2S)-2,3-dimethyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid

10 amide and (S)- 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-((2R)-2,3-dimethyl-



2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid
amide







5



K (S)- 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-((4*S*)-3,4-dimethyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid
amide and (S)- 3-[(5-chloro-thiophen-2-yl)-carbonylamino]-*N*-((4*R*)-3,4-dimethyl-2,3,4,5-tetrahydro-1*H*-benzo[*d*]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid
amide

10





L5 **M**



N (S)- and (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(2,2,3-trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



- O (S)- and (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3,4,4-
 5 trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-
 carboxylic acid amide



- 10 P (S)- and (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3,5,5-
 trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-
 carboxylic acid amide



(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3,5,5-trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

5 was prepared in analogy to the procedures described above.


R_f value: 0.38 (RP-8; methanol/5%NaCl-solution = 6:4)


C₂₃H₂₈ClN₃O₃S (462,01)

Mass spectrum: (M+H)⁺ = 462/464 (chlorine isotopes)


10

Q (S)- and (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-9-fluor-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide






R (S)- and (R)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-8-fluor-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide





5



S (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1S)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and (S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1R)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide

10





The Examples that follow describe the preparation of pharmaceutical

- 5 formulations which contain as active substance any desired compound of general formula I.

Example A – Dry ampoule containing 75 mg of active substance per 10 ml

10 Composition:

Active substance	75.0 mg
Mannitol	50.0 mg
water for injections	ad 10.0 ml

15

Preparation:

Active substance and mannitol are dissolved in water. After packaging the solution is freeze-dried. To produce the solution ready for use for injections, the
20 product is dissolved in water.

Example B – Dry ampoule containing 35 mg of active substance per 2 ml

Composition:

5	Active substance	35.0 mg
	Mannitol	100.0 mg
	water for injections	ad 2.0 ml

Preparation:

- 10 Active substance and mannitol are dissolved in water. After packaging, the solution is freeze-dried.

To produce the solution ready for use for injections, the product is dissolved in water.

- 15 Example C – Tablet containing 50 mg of active substance

Composition:

	(1) Active substance	50.0 mg
20	(2) Lactose	98.0 mg
	(3) Maize starch	50.0 mg
	(4) Polyvinylpyrrolidone	15.0 mg
	(5) Magnesium stearate	2.0 mg
		215.0 mg

Preparation:

(1), (2) and (3) are mixed together and granulated with an aqueous solution of (4). (5) is added to the dried granulated material. From this mixture tablets are pressed, biplanar, faceted on both sides and with a dividing notch on one side. Diameter of the tablets: 9 mm.

Example D – Tablet containing 350 mg of active substance

10 Composition:

(1) Active substance	350.0 mg
(2) Lactose	136.0 mg
(3) Maize starch	80.0 mg
15 (4) Polyvinylpyrrolidone	30.0 mg
(5) Magnesium stearate	4.0 mg
	600.0 mg

Preparation:

20 (1), (2) and (3) are mixed together and granulated with an aqueous solution of (4). (5) is added to the dried granulated material. From this mixture tablets are pressed, biplanar, faceted on both sides and with a dividing notch on one side. Diameter of the tablets: 12 mm.

25 Example E – Capsules containing 50 mg of active substance

Composition:

	(1) Active substance	50.0 mg
5	(2) Dried maize starch	58.0 mg
	(3) Powdered lactose	50.0 mg
	(4) Magnesium stearate	2.0 mg
		160.0 mg

10 Preparation:

(1) is triturated with (3). This trituration is added to the mixture of (2) and (4) with vigorous mixing.

This powder mixture is packed into size 3 hard gelatine capsules in a capsule
15 filling machine.

Example F – Capsules containing 350 mg of active substance

Composition:

20

	(1) Active substance	350.0 mg
	(2) Dried maize starch	46.0 mg
	(3) Powdered lactose	30.0 mg
	(4) Magnesium stearate	4.0 mg
25		430.0 mg

Preparation:

(1) is triturated with (3). This trituration is added to the mixture of (2) and (4) with vigorous mixing.

5

This powder mixture is packed into size 0 hard gelatine capsules in a capsule filling machine.

Example G – Suppositories containing 100 mg of active substance

10

1 suppository contains:


Active substance	100.0 mg
Polyethyleneglycol (M.W. 1500)	600.0 mg
Polyethyleneglycol (M.W. 6000)	460.0 mg
15 Polyethylenesorbitan monostearate	840.0 mg
	2,000.0 mg

Preparation:

The polyethyleneglycol is melted together with polyethylenesorbitan
20 monostearate. At 40°C the ground active substance is homogeneously dispersed in the melt. It is cooled to 38°C and poured into slightly chilled suppository moulds.


CLAIMS

1. Substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I)



in an enantiomeric excess of more than 96% at the carbon atom in position 3 of the tetrahydrofuran ring, wherein

D denotes D¹ a substituted bicyclic ring system of formula (II),



wherein

K¹ and K⁴

each independently of one another denote a -CH₂, -CHR^{7a}, -CR^{7b}R^{7c} or a -C(O) group, and

R^{7a}/R^{7b}/R^{7c}

each independently of one another denote a fluorine atom, a hydroxy, C₁₋₅-alkyloxy, amino, C₁₋₅-alkylamino, di-(C₁₋₅-alkyl)-amino, C₃₋₅-cycloalkyleneimino or C₁₋₅-alkylcarbonylamino group,

a C₁₋₅-alkyl group which may be substituted by 1-3 fluorine atoms,

or

two groups R^{7b}/R^{7c} together with the cyclic carbon atom may form a 3, 4, 5-, 6-

5 or 7-membered saturated carbocyclic group

wherein the methylene groups thereof may be substituted by 1-2 C₁₋₃-alkyl or CF₃- groups, and/or

the methylene groups thereof, if they are not bound to a heteroatom, may be substituted by 1-2 fluorine atoms, and

10

K² and K³

each independently of one another denote a -CH₂-, -CHR^{8a}-, -CR^{8b}R^{8c}- or a -C(O)- group, and

15 R^{8a}/R^{8b}/R^{8c}

each independently of one another denote a C₁₋₅-alkyl group which may be substituted by 1-3 fluorine atoms,

or two groups R^{8b}/R^{8c} together with the cyclic carbon atom may form a 3, 4, 5-,

20 6- or 7-membered saturated carbocyclic group, and

in all there should be no more than four groups selected from among R^{7a}, R^{7b}, R^{7c}, R^{8a}, R^{8b} and R^{8c} in formula (II), and

25 X denotes a NR¹ group, wherein

R^1 denotes a hydrogen atom or a hydroxy, C_{1-3} -alkyloxy, amino, C_{1-3} -alkylamino, di- $(C_{1-3}$ -alkyl)-amino, a C_{1-5} -alkyl, C_{2-5} -alkenyl- CH_2 , C_{2-5} -alkynyl- CH_2 or C_{3-6} -cycloalkyl group,

5 wherein the methylene and methyl groups present in the above-mentioned groups

may additionally be substituted by a C_{1-3} -alkyl, carboxy, C_{1-5} -alkoxycarbonyl group or

by a hydroxy, C_{1-5} -alkyloxy, amino, C_{1-5} -alkylamino, C_{1-5} -dialkylamino or C_{4-7} -

10 cycloalkyleneimino group,

provided that O-C-O or O-C-N or N-C-N-bonds are excluded and/or

one to three hydrogen atoms may be replaced by fluorine atoms, provided that the methylene or methyl groups are not directly bound to a nitrogen atom,

15 and wherein

A^1 denotes either N or CR^{10} ,

A^2 denotes either N or CR^{11} ,


20

A^3 denotes either N or CR^{12} ,


wherein R^{10} , R^{11} and R^{12} each independently of one another represent

a hydrogen, fluorine, chlorine, bromine or iodine atom, or a C₁₋₅-alkyl, CF₃, C₂₋₅-alkenyl, C₂₋₅-alkynyl, a cyano, carboxy, C₁₋₅-alkyloxycarbonyl, hydroxy, C₁₋₃-alkyloxy, CF₃O, CHF₂O, CH₂FO, or


- 5 D denotes D² a group of general formula



wherein A denotes A⁴, a group



- 10 or wherein A denotes A⁵, a group of general formula



15

wherein

m is the number 1 or 2,

X^1 denotes a carbonyl, thiocarbonyl, $C=NR^{9c}$, $C=N-OR^{9c}$, $C=N-NO_2$, $C=N-$
5 CN or sulphonyl group,

X^2 denotes an oxygen atom or a $-NR^{9b}$ group,

X^3 denotes a carbonyl, thiocarbonyl, $C=NR^{9c}$, $C=N-OR^{9c}$, $C=N-NO_2$, $C=N-$
10 CN or sulphonyl group,

X^4 denotes an oxygen or sulphur atom or a $-NR^{9c}$ group,

R^{9a} in each case independently of one another denotes a hydrogen or
15 halogen atom or a C_{1-5} -alkyl, hydroxy, hydroxy- C_{1-5} -alkyl, C_{1-5} -alkoxy, C_{1-5} -alkoxy- C_{1-5} -alkyl, amino, C_{1-5} -alkylamino, di- $(C_{1-5}$ -alkyl)-amino, amino- C_{1-5} -alkyl, C_{1-5} -alkylamino- C_{1-5} -alkyl, di- $(C_{1-5}$ -alkyl)-amino- C_{1-5} -alkyl, aminocarbonyl, C_{1-5} -alkylaminocarbonyl, di- $(C_{1-5}$ -alkyl)-aminocarbonyl or C_{1-5} -alkylcarbonylamino group, while
20 in the previously mentioned substituted 5- to 7-membered groups A^5 the heteroatoms F, Cl, Br, I, O or N optionally introduced with R^{9a} as substituent are not separated by precisely one carbon atom from a heteroatom selected from among N, O, S,

R^{9b} each independently of one another denote a hydrogen atom or a C_{1-5} -alkyl group,


R^{9c} each independently of one another denote a hydrogen atom, a C_{1-5} -alkyl, C_{1-5} -alkylcarbonyl, C_{1-5} -alkyloxycarbonyl or C_{1-5} -alkylsulphonyl group,

R^4 denotes a hydrogen or halogen atom, a C_{1-3} -alkyl or C_{1-3} -alkoxy group, while the hydrogen atoms of the C_{1-3} -alkyl or C_{1-3} -alkoxy group may optionally be wholly or partly replaced by fluorine atoms, a C_{2-3} -alkenyl, C_{2-3} -alkynyl, nitrile, nitro or amino group,

R^5 denotes a hydrogen or halogen atom or a C_{1-3} -alkyl group,

R^3 denotes a hydrogen atom or a C_{1-3} -alkyl group, and

M denotes a thiophene ring according to formula (III),



which is bound to the carbonyl group in formula (I) via the 2-position and which is substituted in the 5-position by R^2 and optionally additionally by R^6 , wherein

R^2 denotes

R^{2a} a hydrogen, fluorine or iodine atom, or

R^{2b} a methoxy, C_{1-2} -alkyl, formyl, NH_2CO , or

R^{2c} a chlorine, bromine or ethynyl group,

- 5 R^6 denotes a hydrogen, fluorine, chlorine, bromine or iodine atom or a C_{1-2} -alkyl or amino group,

wherein, unless otherwise stated, by the term "halogen atom" mentioned
hereinbefore in the definitions is meant an atom selected from among fluorine,
10 chlorine, bromine and iodine,


and wherein the alkyl, alkenyl, alkynyl and alkyloxy groups contained in the
previously mentioned definitions which have more than two carbon atoms may,
unless otherwise stated, be straight-chain or branched and the alkyl groups in
15 the previously mentioned dialkylated groups, for example the dialkylamino
groups, may be identical or different,

and the hydrogen atoms of the methyl or ethyl groups contained in the
foregoing definitions, unless otherwise stated, may be wholly or partly replaced
20 by fluorine atoms,

the tautomers, enantiomers, diastereomers, mixtures and salts thereof.

2. Substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in accordance with claim 1 in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein

D denotes a substituted bicyclic ring system of formula (II),



wherein

K^1 and K^4

each independently of one another denote a $-CH_2$, $-CHR^{7a}$, or a $-CR^{7b}R^{7c}$ group, wherein

$R^{7a}/R^{7b}/R^{7c}$

each independently of one another denote a fluorine atom, a hydroxy, methoxy or C_{1-2} -alkyl group which may be substituted by 1-3 fluorine atoms,

wherein the two groups R^{7b}/R^{7c} cannot both simultaneously be bound to the cyclic carbon atom via a heteroatom, except if $-C(R^{7b}R^{7c})-$ corresponds to a $-CF_2$ group, or

two groups R^{7b}/R^{7c} may form, together with the cyclic carbon atom, a 3-, 4- or 5-membered saturated carbocyclic group, and

K^2 and K^3

each independently of one another represent a $-\text{CH}_2$, $-\text{CHR}^{8a}$, or $-\text{CR}^{8b}\text{R}^{8c}$ group, and

5 $\text{R}^{8a}/\text{R}^{8b}/\text{R}^{8c}$

each independently of one another denote a C_{1-2} -alkyl group which may be substituted by 1-3 fluorine atoms,

or two groups $\text{R}^{8b}/\text{R}^{8c}$ may form, together with the cyclic carbon atom, a 3-, 4-,

10 5-membered carbocyclic group, and

in all in formula (II) there should be no more than four groups selected from among R^{7a} , R^{7b} , R^{7c} , R^{8a} , R^{8b} and R^{8c} , and

15

X denotes an NR^1 group, wherein

R^1 denotes a hydrogen atom or a C_{1-2} -alkyl or C_{3-4} -cycloalkyl group,

20 wherein the methylene and methyl groups present in the above-mentioned groups may additionally be substituted by a methyl group ,

and wherein

25 A^1 denotes CR^{10} ,

A^2 denotes CR^{11} ,

A^3 denotes CR^{12} ,

5


wherein R^{10} , R^{11} and R^{12} each independently of one another represent

a hydrogen, fluorine, chlorine, bromine atom or a Methyl, CF_3 , a cyano,


Methoxy, CF_3O , CHF_2O , CH_2FO - group, or

10

D denotes D^2 a group of general formula





wherein A denotes A^4 , a group



15

or wherein A denotes A^5 , a group of general formula





wherein

5

m is the number 1 or 2,

X^1 denotes a carbonyl or C=N-CN group,

10 X^3 denotes a carbonyl or C=N-CN group,

X^4 denotes an oxygen atom,

R^{9a} in each case independently of one another denotes a hydrogen atom or a


15 C_{1-2} -alkyl group, while

R^4 denotes a hydrogen or fluorine, chlorine or bromine atom, a methyl or a methoxy group,

20 R^5 denotes a hydrogen, fluorine or chlorine atom or a methyl group,

R^3 denotes a hydrogen atom and

M denotes a thiophene ring according to formula (III),



which is bound to the carbonyl group in formula (I) via the 2-position and which is substituted in the 5-position by R^2 and optionally additionally by R^6 , wherein

R^2 denotes


R^{2c} a chlorine, bromine atom or an ethynyl group,

R^6 denotes a hydrogen atom,

the tautomers, diastereomers, mixtures and salts thereof.

3. Substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in accordance with claim 1 or 2 in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein

D denotes a substituted bicyclic ring system of formula (II),




wherein K1, K2, K3, K4, X, A1, A2 and A3 are as defined in claim 1 or 2,

the tautomers, diastereomers, mixtures and salts thereof.

4. Substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in accordance with claim 1 or 2 in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein

D denotes D² a group of general formula




wherein R⁴ and R⁵ are as defined in claim 1 or 2,

the tautomers, diastereomers, mixtures and salts thereof.

5. Substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in accordance with claim 1 or 2 in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein

D denotes D² a group of general formula




or wherein A denotes A⁵, wherein A⁵, R⁴ and R⁵ are as defined in claim 1 or 2, the tautomers, diastereomers, mixtures and salts thereof.

6. Substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in accordance with any one of claim 1 to 5 in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein the amino-tetrahydrofuran carboxylic acid amide moiety has the R-configuration.


7. Substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in accordance with any one of claim 1 to 5 in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein the amino-tetrahydrofuran carboxylic acid amide moiety has the S-configuration.

8. A compound in accordance with any one of claims 1 to 5, which is selected from the following list of compounds and mixtures and salts thereof:



(S)-3-[(5-bromo-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide





(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide



- (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5R)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
- 5 (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((5S)-3,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




- 10 (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1R)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide and
- (3S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-((1S)-1,3-dimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




15

(S)-5-chloro-thiophene-2-carboxylic acid-N-{3-[3-methyl-4-(5-cyanimin-1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(5-oxo-
[1,4]oxazepan-4-yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide




5


(S)-5-chloro-thiophene-2-carboxylic acid-*N*-{3-[4-(5-cyanimin-[1,4]oxazepan-4-
yl)-phenylcarbamoyl]-tetrahydrofuran-3-yl}-amide



(S)-5-bromo-thiophene-2-carboxylic acid-*N*-{3-[3-methyl-4-(3-oxo-morpholin-4-
10 yl)-phenylcarbamoyl]-tetrahydro-thiophen-3-yl}-amide




(S)-3-[(5-chloro-thiophen-2-yl)-carbonylamino]-N-(3,5,5-trimethyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-tetrahydrofuran-3-carboxylic acid amide




9. A method for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (I) in accordance with any one of claims 1 to 8 in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring, wherein the enantiomers are separated via chiral column chromatography.


10. A method for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ia) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



comprising reacting a compound of the general formula (IVa)



with a compound of the general formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



optionally further comprising cleaving protecting groups,


wherein K1, K2, K3, K4, X, A1, A2, A3, R2, R3 and R6 are as defined in claim 1 or 2, and

wherein Q is a hydroxy or C₁₋₄-alkyloxy group, a halogen atom or a C₁₋₅-alkyloxycarbonyloxy or acyloxy group.


11. The method according to claim 10, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (V) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ia) have the R-configuration.

12. The method according to claim 10, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (V) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ia) have the S-configuration.


13. A method for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ib) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



comprising reacting a compound of the general formula (IVb)



with a compound of the general formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



optionally further comprising cleaving protecting groups,


wherein A, R₄, R₅, R₂, R₃ and R₆ are as defined in claim 1 or 2 and

wherein Q is a hydroxy or C₁₋₄-alkyloxy group, a halogen atom or a C₁₋₅-alkyloxycarbonyloxy or acyloxy group.

14. The method according to claim 13, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (V) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ib) have the R-configuration.


15. The method according to claim 13, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (V) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ib) have the S-configuration.

16. A method for the preparation of substituted 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ic) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring




comprising the steps of:


a) reacting a compound of the formula (IV) with a compound of the formula (VI) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring,



and subsequently cleaving the protecting group PG to obtain a compound of the formula (VII) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring; and



b) reacting said compound (VII) of step a) with a compound of formula (VIII)



wherein

Q is a hydroxy or C₁₋₄-alkyloxy group, a halogen atom or a C₁₋₅-alkyloxycarbonyloxy or acyloxy group,

PG is a hydrogen atom or a protective group for the amino function, and


D, R₃, R₂ and R₆ are as defined in claim 1 or 2.

17. The method according to claim 16, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (VI) and of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ic) have the R-configuration.

18. The method according to claim 16, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of the compound of the general formula (VI) and

of the 3-amino-tetrahydrofuran-3-carboxylic acid amides of general formula (Ic) have the S-configuration.

19. A method for preparing a compound of the formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring




by enzymatic resolution of a racemic mixture of said compound of the formula (V),

wherein R² and R⁶ are as defined in claim 1 or 2, and

wherein Q is a straight or substituted C₁₋₁₂-alkyloxy group, or allyloxy or substituted allyloxy group, or C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group.

20. A compound of the formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring,




wherein R² and R⁶ are as defined in claim 1 or 2, and

wherein Q is a hydroxy or C₁₋₁₂-alkyloxy group, or allyloxy or substituted allyloxy group, or a halogen atom or a C₁₋₁₂-alkyloxycarbonyloxy or acyloxy group.


21. The compound of the formula (V) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring according to claim 20, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of said compound of the general formula (V) has the S-configuration.

22. A method for preparing a compound of the formula (VII) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



by chemical resolution of a racemic mixture of said compound of the formula (VII) with a chiral acid, wherein D and R³ are as defined in claim 1 or 2.

23. A compound of the formula (VII) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring



wherein D and R₃ are as defined in claim 1 or 2.

24. The compound of the formula (VII) in an enantiomeric excess of more than 96% at the carbon in position 3 of the tetrahydrofuran ring according to claim 23, wherein the amino-tetrahydrofuran carboxylic acid amide moiety of said compound of the general formula (VII) has the S-configuration.