PRESERVATION SOLUTION FOR PRESERVING EXTRACELLULAR VESICLES/EXOSOMES, METHOD FOR PRESERVING THE SAME, AND MIXED SOLUTION THEREOF

Preservation solution, in particular solution formulation and method capable of long-term preservation of extracellular vesicles/exosomes in a frozen environment.

Due to the challenge of preservation and susceptibility to degradation after purification, the extracellular vesicles/exosomes are often purified according to the amount required for each application. Subsequent modification or drug conjugation, if implemented, facilitates to extend the overall process duration and presents challenges to progress towards industrial development.

To overcome the challenges of difficulty in preservation and susceptibility to degradation of the extracellular vesicles/exosomes after purification, the present invention provides a solution for preserving the extracellular vesicles/exosomes comprising: a carrier solvent selected from the group consisting of polysorbate 80, sucrose, and polyethylene glycol 3350/4000 (PEG-3350/4000); and a stabilizer selected from the group consisting of salts, amino acids, and amino acid salts. The pH of the preservation solution is in the range of 5 to 7.4.

Wherein, the salt is a sodium acetate or a sodium chloride, and the weight percent concentration of the salt is in the range of 0.1 to 2 of the total volume of the preservation solution.

Wherein, the amino acids are selected from the group consisting of histidine, arginine, and glycine, and the weight percent concentration of the amino acid is in the range of 0.5 to 3 of the total volume of the preservation solution.

Wherein, the amino acid salts are selected from the group consisting of histidine hydrochloride, arginine hydrochloride, and glycine hydrochloride, and the weight percent concentration of the amino acid salt is in the range of 0.05 to 0.25 of the total volume of the preservation solution.

Wherein, the weight percent concentration of polysorbate 80 is in the range of 0.01 to 0.03 of the total volume of the preservation solution; the weight percent concentration of sucrose is in the range of 5 to 10 of the total volume of the preservation solution; and the

weight percent concentration of polyethylene glycol 3350/4000 (PEG-3350/4000) is in the range of 1 to 5 of the total volume of the preservation solution.

The present invention also provides a method for preserving the extracellular vesicles/exosomes. The steps include:

adding the extracellular vesicles/exosomes to the preservation solution to form a mixed preservation solution; and

gradually cooling the mixed preservation solution and placing it in a frozen preservation environment.

Wherein, the temperature of the frozen preservation environment may be -20 degrees Celsius or -80 degrees Celsius.

Wherein, after forming the mixed preservation solution, drying to form a dried preservation sample, and then placing the dried preservation sample in the frozen preservation environment.

The present invention further provides a mixed solution, comprising the preservation solution and the extracellular vesicles/exosomes.

Wherein, the extracellular vesicles/exosomes in the mixed preservation solution are in the range of 10^{11} to 10^{13} particles per milliliter.

Wherein, the extracellular vesicles/exosomes are modified vesicles carrying a specific protein.

Wherein, the extracellular vesicles/exosomes are loaded with a drug.

The present invention utilizes the selected carrier solvent and the stabilizer in the preservation solution to configure appropriate preservation conditions and pH according to the different types of extracellular vesicles/exosomes and to enable long-term preservation under frozen conditions.

Additionally, the preservation solution can preserve the extracellular vesicles/exosomes in various preservation environments, which contributes to the commercial development of the product.

Through the configuration of the preservative solution, the present invention enables frozen preservation based on specific liquid or dry preservation forms as needed to achieve long-term preservation of the extracellular vesicles/exosomes. Upon thawing and application, a certain reaction efficiency can be maintained, which contributes to research and application in the medical industry, and enhances time cost and commercial value.

biological activity of the extracellular vesicles, consistent with the above experimental results.

Furthermore, in FIG. 7, a comparative analysis of the size of the extracellular vesicles after dry preservation is performed between Comparative Example 2 and Embodiments 1, 2, and 3. This ensures that the physiological structure of the extracellular vesicles is not affected by the preservation solution and cryopreservation method. As observed in the Figure, after cryopreservation of both Comparative Example 2 and each Embodiment, there is no significant difference in size between the extracellular vesicles. However, the concentration of extracellular vesicles decreases significantly after dry cryopreservation in Comparative Example 2.

In Experiment 3, referring FIG. 8, the present invention then subjected the extracellular vesicles to a drug-carrying process after thawing. By drug testing to confirm that the extracellular vesicles still possess the functionality to carry the drug after dry cryopreservation via the preservation solution.

In this embodiment, Doxorubicin (DOX) is used as the drug. The drug-carrying efficiency was evaluated by detecting the reflected wavelength of the drug to verify whether the extracellular vesicles were carrying the drug. A better drug-carrying efficiency indicates a higher proportion of drug loaded by the extracellular vesicles. The extracellular vesicles with the drug showed more distinct reflected wavelengths between 400 and 550 nm, indicating a higher efficiency in carrying the drug.

In the drug-carrying process provided by the present invention, a cell count of 1011 extracellular vesicles is initially added to the solution having a concentration of 2 mg/mL of Doxorubicin (DOX) and allowed to stand and react. This results in the extracellular vesicles carrying the drug. The uncoated DOX is then removed by centrifugation, leaving the extracellular vesicles ready for further testing.

In this experiment, different groups are established: Control Group 1 with only the drug, Control Group 2 with the extracellular vesicles without drug carrying, Control Group 3 with the drug-loaded extracellular vesicles, and Experiment Group 1 with the drug-loaded extracellular vesicles subjected to cryopreservation in the form of dry preservation. Wherein, Control Group 3 uses the formula provided in Embodiment 1 as the medium for drug carrying of the extracellular vesicles. Similarly, Experiment Group 1 uses the formula provided in Embodiment 1 as the preservation solution to perform the drug-carrying

process prior to cryopreservation. The results show that the drug- loaded extracellular vesicles have a lower drug content after dry preservation.

To demonstrate the effectiveness of Experiment Group 1 after the drug-carrying process, the cryopreservation, and the subsequent reconstitution, the present invention further conducted a test to evaluate the therapeutic effectiveness of the extracellular vesicles carrying the drug via the cryopreservation with the preservation solution.

In Experiment 4, the extracellular vesicles are first subjected to the drug-carrying process and then frozen with the preservation solution (dry preservation form). After thawing, the drug-loaded extracellular vesicles (Doxorubicin) are tested for cytotoxicity against human breast cancer cells to verify the killing effect of the drug-loaded extracellular vesicles.

Referring to FIG. 9, the drug-loaded extracellular vesicles (Control Group 3) and the drug-loaded extracellular vesicles after dry cryopreservation (Experiment Group 1) are separately added to the culture medium of the human breast cancer cells for reaction. Then, the cell viability rate (%) is then measured at 12 hours, 24 hours, and 48 hours before and after the reaction to compare the cytotoxic effects of Control Group 3 and Experiment Group 1. The results indicated that the cytotoxic effect of the drug-loaded extracellular vesicles after cryopreservation is lower than that of the group without cryopreservation, but it still possessed therapeutic efficacy in drug delivery.

In Experiment 5, the extracellular vesicles are taken out after cryopreservation (dry preservation form) and subjected to the drug-carrying process after thawing. Likewise, the cytotoxicity test against the human breast cancer cells is conducted to verify the killing effect of the drug-loaded extracellular vesicles (Doxorubicin).

In FIG. 10, the present invention first uses the group in which the extracellular vesicles previously preserved with cryopreservation in the form of dry preservation are thawed and then combined with the drug (DOX) as the Experiment Group 4, the group in which the extracellular vesicles are not subjected to cryopreservation and combined with the drug as the Control Group 4, and both groups are subjected to the cytotoxicity test against the human breast cancer cells. The results indicate that there is no significant difference in the cytotoxic effect against the human breast cancer cells between the groups where the extracellular vesicles are preserved with cryopreservation and then combined with the drug

after thawing (Experimental Group 4) and Control Group 4. Both groups exhibit excellent cytotoxic effects.

It is noteworthy that compared to the results in FIG. 9, an optimal procedure for preserving the extracellular vesicles using the preservation solution provided in the present invention and performing the drug carrying can be established.

In Experiment 6, as shown in FIG. 11, the preservation solution provided by the present invention is further used to compare the preservation effects of the extracellular vesicles in different preservation environments and to verify the temporal effectiveness of the preservation. The extracellular vesicles are placed in the preservation solution and preserved in different temperature environments, including -80 degrees Celsius, -20 degrees Celsius, 4 degrees Celsius, room temperature, and 40 degrees Celsius. The concentrations of the extracellular vesicles prior to preservation are used as a reference, and the residual concentration percentages of the extracellular vesicles are evaluated for each group after 1, 2, 4, 8, 16, and 26 weeks of preservation.

The results showed the excellent preservation effects of the extracellular vesicles in the freezing environment (at -80°C and -20°C) using the provided preservation solution. The preservation effect could even be extended to over six months (26 weeks), with the residual concentration of the extracellular vesicles reaching 80%. As the preservation temperature increased, the preservation effect of the extracellular vesicles decreased with longer preservation times.

FIGS. 11 and 12 further demonstrate the drug (Doxorubicin) carrying process of the extracellular vesicles preserved in the freezing environment for 12 and 14 weeks using the provided preservation solution. The cytotoxicity tests are also conducted against the human breast cancer cells to verify the efficiency of carrying the drug over an extended preservation period and the killing effect of the extracellular vesicles.

Wherein, the group which directly treats the human breast cancer cells with the drug serves as Control Group 5; the group which treats the human breast cancer cells with the extracellular vesicles without carrying the drug serves as Control Group 6; and the group which does not undergo cryopreservation and carries the drug in the extracellular vesicles serves as Control Group 7. The group of extracellular vesicles loaded with the drug after cryopreservation at -20 degrees Celsius serves as Experiment Group 5, while those loaded with the drug after cryopreservation at -80 degrees Celsius serves as Experiment Group 6.

The human breast cancer cells are co-cultured with each group for the cytotoxicity test. The cell counts are measured on days 1, 2, and 3 and compared to the total cell count of the human breast cancer cells prior to the cytotoxicity test to calculate the cytotoxic death rate (%).

The results showed a significant cytotoxic effect in Experiment Groups 5 and 6 on the day 1. With increasing co-culture time, the cytotoxic death rate (%) of the human breast cancer cells in Experiment Groups 5 and 6 continues to increase. On the days 2 and 3, Experiment Group 6 exhibits cytotoxic results similar to Control Group 7, demonstrating that the preservation solution provided by the present invention can not only preserve the extracellular vesicles/exosomes for a longer period of time in the freezing environment, but also maintain the biological activity of the extracellular vesicles/exosomes.

The present invention discloses that the provided preservation solution, utilizing the selected carrier solvent and stabilizer, can be tailored according to the specific type of extracellular vesicles/exosomes to suit different preservation conditions and pH levels and enable long-term preservation under the freezing environments.

Moreover, the preservation solution can be used to preserve the extracellular vesicles/exosomes in various preservation environments, which contributes to the commercial development of the product.

Through the configuration of the preservation solution, the present invention allows for cryopreservation in liquid or dry form according to specific needs. This allows for long-term preservation of the extracellular vesicles/exosomes while still maintaining the certain reaction efficiency upon thawing for applications, contributing to research and applications in the medical industry while enhancing time efficiency and commercial value.

Claims

1. A use of a preservation solution for preserving extracellular vesicles/exosomes, the solution consisting of:

a carrier consisting of polysorbate 80, wherein the weight percent concentration of polysorbate 80 is 0.01 of the total volume of the preservation solution; and

a stabilizer selected from the group consisting of salts, amino acids, and amino acid salts;

wherein the pH of the preservation solution is in the range of 5 to 7.4

whereby the preservation solution is used in preserving extracellular vesicles/exosomes in a frozen preservation environment at a preservation time up to at least 4 weeks, with a residual concentration of the extracellular vesicles/exosomes reaching 80%.

2. The use of the preservation solution according to claim 1, wherein the salt is a sodium acetate or a sodium chloride.

3. The use of the preservation solution according to claim 1, wherein the amino acids are selected from the group consisting of histidine, arginine, and glycine, and the weight percent concentration of the amino acid is in the range of 0.5 to 3 of the total volume of the preservation solution.

4. The use of the preservation solution according to claim 1, wherein the amino acid salts are selected from the group consisting of histidine hydrochloride, arginine hydrochloride, and glycine hydrochloride, and the weight percent concentration of the amino acid is in the range of 0.05 to 0.25 of the total volume of the preservation solution.

5. A method for preserving extracellular vesicles/exosomes, using the preservation solution according to any one of claims 1 to 4, the steps including:

adding the extracellular vesicles/exosomes to the preservation solution to form a mixed preservation solution; and

gradually cooling the mixed preservation solution and placing the mixed preservation solution in a frozen preservation environment for preservation.

6. The preservation method according to claim 5, wherein after mixing the extracellular vesicles/exosomes with the preservation solution, and gradually cooling the mixed preservation solution, first drying to form a dried preservation sample, and then placing the dried preservation sample in the frozen preservation environment for preservation.

7. A mixed solution comprising the preservation solution according to any one of claims 1 to 4 and extracellular vesicles/exosomes.

8. The mixed solution according to claim 7, wherein the extracellular vesicles/exosomes in the mixed preservation solution are in the range of 10^{11} to 10^{13} particles per milliliter.

9. The mixed solution according to claim 7, wherein the extracellular vesicles/exosomes are modified extracellular vesicles carrying a specific protein.

10. The mixed solution according to any one of claims 7 to 9, wherein the extracellular vesicles/exosomes are loaded with a drug.