

PHOSPHORAMIDATE, AND MONO-, DI-, AND TRI-PHOSPHATE ESTERS OF (1R, CIS)-4-(6-AMINO-9H-PURIN-9-YL)-2-CYCLOPENTENE-1-METHANOL AS ANTIVIRAL AGENTS

5

FIELD OF THE INVENTION

The present invention relates to certain analogs of (1R, cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol and their use in medical therapy.

10



G00002

BACKGROUND OF THE INVENTION

Retroviruses form a sub-group of RNA viruses which, in order to replicate, must first "reverse transcribe" the RNA of their genome into DNA ("transcription" conventionally describes the synthesis of RNA from DNA). Once in the form of DNA, the viral genome may be incorporated into the host cell genome, allowing it to take advantage of the host cell's transcription/translation machinery for the purposes of replication. Once incorporated, the viral DNA is virtually indistinguishable from the host's DNA and, in this state, the virus may persist for the life of the cell.

A species of retrovirus, the Human immunodeficiency virus (HIV) has been reproducibly isolated from patients with AIDS (acquired immunodeficiency syndrome) or with the symptoms that frequently precede AIDS. AIDS is an immunosuppressive or immunodestructive disease that predisposes subjects to fatal opportunistic infections. Characteristically, AIDS is associated with a progressive depletion of T-cells, especially the helper-inducer subset bearing the CD4 surface marker. HIV is cytopathic and appears to preferentially infect and destroy T-cells bearing the CD4 marker, and it is now generally recognized that HIV is the etiological agent of AIDS. Clinical conditions such as AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), Kaposi's sarcoma, thrombocytopenic purpura, AIDS-related neurological conditions, such as AIDS dementia complex, multiple sclerosis or tropical paraparesis, and also anti-HIV antibody-positive and HIV-positive conditions, including such conditions in asymptomatic patients, are also

30



166166

conditions which may be treated by appropriate anti-viral therapy.


- Another RNA virus which has been recognized as the causative agent of an increasingly serious international health problem is the non-A, non-B hepatitis virus. At least 80% of cases of chronic post-transfusional non-A, non-B hepatitis have been shown to be due to the virus now identified as hepatitis C and this virus probably accounts for virtually all cases of post-transfusional hepatitis in clinical settings where blood products are screened for hepatitis B. Whereas approximately half of the cases of acute hepatitis C infection resolve spontaneously over a period of months, the remainder become chronic and in many if not all such cases chronic active hepatitis ensues with the potential for cirrhosis and hepatocellular carcinoma. The structure of the hepatitis C virus genome has been elucidated and the virus has been characterized as a single stranded RNA virus with similarities to flaviviruses.
- Hepatitis B virus (HBV) is a small DNA containing virus which infects humans. It is a member of the class of closely related viruses known as the hepadnaviruses, each member of which selectively infects either mammalian or avian hosts, such as the woodchuck and the duck. Recent insights into the mechanism of replication of the hepadnavirus genome indicate the importance of reverse transcription of an RNA intermediate, suggesting that the reverse transcriptase is a logical chemotherapeutic target. HBV is a viral pathogen of major worldwide importance. The virus is etiologically associated with primary hepatocellular carcinoma and is thought to cause 80% of the world's liver cancer. Clinical effects of infection with HBV range from headache, fever, malaise, nausea, vomiting, anorexia and abdominal pains. Replication of the virus is usually controlled by the immune response, with a course of recovery lasting weeks or months in humans, but infection may be more severe leading to persistent chronic liver disease outlined above.
- U.S. Patent No. 4,916,224 discloses 2',3'-dideoxy-2',3'-didehydro-carbocyclic nucleosides and their use in the treatment of HIV. WO 96/29336 discloses masked monophosphate nucleoside analogues for the treatment of HIV. Wang et al. (*Bioorganic & Medicinal Chemistry Letters* 8, pp. 1585-1588, 1998) disclose the synthesis of L-carbocyclic 2',3'-didehydro-2',3'-dideoxyadenosine and its use in HIV infections.

It has now been discovered that certain phosphoramidates of (1R, cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol are useful for the treatment of viral infections, particularly hepatitis B and retroviral infections, especially HIV. Compounds of the present invention have pharmacokinetic properties which render them advantageous as therapeutic agents.

SUMMARY OF THE INVENTION

10

The present invention relates to compounds of formula (I)



15

(I)

wherein:

- 18 R¹ is hydrogen; C₆₋₁₄aryl; or heteroaryl, optionally substituted with one or more
 20 substituents selected from the group consisting of C₁₋₆alkoxy, nitro, halogen, amino, hydroxy, carboxylate and esters thereof, carboxyalkyl, -CONHR⁶, and -CONR⁶R⁷, wherein R⁶ and R⁷, which may be the same or different, are independently selected from C₁₋₈alkyl, C₁₋₈alkylaryl or C₆₋₁₄aryl;
 25 R² and R³ are independently selected from hydrogen or C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, C₆₋₁₄aryl, or aralkyl wherein each C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, C₆₋₁₄aryl or aralkyl may be optionally substituted with one or more

substituents selected from the group consisting of C₁₋₈alkyl, halo, hydroxy, alkoxy, amino, aminoalkyl, aminodialkyl, -SH, thioalkyl, heterocycle, carboxylate and esters thereof, carboxyalkyl, -CONHR⁶, and -CONR⁶R⁷, wherein R⁶ and R⁷, which may be the same or different, are independently selected from C₁₋₈alkyl, C₁₋₈alkylaryl or C₆₋₁₄aryl; or R² and R³ can together form a 3 to 8-membered ring;

R⁴ is -OR⁸, -NR⁸R⁹ or -SR⁸, where R⁸ and R⁹, which may be the same or different, are independently selected from hydrogen; or C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, aralkyl, heteroaryl, or C₆₋₁₄aryl wherein each C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, aralkyl, heteroaryl, or C₆₋₁₄aryl may be optionally substituted with one or more substituents selected from the group consisting of halo, hydroxy, alkoxy, amino, aminoalkyl, aminodialkyl, -SH, thioalkyl, carboxylate and esters thereof, carboxyalkyl, -CONHR⁶, and -CONR⁶R⁷, wherein R⁶ and R⁷, which may be the same or different, are independently selected from C₁₋₈alkyl, C₁₋₈alkylaryl or C₆₋₁₄aryl;


R⁵ is hydrogen; C₁₋₈alkyl; or C₆₋₁₄aryl; or R² and R⁵ may together form a 5- or 6-membered ring or R³ and R⁵ may together form a 5- or 6-membered ring;

or a pharmaceutically acceptable derivative thereof, and their use in the treatment of viral infections.

DETAILED DESCRIPTION OF THE INVENTION

25

The present invention features compounds of formula (I)



(I)

wherein:

R^1 is hydrogen; C_{6-14} aryl; or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of C_{1-6} alkoxy, nitro, halogen, amino, hydroxy, carboxylate and esters thereof, carboxyalkyl, $-\text{CONHR}^6$, and $-\text{CONR}^6\text{R}^7$, wherein R^6 and R^7 , which may be the same or different, are independently selected from C_{1-6} alkyl, C_{1-6} alkylaryl or C_{6-14} aryl;


R^2 and R^3 are independently selected from hydrogen or C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{5-6} cycloalkenyl, C_{6-14} aryl, or aralkyl wherein each C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{5-6} cycloalkenyl, C_{6-14} aryl or aralkyl may be optionally substituted with one or more substituents selected from the group consisting of C_{1-6} alkyl, halo, hydroxy, alkoxy, amino, aminoalkyl, aminodialkyl, $-\text{SH}$, thioalkyl, heterocycle, carboxylate and esters thereof, carboxyalkyl, $-\text{CONHR}^6$, and $-\text{CONR}^6\text{R}^7$, wherein R^6 and R^7 , which may be the same or different, are independently selected from C_{1-6} alkyl, C_{1-6} alkylaryl or C_{6-14} aryl; or R^2 and R^3 can together form a 3 to 8-membered ring;

R^4 is $-\text{OR}^8$, $-\text{NR}^8\text{R}^9$ or $-\text{SR}^8$, where R^8 and R^9 , which may be the same or different, are independently selected from hydrogen; or C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{5-6} cycloalkenyl, aralkyl, heteroaryl, or C_{6-14} aryl wherein each C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{2-6} alkenyl, C_{5-6} cycloalkenyl, aralkyl, heteroaryl, or C_{6-14} aryl may be optionally substituted with one or more substituents selected from the group consisting of halo, hydroxy, alkoxy, amino, aminoalkyl, aminodialkyl, $-\text{SH}$, thioalkyl, carboxylate and esters thereof, carboxyalkyl, $-\text{CONHR}^6$, and $-\text{CONR}^6\text{R}^7$, wherein R^6 and R^7 , which may be the same or different, are independently selected from C_{1-6} alkyl, C_{1-6} alkylaryl or C_{6-14} aryl;

R^5 is hydrogen; C_{1-6} alkyl; or C_{6-14} aryl; or R^2 and R^5 may together form a 5- or 6-membered ring or R^3 and R^5 may together form a 5- or 6-membered ring;

or a pharmaceutically acceptable derivative thereof, and their use in the treatment of viral infections.

An embodiment of the present invention features compounds of formula (II)



(II)

wherein:

- 5 R¹ is hydrogen; C₆₋₁₄aryl; or heteroaryl, optionally substituted with one or more substituents selected from C₁₋₆alkoxy, nitro, halogen, amino, carboxylate and hydroxy.


R² and R³ are independently selected from hydrogen, C₁₋₆alkyl, C₃₋₈cycloalkyl, C₆₋₁₄aryl, or aralkyl;

10

R⁴ is OR¹⁰, NHR¹⁰ or SR¹⁰, where R¹⁰ is hydrogen, C₁₋₆alkyl, C₃₋₈cycloalkyl, or aralkyl; or NHR¹¹ wherein R¹¹ is C₁₋₆alkyl, aralkyl, or C₆₋₁₄aryl;

15 or a pharmaceutically acceptable derivative thereof, and their use in the treatment of viral infections.

A further aspect of the present invention features a compound of formula (III)




(III)

20

wherein R¹² is

23 JUL 2003



wherein n is 0, 1, or 2, and wherein R¹² is optionally substituted by C₆₋₁₄aryl.

The compounds of the present invention include diastereomers differing in the absolute configuration at phosphorus. Diastereomers may be present as a single isomer or as mixtures of diastereomers.

The term "alkyl" refers to a straight-chain or branched-chain saturated aliphatic hydrocarbon radical containing the specified number of carbon atoms, or where no number is specified, preferably from 1 to about 10, more preferably from 1 to about 8 carbon atoms. Examples of alkyl radicals include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, n-hexyl and the like.

The term "alkenyl," alone or in combination with any other term, refers to a straight-chain or branched-chain mono- or poly-unsaturated aliphatic hydrocarbon radical containing the specified number of carbon atoms, or where no number is specified, preferably from 2- 10 carbon atoms and more preferably, from 2-6 carbon atoms. Examples of alkenyl radicals include, but are not limited to, ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, pentenyl, hexenyl, hexadienyl and the like.

The term "alkoxy" refers to an alkyl ether radical, wherein the term "alkyl" is defined above. Examples of suitable alkyl ether radicals include, but are not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy and the like, with methoxy being preferred.

The term "halo" or "halogen" refers to a radical of fluorine, chlorine, bromine or iodine.

The term "aryl" refers to a carbocyclic aromatic radical (such as phenyl or naphthyl) containing the specified number of carbon atoms, preferably from 6-14 carbon atoms, and

more preferably from 6-10 carbon atoms, optionally substituted with one or more substituents selected from C₁₋₆ alkoxy (for example, methoxy), nitro, halogen (for example chloro), amino, carboxylate and hydroxy. Examples of aryl radicals include, but are not limited to phenyl, naphthyl, indenyl, indanyl, azulenyl, fluorenyl, anthracenyl and the like.

5

The term "heterocycle", alone or in combination with another term, refers to a stable 3-7 membered monocyclic heterocyclic ring or 8-11 membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which may be optionally benzofused if monocyclic. Each heterocycle consists of one or more carbon atoms and from one to four heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. As used herein, the terms "nitrogen and sulfur heteroatoms" include any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen. A heterocyclyl radical may be attached at any endocyclic carbon or heteroatom which results in the creation of a stable structure. Preferred heterocycles include 5-7 membered monocyclic heterocycles and 8-10 membered bicyclic heterocycles. Examples of such groups include imidazolyl, imidazolinoyl, imidazolidinyl, quinolyl, isoquinolyl, indolyl, indazolyl, indazolinolyl, perhydropyridazyl, pyridazyl, pyridyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazinyl, quinoxolyl, piperidinyl, pyranyl, pyrazolinyl, piperazinyl, pyrimidinyl, pyridazinyl, morpholinyl, thiamorpholinyl, furyl, thienyl, triazolyl, thiazolyl, carbolinyl, tetrazolyl, thiazolidinyl, benzofuranoyl, thiamorpholinyl sulfone, oxazolyl, benzoxazolyl, oxopiperidinyl, oxopyrrolidinyl, oxoazepinyl, azepinyl, isoxozolyl, isothiazolyl, furazanyl, tetrahydropyranyl, tetrahydrofuranlyl, thiazolyl, thiadiazolyl, dioxolyl, dioxinyl, oxathioly, benzodioxolyl, dithiolyl, thiophenyl, tetrahydrothiophenyl, sulfolanyl, dioxanyl, dioxolanyl, tetrahydrofurodihydrofuranlyl, tetrahydropyranodihydrofuranlyl, dihydropyranlyl, tetrahydrofurofuranlyl and tetrahydropyranofuranlyl.

25

The term "pharmaceutically acceptable derivative", as used herein, means any pharmaceutically acceptable salt, ester, salt of an ester, or other derivative of a compound of this invention which, upon administration to a recipient, is capable of providing (directly or indirectly) a compound of this invention or an inhibitorily active metabolite or residue thereof. Particularly favored derivatives and prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered


30

to a mammal (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological compartment (e.g., the brain or lymphatic system) relative to the parent species.

- 5 Compounds of formula (I), (II), and (III) and their pharmaceutically acceptable derivatives may hereinafter be referred to as compounds according to the invention.



Preferred compounds of formulae (I) and (II) include the compounds listed in Table 1.

Table 1.



10

Example	R ¹	R ²	R ³	R ⁴	R ⁵
2	Ph	H	Me	OMe	H
3	Ph	H	Me	OEt	H
4	Ph	Me	H	OMe	H
5	Ph	H	Me	OCH ₂ Ph	H
6	Ph	H	CH ₂ Ph	OMe	H
7	Ph	H	CH ₂ Ph	OEt	H
8	H	H	CH ₂ Ph	OH	H
9	Ph	Me	Me	OMe	H
10	Ph	H	CH ₂ Ph(4-O-t-Bu)	OMe	H
11	Ph	H	Me	O-i-Pr	H
12	Ph	H	Me	O-t-Bu	H
13	Ph	H	Me	OCH ₂ -t-Bu	H
14	Ph	H	Me	OCH ₂ - 	H
15	Ph	H	CH ₂ CHMe ₂	OMe	H
16	Ph	H	CH ₂ Ph(4-OMe)	OMe	H
17	Ph	CH ₂ Ph	H	OMe	H

Example	R ¹	R ²	R ³	R ⁴	R ⁵
18	Ph	CH ₂ CHMe ₂	H	OMe	H
19	Ph		H	OMe	H
20	Ph		H	OMe	H
21	Ph	CHMe ₂	H	OMe	H
22	Ph	CH ₂ CO ₂ Me	H	OMe	H
23	Ph	H	R ³ /R ⁵ = (CH ₂) ₃	OMe	-
24	Ph	H	CH(Me)Et	OMe	H
25	Ph	H	n-Pr	OMe	H
25	Ph	H	CH ₂ Ph	O-i-Pr	H
27	Ph	H	CH ₂ Ph	O-t-Bu	H
28	Ph	H	CH ₂ Ph	OCH ₂ -t-Bu	H
29	Ph	H	CH ₂ Ph	OCH ₂ Ph	H
30	(2-CO ₂ Me)Ph	H	Me	OMe	H
31	(4-PhCOCH=CH)Ph	H	Me	OMe	H
32	Ph	H	H	OMe	H
33	Ph	H	i-Pr	OMe	H
34	Ph	H	CH ₂ CH ₂ SMe	OMe	H
35	Ph	Me	Me	OEt	H
36	Ph	Me	Me	O-i-Pr	H
37	Ph	Me	Me	OCH ₂ -t-Bu	H
38	Ph	-	R ² /R ³ = (CH ₂) ₄	OMe	H
39	Ph	-	R ² /R ³ = (CH ₂) ₅	OMe	H
40	Ph	H	CH ₂ CO ₂ Me	OMe	H
41	(4-Cl)Ph	H	Me	OMe	H
42	H	Me	H	NHMe	H

Preferred compounds of the present invention include

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-alaninyl)phosphoramidate;

5 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-alaninyl)phosphoramidate;

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-

- alaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl benzyloxy-L-alaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-phenylalaninyl)phosphoramidate;
 5 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-phenylalaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(L-phenylalaninyl)phosphoramidate Disodium Salt;
 10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy- α , α ,-dimethylglycinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-L-(O-*tert*-butyltyrosinyl)]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy-L-alaninyl)phosphoramidate;
 15 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl *tert*-butoxy)-L-alaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy)-L-alaninyl]phosphoramidate;
 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl cyclopropylmethoxy-L-alaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-leucinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-L-(O-methyltyrosinyl)]phosphoramidate;
 25 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-phenylalaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-leucinyl)phosphoramidate;
 30 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-D-tryptophanyl]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-D-

- tryptophanyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-
 valinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl dimethoxy-D-
 5 aspartyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-
 prolinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-
 isoleucinyl)phosphoramidate;
 10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-
 norvalinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy-L-
 phenylalaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl *tert*-butoxy-L-
 15 phenylalaninyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-
 dimethylpropoxy)-L-phenylalaninyl]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl benzyloxy-L-
 phenylalaninyl)phosphoramidate;
 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[(2-
 carbomethoxy)phenyl methoxy-L-alaninyl]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[4-(3-oxo-3-
 phenylpropenyl)phenyl methoxy-L-alaninyl]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl
 25 methoxyglycinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-
 valinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(methoxy)-L-
 methioninyl]phosphoramidate;
 30 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy- α , α ,-
 dimethylglycinyl)phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy-

- α, α -dimethylglyciny]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy)- α, α -dimethylglyciny]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxycyclopentaneglyciny]phosphoramidate;
 5 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxycyclohexaneglyciny]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl dimethoxy-L-aspartyl)phosphoramidate;
 10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[(4-chloro)phenyl methoxy-L-alaniny]phosphoramidate;
 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[L-(N-methyl)amino)alaniny]phosphoramidate sodium salt;
 and pharmaceutically acceptable derivatives thereof.

15

A further aspect of the present invention features a compound of formula (II) wherein R^1 is H or C_{6-14} aryl, R^2 is C_{1-6} alkyl or aralkyl, R^3 is hydrogen, C_{1-6} alkyl or aralkyl and R^4 is OR^{10} wherein R^{10} is C_{1-6} alkyl or C_{3-8} cycloalkyl.

- 20 A preferred aspect of the present invention features a compound of formula (II) wherein R^1 is C_{6-14} aryl, R^2 is methyl, R^3 is hydrogen, and R^4 is OR^{10} where R^{10} is methyl or ethyl. More preferably, R^1 is phenyl.

- In another aspect of the present invention there is provided compounds of formula (II)
 25 wherein R^1 is hydrogen.

In another aspect of the present invention there is provided compounds of formula (I) and (II) wherein R^2 and R^3 are not both hydrogen.

- 30 When R^2 and R^3 are different, the L-configuration of naturally occurring amino acids is preferred.

More preferred compounds of the present invention include

- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-
 - 5 alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl benzyloxy-L-
 - 10 alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-phenylalaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-phenylalaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(L-phenylalaninyl)
 - 15 phosphoramidate Disodium Salt;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy- α , α ,-dimethylglycyl)phosphoramidate;
 - and pharmaceutically acceptable derivatives thereof.
- 20 Pharmaceutically acceptable salts of the compounds of the present invention include salts of a basic or acidic portion of the molecule. Salts of a basic moiety may be formed by organic carboxylic acids such as acetic, lactic, tartaric, malic, isethionic, lactobionic, and succinic acids, organic sulphonic acids, such as methanesulphonic, ethanesulphonic, benzenesulphonic and p-toluenesulphonic acids and inorganic acids, such as hydrochloric,
- 25 sulphuric, phosphoric and sulphamic acids. Salts of an acidic moiety may be formed by an appropriate base, such as an alkali metal (for example, sodium), an alkaline earth (for example, magnesium, calcium), ammonium and ammonium salts.

- Preferred esters of the compounds according to the invention may be independently
- 30 selected from the following groups: (1) carboxylic acid esters obtained by esterification of the hydroxy groups, in which the non-carbonyl moiety of the carboxylic acid portion of the ester grouping is selected from straight or branched chain alkyl (for example, acetyl, n-


propyl, t-butyl, or n-butyl), alkoxyalkyl (for example, methoxymethyl), aralkyl (for example, benzyl), aryloxyalkyl (for example, phenoxymethyl), aryl (for example, phenyl optionally substituted by, for example, halogen, C₁₋₄alkyl, or C₁₋₄alkoxy or amino); (2) sulfonate esters, such as alkyl- or aralkylsulfonyl (for example, methanesulfonyl); (3) amino acid esters (for example, L-valyl or L-isoleucyl); (4) phosphonate esters and (5) mono-, di- or triphosphate esters. The phosphate esters may be further esterified by, for example, a C₁₋₂₀ alcohol or reactive derivative thereof, or by a 2,3-di (C₆₋₂₄)acyl glycerol.

In such esters, unless otherwise specified, any alkyl moiety present advantageously contains from 1 to 18 carbon atoms, particularly from 1 to 6 carbon atoms, more particularly from 1 to 4 carbon atoms. Any cycloalkyl moiety present in such esters advantageously contains from 3 to 6 carbon atoms. Any aryl moiety present in such esters advantageously comprises a phenyl group.


Esters of carboxylate may include alkyl, cycloalkyl, aralkyl, and aryl esters.

Compounds of formula (I) and (II) may be made by modifications of the procedures described in Biochem. Biophys. Res. Commun. 225:363-369, 1997.

The present invention further includes a process for the preparation of a compound of formula (I) which comprises reaction of a compound of formula (IV)



with a compound of formula (VI)



(VI)

wherein R¹ – R⁵ are as hereinbefore defined for formula (I).

5


The reaction may be carried out in pyridine, pyridine-tetrahydrofuran or acetonitrile in the presence of t-butyl magnesium chloride (Balzarini et al., *Biochem. Biophys. Res. Comm.* 225:363-369 (1996). The phosphochloridate intermediates, compounds of formula (VI), may be prepared according to WO 96/29336, incorporated herein by reference hereto;

10 McGuigan et al, *J. Med. Chem.*, 1996, 39, 1748-1753; and McGuigan et al, *Antiviral Res.*, 1997, 35, 195-204.

Compounds of formula (IV) may be made according to Example 1 or by any method known in the art.


15

The present invention further includes a process for the preparation of a compound of formula (II) which comprises reaction of a compound of formula (IV)



(IV)

20 with a compound of formula (V)



(V)

5 wherein R¹ – R⁴ are as hereinbefore defined for formula (II).

The reaction may be carried out in pyridine, pyridine-tetrahydrofuran or acetonitrile in the presence of t-butyl magnesium chloride (Balzarini et al., *Biochem. Biophys. Res. Comm.* 225:363-369 (1996). The phosphochloridate intermediates, compounds of formula (V),
 10 may be prepared according to WO 96/29336, incorporated herein by reference hereto; McGuigan et al, *J. Med. Chem.*, 1996, 39, 1748-1753; and McGuigan et al, *Antiviral Res.*, 1997, 35, 195-204.

O-monophosphate compounds of formula (III) may be prepared by treating a compound of
 15 formula (IV) with an appropriate phosphorylating agent, e.g. phosphoryl chloride as in M. Yoshikawa, T. Kato and T. Takenishi, *Bulletin Chem. Soc. Japan*, 1969, 42, 3505. The corresponding O-di and O-triphosphates may be prepared by the method of N. C. Mishra and A. D. Broom, *J. Chem. Soc., Chem. Commun.*, 1991, 1276 or by the methods described in "Nucleotide Analogs" K.H. Sheit, John Wiley and Sons, New York 1980, pp.211-215, and
 20 D. E. Hoard and D. G. Ott, *J. Amer. Chem. Soc.* 1965, 87, 1785.

Compounds of formula (III) may also be prepared by any method known in the art.

Separation of isomers may be accomplished by methods known in the art, for example, by high-pressure liquid chromatography with chiral columns, particularly using liquid carbon dioxide as the mobile phase, or by crystallization of salts with chiral acids or bases.

- 5 Phosphate isomers may be separated with Supercritical Fluid Chromatography using a Chiralpak AS column, 25% methanol in carbon dioxide as the eluent, flow rate 2 mL/min, temperature 40°C, and pressure 3000 psi.

- 10 One aspect of the invention features the compounds according to the invention for use in medical therapy, particularly for the treatment or prophylaxis of retroviral infections and hepatitis B virus infections.

- 15 A further aspect of the invention features the compounds according to the invention for use in the manufacture of a medicament for the treatment or prophylaxis of viral infections, particularly for the treatment of retroviral infections, for example HIV infections, and hepatitis B virus infections.

- 20 In a further aspect of the present invention there is provided a method for the treatment of viral infections, for example, retroviral infections, particularly HIV infections, and hepatitis B virus infections in a host comprising administering to said host a therapeutically effective amount of a compound according to the invention.

- 25 Examples of retroviral infections which may be treated or prevented in accordance with the invention include human retroviral infections such as human immunodeficiency virus (HIV), HIV-1, HIV-2 and human T-cell lymphotropic virus (HTLV), for example, HTLV-I or HTLV-II infections. The compounds according to the invention are especially useful for the treatment of AIDS and related clinical conditions such as AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), Kaposi's sarcoma, AIDS-related neurological conditions, such as multiple sclerosis, tropical paraparesis, and AIDS dementia, anti-HIV antibody-positive and HIV-positive conditions and thrombocytopenic purpura.
- 30

The compounds according to the invention are particularly applicable for the treatment of asymptomatic infections or diseases in humans caused by or associated with human retroviruses.

- 5 The compounds according to the invention may be employed in combination with other therapeutic agents for the treatment of the above infections or conditions. Other therapeutic agents may include agents that are effective for the treatment of viral infections or associated conditions such as reverse transcriptase inhibitors, for example, zidovudine or abacavir; (1 alpha, 2 beta, 3 alpha)-9-[2,3-
- 10 bis(hydroxymethyl)cyclobutyl]guanine [(-)BHCG, SQ-34514]; oxetanocin-G (3,4-bis-(hydroxymethyl)-2-oxetanosyl]guanine); acyclic nucleosides (e.g. acyclovir, valaciclovir, famciclovir, ganciclovir, penciclovir); acyclic nucleoside phosphonates (e.g. (S)-1-(3-hydroxy-2-phosphonyl-methoxypropyl)cytosine (HPMPC) or PMEA or PMPA; ribonucleotide reductase inhibitors such as hydroxyurea, 2-acetylpyridine 5-[(2-
- 15 chloroanilino)thiocarbonyl) thiocarbonohydrazone; other 2',3'-dideoxynucleosides such as 2',3'-dideoxycytidine, 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine, 3'-deoxy-2',3'-didehydrothymidine (d4T); protease inhibitors such as saquinavir, indinavir, ritonavir, nelfinavir, amprenavir; oxathiolane nucleoside analogues such as lamivudine, cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine (FTC); 3'-deoxy-3'-fluorothymidine,
- 20 5-chloro-2',3'-dideoxy-3'-fluorouridine, ribavirin, 9-[4-hydroxy-2-(hydroxymethyl)but-1-yl]-guanine (H2G); tat inhibitors such as 7-chloro-5-(2-pyrrolyl)-3H-1,4-benzodiazepin-2-(H)one (Ro5-3335), 7-chloro-1,3-dihydro-5-(1H-pyrrol-2yl)-3H-1,4-benzodiazepin-2-amine (Ro24-7429); interferons such as α -interferon; renal excretion inhibitors such as probenecid; nucleoside transport inhibitors such as dipyridamole; pentoxifylline, N-
- 25 acetylcysteine (NAC), Procysteine, α -trichosanthin, phosphonoformic acid, as well as immunomodulators such as interleukin II or thymosin, granulocyte macrophage colony stimulating factors, erythropoetin, soluble CD4 and genetically engineered derivatives thereof; or non-nucleoside reverse transcriptase inhibitors (NNRTIs) such as nevirapine (BI-
- RG-587), loviride (α -APA) and delavuridine (BHAP), and phosphonoformic acid, and 1,4-
- 30 dihydro-2H-3,1-benzoxazin-2-ones NNRTIs such as (-)-6-chloro-4-cyclopropylethynyl-4-trifluoromethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one (L-743,726 or DMP-266), and quinoxaline NNRTIs such as isopropyl (2S)-7-fluoro-3,4-dihydro-2-ethyl-3-oxo-1(2H)-

quinoxalinecarboxylate (HBY1293). The component compounds of such combination therapy may be administered simultaneously, in either separate or combined formulations, or at different times, for example, sequentially such that a combined effect is achieved.

- 5 Another aspect of the present invention features a method of delivering a compound of formula (III), wherein R^{12} and n are defined as above, into cells by treating said cells with a compound of formula (I) or (II) as defined above. The cells to be treated may be within a human or *ex vivo*, for example, in culture.
- 10 The compounds according to the invention, also referred to herein as the active ingredient, may be administered for therapy by any suitable route including oral, rectal, nasal, topical (including buccal and sublingual), vaginal and parenteral (including subcutaneous, intramuscular, intravenous and intradermal). It will be appreciated that the preferred route will vary with the condition and age of the recipient, the nature of the infection and
- 15 the chosen active ingredient.

The amounts required of the active ingredient will depend upon a number of factors including the severity of the condition to be treated and the identity of the recipient and will ultimately be at the discretion of the attendant physician or veterinarian. In general

20 however, for each of these utilities and indications, a suitable effective dose of a compound of formula (I) will be in the range of 0.01 to 200 mg per kilogram body weight of recipient per day, advantageously in the range of 1 to 100 mg per kilogram body weight per day.

- 25 The desired dose is preferably presented as one, two, three or four or more subdoses administered at appropriate intervals throughout the day. These sub-doses may be administered in unit dosage forms, for example, containing about 0.5 to 2000 mg, preferably about 5, 25, 50, 150, 200, or 250 mg of active ingredient per unit dose form.
- 30 A further aspect of the present invention features a patient pack comprising at least one active ingredient selected from a compound of formula (I), (II), and (III) and an information insert containing directions on the use of the compound.

While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical composition. A further aspect of the present invention features pharmaceutical compositions comprising a compound of formula (I), (II) or (III) or
5 a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier therefor.

The compositions of the present invention comprise at least one active ingredient, as defined above, together with one or more pharmaceutically acceptable carriers thereof and
10 optionally other therapeutic agents. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipient thereof. Compositions include those suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration.

15 The compositions may conveniently be presented in unit dosage form prepared by any of the methods well known in the art of pharmacy. Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and
20 intimately bringing in to association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets, sachets of granules or tablets (such as a
25 swallowable, dispersible or chewable tablet) each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

30 A tablet may be made by compression or moulding optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally

mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored any may be formulated so as to provide slow or controlled release of the active ingredient therein. Tablets may be enteric coated.

Compositions suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Compositions for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate.

Compositions suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Compositions suitable for parenteral administration include aqueous and non-aqueous isotonic sterile injection solution which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The compositions may be presented in unit-dose or multidose sealed containers, for example, ampoules and vial, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

The active ingredient may also be presented in a composition comprising micrometer- to nanometer-size particles of active ingredient.

Preferred unit dosage compositions are those containing a daily dose or unit daily sub-dose (as herein above recited) or an appropriate fraction thereof, of the active ingredient.

It should be understood that in addition to the ingredients particularly mentioned above
5 the composition of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents or taste masking agents.

A further aspect of the invention relates to kits to be used in the treatment of patients
10 suffering from viral infections. These kits include one or more oral dosages of a compound of formula (I), (II), or (III) and may include one or more additional therapeutic agents. By way of illustration, a kit of the invention may include one or more tablets, capsules, caplets, gelcaps or liquid formulations containing a compound of formula (I) and one or more tablets, capsules, caplets, gelcaps or liquid formulations containing a compound of
15 formula (I) in dosage amounts within the ranges described above. The kits may include as an insert printed dosing information for the co-administration of the agents.

The following examples are intended for illustration only and are not intended to limit the scope of the invention in any way.

20

Example 1

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol

25

(a) (1S, 4R)- tert-Butyl 3-oxo-2-azabicyclo[2.2.1]hept-5-ene-2-carboxylate

(+)-2-Azabicyclo[2.2.1]hept-5-en-3-one (Chiroscience, Cambridge, England; 54.565 g, 0.500 mole) was dissolved in dry tetrahydrofuran (350 ml). Di-tert-butyl carbonate (Aldrich, 114.87 g, 0.510 mole as 97%) and 4-dimethylaminopyridine (Aldrich, 600 mg) were added to the stirred mixture. The resulting solution was stirred at ambient
30 temperature for 2 hours. Solvent was evaporated under reduced pressure and the residual orange solid was crystallized from toluene-hexanes to give title compound as

white crystals (95.72 g, 91%), m.p. 85-86°C; ¹H-NMR (CDCl₃) δ 1.50 (s, 9H), 2.24 (app AB q, J = 8.4 Hz, 2H), 3.39 (br s, 1H), 4.96 (m, 1H), 6.66 (m, 1H), 6.89 (m, 1H).

Anal. Calcd. for C₁₁H₁₅NO₃: C, 63.14; H, 7.21; N, 6.69. Found: C, 63.20; H, 7.26; N, 6.65.

5

(b) (1S,cis)-tert-Butyl N-[4-(hydroxymethyl)-2-cyclopenten-1-yl]carbamate

A solution of (1S, 4R)- tert-butyl 3-oxo-2-azabicyclo[2.2.1]hept-5-ene-2-carboxylate (part a of this example, 95.50 g, 0.456 mole) in tetrahydrofuran (500 ml)-water (50 ml) was added over 10 minutes to a vigorously stirred solution of sodium borohydride (Aldrich, 21.96 g, 0.580 mole as 99%) in water (100 ml). The temperature was maintained below 35°C. After 2 hours, the solution was cooled to maintain the temperature below 25°C while concentrated hydrochloric acid (50 ml) was added over 10 minutes. Additional water (100 ml) was added to dissolve solid and the solution was extracted with toluene (4x300 ml). The combined organic layers were washed with 9:1 saturated sodium sulfate/saturated sodium carbonate (200 ml) and dried (sodium sulfate). Evaporation of solvents under reduced pressure left a colorless syrup which crystallized on stirring with hexanes (200 ml) to provide title compound as a fine white powder (87.16 g, 90%), m.p. 72-73°C; ¹H-NMR (DMSO-d₆) δ 6.78 (d, J = 7.6 Hz, 1H), 5.80 and 5.60 (two m, 2H), 4.58 (t, J = 5.25 Hz, 2H), 4.45 (m, 1H), 3.35 (m overlapping water), 2.60 (m, 1H), 2.30 (m, 1H), 1.38 (s, 9H), 1.20 (m, 1H); [α]_D²⁰ +2.80° (c 5.0, methanol).

20

Anal. Calcd. for C₁₁H₁₉NO₃: C, 61.95; H, 8.98; N, 6.57. Found: C, 61.87; H, 8.97; N, 6.55.

25

(c) (1R,cis)-4-Amino-2-cyclopentene-1-methanol hydrochloride

(1S,cis)-tert-Butyl N-[4-(hydroxymethyl)-2-cyclopenten-1-yl]carbamate (part b of this example, 10.66 g, 50.0 mmol) was refluxed in absolute ethanol (25 mL) with concentrated hydrochloric acid (5.0 mL, 60 mequiv) for 2.5 hours. Evaporation of volatiles left title compound as white solid; mass spectrum (ES): 114 (M+1); ¹H-NMR (DMSO-d₆) δ 7.9 (m, 3H), 6.03 and 5.75 (two m, 2H), 4.11 (m, 1H), 3.41 (d, J = 5.4 Hz,

30

2H), 2.8 (m, 1H), 2.36 (m, 1H), 1.4 (m, 1H). This solid was used immediately in the following example.

(d) (1R,cis)-4-[(5-Amino-6-chloro-4-pyrimidinyl)amino]-2-cyclopentene-1-methanol

5

A solution of (+)-(1R,cis)-4-amino-2-cyclopentene-1-methanol hydrochloride (from deblocking of 10.66 g, 50.0 mmoles of (+)-(1S,cis)-tert-butyl N-[4-(hydroxymethyl)-2-cyclopenten-1-yl]carbamate as described in part c of this example), 5-amino-4,6-dichloropyrimidine (Aldrich, 16.40 g, 0.100 mole), and triethylamine (15.2 g, 0.150 mole) in 1-butanol (25 mL) was refluxed under nitrogen for 18 hours. The solution was cooled and 1 N sodium hydroxide (100 mL) was added. Volatiles were evaporated under reduced pressure and the residual solid was chromatographed on silica gel. Title compound eluted with 5% methanol-chloroform as a pale yellow glass (10.8 g). Crystallization of such a sample from ethyl acetate gave title compound as white needles, m.p. 144-146°C; ¹H-NMR (DMSO-d₆) δ 7.75 (s, 1H), 6.76 (d, J = 6.8 Hz, 1H), 5.93 and 5.82 (two m, 2H), 5.11 (m, 3H), 4.66 (t, J = 5.3 Hz, 1H), 3.40 (br t, J = 6.1 Hz, 2H), 2.75 (m, 1H), 2.20 (m, 1H), 1.38 (m, 1H).

10

15

Anal. Calcd. for C₁₀H₁₃N₄ClO: C, 49.90; H, 5.44; N, 23.28. Found: C, 49.92; H, 5.57; N, 23.10.

20

(e) (1R,cis)-4-[6-Chloro-9H-purin-9-yl]-2-cyclopentene-1-methanol

25

(1R,cis)-4-[(5-Amino-6-chloro-4-pyrimidinyl)amino]-2-cyclopentene-1-methanol (from part d of this example, 9.63 g, 40.0 mmol), triethylorthoformate (150 mL), and concentrated hydrochloric acid (14 mL) were stirred for 3 hours. Volatiles were evaporated and the residual solid was partitioned between chloroform (300 mL) and saturated aqueous sodium carbonate (100 mL). The aqueous layer was extracted with chloroform (2x100 mL). The combined chloroform layers were dried (sodium sulfate). Volatiles were evaporated under reduced pressure and the residual yellow glass was chromatographed on silica gel. Elution with ethyl acetate gave title compound as white needles from ethyl acetate (7.45 g, 74%), m.p. 121-124°C; ¹H-NMR (DMSO-d₆)

30

δ 8.81 (s, 1H), 8.64 (s, 1H), 6.24 and 6.21 (two m, 2H), 5.75 (m, 1H), 4.75 (t, $J = 5.4$ Hz, 1H), 3.34 (m, 2H), 2.95 (m, 1H), 2.75 (m, 1H), 1.75 (m, 1H).

Anal. Calcd. for $C_{11}H_{11}N_4ClO$: C, 52.70; H, 4.42; N, 22.35; Cl, 14.14. Found: C, 52.81; H, 4.46; N, 22.31; 14.22.

5

(f) (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol

(1R,cis)-4-(6-Chloro-9H-purin-9-yl)-2-cyclopentene-1-methanol (from part e of this example, 2.00 g, 7.98 mmol) was stirred in liquid ammonia (50 mL) in a Parr bomb at 10 25°C for 18 hours. Evaporation of volatiles and crystallization of the residual solid from methanol-acetonitrile gave title compound as white prisms (1.61 g, 87%), m.p. 195-200°C; 1H -NMR (DMSO- d_6) δ 8.15 (s, 1H), 8.06 (s, 1H), 7.21 (br s, 2H), 6.15 and 5.95 (two m, 2H), 5.61 (m, 1H), 4.76 (t, $J = 5.4$ Hz, 1H), 3.48 (t, $J = 5.5$ Hz, 2H), 2.92 (m, 1H), 2.71 (m, 1H), 1.67 (m, 1H); $[\alpha]_{589}^{20} +4.5^\circ$ (c 0.5, methanol).

15 Anal. Calcd. for $C_{11}H_{13}N_5O$: C, 57.13; H, 5.67; N, 30.28. Found: C, 57.25; H, 5.67; N, 30.33.

Example 2

20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-alaninyl)phosphoramidate

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (part f of example 1, 925 mg, 4.00 mmol) was stirred in dry pyridine (100 ml) and *tert*-butyl magnesium chloride (Aldrich, 1 M in tetrahydrofuran, 4.3 ml) was added. After 15 minutes, a solution of 25 phenyl(methoxy-L-alaninyl)phosphorochloridate (prepared as described by McGuigan, C. et al., *J. Med. Chem.* 1993, 36: 1048-1052) (2.22 g, 8.00 mmol) in tetrahydrofuran (10 ml) was added. After 24 hours, additional *tert*-butyl magnesium chloride (4.4 ml) and phenyl(methoxy-L-alaninyl)phosphorochloridate (2.22 g) were added and stirring continued for an additional 24 hours. Volatiles were removed and the residual gummy 30 solid was partitioned between chloroform (200 ml) and water (50 ml). The chloroform layer was dried (sodium sulfate) and concentrated to a colorless glass. The glass was chromatographed on silica gel. Title compound was eluted with 5% methanol-chloroform.

Evaporation of a methanol solution left the title compound as a white solid foam (1.02 g, 54%); high resolution mass spectrum calcd for $C_{21}H_{26}N_6O_5P$ (M+H)⁺ m/z 473.1702, found 473.1707; ¹H-NMR (CDCl₃) δ 8.39 (s, 1H), 8.04 and 7.99 (both s, 1H), 7.2 (m, 5H), 6.50 (m, 1H), 6.2 (m, 1H), 6.0 (m, 1H), 5.79 (m, 1H), 4.3-3.95 (m, 3H), 3.74 and 3.71 (two s, 3H), 3.25 (m, 1H), 2.95 (m, 1H), 1.80 (m, 1H), 1.38 and 1.37 (two d, each J = 7.0 Hz, 3H); ³¹P-NMR (CDCl₃): 3.12, 2.80.

Anal. Calcd. for $C_{21}H_{25}N_6O_5P \cdot 0.19 CH_3OH \cdot 0.35 H_2O$: C, 52.49; H, 5.50; N, 17.73. Found: C, 52.51; H, 5.49; N, 17.35.

10 Example 3

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-alaninyl)phosphate

In the same manner as Example 2, (1R,cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (part f of example 1) was reacted with phenyl(ethoxy-L-alaninyl)phosphorochloridate (prepared as described by McGuigan, C. et al., *J. Med. Chem.* 1993, 36: 1048-1052). Title compound was eluted with 5% methanol-chloroform.

Evaporation of a methanol solution left the title compound as a white solid foam; high resolution mass spectrum calcd for $C_{22}H_{28}N_6O_5P$ (M+H)⁺ m/z 487.1859, found 187.1841; ¹H-NMR(DMSO-d₆) δ 8.16 (s, 1H), 8.05 and 8.01 (both s, total 1H), 7.36 (m, 2H), 7.24 (br s, 2H), 7.19 (m, 3H), 6.13 (m, 1H), 6.02 (m, 2H), 5.63 (m, 1H), 4.03 (m overlapping q, J = 7.0 Hz, 4H), 3.77 (m, 1H), 3.15 (m, 1H), 2.75 (m, 1H), 1.70 (m, 1H), 1.21 (d, J = 7.1 Hz, 3H), 1.14 and 1.12 (two t, J = 7.0, 7.0 Hz, total 3H); ³¹P-NMR (DMSO-d₆) 4.23, 3.88.

25 Example 4

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(methoxy)-D-alaninyl]phosphoramidate

In the same manner as Example 2, (1R,cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (part f of example 1, 200 mg, 0.860 mmol) was reacted with phenyl(methoxy-D-alaninyl)phosphorochloridate (prepared as described by McGuigan, C. et al., *J. Med. Chem.* 1993, 36: 1048-1052). Title compound was eluted with 5% methanol-chloroform.

Evaporation of a methanol solution left the title compound as a white solid foam (290 mg, 71%); ¹H-NMR (DMSO-d₆) δ 8.13 (s, 1H), 8.02 and 8.01 (both s, total 1H), 7.36 (m, 1H), 7.34 (m, 1H), 7.19 (m, 5H), 6.13 (m, 1H), 6.02 (m, 2H), 5.63 (m, 1H), 4.1 (m, 2H), 3.8 (m, 1H), 3.57 (s, 3H), 3.12 (m, 1H), 2.72 (m, 1H), 1.68 (m, 1H), 1.20 (m, 3H); ³¹P-NMR (DMSO-d₆) 3.96, 3.625.

Anal. Calcd. for C₂₁H₂₅N₆O₅P·0.48 H₂O: C, 52.43; H, 5.44; N, 17.47. Found: C, 52.43; H, 5.43; N, 17.43.

Example 5

10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(benzyloxy)-L-alaninyl]phosphoramidate

In the same manner as Example 2, (1R,cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (part f of example 1, 200 mg, 0.860 mmol) was reacted with phenyl(benzyloxy-L-alaninyl)phosphorochloridate (prepared as described by McGuigan, C. et al., *J. Med. Chem.* 1993, 36: 1048-1052). Title compound was eluted with 5% methanol-chloroform. Evaporation of a methanol solution left the title compound as a white solid foam (270 mg, 57%); ¹H-NMR (DMSO-d₆) δ 8.13 (s, 1H), 8.01 and 7.985 (both s, total 1H), 7.2-7.4 (m, 12H), 6.02 (m, 3H), 5.59 (m, 1H), 5.06 (m, 2H), 4.03 (m, 2H), 3.83 (m, 1H), 3.05 (m, 1H), 2.65 (m, 1H), 1.62 (m, 1H), 1.22 (d, J = 7.0 Hz, 3H); ³¹P-NMR (DMSO-d₆) 4.00, 3.55.

Anal. Calcd. for C₂₇H₂₉N₆O₅P·0.47 H₂O: C, 58.22; H, 5.42; N, 15.09. Found: C, 58.22; H, 5.44; N, 14.84.

Example 6

25 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(methoxy)-L-phenylalaninyl]phosphoramidate

In the same manner as Example 2, (1R,cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (part f of example 1, 500 mg, 2.16 mmol) was reacted with phenyl(methoxy-L-phenylalaninyl)phosphorochloridate (prepared as described by McGuigan, C. et al., *J. Med. Chem.* 1993, 36: 1048-1052). Title compound was eluted with 5% methanol-chloroform. Evaporation of a methanol solution left the title compound as a white solid foam (500 mg,

42%); $^1\text{H-NMR}$ (DMSO-d_6) δ 8.14 and 8.13 (both s, total 1H), 8.01 and 7.985 (both s, total 1H), 7.2-7.4 (m, 5H), 6.95-7.05 (m, 5H), 7.00 and 6.98 (both br s, total 2H), 6.1 (m, 1H), 5.975 (m, 2H), 5.58 (m, 1H), 3.55-4.0 (m, 3H), 3.57 (s, 3H), 2.95 (m, 2H), 2.7 (m, 2H), 1.55 (m, 1H); $^{31}\text{P-NMR}$ (DMSO-d_6) 3.605, 3.25.

5 Anal. Calcd. for $\text{C}_{27}\text{H}_{29}\text{N}_6\text{O}_5\text{P}\cdot 0.38\text{H}_2\text{O}\cdot 0.16\text{CH}_3\text{CN}$: C, 58.39; H, 5.42; N, 15.35. Found: C, 58.39; H, 5.41; N, 15.36.

Example 7

10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(ethoxy)-L-phenylalaninyl]phosphoramidate

In the same manner as Example 2, (1R,cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (part f of example 1, 150 mg, 0.27 mmol) was reacted with phenyl(methoxy-L-phenylalaninyl)phosphorochloridate (prepared as described by McGuigan, C. et al., *J. Med. Chem.* 1993, 36: 1048-1052). Title compound was eluted with 5% methanol-chloroform. Evaporation of a methanol solution left the title compound as a white solid foam (120 mg, 80%); $^1\text{H-NMR}$ (CDCl_3) δ 8.38 (two s, 1H), 7.83 (two s, 1H), 7.0-7.3 (m, 10H), 6.25 (broad s, 2H), 6.10 (m, 1H), 5.92 (m, 1H), 5.70 (m, 1H), 3.7-4.3 (m, 5H), 2.6-3.2 (m, 5H), 1.60 (m, 1H), 1.17 (two t, 3H); $^{31}\text{P-NMR}$ (CDCl_3) δ 3.86. Mass spectrum calcd for $\text{C}_{28}\text{H}_{31}\text{N}_6\text{O}_5\text{P}$ ($\text{M}+\text{H}$) $^+$ m/z 563, found 563.

Anal. Calcd. for $\text{C}_{28}\text{H}_{31}\text{N}_6\text{O}_5\text{P}\cdot 2/3\text{H}_2\text{O}$ C, 58.53; H, 5.67; N, 14.63. Found: C, 58.77; H, 5.51; N, 14.65.

25 Example 8

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(L-phenylalaninyl)phosphoramidate Disodium Salt

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(methoxy)-L-phenylalaninyl]phosphoramidate (0.060 g, 0.11 mmoles) was suspended in solution of triethylamine (2 mL) and deionized water (2 mL) and stirred at room temperature for 18 hours. The volatiles were removed by spin evaporation *in vacuo* and the residue was

dissolved in water (20 mL), extracted with dichloromethane (2x20 mL), and purified by anion exchange chromatography on a Sep-Pak® Vac 35cc Accell™ Plus QMA cartridge (Waters Corp., P/N WAT054725) (HCO₃⁻ form) with an aqueous ammonium bicarbonate buffer (0 - 0.5 M gradient, 1 L). The appropriate fractions were combined and the volatiles
5 were removed by spin evaporation *in vacuo*. The residue was twice dissolved in deionized water and spin evaporated *in vacuo* to give the title compound as an ammonium salt. This salt was dissolved in deionized water and passed through a Sep-Pak® Vac 20cc Accell™ Plus CM cartridge (Waters Corp., P/N WAT054675) (Na⁺ form) using deionized water.

Lyophilization of the appropriate fractions left the title compound disodium salt 3.2
10 hydrate as a white solid (35 mg, 51%); ¹H-NMR (D₂O) δ 8.02 (s, 1H), 7.96 (s, 1H), 6.98 - 7.10 (m, 3H), 6.84 - 6.92 (m, 2H), 6.14 - 6.20 (m, 1H), 5.88 - 5.90 (m, 1H), 5.42 - 5.50 (m, 1H), 4.4 - 5.0 (br m, 3H + HOD), 3.58 - 3.66 (m, 1H), 3.36 - 3.48 (m, 2H), 2.94 - 3.06 (bm, 1H), 2.64 - 2.78 (m, 1H), 2.40 (d, J = 6.6 Hz, 2H), 1.54 - 1.64 (m, 1H); ³¹P-NMR (D₂O) 7.8. MS (ES⁻) *m/e* 457 (MH⁻).
15 Anal. Calcd. for C₂₀H₂₁N₆Na₂O₅P 3.2 H₂O: C, 42.89; H, 4.93; N, 15.01. Found: C, 42.92; H, 4.58; N, 14.70.

Example 9

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl(methoxy)-
20 α,α-dimethylglyciny)phosphoramidate

In the same manner as Example 2, (1R,cis)-4-(6-amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (part f of example 1, 200 mg, 0.86 mmol) was reacted with phenyl(methoxy-
α,α-alaniny)phosphorochloridate (prepared as described by McGuigan, C. et al., *J. Med.*
25 *Chem.* 1993, 36: 1048-1052). Title compound was eluted with 5% methanol-chloroform. Evaporation of a methanol solution left the title compound as a white solid foam (200 mg, 48%); ¹H-NMR (DMSO-d₆) δ 8.15 (s, 1H), 8.03 and 8.01 (both s, total 1H), 7.1-7.4 (m, 7H), 6.10 (m, 1H), 6.00 (m, 1H), 5.89 (m, 1H), 5.625 (m, 1H), 4.08 (m, 2H), 3.55 (s, 3H), 3.12 (m, 1H), 2.74 (m, 1H), 1.70 (m, 1H), 1.35 (m, 6H); ³¹P-NMR (DMSO-d₆) 2.43, 2.39.

30 Anal. Calcd. for C₂₂H₂₇N₆O₅P·0.53 H₂O·0.03 CH₃CN: C, 53.29; H, 5.71; N, 16.99. Found: C, 53.29; H, 5.67; N, 16.99.

The following general procedures were used in the preparation of compounds of Examples 10 – 41.

Standard Procedure for Phosphorochloridate Preparation

- 5 Dry triethylamine (2.0 mol equiv.) in dry dichloromethane (40 ml) was added dropwise to a stirred solution of phenyl dichlorophosphate (1.0 mol equiv.) and the appropriate amino acid ester salt (1.0 mol equiv.) in dry dichloromethane (40 ml), at -78 °C under nitrogen. Following the addition, the reaction mixture was allowed to warm slowly to room temperature and stirred overnight. The solvent was removed under reduced pressure and
10 the crude residue was resuspended in dry diethyl ether or THF, and filtered under nitrogen. The solvent was removed under reduced pressure to leave the crude product as an oil.

All crude phosphorochloridates were used as solutions in dry THF or dry acetonitrile in subsequent coupling reactions.

15

Standard Procedure 1 for Phosphoramidate Preparation

- 20 To (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (200 mg, 0.87 mmol) suspended in anhydrous acetonitrile (15 ml) under a nitrogen atmosphere, was added 1M t-butyl magnesium chloride dissolved in tetrahydrofuran (1.73 ml, 1.74 mmol). After 15 minutes, phosphorochloridate (2.61 mmol) dissolved in acetonitrile (15ml) was added dropwise over 1 minute and the reaction mixture allowed to stir for a further 4 hours.
25 Following the removal of volatiles *in vacuo*, the product was purified by column chromatography (silica) eluting with 4-5% MeOH in chloroform or dichloromethane.

Standard Procedure 2 for Phosphoramidate Preparation

- 30 To (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (200 mg, 0.87 mmol) suspended in anhydrous tetrahydrofuran (15 ml) under a nitrogen atmosphere, was added 1M t-butyl magnesium chloride dissolved in tetrahydrofuran (1.73 ml, 1.74 mmol). After 15

minutes, phosphorochloridate (2.61 mmol) dissolved in acetonitrile (15ml) was added dropwise over 1 minute and the reaction mixture allowed to stir for a further 4 hours. Following the removal of volatiles *in vacuo*, the product was purified by column chromatography (silica) eluting with 4-5% MeOH in chloroform or dichloromethane.

5

Standard Procedure 3 for Phosphoramidate Preparation

To (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (200 mg, 0.87 mmol) suspended in anhydrous pyridine (15 ml) under a nitrogen atmosphere, was added 1M t-butyl magnesium chloride dissolved in tetrahydrofuran (1.73 ml, 1.74 mmol). After 15 minutes, phosphorochloridate (2.61 mmol) dissolved in acetonitrile (15ml) was added dropwise over 1 minute and the reaction mixture allowed to stir for a further 4 hours. Following the removal of volatiles *in vacuo*, the product was purified by column chromatography (silica) eluting with 4-5% MeOH in chloroform or dichloromethane.

15

Example 10

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-L-(O-tert-butyltyrosinyl)]phosphoramidate

20 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (60 %); δ_P 3.98, 4.04; δ_H 1.32 (9H, s), 1.64 (1H, m), 2.83 (1H, m), 2.95 (2H, m), 3.08 (1H, m), 3.61, 3.66 (3H, s), 4.07 (4H, m), 5.73 (1H, m), 5.94 (1H, m), 6.11 (1H, m), 6.39 (2H, br s), 6.88 (2H, d), 6.98 (2H, d), 7.16 (2H, m), 7.29 (3H, m), 7.84, 7.88 (1H, s), 8.37, 8.38 (1H, s); m/z (ES+) 643.2405 (MH⁺, C₃₁H₃₇N₆O₆NaP requires 643.2410).

25

Example 11

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy-L-alaninyl)phosphoramidate

30 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (89 %); δ_P 3.97, 4.20; δ_H 1.14 (6H, m), 1.31 (3H, m), 1.63 (1H, m), 2.79 (1H, m), 3.12 (1H, m), 4.05 (3H, m), 4.92 (1H, m), 5.66 (1H, m), 5.87 (1H, m), 6.08 (1H, m), 6.22 (2H, br s),

7.06 (2H, m), 7.19 (3H, m), 7.78, 7.82 (1H, s), 8.29 (1H, s); m/z (ES+) 523.1832 (MH⁺, C₂₃H₂₉N₆O₅NaP requires 523.1835).

Example 12

- 5 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl *tert*-butoxy-L-alaninyl)phosphoramidate

Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (59 %); δ_F 4.10, 4.27; δ_H 1.24 (3H, m), 1.35 (9H, s), 1.54 (1H, m), 2.79 (1H, m), 3.13 (1H, m), 3.80 (1H, m), 4.09 (3H, m), 5.66 (1H, m), 5.87 (1H, m), 6.08 (1H, m), 6.22 (2H, br s), 7.15
10 (5H, m), 7.78, 7.82 (1H, s), 8.29 (1H, s); m/z (ES+) 537.2001 (MH⁺, C₂₄H₃₁N₆O₅NaP requires 537.1991).

Example 13

- 15 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy)-L-alaninyl]phosphoramidate

Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (66 %); δ_F 3.89, 4.18; δ_H 0.94 (9H, d), 1.41 (3H, m), 1.73 (1H, m), 2.89 (1H, m), 3.23
20 (1H, m), 3.83 (2H, m), 4.14 (4H, m), 5.77 (1H, m), 5.98 (1H, m), 6.20 (3H, m), 7.18 (2H, m), 7.32 (3H, m), 7.88, 7.94 (1H, s), 8.38 (1H, s); m/z (ES+) 551.2145 (MH⁺, C₂₅H₃₃N₆O₅NaP requires 551.2148).

Example 14

- 25 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl cyclopropyl methoxy-L-alaninyl)phosphoramidate

Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (59 %); δ_F 3.99, 4.21; δ_H 0.14 (2H, m), 0.45 (2H, m), 0.99 (1H, m), 1.28 (3H, m), 1.61
30 (1H, m), 2.76 (1H, m), 3.10 (1H, m), 3.84 (2H, m), 4.06 (2H, m), 4.50 (1H, m), 5.64 (1H, m), 5.85 (1H, m), 6.06 (1H, m), 6.42 (2H, br s), 7.13 (5H, m), 7.77, 7.80 (1H, s), 8.27 (1H, s); m/z (ES+) 535.1834 (MH⁺, C₂₄H₂₉N₆O₅NaP requires 535.1835).

Example 15

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-leuciny)phosphoramidate

- 5 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (79 %); δ_P 4.14, 4.37; δ_H 0.78 (6H, m), 1.39 (2H, m), 1.58 (2H, m), 2.77 (1H, m), 3.10 (1H, m), 3.56, 3.59 (3H, s), 4.02 (4H, m), 5.65 (1H, m), 5.86 (1H, m), 6.00 (1H, m), 6.26 (2H, br s), 7.06 (2H, m), 7.21 (3H, m), 7.77, 7.84 (1H, s), 8.28 (1H, s); m/z (ES+) 537.2000 (MH⁺, C₂₄H₃₁N₆O₅NaP requires 537.1991).

10

Example 16

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-L-(O-methyltyrosinyl)]phosphoramidate

- 15 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (89 %); δ_P 4.07; δ_H 1.62 (1H, m), 2.81 (1H, m), 2.93 (2H, m), 3.12 (1H, m), 3.64, 3.68 (3H, s), 3.74 (3H, m), 4.09 (3H, m), 4.47 (1H, m), 5.73 (1H, m), 5.94 (1H, m), 6.11 (1H, m), 6.43 (2H, br s), 6.75 (2H, d), 6.98 (2H, d), 7.16 (3H, m), 7.27 (2H, m), 7.83, 7.85 (1H, s), 8.36, 8.37 (1H, s).

20

Example 17

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-phenylalaninyl)phosphoramidate

- 25 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (24 %), δ_P 3.92, 4.04; δ_H 1.7 (1H, m), 2.8 (1H, m), 3.1 (1H, m), 3.2 (1H, m), 3.7 (3H, d), 4.1 (2H, m), 4.6 (1H, q), 5.7 (1H, m), 5.9 (1H, m), 6.1 (1H, m), 6.5 (2H, d), 7.2 (10H, m), 7.85 (1H, d), 8.4 (1H, d).

30

Example 18

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-leuciny)phosphoramidate

- 5 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (28 %), δ_p 4.27, 4.36; δ_H 0.8 (6H, m), 1.5 (2H, m), 5.9 (1H, m), 6.05 (1H, m), 6.55 (2H, s), 7.2 (5H, m), 7.8 (1H, d), 8.23 (1H, d).

Example 19

- 10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-tryptophanyl)phosphoramidate (diastereomeric mixture)

- Synthesis according to standard procedure 1. Elution from a silica gel column with 5% methanol in methylene chloride gave a 1.15:1 mixture of high:low R_f isomers (from ^{31}P -
15 NMR), isolated as a brittle white foam (23 %); δ_p 4.13, 4.23; δ_H 1.5 (1H, m), 2.65 (1H, m), 3.2 (2H, m), 3.62 (3H, d), 3.95 (2H, m), 4.35 (1H, m), 5.67 (1H, m), 5.9 (1H, m), 6.0 (1H, m), 6.4 (2H, s), 7.2 (9H, m), 7.5 (1H, d), 7.73 (1H, s), 8.35 (1H, s), 9.05 (1H, s).

Example 20

- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-tryptophanyl)phosphoramidate (single diastereomer)

- Synthesis according to standard procedure 1. Elution from a silica gel column with 5% methanol in methylene chloride gave initial fractions containing the pure higher R_f isomer
25 of the mixture described in Example 19 (from ^{31}P -NMR), isolated as a brittle white foam (34 %); δ_p 4.02; δ_H 1.5 (1H, m), 2.7 (1H, m), 3.2 (2H, m), 3.6 (3H, d), 4 (2H, m), 4.3 (1H, m), 5.7 (1H, m), 5.85 (1H, m), 6 (1H, m), 6.5 (2H, s), 7.3 (9H, m), 7.5 (1H, d), 7.8 (1H, s), 8.4 (1H, s), 9.2 (1H, s).

Example 21

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-valinyl)phosphoramidate

- 5 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (26 %), δ_P 4.76, 4.85; δ_H 0.9 (6H, m), 1.8 (1H, m), 2.1 (1H, m), 3.25 (1H, m), 3.7 (3H, d), 3.9 (1H, m), 4.2 (2H, m), 5.8 (1H, m), 6 (1H, m), 6.2 (1H, m), 6.3 (2H, s), 7.25 (5H, m), 7.9 (1H, d), 8.4 (1H, s).

10 Example 22

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl dimethoxy-D-aspartyl)phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
15 foam (28 %), δ_P 3.79, 4.24; δ_H 1.7 (1H, m), 2.8 (3H, m), 3.1 (1H, m), 3.6 (3H, d), 3.7 (3H, d), 4.1 (2H, m), 4.4 (1H, m), 5.5 (1H, m), 5.9 (1H, m), 6.1 (1H, m), 6.3 (2H, s), 7.15 (5H, m), 7.8 (1H, d), 8.3 (1H, d)

Example 23

- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-prolinyl)phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
) foam (19%); δ_P 2.89, 3.04; δ_H 1.68-2.19 (5H, br m), 2.94 (1H, m), 3.32 (2H, m), 3.42 (1H, m),
25 3.61, 3.74 (3H, s), 4.12, 4.30 (m, 5'-H), 5.70 (2H, br s), 5.81 (1H, m), 5.98 (1H, m), 6.19, 6.25 (1H, m), 7.21 (2H, m), 7.35 (3H, m), 7.88, 8.02 (1H, s), 8.43 (1H, s).

Example 24

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-isoleuciny)phosphoramidate

- 5 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (45%); δ_P 4.55, 4.72; δ_H 0.92 (6H, m), 1.16 (1H, m), 1.42 (1H, m), 1.77 (2H, m), 2.99 (1H, m), 3.22 (1H, m), 3.62, 3.66 (3H, s), 3.84 (2H, m), 4.22 (2H, m), 5.74 (1H, m), 5.99 (3H, m), 6.17 (1H, m), 7.11 (2H), 7.29 (3H, m), 7.86, 7.92 (1H, s), 8.36 (1H, s).

10 **Example 25**

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-norvaliny)phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
15 foam (73%); δ_P 4.10, 4.35; δ_H 0.88 (3H, t, J 8.4), 1.32 (2H, m), 1.71 (3H, m), 2.91 (1H, m), 3.20 (1H, m), 3.65, 3.68 (3H, s), 3.87 (1H, m), 4.03 (2H, m), 4.19 (1H, m), 5.74 (1H, m), 5.93 (1H, m), 6.05 (2H, br s), 6.13 (1H, m), 7.14 (2H, m), 7.28 (3H, m), 7.83, 7.90 (1H, s), 8.34 (1H, s).

20 **Example 26**

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl *i*-propoxy-L-phenylalaniny)phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
25 foam (76%); δ_P 3.97; δ_H 1.20 (6H, m), 1.66 (1H, m), 2.82 (1H, m), 2.98 (2H, m), 3.11 (1H, m), 3.79-4.06 (4H, br m), 4.99 (1H, m), 5.72 (1H, m), 5.96 (1H, m), 6.08 (3H, m), 7.23 (10H, m), 7.87 (1H, s), 8.34 (1H, s).

Example 27

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl *tert*-butoxy-L-phenylalaninyl)phosphoramidate

- 5 Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (51%); δ_P 4.14, 4.09; δ_H 1.31, 1.34 (9H, s), 1.56 (1H, m), 2.75 (1H, m), 2.90 (2H, m), 3.20 (1H, m), 3.94 (4H, m), 5.63 (1H, m), 5.86 (1H, m), 6.02 (3H, m), 7.13 (10H, m), 7.73 (1H, s), 8.29 (1H, s).

10 Example 28

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy)-L-phenylalaninyl]phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
15 foam (23%); δ_P 3.94, 3.99; δ_H 0.85, 0.92 (9H, s), 1.62 (1H, m), 2.80 (1H, m), 3.03 (2H, m), 3.20 (1H, m), 3.79 (3H, m), 4.02 (2H, m), 4.29 (1H, m), 5.73 (1H, m), 5.92 (1H, m), 6.02 (2H, br s), 6.09 (1H, m), 7.18 (10H, m), 7.81, 7.83 (1H, s), 8.36, 8.37 (1H, s).

Example 29

- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl benzyloxy-L-phenylalaninyl)phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
foam (53%); δ_P 3.84, 3.90; δ_H 1.56 (1H, m), 2.81 (1H, m), 3.02 (3H, m), 3.92 (3H, m), 4.27
25 (1H, m), 5.11 (2H, m), 5.69 (1H, m), 5.85 (1H, m), 6.03 (3H, br s), 6.94 (2H, m), 7.13 (13H, m), 7.77, 7.78, 7.80 (1H, s), 8.34 (1H, s).

Example 30

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[(2-carbomethoxy)phenyl methoxy-L-alaninyl]phosphoramidate

- 5 Synthesis according to standard procedure 3. Title compound isolated as a brittle white foam (39%); δ_P 3.95, 4.05; δ_H 1.28 (6H, m), 1.69 (1H, m), 2.80 (1H, m), 3.15 (1H, m), 3.52, 3.60 (3H, s), 4.18 (5H, m), 5.70 (1H, m), 5.88 (1H, m), 7.16 (1H, m), 7.44 (2H, m), 7.80 (1H, m), 7.87, 7.92 (1H, s), 8.28 (1H, s); MS (E/I) 567.1722 (calc. 567.1733).

10 Example 31

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[4-(3-oxo-3-phenylpropenyl)phenyl methoxy-L-alaninyl]phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
15 foam (74%); δ_P 3.68, 3.97; δ_H 1.40 (3H, m), 1.78 (1H, m), 2.93 (1H, m), 3.25 (1H, br, s), 3.75 (3H, s), 4.27 (3H, m), 5.78 (1H, m), 6.10 (1H, m), 6.19 (1H, m), 7.25 (1H, m), 7.32 (2H, m), 7.58 (4H, m), 7.90 (1H, s), 8.05 (2H, s), 8.50 (1H, s); MS (E/I) 625.1956 (calc. 625.1940).

Example 32

- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy glyciny)phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
foam (38%); δ_P 4.51; δ_H 1.72 (1H, m), 2.82 (1H, m), 3.20 (1H, m), 3.70 (3H, s), 4.00 (2H, m),
25 4.20 (2H, m), 5.68 (1H, m), 5.90 (1H, m), 6.15 (1H, m), 7.20 (5H, m), 7.99 (1H, s), 8.31 (1H, s);
MS (EI) 481.1373 (calc. 481.1365).

Example 33

- 30 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-valinyl)phosphoramidate

Synthesis according to standard procedure 1. Title compound isolated as a brittle white foam (86%); δ_P 4.64, 4.84; δ_H 0.83 (6H, m), 1.65 (1H, m), 1.95 (1H, m), 2.82 (1H, m), 3.15

(1H, m), 3.60, 3.65 (3H, s), 3.78 (1H, m), 4.11 (2H, m), 5.68 (1H, m), 5.89 (1H, m), 6.12 (1H, m), 7.10 (2H, m), 7.22 (3H, m), 7.84, 7.92 (1H, s), 8.29 (1H, s); MS (EI) 523.1855 (calc. 523.1835).

5 Example 34

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-methioninyl)phosphoramidate

Synthesis according to standard procedure 1. Title compound isolated as a brittle white
10 foam (96%); δ_P 3.79, 4.19; δ_H 1.70 (1H, m), 1.85 (2H, m), 1.97 (3H, s), 2.39 (2H, t), 2.81 (1H, m), 3.15 (1H, m), 3.64, 3.67 (3H, s), 3.91 (1H, m), 4.13 (2H, m), 5.68 (1H, m), 5.89 (1H, m), 6.12 (1H, m), 7.12 (2H, m), 7.28 (3H, m), 7.90, 8.00 (1H, s), 8.29 (1H, s); MS (E/I) 555.1567 (calc. 555.1555).

15 Example 35

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy- α,α -dimethylglycinyl)phosphoramidate

Synthesis according to standard procedure 1. Title compound isolated as a brittle white
20 foam (91%); δ_P 2.51; δ_H 1.30 (3H, t), 1.57 (6H, m), 1.73 (1H, m), 2.88 (1H, m), 3.22 (1H, m), 4.19 (4H, m), 5.74 (1H, m), 5.95 (1H, m), 6.19 (1H, m), 7.23 (5H, m), 7.92, 7.96 (1H, s), 8.35 (1H, s).

Example 36

25 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy- α,α -dimethylglycinyl)phosphoramidate

Synthesis according to standard procedure 1. Title compound isolated as a brittle white
foam (21%); δ_P 2.56; δ_H 1.30 (6H, m), 1.58 (6H, m), 1.78 (1H, m), 2.90 (1H, m), 3.24 (1H, m),
30 4.08 (1H, m), 4.20 (2H, m), 5.07 (1H, m), 5.65 (2H, br, s), 6.00 (1H, m), 6.20 (1H, m), 7.26 (5H, m), 7.83, 7.89 (1H, s), 8.43 (1H, s).

Example 37

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy- α,α -dimethylglyciny)]phosphoramidate

- 5 Synthesis according to standard procedure 2. Title compound isolated as a brittle white foam (38%); δ_P 2.42, 2.44; δ_H 0.99 (9H, s), 1.61 (6H, dd), 1.74 (1H, m), 2.90 (1H, m), 3.23 (1H, m), 3.86 (2H, s), 4.20 (2H, m), 5.78 (1H, m), 6.00 (1H, m), 6.20 (1H, m), 7.28 (5H, m), 7.86, 7.92 (1H, s), 8.40 (1H, s).

10 Example 38

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy cyclopentaneglyciny)]phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
15 foam (54%); δ_P 2.94; δ_H 1.67 (1H, m), 2.04 (8H, m), 2.91 (1H, m), 3.24 (1H, m), 3.75 (3H, s), 4.25 (2H, m), 5.80 (1H, m), 6.02 (1H, m), 6.22 (1H, m), 7.25 (5H, m), 7.90, 8.00 (1H, s), 8.45 (1H, s); MS (EI) 535.1816 (calc. 535.1835).

Example 39

- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy cyclohexaneglyciny)]phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
foam (44%); δ_P 2.88; δ_H 1.64 (10H, m), 1.71 (1H, m), 2.85 (1H, m), 3.20 (1H, m), 3.71 (3H, s),
25 4.19 (2H, m), 5.76 (1H, m), 5.98 (1H, m), 6.17 (1H, m), 7.25 (5H, m), 7.85, 7.95 (1H, s), 8.42 (1H, s); MS (EI) 549.2008 (calc. 549.1991).

Example 40

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl dimethoxy-L-aspartyl)phosphoramidate

- 5 Synthesis according to standard procedure 2. Title compound isolated as a brittle white foam (34%); δ_P 3.78, 4.38; δ_H 1.74 (1H, m), 2.85 (1H, m), 3.22 (1H, m), 3.50 (2H, s), 3.65 (3H, s), 3.75 (3H, s), 4.13 (1H, m), 4.26 (2H, m), 5.71 (1H, m), 5.92 (1H, m), 6.23 (1H, m), 7.16 (5H, m), 8.00 (1H, m), 8.13, 8.21 (1H, s).

10 Example 41

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[(4-chlorophenyl) methoxy-L-alaninyl]phosphoramidate

- Synthesis according to standard procedure 1. Title compound isolated as a brittle white
15 foam (28 %), δ_P 3.88, 4.16; δ_H [CDCl₃] 1.3 (3H, m), 1.7 (1H, m), 2.9 (1H, m), 3.2 (1H, m), 3.7 (3H, d), 4.2 (2H, m), 5.7 (1H, m), 6 (1H, m), 6.2 (1H, m), 7.3 (4H, m), 7.95 (1H, d), 8.4 (1H, d).

Example 42

- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(L-(methylamino)alaninyl)phosphoramidate

- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(methoxy)-L-alaninyl]phosphoramidate (Example 2, 60 mg, 0.127 mmoles) was suspended in solution of 40% methylamine (2 mL) and stirred at room temperature for 4 hours. The volatiles were
25 removed by spin evaporation *in vacuo* and the residue was dissolved in water (20 mL), extracted with dichloromethane (2x20 mL) and purified by anion exchange chromatography on a Sep-Pak® Vac 35cc Accell™ Plus QMA cartridge (Waters Corp., P/N WAT054725) (HCO₃⁻ form) with an aqueous ammonium bicarbonate buffer (0 - 0.5 M gradient, 1 L). The appropriate fractions were combined and the volatiles were removed by
30 spin evaporation *in vacuo*. The residue was twice dissolved in deionized water and spin evaporated *in vacuo* to give the title compound as an ammonium salt. This salt was dissolved in deionized water and passed through a Sep-Pak® Vac 20cc Accell™ Plus CM

cartridge (Waters Corp., P/N WAT054675) (Na⁺ form) using deionized water. Lyophilization of the appropriate fractions left the title compound sodium salt as a white solid (26 mg, 52%); ¹H-NMR (D₂O) δ 7.92 (s, 1H), 7.91 (s, 1H), 6.09 - 6.15 (m, 1H), 5.83 - 5.88 (m, 1H), 5.30 - 5.40 (m, 1H), 3.62 (t, J = 5.3 Hz, 1H), 3.28 - 3.48 (m, 2H), 2.92 - 3.04 (bm, 1H), 2.58 - 2.71 (m, 1H), 2.44 - 2.51 (m, 3H), 1.36 - 1.46 (m, 1H), 1.01 (d, J = 6.8 Hz, 3H); ³¹P-NMR (D₂O) 6.55. Mass spectrum (ES⁻) *m/e* 394 (M-H).

Example 43

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl)phosphate

This compound was prepared in an analogous manner to Example 42 with the replacement of phosphoryl chloride by phenyl dichlorophosphate. Removal of the volatiles by evaporation *in vacuo* gave the ammonium salt of the title molecule as a hygroscopic solid (99 mg, 57%); ¹H-NMR (D₂O) δ 7.92 (s, 1H), 7.75 (s, 1H), 6.82 - 6.92 (m, 2H), 6.62 - 6.76 (m, 3H), 6.08 - 6.16 (m, 1H), 5.81 - 5.88 (m, 1H), 5.16 - 5.25 (m, 1H), 3.79 - 3.88 (m, 1H), 3.61 - 3.72 (m, 1H), 3.2.95 - 3.3.06 (bm, 1H), 2.42 - 2.58 (m, 1H), 1.18 - 1.30 (m, 1H); ³¹P-NMR (D₂O) -4.56 (m, 1P), Mass spectrum (ES⁻) *m/e* 386 (M-H).

Example 44

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-monophosphate

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (Example 1, 200 mg, 0.865 mmol) was dissolved in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (2 mL). Phosphoryl chloride (0.24 mL, 0.26 mmol) was added to the stirred, cooled (0° C.) solution. After 1 minute, 1.0 M sodium hydrogen carbonate (3.3 mL, 3.3 mmol) was added and stirring was continued at 0° C. for 30 minutes and then at 25° C. for 1 hour. The reaction solution was diluted to 125 mL with deionized water and applied to a 1.1 x 7.0 cm DEAE Sepahadex A25 (Aldrich) ion exchange chromatography column which had been washed with 1.0 M ammonium bicarbonate buffer and then equilibrated with deionized water. The title compound was eluted with a 0 to 0.5 M gradient (2 L) of ammonium bicarbonate. The appropriate fractions were combined and the volatiles removed by evaporation *in vacuo*. The residue was redissolved in deionized water (20 mL) and evaporated *in vacuo*.

three times. Lyophilization of the residue from water gave the ammonium salt of the title compound (270 mg, 90% as mono ammonium salt, monohydrate); $^1\text{H-NMR}$ (D_2O) δ 8.10 (s, 2H), 6.14 – 6.20 (m, 1H), 5.84 – 5.90 (m, 1H), 5.48 – 5.58 (m, 1H), 3.71 – 3.86 (m, 2H), 3.02 – 3.14 (br m, 1H), 2.66 – 2.80 (m, 1H), 1.56 – 1.66 (m, 1H); $^{31}\text{P-NMR}$ (D_2O) 0.62. Mass spectrum (ES^-) m/e 310 (M-H).

Example 45

(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-diphosphate

The title molecule was prepared by a modification of the method of Broom (Mishra, N. C. and Broom, A. D., J. Chem. Soc., Chem. Commun., 1991, 1276). (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol (Example 1, 231 mg, 1.0 mmol) was dissolved in trimethyl phosphate (5 mL). Phosphoryl chloride (0.27 mL, 3.0 mmol) was added to the stirred, cooled (0°C) solution. After 3 hours, a solution of 86% phosphoric acid (1.14 g, 10 mmol) and tri-*n*-butylamine (2.4 mL, 10 mmol) in dimethylformamide (15 mL) was added to the cold (0°C) stirring solution, followed immediately by tri-*n*-butylamine (3 mL). After 1 minute, 1 M triethylammonium hydrogen carbonate buffer (100 mL) was added and stirring was continued for 30 minutes. The reaction solution was diluted to 3 L with deionized water and applied to a 1.1 x 7.0 cm DEAE Sephadex A25 (Aldrich) ion exchange chromatography column which had been washed with 1.0 M ammonium bicarbonate and preequilibrated with deionized water. The title compound was eluted with a 0 to 0.5 M gradient (4 L) of ammonium bicarbonate. The appropriate fractions were combined and the volatiles removed by evaporation *in vacuo*, redissolved in water and evaporated again to give the ammonium salt of the title compound (0.37 mmol, 37%); UV (0.1 M HCl) λ_{max} = 260 nm. UV purity (254 nm detection) was 100% by analytical strong anion exchange HPLC (Whatman Partisil 5, SAX RAC II, 0.05 M to 0.95 M gradient with ammonium phosphate buffer (pH 5.5), 5% methanol). $^1\text{H-NMR}$ (D_2O) δ 8.06 (s, 1H), 8.05 (s, 1H), 6.13 – 6.21 (m, 1H), 5.82 – 5.88 (m, 1H), 5.45 – 5.53 (m, 1H), 3.76 – 3.90 (m, 2H), 3.02 – 3.16 (br m, 1H), 2.64 – 2.78 (m, 1H), 1.52 – 1.64 (m, 1H); $^{31}\text{P-NMR}$ (D_2O) -9.90 (d, 1P, $J=20.3$ Hz), -10.84 (d, 1P, $J=20.3$). Mass spectrum (ES^-) m/e 390 (M-H).

Example 46(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-triphosphate

Continued elution of the column described in Example 45 gave, on evaporation, the
5 ammonium salt of the title compound (11.4 mmoles, 11.4%). UV (0.1 M HCl) λ max = 260
nm. UV purity (254 nm detection) was 98% by analytical strong anion exchange HPLC
(Whatman Partisil 5, SAX RAC II, 0.05 M to 0.95 M gradient with ammonium phosphate
buffer (pH 5.5), 5% methanol). ¹H-NMR (D₂O) δ 8.10 (s, 2H), 6.18 – 6.23 (m, 1H), 5.84 –
5.90 (m, 1H), 5.41 – 5.59 (m, 1H), 3.82 – 3.98 (m, 2H), 3.04 – 3.18 (br m, 1H), 2.54 – 2.66 (m,
10 1H), 1.52 – 1.64 (m, 1H); ³¹P-NMR (D₂O) –8.38 (m, 1P), 10.85 (d, 1P, J=19.3 Hz), –22.73 (m,
1P). Mass spectrum (ES[–]) *m/e* 470 (M–H).

Example 47Anti-HIV Activity

15

Compounds according to the invention were tested for anti-HIV activity in MT₄ cells
according to the method described by Averett, D.R., *J. Virol. Methods*, 23, 1989, 263–276.
Activity of the compounds was in the range of IC₅₀ 0.009 – 2.1 μ M.

20 Example 48Anti-Hepatitis B Virus Activity

Compounds were tested for anti- Hepatitis B Virus activity according to the method
described by Jansen, R. et al., *Antimicrobial Agents and Chemotherapy*, Vol. 37, No. 3, pp.
25 441–447, 1993, and results are shown in the table below. IC₅₀ values of the compounds
according to the invention demonstrated improved activity by as much as 500-fold over
that of the corresponding nucleoside analog, (1R,cis)-4-(6-amino-9H-purin-9-yl)-2-
cyclopentene-1-methanol.

30

Example	IC ₅₀ (mM) vs. HBV	Selectivity Index
2	0.018	1300
4	0.033	6100
5	0.0075	1250
6	0.0045	2000
7	0.0032	5000
8	0.11	>1800
9	0.002	5000
1	0.98	>200

Example 49: Tablet Formulation

The following formulations A, B and C are prepared by wet granulation of the ingredients with a solution of povidone, followed by addition of magnesium stearate and compression.

5 Formulation A

	<u>mg/tablet</u>
Active Ingredient	250
Lactose B.P.	210
Povidone B.P.	15
10 Sodium Starch Glycollate	20
Magnesium Stearate	5
	500

Formulation B

	<u>mg/tablet</u>
Active Ingredient	250
Lactose B.P.	150
5 Avicel PH 101	60
Povidone B.P.	15
Sodium Starch Glycollate	20
Magnesium Stearate	5
10	500

Formulation C

	<u>mg/tablet</u>
Active Ingredient	250
15 Lactose B.P.	200
Starch	50
Povidone	5
Magnesium Stearate	4
20	359

The following formulations, D and E, are prepared by direct compression of the admixed ingredients. The lactose in formulation E is of the direct compression type (Dairy Crest-"Zeparox").

25

Formulation D

	<u>mg/tablet</u>
Active Ingredient	250
Pregelatinized Starch NF15	150
30	
	400

Formulation E

	<u>mg/tablet</u>
Active Ingredient	250
Lactose B.P.	150
5 Avicel	100
	500

Formulation F (Controlled Release Formulation)

10

The formulation is prepared by wet granulation of the ingredients with a solution of povidone followed by the addition of magnesium stearate and compression.

	<u>mg/tablet</u>
15 Active Ingredient	500
Hydroxypropylmethylcellulose	112
(Methocel K4M Premium)	
Lactose B.P.	53
Povidone B.P.	28
20 Magnesium Stearate	7
	700

Drug release takes place over a period of about 6-8 hours and is complete after 12 hours.

25

Example 50: Capsule FormulationsFormulation A

A capsule formulation is prepared by admixing the ingredients of formulation D in

30 Example 49 above and filling into a two-part hard gelatin capsule. Formulation B (infra) is prepared in a similar manner.

Formulation B

	<u>mg/capsule</u>
Active Ingredient	250
Lactose B.P.	143
5 Sodium Starch Glycollate	25
Magnesium Stearate	2
	420

10 Formulation C

	<u>mg/capsule</u>
Active Ingredient	250
Macrogel 4000 B.P.	350
15	600

Capsules of formulation C are prepared by melting the Macrogel 4000 B.P., dispersing the active ingredient in the melt and filling the melt into a two-part hard gelatin capsule.

20 Formulation D

	<u>mg/capsule</u>
Active Ingredient	250
Lecithin	100
Arachis Oil	100
25	450

Capsules of formulation D are prepared by dispersing the active ingredient in the lecithin and arachis oil and filling the dispersion into soft, elastic gelatin capsules.

Formulation E

	<u>mg/capsule</u>
Active Ingredient	150.0
Vitamin E TPGS	400.0
5 Polyethylene Glycol 400 NF	200.5
Propylene Glycol USP	39.5

Four (4) kilograms (kg) of Vitamin E TPGS (obtained from Eastman Chemical Co.) was heated at 50°C until liquefied. To the liquefied Vitamin E TPGS, 2.005 kg of polyethylene glycol 400 (PEG400) (low aldehyde, <10 ppm, obtained from Union Carbide or Dow Chemical Co.) heated to 50°C was added and mixed until a homogeneous solution was formed. The resultant solution was heated to 65°C. 1.5 kg of active ingredient was dissolved in the liquefied solution of Vitamin E TPGS and PEG 400. 0.395 kg of propylene glycol at room temperature was added and mixed until a homogenous solution was formed. The solution was cooled to 28-35°C. The solution was then de-gassed. The mixture was preferably encapsulated at 28-35°C at a fill weight equivalent to 150 mg of volatiles-free compound, into Size 12 oblong, white opaque soft gelatin capsules using a capsule filling machine. The capsule shells were dried to a constant fill moisture of 3-6% water and a shell hardness of 7-10 Newtons, and placed in a suitable container.

20

Formulation F (Controlled Release Capsule)

The following controlled release capsule formulation is prepared by extruding ingredients a, b, and c using an extruder, followed by spheronization of the extrudate and drying. The dried pellets are then coated with release-controlling membrane (d) and filled into a two-piece, hard gelatin capsule.

25

30

		<u>mg/capsule</u>
	(a) Active Ingredient	250
	(b) Microcrystalline Cellulose	125
	(c) Lactose B.P.	125
5	(d) Ethyl Cellulose	13
		513

Example 51: Injectable Formulation

10

Formulation A

		<u>mg</u>
	Active Ingredient	200
	Hydrochloric Acid Solution 0.1M or	
15	Sodium Hydroxide Solution 0.1M q.s. to pH	4.0 to 7.0
	Sterile water q.s. to	10 ml

The active ingredient is dissolved in most of the water (35° - 40° C) and the pH adjusted to between 4.0 and 7.0 with the hydrochloric acid or the sodium hydroxide as appropriate.

20 The batch is then made up to volume with water and filtered through a sterile micropore filter into a sterile 10 ml amber glass vial (type 1) and sealed with sterile closures and overseals.

Formulation B

25

	Active Ingredient	125 mg
	Sterile, Pyrogen-free, pH 7 Phosphate	
	Buffer, q.s. to	25 ml

30

Example 52: Intramuscular Injection

	Active Ingredient	200 mg
	Benzyl Alcohol	0.10 g
5	Glycofurol 75	1.45 g
	Water for injection q.s. to	3.00 ml

The active ingredient is dissolved in the glycofurol. The benzyl alcohol is then added and dissolved, and water added to 3 ml. The mixture is then filtered through a sterile
 10 micropore filter and sealed in sterile 3ml amber glass vials (type 1).

Example 53: Syrup

	Active Ingredient	250 mg
15	Sorbitol Solution	1.50 g
	Glycerol	2.00 g
	Sodium Benzoate	0.005 g
	Flavor, Peach 17.42.3169	0.0125 ml
	Purified Water q.s. to	5.00 ml

20

The active ingredient is dissolved in a mixture of the glycerol and most of the purified water. An aqueous solution of the sodium benzoate is then added to the solution, followed by addition of the sorbitol solution and finally the flavor. The volume is made up with purified water and mixed well.

25

Example 54: Suppository

		<u>mg/capsule suppository</u>
	Active Ingredient	250
	Hard Fat, B.P. (Witepsol H15-Dynamit Nobel)	1770
30		
		2020

One-fifth of the Witepsol H15 is melted in a steam-jacketed pan at 45°C maximum. The active ingredient is sifted through a 200µm sieve and added to the molten base with mixing, using a Silverson fitted with a cutting head, until a smooth dispersion is achieved. Maintaining the mixture at 45° C, the remaining Witepsol H15 is added to the suspension
 5 and stirred to ensure a homogenous mix. The entire suspension is passed through a 250µm stainless steel screen and, with continuous stirring, is allowed to cool to 45° C. At a temperature of 38° C to 40° C, 2.02 g of the mixture is filled into suitable, 2ml plastic molds. The suppositories are allowed to cool to room temperature.

10 Example 55: Pessaries

	<u>mg/pessary</u>
) Active Ingredient	250
Anhydrate Dextrose	380
Potato Starch	363
15 Magnesium Stearate	7
	1000

The above ingredients are mixed directly.

20

Example 56: Acid Stability

Compounds according to the invention were tested for their stability towards acid-mediated hydrolytic decomposition employing a test designed to simulate stomach
 25 conditions. Each compound was incubated at an initial concentration of 0.3 mg/ml in dilute hydrochloric acid at pH 1 at 37°C. HPLC was run immediately for t=0 and at intervals up to approximately 24 hours. The half-life of title compound from Example 7 was 76 hours under these conditions. Comparative phosphoramidates of 2',3'-dideoxy-adenosine (Compound 1093) and 2',3'-didehydro-2',3'-dideoxy-adenosine (Compound
 30 1001), described in PCT/GB96/00580, were significantly less stable at pH1. Compound 1001 was completely decomposed in <1 minute at pH 1 (25°C). Compound 1093 was completely decomposed after 13 hours at pH 1 (25°C).

Example 57: Biological Stability


Title compound of Example 7 and phosphoramidates of 2',3'-dideoxy-adenosine
5 (Compound 1093) and 2',3'-didehydro-2',3'-dideoxy-adenosine (Compound 1001),
described in PCT/GB96/00580, were tested for their stability towards biological
decomposition. Each compound was incubated in normal heparized human plasma at
37°C. At selected time points duplicate samples were removed and deproteinated by
acetonitrile extraction. Drug concentrations were then determined by LC/MS/MS analysis
10 using standard methods. Half-lives were calculated and are shown in the table below.

<u>Compound</u>	<u>Half-life (hours)</u>
Example 7	53
1001	4.6
1093	4.2

15 The half-life in human plasma of compound of the present invention is more than 10-fold
greater than those of compounds 1001 and 1093.

Claims

1. A compound of formula (I)



(I)

wherein:

- 10 R¹ is hydrogen; C₆₋₁₄aryl; or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of C₁₋₆alkoxy, nitro, halogen, amino, hydroxy, carboxylate and esters thereof, carboxyalkyl, -CONHR⁶, and -CONR⁶R⁷, wherein R⁶ and R⁷, which may be the same or different, are independently selected from C₁₋₈alkyl, C₁₋₈alkylaryl or C₆₋₁₄aryl;
- 15 R² and R³ are independently selected from hydrogen or C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, C₆₋₁₄aryl, or aralkyl wherein each C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, C₆₋₁₄aryl or aralkyl may be optionally substituted with one or more substituents selected from the group consisting of C₁₋₈alkyl, halo, hydroxy, alkoxy, amino, aminoalkyl, aminodialkyl, -SH, thioalkyl, heterocycle, carboxylate and esters thereof,
- 20 carboxyalkyl, -CONHR⁶, and -CONR⁶R⁷, wherein R⁶ and R⁷, which may be the same or different, are independently selected from C₁₋₈alkyl, C₁₋₈alkylaryl or C₆₋₁₄aryl; or R² and R³ can together form a 3 to 8-membered ring;
- 25 R⁴ is -OR⁸, -NR⁸R⁹ or -SR⁸, where R⁸ and R⁹, which may be the same or different, are independently selected from hydrogen; or C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, aralkyl, heteroaryl, or C₆₋₁₄aryl wherein each C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₅₋₈cycloalkenyl, aralkyl, heteroaryl, or C₆₋₁₄aryl may be optionally substituted with


one or more substituents selected from the group consisting of halo, hydroxy, alkoxy, amino, aminoalkyl, aminodialkyl, -SH, thioalkyl, carboxylate and esters thereof, carboxyalkyl, -CONHR⁶, and -CONR⁶R⁷, wherein R⁶ and R⁷, which may be the same or different, are independently selected from C₁₋₈alkyl, C₁₋₈alkylaryl or C₆₋₁₄aryl;

5

R⁵ is hydrogen; C₁₋₈alkyl; or C₆₋₁₄aryl; or R² and R⁵ may together form a 5- or 6-membered ring or R³ and R⁵ may together form a 5- or 6-membered ring;

10 or a pharmaceutically acceptable derivative thereof, and their use in the treatment of viral infections.

2. A compound of formula (II)



15

wherein:

R¹ is hydrogen; C₆₋₁₄aryl; or heteroaryl, optionally substituted with one or more substituents selected from C₁₋₆alkoxy, nitro, halogen, amino, carboxylate and hydroxy.


20

R² and R³ are independently selected from hydrogen, C₁₋₆alkyl, C₃₋₈cycloalkyl, C₆₋₁₄aryl, or aralkyl;

25 R⁴ is OR¹⁰, NHR¹⁰ or SR¹⁰, where R¹⁰ is hydrogen, C₁₋₆alkyl, C₃₋₈cycloalkyl, or aralkyl; or NHR¹¹ wherein R¹¹ is C₁₋₆alkyl, aralkyl, or C₆₋₁₄aryl;

or a pharmaceutically acceptable derivative thereof.


3. A compound of formula (II) according to claim 2 wherein R^1 is H or C_{6-14} aryl, R^2 is C_{1-6} alkyl or aralkyl, R^3 is hydrogen, C_{1-6} alkyl or aralkyl and R^4 is OR^{10} wherein R^{10} is C_{1-6} alkyl or C_{3-8} cycloalkyl.
4. A compound of formula (II) according to claim 2 wherein R^1 is C_{6-14} aryl, R^2 is methyl, R^3 is hydrogen, and R^4 is OR^{10} where R^{10} is methyl or ethyl.
5. A compound of formula (II) according to claim 2 wherein R^1 is hydrogen.
6. A compound of formula (II) according to claim 2 wherein R^1 is C_{6-14} aryl.
7. A compound of formula (I) according to claim 1 wherein R^2 and R^3 are not both hydrogen.
8. A compound of formula (II) according to claim 2, which is (1R,4S)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-alaninyl)phosphate or a pharmaceutically acceptable derivative thereof.
9. A compound of formula (III)



(III)

29 JUL 2003

wherein R¹² is



wherein n is 0, 1, or 2, and wherein R¹² is optionally substituted by C₆₋₁₄aryl.

10. A compound as claimed in claim 1 selected from the group consisting of
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl benzyloxy-L-alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-phenylalaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-phenylalaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(L-phenylalaninyl)phosphoramidate Disodium Salt;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy- α , α ,-dimethylglycinyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-L-(O-tert-butyltyrosinyl)]phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy-L-alaninyl)phosphoramidate;
 - (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl tert-butoxy-L-alaninyl)phosphoramidate;

- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy)-L-alaninyl]phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl cyclopropylmethoxy-L-alaninyl)phosphoramidate;
- 5 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-leucinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-L-(O-methyltyrosinyl)]phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-phenylalaninyl)phosphoramidate;
- 10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-leucinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-D-tryptophanyl]phosphoramidate;
- 15 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl methoxy-D-tryptophanyl]phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-valinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl dimethoxy-D-aspartyl)phosphoramidate;
- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-prolinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-isoleucinyl)phosphoramidate;
- 25 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-norvalinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy-L-phenylalaninyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl *tert*-butoxy-L-phenylalaninyl)phosphoramidate;
- 30 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy)-L-phenylalaninyl]phosphoramidate;


- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl benzyloxy-L-phenylalaninyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[(2-carbomethoxy)phenyl methoxy-L-alaninyl]phosphoramidate;
- 5 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[4-(3-oxo-3-phenylpropenyl)phenyl methoxy-L-alaninyl]phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxyglycinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-valinyl)phosphoramidate;
- 10 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl(methoxy)-L-methioninyl]phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy- α , α -dimethylglycinyl)phosphoramidate;
- 15 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl isopropoxy- α , α -dimethylglycinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[phenyl (2,2-dimethylpropoxy)- α , α -dimethylglycinyl]phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-cyclopentaneglycinyl)phosphoramidate;
- 20 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-cyclohexaneglycinyl)phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl dimethoxy-L-aspartyl)phosphoramidate;
- 25 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[(4-chloro)phenyl methoxy-L-alaninyl]phosphoramidate;
- (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-[L-(N-methyl)amino)alaninyl]phosphoramidate sodium salt;
- and pharmaceutically acceptable derivatives thereof.

11. A compound as claimed in claim 10 selected from the group consisting of
(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-alaninyl)phosphoramidate;
5 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-alaninyl)phosphoramidate;
(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-D-alaninyl)phosphoramidate;
(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl benzyloxy-L-
10 alaninyl)phosphoramidate;
(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy-L-phenylalaninyl)phosphoramidate;
(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-phenylalaninyl)phosphoramidate;
15 (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(L-phenylalaninyl)phosphoramidate Disodium Salt;
(1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl methoxy- α , α -dimethylglyciny)phosphoramidate;
and pharmaceutically acceptable derivatives thereof.
- 20
12. A compound according to any of claims 1 to 11 in the form of a single isomer.
13. A compound according to any of claims 1 to 11 in the form of a mixture of diastereomers.
- 25
14. A method of treating a virus infection in a human comprising administering to said human an effective anti-virus treatment amount of a compound of formula (I), (II) or (III) according to any of claims 1 to 13 or a pharmaceutically acceptable derivative thereof.
- 30
15. The method according to claim 14 wherein the virus is selected from Human Immunodeficiency Virus and hepatitis B virus.


16. A method of treating a hepatitis B virus infection in a human comprising administering to said human an effective anti-hepatitis B treatment amount of (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol O-(phenyl ethoxy-L-alaninyl)phosphate or of (1R,cis)-4-(6-Amino-9H-purin-9-yl)-2-cyclopentene-1-methanol
5 O-(phenyl ethoxy-L-phenylalaninyl)phosphate or a pharmaceutically acceptable derivative thereof.
17. A compound according to any of claims 1 to 13 wherein the pharmaceutically acceptable derivative is a salt.
- 10 18. A pharmaceutical composition comprising an effective anti-viral amount of a compound of formula (I), (II), or (III) or a pharmaceutically acceptable derivative thereof together with a pharmaceutically acceptable carrier therefor.
- 15 19. The pharmaceutical composition according to claim 18, further comprising an antiviral agent other than a compound of formula (I), (II), or (III).
20. A pharmaceutical composition according to claim 18 or 19 in the form of a tablet or capsule.
- 20 21. A pharmaceutical composition according to claim 18 or 19 in the form of a solution, suspension, or syrup.
- 22. Use of a compound of formula (I), (II) or (III) according to any of claims 1 – 13 in
25 the manufacture of a medicament for the treatment or prophylaxis of a virus infection.
23. A compound of formula (I), (II), or (III) as claimed in any of claims 1 to 13 for use in medical therapy.

29 JUL 2003

24. A method of delivering a compound of formula (III)



wherein R¹² is




wherein n is 0, 1, or 2, and wherein R¹² is optionally substituted by C₆₋₁₄aryl,


into cells by treating said cells with a compound of formula (I) or (II) as defined in any of claims 1 to 8.

25. A patient pack comprising at least one active ingredient selected from a compound of formula (I), (II) or (III) according to any of claims 1 to 13.

26. A process for the preparation of a compound of formula (I) as defined in claim 1, said process comprising the reaction of a compound of formula (IV)



with a compound of formula (VI)




(VI)

wherein R¹ – R⁵ are as hereinbefore defined for formula (I).

5


27. A process as claimed in claim 26 wherein the solvent for the reaction is pyridine, pyridine-tetrahydrofuran or acetonitrile in the presence of t-butyl magnesium chloride.

10 28. A process for the preparation of a compound of formula (II) as defined in claim 2, said process comprising the reaction of a compound of formula (IV)



(IV)

with a compound of formula (V)



(V)

15

wherein $R^1 - R^4$ are as hereinbefore defined for formula (II).

5 29. A process as claimed in claim 28 wherein the solvent for the reaction is pyridine, pyridine-tetrahydrofuran or acetonitrile in the presence of t-butyl magnesium chloride.

10 30. A process for the preparation of a compound of formula (III), as defined in claim 9, said process comprising treating a compound of formula (IV) as hereinbefore defined with an appropriate phosphorylating agent.

15 31. A process as claimed in claim 30 wherein the phosphorylating agent is phosphoryl chloride.

32. A process as claimed in any of claims 26 to 31 wherein the compound obtained is in the form of a single isomer, a mixture of diastereomers, or a salt.

20