USE OF DES-ASPARTATE-ANGIOTENSIN I

Field of the invention

The present invention relates to the use of des-aspartate-angiotensin I (des-asp-ang I), its derivative, a functional part and/or analogue thereof in medicine.

Background of the invention

Viral respiratory tract infection is the most common illness in humans and the economic cost for non-influenza-related viral respiratory tract infection approaches $40 billion in the US (Fendrick AM *et al*, 2003), while influenza causes tens of millions of day of restricted activity, bed disability and work loss (Sullivan KM *et al*, 1993). There is also the ever present threat of a pandemic flu outbreak and the apparent unstoppable AIDS virus spreading far and wide. With the ability of viruses to mutate by genetic drift and genetic shift, new viruses that are not susceptible to existing vaccines will continue to come into being. There is, therefore, a need to develop more effective antiviral drugs.

Another medical condition, prolonged or chronic hyperglycemia, wherein blood glucose levels are elevated from long periods, produces many detrimental effects. This condition is exemplified by diabetes. Diabetes exacts a huge toll in money and human suffering, accounting for more than 100 billion dollars of healthcare costs annually in the United States (DiRamond J, 2003). The number of cases worldwide is estimated at 150 million with an equal number of undiagnosed cases in the First World countries and eight times more undiagnosed cases in the Third World (Levitt NS *et al*, 1993). The disease is characterised by high blood sugar (glucose) resulting from the defects in either insulin secretion or decreased sensitivity of the body's cells to the action of

insulin, leading to a condition of hyperglycemia. Type I diabetes is currently treated by preprandial administration of exogenous insulin and dietary restriction. Current therapy of type 2 diabetes includes lifestyle modifications and the use of a variety of pharmacological agents that aim to increase insulin secretion, decrease hepatic glucose production, and/or increase insulin actions. Despite these approaches, good glycemic control is beyond the reach of most diabetic individuals, and the state of prolonged or chronic hyperglycemia can cause cardiovascular diseases, stroke, blindness, kidney failure, neurological dysfunction, necrosis and gangrene of extremities. The search for better and more specific agents with physiological properties for hyperglycemia is highly warranted.

Accordingly, any new and improved treatment of viral infections and/or conditions relating or due to hyperglycemia will be welcome.

The compound des-aspartate-angiotensin I is an endogenous angiotensin peptide (Sim Mk *et al*, 1994, a,b). It is formed or derived from angiotensin I by a specific aminopeptidase present in blood vessels and the hypothalamus (Sim MK, Qui XS, 1994). Previous studies suggested the use of des-aspartate-angiotensin I in cardiovascular and renal actions (Sim US. 5773415, 6100237, 6589938B2 and US2003/0086920). However, no further uses are taught or suggested for des-aspartate-angiotensin I.

Summary of the invention

The present invention addresses the problems above, and in particular, provides new and effective means of treatment and/or prophylaxis of viral infections and hyperglycemia-related conditions. In particular, the present inventor has surprisingly found that at least one derivative of angiotensin I may be used in inducing hypoglycemia and/or for reducing hyperglycemia and/or for

the treatment of hyperglycemia-related conditions as well as in the treatment and/or prophylaxis of viral infections. More in particular, the at least one derivative of angiotensin I is des-aspartate-angiotensin I, its derivative, a functional part, and/or an analogue thereof.

Accordingly, in one aspect, the present invention provides a method for the treatment and/or prophylaxis of at least one viral infection comprising administering to a subject des-aspartate-angiotensin I, its derivative, a functional part and/or an analogue thereof.

In another aspect, the present invention provides a method of inducing hypoglycaemia and/or reducing hyperglycaemia in a subject, the method comprising administering to the subject des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

There is also provided a method of treatment and/or prophylaxis of at least one hyperglycaemia-related condition comprising administering to a subject des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, provided the at least one hyperglycaemia-related condition is not a renal-related disorder.

There is also provided des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, and/or a pharmaceutical composition thereof, for use in the treatment and/or prophylaxis of at least one viral infection.

There is also provided des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, and/or a pharmaceutical composition thereof, for use in the induction of hypoglycaemia and/or reduction of hyperglycaemia. There is also provided des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, and/or a pharmaceutical composition thereof, for

use in the treatment and/or prophylaxis of at least one hyperglycaemia-related condition, provided the at least one hyperglycaemia-related condition is not a renal-related disorder.

There is also provided the use of des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof for the preparation of a medicament for the treatment and/or prophylaxis of at least one viral infection; for the induction and/or reduction of hyperglycaemia; as well as for the treatment and/or prophylaxis of at least one hyperglycaemia-related condition, provided the at least one hyperglycaemia-related condition is not a renal-related disorder.

In another aspect, the present invention provides a kit for the treatment and/or prophylaxis of at least one viral infection comprising des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

In another aspect, the present invention provides a kit for the induction of hypoglycaemia and/or reduction of hyperglycaemia comprising des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof. There is also provided a kit for the treatment and/or prophylaxis of at least one hyperglycaemia-related condition provided the at least one hyperglycaemia-related condition is not a renal-related disorder, the kit comprising des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

Under the relevant aspects above, the at least one viral infection may be a respiratory viral infection. In particular, the viral infection may be an influenza virus infection such as an influenza A infection.

The at least one hyperglycaemia-related condition may be type I diabetes, obesity, and/or bulimia nervosa.

The des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, may be administered in a pharmaceutically and/or therapeutically effective amount. The des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, may comprise at least one pharmaceutically-acceptable carrier, excipient, diluent and/or adjuvant.

The kits may further comprise information and/or instructions pertaining to their use.

Brief description of the figures and tables

Figures:

Figure 1 shows the effect of des-aspartate-angiotensin I on weight loss and survival of influenza A virus-infected female BALB/c mice. Six to 7 week-old mice were intranasally infected with 50 μL passage-6 lung homogenate (equivalent to 2.5 x 105 TCID50 of influenza A in MDCK cells). Group of 5-9 mice were then orally administered with one of the following doses of des-aspartate-angiotensin I:- 300 nmole/kg/dat for 8 days (n = 5), 600 nmole/kg/day for 8 days (n = 9), and 120 nmole/kg administered on Day 2 post-infection (n = 9, only one dose was administered). The control group (n = 8) was similarly administered water for 8 days. The weight and survival of the mice were recorded daily for 16 days.

Figures 2 to 6 show the effect of des-aspartate-angiotensin I on blood glucose profile animal models of diabetes.

Figure 2 shows the effect of des-aspartate-angiotensin I on blood glucose profile in diabetic KK-Ay mice. The diabteic KKAy mice were divided into four groups consisting of 6 animals per group. Animals in the control group were intraperitoneally administered 0.1 mL saline. Animals in the second, third, and

fourth groups were similarly administered with 100, 200, 400 nmole/kg des-aspartate-angiotensin I, respectively. Treatment with saline and des-aspartate-angiotensin I was carried out daily for 4 weeks. Following this, animals were fasted overnight for 16 hours and oral glucose tolerance test was performed as follows: blood was withdrawn from the orbital sinus for blood glucose determination (time of withdrawal was designated as 0 time), animals were then orally administered glucose (2g/kg), and blood was withdrawn at 30, 60 and 120 min for blood glucose determination. The blood was allowed to clot and blood glucose was measured as serum glucose using a commercial glucose kit from Thermo Electron Corporation, Australia. *Significantly different from the corresponding values of the control untreated mice ($p < 0.05$, ANOVA followed by post hoc Tukey test).

Figure 3 shows the effect of orally administered des-aspartate-angiotensin II on blood glucose profile in diabetic GK rats at 4 and 6 weeks of treatment. The experiment was similarly carried as for KKAy mice. Control animals were administered (by gavage) 0.2 mL of water and groups 1 and 2 animals were similarly administered 400 and 600 nmole/kg des-aspartate-angiotensin I in 0.2 mL water, respectively, for a period 8 weeks. Oral glucose tolerance test (determined at 30 min) was carried out after 4 and 6 weeks of treatment. N = 5 to 6. *Significantly different from the corresponding values of the control untreated rat ($p < 0.05$, ANOVA followed by post hoc Tukey test). Data obtained after 8 week of treatment are given in another figure, Fig. 4. *Note*: The data show that orally administered DAA-I exerts hypoglycaemic action in diabetic GK rats after 6 weeks of treatment.

Figure 4 shows the effect of orally administered des-aspartate-angiotensin I on blood glucose profile in diabetic GK rats at 8 weeks of treatment. This figure gives the data of a full oral glucose tolerance test conducted at 8 weeks of the same experiment described in Fig. 3. *Significantly different from the value of the corresponding control ($p<0.05$, ANOVA post hoc Tukey test).

Figure 5 shows the Effects of des-aspartate-angiotesnin I on insulin-induced translocation of GLUT-4 in skeletal muscle of diabetic GK rats. Upper panel: Representative Western blot of plasma membrane GLUT4 in skeletal muscle of des-aspartate-angiotensin I treated and non treated diabetic GK rats. Lower panel: Relative (to the non insulin stimulated samples) levels of plasma membrane GLUT4 in skeletal muscle of des-aspartate-angiotensin I treated and control diabetic GK rats. N =3 for each histogram. *Significantly different from the value of the corresponding control ($p<0.05$, ANOVA post hoc Tukey test).

Figure 6 shows the effect of des-aspartate-angiotesnin I on serum insulin level of diabetic GK rats. Serum from 8 weeks treated animals (as described in EXAMPLE 9) were assayed for insulin concentration by the "Ultra Senstive Rat Insulin ELISA kit' (Crystal Chem Inc., IL, USA). The values were from serum taken at 30 min of the oral glucose tolerance test. There were no significant difference between the values of the control and treated group.

Tables:

Table 1 shows some examples of unnatural amino acids contemplated by the present invention .

Tables 2 and 3 show the effect of des-aspartate-angiotensin I in animal models of influenza.

Table 2 shows the percentage weight change and survival of influenza A virus-infected BALB/c mice treated with orally-administered des-aspartate-angiotensin I

Table 3 shows the percentage weight change and survival of influenza A virus-infected BALB/c mice treated with intraperitoneally-administered des-aspartate-angiotensin I

Tables 4 and 5 show the effects of des-aspartate-angiotensin I in control of blood glucose levels in animal models of diabetes.

Table 4 shows the data of the oral glucose tolerance test carried out in diabetic KKAy mice that were treated with intraperitoneally-administered des-aspartate-angiotensin I

Table 5 shows data of oral the glucose tolerance test carried out in diabetic KKAy mice that were treated with orally-administered des-aspartate-angiotensin I.

Detailed description of the invention

Bibliographic references mentioned in the present specification are for convenience listed in the form of a list of references and added at the end of the examples. The whole content of such bibliographic references is herein incorporated by reference.

Des-aspartate angiotensin I have been previously described for use in treatment of certain disorders. For example, the effect of des-aspartate angiotensin I in segmental glomerulosclerosis rat model resembling renal lesions in humans arising from various disorders including those arising from diabetus mellitus was previously determined (US2003/0086920).

However, the prior art does not teach or suggest use of des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof for the treatment and/or prophylaxis of viral infections and/or for the induction of hypoglycaemia and/or reduction of hyperglycemia-related conditions. In particular, the present invention relates to the use of des-asparatate-angiotensin I in the treatment of hyperglicemia-related condition(s), excluding renal-related disorders. In particular, the condition(s) treated according to the present invention do not include renal-related disorders described in US2003/0086920.

The present invention relates to the use of des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof for the treatment and/or prophylaxis of viral infections and/or for the induction of hypoglycaemia and/or the reduction of hyperglycaemia. In particular, the present invention relates to the use of des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof for the treatment and/or prophylaxis of influenza A infection and hyperglycaemia-related conditions as type I diabetes, obesity, and/or bulimia nervosa.

While researching the effects of des-aspartate-angiotensin I on the cardiopulmonary functions of influenza A virus-infected mice, it was surprisingly discovered that des-aspartate-angiotensin I increased the survival of the animals.

Similarly, it was surprisingly discovered that des-aspartate-angiotensin I exerts marked hypoglycaemic effects in animal models of hyperglycaemia. Accordingly, the present invention provides new treatments, prophylaxes and/or pharmaceutical compositions for viral infections and/or hyperglycemia-related conditions using its des-aspartate-angiotensin I, its derivative, functional part, and/or analogue thereof.

Treatment of viral infections

The present invention provides a method for the treatment and/or prophylaxis of at least one viral infection and its symptoms comprising administering to a subject des-aspartate-angiotensin I, its derivative, a functional part and/or an analogue thereof. In particular, the des-aspartate-angiotensin I may be administered in a therapeutically and/or pharmaceutically effective amount. More in particular, the method may comprise administering to a subject an therapeutically and/or pharmaceutically effective amount of at least one derivative of angiotensin I, according to the present invention, for a time and

under conditions sufficient for the onset of the viral infection and/or its symptoms to be prevented, inhibited and/or delayed or the symptoms of the viral infection to be ameliorated. The derivative may be des-aspartate-angiotensin I, or a derivative, homologue and/or analogue thereof.

The at least one viral infection may be a respiratory viral infection. In particular, the viral infection may be an influenza virus infection such as an influenza A infection.

Induction of hypoglycaemia/reduction of hyperglycemia

In another aspect, the present invention also provides a method of inducing hypoglycaemia and/or reducing hyperglycemia in a subject, the method comprising administering to the subject des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

In particular, the present invention provides a method of treatment and/or prophylaxis of at least one hyperglycaemia-related condition comprising administering to a subject des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, wherein the at least one hyperglycaemia-related condition excludes renal-related disorders.

In particular, the present invention provides a method for the treatment and/or prophylaxis of at least one of the following states or conditions: hyperglycaemia, particularly prolonged or uncontrolled hyperglycaemic conditions as in type I diabetes, obesity, bulimia nervosa excluding renal disorders arising from hyperglycaemia as in Type 2 diabetes (diabetes mellitus), the method comprising administering to a subject at least one derivative of angiotensin I. In particular, the at least one derivative of angiotensin I is des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof. In particular, the angiotensin I derivative is administered for a time and under

conditions sufficient for the condition and/or its symptoms to be prevented, inhibited, delayed and/or ameliorated.

The present invention also provides des-aspartate-angiotensin I or a pharmaceutical composition threfore for use in the treatment and/or prophylaxis of hyperglcaemia, in particular prolonged or uncontrolled hyperglycaemic conditions as in type I diabetes, obesity, bulimia nervosa excluding renal disorders arising from hyperglycemia as in Type 2 diabetes mellitus, The pharmceutical composition may comprise at least one derivative of angiotensin I according to the present invention and, optionally, a pharmaceutically acceptable carrier, excipient, adjuvant and/or diluent.

The compositions may be administered in a therapeutically and/or pharmaceutically effective dose, and/or for a time and under conditions sufficient for its symptoms to be prevented, inhibited or delayed and/or ameliorated.

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising, will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

An "effective amount" refers to an amount effective, at dosages and for periods of time necessary to achieve the desired therapeutic result, such as to prevent, inhibit and/or delay the onset hyperglycaemia-related disorders and/or ameliorate the symptoms of hyperglycaemia-related disorders. While the effective amount may vary according to various factors such as the disease state, age, sex, and weight of the individual.

A derivative of des-aspartate-angiotensin I includes any mutant, fragment, part and/or portion of angiotensin I or des-aspartate-angiotensin I including

molecules comprising single or multiple amino acid substitutions, deletions and/or additions to angiotensin I or to des-aspartate-angiotensin I. The processes of substitutions, deletions and/or additions may result in a derivative or peptide that has more amino acids than the original angiotensin I or des-aspartate-angiotensin I. The processes of substitutions, deletions and/or additions may result in a derivative or peptide that has more amino acids than the original angiotensin I or des-aspartate angiotensin I

The preferred derivative in accordance with the present invention is des-aspartate-angiotensin I or a derivative, homologue and/or analogue thereof. Reference to a homologue and/or an analogue includes a mimotope or peptide and/or analogue mimetic. Such derivatives, homologues and/or analogues may function in place of des-aspartate-angiotensin I and/or its equivalent or they may act as an agonist thereof or, when necessary, an antagonist thereof.

As stated above, a derivative includes single or multiple amino acid substitutions, additions and/or deletions to des-aspartate-angiotensin I and/or angiotensin I. The final derivative is a peptide having similar functions and consists of no less than two amino acid sequence of the original angiotensin I or des-aspartate-angiotensin.

Amino acid insertional derivatives of des-Asp-angiotensin I include amino and/or carboxyl terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Insertional amino acid sequence variants are those in which one or more amino acid residues are introduced into a predetermined site in the des-aspartate-angiotensin I although random insertion is also possible with suitable screening of the resulting product.

Deletional variants are characterized by the removal of one or more amino acids from the sequence. Substitutional amino acid variants are those in which at least one residue in the sequence has been removed and a different residue inserted in its place.

Where the des-aspartate-angiotensin I is derivatized by amino acid substitution, the amino acids are generally replaced by other amino acids having like properties, such as hydrophobicity, hydrophilicity, electronegativity, bulky side chains and the like. Amino acid substitutions are either of single or multiple residues. Amino acid insertions will usually be in the order of about 1-7 amino acid residues and deletions will range from about 1-7 residues. Preferably, deletions or insertions are made in adjacent pairs, i.e. a deletion of two residues or insertion of two residues.

Homologues include functionally, structurally or stereochemically similar polypeptides from, for example, other sources such as for livestock animals, laboratory test animals or primates. The similar peptides may also be homologues of des-aspartate-angiotensin I.

Analogues and mimetics include molecules which contain non-naturally occurring amino acids as well as molecules which do not contain amino acids but nevertheless behave functionally the same as the des-aspartate-angiotensin I. Analogues contemplated herein include modifications to side chains, incorporation of unnatural amino acids and/or their derivatives during peptide synthesis and the use of crosslinkers and other methods which impose conformational constraints on the peptide molecule or their analogues.

Examples of incorporating unnatural amino acids and derivatives during peptide synthesis include, but are not limited to, use of norleucine, 4-amino butyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, ornithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid, 2-thienyl alanine and/or D-isomers of amino acids.

Crosslinkers can be used, for example, to stabilize 3D conformations, using homo-bifunctional crosslinkers such as the bifunctional imido esters having (CH.2)n spacer groups with n=1 to n=6, glutaraldehyde, N-hydroxysuccinimide esters and heterobifunctional reagents which usually contain anaminoreactive

moiety such as N-hydroxysuccinimide and another group specific-reactive moiety.

All these types of modifications may be important to stabilize the subject des-Asp-angiotensin I. This may be important if used, for example, in the manufacture of a vaccine or therapeutic composition or in detection assays. Examples of unnatural amino acids contemplated by the present invention are presented in Table 1.

Table 1

Non-conventional amino acid	Code	Non-conventional amino acid	Code
α-aminobutyric acid	Abu	L-N-methylalanine	Nmala
α-amino-α-methylbutyrate	Mgabu	L-N-methylarginine	Nmarg
aminocyclopropane-carboxylate	Cpro	L-N-methylasparagine	Nmasn
		L-N-methylaspartic acid	Nmasp
aminoisobutyric acid	Aib	L-N-methylcysteine	Nmcys
aminonorbornyl-carboxylate	Norb	L-N-methylglutamine	Nmgln
		L-N-methylglutamic acid	Nmglu
cyclohexylalanine	Chexa	L-N-methylhistidine	Nmhis
cyclopentylalanine	Cpen	L-N-methylisolleucine	Nmile
D-alanine	Dal	L-N-methylleucine	Nmleu
D-arginine	Darg	L-N-methyllysine	Nmlys
D-aspartic acid	Dasp	L-N-methylmethionine	Nmmet
D-cysteine	Dcys	L-N-methylnorleucine	Nmnle
D-glutamine	Dgln	L-N-methylnorvaline	Nmnva
D-glutamic acid	Dglu	L-N-methylornithine	Nmorn

	D-histidine	Dhis	L-N-methylphenylalanine	Nmphe
	D-isoleucine	Dile	L-N-methylproline	Nmpro
	D-leucine	Dleu	L-N-methylserine	Nmser
	D-lysine	Dlys	L-N-methylthreonine	Nmthr
5	D-methionine	Dmet	L-N-methyltryptophan	Nmtrp
	D-ornithine	Dorn	L-N-methyltyrosine	Nmtyr
	D-phenylalanine	Dphe	L-N-methylvaline	Nmval
	D-proline	Dpro	L-N-methylethylglycine	Nmetg
	D-serine	Dser	L-N-methyl-t-butylglycine	Nmtbug
10	D-threonine	Dthr	L-norleucine	Nle
	D-tryptophan	Dtrp	L-norvaline	Nva
	D-tyrosine	Dtyr	α-methyl-aminoisobutyrate	Maib
	D-valine	Dval	α-methyl-α-aminobutyrate	Mgabu
	D-α-methylalanine	Dmala	α-methylcyclohexylalanine	Mchexa
15	D-α-methylarginine	Dmarg	α-methylcylcopentylalanine	Mcpen
	D-α -methylasparagine	Dmasn	α-methyl-α-napthylalanine	Manap
	D-α-methylaspartate	Dmasp	α-methylpenicillamine	Mpen
	D-α -methylcysteine	Dmcys	N-(4-aminobutyl)glycine	Nglu
	D-α-methylglutamine	Dmgln	N-(2-aminoethyl)glycine	Naeg
20	D-α-methylhistidine	Dmhis	N-(3-aminopropyl)glycine	Norn
	D-α -methylisoleucine	Dmile	N-amino-α-methylbutyrate	Nmaabu
	D-α-methylleucine	Dmleu	α-napthylalanine	Anap
	D-α-methyllysine	Dmlys	N-benzylglycine	Nphe
	D-α-methylmethionine	Dmmet	N-(2-carbamylethyl)glycine	Ngln
25	D-α-methylornithine	Dmorn	N-(carbamylmethyl)glycine	Nasn
	D-α-methylphenylalanine	Dmphe	N-(2-carboxyethyl)glycine	Nglu
	D-α-methylproline	Dmpro	N-(carboxymethyl)glycine	Nasp

	D-α-methylserine	Dmser	N-cyclobutylglycine	Ncbut
	D-α-methylthreonine	Dmthr	N-cycloheptylglycine	Nchep
	D-α-methyltryptophan	Dmtrp	N-cyclohexylglycine	Nchex
	D-α-methyltyrosine	Dmty	N-cyclodecylglycine	Ncdec
5	D-α-methylvaline	Dmval	N-cylcododecylglycine	Ncdod
	D-N-methylalanine	Dnmala	N-cyclooctylglycine	Ncoct
	D-N-methylarginine	Dnmarg	N-cyclopropylglycine	Ncpro
	D-N-methylasparagine	Dnmasn	N-cycloundecylglycine	Ncund
	D-N-methylaspartate	Dnmasp	N-(2,2-diphenylethyl)glycine	Nbhm
10	D-N-methylcysteine	Dnmcys	N-(3,3-diphenylpropyl)glycine	Nbhe
	D-N-methylglutamine	Dnmgln	N-(3-guanidinopropyl)glycine	Narg
	D-N-methylglutamate	Dnmglu	N-(1-hydroxyethyl)glycine	Nthr
	D-N-methylhistidine	Dnmhis	N-(hydroxyethyl))glycine	Nser
	D-N-methylisoleucine	Dnmile	N-(imidazolylethyl))glycine	Nhis
15	D-N-methylleucine	Dnmleu	N-(3-indolylyethyl)glycine	Nhtrp
	D-N-methyllysine	Dnmlys	N-methyl-α-aminobutyrate	Nmgabu
	N-methylcyclohexylalanine	Nmchexa	D-N-methylmethionine	Dnmmet
	D-N-methylornithine	Dnmorn	N-methylcyclopentylalanine	Nmcpen
	N-methylglycine	Nala	D-N-methylphenylalanine	Dnmphe
20	N-methylaminoisobutyrate	Nmaib	D-N-methylproline	Dnmpro
	N-(1-methylpropyl)glycine	Nile	D-N-methylserine	Dnmser
	N-(2-methylpropyl)glycine	Nleu	D-N-methylthreonine	Dnmthr
	D-N-methyltryptophan	Dnmtrp	N-(1-methylethyl)glycine	Nval
	D-N-methyltyrosine	Dnmtyr	N-methyla-napthylalanine	Nmanap
25	D-N-methylvaline	Dnmval	N-methylpenicillamine	Nmpen
	α-aminobutyric acid	Gabu	N-(p-hydroxyphenyl)glycine	Nhtyr
	L-t-butylglycine	Tbug	N-(thiomethyl)glycine	Ncys

	L-ethylglycine	Etg	penicillamine	Pen
	L-homophenylalanine	Hphe	L-α-methylalanine	Mala
	L- -methylarginine	Marg	L-α-methylasparagine	Masn
	L- -methylaspartate	Masp	L-α-methyl-t-butylglycine	Mtbug
5	L- -methylcysteine	Mcys	L-methylethylglycine	Metg
	L- -methylglutamine	Mgln	L-α-methylglutamate	Mglu
	L- -methylhistidine	Mhis	L-α-methylhomophenylalanine	Mhphe
	L- -methylisoleucine	Mile	N-(2-methylthioethyl)glycine	Nmet
	L- -methylleucine	Mleu	L-α-methyllysine	Mlys
10	L- -methylmethionine	Mmet	L-α-methylnorleucine	Mnle
	L- -methylnorvaline	Mnva	L-α-methylornithine	Morn
	L- -methylphenylalanine	Mphe	L-α-methylproline	Mpro
	L- -methylserine	Mser	L-α-methylthreonine	Mthr
	L- -methyltryptophan	Mtrp	L-α-methyltyrosine	Mtyr
15	L- -methylvaline	Mval	L-N-methylhomophenylalanine	Nmhphe
	N-(N-(2,2-diphenylethyl) carbamylmethyl)glycine	Nnbhm	N-(N-(3,3-diphenylpropyl) carbamylmethyl)glycine	Nnbhe
	1-carboxy-1-(2,2-diphenyl-ethylamino)cyclopropane	Nmbc		

20

Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be limiting to the present invention.

EXAMPLES

Standard molecular biology techniques known in the art and not specifically described were generally followed as described in Sambrook and Russel, Molecular Cloning: A Laboratory Manual, Cold Springs Harbor Laboratory, New York (2001).

Example 1 - Source of Materials

Des-aspartate-angiotensin was purchased from Bachem (Dubendorf, Switzerland). Des-aspartate-angiotensin I can be prepared by techniques well known in the art. Influenza A/Aichi/2/68 virus and Madin-Darby Canine Kidney cells were purchased from ATCC. The virus has a titer of 106.75 CEID50/0.2 ml in 2 days on 11 day old SPF CE. Six to 7 weeks old female BALB/c mice were obtained from the Animal Center, National University of Singapore. Six to 7 weeks old diabetic KKAy mice were purchased from CLEA, Japan.

Example 2 - Development of a Mouse Model for Influenza A Virus

Two BALB/c mice were intranasally infected with 50 µL of influenza A virus. Two days later, the infected mice were sacrificed by cervical dislocation and their lungs were removed. 0.3 g of lung tissue was homogenized in 1mL of PBS containing 1000 units/L penicillin and 10 µg/L streptomycin. The lung homogenate was cleared of connective tissue by centrifuging at 5,000 rpm for 5 minutes on a bench-top centrifuge. This homogenate was labeled as passage-1 lung homogenate. A second batch of 2 BALB/c mice was infected with 50 µL of passage-1 lung homogenate. The process of passaging was repeated and the virus virulence and titer in each passage of lung homogenate was monitored by assaying its infectivity in Madin-Darby Canine Kidney (MDCK) cells, which is a host cell for influenza A virus. The virulence and titer of the influenza A virus

increased progressively with each passage. Fifty µL of passage-6 lung homogenate, equivalent to 2.5 x 105 TCID50 of the virus in MDCK cells, were then used to infect 6-7 weeks old female BALB/c mice for the study of the antiviral action of des-aspartate-angiotensin I.

Example 3 - Monitoring and Treatment (with Orally-Administerd Des-aspartate-angiotensin I) of Influenza A Virus-Infected BALB/c Mice

BALB/c mice when infected with passage-6 lung homogenate lost weight and suffered fatality. The following protocol described by Johasson et al was used to study the effect of des-aspartate-angiotensin on these parameters. Six to 7 weeks old female BALB/c mice were intranasally infected with 50 µL passage-6 lung homogenate. Groups of 5-9 mice were orally administered with one of the following dose of des-aspartate-angiotensin I:- 300 nmole/kg daily for 8 days, 600 nmole/kg daily for 8 days, and 1200 nmole/kg on day 2 post virus infection (i.e. only one dose was administered). A group of 8 mice were similarly administered water for 8 days and served as the control group. The weight and survival of the mice were recorded daily for 16 days.

Example 4 - Effect of Orally-Administered Des-aspartate-angiotensin I on Virus-Infected BALB/c Mice

The results of the study are summarized in Table 2 and depicted in Fig. 1. The daily weight of each surviving animal was expressed as a percentage of its weight at Day 0 (i.e. day of virus inoculation). The data of the control and each treatment group were expressed as the mean ± SEM. Significant differences were determined by one way ANOVA and post hoc Tukey test. The accepted level of significance was $p < 0.05$. Des-aspartate-angiotensin I is an antiviral agent in term of its ability to attenuate the weight loss and survival of the virus-

infected animals. Effective antiviral activity was observed with a dose of 600 nmole/kg administered daily for 8 days, and a single dose of 1200 nmole/kg given on Day 2.

Example 5 - Monitoring and Treatment (with Intraperitoneally-administered Des-aspartate-angiotensin I) of Influenza A Virus-infected BALB/c Mice

Thirty BALB/c mice were infected with 55 µL passage-6 lung homogenate and the weight of each animal was monitored daily as described in EXAMPLES 3 and 4. Animals were intraperitoneally administered with either 60 nmole/kg des-aspartate-angiotensin I in 0.1 ml of phosphate buffer saline (PBS) or 0.1 ml PBS on the day when their weight loss was 15% or greater during the first post-infection week. This treatment was continued for 7 days. Of the 30 animals, 14 animals lost more than 15% of their weight during the first post-infection week. Hence, 7 animals were randomly chosen for des-aspartate-angiotensin I treatment and 7 animals for vehicle treatment. The weight of the treated animals were monitored till either the animal died or regained 100% of their initial weight. Surviving animals were euthanized and their lung homogenate was prepared and assayed for influenza A virus as described in Example 2.

Post Infection Period (in Days)	Average Percentage Weight Change (weight of individual animals at Day 0 was taken as 100%)			
	Control (n =8)	300 nmole/kg/day for 16 days (n = 5)	600 nmole/kg/day for 16 days (n = 9)	1200 nmole/kg given as a single dose on Day 2 (n = 9)
0	100	100	100	100
1	102.4 ± 2.2	102.1 ± 1.5	98.7 ± 1.3	103.5 ± 0.5
2	93.4 ± 1.4	96.2 ± 2.0	95.8 ± 1.4	96.8 ± 0.6
3	88.7 ± 1.0	90.8 ± 2.4	92.0 ± 1.7	**95.1 ± 0.8***
4	86.4 ± 1.3	85.8 ± 2.9	**92.4 ± 1.8***	**96.7 ± 1.2***
5	84.1 ± 2.3	83.2 ± 3.4	**94.8 ± 1.8***	**96.3 ± 2.1***
6	83.3 ± 3.6	94.1# **(3)**	**95.5 ± 2.1***	**95.7 ± 3.1***
7	84.1 ± 4.6	95.7#	**97.8 ± 2.6***	**97.8 ± 4.0***
8	91.0 ± 5.4 **(2)**	97.9#	96.7 ± 2.9	98.3 ± 4.3
9	91.8 ± 6.2	98.7#	96.3 ± 3.5	101.1 ± 4.7
10	93.4 ± 7.1	98.1#	95.6 ± 4.0	101.1 ± 4.8
11	101.6 ± 2.8 **(1)**	97.4#	95.4 ± 4.5	105.9 ± 1.3 **(1)**
12	102.4 ± 3.1	96.9#	95.1 ± 5.0	106.1 ± 1.3
13	99.9 ± 5.8	96.8#	95.5 ± 5.2	108.1 ± 1.2
14	103.0 ± 3.9	98.2#	99.1 ± 3.4	107.2 ± 1.3
15	105.4 ± 3.8	99.6#	100.3 ± 4.2	109.4 ± 1.2
16	104.4 ± 3.5	102.4#	97.0 ± 4.8	108.4 ± 1.5

Table 2 *Significantly different from the corresponding control value ($p < 0.05$, ANOVA post hoc Tukey test) Bold numbers within brackets indicate number of animals that died on the day of recording. #Average value from two surviving mice i.e. no SEM.

Example 6 - Effect of Intraperitoneally-administered Des-aspartate-angiotensin I on Virus-infected BALB/c Mice

The data of the study are summarized in Table 3. The weight of each animal was expressed as a percentage of its weight at Day 0. All mice that were treated with saline died within a week. In the des-aspartate-angiotensin I treated

mice, only 1 mouse died. Six animals survived and regained their original weight. No influenza A virus was detected in the lung homogenate of the surviving mice. The results show that des-aspartate-angiotensin I exerts effective antiviral action in preventing influenza A virus-infected mice from dying.

Example 7 - Effect of Intraperitoneally-Administered Des-aspartate-angiotensin I on Blood Glucose Profile in Diabetic KKAy Mice

The diabteic KKAy mice were divided into four groups consisting of 6 animals per group. Animals in the control group were intraperitoneally administered 0.1 mL saline. Animals in the second, third, and fourth groups were similarly administered with 100, 200, 400 nmole/kg des-aspartate-angiotensin I, respectively in 0.1 ml of PBS. Treatment with saline and des-aspartate-angiotensin I was carried out daily for 4 weeks. Following this, animals were fasted overnight for 16 hours and oral glucose tolerance test was performed as follows: blood was withdrawn from the orbital sinus for blood glucose determination (time of withdrawal was designated as 0 time), animals were then orally administered glucose (2g/kg), and blood was withdrawn at 30, 60 and 120 min for blood glucose determination. The blood was allowed to clot and blood glucose was measured as serum glucose using a commercial glucose kit from Thermo Electron Corporation, Australia. Table 3 shows that des-aspartate-angiotensin I, administered at doses of 200 and 400 nmole/kg/day for 4 weeks, significantly lowered the blood glucose level at 30 and 60 min of the oral glucose tolerance test. The data, graphically displayed in Fig. 2, show that des-aspartate-angiotensin I possesses hypoglycaemic action and can be use to treat the symptoms of hyperglycaemia.

Example 8 - Dose-Range Finder Experiment for Orally Administered Des-aspartate-angiotensin I

A dose-range finder experiment to determine the hypoglycaemic dose for orally administered des-aspartate-angiotensin I was also carried out. In this experiment, the animals in each of the three treatment groups of 7 diabetic KKAy mice were orally administered (by gavage) 200, 400, and 600 nmole/kg/day des-aspartate-angiotensin I in volume of 0.1 mL, respectively. Animals in the control group were similarly administered 0.1 mL water. The treatment period was 4 weeks. Oral glucose tolerance test was carried out as described in Example 5, except that blood was sampled at 0 and 30 min. The data, presented in Table 4, show that des-aspartate-angiotensin I is an effective oral hyperglycemic agent at an oral dose of 600 nmole/kg.

Example 9 - Effect of Orally-Administered Des-aspartate-angiotensin I on Blood Glucose Profile in Diabetic KKAy Mice

In this experiment, the animals in each of the three treatment groups of 7 diabetic KKAy mice were orally administered (by gavage) 200, 400, and 600 nmole/kg/day des-aspartate-angiotensin I in volume of 0.1 mL water, respectively. Animals in the control group were similarly administered 0.1 mL water. The treatment period was 4 weeks. Oral glucose tolerance test was carried out as described in EXAMPLE 7, except that blood was sampled at 0 and 30 min. The data, presented in Table 5, show that des-aspartate-angiotensin I exerts significant hypoglycaemic action at an oral dose of 600 nmole/kg.

Example 10 - Effect of Orally-administered Des-aspartate-angiotensin I on Blood Glucose Profile in Diabetic GK Rats

Diabetic GK rats were divided into three groups, consisting of six animals in each group. Animals in groups 1 and 2 were administered, by gavage, one of the following: 400 or 600 nmole/kg of des-aspartate-angiotensin I in a volume of 0.2 mL water daily for up to 8 weeks. Control animals in group 3 were similarly administered 0.2 mL water. Thirty min oral glucose tolerance test was performed fortnightly as a spot test to detect significant hypoglycaemic action (if any) that arose from des-aspartate-angiotensin I treatment. As shown in Fig. 3, significant hypoglycaemic action was detected at 6 weeks of treatment. Animals were treated for a further 2 weeks and a full oral glucose tolerance test was performed thereafter. The data, presented in Fig. 4, show that the hypoglycaemic action of des-aspartate-angiotensin I was also significantly expressed at 8 weeks of treatment. The results of this study show that des-aspartate-angiotensin I is an effective oral hypoglycaemic agent and can be used to treat the hyperglycemia in diabetes.

Example 11 - Effect of Orally-Administered Des-aspartate-angiotensin I on Insulin-Induced Translocation of Glucose Transporter-4 (GLU4) to Plasma Membrane in Skeletal Muscles of Diabetic GK Rats

The diabetic GK rats described in EXAMPLE 9 were similarly treated for another week. At 9 weeks of treatment, animals were fasted overnight and half the animals in each group were intraperitoneally administered with PBS and the other half with insulin (0.5 U/ kg). After 30 minutes, the animals were sacrificed by cervical dislocation. Hind limb skeletal muscles were rapidly excised and frozen in liquid nitrogen. Muscle samples were stored at -80 oC until used. The hindlimb muscles were subsequently thawed over an ice pack and plasma membranes were prepared as described by Dombrowaski et al .

Thirty µg of protein from the 25 % interface of the discontinued sucrose gradient was subjected to SDS-PAGE, transferred to PVDF sheets and incubated with anti-GLUT4 polyclonal primary antibody (1: 800), washed and followed by incubation with anti-goat secondary antibody (1: 10,000). The bands were detected at 49 kDa. The data, presented in Fig. 5, show that des-aspartate-angiotensin I significantly enhances the insulin-induced translocation of GLUT4 to the plasma membrane in skeletal muscle of diabetic GK rats. As insulin resistance is the hallmark of diabetes, it is concluded that the des-aspartate-angiotensin I exerts marked hypoglycaemic action in diabetic GK rats by overcoming insulin resistance in these animals. This is especially so as serum level of insulin was not significantly affected in the same diabetic GK rats that were treated with 600 nmole/kg of des-aspartate-angiotensin I (see Fig. 6 for details).

Example 12 – Kit

The present invention provides a kit comprising des-aspartate-angiotensin I, its derivative, a functional part and/or an analogue thereof for the treatment and/or prophylaxis of at least one viral infection and/or for the induction of hypoglycaemia and/or reduction of hyperglycaemia. There is also provided a kit for the treatment and/or prophylaxis of a hyperglycaemia-related condition excluding renal-related disorders. The kit may further comprise information and/or instruction pertaining to its use.

Table 3: Percentage Weight Change of Individual Mouse

Post Infection Period (in Days)	Saline treated mice							Des-aspartate-angiotensin I treated mice						
	1	2	3	4	5	6	7	1	2	3	4	5	6	7
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.4	98.1	98.7	99.2	99.5	99.3	94.8	99.4	100.4	99.0	98.4	99.3	98.5	98.9
2	96.5	97.1	97.5	97.9	98.2	96.6	87.8	99.3	99.8	96.3	97.1	96.6	97.3	95.8
3	89.8	96.8	96.4	96.8	95.8	93.9	86.0	99.4	99.0	90.3	94.6	88.6	94.7	93.9
4	88.2	93.2	93.1	86.0	85.7	87.5	**81.5**	97.5	96.6	85.7	86.2	**83.3**	92.1	90.8
5	87.7	86.5	90.9	**80.7**	**82.0**	**84.7**	**80.1**	90.3	91.7	**82.8**	85.7	**81.0**	90.1	89.0
6	81.8	**82.9**	86.7	**76.8**	**78.7**	**83.4**	**78.6**	87.5	87.4	**80.8**	**82.7**	**78.2**	**85.2**	88.8
7	**80.9**	**81.1**	**84.1**	**69.8**	**77.4**	**78.4**	**73.9**	**85.0**	**83.8**	**80.7**	**76.8**	**73.9**	**82.8**	**84.4**
8	**74.6**	**79.4**	**81.5**	**65.4**	**74.9**	**76.3**	**69.4**	**86.2**	**80.4**	**79.7**	**72.9**	**71.2**	**83.2**	**82.4**
9	**71.8**	**75.3**	**80.1**	died	**72.7**	**74.0**	**65.1**	**90.3**	**78.8**	**81.4**	**69.9**	**71.8**	**80.5**	**78.6**
10	**70.8**	**70.6**	**76.4**		**69.4**	**71.0**	**61.0**	**93.0**	**77.5**	**80.5**	**69.0**	**73.2**	**82.3**	**77.7**
11	**69.6**	died	**72.6**		**68.2**	**68.9**	died	**99.3**	**77.1**	**82.1**	**67.7**	75.8	**88.9**	**74.5**
12	**68.8**		died		died	died		**101.2**	**78.9**	80.0	**64.9**	80.1	**88.9**	**74.5**
13	died								**80.8**	76.6	63.0	82.8	97.4	**72.3**
14									82.6	75.2	61.3	86.3	102.8	74.0
15									84.3	83.0	58.9	90.1		76.7
16									89.4	90.3	58.9	94.6		78.6
17									94.8	93.5	58.7	99.3		81.2
18									98.7	99.1	died	101.0		83.2
19									100.9	100.8				86.8
20														91.8
21														95.7
22														98.4
23														100.1

Table 4

Dose of des-aspartate-angiotensin I	Serum glucose level at different sampling time (nmole/dL)			
	0 min	30 min	60 min	120 min
Control	8.3 ± 0.6	27.3 ± 1.9	22.2 ± 1.3	12.2 ± 1.3
100 nmole/kg	7.9 ± 0.9	**21.2 ± 1.6***	**13.4 ± 1.3***	9.9 ± 0.5
200 nmole/kg	6.9 ± 0.9	**16.0 ± 1.4***	**14.5 ± 0.9***	8.7 ± 1.2
400 nmole/kg	6.5 ± 0.6	**15.4 ± 1.6***	**13.0 ± 1.1***	8.8 ± 0.6

*Significantly different from the corresponding value of the Control ($p < 0.05$, ANOVA post hoc Tukey test)

Table 5

Dose of des-aspartate-angiotensin I	Serum glucose level at different sampling time (nmole/dL)	
	0 min	30 min
Control	9.0 ± 1.0	33.7 ± 1.7
200 nmole/kg	8.9 ± 1.0	37.7 ± 2.6
400 nmole/kg	9.1 ± 1.6	26.0 ± 3.6
600 nmole/kg	8.9 ± 0.7	**17.5 ± 1.9***

*Significantly different from the corresponding value of the Control ($p < 0.05$, ANOVA post hoc Tukey test)

References

DiRamond J. The double puzzle of diabetes. Nature 2003; 423: 599-602.

Dombrowski *et al* A new procedure for the isolation of plasma membranes, T tubules, and internal membranes from skeletal muscle. Am J Physiol 1996; 270:E667-E676.

Fendrick *et al*. The economic burden of non-influenza-related viral respiratory tract infection in the United States. Arch Intern Med 2003; 163:487-494.

Levitt NS *et al*. The prevalence and identification of risk factors for NIDDM in urban Africans in Cape Town, South Africa. Diabetes Care 1993; 16:601-607.

Johasson *et al*. Infection-permissive immunization with influenza virus neuraminidase prevents weight loss in infected mice. Vaccine 1993; 11:1037-1039.

Sim MK *et al*. Degradation of angiotensin I in the endothelium and smooth muscle of the rat aorta. Biochem Pharmacol 1994(a); 45:1524-1527.

Sim MK *et al*. Degradation of angiotensin I to [des-Asp1]angiotensin I by a novel aminopeptidase in the rat hypothalamus. Biochem Pharmacol 1994(b); 48:1043-1046.

Sim MK, Qui XS. Angiotensins in plasma of hypertensive rats and human. Regul Peptides 2003; 111:179-182.

Sullivan KM, *et al*. Estimates of the US health impact of influenza. Am J Public Health 1993; 83:1712-1716.

US 5,773,416

US 6,100,237

US 6,589,938 B2

US 2003/0086920

Claims

1. A method for the treatment and/or prophylaxis of at least one viral infection comprising administering to a subject des-aspartate-angiotensin I, its derivative, a functional part and/or an analogue thereof.

2. The method according to claim 1, wherein the at least one viral infection is a respiratory viral infection.

3. The method according to any one of claims 1 or 2, wherein the at least one viral infection is an influenza virus infection.

4. The method according to any one of claims 1 to 3, wherein the at least one viral infection is influenza A infection.

5. A method of inducing hypoglycaemia and/or reducing hyperglycaemia in a subject, the method comprising administering to the subject des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

6. A method of treatment and/or prophylaxis of at least one hyperglycaemia-related condition comprising administering to a subject des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, provided the at least one hyperglycaemia-related condition is not a renal-related disorder.

7. The method according to claim 6, wherein at least one hyperglycaemia-related condition is type I diabetes, obesity, and/or bulimia nervosa.

8. The method according to any one of claims 1 to 7, wherein the des-aspartate-angiotensin I, its derivative, a functional part and/or analogue

thereof, is administered in a pharmaceutically and/or therapeutically effective amount.

9. The method according to any one of claims 1 to 8, wherein the des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, is administered in combination with at least one pharmaceutically-acceptable carrier, excipient, diluent and/or adjuvant.

10. Des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, for use in the treatment and/or prophylaxis of at least one viral infection.

11. The use according to claim 10, wherein the at least one viral infection is a respiratory viral infection.

12. The use according to any one of claims 10 or 11, wherein the at least one viral infection is an influenza virus infection.

13. The use according to any one of claims 10 to 12, wherein the at least one viral infection is influenza A infection.

14. Des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, for use in the induction of hypoglycaemia and/or reduction of hyperglycaemia.

15. Des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, for use in the treatment and/or prophylaxis of at least one hyperglycaemia-related condition, provided the at least one hyperglycaemia-related condition is not a renal-related disorder.

16. The use according to claim 15, wherein at least one hyperglycaemia-related condition is type I diabetes, obesity, and/or bulimia nervosa.

17. The use according to any one of claims 10 to 16, wherein the des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, is administered in a pharmaceutically and/or therapeutically effective amount.

18. Use of des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof for the preparation of a medicament for the treatment and/or prophylaxis of at least one viral infection.

19. The use according to claim 18, wherein the at least one viral infection is a respiratory viral infection.

20. The use according to any one of claims 18 or 19, wherein the at least one viral infection is an influenza virus infection.

21. The method according to any one of claims 18 to 20, wherein the at least one viral infection is influenza A infection.

22. Use of des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, for the preparation of a medicament for the induction and/or reduction of hyperglycaemia.

23. Use of des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, for the preparation of a medicament for the treatment and/or prophylaxis of the at least one hyperglycaemia-related condition, provided the at least one hyperglycaemia-related condition is not a renal-related disorder.

24. The use according to claim 23, wherein the at least one hyperglycemia-related condition is type I diabetes, obesity, and/or bulimia nervosa.

25. The use according to any one of claims 18 to 24, wherein the des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof, is administered in a pharmaceutically and/or therapeutically effective amount.

26. The use according to any one of claims 18 to 25, wherein the medicament, further comprises at least one pharmaceutically acceptable carrier, excipient, diluent and/or adjuvant.

27. A kit for the treatment and/or prophylaxis of at least one viral infection comprising des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

28. The kit according to claim 27, wherein the at least one viral infection is a respiratory viral infection.

29. The kit according to any one of claims 27 or 28, wherein the at least one viral infection is an influenza virus infection.

30. The kit according to any one of claims 27 to 29, wherein the at least one viral infection is influenza A infection.

31. A kit for the induction of hypoglycaemia and/or reduction of hyperglycaemia comprising des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

32. A kit for the treatment and/or prophylaxis of at least one hyperglycaemia-related condition provided the at least one hyperglycaemia-related condition

is not a renal-related disorder, the kit comprising des-aspartate-angiotensin I, its derivative, a functional part and/or analogue thereof.

33. The kit according to claim 32, wherein at least one, hyperglycemia-related condition is type I diabetes, obesity, and/or bulimia nervosa.