CLAIMS

1. Use of *Enterococcus faecalis* that is killed in the manufacture of a medicament for reducing incidence of and/or treating *Clostridium difficile* infection.

2. The use of claim 1, wherein the *Enterococcus faecalis* is EF-2001 strain.

3. The use of any one of claims 1 to 2, wherein the *Enterococcus faecalis* that is killed is to be orally administered.

4. The use of claim 3, wherein the *Enterococcus faecalis* that is killed is to be orally administered in an amount of 1×10^8 - 1×10^{11} CFU/kg body weight per dose.

5. The use of claim 4, *Enterococcus faecalis* that is killed is to be orally administered one or more times per day in an amount of 1×10^8 - 1×10^{11} CFU/kg body weight per dose.

6. The use of claim 5, wherein the *Enterococcus faecalis* that is killed is to be orally administered one to five times per day in an amount of 1×10^8 - 1×10^{11} CFU/kg body weight per dose.

7. The use of claim 3, wherein the bacterium belonging to the *Enterococcus faecalis* that is killed is to be orally administered in an amount of 1×10^9 - 5×10^{10} CFU/kg body weight per dose.

8. The use of claim 7, wherein the *Enterococcus faecalis* that is killed is to be orally administered one or more times per day in an amount of 1×10^9 - 5×10^{10} CFU/kg body weight per dose.

9. The use of claim 8, wherein the *Enterococcus faecalis* that is killed is to be orally administered one to five times per day in an amount of 1×10^9 - 5×10^{10} CFU/kg body weight per dose.

10. The use of claim 3, wherein the *Enterococcus faecalis* that is killed is to be orally administered in an amount of 1.2 x 10^{10} or more CFU/kg body weight per dose.

11. The use of claim 10, wherein the *Enterococcus faecalis* that is killed is to be orally administered one or more times per day in an amount of 1.2 x 10^{10} or more CFU/kg body weight per dose.

12. The use of claim 11, wherein the *Enterococcus faecalis* that is killed is to be orally administered one to five times per day in an amount of 1.2 x 10^{10} or more CFU/kg body weight per dose.

13. The use of any one of claims 3 to 12, wherein the *Enterococcus faecalis* that is killed is to be orally administered for seven or more days.

14. The use of any one of claims 1 to 13, which is to be used for prevention of *Clostridium difficile* infection.

15. The use of any one of claims 1 to 13, which is to be used for treatment of *Clostridium difficile* infection.