

Specification

Drugs for the Diagnosis of Tissue Reproductive Activity or
the Treatment of Proliferative Diseases

Technical Field

5 The present invention relates to use of radiolabeled nucleoside derivatives for diagnosis of tissue proliferation activity and treatment of proliferative diseases.

Background Art

 If proliferation activity of tumor cells can be determined non-invasively by image diagnosis, it will be help for evaluation of growth rate and malignancy of the
10 tumor. Detection of the most rapidly growing regions of a tumor by image diagnosis will be useful in preparing plans for radiation fields in radiotherapy and identifying suitable portions for biopsy. Such methods will permit an early and accurate evaluation of therapeutic effects, which is difficult to identify by CT- or MRI-based anatomical evaluation or PET-based measurement of glucose-metabolic changes.
15 Particularly, they will be useful for an early assessment of therapeutic effects of anticancer agents that may cause strong side effects.

 In order to solve these clinically important problems, use of 5-iodo-deoxyuridine labeled with a radioactive iodine and thymidine labeled with carbon-11 which is a positron-emitter, have been studied (Tjuvajev JG et al., J. Nucl. Med. 35, pp.1407-1417 (1994); Blasberg RG et al, Cancer Res. 60, pp.624-635 (2000); Martiat
20 Ph et al., J. Nucl. Med. 29, pp.1633-1637 (1998); Eary JF et al., Cancer Res. 59, pp. 615-621 (1999); U.S. Patent No. 5,094,835; U.S. Patent No. 5,308,605). It is considered that these radiolabeled compounds are taken into cells as precursors for DNA synthesis required for cell division of rapidly-growing tumors, and then
25 phosphorylated by thymidine kinase, followed by incorporation into DNA, to reflect

proliferation activity of the tumor. These radiolabeled compounds, however, are decomposed rapidly *in vivo*, making it difficult to perform non-invasive evaluation of the proliferation activity of the tumor. The method using carbon-11-labeled thymidine, in particular, requires very complicated mathematical model analysis, and cannot
5 become popular as a diagnostic technique of nuclear medicine imaging.

The rapid metabolic decomposition of these radiolabeled compounds *in vivo* is considered to be due to cleavage of C-N glycosidic bonds by thymidine phosphorylase and instability of the labels *in vivo*. If the C-N glycosidic bonds are cleaved, the compound loses its affinity to tumors, thereby decreasing in accumulation of
10 radioactivity in tumors, while the radioactive metabolites increase background radioactivity, thereby making imaging of the tumors difficult.

To solve these problems, radiolabeled compounds with metabolic stability have been synthesized by introducing fluorine atoms, which are high in electronegativity, to the 2' or 3' position in certain nucleosides, and have been studied for imaging of tumors.
15 Thus, 3'-deoxy-3'-fluorothymidine that contains fluorine 18, a positron emitter, at the 3' position shows a high stability *in vivo* and an accumulation in tumor tissue (Shields AF et al., Nature Med. 4, pp.1334-1336 (1998)). Though this radiolabeled compound is stable *in vivo*, it is a radio-labeled compound with a short-life positron emitter, and therefore a cyclotron is required in the hospital, limiting the usage of the compound.
20 For this radiolabeled compound, the major process responsible for its accumulation in cells is the phosphorylation caused by thymidine kinase that is an index of DNA synthesis, and thus it does not serve as an agent that essentially reflects DNA synthesis.

A derivative of 5-iododeoxyuridine, in which fluorine is introduced to the 3' position in the same manner as above to increase its stability *in vivo*, has recently been
25 reported. Though stable *in vivo*, however, this radiolabeled compound was high in

retention in blood and failed to show a significant accumulation in a tumor compared to 5-iododeoxyuridine (Choi SR et al., J. Nucl. Med. 41, p. 233 (2000)).

2'-fluoro-5-iodoarabinouridine, in which fluorine is introduced to the 2' position, shows a high stability *in vivo*, and has been used for identification of introduction and expression *in vivo* of a vector for gene therapy, utilizing a phosphorylation reaction specific to thymidine kinase of human herpesvirus. It has also been applied to image diagnosis for virus infection, based on the high specificity to the viral thymidine kinase (Tjuvajev JG et al., Cancer Res. 56, pp.4087-95 (1996); Tjuvajev JG et al., Cancer Res. 58, pp.4333-4441 (1998); Wiebe LI et al., Nucleosides
10 Nucleotides 18, 1065-1076 (1999); Gambhir SS et al., Nucl. Med. Biol. 26, pp.481-490 (1999); Haubner R et al., Eur. J. Nucl. Med. 27, pp.283-291 (2000); Tjuvajev JG et al. Cancer Res. 59, 5186-193 (1999); Bengel FM et al., Circulation 102, pp.948-950 (2000)).


In view of the above situation, the present invention aims to provide a radiolabeled compounds that are practically useful in clinical fields, stable *in vivo*, and able to retain in cells after being phosphorylated by thymidine kinase of mammals, or reflect the DNA synthesis activity after being incorporated in DNA, particularly those compounds which are labeled with a single-photon emitter to achieve a wide spectrum of use, and also aims to provide methods for diagnosis of tissue proliferation activity and for treatment of proliferative disease, utilizing agents that contain said radiolabeled
20 compounds.

Disclosure of the Invention

To achieve the above-mentioned objectives, the present inventors have synthesized a variety of radiolabeled compounds and have intensively studied to see if
25 they are useful for image evaluation of tissue proliferation activity. As a result, the

inventors have found that radiolabeled compounds as represented by the following formula can serve for diagnosis of tissue proliferation activity or treatment of proliferative disease, and have completed the present invention.

Specifically, the present invention provides an agent for diagnosis of tissue proliferation activity or for treatment of proliferative disease, which comprises, as an active ingredient, a radiolabeled compound as represented by the following formula or a pharmaceutically acceptable salt thereof:



wherein R₁ denotes hydrogen, or a linear- or branched-chain alkyl group having 1-8 carbon atoms; R₂ denotes hydrogen, hydroxyl, or a halogen substituent; R₃ denotes hydrogen or fluorine substituent, R₄ denotes oxygen, sulfur or a methylene substituent, and R₅ denotes a radioactive halogen substituent, excluding the case where R₁, R₂ and R₃ are hydrogen, R₄ being oxygen, and R₅ being radioactive fluorine, bromine, iodine, or astatine; the case where R₁ and R₃ are hydrogen, R₂ being fluorine, R₄ being oxygen, and R₅ being radioactive bromine or iodine; and the case where R₁ and R₂ are hydrogen, R₃ being fluorine, R₄ being oxygen, and R₅ being radioactive bromine or iodine.

The radiolabeled compounds of the present invention are stable *in vivo*, and can retain in cells after being phosphorylated by mammalian thymidine kinase or reflect the

DNA synthesis activity after being incorporated in DNA. Therefore, they realize effective diagnosis of tissue proliferation activity and treatment of proliferative disease, and are particularly useful as diagnostic radioactive imaging agents for diagnosis of tissue proliferation activity or as radioactive therapeutic agents for treatment of proliferative disease in accordance with internal radiotherapy, local radiotherapy or the like.

Thus, according to another aspect of the present invention, there are provided methods for diagnosis of tissue proliferation activity, which comprise administering an effective amount of a radiolabeled compound as represented by the above formula or a pharmaceutically acceptable salt thereof to a mammal, followed by imaging *in vivo* distribution thereof, and methods for treatment of proliferative disease, which comprises administering an effective amount of said radiolabeled compound or salt to a mammal. Herein, mammal includes human beings.

In the present invention, the radiolabeled compounds as represented by the above formula include salts thereof, or may be in a form of a hydrate or solvate of these. Such salts include pharmaceutically acceptable salts, for example, one formed with a mineral acid such as hydrochloric acid and sulfuric acid or with an organic acid such as acetic acid. As such a hydrate or solvate, mention may be made of the present radiolabeled compounds or salts thereof to which water molecules or solvent molecules are attached. Furthermore, the compounds of the present invention include their various isomers such as tautomers.

In the above formula, the linear- or branched-chain alkyl group having 1-8 carbon atoms as represented by R_1 includes, for example, methyl group, ethyl group, propyl group, t-butyl group, and n-hexyl group, of which methyl group is preferable. The halogen-substituent as represented by R_2 preferably includes fluorine, chlorine, and

bromine. R_4 is preferably oxygen or sulfur, of which sulfur is particularly preferable.

The radioactive halogen-substituent as represented by R_5 in the above formula includes F-18, Cl-36, Br-75, Br-76, Br-77, Br-82, I-123, I-124, I-125, I-131, and At-211, of which F-18, Br-76, I-123, and I-124 are preferable for diagnostic purposes while Br-
5 77, I-125, I-131, and At-211 are preferable for therapeutic purposes.

Preferred compounds as represented by the above formula include those wherein R_1 is hydrogen or methyl, R_2 is hydrogen or a halogen-substituent, R_3 is hydrogen, and R_4 is oxygen or sulfur, particularly preferably those wherein R_1 , R_2 and R_3 are hydrogen, R_4 is sulfur, R_5 is a radioactive halogen-substituent selected from F-18,
10 I-123, I-125, and I-131.

Certain 4'-thio nucleic acid derivatives as represented by the above formula (where R_5 is a non-radioactive halogen-substituent) have been reported to be resistant to bacterial thymidine phosphorylase as a result of studies on antiviral agents (Dyson MR et al., J. Med. Chem. 34, pp.2782-2786 (1991); Rahim SG et al., J. Med. Chem. 39,
15 pp.789-795 (1996)). It has also been known that certain 5-iodine- and 5-methyl-4'-sulfur substitution products inhibit phosphorylation of thymidine by human thymidine kinase (Strosselli S et al., Biochem J. 334, pp.15-22 (1998)). The chemical structures of these compounds with sulfur at the 4' position and their use as an antiviral agent are already known (International Publication WO9101326, International Publication
20 WO9104982, Japanese Patent Laid-Open No. HEI 10-087687), but neither the corresponding radiolabeled compounds nor their use as a radioactive diagnostic imaging agent or radioactive therapeutic agent have been known.

The chemical structures of certain compounds with a substituent at the 1' position as represented by the above formula (where R_5 is a non-radioactive substituent)
25 and production methods thereof have already been known (Japanese Patent Laid-Open

No. HEI 07-109289). However, neither the corresponding radiolabeled compounds nor their use as a radioactive diagnostic imaging agent or radioactive therapeutic agent have been known.

The compounds as represented by the above formula can be used for various
5 diagnoses of tissue proliferation activity and treatment for proliferative diseases by virtue of their *in vivo* stability and their capability for retention in cells or capability for being incorporated in DNA.

Such diagnoses of tissue proliferation activity include, for example, diagnosis of hyperplasia, regeneration, transplantation or viral infection accompanied by abnormal
10 proliferation .

The diagnosis of hyperplasia accompanied by abnormal proliferation includes, for example, diagnosis of hyperplastic inflammation, benign tumors, or malignant tumors. The diagnosis of the hyperplastic inflammation includes, for example, diagnoses concerning activity of chronic rheumatoid arthritis and determination of
15 therapeutic effects. The diagnosis of the benign tumors includes, for example, diagnoses concerning localization, activity and determination of therapeutic effects. The diagnosis of the malignant tumors includes, for example, diagnoses concerning localization, progress, malignancy and determination of therapeutic effects, of primary and metastatic malignant tumors. Benign tumors include, for example, prostatic
20 hyperplasia, endometrium hyperplasia (cystic hyperplasia, adenomyosis uteri, hystero myoma), ovarian tumor (cystadenoma), mammary gland (mastopathy, mammary gland fibroadenoma), pituitary adenoma, craniopharyngioma, thyroid adenoma, adrenocortical adenoma and pheochromocytoma. Malignant tumors include, for example, malignant lymphoma (Hodgkin's disease, non-Hodgkin lymphoma),
25 pharyngeal cancer, lung cancer, esophagus cancer, gastric cancer, colon cancer, hepatic

cancer, pancreatic cancer, nephric tumor (nephric cancer, nephroblastoma), bladder tumor, prostatic cancer, testicular tumor, uterine cancer, ovarian cancer, breast cancer, thyroid cancer, neuroblastoma, brain tumor (primary brain tumor, metastatic brain tumor), rhabdomyosarcoma, bone tumor (osteosarcoma, metastatic bone tumor),

5 Kaposi's sarcoma, and malignant melanoma.

The diagnosis of regeneration accompanied by abnormal proliferation is exemplified by diagnosis of function of physiological regeneration of blood and diagnosis of pathological regeneration resulting from pathological loss of blood cells, such as evaluation of physiological hematopoietic functions of bone marrow during

10 treatment with anti-cancer drugs and diagnosis of pathological functions of the bone marrow in patients suffering from hypoplastic anemia.

The diagnosis of transplantation accompanied by abnormal proliferation is exemplified by diagnosis of blood cancer patients undergoing bone marrow transplantation or very high-dose chemotherapy using an anticancer agent, such as

15 diagnosis of take or proliferation of transplanted bone marrow cells in bone marrow transplantation.

The diagnosis of viral infection accompanied by abnormal proliferation includes, for example, diagnosis of virus-infected portions and proliferation thereof in infectious diseases caused by Type I or Type II herpes simplex virus, varicella-zoster

20 herpes virus, cytomegalovirus, Epstein-Barr virus, or human immunodeficiency virus, particularly infectious diseases of central nervous system (e.g., viral-infectious cerebritis, meningitis, etc.) caused by Type I or Type II herpes simplex virus or human immunodeficiency virus.

The treatment for proliferative diseases is exemplified by treatment of

25 malignant tumors or viral infection accompanied by abnormal proliferation. Such

malignant tumors include, for example, malignant lymphoma (Hodgkin's disease, non-Hodgkin lymphoma), pharyngeal cancer, lung cancer, liver cancer, bladder tumor, rectal cancer, prostatic cancer, uterine cancer, ovarian cancer, breast cancer, brain tumor (primary brain tumor, metastatic brain tumor), and malignant melanoma. Such a viral
5 infection includes infectious diseases of central nervous system caused by Type I or Type II herpes simplex virus or human immunodeficiency virus, particularly viral encephalitis or meningitis.


Methods for labeling the compounds represented by the above formula at the "5" position with a radioactive halogen may be known methods, such as methods using
10 isotope exchange reaction, and a method using a 5-chloromercuri compound in which mercury is introduced into the "5" position of the compound or a 5-hydrogen compound in which there is no substitution at the "5" position of the compound. The method using the 5-chloromercuri compound is already known as an iodo-labeling method for producing 5-iodo-2'-deoxyuridine (United States Patent No. 4,851,520; Baranowska-
15 Kortylewicz J et al., Appl. Radiat. Isot. 39, p.335 (1988)). This method is, however, disadvantageous for producing pharmaceuticals labeled with a short half-time radioactive nuclide due to side reactions (formation of "5-chloro" compounds, demercurization reaction), a long reaction time (6 hours), and formation of inorganic mercury compounds. The method using a 5-hydrogen compound is already known as
20 a method for producing 5-iodo-2'-deoxyuridine from 2'-deoxyuridine (Knaus EE et al., Appl. Radiat. Isot. 37, p.901 (1986); Fin RD et al., J. Label. Comds. Radiopharm. 40, p.103 (1997)). This method, however, requires heating at 65-115°C, and therefore, it is not suitable for use with compounds that are easily decomposed under heating conditions and cannot be said to be an ideal labeling method, considering the properties
25 of radioactive halogen atoms which preferably should not involve heating operations

during the labeling reaction. Further, the radiolabeling method using isotope exchange reaction is also unsuitable for producing pharmaceuticals that must be maintained at a certain level of quality, because the method is not able to produce carrier-free labeled compounds and is difficult to control variation of specific activity among different
5 labeling runs.

Another useful method for labeling the compounds represented by the above formula at the "5" position with a radioactive halogen is to allow a compound (5-trialkyltin compound), in which the pyrimidine base is substituted by a trialkylstannyl group at the "5" position as represented by Formula 11 in Fig.1, Formula 21 in Fig.2,
10 Formula 28 in Fig.3, Formula 40 in Fig.4, Formula 50 in Fig.5 or Formula 58 in Fig.6, to react with 0.1N sodium hydroxide solution of a radioactive halogen in an appropriate solvent such as chloroform, so that the trialkylstannyl group at the "5" position is converted into a radioactive halogen-substituent. This labeling method, which uses a 5-trialkyltin compound, is preferable as it does not suffer such problems as with the
15 above three labeling methods. Specifically, this method requires only a relatively short reaction time, and it does not produce "5-chloro" compounds or need heating as the reaction readily proceeds at room temperature. The resulting labeled compounds are free of carriers, and if a lower specific activity is desired, a labeled compound with a fixed specific activity can be readily prepared by adding a carrier. This method is also
20 featured in that purification after the reaction is easy to operate. Specifically, 5-trialkyltin compounds are largely different from the corresponding radioactive halogen-labeled compounds in terms of overall molecular polarity as the electrical properties at the "5" position differ between them. Owing to the difference in the molecular polarity, labeled compounds and unreacted precursors can be separated easily by using a
25 commercial reverse-phase silica gel cartridge after the labeling reaction. This permits


elimination of the need of troublesome high performance liquid chromatographic purification.

Thus, according to another aspect of the present invention, there is provided a method for producing a radiolabeled compound as represented by the following
5 formula:



wherein R_1 denotes hydrogen or a linear- or branched-chain alkyl groups having 1-8 carbon atoms, R_2 denotes hydrogen, hydroxyl or a halogen substituent, R_3 denotes
10 hydrogen or fluorine substituent, R_4 denotes oxygen, sulfur or a methylene substituent, and R_5 denotes a radioactive halogen substituent;

comprising reacting a nucleoside derivative as represented by the following formula:




wherein R_1 denotes hydrogen or a linear- or branched-chain alkyl groups having 1-8 carbon atoms, R_2 denotes hydrogen, hydroxyl or a halogen substituent, R_3 denotes hydrogen or fluorine substituent, R_4 denotes oxygen, sulfur or a methylene substituent, and R_5 denotes a trialkylstannyl group,

with an alkaline solution of a radioactive halogen in a solvent, whereby the trialkylstannyl group of R_5 is converted into the radioactive halogen substituent.

The 5-trialkyltin compounds as represented by Formula 11 in Fig.1, Formula 21 in Fig.2, Formula 28 in Fig.3, Formula 40 in Fig.4, Formula 50 in Fig.5, and Formula 58 in Fig.6 are novel compounds which are useful intermediates for producing the radiolabeled compounds of the present invention.

Thus, according to another aspect of the present invention, there is provided a compound as represented by the following formula:



wherein R_1 denotes hydrogen or a linear- or branched-chain alkyl groups having 1-8 carbon atoms, R_2 denotes hydrogen, hydroxyl or a halogen substituent, R_3 denotes hydrogen or fluorine substituent, R_4 denotes oxygen, sulfur or a methylene substituent, and R_5 denotes a trialkylstannyl group.

In the above formula, the linear- or branched-chain alkyl groups having 1-8 carbon atoms as represented by R_1 include, for example, methyl group, ethyl group, propyl group, t-butyl group, and n-hexyl group, of which methyl group is preferred. The halogen-substituent as represented by R_2 preferably includes fluorine, chlorine and bromine. R_4 is preferably oxygen or sulfur. The trialkylstannyl group as represented by R_5 includes trimethylstannyl group, triethylstannyl group and tributylstannyl group.

Preferred compounds as represented by the above formula include those wherein R_1 is hydrogen or methyl, R_2 is hydrogen or a halogen-substituent, R_3 is hydrogen, and R_4 is oxygen or sulfur.

As seen from Figs.1-6, 5-trialkyltin compounds can generally be synthesized by providing their corresponding halogen-containing compound (as represented by Formula 10 in Fig.1, Formula 20 in Fig.2, Formula 27 in Fig.3, Formula 39 in Fig.4, Formula 49 in Fig.5, or Formula 57 in Fig.6) as starting materials, reacting the

compound with bis(trialkyltin) and bis(triphenylphosphine)palladium chloride in anhydrous 1,4-dioxane under heat at reflux in an argon atmosphere, followed by purification.

Compound 10 (ITDU) in Fig.1 can be synthesized by a known method
5 (Formulae 1-8: Dyson, MR et al., Carbo. Res. 216, p.237 (1991), and Formulae 8-10: Oivanen, M et al., J. Chem. Soc., Perkin Trans. 2, p.2343 (1998)). Specifically, 2-deoxy-D-erythro-pentose (Compound 1) is reacted with a 1% hydrochloric acid-methanol solution to produce Compound 2, which is then reacted with sodium hydride, tetrabutylammonium iodide, and benzyl bromide to produce Compound 3, in which
10 hydroxyl groups are protected. The compound is reacted with α -toluenethiol and concentrated hydrochloric acid to produce Compound 4, which is then reacted with triphenylphosphine, benzoic acid, and diethylazodicarboxylate to produce Compound 5. Sodium methoxide is then used to remove the benzoyl group from Compound 5 to produce Compound 6, followed by its conversion into Compound 7 with
15 methanesulfonyl chloride. A ring is formed with sodium iodide and barium carbonate to produce Compound 8, which is reacted with 5-iodouracil in the presence of bistrimethylsilylacetamide and then with N-iodosuccinimide to produce Compound 9. Subsequently, Compound 9 is deprotected with titanichloride to produce Compound 10.

20 Compound 20 (ITAU) in Fig.2 can be synthesized by a known method (Formulae 13-17: Yoshimura Y et al., J. Org. Chem.61, p.822 (1996) and Formula 17-20: Yoshimura Y et al., J. Med. Chem. 40, p.2177 (1997)). Specifically, 1,2;5,6-di-O-isopropylidene glucose (Compound 13) is reacted with sodium hydride and benzyl bromide to produce a 3-benzyl compound, which is subsequently reacted with
25 hydrochloric acid, aqueous sodium periodate solution, and sodium borohydride to

produce Compound 14, which is then converted with hydrogen chloride into Compound 15. The compound is then reacted with mesyl chloride and sodium sulfide to produce Compound 16, which is reacted with hydrochloric acid and sodium borohydride successively to produce Compound 17. Hydroxyl groups are protected with sodium
5 hydride and benzyl bromide (Compound 18), and the resulting compound is converted to Compound 19 with m-chloroperbenzoic acid (m-CPBA) and acetic anhydride. It is further reacted with 5-iodouracil in the presence of 1,1,1,3,3,3-hexamethylene disilazane (HMDS) to produce a glycosylated compound, which is then reacted with boron chloride to produce Compound 20.

10 Compound 27 in Fig.3 can be produced as follows. Compound 17 shown in Fig.2 is used as a starting material, which is reacted with t-butyldimethylsilyl chloride (TBDMSCl) in dimethylformamide (DMF) in the presence of imidazole to protect the hydroxyl group at the "5" position with a silyl group to produce Compound 23. Trifluoromethanesulfonic acid anhydride (Trf_2O) is added thereto in pyridine to produce
15 Compound 24 in which the hydroxyl group at the "2" position is trifluoromethanesulfonylated. The compound is reacted with potassium fluoride, along with Kryptofix (registered trademark) 222 and potassium carbonate, in acetonitrile, to produce a fluoride compound (Compound 25) in which the substituent at the "2" position is stereochemically reversed. The compound is reacted with m-
20 chloroperbenzoic acid (m-CPBA) in methylene chloride and further treated with acetic anhydride to produce Compound 26. This is reacted with the product resulting from a reaction of 5-iodouracil and 1,1,1,3,3,3-hexamethylene disilazane (HMDS), and with trifluoromethanesulfonic acid trimethylsilyl (TMSOTf). The resulting product is further treated with boron chloride in methylene chloride to produce Compound 27.

25 Compound 39 (FIAU) in Fig.4 can be synthesized by a known method

(Formulae 30-37: Reichman U et al., Carbohydrate Res. 42, p.233 (1975) and Formulae 37-39: Asakura J et al., J. Org. Chem. 55, p.4928 (1990)). Specifically, Compound 31, which has been synthesized in four steps from 1,2:5,6-di-O-isopropylidene glucose (Compound 30), is treated with a cation exchange resin (Amberlite IR-120) to produce
5 Compound 32, which is then reacted with potassium periodate to produce Compound 33. This is then reacted with sodium methoxide to produce Compound 34, followed by acetylation of hydroxyl groups to produce Compound 35. The compound is treated with a hydrogen bromide-acetic acid solution to produce Compound 36, followed by condensation with an uracil derivative to produce Compound 37. It is subsequently
10 reacted with diammonium cerium(III) sulfate (CAN) to produce Compound 38, followed by deprotection of hydroxyl groups with sodium methoxide to produce Compound 39.

Compound 49 (FITAU) in Fig.5 can be synthesized by a known method (Formulae 42-46: Yoshimura Y et al., J. Org. Chem. 62, p.3140 (1997) and Formulae
15 46-49: Yoshimura Y et al., Bioorg. Med. Chem. 8, p.1545 (2000)). Specifically, Compound 43, which has been synthesized in nine steps from 1,2:5,6-di-O-isopropylidene glucose (Compound 42), is reacted with diethylaminosulfur trifluoride (DAST) to produce Compound 44, which is then reacted with *m*-chloroperbenzoic acid (*m*-CPBA) to produce Compound 45. This is subsequently reacted with acetic
20 anhydride to produce Compound 46, which is reacted with trifluoromethanesulfonic acid trimethylsilyl (TMSOTf) to cause condensation with a 5-iodouracil derivative to produce Compound 47. Finally, the two protective hydroxyl groups are removed to produce Compound 49.

Compound 57 (IMBAU) in Fig.6 can be synthesized by a known method
25 (Formulae 52-54: Itoh Y et al., J. Org. Chem. 60, p.656 (1995) and Formulae 55-56:

Asakura J et al., J. Org. Chem. 55, p.4928 (1990)), combined with known reactions for protection and deprotection of hydroxyl groups (Formulae 54-55 and Formula 56-57). Specifically, 1-[3,5-bis-O-(tert-butyldimethylsilyl)-2-deoxy-D-erythro-pento-1-enofuranosyl]uracil (Compound 52) is reacted with pivalic acid and bromosuccinimide (NBS) to produce Compound 53, which is then reacted with trimethylaluminum to produce Compound 54. The protection groups for hydroxyl groups are converted from tert-butyldimethylsilyl to acetyl, followed by reaction with diammonium cerium(III) sulfate (CAN) to produce Compound 56. Finally, the protection groups in Compound 56 are removed with ammonia to produce Compound 57.

For radiolabeled compounds of the present invention, appropriate doses and routes of administration should be selected depending upon target diseases and objectives, but if they are used as an agent for diagnosis of tissue proliferation activity, a radioactivity in the range of 37MBq to 740MBq, preferably 111MBq to 370MBq is administered. Usually, they are administered intravenously, but in some cases, other routes of administration including arterial or intraperitoneal administration and direct administration to a tumor or other affected portions may be used.

If they are used as an agent for treatment of proliferative disease, a radioactivity in the range of 37MBq to 7400MBq, preferably 185MBq to 3700MBq, is administered. Usually, they are administered intravenously, but in some cases, other routes of administration including arterial or intraperitoneal administration and direct administration to a tumor or other affected portions may be used. Furthermore, if they are used for therapeutic purposes, the above dose may be administered several times at appropriate intervals.

The agent for diagnosis of tissue proliferation activity of the present invention can serve for whole-body or local scintigraphy and whole-body or local SPECT

imaging by use of nuclides for SPECT. Using nuclides for PET, they can also be applied for whole-body or local PET imaging.

The agent for diagnosis of tissue proliferation activity of the present invention can serve for quantitative determination of local proliferative activity based on
5 appropriate model analysis. Furthermore, if non-proliferation tissue is used as a control, local proliferative activity can be defined easily in a semi-quantitative way.

The agent for treatment of proliferative disease of the present invention, when a beta-emitter such as I-131 is used therein, can serve to decrease large tumors of 1cm or more in diameter, depending on the range of the ray. When an alpha-emitter such as
10 At-211 is used, they can work on small lesions of 0.1mm or less in diameter more effectively than beta-emitter, and therefore, they are expected to serve for treatment of micrometastasis over the body. Furthermore, nuclides that emit Auger electrons, such as I-125, can have antitumor effects due to DNA breakage, only after labeled compounds have gathered around the DNAs. Therefore, suitable label nuclides for
15 treatment of systemic tumor foci including metastatic ones include alpha-emitter such as At-211, and beta-emitter such as I-131 that can have effect on portions around the foci depending on the range. The most effective method is the cocktail therapy which uses a mixture of a compound labeled with an alpha-emitter and a compound labeled with a beta-emitter.

20 For treatment by local administration, compounds labeled with nuclides that emit Auger electrons, such as I-125, are particularly effective for brain tumor that is difficult to remove completely by surgical operation, and residual tumor from malignant melanoma, and in view of functional preservation, breast cancer, rectal cancer, prostatic cancer, and malignant mouth tumor, because they do no harm on portions other than
25 pathologically proliferating cells owing to the properties of rays emitted therefrom.

Technique for local administration includes, for example, an administration into intracavitary foci such as colon cancer by use of an endoscope, a direct administration to foci affected by brain tumor during craniotomy, and an administration by use of a catheter into an artery relevant to an affected organ such as liver affected by cancer.

5 **Brief Description of the Drawings**

Fig.1 illustrates a synthetic pathway for a compound of the present invention.

Fig.2 illustrates another synthetic pathway for a compound of the present invention.

10 Fig.3 illustrates a third synthetic pathway for a compound of the present invention.

Fig.4 illustrates a fourth synthetic pathway for a compound of the present invention (5-trialkyltin compounds) and [I-125] FIAU produced therefrom.

Fig.5 illustrates a fifth synthetic pathway for a compound of the present invention.

15 Fig.6 illustrates the other synthetic pathway for a compound of the present invention.

Fig.7 illustrates a diagram showing *in vivo* label stability of [I-125] ITDU and [I-125] ITAU measured in Example 16, along with [I-125] IUR as a control.

20 Fig.8 illustrates a diagram showing *in vivo* label stability of [I-125] ITDU, [I-125] ITAU, [I-125] FITAU and [I-125] IMBAU measured in Example 17, along with [I-125] IUR (highly decomposable) as a control.

Fig.9 illustrates a diagram showing *in vivo* distribution of [I-125] ITDU in normal mice measured in Example 18.

25 Fig.10 illustrates a diagram showing *in vivo* accumulation of [I-125] ITDU, [I-125] ITAU, [I-125] FITAU and [I-125] IMBAU, along with [I-125] IUR as a control, in

proliferating tissue measured in Example 18.

Fig.11 illustrates a photograph (biological morphology) showing a scintigram of Walker tumor observed in Example 19.

Examples

5 The present invention will be described in detail below with reference to examples, but is not limited to these examples.

Example 1

Synthesis of 5-trimethylstannyl-4'-thio-2'-deoxyuridine (Compound 11)

As shown in Fig.1, benzyl-3,5-di-O-benzyl-2-deoxy-1,4-dithio- α,β -D-erythro-
10 pentofuranoside (Compound 8) was synthesized, using 2-deoxy-D-erythro-pentose (Compound 1) as starting material, according to the method of Dyson MR et al. (Carbo. Res. 216, p.237 (1991)). Further, 5-iodo-4'-thio-2'-deoxyuridine (ITDU: Compound 10) was produced from Compound 8 according to the method of Oivanen M et al. (J. Chem. Soc., Perkin Trans. 2, p.2343 (1998)). Compound 10 was then used as a
15 starting material to produce 5-trimethylstannyl-4'-thio-2'-deoxyuridine (Compound 11) according to the following procedure.

Compound 10 (9.5mg, 0.026mmol), bis(trimethyltin) (17.3mg, 0.052mmol) and bis(triphenylphosphine)palladium(II) chloride (5mg) were dissolved in anhydrous 1,4-dioxane (3mL) under argon atmosphere, and after heating at reflux for 3 hours,
20 concentrated under reduced pressure. The residue was purified by silica gel thin layer chromatography (chloroform-methanol, 6:1) to produce the target Compound 11 (6.9mg, 65%).

¹H NMR (270MHz, CD₃OD) δ 0.26 (s, 9H, CH₃Sn), 2.26 (ddd, 1H, J=4.6, 7.9, 13.2 Hz, 1H, H-2'), 2.27 (ddd, J=4.6, 6.6, 13.4 Hz, 1H, H-2'), 3.41 (m, 1H, H-4'), 3.71 (dd, J=5.9, 11.2 Hz, 1H, H-5'), 3.80 (dd, J=4.6, 11.2 Hz, 1H, H-5'), 4.47 (q, J=4.0 Hz, 1H, H-3'),
25

6.41 (t, J=7.2 Hz, 1H, H-1'), 7.93 (s, 1H, H-5).

Example 2

Synthesis of [I-125]-5-iodo-4'-thio-2'-deoxyuridine ([I-125] ITDU: Compound 12)

To 0.1N sodium hydroxide solution (50 μ L) of [I-125]-sodium iodide (33MBq),
5 water (1mL) and chloroform (1mL) were added, and then chloroform solution (4.7 μ L)
of iodine (60 μ g, 0.47 μ mol) was added, and shaken for 10 seconds. After removing
only the aqueous layer, ethyl acetate solution (100 μ L) of Compound 11 (100 μ g,
0.25 μ mol) was added, and the resulting solution was left to stand at room temperature
for 2 hours. One drop of 1N sodium thiosulfate solution was added, and chloroform
10 was evaporated. After adding water (1mL), the solution was passed through a Sep-Pak
Plus QMA cartridge column. The column was washed with water (0.5mL $\times$ 2), and the
resulting aqueous solution was combined to produce I-125-labeled Compound 12
(7.3MBq, 22%).

Example 3

Synthesis of [I-123]-5-iodo-4'-thio-2'-deoxyuridine ([I-123] ITDU: Compound 12)

To 0.1% ammonium iodide solution (1mL) containing [I-123]-ammonium
iodide (2.0GBq), 1N hydrochloric acid (0.1mL) and chloroform (1mL) were added, and
then chloroform solution (4.7 μ L) of iodine (60 μ g, 0.47 μ mol) was added, and shaken for
10 seconds. After removing only the aqueous layer, ethyl acetate solution (100 μ L) of
20 Compound 11 (100 μ g, 0.25 μ mol) was added and left to stand at room temperature for 2
hours. One drop of 1N sodium thiosulfate solution was added, and chloroform was
evaporated. After adding water (1mL), the solution was passed through a Sep-Pak
Plus QMA cartridge column. The column was washed with water (0.5mL $\times$ 2), and the
resulting aqueous solution was combined to produce I-125-labeled Compound 12
25 (228MBq, 15%).

Example 4

Synthesis of 5-trimethylstannyl-1-(4-thio-D-arabinofuranosyl)uracil (Compound 21)

As shown in Fig.2, 1,4-anhydro-3-O-benzyl-4-thio- α -D-arabitol (Compound 17) was synthesized from 1,2,5,6-di-O-isopropylidene glucose (Compound 13) according to the method of Yoshimura Y et al. (J. Org. Chem. 61, p.822 (1996)). Then, 5-iodo-1-(4-thio-D-arabinofuranosyl)uracil (ITAU: Compound 20) was produced from Compound 17 according to the method of Yoshimura Y et al. (J. Med. Chem.40, p.2177 (1997)). This Compound 20 was used as a starting material to produce 5-trimethylstannyl-1-(4-thio-D-arabinofuranosyl)uracil (Compound 21) by the following procedure.

Compound 20 (4.0mg, 0.010mmol), bis(trimethyltin) (6.6mg, 0.020mmol) and bis(triphenylphosphine)palladium(II) chloride (5mg) were dissolved in anhydrous 1,4-dioxane (5mL) in an argon atmosphere, and after heating at reflux for 4 hours, concentrated under a reduced pressure. The residue was purified by silica gel thin layer chromatography (25% methanol/chloroform) to produce the target Compound 21 (2.3mg, 55%).

¹H NMR (270MHz, CD₃OD) δ 0.7 (s, 9H), 3.55-3.67 (m, 1H), 3.77-3.95 (m, 2H), 4.07 (t, J=5.9 Hz, 1H), 4.16 (t, J=5.9, 1H), 6.28 (d, J=5.3 Hz, 1H), 8.03 (s, 1H).

Example 5

Synthesis of [I-125]-5-iodo-1-(4-thio-D-arabinofuranosyl)uracil ([I-125] ITAU: Compound 22)

To 0.1N sodium hydroxide solution (50 μ L) of [I-125]-sodium iodide (67MBq), water (1mL) and chloroform (1mL) were added, and then chloroform solution (4.7 μ L) of iodine (60 μ g, 0.47 μ mol) was added, and shaken for 10 seconds. After removing only the aqueous layer, ethyl acetate solution (100 μ L) of Compound 21 (100 μ g,

0.24 μ mol) was added, and the resulting solution was left to stand at room temperature for 2 hours. One drop of 1N sodium thiosulfate solution was added, and chloroform was evaporated. After adding water (1mL), the solution was passed through a Sep-Pak Plus QMA cartridge column. The column was washed with water (0.5mL $\times$ 2), and the
5 resulting aqueous solution was combined to produce I-125-labeled Compound 22 (17.3MBq, 26%).

Example 6

Synthesis of 5-trimethylstannyl-1-(2-deoxy-2-fluoro- β -D-arabinopentofuranosyl)uracil (Compound 40)

10 As shown in Fig.4, 1-(3,5-di-O-acetyl-2-deoxy-2-fluoro- β -D-arabinopentofuranosyl)uracil (Compound 37) was synthesized from 1,2:5,6-di-O-isopropylidene glucose (Compound 30) according to the method of Reichman U et al. (Carbohydrate Res. 42, p.233 (1975)). Further, 5-iodo-1-(2-deoxy-2-fluoro- β -D-arabinopentofuranosyl)uracil (Compound 39) was produced from Compound 37
15 according to the method of Asakura J et al. (J. Org. Chem.55, p.4928 (1990)). This compound was used as starting material to produce 5-trimethylstannyl-1-(2-deoxy-2-fluoro- β -D-arabinopentofuranosyl)uracil (Compound 40) by the following procedure.

Compound 39 (5.0mg, 0.013mmol), bis(trimethyltin) (20.5mg, 0.063mmol) and bis(triphenylphosphine)palladium(II) chloride (6.2mg) were dissolved in anhydrous
20 1,4-dioxane (3mL) in an argon atmosphere, and after heating at reflux for 2 hours, concentrated under a reduced pressure. The residue was purified by silica gel thin layer chromatography (chloroform-methanol, 6:1) to produce the target Compound 40 (3.6mg, 66%).

¹H-NMR(500MHz, CD₃OD) δ 0.25 (s, 9H, CH₃Sn), 3.72 (dd, J=5.0, 12.0Hz, 1H, H-5'),
25 3.79-3.91 (m, H-4'), 4.33 (ddd, J=3.0, 5.0, 18.5 Hz, 1H, H-3'), 5.02 (td, J=4.0, 53.0 Hz,

1H, H-2'), 6.25 (dd, J=4.5, 16.0 Hz, 1H, H-1'), 7.56 (s, 1H, H-5).

Example 7

Synthesis of [I-125]-5-iodo-1-(2-deoxy-2-fluoro- β -D-arabinopentofuranosyl)uracil ([I-125] FIAU: Compound 41)

5 First, 0.1N sodium hydroxide solution of [I-125]-sodium iodide (80MBq) was distilled off, followed by addition of methanol (1mL), addition of methanol solution (4.8 μ L) of iodine (61 μ g, 0.48 μ mol), and shaking for 10 seconds. Then, methanol solution (100 μ L) of Compound 40 (100 μ g, 0.24 μ mol) was added, and the solution was left to stand at room temperature for 2 hours. One drop of 1N sodium thiosulfate
10 solution was added, and methanol was evaporated. After adding water (1mL), the solution was passed through a Sep-Pak Plus QMA cartridge column. The column was washed with water (1.0mL), and the resulting aqueous solution was combined to obtain I-125-labeled Compound 41 (9.5MBq, 12%).

Example 8

15 Synthesis of 5-trimethylstannyl-1-(2-deoxy-2-fluoro-4-thio- β -D-arabinopentofuranosyl)uracil (Compound 50)

As shown in Fig.5, 1-O-acetyl-3-O-benzyl-5-O-(tert-butyldiphenylsilyl)-2-deoxy-2-fluoro-4-thio-D-arabinopentofuranose (Compound 46) was synthesized from 1,2:5,6-di-O-isopropylidene glucose (Compound 42) according to the method of Yoshimura
20 Y et al. (J. Org. Chem.62, p.3140 (1997)). Further, Compound 46 was used to produce 5-iodo-1-(2-deoxy-2-fluoro-4-thio- β -D-arabinopentofuranosyl)uracil (Compound 49) according to the method of Yoshimura Y et al. (Bioorg. Med. Chem.8, p.1545 (2000)). This compound was then used as a starting material to produce 5-trimethylstannyl-1-(2-deoxy-2-fluoro-4-thio- β -D-arabinopentofuranosyl)uracil (Compound 50) by the
25 following procedure.

Compound 49 (5.0mg, 0.013mmol), bis(trimethyltin) (16.9mg, 0.052mmol) and bis(triphenylphosphine)palladium(II) chloride (6.0mg) were dissolved in anhydrous 1,4-dioxane (3mL) in an argon atmosphere, and after heating at reflux for 3.5 hours, concentrated under reduced pressure. The residue was purified by silica gel thin layer chromatography (chloroform-methanol, 6:1) to produce the target Compound 50 (1.9mg, 35%).

¹H-NMR (500MHz, CD₃OD) δ 0.26 (s, 9H, CH₃Sn), 3.61-3.68 (m, 1H, H-5'), 3.80-3.81 (m, H-4'), 4.37 (td, J=6.0, 12.0 Hz, 1H, H-3'), 4.97 (td, J=5.5, 49.0 Hz, 1H, H-2'), 6.46 (dd, J=5.5, 11.5 Hz, 1H, H-1'), 7.99 (s, 1H, H-5).

10 Example 9

Synthesis of [I-125]-5-iodo-1-(2-deoxy-2-fluoro-4-thio-β-D-arabinopentofuranosyl)uracil ([I-125] FITAU: Compound 51)

First, 0.1N sodium hydroxide solution of [I-125]-sodium iodide (45MBq) was distilled off, followed by addition of methanol (1mL), addition of methanol solution (4.8μL) of iodine (61μg, 0.48μmol), and shaking for 10 seconds. Then, methanol solution (100μL) of Compound 50 (100μg, 0.24μmol) was added, and the resulting solution was left to stand at room temperature for 2 hours. One drop of 1N sodium thiosulfate solution was added, and methanol was evaporated. After adding water (1mL), the solution was passed through a Sep-Pak Plus QMA cartridge column. The column was washed with water (1.0mL), and the resulting aqueous solution was combined to obtain I-125-labeled Compound 51 (3.5MBq, 7.8%).

Example 10

Synthesis of 5-trimethylstannyl-1-methyl(2-deoxy-2-bromo- β -D-arabinopentofuranosyl)uracil ([I-125] IMBAU: Compound 58)

As shown in Fig.6, 1-[2-bromo-3,5-bis-O-(tert-butyldimethylsilyl)-2-deoxy-1-C-methyl- β -D-arabinofuranosyl]uracil (Compound 54) was produced from 1-[3,5-bis-O-(tert-butyldimethylsilyl)-2-deoxy-D-erythro-pento-1-enofuranosyl]uracil (Compound 52) according to the method of Itoh Y et al. (J. Org. Chem. 60, p.656 (1995)). Further, 5-iodo-1-methyl(2-deoxy-2-bromo- β -D-arabinopentofuranosyl)uracil (Compound 57) was produced from Compound 54 according to the method of Asakura J et al. (J. Org. Chem. 55, p.4928 (1990)). This compound was used as starting material to produce 5-trimethylstannyl-1-methyl(2-deoxy-2-bromo- β -D-arabinopentofuranosyl)uracil (Compound 58) by the following procedure.

Compound 57 (4.9mg, 0.011mmol), bis(trimethyltin) (16.0mg, 0.049mmol) and bis(triphenylphosphine)palladium(II) chloride (5mg) were dissolved in anhydrous 1,4-dioxane (3mL) in an argon atmosphere, and after heating at reflux for 2.5 hours, concentrated under a reduced pressure. The residue was purified by silica gel thin layer chromatography (chloroform-methanol, 6:1) to produce the target Compound 58 (3.8mg, 72%).

^1H NMR (500MHz, CD_3OD) δ 0.25 (s, 9H, CH_3Sn), 1.95 (s, 3H, $1'\text{-CH}_3$), 3.61-3.70 (m, 2H, H-5'), 4.08-4.11 (m, 1H, H-4'), 4.53 (d, $J=3.0$ Hz, 1H, H-5'), 4.79 (s, 1H, H-2'), 7.75 (s, 1H, H-5).

Example 11

Synthesis of [I-125]-5-iodo-1-methyl-(2-deoxy-2-bromo- β -D-arabinopentofuranosyl)uracil ([I-125] IMBAU: Compound 59)

First, 0.1N sodium hydroxide solution of [I-125]-sodium iodide (62MBq) was

distilled off, followed by addition of methanol (1mL), addition of methanol solution (4.0μL) of iodine (51μg, 0.40μmol), and shaking for 10 seconds. Then, methanol solution (100μL) of Compound 58 (100μg, 0.20μmol) was added, and the resulting solution was left to stand at room temperature for 2 hours. One drop of 1N sodium
5 thiosulfate solution was added, and methanol was evaporated. After adding water (1mL), the solution was passed through a Sep-Pak Plus QMA cartridge column. The column was washed with water (1.0mL), and the resulting aqueous solution was combined to obtain I-125-labeled Compound 59 (8.3MBq, 13%).

Example 12

10 Test for *in vitro* phosphorylation activity of [I-125] ITDU and [I-125] ITAU

The phosphorylation activity of a labeled compound by thymidine kinase was determined using a crude enzyme extracted from a mouse's lung cancer cell strain LL/2. Liquid enzyme was extracted from a LL/2 mouse's lung cancer cell strain in the logarithmic growth phase according to the method of Wolcott RM and Colacino JM
15 (Anal. Biochem 178, p.38-40 (1989)). To a reaction liquid containing ATP, which is a phosphate donor, 2nmole of the label compound and the liquid enzyme were added and reacted at 37°C for a fixed period of time. The reaction was stopped by adding 1mL of a 100mM lanthanum chloride/5mM triethanolamine solution. Phosphorylated material
20 by measuring a radioactivity of the resulting precipitate with an automatic well-type gamma counter (ARC-380, Aloka Co., Ltd.). Results are shown in Table 1 from which phosphorylation activity attributed to thymidine kinase was confirmed in both [I-125] ITDU and [I-125] ITAU.

Table 1 : Phosphorylation activity of iodo-labeled nucleoside derivatives (n=3)

Iodo-labeled nucleosides	Phosphorylated material production rate (p mole/mg protein/h)
[I-125]ITDU	1182.7±100.1
[I-125]ITAU	13.6±6.6

Example 13Test for *in vitro* metabolic stability of [I-125] ITDU and [I-125] ITAU

To evaluate metabolic stability of glycosidic bond, decomposition reactivity for *E. coli*-originating thymidine phospholylase was studied. To the reaction liquid, 2nmole of the labeled compound and 9 units of a liquid enzyme (Sigma Corporation) were added and reacted at 25°C for a fixed period of time, and the reaction was stopped by treatment in a boiling water bath for 3 minutes. The reaction liquid was subjected to centrifugal separation, and the supernatant was applied over a thin layer silica gel plate along with an authentic standard (5-iodouracil: IU) and a non-labeled parent compound. It was developed with a mixture of chloroform and isopropyl alcohol (3:1), and then autoradiography was measured with a bioimaging analyzer (BAS-1500, Fuji Photo Film Co., Ltd.). The area of interest was set to peak components of the R_f value corresponding to the authentic standard, and the amount of the resulting metabolite was calculated from its proportion in percentage. Results are shown in Table 2 which indicates that [I-125] ITDU and [I-125] ITAU are stabler than 5-iododeoxyuridine ([I-125] IUR).

Table 2 : C-N glycosidic bond cleavage activity for iodo-labeled nucleoside derivatives (n=3)

Iodo-labeled nucleosides	5-iodouracil production rate* (relative activity)
[I-125]IUR	138606.2±14902.3 (1.00)
[I-125]ITDU	4075.9±736.4 (0.03)
[I-125]ITAU	524.3±373.8(<0.01)

* p mole/units/30 min

Example 14

Evaluation of *in vitro* stability of metabolism of various radioactive-iodine-labeled nucleoside derivatives by thymidine phospholylase

To evaluate the metabolic stability of glycosidic bond in various radioactive iodine-labeled nucleoside derivatives, their decomposition reactivity for *E. coli*-originating thymidine phospholylase was studied. To the reaction liquid, 0.5-12.0nmol of the labeled compound and 0.0009-9.0 units of a liquid enzyme (Sigma Corporation) was added and reacted at 25°C for a fixed period of time, followed by treatment in a boiling water bath for 3 minutes to stop the reaction. As [I-125] IBMAU was unstable under heat treatment, the reaction liquid was cooled with ice to stop the reaction. The reaction liquid was subjected to centrifugal separation, and the supernatant was applied over a silica gel plate along with an authentic standard

(5-iodouracil: IU) and a non-labeled parent compound. It was developed with a mixture of chloroform and isopropyl alcohol (3:1), and then the autoradiogram was measured with a bioimaging analyzer (BAS-1500, Fuji Photo Film Co., Ltd.). The area of interest was set to peak components of the R_f value corresponding to the authentic standard, and the amount of the resulting metabolite was calculated from its proportion in percentage. In the case of [I-125] FITAU and [I-125] IMBAU, a reversed phase silica gel plate was used, and after development with a mixture of methanol and water (3:7), an autoradiogram was

measured with a bioimaging analyzer (BAS-1500, Fuji Photo Film Co., Ltd.) similarly to the [I-125] IUR and others. Results of analysis are shown in Table 3 which indicates that [I-125] FITAU and [I-125] IMBAU are still stabler than [I-125] IUR.

Table 3: C-N glycosidic linkage cleavage activity of iodo-labeled nucleoside derivatives by thymidine phosphorylase for (n=3)

Iodo-labeled nucleosides	5-IU production rate (pmol/units/0.5h)	Relative activity
[I-125]IUR	138606.2±14902.3	1.00
[I-125]ITdU	3778.7±692.0	0.03
[I-125]ITAU	514.8±367.0	<0.01
[I-125]FITAU	0.5±0.1	<0.00001
[I-125]IMBAU	0.0	0.0

Example 15

Test of thymidine kinase-dependent incorporation into cells using thymidine kinase-deficient cells

Thymidine kinase-dependent incorporation of labeled compounds into cells was studied based on difference in incorporation between thymidine kinase-deficient cell strains L-M (TK-) and their parent L-M cells. L-M and L-M (TK-) cells in the logarithmic growth phase were planted on 24-well plates, each carrying 2.0×10^5 cells, and cultured overnight. Then 2nmol of a radioactive iodine-labeled nucleoside derivative was added and allowed to be incorporated in the cells for one hour. The cells were washed three times with an ice-cooled phosphate buffer solution, and dissolved in 0.1N NaOH, followed by determination of degree of radioactivity incorporated in the cells using an automatic well-type gamma counter (ARC-380 or ARC-300, Aloka Co., Ltd.). Measurements were analyzed to make evaluations based on the amount of the incorporated label molecules per unit weight of cellular proteins. Results are shown in Table 4 which indicates that [I-125] ITdU and [I-125] FITAU were

incorporated in cells in a thymidine kinase dependent way as in the case of the [I-125] IUR as a control.

5 Table 4: Incorporation of radioactive iodine-labeled nucleosides in L-M and L-M (TK-) cells

Iodo-labeled nucleosides	Incorporation (pmol/mg protein/h)		(L-M)/ {L-M(TK-)}
	L-M	L-M (TK-)	
[I-125] IUR	77.80±7.45	27.86±2.94	2.79*
[I-125] ITdU	10.90±1.48	3.94±0.63	2.77*
[I-125] ITAU	1.68±0.28	1.20±0.20	1.40**
[I-125] FITAU	0.34±0.05	0.21±0.05	1.62***

*p<0.0005, **p<0.05, ***p<0.01(T-test)

Example 16

Test for *in vivo* label stability of [I-125] ITDU and [I-125] ITAU

10 To evaluate *in vivo* label stability of [I-125] ITDU and [I-125] ITAU, tests were conducted to study the accumulation of free iodine in the thyroid gland in normal mice. A 370KBq portion of each labeled compound was injected in each of 10 week old normal mice into its tail vein, and three animals were sacrificed and anatomized at appropriate intervals. For a control, *in vivo* distribution of [I-125] IUR was also
15 observed. Incorporation of radioactivity in the thyroid gland was measured with an automatic well-type gamma counter (ARC-300, Aloka Co., Ltd.). Incorporated radioactivity in tissue was calculated as the administrated dose per gram of the tissue per unit time, and represented in percentage, as shown in Fig.7. Results indicate that the accumulated radioactivity from [I-125] ITDU and [I-125] ITAU in the thyroid gland
20 was significantly smaller than that from the control [I-125] IUR, proving that the *in vivo* label stability of the agents is high.

Example 17*In vivo* label stability of radioaction iodine-labeled nucleoside derivatives

To evaluate *in vivo* stability of deiodination against each radioactive iodine-labeled nucleoside derivative, tests were conducted to study accumulation of free iodine in the thyroid gland of normal mice. A 185KBq of each labeled compound was injected in each of 10 week old normal mice (C57BL/6) into the tail vein, and three animals were sacrificed and anatomized at intervals longer than in Example 16. Incorporation of radioactivity in the thyroid gland was measured with an automatic well-type gamma counter (ARC-300, Aloka Co., Ltd.). Incorporated radioactivity in tissue (%ID) was calculated as the administrated per gram of the tissue, and represented in percentage, as shown in Fig.8. Results indicate that the accumulated radioactivity from [I-125] ITDU, [I-125] ITAU, [I-125] FITAU and [I-125] IMBAU in the thyroid gland was significantly smaller than that from the control [I-125] IUR (highly metabolizable substance), proving that the *in vivo* label stability of the agents is high.

Example 18*In vivo* distribution of [I-125] ITDU in normal mice

A 370KBq of [I-125] ITDU was injected in each of 10 week old normal mice into the tail vein, and three animals were sacrificed and anatomized at appropriate intervals. Incorporation of radioactivity in each tissue sample was measured with an automatic well-type gamma counter (ARC-300, Aloka Co., Ltd.). Incorporated radioactivity in tissue was calculated as the administrated dose per gram of the tissue, and represented in percentage, as shown in Fig.9. Results indicate that the accumulated radioactivity in proliferating tissues, namely the thymus and the small intestine, was certainly higher than that in non-proliferating tissues, namely the brain,

To evaluate the accumulation of each radioactive iodine-labeled nucleoside derivative in proliferating tissues, tests were conducted to study *in vivo* distribution in normal mice. A 185MBq of each labeled compound was injected in each of 10 week old normal mice (C57BL/6) through its tail vein, and three animals were sacrificed and anatomized at appropriate intervals. Incorporation of radioactivity in each tissue sample was measured with an automatic well-type gamma counter (ARC-300, Aloka Co., Ltd.). Incorporated radioactivity in tissue was calculated as the administrated dose per unit weight of the tissue, and represented in percentage (%ID/g). As shown in Fig.10, results indicate that [I-125] IUR (positive control) and [I-125] ITDU have accumulated in large amounts particularly in the thymus which is a proliferating tissue in normal young mice.

Example 19

Scintigraphy of Walker tumor using [I-123] ITDU


Malignant tumor, a typical proliferative disease, was observed by scintigraphy. Walker tumor cells were transplanted subcutaneously in the right inguinal region of Wistar rats. After the transplantation, 37MBq of [I-123] ITDU was injected into the tail vein of rats that suffered a palpable tumor of about 20mm that was suitable for scintigraphy. Each tumor-transplanted rat was anesthetized with Ravonal four hours after the administration of a drug. Then the rat was fixed in the face-up position and observed statically with gamma-camera imaging equipment (GCA-90B, Toshiba Corporation). Imaging was performed using a high-resolution medium-energy collimator to obtain images for 10 minutes with a resolution of 256×256. Results are illustrated in Fig.11 which shows that [I-123] ITDU serves for clear imaging of transplanted tumors (indicated by an arrow) in Wistar rats.

Industrial Applicability

The radiolabeled compounds of the present invention are stable *in vivo*, and they either retain in cells after being phosphorylated by mammal thymidine kinase or are incorporated in DNA to reflect the DNA synthesis activity, thus serving for
5 diagnosis of tissue proliferation activity and treatment of proliferative diseases, particularly as radioactive diagnostic imaging agents for tissue proliferation activity diagnosis and as radioactive therapeutic agents for proliferative disease treatment by internal radiotherapy, local radiotherapy and the like.

Claims

1. (amended) An agent for diagnosis of tissue proliferation activity
which comprises, as an active ingredient, a
radiolabeled compound as represented by the following formula or a pharmaceutically
acceptable salt thereof:



wherein R₁ denotes hydrogen, or a linear- or branched-chain alkyl group having 1-8 carbon atoms; R₂ denotes hydrogen, hydroxyl, or a halogen substituent; R₃ denotes hydrogen or fluorine substituent, R₄ denotes oxygen, sulfur or a methylene substituent, and R₅ denotes a radioactive halogen substituent, excluding the case where R₁ is hydrogen, R₄ being oxygen.

2. An agent according to Claim 1, wherein said radioactive halogen-substituent in R₅ is selected from the group consisting of F-18, Cl-36, Br-75,

Br-76, Br-77, Br-82, I-123, I-124, I-125, I-131 and At-211.

3. An agent according to Claim 1, wherein R_4 is sulfur.
4. An agent according to Claim 1, wherein R_1 is hydrogen or methyl group, R_2 is hydrogen or a halogen-substituent, R_3 is hydrogen, and R_4 is oxygen or sulfur.
5. An agent according to Claim 1, wherein R_1 , R_2 and R_3 are each hydrogen, R_4 is sulfur, and said radioactive halogen-substituent in R_5 is selected from the group consisting of F-18, I-123, I-125 and I-131.
6. An agent for diagnosis of tissue proliferation activity according to Claim 1, wherein said diagnosis of tissue proliferation activity is diagnosis of hyperplasia, regeneration, transplantation or viral infection accompanied by abnormal proliferation.
7. An agent for diagnosis of tissue proliferation activity according to Claim 6, wherein said diagnosis of hyperplasia accompanied by abnormal proliferation is diagnosis of hyperplastic inflammation, benign tumor or malignant tumor.
8. An agent for diagnosis of tissue proliferation activity according to Claim 7, wherein said diagnosis of hyperplastic inflammation is diagnosis concerning activity of chronic rheumatoid arthritis or determination of therapeutic effects.
9. An agent for diagnosis of tissue proliferation activity according to Claim 7, wherein said diagnosis of benign tumor is diagnosis concerning localization, activity or determination of therapeutic effects.
10. An agent for diagnosis of tissue proliferation activity according to Claim 7, wherein said diagnosis of malignant tumor is diagnoses concerning localization, progress, malignancy or determination of therapeutic effects, of primary and metastatic malignant tumors.
11. An agent for diagnosis of tissue proliferation activity according to Claim 6, wherein said diagnosis of regeneration accompanied by abnormal proliferation is


evaluation of physiological hematopoietic functions of bone marrow during treatment with anti-cancer drugs, or diagnosis of pathological functions of the bone marrow in patients suffering from hypoplastic anemia.

12. An agent for diagnosis of tissue proliferation activity according to Claim 6, wherein said diagnosis of transplantation accompanied by abnormal proliferation is diagnosis of take or proliferation of transplanted bone marrow cells in bone marrow transplantation.

13. An agent for diagnosis of tissue proliferation activity according to Claim 6, wherein said diagnosis of viral infection accompanied by abnormal proliferation is diagnosis of virus-infected portions and proliferation thereof in infectious diseases caused by Type I or Type II herpes simplex virus, varicella-zoster herpes virus, cytomegalovirus, Epstein-Barr virus, or human immunodeficiency virus.

14. A nucleoside derivative as represented by the following formula:

- 38 -



wherein R_1 denotes hydrogen or a linear- or branched-chain alkyl groups having 1-8 carbon atoms, R_2 denotes hydrogen, hydroxyl or a halogen substituent, R_3 denotes hydrogen or fluorine substituent, R_4 denotes oxygen, sulfur or a methylene substituent, and R_5 denotes a trialkylstannyl group, excluding the case where R_1 is hydrogen, R_4 being oxygen.


15. A nucleoside derivative according to Claim 14, wherein R_4 is sulfur.

16. A nucleoside derivatives according to Claim 14, wherein R_1 is hydrogen or methyl group, R_2 is hydrogen or a halogen-substituent, R_3 is hydrogen, and R_4 is oxygen or sulfur.

17. A nucleoside derivatives according to Claim 14, wherein the trialkylstannyl group in R_5 is a trimethylstannyl group, a triethylstannyl group or a tributylstannyl group.

18. The use of radiolabeled compound in the manufacture of a medicament suitable for administration to a mammalian subject to diagnose tissue proliferation activity; said radiolabeled compound being represented by the following formula or a pharmaceutically acceptable salt thereof:


- 39 -



wherein R₁ denotes hydrogen, or a linear- or branched-chain alkyl group having 1-8 carbon atoms; R₂ denotes hydrogen, hydroxyl, or a halogen substituent; R₃ denotes hydrogen or fluorine substituent, R₄ denotes oxygen, sulfur or a methylene substituent, and R₅ denotes a radioactive halogen substituent, excluding the case where R₁ is hydrogen, R₄ being oxygen.


- 40 -

19. A method to producing a radiolabeled compound as represented by the following formula:



wherein R₁ denotes hydrogen or a linear- or branched-chain alkyl groups having 1-8 carbon atoms, R₂ denotes hydrogen, hydroxyl or a halogen substituent, R₃ denotes hydrogen or fluorine substituent, R₄ denotes oxygen, sulfur or a methylene substituent, and R₅ denotes a radioactive halogen substituent, excluding the case where R₁ is hydrogen, R₄ being oxygen;

comprising reacting a nucleoside derivative as represented by the following formula:



- 41 -

wherein R_1 denotes hydrogen or a linear- or branched-chain alkyl groups having 1-8 carbon atoms, R_2 denotes hydrogen, hydroxyl or a halogen substituent, R_3 denotes hydrogen or fluorine substituent, R_4 denotes oxygen, sulfur or a methylene substituent, and R_5 denotes a trialkylstannyl group, excluding the case where R_1 is hydrogen, R_4 being oxygen, with an alkaline solution of a radioactive halogen in a solvent, whereby the trialkylstannyl group of R_5 is replaced with a radioactive halogen substituent.