group.

[0120] Examples of hydrocarbylidene groups for R^{75} include ethylidene, propylidene, butylidene, 1-methylethylidene, 1-methylpropylidene and 1,3-dimethylbutylidene groups. The number of carbon atoms of a hydrocarbylidene group is preferably 2 to 20 and more preferably 2 to 6.

[0121] Examples of groups represented by formula (7-Y) include acyclic amino groups such as ethylideneamino, 1-methylpropylideneamino, 1,3-dimethylbutylideneamino, 1-methylethylideneamino and 4-N,N-dimethylaminobenzylideneamino groups.

[0122] In formula (7), the substituted amino groups for X^7, X^8 and X^9 are preferably acyclic amino groups, more preferably dialkylamino groups, even more preferably dimethylamino, diethylamino, di(n-propyl)amino or di(n-butyl)amino groups, and most preferably dimethylamino or diethylamino groups.

[0123] In formula (7), at least one of X^7, X^8 and X^9 is a substituted amino group, preferably two or more of X^7, X^8 and X^9 are substituted amino groups, and more preferably two of X^7, X^8 and X^9 are substituted amino groups.

[0124] Among compounds represented by formula (7), compounds wherein R^{71} is hydrogen and one of X^7, X^8 and X^9 is a dialkylamino group include the following compounds.

[0125] Examples of compounds wherein s in formula (7) is 0 include (dimethylamino)dimethylvinylsilane,
(diethylamino)dimethylvinylsilane,