

NOVEL PEPTIDE

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a novel peptide, and more particularly, to a novel peptide for promoting regeneration of dentin or dental pulp tissues and treating dentin hypersensitivity, a polynucleotide encoding the peptide, an expression vector including the polynucleotide, a pharmaceutical composition for preventing or treating dentin or dental pulp diseases including the peptide, a quasi-drug composition for preventing or improving dentin or dental pulp diseases including the peptide, and a health functional food composition for preventing or improving dentin or dental pulp diseases including the peptide.

2. Description of the Related Art

Dental pulp is a richly innervated and vascularized soft connective tissue that occupies the pulp chamber inside a tooth and extends to the outer surface of the dentin. Disorders occurring in the dental pulp are called dental pulp diseases.

There are many causes of dental pulp diseases, but in most cases, dental pulp diseases are caused by a bacterial

infection due to dental caries, or infections in the dental pulp through perforation, fracture, cracks, or periodontal pocket. External wound, abrasion, tooth cracks, or friction or heat from dental equipment may also cause dental pulp diseases. The pulpitis caused by bacterial infection may lead to root apex and periodontal diseases. Dental pulp diseases successively progress to pulp hyperemia, pulpitis, and pulp necrosis. Pulp necrosis may lead to periapical diseases or disorders to the entire tooth, because death of the dental pulp prevents the blood supply to the dental pulp and thus the entire pulp tissue is lost.

For treatment of the pulp or periapical diseases, pulp capping materials and pulp canal filling materials are used, and calcium hydroxides, MTA (Mineral Trioxide Aggregate), Gutta-percha, etc., has been generally used. MTA shows therapeutic effects because it has a leakage sealing ability and biocompatibility. However, use of MTA is hampered due to its relatively high cost as a dental repair material and discoloration leading to an esthetic problem. Gutta-percha is relatively low cost and has good flow characteristics. However, it is not a physiologically acceptable method which causes a loss of viability of the pulp. Up to now, conservative treatments for dentin and pulp diseases have problems of the weak or brittle teeth or reinfection.

Therefore, many studies have been actively conducted to develop therapeutic agents capable of effectively treating dentin or pulp diseases. For example, Korean Patent Publication No. 2012-0089547 discloses a composition for forming hard tissues or regenerating dentin or pulp tissues, including ameloblasts, apical bud cells, or cultures thereof as an active ingredient, and Korean Patent Publication No. 2009-0033643 discloses novel tooth stem cells derived from tooth follicles and a method of culturing the same.

The present inventors have made many efforts to develop an agent capable of more effectively treating dentin or dental pulp diseases, and as a result, they developed peptides showing effects of promoting regeneration of dentin or dental pulp tissues and treating dentin hypersensitivity, thereby completing the present invention.

SUMMARY OF THE INVENTION

Embodiments of the present inventive concepts may provide a new peptide for promoting regeneration of dentin or dental pulp tissues and treating dentin or dental pulp diseases.

Embodiments of the present inventive concepts may also provide a polynucleotide encoding the peptide.

Embodiments of the present inventive concepts may also provide an expression vector including the polynucleotide.

Embodiments of the present inventive concepts may also provide a pharmaceutical composition for preventing or treating dentin or dental pulp diseases including the peptide.

Embodiments of the present inventive concepts may also provide a quasi-drug composition for preventing or improving dentin or dental pulp diseases including the peptide.

Embodiments of the present inventive concepts may also provide a health functional food composition for preventing or improving dentin or dental pulp diseases including the peptide.

Embodiments of the present inventive concepts may also provide a method of preventing or treating dentin or dental pulp diseases, the method including administering the composition including the peptide to a subject.

Embodiments of the present inventive concepts may also provide a method of promoting regeneration of dentin or dental pulp tissues, the method including administering the composition including the peptide to a subject.

Embodiments of the present inventive concepts may also provide use of a peptide including an amino acid sequence of the following Formula 1 or a composition including the peptide in promoting regeneration of dentin or dental pulp

tissues, in preventing or treating dentin hypersensitivity, and in preventing or treating dentin or dental pulp diseases:

K-Y-R1-R2-R3-R4-R5-R6-R7-R8 (Formula 1)

wherein R1 is arginine(R), lysine(K) or glutamine(Q);

R2 is arginine(R) or glutamine(Q);

R3, R4, and R5 are arginine(R) or lysine(K), respectively;

R6 is asparagine(N) or serine(S); and

R7 and R8 are lysine(K) or tyrosine(Y), respectively.

Embodiments of the present inventive concepts may also provide use of a peptide including any one amino acid sequence of SEQ ID NOS: 1 to 96 or a composition including the peptide in promoting regeneration of dentin or dental pulp tissues, in preventing or treating dentin hypersensitivity, and in preventing or treating dentin or dental pulp diseases.

Effect of the invention

A peptide for promoting regeneration of dentin or dental pulp tissues and treating dentin hypersensitivity of the present invention exhibits excellent effects of promoting regeneration of dentin or dental pulp tissues, and therefore, it may be widely applied to development of a variety of agents for preventing or treating dentin or dental pulp-related diseases.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A is a graph showing the results of comparing the effects of the peptides of the respective groups of the present invention on expression of DSPP which is an odontoblast differentiation marker gene.

FIG. 1B is a graph showing the results of comparing the expression levels of the odontoblast differentiation marker Dspp gene in MDPC-23 cells treated with the peptides of the present invention.

FIG. 1C is a graph showing the results of comparing expression levels of odontoblast differentiation marker genes, Dspp, Dmp1, and Nestin in MDPC-23 cells treated with peptides of Group 11 and Group 12 of the present invention.

FIG. 1D is a graph showing the results of evaluating cytotoxicity of the peptides of the present invention on dental pulp cells.

FIG. 2 shows microscopic images of dentin/dental pulp-like tissues which were formed *in vivo* for 6 weeks after transplantation of implants including dental pulp cells and various components, in which A to C are microscopic images at 6 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 200 μm , B 100 μm , C 50 μm); D to F are microscopic images at 6 weeks after transplantation of an implant including dental

pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 200 μm , E 100 μm , F 50 μm); G to I are microscopic images at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 200 μm , H 100 μm , I 50 μm); and J to L are microscopic images at 6 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 200 μm , K 100 μm , L 50 μm).

FIG. 3 shows microscopic images of collagen production levels in dentin/dental pulp-like tissues which were formed *in vivo* for 6 weeks after transplantation of implants including dental pulp cells and various components, in which A to C show the results at 6 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 500 μm , B 200 μm , C 50 μm); D to F show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 500 μm , E 200 μm , F 50 μm); G to I show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 500 μm , H 200 μm , I 50 μm); and J to L show the results at 6 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 500 μm , K 200 μm , L 50 μm).

FIG. 4 shows immunostaining images showing the results of evaluating expression levels of DSP which is an odontoblast-specific differentiation marker gene and BSP which is an osteoblast-specific differentiation marker gene in dentin/dental pulp-like tissues which were formed *in vivo* for 6 weeks after transplantation of implants including dental pulp cells and various components, in which A and E show the results at 6 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only; B and F show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11; C and G show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12; and D and H show the results at 6 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2. Arrows of A, B, C and D indicate DSP-expressing regions in the newly formed dentin-like tissues, and arrowheads of E, F, G and H indicate BSP-expressing regions in the newly formed bone-like tissues. Scale bar is 50 μ m.

FIG. 5 shows microscopic images of dentin/dental pulp-like tissues which were formed *in vivo* for 12 weeks after transplantation of implants including dental pulp cells and various components, in which A to C are microscopic images at 12 weeks after transplantation of a control implant

including dental pulp cells and HA/TCP only (scale bar: A 500 μm , B 200 μm , C 50 μm); D to F are microscopic images at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 500 μm , E 200 μm , F 50 μm); G to I are microscopic images at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 500 μm , H 200 μm , I 50 μm); and J to L are microscopic images at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 500 μm , K 200 μm , L 50 μm).

FIG. 6 shows microscopic images of collagen production levels in the dentin/dental pulp-like tissues which were formed *in vivo* for 12 weeks after transplantation of implants including dental pulp cells and various components, in which A to C show the results at 12 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 500 μm , B 200 μm , C 50 μm); D to F show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 500 μm , E 200 μm , F 50 μm); G to I show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 500 μm , H

200 μm , I 50 μm); and J to L show the results at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 500 μm , K 200 μm , L 50 μm).

FIG. 7 shows immunostaining images showing the results of evaluating expression levels of DSP which is an odontoblast-specific differentiation marker gene and BSP which is an osteoblast-specific differentiation marker gene in the dentin/dental pulp-like tissues which were formed *in vivo* for 12 weeks after transplantation of implants including dental pulp cells and various components, in which A and E show the results at 12 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only; B and F show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11; C and G show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12; and D and H show the results at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2. Arrows of A, B, C and D indicate DSP-expressing regions in the newly formed dentin-like tissues, and arrowheads of E, F, G and H indicate BSP-expressing regions in the newly formed bone-like tissues. Scale bar is 50 μm .

FIG. 8 shows scanning microscopic images of the internal structures of the dentin/dental pulp-like tissues which were formed *in vivo* at 12 weeks after transplantation of implants including dental pulp cells and various components, in which A and B are the results at 12 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 50 μ m, B 20 μ m); C and D are the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (SEQ ID NO: 96) (scale bar: C 50 μ m, D 20 μ m); and E and F are the results at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: E 50 μ m, F 20 μ m).

FIG. 9 shows microscopic images and scanning microscopic images showing the results of analyzing odontoblasts/dental pulp-like tissues which were formed after transplantation of implants including the peptides of the present invention into the root canal spaces of the human teeth, in which A and B show the results of staining tissues transplanted with a control implant including dental pulp cells and HA/TCP only (scale bar: A 500 μ m, B 50 μ m); C, D and D' show the results of staining tissues transplanted with an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (SEQ ID NO: 96) (scale bar: A 500 μ m, B

50 μm); D is magnification of box 1 in C, D' is magnification of box 2 in C (scale bar: C 500 μm , D 50 μm , D' 50 μm); E and F show scanning microscopic images of tissues transplanted with a control implant including dental pulp cells and HA/TCP only (scale bar: E 50 μm , F 10 μm); G and H show scanning microscopic images of tissues transplanted with an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (SEQ ID NO: 96) (scale bar: G 50 μm , H 10 μm); P indicates regenerated dental pulp, De indicates existing dentinal wall, DT indicates existing dentinal tubules, Od indicates odontoblasts, OP indicates odontoblast processes, and arrows in D indicate regenerated odontoblast-like cells with odontoblast processes.

FIG. 10 shows microscopic images of histological analysis of the effects of the novel peptides on physiologic dentin (tertiary dentin) formation in a shallow cavity model among indirect pulp capping canine models, in which A to C are microscopic images showing the result of a control group, in which none of the materials was applied to the inlet of dentinal tubule in the exposed dentin (scale bar: A 500 μm , B 100 μm , C 50 μm); D to F are microscopic images showing the result of applying the peptide of Group 11 (1.5 μg) to the inlet of dentinal tubule in the exposed dentin (scale bar: D 500 μm , E 100 μm , F 50 μm); and G to I are microscopic images showing the result of applying the

peptide of Group 12 (1.5 μg) to the inlet of dentinal tubule in the exposed dentin (scale bar: G 500 μm , H 100 μm , I 50 μm). P indicates pulp, D indicates dentin, and TD indicates newly formed physiologic (tertiary) dentin.

FIG. 11 shows microscopic images of histological analysis of the effects of the novel peptides on physiologic dentin (tertiary dentin) formation in a deep cavity model among indirect pulp capping canine models, in which A to C are microscopic images showing the result of a control group, in which the inlet of dentinal tubule in the exposed dentin was treated with none of the materials and filled with a dental restorative material, GI cement (scale bar: A 500 μm , B 100 μm , C 50 μm); D to F are microscopic images showing the result of filling the inlet of dentinal tubule in the exposed dentin with GI cement after being treated with the peptide of Group 11 (1.5 μg) (scale bar: D 500 μm , E 100 μm , F 50 μm); and G to I are microscopic images showing the results of filling the inlet of dentinal tubule in the exposed dentin with GI cement after being treated with the peptide of Group 12 (1.5 μg) (scale bar: G 500 μm , H 100 μm , I 50 μm). P indicates pulp, D indicates dentin, and TD indicates newly formed physiologic (tertiary) dentin.

FIG. 12 shows microscopic images of histological analysis of the effects of the novel peptides on physiologic dentin (tertiary dentin) formation in direct pulp capping

models, in which A to C are microscopic images showing the result of a control group which was treated with none of the materials and then treated with GI cement (scale bar: A 200 μm , B 100 μm , C 50 μm); D to F are microscopic images showing the result of a positive control group which was treated with none of the materials, sealed with MTA, and then covered with GI cement (scale bar: D 500 μm , E 200 μm , F 50 μm); G to I are microscopic images showing the result of the groups which were treated with the peptide of Group 11 (1.5 μg), sealed with MTA, and then covered with GI cement (scale bar: G 200 μm , H 100 μm , I 50 μm); J to L are microscopic images showing the results of the groups which were treated with the peptide of Group 12 (1.5 μg), sealed with MTA, and then covered with GI cement (scale bar: J 500 μm , K 100 μm , L 50 μm). D indicates dentin, OD indicates osteodentin, and TD indicates newly formed physiologic (tertiary) dentin.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present inventors conducted many studies to develop an agent capable of more effectively treating dentin or dental pulp diseases, and as a result, they developed a novel peptide consisting of 10 amino acids.

The newly developed peptide was prepared by substitution of a part of an amino acid sequence of a

peptide which may exhibit a therapeutic effect on dentin or dental pulp diseases, and it was confirmed that the newly developed peptide may increase expression levels of Dspp, Dmp1 and Nestin genes which are odontoblast differentiation marker genes, thereby showing an effect of promoting dentin regeneration without any cytotoxicity against dental pulp cells.

Further, an implant including the peptide together with dental pulp cells was prepared, and the prepared implant was transplanted into a subcutaneous tissue of an immunocompromised mouse, and after 6 weeks to 12 weeks, the transplanted tissue was analyzed. As a result, it was found that a dentin/pulp-like tissue having the most similar morphology to a dentin/dental pulp tissue *in vivo* was formed, a production level of collagen was increased, an expression level of DSP which is an odontoblast-specific differentiation marker was increased, whereas an expression level of BSP which is an osteoblast-specific differentiation marker was not markedly increased.

Furthermore, morphology of the transplanted tissue was examined under a scanning electron microscope, and as a result, odontoblast-like cells exhibited a palisade arrangement on the existing dentinal wall, and their cytoplasmic processes, with lengthened nuclei, extended toward existing dentinal tubules.

Effects of the peptide in indirect and direct pulp capping canine models were examined, and as a result, the same physiologic dentin as observed in the natural human tooth dentin was formed by the novel peptide.

Therefore, it can be seen that the peptide of the present invention may exhibit effects of promoting regeneration of dentin or dental pulp and treating dentin hypersensitivity. The peptide of the present invention having these effects has never been reported so far, and it was first developed by the present inventors.

In an aspect to achieve the above-described objects, the present invention provides a peptide for promoting regeneration of dentin or dental pulp and treating dentin or dental pulp diseases, the peptide including an amino acid sequence of the following Formula 1:

K-Y-R1-R2-R3-R4-R5-R6-R7-R8 (Formula 1)

wherein R1 is arginine(R), lysine(K) or glutamine(Q);

R2 is arginine(R) or glutamine(Q);

R3, R4, and R5 are arginine(R) or lysine(K), respectively;

R6 is asparagine(N) or serine(S); and

R7 and R8 are lysine(K) or tyrosine(Y), respectively.

The term "dentin", as used herein, is also called dentine, and refers to a yellowish white hard tissue that makes up most of a tooth. The dentin is not exposed to the

surface of the tooth, because it is covered by enamel in the tooth crown and cementum in the root. However, dentin exposure may occur at the apical end or the occlusal surface of the tooth crown as the enamel wears with aging. The dentin is a kind of bone tissue, but it is distinguished from a general bone tissue in that the cell bodies of the dentin stay in the dental pulp while their processes extend into the dentinal tubules.

The term "dental pulp tissue", as used herein, is also called "dental pulp", and refers to a soft connective tissue occupying the pulp chamber inside a tooth. Anatomically, the dental pulp has numerous nerves and blood vessels distributed therein, and it extends to the outer surface of the dentin.

The peptide of the present invention is characterized in that it may increase expression levels of Dspp, Dmp1 and Nestin genes which are odontoblast differentiation marker genes without cytotoxicity, and when the peptide is transplanted together with dental pulp cells, the dental pulp cells form a dentin/dental pulp-like tissue.

The peptide of the present invention includes peptide variants thereof having a sequence including one or more amino acid residues different from those of the amino acid sequence of the peptide of the present invention, as long as it may promote regeneration of dentin or dental pulp and

exhibit a therapeutic effect on dentin or dental pulp diseases.

Amino acid exchanges in proteins and polypeptides, which do not generally alter the molecular activity, are known in the art. The most commonly occurring exchanges are amino acid residues Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Thy/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, Asp/Gly, in both directions. The peptide may include peptides that have improved structural stability against heat, pH, etc., or improved ability to promote regeneration of dentin or dental pulp due to alteration or modification of the amino acid sequence.

For example, although glutamine which is an acidic amino acid at position 3 of the peptide of SEQ ID NO: 1 of the present invention is substituted with a basic amino acid, lysine or arginine, the effects of the peptide of the present invention may be obtained as it is; although arginine which is a basic amino acid at position 4 or 5 of the peptide of SEQ ID NO: 1 is substituted with an acidic amino acid glutamine or a basic amino acid lysine, the effects of the peptide of the present invention may be obtained as it is; although lysine which is a basic amino acid at position 6, 7, or 9 of the peptide of SEQ ID NO: 1 is substituted with a basic amino acid arginine or an

aromatic amino acid tyrosine, the effects of the peptide of the present invention may be obtained as it is; although asparagine which is an acidic amino acid at position 8 of the peptide of SEQ ID NO: 1 is substituted with a neutral amino acid serine, the effects of the peptide of the present invention may be obtained as it is; and although tyrosine which is an aromatic amino acid at position 10 of the peptide of SEQ ID NO: 1 is substituted with a basic amino acid lysine, the effects of the peptide of the present invention may be obtained as it is.

As such, although the acidic amino acids, basic amino acids, or aromatic amino acids constituting the peptide of the present invention are substituted with amino acids having the same properties, or substituted with different acidic amino acids, basic amino acids, neutral amino acids, or aromatic amino acids, respectively, the effects of the peptide of the present invention may be obtained as it is. Therefore, it is apparent that a peptide variant having a sequence including one or more amino acid residues different from those of the amino acid sequence constituting the peptide of the present invention is also included in the scope of the peptide of the present invention.

Further, although arbitrary amino acids are added at the N-terminus or C-terminus of the peptide of the present invention, the effects of the peptide of the present

invention may be obtained as it is. Therefore, a peptide prepared by adding arbitrary amino acids at the N-terminus or C-terminus of the peptide of the present invention is also included in the scope of the peptide of the present invention. For example, a peptide prepared by adding 1 to 300 amino acids at the N-terminus or C-terminus of the peptide of the present invention may be exemplified, for another example, a peptide prepared by adding 1 to 100 amino acids at the N-terminus or C-terminus of the peptide of the present invention may be exemplified, and for still another example, a peptide prepared by adding 1 to 24 amino acids at the N-terminus or C-terminus of the peptide of the present invention may be exemplified.

The peptide of the present invention may be chemically modified or protected with an organic group at the N-terminus and/or C-terminus, or may be modified by adding amino acids at the peptide terminus in order to protect the peptide from protease *in vivo* and to increase stability thereof. In particular, since chemically synthesized peptides have charged N-terminus and C-terminus, N-terminal acetylation, N-terminal methylation, or/and C-terminal amidation may be performed, or D-amino acid introduction, peptide bond modification such as $\text{CH}_2\text{-NH}$, $\text{CH}_2\text{-S}$, $\text{CH}_2\text{-S=O}$, $\text{CH}_2\text{-CH}_2$, backbone modification, or side-chain modification may be included in order to remove the charge, but is not

limited thereto. Methods of preparing peptidomimetic compounds are well known in the art, for example, referring to a description in Quantitative Drug Design, C.A. Ramsden Gd., Choplin Pergamon Press (1992).

The term "backbone modification", as used herein, refers to direct modification of amino acids constituting a peptide backbone with amino acid analogs, in which the backbone (main chain) refers to a main chain- or ring-shaped framework of amino acids constituting a peptide. The amino acid analog refers to an amino acid modified by substitution of hydrogen atoms on the nitrogen or α -carbon of the amino acid backbone.

The term "side-chain modification", as used herein, refers to modification of side-chains of amino acids by using a chemical material, in which the side-chains of amino acids refer to atomic groups branched from a main chain- or ring-shaped framework of amino acids constituting a peptide. Examples of the peptide side-chain modification may include amino group modification such as reductive alkylation; amidation with methyl acetimidate; alkylation with acetic anhydride; carbamylation of amino groups with cyanate; trinitrobenzylation of amino acids with 2,4,6-trinitrobenzene sulfonic acid (TNBS); alkylation of amino groups with succinic anhydride; and pyridoxylation with pyridoxal-5-phosphate followed by reduction with NaBH_4 .

Further, the peptide of the present invention may be used alone, or in a combination with various carriers approved as a drug, such as an organic solvent. In order to improve stability and efficacy, the peptide of present invention may be also used by including carbohydrates such as glucose, sucrose, or dextran, antioxidants such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins, other stabilizers, etc.

According to an embodiment of the present invention, 96 kinds of peptides corresponding to Formula 1 of the present invention were synthesized, and effects of the synthesized peptides on an expression level of Dspp gene which is an odontoblast differentiation marker gene were examined. As a result, it was confirmed that all mRNA levels of the odontoblast differentiation marker Dspp gene in MDPC-23 cells which were treated with 96 kinds of the peptides were 1.3 times higher than an mRNA level of the Dspp gene which was measured in MDPC-23 cells (control group) which were treated with none of the peptides of the present invention (Tables 13 to 24).

As reported up to now, it is known that as the mRNA level of DSPP is increased, odontoblast differentiation and dentin regeneration are promoted, and therefore, it can be seen that 96 kinds of the peptides showing the effect of increasing the mRNA level of Dspp gene may exhibit the

effect of promoting odontoblast differentiation and dentin regeneration (Taduru Sreenath et al., THE JOURNAL OF BIOLOGICAL CHEMISTRY, Vol. 278, No. 27, Issue of July 4, pp. 24874-24880, 2003; William T. Butler et al, Connective Tissue Research, 44(Suppl. 1): 171-178, 2003).

In another aspect, the present invention provides a polynucleotide encoding the peptide.

The polynucleotide may be modified by substitution, deletion, or insertion of one or more bases, or a combination thereof. When the nucleotide sequence is prepared by chemical synthesis, a synthetic method widely known in the art, for example, a method described in a literature (Engels and Uhlmann, Angew Chem IntEd Engl., 37:73-127, 1988) may be used, and the nucleotide sequence may be synthesized by triester, phosphite, phosphoramidite and H-phosphate methods, PCR and other autoprimers methods, oligonucleotide synthesis on solid supports, etc. For example, the polynucleotide encoding the peptide of the present invention may include a nucleotide sequence of SEQ ID NO: 4.

In still another aspect, the present invention provides an expression vector including the polynucleotide, a transformant including the expression vector, and a method of preparing the peptide by using the transformant.

The term "expression vector", as used herein, refers to a recombinant vector capable of expressing a target peptide in a host cell, and refers to a genetic construct including essential regulatory elements which are operably linked to express a gene insert. The expression vector includes expression regulatory sequences such as an initiation codon, a stop codon, a promoter, an operator, etc. The initiation and stop codons are generally considered as part of a nucleotide sequence encoding a polypeptide and are necessary to be functional in an individual to whom a genetic construct has been administered, and must be in frame with the coding sequence. The promoter of the vector may be constitutive or inducible.

The term "operably linked", as used herein, refers to a functional linkage between a nucleic acid expression control sequence and a nucleotide sequence encoding a target protein or RNA in such a manner as to allow general functions. For example, a promoter may be operably linked to a nucleotide sequence encoding a protein or RNA to influence expression of the coding sequence. The operable linkage to the expression vector may be prepared by using a genetic recombinant technique well known in the art, and site-specific DNA cleavage and ligation may be carried out by using enzymes generally known in the art.

Further, the expression vector may include signal sequences for discharge of the peptide in order to promote isolation of the peptide from a cell culture. Specific initiation signals may also be required for efficient translation of inserted nucleotide sequences. These signals include ATG initiation codon and adjacent sequences. In some cases, exogenous translational control signals, including ATG initiation codon, should be provided. These exogenous translational control signals and initiation codons may be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by introduction of appropriate transcription or translation enhancer elements.

In addition, the expression vector may further include a protein tag that may be optionally removed by endopeptidase in order to facilitate detection of the peptide.

The term "tag", as used herein, refers to a molecule which exhibits a quantifiable activity or characteristic. The tag may include fluorescent molecules including chemical fluoescers such as fluorescein and polypeptide fluoescers such as green fluorescent protein (GFP) or related proteins; and epitope tags such as a Myc tag, a Flag tag, a His tag, a leucine tag, an IgG tag, a streptavidin tag, etc. In particular, if an epitope tag is used, a peptide tag consisting of preferably 6 or more amino acid residues, and

more preferably, about 8 to 50 amino acid residues may be used.

In the present invention, the expression vector may include a nucleotide sequence encoding the above-described peptide for promoting regeneration of dentin or dental pulp and treating dentin or dental pulp diseases of the present invention. The vector used herein is not specifically limited, as long as it is able to produce the peptide. Preferably, the vector may be plasmid DNA, phage DNA, etc. More preferably, the vector may be a commercially developed plasmid (pUC18, pBAD, pIDTSAMRT-AMP, etc.), an *E. coli*-derived plasmid (pYG601BR322, pBR325, pUC118, pUC119, etc.), a *Bacillus subtilis*-derived plasmid (pUB110, pTP5, etc.), a yeast-derived plasmid (YEp13, YEp24, YCp50, etc.), a phage DNA (Charon4A, Charon21A, EMBL3, EMBL4, λ gt10, λ gt11, λ ZAP, etc.), an animal virus vector (retrovirus, adenovirus, vaccinia virus, etc.), an insect virus (baculovirus, etc.), or the like. For the expression vector, a host cell most suitable for the intended use is preferably selected and used, because the expression level and modification of protein vary depending on the kind of host cell.

The transformant of the present invention may be prepared by transformation of a host with the expression vector of the present invention, and the transformant may express the polynucleotide in the expression vector, thereby

producing the peptide. The transformation may be performed by various methods. The transformation method is not particularly limited, as long as it may produce the peptide. CaCl_2 precipitation, a Hanahan method that is an improved CaCl_2 precipitation method by using DMSO (dimethyl sulfoxide) as a reducing agent, electroporation, calcium phosphate precipitation, protoplast fusion, agitation using silicon carbide fiber, Agrobacterium-mediated transformation, PEG-mediated transformation, dextran sulfate-, lipofectamine- or desiccation/inhibition-mediated transformation, etc. may be used. The host used in the preparation of the transformant is not particularly limited, as long as it may produce the peptide of the present invention. The host may be bacterial cells such as *E. coli*, *Streptomyces*, *Salmonella typhimurium*, etc.; yeast cells such as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, etc.; fungal cells such as *Pichia pastoris*, etc.; insect cells such as Drosophila, Spodoptera Sf9 cells, etc.; animal cells such as CHO, COS, NSO, 293, Bowes melanoma cells, etc.; and plant cells.

The transformant may be used in a method of producing the peptide for promoting regeneration of dentin or dental pulp and treating dentin hypersensitivity of the present invention. Specifically, the method of producing the peptide for promoting regeneration of dentin or dental pulp and treating dentin or dental pulp diseases of the present

invention may include (a) culturing the transformant to obtain a culture; and (b) recovering the peptide of the present invention from the culture.

The term "culturing", as used herein, refers to a method of allowing a microorganism to grow under artificially controlled environmental conditions. In the present invention, the method of culturing the transformant may be performed by a method widely known in the art. Specifically, the culturing is not particularly limited, as long as it may express and produce the peptide for promoting regeneration of dentin or dental pulp and treating dentin or dental pulp diseases of the present invention, and the culturing may be performed by a batch process, a fed-batch process, or a repeated fed batch process.

A medium used in the culturing includes appropriate carbon sources, nitrogen sources, amino acids, vitamins, etc. and should satisfy the requirements of a specific strain in a suitable manner while adjusting temperature, pH, etc. under aerobic conditions. Applicable carbon sources may include, in addition to mixed sugars of glucose and xylose as a main carbon source, sugars and carbohydrates such as sucrose, lactose, fructose, maltose, starch, and cellulose, oils and fats such as soybean oil, sunflower oil, castor oil, coconut oil, etc., fatty acids such as palmitic acid, stearic acid, or linoleic acid, alcohols such as glycerol or

ethanol, and organic acids such as acetic acid. These substances may be used alone or in combination. Applicable nitrogen sources may include inorganic nitrogen sources such as ammonia, ammonium sulfate, ammonium chloride, ammonium acetate, ammonium phosphate, ammonium carbonate, or ammonium nitrate; amino acids such as glutamic acid, methionine, or glutamine; and organic nitrogen sources such as peptone, NZ-amine, meat extract, yeast extract, malt extract, corn steep liquor, casein hydrolysate, fish meal or digested products thereof, defatted soybean cake or digested products thereof, etc. These nitrogen sources may be used alone or in combination. The medium may include, as phosphorus sources, potassium phosphate monobasic, potassium phosphate dibasic, and corresponding sodium-containing salts. Applicable phosphorus sources may include potassium dihydrogen phosphate, dipotassium hydrogen phosphate, or corresponding sodium-containing salts. In addition, inorganic compounds may include sodium chloride, calcium chloride, iron chloride, magnesium sulfate, iron sulfate, manganese sulfate and calcium carbonate. In addition to the above materials, essential growth materials such as amino acids and vitamins may be used.

Further, appropriate precursors may be used in the culture medium. During culturing, the above-described materials may be appropriately added to the culture in a

batch, fed-batch, or continuous manner, but are not limited thereto. The pH of the culture may be adjusted by appropriately using a basic compound such as sodium hydroxide, potassium hydroxide, or ammonia, or an acidic compound such as phosphoric acid or sulfuric acid.

In addition, formation of bubbles may be inhibited by using an antifoaming agent such as fatty acid polyglycol ester. In order to maintain an aerobic state, oxygen or oxygen-containing gas (e.g., air) may be injected into the culture. The temperature of the culture is generally 27°C to 37°C, preferably 30°C to 35°C. Culturing is continued until the desired level of the peptide production will be obtained. This is achieved within 10 hours to 100 hours.

In addition, the recovering of the peptide from the culture may be performed by a method known in the art. Specifically, the recovering method is not particularly limited, as long as it may recover the produced peptide. Preferably, a method such as centrifugation, filtration, extraction, spraying, drying, evaporation, precipitation, crystallization, electrophoresis, fractional dissolution (e.g., ammonium sulfate precipitation), chromatography (e.g., ion exchange, affinity, hydrophobic, and size exclusion), etc. may be used.

In still another aspect, the present invention provides a pharmaceutical composition for preventing or treating dentin or dental pulp disease including the peptide.

As described above, when the peptide for promoting regeneration of dentin or dental pulp and treating dentin hypersensitivity of the present invention is transplanted into the body, together with dental pulp cells, formation of dentin/dental pulp-like tissue by the dental pulp cells may be promoted, and when the peptide is applied to the damaged dentin or dental pulp site, the same physiologic dentin as observed in the natural human tooth dentin may be formed. Therefore, the peptide may be used as an active ingredient of the pharmaceutical composition for treating dentin or dental pulp diseases which are caused by damage to dentin or dental pulp.

The peptide included in the pharmaceutical composition may be used in a single form of the peptide or in a polypeptide form of 2 or more repeats of the peptide, and the peptide may be also used in a complex form of a drug having a therapeutic effect on dentin or dental pulp diseases linked at the N-terminus or C-terminus of the peptide.

The term "dentin or dental pulp diseases", as used herein, refer to all diseases caused by damaged dental pulp

tissue and dentin linked to the dental pulp, due to damage to the dentin and dental pulp tissues.

In the present invention, the dentin or dental pulp diseases are not particularly limited, as long as the peptide of the present invention exhibits the therapeutic effects on the diseases, and the dentin or dental pulp diseases may include, for example, dentin hypersensitivity, pulp hyperemia, pulpitis, pulp degeneration, pulp necrosis, gangrenous pulp, etc.

The term "preventing", as used herein, means all actions by which the occurrence of dentin or dental pulp diseases is restrained or retarded by administration of the pharmaceutical composition for preventing or treating dentin or dental pulp diseases including the peptide of the present invention.

The term "treating", as used herein, means all actions by which dentin or dental pulp diseases are treated by promoting regeneration of dentin or dental pulp by administering the pharmaceutical composition including the peptide of the present invention as an active ingredient to a subject in need of treatment of dentin or dental pulp diseases.

The pharmaceutical composition of the present invention may be prepared in the form of a pharmaceutical composition for treating dentin or dental pulp diseases further

including, in addition to the peptide, an appropriate carrier (natural or non-natural carrier), excipient, or diluent commonly used in the preparation of pharmaceutical compositions. Specifically, the pharmaceutical composition may be formulated according to a common method in the form of a sterile injectable solution which may be administered to dentin or dental pulp disease-induced site. In the present invention, the carrier, excipient, and diluent which may be included in the pharmaceutical composition may include lactose, dextrose, sucrose, sorbitol, mannitol, xylitol, erythritol, maltitol, starch, acacia rubber, alginate, gelatin, calcium phosphate, calcium silicate, cellulose, methylcellulose, microcrystalline cellulose, polyvinylpyrrolidone, water, methylhydroxybenzoate, propylhydroxybenzoate, talc, magnesium stearate, mineral oils, collagen, etc. Upon formulation, commonly used diluents or excipients such as a filler, an extender, a binder, a wetting agent, a disintegrant, a surfactant, etc. may be used. In particular, a sterilized aqueous solution, a non-aqueous solvent, a suspension, an emulsion, a freeze-dried preparation, a suppository, an ointment (e.g., pulp liner, etc.) may be included. As non-aqueous solvents or suspensions, propylene glycol, polyethylene glycol, plant oils such as olive oil, injectable esters such as ethyl oleate, etc. may be used. As a base of the suppositories,

witepsol, Macrogol, Tween 61, cacao butter, laurin fat, glycerogelatin, etc. may be used.

A content of the peptide in the pharmaceutical composition of the present invention is not particularly limited, but the peptide may be included in an amount of 0.0001% by weight to 50% by weight, more preferably, 0.01% by weight to 20% by weight, based on the total weight of the final composition.

The pharmaceutical composition of the present invention may be administered in a pharmaceutically effective amount. The term "pharmaceutically effective amount", as used herein, refers to an amount sufficient to treat or prevent diseases, at a reasonable benefit/risk ratio applicable to any medical treatment or prevention. An effective dosage level may be determined depending on factors including severity of the disease, drug activity, a patient's age, body weight, health conditions, sex, sensitivity to the drug, administration time, administration route, and excretion rate of the composition of the present invention, duration of treatment, drugs blended with or co-administered with the composition of the present invention, and other factors known in the medical field. The pharmaceutical composition of the present invention may be administered individually or in combination with a known pharmaceutical composition for treating dentin or pulp diseases. It is important to administer the

composition in a minimum amount that may exhibit a maximum effect without causing side effects, in view of all the above-described factors.

An administration dose of the pharmaceutical composition of the present invention may be determined by those skilled in the art, in view of purpose of use, severity of the disease, a patient's age, body weight, sex, and medical history, a kind of a material used as an active ingredient, etc. For example, the pharmaceutical composition of the present invention may be administered at a dose of about 0.1 ng/kg to about 100 mg/kg, and preferably, about 1 ng/kg to about 10 mg/kg per adult, and administration frequency of the composition of the present invention is not particularly limited, but the composition may be administered once a day or several times a day in divided doses. The administration dose does not limit the scope of the present invention in any aspect.

In still another aspect, the present invention provides a method of treating dentin or dental pulp diseases, the method including administering the pharmaceutically effective amount of the pharmaceutical composition to a subject having dentin or dental pulp diseases, excluding humans.

The term "subject, as used herein, may include mammals including rats, livestock, etc. in need of treatment of

dentin or dental pulp diseases without limitation, and humans are excluded from the subjects having the above diseases.

The pharmaceutical composition for treating dentin or dental pulp diseases of the present invention may be administered via any general route, as long as the pharmaceutical composition is able to reach a target tissue. The pharmaceutical composition may be administered, but is not particularly limited to, via intraoral administration, intraoral injection, etc., depending on the purpose.

In still another aspect, the present invention provides a quasi-drug composition for preventing or improving dentin or dental pulp diseases including the peptide.

The term "improving", as used herein, means all actions that at least reduce a parameter related to the conditions to be treated, for example, the degree of symptom.

In the present invention, the improving is to be interpreted as all actions by which symptoms of dentin or dental pulp diseases have taken a turn for the better or been modified favorably by promoting regeneration of dentin or dental pulp by administering the pharmaceutical composition including the peptide of the present invention as an active ingredient to a subject in need of treatment of dentin or dental pulp diseases.

The term "quasi-drug", as used herein, refers to an article having a milder action than drugs, among articles being used for the purpose of diagnosis, treatment, improvement, alleviation, handling, or prevention of human or animal diseases. For example, according to Pharmaceutical Affairs Law, the quasi-drugs are those, excluding articles used as drugs, including articles made from fiber or rubber which are used for the purpose of treating or preventing human or animal diseases, articles, other than a tool or a machine, or an analogue thereof, which have a mild action on or have no direct influence on the human body, and articles which are used for the purpose of disinfection or pest control for the prevention of infectious diseases.

In the present invention, a kind or formulation of the quasi-drug composition including the peptide is not particularly limited, but the quasi-drug composition may be, for example, oral antiseptic mouthwashes, oral hygiene products, toothpastes, floss, oral ointments, etc.

In still another aspect, the present invention provides a health functional food composition for preventing or improving dentin or dental pulp diseases including the peptide.

The term "food", as used herein, includes meats, sausages, breads, chocolates, candies, snacks, confectionery, pizzas, ramen noodles, other noodles, gums, dairy products

including ice-creams, various soups, beverages, teas, drinks, alcoholic beverages, and vitamin complexes, health functional foods, health foods, etc., and the food includes all foods in the ordinary acceptance of the term.

The term "functional food", as used herein, is the term identical to the food for special health use (FoSHU), and refers to a food having high medical, medicinal effects, which is processed so as to efficiently exhibit the biologically modulating function as well as to supply nutrients. Here, the term "functional" indicates a useful effect for human health, such as regulation of nutrients for the structure and function of the human body, physiological action, etc. The food of the present invention may be prepared according to a method commonly employed in the art, and raw materials and ingredients commonly used in the art may be added upon preparing the food. In addition, a formulation of the food is not limited, as long as the formulation is accepted as a food. The food composition of the present invention may be prepared as a variety of formulations. Since the food is used as raw materials, unlike general drugs, the food composition lacks side effects which may occur when a drug is taken for a long period of time, and may have excellent portability. Therefore, the food of the present invention may be taken as

a supplement for enhancing the effects of preventing or improving dentin or dental pulp diseases.

The health food means a food having effects of actively maintaining or promoting health conditions, as compared with general foods, and the health supplement food means a food for supplementing health. If necessary, the health functional food, health food, and health supplement food may be interchangeably used.

Specifically, the health functional food is a food prepared by adding the peptide of the present invention to food materials such as beverages, teas, spices, gums, confectionery, etc., or prepared as a capsule, a powder, a suspension, etc. The health functional food means that it takes a certain effect on health when consumed, but unlike general drugs, the health functional food has an advantage of having no side effects that may occur when a drug is taken for a long time, because it uses a food as a raw material.

Since the food composition of the present invention is routinely ingested, the food composition is expected to show high efficacy on prevention or improvement of dentin or dental pulp diseases, and thus it may be very usefully applied.

The food composition may further include a physiologically acceptable carrier. A kind of the carrier is

not particularly limited. Any carrier may be used, as long as it is commonly used in the art.

Further, the food composition may further include additional ingredients that are commonly used in food compositions so as to improve smell, taste, vision, etc. For example, the food composition may include vitamins A, C, D, E, B1, B2, B6, B12, niacin, biotin, folate, pantothenic acid, etc. Additionally, the food composition may also include minerals such as Zn, Fe, Ca, Cr, Mg, Mn, Cu, etc. Additionally, the food composition may also include amino acids such as lysine, tryptophan, cysteine, valine, etc. Additionally, the food composition may also include food additives, such as preservatives (potassium sorbate, sodium benzoate, salicylic acid, sodium dehydroacetate, etc.), disinfectants (bleaching powder, higher bleaching powder, sodium hypochlorite, etc.), antioxidants (butylhydroxyanisole (BHA), butylhydroxytoluene (BHT), etc.), coloring agents (tar color, etc.), color-developing agents (sodium nitrite, etc.), bleaching agents (sodium sulfite), seasonings (monosodium glutamate (MSG), etc.), sweeteners (dulcin, cyclamate, saccharin, sodium, etc.), flavors (vanillin, lactones, etc.), swelling agents (alum, potassium D-bitartrate, etc.), fortifiers, emulsifiers, thickeners (adhesive pastes), film-forming agents, gum base agents, antifoaming agents, solvents, improvers, etc. The additives

may be selected and used in an appropriate amount according to the food types.

The peptide of the present invention may be added as it is, or may be used in conjunction with other foods or food ingredients according to a common method, or may be used appropriately according to a common method. Mixing amounts of the active ingredient may be suitably determined depending upon the purpose of use (prophylactic, health or therapeutic treatment). Generally, upon production of a food or a beverage, the food composition of the present invention may be added in an amount of 50 parts by weight or less, specifically 20 parts by weight or less, based on the total weight of the food or the beverage. However, when prolonged intake is intended for the purpose of health and hygiene, the food composition may be included in an amount below the above range. In addition, since there is no safety problem, the active ingredient may be used in an amount above the above range.

The food composition of the present invention may be used as, for example, a health beverage composition, and in this case, the health beverage composition may further include various flavors or natural carbohydrates, as in common beverages. The natural carbohydrates may include monosaccharides such as glucose and fructose; disaccharides such as maltose and sucrose; polysaccharides such as dextrin

and cyclodextrin; and sugar alcohols such as xylitol, sorbitol and erythritol. The sweeteners may be natural sweeteners such as thaumatin or a stevia extract; or synthetic sweeteners such as saccharin or aspartame. The natural carbohydrate may be generally used in an amount of about 0.01 g to 0.04 g, and specifically, about 0.02 g to 0.03 g, based on 100 mL of the health beverage composition of the present invention

In addition, the health beverage composition may include various nutrients, vitamins, minerals, flavors, colorants, pectic acid and salts thereof, alginic acid and salts thereof, organic acids, protective colloidal thickeners, pH modifiers, stabilizers, antiseptics, glycerin, alcohols, carbonating agents, etc. Moreover, the health beverage composition may include the fruit flesh used to prepare natural fruit juices, fruit juice beverages, or vegetable beverages. These ingredients may be used individually or in combination. A proportion of the additives is not critical, but is generally selected from 0.01 parts by weight to 0.1 parts by weight per 100 parts by weight of the health beverage composition of the present invention.

The food composition of the present invention may include the peptide of the present invention in a variety of % by weight, as long as it may exhibit the effect of

preventing or improving dentin or dental pulp diseases. Specifically, the peptide of the present invention may be included in an amount of 0.00001% by weight to 100% by weight or 0.01% by weight to 80% by weight, based on the total weight of the food composition, but is not limited thereto.

In still another aspect, the present invention provides a method of preventing or treating dentin or dental pulp diseases, the method including administering the composition including the peptide to a subject.

In still another aspect, the present invention provides a method of promoting regeneration of dentin or dental pulp tissues, the method including administering the composition including the peptide to a subject.

In still another aspect, the present invention provides use of a peptide including an amino acid sequence of the following Formula 1 or a composition including the peptide in promoting regeneration of dentin or dental pulp tissues, in preventing or treating dentin hypersensitivity, and in preventing or treating dentin or dental pulp diseases:

K-Y-R1-R2-R3-R4-R5-R6-R7-R8 (Formula 1)

wherein R1 is arginine(R), lysine(K) or glutamine(Q);

R2 is arginine(R) or glutamine(Q);

R3, R4, and R5 are arginine(R) or lysine(K), respectively;

R6 is asparagine(N) or serine(S); and

R7 and R8 are lysine(K) or tyrosine(Y), respectively.

In still another aspect, the present invention provides use of a peptide including any one amino acid sequence of SEQ ID NOS: 1 to 96 or a composition including the peptide in promoting regeneration of dentin or dental pulp tissues, in preventing or treating dentin hypersensitivity, and in preventing or treating dentin or dental pulp diseases.

Hereinafter, the present invention will be described in more detail with reference to Examples. However, these Examples are for illustrative purposes only, and the scope of the present invention is not intended to be limited by these Examples.

Example 1: Synthesis of peptides for promoting generation of dentin or dental pulp tissues and treating dentin hypersensitivity

The present inventors synthesized a peptide (SEQ ID NO: 1) showing the effect of promoting regeneration of dentin or dental pulp tissues by a 9-fluorenylmethyloxycarbonyl (Fmoc) method, and they synthesized peptides of respective groups (Tables 1 to 12) by substituting the amino acids of the synthesized peptide.

N-KYQRRKKNKY-C (SEQ ID NO: 1)

First, peptides of Group 1 were synthesized by using the peptide of SEQ ID NO: 1 or by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine (Table 1).

[Table 1]

Peptides of Group 1

SEQ ID NO:	Amino acid sequence(N-C)
1	KYQRRKKNKY
2	KYQRRKRNKY
3	KYQRRRKKNKY
4	KYQRRRRNKY
5	KYQRKKKKNKY
6	KYQRKRKNKY
7	KYQRKKRNKY
8	KYQRKRRNKY

Next, peptides of Group 2 were synthesized by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine or by substituting an amino acid at position 8 of the peptide of SEQ ID NO: 1 with serine (Table 2).

[Table 2]

Peptides of Group 2

SEQ ID NO:	Amino acid sequence(N-C)
9	KYQRRKSKY

10	KYQRRKRSKY
11	KYQRRRKSKY
12	KYQRRRRSKY
13	KYQRKKKSKY
14	KYQRKRKSKY
15	KYQRKKRSKY
16	KYQRKRRSKY

Next, peptides of Group 3 were synthesized by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine or by substituting an amino acid at position 9 of the peptide of SEQ ID NO: 1 with tyrosine (Table 3).

[Table 3]

Peptides of Group 3

SEQ ID NO:	Amino acid sequence(N-C)
17	KYQRRKKNYK
18	KYQRRKRNYK
19	KYQRRRKNYK
20	KYQRRRRNYK
21	KYQRKKKNYK
22	KYQRKRKNYK
23	KYQRKKRNYK
24	KYQRKRRNYK

Next, peptides of Group 4 were synthesized by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine, by substituting an amino acid at position 8 of the peptide of SEQ ID NO: 1 with serine, by substituting an amino acid at position 9 of the peptide of SEQ ID NO: 1 with tyrosine, or by substituting an amino acid at position 10 of the peptide of SEQ ID NO: 1 with lysine (Table 4).

[Table 4]

Peptides of Group 4

SEQ ID NO:	Amino acid sequence(N-C)
25	KYQRRKKS ^Y YK
26	KYQRRKR ^S SYK
27	KYQRRRRK ^S SYK
28	KYQRRRRR ^S SYK
29	KYQRKKK ^S SYK
30	KYQRKRK ^S SYK
31	KYQRKKR ^S SYK
32	KYQRKR ^R SYK

Next, peptides of Group 5 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with arginine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, or by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine (Table 5).

[Table 5]

Peptides of Group 5

SEQ ID NO:	Amino acid sequence(N-C)
33	KYRQRKKNKY
34	KYRQRKRKNKY
35	KYRQRRKNKY
36	KYRQRRRNKY
37	KYRQKKKNKY
38	KYRQKRKNKY
39	KYRQKKRNKY
40	KYRQKRRNKY

Next, peptides of Group 6 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with arginine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine, or by substituting an amino acid at position 8 of the peptide of SEQ ID NO: 1 with serine (Table 6).

[Table 6]

Peptides of Group 6

SEQ ID NO:	Amino acid sequence(N-C)
41	KYRQRKKSKY
42	KYRQRKRISKY

43	KYRQRRKSKY
44	KYRQRRRSKY
45	KYRQKKKSKY
46	KYRQKRKSKY
47	KYRQKKRSKY
48	KYRQKRRSKY

Next, peptides of Group 7 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with arginine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine, by substituting an amino acid at position 9 of the peptide of SEQ ID NO: 1 with tyrosine, or by substituting an amino acid at position 10 of the peptide of SEQ ID NO: 1 with lysine (Table 7).

[Table 7]

Peptides of Group 7

SEQ ID NO:	Amino acid sequence(N-C)
49	KYRQRKKNYK
50	KYRQRKRNYK
51	KYRQRRKNYK
52	KYRQRRRNYK
53	KYRQKKKNYK

54	KYRQKRKNYK
55	KYRQKKRNYK
56	KYRQKRRNYK

Next, peptides of Group 8 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with arginine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine, by substituting an amino acid at position 8 of the peptide of SEQ ID NO: 1 with serine, by substituting an amino acid at position 9 of the peptide of SEQ ID NO: 1 with tyrosine, or by substituting an amino acid at position 10 of the peptide of SEQ ID NO: 1 with lysine (Table 8).

[Table 8]

Peptides of Group 8

SEQ ID NO:	Amino acid sequence(N-C)
57	KYRQRKKSYK
58	KYRQRKRSYK
59	KYRQRRKSYK
60	KYRQRRRSYK
61	KYRQKKKSYK
62	KYRQKRKSYK
63	KYRQKKRSYK

64	KYRQKRRSYK
----	------------

Next, peptides of Group 9 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with lysine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, or by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine (Table 9).

[Table 9]

Peptides of Group 9

SEQ ID NO:	Amino acid sequence(N-C)
65	KYKQRKKNKY
66	KYKQRKRKNKY
67	KYKQRRKNKY
68	KYKQRRRNKY
69	KYKQKKKNKY
70	KYKQKRKNKY
71	KYKQKKRNKY
72	KYKQKRRNKY

Next, peptides of Group 10 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with lysine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine, or by

substituting an amino acid at position 8 of the peptide of SEQ ID NO: 1 with serine (Table 10).

[Table 10]

Peptides of Group 10

SEQ ID NO:	Amino acid sequence(N-C)
73	KYKQRKKSKY
74	KYKQRKRSKY
75	KYKQRRKSKY
76	KYKQRRRSKY
77	KYKQKKKSKY
78	KYKQKRKSKY
79	KYKQKKRSKY
80	KYKQKRRSKY

Next, peptides of Group 11 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with lysine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine, by substituting an amino acid at position 9 of the peptide of SEQ ID NO: 1 with tyrosine, or by substituting an amino acid at position 10 of the peptide of SEQ ID NO: 1 with lysine (Table 11).

[Table 11]

Peptides of Group 11

SEQ ID NO:	Amino acid sequence(N-C)
81	KYKQRKKNYK
82	KYKQRKRNYK
83	KYKQRRKNYK
84	KYKQRRRNYK
85	KYKQKKKNYK
86	KYKQKRKNYK
87	KYKQKKRNYK
88	KYKQKRRNYK

Lastly, peptides of Group 12 were synthesized by substituting an amino acid at position 3 of the peptide of SEQ ID NO: 1 with lysine, by substituting an amino acid at position 4 of the peptide of SEQ ID NO: 1 with glutamine, by substituting any amino acid at positions 5 to 7 of the peptide of SEQ ID NO: 1 with lysine or arginine, by substituting an amino acid at position 8 of the peptide of SEQ ID NO: 1 with serine, by substituting an amino acid at position 9 of the peptide of SEQ ID NO: 1 with tyrosine, or by substituting an amino acid at position 10 of the peptide of SEQ ID NO: 1 with lysine (Table 12).

[Table 12]

Peptides of group 12

SEQ ID NO:	Amino acid sequence(N-C)
89	KYKQRKKSYK

90	KYKQRKRSYK
91	KYKQRRKSYK
92	KYKQRRRSYK
93	KYKQKKKSYK
94	KYKQKRKSYK
95	KYKQKKRSYK
96	KYKQKRRSYK

Example 2: Investigation of efficacy of peptides for promoting regeneration of dentin or dental pulp tissues and treating dentin hypersensitivity by using odontoblasts

Example 2-1: Effects of peptides for promoting regeneration of dentin or dental pulp tissues and treating dentin hypersensitivity on DSPP (dentin sialophosphoprotein) promoter activity

First, MDPC-23 cells which are mouse-derived odontoblasts were cultured in DMED medium supplemented with 10% FBS under conditions of 5% CO₂ and 37°C.

Next, the cultured MDPC-23 cells were seeded on a 24-well plate at a density of 5 X 10⁴ cells per well, and cultured for 24 hours. Then, the cultured cells were transfected with a recombinant vector constructed by introducing a DSPP promoter and a luciferase gene into a pGL3 vector, by using Lipofectamine Plus™ reagent. The transfected MDPC-23 cells were treated with the peptides of

Groups 1 to 12 synthesized in Example 1, respectively and cultured for 48 hours. Luciferase activity in each of the transfected MDPC-23 cells was measured, and the calculated mean levels of the respective groups were compared with each other (FIG. 1A). In this regard, transfected MDPC-23 cells which were treated with none of the peptides of the present invention were used as a control group.

FIG. 1A is a graph showing the results of comparing the effects of the peptides of the respective groups of the present invention on expression of DSPP which is an odontoblast differentiation marker gene. As shown in FIG. 1A, overall luciferase activity levels of the respective peptides of the present invention were about 1.3 times higher than that of the control group, but there were differences between groups. It was confirmed that the peptides of Group 12 showed the highest level of luciferase activity, and the peptides of Group 11 showed the next highest level of luciferase activity.

Therefore, it can be seen that the peptides of the present invention exhibit the effect of activating the Dspp promoter.

Example 2-2: Effects of peptides for promoting regeneration of dentin or dental pulp tissues and treating dentin or dental pulp diseases on expression level of odontoblast differentiation marker DSPP gene

The MDPC-23 cells cultured in Example 2-1 were treated with the peptides of the respective groups which were synthesized in Example 1, and cultured for 48 hours. Then, mRNA levels of an odontoblast differentiation marker Dspp gene expressed in the MDPC-23 cells were measured, and a ratio of the measured Dspp mRNA level relative to a Dspp mRNA level measured in a control group was calculated, respectively (Tables 13 to 24). Further, mean values of the Dspp mRNA levels measured in the peptides of respective groups were compared between the groups (FIG. 1B). In this regard, MDPC-23 cells which were treated with none of the peptides of the present invention were used as a control group.

The expression levels of the Dspp gene were measured by RT-PCR and real time PCR analysis: In detail, total RNA was extracted from the MDPC-23 cells with TRIzol reagent. 2 μ g of the total RNA, 1 μ l of reverse transcriptase, and 0.5 μ g of oligo (dT) were used to synthesize cDNA. The synthesized cDNA were used in real-time polymerase chain reaction. The real-time polymerase chain reaction was performed on an ABI PRISM 7500 sequence detection system (Applied Biosystems) using the following primers and a SYBR GREEN PCR Master Mix (Takara, Japan). The real-time polymerase chain reaction was performed under conditions of 94°C, 1 min; 95°C, 15 sec - 60°C, 34 sec for 40 cycles. Results were analyzed by a

comparative cycle threshold (CT) method. In this regard, Gapdh gene was used as an internal control. The experiments were performed in triplicate, and then mean values and standard deviations thereof were taken as measured values.

Dspp_R: 5'-CTGTTGCTAGTGGTGCTGTT-3'(SEQ ID NO: 97)

Dmp1_F: 5'-CATTCTCCTTGTGTTCTTTGGG-3'(SEQ ID NO: 98)

Gapdh_F: 5'-AGGTCGGTGTGAACGGATTTG-3'(SEQ ID NO: 99)

Gapdh_R: 5'-TGTAGACCATGTAGTTGAGGTCA-3'(SEQ ID NO: 100)

[Table 13]

Effects of peptides of Group 1 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
1	1.754	0.132
2	1.646	0.092
3	1.464	0.221
4	1.855	0.102
5	1.639	0.057
6	1.746	0.091
7	1.864	0.132
8	1.639	0.032

[Table 14]

Effects of peptides of Group 2 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
9	1.854	0.032
10	1.746	0.052
11	1.639	0.201
12	1.548	0.027
13	1.685	0.077
14	1.846	0.141
15	1.964	0.279
16	1.739	0.092

[Table 15]

Effects of peptides of Group 3 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
17	2.117	0.209
18	2.319	0.092
19	1.931	0.102
20	2.553	0.099
21	1.893	0.132
22	2.412	0.072
23	2.171	0.281
24	2.212	0.111

[Table 16]

Effects of peptides of Group 4 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
25	2.371	0.089
26	2.193	0.052
27	1.993	0.202
28	2.453	0.192
29	1.883	0.101
30	2.512	0.209
31	2.371	0.139
32	2.219	0.302

[Table 17]

Effects of peptides of Group 5 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
33	1.712	0.091
34	1.931	0.172
35	1.983	0.102
36	2.319	0.292
37	1.597	0.301

38	2.116	0.211
39	1.712	0.191
40	2.219	0.212

[Table 18]

Effects of peptides of Group 6 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
41	1.546	0.091
42	1.586	0.103
43	1.669	0.095
44	1.793	0.203
45	1.532	0.310
46	1.887	0.077
47	1.697	0.009
48	1.558	0.201

[Table 19]

Effects of peptides of Group 7 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
49	1.923	0.192

50	1.887	0.007
51	1.601	0.082
52	2.019	0.135
53	1.592	0.222
54	1.437	0.341
55	1.663	0.094
56	1.701	0.109

[Table 20]

Effects of peptides of Group 8 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
57	2.039	0.082
58	1.998	0.172
59	1.792	0.007
60	2.107	0.201
61	2.301	0.019
62	1.672	0.308
63	1.769	0.085
64	1.967	0.039

[Table 21]

Effects of peptides of Group 9 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
65	1.723	0.072
66	1.627	0.291
67	1.777	0.027
68	1.432	0.410
69	2.011	0.081
70	1.927	0.105
71	1.879	0.060
72	2.011	0.009

[Table 22]

Effects of peptides of Group 10 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
73	2.035	0.021
74	2.011	0.063
75	1.997	0.059
76	2.351	0.109
77	1.729	0.111
78	2.635	0.091
79	2.231	0.077
80	1.837	0.201

[Table 23]

Effects of peptides of Group 11 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
81	3.092	0.152
82	3.361	0.098
83	3.572	0.209
84	3.702	0.301
85	3.670	0.088
86	3.705	0.137
87	3.888	0.072
88	4.021	0.301

[Table 24]

Effects of peptides of Group 12 on mRNA level of Dspp gene

SEQ ID NO:	mRNA level of Dspp gene	
	Mean	Standard deviation
89	4.211	0.413
90	4.811	0.302
91	4.362	0.182
92	4.211	0.287
93	4.525	0.250

94	3.836	0.099
95	4.620	0.401
96	5.211	0.371

As shown in Tables 13 to 24, when the peptides of the present invention were treated, overall mRNA levels of the Dspp gene which is an odontoblast differentiation marker were 1.3 times higher than the mRNA level of Dspp gene measured in MDPC-23 cells (control group) treated with none of the peptides of the present invention.

In particular, all peptides of Group 11 showed 3 times higher Dspp mRNA levels and all peptides of Group 12 showed 3.8 times higher Dspp mRNA levels, as compared with the control group.

Moreover, FIG. 1B is a graph showing the results of comparing the expression levels of the odontoblast differentiation marker Dspp gene in the MDPC-23 cells treated with the peptides of the present invention. As shown in FIG. 1B, when the peptides of the present invention were treated, the mRNA levels of the odontoblast differentiation marker Dspp gene were increased, and similar to those of FIG. 1A, the mRNA levels were about 1.3 times higher than the Dspp mRNA level measured in the control group.

Example 2-3: Effects of peptides for promoting regeneration of dentin or dental pulp tissues and treating

dentin or dental pulp diseases on expression levels of odontoblast differentiation marker genes, Dspp, Dmp1 and Nestin

The results of Example 2-2 showed that the peptides of the present invention may increase the Dspp mRNA level, and in particular, the peptides of Group 11 and Group 12 may increase the Dspp mRNA level at least three times or higher.

Accordingly, it was examined whether the peptides of Group 11 and Group 12 may increase mRNA levels of other odontoblast differentiation marker genes, Dmp1 and Nestin.

Briefly, experiments were performed in the same manner as in Example 2-2, except that the following primers were used and the peptides of Group 11 and Group 12 were used as peptides. The effects of the peptides of the present invention on expression levels of Dmp1 and Nestin genes were measured, and the calculated mean values were compared between the groups (FIG. 1C). In this regard, MDPC-23 cells which were treated with none of the peptides of the present invention were used as a control group, and the mRNA level of the Dspp gene was used as a comparative group.

Dmp1_F: 5'-CATTCTCCTTGTGTTCTTTGGG-3'(SEQ ID NO: 101)

Dmp1_R: 5'-TGTGGTCACTATTTGCCTGTG-3'(SEQ ID NO: 102)

Nestin_F: 5'-CCCTGAAGTCGAGGAGCTG-3'(SEQ ID NO: 103)

Nestin_R: 5'-CTGCTGCACCTCTAAGCGA-3'(SEQ ID NO: 104)

FIG. 1C is a graph showing the results of comparing the expression levels of the odontoblast differentiation marker genes, Dspp, Dmp1 and Nestin in the MDPC-23 cells treated with the peptides of Group 11 and Group 12 of the present invention. As shown in FIG. 1C, when the peptides of the present invention were treated, all mRNA levels of the odontoblast differentiation marker genes, Dspp, Dmp1 and Nestin were increased, and there were differences in the increased levels between the genes. The expression levels increased by the peptides of Group 12 were higher than the expression levels increased by the peptides of Group 11.

The above differentiation marker genes are known as genes involved in odontoblast differentiation and dentin mineralization, which infers that the peptides of the present invention may exhibit the effect of promoting dentin regeneration.

Example 2-4: Test of cytotoxicity of peptides for promoting regeneration of dentin or dental pulp tissues and treating dentin or dental pulp diseases in dental pulp cells

First, human dental pulp stem cells were separated from wisdom teeth of 10 adults (aged 18-22) at School of Dentistry, Seoul National University. In detail, all experiments were performed after the approval of Institutional Review Board and the informed consent from patients. Wisdom teeth were fractured according to a method

of Jung HS et al. (J Mol Histol.(2011)) to expose the dental pulps, and dental pulp tissues were separated with forceps. Each of the separated dental pulp tissues was cut into small pieces with a razor blade, put in a 60-mm dish, covered with a cover slip, and then cultured in a DMEM medium to obtain cultured dental pulp cells.

Next, the obtained dental pulp cells were seeded on a 96-well plate at a density of 3×10^3 cells per well, and cultured for 24 hours. Then, the cells were treated with the peptide of Group 11 or 12 at a concentration of 10 µg/ml or 50 µg/ml, and further cultured for 1 day, 3 days, or 5 days. After completion of the culture, the cultured cells were washed with PBS, 20 µl of MTT solution was added thereto, and then allowed to react at 37°C for 4 hours. After completion of the reaction, the MTT solution was removed, and 100 µl of DMSO was added thereto, and absorbance at 540 nm was measured (FIG. 1D). In this regard, dental pulp cells which were cultured without treatment of the peptides were used as a control group.

FIG. 1D is a graph showing the results of evaluating cytotoxicity of the peptides of the present invention on dental pulp cells. As shown in FIG. 1D, although the peptides of the present invention were treated, the dental pulp cells showed the same level of viability as the control group.

Example 3: Effects of peptides for promoting regeneration of dentin or dental pulp tissues and treating dentin or dental pulp diseases on formation of dentin/dental pulp-like tissues *in vivo*

Example 3-1: Morphological analysis of transplanted tissues derived from animals raised for 6 weeks

100 mg of a dentin substitute was added to 2×10^6 of dental pulp cells, and mixed with 0.5% fibrin gel to prepare an implant. In this regard, hydroxy apatite/tricalcium phosphate (HA/TCP) ceramic powder (Zimmer, USA) was used as the dentin substitute, and the fibrin gel was prepared by including 10 μ g of the peptide of Group 11 (SEQ ID NO: 87), the peptide of Group 12 (SEQ ID NO: 96), or 2 μ g of BMP-2. The prepared implant was transplanted into the subcutaneous tissues of immunocompromised mice (NIH-bg-nu-xid; Harlan Laboratories, Indianapolis, IN), and the mice were raised for 6 weeks. Then, dentin/dental pulp-like tissues formed from the implants were removed. The removed tissues were fixed in 4% paraformaldehyde, decalcified in 10% EDTA (pH 7.4), embedded in paraffin, and stained with hematoxylin-eosin (H-E) (Vector Labs) to evaluate levels of the dentin/dental pulp-like tissues (FIG. 2). In this regard, an implant including none of the peptides of the present invention was used as a control group.

FIG. 2 shows microscopic images of dentin/dental pulp-like tissues which were formed *in vivo* for 6 weeks after transplantation of the implants including dental pulp cells and various components, in which A to C are microscopic images at 6 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 200 μm , B 100 μm , C 50 μm); D to F are microscopic images at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 200 μm , E 100 μm , F 50 μm); G to I are microscopic images at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 200 μm , H 100 μm , I 50 μm); and J to L are microscopic images at 6 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 200 μm , K 100 μm , L 50 μm).

As shown in FIG. 2, when the implant including the peptide of the present invention or not was transplanted (A to I), generation of dentin/dental pulp-like tissue was observed at the periphery of HA/TCP particles. However, when the comparative implant including rhBMP-2 was transplanted (J to L), generation of bone-like mineralized tissue and bone marrow-like tissue was observed at the periphery of HA/TCP particles.

Moreover, dentin/dental pulp-like tissues which were the most similar to the *in vivo* dentin-dental pulp tissue were generated at a relatively high level upon transplantation of the implants including the peptides of the present invention (D to I), as compared with transplantation of the implants including none of the peptides of the present invention (A to C).

Example 3-2: Collagen staining analysis of transplanted tissues derived from animals raised for 6 weeks

It is known that collagen is the most abundant organic matrix in dentin and bone, and allows mineral deposition to play an important role in dentin regeneration.

Accordingly, to examine a collagen production level in each of the tissues removed in Example 3-1, the above tissues were subjected to collagen staining (Masson's Trichrome Staining) by using a Masson's Trichrome Stain Kit (Cat. 25088-100) of Polysciences (FIG. 3). In this regard, a tissue transplanted with an implant including none of the peptides of the present invention was used as a control group.

FIG. 3 shows microscopic images of collagen production levels in the dentin/dental pulp-like tissues which were formed *in vivo* for 6 weeks after transplantation of the implants including dental pulp cells and various components, in which A to C show the results at 6 weeks after

transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 500 μ m, B 200 μ m, C 50 μ m); D to F show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 500 μ m, E 200 μ m, F 50 μ m); G to I show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 500 μ m, H 200 μ m, I 50 μ m); and J to L show the results at 6 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 500 μ m, K 200 μ m, L 50 μ m).

As shown in FIG. 3, transplantation of the implants including the peptides of the present invention showed the increased collagen production levels, as compared with transplantation of the control implant.

Example 3-3: Immunostaining analysis of transplanted tissues derived from animals raised for 6 weeks

To evaluate expression levels of DSP which is an odontoblast-specific differentiation marker gene and BSP which is an osteoblast-specific differentiation marker gene in each of the tissues removed in Example 3-1, immunostaining analysis was performed.

Briefly, the removed tissues were fixed in 4% paraformaldehyde, decalcified in 10% EDTA (pH 7.4), embedded

in paraffin, and then immunostained with anti-DSP and anti-BSP antibodies at a dilution of 1:150 as primary antibodies, and reacted with biotin-labeled goat anti-rabbit IgG (Vector Labs) as secondary antibodies to determine DSP and BSP levels (FIG. 4). In this regard, the tissue transplanted with an implant including none of the peptides of the present invention was used as a control group.

FIG. 4 shows immunostaining images showing the results of evaluating the expression levels of DSP which is an odontoblast-specific differentiation marker gene and BSP which is an osteoblast-specific differentiation marker gene in the dentin/dental pulp-like tissues which were formed *in vivo* for 6 weeks after transplantation of the implants including dental pulp cells and various components, in which A and E show the results at 6 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only; B and F show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11; C and G show the results at 6 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12; and D and H show the results at 6 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2, arrows of A, B, C and D indicate DSP-expressing regions in the newly formed dentin-

like tissues, and arrowheads of E, F, G and H indicate BSP-expressing regions in the newly formed bone-like tissues. Scale bar is 50 μ m.

As shown in FIG. 4, transplantation of the control implant showed a low DSP expression level in the newly formed dentin/dental pulp-like tissue (A), whereas transplantation of the implant including the peptide of Group 11 or 12 showed a relatively high DSP expression level in the newly formed mineralized tissue (B and C). However, transplantation of the implant including rhBMP2 showed little DSP expression (D).

Further, transplantation of the control implant (E), the implant (F) including the peptide of Group 11, or the implant (G) including the peptide of Group 12 showed a low BSP expression level in the formed mineralized tissue, whereas transplantation of the implant (H) including rhBMP2 showed a relatively very high BSP expression level in the formed mineralized tissue and the osteoblast-like cells entrapped in the mineralized tissue.

Example 3-4: Morphological analysis of transplanted tissues derived from animals raised for 12 weeks

The levels of dentin/dental pulp-like tissues were evaluated in the same manner as in Example 3-1, except implant-transplanted mice were raised for 12 weeks (FIG. 5).

FIG. 5 shows microscopic images of dentin/dental pulp-like tissues which were formed *in vivo* for 12 weeks after transplantation of the implants including dental pulp cells and various components, in which A to C are microscopic images at 12 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 500 μm , B 200 μm , C 50 μm); D to F are microscopic images at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 500 μm , E 200 μm , F 50 μm); G to I are microscopic images at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 500 μm , H 200 μm , I 50 μm); and J to L are microscopic images at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 500 μm , K 200 μm , L 50 μm).

As shown in FIG. 5, when mice were raised for 12 weeks after transplantation of the implant including the peptide of the present invention or not (A to I), generation of dentin/dental pulp-like tissue and odontoblast processes was observed at the periphery of HA/TCP particles, like in the mice raised for 6 weeks after transplantation. However, when the comparative implant including rhBMP-2 was transplanted (J to L), generation of the bone-like tissue and bone

marrow-like tissue having cells entrapped inside the matrix was observed at the periphery of HA/TCP particles.

Moreover, odontoblasts and dentin/dental pulp complexes which were highly similar to those *in vivo* were generated upon transplantation of the implants including the peptides of the present invention (D to I), as compared with transplantation of the implants including none of the peptides of the present invention (A to C).

Example 3-5: Collagen staining analysis of transplanted tissues derived from animals raised for 12 weeks

In order to examine a collagen production level in each of the tissues removed in Example 3-4, the above tissues were subjected to collagen staining (Masson's Trichrome Staining) by using a Masson's Trichrome Stain Kit (Cat. 25088-100) of Polysciences (FIG. 6). In this regard, a tissue transplanted with an implant including none of the peptides of the present invention was used as a control group.

FIG. 6 shows microscopic images of collagen production levels in the dentin/dental pulp-like tissues which were formed *in vivo* for 12 weeks after transplantation of the implants including dental pulp cells and various components, in which A to C show the results at 12 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 500 μm , B 200 μm , C 50

μm); D to F show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11 (scale bar: D 500 μm, E 200 μm, F 50 μm); G to I show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (scale bar: G 500 μm, H 200 μm, I 50 μm); and J to L show the results at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: J 500 μm, K 200 μm, L 50 μm).

As shown in FIG. 6, the increased collagen production levels were observed at 12 weeks after transplantation of the implants including the peptides of the present invention, like at 6 weeks after transplantation thereof, as compared with transplantation of the control implant.

Example 3-6: Immunostaining analysis of transplanted tissues derived from animals raised for 12 weeks

Immunostaining analysis was performed in the same manner as in Example 3-3, except that the tissues removed in Example 3-4 were used instead of the tissues removed in Example 3-1 (FIG. 7). In this regard, a tissue transplanted with an implant including none of the peptides of the present invention was used as a control group.

FIG. 7 shows immunostaining images showing the results of evaluating the expression levels of DSP which is an

odontoblast-specific differentiation marker gene and BSP which is an osteoblast-specific differentiation marker gene in the dentin/dental pulp-like tissues which were formed *in vivo* for 12 weeks after transplantation of the implants including dental pulp cells and various components, in which A and E show the results at 12 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only; B and F show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 11; C and G show the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12; and D and H show the results at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2, arrows of A, B, C and D indicate DSP-expressing regions in the newly formed dentin-like tissues, and arrowheads of E, F, G and H indicate BSP-expressing regions in the newly formed bone-like tissues. Scale bar is 50 μ m.

As shown in FIG. 7, low DSP expression levels were observed in the newly formed dentin/dental pulp-like tissues at 12 weeks after transplantation of the control implant (A) or the implant including rhBMP2 (D), whereas relatively very high DSP expression levels were observed in the newly formed mineralized tissues at 12 weeks after transplantation of the

implant (B and C) including the peptide of Group 11 or 12, similar to those at 6 weeks after transplantation.

Further, transplantation of the control implant (E), the implant (F) including the peptide of Group 11, or the implant (G) including the peptide of Group 12 showed a low BSP expression level in the formed mineralized tissue, whereas transplantation of the implant (H) including rhBMP2 showed a relatively very high BSP expression level in the formed mineralized tissue and the osteoblast-like cells entrapped in the mineralized tissue.

Accordingly, it can be seen that the peptides of the present invention exhibit the effect of promoting regeneration of dentin/dental pulp complex.

Example 3-7: Analysis of differentiation of transplanted tissue into odontoblasts

A scanning electron microscope was used to examine whether the dental pulp cells included in the implant were differentiated into odontoblasts in the tissues removed in Example 3-4 (FIG. 8). In this regard, a tissue transplanted with an implant including none of the peptides of the present invention was used as a control group.

Briefly, each tissue was immersed and fixed in 2.5% glutaraldehyde/0.1 M cacodylate buffer for 30 minutes, and each fixed tissue was immersed and reacted in a cacodylate buffer containing 1% osmium tetroxide for 1 hour. After

dehydration of the tissues in ethanol and drying, the dried tissues were coated with gold, and visualized under a scanning electron microscope (S-4700, HITACHI, Tokyo, Japan).

FIG. 8 shows scanning microscopic images of the internal structures of the dentin/dental pulp-like tissues which were formed *in vivo* at 12 weeks after transplantation of the implants including dental pulp cells and various components, in which A and B are the results at 12 weeks after transplantation of a control implant including dental pulp cells and HA/TCP only (scale bar: A 50 μm , B 20 μm); C and D are the results at 12 weeks after transplantation of an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (SEQ ID NO: 96) (scale bar: C 50 μm , D 20 μm); and E and F are the results at 12 weeks after transplantation of a comparative implant including dental pulp cells, HA/TCP, and rhBMP-2 (scale bar: E 50 μm , F 20 μm).

As shown in FIG. 8, when the control implants (A and B) were transplanted, odontoblast-like cells having odontoblast processes were partially observed at the periphery of the formed hard tissue, and when the implants (C and D) including the peptides of the present invention were transplanted, numerous odontoblast-like cells were observed at the periphery of the formed hard tissue, and odontoblast processes also extended toward the formed hard tissue.

However, when the comparative implants (E and F) including rhBMP-2 were transplanted, cuboidal cells attached to the surface of the formed hard tissue were observed, indicating typical characteristics of osteoblasts.

Therefore, it can be seen that the peptides of the present invention more effectively form odontoblasts.

Example 4: Effects of peptides for promoting regeneration of dentin or dental pulp tissues and treating dentin hypersensitivity in human tooth

Reportedly, the dentinal wall and the empty pulp cavity of a natural tooth provide the specific local environment for the regeneration of dentin/pulp-like tissues by dental pulp cells (Huang GT, et al. (2010) Tissue engineering. Part A 16(2):605-615). Therefore, formation of dentin/pulp-like tissues in root canal spaces was evaluated.

In detail, among the implants prepared in Example 3-1, a comparative implant or an implant including the peptide of Group 12 (SEQ ID NO: 96) at a concentration of 10 µg/ml was transplanted into the root canal spaces of human teeth for 6 weeks, and dentin/dental pulp-like tissues formed in the respective implants were subjected to hematoxylin-eosin (H-E) staining according to the method of Example 3-1, and photographed by a scanning electron microscope according to the method of Example 3-7 (FIG. 9).

FIG. 9 shows microscopic images and scanning microscopic images showing the results of analyzing odontoblasts/dental pulp-like tissues which were formed after transplantation of the implants including the peptides of the present invention into the root canal spaces of the human teeth, in which A and B show the results of staining the tissues transplanted with a control implant including dental pulp cells and HA/TCP only (scale bar: A 500 μm , B 50 μm); C, D and D' show the results of staining the tissues transplanted with an implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (SEQ ID NO: 96) (scale bar: A 500 μm , B 50 μm); D is magnification of box 1 in C, D' is magnification of box 2 in C (scale bar: C 500 μm , D 50 μm , D' 50 μm); E and F show scanning microscopic images of the tissues transplanted with the control implant including dental pulp cells, HA/TCP only (scale bar: E 50 μm , F 10 μm); G and H show scanning microscopic images of the tissues transplanted with the implant including dental pulp cells, HA/TCP, and the peptide of Group 12 (SEQ ID NO: 96) (scale bar: G 50 μm , H 10 μm); P indicates the regenerated dental pulp, De indicates existing dentinal wall, DT indicates existing dentinal tubules, Od indicates odontoblasts, OP indicates odontoblast processes, and arrows in D indicate regenerated odontoblast-like cells with odontoblast processes.

As shown in FIG. 9, vascularized pulp-like tissues were formed inside all the root canals which were transplanted with the control implant and the implant including the peptide of Group 12.

However, when the implant including the peptide of Group 12 of the present invention was transplanted, odontoblast-like cells exhibited a palisade arrangement on the existing dentinal wall, and their cytoplasmic processes, with lengthened nuclei, extended toward existing dentinal tubules, and newly formed dentin was observed in the existing dentinal wall (C, D and D').

In particular, as shown in scanning microscopic images (E to H), transplantation of the implants (G and H) including the peptide of Group 12 showed that odontoblast-like cells exhibited a palisade arrangement on the existing dentinal wall, and their cytoplasmic processes, with lengthened nuclei, extended toward existing dentinal tubules, as compared with transplantation of the control implants (E and F).

Example 5: Effects of novel peptides on physiologic dentin (tertiary dentin) formation in indirect pulp capping canine model

In order to examine effects of the novel peptides on physiologic dentin (tertiary dentin) formation in indirect

pulp capping models, their dentin regeneration capabilities were evaluated.

In detail, to establish dentin-damaged canine models by indirect pulp capping, a part of enamel in the cervical part was removed from the premolars of 12-month-old adult dogs by using a dental bur. The dentin was shallowly removed according to the exposure degree of the dentin to form a shallow cavity, thereby establishing a shallow cavity model. The dentin was deeply removed such that the dental pulp was seen through, but not exposed to the outside, thereby establishing a deep cavity model. The models were treated with the control group, the peptide of Group 11, or the peptide of Group 12. 3 weeks later, the adult dogs were euthanized, and teeth were extracted therefrom.

The extracted teeth were fixed in 4% paraformaldehyde for 24 hours, washed with PBS (pH 7.4) several times, and then decalcified with 10% formic acid. The decalcified teeth were embedded in paraffin to prepare tissue sections. The tissue sections were stained with hematoxylin/eosin (H&E), and dentin regeneration capabilities were evaluated by histological analysis.

FIG. 10 shows microscopic images of histological analysis of the effects of the novel peptides on physiologic dentin (tertiary dentin) formation in the shallow cavity model among the indirect pulp capping canine models, in

which A to C are microscopic images showing the result of the control group, in which none of the materials was applied to the inlet of dentinal tubule in the exposed dentin (scale bar: A 500 μm , B 100 μm , C 50 μm); D to F are microscopic images showing the result of applying the peptide of Group 11 (1.5 μg) to the inlet of dentinal tubule in the exposed dentin (scale bar: D 500 μm , E 100 μm , F 50 μm); and G to I are microscopic images showing the result of applying the peptide of Group 12 (1.5 μg) to the inlet of dentinal tubule in the exposed dentin (scale bar: G 500 μm , H 100 μm , I 50 μm). P indicates pulp, D indicates dentin, and TD indicates newly formed physiologic (tertiary) dentin.

In the control group, no change was observed in the damaged dentin (FIG. 10A-C). However, in the groups treated with the peptides of Group 11 (FIG. 10D-F) or Group 12 (FIG. 10G-I), regeneration of physiologic tertiary dentin (TD) which was continuous with the odontoblast processes of the original dentin structure was observed beneath the remaining dentin in the damaged dentin.

FIG. 11 shows microscopic images of histological analysis of the effects of the novel peptides on physiologic dentin (tertiary dentin) formation in the deep cavity model among the indirect pulp capping models, in which A to C are microscopic images showing the result of the control group, in which the inlet of dentinal tubule in the exposed dentin

was treated with none of the materials and filled with a dental restorative material, GI cement (Glass Ionomer Cement, Fuji II LC, GC America Inc., Alsip, IL, USA) (scale bar: A 500 μm , B 100 μm , C 50 μm); D to F are microscopic images showing the result of filling the inlet of dentinal tubule in the exposed dentin with GI cement after being treated with the peptide of Group 11 (1.5 μg) (scale bar: D 500 μm , E 100 μm , F 50 μm); and G to I are microscopic images showing the result of filling the inlet of dentinal tubule in the exposed dentin with GI cement after being treated with the peptide of Group 12 (1.5 μg) (scale bar: G 500 μm , H 100 μm , I 50 μm). P indicates pulp, D indicates dentin, and TD indicates newly formed physiologic (tertiary) dentin.

The deep cavity models showed the similar results to the shallow cavity models. In the control group, in which the damaged dentin lesion was treated with GI cement which is a dental restorative material, no change was observed (FIG. 11A-C). However, in the groups treated with the peptides of Group 11 (FIG. 11D-F) or Group 12 (FIG. 11G-I) together with GI cement, regeneration of physiologic tertiary dentin (TD) which was continuous with the odontoblast processes of the original dentin structure was observed beneath the remaining dentin in the damaged dentin.

Example 6: Effects of novel peptides on physiologic dentin (tertiary dentin) formation in direct pulp capping canine model

In order to examine effects of the novel peptides on physiologic dentin (tertiary dentin) formation in direct pulp capping models, their dentin regeneration capabilities were evaluated.

In detail, to establish dentin-damaged canine models by direct pulp capping, the enamel and dentin in the cervical part were removed from the premolars of 12-month-old adult dogs by using a dental bur so that the dental pulp was exposed to the outside. The models were treated with the control group, the peptide of Group 11, or the peptide of Group 12. 3 weeks later, the adult dogs were euthanized and teeth were extracted therefrom.

The extracted teeth were fixed in 4% paraformaldehyde for 24 hours, washed with PBS (pH 7.4) several times, and then decalcified with 10% formic acid. The decalcified teeth were embedded in paraffin to prepare tissue sections. The tissue sections were stained with hematoxylin/eosin (H&E), and dentin regeneration capabilities were evaluated by histological analysis.

FIG. 12 shows microscopic images of histological analysis of the effects of the novel peptides on physiologic dentin (tertiary dentin) formation in the direct pulp

capping models, in which A to C are microscopic images showing the results of the control group, in which the exposed dental pulp was treated with none of the materials and then treated with GI cement (scale bar: A 200 μm , B 100 μm , C 50 μm); D to F are microscopic images showing the results of the positive control group, in which the exposed dental pulp was treated with none of the materials, sealed with a dental restorative material, MTA (Mineral trioxide aggregate, ProRoot MTA, Dentsply Tulsa Dental, Tulsa, OK, USA), and then covered with GI cement (scale bar: D 500 μm , E 200 μm , F 50 μm); G to I are microscopic images showing the results of the groups, in which the exposed dental pulp was treated with the peptide of Group 11 (1.5 μg), sealed with MTA, and then covered with GI cement (scale bar: G 200 μm , H 100 μm , I 50 μm); J to L are microscopic images showing the results of the groups, in which the exposed dental pulp was treated with the peptide of Group 12 (1.5 μg), sealed with MTA, and then covered with GI cement (scale bar: J 500 μm , K 100 μm , L 50 μm). D indicates dentin, OD indicates osteodentin, and TD indicates newly formed physiologic (tertiary) dentin.

In the control group treated with GI cement, no change was observed in the dental pulp of the damaged dentin (FIG. 12A-C). MTA is a dental restorative material widely used in dentin formation and regeneration. However, hard tissues

formed by MTA are known to have cells entrapped in the mineralized tissues and to exhibit osteodentin which is similar to bone having no dentinal tubule which is an important component of the dentin. The present results also showed that osteodentin (OD) with cells entrapped in the mineralized tissue along the remaining original dentin on and beneath the exposed pulp was observed in the groups treated with MTA (FIG. 12D-F). However, in the groups treated with MTA and the peptides of Group 11 (FIG. 12G-I) or Group 12 (FIG. 12J-L), new physiologic dentin (TD) which was continuous with the odontoblast processes of the original dentin was formed along the remaining original dentin on and beneath the exposed pulp.

The results of Examples taken together, it can be seen that the peptides of the present invention increase expression levels of *Dspp*, *Dmp1* and *Nestin* genes which are odontoblast-specific differentiation marker genes, and promote formation of odontoblasts when the peptides are transplanted, together with dental pulp cells, into the body, and in particular, the dental pulp cells promote formation of dentin/dental pulp-like tissue when the peptides are transplanted into root canal spaces. It can be also seen that the same physiologic dentin as observed in the natural human tooth dentin is formed by the novel peptides.

Accordingly, it can be seen that the peptides of the present invention may be used in the treatment of dentin or dental pulp diseases.

This study was supported by the Korea Evaluation Institute of Industrial Technology (KEIT) funded by the Ministry of Trade, Industry & Energy in 2017.

WHAT IS CLAIMED IS:

1. A peptide for promoting regeneration of dentin or dental pulp tissues and treating dentin or dental pulp diseases, comprising an amino acid sequence of the following Formula 1:

K-Y-R1-R2-R3-R4-R5-R6-R7-R8 (Formula 1)

wherein R1 is arginine(R), lysine(K) or glutamine(Q);

R2 is arginine(R) or glutamine(Q);

R3, R4, and R5 are arginine(R) or lysine(K), respectively;

R6 is asparagine(N) or serine(S); and

R7 and R8 are lysine(K) or tyrosine(Y), respectively.

2. The peptide of claim 1, wherein the peptide consists of any one amino acid sequence of SEQ ID NOS: 1 to 96.

3. The peptide of claim 1, wherein the peptide is subjected to N- or C-terminal acetylation, amidation, or methylation; D-amino acid introduction; peptide bond modification including CH₂-NH, CH₂-S, CH₂-S=O, CH₂-CH₂; backbone modification; or side-chain modification.

4. A polynucleotide encoding the peptide of any one of claims 1 to 3.

5. An expression vector comprising the polynucleotide of claim 4.

6. A pharmaceutical composition for preventing or treating dentin or dental pulp diseases, comprising the peptide of any one of claims 1 to 3.

7. The pharmaceutical composition of claim 6, comprising a polypeptide which is prepared by linking repeats of the peptide.

8. The pharmaceutical composition of claim 6, comprising a complex of the peptide linked to a drug having a therapeutic effect on dentin or dental pulp diseases.

9. The pharmaceutical composition of claim 6, further comprising a pharmaceutically acceptable carrier, excipient, or diluent.

10. The pharmaceutical composition of claim 6, wherein the dentin or dental pulp diseases are dentin hypersensitivity, pulp hyperemia, pulpitis, pulp degeneration or pulp necrosis, and gangrenous pulp.

11. A quasi-drug composition for preventing or improving dentin or dental pulp diseases, comprising the peptide of any one of claims 1 to 3.

12. The quasi-drug composition of claim 11, wherein the quasi-drug composition is oral antiseptic mouthwashes, oral hygiene products, toothpastes, floss, or oral ointments.

13. A health functional food composition for preventing or improving dentin or dental pulp diseases, comprising the peptide of any one of claims 1 to 3.

14. The health functional food composition of claim 13, wherein the health functional food composition is used in the preparation of beverages, teas, spices, gums, or confectionery.

15. A method of preventing or treating dentin or dental pulp diseases, the method comprising administering to a subject a composition comprising the peptide of any one of claims 1 to 3.

16. A method of promoting regeneration of dentin or dental pulp tissues, the method comprising administering to

a subject a composition comprising the peptide of any one of claims 1 to 3.