

POLYMORPHIC AND AMORPHOUS FORMS OF (R)-2-HYDROXY-2-METHYL-4-(2,4,5-TRIMETHYL-3,6-DIOXOCYCLOHEXA-1,4-DIENYL)BUTANAMIDE

[0001] The application claims priority to, and the benefit of, U.S. Provisional Patent Application No. 62/092,743, filed December 16, 2014, entitled POLYMORPHIC AND AMORPHOUS FORMS OF (R)-2-HYDROXY-2-METHYL-4-(2,4,5-TRIMETHYL-3,6-DIOXOCYCLOHEXA-1,4-DIENYL)BUTANAMIDE, and U.S. Provisional Patent Application No. 62/133,276, filed March 13, 2015, entitled POLYMORPHIC AND AMORPHOUS FORMS OF (R)-2-HYDROXY-2-METHYL-4-(2,4,5-TRIMETHYL-3,6-DIOXOCYCLOHEXA-1,4-DIENYL)BUTANAMIDE, the contents of both of which are herein incorporated by reference in their entirety for all purposes.

TECHNICAL FIELD

[0002] The application discloses compositions and methods useful for treatment or suppression of diseases, developmental delays and symptoms related to oxidative stress disorders. Examples of such disorders include mitochondrial disorders, impaired energy processing disorders, neurodegenerative diseases and diseases of aging. The application further discloses methods of making such compositions.

BACKGROUND

[0003] Oxidative stress is caused by disturbances to the normal redox state within cells. An imbalance between routine production and detoxification of reactive oxygen species such as peroxides and free radicals can result in oxidative damage to the cellular structure and machinery. The most important source of reactive oxygen species under normal conditions in aerobic organisms is probably the leakage of activated oxygen from mitochondria during normal oxidative respiration. Impairments associated with this process are suspected to contribute to mitochondrial disease, neurodegenerative disease, and diseases of aging.

[0004] Mitochondria are organelles in eukaryotic cells, popularly referred to as the “powerhouse” of the cell. One of their primary functions is oxidative phosphorylation. The molecule adenosine triphosphate (ATP) functions as an energy “currency” or energy carrier in

the cell, and eukaryotic cells derive the majority of their ATP from biochemical processes carried out by mitochondria. These biochemical processes include the citric acid cycle (the tricarboxylic acid cycle, or Krebs cycle), which generates reduced nicotinamide adenine dinucleotide ($\text{NADH} + \text{H}^+$) from oxidized nicotinamide adenine dinucleotide (NAD^+), and oxidative phosphorylation, during which $\text{NADH} + \text{H}^+$ is oxidized back to NAD^+ . The citric acid cycle also reduces flavin adenine dinucleotide, or FAD, to FADH_2 ; FADH_2 also participates in oxidative phosphorylation.

[0005] The electrons released by oxidation of $\text{NADH} + \text{H}^+$ are shuttled down a series of protein complexes (Complex I, Complex II, Complex III, and Complex IV) known as the mitochondrial respiratory chain. These complexes are embedded in the inner membrane of the mitochondrion. Complex IV, at the end of the chain, transfers the electrons to oxygen, which is reduced to water. The energy released as these electrons traverse the complexes is used to generate a proton gradient across the inner membrane of the mitochondrion, which creates an electrochemical potential across the inner membrane. Another protein complex, Complex V (which is not directly associated with Complexes I, II, III and IV) uses the energy stored by the electrochemical gradient to convert ADP into ATP.

[0006] When cells in an organism are temporarily deprived of oxygen, anaerobic respiration is utilized until oxygen again becomes available or the cell dies. The pyruvate generated during glycolysis is converted to lactate during anaerobic respiration. The buildup of lactic acid is believed to be responsible for muscle fatigue during intense periods of activity, when oxygen cannot be supplied to the muscle cells. When oxygen again becomes available, the lactate is converted back into pyruvate for use in oxidative phosphorylation.

[0007] Oxygen poisoning or toxicity is caused by high concentrations of oxygen that may be damaging to the body and increase the formation of free-radicals and other structures such as nitric oxide, peroxynitrite, and trioxidane. Normally, the body has many defense systems against such damage but at higher concentrations of free oxygen, these systems are eventually overwhelmed with time, and the rate of damage to cell membranes exceeds the capacity of systems which control or repair it. Cell damage and cell death then results.

[0008] Qualitative and/or quantitative disruptions in the transport of oxygen to tissues result in energy disruption in the function of red cells and contribute to various diseases such as haemoglobinopathies. Haemoglobinopathy is a kind of genetic defect that results in abnormal structure of one of the globin chains of the hemoglobin molecule. Common

haemoglobinopathies include thalassemia and sickle-cell disease. Thalassemia is an inherited autosomal recessive blood disease. In thalassemia, the genetic defect results in reduced rate of synthesis of one of the globin chains that make up hemoglobin. While thalassemia is a quantitative problem of too few globins synthesized, sickle-cell disease is a qualitative problem of synthesis of an incorrectly functioning globin. Sickle-cell disease is a blood disorder characterized by red blood cells that assume an abnormal, rigid, and sickle shape. Sickling decreases the cells' flexibility and results in their restricted movement through blood vessels, depriving downstream tissues of oxygen.

[0009] Mitochondrial dysfunction contributes to various disease states. Some mitochondrial diseases are due to mutations or deletions in the mitochondrial genome. If a threshold proportion of mitochondria in the cell is defective, and if a threshold proportion of such cells within a tissue have defective mitochondria, symptoms of tissue or organ dysfunction can result. Practically any tissue can be affected, and a large variety of symptoms may be present, depending on the extent to which different tissues are involved. Some examples of mitochondrial diseases are Friedreich's ataxia (FRDA), Leber's Hereditary Optic Neuropathy (LHON), mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke (MELAS), Myoclonus Epilepsy Associated with Ragged-Red Fibers (MERRF) syndrome, Leigh's syndrome, and respiratory chain disorders. Most mitochondrial diseases involve children who manifest the signs and symptoms of accelerated aging, including neurodegenerative diseases, stroke, blindness, hearing impairment, vision impairment, diabetes, and heart failure.

[0010] Friedreich's ataxia is an autosomal recessive neurodegenerative and cardiodegenerative disorder caused by decreased levels of the protein Frataxin. The disease causes the progressive loss of voluntary motor coordination (ataxia) and cardiac complications. Symptoms typically begin in childhood, and the disease progressively worsens as the patient grows older; patients eventually become wheelchair-bound due to motor disabilities.

[0011] Leber's Hereditary Optic Neuropathy (LHON) is a disease characterized by blindness which occurs on average between 27 and 34 years of age. Other symptoms may also occur, such as cardiac abnormalities and neurological complications.

[0012] Mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke (MELAS) can manifest itself in infants, children, or young adults. Strokes, accompanied by vomiting and seizures, are one of the most serious symptoms; it is postulated that the metabolic impairment of

mitochondria in certain areas of the brain is responsible for cell death and neurological lesions, rather than the impairment of blood flow as occurs in ischemic stroke.

[0013] Myoclonus Epilepsy Associated with Ragged-Red Fibers (MERRF) syndrome is one of a group of rare muscular disorders that are called mitochondrial encephalomyopathies.

Mitochondrial encephalomyopathies are disorders in which a defect in the genetic material arises from a part of the cell structure that releases energy (mitochondria). This can cause a dysfunction of the brain and muscles (encephalomyopathies). The mitochondrial defect as well as “ragged-red fibers” (an abnormality of tissue when viewed under a microscope) are always present. The most characteristic symptom of MERRF syndrome is myoclonic seizures that are usually sudden, brief, jerking, spasms that can affect the limbs or the entire body, difficulty speaking (dysarthria), optic atrophy, short stature, hearing loss, dementia, and involuntary jerking of the eyes (nystagmus) may also occur.

[0014] Leigh’s syndrome is a rare inherited neurometabolic disorder characterized by degeneration of the central nervous system where the symptoms usually begin between the ages of 3 months to 2 years and progress rapidly. In most children, the first signs may be poor sucking ability and loss of head control and motor skills. These symptoms may be accompanied by loss of appetite, vomiting, irritability, continuous crying, and seizures. As the disorder progresses, symptoms may also include generalized weakness, lack of muscle tone, and episodes of lactic acidosis, which can lead to impairment of respiratory and kidney function. Heart problems may also occur.

[0015] Co-Enzyme Q10 Deficiency is a respiratory chain disorder, with syndromes such as myopathy with exercise intolerance and recurrent myoglobin in the urine manifested by ataxia, seizures or mental retardation and leading to renal failure (Di Mauro et al., (2005) *Neuromusc. Disord.*, 15:311-315), “Childhood-onset cerebellar ataxia and cerebellar atrophy,” (Masumeci et al., (2001) *Neurology* 56:849-855 and Lamperti *et al.*, (2003) 60:1206:1208); and infantile encephalomyopathy associated with nephrosis. Biochemical measurement of muscle homogenates of patients with CoQ10 deficiency showed severely decreased activities of respiratory chain complexes I and II + III, while complex IV (COX) was moderately decreased (Gempel *et al.*, (2007) *Brain*, 130(8):2037-2044).

[0016] Complex I Deficiency or NADH dehydrogenase NADH-CoQ reductase deficiency is a respiratory chain disorder, with symptoms classified by three major forms: (1) fatal infantile multisystem disorder, characterized by developmental delay, muscle weakness, heart disease,

congenital lactic acidosis, and respiratory failure; (2) myopathy beginning in childhood or in adult life, manifesting as exercise intolerance or weakness; and (3) mitochondrial encephalomyopathy (including MELAS), which may begin in childhood or adult life and consists of variable combinations of symptoms and signs, including ophthalmoplegia, seizures, dementia, ataxia, hearing loss, pigmentary retinopathy, sensory neuropathy, and uncontrollable movements.

[0017] Complex II Deficiency or Succinate dehydrogenase deficiency is a respiratory chain disorder with symptoms including encephalomyopathy and various manifestations, including failure to thrive, developmental delay, hypotonia, lethargy, respiratory failure, ataxia, myoclonus and lactic acidosis.

[0018] Complex III Deficiency or Ubiquinone-cytochrome C oxidoreductase deficiency is a respiratory chain disorder with symptoms categorized in four major forms: (1) fatal infantile encephalomyopathy, congenital lactic acidosis, hypotonia, dystrophic posturing, seizures, and coma; (2) encephalomyopathies of later onset (childhood to adult life): various combinations of weakness, short stature, ataxia, dementia, hearing loss, sensory neuropathy, pigmentary retinopathy, and pyramidal signs; (3) myopathy, with exercise intolerance evolving into fixed weakness; and (4) infantile histiocytoid cardiomyopathy.

[0019] Complex IV Deficiency or Cytochrome C oxidase deficiency is a respiratory chain disorder with symptoms categorized in two major forms: (1) encephalomyopathy, where patients typically are normal for the first 6 to 12 months of life and then show developmental regression, ataxia, lactic acidosis, optic atrophy, ophthalmoplegia, nystagmus, dystonia, pyramidal signs, respiratory problems and frequent seizures; and (2) myopathy with two main variants: (a) Fatal infantile myopathy-may begin soon after birth and accompanied by hypotonia, weakness, lactic acidosis, ragged-red fibers, respiratory failure, and kidney problems; and (b) Benign infantile myopathy- may begin soon after birth and accompanied by hypotonia, weakness, lactic acidosis, ragged-red fibers, respiratory problems, but (if the child survives) followed by spontaneous improvement.

[0020] Complex V Deficiency or ATP synthase deficiency is a respiratory chain disorder including symptoms such as slow, progressive myopathy.

[0021] CPEO or Chronic Progressive External Ophthalmoplegia Syndrome is a respiratory chain disorder including symptoms such as visual myopathy, retinitis pigmentosa, or dysfunction of the central nervous system.

[0022] Kearns-Sayre Syndrome (KSS) is a mitochondrial disease characterized by a triad of features including: (1) typical onset in persons younger than age 20 years; (2) chronic, progressive, external ophthalmoplegia; and (3) pigmentary degeneration of the retina. In addition, KSS may include cardiac conduction defects, cerebellar ataxia, and raised cerebrospinal fluid (CSF) protein levels (*e.g.*, >100 mg/dL). Additional features associated with KSS may include myopathy, dystonia, endocrine abnormalities (*e.g.*, diabetes, growth retardation or short stature, and hypoparathyroidism), bilateral sensorineural deafness, dementia, cataracts, and proximal renal tubular acidosis.

[0023] Maternally inherited diabetes and deafness (MIDD) is a mitochondrial disorder characterized by maternally transmitted diabetes and sensorineural deafness. In most cases, MIDD is caused by a point mutation in the mitochondrial gene MT-TL1, encoding the mitochondrial tRNA for leucine, and in rare cases in MT-TE and MT-TK genes, encoding the mitochondrial tRNAs for glutamic acid, and lysine, respectively.

[0024] In addition to congenital disorders involving inherited defective mitochondria, acquired mitochondrial dysfunction contributes to diseases, particularly neurodegenerative disorders associated with aging like Parkinson's, Alzheimer's, and Huntington's Diseases. The incidence of somatic mutations in mitochondrial DNA rises exponentially with age; diminished respiratory chain activity is found universally in aging people. Mitochondrial dysfunction is also implicated in excitotoxic, neuronal injury, such as that associated with cerebrovascular accidents, seizures and ischemia.

[0025] Some of the above diseases appear to be caused by defects in Complex I of the respiratory chain. Electron transfer from Complex I to the remainder of the respiratory chain is mediated by the compound coenzyme Q (also known as Ubiquinone). Oxidized coenzyme Q (CoQ_{ox} or Ubiquinone) is reduced by Complex I to reduced coenzyme Q (CoQ_{red} or Ubiquinol). The reduced coenzyme Q then transfers its electrons to Complex III of the respiratory chain, where it is re-oxidized to CoQ_{ox} (Ubiquinone). CoQ_{ox} can then participate in further iterations of electron transfer.

[0026] Very few treatments are available for patients suffering from these mitochondrial diseases. Recently, the compound Idebenone has been proposed for treatment of Friedreich's ataxia. While the clinical effects of Idebenone have been relatively modest, the complications of mitochondrial diseases can be so severe that even marginally useful therapies are preferable to the untreated course of the disease. Another compound, MitoQ, has been proposed for treating

mitochondrial disorders (see U.S. Patent No. 7,179,928); clinical results for MitoQ have not yet been reported. Administration of coenzyme Q10 (CoQ10) and vitamin supplements has shown only transient beneficial effects in individual cases of KSS. CoQ10 supplementation has also been used for the treatment of CoQ10 deficiency with mixed results.

[0027] Oxidative stress is suspected to be important in neurodegenerative diseases such as Motor Neuron Disease, Amyotrophic Lateral Sclerosis (ALS), Creutzfeldt-Jakob disease, Machado-Joseph disease, Spino-cerebellar ataxia, Multiple sclerosis (MS), Parkinson's disease, Alzheimer's disease, and Huntington's disease. Oxidative stress is thought to be linked to certain cardiovascular disease and also plays a role in the ischemic cascade due to oxygen reperfusion injury following hypoxia. This cascade includes both strokes and heart attacks.

[0028] Damage accumulation theory, also known as the free radical theory of aging, invokes random effects of free radicals produced during aerobic metabolism that cause damage to DNA, lipids and proteins and accumulate over time. The concept of free radicals playing a role in the aging process was first introduced by Himan D (1956), "Aging –A theory based on free-radical and radiation chemistry," *J. Gerontol.* 11, 298-300.

[0029] According to the free radical theory of aging, the process of aging begins with oxygen metabolism (Valko et al, (2004), "Role of oxygen radicals in DNA damage and cancer incidence," *Mol. Cell. Biochem.*, 266, 37-56). Even under ideal conditions some electrons "leak" from the electron transport chain. These leaking electrons interact with oxygen to produce superoxide radicals, so that under physiological conditions, about 1-3% of the oxygen molecules in the mitochondria are converted into superoxide. The primary site of radical oxygen damage from superoxide radical is mitochondrial DNA (mtDNA) (Cadenas *et al.*, (2000) Mitochondrial free radical generation, oxidative stress and aging, *Free Radic. Res*, 28, 601-609). The cell repairs much of the damage done to nuclear DNA (nDNA) but mtDNA repair seems to be less efficient. Therefore, extensive mtDNA damage accumulates over time and shuts down mitochondria causing cells to die and the organism to age.

[0030] Some of the diseases associated with increasing age are cancer, diabetes mellitus, hypertension, atherosclerosis, ischemia/reperfusion injury, rheumatoid arthritis, neurodegenerative disorders such as dementia, Alzheimer's and Parkinson's. Diseases resulting from the process of aging as a physiological decline include decreases in muscle strength, cardiopulmonary function, vision and hearing as well as wrinkled skin and graying hair.

[0031] The ability to adjust biological production of energy has applications beyond the diseases described above. Various other disorders can result in suboptimal levels of energy biomarkers (sometimes also referred to as indicators of energetic function), such as ATP levels. Treatments for these disorders are also needed, in order to modulate one or more energy biomarkers to improve the health of the patient. In other applications, it can be desirable to modulate certain energy biomarkers away from their normal values in an individual that is not suffering from disease. For example, if an individual is undergoing an extremely strenuous undertaking, it can be desirable to raise the level of ATP in that individual.

[0032] Certain polymorphic or amorphous forms of a drug can have advantageous characteristics versus other forms; for example, increased stability, increased solubility, better handling properties, lack of associated toxic solvents, and increased purity.

[0033] Example 16 of PCT Application No. PCT/US2008/082374, published as WO 2009/061744 on May 14, 2009, describes a synthesis for racemic 2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide; this Example does not specifically describe the synthesis of any particular polymorphic or amorphous form for (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide or any particular stereoisomer thereof.

SUMMARY

[0034] In one aspect, the present disclosure refers to a polymorph of an anhydrate or a hydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is selected from the group consisting of Form I, Form III and Form II; wherein a powder X-ray diffraction pattern for polymorph Form I comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 17.03, and 17.26; wherein a powder X-ray diffraction pattern for polymorph Form III comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 14.02, 15.23, and 21.10; wherein a powder X-ray diffraction pattern for polymorph Form II comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, and 19.33; and wherein the XRPD pattern is generated at 25 °C. In one example, there is disclosed a polymorph of an anhydrate, a hydrate, or a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is selected from the group consisting of Forms I-VI as described in Table A or Tables 2-7 respectively. In another aspect, the present disclosure refers to a composition comprising the polymorph as disclosed herein, wherein at least about 95% by mole of the composition is the

polymorph Form I, exclusive of any solvents, carriers or excipients. In yet another aspect, the present disclosure refers to a pharmaceutical composition comprising the polymorph as disclosed herein, or the composition as disclosed herein, and a pharmaceutically acceptable carrier. In another aspect, the present disclosure refers to a composition comprising the polymorph as disclosed herein, wherein the composition is essentially free of Forms II-VI, wherein a powder X-ray diffraction pattern for polymorph Form V comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, and 15.45; wherein a powder X-ray diffraction pattern for polymorph Form III comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 14.02, 15.23, and 21.10; wherein a powder X-ray diffraction pattern for polymorph Form II comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, and 19.33; wherein a powder X-ray diffraction pattern for polymorph Form IV comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 12.97, and 13.20; and wherein a powder X-ray diffraction pattern for polymorph Form VI comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.91, and 12.94; and wherein the XRPD pattern is generated at 25 °C. In a further aspect, the present disclosure refers to a composition comprising the polymorph as disclosed herein, wherein at least about 95% by mole of the composition is the

polymorph Form I, exclusive of any solvents, carriers or excipients. In one example, there is disclosed a polymorph of an anhydrate, a hydrate, or a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is selected from the group consisting of Form I, Form II, Form III, Form IV, Form V, and Form VI as described herein.

[0035] In another example, there is disclosed a polymorph of an anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is Form I as described herein. In some examples, the polymorph has a powder x-ray diffraction pattern substantially as shown in Figure 10. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 17.03, and 17.26. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 17.03, and 17.26. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 15.33, 17.03, and 17.26. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 15.33, 17.03, 17.26, and 18.72. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, where in the angular positions may vary by ± 0.2 : 12.06, 15.33, 17.03, 17.26, and 18.72. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 7.67, 10.75, 12.06, 15.33, 16.41, 17.03, 17.26, 18.72, 20.04, and 23.92. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 7.67, 10.75, 12.06, 15.33, 16.41, 17.03, 17.26, 18.72, 20.04, 20.64, 20.91, 21.14, 22.58, 23.13, 23.92, 24.19, 24.53, 27.21, and 27.56. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 5.48, 7.67, 10.75, 12.06, 15.33, 16.41, 17.03, 17.26, 17.71, 17.94, 18.40, 18.72, 19.51, 20.04, 20.64, 20.91, 21.14, 21.55, 21.91, 22.25, 22.58, 23.13, 23.41, 23.92, 24.19, 24.53, 25.64, 26.13, 26.34, 27.21, 27.56, 28.01, 29.04, and 29.46. In some examples, including any of the foregoing examples, the angular positions

may vary by ± 0.1 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.05 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.02 . In some examples, including any of the foregoing examples, the polymorph is isolated. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein the composition is essentially free of Forms II-VI, wherein Forms II-VI are described in Table A or Tables 3-7 respectively. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein at least about 95% of the composition is the polymorph, exclusive of any solvents, carriers or excipients.

[0036] In another example, there is disclosed a polymorph of an anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is Form V as described herein. In some examples, the polymorph has a powder x-ray diffraction pattern substantially as shown in a) or b) of Figure 30. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, and 15.45. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, and 15.45. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, 15.45, and 23.96. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, 14.80, 15.45, 23.96. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, 12.93, 15.45, and 26.05. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, 12.93, 14.80, 15.45, 16.53, 23.96, 24.54, and 26.05. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, 12.93, 14.80, 15.45, 16.10, 16.34, 16.53, 20.18, 22.52, 22.86, 23.96, 24.54, and 26.05. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the

angular positions may vary by ± 0.2 : 6.91, 7.72, 9.61, 11.49, 11.86, 12.93, 13.19, 13.87, 14.80, 15.45, 16.10, 16.34, 16.53, 17.14, 17.85, 19.12, 19.85, 20.18, 21.00, 22.06, 22.52, 22.86, 23.09, 23.96, 24.54, 25.26, 26.05, and 26.90. In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.1 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.05 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.02 . In some examples, including any of the foregoing examples, the polymorph is isolated. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein the composition is essentially free of Forms I-IV and-VI, wherein Forms I-IV and-VI are described in Table A or Tables 2, or 4-7 respectively. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein at least about 95% of the composition is the polymorph, exclusive of any solvents, carriers or excipients.

[0037] In another example, there is described a polymorph of a hydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is Form III as described herein. In some examples, the polymorph has a powder x-ray diffraction pattern substantially as shown in a) or b) of Figure 20. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 14.02, 15.23, and 21.10. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 14.02, 15.23, and 21.10. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.16, 14.02, 15.23, and 21.10. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.16, 13.74, 14.02, 15.23, and 21.10. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.16, 14.02, 15.23, 21.10, and 22.69. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.16, 11.81, 13.74, 14.02, 15.23, 21.10, 22.69, and 23.90. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions,

wherein the angular positions may vary by ± 0.2 : 9.16, 11.81, 13.74, 14.02, 15.23, 17.35, 21.10, 22.69, 23.15, 23.90, and 26.10. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.16, 11.53, 11.81, 12.68, 12.93, 13.74, 14.02, 15.23, 16.53, 17.35, 17.98, 18.54, 19.09, 20.23, 21.10, 21.93, 22.69, 23.15, 23.50, 23.90, 24.65, 25.09, 25.46, 25.79, 26.10, 27.79, 28.22, 28.93, and 29.33. In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.1 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.05 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.02 . In some examples, including any of the foregoing examples, the polymorph is isolated. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein the composition is essentially free of Forms I, II, IV, V, and VI, wherein Forms I, II, IV, V, and VI are described in Table A or Tables 2-3 and 5-7 respectively. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein at least about 95% of the composition is the polymorph, exclusive of any solvents, carriers or excipients.

[0038] In another example, there is described a polymorph of a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is Form II as described herein. In some examples, the polymorph has a powder x-ray diffraction pattern substantially as shown in a) or b) of Figure 15. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, and 19.33. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, 19.33. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, 10.85, and 19.33. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : . 9.63, 11.33, 10.85, 19.33, and 17.3. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 10.85, 11.33, 13.47, and 19.33. In some examples, a powder X-ray diffraction

pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 5.76, 8.04, 9.63, 10.85, 11.33, 11.97, 13.47, 14.75, 17.37, 17.71, and 19.33. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 5.76, 8.04, 9.63, 10.85, 11.33, 11.97, 13.47, 14.75, 16.42, 16.89, 17.37, 17.71, 19.33, 22.89, and 24.59. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 5.76, 6.72, 7.57, 8.04, 9.63, 10.85, 11.33, 11.97, 12.38, 13.13, 13.47, 14.75, 15.28, 16.42, 16.89, 17.37, 17.71, 18.17, 18.66, 19.33, 20.01, 20.29, 20.67, 20.90, 21.36, 21.54, 21.80, 22.55, 22.89, 23.27, 23.54, 23.87, 24.35, 24.59, 24.87, 25.29, 25.55, 25.89, 26.44, 27.49, 28.01, 28.39, and 29.17. In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.1 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.05 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.02 . In some examples, including any of the foregoing examples, the polymorph is isolated. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein the composition is essentially free of Forms I, and III-VI, wherein Forms I, and III-VI are described in Table A or Tables 2-4 and 6-7 respectively. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein at least about 95% of the composition is the polymorph, exclusive of any solvents, carriers or excipients.

[0039] In another example, there is described a polymorph of a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is Form IV as described herein. In some examples, the polymorph has a powder x-ray diffraction pattern substantially as shown in a), b), or c) of Figure 25. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 12.97, and 13.20. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 12.97, 13.20. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 8.76, 12.97, and 13.20. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following

angular positions, wherein the angular positions may vary by ± 0.2 : 0.2: 4.31, 8.76, 12.97, 13.20, 16.66. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 8.76, 12.97, 13.20, and 16.66. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 7.94, 8.76, 12.97, 13.20, 16.66, 17.33, and 20.57. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 7.94, 8.76, 12.97, 13.20, 15.08, 16.66, 17.33, 19.09, 20.57, and 21.58. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 5.77, 6.28, 7.53, 7.94, 8.76, 9.39, 9.87, 10.54, 11.07, 11.68, 12.02, 12.28, 12.97, 13.20, 13.52, 14.40, 15.08, 15.90, 16.66, 16.96, 17.33, 17.59, 18.77, 19.09, 19.74, 20.27, 20.57, 21.09, 21.58, 22.81, 23.23, 24.01, 24.65, and 25.60. In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.1 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.05 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.02 . In some examples, including any of the foregoing examples, the polymorph is isolated. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein the composition is essentially free of Forms I-III and V-VI, wherein Forms I-III and V-VI are described in Table A or Tables 2-5 and 7 respectively. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein at least about 95% of the composition is the polymorph, exclusive of any solvents, carriers or excipients.

[0040] In another example, there is described a polymorph of a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is Form VI as described herein. In some examples, the polymorph has a powder x-ray diffraction pattern substantially as shown in a) of Figure 33. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.91, and 12.94. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.91, and 12.94. In some examples, a powder X-ray diffraction pattern for the polymorph comprises

characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.91, 12.94, and 15.71. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.91, 12.94, 15.71, and 19.13. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.41, 9.91, 12.94, and 13.29. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 8.85, 9.41, 9.91, 12.94, 13.29, 16.67, and 19.13. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.39, 6.27, 8.85, 9.41, 9.91, 11.32, 12.94, 13.29, 14.03, 16.67, 19.13, 20.76, and 22.06. In some examples, a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.39, 6.27, 7.00, 8.62, 8.85, 9.41, 9.91, 11.32, 11.50, 12.25, 12.56, 12.94, 13.29, 14.03, 14.82, 15.10, 15.44, 15.71, 16.01, 16.67, 16.91, 17.33, 17.59, 18.33, 18.75, 19.13, 20.25, 20.76, 21.68, 22.06, 22.27, 22.61, 22.94, 24.01, 24.33, 24.65, 25.48, 26.05, 28.63, and 29.18. In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.1 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.05 . In some examples, including any of the foregoing examples, the angular positions may vary by ± 0.02 . In some examples, including any of the foregoing examples, the polymorph is isolated. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein the composition is essentially free of Forms I-V, wherein Forms I-V are described in Table A or Tables 2-6 respectively. In some examples, including any of the foregoing examples, the polymorph is present in a composition, wherein at least about 95% of the composition is the polymorph, exclusive of any solvents, carriers or excipients.

[0041] In another example, there is described a composition comprising amorphous (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide. In some examples, the amorphous form is isolated. In some examples, the composition is essentially free of Forms I-VI, wherein Forms I-VI are described in Table A or Tables 2-7 respectively. In some examples, including any of the foregoing examples, at least about 95% of the composition is

amorphous (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, exclusive of any solvents, carriers or excipients.

[0042] In another example, there is described a pharmaceutical composition comprising a polymorphic or amorphous form of an anhydrate, a hydrate, or a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, or composition comprising such form, as described herein, including any of the foregoing or hereafter examples, and a pharmaceutically acceptable carrier. In some examples, the form is polymorph Form I. In some examples, the form is polymorph Form II. In some examples, the form is polymorph Form III. In some examples, the form is polymorph Form IV. In some examples, the form is polymorph Form V. In some examples, the form is polymorph Form VI. In some examples, the form is amorphous. In some examples, the pharmaceutical composition has an HPLC purity of more than about 95% for the anhydrate, hydrate, or solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, exclusive of any solvents, carriers or excipients. In some examples, the pharmaceutical composition has an HPLC purity of more than about 99% for the anhydrate, hydrate, or solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, exclusive of any solvents, carriers or excipients. In some examples, the pharmaceutical composition has an HPLC purity of more than about 99.9% for the anhydrate, hydrate, or solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, exclusive of any solvents, carriers or excipients. HPLC purity % refers to the proportional area of a given compound's HPLC peak with respect to the area of all peaks in a given HPLC spectrum. HPLC % is calculated by dividing the area of a compound peak by the area of all peaks, in a HPLC spectrum, and multiplying this quotient by one-hundred.

[0043] In another example, there is described a pharmaceutical composition comprising an active agent and a pharmaceutically acceptable carrier, wherein the active agent consists of, or consists essentially of, a polymorphic or amorphous form of an anhydrate, a hydrate, or a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide as described herein. In some examples, the form is polymorph Form I. In some examples, the form is polymorph Form II. In some examples, the form is polymorph Form III. In some examples, the form is polymorph Form IV. In some examples, the form is polymorph Form V. In some examples, the form is polymorph Form VI. In some examples, the form is amorphous.

[0044] In another examples, there is described a method of treating or suppressing an oxidative stress disorder, modulating one or more energy biomarkers, normalizing one or more energy biomarkers, or enhancing one or more energy biomarkers, comprising administering to an individual in need thereof a therapeutically effective amount or effective amount of a polymorphic or amorphous form of an anhydrate, a hydrate, or a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, or composition comprising such form, as described herein, including any of the foregoing or hereafter examples. The method can use any individual polymorphic or amorphous form as described herein, or a combination of such. In some examples, the form is polymorph Form I. In some examples, the form is polymorph Form II. In some examples, the form is polymorph Form III. In some examples, the form is polymorph Form IV. In some examples, the form is polymorph Form V. In some examples, the form is polymorph Form VI. In some examples, the form is amorphous. In some examples, including any of the foregoing examples, the anhydrate, hydrate, or solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide is administered in a pharmaceutical composition comprising the polymorphic or amorphous form and a pharmaceutically acceptable carrier. In some examples, including any of the foregoing examples, the pharmaceutical composition comprises an active agent consisting essentially of the polymorphic or amorphous form of the anhydrate, hydrate, or solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide. In one aspect, the present disclosure refers to use of the polymorph as disclosed herein, the composition as disclosed herein, or the pharmaceutical composition as disclosed herein in the manufacture of a medicament for treating or suppressing an oxidative stress disorder. In some examples, including any of the foregoing examples, the method is a method of treating or suppressing an oxidative stress disorder. In some examples, including any of the foregoing examples, the method is a method of treating an oxidative stress disorder. In some examples, including any of the foregoing examples, the method is a method of suppressing an oxidative stress disorder. In some examples, including any of the foregoing examples, the oxidative stress disorder is selected from the group consisting of: a mitochondrial disorder; an inherited mitochondrial disease; Alpers Disease; Barth syndrome; a Beta-oxidation Defect; Carnitine-Acyl-Carnitine Deficiency; Carnitine Deficiency; a Creatine Deficiency Syndrome; Co-Enzyme Q10 Deficiency; Complex I Deficiency; Complex II Deficiency; Complex III

Deficiency; Complex IV Deficiency; Complex V Deficiency; COX Deficiency; chronic progressive external ophthalmoplegia (CPEO); CPT I Deficiency; CPT II Deficiency; Friedreich's Ataxia (FA); Glutaric Aciduria Type II; Kearns-Sayre Syndrome (KSS); Lactic Acidosis; Long-Chain Acyl-CoA Dehydrogenase Deficiency (LCAD); LCHAD; Leigh's syndrome; Leigh-like Syndrome; Leber's Hereditary Optic Neuropathy (LHON); Lethal Infantile Cardiomyopathy (LIC); Luft Disease; Multiple Acyl-CoA Dehydrogenase Deficiency (MAD); Medium-Chain Acyl-CoA Dehydrogenase Deficiency (MCAD); Mitochondrial Myopathy, Encephalopathy, Lactacidosis, Stroke (MELAS); Myoclonic Epilepsy with Ragged Red Fibers (MERRF); Mitochondrial Recessive Ataxia Syndrome (MIRAS); Mitochondrial Cytopathy, Mitochondrial DNA Depletion; Mitochondrial Encephalopathy; Mitochondrial Myopathy; Myoneurogastrointestinal Disorder and Encephalopathy (MNGIE); Neuropathy, Ataxia, and Retinitis Pigmentosa (NARP); Pearson Syndrome; Pyruvate Carboxylase Deficiency; Pyruvate Dehydrogenase Deficiency; a POLG Mutation; a Respiratory Chain Disorder; Short-Chain Acyl-CoA Dehydrogenase Deficiency (SCAD); SCHAD; Very Long-Chain Acyl-CoA Dehydrogenase Deficiency (VLCAD); a myopathy; cardiomyopathy; encephalomyopathy; a neurodegenerative disease; Parkinson's disease; Alzheimer's disease; amyotrophic lateral sclerosis (ALS); a motor neuron disease; a neurological disease; epilepsy; an age-associated disease; macular degeneration; diabetes; metabolic syndrome; cancer; brain cancer; a genetic disease; Huntington's Disease; a mood disorder; schizophrenia; bipolar disorder; a pervasive developmental disorder; autistic disorder; Asperger's syndrome; childhood disintegrative disorder (CDD); Rett's disorder; PDD-not otherwise specified (PDD-NOS); a cerebrovascular accident; stroke; a vision impairment; optic neuropathy; dominant inherited juvenile optic atrophy; optic neuropathy caused by a toxic agent; glaucoma; Stargardt's macular dystrophy; diabetic retinopathy; diabetic maculopathy; retinopathy of prematurity; ischemic reperfusion-related retinal injury; oxygen poisoning; a haemoglobinopathy; thalassemia; sickle cell anemia; seizures; ischemia; renal tubular acidosis; attention deficit/hyperactivity disorder (ADHD); a neurodegenerative disorder resulting in hearing or balance impairment; Dominant Optic Atrophy (DOA); Maternally inherited diabetes and deafness (MIDD); chronic fatigue; contrast-induced kidney damage; contrast-induced retinopathy damage; Abetalipoproteinemia; retinitis pigmentosum; Wolfram's disease; Tourette syndrome; cobalamin c defect; methylmalonic aciduria; glioblastoma; Down's syndrome; acute tubular necrosis; a muscular dystrophy; a leukodystrophy; Progressive Supranuclear Palsy; spinal muscular atrophy; hearing

loss; noise induced hearing loss; traumatic brain injury; Juvenile Huntington's Disease; Multiple Sclerosis; NGLY1; Multiple System Atrophy; Adrenoleukodystrophy; and Adrenomyeloneuropathy. In some examples, the oxidative stress disorder is Multiple System Atrophy. In some examples, including any of the foregoing examples, the oxidative stress disorder is cancer. In some examples, including any of the foregoing examples, the oxidative stress disorder is bipolar disorder. In some examples, the oxidative stress disorder is schizophrenia. In some examples, including any of the foregoing examples, the oxidative stress disorder is an age-associated disease. In some examples, including any of the foregoing examples, the oxidative stress disorder is Huntington's Disease. In some examples, including any of the foregoing examples, the oxidative stress disorder is Alzheimer's disease. In some examples, including any of the foregoing examples, the oxidative stress disorder is amyotrophic lateral sclerosis (ALS). In some examples, including any of the foregoing examples, the oxidative stress disorder is epilepsy. In some examples, including any of the foregoing examples, the oxidative stress disorder is Parkinson's disease. In some examples, including any of the foregoing examples, the oxidative stress disorder is seizures. In some examples, including any of the foregoing examples, the oxidative stress disorder is stroke. In some examples, including any of the foregoing examples, the oxidative stress disorder is a mitochondrial disorder. In some examples, including any of the foregoing examples, the oxidative stress disorder is an inherited mitochondrial disease. In some examples, including any of the foregoing examples, the oxidative stress disorder is Friedreich's Ataxia (FA). In some examples, including any of the foregoing examples, the oxidative stress disorder is Kearns-Sayre Syndrome (KSS). In some examples, including any of the foregoing examples, the oxidative stress disorder is Leigh Syndrome or Leigh-like Syndrome. In some examples, including any of the foregoing examples, the oxidative stress disorder is Leber's Hereditary Optic Neuropathy (LHON). In some examples, including any of the foregoing examples, the oxidative stress disorder is Mitochondrial Myopathy, Encephalopathy, Lactacidosis, Stroke (MELAS). In some examples, including any of the foregoing examples, the oxidative stress disorder is Myoclonic Epilepsy with Ragged Red Fibers (MERRF). In some examples, including any of the foregoing examples, the oxidative stress disorder is macular degeneration. In some examples, including any of the foregoing examples, the oxidative stress disorder is brain cancer. In some examples, including any of the foregoing examples, the oxidative stress disorder is autistic disorder. In some examples, including any of the foregoing examples, the oxidative stress disorder is Rett's

disorder. In some examples, including any of the foregoing examples, the oxidative stress disorder is Maternally inherited diabetes and deafness (MIDD). In some examples, including any of the foregoing examples, the oxidative stress disorder is chronic fatigue. In some examples, including any of the foregoing examples, the oxidative stress disorder is contrast-induced kidney damage. In some examples, including any of the foregoing examples, the oxidative stress disorder is contrast-induced retinopathy damage. In some examples, including any of the foregoing examples, the oxidative stress disorder is cobalamin c defect. In some examples, including any of the foregoing examples, the method is a method for modulating one or more energy biomarkers, normalizing one or more energy biomarkers, or enhancing one or more energy biomarkers, wherein the one or more energy biomarkers are selected from the group consisting of: lactic acid (lactate) levels, either in whole blood, plasma, cerebrospinal fluid, or cerebral ventricular fluid; pyruvic acid (pyruvate) levels, either in whole blood, plasma, cerebrospinal fluid, or cerebral ventricular fluid; lactate/pyruvate ratios, either in whole blood, plasma, cerebrospinal fluid, or cerebral ventricular fluid; total, reduced or oxidized glutathione levels, or reduced/oxidized glutathione ratio either in whole blood, plasma, lymphocytes, cerebrospinal fluid, or cerebral ventricular fluid; total, reduced or oxidized cysteine levels, or reduced/oxidized cysteine ratio either in whole blood, plasma, lymphocytes, cerebrospinal fluid, or cerebral ventricular fluid; phosphocreatine levels, NADH (NADH +H⁺) levels; NADPH (NADPH+H⁺) levels; NAD levels; NADP levels; ATP levels; reduced coenzyme Q (CoQ_{red}) levels; oxidized coenzyme Q (CoQ_{ox}) levels; total coenzyme Q (CoQ_{tot}) levels; oxidized cytochrome C levels; reduced cytochrome C levels; oxidized cytochrome C/reduced cytochrome C ratio; acetoacetate levels, β hydroxy butyrate levels, acetoacetate/ β hydroxy butyrate ratio, 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels; levels of reactive oxygen species; levels of oxygen consumption (VO₂); levels of carbon dioxide output (VCO₂); respiratory quotient (VCO₂/VO₂); exercise tolerance; and anaerobic threshold. Energy biomarkers can be measured in whole blood, plasma, cerebrospinal fluid, cerebroventricular fluid, arterial blood, venous blood, or any other body fluid, body gas, or other biological sample useful for such measurement. In some examples, including any of the foregoing examples, the levels are modulated to a value within about 2 standard deviations of the value in a healthy subject. In some examples, including any of the foregoing examples, the levels are modulated to a value within about 1 standard deviation of the value in a healthy subject. In some examples, including any of the foregoing examples, the levels in a subject are changed by at least about 10% above

or below the level in the subject prior to modulation. In some examples, including any of the foregoing examples, the levels are changed by at least about 20% above or below the level in the subject prior to modulation. In some examples, including any of the foregoing examples, the levels are changed by at least about 30% above or below the level in the subject prior to modulation. In some examples, including any of the foregoing examples, the levels are changed by at least about 40% above or below the level in the subject prior to modulation. In some examples, including any of the foregoing examples, the levels are changed by at least about 50% above or below the level in the subject prior to modulation. In some examples, including any of the foregoing examples, the levels are changed by at least about 75% above or below the level in the subject prior to modulation. In some examples, including any of the foregoing examples, the levels are changed by at least about 100% above or at least about 90% below the level in the subject prior to modulation. In some examples, including any of the foregoing examples, the subject or subjects in which a method of treating or suppressing an oxidative stress disorder, modulating one or more energy biomarkers, normalizing one or more energy biomarkers, or enhancing one or more energy biomarkers is performed is/are selected from the group consisting of subjects undergoing strenuous or prolonged physical activity; subjects with chronic energy problems; subjects with chronic respiratory problems; pregnant females; pregnant females in labor; neonates; premature neonates; subjects exposed to extreme environments; subjects exposed to hot environments; subjects exposed to cold environments; subjects exposed to environments with lower-than-average oxygen content; subjects exposed to environments with higher-than-average carbon dioxide content; subjects exposed to environments with higher-than-average levels of air pollution; airline travelers; flight attendants; subjects at elevated altitudes; subjects living in cities with lower-than-average air quality; subjects working in enclosed environments where air quality is degraded; subjects with lung diseases; subjects with lower-than-average lung capacity; tubercular patients; lung cancer patients; emphysema patients; cystic fibrosis patients; subjects recovering from surgery; subjects recovering from illness; elderly subjects; elderly subjects experiencing decreased energy; subjects suffering from chronic fatigue; subjects suffering from chronic fatigue syndrome; subjects undergoing acute trauma; subjects in shock; subjects requiring acute oxygen administration; subjects requiring chronic oxygen administration; subjects requiring organ visualization via contrast solution; or other subjects with acute, chronic, or ongoing energy demands who can benefit from enhancement of energy biomarkers.

[0045] In another example, there is described the use of a polymorphic or amorphous form of an anhydrate, a hydrate, or a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide as described herein, including any of the foregoing or hereafter described examples, for treating or suppressing an oxidative stress disorder. In some examples, the form is polymorph Form I. In some examples, the form is polymorph Form II. In some examples, the form is polymorph Form III. In some examples, the form is polymorph Form IV. In some examples, the form is polymorph Form V. In some examples, the form is polymorph Form VI. In some examples, the form is amorphous. In another example, there is described the use of a polymorphic or amorphous form of an anhydrate, a hydrate, or a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide as described herein, including any of the foregoing or hereafter described examples, in the manufacture of a medicament for use in treating or suppressing an oxidative stress disorder. In some examples, the form is polymorph Form I. In some examples, the form is polymorph Form II. In some examples, the form is polymorph Form III. In some examples, the form is polymorph Form IV. In some examples, the form is polymorph Form V. In some examples, the form is polymorph Form VI. In some examples, the form is amorphous.

[0046] For all compositions described herein, and all methods using a composition described herein, the compositions can either comprise the listed components or steps, or can “consist essentially of” the listed components or steps. When a composition is described as “consisting essentially of” the listed components, the composition contains the components listed, and may contain other components which do not substantially affect the condition being treated, but do not contain any other components which substantially affect the condition being treated other than those components expressly listed; or, if the composition does contain extra components other than those listed which substantially affect the condition being treated, the composition does not contain a sufficient concentration or amount of the extra components to substantially affect the condition being treated. When a method is described as “consisting essentially of” the listed steps, the method contains the steps listed, and may contain other steps that do not substantially affect the condition being treated, but the method does not contain any other steps which substantially affect the condition being treated other than those steps expressly listed. As a non-limiting specific example, when a composition is described as ‘consisting essentially of’ a component, the composition may additionally contain any amount of pharmaceutically acceptable carriers, vehicles, or diluents and other such components which do not substantially affect the condition being treated.

[0047] The present disclosure refers to a process for the preparation of a polymorph of anhydrous (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the process comprises the steps: (a) contacting (R)-2-hydroxy-2-methyl-

4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide with a liquid comprising IPA; and (b) removing the liquid. In another example, there is described a process for the preparation of polymorph Form I of an anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the process comprises the steps: (a) contacting (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide with a liquid comprising IPA; and (b) removing the liquid. In some examples, step (a) comprises dissolving the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in the liquid. In some examples, step (a) comprises slurrying the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in the liquid. In some examples, the slurrying in step (a) may be performed for at least about 24 hours. In some examples, including any of the foregoing examples, the liquid is 100% IPA. In some examples, including any of the foregoing examples, the liquid is 98% IPA/2% water (v/v). In some examples, including any of the foregoing examples, the process further comprises step (a)(i): adding heptane to the liquid. In some examples, including any of the foregoing embodiments, step (b) comprises filtering the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide. In some examples, including any of the foregoing examples, the mixture in step (a) or step (a)(i) may be seeded with Form I crystals. In some examples, including any of the foregoing examples, the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in step (a) is at least about 95% pure. In various examples, including any of the foregoing examples, the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in step (a) is at least about 96%, at least about 97%, at least about 98%, at least about 99%, at least about 99.5%, at least about 99.9% pure. In another example, there is described an anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide prepared by the above described process.

In another example, there is described a process for the preparation of polymorph Form II of an anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the process comprises the steps: (a) dissolving the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in EtOAc, (b) rapidly cooling the mixture from (a), and (c) isolating the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide. In some examples, the initial (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide is Form I. In some examples, including any of the foregoing examples, step (a) is at about 60°C. In some examples, including any of the foregoing examples, step (b) comprises rapidly cooling the mixture in an ice bath. In some examples, including any of the foregoing examples, the mixture in step (b) may be seeded with Form II

crystals. In another example, there is described an anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide prepared by the above described process.

[0048] In another example, there is described a process for the preparation of polymorph Form III of a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the process comprises the steps: (a) combining (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and 0.5% MC/2% Tween 80 in water to create a slurry; (b) slurrying the mixture from (a), and (c) removing the 0.5% MC/2% Tween 80 in water. As used herein “MC” refers to methyl cellulose and “Tween 80” refers to a commercially available polysorbate nonionic surfactant. In some examples, the initial (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide is Form I. In some examples, including any of the foregoing examples, step (b) is performed for at least about 24 hours. In some examples, including any of the foregoing examples, step (b) is at room temperature. In some examples, including any of the foregoing examples, the mixture in step (b) may be seeded with Form III crystals. In another example, there is described a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide prepared by the above described process.

[0049] In another example, there is described a process for the preparation of polymorph Form III of a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the process comprises the steps: (a) combining (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and 0.5% MC in water to create a slurry; (b) slurrying the mixture from (a), and (c) removing the 0.5% MC in water. In some examples, the initial (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide is Form I, II, IV, V or VI. In some examples, including any of the foregoing examples, step (b) is performed at room temperature. In some examples, including any of the foregoing examples, step (b) may be performed for at least about 7 days. In some examples, including any of the foregoing examples, the mixture in step (b) may be seeded with Form III crystals. In another example, there is described a solvate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide prepared by the above described process.

The present disclosure refers to a polymorph produced by the process comprising the steps: (a) contacting (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide with a liquid comprising IPA; and (b) removing the liquid.

BRIEF DESCRIPTION OF THE FIGURES

[0050] Figure 1 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide short term slurry experiments, a) Pattern C, from slurry in 0.5% Methyl Cellulose/2% Tween 80, b) Pattern B, from slurry in tetrahydrofuran (THF), and c) starting material, Pattern A.

[0052] Figure 2 shows a XRPD overlay of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide sample lots, a) starting material, Pattern A, and b) Pattern D following evaporative crystallization from 2-methyltetrahydrofuran (2-MeTHF).

[0053] Figure 3 shows a XRPD overlay of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide crystallization experiments, a) starting material, Pattern A, b) Pattern D following evaporative crystallization from 2-MeTHF (Example 4), c) from evaporative crystallization in 2-MeTHF (Example 5, fast cooling), and d) from evaporative crystallization in 2-MeTHF (Example 5, slow cooling).

[0054] Figure 4 shows a XRPD stack plot of all (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide forms, a) starting material, Pattern A, b) Pattern B, c) Pattern C, d) Pattern D, e) Pattern E from fast cooling crystallization in ethyl acetate (EtOAc), and f) Pattern F, from fast cooling crystallization in 2-MeTHF.

[0055] Figure 5 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide forms, a) Pattern E from fast cooling crystallization in EtOAc (Example 5), and b) from scale-up fast cooling crystallization in EtOAc (Example 6).

[0056] Figure 6 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide forms, a) Pattern B from slurry in THF (Example 3), and b) from scale-up slurry in THF (Example 6).

[0057] Figure 7 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide forms, a) Pattern C from slurry in 0.5% Methyl Cellulose/2% Tween 80 (Example 3), and b) from scale-up slurry in 0.5% Methyl Cellulose/2% Tween 80 (Example 6).

[0058] Figure 8 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide samples, a) following 24 hours of competitive slurry in 0.5% Methyl Cellulose in water, b) starting material, Pattern A, c) Pattern C, and d) following 7 days of competitive slurry in 0.5% Methyl Cellulose in water.

[0059] Figure 9 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide samples, a) following 24 hours of competitive slurry in IPA, b) following 7 days of competitive slurry in IPA, c) starting material, Pattern A, d) following 24 hours of competitive slurry in isopropanol (IPA)/2% water, e) following 7 days of competitive slurry in IPA/2% water.

- [0060] Figure 10 shows a XRPD of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide starting material, Pattern A.
- [0061] Figure 11 shows an optical microscopy image of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide starting material, Pattern A.
- [0062] Figure 12 shows a DSC thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide starting material, Pattern A.
- [0063] Figure 13 shows a TGA thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide starting material, Pattern A.
- [0064] Figure 14 shows a moisture sorption-desorption plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide starting material, Pattern A.
- [0065] Figure 15 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern E isolated from single solvent fast cooling experiments, a) from 50 mg scale and b) from 300 mg scale-up.
- [0066] Figure 16 shows an optical microscopy image of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern E.
- [0067] Figure 17 shows a DSC thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern E.
- [0068] Figure 18 shows a TGA thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern E.
- [0069] Figure 19 shows a moisture sorption-desorption plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern E.
- [0070] Figure 20 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern C from slurry in 0.5% Methyl Cellulose/2% Tween 80, a) from 50 mg scale and b) from 300 mg scale-up.
- [0071] Figure 21 shows an optical microscopy image of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern C.
- [0072] Figure 22 shows a DSC thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern C.
- [0073] Figure 23 shows a TGA thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern C.
- [0074] Figure 24 shows a moisture sorption-desorption plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern C.

[0075] Figure 25 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern B from slurry in THF, a) from 50 mg scale and b) from 300 mg scale-up.

[0076] Figure 26 shows an optical microscopy image of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern B.

[0077] Figure 27 shows a DSC thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern B.

[0078] Figure 28 shows a TGA thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern B.

[0079] Figure 29 shows a moisture sorption-desorption plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern B.

[0080] Figure 30 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern D from evaporative crystallization in 2-MeTHF, a) from Example 4, b) Example 5 (fast cooling), c) Example 5 (slow cooling) and d) starting material Pattern A.

[0081] Figure 31 shows a DSC thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern D.

[0082] Figure 32 shows a TGA thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern D.

[0083] Figure 33 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern F from single solvent crystallization in 2-MeTHF, a) from Example 6, b) following moisture sorption analysis (starting with Pattern F from Example 6) and c) Pattern E.

[0084] Figure 34 shows an optical microscopy image of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern F.

[0085] Figure 35 shows a DSC thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern F.

[0086] Figure 36 shows a TGA thermogram of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern F.

[0087] Figure 37 shows a moisture sorption-desorption plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern F.

[0088] Figure 38 shows (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Polymorphic Form-Interrelations.

[0089] Figure 39 shows a XRPD stack plot of Edison (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Polymorphic Forms.

[0090] Figure 40 shows a ^1H NMR spectrum of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide starting material, Pattern A

[0091] Figure 41 shows a ^1H NMR spectrum of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern B.

[0092] Figure 42 shows a ^1H NMR spectrum of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Pattern C

[0093] Figure 43 shows a ^1H NMR spectrum of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern D.

[0094] Figure 44 shows a ^1H NMR spectrum of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern E

[0095] Figure 45 shows a ^1H NMR spectrum of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, Pattern F.

DETAILED DESCRIPTION

[0096] The invention embraces polymorphic and amorphous forms of anhydrides, hydrates, and solvates of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide useful in treating or suppressing diseases, developmental delays and symptoms related to oxidative stress such as mitochondrial disorders, impaired energy processing disorders, neurodegenerative diseases and diseases of aging, and methods of using such compositions for treating or suppressing an oxidative stress disorder, or for modulating, normalizing, or enhancing one or more (e.g. one, two, three, or more) energy biomarkers. The invention further embraces methods for producing such polymorphic and amorphous forms.

[0097] The abbreviations used herein have their conventional meaning within the chemical and biological arts, unless otherwise specified.

[0098] Reference to “about” a value or parameter herein includes (and describes) variations that are directed to that value or parameter per se. For example, description referring to “about X” includes description of “X”.

[0099] The terms “a” or “an,” as used in herein means one or more, unless context clearly dictates otherwise.

[00100] By “subject,” “individual,” or “patient” is meant an individual organism, preferably a vertebrate, more preferably a mammal, most preferably a human.

[00101] (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide may exist in anhydrate, hydrate, and solvate forms. Unless otherwise specified or clear from context, as used herein the term “(R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide” encompasses anhydrate, hydrate, and solvate forms of the compound.

[00102] The term “substantially as shown in” when referring, for example, to an XRPD pattern, a DSC thermogram, or a TGA graph includes a pattern, thermogram or graph that is not necessarily identical to those depicted herein, but that falls within the limits of experimental error or deviations when considered by one of ordinary skill in the art. For example, in an XRPD pattern, the relative intensity of the peaks in the diffraction pattern can vary, e.g. due to sample preparation conditions. In addition, changes in temperature (when generating the XRPD data) can affect the shape and location of peaks. The XRPD patterns given herein were generated at room temperature (~25° C). In some embodiments, the XRPD pattern is generated at about 15° C to about 30° C. In some embodiments, the XRPD pattern is generated at about 20° C to about 30° C. In some embodiments, the XRPD pattern is generated at about 23° C to about 27° C. In some embodiments, the XRPD pattern is generated at about 24° C to about 26° C. In some embodiments, the XRPD pattern is generated at about 25° C.

[00103] Similarly, when describing a polymorph by characteristic peaks (e.g. angular positions of the peaks), it is to be understood that the location of the peaks may vary depending on sample preparation, temperature, etc. The characteristic XRPD peaks given herein were generated at room temperature (~25° C). In some embodiments, the XRPD data is generated at about 15° C to about 30° C. In some embodiments, the XRPD data is generated at about 20° C to about 30° C. In some embodiments, the XRPD data is generated at about 23° C to about 27° C. In some embodiments, the XRPD data is generated at about 24° C to about 26° C. In some embodiments, the XRPD data is generated at about 25° C.

[00104] A polymorph or amorphous form that is “isolated” is used herein to refer to a form that is at least 90% that particular form (i.e. less than 10% of the material is comprises of other forms

or other compounds, including but not limited to (S)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

[00105] A polymorph composition that is “essentially free of” a particular component(s) indicates that the composition contains less than about 5% of the particular component(s). As a non-limiting example, a polymorph composition that is essentially free of polymorph Form II indicates a composition that contains less than about 5% of Form II. In some embodiments, “essentially free of” indicates that the composition contains less than about 4%, less than about 3%, less than about 2%, or less than about 1% of the particular component(s), or wherein the particular component(s) are not present within the limit of detection.

[00106] “Angular positions” indicates Angle, 2 theta.

[0100] “Treating” a disorder with the compounds, compositions, and methods discussed herein is defined as administering one or more of the compounds or compositions discussed herein, with or without additional therapeutic agents, in order to reduce or eliminate either the disorder or one or more symptoms of the disorder, or to retard the progression of the disorder or of one or more symptoms of the disorder, or to reduce the severity of the disorder or of one or more symptoms of the disorder. “Suppression” of a disorder with the compounds, compositions, and methods discussed herein is defined as administering one or more of the compounds or compositions discussed herein, with or without additional therapeutic agents, in order to suppress the clinical manifestation of the disorder, or to suppress the manifestation of adverse symptoms of the disorder. The distinction between treatment and suppression is that treatment occurs after adverse symptoms of the disorder are manifest in a subject, while suppression occurs before adverse symptoms of the disorder are manifest in a subject. Suppression may be partial, substantially total, or total. Because some of the disorders are inherited, genetic screening can be used to identify patients at risk of the disorder. The compounds, compositions, and methods of the invention can then be administered to asymptomatic patients at risk of developing the clinical symptoms of the disorder, in order to suppress the appearance of any adverse symptoms.

[0101] “Therapeutic use” of the compounds and compositions discussed herein is defined as using one or more of the compounds or compositions discussed herein to treat or suppress a disorder, as defined above. An “effective amount” of a compound or composition is an amount of the compound or composition sufficient to modulate, normalize, or enhance one or more energy biomarkers (where modulation, normalization, and enhancement are defined

below). A “therapeutically effective amount” of a compound or composition is an amount of the compound or composition, which, when administered to a subject, is sufficient to reduce or eliminate either a disorder or one or more symptoms of a disorder, or to retard the progression of a disorder or of one or more symptoms of a disorder, or to reduce the severity of a disorder or of one or more symptoms of a disorder, or to suppress the clinical manifestation of a disorder, or to suppress the manifestation of adverse symptoms of a disorder. A therapeutically effective amount can be given in one or more administrations. An “effective amount” of a compound or composition embraces both a therapeutically effective amount, as well as an amount effective to modulate, normalize, or enhance one or more energy biomarkers in a subject.

[0102] “Modulation” of, or to “modulate,” an energy biomarker means to change the level of the energy biomarker towards a desired value, or to change the level of the energy biomarker in a desired direction (e.g., increase or decrease). Modulation can include, but is not limited to, normalization and enhancement as defined below.

[0103] “Normalization” of, or to “normalize,” an energy biomarker is defined as changing the level of the energy biomarker from a pathological value towards a normal value, where the normal value of the energy biomarker can be 1) the level of the energy biomarker in a healthy person or subject, or 2) a level of the energy biomarker that alleviates one or more undesirable symptoms in the person or subject. That is, to normalize an energy biomarker which is depressed in a disease state means to increase the level of the energy biomarker towards the normal (healthy) value or towards a value which alleviates an undesirable symptom; to normalize an energy biomarker which is elevated in a disease state means to decrease the level of the energy biomarker towards the normal (healthy) value or towards a value which alleviates an undesirable symptom.

[0104] “Enhancement” of, or to “enhance,” energy biomarkers means to intentionally change the level of one or more energy biomarkers away from either the normal value, or the value before enhancement, in order to achieve a beneficial or desired effect. For example, in a situation where significant energy demands are placed on a subject, it may be desirable to increase the level of ATP in that subject to a level above the normal level of ATP in that subject. Enhancement can also be of beneficial effect in a subject suffering from a disease or pathology such as e.g. a mitochondrial disorder, in that normalizing an energy biomarker may not achieve the optimum outcome for the subject; in such cases, enhancement of one or more

energy biomarkers can be beneficial, for example, higher-than-normal levels of ATP, or lower-than-normal levels of lactic acid (lactate) can be beneficial to such a subject.

[0105] By modulating, normalizing, or enhancing the energy biomarker Coenzyme Q is meant modulating, normalizing, or enhancing the variant or variants of Coenzyme Q which is predominant in the species of interest. For example, the variant of Coenzyme Q which predominates in humans is Coenzyme Q10. If a species or subject has more than one variant of Coenzyme Q present in significant amounts (i.e., present in amounts which, when modulated, normalized, or enhanced, can have a beneficial effect on the species or subject), modulating, normalizing, or enhancing Coenzyme Q can refer to modulating, normalizing or enhancing any or all variants of Coenzyme Q present in the species or subject.

[0106] By “respiratory chain disorder” is meant a disorder which results in the decreased utilization of oxygen by a mitochondrion, cell, tissue, or individual, due to a defect or disorder in a protein or other component contained in the mitochondrial respiratory chain. By “protein or other component contained in the mitochondrial respiratory chain” is meant the components (including, but not limited to, proteins, tetrapyrroles, and cytochromes) comprising mitochondrial complex I, II, III, IV, and/or V. “Respiratory chain protein” refers to the protein components of those complexes, and “respiratory chain protein disorder” is meant a disorder which results in the decreased utilization of oxygen by a mitochondrion, cell, tissue, or individual, due to a defect or disorder in a protein contained in the mitochondrial respiratory chain.

[0107] The terms “Parkinson’s”, (also called “Parkinsonism” and “Parkinsonian syndrome”) (“PD”) is intended to include not only Parkinson’s disease but also drug-induced Parkinsonism and post-encephalitic Parkinsonism. Parkinson’s disease is also known as paralysis agitans or shaking palsy. It is characterized by tremor, muscular rigidity and loss of postural reflexes. The disease usually progresses slowly with intervals of 10 to 20 years elapsing before the symptoms cause incapacity. Due to their mimicry of effects of Parkinson’s disease, treatment of animals with methamphetamine or MPTP has been used to generate models for Parkinson’s disease. These animal models have been used to evaluate the efficacy of various therapies for Parkinson’s disease.

[0108] The term “Friedreich’s ataxia” is intended to embrace other related ataxias, and is also sometimes referred to as hereditary ataxia, familial ataxia, or Friedreich’s tabes.


[0109] The term “ataxia” is an aspecific clinical manifestation implying dysfunction of parts of the nervous system that coordinate movement, such as the cerebellum. People with ataxia have problems with coordination because parts of the nervous system that control movement and balance are affected. Ataxia may affect the fingers, hands, arms, legs, body, speech, and eye movements. The word ataxia is often used to describe a symptom of incoordination which can be associated with infections, injuries, other diseases, or degenerative changes in the central nervous system. Ataxia is also used to denote a group of specific degenerative diseases of the nervous system called the hereditary and sporadic ataxias. Ataxias are also often associated with hearing impairments.

[0110] There are three types of ataxia, cerebellar ataxia, including vestibulo-cerebellar dysfunction, spino-cerebellar dysfunction, and cerebro-cerebellar dysfunction; sensory ataxia; and vestibular ataxia. Examples of the diseases which are classifiable into spino-cerebellar ataxia or multiple system atrophy are hereditary olivo-ponto-cerebellar atrophy, hereditary cerebellar cortical atrophy, Friedreich’s ataxia, Machado-Joseph diseases, Ramsay Hunt syndrome, hereditary dentatorubral-pallidoluyasian atrophy, hereditary spastic paraplegia, Shy-Drager syndrome, cortical cerebellar atrophy, striato-nigral degeneration, Marinesco-Sjogren syndrome, alcoholic cortical cerebellar atrophy, paraneoplastic cerebellar atrophy associated with malignant tumor, toxic cerebellar atrophy caused by toxic substances, Vitamin E deficiency due to mutation of a Tocopherol transfer protein (aTTP) or lipid absorption disorder such as Abetalipoproteinemia, cerebellar atrophy associated with endocrine disturbance and the like.

[0111] Examples of ataxia symptoms are motor ataxia, trunk ataxia, limb ataxia and the like, autonomic disturbance such as orthostatic hypotension, dysuria, hypohidrosis, sleep apnea, orthostatic syncope and the like, stiffness of lower extremity, ocular nystagmus, oculomotor nerve disorder, pyramidal tract dysfunction, extrapyramidal symptoms (postural adjustment dysfunction, muscular rigidity, akinesia, tremors), dysphagia, lingual atrophy, posterior funiculus symptom, muscle atrophy, muscle weakness, deep hyperreflexia, sensory disturbance, scoliosis, kyphoscoliosis, foot deformities, anarthria, dementia, manic state, decreased motivation for rehabilitation and the like.

Polymorphic and Amorphous Forms of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide

[0112] Provided herein are various crystalline and amorphous forms of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide:



(R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamide

[0113] and methods for producing such forms, and methods for using such forms.

[0114] Table 1 below provides a summary of certain polymorphic forms of the invention of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

Table 1: Summary of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide Polymorphic Form Characterization

XRPD [Pattern] (Conditions)	Form Designation	DSC [°C]	TGA [% Wt. loss]	¹ H NMR [Residual Solvent]	KF [Wt% water]	Optical Microscopy
Crystalline [Pattern A] (Starting material)	Form I (Anhydrate)	152.9	0.0	Consistent with structure [0.28 wt% IPA]	0.1	Birefringent
Crystalline [Pattern B] (RT Slurry in THF)	Form IV (THF Solvate)	70.5, 89.1, 149.7	4.7	Consistent with Structure [6.9 wt% THF]	0.3	Birefringent
Crystalline [Pattern C] (RT Slurry in 0.5%MC/ 2% Tween 80)	Form III (Hydrate)	72.0, 150.7	2.5, 2.3	Consistent with structure	4.3	Birefringent
Crystalline [Pattern D] (Evap Cryst. In 2- MeTHF)	Form V (2-MeTHF Solvate)	67.2, 92.2, 150.6	2.7, 5.3	Consistent with Structure [6.1 wt% 2- MeTHF]	-	Birefringent
Crystalline [Pattern E] (Single Solvent fast cooling cryst. in EtOAc)	Form II (Anhydrate)	133.9, 151.3	0.4	Consistent with Structure [0.4 wt.% EtOAc]	0.1	Birefringent
Crystalline [Pattern F] (Scale up, Single Solvent fast cooling cryst. in 2-MeTHF)	Form VI (2-MeTHF Solvate)	93.2, 135.2, 151.0	1.1, 0.2	Consistent with Structure [3.9 wt.% 2- MeTHF]	0.1	Birefringent

[0115] Figure 38 provides a chart showing interrelations between the various polymorphic forms for (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

Figure 39 shows a XRPD stack plot of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide polymorphic forms.

[0116] Table A provides various embodiments of angular positions of certain characteristic peaks in powder x-ray diffraction for the polymorphic forms of the invention. In some embodiments, the angular positions may vary by ± 0.2 . In some embodiments, the angular positions may vary by ± 0.1 . In some embodiments, the angular positions may vary by ± 0.05 . In some embodiments, the angular positions may vary by ± 0.02 .

Table A. Angular Positions of Certain Characteristic Peaks in Powder X-ray Diffraction for Forms I-VI

Form	Angular Positions of Certain Characteristic Peaks (Angle, 2 theta) (all ± 0.2)
I	Embodiment 1. 12.06, 17.03, 17.26 Embodiment 2. 12.06, 15.33, 17.03, 17.26, 18.72 Embodiment 3. 7.67, 10.75, 12.06, 15.33, 16.41, 17.03, 17.26, 18.72, 20.04, 23.92 Embodiment 4. 7.67, 10.75, 12.06, 15.33, 16.41, 17.03, 17.26, 18.72, 20.04, 20.64, 20.91, 21.14, 22.58, 23.13, 23.92, 24.19, 24.53, 27.21, 27.56 Embodiment 5. 5.48, 7.67, 10.75, 12.06, 15.33, 16.41, 17.03, 17.26, 17.71, 17.94, 18.40, 18.72, 19.51, 20.04, 20.64, 20.91, 21.14, 21.55, 21.91, 22.25, 22.58, 23.13, 23.41, 23.92, 24.19, 24.53, 25.64, 26.13, 26.34, 27.21, 27.56, 28.01, 29.04, 29.46 Embodiment 6. 12.06, 15.33, 17.03, 17.26. Embodiment 7. 12.06, 15.33, 17.03, 17.26, 18.72, 23.92. Embodiment 8. 12.06, 15.33, 17.03, 17.26, 18.72, 23.92, 16.41 Embodiment 9. 12.06, 15.33, 17.03, 17.26, 18.72, 23.92, 16.41, 10.75 Embodiment 10. 12.06, 15.33, 17.03, 17.26, 18.72, 23.92, 16.41, 10.75, 20.64
V	Embodiment 1. 9.61, 11.49, 15.45 Embodiment 2. 9.61, 11.49, 12.93, 15.45, 23.96, 26.05 Embodiment 3. 9.61, 11.49, 12.93, 14.80, 15.45, 16.53, 23.96, 24.54, 26.05 Embodiment 4. 9.61, 11.49, 12.93, 14.80, 15.45, 16.10, 16.34, 16.53, 20.18, 22.52, 22.86, 23.96, 24.54, 26.05 Embodiment 5. 6.91, 7.72, 9.61, 11.49, 11.86, 12.93, 13.19, 13.87, 14.80, 15.45,

	16.10, 16.34, 16.53, 17.14, 17.85, 19.12, 19.85, 20.18, 21.00, 22.06, 22.52, 22.86, 23.09, 23.96, 24.54, 25.26, 26.05, 26.90 Embodiment 6. 9.61, 11.49, 12.93, 15.45 Embodiment 7. 9.61, 11.49, 12.93, 15.45, 23.96 Embodiment 8. 9.61, 11.49, 12.93, 15.45, 14.80 Embodiment 9. 9.61, 11.49, 12.93, 15.45, 7.72 Embodiment 10. 9.61, 11.49, 12.93, 15.45, 7.72, 16.53
III	Embodiment 1. 14.02, 15.23, 21.10 Embodiment 2. 9.16, 14.02, 15.23, 21.10, 22.69 Embodiment 3. 9.16, 11.81, 13.74, 14.02, 15.23, 21.10, 22.69, 23.90 Embodiment 4. 9.16, 11.81, 13.74, 14.02, 15.23, 17.35, 21.10, 22.69, 23.15, 23.90, 26.10 Embodiment 5. 9.16, 11.53, 11.81, 12.68, 12.93, 13.74, 14.02, 15.23, 16.53, 17.35, 17.98, 18.54, 19.09, 20.23, 21.10, 21.93, 22.69, 23.15, 23.50, 23.90, 24.65, 25.09, 25.46, 25.79, 26.10, 27.79, 28.22, 28.93, 29.33 Embodiment 6. 9.16, 14.02, 15.23, 21.10 Embodiment 7. 9.16, 13.74, 14.02, 15.23, 21.10 Embodiment 8. 9.16, 11.81, 13.74, 14.02, 15.23, 21.10 Embodiment 9. 9.16, 11.81, 13.74, 14.02, 15.23, 21.10, 23.90 Embodiment 10. 9.16, 11.81, 13.74, 14.02, 15.23, 21.10, 22.69, 23.90
II	Embodiment 1. 9.63, 11.33, 19.33 Embodiment 2. 9.63, 10.85, 11.33, 13.47, 19.33 Embodiment 3. 5.76, 8.04, 9.63, 10.85, 11.33, 11.97, 13.47, 14.75, 17.37, 17.71, 19.33 Embodiment 4. 5.76, 8.04, 9.63, 10.85, 11.33, 11.97, 13.47, 14.75, 16.42, 16.89, 17.37, 17.71, 19.33, 22.89, 24.59 Embodiment 5. 5.76, 6.72, 7.57, 8.04, 9.63, 10.85, 11.33, 11.97, 12.38, 13.13, 13.47, 14.75, 15.28, 16.42, 16.89, 17.37, 17.71, 18.17, 18.66, 19.33, 20.01, 20.29, 20.67, 20.90, 21.36, 21.54, 21.80, 22.55, 22.89, 23.27, 23.54, 23.87, 24.35, 24.59, 24.87, 25.29, 25.55, 25.89, 26.44, 27.49, 28.01, 28.39, 29.17 Embodiment 6. 9.63, 11.33, 10.85 Embodiment 7. 9.63, 11.33, 10.85, 19.33

	Embodiment 8. 9.63, 11.33, 10.85, 19.33, 17.37 Embodiment 9. 9.63, 11.33, 10.85, 19.33, 17.37, 13.47 Embodiment 10. 9.63, 11.33, 10.85, 19.33, 17.37, 13.47, 11.97
IV	Embodiment 1. 4.31, 12.97, 13.20 Embodiment 2. 4.31, 8.76, 12.97, 13.20, 16.66 Embodiment 3. 4.31, 7.94, 8.76, 12.97, 13.20, 16.66, 17.33, 20.57 Embodiment 4. 4.31, 7.94, 8.76, 12.97, 13.20, 15.08, 16.66, 17.33, 19.09, 20.57, 21.58 Embodiment 5. 4.31, 5.77, 6.28, 7.53, 7.94, 8.76, 9.39, 9.87, 10.54, 11.07, 11.68, 12.02, 12.28, 12.97, 13.20, 13.52, 14.40, 15.08, 15.90, 16.66, 16.96, 17.33, 17.59, 18.77, 19.09, 19.74, 20.27, 20.57, 21.09, 21.58, 22.81, 23.23, 24.01, 24.65, 25.60 Embodiment 6. 12.97, 13.20, 8.76 Embodiment 7. 12.97, 13.20, 8.76, 16.66 Embodiment 8. 12.97, 13.20, 8.76, 16.66, 4.31 Embodiment 9. 12.97, 13.20, 8.76, 16.66, 4.31, 17.33 Embodiment 10. 12.97, 13.20, 8.76, 16.66, 4.31, 17.33, 20.57
VI	Embodiment 1. 6.27, 9.91, 12.94 Embodiment 2. 6.27, 9.41, 9.91, 12.94, 13.29 Embodiment 3. 6.27, 8.85, 9.41, 9.91, 12.94, 13.29, 16.67, 19.13 Embodiment 4. 4.39, 6.27, 8.85, 9.41, 9.91, 11.32, 12.94, 13.29, 14.03, 16.67, 19.13, 20.76, 22.06 Embodiment 5. 4.39, 6.27, 7.00, 8.62, 8.85, 9.41, 9.91, 11.32, 11.50, 12.25, 12.56, 12.94, 13.29, 14.03, 14.82, 15.10, 15.44, 15.71, 16.01, 16.67, 16.91, 17.33, 17.59, 18.33, 18.75, 19.13, 20.25, 20.76, 21.68, 22.06, 22.27, 22.61, 22.94, 24.01, 24.33, 24.65, 25.48, 26.05, 28.63, 29.18 Embodiment 6. 6.27, 9.91, 12.94, 15.71 Embodiment 7. 6.27, 9.91, 12.94, 15.71, 19.13 Embodiment 8. 6.27, 9.91, 12.94, 15.71, 16.91, 19.13 Embodiment 9. 6.27, 9.41, 9.91, 12.94, 15.71, 16.91, 19.13 Embodiment 10. 6.27, 8.85, 9.41, 9.91, 12.94, 15.71, 16.91, 19.13

[0117] Tables 2-7 provide additional embodiments of angular positions of characteristic peaks in powder x-ray diffraction for the polymorphic forms of the invention. In some embodiments, a polymorphic form is characterized by the angular positions of characteristics peaks shown in Table A. In some embodiments, the polymorphic form is characterized by 3 or more (e.g. 4, 5, 6, 7, 8, 9, 10, or more than 10) angular positions of characteristic peaks in powder x-ray diffraction as shown in Tables 2-7 below. In some embodiments, the angular positions may vary by ± 0.2 . In some embodiments, the angular positions may vary by ± 0.1 . In some embodiments, the angular positions may vary by ± 0.05 . In some embodiments, the angular positions may vary by ± 0.02 .

[0118] As a non-limiting example, polymorph Form I may be characterized by 3, 4, 5, 6, 7, 8, 9, 10, or more angular positions as shown in Table 2.

Table 2. Angular Positions of Characteristic Peaks in Powder X-ray Diffraction for Pattern A (Anhydrate Form I)

[0119] Polymorph Form I may be characterized by 3, 4, 5, 6, 7, 8, 9, 10, or more angular positions as shown below:

Angle, 2 theta	d spacing, Å		Angle, 2 theta	d spacing, Å
5.48	16.1		21.55	4.1
7.67	11.5		21.91	4.1
10.75	8.2		22.25	4.0
12.06	7.3		22.58	3.9
15.33	5.8		23.13	3.8
16.41	5.4		23.41	3.8
17.03	5.2		23.92	3.7
17.26	5.1		24.19	3.7
17.71	5.0		24.53	3.6
17.94	4.9		25.64	3.5
18.40	4.8		26.13	3.4
18.72	4.7		26.34	3.4
19.51	4.5		27.21	3.3
20.04	4.4		27.56	3.2
20.64	4.3		28.01	3.2
20.91	4.2		29.04	3.1
21.14	4.2		29.46	3.0

[0120] In some examples, polymorph Form I is characterized by at least 3 or more angular positions. In certain examples, these angular positions include 12.06, 17.03, and 17.26 ± 0.2 .

In some examples, polymorph Form I is characterized by at least 4 or more angular positions. In certain examples, these at least four angular positions include 12.1, 17.0, 17.3, and 15.33 ± 0.2 . In certain examples, these at least four angular positions include 12.1, 17.0, 17.3, 15.33, and 18.72 ± 0.2 .

Table 3. Angular Positions of Characteristic Peaks in Powder X-ray Diffraction for Pattern D (2-MeTHF Solvate Form V)

[0121] Polymorph Form V may be characterized by 3, 4, 5, 6, 7, 8, 9, 10, or more angular positions as shown below:

Angle, 2 theta	d spacing, Å		Angle, 2 theta	d spacing, Å
6.91	12.8		17.85	5.0
7.72	11.4		19.12	4.6
9.61	9.2		19.85	4.5
11.49	7.7		20.18	4.4
11.86	7.5		21.00	4.2
12.93	6.8		22.06	4.0
13.19	6.7		22.52	3.9
13.87	6.4		22.86	3.9
14.80	6.0		23.09	3.8
15.45	5.7		23.96	3.7
16.10	5.5		24.54	3.6
16.34	5.4		25.26	3.5
16.53	5.4		26.05	3.4
17.14	5.2		26.90	3.3

Table 4. Angular Positions of Characteristic Peaks in Powder X-ray Diffraction for Pattern C (Hydrate Form III)

[0122] Polymorph Form III may be characterized by 3, 4, 5, 6, 7, 8, 9, 10, or more angular positions as shown below:

Angle, 2 theta	d spacing, Å		Angle, 2 theta	d spacing, Å
9.16	9.6		21.93	4.0
11.53	7.7		22.69	3.9
11.81	7.5		23.15	3.8
12.68	7.0		23.50	3.8
12.93	6.8		23.90	3.7
13.74	6.4		24.65	3.6
14.02	6.3		25.09	3.5

15.23	5.8		25.46	3.5
16.53	5.4		25.79	3.5
17.35	5.1		26.10	3.4
17.98	4.9		27.79	3.2
18.54	4.8		28.22	3.2
19.09	4.6		28.93	3.1
20.23	4.4		29.33	3.0
21.10	4.2			

Table 5. Angular Positions of Characteristic Peaks in Powder X-ray Diffraction for Pattern E (Anhydrate II)

[0123] Polymorph Form II may be characterized by 3, 4, 5, 6, 7, 8, 9, 10, or more angular positions as shown below:

Angle, 2 theta	d spacing, Å		Angle, 2 theta	d spacing, Å
5.76	15.3		20.67	4.3
6.72	13.2		20.90	4.2
7.57	11.7		21.36	4.2
8.04	11.0		21.54	4.1
9.63	9.2		21.80	4.1
10.85	8.1		22.55	3.9
11.33	7.8		22.89	3.9
11.97	7.4		23.27	3.8
12.38	7.1		23.54	3.8
13.13	6.7		23.87	3.7
13.47	6.6		24.35	3.7
14.75	6.0		24.59	3.6
15.28	5.8		24.87	3.6
16.42	5.4		25.29	3.5
16.89	5.2		25.55	3.5
17.37	5.1		25.89	3.4
17.71	5.0		26.44	3.4
18.17	4.9		27.49	3.2
18.66	4.8		28.01	3.2
19.33	4.6		28.39	3.1
20.01	4.4		29.17	3.1
20.29	4.4			

Table 6. Angular Positions of Characteristic Peaks in Powder X-ray Diffraction for Pattern B (THF Solvate Form IV)

[0124] Polymorph Form IV may be characterized by 3, 4, 5, 6, 7, 8, 9, 10, or more angular positions as shown below:

Angle, 2 theta	d spacing, Å		Angle, 2 theta	d spacing, Å
4.31	20.5		15.90	5.6
5.77	15.3		16.66	5.3
6.28	14.1		16.96	5.2
7.53	11.7		17.33	5.1
7.94	11.1		17.59	5.0
8.76	10.1		18.77	4.7
9.39	9.4		19.09	4.6
9.87	9.0		19.74	4.5
10.54	8.4		20.27	4.4
11.07	8.0		20.57	4.3
11.68	7.6		21.09	4.2
12.02	7.4		21.58	4.1
12.28	7.2		22.81	3.9
12.97	6.8		23.23	3.8
13.20	6.7		24.01	3.7
13.52	6.5		24.65	3.6
14.40	6.1		25.60	3.5
15.08	5.9			

Table 7. Angular Positions of Characteristic Peaks in Powder X-ray Diffraction for Pattern F (2-MeTHF Solvate Form VI)

[0125] Polymorph Form VI may be characterized by 3, 4, 5, 6, 7, 8, 9, 10, or more angular positions as shown below:

Angle, 2 theta	d spacing, Å		Angle, 2 theta	d spacing, Å
4.39	20.1		16.91	5.2
6.27	14.1		17.33	5.1
7.00	12.6		17.59	5.0
8.62	10.3		18.33	4.8
8.85	10.0		18.75	4.7
9.41	9.4		19.13	4.6
9.91	8.9		20.25	4.4
11.32	7.8		20.76	4.3
11.50	7.7		21.68	4.1
12.25	7.2		22.06	4.0
12.56	7.0		22.27	4.0

12.94	6.8		22.61	3.9
13.29	6.7		22.94	3.9
14.03	6.3		24.01	3.7
14.82	6.0		24.33	3.7
15.10	5.9		24.65	3.6
15.44	5.7		25.48	3.5
15.71	5.6		26.05	3.4
16.01	5.5		28.63	3.1
16.67	5.3		29.18	3.1

[0126] Certain polymorphic or amorphous forms of a drug can have advantageous characteristics versus other forms, which can affect the desirability of the drug from a pharmaceutical and/or manufacturing perspective, for example: increased stability, increased solubility, better handling properties, lack of associated undesired solvents (e.g. solvates with toxic solvents), increased purity, better particle size and/or distribution, improved bulk density, and ease of manufacture.

[0127] Forms I, II, III and the amorphous form are anhydrides or hydrates, are advantageously are not solvates with undesired solvents (e.g. THF and 2-MeTHF).

[0128] Forms I-IV and VI were shown to have good solubilities in water (each >1.3 mg/ml), with Form I having the highest aqueous solubility at 1.74 mg/ml (Example 10). Furthermore, Form I was shown to be soluble in a variety of polar and non-polar solvents (Example 2), indicating an ability to be administered using a variety of solvents. In some embodiments, it is advantageous for a drug to have a physiological log D close to zero; solubility in polar and non-polar solvents thus indicates a more favorable physiological log D. Form I was further shown to have increased solubility in a simple detergent (0.5% MC/2%Tween 80) (Example 2); such solubility in simple detergents may be advantageous, as these conditions may mimic gut conditions for oral administration of the drug. Form III (Hydrate), is formed under simple detergent conditions, and thus Form III may be the form of the drug that will be produced in the gut.

[0129] Forms I, II, and III advantageously demonstrated stability to elevated humidity (Example 8). Form I was also tested by grinding, and showed stability to grinding (Example 9). As shown in the examples, the experiments performed indicated that Form I was highly stable.

[0130] Forms that are non-hygroscopic are easier to handle from a manufacturing perspective; as shown in Example 11, Forms I, II, IV, and VI were non-hygroscopic.

[0131] Certain particle shapes and sizes may be advantageous: particles that are closer to spherical may be preferred, with plates and needles less preferred. As shown in Example 11, Forms I, II, and III had more favorable shapes, whereas Form VI was plate shaped, and Form IV needle-shaped. Regarding particle size, smaller, more homogenous sizes may be preferred. Smaller particles may have increased bioavailability, may be easier to dissolve, and may be easier to handle due to decreased drying times. Furthermore, smaller particles may not require a micronizing step that may be required for larger particles. As shown in the Figures, Forms I-III had more favorable particle sizes than IV and VI.

[0132] Higher melting points may indicate a form with improved handling characteristics (e.g. easier to dry and process) and more thermal stability. Form I had the highest melting point of Forms I-VI. Furthermore, a single peak may in some embodiments be preferred, as multiple peaks may indicate conversion to a different form. Form I had a single DSC peak, all others had two or three.

[0133] The various forms (polymorphs and amorphous) of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide may also be utilized as intermediates in making a desired form. As a non-limiting example, if a preferred synthetic method for making (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide results in a non-preferred form, the non-preferred form may be utilized as an intermediate to make the desired form.

Diseases amenable to treatment or suppression with compositions and methods of the invention

[0134] A variety of disorders/diseases are believed to be caused or aggravated by oxidative stress affecting normal electron flow in the cells, such as mitochondrial disorders, impaired energy processing disorders, neurodegenerative diseases and diseases of aging, and can be treated or suppressed using the polymorphic and amorphous forms of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and methods of the invention.

[0135] Non-limiting examples of oxidative stress disorders include, for example, mitochondrial disorders (including inherited mitochondrial diseases) such as Alpers Disease, Barth syndrome, Beta-oxidation Defects, Carnitine-Acyl-Carnitine Deficiency, Carnitine Deficiency, Creatine Deficiency Syndromes, Co-Enzyme Q10 Deficiency, Complex I Deficiency, Complex II Deficiency, Complex III Deficiency, Complex IV Deficiency,

Complex V Deficiency, COX Deficiency, chronic progressive external ophthalmoplegia (CPEO), CPT I Deficiency, CPT II Deficiency, Friedreich's Ataxia (FA), Glutaric Aciduria Type II, Kearns-Sayre Syndrome (KSS), Lactic Acidosis, Long-Chain Acyl-CoA Dehydrogenase Deficiency (LCAD), LCHAD, Leigh Disease or Syndrome, Leigh-like Syndrome, Leber's Hereditary Optic Neuropathy (LHON, also referred to as Leber's Disease, Leber's Optic Atrophy (LOA), or Leber's Optic Neuropathy (LON)), Lethal Infantile Cardiomyopathy (LIC), Luft Disease, Multiple Acyl-CoA Dehydrogenase Deficiency (MAD), Medium-Chain Acyl-CoA Dehydrogenase Deficiency (MCAD), Mitochondrial Myopathy, Encephalopathy, Lactacidosis, Stroke (MELAS), Myoclonic Epilepsy with Ragged Red Fibers (MERRF), Mitochondrial Recessive Ataxia Syndrome (MIRAS), Mitochondrial Cytopathy, Mitochondrial DNA Depletion, Mitochondrial Encephalopathy, Mitochondrial Myopathy, Myoneurogastrointestinal Disorder and Encephalopathy (MNGIE), Neuropathy, Ataxia, and Retinitis Pigmentosa (NARP), Pearson Syndrome, Pyruvate Carboxylase Deficiency, Pyruvate Dehydrogenase Deficiency, POLG Mutations, Respiratory Chain Disorder, Short-Chain Acyl-CoA Dehydrogenase Deficiency (SCAD), SCHAD, Very Long-Chain Acyl-CoA Dehydrogenase Deficiency (VLCAD); myopathies such as cardiomyopathy and encephalomyopathy; neurodegenerative diseases such as Parkinson's disease, Alzheimer's disease, and amyotrophic lateral sclerosis (ALS, also known as Lou Gehrig's disease); motor neuron diseases; neurological diseases such as epilepsy; age-associated diseases, particularly diseases for which CoQ10 has been proposed for treatment, such as macular degeneration, diabetes (e.g. Type 2 diabetes mellitus), metabolic syndrome, and cancer (e.g. brain cancer); genetic diseases such as Huntington's Disease (which is also a neurological disease); mood disorders such as schizophrenia and bipolar disorder; pervasive developmental disorders such as autistic disorder, Asperger's syndrome, childhood disintegrative disorder (CDD), Rett's disorder, and PDD-not otherwise specified (PDD-NOS); cerebrovascular accidents such as stroke; vision impairments such as those caused by neurodegenerative diseases of the eye such as optic neuropathy, Leber's hereditary optic neuropathy, dominant inherited juvenile optic atrophy, optic neuropathy caused by toxic agents, glaucoma, age-related macular degeneration (both "dry" or non-exudative macular degeneration and "wet" or exudative macular degeneration), Stargardt's macular dystrophy, diabetic retinopathy, diabetic maculopathy, retinopathy of prematurity, or ischemic reperfusion-related retinal injury; disorders caused by energy impairment include diseases

due to deprivation, poisoning or toxicity of oxygen, and qualitative or quantitative disruption in the transport of oxygen such as haemoglobinopathies, for example thalassemia or sickle cell anemia; other diseases in which mitochondrial dysfunction is implicated such as excitotoxic, neuronal injury, such as that associated with seizures, stroke and ischemia; and other disorders including renal tubular acidosis; attention deficit/hyperactivity disorder (ADHD); neurodegenerative disorders resulting in hearing or balance impairment; Dominant Optic Atrophy (DOA); Maternally inherited diabetes and deafness (MIDD); chronic fatigue; contrast-induced kidney damage; contrast-induced retinopathy damage; Abetalipoproteinemia; retinitis pigmentosa; Wolfram's disease; Tourette syndrome; cobalamin c defect; methylmalonic aciduria; glioblastoma; Down's syndrome; acute tubular necrosis; muscular dystrophies; leukodystrophies; Progressive Supranuclear Palsy; spinal muscular atrophy; hearing loss (e.g. noise induced hearing loss); traumatic brain injury; Juvenile Huntington's Disease; Multiple Sclerosis; NGLY1; Multiple System Atrophy; Adrenoleukodystrophy; and Adrenomyeloneuropathy. It is to be understood that certain specific diseases or disorders may fall within more than one category; for example, Huntington's Disease is a genetic disease as well as a neurological disease. Furthermore, certain oxidative stress diseases and disorders may also be considered mitochondrial disorders.

[0136] For some disorders amenable to treatment with compounds and methods of the invention, the primary cause of the disorder is due to a defect in the respiratory chain or another defect preventing normal utilization of energy in mitochondria, cells, or tissue(s). Non-limiting examples of disorders falling in this category include inherited mitochondrial diseases, such as Myoclonic Epilepsy with Ragged Red Fibers (MERRF), Mitochondrial Myopathy, Encephalopathy, Lactacidosis, and Stroke (MELAS), Leber's Hereditary Optic Neuropathy (LHON, also referred to as Leber's Disease, Leber's Optic Atrophy (LOA), or Leber's Optic Neuropathy (LON)), Leigh Disease or Leigh Syndrome, Kearns-Sayre Syndrome (KSS), and Friedreich's Ataxia (FA). For some disorders amenable to treatment with compounds and methods of the invention, the primary cause of the disorder is not due to respiratory chain defects or other defects preventing normal utilization of energy in mitochondria, cells, or tissue(s); non-limiting examples of disorders falling in this category include stroke, cancer, and diabetes. However, these latter disorders are particularly aggravated by energy impairments, and are particularly amenable to treatment with compounds of the invention in

order to ameliorate the condition. Pertinent examples of such disorders include ischemic stroke and hemorrhagic stroke, where the primary cause of the disorder is due to impaired blood supply to the brain. While an ischemic episode caused by a thrombosis or embolism, or a hemorrhagic episode caused by a ruptured blood vessel, is not primarily caused by a defect in the respiratory chain or another metabolic defect preventing normal utilization of energy, oxidative stress plays a role in the ischemic cascade due to oxygen reperfusion injury following hypoxia (this cascade occurs in heart attacks as well as in strokes). Accordingly, treatment with compounds and methods of the invention will mitigate the effects of the disease, disorder or condition. Modulating one or more energy biomarkers, normalizing one or more energy biomarkers, or enhancing one or more energy biomarkers can also prove beneficial in such disorders both as a therapeutic measure and a prophylactic measure. For example, for a patient scheduled to undergo non-emergency repair of an aneurysm, enhancing energy biomarkers before and during the pre-operative can improve the patient's prognosis should the aneurysm rupture before successful repair.

[0137] The term “oxidative stress disorder” or “oxidative stress disease” encompass both diseases caused by oxidative stress and diseases aggravated by oxidative stress. The terms “oxidative stress disorder” or “oxidative stress disease” encompass both diseases and disorders where the primary cause of the disease is due to a defect in the respiratory chain or another defect preventing normal utilization of energy in mitochondria, cells, or tissue(s), and also diseases and disorders where the primary cause of the disease is not due to a defect in the respiratory chain or another defect preventing normal utilization of energy in mitochondria, cells, or tissue(s). The former set of diseases can be referred to as “primary oxidative stress disorders,” while the latter can be referred to as “secondary oxidative stress disorders.” It should be noted that the distinction between “diseases caused by oxidative stress” and “diseases aggravated by oxidative stress” is not absolute; a disease may be both a disease caused by oxidative stress and a disease aggravated by oxidative stress. The boundary between “primary oxidative stress disorder” and a “secondary oxidative stress disorder” is more distinct, provided that there is only one primary cause of a disease or disorder and that primary cause is known.

[0138] Bearing in mind the somewhat fluid boundary between diseases caused by oxidative stress and diseases aggravated by oxidative stress, mitochondrial diseases or disorders and impaired energy processing diseases and disorders tend to fall into the category of diseases

caused by oxidative stress, while neurodegenerative disorders and diseases of aging tend to fall into the category of diseases aggravated by oxidative stress. Mitochondrial diseases or disorders and impaired energy processing diseases and disorders are generally primary oxidative stress disorders, while neurodegenerative disorders and diseases of aging may be primary or secondary oxidative stress disorders

Clinical assessment of oxidative stress and efficacy of therapy

[0139] Several readily measurable clinical markers are used to assess the metabolic state of patients with oxidative stress disorders. These markers can also be used as indicators of the efficacy of a given therapy, as the level of a marker is moved from the pathological value to the healthy value. These clinical markers include, but are not limited to, energy biomarkers such as lactic acid (lactate) levels, either in whole blood, plasma, cerebrospinal fluid, or cerebral ventricular fluid; pyruvic acid (pyruvate) levels, either in whole blood, plasma, cerebrospinal fluid, or cerebral ventricular fluid; lactate/pyruvate ratios, either in whole blood, plasma, cerebrospinal fluid, or cerebral ventricular fluid; total, reduced or oxidized glutathione levels, or reduced/oxidized glutathione ratio either in whole blood, plasma, lymphocytes, cerebrospinal fluid, or cerebral ventricular fluid; total, reduced or oxidized cysteine levels, or reduced/oxidized cysteine ratio either in whole blood, plasma, lymphocytes, cerebrospinal fluid, or cerebral ventricular fluid; phosphocreatine levels, NADH (NADH +H⁺) or NADPH (NADPH+H⁺) levels; NAD or NADP levels; ATP levels; anaerobic threshold; reduced coenzyme Q (CoQ_{red}) levels; oxidized coenzyme Q (CoQ_{ox}) levels; total coenzyme Q (CoQ_{tot}) levels; oxidized cytochrome C levels; reduced cytochrome C levels; oxidized cytochrome C/reduced cytochrome C ratio; acetoacetate levels, β -hydroxy butyrate levels, acetoacetate/ β -hydroxy butyrate ratio, 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels; levels of reactive oxygen species; and levels of oxygen consumption (VO₂), levels of carbon dioxide output (VCO₂), and respiratory quotient (VCO₂/VO₂). Several of these clinical markers are measured routinely in exercise physiology laboratories, and provide convenient assessments of the metabolic state of a subject. In one embodiment of the invention, the level of one or more energy biomarkers in a patient suffering from an oxidative stress disorder, such as Friedreich's ataxia, Leber's hereditary optic neuropathy, MELAS, KSS or CoQ10 deficiency, is improved to within two standard deviations of the average level in a healthy subject. In another embodiment of the invention, the level of one or more of

these energy biomarkers in a patient suffering from an oxidative stress disorder, such as Friedreich's ataxia, Leber's hereditary optic neuropathy, MELAS, KSS or CoQ10 deficiency is improved to within one standard deviation of the average level in a healthy subject. Exercise intolerance can also be used as an indicator of the efficacy of a given therapy, where an improvement in exercise tolerance (i.e., a decrease in exercise intolerance) indicates efficacy of a given therapy.

- [0140] Several metabolic biomarkers have already been used to evaluate efficacy of CoQ10, and these metabolic biomarkers can be monitored as energy biomarkers for use in the methods of the current invention. Lactate, a product of the anaerobic metabolism of glucose, is removed by reduction to pyruvate in an aerobic setting or by oxidative metabolism, which is dependent on a functional mitochondrial respiratory chain. Dysfunction of the respiratory chain may lead to inadequate removal of lactate and pyruvate from the circulation and elevated lactate/pyruvate ratios are observed in mitochondrial cytopathies (see Scriver CR, *The metabolic and molecular bases of inherited disease*, 7th ed., New York: McGraw-Hill, Health Professions Division, 1995; and Munnich et al., *J. Inherit. Metab. Dis.* 15(4):448-55 (1992)). Blood lactate/pyruvate ratio (Chariot et al., *Arch. Pathol. Lab. Med.* 118(7):695-7 (1994)) is, therefore, widely used as a noninvasive test for detection of mitochondrial cytopathies (see again Scriver CR, *The metabolic and molecular bases of inherited disease*, 7th ed., New York: McGraw-Hill, Health Professions Division, 1995; and Munnich et al., *J. Inherit. Metab. Dis.* 15(4):448-55 (1992)) and toxic mitochondrial myopathies (Chariot et al., *Arthritis Rheum.* 37(4):583-6 (1994)). Changes in the redox state of liver mitochondria can be investigated by measuring the arterial ketone body ratio (acetoacetate/3-hydroxybutyrate: AKBR) (Ueda et al., *J. Cardiol.* 29(2):95-102 (1997)). Urinary excretion of 8-hydroxy-2'-deoxyguanosine (8-OHdG) often has been used as a biomarker to assess the extent of repair of ROS-induced DNA damage in both clinical and occupational settings (Erhola et al., *FEBS Lett.* 409(2):287-91 (1997); Honda et al., *Leuk. Res.* 24(6):461-8 (2000); Pilger et al., *Free Radic. Res.* 35(3):273-80 (2001); Kim et al. *Environ Health Perspect* 112(6):666-71 (2004)).
- [0141] Magnetic resonance spectroscopy (MRS) has been useful in the diagnoses of mitochondrial cytopathy by demonstrating elevations in cerebrospinal fluid (CSF) and cortical white matter lactate using proton MRS (¹H-MRS) (Kaufmann et al., *Neurology* 62(8):1297-302 (2004)). Phosphorous MRS (³¹P-MRS) has been used to demonstrate low levels of cortical phosphocreatine (PCr) (Matthews et al., *Ann. Neurol.* 29(4):435-8 (1991)),

and a delay in PCr recovery kinetics following exercise in skeletal muscle (Matthews et al., *Ann. Neurol.* 29(4):435-8 (1991); Barbiroli et al., *J. Neurol.* 242(7):472-7 (1995); Fabrizi et al., *J. Neurol. Sci.* 137(1):20-7 (1996)). A low skeletal muscle PCr has also been confirmed in patients with mitochondrial cytopathy by direct biochemical measurements.

[0142] Exercise testing is particularly helpful as an evaluation and screening tool in mitochondrial myopathies. One of the hallmark characteristics of mitochondrial myopathies is a reduction in maximal whole body oxygen consumption (VO₂max) (Taivassalo et al., *Brain* 126(Pt 2):413-23 (2003)). Given that VO₂max is determined by cardiac output (Q_c) and peripheral oxygen extraction (arterial-venous total oxygen content) difference, some mitochondrial cytopathies affect cardiac function where delivery can be altered; however, most mitochondrial myopathies show a characteristic deficit in peripheral oxygen extraction (A-V O₂ difference) and an enhanced oxygen delivery (hyperkinetic circulation) (Taivassalo et al., *Brain* 126(Pt 2):413-23 (2003)). This can be demonstrated by a lack of exercise induced deoxygenation of venous blood with direct AV balance measurements (Taivassalo et al., *Ann. Neurol.* 51(1):38-44 (2002)) and non-invasively by near infrared spectroscopy (Lynch et al., *Muscle Nerve* 25(5):664-73 (2002); van Beekvelt et al., *Ann. Neurol.* 46(4):667-70 (1999)).

[0143] Several of these energy biomarkers are discussed in more detail as follows. It should be emphasized that, while certain energy biomarkers are discussed and enumerated herein, the invention is not limited to modulation, normalization or enhancement of only these enumerated energy biomarkers.

[0144] Lactic acid (lactate) levels: Mitochondrial dysfunction typically results in abnormal levels of lactic acid, as pyruvate levels increase and pyruvate is converted to lactate to maintain capacity for glycolysis. Mitochondrial dysfunction can also result in abnormal levels of NADH +H⁺, NADPH+H⁺, NAD, or NADP, as the reduced nicotinamide adenine dinucleotides are not efficiently processed by the respiratory chain. Lactate levels can be measured by taking samples of appropriate bodily fluids such as whole blood, plasma, or cerebrospinal fluid. Using magnetic resonance, lactate levels can be measured in virtually any volume of the body desired, such as the brain.

[0145] Measurement of cerebral lactic acidosis using magnetic resonance in MELAS patients is described in Kaufmann et al., *Neurology* 62(8):1297 (2004). Values of the levels of lactic acid in the lateral ventricles of the brain are presented for two mutations resulting in MELAS,

A3243G and A8344G. Whole blood, plasma, and cerebrospinal fluid lactate levels can be measured by commercially available equipment such as the YSI 2300 STAT Plus Glucose & Lactate Analyzer (YSI Life Sciences, Ohio).

[0146] NAD, NADP, NADH and NADPH levels: Measurement of NAD, NADP, NADH (NADH +H⁺) or NADPH (NADPH+H⁺) can be measured by a variety of fluorescent, enzymatic, or electrochemical techniques, e.g., the electrochemical assay described in US 2005/0067303.

[0147] GSH, GSSG, Cys, and CySS levels: Briefly, plasma levels of GSH, GSSG, Cys, and CySS are used to calculate the in vivo E_h values. Samples are collected using the procedure of Jones et al (2009 Free Radical Biology & Medicine 47(10) pp 1329-1338), and bromobimane is used to alkylate free thiols and HPLC and either electrochemical or MSMS to separate, detect, and quantify the molecules. As described in more detail in PCT Application No. PCT/US2013/058568, a method was developed for different experimental parameters to analyze the most common monothiols and disulfide (cystine, cysteine, reduced (GSH) and oxidized glutathione (GSSG)) present in human plasma, and using Bathophenanthroline disulfonic acid as the internal standard (IS). Complete separation of all the targets analytes and IS at 35°C on a C18 RP column (250mm×4.6mm, 3 micron) was achieved using 0.2% TFA:Acetonitrile as a mobile phase pumped at the rate of 0.6 ml min⁻¹ using electrochemical detector in DC mode at the detector potential of 1475 mV.

[0148] Oxygen consumption (vO₂ or VO₂), carbon dioxide output (vCO₂ or VCO₂), and respiratory quotient (VCO₂/VO₂): vO₂ is usually measured either while resting (resting vO₂) or at maximal exercise intensity (vO₂ max). Optimally, both values will be measured. However, for severely disabled patients, measurement of vO₂ max may be impractical. Measurement of both forms of vO₂ is readily accomplished using standard equipment from a variety of vendors, e.g. Korr Medical Technologies, Inc. (Salt Lake City, Utah). VCO₂ can also be readily measured, and the ratio of VCO₂ to VO₂ under the same conditions (VCO₂/VO₂, either resting or at maximal exercise intensity) provides the respiratory quotient (RQ).

[0149] Oxidized Cytochrome C, reduced Cytochrome C, and ratio of oxidized Cytochrome C to reduced Cytochrome C: Cytochrome C parameters, such as oxidized cytochrome C levels (Cyt C_{ox}), reduced cytochrome C levels (Cyt C_{red}), and the ratio of oxidized cytochrome C/reduced cytochrome C ratio (Cyt C_{ox})/(Cyt C_{red}), can be measured by in vivo near infrared

spectroscopy. See, e.g., Rolfe, P., “In vivo near-infrared spectroscopy,” *Annu. Rev. Biomed. Eng.* 2:715-54 (2000) and Strangman et al., “Non-invasive neuroimaging using near-infrared light” *Biol. Psychiatry* 52:679-93 (2002).

[0150] Exercise tolerance/Exercise intolerance: Exercise intolerance is defined as “the reduced ability to perform activities that involve dynamic movement of large skeletal muscles because of symptoms of dyspnea or fatigue” (Piña et al., *Circulation* 107:1210 (2003)). Exercise intolerance is often accompanied by myoglobinuria, due to breakdown of muscle tissue and subsequent excretion of muscle myoglobin in the urine. Various measures of exercise intolerance can be used, such as time spent walking or running on a treadmill before exhaustion, time spent on an exercise bicycle (stationary bicycle) before exhaustion, and the like. Treatment with the compounds, compositions, or methods of the invention can result in about a 10% or greater improvement in exercise tolerance (for example, about a 10% or greater increase in time to exhaustion, e.g. from 10 minutes to 11 minutes), about a 20% or greater improvement in exercise tolerance, about a 30% or greater improvement in exercise tolerance, about a 40% or greater improvement in exercise tolerance, about a 50% or greater improvement in exercise tolerance, about a 75% or greater improvement in exercise tolerance, or about a 100% or greater improvement in exercise tolerance. While exercise tolerance is not, strictly speaking, an energy biomarker, for the purposes of the invention, modulation, normalization, or enhancement of energy biomarkers includes modulation, normalization, or enhancement of exercise tolerance.

[0151] Similarly, tests for normal and abnormal values of pyruvic acid (pyruvate) levels, lactate/pyruvate ratio, ATP levels, anaerobic threshold, reduced coenzyme Q (CoQ_{red}) levels, oxidized coenzyme Q (CoQ_{ox}) levels, total coenzyme Q (CoQ_{tot}) levels, oxidized cytochrome C levels, reduced cytochrome C levels, oxidized cytochrome C/reduced cytochrome C ratio, GSH and cysteine reduced, oxidized, total levels and ratio, acetoacetate levels, β -hydroxy butyrate levels, acetoacetate/ β -hydroxy butyrate ratio, 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels, and levels of reactive oxygen species are known in the art and can be used to evaluate efficacy of the compounds, compositions, and methods of the invention. (For the purposes of the invention, modulation, normalization, or enhancement of energy biomarkers includes modulation, normalization, or enhancement of anaerobic threshold.)

[0152] Table 8, following, illustrates the effect that various dysfunctions can have on biochemistry and energy biomarkers. It also indicates the physical effect (such as a disease

symptom or other effect of the dysfunction) typically associated with a given dysfunction. It should be noted that any of the energy biomarkers listed in the table, in addition to energy biomarkers enumerated elsewhere, can also be modulated, enhanced, or normalized by the compounds, compositions, and methods of the invention. RQ = respiratory quotient; BMR = basal metabolic rate; HR (CO) = heart rate (cardiac output); T = body temperature (preferably measured as core temperature); AT = anaerobic threshold; pH = blood pH (venous and/or arterial).

Table 8

Site of Dysfunction	Biochemical Event	Measurable Energy Biomarker	Physical Effect
Respiratory Chain	↑ NADH	Δ lactate, Δ lactate: pyruvate ratio; and Δ acetoacetate: β-hydroxy butyrate ratio	Metabolic dyscrasia & fatigue
Respiratory Chain	↓ H ⁺ gradient	Δ ATP	Organ dependent dysfunction
Respiratory Chain	↓ Electron flux	Δ VO ₂ , RQ, BMR, ΔT, AT, pH	Metabolic dyscrasia & fatigue
Mitochondria & cytosol	↓ ATP, ↓ VO ₂	Δ Work, ΔHR (CO)	Exercise intolerance
Mitochondria & cytosol	↓ ATP	Δ PCr	Exercise intolerance
Respiratory Chain	↓ Cyt C _{Ox} /Red	Δ λ ~700 – 900 nm (Near Infrared Spectroscopy)	Exercise intolerance
Intermediary metabolism	↓ Catabolism	Δ C14-Labeled substrates	Metabolic dyscrasia & fatigue
Respiratory Chain	↓ Electron flux	Δ Mixed Venous VO ₂	Metabolic dyscrasia & fatigue
Mitochondria & cytosol	↑ Oxidative stress	Δ Tocopherol & Tocotrienols, CoQ10, docosahexaenoic acid	Uncertain
Mitochondria & cytosol	↑ Oxidative stress	Δ Glutathionered	Uncertain
Mitochondria & cytosol	Nucleic acid oxidation	Δ8-hydroxy 2-deoxy guanosine	Uncertain
Mitochondria & cytosol	Lipid oxidation	Δ Isoprostane(s), eicosanoids	Uncertain
Cell membranes	Lipid oxidation	Δ Ethane (breath)	Uncertain
Cell membranes	Lipid oxidation	Δ Malondialdehyde	Uncertain

[0153] Treatment of a subject afflicted by an oxidative stress disorder in accordance with the methods of the invention may result in the inducement of a reduction or alleviation of symptoms in the subject, e.g., to halt the further progression of the disorder.

[0154] Partial or complete suppression of the oxidative stress disorder can result in a lessening of the severity of one or more of the symptoms that the subject would otherwise experience. For example, partial suppression of MELAS could result in reduction in the number of stroke-like or seizure episodes suffered.

[0155] Any one or any combination of the energy biomarkers described herein provide conveniently measurable benchmarks by which to gauge the effectiveness of treatment or suppressive therapy. Additionally, other energy biomarkers are known to those skilled in the art and can be monitored to evaluate the efficacy of treatment or suppressive therapy.

Use of compounds or compositions for modulation of energy biomarkers

[0156] In addition to monitoring energy biomarkers to assess the status of treatment or suppression of oxidative stress diseases, the compounds or compositions of the invention can be used in subjects or patients to modulate one or more energy biomarkers. Modulation of energy biomarkers can be done to normalize energy biomarkers in a subject, or to enhance energy biomarkers in a subject.

[0157] Normalization of one or more energy biomarkers is defined as either restoring the level of one or more such energy biomarkers to normal or near-normal levels in a subject whose levels of one or more energy biomarkers show pathological differences from normal levels (i.e., levels in a healthy subject), or to change the levels of one or more energy biomarkers to alleviate pathological symptoms in a subject. Depending on the nature of the energy biomarker, such levels may show measured values either above or below a normal value. For example, a pathological lactate level is typically higher than the lactate level in a normal (i.e., healthy) person, and a decrease in the level may be desirable. A pathological ATP level is typically lower than the ATP level in a normal (i.e., healthy) person, and an increase in the level of ATP may be desirable. Accordingly, normalization of energy biomarkers can involve restoring the level of energy biomarkers to within about at least two standard deviations of normal in a subject, more preferably to within about at least one standard deviation of normal

in a subject, to within about at least one-half standard deviation of normal, or to within about at least one-quarter standard deviation of normal.

[0158] Enhancement of the level of one or more energy biomarkers is defined as changing the extant levels of one or more energy biomarkers in a subject to a level which provides beneficial or desired effects for the subject. For example, a person undergoing strenuous effort or prolonged vigorous physical activity, such as mountain climbing, could benefit from increased ATP levels or decreased lactate levels. As described above, normalization of energy biomarkers may not achieve the optimum state for a subject with an oxidative stress disease, and such subjects can also benefit from enhancement of energy biomarkers. Examples of subjects who could benefit from enhanced levels of one or more energy biomarkers include, but are not limited to, subjects undergoing strenuous or prolonged physical activity, subjects with chronic energy problems, or subjects with chronic respiratory problems. Such subjects include, but are not limited to, pregnant females, particularly pregnant females in labor; neonates, particularly premature neonates; subjects exposed to extreme environments, such as hot environments (temperatures routinely exceeding about 85-86 degrees Fahrenheit or about 30 degrees Celsius for about 4 hours daily or more), cold environments (temperatures routinely below about 32 degrees Fahrenheit or about 0 degrees Celsius for about 4 hours daily or more), or environments with lower-than-average oxygen content, higher-than-average carbon dioxide content, or higher-than-average levels of air pollution (airline travelers, flight attendants, subjects at elevated altitudes, subjects living in cities with lower-than-average air quality, subjects working in enclosed environments where air quality is degraded); subjects with lung diseases or lower-than-average lung capacity, such as tubercular patients, lung cancer patients, emphysema patients, and cystic fibrosis patients; subjects recovering from surgery or illness; elderly subjects, including elderly subjects experiencing decreased energy; subjects suffering from chronic fatigue, including chronic fatigue syndrome; subjects undergoing acute trauma; subjects in shock; subjects requiring acute oxygen administration; subjects requiring chronic oxygen administration; or other subjects with acute, chronic, or ongoing energy demands who can benefit from enhancement of energy biomarkers.

[0159] Accordingly, when an increase in a level of one or more energy biomarkers is beneficial to a subject, enhancement of the one or more energy biomarkers can involve increasing the level of the respective energy biomarker or energy biomarkers to about at least one-quarter

standard deviation above normal, about at least one-half standard deviation above normal, about at least one standard deviation above normal, or about at least two standard deviations above normal. Alternatively, the level of the one or more energy biomarkers can be increased by about at least 10% above the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 20% above the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 30% above the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 40% above the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 50% above the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 75% above the subject's level of the respective one or more energy biomarkers before enhancement, or by about at least 100% above the subject's level of the respective one or more energy biomarkers before enhancement.

[0160] When a decrease in a level of one or more energy biomarkers is desired to enhance one or more energy biomarkers, the level of the one or more energy biomarkers can be decreased by an amount of about at least one-quarter standard deviation of normal in a subject, decreased by about at least one-half standard deviation of normal in a subject, decreased by about at least one standard deviation of normal in a subject, or decreased by about at least two standard deviations of normal in a subject. Alternatively, the level of the one or more energy biomarkers can be decreased by about at least 10% below the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 20% below the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 30% below the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 40% below the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 50% below the subject's level of the respective one or more energy biomarkers before enhancement, by about at least 75% below the subject's level of the respective one or more energy biomarkers before enhancement, or by about at least 90% below the subject's level of the respective one or more energy biomarkers before enhancement.

[0161] The compounds or compositions of the invention can also be used in research applications. They can be used in in vitro, in vivo, or ex vivo experiments to modulate one or more energy biomarkers in an experimental system. Such experimental systems can be cell samples, tissue samples, cell components or mixtures of cell components, partial organs, whole organs, or organisms. Any one or more of the compounds or compositions can be used in experimental systems or research applications. Such research applications can include, but are not limited to, use as assay reagents, elucidation of biochemical pathways, or evaluation of the effects of other agents on the metabolic state of the experimental system in the presence/absence of one or more compounds or compositions of the invention.

[0162] Additionally, the compounds or compositions of the invention can be used in biochemical tests or assays. Such tests can include incubation of one or more compounds or compositions of the invention with a tissue or cell sample from a subject to evaluate a subject's potential response (or the response of a specific subset of subjects) to administration of said one or more compounds or compositions, or to determine which compound or composition of the invention produces the optimum effect in a specific subject or subset of subjects. One such test or assay would involve 1) obtaining a cell sample or tissue sample from a subject in which modulation of one or more energy biomarkers can be assayed; 2) administering one or more compounds or compositions of the invention to the cell sample or tissue sample; and 3) determining the amount of modulation of the one or more energy biomarkers after administration of the one or more compounds or compositions, compared to the status of the energy biomarker prior to administration of the one or more compounds or compositions. Another such test or assay would involve 1) obtaining a cell sample or tissue sample from a subject in which modulation of one or more energy biomarkers can be assayed; 2) administering at least two compounds or compositions of the invention to the cell sample or tissue sample; 3) determining the amount of modulation of the one or more energy biomarkers after administration of the at least two compounds or compositions, compared to the status of the energy biomarker prior to administration of the at least two compounds or compositions, and 4) selecting a compound or composition for use in treatment, suppression, or modulation based on the amount of modulation determined in step 3.

[0163] In certain embodiments, provided herein are methods for the use of a polymorph of Form I-VI for treating or protecting against injury or damage caused by radiation exposure and methods of using such compounds for treating or for protecting against injury or damage

caused by radiation exposure. In certain embodiments, the methods for treating or for protecting against injury or damage caused by radiation exposure, comprise administering to a cell or cells, a tissue or tissues, or a subject in need thereof, a therapeutically effective amount or a prophylactically effective amount of a polymorph of Form I-VI or composition disclosed herein. In one embodiment, the polymorph of Form I-VI are used therapeutically during, after, or during and after radiation exposure. In another embodiment, a polymorph of Form I-VI are used prophylactically prior to radiation exposure. In another embodiment, a polymorph of Form I-VI is administered concurrently with radiation exposure. In another embodiment, the one or more compounds are administered after radiation exposure.

[0164] In certain embodiments, provided herein are methods for the use of a polymorph of Form I-VI for treating against injury or damage caused by radiation exposure and methods of using such compounds for treating or for protecting against injury or damage caused by radiation exposure. In certain embodiments, the methods for treating an injury or damage caused by radiation exposure, comprise administering to a cell or cells, a tissue or tissues, or a subject in need thereof, a therapeutically effective amount or a prophylactically effective amount of a polymorph of Form I-VI or composition disclosed herein. In one embodiment, the polymorph of Form I-VI are used therapeutically during, after, or during and after radiation exposure. In another embodiment, a polymorph of Form I-VI are used prophylactically prior to radiation exposure. In another embodiment, a polymorph of Form I-VI is administered concurrently with radiation exposure. In another embodiment, the one or more compounds are administered after radiation exposure.

Pharmaceutical formulations

[0165] The compounds or compositions described herein can be formulated as pharmaceutical compositions by formulation with additives such as pharmaceutically acceptable excipients, pharmaceutically acceptable carriers, and pharmaceutically acceptable vehicles. Suitable pharmaceutically acceptable excipients, carriers and vehicles include processing agents and drug delivery modifiers and enhancers, such as, for example, calcium phosphate, magnesium stearate, talc, monosaccharides, disaccharides, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, dextrose, hydroxypropyl- β -cyclodextrin, polyvinylpyrrolidinone, low melting waxes, ion exchange resins, and the like, as well as combinations of any two or more thereof. Other suitable pharmaceutically acceptable

excipients are described in “Remington’s Pharmaceutical Sciences,” Mack Pub. Co., New Jersey (1991), and “Remington: The Science and Practice of Pharmacy,” Lippincott Williams & Wilkins, Philadelphia, 20th edition (2003) and 21st edition (2005), incorporated herein by reference.

[0166] A pharmaceutical composition can comprise a unit dose formulation, where the unit dose is a dose sufficient to have a therapeutic or suppressive effect or an amount effective to modulate, normalize, or enhance an energy biomarker. The unit dose may be sufficient as a single dose to have a therapeutic or suppressive effect or an amount effective to modulate, normalize, or enhance an energy biomarker. Alternatively, the unit dose may be a dose administered periodically in a course of treatment or suppression of a disorder, or to modulate, normalize, or enhance an energy biomarker.

[0167] Pharmaceutical compositions containing the compounds or compositions of the invention may be in any form suitable for the intended method of administration, including, for example, a solution, a suspension, or an emulsion. Liquid carriers are typically used in preparing solutions, suspensions, and emulsions. Liquid carriers contemplated for use in the practice of the present invention include, for example, water, saline, pharmaceutically acceptable organic solvent(s), pharmaceutically acceptable oils or fats, and the like, as well as mixtures of two or more thereof. The liquid carrier may contain other suitable pharmaceutically acceptable additives such as solubilizers, emulsifiers, nutrients, buffers, preservatives, suspending agents, thickening agents, viscosity regulators, stabilizers, and the like. Suitable organic solvents include, for example, monohydric alcohols, such as ethanol, and polyhydric alcohols, such as glycols. Suitable oils include, for example, soybean oil, coconut oil, olive oil, safflower oil, cottonseed oil, and the like. For parenteral administration, the carrier can also be an oily ester such as ethyl oleate, isopropyl myristate, and the like. Compositions of the present invention may also be in the form of microparticles, microcapsules, liposomal encapsulates, and the like, as well as combinations of any two or more thereof.

[0168] Time-release or controlled release delivery systems may be used, such as a diffusion controlled matrix system or an erodible system, as described for example in: Lee, “Diffusion-Controlled Matrix Systems”, pp. 155-198 and Ron and Langer, “Erodible Systems”, pp. 199-224, in “Treatise on Controlled Drug Delivery”, A. Kydonieus Ed., Marcel Dekker, Inc., New York 1992. The matrix may be, for example, a biodegradable material that can degrade

spontaneously in situ and in vivo for, example, by hydrolysis or enzymatic cleavage, e.g., by proteases. The delivery system may be, for example, a naturally occurring or synthetic polymer or copolymer, for example in the form of a hydrogel. Exemplary polymers with cleavable linkages include polyesters, polyorthoesters, polyanhydrides, polysaccharides, poly(phosphoesters), polyamides, polyurethanes, poly(imidocarbonates) and poly(phosphazenes).

[0169] The compounds or compositions of the invention may be administered enterally, orally, parenterally, sublingually, by inhalation (e.g. as mists or sprays), rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. For example, suitable modes of administration include oral, subcutaneous, transdermal, transmucosal, iontophoretic, intravenous, intraarterial, intramuscular, intraperitoneal, intranasal (e.g. via nasal mucosa), subdural, rectal, gastrointestinal, and the like, and directly to a specific or affected organ or tissue. For delivery to the central nervous system, spinal and epidural administration, or administration to cerebral ventricles, can be used. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques. The compounds or compositions are mixed with pharmaceutically acceptable carriers, adjuvants, and vehicles appropriate for the desired route of administration. Oral administration is a preferred route of administration, and formulations suitable for oral administration are preferred formulations. The compounds described for use herein can be administered in solid form, in liquid form, in aerosol form, or in the form of tablets, pills, powder mixtures, capsules, granules, injectables, creams, solutions, suppositories, enemas, colonic irrigations, emulsions, dispersions, food premixes, and in other suitable forms. The compounds or compositions can also be administered in liposome formulations. The compounds can also be administered as prodrugs, where the prodrug undergoes transformation in the treated subject to a form which is therapeutically effective. Additional methods of administration are known in the art.

[0170] In some embodiments of the invention, especially those embodiments where a formulation is used for injection or other parenteral administration including the routes listed herein, but also including embodiments used for oral, gastric, gastrointestinal, or enteric administration, the formulations and preparations used in the methods of the invention are

sterile. Sterile pharmaceutical formulations are compounded or manufactured according to pharmaceutical-grade sterilization standards (United States Pharmacopeia Chapters 797, 1072, and 1211; California Business & Professions Code 4127.7; 16 California Code of Regulations 1751, 21 Code of Federal Regulations 211) known to those of skill in the art.

[0171] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions, may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in propylene glycol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[0172] Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

[0173] Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, cyclodextrins, and sweetening, flavoring, and perfuming agents.

[0174] The compounds or compositions of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multilamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are

the phospholipids and phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.W., p. 33 et seq (1976).

[0175] The invention also provides articles of manufacture and kits containing materials useful for treating or suppressing oxidative stress disorders. The invention also provides kits comprising any one or more of the compounds or compositions as described herein. In some embodiments, the kit of the invention comprises a suitable container.

[0176] In other aspects, the kits may be used for any of the methods described herein, including, for example, to treat an individual with a mitochondrial disorder, or to suppress a mitochondrial disorder in an individual.

[0177] The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host to which the active ingredient is administered and the particular mode of administration. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, body area, body mass index (BMI), general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination, and the type, progression, and severity of the particular disease undergoing therapy. The pharmaceutical unit dosage chosen is usually fabricated and administered to provide a defined final concentration of drug in the blood, tissues, organs, or other targeted region of the body. The therapeutically effective amount or effective amount for a given situation can be readily determined by routine experimentation and is within the skill and judgment of the ordinary clinician.

[0178] Examples of dosages which can be used are a therapeutically effective amount or effective amount within the dosage range of about 0.1 mg/kg to about 300 mg/kg body weight, or within about 1.0 mg/kg to about 100 mg/kg body weight, or within about 1.0 mg/kg to about 50 mg/kg body weight, or within about 1.0 mg/kg to about 30 mg/kg body weight, or within about 1.0 mg/kg to about 10 mg/kg body weight, or within about 10 mg/kg to about 100 mg/kg body weight, or within about 50 mg/kg to about 150 mg/kg body weight, or within about 100 mg/kg to about 200 mg/kg body weight, or within about 150 mg/kg to about 250 mg/kg body weight, or within about 200 mg/kg to about 300 mg/kg body weight, or within about 250 mg/kg to about 300 mg/kg body weight. Compounds or compositions of

the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided dosage of two, three or four times daily.

[0179] While the compounds or compositions of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more other agents used in the treatment or suppression of disorders. Representative agents useful in combination with the compounds or compositions of the invention for the treatment or suppression of mitochondrial diseases include, but are not limited to, Coenzyme Q, vitamin E, idebenone, MitoQ, vitamins, NAC, and antioxidant compounds.

[0180] When additional active agents are used in combination with the compounds or compositions of the present invention, the additional active agents may generally be employed in therapeutic amounts as indicated in the Physicians' Desk Reference (PDR) 53rd Edition (1999), or such therapeutically useful amounts as would be known to one of ordinary skill in the art.

[0181] The compounds or compositions of the invention and the other therapeutically active agents can be administered at the recommended maximum clinical dosage or at lower doses. Dosage levels of the active compounds in the compositions of the invention may be varied so as to obtain a desired therapeutic response depending on the route of administration, severity of the disease and the response of the patient. When administered in combination with other therapeutic agents, the therapeutic agents can be formulated as separate compositions that are given at the same time or different times, or the therapeutic agents can be given as a single composition.

[0182] The invention will be further understood by the following nonlimiting examples.

Preparation of Compositions of the Invention

[0183] The compositions of this invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

Synthetic Reaction Parameters

[0184] Solvents employed in synthesis of the compounds and compositions of the invention include, for example, methanol (“MeOH”), acetone, water, acetonitrile, 1,4-dioxane, dimethylformamide (“DMF”), benzene, toluene, xylene, tetrahydrofuran (“THF”), chloroform, methylene chloride (or dichloromethane, (“DCM”)), diethyl ether, pyridine, 2-methyl-tetrahydrofuran (“2-MeTHF”), dimethylacetamide (“DMA”), ethyl acetate (“EtOAc”), ethanol (“EtOH”), isopropyl alcohol (“IPA”), isopropyl acetate (“IPAc”), methyl cellulose (“MC”), acetonitrile (“MeCN”), methanol (MeOH), methyl tert-butyl ether (“MTBE”), phosphate buffered saline (“PBS”), tetrahydrofuran (“THF”), and the like, as well as mixtures thereof.

[0185] The term “q.s.” means adding a quantity sufficient to achieve a stated function, e.g., to bring a solution to the desired volume (i.e., 100%).

[0186] The compounds and compositions herein are synthesized by an appropriate combination of generally well-known synthetic methods. Techniques useful in synthesizing the compounds and compositions herein are both readily apparent and accessible to those of skill in the relevant art in light of the teachings described herein. The discussion below is offered to illustrate certain of the diverse methods available for use in assembling the compounds and compositions herein. However, the discussion is not intended to define the scope of reactions or reaction sequences that are useful in preparing the compounds and compositions herein.

[0187] Other methods for producing the compounds and compositions of the invention will be apparent to one skilled in the art in view of the teachings herein.

EXAMPLES

Example 1. Synthesis of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide (Form I)

Example 1A. Extraction of (1S, 2S)-(+)-Pseudoephedrine free base.

[0188] To a suspension of (1S, 2S)-(+)-Pseudoephedrine hydrochloride salt (300 g, Spectrum) in 2-MeTHF (1.5 L, 5 vol) was added 20% Aq NaOH solution (750 mL, 2.5 vol) and the mixture was stirred for 30 min (some solids remained undissolved) and transferred to a separatory funnel. The lower aqueous layer was drained along with solids that remained at the


interphase and back extracted with 2-MeTHF (750 mL, 2.5 vol), the undissolved solids completely dissolved to form two clear layers. The combined organic layers were evaporated to dryness on rotavapor and the solids obtained were dried in a vacuum oven at 50 °C overnight to afford 240.3 g of free base as a white solid (97.7% recovery).

Example 1B. Precipitation of (1*S*,2*S*)-Pseudoephedrine from 2-MeTHF/heptane

[0189] (1*S*,2*S*)-pseudoephedrine (Sigma-Aldrich, sku#212464, 8.2 g) was dissolved at 50 °C in 2-MeTHF (41 ml, 5 vol). The resulting solution was diluted with heptane (82 ml, 10 vol) and the resulting suspension was stirred at room temperature overnight. The crystallized (1*S*,2*S*)-pseudoephedrine was filtered off and dried overnight at 40 ° under vacuum affording 6.4 g (78%) of white crystalline material. Filtrate was discarded to general waste.

[0190] Relatively low (78%) crystallization yield prompted an additional crystallization experiment with higher heptane to 2-MeTHF ratio. Crystalline (1*S*,2*S*)-pseudoephedrine obtained in the experiment above was dissolved at 50 °C in 2-MeTHF (32 ml, 5 vol). The resulting solution was diluted with heptane (32 ml, 5 vol) and the resulting suspension was chased with heptane (3×50 ml) on rotary evaporator until molar ratio of 2-MeTHF to heptane became lower than 6% by NMR. The resulting suspension was filtered off and the product dried overnight at 40 ° under vacuum affording 6.3 g (98%) of white crystalline material.


Example 1C. Chiral resolution of Trolox using (1*S*, 2*S*)-(+)-Pseudoephedrine.



[0191] Racemic Trolox (316.6 g, 1.27 mol) and (1*S*, 2*S*)-(+)-Pseudoephedrine free base described in Example 1A (240.0 g, 1.46 mol) were charged to a 4L jacketed reactor equipped with an overhead stirrer, temperature probe and a nitrogen purge. Ethyl acetate (EtOAc, 1585 mL, 5 vol) was charged and the slurry was heated to 50°C resulting in clear solution. (Premature (prior to complete dissolution of *rac*-trolox) precipitation of the (*R*)-trolox-


pseudoephedrine salt was occasionally observed at 40°. If premature precipitation takes place the reaction mixture was heated (usually to reflux temperature) to achieve complete dissolution.) The reaction mixture was cooled overnight to room temperature at which time massive precipitation was observed. The mixture was cooled to 10 °C over 30 min and held at this temperature for 1 h. The solids formed were collected by filtration, the wet cake was washed with EtOAc (1.9 L, 6 Vol) and the filter cake was dried in a vacuum oven at 25-30 °C to constant weight to afford 188.1 g (71.3% based on (*R*)-trolox) of a white solid. Chiral HPLC data indicated nearly 100% enantiomeric purity.

Example 1D. Recovery of (*R*)-Trolox from its salt with (1*S*, 2*S*)-(+)-Pseudoephedrine.



[0192] The resulting (*R*)-Trolox PE salt (187.3 g, 0.45 mol) was charged to a 2L round-bottom flask followed by 2-MeTHF (570 ml, 3 vol.) to form a slurry. Hydrochloric acid (2.5 M, 325 ml, 0.81mol, 1.75 eq) was added portionwise while maintaining temperature below 25°C. The trolox-PE salt was dissolved and (*R*)-Trolox was extracted into organic phase. Small black rag was observed in the interface and was kept with the aqueous. The aqueous phase was additionally extracted with 2-MeTHF (2×200 ml). The combined organic layer was then washed with 15% NaCl (200 ml) followed by water (200 ml). The organic layer was dried over anhydrous sodium sulfate (150 g), filtered and evaporated to dryness to afford white solid which was dried under vacuum oven at 30 °C to constant weight of 128.3 g, which is an overstoichiometric amount.

Example 1E. Preparation of (*R*)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxamide.




[0193] CDI (Sigma-Aldrich) (188 g, 1.16 mol) was charged to a 3-neck 2L RBF equipped with an overhead stirrer, nitrogen inlet and temperature probe. 2-MeTHF (290 mL) was added to give a stirrable slurry followed by slow addition of (*R*)-trolox (126.0 g, 504 mmol) in 2-MeTHF (500 ml) at below 30 °C. A slightly exothermic reaction accompanied by CO₂ evolution was observed. Outgassing started after addition of approximately one third of (*R*)-trolox. Complete dissolution of the starting materials was observed in approximately 15 min.

[0194] The content of this flask was slowly added to a pre-cooled to 5 °C 28-30% aqueous ammonia (380 ml) maintaining temperature below 30 °C. The resulting biphasic suspension was stirred at room temperature and monitored by HPLC. The reaction was found to be complete at 36 h and was further processed after 48h.

[0195] The reaction mixture was acidified to pH 1-2 with sulfuric acid (1:4 v/v) (850 ml) maintaining the temperature ≤ 28 °C, reaction was highly exothermic. The aqueous layer (pH=1) was removed and the organic layer was washed with NaCl (15% aqueous w/v, 250 mL), NaHCO₃ (1 M, 250 mL), NaCl (15% aqueous w/v, 250 mL) and water (250 ml). The majority of the organic layer was used for the subsequent steps.

Example 1F. Preparation of (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.



[0196] A solution of (*R*)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxamide (708 ml) which contains ~ 0.39 mole of the intermediate amide and water (126 ml) were charged to a 2L 3N RBF equipped with an overhead stirrer and a thermocouple.

[0197] A stock solution of FeCl₃ × 6H₂O (480 g, 1.78 mol) in water (336 ml) was divided into 4 equal parts (204 g each) and one-fourth of the iron(III) chloride solution was added to the reaction flask. A weak (~3 °C) exotherm was observed, the color of the organic layer turned nearly black then lightened to dark-brown. The biphasic reaction mixture was vigorously stirred for 40 min at room temperature. After removal of the lightly colored aqueous phase another portion of the iron(III) chloride solution was added and stirred for 40 min. The operation was repeated one more time and the organic phase was stored overnight at room temperature. The fourth treatment with FeCl₃ × 6H₂O was performed next morning. Nearly

complete (99.44%) conversion of (R)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxamide to (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide was observed. Initial iron extraction was performed with 1M trisodium citrate solution (2×350 ml); the AUC% of (R)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxamide increased to 0.84%. pH of the organic phase remained highly acidic (pH=1). A 1ml aliquot of the organic phase was treated with 1M NaHCO₃ resulting in massive precipitation of red Fe(OH)₃. Based on this observation one more trisodium citrate wash (175 ml) was performed (0.74% (R)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxamide). The repeat testing of the 1 ml aliquot with 1M NaHCO₃ gave no precipitation in the aqueous layer and the color of the aqueous layer was yellow, not red, indicating complete or nearly complete iron removal.

[0198] The organic layer was heated to 40 °C to prevent premature precipitation of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and washed with 1M sodium bicarbonate solution (175 ml). The phase split was not immediate but was complete in 15 min forming two clear yellow layers. The organic layer (0.30% (R)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxamide) was additionally washed with water (350 ml) giving 0.22% (R)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxamide. Evaporation of the organic layer gave 96 g of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

[0199] The combined bicarbonate/water layers were back extracted with 2×250 ml of 2-MeTHF. Evaporation of these extracts separately gave 4.0 and 0.9 g of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

[0200] The combined solids (100.9 g – crude (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, 84% yield based on (R)-trolox-pseudoephedrine salt) were dissolved in isopropanol (600 ml) at 70 °C and the resulting yellow solution was charged to a 2L 3N RBF equipped with an overhead stirrer a heating mantle and a thermocouple.

[0201] Heptane (600 ml) was added, no precipitation was observed. The reaction mixture was reheated to 55 °C and slowly cooled down to room temperature. Seeds of the desired polymorph (0.2 g) were added and the reaction mixture was stirred overnight at room temperature. Massive precipitation was observed overnight. The reaction mixture was cooled to 7 °C and stirred for additional 8 hours. The product was filtered, washed with isopropanol-

heptane 1:1 v/v (2×75 ml) and dried over the weekend at 40 °C. Yield 69.4 g of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide (58% based on (R)-trolox – pseudoephedrine salt). XRPD data for the product corresponded to the desired Form I.

Example 2. Solubility Measurement of Pattern A

[0202] (R)-Trolox was produced from racemic Trolox via methylbenzyl amine double resolution in a manner similar to that described in Example 29 of U.S. Pat. No. 4,026,907. This (R)-Trolox was used to synthesize the (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide starting material. This starting material, designated as Pattern A, was used for the solubility measurement.

[0203] Excess amount of solid was slurried in 17 solvent systems having diverse properties for minimum of 3 days. The slurry was centrifuged and the clear solution was used for gravimetric method. The compound showed elevated solubility in MeOH, EtOH at ambient temperature, and IPA, acetone, MeOH, EtOH and 2-MeTHF at 50 °C. Moderate solubility was observed in EtOAc, THF, IPA, acetone, 2-MeTHF, MeCN, 0.5% Methyl Cellulose/ 2% Tween 80, IPAc and 4% DMA in PBS at ambient and 50 °C. Limited solubility was observed from heptane, toluene, MTBE, water and 0.5% Methyl Cellulose in water at ambient and elevated temperatures. Table 9 presents the measured solubility data. A ±10% error is expected.

Table 9: Solubility of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in various solvent systems (Starting material, Pattern A used)

Solvent	mg/mL at 25 °C	mg/mL at 50 °C
Heptane	6	6
Toluene	3	<18
MTBE	10	16
EtOAc	22	38
THF	28.	33
IPA	20	53
Acetone	39	90
EtOH	40	>121
MeOH	>101	>112
2-MeTHF	35	59
MeCN	20	38

Water	3	3
¹ Water	2	6
0.5% MC/2% Tween 80*	26	30
IPAc	13	16
0.5% MC in Water*	7	4
4% DMA in PBS*	10	13

* Part of the concentration relates to solvent constituents

[0204] As shown in Table 9, Pattern A is soluble in a variety of polar and non-polar solvents, and in addition has increased solubility in a simple detergent (0.5% MC/2% Tween 80).

Example 3. Short Term Slurry Experiments

[0205] Short term slurry experiments of the starting material, Pattern A, were performed for a minimum of 3 days in 17 different solvent systems having diverse properties at two different temperatures (25 and 50 °C). A 48 position Chemglass reaction block was used for heating and stirring the slurries which were in 2 mL HPLC vials. After the due time, vials were centrifuged and the wet solids were used for X-ray diffraction. Table 10 shows the results of the slurry experiments. These results demonstrate that Pattern A solid was relatively stable if slurried in most of these solvents for a short period of time. However, slurrying Pattern A solid in THF and 0.5% Methyl Cellulose/2% Tween 80 resulted in two new X-ray patterns designated as Patterns B and C respectively (Figure 1).

Table 10: Summary of a minimum 3 days slurry of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide (Starting material, Pattern A used)

Solvent	Starting Form	25 °C		50 °C
		Resulting Form, Wet	Resulting Form, Dry	Resulting Form, Wet
Heptane	A	A		A
Toluene	A	A		A
MTBE	A	A		A
EtOAc	A	A		A
THF	A	B	B	-
IPA	A	A		A
Acetone	A	A		A
EtOH	A	A		-
MeOH	A	-		-
2-MeTHF	A	A		A
MeCN	A	A		A
Water	A	A		A

¹ H ₂ O	A	A		A
0.5% MC/2% Tween 80	A	C	C	A
IPAc	A	A		A
0.5% MC in Water	A	A		A
4% DMA in PBS	A	A		A

Example 4. Evaporative Crystallization Experiments

[0206] Evaporative crystallization experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide were performed using the samples generated during the gravimetric solubility determination (Example 2). XRPD analysis of most samples afforded Pattern A. However XRPD analysis of solids isolated from 2-MeTHF at 25 °C was found to afford a unique crystalline pattern, designated Pattern D as shown in Figure 2. XRPD analysis of samples from IPAc, 0.5% MC in water and 4% DMA in PBS mostly yielded amorphous patterns with the exception of IPAc at 50 °C which afforded a crystalline pattern consistent with Pattern A. All results are summarized in Table 11.

Table 11: Summary of evaporative crystallization experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide (Starting material, Pattern A used)

Solvent	Starting Form	25 °C		50 °C
		Resulting Form, Wet	Resulting Form, Dry	Resulting Form, Wet
Heptane	A	-		-
Toluene	A	A		A
MTBE	A	Oil		Amorphous
EtOAc	A	A		A
THF	A	Amorphous		A
IPA	A	Amorphous		Amorphous
Acetone	A	A		A
EtOH	A	A		A
MeOH	A	A		A
2-MeTHF	A	D	NA	A
MeCN	A	A		A
Water	A	A		A
¹ H ₂ O	A	A		A
0.5% MC/2% Tween 80	A	Oil		Oil
IPAc	A	Amorphous		A
0.5% MC in Water	A	Amorphous		-
4% DMA in PBS	A	Amorphous		Amorphous

NA- Drying not performed due to sample dried overnight

Example 5. Crystallization Experiments

[0207] Fast and slow cooling single solvent crystallization experiments were performed in Toluene, EtOAc, IPA, acetone, EtOH, 2-MeTHF, and IPA with 2% water (Table 12). A 48 position Chemglass reaction block was used for heating and stirring which were performed in 4 mL vials. Each vial was charged with 50-80 mg of starting material (Pattern A) fitted with a magnetic stir bar. Primary solvent was added and heated with stirring until dissolution achieved. Once fully dissolved the sample was slow cooled by radiative cooling or crashed cooled with use of an ice bath followed over night equilibration with stirring. For Binary solvent crystallizations, anti-solvent (Heptane) was added in two methods (Table 13). In method one, the anti-solvent was added drop wise to the sample solution until slight precipitation was observed. Method two used a reverse addition of the sample solution to a heated anti-solvent in a 2:1 ratio before being allowed to cool. Samples which afforded solids after overnight equilibration were isolated by filtration and samples that did not precipitate were evaporated under a gentle stream of nitrogen. All samples were dried overnight in a vacuum oven at ambient conditions and analyzed by XRPD to check for form change. All experimental details and results are summarized in Tables 12-13.

[0208] XRPD analysis of all isolated solids mostly afforded crystalline patterns consistent with the starting material, Pattern A. However, single solvent crystallizations performed in 2-MeTHF with fast and slow cooling profiles, were not observed to afford solids upon cooling and were evaporated to dryness under nitrogen. These evaporated solids were found to afford XRPD patterns consistent with previously observed Pattern D (Figure 3). Fast cooling crystallization performed in EtOAc, yielded unique crystalline solids by XRPD which were compared to all known forms and designated as Pattern E (Figure 4). Binary solvent crystallizations with fast and slow cooling profiles performed in EtOH/Heptane and Acetone/Heptane afforded XRPD patterns consistent with Pattern A.

Table 12: Summary of single solvent crystallization experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide (Starting material, Pattern A used)

Starting Material (Pattern A)	Primary Solvent		Temp (C)	Cooling Rate	Isolation	XRPD [Pattern]
	Solvent	Vol (mL)				

(mg)						
49.7	Toluene	2.75	60	Fast	NA	-
52.3	EtOAc	1	60	Fast	Filter	Crystalline [Pattern E]
55.9	IPA	1	60	Fast	Filter	Crystalline [Pattern A]
50.9	Acetone	0.5	60	Fast	Filter	Crystalline [Pattern A]
82.0	EtOH	0.5	60	Fast	Filter	Crystalline [Pattern A]
51.1	2-MeTHF	1	60	Fast	Evap	Crystalline [Pattern D]
48.9	IPA with 2% water	0.5	60	Fast	Filter	Crystalline [Pattern A]
50.2	Toluene	2.75	60	Slow	NA	-
49.2	EtOAc	1	60	Slow	Filter	Crystalline [Pattern A]
54.9	IPA	1	60	Slow	Filter	Crystalline [Pattern A]
51.9	Acetone	0.5	60	Slow	Filter	Crystalline [Pattern A]
82.7	EtOH	0.5	60	Slow	Filter	Crystalline [Pattern A]
54.6	2-MeTHF	1	60	Slow	Evap	Crystalline [Pattern D]
54	IPA with 2% water	0.5	60	Slow	Filter	Crystalline [Pattern A]

Table 13: Summary of binary solvent crystallization experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide (Starting material, Pattern A used)

Starting Material (Pattern A) (mg)	Primary Solvent		Temp (°C)	Anti- Solvent		Rate of Anti-Solvent Addition	Isolation	XRPD [Pattern]
	Solvent	Vol (mL)		Solvent	Vol (mL)			
84.5	EtOH	0.5	60	Heptane	2.5	Slow	Filter	Crystalline [Pattern A]
53.7	Acetone	0.5	60	Heptane	1.5	Slow	Filter	Crystalline [Pattern A]
82.2	EtOH	0.5	60	Heptane	1.0	Fast	Filter	Crystalline [Pattern A]
56.1	Acetone	0.5	60	Heptane	1.0	Fast	Filter	Crystalline [Pattern A]

Example 6. Scale-Up Experiments

[0209] Scale-up experiments were performed on a 300 mg scale by single solvent fast cooling crystallizations in 2-MeTHF, EtOAc, and slurries in THF and 0.5% Methyl Cellulose/2 % Tween 80 in an attempt to isolate previously observed Patterns D, E, B and C respectively for further characterization. Experimental details and results are summarized in Tables 14-15.

[0210] Fast cooling single solvent crystallization experiments were performed in 2-MeTHF and EtOAc (Table 14). A 48 position Chemglass reaction block was used for heating and stirring

which were performed in 4 mL vials. Each vial was charged with approximately 300 mg of starting material fitted with a magnetic stir bar. 6 mL of primary solvent was added and heated to 60 °C with stirring until dissolution achieved. Once fully dissolved the sample was (crashed cooled) transferred to an ice bath and seeded with a spatula tip full of Pattern D or E, followed by overnight equilibration at room temperature with stirring. Samples which afforded solids after overnight equilibration were isolated by filtration and samples that did not precipitate were evaporated under a gentle stream of nitrogen. All samples were dried overnight under vacuum at ambient conditions and XRPD analysis was performed to check for form change.

[0211] Slurry experiments were performed in THF and 0.5% Methyl Cellulose/2 % Tween 80 (Table 15). A 48 position Chemglass reaction block was used for heating and stirring which were performed in 4 mL vials. Each vial was charged with approximately 300 mg of starting material fitted with a magnetic stir bar. Slurry solvent was added up to 2 mL at ambient conditions and allowed to equilibrate for 30 minutes before adding a spatula tip full of Pattern B or C.

[0212] XRPD analysis of solids isolated from single solvent crystallizations performed in 2-MeTHF with fast and cooling profiles, afforded a unique XRPD pattern, designated as Pattern F (Figure 4). Fast cooling crystallization performed in EtOAc yielded crystalline solids consistent with Pattern E by XRPD (Figure 5). Slurry experiments performed in THF and 0.5% Methyl Cellulose/2 % Tween 80, were found to afford Patterns B and C respectively following 24 hours of equilibration (Figure 6-7).

Table 14: Summary of single solvent crystallization scale-up experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

Starting material (Pattern A) (mg)	Primary Solvent		Temp (°C)	Cooling Rate	Seeding upon cooling [mg]	Isolation	XRPD [Pattern]
	Solvent	Vol (mL)					
312.0	2-MeTHF	6.0	60	Fast	[~10]	Evap	Crystalline [Pattern F]
300.7	EtOAc	6.0	60	Fast	[~10]	Filter	Crystalline [Pattern E]

Table 15: Summary of slurry scale-up experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

Starting material (Pattern A) (mg)	Primary Solvent		Slurry Temp. (°C)	Slurry Formation	Seeding upon Slurry [mg]	XRPD [Pattern]
	Solvent	Vol (mL)				24 hr
305.7	THF	2	Ambient	Yes	[~10]	Crystalline [Pattern B]
308.3	0.5% MC/2% Tween 80 in water	2	Ambient	Yes	[~10]	Crystalline [Pattern C]

Example 7. Competitive Slurries

[0213] Competitive slurry experiments of Patterns A, B, C, D, E and F were initiated in IPA, IPA/2% water and 0.5% Methyl Cellulose in water at ambient conditions as summarized in Table 16. Approximately 100 mg of the starting material, Pattern A, was added to a glass vial fitted with a magnetic stir bar. Solvent was added to the vial and allowed to slurry with Pattern A for 15 minutes before approximately 10-20 mg of each relative Pattern (B, C, D, E and F) was added to each vial. The samples were allowed to equilibrate with stirring and following 24 hours or equilibration, XRPD analysis showed solids isolated from slurry in 0.5% Methyl Cellulose in water was a mixture of Patterns A/C (Figure 8). All other solids isolated from IPA and IPA/2% water were consistent with the starting material, Pattern A (Figure 9). However, following 7 days of equilibration full conversion to Pattern C was observed from slurry in 0.5% Methyl Cellulose in water (Figure 8), where as solids isolated from competitive slurry in IPA and IPA/2% water were consistent with Pattern A (Figure 9).

Table 16: A summary of competitive slurry experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in 2 mL of solvent.

Solvent	Pattern (mg)						XRPD	
	Pattern A	Pattern B	Pattern C	Pattern D	Pattern E	Pattern F	24 hrs	7 days
IPA	101.9*	~ 10	~ 10	~ 10	~ 10	~ 10	A	A
IPA:water (98:2)	107.3*	~10	~10	~10	~10	~10	A	A
0.5%	103.9*	~10	~10	~10	~10	~10	A/C	C

MC/water								
----------	--	--	--	--	--	--	--	--

*- Used to saturate the solvent and make a thin slurry of Pattern A

Example 8. Elevated Humidity Studies

[0214] 7 day elevated aqueous humidity experiments were performed on all Patterns A, B, C, D, E and F at ambient conditions at > 95 %RH (Table 17). Approximately 30 mg of each pattern was weighed into a 4 ml glass vial. The uncovered 4 ml vial was inserted into a 20 ml scintillation vial half filled with water and capped. Following 24 hours of equilibration, visual inspection was performed to check for changes in physical appearance, however no change was observed. Following 7 days of equilibration at elevated humidity, the sample showed no physical changes and was analyzed by XRPD to check for form. Patterns A, C and E showed no change in form after 7 days of equilibration; however Pattern B was converted to a mixture of Patterns A/B. Pattern D was found to convert to a mixture of Patterns D/B and Pattern F was converted to Pattern E at > 95 %RH. All experimental details and results are summarized in Table 17.

Table 17: Summary of 7 day elevated aqueous humidity experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide

[mg]	Storage Conditions	Initial Pattern	Visual Observations		XRPD [Pattern]
			1 day	7 day	7 day
~ 30	> 95 %RH at Ambient	A	Yellow solids	Yellow Solids	Crystalline [Pattern A]
~ 30	> 95 %RH at Ambient	B	Yellow solids	Yellow solids	Crystalline [Mixture of Patterns A/B]
~ 30	> 95 %RH at Ambient	C	Yellow solids	Yellow solids	Crystalline [Pattern C]
~ 30	> 95 %RH at Ambient	D	Yellow solids	Yellow solids	Crystalline [Mixture of Patterns D/B]
~ 30	> 95 %RH at Ambient	E	Yellow solids	Yellow solids	Crystalline [Pattern E]
~ 30	> 95 %RH at Ambient	F	Yellow solids	Yellow solids	Crystalline [Pattern E]

Example 9. Grinding Experiments

[0215] Grinding experiments of the starting material were performed by dry grinding and solvent drop grinding in IPA and water utilizing a mortar and pestle (Table 18). Following light grinding pattern A, no change in the crystal form was observed by XRPD analysis.

Table 18: Summary of Grinding Experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide

Sample Wt. (mg)	Solvent	Grinding	Conditions	XRPD
30-50	-	yes	RT	Pattern A
30-50	IPA (2 drops)	yes	RT	Pattern A
30-50	H2O (2 drops)	yes	RT	Pattern A

Example 10. Aqueous Solubility

[0216] Aqueous solubility of Patterns A, B, C, E and F was performed using an Agilent HPLC system. Approximately 10-20 mg of each form was charged to a 2 mL glass vial loaded with a magnetic stir bar and added 2 mL of water. The samples were allowed to stir overnight at ambient conditions. Following 24 hours of equilibration the samples were centrifuged and decanted into HPLC vials. A calibration curve was generated based on Pattern A in MeOH at 0.05, 0.1, 0.5 and 1.0 mg/mL. Following injection of the standard curve, the samples were run as is. All experimental details and results are summarized in Table 19.

Table 19: Summary of Aqueous Solubility experiments of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide

(mg)	Water (mL)	Pattern	Designated Form	% AUC	Solubility (mg/mL)
15.0	2	F	2-MeTHF Solvate Form VI	6813	1.36
12.1	2	E	Anhydrate Form II	6602	1.32
12.3	2	B	THF Solvate Form IV	6514	1.30
13.5	2	C	Hydrate Form III	6646	1.32
21.5	2	A	Anhydrate Form I	8729	1.74

Example 11. Characterization of Forms

[0217] Solid form characterization of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, polymorphic forms was completed by XRPD, DSC, TGA, ¹H NMR, Karl Fischer, optical microscopy, and moisture sorption. Results are summarized in Table 1.

Pattern A (Anhydrate, Form I)

- [0218] Utilizing the starting material as noted in Example 2, XRPD analysis of the yellow colored starting material was found to afford a crystalline pattern, designated as Pattern A (Figure 10). The crystallinity observed by XRPD was confirmed by the exhibition of birefringence observed by optical microscopy. The morphology of the crystals was determined to be irregularly shaped with some aggregation as shown in Figure 11.
- [0219] Thermal analysis by DSC showed a single endothermic event at peak of 152.9 °C, followed by degradation after 200 °C (Figure 12).
- [0220] TGA analysis showed no weight loss between 45-160 °C, however weight loss due to decomposition was observed from 160-300 °C (Figure 13). Minimal moisture content was confirmed by Karl Fischer analysis which showed the materials to contain approximately 0.12 wt% water.
- [0221] Further analysis by ¹H NMR showed the starting material to be consistent with structure of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and to contain 0.28 wt% residual IPA. See Figure 40.
- [0222] Moisture sorption analysis of the starting material was performed by equilibrating the sample at 25 °C and 50% RH to simulate ambient lab conditions. Humidity then decreased to 0% RH, increased from 0 to 95% RH, reduced from 95 to 0% RH, increased from 0 to 95 %RH and then decreased from 95 to 50% RH. Each point represents the estimated asymptotic weight for each humidity or weight. The starting material was found to non-hygroscopic, adsorbing 0.1% water at 90 %RH. No hysteresis was observed upon desorption (Figure 14). XRPD analysis of sample following moisture sorption analysis was found to be consistent with the starting material, Pattern A.

Pattern E (Anhydrate, Form II)

- [0223] Pattern E (Anhydrate, Form II) was observed during single solvent crystallizations (Examples 5 and 6) using a fast cooling profile at the 50 mg scale and again at the 300 mg scale-up. XRPD analysis of the solids were found to afford a unique crystalline pattern, designated as Pattern E (Figure 15). The crystallinity observed by XRPD was confirmed by the exhibition of birefringence observed by optical microscopy. The morphology of the

crystals was determined to be irregularly shaped with some aggregation as shown in Figure 16.

[0224] Thermal analysis of the 50 mg scale lot by DSC showed a two endothermic events at peak of 133.9 °C and 151.3 °C, followed by degradation after 200 °C (Figure 17).

[0225] TGA analysis of the 50 mg scale lot, showed a 0.4% weight loss between 120-140 °C, likely attributed to the loss of EtOAc, followed by decomposition (Figure 18). Minimal moisture content was confirmed by Karl Fischer analysis which showed the materials to contain approximately 0.1 wt% water.

[0226] Further analysis of Pattern E (from the 300 mg scale lot) by ¹H NMR showed the material to be consistent with structure of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and to contain 0.4 wt% residual EtOAc. See Figure 44.

[0227] Moisture sorption analysis of Pattern E (from the 300 mg scale lot) was performed by equilibrating the sample at 25 °C and 50% RH to simulate ambient lab conditions. Humidity was then decreased to 0% RH, increased from 0 to 95% RH, reduced from 95 to 0% RH, increased from 0 to 95 %RH and then decreased from 95 to 50% RH. Each point represents the estimated asymptotic weight for each humidity or weight. Pattern E was found to non-hygroscopic, adsorbing 0.2% water at 95 %RH. No hysteresis was observed upon desorption (Figure 19). XRPD analysis of sample following moisture sorption analysis was found to be consistent with Pattern E.

Pattern C (Hydrate, Form III)

[0228] Pattern C (Hydrate, Form III) was observed during short term slurry in 0.5% Methyl Cellulose/2% Tween 80 at the 50 mg scale and again at the 300 mg scale-up (Examples 3 and 6, respectively). XRPD analysis of the solids were found to afford a unique crystalline pattern, designated as Pattern C (Figure 20). The crystallinity observed by XRPD was confirmed by the exhibition of birefringence observed by optical microscopy. The morphology of the crystals was determined to be irregularly shaped with some aggregation as shown in Figure 21.

[0229] Thermal analysis of the 50 mg lot by DSC showed two endothermic events at peaks of 72 °C and 150.7 °C, followed by degradation after 200 °C (Figure 22).

[0230] TGA analysis of the 50 mg lot showed a 2.5% weight loss between 20-60 °C, followed by a 2.3% wt. loss from 60-125 °C, likely attributed to dehydration, followed by decomposition (Figure 23). Moisture content was confirmed by Karl Fischer analysis which showed the materials to contain approximately 4.3 wt% water, slightly lower than a monohydrate.

[0231] Further analysis of Pattern C (300 mg lot) by ¹H NMR showed the material to be consistent with structure of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide. See Figure 42.

[0232] Moisture sorption analysis of Pattern C (300 mg lot) was performed by equilibrating the sample at 25 °C and 50% RH to simulate ambient lab conditions. Humidity was then decreased to 0% RH, increased from 0 to 95% RH, reduced from 95 to 0% RH, increased from 0 to 95 %RH and then decreased from 95 to 50% RH. Each point represents the estimated asymptotic weight for each humidity or weight. Pattern C was found to be slightly hygroscopic, adsorbing 2% water at 95 %RH. This increased the total water to about 6% which is the water content of mono-hydrate. However, upon reducing the relative humidity, the solid lost its water. Therefore this could be a channel hydrate. No hysteresis was observed upon desorption (Figure 24). XRPD analysis of sample following moisture sorption analysis was found to be consistent with Pattern C.

Pattern B (THF Solvate, Form IV)

[0233] Pattern B (THF Solvate, Form IV) was observed during short term slurry in THF at the 50 mg scale and again at the 300 mg scale-up (Examples 3 and 6 respectively). XRPD analysis of the solids were found to afford a unique crystalline pattern, designated as Pattern B (Figure 25). The crystallinity observed by XRPD was confirmed by the exhibition of birefringence observed by optical microscopy. The morphology of the crystals was determined to be needle shaped with some aggregation as shown in Figure 26.

[0234] Thermal analysis of the 50 mg lot, by DSC showed three endothermic events at peak of 70.5, 89.1°C and 149.7 °C, followed by degradation after 200 °C (Figure 27).

[0235] TGA analysis of the 300 mg lot, showed a 4.7% weight loss between 25-115 °C, likely attributed to the loss of THF, followed by decomposition (Figure 28). Moisture content was confirmed by Karl Fischer analysis which showed the materials to contain approximately 0.3 wt% water.

[0236] Further analysis of Pattern B (300 mg lot) by ¹H NMR showed the material to be consistent with structure of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and contain 6.9 wt% residual THF. See Figure 41.

[0237] Moisture sorption analysis of Pattern B (300 mg lot) was performed by equilibrating the sample at 25 °C and 50% RH to simulate ambient lab conditions. Humidity then decreased to 0% RH, increased from 0 to 95% RH, reduced from 95 to 0% RH, increased from 0 to 95 %RH and then decreased from 95 to 50% RH. Each point represents the estimated asymptotic weight for each humidity or weight. Pattern B was found to be non-hygroscopic, showing weight loss likely due to release of residual THF (Figure 29). The major weight loss at the beginning was due to loss of solvent. XRPD analysis of sample following moisture sorption analysis was found to be consistent with Pattern A.

Pattern D (2-MeTHF Solvate, Form V)

[0238] Pattern D (2-MeTHF Solvate, Form V) was observed during evaporative crystallizations in 2-MeTHF (Examples 4 and 5). XRPD analysis of the solids were found to afford a unique crystalline pattern, designated as Pattern D (Figure 30).

[0239] Thermal analysis of the Pattern D from Example 5 (slow cooling), by DSC showed four endothermic events at peaks of 67.2, 92.2, 132.6 and 150.6 °C, followed by degradation after 220 °C (Figure 31).

[0240] TGA analysis of Example 5 slow cooling lot, showed a 2.7% weight loss between 40-60 °C followed by a 5.3% weight loss from 60-115 °C, likely attributed to the loss of 2-MeTHF, followed by decomposition (Figure 32).

[0241] Further analysis of Pattern D (Example 5 slow cooling lot) by ¹H NMR showed the material to be consistent with structure of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and contain 6.1 wt% residual 2-MeTHF. See Figure 43.

Pattern F (2-MeTHF Solvate, Form VI)

[0242] Pattern F (2-MeTHF Solvate, Form VI) was observed during single solvent crystallization scale-up experiments 2-MeTHF at the 300 mg scale (Example 6). XRPD analysis of the solids were found to afford a unique crystalline pattern, designated as Pattern F (Figure 33). The crystallinity observed by XRPD was confirmed by the exhibition

of birefringence observed by optical microscopy. The morphology of the crystals was determined to be plate shaped with some aggregation as shown in Figure 34.

[0243] Thermal analysis of Pattern F (from Example 6) by DSC showed three endothermic events at peaks of 93.2, 135.2 °C and 151.0 °C, followed by degradation after 220 °C (Figure 35).

[0244] TGA analysis of Pattern F (from Example 6), showed a 1.1% weight loss between 30-110 °C, followed by a 0.2% weight loss from 110-160 °C, likely attributed to the loss of 2-MeTHF, followed by decomposition (Figure 36). Moisture content was confirmed by Karl Fischer analysis which showed the materials to contain approximately 0.1 wt% water.

[0245] Further analysis of Pattern F (from Example 6) by ¹H NMR showed the material to be consistent with structure of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide and contain 3.9 wt% residual 2-MeTHF. See Figure 45.

[0246] Moisture sorption analysis of Pattern F (from Example 6) was performed by equilibrating the sample at 25 °C and 50% RH to simulate ambient lab conditions. Humidity was then decreased to 0% RH, increased from 0 to 95% RH, reduced from 95 to 0% RH, increased from 0 to 95 %RH and then decreased from 95 to 50% RH. Each point represents the estimated asymptotic weight for each humidity or weight. Pattern F was found to be non-hygroscopic, showing weight loss likely due to release of residual 2- MeTHF (Figure 37). The major weight loss at the beginning is due to loss of solvent. XRPD analysis of the sample following moisture sorption analysis was found to be consistent with Pattern E (Figure 15).

Example 12. Screening Compounds of the Invention in Human Dermal Fibroblasts from Friedreich's Ataxia Patients

[0247] An initial screen was performed to identify compounds effective for the amelioration of redox disorders. Test samples were tested for their ability to rescue FRDA fibroblasts stressed by addition of L-buthionine-(S,R)-sulfoximine (BSO), as described in Jauslin et al., Hum. Mol. Genet. 11(24):3055 (2002), Jauslin et al., FASEB J. 17:1972-4 (2003), and International Patent Application WO 2004/003565. Human dermal fibroblasts from Friedreich's Ataxia patients have been shown to be hypersensitive to inhibition of the de novo synthesis of glutathione (GSH) with L-buthionine-(S,R)-sulfoximine (BSO), a specific inhibitor of GSH synthetase (Jauslin et al., Hum. Mol. Genet. 11(24):3055 (2002)).

[0248] MEM (a medium enriched in amino acids and vitamins, catalog no. 1-31F24-I) and Medium 199 (M199, catalog no. 1-21F22-I) with Earle's Balanced Salts, without phenol red,

were purchased from Bioconcept. Fetal Calf Serum was obtained from PAA Laboratories. Basic fibroblast growth factor and epidermal growth factor were purchased from PeproTech. Penicillin-streptomycin-glutamine mix, L-buthionine (S,R)-sulfoximine, and insulin from bovine pancreas were purchased from Sigma. Calcein AM was purchased from Anaspec. Cell culture medium was made by combining 125 ml M199 EBS, 50 ml Fetal Calf Serum, 100 U/ml penicillin, 100 microgram/ml streptomycin, 2 mM glutamine, 10 microgram/ml insulin, 10 ng/ml EGF, and 10 ng/ml bFGF; MEM EBS was added to make the volume up to 500 ml. During the course of the experiments, this solution was stored at +4°C. The cells were obtained from the Coriell Cell Repositories (Camden, NJ; repository number GM04078) and grown in 10 cm tissue culture plates. Every third day, they were split at a 1:3 ratio.

[0249] The test samples were supplied in 1.5 ml glass vials. The compounds were diluted with DMSO, ethanol or PBS to result in a 5 mM stock solution. Once dissolved, they were stored at -20°C.

[0250] Test samples were screened according to the following protocol:

[0251] A culture with FRDA fibroblasts was started from a 1 ml vial with approximately 500,000 cells stored in liquid nitrogen. Cells were propagated in 10 cm cell culture dishes by splitting every third day in a ratio of 1:3 until nine plates were available. Once confluent, fibroblasts were harvested. For 54 micro titer plates (96 well-MTP) a total of 14.3 million cells (passage eight) were re-suspended in 480 ml medium, corresponding to 100 microliters medium with 3,000 cells/well. The remaining cells were distributed in 10 cm cell culture plates (500,000 cells/plate) for propagation. The plates were incubated overnight at 37°C in a atmosphere with 95% humidity and 5% CO₂ to allow attachment of the cells to the culture plate.

[0252] 10% DMSO (242.5 microliters) was added to a well of the microtiter plate. The test compounds were unfrozen, and 7.5 microliters of a 5 mM stock solution was dissolved in the well containing 242.5 microliters of 10% DMSO, resulting in a 150 micromolar master solution. Serial dilutions from the master solution were made. The period between the single dilution steps was kept as short as possible (generally less than 30 seconds). At least 4 hours after attachment into MTP, cells were then treated with the various compound dilutions.

[0253] Plates were kept overnight in the cell culture incubator. The next day, a solution containing BSO was added to the wells, in a manner similar to that described in Jauslin et al., Hum. Mol. Genet. 11(24):3055 (2002), Jauslin et al., FASEB J. 17:1972-4 (2003), and

International Patent Application WO 2004/003565. Forty-eight hours later, three plates were examined under a phase-contrast microscope to verify that the cells in the negative control (wells E1-H1) were clearly dead. The medium from all plates was discarded, and the remaining liquid was removed by gently tapping the plate inversed onto a paper towel. The plates were washed twice with 100uL of PBS containing Calcium and Magnesium.

[0254] 100 microliters of PBS +Ca +Mg containing 1.2 micromolar Calcein AM were then added to each well. The plates were incubated for 30 minutes at 37C. After that time fluorescence (excitation/emission wavelengths of 485 nm and 525 nm, respectively) was read on a Gemini fluorescence reader. Data was imported into Microsoft Excel (EXCEL is a registered trademark of Microsoft Corporation for a spreadsheet program) and ExcelFit was used to calculate the EC50 concentration for each compound.

[0255] The compounds were tested three times, i.e., the experiment was performed three times, the passage number of the cells increasing by one with every repetition.

[0256] The solvents (DMSO, ethanol, PBS) neither had a detrimental effect on the viability of non-BSO treated cells nor did they have a beneficial influence on BSO-treated fibroblasts even at the highest concentration tested (1%). None of the compounds showed auto-fluorescence. The viability of non-BSO treated fibroblasts was set as 100%, and the viability of the BSO- and compound-treated cells was calculated as relative to this value.

[0257] The following table summarizes the EC50 for (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

Disorder	EC50
Friedrich's Ataxia	+++

+++ indicates less than 100 nM

Example 13. Screening (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in Fibroblasts from Patients Having Various Oxidative Stress Disorders

[0258] (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide was tested using a screen similar to the one described in Example 12, and substituting FRDA cells with cells from patients having other oxidative stress disorders.

[0259] The following table summarizes the EC₅₀ for (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide for various disorders.

Disorder	Cell Line Tested	EC ₅₀
Leigh Syndrome	Coriell Cell Repositories (Camden, NJ; repository number GMOI503A)	+++
Leber's Hereditary Optic Neuropathy (LHON)	Coriell Cell Repositories (Camden, NJ; repository number GM03858)	+++
Parkinson's Disease	Coriell Cell Repositories (Camden, NJ; repository number AG20439)	++
Huntington's Disease	Coriell Cell Repositories (Camden, NJ; repository number GM 04281)	+++
Rett's Disorder	Coriell Cell Repositories (Camden, NJ; repository number GM-17567)	+++
CoQ10 Deficiency	From patients having a CoQ2 mutation	+++
Amyotrophic Lateral Sclerosis (ALS)	Coriell Cell Repositories (Camden, NJ; repository number ND29523)	+++

+++ indicates less than 100 nM; ++ indicates 100-500 nM

Example 14. Screening (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide for Protection from Cisplatin-Induced Ototoxicity

[0260] The conditionally immortalized auditory HEI-OC1 cells from long-term cultures of transgenic mice ImmortomouseTM cochleas as described in Kalinec, G. et al., *Audiol.*

Nerootol. 2003; 8, 177-189/. were maintained in high glucose Dulbecco's modified Eagle medium (DMEM) containing 10% FBS under permissive conditions, 33°C, 10%CO₂. Cells were pretreated overnight with (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, and apoptosis was detected by caspase3/7 activity after 24 hours of 50 uM cisplatin incubation. Cells incubated in diluent alone were the controls.

[0261] The following table summarizes the EC₂₅ for (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide.

Disorder	EC ₂₅
Cisplatin-induced ototoxicity of auditory cells	+++

+++ indicates less than 100 micromolar

Example 15. Screening Polymorphic and Amorphous Compositions of the Invention in Fibroblasts from Patients

[0262] Polymorphic and amorphous compositions of the invention are tested using screens similar to the ones described in Examples 12-14, substituting the polymorphic or amorphous form for (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide. Polymorphic and amorphous compositions of the invention are also tested using screens similar to the ones described in Examples 12-14, and where appropriate substituting the FRDA cells or other cell lines with cells obtained from patients having an oxidative stress disorder described herein (e.g. MERRF, MELAS, KSS, Alzheimer's disease, a pervasive development disorder (such as autism), etc). The compositions are tested for their ability to rescue human dermal fibroblasts from these patients from oxidative stress or for their ability to protect cells from cisplatin-induced toxicity.

Example 16. Administration of compositions of the invention

[0263] A composition of the invention is presented in a capsule containing 300 mg of (*R*)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide in a pharmaceutically acceptable carrier. A capsule is taken orally, once a day.

[0264] The disclosures of all publications, patents, patent applications and published patent applications referred to herein by an identifying citation are hereby incorporated herein by reference in their entirety.

[0265] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is apparent to those skilled in the art that certain minor changes and modifications will be practiced. Therefore, the description and examples should not be construed as limiting the scope of the invention.

CLAIMS

What is claimed is:

1. A polymorph of an anhydrate or a hydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, wherein the polymorph is selected from the group consisting of Form I, Form III and Form II; wherein a powder X-ray diffraction pattern for polymorph Form I comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 17.03, and 17.26; wherein a powder X-ray diffraction pattern for polymorph Form III comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 14.02, 15.23, and 21.10; wherein a powder X-ray diffraction pattern for polymorph Form II comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, and 19.33; and wherein the XRPD pattern is generated at 25 °C.

2. The polymorph of claim 1, wherein the polymorph is Form I, wherein a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 17.03, and 17.26.

3. The polymorph of claim 1, wherein the polymorph is Form I, wherein a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 12.06, 15.33, 17.03, 17.26, and 18.72.

4. The polymorph of claim 1, wherein the polymorph is Form I, wherein the polymorph has a powder x-ray diffraction pattern substantially as shown in Figure 10.

5. A composition comprising the polymorph of any one of claims 2-4, wherein the composition is essentially free of Forms II-VI, wherein a powder X-ray diffraction pattern for polymorph Form V comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, and 15.45; wherein a powder X-ray diffraction pattern for polymorph Form III comprises characteristic peaks at least at the

following angular positions, wherein the angular positions may vary by ± 0.2 : 14.02, 15.23, and 21.10; wherein a powder X-ray diffraction pattern for polymorph Form II comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, and 19.33; wherein a powder X-ray diffraction pattern for polymorph Form IV comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 12.97, and 13.20; and wherein a powder X-ray diffraction pattern for polymorph Form VI comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.91, and 12.94; and wherein the XRPD pattern is generated at 25 °C.

6. A composition comprising the polymorph of any one of claims 2-4, wherein at least about 95% by mole of the composition is the polymorph Form I, exclusive of any solvents, carriers or excipients.

7. A pharmaceutical composition comprising the polymorph of any one of claims 2 to 4, or the composition of any one of claims 5-6, and a pharmaceutically acceptable carrier.

8. The pharmaceutical composition of claim 7, wherein the pharmaceutical composition has an HPLC purity of more than about 95% for the anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, exclusive of any solvents, carriers or excipients.

9. The pharmaceutical composition of claim 7, wherein the pharmaceutical composition has an HPLC purity of more than about 99% for the anhydrate of (R)-2-hydroxy-2-methyl-4-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dienyl)butanamide, exclusive of any solvents, carriers or excipients.

10. Use of the polymorph of any one of claims 2-4, the composition of any one of claims 5-6, or the pharmaceutical composition of any one of claims 7-9 in the manufacture of a medicament for treating or suppressing an oxidative stress disorder.

11. The use of claim 10, wherein the medicament is for treating or suppressing an oxidative stress disorder selected from the group consisting of: a mitochondrial disorder; an inherited mitochondrial disease; Alpers Disease; Barth syndrome; a Beta-oxidation Defect; Carnitine-Acyl-Carnitine Deficiency; Carnitine Deficiency; a Creatine Deficiency Syndrome; Co-Enzyme Q10 Deficiency; Complex I Deficiency; Complex II Deficiency; Complex III Deficiency; Complex IV Deficiency; Complex V Deficiency; COX Deficiency; chronic progressive external ophthalmoplegia (CPEO); CPT I Deficiency; CPT II Deficiency; Friedreich's Ataxia (FA); Glutaric Aciduria Type II; Kearns-Sayre Syndrome (KSS); Lactic Acidosis; Long-Chain Acyl-CoA Dehydrogenase Deficiency (LCAD); LCHAD; Leigh Syndrome; Leigh-like Syndrome; Leber's Hereditary Optic Neuropathy (LHON); Lethal Infantile Cardiomyopathy (LIC); Luft Disease; Multiple Acyl-CoA Dehydrogenase Deficiency (MAD); Medium-Chain Acyl-CoA Dehydrogenase Deficiency (MCAD); Mitochondrial Myopathy, Encephalopathy, Lactacidosis, Stroke (MELAS); Myoclonic Epilepsy with Ragged Red Fibers (MERRF); Mitochondrial Recessive Ataxia Syndrome (MIRAS); Mitochondrial Cytopathy, Mitochondrial DNA Depletion; Mitochondrial Encephalopathy; Mitochondrial Myopathy; Myoneurogastrointestinal Disorder and Encephalopathy (MNGIE); Neuropathy, Ataxia, and Retinitis Pigmentosa (NARP); Pearson Syndrome; Pyruvate Carboxylase Deficiency; Pyruvate Dehydrogenase Deficiency; a POLG Mutation; a Respiratory Chain Disorder; Short-Chain Acyl-CoA Dehydrogenase Deficiency (SCAD); SCHAD; Very Long-Chain Acyl-CoA Dehydrogenase Deficiency (VLCAD); a myopathy; cardiomyopathy; encephalomyopathy; a neurodegenerative disease; Parkinson's disease; Alzheimer's disease; amyotrophic lateral sclerosis (ALS); a motor neuron disease; a neurological disease; epilepsy; an age-associated disease; macular degeneration; diabetes; metabolic syndrome; cancer; brain cancer; a genetic disease; Huntington's Disease; a mood disorder; schizophrenia; bipolar disorder; a pervasive developmental disorder; autistic disorder; Asperger's syndrome; childhood disintegrative disorder (CDD); Rett's disorder; PDD-not otherwise specified (PDD-NOS); a cerebrovascular accident; stroke; a vision impairment; optic neuropathy; dominant inherited juvenile optic atrophy; optic neuropathy caused by a toxic agent; glaucoma; Stargardt's macular dystrophy; diabetic retinopathy; diabetic maculopathy; retinopathy of prematurity; ischemic reperfusion-related retinal injury; oxygen poisoning; a haemoglobinopathy; thalassemia; sickle cell anemia; seizures; ischemia; renal tubular acidosis; attention deficit/hyperactivity disorder (ADHD); a neurodegenerative disorder resulting in hearing or balance impairment; Dominant

Optic Atrophy (DOA); Maternally inherited diabetes and deafness (MIDD); chronic fatigue; contrast-induced kidney damage; contrast-induced retinopathy damage; Abetalipoproteinemia; retinitis pigmentosa; Wolfram's disease; Tourette syndrome; cobalamin c defect; methylmalonic aciduria; glioblastoma; Down's syndrome; acute tubular necrosis; a muscular dystrophy; a leukodystrophy; Progressive Supranuclear Palsy; spinal muscular atrophy; hearing loss; noise induced hearing loss; traumatic brain injury; Juvenile Huntington's Disease; Multiple Sclerosis; NGLY1; Multiple System Atrophy; Adrenoleukodystrophy; and Adrenomyeloneuropathy.

12. The use of claim 11, wherein the oxidative stress disorder is amyotrophic lateral sclerosis (ALS).

13. The use of claim 11, wherein the oxidative stress disorder is Parkinson's disease.

14. The polymorph of claim 1, wherein the polymorph is Form I, having a differential scanning calorimetry (DSC) thermogram substantially as shown in Figure 12.

15. The polymorph of claim 14, wherein the DSC thermogram has a single endothermic peak at about 152.9 °C.

16. The polymorph of claim 1, wherein the polymorph is Form I, having a thermogravimetric analysis (TGA) thermogram substantially as shown in Figure 13.

17. The polymorph of claim 1, wherein the polymorph is Form I, having a ¹H NMR spectrum substantially as shown in Figure 40.

18. A composition comprising the polymorph of any one of claims 14-17, wherein the composition is essentially free of Forms II-VI wherein a powder X-ray diffraction pattern for polymorph Form V comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.61, 11.49, and 15.45; wherein a powder X-ray diffraction pattern for polymorph Form III comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 14.02, 15.23, and 21.10; wherein a powder X-ray diffraction pattern for polymorph Form II comprises

characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 9.63, 11.33, and 19.33; wherein a powder X-ray diffraction pattern for polymorph Form IV comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.31, 12.97, and 13.20; and wherein a powder X-ray diffraction pattern for polymorph Form VI comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 6.27, 9.91, and 12.94; and wherein the XRPD pattern is generated at 25 °C.

19. A composition comprising the polymorph of any one of claims 14-17, wherein at least about 95% by mole of the composition is the polymorph Form I, exclusive of any solvents, carriers or excipients.

20. The polymorph of claim 1, wherein the DSC thermogram has a single endothermic peak.

21. The use of claim 11, wherein the oxidative stress disorder is Huntington's Disease.