

EXTRACT OF AN HERBAL COMPOSITION AS ANTIMICROBIAL AND/OR ANTIBIOFILM AGENT

FIELD OF THE INVENTION

The invention relates to an extract of an herbal composition comprising at least two different dried plants useful as antimicrobial and/or antibiofilm agent. It has also been found that in such extract, the active ingredients exert their biological effects in a synergistic manner.

BACKGROUND OF THE INVENTION

Like any living organism, plants contain numerous chemical entities, comprising low molecular weight chemicals, polypeptides, poly-saccharides, enzymes. A plant extract is thus a trivial example of naturally occurring mix of chemical entities.

The biological and therapeutic activities of many chemical entities mixes obtained from plants is of secular knowledge and is compiled in numerous manuscripts (see for example, Encyclopedia of Herbal Medicine: The Definitive Home Reference Guide to 550 Key Herbs with all their Uses as Remedies for Common Ailments Hardcover – November 15, 2000, by Andrew Chevallier).

Modern science allows to observe biological and therapeutic effects of chemical entities combinations obtained from plants on a large number of organisms.

For instance, it is described in Scheggi (Scheggi S. et al, Pharm. Biol. 2016 Jan 5:1-11. [Epub ahead of print]) how a rat model allows to test the effect against depression of chemical entities combinations obtained from plants.

In Zhipeng QU (Oncotarget. 2016 Sep 1 Epub) it is shown how it is possible to identify candidate anti-cancer molecular mechanisms of Compound Kushen Injection using functional genomics. Compound Kushen Injection (CKI) has been clinically used in China for over 15 years to treat various types of solid tumours. However, because such Traditional Chinese Medicine (TCM) preparations are complex mixtures of plant secondary metabolites, it is essential to explore their underlying molecular mechanisms in a systematic fashion. This study shows that CKI inhibited MCF-7 cell proliferation and induced apoptosis in a dose-dependent fashion. Multiple pathways were found to be perturbed and the cell cycle was identified as the potential primary target pathway of CKI in MCF-7 cells. CKI may also induce apoptosis in MCF-7 cells via a p53 independent mechanism. In addition, they identified novel lncRNAs and showed that many of them might be expressed as a response to CKI treatment.

The effect of chemical entities mixtures obtained for instance from plants, fungi or bacteria on the growth of microorganisms such as bacteria and fungi has been described many times, for instance in Venkatadri (Venkatadri B. *et al.*, Indian J. Pharm. Sci. 2015 Nov-Dec;77(6):788-91).

The effect of chemical entities combinations obtained for instance from plants, fungi or bacteria on the formation of biofilms (communities of microorganisms embedded in a polymeric matrix attached to a surface) has also been described many times, for instance in Süntar (Süntar I. et al., Pharm. Biol. 2016 Jun;54(6):1065-70Epub 2015 Oct 29).

Shimizu M. *et al.* (Antimicrob Agents Chemother. 2001 Nov;45(11):3198-201) found that an extract of *Arctostaphylos uva-ursi* markedly reduced the MICs (Minimum Inhibitory Concentrations) of beta-lactam antibiotics, such as oxacillin and cefmetazole, against methicillin-resistant *Staphylococcus aureus*. The same group also isolated the effective compound and identified it as corilagin.

Another group (Denev P. *et al.*, Acta Biochim. Pol. 2014;61(2):359-67. Epub 2014 Jun 18) studied the antioxidant, antimicrobial and neutrophil-modulating activities of extracts from six medicinal plants--

blackberry (*Rubus fruticosus*) leaves, chokeberry (*Aronia melanocarpa*) leaves, hawthorn (*Crataegus monogyna*) leaves, lady's mantle (*Alchemilla glabra*) aerial parts, meadowsweet (*Filipendula ulmaria*) aerial parts and raspberry (*Rubus idaeus*) leaves. The antimicrobial activity of the investigated extracts against 11 human pathogens was investigated using three different methods and the results of this study allowed concluding that meadowsweet and blackberry leaves extracts had the highest antimicrobial effect and the lowest minimal inhibiting concentrations (MICs) against the microorganisms tested.

Yam T. *et al.* (FEMS Microbiol Lett. 1997 Jul 1;152(1):169-74) focused their study on the microbiological activity of whole and fractionated crude extracts of tea (*Camellia sinensis*), and of tea components. Aqueous extracts of teas (*Camellia sinensis*) of different types and from various sources were tested and found to inhibit a wide range of pathogenic bacteria, including methicillin-resistant *Staphylococcus aureus*. Tea extracts were bactericidal to staphylococci and *Yersinia enterocolitica* at well below 'cup of tea' concentrations. Activity was confined to one of four fractions obtained from a green tea extract by partition chromatography. Testing of pure tea compounds and closely related chemicals suggested that the antibacterial activity of extracts of green tea can be explained by its content of epigallocatechin, epigallocatechin gallate and epicatechin gallate. In black tea extracts, theaflavin and its gallates are additional antibacterially active components.

WO 2009/031041 reports a composition comprising (a) an antimicrobial compound having a given chemical structural formula, (b) an antimicrobial material selected from lanthionine bacteriocins, tea [*Camellia sinensis*] extract, hop [*Humulus lupulus* L] extract, grape skin extract, grape seed extract, Uva Ursi [*Arctostaphylos uva-ursi*] extract and combinations thereof.

WO 99/20289 reports a clear herbal extract solution comprising a concentrated herbal extract and a fill liquid. The concentrated herbal extract is reported to be selected from a long list of plants extracts, where such list includes *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi*. However, WO 99/20289 does not mention any specific therapeutic or medical use of the claimed herbal solution.

US 2015/0056255 reports a product for medicinal, cosmetic, coloring, or dermatologic use, the product comprising a layer of a fibrous plant product and a plant extract applied to the layer of the fibrous plant product. The plant extract is said to comprise substances from one or more specific parts of one or more plants, wherein the plants are selected from at least one of a long list of plants.

Azanida NN (Med. Aromat. Plants, 4:3, p.1-6, 2015) published a review, which reports a description and a comparison of the most commonly used methods for preparing medicinal plant extracts. The authors conclude their article by stating that "no universal extraction method is the ideal method and each extraction procedure is unique to the plants" (see "*Conclusions*", lines 5-7).

Selman A. Wachsman *et al.* (Bacteriology, 31, p.157-164, 1945) reports that different strains of the same bacterial species may vary greatly in their resistance to streptomycin. Different *Staphylococcus aureus* bacterial strains are used in such paper to test the activity of streptomycin. However, no plant extract is tested on the same strains and therefore no clear guidance can be drawn from such paper.

Biological effects such as contemplated in the previous examples can be induced either by a single chemical entity, such as in standard drugs, or by a synergistic effects of different chemical entities, such as in extracts. In some cases, the biological effect is nearly fully attributable to a single chemical entity contained in the extract. In this case, the extract can be replaced by the purified active chemical entity.

Biological effects of a chemical entity can be either due to chemical reactions (transformations) with the chemical entities constituting the organism or be due to the transient or permanent binding to the chemical entities, in particular the macromolecules, constituting the organism. Such binding will modulate the activity of the organism's chemical entities, in particular the macromolecules, e.g., increase or decrease the activity of an enzyme.

Unfortunately, chemical entities have never a single effect on an organism. In particular, the interactions with the macromolecular constituents are numerous. As a matter of fact, it is well established that any chemical entity, whether natural or resulting from chemical synthesis, will have deleterious effects on organisms (humans, animals, plants, micro-flora, etc.) in a dose related fashion. Eventually, a dose related positive or therapeutic effect can be observed also. It is obvious that for the chemical entity to be of industrial interest, the dose required to observe the positive or therapeutic effect needs to be lower than the dose for which the deleterious effects becomes unacceptable.

Chemical entities differing in their chemical composition or structure will have different affinities (or binding constants) with the different macromolecules constituting the organism, and thus different modes of actions resulting in the observed deleterious, positive and therapeutic effects.

The macromolecules involved in deleterious effects and those involved in positive or therapeutic effects are different in many cases. Chemical entities differing in their chemical composition will in most of the cases have very different affinities (or binding constants) with the numerous different macromolecules constituting the organism.

In most cases, different chemical entities will have different affinities (or binding constants) with a given macromolecule. In rare cases, different chemical entities will have similar affinities (or binding constants) with the same macromolecule or sets of macromolecules involved in a positive or therapeutic effect. Among these cases, it will be rare that these different chemical entities will have comparable affinities (or binding constants) with the macromolecules involved in given deleterious effect.

Mixes of different chemical entities may thus have a synergistic positive or therapeutic effect without a cumulative given deleterious effect. Thus, for a given positive or therapeutic effect, the dose required for each individual different chemical entity in a mix will be lower than the dose required by each of the chemical entity considered individually to obtain that given positive or therapeutic effect. In those cases where the deleterious effects induced by each different chemical compound is different or has a different macromolecular aetiology, each deleterious effect of the mix will be lower than the deleterious effect of the chemical entity considered individually because it is used at a lower dose in the mix.

Note that the synergistic situation of the simultaneous administration of reduced doses of different chemical entities is different from the co-administration of different drugs (chemical entities) having different therapeutic effects administered at their standard doses. In the latter situation, increased side effects due to drug-interactions are well known. One of the cause of side effects is the possible chemical interaction of the different drugs (chemical entities). In the synergistic situation, each chemical entity is present at a reduced concentration and thus with much lower rates of chemical interactions. For illustration purposes, a synergistic mix of 10 different chemical entities (drugs) administrated each at $1/10^{\text{th}}$ of the concentration (dose) required for a chemical entity (drug) administrated individually, the rate of inter-drug chemical reactions will be reduced by a factor of 100 as implied by the universal law of action of mass.

It is therefore of interest to use synergistic mixes of different chemical entities in order to obtain a given positive or therapeutic effect with reduced deleterious effects, or to obtain an increased positive or therapeutic effect with a given level of different deleterious effects. Indeed, several extracts are known have synergistic activities relative to isolated chemical entities (see for instance, Deharo E. *et al.*, Malar. J. 2011 Mar 15;10 Suppl 1:S5).

Natural extracts composed of a mix of synergistic chemical entities can be discovered for example by means of systematic testing numerous natural extracts of single plant powders or herbal compositions containing at least two or more different plant powders.

DESCRIPTION OF THE INVENTION

It has now been found that an extract of an herbal composition comprising at least two dried plants selected among *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi* is useful as antimicrobial and/or antibiofilm agent in the treatment or prevention of microbial infections caused by bacteria, such as for example *Escherichia*, *Klebsiella*, *Listeria*, *Pseudomonas*, *Salmonella*, *Streptococcus* or *Staphylococcus*, or by fungi, such as for example *Candida*. It has also been found that in such extract, the active ingredients exert their biological effects in a synergistic manner.

Therefore, one object of the present invention is a method for the treatment or prevention of the above mentioned microbial infections comprising administering an effective amount of an extract of an herbal composition comprising at least two dried plants selected among *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi*, a lyophilisate of said extract, or a pharmaceutical or veterinary composition comprising said extract or said lyophilisate, to a subject in the need thereof.

Another object of the present invention is an extract of an herbal composition, comprising at least two dried plants selected among *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi* and at least one plant selected among *Vitis vinifera* var. *tinctoria*, *Eugenia caryophyllus* and *Desmodium adscendens*.

Said herbal composition may further comprise one or more additional dried plants selected among:

Achillea millefolium, *Acorus calamus*, *Agrimonia eupatoria*, *Agropyrum repens*, *Agropyrum repens*, *Alchemilla vulgaris*, *Alkanna tinctoria*, *Althaea officinalis*, *Anethum graveolens*, *Angelica archangelica*, *Arbutus unedo*, *Arnica montana*, *Artemisia pontica*, *Artemisia vulgaris*, *Asparagus officinalis*, *Asparagus officinalis*, *Asperula odorata*, *Betula pendula*, *Borrigo officinalis*, *Buxus sempervirens*, *Calamintha officinalis*, *Calendula officinalis*, *Calluna vulgaris*, *Carum carvi*, *Cassia angustifolia*, *Centaurea cyanus*, *Centaureum erythraea*, *Centella asiatica*, *Cetraria islandica*, *Chamaemelum nobile*, *Chamomilla recutita*, *Chrysanthellum americanum*, *Cichorium endivia*, *Cichorium intybus*, *Cinnamomum zeylanicum*, *Citrus aurantium*, *Combretum micranthum*, *Crataegus oxyacantha*, *Cuminum cyminum*, *Cupressus sempervirens*, *Curcuma zedoaria*, *Cynara scolymus*, *Cytisus scoparius*, *Elettaria cardamomum*, *Eleutherococcus senticosus*, *Epilobium parviflorum*, *Erysimum officinale*, *Eucalyptus globulus*, *Eupatorium cannabinum*, *Foeniculum vulgare*, *Fraxinus excelsior*, *Fucus vesiculosus*, *Fumaria officinalis*, *Galium odoratum*, *Gentiana lutea*, *Geranium robertianum*, *Ginkgo biloba*, *Glechoma hederacea*, *Glycyrrhiza glabra*, *Handroanthus impetiginosus*, *Harpagophytum procumbens*, *Hieracium pilosella*, *Humulus lupulus*, *Hypericum perforatum*, *Hyssopus officinalis*, *Illicium verum*, *Inula helenium*, *Juglans regia*, *Juniperus communis*, *Lamium album*, *Lavandula angustifolia*, *Levisticum officinale*, *Lippia citriodora*, *Lotus corniculatus*, *Lythrum salicaria*, *Malva sylvestris*, *Marrubium vulgare*, *Medicago sativa*, *Melissa officinalis*, *Mentha × piperita*, *Morus nigra*, *Myrtus communis*, *Olea europaea*, *Origanum majorana*, *Panax ginseng*, *Papaver rhoeas*, *Parietaria officinalis*, *Passiflora incarnata*, *Petroselinum crispum*, *Peumus boldus*, *Phaseolus vulgaris*, *Pimpinella*

anisum, Plantago lanceolata, Plantago ovata, Potentilla anserina, Quercus robur, Rhamnus frangula, Rheum palmatum, Rosa centifolia, Rosmarinus officinalis, Rubia tinctorum, Rubus idaeus, Salix alba, Salvia officinalis, Sambucus nigra, Satureja montana, Silybum marianum, Solanum dulcamara, Tabebuia impetiginosa, Tanacetum vulgare, Taraxacum officinalis, Thymus serpyllum, Thymus vulgaris,
 5 *Tilia tomentosa, Tilia cordata, Trigonella foenum-graecum, Tussilago farfara, Vaccinium myrtillus, Valeriana officinalis, Verbascum thapsus, Verbena officinalis, Viscum album, Zea mays and Zingiber officinale.*

A further object of the invention is an extract of an herbal composition comprising from three to seven dried plants selected among *Filipendula ulmaria, Camellia sinensis, Arctostaphylos uva-ursi, Rheum palmatum, Rosmarinus officinalis, Vitis vinifera tinctoria, Desmodium adscendes, Eugenia caryophyllus*
 10 *and Eucalyptus globulus*, wherein at least two of such plants are selected among *Filipendula ulmaria, Camellia sinensis* and *Arctostaphylos uva-ursi* and at least one of such plants is selected among *Vitis vinifera tinctoria, Desmodium adscendes* and *Eugenia caryophyllus*.

Another object of the invention is an extract of an herbal composition comprising the following dried
 15 plants: *Rosmarinus officinalis, Filipendula ulmaria, Camellia sinensis, Vitis vinifera tinctoria* and *Arctostaphylos uva-ursi*.

Another object of the invention is an extract of an herbal composition comprising the following dried plants: *Rheum palmatum, Filipendula ulmaria, Camellia sinensis, Vitis vinifera tinctoria* and *Arctostaphylos uva-ursi*.

Another object of the invention is an extract of an herbal composition comprising the following dried
 20 plants: *Rheum palmatum, Rosmarinus officinalis, Filipendula ulmaria, Camellia sinensis, Vitis vinifera tinctoria, Eugenia caryophyllus* and *Arctostaphylos uva-ursi*.

A further object of the invention is an extract of an herbal composition comprising the following dried
 25 plants: *Rheum palmatum, Rosmarinus officinalis, Filipendula ulmaria, Camellia sinensis, Vitis vinifera tinctoria, Eucalyptus globulus, Arctostaphylos uva-ursi, Mentha spicata* and *Rubia tinctorum*.

Another object of the invention is an extract of an herbal composition comprising at least fourteen or
 fifteen dried plants selected among the following sixteen: *Rheum palmatum, Rosmarinus officinalis, Filipendula ulmaria, Satureja montana, Valeriana officinalis, Camellia sinensis, Vitis vinifera tinctoria, Fucus vesiculosus, Foeniculum vulgare, Arctostaphylos uva-ursi, Arbutus unedo, Eugenia caryophyllus,*
 30 *Juniperus communis, Combretum micranthum, Lippia citrodora* and *Tanacetum vulgare*

Another object of the invention is an extract of an herbal composition comprising the following dried
 plants: *Rheum palmatum, Rosmarinus officinalis, Filipendula ulmaria, Satureja montana, Valeriana officinalis, Camellia sinensis, Vitis vinifera tinctoria, Fucus vesiculosus, Foeniculum vulgare, Arctostaphylos uva-ursi, Arbutus unedo, Eugenia caryophyllus, Juniperus communis, Combretum micranthum* and *Lippia citrodora*.
 35

Another object of the invention is an extract of an herbal composition comprising the following dried
 plants: *Rheum palmatum, Rosmarinus officinalis, Filipendula ulmaria, Satureja montana, Valeriana officinalis, Camellia sinensis, Vitis vinifera tinctoria, Fucus vesiculosus, Foeniculum vulgare, Arctostaphylos uva-ursi, Arbutus unedo, Eugenia caryophyllus, Juniperus communis, Combretum micranthum, Lippia citrodora* and *Tanacetum vulgare*.
 40

All the above-mentioned herbal compositions are also an object of the invention, as intermediates in the preparation of the corresponding extracts.

The extract may constitute the active ingredient of a food supplement, a nutraceutical, pharmaceutical or cosmetic composition or a functional food or a food additive.

The extract of the invention may be used as such or in a dried or lyophilized form.

5 A food supplement, a nutraceutical, pharmaceutical, veterinary or cosmetic composition or a functional food or a food additive containing the extract defined above together with suitable excipients are further objects of the invention.

10 Such food supplement, a nutraceutical, pharmaceutical, veterinary or cosmetic composition or a functional food or a food additive may be formulated in any form suitable, for example, for oral administration, including solid forms such as powders, granules, capsules, pills, tablets (normal or controlled release), chewing gums, or liquid forms such as syrups, drops, elixirs, solutions and suspensions in general.

15 Furthermore, the extract according to the invention can be incorporated into other formulations, such as creams, ointments, gels, milks, pastes, creams based on water, emulsions, serums, masks, balms, fluids, sprays, suppositories, vaginal suppositories, transdermal patches and toothpastes, periodontal gels, mouthwashes.

According to the present invention, the expression "herbal composition" refers to a powder containing one or more different grinded dried plants. The powder of a single dried plant may have been obtained from the whole plant or from a specific part of it, for example, flowers, buds, fruits, stems, leaves, seeds, roots or other.

20 According to the invention, an extract may be obtainable by simple incubation of an herbal composition with an extraction solvent (with or without the addition of a sugar) at room temperature.

25 Suitable extraction solvents according to the invention are water and all non-aqueous solvents authorized for use in the production of foodstuffs and food ingredients. For example, according to Directive 2009/32/EC a list of such non-aqueous solvents include propane, butane, ethyl acetate, ethanol, carbon dioxide, acetone, nitrous oxide. According to specific embodiments of the invention, the extraction solvent is selected among water, ethanol, ethyl acetate and any mixture thereof.

According to another alternative embodiment, an extract may be obtainable by incubation of an herbal composition with an extraction solvent (with or without the addition of a sugar) and subsequent heating, with or without centrifugation and filtration.

30 According to an alternative embodiment, a water extract may be obtainable by incubation of an herbal composition with water (with or without the addition of a sugar) and subsequent heating (for example by heating the resulting liquid preparation at a temperature ranging from 60°C to 134°C, preferably from 119° to 121°C, for a time ranging from 5 to 60 minutes, preferably from 20 to 30 minutes).

35 With the term "sugar", we herein intend to mean a pharmaceutical or alimentary grade sugar, such as for example dextrose, lactose, anhydrous lactose, mannitol, sorbitol, maltose, galactose or sucrose.

According to a particular embodiment of the invention, such sugar is sucrose.

Said sugar may be used at concentrations varying from 1 to 100 g/l, optionally from 50 to 75 g/l.

40 With the term "food supplement" it is here meant a food product intended to supplement the common diet and which is a concentrated source of nutrients, such as vitamins and minerals, or other substances with a nutritional or physiological effect, in unit dose form (see also Directive 2002/46 / EC of 10 June 2002).

The expression "functional food" means to include foods characterized by additional effects due to the presence of components (generally non-nutrients) naturally present or added, which interact more or less selectively with one or more physiological functions of the organism (biomodulation) leading to a positive effect on maintaining health and / or disease prevention. A food can be considered 'functional', if it is sufficiently demonstrated its beneficial influence on one or more functions of the body, in addition to adequate nutritional effects, so as to be relevant to a state of well-being and health, or for risk reduction of an illness. The beneficial effects could include both in maintaining that in promoting a state of well-being or health and / or a reduction in the risk of a disease process or disease (see AT Diplock et al: Scientific concepts of functional foods in Europe: Consensus Document, British Journal of Nutrition 1999, 81 (Suppl. 1), S1-S27).

According to the present invention with the expression, "nutraceutical composition" it is intended to mean a product, which not only supplement the diet, but should also aid in the prevention and/or treatment of disease and/or disorder. This expression was coined in 1989 by Stephen L. DeFelice (MD, founder and chairperson of the Foundation for Innovation in Medicine (FIM)) joining the concepts of "nutrition" and "pharmaceutical".

With the term "excipient", it is intended here to refer to conventional excipients, i.e. inert compounds towards the active ingredient of a composition.

Excipients may be divided into various functional classifications, depending on the role that they are intended to play in the resultant formulation, e.g. in solid dosage forms they may act as diluents (e.g. lactose, microcrystalline cellulose), disintegrants (e.g. sodium starch glycolate, croscarmellose sodium), binders (e.g. PVP, HPMC), lubricants (e.g. magnesium stearate), glidants (e.g. colloidal SiO₂) or flavoring agents (e.g. peppermint, lemon oils). In liquid dosage forms they may act as solvents, co-solvents or solubilizers (e.g. propylene glycol, glycerol, ethanol, sorbitol, dextrose), buffering agents (e.g. sodium phosphate), antimicrobial agents (e.g. potassium sorbate), wetting agents (e.g. polysorbates or Tweens), anti-foaming agents (e.g. polydimethylsiloxane-silicon dioxide), thickening agents (e.g. methylcellulose or hydroxyethylcellulose) or sweetening agents (e.g. sorbitol, acesulfame-K).

As already said, the extract of the invention is an antimicrobial and/or antibiofilm agent and it is advantageously used in the treatment, the prevention and/or the diagnosis of microbial infections in humans and animals.

Another object of the invention is therefore the use of the extract of the invention in the manufacture of a medicament for the treatment, prevention or diagnosis of microbial infections. Such medicament may be useful for humans of animals.

According to the present invention, the expressions "treatment" or "treating" are intended to activities designed to cure, mitigate the symptoms or delay the progression of a disease or a pathological condition.

The expressions "prevention" or "preventing" are intended to refer to activities designed to minimize the incidence or effects of a disease or a pathological condition. For example in the case of microbial infections, prevention specifically herein means to prevent the adhesion and colonization of pathogenic microbes to cells, tissues and organs.

The expressions "diagnosis" or "diagnosing" are intended to refer to activities designed to identify a series of physiological parameters in order to assess the presence and the extent of a determined disease or pathological condition.

The expression “antimicrobial agent” will here be used to mean an agent that is able to inhibit the growth of microbial cells. If the microbial cells are pathogenic microbial cells, the antimicrobial agent can be used as active ingredient of a pharmaceutical or veterinary composition for the treatment of microbial infections in humans or animals.

- 5 The expression “antibiofilm agent” will here be used to mean an agent that is able to inhibit the formation of biofilm by microbial cells and/or to facilitate the dispersion of preformed biofilms. In contrast to conventional antimicrobial agents, an antibiofilm agent does not directly affect microbial survival and thus the expectation is that resistance to these agents will not readily occur.

10 According to the present invention, the term “synergistic” is intended as referred to a biological effect or activity (herein antimicrobial or antibiofilm), presented by the extract of an herbal composition of the invention (i.e. comprising at least two dried plants selected among *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi*), which is greater than the sum of the effects or activities of the extracts of the corresponding single dried plants.

15 Such greater biological effect or activity may be illustrated, for example, in terms of inhibition of planktonic growth (antimicrobial activity), in terms of biofilm formation inhibition (antibiofilm activity) or in terms of number of strains, in which one or both these activities can be identified.

20 According to a particular embodiment of the invention, the microbial infections are caused by bacteria, such as for example *Escherichia coli*, *Klebsiella pneumoniae*, *Listeria innocua*, *Listeria monocytogenes*, *Pseudomonas aeruginosa*, *Salmonella enterica subsp. enterica serovar Enteritidis*, *Salmonella enterica subsp. enterica serovar Typhimurium*, *Streptococcus pneumoniae*, *Streptococcus equi subsp. zooepidemicus*, *Streptococcus pyogenes*, or by fungi, such as for example *Candida albicans*.

According to another particular embodiment of the invention, the microbial infections are caused by *Staphylococcus aureus*, *Staphylococcus epidermidis* or *Staphylococcus pseudintermedius*.

25 *Staphylococcus aureus*, a commensal Gram-positive bacterium is a leading pathogen that may cause a wide variety of diseases in human ranging from skin and soft tissue infections to severe septicemia, toxic choc or pneumonia. *Staphylococcus aureus* is an important cause of hospital-associated infections, and many hospital strains of *Staphylococcus aureus* have acquired resistant to antibiotics, most particularly to methicillin. There is no vaccine available yet, but important efforts are conducted to develop active and passive immunotherapy against various targets of *Staphylococcus aureus*.

30 In order to evaluate such biological activities for the extract of the invention and for comparative samples, many different strains have been used and will be mentioned in the Examples' section.

In particular, the following bacterial strains have been used to test both the antimicrobial and antibiofilm activity of the extracts of the invention.

35 ATCC 25923, also known as Seattle 1945, is a *Staphylococcus aureus* strain widely used in academic research (it is cited in 627 publications as of October 2016 (see NIH-NCBI-Pubmed)). Its antibiotic resistance behavior has been thoroughly characterized (see, for instance, Reimer L.G. et al., Antimicrob. Agents Chemother. 1981 Jun;19(6):1050-5).

40 ATCC 29213 is also a *Staphylococcus aureus* strain widely used in academic research (cited in 327 publications as of October 2016 (see NIH-NCBI-Pubmed)). It has been used in biofilm research (see, for instance, Ceri H. et al., J. Clin. Microbiol. 1999 Jun;37(6):1771-6).

ATCC 1228 is a *Staphylococcus epidermidis* strain (cited in 101 publications as of October 2016 (see NIH-NCBI-Pubmed)). The Code of Federal Regulations (FDA) lists such strain as the preferred assay organism for neomycin (see also Robertson J.H. et al., Appl. Microbiol. 1971 Dec;22(6):1164-5).

ATCC 43300 is a *Staphylococcus aureus* strain isolated from a clinical study (cited in 81 publications as of October 2016 (see NIH-NCBI-Pubmed)), including Lubenko et al., J. Antimicrob. Chemother. 2008 Nov;62(5):1065-9).

Staphylococcus aureus strain Newman was isolated in 1952 from a human infection and has been used extensively in animal models of staphylococcal infections due to its robust virulence phenotype (see Baba T. et al., Journal of bacteriology, 190 :300-310 (2008)). In contrast to hospital-acquired MRSA (Methicillin-resistant *Staphylococcus aureus*), *Staphylococcus aureus* Newman carries a harbors only small a small number of insertion sequences (IS) and lacks known antibiotic resistance determinants.

Other strains used in the examples are also commonly used in academic research or in industry for quality control (QC). Most of them are also methicillin resistant.

For reference to ATCC 33591, see Reyes N. et al., J. Antimicrob. Chemother. 2006 Aug;58(2):462-5, Epub 2006 May 30. For reference to ATCC 33592, see Teka A. et al., BMC Complement Altern Med. 2015 Aug 18;15:286. For reference to NCTC 12493, see Carey B.E. et al., J. Antimicrob. Chemother. 2006 Aug;58(2):455-7, Epub 2006 Jun 20. For reference to ATCC 700698, see Kirker K.R. et al., Int. J. Antimicrob Agents, 2015 Oct;46(4):451-5, Epub 2015 Jul 9. For reference to ATCC 700699, see Sy C.L. et al., J. Clin. Microbiol. 2016 Mar;54(3):565-8, Epub 2015 Dec 16. For reference to ATCC 9144, see Carson C.F. et al., Antimicrob. Agents Chemother. 2002 Jun;46(6):1914-20. For reference to ATCC BAA-44, see Monecke S. et al., PLoS One. 2011 Apr 6;6(4):e17936. For reference to ATCC 14990, see Bernardo T.H. et al., Scientific World Journal. 2015;751791, Epub 2015 Jun 3). For reference to ATCC 49444, see Ramsey KJ et al., J Food Prot. (2010). For reference to ATCC 60193, see Petrikaite V et al., Medicina (Kaunas). 2007;43(8):657-63. For reference to ATCC 11775, see Usman H et al., Afr J Tradit Complement Altern Med. 2007 Jun 10;4(4):488-94. For reference to ATCC 13883, see Supardy NA et al., J Microbiol Biotechnol. 2012 Jun;22(6):872-81. For reference to ATCC 33090, see Le Marc Y et al., Int J Food Microbiol. 2002 Mar;73(2-3):219-37. For reference to ATCC 19115, see Vodnar DC., Chem Cent J. 2012 Jul 28;6(1):74. doi: 10.1186/1752-153X-6-74. For reference to ATCC 27853, see Reimer LG et al., Antimicrob Agents Chemother. 1981 Jun;19(6):1050-5. For reference to ATCC 13076, see Fioretto F et al., Braz J Med Biol Res. 2005 Aug;38(8):1259-65. Epub 2005 Jul 30. For reference to ATCC 13311, see Leguérinel I et al., Int J Food Microbiol. 2007 May 1;116(1):88-95. Epub 2007 Jan 13. For reference to ATCC 27336, see Chen S et al., Appl Environ Microbiol. 2013 Jul;79(13):4015-23. doi: 10.1128/AEM.00704-13. Epub 2013 Apr 19. For reference to ATCC 43079, see Farrow JA and Collins MD., Syst. Appl. Microbiol. 5: 483-493, 1984. For reference to ATCC BAA-1323, see Svensson Mikael D et al., Microbiology 148: 3933-3945, 2002.

The food supplement, the nutraceutical, pharmaceutical or cosmetic composition of the invention may be suitably formulated for different routes of administration. Such routes of administrations include: auricular, buccal, cutaneous, dental, nasal, ophthalmic, oral, oropharyngeal, rectal, respiratory (inhalation), sublingual, topical or vaginal.

Generally, the compositions comprising the extracts of the invention are administered in a "therapeutically effective amount".

For the aim of the present description and of the following claims, the term "comprising" also includes the terms "which essentially consists of" or "which consists of".

The amount of the composition actually administered will typically be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual composition administered, the age, body weight, and response of the individual subject, the severity of the subject's symptoms and the like. For any composition, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rats, guinea pigs, rabbits, dogs, or pigs.

The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. In calculating the Human Equivalent Dose (HED) it is recommended to use the conversion table provided in Guidance for Industry and Reviewers document (2005, U.S. Food and Drug Administration, Rockville, Maryland, USA).

According to the invention, the dose of the extract to be administered to humans may be regulated referring to the herbal composition used to prepare it and, consequently, the amount of grinded single dried plants present in the herbal composition. Usually, from 0.001 to 500 g of each grinded dried plant is used to prepare the final extract, optionally from 0.01 to 100 g. Such extract may be used as such or in a lyophilized, dried or freeze dried form in the final composition for human or animal use or it may be further diluted.

The precise effective dose for a human subject will depend upon the severity of the disease or condition, general health of the subject, age, weight and gender of the subject, diet, time and frequency of administration, other simultaneous therapies, reaction sensitivities, and tolerance/response to therapy. This amount can be determined by routine experimentation and is within the judgement of the clinician.

Dosage treatment may be a single dose schedule or a multiple dose schedule.

Another object of the invention is the process for preparing the extract of the invention. Such process schematically comprises the following steps:

- a) chopping or grinding separately or in a mixture at least two single dried plants to obtain corresponding plant powders;
- b) mixing at least two different plant powders to obtain an herbal composition;
- c) adding an extraction solvent to said herbal composition to obtain a corresponding liquid preparation (or extract);
- d) incubating said liquid preparation at room temperature for a time ranging between 5 and 15 minutes; optionally,
- e) heating said liquid preparation at a temperature ranging from 60°C to 134°C for a time ranging from 5 to 60 minutes; optionally,
- f) centrifuging, collecting the supernatant and filtering the collected supernatant; and, optionally,
- g) concentrating and/or drying or freeze drying the liquid preparation obtained from step e) or f).

According to the present application the word "lyophilisate" will refer to both freeze-dried (dehydrated) and dried extracts according to the invention.

The single dried plants are commercial products that may be purchased as part of the dried plants collections of pharmacies or herbal shops. Lists of dried plants collections are available at each

pharmacy of herbal shop of choice and the dried plants available in such collections may vary from country to country.

For example, Belgium, France and Italy, decided in 2012 to join forces to develop a common list of plants, whose use could be allowed, provided that manufacturers meet the quality requirements of European law (EC Regulation 852/2004). Thanks to the hard work of three recognized experts (Robert Anton, Luke Delmulle and Mauro Serafini) the project resulted in the establishment of a list of 1,029 plants and 11 mushrooms (see list BelFrit at

http://www.economie.gouv.fr/files/files/directions_services/dgccrf/imgs/breve/2014/documents/harmonized_list_Section_A.pdf)

The invention will now be described by making reference to the following non-limiting Examples.

EXAMPLES

List of plants used in the Examples

The plants and parts of plants in the following list have been used in the different Examples. For the sake of conciseness, each plant is identified by a code used throughout the Examples. The weight corresponding to 200 µl (volume of the calibrated spoon) of plant powder is also indicated (Table 0-1). The dried plants of pharmaceutical grade PA00 to PB12 were obtained from “Pharmacie Fontgèze” (Clermont-Ferrand, France) and the dried plants of pharmaceutical grade PB13 to PD08 were obtained from “Pharmacie St Herem” (Clermont-Ferrand, France). When indicated in the Example, plants were also obtained from “Herboristerie Cailleau” (Chemillé, France) or from Siccarapam (Aubiat, France).

Table 0-1: List of plants with associated codes and weights of dried powders for one calibrated spoon

Plant Code	Botanic Name	Part used	Weight (g)	Plant Code	Botanic Name	Part used	Weight (g)
PA00	<i>Sambucus nigra</i>	flower	0.0684	PB32	<i>Tabebuia impetiginosa</i>	bark	0.0775
PA01	<i>Plantago afra</i>	seed	0.1357	PB33	<i>Glechoma hederacea</i>	aerial part	0.0601
PA02	<i>Equisetum arvense</i>	aerial part	0.0578	PB35	<i>Cetraria islandica</i>	aerial part	0.0776
PA05	<i>Aloysia citrodora</i>	leaf	0.0712	PB36	<i>Rubus idaeus</i>	leaf	0.0553
PA06	<i>Glycyrrhiza glabra</i>	root	0.0774	PB38	<i>Fraxinus excelsior</i>	leaf	0.1241
PA07	<i>Camellia sinensis</i>	leaf	0.07	PB39	<i>Lavandula angustifolia</i>	flower	0.0207
PA08	<i>Solidago virg aurea</i>	aerial part	0.0394	PB41	<i>Cassia angustifolia</i>	fruit	0.0568
PA10	<i>Taraxacum officinale</i>	root	0.0585	PB42	<i>Fragaria vesca</i>	leaf	0.0692
PA11	<i>Rheum palmatum</i>	root and rhizome	0.1016	PB45	<i>Aspalathus linearis</i>	leaf	0.0684
PA12	<i>Rosmarinus officinalis</i>	leaf	0.0545	PB46	<i>Trigonella foenum-graecum</i>	seed	0.1342
PA13	<i>Filipendula ulmaria</i>	flower	0.095	PB47	<i>Foeniculum vulgare</i>	fruit	0.0818
PA14	<i>Satureja montana</i>	leaf	0.0619	PB48	<i>Alkanna tinctoria</i>	root	0.0606
PA15	<i>Calendula officinalis</i>	petal	0.0383	PB49	<i>Mentha spicata</i>	leaf	0.0694

PA16	<i>Salvia officinalis</i>	leaf	0.0415	PB50	<i>Eupatorium cannabinum</i>	aerial part	0.0529
PA17	<i>Passiflora incarnata</i>	aerial part	0.0423	PB51	<i>Echinacea purpurea</i>	root	0.0915
PA18	<i>Plantago lanceolata</i>	leaf	0.0626	PB55	<i>Eucalyptus globulus</i>	leaf	0.0666
PA19	<i>Tussilago farfara</i>	inflorescence	0.0494	PB57	<i>Rosaxcentifolia</i>	bud	0.0819
PA20	<i>Valeriana officinalis</i>	root	0.1142	PB58	<i>Solanum dulcamara</i>	stem	0.0709
PA21	<i>Camellia sinensis</i>	leaf	0.135	PB60	<i>Arbutus unedo</i>	leaf	0.0891
PA22	<i>Vitis vinifera</i> var. <i>tinctoria</i>	leaf	0.0762	PB61	<i>Borrigo officinalis</i>	aerial part	0.0754
PA23	<i>Thymus vulgaris</i>	leaf	0.0373	PB76	<i>Eleutherococcus senticosus</i>	root	0.0571
PA24	<i>Cassia angustifolia</i>	leaflet	0.0648	PB77	<i>Cupressus sempervirens</i>	nut	0.0862
PA25	<i>Tilia cordata</i>	sapwood	0.0391	PB78	<i>Curcuma longa</i>	root	0.0946
PA26	<i>Thymus serpyllum</i>	aerial part	0.0442	PB79	<i>Cuminum cyminum</i>	fruit	0.1109
PA27	<i>Tilia tomentosa</i>	bract	0.0775	PB81	<i>Salvia officinalis</i>	leaf	0.0536
PA28	<i>Rubus idaeus</i>	leaf	0.0463	PB82	<i>Salix alba</i>	bark	0.0522
PA29	<i>Fucus vesiculosus</i>	lamina	0.1702	PB83	<i>Saponaria officinalis</i>	aerial part	0.0602
PA30	<i>Lavandula angustifolia</i>	flower	0.0216	PB85	<i>Lythrum salicaria</i>	aerial part	0.0581
PA31	<i>Trigonella foenum-graecum</i>	seed	0.1043	PB86	<i>Citrus limon</i> (L) <i>Burn.f.</i>	peel	0.098
PA32	<i>Lotus corniculatus</i>	aerial part	0.0497	PB88	<i>Centella asiatica</i>	aerial part	0.0685
PA33	<i>Handroanthus impetiginosus</i>	bark	0.0814	PB92	<i>Plantago ovata</i>	seed	0.1266
PA34	<i>Viscum album</i>	leaf	0.0783	PB97	<i>Agropyrum repens</i>	rhizome	0.0555
PA35	<i>Fumaria officinalis</i>	flowering top	0.0661	PB98	<i>Cichorium intybus</i>	leaf	0.0715
PA36	<i>Humulus lupulus</i>	fruit	0.0307	PB99	<i>Potentilla anserina</i>	leaf	0.0298
PA37	<i>Fraxinus excelsior</i>	leaf	0.0647	PC01	<i>Quercus robur</i>	bark	0.06
PA38	<i>Cytisus scoparius</i>	flower	0.0253	PC02	<i>Chelidonium majus</i>	aerial part	0.0608
PA39	<i>Foeniculum vulgare</i>	fruit	0.081	PC04	<i>Prunus cerasus</i>	peduncle	0.0199
PA40	<i>Origanum majorana</i>	leaf	0.0345	PC06	<i>Asparagus officinalis</i>	root	0.0812
PA41	<i>Citrus aurantium</i> var. <i>dulcis</i>	leaf	0.0574	PC08	<i>Medicago sativa</i>	seed	0.113
PA44	<i>Hyssopus officinalis</i>	flowering top	0.0677	PC11	<i>Juglans regia</i>	leaf	0.0641
PA45	<i>Mentha</i> × <i>piperita</i>	leaf	0.07	PC12	<i>Olea europaea</i>	leaf	0.0561
PA46	<i>Melissa officinalis</i>	leaf	0.0747	PC14	<i>Myrtus communis</i>	leaf	0.0716
PA47	<i>Fragaria vesca</i>	whole plant	0.0692	PC15	<i>Trifolium rubens</i>	flower	0.0484
PA48	<i>Phaseolus vulgaris</i>	fruit	0.0608	PC17	<i>Curcuma zedoaria</i>	rhizome	0.077
PA51	<i>Marrubium vulgare</i>	aerial part	0.0457	PC19	<i>Vitis vinifera</i> var. <i>tinctoria</i>	leaf	0.0695
PA52	<i>Malva sylvestris</i>	flower	0.0355	PC20	<i>Eugenia caryophyllus</i>	clove	0.0852
PA53	<i>Citrus aurantium</i> var. <i>dulcis</i>	petal	0.0851	PC21	<i>Althaea officinalis</i>	root	0.0642

PA55	<i>Ginkgo biloba</i>	leaf	0.0701	PC22	<i>Alpinia officinarum</i>	rhizome	0.0658
PA57	<i>Lamium album</i>	flower	0.0591	PC23	<i>Ginkgo biloba</i>	leaf	0.0775
PA60	<i>Olea europaea</i>	leaf	0.0572	PC24	<i>Panax ginseng</i>	lateral root	0.1782
PA61	<i>Ilex paraguariensis</i>	leaf	0.0598	PC25	<i>Zingiber officinale</i>	rhizome	0.078
PA62	<i>Artemisia vulgaris</i>	leaf	0.0447	PC26	<i>Juniperus communis</i>	fruit	0.0505
PA67	<i>Epilobium parviflorum</i>	aerial part	0.0549	PC27	<i>Geranium robertianum</i>	aerial part	0.0639
PA69	<i>Chamomilla recutita</i>	inflorescence	0.0399	PC28	<i>Gentiana lutea</i>	root	0.0992
PA70	<i>Sisymbrium officinale</i>	aerial part	0.0334	PC29	<i>Rubia tinctorium</i>	root	0.0868
PA72	<i>Asparagus officinalis</i>	root	0.0523	PC32	<i>Jasminum officinale</i>	flower	0.0616
PA73	<i>Pimpinella anisum</i>	fruit	0.0993	PC33	<i>Combretum micranthum</i>	leaf	0.0769
PA74	<i>Ribes nigrum</i>	fruit	0.0769	PC34	<i>Satureja montana</i>	leaf	0.0715
PA75	<i>Crataegus oxyacantha</i>	flowering top	0.0656	PC35	<i>Cassia angustifolia</i>	leaflet	0.0881
PA76	<i>Chamaemelum nobile</i>	inflorescence	0.0193	PC36	<i>Verbena officinalis</i>	aerial part	0.0603
PA79	<i>Eucalyptus globulus</i>	leaf	0.0703	PC37	<i>Lippia citrodora</i>	leaf	0.0813
PA80	<i>Prunus cerasus</i>	peduncle	0.0141	PC38	<i>Veronica officinalis</i>	complete	0.0781
PA81	<i>Acorus calamus</i>	rhizome	0.0631	PC41	<i>Calendula officinalis</i>	petal	0.0523
PA82	<i>Calluna vulgaris</i>	flower	0.0493	PC42	<i>Sambucus nigra</i>	bark	0.0593
PA83	<i>Papaver rhoeas</i>	petal	0.0278	PC45	<i>Solidago gigantea</i>	aerial part	0.0612
PA85	<i>Cuminum cyminum</i>	fruit	0.102	PC49	<i>Tanacetum vulgare</i>	aerial part	0.0577
PA86	<i>Centaureum erythraea</i>	aerial part	0.0591	PC50	<i>Galium odoratum</i>	aerial part	0.041
PA88	<i>Vaccinium myrtillus</i>	fruit	0.0939	PC51	<i>Rhamnus frangula</i>	bark	0.0726
PA89	<i>Vaccinium myrtillus</i>	leaf	0.0711	PC52	<i>Betula alba</i>	bark	0.0806
PA90	<i>Angelica archangelica</i>	fruit	0.0388	PC53	<i>Verbascum thapsus</i>	bark	0.0435
PA92	<i>Borago officinalis</i>	flower	0.067	PC54	<i>Peumus boldus</i>	leaf	0.0695
PA93	<i>Alchemilla vulgaris</i>	aerial part	0.0588	PC55	<i>Centaurea cyanus</i>	petal	0.0456
PA94	<i>Artemisia pontica</i>	flowering top	0.431	PC57	<i>Illicium verum</i>	fruit	0.1027
PA95	<i>Illicium verum</i>	fruit	0.0793	PC58	<i>Inula helenium</i>	root	0.0904
PA96	<i>Chrysanthellum americanum</i>	aerial part	0.0768	PC60	<i>Cinnamomum zeylanicum</i>	bark	0.1374
PA97	<i>Harpagophytum procumbens</i>	root	0.0853	PC61	<i>Chamaemelum nobile</i>	flower	0.0316
PA98	<i>Betula pendula</i>	leaf	0.0633	PC62	<i>Calamintha officinalis</i>	aerial part	0.0726
PA99	<i>Plantago lanceolata</i>	aerial part	0.0718	PC63	<i>Arctostaphylosuva-ursi</i>	leaf	0.088
PB00	<i>Agropyrum repens</i>	rhizome	0.0646	PC64	<i>Ononis spinosa</i>	root	0.0832
PB01	<i>Carum carvi</i>	fruit	0.1089	PC65	<i>Calluna vulgaris</i>	flower	0.0446
PB02	<i>Buxus sempervirens</i>	leaf	0.0631	PC66	<i>Morus nigra</i>	leaf	0.1032

PB03	<i>Cichorium endivia</i>	root	0.1513	PC67	<i>Hypericum perforatum</i>	flowering top	0.0829
PB04	<i>Rosa canina</i>	fruit	0.1227	PC68	<i>Achillea millefolium</i>	flowering top	0.0332
PB05	<i>Silybum marianum</i>	aerial part	0.0502	PC70	<i>Mentha × piperita</i>	leaf	0.071
PB06	<i>Asperula odorata</i>	aerial part	0.062	PC71	<i>Melissa officinalis</i>	leaf	0.0847
PB07	<i>Desmodium adscendens</i>	leaf	0.0493	PC72	<i>Cochlearia officinalis</i>	flowers	0.0497
PB08	<i>Combretum micranthum</i>	leaf	0.0709	PC73	<i>Malva sylvestris</i>	flower	0.0615
PB09	<i>Peumus boldus</i>	foliole	0.0574	PC74	<i>Cynara scolymus</i>	leaf	0.0458
PB10	<i>Rhamnus frangula</i>	bark	0.0682	PC75	<i>Arnica montana</i>	flower	0.0254
PB11	<i>Arctium lappa</i>	root	0.1166	PC76	<i>Artemisia vulgaris</i>	aerial part	0.0289
PB12	<i>Arctostaphylos uva-ursi</i>	leaf	0.122	PC78	<i>Angelica archangelica</i>	complete	0.0909
PB13	<i>Cichorium intybus</i>	root	0.0924	PC79	<i>Anethum graveolens</i>	fruit	0.1335
PB14	<i>Fragaria vesca</i>	rhizome	0.0529	PC82	<i>Ballota nigra</i>	aerial part	0.0368
PB15	<i>Citrus aurantium var. dulcis.</i>	peel	0.0817	PC83	<i>Origanum majorana</i>	leaf	0.0539
PB16	<i>Elettaria cardamomum</i>	fruit	0.0606	PC84	<i>Zea mays</i>	stigma	0.0632
PB17	<i>Althaea officinalis</i>	leaf	0.0302	PC86	<i>Agrimonia eupatoria</i>	aerial part	0.0721
PB18	<i>Malva sylvestris</i>	leaf	0.0809	PC88	<i>Levisticum officinale</i>	aerial part	0.0848
PB19	<i>Althaea officinalis</i>	leaf	0.0322	PC94	<i>Phaseolus vulgaris</i>	fruit	0.0686
PB20	<i>Tilia cordata</i>	flowering top	0.0596	PC97	<i>Hieracium pilosella</i>	aerial part	0.0406
PB22	<i>Betula pendula</i>	leaf	0.0609	PC98	<i>Parietaria officinalis</i>	aerial part	0.0349
PB24	<i>Tilia cordata</i>	sapwood	0.0368	PC99	<i>Passiflora incarnata</i>	aerial part	0.0817
PB25	<i>Fucus vesiculosus</i>	lamina	0.1119	PD00	<i>Rumex patientia</i>	root	0.1208
PB26	<i>Borago officinalis</i>	flower	0.0342	PD02	<i>Petroselinum crispum</i>	fruit	0.1084
PB27	<i>Vaccinium myrtillus</i>	fruit	0.1553	PD05	<i>Urtica dioica</i>	aerial part	0.0584
PB28	<i>Verbascum thapsus</i>	flower	0.045	PD06	<i>Cola acuminata</i>	nut	0.1556
PB29	<i>Citrus aurantium</i>	petal	0.0477	PD07	<i>Capsella bursa-pastoris</i>	aerial part	0.0535
PB31	<i>Galega officinalis</i>	aerial part	0.0678	PD08	<i>Centaurea erythraea</i>	aerial part	0.058

Method for preparing the herbal compositions (Method A)

Each dried plant was grinded using a coffee bean grinder (Type 8100, SEB, France). The obtained powder (plant powder) was sieved using a 2 mm mesh sieve, and conserved in 60 ml containers (Labbox, France).

Mixtures of different grinded dried plants (or plant powders) were prepared by collecting a fixed volume of plant powder with a calibrated spoon allowing to collect 200 µL of powder. The weight of the collected volume is dependent on the specific plant powder (see Table 0-1).

Such mixes or mixtures of different grinded dried plants (or plant powders) as well as single dried plants will herein be called also as "herbal compositions".

Method for preparing the water extracts (also called in this Examples section as “processed samples”, “extracts” or “mixes extracts”) (Method B)

Such herbal compositions were placed into 50 ml falcon tubes (Falcon, USA) and 20 ml or 40 ml (as indicated in Examples) of alimentary grade spring water (Volvic, France) solution containing 100g/l of alimentary grade sucrose (Daddy, France) were added. The choice of alimentary grade sucrose and water rather than their scientific grade counterparts was dictated by the objective of obtaining an edible composition entering the class of “flavourings” that may be directly consumed under the Regulation (EC) No 1334/2008 of 16 December 2008 on flavourings and certain food ingredients with flavouring properties for use in and on foods.

Each falcon tube containing such liquid preparations was agitated using a vortex mixer (ThermoFisher, USA) for 10 seconds, incubated at room temperature for 10 min and placed in an autoclave (MED 12, JP Selecta, Spain) and underwent a 20 minutes sterilization cycle at 119° or 121°C, as indicated. After cooling down, the sterilized preparation was kept at -20°C until use.

The day of manipulation, the preparation was defrosted at 37°C. The preparation was centrifuged at 4000 rpm during 10 min (Multifuge 35-R, Heraeus, Germany) or incubated at room temperature for decantation (as indicated in Examples). 2 mL of the supernatant was collected with a 10 mL syringe and filtered using 0.2µm filters (Minisart Syringe Filters, Sartorius, Germany) and collected onto a 2mL Eppendorf tube. The remaining sterilized preparation is kept at -20°C for further use.

The samples so obtained will herein be called also as “processed samples” and will be identified by the same name or code given to the corresponding herbal composition (for example the name “**Mix[2P]-1**” will be used to identify an herbal composition and the corresponding extract (or processed sample), for which a biological activity is determined).

Method for determining the antimicrobial and antibiofilm activity (Method C)

The biological activity of each processed sample was determined by using the strains indicated in each experiment (or Example). For each measurement, each bacteria stored at -80°C on cryobeads (Technical Service, UK) is plated on a TS agar plate (Tryptic Soy Broth, Sigma, USA; Agar, Fisherbrand, USA) and incubated at 37°C overnight. Several colonies are collected using a inoculation loop and suspended in saline solution. The optical density is measured and a dilution is performed in TS broth to obtain a concentration of $2.10^{5.5}$ bacteria/mL for *Staphylococcus aureus* and *Staphylococcus pseudintermedius* strains, 2.10^5 bacteria/mL for *Staphylococcus epidermidis*.

The biological activity of each processed sample was measured on 96 wells plates, using the strains indicated in each experiment (or Example) and a control without bacteria. Each well contains 190µL of TS broth inoculated with $10^{5.5}$ bacteria/mL for *Staphylococcus aureus* and *Staphylococcus pseudintermedius* strains, 10^5 bacteria/mL for *Staphylococcus epidermidis* (or at the concentration indicated in the Examples for all other species) or no bacteria for the control and diluted processed sample or negative control (sucrose solution at 100g/l without any plants) at 1:10, 1:20, 1:40 and 1:80 final concentration or at 1:20, 1:63 and 1:200 (as indicated in Examples).

Each plate was incubated for 17 hours at 37°C in an incubator (HettCube 400, Hettich, Germany).

The measurements were performed based on analyzing with the ImageJ software (NIH, USA) (but any other equivalent software may be used) images of the plates obtained using a commercial document scanner (V 220, Epson, Japan). Images of the plate positioned on the scanning area are acquired at 600dpi using the software provided with the scanner. For each plate, 3 images were acquired: the first, before incubation (T0), the second after 17h incubation (Tf), and the third after Crystal Violet (CV)

staining as described in Optimization of microtitre plate assay for the testing of biofilm formation ability in different *Salmonella* serotypes (Agarwal, R.K. et al., International Food Research Journal 18(4):1493-1498, 2011).

For each well of interest, from respectively the first, second and the third image, circular regions of diameter, respectively, 14, 14 and 76 pixels, centered relative to the well, were characterized using an ImageJ macro to calculate, respectively, the mean M1 and M2 of the 1% less intense pixels of the region and the mean M3 of all the pixels of the region. The turbidity of the bacterial solution in the well was deduced from M1 and M2 using a reference curve obtained in an independent experiment where the mean of the 1% less intense pixels of the region 14 pixels diameter region was measured from wells containing bacterial solutions of known turbidity. The optical density of the Crystal Violet solution in the well was deduced from M3 using a reference curve obtained in an independent experiment where the mean of the pixels of the region 76 pixels diameter region was measured from wells containing Crystal Violet solutions of known optical density.

For each processed sample and a given bacterium, the MIC was determined from the highest inhibitory dilution (HID) among the 4 tested (1:10, 1:20, 1:40, 1:80) of processed sample which prevents the appearance of visible growth within 17h. "0" was assigned when none of the tested dilutions was preventing bacterial growth.

For each well containing processed sample and a given bacterium, the inhibition percentage of bacteria planktonic growth was established according to the formula below:

$$\text{Planktonic growth inhibition (\%)} = \left(1 - \frac{Tf(\text{processed sample}) - T0(\text{processed sample})}{Tf(\text{negative control}) - T0(\text{negative control})}\right) * 100$$

where Tf(processed sample) is the turbidity from the second image of the sample well, T0(processed sample) is the turbidity of the first image of the same well, Tf(negative control) is the turbidity from the second image of a well containing the same bacteria without processed sample, T0(negative control) is the turbidity of the first image of the same well.

For each well containing processed sample and a given bacterium, the percentage of inhibition of biofilm formation (IBF) was established according to the following formula:

$$\text{Biofilm formation inhibition (\%)} = \left(1 - \frac{CV(\text{processed sample}) - CV(\text{control sample})}{CV(\text{negative control}) - CV(\text{control sample})}\right) * 100$$

where CV(processed sample) is the OD from the third image of the sample well, CV(negative control) is the OD from the third image of a well containing the same bacteria without processed sample, CV(control sample) is the OD of the third image of a well containing the same sample or the negative control at the same dilution without any bacteria. A negative value of IBF means that the tested mixture is promoting the formation of biofilm.

Example 1

In this Example we illustrate simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus* of the disclosed composition of 2 plants among 3: *Filipendula ulmaria*, *Camellia sinensis*, *Arctostaphylos uva-ursi*. The dried plants of pharmaceutical grade were obtained from "Herboristerie Cailleau" (Chemillé, France).

Starting from such three different plant powders six herbal compositions were prepared according to the Method A reported before, whose content is reported in Table 1-1.

Table 1-1: herbal compositions of processed samples

Herbal compositions	PA13 (nb spoon)	PA21 (nb spoon)	PB12 (nb spoon)
PA13	1		
PA21		1	
PB12			1
Mix[2P]-1	1	1	
Mix[2P]-2		1	1
Mix[2P]-3	1		1

As shown in Table 1-1 the first three herbal compositions only contain a single plant powder, while the last three compositions contain two of the three different plant powders.

Starting from each herbal composition reported in Table 1-1 corresponding processed samples were prepared according to the Method B reported before using 20mL of water containing 100g/l of sucrose for water extraction and including the centrifugation step before filtration.

The bacteria planktonic growth and antibiofilm formation inhibition percentage of the processed samples of the 2 plant mixes and the associated separated plants of this Example 1 at 1:10 dilution were determined by using the following *Staphylococcus aureus* strains: ATCC 25923 and ATCC 29213, according to the Method C reported before and are given in Table 1-2.

Table 1-2: planktonic growth and biofilm formation inhibition (IBF) of 2 plants mixture processed sample and single plants processed sample at 1:10 dilution

		PA13	PA21	Mix[2P]-1
ATCC 25923	Planktonic growth inhibition	20%	26%	82%
	Biofilm formation inhibition	32%	44%	78%
ATCC 29213	Planktonic growth inhibition	16%	27%	100%
	Biofilm formation inhibition	33%	17%	85%

		PA21	PB12	Mix[2P]-2
ATCC 25923	Planktonic growth inhibition	26%	17%	104%
	Biofilm formation inhibition	44%	51%	88%
ATCC 29213	Planktonic growth inhibition	27%	8%	100%
	Biofilm formation inhibition	17%	28%	83%

		PA13	PB12	Mix[2P]-3
ATCC 25923	Planktonic growth inhibition	20%	17%	82%
	Biofilm formation inhibition	32%	51%	93%
ATCC 29213	Planktonic growth inhibition	16%	8%	66%
	Biofilm formation inhibition	33%	28%	78%

PA13, PA21 and PB12 have a known antibiotic effect. Limited inhibition of planktonic growth (< 27%) and biofilm formation (< 51%) were measured at 1:10 dilution, whereas, the 3 aromatic preparations of 2 plant mix extract show antimicrobial effects (> 66%) and antibiofilm effects (>78%) on both tested strains, ATCC 25923 and ATCC 29213.

For all Mixes (Mix[2P]-1, Mix[2P]-2, Mix[2P]-3), a synergistic effect of the 2 plants is observed for antimicrobial effects on the 2 strains tested.

Comparative Example 1

As a comparative Example, we illustrate the activity of some processed samples, which have been found to have none or poor antimicrobial and antibiofilm activities against *Staphylococcus aureus* and which were obtained from corresponding herbal compositions containing 2 different plant powders.

- 5 Five processed samples were prepared using 2 additional plants powders: PA74 (*Ribes nigrum* fruit powder) and PC72 (*Cochlearia officinalis* flowers powder).

The single dried plant powders were treated as described in Example 1 and five herbal compositions were prepared following the protocol reported in Method A described before. Such five herbal compositions had the content reported in Table 1-3.

Table 1-3: herbal compositions of processed samples

Herbal Compositions	PA74 (nb spoon)	PC72 (nb spoon)	PA13 (nb spoon)	PA21 (nb spoon)
PA74	1			
PC72		1		
Mix 1	1	1		
Mix 2	1		1	
Mix 3		1		1

- 10 Each herbal composition was treated following the protocol reported in Method B to arrive at the corresponding processed sample using 20mL of water containing 100g/l of sucrose for water extraction and including the centrifugation step before filtration.

The biological activity of each processed sample was determined by using the following *Staphylococcus aureus* strains: ATCC 25923 and ATCC 29213, according to Method C described before.

- 15 The bacteria planktonic growth and antibiofilm formation inhibition percentage of the processed samples of the 2 plant mixes and the associated separated plants at 1:10 dilution are given in Table 1-4.

Table 1-4: planktonic growth and biofilm formation inhibition (IBF) of 2 plants mixture processed sample and single plants processed sample at 1:10 dilution

		PC72	PA74	Mix 1
ATCC 25923	Planktonic growth inhibition	-9%	0%	-3%
	Biofilm formation inhibition	12%	36%	24%
ATCC 29213	Planktonic growth inhibition	-10%	-7%	-6%
	Biofilm formation inhibition	13%	5%	-2%
		PA13	PA74	Mix 2
ATCC 25923	Planktonic growth inhibition	20%	0%	19%
	Biofilm formation inhibition	32%	36%	48%
ATCC 29213	Planktonic growth inhibition	16%	-7%	13%
	Biofilm formation inhibition	33%	5%	37%
		PA21	PC72	Mix 3
ATCC 25923	Planktonic growth inhibition	26%	-9%	25%
	Biofilm formation inhibition	44%	12%	51%
ATCC 29213	Planktonic growth inhibition	27%	-10%	18%
	Biofilm formation inhibition	17%	13%	29%

PC72 and PA74 and the associated Mix 1 showed no antimicrobial activity and limited antibiofilm activity on both tested strains at 1:10 dilution. In each case, no synergistic effect was observed.

For all mixes containing PA13 and PA21 (Mix 2 and 3), no synergistic effect was observed for antimicrobial effects on the 2 strains tested at 1:10 dilution, when such mixes contained also PC72 or PA74.

It is interesting to note that both *Ribes nigrum* and *Cochlearia officinalis* were known to have antimicrobial effect.

This comparative Example illustrates that not any random combination between single dried plants having known antimicrobial and/or antibiofilm activity allows to obtain a water extract, in which a synergistic activity is observed.

Example 2

In this Example we illustrate simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus* of different water extracts obtained from the herbal compositions containing 2 plants among the 3 reported in Example 1, when such water extracts are prepared according to different methods.

The dried plants of pharmaceutical grade were obtained from “Herboristerie Cailleau” (Chemillé, France).

Twenty herbal compositions, whose content is reported in Table 2-1 were prepared according to the Method A reported before.

Table 2-1: herbal compositions of processed samples

Herbal compositions	PA13 (nb spoon)	PA21 (nb spoon)	PB12 (nb spoon)
Mix 1		1	1
Mix 2	1		1

For each herbal composition, 20 liquid (water) preparations were prepared as follow:

The collected powders were placed into twenty 50 ml falcon tubes (Falcon, USA). 20 ml of alimentary grade spring water (Volvic, France) solution containing 0, 25, 50, 75 or 100g/l of alimentary grade sucrose (Daddy, France) were added to 4 falcon tubes per condition. Each falcon tube containing such liquid preparations was agitated using a vortex mixer (ThermoFisher, USA) for 10 seconds, incubated at room temperature for 10 minutes, then 4 different heating treatments were applied to each condition:

- Cold infusion (no heating before centrifugation and filtration):

after incubation at room temperature, the preparations were centrifuged at 4,000 rpm during 10 minutes (Multifuge 35-R, Heraeus, Germany), 2 mL of the supernatant was collected with a 10 mL syringe and filtered using 0.2 µm filters (Minisart Syringe Filters, Sartorius, Germany) and collected onto a 2mL Eppendorf tube and kept at -20°C until use. The day of manipulation, the preparation was defrosted at 37°C.

- Heating at 60°C for 30 minutes:

after incubation at room temperature, the preparations were placed in an autoclave (VWR, USA) and underwent a 20 minutes cycle at 60°C. After cooling down, the treated preparations were centrifuged at 4000 rpm during 10 min (Multifuge 35-R, Heraeus, Germany), 2 mL of the

supernatant was collected with a 10 mL syringe and filtered using 0.2µM filters (Minisart Syringe Filters, Sartorius, Germany) and collected onto a 2mL Eppendorf tube and kept at -20°C until use. The day of manipulation, the preparation was defrosted at 37°C.

- 5 - Heating at 119°C for 20 minutes:
 after incubation at room temperature, the preparations were placed in an autoclave (VWR, USA) and underwent a 20 minutes sterilization cycle at 119°C. After cooling down, the treated preparations were kept at -20°C until use. The day of manipulation, the preparation was defrosted at 37°C, centrifuged at 4000 rpm during 10 min (Multifuge 35-R, Heraeus, Germany),
 10 1 mL of the supernatant was collected with a 10 mL syringe and filtered using 0.2µM filters (Minisart Syringe Filters, Sartorius, Germany) and collected onto a 2mL Eppendorf tube.

These samples will herein called “processed sample”.

- 15 The biological activity of each processed sample was determined by using *Staphylococcus aureus* strain ATCC 25923, according to the previously described Method C.

The bacteria planktonic growth percentage of 20 processed samples, prepared starting from the herbal compositions after each of the different heating treatment presented before, at 1:10 dilution are given in Table 2-2.

20 **Table 2-2: planktonic growth of ATCC 25923 with 20 processed samples at 1:10 dilution prepared starting from the herbal compositions (Mix 1 and Mix 2 of Table 2-1)**

Temperature treatment	Sucrose concentration (g/l)	Mix 1 Processed samples	Mix 2 Processed samples
Cold infusion	100	-4%	48%
	75	-3%	52%
	50	-4%	31%
	25	-4%	31%
	0	1%	0%
60°C 30min	100	0%	82%
	75	4%	83%
	50	1%	83%
	25	1%	86%
	0	-4%	16%
119°C 20min	100	103%	92%
	75	103%	91%
	50	70%	94%
	25	103%	99%
	0	96%	99%

All processed samples prepared from Mix 1 showed antimicrobial effects on ATCC 25923, only after temperature treatment at 119°, whereas Mix 2 showed limited antimicrobial effects (31 to 52% of inhibition) after cold infusion for all concentration of sucrose except for the condition without sucrose

and after all tested temperature treatments (> 82% of inhibition) except for the condition with no sucrose addition treated at 60°C, for which the processed sample showed only 16% of planktonic growth inhibition. The temperature treatments enhanced the antimicrobial effects of Mix 2. The optimal temperature treatment among tested treatments for Mix 1 and Mix 2 was 119°C for 20 minutes.

Sucrose concentration influence on antimicrobial effect depends on the temperature treatment applied. For Mix 1 treated at 119°C during 20min, the same antimicrobial effect was obtained with 100, 75, 25 and 0 g/l of sucrose, lower activity was observed with 50 g/l. For Mix 2, the same antimicrobial effect was obtained regardless of sucrose concentration for 119°C temperature treatment.

The biofilm formation inhibition (IBF) percentage of 20 processed samples, prepared starting from the herbal compositions after each of the different heating treatment presented before, at 1:10 dilution are given in Table 2-3.

Table 2-3: biofilm formation inhibition (IBF) of ATCC 25923 with 20 processed samples at 1:10 dilution prepared starting from the herbal compositions (Mix 1 and Mix 2 of Table 2-1)

Temperature treatment	Sucrose concentration (g/l)	Mix 1 Processed samples	Mix 2 Processed samples	Temperature treatment	Sucrose concentration (g/l)	Mix 1 Processed samples	Mix 2 Processed samples
Cold infusion	100	34%	74%	119°C 20min	100	91%	89%
	75	29%	72%		75	91%	88%
	50	22%	60%		50	75%	89%
	25	30%	55%		25	92%	87%
	0	54%	50%		0	90%	89%
60°C 30min	100	48%	91%				
	75	34%	94%				
	50	26%	91%				
	25	26%	88%				
	0	48%	78%				

In all tested conditions, processed samples showed some antibiofilm effect.

For Mix 1, limited antibiofilm activity was observed for all concentration of sucrose with cold infusion and with 60°C temperature treatment (22 to 55% of inhibition). High antibiofilm activity (> 90% of inhibition) was observed with 119°C temperature treatment for all concentrations of sucrose except for 50 g/l which showed 75% of inhibition.

For Mix 2, equivalent antibiofilm activities were obtained regardless of sucrose concentrations for 60°C and 119°C temperature treatments (> 77% of inhibition). After cold infusion, Mix 2 showed medium to good antibiofilm effect (from 50% to 74% of inhibition) depending on the sucrose concentrations: the more the sucrose concentration was high, the more the preparation was active.

Example 3

In this Example we illustrate simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus* and *Staphylococcus epidermidis* of 12 mixtures containing 2 to 7 plants.

The dried plants of pharmaceutical grade were obtained from “Herboristerie Cailleau” (Chemillé, France).

Herbal compositions comprising from 2 to 7 plant powders were prepared by collecting a fixed volume of plant powder with a calibrated spoon allowing to collect 200 µL of powder

- 5 Twelve herbal composition were prepared as below (Table 3-1) according to the previously described Method A.

Table 3-1: herbal compositions of processed samples

Herbal compositions	PA13 (nb spoon)	PA21 (nb spoon)	PB12 (nb spoon)	PA11 (nb spoon)	PA12 (nb spoon)	PA22 (nb spoon)	PA79 (nb spoon)
Mix[2P]-1	1	1					
Mix[2P]-2		1	1				
Mix[2P]-3	1		1				
Mix[3P]-1	1	1	1				
Mix[4P]-1	1	1	1	1			
Mix[4P]-2	1	1	1		1		
Mix[4P]-3	1	1	1			1	
Mix[5P]-1	1	1	1		1	1	
Mix[5P]-2	1	1	1	1		1	
Mix[5P]-3	1	1	1	1	1		
Mix[6P]-1	1	1	1	1	1	1	
Mix[7P]-1	1	1	1	1	1	1	1

Twelve corresponding processed samples were prepared starting from each herbal composition according to the previously described Method B using 20mL of water containing 100g/l of sucrose for water extraction and including the centrifugation step before filtration.

10

The biological activity of the processed sample was determined by using the following *Staphylococcus aureus* strains: ATCC 25923, ATCC 29213, NCTC 12493, ATCC 33591, ATCC 33592, ATCC 43300, ATCC 700698, ATCC 700699, ATCC 9144 and ATCC BAA-44 and the following *Staphylococcus epidermidis* strains: ATCC 12228, ATCC 700296, ATCC 49461 and ATCC 14990, according to the previously described Method C.

15

The minimum inhibitory concentration (MIC) observed with the 12 processed samples is given in Table 3-2 in the form of HID, as well as the average efficacy over all the strains of *Staphylococcus aureus* and *Staphylococcus epidermidis*.

20 **Table 3-2: HID of the 12 processed samples on 10 strains of *Staphylococcus aureus* and 4 strains of *Staphylococcus epidermidis***

		Mix[2P]-1	Mix[2P]-2	Mix[2P]-3	Mix[3P]-1	Mix[4P]-1	Mix[4P]-2	Mix[4P]-3	Mix[5P]-1	Mix[5P]-2	Mix[5P]-3	Mix[6P]-1	Mix[7P]-1
<i>S. aureus</i>	ATCC 25923	10	10	10	10	10	10	10	10	10	20	10	20
	ATCC 29213	10	10	0	10	10	10	10	10	10	10	10	10
	NCTC 12493	10	10	10	10	10	20	20	20	20	20	20	20
	ATCC 33591	10	10	10	10	10	10	10	20	20	20	20	20

	ATCC 33592	10	10	10	10	10	10	10	20	20	20	10	20
	ATCC 43300	10	10	0	10	10	10	10	10	10	10	10	20
	ATCC 700698	10	10	10	10	10	10	10	20	20	20	10	20
	ATCC 700699	10	10	10	20	20	20	20	20	20	20	20	20
	ATCC 9144	10	10	10	10	10	10	10	20	20	20	10	20
	ATCC BAA-44	10	10	10	10	10	20	20	20	20	20	20	20
	Average	10.0	10.0	8.0	11.0	11.0	13.0	13.0	17.0	17.0	18.0	14.0	19.0
<i>S. epidermidis</i>	ATCC 12228	10	10	10	20	20	20	20	20	20	20	20	40
	ATCC 700296	20	20	20	20	20	20	20	20	20	20	20	40
	ATCC 49461	10	10	20	20	20	20	20	20	20	20	20	20
	ATCC 14990	10	20	10	20	20	20	20	20	20	20	20	20
	Average	12.5	15	15	20	20	20	20	20	20	20	20	30
	Total average	10.7	11.4	10.0	13.6	13.6	15.0	15.0	17.9	17.9	18.6	15.7	22.1

Since in this Example observed HID are in most case between 1:10 and 1:20, 1:20 dilution was chosen as the dilution of reference for the calculation of bacteria planktonic growth inhibition percentage and biofilm formation inhibition percentage. The bacteria planktonic growth inhibition percentage of the 12 processed samples at 1:20 dilution is given in Table 3-3, as well as the average efficacy over all the strains of *Staphylococcus aureus* and *Staphylococcus epidermidis*.

5

Table 3-3: planktonic growth inhibition of the 12 processed samples on 10 strains of *Staphylococcus aureus* and 4 strains of *Staphylococcus epidermidis* at 1:20 dilution

		Mix[2P]-1	Mix[2P]-2	Mix[2P]-3	Mix[3P]-1	Mix[4P]-1	Mix[4P]-2	Mix[4P]-3	Mix[5P]-1	Mix[5P]-2	Mix[5P]-3	Mix[6P]-1	Mix[7P]-1
<i>S. aureus</i>	ATCC 25923	12%	12%	9%	11%	20%	11%	20%	34%	41%	35%	21%	96%
	ATCC 29213	11%	11%	7%	11%	19%	11%	11%	33%	33%	33%	29%	45%
	NCTC 12493	19%	8%	5%	25%	21%	101%	97%	98%	108%	108%	96%	100%
	ATCC 33591	12%	8%	8%	12%	19%	16%	34%	104%	107%	111%	83%	98%
	ATCC 33592	19%	15%	12%	18%	26%	23%	26%	86%	105%	98%	38%	100%
	ATCC 43300	18%	11%	11%	15%	18%	18%	18%	33%	34%	37%	16%	97%
	ATCC 700698	11%	8%	8%	11%	11%	11%	16%	107%	97%	73%	40%	98%
	ATCC 700699	43%	36%	38%	87%	98%	99%	99%	100%	108%	112%	116%	98%
	ATCC 9144	12%	7%	8%	12%	24%	17%	24%	80%	97%	73%	43%	94%
	ATCC BAA-44	30%	11%	12%	32%	28%	83%	102%	100%	107%	110%	114%	103%
	Average	19%	13%	12%	23%	28%	39%	45%	78%	84%	79%	60%	93%
<i>S. epidermidis</i>	ATCC 12228	49%	49%	66%	94%	99%	99%	98%	99%	99%	104%	114%	97%
	ATCC 700296	87%	83%	78%	94%	99%	99%	95%	101%	98%	102%	105%	92%
	ATCC 49461	68%	68%	70%	93%	101%	101%	97%	102%	102%	106%	112%	93%
	ATCC 14990	38%	71%	61%	92%	103%	100%	95%	102%	102%	106%	107%	93%
	Average	61%	68%	69%	94%	100%	100%	96%	101%	100%	104%	109%	94%
	Total average	31%	28%	28%	43%	49%	56%	59%	84%	88%	86%	74%	93%

The biofilm formation inhibition (IBF) percentage of the 12 processed samples at 1:20 dilution is given in Table 3-4, as well as the average efficacy over all the strains of *Staphylococcus aureus* and *Staphylococcus epidermidis*. The average efficacy of bacteria planktonic growth inhibition and biofilm formation inhibition percentage over all the tested strains is a good indicator of antimicrobial and antibiofilm efficacy on different bacterial strains.

Table 3-4: biofilm formation inhibition (IBF) of the 12 processed samples on 10 strains of *Staphylococcus aureus* and 4 strains of *Staphylococcus epidermidis* at 1:20 dilution

		Mix[2P]-1	Mix[2P]-2	Mix[2P]-3	Mix[3P]-1	Mix[4P]-1	Mix[4P]-2	Mix[4P]-3	Mix[5P]-1	Mix[5P]-2	Mix[5P]-3	Mix[6P]-1	Mix[7P]-1
<i>S. aureus</i>	ATCC 25923	37%	14%	22%	18%	29%	36%	35%	51%	41%	40%	38%	90%
	ATCC 29213	49%	12%	38%	27%	33%	38%	25%	35%	38%	34%	38%	41%
	NCTC 12493	49%	-6%	47%	36%	35%	88%	87%	92%	92%	89%	47%	93%
	ATCC 33591	38%	-21%	30%	16%	20%	25%	19%	92%	91%	89%	57%	93%
	ATCC 33592	51%	-10%	26%	25%	43%	47%	31%	60%	78%	70%	43%	91%
	ATCC 43300	33%	-19%	24%	2%	18%	26%	2%	-15%	-22%	-18%	-19%	68%
	ATCC 700698	27%	-8%	20%	10%	25%	23%	13%	62%	61%	39%	24%	88%
	ATCC 700699	61%	34%	63%	64%	79%	78%	86%	89%	87%	86%	87%	90%
	ATCC 9144	33%	-10%	22%	16%	21%	10%	16%	37%	59%	28%	0%	81%
	ATCC BAA-44	50%	22%	47%	40%	39%	48%	89%	92%	90%	87%	91%	93%
	Average	43%	1%	34%	25%	34%	42%	40%	59%	62%	54%	41%	83%
<i>S. epidermidis</i>	ATCC 12228	62%	-32%	78%	84%	95%	92%	92%	92%	91%	93%	96%	96%
	ATCC 700296	61%	39%	78%	73%	85%	86%	84%	80%	75%	76%	79%	87%
	ATCC 49461	18%	15%	72%	84%	90%	90%	90%	88%	88%	86%	90%	94%
	ATCC 14990	68%	39%	68%	88%	94%	91%	92%	90%	90%	89%	91%	90%
	Average	52%	15%	74%	82%	91%	90%	89%	87%	86%	86%	89%	92%
Total average		45%	5%	45%	42%	50%	56%	54%	67%	69%	63%	54%	85%

2-plants mixes extracts showed full antimicrobial activity at 1:10 dilution on the 10 *Staphylococcus aureus* strains tested except for the Mix[2P]-3 which had no antimicrobial at the tested dilution on 2 of the 10 *Staphylococcus aureus* strains tested (ATCC 29213 and ATCC 43300) (see Table 3-2).

The three 2-plant mixes extracts showed different antimicrobial activity on the 4 *Staphylococcus epidermidis* strains tested. Mix[2P]-1 had an HID of 1:10 on 3 strains (ATCC 12228, ATCC 49461 and ATCC 14990) and an HID of 1:20 on ATCC 700296 strain. Mix[2P]-2 had an HID of 1:10 on 2 strains (ATCC 12228 and ATCC 49461) and an HID of 1:20 on the 2 other strains (ATCC 700296 and ATCC 14990). Whereas Mix[2P]-3 had an HID of 1:10 on 2 strains (ATCC 12228 and ATCC 14990) and an HID of 1:20 on the 2 other strains (ATCC 700296 and ATCC 49461) (see Table 3-2).

Mix[3P]-1 obtained from the 3 plants which are included in Mix[2P]-1, Mix[2P]-2 and Mix[2P]-3, showed a wider antimicrobial activity than the 3 two-plant mixes (HID of 1:10 on 9 *Staphylococcus aureus* strains and HID of 1:20 on ATCC 700699, HID of 1:20 on the 4 *Staphylococcus epidermidis* tested) (see Table 3-2). The gain of antimicrobial activity is confirmed by the calculation of the planktonic growth inhibition percentage of each plant mixes (see Table 3-3).

3 different plants, PA11, PA12 and PA22, were added to the 3-plant mix Mix[3P]-1 to obtain three 4 plant-mixes extracts, Mix[4P]-1, Mix[4P]-2 and Mix[4P]-3 respectively. These 4-plant mixes extracts

showed same HID or better HID than the 3-plant mix extract Mix[3P]-1 (see Table 3-2), and higher planktonic growth inhibition percentage: an average of 28% to 45% is obtained with 4-plant mixes extracts on the *10 Staphylococcus aureus* strains compare to 23% with the 3-plant mix extract (see Table 3-3). They also showed higher biofilm formation inhibition percentage: an average of 34% to 42% is obtained with 4-plant mixes extracts on the *10 Staphylococcus aureus* strains compare to 25% with the 3-plant mix extract (see Table 3-4).

5-plant mixes were prepared by adding 2 plants among the 3 selected additional plants. All the three 5-plant mixes extracts showed higher antimicrobial and antibiofilm activity than the 4-plant mixes extracts (see Tables 3-2, 3-3 and 3-4).

Mix[6P]-1 consists in the 6 plants which were used for the preparation of the previously tested processed samples. Interestingly, the 6-plant mix extract showed less antimicrobial and antibiofilm activity than 5-plant mixes extracts but higher antimicrobial effect and higher to equivalent antibiofilm effects than 4-plant mixes extracts. Antimicrobial and antibiofilm activity was recovered and enhanced by the addition of a new plant (PA79) (see Tables 3-2, 3-3 and 3-4).

Increasing the number of plants in the mixes allowed to prepare corresponding water extracts having enhanced antimicrobial and antibiofilm activity and increased the number of strains, in which one or both these activities can be identified.

Example 4

In this Example we illustrate simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus* of 196 water extract obtained from herbal compositions including at least 2 plants among the 3 plants described in Example 1 and additional plants to form mixtures of 3 to 10 plants.

The dried plants of pharmaceutical grade were obtained from "Pharmacie Fontgèze" (Clermont-Ferrand, France, plants PA00 to PB12) and "Pharmacie St Herem" (Clermont-Ferrand, France, plants PB13 to PD08).

The 4 reference herbal compositions Mix[2P]-1, Mix[2P]-2, Mix[2P]-3, Mix[3P]-1 and the 196 new herbal compositions were prepared according to the previously described Method A and had the content reported in List 4-1.

List 4-1: herbal compositions of processed samples

Mix[2P]-1: PA13; PA21;
 Mix[2P]-2: PA21; PB12;
 Mix[2P]-3: PA13; PB12;
 Mix[3P]-1: PA13; PA21; PB12;
 Mix[3P]-2: PA11; PA13; PA21;
 Mix[3P]-3: PA21; PB12; PC63;
 Mix[3P]-4: PA13; PB12; PC63;
 Mix[3P]-5: PA13; PA21; PC63;
 Mix[3P]-6: PA13; PA21; PB12;
 Mix[3P]-7: PA21; PB12; PC29;
 Mix[3P]-8: PA13; PB12; PC29;
 Mix[3P]-9: PA13; PA21; PB12;
 Mix[4P]-1: PA11; PA13; PA21; PB12;
 Mix[4P]-2: PA12; PA13; PA21; PB12;

- Mix[4P]-3: PA13; PA21; PA22; PB12;
 Mix[4P]-4: PA11; PA13; PA21; PA22;
 Mix[4P]-5: PA13; PA21; PB12; PC63;
 Mix[4P]-6: PA11; PA12; PA21; PB12;
 5 Mix[4P]-7: PA11; PA12; PA13; PB12;
 Mix[4P]-8: PA11; PA13; PA21; PB12;
 Mix[4P]-9: PA12; PA13; PA21; PB12;
 Mix[4P]-10: PA13; PA21; PB12; PC29;
 Mix[5P]-1: PA12; PA13; PA21; PA22; PB12;
 10 Mix[5P]-2: PA11; PA13; PA21; PA22; PB12;
 Mix[5P]-3: PA11; PA12; PA13; PA21; PB12;
 Mix[5P]-4: PA11; PA12; PA21; PA22; PB12;
 Mix[5P]-5: PA11; PA12; PA13; PA22; PB12;
 Mix[5P]-6: PA11; PA12; PA13; PA21; PA22;
 15 Mix[5P]-7: PA13; PA21; PA22; PB12; PC63;
 Mix[5P]-8: PA13; PA21; PB12; PC29; PC63;
 Mix[6P]-1: PA11; PA12; PA13; PA21; PA22; PB12;
 Mix[6P]-2: PA12; PA13; PA21; PA22; PB12; PC63;
 Mix[6P]-3: PA11; PA13; PA21; PA22; PB12; PC63;
 20 Mix[6P]-4: PA11; PA12; PA13; PA21; PB12; PC63;
 Mix[6P]-5: PA12; PA13; PA21; PA22; PB12; PC63;
 Mix[6P]-6: PA11; PA13; PA21; PA22; PB12; PC63;
 Mix[6P]-7: PA11; PA12; PA21; PA22; PB12; PC63;
 Mix[6P]-8: PA11; PA12; PA13; PA22; PB12; PC63;
 25 Mix[6P]-9: PA11; PA12; PA13; PA21; PB12; PC63;
 Mix[6P]-10: PA11; PA12; PA13; PA21; PA22; PC63;
 Mix[6P]-11: PA11; PA12; PA13; PA21; PA22; PB12;
 Mix[7P]-1: PA11; PA12; PA13; PA21; PA22; PB12; PA79;
 M[7P]-2: PA11; PA12; PA13; PA21; PA22; PB12; PB49;
 30 M[7P]-3: PA11; PA12; PA13; PA21; PA22; PB12; PC29;
 M[7P]-4: PA11; PA12; PA13; PA21; PA22; PB12; PC63;
 M[9P]-11: PA12; PA13; PA21; PA22; PA79; PB12; PB49; PC29; PC63;
 M[9P]-12: PA11; PA13; PA21; PA22; PA79; PB12; PB49; PC29; PC63;
 M[9P]-13: PA11; PA12; PA21; PA22; PA79; PB12; PB49; PC29; PC63;
 35 M[9P]-14: PA11; PA12; PA13; PA22; PA79; PB12; PB49; PC29; PC63;
 M[9P]-15: PA11; PA12; PA13; PA21; PA79; PB12; PB49; PC29; PC63;
 M[9P]-16: PA11; PA12; PA13; PA21; PA22; PB12; PB49; PC29; PC63;
 M[9P]-17: PA11; PA12; PA13; PA21; PA22; PA79; PB49; PC29; PC63;
 M[9P]-18: PA11; PA12; PA13; PA21; PA22; PA79; PB12; PC29; PC63;
 40 M[9P]-19: PA11; PA12; PA13; PA21; PA22; PA79; PB12; PB49; PC63;
 M[9P]-20: PA11; PA12; PA13; PA21; PA22; PA79; PB12; PB49; PC29;
 M[10P]-40: PA11; PA13; PA14; PA20; PA21; PA23; PA67; PB12; PC27; PC60;
 M[10P]-44: PA13; PA40; PA41; PA62; PA75; PB18; PC22; PC63; PC64; PD05;
 M[10P]-71: PA21; PA85; PB14; PB18; PB85; PC19; PC61; PC63; PD05; PD08;
 45 M[10P]-75: PA02; PA10; PA12; PA13; PA18; PA21; PA27; PA79; PB86; PC26;
 M[10P]-84: PA10; PA21; PA27; PB12; PB15; PC26; PC68; PD05; PD06; PD08;
 M[10P]-98: PA11; PA12; PA13; PA14; PA20; PA21; PA27; PB16; PB18; PD06;
 M[10P]-102: PA12; PA13; PA19; PA21; PA41; PA79; PB10; PC26; PC61; PC64;

M[10P]-103: PA13; PA17; PA21; PA24; PB08; PB49; PB85; PC15; PC29; PD05;
 M[10P]-105: PA10; PA11; PA12; PA18; PA19; PA21; PA22; PB12; PC63; PD08;
 M[10P]-109: PA10; PA12; PA13; PA21; PA27; PA75; PB14; PB16; PC45; PC63;
 M[10P]-112: PA08; PA13; PA15; PA17; PA18; PA21; PA27; PB18; PB41; PB81;
 5 M[10P]-115: PA13; PA20; PA21; PA22; PA62; PA79; PB08; PB19; PC15; PC61;
 M[10P]-119: PA11; PA13; PA14; PA24; PB12; PB41; PB49; PC51; PD06; PD08;
 M[10P]-123: PA10; PA13; PA15; PA20; PA21; PA75; PB12; PB18; PB49; PC63;
 M[10P]-124: PA08; PA13; PA21; PA24; PA27; PB08; PB41; PB49; PC64; PD05;
 M[10P]-125: PA13; PA14; PA19; PA21; PA75; PB12; PC15; PC26; PC61; PC63;
 10 M[10P]-129: PA08; PA12; PA18; PA21; PA41; PB08; PB85; PC63; PC64; PD08;
 M[10P]-130: PA12; PA13; PA21; PA24; PA27; PB10; PB41; PB49; PC15; PC61;
 M[10P]-132: PA12; PA20; PA21; PA24; PC15; PC26; PC61; PC63; PC64; PD08;
 M[10P]-134: PA21; PA22; PA24; PA27; PB10; PB12; PB49; PC15; PC29; PD05;
 M[10P]-136: PA10; PA11; PA12; PA13; PA18; PA21; PB16; PB18; PC26; PC29;
 15 M[10P]-140: PA13; PA15; PA21; PA24; PA62; PA79; PB10; PB12; PC63; PC64;
 M[10P]-142: PA11; PA12; PA13; PA19; PA21; PB12; PB85; PC15; PC26; PC64;
 M[10P]-144: PA13; PA15; PA21; PA24; PA75; PB49; PC51; PC61; PC64; PD05;
 M[10P]-145: PA12; PA13; PA17; PA20; PA21; PA79; PB08; PB16; PB85; PC63;
 M[10P]-148: PA10; PA13; PA20; PA24; PB08; PB12; PB49; PC26; PC51; PC64;
 20 M[10P]-149: PA13; PA14; PA15; PA19; PA21; PB08; PB10; PB41; PC15; PC61;
 M[10P]-150: PA10; PA12; PA13; PA14; PA21; PA62; PB10; PC26; PC64; PD06;
 M[10P]-151: PA10; PA13; PA21; PA22; PA24; PA41; PB49; PC15; PC61; PC63;
 M[10P]-152: PA11; PA12; PA13; PA15; PA21; PA27; PA75; PB12; PB49; PB85;
 M[10P]-154: PA11; PA12; PA13; PA21; PA22; PA79; PB12; PB49; PC29; PC63;
 25 M[10P]-156: PA15; PA21; PA22; PB10; PB85; PC26; PC29; PC61; PC63; PC64;
 M[10P]-157: PA11; PA13; PA15; PA19; PA21; PB18; PB41; PC15; PC51; PD08;
 M[10P]-158: PA10; PA17; PA21; PA24; PA27; PA41; PB49; PC26; PC61; PC63;
 M[10P]-160: PA10; PA12; PA13; PA15; PA18; PA20; PA21; PA24; PB12; PC26;
 M[10P]-162: PA12; PA13; PA21; PA27; PA79; PB12; PB16; PB18; PC63; PC64;
 30 M[10P]-164: PA12; PA13; PA15; PA19; PA21; PA27; PB08; PB12; PB49; PD05;
 M[10P]-165: PA13; PA15; PA22; PA75; PA79; PB12; PB18; PC15; PC26; PC29;
 M[10P]-167: PA12; PA13; PA75; PB08; PB12; PB85; PC15; PC51; PC61; PC63;
 M[10P]-168: PA11; PA13; PA17; PA21; PA27; PA62; PA79; PB16; PB49; PC64;
 M[10P]-169: PA12; PA17; PA21; PA24; PA75; PA79; PB10; PB12; PB85; PC15;
 35 M[10P]-170: PA11; PA13; PA15; PA20; PA21; PA22; PC26; PC29; PC63; PC64;
 M[10P]-172: PA13; PA20; PA21; PA75; PB10; PB12; PB18; PB49; PC29; PC61;
 M[10P]-173: PA10; PA12; PA13; PA14; PA19; PA20; PA21; PA79; PC15; PC63;
 M[10P]-174: PA11; PA12; PA13; PA27; PA62; PB12; PB16; PB49; PB85; PC26;
 M[10P]-175: PA11; PA12; PA13; PA21; PA22; PA75; PB10; PB49; PC15; PC63;
 40 M[10P]-176: PA11; PA13; PA21; PA24; PA62; PB12; PC15; PC26; PC61; PC64;
 M[10P]-178: PA11; PA14; PA15; PA18; PA19; PA21; PA22; PB16; PC15; PC63;
 M[10P]-180: PA08; PA11; PA12; PA13; PA15; PA17; PA21; PA62; PB12; PC63;
 M[10P]-181: PA13; PA19; PA21; PA22; PA79; PB08; PB10; PB18; PC15; PC29;
 M[10P]-182: PA11; PA15; PA20; PA21; PA22; PA75; PB12; PC29; PC63; PC64;
 45 M[10P]-183: PA11; PA12; PA13; PA27; PA79; PB16; PB18; PB49; PC15; PC63;
 M[10P]-184: PA11; PA13; PA15; PA20; PA21; PA22; PA27; PA75; PB12; PC29;
 M[10P]-186: PA11; PA15; PA19; PA20; PA21; PA22; PA27; PB10; PB12; PB85;
 M[10P]-188: PA11; PA12; PA13; PA19; PA21; PA75; PA79; PB49; PC15; PC63;

M[10P]-190: PA11; PA13; PA21; PA22; PA27; PA75; PB10; PB12; PB49; PC63;
 M[10P]-191: PA13; PA19; PA20; PA21; PA79; PB10; PB18; PB85; PC29; PC61;
 M[10P]-192: PA12; PA13; PA19; PA21; PA27; PA75; PB12; PB49; PC63; PC64;
 M[10P]-193: PA13; PA15; PA20; PA21; PA79; PB16; PC15; PC26; PC29; PC61;
 5 M[10P]-194: PA11; PA17; PA21; PA22; PA27; PB12; PB49; PC15; PC63; PC64;
 M[10P]-196: PA11; PA12; PA13; PA15; PA21; PA75; PB10; PB18; PC51; PC63;
 M[10P]-197: PA11; PA12; PA13; PA15; PA21; PA24; PA79; PB49; PC61; PC64;
 M[10P]-198: PA12; PA15; PA21; PA22; PB12; PB16; PB85; PC15; PC26; PC29;
 M[10P]-199: PA11; PA12; PA13; PA15; PA21; PA75; PB12; PC15; PC63; PC64;
 10 M[10P]-201: PA11; PA13; PA20; PA21; PA22; PA27; PA75; PB12; PB49; PC64;
 M[10P]-202: PA11; PA21; PA75; PB10; PB12; PB61; PC26; PC29; PC42; PC63;
 M[10P]-205: PA11; PA13; PA19; PA21; PA22; PA27; PA75; PB12; PB41; PC63;
 M[10P]-207: PA11; PA13; PA21; PA75; PB10; PB12; PB41; PB51; PC45; PC63;
 M[10P]-209: PA12; PA13; PA21; PA24; PA75; PB12; PB16; PB49; PB60; PC15;
 15 M[10P]-210: PA13; PA15; PA20; PA22; PA27; PB08; PB10; PB12; PB18; PC63;
 M[10P]-212: PA11; PA13; PA21; PA22; PA27; PA75; PB12; PB16; PC15; PC63;
 M[10P]-214: PA11; PA12; PA15; PA21; PA22; PA75; PB12; PB49; PC15; PC61;
 M[10P]-215: PA11; PA12; PA13; PA21; PA25; PB12; PC02; PC29; PC51; PC64;
 M[10P]-216: PA11; PA12; PA13; PA21; PA27; PA75; PA85; PB18; PB49; PC49;
 20 M[10P]-217: PA18; PA20; PA21; PA22; PA75; PB10; PB12; PB16; PB85; PC64;
 M[10P]-218: PA10; PA13; PA15; PA20; PA22; PA27; PB18; PB49; PC51; PC63;
 M[10P]-219: PA12; PA13; PA15; PA19; PA21; PA22; PA75; PB10; PB12; PC26;
 M[10P]-220: PA11; PA12; PA13; PA19; PA21; PA22; PA27; PB12; PB49; PC63;
 M[10P]-221: PA13; PA15; PA19; PA20; PA21; PA41; PA75; PB12; PC15; PC60;
 25 M[10P]-222: PA05; PA13; PA21; PA22; PA75; PA79; PB49; PC26; PC29; PC63;
 M[10P]-223: PA11; PA12; PA13; PA22; PA25; PA75; PB12; PB49; PC61; PC64;
 M[10P]-225: PA12; PA13; PA21; PA27; PB10; PB12; PC29; PC63; PC64; PD08;
 M[10P]-226: PA11; PA12; PA21; PA39; PA79; PB18; PB49; PB85; PC61; PC63;
 M[10P]-227: PA11; PA12; PA15; PA20; PA21; PA22; PB12; PB42; PC15; PC26;
 30 M[10P]-229: PA13; PA20; PA62; PA79; PB12; PB46; PB85; PC29; PC64; PD08;
 M[10P]-230: PA11; PA12; PA13; PA21; PA22; PA75; PB10; PB12; PB58; PC61;
 M[10P]-232: PA12; PA13; PA15; PA22; PA27; PA83; PB12; PB49; PC26; PC63;
 M[10P]-235: PA12; PA13; PA15; PA20; PA21; PA22; PC29; PC51; PC63; PC65;
 M[10P]-236: PA12; PA13; PA20; PA21; PA27; PA75; PB50; PB85; PC29; PC64;
 35 M[10P]-237: PA11; PA13; PA15; PA21; PA27; PA75; PB10; PB12; PC63; PC70;
 M[10P]-239: PA10; PA11; PA12; PA20; PA21; PB10; PB41; PB49; PC63; PD08;
 M[10P]-240: PA11; PA12; PA15; PA21; PA27; PA79; PB46; PC29; PC51; PC63;
 M[10P]-241: PA11; PA14; PA20; PA21; PA22; PA75; PB12; PB18; PC15; PC29;
 M[10P]-242: PA11; PA12; PA13; PA21; PA22; PA79; PB18; PB49; PC51; PC63;
 40 M[10P]-244: PA11; PA12; PA13; PA20; PA22; PA27; PB10; PB12; PC63; PC64;
 M[10P]-246: PA11; PA12; PA13; PA21; PA61; PA79; PB10; PB12; PB18; PB85;
 M[10P]-247: PA11; PA12; PA21; PA27; PB16; PB49; PC21; PC51; PC61; PC63;
 M[10P]-248: PA11; PA13; PA14; PA19; PA21; PA75; PA79; PB14; PB49; PC15;
 M[10P]-250: PA11; PA12; PA15; PA20; PA21; PA41; PA75; PB49; PC26; PC63;
 45 M[10P]-251: PA12; PA13; PA21; PA22; PA27; PA79; PB10; PB12; PB13; PB16;
 M[10P]-252: PA11; PA13; PA20; PA21; PA44; PB08; PB12; PB18; PB88; PD06;
 M[10P]-253: PA10; PA11; PA12; PA13; PA15; PA21; PA22; PA27; PA79; PC63;
 M[10P]-255: PA11; PA13; PA21; PA75; PB12; PB18; PB49; PC15; PC26; PC63;

- M[10P]-256: PA11; PA12; PA13; PA21; PA22; PA23; PA75; PA79; PC29; PC64;
M[10P]-257: PA13; PA20; PA21; PA27; PB12; PB49; PB83; PC29; PC63; PD00;
M[10P]-258: PA12; PA13; PA20; PA21; PA22; PA75; PB10; PB12; PC15; PC63;
M[10P]-262: PA11; PA12; PA21; PA31; PA75; PB49; PB85; PC61; PC63; PC64;
5 M[10P]-264: PA21; PA24; PA27; PA31; PA79; PB10; PB18; PB49; PC15; PC63;
M[10P]-267: PA11; PA12; PA21; PA22; PA75; PC29; PC51; PC52; PC61; PC63;
M[10P]-268: PA11; PA12; PA13; PA15; PA27; PA75; PB10; PB12; PB45; PC64;
M[10P]-269: PA13; PA20; PA21; PA22; PA79; PB10; PB11; PB49; PC61; PC63;
M[10P]-270: PA11; PA12; PA13; PA15; PA21; PA75; PB12; PB14; PB18; PC49;
10 M[10P]-271: PA11; PA13; PA20; PA22; PA27; PA45; PB08; PB12; PB49; PC64;
M[10P]-272: PA13; PA21; PA24; PA79; PB10; PB12; PB41; PC15; PC29; PC63;
M[10P]-273: PA11; PA15; PA21; PA22; PA27; PA75; PB12; PB14; PC64; PD00;
M[10P]-274: PA12; PA13; PA21; PA22; PB12; PB49; PC26; PC51; PC63; PC64;
M[10P]-278: PA13; PA15; PA21; PA22; PA75; PA79; PB10; PB12; PB49; PC63;
15 M[10P]-280: PA11; PA12; PA15; PA21; PB08; PB12; PB18; PB49; PB58; PC51;
M[10P]-281: PA11; PA12; PA20; PA21; PA22; PA27; PA29; PA75; PB16; PC63;
M[10P]-282: PA10; PA13; PA21; PA22; PA27; PA79; PB12; PC15; PC29; PC61;
M[10P]-284: PA11; PA13; PA20; PA21; PA22; PA27; PA95; PC15; PC29; PC63;
M[10P]-285: PA11; PA13; PA14; PA21; PA79; PB10; PB12; PB18; PB49; PB82;
20 M[10P]-286: PA11; PA12; PA13; PA14; PA20; PA21; PA79; PC51; PC64; PC82;
M[10P]-287: PA12; PA13; PA19; PA20; PA75; PB10; PB12; PC26; PC63; PD08;
M[10P]-289: PA12; PA13; PA15; PA19; PA20; PA21; PB10; PB12; PC29; PC51;
M[10P]-290: PA11; PA12; PA13; PA22; PA27; PA75; PB85; PC29; PC63; PD07;
M[10P]-292: PA12; PA13; PA15; PA75; PA79; PB10; PB12; PC29; PC45; PC63;
25 M[10P]-293: PA00; PA11; PA12; PA13; PA21; PA27; PA75; PB18; PB85; PC26;
M[10P]-294: PA12; PA13; PA15; PA21; PA22; PA79; PB12; PB49; PC51; PC63;
M[10P]-296: PA11; PA13; PA15; PA17; PA21; PA62; PA79; PB10; PB12; PC63;
M[10P]-300: PA12; PA13; PA17; PA48; PB10; PB12; PB16; PC26; PC29; PC63;
M[10P]-301: PA11; PA12; PA13; PA20; PA21; PA75; PB08; PC15; PC35; PC51;
30 M[10P]-302: PA11; PA12; PA13; PA21; PA70; PB12; PB16; PB18; PC26; PC63;
M[10P]-303: PA11; PA12; PA15; PA21; PA75; PA79; PB10; PB12; PC29; PC64;
M[10P]-307: PA12; PA13; PA21; PB10; PB12; PB18; PB78; PC51; PC63; PD08;
M[10P]-308: PA17; PA21; PA22; PA27; PB12; PB49; PB85; PC15; PC26; PC29;
M[10P]-309: PA13; PA22; PA27; PA53; PA62; PA75; PB10; PC51; PC63; PD06;
35 M[10P]-311: PA12; PA15; PA19; PA21; PA79; PB12; PB49; PC26; PC63; PC83;
M[10P]-312: PA00; PA11; PA13; PA21; PA22; PA29; PA75; PB10; PB85; PC29;
M[10P]-313: PA13; PA15; PA21; PA22; PA27; PA75; PB49; PC51; PC61; PC63;
M[10P]-314: PA08; PA12; PA13; PA21; PB10; PB12; PB49; PC15; PC32; PC64;
M[10P]-315: PA11; PA13; PA21; PA47; PA75; PB16; PB83; PC15; PC51; PC63;
40 M[10P]-317: PA11; PA12; PA19; PA21; PA22; PA79; PB10; PB16; PC29; PC63;
M[10P]-319: PA12; PA13; PA22; PA75; PA79; PA98; PB08; PB10; PB85; PC63;
M[10P]-321: PA11; PA12; PA13; PA20; PA21; PA22; PB49; PB85; PC55; PC63.

The corresponding water extracts (or processed samples) of such herbal compositions were prepared according to the previously described Method B using 20mL of water containing 100g/l of sucrose for water extraction and including centrifugation step before filtration.

The biological activity of the processed sample was determined by using the following *Staphylococcus aureus* strains: ATCC 25923, ATCC 49476, ATCC 6538, ATCC 51740, ATCC 29213 and ATCC 14775, , with

the following final dilutions of processed samples: 1:10, 1:20, 1:40 and 1:80 for processed samples containing less than 10 spoons of plant powder (M[2P] to M[9P]) or 1:20, 1:63 and 1:200 for processed samples containing 10 spoons of plant powder (M[10P]), according to the previously described Method C.

- 5 The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from the 2 plants of Mix[2P]-1 at 1:20 dilution is given in Table 4-1.

Table 4-1: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[2P]-1 at 1:20 dilution

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
M[10P]-124	28%	47%
M[10P]-236	24%	50%
M[10P]-193	27%	54%
M[10P]-286	45%	38%
Mix[5P]-6	39%	45%
M[10P]-197	32%	55%
M[10P]-112	40%	49%
M[10P]-157	41%	52%
M[10P]-136	39%	55%
M[10P]-144	45%	52%
M[10P]-191	39%	57%
M[10P]-168	44%	53%
M[10P]-115	44%	54%
M[10P]-98	48%	51%
M[10P]-102	51%	54%
M[10P]-150	49%	59%
M[10P]-103	51%	58%
M[10P]-130	57%	61%
M[10P]-149	56%	63%
M[10P]-181	65%	70%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
M[10P]-75	23%	40%
M[10P]-216	22%	41%
M[10P]-256	19%	47%
M[10P]-248	14%	48%

(C)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
Mix[3P]-2	27%	26%
Mix[4P]-4	27%	29%
M[10P]-293	28%	32%
M[10P]-301	28%	28%
M[10P]-312	40%	37%

20 processed samples obtained from herbal compositions containing 5 or 10 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[2P]-1 (Table 4-1 A). Four 10 plant mixes extracts were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 4-1 B). Five processed samples containing 3, 4 or 10 plants were found to have higher antimicrobial activity and equivalent or lower antibiofilm activity (Table 4-1 C).

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from herbal compositions containing the 2 plants of Mix[2P]-2 at 1:20 dilution is given in Table 4-2.

Table 4-2: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[2P]-2 at 1:20 dilution. (A) Mixes extracts showing higher antimicrobial and antibiofilm activity. (B) Mixes extracts showing antibiofilm activity and equivalent or lower antimicrobial activity. (C) Mixes extracts showing antimicrobial activity and equivalent or lower antibiofilm activity.

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-2	12%	-3%
M[10P]-308	16%	13%
Mix[3P]-7	23%	9%
M[10P]-264	15%	29%
M[10P]-317	36%	9%
M[10P]-217	17%	28%
M[10P]-280	21%	27%
Mix[6P]-7	36%	16%
M[10P]-267	17%	36%
Mix[3P]-3	27%	28%
M[10P]-247	16%	40%
M[10P]-214	23%	33%
M[10P]-303	35%	22%
M[10P]-281	17%	42%
M[10P]-273	19%	41%
Mix[5P]-4	34%	27%
M[10P]-198	16%	46%
M[10P]-241	26%	38%
M[10P]-132	23%	41%
M[10P]-239	28%	44%
M[10P]-262	25%	48%
M[10P]-129	28%	46%
M[10P]-186	31%	44%
M[10P]-311	40%	38%
M[10P]-134	32%	48%
M[10P]-182	40%	44%
M[10P]-178	39%	45%
M[10P]-202	37%	50%
M[10P]-156	40%	48%
M[10P]-194	41%	49%
M[10P]-105	47%	44%
M[10P]-240	42%	61%
M[10P]-169	53%	57%
M[9P]-13	100%	86%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-2	12%	-3%
M[10P]-84	9%	27%
M[10P]-158	12%	30%
M[10P]-071	11%	35%
M[10P]-227	12%	42%
M[10P]-250	11%	44%
M[10P]-226	13%	51%

(C)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-2	12%	-3%
Mix[4P]-6	19%	-5%

33 processed samples obtained from herbal compositions containing 3, 5, 9 or 10 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[2P]-2 (Table 4-2 A). Six 10 plant mixes extracts were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 4-2 B). One 4 plant mix extract was found to have higher antimicrobial activity and equivalent antibiofilm activity (Table 4-2 C).

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from herbal compositions containing the 2 plants of Mix[2P]-3 at 1:20 dilution is given in Table 4-3.

Table 4-3: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[2P]-3 at 1:20 dilution. (A) Mixes extracts showing higher antimicrobial and antibiofilm activity. (B) Mixes extracts showing antimicrobial activity and equivalent or lower antibiofilm activity.

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-3	12%	28%
M[10P]-44	15%	35%
M[10P]-271	13%	47%
M[10P]-218	19%	42%
M[10P]-268	16%	46%
Mix[3P]-4	17%	47%
M[10P]-210	22%	42%
M[10P]-165	22%	42%
Mix[5P]-5	29%	36%
M[10P]-287	34%	31%
M[10P]-148	28%	41%
M[10P]-290	31%	39%
M[10P]-229	15%	56%
M[6P]-8	34%	40%
M[10P]-319	38%	37%
M[10P]-244	21%	56%
M[10P]-223	23%	55%
M[10P]-232	25%	61%
M[10P]-119	42%	48%
M[10P]-183	39%	51%
M[10P]-292	47%	48%
M[10P]-174	47%	59%
M[10P]-167	80%	71%
M[9P]-14	90%	83%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-3	12%	28%
Mix[4P]-7	17%	5%
Mix[3P]-8	19%	14%
M[10P]-300	20%	15%
M[10P]-309	33%	2%

23 processed samples obtained from herbal compositions containing 3, 5, 9 or 10 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[2P]-3 (Table 4-3 A). Four processed samples obtained from herbal compositions containing 3, 4 or 10 plants were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 4-3 B).

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from herbal compositions containing the 3 plants of Mix[3P]-1 at 1:20 dilution is given in Table 4-4.

Table 4-4: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[3P]-1 at 1:20 dilution. (A) Mixes extracts showing higher antimicrobial and antibiofilm activity. (B) Mixes extracts showing antibiofilm activity and equivalent or lower antimicrobial activity. (C) Mixes extracts showing antimicrobial activity and equivalent or lower antibiofilm activity.

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[3P]-1	31%	15%
Mix[4P]-8	37%	20%
Mix[6P]-10	33%	25%
M[10P]-314	34%	24%
Mix[4P]-2	37%	22%
Mix[6P]-11	35%	27%
Mix[4P]-3	41%	24%
M[10P]-315	39%	27%
M[10P]-284	41%	31%
M[10P]-321	37%	39%
M[10P]-109	36%	43%
M[10P]-302	44%	35%
Mix[5P]-8	43%	40%
M[6P]-9	51%	35%
M[10P]-160	35%	52%
M[10P]-212	43%	45%
M[10P]-274	38%	51%
M[10P]-246	34%	56%
M[10P]-205	42%	48%
M[10P]-285	52%	39%
M[10P]-201	37%	54%
M[10P]-222	32%	59%
Mix[4P]-10	52%	40%
M[10P]-230	38%	54%
M[10P]-220	43%	50%
Mix[5P]-2	51%	42%
Mix[5P]-1	50%	44%
M[10P]-215	38%	55%
Mix[6P]-6	56%	38%
M[10P]-235	40%	54%
M[10P]-207	43%	53%
Mix[5P]-3	51%	48%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[3P]-1	31%	15%
Mix[3P]-6	16%	20%
Mix[4P]-9	30%	21%
Mix[3P]-9	19%	23%
M[10P]-282	24%	26%
Mix[3P]-5	29%	36%
M[10P]-269	14%	37%
M[10P]-278	18%	40%
M[10P]-219	30%	42%
M[10P]-255	20%	43%
M[10P]-270	15%	43%
M[10P]-251	13%	44%
M[10P]-258	20%	44%
M[10P]-209	21%	45%
M[10P]-221	17%	49%
M[10P]-253	21%	49%
M[10P]-252	18%	50%
M[10P]-257	18%	51%
M[10P]-242	29%	54%

(C)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[3P]-1	31%	15%
Mix[4P]-1	37%	15%
M[10P]-313	38%	12%

M[10P]-307	58%	43%
M[10P]-289	60%	41%
M[10P]-272	40%	62%
M[10P]-184	48%	55%
M[10P]-151	48%	57%
M[10P]-123	51%	55%
M[6P]-1	60%	50%
M[6P]-5	61%	51%
M[7P]-3	61%	54%
M[10P]-192	57%	60%
M[10P]-164	58%	59%
M[7P]-1	62%	57%
M[10P]-296	63%	60%
M[6P]-4	73%	51%
Mix[5P]-7	64%	60%
M[6P]-2	68%	57%
M[10P]-225	60%	66%
M[10P]-188	65%	61%
Mix[4P]-5	61%	66%
M[10P]-125	66%	62%
M[10P]-173	65%	67%
M[10P]-237	69%	63%
M[7P]-3	71%	64%
M[10P]-294	71%	65%
M[10P]-199	70%	69%
M[10P]-142	71%	69%
Mix[6P]-3	72%	68%
M[10P]-145	72%	69%
M[10P]-180	72%	70%
M[10P]-140	72%	71%
M[10P]-162	73%	70%
M[10P]-176	74%	70%
M[10P]-172	72%	73%
M[10P]-152	71%	74%
M[10P]-40	77%	68%
M[7P]-4	72%	76%
M[10P]-170	80%	74%
M[10P]-196	78%	77%
M[10P]-190	82%	76%
M[10P]-154	80%	79%
M[10P]-175	84%	77%
M[9P]-20	93%	81%
M[9P]-17	95%	84%
M[9P]-19	95%	86%
M[9P]-11	96%	91%

M[9P]-18	98%	89%
M[9P]-12	103%	91%
M[9P]-15	106%	88%
M[9P]-16	106%	91%

80 processed samples obtained from herbal compositions containing 4, 5, 6, 7, 9 or 10 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[3P]-1 (Table 4-4 A). 18 processed samples obtained from herbal compositions containing 3, 4, or 10 plants were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 4-4 B). Two processed samples obtained from herbal compositions 4 and 10 plants were found to have higher antimicrobial activity and equivalent antibiofilm activity (Table 4-4 C).

Comparative Example 2

As a comparative Example, we illustrate several 3 or 10 plants mixes extracts obtained from herbal compositions containing only one plant among the 3 plants of Example 1 (PA13, PA21, PB12), which have been found to have none or poor antimicrobial and antibiofilm activities against *Staphylococcus aureus*.

Ten herbal compositions, whose content is reported in List 4-8, were prepared according to the previously described Method A.

List 4-8 : herbal compositions of processed samples

M[3P]-31: PA21; PA05; PB06;
M[3P]-35: PA21; PA76; PB50;
M[3P]-37: PA21; PB06; PB50;
M[3P]-43: PB12; PA05; PC34;
M[3P]-45: PB12; PA76; PB50;
M[10P]-65: PA01; PA08; PA25; PA29; PA37; PB12; PB50; PC15; PC37; PD05;
M[10P]-70: PA10; PA16; PA21; PA25; PA37; PA75; PB18; PB45; PC04; PC57;
M[10P]-72: PA08; PA21; PA62; PA70; PA85; PB19; PB31; PB45; PB49; PC26;
M[10P]-89: PA10; PA13; PA18; PA19; PA37; PA41; PA75; PB49; PC21; PC67;
M[10P]-306: PA15; PA21; PA22; PA62; PA75; PB18; PB49; PB85; PC49; PC61;

Ten water extracts (or processed samples) were prepared starting from the corresponding herbal compositions.

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains for such processed samples at 1:20 dilution were determined according to the previously reported Method C and are given in Table 4-5.

Table 4-5: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed samples at 1:20 dilution.

	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
Mix[2P]-2	12%	-3%
Mix[2P]-3	12%	28%

M[3P]-45	-4%	-5%
M[3P]-35	0%	-8%
M[3P]-37	2%	-9%
M[3P]-31	2%	-9%
M[3P]-43	1%	-1%
M[10P]-72	3%	1%
M[10P]-70	1%	15%
M[10P]-306	10%	6%
M[10P]-65	3%	17%
M[10P]-89	10%	27%

All tested processed samples, obtained from herbal compositions composed of 3 or 10 plants, which included one plant among the 3 plants of Example 1 (PA13, PA21, PB12), showed lower antimicrobial activity than the three 2 plant mixes extracts of reference Mix[2P]-1 (obtained from PA13 and PA21), Mix[2P]-2 (obtained from PA21 and PB12) and Mix[2P]-3 (obtained from PA13 and PB12) and lower antibiofilm activity than Mix[2P]-1 and Mix[2P]-3.

Example 5

In this Example, we illustrate simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus* of 390 mixtures extracts obtained from herbal compositions including at least 2 plants among the 3 plants described in Example 1 (cf Table 1-1) and additional plants to assemble mixtures of 9 to 20 plants.

The dried plants of pharmaceutical grade were obtained from “Pharmacie Fontgèze” (Clermont-Ferrand, France, plants PA00 to PB12) and “Pharmacie St Herem” (Clermont-Ferrand, France, plants PB13 to PD02).

Herbal compositions containing from 9 to 20 plant powders were prepared according to the previously described Method A.

The 4 reference herbal compositions Mix[2P]-1, Mix[2P]-2, Mix[2P]-3, Mix[3P]-1 and 390 new herbal compositions were prepared having the content showed below (List 5-3).

List 5-3 : herbal compositions of processed samples

Mix[2P]-1: PA13; PA21;

Mix[2P]-2: PA21; PB12;

Mix[2P]-3: PA13; PB12;

Mix[3P]-1: PA13; PA21; PB12;

M[09P]-1: PA13; PA14; PA20; PA21; PA23; PA67; PB12; PC27; PC60;

M[09P]-2: PA11; PA14; PA20; PA21; PA23; PA67; PB12; PC27; PC60;

M[09P]-3: PA11; PA13; PA20; PA21; PA23; PA67; PB12; PC27; PC60;

M[09P]-4: PA11; PA13; PA14; PA21; PA23; PA67; PB12; PC27; PC60;

M[09P]-6: PA11; PA13; PA14; PA20; PA23; PA67; PB12; PC27; PC60;

M[09P]-6: PA11; PA13; PA14; PA20; PA21; PA67; PB12; PC27; PC60;

M[09P]-7: PA11; PA13; PA14; PA20; PA21; PA23; PB12; PC27; PC60;

M[09P]-8: PA11; PA13; PA14; PA20; PA21; PA23; PA67; PC27; PC60;

M[09P]-9: PA11; PA13; PA14; PA20; PA21; PA23; PA67; PB12; PC60;

M[09P]-10: PA11; PA13; PA14; PA20; PA21; PA23; PA67; PB12; PC27;

M[09P]-21: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12;

- M[10P]-41: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20;
M[10P]-303: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33;
M[11P]-1: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA11;
M[11P]-2: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA14;
5 M[11P]-3: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA25;
M[11P]-4: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA26;
M[11P]-5: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA28;
M[11P]-6: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA29;
M[11P]-7: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA67;
10 M[11P]-8: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PA75;
M[11P]-9: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49;
M[11P]-10: PA13; PA20; PA21; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC60;
M[11P]-11: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39;
M[11P]-12: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21;
15 M[12P]-1: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39; PA22;
M[12P]-2: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21; PA29;
M[13P]-1: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39; PA22; PA20;
M[13P]-2: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21; PA29; PC20;
M[14P]-1: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39; PA22; PA20; PC26;
20 M[14P]-2: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21; PA29; PC20; PC49;
M[15P]-1: PA10; PA11; PA13; PA15; PA20; PA21; PA23; PA26; PA33; PB08; PB12; PB60; PB88; PC12;
PC20;
M[15P]-5: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39; PA22; PA20; PC26;
PC20;
25 M[16P]-1: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PC11; PC12; PC20; PC51;
PC55; PC67;
M[16P]-2: PA11; PA12; PA13; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33;
PC37; PC49;
M[16P]-3: PA11; PA12; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33;
30 PC37; PC49;
M[16P]-5: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39; PA22; PA20; PC26;
PC20; PA12;
M[16P]-6: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21; PA29; PC20; PC49;
PA14; PA14;
35 M[17P]-1: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PC11; PC11; PC12; PC20;
PC51; PC55; PC67;
M[17P]-2: PA11; PA12; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26;
PC33; PC37; PC49;
M[17P]-3: PA11; PA12; PA13; PA13; PA13; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26;
40 PC33; PC37; PC49;
M[17P]-5: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PC11; PC11; PC12; PC20;
PC51; PC55; PC67;
M[17P]-4: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA33; PB08; PB12; PB12; PB60;
PB88; PC12; PC20;
45 M[17P]-6: PA10; PA11; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11;
PC52; PC60; PC71;
M[17P]-8: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21; PA29; PC20; PC49;
PA14; PA14; PA22;

- M[17P]-9: PA13; PA13; PA13; PC49; PC37; PA11; PA20; PA22; PA12; PC26; PB60; PA39; PC33; PC20; PA14; PA14; PA21;
- M[18P]-1: PA10; PA11; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;
- 5 M[18P]-2: PA11; PA11; PA12; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[18P]-3: PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- 10 M[18P]-4: PA11; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[18P]-5: PA11; PA12; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[18P]-6: PA11; PA12; PA13; PA13; PA13; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- 15 M[18P]-7: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[18P]-9: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[18P]-9: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- 20 M[18P]-10: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[18P]-11: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- 25 M[18P]-12: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB60; PC20; PC26; PC33; PC37; PC49;
- M[18P]-13: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PC20; PC26; PC33; PC37; PC49;
- M[18P]-14: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC26; PC33; PC37; PC49;
- 30 M[18P]-15: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC33; PC37; PC49;
- M[18P]-16: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC37; PC49;
- 35 M[18P]-17: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC49;
- M[18P]-18: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37;
- 40 M[18P]-23: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;
- M[18P]-19: PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[18P]-20: PA10; PA10; PA11; PA13; PA15; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- 45 M[18P]-21: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[18P]-22: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB60; PB88; PC12; PC20;

- M[18P]-24: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC12; PC20; PC51; PC55; PC67;
- M[18P]-25: PA10; PA11; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;
- 5 M[18P]-31: PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[18P]-33: PA11; PA11; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- 10 M[18P]-34: PA11; PA11; PA13; PA13; PA13; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[18P]-34: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[18P]-35: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39; PA22; PA20; PC26; PC20; PA12; PA14; PA14;
- 15 M[18P]-36: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21; PA29; PC20; PC49; PA14; PA14; PA22; PA20;
- M[18P]-37: PA13; PA13; PA13; PC49; PC37; PA11; PA20; PA22; PA12; PC26; PB60; PA39; PC33; PC20; PA14; PA14; PA21; PA29;
- M[19P]-1: PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- 20 M[19P]-2: PA11; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-3: PA11; PA13; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- 25 M[19P]-4: PA11; PA13; PA14; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-5: PA11; PA13; PA14; PA20; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-6: PA11; PA13; PA14; PA20; PA21; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- 30 M[19P]-7: PA11; PA13; PA14; PA20; PA21; PA25; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-8: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- 35 M[19P]-9: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-10: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-11: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- 40 M[19P]-12: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-13: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- 45 M[19P]-14: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB33; PB60; PB88; PC20; PC49; PC60;
- M[19P]-15: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB60; PB88; PC20; PC49; PC60;

- M[19P]-16: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB88; PC20; PC49; PC60;
- M[19P]-17: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PC20; PC49; PC60;
- 5 M[19P]-18: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC49; PC60;
- M[19P]-19: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC60;
- 10 M[19P]-20: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49;
- M[19P]-21: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC12; PC20; PC51; PC55; PC67;
- M[19P]-22: PA10; PA11; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;
- 15 M[19P]-23: PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[19P]-24: PA11; PA11; PA12; PA13; PA13; PA13; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
- M[19P]-25: PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- 20 M[19P]-26: PA10; PA10; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[19P]-27: PA10; PA10; PA11; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- 25 M[19P]-28: PA10; PA10; PA11; PA13; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[19P]-29: PA10; PA10; PA11; PA13; PA15; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[19P]-30: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- 30 M[19P]-31: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[19P]-32: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- 35 M[19P]-33: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[19P]-34: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB12; PB12; PB60; PB88; PC12; PC20;
- M[19P]-35: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB60; PB88; PC12; PC20;
- 40 M[19P]-36: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB88; PC12; PC20;
- M[19P]-37: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PC12; PC20;
- 45 M[19P]-38: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC20;
- M[19P]-39: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12;

M[19P]-40: PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;

M[19P]-41: PA00; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;

5 M[19P]-42: PA00; PA10; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;

M[19P]-44: PA00; PA10; PA20; PA21; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;

10 M[19P]-45: PA00; PA10; PA20; PA21; PA23; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11;
PC12; PC20; PC51; PC55; PC67;

M[19P]-46: PA00; PA10; PA20; PA21; PA23; PA24; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;

M[19P]-47: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;

15 M[19P]-48: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB12; PB12; PB12; PB12; PC11; PC11;
PC12; PC20; PC51; PC55; PC67;

M[19P]-49: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;

M[19P]-50: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11;
20 PC12; PC20; PC51; PC55; PC67;

M[19P]-51: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC20; PC51; PC55; PC67;

M[19P]-52: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC51; PC55; PC67;

25 M[19P]-53: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11;
PC11; PC12; PC20; PC55; PC67;

M[19P]-54: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC67;

30 M[19P]-55: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11;
PC11; PC12; PC20; PC51; PC55;

M[19P]-56: PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;

M[19P]-57: PA10; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;

35 M[19P]-58: PA10; PA11; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33;
PB77; PC11; PC52; PC60; PC71:

M[19P]-59: PA10; PA11; PA13; PA13; PA13; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71:

40 M[19P]-60: PA10; PA11; PA13; PA13; PA13; PA14; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33;
PB77; PC11; PC52; PC60; PC71:

M[19P]-62: PA10; PA11; PA13; PA13; PA14; PA18; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71:

M[19P]-62: PA10; PA11; PA13; PA13; PA14; PA18; PA21; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71:

45 M[19P]-63: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA67; PB00; PB07; PB12; PB33;
PB77; PC11; PC52; PC60; PC71;

M[19P]-64: PA10; PA11; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PB00; PB07; PB12; PB33;
PB77; PC11; PC52; PC60; PC71:

- M[19P]-65: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;
- M[19P]-66: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB12; PB33; PB77; PC11; PC52; PC60; PC71;
- 5 M[19P]-67: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB33; PB77; PC11; PC52; PC60; PC71;
- M[19P]-68: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB77; PC11; PC52; PC60; PC71;
- 10 M[19P]-69: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PC11; PC52; PC60; PC71;
- M[19P]-70: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC52; PC60; PC71;
- M[19P]-71: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC60; PC71;
- 15 M[19P]-72: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC71;
- M[19P]-73: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60;
- M[19P]-89: PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- 20 M[19P]-91: PA11; PA11; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[19P]-92: PA11; PA11; PA13; PA13; PA13; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- 25 M[19P]-93: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[19P]-93: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[19P]-95: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- 30 M[19P]-96: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[19P]-96: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- 35 M[19P]-98: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB28; PB38; PB60; PC37; PC52;
- M[19P]-100: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB38; PB60; PC37; PC52;
- M[19P]-100: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB60; PC37; PC52;
- 40 M[19P]-101: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PC37; PC52;
- M[19P]-102: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC52;
- 45 M[19P]-103: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37;
- M[19P]-104: PC49; PC37; PA13; PA13; PA13; PA11; PA29; PA21; PB12; PC33; PA39; PA22; PA20; PC26; PC20; PA12; PA14; PA14; PB60;

- M[19P]-105: PA11; PC37; PC33; PA39; PB60; PC26; PB12; PA13; PA13; PA13; PA21; PA29; PC20; PC49; PA14; PA14; PA22; PA20; PA12;
- M[19P]-106: PA13; PA13; PA13; PC49; PC37; PA11; PA20; PA22; PA12; PC26; PB60; PA39; PC33; PC20; PA14; PA14; PA21; PA29; PB12;
- 5 M[20P]-29: PA10; PA11; PA12; PA13; PA14; PA15; PA16; PA17; PA18; PA19; PA20; PA21; PA22; PA23; PA24; PA25; PA26; PA27; PA28; PA29;
- M[20P]-240: PA21; PA24; PA26; PA33; PA37; PA39; PA67; PB00; PB10; PB12; PB26; PB28; PB33; PB82; PB88; PC20; PC21; PC52; PC60; PC97;
- M[20P]-296: PA00; PA11; PA13; PA15; PA21; PA22; PA24; PA26; PA28; PA33; PA39; PA94; PB08; PB12;
- 10 PB20; PB28; PB33; PB99; PC26; PC52;
- M[20P]-297: PA10; PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA29; PA33; PA57; PA67; PA69; PB07; PB12; PB60; PC20; PC37; PC49; PC98;
- M[20P]-335: PA00; PA10; PA11; PA21; PA26; PA27; PA31; PA34; PA67; PA69; PA81; PA89; PB01; PB10; PB12; PB25; PB58; PC11; PC86; PC97;
- 15 M[20P]-340: PA11; PA13; PA15; PA18; PA21; PA26; PA34; PA36; PA69; PA79; PA95; PB00; PB05; PB07; PB25; PB48; PB60; PC35; PC74; PC84;
- M[20P]-349: PA10; PA11; PA13; PA16; PA20; PA33; PA69; PA94; PB07; PB08; PB12; PC01; PC27; PC28; PC49; PC50; PC53; PC55; PC60; PC67;
- M[20P]-371: PA05; PA14; PA15; PA21; PA22; PA23; PA25; PA27; PA32; PA34; PA35; PA57; PA72; PA89;
- 20 PB12; PB39; PB88; PC20; PC55; PC74;
- M[20P]-374: PA06; PA13; PA21; PA24; PA25; PA26; PA27; PA57; PB12; PB24; PB28; PB29; PB35; PB38; PB61; PC11; PC37; PC54; PC74; PC86;
- M[20P]-375: PA11; PA15; PA19; PA20; PA21; PA22; PA28; PA29; PA67; PA81; PB10; PB12; PB26; PB28; PC23; PC52; PC58; PC60; PC73; PC98;
- 25 M[20P]-377: PA07; PA10; PA11; PA13; PA14; PA21; PA22; PA24; PA29; PA32; PA33; PA35; PA57; PB10; PB28; PB36; PB82; PC49; PC52; PC94;
- M[20P]-380: PA13; PA15; PA16; PA21; PA24; PA28; PA33; PA38; PA40; PA72; PB12; PB26; PB27; PB82; PC14; PC33; PC55; PC57; PC58; PC98;
- M[20P]-382: PA00; PA12; PA13; PA16; PA21; PA24; PA25; PA26; PA28; PA29; PA35; PA57; PA75; PB33;
- 30 PB88; PC14; PC24; PC36; PC58; PC66;
- M[20P]-383: PA10; PA21; PA22; PA23; PA25; PA27; PA81; PA85; PB12; PB26; PB28; PB39; PB77; PB82; PC01; PC17; PC37; PC50; PC52; PC67;
- M[20P]-386: PA13; PA14; PA28; PA29; PA33; PA35; PA40; PA57; PA67; PA79; PA88; PB00; PB01; PB07; PB12; PB33; PB77; PB85; PC50; PC66;
- 35 M[20P]-387: PA00; PA05; PA10; PA21; PA23; PA24; PA29; PA40; PA94; PA99; PB12; PB82; PB88; PB99; PC11; PC20; PC24; PC34; PC37; PC61;
- M[20P]-388: PA12; PA13; PA14; PA15; PA20; PA21; PA26; PA29; PA35; PA40; PA67; PA89; PA98; PB12; PB33; PC01; PC14; PC58; PC68; PC88;
- M[20P]-392: PA11; PA14; PA21; PA23; PA26; PA29; PA30; PA57; PA81; PA85; PA94; PB03; PB07; PB12;
- 40 PB29; PC33; PC51; PC53; PC71; PC73;
- M[20P]-396: PA05; PA14; PA15; PA17; PA21; PA28; PA29; PA35; PA37; PA39; PA40; PA69; PA72; PA89; PB10; PB12; PC34; PC67; PC71; PC97;
- M[20P]-401: PA13; PA15; PA16; PA23; PA26; PA29; PA33; PA34; PA85; PB12; PB26; PB33; PC21; PC50; PC52; PC53; PC55; PC58; PC67; PC71;
- 45 M[20P]-403: PA13; PA14; PA20; PA22; PA24; PA26; PA33; PA35; PB08; PB12; PB26; PB33; PB50; PB61; PC21; PC27; PC37; PC52; PC54; PC58;
- M[20P]-407: PA20; PA21; PA22; PA25; PA29; PA48; PA51; PA57; PA67; PA75; PA94; PB07; PB12; PC21; PC34; PC41; PC51; PC53; PC55; PC97;

- M[20P]-408: PA11; PA12; PA13; PA15; PA16; PA20; PA21; PA22; PA25; PA26; PA27; PA28; PA29; PB10; PB26; PC20; PC21; PC24; PC25; PC58;
- M[20P]-409: PA10; PA15; PA21; PA28; PA35; PA69; PA89; PA94; PA98; PB07; PB12; PB28; PB35; PB36; PC11; PC23; PC49; PC52; PC73; PC98;
- 5 M[20P]-416: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
- M[20P]-418: PA10; PA11; PA12; PA13; PA21; PA22; PA23; PA24; PA26; PA39; PB07; PB12; PB38; PC11; PC12; PC20; PC23; PC24; PC49; PC55;
- 10 M[20P]-420: PA10; PA14; PA21; PA22; PA23; PA29; PA30; PA33; PA37; PB01; PB12; PB26; PB33; PB99; PC20; PC21; PC36; PC37; PC73; PC98;
- M[20P]-421: PA13; PA15; PA21; PA25; PA26; PA29; PA39; PA40; PA75; PA94; PB20; PB55; PB82; PC06; PC20; PC24; PC49; PC71; PC74; PC99;
- M[20P]-422: PA10; PA11; PA13; PA18; PA21; PA23; PA25; PA29; PA33; PA57; PA70; PB01; PB07; PB12; PB88; PC11; PC27; PC33; PC55; PC97;
- 15 M[20P]-427: PA06; PA13; PA15; PA21; PA24; PA25; PA27; PA29; PA33; PA75; PA98; PB00; PB28; PB60; PC11; PC20; PC26; PC50; PC55; PC74;
- M[20P]-429: PA11; PA19; PA21; PA23; PA26; PA33; PA40; PA75; PB00; PB03; PB12; PB60; PC11; PC21; PC24; PC26; PC36; PC37; PC68; PC98;
- M[20P]-430: PA00; PA07; PA13; PA23; PA26; PA27; PA34; PA40; PA75; PB07; PB12; PB18; PB28; PB88; PC20; PC41; PC66; PC67; PC71; PC97;
- 20 M[20P]-431: PA00; PA06; PA11; PA12; PA13; PA15; PA18; PA21; PA22; PA25; PA28; PA33; PA81; PB09; PB32; PB33; PB38; PC11; PC33; PC52;
- M[20P]-432: PA12; PA14; PA19; PA21; PA24; PA26; PA33; PA67; PB05; PB09; PB12; PB28; PB32; PB46; PB60; PC20; PC49; PC58; PC71; PC73;
- 25 M[20P]-435: PA10; PA13; PA14; PA15; PA21; PA28; PA33; PA40; PA55; PA57; PB07; PB12; PB58; PB88; PB99; PC12; PC21; PC29; PC66; PC67;
- M[20P]-439: PA10; PA13; PA16; PA21; PA24; PA27; PA29; PA34; PA40; PA60; PA75; PB12; PB27; PB35; PB38; PC25; PC26; PC36; PC51; PC55;
- M[20P]-443: PA00; PA10; PA13; PA15; PA17; PA20; PA29; PA33; PA35; PA40; PB07; PB12; PB26; PB27; PB82; PC12; PC24; PC50; PC60; PD02;
- 30 M[20P]-444: PA11; PA13; PA14; PA18; PA21; PA24; PA28; PA39; PA40; PA57; PA60; PA67; PA88; PB20; PB85; PB88; PC06; PC52; PC57; PC58;
- M[20P]-445: PA11; PA13; PA17; PA20; PA21; PA29; PA30; PA33; PA35; PA40; PA57; PB00; PB12; PB28; PB60; PC11; PC20; PC35; PC50; PC51;
- 35 M[20P]-446: PA13; PA15; PA16; PA21; PA22; PA23; PA25; PA26; PA27; PA29; PA39; PA67; PA75; PB12; PB26; PB27; PC36; PC49; PC60; PC98;
- M[20P]-447: PA11; PA13; PA17; PA19; PA24; PA28; PA30; PA33; PA39; PA40; PA57; PA92; PB07; PB08; PB12; PB33; PB61; PB85; PC58; PC94;
- M[20P]-449: PA10; PA11; PA13; PA14; PA15; PA20; PA21; PA24; PA28; PA40; PA67; PA75; PA95; PB12; PB20; PB35; PC14; PC20; PC52; PC60;
- 40 M[20P]-453: PA10; PA13; PA15; PA18; PA21; PA25; PA29; PA34; PA35; PA67; PA93; PB06; PB07; PB09; PB12; PB20; PB26; PB35; PB99; PC55;
- M[20P]-455: PA00; PA00; PA11; PA11; PA13; PA14; PA18; PA20; PA25; PA28; PA33; PA70; PA75; PB00; PB12; PC20; PC23; PC26; PC67; PC67;
- 45 M[20P]-456: PA14; PA21; PA33; PA34; PA39; PB00; PB07; PB07; PB12; PB27; PB28; PB38; PB88; PC20; PC23; PC26; PC49; PC58; PC67; PC71;
- M[20P]-457: PA00; PA10; PA11; PA12; PA13; PA13; PA14; PA29; PA29; PA57; PA67; PB02; PB07; PB09; PB12; PB38; PC11; PC60; PC97; PC98;

- M[20P]-458: PA10; PA11; PA11; PA13; PA13; PA18; PA21; PA26; PA33; PA57; PA75; PA98; PB12; PB33; PB33; PB38; PB82; PC20; PC55; PC73;
- M[20P]-459: PA00; PA06; PA21; PA26; PA26; PA29; PA33; PA75; PB10; PB10; PB12; PB18; PB32; PB85; PC20; PC24; PC49; PC55; PC66; PC98;
- 5 M[20P]-461: PA06; PA21; PA29; PA29; PA33; PA33; PA39; PA57; PA67; PB07; PB07; PB12; PB26; PB32; PB32; PB85; PC20; PC27; PC52; PC58;
- M[20P]-465: PA11; PA11; PA13; PA16; PA21; PA22; PA27; PA29; PA33; PA69; PB07; PB26; PB28; PB32; PB77; PB88; PB88; PC23; PC60; PC73;
- 10 M[20P]-467: PA00; PA06; PA21; PA21; PA23; PA26; PA33; PA33; PA40; PA67; PB12; PB38; PB88; PC11; PC23; PC23; PC27; PC55; PC71; PC97;
- M[20P]-469: PA00; PA12; PA13; PA13; PA13; PA21; PA29; PA33; PA33; PA34; PA57; PB07; PC11; PC11; PC20; PC23; PC23; PC55; PC74; PC97;
- M[20P]-470: PA13; PA14; PA21; PA21; PA22; PA23; PA24; PA24; PA27; PA29; PA33; PA67; PA75; PB07; PB07; PB33; PB33; PB77; PC37; PC52;
- 15 M[20P]-472: PA10; PA10; PA11; PA13; PA21; PA25; PA25; PA28; PA33; PA39; PB07; PB12; PB88; PC12; PC12; PC20; PC20; PC24; PC55; PC71;
- M[20P]-474: PA00; PA13; PA20; PA21; PA26; PA28; PA29; PA33; PA57; PA67; PA81; PA81; PA98; PB02; PB12; PB38; PB60; PB88; PC11; PC20;
- M[20P]-475: PA11; PA12; PA12; PA13; PA28; PA28; PA29; PA29; PA32; PA52; PA52; PA75; PA81; PB12; PB20; PB33; PB60; PC21; PC49; PC51;
- 20 M[20P]-478: PA15; PA16; PA21; PA21; PA21; PA23; PA25; PA32; PA34; PA39; PB12; PB77; PB85; PB88; PB88; PB88; PB88; PC11; PC33; PC37;
- M[20P]-479: PA10; PA11; PA13; PA13; PA18; PA20; PA21; PA28; PA32; PA40; PA67; PA69; PB07; PB07; PB10; PB12; PB20; PC55; PC60; PC98;
- 25 M[20P]-480: PA00; PA06; PA13; PA22; PA23; PA26; PA26; PA27; PA29; PA40; PA81; PB07; PB10; PB10; PB12; PB12; PB18; PB32; PC49; PC71;
- M[20P]-481: PA11; PA13; PA26; PA29; PA29; PA35; PA69; PA75; PA75; PB02; PB12; PB33; PC20; PC23; PC23; PC37; PC41; PC50; PC52; PC98;
- M[20P]-482: PA06; PA10; PA11; PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA20; PA25; PA26; PA28; PA39; PA67; PB12; PC11; PC24; PC74;
- 30 M[20P]-484: PA11; PA13; PA23; PA28; PA29; PA29; PA35; PA39; PA40; PA60; PA75; PA98; PB02; PB12; PB38; PB38; PB82; PC20; PC35; PC50;
- M[20P]-485: PA13; PA14; PA14; PA21; PA21; PA21; PA26; PA26; PA26; PA26; PA29; PA29; PA52; PA57; PB00; PC11; PC11; PC21; PC51; PC67;
- 35 M[20P]-486: PA11; PA13; PA23; PA24; PA35; PA35; PA35; PA69; PA69; PA75; PB07; PB12; PB27; PC20; PC20; PC21; PC23; PC50; PC55; PC73;
- M[20P]-490: PA10; PA11; PA11; PA14; PA21; PA24; PA26; PA26; PA39; PA40; PA69; PA75; PB00; PB01; PB12; PB12; PB18; PB28; PB60; PC20;
- M[20P]-493: PA06; PA14; PA20; PA21; PA21; PA28; PA29; PA33; PA69; PA81; PA98; PB12; PB38; PB60; PC11; PC12; PC49; PC55; PC71; PD02;
- 40 M[20P]-494: PA06; PA10; PA11; PA11; PA13; PA13; PA15; PA23; PA26; PA35; PB01; PB12; PB60; PC12; PC20; PC37; PC41; PC49; PC49; PC52;
- M[20P]-495: PA10; PA10; PA13; PA14; PA18; PA20; PA21; PA26; PA29; PA29; PA57; PB02; PC11; PC11; PC20; PC20; PC24; PC33; PC49; PC49;
- 45 M[20P]-496: PA00; PA18; PA21; PA24; PA26; PA27; PA29; PA39; PA39; PA52; PA69; PA70; PA98; PB00; PB02; PB12; PC12; PC24; PC55; PC67;
- M[20P]-497: PA11; PA11; PA13; PA13; PA15; PA21; PA23; PA24; PA25; PA26; PA28; PA39; PA52; PA67; PB00; PB12; PB27; PB28; PC12; PC37;

- M[20P]-499: PA13; PA21; PA21; PA21; PA29; PA30; PA69; PA75; PB07; PB07; PB07; PB60; PC20; PC20; PC21; PC37; PC37; PC49; PC55; PD02;
- M[20P]-503: PA06; PA06; PA10; PA10; PA13; PA13; PA20; PA21; PA25; PA26; PA29; PA33; PA39; PA70; PB28; PC11; PC20; PC23; PC49; PC97;
- 5 M[20P]-506: PA11; PA16; PA21; PA23; PA29; PA39; PA57; PB12; PB27; PB32; PB88; PB88; PB88; PC12; PC20; PC21; PC37; PC71; PD02; PD02;
- M[20P]-510: PA13; PA21; PA25; PA26; PA27; PA33; PA39; PA40; PA69; PA81; PB01; PB28; PB60; PB85; PB88; PC27; PC41; PC41; PC50; PC55;
- 10 M[20P]-514: PA11; PA12; PA13; PA13; PA21; PA21; PA28; PA29; PA33; PA35; PA39; PA40; PA40; PA67; PA67; PB12; PC14; PC20; PC41; PC55;
- M[20P]-516: PA07; PA10; PA13; PA14; PA17; PA21; PA28; PA29; PA39; PA40; PA40; PA69; PB07; PB12; PC21; PC21; PC27; PC37; PD02; PD02;
- M[20P]-520: PA11; PA11; PA12; PA13; PA14; PA15; PA29; PA33; PA33; PA57; PA67; PA70; PA98; PB12; PC20; PC23; PC33; PC52; PC55; PD02;
- 15 M[20P]-523: PA10; PA11; PA18; PA20; PA21; PA33; PA39; PA67; PA67; PA69; PB07; PB12; PB28; PB88; PB88; PC21; PC33; PC35; PC49; PC50;
- M[20P]-524: PA00; PA11; PA11; PA12; PA13; PA20; PA29; PA40; PB07; PB12; PB28; PB38; PB60; PB60; PB77; PC20; PC20; PC49; PC49; PC98;
- M[20P]-525: PA11; PA11; PA13; PA15; PA21; PA26; PA26; PA27; PA28; PA29; PA29; PA30; PA40; PA52; 20 PA75; PB07; PB07; PC11; PC33; PC37;
- M[20P]-527: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26; PA26; PA26; PA33; PB08; PB12; PB12; PB60; PB88; PC12; PC20;
- M[20P]-532: PA14; PA20; PA20; PA21; PA22; PA25; PA26; PA27; PA28; PA29; PA33; PA40; PA75; PB00; PB12; PB12; PB60; PB85; PC12; PC51;
- 25 M[20P]-533: PA13; PA21; PA26; PA27; PA27; PA28; PA32; PA60; PA70; PA75; PB18; PB32; PB38; PC23; PC41; PC50; PC50; PC51; PC52; PC97;
- M[20P]-535: PA11; PA21; PA26; PA28; PA29; PA29; PA29; PA34; PA57; PB07; PB12; PB12; PB12; PB33; PB60; PC20; PC20; PC51; PC55; PC67;
- M[20P]-536: PA00; PA10; PA11; PA12; PA12; PA13; PA14; PA23; PA29; PA29; PA35; PA40; PA81; PB07; 30 PB07; PB12; PB27; PB33; PB60; PC20;
- M[20P]-538: PA21; PA23; PA25; PA26; PA27; PA28; PA75; PB07; PB12; PB27; PB32; PB88; PC11; PC11; PC12; PC20; PC33; PC49; PC55; PC97;
- M[20P]-541: PA10; PA11; PA13; PA20; PA24; PA28; PA29; PA67; PA70; PB10; PB12; PB77; PC20; PC20; PC23; PC41; PC51; PC52; PC52; PC97;
- 35 M[20P]-548: PA10; PA13; PA14; PA14; PA22; PA27; PA27; PA28; PA29; PA29; PA33; PA39; PB10; PB12; PB60; PB88; PC20; PC37; PC50; PC55;
- M[20P]-551: PA12; PA13; PA13; PA20; PA22; PA23; PA24; PA24; PA26; PA33; PA33; PA39; PA67; PB07; PB07; PB12; PB33; PC20; PC55; PC98;
- M[20P]-552: PA06; PA11; PA11; PA13; PA14; PA14; PA20; PA21; PA21; PA21; PA24; PA35; PB00; PB02; 40 PB12; PB60; PB60; PC12; PC21; PC98;
- M[20P]-555: PA00; PA10; PA11; PA18; PA20; PA20; PA21; PA25; PA26; PA26; PA39; PA39; PA39; PA89; PB12; PB12; PC20; PC20; PC49; PC74;
- M[20P]-557: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12; PB12; PB12; PB12; PC11; PC11; PC12; PC20; PC51; PC55; PC67;
- 45 M[20P]-559: PA00; PA11; PA12; PA13; PA20; PA22; PA24; PA25; PA25; PA26; PA28; PA57; PA70; PA70; PB07; PB12; PB88; PB88; PC11; PC60;
- M[20P]-560: PA10; PA10; PA11; PA12; PA13; PA14; PA20; PA25; PA28; PA28; PA29; PA67; PB12; PB38; PB60; PC24; PC35; PC41; PC51; PC97;

- M[20P]-563: PA12; PA13; PA20; PA20; PA21; PA24; PA25; PA25; PA29; PA29; PA33; PA81; PB33; PB60; PB60; PB88; PB88; PB88; PC11; PC41;
- M[20P]-564: PA07; PA13; PA13; PA21; PA26; PA37; PA39; PA67; PA67; PB02; PB07; PB07; PB12; PB12; PB28; PC12; PC37; PC97; PC97; PC97;
- 5 M[20P]-565: PA07; PA11; PA13; PA20; PA29; PA29; PA32; PA75; PA95; PA98; PB02; PB12; PB32; PB33; PB60; PB85; PC20; PC24; PC35; PC41;
- M[20P]-567: PA10; PA13; PA14; PA16; PA21; PA26; PA28; PA75; PA81; PB07; PB07; PB12; PB88; PC23; PC24; PC24; PC33; PC52; PC55; PC67;
- M[20P]-569: PA13; PA14; PA21; PA26; PA27; PA39; PA57; PA67; PB02; PB07; PB12; PB28; PC12; PC21; 10 PC49; PC55; PC55; PC67; PC67; PC98;
- M[20P]-572: PA13; PA14; PA21; PA27; PA29; PA33; PA67; PA69; PA79; PB00; PB01; PB12; PB33; PB38; PC37; PC41; PC49; PC51; PC55; PD02;
- M[20P]-573: PA10; PA11; PA13; PA13; PA13; PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;
- 15 M[20P]-574: PA13; PA21; PA22; PA23; PA25; PA29; PA35; PA67; PB01; PB12; PB12; PB35; PB88; PC24; PC37; PC37; PC51; PC52; PC52; PC66;
- M[20P]-575: PA11; PA11; PA13; PA13; PA13; PA14; PA14; PA26; PA26; PA28; PA52; PA67; PB00; PB12; PB27; PB28; PB38; PB60; PC37; PC52;
- M[20P]-576: PA11; PA11; PA12; PA13; PA13; PA13; PA14; PA14; PA20; PA21; PA22; PA29; PA39; PB12; 20 PB60; PC20; PC26; PC33; PC37; PC49;
- M[20P]-577: PA06; PA13; PA13; PA14; PA22; PA22; PA23; PA26; PA28; PA69; PA70; PB07; PB12; PB12; PB32; PC12; PC20; PC24; PC24; PC98;
- M[20P]-579: PA12; PA13; PA13; PA13; PA14; PA26; PA26; PA28; PA57; PA81; PB02; PB07; PB12; PC11; PC12; PC20; PC51; PC55; PD02; PD02;
- 25 M[20P]-580: PA00; PA11; PA12; PA13; PA20; PA21; PA21; PA25; PA26; PA26; PA33; PA75; PA75; PA75; PB28; PB88; PC20; PC76; PC98; PC98;
- M[20P]-581: PA10; PA11; PA13; PA21; PA22; PA25; PA29; PA67; PB07; PB60; PB88; PB88; PC20; PC21; PC24; PC37; PC41; PC55; PC66; PC98;
- M[20P]-582: PA10; PA11; PA12; PA13; PA20; PA21; PA23; PA25; PA39; PA75; PB07; PB07; PB38; PB38; 30 PB60; PB60; PC20; PC24; PC41; PC66;
- M[20P]-583: PA11; PA13; PA14; PA23; PA25; PA25; PA26; PA26; PA28; PA67; PB12; PB18; PB88; PC24; PC24; PC37; PC41; PC49; PC55; PC60;
- M[20P]-584: PA00; PA14; PA21; PA26; PA26; PA28; PA39; PA39; PA67; PA75; PB12; PB12; PB27; PB33; PB60; PC11; PC41; PC49; PC55; PC97;
- 35 M[20P]-588: PA07; PA13; PA14; PA21; PA26; PA26; PA26; PA26; PA32; PA67; PA67; PA75; PB12; PB27; PB33; PB60; PC20; PC20; PC23; PC24;
- M[20P]-589: PA13; PA13; PA22; PA25; PA25; PA27; PA29; PA29; PA35; PA39; PA52; PA75; PB12; PB32; PB38; PB85; PB88; PC11; PC23; PC98;
- M[20P]-590: PA11; PA13; PA13; PA14; PA25; PA29; PA33; PA35; PA35; PA35; PA39; PB02; PB12; PC11; 40 PC20; PC20; PC21; PC27; PC55; PC55;
- M[20P]-592: PA10; PA10; PA11; PA11; PA13; PA14; PA21; PA21; PA26; PA26; PA27; PA29; PA30; PA75; PB07; PB07; PB12; PB38; PC20; PC24;
- M[20P]-594: PA13; PA21; PA21; PA21; PA23; PA27; PA28; PA39; PA60; PA69; PA70; PB02; PB09; PB12; PB12; PB88; PC24; PC49; PC97; PC98;
- 45 M[20P]-595: PA11; PA13; PA22; PA23; PA25; PA28; PA28; PA28; PA29; PA29; PA39; PA67; PA69; PB12; PB60; PC12; PC20; PC24; PC49; PC55;
- M[20P]-596: PA11; PA11; PA13; PA13; PA14; PA21; PA22; PA23; PA26; PA26; PA67; PB12; PB60; PC11; PC20; PC20; PC37; PC50; PC55; PC60;

- M[20P]-599: PA13; PA13; PA14; PA16; PA26; PA29; PA75; PA75; PB07; PB12; PB60; PB60; PB60; PB60; PC12; PC20; PC20; PC37; PC37; PC52;
- M[20P]-601: PA11; PA11; PA13; PA13; PA13; PA23; PA26; PA29; PA29; PA35; PB02; PB10; PB12; PB33; PC20; PC21; PC41; PC49; PC60; PC67;
- 5 M[20P]-602: PA00; PA10; PA10; PA14; PA14; PA14; PA21; PA25; PA26; PA29; PA29; PB07; PB07; PB07; PB12; PB12; PC20; PC52; PC55; PD02;
- M[20P]-603: PA11; PA13; PA16; PA20; PA21; PA23; PA23; PA26; PA26; PA28; PA33; PA39; PA39; PA67; PB60; PB60; PC20; PC23; PC49; PC55;
- 10 M[20P]-604: PA10; PA13; PA14; PA14; PA18; PA21; PA22; PA23; PA26; PA28; PA28; PA29; PB12; PB33; PC11; PC11; PC51; PC51; PC60; PC71;
- M[20P]-605: PA11; PA11; PA11; PA11; PA13; PA13; PA13; PA20; PA25; PA29; PA39; PA39; PA69; PA81; PB07; PB12; PB88; PC20; PC41; PC55;
- M[20P]-607: PA10; PA11; PA11; PA13; PA13; PA13; PA22; PA22; PA26; PA33; PA35; PA75; PA75; PA81; PB12; PB88; PC20; PC49; PC55; PC97;
- 15 M[20P]-608: PA00; PA13; PA13; PA13; PA18; PA26; PA27; PA29; PA35; PA57; PA69; PA75; PA98; PB12; PB60; PC11; PC20; PC20; PC55; PC60;
- M[20P]-609: PA11; PA14; PA21; PA22; PA23; PA26; PA34; PA52; PA67; PA69; PA69; PB07; PB12; PB12; PB27; PB77; PC12; PC20; PC55; PC97;
- M[20P]-612: PA11; PA13; PA14; PA14; PA14; PA21; PA22; PA23; PA52; PA81; PB00; PB07; PB12; PB33; PB38; PB60; PC12; PC24; PC33; PC37;
- 20 M[20P]-613: PA10; PA13; PA13; PA14; PA21; PA21; PA22; PA23; PA26; PA28; PA28; PA29; PB12; PB12; PB27; PB60; PC20; PC21; PC21; PC60;
- M[20P]-616: PA13; PA26; PA28; PA29; PA39; PA57; PA67; PB07; PB12; PB38; PB60; PB77; PB77; PB88; PC33; PC49; PC52; PC52; PC98; PC98;
- 25 M[20P]-618: PA11; PA12; PA13; PA13; PA14; PA20; PA21; PA22; PA23; PA26; PA29; PA67; PA79; PB88; PC11; PC20; PC20; PC37; PC41; PC55;
- M[20P]-620: PA12; PA13; PA20; PA22; PA22; PA23; PA26; PA26; PA28; PA39; PA69; PA95; PB12; PB60; PB77; PC20; PC21; PC52; PC60; PC98;
- M[20P]-621: PA10; PA13; PA14; PA24; PA24; PA26; PA27; PA29; PA32; PA35; PA67; PB12; PB12; PB12; PB27; PB33; PB33; PB38; PB38; PB38;
- 30 M[20P]-622: PA10; PA12; PA14; PA14; PA15; PA20; PA21; PA26; PA27; PA32; PA40; PB10; PB12; PB12; PB88; PC20; PC20; PC21; PC52; PC60;
- M[20P]-623: PA10; PA11; PA13; PA14; PA20; PA24; PA25; PA26; PA26; PA28; PA39; PA81; PB09; PB12; PB28; PB88; PC24; PC41; PC52; PC98;
- 35 M[20P]-626: PA11; PA11; PA13; PA13; PA20; PA21; PA21; PA25; PA27; PA27; PA28; PA39; PA39; PA40; PA57; PB12; PB60; PC20; PC37; PC41;
- M[20P]-631: PA10; PA13; PA20; PA21; PA21; PA23; PA23; PA26; PA29; PA39; PA39; PA67; PB10; PB38; PB60; PC11; PC20; PC37; PC49; PC66;
- M[20P]-632: PA10; PA11; PA18; PA21; PA22; PA23; PA28; PA29; PA33; PA34; PA69; PB07; PB12; PB60; PC20; PC20; PC20; PC37; PC41; PC55;
- 40 M[20P]-633: PA10; PA11; PA11; PA13; PA20; PA34; PA39; PB00; PB07; PB12; PB12; PB60; PB88; PB88; PC11; PC12; PC23; PC35; PC49; PC52;
- M[20P]-634: PA13; PA13; PA13; PA21; PA22; PA22; PA25; PA25; PA28; PA28; PA29; PA67; PA81; PB00; PB12; PB12; PB27; PB33; PC11; PC98;
- 45 M[20P]-635: PA00; PA14; PA17; PA20; PA21; PA23; PA26; PA27; PA75; PA81; PB00; PB12; PB82; PC11; PC20; PC26; PC51; PC52; PC55; PC71;
- M[20P]-638: PA00; PA14; PA17; PA21; PA22; PA24; PA28; PA33; PA52; PA57; PA75; PB00; PB07; PB12; PB28; PB88; PC20; PC37; PC41; PC97;

- M[20P]-649: PA06; PA10; PA11; PA12; PA13; PA15; PA16; PA20; PA24; PA30; PA33; PA98; PB10; PB12; PB28; PB32; PB60; PC11; PC20; PC66;
- M[20P]-654: PA00; PA11; PA15; PA20; PA21; PA26; PA32; PA34; PA35; PA39; PA57; PB10; PB12; PB18; PC11; PC20; PC21; PC50; PC55; PC66;
- 5 M[20P]-655: PA11; PA13; PA14; PA20; PA22; PA26; PA28; PA30; PA40; PA52; PA57; PA81; PB12; PB32; PB85; PC41; PC50; PC55; PC66; PC71;
- M[20P]-658: PA10; PA11; PA12; PA13; PA14; PA15; PA17; PA20; PA21; PA28; PA29; PA52; PA67; PB10; PB12; PB18; PB60; PC20; PC41; PC66;
- 10 M[20P]-659: PA00; PA11; PA21; PA22; PA24; PA26; PA27; PA28; PA29; PA33; PA35; PA81; PB07; PB12; PB18; PB28; PB85; PC37; PC49; PC74;
- M[20P]-663: PA06; PA13; PA20; PA25; PA26; PA27; PA28; PA40; PA67; PA81; PB00; PB12; PB18; PB33; PB61; PC20; PC24; PC37; PC41; PC52;
- M[20P]-666: PA06; PA13; PA21; PA23; PA27; PA28; PA30; PA40; PA75; PB00; PB07; PB10; PB12; PB18; PB28; PB33; PC20; PC49; PC52; PC71;
- 15 M[20P]-668: PA10; PA16; PA17; PA21; PA24; PA27; PA28; PA33; PA40; PA57; PA75; PA79; PB07; PB08; PB12; PB33; PB88; PC20; PC35; PC97;
- M[20P]-670: PA11; PA12; PA13; PA14; PA20; PA23; PA27; PA28; PA35; PA39; PA67; PA79; PA81; PA89; PB10; PB12; PB18; PC20; PC21; PC51;
- M[20P]-678: PA06; PA11; PA12; PA13; PA21; PA23; PA29; PA35; PA39; PA40; PB00; PB07; PB10; PB12; PB33; PB60; PB77; PC11; PC67; PC76;
- 20 M[20P]-681: PA07; PA10; PA11; PA21; PA24; PA26; PA30; PA35; PA39; PA57; PA75; PB07; PB12; PB28; PB88; PC11; PC49; PC55; PC58; PC66;
- M[20P]-686: PA00; PA10; PA13; PA22; PA24; PA27; PA28; PA33; PA39; PA40; PA69; PB12; PB32; PB38; PB85; PB88; PC20; PC55; PC60; PC66;
- 25 M[20P]-692: PA07; PA10; PA13; PA21; PA22; PA26; PA34; PA39; PA40; PA81; PB12; PB18; PB32; PB33; PB88; PC35; PC37; PC52; PC55; PC76;
- M[20P]-694: PA07; PA11; PA13; PA14; PA15; PA16; PA21; PA27; PA28; PA29; PA35; PA39; PA40; PA75; PB12; PB33; PB88; PC50; PC51; PC52;
- M[20P]-697: PA00; PA13; PA16; PA19; PA21; PA22; PA25; PA27; PA28; PA39; PA79; PA81; PB00; PB12; PB32; PB60; PB99; PC51; PC58; PC76;
- 30 M[20P]-703: PA00; PA13; PA14; PA15; PA24; PA26; PA33; PA35; PA52; PA75; PA81; PB07; PB12; PB28; PB85; PC20; PC21; PC35; PC41; PD02;
- M[20P]-705: PA11; PA13; PA30; PA33; PA40; PA67; PA75; PB07; PB12; PB28; PB35; PB88; PC20; PC21; PC24; PC35; PC37; PC41; PC50; PC55;
- 35 M[20P]-708: PA07; PA13; PA23; PA25; PA27; PA29; PA30; PA33; PA52; PA57; PA75; PB00; PB08; PB12; PB38; PB60; PB77; PC50; PC76; PD02;
- M[20P]-709: PA11; PA13; PA17; PA20; PA21; PA23; PA26; PA27; PA28; PA40; PA69; PA81; PA92; PB00; PB07; PB12; PC20; PC41; PC52; PC66;
- M[20P]-712: PA12; PA13; PA15; PA20; PA22; PA26; PA27; PA29; PA35; PA52; PA81; PB07; PB10; PB12; PB33; PB88; PC11; PC20; PC41; PC52;
- 40 M[20P]-714: PA07; PA10; PA11; PA16; PA21; PA26; PA33; PA40; PA67; PA81; PA98; PB00; PB07; PB12; PB28; PC11; PC41; PC49; PC50; PC98;
- M[20P]-716: PA11; PA13; PA15; PA19; PA22; PA26; PA33; PA37; PA39; PA57; PB07; PB12; PB28; PB33; PB88; PB99; PC14; PC23; PC35; PC67;
- 45 M[20P]-718: PA12; PA13; PA15; PA16; PA20; PA22; PA25; PA26; PA39; PA40; PA92; PB07; PB12; PB27; PB35; PB38; PB99; PC11; PC27; PC41;
- M[20P]-721: PA19; PA20; PA21; PA24; PA25; PA27; PA29; PA30; PA55; PA57; PB12; PB32; PB58; PB60; PB88; PB99; PC41; PC50; PC60; PC67;

- M[20P]-723: PA12; PA13; PA16; PA17; PA21; PA28; PA29; PA33; PA34; PA40; PA60; PA89; PB28; PB60; PB61; PC14; PC24; PC41; PC51; PC74;
M[20P]-724: PA13; PA23; PA25; PA35; PA57; PA67; PB07; PB12; PB32; PB33; PB88; PB99; PC12; PC20; PC23; PC49; PC50; PC66; PC67; PC94;
- 5 M[20P]-725: PA00; PA13; PA14; PA20; PA22; PA25; PA26; PA28; PA33; PA40; PA57; PA81; PB12; PB16; PB33; PB88; PC11; PC24; PC51; PC74;
M[20P]-733: PA11; PA13; PA15; PA21; PA28; PA30; PA33; PA40; PB12; PB32; PB33; PB99; PC11; PC12; PC20; PC60; PC73; PC74; PC94; PD02;
- 10 M[20P]-737: PA00; PA11; PA13; PA15; PA17; PA21; PA24; PA28; PA29; PA30; PA67; PB07; PB27; PB28; PB77; PC20; PC21; PC41; PC51; PC94;
M[20P]-740: PA11; PA13; PA14; PA15; PA22; PA23; PA28; PA33; PA39; PA57; PA75; PA89; PB02; PB12; PB28; PB33; PC12; PC21; PC24; PC55;
- M[20P]-745: PA11; PA21; PA22; PA24; PA26; PA27; PA28; PA29; PA57; PA89; PB07; PB12; PB28; PB33; PC08; PC11; PC41; PC66; PC67; PC98;
- 15 M[20P]-746: PA20; PA21; PA25; PA26; PA28; PA33; PA39; PA40; PA81; PB06; PB07; PB12; PB32; PB60; PC20; PC21; PC24; PC37; PC50; PC55;
M[20P]-748: PA11; PA13; PA14; PA17; PA23; PA26; PA32; PA57; PA60; PA81; PB07; PB12; PB28; PB32; PB33; PB85; PB99; PC20; PC21; PC35;
- M[20P]-752: PA11; PA13; PA18; PA26; PA28; PA29; PA33; PA35; PA39; PA67; PA75; PB07; PB12; PB32; PB88; PB99; PC12; PC50; PC54; PC62;
- 20 M[20P]-753: PA11; PA12; PA13; PA14; PA15; PA29; PA32; PA40; PA57; PA81; PB00; PB12; PB33; PB60; PC20; PC21; PC49; PC52; PC66; PD02;
M[20P]-757: PA11; PA13; PA21; PA22; PA23; PA27; PA34; PA35; PA40; PA57; PA67; PB10; PB12; PB22; PB58; PC11; PC14; PC23; PC55; PC98;
- 25 M[20P]-758: PA10; PA11; PA13; PA20; PA21; PA23; PA28; PA29; PA30; PA40; PA69; PA75; PA81; PB12; PC06; PC20; PC41; PC52; PC57; PC74;
M[20P]-764: PA11; PA13; PA22; PA29; PA35; PA40; PA57; PA73; PA75; PA81; PB08; PB12; PB33; PB82; PC11; PC20; PC21; PC29; PC41; PC52;
- M[20P]-765: PA10; PA11; PA13; PA15; PA26; PA27; PA30; PA35; PA40; PA89; PB12; PB22; PB58; PB60; PC34; PC37; PC55; PC58; PC67; PD02;
- 30 M[20P]-766: PA11; PA13; PA17; PA22; PA24; PA25; PA29; PA30; PA33; PA34; PA35; PA81; PA92; PB00; PB12; PB82; PC24; PC50; PC74; PC97;
M[20P]-767: PA11; PA21; PA22; PA28; PA29; PA33; PA37; PB12; PB60; PB85; PB88; PB99; PC14; PC20; PC21; PC41; PC51; PC55; PC57; PC98;
- 35 M[20P]-771: PA10; PA11; PA13; PA14; PA22; PA26; PA27; PA28; PA29; PA35; PA86; PA89; PB07; PB12; PB28; PB88; PB99; PC12; PC21; PC50;
M[20P]-777: PA11; PA14; PA15; PA20; PA21; PA24; PA25; PA28; PA29; PA33; PA75; PB12; PB85; PB99; PC21; PC51; PC55; PC62; PC67; PD02;
- M[20P]-779: PA15; PA19; PA20; PA21; PA22; PA23; PA24; PA25; PA29; PA30; PA57; PA67; PA75; PA89; PB07; PB12; PB32; PC20; PC51; PC74;
- 40 M[20P]-781: PA10; PA15; PA21; PA22; PA39; PA57; PA75; PB08; PB09; PB12; PB28; PB35; PB58; PB60; PC12; PC33; PC37; PC66; PC67; PC74;
M[20P]-782: PA10; PA11; PA12; PA13; PA14; PA20; PA26; PA30; PA75; PA89; PA92; PB05; PB07; PB12; PB32; PB82; PB99; PC51; PC58; PC73;
- 45 M[20P]-785: PA16; PA17; PA21; PA28; PA29; PA30; PA32; PA33; PA55; PB07; PB12; PB77; PB85; PB88; PB99; PC12; PC20; PC35; PC67; PC98;
M[20P]-786: PA11; PA13; PA14; PA18; PA19; PA21; PA22; PA27; PA28; PA29; PA33; PA40; PA52; PA57; PA81; PB28; PC11; PC20; PC21; PC50;

- M[20P]-787: PA10; PA11; PA13; PA22; PA23; PA35; PB07; PB12; PB22; PB85; PB99; PC11; PC20; PC41; PC49; PC52; PC53; PC55; PC73; PD02;
- M[20P]-789: PA10; PA11; PA13; PA22; PA23; PA27; PA40; PA57; PA75; PA81; PB07; PB12; PB27; PB60; PB88; PC11; PC20; PC23; PC35; PC41;
- 5 M[20P]-791: PA10; PA11; PA13; PA15; PA21; PA33; PA35; PA40; PA57; PA69; PA85; PA98; PA99; PB07; PB12; PB20; PB99; PC12; PC20; PC58;
- M[20P]-793: PA11; PA13; PA14; PA15; PA19; PA26; PA29; PA55; PA67; PA89; PB12; PB33; PB88; PC14; PC20; PC21; PC34; PC37; PC67; PC97;
- 10 M[20P]-794: PA21; PA23; PA28; PA29; PA34; PA35; PA57; PA75; PA81; PB07; PB12; PB18; PB32; PB58; PC11; PC14; PC41; PC55; PC62; PC74;
- M[20P]-795: PA10; PA13; PA20; PA23; PA27; PA29; PA57; PA60; PA69; PB00; PB07; PB12; PB18; PB32; PB85; PB88; PC20; PC66; PC74; PD02;
- M[20P]-803: PA11; PA13; PA15; PA22; PA25; PA26; PA29; PA30; PA33; PA40; PA75; PB07; PB12; PB28; PC21; PC23; PC34; PC55; PC71; PC73;
- 15 M[20P]-804: PA11; PA12; PA13; PA20; PA22; PA23; PA30; PA33; PA34; PA79; PB12; PB18; PB58; PB60; PB85; PB88; PC20; PC50; PC67; PC68;
- M[20P]-806: PA06; PA10; PA11; PA13; PA19; PA20; PA21; PA23; PA25; PA28; PA52; PA73; PA75; PA98; PB12; PB33; PB82; PB85; PB88; PC11;
- M[20P]-816: PA11; PA13; PA19; PA25; PA28; PA29; PA33; PA57; PA81; PA89; PB00; PB10; PB12; PB26; PB28; PB99; PC11; PC12; PC76; PC97;
- 20 M[20P]-819: PA10; PA11; PA13; PA15; PA26; PA34; PA60; PA81; PA92; PB07; PB12; PB38; PB85; PC01; PC14; PC20; PC52; PC58; PC66; PC67;
- M[20P]-823: PA00; PA11; PA12; PA13; PA14; PA17; PA27; PA29; PA40; PA60; PA75; PA79; PA89; PB07; PB12; PB99; PC20; PC52; PC66; PC71;
- 25 M[20P]-830: PA13; PA20; PA25; PA26; PA28; PA40; PA75; PA89; PB00; PB07; PB12; PB32; PB88; PC06; PC12; PC20; PC41; PC52; PC76; PC98;
- M[20P]-831: PA11; PA15; PA21; PA25; PA33; PB12; PB28; PB32; PB33; PB38; PB60; PB77; PB88; PC20; PC23; PC35; PC37; PC49; PC58; PC66;
- M[20P]-841: PA10; PA11; PA13; PA18; PA22; PA28; PA29; PA30; PA35; PA40; PA55; PA75; PB07; PB12; PB28; PB32; PB33; PC41; PC49; PC60;
- 30 M[20P]-844: PA11; PA15; PA21; PA24; PA26; PA29; PA33; PA40; PA69; PA81; PA98; PB12; PB28; PB61; PC49; PC67; PC71; PC73; PC74; PD02;
- M[20P]-845: PA11; PA13; PA14; PA15; PA21; PA30; PA34; PA39; PA60; PA79; PA81; PA89; PB07; PB18; PC14; PC20; PC21; PC34; PC68; PC97;
- 35 M[20P]-846: PA11; PA13; PA25; PA27; PA29; PA32; PA33; PA35; PA37; PA39; PA40; PB08; PB12; PB33; PB88; PC24; PC41; PC52; PC58; PC62;
- M[20P]-847: PA10; PA19; PA20; PA21; PA28; PA29; PA30; PA67; PA75; PA89; PB00; PB12; PB28; PB33; PB60; PC12; PC55; PC73; PC74; PD02;
- M[20P]-848: PA07; PA11; PA13; PA15; PA20; PA22; PA26; PA27; PA30; PA40; PA69; PA75; PB12; PB32; PB85; PB88; PC34; PC37; PC66; PC68;
- 40 M[20P]-849: PA00; PA13; PA25; PA27; PA28; PA29; PA34; PA40; PA57; PA81; PB00; PB12; PB18; PB77; PC20; PC24; PC35; PC41; PC50; PC51;
- M[20P]-852: PA00; PA11; PA13; PA15; PA19; PA22; PA26; PA35; PA89; PB00; PB12; PB58; PB77; PC12; PC20; PC29; PC62; PC71; PC94; PC98;
- 45 M[20P]-853: PA00; PA11; PA21; PA24; PA25; PA33; PA57; PA60; PB07; PB12; PB28; PB32; PB60; PB82; PB88; PC24; PC41; PC50; PC51; PC58;
- M[20P]-857: PA00; PA11; PA13; PA26; PA28; PA75; PA76; PA79; PB12; PB28; PB60; PB88; PC11; PC12; PC20; PC37; PC50; PC52; PC67; PD02;

M[20P]-873: PA10; PA13; PA26; PA27; PA30; PA34; PA85; PA98; PB05; PB07; PB12; PB28; PB29; PB32; PB60; PC14; PC20; PC26; PC41; PC76;

M[20P]-880: PA11; PA13; PA17; PA20; PA22; PA24; PA26; PA27; PA29; PA38; PA57; PB07; PB12; PB20; PB29; PB97; PC24; PC37; PC51; PC71;

5 M[20P]-888: PA10; PA12; PA15; PA16; PA21; PA27; PA28; PA33; PA44; PB00; PB12; PB13; PB27; PC11; PC20; PC51; PC58; PC73; PC75; PD02;

M[20P]-890: PA00; PA11; PA13; PA15; PA16; PA25; PA34; PA75; PA82; PA86; PB00; PB12; PB16; PB27; PB33; PC20; PC49; PC50; PC57; PC98;

10 M[20P]-894: PA12; PA18; PA21; PA24; PA26; PA27; PA34; PA57; PA67; PA82; PB01; PB12; PB33; PB76; PB98; PB99; PC11; PC24; PC33; PC83;

M[20P]-902: PA12; PA13; PA21; PA26; PA29; PA39; PA40; PA57; PA96; PB00; PB07; PB12; PB27; PC20; PC50; PC53; PC61; PC62; PC75; PC94;

M[20P]-904: PA06; PA13; PA20; PA23; PA25; PA27; PA35; PA46; PA52; PA60; PA81; PA90; PA97; PB03; PB12; PB57; PB99; PC20; PC52; PC73;

15 M[20P]-907: PA00; PA07; PA14; PA21; PA24; PA25; PA45; PA75; PA83; PA97; PB12; PB26; PB32; PB60; PB92; PC06; PC20; PC35; PC37; PD02;

M[20P]-909: PA12; PA14; PA20; PA21; PA29; PA39; PA72; PA93; PB12; PB22; PB28; PB32; PC21; PC23; PC24; PC41; PC52; PC73; PC79; PD02.

20 Note: if in the previous and in the following analogous Lists a plant code occurs more than one time in the same herbal composition, it means that more than one calibrated spoon of the corresponding grinded dried plant was used.

Starting from the herbal compositions listed in List 6-3 corresponding water extracts (or processed samples) were prepared according to the previously described Method B using 40mL of water containing 100g/l of sucrose for water extraction and including the decantation step before filtration.

25 The biological activity of each processed sample was determined according to the previously described Method C by using the following *Staphylococcus aureus* strains: ATCC 25923, ATCC 49476, ATCC 6538, ATCC 51740, ATCC 29213 and ATCC 14775, with the following final dilutions of processed samples: 1:20, 1:63 and 1:200.

30 The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from herbal compositions including the 2 plants of Mix[2P]-1 at 1:20 dilution is given in Table 5-1.

Table 5-1: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[2P]-1 at 1:20 dilution. (A) Mixes extracts showing higher antimicrobial and antibiofilm activity. (B) Mixes extracts showing antibiofilm activity and equivalent or lower antimicrobial activity. (C) Mixes extracts showing antimicrobial activity and equivalent or lower antibiofilm activity

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
M[20P]-563	26%	40%
M[20P]-495	25%	43%
M[20P]-631	27%	42%
M[20P]-465	24%	50%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
M[20P]-723	11%	40%
M[20P]-786	20%	40%
M[20P]-737	22%	40%
M[20P]-340	11%	40%

M[20P]-408	26%	50%
M[09P]-8	31%	48%
M[19P]-13	31%	50%
M[18P]-12	40%	50%
M[20P]-581	40%	50%
M[20P]-469	40%	55%
M[20P]-485	43%	52%
M[20P]-427	37%	66%
M[20P]-582	51%	54%
M[20P]-431	44%	65%
M[20P]-618	54%	57%
M[20P]-580	61%	59%
M[20P]-421	53%	70%
M[20P]-499	60%	67%
M[20P]-525	86%	77%
M[20P]-29	105%	93%

M[20P]-382	13%	41%
M[20P]-444	18%	42%
M[20P]-603	12%	42%
M[20P]-470	19%	42%
M[20P]-503	12%	46%
M[19P]-67	18%	49%
M[20P]-533	21%	50%
M[20P]-845	16%	51%
M[20P]-377	21%	51%
M[20P]-510	7%	55%

(C)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
M[17P]-9	27%	29%
M[18P]-37	29%	36%

20 processed samples obtained from herbal compositions containing 9, 18, 19 or 20 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[2P]-1 (Table 5-1 A). Fourteen 20 plant mixes extracts were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 5-1 B). Two processed samples obtained from herbal compositions containing 17 and 18 plants were found to have higher antimicrobial activity and equivalent or lower antibiofilm activity (Table 5-1 C).

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from herbal compositions including the 2 plants of Mix[2P]-2 at 1:20 dilution is given in Table 5-2.

Table 5-2: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[2P]-2 at 1:20 dilution. (A) Mixes extracts showing higher antimicrobial and antibiofilm activity. (B) Mixes extracts showing antimicrobial activity and equivalent or lower antibiofilm activity.

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-2	12%	-3%
M[17P]-5	17%	17%
M[20P]-654	16%	20%
M[16P]-1	14%	22%
M[17P]-1	16%	22%
M[20P]-779	14%	29%
M[20P]-746	15%	28%
M[20P]-635	17%	26%
M[20P]-844	15%	29%
M[20P]-831	17%	31%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-2	12%	-3%
M[20P]-638	11%	13%
M[20P]-794	6%	16%
M[20P]-909	0%	19%
M[20P]-538	8%	22%
M[20P]-781	6%	24%
M[19P]-27	7%	26%
M[20P]-681	11%	24%
M[16P]-3	8%	28%
M[20P]-721	2%	28%

M[20P]-894	18%	30%
M[20P]-888	23%	25%
M[20P]-407	19%	31%
M[20P]-907	21%	31%
M[20P]-335	19%	33%
M[20P]-714	17%	35%
M[20P]-668	15%	38%
M[20P]-584	14%	39%
M[20P]-777	16%	37%
M[20P]-853	20%	34%
M[20P]-785	16%	39%
M[20P]-622	21%	34%
M[09P]-2	17%	39%
M[19P]-46	18%	37%
M[20P]-387	13%	43%
M[20P]-767	18%	39%
M[18P]-23	24%	33%
M[20P]-745	20%	37%
M[20P]-478	22%	36%
M[19P]-45	16%	42%
M[19P]-49	19%	40%
M[20P]-467	17%	42%
M[20P]-396	21%	38%
M[20P]-392	17%	42%
M[19P]-44	20%	42%
M[19P]-41	23%	40%
M[19P]-48	18%	46%
M[19P]-47	17%	46%
M[20P]-490	20%	44%
M[19P]-21	29%	35%
M[20P]-632	24%	40%
M[19P]-54	19%	45%
M[20P]-461	19%	46%
M[19P]-42	24%	41%
M[20P]-506	17%	49%
M[19P]-55	21%	46%
M[20P]-459	26%	42%
M[20P]-375	19%	52%
M[19P]-2	26%	49%
M[19P]-40	28%	48%
M[20P]-523	28%	51%
M[20P]-420	34%	56%
M[20P]-429	36%	59%
M[20P]-535	45%	55%
M[20P]-532	42%	60%

M[20P]-493	7%	28%
M[17P]-6	9%	29%
M[20P]-659	10%	30%
M[20P]-409	10%	30%
M[20P]-602	4%	31%
M[20P]-456	9%	31%
M[20P]-383	5%	32%
M[20P]-847	13%	33%
M[20P]-496	6%	34%
M[19P]-51	10%	34%
M[20P]-371	7%	36%
M[20P]-609	12%	36%
M[19P]-52	9%	36%
M[19P]-50	12%	36%
M[18P]-24	11%	36%
M[19P]-53	10%	40%

M[20P]-432	54%	72%
M[20P]-555	68%	65%
M[20P]-240	67%	78%
M[20P]-557	86%	78%

58 processed samples obtained from herbal compositions containing 9, 16, 17, 18, 19 or 20 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[2P]-2 (Table 5-2 A). 25 processed samples obtained from herbal compositions containing 16, 17, 19 or 20 plants were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 5-2 B).

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from herbal compositions including the 2 plants of Mix[2P]-3 at 1:20 dilution is given in Table 5-3.

Table 5-3: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[2P]-3 at 1:20 dilution. (A) Mixes extracts showing higher antimicrobial and antibiofilm activity. (B) Mixes extracts showing antimicrobial activity and equivalent or lower antibiofilm activity.

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-3	12%	28%
M[20P]-649	16%	29%
M[20P]-857	15%	32%
M[20P]-703	13%	36%
M[20P]-486	15%	35%
M[20P]-766	13%	37%
M[20P]-789	14%	36%
M[20P]-725	14%	37%
M[20P]-536	20%	32%
M[20P]-753	14%	38%
M[20P]-401	14%	38%
M[20P]-765	14%	38%
M[20P]-848	14%	39%
M[20P]-782	14%	40%
M[20P]-589	15%	39%
M[20P]-607	14%	40%
M[20P]-541	20%	34%
M[20P]-447	15%	39%
M[20P]-601	17%	38%
M[20P]-494	13%	43%
M[20P]-787	15%	40%
M[20P]-748	15%	41%
M[09P]-6	15%	41%
M[20P]-480	18%	38%
M[20P]-633	15%	41%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-3	12%	28%
M[20P]-599	11%	32%
M[20P]-621	10%	32%
M[20P]-740	12%	33%
M[20P]-830	7%	33%
M[20P]-623	12%	34%
M[20P]-708	8%	35%
M[20P]-880	12%	35%
M[20P]-716	8%	35%
M[20P]-803	4%	36%
M[20P]-686	13%	36%
M[20P]-663	5%	36%
M[20P]-873	7%	36%
M[20P]-608	10%	37%
M[20P]-752	10%	37%
M[20P]-443	8%	37%
M[19P]-30	3%	37%
M[20P]-712	12%	37%
M[20P]-616	7%	38%
M[20P]-724	4%	38%
M[20P]-795	11%	38%
M[20P]-403	13%	38%
M[20P]-595	7%	38%
M[20P]-771	9%	39%
M[20P]-718	5%	39%

M[20P]-849	14%	42%
M[20P]-793	16%	42%
M[18P]-9	17%	41%
M[20P]-852	13%	45%
M[20P]-819	20%	39%
M[20P]-655	15%	44%
M[20P]-455	15%	44%
M[20P]-846	16%	44%
M[20P]-481	18%	41%
M[20P]-484	17%	42%
M[20P]-590	26%	34%
M[20P]-890	19%	42%
M[20P]-670	21%	40%
M[20P]-705	23%	39%
M[20P]-816	22%	39%
M[19P]-91	15%	48%
M[20P]-457	22%	43%
M[20P]-551	24%	41%
M[19P]-100	16%	50%
M[20P]-583	23%	43%
M[20P]-823	22%	44%
M[19P]-100	16%	50%
M[19P]-98	17%	50%
M[19P]-101	14%	54%
M[19P]-93	14%	53%
M[19P]-102	13%	54%
M[19P]-5	19%	49%
M[19P]-89	17%	51%
M[19P]-103	13%	55%
M[19P]-96	18%	52%
M[20P]-475	25%	45%
M[20P]-548	30%	42%
M[19P]-96	19%	53%
M[20P]-605	22%	50%
M[20P]-577	29%	44%
M[20P]-482	27%	46%
M[19P]-95	19%	55%
M[20P]-904	21%	55%
M[20P]-559	32%	47%
M[20P]-430	29%	71%
M[20P]-520	62%	67%
M[20P]-579	75%	66%
M[20P]-560	72%	70%
M[20P]-565	73%	73%
M[20P]-524	81%	79%

M[20P]-620	12%	39%
M[20P]-841	6%	40%
M[20P]-804	8%	40%
M[20P]-386	6%	41%
M[20P]-764	11%	42%
M[18P]-33	11%	42%
M[20P]-349	9%	43%
M[19P]-92	10%	46%
M[18P]-34	11%	47%
M[19P]-93	12%	50%
M[18P]-31	12%	51%
M[18P]-34	12%	51%
M[19P]-62	10%	56%

M[20P]-575	85%	76%
------------	-----	-----

70 processed samples obtained from herbal compositions containing 9, 19 or 20 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[2P]-3 (Table 5-3 A). 37 processed samples containing 18 to 20 plants were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 5-3 B).

- 5 The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from herbal compositions including the 3 plants of Mix[3P]-1 at 1:20 dilution is given in Table 5-4.

Table 5-4: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on Mix[3P]-1 at 1:20 dilution. (A) Mixes extracts showing higher antimicrobial and antibiofilm activity. (B) Mixes extracts showing antimicrobial activity and equivalent or lower antibiofilm activity.

(A)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[3P]-1	31%	15%
M[18P]-2	35%	42%
M[16P]-5	36%	41%
M[14P]-2	34%	44%
M[16P]-6	35%	44%
M[09P]-9	33%	47%
M[09P]-1	34%	45%
M[18P]-7	35%	45%
M[17P]-3	36%	44%
M[09P]-7	35%	49%
M[20P]-594	40%	46%
M[09P]-6	37%	51%
M[20P]-658	45%	43%
M[19P]-3	34%	54%
M[18P]-11	40%	49%
M[19P]-104	43%	46%
M[09P]-3	46%	44%
M[09P]-4	46%	45%
M[19P]-19	34%	57%
M[18P]-13	42%	50%
M[18P]-10	43%	49%
M[18P]-9	42%	51%
M[18P]-17	45%	51%
M[20P]-634	43%	56%
M[20P]-596	44%	55%
M[18P]-15	47%	52%
M[20P]-449	47%	56%
M[09P]-10	54%	50%
M[20P]-588	48%	56%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
Mix[3P]-1	31%	15%
M[20P]-692	11%	22%
M[20P]-612	6%	25%
M[18P]-22	4%	27%
M[20P]-388	9%	27%
M[20P]-380	12%	28%
M[11P]-12	22%	29%
M[19P]-33	8%	30%
M[19P]-35	6%	30%
M[17P]-2	12%	30%
M[20P]-678	10%	31%
M[19P]-22	17%	31%
M[20P]-552	16%	32%
M[17P]-4	9%	32%
M[20P]-806	12%	32%
M[12P]-1	19%	33%
M[14P]-1	18%	33%
M[19P]-28	6%	33%
M[12P]-2	22%	34%
M[18P]-1	16%	34%
M[19P]-36	6%	34%
M[20P]-479	23%	34%
M[15P]-1	23%	34%
M[09P]-21	22%	34%
M[20P]-446	10%	34%
M[19P]-34	7%	34%
M[19P]-106	29%	35%
M[10P]-41	8%	35%
M[13P]-1	22%	35%

M[20P]-458	50%	56%
M[19P]-105	49%	56%
M[13P]-2	51%	56%
M[19P]-11	43%	67%
M[18P]-16	54%	59%
M[20P]-569	55%	60%
M[20P]-296	52%	62%
M[18P]-18	56%	61%
M[20P]-516	61%	68%
M[20P]-592	70%	71%
M[20P]-626	71%	71%
M[19P]-23	72%	72%
M[20P]-497	72%	77%
M[20P]-572	74%	78%
M[20P]-574	76%	79%
M[20P]-527	76%	80%
M[20P]-297	83%	81%
M[20P]-422	81%	83%
M[20P]-564	86%	82%
M[20P]-418	98%	87%
M[20P]-416	99%	89%
M[20P]-576	104%	86%
M[20P]-573	104%	93%

M[18P]-5	18%	35%
M[19P]-39	1%	35%
M[11P]-3	12%	35%
M[20P]-439	19%	36%
M[11P]-2	13%	36%
M[18P]-20	6%	36%
M[20P]-757	18%	36%
M[10P]-303	25%	36%
M[11P]-11	29%	37%
M[11P]-6	9%	37%
M[19P]-37	5%	37%
M[18P]-19	10%	37%
M[18P]-4	29%	37%
M[18P]-35	27%	37%
M[18P]-25	11%	37%
M[11P]-5	10%	38%
M[20P]-694	20%	38%
M[11P]-8	10%	38%
M[11P]-9	11%	38%
M[20P]-733	12%	38%
M[11P]-7	10%	38%
M[19P]-31	6%	38%
M[19P]-32	4%	38%
M[19P]-29	6%	38%
M[18P]-6	22%	38%
M[20P]-374	2%	38%
M[11P]-4	12%	39%
M[20P]-791	13%	39%
M[20P]-758	15%	39%
M[19P]-26	6%	39%
M[20P]-697	17%	39%
M[19P]-38	4%	40%
M[19P]-24	28%	40%
M[16P]-2	24%	40%
M[18P]-21	9%	40%
M[15P]-5	29%	40%
M[17P]-8	30%	40%
M[20P]-613	8%	41%
M[18P]-14	27%	41%
M[20P]-604	14%	41%
M[11P]-1	17%	41%
M[18P]-36	29%	41%
M[18P]-3	23%	42%
M[20P]-453	5%	42%
M[19P]-25	9%	42%

M[20P]-709	18%	42%
M[20P]-474	16%	45%
M[20P]-666	22%	45%
M[20P]-902	21%	46%
M[20P]-435	15%	46%
M[20P]-567	30%	47%
M[19P]-4	24%	47%
M[19P]-16	18%	47%
M[19P]-59	10%	48%
M[19P]-17	23%	48%
M[19P]-58	20%	49%
M[19P]-60	13%	49%
M[19P]-15	27%	50%
M[11P]-10	31%	50%
M[19P]-14	27%	50%
M[19P]-6	32%	51%
M[20P]-445	21%	51%
M[19P]-8	30%	51%
M[19P]-18	21%	52%
M[19P]-9	29%	52%
M[19P]-7	30%	52%
M[19P]-10	29%	52%
M[19P]-68	23%	53%
M[19P]-63	16%	53%
M[19P]-69	22%	53%
M[19P]-1	26%	54%
M[19P]-66	24%	54%
M[19P]-12	26%	54%
M[20P]-472	27%	54%
M[19P]-20	31%	54%
M[19P]-70	19%	55%
M[19P]-62	17%	56%
M[19P]-57	21%	56%
M[19P]-65	27%	57%
M[19P]-73	18%	57%
M[20P]-514	30%	58%
M[19P]-56	24%	58%
M[19P]-64	27%	59%
M[19P]-72	20%	60%
M[19P]-71	19%	61%

51 processed samples obtained from herbal compositions containing 9, 16, 17, 18, 19 or 20 plants were found to have higher antibiofilm and antimicrobial activity than the reference mix extract Mix[3P]-1 (Table 5-4 A). 113 processed samples containing 9 to 20 plants were found to have higher antibiofilm activity and equivalent or lower antimicrobial activity (Table 5-4 B).

As a comparative Example, we report several 20 plant mixes extracts containing only one plant among the 3 plants of Example 1 (PA13, PA21, PB12), which have been found to have none or poor antimicrobial and antibiofilm activities against *Staphylococcus aureus*.

Ten herbal compositions having the content reported in List 5-8 were prepared according to the previously described Method A.

List 5-8: herbal compositions of processed samples

M[20P]-39: PB00; PB01; PB02; PB03; PB04; PB05; PB06; PB07; PB08; PB09; PB10; PB11; PB12; PB13; PB14; PB15; PB16; PB17; PB18; PB19;

M[20P]-239: PA00; PA05; PA21; PA23; PA26; PA33; PA39; PA60; PA85; PB00; PB06; PB32; PB79; PC02; PC26; PC34; PC38; PC62; PC78; PC84;

M[20P]-337: PA00; PA10; PA11; PA15; PA18; PA21; PA22; PA24; PA26; PA28; PA52; PA94; PA98; PB10; PB20; PB26; PB29; PB82; PC67; PC97;

M[20P]-362: PA21; PA23; PA25; PA33; PA37; PA57; PA69; PA81; PA92; PB16; PB25; PB26; PB28; PB29; PB78; PC20; PC21; PC71; PC83; PD02;

M[20P]-364: PA15; PA21; PA25; PA28; PA39; PA75; PA80; PA85; PA89; PA98; PB01; PB06; PB24; PB28; PB35; PC24; PC25; PC57; PC75; PC82;

M[20P]-540: PA06; PA10; PA12; PA14; PA20; PA21; PA26; PA27; PA27; PA57; PA57; PA57; PA69; PB07; PB82; PC11; PC20; PC23; PC55; PC98;

M[20P]-647: PA10; PA11; PA24; PA26; PA28; PA33; PA35; PA40; PA57; PA69; PB00; PB07; PB12; PB28; PB33; PB38; PB60; PC55; PC67; PC97;

M[20P]-652: PA00; PA07; PA13; PA15; PA16; PA23; PA26; PA28; PA33; PA39; PA57; PA92; PB28; PB38; PC20; PC21; PC52; PC55; PC74; PC97;

M[20P]-862: PA11; PA15; PA18; PA21; PA25; PA27; PA28; PA29; PA35; PA38; PA52; PA86; PA89; PA92; PB07; PB28; PB88; PC11; PC55; PC98;

M[20P]-872: PA13; PA14; PA16; PA28; PA29; PA30; PA33; PA35; PA39; PA81; PA90; PA92; PB00; PB32; PC12; PC34; PC37; PC67; PC97; PD02.

Ten water extracts (or processed samples) were prepared from the corresponding herbal compositions according to the previously described Method B using 40mL of water containing 100g/l of sucrose for water extraction and including the decantation step before filtration.

For each processed sample the biological activity was determined according to the previously described Method C, with the following final dilutions of processed samples: 1:20, 1:63 and 1:200.

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples at 1:20 dilution is given in Table 5-5.

Table 5-5: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed samples at 1:20 dilution.

	Planktonic growth inhibition	Biofilm formation inhibition
Mix[2P]-1	22%	36%
Mix[2P]-2	12%	-3%
Mix[2P]-3	12%	28%
M[20P]-362	-1%	-6%

M[20P]-364	-5%	6%
M[20P]-039	1%	6%
M[20P]-862	5%	2%
M[20P]-337	0%	9%
M[20P]-872	2%	11%
M[20P]-540	5%	10%
M[20P]-652	9%	7%
M[20P]-239	9%	7%
M[20P]-647	9%	9%

All tested processed samples obtained from herbal compositions containing 20 plants including only one plant among the 3 plants of Example 1 (PA13, PA21, PB12) showed lower antimicrobial activity than the three 2 plant mixes extracts of reference Mix[2P]-1 (obtained from PA13 and PA21), Mix[2P]-2 (obtained from PA21 and PB12) and Mix[2P]-3 (obtained from PA13 and PB12) and lower antibiofilm activity than Mix[2P]-1 and Mix[2P]-3.

Example 6

In this Example, we report the simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Staphylococcus pseudintermedius* of 2 processed samples chosen among the best mixes described in Example 4 and 5.

- 10 The dried plants of pharmaceutical grade were obtained from “Pharmacie Fontgèze” (Clermont-Ferrand, France) and “Pharmacie St Herem” (Clermont-Ferrand, France).

Two herbal preparations containing the dried plant powders reported in the Table 6-1 below, were prepared according to the previously described Method A.

Table 6-1

Herbal compositions	M[20P]-576	M[10P]-154
Plant 1	2 spoons of PA11	1 spoon of PA11
Plant 2	1 spoon of PA12	1 spoon of PA12
Plant 3	3 spoons of PA13	1 spoon of PA13
Plant 4	2 spoons of PA14	1 spoon of PA21
Plant 5	1 spoon of PA20	1 spoon of PA22
Plant 6	1 spoon of PA21	1 spoon of PA79
Plant 7	1 spoon of PA22	1 spoon of PB12
Plant 8	1 spoon of PA29	1 spoon of PB49
Plant 9	1 spoon of PA39	1 spoon of PC29
Plant 10	1 spoon of PB12	1 spoon of PC63
Plant 11	1 spoon of PB60	
Plant 12	1 spoon of PC20	
Plant 13	1 spoon of PC26	
Plant 14	1 spoon of PC33	
Plant 15	1 spoon of PC37	
Plant 16	1 spoon of PC49	

Starting from such herbal compositions 2 corresponding water extracts (or processed samples) were prepared according to the previously described Method B using 40mL for M[20P]-576 and 20mL for M[10P]-154 of water containing 100g/l of sucrose for water extraction and including the centrifugation step before filtration.

5 The biological activity of the processed samples was determined according to the previously described Method C by using the following strains:

- *Staphylococcus aureus* strains: ATCC 25923, ATCC 49476, ATCC 6538, ATCC 51740, ATCC 29213, ATCC 14775, NCTC 12493, ATCC 33591, ATCC 33592, ATCC 43300, ATCC 700698, ATCC 700699, ATCC 9144 and ATCC BAA-44
- 10 - *Staphylococcus epidermidis* strains: ATCC 14990, ATCC 12228, ATCC 700296, ATCC 49134 and ATCC 49461
- *Staphylococcus pseudintermedius* strain: ATCC 49444

The minimum inhibitory concentration (MIC) observed with the 2 processed samples is given in Table 6-2 in the form of HID, as well as the average efficacy over all the strains of *Staphylococcus aureus* and
15 *Staphylococcus epidermidis*.

Table 6-2: HID of the 2 processed samples on 14 strains of *Staphylococcus aureus*, 5 strains of *Staphylococcus epidermidis* and 1 strain of *Staphylococcus pseudintermedius*

		M[10P]-154	M[20P]-576
<i>S. aureus</i>	ATCC 25923	20	20
	ATCC 49476	40	40
	ATCC 6538	10	10
	ATCC 51740	20	40
	ATCC 29213	20	20
	ATCC 14775	20	20
	NCTC 12493	40	40
	ATCC 33591	20	40
	ATCC 33592	20	40
	ATCC 43300	20	40
	ATCC 700698	20	20
	ATCC 700699	40	20
	ATCC 9144	20	20
	ATCC BAA-44	40	20
	Average	25	28
<i>S. epidermidis</i>	ATCC 14990	40	20
	ATCC 12228	40	20
	ATCC 700296	40	20
	ATCC 49461	40	20
	ATCC 49134	80	80
	Average	48	32
<i>S. pseudintermedius</i>	ATCC 49444	40	40
Total average		32	30

M[10P]-154 and M[20P]-576 showed full antimicrobial activity on all tested *Staphylococcus aureus* and *Staphylococcus epidermidis*. M[10P]-154 showed full antimicrobial activity at 1:80 on 1 *Staphylococcus epidermidis* strain, at 1:40 on 4 *Staphylococcus aureus* strains, 4 *Staphylococcus epidermidis* and the tested strain of *Staphylococcus pseudintermedius*, at 1:20 on 9 *Staphylococcus aureus* strains, and at 1:10 on 1 *Staphylococcus aureus* strain.

M[20P]-576 showed full antimicrobial activity at 1:80 on 1 *Staphylococcus epidermidis* strain, at 1:40 on 6 *Staphylococcus aureus* strains and the tested strain of *Staphylococcus pseudintermedius*, at 1:20 on 7 *Staphylococcus aureus* strains and 4 *Staphylococcus epidermidis* and at 1:10 on 1 *Staphylococcus aureus* strain.

The bacteria planktonic growth inhibition percentage and the biofilm formation inhibition (IBF) percentage of the 2 processed samples at 1:20 dilution is given in Table 6-3, as well as the average efficacy over all the strains of *Staphylococcus aureus* and *Staphylococcus epidermidis*.

Table 6-3: planktonic growth inhibition and biofilm formation inhibition (IBF) of the 2 processed samples on 14 strains of *Staphylococcus aureus*, 5 strains of *Staphylococcus epidermidis* and 1 strain of *Staphylococcus pseudintermedius* at 1:20 dilution

		planktonic growth inhibition		biofilm formation inhibition (IBF)	
		M[10P]-154	M[20P]-576	M[10P]-154	M[20P]-576
<i>S. aureus</i>	ATCC 25923	96%	99%	90%	88%
	ATCC 49476	93%	109%	93%	95%
	ATCC 6538	49%	56%	45%	60%
	ATCC 51740	91%	106%	89%	96%
	ATCC 29213	94%	105%	91%	92%
	ATCC 14775	90%	105%	88%	90%
	NCTC 12493	99%	113%	93%	92%
	ATCC 33591	95%	113%	91%	93%
	ATCC 33592	97%	107%	91%	93%
	ATCC 43300	94%	110%	90%	92%
	ATCC 700698	92%	113%	89%	91%
	ATCC 700699	88%	95%	98%	96%
	ATCC 9144	82%	91%	88%	84%
	ATCC BAA-44	90%	101%	93%	94%
	Average	89%	102%	88%	90%
<i>S. epidermidis</i>	ATCC 14990	90%	101%	91%	90%
	ATCC 12228	90%	101%	93%	91%
	ATCC 700296	86%	90%	81%	82%
	ATCC 49461	90%	99%	93%	85%
	ATCC 49134	89%	96%	96%	90%
	Average	89%	98%	91%	88%
<i>S. pseudintermedius</i>	ATCC 49444	92%	103%	95%	93%
Total average		89%	101%	89%	89%

At 1:20 dilution, M[10P]-154 showed an average of 88% of biofilm formation inhibition on all tested *Staphylococcus aureus* strains whereas M[20P]-576 showed an average of 90%. The two mixes showed lower antimicrobial activity on one strain: ATCC 6538 (45% for M[10P]-154, 60% for M[20P]-576). M[10P]-154 showed an average of 91% of planktonic growth inhibition on all tested *Staphylococcus epidermidis* strains whereas M[20P]-576 showed an average of 88%. (cf Table 6-3). The two mixes showed more respectively 95% (M[10P]-154) and 93% (M[20P]-576) of inhibition percentage on the *Staphylococcus pseudintermedius* tested. Thus, M[10P]-154 and M[20P]-576 have demonstrated antibiofilm activity on *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Staphylococcus pseudintermedius* at 1:20 dilution.

Example 7

Solution of Mix[20C]-576 (described in Example 6) used for the following formulations was prepared according to the previously described Method A and B with the exception of the filtering step, which is in this Example performed by suction filtration with a Büchner funnel filled with 5 cm thick hydrophilic cotton layer. For clarity, this cotton filtered solution will be referred as SK576 in the following Example.

An ointment and two creams to be used for topical applications were prepared with an emulsification method as follows:

- Phase A and Phase B were prepared by mixing with orbital shaking in two separate beakers the following ingredients:

	Phase A	Phase B
Ointment	- 73.8g of organic almond oil (Hyteck, "Aroma Zone™", France) - 18.5g of organic beeswax (Hyteck, "Aroma Zone™", France) - 9.7g of organic shea butter (Hyteck, "Aroma Zone™", France) - 1.0 mL of natural tocopherol (Hyteck, "Aroma Zone™", France)	- 29.6g of SK576, prepared as disclosed in Example 6 - 13.5g of organic glycerin (The Naissance Trading & Innovation, "Naissance™", UK) - 2.7g of Ascorbic acid (Hyteck "Aroma Zone™", France)
Cream 1	- 30.4g of organic almond oil (Hyteck, "Aroma Zone™", France): - 8.3g of organic Olivem® 1000 (Hyteck, "Aroma Zone™", France) - 3.1g of menthol (Hyteck "Aroma Zone™", France) - 1.0mL of natural tocopherol (Hyteck, "Aroma Zone™", France)	- 51.5g of SK576, prepared as disclosed in Example 6 - 14.9g of organic glycerin (The Naissance Trading & Innovation "Naissance™", UK)
Cream 2	- 14.9g of organic almond oil (Hyteck "Aroma Zone™", France): - 11.8g of organic Olivem® 1000 (Hyteck "Aroma Zone™", France) - 5.1g of organic shea butter (Hyteck "Aroma Zone™", France) - 1.0mL of natural tocopherol (Hyteck "Aroma Zone™", France)	- 160.0g of SK576, prepared as disclosed in Example 6 - 10g of organic glycerin (The Naissance Trading & Innovation, "Naissance™", UK)

- The two beakers were heated to 70 °C using a warm water bath and maintained at 70°C during 10 min to avoid the crystallization of beeswax.
- Phase B was then slowly poured into Phase A, keeping the beaker in the warm water bath and with constant stirring by a mechanical immersion blender (Moulinex "Infinyforce_hand blender", France).
- The beaker containing the mix of phase A and B was then removed from the warm bath and the ointment was formed by using an immersion "mayonnaise blender" (Moulinex "Infinyforce_hand blender", France).

A gel to be used for topical applications was prepared by

- Filtrating 20.6g of SK576, on an hydrophilic syringe filter (0.22 µm).
- Adding 0.75 g of Xanthan powder (Hyteck "Aroma Zone™", France) and 0.25g of ascorbic acid (Hyteck "Aroma Zone™", France) to the filtered solution and mix with a spatula for 5-10 min such as to form.

A spray solution to be used for topical applications was prepared by:

- Mixing 5.5g of SK576 with 11.9g of absolute ethanol(Sigma-Aldrich, France) in a 20 mL beaker which were stored in a 5 °C refrigerator for 3 H.
- Placing the beaker with the cold mixture in an ice-water bath and gently adding 0.7 g of Kolliphor® P-407 (Sigma-Aldrich, France) under agitation with a magnetic bar and stirrer.
- Following 30 min agitation under ice-water bathing, the solution was filled into a reagent sprayer (CAMAG, Switzerland) for administration.

The protocol of each of these five preparations can be easily scaled to different volumes of preparation by keeping the same relative amount of each ingredient.

Example 8

Biological activity of a water extract of the invention in a mouse model of lethal *Staphylococcus aureus* infection

Introduction

S. aureus strain Newman was isolated in 1952 from a human infection and has been used extensively in animal models of staphylococcal infections due to its robust virulence phenotype.

Female OF1 mice have been extensively used to study the pathogenesis induced by *Staphylococcus aureus*. Early work on the pathogenesis of *Staphylococcus aureus* infections used the intraperitoneal route of challenge to infect and treat mice because of its technical ease.

The aim of this Example was to evaluate the antimicrobial activity of a water extract of the invention in female OF1 mice infected with lethal dose of Newman *Staphylococcus aureus*. The treatment was administered by oral route according to the schedule Q1Dx10, 2 days before infection to 7 days after infection. Monitoring of mortality was performed from D0 to D15.

Vehicle was used as negative control.

Material and Methods

Test substance and controls

Test Substance: water extract M[20C] 576 (obtained starting from 120 g/l of herbal composition, prepared as described in Example 6). It was provided in 11 vials containing 1,200 µl each of water extracts. It was stored at -20°C.

Vehicle: sucrose 10% water solution. It will be stored at -20°C.

- 5 *Bacterial strain:* Newman *Staphylococcus* strain (provided by Dr T. Foster, Trinity College, Dublin, Ireland, Batch No. 14156, LST4).

Animal purchasing and caging

Twenty (20) Crl:OF1 (outbred) female mice, 6 weeks old were purchased from Charles River (L'Arbresle, France).

- 10 The animal care unit was authorized by the French Ministries of Agriculture and Research (Agreement Number B35-288-1). Animal experiments were performed according to ethical guidelines of animal experimentations (Principes d'éthique de l'expérimentation animale, Directive n°86/609 CEE du 24 Nov. 1986, Décret n°87/848 du 19 Oct. 1987, Arrêté d'Application du 19 Avril 1988).

- 15 The animals were maintained in "A2" rooms under controlled conditions of temperature ($22 \pm 3^\circ\text{C}$), humidity ($50 \pm 20\%$), photoperiod (12h light/12h dark), air exchange and low pressure. The air handling system is programmed for 14 air changes an hour, with no recirculation.

Food and water were provided *ad libitum*, being placed in the metal lid on top of the cage.

Treatment and experimental design

Preparation of bacteria - Amplification of bacteria for challenge

- 20 One vial of bacteria was defrosted and 100 µl of inoculum was cultured in 50 ml of Tryptic Soy Broth (TSB, Ref. 43592 or 22902, Sigma). Bacteria were incubated at 37°C during 20h under 110 rpm agitation on horizontal shaker. Bacterial suspension will be diluted at 1/20 (100 µl + 1.9 ml of TSB) and optical density will be read at 680 nm (O.D.680 nm).

- 25 Then, 10 ml of bacteria was centrifuged at 5,200 rpm (3400 g) during 10 min at 4°C. Supernatant was discarded and bacteria was solubilized in 10 ml of DPBS (Ref. BE17-512F, Lonza, France). Bacterial suspension (C0) was diluted at 1/20 (100 µl + 1.9 ml of DPBS) and O.D.680 nm will be measured.

The O.D.680 nm measured before and after centrifugation must be similar.

The concentration of CFU/ml of the bacteria solution was calculated as follow:

$$\text{CFU/ml} = (\text{O.D.680 nm} \times 20 \text{ (dilution factor)}) \times 6.5.108 \text{ (CFU/U.D.O)}$$

- 30 *Mucin preparation*

Mucin (Sigma, France) was prepared at 20% 14-16 days before treatment as described below:

- 35 DPBS (Dulbecco's Phosphate-Buffered Saline) was pre-warmed at 56°C and $\frac{3}{4}$ of the final volume was transferred in a mixer, then hog mucin will be added and mixed. The mixer was washed with the remaining volume of DPBS. Then, the 20% mucin solution was transferred in an Erlenmeyer Flask and was incubated at 37°C under agitation during 1 hour and 30 minutes. The $\frac{3}{4}$ of 20% mucin solution was transferred in a sterile bottle. The remainder $\frac{1}{4}$ of solution was stocked separately. Then the $\frac{3}{4}$ and the $\frac{1}{4}$ solutions was sterilized by autoclaving. Sterile 20% hog mucin solution was stored at +4°C.

The day before the infection, the pH of $\frac{3}{4}$ of 20% mucin solution was adjusted at 7.4 by adding filtered NaOH 30%. If necessary, a part of the $\frac{1}{4}$ remaining solution was used to adjust pH.

Bacteria preparation

According to O.D.680 nm results, bacteria (C0) were diluted in DPBS to reach the concentration of 6.8×10^7 CFU/ml. These suspensions were then diluted at $\frac{1}{2}$ in 20% hog mucin to get the LD80 concentration of 3.4×10^7 CFU/ml in PBS-10% hog mucin. The suspension used for treatment was put in tube containing sterile glass beads (4 mm, Ref. 068502, Dutscher, France). Suspension were not vortexed, but mixed by inverting the tubes. Bacteria were placed on ice before treatments and kept on ice during treatment.

10 *CFU determination*

To determine the exact concentration of the bacterial suspension before and after inoculated into the animals, the bacterial suspension was diluted in 10-fold serial dilutions in DPBS-0.5% Tween 20 (Sigma, France) as described in Table 8-1 in deep wells (50 μ l of bacteria + 450 μ l of DPBS-0.5% Tween 20). Then, 6 x 50 μ l of bacterial dilutions 3, 4, 5 and 6 were seeded on Tryptic Soy Agar plate (Tryptic Soy Agar, Sigma) and placed at 37°C overnight. Colonies were counted to determine the real CFU (Colony-Forming Units) of inoculums.

The numeration was realized on the drop (50 μ l) that contain between 20 and 100 colonies. The shaded boxes correspond to the dilutions seeded on plates.

Table 8-1: Dilutions of the bacterial suspension in order to determine the real CFU (viability) of inoculums. The shaded boxes correspond to the dilutions seeded on plates.

Bacterial overnight suspension in PBS	Dilutions					
	1	2	3	4	5	6
	1/10	1/100	1/10 ³	1/10 ⁴	1/10 ⁵	1/10 ⁶
For example, initial concentration: 3.4×10^7 CFU/ml	3.4×10^6	3.4×10^5	34000	3400	340	34
Number of colonies / 50 μ l (1 drop)	1.7×10^5	1.7×10^4	1.7×10^3	170	17	1.7

Randomization

Before experiment (D-2), 20 healthy OF1 female mice were randomized in 2 groups of 10 mice/group according to body weight criteria, so that the mean body weights of groups was not statistically different.

Treatments

Treatments were performed from D-2 to D7 (D0: day of bacteria inoculation) by oral (PO) administration. Bacterial inoculations were done by intraperitoneal injection (IP).

Inoculation

30 A D0, mice were inoculated with bacterial suspension. Each inoculation consisted in an intraperitoneal injection of 500 μ l of the suspension at 3.4×10^7 CFU/ml (1.7×10^7 CFU /mouse).

Experimental design

Animals were treated with an oral administration of the Test Substance from D-2 to D7. At D0, treatments were performed 1h after bacteria inoculation.

The experimental groups will be defined as described below and summarized in Table 8-2.

- The 10 mice of the group 1 will be treated from D-2 to D7 with the Vehicle by oral administration according to the treatment schedule Q1DX10.
- The 10 mice of the group 2 will be treated from D-2 to D7 with the Test Substance by oral administration according to the treatment schedule Q1DX10.

Table 8-2: Experimental design

Groups	Number of mice	Test substance	Administration route	Administration volume	Treatment schedule	Bacterial model
1	10	Vehicle	PO	10 mL/kg	Q1DX10 (from D-2 to D7)	IP lethal Newman <i>Staphylococcus aureus</i> (3.4x10 ⁷ CFU/ml) 500 µl
2		Test Substance				

10 Monitoring of animals

From D0 to D1, mice were observed twice a day for mortality.

From D2 to D15, mice were observed once a day for mortality.

The body weight of mice was recorded twice a week until the end of the experiment.

- 15 Detailed clinical observations (changes in skin, fur, eyes, mucous membranes, occurrence of secretions and excretions, respiratory function, changes in gait, posture) was monitored twice a week as described in Table 8-3 below for example. Description of this list is not exhaustive and was complemented by other observations, if necessary.

Table 8-3 Clinical observations/ parameters

Parameters	Description	Parameters	Description	Parameters	Description
Mobility/Gait	Stationary	Behavior	Fearful	Fur	Normal
	Reduced		Normal		Spiked
	Normal		Aggressive		Loss
	Excessive	Breathing	Slow		Colored
Carriage	Normal		Normal		Not applicable
	Prostrate		Speed	Mucous membranes	Normal
Paralysis	No	Eyes	Irregular		Abnormal (nose)
	Front right		Normal		Abnormal (mouth)
	Front left		Tearful		Abnormal (nose/mouth)
	Hind right	Skin	Closed	Excretions	Few
	Hind left		Normal		Normal
	Right		Redness		Abundant
	Left		Patch		Dry
	Front		Wound		
	Hind		Hematoma		

Sacrifice of mice

At D15, survived mice were sacrificed by exsanguinations, followed by cervical dislocation, if needed. No autopsy was performed.

Results

- 5 The analysis of mice survival showed the results reported in the following Tables 8-4 and 8-5.

Table 8-4 Number of mice survived per day

Number of survived mice per day									
Groups	N D0	N D1	N D2	N D3	N D4	N D5	N D6	N D7	N D8
Group 1 (Vehicle, PO, Q1Dx10, 10 ml/kg)	10	6	6	5	5	3	2	NA	0
Group 2 (TS pure, PO, Q1Dx10, 10 ml/kg)	10	6	6	6	6	6	4	NA	3

Table 8-5 Percentage of survived mice per day

Percentage of survival									
Groups	N D0	N D1	N D2	N D3	N D4	N D5	N D6	N D7	N D8
Group 1 (Vehicle, PO, Q1Dx10, 10 ml/kg)	100%	60%	60%	50%	50%	30%	20%	NA	0%
Group 2 (TS pure, PO, Q1Dx10, 10 ml/kg)	100%	60%	60%	60%	60%	60%	40%	NA	30%

- 10 NA= Not Available

The treatment allowed to reduce the mortality of mice from 100% to 70%. This animal study thus showed that the claimed antimicrobial and anti-biofilm activities observed *in-vitro* is translated into a therapeutic effect, when the water extract of the invention is administered via oral route to animals.

Example 9

- 15 In this Example we illustrate simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus* of the disclosed composition of 2 plants among 3: *Filipendula ulmaria* PA13, *Camellia sinensis* PA21, *Arctostaphylos uva-ursi* PB12 and at least 1 plant among 3: *Eugenia caryophyllus* PC20, *Vitis vinifera var. tinctorial* PA22, *Desmodium adscendens* PB07. The dried plants of pharmaceutical grade were obtained from “Herboristerie Cailleau” (Chemillé, France).
- 20 Starting from such three different plant powders six herbal compositions were prepared according to the Method A reported before.

List 9-1 : herbal compositions of processed samples

- M[02P]-064: PA13; PA21;
 M[03P]-050: PA13; PA21; PC20;
 25 M[03P]-053: PA13; PA21; PB07;
 M[03P]-056: PA13; PA21; PA22;
 M[02P]-065: PA13; PB12;
 M[03P]-051: PA13; PB12; PC20;
 M[03P]-054: PA13; PB12; PB07;
 30 M[03P]-057: PA13; PB12; PA22;
 M[02P]-066: PA21; PB12;

M[03P]-052: PA21; PB12; PC20;
 M[03P]-055: PA21; PB12; PB07;
 M[03P]-058: PA21; PB12; PA22;
 M[03P]-059: PA13; PB12; PA21;
 5 M[04P]-025: PA13; PB12; PA21; PC20;
 M[04P]-026: PA13; PB12; PA21; PB07;
 M[04P]-027: PA13; PB12; PA21; PA22.

Starting from the herbal compositions listed in List 6-3 corresponding water extracts (or processed samples) were prepared according to the previously described Method B using 40mL of water containing 100g/l of sucrose for water extraction and including the decantation step before filtration.

The biological activity of each processed sample was determined according to the previously described Method C by using the following *Staphylococcus aureus* strains: ATCC 6538, with the following final dilutions of processed samples: 1:5 for the biofilm formation inhibition and 1:10 for the growth inhibition. Note that growth inhibition is close to 100% for each mix at 1:5 dilution, and biofilm formation inhibition is close to 0% for all mixes at 1:10 dilution.

The bacteria planktonic growth inhibition percentage and biofilm formation inhibition percentage over all the tested strain of the processed samples obtained from the different herbal compositions are reported in the following Table 9-1.

Table 9-1: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the *Staphylococcus aureus* strain ATCC® 6538™ of the processed sample based on the compositions given in List 9-1. The processed samples were used at a 1:10 dilution for the planktonic growth inhibition tests and at a dilution of 1:5 for the biofilm formation inhibition.

(A)	Planktonic growth inhibition	Biofilm formation inhibition
M[02P]-064	47%	34%
M[03P]-050	70%	56%
M[03P]-053	55%	35%
M[03P]-056	56%	38%
(C)	Planktonic growth inhibition	Biofilm formation inhibition
M[02P]-066	40%	23%
M[03P]-052	53%	28%
M[03P]-055	59%	27%
M[03P]-058	60%	33%

(B)	Planktonic growth inhibition	Biofilm formation inhibition
M[02P]-065	42%	15%
M[03P]-051	59%	36%
M[03P]-054	54%	22%
M[03P]-057	54%	34%
(D)	Planktonic growth inhibition	Biofilm formation inhibition
M[03P]-059	59%	87%
M[04P]-025	69%	90%
M[04P]-026	64%	90%
M[04P]-027	64%	89%

(A) lists extracts obtained from herbal compositions containing PA13 and PA21, in which the addition of one among PC20, PB07 and PA22 improves the activity; (B) lists extracts obtained from herbal compositions containing PA13 and PB12, in which the addition of one among PC20, PB07 and PA22 improves the activity; (C) lists extracts obtained from herbal compositions containing PA13, PA21 and PB12, in which the addition of one among PC20, PB07 and PA22 improves the activity; (D) lists extracts obtained from herbal compositions containing PA13 and PA21, in which the addition of one among PC20, PB07 and PA22 improves the activity.

Example 10

In this Example, we illustrate simultaneous antimicrobial and antibiofilm activities against *Staphylococcus aureus* of 324 mixtures extracts obtained from the disclosed herbal compositions comprising 2 plants among 3: *Filipendula ulmaria* (PA13), *Camellia sinensis* (PA21), *Arctostaphylos uva-*
 5 *ursi* (PB12) and at least 1 plant among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera var. tinctorial* (PA22), *Desmodium adscendens* (PB07).

For the mixes from M[20C]-2083 to M[20C]-20798, the dried plants of pharmaceutical grade were obtained from “Herboristerie Cailleau” (Chemillé, France). For the mixes from M[20C]-2805 to M[20C]-
 10 2957, the dried plants of pharmaceutical grade were obtained from “Pharmacie Fontgèze” (Clermont-Ferrand, France, plants PA00 to PB12) and “Pharmacie St Herem” (Clermont-Ferrand, France, plants PB13 to PD02).

Herbal compositions containing from 3 to 20 plant powders were prepared according to the previously described Method A.

Starting from the herbal compositions, corresponding water extracts (or processed samples) were
 15 prepared according to the previously described Method B using 40mL of water containing 100g/l of sucrose for water extraction and including the decantation step before filtration.

The biological activity of each processed sample was determined according to the previously described Method C by using the following *Staphylococcus aureus* strains: ATCC 25923, ATCC 49476, ATCC 6538, ATCC 51740, ATCC 29213 and ATCC 14775, with the 1:20 final dilutions of processed samples.

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from the following
 20 herbal compositions of, respectively, List 10-1, List 10-2, List 10-3 and List 10-4 is given in, respectively, Table 10-1, Table 10-2, Table 10-3 and Table 10-4. In each set, the extracts obtained from herbal compositions containing more than 3 or 4 plants were found to exhibit planktonic growth inhibition or
 25 biofilm formation inhibition (or both inhibitions) greater than any of the mixes containing only 3 or 4 plants

List 10-1: Extracts from herbal compositions containing 2 plants among 3: *Filipendula ulmaria* (PA13), *Camellia sinensis* (PA21) and at least one among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera var. tinctorial* (PA22), *Desmodium adscendens* (PB07):

30 M[03P]-050: PA13; PA21; PC20;
 M[03P]-053: PA13; PA21; PB07;
 M[03P]-056: PA13; PA21; PA22;
 M[20C]-2808: PA11; PA12; PA13; PA13; PA18; PA21; PA23; PA24; PA28; PA69; PB07; PB88; PC23; PC24;
 PC49; PC52; PC55 (2 spoons); PC60; PC97;
 35 M[20C]-2820: PA10; PA11 (2 spoons); PA12; PA13; PA14; PA18; PA20; PA21; PA25; PA29; PA33; PA35;
 PA67; PA75; PB07; PB60; PC33; PC35; PC55 ;
 M[20C]-2821: PA11; PA13 (2 spoons); PA19; PA20; PA21; PA22; PA25; PA29; PA33; PA35; PA75; PB07;
 PB36; PC20; PC33 (2 spoons); PC49; PC55; PC97;
 M[20C]-2826: PA11; PA13 (2 spoons); PA20; PA21; PA24; PA25; PA29; PA33; PA34; PA35; PA52; PA70;
 40 PA75; PB07; PC11; PC21; PC33; PC37; PC74;
 M[20C]-2827: PA10; PA11; PA13 (2 spoons); PA20; PA21; PA25; PA26; PA29; PA33; PA35; PA57; PA75;
 PB02; PB07; PB28; PC11; PC23; PC33; PC49;
 M[20C]-2830: PA10; PA11; PA12; PA13; PA14; PA18; PA21; PA22; PA25; PA29; PA32; PA34; PA67; PB07;
 PB60; PB88; PC33; PC35; PC55; PC98;
 45 M[20C]-2831: PA13; PA13; PA19; PA21; PA22; PA22; PA25; PA29; PA32; PA34; PB07; PB36; PB88; PC20;
 PC33; PC33; PC49; PC55; PC97; PC98;

- M[20C]-2833: PA10; PA11; PA13; PA14; PA21; PA22; PA24; PA25; PA26; PA29; PA32; PA34; PB33; PB88; PB88; PC12; PC23; PC27; PC33; PC98;
- M[20C]-2836: PA13; PA13; PA21; PA22; PA24; PA25; PA29; PA32; PA34; PA34; PA52; PA70; PB07; PB88; PC11; PC21; PC33; PC37; PC74; PC98;
- 5 M[20C]-2837: PA10; PA13 (2 spoons); PA21; PA22; PA25; PA26; PA29; PA32; PA34; PA57; PB02; PB07; PB28; PB88; PC11; PC23; PC33; PC49; PC98;
- M[20C]-2838: PA12; PA13 (2 spoons); PA18; PA21; PA21; PA22; PA24; PA25; PA29; PA32; PA34; PB88; PC23; PC33; PC49; PC52; PC55; PC97; PC98;
- 10 M[20C]-2839: PA13 (2 spoons); PA14; PA21; PA22; PA25; PA25; PA26; PA29; PA32; PA34; PA39; PA52; PA57; PB33; PB88; PC33; PC71; PC97; PC98;
- M[20C]-2858: PA12; PA13; PA18; PA20; PA21; PA23; PA24; PA24; PA67; PB07; PB28; PB33; PC12; PC20; PC21; PC23; PC49; PC52; PC55; PC97;
- M[20C]-2861: PA13; PA19; PA21; PA22 (2 spoons); PB07; PB28; PB36; PB60; PC11; PC20; PC20; PC23; PC33; PC49; PC55 (2 spoons); PC60; PC74; PC97;
- 15 M[20C]-2866: PA13; PA21; PA22; PA24; PA34; PA52; PA70; PB07; PB28; PB60; PC11 (2 spoons); PC20; PC21; PC23; PC37; PC55; PC60; PC74 (2 spoons);
- M[20C]-2867: PA10; PA13; PA21; PA22; PA26; PA57; PB02; PB07; PB28; PB28; PB60; PC11 (2 spoons); PC20; PC23; PC23; PC49; PC55; PC60; PC74;
- M[20C]-2868: PA12; PA13; PA18; PA21; PA21; PA22; PA24; PB28; PB60; PC11; PC20; PC23 (2 spoons); PC49; PC52; PC55 (2 spoons); PC60; PC74; PC97;
- 20 M[20C]-2869: PA13; PA14; PA21; PA22; PA25; PA26; PA39; PA52; PA57; PB28; PB33; PB60; PC11; PC20; PC23; PC55; PC60; PC71; PC74; PC97;
- M[20C]-2871: PA10; PA11; PA19; PA21; PA22; PA35; PA57; PB07; PB07; PB36; PB77; PC20 (2 spoons); PC23; PC33; PC49; PC55; PC71; PC97;
- 25 M[20C]-2876: PA10; PA11; PA13; PA21; PA24; PA34; PA35; PA52; PA57; PA70; PB07 (2 spoons); PB77; PC11; PC20; PC21; PC23; PC37; PC71; PC74;
- M[20C]-2877: PA10; PA10; PA11; PA13; PA21; PA26; PA35; PA57; PA57; PB02; PB07; PB28; PB77; PC11; PC20; PC23 (2 spoons); PC49; PC71;
- M[20C]-2878: PA10; PA11; PA12; PA13; PA18; PA21; PA21; PA24; PA35; PA57; PB07; PB77; PC20; PC23; PC23; PC49; PC52; PC55; PC71; PC97;
- 30 M[20C]-2879: PA10; PA11; PA13; PA14; PA21; PA25; PA26; PA35; PA39; PA52; PA57 (2 spoons); PB07; PB33; PB77; PC20; PC23; PC71 (2 spoons); PC97;
- M[20C]-2889: PA10; PA12; PA13; PA14; PA18; PA21; PA26; PA26; PA28; PA29; PA33; PA39; PA75; PB36; PC20; PC52; PC55 (2 spoons); PC60;
- 35 M[20C]-2894: PA11 (2 spoons); PA13; PA21; PA28; PA39; PA39; PA70; PB02; PB02; PB07; PB99 (2 spoons); PC14; PC20; PC49; PC71; PC97; PC98;
- M[20C]-2944: PA10; PA11 (2 spoons); PA13; PA14; PA21; PA28; PA39; PA70; PB02; PB07; PB27; PB99; PC11; PC20; PC23; PC67; PC71; PC97; PC98;
- 40 M[20C]-2954: PA11; PA12; PA13; PA14; PA21; PA23; PA28; PA33; PA39; PA70; PB01; PB02; PB07; PB99; PC20; PC27; PC52; PC71; PC97; PC98.

Table 10-1: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on the extracts of List 10-1 at 1:20 dilution

	Planktonic growth inhibition	Biofilm formation inhibition
M[03C]-050	41%	31%
M[03C]-053	37%	23%
M[03C]-056	34%	21%
M[20C]-2808	66%	30%
M[20C]-2820	54%	27%

	Planktonic growth inhibition	Biofilm formation inhibition
M[20C]-2858	58%	32%
M[20C]-2861	58%	31%
M[20C]-2866	57%	43%
M[20C]-2867	62%	32%
M[20C]-2868	74%	36%

M[20C]-2821	53%	24%
M[20C]-2826	56%	25%
M[20C]-2827	51%	26%
M[20C]-2830	56%	37%
M[20C]-2831	64%	31%
M[20C]-2833	80%	39%
M[20C]-2836	59%	25%
M[20C]-2837	58%	25%
M[20C]-2838	70%	30%
M[20C]-2839	56%	30%

M[20C]-2869	85%	45%
M[20C]-2871	63%	26%
M[20C]-2876	66%	28%
M[20C]-2877	81%	43%
M[20C]-2878	65%	28%
M[20C]-2879	58%	28%
M[20C]-2889	62%	41%
M[20C]-2894	55%	29%
M[20C]-2944	68%	16%
M[20C]-2954	75%	27%

List 10-2: extracts from herbal compositions containing *Filipendula ulmaria* (PA13), *Arctostaphylos uva-ursi* (PB12) and at least one among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera var. tinctoria* (PA22), *Desmodium adscendens* (PB07):

- 5 M[03C]-051: PA13; PB12; PC20;
M[03C]-054: PA13; PB12; PB07;
M[03C]-057: PA13; PB12; PA22;
M[20C]-2083: PA02; PA13; PA28 (2 spoons); PA31; PA36; PA40; PA79; PA93; PB01; PB12; PB17; PB18;
PB31; PB51; PB77; PC20; PC58; PC64; PC66;
- 10 M[20C]-2084: PA13; PA28; PA31; PA36; PA61; PA76; PA76; PA79; PA82; PB01; PB10; PB12; PB17;
PB31; PB57; PC20; PC36; PC41; PC64; PC66;
M[20C]-2085: PA12; PA13; PA14; PA28; PA30; PA31; PA36; PA79; PA94; PB00; PB01; PB08; PB12;
PB17; PB31; PC02; PC14; PC20; PC64; PC66;
M[20C]-2090: PA08; PA13; PA20; PA23; PA26; PA32; PA46; PA52; PA57; PA79; PA95; PB10; PB12;
PB33; PB76; PB82; PC08; PC20; PC21; PD00;
- 15 M[20C]-2091: PA08; PA13; PA26; PA52; PA57; PA72; PA79; PA88; PA92; PA95; PB12; PB60; PB76;
PB83; PC08; PC11; PC20; PC26; PC68; PD00;
M[20C]-2092: PA08; PA13; PA17; PA26; PA52; PA53; PA57; PA79; PA95; PB06; PB12; PB50; PB76;
PC08; PC20; PC20; PC27; PC97; PD00; PD02;
- 20 M[20C]-2100: PA13; PA20; PA23; PA32; PA41; PA46; PA53; PA79; PA80; PA90; PB08; PB10; PB12;
PB33; PB82; PB85; PC20; PC21; PC28; PC75;
M[20C]-2102: PA13; PA17; PA41; PA53; PA53; PA79; PA80; PA90; PB06; PB08; PB12; PB50; PB85; PC20;
PC20; PC27; PC28; PC75; PC97; PD02;
- 25 M[20C]-2104: PA13; PA41; PA53; PA61; PA76; PA76; PA79; PA80; PA82; PA90; PB08; PB10; PB12;
PB57; PB85; PC20; PC28; PC36; PC41; PC75;
M[20C]-2105: PA12; PA13; PA14; PA30; PA41; PA53; PA79; PA80; PA90; PA94; PB00; PB08; PB08;
PB12; PB85; PC02; PC14; PC20; PC28; PC75;
M[20C]-2110: PA00; PA10; PA13; PA20; PA23; PA25; PA30; PA32; PA46; PA52; PA79; PB05; PB10;
PB12; PB33; PB82; PC01; PC20; PC21; PD07;
- 30 M[20C]-2111: PA00; PA10; PA13; PA25; PA30; PA52; PA72; PA79; PA88; PA92; PB05; PB12; PB60;
PB83; PC01; PC11; PC20; PC26; PC68; PD07;
M[20C]-2120: PA11; PA13; PA16; PA20; PA23; PA32; PA44; PA46; PA60; PA67; PA73; PA79; PB10;
PB12; PB33; PB48; PB82; PC17; PC20; PC21;
M[20C]-2122: PA11; PA13; PA16; PA17; PA44; PA53; PA60; PA67; PA73; PA79; PB06; PB12; PB48;
PB50; PC17; PC20; PC20; PC27; PC97; PD02;
- 35 M[20C]-2131: PA05; PA13; PA18; PA37; PA55; PA72; PA72; PA79; PA88; PA92; PA97; PB04; PB12;
PB60; PB83; PC11; PC20; PC26; PC35; PC68;
M[20C]-2135: PA05; PA12; PA13; PA14; PA18; PA30; PA37; PA55; PA72; PA79; PA94; PA97; PB00;
PB04; PB08; PB12; PC02; PC14; PC20; PC35;
- 40 M[20C]-2137: PA05; PA13; PA18; PA27; PA37; PA46; PA55; PA72; PA79; PA82; PA97; PB00; PB04;
PB11; PB12; PC20; PC35; PC45; PC52; PC67;

- M[20C]-2140: PA13; PA14; PA20; PA23; PA32; PA37; PA46; PA79; PA85; PB10; PB12; PB33; PB49; PB82; PB98; PC20; PC21; PC55; PC62; PC98;
- M[20C]-2141: PA13; PA14; PA37; PA72; PA79; PA85; PA88; PA92; PB12; PB49; PB60; PB83; PB98; PC11; PC20; PC26; PC55; PC62; PC68; PC98;
- 5 M[20C]-2142: PA13; PA14; PA17; PA37; PA53; PA79; PA85; PB06; PB12; PB49; PB50; PB98; PC20; PC20; PC27; PC55; PC62; PC97; PC98; PD02;
- M[20C]-2143: PA02; PA13; PA14; PA28; PA37; PA40; PA79; PA85; PA93; PB12; PB18; PB49; PB51; PB77; PB98; PC20; PC55; PC58; PC62; PC98;
- 10 M[20C]-2144: PA13; PA14; PA37; PA61; PA76; PA76; PA79; PA82; PA85; PB10; PB12; PB49; PB57; PB98; PC20; PC36; PC41; PC55; PC62; PC98;
- M[20C]-2150: PA13; PA20; PA23; PA31; PA32; PA46; PA60; PA74; PA79; PB03; PB10; PB12; PB33; PB82; PC20; PC21; PC29; PC37; PC60; PC86;
- M[20C]-2151: PA13; PA31; PA60; PA72; PA74; PA79; PA88; PA92; PB03; PB12; PB60; PB83; PC11; PC20; PC26; PC29; PC37; PC60; PC68; PC86;
- 15 M[20C]-2152: PA13; PA17; PA31; PA53; PA60; PA74; PA79; PB03; PB06; PB12; PB50; PC20; PC20; PC27; PC29; PC37; PC60; PC86; PC97; PD02;
- M[20C]-2153: PA02; PA13; PA28; PA31; PA40; PA60; PA74; PA79; PA93; PB03; PB12; PB18; PB51; PB77; PC20; PC29; PC37; PC58; PC60; PC86;
- M[20C]-2154: PA13; PA31; PA60; PA61; PA74; PA76; PA76; PA79; PA82; PB03; PB10; PB12; PB57; PC20; PC29; PC36; PC37; PC41; PC60; PC86;
- 20 M[20C]-2155: PA12; PA13; PA14; PA30; PA31; PA60; PA74; PA79; PA94; PB00; PB03; PB08; PB12; PC02; PC14; PC20; PC29; PC37; PC60; PC86;
- M[20C]-2163: PA06; PA13; PA17; PA20; PA23; PA72; PA79; PA80; PA92; PB01; PB03; PB10; PB12; PB33; PB57; PB82; PC20; PC62; PC86; PC97;
- 25 M[20C]-2164: PA13; PA17; PA20; PA41; PA79; PA80; PA86; PA92; PB03; PB10; PB12; PC20; PC26; PC27; PC29; PC55; PC60; PC62; PC86; PC97;
- M[20C]-2214: PA13; PA37; PA41; PA79; PA86; PB00; PB10; PB12; PB49; PB82; PC20; PC26; PC27; PC29; PC55; PC60; PC62; PC66; PC86;
- M[20C]-2234: PA13; PA41; PA79; PA85; PA86; PB12; PB31; PB50; PB77; PC14; PC20; PC26; PC27; PC29; PC55; PC55; PC60; PC62; PC98;
- 30 M[20C]-2241: PA10; PA13; PA14; PA23; PA36; PA37; PA46; PA79; PB00; PB10; PB11; PB12; PB17; PB57; PC20; PC27; PC29; PC37; PC64; PC68;
- M[20C]-2242: PA13; PA14; PA17; PA23; PA36; PA37; PA60; PA61; PA76; PA79; PA82; PB00; PB06; PB12; PB49; PC20; PC27; PC29; PC68; PC98;
- 35 M[20C]-2243: PA13; PA14; PA23; PA36; PA37; PA52; PA53; PA73; PA79; PA92; PB00; PB01; PB12; PB18; PB60; PC20; PC20; PC27; PC29; PC68;
- M[20C]-2248: PA13; PA14 (2 spoons); PA23; PA25; PA36; PA36; PA37; PA46; PA79; PA85; PB00; PB12; PC02; PC20; PC27; PC29; PC29; PC55; PC68;
- M[20C]-2249: PA10; PA13; PA14; PA23; PA36; PA37; PA53; PA74; PA79; PA92; PB00; PB12; PC20; PC27; PC29; PC62; PC67; PC68; PC86; PC97;
- 40 M[20C]-2250: PA13; PA23; PA37; PA76; PA79; PB10; PB12; PB33; PB50; PB82; PB82; PB85; PB98; PC01; PC11; PC20; PC45; PC55; PC60; PD02;
- M[20C]-2253: PA13; PA37; PA52; PA53; PA73; PA79; PA92; PB01; PB10; PB12; PB18; PB33; PB50; PB60; PB82; PB85; PC01; PC20; PC20; PC45;
- 45 M[20C]-2254: PA11; PA13; PA17; PA27; PA37; PA79; PB00; PB10; PB12; PB33; PB50; PB82; PB83; PB85; PB92; PB98; PC01; PC20; PC27; PC45;
- M[20C]-2255: PA13; PA36; PA37; PA53; PA72; PA79; PA80; PB03; PB10; PB12; PB33; PB49; PB50; PB82; PB82; PB85; PC01; PC20; PC45; PC68;
- M[20C]-2256: PA13; PA23; PA37; PA53; PA61; PA79; PA85; PB10; PB12; PB33; PB50; PB60; PB82; PB85; PC01; PC11; PC20; PC29; PC45; PD02;
- 50 M[20C]-2257: PA13; PA37; PA37; PA60; PA72; PA79; PB10; PB12; PB33; PB49; PB50; PB82; PB85; PC01; PC08; PC20; PC45; PC68; PC98; PD02;
- M[20C]-2258: PA13; PA14; PA25; PA36; PA37; PA46; PA79; PA85; PB10; PB12; PB33; PB50; PB82; PB85; PC01; PC02; PC20; PC29; PC45; PC55;
- 55 M[20C]-2259: PA10; PA13; PA37; PA53; PA74; PA79; PA92; PB10; PB12; PB33; PB50; PB82; PB85; PC01; PC20; PC45; PC62; PC67; PC86; PC97;

- M[20C]-2261: PA10; PA13; PA23; PA36; PA37; PA46; PA53; PA73; PA79; PA97; PB10; PB11; PB12; PB17; PB57; PC20; PC26; PC29; PC37; PC64;
- M[20C]-2278: PA10; PA13; PA14; PA25; PA28; PA30; PA36; PA37; PA41; PA46; PA79; PA85; PB12; PB50; PB60; PC02; PC20; PC29; PC55; PC66;
- 5 M[20C]-2281: PA10; PA13; PA17; PA46; PA52; PA79; PA82; PA92; PA96; PB10; PB11; PB12; PB17; PB57; PB98; PC20; PC36; PC37; PC64; PC86;
- M[20C]-2321: PA13; PA46; PA53; PA79; PA82; PA92; PB03; PB10; PB11; PB12; PB83; PB85; PC20; PC20; PC21; PC55; PC60; PC86; PC97; PD02;
- 10 M[20C]-2406: PA13; PA37; PA46; PA53; PA79; PA85; PB01; PB05; PB10; PB11; PB12; PB17; PB49; PB98; PC20; PC36; PC37; PC45; PC58; PC67;
- M[20C]-2420: PA10; PA13; PA14; PA36; PA46; PA79; PB10; PB10; PB11; PB11; PB12; PB17; PB17; PB33; PB82; PC20; PC54; PC55; PC68; PC98;
- M[20C]-2441: PA00; PA13; PA14; PA17; PA17; PA37; PA37; PA46; PA74; PA79; PB00; PB03; PB10; PB12; PB49; PC08; PC20; PC35; PC62; PC86;
- 15 M[20C]-2443: PA00; PA13; PA14; PA14; PA17; PA36; PA37; PA79; PB00; PB01; PB10; PB12; PC02; PC08; PC11; PC20; PC29; PC62; PC67; PC86;
- M[20C]-2449: PA00; PA13; PA14; PA17; PA36; PA37; PA60; PA79; PA85; PB00; PB10; PB12; PB82; PB98; PC08; PC20; PC29; PC62; PC86; PC98;
- M[20C]-2451: PA10; PA13; PA17; PA23; PA37; PA46; PA74; PA76; PA79; PB00; PB03; PB08; PB10; PB12; PB49; PB49; PC20; PC35; PC97; PC98;
- 20 M[20C]-2460: PA08; PA13; PA14; PA36; PA36; PA46; PA79; PB10; PB11; PB12; PB17; PB17; PB98; PC20; PC36; PC54; PC62; PC68; PC68; PC86;
- M[20C]-2488: PA10; PA13; PA37; PA46; PA53; PA74; PA79; PA82; PB00; PB00; PB12; PB49; PC20; PC29; PC37; PC45; PC54; PC74; PC86; PC98;
- 25 M[20C]-2545: PA00; PA08; PA10; PA13; PA46; PA73; PA76; PA79; PA90; PB01; PB08; PB10; PB12; PB17; PB33; PB33; PC20; PC55; PC68; PD02;
- M[20C]-2551: PA13; PA14; PA16; PA27; PA36; PA79; PA82; PA92; PA95; PB10; PB12; PB57; PB85; PC11; PC20; PC29; PC29; PC54; PC64; PC98;
- M[20C]-2553: PA13; PA16; PA23; PA60; PA79; PA82; PA95; PB00; PB10; PB10; PB12; PB33; PB49; PB50; PB85; PC02; PC11; PC20; PC29; PC98;
- 30 M[20C]-2555: PA00; PA10; PA13; PA16; PA79; PA82; PA90; PA95; PB08; PB10; PB10; PB12; PB33; PB85; PC11; PC20; PC29; PC68; PC98; PD02;
- M[20C]-2580: PA10; PA13; PA25; PA37; PA40; PA46; PA79; PB11; PB12; PB33; PB49; PB82; PB85; PB98; PC20; PC29; PC29; PC36; PC45; PC97;
- 35 M[20C]-2589: PA13; PA14; PA17; PA40; PA46; PA79; PA80; PB11; PB12; PB33; PB49; PB85; PB98; PC20; PC26; PC29; PC37; PC86; PC97; PD02;
- M[20C]-2622: PA13; PA25; PA36; PA36; PA37; PA46; PA46; PA79; PA82; PB12; PB49; PB82; PC20; PC20; PC54; PC55; PC64; PC68; PC86; PC98;
- M[20C]-2627: PA13; PA14; PA36; PA46; PA76; PA79; PA80; PA96; PB12; PB49; PB49; PC20; PC20; PC45; PC54; PC55; PC64; PC97; PC98; PC98;
- 40 M[20C]-2641: PA05; PA13; PA16; PA27; PA36; PA37; PA41; PA79; PA96; PB05; PB10; PB12; PB82; PB85; PB98; PC11; PC20; PC54; PC68; PD02;
- M[20C]-2651: PA13; PA36; PA41; PA46; PA79; PA92; PB00; PB01; PB05; PB12; PB33; PB82; PB82; PB85; PB98; PC20; PC54; PC55; PC86; PD02;
- 45 M[20C]-2654: PA13; PA36; PA46; PA79; PA92; PB00; PB00; PB01; PB08; PB11; PB12; PB33; PB82; PC01; PC20; PC55; PC55; PC64; PC86; PC86;
- M[20C]-2717: PA13; PA14; PA46; PA79; PB00; PB10; PB12; PB49; PB50; PB77; PB85; PB85; PC02; PC20; PC29; PC64; PC64; PC68; PC98; PD02;
- M[20C]-2728: PA05; PA13; PA16; PA30; PA46; PA60; PA61; PA79; PA85; PB04; PB10; PB12; PB49; PB82; PB82; PB85; PB98; PC11; PC20; PC37;
- 50 M[20C]-2730: PA13; PA36; PA37; PA46; PA61; PA79; PA82; PA85; PB00; PB12; PB17; PB33; PB82; PB85; PC01; PC02; PC20; PC29; PC64; PC98;
- M[20C]-2738: PA05; PA13; PA37; PA46; PA61; PA61; PA79; PA85; PB10; PB12; PB82; PB82; PB85; PB85; PB98; PC02; PC20; PC29; PC37; PC64;
- 55 M[20C]-2788: PA00; PA05; PA13; PA27; PA46; PA61; PA79; PA82; PB00; PB10; PB12; PB82; PB85; PB85; PB98; PC20; PC37; PC37; PC58; PC97;

- M[20C]-2805: PA11; PA13; PA13; PA14; PA22; PA23; PA28; PA33; PA35; PA69; PB07; PB12; PB38; PB88; PB88; PC20; PC24; PC55; PC55; PC60;
- M[20C]-2812: PA11; PA11; PA13; PA20; PA23; PA29; PA30; PA83; PA98; PB02; PB12; PB27; PB38; PC11; PC11; PC20; PC23; PC37; PC51; PC60;
- 5 M[20C]-2815: PA11; PA13; PA14; PA22; PA23; PA29; PA33; PA35; PA98; PB02; PB12; PB27; PB38; PB38; PB88; PC11; PC20; PC23; PC37; PC55;
- M[20C]-2840: PA10; PA11; PA12 (2 spoons); PA13; PA14; PA18; PA28; PA33; PA67; PA67; PB07 (2 spoons); PB12; PB60; PC23; PC35; PC37; PC55; PC67;
- 10 M[20C]-2841: PA12; PA13 (2 spoons); PA19; PA22; PA28; PA33; PA67; PB07 (2 spoons); PB12; PB36; PC20; PC23; PC33; PC37; PC49; PC55; PC67; PC97;
- M[20C]-2842: PA11; PA12; PA13 (2 spoons); PA20; PA28; PA30; PA33; PA67; PA83; PB07; PB12 (2 spoons); PC11; PC20; PC23; PC37; PC51; PC60; PC67;
- M[20C]-2843: PA10; PA11; PA12; PA13; PA14; PA24; PA26; PA28; PA33; PA67; PB07; PB12; PB33; PB88; PC12; PC23; PC23; PC27; PC37; PC67;
- 15 M[20C]-2844: PA12; PA13; PA13; PA15; PA18; PA20; PA28; PA32; PA33; PA67 (2 spoons); PB07; PB12; PB12; PC20; PC23; PC37; PC49; PC55; PC67;
- M[20C]-2845: PA12; PA13 (2 spoons); PA14; PA22; PA28; PA33; PA33; PA35; PA67; PB07; PB12 (2 spoons); PB38; PB88; PC20; PC23; PC37; PC55; PC67;
- M[20C]-2847: PA10; PA12; PA13 (2 spoons); PA26; PA28; PA33; PA57; PA67; PB02; PB07; PB07; PB12; PB28; PC11; PC23; PC23; PC37; PC49; PC67;
- 20 M[20C]-2849: PA12; PA13 (2 spoons); PA14; PA25; PA26; PA28; PA33; PA39; PA52; PA57; PA67; PB07; PB12; PB33; PC23; PC37; PC67; PC71; PC97;
- M[20C]-2852: PA11; PA13; PA20 (2 spoons); PA23; PA24; PA30; PA67; PA83; PB07; PB12; PB28; PB33; PC11; PC12; PC20; PC20; PC21; PC51; PC60;
- 25 M[20C]-2854: PA13; PA15; PA18; PA20 (2 spoons); PA23; PA24; PA32; PA67 (2 spoons); PB07; PB12; PB28; PB33; PC12; PC20 (2 spoons); PC21; PC49; PC55;
- M[20C]-2855: PA13; PA14; PA20; PA22; PA23; PA24; PA33; PA35; PA67; PB07; PB12; PB28; PB33; PB38; PB88; PC12; PC20; PC21; PC55;
- M[20C]-2893: PA00; PA06; PA11; PA12; PA13; PA23; PA26; PA28; PA33; PA39; PB02; PB07 (2 spoons); PB12; PB99; PC14; PC20; PC49; PC98; PD02;
- 30 M[20C]-2897: PA11 (2 spoons); PA13; PA14; PA24; PA29; PA39; PA83; PB02; PB07; PB12; PB60; PB99; PC11; PC14; PC20; PC20; PC49; PC51; PC98;
- M[20C]-2930: PA11; PA13; PA14; PA23; PA26; PA28; PA33; PA35; PA83; PB12; PB28; PB33; PB38; PB77; PB88; PC20; PC23; PC37; PC60; PC60;
- 35 M[20C]-2933: PA00; PA06; PA11; PA12; PA13; PA14; PA23; PA26; PA28 (2 spoons); PA33; PA33; PB07; PB12; PB12; PB28; PB38; PB77; PC60; PD02;
- M[20C]-2935: PA10; PA11; PA13; PA14; PA28; PA29; PA33; PA57; PA67; PB07; PB12; PB28 (2 spoons); PB38; PB77; PB77; PC11; PC20; PC52; PC60;
- M[20C]-2936: PA11 (2 spoons); PA13; PA14; PA18; PA28; PA33; PA57; PA67; PB02; PB12; PB28; PB38 (2 spoons); PC11; PC20; PC27; PC52; PC60;
- 40 M[20C]-2937: PA11; PA11; PA13; PA14 (3 spoons); PA28; PA29; PA33; PA83; PB12 (2 spoons); PB28; PB38; PB60; PB77; PC11; PC20; PC51; PC60;
- M[20C]-2939: PA10; PA11; PA12; PA13 (2 spoons); PA14; PA18; PA26; PA28; PA33 (2 spoons); PA39; PB12; PB28; PB38; PB77; PC20; PC52; PC55; PC60;
- 45 M[20C]-2943: PA00; PA06; PA10; PA11; PA12; PA13; PA14; PA23; PA26; PA28; PA33; PB07 (2 spoons); PB12; PB27; PC11; PC20; PC23; PC67; PD02;
- M[20C]-2947: PA10; PA11 (2 spoons); PA13; PA14 (2 spoons); PA24; PA29; PA83; PB07; PB12; PB27; PB60; PC11 (2 spoons); PC20 (2 spoons); PC23; PC51; PC67;
- M[20C]-2953: PA00; PA06; PA12 (2 spoons); PA13; PA14; PA23 (2 spoons); PA26; PA28; PA33 (2 spoons); PB01; PB07 (2 spoons); PB12; PC20; PC27; PC52; PD02;
- 50 M[20C]-2957: PA11; PA12; PA13; PA14 (2 spoons); PA23; PA24; PA29; PA33; PA83; PB01; PB07; PB12; PB60; PC11; PC20 (2 spoons); PC27; PC51; PC52.

Table 10-2: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on the mixes of List 10-2 at 1:20 dilution

	Planktonic growth inhibition	Biofilm formation inhibition
M[03C]-051	37%	24%
M[03C]-054	30%	13%
M[03C]-057	27%	22%
M[20C]-2083	84%	55%
M[20C]-2084	72%	40%
M[20C]-2085	70%	45%
M[20C]-2090	57%	46%
M[20C]-2091	66%	35%
M[20C]-2092	71%	43%
M[20C]-2100	56%	40%
M[20C]-2102	53%	41%
M[20C]-2104	60%	43%
M[20C]-2105	81%	55%
M[20C]-2110	59%	43%
M[20C]-2111	63%	46%
M[20C]-2120	66%	28%
M[20C]-2122	72%	37%
M[20C]-2131	54%	37%
M[20C]-2135	59%	40%
M[20C]-2137	78%	50%
M[20C]-2140	60%	35%
M[20C]-2141	56%	40%
M[20C]-2142	61%	34%
M[20C]-2143	77%	37%
M[20C]-2144	74%	39%
M[20C]-2150	65%	27%
M[20C]-2151	61%	34%
M[20C]-2152	66%	32%
M[20C]-2153	82%	50%
M[20C]-2154	67%	26%
M[20C]-2155	63%	32%
M[20C]-2163	61%	40%
M[20C]-2164	70%	35%
M[20C]-2214	76%	37%
M[20C]-2234	56%	31%
M[20C]-2241	60%	40%
M[20C]-2242	59%	33%
M[20C]-2243	63%	38%
M[20C]-2248	84%	49%
M[20C]-2249	63%	36%
M[20C]-2250	58%	36%
M[20C]-2253	76%	37%

	Planktonic growth inhibition	Biofilm formation inhibition
M[20C]-2321	55%	30%
M[20C]-2406	58%	31%
M[20C]-2420	55%	34%
M[20C]-2441	52%	36%
M[20C]-2443	58%	33%
M[20C]-2449	55%	32%
M[20C]-2451	53%	28%
M[20C]-2460	54%	37%
M[20C]-2488	54%	27%
M[20C]-2545	52%	29%
M[20C]-2551	52%	29%
M[20C]-2553	52%	27%
M[20C]-2555	58%	39%
M[20C]-2580	58%	44%
M[20C]-2589	56%	35%
M[20C]-2622	57%	40%
M[20C]-2627	58%	39%
M[20C]-2641	53%	36%
M[20C]-2651	53%	33%
M[20C]-2654	53%	35%
M[20C]-2717	55%	30%
M[20C]-2728	56%	24%
M[20C]-2730	55%	27%
M[20C]-2738	52%	26%
M[20C]-2788	56%	25%
M[20C]-2805	71%	25%
M[20C]-2812	54%	31%
M[20C]-2815	52%	29%
M[20C]-2840	55%	31%
M[20C]-2841	51%	31%
M[20C]-2842	53%	35%
M[20C]-2843	58%	34%
M[20C]-2844	76%	42%
M[20C]-2845	46%	33%
M[20C]-2847	61%	41%
M[20C]-2849	76%	32%
M[20C]-2852	77%	32%
M[20C]-2854	57%	27%
M[20C]-2855	66%	35%
M[20C]-2893	61%	37%
M[20C]-2897	66%	36%
M[20C]-2930	81%	52%

M[20C]-2254	55%	25%
M[20C]-2255	60%	29%
M[20C]-2256	58%	30%
M[20C]-2257	59%	25%
M[20C]-2258	81%	39%
M[20C]-2259	60%	29%
M[20C]-2261	52%	25%
M[20C]-2278	53%	25%
M[20C]-2281	56%	26%

M[20C]-2933	60%	32%
M[20C]-2935	65%	34%
M[20C]-2936	68%	41%
M[20C]-2937	78%	41%
M[20C]-2939	77%	44%
M[20C]-2943	67%	29%
M[20C]-2947	71%	40%
M[20C]-2953	66%	38%
M[20C]-2957	71%	45%

List 10-3: herbal compositions of processed samples containing: *Camellia sinensis* (PA21), *Arctostaphylos uva-ursi* (PB12) and at least one among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera* var. *tinctorial* (PA22), *Desmodium adscendens* (PB07):

- 5 M[03C]-052: PA21; PB12; PC20;
M[03C]-055: PA21; PB12; PB07;
M[03C]-058: PA21; PB12; PA22;
M[20C]-2888: PA14; PA15; PA18; PA21; PA21; PA23; PA26; PA28; PA28; PA29; PA39; PA57; PA75; PB07;
PB12; PB36; PC11; PC55; PC55; PC60;
- 10 M[20C]-2913: PA00; PA06; PA12; PA21; PA23; PA26; PA26; PA27; PA28; PA33; PB07; PB07; PB12 (2
spoons); PB33; PC20; PC24; PC41; PC71; PD02;
M[20C]-2918: PA15; PA18; PA21 (2 spoons); PA23; PA26; PA27; PA28; PA57; PB07 (2 spoons); PB12 (2
spoons); PB33; PC11; PC20; PC24; PC41; PC55; PC71;
- 15 M[20C]-2920: PA20; PA21; PA23; PA26; PA28; PA35; PA35; PA67; PA83; PB12; PB33; PB88; PC12; PC20;
PC23; PC37; PC37; PC60 (2 spoons); PC74;
M[20C]-2923: PA00; PA06; PA12; PA20; PA21; PA23; PA26; PA28 (2 spoons); PA33; PA35; PA67; PB07;
PB12 (2 spoons); PC12; PC37; PC60; PC74; PD02;
M[20C]-2928: PA15; PA18; PA20; PA21 (2 spoons); PA23; PA28 (2 spoons); PA35; PA57; PA67; PB07;
PB12 (2 spoons); PC11; PC12; PC37; PC55; PC60; PC74;
- 20 M[20C]-2887: PA11; PA14; PA14; PA21; PA24; PA26; PA28; PA29 (2 spoons); PA39; PA75; PA83; PB12;
PB36; PB60; PC11; PC20; PC51; PC55; PC60;
M[20C]-2914: PA11; PA21 (2 spoons); PA26; PA27; PA28; PA39; PA70; PB02; PB07; PB12; PB33; PB99;
PC20; PC24; PC41; PC71 (2 spoons); PC97; PC98;
- 25 M[20C]-2915: PA10; PA21; PA26; PA27; PA29; PA57; PA67; PB07 (2 spoons); PB12; PB28; PB33; PB77;
PC11; PC20 (2 spoons); PC24; PC41; PC52; PC71;
M[20C]-2916: PA11; PA18; PA21; PA26; PA27; PA57; PA67; PB02; PB07; PB12; PB33; PB38; PC11; PC20;
PC20; PC24; PC27; PC41; PC52; PC71;
- 30 M[20C]-2917: PA11; PA14; PA21; PA24; PA26; PA27; PA29; PA83; PB07; PB12 (2 spoons); PB33; PB60;
PC11; PC20 (2 spoons); PC24; PC41; PC51; PC71;
M[20C]-2925: PA10; PA20; PA21; PA28; PA29; PA35; PA57; PA67 (2 spoons); PB07; PB12; PB28; PB77;
PC11; PC12; PC20; PC37; PC52; PC60; PC74;
- 35 M[20C]-2926: PA11; PA18; PA20; PA21; PA28; PA35; PA57; PA67 (2 spoons); PB02; PB12; PB38; PC11;
PC12; PC20; PC27; PC37; PC52; PC60; PC74;
M[20C]-2927: PA11; PA14; PA20; PA21; PA24; PA28; PA29; PA35; PA67; PA83; PB12; PB12; PB60; PC11;
PC12; PC20; PC37; PC51; PC60; PC74;
M[20C]-2910: PA21; PA23; PA26 (2 spoons); PA27; PA35; PA83; PB07; PB12; PB33 (2 spoons); PB88;
PC20 (2 spoons); PC23; PC24; PC37; PC41; PC60; PC71.

Table 10-3: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on the mixes of List 10-2 at 1:20 dilution

	Planktonic growth inhibition	Biofilm formation inhibition
M[03C]-052	31%	13%
M[03C]-055	29%	10%
M[03C]-058	31%	18%
M[20C]-2888	48%	27%
M[20C]-2913	46%	37%
M[20C]-2918	46%	32%
M[20C]-2920	54%	30%
M[20C]-2923	51%	28%
M[20C]-2928	46%	32%

	Planktonic growth inhibition	Biofilm formation inhibition
M[20C]-2887	46%	8%
M[20C]-2914	29%	38%
M[20C]-2915	30%	35%
M[20C]-2916	39%	40%
M[20C]-2917	40%	41%
M[20C]-2925	41%	36%
M[20C]-2926	47%	36%
M[20C]-2927	41%	41%
M[20C]-2910	46%	39%

List 10-4: extracts from herbal compositions containing: *Filipendula ulmaria* (PA13), *Camellia sinensis* (PA21), *Arctostaphylos uva-ursi* (PB12) and at least one among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera var. tinctorial* (PA22), *Desmodium adscendens* (PB07):

- 5 M[04C]-025: PA13; PA21; PB12; PC20;
M[04C]-026: PA13; PA21; PB12; PB07;
M[04C]-027: PA13; PA21; PB12; PA22;
M[20C]-2898: PA11; PA13; PA15; PA18; PA21; PA23; PA28; PA39; PA57; PB02; PB07 (2 spoons); PB07; PB12; PB99; PC11; PC14; PC20; PC49; PC55; PC98;
- 10 M[20C]-2900: PA10; PA11; PA13; PA21; PA23; PA25; PA26; PA35; PA69; PA83; PB12; PB32; PB33; PB88; PC20 (2 spoons); PC23; PC37; PC55; PC60;
M[20C]-2903: PA00; PA06; PA10; PA11; PA12; PA13; PA21; PA23; PA25; PA26; PA28; PA33; PA69; PB07; PB12 (2 spoons); PB32; PC20; PC55; PD02;
M[20C]-2908: PA10; PA11; PA13; PA15; PA18; PA21 (2 spoons); PA23; PA25; PA28; PA57; PA69; PB07;
- 15 PB12 (2 spoons); PB32; PC11; PC20; PC55 (2 spoons);
M[20C]-2938: PA11; PA13; PA14; PA15; PA18; PA21; PA23; PA28 (2 spoons); PA33; PA57; PB07; PB12 (2 spoons); PB28; PB38; PB77; PC11; PC55; PC60;
M[20C]-2948: PA10; PA11; PA13; PA14; PA15; PA18; PA21; PA23; PA28; PA57; PB07 (2 spoons); PB12; PB27; PC11 (2 spoons); PC20; PC23; PC55; PC67;
- 20 M[20C]-2951: PA12; PA13 (2 spoons); PA14; PA21; PA23; PA24; PA33; PA39; PA67; PB01; PB07; PB12; PB77; PC20; PC27; PC33; PC49; PC52; PC60;
M[20C]-2952: PA12; PA13 (2 spoons); PA14; PA21; PA23; PA25; PA26; PA33; PA37; PA67; PB01; PB07 (2 spoons); PB12; PC20; PC27; PC49; PC52; PC71;
M[20C]-2958: PA12; PA13; PA14; PA15; PA18; PA21; PA23 (2 spoons); PA28; PA33; PA57; PB01; PB07 (2 spoons); PB12; PC11; PC20; PC27; PC52; PC55;
- 25 M[20C]-2822: PA11 (2 spoons); PA13 (2 spoons); PA20 (2 spoons); PA21; PA25; PA29; PA30; PA33; PA35; PA75; PA83; PB12; PC11; PC20; PC33; PC51; PC60;
M[20C]-2824: PA11; PA13 (2 spoons); PA15; PA18; PA20 (2 spoons); PA21; PA25; PA29; PA32; PA33; PA35; PA67; PA75; PB12; PC20; PC33; PC49; PC55;
- 30 M[20C]-2825: PA11; PA13 (2 spoons); PA14; PA20; PA21; PA22; PA25; PA29; PA33 (2 spoons); PA35 (2 spoons); PA75; PB12; PB38; PB88; PC20; PC33; PC55;
M[20C]-2834: PA13 (2 spoons); PA15; PA18; PA20; PA21; PA22; PA25; PA29; PA32 (2 spoons); PA34; PA67; PB12; PB88; PC20; PC33; PC49; PC55; PC98;
- 35 M[20C]-2835: PA13 (2 spoons); PA14; PA21; PA22 (2 spoons); PA25; PA29; PA32; PA33; PA34; PA35; PB12; PB38; PB88 (2 spoons); PC20; PC33; PC55; PC98;
M[20C]-2848: PA12 (2 spoons); PA13 (2 spoons); PA18; PA21; PA24; PA28; PA33; PA67; PB07; PB12; PC23 (2 spoons); PC37; PC49; PC52; PC55; PC67; PC97;
M[20C]-2862: PA11; PA13; PA20; PA21; PA22; PA30; PA83; PB12; PB28; PB60; PC11 (2 spoons); PC20 (2 spoons); PC23; PC51; PC55; PC60 (2 spoons); PC74;

- M[20C]-2864: PA13; PA15; PA18; PA20; PA21; PA22; PA32; PA67; PB12; PB28; PB60; PC11; PC20 (2 spoons); PC23; PC49; PC55 (2 spoons); PC60; PC74;
- M[20C]-2865: PA13; PA14; PA21; PA22 (2 spoons); PA33; PA35; PB12; PB28; PB38; PB60; PB88; PC11; PC20 (2 spoons); PC23; PC55 (2 spoons); PC60; PC74;
- 5 M[20C]-2872: PA10; PA11 (2 spoons); PA13; PA20; PA21; PA30; PA35; PA57; PA83; PB07; PB12; PB77; PC11; PC20 (2 spoons); PC23; PC51; PC60; PC71;
- M[20C]-2874: PA10; PA11; PA13; PA15; PA18; PA20; PA21; PA32; PA35; PA57; PA67; PB07; PB12; PB77; PC20 (2 spoons); PC23; PC49; PC55; PC71;
- 10 M[20C]-2875: A10; A11; A13; A14; A21; A22; A33; A35; A35; A57; B07; B12; B38; B77; B88; C20 (2 spoons); C23; C55; C71;
- M[20C]-2882: PA13; PA14; PA21 (2 spoons); PA25; PA26 (2 spoons); PA28; PA29; PA37; PA39; PA67; PA75; PB07; PB12; PB36; PC49; PC55; PC60; PC71;
- M[20C]-2891: PA11; PA13 (2 spoons); PA21; PA24; PA39; PA39; PA67; PB02; PB07; PB12; PB77; PB99; PC14; PC20; PC33; PC49 (2 spoons); PC60; PC98;
- 15 M[20C]-2892: PA11; PA13 (2 spoons); PA21; PA25; PA26; PA37; PA39; PA67; PB02; PB07 (2 spoons); PB12; PB99; PC14; PC20; PC49 (2 spoons); PC71; PC98;
- M[20C]-2901: PA10; PA11; PA13 (2 spoons); PA21 (2 spoons); PA24; PA25; PA39; PA67; PA69; PB12 (2 spoons); PB32; PB77; PC20; PC33; PC49; PC55; PC60;
- 20 M[20C]-2902: PA10; PA11; PA13 (2 spoons); PA21 (2 spoons); PA25 (2 spoons); PA26; PA37; PA67; PA69; PB07; PB12 (2 spoons); PB32; PC20; PC49; PC55; PC71;
- M[20C]-2904: PA10; PA11 (2 spoons); PA13; PA21 (2 spoons); PA25; PA28; PA39; PA69; PA70; PB02; PB12; PB32; PB99; PC20; PC55; PC71; PC97; PC98;
- M[20C]-2905: PA10 (2 spoons); PA11; PA13; PA21; PA25; PA29; PA57; PA67; PA69; PB07; PB12; PB28; PB32; PB77; PC11; PC20 (2 spoons); PC52; PC55;
- 25 M[20C]-2906: PA10; PA11 (2 spoons); PA13; PA18; PA21; PA25; PA57; PA67; PA69; PB02; PB12; PB32; PB38; PC11; PC20 (2 spoons); PC27; PC52; PC55;
- M[20C]-2907: PA10; PA11 (2 spoons); PA13; PA14; PA21; PA24; PA25; PA29; PA69; PA83; PB12 (2 spoons); PB32; PB60; PC11; PC20 (2 spoons); PC51; PC55;
- 30 M[20C]-2909: A10 (2 spoons); A11; A12; A13 (2 spoons); A18; A21; A25; A26; A33; A39; A69; B12; B32; C20 (2 spoons); C52; C55 (2 spoons);
- M[20C]-2911: PA13; PA21 (2 spoons); PA24; PA26; PA27; PA39; PA67; PB07; PB12 (2 spoons); PB33; PB77; PC20; PC24; PC33; PC41; PC49; PC60; PC71;
- M[20C]-2912: PA13; A21 (2 spoons); PA25; PA26 (2 spoons); PA27; PA37; PA67; PB07 (2 spoons); PB12 (2 spoons); PB33; PC20; PC24; PC41; PC49; PC71 (2 spoons);
- 35 M[20C]-2919: PA10; PA12; PA13; PA18; PA21; PA26 (2 spoons); PA27; PA33; PA39; PB07; PB12; PB33; PC20 (2 spoons); PC24; PC41; PC52; PC55; PC71;
- M[20C]-2922: PA13; PA20; PA21 (2 spoons); PA25; PA26; PA28; PA35; PA37; PA67 (2 spoons); PB07; PB12 (2 spoons); PC12; PC37; PC49; PC60; PC71; PC74;
- M[20C]-2929: PA10; PA12; PA13; PA18; PA20; PA21; PA26; PA28; PA33; PA35; PA39; PA67; PB12; PC12; PC20; PC37; PC52; PC55; PC60; PC74;
- 40 M[20C]-2932: PA11; PA13 (2 spoons); PA14; PA21; PA25; PA26; PA28; PA33; PA37; PA67; PB07; PB12 (2 spoons); PB28; PB38; PB77; PC49; PC60; PC71;
- M[20C]-2941: PA10; PA11; PA13 (2 spoons); PA14; PA21; PA24; PA39; PA67; PB07; PB12; PB27; PB77; PC11; PC20; PC23; PC33; PC49; PC60; PC67;
- 45 M[20C]-2942: PA10; PA11; PA13 (2 spoons); PA14; PA21; PA25; PA26; PA37; PA67; PB07 (2 spoons); PB12; PB27; PC11; PC20; PC23; PC49; PC67; PC71;

Table 10-4: average of planktonic growth inhibition percentage and biofilm formation inhibition percentage efficacy on the 6 tested *Staphylococcus aureus* strains of the processed sample based on the mixes of List 10-4 at 1:20 dilution

	Planktonic growth inhibition	Biofilm formation inhibition
M[04C]-025	57%	18%

	Planktonic growth inhibition	Biofilm formation inhibition
M[20C]-2872	59%	36%

M[04C]-026	48%	16%
M[04C]-027	49%	18%
M[20C]-2898	60%	21%
M[20C]-2900	78%	32%
M[20C]-2903	48%	28%
M[20C]-2908	54%	15%
M[20C]-2938	70%	26%
M[20C]-2948	56%	22%
M[20C]-2951	47%	17%
M[20C]-2952	60%	19%
M[20C]-2958	53%	31%
M[20C]-2822	46%	38%
M[20C]-2824	60%	32%
M[20C]-2825	76%	38%
M[20C]-2834	55%	27%
M[20C]-2835	64%	29%
M[20C]-2848	52%	29%
M[20C]-2862	46%	31%
M[20C]-2864	51%	29%
M[20C]-2865	56%	43%

M[20C]-2874	78%	39%
M[20C]-2875	83%	47%
M[20C]-2882	61%	28%
M[20C]-2891	70%	27%
M[20C]-2892	76%	39%
M[20C]-2901	59%	29%
M[20C]-2902	51%	30%
M[20C]-2904	62%	24%
M[20C]-2905	55%	37%
M[20C]-2906	59%	37%
M[20C]-2907	75%	41%
M[20C]-2909	82%	49%
M[20C]-2911	55%	30%
M[20C]-2912	59%	29%
M[20C]-2919	75%	48%
M[20C]-2922	63%	32%
M[20C]-2929	54%	30%
M[20C]-2932	73%	39%
M[20C]-2941	63%	48%
M[20C]-2942	56%	34%

Comparative Example 4

As a comparative Example, we report several extracts of herbal compositions containing only 1 plant among 3: *Filipendula ulmaria* (PA13), *Camellia sinensis* (PA21), *Arctostaphylos uva-ursi* (PB12) and only 1 among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera* var. *tinctoria* (PA22), *Desmodium adscendens* (PB07).

5 herbal compositions having the content reported in List 10-5 were prepared according to the previously described Method 1.

List 10-5 extracts from herbal compositions containing the following plants

- 10 M[20P]-318: PA15; PA21; PA34; PA35; PA82; PA83; PA85; PA92; PA96; PB03; PB07; PB26; PB30; PB36; PB79; PC11; PC49; PC91; PC92; PD10;
- M[20P]-329: PA10; PA28; PA34; PA36; PA55; PA72; PA83; PA85; PA91; PB03; PB12; PB16; PB28; PB35; PB49; PB92; PC20; PC23; PC64; PC85;
- 15 M[20P]-333: PA04; PA05; PA13; PA16; PA22; PA39; PA51; PA55; PA57; PA72; PA82; PA92; PA98; PB05; PB20; PB36; PB43; PB74; PB77; PC37;
- M[20P]-362: PA21; PA23; PA25; PA33; PA37; PA57; PA69; PA81; PA92; PB16; PB25; PB26; PB28; PB29; PB78; PC20; PC21; PC71; PC83; PD02;
- M[20P]-513: PA12; PA14; PA14; PA20; PA20; PA21; PA22; PA33; PA35; PA52; PA57; PA69; PB00; PB32; PB38; PC24; PC37; PC52; PC55; PC67;
- 20 Herbal compositions containing 20 plant powders were prepared according to the previously described Method A.

Starting from the herbal compositions, corresponding water extracts (or processed samples) were prepared according to the previously described Method B using 40mL of water containing 100g/l of sucrose for water extraction and including the decantation step before filtration.

The biological activity of each processed sample was determined according to the previously described Method C by using the following *Staphylococcus aureus* strains: ATCC 25923, ATCC 49476, ATCC 6538, ATCC 51740, ATCC 29213 and ATCC 14775, with the 1:20 final dilutions of processed samples.

The bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from the herbal compositions of List 10-4 is given in Table 10-5. Each percentage is the average of 2 measurements of the same condition in two different wells. Negative percentages reflect growth promotion and biofilm promotion, instead of the desired inhibition. It is appreciated that such extracts give very poor, if any inhibition as compared to the extracts of Example 10.

Table 10-5: bacteria planktonic growth inhibition percentage average and biofilm formation inhibition percentage average over all the tested strains of the processed samples obtained from the herbal compositions of List 10-4

	Planktonic growth inhibition	Biofilm formation inhibition
M[20C]-318	-10%	0%
M[20C]-329	12%	9%
M[20C]-333	3%	18%
M[20C]-362	22%	3%
M[20C]-513	-21%	21%

Example 11

In this Example, we illustrate antimicrobial activities against 14 different microbial species described in Table 11-1 of 10 mixtures extracts obtained from the herbal compositions disclosed in List 11-1 comprising at least 2 plants among 3: *Filipendula ulmaria* (PA13), *Camellia sinensis* (PA21), *Arctostaphylos uva-ursi* (PB12) and at least 1 plant among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera* var. *tintorial* (PA22), and *Desmodium adscendens* (PB07).

Table 11-1: microbial species and strains used in Example 11

Strain full name	Strain code	Innoculum concentration (bacteria/mL)
<i>Staphylococcus aureus subsp. aureus</i> (MRSA) ATCC® 33592™	B-A49	2.10 ^{5.5}
<i>Staphylococcus epidermidis</i> ATCC® 14990™	B-A08	2.10 ⁵
<i>Staphylococcus pseudintermedius</i> ATCC® 49444™	B-A09	2.10 ⁵
<i>Candida albicans</i> ATCC® 60193™	B-A61	2.10 ^{6.5}
<i>Escherichia coli</i> ATCC® 11775™	B-A67	2.10 ^{5.5}
<i>Klebsiella pneumoniae</i> ATCC® 13883™	B-A68	2.10 ^{5.5}
<i>Listeria innocua</i> ATCC® 33090™	B-A69	2.10 ^{5.5}

<i>Listeria monocytogenes</i> ATCC® 19115™	B-A70	2.10 ^{5.5}
<i>Pseudomonas aeruginosa</i> ATCC® 27853™	B-A71	2.10 ^{5.5}
<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Enteritidis</i> ATCC® 13076™	B-A72	2.10 ^{5.5}
<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Typhimurium</i> ATCC® 13311™	B-A73	2.10 ^{5.5}
<i>Streptococcus pneumoniae</i> ATCC® 27336™	B-A75	2.10 ^{5.5}
<i>Streptococcus equi</i> subsp. <i>zooepidemicus</i> ATCC® 43079™	B-A77	2.10 ^{5.5}
<i>Streptococcus pyogenes</i> ATCC® BAA-1323™	B-B26	2.10 ^{6.5}

List 11-1: extracts from herbal compositions containing: *Filipendula ulmaria* (PA13), *Camellia sinensis* (PA21) and at least one among 3: *Eugenia caryophyllus* (PC20), *Vitis vinifera* var. *tinctoria* (PA22), *Desmodium adscendens* (PB07) used in Example-11:

- 5 M[20P]-297: PA10; PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA29; PA33; PA57; PA67; PA69; PB07; PB12; PB60; PC20; PC37; PC49; PC98;
M[20P]-416: PA11; PA13; PA14; PA20; PA21; PA25; PA26; PA28; PA29; PA67; PA75; PB02; PB12; PB27; PB33; PB60; PB88; PC20; PC49; PC60;
M[20P]-418: PA10; PA11; PA12; PA13; PA21; PA22; PA23; PA24; PA26; PA39; PB07; PB12; PB38; PC11; PC12; PC20; PC23; PC24; PC49; PC55;
10 M[20P]-527: PA10; PA10; PA11; PA13; PA15; PA20; PA20; PA21; PA23; PA26 (3 spoons); PA33; PB08; PB12 (2 spoons); PB60; PB88; PC12; PC20;
M[20P]-557: PA00; PA10; PA20; PA21; PA23; PA24; PA29; PA57; PB07; PB12 (4 spoons); PC11 (2 spoons); PC12; PC20; PC51; PC55; PC67;
15 M[20P]-573: PA10; PA11; PA13 (3 spoons); PA14; PA18; PA21; PA23; PA29; PA67; PB00; PB07; PB12; PB33; PB77; PC11; PC52; PC60; PC71;
M[20P]-576: PA11 (2 spoons); PA12; PA13 (3 spoons); PA14 (2 spoons); PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37; PC49;
M[19P]-576: PA11 (2 spoons); PA12; PA13 (3 spoons); PA14 (2 spoons); PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37 ;
20 M[5P]-1: PA12; PA13; PA21; PA22; PB12;
M[5P]-2: PA11; PA13; PA21; PA22; PB12;
For the mixes M[5P]-1, M[5P]-2 and M[19P]-576, the dried plants of pharmaceutical grade were obtained from “Herboristerie Cailleau” (Chemillé, France). For the other mixes, the dried plants of
25 pharmaceutical grade were obtained from “Pharmacie Fontgèze” (Clermont-Ferrand, France, plants PA00 to PB12) and “Pharmacie St Herem” (Clermont-Ferrand, France, plants PB13 to PD02).

All herbal compositions of 20 plants were prepared according to the previously described Method A.

- Starting from the herbal compositions, corresponding water extracts (or processed samples) were prepared according to the previously described Method B using 40mL of water containing 100g/l of sucrose for water extraction and including the decantation step before filtration.
30

Herbal compositions M[19P]-576, M[5P]-1 and M[5P]-2 were prepared by weighting each dried herb as indicated in Table 11-2

Table 11-2: weight of dried plant (mg) used in the herbal compositions of M[19P]-576, M[5P]-1 and M[5P]-2 used in Example-11

	PA11	PA12	PA13	PA14	PA20	PA21	PA22	PA29	PA39	PB12	PB60	PC20	PC26	PC33	PC37
M[19P]-576	409	145	573	316	277	271	207	380	207	247	206	236	198	187	154
M[5P]-1		903	1574			2237	1263			2022					

M[5P]-2	756		720			1027	577			925					
---------	-----	--	-----	--	--	------	-----	--	--	-----	--	--	--	--	--

Starting from the herbal compositions, corresponding water extracts (or processed samples) were prepared according to the previously described Method B using, using, respectively, 30mL, 40mL and 40mL, of water without sucrose with, respectively, M[19P]-576, M[5P]-1 and M[5P]2, for water extraction and including the decantation step before filtration.

The biological activity of each processed sample was determined according to the previously described Method C by using the following strains at the indicated final dilutions of processed samples is given in Table 11-3.

The microbial planktonic growth inhibition percentage average is given in Table 11-3.

Table 11-3: planktonic growth inhibition percentage on the 14 tested microbial strains of the processed sample of composition disclosed in List 11-1

	M[20C]-297	M[20C]-416	M[20C]-418	M[20C]-527	M[20C]-557	M[20C]-573	M[20C]-576	M[19C]-576	M[5P]-1	M[5P]-2
Dilution	1:10	1:10	1:10	1:10	1:10	1:10	1:10	1:20	1:20	1:20
B-A49	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%
B-A08	100%	99%	100%	100%	100%	98%	98%	100%	100%	100%
B-A09	89%	97%	98%	99%	99%	100%	100%	100%	100%	100%
B-A61	25%	69%	36%	45%	51%	55%	90%	ND	ND	ND
B-A67	45%	52%	59%	66%	62%	52%	39%	34%	55%	38%
B-A68	100%	97%	100%	100%	100%	94%	92%	100%	100%	100%
B-A69	-1%	21%	21%	38%	30%	30%	-1%	38%	100%	38%
B-A70	39%	52%	53%	74%	54%	57%	32%	34%	100%	40%
B-A71	63%	71%	69%	75%	75%	82%	71%	42%	100%	55%
B-A72	47%	53%	59%	63%	58%	58%	32%	39%	62%	42%
B-A73	59%	59%	59%	71%	72%	58%	32%	42%	100%	45%
B-A75	51%	52%	63%	71%	63%	57%	31%	ND	ND	ND
B-A77	40%	47%	64%	68%	59%	58%	40%	100%	100%	100%
B-B26	58%	64%	72%	70%	74%	42%	93%	ND	ND	ND

Note: "ND" stands for "not determined"

Example 12

In this Example, we illustrate antimicrobial activities of a plant mix where the extraction step was realized with the non-aqueous solvents acetone, ethyl-acetate, ethanol, used alone or as binary mixes as indicated in Table 12-1. A comparison with water is provided.

Table 12-1: composition of the binary mixes used in Example 12 for non-aqueous extraction

A)	Binary solvent name	Acetone to Ethanol ratio	B)	Binary solvent name	Ethyl-acetate to Ethanol ratio	C)	Binary solvent name	Ethyl-acetate to Acetone ratio
	AC9ET1	90/10		EA9ET1	90/10		EA9AC1	90/10
	AC8ET2	80/20		EA8ET2	80/20		EA8AC2	80/20
	AC7ET3	70/30		EA7ET3	70/30		EA7AC3	70/30

AC6ET4	60/40	EA6ET4	60/40	EA6AC4	60/40
AC5ET5	50/50	EA5ET5	50/50	EA5AC5	50/50
AC4ET6	40/60	EA4ET6	40/60	EA4AC6	40/60
AC3ET7	30/70	EA3ET7	30/70	EA3AC7	30/70
AC2ET8	20/80	EA2ET8	20/80	EA2AC8	20/80
AC1ET9	10/90	EA1ET9	10/90	EA1AC9	10/90

The M[20P]-576 dried plant mix was composed of following Table 12-2 and placed in a 500mL vial and thoroughly mixed by manual flipping and rolling. The dried herbs were obtained from “Herboristerie Cailleau” (Chemillé, France), except PC49 which was obtained from Siccarapam (Aubiat, France).

5 **Table 12-2: list of dried plant composing the M[20P]-576 used in Example 12**

Dried plant	PA11	PA12	PA13	PA14	PA20	PA21	PA22	PA29	PA39	PB12	PB60	PC20	PC26	PC33	PC37	PC49
Weight (g)	29,7	10,5	41,5	22,9	19,8	19,9	15,0	27,5	15,0	17,8	15,0	17,1	14,4	13,6	11,3	9,3

For each solvent tested, 5g of dried plant mix were placed into a 100ml coloured vial (Schott, Germany) with a magnetic bar and 50ml solvent was added. The liquid suspension was stirred with a magnetic stirrer (RO 15P IKA, Germany) at maximum speed during 48 hours at room temperature. Each suspension/solution was filtered by gravity using a 250mm folded filter (J025106, Prats Dumas, France) placed on a glass funnel (Schott, Germany) itself placed into an evaporation flask. The flask containing the liquid which crossed the filter is placed on a rotary vacuum evaporation device (Rotavapor® R205 + V800 vacuum controller, Büchi, Switzerland) and evaporation is performed at 35°C with vacuum adjusted to be below the solvent’s ebullition point, until all solvent has evaporated. The remaining material is weighted and dissolved into DMSO (Acros organic, USA) qsp 10g/L to form the extracted mix.

Two water extracts were prepared according to the previously described Method B, where the spoons of dried plants were replaced by 2g of the M[20P]-576 dried plant mix of Example 12, using 40mL of water for water extraction and including the decantation step before filtration. The dry residue content of one of the plant mix extract has been determined by weighting the lyophilisation residue obtained with a lyophilisation device (FreeZone™ 2.5, Labconco USA, with RZ6 Vacuubrand, Germany) of the frozen extract.

These dry residue contents were used to deduce the specific activities of the different extracts.

The biological activity of each extracted mix was determined according to the previously described Method C by using the strains indicted in Table 12-3, where the dilution of the extracted mix was: 1:20, 1:40, 1:80, 1:160. The minimal inhibiting concentration (MIC) was noted by visual observation of the wells as the lowest concentration where no growth is observed.

Table 12-3: list of the bacterial strains used in the Example 12

<i>Staphylococcus aureus</i>		<i>Staphylococcus epidermidis</i>	
NCTC 12493™	B-A47	ATCC® 12228™	B-A22
ATCC® 33591™	B-A48	ATCC® 700296™	B-A23
ATCC® 33592™	B-A49	ATCC® 49461™	B-A24
NCTC 43300™	B-A51	ATCC® 14990™	B-A08

The results are compiled in tables 12-4 for the different solvents.

Table 12-4: CMI in mg/L of the extracted mixes using, A) acetone, ethanol-acetate, ethanol and water as a solvent, B), different proportions of acetone and ethanol as binary solvent as indicated in Table 12-2-A, C) different proportions of ethanol-acetate and ethanol as binary solvent as indicated in Table 12-2-B, and D) using different proportions of ethanol-acetate and acetone as binary solvent as indicated in Table 12-2-C.

A)	Acetone	Ethanol-Acetate	Ethanol	Water					
B-A47	500	500	500	725					
B-A48	500	500	500	725					
B-A49	500	500	500	725					
B-A51	500	500	500	725					
B-A22	250	250	250	363					
B-A23	250	250	250	363					
B-A24	250	250	250	363					
B-A08	250	250	250	725					

B)	AC9ET1	AC8ET2	AC7ET3	AC6ET4	AC5ET5	AC4ET6	AC3ET7	AC2ET8	AC1ET9
B-A47	500	500	500	500	500	500	500	500	500
B-A48	250	250	250	250	250	500	500	500	500
B-A49	250	250	250	250	250	500	500	500	500
B-A51	500	500	500	500	500	500	500	500	500
B-A22	250	250	250	250	250	250	250	250	250
B-A23	250	125	125	125	125	250	250	250	250
B-A24	250	250	250	250	250	250	250	250	250
B-A08	250	250	250	250	250	250	250	250	250

C)	EA9ET1	EA8ET2	EA7ET3	EA6ET4	EA5ET5	EA4ET6	EA3ET7	EA2ET8	EA1ET9
B-A47	500	500	500	500	500	500	500	500	500
B-A48	250	250	250	250	250	500	500	500	500
B-A49	250	250	250	250	250	500	500	500	500
B-A51	250	500	500	500	500	500	500	500	500
B-A22	250	250	250	250	250	250	250	250	250
B-A23	250	250	250	250	250	250	250	250	250
B-A24	250	250	250	250	250	250	250	250	250
B-A08	250	250	250	250	250	250	250	250	250

D)	EA9AC1	EA8AC2	EA7AC3	EA6AC4	EA5AC5	EA4AC6	EA3AC7	EA2AC8	EA1AC9
B-A47	500	500	500	500	500	500	500	500	500
B-A48	500	500	500	500	500	500	500	500	500
B-A49	500	500	500	500	500	500	500	500	500
B-A51	500	500	500	500	500	500	500	500	500
B-A22	500	250	250	250	250	250	250	250	250
B-A23	500	250	250	250	250	250	250	250	250
B-A24	500	250	250	250	250	250	250	250	250
B-A08	500	500	250	250	250	250	250	250	250

10 Example 13

In this Example, we illustrate antimicrobial activities of a plant mixes prepared according to the previously described Method B, where the added carbohydrate is chosen among sucrose, dextrose, maltose and galactose.

The M[19P]-576, M[5P]-1 and M[5P]-2 dried plant mix were composed following Table 13-1 and placed in a 500mL vial and thoroughly mixed by manual flipping and rolling. The dried herbs were obtained from “Herboristerie Cailleau” (Chemillé, France), except PC49 which was obtained from Siccarapam (Aubiat, France).

5 **Table 13-1: list of dried plant composing: A) the M[20P]-576 mix, B) the M[5P]-1 mix and C), the M[5P]-2 mix used in Example 13**

A)																
Dried plant	PA11	PA12	PA13	PA14	PA20	PA21	PA22	PA29	PA39	PB12	PB60	PC20	PC26	PC33	PC37	PC49
Weight (g)	29,7	10,5	41,5	22,9	19,8	19,9	15,0	27,5	15,0	17,8	15,0	17,1	14,4	13,6	11,3	9,3

B)						C)					
Dried plant	PA12	PA13	PA21	PA22	PB12	Dried plant	PA11	PA13	PA21	PA22	PB12
Weight (g)	34.1	59.2	84.1	47.5	75.8	Weight (g)	56.9	54	76.9	43.4	69.4

10 For each dried plant mix, extracts were prepared according to the previously described Method B, where the spoons of dried plants were replaced by 2g of the plant mix of Example 13, using 40mL of water and the amounts of carbohydrate indicated in Table13-3 for water extraction and including the decantation step before filtration.

15 The dry residue content of the plant mixes without carbohydrate was determined separately, by weighting the lyophilisation residue obtained with a lyophilisation device (FreeZone™ 2.5, Labconco USA, with RZ6 Vacuubrand, Germany) of a frozen 40mL extract prepared according to the previously described Method B, where the spoons of dried plants were replaced by 2g of the plant mix of Example 13, using 40mL of water for water extraction and including the decantation step before filtration. These dry residue contents were used to deduce the specific activities of the different extracts, where the carbohydrate is considered as an adjuvant and not accounted as part of the active ingredient.

20 The biological activity of each extracted mix was determined according to the previously described Method C by using the strains indicted in Table 13-2, where the dilution of the extracted mix was: 1:20, 1:40, 1:80, 1:160. The minimal inhibiting concentration (MIC) was noted by visual observation of the wells as the lowest concentration where no growth is observed.

Table 13-2: list of the bacterial strains used in the Example 13

<i>Staphylococcus aureus</i>		<i>Staphylococcus epidermidis</i>	
NCTC 12493™	B-A47	ATCC® 12228™	B-A22
ATCC® 33591™	B-A48	ATCC® 700296™	B-A23
ATCC® 33592™	B-A49	ATCC® 49461™	B-A24
NCTC 43300™	B-A51	ATCC® 14990™	B-A08

The results are compiled in tables 13-4.

Table 13-3: MIC in mg/L of the extracted mixes using sucrose, lactose, dextrose, maltose or galactose for: A) the M[20P]-576 mix, B) the M[5P]-1 mix and C), the M[5P]-2 mix

A)	Sucrose	Lactose	Dextrose	Maltose	Galactose
----	---------	---------	----------	---------	-----------

M[20P]-576	100g/L	100g/L	50g/L	25g/L	100g/L	50g/L	25g/L	100g/L	50g/L	25g/L	100g/L	50g/L	25g/L
B-A47	725	>725	>725	725	>725	>725	>725	>725	725	725	725	725	725
B-A48	725	>725	>725	725	>725	>725	>725	725	725	725	725	725	725
B-A49	725	>725	>725	725	>725	>725	>725	725	725	725	725	725	725
B-A51	725	>725	>725	725	>725	>725	>725	725	725	725	725	725	725
B-A22	363	725	725	725	725	725	363	363	363	363	363	363	363
B-A23	363	725	725	363	725	725	363	363	363	363	363	363	363
B-A24	363	725	725	725	725	725	363	363	363	363	363	363	363
B-A08	725	725	725	725	725	725	725	363	363	363	363	363	363

B) M[5P]-1	Sucrose 100g/L	Lactose 100g/L 50g/L 25g/L			Dextrose 100g/L 50g/L 25g/L			Maltose 100g/L 50g/L 25g/L			Galactose 100g/L 50g/L 25g/L		
B-A47	327	327	327	327	327	327	327	327	327	327	327	327	327
B-A48	327	327	327	327	327	327	327	327	327	327	327	327	327
B-A49	327	327	327	327	327	327	327	327	327	327	327	327	653
B-A47	327	327	327	327	327	327	327	327	327	327	327	327	327
B-A22	327	327	327	327	327	327	327	163	163	163	163	327	327
B-A23	163	163	327	163	163	163	327	163	163	163	163	327	163
B-A24	327	327	327	327	327	327	327	163	163	163	163	327	163
B-A08	327	327	327	327	327	327	327	163	327	327	163	327	327

C) M[5P]-2	Sucrose 100g/L	Lactose 100g/L 50g/L 25g/L			Dextrose 100g/L 50g/L 25g/L			Maltose 100g/L 50g/L 25g/L			Galactose 100g/L 50g/L 25g/L		
B-A47	344	344	344	344	344	344	344	344	344	344	344	344	688
B-A48	344	344	344	344	344	344	344	344	344	344	344	344	344
B-A49	344	344	344	344	344	344	344	344	344	344	344	344	688
B-A51	344	344	344	344	344	344	344	688	344	688	344	344	688
B-A22	344	344	344	344	344	344	344	344	344	344	344	344	344
B-A23	344	344	344	344	344	344	344	344	344	344	344	344	344
B-A24	344	344	344	344	344	344	344	344	344	344	344	344	344
B-A08	344	344	344	344	344	344	344	344	344	344	344	344	344

5

Example 14

In this Example, we illustrate the antimicrobial activity enhancement of the M[20P]-576 herbal mix extract towards commonly used antibiotics.

10 **Table 14-1: bacterial species and strains used in Example 14**

Strain full name	Strain code
<i>Staphylococcus aureus subsp. aureus</i> ATCC® 25923™	B-A17
<i>Staphylococcus aureus subsp. aureus</i> ATCC® 29213™	B-A21

List 14-1: herbal compositions of processed sample used in Example 14:

M[19P]-576: PA11; PA12; PA13; PA14; PA20; PA21; PA22; PA29; PA39; PB12; PB60; PC20; PC26; PC33; PC37.

The dried plants of pharmaceutical grade were obtained from “Herboristerie Cailleau” (Chemillé, France).

Herbal compositions M[19P]-576 was prepared by weighting each dried herb as indicated in Table 14-2

5 **Table 14-2: weight of dried plant (mg) used in the herbal compositions of M[19P]-576**

	PA11	PA12	PA13	PA14	PA20	PA21	PA22	PA29	PA39	PB12	PB60	PC20	PC26	PC33	PC37
M[19P]-576	409	145	573	316	277	271	207	380	207	247	206	236	198	187	154

Starting from the herbal composition, the corresponding water extracts (or processed samples) was prepared according to the previously described Method B using 2.06 g of the above herbal composition with 40mL water with 100g/L sucrose for water extraction and including the decantation step before
 10 filtration. In an independent water extract preparation without sucrose, it has been measured that 2.06g/40mL of herbal composition corresponds to 14.5g/L as deduced from the weight of the lyophilized dry herbal extract.

The biological activity of each processed sample was determined according to the previously described Method C by using the following strains at the indicated final dilutions of processed samples is given
 15 in Table 14-3. In Example 14 we calculate the growth of the bacteria in presence of antibiotic and/or herbal mix relative to the growth in the absence of both antibiotic and herbal mix extract.

The bacteria planktonic growth inhibition percentage average is given in Table 14-3.

It has to be noted that the presence of the herbal extract M[19P]-576 has three effects at the concentration used: 1) it reduces the growth of the bacteria at low concentration and absence of
 20 antibiotic, 2) it reduces the increase of bacterial growth induced by low antibiotic concentration relative to the growth in the absence of antibiotics, and 3) it reduces the minimal inhibitory concentration (MIC), of, respectively, vancomycin for B-A17 and of ampicillin and penicillin for B-A21, at M[20C]-576 concentration starting at, respectively, 90.6µg/mL, 181.3µg/mL, and 90.6µg/mL.

25 **Table 14-3: planktonic growth in the presence of different extract concentrations and antibiotic concentrations relative to the growth of the bacteria in the absence of both extract and antibiotic, on two different strains**

	Strain	M[20C]-576 (µg/mL)	Antibiotic concentration (µg/mL)											
			0	3.91	7.81	15.63	31.25	62.50	125	250	500	1000	2000	4000
Ampicillin	B-A17	0	100%	176%	176%	148%	131%	112%	2%	0%	0%	0%	0%	-1%
		90.6	57%	64%	76%	75%	58%	18%	1%	0%	0%	0%	0%	0%
		181.3	54%	52%	53%	52%	31%	7%	-4%	-4%	-4%	-4%	-5%	-1%
	B-A21	0	100%	153%	132%	131%	132%	116%	150%	113%	112%	106%	-1%	0%
		90.6	64%	108%	83%	82%	82%	74%	83%	75%	83%	0%	0%	0%
		181.3	42%	44%	63%	70%	69%	70%	62%	62%	61%	-2%	-3%	-2%
Penicillin	B-A17	0	100%	137%	138%	3%	1%	0%	-1%	0%	0%	0%	0%	0%
		90.6	53%	39%	25%	1%	0%	0%	0%	0%	0%	0%	0%	0%
		181.3	51%	23%	5%	-3%	-3%	-3%	-3%	-2%	-3%	-4%	-4%	-2%
	B-	0	100%	135%	116%	115%	113%	108%	106%	101%	0%	0%	0%	0%

		90.6	65%	104%	83%	83%	84%	80%	74%	26%	0%	0%	0%	0%
		181.3	44%	62%	43%	70%	61%	60%	45%	-3%	-3%	-3%	-3%	-3%
Vancomycin	B-A17	0	100%	135%	118%	114%	117%	115%	116%	118%	128%	117%	0%	0%
		90.6	62%	65%	66%	65%	74%	70%	67%	66%	63%	0%	0%	0%
		181.3	42%	40%	41%	53%	55%	55%	54%	46%	47%	-3%	-3%	-2%
	B-A21	0	100%	118%	117%	113%	117%	110%	102%	102%	102%	0%	0%	0%
		90.6	45%	102%	82%	88%	77%	76%	77%	74%	67%	0%	0%	0%
		181.3	58%	48%	63%	57%	60%	57%	41%	57%	32%	-2%	-3%	0%

Example 15

In this Example, we illustrate how to adapt the temperature in Method B in order to cope with industrial constraints.

- 5 The dried plants of organic grade were obtained from “Herboristerie Cailleau” (Chemillé, France).

Herbal compositions M[5P]-1 was prepared by weighting each dried herb as indicated in Table 15-1

Table 15-1: weight of dried plant (mg) used in the herbal compositions of M[5P]-1

	PA12	PA13	PA21	PA22	PB12
M[5P]-1	395.2	688.8	978.9	552.5	884.6

- 10 A first extract M[5P]-1-A was prepared by incorporating 120g of the herbal composition into 600 mL of water contained in a 2L beaker (Schott, Germany). A second extract M[5P]-1-B was prepared by placing 120g of the herbal composition into a 25x30cm organic grade cotton bag (Ecobags, USA), and the filled bag was placed into 600mL of water contained into a 2L beaker (Schott, Germany). A third extract M[5P]-1-C was prepared by twice placing 60g of the herbal composition into a 25x30cm organic grade cotton bag (Ecobags, USA), and the two filled bags were placed into 600mL of water contained into a single 2L beaker (Schott, Germany). Each, respectively, M[5P]-1-A, M[5P]-1-B and M[5P]-1-C, preparation was left for maceration for 3 hours at room temperature and then placed in its beaker into an autoclave (VWR Vapour Line Eco 25, USA) for 30minutes at, respectively, 121°C, 121°C and 134°C. For each of M[5P]-1-B and M[5P]-1-C, the cotton bags were pressed using a manually operated 5L fruit press (Brouwland, Belgium) and the recovered liquid was placed back into the 2L beaker. Each beaker was left for decantation and the necessary amount of supernatant liquid was used to measure its biological activity.

Table 15-2: bacterial species and strains used in Example 15

Strain full name	Strain code
<i>Staphylococcus aureus subsp. aureus</i> ATCC® 25923™	B-A17
<i>Staphylococcus aureus subsp. aureus</i> ATCC® 29213™	B-A21
<i>Staphylococcus aureus subsp. aureus</i> ATCC® 33592™	B-A49

- 25 The biological activity of each processed sample was determined according to the previously described Method C by using the strains listed in Table 15-2 at final dilutions of 1:40, 1:80, 1:160 and 1:320. The dilution corresponding to the minimal inhibiting concentration (MIC) was noted by visual observation of the wells as the lowest concentration where no growth is observed and is reported into Table 15-3

Table 15-3: dilution corresponding to the minimal inhibiting concentration (MIC) of the extracts realized with different industrial details and temperature

	M[5P]-1-A at 121°	M[5P]-1-B at 121°	M[5P]-1-C at 134°C
B-A17	1/80	1/40	1/80
B-A21	1/80	1/40	1/80
B-A49	1/80	1/40	1/160

It is to be noted that introducing the cotton bag to ease the industrial process reduced slightly the activity of the extract as compared to the unsatisfactory process without bag. Increasing the temperature of the autoclave during the hot extraction phase allowed to recover the full activity.

CLAIMS

- 1) An extract of an herbal composition, comprising at least two dried plants selected among *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi* and at least one plant selected among *Vitis vinifera* var. *tinctoria*, *Eugenia caryophyllus* and *Desmodium adscendens*.
- 2) The extract according to claim 1, wherein the herbal composition further comprises one or more additional dried plants selected among: *Achillea millefolium*, *Acorus calamus*, *Agrimonia eupatoria*, *Agropyrum repens*, *Alchemilla vulgaris*, *Alkanna tinctoria*, *Althaea officinalis*, *Anethum graveolens*, *Angelica archangelica*, *Arbutus unedo*, *Arnica montana*, *Artemisia pontica*, *Artemisia vulgaris*, *Asparagus officinalis*, *Asperula odorata*, *Betula pendula*, *Borrigo officinalis*, *Buxus sempervirens*, *Calamintha officinalis*, *Calendula officinalis*, *Calluna vulgaris*, *Carum carvi*, *Cassia angustifolia*, *Centaurea cyanus*, *Centaureum erythraea*, *Centella asiatica*, *Cetraria islandica*, *Chamaemelum nobile*, *Chamomilla recutita*, *Chrysanthellum americanum*, *Cichorium endivia*, *Cichorium intybus*, *Cinnamomum zeylanicum*, *Citrus aurantium*, *Combretum micranthum*, *Crataegus oxyacantha*, *Cuminum cyminum*, *Cupressus sempervirens*, *Curcuma zedoaria*, *Cynara scolymus*, *Cytisus scoparius*, *Desmodium adscendens*, *Elettaria cardamomum*, *Eleutherococcus senticosus*, *Epilobium parviflorum*, *Erysimum officinale*, *Eucalyptus globulus*, *Eugenia caryophyllus*, *Eupatorium cannabinum*, *Foeniculum vulgare*, *Fraxinus excelsior*, *Fucus vesiculosus*, *Fumaria officinalis*, *Galium odoratum*, *Gentiana lutea*, *Geranium robertianum*, *Ginkgo biloba*, *Glechoma hederacea*, *Glycyrrhiza glabra*, *Handroanthus impetiginosus*, *Harpagophytum procumbens*, *Hieracium pilosella*, *Humulus lupulus*, *Hypericum perforatum*, *Hyssopus officinalis*, *Illicium verum*, *Inula helenium*, *Juglans regia*, *Juniperus communis*, *Lamium album*, *Lavandula angustifolia*, *Levisticum officinale*, *Lippia citriodora*, *Lotus corniculatus*, *Lythrum salicaria*, *Malva sylvestris*, *Marrubium vulgare*, *Medicago sativa*, *Melissa officinalis*, *Mentha × piperita*, *Morus nigra*, *Myrtus communis*, *Olea europaea*, *Origanum majorana*, *Panax ginseng*, *Papaver rhoeas*, *Parietaria officinalis*, *Passiflora incarnata*, *Petroselinum crispum*, *Peumus boldus*, *Phaseolus vulgaris*, *Pimpinella anisum*, *Plantago lanceolata*, *Plantago ovata*, *Potentilla anserina*, *Quercus robur*, *Rhamnus frangula*, *Rheum palmatum*, *Rosa centifolia*, *Rosmarinus officinalis*, *Rubia tinctorum*, *Rubus idaeus*, *Salix alba*, *Salvia officinalis*, *Sambucus nigra*, *Satureja montana*, *Silybum marianum*, *Solanum dulcamara*, *Tabebuia impetiginosa*, *Tanacetum vulgare*, *Taraxacum officinalis*, *Thymus serpyllum*, *Thymus vulgaris*, *Tilia tomentosa*, *Tilia cordata*, *Trigonella foenum-graecum*, *Tussilago farfara*, *Vaccinium myrtillus*, *Valeriana officinalis*, *Verbascum thapsus*, *Verbena officinalis*, *Viscum album*, *Zea mays* and *Zingiber officinale*.
- 3) The extract according to any preceding claim, wherein the herbal composition comprises from three to seven dried plants selected among *Filipendula ulmaria*, *Camellia sinensis*, *Arctostaphylos uva-ursi*, *Rheum palmatum*, *Rosmarinus officinalis*, *Vitis vinifera tinctoria*, *Desmodium adscendes*, *Eugenia caryophyllus* and *Eucalyptus globulus*, wherein at least two of such plants are selected among *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi* and at least one of such plants is selected among *Vitis vinifera tinctoria*, *Desmodium adscendes* and *Eugenia caryophyllus*.
- 4) The extract according to any preceding claim, wherein the herbal composition comprises the following dried plants: *Rosmarinus officinalis*, *Filipendula ulmaria*, *Camellia sinensis*, *Vitis vinifera tinctoria* and *Arctostaphylos uva-ursi*.
- 5) The extract according to any preceding claim, wherein the herbal composition comprises the following dried plants: *Rheum palmatum*, *Filipendula ulmaria*, *Camellia sinensis*, *Vitis vinifera tinctoria* and *Arctostaphylos uva-ursi*.

- 6) The extract according to any preceding claim, wherein the herbal composition comprises the following dried plants: *Rheum palmatum*, *Rosmarinus officinalis*, *Filipendula ulmaria*, *Camellia sinensis*, *Vitis vinifera tinctoria*, *Eugenia caryophyllus* and *Arctostaphylos uva-ursi*.
- 7) The extract according to any preceding claim, wherein the herbal composition comprises the following dried plants: *Rheum palmatum*, *Rosmarinus officinalis*, *Filipendula ulmaria*, *Camellia sinensis*, *Vitis vinifera tinctoria*, *Eucalyptus globulus*, *Arctostaphylos uva-ursi*, *Mentha spicata* and *Rubia tinctorium*.
- 8) The extract according to any preceding claim, wherein the herbal composition comprises at least fourteen or fifteen dried plants selected among the following sixteen: *Rheum palmatum*, *Rosmarinus officinalis*, *Filipendula ulmaria*, *Satureja montana*, *Valeriana officinalis*, *Camellia sinensis*, *Vitis vinifera tinctoria*, *Fucus vesiculosus*, *Foeniculum vulgare*, *Arctostaphylos uva-ursi*, *Arbutus unedo*, *Eugenia caryophyllus*, *Juniperus communis*, *Combretum micranthum*, *Lippia citrodora* and *Tanacetum vulgare*.
- 9) The extract according to any preceding claim, wherein the herbal composition comprises the following dried plants: *Rheum palmatum*, *Rosmarinus officinalis*, *Filipendula ulmaria*, *Satureja montana*, *Valeriana officinalis*, *Camellia sinensis*, *Vitis vinifera tinctoria*, *Fucus vesiculosus*, *Foeniculum vulgare*, *Arctostaphylos uva-ursi*, *Arbutus unedo*, *Eugenia caryophyllus*, *Juniperus communis*, *Combretum micranthum* and *Lippia citrodora*.
- 10) The extract according to any preceding claim, wherein the herbal composition comprises the following dried plants: *Rheum palmatum*, *Rosmarinus officinalis*, *Filipendula ulmaria*, *Satureja montana*, *Valeriana officinalis*, *Camellia sinensis*, *Vitis vinifera tinctoria*, *Fucus vesiculosus*, *Foeniculum vulgare*, *Arctostaphylos uva-ursi*, *Arbutus unedo*, *Eugenia caryophyllus*, *Juniperus communis*, *Combretum micranthum*, *Lippia citrodora* and *Tanacetum vulgare*.
- 11) A lyophilisate of the extract according to any of claims 1 to 10.
- 12) A food supplement, a nutraceutical, pharmaceutical, veterinary or cosmetic composition or a functional food or a food additive comprising the extract according to any of claims 1 to 10, or the lyophilisate of claim 11, as active ingredient, together with at least one excipient.
- 13) The food supplement, the nutraceutical, pharmaceutical, veterinary or cosmetic composition, the functional food, or the food additive according to claim 12, formulated for oral or topic administration.
- 14) The extract according to any of claims 1 to 10, the lyophilisate of claim 11 or the pharmaceutical or veterinary composition according to claims 12 or 13, for use as an antimicrobial and/or antibiofilm agent.
- 15) The extract according to any of claims 1 to 10, the lyophilisate of claim 11 or the pharmaceutical or veterinary composition according to claims 12 or 13, for use as an antimicrobial and/or antibiofilm agent in the treatment, prevention or diagnosis of microbial infections.
- 16) The extract, the lyophilisate or the pharmaceutical or veterinary composition according to claim 15, wherein the infections are caused by *Escherichia*, *Klebsiella*, *Listeria*, *Pseudomonas*, *Salmonella*, *Streptococcus*, *Candida* or *Staphylococcus*.
- 17) The extract, the lyophilisate or the pharmaceutical or veterinary composition according to claim 15, wherein the infections are caused by *Staphylococcus aureus*, *Staphylococcus epidermidis* or *Staphylococcus pseudintermedius*.

- 18) A process for preparing the extract according to any of claims 1 to 10, comprising a) chopping or grinding separately or in a mixture at least two single dried plants to obtain corresponding plant powders; b) mixing at least two different plant powders to obtain an herbal composition; c) adding an extraction solvent to said herbal composition to obtain a corresponding liquid preparation (or extract); d) incubating said liquid preparation at room temperature for a time ranging between 5 and 15 minutes; optionally, e) heating said liquid preparation at a temperature ranging from 60°C to 134°C for a time ranging from 5 to 60 minutes; optionally f) centrifuging, collecting the supernatant and filtering the collected supernatant; and, optionally, g) concentrating and/or drying or freeze drying the liquid preparation obtained from step e) or f).
- 19) The process according to claim 18, where the extraction solvent is water.
- 20) The process according to claim 19, where the water added in step (c) comprises a sugar dissolved in it at a concentration varying from 1 to 100 g/l, optionally from 50 to 75 g/l.
- 21) An herbal composition comprising at least two dried plants selected among *Filipendula ulmaria*, *Camellia sinensis* and *Arctostaphylos uva-ursi* and at least one plant selected among *Vitis vinifera* var. *tinctoria*, *Eugenia caryophyllus* and *Desmodium adscendens*, as intermediate in the process for preparing the extract according to any of claims 1 to 10.
- 22) The herbal composition according to claim 21, further comprising one or more additional dried plants selected among: *Achillea millefolium*, *Acorus calamus*, *Agrimonia eupatoria*, *Agropyrum repens*, *Alchemilla vulgaris*, *Alkanna tinctoria*, *Althaea officinalis*, *Anethum graveolens*, *Angelica archangelica*, *Arbutus unedo*, *Arnica montana*, *Artemisia pontica*, *Artemisia vulgaris*, *Asparagus officinalis*, *Asperula odorata*, *Betula pendula*, *Borrago officinalis*, *Buxus sempervirens*, *Calamintha officinalis*, *Calendula officinalis*, *Calluna vulgaris*, *Carum carvi*, *Cassia angustifolia*, *Centaurea cyanus*, *Centaurium erythraea*, *Centella asiatica*, *Cetraria islandica*, *Chamaemelum nobile*, *Chamomilla recutita*, *Chrysanthellum americanum*, *Cichorium endivia*, *Cichorium intybus*, *Cinnamomum zeylanicum*, *Citrus aurantium*, *Combretum micranthum*, *Crataegus oxyacantha*, *Cuminum cyminum*, *Cupressus sempervirens*, *Curcuma zedoaria*, *Cynara scolymus*, *Cytisus scoparius*, *Desmodium adscendens*, *Elettaria cardamomum*, *Eleutherococcus senticosus*, *Epilobium parviflorum*, *Erysimum officinale*, *Eucalyptus globulus*, *Eugenia caryophyllus*, *Eupatorium cannabinum*, *Foeniculum vulgare*, *Fraxinus excelsior*, *Fucus vesiculosus*, *Fumaria officinalis*, *Galium odoratum*, *Gentiana lutea*, *Geranium robertianum*, *Ginkgo biloba*, *Glechoma hederacea*, *Glycyrrhiza glabra*, *Handroanthus impetiginosus*, *Harpagophytum procumbens*, *Hieracium pilosella*, *Humulus lupulus*, *Hypericum perforatum*, *Hyssopus officinalis*, *Illicium verum*, *Inula helenium*, *Juglans regia*, *Juniperus communis*, *Lamium album*, *Lavandula angustifolia*, *Levisticum officinale*, *Lippia citriodora*, *Lotus comiculatus*, *Lythrum salicaria*, *Malva sylvestris*, *Marrubium vulgare*, *Medicago sativa*, *Melissa officinalis*, *Mentha x piperita*, *Morus nigra*, *Myrtus communis*, *Olea europaea*, *Origanum majorana*, *Panax ginseng*, *Popover rhoeas*, *Parietaria officinalis*, *Passiflora incarnata*, *Petroselinum crispum*, *Peumus boldus*, *Phaseolus vulgaris*, *Pimpinella anisum*, *Plantago lanceolata*, *Plantago ovata*, *Potentilla anserina*, *Quercus robur*, *Rhamnus frangula*, *Rheum palmatum*, *Rosa centifolia*, *Rosmarinus officinalis*, *Rubia tinctorum*, *Rubus idaeus*, *Salix alba*, *Salvia officinalis*, *Sambucus nigra*, *Satureja montana*, *Silybum marianum*, *Solanum dulcamara*, *Tabebuia impetiginosa*, *Tanacetum vulgare*, *Taraxacum officinalis*, *Thymus serpyllum*, *Thymus vulgaris*, *Tilia tomentosa*, *Tilia cordata*, *Trigonella foenumgraecum*, *Tussilago farfara*, *Vaccinium myrtillus*, *Valeriana officinalis*, *Verbascum thapsus*, *Verbena officinalis*, *Viscum album*, *Vitis vinifera* var. *tinctoria*, *Zea mays* and *Zingiber officinale*.
- 23) A process for preparing the food supplement, the nutraceutical, pharmaceutical, veterinary or cosmetic composition, the functional food, or the food additive according to claims 12 or 13, comprising mixing the extract according to any of claims 1 to 10, or the lyophilisate of claim 11, with at least one excipient.