




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C10G 71/00, 35/06, 35/085, 35/095, C10C 5/00		A1	(11) International Publication Number: WO 99/41333
			(43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/US99/02986		(74) Agents: SCUORZO, Linda, M. et al.; Exxon Research and Engineering Company, P.O. Box 390, Florham Park, NJ 07932-0390 (US).	
(22) International Filing Date: 12 February 1999 (12.02.99)			
(30) Priority Data: 60/074,579 13 February 1998 (13.02.98) US		(81) Designated States: AU, CA, JP, SG, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(71) Applicant (for all designated States except US): EXXON RESEARCH AND ENGINEERING COMPANY [US/US]; 180 Park Avenue, P.O. Box 390, Florham Park, NJ 07932-0390 (US).		Published With international search report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): MURPHY, William, John [-/US]; 16445 Coliseum Avenue, Baton Rouge, LA 70816 (US). McVICKER, Gary, Brice [-/US]; 65 Main Street, Califon, NJ 07830 (US). HANTZER, Sylvain [-/US]; 37620 Seven Oaks Avenue, Prairieville, LA 70769 (US). CODY, Ian, Alfred [-/US]; 7942C Wrenwood Boulevard, Baton Rouge, LA 70809 (US). SOLED, Stuart, Leon [-/US]; 21 Cooks Cross Road, Pittstown, NJ 08867 (US). LINEK, Sandra, Jean [-/US]; 5202 Claycut Road, Baton Rouge, LA 70806 (US). BAUMGARTNER, Joseph, Ernest [-/US]; 179 Raritan River Road, Califon, NJ 07830 (US).			

(54) Title: PROCESS FOR MAKING A LUBE BASESTOCK



(57) Abstract

A method for hydroisomerizing a waxy feed to favor one of VI or yield is described. The method uses a unitized pellet powder catalyst comprising a metal hydrogenation component, a first catalytic component selected from 8, 10 and 12 ring molecular sieves and mixtures thereof, and a second amorphous isomerization component which is an amorphous inorganic oxide. Importantly, the ratio of feed and second components are present in a predetermined ratio to provide a preselected acidity favoring one of VI or yield resulting from the hydroisomerization.