BIOLOGICALLY ACTIVE AQUEOUS FRACTION OF AN EXTRACT OBTAINED FROM A MANGROOVE PLANT *SALVADORA PERSICA L.*

Technical Field

The invention relates to biologically active extracts obtained from the plant *Salvadora persica Linneaus* 1753. The invention also provides a process for obtaining the crude extract. Further the invention provides pharmaceutical compositions, exhibiting biological activity, especially tocolytic activity.

Background Art

The associated mangrove plant, *Salvadora persica Linneaus*, 1753, belongs to the order Salvadoracea and are shrubs or small trees with white flowers frequent in degraded mangrove swamps and saline banks all over the west coast of India. Large numbers of marine plants have been examined for bioactive substances. Nazarine, F; Anita F; Rataboli, P.V.; Diniz D'Souza, R. S and Dhume; V.G., 1998 in Indian Journal of marine Sciences, 27: 499-501 have reported promising pharmacological activities in marine organisms from Indian waters.

There are several patents available from all over the world related to processes and compounds from nature for various purposes. Kwak; Wie-jong; Han; Chang-kyun; kum; Hwan-su; An; Jae-suk; Kum; Taek-soo, patented process of extracting and purifying biologically effective ingredients from combined medicinal plant and their extract composition (United States Patent No.5,910,307 published on June 8, 1999). D'Amelio; Frank S; Mirhom; Youssef W. disclosed therapeutic composition and method for treating skin using extract from *Centipeda cunninghami* plant in US Patent No.5,804,206 published on September 8, 1998. Zimmerman; Richard C.; Alber; Randall S.; Todd; James S.; Crews; Philip, isolated compound from the methanolic extract of the eelgrass *Zostrea marina* having significant antifouling aquatic properties (US Patent No.5,607,741 published on March 4, 1997).

Betulinic acid which is prepared from the compound Betulin has many pharmaceutical potentials. Pezzuto John M; Kim; Darrick S.H.L. disclosed methods of manufacturing betulinic acid from betulin (US Patent No.5,804,575 published on 8 September, 1998). The betulinic acid is intensively investigated as a potential therapeutic agent for a variety of diseases. Pezzuto; John, M; Das Gupta, Tapas K; Schmidt; Mary Lou; Kuzmanoff; Konrad Marc; Ling Indeck; Lydia and Kim; Darrick, S.H.L. (US Patent No.5,962,527 dated October 5, 1999). Pisha E, Chai H, Lee IS, Chagwedera TE, Framsworth NR, Cordell GA, Beecher CW, Fong H H, Kinghorn AD, Brown DM, in Nature medicine, 1 pages 1046-

1051 (1995) discloses that betulinic acid has an unexpected selective anti-tumour activity against human melanoma e.g. MEL-1, MEL-2 and MEL-4. In addition Fujioka T, Kashiwada Y, Kilkuskie RE, Cosentino LM, Ballas LM, Jiang JB, Janzen WP, Chen IS, Lee KH. J.Nat.Prod., 57 (2) pages 243-247 (1994) discloses that betulinic acid has anti-HIV activity in H9 lymphocytic cells. These inventors mentioned that the researches directed t betulinic acid as therapeutic agent are hindered because the betulinic acid is available in very limited quantities and at a very high cost. Ramadoss, Sunder, Jaggi, Manu; Siddiqui; Mohammad Jamshed Ahmed in US Patent No.6,048,847 published on April 11, 2000 describes uses of betulinic acid and its derivatives for inhibiting cancer growth and a method of monitoring this. Kang; Raphael K.L.; Zyzak; Li Li; nakatsu; Tetsuo published flavoured product additives in US Patent No.5,948,460, dated 7 September, 1999 in which Ursolic acid was one of the compound amongst a group of three compounds which was added to a flavoured product to reduce aftertaste in the product and enhance its sweetness like in a diet drink. It was also used as a constituent in a preparation for inhibition of skin Tumorigenesis.

Herman, S (US Patent No.5,190,979 published on 2 March, 1993) disclosed lupeol also as a compound which can make pharmacologically active terpene ozonides which have medicinal value.

Objects of the invention

The main object of the present invention is to provide a process for obtaining a crude extract from stem, leaves, and flowers of *Salvadora persica*, a commonly available shrub in mangrove swamps and screening to study its bioactivity.

Another object of the invention is to isolate naturally occurring compounds from the plant *Salvadora persica* and identify their molecular weights, molecular formulae, melting points and their structural formulae, which will be helpful in chemical synthesis of these compounds.

Yet another object of the invention is pharmacological screening of crude extract, its fractions and purified compounds to check that the activities shown by the crude extract and fractions is maintained throughout.

Another object of the invention is to provide pharmaceutical compositions containing extracts from the plant *Salvadora persica* and exhibiting biological activity, especially, tocolytic activity.

Summary of the invention

The present invention seeks to overcome the drawbacks inherent in the prior art by

providing the highly efficient and selective means for processing of active crude extract its fractionation, isolation and purification of the active compounds. Further the invention provides pharmaceutical compositions containing the extract obtained from the plant *Salvadora persica* and useful in relieving pains in the uterine muscles.

Detailed description of the invention

The present invention discloses for the first time the methods of isolation, purification and pharmacological screening of all these above said commercially important compounds from a commonly available plant from mangrove swamps of the west coast of India. The said plant identified as *Salvadora persica,* is an associated mangrove plant. It is a shrub and the twigs with leaves of flowers can be hand picked. Even the crude extract and fractions exhibit therapeutic value. The important point is that the biological activity shown by the extract is maintained in the purified compounds such as Methyl palmitate and Betulin.

This disclosure points towards future potential clinical uses of the extract and fractions for treatment of diseases such as smooth muscle relaxant, bronchial asthma, renal colics and prevention of premature delivery. It further relates to the use of Methyl palmitate and Betulin in motion sickness, abdominal cramps.

The invention provides methods whereby biologically active crude extract of an associated mangrove plant identified as *Salvadora persica Linneaus* 1753 is prepared. *Salvadora persica Linneaus* (Salvadoraceae) are shrubs or small trees with white flowers frequent in degraded mangrove swamps and saline banks all over the west coast of India; west Asia. The process disclosed in the invention further relates to the extraction, fractionation and purification of active constituent metabolites of the said plant. The invention is also concerned with the spectral identification of the compounds such as β-amyrin (non-steroidalpolycyclic triterpene), Betulin, Ursolic acid (triterpenic acid), Methyl palmitate (Aliphatic Ester) and Lupeol (non-steroidalpolycylic triterpene). The invention also deals with molecular formulae, molecular weights, melting points and structural formulae of the said compounds. The invention provides a highly efficient and selective means for processing of active crude extract obtained from *Salvadora persica,* its fractionation, isolation and purification. As used herein, the terms fractionation means separating the crude extract. The term isolation and fractionation means separating the fraction into pure compounds.

The invention further relates to methods of screening pharmacological activities of the said compounds in mammalian tissues. The applicant has found that the crude extract obtained

from the plant *Salvadora persica* can be separated into two fractions, i.e., chloroform and aqueous fractions. The aqueous fraction was found to exhibit tocolytic activity, which is described in detail in the present invention.

Accordingly, the invention provides a process of extracting and purifying biologically useful molecules from an associated mangrove plant which comprises the steps of:

i) collecting and processing the plant plants of *Salvadora persica,*

ii) preparing a crude extract from the plant parts of *Salvadora persica,*

iii) testing the crude extract using methods of pharmacology,

iv) fractionating the crude extract,

v) testing the fractions using methods of pharmacology,

vi) isolating the pure compounds by column chromatography,

vii) testing the pure compounds by using methods of pharmacology, and

viii) identifying the compounds by spectroscopy.

As said earlier, the aqueous fraction of the crude extract obtained from the mangrove plant *Salvadora persica* showed tocolytic activity, while the remaining 2 fractions (butanol and petroleum ether) were inactive.

The invention also provides the identification of the molecules from the spectral data. The molecular formulae of the five compounds is provided from the spectral data. The invention provides molecular weights of the molecules from EIMS. The structural formulae of the compounds is also provided from the spectral data.

Thus, a crude extract was obtained from the plant *Salvadora persica.* The crude extract was tested for its bioactivity and if found promising in terms of its pharmacological activity, it was fractionated using solvents with increasing polarity to obtain fractions such as petroleum ether, chloroform, butanol and aqueous. Each of these were also tested for their pharmacological activity.

The five compounds purified from the extract of the plant were β-amyrin (non - steroidalpolycyclic triterpene), Betulin, Ursolic acid (triterpenic acid), Methyl palmitate (Aliphatic Ester) and Lupeol (non-steroidalpolycyclic triterpene).

β-amyrin was found to be a non-steroidalpolycyclic triterpene with the following details:

Molecular formula	: $C_{30}H_{50}O$
Molecular weight	: 426
Melting point	: 160.degree C

Another molecule found was betulin having:

Molecular formula	: $C_{30}H_{50}O_2$

Molecular weight : 442
Melting point : 255 degree.C

Ursolic acid (triterpenic acid) molecule was also found in the extract. It had:

Molecular formula : $C_{30}H_{48}O_3$
Molecular weight : 456
Melting point : 292 degree.C.

Methyl palmitate (Aliphatic Ester) found in the extract had:

Molecular formula : $C_{16}H_{32}O_2$
Molecular weight : 256
Melting point : 30 degree.C

Lupeol (non-steroidalpolycyclic triterpene) found in the extract had:

Molecular formula : $C_{30}H_{50}O$
Molecular weight : 426
Melting point : 215 degree. C

One of the compounds Betulin which can be used in manufacturing of betulinic acid (US Patent No.5,804,575 published on 8 September, 1998. Betulinic acid is intensively investigated as a potential therapeutic agent for a variety of diseases.

Ursolic acid is added to a flavoured product to reduce aftertaste in the product and enhance its sweetness for example in a diet drink. It was also used as a constituent in a preparation for inhibition of skin Tumorigenesis.

Extracts of some plants show vasoconstrictor and analgesic properties and also contain triterpenoid beta.-amyrin. These compositions for inhibiting the formation of unwanted skin pigmentation combine high tyrosinase blocking capabilities with stability in cosmetic preparations, absence of significant cytotoxic effects and synergy of action (US patent 5,773,014, 5,679,393). Beta.-amyrin and lupeol are used as components for dimethylsterols in medical formulations (US Patent 4,808,574).

Methyl palmitate is compound used in making alcohols as mentioned in US patent No. 6,049,013 published on April, 2000. Lupeol can be used as a component for several remedial medicines, insect repellants, distilleries anti tumour and chemical industries (US Patent No. 4,808,574; 5,962,527; 5,908,628).

Thus, the extract of *Salvadora persica* contained several compounds as listed above. The aqueous fraction of the extract of the said plant *Salvadora persica*, was then tested for its biological activity on guinea pigs. The Applicants, to their surprise found that the aqueous fraction exhibited excellent tocolytic activity.

Accordingly, the invention provides compositions containing the aqueous fraction of the extract obtained from *Salvadora persica*, optionally with conventional additives for relieving pains in the uterine muscles. The composition may contain about 10 µgm of the extract. Thus, the aqueous fraction of the crude extract obtained from the plant *Salvadora persica* is a potential tocolytic agent. The compositions may be formulated in different physical forms, as may be required. The extract may be used as such or with conventional additives, physiologically acceptable carriers, preservatives, buffers, etc. as required.

Additionally, the invention provides a method of treating tocolytic conditions, which comprises administration of therapeutically effective amount of extract obtained from *Salvadora persica* to the subject in need thereof. The extract may be administered at a dosage level in the range of 50 µg/ml to 250µg/ml, in case of normal adults. The exact dosage will vary depending on the patient to be treated and will depend on factors such as requirements of the patient, severity of the condition being treated and the activity of the extract. The determination of optimum dosages for a particular patient is well-known to those skilled in the art.

Detailed description of the accompanying drawings

Fig 1(a) shows the mangrove plant and

Fig 1(b) shows the twig of the associated mangrove plant used .

Fig 2: shows the different fractions obtained from the crude extract of the plant *Salvadora persica.*

Fig.3: shows the structural formula of β-amyrin (non - steroidalpolycyclic triterpene)

Fig.4: Structural formula of Betulin,

Fig 5: Structural formula of Ursolic acid (triterpenic acid),

Fig.6: Structural formula of Methyl palmitate (Aliphatic Ester)

Fig.7: Structural formula of Lupeol (non-steroidalpolycyclic triterpene).

Fig.8 Fractionation chart of extract of *Salvadora persica*

The invention is described in detail and illustrated by the following examples which should not be construed as limitations on the inventive concept embodied herein.

Example 1 : This example provides the Chemicals, Reagents, Apparatus used and their sources.

Name of reagent/chemicals	Company
Aqueous methanol	Sisco Research Laboratories Pvt Ltd.
Petroleum ether	Ranbaxy Fine Chemicals Ltd.
Chloroform	Sisco Research Laboratories Pvt Ltd.

Butanol	Sisco Research Laboratories Pvt Ltd.
Ethyl acetate	Sisco Research Laboratories Pvt Ltd.
Histamine acid phosphate	Blenkinsop & Co. Ltd
Acetylcholine chloride	Hopkin & Williams Ltd.
5-Hydroxytryptamine creatine sulphate	Sigma Chemicals
Barium chloride	Apex Chemicals
Nicotine sulphate	BDH chemicals
Oxytocin	Parke Davis India Ltd.
Prostodin-PGF 2 alpha	Astra IDL Ltd.

APPARATUS:

1. Physiograph .
Company: Biodevices,
Ambala, India.

2. Force Transducer
Model No. T-305
Co.: GRASS,USA

3. Stimulator
Model SS44
Co. Biodevices, Ambala

4. Polygraph
Model 7
Co. GRASS , USA.

5. Force Transducer
Model No. FT-03
Co.: GRASS,USA

6. Organ Bath
Ambala,
India

Example 2 :

Collection of the mangrove plant *Salvadora persica* L from the coast of Goa, a state in India was along Ribandar, near the mouth of the Mandovi estuary, upstream (India). This species is ubiquitous to the coastal areas of Goa and was collected manually from the intertidal banks.

Example 3:

Processing of the collected mangroves were washed first with seawater followed by tap water. The undesired materials were sifted out while washing with tap water to get rid of the salts. The leaves, stems, and flowers of the associated mangrove plant were air dried. After drying, the plant material was cut into small pieces and immersed in the solvent 90% aqueous methanol for a week for extraction. Care was taken to ensure that these were properly soaked/dipped in the solvent and checked for putrefaction.

Example 4:

Extraction and preparation of crude extract was carried out by cold percolation method at room temperature and by solvent evaporation at a water bath (temperature 50°C) under reduced pressure. This helps in protection of any heat labile metabolite present in it. Re-extraction was done twice until the extract was concentrated under vacuum to obtain the crude extract.

Example 5

Fractionation of the crude extract :

The crude extract was partitioned into petroleum ether, chloroform, n-butanol and aqueous fractions using a separating funnel . Petroleum ether was added to the extract in the separating funnel and separated out. Next, chloroform was added to the residue, mixed well and the lower layer separated.. To the residue butanol was added and the top layer represented the butanol fraction and lower layer the aqueous fraction. Extraction of each fraction was done thrice and whenever there was emulsion sodium chloride was added for breaking the emulsion. Sodium sulphate was added to chloroform and butanol fractions to remove traces of water before concentration. All the fractions were concentrated in the same manner as the crude extract. These fractions were tested for the same pharmacological activity as the parent crude extract. Column chromatography for isolation of pure compound was done by repeated column chromatography and thin layer chromatography of the eluents. The TLC revealed compounds such as Beta-amyrin, betulin, ursolic acid and lupeol.

Example 6

To obtain the compounds Beta amyrin and betulin, separation by thin layer chromatography is carried out on 0.25mm thick silica gel plates (Qualigen). The eluent is an 90:10 (v/v) Petroleum ether/ethyl acetate mixture and the spots are developed by spraying with 5% H_2SO_4 solution and fixation by heating at 110.degree.C.

For compounds Ursolic acid and Methyl palmitate separation by thin layer

chromatography were carried out on 0.25 mm thick silica gel plates (Qualigen). The eluent is an 85:15 (v/v) Petroleum ether/ethyl acetate mixture and the spots are developed by spraying with a 5% H_2SO_4 solution and fixation by heating at 110°C.

For the compound lupeol the separations by thin layer chromatography are carried out on 0.25mm thick silica gel plates (Qualigen). The eluent is an 75:25 (v/v) Petroleum ether/ethyl acetate mixture and the spots are developed by spraying with a 5% H_2SO_4 solution and fixation by heating at 110°C.

In the present invention, the active aqueous fraction of the mangrove plant S. persica , was column chromatographed over silica gel for the isolation of the active constituent. Elutes from the column with the same TLC profile were mixed and subjected to pharmacological testing. The active subfractions were further chromatographed till active pure compounds were obtained. Spots on thin layer chromatography (TLC) were visualised by using iodine vapours and spraying with methanolic sulphuric acid.

TLC was done on glass plates (20 X 20 cms) coated with a 0.25mm layer of TLC grade silica gel (Qualigens) activated at 110° C for 1 hour before use. The active aqueous fraction was passed through XAD-column and eluents were treated, as mentioned above, for the isolation and purification of active constituent metabolites.

The five compounds were identified on the basis of spectral data obtained at the Regional Sophisticated Instrumentation Centre (RSIC) by the following spectra :

^{1}HNMR for determining the proton environment of the molecule carried out on Bruker DPX-200 MHz.

Apparatus: Bruker Spectrometer
Model: DPX
Co. Bruker

^{13}CNMR for carbon atoms Bruker DPV 300 MHz.

The compounds were identified from a comparison of their spectral data with those of similar compounds reported in literature.

Mass spectra (EIMS) electron impact Mass spectrometry for determining the molecular weights along with its fragmentation pattern were carried out on Mass spectrometer (E1/CIMS) Model D.300 JEOL,

Apparatus: Mass Spectrometer (EIMS)
Model : D-300
Co. JEOL, Japan

Example 6

Pharmacological testing of pure compounds

Standard drugs used were the following:

Histamine acid phosphate (Blenkinsop & Co. Ltd.) on ileum.

Acetylcholine chloride (Hopkin & Williams Ltd.) on ileum.

5-Hydroxytryptamine creatinine sulphate (sigma Chemicals Co.) on gastrointestinal tract.

Barium chloride (Apex Chemicals) on smooth muscle contraction.

Nicotine sulphate (BDH Chemicals) on intestine as ganglion stimulant.

Oxytocin (Parke Davis India Ltd) on uterus.

Prostodin – PGF_2 (Astra IDL Ltd) on uterus.

All other reagents used were of analytical grade.

Tyrode was used on guinea-pig ileum and de Jalon's solution was used on guinea-pig uterus.

Ringer-Locke physiological solution was used on guinea-pig atria table 3.

All other reagents used were of analytical grade.

Physiological solutions used and various parameters:

All physiological solutions were prepared fresh at the time of the experiment.

pH: The pH of the various physiological salt solutions varied between 7.3 & 7.4. At lower pH the tonus of the preparation tends to decrease and is therefore liable to alter the effect of drugs.

Temperature: In order to get consistent effects it was important to maintain the temperature of the bath solution at a specified level, because if the temperature is decreased below 37°C, the tonus of the intestine is increased, the contractions become smaller and the contraction and relaxation times increased.

Air: Air or oxygen is needed for proper functioning of the tissues. Besides, providing oxygen to the tissues, the stream of gas bubbles also stirred the bath solution thereby facilitating diffusion of drugs added to the bath.

The solution in the bath was changed frequently because prolonged aeration tends to alter the pH.

***In vitro* experiments** :

Female, virgin, guinea pigs weighing around 300 to 350 g, housed under uniform husbandry conditions (temperature 25 ± 1°C) were used. The animals were starved 24 hours prior to the experiment, only water was provided *ad libitum.*

The isolated guinea pig uterus was used to study the tocolytic activity.

For experiments upon isolated guinea pig, the bicornuate uterus was dissected out and freed of fat tissue. One horn was cut-off and kept in a shallow dish containing the physiological solution – de Jalon's fluid which was previously aerated with air. Air was preferred to oxygen as the tissue was thin and saturation was faster. Jalon's fluid comprises:

Glucose	0.5 g
Sodium chloride	9.0 g
Sodium bicarbonate	0.5 g
Potassium chloride	0.42 g
Calcium chloride	0.06 g

All were dissolved in 1000 ml water

The two ends were sutured. The lower end of the uterine strip was tied to a tissue holder and suspended in an organ bath of 10 ml capacity and the upper end being more sensitive, to the lever of the force transducer (FT 03) connected to a Grass Polygraph (Model 7). It was left to stabilize for 30 mins, renewing the physiological solution in the bath every 10 mins. The response of the uterus to different does of the extract (50 & 250 µg/ml) against standard uterine stimulants like oxytocin and PG $F_{2\alpha}$ with a contact period of 60 seconds (was recorded on the polygraph). The tocolytic effect was evaluated by the following formula:

$$\% \text{ Inhibition} = \frac{X - Y}{X} \times 100$$

Wherein: X = Height of standard contraction (mm)

Y = Height of standard contraction in presence of the extract (mm)

The relative in vitro tocolytic effect of *Salvadora persica* crude extract on guinea pig uterus is shown in the Table hereinbelow (mean of 5 readings):

EXTRACT	TISSUE	SPASMOGENS	STANDARD INDUCED CONTRACTION (mm) 48	DOSE 50mg / ml (mm) 25	INHIBITION % DECREASE	DOSE 250 mg/ml 20	INHIBITION % DECREASE
Crade Extract	Guines-pig *Uterus*	Oxy & PG	48	25	48	20	58
Chloroform		Oxy & PG	50	50	0	50	0
BuOH		Oxy & PG	51	48	6	40	21
Aqueous		Oxy & PG	48	35	27	25	48
Fraction-1		Oxy & PG	45	44	2	42	7
Fraction-2		Oxy & PG	48	48	0	45	6
Fraction-3		Oxy & PG	50	42	16	30	40
Fraction-4		Oxy & PG	50	44	12	28	44
Fraction-5		Oxy & PG	46	40	13	26	43
Fraction-6		Oxy & PG	48	40	17	24	50
S fr I		Oxy & PG	49	52	-	52	-
S fr II		Oxy & PG	50	50	0	52	-
S fr III		Oxy & PG	48	40	20	32	40
S fr IV		Oxy & PG	50	45	10	33	34
S fr V		Oxy & PG	50	48	4	45	10
S fr VI		Oxy & PG	52	50	4	50	4

Potentiation

Abbreviations used:

OXT = Oxytocin:PG = PGR 2α

CLAIMS

1. A Pharmaceutical composition useful as a tocolytic agent, said composition comprising an extract obtained from the plant *Salvadora persica* optionally, with conventional additives.

2. A composition as claimed in claim 1 wherein the extract contains compounds selected from β-amyrin, betulin, ursolic acid, methyl palmitate and lupeol.

3. A composition as claimed in claim 1 wherein the extract is an aqueous fraction obtained from the crude extract of *Salvadora persica.*

4. A composition as claimed in claim 1 wherein the additives are selected from therapeutically acceptable additives.

5. A composition as claimed in claim 1 wherein the amount of extract in the composition is 10μgms.

6. A composition as claimed in claim 1 wherein the dosage is of the extract is 3μg/ml to 10μg/ml.

7. A method for relieving pains in the uterine muscles, said method comprising the step of administering a therapeutically effective amount of extract obtained from the plant *Salvadora persica,* optionally with conventional additives to a subject in need thereof.

8. A method as claimed in claim 7 wherein the extract obtained from the plant *Salvadora persica,* is an aqueous extract.

9. A method as claimed in claim 7 wherein 3 to 10 μg/ml. amount of the extract is administered to the subject for a period of 10 minutes.

10. A method as claimed in claim 7 wherein the subject is a human or animal.

11. An aqueous fraction of the crude extract obtained from the plant *Salvadora persica,* and exhibiting tocolytic activity in mammals.

12. A process for the extraction and purification of biologically useful molecules from an associated mangrove plant *Salvadora persica*, which comprises the steps of:

    i) collecting and processing the plant plants of *Salvadora persica,*

    ii) preparing a crude extract from the plant parts of *Salvadora persica,*

iii) testing the crude extract using methods of pharmacology,

iv) fractionating the crude extract,

v) testing the fractions using methods of pharmacology,

vi) isolating the pure compounds by column chromatography,

vii) testing the pure compounds by using methods of pharmacology, and

viii) identifying the compounds by spectroscopy.