

BISPECIFIC ANTIBODY-CAMPTOTHECIN DRUG CONJUGATE AND PHARMACEUTICAL USE THEREOF

TECHNICAL FIELD

5 The present invention relates to the field of biopharmaceuticals, and specifically to an antibody-drug conjugate formed by a bispecific antibody and a camptothecin drug, and to a method of preparation and application of the antibody-drug conjugate. The present invention also relates to
10 linker-drug compounds that can be coupled with Ab to form an antibody-toxin conjugate.

TECHNICAL BACKGROUND

Epidermal Growth Factor Receptor (EGFR) and human epidermal
15 growth factor receptor 3 (also known as HER3 and ErbB3) are receptor protein tyrosine kinases and belong to the epidermal growth factor receptor (EGFR) subfamily of receptor protein tyrosine kinases, which includes EGFR (ErbB-1), HER2/c-neu (ErbB-2), HER3 (ErbB-3) and HER4 (ErbB-4).

20 The epidermal growth factor receptor is a glycoprotein with a molecular weight of 170 kDa that crosses cell membranes and is activated by binding to ligands. Upon activation, EGFR is converted from a monomer to an auto-dimer or forms a heterodimer with other HER family members. Dimer formation activates
25 intracellular kinase pathways that direct the phosphorylation of downstream pathways, including the MAPK, Akt, and JNK pathways, to induce cell proliferation. Studies have shown that high expression of EGFR is associated with tumor cell proliferation, angiogenesis, tumor invasion, etc. EGFR-related
30 signaling pathways play a very important role in the maintenance and growth of epidermal tissues. Especially in breast cancer, malignant glioma and lung cancer, EGFR can promote tumorigenesis. In lung cancer tissues, the EGFR signaling pathway shows a stimulated state, and there is a positive correlation between
35 the expression level of EGFR and the stage of cancer development.

Meanwhile, more and more studies are using EGFR as a biomarker of tumor drug resistance due to the finding of secondary mutations of EGFR under drug stress.

Epidermal growth factor receptor 3 (HER3 or ErbB3) also has the structure of a typical epidermal growth factor receptor, but HER3 lacks the structural domain of the intracellular protein tyrosine kinase and thus cannot autophosphorylate. HER3 can bind to ligand proteins and promote their heterodimerization with other human epidermal growth factor receptor family members to activate receptor-mediated signaling pathways, which not only acts as a signal diversification means, but also plays a role of signal amplification and accelerates tumor progression. Heregulins (glial cell growth factor, neu differentiation factor) can activate intracellular kinase-dependent multistep signaling pathway responses upon binding to the transmembrane receptors HER3 and HER4. Down-regulation of this signaling pathway often leads to Alzheimer's disease, heart failure, atherosclerosis, and cancer, etc. Up-regulation of HER3 expression level can promote tumorigenesis and growth by interacting with receptor tyrosine kinases (RTKs). Also, since HER3 is a heterodimeric molecular chaperone for other EGFR family members, it has the potential to modulate EGFR/HER2 signaling pathway-mediated drug resistance in cancer cells. Studies have also shown that HER3 defeats cancer therapy by activating the PI3K/AKT, MAPK/ERK and JAK/STAT signaling pathways.

Overexpression/dysregulation of EGFR and HER3 has been closely associated with the development of a variety of tumors. EGFR and HER3 have been shown to drive tumor progression in solid tumors such as breast, lung, gastric, and pancreatic cancers. Several studies have shown that high expression of HER3 is associated with clinical failure of EGFR antibodies and inhibitors. Several combination drug therapies targeting EGFR and HER3 are already underway in clinical settings.

Monoclonal antibodies have been widely used in anti-tumor

therapy in recent years, but their efficacy leaves much to be desired. A large number of tumor patients have poor clinical responses, and some patients with clinical responses develop drug resistance after a sustained period of monoclonal antibody therapy, leading to tumor recurrence. A bispecific monoclonal antibody is a monoclonal antibody molecule with two different antigen recognition sequences, which can bind protein molecules of two antigenic epitopes, and thus can achieve, for example, a variety of new and unique anti-tumor mechanisms, such as mediating the killing of tumor cells by immune cells, mediating the killing of tumor cells by toxic small molecules, or blocking the signaling pathways that promote the growth of tumors. The development of bispecific monoclonal antibodies is mainly in consideration of the fact that multiple mediators participate in the pathogenesis of tumors through specific or overlapping mechanisms, and if multiple targets are blocked at the same time, this will produce better therapeutic effects than inhibition of a single target, and at the same time, the action of multiple targets greatly reduces the probability of developing drug resistance. Currently, two bispecific antibodies, Catumaxomab and Blinatumomab, have been approved for marketing in the U.S., and more than fifty bispecific antibody molecules are in clinical trials.


Antibody-Drug Conjugates (ADCs) are molecules with specifically targeted killing effects, obtained by attaching small molecule toxins with cell-killing effects to antibodies, and are mainly used in the treatment of tumors and other diseases. Antibody-drug coupled drugs use antibodies that can specifically bind to proteins on the surface of tumor cells, and therefore have tumor specificity and potential unachievable by conventional drugs. Currently, 12 ADC drugs have received marketing approval worldwide and hundreds of programs are in clinical trials. However, most ADC programs currently on the market or in clinical settings are directed at a single target and cannot achieve the synergistic benefits of dual-target

therapy.

SUMMARY OF THE INVENTION

The inventors, based on a comprehensive understanding of the ADC class of drugs, disclose a bispecific antibody-drug conjugate and a method of preparing the same, a pharmaceutical composition comprising said conjugate and a use of said conjugate or pharmaceutical composition. The present invention also relates to linker-drug compounds that can be coupled to a bispecific antibody to form an antibody-toxin conjugate.

A first aspect of the present invention discloses a ligand-camptothecin derivative conjugate as shown in general formula I or a pharmaceutically acceptable salt or solvate thereof;




I

wherein:

Ab is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets two different epitopes or targets;


20 L_1 is selected without limitation from the group:



preferably L_1 is  ,  ,  ;

preferably L₁ is ,  or  ;


L₂ has the structure shown in formula A below,



formula A

wherein Y is a scaffold selected from C1-C6 alkyl, substituted C1-C6 alkyl, or C3-C8 cycloalkyl; preferably Y is C1-C6 alkyl; Ac is a hydrophilic structural unit; and the carbon No. 2 attached to Y has absolute chirality in the R configuration or the S configuration;

L₃ is present or absent, and when present, L₃ is selected

from PEG hydrophilic units: , o is selected from an integer in the range of 1-10 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10), preferably an integer in the range of 2-8;

L₄ is an enzymatic cutting unit;

L₅ is a linking unit;

in formula I, the No. 1 chiral carbon atom attached to N has absolute chirality in the R configuration or the S configuration;

R is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably R is selected from a hydrogen atom or a C1-C6 alkyl group;

R₁ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered-heteroaryl, or substituted 5-10-membered heteroaryl;

preferably R₁ is selected from a hydrogen atom or a C1-C6 alkyl group;

more preferably R₁ is selected from C1-C6 alkyl;

R₂ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably R₂ is selected from a hydrogen atom, a halogen or a C1-C6 alkyl group;

more preferably R₂ is selected from halogen;

X is selected from -C(O)-CR_aR_b-(CR₃R₄)_m-O-, -C(O)-CR_aR_b-(CR₃R₄)_m-NH- or -C(O)-CR_aR_b-(CR₃R₄)_m-S-;

preferably X is selected from -C(O)-CR_aR_b-(CR₃R₄)_m-O-;

R_a and R_b are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl, a deuterated C1-C6 alkyl, a halogenated C1-C6 alkyl, a C3-C8 cycloalkyl, a C3-C8 cycloalkyl C1-C6 alkyl, a C6-C10 aryl C1-C6 alkyl, a C1-C6 alkoxy C1-C6 alkyl, a 3-7-membered heterocyclic group, a substituted 3-7-membered heterocyclic group, a C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably, R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl, a halo-C1-C6 alkyl, a C3-C8 cycloalkyl C1-C6 alkyl or a C6-C10 aryl C1-C6 alkyl;

alternatively, R_a, R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclyl group, a substituted 3-7-membered heterocyclyl group; preferably R_a, R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group;

R₃, R₄ are identical or different and are each independently a hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, halogenated C1-C6 alkyl, deuterated C1-C6 alkyl, C1-C6 alkoxy, hydroxyl, amino, cyano, nitro, hydroxy C1-C6 alkyl, C3-C8 cycloalkyl, 3-7-membered heterocyclyl, or substituted 3-7-membered heterocyclyl;

preferably, R₃, R₄ are each independently a hydrogen atom or C1-C6 alkyl group;

alternatively, R₃, R₄ and the carbon atoms attached thereto constitute a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclic group, or a substituted 3-7-membered heterocyclic group;

m is selected from integers 0-4 (e.g., 0, 1, 2, 3, or 4), preferably 0, 1; n is selected from integers 1-10 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10).

In some embodiments of the first aspect of the present invention, ligand-camptothecin derivative conjugates as shown in general formula I or pharmaceutically acceptable salts or solvates thereof are disclosed, characterized in that Ab is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets two different epitopes or targets, preferably a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3.

In some embodiments of the first aspect of the present invention, the disclosure of a ligand-camptothecin derivative conjugate as shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that the Ab antibody comprises: an IgG1 heavy chain, a κ light chain, and a single-chain Fv (scFv) structural domain; wherein said single-chain Fv (scFv) structural domain forms a construct with the IgG1 heavy chain or the κ light chain; wherein said IgG1 heavy chain and κ light chain form an IgG portion with binding specificity for EGFR; said scFv structural domain has binding specificity for HER3, and the scFv structural domain is connected by a linker (e.g., having an amino acid sequence of (gly-gly-gly-gly-ser)_n, wherein n is an integer of at least 1, preferably n is an integer of 1 to 10) to the C-terminus or N-terminus of said IgG1 heavy chain or C-terminus or N-terminus of said κ light chain; and wherein the single-chain Fv structural domain has a structural order of N-terminus-heavy chain variable region-joint-light chain variable region-C-

terminus or N-terminus-light chain variable region-joint-heavy chain variable region-C-terminus (e.g., said joint consists of an amino acid sequence of (gly-gly-gly-gly-ser)_m, wherein *m* is an integer of at least 3, preferably *m* is 3, 4, 5, or 6).

5 In some embodiments of the first aspect of the present invention, the disclosure of a ligand-camptothecin derivative conjugate as shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that the κ light chain of the Ab antibody comprises CDRs as shown in
10 SEQ ID NO: 25, SEQ ID NO: 26, and SEQ ID NO: 27, and the IgG1 heavy chain comprises CDRs as shown in SEQ ID NO: 29, SEQ ID NO: 30, and SEQ ID NO: 31, and the single-chain Fv (scFv) structural domain comprises the heavy chain variable region CDRs as shown in SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO:
15 34, and light chain variable region CDRs as shown in SEQ ID NO: 35, SEQ ID NO: 36, and SEQ ID NO: 37.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or
20 solvate thereof, is disclosed, characterized in that the light chain of the Ab antibody comprises a variable region as shown in SEQ ID NO: 28, the IgG1 heavy chain comprises a variable region as shown in SEQ ID NO: 38, and the single chain Fv (scFv) structural domain comprises a heavy chain variable region as
25 shown in SEQ ID NO: 39 and a light chain variable region as shown in SEQ ID NO: 40.

In some embodiments of the first aspect of the present invention, a ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or
30 solvate thereof, is disclosed, characterized in that the light chain amino acid sequence of the Ab antibody is SEQ ID NO: 2, and the amino acid sequence of the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 4.

35 In some embodiments of the first aspect of the present

invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the light chain nucleic acid coding sequence of the Ab antibody is SEQ ID NO: 1, and the nucleic acid coding sequence of the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 3.

In some embodiments of the first aspect of the present invention, a ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the κ light chain of the Ab antibody comprises CDRs as shown in SEQ ID NO: 41, SEQ ID NO: 42, and SEQ ID NO: 43, the IgG1 heavy chain comprises CDRs as shown in SEQ ID NO: 45, SEQ ID NO: 46, and SEQ ID NO: 47, and the single-chain Fv (scFv) structural domain comprises heavy chain variable region CDRs as shown in SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO: 34, and light chain variable region CDRs as shown in SEQ ID NO: 35, SEQ ID NO: 36, and SEQ ID NO: 37.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the light chain of the Ab antibody comprises a variable region as shown in SEQ ID NO: 44, the IgG1 heavy chain comprises a variable region as shown in SEQ ID NO: 48, and the single-chain Fv (scFv) structural domain comprises a heavy chain variable region as shown in SEQ ID NO: 39 and a light chain variable region as shown in SEQ ID NO: 40.

In some embodiments of the first aspect of the present invention, a ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the light chain amino acid sequence of the Ab antibody is SEQ ID NO: 6, and the amino acid sequence of the construct of the heavy chain

of the antibody and the single-chain Fv (scFv) structural domain is SEQ ID NO: 8.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the light chain nucleic acid coding sequence of the Ab antibody is SEQ ID NO: 5, and the nucleic acid coding sequence of the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 7.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the light chain of the Ab antibody comprises a variable region as shown in SEQ ID NO: 49, the IgG1 heavy chain comprises a variable region as shown in SEQ ID NO: 52, and a single-chain Fv (scFv) structural domain comprises a heavy chain variable region as shown in SEQ ID NO: 50 and a light chain variable region as shown in SEQ ID NO: 51.

In some embodiments of the first aspect of the present invention, a ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that an Ab antibody has a heavy chain amino acid sequence of SEQ ID NO: 12, and a construct of the antibody light chain and a single-chain Fv (scFv) structural domain has an amino acid sequence of SEQ ID NO: 10.

In some embodiments of the first aspect of the present invention, a ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody has a heavy chain nucleic acid coding sequence of SEQ ID NO: 11, and a construct of the light chain of the antibody and a single-chain Fv (scFv) structural domain has a nucleic

acid coding sequence of SEQ ID NO: 9.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody has a light chain amino acid sequence of SEQ ID NO: 14, and a construct of the antibody heavy chain and a single-chain Fv (scFv) structural domain has an amino acid sequence of SEQ ID NO: 16.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody has a light chain nucleic acid coding sequence of SEQ ID NO: 13, and a construct of the antibody heavy chain and a single-chain Fv (scFv) structural domain has a nucleic acid coding sequence of SEQ ID NO: 15.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody has a heavy chain amino acid sequence of SEQ ID NO: 20, and a construct of the antibody light chain and a single-chain Fv (scFv) structural domain has an amino acid sequence of SEQ ID NO: 18.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody has a heavy chain nucleic acid coding sequence of SEQ ID NO: 19, and a construct of the light chain of the antibody and a single-chain Fv (scFv) structural domain has a nucleic acid coding sequence of SEQ ID NO: 17.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown

in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the light chain of the Ab antibody comprises a variable region as shown in SEQ ID NO: 53, an IgG1 heavy chain comprises a variable region as shown in SEQ ID NO: 54, and a single-chain Fv (scFv) structural domain comprises a heavy chain variable region as shown in SEQ ID NO: 50 and a light chain variable region as shown in SEQ ID NO: 51.


In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody has a heavy chain amino acid sequence of SEQ ID NO: 24, and the amino acid sequence of a construct of the light chain of the antibody and a single-chain Fv (scFv) structural domain is SEQ ID NO: 22.

In some embodiments of the first aspect of the present invention, the ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody has a heavy chain nucleic acid coding sequence of SEQ ID NO: 23, and a construct of the antibody light chain and a single-chain Fv (scFv) structural domain has a nucleic acid coding sequence of SEQ ID NO: 21.

In some embodiments of the first aspect of the present invention, a ligand-camptothecin derivative conjugate shown in general formula I, or a pharmaceutically acceptable salt or solvate thereof, is disclosed, characterized in that the Ab antibody comprises: two IgG1 heavy chains; two κ light chains; and two single-chain Fv (scFv) structural domains.

In some embodiments of the first aspect of the present invention, ligand-camptothecin derivative conjugates as shown in general formula I or pharmaceutically acceptable salts or solvates thereof are disclosed, characterized in that said X is selected without limitation from the following structures or

isomers thereof:



where the left wavy line is linked to a camptothecin derivative portion and the right wavy line is linked to L₅.

In some embodiments of the first aspect of the present invention, ligand-camptothecin derivative conjugates as shown in general formula I or pharmaceutically acceptable salts or solvates thereof are disclosed, characterized in that L₄ is selected without limitation from peptide residues formed of amino acids;

wherein optionally, said amino acid is further substituted with one or more substituents selected from deuterium atoms, halogens, hydroxyl, cyano, amino, nitro, carboxyl, C1-C6 alkyl, substituted C1-C6 alkyl, C1-C6 alkoxy and C3-C8 cycloalkyl or substituted C3-C8 cycloalkyl;

preferably, said peptide residue is a peptide residue formed from one, two or more amino acids selected from phenylalanine (F), glycine (G), valine (V), lysine (K), citrulline (C), serine (S), glutamic acid (E) or aspartic acid (D);

more preferably, said peptide residue is a tetrapeptide residue consisting of glycine (G)-glycine (G)-phenylalanine (F)-glycine (G).

Particularly preferably, said peptide residue is -GGFG-.

In some embodiments of the first aspect of the present invention, ligand-camptothecin derivative conjugates as shown in general formula I or pharmaceutically acceptable salts or solvates thereof are disclosed, characterized as follows:


L₅ is non-limitatively selected from -NR₅(CR₆R₇)_q- or a chemical bond, q is selected from an integer from 0-6 (e.g., 0, 1, 2, 3, 4, 5, or 6);

R₅, R₆ and R₇ are identical or different and are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl group, a substituted C1-C6 alkyl group, a deuterated C1-C6 alkyl group, a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a C1-C6 alkoxy C1-C6 alkyl group, a 3-7-membered heterocyclyl group, a substituted 3-7-membered heterocyclyl group, a C6-C10 aryl group, a substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;


preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom or a C1-C6 alkyl group;


more preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom.

In certain embodiments, L₁ is selected without limitation from:




In some embodiments of the first aspect of the present invention, ligand-camptothecin derivative conjugates as shown in general formula I or pharmaceutically acceptable salts or solvates thereof are disclosed, characterized in that said linking unit -L₁-L₂-L₃-L₄-L₅- is selected without limitation from the following structures;





preferably



wherein:

Ac is a hydrophilic structural unit;

R₅, R₆ and R₇ are identical or different and are each independently selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom or a C1-C6 alkyl group;


more preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom;

the carbon atom No. 2 attached to N has absolute chirality in the R configuration or the S configuration;

5 the left wavy line is linked to the antibody or its antigen-binding fragment portion, and the right wavy line is connected to the X;

o is selected from integers 1-10 (i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10).

10 The second aspect of the present invention discloses a ligand-camptothecin derivative conjugate as shown in general formula II, or a pharmaceutically acceptable salt or solvate thereof;





II

wherein:

Ab is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3;


20 L₁ is a linking unit that is connected to Ab, selected without limitation from:



preferably, L₁ is: ;

preferably, L₁ is: ;

L₃ is present or absent, and when L₃ is present, L₃ is

selected from , o is selected from integers 1-10 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10), preferably integers 2-8;

5 Ac is a hydrophilic structural unit;

the chiral carbon atoms at position 1, position 2 and position 3 have two chiral configurations, namely the R absolute configuration or the S absolute configuration;

R is selected from hydrogen atom, deuterium atom, halogen,
10 C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, and substituted 5-10-membered heteroaryl;

preferably R is selected from a hydrogen atom or a C1-C6
15 alkyl group;

R₁ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7-membered heterocyclyl, substituted 3-7-membered
20 heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably R₁ is selected from a hydrogen atom or a C1-C6 alkyl group;

more preferably, R₁ is selected from C1-C6 alkyl;

25 R₂ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably R₂ is selected from a hydrogen atom, a halogen or a C1-C6 alkyl group;

more preferably R₂ is selected from halogen;

X is selected from -C(O)-CR_aR_b-(CR₃R₄)_m-O-, -C(O)-CR_aR_b-

$(\text{CR}_3\text{R}_4)_m\text{-NH-}$ or $-\text{C}(\text{O})-\text{CR}_a\text{R}_b-(\text{CR}_3\text{R}_4)_m\text{-S-}$;

preferably X is selected from $-\text{C}(\text{O})-\text{CR}_a\text{R}_b-(\text{CR}_3\text{R}_4)_m\text{-O-}$;

R_a and R_b are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl group, a deuterated C1-C6 alkyl group, a halogenated C1-C6 alkyl group, a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a C1-C6 alkoxy C1-C6 alkyl group, a 3-7-membered heterocyclic group, a substituted 3-7-membered heterocyclic group, a C6-C10 aryl group, a substituted C6-C10 aryl group, a 5-10-membered heteroaryl, a substituted 5-10-membered heteroaryl;

preferably, R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl, a halo-C1-C6 alkyl, a C3-C8 cycloalkyl C1-C6 alkyl or a C6-C10 aryl C1-C6 alkyl;

alternatively, R_a , R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclyl group, a substituted 3-7-membered heterocyclyl group; preferably R_a , R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group;

R_3 , R_4 are identical or different and are independently hydrogen atoms, deuterium atoms, halogens, C1-C6 alkyl, halogenated C1-C6 alkyl, deuterated C1-C6 alkyl, C1-C6 alkoxy, hydroxyl, amino, cyano, nitro, hydroxy C1-C6 alkyl, C3-C8 cycloalkyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, respectively;

preferably, R_3 , R_4 are independently hydrogen atoms or C1-C6 alkyl groups, respectively;


alternatively, R_3 , R_4 and the carbon atoms attached thereto constitute a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclic group, a substituted 3-7-membered heterocyclic group;

m is selected from integers 0-4 (i.e., 0, 1, 2, 3 or 4), preferably 0, 1;

n is selected from the integers 1-10 (e.g., 1, 2, 3, 4, 5,

6, 7, 8, 9, or 10).

In some embodiments of the first and second aspects of the present invention, it is disclosed that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that: said Ac has the structure shown in formula B, as follows,



wherein:


Z is selected without limitation from the group consisting of one or more of a hydrophilic structural carboxyl group, phosphoric acid, polyphosphoric acid, phosphorous acid, sulfonic acid, sulfinic acid, or polyethylene glycol (PEG);


preferably Z is selected from a hydrophilic structural carboxyl group, phosphoric acid or polyethylene glycol (PEG);

Y' is optionally a scaffold connecting the amino group to Z; preferably Y' is a C1-C6 alkylene group (e.g. methylene);

Ac is connected to the 2-position carbon that has been labeled in structural formula I by means of a scaffold Y.

In some embodiments of the first aspect and the second aspect of the present invention, it is disclosed that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that: said Ac is selected without limitation from glycine, (D/L) alanine, (D/L) leucine, (D/L) isoleucine, (D/L) valine, (D/L) phenylalanine, (D/L) proline, (D/L) tryptophan, (D/L) serine, (D/L) tyrosine, (D/L) cysteine, (D/L) cystine, (D/L) arginine, (D/L) histidine, (D/L) methionine, (D/L) asparagine, (D/L) glutamine, (D/L) threonine, (D/L) aspartic acid, (D/L) glutamic acid, natural or unnatural amino acid derivatives or the following structures or isomers thereof,







In some embodiments of the first and second aspects of the present invention, the disclosure of said ligand-camptothecin derivative conjugates or pharmaceutically acceptable salts or solvates thereof is characterized in that: said Ac is selected without limitation from a glycine, phosphoric acid, (D/L) glutamic acid, or polyethylene glycol hydrophilic structure.

In some embodiments of the first and second aspects of the present invention, the disclosure of said ligand-camptothecin derivative conjugates or pharmaceutically acceptable salts or

solvates thereof is characterized in that said



the structure shown in formula d below;



d

wherein:

R is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably R is selected from a hydrogen atom or a C1-C6 alkyl group;

R₁ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably R₁ is selected from a hydrogen atom or a C1-C6 alkyl group;

more preferably R₁ is selected from C1-C6 alkyl;

R₂ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, substituted 5-10-membered heteroaryl;

preferably R₂ is selected from a hydrogen atom, a halogen or a C1-C6 alkyl group;

more preferably R₂ is selected from halogen;

R_a and R_b are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl, a deuterated C1-C6 alkyl, a halogenated C1-C6 alkyl, a C3-C8 cycloalkyl, a C3-C8 cycloalkyl C1-C6 alkyl, a C1-C6 alkoxy C1-C6 alkyl, a 3-7-membered heterocyclic group, a substituted 3-7-membered heterocyclic group, a C6-C10 aryl, a substituted C6-C10 aryl, a 5-10-membered heteroaryl, a substituted 5-10-membered heteroaryl;

preferably, R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl, a halo-C1-C6 alkyl, a C3-C8 cycloalkyl C1-C6 alkyl or a C6-C10 aryl C1-C6 alkyl;


preferably R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl, a halo-C1-C6 alkyl, a C3-C8 cycloalkyl C1-C6 alkyl, a C6-C10 aryl; preferably R_a and R_b are each independently selected from a hydrogen atom, a methyl group, an ethyl group, a trifluoromethyl group, a cyclopropylmethyl group, a phenyl group;


alternatively, R_a , R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclyl group, a substituted 3-7-membered heterocyclyl group; preferably, R_a , R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group (e.g., a C3-C5 cycloalkyl group);

the 1-position chiral carbon atom has two chiral configurations, namely the R absolute configuration or the S absolute configuration;

m is selected from 0 or 1.

In some embodiments of the first and second aspects of the present invention, the disclosure of said ligand-camptothecin derivative conjugate or a pharmaceutically acceptable salt or solvate thereof is characterized in that: said structural formula d is selected without limitation from the following compounds:






In some embodiments of the present invention, L₁ can comprise a succinimide group. In these embodiments, the ligand-drug conjugate can undergo hydrolysis under readily hydrolyzable


conditions, with the site of hydrolysis being the succinimide group of the linker unit. When the ligand contains multiple linker-drugs, the following scenarios can occur with varying degrees of hydrolysis:

- 5 the succinimide groups are completely non-hydrolyzed, i.e., the succinimide groups are all in a closed ring form




incomplete hydrolysis of the succinimide groups, i.e.,

some of the succinimide groups are in a closed ring form




- 10 or , and some of the succinimide groups are in an open



complete hydrolysis of the succinimide groups, i.e., the

succinimide groups are all in an open ring form




- 15 Thus, when multiple L_1 containing succinimide groups are present in the ADC at the same time (i.e., Ab is connected to multiple drug-linkers containing succinimide groups), these succinimide groups may be all in a closed ring form, partially in an open ring form, or all in an open ring form.

- 20 It will be appreciated that the present application, even though the succinimide group appearing in the chemical structural formula of the ADC is in the closed ring form, actually covers the three scenarios of the succinimide being all in a closed ring form, partially in an open ring form, and

preferably R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl, a halo-C1-C6 alkyl, a C3-C8 cycloalkyl C1-C6 alkyl, a C6-C10 aryl; preferably R_a and R_b are each independently selected from a hydrogen atom, a methyl group,
5 an ethyl group, a trifluoromethyl group, a cyclopropylmethyl group, a phenyl group;

alternatively, R_a , R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclyl group, a
10 substituted 3-7-membered heterocyclyl group; preferably, R_a , R_b and the carbon atoms to which they are attached constitute a C3-C8 cycloalkyl group (e.g., a C3-C5 cycloalkyl group);

L_3 is present or absent, and when L_3 is present, it is


selected from , o is selected from an integer from
15 1 to 10 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10);

the 1-position or 2-position chiral carbon atom has two chiralities, namely the R absolute configuration or S absolute configuration;

Ac is a hydrophilic structural unit;

m is selected from 0 or 1.

In some embodiments of the third aspect of the present invention, it is disclosed that said linker-drug compounds or pharmaceutically acceptable salts or solvates thereof are characterized in that: said Ac has the structure shown in
25 formula B below,




wherein:

Z is selected without limitation from the group consisting
30 of one or more of a hydrophilic structural carboxyl group, a phosphoric acid, a polyphosphoric acid, a phosphorous acid, a sulfonic acid, a sulfinic acid, or a polyethylene glycol (PEG);

Y' is an optional scaffold connecting the amino group to Z ; preferably Y' is a C1-C6 alkylene group (e.g. methylene);











Ac is connected to the 2-position carbon that has been labeled in structural formula I by means of scaffold Y.

In some embodiments of the third aspect of the present invention, disclosed are the linker-drug compounds or pharmaceutically acceptable salts or solvates thereof, characterized in that said Ac is selected without limitation from glycine, (D/L) alanine, (D/L) leucine, (D/L) isoleucine, (D/L) valine, (D/L) phenylalanine, (D/L) proline, (D/L) tryptophan, (D/L) serine, (D/L) tyrosine, (D/L) cysteine, (D/L) cystine, (D/L) arginine, (D/L) histidine, (D/L) methionine, (D/L) asparagine, (D/L) glutamine, (D/L) threonine, (D/L) aspartic acid, (D/L) glutamic acid, natural or unnatural amino acid derivatives, or the following structures,



















In some embodiments of the third aspect of the present invention, disclosed are the linker-drug compounds or pharmaceutically acceptable salts or solvates thereof, characterized in that: the Ac is selected without limitation from a glycine, phosphoric acid, (D/L) glutamic acid, or polyethylene glycol hydrophilic structure.

In some embodiments of the third aspect of the present invention, disclosed are the linker-drug compounds or pharmaceutically acceptable salts or solvates thereof, characterized in that said linker-drug compounds are selected without limitation from the following structures or isomers thereof,

















5














5






5






5



or




where: o is selected from the integers 1-10 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10).

10 Said linker-drug compounds or pharmaceutically acceptable salts or solvates thereof disclosed in the third aspect of the

present invention can be used as intermediates for coupling with the ligand Ab to form ligand-camptothecin derivative conjugates of formula I or formula II described in the first aspect and the second aspect.

5 A fourth aspect of the present invention discloses a method of preparing a ligand-camptothecin derivative conjugate or a pharmaceutically acceptable salt or solvate thereof as shown in general formula I or general formula II as described in the first and second aspects, characterized in that the
10 method comprises the following steps,



a ligand-camptothecin derivative conjugate as shown in general formula I or general formula II is obtained by a
15 coupling reaction of a reduced antibody or antigen-binding fragment thereof with a linker-drug compound;


the chiral carbon atom at position 1, position 2, or position 3 has absolute chirality in the R configuration or S configuration;















20 Ab, L₁, L₂, L₃, L₄, L₅, X, R, R₁, R₂ and n are as previously described.















The present application also relates to the use of the linker-drug compounds or pharmaceutically acceptable salts or solvates thereof, disclosed and described in the third aspect,
25 as intermediates in the preparation of ligand-camptothecin derivative conjugates or pharmaceutically acceptable salts or















solvates thereof. In certain embodiments, said ligand-camptothecin derivative conjugates or pharmaceutically acceptable salts or solvates thereof are the ligand-camptothecin derivative conjugates or pharmaceutically acceptable salts or solvates thereof disclosed in the first, second and fourth aspects of the present invention. In certain embodiments, said preparation is carried out according to the method of preparation disclosed in the fourth aspect.















In some embodiments of the first aspect, the second aspect and the fourth aspect of the present invention, the disclosure of said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is selected without limitation from the following structures, or succinimide open-ring structures thereof, or isomers thereof,















































- 5 wherein:
- SI-1x6.4 is a bispecific antibody or its antigen-binding fragment simultaneously targeting EGFR and HER3;
- n is selected from the integers 1-10 (i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10).
- 10 In some embodiments of the first aspect, the second aspect and the fourth aspect of the present invention, the disclosure of said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is selected without limitation
- 15 from the following structures or succinimide open-ring structures thereof or isomers thereof,




ADC-107




ADC-108




ADC-109




ADC-110




ADC-111




ADC-111a




ADC-111b




ADC-112




ADC-112a




ADC-112b




ADC-113




ADC-113a


















ADC-113b


















ADC-114

























or




wherein:

SI-1x4 is a bispecific antibody or its antigen-binding fragment simultaneously targeting EGFR and HER3;

n is selected from the integers 1-10 (i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10).

In some embodiments of the first aspect, the second aspect and the fourth aspect of the present invention, the disclosure of said ligand-camptothecin derivative conjugate, or a pharmaceutically

acceptable salt or solvate thereof, is characterized in that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,




wherein:

SI-1x22 is a bispecific antibody or an antigen-binding fragment thereof simultaneously targeting EGFR and HER3;

n is selected from the integers 1-10 (e.g., 1, 2, 3, 4, 5, 6,

7, 8, 9, or 10).

In some embodiments of the first aspect, the second aspect and the fourth aspect of the present invention, the disclosure of said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,




wherein:


SI-1×24 is a bispecific antibody or its antigen-binding fragment simultaneously targeting EGFR and HER3;

n is selected from the integers 1-10 (i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10).

5 In some embodiments of the first aspect, the second aspect and the fourth aspect of the present invention, the disclosure of said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is selected without limitation
10 from the following structures or succinimide open-ring structures thereof or isomers thereof,



15





wherein:

SI-1x25 is a bispecific antibody or an antigen-binding fragment thereof simultaneously targeting EGFR and HER3;

n is selected from the integers 1-10 (i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10).

In some embodiments of the first aspect, the second aspect and the fourth aspect of the present invention, the disclosure of said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is characterized in that said ligand-camptothecin derivative conjugate, or a pharmaceutically acceptable salt or solvate thereof, is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,





wherein:

SI-1×26 is a bispecific antibody or its antigen-binding
 5 fragment simultaneously targeting EGFR and HER3;

n is selected from the integers 1-10 (i.e., 1, 2, 3, 4, 5, 6,
 7, 8, 9, or 10).

In some embodiments of the first aspect, the second aspect and
 the third aspect of the present invention, the disclosure of said
 10 ligand-camptothecin derivative conjugate or pharmaceutically
 acceptable salt or solvate thereof or linker-drug compound or
 pharmaceutically acceptable salt or solvate thereof is
 characterized in that said pharmaceutically acceptable salt
 comprises a sodium salt, a potassium salt, a calcium salt or a
 15 magnesium salt formed with an acidic functional group in the
 structural formula, and acetate, trifluoroacetate, citrate,
 oxalate, tartrate, malate, nitrate, chloride, bromide, iodide,
 sulfate, bisulfate, phosphate, lactate, oleate, ascorbate,
 salicylate, formate, glutamate, methanesulfonate, ethanesulfonate,
 20 benzenesulfonate or p-toluenesulfonate formed with a basic
 functional group in the structure.

A fifth aspect of the present invention discloses a
 pharmaceutical composition comprising a ligand-camptothecin
 derivative conjugate described in the first aspect and the second
 25 aspect or a pharmaceutically acceptable salt or solvate thereof or
 a linker-drug compound described in the third aspect or a
 pharmaceutically acceptable salt or solvate thereof, and

optionally a pharmaceutically acceptable carrier.

A sixth aspect of the present invention discloses a pharmaceutical preparation comprising a ligand-camptothecin derivative conjugate described in the first aspect and the second
5 aspect, or a pharmaceutically acceptable salt or solvate thereof, or a linker-drug compound described in the third aspect, or a pharmaceutically acceptable salt or solvate thereof.

A seventh aspect of the present invention discloses the use of the following in the preparation of drugs for the treatment or
10 prevention of cancers or tumors: the ligand-camptothecin derivative conjugates described in the first aspect and the second aspect or pharmaceutically acceptable salts or solvates thereof, or the linker-drug compounds described in the third aspect or pharmaceutically acceptable salts or solvates thereof, or the
15 pharmaceutical compositions described in the fifth aspect and/or the pharmaceutical preparations described in the sixth aspect;

alternatively, the use of the following in the treatment or prevention of cancer or tumors: the ligand-camptothecin derivative conjugates described in the first aspect and the second aspect or
20 pharmaceutically acceptable salts or solvates thereof, or the linker-drug compounds described in the third aspect or pharmaceutically acceptable salts or solvates thereof, or the pharmaceutical compositions described in the fifth aspect and/or the pharmaceutical preparations described in the sixth aspect;

preferably, the cancer or tumor expresses EGFR and/or HER3;

more preferably, the cancer or tumor is selected from adenocarcinoma, ovarian cancer, cervical cancer, uterine cancer, prostate cancer, renal cancer, urethral cancer, bladder cancer, liver cancer, gastric cancer, endometrial cancer, salivary gland
30 cancer, esophageal cancer, lung cancer, colon cancer, rectal cancer, colorectal cancer, bone cancer, skin cancer, thyroid cancer, pancreatic cancer, melanoma, glioma, neuroblastoma, glioblastoma multiforme, sarcoma, lymphoma and leukemia, and other solid tumors or blood tumors.

An eighth aspect of the present invention discloses a method

of treating or preventing cancer or tumors, the method comprising administering to a subject in need thereof a prophylactically or therapeutically effective amount of a ligand-camptothecin derivative conjugate described in the first or second aspect or a pharmaceutically acceptable salt or solvate thereof or a linker-drug compound described in the third aspect or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition as described in the fifth aspect and/or a pharmaceutical preparation described in the sixth aspect;

preferably, the cancer or tumor expresses EGFR and/or HER3;

more preferably, the cancer or tumor is selected from adenocarcinoma, ovarian cancer, cervical cancer, uterine cancer, prostate cancer, renal cancer, urethral cancer, bladder cancer, liver cancer, gastric cancer, endometrial cancer, salivary gland cancer, esophageal cancer, lung cancer, colon cancer, rectal cancer, colorectal cancer, bone cancer, skin cancer, thyroid cancer, pancreatic cancer, melanoma, glioma, neuroblastoma, glioblastoma multiforme, sarcoma, lymphoma and leukemia, and other solid tumors or blood tumors.

In the above aspects of the present invention and its embodiments,

"C1-C6 alkyl" and "C1-C6 alkyl" in various composite groups involving "C1-C6 alkyl" (e.g., "substituted C1-C6 alkyl", "deuterated C1-C6 alkyl") may be replaced with "C1-C20 alkyl", "C1-C12 alkyl" or "C1-C10 alkyl";

"C3-C8 cycloalkyl" and "C3-C8 cycloalkyl" in various composite groups involving "C3-C8 cycloalkyl" may be replaced by "C3-C20 cycloalkyl" or "C3-C10 cycloalkyl";

"C1-C6 alkoxy" and "C1-C6 alkoxy" in various composite groups involving same can be replaced with "C1-C20 alkoxy", "C1-C12 alkoxy" or "C1-C10 alkoxy";

"C6-C10 aryl" and "C6-C10 aryl" in various composite groups involving same can be replaced with "C6-C12 aryl";

"3-7-membered heterocyclic group" and "3-7-membered heterocyclic group" in various composite groups involving same may

be replaced with "3-20-membered heterocyclic group", "3-12-membered heterocyclic group" or "3-10-membered heterocyclic group".

Beneficial Effects

The EGFR/HER3 bispecific antibody-drug conjugate provided by the present invention is a bispecific antibody ADC directed against both EGFR/HER3 targets, with good molecular stability and good preclinical efficacy, and is expected to have excellent clinical therapeutic effects.

Description of Drawings

Figure 1A illustrates SEC-HPLC detection/measurement of ADC-5 aggregation.

Figure 1B illustrates SEC-HPLC detection/measurement of ADC-6 aggregation.

Figure 1C illustrates SEC-HPLC detection/measurement of ADC-64 aggregation.

Figure 1D illustrates SEC-HPLC detection/measurement of ADC-DS aggregation.

Figure 1E illustrates SEC-HPLC detection/measurement of ADC-108 aggregation.

Figure 1F illustrates SEC-HPLC detection/measurement of ADC-112 aggregation.

Figure 1G illustrates SEC-HPLC detection/measurement of ADC-215 aggregation.

Figure 1H illustrates SEC-HPLC detection/measurement of ADC-219 aggregation.

Figure 1I illustrates SEC-HPLC detection/measurement of ADC-227 aggregation.

Figure 1J illustrates SEC-HPLC detection/measurement of ADC-235 aggregation.

Figure 2A illustrates RP-HPLC detection/measurement of ADC-5 drug-antibody coupling ratio (DAR).

Figure 2B illustrates RP-HPLC detection/measurement of ADC-6 drug-antibody coupling ratio (DAR).

Figure 2C illustrates RP-HPLC detection/measurement of ADC-10

drug-antibody coupling ratio (DAR).

Figure 2D illustrates RP-HPLC detection/measurement of ADC-12 drug-antibody coupling ratio (DAR).

Figure 2E illustrates RP-HPLC detection/measurement of ADC-64 drug-antibody coupling ratio (DAR).

Figure 2F illustrates RP-HPLC detection/measurement of ADC-108 drug-antibody coupling ratio (DAR).

Figure 2G illustrates RP-HPLC detection/measurement of ADC-112 drug-antibody coupling ratio (DAR).

Figure 2H illustrates RP-HPLC detection/measurement of ADC-215 drug-antibody coupling ratio (DAR).

Figure 2I illustrates RP-HPLC detection/measurement of ADC-219 drug-antibody coupling ratio (DAR).

Figure 2J illustrates RP-HPLC detection/measurement of ADC-227 drug-antibody coupling ratio (DAR).

Figure 2K illustrates RP-HPLC detection/measurement of ADC-235 drug-antibody coupling ratio (DAR).

Figure 2L illustrates RP-HPLC detection/measurement of ADC-243 drug-antibody coupling ratio (DAR).

Figure 3A illustrates that ADC-112 and the SI-1×4 antibody maintain affinity for both the antigens EGFR and HER3-Fc.

Figure 3B illustrates that ADC-6 and the SI-1×6.4 antibody maintain affinity for both the antigens EGFR and HER3-Fc.

Figure 3C illustrates that ADC-219 and the SI-1×22 antibody maintain affinity for both the antigens EGFR and HER3-Fc.

Figure 3D illustrates that ADC-227 and the SI-1×24 antibody maintain affinity for both the antigens EGFR and HER3-Fc.

Figure 3E illustrates that ADC-235 and the SI-1×25 antibody maintain affinity for both the antigens EGFR and HER3-Fc.

Figure 3F illustrates that ADC-243 and the SI-1×26 antibody maintain affinity for both the antigens EGFR and HER3-Fc.

Figure 4A illustrates the in vitro efficacy of six naked antibodies as well as six ADCs in A431.

Figure 4B illustrates the in vitro efficacy of six naked antibodies as well as six ADCs in BXP-3.

Figure 4C illustrates the in vitro efficacy of six naked antibodies as well as six ADCs in FaDu.

Figure 4D illustrates the in vitro efficacy of six naked antibodies as well as six ADCs in HARA-B.

5 Figure 4E illustrates the in vitro efficacy of six naked antibodies as well as six ADCs in HCC827.

Figure 4F illustrates the in vitro efficacy of the six naked antibodies as well as the six ADCs in SW620.

10 Figure 5A illustrates the results of in vivo efficacy experiments of ADC-6 and SI-1x6.4 naked antibody in the A431 single-tumor model.

Figure 5B illustrates the results of in vivo efficacy experiments of ADC-6, SI-1x6.4 naked antibody, Cetuximab (Cet) and ADC-214 in the SW620 single-tumor model.

15 Figure 5C illustrates the results of in vivo efficacy experiments of ADC-6 and SI-1x6.4 naked antibody in A431+SW620 heterogeneous tumors.

20 Figure 6A illustrates the results of in vivo efficacy experiments of ADC-6, ADC-219, ADC-235, ADC-227 and ADC-112 in the A431 single-tumor model.

Figure 6B illustrates the results of in vivo efficacy experiments of ADC-6, ADC-219, ADC-235, ADC-227 and ADC-112 in the BXP-3 single-tumor model.

25 Figure 7A illustrates the in vitro efficacy of SI-1x6.4, Cetuximab, ADC-6, ADC-214 and d3 on the human poorly differentiated lung cancer squamous cell carcinoma cell line Oka-c-1.

Figure 7B illustrates the in vitro efficacy of SI-1x6.4, Cetuximab, ADC-6, ADC-214 and d3 on human lung squamous cell carcinoma cells SK-MES-1. Specific embodiments

30 Abbreviations and definitions

Unless otherwise indicated, the following terms and phrases, as used herein, are intended to have the meanings set forth below. When a trade name is used herein, unless otherwise indicated in the context, the trade name includes the product formula, generic
35 drug and active ingredient of said trade name product.

Unless stated to the contrary, terms used in the claims and specification herein have the meanings set forth below.

The term "ligand" refers to a macromolecular compound that is able to recognize and bind to an antigen or receptor associated with a target cell. The ligand serves to present the drug to a target cell population bound to the ligand; these ligands include, but are not limited to, protein-like hormones, lectins, growth factors, antibodies, or other molecules capable of binding to cells. In embodiments of the present invention, the ligand is denoted as Ab, and the ligand may form a linkage bond with a linkage unit via a heteroatom on the ligand, and is preferably an antibody or an antigen-binding fragment thereof, said antibody being selected from chimeric, humanized, fully human, or murine antibodies; preferably monoclonal antibodies.

A ligand unit is a targeting agent that binds specifically to a target part. Said ligand is capable of specifically binding to a cellular component or binding to a cellular component or binding to other target molecules of interest. The target part or target is typically on the surface of the cell. In some aspects, the ligand unit serves to deliver the drug unit to a specific target cell population with which the ligand unit interacts. Ligands include, but are not limited to, proteins, polypeptides and peptides, and non-proteins such as sugars. Suitable ligand units include, for example, antibodies, such as full-length (intact) antibodies and antigen-binding fragments thereof. In embodiments where the ligand unit is a non-antibody targeting reagent, it may be a peptide or polypeptide, or a non-protein molecule. Examples of such targeting reagents include interferons, lymphokines, hormones, growth factors and colony stimulating factors, vitamins, nutrient transporter molecules, or any other cell-binding molecule or substance. In some embodiments, the linker is covalently attached to the sulfur atom of the ligand. In some aspects, the sulfur atom is a sulfur atom of a cysteine residue, and forms an interchain disulfide bond of the antibody. In another aspect, the sulfur atom is a sulfur atom of a cysteine residue that has been

introduced into the ligand unit, and forms an interchain disulfide bond of the antibody. In another aspect, the sulfur atom is a sulfur atom of a cysteine residue that has been introduced into the ligand unit (e.g., by site-directed mutagenesis or chemical reaction). In other aspects, the linker-bound sulfur atom is selected from a cysteine residue that forms the interchain disulfide bond of the antibody or a cysteine residue that has been introduced into the ligand unit (e.g., by site-directed mutagenesis or chemical reaction). In some embodiments, the system is numbered according to the EU index in Kabat {[Kabat E.A et al, (1991)] "Sequences of proteins of Immunological Interest", Fifth Edition, NIH Publication 91-3242}.

As used herein, "antibody" or "antibody unit" includes, to the extent thereof, any part of an antibody structure. This unit may bind, reactively associate, or complex with a receptor, antigen, or other receptor unit possessed by the target cell population. The antibody may be any protein or protein-like molecule, and can bind, complex, or react with a portion of the cell population to be treated or biologically modified. The antibodies forming the antibody-drug conjugates in the present invention maintain their original antigen-binding capacity from the wild state. Thus, the antibodies of the present invention are capable of binding exclusively to antigens. Antigens involved include, for example, tumor-associated antigens (TAA), cell surface receptor proteins and other cell surface molecules, cell survival regulators, cell proliferation regulators, molecules associated with tissue growth and differentiation (such as are known or foreseen to be functional), lymphokines, cytokines, molecules participating in the regulation of the cell cycle, molecules participating in angiogenesis, and molecules associated with angiogenesis (such as are known or foreseen to be functional). Tumor-associated factors may be cluster differentiation factors (e.g., CD proteins).

Antibodies applied in antibody drug conjugates include, but are not limited to, antibodies directed against cell surface receptors and tumor-associated antigens. Such tumor-associated antigens are

well known in the industry and can be prepared by methods and information for antibody preparation well known in the industry. To develop effective cellular level targets that can be used in cancer diagnosis and therapy, researchers seek to find transmembrane or other tumor-associated peptides. These targets can be specifically expressed on the surface of one or more cancer cells with little or no expression on the surface of one or more non-cancer cells. Typically, such tumor-associated polypeptides are more overexpressed on the surface of cancer cells than on the surface of non-cancer cells. Confirmation of such tumor-associated factors can greatly enhance the specific targeting properties of antibody-based cancer therapies. For convenience, information related to antigens known to the industry is labeled below, including name, other names, and gene bank accession number. Nucleic acid and protein sequences corresponding to the tumor-associated antigens can be found in publicly available databases, such as Genbank. The tumor-associated antigens corresponding to antibody targeting include all amino acid sequence variants and isoforms having at least 70%, 80%, 85%, 90%, or 95% homology with the sequences confirmed in the references, or possessing biological properties and characteristics that are identical to those of the tumor-associated antigen sequences in the cited literature.

The term "inhibit" or "inhibition" means that a detectable amount is reduced or completely prevented.

The term "cancer" refers to a physiological condition or disease characterized by dysregulated cell growth. "Tumor" includes cancer cells.

The term "autoimmune disease" refers to diseases or disorders that originate in tissues or proteins that target an individual's own body.

The term "drug" refers to cytotoxic drugs, denoted by d, which are chemical molecules that have a strong ability to disrupt normal growth in tumor cells. Cytotoxic drugs can in principle kill tumor cells at sufficiently high concentrations, but due to a lack of

specificity, they will also cause apoptosis of normal cells while killing tumor cells, leading to serious side effects. The term includes toxins, such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, radioisotopes (e.g., radioisotopes of At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and Lu¹⁷⁶), toxic drugs, chemotherapeutic drugs, antibiotics and nucleolytic enzymes, preferably toxic drugs.

The term "linker" or "linker fragment" or "linker unit" refers to a fragment or bond of a chemical structure that is attached to a ligand at one end and to the drug at the other end, or may be attached to other connectors and then to the drug.

Connectors, including extensions, spacers, and amino acid units, can be synthesized by methods known in the art, such as those described in US2005-0238649A1. The connectors can be "cleavable connectors" that facilitate the release of the drug in the cell. For example, acid-unstable connectors (e.g., hydrazone), protease-sensitive (e.g., peptidase-sensitive) connectors, photo-unstable connectors, dimethyl connectors, or disulfide-containing connectors can be used (Chari et al. Cancer Research 52:127-131, 1992); U.S. Patent No. 5, 208, 020.

According to the mechanism of intracellular drug release, as used herein, "linkers" or "linkers of antibody-drug conjugates" can be categorized into two types: unbreakable linkers and breakable linkers. For antibody-drug conjugates containing an unbreakable linker, the mechanism of drug release is as follows: after the conjugate binds to the antigen and is endocytosed by the cell, the antibody is cleaved enzymatically in lysosomes, releasing an active molecule consisting of the small-molecule drug, the linker, and amino acid residues of the antibody. The resulting change in the structure of the drug molecule does not diminish its cytotoxicity, but because the active molecule is electrically charged (amino acid residues), it cannot penetrate neighboring cells. Therefore, such active drugs cannot kill neighboring tumor cells that do not express the targeted antigen (antigen-negative cells) (bystander effect) (Ducry et al., 2010, Bioconjugate Chem.

21: 5-13). For antibody-drug conjugates containing a breakable linker, the mechanism of drug release is that after the conjugate binds to the antigen and is endocytosed by the cell, the conjugate breaks and releases the active ingredient (the small-molecule drug itself) in the target cell. Breakable linkers are mainly categorized into: chemical-sensitive linkers and enzyme-sensitive linkers. Chemically sensitive linkers can be selectively broken due to differences in the properties of the plasma and cytoplasm or tumor microenvironment. Such properties include pH, glutathione concentration, etc. pH-sensitive linkers, which are relatively stable in the neutral or weakly alkaline environment of blood (pH 7.3-7.5), will however be hydrolyzed within the weakly acidic tumor microenvironment (pH 5.0-6.5) and lysosomes (pH 4.5-5.0), e.g., hydrazones, carbonates, acetals, and ketals. Due to the limited plasma stability of acid-breakable linkers, antibody-drug conjugates based on such linkers typically have a short half-life (2-3 days). This short half-life has somewhat limited the use of pH-sensitive linkers in the new generation of antibody-drug conjugates. Glutathione-sensitive linkers are also known as disulfide-bond linkers. Drug release is based on the difference between the high intracellular glutathione concentration (millimolar range) and the relatively low glutathione concentration in the blood (micromolar range). This is particularly true for tumor cells, whose low oxygen content leads to enhanced reductase activity and thus to higher glutathione concentrations. Disulfide bonds are thermodynamically stable and therefore have better stability in plasma. Enzyme-unstable linkers, such as peptide linkers, provide better control of drug release. Peptide linkers can be effectively severed by lysosomal proteases such as cathepsins (Cathepsin B). This peptide linkage is thought to be very stable in the plasma circulation due to the unfavorable extracellular pH and serum protease inhibitors resulting in proteases that are normally inactive outside the cell. In view of the high plasma stability and good intracellular break selectivity and effectiveness, enzyme-unstable linkers are widely

used as breakable linkers for antibody-drug conjugates.

The term "antibody-drug conjugate" refers to the attachment of an antibody to a biologically active drug by means of a stable linkage unit. In the context of the present invention, "ligand-
5 drug conjugate", preferably antibody-drug conjugate (ADC), refers to the attachment of a monoclonal antibody or antibody fragment to a biologically active toxic drug by means of a stable linkage unit.

The three-letter codes and single-letter codes for amino acids used in this disclosure are as described in *J. boil. Chem.* 1968,
10 243, 3558.

The term "alkyl" refers to a saturated aliphatic hydrocarbon group, which is a straight or branched chain group containing 1 to 20 carbon atoms (i.e., "C1-C20 alkyl"), preferably an alkyl group containing 1 to 12 carbon atoms (i.e., "C1-C12 alkyl"), more
15 preferably an alkyl group containing 1 to 10 carbon atoms (i.e., "C1-C10 alkyl"), and most preferably an alkyl group containing 1 to 6 carbon atoms (i.e., "C1-C6 alkyl"). Non-limiting examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, sec-butyl, n-pentyl, 1,1-dimethylpropyl, 1,2-
20 dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, 2-methylbutyl, 3-methylbutyl, n-hexyl, 1-ethyl-2-methylpropyl, 1,1,2-trimethylpropyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 1,3-dimethylbutyl, 2-ethylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,3-dimethylbutyl, n-heptyl, 2-
25 methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 2,2-dimethylpentyl, 3,3-dimethylpentyl, 2-ethylpentyl, 3-ethylpentyl, n-octyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 2,2-dimethylhexyl, 3,3-dimethylhexyl, 4,4-dimethylhexyl, 2-ethylhexyl, 3-ethylhexyl, 4-ethylhexyl, 2-methyl-2-ethylpentyl, 2-methyl-3-
30 ethylpentyl, n-nonyl, 2-methyl-2-ethylhexyl, 2-methyl-3-ethylhexyl, 2,2-diethylpentyl, n-decyl, 3,3-diethylhexyl, 2,2-diethylhexyl, and their various branched isomers, and the like. More preferred are lower alkyl groups containing 1 to 6 carbon
35 atoms, and non-limiting embodiments include methyl, ethyl, n-

propyl, isopropyl, n-butyl, isobutyl, tert-butyl, sec-butyl, n-pentyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, 2-methylbutyl, 3-methylbutyl, n-hexyl, 1-ethyl-2-methylpropyl, 1,1,2-trimethylpropyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 1,3-dimethylbutyl, 2-ethylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,3-dimethylbutyl, and the like. The alkyl group may be substituted or non-substituted, and when substituted, the substituent group may be substituted at any available point of attachment, said substituent group preferably being one or more of the following groups, independently selected from alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halogen, mercapto, hydroxy, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkylthio, heterocycloalkylthio, oxo.

The term "substituted alkyl" means that a hydrogen in the alkyl group has been replaced by a substituent group. Unless otherwise indicated in the text, the substituent group of the alkyl group may be a variety of groups selected from the following group: - halogen, -OR', -NR'R'', -SR', -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NR'S(O)₂R'', -CN and -NO₂, the number of substituents is from 0 to (2m'+1), where m' is the total number of carbon atoms in the group. R', R'' and R''' each independently refer to hydrogen, unsubstituted C₁₋₈ alkyl, unsubstituted C₆₋₁₂ aryl (or C₆₋₁₀ aryl), C₆₋₁₂ aryl (or C₆₋₁₀ aryl) substituted by 1-3 halogens, unsubstituted C₁₋₈ alkyl, C₁₋₈ alkoxy or C₁₋₈ thioalkoxy, or unsubstituted C₆₋₁₂ aryl (or C₆₋₁₀ aryl)-C₁₋₄ alkyl. When R' and R'' are attached to the same nitrogen atom, they may form a 3-, 4-, 5-, 6-, or 7-membered ring together with that nitrogen atom. For example, -NR'R'' includes 1-pyrrolidinyl and 4-morpholinyl.

The term "alkylene" means a saturated straight or branched aliphatic hydrocarbon group, having two residues derived by

removing two hydrogen atoms from the same carbon atom or two different carbon atoms of a parent alkane, and is a straight or branched group comprising 1 to 20 carbon atoms, preferably an alkylene group containing 1 to 12 carbon atoms, more preferably 1 to 6 carbon atoms. Non-limiting examples of alkylene groups include, but are not limited to, methylene (-CH₂-), 1,1-ethylidene (-CH(CH₃)-), 1,2-ethylidene (-CH₂CH₂-), 1,1-propylidene (-CH(CH₂CH₃)-), 1,2-propylidene (-CH₂CH(CH₃)-), 1,3-propylidene (-CH₂CH₂CH₂-), 1,4-butylidene (-CH₂CH₂CH₂CH₂-), and 1,5-butylidene (-CH₂CH₂CH₂CH₂CH₂-), among others. The alkylene group may be substituted or non-substituted, and when substituted, the substituent may be substituted at any available point of attachment, said substituent preferably being independently and optionally substituted with one or more substituents selected from alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halo, mercapto, hydroxyl, nitro, cyano, cycloalkyl, heterocyclo, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkylthio, heterocycloalkylthio and oxo.

The term "alkoxy" refers to -O-(alkyl) and -O-(cycloalkyl), wherein alkyl or cycloalkyl is defined as above. Non-limiting examples of C1-C6 alkoxy include: methoxy, ethoxy, propoxy, butoxy, cyclopropoxy, cyclobutoxy, cyclopentyloxy, cyclohexyloxy. The alkoxy group may be optionally substituted or non-substituted, and when substituted, the substituent is preferably one or more of the following groups independently selected from alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halogen, mercapto, hydroxyl, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkylthio, heterocycloalkylthio.

The term "cycloalkyl" refers to a saturated or partially unsaturated monocyclic or polycyclic cyclic hydrocarbon substituent, wherein the cycloalkyl ring comprises from 3 to 20 carbon atoms (i.e. "C3-C20 cycloalkyl"), preferably comprising from 3 to 12 carbon atoms (i.e. "C3-C12 cycloalkyl"), more preferably comprising 3 to 10 carbon atoms (i.e., "C3-C10

cycloalkyl"), most preferably comprising 3 to 8 carbon atoms (i.e., "C3-C8 cycloalkyl"). Non-limiting examples of monocyclic cycloalkyl groups (e.g., "C3-C8 cycloalkyl") include cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl, cycloheptyl, cycloheptatrienyl, cyclooctyl, and the like; and polycyclic cycloalkyl groups include cycloalkyl groups of spirocycles, fused rings, and bridged rings.

The term "heterocyclyl" means a saturated or partially unsaturated monocyclic or polycyclic cyclic hydrocarbon substituent comprising from 3 to 20 ring atoms (i.e., a "3-20-membered heterocyclyl"), wherein one or more of the ring atoms is a heteroatom selected from nitrogen, oxygen or S(O)_m (wherein _m is an integer from 0 to 2), but excluding ring portions of -O-O-, -O-S- or -S-S-), with the remaining ring atom(s) being carbon. Preferably, it comprises 3 to 12 ring atoms (i.e., a "3-12 membered heterocyclic group"), 1 to 4 of which are heteroatoms; more preferably, the cycloalkyl ring comprises 3 to 10 ring atoms (i.e., a "3-10 membered heterocyclic group"). Non-limiting examples of monocyclic heterocyclic groups (e.g., 3-7-membered heterocyclic groups) include pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, homopiperazinyl, and the like. Polycyclic heterocyclic groups include spiro, fused and bridged heterocyclic groups.

The term "cycloalkylalkyl" means that the alkyl group is substituted with one or more cycloalkyl groups, preferably with one cycloalkyl group, wherein alkyl is defined as above and wherein cycloalkyl is defined as above, for example, C3-C8 cycloalkyl C1-C6 alkyl.

The term "haloalkyl" means an alkyl group substituted with one or more halogens, wherein the alkyl group is as defined above, for example, halo C1-C6 alkyl.

The term "deuteroalkyl" means an alkyl group substituted with one or more deuterium atoms, wherein the alkyl group is as defined above, for example, deutero C1-C6 alkyl.

The term "C6-C12 aryl" refers to carbocyclic aromatic system

groups having 6-12 carbon atoms.

The term "C6-C10 aryl" refers to carbocyclic aromatic system groups having 6-10 carbon atoms, such as phenyl, naphthyl, etc.

The term "5-10-membered heteroaryl" refers to aromatic
5 heterocyclic rings, typically 5-, 6-, 7-, 8-, 9- and 10-membered
heterocyclic rings having 1 to 3 heteroatoms selected from N,
O, or S; the heteroaryl ring can optionally be further fused or
attached to aromatic and non-aromatic carbon rings and
heterocyclic rings. Non-limiting examples of said 5- to 10-
10 membered heteroaryl rings are, for example, pyridinyl, pyrazinyl,
pyrimidinyl, pyridazinyl, indolyl, imidazolyl, thiazolyl,
isothiazolyl, thioxazolyl, pyrrolyl, phenyl-pyrrolyl, furanyl,
phenyl-furanyl, oxazolyl, isoxazolyl, pyrazolyl, thiophenyl,
benzofuranyl, benzothiophenyl, benzo 1,3-dioxolane
15 (benzodioxole), isodihydroindolyl, benzimidazolyl, indazolyl,
quinolinyl, isoquinolinyl, 1,2,3-triazolyl, 1-phenyl-1,2,3-
triazolyl, 2,3-dihydroindolyl, 2,3-dihydrobenzofuranyl, 2,3-
dihydrobenzothiophenyl, benzopyranyl, 2,3-dihydrobenzoxazinyl,
2,3-dihydroquinoxalinyl, and others.

The term "substituted C6-C10 aryl" or "substituted 5-10
20 membered heteroaryl" or "substituted 3-7 membered heterocyclyl"
means that a hydrogen in the aryl or heteroaryl or heterocyclic
group is replaced by a substituent group. Unless otherwise stated
in the text, the substituent of the aryl or heteroaryl or
25 heterocyclyl group may be a variety of groups selected from the
following group: -halogen, -OR', -NR'R'', -SR', -SiR'R''R''', -
OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -
NR'-C(O)NR''R''', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-
C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NR'S(O)₂R'', -CN and
30 -NO₂, with substituent numbers from 0 to (2m'+1), where m' is the
total number of carbon atoms in the group. R', R'' and R''' each
independently refer to hydrogen, unsubstituted C₁₋₈ alkyl,
unsubstituted C6-C12 aryl (or C6-C10 aryl), C6-C12 aryl (or C6-
C10 aryl) substituted by 1-3 halogens, unsubstituted C₁₋₈ alkyl,
35 C₁₋₈ alkoxy or C₁₋₈ thioalkoxy, or unsubstituted C6-C12 aryl (or C6-

C₁₀ aryl)-C₁₋₄alkyl. When R' and R'' are attached to the same nitrogen atom, they may form a 3-, 4-, 5-, 6-, or 7-membered ring with that nitrogen atom. For example, -NR'R'' includes 1-pyrrolidinyl and 4-morpholinyl.

5 The term "hydroxyl" refers to the -OH group.

The term "halogen" means fluorine, chlorine, bromine, or iodine.

The term "amino" means -NH₂. The term "nitro" means -NO₂.

The term "amide group" means -C(O)N(alkyl) or (cycloalkyl), wherein alkyl and cycloalkyl are as defined above.

10 The term "carboxylate group" means -C(O)O(alkyl) or (cycloalkyl), wherein alkyl and cycloalkyl are as defined above.

The present invention also includes various deuterated forms of formula I. Each of the available hydrogen atoms attached to the carbon atoms may be independently replaced by a deuterium atom.

15 A person skilled in the art can synthesize the deuterated form of formula I with reference to the relevant literature. Commercially available deuterated starting materials may be used in the preparation of deuterated forms of formula I, or they may be synthesized by conventional techniques using deuterated reagents, non-limiting examples of deuterated reagents including:
20 deuteroborane, trideuteroborane tetrahydrofuran solution, deuterated lithium-aluminum hydride, deuterated ethyl iodide, and deuterated methyl iodide, and the like.

The term "antibody" refers to immunoglobulins, which are
25 tetrapeptide chain structures consisting of two identical heavy chains and two identical light chains linked by interchain disulfide bonds. Immunoglobulins differ in the composition and order of amino acids in the constant region of the heavy chain, and therefore differ in their antigenicity. Accordingly,
30 immunoglobulins can be categorized into five classes, or isoforms of immunoglobulins, i.e., IgM, IgD, IgG, IgA, and IgE, whose corresponding heavy chains are μ , δ , γ , α , and ϵ chains, respectively. The same class of Ig can be further divided into different subclasses according to differences in the amino acid
35 composition of its hinge region and the number and positions of

disulfide bonds of the heavy chain, e.g. IgG can be divided into IgG1, IgG2, IgG3 and IgG4. The light chains are divided into κ -chains or λ -chains by the differences in the constant region. Each of the five classes of Ig may have a κ chain or a λ chain. The antibodies described in the present invention are preferably specific antibodies against cell surface antigens on target cells, and non-limiting embodiments are the following antibodies: one or more of anti-EGFRvIII antibody, anti-DLL-3 antibody, anti-PSMA antibody, anti-CD70 antibody, anti-MUC16 antibody, anti-ENPP3 antibody, anti-TDGF1 antibody, anti-ETBR antibody, anti-MSLN antibody, anti-TIM-1 antibody, anti-LRRC15 antibody, anti-LIV-1 antibody, anti-CanAg/AFP antibody, anti-cladin 18.2 antibody, anti-Mesothelin antibody, anti-HER2 (ErbB2) antibody, anti-EGFR antibody, anti-c-MET antibody, anti-SLITRK6 antibody, anti-KIT/CD117 antibody, anti-STEAP1 antibody, anti-SLAMF7/CS1 antibody, anti-NaPi2B/SLC34A2 antibody, anti-GPNMB antibody, anti-HER3(ErbB3) antibody, anti-MUC1/CD227 antibody, anti-AXL antibody, anti-CD166 antibody, anti-B7-H3(CD276) antibody, anti-PTK7/CCK4 antibody, anti-PRLR antibody, anti-EFNA4 antibody, anti-5T4 antibody, anti-NOTCH3 antibody, anti-Nectin 4 antibody, anti-TROP-2 antibody, anti-CD142 antibody, anti-CA6 antibody, anti-GPR20 antibody, anti-CD174 antibody, anti-CD71 antibody, anti-EphA2 antibody, anti-LYPD3 antibody, anti-FGFR2 antibody, anti-FGFR3 antibody, anti-FR α antibody, anti-CEACAMs antibody, anti-GCC antibody, anti-Integrin Av antibody, anti-CAIX antibody, anti-P-cadherin antibody, anti-GD3 antibody, anti-Cadherin 6 antibody, anti-LAMP1 antibody, anti-FLT3 antibody, anti-BCMA antibody, anti-CD79b antibody, anti-CD19 antibody, anti-CD33 antibody, anti-CD56 antibody, anti-CD74 antibody, anti-CD22 antibody, anti-CD30 antibody, anti-CD37 antibody, anti-CD138 antibody, anti-CD352 antibody, anti-CD25 antibody or anti-CD123 antibody.

The term "solvate" or "solvent compound" refers to the formation of a pharmaceutically usable solvate from the ligand-drug conjugate of the present invention with one or more solvent molecules, non-limiting examples of solvent molecules including

water, ethanol, acetonitrile, isopropanol, DMSO and ethyl acetate.

The term "drug load" refers to the average number of cytotoxic drugs loaded on each antibody in formula I. It can also be expressed as the ratio of drug amount to antibody amount, and the drug load can range from 0-12, preferably 1-10 cytotoxic drugs (D) connected to each antibody (Ab). In embodiments of the invention, the drug load is denoted as n , which exemplarily may be the mean value of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10. The average number of drugs per ADC molecule after the coupling reaction can be identified by conventional methods such as UV/visible spectroscopy, mass spectrometry, ELISA tests and HPLC characteristics.

In one embodiment of the present invention, the cytotoxic drug is coupled to the open cysteine sulfhydryl-SH and/or the sulfhydryl-SH of the site-directed mutagenesis cysteine residue between the antibody chains by a linker unit, and generally the number of drug molecules that can be coupled to the antibody in the coupling reaction will be less than or equal to a theoretical maximum.

The loading of ligand cytotoxic drug conjugates can be controlled by the following non-limiting methods, including:

- (1) controlling the molar ratio of the linking reagent to the monoclonal antibody;
- (2) control of reaction time and temperature;
- (3) selection of different reaction reagents.

For the preparation of conventional pharmaceutical compositions, see the Chinese Pharmacopoeia.

The term "pharmaceutically acceptable salt" or "pharmaceutically usable salt" means a salt of a ligand-drug conjugate of the present invention, or a salt of a compound described in the present invention; salts of this kind are safe and efficacious when used in a mammal, and are biologically active as desired. The ligand-drug conjugates of the present invention contain at least one carboxyl group, so they can form salts with bases, and non-limiting examples of pharmaceutically acceptable salts include sodium, potassium, calcium, or magnesium salts, etc.

The term "pharmaceutically acceptable salt" or "pharmaceutically usable salt" means a salt of an antibody-drug conjugate of the present invention, or a salt of a compound described herein; salts of this kind are safe and efficacious when
5 used in a mammal, and are biologically active as desired. The ligand-drug conjugates of the present invention contain at least one amino group and can therefore form salts with acids, non-limiting examples of pharmaceutically acceptable salts including: hydrochloride, hydrobromide, hydriodate, sulfate, bisulfate,
10 citrate, acetate, succinate, ascorbate, oxalate, nitrate, phosphate, hydrogen phosphate, dihydrogen phosphate, salicylate, hydrogen citrate, tartrate, maleate, fumarate, formate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate.

15 "Acidic amino acid" refers to amino acids with an isoelectric point of less than 7. Acidic amino acid molecules tend to have one or more acidic groups, such as carboxyl groups, which can be effectively ionized into negative ionic forms in the structure to increase hydrophilicity. Acidic amino acids can be natural or
20 unnatural.


The term "natural amino acids" refers to biologically synthesized amino acids. Natural amino acids are generally of the L-type, but there are a few exceptions, such as glycine, including natural and synthesized by organisms.

25 "Unnatural amino acids" means amino acids obtained by synthetic means.

The invention is further elaborated below in connection with specific embodiments which, it should be understood, are intended only to illustrate the invention, and are not intended to limit
30 the scope of the invention. Test methods for which specific conditions are not indicated in the following embodiments are generally in accordance with conventional conditions or in accordance with conditions recommended by the manufacturer. All percentages, proportions, ratios or parts are by weight unless
35 otherwise indicated.

Example 1


Synthesis of compound M1:



In a 5000 mL single-necked flask, N-fluorenylmethoxycarbonyl-glycine-glycine (100 g, 282 mmol, 1.0 eq), lead tetraacetate (175 g, 395 mmol, 1.4 eq), 2000 mL of dry tetrahydrofuran and 670 mL of toluene were added, stirred uniformly, protected by nitrogen, heated to 85°C and reacted for 2.5 h. The reaction was monitored by TLC, and after the starting materials had finished reacting, cooling to room temperature and filtration were performed, the filtrate was concentrated under reduced pressure, and the residue was purified by column chromatography to obtain compound M1 (87 g); LC-MS: $[\text{M}+\text{NH}_4]^+ = 386.0$.

Example 2

Synthesis of compound M3:




In a 1000 mL single-necked flask, compound SM-2 (synthesized according to the method published in patent CN108452321A) (40 g, 96 mmol, 1.0 eq), triethylamine (26.7 mL, 2.0 eq), and toluene (400 mL) were added, heated to 120°C , and a reaction was carried out by refluxing for 2 h. When TLC monitoring indicated that the reaction was essentially complete, cooling to 50°C was performed, and solvent was removed by spinning under reduced pressure. Dissolution with ethyl acetate (150 mL) and water (40 mL) was performed, the pH was adjusted to 2-3 with 1M HCl under ice bath stirring, and liquid separation was performed. The aqueous layer was extracted once more with ethyl acetate, the organic layers were combined, and drying was performed by adding anhydrous sodium sulfate. After filtration, a light yellow oily crude product was obtained by concentration, and

the crude product was purified by column chromatography (DCM: MeOH = 40:1) to obtain compound M2 (26.6 g); LC-MS: $[M+H]^+ = 399.3$.

In a 1000 mL single-necked flask, compound M2 (26.5 g, 60.5 mmol, 1.0 eq), pentafluorophenol (12.2 g, 66.5 mmol, 1.1 eq), DCC (13.7 g, 66.5 mmol, 1.1 eq), and THF (300 mL) were added, and a reaction was carried out at room temperature for 30 min (monitored by TLC). The insoluble material was filtered out. The reaction solution was purified directly by preparative LC, and the prepared solution was concentrated at 35°C in a water bath under reduced pressure by pumping water to remove acetonitrile, and lyophilized to obtain compound M3 (31.5 g) with 64% yield; LC-MS: $[M+H]^+ = 565.1$.

Example 3


Synthesis of the compound ent-M3:



Referring to the synthetic route of Example 2, compound ent-M3 (27.8 g) was obtained; LC-MS: $[M+H]^+ = 565.2$.

Example 4

Synthesis of compound 1:



Step 1: Compound 1a

In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol), 100 mL THF, *p*-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, perform dropwise addition of benzyl hydroxyacetate (5.4 g, 32.6 mmol), then naturally warm to room temperature for a reaction (the reaction lasts for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO₃ solution was added, extracted with ethyl acetate, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, filtered, concentrated, and the residue was purified by silica gel column (PE:EA=10:1-5:1-1:1) to obtain 1a (4 g) at 52% yield; LC-MS: [M+H]⁺ = 475.18.

Step 2: Compound 1b

In a 25 mL single-necked flask, add 1a (2 g, 4.2 mmol), 10 mL DMF, stir at 0°C, add DBU (766 mg, 5.04 mmol), and react for 1 h. After the completion of Fmoc deprotection as detected by TLC, the reaction solution was set aside ready for use.

M4 (prepared according to the method published in patent CN111051330 A) (1.73 g, 4.2 mmol), PyBOP (2.61 g, 5.04 mmol), HOBT (680 mg, 5.04 mmol) and 10 mL of DMF were added to another 25 mL single-necked flask, DIPEA (830 μL, 5.04 mmol) was added

in an ice-water bath, stirring was continued for 30 min, the above reaction solution was added to the reaction flask, and the temperature was raised to room temperature for a reaction. When HPLC monitoring indicated the end of the reaction, the reaction solution was purified by preparative LC to obtain a product preparation liquid, the preparation liquid was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, filtered, and the filtrate was concentrated under reduced pressure to obtain solid 1b (1.7 g) at 63% yield; LCMS: $[M+H]^+ = 648.26$.

Step 3: Compound 1c

1b (900 mg, 1.39 mmol) was added to a 25 mL single-necked flask, and after dissolution with 15 mL of DMF, 900 mg of 5% Pd/C was added, and a hydrogenation reaction took place for 2 h. After the completion of the reaction, filtering was performed to obtain the filtrate, which was used directly for the next step of the reaction without purification.

Step 4: Compound 1d

The crude product 1c was placed in an ice-water bath, DIPEA (235 μ L, 1.39 mmol) was added, and compound M3 (784 mg, 1.39 mmol) was added, then the temperature was raised to room temperature to react for 1 h. When the reaction was complete as indicated by HPLC monitoring, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 1d (504 mg); LC-MS: $[M+H]^+ = 804.4$.

Step 5: Compound 1e


Add 1d (500mg, 0.62mmol), M5 (310mg, 0.62mmol), PyBOP (448mg, 0.86mmol), HOBt (116mg, 0.86mmol) and 15mL DMF into a 50mL single-necked flask, add DIPEA (378 μ L, 2.29 mmol) in an ice-water bath, raise to room temperature and react for 2h. After completion of the reaction was indicated by monitoring by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid of compound 1e, which was lyophilized to obtain 1e (210 mg); LC-MS: $[M+H]^+ = 1221.6$.

Step 6: Compound 1

1e (100 mg, 0.081 mmol), zinc bromide (368 mg, 1.63 mmol) and 5 mL of nitromethane were added to a 25 mL single-necked flask and the reaction was carried out at 40°C for 1 h. After the completion of the reaction was indicated by monitoring by HPLC, concentration under reduced pressure was performed to remove the solvent, and a crude product was obtained. The crude product was purified by HPLC to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 1 (60 mg); LC-MS: $[M+H]^+ = 1065.3$.

Example 5


Synthesis of compound 2:



Referring to the synthetic route of Example 4, compound 2 (51 mg) was obtained; LC-MS: $[M+H]^+ = 1065.3$.

Example 6

Synthesis of compound 3:



Step 1: Compound 3a

In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol), 100 mL THF, *p*-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, perform dropwise addition of benzyl 2-hydroxy-2-methylpropionate (6.3 g, 32.6 mmol), then perform natural warming to room temperature for a reaction (the reaction lasts for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO₃ solution was added, followed by extraction with ethyl acetate, washing with saturated sodium chloride solution, drying with anhydrous sodium sulfate, filtering and concentration, and the residue was purified by silica gel column (PE:EA=10:1-5:1-2:1) to obtain 3a (4.2 g) at 52% yield; LC-MS: [M+H]⁺ = 503.3.

Step 2: Compound 3b

In a 25 mL single-necked flask, add 3a (2 g, 4.0 mmol) and 10 mL DMF, stir at 0°C, add DBU (760 mg, 5.0 mmol), and react for 1 h. After the completion of Fmoc deprotection as detected by TLC, the reaction solution was set aside ready for use.

Add M4 (1.65 g, 4.0 mmol), PyBOP (2.59 g, 5.0 mmol), HOBT (675 mg, 5.0 mmol) and 10 mL of DMF into another 25 mL single-necked flask, and add DIPEA (823 μ L, 5.04 mmol) in an ice-water

bath, continue stirring for 30 min, and then add the above reaction solution to the reaction flask and raise to room temperature for a reaction. After the completion of the reaction as indicated by monitoring by HPLC, the reaction solution was purified by preparative LC to obtain a product preparation liquid, which was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate and filtered, and the filtrate was concentrated under reduced pressure to obtain the solid 3b (1.4 g) at 53% yield; LC-MS: $[M+H]^+ = 676.2$.

Step 3: Compound 3c

Add 3b (700 mg, 1.04 mmol) to a 25 mL single-necked flask, dissolve in 10 mL of DMF, add 700 mg of 5% Pd/C, and perform a hydrogenation reaction for 1.5 h. After the completion of the reaction, filtering was performed to obtain a filtrate, which was used directly for the next step of the reaction without purification.

Step 4: Compound 3d

Place the crude product 3c in an ice-water bath, add DIPEA (210 μ L, 1.25 mmol) and then compound M3 (704 mg, 1.25 mmol), then raise to room temperature and react for 1 h. After the completion of the reaction as indicated by HPLC monitoring, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to give 3d (486 mg); LC-MS: $[M-H]^- = 830.5$.

Step 5: Compound 3e


Add 3d (300 mg, 0.36 mmol), M5 (180 mg, 0.36 mmol), PyBOP (260 mg, 0.5 mmol), HOBt (67 mg, 0.5 mmol) and 10 mL of DMF to a 50 mL single-necked flask, add DIPEA (219.5 μ L, 1.33 mmol) in an ice-water bath, raise to room temperature and react for 3 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to give the preparation liquid of compound 3e, which was lyophilized to obtain 3e (157 mg); LC-MS: $[M+H]^+ = 1249.6$.

Step 6: Compound 3

Add 3e (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane to a 25 mL single-necked flask, and react at 40°C for 1 h. After the completion of the reaction as detected by HPLC, perform concentration under reduced pressure to remove the solvent, to obtain a crude product. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 3 (64 mg); LC-MS: $[M+H]^+ = 1093.1$.

Example 7


Synthesis of compound 4:



Referring to the synthetic route of Example 6, compound 4 (60 mg) was obtained; LC-MS: $[M+H]^+ = 1093.2$.

Example 8

Synthesis of compound 5A:



Step 1: Compound 5a

Add M1 (500 mg, 1.4 mmol, 1.0 eq), p-toluenesulfonic acid monohydrate (26 mg, 0.1 mmol, 0.1 eq) and 10 mL of THF in a 25 mL single-necked flask, stir well, then lower to 0°C, and then slowly add L-lactic acid benzyl ester (1.2 g, 7.0 mmol, 5 eq), and then raise to room temperature for a reaction. TLC monitoring was performed, and at the end of the reaction, saturated NaHCO₃ solution was added, followed by extraction with ethyl acetate, drying over anhydrous sodium sulfate, filtering and concentration, and the residue was purified by reverse-phase column to give 5a (400 mg);

LC-MS: [M+NH₄]⁺ = 506.2.

¹H NMR (400 Mz, CDCl₃/CD₃OD): 1.39 (3H, d, J = 6.8 Hz), 3.78 (2H, t, J = 4.0 Hz), 4.17-4.27 (2H, m), 4.42 (2H, d, J = 4.0 Hz), 4.72-4.85 (2H, m), 5.11-5.58 (2H, m), 5.43 (1H, s), 7.06 (1H, t, J = 8.0 Hz), 7.25-7.33 (6H, m), 7.38 (2H, t, J = 8.0 Hz), 7.57 (2H, d, J = 8.0 Hz), 7.75 (2H, d, J = 8.0 Hz).

Step 2: Compound 5b

Add Compound 5a (400 mg, 0.8 mmol, 1.0 eq) and 4 mL of DMF to a 25 mL single-necked flask, stir well, and then lower to 0°C before slowly adding DBU (137 mg, 0.9 mmol, 1.1 eq). After the completion of the addition, raise to room temperature for a reaction. The reaction was monitored by TLC, and at the end of the reaction, the reaction solution was recorded as reaction solution①;

To another 25 mL single-necked flask, add M4 (372 mg, 0.9 mmol, 1.1 eq), PyBOP (852 mg, 1.6 mmol, 2.0 eq) and 3 mL of DMF, stir at room temperature for 5 min, and add reaction solution①. The reaction was carried out at room temperature and monitored by HPLC. When the reaction was completed, the reaction solution was purified through HPLC to yield compound 5b (326 mg); LC-MS: [M+NH₄]⁺ = 679.2.

Step 3: Compound 5c

Add 5b (4.0 g, 6.05 mmol, 1.0 eq) to a 100 mL single-necked flask, dissolve in DMF (60 mL), then add 5% Pd/C (4 g), and perform a hydrogenation reaction at room temperature for 4 h

(HPLC was used to monitor the progress of the reaction). The Pd/C was filtered, and the filtrate was placed directly in an ice-water bath (about 0°C) without being concentrated, ready for use.

5 Step 4: Compound 5d

Place the crude product 5c in an ice-water bath, add DIPEA (1.1 mL, 1.1 eq) and then compound M3 (3.4 g, 6.05 mmol), and then raise to room temperature and react for 2 h. After the completion of the reaction as detected by HPLC, the reaction
10 solution was purified by HPLC to obtain a preparative liquid, which was lyophilized to obtain 5d (3.15 g); LC-MS: $[M-H]^- = 816.3$.

Step 5: Compound 5e

Add 5d (2.07g, 2.53mmol, 1.0eq), M5 (1.35g, 2.53mmol, 1.0eq),
15 PyBOP (1.98g, 3.79mmol, 1.5eq), HOBt (0.51g, 3.79mmol, 1.5eq) and DMF (40mL) to a 100 mL single-necked flask, add DIPEA (1.05mL, 1.5eq) in an ice-water bath, raise to room temperature and react for 2h (monitored by HPLC). The reaction solution was directly purified by preparative LC, and the preparation
20 liquid was concentrated in a water bath under reduced pressure by a water pump at 35°C to remove acetonitrile and freeze-dried to obtain compound 5e (1.92g), with a yield of 61%; LC-MS: $[M+H]^+ = 1235.4$.

Step 6: Compound 5A


25 To a 100 mL single-necked flask, add compound 5e (1.0 g, 0.8 mmol, 1.0 eq) and 35 mL nitromethane, and after dissolution, add zinc bromide (3.64 g, 16 mmol, 20.0 eq), react for 30 min in an oil bath at 40°C (stabilized in advance by preheating), and concentrate at 45°C in a water bath under reduced pressure
30 by a water pump to remove the nitromethane, yielding a yellow residue solid (monitored by HPLC). After acid preparation, the preparation liquid of compound 5A was obtained, and was concentrated at 35°C in a water bath under reduced pressure by a water pump to remove acetonitrile by spinning, and lyophilized
35 to obtain compound 5A (786 mg) at a yield of 90%.

LC-MS: $[M+H]^+ = 1079.4$;

^1H NMR (400 MHz, DMSO- d_6) δ 9.39 - 9.02 (m, 1H), 8.70 (t, J = 6.5 Hz, 1H), 8.64 (t, J = 5.7 Hz, 1H), 8.56 (d, J = 8.8 Hz, 1H), 8.34 (t, J = 5.7 Hz, 1H), 8.16 (d, J = 8.2 Hz, 1H), 8.01 (t, J = 5.5 Hz, 1H), 7.71 (d, J = 10.9 Hz, 1H), 7.30 (s, 1H), 7.28 - 7.15 (m, 4H), 7.14 (s, 2H), 5.53 (dd, J = 14.5, 6.4 Hz, 1H), 5.49 - 5.34 (m, 2H), 5.22 (d, J = 18.8 Hz, 1H), 5.09 (d, J = 18.7 Hz, 1H), 5.03 (dd, J = 9.6, 3.9 Hz, 1H), 4.73 (dd, J = 9.9, 6.9 Hz, 1H), 4.59 (dd, J = 10.1, 6.5 Hz, 1H), 4.49 (ddd, J = 13.2, 8.6, 4.4 Hz, 1H), 4.14 (dd, J = 13.3, 6.6 Hz, 2H), 3.93 (s, 2H), 3.84 (dd, J = 16.5, 6.3 Hz, 1H), 3.76 (dd, J = 16.9, 5.7 Hz, 2H), 3.70 (dd, J = 5.2 Hz, 2H), 3.60 (dd, J = 16.7, 5.4 Hz, 1H), 3.52 (dd, J = 16.4, 5.1 Hz, 1H), 3.45 (dd, J = 12.8, 10.1 Hz, 1H), 3.25 - 3.15 (m, 1H), 3.14 - 3.05 (m, 1H), 3.01 (dd, J = 13.7, 4.1 Hz, 1H), 2.73 (dd, J = 13.5, 9.8 Hz, 1H), 2.54 - 2.47 (m, 1H), 2.33 (s, 2H), 2.17 (d, J = 5.5 Hz, 2H), 1.91 - 1.79 (m, 2H), 1.33 (d, J = 6.6 Hz, 2H), 0.87 (t, J = 7.3 Hz, 2H).

Example 9

Synthesis of compound 5B:



Step 1: Compound 5d-1

Add compound 5b (300mg, 0.45mmol, 1.0eq) and DMF (3mL) to a 25mL single-necked flask, stir to dissolve, add 5% Pd/C (300mg), perform hydrogen replacement three times, perform a hydrogenation reaction for 2h, and monitor for the end of the

reaction by HPLC. After the reaction is over, remove Pd/C by filtration, cool the filtrate to 0-5°C, add DIPEA (65mg, 0.5mmol, 1.1eq), then add ent-M3 (255mg, 0.45mmol) to the filtrate, then raise to 20±5°C and react for 1 h, and monitor for the end of the reaction by HPLC. After the completion of the reaction, preparative purification was performed by HPLC, and the product preparation liquid was collected and lyophilized to obtain compound 5d-1 (200 mg), with a yield of 54%; LC-MS: [M-H]⁻ = 816.3.

Step 2: Compound 5e-1


Add compound 5d-1 (200mg, 0.24mmol, 1.0eq), M5 (127mg, 0.24mmol, 1.0eq), PyBOP (187mg, 0.36mmol, 1.2eq), HOBt (48mg, 0.36mmol, 1.2 eq) and DMF (6mL) into a 25mL single-necked flask, lower the temperature in an ice-water bath to 0-5°C, and add DIPEA (62mg, 0.48mmol, 2.0eq). After the addition is completed, raise to 20±5°C and react for 2h, and use HPLC to monitor for the completion of the reaction. The reaction solution underwent preparative purification by HPLC, and the product preparation liquid was collected and lyophilized to obtain compound 5e-1 (162.8 mg); LC-MS: [M+H]⁺ =1235.4.

Step 3: Compound 5B

Sequentially add compound 5e-1 (110 mg, 0.089 mmol, 1.0 eq), ZnBr₂ (400 mg, 1.78 mmol, 20.0 eq) and CH₃NO₂ (10 mL) into a 25 mL single-necked flask. After the addition, raise to 40°C and react for 0.5 h, and then stop the reaction. The reaction solution was dried directly at 45°C by spin-drying under reduced pressure to obtain a yellow solid. Samples were taken and the reaction was monitored by HPLC. The spin-dried solid was directly purified by HPLC preparation, and the product preparation liquid was collected and lyophilized to obtain compound 5B (73.4 mg) at 76.5% yield; LC-MS: [M+H]⁺ = 1079.4.

Example 10


Preparation of compound 6A:



Referring to the synthetic route of Example 8, compound 6A (71 mg) was obtained; LC-MS: $[M+H]^+ = 1079.4$.

Example 11


5 Preparation of compound 6B:



Referring to the synthetic route of Example 9, compound 6B (59 mg) was obtained; LC-MS: $[M+H]^+ = 1079.4$.

Example 12

10 Preparation of compounds 7A and 7B:



Step 1: Compound 7a

In a 250 mL single-necked flask, add M1 (10 g, 27.1 mmol),

benzyl 3,3,3-trifluorolactate (prepared according to the method published in patent W02020063673A1) (12.7 g, 54.3 mmol), zinc acetate (9.96 g, 54.3 mmol) and 100 mL of toluene, heat to 100°C and react for 4 h. When the reaction is completed, reduce to room temperature, filter to remove insoluble material, and concentrate the filtrate to obtain a crude product. The crude product was purified by silica gel column chromatography (PE:EA=10:1-5:1-2:1) to obtain 5.15 g of the target material, yield 35.1%; LC-MS: $[M+H]^+ = 543.17$.

Step 2: Compound 7b

Add 7a (5 g, 9.2 mmol) and 15 mL of DMF in a 50 mL single-necked flask, dissolve until clear, and then add DBU (1.68 g, 11 mmol) in an ice-water bath and react for 1 h, and record the reaction solution as reaction solution ①.

Take another 50 mL single-necked flask, add M4 (3.8 g, 9.2 mmol), PyBOP (5.75 g, 11 mmol), HOBT (1.49 g, 11 mmol) and 10 mL DMF. When dissolved, add DIPEA (1.82 mL, 11 mmol) in an ice-water bath, and continue the reaction for 30 min, then add reaction solution ①, and raise to room temperature and react for 2 h. The reaction progress was monitored by HPLC. After the completion of the reaction, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid. The preparation liquid was extracted with dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, filtered, and concentrated to obtain 4.1 g of solid, with a yield of 62.3%; LC-MS: $[M+H]^+ = 716.25$.

Step 3: Compound 7d


Add 7b (900 mg, 1.26 mmol) in a 25 mL single-necked flask, and after dissolution with 15 mL of DMF, add 900 mg of 5% Pd/C, and perform a hydrogenation reaction for 2 h. After the completion of the reaction, perform filtration, place the filtrate in an ice-water bath, add DIPEA (228 μ L, 1.38 mmol), and then add M3 (712 mg, 1.26 mmol), and then raise to room temperature and react for 1 h. When the reaction was completed

as detected by HPLC, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid, which was lyophilized to obtain 525 mg of a product at 47.9% yield; LC-MS: $[M-H]^- = 870.33$.

5 Step 4: Compound 7e


Add 7d (500 mg, 0.57 mmol), M5 (305 mg, 0.57 mmol), PyBOP (448 mg, 0.86 mmol), HOBT (116 mg, 0.86 mmol) and 15 mL of DMF to a 50 mL single-necked flask, add DIPEA (378 μ L, 2.29 mmol) in an ice-water bath, raise to room temperature and react for 2 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by high performance liquid chromatography to obtain the preparation liquids of compound 7e-1 and compound 7e-2, and the preparation liquids were lyophilized to obtain 150 mg of compound 7e-1, LC-MS: $[M+H]^+ = 1289.46$, and 220 mg of compound 7e-2, LC-MS: $[M+H]^+ = 1289.46$, respectively.

Step 5: Compound 7A



Add 7e-1 (100 mg, 0.077 mmol), zinc bromide (349 mg, 1.55 mmol) and 5 mL of nitromethane to a 25 mL single-necked flask, and react at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 52 mg of solid; TOF result: 1133.3613.


Step 6: Compound 7B



Add 7e-2 (100 mg, 0.077 mmol), zinc bromide (349 mg, 1.55 mmol) and 5 mL of nitromethane to a 25 mL single-necked flask and react at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 63 mg of solid; TOF result: 1133.3668.

Example 13

Synthesis of compounds 8A and 8B:



Step 1: Compound 8d


In a 25 mL single-necked flask, 7c (900 mg, 1.83 mmol) was added, 20 mL of DMF was used to dissolve, then DIPEA (303 μ L, 1.83 mmol) was added, and ent-M3 (1034 mg, 1.83 mmol) was added, and then the temperature was raised to room temperature to react for 1 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by high-performance liquid chromatography to obtain a preparation liquid, which was lyophilized to give 613 mg of a product at 38.5% yield; LC-MS: $[M-H]^- = 870.32$.

Step 2: Compound 8e-1 and Compound 8e-2

Add 8d (500 mg, 0.57 mmol), M5 (305 mg, 0.57 mmol), PyBOP (448 mg, 0.86 mmol), HOBT (116 mg, 0.86 mmol) and 15 mL of DMF into a 50 mL single-necked flask, add DIPEA (378 μ L, 2.29 mmol) in an ice-water bath, and raise to room temperature to react for 2 h. After the completion of the reaction as detected by


HPLC, the reaction solution was purified by high performance liquid chromatography to obtain the preparation liquids of compound 8e-1 and compound 8e-2, and the preparation liquids were lyophilized to obtain 140 mg of compound 8e-1 and 210 mg of compound 8e-2, respectively. LC-MS of compound 8e-1: $[M+H]^+ = 1289.47$; LC-MS of compound 8e-2: $[M+H]^+ = 1289.47$.

Step 3: Compound 8A



Add Compound 8e-1 (100 mg, 0.077 mmol), zinc bromide (349 mg, 1.55 mmol) and 5 mL of nitromethane into a 25 mL single-necked flask, and react at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 50 mg of solid; TOF result: 1133.3623.


Step 4: Compound 8B



Add Compound 8e-2 (100 mg, 0.077 mmol), zinc bromide (349 mg, 1.55 mmol) and 5 mL of nitromethane into a 25 mL single-necked flask and react at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 58 mg of a solid; TOF result: 1133.3653.

Example 14

Synthesis of compound 9A:



Step 1: Compound 9a

5 In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol), 100 mL THF and p-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, add benzyl 2-hydroxy-2-cyclopropyl acetate dropwise (prepared according to the method published in patent US20050020645 A1) (6.3 g, 32.6 mmol), then naturally
 10 warm up to room temperature to react (the reaction lasts for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO₃ solution was added, followed by extraction with ethyl acetate, washing with saturated sodium chloride solution, drying with anhydrous sodium sulfate, filtering and
 15 concentration, and the residue was purified by silica gel column (PE:EA=10:1-5:1-2:1) to give 9a (3.7 g) at 45% yield; LC-MS: [M+H]⁺ = 501.5.

Step 2: Compound 9b

20 In a 25 mL single-necked flask, add 9a (2 g, 4.0 mmol) and 10 mL DMF, stir at 0°C, add DBU (760 mg, 5.0 mmol), and react for 1 h. After the completion of TLC-monitored Fmoc deprotection, the reaction solution was set aside, ready for use.

Add M4 (1.65 g, 4.0 mmol), PyBOP (2.59 g, 5.0 mmol), HOBt

(675 mg, 5.0 mmol) and 10 mL of DMF into another 25 mL single-necked flask, add DIPEA (823 μ L, 5.04 mmol) in an ice-water bath, continue stirring for 30 min, and then add the above reaction solution to the reaction flask and raise to room temperature to react. After completion of the reaction as detected by HPLC, the reaction solution was purified by preparative LC to obtain a product preparation liquid, which was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, and filtered, and the filtrate was concentrated under reduced pressure to obtain 1.5 g of a solid at 56% yield; LC-MS: $[M+H]^+ = 674.7$.

Step 3: Compound 9c

9b (900 mg, 1.3 mmol) was added into a 25 mL single-necked flask, and after dissolution with 10 mL of DMF, 900 mg of 5% Pd/C was added, and a hydrogenation reaction was carried out for 1.5 h. After the completion of the reaction, filtration was performed to obtain the filtrate, which was used directly for the next step of the reaction without purification.

Step 4: Compound 9d

Place the crude product 9c in an ice-water bath, add DIPEA (223 μ L, 1.3 mmol) and then compound M3 (750 mg, 1.3 mmol), then raise to room temperature and react for 1 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to give a preparation liquid, which was lyophilized to give 9d (529 mg); LC-MS: $[M-H]^- = 828.4$.

Step 5: Compound 9e

Add 9d (500 mg, 0.6 mmol), M5 (300 mg, 0.6 mmol), PyBOP (416 mg, 0.8 mmol), HOBt (108 mg, 0.5 mmol) and 15 mL of DMF into a 50 mL single-necked flask, add DIPEA (351 μ L, 2.13 mmol) in an ice-water bath, raise to room temperature and react for 3 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by high performance liquid chromatography to obtain the preparative liquid of compound 9e, which was lyophilized to obtain 9e (257 mg); LC-MS: $[M+H]^+ =$


1247.5.

Step 6: Compound 9A

Add 9e (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane to a 25 mL single-necked flask and carry out a reaction at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 9A (55 mg); LC-MS: $[M+H]^+ = 1091.3$.

Example 15


Synthesis of compound 9B:



Referring to the synthetic route of Example 14, compound 9B (44 mg) was obtained; LC-MS: $[M+H]^+ = 1091.3$.

Example 16

Synthesis of compound 10A:



Step 1: Compound 10a

In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol), 100 mL THF and p-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, add benzyl 3-hydroxy-2-cyclopropylpropionate dropwise (prepared with reference to the method published in patent W02013187496A1) (6.7 g, 32.6 mmol), then naturally warm up to room temperature to react (react for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO₃ solution was added, followed by extraction with ethyl acetate, washing with saturated sodium chloride solution, drying with anhydrous sodium sulfate, filtering and concentration, and the residue was purified by silica gel column (PE:EA=10:1-5:1-2:1) to give 10a (4.9 g) at 58% yield; LC-MS: [M+H]⁺ = 515.4.

Step 2: Compound 10b

In a 25 mL single-necked flask, add 10a (4 g, 7.8 mmol) and 10 mL DMF, stir at 0°C, add DBU (1.2 g, 8.0 mmol), and react for 1 h. After the completion of TLC-monitored Fmoc deprotection, set aside the reaction solution, ready for use.

Add M4 (3.3 g, 8.0 mmol), PyBOP (5.2 g, 10.0 mmol), HOBT (1.35 g, 10.0 mmol) and 10 mL of DMF into another 25 mL single-necked flask, and add DIPEA (1.65 mL, 10.1 mmol) in an ice-water bath, continue stirring for 50 min, and then add the above reaction solution to the reaction flask, raise to room temperature and react. After the completion of the reaction as detected by HPLC, the reaction solution was purified by preparative LC to obtain a product preparation liquid, which was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate and filtered, and the filtrate was concentrated under reduced pressure to obtain 2.3 g of a solid at 42% yield; LC-MS: [M+H]⁺ = 688.8.

Step 3: Compound 10c

Add 10b (1.0 g, 1.45 mmol) into a 25 mL single-necked flask, dissolve in 15 mL of DMF until clear, then add 1.0 g of 5% Pd/C, and perform a hydrogenation reaction for 1.5 h. After the

completion of the reaction, filtration was performed to obtain a filtrate, which was used directly for the next step of the reaction without purification.

Step 4: Compound 10d

5 The crude product 10c was placed in an ice-water bath, DIPEA (258 μ L, 1.5 mmol) was added, and compound M3 (837 mg, 1.45 mmol) was then added, and then the temperature was raised to room temperature to react for 1 h. Upon completion of the reaction as detected by HPLC, the reaction solution was purified
10 by HPLC to obtain a preparation liquid, which was lyophilized to obtain 10d (499 mg); LC-MS: $[M-H]^- = 842.4$.

Step 5: Compound 10e


10d (400 mg, 0.48 mmol), M5 (240 mg, 0.48 mmol), PyBOP (250 mg, 0.48 mmol), HOBT (104 mg, 0.48 mmol) and 15 mL of DMF were
15 added into a 50 mL single-necked flask, DIPEA (330 μ L, 2.0 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 3 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to give the preparation liquid of compound 10e,
20 which was lyophilized to obtain 10e (188 mg); LC-MS: $[M+H]^+ = 1261.5$.

Step 6: Compound 10A

10e (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane were added to a 25 mL single-necked flask
25 and a reaction was carried out at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product
30 preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 10A (61 mg); LC-MS: $[M+H]^+ = 1105.4$.

Example 17


Synthesis of compound 10B:



Referring to the synthetic route of Example 16, compound 10B (75 mg) was obtained; LC-MS: $[M+H]^+ = 1105.4$.

Example 18

5 Synthesis of compound 11A:



Step 1: Compound 11a

In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol), 100 mL THF and p-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, add benzyl 2-hydroxy-2-cyclobutylacetate dropwise (synthesized by the method published in the literature *Journal of Medicinal Chemistry*, 2013, 56(13), 5541-5552) (6.7 g, 32.6 mmol), then naturally warm up to room temperature and react (react for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO_3 solution was added, followed by extraction with ethyl acetate, washing with saturated sodium chloride solution, drying with anhydrous sodium sulfate, filtering and concentration, and the

residue was purified by silica gel column (PE:EA=10:1-5:1-2:1) to give 11a (5.1 g) at 62% yield; LC-MS: $[M+H]^+ = 515.7$.

Step 2: Compound 11b

In a 25 mL single-necked flask, add 11a (4 g, 7.8 mmol) and 10 mL DMF, stir at 0°C, add DBU (1.2 g, 8.0 mmol), and react for 1 h. After the completion of TLC-monitored Fmoc deprotection, set aside the reaction solution, ready for use.

Add M4 (3.3 g, 8.0 mmol), PyBOP (5.2 g, 10.0 mmol), HOBt (1.35 g, 10.0 mmol) and 10 mL of DMF into another 25 mL single-necked flask, and add DIPEA (1.63 mL, 10.0 mmol) in an ice-water bath, continue stirring for 40 min, and then add the above reaction solution to the reaction flask, and raise to room temperature to react. After the completion of the reaction as detected by HPLC, the reaction solution was purified by preparative LC to obtain a product preparation liquid, which was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate and filtered, and the filtrate was concentrated under reduced pressure to obtain 2.3 g of a solid at 42% yield; LC-MS: $[M+H]^+ = 688.3$.

Step 3: Compound 11c

Add 11b (2.0 g, 2.9 mmol) into a 25 mL single-necked flask, and after dissolution with 25 mL of DMF, add 2.0 g of 5% Pd/C, and carry out a hydrogenation reaction for 3 h. After the completion of the reaction, it was filtered to obtain a filtrate, which was used directly for the next step of the reaction without purification.

Step 4: Compound 11d

The crude product 11c was placed in an ice-water bath, DIPEA (516 μ L, 3.0 mmol) was added, and then compound M3 (1.7 g, 2.9 mmol) was added, and then the temperature was raised to room temperature to react for 2 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to give 11d (934 mg); LC-MS: $[M-H]^- = 842.4$.

Step 5: Compound 11e


Add 11d (800 mg, 0.96 mmol), M5 (480 mg, 0.96 mmol), PyBOP (500 mg, 0.96 mmol), HOBT (208 mg, 0.96 mmol) and 30 mL of DMF into a 50 mL single-necked flask, add DIPEA (660 μ L, 4.0 mmol) in an ice-water bath, raise to room temperature and react for 4 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to give the preparation liquid of compound 11e, which was lyophilized to obtain 11e (401 mg); LC-MS: $[M+H]^+ = 1261.4$.

Step 6: Compound 11A

11e (150 mg, 0.12 mmol), zinc bromide (532 mg, 2.4 mmol) and 10 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 11A (86 mg); LC-MS: $[M+H]^+ = 1105.4$.

Example 19


Synthesis of compound 11B



Referring to the synthetic route of Example 18, compound 11B (50 mg) was obtained. LC-MS: $[M+H]^+ = 1105.4$.

Example 20

Synthesis of compound 12A:



Step 1: Compound 12a

In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol), 100 mL THF and p-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, add benzyl 3-hydroxy-2-cyclobutylpropionate dropwise (prepared according to the method published in patent W02009011285A1) (7.2 g, 32.6 mmol). Then warm up to room temperature naturally to react (react for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO₃ solution was added, followed by extraction with ethyl acetate, washing with saturated sodium chloride solution, drying with anhydrous sodium sulfate, filtering and concentration, and the residue was purified by silica gel column (PE:EA=10:1-5:1-2:1) to give 12a (4.5 g) at 52% yield; LC-MS: [M+H]⁺ = 529.4.

Step 2: Compound 12b

In a 25 mL single-necked flask, add 12a (4 g, 7.6 mmol) and 10 mL DMF, stir at 0°C, add DBU (1.2 g, 8.0 mmol), and react for 1 h. After the completion of TLC-monitored Fmoc deprotection, set aside the reaction solution, ready for use.

Add M4 (3.2 g, 7.6 mmol), PyBOP (4.7 g, 9.0 mmol), HOBT (1.22 g, 9.0 mmol) and 10 mL of DMF in another 25 mL single-necked flask, add DIPEA (1.49 mL, 0.9 mmol) in an ice-water

bath, and continue stirring for 30 min, then add the above reaction solution to the reaction flask and raise the temperature to room temperature to react. After the completion of the reaction as detected by HPLC, the reaction solution was purified by preparative LC to obtain a product preparation liquid, which was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate and filtered, and the filtrate was concentrated under reduced pressure to obtain 2.0 g of a solid at 37% yield; LC-MS: $[M+H]^+ = 702.8$.

Step 3: Compound 12c

Add 12b (1.0 g, 1.43 mmol) into a 25 mL single-necked flask, and dissolve in 15 mL of DMF until clear, add 1.0 g of 5% Pd/C, and carry out a hydrogenation reaction for 1.5 h. After the completion of the reaction, it was filtered to obtain a filtrate, which was used directly for the next step of the reaction without purification.

Step 4: Compound 12d

Place the crude product 12c in an ice-water bath, add DIPEA (258 μ L, 1.5 mmol) and then compound M3 (825 mg, 1.43 mmol), the raise to room temperature and react for 1 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 12d (522 mg); LC-MS: $[M-H]^- = 856.4$.

Step 5: Compound 12e


Add 12d (400 mg, 0.47 mmol), M5 (240 mg, 0.47 mmol), PyBOP (250 mg, 0.47 mmol), HOBt (101 mg, 0.47 mmol) and 15 mL of DMF into a 50 mL single-necked flask, add DIPEA (330 μ L, 2.0 mmol) in an ice-water bath, raise to room temperature and react for 3 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to give a preparation liquid of compound 12e, which was lyophilized to obtain 12e (198 mg); LC-MS: $[M+H]^+ = 1275.4$.

Step 6: Compound 12A

Add 12e (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane to a 25 mL single-necked flask and carry out a reaction at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 12A (55 mg); LC-MS: $[M+H]^+ = 1119.4$.

Example 21


Synthesis of compound 12B:



Referring to the synthetic route of Example 20, compound 12B (50 mg) was obtained; LC-MS: $[M+H]^+ = 1119.4$.

Example 22

Synthesis of compound 13A:



Step 1: Compound 13a

In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol),

100 mL THF and p-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, add benzyl 2-hydroxy-2-cyclopentylacetate dropwise (synthesized by the method published in the literature, *Journal of Medicinal Chemistry*, 2013, 56(13), 5541-5552) (7.2 g, 32.6 mmol), then naturally warm up to room temperature to react (react for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO₃ solution was added, followed by extraction with ethyl acetate, washing with saturated sodium chloride solution, drying with anhydrous sodium sulfate, filtering and concentration, and the residue was purified by silica gel column (PE:EA=10:1-5:1-2:1) to give 13a (4.6 g) at 53% yield; LC-MS: [M+H]⁺ = 529.5.

Step 2: Compound 13b

In a 25 mL single-necked flask, add 13a (4 g, 7.6 mmol) and 10 mL DMF, stir at 0°C, add DBU (1.17 g, 7.8 mmol), and react for 1 h. After the completion of TLC-monitored Fmoc deprotection, set aside the reaction solution, ready for use.

Add M4 (3.14 g, 7.6 mmol), PyBOP (4.42 g, 8.5 mmol), HOBt (1.15 g, 8.5 mmol) and 10 mL of DMF into another 25 mL single-necked flask, add DIPEA (1.39 mL, 0.85 mmol) in an ice-water bath, continue stirring for 30 min, and then add the above reaction solution to the reaction flask, and raise to room temperature to react. After the completion of the reaction as detected by HPLC, the reaction solution was purified by preparative LC to obtain a product preparation liquid, which was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate and filtered, and the filtrate was concentrated under reduced pressure to obtain 2.1 g of a solid at 39% yield; LC-MS: [M+H]⁺ = 702.8.

Step 3: Compound 13c

13b (1.5 g, 1.87 mmol) was added to a 25 mL single-necked flask, and after dissolution with 25 mL of DMF, 1.5 g of 5% Pd/C was added, and a hydrogenation reaction was carried out

for 3 h. After completion of the reaction, filtration was performed, and a filtrate was obtained, and used directly for the next step of the reaction without purification.

Step 4: Compound 13d

5 The crude product 13c was placed in an ice-water bath, DIPEA (333 μ L, 1.93 mmol) was added, and compound M3 (1.1 g, 1.87 mmol) was then added, before raising to room temperature to react for 1 h. Upon completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a
10 preparation liquid, which was lyophilized to give 13d (519 mg); LC-MS: $[M-H]^- = 856.6$.

Step 5: Compound 13e


Add 13d (400 mg, 0.47 mmol), M5 (240 mg, 0.48 mmol), PyBOP (250 mg, 0.48 mmol), HOBT (103 mg, 48 mmol) and 15 mL of DMF to
15 a 50 mL single-necked flask, add DIPEA (330 μ L, 2.0 mmol) in an ice-water bath, and raise to room temperature to react for 4 h. Upon completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquid of compound 13e, which was lyophilized to obtain 13e
20 (187 mg); LC-MS: $[M+H]^+ = 1275.5$.

Step 6: Compound 13A

Add 13e (100 mg, 0.08 mmol), zinc bromide (355 mg, 0.16 mmol) and 5 mL of nitromethane into a 25 mL single-necked flask and carry out a reaction at 40°C for 1 h. Upon completion of the
25 reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid
30 compound 13A (60 mg); LC-MS: $[M+H]^+ = 1119.6$.

Example 23


Synthesis of compound 13B:



Referring to the synthetic route of Example 22, compound 13B (51 mg) was obtained; LC-MS: $[M+H]^+ = 1119.6$.

Example 24

5 Synthesis of compound 14A:



Step 1: Compound 14a

In a 250 mL single-necked flask, add M1 (6 g, 16.3 mmol), 100 mL THF and p-toluenesulfonic acid monohydrate (0.31 g, 1.63 mmol), stir and cool to 0°C, add benzyl 3-hydroxy-2-cyclopentylpropionate dropwise (synthesized by the method published in patent W02009011285A1) (7.6 g, 32.6 mmol), then naturally warm up to room temperature and react (react for about 2-4 h), with TLC monitoring. At the end of the reaction, saturated NaHCO₃ solution was added, followed by extraction with ethyl acetate, washing with saturated sodium chloride solution, drying with anhydrous sodium sulfate, filtering and concentration, and the residue was purified by silica gel column (PE:EA=10:1-5:1-2:1) to give 14a (4.4 g) at 49% yield; LC-MS: $[M+H]^+ = 543.6$.

Step 2: Compound 14b

In a 25 mL single-necked flask, add 14a (4 g, 7.4 mmol) and 10 mL DMF, stir at 0°C, add DBU (1.2 g, 8.0 mmol), and react for 1 h. After the completion of Fmoc deprotection as detected by TLC, set aside the reaction solution, ready for use.

Add M4 (3.1 g, 7.4 mmol), PyBOP (4.6 g, 8.8 mmol), HOBT (1.19 g, 8.8 mmol) and 10 mL of DMF into another 25 mL single-necked flask, add DIPEA (1.49 mL, 9.0 mmol) in an ice-water bath, and continue stirring for 30 min, then add the above reaction solution to the reaction flask and raise the temperature to room temperature to react. After the completion of the reaction as detected by HPLC, the reaction solution was purified by preparative LC to obtain a product preparation liquid, which was extracted by dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate and filtered, and the filtrate was concentrated under reduced pressure to obtain 2.6 g of a solid at 49% yield; LC-MS: $[M+H]^+ = 716.4$.

Step 3: Compound 14c

In a 25 mL single-necked flask, 14b (1.0 g, 1.4 mmol) was added, and after dissolution with 15 mL of DMF, 1.0 g of 5% Pd/C was added, and a hydrogenation reaction was carried out for 1.5 h. After completion of the reaction, filtration was performed, and a filtrate was obtained, and used directly for the next step of the reaction without purification.

Step 4: Compound 14d

The crude product 14c was placed in an ice-water bath, DIPEA (248 μ L, 1.5 mmol) was added, then compound M3 (808 mg, 1.4 mmol) was added, then the temperature was raised to room temperature to react for 1 h. When the reaction was completed as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 14d (500 mg); LC-MS: $[M-H]^- = 870.5$.

Step 5: Compound 14e

Add 14d (400 mg, 0.46 mmol), M5 (235 mg, 0.46 mmol), PyBOP


(245 mg, 0.46 mmol), HOBt (99 mg, 0.46 mmol) and 15 mL of DMF into a 50 mL single-necked flask, add DIPEA (331 μ L, 2.0 mmol) in an ice-water bath, raise to room temperature and react for 3 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to give the preparation liquid of compound 14e, which was lyophilized to obtain 14e (146 mg); LC-MS: $[M+H]^+ = 1289.5$.

Step 6: Compound 14A

14e (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 14A (52 mg); LC-MS: $[M+H]^+ = 1133.4$.

Example 25


Synthesis of compound 14B:



Referring to the synthetic route of Example 24, compound 14B (48 mg) was obtained; LC-MS: $[M+H]^+ = 1133.4$.

Example 26

Synthesis of compounds 15A and 15B:



Step 1: Compound 15a

In a 250 mL single-necked flask, M1 (10 g, 27.1 mmol), benzyl 2-hydroxybutyrate (prepared by the method published in the literature *Chemical Communications*, 2019, 55(53), 7699-7702) (10.5 g, 54.3 mmol), zinc acetate (9.96 g, 54.3 mmol) and 100 mL of toluene were added, heated to 100°C and reacted for 4 h. After the reaction was completed, the temperature was reduced to room temperature, filtration was performed to remove insoluble material, and the filtrate was concentrated to obtain a crude product. The crude product was purified by silica gel column chromatography (PE:EA=10:1-5:1-2:1) to obtain 5.67 g of the target material at 42% yield; LC-MS: [M+H]⁺ = 503.5.

Step 2: Compound 15b

Add 15a (5 g, 9.95 mmol) and 15 mL of DMF in a 50 mL single-necked flask, and after dissolution, add DBU (1.68 g, 11 mmol) in an ice-water bath and react for 1 h. The reaction solution was recorded as reaction solution ①.

In another 50 mL single-necked flask, M4 (4.1 g, 10.0 mmol), PyBOP (5.75 g, 11 mmol), HOBt (1.49 g, 11 mmol) and 10 mL DMF were added, and after dissolution, DIPEA (1.82 mL, 11 mmol) was added in an ice-water bath, and a reaction was continued for 40 min, then reaction solution ① was added, and the temperature was raised to room temperature to react for 2 h. The reaction

progress was monitored by HPLC. After completion of the reaction, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid. The preparation liquid was extracted with dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, filtered, and concentrated to obtain 4.6 g of solid, at a yield of 68%; LC-MS: $[M+H]^+ = 676.7$.


Step 3: Compound 15d

Add 15b (2.0 g, 2.96 mmol) in a 25 mL single-necked flask, dissolve in 15 mL of DMF, then add 2.0 g of 5% Pd/C, and perform a hydrogenation reaction for 2 h. After the completion of the reaction, filtration was carried out, and the filtrate was placed in a bath of ice-water, and DIPEA (496 μ L, 3.0 mmol) was added, followed by M3 (1.7 g, 2.96 mmol), and then the temperature was raised to room temperature to react for 1 h. When the reaction was complete as detected by HPLC, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid, which was lyophilized to give 1120.0 mg of a product at 45% yield; LC-MS: $[M-H]^- = 830.3$.

Step 4: Compound 15e


Add 15d (500 mg, 0.60 mmol), M5 (321 mg, 0.60 mmol), PyBOP (469 mg, 0.90 mmol), HOBT (121 mg, 0.90 mmol) and 15 mL of DMF into a 50 mL single-necked flask, add DIPEA (446 μ L, 2.7 mmol) in an ice-water bath, raise to room temperature and react for 2 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compound 15e-1 and compound 15e-2, and the preparation liquids were lyophilized to obtain 138 mg of compound 15e-1, LC-MS: $[M+H]^+ = 1249.5$, and 140 mg of compound 15e-2, LC-MS: $[M+H]^+ = 1249.5$, respectively.

Step 5: Compound 15A



Add 15e-1 (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane into a 25 mL single-necked flask and react at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 59 mg of a solid; LC-MS: $[M+H]^+ = 1093.4$.


Step 6: Compound 15B




Add 15e-2 (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane into a 25 mL single-necked flask and react at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 60 mg of solid; LC-MS: $[M+H]^+ = 1093.4$.

Example 27

Synthesis of compounds 16A and 16B:




Referring to the synthetic route of Example 26, compound 16A (55 mg) was obtained; LC-MS: $[M+H]^+ = 1093.4$.



Referring to the synthetic route of Example 26, compound 16B (54 mg) was obtained; LC-MS: $[M+H]^+ = 1093.4$.

Example 28

5 Synthesis of compounds 17A and 17B:



Step 1: Compound 17a

Add M1 (10 g, 27.1 mmol), benzyl 2-hydroxy-phenylpropionate (synthesized by the method published in the literature *Nature Communications*, 2020. 11(1), 56.) (14.7 g, 54.3 mmol), zinc acetate (9.96 g, 54.3 mmol) and 100 mL of toluene into a 250 mL single-necked flask, heat to 100°C and react for 4 h. After the reaction was completed, the temperature was lowered to room temperature, filtration was performed to remove insoluble material, and the filtrate was concentrated to obtain a crude product. The crude product was purified by silica gel column chromatography (PE:EA=10:1-5:1-2:1) to obtain 6.13 g of the target material at 40% yield; LC-MS: $[M+H]^+ = 565.6$.

Step 2: Compound 17b

In a 50 mL single-necked flask, add 17a (5 g, 8.86 mmol) and 15 mL of DMF, and after dissolution, add DBU (1.53 g, 10 mmol) in an ice-water bath and react for 1 h, and record the reaction solution as reaction solution ①.

5 In another 50 mL single-necked flask, add M4 (3.6 g, 8.86 mmol), PyBOP (5.23 g, 10 mmol), HOBt (1.36 g, 10 mmol) and 10 mL of DMF, and after dissolution, add DIPEA (1.65 mL, 10 mmol) in an ice-water bath, and continue reacting for 30 min, then add reaction solution ①, raise to room temperature and react
10 for 2 h. The reaction progress was monitored by HPLC. After completion of the reaction, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid. The preparation liquid was extracted with dichloromethane, washed with saturated sodium chloride solution,
15 dried with anhydrous sodium sulfate, filtered, and concentrated to obtain 5.0 g of solid at 77% yield; LC-MS: $[M+H]^+ = 738.3$.

Step 3: Compound 17d


Add 17b (3.0 g, 4.07 mmol) in a 25 mL single-necked flask, and after dissolution with 15 mL of DMF, add 3.0 g of 5% Pd/C,
20 and perform a hydrogenation reaction for 2 h. When the reaction was completed, filtration was performed, the filtrate was placed in an ice-water bath, and DIPEA (744 μ L, 4.5 mmol) was added, followed by M3 (2.34 g, 4.07 mmol), and then the temperature was raised to room temperature to react for 1 h. When the
25 reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to give 1.2 g of a product at 33% yield; LC-MS: $[M-H]^- = 892.4$.

Step 4: Compound 17e

30 17d (500 mg, 0.56 mmol), M5 (300 mg, 0.56 mmol), PyBOP (438 mg, 0.84 mmol), HOBt (113 mg, 0.84 mmol) and 15 mL of DMF were added to a 50 mL single-necked flask, DIPEA (330 μ L, 2.0 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 2 h. After the reaction was
35 complete as detected by HPLC, the reaction solution was purified


by HPLC to obtain the preparation liquids of compound 17e-1 and compound 17e-2, and the preparation liquids were lyophilized to obtain 156 mg of compound 17e-1, LC-MS: $[M+H]^+ = 1311.4$, and 150 mg of compound 17e-2, LC-MS: $[M+H]^+ = 1311.7$, respectively.

5 Step 5: Compound 17A



17e-1 (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 43 mg of solid; LC-MS: $[M+H]^+ = 1155.4$.


Step 6: Compound 17B




17e-2 (100 mg, 0.08 mmol), zinc bromide (360 mg, 1.6 mmol) and 5 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 40 mg of solid; LC-MS: $[M+H]^+ = 1155.4$.

Example 29

Synthesis of compounds 18A and 18B:




Referring to the synthetic route of Example 28, compound 18A (54 mg) was obtained; LC-MS: $[M+H]^+ = 1155.4$.



5 Referring to the synthetic route of Example 28, compound 18B (55 mg) was obtained; LC-MS: $[M+H]^+ = 1155.4$.

Example 30

Synthesis of compounds 19A and 19B:



10 Step 1: Compound 19a

Add M1 (10 g, 27.1 mmol), benzyl 2-cyclopropyl-2-hydroxyacetate (prepared according to the method published in patent W02020244657A1) (11.2 g, 54.3 mmol), zinc acetate (9.96 g, 54.3 mmol) and 100 mL of toluene in a 250 mL single-necked flask, heat to 100°C and react for 4 h. When the reaction was completed, the temperature was reduced to room temperature, filtering was performed to remove insoluble material, and the filtrate was concentrated to obtain a crude product. The crude

product was purified by silica gel column chromatography (PE:EA=10:1-5:1-2:1) to obtain 4.97 g of the target material at 36% yield; LC-MS: $[M+H]^+ = 515.2$.

Step 2: Compound 19b

5 Add 19a (4 g, 7.8 mmol) and 10 mL of DMF in a 50 mL single-necked flask, and after dissolution, add DBU (1.42 g, 9.3 mmol) in an ice-water bath and react for 1 h. The reaction solution was recorded as reaction solution ①.

10 In another 50 mL single-necked flask, M4 (3.2 g, 7.8 mmol), PyBOP (4.5 g, 8.6 mmol), HOBt (1.16 g, 8.6 mmol) and 10 mL of DMF were added, and after dissolution, DIPEA (1.65 mL, 10 mmol) was added in an ice-water bath, and the reaction was continued for 30 min, then reaction solution ① was added, and the temperature was raised to room temperature to react for 2 h.

15 The reaction process was monitored by HPLC, and after completion of the reaction, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid. The preparation liquid was extracted with dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, filtered, and concentrated to obtain 4.2 g of solid, with a yield of 78%; LC-MS: $[M+H]^+ = 688.3$.


Step 3: Compound 19d

25 In a 25 mL single-necked flask, 19b (1000 mg, 1.45 mmol) was added, and after dissolution with 15 mL of DMF, 1000 mg of 5% Pd/C was added, and a hydrogenation reaction was performed for 2 h. When the reaction was completed, filtering was performed, the filtrate was placed in an ice-water bath, and DIPEA (248 μ L, 1.5 mmol) was added, followed by M3 (720 mg, 1.45 mmol), and then the temperature was raised to room temperature to react for 1 h. When the reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, and the preparation liquid was lyophilized to obtain 503 mg of product at 41% yield; LC-MS: $[M-H]^- = 842.3$.

35 Step 4: Compounds 19e-1 and 19e-2


19d (500 mg, 0.59 mmol), M5 (317 mg, 0.59 mmol), PyBOP (339 mg, 0.65 mmol), HOBt (88 mg, 0.86 mmol) and 10 mL of DMF were added to a 50 mL single-necked flask, and DIPEA (292 μ L, 1.77 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 2 h. After the reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compound 19e-1 and compound 19e-2, and the preparation liquids were lyophilized to obtain 112 mg of compound 19e-1, LC-MS: $[M+H]^+ = 1261.5$, and 131 mg of compound 19e-2, LC-MS: $[M+H]^+ = 1261.5$, respectively.

Step 5: Compound 19A



19e-1 (100 mg, 0.079 mmol), zinc bromide (357 mg, 1.59 mmol) and 5 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure, to obtain a crude product. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 55 mg of solid; LC-MS: $[M+H]^+ = 1105.4$.

Step 6: Compound 19B




19e-2 (100 mg, 0.079 mmol), zinc bromide (357 mg, 1.59 mmol) and 5 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was completed as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude

product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain 58 mg of a solid; LC-MS: $[M+H]^+ = 1105.4$.

5

Example 31

Synthesis of compounds 20A and 20B:



Step 1: Compound 20a

Add M1 (10 g, 27.1 mmol), benzyl 2-hydroxy-cyclopropylpropionate (synthesized by the method published in patent W02020063676A) (12.0 g, 54.3 mmol), zinc acetate (9.96 g, 54.3 mmol) and 100 mL of toluene in a 250 mL single-necked flask, heat to 100°C and react for 4 h. When the reaction is completed, reduce to room temperature. Filter to remove insoluble material, and the filtrate is concentrated to obtain a crude product. The crude product was purified by silica gel column chromatography (PE:EA=10:1-5:1-2:1) to obtain 5.09 g of the target material; LC-MS: $[M+H]^+ = 529.2$.

Step 2: Compound 20b

20a (4 g, 7.6 mmol) and 10 mL of DMF were added to a 50 mL single-necked flask, and after dissolution, DBU (1.39 g, 9.1 mmol) was added in an ice-water bath and a reaction was carried out for 1 h, and the reaction solution was recorded as reaction solution ①.

In another 50 mL single-necked flask, M4 (3.12 g, 7.6 mmol), PyBOP (4.5 g, 8.6 mmol), HOBt (1.16 g, 8.6 mmol) and 10 mL of

DMF were added, and after dissolution, DIPEA (1.65 mL, 10 mmol) was added in an ice-water bath and a reaction was continued for 30 min, and then the reaction solution ① was added, and the temperature was raised to room temperature to react for 2 h.

HPLC was performed to monitor the reaction progress, and when the reaction was complete, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid. The preparation liquid was extracted with dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, filtered, and concentrated to obtain 4.5 g of solid, with a yield of 84%; LC-MS: $[M+H]^+ = 702.3$.

Step 3: Compound 20d


Add 20b (1000 mg, 1.42 mmol) in a 25 mL single-necked flask, and after dissolution with 15 mL of DMF, add 1000 mg of 5% Pd/C, and perform a hydrogenation reaction for 2 h. After the reaction was completed, filtration was performed, the filtrate was placed in an ice-water bath, and then DIPEA (248 μ L, 1.5 mmol) was added, followed by M5 (708 mg, 1.42 mmol), and the temperature was raised to room temperature to react for 1 h. When the reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 443 mg of product at 36% yield; LC-MS: $[M-H]^- = 856.4$.

Step 4: Compounds 20e-1 and 20e-2

20d (400 mg, 0.47 mmol), exatecan mesylate (250 mg, 0.47 mmol), PyBOP (223 mg, 0.56 mmol), HOBt (83 mg, 0.56 mmol) and 10 mL of DMF were added to a 50 mL single-necked flask, DIPEA (248 μ L, 1.5 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 2 h. After the reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compound 20e-1 and compound 20e-2, which were lyophilized to obtain 103 mg of compound 20e-1, LC-MS: $[M+H]^+ = 1275.5$, and 103 mg of compound 20e-2, LC-MS: $[M+H]^+ = 1275.5$,


respectively.

Step 5: Compound 20A



In a 25 mL single-necked flask, 8A (100 mg, 0.078 mmol),
5 zinc bromide (352 mg, 1.57 mmol) and 5 mL of nitromethane were
added, and a reaction was carried out at 40°C for 1 h. After
the completion of the reaction as detected by HPLC, the solvent
was removed by concentration under reduced pressure and a crude
product was obtained. The crude product was purified by high
10 performance liquid chromatography to obtain a product
preparation liquid, and the preparation liquid was lyophilized
to obtain 51 mg of solid; LC-MS: $[M+H]^+ = 1119.4$.


Step 6: Compound 20B



15 20e-2 (100 mg, 0.079 mmol), zinc bromide (357 mg, 1.59 mmol)
and 5 mL of nitromethane were added to a 25 mL single-necked
flask and a reaction was carried out at 40°C for 1 h. After the
reaction was complete as detected by HPLC, the solvent was
removed by concentration under reduced pressure, and a crude
20 product was obtained. The crude product was purified by high
performance liquid chromatography to obtain a product
preparation liquid, and the preparation liquid was lyophilized
to obtain 47 mg of a solid; LC-MS: $[M+H]^+ = 1119.4$.

Example 32

25 Synthesis of compound 21:



Step 1: Compound SM3-1

Add 77087-60-6 (100 g, 458 mmol), maleic acid (53.4 g, 460 mmol), TEA (64 mL, 460 mmol) and 1000 mL toluene in a 2000 mL single-necked flask and heat to 100°C to react for 5 h. After the reaction was completed, the temperature was lowered to room temperature, and then filtering was performed to remove insoluble material, and the filtrate was concentrated to obtain a crude product. The crude product was purified by silica gel column chromatography (PE:EA=100:1-50:1-20:1) to obtain 75.6 g of the target material; LC-MS: [M+H]⁺ = 299.1.

Step 2: Compound (R)-tert-butyl 2-hydroxy-1,5-glutarate

Add 172793-31-6 (100 g, 338 mmol) and 1000 mL water in a 2000 mL single-necked flask, then sequentially add sodium nitrite (35 g, 507 mmol) and concentrated sulfuric acid (32 mL, 35 mmol), and slowly raise the temperature to room temperature to react for 24 h. After the reaction was completed, extraction was performed with 500 mL of ethyl acetate three times, and the organic phase was dried with anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to remove the solvent, and a crude product was obtained. The crude product was purified by silica gel column chromatography (PE:EA=50:1-30:1-2:1) to obtain 91.2 g of the target material; LC-MS: [M+H]⁺ =

261.4.

Step 3: Compound SM3

Add (*R*)-tert-butyl 2-hydroxy-1,5-glutarate (50 g, 192 mmol) and 1000 mL of anhydrous tetrahydrofuran into a 2000 mL single-necked flask, cool down the temperature to 0°C in an ice-water bath, then add PPh₃ (87.7 g, 288 mmol), DEAD (50.2 g, 288 mmol) and SM3-1 (57.3, 192 mmol) in sequence. The temperature was slowly raised to room temperature to react for 13 h. After completion of the reaction, the insoluble material was removed by filtration and the filtrate was concentrated to obtain a crude product. The crude product was purified by silica gel column chromatography (PE:EA=50:1-30:1-1:1) to obtain 68.6 g of product.

The above product was dissolved in 500 mL of methanol, cooled to 0°C in an ice-water bath, NaOH (64 mL, 190 mmol, 3M/L) was added dropwise at this temperature, and a reaction was performed for 12 h while maintaining this temperature. Next, the pH was adjusted to 3 by the addition of HCl (6 M/L), extraction was performed five times with 500 mL of dichloromethane, drying was performed with anhydrous sodium sulfate, followed by filtering, the filtrate was concentrated under reduced pressure, and the crude product obtained was purified by column chromatography (DCM/MeOH=50/1-20/1-2/1), to obtain 50.4 g of SM3; LC-MS: [M-H]⁻ = 525.5.

Step 4: Compound M6

In a 2000 mL single-necked flask, compound SM3 (50 g, 95 mmol, 1.0 eq), pentafluorophenol (19.2 g, 104.5 mmol, 1.1 eq), DCC (21.5 g, 104.5 mmol, 1.1 eq) and THF (600 mL) were added, and a reaction was carried out at room temperature for 1 h (monitored by TLC), and insoluble material was filtered off. The reaction solution was purified directly by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure by water pump to remove acetonitrile, and lyophilized to obtain compound M6 (51.9 g) at 79% yield; LC-MS: [M+H]⁺ = 693.3.

Step 5: Compound 21a

In a 25 mL single-necked flask, 1c (1 g, 2.36 mmol) was added, and after dissolution with 25 mL DMF, DIPEA (430 μ L, 2.6 mmol) was added, then M6 (1177 mg, 2.36 mmol) was added, and then the temperature was raised to room temperature to react for 1 h. The completion of the reaction was detected by HPLC, and the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid, which was lyophilized to yield 555 mg of product; LC-MS: $[M-H]^- = 931.0$.

Step 6: Compound 21b


In a 100 mL single-necked flask, 21a (500 mg, 0.54 mmol), exatecan mesylate M5 (285 mg, 0.54 mmol), PyBOP (239 mg, 0.6 mmol), HOBt (239 mg, 0.6 mmol) and 10 mL of DMF were added, DIPEA (248 μ L, 1.5 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 2 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquid of compound 21b, which was lyophilized to give 231 mg of the compound; LC-MS: $[M+H]^+ = 1349.5$.

Step 7: Compound 21

Compound 21b (200 mg, 0.1488 mmol), zinc bromide (665 mg, 2.96 mmol) and 10 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain the product preparation liquid, and the preparation liquid was lyophilized to obtain 103 mg of solid; LC-MS: $[M+H]^+ = 1137.5$.

Example 33


Synthesis of compound 22:



Using compounds M6 and 3c as starting materials and referring to the synthetic route of Example 32, compound 22 (91 mg) was obtained; LC-MS: $[M+H]^+ = 1165.5$.

Example 34


5 Synthesis of compounds 23 and 24:



Using compounds M6 and 5c as starting materials and referring to the synthetic route of Example 32, 102 mg of compound 23 was obtained, LC-MS: $[M+H]^+ = 1151.4$; 99 mg of
10 compound 24 was obtained, LC-MS: $[M+H]^+ = 1151.4$.

Example 35


Synthesis of compounds 25 and 26:



Using compounds M6 and 7c as starting materials and referring to the synthetic route of Example 32, 83 mg of
15 compound 25 was obtained, LC-MS: $[M+H]^+ = 1205.7$; 80 mg of compound 26 was obtained, LC-MS: $[M+H]^+ = 1205.7$.

Example 36


Synthesis of compounds 27 and 28:



Using compounds M6 and 19c as starting materials and referring to the synthetic route of Example 32, 100 mg of compound 27 was obtained, LC-MS: $[M+H]^+ = 1177.5$; 101 mg of
25 compound 28 was obtained, LC-MS: $[M+H]^+ = 1177.5$.

Example 37

Synthesis of compound 29:



Step 1: Compound SM4-1

In a 5000 mL single-necked flask, maleic acid (50 g, 431 mmol, 1.0 eq), 114559-25-0 (110 g, 431 mmol, 1 eq), TEA (263 g, 2.16 mol, 5 eq) and toluene (2000 mL) were added, heating was performed to react under reflux for 5 h (monitored by TLC), and insoluble material was filtered off. The reaction solution was directly subjected to rotary distillation under reduced pressure to remove solvent, and the residue was subjected to silica gel column chromatography (PE/EA=50/1-20/1-1/1) to obtain SM4-1 (64.7 g) at 50% yield; LC-MS: $[M+H]^+ = 299.2$.

Step 2: Compound SM4-2

SM4-1 (64 g, 215 mmol) was added into a 2000 mL single-necked flask, and after dissolution with 1000 mL of DMF, DIPEA (71 mL, 430 mmol) was added, and then nonaethylene glycol monomethyl ether methanesulfonate (111.5 g, 220 mmol) was added, and then the temperature was raised to room temperature to react for 2 h. The completion of the reaction was detected by HPLC, and the reaction solution was purified by silica gel column chromatography (PE/EA=50/1-20/1-1/1) to obtain 59.9 g of product; LC-MS: $[M+H]^+ = 709.4$.

Step 3: Compound SM4

SM4-2 (59 g, 83 mmol) was added into a 2000 mL single-necked flask, and after dissolution with 1000 mL MeOH, K₂CO₃ (11.75 g, 85 mmol) was added, and then a reaction was carried out at room temperature for 4 h. When the reaction was complete as detected by HPLC, insoluble materials were removed by filtration, and the reaction solution was purified directly by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure by water pump to remove acetonitrile, and then lyophilized to obtain compound SM4 (27 g); LC-MS: [M-H]⁻ = 693.5.

Step 4: Compound M7

In a 500 mL single-necked flask, compound SM4 (25 g, 36 mmol, 1.0 eq), pentafluorophenol (7.3 g, 40 mmol, 1.1 eq), DCC (8.2 g, 40 mmol, 1.1 eq) and THF (200 mL) were added, a reaction was carried out at room temperature for 1 h (monitored by TLC), and insoluble material was filtered off. The reaction solution was purified directly by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure by water pump to remove acetonitrile, and lyophilized to obtain compound M7 (23.3 g) at 93% yield; LC-MS: [M+H]⁺ = 695.8.

Step 5: Compound 29a

In a 25 mL single-necked flask, 1c (1 g, 2.36 mmol) was added, and after dissolution with 25 mL DMF, DIPEA (430 uL, 2.6 mmol) was added, then M7 (1640 mg, 2.36 mmol) was added, and then the temperature was raised to room temperature to react for 1 h. The completion of the reaction was detected by HPLC, and the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid, which was lyophilized to obtain 609 mg of product; LC-MS: [M-H]⁻ = 1098.5.

Step 6: Compound 29b

29a (500 mg, 0.45 mmol), exatecan mesylate M5 (240 mg, 0.45 mmol), PyBOP (215 mg, 0.54 mmol), HOBT (215 mg, 0.54 mmol) and 10 mL of DMF were added to a 100 mL single-necked flask, DIPEA (248 uL, 1.5 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 2 h.


After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquid of compound 29b, and the preparation liquid was lyophilized to obtain 187 mg of the compound; LC-MS: $[M+H]^+ = 1517.6$.

Step 7: Compound 29

Compound 29b (150 mg, 0.988 mmol), zinc bromide (223 mg, 0.988 mmol) and 10 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain the product preparation liquid, and the preparation liquid was lyophilized to obtain 114 mg of solid; LC-MS: $[M+H]^+ = 1517.9$.

Example 38


Synthesis of compound 30:



Using compounds M7 and 3c as starting materials and referring to the synthetic route of Example 37, compound 30 (125 mg) was obtained; LC-MS: $[M+H]^+ = 1445.6$.

Example 39


Synthesis of compounds 31 and 32:



Using compounds M7 and 5c as starting materials and referring to the synthetic route of Example 37, 61 mg of compound 31 was obtained, LC-MS: $[M+H]^+ = 1431.7$; 63 mg of compound 32 was obtained, LC-MS: $[M+H]^+ = 1431.7$.

Example 40


Synthesis of compounds 33 and 34:



Using compounds M7 and 7c as starting materials, referring to the synthetic route of Example 37, 60 mg of compound 33 was obtained, LC-MS: $[M+H]^+ = 1485.6$; 58 mg of compound 34 was obtained, LC-MS: $[M+H]^+ = 1485.6$.

Example 41


Synthesis of compounds 35 and 36:



Using compounds M7 and 19c as starting materials and referring to the synthetic route of Example 37, 102 mg of compound 35 was obtained, LC-MS: $[M+H]^+ = 1457.8$; 102 mg of compound 36 was obtained, LC-MS: $[M+H]^+ = 1457.8$.

Example 42

Synthesis of compound 37:



Step 1: Compound SM5-1

In a 2000 mL single-necked flask, compound 16947-84-5 (100 g, 295 mmol, 1.0 eq), DIPEA (50 mL, 300 mmol), benzyl bromide (51.3 g, 300 mmol) and THF (1000 mL) were added, and reacted at room temperature for 12 h (monitored by TLC), and the insoluble material was filtered off. The solvent was directly removed from the reaction solution by rotary distillation under reduced pressure, and the residue was subjected to silica gel column chromatography (PE/EA=50/1-20/1-2/1) to obtain SM5-1 (110.1 g) at 87% yield; LC-MS: $[M+H]^+ = 429.2$.

Step 2: Compound SM5-2

In a 2000 mL single-necked flask, compound SM5-1 (100 g, 233.4 mmol, 1.0eq) and THF (1000 mL) were added, cooled to 0°C in an ice-water bath, NaH (37.4 g, 933.5 mmol) and MeI (132.5 g, 933.5 mmol) were added in batches, and a reaction was maintained at 0°C for 24 h (monitored by TLC). The reaction was quenched by adding 500 mL of saturated NH_4Cl aqueous solution, extraction was performed three times with 500 mL of ethyl acetate, followed by drying of the organic phase with anhydrous sodium sulfate, and filtering. The solvent was directly removed from the filtrate by rotary distillation under reduced pressure, and the residue was subjected to silica gel column chromatography (PE/EA=100/1-50/1-10/1) to obtain SM5-2 (37.1 g); LC-MS: $[M+H]^+ = 443.3$.

Step 3: Compound SM5 (cf. *Org. Lett.*, 2006, 8, 3387-3390.)

In a 1000 mL single-necked flask, compound SM5-2 (35 g, 79 mmol, 1.0eq) and DCE (500 mL) were added, palladium diacetate (180 mg, 0.8 mmol), I_2 (20 g, 79 mmol) and iodobenzene diacetate (40.8 g, 126.4 mmol) were added sequentially, and the temperature was raised to 60°C to react for 40 h (monitored by TLC). The reaction was quenched by adding 500 mL of saturated aqueous sodium thiosulfate solution, extraction was performed three times with 500 mL of dichloromethane, followed by drying of the organic phase with anhydrous sodium sulfate, and filtering. The solvent was directly removed from the filtrate by rotary distillation under reduced pressure, and the residue

was subjected to silica gel column chromatography (PE/EA=100/1-50/1-10/1) to obtain SM5 (28 g); LC-MS: $[M+H]^+ = 501.3$.

Step 4: Compound SM6

In a 500 mL single-necked flask, compound SM5 (25 g, 50 mmol, 1.0 eq), di-tert-butyl potassium phosphate (13.66 g, 55 mmol, 1.1 eq), p-toluenesulfonic acid monohydrate (951 mg, 5 mmol, 0.1 eq) and THF (200 mL) were added, and the reaction was carried out at room temperature for 1 h (monitored by TLC), and insoluble material was filtered off. The reaction solution was purified directly by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure by water pump to remove acetonitrile, and lyophilized to obtain compound SM6 (15.1 g) at 46% yield; LC-MS: $[M+H]^+ = 651.4$.

Step 5: Compound SM7

SM6 (15 g, 23 mmol) and 100 mL of DMF were added into a 250 mL single-necked flask, and after dissolving, 15 g of 5% Pd/C was added in an ice-water bath, the atmosphere in the system was replaced by hydrogen three times, and a reaction was carried out at room temperature for 12 h. The Pd/C was removed by filtration, and then the solvent was removed by rotary distillation under reduced pressure with an oil pump, leaving a crude product ready for use.

In another 250 mL single-necked flask, add the above crude product and 100 mL of toluene, triethylamine (6.4 mL, 46 mmol) and maleic anhydride (2.4 g, 24 mmol), and after dissolution, raise to 100°C to react for 2 h. Use HPLC to monitor the progress of the reaction, and when the reaction is completed, purify the reaction solution by HPLC, to obtain a preparation liquid. The preparation liquid was extracted with dichloromethane, washed with saturated sodium chloride solution, dried with anhydrous sodium sulfate, filtered, and concentrated to obtain 4.2 g of a solid at 36% yield; LC-MS: $[M+H]^+ = 507.3$.

Step 6: Compound M8

In a 100 mL single-necked flask, compound SM7 (4 g, 7.9 mmol, 1.0 eq), pentafluorophenol (1.6 g, 8.7 mmol, 1.1 eq), DCC (1.8

g, 8.7 mmol, 1.1 eq) and THF (60 mL) were added, and the reaction was carried out at room temperature for 1 h (monitored by TLC), and insoluble material was filtered off. The reaction solution was purified directly by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure by water pump to remove acetonitrile, and lyophilized to obtain compound M8 (3.7 g) at 70% yield; LC-MS: $[M+H]^+ = 673.2$.

Step 7: Compound 37a

In a 25 mL single-necked flask, 1c (1 g, 2.36 mmol) was added, and after dissolution with 25 mL of DMF, DIPEA (430 μ L, 2.6 mmol) was added, then M8 (1.2 g, 2.36 mmol) was added, and then the temperature was raised to room temperature to react for 1 h. The completion of the reaction was detected by HPLC, and the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid, which was lyophilized to obtain 488 mg of product; LC-MS: $[M-H]^- = 911.0$.

Step 8: Compound 37b

37a (400 mg, 0.44 mmol), exatecan mesylate M5 (235 mg, 0.44 mmol), PyBOP (199 mg, 0.5 mmol), HOBT (69 mg, 0.5 mmol) and 10 mL of DMF were added to a 100 mL single-necked flask, DIPEA (218 μ L, 1.32 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 2 h. After the reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquid of compound 37b, which was lyophilized to give 201 mg of the compound; LC-MS: $[M+H]^+ = 1329.6$.


Step 9: Compound 37

Compound 37b (130 mg, 0.098 mmol), zinc bromide (221 mg, 0.98 mmol) and 10 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain the product

preparation liquid, and the preparation liquid was lyophilized to obtain 96 mg of solid; LC-MS: $[M+H]^+ = 1117.4$.

Example 43

Synthesis of compound 38:




5

Using compounds M8 and 3c as starting materials and referring to the synthetic route of Example 42, compound 38 (51 mg) was obtained; LC-MS: $[M+H]^+ = 1145.6$.

Example 44

10 Synthesis of compounds 39 and 40:




Using compounds M8 and 5c as starting materials and referring to the synthetic route of Example 42, 57 mg of compound 39 was obtained, LC-MS: $[M+H]^+ = 1131.4$; 60 mg of compound 40 was obtained, LC-MS: $[M+H]^+ = 1131.4$.

15

Example 45

Synthesis of compounds 41 and 42:




Using compounds M7 and 7c as starting materials, referring to the synthetic route of Example 42, 44 mg of compound 41 was obtained, LC-MS: $[M+H]^+ = 1185.3$; 44 mg of compound 42 was obtained, LC-MS: $[M+H]^+ = 1185.3$.

20

Example 46


Synthesis of compounds 43 and 44:



Using compounds M8 and 19c as starting materials and referring to the synthetic route of Example 42, 62 mg of compound 43 was obtained, LC-MS: $[M+H]^+ = 1157.4$; 59 mg of compound 44 was obtained, LC-MS: $[M+H]^+ = 1157.4$.

Example 47 (comparative example)


Synthesis of compound 45:



Compound 45 was synthesized by the method provided in Example 58 of patent "CN104755494A".

Example 48

Synthesis of compound 46:



Step 1: Compound 46a

1d (500 mg, 0.62 mmol), M9 (310 mg, 0.62 mmol), PyBOP (448 mg, 0.86 mmol), HOBT (116 mg, 0.86 mmol) and 15 mL of DMF were added to a 50 mL single-necked flask, DIPEA (378 μ L, 2.29 mmol) was added in an ice-water bath, and the temperature was raised to room temperature to react for 2 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by high performance liquid chromatography to obtain a preparation liquid, which was lyophilized to give 46a (210 mg);


LC-MS: $[M+H]^+ = 1221.6$.

Step 2: Compound 46

46a (200 mg, 0.162 mmol), zinc bromide (736 mg, 3.26 mmol) and 10 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain the product preparation liquid, and the preparation liquid was lyophilized to obtain solid compound 46 (120 mg); LC-MS: $[M+H]^+ = 1065.3$.

Example 49


Synthesis of compound 47:



Referring to the synthetic route of Example 48, compound 47 (81 mg) was obtained; LC-MS: $[M+H]^+ = 1065.3$.

Example 50

Synthesis of compound 48A:



Step 1: Compound 48a

Add 5d (1.66g, 2.02mmol, 1.0eq), M9 (1.08g, 2.02mmol, 1.0eq), PyBOP (1.58g, 3.03mmol, 1.5eq), HOBT (0.41g, 3.03mmol, 1.5eq) and DMF (40mL) to a 100 mL single-necked flask. DIPEA (0.84mL, 1.5eq) was added in an ice-water bath, and the temperature was


raised to room temperature to react for 2h (monitored by HPLC). The reaction solution was directly purified by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure by water pump to remove acetonitrile, and freeze-dried to obtain compound 48a (1.54g), with a yield of 61%; LC-MS: $[M+H]^+ = 1235.4$.

Step 6: Compound 48A

Add compound 48a (1.0g, 0.8mmol, 1.0eq) and 35mL nitromethane into a 100mL single-necked flask, and after dissolving, add zinc bromide (3.64g, 16mmol, 20.0eq), react in an oil bath at 40°C (preheated to stabilize) for 30 min, and perform concentration in a water bath at 45°C under reduced pressure with a water pump to remove nitromethane, to obtain a yellow residue solid (monitored by HPLC). A preparation liquid was obtained by acid preparation. The preparation liquid was concentrated at 35°C in a water bath under reduced pressure by a water pump to remove acetonitrile by spinning, and was lyophilized to obtain compound 48A (786 mg) with a yield of 90%.

Example 51

Synthesis of compound 48B:



Step 1: Compound 48b

Compound 5d-1 (200 mg, 0.24 mmol, 1.0 eq), M9 (127 mg, 0.24 mmol, 1.0 eq), PyBOP (187 mg, 0.36 mmol, 1.2 eq), HOBT (48 mg, 0.36 mmol, 1.2 eq), and DMF (6 mL) were added to a 25 mL single-necked flask. The temperature was lowered to 0-5°C in an ice-water bath, DIPEA (62 mg, 0.48 mmol, 2.0 eq) was added, and then the temperature was raised to 20 ± 5°C to react for 2 h.


The end of the reaction was detected by HPLC. The reaction solution was purified directly by HPLC preparation, and the product preparation liquid was collected and lyophilized to obtain compound 48b (150.2 mg); LC-MS: $[M+H]^+ = 1235.4$.

5 Step 2: Compound 48B

Compound 48b (100 mg, 0.081 mmol, 1.0 eq), $ZnBr_2$ (364 mg, 1.62 mmol, 20.0 eq) and CH_3NO_2 (10 mL) were added to a 25 mL single-necked flask in sequence, and then the temperature was raised to 40°C to react for 0.5 h. The reaction was stopped, and the reaction solution was directly spin-dried at 45°C under reduced pressure to obtain a yellow solid, taking samples to monitor the reaction by HPLC. The spin-dried solid was directly purified by HPLC preparation, and the product preparation liquid was collected and lyophilized to obtain compound 48B (70.0 mg); LC-MS: $[M+H]^+ = 1079.4$.

Example 52


Synthesis of compound 49A:



Referring to the route of Example 50, compound 49A (71 mg) was obtained; LC-MS: $[M+H]^+ = 1079.4$.

Example 53


Preparation of compound 49B:



Referring to the synthetic route of Example 51, compound 49B (65 mg) was obtained; LC-MS: $[M+H]^+ = 1079.4$.

Example 54


Synthesis of compounds 50A and 50B:



Step 1: Compounds 50a and 50b


Add 7d (500mg, 0.57mmol), M9 (305mg, 0.57mmol), PyBOP (448mg, 0.86mmol), HOBT (116mg, 0.86mmol) and 15mL DMF into a 50mL single-necked flask, add DIPEA (378uL, 2.29 mmol) in an ice-water bath, raise to room temperature and react for 2h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compound 50a and compound 50b, and the preparation liquids were freeze-dried respectively to obtain 170 mg of compound 50a, LC-MS: $[M+H]^+ = 1289.46$, and 202 mg of compound 50b, LC-MS: $[M+H]^+ = 1289.46$.

Step 2: Compound 50A



Add 50a (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was freeze-dried to obtain 44 mg of solid.


Step 3: Compound 50B



Add 50b (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was freeze-dried to obtain 45 mg of solid.

Example 55


10 Synthesis of compounds 51A and 51B:



Step 1: Compound 51a and compound 51b


Add 8d (500mg, 0.57mmol), M9 (305mg, 0.57mmol), PyBOP (448mg, 0.86mmol), HOBT (116mg, 0.86mmol) and 15mL DMF into a 50mL single-necked flask, add DIPEA (378uL, 2.29 mmol) in an ice-water bath, raise to room temperature and react for 2h. After the reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain preparation liquids of compound 51a and compound 51b, which were respectively lyophilized to obtain 190 mg of compound 51a and 186 mg of compound 51b. LC-MS of compound 51a: $[M+H]^+ = 1289.47$; LC-MS of compound 51b: $[M+H]^+ = 1289.47$.

Step 2: Compound 51A



Add compound 51a (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 39 mg of solid.


Step 3: Compound 51B



Add compound 51b (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was freeze-dried to obtain 60 mg of solid.

Example 56

Synthesis of compound 52A:



Step 1: Compound 52a

Add 11d (800mg, 0.96mmol), M9 (480mg, 0.96mmol), PyBOP (500mg, 0.96mmol), HOBT (208mg, 0.96mmol) and 30mL DMF into a 50mL


single-necked flask, add DIPEA (660uL, 4.0 mmol) in an ice-water bath, raise to room temperature and react for 4h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid of compound 52a, which was lyophilized to obtain 52a (388 mg); LC-MS: $[M+H]^+ = 1261.4$.

Step 2: Compound 52A

Add 52a (150mg, 0.12mmol), zinc bromide (532mg, 2.4mmol) and 10mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 52A (79 mg); LC-MS: $[M+H]^+ = 1105.4$.

Example 57


Synthesis of compound 52B



Referring to the synthetic route of Example 56, compound 52B (50 mg) was obtained. LC-MS: $[M+H]^+ = 1105.4$.

Example 58

Synthesis of compound 53A:



Step 1: Compound 53a

Add 12d (400mg, 0.47mmol), M9 (240mg, 0.47mmol), PyBOP (250mg,


0.47mmol), HOBt (101mg, 0.47mmol) and 15mL DMF into a 50mL single-necked flask, add DIPEA (330uL, 2.0 mmol) in an ice-water bath, raise to room temperature and react for 3h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid of compound 53a, which was lyophilized to obtain 53a (200 mg); LC-MS: $[M+H]^+ = 1275.4$.

Step 2: Compound 53A

Add 53a (100mg, 0.08mmol), zinc bromide (360mg, 1.6mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 53A (51 mg); LC-MS: $[M+H]^+ = 1119.4$.

Example 59


Synthesis of compound 53B:



Referring to the synthetic route of Example 58, compound 53B (50 mg) was obtained; LC-MS: $[M+H]^+ = 1119.4$.

Example 60


Synthesis of compounds 54A and 54B:



Step 1: Compounds 54a and 54b


Add 19d (500mg, 0.59mmol), M9 (317mg, 0.59mmol), PyBOP (339mg, 0.65mmol), HOBt (88mg, 0.86mmol) and 10mL DMF into a 50mL single-necked flask, add DIPEA (292uL, 1.77 mmol) in an ice-water bath, raise to room temperature and react for 2h. After the completion of reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compounds 54a and 54b. The preparation solutions were lyophilized separately to obtain 103 mg of compound 54a, LC-MS: $[M+H]^+ = 1261.5$, and 111 mg of compound 54b, LC-MS: $[M+H]^+ = 1261.5$.

Step 2: Compound 54A



Add 54a (100mg, 0.079mmol), zinc bromide (357mg, 1.59mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 61 mg of solid; LC-MS: $[M+H]^+ = 1105.4$.


Step 3: Compound 54B



Add 54b (100mg, 0.079mmol), zinc bromide (357mg, 1.59mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 57 mg of solid; LC-MS: $[M+H]^+ = 1105.4$.

Example 61


Synthesis of compounds 55A and 55B:



Step 1: Compounds 55a and 55b


Add 20d (400mg, 0.47mmol), M9 (250mg, 0.47mmol), PyBOP (223mg, 0.56mmol), HOBT (83mg, 0.56mmol) and 10mL DMF into a 50mL single-necked flask, add DIPEA (248uL, 1.5 mmol) in an ice-water bath, raise to room temperature and react for 2h. After the reaction was complete as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compounds 55a and 55b. The preparation liquids were separately freeze-dried to obtain 100 mg of compound 55a, LC-MS: $[M+H]^+ = 1275.5$, and 101 mg of compound 55b, LC-MS: $[M+H]^+ = 1275.5$.

Step 2: Compound 55A



Add 55a (100mg, 0.078mmol), zinc bromide (352mg, 1.57mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 42 mg of solid; LC-MS: $[M+H]^+ = 1119.4$.


Step 3: Compound 55B



Add 55b (100mg, 0.079mmol), zinc bromide (357mg, 1.59mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 45 mg of solid; LC-MS: $[M+H]^+ = 1119.4$.

Example 62


10 Synthesis of compound 56:



Referring to the synthetic route of Example 60, compound 56 (50 mg) was obtained; LC-MS: $[M+H]^+ = 1119.3$.

Example 63


15 Synthesis of compound 57:



Referring to the synthetic route of Example 61, compound 57 (50 mg) was obtained; LC-MS: $[M+H]^+ = 1119.4$.

Example 64

20 Synthesis of compound 58:



Step 1: Synthesis of compound 58a

Add 400 mL of DMF to exatecan mesylate M5 (15 g, 28 mol, prepared by the method disclosed in patent application "EP0737683A1"), cool in an ice-water bath to 0°C, add triethylamine dropwise, and adjust the pH to 7 - 8. Benzyl bromide (9.6 g, 56 mmol) was added dropwise in an ice-water bath, the temperature was raised to room temperature (25 °C) to react for 1 hour, completion of the reaction was detected by TLC, the reaction solution was concentrated under reduced pressure, the crude product obtained was purified by preparative high-performance liquid chromatography (acetonitrile/purified water system), the target peak was collected, the acetonitrile was removed under reduced pressure, then lyophilization was performed, to obtain about 11 g of compound 58a, a yellow solid, with a yield of about 74%, MS m/z: [M+H]⁺ 526.3.

Step 2: Synthesis of compound 58b

At room temperature, compound 58a (11 g, 21 mol) and 120 mL of formic acid for dissolution were sequentially added in a 250 mL single-necked flask, 30 mL of formaldehyde (40% aqueous solution) was added to the resulting bright yellow solution, and the temperature was raised to 50°C to react for 1 h. When the reaction was complete as detected by TLC, the temperature was cooled down to room temperature, and the reaction solution was purified by preparative high-performance liquid chromatography (acetonitrile/purified water system), and the target peak was collected. After removing acetonitrile under reduced pressure, lyophilization was performed to give about 4.5 g of compound 58b, a yellow powdery solid, yield about 40%,


MS m/z: $[M+H]^+$ 540.6.

Step 3: Synthesis of compound 58:

At room temperature, add compound 58b (2.3 g, 4.3 mol) in a 250 mL single-necked flask, add 100 mL of DMF to dissolve, add 2.3 g of 5% Pd/C to the resulting bright yellow solution, use a hydrogen balloon to replace the atmosphere in the system, and maintain the reaction at room temperature for 1.5 hours. Completion of the reaction was then detected by HPLC, Pd/C was removed by filtration, and the resulting reaction solution was concentrated, and then purified by preparative high-performance liquid chromatography (acetonitrile/purified water system). The target peak was collected, and after removing acetonitrile under reduced pressure, lyophilization was performed to obtain about 1.0 g of compound 58, a yellow powdery solid with about 52% yield, MS m/z: $[M+H]^+$ 450.5.

Example 65

Synthesis of compound 59:



Step 1: Compound 59a

Add 1d (500 mg, 0.62 mmol), 58 (279 mg, 0.62 mmol), PyBOP (448 mg, 0.86 mmol), HOBT (116 mg, 0.86 mmol) and 15 mL of DMF into a 50 mL single-necked flask, add DIPEA (378 μ L, 2.29 mmol) in an ice-water bath, raise to room temperature and react for 2h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 59a (166 mg); LC-MS: $[M+H]^+$ = 1235.6.


Step 2: Compound 59

59a (100 mg, 0.081 mmol), zinc bromide (368 mg, 1.63 mmol)

and 10 mL of nitromethane were added to a 25 mL single-necked flask and a reaction was carried out at 40°C for 1 h. After the reaction was complete as detected by HPLC, the solvent was removed by concentration under reduced pressure, and a crude product was obtained. The crude product was purified by high performance liquid chromatography to obtain a product preparation liquid, and the preparation liquid was lyophilized to obtain the solid compound 59 (43 mg); LC-MS: $[M+H]^+ = 1079.3$.

Example 66


10 Synthesis of compound 60:



Referring to the synthetic route of Example 65, compound 60 (40 mg) was obtained; LC-MS: $[M+H]^+ = 1079.3$.

Example 67

15 Synthesis of compound 61:



Step 1: Compound 61a

Add 5d (1.66 g, 2.02 mmol, 1.0 eq), 58 (0.91 g, 2.02 mmol, 1.0 eq), PyBOP (1.58 g, 3.03 mmol, 1.5 eq), HOBT (0.41 g, 3.03 mmol, 1.5 eq), and DMF (40 mL) in a 100 mL single-necked flask, add DIPEA (0.84 mL, 1.5 eq) in an ice-water bath, and raise the temperature to room temperature to react for 2 h (monitored by HPLC). The reaction solution was purified directly by preparative LC, the preparation liquid was concentrated at 35°C


in a water bath under reduced pressure with a water pump to remove acetonitrile, and lyophilized to obtain compound 61a (1.21 g); LC-MS: $[M+H]^+ = 1249.4$.

Step 2: Compound 61

5 Add compound 61a (1.0g, 0.8mmol, 1.0eq) and 35mL nitromethane into a 100mL single-necked flask, and after dissolving, add zinc bromide (3.64g, 16mmol, 20.0eq), react in an oil bath at 40°C (preheated to stabilize) for 30 min, and concentrate at 45°C in a water bath under reduced pressure with
10 a water pump to remove nitromethane, to obtain a yellow residue solid (monitored by HPLC). After acid preparation, a preparation liquid was obtained. The preparation liquid was concentrated at 35°C in a water bath under reduced pressure with a water pump to remove acetonitrile by spinning, and freeze-dried to
15 obtain compound 61 (786 mg). LC-MS: $[M+H]^+ = 1093.6$.

Example 68

Synthesis of compound 62:



Step 1: Compound 62a


20 Add compound 5d-1 (200mg, 0.24mmol, 1.0eq), 58 (110.3mg, 0.24mmol, 1.0eq), PyBOP (187mg, 0.36mmol, 1.2eq), HOBT (48mg, 0.36mmol, 1.2eq) and DMF (6mL) to a 25 mL single-necked flask, lower to 0-5°C in an ice-water bath, and add DIPEA (62mg, 0.48mmol, 2.0eq). Then raise to 20±5°C to react for 2h, and
25 use HPLC to monitor for the end of the reaction. The reaction solution was directly purified by HPLC preparation, and the product preparation liquid was collected and freeze-dried to obtain compound 62a (120.9 mg); LC-MS: $[M+H]^+ = 1249.4$.

Step 2: Compound 62

Add compound 62a (100mg, 0.081mmol, 1.0eq), ZnBr₂ (364mg, 1.62mmol, 20.0eq) and CH₃NO₂ (10mL) sequentially into a 25mL single-necked flask. After the addition is complete, raise the temperature to 40°C and react for 0.5h. The reaction was stopped, and the reaction solution was directly spin-dried under reduced pressure at 45°C to obtain a yellow solid, which was sampled to monitor the reaction by HPLC. The spin-dried solid was directly purified by HPLC preparation, and the product preparation liquid was collected and lyophilized to obtain compound 62 (61 mg); LC-MS: [M+H]⁺ = 1093.4.

Example 69


Preparation of compound 63:



Referring to the route of Example 67, compound 63 (60 mg) was obtained; LC-MS: [M+H]⁺ = 1093.4.

Example 70


Preparation of compound 64:



Referring to the synthetic route of Example 68, compound 64 (65 mg) was obtained; LC-MS: [M+H]⁺ = 1093.4.

Example 71


Preparation of compounds 65A and 65B:



Step 1: Compounds 65a and 65b


Add 7d (500mg, 0.57mmol), 58 (256.8mg, 0.57mmol), PyBOP (448mg, 0.86mmol), HOBT (116mg, 0.86mmol) and 15mL DMF into a 50mL one-necked flask, add DIPEA (378uL, 2.29mmol) in an ice-water bath, raise to room temperature and react for 2h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compounds 65a and 65b. The preparation liquids were lyophilized separately to obtain 155 mg of compound 65a, LC-MS: $[M+H]^+ = 1303.4$, and 158 mg of compound 65b, LC-MS: $[M+H]^+ = 1303.6$.

Step 2: Compound 65A



Add 50a (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 49 mg of solid.


Step 3: Compound 65B



Add 65b (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 47 mg of solid.

Example 72


Synthesis of compounds 66A and 66B:



Step 1: Compound 66a and Compound 66b

Add 8d (500mg, 0.57mmol), 58 (256.8mg, 0.57mmol), PyBOP (448mg, 0.86mmol), HOBT (116mg, 0.86mmol) and 15mL DMF into a 50mL one-necked flask, add DIPEA (378uL, 2.29mmol) in an ice-water bath, raise to room temperature and react for 2h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compound 66a and compound 66b, and the preparation liquids were separately lyophilized to obtain 160 mg of compound 66a and 160 mg of compound 66b. LC-MS of compound 66a: $[M+H]^+ = 1303.7$; LC-MS of compound 66b: $[M+H]^+ = 1303.6$.


Step 2: Compound 66A



Add compound 66a (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask

and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 57 mg of solid, LC-MS: $[M+H]^+ = 1147.5$.


Step 3: Compound 66B



Add compound 66b (100mg, 0.077mmol), zinc bromide (349mg, 1.55mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 57 mg of solid, LC-MS: $[M+H]^+ = 1147.5$.

Example 73

Synthesis of compound 67A:



Step 1: Compound 67a

Add 11d (800mg, 0.96mmol), 58 (432.5mg, 0.96mmol), PyBOP (500mg, 0.96mmol), HOBT (208mg, 0.96mmol) and 30mL DMF into a 50mL single-necked flask, add DIPEA (660 uL, 4.0 mmol) in an ice-water bath, raise to room temperature and react for 4h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation


liquid of compound 67a, which was lyophilized to obtain 67a (402 mg); LC-MS: $[M+H]^+ = 1275.4$.

Step 2: Compound 67A

Add 67a (100mg, 0.78mmol), zinc bromide (356mg, 1.57mmol) and 10mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 67A (47 mg); LC-MS: $[M+H]^+ = 1119.5$.

Example 74


Synthesis of compound 67B:



Referring to the synthetic route of Example 73, compound 67B (50 mg) was obtained. LC-MS: $[M+H]^+ = 1119.4$.

Example 75

Synthesis of compound 68A:



Step 1: Compound 68a

Add 12d (400mg, 0.47mmol), 58 (211.7mg, 0.47mmol), PyBOP (250mg, 0.47mmol), HOBT (101mg, 0.47mmol) and 15mL DMF into a 50mL single-necked flask, add DIPEA (330uL, 2.0mmol) in an ice-water bath, and raise to room temperature and react for 3h. After the completion of the reaction as detected by HPLC, the


reaction solution was purified by HPLC to obtain a preparation liquid of compound 68a, which was lyophilized to obtain 68a (177 mg); LC-MS: $[M+H]^+ = 1289.4$.

Step 2: Compound 68A

5 Add 68a (100mg, 0.08mmol), zinc bromide (360mg, 1.6mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was
10 purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 68A (45 mg); LC-MS: $[M+H]^+ = 1133.4$.

Example 76


Synthesis of compound 68B:



Referring to the synthetic route of Example 75, compound 68B (50 mg) was obtained; LC-MS: $[M+H]^+ = 1133.4$.

Example 77

Synthesis of compounds 69A and 68B:




Step 1: Compounds 69a and 69b

Add 19d (500mg, 0.59mmol), 58 (266mg, 0.59mmol), PyBOP (339mg, 0.65mmol), HOBT (88mg, 0.86mmol) and 10mL DMF into a 50mL single-necked flask, add DIPEA (292uL, 1.77 mmol) in an
25 ice-water bath, raise to room temperature and react for 2h.


After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compounds 69a and 69b. The preparation solutions were freeze-dried separately to obtain 109 mg of compound 69a, LC-MS: $[M+H]^+ = 1275.5$, and 111 mg of compound 69b, LC-MS: $[M+H]^+ = 1275.7$.

Step 2: Compound 69A



Add 69a (100mg, 0.078mmol), zinc bromide (352mg, 1.56mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 53 mg of solid; LC-MS: $[M+H]^+ = 1119.4$.


Step 3: Compound 69B



Add 69b (100mg, 0.078mmol), zinc bromide (352mg, 1.56mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 54 mg of solid; LC-MS: $[M+H]^+ = 1119.4$.

Example 78


Synthesis of compounds 70A and 70B:



Step 1: Compounds 70a and 70b


Add 20d (400mg, 0.47mmol), 58 (211.7mg, 0.47mmol), PyBOP (223mg, 0.56mmol), HOBT (83mg, 0.56mmol) and 10mL DMF into a 50mL single-necked flask, add DIPEA (248uL, 1.5mmol) in an ice-water bath, raise to room temperature and react for 2h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compounds 70a and 70b. The preparation liquids were freeze-dried separately to obtain 106 mg of compound 70a, LC-MS: $[M+H]^+ = 1289.5$, and 101 mg of compound 70b, LC-MS: $[M+H]^+ = 1289.4$.

Step 2: Compound 70A



Add 70a (100mg, 0.078mmol), zinc bromide (352mg, 1.57mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 39 mg of solid; LC-MS: $[M+H]^+ = 1133.4$.


Step 3: Compound 70B



Add 70b (100mg, 0.078mmol), zinc bromide (352mg, 1.56mmol) and 5mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain 35 mg of solid; LC-MS: $[M+H]^+ = 1133.4$.

Example 79


Synthesis of compound 71:



Referring to the synthetic route of Example 78, compound 71 (30 mg) was obtained; LC-MS: $[M+H]^+ = 1133.3$.


Example 80

Synthesis of compound 72:



Referring to the synthetic route of Example 78, compound 72 (33 mg) was obtained; LC-MS: $[M+H]^+ = 1133.4$.

Example 81



Synthesis of compound M11:


In a 100 mL single-necked flask, add compound M3 (11.0 g, 19.5 mmol, 1.0 eq), DIPEA (2.8 g, 21.4 mmol, 1.1 eq), 27-amino-4,7,10,13,16,19,22,25-octa-oxaheptacosanoic acid (9.7 g, 20.5

mmol, 1.05 eq) and DMF (60 mL), and react at room temperature (monitored by TLC) for 20 minutes. The reaction solution was purified directly by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure with a water pump to remove acetonitrile, and lyophilized to obtain compound M10 (13.2 g) in a yield of 78%; LC-MS: $[M+H]^+ = 866.5$.

Add compound M10 (13.0 g, 15 mmol, 1.0 eq), pentafluorophenol (3 g, 16.5 mmol, 1.1 eq), DCC (3.4 g, 16.5 mmol, 1.1 eq) and THF (30 mL) in a 100 mL single-necked flask, react at room temperature for 30 min (monitored by TLC), and filter off the insoluble material. The reaction solution was directly purified by preparative LC, and the preparation liquid was concentrated at 35°C in a water bath under reduced pressure with a water pump to remove acetonitrile, and lyophilized to obtain compound M11 (14.2 g) in 92% yield; LC-MS: $[M+H]^+ = 1032.5$.

Example 82

Synthesis of compound 73:



Step 1: Synthesis of compound 73a

Add 10 mL of DMF to M11 (1 g, 0.79 mol), cool in an ice-water bath to 0°C, add compound 1c (334 mg, 0.79 mol) and DIPEA (154 mg, 1.19 mol), and maintain the conditions to react for 1

h. TLC monitors for the completion of the reaction. The reaction solution was purified by preparative high-performance liquid chromatography (acetonitrile/purified water system), and the target peak was collected. After removing acetonitrile under reduced pressure, freeze-drying was performed to obtain about 1.2 g of compound 73a, MS m/z: $[M+H]^+ = 1271.9$.

Step 2: Synthesis of compound 73b


Add 73a (1.2g, 0.94mmol), M5 (500mg, 0.94mmol), PyBOP (625mg, 1.2mmol), HOBt (162mg, 1.2mmol) and 15mL DMF into a 25mL single-necked flask, add DIPEA (310mg, 2.4mmol) in an ice-water bath, raise to room temperature and react for 2h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 73b (709 mg); LC-MS: $[M+H]^+ = 1720.8$.

Step 3: Synthesis of compound 73:

Add 73b (200mg, 0.116mmol), zinc bromide (523mg, 2.32mmol) and 10mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 73 (88 mg); LC-MS: $[M+H]^+ = 1532.6$.

Example 83


Synthesis of compound 74:



Referring to the synthetic route of Example 82, compound 74 (90 mg) was obtained; LC-MS: $[M+H]^+ = 1532.6$.

Example 84

Synthesis of compound 75:



Step 1: Synthesis of compound 75a

Add 10 mL of DMF to M11 (1 g, 0.79 mol), cool in an ice-water bath to 0°C, add compound 5c (345 mg, 0.79 mol) and DIPEA (154 mg, 1.19 mol), and maintain the conditions to react for 1 h. TLC monitors for completion of the reaction. The reaction solution was purified by preparative high-performance liquid chromatography (acetonitrile/purified water system), and the target peak was collected. After removing acetonitrile under reduced pressure, lyophilization was performed to obtain 0.9g of compound 75a, MS m/z: [M+H]⁺ = 1285.6.

Step 2: Synthesis of compound 75b

Add 75a (700mg, 0.54mmol), M5 (289mg, 0.54mmol), PyBOP (313mg, 0.6mmol), HOBT (81mg, 0.6mmol) and 10mL DMF into a 25mL single-necked flask, add DIPEA (155mg, 1.2 mmol) in an ice-water bath, raise to room temperature and react for 2h. After completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain a preparation liquid, which was lyophilized to obtain 75b (304 mg); LC-MS: [M+H]⁺ = 1734.8.


Step 3: Synthesis of compound 75:

Add 75b (200mg, 0.116mmol), zinc bromide (523mg, 2.32mmol)

and 10mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 75 (96 mg); LC-MS: $[M+H]^+ = 1546.6$.

Example 85


Synthesis of compound 76:



Referring to the synthetic route of Example 84, compound 76 (92 mg) was obtained; LC-MS: $[M+H]^+ = 1546.5$.

Example 86


Synthesis of compound 77:



Referring to the synthetic route of Example 84, compound 77 (87 mg) was obtained; LC-MS: $[M+H]^+ = 1546.5$.

Example 87


Synthesis of compound 78:



Referring to the synthetic route of Example 84, compound 78 (94 mg) was obtained; LC-MS: $[M+H]^+ = 1546.7$.

Example 88

Synthesis of compounds 79 and 80:




Step 1: Synthesis of compound 79a

Add 10 mL of DMF to M11 (1 g, 0.79 mol), cool to 0°C in an ice-water bath, add compound 20c (377 mg, 0.79 mol) and DIPEA (154 mg, 1.19 mol), and maintain the conditions to react for 1 h. TLC monitors for completion of the reaction. The reaction solution was purified by preparative high-performance liquid chromatography (acetonitrile/purified water system), and the target peak was collected. After removing acetonitrile under reduced pressure, lyophilization was performed to obtain 783 mg of compound 79a, MS m/z: $[M+H]^+ = 1325.8$.

Step 2: Synthesis of compounds 79b-1 and 79b-2


Add 79a (600 mg, 0.45 mmol), M5 (240 mg, 0.45 mmol), PyBOP (261 mg, 0.5 mmol), HOBT (68 mg, 0.5 mmol) and 10 mL of DMF to a 25 mL single-necked flask, add DIPEA (130 mg, 1 mmol) in an ice-water bath, and raise to room temperature to react for 2 h. After the completion of the reaction as detected by HPLC, the reaction solution was purified by HPLC to obtain the preparation liquids of compounds 79b-1 and 79b-2, and the preparation liquids were lyophilized to obtain 79b-1 (124 mg); LC-MS: $[M+H]^+ = 1743.0$; and obtain 79b-2 (122 mg); LC-MS: $[M+H]^+ = 1743.0$.

Step 3: Synthesis of compound 79



Add 79b-1 (100mg, 0.057mmol), zinc bromide (258mg, 1.15mmol) and 10mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 79 (30 mg); LC-MS: [M+H]⁺ = 1586.9.


Step 4: Synthesis of compound 80



Add 79b-2 (100mg, 0.057mmol), zinc bromide (258mg, 1.15mmol) and 10mL nitromethane into a 25mL single-necked flask and react at 40°C for 1h. After the completion of the reaction as detected by HPLC, the solvent was removed by concentration under reduced pressure to obtain a crude product. The crude product was purified by HPLC to obtain a product preparation liquid, which was lyophilized to obtain solid compound 80 (33 mg); LC-MS: [M+H]⁺ = 1587.0.

Example 89


Synthesis of compound 81:



Referring to the synthetic route of Example 88, compound 81 (24 mg) was obtained; LC-MS: [M+H]⁺ = 1586.9.

5 Example 90

Synthesis of compound 82:



Referring to the synthetic route of Example 88, compound 82 (29 mg) was obtained; LC-MS: [M+H]⁺ = 1586.9.

10 Example 91

1) Expression and purification of SI-1×6.4 antibody:

Expi293 (Shanghai OPM Biosciences Co., Ltd.) suspension cells were used to express SI-1×6.4 antibody. The day before transfection, cells were inoculated at a density of 0.9×10⁶ cells/mL in a 1 L shake flask containing 300 mL of OPM-293 CD05 Medium (81075-001, Shanghai OPM Biosciences Co., Ltd.), culturing was performed overnight at 37°C, 5% CO₂, and 120 rpm in a cell culture shaker. On the next day, the antibody expression plasmid was transfected with PEI-MAX, wherein the mass ratio of plasmid:PEI-MAX was 1:3. OPM-293 ProFeed supplement was added at 5% (v/v) on the first day after transfection, and then again at 5% (v/v) on the third day after transfection, and centrifugation was performed to collect the supernatant on the sixth day after transfection.

25 The collected cell expression supernatant was eluted with 0.05 M sodium acetate (pH 3.6) in a Protein A affinity chromatography column (UniMab 50, Suzhou Nanomicro Technology Co., Ltd.), and the captured antibody was adjusted to pH 7.0

with 1 M Tris-HCl (pH 8.8) at 0.7/10 (v/v), and then passed through a gel filtration chromatography column SEC (Superdex 200, GE) to remove impurities such as polymers, while the antibody buffer was replaced with 20 mM PB (pH 6.5).

5 Antibody SI-1 × 6.4:

Light chain nucleic acid coding sequence

SEQ ID NO: 1

GACATCTTGCTGACTCAGTCTCCAGTCATCCTGTCTGTGAGTCCAGGAGAAAGAGTC
AGTTTCTCCTGCAGGGCCAGTCAGAGTATTGGCACAAACATACACTGGTATCAGCAA
AGAACAAATGGTTCTCCAAGGCTTCTCATAAAGTATGCTTCTGAGTCTATCTCTGGGA
TTCCTTCCAGGTTTAGTGGCAGTGGATCAGGGACAGATTTTACTCTTAGCATCAACAG
TGTGGAGTCTGAAGATATTGCAGATTATTACTGTCAACAAAATAATAACTGGCCAACC
ACGTTCCGGTGTCTGGGACCAAGCTGGAGCTGAAACGTACGGTGGCTGCACCATCTGTC
TTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAACTGCCTCTGTTGTGTGCC
TGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAACGCCC
TCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACC
TACAGCCTCAGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGT
CTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAA
CAGGGGAGAGTGTTAG

Light chain amino acid sequence

10 SEQ ID NO: 2

DILLTQSPVILSVSPGERVSFSCRASOSIGTNIHWYQORTNGSPRLLIK~~Y~~ASESISGIPSRFSGSGS
GTDFTL~~S~~INSVESEDIADYYCOONNNWPTTFGAGTKLELKRTVAAPSVFIFPPSDEQLKSGTA
SVVCLLN~~N~~FYPREAKVQWKVDNALQSGNSQESVTEQDSK~~D~~STYSLSS~~T~~LTLSKADY~~E~~K
HKVYACEVTHOGLSSPVTKSFNRGEC

wherein the variable region is:

SEQ ID NO: 28

DILLTQSPVILSVSPGERVSFSCRASOSIGTNIHWYQORTNGSPRLLIK~~Y~~ASESISGIPSRFSGSGS
GTDFTL~~S~~INSVESEDIADYYCOONNNWPTTFGAGTKLELK

15 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 25), CDR2 (SEQ ID NO: 26), and CDR3 (SEQ ID NO: 27), respectively.

Nucleic acid coding sequence for the construct of the heavy chain and single-chain Fv (scFv) structural domain

20 SEQ ID NO: 3

CAGGTGCAGCTGAAGCAGTCAGGACCTGGCCTAGTGCAGCCCTCACAGAGCCTGTC
CATCACCTGCACAGTCTCTGGTTTCTCATTAATACTATGGTGTACACTGGGTTCGCC
AGTCTCCAGGAAAGGGTCTGGAGTGGCTGGGAGTGATATGGAGTGGTGGAACACA
GACTATAATACACCTTTTACATCCAGACTGAGCATCAACAAGGACAATTCCAAGAGCC
AAGTTTTCTTTAAAATGAACAGTCTGCAATCTAATGACACAGCCATATATTACTGTGCC
AGAGCCCTCACCTACTATGATTACGAGTTTGCTTACTGGGGCCAAGGGACTCTGGTCA
CTGTCTCTAGCGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAA
GAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCG
AACCGGTGACGGTGTCTGGAACTCAGGCGCCCTGACCAGCGGCGTGCACACCTTC
CCGGCTGTCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCC
TCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAAC
ACCAAGGTGGACAAGAGAGTTGAGCCCAATCTTGTGACAAAACCTCACACATGCCC
ACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCCTCTTCCCCCAA
ACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGA
CGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGG
TGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTG
GTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC
AAGGTCTCCAACAAAGCCCTCCCAGCCCCATCGAGAAAACCATCTCCAAAGCCAA
AGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGA
CCAAGAACCAGGTACGCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCG
CCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCTCCC
GTGCTGGACTCCGACGGCTCCTTCTTCTCTATAGCAAGCTCACCGTGGACAAGAGC
AGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAAC
CACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTGGCGGTGGAGGGTCCGGCGG
TGGTGGATCACAGGTGCAATTGCAGGAGTCGGGGGGAGGCCTGGTCAAGCCTGGAG
GGTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTACCTTTAGTAGTTATTGGATGAG
CTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGGCCAACATAAACC
ATGGAAGTGCAGTACTATGTGGAATCTGTGAAGGGCCGATTACCATCTCCAGAG
ACGACGCCAAGAATCACTGTATCTGCAAATGAACAGCCTGAGAGCTGAGGACACG
GCTGTGTATTACTGTGCGAGAGATCGTGGGGTGGGCTACTTCGATCTCTGGGGCCGTG
GCACCCTGGTCACCGTCTCGAGCGGTGGAGGCGGTTTCAGGCGGAGGTGGTTCCGGC
GGTGGCGGCTCCCAGTCTGCCCTGACTCAGCCTGCCTCCGTGTCTGGGTCTCCTGGA
CAGTCGATCACCATCTCCTGCACTGGAACCAGCAGTGACGTTGGTGGTTATAACTTTG
TCTCCTGGTACCAACAACACCCAGGCAAAGCCCCCAAACCTCATGATCTATGATGTCAG
TGATCGGCCCTCAGGGGTGTCTGATCGCTTCTCCGGCTCCAAGTCTGGCAACACGGC
CTCCCTGATCATCTCTGGCCTCCAGGCTGACGACGAGGCTGATTATTACTGCAGCTCA
TATGGGAGCAGCAGCACTCATGTGATTTTCGGCGGAGGGACCAAGGTGACCGTCCTA
TAA

Amino acid sequence for the construct of heavy chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 4

QVQLKQSGPGLVQPSSLSITCTVSGFSLTNYGVHWVROSPGKGLEWLGITWSGGNTDYNTP
ITSRISINKDNSKSOVFFKMNSLQSNDAIYYCARALYYDYFAYWGOGLTVTVSSASTKGPS
VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV
VTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKHTCPPCPAPELLGGPSVFLFPPK
PKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVS
VLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF
SCSVMHREALHNHYTQKSLSLSPGSGGGGSGGGGSQVQLQESGGGLVKPGGSLRLSCAASGF
TESSYWMASWVROAPGKGLEWVANINRDGSASYVDSEYKGRFTISRDDAKNSLYLOMNSLRAE
DAIYYCARDRGVGYFDLWGRGLTVTVSSGGGGSGGGGSGGGGSQSALTQPAVSGSPGOSI
TISCTGTSSDFGGYNEFVWYQOHPGKAPKLMYDPSDRPSGVSDRFSGSKSGNTASLIISGLOA
DDEADYYCSSYGSSTHYIFGGGTKVTVL

where the variable region of the heavy chain is:

SEQ ID NO: 38

QVQLKQSGPGLVQPSQLSLITCTVSGFSLTNYGLHWVRQSPGKGLEWLGFWSGGNIDYNTPTSRLSINKDNSKSVFFKMNSLQSNDAIYYCARALTYDYEFAYWGQGLTVTVSS

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 29), CDR2 (SEQ ID NO: 30), and CDR3 (SEQ ID NO: 31), respectively.

The variable region of the heavy chain in the structural domain of the single chain Fv (scFv) is:

SEQ ID NO: 39

QVQLQESGGGLVKPGGSLRLSCAASGFTFSSYWMSWVRQAPGKGLEWVANINRDGSASYYYDSVKGRFTISRDDAKNSLYLQMNSLRAEDTAVYYCARDRCVGYFDLWGRGTLTVTVSS

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 32), CDR2 (SEQ ID NO: 33), and CDR3 (SEQ ID NO: 34), respectively.

The variable region of the light chain in the single-chain Fv (scFv) structural domain is:

SEQ ID NO: 40

QSALTQPASVSGSPGQSITISCTCTSSDVGGYNFTSWYQQHPGKAPKLMYDFNSDRPSGVSDRFSGSKSGNTASLISGLQADDEADYYCSSYGSSSTHEIFGGGTKVTVL

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 35), CDR2 (SEQ ID NO: 36), and CDR3 (SEQ ID NO: 37), respectively.

2) Expression and purification of SI-1×22 antibody: a similar method was followed for the expression and purification of SI-1×22 antibody.

Antibody SI-1×22:

Nucleic acid coding sequence for the construct of light chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 9

GACATCTTGCTGACTCAGTCTCCAGTCATCCTGTCTGTGAGTCCAGGAGAAAGAGTC
 AGTTTCTCCTGCAGGGCCAGTCAGAGTATTGGCACAAACATACTGGTATCAGCAA
 AGAACAAATGGTTCTCCAAGGCTTCTCATAAAGTATGCTTCTGAGTCTATCTCTGGG
 ATTCCTTCCAGGTTTGTAGTGGCAGTGGATCAGGGACAGATTTTACTCTTAGCATCAAC
 AGTGTGGAGTCTGAAGATATTGCAGATTATTACTGTCAACAAAATAATAACTGGCCA
 ACCACGTTTCGGTTGTGGGACCAAGCTGGAGCTGAAACGTACGGTGGCTGCACCATC
 TGTCTTCATCTTCCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTG
 TGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAA
 CGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACA
 GCACCTACAGCCTCAGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACAC
 AAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAG
 CTTCAACAGGGGAGAGTGTGGTGGCGGCGGAAGTGGCGGTGGAGGATCCGGCGGTG
 GTGGATCACAGGTGCAGCTGCAGGAGTCGGGGGGAGGCCTGGTCAAGCCTGGAGGG
 TCCCTGAGTCTCTCCTGTGCAGCCTCTGGATTACCTTTAGTAGTTATTGGATGAGCT
 GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGGCCAAACATAAACC GCGAT
 GGAAGTGGCAGTTACTATGTGGACTCTGTGAAGGGCCGATTACCATCTCCAGAGA
 CGACGCCAAGAATACTACTGTATCTGCAAATGAACAGCCTGAGAGCTGAGGACACGG
 CTGTGTATTACTGTGCGAGAGATCGTGGGGTGGGCTACTTCGATCTCTGGGGCCGTG
 GCACCCTGGTCACCGTCTCGAGCGGTGGAGGCGGTTTCAGGCGGAGGTGGTTCCGGC
 GGTGGCGGCTCCAGTCTGCCCTGACTCAGCCTGCCTCCGTGTCTGGGTCTCCTGGA
 CAGTCGATCACCATCTCCTGCACTGGAACCAGCAGTGACGTTGGTGGTTATAACTTT
 GTCTCCTGGTACCAACAACACCCAGGCAAAGCCCCCAAATCATGATCTATGATGTC
 AGTGATCGGCCCTCAGGGGTGTCTGATCGCTTCTCCGGCTCCAAGTCTGGCAACACG
 GCCTCCCTGATCATCTCTGGCCTCCAGGCTGACGACGAGGCTGATTATTACTGCAGC
 TCATATGGGAGCAGCAGCACTCATGTGATTTTCGGCGGAGGGACCAAGGTGACCGT
 CCTATAA

Amino acid sequence of construct of light chain and single-chain Fv (scFv) structural domain

5 SEQ ID NO: 10

DILLTQSPVILSVSPGERVSFSCRASQSIGTNIHWYQOORTNGSPRLLIKYASESISGIPSRFSGSGS
GTDFTLINSVESEDIADYYCOQNNNWPTTFGCGTKLELKRTVAAPSVFIFPPSDEQLKSGTA
 SVVCLLN NFYPREAKVQWKVDNALQSGNSQESVTEQDSK DSTYLSSTLTLSKADYEK
 HKVYACEVTHQGLSSPVTKSFNRGECGGGSGGGGSGGGGSGVQLVQSGGGLVFPKPKGS
LSLSCAASGFTFSYWMSSWVROAPGKGLEWYANINRDGSASYVDSEIKGRFTISRDDAKNSLY
LOMNSLR AEDTAVYYCARDRGVGYFDLWGRGTLVTVSSGGGSGGGGSGGGGSGQSALTGP
ASVSGSPGQSITISCTGTSSDYGGYNFYSWYQOHPGKAPKLMYDYSRPSGVSDRFSGSKSG
NTASLIISGLQADDEADYYCSSYGSSTHTVIFGGGTKVTYL

where the variable region of the light chain is:

SEQ ID NO: 49

DILLTQSPVILSVSPGERVSFSCRASQSIGTNIHWYQOORTNGSPRLLIKYASESISGIPSRFSGSGS
GTDFTLINSVESEDIADYYCOQNNNWPTTFGCGTKLELK

10 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 25), CDR2 (SEQ ID NO: 26), and CDR3 (SEQ ID NO: 27), respectively.

The variable region of the heavy chain in the structural domain of the single-chain Fv (scFv) is:

15 SEQ ID NO: 50

QVQLQESGGGLVKPGGSLSLSCAASGFTFSSYWMMSWVRQAPGKGLEWVANINRDGSASYVD
SVKGRFTISRDDAKNSLYLOMNSLRAEDTAVYYCARDRGVGYFDLWGRGTLVTVSS

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 32), CDR2 (SEQ ID NO: 33), and CDR3 (SEQ ID NO: 34), respectively.

5 The variable region of the light chain in the single-chain Fv (scFv) structural domain is:

SEQ ID NO: 51

QSALTQPASVSGSPGQSITISCTGTSSDFGGYNEYSWYQQHPGKAPKLMYDVS
DRPSGVSDRFSGSKSGNTASLIHSGHQADDEADYYCSNYGSSSTHVF
FGGGTKVTVL

10 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 35), CDR2 (SEQ ID NO: 36), and CDR3 (SEQ ID NO: 37), respectively.

Nucleic acid coding sequence for heavy chains

SEQ ID NO: 11

CAGGTGCAGCTGAAGCAGTCAGGACCTGGCCTAGTGCAGCCCTCACAGAGCCTGTC
CATCACCTGCACAGTCTCTGGTTTCTCATTAACTAACTATGGTGTACACTGGGTTCGC
CAGTCTCCAGGAAAGTGCCTGGAGTGGCTGGGAGTGATATGGAGTGGTGGAAACAC
AGACTATAATACACCTTTACATCCAGACTGAGCATCAACAAGGACAATTCCAAGA
GCCAAGTTTTCTTTAAAATGAACAGTCTGCAATCTAATGACACAGCCATATATTACT
GTGCCAGAGCCCTCACCTACTATGATTACGAGTTTGCTTACTGGGGCCAAGGGACTC
TGGTCACTGTCTCTGCTGCTAGCACCAAGGGGCCATCGGTCTTCCCCCTGGCACCCCT
CCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTAC
TTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGCGCCCTGACCAGCGGCGTGCA
CACCTTCCCGGTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGAC
CGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAGCC
CAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGTGACAAAACCTACA
CATGCCACACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCTCTTCC
CCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTG
GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGG
CGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACG
TACCGTGTGGTCAGCGTCTCACCCTGCTGACCAAGGACTGGCTGAATGGCAAGGA
GTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCATCGAGAAAACCATCT
CCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGG
GATGAGCTGACCAAGAACCAGGTGACCTGACCTGCCTGGTCAAAGGCTTCTATCCC
AGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGA
CCACGCCTCCCGTGTGCTGGACTCCGACGGCTCCTTCTCCTCTATAGCAAGCTACCGT
GGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGG
CTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAA

Amino acid sequence of the heavy chain

SEQ ID NO: 12

OVOLKQSGPGLVQPSQSLSITCTVSGFSLTNYGVHWVROSPGKCLEWLGVWSGGNTDYNTP
FTSRLSINKDNSKSOVFFKMNSLQSNDTAIYYCARALTYDYEFAYWGOGTLVTVSAASTKGPS
 VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSS
 VVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFP
 PKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV
 VSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKN
 QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQG
 NVFSCSVMHEALHNHYTQKSLSLSPG

where the variable region of the heavy chain is:

SEQ ID NO: 52

OVOLKQSGPGLVQPSQSLSITCTVSGFSLTNYGVHWVROSPGKCLEWLGVWSGGNTDYNTP
FTSRLSINKDNSKSOVFFKMNSLQSNDTAIYYCARALTYDYEFAYWGOGTLVTVSA

- 5 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 29), CDR2 (SEQ ID NO: 30), and CDR3 (SEQ ID NO: 31), respectively.

(3) Expression and purification of SI-1×24 antibody: a similar method was followed for the expression and

- 10 purification of SI-1×24 antibody.

Antibody SI-1×24:

Light chain nucleic acid coding sequence

SEQ ID NO: 13

GACATCTTGCTGACTCAGTCTCCAGTCATCCTGTCTGTGAGTCCAGGAGAAAGAGTC
 AGTTTCTCCTGCAGGGCCAGTCAGAGTATTGGCACAAACATACACTGGTATCAGCAA
 AGAACAAATGGTTCTCCAAGGCTTCTCATAAAGTATGCTTCTGAGTCTATCTCTGGG
 ATTCCTTCCAGGTTTAGTGGCAGTGGATCAGGGACAGATTTTACTCTTAGCATCAAC
 AGTGTGGAGTCTGAAGATATTGCAGATTATTACTGTCAACAAAATAATAACTGGCCA
 ACCACGTTCCGGTGCTGGGACCAAGCTGGAGCTGAAACGTACGGTGGCTGCACCATC
 TGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAACCTGCCTCTGTTGTG
 TGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAA
 CGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACA
 GCACCTACAGCCTCAGCAGCACCCTGACGCTGAGCAAAGCAGACTACGAGAAACAC
 AAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAG
 CTTCAACAGGGGAGAGTGTTAG

- 15 Light chain amino acid sequence

SEQ ID NO: 14

DILLTQSPVILSVSPGERVSFSCRASOSIGTNIHWYOORTNGSPRLLIKYASESISGIPSRFSGSGS
GTDFTLSINSVEEDIADYYCOQNNNWPTTFGAGTKLELKRTVAAPSVFIPPSDEQLKSGTA
 SVVCLLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSSLTLSKADYEK
 HKVYACEVTHQGLSSPVTKSFNRGEC

wherein the variable region is:

SEQ ID NO: 28

DILLTQSPVILSVSPGERVSFSCRASOSIGTNIHWYOORTNGSPRLLIKYASESISGIPSRFSGSGS
GTDFTLSINSVEEDIADYYCOQNNNWPTTFGAGTKLEL

- 20 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 25), CDR2 (SEQ ID NO: 26), and CDR3 (SEQ ID NO: 27),

respectively.

Nucleic acid coding sequence for the construct of heavy chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 15

CAGGTGCAGCTGCAGGAGTCGGGGGGAGGCCTGGTCAAGCCTGGAGGGTCCCTGAG
ACTCTCCTGTGCAGCCTCTGGATTACCTTTAGTAGTTATTGGATGAGCTGGGTCCGC
CAGGCTCCAGGGAAGGGGCTGGAGTGGGTGGCCAACATAAACC GCGATGGAAGTGC
GAGTTACTATGTGGACTCTGTGAAGGGCCGATTACCATCTCCAGAGACGACGCCAA
GAACTCACTGTATCTGCAAATGAACAGCCTGAGAGCTGAGGACACGGCTGTGTATTA
CTGTGCGAGAGATCGTGGGGTGGGCTACTTCGATCTCTGGGGCCGTGGCACCCCTGGT
CACCGTCTCGAGCGGTGGAGGCGGTT CAGGCGGAGGTGGTTCCGGCGGTGGCGGCT
CCCAGTCTGCCCTGACTCAGCCTGCCTCCGTGTCTGGGTCTCCTGGACAGTCGATCAC
CATCTCCTGCACTGGAACCAGCAGTGACGTTGGTGGTTATAACTTTGTCTCCTGGTAC
CAACAACACCCAGGCAAAGCCCCCAAAC TCA TGATCTATGATGTCAGTGATCGGCC
TCAGGGGTGTCTGATCGCTTCTCCGGCTCCAAGTCTGGCAACACGGCCTCCCTGATC
ATCTCTGGCCTCCAGGCTGACGACGAGGCTGATTATTACTGCAGCTCATATGGGAGCA
GCAGCACTCATGTGATTTTCGGCGGAGGGACCAAGGTGACCGTCCTAGCGCGGTGGAG
GATCCGGCGGTGGTGGATCACAGGTGCAGCTGAAGCAGTCAGGACCTGGCCTAGTG
CAGCCCTCACAGAGCCTGTCCATCACCTGCACAGTCTCTGGTTTCTCATTAATACTA
ATGGTGTACACTGGGTTCGCCAGTCTCCAGGAAAGGGTCTGGAGTGGCTGGGAGTGA
TATGGAGTGGTGGAAACACAGACTATAATACACCTTTCACATCCAGACTGAGCATCAA
CAAGGACAATTCCAAGAGCCAAGTTTTCTTTAAATGAACAGTCTGCAATCTAATGAC
ACAGCCATATATTACTGTGCCAGAGCCCTCACCTACTATGATTACGAGTTTGCTTACTG
GGGCCAAGGGACTCTGGTCACTGTCTCTAGCGCTAGCACCAAGGGCCCATCGGTCTT
CCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCT
GGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTGGAAC TACAGGCGCCCTGA
CCAGCGGCGTGCACACCTTCCCGGCTGTCTTACAGTCCTCAGGACTCTACTCCCTCA
GCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACG
TGAATCACAAGCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGT
GACAAAAC TCAACATGCCACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTC
AGTCTTCTCTTCCCCC AAAACCAAGGACACCCTCATGATCTCCCGGACCCCTGA
GGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACT
GGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAG
TACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTG
AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGA
GAAAACCATCTCCAAAGCCAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGC
CCCCATCCCGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAG
GCTTCTATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC
AACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTATAGCA
AGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTG
ATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGT
TAA

5

Amino acid sequence of the construct of the heavy chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 16

QVQLOESGGGLVKPGGSLRLSCAASGFTFSSYWMSWVROAPGKGLEWVANINRDGSASTYVD
SIKGRFTISRDDAKNSLYLQMNSLRAEDTAVYYCARDRGFGYFDLWGRGTLTVSSGGGGSG
GGGSGGGGSOSALTOPASVSGSPGOSITISCTGTSSDFGGYNFVSWYQOHPGKAPKLMYDVS
DRPSGVSDRFSGSKSGNTASLIISGLQADDEADYYCSSYGSSSTHIFGGGTKVTVLGGGGSGG
GGSQVQLKQSGPGLVQPSQSLSITCTVSGFSLTNYGVHWVROSPGKGLEWLGVWSSGGNTDY
NTPFTSRLSINKDNSKSQVFFKMNSLQSNDAIYYCARALTYDYEFAYWGQGLTVTVSSASTK
GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLYSL
SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLF
PPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR
VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKISKAKGQPREPQVYTLPPSRDELTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQ
GNVFSCSVMHEALHNHYTQKSLSLSPG

where the variable region of the heavy chain is:

SEQ ID NO: 38

QVQLKQSGPGLVQPSQSLSITCTVSGFSLTNYGVHWVROSPGKGLEWLGVWSSGGNTIDYNT
FTSRLSINKDNSKSQVFFKMNSLQSNDAIYYCARALTYDYEFAYWGQGLTVTVSS

- 5 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 29), CDR2 (SEQ ID NO: 30), and CDR3 (SEQ ID NO: 31), respectively.

The variable region of the heavy chain in the structural domain of the single-chain Fv (scFv) is:

10 SEQ ID NO: 39

QVQLQESGGGLVKPGGSLRLSCAASGFTFSSYWMSWVROAPGKGLEWVANINRDGSASTYVD
SVKGRFTISRDDAKNSLYLQMNSLRAEDTAVYYCARDRGFGYFDLWGRGTLTVSS

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 32), CDR2 (SEQ ID NO: 33), and CDR3 (SEQ ID NO: 34), respectively.

- 15 The variable region of the light chain in the single-chain Fv (scFv) structural domain is:

SEQ ID NO: 40

QSALTQPASVSGSPGQSITISCTGTSSDFGGYNFVSWYQOHPGKAPKLMYDVSDRPSGVSDRF
SGSKSGNTASLIISGLQADDEADYYCSSYGSSSTHIFGGGTKVTVL

- 20 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 35), CDR2 (SEQ ID NO: 36), and CDR3 (SEQ ID NO: 37), respectively.

Example 92

1) Expression and purification of SI-1×4 antibody:

- (Shanghai OPM Biosciences Co., Ltd.) suspension cells were
 25 used to express SI-1×4 antibody. The day before transfection, cells were inoculated at a density of 0.9×10^6 cells/mL in a 1L shake flask containing 300 mL of OPM-293 CD05 Medium (81075-

001, Shanghai OPM Biosciences Co., Ltd.), and cultured overnight at 37°C, 5% CO₂, and 120 rpm in a cell culture shaker. On the next day, the antibody expression plasmid was transfected with PEI-MAX, wherein the mass ratio of plasmid:PEI-MAX was 1:3.

OPM-293 ProFeed supplement was added at 5% (v/v) on the first day after transfection, and then again at 5% (v/v) on the third day after transfection, and then centrifugation was performed to collect the supernatant on the sixth day after transfection.

The collected cell expression supernatant was collected and eluted with 0.05 M sodium acetate (pH 3.6) in a Protein A affinity chromatography column (UniMab 50, Suzhou Nanomicro Technology Co., Ltd.), and the captured antibody was adjusted to pH 7.0 with 1 M Tris-HCl (pH 8.8) at 0.7/10 (v/v), and then passed through a gel filtration chromatography column SEC (Superdex 200, GE) to remove impurities such as polymers, while the antibody buffer was replaced with 20 mM PB (pH 6.5).

Antibody SI-1×4:

Light chain nucleic acid coding sequence

SEQ ID NO: 5

GATATTCAAATGACTCAATCTCCTTCTTCTCTTTCTGCTTCTGTTGGTGATCGTGTTACT
ATTACTTGTGCTTCTTCTCAAAATATTGTTCAATGTAATACTTATCTTGATTGGT
ATCAACAACTCCTGGTAAAGCTCCTAACTTCTTATTTATAAAGTTTCTAATCGTTTT
TCTGGTGTTCCTTCTCGTTTTTCTGGTCTGGTCTGGTACTGATTTTACTTTTACTATT
TCTTCTCTTCAACCTGAAGATATTGCTACTTATTATTGTTTTCAATATTCTCATGTTTCCTT
GGACTTTTGGTCAAGGTACTAACTTCAAATTACTCGTACGGTGGCTGCACCATCTGT
CTTCATCTTCCCAGCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGC
CTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAACGCC
CTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCAC
CTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAG
TCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCA
ACAGGGGAGAGTGTTAG

Light chain amino acid sequence

SEQ ID NO: 6

DIOMTQSPSSLSASVGDRTITCRSSONIVHSNGNTYLDWYQQTPGKAPKLLIYKYSNRESGVP
SRFSGSGSGTDFTFTISLQPEDIAITYCFQYSHVPWTFGQGTKLOITRTVAAPSVFIFPPSDEQ
LKSGTASVVCLLNFPYREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSSLTLSK
ADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

wherein the variable region is:

SEQ ID NO: 44

DIOMTQSPSSLSASVGDRTITCRSSONIVHSNGNTYLDWYQQTPGKAPKLLIYKYSNRESGVP
SRFSGSGSGTDFTFTISLQPEDIAITYCFQYSHVPWTFGQGTKLOIT

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 41), CDR2 (SEQ ID NO: 42), and CDR3 (SEQ ID NO: 43), respectively.

- 5 Nucleic acid coding sequence for the construct of the heavy chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 7

CAGGTGCAGCTGCAGCAGAGCGGCGCCGAGGTGAAGAAGCCCGGCAGCAGCGTGA
AGGTGAGCTGCAAGGCCAGCGGCTACACCTTCACCAACTACTACATCTACTGGGTGC
GGCAGGCCCCCGGCCAGGGCCTGGAGTGGATCGGCGGCATCAACCCCAACAGCGGC
GGCAGCAACTTCAACGAGAAGTTCAAGACCCGGGTGACCATCACCGCCGACGAGAG
CAGCACCACCGCCTACATGGAGCTGAGCAGCCTGCGGAGCGAGGACACCGCCTTCTA
CTTCTGCACCCCGGCAGGGCCTGTGGTTCGACAGCGACGGCCGGGGCTTCGACTTCTG
GGGCCAGGGCACCAACCGTGACCGTGAGCAGCGCTAGCACCAAGGGCCCATCGGTCT
TCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCC
TGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCGCCCTGA
CCAGCGGCGTGACACACCTTCCCGGCTGTCTACAGTCCTCAGGACTCTACTCCCTCA
GCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACG
TGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGT
GACAAAACCTCACACATGCCACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTC
AGTCTTCTCTTCCCCCAAAACCCAAGGACACCTCATGATCTCCCGGACCCCTGA
GGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCTGAGGTCAAGTTCAACT
GGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAG
TACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTG
AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGA
GAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGC
CCCCATCCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAG
GCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC
AACTACAAGACCACGCCTCCCGTGTCTGGACTCCGACGGCTCCTTCTTCTCTATAGCA
AGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTG
ATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGT
GGCGGTGGAGGGTCCGGCGGTGGTGGATCACAGGTGCAATTGCAGGAGTCGGGGGG
AGGCCTGGTCAAGCCTGGAGGGTCCCTGAGACTCTCCTGTGCAGCCTCTGGATTAC
CTTAGTAGTTATTGGATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTG
GGTGGCCAACATAAACCGCGATGGAAGTGCGAGTTACTATGTGGACTCTGTGAAGGG
CCGATTACCATCTCCAGAGACGACGCCAAGAACTCACTGTATCTGCAAATGAACAG
CCTGAGAGCTGAGGACACGGCTGTGTATTACTGTGCGAGAGATCGTGGGGTGGGCTA
CTTCGATCTCTGGGGCCGTGGCACCCCTGGTACCGTCTCGAGCGGTGGAGGCGGTTT
AGGCGGAGGTGGTTCCGGCGGTGGCGGCTCCCAGTCTGCCCTGACTCAGCCTGCCTC
CGTGTCTGGGTCTCCTGGACAGTCGATCACCATCTCCTGCACTGGAACCAGCAGTGA
CGTTGGTGGTTATAACTTTGTCTCCTGGTACCAACAACACCCAGGCAAAGCCCCAA
ACTCATGATCTATGATGTCAGTGATCGGCCCTCAGGGGTGTCTGATCGCTTCTCCGGC
TCCAAGTCTGGCAACACGGCCTCCCTGATCATCTCTGGCCTCCAGGCTGACGACGAG
CCTGATTATTACTGCAGCTCATATGGGAGCAGCAGCACTCATGTGATTTTCGGCGGAG
GGACCAAGGTGACCGTCCTATAA

- 10 Amino acid sequence of the construct of heavy chain and single chain Fv (scFv) structural domain

SEQ ID NO: 8

Nucleic acid coding sequence for the construct of light chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 17

CAGGTGCAGCTGCAGGAGTCGGGGGGAGGCCTGGTCAAGCCTGGAGGGTCCCTGAG
TCTCTCCTGTGCAGCCTCTGGATTACCTTTAGTAGTTATTGGATGAGCTGGGTCCGCC
AGGCTCCAGGGAAGGGGCTGGAGTGGGTGGCCAACATAAACC GCGATGGAAGTGCG
AGTTACTATGTGGACTCTGTGAAGGGCCGATTACCATCTCCAGAGACGACGCCAAG
AACTCACTGTATCTGCAAATGAACAGCCTGAGAGCTGAGGACACGGCTGTGTATTAC
TGTGCGAGAGATCGTGGGGTGGGCTACTTCGATCTCTGGGGCCGTGGCACCCTGGTC
ACCGTCTCGAGCGGTGGAGGCGGTTAGGCGGAGGTGGTTCCGGCGGTGGCGGCTC
CCAGTCTGCCCTGACTCAGCCTGCCTCCGTGTCTGGGTCTCCTGGACAGTCGATCACC
ATCTCCTGCACTGGAACCAGCAGTGACGTTGGTGGTTATAACTTTGTCTCCTGGTACC
AACAACACCCAGGCAAAGCCCCCAAACATCATGATCTATGATGTCAGTGATCGGCCCT
CAGGGGTGTCTGATCGCTTCTCCGGCTCCAAGTCTGGCAACACGGCCTCCCTGATCAT
CTCTGGCCTCCAGGCTGACGACGAGGCTGATTATTACTGCAGCTCATATGGGAGCAGC
AGCACTCATGTGATTTTCGGCGGAGGGACCAAGGTGACCGTCCTAGGCGGTGGAGG
ATCCGGCGGTGGTGGATCAGACATCTTGCTGACTCAGTCTCCAGTCATCCTGTCTGTG
AGTCCAGGAGAAAGAGTCAGTTTCTCCTGCAGGGCCAGTCAGAGTATTGGCACAAA
CATACTGGTATCAGCAAAGAACAATGGTTCTCCAAGGCTTCTCATAAAGTATGCT
TCTGAGTCTATCTCTGGGATTCTTCCAGGTTTAGTGGCAGTGGATCAGGGACAGATT
TTACTCTTAGCATCAACAGTGTGGAGTCTGAAGATATTGCAGATTATTACTGTCAACA
AAATAATAACTGGCCAACCACGTTCCGGTTGTGGGACCAAGCTGGAGCTGAAACGTAC
GGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGA
ACTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGGCCAAAGTACAGT
GGAAGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAG
GACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGA
CTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCC
CGTCACAAAGAGCTTCAACAGGGGAGAGTGTTAG

5

Amino acid sequence of construct of light chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 18

QVLOESGGGLVKPGGSLSLSCAASGFTFSSYWMSWVROAPGKGLEWVANINRDGSASYYID
SVKGRFTISRDDAKNSLYLQMNSLRAEDTAVYYCARDRGVGYFDLWGRGTLVTVSSGGGGSG
GGSGGGGSQSALTOPASVSGSPGOSITISCTGTSSDVGGYNEFYSWYQQHPGKAPKLMYDES
DRPSGVSDRPSGSKSGNTASLIHSGLOADDEADYYCSSYGSSTHVTEGGGTKVTVLGGGGSGG
GGSDILLTOSPVILSVSPGERVSFSCRASOSIGNIHWYQORTNGSPRLIKYASESISGIPSRFSG
SGSGTDFTLSINSVESEDIADYYCOQNNNWPTTFGCGTKLELKRTVAAPSVFIEPPSDEQLKS
GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKAD
YEKHKVYACEVTHQGLSSPVTKSFNRGEC

10 where the variable region of the light chain is:

SEQ ID NO: 49

DILLTQSPVILSVSPGERVSFSCRASOSIGNIHWYQORTNGSPRLIKYASESISGIPSRFSGSGS
GTDFTLSINSVESEDIADYYCOQNNNWPTTFGCGTKLELK

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 25), CDR2 (SEQ ID NO: 26), and CDR3 (SEQ ID NO: 27),
15 respectively.

The variable region of the heavy chain in the structural domain of the single-chain Fv (scFv) is:

SEQ ID NO: 50

5 QVQLQESGGGLV¹PKPGSLS²LS³CAASGFTFS⁴SYWMS⁵WVRQAPGKGL⁶EWVANINRDK⁷ISASY⁸YVD⁹
SI¹⁰KGRFTISRDDAKNSLYLQMNSLRAEDTAVYYC¹¹ARDRGV¹²GYFDL¹³WGRGTLVTVSS¹⁴

where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 32), CDR2 (SEQ ID NO: 33), and CDR3 (SEQ ID NO: 34), respectively.

10 The variable region of the light chain in the single-chain Fv (scFv) structural domain is:

SEQ ID NO: 51

QSALTQPASVSGSPGQSITISC¹TGTSSDVGGYNF²SWYQQHPGKAPKLMYD³SDRPSGVSDRF⁴
SGSKSGNTASLIISGLQADDEADYYC⁵SSYGSS⁶THVIFGGGTKVTVL⁷

15 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 35), CDR2 (SEQ ID NO: 36), and CDR3 (SEQ ID NO: 37), respectively.

Nucleic acid coding sequence for the heavy chain

SEQ ID NO: 19

CAGGTGCAGCTGAAGCAGTCAGGACCTGGCCTAGTGCAGCCCTCACAGAGCCTGTC
CATCACCTGCACAGTCTCTGGTTTCTCATTAATACTAATGGTGTACACTGGGTTCGC
CAGTCTCCAGGAAAGTGCCTGGAGTGGCTGGGAGTGATATGGAGTGGTGGAACAC
AGACTATAATACACCTTTCACATCCAGACTGAGCATCAACAAGGACAATTCCAAGA
GCCAAGTTTCTTTAAATGAACAGTCTGCAATCTAATGACACAGCCATATATTACT
GTGCCAGAGCCCTCACCTACTATGATTACGAGTTTGCTTACTGGGGCCAAGGGACTC
TGGTCACTGTCTGTCTGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCT
CCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTAC
TTCCCCGAACCGGTGACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCA
CACCTTCCCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGAC
CGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCC
CAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGTGACAAAACCTCACA
CATGCCCACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCTCTTCC
CCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTG
GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGG
CGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACG
TACCGTGTGGTCAGCGTCTCACCGTCTGCACCAGGACTGGCTGAATGGCAAGGA
GTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGAGAAAACCATCT
CCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGG
GATGAGCTGACCAAGAACCAGGTACGCTGACCTGCCTGGTCAAAGGCTTCTATCCC
AGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGA
CCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTATAGCAAGCTCACCGT
GGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGG
CTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAA

20 Amino acid sequence of the heavy chain

SEQ ID NO: 20

OVOLKQSGPGLVOPQSLSITCTVSGFSLTNYGVHWVROSPGKCLEWLGVIWSGGNTDYNIP
ETSRLSINKDNSKSOVTFKMNSLOSNDTAIYYCARALTYDYEFAYWGQGLVTVSAASTKGPS
VFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSS
VVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPP
PKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV
VSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQG
NVFSCSVMHREALHNHYTQKSLSLSPG

where the variable region of the heavy chain is:

5 SEQ ID NO: 52

QVQLKQSGPGLVQPSQSLITCTVSGFSLTNYGVHWVROSPGKCLEWLGVIWSGGNTDYNIP
ETSRLSINKDNSKSOVTFKMNSLOSNDTAIYYCARALTYDYEFAYWGQGLVTVSA

where the underlined portions are, from left to right, CDR1
(SEQ ID NO: 29), CDR2 (SEQ ID NO: 30), and CDR3 (SEQ ID NO: 31),
respectively.

10

(3) Expression and purification of SI-1×26 antibody: a
similar method was followed for the expression and
purification of SI-1×26 antibody.

Antibody SI-1×26:

15 Nucleic acid coding sequence for the construct of light
chain and single-chain Fv (scFv) structural domain

SEQ ID NO: 21

SEQ ID NO: 50

QVQLQESGGGLVKPGGSLSLSCAASGFTFSSYWMNWVRQAPGKGLEWVANINRDGSASYVVD
SVKGRFTISRDDAKNSLYLQMNSLRAEDTAVYYCARDRGVGYFDLWGRGTLTVSS

where the underlined portions are, from left to right, CDR1
(SEQ ID NO: 32), CDR2 (SEQ ID NO: 33), and CDR3 (SEQ ID NO: 34),
5 respectively.

The variable region of the light chain in the single-chain
Fv (scFv) structural domain is:

SEQ ID NO: 51

QSALTQPASVSGSPGQSIIISCTGTSSDVGGYNFTSWYQQHPGKAPKLMYDVSDRPSGVSDRF
SGSKSGNTASLIISGLQADDEADYYCNSYGSSSIHVIFGGGTKVTVL

10 where the underlined portions are, from left to right, CDR1
(SEQ ID NO: 35), CDR2 (SEQ ID NO: 36), and CDR3 (SEQ ID NO: 37),
respectively.

Nucleic acid coding sequence for the heavy chain

15 SEQ ID NO: 23

CAGGTGCAGCTGCAGCAGAGCGGCGCCGAGGTGAAGAAGCCCCGGCAGCAGCGTGA
AGGTGAGCTGCAAGGCCAGCGGCTACACCTTCACCAACTACTACATCTACTGGGTGC
GGCAGGCCCGGCCAGTGTCTGGAGTGGATCGGCGGCATCAACCCACCAGCGGC
GGCAGCAACTTCAACGAGAAGTTCAAGACCCGGGTGACCATCACCGCCGACGAGAG
CAGCACCACCGCCTACATGGAGCTGAGCAGCCTGCGGAGCGAGGACACCGCCTTCTA
CTTCTGCACCCGGCAGGGCCTGTGGTTCGACAGCGACGGCCGGGGCTTCGACTTCTG
GGGCCAGGGCACCACCGTGACCGTGAGCAGCGCTAGCACCAAGGGCCCATCGGTCT
TCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCC
TGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCGCCCTGA
CCAGCGGCGTGACACCTTCCCGGCTGTCTACAGTCCTCAGGACTCTACTCCCTCA
GCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACG
TGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGT
GACAAAACCTCACACATGCCACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTC
AGTCTTCTCTTCCCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGA
GGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACT
GGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAG
TACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTG
AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGA
GAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCTGC
CCCCATCCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAG
GCTTCTATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC
AACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTATAGCA
AGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTG
ATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGT
TAA

Amino acid sequence of the heavy chain

SEQ ID NO: 24

QVOLOQSGAEVKKPGSSVKVSCKASGYTFITNYIYWVROAPGQCLEWIGGINPTSGGSNENE
KFKTRVTITADESSTIAYMELSSLRSEDTAFYFCTROGLWFDSDGGRGFEFWGQGTITVTVSSAS
 TKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL
 YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPS
 VFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYN
 STYRVSVLTIVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRD
 ELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSR
 WQQGNVFSCSVMHEALHNHYTQKSLSLSPG

where the variable region of the heavy chain is:

SEQ ID NO: 54

QVOLOQSGAEVKKPGSSVKVSCKASGYTFITNYIYWVROAPGQCLEWIGGINPTSGGSNENE
KFKTRVTITADESSTIAYMELSSLRSEDTAFYFCTROGLWFDSDGGRGFEFWGQGTITVTVSS

5 where the underlined portions are, from left to right, CDR1 (SEQ ID NO: 45), CDR2 (SEQ ID NO: 46), and CDR3 (SEQ ID NO: 47), respectively.

Example 93

1) A sample of SI-1×6.4 antibody-drug conjugate was prepared
 10 by coupling SI-1×6.4 antibody with payload:

After cellular expression and purification by Protein A affinity chromatography and molecular sieve chromatography, the SI-1×6.4 antibody was replaced in 20 mM PB, pH 6.5 buffer, and the SI-1×6.4 antibody was concentrated or diluted to a protein
 15 concentration of 3 mg/mL. The payload was a white powder, which was dissolved to 20 mg/mL using DMA and set aside. To open the interchain disulfide bonds of SI-1×6.4 antibody, 20-fold TECP was first added according to the molecular ratio and a reaction was performed at room temperature for 3 h. Then 20-fold payload
 20 solution was added according to the molecular ratio and a reaction was performed at room temperature for 1 h. After the reaction was completed, the payload that was not coupled to SI-1×6.4 was removed by ultrafiltration using a 30 KDa ultrafiltration centrifuge tube, and the SI-1×6.4 antibody-drug
 25 conjugate sample was thus obtained.

2) A sample of SI-1×22 antibody-drug conjugate was prepared by coupling SI-1×22 antibody with payload following a similar method.

3) A sample of SI-1×24 antibody-drug conjugate was
 30 prepared by coupling SI-1×24 antibody with payload following a similar method.

5
10

15

20


25

Example 95

ADC-1


30

182




Example 97

ADC-3 was prepared according to the general-purpose coupling method of Example 93,




Example 98

ADC-4 was prepared according to the general-purpose coupling method of Example 93,




Example 99

ADC-5 was prepared according to the general-purpose coupling method of Example 93,




Example 100

ADC-6 was prepared according to the general-purpose coupling method of Example 93,




Example 105

ADC-11 was prepared according to the general-purpose coupling method of Example 93,




Example 106

ADC-12 was prepared according to the general-purpose coupling method of Example 93,




Example 107

ADC-13 was prepared according to the general-purpose coupling method of Example 93,




Example 108


ADC-14 was prepared according to the general-purpose coupling method of Example 93,




ADC-15 was prepared according to the general-purpose coupling method of Example 93,




ADC-16 was prepared according to the general-purpose coupling method of Example 93,




ADC-17 was prepared according to the general-purpose coupling method of Example 93,



ADC-18 was prepared according to the general-purpose coupling method of Example 93,




ADC-19 was prepared according to the general-purpose coupling method of Example 93,




Example 114

ADC-20 was prepared according to the general-purpose coupling method of Example 93,




Example 115

ADC-21 was prepared according to the general-purpose coupling method of Example 93,




Example 116

ADC-22 was prepared according to the general-purpose coupling method of Example 93,




Example 117

ADC-23 was prepared according to the general-purpose coupling method of Example 93,



Example 118


ADC-24 was prepared according to the general-purpose coupling method of Example 93,



5


Example 119

ADC-25 was prepared according to the general-purpose coupling method of Example 93,




Example 120

10 ADC-26 was prepared according to the general-purpose coupling method of Example 93,




Example 121

15 ADC-27 was prepared according to the general-purpose coupling method of Example 93,v




Example 122

ADC-28 was prepared according to the general-purpose coupling method of Example 93,




Example 123

ADC-29 was prepared according to the general-purpose coupling method of Example 93,




Example 124

ADC-30 was prepared according to the general-purpose coupling method of Example 93,




Example 125

ADC-31 was prepared according to the general-purpose coupling method of Example 93,




Example 126

ADC-32 was prepared according to the general-purpose coupling method of Example 93,




Example 127

ADC-33 was prepared according to the general-purpose coupling method of Example 93,



Example 128


5 ADC-34 was prepared according to the general-purpose coupling method of Example 93,



Example 129


ADC-35 was prepared according to the general-purpose coupling method of Example 93,

10



Example 130


ADC-36 was prepared according to the general-purpose coupling method of Example 93,



15


Example 131

ADC-37 was prepared according to the general-purpose coupling method of Example 93,




Example 132

ADC-38 was prepared according to the general-purpose coupling method of Example 93,




Example 133

ADC-39 was prepared according to the general-purpose coupling method of Example 93,




Example 134

ADC-40 was prepared according to the general-purpose coupling method of Example 93,




Example 135

ADC-41 was prepared according to the general-purpose coupling method of Example 93,




Example 136

ADC-42 was prepared according to the general-purpose coupling method of Example 93,




Example 137

ADC-43 was prepared according to the general-purpose coupling method of Example 93,




Example 138

ADC-44 was prepared according to the general-purpose coupling method of Example 93,




Example 139

ADC-45 was prepared according to the general-purpose coupling method of Example 93,




Example 140

ADC-46 was prepared according to the general-purpose coupling method of Example 93,




Example 141

ADC-47 was prepared according to the general-purpose coupling method of Example 93,




Example 142

ADC-48 was prepared according to the general-purpose coupling method of Example 93,




Example 143

ADC-49 was prepared according to the general-purpose coupling method of Example 93,




Example 144

ADC-50 was prepared according to the general-purpose coupling method of Example 93,




Example 145

ADC-51 was prepared according to the general-purpose coupling method of Example 93,



Example 146

ADC-52 was prepared according to the general-purpose coupling method of Example 93,



Example 147

ADC-53 was prepared according to the general-purpose coupling method of Example 93,



ADC-54 was prepared according to the general-purpose coupling method of Example 93,




ADC-55 was prepared according to the general-purpose coupling method of Example 93,



ADC-56 was prepared according to the general-purpose coupling method of Example 93,




ADC-57 was prepared according to the general-purpose coupling method of Example 93,




Example 152

ADC-58 was prepared according to the general-purpose coupling method of Example 93,




Example 153

ADC-59 was prepared according to the general-purpose coupling method of Example 93,




Example 154

ADC-60 was prepared according to the general-purpose coupling method of Example 93,




Example 155

ADC-61 was prepared according to the general-purpose coupling method of Example 93,




Example 156

ADC-62 was prepared according to the general-purpose coupling method of Example 93,




Example 157

ADC-63 was prepared according to the general-purpose coupling method of Example 93,




Example 158

ADC-64 was prepared according to the general-purpose coupling method of Example 93,




Example 159

ADC-65 was prepared according to the general-purpose coupling method of Example 93,




Example 160

ADC-66 was prepared according to the general-purpose coupling method of Example 93,




Example 161

ADC-67 was prepared according to the general-purpose coupling method of Example 93,




Example 162

ADC-68 was prepared according to the general-purpose coupling method of Example 93,




Example 163

ADC-69 was prepared according to the general-purpose coupling method of Example 93,




Example 164

ADC-70 was prepared according to the general-purpose coupling method of Example 93,




Example 165

ADC-71 was prepared according to the general-purpose coupling method of Example 93,




Example 166

ADC-72 was prepared according to the general-purpose coupling method of Example 93,




Example 167

ADC-73 was prepared according to the general-purpose coupling method of Example 93,



Example 168


ADC-74 was prepared according to the general-purpose coupling method of Example 93,



5


Example 169

ADC-75 was prepared according to the general-purpose coupling method of Example 93,




Example 170

10 ADC-76 was prepared according to the general-purpose coupling method of Example 93,




Example 171

15 ADC-77 was prepared according to the general-purpose coupling method of Example 93,




Example 172

ADC-78 was prepared according to the general-purpose coupling method of Example 93,




Example 173

ADC-79 was prepared according to the general-purpose coupling method of Example 93,




Example 174

ADC-80 was prepared according to the general-purpose coupling method of Example 93,




Example 175

ADC-81 was prepared according to the general-purpose coupling method of Example 93,




Example 176

ADC-82 was prepared according to the general-purpose coupling method of Example 93,




Example 177

ADC-83 was prepared according to the general-purpose coupling method of Example 93,




Example 178

ADC-84 was prepared according to the general-purpose coupling method of Example 93,




Example 179

ADC-85 was prepared according to the general-purpose coupling method of Example 93,




Example 180

ADC-86 was prepared according to the general-purpose coupling method of Example 93,




Example 181

ADC-87 was prepared according to the general-purpose coupling method of Example 93,




Example 182

ADC-88 was prepared according to the general-purpose coupling method of Example 93,




Example 183

ADC-89 was prepared according to the general-purpose coupling method of Example 93,




Example 184

ADC-90 was prepared according to the general-purpose coupling method of Example 93,



Example 185


ADC-91 was prepared according to the general-purpose coupling method of Example 93,



5


Example 186

ADC-92 was prepared according to the general-purpose coupling method of Example 93,




Example 187

10 ADC-93 was prepared according to the general-purpose coupling method of Example 93,




Example 188

15 ADC-94 was prepared according to the general-purpose coupling method of Example 93,




Example 189

ADC-95 was prepared according to the general-purpose coupling method of Example 93,




Example 190

ADC-96 was prepared according to the general-purpose coupling method of Example 93,




Example 191

ADC-97 was prepared according to the general-purpose coupling method of Example 93,




Example 192

ADC-98 was prepared according to the general-purpose coupling method of Example 93,




Example 193

ADC-99 was prepared according to the general-purpose coupling method of Example 93,



Example 194


ADC-100 was prepared according to the general-purpose coupling method of Example 93,



5

Example 195


ADC-101 was prepared according to the general-purpose coupling method of Example 93,



10


Example 196

ADC-102 was prepared according to the general-purpose coupling method of Example 93,




Example 197


15 ADC-103 was prepared according to the general-purpose coupling method of Example 93,




ADC-104 was prepared according to the general-purpose coupling method of Example 93,




ADC-105 was prepared according to the general-purpose coupling method of Example 93,




ADC-106 was prepared according to the general-purpose coupling method of Example 93,



ADC-DS was prepared from compound 45 according to the general-purpose coupling method of Example 93,




ADC-107 was prepared according to the general-purpose coupling method of Example 94,




Example 203

ADC-108 was prepared according to the general-purpose coupling method of Example 94,




Example 204

ADC-109 was prepared according to the general-purpose coupling method of Example 94,




Example 205

ADC-110 was prepared according to the general-purpose coupling method of Example 94,




Example 206

ADC-111 was prepared according to the general-purpose coupling method of Example 94,




Example 207

ADC-112 was prepared according to the general-purpose coupling method of Example 94,




Example 208

ADC-113 was prepared according to the general-purpose coupling method of Example 94,




Example 209

ADC-114 was prepared according to the general-purpose coupling method of Example 94,




Example 210

ADC-115 was prepared according to the general-purpose coupling method of Example 94,




Example 211

ADC-116 was prepared according to the general-purpose coupling method of Example 94,




Example 212

ADC-117 was prepared according to the general-purpose coupling method of Example 94,




Example 213

ADC-118 was prepared according to the general-purpose coupling method of Example 94,




Example 214

ADC-119 was prepared according to the general-purpose coupling method of Example 94,



Example 215


ADC-120 was prepared according to the general-purpose coupling method of Example 94,



5


Example 216

ADC-121 was prepared according to the general-purpose coupling method of Example 94,




Example 217

10 ADC-122 was prepared according to the general-purpose coupling method of Example 94,



Example 218

15 ADC-123 was prepared according to the general-purpose coupling method of Example 94,




Example 219

ADC-124 was prepared according to the general-purpose coupling method of Example 94,

Example 224


ADC-129 was prepared according to the general-purpose coupling method of Example 94,



5


Example 225

ADC-130 was prepared according to the general-purpose coupling method of Example 94,




Example 226

10 ADC-131 was prepared according to the general-purpose coupling method of Example 94,




Example 227

15 ADC-132 was prepared according to the general-purpose coupling method of Example 94,




Example 228

ADC-133 was prepared according to the general-purpose coupling method of Example 94,




Example 229

ADC-134 was prepared according to the general-purpose coupling method of Example 94,




Example 230

ADC-135 was prepared according to the general-purpose coupling method of Example 94,




Example 231

ADC-136 was prepared according to the general-purpose coupling method of Example 94,




Example 232

ADC-137 was prepared according to the general-purpose coupling method of Example 94,




Example 233

ADC-138 was prepared according to the general-purpose coupling method of Example 94,




Example 234

5 ADC-139 was prepared according to the general-purpose coupling method of Example 94,




Example 235

10 ADC-140 was prepared according to the general-purpose coupling method of Example 94,




Example 236

ADC-141 was prepared according to the general-purpose coupling method of Example 94,




Example 237

ADC-142 was prepared according to the general-purpose coupling method of Example 94,




Example 238

ADC-143 was prepared according to the general-purpose coupling method of Example 94,




Example 239

ADC-144 was prepared according to the general-purpose coupling method of Example 94,




Example 240

ADC-145 was prepared according to the general-purpose coupling method of Example 94,




Example 241

ADC-146 was prepared according to the general-purpose coupling method of Example 94,




Example 242

ADC-147 was prepared according to the general-purpose coupling method of Example 94,




Example 243

ADC-148 was prepared according to the general-purpose coupling method of Example 94,




Example 244

ADC-149 was prepared according to the general-purpose coupling method of Example 94,



Example 245



ADC-150 was prepared according to the general-purpose coupling method of Example 94,




ADC-151 was prepared according to the general-purpose coupling method of Example 94,



ADC-152 was prepared according to the general-purpose coupling method of Example 94,




ADC-155 was prepared according to the general-purpose coupling method of Example 94,




Example 251

ADC-156 was prepared according to the general-purpose coupling method of Example 94,




Example 252

ADC-157 was prepared according to the general-purpose coupling method of Example 94,




Example 253

ADC-158 was prepared according to the general-purpose coupling method of Example 94,




Example 254

ADC-159 was prepared according to the general-purpose coupling method of Example 94,




Example 255

ADC-160 was prepared according to the general-purpose coupling method of Example 94,




Example 256

ADC-161 was prepared according to the general-purpose coupling method of Example 94,




Example 257

ADC-162 was prepared according to the general-purpose coupling method of Example 94,




Example 258

ADC-163 was prepared according to the general-purpose coupling method of Example 94,




Example 259

ADC-164 was prepared according to the general-purpose coupling method of Example 94,




Example 260

ADC-165 was prepared according to the general-purpose coupling method of Example 94,




Example 261

ADC-166 was prepared according to the general-purpose coupling method of Example 94,




Example 262

ADC-167 was prepared according to the general-purpose coupling method of Example 94,




Example 267

ADC-172 was prepared according to the general-purpose coupling method of Example 94,




Example 268

ADC-173 was prepared according to the general-purpose coupling method of Example 94,




Example 269

ADC-174 was prepared according to the general-purpose coupling method of Example 94,




Example 270

ADC-175 was prepared according to the general-purpose coupling method of Example 94,



Example 271


ADC-176 was prepared according to the general-purpose coupling method of Example 94,



5

Example 272


ADC-177 was prepared according to the general-purpose coupling method of Example 94,



10


Example 273

ADC-178 was prepared according to the general-purpose coupling method of Example 94,




Example 274

15 ADC-179 was prepared according to the general-purpose coupling method of Example 94,



Example 275


ADC-180 was prepared according to the general-purpose coupling method of Example 94,



5


Example 276

ADC-181 was prepared according to the general-purpose coupling method of Example 94,




Example 277

10 ADC-182 was prepared according to the general-purpose coupling method of Example 94,




Example 278

15 ADC-183 was prepared according to the general-purpose coupling method of Example 94,




Example 279

ADC-184 was prepared according to the general-purpose coupling method of Example 94,




Example 280

ADC-185 was prepared according to the general-purpose coupling method of Example 94,




Example 281

ADC-186 was prepared according to the general-purpose coupling method of Example 94,




Example 282

ADC-187 was prepared according to the general-purpose coupling method of Example 94,




Example 283

ADC-188 was prepared according to the general-purpose coupling method of Example 94,




Example 284

ADC-189 was prepared according to the general-purpose coupling method of Example 94,




Example 285

ADC-190 was prepared according to the general-purpose coupling method of Example 94,




Example 286

ADC-191 was prepared according to the general-purpose coupling method of Example 94,




Example 287

ADC-192 was prepared according to the general-purpose coupling method of Example 94,




Example 288

ADC-193 was prepared according to the general-purpose coupling method of Example 94,




Example 289

ADC-194 was prepared according to the general-purpose coupling method of Example 94,




Example 290

ADC-195 was prepared according to the general-purpose coupling method of Example 94,




Example 291

ADC-196 was prepared according to the general-purpose coupling method of Example 94,



Example 292


ADC-197 was prepared according to the general-purpose coupling method of Example 94,



5


Example 293

ADC-198 was prepared according to the general-purpose coupling method of Example 94,




Example 294

10 ADC-199 was prepared according to the general-purpose coupling method of Example 94,




Example 295

15 ADC-200 was prepared according to the general-purpose coupling method of Example 94,




Example 296

ADC-201 was prepared according to the general-purpose coupling method of Example 94,




Example 297

ADC-202 was prepared according to the general-purpose coupling method of Example 94,




Example 298

ADC-203 was prepared according to the general-purpose coupling method of Example 94,




Example 299

ADC-204 was prepared according to the general-purpose coupling method of Example 94,




Example 300

ADC-205 was prepared according to the general-purpose coupling method of Example 94,




Example 301

ADC-206 was prepared according to the general-purpose coupling method of Example 94,




Example 302

ADC-207 was prepared according to the general-purpose coupling method of Example 94,




Example 303

ADC-208 was prepared according to the general-purpose coupling method of Example 94,




Example 304


ADC-209 was prepared according to the general-purpose coupling method of Example 94,




ADC-210 was prepared according to the general-purpose coupling method of Example 94,




ADC-211 was prepared according to the general-purpose coupling method of Example 94,




ADC-212 was prepared according to the general-purpose coupling method of Example 94,



ADC-213 was prepared from compound 45 according to the general-purpose coupling method of Example 94,




ADC-214 was prepared from compound 5A according to the general-purpose coupling method of Example 93,



Example 310


ADC-215 of HER3 was prepared from compound 5A according to the general-purpose coupling method of Example 93,




where the H3 antibody is the portion of the SI-1×6.4 antibody knockout targeting EGFR.

Example 311


ADC-216 to ADC-223 were prepared according to the general-purpose coupling method of Example 93,






Example 312

ADC-224 to ADC-231 were prepared according to the general-purpose coupling method of Example 93,



Example 313


ADC-232 to ADC-239 were prepared according to the general-purpose coupling method of Example 94,




5

Example 314

ADC-240 to ADC-247 were prepared according to the general-purpose coupling method of Example 94,





Example 315

The monomer rate was determined using the SEC-HPLC method:

Chromatography column: Biocore SEC-300 5 μ m, 4.6×300 mm

Manufacturer: NanoChrom, Item No.: B213-050030-04630S

5 Mobile phase: 50 mM PB + 300 mM NaCl + 200 mM Arg + 5% IPA,
pH = 6.5

Table 1: Methodological parameters

Parameters	Settings
Flow rate	0.3 mL/min
Wavelengths	214 nm and 280 nm
Column temperature	30°C
Sample plate temperature	room temperature
Injection volume	20 μ g
Maximum pressure	150 bar/15 MPa/2175 PSI
Gradient	isocratic
Operating time	20 minutes

10 Table 2. Monomer rate data for the disclosed ligand-drug conjugate (ADC) disclosed in the present application

Molecule name	Degradation %	Aggregate %	Monomer rate %
ADC-1	0.01	1.60	98.39
ADC-2	0.0	1.51	98.49
ADC-5	0.0	2.43	97.57
ADC-6	0.0	2.42	97.58
ADC-7	0.12	2.16	97.72
ADC-8	0.07	1.37	98.56
ADC-10	0.05	1.35	98.60
ADC-11	0.11	1.51	98.38
ADC-12	0.05	1.62	98.33
ADC-13	0.08	1.30	98.62
ADC-14	0.02	1.49	98.49

ADC-48	0.06	1.71	98.23
ADC-52	0.05	1.56	98.39
ADC-DS	0.0	2.82	97.18
ADC-64	0.0	1.16	98.84
ADC-68	0.01	0.44	98.55
ADC-75	0.0	1.61	98.39
ADC-81	0.2	1.59	98.39
ADC-87	0.07	1.30	98.63
ADC-96	0.08	1.32	98.60
ADC-102	0.05	1.31	98.64
ADC-108	0.0	6.36	93.64
ADC-112	1.2	2.81	95.99
ADC-113	0.03	1.22	98.75
ADC-120	0.0	1.73	98.27
ADC-122	0.01	1.44	98.55
ADC-129	0.13	2.54	97.33
ADC-130	0.37	2.18	97.45
ADC-131	0.60	1.00	98.40
ADC-132	0.03	2.22	97.75
ADC-133	0.53	1.37	98.10
ADC-145	0.11	1.24	98.65
ADC-158	0.98	2.89	96.13
ADC-160	0.10	1.45	98.45
ADC-172	0.65	1.31	98.04
ADC-179	0.1	1.06	98.84
ADC-188	0.11	1.33	98.56
ADC-193	0.11	1.51	98.38
ADC-206	0.09	1.32	98.59
ADC-214	0.03	1.18	98.79
ADC-215	0.0	3.01	96.99
ADC-219	0.0	7.35	92.65
ADC-227	0.06	4.5	95.44
ADC-235	0.1	6.83	93.07

CONCLUSION: The ADC disclosed in the present invention is

characterized by a low degradation rate and a low aggregation rate, and has the excellent property of a high monomer rate.

Example 316

The drug antibody ratio DAR was determined using the RP-HPLC method:

Chromatography column name: Proteomix RP-1000 4.6×100mm 5µm 1000A Manufacturer: Sepax

Table 3: Methodological parameters

Parameters	Settings			
Mobile phase	A: 0.1% TFA aqueous solution; B: 0.1% TFA acetonitrile solution			
Flow rate	0.5 mL/min			
Wavelength	214 nm and 280 nm			
Column temperature	65°C			
Sample plate temperature	room temperature			
Injection volume	25 ug			
Maximum pressure	100bar/10MPa/1450PSI			
Gradient	Time (min)	Flow rate (mL/min)	Mobile phase A (%)	Mobile phase B (%)
	0.0	0.5	75	25
	3	0.5	75	25
	28	0.5	50	50
	30	0.5	5	95
	32	0.5	5	95
	33	0.5	75	25
	40	0.5	75	25

Table 4: Detailed data on ADC drug-antibody coupling ratio (DAR)

Sample Name	Batch number	RP - DAR	Sample Name	Batch number	RP - DAR
ADC-5	20201028	7.65	ADC-120	20200708	7.83
ADC-6	20200923	6.99	ADC-122	20200708	6.97
ADC-10	20200104	7.38	ADC-137	20191125	7.44
ADC-12	20191125	7.61	ADC-140	20200708	7.52
ADC-18	20201028	7.71	ADC-142	20201028	7.50
ADC-23	20200104	7.19	ADC-143	20201028	7.61
ADC-24	20200104	7.09	ADC-159	20200708	7.28
ADC-36	20200708	7.44	ADC-160	20200708	7.89
ADC-41	20201028	7.81	ADC-168	20201028	7.24
ADC-42	20201028	7.55	ADC-169	20201028	7.89
ADC-55	20200708	7.54	ADC-170	20201028	7.55
ADC-59	20200708	7.57	ADC-171	20201028	6.91
ADC-64	20201028	7.42	ADC-176	20200708	7.32
ADC-70	20200104	7.60	ADC-183	20200708	7.18
ADC-71	20200104	7.50	ADC-184	20200708	7.49
ADC-72	20200104	7.42	ADC-185	20200708	7.30
ADC-88	20200708	7.31	ADC-	20200708	7.77

			186		
ADC-96	20191125	7.75	ADC-190	20200708	7.26
ADC-100	20200708	7.56	ADC-199	20191125	7.42
ADC-108	20200708	7.62	ADC-214	20201028	7.85
ADC-112	20201116	7.46	ADC-215	20211013	7.38
ADC-219	20211015	7.14	ADC-227	20211013	7.46
ADC-235	20211013	7.26	ADC-243	20211027	7.38

CONCLUSION: The ADC disclosed in the present invention has the excellent property of high DAR value, which can significantly increase the concentration of the drug at the target site location at the same dose of ADC drug administered.

Example 317

ADC maintained the affinity of the corresponding original bispecific antibodies SI-1×6.4, SI-1×4, SI-1×22, SI-1×24, SI-1×25 and SI-1×26 for EGFR and HER3:

The relative affinities of SI-1×6.4 versus ADC-6 and SI-1×4 versus ADC-112 for EGFR and HER3 were compared by double-antigen sandwich ELISA. The specific steps were as follows:

Recombinant EGFR-His*6 antigen-coated plates were closed with 1% bovine serum protein; then SI-1×6.4, ADC-6, SI-1×4, and ADC-112 were diluted, respectively, and then diluted with a starting concentration of 5,000 ng/mL in successive 3-fold gradients, for a total of 11 concentrations; the samples were incubated on the coated ELISA plates for a certain period of time, followed by biotin-labeled HER3-Fc antigen incubation, followed by streptavidin HRP-labeled incubation; finally, TMB color development was performed, followed by sulfuric acid solution termination, and the absorbance value at 450 nm was

detected on a microplate reader. The assay results were plotted against concentration for 0D450 nm.

CONCLUSION: As shown in the accompanying Figures 3A and 3B, after coupling, ADC-6 and ADC-112 maintained similar affinities to those of SI-1×6.4 and SI-1×4, respectively, with no significant difference in EC50 values; indicating that coupling of toxins to SI-1×6.4 and SI-1×4 did not affect their affinity for the antigen.

Similarly, using a similar testing method to that described above, the results are shown in FIG. 3C, FIG. 3D, FIG. 3E, and FIG. 3F; after coupling, ADC-219, ADC-227, ADC-235, and ADC-243 retained similar affinities to those of SI-1×22, SI-1×24, SI-1×25, and SI-1×26, respectively, with no significant difference in the EC50 values; indicating that coupling of toxins to SI-1×22, SI-1×24, SI-1×25 and SI-1×26 did not affect their affinity for the antigen.

Example 318

In vitro pharmacodynamic assays:

A variety of human-derived tumor cell lines (human epidermal cancer cells A431, human *in situ* pancreatic adenocarcinoma cells BXPC-3, human pharyngeal squamous cell carcinoma cells FaDu, human lung cancer squamous cell carcinoma cell line HARA-B, human non-small cell lung cancer cells HCC827, and human colon cancer cells SW620) were utilized as experimental models for evaluating *in vitro* efficacy of the ADC-coupled drug in the present invention. A certain number of tumor cells were inoculated in a 96-well plate, and the gradient-diluted test antibody and the corresponding ADC drug were added to the cells, followed by 5 days of treatment; the cell viability was measured using Alamar Blue or MTS, and the inhibitory effect of the test antibody and ADC on the tumor cell lines was evaluated by calculating the IC50. The starting concentration of the antibody drug was 500 nM, and the dilution was 7-fold, totaling 8 concentration points, and the treatment was carried out for 5

days. The final calculation method was survival rate = (experimental group - blank) / (control group - blank group) × 100%, then Graph Pad Prism was used for curve fitting, to calculate the half inhibitory concentration (IC₅₀) as well as Efficacy (%).

Table 5: *In vitro* efficacy of six naked antibodies (SI-1X4, SI-1×6.4, SI-1X22, SI-1X24, SI-1X25, SI-1X26) and six ADCs (ADC-112, ADC-6, ADC-219, ADC-227, ADC-235, ADC-243) in A431

A431			
Groups	IC ₅₀ (nM)	Groups	Efficacy (%)
ADC-112	4.95	ADC-112	85.63
ADC-6	0.98	ADC-6	87.88
ADC-219	0.75	ADC-219	89.37
ADC-227	0.77	ADC-227	86.13
ADC-235	0.81	ADC-235	86.86
ADC-243	3.61	ADC-243	89.63
SI-1X4	>500	SI-1X4	31.99
SI-1X6.4	6.98	SI-1X6.4	62.18
SI-1X22	4.47	SI-1X22	69.01
SI-1X24	6.23	SI-1X24	69.98
SI-1X25	5.54	SI-1X25	75.89
SI-1X26	>500	SI-1X26	44.63

Table 6: *In vitro* efficacy of six naked antibodies (SI-1X4, SI-1×6.4, SI-1X22, SI-1X24, SI-1X25, SI-1X26) and six ADCs (ADC-112, ADC-6, ADC-219, ADC-227, ADC-235, ADC-243) in BXP-3

BxPC-3			
Groups	IC ₅₀ (nM)	Groups	Efficacy (%)
ADC-112	46.75	ADC-112	61.96
ADC-6	12.29	ADC-6	61.02

ADC-219	16.58	ADC-219	65.64
ADC-227	32.48	ADC-227	68.85
ADC-235	15.69	ADC-235	68.24
ADC-243	16.69	ADC-243	65.96
SI-1X4	>500	SI-1X4	N/A
SI-1X6.4	>500	SI-1X6.4	N/A
SI-1X22	>500	SI-1X22	N/A
SI-1X24	>500	SI-1X24	N/A
SI-1X25	>500	SI-1X25	N/A
SI-1X26	>500	SI-1X26	N/A

Table 7: *In vitro* efficacy of six naked antibodies (SI-1X4, SI-1x6.4, SI-1X22, SI-1X24, SI-1X25, SI-1X26) and six ADCs (ADC-112, ADC-6, ADC-219, ADC-227, ADC-235, ADC-243) in FaDu

FaDu			
Groups	IC ₅₀ (nM)	Groups	Efficacy (%)
ADC-112	1.01	ADC-112	74.09
ADC-6	0.18	ADC-6	74.31
ADC-219	0.21	ADC-219	74.68
ADC-227	0.25	ADC-227	72.44
ADC-235	0.28	ADC-235	74.09
ADC-243	0.40	ADC-243	72.63
SI-1X4	>500	SI-1X4	23.34
SI-1X6.4	>500	SI-1X6.4	38.92
SI-1X22	>500	SI-1X22	39.45
SI-1X24	>500	SI-1X24	36.18
SI-1X25	>500	SI-1X25	41.65
SI-1X26	>500	SI-1X26	34.43

5

Table 8: *In vitro* efficacy of six naked antibodies (SI-1X4, SI-1x6.4, SI-1X22, SI-1X24, SI-1X25, SI-1X26) as well as six ADCs (ADC-112, ADC-6, ADC-219, ADC-227, ADC-235, ADC-243) in HARA-

B.

HARA - B			
Groups	IC ₅₀ (nM)	Groups	Efficacy (%)
ADC-112	14.73	ADC-112	60.67
ADC-6	3.89	ADC-6	73.43
ADC-219	2.37	ADC-219	73.4
ADC-227	2.45	ADC-227	71.07
ADC-235	3.18	ADC-235	73
ADC-243	9.93	ADC-243	64.77
SI-1X6.4	>500	SI-1X6.4	39.27
SI-1X22	>500	SI-1X22	43.36
SI-1X24	>500	SI-1X24	43.65
SI-1X25	>500	SI-1X25	44.73
SI-1X26	>500	SI-1X26	36.55
SI-1X4	>500	SI-1X4	44.32

Table 9: *In vitro* efficacy of six naked antibodies (SI-1X4, SI-1x6.4, SI-1X22, SI-1X24, SI-1X25, SI-1X26) and six ADCs (ADC-112, ADC-6, ADC-219, ADC-227, ADC-235, ADC-243) in HCC827.

HCC827			
Groups	IC ₅₀ (nM)	Groups	Efficacy (%)
ADC-112	2.35	ADC-112	73.10
ADC-6	1.06	ADC-6	71.31
ADC-219	0.65	ADC-219	71.21
ADC-227	0.51	ADC-227	66.61
ADC-235	0.43	ADC-235	68.22
ADC-243	2.11	ADC-243	69.56
SI-1X4	>500	SI-1X4	18.52
SI-1X6.4	2.08	SI-1X6.4	60.02

SI-1X22	1.42	SI-1X22	63.69
SI-1X24	1.48	SI-1X24	66.70
SI-1X25	1.32	SI-1X25	69.43
SI-1X26	>500	SI-1X26	28.44

Table 10: *In vitro* efficacy of six naked antibodies (SI-1X4, SI-1X6.4, SI-1X22, SI-1X24, SI-1X25, SI-1X26) and six ADCs (ADC-112, ADC-6, ADC-219, ADC-227, ADC-235, ADC-243) in SW620.

SW620			
Groups	IC ₅₀ (nM)	Groups	Efficacy (%)
ADC-112	5.82	ADC-112	70.72
ADC-6	4.51	ADC-6	75.35
ADC-219	4.20	ADC-219	75.56
ADC-227	5.78	ADC-227	70.76
ADC-235	4.69	ADC-235	73.94
ADC-243	3.91	ADC-243	69.68
SI-1X4	>500	SI-1X4	4.02
SI-1X6.4	>500	SI-1X6.4	0.24
SI-1X22	>500	SI-1X22	3.45
SI-1X24	>500	SI-1X24	5.61
SI-1X25	>500	SI-1X25	5.92
SI-1X26	>500	SI-1X26	6.47

5

CONCLUSION: In human epidermal cancer cells A431, human *in situ* pancreatic adenocarcinoma cells BXPc-3, human pharyngeal squamous cell carcinoma cells FaDu, human lung cancer squamous cell carcinoma cell line HARA-B, human non-small-cell lung cancer cells HCC827, and human colon cancer cells SW620, six ADCs (ADC-112, ADC-6, ADC-219, ADC-227, ADC-235, ADC-243) all

10

showed more sensitive efficacy and stronger tumor cell growth inhibition relative to their respective corresponding naked antibodies (SI-1X4, SI-1X6.4, SI-1X22, SI-1X24, SI-1X25, SI-1X26) (Figure 4A - Figure 4F and Table 5 - Table 10).

Example 319

In vivo efficacy testing:

In the present invention, BALB/c nude mice were subcutaneously inoculated with a variety of human tumor cell lines (A431, SW620, A431+SW620) as experimental models to evaluate the *in vivo* efficacy of ADC-coupled drugs. A certain number of tumor cell lines were inoculated subcutaneously in BALB/c nude mice, and when the tumor volume grew to 150-300 mm³, the antibodies and corresponding ADC drugs were injected into the tail vein; the drugs were administered once a week four times, with continuous observation, and tumor measurements were performed twice a week to evaluate the inhibitory effect of the test antibodies and ADCs on the tumor cell lines.

CONCLUSION:

In the A431 single-tumor model with high expression of EGFR, 10 mg/kg of ADC-6 (DAR=8) exhibited stronger tumor inhibition relative to SI-1X6.4 naked antibody (Figure 5A).

In the SW620 single-tumor model with low EGFR expression, 10 and 15 mg/kg of ADC-6 also exhibited stronger tumor suppression relative to 15 mg/kg of SI-1X6.4 naked antibody; and was stronger than the tumor suppression of Cet-ADC (ADC-214, DAR=8) at a dose corresponding to the molecular molar concentration (Figure 5B).

In A431+SW620 heterogeneous tumors, 5, 15, and 30 mg/kg of ADC-6 similarly demonstrated stronger tumor inhibition relative to 30 mg/kg of SI-1X6.4 naked antibody (Figure 5C).

Example 320

In vivo efficacy testing:

BALB/c Nude mice subcutaneously inoculated with human epidermal cancer cells A431 and human pancreatic cancer cells

BxPC3 were utilized as experimental models in the present invention to evaluate the *in vivo* efficacy of ADC-coupled drugs. A certain number of tumor cell lines were inoculated subcutaneously on the right shoulder side of female BALB/c Nude mice, and when the average tumor volume grew to 180-250 mm³, the corresponding ADC drugs were injected into the tail vein once a week for four consecutive administrations under continuous observation, and the tumor volume was measured twice a week to evaluate the inhibitory effect of the test ADC drugs on the growth of the tumors.

$$\% \text{ change to } D_0 = (D_n - D_0) / D_0 * 100;$$

$$\% \text{ TGI} = 1 - [\text{changes of tumor volume in treatment group} / \text{changes of tumor volume in control group}] \times 100.$$

In the EGFR high-expressing A431 single-body tumor model, 42 days after the first administration, each test ADC drug caused significant tumor growth inhibition ($P < 0.05$) in the BALB/c-Nude mouse subcutaneous graft tumor model of human epidermal cancer cells A431, wherein ADC-6, ADC-219, ADC-235, ADC-227 and ADC-112 had comparable tumor-suppressing effects, all of them showing strong tumor-suppressing effects (Figure 6A).

In the BxPC3 single-body tumor model with medium expression of EGFR, 35 days after the first administration, each test ADC drug caused significant tumor growth inhibition ($P < 0.05$) in the human epidermal cancer cell BxPC3 BALB/c-Nude mouse subcutaneous graft tumor model, wherein ADC-6, ADC-219, ADC-235, ADC-227 and ADC-112 had comparable tumor-suppressing effects, all showing strong tumor-suppressing effects (Figure 6B).

Example 321

In vitro efficacy of Payload compound **5A**:

1) Experimental materials:

Cells: Cells for testing were obtained from the Cell Bank of the Chinese Academy of Sciences;

Cell culture medium DMEM: Gibco;

FBS: BIOWEST.

2) Preparation of culture medium:

Growth medium (with 10% FBS, Penicillin/streptomycin (100 U/mL);

5 Detection medium (with 1% FBS, Penicillin/streptomycin (100 U/mL).

3) Operation:

Turn on the UV light of the biosafety cabinet 30 min in advance, and then turn on the ventilation for 3 min. Put the
10 growth medium, detection medium, D-PBS and pancreatin into a 37°C constant-temperature water bath to preheat, and then sterilize the surface with alcohol, and put into the biosafety cabinet. Place cells with about 80% confluence in the biosafety cabinet, draw off the old medium, rinse with D-PBS, draw and
15 discard, digest with pancreatin for 2-3 min, and then the growth medium for neutralization, and centrifuge at 1200 rpm for 3 min. Draw off the centrifugation supernatant, mix evenly with 4 mL of detection medium, and collect 100 uL for counting (wherein 50 uL of cell fluid is taken out, 50 uL of Trypan Blue Stain is
20 added and mixed evenly, followed by counting). Plate spreading was performed according to a pre-optimized cell spreading density; 80ul/well was spread in a 96-well plate, with only 80 uL of detection medium being added to wells E11 and F11, and 150 uL of DPBS being added to the edge wells. 24 h after
25 spreading the plate, diluted antibody was added, 20 uL per well, and a control was set up; only 20 uL of detection medium was added to the 11th column, and 2 duplicate wells were set up for each concentration; after addition, uniform mixing was performed on a cell vortex mixer, at 550 rpm for 3 min.

30 Dilution of solution: Use detection medium to prepare 300 uL of a test sample solution with a starting concentration of 5 uM in the first column of a V-shaped 96-well plate, add 240 uL of detection medium to the second to 10th columns respectively, take 60 uL from the evenly mixed first column and
35 add it to the second column, mix uniformly up and down with a

multi-channel pipette 10 times, discard the pipette tips, and perform operations for the next 7 concentrations in turn.

4) Detection:

After 4 days, take out the MTS reagent, thaw at room temperature in the dark, then mix thoroughly and evenly using a vortex mixer, then add 20 μ L of CellTiter One Solution Reagen MTS reagent per 100 μ L of cell culture volume along the side wall of the wells in the biosafety cabinet, gently tap the plate surface to mix the MTS solution evenly, and then put into a cell culture incubator and leave to incubate for 2 h in the dark. At the end of the reaction, take out the 96-well plate and measure the OD490 nm absorbance value in the microplate reader, and record, organize, analyze and store the data.

5) Results:

Table 11 illustrates that compound **5A** (Payload) showed good inhibition of the following solid tumor cells and hematoma cells. Table 11: *In vitro* inhibitory effect of compound **5A** on human non-small cell lung adenocarcinoma cells H1975, human non-small cell lung cancer cells HCC827, human epidermal cancer cells A431, human gastric cancer cells NCI-N87, human *in situ* pancreatic adenocarcinoma cells BxPC-3, human epidermal cancer cells A431 + human colon cancer cells SW620, human breast cancer cells ZR-75-1, human plasma cell leukemia cells H929, human multiple myeloma cells RPMI8226, human leukemia cells JJN-3, human breast cancer cells MDA-MB-361 and human breast cancer cells SK-BR-3.

Group	IC ₅₀ (nM)	Efficacy%
H1975	970.33	76.78
HCC827	655.60	80.57
A431	438.77	84.97
NCI-N87	1023.92	63.93
BxPC-3	320.09	64.4
A431+SW620	604.87	68.26
ZR-75-1	557.95	54.79
H929	23.94	99.03

RPMI8226	147.00	93.61
JJN-3	36.11	80.59
MDA-MB-361	572.13	62.34
SK-BR-3	732.43	70.9

Example 322: *In vitro* efficacy data for **ADC-6** and **ADC-214**:

The *in vitro* efficacy of ADC-coupled drugs was evaluated using an experimental model of two human tumor cell lines (human
5 poorly differentiated lung cancer squamous cell carcinoma cell line Oka-c-1, human lung squamous cell carcinoma cells SK-MES-1).

Inoculate a certain number of tumor cells in a 96-well plate, add the gradient-diluted test antibody and the corresponding
10 ADC drug to the cells, treat for 5 days, measure the cell viability using Alamar Blue or MTS, and evaluate the inhibitory effect of the test antibody, the ADC and the small molecule drug d₃ on the tumor cell lines by calculating the IC₅₀. The starting concentration of the antibody drug was 500 nM, and the
15 dilution was 7-fold, totaling 7 concentration points, and the treatment was carried out for 5 days. The final calculation method was survival rate = (experimental group - blank) / (control group - blank group) × 100%, followed by curve fitting using Graph Pad Prism, and calculation of the half inhibitory
20 concentration (IC₅₀). The results are shown in Figures 7A and 7B and Table 12.

Table 12: *In vitro* efficacy of SI-1×6.4, Cetuximab, ADC-6, ADC-214 and d₃ on human poorly differentiated lung cancer
25 squamous cell carcinoma cell line Oka-c-1 and human lung squamous cell carcinoma cells SK-MES-1.

Group	IC ₅₀ (nM)
-------	-----------------------

	SI-1×6.4	Cetuximab	ADC-6	ADC-214	d ₃
Oka-c-1	0.033	0.015	0.013	0.017	0.846
SK-MES-1	0.227	0.289	0.097	0.542	2.044

Claims


5



wherein:

10


L_1 is selected without limitation from:



15




L₂ has the structure shown in formula A below:



formula A

wherein Y is a scaffold selected from C1-C6 alkyl, substituted C1-C6 alkyl, or C3-C8 cycloalkyl; preferably Y is C1-C6 alkyl; Ac is a hydrophilic structural unit; and the carbon No. 2 attached to Y has absolute chirality in the R configuration or the S configuration;

L₃ is present or absent, and when present, L₃ is selected



from PEG hydrophilic units: , o is selected

from integers 1-10, preferably integers 2-8;

L₄ is an enzymatic cutting unit;

L₅ is a linking unit;

the chiral carbon atom No. 1 attached to N in formula I has absolute chirality in the R configuration or the S configuration;

R is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, and substituted 5-10-membered heteroaryl;

preferably R is selected from a hydrogen atom or a C1-C6 alkyl group;

R₁ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7 membered heterocyclyl, substituted 3-7 membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10 membered heteroaryl, and substituted 5-10 membered heteroaryl;

preferably R₁ is selected from a hydrogen atom or a C1-C6 alkyl group;

more preferably R₁ is selected from C1-C6 alkyl;

R₂ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl,

C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7 membered heterocyclyl, substituted 3-7 membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10 membered heteroaryl, and substituted 5-10 membered heteroaryl;

preferably R_2 is selected from a hydrogen atom, a halogen or a C1-C6 alkyl group;

more preferably R_2 is selected from halogen;

X is selected from $-C(O)-CR_aR_b-(CR_3R_4)_m-O-$, $-C(O)-CR_aR_b-(CR_3R_4)_m-NH-$ or $-C(O)-CR_aR_b-(CR_3R_4)_m-S-$;

preferably X is selected from $-C(O)-CR_aR_b-(CR_3R_4)_m-O-$;

R_a and R_b are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl, a deuterated C1-C6 alkyl, a halogenated C1-C6 alkyl, a C3-C8 cycloalkyl, a C3-C8 cycloalkyl C1-C6 alkyl, a C6-C10 aryl C1-C6 alkyl, a C1-C6 alkoxy C1-C6 alkyl, a 3-7 membered heterocyclic group, a substituted 3-7 membered heterocyclic group, a C6-C10 aryl, substituted C6-C10 aryl, 5-10 membered heteroaryl, and substituted 5-10 membered heteroaryl;

preferably, R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl, a halo-C1-C6 alkyl, a C3-C8 cycloalkyl C1-C6 alkyl, or a C6-C10 aryl C1-C6 alkyl;

alternatively, R_a , R_b and the carbon atoms to which they are connected form a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclic group, or a substituted 3-7-membered heterocyclic group; preferably R_a , R_b and the carbon atoms to which they are connected form a C3-C8 cycloalkyl group;

R_3 , R_4 are identical or different and are each independently a hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, halogenated C1-C6 alkyl, deuterated C1-C6 alkyl, C1-C6 alkoxy, hydroxyl, amino, cyano, nitro, hydroxy C1-C6 alkyl, C3-C8 cycloalkyl, 3-7-membered heterocyclyl, or substituted 3-7-membered heterocyclyl;

preferably, R_3 , R_4 are each independently a hydrogen atom or C1-C6 alkyl group;

alternatively, R₃, R₄ and the carbon atoms connected thereto form a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclic group, or a substituted 3-7-membered heterocyclic group;

5 m is selected from integers 0-4, preferably 0, 1; n is selected from integers 1-10.

2. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in claim 1, or a pharmaceutically acceptable salt or solvate thereof, characterized in that Ab is
10 a bispecific antibody or antigen-binding fragment thereof which simultaneously targets two different epitopes or targets, preferably a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3.

3. The ligand-camptothecin derivative conjugate as shown in
15 general formula I as claimed in claim 1 or 2, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the Ab antibody comprises: an IgG1 heavy chain, a κ light chain, and a single-chain Fv (scFv) structural domain; wherein the single-chain Fv (scFv) structural domain
20 forms a construct with the IgG1 heavy chain or the κ light chain; wherein the IgG1 heavy chain and the κ light chain form an IgG portion with binding specificity for EGFR; the scFv structural domain has binding specificity for HER3, and the scFv structural domain is linked via a linker (e.g., having an amino acid
25 sequence of (gly-gly-gly-gly-ser)_n, wherein n is an integer of at least 1, and preferably n is an integer of 1 to 10) to a C-terminus or N-terminus of the IgG1 heavy chain or a C-terminus or N-terminus of the κ light chain; and wherein the single-chain
30 Fv structural domain has a structural order of N-terminus-heavy chain variable region-joint-light chain variable region-C-terminus, or N-terminus-light chain variable region-joint-heavy chain variable region-C-terminus (e.g., the joint consists of
an amino acid sequence of (gly-gly-gly-gly-ser)_m, wherein m is an integer of at least 3, and preferably m is 3, 4, 5, or 6).

35 4. The ligand-camptothecin derivative conjugate as shown in

general formula I as claimed in any one of claims 1 - 3, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the κ light chain of the Ab antibody comprises CDRs as shown in SEQ ID NO: 25, SEQ ID NO: 26, and
5 SEQ ID NO: 27, the IgG1 heavy chain comprises CDRs as shown in SEQ ID NO: 29, SEQ ID NO: 30, and SEQ ID NO: 31, and the single-chain Fv (scFv) structural domain comprises heavy chain variable region CDRs as shown in SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO: 34, and light chain variable region CDRs as shown in SEQ
10 ID NO: 35, SEQ ID NO: 36, and SEQ ID NO: 37.

5. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1 - 4, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain of the Ab antibody
15 comprises a variable region as shown in SEQ ID NO: 28, the IgG1 heavy chain comprises a variable region as shown in SEQ ID NO: 38, and the single-chain Fv (scFv) structural domain comprises a heavy chain variable region as shown in SEQ ID NO: 39 and a light chain variable region as shown in SEQ ID NO: 40.

20 6. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1 - 5, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain amino acid sequence of the Ab antibody is SEQ ID NO: 2, and the amino acid sequence of
25 the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 4.

7. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1 - 6, or a pharmaceutically acceptable salt or solvate thereof,
30 characterized in that the light chain nucleic acid coding sequence of the Ab antibody is SEQ ID NO: 1, and the nucleic acid coding sequence of the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 3.

35 8. The ligand-camptothecin derivative conjugate as shown in

general formula I as claimed in any one of claims 1 - 3, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the κ light chain of the Ab antibody comprises CDRs as shown in SEQ ID NO: 41, SEQ ID NO: 42, and
5 SEQ ID NO: 43, the IgG1 heavy chain comprises CDRs as shown in SEQ ID NO: 45, SEQ ID NO: 46, and SEQ ID NO: 47, and the single-chain Fv (scFv) structural domain comprises heavy chain variable region CDRs as shown in SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO: 34, and light chain variable region CDRs as shown in SEQ
10 ID NO: 35, SEQ ID NO: 36, and SEQ ID NO: 37.

9. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1 - 3 and 8, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain of the Ab antibody
15 comprises a variable region as shown in SEQ ID NO: 44, the IgG1 heavy chain comprises a variable region as shown in SEQ ID NO: 48, and the single-chain Fv (scFv) structural domain comprises a heavy chain variable region as shown in SEQ ID NO: 39 and a light chain variable region as shown in SEQ ID NO: 40.

20 10. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1 - 3 and 8 - 9, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain amino acid sequence of the Ab antibody is SEQ ID NO: 6, and the amino acid sequence of
25 the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 8.

11. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1 - 3 and 8 - 10, or a pharmaceutically acceptable salt or solvate thereof,
30 characterized in that the light chain nucleic acid coding sequence of the Ab antibody is SEQ ID NO: 5, and the nucleic acid coding sequence of the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 7.

35 12. The ligand-camptothecin derivative conjugate as shown in

general formula I as claimed in any one of claims 1-3 and 4, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain of the Ab antibody comprises a variable region as shown in SEQ ID NO: 49, the IgG1 heavy chain comprises a variable region as shown in SEQ ID NO: 52, and the single chain Fv (scFv) structural domain comprises a heavy chain variable region as shown in SEQ ID NO: 50 and a light chain variable region as shown in SEQ ID NO: 51.

13. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-3, 4 and 12, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the heavy chain amino acid sequence of the Ab antibody is SEQ ID NO: 12, and the amino acid sequence of the construct of the antibody light chain and the single chain Fv (scFv) structural domain is SEQ ID NO: 10.

14. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-3, 4 and 12-13, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the heavy chain nucleic acid coding sequence of the Ab antibody is SEQ ID NO: 11, and the nucleic acid coding sequence of the construct of the antibody light chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 9.

15. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-3, 4 and 5, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain amino acid sequence of the Ab antibody is SEQ ID NO: 14, and the amino acid sequence of the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 16.

16. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-3, 4, 5 and 15, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain nucleic acid coding sequence of the Ab antibody is SEQ ID NO: 13, and the nucleic

acid coding sequence of the construct of the antibody heavy chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 15.

17. The ligand-camptothecin derivative conjugate as shown in
5 general formula I as claimed in any one of claims 1-3, 4 and 12, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the heavy chain amino acid sequence of the Ab antibody is SEQ ID NO: 20, and the amino acid sequence of the construct of the antibody light chain and the single-
10 chain Fv (scFv) structural domain is SEQ ID NO: 18.

18. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-3, 4, 12 and 17, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the heavy chain nucleic acid coding
15 sequence of the Ab antibody is SEQ ID NO: 19, and the nucleic acid coding sequence of the construct of the antibody light chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 17.

19. The ligand-camptothecin derivative conjugate as shown in
20 general formula I as claimed in any one of claims 1-3 and 8, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the light chain of the Ab antibody comprises a variable region as shown in SEQ ID NO: 53, the IgG1 heavy chain comprises a variable region as shown in SEQ ID NO:
25 54, and the single-chain Fv (scFv) structural domain comprises a heavy chain variable region as shown in SEQ ID NO: 50 and a light chain variable region as shown in SEQ ID NO: 51.


20. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-3, 8 and 19,
30 or a pharmaceutically acceptable salt or solvate thereof, characterized in that the heavy chain amino acid sequence of the Ab antibody is SEQ ID NO: 24, and the amino acid sequence of the construct of the antibody light chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 22.

35 21. The ligand-camptothecin derivative conjugate as shown in

general formula I as claimed in any one of claims 1-3, 8 and 19-20, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the heavy chain nucleic acid coding sequence of the Ab antibody is SEQ ID NO: 23, and the nucleic acid coding sequence of the construct of the antibody light chain and the single-chain Fv (scFv) structural domain is SEQ ID NO: 21.

22. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-21, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the Ab antibody comprises: two IgG1 heavy chains; two κ light chains; and two single-chain Fv (scFv) structural domains.

23. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-22, or a pharmaceutically acceptable salt or solvate thereof, characterized in that said X is selected without limitation from the following structures or isomers thereof:



where the left wavy line is connected to a camptothecin derivative portion, and the right wavy line is connected to L_5 .

24. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-23, or a pharmaceutically acceptable salt or solvate thereof, characterized in that L₄ is selected without limitation from peptide residues formed of amino acids,

wherein optionally, the amino acid is further substituted with one or more substituents selected from deuterium atoms, halogens, hydroxyl, cyano, amino, nitro, carboxyl, C1-C6 alkyl, substituted C1-C6 alkyl, C1-C6 alkoxy and C3-C8 cycloalkyl or

substituted C3-C8 cycloalkyl;

preferably, the peptide residue is a peptide residue formed from one, two or more amino acids selected from phenylalanine (F), glycine (G), valine (V), lysine (K), citrulline (C), serine (S), glutamic acid (E) or aspartic acid (D);

more preferably, the peptide residue is a tetrapeptide residue consisting of glycine (G) - glycine (G) - phenylalanine (F) - glycine (G).

25. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-24, or a pharmaceutically acceptable salt or solvate thereof, characterized in that:


L₅ is selected without limitation from -NR₅(CR₆R₇)_q- or a chemical bond, and q is selected from integers 0-6;


R₅, R₆ and R₇ are identical or different, and are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl group, a substituted C1-C6 alkyl group, a deuterated C1-C6 alkyl group, a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a C1-C6 alkoxy C1-C6 alkyl group, a 3-7 membered heterocyclic group, a substituted 3-7 membered heterocyclic group, a C6-C10 aryl group, a substituted C6-C10 aryl group, a 5-10 membered heteroaryl group, and a substituted 5-10 membered heteroaryl group;

preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom or a C1-C6 alkyl group;


more preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom.

26. The ligand-camptothecin derivative conjugate as shown in general formula I as claimed in any one of claims 1-25, or a pharmaceutically acceptable salt or solvate thereof, characterized in that the linking unit $-L_1-L_2-L_3-L_4-L_5-$ is selected without limitation from the following structures;





preferably:



wherein:

Ac is a hydrophilic structural unit;

R₅, R₆ and R₇ are identical or different, and are each independently selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6

alkyl, C3-C8 cycloalkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, 3-7-membered heterocyclyl, substituted 3-7-membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, and substituted 5-10-membered heteroaryl;

preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom or a C1-C6 alkyl group;


more preferably, R₅, R₆ and R₇ are each independently selected from a hydrogen atom;

the carbon atom No. 2 connected to N has absolute chirality in the R configuration or the S configuration;

the left wavy line is linked to the antibody or its antigen-binding fragment portion, and the right wavy line is linked to the X;

o is selected from integers 1-10.

27. A ligand-camptothecin derivative conjugate as shown in general formula II, or a pharmaceutically acceptable salt or solvate thereof;





II

wherein:

Ab is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3;

L₁ is a linking unit connected to Ab, and is selected without limitation from:






L₁ is preferably ;

preferably L₁ is or ;

L₃ is present or absent, and when L₃ is present, L₃ is



5 selected from , o is selected from integers 1-10, preferably integers 2-8;

Ac is a hydrophilic structural unit;

the 1-position, 2-position and 3-position chiral carbon atoms have two chiral configurations, the R-absolute configuration or the S-absolute configuration;

R is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, and substituted 5-10-membered heteroaryl;

preferably R is selected from a hydrogen atom or a C1-C6 alkyl group;

R₁ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7 membered heterocyclyl, substituted 3-7 membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10 membered heteroaryl, and substituted 5-10 membered heteroaryl;

preferably R₁ is selected from a hydrogen atom or a C1-C6 alkyl group;

more preferably R₁ is selected from C1-C6 alkyl;

R₂ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl,

C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7 membered heterocyclyl, substituted 3-7 membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10 membered heteroaryl, and substituted 5-10 membered heteroaryl;

preferably R_2 is selected from a hydrogen atom, a halogen or a C1-C6 alkyl group;

more preferably R_2 is selected from halogen;

X selected from $-C(O)-CR_aR_b-(CR_3R_4)_m-O-$, $-C(O)-CR_aR_b-(CR_3R_4)_m-NH-$ or $-C(O)-CR_aR_b-(CR_3R_4)_m-S-$;

preferably X is selected from $-C(O)-CR_aR_b-(CR_3R_4)_m-O-$;

R_a and R_b are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl, a deuterated C1-C6 alkyl, a halogenated C1-C6 alkyl, a C3-C8 cycloalkyl, a C3-C8 cycloalkyl C1-C6 alkyl, a C1-C6 alkoxy C1-C6 alkyl, a 3-7-membered heterocyclic group, a substituted 3-7-membered heterocyclic group, a C6-C10 aryl, a substituted C6-C10 aryl, a 5-10-membered heteroaryl, and a substituted 5-10-membered heteroaryl;

preferably, R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl, a halo-C1-C6 alkyl, a C3-C8 cycloalkyl C1-C6 alkyl or a C6-C10 aryl C1-C6 alkyl;

alternatively, R_a , R_b and the carbon atoms to which they are attached form a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclic group, or a substituted 3-7-membered heterocyclic group; preferably R_a , R_b and the carbon atoms to which they are attached form a C3-C8 cycloalkyl group;

R_3 and R_4 are identical or different, and are each independently a hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, halogenated C1-C6 alkyl, deuterated C1-C6 alkyl, C1-C6 alkoxy, hydroxyl, amino, cyano, nitro, hydroxy C1-C6 alkyl, C3-C8 cycloalkyl, 3-7-membered heterocyclyl, or substituted 3-7-membered heterocyclyl;


preferably, R_3 and R_4 are each independently a hydrogen atom or C1-C6 alkyl group;

alternatively, R₃, R₄ and the carbon atoms attached thereto form a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclic group, or a substituted 3-7-membered heterocyclic group;

m is selected from the integers 0-4, preferably 0, 1;

n is selected from the integers 1-10.

28. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-27, characterized in that: the Ac has the structure shown in formula B below,



wherein:


Z is selected without limitation from the group consisting of one or more of a hydrophilic structural carboxyl group, phosphoric acid, polyphosphoric acid, phosphorous acid, sulfonic acid, sulfinic acid, or polyethylene glycol (PEG);

preferably Z is selected from a hydrophilic structural carboxyl group, phosphoric acid or polyethylene glycol (PEG);

Y' is an optional scaffold connecting the amino group to Z; preferably Y' is C1-C6 alkyl;


Ac is linked to the 2-position carbon marked in structural formula I via scaffold Y.

29. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-28, characterized in that: the Ac is selected without limitation from glycine, (D/L) alanine, (D/L) leucine, (D/L) isoleucine, (D/L) valine, (D/L) phenylalanine, (D/L) proline, (D/L) tryptophan, (D/L) serine, (D/L) tyrosine, (D/L) cysteine, (D/L) cystine, (D/L) arginine, (D/L) histidine, (D/L) methionine, (D/L) asparagine, (D/L) glutamine, (D/L) threonine, (D/L) aspartic acid, (D/L) glutamic acid, natural or unnatural amino acid derivatives or the following structures or isomers thereof,




30. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-29, characterized in that: the Ac is selected without limitation from a glycine, phosphoric acid, (D/L) glutamic acid or polyethylene glycol hydrophilic structure.

31. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-30, characterized in that the



has the structure shown in formula d below;



d

wherein:

R is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, and substituted 5-10-membered heteroaryl;

preferably R is selected from a hydrogen atom or a C1-C6 alkyl group;

R₁ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7 membered heterocyclyl, substituted 3-7 membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10 membered heteroaryl, and substituted 5-10 membered heteroaryl;

preferably, R₁ is selected from a hydrogen atom or a C1-C6 alkyl group;

more preferably, R₁ is selected from C1-C6 alkyl;

R₂ is selected from hydrogen atom, deuterium atom, halogen, C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, carboxyl, 3-7 membered heterocyclyl, substituted 3-7 membered heterocyclyl, C6-C10 aryl, substituted C6-C10 aryl, 5-10 membered heteroaryl, and substituted 5-10 membered heteroaryl;

preferably, R₂ is selected from a hydrogen atom, a halogen or a C1-C6 alkyl group;

more preferably, R₂ is selected from halogen;

R_a and R_b are each independently selected from a hydrogen atom, a deuterium atom, a halogen, a C1-C6 alkyl, a deuterated C1-C6 alkyl, a halogenated C1-C6 alkyl, a C3-C8 cycloalkyl, a C3-C8 cycloalkyl C1-C6 alkyl, a C1-C6 alkoxy C1-C6 alkyl, a 3-7-membered heterocyclic group, a substituted 3-7-membered heterocyclic group, a C6-C10 aryl, a substituted C6-C10 aryl, a 5-10-membered heteroaryl, and a substituted 5-10-membered heteroaryl;

preferably, R_a and R_b are each independently selected from a hydrogen atom, a C1-C6 alkyl group, a halo-C1-C6 alkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, and a C6-C10 aryl group;


alternatively, R_a, R_b and the carbon atoms to which they are connected form a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl group, a 3-7-membered heterocyclic group, or a substituted 3-7-membered heterocyclic group; preferably R_a, R_b


and the carbon atoms to which they are linked form a C3-C8 cycloalkyl group;

the 1-position chiral carbon atom has two chiral configurations, the R absolute configuration or the S absolute configuration;


m is selected from 0 or 1.

32. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-31, characterized in that: the structural formula d is selected without limitation from the following compounds:





33. A linker-drug compound or a pharmaceutically acceptable salt or solvate thereof, characterized by having the structure shown in formula III below,



III

wherein:


R is selected from hydrogen atom, deuterium atom, halogen,
 5 C1-C6 alkyl, substituted C1-C6 alkyl, deuterated C1-C6 alkyl,
 C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy C1-C6 alkyl, C6-C10
 aryl, substituted C6-C10 aryl, 5-10-membered heteroaryl, and
 substituted 5-10-membered heteroaryl;


R_a is selected from hydrogen atom, deuterium atom, halogen,
 10 C1-C6 alkyl, deuterated C1-C6 alkyl, halogenated C1-C6 alkyl,
 C3-C8 cycloalkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy
 C1-C6 alkyl, 3-7 membered heterocyclic group, substituted 3-7
 membered heterocyclic group, C6-C10 aryl, substituted C6-C10
 aryl, 5-10 membered heteroaryl, and substituted 5-10-membered
 15 heteroaryl;

R_b is selected from hydrogen atom, deuterium atom, halogen,
 C1-C6 alkyl, deuterated C1-C6 alkyl, halogenated C1-C6 alkyl,
 C3-C8 cycloalkyl, C3-C8 cycloalkyl C1-C6 alkyl, C1-C6 alkoxy
 C1-C6 alkyl, 3-7 membered heterocyclic group, substituted 3-7
 20 membered heterocyclic group, C6-C10 aryl, substituted C6-C10
 aryl, 5-10 membered heteroaryl, and substituted 5-10-membered
 heteroaryl;

alternatively, R_a, R_b and the carbon atoms attached thereto
 form a C3-C8 cycloalkyl group, a C3-C8 cycloalkyl C1-C6 alkyl
 25 group, a 3-7-membered heterocyclic group, or a substituted 3-7-
 membered heterocyclic group;

L₃ is present or absent, and when L₃ is present, it is



selected from , o is selected from integers 1 to
 10;

the 1-position or 2-position chiral carbon atom has two chiralities, the R absolute configuration or S absolute configuration;

Ac is a hydrophilic structural unit;

m is selected from 0 or 1.

34. The linker-drug compound or pharmaceutically acceptable salt or solvate thereof as claimed in claim 33, characterized in that: the Ac has the structure shown in formula B below,



B

wherein:


Z is selected without limitation from the group consisting of one or more of a hydrophilic structural carboxyl group, phosphoric acid, polyphosphoric acid, phosphorous acid, sulfonic acid, sulfinic acid, or polyethylene glycol (PEG);

Y' is an optional scaffold linking the amino group and Z;

Ac is connected to the 2-position carbon labeled in structural formula I by means of scaffold Y;


preferably, the linker-drug compound or the pharmaceutically acceptable salt or solvate thereof is used to couple with the ligand Ab to form the ligand-camptothecin derivative conjugate of formula I or formula II as claimed in any one of claims 1-32.











35. The linker-drug compound or pharmaceutically acceptable salt or solvate thereof as claimed in claim 33 or 34, characterized in that: the Ac is selected without limitation from glycine, (D/L) alanine, (D/L) leucine, (D/L) isoleucine, (D/L) valine, (D/L) phenylalanine, (D/L) proline, (D/L) tryptophan, (D/L) serine, (D/L) tyrosine, (D/L) cysteine, (D/L) cystine, (D/L) arginine, (D/L) histidine, (D/L) methionine, (D/L) asparagine, (D/L) glutamine, (D/L) threonine, (D/L) aspartic acid, (D/L) glutamic acid, natural or unnatural amino acid derivatives or the following structures,





36. The linker-drug compound or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-35, characterized in that: Ac is selected without limitation from a glycine, phosphoric acid, (*D/L*) glutamic acid or polyethylene glycol hydrophilic structure.












37. The linker-drug compound or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-36, characterized in that the linker-drug compound is selected without limitation from the following structures or isomers thereof,















5
















5




5



5





or



5 wherein: o is selected from the integers 1-10.

38. A method for preparing a ligand-camptothecin derivative conjugate as shown in general formula I or general formula II or a pharmaceutically acceptable salt or solvate thereof, characterized by comprising the following steps,














a ligand-camptothecin derivative conjugate as shown in general formula I or general formula II is obtained by a coupling reaction of a reduced antibody or antigen-binding fragment thereof with a linker-drug compound;









the 1-position, 2-position, or 3-position chiral carbon atom has absolute chirality in the R-configuration or S-configuration;


Ab, L₁, L₂, L₃, L₄, L₅, X, R, R₁, R₂ and n are as described in any one of claims 1-37.


39. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or the method as claimed in claim 38, characterized in that: the ligand-camptothecin derivative conjugate or the pharmaceutically acceptable salt or solvate thereof is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,

















































5 wherein:

SI-1x6.4 is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3, preferably as described in any one of claims 2-7 and 22;

n is selected from the integers 1-10.


10 40. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or the method as claimed in claim 38, characterized in that: the ligand-camptothecin derivative conjugate or the pharmaceutically acceptable salt or solvate thereof is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,

15




ADC-107

/




ADC-108

/




ADC-109

/




ADC-110

/




ADC-111

/




ADC-111a

/




ADC-111b

/




ADC-112

/




ADC-112a

/




ADC-112b


/





ADC-113


/









































5 wherein:

SI-1x4 is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3, preferably as described in any one of claims 2-3, 8-11 and 22;

n is selected from the integers 1-10.

10 41. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or the method as claimed in claim 38, characterized in that: the ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof

15 is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,




wherein:

SI-1x22 is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3, preferably as described in any one of claims 2-3, 4, 12-14 and 22;

n is selected from the integers 1-10.

42. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or the method as claimed in claim 38, characterized in that: the ligand-camptothecin derivative conjugate or the pharmaceutically acceptable salt or solvate

thereof is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,




wherein:

10 SI-1x24 is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3, preferably as described in any one of claims 2-5, 15, 16 and 22;

n is selected from the integers 1-10.

43. The ligand-camptothecin derivative conjugate or
15 pharmaceutically acceptable salt or solvate thereof as claimed

in any one of claims 1-32 or the method as claimed in claim 38, characterized in that: the ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof is selected without limitation from the following structures or




wherein:

SI-1x25 is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3, preferably as described in any one of claims 2-3, 4, 12, 17, 18 and 22;

n is selected from the integers 1-10.

44. The ligand-camptothecin derivative conjugate or

pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or the method as claimed in claim 38, characterized in that: the ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof is selected without limitation from the following structures or succinimide open-ring structures thereof or isomers thereof,



SI-1x26 is a bispecific antibody or antigen-binding fragment thereof that simultaneously targets EGFR and HER3, preferably as described in any one of claims 2-3, 8, 19-21 and 22;

n is selected from the integers 1-10.

45. The ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or 39-44 or the linker-drug compound
5 or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-37, characterized in that: the pharmaceutically acceptable salt comprises a sodium salt, a potassium salt, a calcium salt or a magnesium salt formed with an acidic functional group in the structural formula, and
10 acetate, trifluoroacetate, citrate, oxalate, tartrate, malate, nitrate, chloride, bromide, iodide, sulfate, bisulfate, phosphate, lactate, oleate, ascorbate, salicylate, formate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate or p-toluenesulfonate formed with a basic functional group in
15 the structure.

46. A pharmaceutical composition, comprising the ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or 39-44 or the linker-drug compound or
20 pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-37, and optionally a pharmaceutically acceptable carrier.

47. A pharmaceutical preparation, comprising the ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or 39-44 or the linker-drug compound or
25 pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-37.

48. The use of the ligand-camptothecin derivative conjugate
30 or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or 39-44 or the linker-drug compound or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-37, or the pharmaceutical composition as claimed in claim 46 and/or the pharmaceutical
35 preparation as claimed in claim 47, in the preparation of a drug

for the treatment or prevention of cancer or tumors;

alternatively, the use of the ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or 39-44 or the linker-drug compound or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-37, or the pharmaceutical composition as claimed in claim 46 and/or the pharmaceutical preparation as claimed in claim 47, for the treatment or prevention of cancer or tumors;

preferably, the cancer or tumor expresses EGFR and/or HER3;

more preferably, the cancer or tumor is selected from adenocarcinoma, ovarian cancer, cervical cancer, uterine cancer, prostate cancer, renal cancer, urethral cancer, bladder cancer, liver cancer, gastric cancer, endometrial cancer, salivary gland cancer, esophageal cancer, lung cancer, colon cancer, rectal cancer, colorectal cancer, bone cancer, skin cancer, thyroid cancer, pancreatic cancer, melanoma, glioma, neuroblastoma, glioblastoma multiforme, sarcoma, lymphoma and leukemia, and other solid tumors or blood tumors.

49. A method for treating or preventing cancer or tumors, the method comprising administering to a subject in need thereof a prophylactically or therapeutically effective amount of the ligand-camptothecin derivative conjugate or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 1-32 or 39-44 or the linker-drug compound or pharmaceutically acceptable salt or solvate thereof as claimed in any one of claims 33-37, or the pharmaceutical composition as claimed in claim 46 and/or the pharmaceutical preparation as claimed in claim 47;

preferably, the cancer or tumor expresses EGFR and/or HER3;

more preferably, the cancer or tumor is selected from adenocarcinoma, ovarian cancer, cervical cancer, uterine cancer, prostate cancer, renal cancer, urethral cancer, bladder cancer, liver cancer, gastric cancer, endometrial cancer, salivary gland cancer, esophageal cancer, lung cancer, colon cancer, rectal

cancer, colorectal cancer, bone cancer, skin cancer, thyroid cancer, pancreatic cancer, melanoma, glioma, neuroblastoma, glioblastoma multiforme, sarcoma, lymphoma and leukemia, and other solid tumors or blood tumors.

5

10

15

20

25

30

35

40

45