
Claims

1. A compound, or pharmaceutically acceptable salt thereof, having the structure of Formula I:



Formula I

wherein the dotted lines represent zero, one, two, three, or four non-adjacent double bonds;

A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 10-membered heteroarylene;

B is -CH(R^9)- or >C=C$R^9R^{9'}$ where the carbon is bound to the carbonyl carbon of -N(R^{11})C(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

G is optionally substituted C_1-C_4 alkylene, optionally substituted C_1-C_4 alkenylene, optionally substituted C_1-C_4 heteroalkylene, -C(O)O-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, -C(O)NH-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, optionally substituted C_1-C_4 heteroalkylene, or 3 to 8-membered heteroarylene;

L is absent or a linker;

W is a cross-linking group comprising a carbodiimide, an oxazoline, a thiazoline, a chloroethyl urea, a chloroethyl thiourea, a chloroethyl carbamate, a chloroethyl thiocarbamate, an aziridine, a trifluoromethyl ketone, a boronic acid, a boronic ester, an *N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), an iso-EEDQ or other EEDQ derivative, an epoxide, an oxazolium, or a glycal;

X^1 is optionally substituted C_1-C_2 alkylene, NR, O, or S(O)$_n$;

X^2 is O or NH;

X^3 is N or CH;

n is 0, 1, or 2;

R is hydrogen, cyano, optionally substituted C_1-C_4 alkyl, optionally substituted C_2-C_4 alkenyl, optionally substituted C_2-C_4 alkynyl, C(O)R', C(O)OR', C(O)N(R')$_2$, S(O)R', S(O)$_2$R', or S(O)$_2$N(R')$_2$;

each R' is, independently, H or optionally substituted C_1-C_4 alkyl;

Y^1 is C, CH, or N;

Y^2, Y^3, Y^4, and Y^7 are, independently, C or N;

Y^5 is CH, CH_2 or N;

Y^6 is C(O), CH, CH_2, or N;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl, or

R^1 and R^2 combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

R^2 is absent, hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 7-membered heterocycloalkyl, optionally substituted 6-membered aryl, optionally substituted 5 or 6-membered heteroaryl; R^3 is absent, or

R^2 and R^3 combine with the atom to which they are attached to form an optionally substituted 3 to 8-membered cycloalkyl or optionally substituted 3 to 14-membered heterocycloalkyl;

R^4 is absent, hydrogen, halogen, cyano, or methyl optionally substituted with 1 to 3 halogens;

R^5 is hydrogen, C_1-C_4 alkyl optionally substituted with halogen, cyano, hydroxy, or C_1-C_4 alkoxy, cyclopropyl, or cyclobutyl;

R^6 is hydrogen or methyl; R^7 is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl, or

R^6 and R^7 combine with the carbon atoms to which they are attached to form an optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^8 is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

R^7 and R^8 combine with the carbon atom to which they are attached to form $C=CR^{7'}R^{8'}$; C=N(OH), C=N(O-C_1-C_3 alkyl), C=O, C=S, C=NH, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{7a} and R^{8a} are, independently, hydrogen, halo, optionally substituted C_1-C_3 alkyl, or combine with the carbon to which they are attached to form a carbonyl;

$R^{7'}$ is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl; $R^{8'}$ is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is hydrogen, F, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl, or

R^9 and L combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

$R^{9'}$ is hydrogen or optionally substituted C_1-C_6 alkyl;


R^{10} is hydrogen, halo, hydroxy, C_1-C_3 alkoxy, or C_1-C_3 alkyl;

R^{10a} is hydrogen or halo;

R^{11} is hydrogen or C_1-C_3 alkyl; and

R^{34} is hydrogen or C_1-C_3 alkyl.

2. The compound, or pharmaceutically acceptable salt thereof, of claim 1, wherein the compound has the structure of Formula Ia:



Formula Ia

wherein the dotted lines represent zero, one, two, three, or four non-adjacent double bonds;

A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -N(R^{11})C(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

L is absent or a linker;

W is a cross-linking group comprising a carbodiimide, an oxazoline, a thiazoline, a chloroethyl urea, a chloroethyl thiourea, a chloroethyl carbamate, a chloroethyl thiocarbamate, an aziridine, a trifluoromethyl ketone, a boronic acid, a boronic ester, an *N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), an iso-EEDQ or other EEDQ derivative, an epoxide, an oxazolium, or a glycal;

X^2 is O or NH;

X^3 is N or CH;

n is 0, 1, or 2;

R is hydrogen, cyano, optionally substituted C_1-C_4 alkyl, optionally substituted C_2-C_4 alkenyl, optionally substituted C_2-C_4 alkynyl, C(O)R', C(O)OR', C(O)N(R')$_2$, S(O)R', S(O)$_2$R', or S(O)$_2$N(R')$_2$;

each R' is, independently, H or optionally substituted C_1-C_4 alkyl;

Y^1 is C, CH, or N;

Y^2, Y^3, Y^4, and Y^7 are, independently, C or N;

Y^5 and Y^6 are, independently, CH or N;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl;

R^2 is hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 7-membered heterocycloalkyl, optionally substituted 6-membered aryl, optionally substituted 5 or 6-membered heteroaryl; R^3 is absent, or

R^2 and R^3 combine with the atom to which they are attached to form an optionally substituted 3 to 8-membered cycloalkyl or optionally substituted 3 to 14-membered heterocycloalkyl;

R^4 is absent, hydrogen, halogen, cyano, or methyl optionally substituted with 1 to 3 halogens;

R^5 is hydrogen, C_1-C_4 alkyl optionally substituted with halogen, cyano, hydroxy, or C_1-C_4 alkoxy, cyclopropyl, or cyclobutyl;

R^6 is hydrogen or methyl; R^7 is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl, or

R^6 and R^7 combine with the carbon atoms to which they are attached to form an optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^8 is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

R^7 and R^8 combine with the carbon atom to which they are attached to form $C=CR^{7'}R^{8'}$; C=N(OH), C=N(O-C_1-C_3 alkyl), C=O, C=S, C=NH, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

$R^{7'}$ is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl; $R^{8'}$ is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or


$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{10} is hydrogen, hydroxy, C_1-C_3 alkoxy, or C_1-C_3 alkyl; and

R^{11} is hydrogen or C_1-C_3 alkyl.

3. The compound, or pharmaceutically acceptable salt thereof, of any one of claim 1 or 2, wherein the compound has the structure of Formula Ib:



Formula Ib

wherein the dotted lines represent zero, one, two, three, or four non-adjacent double bonds;

A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -NHC(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

L is absent or a linker;

W is a cross-linking group comprising a carbodiimide, an oxazoline, a thiazoline, a chloroethyl urea, a chloroethyl thiourea, a chloroethyl carbamate, a chloroethyl thiocarbamate, an aziridine, a trifluoromethyl ketone, a boronic acid, a boronic ester, an *N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), an iso-EEDQ or other EEDQ derivative, an epoxide, an oxazolium, or a glycal;

n is 0, 1, or 2;

R is hydrogen, cyano, optionally substituted C_1-C_4 alkyl, optionally substituted C_2-C_4 alkenyl, optionally substituted C_2-C_4 alkynyl, C(O)R', C(O)OR', C(O)N(R')$_2$, S(O)R', S(O)$_2$R', or S(O)$_2$N(R')$_2$;

each R' is, independently, H or optionally substituted C_1-C_4 alkyl;

Y^1 is C, CH, or N;

Y^2, Y^3, Y^4, and Y^7 are, independently, C or N;

Y^5 and Y^6 are, independently, CH or N;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl;

R^2 is hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 7-membered heterocycloalkyl, optionally substituted 6-membered aryl, optionally substituted 5 or 6-membered heteroaryl; R^3 is absent, or

R^2 and R^3 combine with the atom to which they are attached to form an optionally substituted 3 to 8-membered cycloalkyl or optionally substituted 3 to 14-membered heterocycloalkyl;

R^4 is absent, hydrogen, halogen, cyano, or methyl optionally substituted with 1 to 3 halogens;

R^5 is hydrogen, C_1-C_4 alkyl optionally substituted with halogen, cyano, hydroxy, or C_1-C_4 alkoxy, cyclopropyl, or cyclobutyl;

R^6 is hydrogen or methyl; R^7 is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl, or

R^6 and R^7 combine with the carbon atoms to which they are attached to form an optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^8 is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

R^7 and R^8 combine with the carbon atom to which they are attached to form $C=CR^{7'}R^{8'}$; C=N(OH), C=N(O-C_1-C_3 alkyl), C=O, C=S, C=NH, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;


$R^{7'}$ is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl; $R^{8'}$ is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl; and

R^{10} is hydrogen, hydroxy, C_1-C_3 alkoxy, or C_1-C_3 alkyl.

4. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 3, wherein the compound has the structure of Formula Ic:



Formula Ic

wherein A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -NHC(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

L is absent or a linker;

W is a cross-linking group comprising a carbodiimide, an oxazoline, a thiazoline, a chloroethyl urea, a chloroethyl thiourea, a chloroethyl carbamate, a chloroethyl thiocarbamate, an aziridine, a trifluoromethyl ketone, a boronic acid, a boronic ester, an *N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), an iso-EEDQ or other EEDQ derivative, an epoxide, an oxazolium, or a glycal;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl;

R^2 is hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 7-membered heterocycloalkyl, optionally substituted 6-membered aryl, optionally substituted 5 or 6-membered heteroaryl; R^3 is absent, or

R^2 and R^3 combine with the atom to which they are attached to form an optionally substituted 3 to 8-membered cycloalkyl or optionally substituted 3 to 14-membered heterocycloalkyl;

R^5 is hydrogen, C_1-C_4 alkyl optionally substituted with halogen, cyano, hydroxy, or C_1-C_4 alkoxy, cyclopropyl, or cyclobutyl;

R^6 is hydrogen or methyl; R^7 is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl, or

R^6 and R^7 combine with the carbon atoms to which they are attached to form an optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^8 is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

R^7 and R^8 combine with the carbon atom to which they are attached to form C=C$R^{7'}R^{8'}$; C=N(OH), C=N(O-C_1-C_3 alkyl), C=O, C=S, C=NH, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;


$R^{7'}$ is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl; $R^{8'}$ is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl; and

R^{10} is hydrogen, hydroxy, C_1-C_3 alkoxy, or C_1-C_3 alkyl.

5. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 4, wherein the compound has the structure of Formula Id:



Formula Id

wherein A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -NHC(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

L is absent or a linker;

W is a cross-linking group comprising a carbodiimide, an oxazoline, a thiazoline, a chloroethyl urea, a chloroethyl thiourea, a chloroethyl carbamate, a chloroethyl thiocarbamate, an aziridine, a trifluoromethyl ketone, a boronic acid, a boronic ester, an *N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), an iso-EEDQ or other EEDQ derivative, an epoxide, an oxazolium, or a glycal;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl;

R^2 is C_1-C_6 alkyl or 3 to 6-membered cycloalkyl;

R^7 is C_1-C_3 alkyl;


R^8 is C_1-C_3 alkyl; and

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl.

6. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 5, wherein R^1 is 5 to 10-membered heteroaryl.

7. The compound, or pharmaceutically acceptable salt thereof, of claim 6, wherein R^1 is optionally substituted 6-membered aryl or optionally substituted 6-membered heteroaryl.

8. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 7, wherein the compound has the structure of Formula Ie:



Formula Ie

wherein A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -NHC(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

L is absent or a linker;

W is a cross-linking group comprising a carbodiimide, an oxazoline, a thiazoline, a chloroethyl urea, a chloroethyl thiourea, a chloroethyl carbamate, a chloroethyl thiocarbamate, an aziridine, a trifluoromethyl ketone, a boronic acid, a boronic ester, an *N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), an iso-EEDQ or other EEDQ derivative, an epoxide, an oxazolium, or a glycal;

R^2 is C_1-C_6 alkyl or 3 to 6-membered cycloalkyl;

R^7 is C_1-C_3 alkyl;

R^8 is C_1-C_3 alkyl; and

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl

X^e and X^f are, independently, N or CH; and

R^{12} is optionally substituted C_1-C_6 alkyl or optionally substituted C_1-C_6 heteroalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl.

9. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 8, wherein A is optionally substituted 6-membered arylene.

10. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 9, wherein B is -CHR^9-.

11. The compound, or pharmaceutically acceptable salt thereof, of claim 10, wherein R^9 is optionally substituted C_1-C_6 alkyl or optionally substituted 3 to 6-membered cycloalkyl.

12. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 9, wherein B is optionally substituted 6-membered arylene.

13. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 12, wherein the linker is the structure of Formula II:


$$A^1\text{-}(B^1)_f\text{-}(C^1)_g\text{-}(B^2)_h\text{-}(D^1)\text{-}(B^3)_i\text{-}(C^2)_j\text{-}(B^4)_k\text{–}A^2$$

Formula II

where A^1 is a bond between the linker and B; A^2 is a bond between W and the linker; B^1, B^2, B^3, and B^4 each, independently, is selected from optionally substituted C_1-C_2 alkylene, optionally substituted C_1-C_3 heteroalkylene, O, S, and NR^N; R^N is hydrogen, optionally substituted C_1-C_4 alkyl, optionally substituted C_2-C_4 alkenyl, optionally substituted C_2-C_4 alkynyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted C_1-C_7 heteroalkyl; C^1 and C^2 are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; f, g, h, i, j, and k are each, independently, 0 or 1; and D^1 is optionally substituted C_1-C_{10} alkylene, optionally substituted C_2-C_{10} alkenylene, optionally substituted C_2-C_{10} alkynylene, optionally substituted 3 to 14-membered heterocycloalkylene, optionally substituted 5 to 10-membered heteroarylene, optionally substituted 3 to 8-membered cycloalkylene, optionally substituted 6 to 10-membered arylene, optionally substituted C_2-C_{10} polyethylene glycolene, or optionally substituted C_1-C_{10} heteroalkylene, or a chemical bond linking A^1-$(B^1)_f$-$(C^1)_g$-$(B^2)_h$- to -$(B^3)_i$-$(C^2)_j$-$(B^4)_k$–A^2.

14. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 13, wherein the linker is acyclic.

15. The compound, or a pharmaceutically acceptable salt thereof, of claim 14, wherein the linker has the structure of Formula IIa:



Formula IIa

wherein X^a is absent or N;


R^{14} is absent, hydrogen or optionally substituted C_1-C_6 alkyl; and

L^2 is absent, -SO_2-, optionally substituted C_1-C_4 alkylene or optionally substituted C_1-C_4 heteroalkylene,

wherein at least one of X^a, R^{14}, or L^2 is present.

16. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 13, wherein the linker is or a comprises a cyclic group.

17. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 13 or 16, wherein the linker has the structure of Formula IIb:



Formula IIb

wherein o is 0 or 1;


R^{15} is hydrogen or optionally substituted C_1-C_6 alkyl;

Cy is optionally substituted 3 to 8-membered cycloalkylene, optionally substituted 3 to 8-membered heterocycloalkylene, optionally substituted 6-10 membered arylene, or optionally substituted 5 to 10-membered heteroarylene; and

L^3 is absent, -SO_2-, optionally substituted C_1-C_4 alkylene or optionally substituted C_1-C_4 heteroalkylene.

18. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 17, wherein W comprises a carbodiimide.

19. The compound, or a pharmaceutically acceptable salt thereof, of claim 18, wherein W has the structure of Formula IIIa:




Formula IIIa

wherein R^{14} is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 6 to 10-membered aryl, optionally substituted 3 to 14-membered heterocycloalkyl, or optionally substituted 5 to 10-membered heteroaryl.

20. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 17, wherein W comprises an oxazoline or thiazoline.

21. The compound, or a pharmaceutically acceptable salt thereof, of claim 20, wherein W has the structure of Formula IIIb:



Formula IIb

wherein X^1 is O or S;


X^2 is absent or NR^{19};

R^{15}, R^{16}, R^{17}, and R^{18} are, independently, hydrogen or optionally substituted C_1-C_6 alkyl; and

R^{19} is hydrogen, C(O)(optionally substituted C_1-C_6 alkyl), optionally substituted C_1-C_6 alkyl, optionally substituted 6 to 10-membered aryl, optionally substituted 3 to 14-membered heterocycloalkyl, or optionally substituted 5 to 10-membered heteroaryl.

22. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 17, wherein W comprises a chloroethyl urea, a chloroethyl thiourea, a chloroethyl carbamate, or a chloroethyl thiocarbamate.

23. The compound, or a pharmaceutically acceptable salt thereof, of claim 22, wherein W has the structure of Formula IIIc:



Formula IIIc

wherein X^3 is O or S;


X^4 is O, S, NR^{26};

R^{21}, R^{22}, R^{23}, R^{24}, and R^{26} are, independently, hydrogen or optionally substituted C_1-C_6 alkyl; and

R^{25} is hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted 6 to 10-membered aryl, optionally substituted 3 to 14-membered heterocycloalkyl, or optionally substituted 5 to 10-membered heteroaryl.

24. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 17, wherein W comprises an aziridine.

25. The compound, or a pharmaceutically acceptable salt thereof, of claim 24, wherein W has the structure of Formula IIId1, Formula IIId2, Formula IIId3, or Formula IIId4:



Formula IIId1 Formula IIId2 Formula IIId3 Formula IIId4

wherein X^5 is absent or NR^{30};

Y is absent or C(O), C(S), S(O), SO_2, or optionally substituted C_1-C_3 alkylene;

R^{27} is hydrogen, $-C(O)R^{32}$, $-C(O)OR^{32}$, $-SO_2R^{33}$, $-SOR^{33}$, optionally substituted C_1-C_6 alkyl, optionally substituted 6 to 10-membered aryl, optionally substituted 3 to 14-membered heterocycloalkyl, or optionally substituted 5 to 10-membered heteroaryl;

R^{28} and R^{29} are, independently, hydrogen, CN, $C(O)R^{31}$, CO_2R^{31}, $C(O)R^{31}R^{31}$ optionally substituted C_1-C_6 alkyl, optionally substituted 3 to 10-membered cycloalkyl, optionally substituted 6 to 10-

membered aryl, optionally substituted 3 to 14-membered heterocycloalkyl, or optionally substituted 5 to 10-membered heteroaryl;

each R^{31} is, independently, hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted 6 to 10-membered aryl, optionally substituted 3 to 14-membered heterocycloalkyl, or optionally substituted 5 to 10-membered heteroaryl;

R^{30} is hydrogen or optionally substituted C_1-C_6 alkyl; and

R^{32} and R^{33} are, independently, hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted 6 to 10-membered aryl, optionally substituted 3 to 14-membered heterocycloalkyl, or optionally substituted 5 to 10-membered heteroaryl.

26. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 17, wherein W comprises an epoxide.

27. A compound, or a pharmaceutically acceptable salt thereof, of Table 1 or Table 2.

28. A pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 27 and a pharmaceutically acceptable excipient.

29. A conjugate, or salt thereof, comprising the structure of Formula IV:


M-L-P

Formula IV

wherein L is a linker;

P is a monovalent organic moiety; and

M has the structure of Formula V:



Formula V

wherein the dotted lines represent zero, one, two, three, or four non-adjacent double bonds;

A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is -CH(R^9)- or >C=C$R^9R^{9'}$ where the carbon is bound to the carbonyl carbon of -N(R^{11})C(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

G is optionally substituted C_1-C_4 alkylene, optionally substituted C_1-C_4 alkenylene, optionally substituted C_1-C_4 heteroalkylene, -C(O)O-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, -C(O)NH-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, optionally substituted C_1-C_4 heteroalkylene, or 3 to 8-membered heteroarylene;

X^1 is optionally substituted C_1-C_2 alkylene, NR, O, or S(O)$_n$;

X^2 is O or NH;

X^3 is N or CH;

n is 0, 1, or 2;

R is hydrogen, cyano, optionally substituted C_1-C_4 alkyl, optionally substituted C_2-C_4 alkenyl, optionally substituted C_2-C_4 alkynyl, C(O)R', C(O)OR', C(O)N(R')$_2$, S(O)R', S(O)$_2$R', or S(O)$_2$N(R')$_2$;

each R' is, independently, H or optionally substituted C_1-C_4 alkyl;

Y^1 is C, CH, or N;

Y^2, Y^3, Y^4, and Y^7 are, independently, C or N;

Y^5 is CH, CH_2, or N;

Y^6 is C(O), CH, CH_2, or N;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl, or

R^1 and R^2 combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

R^2 is absent, hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 7-membered heterocycloalkyl, optionally substituted 6-membered aryl, optionally substituted 5 or 6-membered heteroaryl; R^3 is absent, or

R^2 and R^3 combine with the atom to which they are attached to form an optionally substituted 3 to 8-membered cycloalkyl or optionally substituted 3 to 14-membered heterocycloalkyl;

R^4 is absent, hydrogen, halogen, cyano, or methyl optionally substituted with 1 to 3 halogens;

R^5 is hydrogen, C_1-C_4 alkyl optionally substituted with halogen, cyano, hydroxy, or C_1-C_4 alkoxy, cyclopropyl, or cyclobutyl;

R^6 is hydrogen or methyl; R^7 is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl, or

R^6 and R^7 combine with the carbon atoms to which they are attached to form an optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^8 is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

R^7 and R^8 combine with the carbon atom to which they are attached to form $C=CR^{7'}R^{8'}$; C=N(OH), C=N(O-C_1-C_3 alkyl), C=O, C=S, C=NH, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{7a} and R^{8a} are, independently, hydrogen, halo, optionally substituted C_1-C_3 alkyl, or combine with the carbon to which they are attached to form a carbonyl;

$R^{7'}$ is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl; $R^{8'}$ is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxyl, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is hydrogen, F, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl, or

R^9 and L combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

$R^{9'}$ is hydrogen or optionally substituted C_1-C_6 alkyl;

R^{10a} is hydrogen or halo;

R^{11} is hydrogen or C_1-C_3 alkyl; and

R^{34} is hydrogen or C_1-C_3 alkyl.

30. A compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 27 or a pharmaceutical composition of claim 28 for use as a medicament.

~~30~~31. ~~A method of treating cancer in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount~~Use of a compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 27 or a pharmaceutical composition of claim 28, in the manufacture of a medicament for the treatment of cancer.

~~31~~32. The ~~method~~ use of claim ~~30~~31, wherein the cancer is pancreatic cancer, non-small cell lung cancer, colorectal cancer or endometrial cancer.

~~32~~33. The ~~method~~ use of claim ~~30~~31 or ~~31~~32, wherein the cancer comprises a Ras mutation.

~~33~~34. The ~~method~~ use of claim ~~32~~33, wherein the Ras mutation is K-Ras G12D or K-Ras G13D.

~~34~~35. ~~A method of treating a Ras protein-related disorder in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount~~Use of a compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 27 or a pharmaceutical composition of claim 28, in the manufacture of a medicament for the treatment of a Ras protein related disorder.

~~35~~36. The ~~method or~~ use of any one of claims ~~30~~ 31 to ~~34~~35, wherein ~~the method further comprises administering~~ an additional anticancer therapy is to be administered.

~~36~~37. The ~~method~~ use of claim ~~35~~36, wherein the additional anticancer therapy is an EGFR inhibitor, a second Ras inhibitor, a SHP2 inhibitor, a SOS1 inhibitor, a Raf inhibitor, a MEK inhibitor, an ERK inhibitor, a PI3K inhibitor, a PTEN inhibitor, an AKT inhibitor, an mTORC1 inhibitor, a BRAF inhibitor, a PD-L1 inhibitor, a PD-1 inhibitor, a CDK4/6 inhibitor, a HER2 inhibitor, or a combination thereof.

~~37~~38. The ~~method~~ use of claim ~~35~~ 36 or ~~36~~37, wherein the additional anticancer therapy is a SHP2 inhibitor.