

TOPICAL FOAM COMPOSITION

FIELD OF THE INVENTION

The present invention relates to novel aerosol foam compositions
5 containing a pharmaceutically active agent for the treatment of skin disease.

BACKGROUND OF THE INVENTION

There are many challenges in topical drug delivery. The topical
composition should deliver the active agent to the site of treatment, have desirable
10 sensory characteristics, not leave an unpleasant residue on the surface of the
skin, and not cause irritation or discomfort. Aerosol foams are known to be
suitable for this purpose.

US Patent No. 7,387,807 to Callaghan et al. discloses a topical
composition comprising an extract of feverfew, pharmaceutical excipients,
15 cosmetic agents, and other biologically active substances, such as retinoids. The
composition may be in the form of an emulsion such as a cream or lotion.

US Patent No. 7,326,408 to Angel et al. discloses a composition for the
topical treatment of acne comprising one or more sunscreen agents and an
antibacterial medication, where the composition may be in the form of a gel,
20 spray, foam, lotion or other form suitable for application to the skin. The
composition may further comprise an additional anti-acne medication, such as a
retinoid (e.g. tretinoin, adapalene or tazarotene).

US Patent No. 6,730,288 to Abram describes an aerosol foam composition
including an effective amount of a pharmaceutically active ingredient, an occlusive
25 agent, an aqueous solvent, an organic cosolvent, the pharmaceutically active
ingredient being insoluble in both water and the occlusive agent, and the
occlusive agent being present in an amount sufficient to form an occlusive layer
on the skin, in use.

US published application No. 2006/0292080 to Abram et al. describes an
30 oil in water emulsion foam comprising: a vitamin or analogue thereof solubilized in
the water phase and a stabilizer solubilized in the oil phase, an emulsifier, an
occlusive agent, and an organic co-solvent, and no description of the particle size
of the oil phase.

US published application No. 2006/0057168 to Larm et al. describes a
35 process for the preparation of an oil in water microemulsion or sub-micron
emulsion composition, in particular an oil in water microemulsion or sub-micron
emulsion foam composition. These submicron or micro emulsion foams have a

comparatively high oil content and require both a hydrophilic and a hydrophobic surfactant.

The present invention is directed to low oil and low surfactant submicron emulsions or microemulsion aerosol foams containing a pharmaceutically active agent, which are cosmetically elegant, chemically and physically stable, well tolerated, easy to formulate, and suitable for application to the face.

BRIEF SUMMARY OF THE INVENTION

The present invention is directed to an oil in water emulsion aerosol foam composition having an oil phase and a water phase, said composition comprising a pharmaceutically active agent, water, oil, an oil miscible organic solvent, a surfactant, and a propellant. The present invention provides for a low oil and a low surfactant content in an oil in water emulsion aerosol foam composition.

In one embodiment, the present invention provides an oil in water emulsion aerosol foam composition comprising an oil phase and a water phase, said composition comprising:

a pharmaceutically active agent,
water,
an oil present in an amount of less than about 10% by weight,
an oil miscible organic solvent,
a surfactant component comprising a hydrophilic surfactant, in an amount from about 0.1% to about 10% by weight, and
a propellant; and wherein the pharmaceutically active agent is solubilized in the oil phase of the composition, and wherein the particle size of the oil phase is less than about 1000 nm.

According to another embodiment, the present invention provides a process for the preparation of an oil in water submicron or micro emulsion aerosol foam composition, comprising a pharmaceutically active agent, water, an oil present in an amount of less than about 10% by weight, an oil miscible organic solvent, a surfactant component comprising a hydrophilic surfactant in an amount from about 0.1% to about 10% by weight, and a propellant, the process comprising:

a) admixing the pharmaceutically active agent, a first aliquot of water, oil, oil miscible organic solvent and surfactant component to form an oil in water emulsion,

- b) heating the oil in water emulsion of step (a) to a phase inversion temperature wherein the oil in water emulsion forms a water in oil emulsion,
- 5 c) cooling the water in oil emulsion to below the phase inversion temperature to form a submicron or micro oil in water emulsion,
- d) adding a second aliquot of water to cool the submicron or micro oil in water emulsion,
- 10 e) actuating a sample of the submicron or micro oil in water emulsion with a propellant to form an oil in water submicron or micro emulsion aerosol foam.

According to an embodiment, the present invention provides a product produced by this process.

15 According to a further embodiment, the present invention provides a method of treating a skin disease, disorder or condition, comprising administering to the skin of a patient requiring such treatment an effective amount of a composition of the present invention.

20 According to yet another embodiment, the present invention relates to the use of the compositions described herein for the preparation of a medicament for the treatment of a skin disease, disorder or condition.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figure 1 is a schematic flow diagram showing one preferred embodiment of the present invention in which a low oil and low surfactant content, oil in water submicron or micro emulsion aerosol foam composition (730/2/1) is prepared. The method is described in detail in Example 1.

30 Figure 2 shows the distribution of tazarotene within the epidermis after application of tazarotene formulations of the present invention (706/8/2 and 706/8/3) compared to commercially available tazarotene cream and gel formulations (i.e. TAZORAC® cream and gel).

35 Figure 3 shows the distribution of tazarotene within the dermis after application of tazarotene formulations of the present invention (706/8/2 and 706/8/3) compared to commercially available tazarotene cream and gel formulations (i.e. TAZORAC® cream and gel).

Figure 4 shows the compatibility of high HLB ethoxylated fatty alcohol ether surfactant with tazarotene (tazarotene stability). A comparison is made against various low HLB surfactants.

5 Figure 5 shows the compatibility of high HLB ethoxylated fatty alcohol ether surfactant with tazarotene (tazarotene sulfoxide formation). A comparison is made against various low HLB surfactants.

Figure 6 shows the particle size distribution of the oil phase of the present submicron or micro emulsions (730/2/1).

10 DETAILED DESCRIPTION

The present invention is directed to an oil in water emulsion aerosol foam composition having an oil phase and a water phase, said composition comprising a pharmaceutically active agent, water, oil, an oil miscible organic solvent, a
15 surfactant, and a propellant. In an embodiment, the present invention provides low oil and low surfactant content, oil in water emulsion aerosol foam compositions. According to a further embodiment, the compositions are formulated as a submicron emulsion or microemulsion.

Thus, according to an embodiment, the present invention provides an oil in
20 water emulsion aerosol foam composition comprising an oil phase and a water phase, said composition comprising:
a pharmaceutically active agent,
water,
an oil present in an amount of less than about 10% by weight,
25 an oil miscible organic solvent,
a surfactant component comprising a hydrophilic surfactant, in an amount from about 0.1% to about 10% by weight, and
a propellant; and wherein the pharmaceutically active agent is solubilized in the oil phase of the composition, and wherein the particle size of the oil phase is
30 less than about 1000 nm.

Together, the oil and oil miscible organic solvent comprise the oil phase of the composition, along with any oil miscible excipients.

According to an embodiment, the mean particle size of the oil phase is about 100 nm.

35 In one embodiment, the present compositions are substantially free or free from a water miscible organic solvent, such as propylene glycol.

In one alternative embodiment, the pharmaceutically active agent is solubilized in the water phase of the composition.

The characteristics of oil particles in their delivery of pharmaceutically active agents across the skin barrier are summarised in Table 1 below:

5

Table 1

Particle size	Description	Characteristics
<10,000 nm	Emulsion	Blue-white, milky liquid, reasonable physical stability. Particles reside on skin surface → percutaneous delivery.
100-1000 nm	Submicron emulsion	Bluish, translucent liquid. Enhanced physical stability. Particles reside on skin surface → enhanced percutaneous delivery.
10-100 nm	Microemulsion	Translucent-transparent liquid. Excellent physical stability. Particles reside on skin surface → enhanced percutaneous delivery.
1-100 nm	Nanoemulsion	Translucent-transparent liquid. Excellent physical stability. Particles reside on skin surface, within stratum corneum and in hair follicles → optimal percutaneous delivery.

Oil Component

10

Suitably, the oil is present in the composition in an amount from about 1% to about 9% by weight. In another embodiment the oil is present in an amount from about 3% to about 8% by weight, such as about 3%, 4%, 5%, 6%, 7% or 8% by weight.

15

The oil is in the discontinuous phase of the oil in water emulsion system. In an embodiment, the oil is a hydrocarbon. Suitably, the hydrocarbon is selected from an aromatic compound, or a linear, branched or cyclic alkane or alkene, or mixtures thereof.

20

According to an embodiment, the aromatic compound is selected from the group consisting of azulene, chamazulene and cyclohexylidene-diphenylmethane, and mixtures thereof.

25

According to a further embodiment, the linear, branched or cyclic alkane or alkene is selected from the group consisting of isoparaffin, didecene, diethylhexylcyclohexane, eicosane, isododecane, isoeicosane, isohexadecane, longifolene, mineral oil, paraffin, pentahydrosqualene, petrolatum, squalane, squalene, tetradecene, derivatives thereof, and mixtures thereof.

According to an embodiment, the oil is mineral oil. In one embodiment, the mineral oil is present in an amount from about 1% to about 9% by weight. In another embodiment the mineral oil is present in about 3% to about 8% by weight, such as about 3%, 4%, 5%, 6%, 7% or 8% by weight.

5 In another embodiment, the oil is a vegetable oil. Suitably, the vegetable oil is selected from palm oil, soybean oil, rapeseed oil, sunflower oil, peanut oil, corn oil, olive oil, coconut oil, cottonseed oil, linseed oil, grapeseed oil, hazelnut oil or sesame oil, and mixtures thereof.

10 Surfactant Component

The present topical foam compositions comprise a surfactant component. Suitably, the surfactant is present in the composition in an amount from about 1% to about 8% by weight. In another embodiment the surfactant is present in an amount from about 2% to about 6% by weight, such as about 2%, 3%, 4%, 5% or
15 6% by weight.

A surfactant's hydrophilic/lipophilic balance (HLB) describes the surfactant's affinity toward water or oil. The HLB scale ranges from 1 (totally lipophilic) to 20 (totally hydrophilic), with 10 representing an equal balance of both characteristics. Lipophilic surfactants tend to form water-in-oil (w/o) emulsions,
20 and hydrophilic surfactants tend to form oil-in-water (o/w) emulsions. The HLB of a blend of two surfactants equals the weight fraction of surfactant A times its HLB value plus the weight fraction of surfactant B times its HLB value (weighted average).

In one embodiment the surfactant component comprises a hydrophilic
25 surfactant. In an another embodiment, the surfactant component consists of a single hydrophilic surfactant, and in another embodiment, the surfactant component comprises more than one surfactant and the weighted average of their HLB values is between about 10 and about 20. In another embodiment, the surfactant component consists of non-ionic surfactants.

30 Suitable non-ionic surfactants include but are not limited to ethoxylated fatty alcohol ethers, PEG derivatives, ethoxylated fatty acids, propylene glycol esters, fatty alcohols, glycerol esters and derivatives, polymeric ethers and sorbitan esters, and mixtures thereof.

Exemplary ethoxylated fatty alcohol ethers include steareth-2, steareth-10,
35 steareth-20, cetareth-2, cetareth-3, cetareth-5, cetareth-6, cetareth-10, cetareth-12, cetareth-15, cetareth-20, cetareth-21, cetareth-22, cetareth-25, cetareth-30, cetareth-31, cetareth-32, cetareth-33, laureth-3, laureth-4,

laureth-5, laureth-9, laureth-10, laureth-12, laureth-15, laureth-20, laureth-21, laureth-22, laureth-23, nonoxynol-9, oleth-2, oleth-5, oleth-10 and oleth-20.

In one embodiment the hydrophilic ethoxylated fatty alcohol ether is selected from the group consisting of steareth-10, steareth-20, cetareth-10, cetareth-12, cetareth-15, cetareth-20, cetareth-21, cetareth-22, cetareth-25, cetareth-30, cetareth-31, cetareth-32, cetareth-33, cetareth-6, laureth-5, laureth-9, laureth-10, laureth-12, laureth-15, laureth-20, laureth-21, laureth-22, laureth-23, nonoxynol-9, oleth-10 and oleth-20.

In another embodiment the wherein the hydrophilic ethoxylated fatty alcohol ether is Macrogol Cetostearyl Ether 12 (cetareth-12). In one embodiment the Macrogol Cetostearyl Ether 12 is present in the composition in an amount from about 1% to about 8% by weight. In another embodiment the Macrogol Cetostearyl Ether 12 is present in an amount from about 2% to about 6% by weight, such as about 2%, 3%, 4%, 5% or 6% by weight.

Exemplary PEG derivatives include PEG-7 hydrogenated castor oil, PEG-25 hydrogenated castor oil, PEG-30 castor oil, PEG-31 castor oil, PEG-32 castor oil, PEG-33 castor oil, PEG-34 castor oil, PEG-35 castor oil, PEG-40 hydrogenated castor oil, PEG-50 castor oil and PEG-60 hydrogenated castor oil.

Exemplary ethoxylated fatty acids include PEG-5 oleate, PEG-6 oleate, PEG-10 oleate, PEG-6 stearate, PEG-8 stearate and PEG-9 stearate, PEG-20 stearate, PEG-40 stearate, PEG-41 stearate, PEG-42 stearate, PEG-43 stearate, PEG-44 stearate, PEG-45 stearate, PEG-46 stearate, PEG-47 stearate, PEG-48 stearate, PEG-49 stearate, PEG-50 stearate and PEG-100 stearate.

Exemplary propylene glycol esters include propylene glycol palmitate and propylene glycol stearate.

Exemplary fatty alcohols include cetyl alcohol and stearyl alcohol.

Exemplary glyceryl esters and derivatives include glyceryl behenate, glyceryl dibehenate, glyceryl dioleate, glyceryl distearate, glyceryl linoleate, glyceryl oleate, glyceryl stearate, PEG-23 glyceryl cocoate, PEG-6 caprylic/capric glycerides, PEG-7 glyceryl cocoate, polyglyceryl-10 diisostearate, polyglyceryl-2 diisostearate, polyglyceryl-3 diisostearate and polyglyceryl-6 diisostearate.

Exemplary polymeric ethers include poloxamer 124, poloxamer 182, poloxamer 184, poloxamer 188, poloxamer 237, poloxamer 331, poloxamer 338 and poloxamer 407.

Exemplary sorbitan derivatives include polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80, sorbitan laurate, sorbitan oleate, sorbitan palmitate, sorbitan sesquioleate, sorbitan stearate and sorbitan trioleate.

In one embodiment, the surfactant component comprises one or more hydrophilic non-ionic surfactants and is substantially free, or free of lipophilic surfactant. According to an embodiment, the surfactant component is substantially free, or free of fatty alcohol.

5 In one embodiment, the surfactant component comprises a hydrophilic ethoxylated fatty alcohol ether. In another embodiment, the surfactant component comprises a hydrophilic ethoxylated fatty alcohol ether and is substantially free, or free of lipophilic surfactant.

10 Pharmaceutical Active Agent

Suitably, the pharmaceutically acceptable active agent for use in the compositions herein is selected from retinoids, retinoic acid metabolic blocking agents (RAMBAs), immune response modifier compounds, vitamin D analogues, corticosteroids, antihistamines, antibacterial agents, antifungal agents, antiviral
15 agents, cytotoxic agents, psoralens, antialopecia agents, anti-androgens, antipruritic agents, keratolytic agents, tars, dithranol, antiseptics, sunscreens, anaesthetics, analgesics, skin conditioning agents and nutritional agents, salts thereof, derivatives thereof and mixtures thereof. In one embodiment, the compositions may comprise more than one pharmaceutically active agent, salt or
20 derivative thereof. Suitable concentration ranges for the pharmaceutically active agent range from about 0.001% to about 30% by weight, depending on the nature of the active agent or combination of active agents.

In one embodiment, the pharmaceutically active agent is a retinoid. Examples of suitable retinoids include, but are not limited to, tazarotene, tretinoin, isotretinoin, acitretin, etretinate, adapalene, bexarotene, alitretinoin, retinol, retinal, retinyl palmitate, retinyl acetate, retinyl propionate, retinyl linoleate, ethyl 5-(2-(4,4-
25 dimethylthiochroman-6-yl)ethynyl)thiophene-2-carboxylate, 6-(2-(4,4-dimethylthiochroman-6-yl)-ethynyl)-3-pyridylmethanol and 6-(2-(4,4-dimethylthiochroman-6-yl)-ethynyl) pyridine-3-carbaldehyde, salts thereof, derivatives thereof and mixtures thereof. In one embodiment, the retinoid is
30 tazarotene. In an alternative embodiment, the retinoid is tretinoin. In another embodiment, the composition comprises a retinoid in combination with a second pharmaceutically active agent. In one embodiment the combination is tazarotene and a second pharmaceutically active agent. In another embodiment the
35 combination is tretinoin and a second pharmaceutically active agent.

Suitably, one combination of the retinoid is with a corticosteroid, such as clobetasol propionate; or in combination with a vitamin D analogue such as

calcipotriene; or in combination with an antibacterial such as clindamycin or a pharmaceutically acceptable salt thereof (e.g. clindamycin phosphate).

Alternatively, in an embodiment, the present compositions comprise tretinoin in combination with an antibacterial agent, such as clindamycin or a pharmaceutically acceptable salt thereof (e.g. clindamycin phosphate).

Suitable concentration ranges for the retinoid in the composition include, for example, about 0.001% to about 5% by weight. In one embodiment the retinoid is present in an amount from about 0.01% to about 1%. In another embodiment the retinoid is present in an amount from about 0.025% to about 0.5%. In one embodiment when the retinoid is tazarotene, it is present in an amount from about 0.05% or 0.1% by weight. In another embodiment when the retinoid is tretinoin, it is present in an amount from about 0.025%, 0.05% or 0.1%.

A suitable retinoic acid metabolic blocking agents (RAMBAs) for use herein as a pharmaceutically acceptable active agent is rambazole.

Suitable immune response modifier compounds, immunosuppressant agents, immunoregulating agents and immunomodulators for use herein include chemically or biologically-derived agents that modify the immune response or the functioning of the immune system (by the stimulation of antibody formation or the inhibition of white blood cell activity). Exemplary agents or compounds include, but are not limited to cyclic peptides (such as cyclosporine), tacrolimus, tresperimus, pimecrolimus, sirolimus (rapamycin), verolimus, laflunimus, laquinimod, mycophenolic acid, and imidazoquinoline amines such as imiquimod, salts thereof, derivatives thereof, and mixtures thereof.

Suitable vitamin D analogues include, but are not limited to, calcidiol, calcitriol, calcipotriene, paricalcitol, 22-oxacalcitriol, dihydrotachysterol, calciferol, salts thereof, derivatives thereof, and mixtures thereof.

Suitable corticosteroids include, but are not limited to, alclometasone dipropionate, amcinonide, beclomethasone dipropionate, betamethasone benzoate, betamethasone dipropionate, betamethasone valerate, budesonide, clobetasol propionate, clobetasone butyrate, cortisone acetate, desonide, desoximetasone, diflorasone diacetate, diflucortolone valerate, fluoclorolone acetonide, flumethasone pivalate, fluocinolone acetonide, fluocinonide, fluocortin butyl, fluocortolone, fluprednidene acetate, flurandrenolide, flurandrenolone, fluticasone propionate, halcinonide, halobetasol propionate, hydrocortisone, hydrocortisone acetate, hydrocortisone butyrate, hydrocortisone propionate, hydrocortisone valerate, methylprednisolone acetate, mometasone furoate,

pramoxine hydrochloride, prednisone acetate, prednisone valerate, triamcinolone acetonide, prednicarbate, salts thereof, derivatives thereof, and mixtures thereof.

Suitable antihistamines include, but are not limited to, cetirizine, vapitadine, diphenhydramine, triprolidine, pyrillamine, chlorcyclizine, promethazine,
5 carbinoxamine, tripelennamine, brompheniramine, hydroxyzine, terfenadine, chlorpheniramine, salts thereof, derivatives thereof, and mixtures thereof.

Suitable antibacterial agents include, but are not limited to, gentamicin, neomycin, streptomycin, cefpodoxime proxetil, clindamycin, lincomycin, erythromycin, bacitracin, gramicidin(s), vancomycin, doxycycline, minocycline,
10 oxytetracycline, tetracycline, fosfomycin, fusidic acid, mupirocin, sulfacetamide, metronidazole and dapsone, salts thereof, derivatives thereof, and mixtures thereof.

Suitable antifungal agents include, but are not limited to, those selected from the group consisting of echinocandins such as anidulafunin, caspofungin and
15 micafungin; polyenes such as amphotericin B, candicidin, filipin, fungichromin, hachimycin, hamycin, lucensomycin, mepartricin, natamycin, nystatin, pecilocin, perimycin; allylamines such as butenafine, naftifine and terbinafine; imidazoles such as bifonazole, butoconazole, chlormidazole, cloconazole, clotrimazole, econazole, enilconazole, fenticonazole, flutrimazole, isoconazole, ketoconazole,
20 lanoconazole, miconazole, neticonazole, omoconazole, oxiconazole nitrate, sertaconazole, sulconazole and tioconazole; thiocarbamates such as liranaftate, tolclate, tolindate and tolnafate; triazoles such as albaconazole, fluconazole, itraconazole, posaconazole, ravuconazole, saperconazole, terconazole and voriconazole; and other antifungal agents such as acrisorcin, amorolfine,
25 biphenamine, bromosalicylchloranilide, buclosamide, calcium propionate, chlorphenesin, ciclopirox, cloxyquin, coparaffinate, exalamide, flucytosine, haloprogin, hexetidine, loflucarban, nifuratel, potassium iodide, propionic acid, pyrrhione, salicylanilide, sodium propionate, sulbentine, tenonitroazole, triacetin, undecylenic acid, zinc propionate, griseofulvin, oligomycins, pyrrolnitrin, siccanin,
30 viridian, salts thereof, derivatives thereof, and mixtures thereof.

Suitable antivirals include, but are not limited to, acyclovir, desciclovir, carbovir, famciclovir, foscarnet sodium, ganciclovir, interferons, penciclovir, valaciclovir, salts thereof, derivatives thereof, and mixtures thereof.

Suitable cytotoxic agents include, but are not limited to, azathioprine,
35 cyclophosphamide, cyclosporine, methotrexate, hydroxyurea, thalidomide, bleomycin, fluorouracil, salts thereof, derivatives thereof, and mixtures thereof.

An exemplary psoralen is methoxsalen.

Suitable anti-androgens include, but are not limited to, spironolactone, cyproterone, flutamide, finasteride, salts thereof, derivatives thereof, and mixtures thereof.

An exemplary antialopecia agent is minoxidil.

5 Suitable antipruritics include, but are not limited to, calamine, camphor and menthol, salts thereof, derivatives thereof, and mixtures thereof.

Suitable keratolytic agents include, but are not limited to, benzoyl peroxide, salicylic acid, urea, resorcinol, sulphur, salts thereof, derivatives thereof, and mixtures thereof.

10 Suitable tars include, but are not limited to, coal tar, pine tar, wood tar, salts thereof, derivatives thereof, and mixtures thereof.

Suitable antiseptics include, but are not limited to, hydrogen peroxide, chlorhexidine, cetrimide, povidone iodine, triclosan, salts thereof, derivatives thereof, and mixtures thereof.

15 Suitable sunscreens include, but are not limited to, aminobenzoic acid, avobenzene, bemotrizinol, bisoctrizole, β -carotene, cinoxate, 4-(dimethylamino)benzoic acid, dioxybenzone, drometrizole, ecamsule, ensulizole, ethylhexyl triazone, homosalate, lawsone, menthyl anthranilate, 4-methylbenzylidene camphor, mexenone, octabenzene, octocrylene, octyl
20 methoxycinnamate, octyl salicylate, oxybenzone, sulisobenzene, zinc oxide, titanium dioxide, salts thereof, derivatives thereof, and mixtures thereof.

Suitable anaesthetics and analgesics include, but are not limited to, benzocaine, lidocaine, prilocaine and choline salicylate, salts thereof, derivatives thereof, and mixtures thereof.

25 Suitable skin-conditioning agents include, but are not limited to, hydrocarbon oils and waxes, silicones, fatty acid derivatives, cholesterol, di- and tri-glycerides, vegetable oils, acetoglyceride esters, alkyl esters, alkenyl esters, lanolin, milk tri-glycerides, wax esters, beeswax, sterols, phospholipids, derivatives thereof, and mixtures thereof.

30 Exemplary nutritional agents include vitamins, essential amino acids, essential fats, antioxidants, salts thereof, derivatives thereof, and mixtures thereof.

Water Phase

35 The topical foam compositions of the present invention also comprise water. Water forms the continuous phase of the emulsion system. In an embodiment, the foam compositions comprise water in an amount from about 65% to about 90% by weight. In another embodiment the water is present in an

amount from about 70% to about 85% by weight. In another embodiment the water is present at about 77% by weight.

Oil miscible organic solvent

5 The present topical foam compositions comprise an oil miscible organic solvent to facilitate solubilisation of the active agent in the oil. Together, the oil miscible organic solvent and oil comprise the oil phase of the composition.

 The oil miscible organic solvent is present in the composition in an amount from about 1% to about 20% by weight. In one embodiment the oil miscible
10 organic solvent is present in an amount from about 3% to about 15% by weight. In another embodiment the oil miscible organic solvent is present in an amount of about 5% by weight.

 Suitable, non-limiting examples of oil miscible organic solvents include alcohols such as caprylic alcohol, decyl alcohol, dodecylhexadecanol,
15 dodecyltetradecanol, hexyl alcohol, hexyldecanol, hexyldecyloctadecanol, isocetyl alcohol, isostearyl alcohol, lauryl alcohol, myristyl alcohol, octyldecanol, octyldodecanol, oleyl alcohol, tridecyl alcohol; esters such as butyl stearate, C12-15 alkyl benzoate, C12-15 alkyl lactate, caprylic/capric triglyceride, cetearyl ethylhexanoate, cetearyl isononanoate, cetyl octanoate, cetyl palmitate, coco-
20 caprylate/caprinate, cocoglycerides, decyl oleate, dibutyl adipate, dicaprylyl carbonate, diethylhexyl adipate, di-ethylhexyl succinate, diisopropyl adipate, dioctyl malate, di-PPG-2 myreth-10 adipate, di-PPG-3 myristyl ether adipate, ethyl oleate, ethylhexyl cocoate, ethylhexyl hydroxystearate, ethylhexyl palmitate, ethylhexyl pelargonate, ethylhexyl stearate, hexyl laurate, hexyldecyl laurate,
25 hexyldecyl stearate, isocetyl stearate, isocetyl stearyl stearate, isodecyl oleate, isopropyl myristate, isopropyl palmitate, isostearyl neopentanoate, isotridecyl isononanoate, lauryl lactate, myristyl lactate, myristyl myristate, octyldodecyl stearyl stearate, oleyl erucate, oleyl oleate, pentaerythrityl tetracaprylate/caprinate, pentaerythrityl tetraisostearate, PPG-2 myristyl ether propionate, propylene glycol
30 dicaprylate/ dicaprinate, propylene glycol isostearate, propylheptyl caprylate, stearyl octanoate; ethers such as dicaprylyl ether, PPG-10 cetyl ether, PPG-11 stearyl ether, PPG-14 butyl ether, PPG-15 stearyl ether, PPG-3 hydrogenated castor oil, PPG-3 myristyl ether; carboxylic acids such as C10-40 hydroxyalkyl acid, C10-40 isoalkyl acid, C32-36 isoalkyl acid, capric acid, caproic acid, caprylic acid, coconut
35 acid, lauric acid, linoleic acid, linolenic acid, linseed acid, myristic acid, oleic acid, ricinoleic acid and lanolin.

According to an embodiment, the oil miscible organic solvent is selected from the group consisting of diisopropyl adipate, isopropyl myristate, dodecanol and caprylic / capric triglycerides.

5 In one embodiment, the oil miscible organic solvent is diisopropyl adipate (also referred to herein as DIPA). In one embodiment the DIPA is present in the composition in an amount from about 1% to about 20% by weight. In another embodiment the DIPA is present in an amount from about 3% to about 15% by weight. In another embodiment the DIPA is present in an amount of about 5% by weight.

10

According to a further embodiment, the present compositions comprise a second pharmaceutically acceptable active agent. In one embodiment, the second active agent is solubilized in the water phase (i.e. the continuous phase). This may be achieved by dissolving the second active agent in an aliquot of water or an aliquot of a water miscible organic solvent, and subsequently solubilising the aliquot in the composition.

15 According to an embodiment, the water miscible organic solvent is present in an amount from about 1% to about 20% by weight. According to a further embodiment, the water miscible organic solvent is present in an amount of about 10% by weight.

Suitably, water miscible organic solvents include, but are not limited to, alcohols, including amyl alcohol, benzyl alcohol, cyclohexanedimethanol, diacetone alcohol, ethyl alcohol, hexyl alcohol, isobutyl alcohol, isopropyl alcohol, methyl alcohol, n-butyl alcohol, propyl alcohol, t-butyl alcohol, tetrahydrofurfuryl alcohol; carboxylic acids, including acetic acid; diols, including 1,2-hexanediol, butylene glycol, diethylene glycol, dipropylene glycol, ethyl hexanediol, ethylene glycol, hexylene glycol, pentylene glycol, propylene glycol, tetraethylene glycol, triethylene glycol, tripropylene glycol; and polyols including polyethylene glycol, butanetriol, glycerol and 1,2,6-hexanetriol.

25 30 In one embodiment, the water miscible organic solvent is propylene glycol.

Propellant

It is recognized that the present topical foam compositions must utilize a propellant in order to produce the foam upon application. The propellant may be any suitable liquefied gas or mixture thereof, such as a hydrocarbon, a chlorofluorocarbon, dimethyl ether, hydrofluorocarbons and a mixture thereof.

35

Other suitable propellants include compressed gases such as nitrogen, carbon dioxide, nitrous oxide and air. In a preferred embodiment, the propellant is a mixture of hydrocarbons. In a further preferred embodiment, the mixture of hydrocarbons is a mixture of propane, n-butane and isobutane.

5 The propellant is present in an amount from about 1% to about 20% by weight, or about 3% to about 15% by weight. In one embodiment, the propellant is present in an amount from about 5% to about 10% by weight, such as about 5%, 6%, 7%, 8%, 9% or 10% by weight. The propellant may be introduced into the composition at the time of filling, utilizing a pressurized container such as a
10 standard aerosol dispenser.

 When the composition is released from the pressurized container, the composition is an aerosol foam (also known as a mousse). According to one embodiment, the aerosol foam is homogeneous. In another embodiment, the aerosol foam breaks easily with shear, such as gentle mechanical action e.g.
15 rubbing or spreading.

 In another embodiment the propellant is absent from the composition. According to such an embodiment, the composition may be expelled from its container by mechanical means, such as by a pump action or a squeezing action on the container.

20 Suitable pressurized containers for use herein include aluminium, tin-plate and glass containers.

 In one embodiment, the pressurized container is a one-piece aluminium container in which the inner surface is lined with a chemically inert lining. One suitable inner surface lining for use herein is polyamide-imide (PAM), such as that
25 supplied by Exal Corporation, of Youngstown, Ohio. The container may be fitted with an upright-use or inverted-use valve and a conventional foam spout actuator. Alternatively, the container may be fitted with a metered-dose valve.

Dermatologically acceptable excipients

30 According to an embodiment, the compositions may further comprise one or more dermatologically acceptable excipients. Non-limiting examples of such dermatologically acceptable excipients include diluents, suspending agents, adjuvants, preservatives, colorants, emollients, pH adjusting agents (including buffers), thickeners, humectants, fragrances, stabilizers, chelating agents,
35 anticaking agents, viscosity increasing agents, solubilizers, plasticizers, penetration enhancing agents, film forming agents, antioxidants, wetting agents, foam boosters or any mixture of these components.

In one embodiment, the one or more dermatologically acceptable excipients comprise a preservative, an antioxidant and a pH adjusting agent.

Preservative

5 The present topical aerosol foam compositions may additionally comprise a preservative. The preservative is present in the composition in an amount from about 0.01% to about 2% by weight. In one embodiment the preservative is present in an amount from about 0.1% to about 1% by weight. In another embodiment the preservative is present in an amount of about 0.3% by weight.

10 Suitable preservatives include, but are not limited to benzyl alcohol, diazolidinyl urea, methyl paraben, ethyl paraben, propyl paraben, butyl paraben, phenoxyethanol, sorbic acid and salts thereof such as potassium sorbate, benzoic acid and salts thereof such as sodium benzoate, and mixtures thereof.

15 According to an embodiment, the preservative is a combination of sorbic acid and potassium sorbate.

Antioxidant

The present topical aerosol foam compositions may further comprise an antioxidant. The antioxidant is present in the composition in an amount from about 0.001% to about 1% by weight. In one embodiment the antioxidant is present from about 0.05% to about 0.5% by weight. In another embodiment the antioxidant is present in an amount of about 0.1% by weight.

25 Suitable antioxidants include, but are not limited to, butylated hydroxytoluene (BHT), butylated hydroxyanisole, tocopherol, propyl gallate, vitamin E TPGS, derivatives thereof, and mixtures thereof. In one embodiment, the antioxidant is BHT.

pH Adjusting Agent

30 The present topical aerosol foam compositions may further comprise a pH adjusting agent to aid in stabilizing the active agent. According to an embodiment, the pH adjusting agent is present in an amount from about 0.01% to about 10% by weight. In one embodiment, the pH adjusting agent is a base. Suitable pH adjusting bases include but are not limited to bicarbonates, carbonates and hydroxides (such as alkali or alkaline earth metal hydroxides, as well as transition metal hydroxides). The pH adjusting agent may also be an acid, an acid salt, or mixtures thereof. The pH adjusting agent may also be a buffer. Suitable buffers include, but are not limited to citrate/ citric acid, acetate/ acetic acid, phosphate/

phosphoric acid, formate/ formic acid, propionate/ propionic acid, lactate/ lactic acid, carbonate/ carbonic acid, ammonium/ ammonia, edentate/ edetic acid, derivatives thereof, and combinations thereof. According to an embodiment, the pH adjusting agent is a citrate/citric acid buffer. According to an embodiment, the
5 citrate / citric acid buffer is present in an amount from about 0.02% to about 2% by weight.

In another embodiment, the present invention relates to an oil in water emulsion aerosol foam composition comprising: tazarotene, water, an oil in an
10 amount from about 3% to about 8% by weight, an oil miscible organic solvent, a surfactant component comprising a non-ionic hydrophilic surfactant in an amount from about 2% to about 6% by weight, and a hydrocarbon propellant, wherein the tazarotene is solubilized in the oil phase of the composition and wherein the particle size of the oil phase is less than about 1000 nm.

15 According to an embodiment, the mean particle size of the oil phase of the tazarotene aerosol foam is about 100 nm.

The tazarotene foam formulation exemplified by these embodiments is particularly suited for application to small regions of the face for the treatment of acne. In one embodiment, the tazarotene aerosol foam formulation is dispensed
20 from a metered dose actuator. In another alternative embodiment, the tazarotene aerosol foam composition is formulated as a macro emulsion (i.e. the particle size of the oil phase is greater than 1000nm).

In another embodiment, a second pharmaceutically active agent is solubilized in the composition. In another embodiment the second
25 pharmaceutically active agent is solubilized in the water phase of the composition. This may be achieved by dissolving the further pharmaceutically active agent in an aliquot of water which is subsequently solubilized in the composition.

Alternatively, the second pharmaceutically active agent is dissolved in an aliquot of water miscible organic solvent which is subsequently solubilized in the
30 composition. In another embodiment the second pharmaceutically active agent is in a composition comprising the retinoid tazarotene.

Definitions

As used herein, the terms "administering" and "administered," refer to any
35 method which delivers the composition to a subject in such a manner as to provide a therapeutic effect.

As used herein, the term "derivative(s) thereof" refers to prodrugs, solvates, hydrates, esters and acids of the pharmaceutically active agent.

As used herein, the phrase an "effective amount" of an active agent or ingredient, or pharmaceutically active agent or ingredient, which are synonymous
5 herein, refers to an amount of the pharmaceutically active agent sufficient to have a therapeutic effect upon administration. An effective amount of the active agent may, will, or is expected to cause relief of symptoms. Effective amounts of the active agent will vary with the particular disease or diseases being treated, the severity of the disease, the duration of the treatment, and the specific components
10 of the composition being used.

As used herein, the term "fatty alcohol" refers to an alcohol having an aliphatic chain from about 9 to about 22 carbon atoms long.

As used herein, an "emulsion" refers to a mixture of two or more immiscible (unblendable) liquids wherein the particle size of the dispersed phase (i.e. oil in
15 the case of an oil in water emulsion) is less than about 10,000 nm.

As used herein, a "submicron emulsion" refers to a mixture of two or more immiscible (unblendable) liquids wherein the particle size of the dispersed phase (i.e. oil in the case of an oil in water emulsion) is in the range from about 100 nm to about 1000 nm.

As used herein, a "microemulsion" refers to a mixture of two or more immiscible (unblendable) liquids wherein the particle size of the dispersed phase (i.e. oil in the case of an oil in water emulsion) is in the range from about 10 nm to about 100 nm.

As used herein, "nanoemulsion" refers to a mixture of two or more
25 immiscible (unblendable) liquids wherein the particle size of the dispersed phase (i.e. oil in the case of an oil in water emulsion) is in the range from about 1 nm to 100 nm.

As used herein, the term "phase inversion temperature" refers to a temperature where an oil in water emulsion inverts to a water in oil emulsion (or
30 vice versa).

As used herein, a "pH adjusting agent" refers to a specific pH adjusting agent or agents, including but not limited to, a buffer, a base or an acid, salts thereof and mixtures thereof, added to a composition.

The phrase "dermatologically acceptable excipient" as used herein refers to
35 any inactive ingredient present in the herein described compositions.

As used herein, the phrase "pharmaceutically acceptable salts" refers to salts that are pharmaceutically acceptable and that possess the desired

pharmacological activity of the parent compound. Such salts include: (1) acid addition salts, formed with acids such as, for example, acetic acid, benzoic acid, citric acid, gluconic acid, glutamic acid, glutaric acid, glycolic acid, hydrochloric acid, lactic acid, maleic acid, malic acid, malonic acid, mandelic acid, phosphoric acid, propionic acid, sorbic acid, succinic acid, sulfuric acid, tartaric acid, naturally and synthetically derived amino acids, and mixtures thereof; or (2) salts formed when an acidic proton present in the parent compound is either (i) replaced by a metal ion e.g. an alkali metal ion, an alkaline earth metal ion, or an aluminium ion; or (ii) protonates an organic base such as, for example, ethanolamine, diethanolamine, triethanolamine, tromethamine and N-methylglucamine.

As used herein, a "subject", "individual" or "patient" refers to any subject, particularly a human, for whom therapy is desired.

As used herein, a "treatment" or "treating" of a disease, disorder or condition encompasses alleviation of at least one symptom thereof, a reduction in the severity thereof, or the delay, prevention or inhibition of the progression thereof. Treatment need not mean that the disease, disorder or condition is totally cured. A useful composition herein need only to reduce the severity of a disease, disorder or condition, reduce the severity of symptoms associated therewith, provide improvement to a patient's quality of life, or delay, prevent or inhibit the onset of a disease, disorder or condition.

Any concentration range, percentage range or ratio range recited herein is to be understood to include concentrations, percentages or ratios of any integer within that range and fractions thereof, such as one tenth and one hundredth of an integer, unless otherwise indicated.

It should be understood that the terms "a" and "an" as used above and elsewhere herein refer to "one or more" of the enumerated components. It will be clear to one of ordinary skill in the art that the use of the singular includes the plural unless specifically stated otherwise. Therefore, the terms "a," "an" and "at least one" are used interchangeably in this application.

Throughout the application, descriptions of various embodiments use "comprising" language, however in some specific instances, an embodiment can alternatively be described using the language "consisting essentially of" or "consisting of".

For the purposes of better understanding the present teachings and in no way limiting their scope, unless otherwise indicated, all numbers expressing quantities, percentages or proportions, and other numerical values used in the

specification and claims, are to be understood as being modified in all instances by the term "about."

As used herein, the term "substantially free" of a specified component refers to a composition with less than about 1% of the specified component.

5 Other terms used herein are intended to be defined by their well-known meanings in the art.

Process of Preparation

10 In one embodiment, the present invention provides a process for the preparation of an oil in water submicron or micro emulsion aerosol foam composition, comprising a pharmaceutically active agent, water, an oil present in an amount of less than about 10% by weight, an oil miscible organic solvent, a surfactant component comprising a hydrophilic surfactant in an amount from about 0.1% to about 10% by weight, and a propellant, the process comprising:

- 15
- a) admixing the pharmaceutically active agent, a first aliquot of water, oil, oil miscible organic solvent and surfactant component to form an oil in water emulsion,
 - 20 b) heating the oil in water emulsion of step (a) to a phase inversion temperature wherein the oil in water emulsion forms a water in oil emulsion,
 - c) cooling the water in oil emulsion to below the phase inversion temperature to form a submicron or micro oil in water emulsion,
 - 25 d) adding a second aliquot of water to cool the submicron or micro oil in water emulsion,
 - e) actuating a sample of the submicron or micro oil in water emulsion with a propellant to form an oil in water submicron or micro emulsion aerosol foam.

30 According to an embodiment, the first aliquot of water comprises about 10% to about 40% by weight of the total water content of the submicron or micro oil in water emulsion aerosol foam and the second aliquot of water comprises about 90% to about 60% by weight of the total water content of the submicron or micro oil in water emulsion aerosol foam.

35 According to an embodiment, the first aliquot of water comprises about 10% to about 40% by weight of the total water content of the submicron or micro oil in water emulsion aerosol foam and the second aliquot of water comprises

about 90% to about 60% by weight of the total water content of the submicron or micro oil in water emulsion aerosol foam.

According to another embodiment, the first aliquot of water comprises about 15% to about 35% of the total water content of the submicron or micro oil in water emulsion aerosol foam and the second aliquot of water comprises about 85% to about 65% of the total water content of the submicron or micro oil in water emulsion aerosol foam.

According to a further embodiment, the first aliquot of water comprises about 20% to about 30% of the total water content of the submicron or micro oil in water emulsion aerosol foam and the second aliquot of water comprises about 80% to about 70% of the total water content of the submicron or micro oil in water emulsion aerosol foam. According to yet a further embodiment, the first aliquot of water comprises about 25% of the total water content of the submicron or micro oil in water emulsion aerosol foam and the second aliquot of water comprises about 75% of the total water content of the submicron or micro oil in water emulsion aerosol foam.

The presently described process provides for the preparation of a low oil and low surfactant submicron or micro emulsion aerosol foam, where a concentrated submicron or micro emulsion is prepared (comprising the first aliquot of water) and is subsequently diluted (with the second aliquot of water) to ultimately afford the desired low oil and low surfactant submicron or micro emulsion. By initially preparing a concentrated submicron or micro emulsion, a sufficient concentration of surfactant is present in the composition to stabilize the submicron or micro sized oil phase. The dilution step then provides the desired low oil and low surfactant submicron or micro emulsion. It is thought that the addition of the second aliquot of water (which is at a temperature substantially below the phase inversion temperature) serves to fix the submicron or micro emulsion structure of the composition and, of course, dilutes the composition to provide the low oil and low surfactant composition.

According to an embodiment, a second pharmaceutically active agent is solubilized in the water phase of the composition. This may be achieved by dissolving the second pharmaceutically active agent in an aliquot of water or water miscible organic solvent, which is subsequently solubilized in the composition.

In one embodiment, the second pharmaceutically active agent dissolved in an aliquot of water or water miscible organic solvent is added to the submicron or micro oil in water emulsion following step (d)). That is, the second

pharmaceutically active agent is solubilized in the water phase of the composition after formation of the submicron or micro emulsion.

Another aspect of the invention is the product produced by this process.

5 **Methods of Treatment**

The emulsion aerosol foam compositions of the present invention are cosmetically elegant and suitable for application to the face for treating a skin disorder or condition. The compositions are easily spread, non-greasy, non-drying and leave minimal residue on the skin.

10 The present invention provides for a method of treating a skin disease, disorder or condition, comprising administering to the skin of a patient requiring such treatment an effective amount of a composition of the present invention. In one embodiment, the skin disease, disorder or condition is acne. In another embodiment, the skin disease, disorder or condition is psoriasis.

15 The present invention also relates to the use of the compositions as described herein for the preparation of a medicament for the treatment of a skin disease, disorder or condition.

The present invention also relates to a method of treating a skin disease, disorder or condition by administering to the skin of a patient requiring such
20 treatment an effective amount of a composition of the present invention.

Exemplary, non-limiting, skin diseases, disorders or conditions treatable by the present compositions include acne, rosacea, dermatitis, psoriasis and fungal disorders. According to an embodiment, the skin disease, disorder or condition is acne. According to another embodiment, the skin disease, disorder or condition is
25 psoriasis.

In an embodiment, the present compositions are used in combination with a suitable additional pharmaceutical dosage form. The additional pharmaceutical dosage form is administered to a patient either prior to, concomitantly with, or after the compositions described herein.

30 In one embodiment in this regard, the present composition and the additional pharmaceutical dosage form are administered to a patient at the same time. In an alternative embodiment, one of the present compositions and the additional pharmaceutical dosage form is administered in the morning and the other is administered in the evening.

35 In another embodiment, the additional pharmaceutical dosage form can be a suitable oral pharmaceutical. In this regard, the present composition can be

applied to the target area of the patient, prior to, concomitantly with, or after ingestion of the oral medication.

According to an embodiment, the present composition comprises a retinoid and the oral pharmaceutical dosage form comprises a pharmaceutically active agent selected from the group consisting of an antibiotic, a contraceptive, a retinoid such as isotretinoin, and a nutritional agent.

Furthermore, the foam compositions of the present invention may be used with other adjunct therapies and treatments, such as pre-washing with soaps or cleansers. However, care is needed in this regard, since antibacterial soaps and abrasive soaps may increase irritation.

EXAMPLES

The following examples are illustrative of preferred embodiments herein and are not to be construed as limiting the present invention thereto. All percentages are based on the percent by weight of the final delivery system or formulation prepared, unless otherwise indicated and all totals equal 100% by weight.

Tables 2a, 2b, 2c and 2d

Oil in water submicron / micro emulsion foam formulations – tazarotene (0.1%)

706/8/2 (Table 2a) – absent propellant*

Component		% w/w
	BHT	0.10
	Macrogol Cetostearyl Ether 12 (Ceteareth 12)	5.00
	Light Mineral Oil	6.00
	Sorbic Acid	0.40
	DIPA	5.00
	Tazarotene	0.10
	Citric Acid Anhydrous	0.03
	Potassium Citrate Monohydrate	0.57
	Deionized Water	82.80
	Total	100.000

706/8/3 (Table 2b) – absent propellant*

Component	% w/w
BHT	0.10
Macrogol Cetostearyl Ether 12 (Cetareth 12)	5.00
Light Mineral Oil	8.00
Sorbic Acid	0.40
DIPA	7.00
Tazarotene	0.10
Citric Acid Anhydrous	0.03
Potassium Citrate Monohydrate	0.54
Deionized Water	78.83
Total	100.000

*The formulations described in Tables 2a & 2b were dispensed as a foam following the addition of approximately 7 to 8 grams of AP70 propellant per 100 grams of aerosol base.

730/2/1 (Table 2c)

Component	Function	% w/w	% w/w
BHT	Antioxidant	0.100	0.093
Macrogol Cetostearyl Ether 12 (Cetareth 12)	Surfactant	5.000	4.635
Light Mineral Oil	Oil	6.000	5.562
Sorbic Acid	Preservative	0.150	0.139
Potassium Sorbate	Preservative	0.200	0.185
Diisopropyl adipate	Organic Solvent	5.000	4.635
Tazarotene	Active ingredient	0.100	0.093
Citric Acid Anhydrous	Buffer	0.037	0.034
Potassium Citrate Monohydrate	Buffer	0.103	0.096
Deionized Water	Water	83.310	77.228
Propellant AP70 (propane, butane & isobutane)	Propellant	-	7.300
Total		100.000	100.000

SFW0260-02 (Table 2d)

Component	Function	% w/w	% w/w
BHT	Antioxidant	0.100	0.093
Macrogol Cetostearyl Ether 12 (Cetareth 12)	Surfactant	5.000	4.648
Light Mineral Oil	Oil	6.000	5.578
Sorbic Acid	Preservative	0.150	0.139
Potassium Sorbate	Preservative	0.200	0.186
Diisopropyl adipate	Organic Solvent	5.000	4.648
Tazarotene	Active ingredient	0.100	0.093
Citric Acid Anhydrous	Buffer	0.037	0.034
Potassium Citrate Monohydrate	Buffer	0.103	0.096
Deionized Water	Water	83.310	77.445
Propellant AP70 (propane, butane & isobutane)	Propellant	-	7.040
Total		100.000	100.000

5 **Example 1**

Method of preparing submicron / micro emulsion foam formulation – tazarotene

Procedure: aerosol base production (730/2/1):

- Phase 1 (Oil phase): BHT, cetareth-12, mineral oil and sorbic acid
- 10 Phase 2 (Active phase): tazarotene dissolved in diisopropyl adipate (DIPA)
- Phase 3 (Buffer phase): deionized water (about 22.8% of the total water content of aerosol base), citric acid, potassium citrate and potassium sorbate
- Phase 4 (Water phase): deionized water (about 77.2% of the total water content of aerosol base)
- 15

1. Dissolve tazarotene in diisopropyl adipate in a suitable stainless steel container at ambient temperature (Phase 2).
 2. Prepare buffer solution at ambient temperature in a stainless steel container (Phase 3).
 3. Add oil phase ingredients to the main batch vessel (Phase 1).
 4. Combine Phase 1, Phase 2 and Phase 3 in the main batch vessel. Commence low heating with stirring. Continue heating to the phase inversion temperature range of 70°C to 80°C until a water in oil emulsion occurs.
 5. Commence cooling to below the phase inversion temperature. When the emulsion becomes an oil in water submicron / micro emulsion, add ambient temperature Phase 4 deionized water at a moderate rate. Stir until uniform. Maintain batch at 25°C-30°C.
 6. Test pH of the aerosol base.
 7. Adjust pH to a range of 4.70 to 5.50 with 1N potassium hydroxide solution or with 10% citric acid solution.
 8. Adjust to 100% with deionized water (to account for evaporative water loss).
- The abovementioned procedure is represented in schematic form in Figure 1.

Alternative procedure – aerosol base production (SFW0260-02):

At ambient temperature, dissolve tazarotene in diisopropyl adipate in a side vessel, while stirring.

Add mineral oil and BHT to the side vessel, while stirring until the solution is homogenous.

At ambient temperature, add deionized water (about 28.8% of total water content of aerosol base), citric acid and potassium citrate to a main batch vessel, while stirring until all components are fully dissolved.

Add potassium sorbate to the main batch vessel, while stirring until the potassium sorbate is fully dissolved.

Continue to stir the contents of the main batch vessel and add the contents of the side vessel (tazarotene, DIPA, mineral oil and BHT) to the main batch vessel, followed by addition of sorbic acid and cetareth-12.

Commence heating and continue heating until the phase inversion temperature is reached i.e. where the oil in water emulsion inverts to a water in oil

emulsion (as measured by conductivity or visual observation). The phase inversion temperature is in the range of about 70°C to about 80°C.

- 5 Cool the main batch vessel to below the phase inversion temperature (approximately 69°C) to facilitate the formation of an oil in water submicron / micro emulsion.

Add ambient temperature deionized water (about 71.2% of total water content of aerosol base). Continue stirring and maintain main batch vessel at around 25°C – 30°C.

- 10 Test the pH and adjust as necessary to a range of 4.70 to 5.50. Adjust with deionized water to account for evaporative water loss.

Aerosol Filling and Crimping

- 15 The aerosol base is filled into suitable aluminium aerosol containers, suitable valves are inserted, vacuum crimped and gassed with a suitable propellant.

Table 3

Submicron / micro emulsion foam - tazarotene (0.1%) plus calcipotriene (0.005%)

Item	Ingredient	Function	%w/w
	First active phase		
1	Diisopropyl adipate	Organic solvent	5.000
2	Tazarotene	Active ingredient	0.100
	Hydrocarbon oil phase		
3	BHT	Antioxidant	0.100
4	Macrogol Cetostearyl Ether 12 (Ceteareth 12)	Surfactant	5.000
5	Mineral oil (light)	Hydrocarbon	6.000
6	Tocopherol	Antioxidant	0.002
	First water phase		
7	Water (deionized)	Aqueous solvent	19.000
8	EDTA, Na ₂	Buffer	0.060
9	Disodium phosphate	Buffer	0.080
	Second water phase		
10	Water (deionized)	Aqueous solvent	54.653
	Second active phase		
11	Propylene glycol	Organic solvent	10.000
12	Calcipotriene	Active ingredient	0.005
	Total		100.000
	Finished product		
Items	Ingredient	Function	%w/w
1 to 12	Aerosol base	Concentrate	92.7
13	Hydrocarbon propellant AP70	Propellant	7.3
	Total		100.0

Example 2**Method of preparing submicron / micro emulsion foam –**

5 **tazarotene (0.1%) plus calcipotriene (0.005%)**

Aerosol base:

- 10 1. Add BHT (Item 3), cetareth-12 (Item 4), mineral oil (Item 5) and tocopherol (Item 6) to the main mixing vessel.
2. Prepare the First active phase by adding diisopropyl adipate (Item 1) and tazarotene (Item 2) to a small mixing vessel and mix until tazarotene has dissolved.
3. Add the First active phase solution (Items 1 and 2) to the main mixing vessel.
- 15 4. Into a separate mixing vessel prepare First water phase by adding water (Item 7).
5. Start stirring First water phase and add EDTA, Na₂ (Item 8) and disodium phosphate (Item 9) to water and mix until dissolved.
- 20 6. Add First water phase (Items 7 to 9) to the main mixing vessel.
7. While monitoring the conductivity of the contents of the main mixing vessel (Items 1 to 9) commence heating, to approximately 80°C, and start stirring.
8. As the temperature increases the mixture inverts from an oil in water emulsion to a water in oil emulsion and a corresponding decrease in the conductivity is observed.
- 25 9. When the temperature reaches approximately 80°C stop heating and commence slow cooling of the contents of the main mixing vessel while maintaining stirring.
10. At approximately 70°C the mixture inverts from a water in oil emulsion to an oil in water emulsion and a corresponding increase in the conductivity is observed. The appearance of the mixture also changes from opaque to translucent.
- 30 11. At this temperature, when the mixture is translucent, add the Second water phase (Item 10), at ambient temperature, to the main mixing vessel and continue stirring. The temperature of the mixture will decrease to approximately 40°C following the addition of the Second water phase.
- 35

12. While stirring the mixture, continue cooling until the temperature range decreases to approximately 25°C to 30°C.
13. Prepare the Second active phase by adding propylene glycol (Item 11) and calcipotriene (Item 12) to a small mixing vessel and mix until calcipotriene has dissolved.
14. Transfer the Second active phase (Items 11 and 12) to the main mixing vessel and stir until the mixture is uniform.

Finished product:

1. Add Aerosol base (Items 1 to 12) to an empty aerosol container.
2. Secure an aerosol valve onto the aerosol container.
3. Add Propellant (Item 13) to the aerosol container.
4. Test the aerosol container to confirm that there is no leakage.
5. Place an actuator onto the aerosol valve.
6. Shake the aerosol container prior to dispensing at room temperature.

Table 4

Submicron / micro emulsion foam - clindamycin phosphate (1%) plus tretinoin (0.05%)

Item	Ingredient	Function	%w/w
First active phase			
1	Diisopropyl adipate	Organic solvent	10.000
2	Tretinoin	Active ingredient	0.050
Hydrocarbon oil phase			
3	BHT	Antioxidant	0.100
4	Macrogol Cetostearyl Ether 12 (Ceteareth 12)	Surfactant	5.000
5	Mineral oil (light)	Hydrocarbon	6.000
First water phase			
6	Water (deionized)	Aqueous solvent	22.000
7	Anhydrous citric acid	Buffer	0.037

8	Potassium citrate monohydrate	Buffer	0.103
	Second water phase		
9	Water (deionized)	Aqueous solvent	32.960
	Second active phase		
10	Water (deionized)	Aqueous solvent	20.000
11	Clindamycin phosphate (80% active)	Active ingredient	1.250
	Preservative phase		
12	Propylene glycol		2.000
13	Benzyl alcohol		0.500
	Total		100.000
	Finished product		
Items	Ingredient	Function	%w/w
1 to 13	Aerosol base	Concentrate	91.1
14	Hydrocarbon propellant AP70	Propellant	8.9
	Total		100.0

Example 3

5 **Method of preparing submicron / micro emulsion foam formulation – clindamycin phosphate (1%) plus tretinoin (0.05%)**

Aerosol base:

- 10 1. Add BHT (Item 3), cetareth-12 (Item 4) and mineral oil (Item 5) to the main mixing vessel.

2. Prepare the First active phase by adding (Item 1) and tretinoin (Item 2) to a small mixing vessel and mix until dissolved.
3. Add the First active phase solution (Items 1 and 2) to the main mixing vessel.
- 5 4. Into a separate mixing vessel prepare First water phase by adding water (Item 6).
5. Start stirring First water phase and add anhydrous citric acid (Item 7) and potassium citrate monohydrate (Item 8) to water and mix until dissolved.
- 10 6. Add First water phase (Items 6 to 8) to the main mixing vessel.
7. While monitoring the conductivity of the contents of the main mixing vessel (Items 1 to 8) commence heating, to approximately 80°C, and start stirring.
8. As the temperature increases the mixture inverts from an oil in water emulsion to a water in oil emulsion and a corresponding decrease in the conductivity is observed.
- 15 9. After the mixture inverts from an oil in water emulsion to a water in oil emulsion stop heating and commence slow cooling of the contents of the main mixing vessel while maintaining stirring.
- 20 10. With cooling, the mixture inverts from a water in oil emulsion back to an oil in water emulsion and a corresponding increase in the conductivity is observed. The appearance of the mixture also changes from opaque to translucent.
11. At this temperature, when the mixture is translucent, add the Second water phase (Item 9), at ambient temperature, to the main mixing vessel and continue stirring. The temperature of the mixture will decrease following the addition of the Second water phase.
- 25 12. While stirring the mixture, continue cooling until the temperature range decreases to approximately 25°C to 30°C.
- 30 13. Prepare the Second active phase by adding water (Item 10) and clindamycin phosphate (Item 11) to a small mixing vessel and mix until clindamycin phosphate has dissolved.
14. Transfer the Second active phase (Items 10 and 11) to the main mixing vessel and stir until the mixture is uniform.
- 35 15. Prepare the Preservative phase by adding propylene glycol (Item 12) and benzyl alcohol (Item 13) to a small mixing vessel and mix until uniform.

16. Transfer the Preservative phase (Items 12 and 13) to the main mixing vessel and stir until the mixture is uniform.

Finished product:

- 5 1. Add Aerosol base (Items 1 to 13) to an empty aerosol container.
2. Secure an aerosol valve onto the aerosol container.
3. Add Propellant (Item 14) to the aerosol container.
4. Test the aerosol container to confirm that there is no leakage.
5. Place an actuator onto the aerosol valve.
- 10 6. Shake the aerosol container prior to dispensing at room temperature.

The formulations described in tables 5 to 9 are further illustrative of the present invention.

15

Table 5

Submicron / micro emulsion foam - tazarotene (0.1%) plus clobetasol propionate (0.05%)

Item	Ingredient	Function	%w/w
	First active phase		
1	Diisopropyl adipate	Organic solvent	5.000
2	Tazarotene	Active ingredient	0.100
	Hydrocarbon oil phase		
3	BHT	Antioxidant	0.100
4	Macrogol Cetostearyl Ether 12 (Cetareth 12)	Surfactant	5.000
5	Mineral oil (light)	Hydrocarbon	2.000
6	Sorbic acid	Preservative	0.200
7	Petrolatum	Hydrocarbon	2.000
	First water phase		
8	Water (deionized)	Aqueous solvent	22.030
9	Anhydrous citric acid	Buffer	0.040
10	Potassium citrate monohydrate	Buffer	0.100
11	Potassium sorbate	Preservative	0.270

5		Second water phase		
	12	Water (deionized)	Aqueous solvent	53.110
		Second active phase		
	13	Propylene glycol	Organic solvent	10.000
	14	Clobetasol propionate	Active ingredient	0.050
10		Total		100.000
		Finished product		
	Items	Ingredient	Function	%w/w
	1 to 14	Aerosol base	Concentrate	92.7
	15	Hydrocarbon propellant AP70	Propellant	7.3
15		Total		100.0

Table 6

Submicron / micro emulsion foam – Vitamin D3 (0.02%)

5

Item	Ingredient	Function	%w/w
	Active phase		
1	Diisopropyl adipate	Organic solvent	2.000
2	Vitamin D3	Active ingredient	0.020
	Hydrocarbon oil phase		
3	Macrogol Cetostearyl Ether 12 (Cetareth 12)	Surfactant	3.000
4	Mineral oil (light)	Oil	3.000
5	Tocopherol	Antioxidant	0.002
	First water phase		
6	Water (deionized)	Aqueous solvent	10.000
7	Citric acid	Buffer	q.s.
8	Potassium citrate	Buffer	q.s.
	Second water phase		
9	Water (deionized)	Aqueous solvent	to 100.000
10	Methyl paraben	Preservative	q.s.
	Total		100.000
	Finished product		
Items	Ingredient	Function	%w/w
1 to 10	Aerosol base	Concentrate	92.7
11	Hydrocarbon propellant AP70	Propellant	7.3
	Total		100.0

Table 7**Submicron / micro emulsion foam – Salicylic acid (2%)**

5

Item	Ingredient	Function	%w/w
	Active phase		
1	Diisopropyl adipate	Organic solvent	9.000
2	Salicylic acid	Active ingredient	2.000
	Hydrocarbon oil phase		
3	Macrogol Cetostearyl Ether 12 (Cetareth 12)	Surfactant	6.000
4	Mineral oil (light)	Oil	8.000
	First water phase		
5	Water (deionized)	Aqueous solvent	26.000
6	Citric acid	Buffer	q.s.
7	Potassium citrate	Buffer	q.s.
	Second water phase		
8	Water (deionized)	Aqueous solvent	to 100.000
9	Benzyl alcohol	Preservative	q.s.
	Total		100.000
	Finished product		
Items	Ingredient	Function	%w/w
1 to 9	Aerosol base	Concentrate	92.7
10	Hydrocarbon propellant AP70	Propellant	7.3
	Total		100.0

Table 8

Submicron / micro emulsion foam – Adapalene 0.1%

5

Item	Ingredient	Function	%w/w
	Active phase		
1	Diisopropyl adipate	Organic solvent	3.000
2	Adapalene	Active ingredient	0.100
	Hydrocarbon oil phase		
3	BHT	Antioxidant	0.100
4	Macrogol Cetostearyl Ether 20 (Cetareth 20)	Surfactant	3.000
5	Mineral oil (light)	Oil	4.000
	First water phase		
6	Water (deionized)	Aqueous solvent	15.000
7	Citric acid	Buffer	q.s.
8	Potassium citrate	Buffer	q.s.
	Second water phase		
9	Water (deionized)	Aqueous solvent	to 100.000
10	Benzyl alcohol	Preservative	q.s.
	Total		100.000
	Finished product		
Items	Ingredient	Function	%w/w
1 to 9	Aerosol base	Concentrate	92.7
10	Hydrocarbon propellant AP70	Propellant	7.3
	Total		100.0

Table 9

Submicron / micro emulsion foam - clobetasol propionate (0.05%)

5

Item	Ingredient	Function	%w/w
	Active phase		
1	Diisopropyl adipate	Organic solvent	5.00
2	Clobetasol propionate	Active ingredient	0.05
	Hydrocarbon oil phase		
3	Macrogol Cetostearyl Ether 20 (Cetareth 20)	Surfactant	5.00
4	Mineral oil (light)	Oil	5.00
5	BHT	Antioxidant	0.10
	First water phase		
6	Water (deionized)	Water	20.00
7	Citric acid	Buffer	q.s.
8	Potassium citrate	Buffer	q.s.
	Second water phase		
9	Water (deionized)	Water	to 100%
10	Methyl paraben	Preservative	q.s.
	Total		100.00
	Finished product		
Items	Ingredient	Function	%w/w
1 to 10	Aerosol base	Concentrate	92.7
11	Hydrocarbon propellant AP70	Propellant	7.3
	Total		100.0

Example 4

Skin Penetration Study

5 A skin penetration study was performed, comparing the topical delivery of the tazarotene submicron / micro emulsion foam (base) formulations of Examples 2a and 2b to TAZORACTM cream and gel formulations using *in-vitro* skin distribution assays.

10 Split-thickness skin (~0.50 mm) sections were mounted in flow-through diffusion cells and test agents were applied at a finite dose of 15.63 mg/cm² on 3 skin donors with 5 replicates each.

The tissues were collected at various time points (t=0, 2, 6 hrs), washed, tape stripped, and split into epidermis and dermis. The distribution of tazarotene within the epidermis, dermis, and wash were quantified by LC/MS/MS with a 50 pg/mL LOQ.

15 As shown in Figures 2 and 3, the skin penetration of the tazarotene foam (base) formulations in the epidermis and dermis was similar to the commercial comparators (i.e. TAZORAC[®] cream and gel).

Example 5

20

Compatibility of surfactants with tazarotene

An accelerated stability study (using HPLC) was conducted at 50°C to measure the compatibility of tazarotene with various surfactants. Figures 4 and 5 show that tazarotene was most stable in a hydrophilic ethoxylated fatty alcohol ether surfactant, such as Cetareth – 12, in contrast to various lipophilic surfactants.

Example 6

Measurement of particle size distribution

30 The particle size distribution of the oil phase of the submicron / micro emulsions (aerosol foam base) of the present invention was measured using a Malvern Mastersizer 2000 instrument. Figure 6 illustrates the particle size distribution of the oil phase of a preferred formulation (730/2/1). The mean particle size was 0.097 µm, that is, about 0.1 µm (i.e. about 100 nm).

CLAIMS

What is claimed is:

1. An oil in water emulsion aerosol foam composition comprising an oil phase and a water phase, said composition comprising:
 - i) tazarotene,
 - ii) water in an amount from 65% to 90% by weight,
 - iii) an oil present in an amount of less than 10% by weight,
 - iv) an oil miscible organic solvent,
 - v) a surfactant component comprising a hydrophilic surfactant, in an amount from 0.1% to 10% by weight, and
 - vi) a propellant; andwherein the tazarotene is solubilized in the oil phase of the composition, and wherein the particle size of the oil phase is less than 1000 nm.
2. The aerosol foam composition according to claim 1, further comprising a corticosteroid.
3. The aerosol foam composition according to claim 1, further comprising a vitamin D analogue.
4. The aerosol foam composition according to any one of claims 1 to 3, wherein the composition comprises water in an amount from 70% to 85% by weight.
5. The aerosol foam composition according to any one of claims 1 to 4, wherein the oil is selected from the group consisting of azulene, chamazulene, isoparaffin, linear alpha olefins, cyclohexylidenediphenyl methane, didecene, diethylhexylcyclohexane, eicosane, isododecane, isoeicosane, isohexadecane, longifolene, mineral oil, paraffin, pentahydrosqualene, petrolatum, squalane, squalene, tetradecene, and mixtures thereof.
6. The aerosol foam composition according to claim 5, wherein the oil is mineral oil.
7. The aerosol foam composition according to any one of claims 1 to 6, wherein the oil is present in an amount from 1% to 9% by weight.
8. The aerosol foam composition according to claim 7, wherein the oil is present in an amount from 3% to 8% by weight.

9. The aerosol foam composition according to any one of claims 1 to 8, wherein the oil miscible organic solvent is selected from the group consisting of diisopropyl adipate, isopropyl myristate, octyl dodecanol and caprylic / capric triglyceride, and mixtures thereof.
10. The aerosol foam composition according to any one of claims 1 to 9, wherein the surfactant component comprises a hydrophilic ethoxylated fatty alcohol ether.
11. The aerosol foam composition according to claim 10, wherein the surfactant component comprises a hydrophilic ethoxylated fatty alcohol ether and is substantially free, or free of lipophilic surfactant.
12. The aerosol foam composition according to claim 10 or 11, wherein the hydrophilic ethoxylated fatty alcohol ether is selected from the group consisting of steareth-10, steareth-20, cetareth-6, cetareth-10, cetareth-12, cetareth-15, cetareth-20, cetareth-21, cetareth-22, cetareth-25, cetareth-30, cetareth-31, cetareth-32, cetareth-33, laureth-5, laureth-9, laureth-10, laureth-12, laureth-15, laureth-20, laureth-21, laureth-22, laureth-23, nonoxynol-9, oleth-10, oleth-20, and mixtures thereof.
13. The aerosol foam composition according to claim 12, wherein the hydrophilic ethoxylated fatty alcohol ether is cetareth-12.
14. The aerosol foam composition according to any one of claims 1 to 13, wherein the surfactant component is present in an amount from 1% to 8% by weight.
15. The aerosol foam composition according to claim 14, wherein the surfactant component is present in an amount from 2% to 6% by weight.
16. The aerosol foam composition according to any one of claims 1 to 15, wherein the propellant comprises one or more hydrocarbon propellants.
17. An oil in water emulsion aerosol foam composition comprising an oil phase and a water phase, said composition comprising:
- i) tazarotene;
 - ii) water in an amount from 70% to 85% by weight;
 - iii) an oil present in an amount from 3% to 8% by weight;
 - iv) an oil miscible organic solvent;

- v) a surfactant component comprising a hydrophilic surfactant, in an amount from 2% to 6% by weight; and
 - vi) a hydrocarbon propellant; and
- wherein the tazarotene is solubilized in the oil phase of the composition, and wherein the particle size of the oil phase is less than 1000 nm.

18. The use of a composition according to any one of claims 1 to 17 for the manufacture of a medicament for the treatment of a skin disease, disorder or condition.

19. A process for the preparation of an oil in water submicron or micro emulsion aerosol foam composition, comprising tazarotene, water in an amount from 65% to 90% by weight, an oil present in an amount of less than 10% by weight, an oil miscible organic solvent, a surfactant component comprising a hydrophilic surfactant in an amount from 0.1% to 10% by weight, and a propellant, the process comprising:

- a) admixing the tazarotene, a first aliquot of water, oil, oil miscible organic solvent and surfactant component to form an oil in water emulsion,
- b) heating the oil in water emulsion of step (a) to a phase inversion temperature wherein the oil in water emulsion forms a water in oil emulsion,
- c) cooling the water in oil emulsion to below the phase inversion temperature to form an oil in water submicron or micro emulsion,
- d) adding a second aliquot of water to cool the oil in water submicron or micro emulsion, and
- e) actuating a sample of the oil in water submicron or micro emulsion with a propellant to form an oil in water submicron or micro emulsion aerosol foam.

20. The process according to claim 19, wherein the first aliquot of water comprises 10% to 40% by weight of the total water content of the submicron or micro emulsion aerosol foam and the second aliquot of water comprises 90% to 60% by weight of the total water content of the submicron or micro emulsion aerosol foam.