

Imidazopyrazinones as PDE1 inhibitors

FIELD OF THE INVENTION

The present invention provides compounds that are PDE1 enzyme inhibitors and their use as a medicament, in particular for the treatment of neurodegenerative disorders and psychiatric disorders. The present invention also provides pharmaceutical compositions comprising compounds of the invention and methods of treating disorders using the compounds of the invention.

BACKGROUND OF THE INVENTION

Throughout this application, various publications are referenced in full. The disclosures of these publications are hereby incorporated by reference into this application to describe more fully the state of the art to which this invention pertains.

The second messenger cyclic Nucleotides (cNs), cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP) play a major role in intracellular signal transduction cascade, by regulating cN-dependent protein kinases (PKA and PKG), EPACs (Exchange Protein Activated by cAMP), phosphoprotein phosphatases, and/or cN-gated cation channels. In neurons, this includes the activation of cAMP- and cGMP-dependent kinases and subsequent phosphorylation of proteins involved in acute regulation of synaptic transmission as well as in neuronal differentiation and survival. Intracellular concentrations of cAMP and cGMP are strictly regulated by the rate of biosynthesis by cyclases and by the rate of degradation by phosphodiesterases (PDEs, EC 3.1.4.17). PDEs are bimetallic hydrolases that inactivate cAMP/cGMP by catalytic hydrolysis of the 3'-ester bond, forming the inactive 5'-monophosphate. Since PDEs provide the only means of degrading the cyclic nucleotides cAMP and cGMP in cells, PDEs play an essential role in cyclic nucleotide signalling. The catalytic activities of PDEs provide for breakdown of cNs over a spectrum of cN-concentrations in all cells, and their varied regulatory mechanisms provide for integration and crosstalk with myriads of signalling pathways. Particular PDEs are targeted to discrete compartments within cells where they control cN level and sculpt microenvironments for a variety of cN signalosomes (Sharron H. Francis, Mitsi A. Blount, and Jackie D. Corbin. *Physiol Rev* 2011, 91: 651-690).

On the basis of substrate specificity, the PDE families can be divided into three groups: 1) The cAMP-specific PDEs, which include PDE4, PDE7, and PDE8, 2) the cGMP-selective

enzymes PDE5 and PDE9, and 3) the dual-substrate PDEs, PDE1, PDE2, PDE3, as well as PDE10 and PDE11.

Previously named calmodulin-stimulated PDE (CaM-PDE), PDE1 is unique in that it is Ca^{2+} -dependently regulated via calmodulin (CaM, a 16 kDa Ca^{2+} -binding protein) complexed with four Ca^{2+} (for review, Sharron H. Francis, Mitsi A. Blount, and Jackie D. Corbin. *Physiol Rev* 2011, 91: 651–690). Thus, PDE1 represents an interesting regulatory link between cyclic nucleotides and intracellular Ca^{2+} . The PDE1 family is encoded by three genes: PDE1A (mapped on human chromosome 2q32), PDE1B (human chromosome location, hcl: 12q13) and PDE1C (hcl: 7p14.3). They have alternative promoters and give rise to a multitude of proteins by alternative splicing which differ in their regulatory properties, substrate affinities, specific activities, activation constants for CaM, tissue distribution and molecular weights. More than 10 human isoforms are identified. Their molecular weights vary from 58 to 86 kDa per monomer. The N-terminal regulatory domain that contains two Ca^{2+} /CaM binding domains and two phosphorylation sites differentiate their corresponding proteins and modulate their biochemical functions. PDE1 is a dual substrate PDE and the PDE1C-subtype has equal activity towards cAMP and cGMP ($K_m \approx 1\text{--}3 \mu\text{M}$), whereas the subtypes PDE1A and PDE1B have a preference for cGMP (K_m for cGMP $\approx 1\text{--}3 \mu\text{M}$ and for cAMP $\approx 10\text{--}30 \mu\text{M}$).

The PDE1 subtypes are highly enriched in the brain and located especially in the striatum (PDE1B), hippocampus (PDE1A) and cortex (PDE1A) and this localization is conserved across species (Amy Bernard et al. *Neuron* 2012, 73, 1083–1099). In the cortex, PDE1A is present mainly in deep cortical layers 5 and 6 (output layers), and used as a specificity marker for the deep cortical layers. PDE1 inhibitors enhance the levels of the second messenger cNs leading to enhanced neuronal excitability.

Thus, PDE1 is a therapeutic target for regulation of intracellular signaling pathways, preferably in the nervous system and PDE1 inhibitors can enhance the levels of the second messengers cAMP/cGMP leading to modulation of neuronal processes and to the expression of neuronal plasticity-related genes, neurotrophic factors, and neuroprotective molecules. These neuronal plasticity enhancement properties together with the modulation of synaptic transmission make PDE1 inhibitors good candidates as therapeutic agents in many neurological and psychiatric conditions. The evaluation of PDE1 inhibitors in animal models (for reviews see e.g. Blokland et al. *Expert Opinion on Therapeutic Patents* (2012), 22(4), 349–354; and Medina, A. E. *Frontiers in Neuropharmacology* (2011), 5(Feb.), 21) has suggested the potential for the therapeutic use of PDE1 inhibitors in neurological disorders, like e.g. Alzheimer's, Parkinson's and Huntington's Diseases and in psychiatric disorders like

e.g. Attention Deficit hyperactivity Disorder (ADHD), restless leg syndrome, depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS). There have also been patent applications claiming that PDE1 inhibitors are useful in diseases that may be alleviated by the enhancement of progesterone-signaling such as female sexual dysfunction (e.g. WO 2008/070095).

WO 2013/053690 A1 discloses imidazopyrazinones that are inhibitors of the PDE9 enzyme.


The compounds of the invention may offer alternatives to current marketed treatments for neurodegenerative and/or psychiatric disorders, treatments which are not efficacious in all patients. Hence, there remains a need for alternative methods of treatment of such diseases.

SUMMARY OF THE INVENTION

PDE1 enzymes are expressed in the Central Nervous System (CNS), making this gene family an attractive source of new targets for the treatment of psychiatric and neurodegenerative disorders.

The objective of the present disclosure is to provide compounds that are PDE1 inhibitors, and as such are useful to treat neurodegenerative disorders and psychiatric disorders. Preferably, said compounds are at least a ten-fold stronger as PDE1 inhibitors than as PDE9 inhibitors in order to prevent potentially unwanted effects associated with PDE9 inhibition.

In an aspect, there is provided a compound according to formula (I)



(I)

wherein

n is 0 or 1;

q is 0 or 1;

R1 is selected from the group consisting of benzyl, indanyl, indoline and 5-membered heteroaryls; all of which can be substituted with a substituent selected from the group consisting of halogen and C₁-C₃ alkyl; or

5 R1 is selected from the group consisting of saturated monocyclic rings containing 4-6 carbon atoms and 1-2 nitrogen atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine and sulfonamide; or

R1 is selected from the group consisting of lactams containing 4-6 carbon atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

10 R1 is selected from the group consisting of bicyclic ethers such as, 7-oxabicyclo[2.2.1]heptane; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

15 R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl and tetrahydropyranyl; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine, hydroxy, cyano or methoxy; or

R1 is a linear or branched C₁-C₃ alkyl, which is substituted with a substituent selected from phenyl and 5-membered heteroaryl, wherein said 5-membered heteroaryl can be substituted with one or more C₁-C₃ alkyls; or

20 R1 is selected from the group consisting of morpholine, tetrahydrofuran-3-amine, hexahydro-2H-furo[3,2-b]pyrrole and homomorpholine; all of which can be substituted with one or more substituents selected from the group consisting of C₁-C₃ alkyl;

25 R2 is selected from the group consisting of hydrogen, linear or branched C₁-C₈ alkyl, phenyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, benzo[d][1,3]dioxolyl, tetrahydrofuranyl and tetrahydropyranyl; or

R2 is phenyl or pyridyl substituted with one or more substituents selected from the group consisting of hydroxyl, amino, cyano, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₃-C₅ cycloalkoxy, C₃-C₅ cycloalkyl-methoxy, C₁-C₃ fluoroalkoxy, and -NC(O)CH₃; or


R2 is a 5-membered heteroaryl which can be substituted with one or more C₁-C₃ alkyl;

R3 is selected from the group consisting of halogen, C₁-C₅ alkyl, C₃-C₅ cycloalkyl and phenyl; or

R3 is selected from the group consisting of phenyl substituted one or more times with C₁-C₃ alkyl; methyl substituted one, two or three times with fluorine; ethyl substituted one, two or three times with fluorine;

R4 is hydrogen.

In another aspect, there is provided a pharmaceutical composition comprising therapeutically effective amount of a compound according to formula (I), and one or more pharmaceutically acceptable carriers, diluents and excipients



(I)

wherein

n is 0 or 1;

q is 0 or 1;

R1 is selected from the group consisting of benzyl, indanyl, indoline and 5-membered heteroaryls; all of which can be substituted with a substituent selected from the group consisting of halogen and C₁-C₃ alkyl; or

R1 is selected from the group consisting of saturated monocyclic rings containing 4-6 carbon atoms and 1-2 nitrogen atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine and sulfonamide; or

R1 is selected from the group consisting of lactams containing 4-6 carbon atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

R1 is selected from the group consisting of bicyclic ethers such as, 7-oxabicyclo[2.2.1]heptane; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

5 R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl and tetrahydropyranyl; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine, hydroxy, cyano or methoxy; or

10 R1 is a linear or branched C₁-C₃ alkyl, which is substituted with a substituent selected from phenyl and 5-membered heteroaryl, wherein said 5-membered heteroaryl can be substituted with one or more C₁-C₃ alkyls; or

R1 is selected from the group consisting of morpholine, tetrahydrofuran-3-amine, hexahydro-2H-furo[3,2-b]pyrrole and homomorpholine; all of which can be substituted with one or more substituents selected from the group consisting of C₁-C₃ alkyl;

15 R2 is selected from the group consisting of hydrogen, linear or branched C₁-C₈ alkyl, phenyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, benzo[d][1,3]dioxolyl, tetrahydrofuranyl and tetrahydropyranyl; or

R2 is phenyl or pyridyl substituted with one or more substituents selected from the group consisting of hydroxyl, amino, cyano, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₃-C₅ cycloalkoxy, C₃-C₅ cycloalkyl-methoxy, C₁-C₃ fluoroalkoxy, and -NC(O)CH₃; or

20 R2 is a 5-membered heteroaryl which can be substituted with one or more C₁-C₃ alkyl;

R3 is selected from the group consisting of hydrogen, halogen, C₁-C₅ alkyl, C₃-C₅ cycloalkyl and phenyl; or

25 R3 is selected from the group consisting of phenyl substituted one or more times with C₁-C₃ alkyl; methyl substituted one, two or three times with fluorine; ethyl substituted one, two or three times with fluorine;

R4 is hydrogen;

and tautomers and pharmaceutically acceptable addition salts thereof;

with the proviso that R2 and R3 cannot be hydrogen at the same time;


with the proviso, that the compound of formula (I) is not one of the three following compounds:

3-methyl-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

7-butyl-3-methylimidazo[1,5-a]pyrazin-8(7*H*)-one; and

5 7-(4-methoxybenzyl)-3-methylimidazo[1,5-a]pyrazin-8(7*H*)-one.

Accordingly, the present disclosure relates to compounds of formula (I)



(I)

wherein

n is 0 or 1;

q is 0 or 1;

R1 is selected from the group consisting of benzyl, indanyl, indoline and 5-membered heteroaryls; all of which can be substituted with a substituent selected from the group consisting of halogen and C₁-C₃ alkyl; or

R1 is selected from the group consisting of saturated monocyclic rings containing 4-6 carbon atoms and 1-2 nitrogen atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine and sulfonamide; or

R1 is selected from the group consisting of lactams containing 4-6 carbon atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

R1 is selected from the group consisting of bicyclic ethers such as, 7-oxabicyclo[2.2.1]heptane; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl and tetrahydropyranyl; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine, hydroxy, cyano or methoxy; or

R1 is a linear or branched C₁-C₃ alkyl, which is substituted with a substituent selected from phenyl and 5-membered heteroaryl, wherein said 5-membered heteroaryl can be substituted with one or more C₁-C₃ alkyls; or

R1 is selected from the group consisting of morpholine, tetrahydrofuran-3-amine, hexahydro-2H-furo[3,2-b]pyrrole and homomorpholine; all of which can be substituted with one or more substituents selected from the group consisting of C₁-C₃ alkyl;

R2 is selected from the group consisting of hydrogen, linear or branched C₁-C₈ alkyl, phenyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, benzo[d][1,3]dioxolyl, tetrahydrofuranyl and tetrahydropyranyl; or

R2 is phenyl or pyridyl substituted with one or more substituents selected from the group consisting of hydroxyl, amino, cyano, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₃-C₅ cycloalkoxy, C₃-C₅ cycloalkyl-methoxy, C₁-C₃ fluoroalkoxy, and -NC(O)CH₃; or

R2 is a 5-membered heteroaryl which can be substituted one or more times with C₁-C₃ alkyl;

R3 is selected from the group consisting of hydrogen, halogen, C₁-C₅ alkyl, C₃-C₅ cycloalkyl and phenyl; or

- 5 R3 is selected from the group consisting of phenyl substituted one or more times with C₁-C₃ alkyl; methyl substituted one, two or three times with fluorine; ethyl substituted one, two or three times with fluorine;

R4 is hydrogen;

and tautomers and pharmaceutically acceptable addition salts thereof;

- 10 with the proviso that R2 and R3 cannot be hydrogen at the same time;

with the proviso, that the compound of formula (I) is not one of the three following compounds:

3-methyl-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

7-butyl-3-methylimidazo[1,5-a]pyrazin-8(7*H*)-one; and

- 15 7-(4-methoxybenzyl)-3-methylimidazo[1,5-a]pyrazin-8(7*H*)-one.

Reference to Compound I includes the free base of Compound I, pharmaceutically acceptable salts of Compound I, such as acid addition salts of Compound I, racemic mixtures of Compound I, or the corresponding enantiomer and/or optical isomer of Compound I, and polymorphic and amorphous forms of Compound I as well as tautomeric

- 20 forms of Compound I. Furthermore, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of present disclosure.

In one embodiment, the present disclosure relates to a compound according to formula (I) for use in therapy.

25

In one embodiment, the present disclosure relates to a compound according to formula (I), for use in the treatment of a neurodegenerative disorder, selected from the group consisting of Alzheimer's Disease, Parkinson's Disease and Huntington's Disease or for the treatment of a psychiatric disorder such as Attention Deficit hyperactivity Disorder (ADHD),

depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS), or another brain disease like restless leg syndrome.

In one embodiment, the present disclosure relates to a pharmaceutical composition comprising a compound according formula (I), and one or more pharmaceutically acceptable carrier or excipient.

5 In one embodiment, the present disclosure relates to a method for the treatment of a neurodegenerative disorder, selected from the group consisting of Alzheimer's Disease, Parkinson's Disease and Huntington's Disease or for the treatment of a psychiatric disorder such as Attention Deficit hyperactivity Disorder (ADHD), depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS), or
10 another brain disease like restless leg syndrome, which method comprises the administration of a therapeutically effective amount of a compound according to formula (I) to a patient in need thereof.

DETAILED DESCRIPTION OF THE INVENTION


EMBODIMENTS OF THE INVENTION

15 The following notation is applied: an embodiment of the invention is identified as E_i, where i is an integer indicating the number of the embodiment. An embodiment E_i' specifying a specific embodiment a previously listed embodiment E_i is identified as E_i'(E_i), e.g. E₂(E₁) means "in an embodiment E₂ of embodiment E₁".

Where an embodiment is a combination of two embodiments the notation is similarly
20 E_i''(E_i' and E_i), e.g. E₃(E₂ and E₁) means "in an embodiment E₃ of any of embodiments E₂ and E₁"

Where an embodiment is a combination of more than two embodiments the notation is similarly E_i''' (E_i, E_i' and E_i''), e.g. E₄(E₁, E₂ and E₃) means "in an embodiment E₄ of any of embodiments E₁, E₂ and E₃"

25 In a first embodiment E₁, the present invention relates to compounds of formula (I)



(I)

wherein

n is 0 or 1;

5 q is 0 or 1;

R1 is selected from the group consisting of benzyl, indanyl, indoline and 5-membered heteroaryls; all of which can be substituted with a substituent selected from the group consisting of halogen and C₁-C₃ alkyl; or

10 R1 is selected from the group consisting of saturated monocyclic rings containing 4-6 carbon atoms and 1-2 nitrogen atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine and sulfonamide; or

R1 is selected from the group consisting of lactams containing 4-6 carbon atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

15 R1 is selected from the group consisting of bicyclic ethers such as, 7-oxabicyclo[2.2.1]heptane; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

20 R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl and tetrahydropyranyl; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine, hydroxy, cyano or methoxy; or

R1 is a linear or branched C₁-C₃ alkyl, which is substituted with a substituent selected from phenyl and 5-membered heteroaryl, wherein said 5-membered heteroaryl can be substituted with one or more C₁-C₃ alkyls; or

25 R1 is selected from the group consisting of morpholine, tetrahydrofuran-3-amine, hexahydro-2H-furo[3,2-b]pyrrole and homomorpholine; all of which can be substituted with one or more substituents selected from the group consisting of C₁-C₃ alkyl;

30 R2 is selected from the group consisting of hydrogen, linear or branched C₁-C₈ alkyl, phenyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, benzo[d][1,3]dioxolyl, tetrahydrofuranyl and tetrahydropyranyl; or

R2 is phenyl or pyridyl substituted with one or more substituents selected from the group consisting of hydroxyl, amino, cyano, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₃-C₅ cycloalkoxy, C₃-C₅ cycloalkyl-methoxy, C₁-C₃ fluoroalkoxy, and -NC(O)CH₃; or

R2 is a 5-membered heteroaryl which can be substituted with one or more C₁-C₃ alkyl;

- 5 R3 is selected from the group consisting of hydrogen, halogen, C₁-C₅ alkyl, C₃-C₅ cycloalkyl and phenyl; or

R3 is selected from the group consisting of phenyl substituted one or more times with C₁-C₃ alkyl; methyl substituted one, two or three times with fluorine; ethyl substituted one, two or three times with fluorine;

- 10 R4 is hydrogen;

and tautomers and pharmaceutically acceptable addition salts thereof;

with the proviso that R2 and R3 cannot be hydrogen at the same time;

with the proviso, that the compound of formula (I) is not one of the three following compounds:

- 15 3-methyl-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-a]pyrazin-8(7H)-one;
7-butyl-3-methylimidazo[1,5-a]pyrazin-8(7H)-one; and
7-(4-methoxybenzyl)-3-methylimidazo[1,5-a]pyrazin-8(7H)-one.

E2(E1)

- 20 n is 0 or 1;

q is 0 or 1;

R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl, and tetrahydropyranyl;

- 25 R2 is selected from the group consisting of, linear or branched C₁-C₈ alkyl, phenyl, and saturated monocyclic C₃-C₈ cycloalkyl; or

R2 is selected from the group consisting of phenyl substituted with one or more times with one or more substituents selected from the group consisting of halogen, C₁-C₃ alkyl and methoxy;

E28(E1 to E22) a method of treating a subject suffering from neurodegenerative disorder, selected from the group consisting of Alzheimer's Disease, Parkinson's Disease and Huntington's Disease or for the treatment of a psychiatric disorder such as Attention Deficit hyperactivity Disorder (ADHD), depression, anxiety, narcolepsy, cognitive impairment and
 5 cognitive impairment associated with schizophrenia (CIAS) or another brain disease like restless leg syndrome, which method comprises administering to said subject an amount of a compound of formula I effective in inhibiting PDE1.

E29 (E1 to E22) Use of a compound of any of embodiments (E1) to (E21) in the manufacture of a medicament for use in the treatment of a neurodegenerative disorder, selected from the
 10 group consisting of Alzheimer's Disease, Parkinson's Disease and Huntington's Disease or for the treatment of a psychiatric disorder such as Attention Deficit hyperactivity Disorder (ADHD), depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS) or another brain disease like restless leg syndrome.

15 DEFINITIONS

PDE1 ENZYMES

The PDE1 isozyme family includes numerous splice variant PDE1 isoforms. It has three subtypes, PDE1A, PDE1B and PDE1C which divide further into various isoforms. In the context of the present invention PDE1 and PDE1 enzymes are synonymous and refer to
 20 PDE1A, PDE1B and PDE1C enzymes as well as their isoforms unless otherwise specified.

SUBSTITUENTS

As used in the context of the present invention, the terms "halo" and "halogen" are used interchangeably and refer to fluorine, chlorine, bromine or iodine.

25 A given range may interchangeably be indicated with "-" (dash) or "to", e.g. the term "C₁-C₃ alkyl" is equivalent to "C₁ to C₃ alkyl".

The terms "C₁-C₃ alkyl", "C₁-C₄ alkyl", "C₁-C₅ alkyl", "C₁-C₆ alkyl", "C₁-C₇ alkyl" and "C₁-C₈ alkyl" refer to a linear (i.e. unbranched) or branched saturated hydrocarbon having from one up to eight carbon atoms, inclusive. Examples of such groups include, but are not limited to,
 30 methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, 2-methyl-1-butyl, n-hexyl, n-heptyl and n-octyl.

The term saturated monocyclic C₃-C₈ cycloalkyl refers to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. The term C₃-C₅ cycloalkyl refers to cyclopropyl, cyclobutyl and cyclopentyl.

The term "C₁-C₃ alkoxy" refers to a moiety of the formula -OR', wherein R' indicates C₁-C₃ alkyl as defined above. The term "C₃-C₅ cycloalkoxy" refers to a moiety of the formula -OR', wherein R' indicates C₃-C₅ cycloalkyl as defined above. The term "C₃-C₅ cycloalkyl-methoxy" refers to a moiety of the formula -OCH₂R', wherein R' indicates C₃-C₅ cycloalkyl as defined above. C₁-C₃ fluoroalkoxy refers to a C₁-C₃ alkoxy substituted with one or more fluorine.

5-membered heteroaryls are defined as 5 membered aromatic rings containing at least one atom selected from nitrogen, sulfur and oxygen. Examples include, but are not limited to thiazole, thiophene and isoxazole.

The term "lactams containing 4-6 carbon atoms" refers to pyrrolidin-2-one, piperidin-2-one or azepan-2-one

15 ISOMERIC FORMS

Where compounds of the present invention contain one or more chiral centers reference to any of the compounds will, unless otherwise specified, cover the enantiomerically or diastereomerically pure compound as well as mixtures of the enantiomers or diastereomers in any ratio.

20 The above also applies where compounds of the invention contain more than two chiral centers.

PDE1 INHIBITORS AND PDE9 INHIBITORS

In the context of the present invention a compound is considered to be a PDE1 inhibitor if the amount required to reach the IC₅₀ level of one or more of the three PDE1 isoforms is 10 micro molar or less, preferably less than 9 micro molar, such as 8 micro molar or less, such as 7 micro molar or less, such as 6 micro molar or less, such as 5 micro molar or less, such as 4 micro molar or less, such as 3 micro molar or less, more preferably 2 micro molar or less, such as 1 micro molar or less, in particular 500 nM or less. In preferred embodiments the required amount of PDE1 inhibitor required to reach the IC₅₀ level of PDE1B is 400nM or

This invention further provides a method of treating a neurodegenerative disorder in a mammal, including a human, which method comprises administering to said mammal an amount of a compound of formula (I) effective in inhibiting PDE1.

This invention also provides a method of treating a subject suffering from a psychiatric disorder, which method comprises administering to the subject a therapeutically effective amount of a compound of formula (I). Examples of psychiatric disorders that can be treated according to the present invention include Attention Deficit hyperactivity Disorder (ADHD), depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS).

This invention also provides a method of treating a subject suffering from another brain disorder such as restless leg syndrome.

Further, the invention is directed to the use of a compound of formula (I) in the manufacture of a medicament for the treatment of a neurodegenerative disorder, such as Alzheimer's Disease, Parkinson's Disease and Huntington's Disease or for the treatment of a psychiatric disorder such as Attention Deficit hyperactivity Disorder (ADHD), depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS).

Further, the invention is directed to the use of a compound of formula (I) in the manufacture of a medicament for the treatment of another brain disease, such as restless leg syndrome

The invention is also directed to a compound of formula (I) for use as a medicine. In a specific embodiment the compound of formula (I) is for use in the treatment of a neurodegenerative disorder, such as Alzheimer's Disease, Parkinson's Disease and Huntington's Disease or for the treatment of a psychiatric disorder such as Attention Deficit hyperactivity Disorder (ADHD), depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS) or for the treatment of another brain disease like restless leg syndrome.

All references, including publications, patent applications and patents, cited herein are hereby incorporated by reference in their entirety and to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety (to the maximum extent permitted by law).

Headings and sub-headings are used herein for convenience only, and should not be construed as limiting the invention in any way.

COMPOUNDS OF THE INVENTION

Table 1: Compounds of the invention

Example	Name	PDE1A, IC ₅₀ (nM)	PDE1B, IC ₅₀ (nM)	PDE1C, IC ₅₀ (nM)	% inhibition of PDE9 at 10 microM
1	7-(3-Fluorobenzyl)-3-propylimidazo[1,5-a]pyrazin-8(7H)-one	801	790	300	-12
2	6-Benzyl-7-(3-fluorobenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	51	14	11	-14
3	6-Benzyl-7-(cyclohexylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	8	12	6	-6
4	7-(Cyclohexylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	11	25	7	-5
5	7-(3-Fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	108	120	14	18
6	3-Cyclopropyl-7-(3-fluorobenzyl)imidazo[1,5-a]pyrazin-8(7H)-one	68% (at 2 μ M)	81% (at 2 μ M)	290	25
7	7-(Cyclopentylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one	60% (at 2 μ M)	85% (at 2 μ M)	230	1
8	7-(Cyclohexylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one	252	67	100	9
9	7-(3-Fluorobenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	68% (at 2 μ M)	90% (at 2 μ M)	170	2
10	7-(Cyclopentylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	73% (at 2 μ M)	88% (at 2 μ M)	79	8
11	7-(Cyclohexylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	113	72	43	18

25	6-Methyl-7-(3-methylbenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	63	57	9	-42
26	7-(3-fluorobenzyl)-6-methyl-3-(4-methyltetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	234	218	47	-2
27	4-(7-(3-fluorobenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)tetrahydro-2H-pyran-4-carbonitrile	603	699	103	5
28	7-(3-fluorobenzyl)-3-(4-methoxytetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one	663	737	72	13
29	7-(3-fluorobenzyl)-3-(4-fluorotetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one	493	249	50	16
30, isomer 1	7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-2-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 1	20% (at 1 μ M)	615	225	-11
30, isomer 2	7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-2-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 2	27% (at 1 μ M)	40% (at 1 μ M)	215	28
31, isomer 1	7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 1	337	106	27	-1
31, isomer 2	7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 2	347	138	31	5
32, isomer 1	7-(3-fluorobenzyl)-6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 1	257	122	22	-1
32, isomer 2	7-(3-fluorobenzyl)-6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 2	401	170	40	16
33	7-(3-fluorobenzyl)-6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one	94	52	7	-2
34	3-(2,2-difluorocyclopropyl)-7-(3-fluorobenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one	331	359	93	0

35, isomer 1	7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 1	193	75	19	-11
35, isomer 2	7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 2	364	166	41	15
35, isomer 3	7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 3	82	18	8	0
35, isomer 4	7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 4	43% (at 1 μ M)	360	85	5
36, isomer 1	7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 1	152	59	15	3
36, isomer 2	7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 2	282	107	38	11
36, isomer 3	7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 3	77	21	7	-10
36, isomer 4	7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 4	51	520	40% (at 1 μ M)	11
37, isomer 1	7-(3-fluorobenzyl)-3-(<i>cis</i> -2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 1	1140	1180	345	-2
37, isomer 2	7-(3-fluorobenzyl)-3-(<i>cis</i> -2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 2	149	164	55	13
38, isomer 1	7-(3-fluorobenzyl)-3-(<i>trans</i> -2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 1	582	601	118	-1
38, isomer 2	7-(3-fluorobenzyl)-3-(<i>trans</i> -2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, stereoisomer 2	667	667	153	2
39	7-(4-cyclopropoxybenzyl)-6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	413	118	364	23

40	7-(4-(difluoromethoxy)benzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	73	27	52	-25
41	6-methyl-3-(tetrahydro-2H-pyran-4-yl)-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-a]pyrazin-8(7H)-one	27	18	52	-22
42	7-(4-(cyclopropylmethoxy)benzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	50% (at 1 μ M)	155	55% (at 1 μ M)	15
43	7-benzyl-6-ethyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	105	32	9	6
44	6-ethyl-7-(4-methoxybenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	40	5	16	18
45	3-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)benzonitrile	64 (at 2 μ M)	54% (at 1 μ M)	149	6
46	4-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)benzonitrile	809	58% (at 1 μ M)	9% (at 1 μ M)	14
47	N-(4-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)phenyl)acetamide	0% (at 1 μ M)	388	10% (at 1 μ M)	10
48	7-(4-chloro-3-methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	-	54% (at 1 μ M)	34% (at 1 μ M)	10
49	7-(2-ethylbenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	44 (at 2 μ M)	60% (at 1 μ M)	173	1
50	7-(benzo[d][1,3]dioxol-5-ylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	50	16	38	12
51	7-(3-chloro-4-methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	199	36	46	1
52	7-(4-aminobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one	0% (at 1 μ M)	388	10% (at 1 μ M)	-3

53	7-(4-hydroxybenzyl)-6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	401	35	84	-41
54	6-ethyl-7-(3-fluorobenzyl)-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	62	53	12	23
55, isomer 1	7-(4-methoxybenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one, stereoisomer 1	20	10	17	18
55, isomer 2	7-(4-methoxybenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one, stereoisomer 2	86	64	51	-13
55, isomer 3	7-(4-methoxybenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one, stereoisomer 3	13	3	8	27
55, isomer 4	7-(4-methoxybenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one, stereoisomer 4	137	208	131	1
56	7-(4-methoxybenzyl)-6-methyl-3-propylimidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	65	21	43	7
57	7-((6-methoxypyridin-3-yl)methyl)-6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	45	26	84	-5
58	6,7-dimethyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	58% (at 10 μ M)	1767	44% (at 10 μ M)	36
59	7-ethyl-6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	2333	496	1737	-13
60	6-methyl-7-propyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	810	221	423	-13
61	7-isopropyl-6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	4% (at 10 μ M)	50% (at 10 μ M)	47% (at 10 μ M)	4
62	7-isopentyl-6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	1078	558	93	7

74	3-(6-chloro-2,3-dihydro-1 <i>H</i> -inden-1-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	5	2	3	-19
75	7-(4-methoxybenzyl)-6-methyl-3-(3-methyl-5-oxopyrrolidin-3-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	236	297	360	8
76	3-(1-methoxy-2-methylpropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	122	44	126	-14
77	3-isopropyl-7-(4-methoxybenzyl)-6-methylimidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	13	7	20	nd
78	6-methyl-7-((2-methylthiazol-4-yl)methyl)-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	1691	641	251	6
79	6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)-7-(thiophen-3-ylmethyl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	228	112	15	14
80	6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)-7-(thiazol-4-ylmethyl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	72% (at 10 μ M)	1310	496	18
81	7-((3,5-dimethylisoxazol-4-yl)methyl)-6-methyl-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	27% (at 10 μ M)	36% (at 10 μ M)	2305	36
82	6-methyl-7-((5-methylisoxazol-3-yl)methyl)-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	67% (at 10 μ M)	1498	1034	7
83	6-methyl-7-((3-methylisoxazol-5-yl)methyl)-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	4444	2289	1606	-4
84	3-(2,6-dimethyltetrahydro-2 <i>H</i> -pyran-4-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	627	103	337	-5
85	7-(cyclohexylmethyl)-6-methyl-3-propylimidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	120	32	32	7
86	3-(2-hydroxypropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5- <i>a</i>]pyrazin-8(7 <i>H</i>)-one	193	118	214	-11

87	3-(2-fluoropropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	108	124	276	10
88	7-(4-methoxybenzyl)-6-methyl-3-(7-oxazepan-4-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	484	308	548	-32
89	7-(4-methoxybenzyl)-6-methyl-3-(5-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	65% (at 10 μ M)	128	303	1
90	7-(4-methoxybenzyl)-6-methyl-3-(1-(4-methylthiazol-2-yl)ethyl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	34	15	23	10
91	3-(7-(4-methoxybenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)-3-methylpyrrolidine-1-sulfonamide	181	185	417	20
92	6-(cyclopentylmethyl)-7-(4-methoxybenzyl)-3-(tetrahydro-2 <i>H</i> -pyran-4-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	71	11	79	57
93	3-(morpholino)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	96	52	99	23
94	7-(4-methoxybenzyl)-6-methyl-3-((tetrahydrofuran-3-yl)amino)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	2516	465	1231	-5
95	(<i>R</i>)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	258	162	218	9
96	(<i>S</i>)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	48	38	51	nd
97	7-(4-methoxybenzyl)-6-methyl-3-(1,4-oxazepan-4-yl)imidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	202	82	128	-8
98	3-(2,2-dimethylmorpholino)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one	287	102	135	4
99, isomer 1	7-(3-fluorobenzyl)-3-(hexahydro-4 <i>H</i> -furo[3,2- <i>b</i>]pyrrol-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7 <i>H</i>)-one, isomer 1	241	134	27	9

99, isomer 2	7-(3-fluorobenzyl)-3-(hexahydro-4H-furo[3,2-b]pyrrol-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one, isomer 2	47% (at 10 μ M)	760	119	6
100, isomer 1	7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 1	306	223	43	11
100, isomer 2	7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 2	333	93	33	-21

nd means "not determined"

Table 1 lists the IC₅₀ value for inhibition of PDE1 by the compounds of the invention. The IC₅₀ value refers to the concentration (nM) of the compound required to reach 50% inhibition of the PDE1 enzyme at the specified substrate concentration.


- 5 For certain compounds, the inhibition of PDE is listed as % inhibition at a certain concentration.

For comparative purpose, the table also lists % inhibition of PDE9 at 10 μ M.

PDE1 and PDE9 assays are described in the Experimental Section.

10 EXPERIMENTAL SECTION

PREPARATION OF THE COMPOUNDS OF THE INVENTION




(I)

- 15 The compounds of formula (I) may be prepared by methods described below, together with synthetic methods known in the art of organic chemistry, or modifications that are familiar to those of ordinary skill in the art. The starting materials used herein are available commercially or may be prepared by routine methods known in the art, such as those method described in standard reference books such as "Compendium of Organic Synthetic

limited to using; a boronic acid ester, potassium carbonate as the base, a mixture of 1,4-dioxane and water as the solvent and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (Pd(dppf)Cl_2) as the catalyst. Imidazo[1,5-a]pyrazin-8(7H)-ones of formula II are prepared by treating compounds of formula III with an acid exemplified but not limited to hydrochloric acid in a mixture of solvents such as water and methanol.

Imidazo[1,5-a]pyrazin-8(7H)-ones of formula I are prepared by alkylation of II with an alkylating reagent exemplified by but not limited to alkylbromide using a base exemplified by but not limited to potassium carbonate in a solvent such as dimethylformamide.

Method 3:



In brief, compounds of the invention can be prepared from the commercial available methyl 1H-imidazole-5-carboxylate VI (CAS: 17325-26-7). Reaction of methyl 1H-imidazole-5-carboxylate VI with an α -halogenated ketone exemplified but not limited to an α -chloroketone, under the influence of a base exemplified but not limited to potassium carbonate in a solvent such as acetone yields the imidazole V. Treating imidazole V with a brominating reagent exemplified but not limited to N-bromosuccinimide (NBS) in the presence of a radical initiator exemplified by but not limited to azobisisobutyronitrile (AIBN) gives imidazole IV. Compounds of the formula III are formed by treatment imidazole IV with ammonium acetate in a solvent such as 1,4-dioxane. Compounds of the formula II can be prepared from intermediate III using standard cross-coupling reaction conditions exemplified by but not limited to a Suzuki-Miyaura cross-coupling reaction. Such conditions for the cross-coupling reaction are exemplified by but not limited to using; a boronic acid ester, potassium carbonate as the base, a mixture of 1,4-dioxane and water as the solvent and Pd(dppf)Cl_2 as the catalyst. In some examples R_1 contains an unsaturated carbon-carbon bond which can be reduced by hydrogenation under conditions known to the person skilled in the art.

Imidazo[1,5-a]pyrazin-8(7H)-ones of formula I are prepared by alkylation of II with an

alkylating reagent exemplified by but not limited to an alkyl bromide using a base exemplified but not limited to potassium carbonate in a solvent such as dimethylformamide.

Analytical Methods

Analytical LC-MS data were obtained using the methods identified below.

- 5 **Method 1:** An Agilent 1200 LCMS system with ELS detector was used. Column: XBridge ShieldRP18, 5µm, 50x2.1mm; Column temperature: 40 °C; Solvent system: A = water/NH₃*H₂O (99.95:0.05) and B = acetonitrile; Method: Linear gradient elution with A:B = 99:1 to 0:100 in 3.4 minutes and with a flow rate of 0.8 mL/min.

- 10 **Method 2:** A Shimadzu 20 MS instrument equipped with atmospheric pressure photo ionisation ion source and a Shimadzu LC-20AB system was used. Column: MERCK, RP-18e 25-2mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9625:0.0375) and B = acetonitrile /trifluoroacetic acid (99.981:0.019); Method: A linear gradient elution A:B = 95:5 to A:B=5:95 in 0.7 minutes, then A:B=5:95 for 0.4 minutes, then with a linear gradient elution to A:B 95:5 for 0.4 minutes with a constant flow rate of 1.5
15 mL/min.

Method 3: An Agilent 1200 LCMS system with ELS detector was used. Column: Agilent TC-C18 5 µm; 2.1x50mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9:0.1) and B = acetonitrile /trifluoroacetic acid (99.95:0.05); Method: Linear gradient elution with A:B = 99:1 to 0:100 in 4.0 minutes and with a flow rate of 0.8 mL/min.

- 20 **Method 4:** An Agilent 1200 LCMS system with ELS detector was used. Column: Agilent TC-C18 5 µm; 2.1x50mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9:0.1) and B = acetonitrile /trifluoroacetic acid (99.95:0.05); Method: Linear gradient elution with A:B = 90:10 to 0:100 in 4.0 minutes and with a flow rate of 0.8 mL/min.

- 25 **Method 5:** A Waters Acquity UPLC-MS was used. Column: Acquity UPLC BEH C18 1.7µm; 2.1x50mm; Column temperature: 60 °C; Solvent system: A = water/trifluoroacetic acid (99.965:0.035) and B = acetonitrile /water/trifluoroacetic acid (94.965:5:0.035); Method: Linear gradient elution with A:B = 90:10 to 0:100 in 1.0 minutes and with a flow rate of 1.2 mL/minute.

- 30 **Method 6:** A Waters Acquity UPLC-MS was used. Column: Acquity UPLC BEH C18 1.7µm; 2.1x50mm; Column temperature: 60 °C; Solvent system: A = water/formic acid (99.9:0.1) and B = acetonitrile /water/formic acid (94.9:5:0.1); Method: Linear gradient elution with A:B = 90:10 to 0:100 in 1.0 minutes and with a flow rate of 1.2 mL/minute.

Method 7: A Waters Acquity UPLC-MS was used. Column: Acquity UPLC HSS T3 C18 1.8 μ m; 2.1x50mm; Column temperature: 60 °C; Solvent system: A = water/trifluoroacetic acid (99.965:0.035) and B = acetonitrile /water/trifluoroacetic acid (94.965:5:0.035); Method: Linear gradient elution with A:B = 98:02 to 0:100 in 1.0 minutes and with a flow rate of 1.2 mL/min.

- 5 **Method 8:** An Agilent 1200 LCMS system with ELS detector was used. Phenomenex Luna-C18, 5 μ m; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9:0.1) and B = acetonitrile /trifluoroacetic acid (99.95:0.05); Method: Linear gradient elution with A:B = 99:1 to 0:100 in 4.0 minutes and with a flow rate of 0.8 mL/min.

- 10 **Method 9:** An Agilent 1200 LCMS system with ELS detector was used. Column: Xtimate C18 2.1*30mm,3 μ m; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = ater/trifluoroacetic acid (99.9996:0.0004) and B = acetonitrile /trifluoroacetic acid (99.9998:0.0002); Method: Linear gradient elution with A:B = 100:0 to 70:30 in 3.0 minutes and with a flow rate of 0.8 mL/min.

- 15 **Method 10:** An Agilent 1200 LCMS system with ELS detector was used. Column: Xtimate C18 2.1*30mm,3 μ m; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = ater/trifluoroacetic acid (99.9996:0.0004) and B = acetonitrile /trifluoroacetic acid (99.9998:0.0002); Method: Linear gradient elution with A:B = 100:0 to 40:60 in 1.5 minutes and with a flow rate of 1.2 mL/min.

- 20 **Method 11:** An Agilent 1200 LCMS system with ELS detector was used. Column: Waters XBridge ShieldRP18,2.1*50mm,5 μ m; Column temperature: 40 °C; Solvent system: A = water/ammonia (99.95:0.05) and B = acetonitrile; Method: Linear gradient elution with A:B = 95:5 to 0:100 in 4.0 minutes and with a flow rate of 0.8 mL/min.

Method 12: An Agilent 1100 LCMS system with ELS detector was used. Column: YMC ODS-AQ 5 μ m; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9:0.1) and B = acetonitrile /trifluoroacetic acid (99.95:0.05); Method: Linear gradient elution with A:B = 99:1 to 5:95 in 3.5 minutes and with a flow rate of 0.8 mL/min.

- 25 **Method 13:** An Agilent 1200 LCMS system with ELS detector was used. Phenomenex Luna-C18, 5 μ m; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9:0.1) and B = acetonitrile /trifluoroacetic acid (99.95:0.05); Method: Linear gradient elution with A:B = 99:1 to 0:100 in 4.0 minutes and with a flow rate of 0.8 mL/min.

- 30 **Method 14:** An Agilent 1200 LCMS system with ELS detector was used. Column: Xtimate C18 2.1*30mm,3 μ m; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = ater/trifluoroacetic acid (99.9996:0.0004) and B = acetonitrile / trifluoroacetic acid (99.9998:0.0002); Method: Linear gradient elution with A:B = 100:0 to 40:60 in 6.0 minutes and with a flow rate of 0.8 mL/min.

Method 15: An Agilent 1200 LCMS system with ELS detector was used. Column: MERCK,RP-18e 25-2mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9996:0.0004) and B = acetonitrile /trifluoroacetic acid (99.9998:0.0002); Method: Linear gradient elution with A:B = 95:5 to 5:95 in 0.7 minutes and with a flow rate of 1.5 mL/min.

- 5 **Method 16:** An Agilent 1200 LCMS system with ELS detector was used. Column: Xtimate C18 2.1*30mm,3µm; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9996:0.0004) and B = acetonitrile /trifluoroacetic acid (99.9998:0.0002); Method: Linear gradient elution with A:B = 100:0 to 40:60 in 0.9 minutes and with a flow rate of 1.2 mL/min.

- 10 **Method 17:** An Agilent 1200 LCMS system with ELS detector was used. Phenomenex Luna-C18, 5µm; 2.0x50mm; Column temperature: 50 °C; Solvent system: A = water/trifluoroacetic acid (99.9:0.1) and B = acetonitrile /trifluoroacetic acid (99.95:0.05); Method: Linear gradient elution with A:B = 90:10 to 0:100 in 4.0 minutes and with a flow rate of 0.8 mL/min.

- 15 **Method 18:** An Agilent 1200 LCMS system with ELS detector was used. Column: Waters XBridge ShieldRP18,2.1*50mm,5µm; Column temperature: 40 °C; Solvent system: A = water/ammonia (99.95:0.05) and B = acetonitrile; Method: Linear gradient elution with A:B = 85:15 to 0:100 in 3.4 minutes and with a flow rate of 0.8 mL/min.

- 20 **Method 19:** A Waters Acquity UPLC-MS was used. Column: Acquity UPLC BEH C18 1.7µm; 2.1x50mm; Column temperature: 60 °C; Solvent system: A = water/formic acid (99.9:0.1) and B = acetonitrile /water/formic acid (94.9:5:0.1); Method: Linear gradient elution with A:B = 98:2 to 0.1:99.9 in 1.0 minutes and with a flow rate of 1.2 mL/minute.


- 25 Preparative LC-MS-purification was performed on a PE Sciex API 150EX instrument with atmospheric pressure chemical ionization. Column: 50 X 20 mm YMC ODS-A with 5 µm particle size; Solvent system: A = water/trifluoroacetic acid (99.965:0.035) and B = acetonitrile /water/trifluoroacetic acid (94.965:5:0.035); Method: Linear gradient elution with A:B = 80:20 to 0:100 in 7 minutes and with a flow rate of 22.7 mL/minute. Fraction collection was performed by split-flow MS detection.

Preparative SFC was performed on a Thar 80 instrument. Exemplified conditions can be, but not limited to: Column AD 250 X 30mm with 20 µm particle size; Column temperature: 38 °C, Mobile phase: Supercritical CO₂/ EtOH(0.2%NH₃H₂O) =45/55.

75°C for 6 hours. The mixture was filtered and concentrated *in vacuo*. The residue was purified by flash chromatography using a gradient of petroleum ether and ethyl acetate to afford *N*-((5-iodo-3-methoxypyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-4-carboxamide 2.20 g (64.7%).


5

6-Iodo-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine:



To a solution of *N*-((5-iodo-3-methoxypyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-4-carboxamide (2 g, 5.30 mmol) in 1,4-dioxane (60 mL) was added phosphoryl chloride (8.13 g, 53.0 mmol) at 0°C. The reaction was stirred at 85°C for 12 hours. The mixture was concentrated *in vacuo*. The residue was diluted with dichloromethane (100 mL) and ice-water (60 mL), followed by saturated aqueous NaHCO₃ (30 mL). The organic phase was separated and the water phase was extracted with dichloromethane (3×20 mL). The combined organic phases were dried and concentrated *in vacuo*. The residue was purified by flash chromatography using a gradient of petroleum ether and ethyl acetate to yield 6-iodo-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine 500 mg (23.6%).


6-Benzyl-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine:



A mixture of 6-iodo-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine (500 mg, 1.39 mmol), 2-benzyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (911 mg, 4.18 mmol), Pd(dppf)Cl₂ (51 mg, 0.07 mmol), K₂CO₃ (577 mg, 4.18 mmol) and H₂O (3 mL) in 1,4-dioxane (15 mL) was stirred at 80°C for 12 hrs under an atmosphere of N₂. It was then filtered and the filtrate was concentrated *in vacuo*. The residue was purified by flash chromatography using a gradient of petroleum ether and ethyl acetate to yield 6-benzyl-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine 260 mg (52%).

9.8 mmol) in anhydrous 1,4-dioxane (20 mL) was added phosphoryl chloride (3.4 g, 22 mmol). The reaction was stirred at 80°C for 2 hours. Then the solution was cooled and poured into water (100 mL), pH was adjusted to 8-9 by the addition of saturated aqueous K₂CO₃. The crude mixture was extracted with ethyl acetate (2×100 mL). The combined organic phases were washed with brine (50 mL), dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by flash chromatography using a gradient of petroleum ether and ethyl acetate to yield 8-chloro-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine 2.1 g (90%).

10 **3-(Tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:**



To a solution of 8-chloro-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine (2.1 g, 8.8 mmol) in 1,4-dioxane (20 mL) was added 2 M aq. HCl (10 mL). The solution was stirred at 80°C for 2 hours. The mixture was cooled and pH was adjusted to 8-9 by addition of saturated aqueous K₂CO₃. The crude mixture was concentrated *in vacuo* and the residue was dissolved in methanol (150 mL) and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash chromatography using a mixture of dichloromethane and methanol (10:1) to give 3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 1.6 g (81%).

20 ¹H NMR (DMSO-*d*₆, 400 MHz): δ 10.55 (s, 1H), 7.63 (s, 1H), 7.39 (d, *J* = 6.0 Hz, 1H), 6.60 (s, 1H), 3.91-3.88 (m, 2H), 3.49-3.42 (m, 2H), 3.34-3.29 (m, 1H), 1.82-1.72 (m, 4H).

LC-MS: (*m/z*) 220.1 (MH⁺) *t_R* (minutes, method 3) = 1.37 minutes

***N*-((3-Chloropyrazin-2-yl)methyl)cyclopropanecarboxamide:**

25



flash chromatography using dichloromethane/methanol (10/1) to afford 3-cyclopropylimidazo[1,5-a]pyrazin-8(7*H*)-one 1.9 g (83%).

¹H NMR (DMSO-*d*₆, 400 MHz): δ 10.49 (s, 1H), 7.54 (s, 1H), 7.41 (d, *J* = 5.6 Hz, 1H), 6.61 (d, *J* = 5.6 Hz, 1H), 2.29-2.24 (m, 1H), 0.99-0.89 (m, 4H).

5 LC-MS: (*m/z*) 176.1 (MH⁺) *t*_R (minutes, method 1) = 1.04 minutes


***N*-((5-Bromo-3-methoxypyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-4-carboxamide:**



10 A solution of NaNO₂ (972 mg, 14.09 mmol) in H₂O (100 mL) was added to a stirred solution of *N*-((5-amino-3-methoxypyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-4-carboxamide (2.5 g, 9.4 mmol) in 40% aq. HBr (33 mL) at 0°C. After stirring for 1.5 hrs, CuBr (2.02 g, 14.1 mmol) was added and the mixture was stirred at 70°C for 1 hour. The pH value was adjusted to pH 8 by addition of saturated aqueous NaHCO₃. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by flash chromatography using a gradient of
15 petroleum ether and ethyl acetate to yield *N*-((5-bromo-3-methoxypyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-4-carboxamide 800 mg (25%).

LC-MS: (*m/z*) 331.8 (MH⁺) *t*_R (minutes, method 2) = 0.723 minutes

6-Bromo-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine:




20

To a solution of *N*-((5-bromo-3-methoxypyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-4-carboxamide (800 mg, 2.42 mmol) in 1,4-dioxane (30 mL) was added phosphoryl chloride (3.8 g, 25 mmol) at 0°C. The mixture was heated to 70°C and stirred for 1 hour. The mixture was concentrated *in vacuo* and the residue was diluted with dichloromethane (100 mL) and
25 ice-water (60 mL). The pH value was adjusted to pH 8 by addition of saturated aqueous NaHCO₃. The organic phase was separated and aqueous phase was extracted with

dichloromethane (3×30 mL). The combined organic phases were dried and concentrated *in vacuo*. The residue was purified by flash chromatography using a gradient of dichloromethane and methanol to yield 6-bromo-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine 500 mg (66%).

5 LC-MS: (*m/z*) 313.7 (MH^+) t_R (minutes, method 2) = 0.740 minutes

6-Bromo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:



10 To a solution of 6-bromo-8-methoxy-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazine (200 mg, 0.641 mmol) in dichloromethane (30 mL) was added boron tribromide (1.61 g, 6.41 mmol) at 0°C. The reaction was warmed to 20°C and stirred for 3 hours. The solution was quenched with water (2 mL) at 0°C. The reaction was concentrated *in vacuo* and the residue was purified by flash chromatography using a gradient of dichloromethane and methanol to yield 6-bromo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 130 mg (68%).


15 LC-MS: (*m/z*) 299.7 (MH^+) t_R (minutes, method 2) = 0.730 minutes

2-methoxy-6-methylpyrazine




20 To a suspension of 2-chloro-6-methylpyrazine (24 g, 186.7 mmol) in anhydrous MeOH (240 mL) was added NaOMe (12.1 g, 224 mmol). The mixture was stirred at 60-70°C for 16 hours. The mixture was cooled and filtered. The filtrate was concentrated *in vacuo* to give 2-methoxy-6-methylpyrazine (22 g, 95% yield). ¹H NMR (CDCl₃ 400 MHz): δ 7.98 (s, 1H), 7.94 (s, 1H), 3.91 (s, 3H), 3.40 (s, 3H). LC-MS: t_R = 1.47 min (method 14), *m/z* = 124.8 [$M + H$]⁺.

25 **3-methoxy-5-methylpyrazine 1-oxide**




To a solution of 2-methoxy-6-methylpyrazine (21.3 g, 171.6 mmol) in AcOH (150 mL) was added $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2 \cdot 3\text{H}_2\text{O}$ (31.7 g, 205.9 mmol). The mixture was stirred at 80°C for 16 hours. The mixture was concentrated *in vacuo* and diluted with 2 M aq. NaOH (300 mL). The mixture was extracted with EtOAc (200 mL $\times$ 4). The organic layer was washed with brine (100 mL), dried over Na_2SO_4 and concentrated in vacuo to give 3-methoxy-5-methylpyrazine 1-oxide (14.4 g, 60% yield).

3-methoxy-5-methylpyrazine-2-carbonitrile



To a mixture of 3-methoxy-5-methylpyrazine 1-oxide (10 g, 71.4 mmol) in MeCN (200 mL) was added TMS-CN (24.8 g, 249.8 mmol) and triethylamine (36.1 g, 356.8 mmol), ZnBr_2 (32.1 g, 142.7 mmol). The mixture was stirred at 85-90°C for 16 hours. The mixture was concentrated in vacuo. The residue was diluted with DCM (500 mL) and filtered. The filtrate was washed with water (300 mL) and brine (200 mL). The organic layer was dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=5:1) to give 3-methoxy-5-methylpyrazine-2-carbonitrile (4.1 g, 38% yield).

tert-butyl ((3-methoxy-5-methylpyrazin-2-yl)methyl)carbamate




To a solution of 3-methoxy-5-methylpyrazine-2-carbonitrile (6.22 g, 41.7 mmol) in MeOH (100 mL) was added $(\text{Boc})_2\text{O}$ (13.65 g, 62.6 mmol) and Raney Ni (2.0 g). The mixture was stirred at 20-25°C under H_2 (45 psi) for 16 hours. The mixture was filtered and the filtrate was concentrated in vacuo. The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=5:1) to give *tert*-butyl ((3-methoxy-5-methylpyrazin-2-

yl)methyl)carbamate (7.7 g, 72% yield).

LC-MS: t_R = 0.70 min (method 15), m/z = 254.0 $[M + H]^+$.

(3-methoxy-5-methylpyrazin-2-yl)methanamine



To a solution of *tert*-butyl ((3-methoxy-5-methylpyrazin-2-yl)methyl)carbamate (7.7 g, 30.3 mmol) in THF (50 mL) was added TFA (20 mL). The mixture was stirred at 80°C for 2 hours. The mixture was concentrated in vacuo. The residue was diluted with 2 M aq. NaOH (200 mL), extracted with DCM (100 mL $\times$ 2). The organic layer was washed with brine (50 mL),
 10 dried over Na_2SO_4 and concentrated in vacuo to give (3-methoxy-5-methylpyrazin-2-yl)methanamine (2.5 g, 54% yield).


^1H NMR ($CDCl_3$ 400 MHz): δ 7.90 (s, 1H), 3.96 (s, 3H), 3.93 (s, 2H), 2.42 (s, 3H), 1.69 (s, 2H).

LC-MS: t_R = 0.73 min(method 16), m/z = 154.2 $[M + H]^+$.

15

COMPOUNDS OF THE INVENTION:

Example 1



7-(3-Fluorobenzyl)-3-propylimidazo[1,5-a]pyrazin-8(7H)-one:


20 To a solution of 3-propylimidazo[1,5-a]pyrazin-8(7H)-one (1.2 g, 6.8 mmol) in DMF (10 mL) was added potassium carbonate (1.4 g, 10 mmol) and 1-(bromomethyl)-3-fluorobenzene (1.54 g, 8.13 mmol). The mixture was stirred at 60-70°C for 2 hrs and then cooled to room temperature. To the reaction was added water (75 mL) and it was extracted with ethyl acetate (2 $\times$ 50mL). The combined organic phases were washed with brine, dried and

concentrated *in vacuo*. The residue purified by flash chromatography to yield 1.5 g (78%) of 7-(3-fluorobenzyl)-3-propylimidazo[1,5-a]pyrazin-8(7*H*)-one.

¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.70 (s, 1H), 7.46-7.37 (m, 2H), 7.17-7.09 (m, 3H), 7.00 (d, *J*=6.0 Hz, 1H), 5.02 (s, 2H), 2.83 (t, 2H), 1.75-1.66 (m, 2H), 0.91 (t, 3H).

5 LC-MS: (*m/z*) 286.1 (MH⁺) *t*_R (minutes, method 3) = 2.19 minutes

Example 2



6-Benzyl-7-(3-fluorobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

10

A mixture of 6-benzyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 0.647 mmol), 1-(bromomethyl)-3-fluorobenzene (159 mg, 840 μ mol) and potassium carbonate (179 mg, 1.29 mmol) in DMF (6 mL) was stirred at 60°C for 12 hours. The reaction mixture was concentrated *in vacuo* and the residue was diluted

15


with dichloromethane (20 mL) and washed with water (3 $\times$ 5 mL). The combined organic phases were dried, filtered and concentrated *in vacuo*. The residue was purified by preparative TLC, eluting with petroleum ether and ethyl acetate 1:2, to yield 6-benzyl-7-(3-fluorobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 80 mg (28%).

20

¹H NMR (CDCl₃, 400 MHz): δ 7.97 (s, 1 H), 7.39-7.30 (m, 4 H), 7.17 (d, *J*=7.53 Hz, 2 H), 6.94-6.92 (m, 2 H), 6.83 (d, *J*=9.54 Hz, 1 H), 6.74 (s, 1 H), 5.06 (s., 2 H), 4.12 (d, *J*=12.05 Hz, 2 H), 3.74 (s, 2 H), 3.58 - 3.52 (m, 2 H), 3.03 - 3.09 (m, 1 H), 2.19 - 2.09 (m, 2 H), 1.89 (d, *J*=13.55 Hz, 2 H).

LC-MS: (*m/z*) 418.2 (MH⁺) *t*_R (minutes, method 3) = 2.69 minutes

25 Example 3




6-Benzyl-7-(cyclohexylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

A mixture of 6-benzyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (60 mg, 0.19 mmol), (bromomethyl)cyclohexane (52 mg, 0.29 mmol) and potassium carbonate (54 mg, 0.39 mmol) in DMF (10 mL) was stirred at 75°C for 12 hours. The reaction mixture was concentrated *in vacuo*. The residue was diluted with dichloromethane (20 mL) and washed with water (3×5 mL). The combined organic phases were dried, filtered and concentrated *in vacuo*. The residue was purified by preparative TLC, eluting with petroleum ether and ethyl acetate 1:2, to yield 6-benzyl-7-(cyclohexylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 15 mg (8.1%).

¹H NMR (CDCl₃, 400 MHz): δ 7.87 (s, 1 H), 7.42-7.30 (m, 3 H), 7.19 (d, *J*=7.34 Hz, 2 H), 6.65 (s, 1 H), 4.11 (d, *J*=11.25 Hz, 2 H), 3.90 (s, 2 H), 3.66 (d, *J*=6.36 Hz, 2 H), 3.54 (t, *J*=10.76 Hz, 2 H), 3.05-2.99 (m, 1 H), 2.18 - 2.04 (m, 2 H), 1.86 (d, *J*=13.94 Hz, 2 H), 1.63 - 1.77 (m, 7 H), 1.18-1.15 (m, 2 H), 1.04-1.01 (m, 2 H).

LC-MS: (*m/z*) 406.2 (MH⁺) *t_R* (minutes, method 4) = 2.38 minutes

Example 4



7-(Cyclohexylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (50 mg, 0.21 mmol), (bromomethyl)cyclohexane (57 mg, 0.32 mmol) and potassium carbonate (59 mg, 0.43 mmol) in DMF (2 mL) was stirred at 60°C for 12 hours. The reaction mixture was concentrated *in vacuo*. The residue was diluted with dichloromethane (20 mL) and washed

with water (3×5 mL). The combined organic phases were dried, filtered and concentrated *in vacuo*. The residue was purified by preparative LC-MS to yield 7-(cyclohexylmethyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 20 mg (28%),

¹H NMR (CDCl₃, 400 MHz): δ 7.85 (s, 1 H), 6.71 (s, 1 H), 4.13 (d, *J*=11.04 Hz, 2 H), 3.78 (d, *J*=7.03 Hz, 2 H), 3.66-3.53 (m, 2 H), 3.11-3.05 (m, 1 H), 2.27 (s, 3 H), 2.20-2.05 (m, 2 H), 1.88 (d, *J*=12.05 Hz, 2 H), 1.80-1.64 (m, 6 H), 1.25-1.13 (m, 3 H), 1.12-0.99 (m, 2 H).

LC-MS: (*m/z*) 330.2 (MH⁺) *t_R* (minutes, method 3) = 2.49 minutes

Example 5




7-(3-Fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (50 mg, 0.21 mmol), 1-(bromomethyl)-3-fluorobenzene (60 mg, 0.32 mmol) and K₂CO₃ (59 mg, 0.43 mmol) in DMF (2 mL) was stirred at 60°C for 12 hours. The reaction mixture was concentrated *in vacuo*. The residue was diluted with dichloromethane (20 mL) and washed with water (3×5 mL). The organic layer was dried and concentrated *in vacuo*. The residue was purified by preparative LC-MS to give 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 20 mg (27%).

¹H NMR (CDCl₃, 400 MHz): δ 7.94 (s, 1 H), 7.34-7.29 (m, 1 H), 7.03-6.94 (m, 2 H), 6.90 (d, *J*=9.70 Hz, 1 H), 6.77 (s, 1 H), 5.23 (s, 2 H), 4.14 (d, *J*=10.14 Hz, 2 H), 3.59 (td, *J*=11.69, 1.76 Hz, 2 H), 3.15-3.05 (m, 1 H), 2.18 (s, 3 H), 2.17-2.08 (m, 2 H), 1.90 (d, *J*=13.45 Hz, 2 H).

LC-MS: (*m/z*) 342.2 (MH⁺) *t_R* (minutes, method 3) = 2.34 minutes

Example 6




3-Cyclopropyl-7-(3-fluorobenzyl)imidazo[1,5-a]pyrazin-8(7H)-one:

To a solution of 3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one (300 mg, 1.71 mmol) in anhydrous DMF (5 mL) was added K_2CO_3 (355 mg, 2.57 mmol) and 1-(bromomethyl)-3-fluorobenzene (388 mg, 2.05 mmol). The mixture was stirred at 65°C for 16 hours. The mixture was filtered and the filtrate was purified by preparative-HPLC to yield 3-cyclopropyl-7-(3-fluorobenzyl)imidazo[1,5-a]pyrazin-8(7H)-one 280 mg (58%).

^1H NMR (DMSO- d_6 , 400 MHz): δ 7.59-7.56 (m, 2H), 7.36-7.33 (m, 1H), 7.13-7.09 (m, 3H), 6.98 (d, $J=6.0$ Hz, 1H), 4.98 (s, 2H), 2.29-2.24 (m, 1H), 1.00-0.90 (m, 4H).

LC-MS: (m/z) 284.1 (MH^+) t_R (minutes, method 3) = 2.23 minutes

Example 7




7-(Cyclopentylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one:

To a solution of 3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one (300 mg, 1.71 mmol) in anhydrous DMF (5 mL) was added K_2CO_3 (355 mg, 2.57 mmol) and (bromomethyl)cyclopentane (335 mg, 2.05 mmol). The reaction was stirred at 65°C for 16 hours. The mixture was filtered and the filtrate was purified by preparative LC-MS to yield 7-(cyclopentylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one 210 mg (47%).

^1H NMR (DMSO- d_6 , 400 MHz): δ 7.53 (s, 1H), 7.51 (d, $J = 6.0$ Hz, 1H), 6.90 (d, $J = 6.0$ Hz, 1H), 3.69 (d, $J = 7.6$ Hz, 2H), 2.29-2.24 (m, 2H), 1.58-1.45 (m, 6H), 1.22-1.19 (m, 2H), 1.00-0.89 (m, 4H).


LC-MS: (m/z) 258.2 (MH^+) t_R (minutes, method 3) = 2.24 minutes

Example 8**7-(Cyclohexylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one:**

To a solution of 3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one (300 mg, 1.71 mmol) in anhydrous DMF (5 mL) was added K_2CO_3 (355 mg, 2.57 mmol) and (bromomethyl)cyclohexane (363 mg, 2.05 mmol). The reaction was stirred at 65°C for 16 hours. The reaction mixture was filtered and the filtrate was purified by preparative LC-MS to yield 7-(cyclohexylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one 195 mg (42%).

^1H NMR (DMSO- d_6 , 400 MHz): δ 7.53-7.49 (m, 2H), 6.84 (d, J = 6.0 Hz, 1H), 3.59 (d, J = 7.6 Hz, 2H), 2.27-2.25 (m, 1H), 1.68-1.55 (m, 6H), 1.10-0.89 (m, 9H).

LC-MS: (m/z) 272.2 (MH^+) t_R (minutes, method 3) = 2.40 minutes


Example 9**7-(3-Fluorobenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**

To a solution of 3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (200 mg, 0.91 mmol) in anhydrous DMF (5 mL) was added K_2CO_3 (189 mg, 1.37 mmol) and 1-(bromomethyl)-3-fluorobenzene (207 mg, 1.10 mmol). The reaction mixture was stirred at 60°C for 16 hours. The mixture was filtered and the filtrate was purified by preparative LC-MS to afford 7-(3-fluorobenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 190 mg (64%).

^1H NMR (DMSO- d_6 , 400 MHz): δ 7.68 (s, 1H), 7.54 (d, J = 6.0 Hz, 1H), 7.36-7.33 (m, 1H), 7.14-7.09 (m, 3H), 6.99 (d, J = 6.0 Hz, 1H), 4.99 (s, 2H), 3.91-3.88 (m, 2H), 3.48-3.42 (m, 2H), 3.30-3.29 (m, 1H), 1.81-1.73 (m, 4H).

LC-MS: (*m/z*) 328.1 (MH^+) t_R (minutes, method 1) = 1.92 minutes

Example 10




5 **7-(Cyclopentylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**

To a solution of 3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (400 mg, 1.82 mmol) in anhydrous DMF (5 mL) was added K_2CO_3 (503 mg, 3.64 mmol) and (bromomethyl)cyclopentane (445 mg, 2.73 mmol). The reaction mixture was stirred at 60°C for 16 hours. The mixture was filtered and the filtrate was purified by preparative LC-MS to afford 7-(cyclopentylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 290 mg (53%).

^1H NMR ($DMSO-d_6$, 400 MHz): δ 7.62 (s, 1H), 7.49 (d, J = 6.0 Hz, 1H), 6.90 (d, J = 6.0 Hz, 1H), 3.91-3.88 (m, 2H), 3.69 (d, J = 7.6 Hz, 2H), 3.48-3.43 (m, 2H), 3.42-3.31 (m, 1H), 2.27-2.24 (m, 1H), 1.78-1.73 (m, 4H), 1.58-1.44 (m, 6H), 1.21-1.20 (m, 2H).

15 LC-MS: (*m/z*) 302.2 (MH^+) t_R (minutes, method 3) = 2.29 minutes

Example 11



7-(Cyclohexylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


20 To a solution of 3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (300 mg, 1.37 mmol) in anhydrous DMF (5 mL) was added K_2CO_3 (379 mg, 2.74 mmol) and (bromomethyl)cyclohexane (364 mg, 2.06 mmol). The reaction mixture was stirred at 60°C for 16 hours. The mixture was filtered and the filtrate was purified by preparative LC-MS to yield 7-(cyclohexylmethyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-

a]pyrazin-8(7*H*)-one 240 mg (55%).

¹H NMR (DMSO-*d*₆, 400 Mhz): δ 7.62 (s, 1H), 7.47 (d, *J* = 6.0 Hz, 1H), 6.84 (d, *J* = 6.2 Hz, 1H), 3.91-3.88 (m, 2H), 3.60 (d, *J* = 7.2 Hz, 2H), 3.48-3.42 (m, 2H), 3.30-3.29 (m, 1H), 1.78-1.52 (m, 10H), 1.10-1.06 (m, 3H), 0.93-0.91 (m, 2H).

5 LC-MS: (*m/z*) 316.2 (MH⁺) *t*_R (minutes, method 3) = 2.44 minutes.

Example 12



7-(Cycloheptylmethyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:


10 To a solution of 3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (300 mg, 1.37 mmol) in anhydrous DMF (4 mL) was added K₂CO₃ (568 mg, 4.11 mmol) and cycloheptylmethyl methanesulfonate (565 mg, 2.74 mmol). The reaction mixture was stirred at 95°C for 16 hours. The mixture was filtered and the filtrate was purified by preparative LC-MS to afford 7-(cycloheptylmethyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-
15 a]pyrazin-8(7*H*)-one 140 mg (30%).

¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.62 (s, 1H), 6.48 (d, *J* = 5.6 Hz, 1H), 6.87 (d, *J* = 6.0 Hz, 1H), 3.91-3.88 (m, 2H), 3.59 (d, *J* = 7.6 Hz, 2H), 3.48-3.42 (m, 2H), 3.31-3.26 (m, 1H), 1.90-1.73 (m, 5H), 1.57-1.43 (m, 10H), 1.13-1.11 (m, 2H).

LC-MS: (*m/z*) 330.2 (MH⁺) *t*_R (minutes, method 3) = 2.58 minutes

20

Example 13



7-(Cycloheptylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one:


To a solution of 3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one (300 mg, 1.71 mmol) in anhydrous DMF (4 mL) was added K_2CO_3 (709 mg, 5.1

- 5 mmol) and cycloheptylmethyl methanesulfonate (706 mg, 3.42 mmol). The reaction mixture was stirred at 95°C for 16 hours. The mixture was filtered and the filtrate was purified by preparative LC-MS to afford 7-(cycloheptylmethyl)-3-cyclopropylimidazo[1,5-a]pyrazin-8(7H)-one 115 mg (24%).

^1H NMR (DMSO- d_6 , 400 MHz): δ 7.55-7.52 (m, 2H), 6.89 (d, J = 6.0 Hz, 1H), 3.61 (d, J = 7.6 Hz, 2H), 2.32-2.27 (m, 1H), 1.91 (brs, 1H), 1.91-1.02 (m, 12H), 1.01-0.91 (m, 4H).

LC-MS: (m/z) 286.2 (MH^+) t_R (minutes, method 3) = 2.54 minutes.

Example 14



- 15 **7-(4-Chlorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**


A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-4-chlorobenzene (132 mg, 0.643 mmol) and Cs_2CO_3 (280 mg, 0.857 mmol) in DMF (2.0 mL) was stirred at 70°C for 12 hours. The reaction mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to afford 7-(4-chlorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 53 mg (34%).

^1H NMR ($CDCl_3$, 400 MHz): δ 7.93 (s, 1H), 7.30 (d, J = 8.4 Hz, 2H), 7.16 (d, J = 8.4 Hz, 2H),

6.76 (s, 1H), 5.20 (s, 2H), 4.13 (d, $J = 10.8$ Hz, 2H), 3.62 - 3.56 (m, 2H), 3.12 - 3.05 (m, 1H), 2.18 (s, 3H), 2.16 - 2.09 (m, 2H), 1.89 (d, $J = 13.2$ Hz, 2H).

LC-MS: (m/z) 358.1 (MH^+) t_R (minutes, method 3) = 2.46 minutes

5 Example 15




6-Bromo-7-(3-fluorobenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

A mixture of 6-bromo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.335 mmol), 1-(bromomethyl)-3-fluorobenzene (95 mg, 0.50 mmol) and K_2CO_3 (93 mg, 0.67 mmol) in DMF (2.0 mL) was stirred at 60°C for 12 hours. The reaction mixture was concentrated *in vacuo*. The residue was purified by preparative LC-MS to give 6-bromo-7-(3-fluorobenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 30 mg (22%).

^1H NMR ($CDCl_3$, 400 MHz): δ 7.95 (s, 1H), 7.34 - 7.27 (m, 1H), 7.18 (s, 1H), 7.11 (d, $J = 7.6$ Hz, 1H), 7.03 - 6.99 (m, 2H), 5.39 (s, 2H), 4.13 (d, $J = 12.0$ Hz, 2H), 3.62 - 3.56 (m, 2H), 3.11 - 3.05 (m, 1H), 2.18 - 2.08 (m, 2H), 1.88 (d, $J = 14.0$ Hz, 2H).

LC-MS: (m/z) 408.0 (MH^+) t_R (minutes, method 3) = 2.65 minutes

Example 16




7-Benzyl-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 0.429 mmol), (bromomethyl)benzene (110 mg, 0.643 mmol) and K₂CO₃ (119 mg, 0.857 mmol) in DMF (1.0 mL) was stirred at 60°C for 3 hours. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to yield 7-benzyl-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 39 mg (28%).

¹H NMR (CDCl₃, 400 MHz): δ 7.93 (s, 1H), 7.35 - 7.29 (m, 3H), 7.21 (d, *J* = 7.6 Hz, 2H), 6.75 (s, 1H), 5.25 (s, 2H), 4.13 (d, *J* = 11.6 Hz, 2H), 3.61 - 3.56 (m, 2H), 3.13 - 3.06 (m, 1H), 2.19 (s, 3H), 2.15 - 2.08 (m, 2H), 1.89 (d, *J* = 13.2 Hz, 2H).

LC-MS: (*m/z*) 324.2 (MH⁺) *t_R* (minutes, method 3) = 2.25 minutes

Example 17




7-(2-Fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-2-fluorobenzene (122 mg, 0.643 mmol) and Cs₂CO₃ (279 mg, 0.857 mmol) in DMF (2.0 mL) was stirred at 70°C for 1 hour. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to afford 7-(2-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 49 mg (33%).

¹H NMR (CDCl₃, 400 MHz): δ 7.93 (s, 1H), 7.27 - 7.23 (m, 1H), 7.11 - 7.06 (m, 3H), 6.77 (s, 1H), 5.28 (s, 2H), 4.13 (d, *J* = 10.4 Hz, 2H), 3.62-3.56 (m, 2H), 3.12 - 3.07 (m, 1H), 2.19 (s, 3H), 2.16 - 2.10 (m, 2H), 1.89 (d, *J* = 12.0 Hz, 2H).

LC-MS: (*m/z*) 342.1 (MH⁺) *t_R* (minutes, method 3) = 2.30 minutes

Example 18




7-(3-Chlorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-3-chlorobenzene (132 mg, 0.643 mmol) and K₂CO₃ (119 mg, 0.857 mmol) in DMF (1.0 mL) was stirred at 65°C for 12 hours. To the mixture was added Cs₂CO₃ (280 mg, 0.857 mmol) and the reaction was stirred at 80°C for another 1 hour. The reaction mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to afford 7-(3-chlorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 49 mg (32%).

¹H NMR (CDCl₃, 400 MHz): δ 7.92 (s, 1H), 7.25 - 7.24 (m, 2H), 7.16 (s, 1H), 7.10 - 7.07 (m, 1H), 6.75 (s, 1H), 5.19 (s, 2H), 4.11 (d, *J* = 10.4 Hz, 2H), 3.60 - 3.54 (m, 2H), 3.12 - 3.05 (m, 1H), 2.16 (s, 3H), 2.14 - 2.07 (m, 2H), 1.87 (d, *J* = 12.0 Hz, 2H).

LC-MS: (*m/z*) 358.1 (MH⁺) *t_R* (minutes, method 3) = 2.44 minutes

Example 19



7-(2-Chlorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-2-chlorobenzene (132 mg, 0.643 mmol) and Cs₂CO₃ (279 mg, 0.857 mmol) in DMF (2.0 mL) was stirred at 80°C for 12 hours. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to yield 7-(2-chlorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 55 mg (36%).

^1H NMR (CDCl_3 , 400 MHz): δ 7.95 (s, 1H), 7.42 - 7.40 (m, 1H), 7.25 - 7.20 (m, 2H), 6.93 (d, $J = 7.2$ Hz, 1H), 6.80 (s, 1H), 5.32 (s, 2H), 4.14 (d, $J = 10.8$ Hz, 2H), 3.63 - 3.57 (m, 2H), 3.14 - 3.07 (m, 1H), 2.20 - 2.10 (m, 2H), 2.13 (s, 3H), 1.91 (d, $J = 13.2$ Hz, 2H).

LC-MS: (m/z) 358.1 (MH^+) t_R (minutes, method 3) = 2.46 minutes

5

Example 20



7-(3-Methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


- 10 A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-3-methoxybenzene (129 mg, 0.643 mmol) and Cs_2CO_3 (280 mg, 0.857 mmol) in DMF (1.0 mL) was stirred at 80°C for 12 hours. The reaction mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to yield 7-(3-methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 68 mg (45%).

^1H NMR (CDCl_3 , 400 MHz): δ 7.92 (s, 1H), 7.27 - 7.23 (m, 1H), 6.82 - 6.77 (m, 2H), 6.74 (s, 2H), 5.21 (s, 2H), 4.13 (d, $J = 10.8$ Hz, 2H), 3.78 (s, 3H), 3.62 - 3.56 (m, 2H), 3.12 - 3.08 (m, 1H), 2.19 (s, 3H), 2.19 - 2.10 (m, 2H), 1.89 (d, $J = 13.2$ Hz, 2H).

LC-MS: (m/z) 354.2 (MH^+) t_R (minutes, method 3) = 2.28 minutes

20

Example 21




6-Methyl-7-(2-methylbenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-2-methylbenzene (119 mg, 0.643 mmol) and K₂CO₃ (119 mg, 0.857 mmol) in DMF (2.0 mL) was stirred at 60°C for 12 hours. To the mixture was added additionally Cs₂CO₃ (280 mg, 0.86 mmol) and the reaction stirred at 70°C for another 13 hours. The reaction mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to afford 6-methyl-7-(2-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 23 mg (16%).

¹H NMR (CDCl₃, 400 MHz): δ 7.93 (s, 1H), 7.21 - 7.12 (m, 3H), 6.80 (s, 1H), 6.78 (d, *J* = 7.6 Hz, 1H), 5.18 (s, 2H), 4.15 (d, *J* = 10.8 Hz, 2H), 3.63 - 3.57 (m, 2H), 3.15 - 3.09 (m, 1H), 2.40 (s, 3H), 2.20 - 2.13 (m, 2H), 2.11 (s, 3H), 1.92 (d, *J* = 13.2 Hz, 2H).

LC-MS: (*m/z*) 338.2 (MH⁺) *t_R* (minutes, method 3) = 2.37 minutes

Example 22




6-Methyl-7-(4-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-4-methylbenzene (119 mg, 0.643 mmol) and Cs₂CO₃ (280 mg, 0.857 mmol) in DMF (2.0 mL) was stirred at 70°C for 12 hours. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to afford 6-methyl-7-(4-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 68 mg (47%).

¹H NMR (CDCl₃, 400 MHz): δ 7.92 (s, 1H), 7.14 - 7.09 (m, 4H), 6.74 (s, 1H), 5.20 (s, 2H), 4.13 (d, *J* = 10.4 Hz, 2H), 3.62 - 3.56 (m, 2H), 3.11 - 3.06 (m, 1H), 2.32 (s, 3H), 2.19 (s, 3H), 2.19 - 2.08 (m, 2H), 1.88 (d, *J* = 13.2 Hz, 2H).

LC-MS: (*m/z*) 338.2 (MH⁺) *t_R* (minutes, method 3) = 2.41 minutes.

Example 23




7-(4-Methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.43 mmol), 1-(bromomethyl)-4-methoxybenzene (129 mg, 0.643 mmol) and Cs₂CO₃ (280 mg, 0.857 mmol) in DMF (2.0 mL) was stirred at 70°C for 12 hours. The mixture was filtered and the filtrate was concentrated in vacuo. The residue was purified by preparative LC-MS to afford 7-(4-methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 60 mg (39%).

¹H NMR (CDCl₃, 400 MHz): δ 7.93 (s, 1H), 7.17 (d, *J* = 8.4 Hz, 2H), 6.86 (d, *J* = 8.4 Hz, 2H), 6.73 (s, 1H), 5.17 (s, 2H), 4.13 (d, *J* = 10.0 Hz, 2H), 3.79 (s, 3H), 3.58 (t, *J* = 12.0 Hz, 2H), 3.11 - 3.05 (m, 1H), 2.21 (s, 3H), 2.18 - 2.08 (m, 2H), 1.88 (d, *J* = 13.6 Hz, 2H).

LC-MS: (*m/z*) 354.2 (MH⁺) *t_R* (minutes, method 3) = 2.26 minutes.

Example 24



7-(4-Fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-4-fluorobenzene (122 mg, 0.643 mmol) and Cs₂CO₃ (279 mg, 0.857 mmol) in DMF (2.0 mL) was stirred at 70°C for 12 hours. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to yield 7-(4-fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one 30 mg (21%).

¹H NMR (CDCl₃, 400 MHz): δ 7.93 (s, 1H), 7.22 - 7.19 (m, 2H), 7.04 - 7.00 (m, 2H), 6.75 (s,

¹H), 5.20 (s, 2H), 4.13 (d, *J* = 11.2 Hz, 2H), 3.62 - 3.60 (m, 2H), 3.17 - 3.06 (m, 1H), 2.19 (s, 3H), 2.14 - 2.08 (m, 2H), 1.88 (d, *J* = 13.6 Hz, 2H).

LC-MS: (*m/z*) 342.1 (MH⁺) *t*_R (minutes, method 3) = 2.30 minutes

5 Example 25




6-Methyl-7-(3-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 0.429 mmol), 1-(bromomethyl)-3-methylbenzene (119 mg, 0.643 mmol) and Cs₂CO₃ (280 mg, 0.857 mmol) in DMF (1 mL) was stirred at 80°C for 12 hours. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was purified by preparative LC-MS to afford 6-methyl-7-(3-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one 62 mg (43%).

¹H NMR (CDCl₃, 400 MHz): δ 7.93 (s, 1H), 7.21 (t, *J* = 7.2 Hz, 1H), 7.08 (d, *J* = 7.6 Hz, 1H), 7.02 - 6.98 (m, 2H), 6.74 (s, 1H), 5.21 (s, 2H), 4.13 (d, *J* = 10.8 Hz, 2H), 3.62 - 3.56 (m, 2H), 3.12 - 3.07 (m, 1H), 2.33 (s, 3H), 2.19 (s, 3H), 2.19 - 2.09 (m, 2H), 1.89 (d, *J* = 13.2 Hz, 2H).

LC-MS: (*m/z*) 338.1 (MH⁺) *t*_R (minutes, method 3) = 2.40 minutes.

20 Example 26:



7-(3-fluorobenzyl)-6-methyl-3-(4-methyltetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (150 mg, 0.98 mmol) and 4-methyltetrahydro-2H-pyran-4-carboxylic acid (212 mg, 1.5 mmol) in DCM (6 mL) was added HATU (670 mg, 1.8 mmol) and Et₃N (198 mg, 1.96 mmol). The mixture was stirred at 20-25°C for 1 hour. The mixture was diluted with DCM (50 mL), washed with water (30 mL) and brine (30 mL). The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=3:1) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-4-methyltetrahydro-2H-pyran-4-carboxamide (250 mg, 91% yield).

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-4-methyltetrahydro-2H-pyran-4-carboxamide (300 mg, 1.07 mmol) in dioxane (5 mL) was added POCl₃ (660 mg, 4.3 mmol). The solution was stirred at 80-90°C for 3 hours. The mixture was concentrated in vacuo, diluted with DCM (50 mL) and slowly added into water (30 mL). The organic layer was washed with brine (20 mL) and dried over Na₂SO₄, concentrated in vacuo to give 8-methoxy-6-methyl-3-(4-methyltetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazine (250 mg, 89% yield).

Step 3: To a solution of 8-methoxy-6-methyl-3-(4-methyltetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazine (200 mg, 0.77 mmol) in dioxane (10 mL) was added 2M HCl(aq) (10 mL). The solution was stirred at 80-90°C for 1 hour. The mixture was cooled and added saturated aqueous NaHCO₃ (100 mL), extracted with DCM (100 mL × 2). The organic layer was washed with brine (50 mL), dried over Na₂SO₄ and concentrated in vacuo to give 6-methyl-3-(4-methyltetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (160 mg, 68% yield).

LC-MS: *t*_R = 0.89 min (method 10), *m/z* = 248.3 [M + H]⁺.

Step 4: To a solution of 6-methyl-3-(4-methyltetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (200 mg, 0.81 mmol) and 1-(bromomethyl)-3-fluorobenzene (183 mg, 0.97 mmol) in anhydrous DMF (5 mL) was added K₂CO₃ (168 mg, 1.21 mmol). The mixture was stirred at 60-70°C for 16h. The mixture was cooled and diluted with water (20 mL), extracted with EtOAc(30 mL × 2). The organic layer was washed with brine (20 mL), dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=1:2) to give 7-(3-fluorobenzyl)-6-methyl-3-(4-methyltetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (90 mg, 31% yield) as a off-white solid.

¹H NMR (CDCl₃ 400 MHz): δ 7.94 (s, 1H), 7.33-7.28 (m, 1H), 7.02-6.91 (m, 4H), 5.22 (s, 2H), 3.85-3.80 (m, 2H), 3.73-3.67 (m, 2H), 2.45-2.41 (m, 2H), 2.17 (s, 3H), 1.86-1.79 (m, 2H),

mg, 92% yield) as a off-white solid.


LC-MS: $t_R = 0.98$ min (method 10), $m/z = 259.2$ $[M + H]^+$.

Step 4: To a solution of 4-(6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)tetrahydro-2H-pyran-4-carbonitrile (70 mg, 0.27 mmol) in anhydrous DMF (5 mL) was added 1-(bromomethyl)-3-fluoro-benzene (77 mg, 0.41 mmol) and K_2CO_3 (75 mg, 0.54 mmol). The mixture was stirred at 70-80°C for 2h. The mixture was cooled and filtered. The filtrate was purified by preparative LC-MS to give 4-(7-(3-fluorobenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)tetrahydro-2H-pyran-4-carbonitrile (65 mg, 65% yield).

^1H NMR ($CDCl_3$ 400 MHz): δ 7.97 (s, 1H), 7.34-7.31 (m, 1H), 7.16 (s, 1H), 7.02-6.92 (m, 3H), 5.26 (s, 2H), 4.13-4.10 (m, 2H), 3.97-3.91 (m, 2H), 2.50-2.43 (m, 2H), 2.35-2.32 (m, 2H), 2.24 (s, 3H).

LC-MS: $t_R = 2.69$ min (method 3), $m/z = 367.1$ $[M + H]^+$.

Example 28:



7-(3-fluorobenzyl)-3-(4-methoxytetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: To a solution of 3-bromo-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (500 mg, 2.2 mmol) and 1-(bromomethyl)-3-fluorobenzene (497 mg, 2.6 mmol) in DMF (5 mL) was added K_2CO_3 (605 mg, 4.4 mmol). The mixture was stirred at 60°C for 12 hours. The mixture was diluted with water (20 mL) and extracted with EtOAc (10 mL $\times$ 3). The combine organic layer was washed with water (10 mL $\times$ 2); dried over Na_2SO_4 and evaporated under vacuum to give 3-bromo-7-(3-fluorobenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (420 mg, 57% yield).

Step 2: To a solution of 3-bromo-7-(3-fluorobenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (200 mg, 0.6 mmol) in THF (10 mL) was added n-BuLi (0.31 mL, 0.77 mmol) at -78°C. The mixture was stirred at -78°C for 30 minutes. tetrahydro-4H-pyran-4-one (77 mg, 0.77 mmol) was added at -78°C. The mixture was stirred at -78°C for 1 hour. The mixture was quenched


with saturated aqueous NH_4Cl (0.5 mL) and evaporated under vacuum. The residue was dissolved in DCM (20 mL) and washed with water (10 mL). The organic layer was dried over Na_2SO_4 and evaporated. The residue was washed with EtOAc (3 mL) and filtered. The filter cake was dried under vacuum to give 7-(3-fluorobenzyl)-3-(4-hydroxytetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 47% yield).

Step 3: To a solution of 7-(3-fluorobenzyl)-3-(4-hydroxytetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (80 mg, 0.22 mmol) in THF (5 mL) was added NaH (60% in mineral oil, 13.4 mg, 0.36 mmol) at 0°C . The mixture was stirred at 20°C for 30 minutes. MeI (64 mg, 0.45 mmol) was added at 0°C . The mixture was stirred at 20°C for 11.5 hours. The mixture was quenched with saturated aqueous NH_4Cl (0.5 mL) and evaporated under vacuum. The residue was dissolved in DCM (10 mL) and washed with water (4 mL). The organic layer was dried over Na_2SO_4 and evaporated. The residue was purified by preparative TLC (EtOAc) to give 7-(3-fluorobenzyl)-3-(4-methoxytetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (25 mg, 30% yield).

^1H NMR (CDCl_3 , 400 MHz): δ 7.93 (s, 1H), 7.36 (s, 1H), 7.31 (dd, $J = 8.0$ Hz, $J = 14.0$ Hz, 1H), 7.03-6.92 (m, 3H), 5.23 (s, 2H), 3.91-3.80 (m, 4H), 3.07 (s, 3H), 2.38-2.31 (m, 2H), 2.17 (s, 3H), 2.14-2.10 (m, 2H).

LC-MS: $t_R = 2.73\text{min}$ (method 3), $m/z = 372.1$ $[\text{M} + \text{H}]^+$.

Example 29:



7-(3-fluorobenzyl)-3-(4-fluorotetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: To a solution of 3-bromo-7-(3-fluorobenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 0.3 mmol) in dry THF (5 mL) was added $n\text{-BuLi}$ (2.5 M, 0.15 mL) (2.5 M in $n\text{-hexane}$) dropwise. The mixture was stirred at -78°C for 0.5 hours. Then tetrahydro-4H-pyran-4-one (45 mg, 0.45 mmol) was added to the mixture. The mixture was stirred at -78°C for 2 hours. The mixture was quenched with saturated aqueous NH_4Cl (2 mL). The mixture

was extracted with DCM (20 mL × 2). The combined organic layer was washed with H₂O (10 mL), brine (10 mL), dried over Na₂SO₄, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (1%~10% MeOH in DCM) to give 7-(3-fluorobenzyl)-3-(4-hydroxytetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (40 mg, 38% yield).


LC-MS: t_R = 0.671 min (method 2), m/z = 358.1 [M + H]⁺.

Step 2: To a solution of 7-(3-fluorobenzyl)-3-(4-fluorotetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (40 mg, 111.9 μmol) in dry DCM (4 mL) was added DAST (diethylaminosulfur trifluoride) (28 mg, 170 μmol) at 0°C. The mixture was stirred at 0°C for 2 hours. Water (10 mL) was added to the mixture. The mixture was extracted with DCM (20 mL × 2). The combined organic layer was washed with H₂O (20 mL), dried over Na₂SO₄, filtered and concentrated to give the crude product. The crude product was purified by preparative TLC (DCM/MeOH=10/1) to give 7-(3-fluorobenzyl)-3-(4-fluorotetrahydro-2H-pyran-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (12.83 mg, 32% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.93 (s, 1H), 7.34-7.28 (m, 1H), 7.21 (s, 1H), 7.01-6.98 (m, 2H), 6.92 (d, J = 9.6 Hz, 1H), 5.24 (s, 2H), 4.00-3.89 (m, 4H), 2.57-2.42 (m, 2H), 2.23-2.18 (m, 5H).

LC-MS: t_R = 2.753 min (method 3), m/z = 360.1 [M + H]⁺.

Example 30:



7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-2-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 1 and 2:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (200 mg, 1.31 mmol) in DCM (8 mL) was added tetrahydro-2H-pyran-2-carboxylic acid (255 mg, 1.96 mmol) and HATU (894 mg, 2.35 mmol), Et₃N (264 mg, 2.61 mmol). The solution was stirred at 20-25°C for 1 hour. The mixture was diluted with water (30 mL), extracted with DCM (40 mL × 2). The organic layer was washed with brine (20 mL), dried over Na₂SO₄ and concentrated in vacuo.

The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=1:1) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-2-carboxamide (250 mg, 72% yield). LC-MS: t_R = 0.70 min (method 2), m/z = 266.2 $[M + H]^+$.

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)tetrahydro-2*H*-pyran-2-carboxamide (250 mg, 0.94 mmol) in dioxane (8 mL) was added POCl₃ (480 mg, 3.13 mmol). The solution was stirred at 90°C for 2h. The mixture was cooled and concentrated in vacuo. The residue was diluted with DCM (50 mL), washed with saturated aqueous NaHCO₃ (aq) (50 mL) and brine (50 mL). The organic layer was dried over Na₂SO₄ and concentrated in vacuo to give 8-methoxy-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazine (200 mg, 86% yield).

Step 3: To a solution of 8-methoxy-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazine (270 mg, 1.09 mmol) in dioxane (8 mL) was added 2M HCl(aq) (4 mL). The solution was stirred at 90°C for 1 hour. The mixture was concentrated in vacuo and added saturated aqueous NaHCO₃ (50 mL). The mixture was extracted with DCM (50 mL × 2). The organic layer was washed with brine (50 mL) and concentrated in vacuo to give 6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 79% yield).

Step 4: To a solution of 6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 0.86 mmol) in anhydrous DMF (10 mL) was added K₂CO₃ (237 mg, 1.71 mmol) and 1-(bromomethyl)-3-fluorobenzene (243 mg, 1.29 mmol). The mixture was stirred at 80°C for 24h. The mixture was cooled and diluted with water (100 mL), extracted with EtOAc (50 mL × 3). The organic layer was washed with brine (50 mL), dried over Na₂SO₄ and concentrated in vacuo to give 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (130 mg, 44% yield).

Step 5: 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (130 mg, 380.8 μmol) was purified by SFC.

7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one, **stereoisomer 1** (35 mg, 27% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.91 (s, 1H), 7.32-7.28 (m, 1H), 7.19(s, 1H), 6.98-6.94 (m, 2H), 6.89 (d, J = 9.6 Hz, 1H), 5.23 (s, 2H), 4.76 (t, J = 6.4 Hz, 1H), 4.06 (d, J = 10.8 Hz, 1H), 3.67 (t, J = 10.8 Hz, 1H), 2.23 (s, 3H), 2.17-2.05 (m, 3H), 1.74-1.68 (m, 3H).


LC-MS: t_R = 2.33 min (method 3), m/z = 342.1 $[M + H]^+$. SFC: t_R = 5.478 min, ee% = 99.90%. $[\alpha]_D^{20}$ +16.00 (c = 0.10, CHCl₃).

7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one, stereoisomer **2** (33 mg, yield: 35%) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.92 (s, 1H), 7.32-7.28 (m, 1H), 7.19(s, 1H), 6.98-6.94 (m, 2H), 6.89 (d, *J* = 9.2 Hz, 1H), 5.23 (s, 2H), 4.77 (t, *J* = 6.8 Hz, 1H), 4.06 (d, *J* = 10.8 Hz, 1H), 3.67 (t, *J* = 10.8 Hz, 1H), 2.17 (s, 3H), 2.12-2.02 (m, 3H), 1.74-1.68 (m, 3H).

LC-MS: *t*_R = 2.33 min (method 3), *m/z* = 342.1 [M + H]⁺. SFC: *t*_R = 5.789 min, ee% = 98.92%. [α]_D²⁰ -23.33 (c = 0.10, CHCl₃).

Example 31:



7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one, stereoisomer **1** and **2**:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (200 mg, 1.3 mmol) in dry DCM (10 mL) was added tetrahydrofuran-3-carboxylic acid (228 mg, 2.0 mmol), Et₃N (265 mg, 2.6 mmol) and HATU (747 mg, 2.0 mmol). The mixture was stirred at 15°C for 16 hours. Water (10 mL) was added to the mixture. The organic layer was washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (10%~100% ethyl acetate in petroleum ether) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)tetrahydrofuran-3-carboxamide (200 mg, 61% yield).

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)tetrahydrofuran-3-carboxamide (300 mg, 1.19 mmol) in dioxane (5 mL) was added POCl₃ (366 mg, 2.39 mmol). The mixture was heated at 90°C for 2 hours. The mixture was cooled to 15°C and adjusted to pH=8 by saturated aqueous NaHCO₃. The aqueous layer was extracted with DCM (20 mL × 2). The combined organic layer was washed with H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give 8-methoxy-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazine (300 mg).

Step 3: To a solution of 8-methoxy-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazine (300 mg, 1.29 mmol) in dioxane (5 mL) was added 2 N HCl (2 mL). The mixture was heated at 90°C for 1 hour. The mixture was cooled to 15°C and extracted with DCM (20 mL × 2). The combined organic layer was washed with H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give 6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one (210 mg, 74% yield).

Step 4: To a solution of 6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one (200 mg, 912.24 μmol) in dry DMF (5 mL) was added 1-(bromomethyl)-3-fluorobenzene (259 mg, 1.37 mmol) and K₂CO₃ (252 mg, 1.82 mmol). The mixture was stirred at 60°C for 16 hours. The mixture was concentrated and the residue was dissolved in DCM (20 mL) and H₂O (10 mL). The aqueous layer was extracted with DCM (20 mL × 2). The combined organic layer was washed with H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (10%~100% ethyl acetate in petroleum ether) to give 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one (40 mg, 13% yield).

Step 5: 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one (40 mg, 122.2 μmol) was purified by SFC to give 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one, **stereoisomer 1** (16.43 mg, 41% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.90 (s, 1H), 7.31-7.28 (m, 1H), 6.98-6.93 (m, 2H), 6.89 (d, *J* = 9.2 Hz, 1H), 6.79 (s, 1H), 5.21 (s, 2H), 4.19-4.17 (m, 1H), 4.12-4.09 (m, 1H), 4.05-3.95 (m, 2H), 3.66-3.62 (m, 1H), 2.40 (q, *J* = 7.2 Hz, 2H), 2.16 (s, 3H).

LC-MS: *t*_R = 1.964 min (method 3), *m/z* = 328.0 [M + H]⁺. SFC: *t*_R = 4.503 min, ee% = 99.8%; [α]_D²⁰ +14.7 (c = 0.10, DCM).

7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one, **stereoisomer 2** (15.58 mg, 38% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.90 (s, 1H), 7.29-7.27 (m, 1H), 6.98-6.92 (m, 2H), 6.89 (d, *J* = 9.6 Hz, 1H), 6.79 (s, 1H), 5.21 (s, 2H), 4.19 (t, *J* = 8.4 Hz, 1H), 4.11-4.09 (m, 1H), 4.05-4.03 (m, 1H), 3.97-3.95 (m, 1H), 3.66-3.62 (m, 1H), 2.38 (q, *J* = 7.6 Hz, 2H), 2.16 (s, 3H).

LC-MS: *t*_R = 1.957 min (method 3), *m/z* = 328.0 [M + H]⁺. SFC: *t*_R = 4.779 min, ee% = 96%; [α]_D²⁰ -14.0 (c = 0.10, DCM).

Step 4: To a solution of 6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (260 mg, 1.1 mmol) in anhydrous DMF (10 mL) was added K_2CO_3 (308 mg, 2.2 mmol) and 1-(bromomethyl)-3-fluorobenzene (316 mg, 1.7 mmol). The mixture was stirred at 70-80°C for 2h. The mixture was cooled and diluted with water (50 mL), extracted with
 5 EtOAc (50 mL $\times$ 2). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated in vacuo to give 7-(3-fluorobenzyl)-6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (160 mg, 42% yield).

Step 5: 7-(3-fluorobenzyl)-6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (160 mg, 0.47 mmol) was separated by SFC to give 7-(3-fluorobenzyl)-6-methyl-
 10 3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, **stereoisomer 1** (37 mg, 23% yield) as a off-white solid. ^1H NMR ($CDCl_3$ 400 MHz): δ 7.88 (s, 1H), 7.32-7.26 (m, 1H), 6.94-6.82 (m, 4H), 5.21 (s, 2H), 4.33 (d, J = 8.8 Hz, 1H), 4.07-4.04 (m, 1H), 4.00-3.98 (m, 1H), 3.87 (d, J = 8.8 Hz, 1H), 2.62-2.57 (m, 1H), 2.16 (s, 3H), 2.14-2.09 (m, 1H), 1.60 (s, 3H), . LC-MS: t_R = 2.47 min (method 3), m/z = 342.1 $[M + H]^+$. **SFC**: t_R = 4.91 min, ee% >
 15 99%. $[\alpha]_D^{20}$ = +5.0 (c = 0.10, MeOH).


7-(3-fluorobenzyl)-6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, **stereoisomer 2** (44 mg, 27% yield) as a off-white solid.

^1H NMR ($CDCl_3$ 400 MHz): δ 7.82 (s, 1H), 7.24-7.20 (m, 1H), 6.94-6.82 (m, 4H), 5.15 (s, 2H), 4.27 (d, J = 8.8 Hz, 1H), 4.01-3.99 (m, 1H), 3.96-3.92 (m, 1H), 3.81 (d, J = 8.8 Hz, 1H), 2.57-
 20 2.51 (m, 1H), 2.10 (s, 3H), 2.09-2.04 (m, 1H), 1.54 (s, 3H).

LC-MS: t_R = 2.47 min (method 3), m/z = 342.1 $[M + H]^+$.

SFC: t_R = 5.33 min, ee% > 99%. $[\alpha]_D^{20}$ = -3.0 (c = 0.10, MeOH).

Example 33:



7-(3-fluorobenzyl)-6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: To a solution of 1-methylcyclopropane-1-carboxylic acid (500 mg, 4.99 mmol) in DCM (2 mL) was added $(COCl)_2$ (3.17 g, 24.95 mmol). The solution was stirred at 40°C for

2h. The reaction mixture was concentrated in vacuo to give 1-methylcyclopropane-1-carbonyl chloride (500 mg, 85% yield).

Step 2: A solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (100 mg, 0.65 mmol) in anhydrous DCM (3 mL) was cooled to 0°C. Then a solution of 1-methylcyclopropane-1-carbonyl chloride (100 mg, 0.85 mmol) in anhydrous DCM (2 mL) was added dropwise and stirred at 0°C for 15 min. The mixture was diluted with DCM (20 mL), washed with saturated aqueous NaHCO₃ (20 mL), brine (20 mL) and dried over Na₂SO₄. The organic layer was concentrated in vacuo to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-1-methylcyclopropane-1-carboxamide (120 mg, 78% yield).

LC-MS: t_R = 0.66 min (method 2), m/z = 236.1 [M + H]⁺.

Step 3: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-1-methylcyclopropane-1-carboxamide (120 mg, 0.51 mmol) in dioxane (5 mL) was added POCl₃ (590 mg, 3.85 mmol). The solution was stirred at 80-90°C for 2h. The mixture was cooled and slowly added into water (50 mL), extracted with EtOAc (30 mL × 2). The organic layer was washed with brine (20 mL), dried over Na₂SO₄ and concentrated in vacuo to give 8-methoxy-6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-*a*]pyrazine (80 mg, 72% yield).

Step 4: To a solution of 8-methoxy-6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-*a*]pyrazine (80 mg, 0.37 mmol) in dioxane (5 mL) was added 2M HCl (aq) (2 mL). The solution was stirred at 80-90°C for 1 hour. The mixture was concentrated in vacuo and added saturated aqueous NaHCO₃ (50 mL). The mixture was extracted with DCM (50 mL × 2). The organic layer was washed with brine (20 mL) and concentrated in vacuo to give 6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (70 mg, 94% yield).

Step 5: To a solution of 6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (70 mg, 0.34 mmol) in anhydrous DMF (4 mL) was added 1-(bromomethyl)-3-fluoro-benzene (98 mg, 0.52 mmol) and K₂CO₃ (95 mg, 0.69 mmol). The mixture was stirred at 70-80°C for 1 hour. The mixture was cooled and filtered. The filtrate was purified by preparative LC-MS to give 7-(3-fluorobenzyl)-6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (35 mg, 32% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.84 (s, 1H), 7.33-7.39 (m, 1H), 7.01-6.89 (m, 4H), 5.23 (s, 2H), 2.20 (s, 3H), 1.60 (s, 3H), 1.11-1.09 (m, 2H), 0.88-0.85 (m, 2H).


LC-MS: t_R = 2.34 min (method 3), m/z = 312.1 [M + H]⁺.

2h. The mixture was cooled and filtered. The filtrate was purified by preparative LC-MS to give 3-(2,2-difluorocyclopropyl)-7-(3-fluorobenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (40 mg, 27% yield) as a white solid.

¹H NMR (CDCl₃ 400 MHz): δ 7.90 (s, 1H), 7.30-7.28 (m, 1H), 6.98-6.84 (m, 4H), 5.28-5.16 (m, 2H), 2.80-2.73 (m, 1H), 2.38-2.37 (m, 1H), 2.19 (s, 3H), 2.05-2.04 (m, 1H).

LC-MS: *t*_R = 2.67 min (method 3), *m/z* = 334.1 [M + H]⁺.

Example 35:



10 7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 1, 2, 3 and 4:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (100 mg, 652.8 μmol) and 2-methylcyclopropane-1-carboxylic acid (98 mg, 979.2 μmol) in DCM (5 mL) was added HATU (446.8 mg, 1.2 mmol) and triethylamine (132.1 mg, 1.3 mmol). The mixture was stirred at 24°C for 16h. The mixture was diluted with DCM (20 mL) and washed with water (15 mL). The aqueous layer was extracted with DCM (2*30 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by preparative TLC (ethyl acetate) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methylcyclopropane-1-carboxamide (150 mg, 95% yield).

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methylcyclopropane-1-carboxamide (150 mg, 636 μmol) in dioxane (5 mL) was added POCl₃ (400 mg, 2.6 mmol). The mixture was stirred at 90°C for 2h. The mixture was cooled down to 25°C, neutralized with saturated aqueous NaHCO₃ and extracted with ethyl acetate (3 × 30 mL). The combined organic layer was washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated to give the crude 8-methoxy-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazine (130 mg, 94% yield). This crude product was used directly for the next step.

Step 3: A solution of 8-methoxy-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazine (120 mg, 552.3 μmol) in dioxane (5 mL) and HCl (2 M, 2 mL) was stirred at 80°C for 1 hour. The mixture was cooled to 25°C, neutralized with saturated aqueous NaHCO₃ and extracted with

ethyl acetate (3 × 20 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give the crude 6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one (100 mg, 89% yield). This crude product was used directly for the next step.

- 5 Step 4: To a solution of 6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one (100 mg, 492.0 μmol) and 1-(bromomethyl)-3-fluorobenzene (139.5 mg, 738.0 μmol) in DMF (5 mL) was added K₂CO₃ (136 mg, 984 μmol). The mixture was stirred at 60-70°C for 16h. The mixture was cooled to 25°C, diluted with water (15 mL), extracted with ethyl acetate (3 × 30 mL). The combined organic layer was washed with brine (30 mL), dried over
10 Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography on silica gel (eluent of 0%~50% ethyl acetate in petroleum ether) to give 7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one (80 mg, 52% yield).

Step 5: 7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one (100 mg, 321.2 μmol) was purified by SFC.

- 15 7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one, **stereoisomer 1** (35 mg, 27% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.83 (s, 1H), 7.33-7.29 (m, 1H), 7.00-6.94(m, 2H), 6.91-6.89 (m, 2H), 5.23 (s, 2H), 2.19 (s, 3H), 1.59-1.55 (m, 1H), 1.54-1.50 (m, 1H), 1.35-1.30 (m, 1H), 1.27 (d, *J* = 6.0 Hz, 3H), 0.89-0.88 (m, 1H).

- 20 LC-MS: *t*_R = 2.03 min (method 3), *m/z* = 312.1 [M + H]⁺. SFC: *t*_R = 4.466 min, ee% > 99%. [α]_D²⁰ +29.3 (c = 0.10, CHCl₃).

7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one, **stereoisomer 2** (26 mg, 26% yield) was obtained.

- 25 ¹H NMR (CDCl₃ 400 MHz): δ 7.84 (s, 1H), 7.32-7.29 (m, 1H), 7.00-6.96(m, 2H), 6.90-6.89 (m, 2H), 5.23 (s, 2H), 2.19 (s, 3H), 1.62-1.60 (m, 1H), 1.55-1.53 (m, 1H), 1.36-1.34 (m, 1H), 1.27 (d, *J* = 5.6 Hz, 3H), 0.91-0.89 (m, 1H).

LC-MS: *t*_R = 2.02 min (method 3), *m/z* = 312.1 [M + H]⁺. SFC: *t*_R = 5.227 min, ee% > 99%. [α]_D²⁰ -15.0 (c = 0.10, CHCl₃).

- 30 7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one, **stereoisomer 3** (8.0 mg, 8% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.89 (s, 1H), 7.33-7.30 (m, 1H), 7.01-6.91(m, 4H), 5.29 (d, *J*= 16.0 Hz, 1H), 5.17 (d, *J*= 16.0 Hz, 1H), 2.19 (s, 3H), 2.01-1.97(m, 1H), 1.39-1.27 (m, 1H), 1.23-1.20 (m, 2H), 0.91 (d, *J*= 6.0 Hz, 3H).

LC-MS: *t*_R = 1.98 min (method 3), *m/z* = 312.1 [M + H]⁺. SFC: *t*_R = 6.995 min, ee% > 99%.


5 [α]_D²⁰ +37.0 (c = 0.10, CHCl₃).

7-(3-fluorobenzyl)-6-methyl-3-(2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7*H*)-one, **stereoisomer 4** (9.0 mg, 9% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.86 (s, 1H), 7.29-7.26 (m, 1H), 6.99-6.88(m, 4H), 5.27 (d, *J*= 16.0 Hz, 1H), 5.15 (d, *J*= 16.0 Hz, 1H), 2.17 (s, 3H), 1.97-1.95(m, 1H), 1.58-1.53 (m, 1H),
10 1.23-1.20 (m, 2H), 0.91 (d, *J*= 6.0 Hz, 3H).

LC-MS: *t*_R = 1.98 min (method 3), *m/z* = 312.1 [M + H]⁺. SFC: *t*_R = 8.704 min, ee% > 99%.
[α]_D²⁰ -66.7 (c = 0.10, CHCl₃).

Example 36:



15 **7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one, stereoisomer 1, 2, 3 and 4:**

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (300 mg, 2.0 mmol), 2-methyltetrahydrofuran-3-carboxylic acid (382 mg, 2.9 mmol) in DCM (10 mL) was added
20 HATU (1.3 g, 3.5 mmol) and triethylamine (396 mg, 3.9 mmol). The mixture was stirred at 24°C for 16h. The mixture was diluted with DCM (30 mL) and washed with water (20 mL). The aqueous layer was extracted with DCM (2 × 30 mL). The combined organic layer was washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography on silica gel (0%~70% ethyl acetate in petroleum ether)
25 to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methyltetrahydrofuran-3-carboxamide (350 mg, 62% yield).

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methyltetrahydrofuran-3-carboxamide (350 mg, 1.3 mmol) in dioxane (5 mL) was added

POCl₃ (720 mg, 4.7 mmol). The mixture was stirred at 90°C for 2h. The mixture was cooled to 25°C, neutralized with saturated aq. NaHCO₃ and extracted with ethyl acetate (2 × 30 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give crude 8-methoxy-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazine (320 mg). The crude was used directly for the next step.

Step 3: A solution of 8-methoxy-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazine (320 mg, 1.3 mmol) in dioxane (5 mL) and HCl (2 M, 2 mL) was stirred at 80°C-90°C for 21.5 hours. The mixture was cooled down to 25°C, neutralized with saturated aq. NaHCO₃, extracted with DCM (3 × 30 mL). The combined organic layer was washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated to give crude 6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (300 mg). The crude product was used directly for the next step.

Step 4: To a solution of 6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (250 mg, 1.1 mmol) in DMF (8 mL) was added 1-(bromomethyl)-3-fluorobenzene (304 mg, 1.6 mmol) and K₂CO₃ (296 mg, 2.1 mmol). The mixture was stirred at 60-70°C for 16h. The mixture was cooled down to 25°C and diluted with water (15 mL), extracted with ethyl acetate (3 × 30 mL). The combined organic layer was washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography on silica gel (eluent of 0%~30% ethyl acetate in petroleum ether) to give 7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (170 mg, 47% yield).

Step 5: 7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (220 mg, 644 μmol) was purified by SFC.

7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, **stereoisomer 1** (20 mg, 9% yield) .

¹H NMR (CDCl₃ 400 MHz): δ 7.95 (s, 1H), 7.34-7.28 (m, 1H), 7.01-6.89 (m, 3H), 6.79 (s, 1H), 5.23 (s, 2H), 4.35-4.30 (m, 1H), 4.13-4.07 (m, 2H), 3.12-3.06 (m, 1H), 2.50-2.37 (m, 2H), 2.19 (s, 3H), 1.35 (d, *J* = 6.0 Hz, 3H).

LC-MS: *t*_R = 2.12 min (method 3), *m/z* = 342.1 [M + H]⁺.

SFC: *t*_R = 4.812 min, ee% > 99%. [*α*]_D²⁰ -24.3 (c = 0.10, CHCl₃).

7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, **stereoisomer 2** (10 mg, 5% yield) .

^1H NMR (CDCl_3 400 MHz): δ 7.93 (s, 1H), 7.31-7.26 (m, 1H), 6.99-6.87 (m, 3H), 6.76 (s, 1H), 5.21 (s, 2H), 4.32-4.27 (m, 1H), 4.10-4.04 (m, 2H), 3.10-3.03 (m, 1H), 2.48-2.35 (m, 2H), 2.17 (s, 3H), 1.33 (d, J = 6.4 Hz, 3H).

LC-MS: t_R = 2.07 min (method 3), m/z = 342.1 $[\text{M} + \text{H}]^+$.

5 SFC: t_R = 5.088 min, ee% = 97.9%. $[\alpha]_D^{20}$ +10.3 (c = 0.10, CHCl_3).

7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, **stereoisomer 3** (38.0 mg, 17% yield).


10 ^1H NMR (CDCl_3 400 MHz): δ 7.95 (s, 1H), 7.31-7.29 (m, 1H), 7.01-6.89 (m, 3H), 6.83 (s, 1H), 5.29 (d, J = 16.4 Hz, 1H), 5.17 (d, J = 16.4 Hz, 1H), 4.35-4.28 (m, 2H), 3.91-3.87 (m, 1H), 3.71-3.67 (m, 1H), 2.72-2.67 (m, 1H), 2.44-2.40 (m, 1H), 2.18 (s, 3H), 0.90 (d, J = 6.4 Hz, 3H).

LC-MS: t_R = 2.02 min (method 3), m/z = 342.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 5.516 min, ee% > 99%. $[\alpha]_D^{20}$ +46.3 (c = 0.10, CHCl_3).

15 7-(3-fluorobenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, **stereoisomer 4** (30.0 mg, 14% yield). ^1H NMR (CDCl_3 400 MHz): δ 7.95 (s, 1H), 7.33-7.30 (m, 1H), 7.01-6.90 (m, 3H), 6.83 (s, 1H), 5.29 (d, J = 16.4 Hz, 1H), 5.17 (d, J = 16.4 Hz, 1H), 4.37-4.28 (m, 2H), 3.91-3.87 (m, 1H), 3.71-3.69 (m, 1H), 2.72-2.68 (m, 1H), 2.45-2.40 (m, 1H), 2.18 (s, 3H), 0.90 (d, J = 6.8 Hz, 3H).

20 LC-MS: t_R = 1.98 min (method 3), m/z = 342.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 6.304 min, ee% > 99%. $[\alpha]_D^{20}$ -47.0 (c = 0.10, CHCl_3).

Example 37:



25 7-(3-fluorobenzyl)-3-(c/s-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one, **stereoisomer 1 and 2**:

Step 1: A solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (400 mg, 2.1 mmol) and triethylamine (662 mg, 6.5 mmol) in DCM (8 mL) was cooled to 0°C, 2-

fluorocyclopropane-1-carbonyl chloride (251 mg, 2.1 mmol) was added dropwise and the mixture was stirred at 0°C for 0.5h. The mixture was diluted with water (10 mL), extracted with DCM (20 mL × 2). The organic layer was washed with brine (20 mL), dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by preparative TLC (petroleum

- 5 ether:ethyl acetate=1:1) to give *cis*-2-fluoro-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)cyclopropane-1-carboxamide (200 mg, 40% yield) and (1*S*,2*R*)-2-fluoro-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)cyclopropane-1-carboxamide (200 mg, 40% yield) all.

- Step 2: To a solution of *cis*-2-fluoro-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)cyclopropane-1-carboxamide (260 mg, 1.1 mmol) in dioxane (10 mL) was added
10 POCl₃ (500 mg, 3.3 mmol). The solution was stirred at 80-90°C for 2h. The mixture was concentrated in vacuo and diluted with NaHCO₃ (30 mL), extracted with DCM (50 mL × 2). The organic layer was washed with brine (30 mL), dried over Na₂SO₄ and concentrated in vacuo to give 3-(*cis*-2-fluorocyclopropyl)-8-methoxy-6-methylimidazo[1,5-*a*]pyrazine (220 mg, 91% yield).

- 15 Step 3: A solution of 3-(*cis*-2-fluorocyclopropyl)-8-methoxy-6-methylimidazo[1,5-*a*]pyrazine (220 mg, 994 μmol) in 2N HCl (aq) (5 mL) and dioxane (10 mL) was stirred at 80-90°C for 1 hour. The mixture was concentrated in vacuo. The residue was diluted with NaHCO₃(aq) (30 mL), extracted with DCM (30 mL × 3). The organic layer was washed with brine (30 mL), dried over Na₂SO₄ and concentrated in vacuo to give 3-(*cis*-2-fluorocyclopropyl)-6-
20 methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 97% yield).

- Step 4: To a solution of 3-(*cis*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 965 μmol) in anhydrous DMF (10 mL) was added K₂CO₃ (133 mg, 965 μmol) and 1-(bromomethyl)-3-fluorobenzene (274 mg, 1.5 mmol). The mixture was stirred at 60-70°C for 16h and 80°C for 21h. The mixture was cooled and filtered and the filtrate was purified by
25 preparative LC-MS to give 7-(3-fluorobenzyl)-3-(*cis*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (80 mg, 26% yield).

Step 5: 7-(3-fluorobenzyl)-3-(*cis*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (80 mg, 253 μmol) was separated by SFC.

- 7-(3-fluorobenzyl)-3-(*cis*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one,
30 **stereoisomer 1** (25 mg, 30% yield).


¹H NMR (CDCl₃ Varian_H_400 MHz): δ 7.92 (s, 1H), 7.32-7.26 (m, 1H), 7.00-6.88 (m, 4H), 5.29-5.15 (m, 2H), 5.02-4.84 (m, 1H), 2.18 (s, 3H), 2.12-1.99 (m, 2H), 1.44-1.25 (m, 1H).

LC-MS: t_R = 2.00 min (method 8), m/z = 316.1 $[M + H]^+$. SFC: t_R = 6.53 min, ee% > 99%. $[\alpha]_D^{20}$ -84.00 (c = 0.10, MeOH).

7-(3-fluorobenzyl)-3-(*cis*-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one, **stereoisomer 2** (15 mg, 18% yield). ^1H NMR ($CDCl_3$ Varian_H_400 MHz): 7.93 (s, 1H),
 5 7.33-7.30 (m, 1H), 7.00-6.90 (m, 4H), 5.31-5.17 (m, 2H), 5.04-4.86 (m, 1H), 2.20 (s, 3H),
 2.12-2.00 (m, 2H), 1.46-1.42 (m, 1H).

LC-MS: t_R = 1.99 min (method 8), m/z = 316.1 $[M + H]^+$. SFC: t_R = 5.06 min, ee% > 99%. $[\alpha]_D^{20}$ +75.00 (c = 0.10, MeOH).

10 Example 38:



7-(3-fluorobenzyl)-3-(*trans*-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one, stereoisomer 1 and 2:

Step 1: To a solution of *trans*-2-fluoro-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)cyclopropane-1-carboxamide (280 mg, 1.1 mmol) in dioxane (10 mL) was added
 15 $POCl_3$ (540 mg, 3.5 mmol). The solution was stirred at 80-90°C for 2h. The mixture was concentrated in vacuo and diluted with $NaHCO_3$ (30 mL), extracted with DCM (50 mL $\times$ 2). The organic layer was washed with brine (30 mL), dried over Na_2SO_4 and concentrated in vacuo to give 3-(*trans*-2-fluorocyclopropyl)-8-methoxy-6-methylimidazo[1,5-a]pyrazine (240
 20 mg, 93% yield).

Step 2: A solution of 3-(*trans*-2-fluorocyclopropyl)-8-methoxy-6-methylimidazo[1,5-a]pyrazine (240 mg, 1.1 mmol) in 2N HCl (aq) (5 mL) and dioxane (10 mL) was stirred at 80-90°C for 1 hour. The mixture was concentrated in vacuo. The residue was diluted with $NaHCO_3$ (aq) (30 mL), extracted with DCM (30 mL $\times$ 3). The organic layer was washed with brine (30 mL),
 25 dried over Na_2SO_4 and concentrated in vacuo to give 3-(*trans*-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one (220 mg, 98% yield).

Step 3: To a solution of 3-(*trans*-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one (220 mg, 1.1 mmol) in anhydrous DMF (10 mL) was added K_2CO_3 (293 mg, 2.1 mmol)

and 1-(bromomethyl)-3-fluorobenzene (300 mg, 1.6 mmol). The mixture was stirred at 60-70°C for 16h and 80°C for 21h. The mixture was cooled and filtered and the filtrate was purified by preparative LC-MS to give 7-(3-fluorobenzyl)-3-(*trans*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (250 mg, 77% yield).

- 5 Step 4: 7-(3-fluorobenzyl)-3-(*trans*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (250 mg, 793 μ mol) was separated by SFC.

7-(3-fluorobenzyl)-3-(*trans*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one, **stereoisomer 1** (15 mg, 17% yield).

10 ^1H NMR (CDCl_3 400 MHz): δ 7.81 (s, 1H), 7.31-7.27 (m, 1H), 6.98-6.86 (m, 4H), 5.22 (s, 2H), 5.01-4.84 (m, 1H), 2.43-2.38 (m, 1H), 2.19 (s, 3H), 1.72-1.65 (m, 1H), 1.58-1.55 (m, 1H).


LC-MS: t_R = 2.24 min (method 8), m/z = 316.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 2.83 min, ee% > 99%. $[\alpha]_D^{20}$ +29.00 (c = 0.10, MeOH).

15 7-(3-fluorobenzyl)-3-(*trans*-2-fluorocyclopropyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one, **stereoisomer 2** (45 mg, 17% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.81 (s, 1H), 7.29-7.28 (m, 1H), 6.98-6.86 (m, 4H), 5.22 (s, 2H), 5.01-4.84 (m, 1H), 2.45-2.38 (m, 1H), 2.19 (s, 3H), 1.71-1.64 (m, 1H), 1.59-1.54 (m, 1H).

20 LC-MS: t_R = 2.23 min (method 8), m/z = 316.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 3.69 min, ee% > 99%. $[\alpha]_D^{20}$ -31 (c = 0.10, MeOH).

Example 39:



25 **7-(4-cyclopropoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:**

Step 1: A solution of 1-bromo-4-cyclopropoxybenzene (900 mg, 4.22 mmol) in THF (20 mL, anhydrous) was added n-BuLi (2.5 M, 2.6 mL) at -78°C and stirred at -78°C for 2 hours under N₂. Then, thereto was added dropwise DMF (926 mg, 12.66 mmol, anhydrous) at -78°C and it was stirred for 2 hours. The solution was quenched with NH₄Cl (aq. 1mL) at -78°C and stirred at 0°C for 0.5 hour. The mixture was diluted with ethyl acetate (10 mL). The mixture was filtered and the filtrate was concentrated under vacuum. The residue was diluted with ethyl acetate (30 mL), washed with brine (3 × 15 mL). The organic layer was dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum to give 4-cyclopropoxybenzaldehyde (700 mg). LC-MS: *t_R* = 0.800 min (method 2), *m/z* = 162.8 [M + H]⁺.


Step 2: A solution of 4-cyclopropoxybenzaldehyde (700 mg) in MeOH (15 mL, anhydrous) was added NaBH₄ (319 mg, 8.44 mmol) at 0°C and stirred at 0°C for 1 hour. The solution was quenched with saturated aqueous NH₄Cl (aq. 0.5mL). The mixture was concentrated under vacuum. The residue was purified by silica gel chromatography (petroleum ether/ethyl acetate=10/1, 1/1) to give (4-cyclopropoxyphenyl)methanol (540 mg, 78% yield).

Step 3: A solution of (4-cyclopropoxyphenyl)methanol (500 mg, 3.1 mmol) and Et₃N (617 mg, 6.1 mmol) in anhydrous DCM (10 mL) was cooled to 0°C, then MsCl (1.41 g, 12.3 mmol) was added dropwise. The solution was stirred at 0°C for 0.5h. The mixture was diluted with water (50 mL), extracted with DCM (50 mL × 2). The organic layer was washed with brine (30 mL), dried over Na₂SO₄ and concentrated in vacuo to afford 4-cyclopropoxybenzyl methanesulfonate (600 mg, 81% yield).

Step 4: A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (250 mg, 1.07 mmol), 4-cyclopropoxybenzyl methanesulfonate (311 mg, 1.28 mmol) and Cs₂CO₃ (698 mg, 2.14 mmol) in DMF (6.0 mL, anhydrous) was stirred at 60°C for 12 hours. The mixture was filtered and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (petroleum ether/ethyl acetate=10/1, 0/1) to give 7-(4-cyclopropoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (320 mg, 77% yield).

¹H NMR (CDCl₃, 400 MHz): δ 7.92 (s, 1H), 7.15 (d, *J* = 8.8 Hz, 2H), 6.99 (d, *J* = 8.8 Hz, 2H), 6.73 (s, 1H), 5.17 (s, 2H), 4.12 (d, *J* = 10.4 Hz, 2H), 3.71 - 3.69 (m, 1H), 3.67 - 3.55 (m, 2H), 3.09 - 3.07 (m, 1H), 2.21 (s, 3H), 2.18 - 2.03 (m, 2H), 1.88 (d, *J* = 13.2 Hz, 2H), 0.79 - 0.71 (m, 4H).

LC-MS: *t_R* = 2.154 min (method 3), *m/z* = 380.1 [M + H]⁺.


Example 40:

7-(4-(difluoromethoxy)benzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one

A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (400 mg, 1.7 mmol), 1-(bromomethyl)-4-(difluoromethoxy)benzene (608 mg, 2.6 mmol), Cs₂CO₃ (1.11 g, 3.4 mmol) in DMF (50 mL) was stirred at 60 °C for 12 hours. The reaction mixture was concentrated. The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=5:1~0:1) to give 7-(4-(difluoromethoxy)benzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (360 mg, yield: 52%).

¹H NMR (CDCl₃, 400 MHz): δ 7.91 (s, 1H), 7.20 (d, *J* = 9.2 Hz, 2H), 7.07 (d, *J* = 8.0 Hz, 2H), 6.73 (s, 1H), 6.46 (t, *J* = 73.6 Hz, 1H), 5.19 (s, 2H), 4.11 (d, *J* = 10.8 Hz, 2H), 3.56 (td, *J* = 12.0, 2.0 Hz, 2H), 3.10 - 3.04 (m, 1H), 2.17 - 2.06 (m, 5H), 1.86 (d, *J* = 13.2 Hz, 2H).

LC-MS: *t*_R = 1.85 min (method 3), *m/z* = 390.1 [M + H]⁺.

Example 41:

6-methyl-3-(tetrahydro-2H-pyran-4-yl)-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-a]pyrazin-8(7H)-one:


A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (400 mg, 1.7 mmol), 1-(bromomethyl)-4-(trifluoromethoxy)benzene (654 mg, 2.6 mmol) and

Cs_2CO_3 (1.11 g, 3.4 mmol) in DMF (50 mL) was stirred at 60 °C for 12h. The reaction mixture was concentrated. The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=5:1~0:1) to give 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (300 mg, yield: 41%).

- 5 ^1H NMR (CDCl_3 , 400 MHz): δ 7.91 (s, 1H), 7.25 - 7.22 (m, 2H), 7.17 - 7.15 (m, 2H), 6.74 (s, 1H), 5.21 (s, 2H), 4.11 (d, J = 10.4 Hz, 2H), 3.56 (td, J = 11.2, 2.0 Hz, 2H), 3.10 - 3.04 (m, 1H), 2.17 - 2.06 (m, 5H), 1.86 (d, J = 13.6 Hz, 2H).

LC-MS: t_R = 2.05 min (method 3), m/z = 408.1 $[\text{M} + \text{H}]^+$.

10 Example 42:



7-(4-(cyclopropylmethoxy)benzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

- Step 1: A mixture of 4-hydroxybenzaldehyde (1.0 g, 8.19 mmol), (bromomethyl)cyclopropane (1.33 g, 9.83 mmol) and K_2CO_3 (2.26 g, 16.38 mmol) in DMF (10.0 mL, anhydrous) was stirred at 20°C for 12 hours. The solution was diluted with water (20 mL). The aqueous phase was extracted with ethyl acetate (60 mL $\times$ 3). The combined organic phase was washed with brine (20 mL $\times$ 1), dried over anhydrous Na_2SO_4 , filtered and concentrated in vacuum. The residue was purified by silica gel chromatography (petroleum ether/ethyl acetate=10/1, 1/1) to give 4-(cyclopropylmethoxy)benzaldehyde (1.3 g, 87% yield).

- Step 2: A solution of 4-(cyclopropylmethoxy)benzaldehyde (1.3 g, 7.4 mmol) in MeOH (30 mL) was cooled to 0°C, then NaBH_4 (558 mg, 14.8 mmol) was added and the mixture was stirred at 0°C for 1 hour. The mixture was quenched with sat. brine (aq) (50 mL), extracted with EtOAc (50 mL $\times$ 2). The organic layer was washed with brine, dried over Na_2SO_4 and concentrated in vacuo to afford (4-(cyclopropylmethoxy)phenyl)methanol (1.21 g, 92% yield).

Step 3. A solution of (4-(cyclopropylmethoxy)phenyl)methanol (800 mg, 4.5 mmol) and Et_3N (907 mg, 9.0 mmol) in DCM (10 mL) was cooled to 0°C, then MsCl (617 mg, 5.4 mmol) was


added dropwise. The solution was stirred at 0°C for 0.5h. The mixture was diluted with water (50 mL), extracted with DCM (50 mL × 2). The organic layer was washed with brine (30 mL), dried over Na₂SO₄ and concentrated in vacuo to afford 4-(cyclopropylmethoxy)benzyl methanesulfonate (760 mg, 66% yield).

- 5 Step 4: A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (300 mg, 1.29 mmol), 4-(cyclopropylmethoxy)benzyl methanesulfonate (429 mg, 1.68 mmol) and Cs₂CO₃ (841 mg, 2.58 mmol) in DMF (6 mL, anhydrous) was stirred at 60°C for 12 hours. The mixture was filtered and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (column height: 250 mm, diameter: 100 mm, 100-
10 200 mesh silica gel, Petroleum ether/Ethyl acetate=10/1, 0/1) to give 7-(4-(cyclopropylmethoxy)benzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (360 mg, 68% yield).

¹H NMR (CDCl₃, 400 MHz): δ 7.92 (s, 1H), 7.14 (d, *J* = 8.8 Hz, 2H), 6.85 (d, *J* = 9.2 Hz, 2H), 6.73 (s, 1H), 5.16 (s, 2H), 4.12 (d, *J* = 10.8 Hz, 2H), 3.77 (d, *J* = 6.8 Hz, 2H), 3.61 - 3.55 (m, 2H), 3.10 - 3.04 (m, 1H), 2.20 - 2.09 (m, 5H), 1.88 (d, *J* = 13.2 Hz, 2H), 1.27 - 1.24 (m, 1H), 0.66 - 0.61 (m, 2H), 0.35 - 0.33 (m, 2H).

LC-MS: *t*_R = 2.238 min (method 3), *m/z* = 394.1 [M + H]⁺.

Example 43:



20

7-benzyl-6-ethyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:


- A mixture of 6-ethyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 404 μmol), bromomethylbenzene (83 mg, 485.26 μmol) and Cs₂CO₃ (264 mg, 809 μmol) in DMF (3.0 mL) was stirred at 60°C for 16 hours. The reaction mixture was filtered and the
25 filtrate was concentrated. The residue was purified by HPLC to give 7-benzyl-6-ethyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (56.0 mg, 41.0% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.91 (s, 1H), 7.32 - 7.26 (m, 3H), 7.22 - 7.15 (m, 2H), 6.69 (s, 1H), 5.25 (s, 2H), 4.12 (d, J = 10.8 Hz, 2H), 3.62 - 3.56 (m, 2H), 3.14 - 3.07 (m, 1H), 2.53 - 2.48 (m, 2H), 2.17 - 2.09 (m, 2H), 1.89 (d, J = 13.2 Hz, 2H), 1.22 (t, J = 7.2 Hz, 3H).

LC-MS: t_R = 2.042 min (method 3), m/z = 338.2 $[\text{M} + \text{H}]^+$.

5

Example 44:




6-ethyl-7-(4-methoxybenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

- 10 A mixture of 6-ethyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 404 μmol), 1-(bromomethyl)-4-methoxy-benzene (98 mg, 485 μmol) and Cs_2CO_3 (264 mg, 809 μmol) in DMF (5.0 mL) was stirred at 60°C for 16 hours. The reaction mixture was filtered and the filtrate was concentrated. The residue was purified by HPLC to give 6-ethyl-7-(4-methoxybenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (65 mg, 42.3% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.91 (s, 1H), 7.12 (d, J = 8.0 Hz, 2H), 6.83 (d, J = 7.6 Hz, 2H), 6.68 (s, 1H), 5.18 (s, 2H), 4.12 (d, J = 11.6 Hz, 2H), 3.77 (s, 3H), 3.61 - 3.55 (m, 2H), 3.12 - 3.07 (m, 1H), 2.56 - 2.51 (m, 2H), 2.16 - 2.08 (m, 2H), 1.88 (d, J = 13.2 Hz, 2H), 1.22 (t, J = 7.2 Hz, 3H).

- 20 LC-MS: t_R = 2.050 min (method 3), m/z = 368.2 $[\text{M} + \text{H}]^+$.

Example 45:




3-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)benzonitrile:

A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 428.69 μmol), 3-(bromomethyl)benzonitrile (126 mg, 643.03 μmol) and Cs_2CO_3 (279 mg, 858 μmol) in DMF (3 mL) was stirred at 70°C for 6 hours. The mixture was filtered and the filtrate was concentrated under vacuum. The residue was purified by preparative LC-MS to afford 3-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)benzonitrile (35 mg, 23% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.95 (s, 1H), 7.59 (d, J = 6.8 Hz, 1H), 7.51 - 7.45 (m, 3H), 6.80 (s, 1H), 5.25 (s, 2H), 4.14 (d, J = 10.8 Hz, 2H), 3.63 - 3.57 (m, 2H), 3.14 - 3.08 (m, 1H), 2.19 - 2.10 (m, 5H), 1.91 (d, J = 13.2 Hz, 2H).

LC-MS: t_R = 2.164 min (method 3), m/z = 349.1 $[\text{M} + \text{H}]^+$.

Example 46:



4-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)benzonitrile:


A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 428.69 μmol), 4-(bromomethyl)benzonitrile (126 mg, 643.04 μmol) and Cs_2CO_3 (279 mg, 858 μmol) in DMF (3 mL) was stirred at 60°C for 2 hours. The mixture was filtered and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (petroleum ether/ethyl acetate=10/1, 1/5) to afford 4-((6-methyl-8-oxo-3-

(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-7(8*H*)-yl)methyl)benzonitrile (65 mg, 43% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.94 (s, 1H), 7.64 (d, *J* = 8.0 Hz, 2H), 7.32 (d, *J* = 8.4 Hz, 2H), 6.79 (s, 1H), 5.28 (s, 2H), 4.14 (d, *J* = 11.2 Hz, 2H), 3.62 - 3.56 (m, 2H), 3.13 - 3.07 (m, 1H),
 5 2.19 - 2.09 (m, 5H), 1.89 (d, *J* = 13.2 Hz, 2H).

LC-MS: *t_R* = 2.160 min (method 3), *m/z* = 349.2 [M + H]⁺.

Example 47:



10 ***N*-(4-((6-methyl-8-oxo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-7(8*H*)-yl)methyl)phenyl)acetamide:**

Step 1: A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 858 μmol), 1-(bromomethyl)-4-nitrobenzene (278 mg, 1.29 mmol) and Cs₂CO₃ (559 mg, 1.71 mmol) in DMF (4 mL) was stirred at 60°C for 2 hours. The mixture was filtered and
 15 the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (petroleum ether/ethyl acetate=10/1, 1/5) to afford 6-methyl-7-(4-nitrobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (270 mg, 85% yield). LC-MS: *t_R* = 0.612 min (method 2), *m/z* = 368.8 [M + H]⁺.

Step 2: A mixture of 6-methyl-7-(4-nitrobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-
 20 *a*]pyrazin-8(7*H*)-one (200 mg, 542.90 μmol), Fe (152 mg, 2.71 mmol), NH₄Cl (88 mg, 1.63 mmol) and MeOH (10 mL) in H₂O (10 mL) was stirred at 70°C for 4 hours. The mixture was filtered and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (dichloromethane /methanol=1/0, 15/1) to afford 7-(4-aminobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (170 mg, 93% yield).
 25 LC-MS: *t_R* = 0.287 min (method 2), *m/z* = 338.9 [M + H]⁺.


Step 3: A mixture of 7-(4-aminobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (120 mg, 354.61 μmol), acetyl acetate (60 mg, 588.65 μmol) and

triethylamine (144 mg, 1.42 mmol) in dioxane (10 mL) was stirred at 90°C for 6 hours. The solution was quenched with water (2 mL) and stirred at 60°C for 2 hours. Then it was concentrated under vacuum. The residue was purified by silica gel chromatography (dichloromethane / methanol=1/0, 15/1) to afford *N*-(4-((6-methyl-8-oxo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-7(8*H*)-yl)methyl)phenyl)acetamide (120 mg, 86.52% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.92 (s, 1H), 7.44 (d, *J* = 8.4 Hz, 2H), 7.17 (d, *J* = 8.4 Hz, 2H), 6.74 (s, 1H), 5.19 (s, 2H), 4.13 (d, *J* = 10.4 Hz, 2H), 3.59 (t, *J* = 10.0 Hz, 2H), 3.12 - 3.06 (m, 1H), 2.19 - 2.09 (m, 8H), 1.88 (d, *J* = 13.2 Hz, 2H).

LC-MS: *t*_R = 1.593 min (method 3), *m/z* = 381.1 [M + H]⁺.

Example 48:




7-(4-chloro-3-methoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

To a suspension of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 429 μmol) in dry DMF (2 mL) was added Cs₂CO₃ (279 mg, 858 μmol) and 4-(bromomethyl)-1-chloro-2-methoxybenzene (151 mg, 643 μmol). The mixture was purged with N₂ for 2 min and heated at 60°C for 16 hours. The mixture was concentrated. DCM (30 mL) was added to the residue. It was filtered and the filter cake was washed with DCM (20 mL). The filtrate was concentrated and purified by flash chromatography on silica gel (10%~100% ethyl acetate in petroleum ether) to give 7-(4-chloro-3-methoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (99.01 mg, 60% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.93 (s, 1H), 7.30 (d, *J* = 8.4 Hz, 1H), 6.85 (d, *J* = 1.6 Hz, 1H), 6.75 (s, 1H), 6.72 - 6.69 (m, 1H), 5.17 (s, 2H), 4.14 - 4.11 (m, 2H), 3.86 (s, 3H), 3.61 - 3.55 (m, 2H), 3.10 - 3.06 (m, 1H), 2.20 (s, 3H), 2.18 - 2.11 (m, 2H), 1.89 - 1.86 (m, 2H).

LC-MS: *t*_R = 2.466 min (method 3), *m/z* = 388.1 [M + H]⁺.

Example 49:

7-(2-ethylbenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one :


- 5 Step 1: A solution of (2-ethylphenyl)methanol (500 mg, 3.67 mmol) and triethylamine (742 mg, 7.34 mmol) in DCM (5 mL) was added MsCl (1.0 g, 8.73 mmol) at 0°C and stirred at 0°C for 0.5 hour. Then it was stirred at 20°C for 1 hour. The mixture was quenched with water (0.5 mL), and diluted with DCM (10 mL). The mixture was added NaHCO₃ (aq.) until pH=8. The organic layer was washed with water (3 × 5 mL), dried with anhydrous Na₂SO₄, filtered and concentrated under vacuum to give 2-ethylbenzyl methanesulfonate (600 mg), which
- 10 was used into the next step without further purification.

- Step 2: A mixture of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (150 mg, 643 μmol), 2-ethylbenzyl methanesulfonate (276 mg) and Cs₂CO₃ (419 mg, 1.29 mmol) in DMF (5 mL) was stirred at 60°C for 12 hours. The mixture was filtered and the
- 15 filtrate was concentrated under vacuum. The residue was purified by preparative LC-MS to afford 7-(2-ethylbenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (90 mg, 40% yield).

- ¹H NMR (CDCl₃ 400 MHz): δ 7.93 (s, 1H), 7.25 - 7.21 (m, 2H), 7.14 - 7.10 (m, 1H), 6.81 - 6.77 (m, 2H), 5.25 (s, 2H), 4.14 (d, *J* = 11.2 Hz, 2H), 3.60 (t, *J* = 12.0 Hz, 2H), 3.15 - 3.09 (m, 1H), 2.78 - 2.72 (m, 2H), 2.21 - 2.13 (m, 5H), 1.91 (d, *J* = 13.6 Hz, 2H), 1.31 (t, *J* = 7.6 Hz, 3H).
- 20

LC-MS: *t*_R = 2.533 min (method 3), *m/z* = 352.2 [M + H]⁺.

Example 50:




7-(benzo[d][1,3]dioxol-5-ylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

To a suspension of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 429 μ mol) in dry DMF (2 mL) was added Cs_2CO_3 (279 mg, 858 μ mol) and 5-(bromomethyl)benzo[d][1,3]dioxole (138 mg, 643 μ mol). The mixture was bubbled with N_2 for 2 min and heated at 60°C for 16 hours. The mixture was concentrated. DCM (30 mL) was added to the residue. It was filtered and the filter cake was washed with DCM (20 mL). The filtrate was concentrated and purified by flash chromatography on silica gel (10%~100% ethyl acetate in petroleum ether) to give 7-(benzo[d][1,3]dioxol-5-ylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (81.74 mg, 52% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.92 (s, 1H), 6.76 - 6.69 (m, 4H), 5.95 - 5.93 (m, 2H), 5.13 (s, 2H), 4.14 - 4.11 (m, 2H), 3.61 - 3.55 (m, 2H), 3.10 - 3.05 (m, 1H), 2.21 (s, 3H), 2.20 - 2.10 (m, 2H), 1.90 - 1.86 (m, 2H).

LC-MS: t_R = 2.245 min (method 3), m/z = 368.2 $[\text{M} + \text{H}]^+$.

Example 51:



7-(3-chloro-4-methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 427 μ mol), 4-(bromomethyl)-2-chloro-1-methoxybenzene (151 mg, 643 μ mol) and Cs_2CO_3 (279 mg, 858 μ mol) in DMF (3 mL) was stirred at 60°C for 2 hours. The mixture was filtered and the filtrate was concentrated under vacuum. The residue was purified by silica

gel chromatography (detroleum ether/ethyl acetate=10/1, 0/1) to afford 7-(3-chloro-4-methoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (110 mg, 65% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.93 (s, 1H), 7.24 (d, *J* = 0.8 Hz, 1H), 7.12 - 7.10 (m, 1H), 6.88 (d, *J* = 8.4 Hz, 1H), 6.75 (s, 1H), 5.15 (s, 2H), 4.13 (d, *J* = 10.4 Hz, 2H), 3.88 (s, 3H), 3.62 - 3.56 (m, 2H), 3.12 - 3.06 (m, 1H), 2.20 - 2.09 (m, 5H), 1.89 (d, *J* = 13.2 Hz, 2H).

LC-MS: *t*_R = 2.414 min (method 3), *m/z* = 388.1 [M + H]⁺.

Example 52:




7-(4-aminobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one :

A solution of *N*-(4-((6-methyl-8-oxo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-7(8*H*)-yl)methyl)phenyl)acetamide (100 mg, 263 μmol), NaOH (63 mg, 1.58 mmol) and MeOH (1 mL) in H₂O (1 mL) was stirred at 90°C for 12 hours. The solution was added KHSO₄ (aq.) until pH=7 and concentrated under vacuum. The residue was purified by preparative TLC (DCM: MeOH = 10:1). to afford 7-(4-aminobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (50 mg, 56% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.91 (s, 1H), 7.04 (d, *J* = 8.4 Hz, 2H), 6.71 (s, 1H), 6.63 (d, *J* = 8.4 Hz, 2H), 5.12 (s, 2H), 4.13 (d, *J* = 12.0 Hz, 2H), 3.67 (brs, 2H), 3.61 - 3.55 (m, 4H), 3.10 - 3.05 (m, 1H), 2.21 (s, 3H), 2.17 - 2.08 (m, 2H), 1.88 (d, *J* = 13.2 Hz, 2H).

LC-MS: *t*_R = 1.273 min (method 3), *m/z* = 339.1 [M + H]⁺.

Example 53:




7-(4-hydroxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

A solution of 7-(4-methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (2.0 g, 5.66 mmol) in DCM (32 mL) was added BBr₃ (4.3 g, 16.98 mmol) at 0°C and stirred at 20°C for 3 hours. The solution was quenched with H₂O (5 mL) at 0°C. The mixture was stirred at 0°C for 1 hour and then to it was added NaHCO₃ (saturated aqueous) until pH=6. The mixture was concentrated under vacuum. The residue was diluted with DCM (20 mL) and MeOH (2 mL). Then it was filtered and the filtrate was concentrated under vacuum. The residue was added into KOH (60 mL, 2 M, aq.) at 20°C and stirred at 50°C for 1 hour. The solution was added KHSO₄ (saturated aqueous) until pH=6, the mixture was concentrated under vacuum. The residue was purified by silica gel chromatography (dichloromethane/ methanol=1/0, 15/1) to afford 7-(4-hydroxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (460 mg, 24% yield).

¹H NMR (DMSO Varian_H_400 MHz): δ 9.39 (s, 1H), 7.70 (s, 1H), 7.46 (s, 1H), 6.99 (d, *J* = 8.4 Hz, 2H), 6.70 (d, *J* = 8.4 Hz, 2H), 5.06 (s, 2H), 3.96 - 3.93 (m, 2H), 3.52 - 3.45 (m, 2H), 3.29 - 3.16 (m, 1H), 2.15 (s, 3H), 1.83 - 1.77 (m, 4H).

LC-MS: *t_R* = 1.62 min (method 8), *m/z* = 340.1 [M + H]⁺.

Example 54:



6-ethyl-7-(3-fluorobenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


Step 1: A mixture of methyl 1H-imidazole-5-carboxylate (1.6 g, 13 mmol), 1-bromobutan-2-one (2.0 g, 13 mmol) and K₂CO₃ (3.5 g, 25 mmol) in acetone (20 mL) was stirred at 40°C for

fluorobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (150 mg, 34% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.94 (s, 1H), 7.32 - 7.27 (m, 1H), 6.98 - 6.94 (m, 2H), 6.86 (d, *J* = 9.6 Hz, 1H), 6.72 (s, 1H), 5.25 (s, 2H), 4.14 (d, *J* = 11.2 Hz, 2H), 3.64 - 3.58 (m, 2H), 3.15 - 3.09 (m, 1H), 2.53 - 2.48 (m, 2H), 2.18 - 2.13 (m, 2H), 1.91 (d, *J* = 13.2 Hz, 2H), 1.25 (t, *J* = 7.2 Hz, 2H).

LC-MS: *t*_R = 2.475 min (method 3), *m/z* = 356.1 [M + H]⁺.

Example 55:



7-(4-methoxybenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one, stereoisomers 1,2,3 and 4:

Step 1: To a cooled (0°C) solution of 2-methyltetrahydrofuran-3-carboxylic acid (110 mg, 845 μmol) in dry DCM (2 mL) was added oxalyl dichloride (107 mg, 845 μmol) dropwise. Then one drop of DMF was added and the mixture was stirred at 26°C for 1 hour. The solution of 2-methyltetrahydrofuran-3-carbonyl chloride (126 mg) in DCM (2 mL) was directly used for the next step.

Step 2: To a cooled (0°C) solution of 2-methyltetrahydrofuran-3-carbonyl chloride (160 mg, 844 μmol, HCl) in dry DCM (5 mL) was added triethylamine (256 mg, 2.53 mmol) and (3-methoxy-5-methylpyrazin-2-yl)methanamine (125 mg, 843.70 μmol) in DCM (2 mL) dropwise. The mixture was stirred at 26°C for 1 hour. LCMS showed the reaction was completed. Water (5 mL) was added to the mixture. The mixture was extracted with DCM (30 mL × 2). The combined organic layer was washed with H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated. The crude product was purified by flash chromatography on silica gel (10~50% ethyl acetate in petroleum ether) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methyltetrahydrofuran-3-carboxamide (100 mg, 45% yield) as a light yellow oil.

Step 3: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methyltetrahydrofuran-3-carboxamide (150 mg, 565 μmol) in dry dioxane (5 mL) was added POCl_3 (173 mg, 1.13 mmol). The mixture was heated at 80°C for 2 hours. The mixture was cooled to 26°C and the brown solution of 8-methoxy-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazine (140 mg) in dioxane (5 mL) was directly used for the next step.

Step 4: To a solution of 8-methoxy-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazine (140 mg, 566 μmol) in dioxane (5 mL) was added 2 N HCl (2 M, 2 mL). The mixture was heated at 80°C for 1 hour. The mixture was cooled to 25°C , adjusted to pH=7 by saturated aqueous NaHCO_3 and extracted with DCM (20 mL $\times$ 2). The combined organic phases were washed with H_2O (20 mL), brine (20 mL), dried over Na_2SO_4 , filter and concentrated to give 6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (131 mg).

Step 5: To a solution of 6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (131 mg, 562 μmol) in dry DMF (5 mL) was added 1-(bromomethyl)-4-methoxybenzene (169 mg, 842 μmol) and Cs_2CO_3 (366 mg, 1.12 mmol). The mixture was heated at 60°C for 16 hours. The mixture was concentrated and water (10 mL) was added. The mixture was extracted with DCM (30 mL $\times$ 2). The combined organic layer was washed with H_2O (30 mL $\times$ 2), brine (30 mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by flash chromatography on silica gel (10%~100% ethyl acetate in petroleum ether) to give 7-(4-methoxybenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (90 mg, 45% yield).

7-(4-methoxybenzyl)-6-methyl-3-(2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 283 μmol) was purified by SFC.

Stereoisomer 1: (9.4 mg, 9% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.92 (s, 1H), 7.15 (d, J = 8.8 Hz, 2H), 6.84 (d, J = 8.4 Hz, 2H), 6.74 (s, 1H), 5.16 (s, 2H), 4.30-4.26 (m, 1H), 4.12-4.04 (m, 2H), 3.78 (s, 3H), 3.09-3.03 (m, 1H), 2.46-2.34 (m, 2H), 2.20 (s, 3H), 1.32 (d, J = 6.0 Hz, 3H).

LC-MS: t_R = 2.074 min (method 13), m/z = 354.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 5.177 min, ee% > 99%. $[\alpha]_D^{20}$ -26 (c = 0.10, DCM).

Stereoisomer 2: (7.3 mg, 7% yield) .

^1H NMR (CDCl_3 400 MHz): δ 7.92 (s, 1H), 7.15 (d, J = 8.8 Hz, 2H), 6.84 (d, J = 8.4 Hz, 2H), 6.73 (s, 1H), 5.16 (s, 2H), 4.30-4.26 (m, 1H), 4.10-4.02 (m, 2H), 3.78 (s, 3H), 3.09-3.03 (m, 1H), 2.44-2.35 (m, 2H), 2.20 (s, 3H), 1.32 (d, J = 6.0 Hz, 3H).

LC-MS: t_R = 2.072 min (method 13), m/z = 354.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 5.458 min, ee% = 99.7%. $[\alpha]_D^{20}$ +24 (c = 0.10, DCM).

Stereoisomer 3: (39.7 mg, 40% yield) .

^1H NMR (CDCl_3 400 MHz): δ 7.92 (s, 1H), 7.15 (d, J = 8.4 Hz, 2H), 6.84 (d, J = 8.4 Hz, 2H), 6.78 (s, 1H), 5.22-5.10 (m, 2H), 4.31-4.26 (m, 2H), 3.88-3.82 (m, 1H), 3.77 (s, 3H), 3.70-3.60 (m, 1H), 2.71-2.67 (m, 1H), 2.43-2.40 (m, 1H), 2.19 (s, 3H), 0.87 (d, J = 4.4 Hz, 3H).


LC-MS: t_R = 1.983 min (method 13), m/z = 354.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 5.932 min, ee% = 98.8%. $[\alpha]_D^{20}$ +48 (c = 0.10, DCM).

Stereoisomer 4: (26.4 mg, 26% yield) .

^1H NMR (CDCl_3 400 MHz): δ 7.92 (s, 1H), 7.15 (d, J = 8.8 Hz, 2H), 6.84 (d, J = 8.4 Hz, 2H), 6.78 (s, 1H), 5.22-5.09 (m, 2H), 4.32-4.26 (m, 2H), 3.87-3.85 (m, 1H), 3.77 (s, 3H), 3.68-3.64 (m, 1H), 2.70-2.65 (m, 1H), 2.42-2.37 (m, 1H), 2.19 (s, 3H), 0.87 (d, J = 6.4 Hz, 3H).

LC-MS: t_R = 1.983 min (method 13), m/z = 354.1 $[\text{M} + \text{H}]^+$. SFC: t_R = 6.570 min, ee% = 99.6%. $[\alpha]_D^{20}$ -64 (c = 0.10, DCM).

Example 56:



7-(4-methoxybenzyl)-6-methyl-3-propylimidazo[1,5-a]pyrazin-8(7H)-one:


To a solution of 3-bromo-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (500 mg, 1.44 mmol, 1 eq) in THF (10 mL) was added Ni(dppp)Cl_2 (327.83 mg, 604.80 μmol , 0.42 eq) at 0°C over 10 min, then at -78°C for 10 min. propylmagnesium bromide (1 M, 3 mL, 2.1 eq) was added dropwise at 0°C . The resulting mixture was stirred at 0°C for 1.5 hour. The reaction was quenched by saturated NH_4Cl aqueous solution (5 mL). The residue

was purified by silica column chromatography (Gradient: 0~50, EtOAc in PE with 1% triethylamine) to give 220 mg of crude product. The crude product was purified by preparative TLC (PE:EtOAc=1:1 with 1% triethylamine). The residue was washed with hexane (3 mL) and filtered and the filter cake was dried under vacuum to give 7-(4-methoxybenzyl)-6-methyl-3-propylimidazo[1,5-a]pyrazin-8(7H)-one (82 mg, 18% yield).

¹HNMR (CDCl₃, 400 MHz): δ 7.90 (s, 1H), 7.16 (d, *J* = 8.8, 2H), 6.85 (d, *J* = 8.8, 2H), 6.68 (s, 1H), 5.17 (s, 2H), 3.78 (s, 3H), 2.82 (t, *J* = 7.2, 2H), 2.19 (s, 3H), 1.90-1.81 (m, 2H), 1.03 (t, *J* = 7.2, 3H).

LC-MS: tR = 1.927 min (method 13), m/z = 312.1 [M + H]⁺.

Example 57:




7-((6-methoxypyridin-3-yl)methyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

Into a vial was added 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (200mg, 0.86 mmol), 5-(chloromethyl)-2-methoxypyridine (162 mg, 1.03 mmol), cesium carbonate (559 mg, 1.72 mmol) and sodium iodide (154 mg, 1.03 mmol) in DMF (9.44 g, 10 ml, 129 mmol). The reaction was stirred overnight at 70°C. To the reaction was added ethylacetate, and it was filtered and concentrated. The reaction was purified by chromatography on silicagel to obtain 7-((6-methoxypyridin-3-yl)methyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (127 mg, 0.358 mmol) in 42% yield.

¹H NMR (600 MHz, CDCl₃) δ 8.06 (dd, *J* = 2.5, 0.8 Hz, 1H), 7.91 (d, *J* = 0.6 Hz, 1H), 7.55 (dd, *J* = 8.6, 2.5 Hz, 1H), 6.74 (t, *J* = 1.1 Hz, 1H), 6.71 (dd, *J* = 8.6, 0.7 Hz, 1H), 5.15 (s, 2H), 4.12 (m, 2H), 3.91 (s, 3H), 3.58 (td, *J* = 11.7, 2.2 Hz, 2H), 3.07 (tt, *J* = 11.4, 3.9 Hz, 1H), 2.25 (d, *J* = 1.2 Hz, 3H), 2.12 (dtd, *J* = 13.7, 11.6, 4.3 Hz, 2H), 1.87 (ddd, *J* = 13.5, 4.2, 2.1 Hz, 2H).

LC-MS: t_R = 0.38 min (method 6), m/z = 355.2 $[M + H]^+$.

Example 58:




5 6,7-dimethyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

To a solution of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 428.69 μ mol, 1 eq) in DMSO (2 mL) was added Cs_2CO_3 (139.68 mg, 428.69 μ mol, 1 eq) and methyl iodide (121.70 mg, 858 μ mol, 53.38 μ L, 2 eq). The mixture was stirred at 50 °C for 12 hour. The reaction mixture was diluted with H_2O (25 mL) and extracted with DCM (50 mL $\times$ 2). The combined organic layers were washed with brine (15 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (ethyl acetate). 6,7-dimethyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (73 mg, 68% yield, 98% purity) was obtained.

^1H NMR (CDCl_3 400 MHz): δ 7.87 (s, 1H), 6.76 (s, 1H), 4.13 (d, J = 11.2 Hz, 2H), 3.59 (t, J = 11.2 Hz, 2H), 3.46 (s, 3H), 3.11-3.05 (m, 1H), 2.28 (s, 3H), 2.16-2.08 (m, 2H), 1.88 (d, J = 14.0 Hz, 2H).

LC-MS: t_R = 1.300 min (method 13), m/z = 248.1 $[M + H]^+$.

Example 59:




7-ethyl-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

To a solution of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 429 μ mol, 1 *eq*) in anhydrous DMF (2 mL) was added K_2CO_3 (119 mg, 858 μ mol, 2 *eq*) and iodoethane (134 mg, 858 μ mol, 69 μ L, 2 *eq*). The mixture was stirred at 50°C for 12
 5 hour. The reaction mixture was concentrated under reduced pressure. The residue was diluted with H_2O (20 mL) and extracted with EA (40 mL $\times$ 2). The combined organic layers were washed with brine (15 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford a residue. 7-ethyl-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (40 mg, 34% yield, 96% purity) was obtained.

10 ^1H NMR ($CDCl_3$ 400 MHz): δ 7.85 (s, 1H), 6.73 (s, 1H), 4.12 (d, J = 10.8 Hz, 2H), 4.03-3.99 (m, 2H), 3.58 (t, J = 10.8 Hz, 2H), 3.10-3.04 (m, 1H), 2.30 (s, 3H), 2.16-2.07 (m, 2H), 1.87 (d, J = 13.2 Hz, 2H), 1.29 (t, J = 6.8 Hz, 3H).

LC-MS: t_R = 1.490 min (method 11), m/z = 262.1 $[M + H]^+$.

15 Example 60:



6-methyl-7-propyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:


To a solution of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100 mg, 429 μ mol, 1 *eq*) in anhydrous DMF (2 mL) was added K_2CO_3 (119 mg, 858 μ mol, 2 *eq*) and 1-bromopropane (105 mg, 858 μ mol, 78.11 μ L, 2 *eq*). The mixture was stirred at 50 °C
 20 for 12 hours. The reaction mixture was concentrated under reduced pressure to remove DMF. The residue was diluted with H_2O (20 mL) and extracted with EA (40 mL $\times$ 2). The combined organic layers were washed with brine (15 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (ethyl
 25 acetate). 6-methyl-7-propyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (45 mg, 37% yield, 98% purity) was obtained.

^1H NMR (CDCl_3 , 400 MHz): δ 7.85 (s, 1H), 6.72 (s, 1H), 4.12 (d, J = 10.4 Hz, 2H), 3.88 (t, J = 8.0 Hz, 2H), 3.62-3.56 (m, 2H), 3.10-3.07 (m, 1H), 2.28 (s, 3H), 2.14-2.09 (m, 2H), 1.87 (d, J = 13.2 Hz, 2H), 1.73-1.67 (m, 2H), 1.00 (t, J = 7.2 Hz, 3H).

LC-MS: t_R = 1.666 min (method 11), m/z = 276.1 $[\text{M} + \text{H}]^+$.

5

Example 61:




7-isopropyl-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

To a solution of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (350 mg, 1.50 mmol, 1 eq) in anhydrous DMF (4 mL) was added Cs_2CO_3 (978 mg, 3 mmol, 2 eq) and 2-iodopropane (510 mg, 3 mmol, 300 μL , 2 eq). The mixture was stirred at 50 $^\circ\text{C}$ for 12 hours. The reaction mixture was concentrated under reduced pressure to remove DMF. The residue was diluted with H_2O (15 mL) and extracted with EtOAc (40 mL $\times$ 2). The combined organic layers were washed with brine (10 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (ethyl acetate). 7-isopropyl-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (68 mg, 16% yield, 96% purity) was obtained.

^1H NMR (CDCl_3 , 400 MHz): δ 7.81 (s, 1H), 6.67 (s, 1H), 4.40 (m, 1H), 4.12 (d, J = 11.6 Hz, 2H), 3.58 (t, J = 11.3 Hz, 2H), 3.09-3.03 (m, 1H), 2.27 (s, 3H), 2.14-2.06 (m, 2H), 1.86 (d, J = 13.2 Hz, 2H), 1.62-1.61 (m, 6H).

LC-MS: t_R = 1.576 min (method 13), m/z = 276.1 $[\text{M} + \text{H}]^+$.


20

Example 62:**7-isopentyl-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**

- 5 To a solution of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 428.69 μ mol, 1 eq) in anhydrous DMF (2 mL) was added K_2CO_3 (118.50 mg, 858 μ mol, 2 eq) and 1-bromo-3-methylbutane (129.50 mg, 858 μ mol, 108 μ L, 2 eq). The mixture was stirred at 50 $^{\circ}C$ for 12 hours. The reaction mixture was concentrated under reduced pressure to remove DMF. The residue was diluted with H_2O (20 mL) and extracted with EtOAc(40 mL
- 10 $\times$ 2). The combined organic layers were washed with brine(15 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (ethyl acetate). 7-isopentyl-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (65 mg, 50% yield) was obtained.

- ^1H NMR ($CDCl_3$ 400 MHz): δ 7.84 (s, 1H), 6.73 (s, 1H), 4.12 (d, J = 10.4 Hz, 2H), 3.93 (t, J = 8.0 Hz, 2H), 3.61-3.55 (m, 2H), 3.08-3.04 (m, 1H), 2.28 (s, 3H), 2.10-2.09 (m, 2H), 1.87 (d, J = 14.0 Hz, 2H), 1.74-1.69 (m, 1H), 1.56-1.52 (m, 2H), 1.00-0.98 (m, 6H).
- 15

LC-MS: t_R = 1.968 min (method 13), m/z = 304.2 $[M + H]^+$.


Example 63:

7-(cyclopentylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

To a mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (150 mg, 0.64 mmol) and (bromomethyl)cyclopentane (157 mg, 0.96 mmol) in DMSO (2 mL) was added Cs₂CO₃ (419 mg, 1.29 mmol). The mixture was stirred at 60 °C for 12 hours. The mixture was diluted with water (10 mL) and extracted with DCM (5 mL × 3). The combine organic layer was washed with water (5 mL × 2) and dried over Na₂SO₄. The organic layers was evaporated under vacuum. The residue was purified by preparative TLC (ethyl acetate) to give 7-(cyclopentylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (95 mg, 46% yield). ¹HNMR (CDCl₃, 400 MHz): δ 7.85 (s, 1H), 6.72 (s, 1H), 4.14-4.11 (m, 2H), 3.91 (d, *J* = 7.6 Hz, 2H), 3.61-3.56 (m, 2H), 3.12-3.05 (m, 1H), 2.29 (s, 3H), 2.27-2.25 (m, 1H), 2.14-2.10 (m, 2H), 1.90-1.86 (m, 2H), 1.71-1.54 (m, 6H), 1.34-1.32 (m, 2H).

LC-MS: tR = 1.98 min (method 13), m/z = 316.2 [M + H]⁺.

Example 64:




2-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)benzonitrile:

A mixture of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (100 mg, 429 μmol), 2-(bromomethyl)benzonitrile (126 mg, 643 μmol) and Cs₂CO₃ (279 mg, 857 μmol) in DMF (3.0 mL) was stirred at 70 °C for 6 hours. The mixture was filtered and the filtrate was concentrated under vacuum. The residue was purified by silica gel chromatography (Petroleum ether/Ethyl acetate=10/1, 0/1) to afford 2-((6-methyl-8-oxo-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8H)-yl)methyl)benzonitrile (92 mg, 59% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.96 (s, 1H), 7.72 (d, $J = 7.6$ Hz, 1H), 7.55 (t, $J = 7.2$ Hz, 1H), 7.40 (t, $J = 7.2$ Hz, 1H), 7.19 (d, $J = 8.0$ Hz, 1H), 6.80 (s, 1H), 5.45 (s, 2H), 4.14 (d, $J = 12.0$ Hz, 2H), 3.63 - 3.57 (m, 2H), 3.14 - 3.08 (m, 1H), 2.20 - 2.09 (m, 5H), 1.90 (d, $J = 13.2$ Hz, 2H).

5 LC-MS: $t_R = 2.202$ min (method 3), $m/z = 349.1$ $[\text{M} + \text{H}]^+$.

Example 65:



7-(cycloheptylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

10

Step 1: To a solution of cycloheptanecarboxylic acid (500 mg, 3.52 mmol) in THF (40 mL) was added LiAlH_4 (401 mg, 10.6 mmol) in portions at 0°C . The mixture was stirred at 65°C for 3 hours. The reaction was quenched with H_2O (0.4 mL) and 10% NaOH (0.4 mL, aq). To the mixture was added Na_2SO_4 . The mixture was filtered. The filtrate was concentrated. The residue was purified by flash silica gel chromatography to give cycloheptylmethanol (383 mg, 85% yield).

15

^1H NMR (CDCl_3 400 MHz): δ 3.43 (d, $J = 6.4$ Hz, 2H), 1.80 - 1.62 (m, 5H), 1.56 - 1.40 (m, 4H), 1.31 (br, s, 1H), 1.84 (s, 3H), 1.10 - 1.22 (m, 2H).

Step 2: To a solution of cycloheptylmethanol (313 mg, 2.44 mmol) and triethylamine (494 mg, 4.88 mmol) in DCM (5 mL) was added MsCl (490 mg, 4.28 mmol) at 0°C and it was stirred at 20°C for 40 min. The solution was washed with NaHCO_3 (saturated aqueous 5 mL $\times 4$), water (5 mL $\times 2$), brine (3 mL) and then was dried over Na_2SO_4 , filtered and concentrated to give cycloheptylmethyl methanesulfonate (381 mg) which was used in the next step directly without further purification.

20

Step 3: To a solution of 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (150 mg, 643 μmol) and cycloheptylmethyl methanesulfonate (159 mg, 772 μmol) in DMF (3 mL) was added Cs_2CO_3 (419 mg, 1.29 mmol). The mixture was stirred at 60°C for 6

25


hours. The mixture was diluted with DCM (20 mL) and washed with water (5 mL × 2), brine (10 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash silica gel chromatography and then was purified by preparative LC-MS to give 7-

(cycloheptylmethyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (25.15 mg, 11% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.84 (s, 1H), 6.71 (s, 1H), 4.12 (d, *J* = 5.2 Hz, 2H), 3.77 (d, *J* = 3.6 Hz, 2H), 3.65 - 3.52 (m, 2H), 3.15 - 3.02 (m, 1H), 2.26 (s, 3H), 2.17 - 2.07 (m, 2H), 2.07 - 2.98 (m, 1H), 1.87 (d, *J* = 6.6 Hz, 2H), 1.63 - 1.75 (m, 4H), 1.55 - 1.62 (m, 2H), 1.54 - 1.45 (m, 2H), 1.44 - 1.33 (m, 2H), 1.30 - 1.18 (m, 2H).

LC-MS: *t*_R = 2.227 min (method 13), *m/z* = 344.2 [M + H]⁺.

Example 66:




6-methyl-7-((4-methylcyclohexyl)methyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one, *cis* and *trans*:

Step 1: To a solution of (4-methylcyclohexyl)methanol (400 mg, 3.12 mmol, 1 eq) in anhydrous DCM (20 mL) was added triethylamine (631 mg, 6.24 mmol, 865 μL, 2 eq). The mixture was dropwise added methanesulfonyl chloride (465 mg, 4.06 mmol, 314 μL, 1.30 eq) at 0 °C and it was stirred at 0 °C for 1 hour. The reaction mixture was washed with water (5mL × 3), brine (3mL), dried and concentrated. (4-methylcyclohexyl)methyl methanesulfonate (700 mg) was obtained.

Step 2: To a solution of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (260 mg, 1.11 mmol, 1 eq) and (4-methylcyclohexyl)methyl methanesulfonate (572 mg, 2.78 mmol, 2.50 eq) in anhydrous DMF (6 mL) was added Cs₂CO₃ (723 mg, 2.22 mmol, 2 eq). The mixture was stirred at 60 °C for 12 hours. The reaction mixture was concentrated under reduced pressure to remove DMF. The residue was diluted with H₂O (3 mL) and extracted with DCM (10 mL × 2). The combined organic layers were dried over Na₂SO₄

, filtered and concentrated. The residue was purified by preparative TLC (SiO₂, EA). 6-methyl-7-((4-methylcyclohexyl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (175 mg, 45% yield) was obtained.




5 ***trans*-6-methyl-7-((4-methylcyclohexyl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**

6-methyl-7-((4-methylcyclohexyl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (160 mg, 466 μ mol, 1 eq) was purified by SFC to give 6-methyl-7-(((*trans*)-4-methylcyclohexyl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (60 mg, 37% yield, 98% purity).

¹H NMR (CDCl₃ 400 MHz): δ 7.84 (s, 1H), 6.71 (s, 1H), 4.12 (d, *J* = 10.8 Hz, 2H), 3.79-3.77 (m, 2H), 3.61-3.55 (m, 2H), 3.09-3.05 (m, 1H), 2.26 (s, 3H), 2.14-2.07 (m, 2H), 1.88 (d, *J* = 10.8 Hz, 2H), 1.69-1.67 (m, 5H), 1.34-1.32 (m, 2H), 1.11-1.09 (m, 2H), 0.91-0.86 (m, 4H).

LC-MS: *t*_R = 2.256 min (method 13), *m/z* = 344.2 [M + H]⁺.




15 ***cis*-6-methyl-7-((4-methylcyclohexyl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**

6-methyl-7-(((*cis*)-4-methylcyclohexyl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (90 mg, 54% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.85 (s, 1H), 6.72 (s, 1H), 4.12 (d, *J* = 10.8 Hz, 2H), 3.89-3.87 (m, 2H), 3.61-3.55 (m, 2H), 3.09-3.05 (m, 1H), 2.28 (s, 3H), 2.14-2.11 (m, 2H), 1.87 (d, *J* = 14.4 Hz, 2H), 1.50-1.40 (m, 8H), 0.97-0.95 (m, 3H).

LC-MS: $t_R = 2.240$ min (method 13), $m/z = 344.2$ $[M + H]^+$.

Example 67:



5 7-(((*cis*)-4-methoxycyclohexyl)methyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:


To a solution of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (60 mg, 257 μ mol, 1 eq) in dry DMF (5 mL) was added ((*cis*)-4-methoxycyclohexyl)methyl methanesulfonate (74 mg, 334 μ mol, 1.30 eq) and Cs_2CO_3 (168 mg, 514 μ mol, 2 eq). The mixture was heated at 60°C for 16 hours. The mixture was concentrated. DCM (20 mL) and H_2O (10 mL) was added. The mixture was extracted with DCM (20 mL). The organic layer was washed with H_2O (20 mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by preparative TLC (DCM/MeOH=20/1) to give 7-(((*cis*)-4-methoxycyclohexyl)methyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (15 mg, 16% yield).

^1H NMR (CDCl_3 , 400 MHz): δ 7.83 (s, 1H), 6.69 (s, 1H), 4.10 (d, $J = 11.2$ Hz, 2H), 3.75 (d, $J = 7.2$ Hz, 2H), 3.56 (t, $J = 11.2$ Hz, 2H), 3.42 (m, 1H), 3.29 (s, 3H), 3.11-3.03 (m, 1H), 2.24 (s, 3H), 2.15-2.06 (m, 2H), 1.94-1.84 (m, 5H), 1.44-1.37 (m, 6H).

LC-MS: $t_R = 1.887$ min (method 13), $m/z = 360.2$ $[M + H]^+$.

20

Example 68:




7-(((*trans*)-4-methoxycyclohexyl)methyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

To a solution of 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (60 mg, 257 μ mol, 1 eq) in dry DMF (5 mL) was added ((*trans*)-4-methoxycyclohexyl)methyl methanesulfonate (74 mg, 334 μ mol, 1.30 eq) and Cs₂CO₃ (168 mg, 514 μ mol, 2 eq). The mixture was heated at 60°C for 16 hours. The mixture was concentrated. DCM (20 mL) and H₂O (10 mL) were added. The mixture was extracted with DCM (20 mL). The organic layer was washed with H₂O (20 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by preparative TLC (DCM/MeOH=20/1) to give 7-(((*trans*)-4-methoxycyclohexyl)methyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (35 mg, 38% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.84 (s, 1H), 6.71 (s, 1H), 4.11 (d, *J* = 10.4 Hz, 2H), 3.79 (d, *J* = 5.6 Hz, 2H), 3.57 (t, *J* = 10.0 Hz, 2H), 3.33 (s, 3H), 3.10-3.05 (m, 2H), 2.26 (s, 3H), 2.15-2.07 (m, 4H), 1.88-1.84 (m, 2H), 1.75-1.71 (m, 3H), 1.15-1.11 (m, 4H).

LC-MS: *t*_R = 1.831 min (method 13), *m/z* = 360.2 [M + H]⁺.

Example 69:



7-(4-methoxybenzyl)-6-methyl-3-(3-methyltetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (150 mg, 0.79 mmol, 1 eq) and 3-methyltetrahydro-2*H*-pyran-4-carboxylic acid (125 mg, 870 μ mol, 1.1 eq) in DCM (5 mL) was added HATU (451 mg, 1.19 mmol, 1.5 eq) and DIPEA (204 mg, 1.58 mmol, 276 μ L, 2 eq). The mixture was stirred at 25 °C for 18 hours. The mixture was concentrated. The crude product was purified by flash chromatography with petroleum ether:ethyl acetate = 3:1 ~ 2:1. *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-3-methyltetrahydro-2*H*-pyran-4-carboxamide (220 mg, 780 μ mol, 99% yield) was obtained.


Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-3-methyltetrahydro-2*H*-pyran-4-carboxamide (220 mg, 788 μmol , 1 eq) in dioxane (8 mL) was added POCl_3 (242 mg, 1.58 mmol, 146 μL , 2 eq). The mixture was stirred at 80 $^\circ\text{C}$ for 2 hours. The mixture was concentrated. 6-methyl-3-(3-methyltetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8-ol (223 mg, hydrochloride salt) was used in the next step without further purification.

Step 3: To a solution of 6-methyl-3-(3-methyltetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8-ol (170 mg, 0.6 mmol) in DMF (5 mL) was added Cs_2CO_3 (586 mg, 1.80 mmol, 3 eq) and 1-(chloromethyl)-4-methoxybenzene (113 mg, 719 μmol , 98 μL , 1.20 eq). The mixture was stirred at 60 $^\circ\text{C}$ for 18 hours. The reaction mixture was filtered, and the filtrate was concentrated. The crude mixture was purified by preparative LC-MS, and then by preparative TLC with ethyl acetate as eluent. 7-(4-methoxybenzyl)-6-methyl-3-(3-methyltetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (76 mg, 207 μmol , 35% yield) was obtained.

^1H NMR, a mixture of diastereoisomers (CDCl_3 400 MHz): δ 7.96 - 7.94 (m, 1H), 7.18 (d, J = 8.0 Hz, 2H), 6.86 (d, J = 8.0 Hz, 2H), 6.76 (s, 0.73H), 6.71 (s, 0.28H), 5.22 - 5.12 (m, 2H), 4.17 - 4.01 (m, 2H), 3.79 (s, 3H), 3.55 - 3.49 (m, 1H), 3.19 - 3.14 (m, 1H), 2.66 - 2.64 (m, 1H), 2.36 - 2.33 (m, 1H), 2.21 (s, 3H), 2.13 - 2.10 (m, 1H), 1.78 - 1.61 (m, 1H), 0.87 (d, J = 7.2 Hz, 1H), 0.71 (d, J = 6.8 Hz, 2H).

LC-MS: t_R = 2.370 min (method 11), m/z = 368.1 $[\text{M} + \text{H}]^+$.

Example 70:



racemic 7-(4-methoxybenzyl)-6-methyl-3-((1*R*,2*R*,4*S*)-2-methyl-7-oxabicyclo[2.2.1]heptan-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (150 mg, 791 μmol , 1 eq) in dry DCM (5 mL) was added triethylamine (240 mg, 2.37 mmol, 329 μL , 3 eq), racemic (1*R*,2*S*,4*S*)-2-methyl-7-oxabicyclo[2.2.1]heptane-2-carboxylic acid (124 mg, 791 μmol , 1 eq) and HATU (361 mg, 949 μmol , 1.20 eq). The mixture was stirred at

15°C for 16 hours. H₂O (5 mL) was added and the mixture was extracted with DCM (20 mL × 2). The combined organic layer was washed with H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash chromatography on silica gel (0%~50% ethyl acetate in petroleum ether) to give racemic (1*R*,2*S*,4*S*)-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methyl-7-oxabicyclo[2.2.1]heptane-2-carboxamide (200 mg, 87% yield).


Step 2: To a solution of racemic (1*R*,2*S*,4*S*)-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methyl-7-oxabicyclo[2.2.1]heptane-2-carboxamide (150 mg, 515 μmol, 1 eq) in dry dioxane (5 mL) was added POCl₃ (158 mg, 1.03 mmol, 96 μL, 2 eq). The mixture was heated at 80°C for 2 hours. The mixture was cooled to 15°C and poured into water (5 mL). The mixture was adjusted to pH 8 by saturated aqueous NaHCO₃ and extracted with DCM (20 mL × 2). The combined organics were washed with H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give racemic 6-methyl-3-((1*R*,2*R*,4*S*)-2-methyl-7-oxabicyclo[2.2.1]heptan-2-yl)imidazo[1,5-*a*]pyrazin-8-ol (120 mg).

Step 3: To a solution of racemic 6-methyl-3-((1*R*,2*R*,4*S*)-2-methyl-7-oxabicyclo[2.2.1]heptan-2-yl)imidazo[1,5-*a*]pyrazin-8-ol (100 mg, 386 μmol, 1 eq) in dry DMF (5 mL) was added 1-(chloromethyl)-4-methoxy-benzene (72 mg, 463 μmol, 63 μL, 1.20 eq) and Cs₂CO₃ (251 mg, 771 μmol, 2 eq). The mixture was heated at 60°C for 2 hours. The mixture was concentrated. DCM (20 mL) and H₂O (10 mL) was added. The mixture was extracted with DCM (20 mL). The organic layer was washed with H₂O (20 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by preparative LC-MS to give racemic 7-(4-methoxybenzyl)-6-methyl-3-((1*R*,2*R*,4*S*)-2-methyl-7-oxabicyclo[2.2.1]heptan-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (40 mg, 27% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.87 (s, 1H), 7.17 (d, *J* = 8.4 Hz, 2H), 6.84 (d, *J* = 8.8 Hz, 2H), 6.69 (s, 1H), 5.23-5.07 (m, 2H), 4.67 (d, *J* = 4.8 Hz, 2H), 3.78 (s, 3H), 2.71 (d, *J* = 12.0 Hz, 1H), 2.20 (s, 3H), 1.90-1.86 (m, 1H), 1.70-1.66 (m, 2H), 1.57 (s, 3H), 1.39-1.37 (m, 1H), 1.25-1.22 (m, 1H).

LC-MS: *t*_R = 2.228 min (method 13), *m/z* = 380.2 [M + H]⁺.

Example 71:



(S)-7-(4-methoxybenzyl)-6-methyl-3-(1-phenylethyl)imidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (300 mg, 1.58 mmol) and (S)-2-phenylpropanoic acid (261 mg, 1.74 mmol, 237 μ L, 1.10 eq) in DCM (10 mL) was added triethylamine (400 mg, 3.95 mmol, 548 μ L, 2.50 eq) and HATU (901 mg, 2.37 mmol, 1.50 eq). The mixture was stirred at 15°C for 12 hours. Water (50 mL) was added to the solution. The mixture was extracted with DCM (50 mL $\times$ 2). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by flash silica gel chromatography (Eluent of 0~20% Ethyl acetate/petroleum ether). (S)-N-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-phenylpropanamide (450 mg, 1.56 mmol, 99% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.81 (s, 1H), 7.24-7.37 (m, 5H), 6.74 (s, 1H), 4.37-4.52 (m, 2H), 3.92 (s, 3H), 3.67 (q, J = 7.1 Hz, 1H), 2.40 (s, 3H), 1.56 (d, J = 7.2 Hz, 3H).

Step 2: To a solution of (S)-N-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-phenylpropanamide (450 mg, 1.58 mmol, 1 eq) in dioxane (10 mL) was added POCl₃ (485 mg, 3.16 mmol, 294 μ L, 2 eq). The mixture was stirred at 90°C for 12 hours. The reaction mixture was concentrated under reduced pressure to remove dioxane. The residue was quenched by addition H₂O (50 mL) at 0°C, basified by addition saturated aqueous NaHCO₃ (10 mL) and then extracted with EtOAc (50 mL $\times$ 2). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by flash silica gel chromatography (Eluent of 0~100% Ethyl acetate/petroleum ether). (S)-6-methyl-3-(1-phenylethyl)imidazo[1,5-a]pyrazin-8-ol (128 mg, 505 μ mol, 32% yield).


Step 3: To a solution of (S)-6-methyl-3-(1-phenylethyl)imidazo[1,5-a]pyrazin-8-ol (128 mg, 505 μ mol, 1 eq) and 1-(chloromethyl)-4-methoxybenzene (95 mg, 0.61 mmol, 83 μ L, 1.20 eq) in DMF (10 mL) was added Cs₂CO₃ (329.29 mg, 1.01 mmol, 2 eq). The mixture was stirred at 60°C for 3 hours. The reaction mixture was concentrated under reduced pressure. The residue was purified by preparative LC-MS was further purified by SFC. (S)-7-(4-methoxybenzyl)-6-methyl-3-(1-phenylethyl)imidazo[1,5-a]pyrazin-8(7H)-one (76.40 mg, 203

μmol, 40% yield) was obtained.

¹H NMR (CDCl₃, 400 MHz): δ 7.98 (s, 1H), 7.17-7.31 (m, 5H), 7.12 (d, *J* = 8.4 Hz, 2H), 6.82 (d, *J* = 8.4 Hz, 2H), 6.46 (s, 1H), 5.10 (dd, *J* = 15.6 Hz, 2H), 4.26 (q, *J* = 7.2 Hz, 1H), 3.76 (s, 3H), 2.05 (s, 3H), 1.81 (d, *J* = 7.6 Hz, 3H).

- 5 LC-MS: *t*_R = 2.218 min (method 17), *m/z* = 374.2 [M + H]⁺. SFC: *t*_R = 1.641 min, ee% > 99%, [α]_D²⁰ = -51.4 (c=0.11, MeOH).

Example 72:



10 **(*R*)-7-(4-methoxybenzyl)-6-methyl-3-(1-phenylethyl)imidazo[1,5-a]pyrazin-8(7*H*)-one:**

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (200 mg, 1.05 mmol, 1 eq) and (*R*)-2-phenylpropanoic acid (190 mg, 1.27 mmol, 173 μL, 1.20 eq) in DCM (10 mL) was added triethylamine (267 mg, 2.64 mmol, 365 μL, 2.50 eq) and HATU (602 mg, 1.58 mmol, 1.50 eq). The mixture was stirred at 15°C for 12 hours. Water (50 mL) was added to the solution. The mixture was extracted with DCM (50 mL × 2). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by flash silica gel chromatography (Eluent of 0~20% Ethyl acetate/petroleum ether). (*R*)-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-phenylpropanamide (300 mg, 1.02 mmol, 97% yield) was obtained.

- 20 ¹H NMR (CDCl₃, 400 MHz): δ 7.81 (s, 1H), 7.24-7.37 (m, 5H), 6.75 (s, 1H), 4.37-4.52 (m, 2H), 3.91 (s, 3H), 3.67 (q, *J* = 7.2 Hz, 1H), 2.40 (s, 3H), 1.56 (d, *J* = 7.2 Hz, 3H).

- Step 2: To a solution of (*R*)-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-phenylpropanamide (450 mg, 1.58 mmol, 1 eq) in dioxane (10 mL) was added POCl₃ (727 mg, 4.74 mmol, 440 μL, 3 eq). The mixture was stirred at 90°C for 12 hours. The reaction mixture was concentrated under reduced pressure to remove dioxane. The residue was quenched by addition of H₂O (50 mL) at 0°C, basified by addition saturated aqueous NaHCO₃ (10 mL) and then extracted with EtOAc (50 mL × 2). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under


reduced pressure. The residue was purified by flash silica gel chromatography (Eluent of 0~100% Ethyl acetate/petroleum ether). (*R*)-6-methyl-3-(1-phenylethyl)imidazo[1,5-*a*]pyrazin-8-ol (150 mg, 592 μ mol, 37% yield) was obtained.

Step 3: To a solution of (*R*)-6-methyl-3-(1-phenylethyl)imidazo[1,5-*a*]pyrazin-8-ol (100 mg, 395 μ mol, 1 *eq*) and 1-(chloromethyl)-4-methoxybenzene (74 mg, 474 μ mol, 65 μ L, 1.20 *eq*) in DMF (5 mL) was added Cs₂CO₃ (257 mg, 790 μ mol, 2 *eq*). The mixture was stirred at 60°C for 3 hours. The reaction mixture was concentrated under reduced pressure. The residue was purified by preparative LC-MS. (*R*)-7-(4-methoxybenzyl)-6-methyl-3-(1-phenylethyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (140 mg, 95% yield) was obtained.

¹H NMR (CDCl₃, 400 MHz): δ 7.98 (s, 1H), 7.17-7.32 (m, 5H), 7.12 (d, *J* = 8.4 Hz, 2H), 6.82 (d, *J* = 8.4 Hz, 2H), 6.46 (s, 1H), 5.10 (dd, *J* = 16.0 Hz, 2H), 4.26 (q, *J* = 7.2 Hz, 1H), 3.76 (s, 3H), 2.05 (s, 3H), 1.81 (d, *J* = 6.8 Hz, 3H).

LC-MS: *t*_R = 2.184 min (method 18), *m/z* = 374.1 [M + H]⁺. SFC: *t*_R = 1.324 min, ee% = 97.8%, [α]_D²⁰ = +50.7 (c=0.11, MeOH).

Example 73:



3-(1,4-dimethylpiperidin-4-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (150 mg, 791 μ mol) in DCM (5 mL) was added 1,4-dimethylpiperidine-4-carboxylic acid hydrochloride (169 mg, 870 μ mol, 1.10 *eq*, HCl), triethylamine (240 mg, 2.37 mmol, 329 μ L, 3 *eq*) and HATU (361 mg, 949 μ mol, 1.20 *eq*). The mixture was stirred at 15°C for 16 hours. The mixture was concentrated. The residue was purified by preparative TLC (DCM/MeOH=10/1) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-1,4-dimethylpiperidine-4-carboxamide (150 mg, 65% yield).

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-1,4-dimethylpiperidine-4-carboxamide (100 mg, 342 μ mol, 1 *eq*) in dry dioxane (5 mL) was added POCl₃ (105 mg,


684 μmol , 64 μL , 2 *eq*). The mixture was heated at 80°C for 4 hours. The mixture was cooled to 15°C and poured into water (5 mL). The mixture was adjusted to pH 8 by saturated aqueous NaHCO_3 and concentrated. 10% MeOH in DCM (20 mL) was added to the residue and filtered, the filtrate was concentrated to give 3-(1,4-dimethylpiperidin-4-yl)-6-methylimidazo[1,5-a]pyrazin-8-ol (90 mg).

Step 3: To a solution of 3-(1,4-dimethylpiperidin-4-yl)-6-methylimidazo[1,5-a]pyrazin-8-ol (60 mg, 230 μmol , 1 *eq*) in DMF (2 mL) was added 1-(chloromethyl)-4-methoxybenzene (54 mg, 346 μmol , 1.50 *eq*) and Cs_2CO_3 (150 mg, 461 μmol , 2 *eq*). The mixture was heated at 60°C for 2 hours. The mixture was concentrated. DCM (20 mL) and H_2O (10 mL) were added. The mixture was extracted with DCM (20 mL). The organic layer was washed with H_2O (20 mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by preparative LC-MS to give 3-(1,4-dimethylpiperidin-4-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one (20 mg, 23% yield).

^1H NMR (CDCl_3 400 MHz): δ 7.92 (s, 1H), 7.17 (d, J = 8.8 Hz, 2H), 6.94 (s, 1H), 6.84 (d, J = 8.8 Hz, 2H), 5.15 (s, 2H), 3.78 (s, 3H), 2.58-2.56 (m, 2H), 2.49-2.45 (m, 2H), 2.38-2.35 (m, 2H), 2.25 (s, 3H), 2.19 (s, 3H), 1.86-1.80 (m, 2H), 1.42 (s, 3H).

LC-MS: t_R = 1.747 min (method 13), m/z = 381.2 $[\text{M} + \text{H}]^+$.

Example 74:



3-(6-chloro-2,3-dihydro-1*H*-inden-1-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (200 mg, 1.05 mmol) and 6-chloro-2,3-dihydro-1*H*-indene-1-carboxylic acid (206 mg, 1.05 mmol, 1 *eq*) in DCM (5 mL) was added HATU (479 mg, 1.26 mmol, 1.20 *eq*) and DIPEA (407 mg, 3.15 mmol, 550 μL , 3 *eq*). The mixture was stirred at 18°C for 16 hours. The mixture was washed with H_2O (20 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic was dried over Na_2SO_4 and concentrated under vacuum. 6-chloro-*N*-((3-methoxy-5-methylpyrazin-2-

yl)methyl)-2,3-dihydro-1*H*-indene-1-carboxamide (312 mg, 864 μ mol, 82% yield) was obtained.


Step 2: To a solution of 6-chloro-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2,3-dihydro-1*H*-indene-1-carboxamide (0.262 g, 1eq) in dioxane (10 mL) was added POCl₃ (363 mg, 2.37 mmol, 220 μ L, 3 eq). The mixture was stirred at 80°C for 3 hours. The mixture was quenched by H₂O (20 mL) and adjusted pH > 7 by saturated aqueous NaHCO₃. The mixture was extracted with DCM (25 mL $\times$ 3). The combined organic was dried over Na₂SO₄ and concentrated under vacuum. 3-(6-chloro-2,3-dihydro-1*H*-inden-1-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (236 mg) was obtained and directly used to next step.

Step 3: To a solution of 3-(6-chloro-2,3-dihydro-1*H*-inden-1-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (236 mg, 787 μ mol, 1 eq) and 1-(chloromethyl)-4-methoxybenzene (148 mg, 945 μ mol, 1.20 eq) in DMF (12 mL) was added Cs₂CO₃ (513 mg, 1.57 mmol, 2 eq). The mixture was stirred at 60°C for 2.5 hours. The mixture was concentrated under vacuum. The residue was quenched with H₂O (15 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic phases were dried over Na₂SO₄ and concentrated under vacuum. The residue was purified by preparative HPLC. 3-(6-chloro-2,3-dihydro-1*H*-inden-1-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (99 mg, 237 μ mol, 30% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.93 (s, 1H), 7.30-7.18 (m, 4H), 6.98 (s, 1H), 6.88 (d, *J* = 8.8 Hz, 2H), 6.64 (s, 1H), 5.18 (s, 2H), 4.71 (t, *J* = 8.4 Hz 1H), 3.79 (s, 3H), 3.16-3.13 (m, 1H), 3.05-2.96 (m, 1H), 2.64-2.61 (m, 1H), 2.52-2.49 (m, 1H), 2.17 (s, 3H).

LC-MS: *t*_R = 2.358 min (method 17), *m/z* = 420.1 [M + H]⁺.

Example 75:



7-(4-methoxybenzyl)-6-methyl-3-(3-methyl-5-oxopyrrolidin-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

Step 1: To a solution of ethyl 3-methyl-5-oxopyrrolidine-3-carboxylate (200 mg, 1.17 mmol, 1 eq) in THF (4 mL) and H₂O (2 mL) was added LiOH·H₂O (147.06 mg, 3.50 mmol, 3 eq). The mixture was stirred at 20°C for 16 hours. The mixture was acidified to pH=2 by 1 M HCl and extracted with ethyl acetate (20 mL × 3). The combined organic layers were washed with
 5 H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give 3-methyl-5-oxopyrrolidine-3-carboxylic acid (100 mg, 60% yield).

Step 2: To a cooled (0°C) solution of 3-methyl-5-oxopyrrolidine-3-carboxylic acid (100 mg, 0.7 mmol, 1 eq) in DCM (2 mL) was added oxalyl dichloride (98 mg, 768 μmol, 67 μL, 1.10 eq) and one drop of dry DMF was added. The mixture was stirred at 20°C for 1 hour. The
 10 colorless solution of 3-methyl-5-oxopyrrolidine-3-carbonyl chloride (112.89 mg) in DCM (2 mL) was directly used for the next step.

Step 3: To a cooled (0°C) solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (120 mg, 633 μmol, 1 eq, HCl) in dry DCM (3 mL) was added triethylamine (192 mg, 1.90 mmol, 263 μL, 3 eq) and a solution of 3-methyl-5-oxopyrrolidine-3-carbonyl chloride (112 mg, 696
 15 μmol, 1.10 eq) in dry DCM (2 mL) dropwise. The mixture was stirred at 20°C for 1 hour. H₂O (5 mL) was added and the mixture was extracted with DCM (20 mL × 2). The combined organic layer was washed with H₂O (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by preparative TLC (DCM/MeOH=10/1) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-3-methyl-5-oxopyrrolidine-3-carboxamide (75 mg,
 20 42% yield).

Step 4: A solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-3-methyl-5-oxopyrrolidine-3-carboxamide (100 mg, 359 μmol, 1 eq) in Eaton's reagent (7.7 wt% phosphorus pentoxide solution in methanesulfonic acid) (2 mL) was heated at 60°C for 16 hours. The mixture was cooled to 15°C and poured into ice (5 g). The mixture was adjusted to pH=8 by 7 M
 25 NH₃/MeOH and concentrated. 10% MeOH in DCM (20 mL) was added to the residue and filtered, the filtrate was concentrated to give 4-(8-hydroxy-6-methylimidazo[1,5-a]pyrazin-3-yl)-4-methylpyrrolidin-2-one (100 mg).


Step 5: To a solution of 4-(8-hydroxy-6-methylimidazo[1,5-a]pyrazin-3-yl)-4-methylpyrrolidin-2-one (100 mg, 406 μmol, 1 eq) in DMF (5 mL) was added 1-(chloromethyl)-4-methoxybenzene (76 mg, 487 μmol, 66 μL, 1.20 eq) and Cs₂CO₃ (265 mg, 812 μmol, 2 eq).
 30 The mixture was heated at 60°C for 2h. The mixture was concentrated. DCM (20 mL) and H₂O (10 mL) was added. The mixture was extracted with DCM (20 mL). The organic layers were washed with H₂O (20 mL), dried over Na₂SO₄, filtered and concentrated. The mixture

was purified by preparative HPLC to give 7-(4-methoxybenzyl)-6-methyl-3-(3-methyl-5-oxopyrrolidin-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one (40 mg, 27% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.87 (s, 1H), 7.15 (d, *J* = 8.4 Hz, 2H), 6.84 (d, *J* = 8.4 Hz, 2H), 6.65 (s, 1H), 5.93 (brs, 1H), 5.16 (s, 2H), 4.25 (d, *J* = 9.6 Hz, 1H), 3.77 (s, 3H), 3.53 (d, *J* = 10.4 Hz, 1H), 3.00 (d, *J* = 16.8 Hz, 1H), 2.57 (d, *J* = 16.4 Hz, 1H), 2.20 (s, 3H), 1.63 (s, 3H).

LC-MS: *t_R* = 1.764 min (method 11), *m/z* = 367.1 [M + H]⁺.

Example 76:



10 3-(1-methoxy-2-methylpropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one:

Step 1: To a solution of 3-methoxy-2,2-dimethylpropanoic acid (115 mg, 870 μmol, 1 eq) in DCM (5 mL) was added oxalyl dichloride (121 mg, 957 μmol, 84 μL, 1.10 eq) at 0 °C, followed by one drop of DMF. The mixture was stirred at 20 °C for 1 hour. The mixture was directly used to next step.

Step 2: To a solution of 3-methoxy-2,2-dimethylpropanoyl chloride (150 mg, 791 μmol, 1 eq, HCl) and triethylamine (120 mg, 1.19 mmol, 164 μL, 1.50 eq) in DCM (10mL) was added (3-methoxy-5-methylpyrazin-2-yl)methanamine (131 mg, 870 μmol, 1.10 eq) in DCM (5mL). The mixture was stirred at 20°C for 1 hour. The mixture was quenched with H₂O (20 mL) and extracted with DCM (15 mL × 3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. 3-methoxy-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2,2-dimethylpropanamide (142 mg, 522 μmol, 66% yield) was obtained.

Step 3: To a solution of 3-methoxy-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2,2-dimethylpropanamide (102 mg, 382 μmol, 1 eq) in dioxane (5 mL) was added POCl₃ (117 mg, 763 μmol, 71 μL, 2 eq). The mixture was stirred at 80 °C for 2 hours. The mixture was quenched with water (20 mL) and extracted with DCM (20 mL × 3). The combined organic layers were dried with Na₂SO₄ and concentrated under vacuum. 3-(1-methoxy-2-

methylpropan-2-yl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one (45 mg, 191 μ mol, 50% yield) was obtained.


Step 4: To a solution of 3-(1-methoxy-2-methylpropan-2-yl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one (62 mg, 264 μ mol, 1 eq) and 1-(chloromethyl)-4-methoxybenzene (49.52 mg, 316 μ mol, 43.06 μ L, 1.20 eq) in DMF (5 mL) was added Cs₂CO₃ (171.72 mg, 527 μ mol, 2 eq).

The mixture was stirred at 60°C for 3 hours. The mixture washed with H₂O (20 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The residue was purified by preparative HPLC. 3-(1-methoxy-2-methylpropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one (25.15 mg, 71 μ mol, 27% yield) was obtained.

¹H NMR (CDCl₃, 400 MHz): δ 7.91 (s, 1H), 7.20-7.17 (m, 3H), 6.86 (d, *J* = 8.4 Hz, 2H), 5.15 (s, 2H), 3.79 (s, 3H), 3.59 (s, 2H), 3.35 (s, 3H), 2.18 (s, 3H), 1.52 (s, 6H).

LC-MS: *t*_R = 2.050 min (method 13), *m/z* = 356.1 [M + H]⁺.

Example 77:



3-isopropyl-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one:

Step 1: To a cold (0°C) solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (150 mg, 791 μ mol), Et₃N (176 mg, 1.74 mmol, 241 μ L, 2.20 eq) in anhydrous DCM (5 mL) was added isobutyryl chloride (93 mg, 870 μ mol, 91 μ L, 1.10 eq). The solution was stirred at 0°C for 0.5h. The mixture was diluted with water (20 mL), extracted with DCM (20 mL $\times$ 2). The organic layer was washed with brine (20 mL), dried over Na₂SO₄ and concentrated in vacuo. *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)isobutyramide (160 mg, 717 μ mol, 91% yield) was obtained.

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)isobutyramide (160 mg, 717 μ mol, 1 eq) in dioxane (5 mL) was added POCl₃ (220 mg, 1.43 mmol, 133 μ L, 2 eq). The mixture was stirred at 90°C 2 hours. The mixture was concentrated in vacuo. 3-isopropyl-8-methoxy-6-methylimidazo[1,5-a]pyrazine (130 mg, 633 μ mol, 88%

yield) was

obtained.


Step 3: A solution of 3-isopropyl-8-methoxy-6-methylimidazo[1,5-a]pyrazine (130 mg, 633 μmol , 1 eq) in 2M HCl(aq) (4 mL) and dioxane (8 mL) was stirred at 90°C for 2 hours. The mixture was concentrated in vacuo. The residue was purified by silica gel chromatography (DCM:MeOH=10:1). 3-isopropyl-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (120 mg, 628 μmol , 99% yield) was obtained.

Step 4: To a solution of 3-isopropyl-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (120 mg, 628 μmol , 1 eq) in DMF (8 mL) was added 1-(chloromethyl)-4-methoxy-benzene (118 mg, 753 μmol , 103 μL , 1.20 eq) and Cs_2CO_3 (307 mg, 941 μmol , 1.50 eq). The mixture was stirred at 60°C for 16 hours. The mixture was filtered. The filtrate was purified by pre-HPLC (base). 3-isopropyl-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (90 mg, 287 μmol , 46% yield) was obtained.

^1H NMR (CDCl_3 400 MHz): δ 7.90 (s, 1H), 7.16 (d, $J=8.4$ Hz, 1H), 6.85 (d, $J=8.4$ Hz, 1H), 6.71 (s, 1H), 5.16 (s, 2H), 3.78 (s, 2H), 3.18-3.11 (m, 1H), 2.19 (s, 3H), 1.41-1.79 (d, $J=7.2$ Hz, 6H).

LC-MS: t_R = 1.92 min (method 13), m/z = 312.1 $[\text{M} + \text{H}]^+$.

Example 78:



6-methyl-7-((2-methylthiazol-4-yl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

Into a vial was added 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (30mg, 0.13 mmol), cesium carbonate (84 mg, 0.26 mmol), 4-(chloromethyl)-2-methylthiazole (23 mg, 0.15 mmol) and sodium iodide (23 mg, 0.15 mmol) in DMF (2 mL).


The reaction was heated to 70°C, and stirred over night. The mixture was filtered and evaporated and subsequently chromatographed on silicagel to obtain the crude product. Final purification on preparative LC-MS afforded 6-methyl-7-((2-methylthiazol-4-yl)methyl)-3-

(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (3.9 mg, 0.013 mmol) in 10% yield.

¹H NMR (600 MHz, DMSO-*d*₆) δ 8.17 (s, 1H), 7.63 (m, 1H), 7.30 (m, 1H), 5.17 (s, 2H), 3.99 (t, *J* = 3.3 Hz, 1H), 3.97 (t, *J* = 3.1 Hz, 1H), 3.49 (m, 3H), 2.61 (s, 3H), 2.36 (d, *J* = 1.2 Hz, 3H), 1.83 (m, 4H).

LC-MS: *t*_R = 0.39 min (method 5), *m/z* = 344.9 [M + H]⁺.

Example 79:




10 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)-7-(thiophen-3-ylmethyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:

Into a vial was added 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (30mg, 0.129 mmol), cesium carbonate (84 mg, 0.26 mmol), 3-(chloromethyl)thiophene (20 mg, 0.15 mmol) and sodium iodide (23 mg, 0.15 mmol) in DMF (2 mL). The reaction was heated to 70°C, and stirred over night. The mixture was filtered and evaporated and subsequently chromatographed on silicagel to obtain the crude product. Final purification on preparative LC-MS afforded 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)-7-(thiophen-3-ylmethyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (10 mg, 0.0324 mmol) in 25% yield.

¹H NMR (600 MHz, DMSO-*d*₆) δ 8.21 (s, 1H), 7.66 (m, 1H), 7.53 (dd, *J* = 5.0, 2.9 Hz, 1H), 7.35 (dq, *J* = 2.2, 1.0 Hz, 1H), 7.05 (dd, *J* = 5.0, 1.3 Hz, 1H), 5.17 (s, 2H), 3.98 (dt, *J* = 11.4, 3.4 Hz, 2H), 3.49 (m, 3H), 2.26 (d, *J* = 1.2 Hz, 3H), 1.84(m, 4H).

LC-MS: *t*_R = 0.45 min (method 5), *m/z* = 329.9 [M + H]⁺.

Example 80:




6-methyl-3-(tetrahydro-2H-pyran-4-yl)-7-(thiazol-4-ylmethyl)imidazo[1,5-a]pyrazin-8(7H)-one:

Into a vial was added 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (30mg, 0.129 mmol), cesium carbonate (147 mg, 0.450 mmol), 4-(chloromethyl)thiazole hydrochloride (26 mg, 0.15 mmol) and sodium iodide (23 mg, 0.15 mmol) in DMF (2 mL). The reaction was heated to 70°C, and stirred over night. The mixture was filtered and evaporated and subsequently chromatographed on silicagel to obtain the crude product. Final purification on preparative LC-MS afforded 6-methyl-3-(tetrahydro-2H-pyran-4-yl)-7-(thiazol-4-ylmethyl)imidazo[1,5-a]pyrazin-8(7H)-one (7 mg, 0.0153 mmol) in 12% yield.

¹H NMR (600 MHz, DMSO-d₆) δ 9.07 (d, *J* = 1.9 Hz, 1H), 8.12 (s, 1H), 7.63 (m, 1H), 7.57 (dd, *J* = 1.9, 0.9 Hz, 1H), 5.27 (s, 2H), 3.97 (dt, *J* = 11.3, 3.3 Hz, 2H), 3.48 (m, 3H), 2.36 (d, *J* = 1.2 Hz, 3H), 1.84 (m, 4H).

LC-MS: *t_R* = 0.34 min (method 5), *m/z* = 331.0 [M + H]⁺.

Example 81:



7-((3,5-dimethylisoxazol-4-yl)methyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


Into a vial was added 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (25mg, 0.107 mmol), cesium carbonate (70 mg, 0.21 mmol), 4-(chloromethyl)-3,5-dimethylisoxazole (19 mg, 0.129 mmol) and sodium iodide (19 mg, 0.129 mmol) in DMF (1.6 mL). The reaction was heated to 70°C, and stirred over night. The mixture was filtered

and evaporated and subsequently chromatographed on silicagel to obtain the crude product. Final purification on preparative LC-MS afforded 7-((3,5-dimethylisoxazol-4-yl)methyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (11 mg, 0.024 mmol) in 23% yield.

- 5 ¹H NMR (600 MHz, DMSO-*d*₆) δ 8.11 (s, 1H), 7.63 (m, 1H), 5.00 (s, 2H), 3.97 (dt, *J* = 11.3, 3.3 Hz, 2H), 3.46 (m, 3H), 2.27 (s, 3H), 2.24 (d, *J* = 1.2 Hz, 3H), 2.08 (s, 3H), 1.82 (dd, *J* = 7.8, 3.5 Hz, 4H).

LC-MS: *t*_R = 0.38 min (method 5), *m/z* = 343.0 [*M* + *H*]⁺.

10 Example 82:




6-methyl-7-((5-methylisoxazol-3-yl)methyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one:


- Into a vial was added 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (25mg, 0.107 mmol), cesium carbonate (70 mg, 0.21 mmol), 3-(chloromethyl)-5-methylisoxazole (34 mg, 0.13 mmol, 50 %) and sodium iodide (19 mg, 0.13 mmol) in DMF (1.6 mL). The reaction was heated to 70°C, and stirred over night. The mixture was filtered and evaporated and subsequently chromatographed on silicagel to obtain the crude product. Final purification on preparative LC-MS afforded 6-methyl-7-((5-methylisoxazol-3-yl)methyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (9 mg, 0.0194 mmol) in 18% yield.

¹H NMR (600 MHz, DMSO-*d*₆) δ 8.05 (s, 1H), 7.60 (m, 1H), 6.19 (q, *J* = 0.8 Hz, 1H), 5.18 (s, 2H), 3.97 (dt, *J* = 11.3, 3.4 Hz, 2H), 3.45 (m, 3H), 2.37 (d, *J* = 0.9 Hz, 3H), 2.27 (d, *J* = 1.2 Hz, 3H), 1.86 – 1.79 (m, 4H).

- 25 LC-MS: *t*_R = 0.38 min (method 5), *m/z* = 328.9 [*M* + *H*]⁺.

Example 83:**6-methyl-7-((3-methylisoxazol-5-yl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**

- 5 Into a vial was added 6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (25mg, 0.107 mmol), cesium carbonate (70 mg, 0.21 mmol), 5-(chloromethyl)-3-methylisoxazole (17 mg, 0.13 mmol) and sodium iodide (19 mg, 0.13 mmol) in DMF (1.6 mL). The reaction was heated to 70°C, and stirred over night. The mixture was filtered and evaporated and subsequently chromatographed on silicagel to obtain the crude product.
- 10 Final purification on preparative LC-MS afforded 6-methyl-7-((3-methylisoxazol-5-yl)methyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (14 mg, 0.0313 mmol) in 29 % yield.
- ¹H NMR (600 MHz, DMSO-d₆) δ 8.20 (s, 1H), 7.70 (m, 1H), 6.31 (s, 1H), 5.29 (m, 2H), 3.98 (dt, *J* = 11.4, 3.5 Hz, 2H), 3.49 (m, 3H), 2.32 (d, *J* = 1.2 Hz, 3H), 2.19 (s, 3H), 1.85 (m, 4H).
- 15 LC-MS: *t_R* = 0.37 min (method 5), *m/z* = 328.9 [M + H]⁺.

Example 84:**3-(2,6-dimethyltetrahydro-2H-pyran-4-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one:**

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (200 mg, 1.05 mmol, 1 eq) and 2,6-dimethyltetrahydro-2H-pyran-4-carboxylic acid (167 mg, 1.05

mmol, 1 eq) in DCM (10 mL) was added HATU (481 mg, 1.27 mmol, 1.20 eq) and DIPEA (409 mg, 3.16 mmol, 552 μ L, 3 eq). The mixture was stirred at 18°C for 16 hours. The mixture washed with H₂O (20 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2,6-dimethyltetrahydro-2*H*-pyran-4-carboxamide (300 mg, 0.95 mmol, 90% yield) was obtained.


Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2,6-dimethyltetrahydro-2*H*-pyran-4-carboxamide (280 mg, 954 μ mol, 1 eq) in dioxane (10 mL) was added POCl₃ (293 mg, 1.91 mmol, 177 μ L, 2 eq). The mixture was stirred at 80°C for 2 hours. The mixture washed with H₂O (20 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. 3-(2,6-dimethyltetrahydro-2*H*-pyran-4-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (249 mg) was obtained.

Step 3: To a solution of 3-(2,6-dimethyltetrahydro-2*H*-pyran-4-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 765 μ mol, 1 eq) and 1-(chloromethyl)-4-methoxybenzene (156 mg, 995 μ mol, 135 μ L, 1.30 eq) in DMF (10 mL) was added Cs₂CO₃ (500 mg, 1.53 mmol, 2 eq). The mixture was stirred at 60°C for 4 hours. The mixture was concentrated under vacuum. The mixture was washed with H₂O (20 mL) and extracted with DCM (15 mL $\times$ 3). The combined organic layers were washed with H₂O (40 mL $\times$ 3), brine, dried over Na₂SO₄ and concentrated under vacuum. The residue was purified by preparative LC-MS to give 3-(2,6-dimethyltetrahydro-2*H*-pyran-4-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (48 mg, 126 μ mol, 16% yield).

¹H NMR (CDCl₃ 400 MHz): δ 7.91 (s, 1H), 7.16 (d, *J* = 8.4 HZ, 2H), 6.85 (d, *J* = 8.4 HZ, 2H), 6.73 (s, 2H), 5.17 (s, 2H), 3.79 (s, 3H), 3.67-3.63 (m, 2H), 3.14-3.08 (m, 1H), 2.21 (s, 3H), 1.90-1.87 (m, 3H), 1.75-1.62 (m, 2H), 1.28 (d, *J* = 6.0 HZ, 6H).

LC-MS: *t*_R = 2.133 min (method 17), *m/z* = 382.1 [M + H]⁺.

Example 85:



7-(cyclohexylmethyl)-6-methyl-3-propylimidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: To a solution of 3-bromo-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (1 g, 4.39 mmol, 1 eq) and Cs_2CO_3 (2.86 g, 8.78 mmol, 2 eq) in DMF (20 mL) was added (bromomethyl)cyclohexane (1.55 g, 8.78 mmol, 1.22 mL, 2 eq). The mixture was stirred at 60°C for 18 hour. The reaction mixture was filtered and the filtrate was concentrated. The crude mixture was purified by flash chromatography with petroleum ether:ethyl acetate = 5:1 ~ 3:1. 3-bromo-7-(cyclohexylmethyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (1 g, 3.02 mmol, 69% yield) was obtained.

^1H NMR (CDCl_3 , 400 MHz): δ 7.86 (s, 1H), 6.80 (s, 1H), 3.79 (d, J = 7.2 Hz, 2H), 2.29 (s, 3H), 1.75 - 1.66 (m, 6H), 1.22 - 1.04 (m, 5H).

LC-MS: t_R = 0.791 min (method 15), m/z = 325.9 $[\text{M} + \text{H}]^+$.


Step 2: To a solution of 3-bromo-7-(cyclohexylmethyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (200 mg, 617 μmol , 1 eq) and (E)-4,4,5,5-tetramethyl-2-(prop-1-en-1-yl)-1,3,2-dioxaborolane (155 mg, 925 μmol , 1.50 eq) in dioxane (4 mL) and H_2O (1 mL) was added Pd(dppf) Cl_2 (90 mg, 123 μmol , 0.20 eq) and Cs_2CO_3 (402 mg, 1.23 mmol, 2 eq) under a N_2 atmosphere. The mixture was stirred at 90°C for 2 hours under microwave conditions. Water (50 mL) was added and the mixture was extracted with EtOAc (50 mL $\times$ 3), the combined organic layers were washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude mixture was purified by flash chromatography with petroleum ether:ethyl acetate = 1:1. (E)-7-(cyclohexylmethyl)-6-methyl-3-(prop-1-en-1-yl)imidazo[1,5-a]pyrazin-8(7H)-one (150 mg, 504.59 μmol , 82% yield) was obtained.

Step 3: To a solution of (E)-7-(cyclohexylmethyl)-6-methyl-3-(prop-1-en-1-yl)imidazo[1,5-a]pyrazin-8(7H)-one (150 mg, 526 μmol , 1 eq) in EtOAc (30 mL) was added Pd-C (10%, 40 mg, wet) under N_2 . The suspension was degassed under vacuum and purged with H_2 several times. The mixture was stirred under H_2 (15 psi) at 25°C for 18 hours. The mixture was filtered and the residue was washed with EtOAc (20 mL $\times$ 2). The combined organic layers were concentrated. The residue was purified by preparative LC-MS to give 7-(cyclohexylmethyl)-6-methyl-3-propylimidazo[1,5-a]pyrazin-8(7H)-one (95.5 mg, 321 μmol , 61% yield).

^1H NMR (400 MHz): δ 7.83 (s, 1H), 6.66 (s, 1H), 3.78 (d, J = 7.2 Hz, 2H), 2.81 (t, J = 7.6 Hz, 2H), 2.26 (s, 3H), 1.86 - 1.84 (m, 2H), 1.76 - 1.67 (m, 6H), 1.19 - 1.17 (m, 3H), 1.05 - 1.01 (m, 5H).

LC-MS: t_R = 1.68 min (method 17), m/z = 288.3 $[M + H]^+$.

Example 86:



5 3-(2-hydroxypropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: NaH (2.64 g, 66 mmol, 60% in mineral oil, 2.20 eq) was added to 2-hydroxy-2-methylpropanoic acid (3.12 g, 30 mmol, 1 eq) in DMF (30 mL) at 0°C. The mixture was stirred at 20°C for 30 mins. (bromomethyl)benzene (10.26 g, 60 mmol, 7.13 mL, 2 eq) was added to the reaction mixture at 20°C and stirred at 20°C for 16 hours. The mixture was quenched by H₂O (30 mL) and adjusted pH=7 by HCl (1 M, aq). The mixture was extracted with ethyl acetate (30 mL × 3). The combined organic layers were washed by H₂O (20 mL) and brine. The residue was dried over Na₂SO₄ and concentrated under vacuum. ¹H NMR showed the compound was desired product. benzyl 2-(benzyloxy)-2-methylpropanoate (3.98 g, 14 mmol, 47% yield) was obtained.

¹H NMR (CDCl₃, 400 MHz): δ 7.40-7.32 (m, 10H), 5.24 (s, 2H), 4.48 (s, 2H), 1.58 (s, 6H).

Step 2: To a solution of benzyl 2-(benzyloxy)-2-methylpropanoate (2 g, 7.03 mmol, 1 eq) in H₂O (20 mL), THF (20 mL) and MeOH (20 mL) was added NaOH (1.12 g, 27.98 mmol, 3.98 eq). The mixture was stirred at 80°C for 1 hour. The mixture was adjusted pH=2 by aq. HCl (1 M) and extracted with DCM (10 mL × 3). The combined organic layers were concentrated under vacuum. The residue was washed with aq. NaOH (1 M, 5 mL) and extracted with DCM (15 mL × 3). The aqueous solution was adjusted pH=2 by aq. HCl (1 M, aq) and extracted with DCM (10 mL × 3). The combined organic layers were washed with H₂O (15 mL × 2) and brine. The mixture was dried over Na₂SO₄ and concentrated under vacuum. 2-(benzyloxy)-2-methylpropanoic acid (1.36 g, 7 mmol, 100% yield) was obtained.

Step 3: To a solution of 2-(benzyloxy)-2-methylpropanoic acid (500 mg, 2.64 mmol, 1 eq, HCl) in DCM (10 mL) was added DIPEA (1.02 g, 7.92 mmol, 1.38 mL, 3 eq). The mixture was added (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (513 mg, 2.64

mmol, 1 eq) and HATU (1.20 g, 3.17 mmol, 1.20 eq) . The mixture was stirred at 18 °C for 16 hours. The mixture washed with H₂O (20 mL) and extracted with DCM (20 mL × 3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. 2-(benzyloxy)-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methylpropanamide (617 mg, 1.75 mmol, 66% yield) was obtained.

Step 4: To a solution of 2-(benzyloxy)-*N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-2-methylpropanamide (1.40 g, 4.25 mmol, 1 eq) in dioxane (20 mL) was added POCl₃ (1.30 g, 8.50 mmol, 790 µL, 2 eq) . The mixture was stirred at 80°C for 2 hours. The mixture was quenched with H₂O (15 mL) and adjusted pH >7 by saturated aqueous NaHCO₃. The mixture was extracted with DCM (20 mL × 3). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated under vacuum. 3-(2-(benzyloxy)propan-2-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (375 mg, 1.26 mmol, 30% yield) was obtained.


Step 5: To a solution of 3-(2-(benzyloxy)propan-2-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (264 mg, 888 µmol, 1 eq) in DMF (10 mL) was added Cs₂CO₃ (579 mg, 1.78 mmol, 2 eq) and 1-(chloromethyl)-4-methoxybenzene (180.76 mg, 1.15 mmol, 1578 µL, 1.30 eq). The mixture was stirred at 60°C for 4 hours. The mixture was concentrated under vacuum. The mixture was washed with H₂O (25 mL) and extracted with DCM (20mL). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. 3-(2-(benzyloxy)propan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (445 mg, 998 µmol, 64% yield) was obtained.

Step 6: To a solution of 3-(2-(benzyloxy)propan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (345 mg, 826 µmol, 1 eq) in MeOH (60 mL) was added Pd/C (10%, wet) (40 mg). The mixture was stirred at pressure of H₂ (30 psi). The mixture was stirred at 18°C for 8 hours. The mixture was filtered. The filtered solution was concentrated under vacuum. The residue was purified by TLC (petroleum ether:ethyl acetate=1:1). 3-(2-hydroxypropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (98 mg, 292 µmol, 35% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.86 (s, 1H), 7.47 (s, 1H), 7.17 (d, *J* =12 Hz, 2H), 6.85 (d, *J* =8.4 Hz, 2H), 5.17 (s, 2H), 3.79 (s, 3H), 2.216-2.192 (m, 4H), 1.76 (s, 3H).

LC-MS: *t*_R = 1.906 min (method 13), *m/z* = 328.2 [M + H]⁺.

Example 87:




3-(2-fluoropropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one:

To a solution of 3-(2-hydroxypropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (75 mg, 229.09 μmol , 1 eq) in DCM (10 ml) was added DAST (40.6 mg, 252 μmol , 33 μL , 1.10 eq) at -78°C . The mixture was stirred at 18°C for 2 hours. The mixture was quenched with H_2O (10 mL) and extracted with DCM (15 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 and concentrated under vacuum. The residue was purified with TLC (petroleum ether:ethyl acetate=1:1). 3-(2-fluoropropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (68 mg, 206 μmol , 64% yield) was obtained.

^1H NMR (CDCl_3 400 MHz): δ 7.89 (s, 1H), 7.19-7.16 (m, 3H), 6.86 (d, $J=8.4$ Hz, 2H), 5.18 (s, 2H), 3.79 (s, 3H), 2.20 (s, 3H), 1.91 (s, 3H), 1.86 (s, 3H).

LC-MS: t_R = 1.906 min (method 13), m/z = 328.2 $[\text{M} + \text{H}]^+$.

Example 88:



7-(4-methoxybenzyl)-6-methyl-3-(7-oxoazepan-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine hydrochloride (500 mg, 2.64 mmol, 1 eq) and 7-oxoazepane-4-carboxylic acid (456 mg, 2.90 mmol, 1.10 eq) in DCM (45 mL) was added HATU (1.20 g, 3.17 mmol, 1.20 eq) and DIPEA (1.02 g, 7.92 mmol, 1.38 mL, 3 eq). The mixture was stirred at 18°C for 16 hours. The mixture was quenched with H_2O (30 mL) and extracted with DCM (25 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 and concentrated under vacuum. *N*-((3-methoxy-5-

methylpyrazin-2-yl)methyl)-7-oxoazepane-4-carboxamide (493 mg, 1.58 mmol, 60% yield) was obtained.

Step 2: *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-7-oxoazepane-4-carboxamide (463 mg, 1.58 mmol, 1 eq) was added to Eaton's reagent (7.7 wt% phosphorus pentoxide solution in methanesulfonic acid) (3.04 g, 12.77 mmol, 2 mL, 8.08 eq). The mixture was stirred at 60°C for 7 hours. The mixture was added to ice (30 g). The mixture was adjusted pH>7 by NH₃ (MeOH). The mixture was concentrated under vacuum. The residue was washed with DCM : MeOH =10:1. The mixture was filtered. The filtered solution was concentrated under vacuum. 6-methyl-3-(7-oxoazepan-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (45 mg, 165.83 μmol, 11% yield) was obtained.


¹H NMR (MeOD 400 MHz): δ 7.77 (s, 1H), 7.23 (s, 2H), 3.49-3.39 (m, 3H), 2.83-2.79 (m, 2H), 2.55-2.51 (m, 1H), 2.19 (s, 3H), 2.11-2.09 (m, 2H), 1.93-1.86 (m, 2H).

Step 3: To a solution of 6-methyl-3-(7-oxoazepan-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (45 mg, 173 μmol, 1 eq) and 1-(chloromethyl)-4-methoxybenzene (32 mg, 207 μmol, 28 μL, 1.20 eq) in DMF (3 mL) was added Cs₂CO₃ (113 mg, 346 μmol, 2 eq). The mixture was stirred at 60°C for 2 hours. The mixture was washed with H₂O (10 mL) and extracted with DCM (15 mL × 3). The combined organic layers were washed with water (30 mL × 3), dried over Na₂SO₄ and concentrated under vacuum. The residue was purified by TLC (DCM:MeOH=10:1). 7-(4-methoxybenzyl)-6-methyl-3-(7-oxoazepan-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (12 mg, 32 μmol, 18% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.89 (s, 1H), 7.15 (d, *J* = 8.8 Hz, 2H), 6.84 (d, *J* = 8.4 Hz, 2H), 6.70 (s, 1H), 6.36 (brs, 1H), 5.16 (s, 2H), 5.16 (s, 2H), 3.77 (s, 3H), 3.62-3.49 (m, 1H), 3.42-3.30 (m, 1H), 3.20-3.10 (m, 1H), 2.77-2.72 (m, 1H), 2.61-2.55 (m, 1H), 2.21 (s, 3H), 2.10-2.01 (m, 4H).

LC-MS: *t*_R = 1.748 min (method 13), *m/z* = 381.2 [M + H]⁺.

Example 89:



7-(4-methoxybenzyl)-6-methyl-3-(5-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one:

Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (200 mg, 1.05 mmol, 1 eq, HCl) and 5-methyltetrahydrofuran-3-carboxylic acid (137 mg, 1.05 mmol, 1 eq) in DCM (10 mL) was added HATU (481 mg, 1.27 mmol, 1.20 eq) and DIPEA (409 mg, 3.16 mmol, 553 μ L, 3 eq). The mixture was stirred at 18°C for 16 hours. The mixture washed with H₂O (20 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-5-methyltetrahydrofuran-3-carboxamide (211 mg, 795 μ mol, 76% yield) was obtained.

Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)-5-methyltetrahydrofuran-3-carboxamide (191 mg, 720 μ mol, 1 eq) in dioxane (5 mL) was added POCl₃ (221 mg, 1.44 mmol, 134 μ L, 2 eq). The mixture was stirred at 80°C for 3 hours. The mixture washed with H₂O (20 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. 6-methyl-3-(5-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (167 mg) was obtained.

Step 3: To a solution of 6-methyl-3-(5-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (138 mg, 592 μ mol, 1 eq) and 1-(chloromethyl)-4-methoxybenzene (120 mg, 769 μ mol, 1.30 eq) in DMF (10 mL) was added Cs₂CO₃ (386 mg, 1.18 mmol, 2 eq). The mixture was stirred at 60°C for 4 hours. The mixture was concentrated under vacuum. The mixture was washed with H₂O (20 mL) and extracted with DCM (15 mL $\times$ 3). The combined organic layers were washed with H₂O (40 mL $\times$ 3), brine, dried over Na₂SO₄ and concentrated under vacuum. The residue was purified by pre-HPLC(base). 7-(4-methoxybenzyl)-6-methyl-3-(5-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one (10 mg, 28 μ mol, 5% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.90 (s, 1H), 7.16 (d, *J* = 8.4 Hz, 2H), 6.87-6.82 (m, 3H), 5.17 (s, 2H), 4.19-4.14 (m, 3H), 3.79-3.71 (m, 4H), 2.54-2.47 (m, 1H), 2.20 (s, 3H), 2.04-1.96 (m, 1H), 1.40 (s, *J* = 6.0 Hz, 3H).

LC-MS: *t*_R = 2.036 min (method 13), *m/z* = 354.2 [M + H]⁺.


Example 90:

^1H NMR (CDCl_3 , 400 MHz): δ 7.95 (s, 1H), 7.13 (d, J = 8.4 Hz, 2H), 6.91 (s, 1H), 6.82 (d, J = 8.4 Hz, 2H), 6.79 (s, 1H), 5.19-5.06 (m, 2H), 4.76 (q, J = 7.2 Hz, 1H), 3.76 (s, 3H), 2.41 (s, 3H), 2.14 (s, 3H), 1.92 (d, J = 7.2 Hz, 1H).

LC-MS: t_R = 2.532 min (method 11), m/z = 395.1 $[\text{M} + \text{H}]^+$.

5

Example 91:



3-(7-(4-methoxybenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)-3-methylpyrrolidine-1-sulfonamide:

- 10 Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (200 mg, 1.05 mmol, 1 eq, HCl) and 1-((benzyloxy)carbonyl)-3-methylpyrrolidine-3-carboxylic acid (304 mg, 1.16 mmol, 1.10 eq) in DCM (10 mL) was added HATU (479 mg, 1.26 mmol, 1.20 eq) and DIPEA (407 mg, 3.15 mmol, 550 μL , 3 eq). The mixture was stirred at 18°C for 16 hours. The mixture was quenched with H_2O (30 mL) and extracted with DCM (25 mL $\times$ 3). The
- 15 combined organic layers were dried over Na_2SO_4 and concentrated under vacuum. Benzyl 3-(((3-methoxy-5-methylpyrazin-2-yl)methyl)carbamoyl)-3-methylpyrrolidine-1-carboxylate (420 mg, 1.01 mmol, 96% yield) was obtained.

- Step 2: To a solution of benzyl 3-(((3-methoxy-5-methylpyrazin-2-yl)methyl)carbamoyl)-3-methylpyrrolidine-1-carboxylate (390 mg, 979 μmol , 1 eq) in dioxane (15 mL) was added
- 20 POCl_3 (300 mg, 1.96 mmol, 182 μL , 2 eq). The mixture was stirred at 80°C for 3 hours. The solution was quenched with H_2O (20 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 and concentrated under vacuum. Benzyl 3-methyl-3-(6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)pyrrolidine-1-carboxylate (170 mg, 423 μmol , 43% yield) was obtained.

- 25 Step 3: To a solution of benzyl 3-methyl-3-(6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)pyrrolidine-1-carboxylate (290 mg, 791 μmol , 1 eq) and di-*tert*-butyl dicarbonate (207 mg, 950 μmol , 218 μL , 1.20 eq) in MeOH (200 mL) was added Pd/C (10%, wet) (140 mg). The suspension was degassed under vacuum and purged with H_2 several times. The

mixture was stirred under H₂ (30 psi) at 25°C for 5 hours. The mixture was filtered. The filtered solution was concentrated under vacuum. *tert*-butyl 3-methyl-3-(6-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-3-yl)pyrrolidine-1-carboxylate (204 mg, 614 μmol, 78% yield) was obtained.


- 5 Step 4: To a solution of *tert*-butyl 3-methyl-3-(6-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-3-yl)pyrrolidine-1-carboxylate (274 mg, 824 μmol, 1 eq) and 1-(chloromethyl)-4-methoxybenzene (155 mg, 989 μmol, 135 μL, 1.20 eq) in DMF (15 mL) was added Cs₂CO₃ (537 mg, 1.65 mmol, 2 eq). The mixture was stirred at 80°C for 14 hours. The mixture was concentrated under vacuum. The mixture was washed with H₂O (15 mL) and extracted with
10 DCM (15 mL × 3). The combined organic layers were washed with H₂O (30 mL × 2), dried over Na₂SO₄ and concentrated under vacuum. *tert*-butyl 3-(7-(4-methoxybenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-3-yl)-3-methylpyrrolidine-1-carboxylate (291 mg, 537 μmol, 65% yield) was obtained.

- Step 5: To a solution of *tert*-butyl 3-(7-(4-methoxybenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-3-yl)-3-methylpyrrolidine-1-carboxylate (288 mg, 636 μmol, 1
15 eq) in ethyl acetate (4 mL) was added HCl/EtOAc (4 M, 4 mL, 25 eq). The mixture was stirred at 18°C for 4 hours. The mixture was concentrated under vacuum. 7-(4-methoxybenzyl)-6-methyl-3-(3-methylpyrrolidin-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (247 mg, crude, HCl) was obtained. The product was directly used for the next step.

- 20 Step 6: To a solution of 7-(4-methoxybenzyl)-6-methyl-3-(3-methylpyrrolidin-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (250 mg, 709 μmol, 1 eq) and sulfuric diamide (82 mg, 851 μmol, 51 μL, 1.20 eq) in dioxane (20 mL) was added DIPEA (183 mg, 1.42 mmol, 248 μL, 2 eq). The mixture was stirred at 100°C for 24 hours. The mixture was quenched with H₂O (20 mL) and extracted with DCM (20 mL × 3). The combined organic layers were dried over Na₂SO₄ and
25 concentrated under vacuum. The residue was purified pre-HPLC (base). 3-(7-(4-methoxybenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-3-yl)-3-methylpyrrolidine-1-sulfonamide (14 mg, 32 μmol, 5% yield) was obtained.

- ¹H NMR (CDCl₃ 400 MHz): δ 7.91 (s, 1H), 7.16 (d, *J* = 8.4 Hz, 2H), 7.85 (s, 1H), 7.18 (d, *J* = 8.0 Hz, 2H), 6.88-6.82 (m, 3H), 5.17 (s, 2H), 4.79 (m, 1H), 4.30 (d, *J* = 10.4 Hz, 1H), 3.79
30 (s, 3H), 3.52-3.47 (m, 2H), 3.26-3.20 (m, 1H), 2.73-2.64 (m, 1H), 2.23-2.17 (m, 5H), 1.65-1.59 (m, 4H).


LC-MS: *t*_R = 2.055 min (method 13), *m/z* = 432.2 [M + H]⁺.

Example 92:**6-(cyclopentylmethyl)-7-(4-methoxybenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:**

- 5 6-(bromomethyl)-7-(4-methoxybenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (250 mg, 0.439 mmol) was dissolved in THF (20 ml), at -78°C. cyclopentylmagnesium bromide (0.9 ml, 1.8 mmol, 2 molar in ether) was added and the reaction was allowed to warm to room temperature overnight. Cyclopentylmagnesium bromide (0.9 ml, 1.8 mmol, 2 molar in ether) was added. After 1 hour the reaction was
- 10 warmed to room temperature. After two hours, the reaction mixture was quenched with sat. NH₄Cl, extracted with AcOEt and organics were washed with brine. The combined organics layers were dried with Na₂SO₄, filtered and concentrated. The mixture was purified via preparative LC-MS and 6-(cyclopentylmethyl)-7-(4-methoxybenzyl)-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (26 mg, 0.049 mmol) was isolated in 11% yield, as the
- 15 TFA salt.

¹H NMR (500 MHz, Chloroform-*d*) δ 13.26 (bs, 1H), 8.14 (s, 1H), 7.12 (d, *J* = 8.2 Hz, 2H), 6.88 (m, 3H), 5.21 (s, 2H), 4.16 (d, *J* = 11.7 Hz, 2H), 3.80 (s, 3H), 3.60 (t, *J* = 11.8 Hz, 2H), 3.46 (t, *J* = 12.4 Hz, 1H), 2.74 (s, 1H), 2.57 (d, *J* = 7.0 Hz, 2H), 2.25 – 2.07 (m, 2H), 1.90 (m, 4H), 1.65 (m, 4H), 1.23 (dt, *J* = 13.2, 7.1 Hz, 2H).

- 20 LC-MS: *t*_R = 0.69 min (method 5), *m/z* = 422.0 [M + H]⁺.


Example 93:

7-(4-methoxybenzyl)-6-methyl-3-morpholinoimidazo[1,5-a]pyrazin-8(7H)-one:

To a mixture of 3-bromo-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (300 mg, 0.86 mmol) and morpholine (150 mg, 1.72 mmol) in DMSO (5 mL) were added CsF (261 mg, 1.72 mmol) and K₂CO₃ (238 mg, 1.72 mmol). The mixture was stirred at 100°C for 24 hours. The mixture was diluted with water (20 mL) and extracted with DCM (10 mL × 3). The combine organic layers were washed with water (10 mL × 2); dried over Na₂SO₄ and evaporated under vacuum. The residue was purified by preparative TLC (ethyl acetate) to give 7-(4-methoxybenzyl)-6-methyl-3-morpholinoimidazo[1,5-a]pyrazin-8(7H)-one (18 mg, 6% yield).

¹H NMR (CDCl₃, 400 MHz): δ 7.76 (s, 1H), 7.16 (d, *J* = 8.8 Hz, 2H), 6.85 (d, *J* = 8.8 Hz, 2H), 6.63 (s, 1H), 5.15 (s, 2H), 3.89 (t, *J* = 4.8 Hz, 4H), 3.79 (s, 3H), 3.21 (t, *J* = 4.8 Hz, 4H), 2.18 (s, 3H).

LC-MS: *t*_R = 2.03 min (method 13), *m/z* = 355.1 [M + H]⁺.

Example 94:**7-(4-methoxybenzyl)-6-methyl-3-((tetrahydrofuran-3-yl)amino)imidazo[1,5-a]pyrazin-8(7H)-one:**

3-bromo-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (60 mg, 0.172 mmol) and tetrahydrofuran-3-amine (0.04 mL, 0.465 mmol) were mixed in NMP (2 mL) and DIPEA (0.23 mL, 1.317 mmol).

The reaction was heated for 4 hours at 250°C in the microwave oven.


The reaction was purified on silica gel, via preparative LC-MS, and preparative TLC (10% EtOH in ethyl acetate) to give 7-(4-methoxybenzyl)-6-methyl-3-((tetrahydrofuran-3-yl)amino)imidazo[1,5-a]pyrazin-8(7H)-one (2 mg, 0.005 mmol) in 3% yield.

^1H NMR (600 MHz, Dimehtylsulfoxide- d_6) δ 8.50 (m, NH), 7.68 (s, 1H), 7.31 (s, 1H) 7.22 (d, $J = 7.1$ Hz, 2H), 6.88 (d, $J = 7.1$ Hz, 2H), 4.87 (m, 1H), 4.35 (d, $J = 14$ Hz, 2H), 4.11 (m, 1H), 4.02 (m, 1H), 3.92 (m, 1H), 3.86 (m, 1H), 3.71 (s, 3H), 2.60 (m, 1H), 2.5 (m, 1H) 2.31 (s, 3H).

LC-MS: $t_R = 0.54$ min (method 19), $m/z = 355.2$ $[\text{M} + \text{H}]^+$.

5

Example 95:



(*R*)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7*H*)-one:

- 10 3-bromo-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one (80 mg, 0.23 mmol) and (*R*)-3-methylmorpholine (34.7 mg, 0.039 ml, 0.343 mmol) were mixed in NMP (2.0 mL) and DIPEA (0.2 ml, 1.1 mmol) was added.


The reaction was heated for 6.5 hours at 250°C in the microwave oven.

- 15 The reaction was purified directly by preparative LC-MS to give (*R*)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7*H*)-one (10 mg, 0.024 mmol) in 10% yield.

- ^1H NMR (500 MHz, Chloroform- d) δ 8.00 (s, 1H), 7.18 (d, $J = 7.1$ Hz, 2H), 6.90 (d, $J = 7.1$ Hz, 2H), 6.79 (s, 1H), 5.23 (d, $J = 14.0$ Hz, 1H), 5.13 (d, $J = 15.7$ Hz, 1H), 3.94 (m, 1H), 3.85 (m, 1H), 3.81 (s, 3H), 3.60 (m, 1H), 3.48 (dd, $J = 11.7, 7.9$ Hz, 1H), 3.32 (t, $J = 10.3$ Hz, 1H),
20 3.22 (d, $J = 12.2$ Hz, 1H), 2.28 (d, $J = 1.6$ Hz, 3H), 1.29 (td, $J = 7.1, 1.6$ Hz, 1H), 1.00 (dd, $J = 6.4, 1.5$ Hz, 3H).

LC-MS: $t_R = 0.65$ min (method 7), $m/z = 369.1$ $[\text{M} + \text{H}]^+$.

Example 96:




(S)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7H)-one:

- 3-bromo-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (80 mg, 0.230 mmol) and (S)-3-methylmorpholine (0.039 ml, 0.343 mmol) were mixed in NMP (2 mL) and DIPEA (148 mg, 0.2 ml, 1.1 mmol).

The reaction was heated for 6.5 hours at 250°C, in the microwave oven. The reaction was purified directly by preparative LC-MS to give (S)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7H)-one (10 mg, 0.028 mmol) in 12% yield.

- ¹H NMR (500 MHz, Chloroform-*d*) δ 8.00 (s, 1H), 7.18 (d, *J* = 7.1 Hz, 2H), 6.90 (d, *J* = 7.1 Hz, 2H), 6.79 (s, 1H), 5.23 (d, *J* = 14.0 Hz, 1H), 5.13 (d, *J* = 15.7 Hz, 1H), 3.94 (m, 1H), 3.85 (m, 1H), 3.81 (s, 3H), 3.60 (m, 1H), 3.48 (dd, *J* = 11.7, 7.9 Hz, 1H), 3.32 (t, *J* = 10.3 Hz, 1H), 3.22 (d, *J* = 12.2 Hz, 1H), 2.28 (d, *J* = 1.6 Hz, 3H), 1.29 (td, *J* = 7.1, 1.6 Hz, 1H), 1.00 (dd, *J* = 6.4, 1.5 Hz, 3H).
- LC-MS: *t*_R = 0.65 min (method 7), *m/z* = 369.1 [M + H]⁺.

Example 97:



7-(4-methoxybenzyl)-6-methyl-3-(1,4-oxazepan-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one:


- 3-bromo-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (149 mg, 0.428 mmol) and 1,4-oxazepane hydrochloride (100 mg, 0.727 mmol) were mixed in NMP (2.2 mL) and DIPEA (296 mg, 0.4 mL, 2.29 mmol). The reaction was heated for 3 hours at 250°C.

The reaction was purified directly via preparative LC-MS to afford 7-(4-methoxybenzyl)-6-methyl-3-(1,4-oxazepan-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one (27 mg, 0.073 mmol) in 17% yield.

¹H NMR (500 MHz, Chloroform-*d*) δ 7.84 (s, 1H), 7.14 (m, 2H), 6.87 (m, 2H), 6.64 (s, 1H), 5.13 (s, 2H), 3.93 (m, 2H), 3.88 (m, 2H), 3.80 (d, *J* = 1.7 Hz, 3H), 3.73 (m, 4H), 2.22 (s, 3H), 2.09 (m, 2H).

LC-MS: *t*_R = 0.56 min (method 7), *m/z* = 369.1 [M + H]⁺.

Example 98:




3-(2,2-dimethylmorpholino)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one:

To a mixture of 3-bromo-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (500 mg, 1.44 mmol) and 2,2-dimethylmorpholine (331 mg, 2.88 mmol) in DMSO (5 mL) were added CsF (328 mg, 2.88 mmol) and K₂CO₃ (299 mg, 2.88 mmol). The mixture was stirred at 100°C for 24 hours. The mixture was diluted with water (20 mL) and extracted with DCM (10 mL × 3). The combined organic layer was washed with water (10 mL × 2); dried over Na₂SO₄ and evaporated under vacuum. The residue was purified by preparative TLC (ethyl acetate) to give 3-(2,2-dimethylmorpholino)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one (15 mg, 3% yield).

¹H NMR (CDCl₃, 400 MHz): δ 7.75 (s, 1H), 7.15 (d, *J* = 8.8 Hz, 2H), 6.85 (d, *J* = 8.8 Hz, 2H), 6.60 (s, 1H), 5.15 (s, 2H), 3.93 (t, *J* = 4.8 Hz, 2H), 3.79 (s, 3H), 3.16 (t, *J* = 4.8 Hz, 2H), 2.98 (s, 2H), 2.18 (s, 3H), 1.37 (s, 6H).

LC-MS: *t*_R = 2.26 min (method 13), *m/z* = 383.1 [M + H]⁺.

Example 99:



7-(3-fluorobenzyl)-3-(hexahydro-4*H*-furo[3,2-*b*]pyrrol-4-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one, stereoisomer 1 and 2:

Step 1: To a solution of 3-bromo-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (3 g, 13.16 mmol) and 1-(bromomethyl)-3-fluorobenzene (2.99 g, 15.79 mmol) in DMF (50 mL) was added K_2CO_3 (3.64 g, 26.3 mmol). The mixture was stirred at 60°C for 12 hours. The mixture was diluted with water (100 mL) and extracted with EtOAc (50 mL $\times$ 3). The combined organic layer was washed with water (50 mL $\times$ 2); dried over Na_2SO_4 and evaporated under vacuum. The residue was washed with EtOAc (10 mL) and filtrated. The filter cake was dried under vacuum to give 3-bromo-7-(3-fluorobenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (2 g, 45% yield).

Step 2: To a mixture of 3-bromo-7-(3-fluorobenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (1.2 g, 3.57 mmol) and hexahydro-2*H*-furo[3,2-*b*]pyrrole (0.5 oxalac acid salt) (678 mg, 4.28 mmol) in DMSO (20 mL) were added CsF (542 mg, 3.57 mmol) and K_2CO_3 (1.23 g, 8.92 mmol). The mixture was stirred at 120°C for 72 hours. The mixture was diluted with water (100 mL) and extracted with DCM (50 mL $\times$ 3). The combined organic layer was washed with water (50 mL $\times$ 2); dried over Na_2SO_4 and evaporated under vacuum. The residue was purified by prep- HPLC to give compound 3 (200 mg, 15% yield).

Step 3: 7-(3-fluorobenzyl)-3-(hexahydro-4*H*-furo[3,2-*b*]pyrrol-4-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one (200 mg, 0.54 mmol) was purified by SFC to give **stereoisomer 1** (48 mg).

^1H NMR ($CDCl_3$, 400 MHz): δ 7.74 (s, 1H), 7.33 - 7.29 (m, 1H), 7.01 - 6.89 (m, 3H), 6.73 (s, 1H), 5.19 (s, 2H), 4.72 - 4.66 (m, 2H), 3.92 - 3.86 (m, 2H), 3.75 - 3.73 (m, 1H), 3.50 - 3.47 (m, 1H), 2.16 - 2.09 (m, 6H), 1.91 - 1.86 (m, 1H).

LC-MS: t_R = 1.90 min (method 12), m/z = 369.1 $[M + H]^+$. SFC-MS: t_R = 4.44 min, ee% > 99%. $[\alpha]_D^{20}$ +133.00 (c = 0.10, DCM).

stereoisomer 2 (32 mg).


^1H NMR ($CDCl_3$, 400 MHz): δ 7.74 (s, 1H), 7.32 - 7.28 (m, 1H), 7.01 - 6.89 (m, 3H), 6.73 (s,

1H), 5.19 (s, 2H), 4.74 - 4.67 (m, 2H), 3.92 - 3.87 (m, 2H), 3.75 - 3.72 (m, 1H), 3.50 - 3.47 (m, 1H), 2.16 - 2.09 (m, 6H), 1.91 - 1.86 (m, 1H).

LC-MS: t_R = 1.89 min (method 12), m/z = 369.1 $[M + H]^+$. SFC-MS: t_R = 5.71 min, ee% > 99%. $[\alpha]_D^{20}$ -82.00 (c = 0.10, DCM).

5

Example 100:



7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2H-pyran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one, stereoisomer 1 and 2:

- 10 Step 1: To a solution of (3-methoxy-5-methylpyrazin-2-yl)methanamine (150 mg, 979.2 μ mol), tetrahydro-2H-pyran-3-carboxylic acid (127.4 mg, 979.2 μ mol) in DCM (10 mL) was added HATU (670.2 mg, 1.8 mmol) and triethylamine (198.2 mg, 1.9 mmol). The mixture was stirred at 25°C for 16 hours. The mixture was diluted with water (15 mL), extracted with DCM (3 $\times$ 30 mL). The combined organic layer was washed with brine (30 mL), dried over
- 15 Na_2SO_4 , filtered and concentrated. The residue was purified by preparative TLC (EA/MeOH=20/1) to give *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)tetrahydro-2H-pyran-3-carboxamide (130 mg, 50% yield).

- Step 2: To a solution of *N*-((3-methoxy-5-methylpyrazin-2-yl)methyl)tetrahydro-2H-pyran-3-carboxamide (130 mg, 490 μ mol) in dioxane (3 mL) was added $POCl_3$ (1.28 g, 490 μ mol).
- 20 The mixture was stirred at 80°C for 3h. The mixture was cooled down to 25°C and concentrated. The residue was neutralized by saturated aq. $NaHCO_3$, extracted with ethyl acetate (2 $\times$ 20 mL). The combined organic layer was washed with brine (20 mL), dried over Na_2SO_4 , filtered and concentrated to give crude 8-methoxy-6-methyl-3-(tetrahydro-2H-pyran-3-yl)imidazo[1,5-a]pyrazine (120 mg, 99% yield). The crude was used directly for the next
- 25 step.

Step 3: To a solution of 8-methoxy-6-methyl-3-(tetrahydro-2H-pyran-3-yl)imidazo[1,5-a]pyrazine (120 mg, 485.3 μ mol) in dioxane (3 mL) was added HCl (2 M, 3 mL). The mixture was stirred at 80°C for 3h. The mixture was cooled down to 25°C and concentrated,

neutralized with saturated aq. NaHCO₃, extracted with DCM (3 × 30 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated to give crude 6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (110 mg, 97% yield). The crude product was used directly for the next step.

- 5 Step 4: To a solution of 6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (100.0 mg, 428.7 μmol) and 1-(bromomethyl)-3-fluorobenzene (121.6 mg, 643.0 μmol) in DMF (5 mL) was added K₂CO₃ (118.5 mg, 857.4 μmol). The mixture was stirred at 80°C for 2h. The mixture was cooled down to 25°C and diluted with water (20 mL), extracted with ethyl acetate (3 × 30 mL). The combined organic layer was washed with brine (30 mL), dried
10 over Na₂SO₄, filtered and concentrated. The residue was purified by preparative TLC (PE/EA=1/1) to give 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (80 mg, 54% yield).

7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one (80 mg, 234.3 μmol) was purified by SFC.

- 15 7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one, **stereoisomer 1** (26 mg, 33% yield) was obtained.

¹H NMR (CDCl₃ 400 MHz): δ 7.92 (s, 1H), 7.33-7.27 (m, 1H), 7.00-6.89 (m, 3H), 6.81 (s, 1H), 5.23 (s, 2H), 4.11-4.03 (m, 2H), 3.69 (t, *J* = 11.2 Hz, 1H), 3.57-3.53 (m, 1H), 3.17-3.14 (m, 1H), 2.18-2.11 (m, 5H), 1.85-1.79 (m, 2H).

- 20 LC-MS: *t*_R = 2.06 min (method 3), *m/z* = 342.1 [M + H]⁺. SFC: *t*_R = 5.286 min, ee% > 99%. α_D²⁰ = -3.0 (c = 0.10, CHCl₃).

7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one, **stereoisomer 2** (28 mg, yield: 35%) was obtained.

- 25 ¹H NMR (CDCl₃ 400 MHz): δ 7.92 (s, 1H), 7.31-7.27 (m, 1H), 7.00-6.89 (m, 3H), 6.81 (s, 1H), 5.22 (s, 2H), 4.11-4.03 (m, 2H), 3.68 (t, *J* = 11.2 Hz, 1H), 3.57-3.53 (m, 1H), 3.17-3.14 (m, 1H), 2.18-2.11 (m, 5H), 1.84-1.78 (m, 2H).

LC-MS: *t*_R = 2.06 min (method 3), *m/z* = 342.2 [M + H]⁺. SFC: *t*_R = 6.404 min, ee% > 99%. α_D²⁰ = +3.0 (c = 0.10, CHCl₃).

30 IN VITRO TESTING

PDE1 INHIBITION ASSAY

PDE1A, PDE1B and PDE1C assays were performed as follows: the assays was performed in 60 μ L samples containing a fixed amount of the PDE1 enzyme¹ (sufficient to convert 20–25% of the cyclic nucleotide substrate), a buffer (50 mM HEPES pH 7.6; 10 mM $MgCl_2$; 0.02% Tween20), 0.1 mg/ml BSA, 15 nM tritium labelled cAMP and varying amounts of inhibitors. Reactions were initiated by addition of the cyclic nucleotide substrate, and reactions were allowed to proceed for 1 h at room temperature before being terminated through mixing with 20 μ L (0.2 mg) yttrium silicate SPA beads (PerkinElmer). The beads were allowed to settle for 1 h in the dark before the plates were counted in a Wallac 1450 Microbeta counter. The measured signals were converted to activity relative to an uninhibited control (100%) and IC_{50} values were calculated using XIFit (model 205, IDBS).


PDE9 INHIBITION ASSAY

A PDE9 assay may for example, be performed as follows: The assay is performed in 60 μ L samples containing a fixed amount of the relevant PDE enzyme (sufficient to convert 20–25% of the cyclic nucleotide substrate), a buffer (50 mM HEPES pH 7.6; 10mM $MgCl_2$; 0.02% Tween20), 0.1mg/ml BSA, 225 pCi of ^3H -labelled cyclic nucleotide substrate, tritium labeled cAMP to a final concentration of 5 nM and varying amounts of inhibitors. Reactions are initiated by addition of the cyclic nucleotide substrate, and reactions are allowed to proceed for one hr at room temperature before being terminated through mixing with 15 μ L 8 mg/mL yttrium silicate SPA beads (Amersham). The beads are allowed to settle for one hr in the dark before the plates are counted in a Wallac 1450 Microbeta counter. The measured signal can be converted to activity relative to an uninhibited control (100 %) and IC_{50} values can be calculated using the XIfit extension to EXCEL.

In the context of the present invention the assay was performed in 60 μ L assay buffer (50 mM HEPES pH 7.6; 10mM $MgCl_2$; 0.02% Tween20) containing enough PDE9 to convert 20–25% of 10 nM ^3H -cAMP and varying amounts of inhibitors. Following a 1 hr incubation the reactions were terminated by addition of 15 μ L 8 mg/mL yttrium silicate SPA beads (Amersham). The beads were allowed to settle for one hr in the dark before the plates were counted in a Wallac 1450 Microbeta counter.

Claims

1. A compound according to formula (I)



(I)

- 5 wherein

n is 0 or 1;

q is 0 or 1;

- 10 R1 is selected from the group consisting of benzyl, indanyl, indoline and 5-membered heteroaryls; all of which can be substituted with a substituent selected from the group consisting of halogen and C₁-C₃ alkyl; or

R1 is selected from the group consisting of saturated monocyclic rings containing 4-6 carbon atoms and 1-2 nitrogen atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine and sulfonamide; or

- 15 R1 is selected from the group consisting of lactams containing 4-6 carbon atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

R1 is selected from the group consisting of bicyclic ethers such as, 7-oxabicyclo[2.2.1]heptane; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

- 20 R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl and tetrahydropyranyl; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine, hydroxy, cyano or methoxy; or

- 25 R1 is a linear or branched C₁-C₃ alkyl, which is substituted with a substituent selected from phenyl and 5-membered heteroaryl, wherein said 5-membered heteroaryl can be substituted with one or more C₁-C₃ alkyls; or

R1 is selected from the group consisting of morpholine, tetrahydrofuran-3-amine, hexahydro-2H-furo[3,2-b]pyrrole and homomorpholine; all of which can be substituted with one or more substituents selected from the group consisting of C₁-C₃ alkyl;

5 R2 is selected from the group consisting of hydrogen, linear or branched C₁-C₈ alkyl, phenyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, benzo[d][1,3]dioxolyl, tetrahydrofuranyl and tetrahydropyranyl; or

R2 is phenyl or pyridyl substituted with one or more substituents selected from the group consisting of hydroxyl, amino, cyano, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₃-C₅ cycloalkoxy, C₃-C₅ cycloalkyl-methoxy, C₁-C₃ fluoroalkoxy, and -NC(O)CH₃; or

10 R2 is a 5-membered heteroaryl which can be substituted with one or more C₁-C₃ alkyl;

R3 is selected from the group consisting of halogen, C₁-C₅ alkyl, C₃-C₅ cycloalkyl and phenyl; or

15 R3 is selected from the group consisting of phenyl substituted one or more times with C₁-C₃ alkyl; methyl substituted one, two or three times with fluorine; ethyl substituted one, two or three times with fluorine;

R4 is hydrogen;

and tautomers and pharmaceutically acceptable addition salts thereof.

2. The compound according to claim 1, wherein

20 R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl, and tetrahydropyranyl;

R2 is selected from the group consisting of, linear or branched C₁-C₈ alkyl, phenyl, and saturated monocyclic C₃-C₈ cycloalkyl; or

R2 is selected from the group consisting of phenyl substituted with a substituent selected from the group consisting of halogen, C₁-C₃ alkyl and methoxy;

5 R3 is selected from the group consisting of C₁-C₃ alkyl and halogen;

R4 is hydrogen.

3. The compound of claim 1, wherein R1 is selected from tetrahydrofuranyl and tetrahydropyranyl.

10

15

20

25

4. The compound of claim 1, wherein the compound is selected from the group consisting of:

6-Benzyl-7-(3-fluorobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

5 6-Benzyl-7-(cyclohexylmethyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

7-(Cyclohexylmethyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

7-(3-Fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

7-(4-Chlorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

6-Bromo-7-(3-fluorobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

10 7-Benzyl-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

15

20

25

- 7-(2-Fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(3-Chlorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(2-Chlorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(3-Methoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 5 6-Methyl-7-(2-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 6-Methyl-7-(4-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-Methoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-Fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 6-Methyl-7-(3-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 10 7-(3-fluorobenzyl)-6-methyl-3-(4-methyltetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 4-(7-(3-fluorobenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-*a*]pyrazin-3-yl)tetrahydro-2*H*-pyran-4-carbonitrile;
- 7-(3-fluorobenzyl)-3-(4-methoxytetrahydro-2*H*-pyran-4-yl)-6-methylimidazo[1,5-*a*]pyrazin-
- 15 8(7*H*)-one;
- 7-(3-fluorobenzyl)-3-(4-fluorotetrahydro-2*H*-pyran-4-yl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- (*R*)-7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 20 (*S*)-7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-2-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- (*R*)-7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- (*S*)-7-(3-fluorobenzyl)-6-methyl-3-(tetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- (*R*)-7-(3-fluorobenzyl)-6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 25 one;
- (*S*)-7-(3-fluorobenzyl)-6-methyl-3-(3-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

- 7-(3-fluorobenzyl)-6-methyl-3-(1-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one ;
- 3-(2,2-difluorocyclopropyl)-7-(3-fluorobenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-6-methyl-3-((1*R*,2*S*)-2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 5 7-(3-fluorobenzyl)-6-methyl-3-((1*R*,2*R*)-2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-6-methyl-3-((1*S*,2*S*)-2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-6-methyl-3-((1*S*,2*R*)-2-methylcyclopropyl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 10 7-(3-fluorobenzyl)-6-methyl-3-((2*S*,3*R*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-6-methyl-3-((2*S*,3*S*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 15 7-(3-fluorobenzyl)-6-methyl-3-((2*R*,3*R*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-6-methyl-3-((2*R*,3*S*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-3-((1*R*,2*S*)-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one;
- 20 7-(3-fluorobenzyl)-3-((1*R*,2*R*)-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-3-((1*S*,2*S*)-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(3-fluorobenzyl)-3-((1*S*,2*R*)-2-fluorocyclopropyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one;
- 7-(4-cyclopropoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 25 7-(4-(difluoromethoxy)benzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
- 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-a]pyrazin-8(7H)-one;

- 7-(4-(cyclopropylmethoxy)benzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-benzyl-6-ethyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 6-ethyl-7-(4-methoxybenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 5 3-((6-methyl-8-oxo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-7(8*H*)-yl)methyl)benzonitrile;
- 4-((6-methyl-8-oxo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-7(8*H*)-yl)methyl)benzonitrile;
- N*-(4-((6-methyl-8-oxo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-7(8*H*)-yl)methyl)phenyl)acetamide;
- 10 7-(4-chloro-3-methoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(2-ethylbenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(benzo[d][1,3]dioxol-5-ylmethyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 15 7-(3-chloro-4-methoxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-aminobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-hydroxybenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 20 6-ethyl-7-(3-fluorobenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-methoxybenzyl)-6-methyl-3-((2*S*,3*R*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-methoxybenzyl)-6-methyl-3-((2*S*,3*S*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 25 7-(4-methoxybenzyl)-6-methyl-3-((2*R*,3*R*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-methoxybenzyl)-6-methyl-3-((2*R*,3*S*)-2-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

- 7-(4-methoxybenzyl)-6-methyl-3-propylimidazo[1,5-a]pyrazin-8(7*H*)-one;
- 7-((6-methoxypyridin-3-yl)methyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 6,7-dimethyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 5 7-ethyl-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 6-methyl-7-propyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 7-isopropyl-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 7-isopentyl-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 7-(cyclopentylmethyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 10 2-((6-methyl-8-oxo-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-7(8*H*)-yl)methyl)benzonitrile;
- 7-(cycloheptylmethyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 7-(4-methoxybenzyl)-6-methyl-3-(3-methyltetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 15 7-(4-methoxybenzyl)-6-methyl-3-((1*R*,2*R*,4*S*)-2-methyl-7-oxabicyclo[2.2.1]heptan-2-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- (*S*)-7-(4-methoxybenzyl)-6-methyl-3-(1-phenylethyl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- (*R*)-7-(4-methoxybenzyl)-6-methyl-3-(1-phenylethyl)imidazo[1,5-a]pyrazin-8(7*H*)-one;
- 3-(1,4-dimethylpiperidin-4-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one;

20

25

- 3-(6-chloro-2,3-dihydro-1*H*-inden-1-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-methoxybenzyl)-6-methyl-3-(3-methyl-5-oxopyrrolidin-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 5 3-(1-methoxy-2-methylpropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 3-isopropyl-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 6-methyl-7-((2-methylthiazol-4-yl)methyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 10 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)-7-(thiophen-3-ylmethyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)-7-(thiazol-4-ylmethyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-((3,5-dimethylisoxazol-4-yl)methyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 15 6-methyl-7-((5-methylisoxazol-3-yl)methyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 6-methyl-7-((3-methylisoxazol-5-yl)methyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 3-(2,6-dimethyltetrahydro-2*H*-pyran-4-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 20 a]pyrazin-8(7*H*)-one;
- 7-(cyclohexylmethyl)-6-methyl-3-propylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 3-(2-hydroxypropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 3-(2-fluoropropan-2-yl)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-methoxybenzyl)-6-methyl-3-(7-oxoazepan-4-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 25 7-(4-methoxybenzyl)-6-methyl-3-(5-methyltetrahydrofuran-3-yl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;
- 7-(4-methoxybenzyl)-6-methyl-3-(1-(4-methylthiazol-2-yl)ethyl)imidazo[1,5-*a*]pyrazin-8(7*H*)-one;

3-(7-(4-methoxybenzyl)-6-methyl-8-oxo-7,8-dihydroimidazo[1,5-a]pyrazin-3-yl)-3-methylpyrrolidine-1-sulfonamide;

6-(cyclopentylmethyl)-7-(4-methoxybenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

5 3-(morpholino)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one;

7-(4-methoxybenzyl)-6-methyl-3-((tetrahydrofuran-3-yl)amino)imidazo[1,5-a]pyrazin-8(7*H*)-one;

(*R*)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7*H*)-one;

(*S*)-7-(4-methoxybenzyl)-6-methyl-3-(3-methylmorpholino)imidazo[1,5-a]pyrazin-8(7*H*)-one;

10 7-(4-methoxybenzyl)-6-methyl-3-(1,4-oxazepan-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

3-(2,2-dimethylmorpholino)-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one;

7-(3-fluorobenzyl)-3-((3*aS*,6*aS*)-hexahydro-4*H*-furo[3,2-*b*]pyrrol-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one

15 7-(3-fluorobenzyl)-3-((3*aR*,6*aR*)-hexahydro-4*H*-furo[3,2-*b*]pyrrol-4-yl)-6-methylimidazo[1,5-a]pyrazin-8(7*H*)-one;

(*R*)-7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

(*S*)-7-(3-fluorobenzyl)-6-methyl-3-(tetrahydro-2*H*-pyran-3-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

20 or a pharmaceutically acceptable salt of any of these compounds.

5. The compound according to claim 1 which is

7-(Cyclohexylmethyl)-6-methyl-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

or a pharmaceutically acceptable salt thereof.

25 6. The compound according to claim 1 which is

6-Methyl-7-(4-methylbenzyl)-3-(tetrahydro-2*H*-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

or a pharmaceutically acceptable salt thereof.

7. The compound according to claim 1 which is

7-(4-Methoxybenzyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
or a pharmaceutically acceptable salt thereof.

5 8. The compound according to claim 1 which is

7-(benzo[d][1,3]dioxol-5-ylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
or a pharmaceutically acceptable salt thereof.

10 9. The compound according to claim 1 which is

7-((6-methoxypyridin-3-yl)methyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
or a pharmaceutically acceptable salt thereof.

15 10. The compound according to claim 1 which is

7-(cycloheptylmethyl)-6-methyl-3-(tetrahydro-2H-pyran-4-yl)imidazo[1,5-a]pyrazin-8(7H)-one;
or a pharmaceutically acceptable salt thereof.

11. The compound according to claim 1 which is

20 3-isopropyl-7-(4-methoxybenzyl)-6-methylimidazo[1,5-a]pyrazin-8(7H)-one;
or a pharmaceutically acceptable salt thereof.


12. The compound according to claim 1 which is

25 7-(4-methoxybenzyl)-6-methyl-3-(1-(4-methylthiazol-2-yl)ethyl)imidazo[1,5-a]pyrazin-8(7H)-one;

or a pharmaceutically acceptable salt thereof.

13. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of claims 1 to 12 and one or more pharmaceutically acceptable carriers, diluents and excipients.

14. A pharmaceutical composition comprising therapeutically effective amount of a compound according to formula (I), and one or more pharmaceutically acceptable carriers, diluents and excipients



(I)

wherein

n is 0 or 1;

q is 0 or 1;

R1 is selected from the group consisting of benzyl, indanyl, indoline and 5-membered heteroaryls; all of which can be substituted with a substituent selected from the group consisting of halogen and C₁-C₃ alkyl; or

R1 is selected from the group consisting of saturated monocyclic rings containing 4-6 carbon atoms and 1-2 nitrogen atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine and sulfonamide; or

R1 is selected from the group consisting of lactams containing 4-6 carbon atoms; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

R1 is selected from the group consisting of bicyclic ethers such as, 7-oxabicyclo[2.2.1]heptane; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl and fluorine; or

5 R1 is selected from the group consisting of linear or branched C₁-C₈ alkyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, tetrahydrofuranyl and tetrahydropyranyl; all of which can be substituted one or more times with one or more substituents selected from the group consisting of methyl, fluorine, hydroxy, cyano or methoxy; or

10 R1 is a linear or branched C₁-C₃ alkyl, which is substituted with a substituent selected from phenyl and 5-membered heteroaryl, wherein said 5-membered heteroaryl can be substituted with one or more C₁-C₃ alkyls; or

R1 is selected from the group consisting of morpholine, tetrahydrofuran-3-amine, hexahydro-2H-furo[3,2-b]pyrrole and homomorpholine; all of which can be substituted with one or more substituents selected from the group consisting of C₁-C₃ alkyl;

15 R2 is selected from the group consisting of hydrogen, linear or branched C₁-C₈ alkyl, phenyl, saturated monocyclic C₃-C₈ cycloalkyl, oxetanyl, benzo[d][1,3]dioxolyl, tetrahydrofuranyl and tetrahydropyranyl; or

R2 is phenyl or pyridyl substituted with one or more substituents selected from the group consisting of hydroxyl, amino, cyano, halogen, C₁-C₃ alkyl, C₁-C₃ alkoxy, C₃-C₅ cycloalkoxy, C₃-C₅ cycloalkyl-methoxy, C₁-C₃ fluoroalkoxy, and -NC(O)CH₃; or

20 R2 is a 5-membered heteroaryl which can be substituted with one or more C₁-C₃ alkyl;

R3 is selected from the group consisting of hydrogen, halogen, C₁-C₅ alkyl, C₃-C₅ cycloalkyl and phenyl; or

25 R3 is selected from the group consisting of phenyl substituted one or more times with C₁-C₃ alkyl; methyl substituted one, two or three times with fluorine; ethyl substituted one, two or three times with fluorine;

R4 is hydrogen;

and tautomers and pharmaceutically acceptable addition salts thereof;

with the proviso that R2 and R3 cannot be hydrogen at the same time;

30 with the proviso, that the compound of formula (I) is not one of the three following compounds:

3-methyl-7-(4-(trifluoromethoxy)benzyl)imidazo[1,5-a]pyrazin-8(7*H*)-one;

7-butyl-3-methylimidazo[1,5-a]pyrazin-8(7*H*)-one; and

7-(4-methoxybenzyl)-3-methylimidazo[1,5-a]pyrazin-8(7*H*)-one.

15. Use of a compound of any one of claims 1-12 in the manufacture of a medicament for the treatment of a neurodegenerative disorder, selected from the group consisting of Alzheimer's Disease, Parkinson's Disease and Huntington's Disease or for the treatment of a
- 5 psychiatric disorder such as Attention Deficit hyperactivity Disorder (ADHD), depression, anxiety, narcolepsy, cognitive impairment and cognitive impairment associated with schizophrenia (CIAS), or another brain disease like restless leg syndrome.