

Patent Claims

1. A biparatopic antibody comprising a first antigen-binding site that binds to a first epitope on human CCL2 and a second different antigen-binding site that binds a second different epitope on human CCL2, wherein the biparatopic antibody comprises a constant heavy chain domain of human IgG1 isotype
- wherein
- i) said first antigen-binding site comprises
- a VH domain comprising (a) a CDR-H1 comprising the amino acid sequence SHYGXS of SEQ ID NO: 57 wherein X is I or T, (b) a CDR-H2 comprising the amino acid sequence $GX^1IX^2IFX^3TANYAQKFQG$ of SEQ ID NO: 58 wherein X^1 is V, I, or H, X^2 is P or H, and X^3 is H or G, and (c) a CDR-H3 comprising the amino acid sequence YDAHYGELDF of SEQ ID NO: 59;
- and
- a VL domain comprising (d) a CDR-L1 comprising the amino acid sequence RASQHVSDAYLA of SEQ ID NO: 60; (e) a CDR-L2 comprising the amino acid sequence DASDRAE of SEQ ID NO: 61, and (f) a CDR-L3 comprising the amino acid sequence HQYIHLHSFT of SEQ ID NO: 62;
- and
- ii) said second antigen-binding site comprises
- a VH domain comprising (a) a CDR-H1 comprising the amino acid sequence HTYMH of SEQ ID NO: 76, (b) a CDR-H2 comprising the amino acid sequence RIDPXNHNTKFDPKFQG of SEQ ID NO: 77 wherein X is D or E, and (c) a CDR-H3 comprising the amino acid sequence GVFGFFXH of SEQ ID NO: 78 wherein X is D or E; and
- a VL domain comprising (d) a CDR-L1 comprising the amino acid sequence $KAX^1EDIYNRX^2A$ of SEQ ID NO: 79 wherein X^1 is F or T and X^2 is R or L, (e) a CDR-L2 comprising the amino acid sequence GATSLEH of SEQ ID NO: 80, and (f) a CDR-L3 comprising the amino acid sequence QQFXSAPYT of SEQ ID NO: 81 wherein X is W or R.

2. The biparatopic antibody according to claim 1,
wherein

- i) said first antigen-binding site comprises
a VH domain comprising (a) a CDR-H1 comprising the amino acid
sequence SHYGXS of SEQ ID NO: 57 wherein X is I or T , (b) a CDR-
H2 comprising the amino acid sequence GX¹IX²IFX³TANYAQKFQG
of SEQ ID NO: 58 wherein X¹ is V, I, or H, X² is P or H, and X³ is H
or G, (c) a CDR-H3 comprising the amino acid sequence
YDAHYGELDF of SEQ ID NO: 59, (d) a FR-H1 comprising the
amino acid sequence QVQLVQSGAEVKKPGSSVKV SCKASGGTF
of SEQ ID NO:63, (e) a FR-H2 comprising the amino acid sequence
WVRQAPGQGLEWMG of SEQ ID NO:64, (f) a FR-H3 comprising
the amino acid sequence RVTITADESTSTAYMELSSLRSEDTAVY
YCAR of SEQ ID NO:65, and (g) a FR-H4 comprising the amino acid
sequence WGQGTLVTVSS of SEQ ID NO:66;
and
a VL domain comprising (h) a CDR-L1 comprising the amino acid
sequence RASQHVSDAYLA of SEQ ID NO: 60; (i) a CDR-L2
comprising the amino acid sequence DASDRAE of SEQ ID NO: 61,
and (j) a CDR-L3 comprising the amino acid sequence HQYIHLHSFT
of SEQ ID NO: 62, (k) a FR-L1 comprising the amino acid sequence
EIVLTQSPATLSLSPGERATLSC of SEQ ID NO:67, (l) a FR-L2
comprising the amino acid sequence WYQQKPGQAPRLLIY of SEQ
ID NO:68, (m) a FR-L3 comprising the amino acid sequence
GVPARFSGSGSGTDFTLTISLSEPEDFAVYYC of SEQ ID NO:69,
and (n) a FR-L4 comprising the amino acid sequence GQGTKVEIK of
SEQ ID NO:70;
and
- ii) said second antigen-binding site comprises
a VH domain comprising (a) a CDR-H1 comprising the amino acid
sequence HTYMH of SEQ ID NO: 76, (b) a CDR-H2 comprising the
amino acid sequence RIDPXNHNTKFDPKFQG of SEQ ID NO: 77
wherein X is D or E, (c) a CDR-H3 comprising the amino acid
sequence GVFGFFXH of SEQ ID NO:78 wherein X is D or E, (d) a
FR-H1 comprising the amino acid sequence
QVQLVQSGAEVKKPGSSVKV SCKASGLTIS of SEQ ID NO:82, (e)

a FR-H2 comprising the amino acid sequence WVRQAPGQGLEWMG of SEQ ID NO:83, (f) a FR-H3 comprising the amino acid sequence RVTITADTSTSTAYMELSSLRSEDVAVYYCAR of SEQ ID NO:84, and (g) a FR-H4 comprising the amino acid sequence

5

WGQGTTVTVSS of SEQ ID NO:85;

and

a VL domain comprising (h) a CDR-L1 comprising the amino acid sequence KAX¹EDIYNRX²A of SEQ ID NO: 79 wherein X¹ is F or T and X² is R or L, (i) a CDR-L2 comprising the amino acid sequence GATSLEH of SEQ ID NO: 80, (j) a CDR-L3 comprising the amino acid sequence QQFXSAPYT of SEQ ID NO: 81 wherein X is W or R, (k) a FR-L1 comprising the amino acid sequence

10

DIQMTQSPSSLSASVGDRVTITC of SEQ ID NO:86, (l) a FR-L2 comprising the amino acid sequence WYQQKPGKAPKLLIH of SEQ ID NO:87, (m) a FR-L3 comprising the amino acid sequence GVPSRFSGSGSGTDYTLTISSLQPEDFATYYC of SEQ ID NO:88, and (n) a FR-L4 comprising the amino acid sequence FGGGTKVEIK of SEQ ID NO:89.

15

3. The biparatopic antibody according to claim 1 or 2, wherein

20

A) i) said first antigen-binding site comprises

a VH domain comprising the amino acid sequence of SEQ ID NO:71; and a VL domain comprising the amino acid sequence of SEQ ID NO:75; and

25

ii) said second antigen-binding site comprises

a VH domain comprising the amino acid sequence of SEQ ID NO:91; and a VL domain comprising the amino acid sequence of SEQ ID NO:93;

or

30

B) i) said first antigen-binding site comprises

a VH domain comprising the amino acid sequence of SEQ ID NO:71; and a VL domain comprising the amino acid sequence of SEQ ID NO:75; and

ii) said second antigen-binding site comprises
a VH domain comprising the amino acid sequence of SEQ ID NO:90;
and a VL domain comprising the amino acid sequence of SEQ ID
NO:94.

- 5 4. The biparatopic antibody according to anyone of the preceding claims, wherein
the biparatopic antibody binds to human CCL2 with a 10 times higher affinity
at pH 7.4, than at pH 5.8.
- 10 5. The biparatopic antibody according to anyone of the preceding claims, wherein
the biparatopic antibody, comprises a human IgG1 heavy chain constant
domain comprising one or more of the following mutations (Kabat EU
numbering)
- i) Q311R and/or P343R; and/or
- ii) L234Y, L235W, G236N, P238D, T250V, V264I, H268D, Q295L, T307P,
K326T and/or A330K; and/or
- 15 iii) M428L, N434A and/or Y436T; and/or
- iv) Q438R and/or S440E.
6. Isolated nucleic acid encoding the antibody according to any one of the
preceding claims.
7. A host cell comprising the nucleic acid of claim 6.
- 20 8. A method of producing an antibody comprising culturing the host cell of claim
7 so that the antibody is produced.
9. The method of claim 8, further comprising recovering the antibody from the
host cell.
- 25 10. A pharmaceutical formulation comprising the biparatopic antibody according
any one of claims 1 to 5 and a pharmaceutically acceptable carrier.
11. The biparatopic antibody according any one of claims 1 to 5 for use as a
medicament.

12. Use of the biparatopic antibody according any one of claims 1 to 5 in the manufacture of a medicament for treating cancer.