

REPLACED SHEETS		SG
☒	Description : paged 23,68 to paged	
☒	Claim(s) : paged 14 to paged 4	
☐	Drawing(s) : paged to paged	
☒	Abstract : paged 17 to paged 1	
DATE OF RECEIPT		

- 1 -

COSMETIC AND/OR PHARMACEUTICAL COMPOSITIONS

AND METHODS FOR THEIR USE

This invention relates to cosmetic and/or pharmaceutical compositions for use in protecting human or animal tissues, especially epithelial tissues from the damaging effects of contact with heavy metals, and methods for using said compositions.

Many metal ions are required by organisms in trace amounts as essential nutrients. For example, many functions carried out by polypeptides, e.g. enzymatic, structural and immunological functions, require metallic cofactors.

However other metallic ions, in particular ions of the heavy metals, can adversely affect these functions, especially if the ions are present in non-physiological quantities. Thus overexposure to metals in the environment can lead to toxic effects.

Ecological studies conducted in the industrialised countries of the world have shown that the amounts of these metals present in the environment are increasing. This has led to an increase in levels of heavy metals in the tissues of organisms, including man, through ingestion of contaminated foodstuffs and exposure to metals in, for example, the atmosphere.

The effects of an accumulation of heavy metal ions can be extremely dangerous and their toxicity is thought to be due in part to disruption of the tertiary and quaternary structure of proteins, causing reduction in catalytic, (i.e. enzymatic) activity.

The thus disrupted proteins may be antigenic and cause an immune response. In this regard they may be recognised by the body as "foreign polypeptide biotic agents" which cause an autoimmune response (many common allergies are caused by heavy metals, in association with use of detergents or other environmental factors).



A further mechanism that has been shown to be responsible for the toxic effects of metals is the competitive substitution of natural physiological enzymic cofactors by heavy metals when at non-physiological concentrations.

Thus, the control of heavy metal pollutants in the atmosphere is essential if metal-related diseases are to be prevented. It is therefore expedient to seek products that efficiently and selectively block heavy metal ions rendering them incapable of entering the body and thereby prevent them exerting their toxic effects.

The extensive contamination of the environment by heavy metals and their ubiquitous presence in the ecosystem, mean that the skin and the accessible mucous membranes form the largest surface area available for heavy metals to accumulate on and subsequently be absorbed into the body. It is also known that many of the cutaneous allergic manifestations that have until now been attributed to detergents or other causes have now been shown to involve heavy metals in their aetiopathogenesis.

Of special note are copper, cobalt, zinc, manganese, mercury and nickel which present significant health hazards in the mining industry, and lead, from exhaust fumes. It has been postulated that the accumulation of lead in developing children may exert a deleterious effect on memory and powers of concentration.

US-A- 4 725 670 discloses the extraction from plant material of metallothioneins and their use in the form of pharmaceutical preparations for treating acute and chronic heavy metal poisoning, and metal deficiency phenomena. There is no mention however that metallothioneins may be used as protective agents against the effects of heavy metal ion contact.



- 3 -

It has now been surprisingly found that certain polypeptides having a capacity to sequester heavy metals can be used to manufacture cosmetic and/or pharmaceutical compositions which enable the aforementioned problems to be overcome.

Thus according to the present invention there is provided a topical cosmetic and/or pharmaceutical composition for the external protection of human or animal tissues from the toxic effect of contact with heavy metals, said composition comprising a metal sequestering component capable of binding metal ions and a physiologically inert carrier suitable for topical administration, characterised in that said metal sequestering component comprises one or more metal binding peptide having a high proportion of cysteine residues.

It is preferred that said metal-binding peptide comprises ligands capable of forming metal-thiolate clusters with heavy metal ions.

Thus, it is preferred that the metal-binding peptide has between about 10 and 50% cysteine residues, more preferably between about 25 and 45%, based on the total number of residues.

Metal binding peptides having a high proportion of cysteine exist in nature. Examples of such naturally-occurring peptides include the metallothioneins. A sub-class of metallothioneins is the so-called phytochelatins. However many authorities now use the term "metallothionein" to cover both.

In more detail, metallothioneins (MTs) are a group of proteins, first discovered in equine kidney in studies on cadmium accumulation by Margoshes and Vallee in 1957. The apoprotein (devoid of bound metal ions) may be capable of binding more than one type of metal. Since then, MTs have been found to bind cadmium, zinc, copper, mercury, silver, gold, lead and bismuth, amongst others.

MTs have been isolated from the tissues of animals, plants and microorganisms. They are non-globular proteins, located intracellularly mainly in the cytoplasm and have been isolated in mammals in the greatest abundance from parenchymatous tissues of liver, kidney and intestines.

The classification and nomenclature of MTs referred to herein is that recommended by "The Committee on the Nomenclature of Metallothionein" from the First International Meeting on Metallothioneins and other Low Molecular Weight Metal-Binding Proteins in Zurich, July 1978.

Three classes of MT are now recognised, as follows:

Class I

Included in this class are all polypeptides related in primary structure to equine renal MT. This material is characterised by the following features:-

- (i) Molecular weight 6000-7000;
- (ii) High metal content;
- (iii) Characteristic amino acid composition (high cysteine content, no aromatic amino acids);
- (iv) Unique distribution of cysteine residues in the amino acid sequence;
- (v) Spectroscopic features characteristic of metal-thiolate complexes and metal-thiolate clusters.

Class II

This class comprises forms displaying only slight or no evolutionary correspondence in terms of primary structure to mammalian forms. Examples are MTs from sea urchin and *N crassa*. They share however the same ability to form metal-thiolate complexes and metal-thiolate clusters as Class I MTs.

Class III

This class comprises homologous, atypical oligo - and polypeptides of the general structure (Glu-Cys)_nX, where n = 2-8 and X is an amino acid such as glycine or β-alanine. They are often referred to as phytochelatins and have been isolated from plant and fungal tissue.

These MTs may occur as oligomeric structures composed of two or more chains, differing in weight from 500 to 2000 daltons, linked via metal-thiolate bridges and/or via disulphide bonds.

Thus, it can be seen that the term metallothionein encompasses an extremely wide variety of proteins varying in structure and size, but sharing a common capacity to bind metal ions in complexes with cysteine side chains to form discrete metal-thiolate clusters. The amino acid sequences of typical Class I, II and III metallothioneins are shown below in Table I:

Table I: Amino acid sequences of a Class I, II and II
1. metallothionein.

<u>Organism</u>	<u>Amino Acid Sequence</u>
Human (Class I) MT-2	- Ac-MDPNCSAA GDSCTCAGS CKCKE CKCTSCCKKS CCSCCPVGCA KCAQGCICKGASDK CSCCA
<i>Saccharomyces cerevisiae</i> (Class II)	- QNEGHECQCQCGSCKNNEQCQKSCSCPTGCNSDDKPCGNKS EETKKSCCSGK
<i>Rauwolfia serpentina</i>	- (Glu - Cys) _n Gly n = 4 - 8



- 6 -

Polypeptides having the functional characteristics of metallothioneins as described above may be used in accordance with the invention. These may be characterised by having specific sequences within the peptide chain such as Cys-Cys, Cys-X-Cys, and Cys-X-Y-Cys wherein X and Y are residues other than Cys.

Thus the metal-binding peptide used in accordance with the invention may be a metallothionein of Class I, and/or II, and/or III as defined in accordance with the International Convention on nomenclature, although other metal-binding peptides having a high proportion of cysteine residues may be used.

Preferably, the composition according to the invention is substantially free of heavy metal ions, in order that the metal sequestering facility is not impaired. Thus, preferably the metallothionein(s) present is/are apometallothionein(s).

It is preferred that the compositions of the invention are formulated in the form of gels, creams, ointments, or body lotions. Additionally, it is envisaged that the compositions of the invention are formulated to be applied to exposed parts of the body in the form of a barrier cream or cosmetic make-up foundation which may preferably comprise a dermatologically inert colouring agent and/or perfume.

It is most preferred that the compositions are formulated in the form of a film-forming, water-resistant mixture which may comprise for example oils, waxes, silicone oils or other similar inert hydrophobic carrier materials. It is of course desirable that such materials do not interact with sulfhydryl groups of the metal-binding peptide. The compositions are preferably water resistant and preferably are capable of remaining on the skin throughout the normal activities of the day, whilst being capable of being removed simply by washing with a detergent such as soap.



- 7 -

The compositions of the invention preferably contain between about 0.01 and 10 percent by weight of said metal-binding peptide, and most preferably between about 0.1 and 5 percent by weight of said metal-binding peptide.

The present invention further provides a cosmetic method for protection of human or animal tissues from contact with heavy metals, said composition comprising a metal sequestering component capable of binding metal ions and a physiologically inert carrier suitable for topical administration, characterised in that said metal sequestering component comprises one or more metal binding peptide having a high proportion of cysteine residues.

Although metal-binding peptides may be used in accordance with the invention, to obtain a protective effect on human or animal tissues against contact with heavy metal ions primarily by incorporating them into cosmetic and/or pharmaceutical compositions, the protective effect of the invention may be obtained in other ways.

Thus in a further embodiment of the present invention, there is provided a mask for the protection of human tissues from the hazardous effects of environmental, particularly airborne heavy metal ion contamination, said mask comprising a filter medium carrying an adsorbent material comprising a metal sequestering component capable of binding metal ions, characterised in that said metal sequestering agent comprises one or more metal-binding peptide having a high proportion of cysteine residues.

Metal-binding peptides can, according to the invention, also be employed in disposable cartridges for masks for the protection of human epithelial tissues from the hazardous effects of airborne heavy metal ion contact.



The metal-binding peptide used according to the invention may be capable of binding many of the heavy metal ions which may be present in the environment. However it is envisaged that it will be most important to protect against lead, cadmium, chromium, mercury, copper and nickel.

Metal-binding peptides useful in producing the compositions, methods, masks and disposable cartridges of the invention are of widespread occurrence and can be obtained from plants, and/or animals and/or microorganisms.

Although metal-binding peptides may be obtained from any of these sources for use in the invention, those obtained from plants or microorganisms are preferred.

For example EP-A 0,242,799 discloses a method for extracting metallothioneins from plant material and metallothioneins produced according to this method may advantageously be used in producing compositions according to the invention.

It will be appreciated that the use of a product of vegetable or microbial origin belonging to the metallothionein group therefore provides an effective "molecular" method of intercepting heavy metals. Using the method of the invention it is possible to bind the metals with a thiolic bond to a polypeptide structure and the bound metals are then prevented from diffusing into the body. This prevents the metals from exerting their toxic effects (e.g. enzymatic inhibition and immunological reactions).

The plants most suitable for use in producing metallothioneins are those, for example, belonging to the Cruciferae and Caryophyllaceae families.

Metallothionein can have the ability specifically to sequester selected metals and it is especially preferred, in accordance with the invention, to utilise metallothioneins which are adapted to sequester

toxic metals, in preference to metals which serve a useful purpose in metabolism, such as copper or iron.

Thus it is especially preferred to use a metallothionein which has a relatively high capability to sequester chromium, nickel, lead and/or cadmium, but a relatively low ability to sequester copper or iron.

It is possible to prepare metallothioneins that are adapted to sequester predetermined cation species by cultivating plants such as *Eruca sativa*, *Brassica napus*, etc. under glass or in open fields in areas of low contamination by the metals. Growth may then continue in the presence of appropriate salts of the selected heavy metal acting as inducer. Once the metallothioneins formed in response to the applied metal have been isolated, they may be treated to remove the heavy metal, thereby freeing the active centres to re-sequester the same metal when they come into contact with it again.

When incorporated in appropriate pharmaceutical or cosmetic products, these metallothioneins will block the specific metal, for example one involved in the aetiopathogenesis of a specific metal-associated toxic manifestation.

The extraction of metallothioneins from vegetable material is described in EP-A-0 242 799 and US-373.400 and these metallothioneins can be used according to the invention. Useful metallothioneins can be obtained from vegetable matter, yeasts or microorganisms that contain them by means of extraction of vegetable or microbial material homogenates with water.

The homogenates may then be centrifuged to separate the cell residues, and the supernatant, after partial dilution with C_{1-3} aliphatic alcohols or with C_{3-6} aliphatic ketones, may be filtered to eliminate undesired glycoprotein products of high molecular weight and then concentrated to a reduced volume.

The crude metallothioneins (often still containing bound metal) portion may be separated from the aqueous extract after being rendered insoluble by adding C_{1-3} aliphatic alcohols or C_{3-6} aliphatic ketones.

The precipitate can be resolubilized in water and reprecipitated in the same solvents at a predetermined ratio to water of between 40 and 90% so as to enable products with a low molecular weight such as simple sugars, amino acids and inorganic salts to be eliminated completely. The metallothioneins, whose molecular weight can vary between 500 and 8,000 depending on their origin, can then be purified from the precipitate by conventional methods known in the literature for example using Sephadex G-50 columns or by means of ultrafiltration with a suitable cut-off membrane.

After being isolated, the metallothioneins prior to their intended cosmetic or therapeutic use, may be treated with acid resins in an atmosphere of inert gases and in the presence of antioxidants to remove the cation.

In this form they may then be incorporated into preparations for application to the skin in formulations such as aqueous gels, cleansing milks, or simple emulsions. It will be understood that it is preferred that only excipients and surfactants that do not interfere with the sulfhydryl groups of the polypeptide may be used.

As an example of cosmetic treatment in accordance with the invention, the formulations can be applied to the exposed parts of the body such as the face, neck, legs, etc. This is preferably carried out in the mornings when one is going into areas with a high urban traffic density or staying in areas of high pollution.

At the end of the day, normal washing will remove from the skin the residue of the formulation that has retained the heavy metals over the course of the day, preventing them from being absorbed through the

skin. Suitable formulations can be applied to the hands or other parts of the body after prolonged use of detergents in order to cleanse the lipid layer and at the same time pick up any traces of contaminating metals.

The following examples illustrate suitable formulations for the compositions of the invention :

Example I:

Formulation of a gel containing Pb, Cr, Cd and Ni sequestering metallothioneins.

The formulation has the following percentage composition:

°	Metallothioneins	1%
	Beeswax	10%
4.	Cyclosilicone pentamer	53%
	Vaseline	30%
	Stearyl ammonium hectorite	2%
	Hydrogenated castor oil	2%
	Pyrogenic silica	1%
	Perfume	1%

Example II: Emulsion for hands and body.

The formulation has the following percentage composition:

Metallothioneins	0.5%
Polytrimethylsiloxysilicate/dimethicone	3 %
Mineral oil	0.5%
Isopropyl lanolate	3 %
Stearic acid	1 %
Cetyl alcohol	1 %
TEA	1.2%
Perfume	0.2%
Water	89.6%

Example III: body milk A/O

The formulation has the following percentage composition:

Metallothioneins	1 %
Cetyl dimethicone copolyol	5 %
Tetraglyceryl stearate hexyl laurate	3 %
Stearyl dimethicone	6 %
Isopropyl myristate	6 %
Mineral oil	4 %
Triglycerides C8-10	3 %
Glycerine	5 %
Vaseline	3 %
NaCl	2 %
Perfume	0.5 %
Water	61.5 %



CLAIMS

1. A topical cosmetic and/or pharmaceutical composition for the external protection of human or animal tissues from contact with heavy metals, said composition comprising a metal sequestering component capable of binding metal ions and a physiologically inert carrier suitable for topical administration, characterised in that said metal sequestering component comprises one or more metal binding peptide having a high proportion of cysteine residues.

2. A composition according to Claim 1, wherein said metal-binding peptide has between about 20 and 50% cysteine residues based on the total number of residues.

3. A composition according to Claim 1, wherein said metal-binding peptide has between about 25 and 45% cysteine residues based on the total number of residues.

4. A composition according to any preceding claim wherein said metal-binding peptide comprises ligands capable of forming metal-thiolate clusters with heavy metal ions.

5. A composition according to any preceding claim wherein said metal-binding peptide is a metallothionein.

6. A composition according to any preceding claim substantially free of heavy metal ions.

7. A composition according to Claim 6 in which said metal-binding peptide is an apometallothionein.

8. A composition according to any preceding claim wherein said metal-binding peptide is obtained from plants.

9. A composition according to any preceding claim in the form of a gel, cream, ointment, or body lotion.





10. A composition according to any of Claims 1 to 9 in the form of a make-up foundation.

11. A composition according to Claims 9 and 10 wherein said physiologically inert carrier suitable for topical administration includes a water resistant, film-forming substance.

12. A composition according to Claim 11, wherein said physiologically inert carrier includes a wax, an oil or an emulsion of a wax or oil.

13. A composition according to any of Claims 9 to 12 further comprising a dermatologically inert colouring agent.

14. A composition according to any preceding claim comprising between about 0.01 and 10 percent by weight of said metal-binding peptide.

15. A composition according to Claim 14 comprising between about 0.1 and 5 percent by weight said metal-binding peptide.

16. A cosmetic method for protection against heavy metals which comprises applying a composition comprising a metal sequestering component capable of binding metal ions and a physiologically inert carrier suitable for topical administration, characterised in that said metal sequestering component comprises one or more metal-binding peptide having a high proportion of cysteine residues.

17. A method according to Claim 16 wherein the composition is applied as a coating to the skin.

18. A method according to Claims 16 and 17 wherein the composition is applied prior to exposure to heavy metals, and subsequently washed off.

19. A method according to any of Claims 16 to 18 wherein said composition is a composition as defined in any of Claims 1 to 15.





20. A mask for the protection of human epithelial tissues from the hazardous effects of airborne heavy metal ion contact, said mask comprising a filter medium, carrying a composition comprising a metal sequestering component capable of binding metal ions, characterised in that said metal sequestering component comprises one or more metal binding peptide having a high proportion of cysteine residues.

21. A disposable filter cartridge for use in a mask according to claim 20, carrying a composition comprising a metal binding peptide according to any of Claims 2 to 8.

This Sheet Has Been Replaced



COSMETIC AND/OR PHARMACEUTICAL COMPOSITIONS
AND METHODS FOR THEIR USE

A B S T R A C T

Topical cosmetic and pharmaceutical compositions are provided for the external protection of human or animal tissues from contact with heavy metals, and said composition comprising a metal sequestering component capable of binding metal ions and a physiologically inert carrier suitable for topical administration. The said metal sequestering component comprises one or more metal binding peptide having a high proportion of cysteine residues, for example a metallothionein.

This Sheet Has Been Replaced



REPLACEMENT SHEETS SG

- ☒ Description : paged 23.87 to paged
☒ Claim(s) : paged .1. to paged .15.
☐ Drawing(s) : paged to paged
☒ Abstract : paged .1. to paged

- 2 -

DATE OF RECEIPT 11/3/91

A further mechanism that has been shown to be responsible for the toxic effects of metals is the competitive substitution of natural physiological enzymic cofactors by heavy metals when at non-physiological concentrations.

Thus, the control of heavy metal pollutants in the atmosphere is essential if metal-related diseases are to be prevented. It is therefore expedient to seek products that efficiently and selectively block heavy metal ions rendering them incapable of entering the body and thereby prevent them exerting their toxic effects.

The extensive contamination of the environment by heavy metals and their ubiquitous presence in the ecosystem, mean that the skin and the accessible mucous membranes form the largest surface area available for heavy metals to accumulate on and subsequently be absorbed into the body. It is also known that many of the cutaneous allergic manifestations that have until now been attributed to detergents or other causes have now been shown to involve heavy metals in their aetiopathogenesis.

Of special note are copper, cobalt, zinc, manganese, mercury and nickel which present significant health hazards in the mining industry, and lead, from exhaust fumes. It has been postulated that the accumulation of lead in developing children may exert a deleterious effect on memory and powers of concentration.

US-A- 4 725 670 discloses the extraction from plant material of metallothioneins and their use in the form of pharmaceutical preparations for treating acute and chronic heavy metal poisoning, and metal deficiency phenomena. There is no mention however that metallothioneins may be used as protective agents against the effects of external, topical heavy metal ion contact.

It has now been surprisingly found that certain polypeptides having a capacity to sequester heavy metals can be used to manufacture cosmetic and/or pharmaceutical compositions which enable the aforementioned problems to be overcome.

Thus according to the present invention there is provided a cosmetic and/or pharmaceutical composition comprising a metallothionein characterised in that said composition is in the form of a gel, cream, ointment, or body lotion and is for the external protection of human or animal tissues from contact with heavy metals and further comprises a physiologically inert carrier suitable for topical administration.

It is preferred that the metallothionein has between 10 and 50% cysteine residues, more preferably between 25 and 45%, based on the total number of residues.

A sub-class of metallothioneins is the so-called phytochelatins. However many authorities now use the term "metallothionein" to cover both.

In more detail, metallothioneins (Mts) are a group of proteins, first discovered in equine kidney in studies on cadmium accumulation by Margoshes and Vallee in 1957. The apoprotein (devoid of bound metal ions) may be capable of binding more than one type of metal. Since then, Mts have been found to bind cadmium, zinc, copper, mercury, silver, gold, lead and bismuth, amongst others.

Polypeptides having the functional characteristics of metallothioneins as described above may be used in accordance with the invention. These may be characterised by having specific sequences within the peptide chain such as Cys-Cys, Cys-X-Cys, and Cys-X-Y-Cys wherein X and Y are residues other than Cys.

Thus the metallothionein used in accordance with the invention may be a metallothionein of Class I, and/or II, and/or III as defined in accordance with the International Convention on nomenclature.

Preferably, the composition according to the invention is substantially free of heavy metal ions, in order that the metal sequestering facility is not impaired. Thus, preferably the metallothionein(s) present is/are apometallothionein(s).

It is preferred that the compositions of the invention are formulated in the form of gels, creams, ointments, or body lotions. Additionally, it is envisaged that the compositions of the invention are formulated to be applied to exposed parts of the body in the form of a barrier cream or cosmetic make-up foundation which may preferably comprise a dermatologically inert colouring agent and/or perfume.

It is most preferred that the compositions are formulated in the form of a film-forming, water-resistant mixture which may comprise for example oils, waxes, silicone oils or other similar inert hydrophobic carrier materials. It is of course desirable that such materials do not interact with sulfhydryl groups of the metallothionein. The compositions are preferably water resistant and preferably are capable of remaining on the skin throughout the normal activities of the day, whilst being capable of being removed simply by washing with a detergent such as soap.

The compositions of the invention preferably contain between 0.01 and 10 percent by weight of said metallothionein, and most preferably between 0.1 and 5 percent by weight of said metallothionein.

The present invention further provides a cosmetic method for protection of human or animal tissues from contact with heavy metals, said method comprising applying to said tissues a composition comprising a metallothionein capable of binding metal ions and a physiologically inert carrier suitable for topical administration.

Although metallothioneins may be used in accordance with the invention, to obtain a protective effect on human or animal tissues against contact with heavy metal ions primarily by incorporating them into cosmetic and/or pharmaceutical compositions, the protective effect of the invention may be obtained in other ways.

Thus in a further embodiment of the present invention, there is provided a mask for the protection of human tissues from the hazardous effects of environmental, particularly airborne heavy metal ion contamination, said mask comprising a filter medium carrying an absorbent material comprising a metallothionein capable of binding metal ions.

Metallothioneins can, according to the invention, also be employed in disposable cartridges for masks for the protection of human epithelial tissues from the hazardous effects of airborne heavy metal ion contact.

The metallothionein used according to the invention may be capable of binding many of the heavy metal ions which may be present in the environment. However it is envisaged that it will be most important to protect against lead, cadmium, mercury, copper and nickel.

Metallothioneins useful in producing the compositions, methods, masks and disposable cartridges of the invention are of widespread occurrence and can be obtained from plants, and/or animals and/or microorganisms.

Although metallothioneins may be obtained from any of these sources for use in the invention, those obtained from plants or microorganisms are preferred.

For example EP-A O,242,799 discloses a method for extracting metallothioneins from plant material and metallothioneins produced according to this method may advantageously be used in producing compositions according to the invention.

It will be appreciated that the use of a product of vegetable or microbial origin belonging to the metallothionein group therefore provides an effective "molecular" method of intercepting heavy metals. Using the method of the invention it is possible to bind the metals with a thiolic bond to a polypeptide structure and the bound metals are then prevented from diffusing into the body. This prevents the metals from exerting their toxic effects (e.g. enzymatic inhibition and immunological reactions).

The plants most suitable for use in producing metallothioneins are those, for example, belonging to the Cruciferae and Caryophyllaceae families.

Metallothionein can have the ability specifically to sequester selected metals and it is especially preferred, in accordance with the invention, to utilise metallothioneins which are adapted to sequester

Claims

1. A cosmetic and/or pharmaceutical composition comprising a metallothionein characterised in that said composition is in the form of a gel, cream, ointment, or body lotion and is for the external protection of human or animal tissues from contact with heavy metals and further comprises a physiologically inert carrier suitable for topical administration.
2. A composition according to Claim 1 in which said metallothionein is an apometallothionein.
3. A composition according to Claim 1 or 2 wherein said metallothionein is obtained from plants.
4. A composition according to any of claims 1 to 3 in the form of a make-up foundation.
5. A composition according to any preceding claim wherein said physiologically inert carrier suitable for topical administration includes a water resistant, film-forming substance.
6. A composition according to Claim 5, wherein said physiologically inert carrier includes a wax, an oil, an emulsion of a wax or an emulsion of an oil.
7. A composition according to any preceding claim further comprising a dermatologically inert colouring agent.
8. A composition according to any preceding claim comprising between 0.01 and 10 percent by weight of said metallothionein.



9. A composition according to Claim 8 comprising between 0.1 and 5 percent by weight of said metallothionein.
10. A cosmetic method for external protection of human or animal tissues against heavy metals which comprises applying to said tissues a composition according to any of Claims 1-9.
11. A method according to Claim 10 wherein the composition is applied as a coating to the skin.
12. A method according to Claims 10 and 11 wherein the composition is applied prior to exposure to heavy metals, and subsequently washed off.
13. A mask for the protection of human epithelial tissues from the hazardous effects of airborne heavy metal ion contact, said mask comprising a filter, characterised in that said filter comprises a metallothionein.
14. A disposable filter cartridge, for use in a mask according to Claim 13, comprising a composition according to any of Claims 1 to 3.



COSMETIC AND/OR PHARMACEUTICAL COMPOSITIONS
AND METHODS FOR THEIR USE

A B S T R A C T

Topical cosmetic and pharmaceutical compositions are provided for the external protection of human or animal tissues from contact with heavy metals, and said composition comprising a metal sequestering component capable of binding metal ions and a physiologically inert carrier suitable for topical administration. The said metal sequestering component comprises one or more metal binding peptide having a high proportion of cysteine residues, for example a metallothionein.