

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(43) International publication date

31 January 2002 (31.01.2002)

PCT

(10) International publication number

WO 02/08192 A1

(51) International patent classification⁷:

A61K 31/47, A61P 35/00

C07D 215/50,

88690 Uhldingen-Mühlhofen (DE). NICKEL, Bernd;
Alleestrasse 35, 64367 Mühlthal (DE). KUTSCHER,
Bernhard; Stresemannstrasse 9, 63477 Maintal (DE).

(21) International application number:

PCT/EP01/08261

(22) International filing date:

18 July 2001 (18.07.2001)

(25) Language of filing:

German

(26) Language of publication:

German

(30) Data relating to the priority:

100 35 928.6

21 July 2000 (21.07.2000)

DE

(71) Applicant: ZENTARIS AG [DE/DE]; Weismüllerstrasse 45,
60314 Frankfurt (DE).(72) Inventors: EMIG, Peter; Ludwig-Erhard-Strasse 22, 63486
Bruchköbel (DE). GÜNTHER, Eckhard; Wingertstrasse 176,
63477 Maintal (DE). SCHMIDT, Jürgen; Am Roggersberg 20,(81) Designated states (*national*): AU, BG, BR, BY, CN,
CO, CZ, EE, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR,
KZ, LT, LV, MK, MX, NO, NZ, PL, RO, RU, SG, SI,
SK, TR, UA, UZ, YU, ZA.(84) Designated states (*regional*): Eurasian Patent (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), European Patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR).

Published:


- With the International Search Report

For an explanation of the two-letter codes and the other
abbreviations, reference is made to the explanations
("Guidance Notes on Codes and Abbreviations") at the
beginning of each regular edition of the PCT Gazette.

As printed

(54) Title: NOVEL HETEROARYL DERIVATIVES AND THE USE THEREOF AS PHARMACEUTICALS

(54) Bezeichnung: NEUE HETEROARYL-DERIVATE UND DEREN VERWENDUNG ALS ARZNEIMITTEL



(I)


(57) Abstract: The invention relates to novel heteroaryl derivatives of general formula (I), the production thereof and the use of the same as pharmaceuticals, especially for treating tumours.

(57) Zusammenfassung: Die Erfindung betrifft neue Heteroaryl-Derivate der allgemeinen Formel (I), deren Herstellung und Verwendung als Arzneimittel, insbesondere zur Behandlung von Tumoren.

Novel heteroaryl derivatives and the use thereof as
pharmaceuticals

The invention relates to novel heteroaryl derivatives
5 of the formula 1, to their preparation and to their use
as medicaments, in particular for treating tumors.

According to one aspect of the invention, there is provided
quinoline derivatives according to the formula 1




formula 1

in which

15 R, R₁, R₂, R₃ can be attached to any of the quinoline
carbon atoms C₂ to C₈, are identical or
different and independently of one another
denote hydrogen, straight-chain or branched
20 (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, straight-
chain or branched (C₁-C₈)-alkylcarbonyl,
straight-chain or branched
(C₁-C₈)-alkoxy, halogen, aryl-(C₁-C₈)-alkoxy,
25 nitro, amino, mono-(C₁-C₄)-alkylamino, di-
(C₁-C₄)-alkylamino, (C₁-C₈)-alkoxycarbonyl-
amino, (C₁-C₆)-alkoxycarbonylamino-(C₁-C₈)-
alkyl, cyano, straight-chain or branched
cyano-(C₁-C₆)-alkyl, carboxyl, (C₁-C₈)-alkoxy-
30 carbonyl, (C₁-C₄)-alkyl which is substituted
by one or more fluorine atoms,
carboxy-(C₁-C₈)-alkyl
or (C₁-C₈)-alkoxycarbonyl-(C₁-C₆)-alkyl, (C₂-

C_6)-alkenyl, (C_2-C_6) -
 alkynyl,
 straight-chain or branched cyano- (C_1-C_6) -
 alkyl, aryl, where
 5 the aryl radical may be unsubstituted or
 mono- or polysubstituted by identical or
 different substituents from the group
 consisting of halogen, straight-chain or
 branched (C_1-C_8) -alkyl, (C_3-C_7) -cycloalkyl,
 10 carboxyl, straight-chain or branched (C_1-C_8) -
 alkoxy, carbonyl,
 by trifluoromethyl, hydroxyl,
 straight-chain or branched (C_1-C_8) -alkoxy,
 benzyloxy,
 15 nitro, amino, mono- (C_1-C_4) -alkylamino, di- $(C_1-$
 $C_4)$ -alkylamino, cyano, straight-chain or
 branched cyano- (C_1-C_6) -alkyl, where
 additionally R and R_1 or R_2 and R_3 may form a
 fused aromatic 6-membered ring with the
 20 quinoline ring forming an acridine ring which
 for its part may be substituted at any C atom
 ring position by the radicals R, R_1 , R_2 and R_3
 having the meanings mentioned above;

25 Z is oxygen or sulfur, where the radical



30 substituted on the quinoline heterocycle may
 be attached to C atoms C_2-C_8 of the quinoline
 ring skeleton;

P, Q independently of one another represent oxygen or
 geminal hydrogen atoms;

X is nitrogen or C-R₅, where R₅ represents hydrogen or (C₁-C₆)-alkyl;

n, m independently of one another denotes an integer between 0-3, with the proviso that in the case n = 0, X denotes a CR₅R₆ group where R₅ and R₆ independently of one another represent hydrogen or (C₁-C₆)-alkyl and that the nitrogen atom adjacent to the C=Z group is substituted by a hydrogen atom or a (C-C₆)-alkyl group;

R₄ is a straight-chain or branched (C₁-C₂₀)-alkyl radical which may be saturated or unsaturated, with one to three double and/or triple bonds, and which may be unsubstituted or may optionally be substituted at the same or different C atoms by one, two or more aryl, heteroaryl, halogen, cyano, (C₁-C₆)-alkoxycarbonylamino, (C₁-C₆)-alkoxy, amino, mono-(C₁-C₄)-alkylamino or di-(C₁-C₄)-alkylamino; a (C₆-C₁₄)-aryl radical, (C₆-C₁₄)-aryl-(C₁-C₄)-alkyl radical or a (C₂-C₁₀)-heteroaryl or (C₂-C₁₀)-heteroaryl-(C₁-C₄)-alkyl radical which contains one or more heteroatoms selected from the group consisting of N, O and S, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and where the (C₆-C₁₄)-aryl or (C₂-C₁₀)-heteroaryl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, halogen, cyano, (C₁-C₆)-alkoxycarbonylamino, (C₁-C₆)-alkoxy, carboxyl, (C₁-C₈)-alkoxycarbonyl, straight-chain or branched (C₁-C₆)-alkyl which is substituted by one or more fluorine atoms, hydroxyl, straight-chain or branched (C₁-C₈)-

alkoxy, where
adjacent oxygen atoms may also be linked by (C₁-C₂)-alkylene groups,

benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, aryl, which for its part may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, carboxyl, straight-chain or branched (C₁-C₈)-alkoxycarbonyl, by trifluoromethyl, hydroxyl, straight-chain or branched (C₁-C₈)-alkoxy, benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, cyano, straight-chain or branched cyano-(C₁-C₆)-alkyl;

and their structural isomers and stereoisomers, in particular tautomers, diastereomers and enantiomers, and their pharmaceutically acceptable salts, in particular acid addition salts.

Thus, for example, the compounds of the formula (1) according to the invention which have one or more centers of chirality and which are present as racemates can be separated by methods known per se into their optical isomers, i.e. enantiomers or diastereomers. The separation can be carried out by column separation on chiral phases or by recrystallization from an optically active solvent or using an optically active acid or base or by derivatization with an optically active reagent, such as, for example, an optically active alcohol, and subsequent removal of the radical.

Furthermore, the quinoline derivatives of the formula (1) according to the invention can be converted into their salts with inorganic or organic acids, in particular, for pharmaceutical use, into their

physiologically acceptable salts. Acids which are suitable for this purpose are, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, fumaric acid, succinic acid, lactic acid, citric acid, acetic acid, tartaric acid, malic acid, embonic acid, malonic acid, trifluoroacetic acid or maleic acid.

Moreover, the compounds of the formula (1) according to the invention can, if they contain a sufficiently acidic group, such as a carboxyl group, be converted, if desired, into their salts with inorganic or organic bases, in particular, for pharmaceutical use, into their physiologically acceptable salts. Bases which are suitable for this purpose are, for example, sodium hydroxide, potassium hydroxide, calcium hydroxide, lysine, cyclohexylamine, ethanolamine, diethanolamine and triethanolamine.

According to a preferred embodiment, quinoline derivatives of the formula 1 are provided in which R, R₁, R₂, R₃, X, Z, P, Q, n and m have the meanings given above and

R₄ denotes a straight-chain or branched (C₁-C₂₀)-alkyl radical which may be saturated or unsaturated, with one to three double and/or triple bonds, and which may be unsubstituted or optionally substituted on the same or different C atoms by one, two or more aryl, heteroaryl, halogen, (C₁-C₆)-alkoxy, amino, mono-(C₁-C₄)-alkylamino or di-(C₁-C₄)-alkylamino;

a phenyl ring or a naphthyl ring, each of which may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, halogen, carboxyl, (C₁-C₈)-alkoxycarbonyl, straight-chain or

branched (C₁-C₆)-alkyl which is substituted by one or more fluorine atoms, preferably trifluoromethyl, hydroxyl, straight-chain or branched (C₁-C₈)-alkoxy, preferably methoxy or ethoxy, where adjacent oxygen atoms may also be linked by (C₁-C₂)-alkylene groups, preferably a methylene group, benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, aryl, which for its part may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, carboxyl, straight-chain or branched (C₁-C₈)-alkoxycarbonyl, by trifluoromethyl, hydroxyl, straight-chain or branched (C₁-C₈)-alkoxy, preferably methoxy or ethoxy, benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, cyano, straight-chain or branched cyano-(C₁-C₆)-alkyl;

a 2-, 4-, 5- or 6-pyrimidinyl radical or 2-, 4-, 5- or 6-pyrimidinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or

different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-, 5- or 6-pyrimidinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 3-, 4-, 5- or 6-pyridazinyl radical or 3-, 4-, 5- or 6-pyridazinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or

5 different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 3-, 4-, 5- or 6-pyridazinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group
10 consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono-
15 or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 5- or 6-pyrazinyl radical or 2-, 3-, 5- or 6-pyrazinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different

substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 3-, 5- or
25 6-pyrazinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino,
30 hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

35

a 3-, 4-, 5-, 6-, 7-, or 8-cinnolinyl radical or 3-, 4-, 5-, 6-, 7-, or 8-cinnolinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by

identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 3-, 4-, 5-, 6-, 7-, or 8-cinnolinyl radical may be unsubstituted or mono- to pentasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, 5-, 6-, 7-, or 8-quinazolinyl radical or 2-, 4-, 5-, 6-, 7-, or 8-quinazolinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by

identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen and oxo (=O) and the or [sic] 2-, 4-, 5-, 6-, 7-, or 8-quinazolinyl radical may be unsubstituted or mono- to pentasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 5-, 6-, 7-, or 8-quinoxaliny radical 2-, 3-, 5-, 6-, 7-, or 8-quinoxaliny-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-,

3-, 5-, 6-, 7-, or 8-quinoxaliny radical may be unsubstituted or mono- to pentasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 4-, 5-, 6-, 7-, or 8-phthalaziny radical or 1-, 4-, 5-, 6-, 7-, or 8-phthalaziny-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 4-, 5-, 6-, 7-, or 8-phthalaziny radical may be unsubstituted or mono- to pentasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 4-, 5-, 6-, 7- or 8-quinoly radical or 2-, 3-, 4-, 5-, 6-, 7 or 8-quinoly-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 3-, 4-, 5-, 6-, 7- or 8-quinoly radical may be unsubstituted or mono- to hexasubstituted by identical or different

substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, preferably methyl, particularly preferably 2-methyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl radical or 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl radical may be unsubstituted or mono- to hexasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 6-, 8- or 9-[9H]-purinyl radical or 2-, 6-, 8- or 9-[9H]-purinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 6-, 8- or 9-[9H]-purinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino,

mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 6-, 7- or 8-[7H]-purinyl radical or 2-, 6-, 7- or 8-[7H]-purinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 6-, 7- or 8-[7H]-purinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl radical or 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl radical may be unsubstituted or mono- to octasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono-

or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

5 a 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-phenanthridinyl radical or 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-phenanthridinyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by
 10 identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-phenanthridinyl radical may be unsubstituted or mono- to octasubstituted by identical or different
 15 substituents from the group consisting of (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, (C₆-C₁₀)-aryl-(C₁-C₆)-alkoxy, preferably benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by
 20 fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

25 a 2-, 3-, 4-, 5- or 6-pyridyl radical where the 2-, 3-, 4-, 5- or 6-pyridyl radical may be unsubstituted or mono- to tetrasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen,
 30 nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;
 35

a 2-, 3-, 4-, 5- or 6-pyridinyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be

unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 3-, 4-, 5- or 6-pyridinyl radical may be unsubstituted or mono- to tetrasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 4- or 5-thienyl radical or 2-, 3-, 4- or 5-thienyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 3-, 4- or 5-thienyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, or 5-thiazolyl radical or 2-, 4-, or 5-thiazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-, or 5-thiazolyl radical may be unsubstituted or mono- or

disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 3-, 4-, or 5-isothiazolyl radical or 3-, 4-, or 5-isothiazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 3-, 4-, or 5-isothiazolyl radical may be unsubstituted or mono- or disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, 5-, 6-, or 7-benzothiazolyl radical or 2-, 4-, 5-, 6-, or 7-benzothiazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-, 5-, 6-, or 7-benzothiazolyl radical may be unsubstituted or mono- to tetrasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino,

hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 2-, 4-, or 5-imidazolyl radical or 1-, 2-, 4-, or 5-imidazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 4-, or 5-imidazolyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 3-, 4-, or 5-pyrazolyl radical or 1-, 3-, 4- or 5-pyrazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 3-, 4- or 5-pyrazolyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

5 a 1-, 2-, 3-, 4-, or 5-pyrrolyl radical or 1-, 2-,
3-, 4-, or 5-pyrrolyl-(C₁-C₆)-alkyl radical, where
the (C₁-C₆)-alkyl radical may be unsubstituted or
mono- or polysubstituted by identical or different
substituents from the group consisting of (C₁-C₆)-
alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4-
or 5-pyrrolyl radical may be unsubstituted or
mono- to tetrasubstituted by identical or
10 different substituents from the group consisting
of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino,
mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino,
hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-
C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or
15 (C₁-C₆)-alkyl which is mono- or polysubstituted by
fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl
and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

20 a 1-, 3-, or 5-[1.2.4]-triazolyl radical or 1-,
3-, or 5-[1.2.4]-triazolyl-(C₁-C₆)-alkyl radical,
where the (C₁-C₆)-alkyl radical may be
unsubstituted or mono- or polysubstituted by
identical or different substituents from the group
consisting of hydrogen, (C₁-C₆)-alkyl, halogen and
25 oxo (=O) and the 1-, 3-, or 5-[1.2.4]-triazolyl
radical may be unsubstituted or mono- or
disubstituted by identical or different
substituents from the group consisting of (C₁-C₆)-
alkyl, halogen, nitro, amino, mono-(C₁-C₆)-
30 alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-
C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-
alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-
C₆)-alkyl which is mono- or polysubstituted by
fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl
35 and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 4-, or 5-[1.2.3]-triazolyl radical or 1-,
4-, or 5-[1.2.3]-triazolyl-(C₁-C₆)-alkyl radical,
where the (C₁-C₆)-alkyl radical may be

- unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 4-, or 5-[1.2.3]-triazolyl radical may be unsubstituted or mono- or disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;
- a 1- or 5-[1H]-tetrazolyl radical or 1-, or 5-[1H]-tetrazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, or 5-[1H]-tetrazolyl radical may be unsubstituted or substituted by hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;
- a 2- or 5-[2H]-tetrazoyl radical or 2- or 5-[2H]-tetrazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2- or 5-[2H]-tetrazolyl radical may be unsubstituted or substituted by hydrogen, (C₁-C₆)-alkyl, halogen,

nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono-

5 or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, or 6-[1.3.5]-triazinyl radical or 2-,
10 4-, or 6-[1.3.5]-triazinyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen and
15 oxo (=O) and the 2-, 4-, or 6-[1.3.5]-triazinyl radical may be unsubstituted or mono- or disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino,
20 mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-
25 aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, or 5-oxazolyl radical or 2-, 4-, or 5-oxazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono-
30 or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-, or 5-oxazolyl radical may be unsubstituted or mono- or disubstituted by identical or different
35 substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino

or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

5 a 3-, 4-, or 5-isoxazolyl radical or 3-, 4-, or 5-isoxazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-
10 alkyl, halogen and oxo (=O) and the 3-, 4-, or 5-isoxazolyl radical may be unsubstituted or mono- or disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl radical or 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be
25 unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl radical may be unsubstituted or mono- to
30 hexasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl and the isomers, in particular tautomers, diastereomers and

enantiomers, and the pharmaceutically acceptable salts, in particular acid addition salts, thereof.

According to a further embodiment, quinoline derivatives of the formula (1) are provided which are characterized in that R, R₁, R₂, R₃, X, Z, P, Q, n and m have the meanings given above and R₄ represents phenyl which is unsubstituted or substituted by one to five identical or different (C₁-C₆)-alkoxy groups, where adjacent oxygen atoms may also be linked by (C₁-C₂)-alkylene groups.

According to a further embodiment, quinoline derivatives of the formula (1) are provided which are characterized in that R, R₁, R₂, R₃, X, Z, P, Q, n and m have the meanings given above and R₄ represents 3,5-dimethoxyphenyl.


According to a further embodiment, quinoline derivatives of the formula (1) are provided which are characterized in that R₄ has the meanings given above, R, R₁, R₂, R₃ each represent a hydrogen atom, Z represents an oxygen atom and X represents a nitrogen atom, P and Q each represent geminal hydrogen atoms, m is zero and n is the integer 2.

According to a further embodiment, quinoline derivatives of the formula (1) are provided which are characterized in that R, R₁, R₂, R₃ each represent a hydrogen atom, Z represents an oxygen atom, X represents a nitrogen atom, P and Q each represent geminal hydrogen atoms, m is zero, n represents the integer 2 and R₄ represents a 3,5-dimethoxyphenyl radical.

According to yet another aspect of the invention, there is provided a use of quinoline derivatives as defined above in the manufacture of a medicament for treating tumors in mammals.


According to a further aspect of the invention, a process for preparing quinoline derivatives as defined above characterized in that a quinoline carboxylic acid of the formula (2)

21



formula 2

- 5 in which R, R₁, R₂, R₃ have the meanings given above, Z denotes an oxygen or sulfur atom and Y represents a leaving group such as halogen, hydroxyl, (C₁-C₆)-alkoxy, -O-tosyl, -O-mesyl or imidazolyl, is reacted with an amine of the
 10 formula (3)



formula 3


- 15 in which R₄, X, P, Q, m and n have the meanings given above, and
 the desired quinoline derivatives are formed.

Synthesis route:

20

The compounds of the formula 1 can be obtained according to Scheme 1 below:

Scheme 1



5 The starting materials (2) and (3) are either commercially available or can be prepared by procedures known per se. The starting materials (2) and (3) are useful intermediates for preparing the quinoline derivatives of the formula (1) according to the
10 invention.

The solvents and auxiliaries to be used, if appropriate, and the reaction parameters to be used, such as reaction temperature and reaction time, are
15 known to the person skilled in the art owing to his expert knowledge.

The quinoline derivatives of the formula (1) according to the invention are suitable as medicaments, in particular as antitumor agents, for treating mammals,
20 in particular man, but also domestic animals such as horses, cattle, dogs, cats, hares, sheep, poultry and the like.

According to a further aspect, the invention provides quinoline derivatives as described above for use in therapy. According to another aspect, the invention provides the use of quinoline derivatives as described above for the manufacture of a medicament for treating tumors in mammals, in particular man.

Preferably, at least one quinoline derivative of the formula (1) is administered to a mammal in an amount effective for the treatment of the tumor. The therapeutically effective dose of the quinoline derivative according to the invention in question which is to be administered for the treatment depends inter alia on the nature and the stage of the oncosis, the age and the sex of the patient, the type of administration and the duration of the treatment. Administration can take place orally, rectally, buccally (for example sublingually), parenterally (for example subcutaneously, intramuscularly, intradermally or intravenously), topically or transdermally.

According to a further aspect of the invention, medicament

characterized in that it comprises, as active ingredient, at least one quinoline derivative as defined above.

These can be solid, semisolid, liquid or aerosol preparations. Suitable solid preparations are, for example, capsules, powders, granules, tablets. Suitable semisolid preparations are, for example, ointments, creams, gels, pastes, suspensions, oil-in-water and water-in-oil emulsions. Suitable liquid preparations are, for example, sterile aqueous preparations for parenteral administration which are isotonic with the blood of the patient.

The invention is to be illustrated in more detail by the example below, without being restricted to the example.

Working Example**1-(3,5-dimethoxyphenyl)-4-(4-quinolyl-carbonyl)piperazine**

5 2 g (11.5 mmol) of quinoline-4-carboxylic acid were suspended in 80 ml of DMF. With stirring, 1.74 g (17.2 mmol) of N-methylmorpholine and then a solution of 8.95 g (17.2 mmol) of Py-BOP (1-benzotriazolyl-tripyrrolidinophosphonium hexafluorophosphate) and 10 2.56 g (11.5 mmol) of 1-(3,5-dimethoxyphenyl)piperazine in 25 ml of DMF were added to this mixture. The mixture was stirred at RT for 12 h, the DMF was distilled off under reduced pressure and the residue was purified on 15 a silica gel column (Kieselgel 60, from Merck AG, Darmstadt) using the mobile phase dichloromethane/methanol/25 percent ammonia (90:10:1 v/v/v).

Yield: 3.4 g (78.3% of theory)
20 m.p.: 146-148°C

1. Antiproliferative action in various tumor cell lines

25 In a proliferation test, the antiproliferative activity of the substance D-43411 was examined using established tumor cell lines. In the test used, the cellular dehydrogenation activity is determined, which makes it possible to determine the vitality of the cell and, 30 indirectly, the cell count. The cell lines used are the human cervical carcinoma cell lines KB/HeLa (ATCC/CCL17), the murine lymphocyte leukaemia L1210 (ATCC CCL-219), the human breast adenocarcinoma line MCF7 /ATCC HTB22) and the ovary adenocarcinoma line 35 SKOV-3 (ATCC HTB77). These are established cell lines which are very well characterized and were obtained from ATCC and cultured.

The results shown in Tab. 1 demonstrate the highly potent antiproliferative action of D-43411 in the cell lines SKOV-3, L-1210 and HeLa/KB. Owing to the particularly slow growth of the MCF7 line, the effect of D-43411 in the test period of 48 h is only small (18% inhibition at 3.16 $\mu\text{g/ml}$; thus stated as >3.16).

Tab. 1 In-vitro cytotoxicity in tumor cell lines
(values determined from 5 substance concentrations)

D number	Structure	XTT - Assay IC ⁵⁰ [$\mu\text{g/ml}$]				
		MW	SKOV-3	L1210	KB/HeLa	MCF7
D-43411		429	<0.0003	<0.0003	<0.0003	>3.16

2. Method

15 XTT Test for cellular dehydrogenase activity

The adherently growing tumor cell lines HeLa/KB, SKOV-3 and MCF7 and the L1210 leukaemia line, which grows in suspension, were cultivated under standard conditions in an incubator with gas inlet at 37°C, 5% CO₂ and 95% atmospheric humidity. On Test Day 1, the adherent cells are detached using trypsin/EDTA and pelleted by centrifugation. The cell pellet is then resuspended in RPMI culture medium at the appropriate cell count and transferred to a 96-well microtitre plate. The plates are then cultivated overnight in the incubator with gas inlet. The test substances are made up as stock solutions in DMSO and, on Test Day 2, diluted with culture medium to the desired concentrations. The

substances in the culture medium are then added to the cells and incubated in the incubator with gas inlet for 45 h. Cells which have not been treated with test substance serve as control.

5

For the XTT assay, 1 mg/ml of XTT (sodium 3'-[1-(phenylaminocarbonyl)-3,4-tetrazolium]-bis(4-methoxy-6-nitro)benzenesulfonic acid) is dissolved in RPMI-1640 medium without Phenol Red. Additionally, a 0.383 mg/ml solution of PMS (N-methyldibenzopyrazine methyl sulfate) in phosphate-buffered saline (PBS) is prepared. On Test Day 4, 75 µl/well of the XTT-PMS mixture are pipetted onto the cell plates, which by now have been incubated with the test substances for 45 h. To this end, the XTT solution is mixed with the PMS solution in a ratio of 50:1 (v/v) shortly before use. The cell plates are then incubated in the incubator with gas inlet for a further 3 h, and the optical density (OD_{490nm}) is determined in a photometer.

20

Using the OD_{490nm} obtained, the inhibition in percent relative to the control is calculated. The antiproliferative activity is estimated using regression analysis.

25

Example I

Tablet containing 50 mg of active compound

Composition:

	(1) Active compound	50.0 mg
30	(2) Lactose	98.0 mg
	(3) Maize starch	50.0 mg
	(4) Polyvinylpyrrolidone	15.0 mg
	(5) Magnesium stearate	2.0 mg
	Total:	215.0 mg

35

Preparation:

(1), (2) and (3) are mixed and granulated with an aqueous solution of (4). The dried granules are admixed with (5). This mixture is tabletted.

Example II

Capsule containing 50 mg of active compound

Composition:

5	(1) Active compound	50.0 mg
	(2) Maize starch, dried	58.0 mg
	(3) Lactose powder	50.0 mg
	(4) Magnesium stearate	2.0 mg
	Total:	160.0 mg


10

Preparation:

(1) is ground with (3). This ground material is added with vigorous mixing to the mixture of (2) and (4). This powder mixture is, on a capsule filling machine,
15 filled into hard gelatine capsules size 3.

Patent Claims

1. Quinoline derivatives according to the formula 1



formula 1

in which

- 10 R, R₁, R₂, R₃ can be attached to any of the quinoline carbon atoms C₂ to C₈, are identical or different and independently of one another denote hydrogen, straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, straight-chain or branched (C₁-C₈)-alkylcarbonyl, straight-chain or branched (C₁-C₈)-alkoxy, halogen, aryl-(C₁-C₈)-alkoxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, (C₁-C₈)-alkoxycarbonylamino, (C₁-C₆)-alkoxycarbonylamino-(C₁-C₈)-alkyl, cyano, straight-chain or branched cyano-(C₁-C₆)-alkyl, carboxyl, (C₁-C₈)-alkoxy-carbonyl, (C₁-C₄)-alkyl which is substituted by one or more fluorine atoms, carboxy-(C₁-C₈)-alkyl or (C₁-C₈)-alkoxycarbonyl-(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, straight-chain or branched cyano-(C₁-C₆)-alkyl, aryl, where the aryl radical may be unsubstituted or mono- or polysubstituted by identical or


different substituents from the group consisting of halogen, straight-chain or branched (C_1-C_8) -alkyl, (C_3-C_7) -cycloalkyl, carboxyl, straight-chain or branched (C_1-C_8) -alkoxycarbonyl,

by trifluoromethyl, hydroxyl, straight-chain or branched (C_1-C_8) -alkoxy, benzyloxy,

nitro, amino, mono- (C_1-C_4) -alkylamino, di- (C_1-C_4) -alkylamino, cyano, straight-chain or branched cyano- (C_1-C_6) -alkyl, where

additionally R and R_1 or R_2 and R_3 may form a fused aromatic 6-membered ring with the quinoline ring forming an acridine ring which for its part may be substituted at any C atom ring position by the radicals R, R_1 , R_2 and R_3 having the meanings mentioned above;

Z is oxygen or sulfur, where the radical



substituted on the quinoline heterocycle may be attached to C atoms C_2-C_8 of the quinoline ring skeleton;

P, Q independently of one another represent oxygen or geminal hydrogen atoms;

X is nitrogen or $C-R_5$, where R_5 represents hydrogen or (C_1-C_6) -alkyl;

n, m independently of one another denotes an integer between 0-3, with the proviso that in the case $n = 0$, X denotes a CR_5R_6 group where R_5 and R_6 independently of one another represent hydrogen or (C_1-C_6) -alkyl and that the nitrogen atom adjacent

to the C=Z group is substituted by a hydrogen atom or a (C-C₆)-alkyl group;

R₄ is a straight-chain or branched (C₁ - C₂₀)-

5 alkyl radical which may be saturated or unsaturated, with one to three double and/or triple bonds, and which may be unsubstituted or may optionally be substituted at the same or different C atoms by one, two or more aryl, 10 heteroaryl, halogen, cyano, (C₁-C₆)-alkoxy-carbonylamino, (C₁-C₆)-alkoxy, amino, mono-(C₁-C₄)-alkylamino or di-(C₁-C₄)-alkylamino; a (C₆-C₁₄)-aryl radical, (C₆-C₁₄)-aryl-(C₁-C₄)-alkyl radical or a (C₂-C₁₀)-heteroaryl or (C₂-C₁₀)-heteroaryl-(C₁-C₄)- 15 alkyl radical which contains one or more heteroatoms selected from the group consisting of N, O and S, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by

identical or different substituents from the group 20 consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and where the (C₆-C₁₄)-aryl or (C₂-C₁₀)-heteroaryl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of 25 straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, halogen, cyano, (C₁-C₆)-alkoxycarbonylamino, (C₁-C₆)-alkoxy, carboxyl, (C₁-C₈)-alkoxycarbonyl, straight-chain or branched (C₁-C₆)-alkyl which is substituted by one or more 30 fluorine atoms,

hydroxyl, straight-chain or branched (C₁-C₈)-alkoxy, where adjacent oxygen atoms may also be linked by (C₁-C₂)-alkylene groups,

35 benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, aryl, which for its part may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of

straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, carboxyl, straight-chain or branched (C₁-C₈)-alkoxycarbonyl, by trifluoromethyl, hydroxyl, straight-chain or branched (C₁-C₈)-alkoxy, benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, cyano, straight-chain or branched cyano-(C₁-C₆)-alkyl;

and their structural isomers and stereoisomers, in particular tautomers, diastereomers and enantiomers, and their pharmaceutically acceptable salts, in particular acid addition salts.

2. Quinoline derivatives of the formula 1 according to Claim 1, characterized in that R, R₁, R₂, R₃, X, Z, P, Q, n and m have the meanings given in Claim 1 and

R₄ denotes a straight-chain or branched (C₁-C₂₀)-alkyl radical which may be saturated or unsaturated, with one to three double and/or triple bonds, and which may be unsubstituted or optionally substituted on the same or different C atoms by one, two or more aryl, heteroaryl, halogen, (C₁-C₆)-alkoxy, amino, mono-(C₁-C₄)-alkylamino or di-(C₁-C₄)-alkylamino;

a phenyl ring or a naphthyl ring, each of which may be unsubstituted or mono- or poly-substituted by identical or different substituents from the group consisting of straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, halogen, cyano, (C₁-C₆)-alkoxycarbonylamino, (C₁-C₆)-alkoxy, carboxyl, (C₁-C₈)-alkoxycarbonyl, straight-chain or branched (C₁-C₆)-alkyl which is substituted by one or more fluorine atoms, hydroxyl, straight-chain or

branched (C₁-C₈)-alkoxy,

benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, aryl, which for its part may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched (C₁-C₈)-alkyl, (C₃-C₇)-cycloalkyl, carboxyl, straight-chain or branched (C₁-C₈)-alkoxycarbonyl, by trifluoromethyl, hydroxyl, straight-chain or branched (C₁-C₈)-alkoxy,

benzyloxy, nitro, amino, mono-(C₁-C₄)-alkylamino, di-(C₁-C₄)-alkylamino, cyano, straight-chain or branched cyano-(C₁-C₆)-alkyl;

a 2-, 4-, 5- or 6-pyrimidinyl radical or 2-, 4-, 5- or 6-pyrimidinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by

by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-, 5- or 6-pyrimidinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine,

(C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 3-, 4-, 5- or 6-pyridazinyl radical or 3-, 4-, 5- or 6-pyridazinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the

group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 3-, 4-, 5- or 6-pyridazinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 5- or 6-pyrazinyl radical or 2-, 3-, 5- or 6-pyrazinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- [lacu identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 3-, 5- or 6-pyrazinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 3-, 4-, 5-, 6-, 7-, or 8-cinnolinyl radical or 3-, 4-, 5-, 6-, 7-, or 8-cinnolinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-

C₆)-alkyl, halogen and oxo (=O) and the 3-, 4-,
 5-, 6-, 7-, or 8-cinnolinyl radical may be
 unsubstituted or mono- to pentasubstituted by
 identical or different substituents from the
 group consisting of hydrogen, (C₁-C₆)-alkyl,
 halogen, nitro, amino, mono-(C₁-C₆)-alkylamino,
 di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy,
 benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl,
 (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl
 which is mono- or polysubstituted by fluorine,
 (C₆-C₁₀)-aryl and
 (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, 5-, 6-, 7-, or 8-quinazolinyl radical
 or 2-, 4-, 5-, 6-, 7-, or 8-quinazolinyl-(C₁-
 C₄)-alkyl radical, where the (C₁-C₄)-alkyl
 radical may be unsubstituted or mono- or
 polysubstituted by identical or different
 substituents from the group consisting of
 hydrogen, (C₁-C₆)-alkyl, halogen and oxo (=O)
 and the 2-, 4-, 5-, 6-, 7-, or
 8-quinazolinyl radical may be unsubstituted or
 mono- to pentasubstituted by identical or
 different substituents from the group
 consisting of hydrogen, (C₁-C₆)-alkyl, halogen,
 nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-
 C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy,
 benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl,
 (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl
 which is mono- or polysubstituted by fluorine,
 (C₆-C₁₀)-aryl and
 (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 5-, 6-, 7-, or 8-quinoxaliny radical
 2-, 3-, 5-, 6-, 7-, or 8-quinoxaliny-
 (C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl
 radical may be unsubstituted or mono- or
 polysubstituted by identical or different
 substituents from the group consisting of

(C₁-C₆)-alkyl, halogen and oxo (=O) and the
 2-, 3-, 5-, 6-, 7-, or 8-quinoxaliny
 radical may be unsubstituted or mono- to
 pentasubstituted by identical or different
 substituents from the group consisting of
 hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino,
 mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino,
 hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl,
 (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonyl-
 amino or (C₁-C₆)-alkyl which is mono- or
 polysubstituted by fluorine,
 (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-
 (C₁-C₆)-alkyl;

a 1-, 4-, 5-, 6-, 7-, or 8-phthalaziny
 radical or 1-, 4-, 5-, 6-, 7-, or 8-phthalaziny
 (C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl
 radical may be unsubstituted or mono- or
 polysubstituted by identical or different
 substituents from the group consisting of (C₁-
 C₆)-alkyl, halogen and oxo (=O) and the
 1-, 4-, 5-, 6-, 7-, or 8-phthalaziny
 radical may be unsubstituted or mono- to
 pentasubstituted by identical or different
 substituents from the group consisting of
 hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino,
 mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino,
 hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl,
 (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-
 alkoxycarbonylamino or (C₁-C₆)-alkyl which is
 mono- or polysubstituted by fluorine,
 (C₆-C₁₀)-aryl and
 (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 4-, 5-, 6-, 7- or 8-quinoly
 radical or 2-, 3-, 4-, 5-, 6-, 7 or 8-quinoly
 (C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical
 may be unsubstituted or mono- or poly-
 substituted by identical or different

substituents from the group consisting of
(C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-,
3-, 4-, 5-, 6-, 7- or 8-quinolyl radical may be
unsubstituted or mono- to hexasubstituted by
identical or different substituents from the
group consisting of hydrogen, (C₁-C₆)-alkyl,

halogen, nitro, amino, mono-(C₁-C₆)-
alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl,
(C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-
alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or
(C₁-C₆)-alkyl which is mono- or polysubstituted
by fluorine, (C₆-
C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl
radical or 1-, 3-, 4-, 5-, 6-, 7- or 8-
isoquinolyl-(C₁-C₄)-alkyl radical, where the
(C₁-C₄)-alkyl radical may be unsubstituted or
mono- or polysubstituted by identical or
different substituents from the group
consisting of (C₁-C₆)-alkyl, halogen and oxo
(=O) and the 1-, 3-, 4-, 5-, 6-, 7- or 8-
isoquinolyl radical may be unsubstituted or
mono- to hexasubstituted by identical or
different substituents from the group
consisting of hydrogen, (C₁-C₆)-alkyl, halogen,
nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-
C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy,
benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl,
(C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl
which is mono- or polysubstituted by fluorine,
(C₆-C₁₀)-aryl and
(C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 6-, 8- or 9-[9H]-purinyl radical or 2-,
6-, 8- or 9-[9H]-purinyl-(C₁-C₄)-alkyl radical,
where the (C₁-C₄)-alkyl radical may be

unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 6-, 8- or 9-[9H]-purinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine,

(C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 6-, 7- or 8-[7H]-purinyl radical or 2-, 6-, 7- or 8-[7H]-purinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₄)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 6-, 7- or 8-[7H]-purinyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine,

(C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl radical or 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl-(C₁-C₄)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or

mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl radical may be unsubstituted or mono- to octasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-phenanthridinyl radical or 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-phenanthridinyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-phenanthridinyl radical may be unsubstituted or mono- to octasubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, (C₆-C₁₀)-aryl-(C₁-C₆)-alkoxy, preferably benzyl-oxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 4-, 5- or 6-pyridyl radical where the 2-, 3-, 4-, 5- or 6-pyridyl radical may be

unsubstituted or mono- to tetrasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 4-, 5- or 6-pyridinyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 3-, 4-, 5- or 6-pyridinyl radical may be unsubstituted or mono- to tetrasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 3-, 4- or 5-thienyl radical or 2-, 3-, 4- or 5-thienyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 3-, 4- or 5-thienyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl,

halogen, nitro, amino, mono-(C₁-C₆)-alkylamino,
di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy,
benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl,
(C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl
which is mono- or polysubstituted by fluorine,
(C₆-C₁₀)-aryl and
(C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, or 5-thiazolyl radical or 2-, 4-, or
5-thiazolyl-(C₁-C₆)-alkyl radical, where the
(C₁-C₆)-alkyl radical may be unsubstituted or
mono- or polysubstituted by identical or
different substituents from the group
consisting of (C₁-C₆)-alkyl, halogen and oxo
(=O) and the 2-, 4-, or 5-thiazolyl radical may
be unsubstituted or mono- or disubstituted by
identical or different substituents from the
group consisting of hydrogen, (C₁-C₆)-alkyl,
halogen, nitro, amino, mono-(C₁-C₆)-alkylamino,
di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy,
benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl,
(C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl
which is mono- or polysubstituted by fluorine,
(C₆-C₁₀)-aryl and
(C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 3-, 4-, or 5-isothiazolyl radical or 3-, 4-,
or 5-isothiazolyl-(C₁-C₆)-alkyl radical, where
the (C₁-C₆)-alkyl radical may be unsubstituted
or mono- or polysubstituted by identical or
different substituents from the group
consisting of (C₁-C₆)-alkyl, halogen and oxo
(=O) and the 3-, 4-, or 5-isothiazolyl radical
may be unsubstituted or mono- or disubstituted
by identical or different substituents from the
group consisting of hydrogen, (C₁-C₆)-alkyl,
halogen, nitro, amino, mono-(C₁-C₆)-alkylamino,
di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy,
benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl,

(C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl
which is mono- or polysubstituted by fluorine,
(C₆-C₁₀)-aryl and
(C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

5

10

15

20

a 2-, 4-, 5-, 6-, or 7-benzothiazolyl radical
or 2-, 4-, 5-, 6-, or 7-benzothiazolyl-(C₁-C₆)-
alkyl radical, where the (C₁-C₆)-alkyl radical
may be unsubstituted or mono- or
polysubstituted by identical or different
substituents from the group consisting of (C₁-
C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-,
5-, 6-, or 7-benzothiazolyl radical may be
unsubstituted or mono- to tetrasubstituted by
identical or different substituents from the
group consisting of hydrogen, (C₁-C₆)-alkyl,
halogen, nitro, amino, mono-(C₁-C₆)-alkylamino,
di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy,
benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl,
(C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl
which is mono- or polysubstituted by fluorine,
(C₆-C₁₀)-aryl and
(C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

25

30

35

a 1-, 2-, 4-, or 5-imidazolyl radical or 1-,
2-, 4-, or 5-imidazolyl-(C₁-C₆)-alkyl radical,
where the (C₁-C₆)-alkyl radical may be
unsubstituted or mono- or polysubstituted by
identical or different substituents from the
group consisting of (C₁-C₆)-alkyl, halogen and
oxo (=O) and the 1-, 2-, 4-, or 5-imidazolyl
radical may be unsubstituted or mono- to
trisubstituted by identical or different
substituents from the group consisting of
hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino,
mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino,
hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl,
(C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonyl-
amino or (C₁-C₆)-alkyl which is mono- or poly-

substituted by fluorine, preferably trifluoromethyl, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

5 a 1-, 3-, 4-, or 5-pyrazolyl radical or 1-, 3-, 4- or 5-pyrazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or
10 different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 3-, 4- or 5-pyrazolyl radical may be unsubstituted or mono- to trisubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl,
15 halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine,
20 (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1-, 2-, 3-, 4-, or 5-pyrrolyl radical or 1-, 2-, 3-, 4-, or 5-pyrrolyl-(C₁-C₆)-alkyl radical,
25 where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4- or 5-pyrrolyl radical may be unsubstituted or mono- to
30 tetrasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonyl-
35 amino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine,

(C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-
(C₁-C₆)-alkyl;

5 a 1-, 3-, or 5-[1.2.4]-triazolyl radical or 1-,
3-, or 5-[1.2.4]-triazolyl-(C₁-C₆)-alkyl
radical, where the (C₁-C₆)-alkyl radical may be
unsubstituted or mono- or polysubstituted by
identical or different substituents from the
10 group consisting of hydrogen, (C₁-C₆)-alkyl,
halogen and oxo (=O) and the 1-, 3-, or 5-
[1.2.4]-triazolyl radical may be unsubstituted
or mono- or disubstituted by identical or
different substituents from the group
15 consisting of (C₁-C₆)-alkyl, halogen, nitro,
amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-
alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy,
carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-
alkoxycarbonylamino or (C₁-C₆)-alkyl which is
mono- or polysubstituted by fluorine,
20 (C₆-C₁₀)-aryl and
(C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

25 a 1-, 4-, or 5-[1.2.3]-triazolyl radical or 1-,
4-, or 5-[1.2.3]-triazolyl-(C₁-C₆)-alkyl
radical, where the (C₁-C₆)-alkyl radical may be
unsubstituted or mono- or polysubstituted by
identical or different substituents from the
group consisting of (C₁-C₆)-alkyl, halogen and
30 oxo (=O) and the 1-, 4-, or 5-[1.2.3]-triazolyl
radical may be unsubstituted or mono- or
disubstituted by identical or different
substituents from the group consisting of
hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino,
mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino,
35 hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl,
(C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonyl-
amino or (C₁-C₆)-alkyl which is mono- or
polysubstituted by fluorine,

(C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 1- or 5-[1H]-tetrazolyl radical or 1-, or 5-[1H]-tetrazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, or 5-[1H]-tetrazolyl radical may be unsubstituted or substituted by hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2- or 5-[2H]-tetrazoyl radical or 2- or 5-[2H]-tetrazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2- or 5-[2H]-tetrazolyl radical may be unsubstituted or substituted by hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, or 6-[1.3.5]-triazinyl radical or 2-, 4-, or 6-[1.3.5]-triazinyl-(C₁-C₆)-alkyl

radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-, or 6-[1.3.5]-triazinyl radical may be unsubstituted or mono- or disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 2-, 4-, or 5-oxazolyl radical or 2-, 4-, or 5-oxazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 2-, 4-, or 5-oxazolyl radical may be unsubstituted or mono- or disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;

a 3-, 4-, or 5-isoxazolyl radical or 3-, 4-, or 5-isoxazolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or

different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 3-, 4-, or 5-isoxazolyl radical may be unsubstituted or mono- or disubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl;


a 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl radical or 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl-(C₁-C₆)-alkyl radical, where the (C₁-C₆)-alkyl radical may be unsubstituted or mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, halogen and oxo (=O) and the 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl radical may be unsubstituted or mono- to hexasubstituted by identical or different substituents from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, nitro, amino, mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, hydroxyl, (C₁-C₆)-alkoxy, benzyloxy, carboxyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkoxycarbonylamino or (C₁-C₆)-alkyl which is mono- or polysubstituted by fluorine, (C₆-C₁₀)-aryl and (C₆-C₁₀)-aryl-(C₁-C₆)-alkyl.

3. Quinoline derivatives according to Claim 1 or 2, characterized in that R, R₁, R₂, R₃, X, Z, P, Q, n and m have the meanings given above and R₄ represents phenyl which is unsubstituted or substituted by one to five identical or different

(C₁-C₆)-alkoxy groups, where adjacent oxygen atoms may also be linked by (C₁-C₂)-alkylene groups.

4. Quinoline derivatives according to any of Claims 1 to 3, characterized in that R, R₁, R₂, R₃, X, Z, P, Q, n and m have the meanings given above and R₄ represents 3,5-dimethoxyphenyl.
5. Quinoline derivatives according to any of the preceding claims, characterized in that R₄ has the meanings given above, R, R₁, R₂, R₃ each represent a hydrogen atom, Z represents an oxygen atom and X represents a nitrogen atom, P and Q each represent geminal hydrogen atoms, m is zero, n represents the integer 2.
6. Quinoline derivative according to any of Claims 1 to 5, characterized in that R, R₁, R₂, R₃ each represent a hydrogen atom, Z represents an oxygen atom, X represents a nitrogen atom, P and Q each represent geminal hydrogen atoms, m is zero, n represents the integer 2, and R₄ represents a 3,5-dimethoxyphenyl radical.
7. Quinoline derivatives according to any of Claims 1 to 6 for use in therapy.
8. Use of the quinoline derivatives according to any of Claims 1 to 6 for the manufacture of a medicament for treating tumors in mammals.
9. Process for preparing quinoline derivatives according to any of Claims 1 to 6, characterized in that a quinoline carboxylic acid of the formula (2)


48



formula 2

in which R, R₁, R₂, R₃ have the meanings given above, Z denotes an oxygen or sulfur atom and Y represents a leaving group such as halogen, hydroxyl, (C₁-C₆)-alkoxy,

-O-tosyl, -O-mesyl or imidazolyl, is reacted with an amine of the formula (3)



formula 3

in which R₄, X, P, Q, m and n have the meanings given above,

and the desired quinoline derivatives are formed.

10. Medicament, characterized in that it comprises, as active ingredient, at least one quinoline derivative according to any of Claims 1 to 6,