



(51) International Patent Classification:

C12Q 1/37 (2006.01) *G01N* 33/48 (2006.01)
C12Q 1/56 (2006.01)

(21) International Application Number:

PCT/IT2012/000193

(22) International Filing Date:

22 June 2012 (22.06.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant (for all designated States except US): **GENTI-UM S.P.A.** [IT/IT]; Piazza XX Settembre, 2, I-22079 Villa Guardia (CO) (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **IGNONI, Terenzio** [IT/IT]; Via Verdi, 15, I-22070 Solbiate (CO) (IT). **KUMAR, Vijay** [GB/IT]; Via Galilei, 4, I-22070 Casnate (CO) (IT). **ISLAM, Khalid** [GB/IT]; Via San Gottardo, 80, CH-6900 Massagno (CH).

(74) Agents: **PISTOLESI, Roberto** et al.; c/o Dragotti & Associati S.R.L., Via Nino Bixio, 7, I-20129 Milano (IT).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: EUGLOBULIN-BASED METHOD FOR DETERMINING THE BIOLOGICAL ACTIVITY OF DEFIBROTIDE

Figure 1



(57) Abstract: It is disclosed a method for determining the biological activity of defibrinolytic, which comprises the steps of: a) bringing into contact defibrinolytic, mammalian euglobulin and a substrate specific for the plasmin which, by reaction with the plasmin, provides a measurable product; and b) measuring the amount of product formed at successive times, to thereby determine the biological activity of the defibrinolytic. Liquid defibrinolytic formulations are also disclosed, preferably water solutions, having a defined biological activity and, in particular, having an activity of 25 to 35 IU/mg of defibrinolytic, preferably from 27.5 to 32.5 IU/mg and, more preferably, from 28 to 32 IU/mg.