HYBRID HEPATOCYTE GROWTH FACTOR GENE HAVING HIGH EXPRESSION EFFICIENCY OF TWO HETEROTYPES OF HEPATOCYTE GROWTH FACTOR

Field of the Invention

The present invention relates to a highly efficient hybrid Hepatocyte Growth Factor (HGF) gene which simultaneously expresses two heterotypes of HGF.

Background of the Invention

The present invention relates to a hybrid HGF gene prepared by inserting an inherent or foreign intron between exons 4 and 5 in HGF cDNA, which has higher expression efficiency than HGF cDNA and simultaneously expresses two heterotypes of HGF and dHGF (deleted variant HGF).

HGF is a heparin binding glycoprotein called a scatter factor. A gene encoding HGF is located at chromosome 721.1 and comprises 18 exons and 17 introns, having the nucleotide sequence of SEQ ID NO: 1 (Seki T et al., *Gene* 102:213-219, 1991). A transcript of about 6 kb is transcribed from the HGF gene, and then, a polypeptide HGF precursor consisting of 728 amino acids is synthesized therefrom. Simultaneously, a polypeptide of dHGF precursor consisting of 723 amino acids is also synthesized by an alternative splicing of the HGF gene. The biologically inactive precursors may be converted into active forms of disulfide-linked heterodimer by protease in serum. In the heterodimers, the alpha chain having a high molecular weight forms four kringle domains and an N-terminal hairpin loop like a preactivated peptide region of plasminogen. The kringle domains of a triple disulfide-bonded loop structure consisting of about 80 amino acids may play an important role in protein-protein interaction. The low molecular weight beta chain forms an inactive serine protease-like domain. dHGF consisting 723 amino acids is a polypeptide with deletion of five amino acids in the 1st kringle domain of the alpha chain, i.e., F, L, P, S and S.

It has been recently reported that both of HGF and dHGF have several biological functions, e.g., promoting the growth and morphogenesis of epithelial cell, melanocyte and endothelial cell. However, they are different in terms of immunological or biological properties.

For example, HGF shows about 20-fold, 10-fold and 2-fold higher activities than dHGF in promoting DNA synthesis in human umbilical cord venous endothelial cell, arterial smooth muscle cell and NSF-60 (murine myeloblast cell), respectively. dHGF shows about 3-fold and 2-fold higher activities than HGF in promoting DNA synthesis of LLC-PK1(pig kidney epithelial cell), and OK(American opossum kidney epithelial cell) and mouse interstitial cell, respectively. HGF has a 70-fold higher solubility in PBS than dHGF. Several anti-dHGF monoclonal antibodies recognize only dHGF, but not HGF or a reduced form of dHGF, which implies structures of HGF and dHGF are different. Accordingly, the simultaneous synthesis of HGF and dHGF *in vivo* suggests that they biologically interact with each other (Shima, N. et al., *Biochemical and Biophysical Research Communications* 200:808-815, 1994).

HGF secreted from mesoderm-derived cells has various biological functions, e.g., 1) inducing epithelial cells into a tubular structure; 2) stimulating vascularization from endothelial cells *in vitro* and *in vivo*; 3) regeneration of liver and kidney, owing to its anti-apoptosis activity; 4) organogenesis of kidney, ovary and testis; 5) controlling osteogenesis; 6) stimulating the growth and differentiation of erythroid hematopoietic precursor cells; and 7) axon sprouting of neurons (Stella, M. C. and Comoglio, P. M., *The International Journal of Biochemistry & Cell Biology* 31:1357-1362, 1999). Based on these various functions, HGF or a gene encoding HGF may be developed as a therapeutic agent for treating ischemic or liver diseases. Actually, *in vivo*, the HGF may exist as either HGF or dHGF, and therefore, the coexpression of HGF and dHGF is important for maximizing the therapeutic effect. Accordingly, the present inventors have endeavored to develop a hybrid HGF gene which can simultaneously express HGF and dHGF with a high efficiency for gene therapy.

Summary of the Invention

Accordingly, it is a primary object of the present invention to provide a hybrid HGF gene which simultaneously expresses two heterotypes of HGF.

In accordance with one aspect of the present invention, there is provide the hybrid HGF gene having an inherent or foreign intron is inserted between exons 4 and 5 of HGF cDNA.

It is a another object of the present invention to provide a recombinant

vector comprising the hybrid HGF gene and a microorganism transformed with the above vector.

It is a still further object of the present invention to provide a pharmaceutical composition for treating or preventing ischemic or liver dieases, which comprises the HGF gene.

Brief Description of the Drawings

The above and other objects and features of the present invention will become apparent from the following description of the invention, when taken in conjunction with the accompanying drawings which respectively show:

Figure 1 : a schematic diagram of HGF-X prototype illustrating the positions of the gene fragments,

Figure 2 : a process for cloning gene fragments from HepG2 genomic DNA,

Figure 3 : a process for cloning gene fragments from human plcenta cDNA,

Figures 4A and 4B : processes for preparing expression vectors pCK-HGF-X,

Figure 5 : a process for preparing expression vectors pCK-cHGF and pCK-dHGF,

Figure 6 : a process for preparing expression vectors pCP-HGF-X family,

Figure 7 : a process for preparing expression vectors pCP-cHGF and pCP-dHGF,

Figure 8 : gene expression levels of pCP-cHGF, pCP-dHGF and pCP-HGF-X.

Figure 9 : gene expression patterns of pCP-cHGF, pCP-dHGF and pCP-HGF-X observed by electrophoresis on 12% polyacrylamide gel,

Figure 10 : gene expression levels of pCP-cHGF, pCP-dHGF and pCP-HGF-X7, *in vivo*,

Figure 11 : cerebral angiogenesis of two groups of rabbits which were subject to administrating pCP and pCP-HGF-X7, respectively,

Detailed Description of the Invention

The hybrid Hepatocyte Growth Factor (HGF) gene of the present invention comprises cDNA corresponding to the exons 1 to 18, and an inherent or foreign intron inserted between exons 4 and 5 of the cDNA. The intron comprises a fragment of the inherent intro or a recombinant sequence.

An embodiment of the hybrid HGF gene of the present invention comprising the inherent intron is 7112bp long and has the nucleotide sequence of SEQ ID NO: 2. The hybrid HGF gene simultaneously expresses both HGF and dHGF, and has higher expression efficiency than HGF cDNA.

Codon degeneracy enables the hybrid HGF gene of the present invention to be modified or changed in the coding and/or non-coding region without altering the amino acid sequence of the protein and the expression of the gene. Accordingly, polynucleotides which is substantially identical to the hybrid HGF gene of SEQ ID NO:2, and the fragments thereof fall within the scope of the invention. "Substantially identical" means that the sequence homology is not less than 80%, preferably not less than 90%, and more preferably not less than 95%.

A hybrid HGF gene may comprise a fragment of inherent intron optionally having a small recombinant sequence inserted thereinto between exons 4 and 5 of HGF cDNA. Herein, such a hybrid HGF gene comprising a fragment of inherent intron designates "HGF-X". HGFX-6, HGF-X7 and HGF-X8 having the nucleotide sequence of SEQ ID Nos: 19 to 21, respectively, are preferred.

The hybrid HGF gene of the present invention is synthesized and inserted into an expression vector, according to the known genetic engineering methods. Then, the vector can be introduced into an appropriate host cells such as *E.coli* and *yeast*. For example, *Escherichia coli* Top10F' may be transfected with HGF-X7 gene of the present invention. *Escherichia coli* Top10F' pCK-HGFX7 and *Escherichia coli* Top10F' pCP-HGFX7 then obtained were deposited as the accession numbers KCCM-10361 and KCCM-10362, respectively, on March 12, 2002.

By using the transformed cells, the gene of the present invention and the protein encoded thereby may be produced on a large scale.

The vector of the present invention may selectively comprise sequence(s) for regulating gene expression such as promoter or terminator, self-replication sequence and secretory signal, depending on host cells.

Further, the present invention comprises a pharmaceutical compostion for treating or preventing ischemic and liver dieasese, which comprises the hybrid HGF gene or the vector comprising the gene as an active ingredient. Preferably,

the composition is formulated for injection.

The composition of the present invention may further comprise pharmaceutically acceptable carriers. Any of the conventional procedures in the pharmaceutical field may be used to prepare oral formulations such as tablets, capsules, pills, granules, suspensions and solutions; rejection formulations such as solutions, suspensions, or dried powders that may be mixed with distilled water before injection; locally-applicable formulations such as ointments, creams and lotions; and other formulations.

Carriers generally used in the pharmaceutical field may be employed in the composition of the present invention. For example, orally-administered formulations may include binders, emulsifiers, disintegrating agents, excipients, solubilizing agents, dispersing agents, stabilizing agents, suspending agents, coloring agents or spicery. Injection formulations may comprise preservatives, unagonizing agents, solubilizing agents or stabilizing agents. Preparation for local administration may contain bases, excipients, lubricants or preservatives. Any of the suitable formulations known in the art (Remington's Pharmaceutical Science [the new edition], Mack Publishing Company, Eaton PA) may be used in the present invention.

The inventive composition can be clinically administered as various oral and parenteral formulations. A suitable formulation may be prepared using such excipients as additives, enhancers, binders, wetting agents, disintegrating agents and surfactants, or diluents. Solid formulations for oral administration include pills, tablets, dusting powder, granules and capsules. Those solid formulations may be prepared by mixing one or more excipients, e.g. starch, calcium carbonate, sucrose, lactose and gelatin with dibenzylbuthyllacton lignan derivatives. Also, lubricants such as magnesium stearate and talc may be included in the present formulation. Liquid formulations for oral administration include suspension, solution, emulsion and syrup. Those formulations may contain wetting agents, sweeteners, aromatics and preservatives, in addition to general simple diluents such as water and liquid paraffin. Formulations for parenteral administration include sterilized aqueous solution, suspension, emulsion, freeze-dried alternative treatment and suppositories. Water-insoluble excipients and suspending agents comprise vegetable fats such as propylene glycol, polyethylene glycol and olive oil, and injectable esters such as ethyl oleate. Witepsol®, Macrogol®, Tween® 61, cacao fats, laurin fats and glycerogelatins may be used as bases of suppositories.

The inventive composition may be administered orally or via parenteral

routes such as intravenous, intramuscular, subcutaneous, intraabdominal, sternal and arterial injection or infusion, or topically through rectal, intranasal, inhalational or intraocular administration.

It should be understood that the typical daily dose of composition of the present invention ought to be determined in light of various relevant factors including the conditions to be treated, the chosen route of administration, the age, sex and body weight of the individual patient, and the severity of the patient's symptom, and can be administrated in a single dose or in divided dose. Therefore, the daily dose should not be construed as a limitation to the scope of the invention in any way.

The following Examples are given for the purpose of illustration only, and are not intended to limit the scope of the invention.

Example 1: Preparation of hybrid gene constructs encoding human HGF

(1) Cloning of HGF gene fragments obtained from genomic DNA

Human HepG2 cells (ATCC Accession NO: HB-8065) were suspended in TES buffer (10 mM Tris-HCl; 1 mM EDTA; 0.7% SDS) and treated with 400 μg/mℓ of proteinase K at 50℃ for 1 hour. Subsequently, genomic DNA was extracted from the cell suspension by phenol/chloroform extraction and ethanol precipitation according to the conventional method in the art.

In the PCR amplication, the extracted genomic DNA was employed as a template DNA. As primer pairs, the synthetic nuclotides of SEQ ID NOs: 3 and 4 were employed to obtain DNA fragments containing : HGF gene fragment 2 (HGF-F2), SEQ ID NOs: 3 and 5; HGF-F3, SEQ ID NOs: 6 and 7; HGF-F5, SEQ ID NOs: 8 and 7; HGF-F7, SEQ ID NOs: 9 and 7; HGF-F8, SEQ ID NOs: 10 and 7; HGF-F6, respectively (Fig. 1). The PCR amplification mixture was prepared by mixing 1 μℓ of template DNA, 1 μℓ each of primer (10 pmol/μℓ), 10 μℓ of dNTP (10 mM), 3.5 unit of Expand High Fidelity enzyme (Gibco BRL, USA) and 10 μℓ of enzyme buffer solution and adjusted to a final volume of 100 μℓ with distilled water. 30 cycles of the PCR amplification was carried out, each cycle consisting of 1 min at 94℃, 1 min at 55℃ and 30 sec at 72℃. The primers used herein and the amplified gene fragments obtained therefrom are shown in Table 1.

Table 1

5' primer	3' primer	Amplified fragment
gHGF3 (SEQ ID NO: 3)	gHGF4 (SEQ ID NO: 4)	HGF gene fragment 2 (HGF-F2)
gHGF3 (SEQ ID NO: 3)	gHGF10 (SEQ ID NO: 5)	HGF gene fragment 3 (HGF-F3)
gHGF5 (SEQ ID NO: 6)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 5 (HGF-F5)
gHGF12 (SEQ ID NO: 8)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 7 (HGF-F7)
gHGF13 (SEQ ID NO: 9)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 8 (HGF-F8)
gHGF6 (SEQ ID NO: 10)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 6 (HGF-F6)

The amplified HGF-F2 comprised the sequence ranging from 392 to 2247 of human HGF cDNA prototype (HGF-X1; composed of exons 1 to 4-intron 4-exons 5 to 18) of SEQ ID NO: 2; HGF-F3, the sequence ranging from 392 to 727; HGF-5, the sequence ranging from 2229 to 5471; HGF-F6, the sequence ranging from 5117 to 5471; HGF-F7, the sequence ranging from 3167 to 5471; and HGF-F8, the sequence ranging from 4167 to 5471.

The amplified HGF gene fragments were each inserted into the multiple cloning site of pGEM-T easy vector (Promega, WI, USA) to obtain pGEM-T easy-HGF-F2, pGEM-T easy-HGF-F3, pGEM-T easy-HGF-F5, pGEM-T easy-HGF-F6, pGEM-T easy-HGF-F7 and pGEM-T easy-HGF-F8, respectively (Fig. 2). The nucleotide sequences of the amplified HGF gene fragments were confirmed by a sequence analysis.

(2) Cloning of HGF gene fragments obtained from cDNA

In the PCR amplification, human placenta cDNA (Clontech, CA, USA) was employed as a template DNA under the same condition as described in Example 1. As primer pairs, the synthetic oligonucleotides of SEQ ID NOs: 11 and 12, and SEQ ID NOs: 13 and 14 were employed to obtain DNA fragments containing HGF-F1 and HGF-F4, respectively. Further, DNA fragments containing cDNAs of HGF gene (cHGF) and deleted HGF gene (dHGF) were amplified by PCR using synthetic oligonucleotides of SEQ ID NOs: 15 and 16 as a primer pair, respectively. dHGF is a HGF gene with deletion of 5 base sequences.

The primers used herein and the amplified gene fragments obtained therefrom are shown in Table 2.

Table 2

5' primer	3' primer	Amplified product
gHGF1 (SEQ ID NO: 11)	gHGF2 (SEQ ID NO: 12)	HGF gene fragment 1 (HGF-F1)
gHGF8 (SEQ ID NO: 13)	gHGF9 (SEQ ID NO: 14)	HGF gene fragment 4 (HGF-F4)
cHGF5 (SEQ ID NO: 15)	cHGF3 (SEQ ID NO: 16)	HGF gene cDNA (cHGF) dHGF gene cDNA (dHGF)

The amplified HGF-F1 and HGF-F4 comprised the nucleotide sequences ranging from 1 to 402 and from 6533 to 7113 of SEQ ID NO: 2 of human HGF cDNA prototype, respectively. HGF gene cDNA comprised the nucleotide sequence ranging from 1 to 2184 of SEQ ID NO: 1 of human HGF gene, and dHGF gene cDNA has the same sequence as HGF gene cDNA except for the deletion of the sequence ranging from 483 to 495.

The amplified fragments of HGF gene were each inserted into the multiple cloning site of pGEM-T easy vector (Promega, WI, USA) to obtain pGEM-T easy-HGF-F1, pGEM-T easy-HGF-F4, pGEM-T easy-cHGF and pGEM-T easy-dHGF, respectively (Fig. 3). The nucleotide sequences of the human HGF gene fragments, HGF gene cDNA and dHGF gene cDNA were confirmed by sequence analyses.

(3) Preparation of hybrid HGF gene constructs

Hybrid HGF gene constructs of genomic DNA and cDNA were prepared by combining the fragments of HGF gene cloned in steps (1) and (2) as follows (Figs. 4A and 4B).

Plasmid pGEM-T-easy-HGF-F1 was treated with *Hin*dIII/*Bam*HI to obtain HGF-F1. Plasmid pCK (see PCT International Publication NO: WO/0040737) was treated with *Hin*dIII/*Bam*HI, and HGF-F1 was inserted thereinto to obtain pCK-F1. And then, plasmids pGEM-T-easy-HGF-F2 and pGEM-T-easy-HGF-F3 were treated with *Mlu*I/*Bam*HI to obtain HGF-F2 and HGF-F3, respectively. pCK-1 was treated with *Mlu*I/*Bam*HI, and then HGF-F2 and HGF-F3 were inserted thereinto to obtain pCK-F12M and pCKF13M. The *Mlu*I restriction site of vectors pCK-F12M and pCK-F13M was substituted with an *Hga*I restriction site by employing a site-directed mutagenesis kit (Stratagene, CA, USA) to obtain pCK-F12 and pCK-F13, respectively.

Further, plasmid pGEM-T-easy-HGF-F4 was treated with *Bam*HI/*Xba*I

to obtain HGF-F4. pCK-F12 and pCK-F13 were treated with *Bam*HI/*Xba*I, and HGF-F4 was inserted thereinto to obtain pCK-F124 and pCK-F134, respectively. And then, plasmids pGEM-T-easy-HGF-F5, pGEM-T-easy-HGF-F6, pGEM-T-easy-HGF-F7 and pGEM-T-easy-HGF-F8 were treated with *Bam*HI/*Xho*I to obtian HGF-F5, HGF-F6, HGF-F7 and HGF-F8, respectively. pCK-F124 and pCK-F134 were treated with *Bam*HI/*Xho*I, and then HGF-F5, HGF-F6, HGF-F7 and HGF-F8 were inserted thereinto to obtain pCK-F1254 and pCK-F1264, pCK-F1274, pCK-F1284, pCK-F1354, pCK-F1364, pCK-F1374 and pCK-F1384, respectively.

And then, pGEM-T easy-cHGF was treated with *Xho*I to obtain cDNA fragment of HGF gene (HGF-*Xho*I) of about 1100bp. Then, HGF-*Xho*I was inserted into pCK-F1254, pCK-F1264, pCK-F1274, pCK-F1284, pCK-F1354, pCK-F1364, pCK-F1374 and pCK-F1384 to obtain pCK-HGF-X1, pCK-HGF-X2, pCK-HGF-X3, pCK-HGF-X4, pCK-HGF-X5, pCK-HGF-X6, pCK-HGF-X7 and pCK-HGF-X8, respectively. On the other hand, pGEM-T easy-cHGF and pGEM-T easy-dHGF were treated with *Bam*HI to obtain HGF gene cDNA and dHGF gene cDNA. Then, HGF gene cDNA and dHGF gene cDNA were inserted into the *Bam*HI restriction site of pCK to obtain pCK-cHGF and pCK-dHGF, respectively (Fig. 5).

(4) Preparation of an expression vector containing a hybrid HGF gene construct

Plasmid pCDNA3.1 (Invitrogen, USA) was digested with *Nde*I, treated with the Klenow fragment to build blunt ends, and then digested with *Nhe*I to obtain a DNA fragment containing human cytomegalovirus promoter. Plasmids pCK-HGF-X1, pCK-HGF-X2, pCK-HGF-X3, pCK-HGF-X4, pCK-HGF-X5, pCK-HGF-X6, pCK-HGF-X7 and pCK-HGF-X8 were digested with *Sna*BI, treated with the Klenow fragment to make blunt ends and digested with *Nhe*I, and then the above DNA fragment containing human cytomegalovirus promoter was inserted thereinto to obtain pCP-HGF-X1, pCP-HGF-X2, pCP-HGF-X3, pCP-HGF-X4, pCP-HGF-X5, pCP-HGF-X6, pCP-HGF-X7 and pCP-HGF-X8, respectively (Fig. 6).

Plasmid pCDNA3.1 (Invitrogen, USA) was digested with *Nhe*I, treated with the Klenow fragment to make blunt ends and digested with *Nde*I to obtain the DNA fragment containing human cytomegalovirus promoter. pCK-cHGF and pCK-dHGF were digested with *Mlu*I, treated with the Klenow fragment to make blunt ends and digested with *Nde*I, and then the above DNA fragment

containing human cytomegalovirus promoter was inserted thereinto to obtain pCP-cHGF and pCP-dHGF, respectively (Fig. 7).

Example 2: Examination of the expression efficiency of hybrid HGF gene construct and the coexpression of HGF/dHGF

Studies was conducted to examine whether the hybrid HGF gene constructs (HGF-X1 to HGF-X8) obtained in Example 1 can simultaneously express HGF and dHGF and whether there is any difference in the gene expression level between hybrid HGF gene constructs and HGF cDNA.

(1) Gene expression efficiency

First, 5 μg of pCP-HGF-X2, pCP-HGF-X3, pCP-HGF-X6, pCP-HGF-X7 and pCP-HGF-X8 were transfected into 5 × 10^6 cells of 293 cell (ATCC CRL 1573) together with 0.5 μg of DONAI-LacZ (TAKARA SHUZO, Japan) DNA using FuGENE6 (Gibco BRL, MD, USA), according to the manufacturer's instructions. At this time, 5 μg each of pCP-cHGF and pCP-dHGF were used as controls, and DONAI-LacZ DNA was used to calibrate the infection efficiency. 3 hours after transfection, cells were re-fed with a fresh medium and further cultured for 48 hours. The culture solution thus obtained was divided into two parts. One part of the 293 cell culture solution was subjected to RNA extraction, and the other, to measurement of LacZ activity. The LacZ activity was measured using an activity measuring kit (Stratagene, CA, USA) according to the manufacturer's instructions.

In order to compare the gene expression levels, the amount of HGF in the cell culture was measured by an enzyme-linked immunosorbent assay kit (ELISA, R&D System, MN, USA). After calibrating the infection efficiency by the measured LacZ activity, the expression level of HGF-X gene was found to be from 20 to 150-fold higher than those of HGF cDNA and dHGF cDNA (Fig. 8). HGF-X7, in particular, showed the highest gene expression level.

(2) Coexpression of HGF and dHGF

In order to examine coexpression of HGF and dHGF from hybrid HGF gene constructs, total cellular RNAs were extracted from the transfected 293 cells using the Trizol method (Trizol; Gibco BRL, USA) and subjected to RT-

PCR to obtain cDNA. Then, using cDNA as a template DNA, PCR amplification was carried out using synthetic oligonucleotides of SEQ ID NOs: 17 and 18 as a primer pair. The PCR amplification mixture was prepared by mixing 1 ㎕ of the template DNA, 1 ㎕ each of the primer (10 pmol/㎕), 10 ㎕ of dNTP (10 mM), 3.5 unit of Taq polymerase (TAKARA SHUZO, Japan) and 10 ㎕ of enzyme buffer solution and adjusted to a final volume of 100 ㎕ with distilled water. 30 cycles of PCR amplication was conducted, each cycle consisting of 1 min at 94℃, 1 min at 55℃, and 90 sec at 72℃.

The amplified PCR products corresponded to the boundary region between exons 4 and 5 of HGF gene; HGF gene cDNA of 142 bp and dHGF gene cDNA of 127bp, respectively. With no splicing, the PCR product of at least 1 kb in length was amplified; and if alternative splicing occurred, HGF gene cDNA of 142bp and dHGF gene cDNA of 127 bp were simultaneously synthesized and amplified. The amplified PCR products were distinguished by electrophoresis on a 12% polyacrylamide gel.

As shown in Fig. 9, while the bands of 142 bp and 127 bp were detected in the lanes loading HGF gene cDNA and dHGF gene cDNA, respectively, both bands of 142 bp and 127 bp were detected in the lanes loading HGF-X. The above results suggest that HGF and dHGF are simultaneously expressed from hybrid HGF-X gene constructs of the present invention.

Example 3: Comparison of expression levels of HGF-X7, HGF gene cDNA and dHGF gene cDNA (*in vivo*)

100 ㎍ each of pCP-HGF-X7, pCP-cHGF and pCP-dHGF were injected into the enterior tibial muscle of the hind limb of mice with an insulin syringe. After 5 days, the mice were sacrificed and the muscles around the injection spot were removed and smashed in a protein extraction buffer(25 mM Tris-HCl(pH 7.4), 50 mM NaCl, 0.5 % Na-deoxycholate, 2 % NP-40, 0.2 % SDS) to separate total proteins. The amount of the total proteins was measured with a DC protein analysis kit (Bio-Rad Laboratories, CA, USA) and the amount of expressed HGF was determined with an ELISA kit (R&D System) according to the manufacturer's instruction.

As can be seen from the result shown in Fig. 10, the amount of HGF expressed from HGF-X7 is 250-fold higher than that from HGF gene cDNA or

dHGF gene cDNA.

Together with the result of the experiment of Example 2 (*in vivo*), this result demonstrates that the expression efficiency of HGF-X gene is much superior to those of HGF gene cDNA or dHGF gene cDNA.

Example 4: Gene therapy employing HGF-X7 in a rabbit ischemic hind limb model

In order to examine whether HGF-X7 gene is effective in the treatment of ischemic hind limb disease, gene therapy was carried out on a rabbit ischemic hind limb model as follows.

A rabbit ischemic hind limb model, which is a standard animal model for the ischemic limb disease, was prepared by the method described by Takeshita et al., *Journal of Clinical Investigation*, 93: 662(1994). At the day before operation (day 0), each of 30 white rabbits from New Zealand(male, from 3.8 to 4.2 kg) was intramuscularly injected with 5 mg/kg of xylazine and, then, anesthetized by an intramuscular injection of 50 mg/kg of ketamine. Subsequently, the left femoral region of the rabbit was incised and all branches of the femoral artery were separated and tied. The region from the proximal part to the branching point of the saphenous and popliteal arteries was incised to prepare the model. After the operation, 15 mg/kg/day of cefazolin was injected intramuscularly for 5 days and 0.3 mg/day of morphin, for 10 days. 10 days after the operation (day 10), angiography was carried out for the left hind limb where the ischemia was induced, and the degree of arteriogenesis was recorded as a basal level. The rabbits were randomly divided into two groups and injected at four sites in the femoral muscle with 500 ㎍ of plasmid pCP-HGF-X7(experimental group) or 500 ㎍ of plasmid pCP(control), respectively. 40 days after the operation (day 40), angiography was carried out again for the left hind limb and the degree of arteriogenesis at the arteriole level was determined and compared to that of day 10.

As can be seen from the result in Fig. 11, the degree of angiogenesis was significantly enhanced in the experimental group administered with pCP-HGF-X7 as compared with the pCP-administered control group.

This result demonstrates that HGF-X7 gene can be effectively used in the gene therapy of an ischemic disease.

While the invention has been described with respect to the above

specific embodiments, it should be recognized that various modifications and changes may be made to the invention by those skilled in the art which also fall within the scope of the invention as defined by the appended claims.

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To. Viromed Co., Limited
1510-8 BongChen-dong, Kwanak-gu,
Seoul 151-818,
Korea

RECEIPT IN THE CASE OF AN ORIGINAL
Issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : *Escherichia coli* Top10F' pCK-HGFX7	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10361
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: ☐ a scientific description ☐ a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Mar. 12, 2002 (date of the original deposit)[1]	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s) : Date: Mar. 19. 2002.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of I nternational depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

Form BP/4 Sole page

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To. Viromed Co., Limited
1510-8 BongChen-dong, Kwanak-gu,
Seoul 151-818,
Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : *Escherichia coli* Top10F' pCP-HGFX7	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10362
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: ☐ a scientific description ☐ a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Mar. 12. 2002. (date of the original deposit)[1]	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Mar. 19. 2002.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired ; where a deposit made outside the Budapest Treaty after the acquisition of the status of International depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

Form BP/4 Sole page

What is claimed is:

1. A hybrid Hepatocyte Growth Factor (HGF) construct comprising:
an HGF cDNA which comprises HGF exons 1-18, or a degenerate form thereof which do not alter the encoded amino acid sequence; and
HGF intron 4 or a fragment thereof between exons 4 and 5 of the HGF cDNA.

2. The hybrid HGF construct of claim1, wherein the hybrid HGF construct has a nucleotide sequence of SE ID NO:2.

3. The hybrid HGF construct of claim 1, wherein the hybrid HGF construct has a nucleotide sequence not less than 80 % identical to SEQ ID NO:2.

4. The hybrid HGF construct of claim 1, wherein the hybrid HGF construct has a nucleotide sequence not less than 90 % identical to SEQ ID NO:2.

5. The hybrid HGF construct of claim 1, wherein the hybrid HGF construct has a nucleotide sequence not less than 95 % identical to SEQ ID NO:2.

6. The hybrid HGF construct of claim 1, wherein the hybrid HGF construct has a nucleotide sequence selected from the group consisting of SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21.

7. A vector comprising the hybrid construct of claim 1.

8. The vector of claim 7, wherein the vector further comprises one or more sequences for regulating gene expression, a self-replication sequence, or a secretory signal.

9. The vector of claim 8, wherein the vector is selected from the group consisting of: pCK-HGF-X2, pCK-HGF-X3, pCK-HGF- X6, pCK-HGF-X7, pCK-HGF-X8,

pCP-HGF-X2, pCP-HGF-X3, pCP-HGF-X6, pCP-HGF-X7 and pCP-HGF-X8.

10. A cell containing the vector of claim 7.

11. The cell of claim 10, wherein the cell is a yeast cell or an *E. coli* cell.

12. The cell of claim 11, wherein the cell is an *E. coli* Topl0F'pCK-HGFX7 (Deposit No. KCCM-10361) or *E. coli* Topl0F'pCP- HGFX7 (Deposit No. KCCM-10362) cell.

13. A pharmaceutical composition comprising the hybrid HGF construct of claim 1 and a carrier.

14. A use of a hybrid HGF construct of claim 1 for the manufacture of a medicament for preventing or treating an ischemic or liver disease.

15. The use of claim 14, wherein the hybrid HGF construct is in a vector.

16. The use of claim 15, wherein the vector is selected from the group consisting of: pCK-HGF-X2, pCK-HGF-X3, pCK-HGF- X6, pCK-HGF-X7, pCK-HGF-X8, pCP-HGF-X2, pCP-HGF-X3, pCP-HGF-X6, pCP-HGF-X7 and pCP-HGF-X8.

17. The use of claim 14, wherein the hybrid HGF construct is formulated in a pharmaceutical composition suitable for direct injection.

18. A method of simultaneously producing two heterotypes of HGF and dHGF protein comprising the steps of culturing the cell of any one of claims 10 to 12 in a suitable medium and collecting the HGF and dHGF proteins.

SEQUENCE LISTING

<110> VIROMED LIMITED

<120> HYBRID HEPATOCYTE GROWTH FACTOR GENE WHICH HAS A HIGH EXPRESSION EFFICIENCY AND EXPRESSES TWO HETEROTYPES OF HEPATOCYTE GROWTH FACTOR

<130> PCA30320/VML

<160> 21

<170> KopatentIn 1.71

<210> 1
<211> 2187
<212> DNA
<213> Hepatocyte Growth Factor

<400> 1

atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc 60

ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat 120

gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa 180

accaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt 240

ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc 300

ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa 360

aacaaagact acattagaaa ctgcatcatt ggtaaaggac gcagctacaa gggaacagta 420

tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac 480

agctttttgc cttcgagcta tcggggtaaa gacctacagg aaaactactg tcgaaatcct 540

cgaggggaag aaggggacc ctggtgtttc acaagcaatc cagaggtacg ctacgaagtc 600

tgtgacattc ctcagtgttc agaagttgaa tgcatgacct gcaatgggga gagttatcga 660

ggtctcatgg atcatacaga atcaggcaag atttgtcagc gctgggatca tcagacacca 720

caccggcaca aattcttgcc tgaaagatat cccgacaagg gctttgatga taattattgc 780

cgcaatcccg atggccagcc gaggccatgg tgctatactc ttgaccctca cacccgctgg 840

gagtactgtg caattaaaac atgcgctgac aatactatga atgacactga tgttcctttg 900

gaaacaactg aatgcatcca aggtcaagga gaaggctaca ggggcactgt caataccatt 960

tggaatggaa ttccatgtca gcgttgggat tctcagtatc ctcacgagca tgacatgact 1020

cctgaaaatt tcaagtgcaa ggacctacga gaaaattact gccgaaatcc agatgggtct 1080

gaatcaccct ggtgttttac cactgatcca aacatccgag ttggctactg ctcccaaatt 1140

ccaaactgtg atatgtcaca tggacaagat tgttatcgtg ggaatggcaa aaattatatg 1200

ggcaacttat cccaaacaag atctggacta acatgttcaa tgtgggacaa gaacatggaa 1260

gacttacatc gtcatatctt ctgggaacca gatgcaagta agctgaatga gaattactgc 1320

cgaaatccag atgatgatgc tcatggaccc tggtgctaca cgggaaatcc actcattcct 1380

tgggattatt gccctatttc tcgttgtgaa ggtgatacca cacctacaat agtcaattta 1440

gaccatcccg taatatcttg tgccaaaacg aaacaattgc gagttgtaaa tgggattcca 1500

acacgaacaa acataggatg gatggttagt ttgagataca gaaataaaca tatctgcgga 1560

ggatcattga taaaggagag ttgggttctt actgcacgac agtgtttccc ttctcgagac 1620

ttgaaagatt atgaagcttg gcttggaatt catgatgtcc acggaagagg agatgagaaa 1680

tgcaaacagg ttctcaatgt ttcccagctg gtatatggcc ctgaaggatc agatctggtt 1740

ttaatgaagc ttgccaggcc tgctgtcctg gatgattttg ttagtacgat tgatttacct 1800

aattatggat gcacaattcc tgaaaagacc agttgcagtg tttatggctg gggctacact 1860

ggattgatca actatgatgg cctattacga gtggcacatc tctatataat gggaaatgag 1920

aaatgcagcc agcatcatcg agggaaggtg actctgaatg agtctgaaat atgtgctggg 1980

gctgaaaaga ttggatcagg accatgtgag ggggattatg gtggcccact tgtttgtgag 2040

caacataaaa tgagaatggt tcttggtgtc attgttcctg gtcgtggatg tgccattcca 2100

aatcgtcctg gtatttttgt ccgagtagca tattatgcaa aatggataca caaaattatt 2160

ttaacatata aggtaccaca gtcatag 2187

<210> 2
<211> 7112
<212> DNA
<213> hepatocyte growth factor hybrid

<400> 2
atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc 60

ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat 120

gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa 180

accaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt 240

ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc 300

ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa 360

aacaaagact acattagaaa ctgcatcatt ggtaaaggac gcagctacaa gggaacagta 420

tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac 480

aggtaagaac agtatgaaga aaagagatga agcctctgtc ttttttacat gttaacagtc 540

tcatattagt ccttcagaat aattctacaa tcctaaaata acttagccaa cttgctgaat 600

tgtattacgg caaggtttat atgaattcat gactgatatt tagcaaatga ttaattaata 660

tgttaataaa atgtagccaa aacaatatct taccttaatg cctcaatttg tagatctcgg 720

tatttgtgaa ataataacgt aaacttcgtt taaaaggatt cttcttcctg tctttgagaa 780

agtacggcac tgtgcagggg gagaggttga ttgtgaaaaa tcagaggtag atgagaatct 840

tactgagggc tgagggttct ttaaccttgg tggatctcaa cattggttgc acattaaaat 900

cacctgctgc aagcccttga cgaatcttac ttagaagatg acaacacaga acaattaaat 960

cagaatctct ggggagaata gggcaccagt atttttgag ctcccaccat gattccaaag 1020

tgcagccaaa tttgagaacc actgctaaaa gctcaagctt cagattgacc agcttttcca 1080

tctcacctat cgcctaaaga ccaaattgga taaatgtgtt cattacgaca gatgggtact 1140

atttaaagat gagtaaacac aatatactta ggctcgtcag actgagagtt ttaatcatca 1200

ctgaggaaaa acatagatat ctaatactga ctggagtatt agtcaaggct tatttcacac 1260

acaattttat cagaaaccaa agtagtttaa aacagctctc cccttattag taatgcattg 1320

gagggtttac tttaccatgt accttgctga gcactgtacc ttgttaatct catttacttg 1380

taatgagaac cacacagcgg gtagttttat tggttctatt ttacctacat gacaaaactg 1440

aagcataaaa acacttagta agttttcagt gtcatgcaca actaggaagt gacatggcca 1500

gaatataagc ccagtcacca tcactctata acctgcgctt ttaacaactt cagggcatga 1560

cacatttggc cggtcagtag aacccatgct gtgatttgtt tttgcagtgg tggtgatgac 1620

tgccttgttg aatccacttt ttattctatt ccattttggg gacacaattc tgcaagatga 1680

ttcttcatta ggaaacagag atgagttatt gaccaacaca gaaagaaaaa gagtttgttg 1740

ctccacactg ggattaaacc tatgatcttg gcctaattaa cactagctag taagtgtcca 1800

agctgatcat ctctacaaca tttcaataac agaaaacaac aattttcaaa attagttact 1860

tacaattatg tagaaatgcc tctaaaacac agtattttcc ttatattaca aaaacaaaaa 1920

ttataattgg ttttgtcctc ttttgagagt ttgcatggtg ttactccctg catagtgaag 1980

aaaacatttt atttaagtag atggatctaa gtttttcatg aacaaaggaa tgacatttga 2040

aatcaatcct accctagtcc aggagaatgc attagattaa cctagtagag gtcttatttc 2100

accctgagtt ttctatgatc gtgattctct gctggaggag taattgtgaa atagatctct 2160

ctgggaactg gcttcctagt ccaatcagct cttttaccaa tgaacacttc cttgtgatat 2220

agatgtttat ggccgagagg atccagtata ttaataaaat ccctttttgt attcaatgag 2280

ggaaacacat aattttcatc aattagcagc ttattggaat atctgcatga tggtttaaca 2340

cttttaagtg ttgactaaag attaatttta cagaaaatag aaaaagaaat atgtttctgt 2400

ctggaggaat gatttattgt tgacccctaa attgaaatat tttactagtg gcttaatgga 2460

aagatgatga aagatgatga aattaatgta gaagcttaac tagaaaatca ggtgacctga 2520

tatctacatc tgtatccttc attggccacc cagcattcat taatgaatca gatgatggaa 2580

tagatcaagt ttcctaggaa cacagtgaat attaaaagaa aacaaagga gcctagcacc 2640

tagaagacct agtttatatt tcaaagtata tttggatgta acccaatttt aaacatttcc 2700

tcacttgtct ctcttaaagc cttgccaaca gcaaggacag agaaccaaaa atagtgtata 2760

tatgaataaa tgcttattac agaatctgct gactggcaca tgctttgtgt gtaatgggtt 2820

ctcataaaca cttgttgaat gaacacacat aagtgaaaga gcatggctag gcttcatccc 2880

ttggtcaaat atggggtgct aaagaaaagc aggggaaata cattgggaca ctaacaaaaa 2940

aaaacagtta atttaggtaa aagataaaat acaccacaga atgaagaaaa gagatgaccc 3000

agactgctct ttaaccttca tgtcctagag aggtttttga tatgaattgc attcagaatt 3060

gtggaaagga gcccatcttt tctcttcatt ttgattttat taactccaat gggggaattt 3120

tattcgtgtt ttggccatat ctacttttga tttctacatt attctctctt cctttctacc 3180

tgtatttgtc ctaataaatt gttgacttat taattcacta cttcctcaca gcttttttt 3240

ggctttacaa atccactgga aaggtatatg ggtgtatcac tttgtgtatt tcggtgtgca 3300

tgtgtagagg ggacaaaaat cctctctcaa actataaata ttgagtattt gtgtattgaa 3360

catttgctat aactactagg tttcttaaat aatcttaata tataaaatga tatagaaaaa 3420

gggaaattat agttcgtatt attcatctaa gtgaagagat taaaacccag ggagtaaata 3480

aattgtctaa ggactaaggt tgtatactat ttaggtgata gatatggggc aaccgtatgg 3540

gttttatgat taacaaataa acttctcacc actctaccat atcaactttt ccataaaaga 3600

gagctatagt attctttgct taaataaatt tgattagtgc atgacttctt gaaaacatat 3660

aaagcaaaag tcacatttga ttctatcaga aaagtgagta agccatggcc caaacaaaag 3720

atgcattaaa atattctgga atgatggagc taaaagtaag aaaaatgact ttttaaaaaa 3780

gtttactgtt aggaattgtg aaattatgct gaattttagt tgcattataa tttttgtcag 3840

tcatacggtc tgacaacctg tcttatttct atttccccat atgaggaatg ctagttaagt 3900

atggatatta actattacta cttagatgca ttgaagttgc ataatatgga taatacttca 3960

ctggttccct gaaaatgttt agttagtaat aagtctctta cactatttgt tttgtccaat 4020

aatttatatt ttctgaagac ttaactctag aatacactca tgtcaaaatg aaagaatttc 4080

attgcaaaat attgcttggt acatgacgca tacctgtatt tgttttgtgt cacaacatga 4140

aaaatgatgg tttattagaa gtttcattgg gtaggaaaca catttgaatg gtatttacta 4200

agatactaaa atccttggac ttcactctaa ttttagtgcc atttagaact caaggtctca 4260

gtaaaagtag aaataaagcc tgttaacaaa acacaagctg aatattaaaa atgtaactgg 4320

attttcaaag aaatgtttac tggtattacc tgtagatgta tattctttat tatgatcttt 4380

tgtgtaaagt ctggcagaca aatgcaatat ctaattgttg agtccaatat cacaagcagt 4440

acaaaagtat aaaaaagact tggccttttc taatgtgtta aaatacttta tgctggtaat 4500

aacactaaga gtagggcact agaaatttta agtgaagata atgtgttgca gttactgcac 4560

tcaatggctt actattataa accaaaactg ggatcactaa gctccagtca gtcaaaatga 4620

tcaaaattat tgaagagaat aagcaattct gttctttatt aggacacagt agatacagac 4680

tacaaagtgg agtgtgctta ataagaggta gcatttgtta agtgtcaatt actctattat 4740

cccttggagc ttctcaaaat aaccatataa ggtgtaagat gttaaaggtt atggttacac 4800

tcagtgcaca gtaagctaat aggctgagag aagctaaatt acttactggg gtctcacagt 4860

aagaaagtga gctgaagttt cagcccagat ttaactggat tctgggctct ttattcatgt 4920

tacttcatga atctgtttct caattgtgca gaaaaaaggg ggctatttat aagaaaagca 4980

ataaacaaac aagtaatgat ctcaaataag taatgcaaga aatagtgaga tttcaaaatc 5040

agtggcagcg atttctcagt tctgtcctaa gtggccttgc tcaatcacct gctatctttt 5100

agtggagctt tgaaattatg tttcagacaa cttcgattca gttctagaat gtttgactca 5160

gcaaattcac aggctcatct ttctaacttg atggtgaata tggaaattca gctaaatgga 5220

tgttaataaa attcaaacgt tttaaggaca gatgaaaatg acagaatttt aaggtaaaat 5280

atatgaagga atataagata aaggattttt ctaccttcag caaaaacata cccactaatt 5340

agtaaaatta ataggcaaaa aaaagttgca tgctcttata ctgtaatgat tatcatttta 5400

aaactagctt tttgccttcg agctatcggg gtaaagacct acaggaaaac tactgtcgaa 5460

atcctcgagg ggaagaaggg ggaccctggt gtttcacaag caatccagag gtacgctacg 5520

aagtctgtga cattcctcag tgttcagaag ttgaatgcat gacctgcaat ggggagagtt 5580

atcgaggtct catggatcat acagaatcag gcaagatttg tcagcgctgg gatcatcaga 5640

caccacaccg gcacaaattc ttgcctgaaa gatatcccga caagggcttt gatgataatt 5700

attgccgcaa tcccgatggc cagccgaggc catggtgcta tactcttgac cctcacaccc 5760

gctgggagta ctgtgcaatt aaaacatgcg ctgacaatac tatgaatgac actgatgttc 5820

ctttggaaac aactgaatgc atccaaggtc aaggagaagg ctacaggggc actgtcaata 5880

ccatttggaa tggaattcca tgtcagcgtt gggattctca gtatcctcac gagcatgaca 5940

tgactcctga aaatttcaag tgcaaggacc tacgagaaaa ttactgccga aatccagatg 6000

ggtctgaatc accctggtgt tttaccactg atccaaacat ccgagttggc tactgctccc 6060

aaattccaaa ctgtgatatg tcacatggac aagattgtta tcgtgggaat ggcaaaaatt 6120

atatgggcaa cttatcccaa acaagatctg gactaacatg ttcaatgtgg gacaagaaca 6180

tggaagactt acatcgtcat atcttctggg aaccagatgc aagtaagctg aatgagaatt 6240

actgccgaaa tccagatgat gatgctcatg gaccctggtg ctacacggga aatccactca 6300

ttccttggga ttattgccct atttctcgtt gtgaaggtga taccacacct acaatagtca 6360

atttagacca tcccgtaata tcttgtgcca aaacgaaaca attgcgagtt gtaaatggga 6420

ttccaacacg aacaaacata ggatggatgg ttagtttgag atacagaaat aaacatatct 6480

gcggaggatc attgataaag gagagttggg ttcttactgc acgacagtgt ttcccttctc 6540

gagacttgaa agattatgaa gcttggcttg gaattcatga tgtccacgga agaggagatg 6600

agaaatgcaa acaggttctc aatgtttccc agctggtata tggccctgaa ggatcagatc 6660

tggttttaat gaagcttgcc aggcctgctg tcctggatga ttttgttagt acgattgatt 6720

tacctaatta tggatgcaca attcctgaaa agaccagttg cagtgtttat ggctggggct 6780

acactggatt gatcaactat gatggcctat tacgagtggc acatctctat ataatgggaa 6840

atgagaaatg cagccagcat catcgaggga aggtgactct gaatgagtct gaaatatgtg 6900

ctggggctga aaagattgga tcaggaccat gtgaggggga ttatggtggc ccacttgttt 6960

gtgagcaaca taaaatgaga atggttcttg gtgtcattgt tcctggtcgt ggatgtgcca 7020

ttccaaatcg tcctggtatt tttgtccgag tagcatatta tgcaaaatgg atacacaaaa 7080

ttattttaac atataaggta ccacagtcat ag 7112

<210> 3
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF3 primer

<400> 3
gtaaaggacg cgtctacaag ggaacagtat ctat 34

<210> 4
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF4 primer

<400> 4
actggatcct ctcggccata aacatct 27

<210> 5
<211> 34

<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF10 primer

<400> 5
gaagcttagc accatgtggg tgaccaaact cctg 34

<210> 6
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF5 primer

<400> 6
tggccgagag gatccagtat attaata 27

<210> 7
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF7 primer

<400> 7
cccctcgagg atttcgacag tagtttt 27

<210> 8
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF12 primer

<400> 8
gggatccctt cctttctacc tgtatttg 28

<210> 9
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF13 primer

<400> 9
gggatcctgg gtaaacacat ttgaa 25

<210> 10
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF6 primer

<400> 10
gggatcctta tgtttcagac aacttcga 28

<210> 11
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF1 primer

<400> 11
gaagcttgcc accatgtggg tgaccaaact cctg 34

<210> 12
<211> 34
<212> DNA

<213> Artificial Sequence

<220>
<223> gHGF2 primer

<400> 12
gggatccaga acgcgtcctt taccgatgat gcag 34

<210> 13
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF8 primer

<400> 13
gggatccctt ctcgagactt gaaagattat gaagc 35

<210> 14
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> gHGF9 primer

<400> 14
gtctagagcg gccgctatga ctgtggtacc tt 32

<210> 15
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> cHGF5 primer

<400> 15

ggatccacgc gtagcagcac catgtgggtg accaaa 36

<210> 16
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> cHGF3 primer

<400> 16
ggatcctcta gattacttca gctatgactg tggtac 36

<210> 17
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> GHGF5' primer

<400> 17
caaatgtcag ccctggagtt ccatga 26

<210> 18
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> GHGF3' primer

<400> 18
ctggattgct tgtgaaacag ggt 23

<210> 19
<211> 4679
<212> DNA
<213> Artificial Sequence

<220>
<223> HGF-X6 gene

<400> 19

atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc 60

ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat 120

gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa 180

accaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt 240

ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc 300

ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa 360

aacaaagact acattagaaa ctgcatcatc ggtaaaggac gcagctacaa gggaacagta 420

tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac 480

aggtaagaac agtatgaaga aaagagatga agcctctgtc ttttttacat gttaacagtc 540

tcatattagt ccttcagaat aattctacaa tcctaaaata acttagccaa cttgctgaat 600

tgtattacgg caaggtttat atgaattcat gactgatatt tagcaaatga ttaattaata 660

tgttaataaa atgtagccaa aacaatatct taccttaatg cctcaatttg tagatctcgg 720

tatttgtgga tcccttcctt tctacctgta tttgtcctaa taaattgttg acttattaat 780

tcactacttc ctcacagctt ttttttggct ttacaaatcc actggaaagg tatatgggtg 840

tatcactttg tgtatttcgg tgtgcatgtg tagaggggac aaaaatcctc tctcaaacta 900

taaatattga gtatttgtgt attgaacatt tgctataact actaggtttc ttaaataatc 960

ttaatatata aaatgatata gaaaaaggga aattatagtt cgtattattc atctaagtga 1020

agagattaaa acccagggag taaataaatt gtctaaggac taaggttgta tactatttag 1080

gtgatagata tggggcaacc gtatgggttt tatgattaac aaataaactt ctcaccactc 1140

taccatatca acttttccat aaaagagagc tatagtattc tttgcttaaa taaatttgat 1200

tagtgcatga cttcttgaaa acatataaag caaaagtcac atttgattct atcagaaaag 1260

tgagtaagcc atggcccaaa caaaagatgc attaaaatat tctggaatga tggagctaaa 1320

agtaagaaaa atgacttttt aaaaaagttt actgttagga attgtgaaat tatgctgaat 1380

tttagttgca ttataatttt tgtcagtcat acggtctgac aacctgtctt atttctattt 1440

ccccatatga ggaatgctag ttaagtatgg atattaacta ttactactta gatgcattga 1500

agttgcataa tatggataat acttcactgg ttccctgaaa atgtttagtt agtaataagt 1560

ctcttacact atttgttttg tccaataatt tatattttct gaagacttaa ctctagaata 1620

cactcatgtc aaaatgaaag aatttcattg caaaatattg cttggtacat gacgcatacc 1680

tgtatttgtt ttgtgtcaca acatgaaaaa tgatggttta ttagaagttt cattgggtag 1740

gaaacacatt tgaatggtat ttactaagat actaaaatcc ttggacttca ctctaatttt 1800

agtgccattt agaactcaag gtctcagtaa aagtagaaat aaagcctgtt aacaaaacac 1860

aagctgaata ttaaaaatgt aactggattt tcaaagaaat gtttactggt attacctgta 1920

gatgtatatt ctttattatg atcttttgtg taaagtctgg cagacaaatg caatatctaa 1980

ttgttgagtc caatatcaca agcagtacaa aagtataaaa aagacttggc cttttctaat 2040

gtgttaaaat actttatgct ggtaataaca ctaagagtag ggcactagaa attttaagtg 2100

aagataatgt gttgcagtta ctgcactcaa tggcttacta ttataaacca aaactgggat 2160

cactaagctc cagtcagtca aaatgatcaa aattattgaa gagaataagc aattctgttc 2220

tttattagga cacagtagat acagactaca aagtggagtg tgcttaataa gaggtagcat 2280

ttgttaagtg tcaattactc tattatccct tggagcttct caaaataacc atataaggtg 2340

taagatgtta aaggttatgg ttacactcag tgcacaggta agctaatagg ctgagagaag 2400

ctaaattact tactggggtc tcacagtaag aaagtgagct gaagtttcag cccagattta 2460

actggattct gggctcttta ttcatgttac ttcatgaatc tgtttctcaa ttgtgcagaa 2520

aaaaggggggc tatttataag aaaagcaata aacaaacaag taatgatctc aaataagtaa 2580

tgcaagaaat agtgagattt caaaatcagt ggcagcgatt tctcagttct gtcctaagtg 2640

gccttgctca atcacctgct atcttttagt ggagctttga aattatgttt cagacaactt 2700

cgattcagtt ctagaatgtt tgactcagca aattcacagg ctcatctttc taacttgatg 2760

gtgaatatgg aaattcagct aaatggatgt taataaaatt caaacgtttt aaggacagat 2820

gaaaatgaca gaattttaag gtaaaatata tgaaggaata taagataaag gatttttcta 2880

ccttcagcaa aaacataccc actaattagt aaaattaata ggcaaaaaaa agttgcatgc 2940

tcttatactg taatgattat cattttaaaa ctagcttttt gccttcgagc tatcggggta 3000

aagacctaca ggaaaactac tgtcgaaatc ctcgagggga agaaggggga ccctggtgtt 3060

tcacaagcaa tccagaggta cgctacgaag tctgtgacat tcctcagtgt tcagaagttg 3120

aatgcatgac ctgcaatggg gagagttatc gaggtctcat ggatcataca gaatcaggca 3180

agatttgtca gcgctgggat catcagacac cacaccggca caaattcttg cctgaaagat 3240

atcccgacaa gggctttgat gataattatt gccgcaatcc cgatggccag ccgaggccat 3300

ggtgctatac tcttgaccct cacacccgct gggagtactg tgcaattaaa acatgcgctg 3360

acaatactat gaatgacact gatgttcctt tggaaacaac tgaatgcatc caaggtcaag 3420

gagaaggcta caggggcact gtcaatacca tttggaatgg aattccatgt cagcgttggg 3480

attctcagta tcctcacgag catgacatga ctcctgaaaa tttcaagtgc aaggacctac 3540

gagaaaatta ctgccgaaat ccagatgggt ctgaatcacc ctggtgtttt accactgatc 3600

caaacatccg agttggctac tgctcccaaa ttccaaactg tgatatgtca catggacaag 3660

attgttatcg tgggaatggc aaaaattata tgggcaactt atcccaaaca agatctggac 3720

taacatgttc aatgtgggac aagaacatgg aagacttaca tcgtcatatc ttctgggaac 3780

cagatgcaag taagctgaat gagaattact gccgaaatcc agatgatgat gctcatggac 3840

cctggtgcta cacgggaaat ccactcattc cttgggatta ttgccctatt tctcgttgtg 3900

aaggtgatac cacacctaca atagtcaatt tagaccatcc cgtaatatct tgtgccaaaa 3960

cgaaacaatt gcgagttgta aatgggattc caacacgaac aaacatagga tggatggtta 4020

gtttgagata cagaaataaa catatctgcg gaggatcatt gataaaggag agttgggttc 4080

ttactgcacg acagtgtttc ccttctcgag acttgaaaga ttatgaagct tggcttggaa 4140

ttcatgatgt ccacggaaga ggagatgaga aatgcaaaca ggttctcaat gtttcccagc 4200

tggtatatgg ccctgaagga tcagatctgg ttttaatgaa gcttgccagg cctgctgtcc 4260

tggatgattt tgttagtacg attgatttac ctaattatgg atgcacaatt cctgaaaaga 4320

ccagttgcag tgtttatggc tggggctaca ctggattgat caactatgat ggcctattac 4380

gagtggcaca tctctatata atgggaaatg agaaatgcag ccagcatcat cgagggaagg 4440

tgactctgaa tgagtctgaa atatgtgctg gggctgaaaa gattggatca ggaccatgtg 4500

agggggatta tggtggccca cttgtttgtg agcaacataa aatgagaatg gttcttggtg 4560

tcattgttcc tggtcgtgga tgtgccattc caaatcgtcc tggtattttt gtccgagtag 4620

catattatgc aaaatggata cacaaaatta ttttaacata taaggtacca cagtcatag 4679

<210> 20
<211> 3679
<212> DNA
<213> Artificial Sequence

<220>
<223> HGF-X7 gene

<400> 20
atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc 60

ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat 120

gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa 180

accaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt 240

ccattcactt gcaaggcttt gtttttgat aaagcaagaa aacaatgcct ctggttcccc 300

ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa 360

aacaaagact acattagaaa ctgcatcatc ggtaaaggac gcagctacaa gggaacagta 420

tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac 480

aggtaagaac agtatgaaga aaagagatga agcctctgtc ttttttacat gttaacagtc 540

tcatattagt ccttcagaat aattctacaa tcctaaaata acttagccaa cttgctgaat 600

tgtattacgg caaggtttat atgaattcat gactgatatt tagcaaatga ttaattaata 660

tgttaataaa atgtagccaa aacaatatct taccttaatg cctcaatttg tagatctcgg 720

tatttgtgga tcctgggtag gaaacacatt tgaatggtat ttactaagat actaaaatcc 780

ttggacttca ctctaatttt agtgccattt agaactcaag gtctcagtaa aagtagaaat 840

aaagcctgtt aacaaaacac aagctgaata ttaaaaatgt aactggattt tcaaagaaat 900

gtttactggt attacctgta gatgtatatt ctttattatg atcttttgtg taaagtctgg 960

cagacaaatg caatatctaa ttgttgagtc caatatcaca agcagtacaa aagtataaaa 1020

aagacttggc cttttctaat gtgttaaaat actttatgct ggtaataaca ctaagagtag 1080

ggcactagaa attttaagtg aagataatgt gttgcagtta ctgcactcaa tggcttacta 1140

ttataaacca aaactgggat cactaagctc cagtcagtca aaatgatcaa aattattgaa 1200

gagaataagc aattctgttc tttattagga cacagtagat acagactaca aagtggagtg 1260

tgcttaataa gaggtagcat ttgttaagtg tcaattactc tattatccct tggagcttct 1320

caaaataacc atataaggtg taagatgtta aaggttatgg ttacactcag tgcacaggta 1380

agctaatagg ctgagagaag ctaaattact tactggggtc tcacagtaag aaagtgagct 1440

gaagtttcag cccagattta actggattct gggctcttta ttcatgttac ttcatgaatc 1500

tgtttctcaa ttgtgcagaa aaaagggggc tatttataag aaaagcaata aacaaacaag 1560

taatgatctc aaataagtaa tgcaagaaat agtgagattt caaaatcagt ggcagcgatt 1620

tctcagttct gtcctaagtg gccttgctca atcacctgct atcttttagt ggagctttga 1680

aattatgttt cagacaactt cgattcagtt ctagaatgtt tgactcagca aattcacagg 1740

ctcatctttc taacttgatg gtgaatatgg aaattcagct aaatggatgt taataaaatt 1800

caaacgtttt aaggacagat gaaaatgaca gaattttaag gtaaaatata tgaaggaata 1860

taagataaag gatttttcta ccttcagcaa aaacataccc actaattagt aaaattaata 1920

ggcaaaaaaa agttgcatgc tcttatactg taatgattat cattttaaaa ctagcttttt 1980

gccttcgagc tatcggggta aagacctaca ggaaaactac tgtcgaaatc ctcgagggga 2040

agaaggggga ccctggtgtt tcacaagcaa tccagaggta cgctacgaag tctgtgacat 2100

tcctcagtgt tcagaagttg aatgcatgac ctgcaatggg gagagttatc gaggtctcat 2160

ggatcataca gaatcaggca agatttgtca gcgctgggat catcagacac cacaccggca 2220

caaattcttg cctgaaagat atcccgacaa gggctttgat gataattatt gccgcaatcc 2280

cgatggccag ccgaggccat ggtgctatac tcttgaccct cacacccgct gggagtactg 2340

tgcaattaaa acatgcgctg acaatactat gaatgacact gatgttcctt tggaaacaac 2400

tgaatgcatc caaggtcaag gagaaggcta caggggcact gtcaatacca tttggaatgg 2460

aattccatgt cagcgttggg attctcagta tcctcacgag catgacatga ctcctgaaaa 2520

tttcaagtgc aaggacctac gagaaaatta ctgccgaaat ccagatgggt ctgaatcacc 2580

ctggtgtttt accactgatc caaacatccg agttggctac tgctcccaaa ttccaaactg 2640

tgatatgtca catggacaag attgttatcg tgggaatggc aaaaattata tgggcaactt 2700

atcccaaaca agatctggac taacatgttc aatgtgggac aagaacatgg aagacttaca 2760

tcgtcatatc ttctgggaac cagatgcaag taagctgaat gagaattact gccgaaatcc 2820

agatgatgat gctcatggac cctggtgcta cacgggaaat ccactcattc cttgggatta 2880

ttgccctatt tctcgttgtg aaggtgatac cacacctaca atagtcaatt tagaccatcc 2940

cgtaatatct tgtgccaaaa cgaaacaatt gcgagttgta aatgggattc caacacgaac 3000

aaacatagga tggatggtta gtttgagata cagaaataaa catatctgcg gaggatcatt 3060

gataaaggag agttgggttc ttactgcacg acagtgtttc ccttctcgag acttgaaaga 3120

ttatgaagct tggcttggaa ttcatgatgt ccacggaaga ggagatgaga aatgcaaaca 3180

ggttctcaat gtttcccagc tggtatatgg ccctgaagga tcagatctgg ttttaatgaa 3240

gcttgccagg cctgctgtcc tggatgattt tgttagtacg attgatttac ctaattatgg 3300

atgcacaatt cctgaaaaga ccagttgcag tgtttatggc tggggctaca ctggattgat 3360

caactatgat ggcctattac gagtggcaca tctctatata atgggaaatg agaaatgcag 3420

ccagcatcat cgagggaagg tgactctgaa tgagtctgaa atatgtgctg gggctgaaaa 3480

gattggatca ggaccatgtg agggggatta tggtggccca cttgtttgtg agcaacataa 3540

aatgagaatg gttcttggtg tcattgttcc tggtcgtgga tgtgccattc caaatcgtcc 3600

tggtattttt gtccgagtag catattatgc aaaatggata cacaaaatta ttttaacata 3660

taaggtacca cagtcatag 3679

<210> 21
<211> 2729
<212> DNA
<213> Artificial Sequence

<220>
<223> HGF-X8 gene

<400> 21
atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc 60

ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat 120

gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa 180

accaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt 240

ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc 300

ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa 360

aacaaagact acattagaaa ctgcatcatc ggtaaaggac gcagctacaa gggaacagta 420

tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac 480

aggtaagaac agtatgaaga aaagagatga agcctctgtc ttttttacat gttaacagtc 540

tcatattagt ccttcagaat aattctacaa tcctaaaata acttagccaa cttgctgaat 600

tgtattacgg caaggtttat atgaattcat gactgatatt tagcaaatga ttaattaata 660

tgttaataaa atgtagccaa aacaatatct taccttaatg cctcaatttg tagatctcgg 720

tatttgtgga tccttatgtt tcagacaact tcgattcagt tctagaatgt ttgactcagc 780

aaattcacag gctcatcttt ctaacttgat ggtgaatatg gaaattcagc taaatggatg 840

ttaataaaat tcaaacgttt taaggacaga tgaaaatgac agaattttaa ggtaaaatat 900

atgaaggaat ataagataaa ggatttttct accttcagca aaaacatacc cactaattag 960

taaaattaat aggcaaaaaa aagttgcatg ctcttatact gtaatgatta tcattttaaa 1020

actagctttt tgccttcgag ctatcggggt aaagacctac aggaaaacta ctgtcgaaat 1080

cctcgagggg aagaaggggg accctggtgt ttcacaagca atccagaggt acgctacgaa 1140

gtctgtgaca ttcctcagtg ttcagaagtt gaatgcatga cctgcaatgg ggagagttat 1200

cgaggtctca tggatcatac agaatcaggc aagatttgtc agcgctggga tcatcagaca 1260

ccacaccggc acaaattctt gcctgaaaga tatcccgaca agggctttga tgataattat 1320

tgccgcaatc ccgatggcca gccgaggcca tggtgctata ctcttgaccc tcacacccgc 1380

tgggagtact gtgcaattaa aacatgcgct gacaatacta tgaatgacac tgatgttcct 1440

ttggaaacaa ctgaatgcat ccaaggtcaa ggagaaggct acaggggcac tgtcaatacc 1500

atttggaatg gaattccatg tcagcgttgg gattctcagt atcctcacga gcatgacatg 1560

actcctgaaa atttcaagtg caaggaccta cgagaaaatt actgccgaaa tccagatggt 1620

ctgaatcacc ctggtgtttt accactgatc caaacatccg agttggctac tgctcccaaa 1680

ttccaaactg tgatatgtca catggacaag attgttatcg tgggaatggc aaaaattata 1740

tgggcaactt atcccaaaca agatctggac taacatgttc aatgtgggac aagaacatgg 1800

aagacttaca tcgtcatatc ttctgggaac cagatgcaag taagctgaat gagaattact 1860

gccgaaatcc agatgatgat gctcatggac cctggtgcta cacgggaaat ccactcattc 1920

cttgggatta ttgccctatt tctcgttgtg aaggtgatac cacacctaca atagtcaatt 1980

tagaccatcc cgtaatatct tgtgccaaaa cgaaacaatt gcgagttgta aatgggattc 2040

caacacgaac aaacatagga tggatggtta gtttgagata cagaaataaa catatctgcg 2100

gaggatcatt gataaaggag agttgggttc ttactgcacg acagtgtttc ccttctcgag 2160

acttgaaaga ttatgaagct tggcttggaa ttcatgatgt ccacggaaga ggagatgaga 2220

aatgcaaaca ggttctcaat gtttcccagc tggtatatgg ccctgaagga tcagatctgg 2280

ttttaatgaa gcttgccagg cctgctgtcc tggatgattt tgttagtacg attgatttac 2340

ctaattatgg atgcacaatt cctgaaaaga ccagttgcag tgtttatggc tggggctaca 2400

ctggattgat caactatgat ggcctattac gagtggcaca tctctatata atgggaaatg 2460

agaaatgcag ccagcatcat cgagggaagg tgactctgaa tgagtctgaa atatgtgctg 2520

gggctgaaaa gattggatca ggaccatgtg agggggatta tggtggccca cttgtttgtg 2580

agcaacataa aatgagaatg gttcttggtg tcattgttcc tggtcgtgga tgtgccattc 2640

caaatcgtcc tggtattttt gtccgagtag catattatgc aaaatggata cacaaaatta 2700

ttttaacata taaggtacca cagtcatag 2729