PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 :
A61K 31/165, 9/08, 47/10

A1

(11) International Publication Number: WO 00/07588

(43) International Publication Date: 17 February 2000 (17.02.00)

(21) International Application Number: PCT/EP99/05486

(22) International Filing Date: 27 July 1999 (27.07.99)

(30) Priority Data:
MI98A001795 31 July 1998 (31.07.98) IT

(71) Applicant *(for all designated States except US)*: AZIENDE CHIMICHE RIUNITE ANGELINI FRANCESCO A.C.R.A.F. S.P.A. [IT/IT]; Viale Amelia, 70, I-00181 Roma (IT).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: CAVALLO, Giovanni [IT/IT]; Via delle Tartane, 45, I-00122 Ostia (IT). PINZA, Mario [IT/IT]; Via per Cesano Boscone, 24, I-20094 Corsico (IT).

(74) Agents: MARCHI, Massimo et al.; Marchi & Partners S.r.l., Via Pirelli, 19, I-20124 Milano (IT).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: PHARMACEUTICAL COMPOSITION FOR INJECTION BASED ON PARACETAMOL

(57) Abstract

A pharmaceutical composition, characterized in that: a) it comprises i) paracetamol, ii) from 1 to 4 parts by volume of a low molecular weight alcohol for each part by weight of paracetamol, and iii) from 1 to 5 parts by volume of a polyethylene glycol for each part by weight of paracetamol; b) it is substantially anhydrous; and c) it forms a clear solution for injection with 4–10 parts by volume of water for each part by weight of paracetamol.