

PYRAZOLOPYRIDINES AND TRIAZOLOPYRIDINES AS A2A / A2B INHIBITORS**TECHNICAL FIELD**

The present invention provides fused heterocycles and derivatives thereof that
5 modulate the activity of adenosine receptors, such as subtypes A2A and A2B, and are useful
in the treatment of diseases related to the activity of adenosine receptors including, for
example, cancer, inflammatory diseases, cardiovascular diseases and neurodegenerative
diseases.

BACKGROUND

Adenosine is an extracellular signaling molecule that can modulate immune
responses through many immune cell types. Adenosine was first recognized as a physiologic
regulator of coronary vascular tone by Drury and Szent-György (Sachdeva, S. and Gupta, M.
Saudi Pharmaceutical Journal, 2013, 21, 245–253), however it was not until 1970 that Sattin
15 and Rall showed that adenosine regulates cell function via occupancy of specific receptors on
the cell surface (Sattin, A., and Rall, T.W., 1970. *Mol. Pharmacol.* 6, 13–23; Hasko', G., at
al., 2007, *Pharmacol. Ther.* 113, 264–275).

Adenosine plays a vital role in various other physiological functions. It is involved in
the synthesis of nucleic acids and when linked to three phosphate groups, it forms ATP, the
20 integral component of the cellular energy system. Adenosine can be generated by the
enzymatic breakdown of extracellular ATP, or can also be released from injured neurons and
glial cells by passing the damaged plasma membrane (Tautenhahn, M. et al.
Neuropharmacology, 2012, 62, 1756–1766). Adenosine produces various pharmacological
effects, both in the periphery and in the central nervous system (CNS), through an action on
25 specific receptors localized on cell membranes (Matsumoto, T. et al. *Pharmacol. Res.*, 2012,
65, 81–90). Alternative pathways for extracellular adenosine generation have been described.
These pathways include the production of adenosine from nicotinamide dinucleotide (NAD)
instead of ATP by the concerted action of CD38, CD203a and CD73. CD73-independent
production of adenosine can also occur by other phosphates such as alkaline phosphatase or
30 prostate-specific phosphatase.

There are four known subtypes of adenosine receptor in humans including A1, A2A,
A2B and A3 receptors. A1 and A2A are high affinity receptors, whereas A2B and A3 are low
affinity receptors. Adenosine and its agonists can act via one or more of these receptors and
can modulate the activity of adenylate cyclase, the enzyme responsible for increasing cyclic
35 AMP (cAMP). The different receptors have differential stimulatory and inhibitory effects on

this enzyme. Increased intracellular concentrations of cAMP can suppress the activity of immune and inflammatory cells (Livingston, M. et al., *Inflamm. Res.*, 2004, 53, 171–178).

The A2A adenosine receptor can signal in the periphery and the CNS, with agonists explored as anti-inflammatory drugs and antagonists explored for neurodegenerative diseases (Carlsson, J. et al., *J. Med. Chem.*, 2010, 53, 3748–3755). In most cell types the A2A subtype inhibits intracellular calcium levels whereas the A2B potentiates them. The A2A receptor generally appears to inhibit inflammatory response from immune cells (Borrmann, T. et al., *J. Med. Chem.*, 2009, 52(13), 3994–4006).

A2B receptors are highly expressed in the gastrointestinal tract, bladder, lung and on mast cells (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842–857). The A2B receptor, although structurally closely related to the A2A receptor and able to activate adenylate cyclase, is functionally different. It has been postulated that this subtype may utilize signal transduction systems other than adenylate cyclase (Livingston, M. et al., *Inflamm. Res.*, 2004, 53, 171–178). Among all the adenosine receptors, the A2B adenosine receptor is a low affinity receptor that is thought to remain silent under physiological conditions and to be activated in consequence of increased extracellular adenosine levels (Ryzhov, S. et al. *Neoplasia*, 2008, 10, 987–995). Activation of A2B adenosine receptor can stimulate adenylate cyclase and phospholipase C through activation of Gs and Gq proteins, respectively. Coupling to mitogen activated protein kinases has also been described (Borrmann, T. et al., *J. Med. Chem.*, 2009, 52(13), 3994–4006).

In the immune system, engagement of adenosine signaling can be a critical regulatory mechanism that protects tissues against excessive immune reactions. Adenosine can negatively modulate immune responses through many immune cell types, including T-cells, natural-killer cells, macrophages, dendritic cells, mast cells and myeloid-derived suppressor cells (Allard, B. et al. *Current Opinion in Pharmacology*, 2016, 29, 7–16).

In tumors, this pathway is hijacked by tumor micro-environments and sabotages the antitumor capacity of immune system, promoting cancer progression. In the tumor micro-environment, adenosine was mainly generated from extracellular ATP by CD39 and CD73. Multiple cell types can generate adenosine by expressing CD39 and CD73. This is the case for tumor cells, T-effector cells, T-regulatory cells, tumor associated macrophages, myeloid derived suppressive cells (MDSCs), endothelial cells, cancer-associated fibroblast (CAFs) and mesenchymal stromal/stem cells (MSCs). Hypoxia, inflammation and other immune-suppressive signaling in tumor micro-environment can induce expression of CD39, CD73 and subsequent adenosine production. As a result, adenosine level in solid tumors is unusually high compared to normal physiological conditions.

A2A are mostly expressed on lymphoid-derived cells, including T-effector cells, T regulatory cells and nature killing cells. Blocking A2A receptor can prevent downstream immunosuppressive signals that temporarily inactivate T cells. A2B receptors are mainly expressed on monocyte-derived cells including dendritic cells, tumor-associated
5 macrophages, myeloid derived suppressive cells (MDSCs), and mesenchymal stromal/stem cells (MSCs). Blocking A2B receptor in preclinical models can suppress tumor growth, block metastasis, and increase the presentation of tumor antigens.


In terms of safety profile of ADORA2A/ADORA2B (A2A/A2B) blockage, the A2A and A2B receptor knockout (KO) mice are all viable, showing no growth abnormalities and are fertile (Allard, B. et al. *Current Opinion in Pharmacology*, 2016, 29, 7–16). A2A KO
10 mice displayed increased levels of pro-inflammatory cytokines only upon challenge with LPS and no evidence of inflammation at baseline (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842-857). A2B KO mice exhibited normal platelet, red blood, and white cell counts but increased inflammation at baseline (TNF-alpha, IL-6) in naive A2B KO mice (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842-857). Exaggerated production of TNF-alpha
15 and IL-6 was detected following LPS treatment. A2B KO mice also exhibited increased vascular adhesion molecules that mediate inflammation as well leukocyte adhesion/rolling; enhanced mast-cell activation; increased sensitivity to IgE-mediated anaphylaxis and increased vascular leakage and neutrophil influx under hypoxia (Antonioli, L. et al., *Nature Reviews Cancer*, 2013, 13, 842-857).
20

In summary, there is a need to develop new adenosine receptor selective ligands, such as for subtypes A2A and A2B, for the treatment of diseases such as cancer, inflammatory diseases, cardiovascular diseases and neurodegenerative diseases. This application is directed to this need and others.

25

SUMMARY

The present invention relates to, *inter alia*, compounds of Formula (I):



(I),

or pharmaceutically acceptable salts thereof, wherein constituent members are defined
5 herein.

The present invention further provides pharmaceutical compositions comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

10 The present invention further provides methods of inhibiting an activity of an adenosine receptor, comprising contacting the receptor with a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

The present invention further provides methods of treating a disease or a disorder associated with abnormal expression of adenosine receptors, comprising administering to said patient a therapeutically effective amount of a compound of Formula (I), or a
15 pharmaceutically acceptable salt thereof.


The present invention further provides a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for use in any of the methods described herein.

20 The present invention further provides use of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for the preparation of a medicament for use in any of the methods described herein.

DETAILED DESCRIPTION

Compounds

The present application provides, *inter alia*, a compound of Formula (I):



5 or a pharmaceutically acceptable salt thereof, wherein:

X^3 is N or CR^3 ;

X^4 is N or CR^4 ;

Cy^1 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy^1 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^5 substituents;

Cy^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy^2 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^6 substituents;

R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $C(=NR^{e1})R^{b1}$, $C(=NR^{e1})NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl of R^1 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

20 each R^{a1} , R^{b1} , R^{c1} , and R^{d1} is independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl of R^{a1} , R^{b1} , R^{c1} , and R^{d1} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

25 each R^{e1} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, and C_{1-6} haloalkoxy;

each R^{1A} is independently selected from OH, CN, halo, C_{1-3} alkyl, C_{2-3} alkenyl, C_{2-3} alkynyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, C_{1-3} haloalkoxy, amino, C_{1-3} alkylamino, and di(C_{1-3} alkyl)amino;

- R^2 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $C(=NR^{e2})R^{b2}$,
5 $C(=NR^{e2})NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, and $S(O)_2NR^{c2}R^{d2}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^2 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;
10 each R^{a2} , R^{b2} , R^{c2} , and R^{d2} is independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered
15 heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a2} , R^{b2} , R^{c2} , and R^{d2} , are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;
or, any R^{c2} and R^{d2} , attached to the same N atom, together with the N atom to which
20 they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;
each R^{e2} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;
25 each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)NR^{c21}(OR^{a21})$, $C(O)OR^{a21}$, $OC(O)R^{b21}$,
30 $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $C(=NR^{e21})R^{b21}$, $C(=NR^{e21})NR^{c21}R^{d21}$, $NR^{c21}C(=NR^{e21})NR^{c21}R^{d21}$, $NR^{c21}C(=NR^{e21})R^{b21}$, $NR^{c21}S(O)NR^{c21}R^{d21}$, $NR^{c21}S(O)R^{b21}$, $NR^{c21}S(O)_2R^{b21}$, $NR^{c21}S(O)(=NR^{e21})R^{b21}$, $NR^{c21}S(O)_2NR^{c21}R^{d21}$, $S(O)R^{b21}$, $S(O)NR^{c21}R^{d21}$, $S(O)_2R^{b21}$, $S(O)_2NR^{c21}R^{d21}$, $OS(O)(=NR^{e21})R^{b21}$, $OS(O)_2R^{b21}$, SF_3 , $P(O)R^{f21}R^{g21}$, $OP(O)(OR^{h21})(OR^{i21})$,
35 $P(O)(OR^{h21})(OR^{i21})$, and $BR^{j21}R^{k21}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10}

aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2A} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2B} substituents;

5 each R^{a21}, R^{b21}, R^{c21} and R^{d21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a21}, R^{b21}, R^{c21} and R^{d21} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2B} substituents;

or, any R^{c21} and R^{d21}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2B} substituents;

each R^{e21} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

20 each R^{f21} and R^{g21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

25 each R^{h21} and Rⁱ²¹ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

30 each R^{j21} and R^{k21} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j21} and R^{k21} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

35 each R^{2B} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered

- heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a22}, SR^{a22}, NHOR^{a22}, C(O)R^{b22}, C(O)NR^{c22}R^{d22}, C(O)NR^{c22}(OR^{a22}), C(O)OR^{a22}, OC(O)R^{b22}, OC(O)NR^{c22}R^{d22}, NR^{c22}R^{d22}, NR^{c22}NR^{c22}R^{d22}, NR^{c22}C(O)R^{b22}, NR^{c22}C(O)OR^{a22}, NR^{c22}C(O)NR^{c22}R^{d22}, C(=NR^{e22})R^{b22}, C(=NR^{e22})NR^{c22}R^{d22}, NR^{c22}C(=NR^{e22})NR^{c22}R^{d22}, NR^{c22}C(=NR^{e22})R^{b22}, NR^{c22}S(O)NR^{c22}R^{d22}, NR^{c22}S(O)R^{b22}, NR^{c22}S(O)₂R^{b22}, NR^{c22}S(O)(=NR^{e22})R^{b22}, NR^{c22}S(O)₂NR^{c22}R^{d22}, S(O)R^{b22}, S(O)NR^{c22}R^{d22}, S(O)₂R^{b22}, S(O)₂NR^{c22}R^{d22}, OS(O)(=NR^{e22})R^{b22}, OS(O)₂R^{b22}, SF₅, P(O)R^{f22}R^{g22}, OP(O)(OR^{h22})(ORⁱ²²), P(O)(OR^{h22})(ORⁱ²²), and BR^{j22}R^{k22}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2B} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;
- each R^{a22}, R^{b22}, R^{c22} and R^{d22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a22}, R^{b22}, R^{c22} and R^{d22} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;
- or, any R^{e22} and R^{d22}, attached to the same N atom, together with the N atom to which they are attached, form a 4-7 membered heterocycloalkyl group, wherein the 4-7 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;
- each R^{e22} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;
- each R^{f22} and R^{g22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;
- each R^{h22} and Rⁱ²² is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered

heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j22} and R^{k22} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

5 or any R^{j22} and R^{k22} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{2C} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a23}, SR^{a23}, NHOR^{a23}, C(O)R^{b23}, C(O)NR^{c23}R^{d23}, C(O)NR^{c23}(OR^{a23}), C(O)OR^{a23}, OC(O)R^{b23}, OC(O)NR^{c23}R^{d23}, NR^{c23}R^{d23}, NR^{c23}NR^{c23}R^{d23}, NR^{c23}C(O)R^{b23}, NR^{c23}C(O)OR^{a23}, NR^{c23}C(O)NR^{c23}R^{d23}, C(=NR^{e23})R^{b23}, C(=NR^{e23})NR^{c23}R^{d23}, NR^{c23}C(=NR^{e23})NR^{c23}R^{d23}, NR^{c23}C(=NR^{e23})R^{b23}, NR^{c23}S(O)NR^{c23}R^{d23}, NR^{c23}S(O)R^{b23}, NR^{c23}S(O)₂R^{b23}, NR^{c23}S(O)(=NR^{e23})R^{b23}, NR^{c23}S(O)₂NR^{c23}R^{d23}, S(O)R^{b23}, S(O)NR^{c23}R^{d23}, S(O)₂R^{b23}, S(O)₂NR^{c23}R^{d23}, OS(O)(=NR^{e23})R^{b23}, OS(O)₂R^{b23}, SF₅, P(O)R^{f23}R^{g23}, OP(O)(OR^{h23})(ORⁱ²³), P(O)(OR^{h23})(ORⁱ²³), and BR^{j23}R^{k23}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2C} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2D} substituents;

each R^{a23}, R^{b23}, R^{c23} and R^{d23} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a23}, R^{b23}, R^{c23} and R^{d23} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2D} substituents;

or, any R^{e23} and R^{d23}, attached to the same N atom, together with the N atom to which they are attached, form a 4-7 membered heterocycloalkyl group, wherein the 4-7 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2D} substituents;

each R^{e23} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{f23} and R^{g23} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10
 5 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h23} and R^{i23} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered
 10 heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j23} and R^{k23} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j23} and R^{k23} attached to the same B atom, together with the B atom to which
 15 they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{2D} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)-
 20 C_{1-6} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a24} , SR^{a24} , $NHOR^{a24}$, $C(O)R^{b24}$, $C(O)NR^{c24}R^{d24}$, $C(O)NR^{c24}(OR^{a24})$, $C(O)OR^{a24}$, $OC(O)R^{b24}$, $OC(O)NR^{c24}R^{d24}$, $NR^{c24}R^{d24}$, $NR^{c24}NR^{c24}R^{d24}$, $NR^{c24}C(O)R^{b24}$, $NR^{c24}C(O)OR^{a24}$, $NR^{c24}C(O)NR^{c24}R^{d24}$, $C(=NR^{e24})R^{b24}$, $C(=NR^{e24})NR^{c24}R^{d24}$, $NR^{c24}C(=NR^{e24})NR^{c24}R^{d24}$, $NR^{c24}C(=NR^{e24})R^{b24}$, $NR^{c24}S(O)NR^{c24}R^{d24}$, $NR^{c24}S(O)R^{b24}$, $NR^{c24}S(O)_2R^{b24}$, $NR^{c24}S(O)(=NR^{e24})R^{b24}$,
 25 $NR^{c24}S(O)_2NR^{c24}R^{d24}$, $S(O)R^{b24}$, $S(O)NR^{c24}R^{d24}$, $S(O)_2R^{b24}$, $S(O)_2NR^{c24}R^{d24}$, $OS(O)(=NR^{e24})R^{b24}$, $OS(O)_2R^{b24}$, SF_5 , $P(O)R^{f24}R^{g24}$, $OP(O)(OR^{h24})(OR^{i24})$, $P(O)(OR^{h24})(OR^{i24})$, and $BR^{j24}R^{k24}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, and (4-7 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{2D} are each optionally substituted with 1, 2, 3, or 4
 30 independently selected R^{2E} substituents;

each R^{a24} , R^{b24} , R^{c24} and R^{d24} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, and (4-7 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6}
 35

- alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a24}, R^{b24}, R^{c24} and R^{d24} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2E} substituents;
- 5 or, any R^{e24} and R^{d24}, attached to the same N atom, together with the N atom to which they are attached, form a 4-7 membered heterocycloalkyl group, wherein the 4-7 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2E} substituents;
- each R^{e24} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;
- 10 each R^{f24} and R^{g24} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;
- 15 each R^{h24} and Rⁱ²⁴ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;
- each R^{j24} and R^{k24} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;
- 20 or any R^{j24} and R^{k24} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;
- each R^{2E} is independently selected from OH, NO₂, CN, halo, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₃ haloalkyl, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, C₁₋₃ alkoxy, C₁₋₃ haloalkoxy, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxy carbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxy carbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;
- 35 R³ is selected from H, D, halo, OH, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃

- alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;
- R⁴ is selected from H, D, halo, OH, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;
- each R⁵ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a5}, SR^{a5}, NHOR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)NR^{c5}(OR^{a5}), C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(O)NR^{c5}R^{d5}, C(=NR^{e5})R^{b5}, C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})R^{b5}, NR^{c5}S(O)NR^{c5}R^{d5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)(=NR^{e5})R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, S(O)₂NR^{c5}R^{d5}, OS(O)(=NR^{e5})R^{b5}, OS(O)₂R^{b5}, SF₅, P(O)R^{f5}R^{g5}, OP(O)(OR^{h5})(ORⁱ⁵), P(O)(OR^{h5})(ORⁱ⁵), and BR^{j5}R^{k5}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁵ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{5A} substituents;
- each R^{a5}, R^{b5}, R^{c5} and R^{d5} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10

membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a5}, R^{b5}, R^{c5} and R^{d5} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{5A} substituents;

or, any R^{c5} and R^{d5}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{5A} substituents;

each R^{e5} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f5} and R^{g5} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h5} and Rⁱ⁵ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j5} and R^{k5} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy; or any R^{j5} and R^{k5} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{5A} is independently selected from OH, NO₂, CN, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, cyano-C₁₋₆ alkyl, HO-C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, amino, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, thio, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, carbamyl, C₁₋₆ alkylcarbamyl, di(C₁₋₆ alkyl)carbamyl, carboxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxycarbonylamino, C₁₋₆ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₆ alkylaminocarbonyloxy, di(C₁₋₆ alkyl)aminocarbonyloxy, C₁₋₆ alkylsulfonylamino, aminosulfonyl, C₁₋₆ alkylaminosulfonyl, di(C₁₋₆ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₆ alkylaminosulfonylamino, di(C₁₋₆ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₆ alkylaminocarbonylamino, and di(C₁₋₆ alkyl)aminocarbonylamino;

- each R⁶ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a6}, SR^{a6},
 5 NHOR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)NR^{c6}(OR^{a6}), C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)OR^{a6}, NR^{c6}C(O)NR^{c6}R^{d6}, C(=NR^{e6})R^{b6}, C(=NR^{e6})NR^{c6}R^{d6}, NR^{c6}C(=NR^{e6})NR^{c6}R^{d6}, NR^{c6}C(=NR^{e6})R^{b6}, NR^{c6}S(O)NR^{c6}R^{d6}, NR^{c6}S(O)R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)(=NR^{e6})R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, S(O)₂NR^{c6}R^{d6}, OS(O)(=NR^{e6})R^{b6}, OS(O)₂R^{b6}, SF₆, P(O)R^{f6}R^{g6},
 10 OP(O)(OR^{h6})(ORⁱ⁶), P(O)(OR^{h6})(ORⁱ⁶), and BR^{j6}R^{k6}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁶ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{6A} substituents;
 15 or, two R⁶ groups together form an oxo group;
 each R^{a6}, R^{b6}, R^{c6} and R^{d6} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein
 20 the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a6}, R^{b6}, R^{c6} and R^{d6} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{6A} substituents;
 25 or, any R^{c6} and R^{d6}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{6A} substituents;
 each R^{e6} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;
 30 each R^{f6} and R^{g6} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;
 35 heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h6} and R^{i6} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

5 each R^{j6} and R^{k6} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy; or any R^{j6} and R^{k6} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl; and

each R^{6A} is independently selected from OH, NO_2 , CN, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, cyano- C_{1-6} alkyl, HO- C_{1-6} alkyl, C_{1-6} alkoxy- C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, amino, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino, thio, C_{1-6} alkylthio, C_{1-6} alkylsulfinyl, C_{1-6} alkylsulfonyl, carbamyl, C_{1-6} alkylcarbamyl, di(C_{1-6} alkyl)carbamyl, carboxy, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy carbonyl, C_{1-6} alkylcarbonylamino, C_{1-6} alkoxy carbonylamino, C_{1-6} alkylcarbonyloxy, aminocarbonyloxy, C_{1-6} alkylaminocarbonyloxy, di(C_{1-6} alkyl)aminocarbonyloxy, C_{1-6} alkylsulfonylamino, aminosulfonyl, C_{1-6} alkylaminosulfonyl, di(C_{1-6} alkyl)aminosulfonyl, aminosulfonylamino, C_{1-6} alkylaminosulfonylamino, di(C_{1-6} alkyl)aminosulfonylamino, aminocarbonylamino, C_{1-6} alkylaminocarbonylamino, and di(C_{1-6} alkyl)aminocarbonylamino.

In some embodiments, X^3 is N and X^4 is N.

20 In some embodiments, X^3 is CR^3 and X^4 is N.

In some embodiments, X^3 is N and X^4 is CR^4 .

In some embodiments, R^3 is selected from H and C_{1-6} alkyl.

In some embodiments, R^3 is selected from H and methyl.

In some embodiments, R^3 is H.

25 In some embodiments, R^4 is selected from H and C_{1-6} alkyl.

In some embodiments, R^4 is selected from H and methyl.

In some embodiments, R^4 is H.

In some embodiments, Cy^1 is C_{6-10} aryl or 5-10 membered heteroaryl, wherein the C_{6-10} aryl and 5-10 membered heteroaryl are each optionally substituted with 1, 2, 3, or 4 independently selected R^5 substituents.

In some embodiments, Cy^1 is phenyl optionally substituted with 1, 2, 3, or 4 independently selected R^5 substituents; and

each R^5 is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered

heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a5}, SR^{a5}, NHOR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)NR^{c5}(OR^{a5}), C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(O)NR^{c5}R^{d5}, C(=NR^{e5})R^{b5}, C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})R^{b5}, NR^{c5}S(O)NR^{c5}R^{d5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)(=NR^{e5})R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, S(O)₂NR^{c5}R^{d5}, OS(O)(=NR^{e5})R^{b5}, and OS(O)₂R^{b5}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁵ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{5A} substituents.

In some embodiments, Cy¹ is phenyl optionally substituted with 1 or 2 independently selected R⁵ substituents; and

each R⁵ is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and NO₂.

In some embodiments, Cy¹ is cyanophenyl optionally substituted with fluoro.

In some embodiments, Cy¹ is cyanophenyl.

In some embodiments, Cy¹ is 3-benzonitrile.

In some embodiment, Cy¹ is 3-(2-fluorobenzonitrile).

In some embodiments, Cy² is 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, or 4 independently selected R⁶ substituents.

In some embodiments, Cy² is 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, or 4 independently selected R⁶ substituents; and

each R⁶ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a6}, SR^{a6}, NHOR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)NR^{c6}(OR^{a6}), C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, NR^{c6}NR^{c6}R^{d6}, NR^{c6}C(O)R^{b6}, NR^{c6}C(O)OR^{a6}, NR^{c6}C(O)NR^{c6}R^{d6}, C(=NR^{e6})R^{b6}, C(=NR^{e6})NR^{c6}R^{d6}, NR^{c6}C(=NR^{e6})NR^{c6}R^{d6}, NR^{c6}C(=NR^{e6})R^{b6}, NR^{c6}S(O)NR^{c6}R^{d6}, NR^{c6}S(O)R^{b6}, NR^{c6}S(O)₂R^{b6}, NR^{c6}S(O)(=NR^{e6})R^{b6}, NR^{c6}S(O)₂NR^{c6}R^{d6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, S(O)₂R^{b6}, S(O)₂NR^{c6}R^{d6}, OS(O)(=NR^{e6})R^{b6}, and OS(O)₂R^{b6}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10

membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁶ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{6A} substituents,

or, two R⁶ groups together form an oxo group.

5 In some embodiments, Cy² is 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, or 3 independently selected R⁶ substituents; and

each R⁶ is independently selected from halo, C₁₋₆ alkyl, and C₁₋₆ haloalkyl, wherein
10 the C₁₋₆ alkyl is optionally substituted with OH,
or, two R⁶ groups together form an oxo group.

In some embodiments, Cy² is selected from pyrimidinyl, pyrazolyl, pyridinyl, dihydropyridinyl, oxazolyl, quinolinyl, and quinoxalinyl, wherein the pyrimidinyl, pyrazolyl, pyridinyl, dihydropyridinyl, oxazolyl, quinolinyl, and quinoxalinyl are each optionally
15 substituted with 1, 2, or 3 independently selected R⁶ substituents; and

each R⁶ is independently selected from halo, C₁₋₆ alkyl, and C₁₋₆ haloalkyl, wherein the C₁₋₆ alkyl is optionally substituted with OH,
or, two R⁶ groups together form an oxo group.

In some embodiments, Cy² is selected from pyrimidin-4-yl, 1-ethyl-1H-pyrazol-5-yl, pyridin-4-yl, 1-isopropyl-1H-pyrazol-5-yl, 1-methyl-1H-pyrazol-5-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, 3-methylpyridin-4-yl, 4-(hydroxymethyl)-2-methyloxazol-5-yl, 4-methyloxazol-5-yl, oxazol-5-yl, quinolin-6-yl, and quinoxalin-6-yl.

In some embodiments, R¹ is selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, and C₁₋₆ haloalkyl.

25 In some embodiments, R¹ is H.

In some embodiments, R² is selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

In some embodiments, R² is selected from C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-,
35

(5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents; and

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a21}, SR^{a21}, NHOR^{a21}, C(O)R^{b21}, C(O)NR^{c21}R^{d21}, C(O)NR^{c21}(OR^{a21}), C(O)OR^{a21}, OC(O)R^{b21}, OC(O)NR^{c21}R^{d21}, NR^{c21}R^{d21}, S(O)R^{b21}, S(O)NR^{c21}R^{d21}, and S(O)₂R^{b21}.

In some embodiments, R² is phenyl-C₁₋₃ alkyl- or (5-10 membered heteroaryl)-C₁₋₃ alkyl-, wherein the phenyl-C₁₋₃ alkyl- and (5-10 membered heteroaryl)-C₁₋₃ alkyl- are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents; and


each R^{2A} is independently selected from halo, C₁₋₆ alkyl, and C₁₋₆ haloalkyl.

In some embodiments, R² is selected from benzyl, pyridinylmethyl, pyrazolylmethyl, and pyridinylethyl, wherein benzyl, pyridinylmethyl, pyrazolylmethyl, and pyridinylethyl are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents; and

each R^{2A} is independently selected from halo, C₁₋₆ alkyl, and C₁₋₆ haloalkyl.


In some embodiments, R² is selected from pyridin-2-ylmethyl, (1-methyl-1H-pyrazol-3-yl)methyl, (3-fluoropyridin-2-yl)methyl, (6-methylpyridin-2-yl)methyl, 1-(pyridin-2-yl)ethyl, 2-(pyridin-2-yl)ethyl, and 2-fluorobenzyl.

In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is a compound of Formula (II):




or a pharmaceutically acceptable salt thereof, wherein variables R¹, R², Cy¹ and Cy² are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or Formula (II), or a pharmaceutically acceptable salt thereof, is a compound of Formula (IIa):




or a pharmaceutically acceptable salt thereof, wherein variables R^2 , Cy^1 and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or Formula (II), or a pharmaceutically acceptable salt thereof, is a compound of Formula (IIb):




or a pharmaceutically acceptable salt thereof, wherein variables R^2 and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is a compound of Formula (III):




or a pharmaceutically acceptable salt thereof, wherein variables R^1 , R^2 , R^3 , Cy^1 and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or Formula (III), or a pharmaceutically acceptable salt thereof, is a compound of Formula (IIIa):




5 or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^3 , Cy^1 and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or Formula (III), or a pharmaceutically acceptable salt thereof, is a compound of Formula (IIIb):




10 or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^3 and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or a pharmaceutically acceptable salt thereof, is a compound of Formula (IV):




15 or a pharmaceutically acceptable salt thereof, wherein variables R^1 , R^2 , R^4 , Cy^1 and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or Formula (IV), or a pharmaceutically acceptable salt thereof, is a compound of Formula (IVa):




5 or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^4 , Cy^1 and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or Formula (IV), or a pharmaceutically acceptable salt thereof, is a compound of Formula (IVb):



10 or a pharmaceutically acceptable salt thereof, wherein variables R^2 , R^4 , and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments, the compound of Formula (I) or a pharmaceutically acceptable salt thereof, is a compound of Formula (V):



15 or a pharmaceutically acceptable salt thereof,
 wherein n is 0, 1, or 2; and
 variables X^3 , X^4 , R^1 , R^2 , R^5 , and Cy^2 are defined according to the definitions provided herein, such as for compounds of Formula (I).

In some embodiments of the compound of Formula (I), or a pharmaceutically acceptable salt thereof,

X^3 is N or CR^3 ,

X^4 is N or CR^4 ,

5 Cy^1 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy^1 are each optionally substituted with 1, 2, 3, or 4 independently selected R^5 substituents;

Cy^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^6 substituents;

R^1 is selected from H and C_{1-6} alkyl;

R^2 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{6-10} aryl, 15 C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^2 are each optionally substituted with 1, 2, 3, or 4 20 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a21} , SR^{a21} , 25 $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)NR^{c21}(OR^{a21})$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $S(O)R^{b21}$, $S(O)NR^{c21}R^{d21}$, and $S(O)_2R^{b21}$,

each R^{a21} , R^{b21} , R^{c21} and R^{d21} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-; 30

or, any R^{c21} and R^{d21} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group;

R^3 is selected from H, D, halo, OH, CN, NO_2 , C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, 35 C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano- C_{1-3} alkyl, HO- C_{1-3} alkyl, C_{1-3} alkoxy- C_{1-3}

alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃

- 5 alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

R⁴ is selected from H, D, halo, OH, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl,

- 10 C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃
- 15 alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

each R⁵ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl,

- 20 C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a5}, SR^{a5}, NHOR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)NR^{c5}(OR^{a5}), C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, and S(O)₂R^{b5}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆
- 25 alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁵ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{5A} substituents;

each R^{a5}, R^{b5}, R^{c5} and R^{d5} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl,

- 30 C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-,
- 35 , (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-

of R^{a5}, R^{b5}, R^{c5} and R^{d5} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{5A} substituents;

or, any R^{c5} and R^{d5}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{5A} substituents;

each R^{5A} is independently selected from OH, NO₂, CN, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, cyano-C₁₋₆ alkyl, HO-C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, amino, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, thio, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, carbamyl, C₁₋₆ alkylcarbamyl, di(C₁₋₆ alkyl)carbamyl, carboxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxycarbonylamino, C₁₋₆ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₆ alkylaminocarbonyloxy, di(C₁₋₆ alkyl)aminocarbonyloxy, C₁₋₆ alkylsulfonylamino, aminosulfonyl, C₁₋₆ alkylaminosulfonyl, di(C₁₋₆ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₆ alkylaminosulfonylamino, di(C₁₋₆ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₆ alkylaminocarbonylamino, and di(C₁₋₆ alkyl)aminocarbonylamino;

each R⁶ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a6}, SR^{a6}, NHOR^{a6}, C(O)R^{b6}, C(O)NR^{c6}R^{d6}, C(O)NR^{c6}(OR^{a6}), C(O)OR^{a6}, OC(O)R^{b6}, OC(O)NR^{c6}R^{d6}, NR^{c6}R^{d6}, S(O)R^{b6}, S(O)NR^{c6}R^{d6}, and S(O)₂R^{b6}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁶ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{6A} substituents,

or, two R⁶ groups together form an oxo group;

each R^{a6}, R^{b6}, R^{c6} and R^{d6} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-

of R^{a6}, R^{b6}, R^{c6} and R^{d6} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{6A} substituents;

or, any R^{c6} and R^{d6}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{6A} substituents; and

each R^{6A} is independently selected from OH, NO₂, CN, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, cyano-C₁₋₆ alkyl, HO-C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, amino, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, thio, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, carbamyl, C₁₋₆ alkylcarbamyl, di(C₁₋₆ alkyl)carbamyl, carboxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxycarbonylamino, C₁₋₆ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₆ alkylaminocarbonyloxy, di(C₁₋₆ alkyl)aminocarbonyloxy, C₁₋₆ alkylsulfonylamino, aminosulfonyl, C₁₋₆ alkylaminosulfonyl, di(C₁₋₆ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₆ alkylaminosulfonylamino, di(C₁₋₆ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₆ alkylaminocarbonylamino, and di(C₁₋₆ alkyl)aminocarbonylamino.

In some embodiments of the compound of Formula (I), or a pharmaceutically acceptable salt thereof,

X³ is N or CR³;
X⁴ is N or CR⁴;
wherein at least one of X³ and X⁴ is N;
Cy¹ is C₆₋₁₀ aryl optionally substituted with 1, 2, 3, or 4 independently selected R⁵ substituents;

Cy² is selected from 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of Cy² are each optionally substituted with 1, 2, 3, or 4 independently selected R⁶ substituents;

R¹ is selected from H and C₁₋₆ alkyl;

R² is selected from C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R² are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered

heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, and OR^{a21}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2A} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2B} substituents;

R^{a21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl;

each R^{2B} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, and OR^{a22}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2B} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

R^{a22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl;

each R^{2C} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, and OR^{a23}

R^{a23} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl.

R³ is selected from H and C₁₋₆ alkyl;

R⁴ is selected from H and C₁₋₆ alkyl;

each R⁵ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a5}, and NR^{c5}R^{d5};

each R^{a5} , R^{c5} and R^{d5} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

5 or, any R^{c5} and R^{d5} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group;

each R^6 is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a6} , and $NR^{c6}R^{d6}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^6 are each optionally substituted with 1, 2, 3, or 4
15 independently selected R^{6A} substituents,

or, two R^6 groups together form an oxo group;

each R^{a6} , R^{c6} and R^{d6} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

20 or, any R^{c6} and R^{d6} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group; and

each R^{6A} is independently selected from OH, NO_2 , CN, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, cyano- C_{1-6} alkyl, HO- C_{1-6} alkyl, C_{1-6} alkoxy- C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, amino, and C_{1-6} alkylamino.

In some embodiments of the compound of Formula (I), or a pharmaceutically acceptable salt thereof,

X^3 is N or CH;

X^4 is N or CH;

30 wherein at least one of X^3 and X^4 is N;

Cy^1 is phenyl substituted with 1 or 2 independently selected R^5 substituents;

Cy^2 is selected from pyrimidin-4-yl, 1-ethyl-1H-pyrazol-5-yl, pyridin-4-yl, 1-isopropyl-1H-pyrazol-5-yl, 1-methyl-1H-pyrazol-5-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, 3-methylpyridin-4-yl, 4-(hydroxymethyl)-2-methyloxazol-5-yl, 4-methyloxazol-5-yl, oxazol-5-yl, quinolin-6-yl, and quinoxalin-6-yl;

R¹ is H;

R² is selected from pyridin-2-ylmethyl, (1-methyl-1H-pyrazol-3-yl)methyl, (3-fluoropyridin-2-yl)methyl, (6-methylpyridin-2-yl)methyl, 1-(pyridin-2-yl)ethyl, 2-(pyridin-2-yl)ethyl, and 2-fluorobenzyl; and

5 each R⁵ is independently selected from F and CN.

In some embodiments, the compound is the (*S*)-enantiomer of one of the preceding compounds, or a pharmaceutically acceptable salt thereof. In some embodiments, the compound is the (*R*)-enantiomer of one of the preceding compounds, or a pharmaceutically acceptable salt thereof.

10 In some embodiments, 1, 2, 3, 4, 5, 6, 7, or 8 hydrogen atoms, attached to carbon atoms of any “alkyl”, “alkenyl”, “alkynyl”, “aryl”, “phenyl”, “cycloalkyl”, “heterocycloalkyl”, or “heteroaryl” substituents or “-C₁₋₆ alkyl-”, “alkylene”, “alkenylene”, and “alkynylene” linking groups, are each optionally replaced by a deuterium atom.

It is further appreciated that certain features of the invention, which are, for clarity,
15 described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

At various places in the present specification, divalent linking substituents are
20 described. It is specifically intended that each divalent linking substituent include both the forward and backward forms of the linking substituent. For example, -NR(CR'R'')_n- includes both -NR(CR'R'')_n- and -(CR'R'')_nNR-. Where the structure clearly requires a linking group, the Markush variables listed for that group are understood to be linking groups.

The term “n-membered” where n is an integer typically describes the number of ring-
25 forming atoms in a moiety where the number of ring-forming atoms is n. For example, piperidinyl is an example of a 6-membered heterocycloalkyl ring, pyrazolyl is an example of a 5-membered heteroaryl ring, pyridyl is an example of a 6-membered heteroaryl ring, and 1,2,3,4-tetrahydro-naphthalene is an example of a 10-membered cycloalkyl group.

As used herein, the phrase “optionally substituted” means unsubstituted or
30 substituted. The substituents are independently selected, and substitution may be at any chemically accessible position. As used herein, the term “substituted” means that a hydrogen atom is removed and replaced by a substituent. A single divalent substituent, *e.g.*, oxo, can replace two hydrogen atoms. It is to be understood that substitution at a given atom is limited by valency.

As used herein, the phrase “each ‘variable’ is independently selected from” means substantially the same as wherein “at each occurrence ‘variable’ is selected from.”

Throughout the definitions, the term “C_{n-m}” indicates a range which includes the endpoints, wherein n and m are integers and indicate the number of carbons. Examples

5 include C₁₋₃, C₁₋₄, C₁₋₆, and the like.

As used herein, the term “C_{n-m} alkyl”, employed alone or in combination with other terms, refers to a saturated hydrocarbon group that may be straight-chain or branched, having n to m carbons. Examples of alkyl moieties include, but are not limited to, chemical groups such as methyl (Me), ethyl (Et), *n*-propyl (*n*-Pr), isopropyl (iPr), *n*-butyl, *tert*-butyl, isobutyl, *sec*-butyl; higher homologs such as 2-methyl-1-butyl, *n*-pentyl, 3-pentyl, *n*-hexyl, 1,2,2-trimethylpropyl, and the like. In some embodiments, the alkyl group contains from 1 to 6 carbon atoms, from 1 to 4 carbon atoms, from 1 to 3 carbon atoms, or 1 to 2 carbon atoms.

As used herein, “C_{n-m} alkenyl” refers to an alkyl group having one or more double carbon-carbon bonds and having n to m carbons. Example alkenyl groups include, but are not limited to, ethenyl, *n*-propenyl, isopropenyl, *n*-butenyl, *sec*-butenyl, and the like. In some

15 embodiments, the alkenyl moiety contains 2 to 6, 2 to 4, or 2 to 3 carbon atoms.

As used herein, “C_{n-m} alkynyl” refers to an alkyl group having one or more triple carbon-carbon bonds and having n to m carbons. Example alkynyl groups include, but are not limited to, ethynyl, propyn-1-yl, propyn-2-yl, and the like. In some embodiments, the alkynyl moiety contains 2 to 6, 2 to 4, or 2 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkoxy”, employed alone or in combination with other terms, refers to a group of formula -O-alkyl, wherein the alkyl group has n to m carbons. Example alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy (e.g., *n*-propoxy and isopropoxy), butoxy (e.g., *n*-butoxy and *tert*-butoxy), and the like. In some

25 embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “amino” refers to a group of formula -NH₂.

As used herein, the term “aryl,” employed alone or in combination with other terms, refers to an aromatic hydrocarbon group, which may be monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings). The term “C_{n-m} aryl” refers to an aryl group having from n to m ring carbon atoms. Aryl groups include, e.g., phenyl, naphthyl, anthracenyl, phenanthrenyl, indanyl, indenyl, and the like. In some embodiments, aryl groups have from 5 to 10 carbon atoms. In some embodiments, the aryl group is phenyl or naphthyl. In some embodiments, the aryl is phenyl.

As used herein, “halo” refers to F, Cl, Br, or I. In some embodiments, a halo is F, Cl, or Br. In some embodiments, a halo is F or Cl. In some embodiments, a halo is F. In some embodiments, a halo is Cl.

As used herein, “C_{n-m} haloalkoxy” refers to a group of formula –O-haloalkyl having n to m carbon atoms. Example haloalkoxy groups include OCF₃ and OCHF₂. In some
5 to m carbon atoms. Example haloalkoxy groups include OCF₃ and OCHF₂. In some embodiments, the haloalkoxy group is fluorinated only. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} haloalkyl”, employed alone or in combination with other terms, refers to an alkyl group having from one halogen atom to 2s+1 halogen atoms
10 which may be the same or different, where “s” is the number of carbon atoms in the alkyl group, wherein the alkyl group has n to m carbon atoms. In some embodiments, the haloalkyl group is fluorinated only. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms. Example haloalkyl groups include CF₃, C₂F₅, CHF₂, CH₂F, CCl₃, CHCl₂, C₂Cl₅ and the like.

As used herein, the term “thio” refers to a group of formula-SH.

As used herein, the term “carbamyl” refers to a group of formula –C(O)NH₂.

As used herein, the term “carbonyl”, employed alone or in combination with other terms, refers to a –C(O)- group.

As used herein, the term “C_{n-m} alkylamino” refers to a group of formula -NH(alkyl),
20 wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkoxy carbonyl” refers to a group of formula -C(O)O-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkyl carbonyl” refers to a group of formula -C(O)-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkyl carbonylamino” refers to a group of formula -NHC(O)-alkyl, wherein the alkyl group has n to m carbon atoms. In some
30 embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m} alkyl sulfonylamino” refers to a group of formula -NHS(O)₂-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “aminosulfonyl” refers to a group of formula -S(O)₂NH₂.

As used herein, the term “C_{n-m}alkylaminosulfonyl” refers to a group of formula -S(O)₂NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “di(C_{n-m} alkyl)aminosulfonyl” refers to a group of formula -S(O)₂N(alkyl)₂, wherein each alkyl group independently has n to m carbon atoms. In some embodiments, each alkyl group has, independently, 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “aminosulfonylamino” refers to a group of formula -NHS(O)₂NH₂.

As used herein, the term “C_{n-m}alkylaminosulfonylamino” refers to a group of formula -NHS(O)₂NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “di(C_{n-m} alkyl)aminosulfonylamino” refers to a group of formula -NHS(O)₂N(alkyl)₂, wherein each alkyl group independently has n to m carbon atoms. In some embodiments, each alkyl group has, independently, 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “aminocarbonylamino”, employed alone or in combination with other terms, refers to a group of formula -NHC(O)NH₂.

As used herein, the term “C_{n-m}alkylaminocarbonylamino” refers to a group of formula -NHC(O)NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “di(C_{n-m} alkyl)aminocarbonylamino” refers to a group of formula -NHC(O)N(alkyl)₂, wherein each alkyl group independently has n to m carbon atoms. In some embodiments, each alkyl group has, independently, 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m}alkylcarbonyl” refers to a group of formula -C(O)-NH(alkyl), wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m}alkylthio” refers to a group of formula -S-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m}alkylsulfinyl” refers to a group of formula -S(O)-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m}alkylsulfonyl” refers to a group of formula -S(O)₂-alkyl, wherein the alkyl group has n to m carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “cyano-C_{1-n}alkyl” refers to a group of formula -(C_{1-n}alkylene)-CN, wherein the alkyl group has 1 to n carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms, e.g., -(C₁₋₃alkylene)-CN.

As used herein, the term “HO-C_{1-n}alkyl” refers to a group of formula -(C_{1-n}alkylene)-OH, wherein the alkyl group has 1 to n carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms, e.g., -(C₁₋₃alkylene)-OH.

As used herein, the term “C_{1-n}alkoxy-C_{1-n}alkyl” refers to a group of formula -(C_{1-n}alkylene)-O(C_{1-n}alkyl), wherein the alkyl group has 1 to n carbon atoms. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms, e.g., -(C₁₋₆alkylene)-O(C₁₋₆alkyl).

As used herein, the term “carboxy” refers to a group of formula -C(O)OH.

As used herein, the term “di(C_{n-m}-alkyl)amino” refers to a group of formula -N(alkyl)₂, wherein the two alkyl groups each has, independently, n to m carbon atoms. In some embodiments, each alkyl group independently has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “di(C_{n-m}-alkyl)carbonyl” refers to a group of formula -C(O)N(alkyl)₂, wherein the two alkyl groups each has, independently, n to m carbon atoms. In some embodiments, each alkyl group independently has 1 to 6, 1 to 4, or 1 to 3 carbon atoms.

As used herein, the term “C_{n-m}alkylcarbonyloxy” is a group of formula -OC(O)-alkyl, wherein the alkyl group has n to m carbon atoms.

As used herein, “aminocarbonyloxy” is a group of formula -OC(O)-NH₂.

As used herein, “C_{n-m}alkylaminocarbonyloxy” is a group of formula -OC(O)-NH-alkyl, wherein the alkyl group has n to m carbon atoms.

As used herein, “di(C_{n-m}alkyl)aminocarbonyloxy” is a group of formula -OC(O)-N(alkyl)₂, wherein each alkyl group has, independently, n to m carbon atoms.

As used herein “C_{n-m}alkoxycarbonylamino” refers to a group of formula -NHC(O)O(C_{n-m}alkyl), wherein the alkyl group has n to m carbon atoms.

As used herein, “cycloalkyl” refers to non-aromatic cyclic hydrocarbons including cyclized alkyl and alkenyl groups. Cycloalkyl groups can include mono- or polycyclic (e.g., having 2 fused rings) groups, spirocycles, and bridged rings (e.g., a bridged bicycloalkyl group). Ring-forming carbon atoms of a cycloalkyl group can be optionally substituted by oxo or sulfido (e.g., C(O) or C(S)). Also included in the definition of cycloalkyl are moieties that have one or more aromatic rings fused (i.e., having a bond in common with) to the

cycloalkyl ring, for example, benzo or thienyl derivatives of cyclopentane, cyclohexane, and the like. A cycloalkyl group containing a fused aromatic ring can be attached through any ring-forming atom including a ring-forming atom of the fused aromatic ring. Cycloalkyl groups can have 3, 4, 5, 6, 7, 8, 9, or 10 ring-forming carbons (i.e., C₃₋₁₀). In some
 5 embodiments, the cycloalkyl is a C₃₋₁₀ monocyclic or bicyclic cycloalkyl. In some embodiments, the cycloalkyl is a C₃₋₇ monocyclic cycloalkyl. In some embodiments, the cycloalkyl is a C₄₋₇ monocyclic cycloalkyl. In some embodiments, the cycloalkyl is a C₄₋₁₀ spirocycle or bridged cycloalkyl (e.g., a bridged bicycloalkyl group). Example cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl,
 10 cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcamyl, cubane, adamantane, bicyclo[1.1.1]pentyl, bicyclo[2.1.1]hexyl, bicyclo[2.2.1]heptanyl, bicyclo[3.1.1]heptanyl, bicyclo[2.2.2]octanyl, spiro[3.3]heptanyl, and the like. In some embodiments, cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

As used herein, “heteroaryl” refers to a monocyclic or polycyclic (e.g., having 2 fused
 15 rings) aromatic heterocycle having at least one heteroatom ring member selected from N, O, S and B. In some embodiments, the heteroaryl ring has 1, 2, 3, or 4 heteroatom ring members independently selected from N, O, S, and B. In some embodiments, any ring-forming N in a heteroaryl moiety can be an N-oxide. In some embodiments, the heteroaryl is a 5-10
 20 membered monocyclic or bicyclic heteroaryl having 1, 2, 3, or 4 heteroatom ring members independently selected from N, O, S, and B. In some embodiments, the heteroaryl is a 5-10 membered monocyclic or bicyclic heteroaryl having 1, 2, 3, or 4 heteroatom ring members independently selected from N, O, and S. In some embodiments, the heteroaryl is a 5-6 monocyclic heteroaryl having 1 or 2 heteroatom ring members independently selected from
 25 N, O, S, and B. In some embodiments, the heteroaryl is a 5-6 monocyclic heteroaryl having 1 or 2 heteroatom ring members independently selected from N, O, and S. In some embodiments, the heteroaryl group contains 3 to 10, 4 to 10, 5 to 10, 5 to 7, 3 to 7, or 5 to 6 ring-forming atoms. In some embodiments, the heteroaryl group has 1 to 4 ring-forming heteroatoms, 1 to 3 ring-forming heteroatoms, 1 to 2 ring-forming heteroatoms or 1 ring-forming heteroatom. When the heteroaryl group contains more than one heteroatom ring
 30 member, the heteroatoms may be the same or different. Example heteroaryl groups include, but are not limited to, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrrolyl, pyrazolyl, azolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, furyl, thienyl, triazolyl (e.g., 1,2,3-triazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl), tetrazolyl, thiadiazolyl (e.g., 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl), quinolinyl, isoquinolinyl, indolyl, benzothienyl,
 35 benzofuran, benzisoxazole, imidazo[1, 2-b]thiazolyl, purinyl, triazinyl, thieno[3,2-

b]pyridinyl, imidazo[1,2-*a*]pyridinyl, 1,5-naphthyridinyl, 1*H*-pyrazolo[4,3-*b*]pyridinyl, and oxadiazolyl (e.g., 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl).

As used herein, "heterocycloalkyl" refers to monocyclic or polycyclic heterocycles having at least one non-aromatic ring (saturated or partially unsaturated ring), wherein one or more of the ring-forming carbon atoms of the heterocycloalkyl is replaced by a heteroatom selected from N, O, S, and B, and wherein the ring-forming carbon atoms and heteroatoms of a heterocycloalkyl group can be optionally substituted by one or more oxo or sulfido (e.g., C(O), S(O), C(S), or S(O)₂, etc.). Heterocycloalkyl groups include monocyclic and polycyclic (e.g., having 2 fused rings) systems. Included in heterocycloalkyl are monocyclic and polycyclic 3 to 10, 4 to 10, 5 to 10, 4 to 7, 5 to 7, or 5 to 6 membered heterocycloalkyl groups. Heterocycloalkyl groups can also include spirocycles and bridged rings (e.g., a 5 to 10 membered bridged biheterocycloalkyl ring having one or more of the ring-forming carbon atoms replaced by a heteroatom independently selected from N, O, S, and B). The heterocycloalkyl group can be attached through a ring-forming carbon atom or a ring-forming heteroatom. In some embodiments, the heterocycloalkyl group contains 0 to 3 double bonds. In some embodiments, the heterocycloalkyl group contains 0 to 2 double bonds.

Also included in the definition of heterocycloalkyl are moieties that have one or more aromatic rings fused (*i.e.*, having a bond in common with) to the non-aromatic heterocyclic ring, for example, benzo or thienyl derivatives of piperidine, morpholine, azepine, etc. A heterocycloalkyl group containing a fused aromatic ring can be attached through any ring-forming atom including a ring-forming atom of the fused aromatic ring. In some embodiments, the heterocycloalkyl group contains 3 to 10 ring-forming atoms, 4 to 10 ring-forming atoms, 3 to 7 ring-forming atoms, or 5 to 6 ring-forming atoms. In some embodiments, the heterocycloalkyl group has 1 to 4 heteroatoms, 1 to 3 heteroatoms, 1 to 2 heteroatoms or 1 heteroatom. In some embodiments, the heterocycloalkyl is a monocyclic 4 to 6 membered heterocycloalkyl having 1 or 2 heteroatoms independently selected from N, O, S, and B and having one or more oxidized ring members.

In some embodiments, the heterocycloalkyl is a monocyclic or bicyclic 5-10 membered heterocycloalkyl having 1, 2, 3, or 4 heteroatoms independently selected from N, O, S, and B and having one or more oxidized ring members. In some embodiments, the heterocycloalkyl is a monocyclic or bicyclic 5 to 10 membered heterocycloalkyl having 1, 2, 3, or 4 heteroatoms independently selected from N, O, and S and having one or more oxidized ring members. In some embodiments, the heterocycloalkyl is a monocyclic 5 to 6 membered heterocycloalkyl having 1, 2, 3, or 4 heteroatoms independently selected from N, O, and S and having one or more oxidized ring members.

Non-limiting examples of heterocycloalkyl groups include pyrrolidin-2-one, 1,3-isoxazolidin-2-one, pyranyl, tetrahydropyran, oxetanyl, azetidiny, morpholino, thiomorpholino, piperaziny, tetrahydrofuranyl, tetrahydrothienyl, piperidiny, pyrrolidiny, isoxazolidiny, isothiazolidiny, pyrazolidiny, oxazolidiny, thiazolidiny, imidazolidiny, azepanyl, 1,2,3,4-tetrahydroisoquinoline, benzazapene, azabicyclo[3.1.0]hexanyl, diazabicyclo[3.1.0]hexanyl, oxabicyclo[2.1.1]hexanyl, azabicyclo[2.2.1]heptanyl, diazabicyclo[2.2.1]heptanyl, azabicyclo[3.1.1]heptanyl, diazabicyclo[3.1.1]heptanyl, azabicyclo[3.2.1]octanyl, diazabicyclo[3.2.1]octanyl, oxabicyclo[2.2.2]octanyl, azabicyclo[2.2.2]octanyl, azaadamantanyl, diazaadamantanyl, oxa-adamantanyl, azaspiro[3.3]heptanyl, diazaspiro[3.3]heptanyl, oxa-azaspiro[3.3]heptanyl, azaspiro[3.4]octanyl, diazaspiro[3.4]octanyl, oxa-azaspiro[3.4]octanyl, azaspiro[2.5]octanyl, diazaspiro[2.5]octanyl, azaspiro[4.4]nonanyl, diazaspiro[4.4]nonanyl, oxa-azaspiro[4.4]nonanyl, azaspiro[4.5]decanyl, diazaspiro[4.5]decanyl, diazaspiro[4.4]nonanyl, oxa-diazaspiro[4.4]nonanyl and the like. Other examples of heterocycloalkyls include 1,6-dihydropyridiny (e.g., 6-oxo-1,6-dihydropyridiny).

As used herein, “C_{o-p} cycloalkyl-C_{n-m} alkyl-” refers to a group of formula cycloalkyl-alkylene-, wherein the cycloalkyl has o to p carbon atoms and the alkylene linking group has n to m carbon atoms.

As used herein “C_{o-p} aryl-C_{n-m} alkyl-” refers to a group of formula aryl-alkylene-, wherein the aryl has o to p carbon atoms and the alkylene linking group has n to m carbon atoms.

As used herein, “heteroaryl-C_{n-m} alkyl-” refers to a group of formula heteroaryl-alkylene-, wherein alkylene linking group has n to m carbon atoms.

As used herein “heterocycloalkyl-C_{n-m} alkyl-” refers to a group of formula heterocycloalkyl-alkylene-, wherein alkylene linking group has n to m carbon atoms.

As used herein, an “alkyl linking group” is a bivalent straight chain or branched alkyl linking group (“alkylene group”). For example, “C_{o-p} cycloalkyl-C_{n-m} alkyl-”, “C_{o-p} aryl-C_{n-m} alkyl-”, “phenyl-C_{n-m} alkyl-”, “heteroaryl-C_{n-m} alkyl-”, and “heterocycloalkyl-C_{n-m} alkyl-” contain alkyl linking groups. Examples of “alkyl linking groups” or “alkylene groups” include methylene, ethan-1,1-diyl, ethan-1,2-diyl, propan-1,3-diyl, propan-1,2-diyl, propan-1,1-diyl and the like.

At certain places, the definitions or embodiments refer to specific rings (e.g., an azetidine ring, a pyridine ring, etc.). Unless otherwise indicated, these rings can be attached to any ring member provided that the valency of the atom is not exceeded. For example, an

azetidine ring may be attached at any position of the ring, whereas a pyridin-3-yl ring is attached at the 3-position.

As used herein, the term “oxo” refers to an oxygen atom (i.e., =O) as a divalent substituent, forming a carbonyl group when attached to a carbon (e.g., C=O or C(O)), or
5 attached to a nitrogen or sulfur heteroatom forming a nitroso, sulfinyl, or sulfonyl group.

As used herein, the term “independently selected from” means that each occurrence of a variable or substituent, e.g., R^{2A}, are independently selected at each occurrence from the applicable list.

The compounds described herein can be asymmetric (e.g., having one or more
10 stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present disclosure that contain asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically inactive starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many
15 geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present disclosure are described and may be isolated as a mixture of isomers or as separated isomeric forms. In some embodiments, the compound has the (*R*)-configuration. In some embodiments, the
20 compound has the (*S*)-configuration. The Formulas (e.g., Formula (I), (II), etc.) provided herein include stereoisomers of the compounds.

Resolution of racemic mixtures of compounds can be carried out by any of numerous methods known in the art. An example method includes fractional recrystallization using a chiral resolving acid which is an optically active, salt-forming organic acid. Suitable resolving
25 agents for fractional recrystallization methods are, for example, optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or the various optically active camphorsulfonic acids such as β-camphorsulfonic acid. Other resolving agents suitable for fractional crystallization methods include stereoisomerically pure forms of α-methylbenzylamine (e.g., *S* and *R* forms, or
30 diastereomerically pure forms), 2-phenylglycinol, norephedrine, ephedrine, N-methylephedrine, cyclohexylethylamine, 1,2-diaminocyclohexane, and the like.

Resolution of racemic mixtures can also be carried out by elution on a column packed with an optically active resolving agent (e.g., dinitrobenzoylphenylglycine). Suitable elution solvent composition can be determined by one skilled in the art.

Compounds provided herein also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond together with the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Example

5 prototropic tautomers include ketone – enol pairs, amide - imidic acid pairs, lactam – lactim pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, for example, 1H- and 3H-imidazole, 1H-, 2H- and 4H-1,2,4-triazole, 1H- and 2H- isoindole, 2-hydroxypyridine and 2-pyridone, and 1H- and 2H-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by

10 appropriate substitution.

All compounds, and pharmaceutically acceptable salts thereof, can be found together with other substances such as water and solvents (e.g. hydrates and solvates) or can be isolated.

In some embodiments, preparation of compounds can involve the addition of acids or

15 bases to affect, for example, catalysis of a desired reaction or formation of salt forms such as acid addition salts.

In some embodiments, the compounds provided herein, or salts thereof, are substantially isolated. By “substantially isolated” is meant that the compound is at least partially or substantially separated from the environment in which it was formed or detected.

20 Partial separation can include, for example, a composition enriched in the compounds provided herein. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compounds provided herein, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

25 The term “compound” as used herein is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted. Compounds herein identified by name or structure as one particular tautomeric form are intended to include other tautomeric forms unless otherwise specified.

The phrase “pharmaceutically acceptable” is employed herein to refer to those

30 compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The present application also includes pharmaceutically acceptable salts of the

35 compounds described herein. As used herein, “pharmaceutically acceptable salts” refers to

derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

- 5 The pharmaceutically acceptable salts of the present disclosure include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present disclosure can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of
- 10 these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, ethyl acetate, alcohols (e.g., methanol, ethanol, iso-propanol, or butanol) or acetonitrile (ACN) are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and *Journal of Pharmaceutical*
- 15 *Science*, 66, 2 (1977), each of which is incorporated herein by reference in its entirety.

Synthesis

- As will be appreciated by those skilled in the art, the compounds provided herein, including salts and stereoisomers thereof, can be prepared using known organic synthesis
- 20 techniques and can be synthesized according to any of numerous possible synthetic routes, such as those provided in the schemes below.

- The reactions for preparing compounds described herein can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the
- 25 intermediates, or products at the temperatures at which the reactions are carried out, (e.g., temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature). A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.


- 30 The expressions, "ambient temperature" or "room temperature" or "rt" as used herein, are understood in the art, and refer generally to a temperature, e.g., a reaction temperature, that is about the temperature of the room in which the reaction is carried out, for example, a temperature from about 20 °C to about 30 °C.

- Preparation of compounds described herein can involve the protection and
- 35 deprotection of various chemical groups. The need for protection and deprotection, and the

selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., Wiley & Sons, Inc., New York (1999).

Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (e.g., ¹H or ¹³C), infrared spectroscopy, spectrophotometry (e.g., UV-visible), mass spectrometry, or by chromatographic methods such as high performance liquid chromatography (HPLC), liquid chromatography-mass spectroscopy (LCMS), or thin layer chromatography (TLC). Compounds can be purified by those skilled in the art by a variety of methods, including high performance liquid chromatography (HPLC) and normal phase silica chromatography.

Compounds of Formula (I) can be prepared, e.g., using a process as illustrated in the schemes below:




Compounds of formula 1-11 can be prepared *via* the synthetic route outlined in Scheme 1. An appropriate reaction, such as a Mitsunobu reaction, between starting materials 1-1 and 1-2 can be carried out under suitable conditions (such as using DEAD and Ph₃P) to generate intermediate 1-3. A nucleophilic aromatic substitution (S_NAr) reaction of intermediate 1-3 with amine 1-4 (PG is a suitable protecting group, such as 2,4-dimethoxybenzyl) affords compound 1-5. Compound 1-5 can then be cross-coupled with an adduct of formula 1-6, in which M is a boronic acid, boronic ester or an appropriately substituted metal (e.g., M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal), under standard Suzuki cross-coupling conditions (e.g., in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (e.g., in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (e.g., in the presence of a palladium catalyst) to afford the cross-coupling product, which undergoes protecting group removal to generate

intermediate **1-8**. Halogenation of **1-8** with an appropriate reagent, such as *N*-bromosuccinimide (NBS), generates intermediate **1-9**. The final product **1-11** can then be prepared by reacting compound **1-9** with an adduct of formula **1-10** using reaction conditions similar to that described for the preparation of **1-7** from **1-5**. The sequence of the above


5 mentioned synthetic steps can be rearranged, as deemed appropriate, to fit each analogue synthesis.

Scheme 2



Compounds of formula 2-7 can be prepared *via* the synthetic route outlined in Scheme 2. A nucleophilic aromatic substitution (S_NAr) reaction of starting material 2-1 with amine 1-4 (wherein PG is a suitable protecting group, such as 2,4-dimethoxybenzyl) affords compound 2-2. An appropriate reaction, such as a Mitsunobu reaction, between 2-2 and alcohol 1-2 can then be carried out under suitable conditions (such as using DEAD and Ph₃P) to generate intermediate 2-3. Compound 2-3 can be cross-coupled with an adduct of formula 1-6, in which M is a boronic acid, boronic ester or an appropriately substituted metal (*e.g.*, M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal), under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst) to afford the cross-coupling product, which undergoes protecting group removal to generate intermediate 2-5. Halogenation of 2-5 with an appropriate reagent, such as *N*-bromosuccinimide (NBS), generates intermediate 2-6. The final product 2-7 can then be prepared by reacting compound 2-6 with an adduct of formula 1-10 using reaction conditions similar to that described for the preparation of 2-4 from 2-3. The sequence of the above mentioned synthetic steps can be rearranged, as deemed appropriate, to fit each analogue synthesis.

Scheme 3



Compounds of formula **3-8** can be prepared *via* the synthetic route outlined in Scheme 3. A cyclization reaction of starting material **3-1** using appropriate reagents, such as NaNO₂ and HCl, affords bicyclic compound **3-2**. A coupling reaction, such as a Mitsunobu reaction, between **3-2** and alcohol **1-2** can then be carried out under suitable conditions (such as using DEAD and Ph₃P) to generate intermediate **3-3**. A nucleophilic aromatic substitution (S_NAr) reaction of **3-3** with amine **1-4** (PG is a suitable protecting group, such as 2,4-dimethoxybenzyl) affords compound **3-4**. Intermediate **3-4** can then be cross-coupled with an adduct of formula **1-6**, in which M is a boronic acid, boronic ester or an appropriately substituted metal [*e.g.*, M is B(OR)₂, Sn(Alkyl)₃, or Zn-Hal], under standard Suzuki cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst and a suitable base), or standard Stille cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst), or standard Negishi cross-coupling conditions (*e.g.*, in the presence of a palladium catalyst) to afford the cross-coupling product **3-5**, which undergoes protecting group removal to generate **3-6**. Halogenation of **3-6** with an appropriate reagent, such as *N*-bromosuccinimide (NBS), generates compound **3-7**. The final products **3-8** can then be prepared by reacting compound **3-7** with an adduct of formula **1-10** using reaction conditions similar to that described for the preparation of **3-5** from **3-4**. The sequence of the above mentioned synthetic steps can be rearranged, as deemed appropriate, to fit each analogue synthesis.

Methods of Use

The compounds of the present disclosure can modulate the activity of adenosine receptors, such as subtypes A2A and A2B receptors. Accordingly, the compounds, salts or stereoisomers described herein can be used in methods of inhibiting adenosine receptors (e.g.,
5 A2A and/or A2B receptors) by contacting the receptor with any one or more of the compounds, salts, or compositions described herein. In some embodiments, the compounds or salts can be used in methods of inhibiting activity of an adenosine receptor in an individual/patient in need of the inhibition by administering an effective amount of a compound or salt of described herein. In some embodiments, modulating is inhibiting. In
10 some embodiments, the contacting is *in vivo*. In some embodiments, the contacting is *ex vivo* or *in vitro*.

The compounds or salts described herein can be selective. By “selective,” it is meant that the compound binds to or inhibits an adenosine receptor with greater affinity or potency, respectively, compared to at least one other receptor, kinase, etc. The compounds of the
15 present disclosure can also be dual antagonists (i.e., inhibitors) of adenosine receptors, e.g., A2A and A2B adenosine receptors.

Another aspect of the present disclosure pertains to methods of treating an adenosine receptor associated disease or disorder in an individual (e.g., patient) by administering to the individual in need of such treatment a therapeutically effective amount or dose of one or more
20 compounds of the present disclosure or a pharmaceutical composition thereof. An adenosine receptor associated disease or disorder can include any disease, disorder or condition that is directly or indirectly linked to expression or activity of the adenosine receptor, including overexpression and/or abnormal activity levels.

The compounds of the present disclosure are useful in the treatment of diseases
25 related to the activity of adenosine receptors including, for example, cancer, inflammatory diseases, cardiovascular diseases, neurodegenerative diseases, immunomodulatory disorders, central nerve system diseases, and diabetes.

Based on the compelling roles of adenosine, e.g., A2A, A2B, receptors in multiple immunosuppressive mechanisms, developing inhibitors can boost the immune system to
30 suppress tumor progression. Adenosine receptor inhibitors can be used to treat, alone or in combination with other therapies, bladder cancer, lung cancer (e.g., non-small cell lung cancer (NSCLC), lung metastasis), melanoma (e.g., metastatic melanoma), breast cancer, cervical cancer, ovarian cancer, colorectal cancer, pancreatic cancer, esophageal cancer, prostate cancer, kidney cancer, skin cancer, thyroid cancer, liver cancer, uterine cancer, head
35 and neck cancer, and renal cell carcinoma (Antonioli, L. et al., *Nature Reviews Cancer*, 2013,

13, 842-857). *See also*, <https://globenewswire.com/news-release/2017/04/04/954192/0/en/Corvus-Pharmaceuticals-Announces-Interim-Results-from-Ongoing-Phase-I-Ib-Study-Demonstrating-Safety-and-Clinical-Activity-of-Lead-Checkpoint-Inhibitor-CPI-444-in-Patients-with-Adva.html>; Cekic C. et al., *J Immunol*, 2012, 188:198-205; Iannone, R. et al., *Am. J. Cancer Res.* 2014, 4:172-181 (study shows that both A2A and CD73 blockade enhance the antitumor activity of anti-CTLA-4 mAb therapy in a B16F10 murine melanoma model); Iannone, R. et al., *Neoplasia*, 2013, 15:1400-1410 and Beavis PA., et al., *Proc Natl Acad Sci. USA*, 2013, 110:14711-14716 (study shows that A2A and CD73 blockade decreased metastasis in 4T1 breast tumor model with has high CD73 expression). In some embodiments, the prostate cancer is metastatic castrate-resistant prostate carcinoma (mCRPC). In some embodiments, the colorectal cancer is colorectal carcinoma (CRC).

In some embodiments, the disease or disorder is lung cancer (e.g., non-small cell lung cancer), melanoma, pancreatic cancer, breast cancer, head and neck squamous cell carcinoma, prostate cancer, liver cancer, color cancer, endometrial cancer, bladder cancer, skin cancer, cancer of the uterus, renal cancer, gastric cancer, or sarcoma. In some embodiments, the sarcoma is Askin's tumor, sarcoma botryoides, chondrosarcoma, Ewing's sarcoma, malignant hemangioendothelioma, malignant schwannoma, osteosarcoma, alveolar soft part sarcoma, angiosarcoma, cystosarcoma phyllodes, dermatofibrosarcoma protuberans, desmoid tumor, desmoplastic small round cell tumor, epithelioid sarcoma, extraskeletal chondrosarcoma, extraskeletal osteosarcoma, fibrosarcoma, gastrointestinal stromal tumor (GIST), hemangiopericytoma, hemangiosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, lymphosarcoma, malignant peripheral nerve sheath tumor (MPNST), neurofibrosarcoma, rhabdomyosarcoma, synovial sarcoma, or undifferentiated pleomorphic sarcoma.

In some embodiments, the disease or disorder is mesothelioma or adenocarcinoma. In some embodiments, the disease or disorder is mesothelioma. In some embodiments, the disease or disorder is adenocarcinoma.

MDSC (myeloid-derived suppressor cells) are a heterogenous group of immune cells from the myeloid lineage (a family of cells that originate from bone marrow stem cells). MDSCs strongly expand in pathological situations such as chronic infections and cancer, as a result of an altered haematopoiesis. MDSCs are discriminated from other myeloid cell types in which they possess strong immunosuppressive activities rather than immunostimulatory properties. Similar to other myeloid cells, MDSCs interact with other immune cell types including T cells, dendritic cells, macrophages and natural killer cells to regulate their

functions. In some embodiments, the compounds, etc. described herein can be used in methods related to cancer tissue (e.g., tumors) with high infiltration of MDSCs, including Solid tumors with high basal level of macrophage and/or MDSC infiltration.

In some embodiments, the compounds of the disclosure can be used in treating
5 pulmonary inflammation, including bleomycin-induced pulmonary fibrosis and injury related to adenosine deaminase deficiency (Baraldi, et al., *Chem. Rev.*, 2008, 108, 238–263).

In some embodiments, the compounds of the disclosure can be used as a treatment for inflammatory disease such as allergic reactions (e.g., A2B adenosine receptor dependent allergic reactions) and other adenosine receptor dependent immune reactions. Further
10 inflammatory diseases that can be treated by compounds of the disclosure include respiratory disorders, sepsis, reperfusion injury, and thrombosis.

In some embodiments, the compounds of the disclosure can be used as a treatment for cardiovascular disease such as coronary artery disease (myocardial infarction, angina pectoris, heart failure), cerebrovascular disease (stroke, transient ischemic attack), peripheral artery
15 disease, and aortic atherosclerosis and aneurysm. Atherosclerosis is an underlying etiologic factor in many types of cardiovascular disease. Atherosclerosis begins in adolescence with fatty streaks, which progress to plaques in adulthood and finally results in thrombotic events that cause occlusion of vessels leading to clinically significant morbidity and mortality. Antagonists to the A2B adenosine receptor and A2A adenosine receptor may be beneficial in
20 preventing atherosclerotic plaque formation (Eisenstein, A. et al., *J. Cell Physiol.*, 2015, 230(12), 2891-2897).

In some embodiments, the compounds of the disclosure can be used as a treatment for disorders in motor activity; deficiency caused by degeneration of the striatonigral dopamine system; and Parkinson's disease; some of the motivational symptoms of depression (Collins,
25 L. E. et al. *Pharmacol. Biochem. Behav.*, 2012, 100, 498–505.).

In some embodiments, the compounds of the disclosure can be used as a treatment for diabetes and related disorders, such as insulin resistance. Diabetes affects the production of adenosine and the expression of A2B adenosine receptors (A2BRs) that stimulate IL-6 and CRP production, insulin resistance, and the association between A_{2B}R gene single-nucleotide
30 polymorphisms (ADORA2B SNPs) and inflammatory markers. The increased A2BR signaling in diabetes may increase insulin resistance in part by elevating pro-inflammatory mediators. Selective A2BR blockers may be useful to treat insulin resistance (Figler, R. A. et al. *Diabetes*, 2011, 60 (2), 669-679).

It is believed that compounds provided herein, e.g., compounds of Formula (I), or any
35 of the embodiments thereof, may possess satisfactory pharmacological profile and promising

biopharmaceutical properties, such as toxicological profile, metabolism and pharmacokinetic properties, solubility, and permeability. It will be understood that determination of appropriate biopharmaceutical properties is within the knowledge of a person skilled in the art, *e.g.*, determination of cytotoxicity in cells or inhibition of certain targets or channels to
5 determine potential toxicity.

The terms “individual” or “patient”, used interchangeably, refer to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

The phrase “therapeutically effective amount” refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system,
10 animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

As used herein, the term “treating” or “treatment” refers to one or more of (1) inhibiting the disease; *e.g.*, inhibiting a disease, condition or disorder in an individual who is
15 experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, arresting further development of the pathology and/or symptomatology); and (2) ameliorating the disease; *e.g.*, ameliorating a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (*i.e.*, reversing the pathology and/or symptomatology) such as
20 decreasing the severity of disease.

In some embodiments, the compounds of the invention are useful in preventing or reducing the risk of developing any of the diseases referred to herein; *e.g.*, preventing or reducing the risk of developing a disease, condition or disorder in an individual who may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or
25 symptomatology of the disease.

Combination Therapies

I. Immune-checkpoint therapies

In some embodiments, A2A and A2B dual inhibitors provided herein can be used in
30 combination with one or more immune checkpoint inhibitors for the treatment of cancer as described herein. In one embodiment, the combination with one or more immune checkpoint inhibitors as described herein can be used for the treatment of melanoma. Compounds of the present disclosure can be used in combination with one or more immune checkpoint inhibitors. Exemplary immune checkpoint inhibitors include inhibitors against immune
35 checkpoint molecules such as CD20, CD28, CD40, CD122, CD96, CD73, CD47, GITR,

CSF1R, JAK, PI3K delta, PI3K gamma, TAM, arginase, HPK1, CD137 (also known as 4-1BB), ICOS, B7-H3, B7-H4, BTLA, CTLA-4, LAG3, TIM3, VISTA, TIGIT, PD-1, PD-L1 and PD-L2. In some embodiments, the immune checkpoint molecule is a stimulatory checkpoint molecule selected from CD27, CD28, CD40, ICOS, OX40, GITR and CD137. In some embodiments, the immune checkpoint molecule is an inhibitory checkpoint molecule selected from A2AR, B7-H3, B7-H4, BTLA, CTLA-4, IDO, KIR, LAG3, PD-1, TIM3, TIGIT, and VISTA. In some embodiments, the compounds of the disclosure provided herein can be used in combination with one or more agents selected from KIR inhibitors, TIGIT inhibitors, LAIR1 inhibitors, CD160 inhibitors, 2B4 inhibitors and TGFR beta inhibitors.

10 In some embodiments, the A2A and A2B dual inhibitors provided herein can be used in combination with one or more agonists of immune checkpoint molecules, e.g., OX40, CD27, OX40, GITR, and CD137 (also known as 4-1BB).

In some embodiments, the inhibitor of an immune checkpoint molecule is anti-PD1 antibody, anti-PD-L1 antibody, or anti-CTLA-4 antibody.

15 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-1, e.g., an anti-PD-1 monoclonal antibody. In some embodiments, the anti-PD-1 monoclonal antibody is nivolumab, pembrolizumab (also known as MK-3475), durvalumab (Imfinzi®), pidilizumab, SHR-1210, PDR001, MGA012, PDR001, AB122, or AMP-224. In some embodiments, the anti-PD-1 monoclonal antibody is nivolumab or pembrolizumab. In some embodiments, the anti-PD1 antibody is pembrolizumab. In some embodiments, the anti-PD-1 monoclonal antibody is MGA012. In some embodiments, the anti-PD1 antibody is SHR-1210. Other anti-cancer agent(s) include antibody therapeutics such as 4-1BB (e.g. urelumab, utomilumab).

25 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-L1, e.g., an anti-PD-L1 monoclonal antibody. In some embodiments, the anti-PD-L1 monoclonal antibody is BMS-935559, MEDI4736, MPDL3280A (also known as RG7446), or MSB0010718C. In some embodiments, the anti-PD-L1 monoclonal antibody is MPDL3280A or MEDI4736.

30 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-1 and PD-L1, e.g., an anti-PD-1/PD-L1 monoclonal antibody. In some embodiments, the anti-PD-1/PD-L1 is MCLA-136.

In some embodiments, the inhibitor is MCLA-145.

35 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of CTLA-4, e.g., an anti-CTLA-4 antibody. In some embodiments, the anti-CTLA-4 antibody is ipilimumab, tremelimumab, AGEN1884, or CP-675,206.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of LAG3, e.g., an anti-LAG3 antibody. In some embodiments, the anti-LAG3 antibody is BMS-986016, LAG525, or INCAGN2385.

5 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of TIM3, e.g., an anti-TIM3 antibody. In some embodiments, the anti-TIM3 antibody is INCAGN2390, MBG453, or TSR-022.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of GITR, e.g., an anti-GITR antibody. In some embodiments, the anti-GITR antibody is TRX518, MK-4166, INCAGN1876, MK-1248, AMG228, BMS-986156, GWN323, or
10 MEDI1873.

In some embodiments, the inhibitor of an immune checkpoint molecule is an agonist of OX40, e.g., OX40 agonist antibody or OX40L fusion protein. In some embodiments, the anti-OX40 antibody is MEDI0562, MOXR-0916, PF-04518600, GSK3174998, or BMS-986178. In some embodiments, the OX40L fusion protein is MEDI6383.

15 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of CD20, e.g., an anti-CD20 antibody. In some embodiments, the anti-CD20 antibody is obinutuzumab or rituximab.

The compounds of the present disclosure can be used in combination with bispecific antibodies. In some embodiments, one of the domains of the bispecific antibody targets PD-1, PD-L1, CTLA-4, GITR, OX40, TIM3, LAG3, CD137, ICOS, CD3, tumor specific antigens
20 (e.g., CD70) or TGF β receptor.

In some embodiments, the compounds of the disclosure can be used in combination with one or more metabolic enzyme inhibitors. In some embodiments, the metabolic enzyme inhibitor is an inhibitor of IDO1, TDO, or arginase. Examples of IDO1 inhibitors include
25 epacadostat, NLG919, BMS-986205, PF-06840003, IOM2983, RG-70099 and LY338196.

As provided throughout, the additional compounds, inhibitors, agents, etc. can be combined with the present compound in a single or continuous dosage form, or they can be administered simultaneously or sequentially as separate dosage forms.

30 II. Cancer therapies

Cancer cell growth and survival can be impacted by multiple signaling pathways. Thus, it is useful to combine different enzyme/protein/receptor inhibitors, exhibiting different preferences in the targets which they modulate the activities of, to treat such conditions. Targeting more than one signaling pathway (or more than one biological molecule involved in

a given signaling pathway) may reduce the likelihood of drug-resistance arising in a cell population, and/or reduce the toxicity of treatment.

The compounds of the present disclosure can be used in combination with one or more other enzyme/protein/receptor inhibitors or one or more therapies for the treatment of diseases, such as cancer. Examples of diseases and indications treatable with combination therapies include those as described herein.

The compounds of the present disclosure can be used in combination with one or more additional pharmaceutical agents such as, for example, chemotherapeutics, immune-oncology agents, metabolic enzyme inhibitors, chemokine receptor inhibitors, and phosphatase inhibitors, as well as targeted therapies such as Bcr-Abl, Flt-3, EGFR, HER2, JAK, c-MET, VEGFR, PDGFR, c-Kit, IGF-1R, RAF and FAK kinase inhibitors. The one or more additional pharmaceutical agents can be administered to a patient simultaneously or sequentially.

For example, the compounds as disclosed herein can be combined with one or more inhibitors of the following kinases for the treatment of cancer and other diseases or disorders described herein: Akt1, Akt2, Akt3, TGF- β R, PKA, PKG, PKC, CaM-kinase, phosphorylase kinase, MEKK, ERK, MAPK, mTOR, EGFR, HER2, HER3, HER4, INS-R, IGF-1R, IR-R, PDGF α R, PDGF β R, CSF1R, KIT, FLK-II, KDR/FLK-1, FLK-4, flt-1, FGFR1, FGFR2, FGFR3, FGFR4, c-Met, Ron, Sea, TRKA, TRKB, TRKC, FLT3, VEGFR/Flt2, Flt4, EphA1, EphA2, EphA3, EphB2, EphB4, Tie2, Src, Fyn, Lck, Fgr, Btk, Fak, SYK, FRK, JAK, ABL, ALK and B-Raf. Non-limiting examples of inhibitors that can be combined with the compounds of the present disclosure for treatment of cancer and other diseases and disorders described herein include an FGFR inhibitor (FGFR1, FGFR2, FGFR3 or FGFR4, e.g., INCB54828, INCB62079 and INCB63904), a JAK inhibitor (JAK1 and/or JAK2, e.g., ruxolitinib, baricitinib or INCB39110), an IDO inhibitor (e.g., epacadostat, NLG919, or BMS-986205), an LSD1 inhibitor (e.g., INCB59872 and INCB60003), a TDO inhibitor, a PI3K-delta inhibitor (e.g., INCB50797 and INCB50465), a Pim inhibitor, a CSF1R inhibitor, a TAM receptor tyrosine kinases (Tyro-3, Axl, and Mer), a histone deacetylase inhibitor (HDAC) such as an HDAC8 inhibitor, an angiogenesis inhibitor, an interleukin receptor inhibitor, bromo and extra terminal family members inhibitors (for example, bromodomain inhibitors or BET inhibitors such as INCB54329 and INCB57643) and an adenosine receptor antagonist or combinations thereof.

Example antibodies for use in combination therapy include but are not limited to Trastuzumab (e.g. anti-HER2), Ranibizumab (e.g. anti-VEGF-A), Bevacizumab (trade name

Avastin, e.g. anti-VEGF, Panitumumab (e.g. anti-EGFR), Cetuximab (e.g. anti-EGFR), Rituxan (anti-CD20) and antibodies directed to c-MET.

One or more of the following agents may be used in combination with the compounds of the present disclosure and are presented as a non-limiting list: a cytostatic agent, cisplatin,

5 doxorubicin, taxotere, taxol, etoposide, irinotecan, camptostar, topotecan, paclitaxel, docetaxel, epothilones, tamoxifen, 5-fluorouracil, methotrexate, temozolomide, cyclophosphamide, SCH 66336, R115777, L778,123, BMS 214662, IRESSATM(gefitinib), TARCEVATM (erlotinib), antibodies to EGFR, intron, ara-C, adriamycin, cytoxan, gemcitabine, uracil mustard, chlormethine, ifosfamide, melphalan, chlorambucil, pipobroman,

10 triethylenemelamine, triethylenethiophosphoramine, busulfan, carmustine, lomustine, streptozocin, dacarbazine, floxuridine, cytarabine, 6-mercaptopurine, 6-thioguanine, fludarabine phosphate, oxaliplatin, leucovirin, ELOXATINTM (oxaliplatin), pentostatin, vinblastine, vincristine, vindesine, bleomycin, dactinomycin, daunorubicin, doxorubicin, epirubicin, idarubicin, mithramycin, deoxycoformycin, mitomycin-C, L-asparaginase,

15 teniposide 17.alpha.-ethinylestradiol, diethylstilbestrol, testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, testolactone, megestrolacetate, methylprednisolone, methyltestosterone, prednisolone, triamcinolone, chlorotrianisene, hydroxyprogesterone, aminoglutethimide, estramustine, medroxyprogesteroneacetate, leuprolide, flutamide, toremifene, goserelin, carboplatin, hydroxyurea, amsacrine,

20 procarbazine, mitotane, mitoxantrone, levamisole, navelbene, anastrozole, letrozole, capecitabine, reloxafine, droloxafine, hexamethylmelamine, avastin, HERCEPTINTM (trastuzumab), BEXXARTM (tositumomab), VELCADETM (bortezomib), ZEVALINTM (ibritumomab tiuxetan), TRISENOXTM (arsenic trioxide), XELODATM (capecitabine), vinorelbine, porfimer, ERBITUXTM (cetuximab), thiotepa, altretamine, melphalan,

25 trastuzumab, lerozole, fulvestrant, exemestane, ifosfomide, rituximab, C225 (cetuximab), Campath (alemtuzumab), clofarabine, cladribine, aphidicolon, rituxan, sunitinib, dasatinib, tezacitabine, Sml1, fludarabine, pentostatin, triapine, didox, trimidox, amidox, 3-AP, and MDL-101,731.

The compounds of the present disclosure can further be used in combination with

30 other methods of treating cancers, for example by chemotherapy, irradiation therapy, tumortargeted therapy, adjuvant therapy, immunotherapy or surgery. Examples of immunotherapy include cytokine treatment (e.g., interferons, GM-CSF, G-CSF, IL-2), CRS-207 immunotherapy, cancer vaccine, monoclonal antibody, adoptive T cell transfer, Toll receptor agonists, STING agonists, oncolytic virotherapy and immunomodulating small

35 molecules, including thalidomide or JAK1/2 inhibitor and the like. The compounds can be

administered in combination with one or more anti-cancer drugs, such as a chemotherapeutics. Example chemotherapeutics include any of: abarelix, aldesleukin, alemtuzumab, alitretinoin, allopurinol, altretamine, anastrozole, arsenic trioxide, asparaginase, azacitidine, bevacizumab, bexarotene, baricitinib, bleomycin, bortezomib, 5 bortezomib, busulfan intravenous, busulfan oral, calusterone, capecitabine, carboplatin, carmustine, cetuximab, chlorambucil, cisplatin, cladribine, clofarabine, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, dalteparin sodium, daunorubicin, decitabine, denileukin, denileukin diftitox, dexrazoxane, docetaxel, doxorubicin, dromostanolone propionate, eculizumab, epirubicin, erlotinib, estramustine, etoposide phosphate, etoposide, 10 exemestane, fentanyl citrate, filgrastim, floxuridine, fludarabine, fluorouracil, fulvestrant, gefitinib, gemcitabine, gemtuzumab ozogamicin, goserelin acetate, histrelin acetate, ibritumomab tiuxetan, idarubicin, ifosfamide, imatinib mesylate, interferon alfa 2a, irinotecan, lapatinib ditosylate, lenalidomide, letrozole, leucovorin, leuprolide acetate, levamisole, lomustine, meclorethamine, megestrol acetate, melphalan, mercaptopurine, methotrexate, 15 methoxsalen, mitomycin C, mitotane, mitoxantrone, nandrolone phenpropionate, nelarabine, nofetumomab, olaparib, oxaliplatin, paclitaxel, pamidronate, panitumumab, pegaspargase, pegfilgrastim, pemetrexed disodium, pentostatin, pipobroman, plicamycin, procarbazine, quinacrine, rasburicase, rituximab, ruxolitinib, rucaparib, streptozocin, tamoxifen, temozolomide, teniposide, testolactone, thalidomide, thioguanine, thiotepa, topotecan, 20 toremifene, tositumomab, trastuzumab, tretinoin, uracil mustard, valrubicin, vinblastine, vincristine, vinorelbine, vorinostat, niraparib, veliparib, talazoparib, and zoledronate.

Additional examples of chemotherapeutics include proteasome inhibitors (e.g., bortezomib), thalidomide, revlimid, and DNA-damaging agents such as melphalan, doxorubicin, cyclophosphamide, vincristine, etoposide, carmustine, and the like.

25 Example Bcr-Abl inhibitors include imatinib mesylate (GLEEVAC™), nilotinib, dasatinib, bosutinib, and ponatinib, and pharmaceutically acceptable salts. Other example suitable Bcr-Abl inhibitors include the compounds, and pharmaceutically acceptable salts thereof, of the genera and species disclosed in U.S. Pat. No. 5,521,184, WO 04/005281, and U.S. Ser. No. 60/578,491.

30 Example suitable Flt-3 inhibitors include midostaurin, lestaurtinib, linifanib, sunitinib, sunitinib maleate, sorafenib, quizartinib, crenolanib, pacritinib, tandutinib, PLX3397 and ASP2215, and their pharmaceutically acceptable salts. Other example suitable Flt-3 inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 03/037347, WO 03/099771, and WO 04/046120.

Example suitable RAF inhibitors include dabrafenib, sorafenib, and vemurafenib, and their pharmaceutically acceptable salts. Other example suitable RAF inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 00/09495 and WO 05/028444.

5 Example suitable FAK inhibitors include VS-4718, VS-5095, VS-6062, VS-6063, BI853520, and GSK2256098, and their pharmaceutically acceptable salts. Other example suitable FAK inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 04/080980, WO 04/056786, WO 03/024967, WO 01/064655, WO 00/053595, and WO 01/014402.

10 In some embodiments, the compounds of the disclosure can be used in combination with one or more other kinase inhibitors including imatinib, particularly for treating patients resistant to imatinib or other kinase inhibitors.

 In some embodiments, the compounds of the disclosure can be used in combination with a chemotherapeutic in the treatment of cancer, and may improve the treatment response as compared to the response to the chemotherapeutic agent alone, without exacerbation of its toxic effects. In some embodiments, the compounds of the disclosure can be used in combination with a chemotherapeutic provided herein. For example, additional pharmaceutical agents used in the treatment of multiple myeloma, can include, without limitation, melphalan, melphalan plus prednisone [MP], doxorubicin, dexamethasone, and Velcade (bortezomib). Further additional agents used in the treatment of multiple myeloma include Bcr-Abl, Flt-3, RAF and FAK kinase inhibitors. In some embodiments, the agent is an alkylating agent, a proteasome inhibitor, a corticosteroid, or an immunomodulatory agent. Examples of an alkylating agent include cyclophosphamide (CY), melphalan (MEL), and bendamustine. In some embodiments, the proteasome inhibitor is carfilzomib. In some
25 embodiments, the corticosteroid is dexamethasone (DEX). In some embodiments, the immunomodulatory agent is lenalidomide (LEN) or pomalidomide (POM). Additive or synergistic effects are desirable outcomes of combining a PI3K inhibitor of the present disclosure with an additional agent.

 In some embodiments, the compounds of the disclosure can be used in combination
30 with an inhibitor of JAK or PI3K δ .

 The agents can be combined with the present compound in a single or continuous dosage form, or the agents can be administered simultaneously or sequentially as separate dosage forms.

 The compounds of the present disclosure can be used in combination with one or
35 more other inhibitors or one or more therapies for the treatment of infections. Examples of

infections include viral infections, bacterial infections, fungus infections or parasite infections.

In some embodiments, a corticosteroid such as dexamethasone is administered to a patient in combination with the compounds of the disclosure where the dexamethasone is administered intermittently as opposed to continuously.

The compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be combined with another immunogenic agent, such as cancerous cells, purified tumor antigens (including recombinant proteins, peptides, and carbohydrate molecules), cells, and cells transfected with genes encoding immune stimulating cytokines. Non-limiting examples of tumor vaccines that can be used include peptides of melanoma antigens, such as peptides of gp100, MAGE antigens, Trp-2, MARTI and/or tyrosinase, or tumor cells transfected to express the cytokine GM-CSF.

The compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be used in combination with a vaccination protocol for the treatment of cancer. In some embodiments, the tumor cells are transduced to express GM-CSF. In some embodiments, tumor vaccines include the proteins from viruses implicated in human cancers such as Human Papilloma Viruses (HPV), Hepatitis Viruses (HBV and HCV) and Kaposi's Herpes Sarcoma Virus (KHSV). In some embodiments, the compounds of the present disclosure can be used in combination with tumor specific antigen such as heat shock proteins isolated from tumor tissue itself. In some embodiments, the compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be combined with dendritic cells immunization to activate potent anti-tumor responses.

The compounds of the present disclosure can be used in combination with bispecific macrocyclic peptides that target Fe alpha or Fe gamma receptor-expressing effectors cells to tumor cells. The compounds of the present disclosure can also be combined with macrocyclic peptides that activate host immune responsiveness.

In some further embodiments, combinations of the compounds of the disclosure with other therapeutic agents can be administered to a patient prior to, during, and/or after a bone marrow transplant or stem cell transplant. The compounds of the present disclosure can be used in combination with bone marrow transplant for the treatment of a variety of tumors of hematopoietic origin.

The compounds of Formula (I) or any of the formulas as described herein, a compound as recited in any of the claims and described herein, or salts thereof can be used in combination with vaccines, to stimulate the immune response to pathogens, toxins, and self antigens. Examples of pathogens for which this therapeutic approach may be particularly
5 useful, include pathogens for which there is currently no effective vaccine, or pathogens for which conventional vaccines are less than completely effective. These include, but are not limited to, HIV, Hepatitis (A, B, & C), Influenza, Herpes, Giardia, Malaria, Leishmania, Staphylococcus aureus, Pseudomonas Aeruginosa.

Viruses causing infections treatable by methods of the present disclosure include, but
10 are not limit to human papillomavirus, influenza, hepatitis A, B, C or D viruses, adenovirus, poxvirus, herpes simplex viruses, human cytomegalovirus, severe acute respiratory syndrome virus, ebola virus, measles virus, herpes virus (e.g., VZV, HSV-1, HAV-6, HSV-II, and CMV, Epstein Barr virus), flaviviruses, echovirus, rhinovirus, coxsackie virus, cornovirus, respiratory syncytial virus, mumpsvirus, rotavirus, measles virus, rubella virus, parvovirus,
15 vaccinia virus, HTLV virus, dengue virus, papillomavirus, molluscum virus, poliovirus, rabies virus, JC virus and arboviral encephalitis virus.

Pathogenic bacteria causing infections treatable by methods of the disclosure include, but are not limited to, chlamydia, rickettsial bacteria, mycobacteria, staphylococci, streptococci, pneumonococci, meningococci and conococci, klebsiella, proteus, serratia,
20 pseudomonas, legionella, diphtheria, salmonella, bacilli, cholera, tetanus, botulism, anthrax, plague, leptospirosis, and Lyme's disease bacteria.

Pathogenic fungi causing infections treatable by methods of the disclosure include, but are not limited to, Candida (albicans, krusei, glabrata, tropicalis, etc.), Cryptococcus neoformans, Aspergillus (fumigatus, niger, etc.), Genus Mucorales (mucor, absidia,
25 rhizophus), Sporothrix schenkii, Blastomyces dermatitidis, Paracoccidioides brasiliensis, Coccidioides immitis and Histoplasma capsulatum. Pathogenic parasites causing infections treatable by methods of the disclosure include, but are not limited to, Entamoeba histolytica, Balantidium coli, Naegleria fowleri, Acanthamoeba sp., Giardia lamblia, Cryptosporidium sp., Pneumocystis carinii, Plasmodium vivax, Babesia microti, Trypanosoma brucei,
30 Trypanosoma cruzi, Leishmania donovani, Toxoplasma gondi, and Nippostrongylus brasiliensis.

Methods for the safe and effective administration of most of these chemotherapeutic agents are known to those skilled in the art. In addition, their administration is described in the standard literature. For example, the administration of many of the chemotherapeutic
35 agents is described in the "Physicians' Desk Reference" (PDR, e.g., 1996 edition, Medical

Economics Company, Montvale, NJ), the disclosure of which is incorporated herein by reference as if set forth in its entirety.

Pharmaceutical Formulations and Dosage Forms

5 When employed as pharmaceuticals, the compounds of the disclosure can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including transdermal, epidermal, 10 ophthalmic and to mucous membranes including intranasal, vaginal and rectal delivery), pulmonary (*e.g.*, by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral, or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal intramuscular or injection or infusion; or intracranial, *e.g.*, intrathecal or intraventricular, administration. Parenteral administration can 15 be in the form of a single bolus dose, or may be, for example, by a continuous perfusion pump. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

20 This disclosure also includes pharmaceutical compositions which contain, as the active ingredient, the compound of the disclosure or a pharmaceutically acceptable salt thereof, in combination with one or more pharmaceutically acceptable carriers (excipients). In some embodiments, the composition is suitable for topical administration. In making the compositions of the disclosure, the active ingredient is typically mixed with an excipient, 25 diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a 30 liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

 In preparing a formulation, the active compound can be milled to provide the appropriate particle size prior to combining with the other ingredients. If the active compound 35 is substantially insoluble, it can be milled to a particle size of less than 200 mesh. If the active

compound is substantially water soluble, the particle size can be adjusted by milling to provide a substantially uniform distribution in the formulation, *e.g.*, about 40 mesh.

The compounds of the disclosure may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other
5 formulation types. Finely divided (nanoparticulate) preparations of the compounds of the disclosure can be prepared by processes known in the art, *e.g.*, see International App. No. WO 2002/000196.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium
10 silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and
15 flavoring agents. The compositions of the disclosure can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

The compositions can be formulated in a unit dosage form, each dosage containing from about 5 to about 1000 mg (1 g), more usually about 100 to about 500 mg, of the active
20 ingredient. The term “unit dosage forms” refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

In some embodiments, the compositions of the disclosure contain from about 5 to about 50 mg of the active ingredient. One having ordinary skill in the art will appreciate that
25 this embodies compositions containing about 5 to about 10, about 10 to about 15, about 15 to about 20, about 20 to about 25, about 25 to about 30, about 30 to about 35, about 35 to about 40, about 40 to about 45, or about 45 to about 50 mg of the active ingredient.

In some embodiments, the compositions of the disclosure contain from about 50 to about 500 mg of the active ingredient. One having ordinary skill in the art will appreciate that
30 this embodies compositions containing about 50 to about 100, about 100 to about 150, about 150 to about 200, about 200 to about 250, about 250 to about 300, about 350 to about 400, or about 450 to about 500 mg of the active ingredient.

In some embodiments, the compositions of the disclosure contain from about 500 to about 1000 mg of the active ingredient. One having ordinary skill in the art will appreciate
35 that this embodies compositions containing about 500 to about 550, about 550 to about 600,

about 600 to about 650, about 650 to about 700, about 700 to about 750, about 750 to about 800, about 800 to about 850, about 850 to about 900, about 900 to about 950, or about 950 to about 1000 mg of the active ingredient.

Similar dosages may be used of the compounds described herein in the methods and
5 uses of the disclosure.

The active compound can be effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It will be understood, however, that the amount of the compound actually administered will usually be determined by a physician, according to the relevant circumstances, including the condition to be treated, the chosen
10 route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present disclosure. When referring to these
15 preformulation compositions as homogeneous, the active ingredient is typically dispersed evenly throughout the composition so that the composition can be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, about 0.1 to about 1000 mg of the active ingredient of the
20 present disclosure.

The tablets or pills of the present disclosure can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric
25 layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

30 The liquid forms in which the compounds and compositions of the present disclosure can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. In some embodiments, the compositions are administered by the oral or
5 nasal respiratory route for local or systemic effect. Compositions can be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device can be attached to a face mask, tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an appropriate manner.

10 Topical formulations can contain one or more conventional carriers. In some embodiments, ointments can contain water and one or more hydrophobic carriers selected from, for example, liquid paraffin, polyoxyethylene alkyl ether, propylene glycol, white Vaseline, and the like. Carrier compositions of creams can be based on water in combination with glycerol and one or more other components, e.g. glycerinmonostearate, PEG-
15 glycerinmonostearate and cetylstearyl alcohol. Gels can be formulated using isopropyl alcohol and water, suitably in combination with other components such as, for example, glycerol, hydroxyethyl cellulose, and the like. In some embodiments, topical formulations contain at least about 0.1, at least about 0.25, at least about 0.5, at least about 1, at least about 2, or at least about 5 wt % of the compound of the disclosure. The topical formulations can be
20 suitably packaged in tubes of, for example, 100 g which are optionally associated with instructions for the treatment of the select indication, e.g., psoriasis or other skin condition.

The amount of compound or composition administered to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In
25 therapeutic applications, compositions can be administered to a patient already suffering from a disease in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. Effective doses will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the disease, the age, weight and general condition of the patient, and the like.

30 The compositions administered to a patient can be in the form of pharmaceutical compositions described above. These compositions can be sterilized by conventional sterilization techniques, or may be sterile filtered. Aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and
35 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use

of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

The therapeutic dosage of a compound of the present disclosure can vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. The proportion or concentration of a compound of the disclosure in a pharmaceutical composition can vary depending upon a number of factors including dosage, chemical characteristics (e.g., hydrophobicity), and the route of administration. For example, the compounds of the disclosure can be provided in an aqueous physiological buffer solution containing about 0.1 to about 10% w/v of the compound for parenteral administration. Some typical dose ranges are from about 1 µg/kg to about 1 g/kg of body weight per day. In some embodiments, the dose range is from about 0.01 mg/kg to about 100 mg/kg of body weight per day. The dosage is likely to depend on such variables as the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, formulation of the excipient, and its route of administration. Effective doses can be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

The compositions of the disclosure can further include one or more additional pharmaceutical agents such as a chemotherapeutic, steroid, anti-inflammatory compound, or immunosuppressant, examples of which are listed herein.

Labeled Compounds and Assay Methods

Another aspect of the present disclosure relates to labeled compounds of the disclosure (radio-labeled, fluorescent-labeled, etc.) that would be useful not only in imaging techniques but also in assays, both *in vitro* and *in vivo*, for localizing and quantitating A2A and/or A2B receptors in tissue samples, including human, and for identifying A2A and/or A2B antagonists by inhibition binding of a labeled compound. Substitution of one or more of the atoms of the compounds of the present disclosure can also be useful in generating differentiated ADME (Adsorption, Distribution, Metabolism and Excretion.) Accordingly, the present disclosure includes adenosine receptor (e.g., A2A and/or A2B) assays that contain such labeled or substituted compounds.

The present disclosure further includes isotopically-labeled compounds of the disclosure. An “isotopically” or “radio-labeled” compound is a compound of the disclosure where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (i.e.,

naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present disclosure include but are not limited to ^2H (also written as D for deuterium), ^3H (also written as T for tritium), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{18}F , ^{35}S , ^{36}Cl , ^{82}Br , ^{75}Br , ^{76}Br , ^{77}Br , ^{123}I , ^{124}I , ^{125}I and ^{131}I . For example, one or more hydrogen atoms in a compound of the present disclosure can be replaced by deuterium atoms (e.g., one or more hydrogen atoms of a C_{1-6} alkyl group of Formula (I) can be optionally substituted with deuterium atoms, such as $-\text{CD}_3$ being substituted for $-\text{CH}_3$). In some embodiments, alkyl groups in any of the disclosed Formulas, e.g., Formula (I), can be perdeuterated.

One or more constituent atoms of the compounds presented herein can be replaced or substituted with isotopes of the atoms in natural or non-natural abundance. In some embodiments, the compound includes at least one deuterium atom. For example, one or more hydrogen atoms in a compound presented herein can be replaced or substituted by deuterium (e.g., one or more hydrogen atoms of a C_{1-6} alkyl group can be replaced by deuterium atoms, such as $-\text{CD}_3$ being substituted for $-\text{CH}_3$). In some embodiments, the compound includes two or more deuterium atoms. In some embodiments, the compound includes 1-2, 1-3, 1-4, 1-5, or 1-6 deuterium atoms. In some embodiments, all of the hydrogen atoms in a compound can be replaced or substituted by deuterium atoms.

In some embodiments, 1, 2, 3, 4, 5, 6, 7, or 8 hydrogen atoms, attached to carbon atoms of any "alkyl", "alkenyl", "alkynyl", "aryl", "phenyl", "cycloalkyl", "heterocycloalkyl", or "heteroaryl" substituents or " $-\text{C}_{1-6}$ alkyl-", "alkylene", "alkenylene" and "alkynylene" linking groups, as described herein, are each optionally replaced by a deuterium atom.

Synthetic methods for including isotopes into organic compounds are known in the art (Deuterium Labeling in Organic Chemistry by Alan F. Thomas (New York, N.Y., Appleton-Century-Crofts, 1971; The Renaissance of H/D Exchange by Jens Atzrodt, Volker Derdau, Thorsten Fey and Jochen Zimmermann, Angew. Chem. Int. Ed. 2007, 7744-7765; The Organic Chemistry of Isotopic Labelling by James R. Hanson, Royal Society of Chemistry, 2011). Isotopically labeled compounds can be used in various studies such as NMR spectroscopy, metabolism experiments, and/or assays.

Substitution with heavier isotopes, such as deuterium, may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances. (see e.g., A. Kerekes et. al. *J. Med. Chem.* 2011, 54, 201-210; R. Xu et. al. *J. Label Compd. Radiopharm.* 2015, 58, 308-312). In particular, substitution at one or more metabolism sites may afford one or more of the therapeutic advantages.

The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for *in vitro* adenosine receptor labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I or ^{35}S can be useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br or ^{77}Br can be useful.

It is understood that a “radio-labeled” or “labeled compound” is a compound that has incorporated at least one radionuclide. In some embodiments, the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br .

The present disclosure can further include synthetic methods for incorporating radio-isotopes into compounds of the disclosure. Synthetic methods for incorporating radio-isotopes into organic compounds are well known in the art, and an ordinary skill in the art will readily recognize the methods applicable for the compounds of disclosure.

A labeled compound of the disclosure can be used in a screening assay to identify/evaluate compounds. For example, a newly synthesized or identified compound (*i.e.*, test compound) which is labeled can be evaluated for its ability to bind an adenosine receptor by monitoring its concentration variation when contacting with the adenosine receptor, through tracking of the labeling. For example, a test compound (labeled) can be evaluated for its ability to reduce binding of another compound which is known to bind to an adenosine receptor (*i.e.*, standard compound). Accordingly, the ability of a test compound to compete with the standard compound for binding to the adenosine receptor directly correlates to its binding affinity. Conversely, in some other screening assays, the standard compound is labeled and test compounds are unlabeled. Accordingly, the concentration of the labeled standard compound is monitored in order to evaluate the competition between the standard compound and the test compound, and the relative binding affinity of the test compound is thus ascertained.

Kits

The present disclosure also includes pharmaceutical kits useful, for example, in the treatment or prevention of adenosine receptor-associated diseases or disorders (such as, e.g., cancer, an inflammatory disease, a cardiovascular disease, or a neurodegenerative disease) which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound of the disclosure. Such kits can further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as

inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, can also be included in the kit.

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of non-critical parameters which can be changed or modified to yield essentially the same results. The compounds of the Examples have been found to inhibit the activity of an adenosine receptor (e.g., A2A and/or A2B) according to at least one assay described herein.

EXAMPLES


Preparatory LC-MS purifications of some of the compounds prepared were performed on Waters mass directed fractionation systems. The basic equipment setup, protocols, and control software for the operation of these systems have been described in detail in the literature (see *e.g.* "Two-Pump At Column Dilution Configuration for Preparative LC-MS", K. Blom, *J. Combi. Chem.*, 4, 295 (2002); "Optimizing Preparative LC-MS Configurations and Methods for Parallel Synthesis Purification", K. Blom, R. Sparks, J. Doughty, G. Everlof, T. Haque, A. Combs, *J. Combi. Chem.*, 5, 670 (2003); and "Preparative LC-MS Purification: Improved Compound Specific Method Optimization", K. Blom, B. Glass, R. Sparks, A. Combs, *J. Combi. Chem.*, 6, 874-883 (2004)). The compounds separated were typically subjected to analytical liquid chromatography mass spectrometry (LCMS) for purity analysis under the following conditions: Instrument; Agilent 1100 series, LC/MSD, Column: Waters Sunfire™ C₁₈ 5 μm, 2.1 x 50 mm, Buffers: mobile phase A: 0.025% TFA in water and mobile phase B: acetonitrile; gradient 2% to 80% of B in 3 minutes with flow rate 2.0 mL/minute.

Some of the compounds prepared were also separated on a preparative scale by reverse-phase high performance liquid chromatography (RP-HPLC) with MS detector or flash chromatography (silica gel) as indicated in the Examples. Typical preparative reverse-phase high performance liquid chromatography (RP-HPLC) column conditions are as follows:


pH = 2 purifications: Waters Sunfire™ C₁₈ 5 μm, 30 x 100 mm or Waters XBridge™ C₁₈ 5 μm, 30 x 100 mm column, eluting with mobile phase A: 0.1% TFA (trifluoroacetic acid) in water and mobile phase B: acetonitrile; the flow rate was 60 mL/minute, the separating gradient was optimized for each compound using the Compound Specific Method Optimization protocol as described in the literature (see *e.g.* "Preparative LCMS Purification: Improved Compound Specific Method Optimization", K. Blom, B. Glass, R. Sparks, A. Combs, *J. Comb. Chem.*, 6, 874-883 (2004)).

pH = 10 purifications: Waters XBridge™ C₁₈ 5 μm, 30 x 100 mm column, eluting with mobile phase A: 0.1% NH₄OH in water and mobile phase B: acetonitrile; the flow rate was 60 mL/minute, the separating gradient was optimized for each compound using the Compound Specific Method Optimization protocol as described in the literature (see *e.g.* "Preparative LCMS Purification: Improved Compound Specific Method Optimization", K. Blom, B. Glass, R. Sparks, A. Combs, *J. Comb. Chem.*, 6, 874-883 (2004)).

Example 1. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile




Step A. 4,6-dichloro-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridine



5

To a mixture of 4,6-dichloro-1H-pyrazolo[4,3-c]pyridine (1000 mg, 5.32 mmol), pyridin-2-ylmethanol (0.77 mL, 7.98 mmol) and triphenylphosphine (2790 mg, 10.64 mmol) in DCM (10 mL) was added diisopropyl azodicarboxylate (1.57 mL, 7.98 mmol) at 0 °C. The reaction mixture was stirred at 0°C overnight. Direct chromatography on silica gel column
10 afforded the desired product (571 mg, 39%). LC-MS calculated for C₁₂H₉Cl₂N₄: 279.0 (M+H)⁺; found: 279.3 (M+H)⁺.

Step B. 6-chloro-N-(2,4-dimethoxybenzyl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-4-amine




15

To the mixture of 4,6-dichloro-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridine (500 mg, 1.791 mmol), (2,4-dimethoxyphenyl)methanamine (0.30 mL, 1.97 mmol) in 1,4-dioxane (10 mL) was added triethylamine (0.30 mL, 2.15 mmol). The reaction mixture was


stirred at 110 °C overnight. Direct chromatography on silica gel column afforded the desired product (609 mg, 83%). LC-MS calculated for $C_{21}H_{21}ClN_5O_2$: 410.1 (M+H)⁺; found: 410.3 (M+H)⁺.

- 5 *Step C. 3-(4-((2,4-dimethoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile*



- To the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-4-amine (609 mg, 1.486 mmol), (3-cyanophenyl)boronic acid (327 mg, 2.229 mmol) in 1,4-dioxane (10 mL) and water (1 mL) was added cesium carbonate (968 mg, 2.97 mmol). The reaction mixture was de-gassed under N₂ and then chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (117 mg, 0.149 mmol) was added. The mixture was stirred at 120 °C under microwave for 90 minutes. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product. LC-MS calculated for $C_{28}H_{25}N_6O_2$: 477.2 (M+H)⁺; found: 477.3 (M+H)⁺.

- 20 *Step D. 3-(4-amino-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile*




The solution of 3-(4-((2,4-dimethoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile (500 mg, 1.049 mmol) in TFA (5.00 mL) was stirred

at 100 °C for 30 minutes. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO₃ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure.

- 5 The residue was purified on silica gel column to afford the desired product (273 mg, 80%). LC-MS calculated for C₁₉H₁₅N₆: 327.1 (M+H)⁺; found: 327.3 (M+H)⁺.

Step E. 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile




10

The reaction mixture of 3-(4-amino-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile (273 mg, 0.836 mmol) and 1-bromopyrrolidine-2,5-dione (156 mg, 0.878 mmol) in THF (10 mL) was stirred at 0 °C for 30 minutes and quenched with saturated NaHCO₃ aqueous solution. The organic phase was separated, dried over Na₂SO₄, filtered and evaporated under reduced pressure. The resulting residue was purified on silica gel column to afford the desired product (251 mg, 74.0%). LC-MS calculated for C₁₉H₁₄BrN₆: 405.0 (M+H)⁺ and 407.0 (M+H)⁺; found: 405.2 (M+H)⁺ and 407.3 (M+H)⁺.

15

Step F. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile

20




A mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile (25 mg, 0.062 mmol), 4-(tributylstannyl)pyrimidine (34.2 mg, 0.093 mmol), and copper(I) chloride (7.33 mg, 0.074 mmol), lithium chloride (3.14 mg, 0.074 mmol) and tetrakis(triphenylphosphine)palladium(0) (7.13 mg, 6.17 μmol) in THF (1 mL) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction was diluted with

25


methanol and purified with prep-LCMS (pH=2) to give the desired product. LC-MS calculated for $C_{23}H_{17}N_8$: 405.2 (M+H)⁺; found: 405.3 (M+H)⁺.

Example 2. 3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile




To the mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile (20 mg, 0.049 mmol) and 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (21.92 mg, 0.099 mmol) in 1,4-dioxane (2 mL) and water (0.2 mL) was added cesium carbonate (16.08 mg, 0.049 mmol). The resulting mixture was sparged with N_2 for 2 min and then chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (3.88 mg, 4.94 μ mol) was added. The reaction mixture was stirred at 120 °C for 1.5 h and then was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for $C_{24}H_{21}N_8$: 421.2 (M+H)⁺; found: 421.3 (M+H)⁺.

Example 3. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedures described in Example 1, except by using (3-fluoropyridin-2-yl)methanol instead of using (pyridin-2-yl)methanol. LC-MS calculated for $C_{23}H_{16}FN_8$: 423.1 (M+H)⁺; found: 423.3 (M+H)⁺.

Example 4. 3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile



The above title compound was synthesized according to the procedures described in
 5 Example 1, except by using (1-methyl-1H-pyrazol-3-yl)methanol instead of using (pyridin-2-yl)methanol. LC-MS calculated for $C_{22}H_{18}N_9$: 408.2 (M+H)⁺; found: 408.3 (M+H)⁺.


Example 5. 3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile



10


The above title compound was synthesized according to the procedures described in
 Example 1, except by using (6-methylpyridin-2-yl)methanol instead of using (pyridin-2-yl)methanol. LC-MS calculated for $C_{24}H_{19}N_8$: 419.2 (M+H)⁺; found: 419.3 (M+H)⁺.

Example 6. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyridin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedures described in
 Example 2, except by using pyridin-4-ylboronic acid instead of using 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole. LC-MS calculated for $C_{24}H_{18}N_7$: 404.2
 20 (M+H)⁺; found: 404.3 (M+H)⁺.

Example 7. 3-(4-amino-2-(2-fluorobenzyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedures described in
 5 Example 1, except by using (2-fluorophenyl)methanol instead of using (pyridin-2-yl)methanol. LC-MS calculated for $C_{24}H_{17}FN_7$: 422.2 (M+H)⁺; found: 422.3 (M+H)⁺.

Example 8. 3-(7-amino-2-(pyridin-2-ylmethyl)-4-(pyrimidin-4-yl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




10

Step A. 5-chloro-N-(2,4-dimethoxybenzyl)-1H-pyrazolo[3,4-c]pyridin-7-amine



The reaction mixture of 5,7-dichloro-1H-pyrazolo[3,4-c]pyridine (1000 mg, 5.32
 mmol), (2,4-dimethoxyphenyl)methanamine (0.879 mL, 5.85 mmol) and triethylamine (0.890
 15 mL, 6.38 mmol) in 1,4-dioxane (10 mL) was stirred at 110 °C for 3 days. Direct purification
 on silica gel column afforded the desired product (1.35 g, 80%). LC-MS calculated for
 $C_{15}H_{16}ClN_4O_2$: 319.1 (M+H)⁺; found: 319.3 (M+H)⁺.


Step B. 5-chloro-N-(2,4-dimethoxybenzyl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-7-amine



The reaction mixture of 5-chloro-N-(2,4-dimethoxybenzyl)-1H-pyrazolo[3,4-c]pyridin-7-amine (1350 mg, 4.24 mmol), pyridin-2-ylmethanol (555 mg, 5.08 mmol) and triphenylphosphine (2222 mg, 8.47 mmol) in DCM (20 mL) was added diisopropyl azodicarboxylate (1.0 mL, 5.08 mmol) at 0 °C. The reaction was stirred at 0 °C overnight. Direct purification on silica gel column afforded the desired product (535 mg, 31%). LC-MS calculated for $C_{21}H_{21}ClN_5O_2$: 410.1 (M+H)⁺; found: 410.2 (M+H)⁺.

10

Step C. 3-(7-((2,4-dimethoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




To the mixture of 5-chloro-N-(2,4-dimethoxybenzyl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-7-amine (535 mg, 1.305 mmol) and (3-cyanophenyl)boronic acid (384 mg, 2.61 mmol) in 1,4-dioxane (10 mL) and water (1 mL) was added cesium carbonate (851 mg, 2.61 mmol). The mixture was purged with N₂ and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (103 mg, 0.131 mmol) was added. The reaction mixture was stirred at 120 °C under microwave irradiation for 90 minutes. The reaction was quenched with 20 mL

20

of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na_2SO_4 , filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (420 mg, 68%). LC-MS calculated for $\text{C}_{28}\text{H}_{25}\text{N}_6\text{O}_2$:


5 477.2 ($\text{M}+\text{H}$)⁺; found: 477.3 ($\text{M}+\text{H}$)⁺.

Step D. 3-(7-amino-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile



The solution of 3-(7-((2,4-dimethoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (420.3 mg, 0.882 mmol) in TFA (5 mL) was stirred at 100 °C for 30 minutes. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO_3 aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na_2SO_4 , filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (244 mg, 85%). LC-MS calculated for $\text{C}_{19}\text{H}_{15}\text{N}_6$: 327.1 ($\text{M}+\text{H}$)⁺; found: 327.2 ($\text{M}+\text{H}$)⁺.

Step E. 3-(7-amino-4-bromo-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




20

The mixture of 3-(7-amino-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (357 mg, 1.094 mmol) and 1-bromopyrrolidine-2,5-dione (204 mg, 1.149 mmol) in THF (10 mL) was stirred at 0 °C for 1h and then quenched with saturated NaHCO_3 aqueous solution. The organic phase was separated, dried over Na_2SO_4 , filtered and evaporated under reduced pressure. The resulting residue was purified on silica gel column to

25


afford the desired product (314 mg, 71%). LC-MS calculated for $C_{19}H_{14}BrN_6$: 405.0 (M+H)⁺ and 407.0 (M+H)⁺; found: 405.1 (M+H)⁺ and 407.2 (M+H)⁺.

5 *Step F. 3-(7-amino-2-(pyridin-2-ylmethyl)-4-(pyrimidin-4-yl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile*




A mixture of 3-(7-amino-4-bromo-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (40 mg, 0.099 mmol), 4-(tributylstannyl)pyrimidine (109 mg, 0.296 mmol), and copper(I) chloride (11.73 mg, 0.118 mmol), lithium chloride (5.02 mg, 0.118 mmol) and
10 tetrakis(triphenylphosphine)palladium(0) (11.41 mg, 9.87 μ mol) in THF (1 ml) was first purged with N_2 , and then heated and stirred at 90 °C for 2 h. The reaction was diluted with methanol and purified with prep-LCMS (pH 2) to give the desired product. LC-MS calculated for $C_{23}H_{17}N_8$: 405.2 (M+H)⁺; found: 405.3 (M+H)⁺.

15 **Example 9. 3-(7-amino-4-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile**




To the mixture of 3-(7-amino-4-bromo-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (162 mg, 0.400 mmol) and (1-ethyl-1H-pyrazol-5-yl)boronic acid
20 (55.9 mg, 0.400 mmol) 1,4-dioxane (2.0 mL) and water (0.2 mL) was added cesium carbonate (260 mg, 0.799 mmol). The resulting mixture was sparged with N_2 for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (0.040 mmol) was added. The reaction mixture was stirred at 120 °C under microwave irradiation for 90 min. The reaction mixture was diluted with methanol.
25 Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for $C_{24}H_{21}N_8$: 421.2 (M+H)⁺; found: 421.3 (M+H)⁺.

Example 10. 3-(7-amino-2-((6-methylpyridin-2-yl)methyl)-4-(pyrimidin-4-yl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile



The above title compound was synthesized according to the procedures described in
 5 Example 8, except by using (6-methylpyridin-2-yl)methanol instead of using (pyridin-2-yl)methanol. LC-MS calculated for $C_{24}H_{19}N_8$: 419.2 (M+H)⁺; found: 419.3 (M+H)⁺.


Example 11. 3-(7-amino-4-(1-ethyl-1H-pyrazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




10

To the mixture of 3-(7-amino-4-bromo-2-((6-methylpyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (60 mg, 0.143 mmol) and (1-ethyl-1H-pyrazol-5-yl)boronic acid (20.03 mg, 0.143 mmol) in 1,4-dioxane (2.0 mL) and water (0.2 mL) was added cesium carbonate (93 mg, 0.286 mmol). The resulting mixture was sparged with N_2 for
 15 1 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (0.014 mmol) was added. The reaction mixture was stirred at 120 °C under microwave irradiation for 90 min. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC (pH=2) afforded the desired product. LC-MS calculated for $C_{25}H_{23}N_8$: 435.2 (M+H)⁺; found: 435.3 (M+H)⁺.

Example 12. 3-(7-amino-4-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile



Step A. 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole




5

Under N₂ atmosphere, a flask was charged with [Ir(OMe)(1,5-cod)]₂ (33 mg, 0.050 mmol), 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (0.22 mL, 1.50 mmol), and pentane (2.0 mL). The resulting mixture was stirred at room temperature for 10 to 15 min. 4,4'-Di-*tert*-butyl-2,2'-dipyridyl (dtbpy) (26.8 mg, 0.10 mmol) was added and the reaction mixture was stirred for an additional 20 minutes. 4-methyloxazole (83.1 mg, 1.0 mmol) in diethyl ether (2.0 mL) was then added and the resulting mixture was stirred at room temperature until completion. The solvent was removed under reduced pressure, and the crude material was washed with pentane to furnish the desired product. LC-MS calculated for C₁₀H₁₇BNO₃: 210.1 (M+H)⁺; found: 210.1 (M+H)⁺.

15

Step B. 3-(7-amino-4-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




Cesium carbonate (80 mg, 0.247 mmol) was added to the mixture of 3-(7-amino-4-bromo-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (50 mg, 0.123 mmol) and 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (25.8 mg, 0.123 mmol) in 1,4-dioxane (2.0 mL) and water (0.2 mL). The resulting mixture was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-

20

amino-1,1'-biphenyl)]palladium(II)(9.71 mg, 0.012 mmol) was added. The reaction mixture was stirred at 120 °C under microwave irradiation for 90 min. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for C₂₃H₁₈N₇O: 408.2 (M+H)⁺; found: 408.2 (M+H)⁺.


5

Example 13. 3-(7-amino-4-(4-methyloxazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




Cesium carbonate (78 mg, 0.239 mmol) was added to the mixture of 3-(7-amino-4-bromo-2-((6-methylpyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (50 mg, 0.119 mmol), 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (24.93 mg, 0.119 mmol) in 1,4-dioxane (2.0 ml) and water (0.2 ml) and then the reaction was sparged with N₂ for 2 min. After chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (0.012 mmol) was added, the reaction was stirred at 120 °C for 1.5 h under microwave irradiation. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for C₂₄H₂₀N₇O: 422.2 (M+H)⁺; found: 422.2 (M+H)⁺.

Example 14. 3-(7-amino-4-(1-ethyl-1H-pyrazol-5-yl)-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




Step A. 5-chloro-N-(2,4-dimethoxybenzyl)-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-7-amine



To the mixture of 5-chloro-N-(2,4-dimethoxybenzyl)-1H-pyrazolo[3,4-c]pyridin-7-amine (1000 mg, 3.14 mmol), (3-fluoropyridin-2-yl)methanol (0.379 mL, 3.76 mmol), and triphenylphosphine (1646 mg, 6.27 mmol) in DCM (15 mL) was added diisopropyl azodicarboxylate (0.741 mL, 3.76 mmol) at 0 °C. The reaction mixture was stirred at 0 °C overnight. Direct purification on silica gel column afforded the desired product (477 mg, 36%). LC-MS calculated for C₂₁H₂₀ClFN₅O₂: 428.1 (M+H)⁺; found: 428.2 (M+H)⁺.


Step B. 3-(7-((2,4-dimethoxybenzyl)amino)-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile



To the mixture of 5-chloro-N-(2,4-dimethoxybenzyl)-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-7-amine (1150 mg, 2.69 mmol) and (3-cyanophenyl)boronic acid (790 mg, 5.38 mmol) in 1,4-dioxane (10.0 mL) and water (1.00 mL) was added cesium carbonate (1751 mg, 5.38 mmol). The resulting mixture was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (211 mg, 0.269 mmol) was added. The reaction mixture was stirred at 120 °C under microwave irradiation for 90 minutes. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified


on silica gel column to afford the desired product, (647 mg, 49%). LC-MS calculated for $C_{28}H_{24}FN_6O_2$: 495.2 (M+H)⁺; found: 495.2 (M+H)⁺.

Step C. 3-(7-amino-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




The solution of 3-(7-((2,4-dimethoxybenzyl)amino)-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (647.3 mg, 1.309 mmol, 48.7 % yield) in TFA (5 mL) was stirred at 100 °C for 30 min. TFA was evaporated under reduced pressure and then 20 mL of saturated $NaHCO_3$ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na_2SO_4 , filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (374 mg, 83%). LC-MS calculated for $C_{19}H_{14}FN_6$: 345.1 (M+H)⁺; found: 345.3 (M+H)⁺.

Step D. 3-(7-amino-4-bromo-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




The mixture of 3-(7-amino-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (2000 mg, 5.81 mmol) and 1-bromopyrrolidine-2,5-dione (2067 mg, 11.62 mmol) in THF (10 mL) was stirred at 0 °C for 1h. Direct purification on silica gel column afforded the desired product. LC-MS calculated for $C_{19}H_{13}BrFN_6$: 423.0 (M+H)⁺ and 425.0 (M+H)⁺; found: 423.0 (M+H)⁺ and 425.0 (M+H)⁺.

Step E. 3-(7-amino-4-(1-ethyl-1H-pyrazol-5-yl)-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




Cesium carbonate (30.8 mg, 0.095 mmol) was added to the mixture of 3-(7-amino-4-bromo-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (20 mg, 0.047 mmol) and 1-ethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (10.49 mg, 0.047 mmol) in 1,4-dioxane (2.0 mL) and water (2 mL). The reaction was sparged with N₂ for 2 min and then chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (4.73 μmol) was added. The resulting reaction mixture was stirred at 120 °C under microwave irradiation for 1.5 h. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for C₂₄H₂₀FN₈: 439.2 (M+H)⁺; found: 439.3 (M+H)⁺.

Example 15. 3-(7-amino-2-((3-fluoropyridin-2-yl)methyl)-4-(pyrimidin-4-yl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile




The mixture of 3-(7-amino-4-bromo-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile (20 mg, 0.047 mmol), 4-(tributylstannyl)pyrimidine (52.3 mg, 0.142 mmol), copper(I) chloride (5.61 mg, 0.057 mmol), lithium chloride (2.404 mg, 0.057 mmol) and tetrakis(triphenylphosphine)palladium(0) (5.46 mg, 4.73 μmol) in THF (1 ml) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction was diluted with methanol and purified with prep-LCMS (pH=2) to give the final desired product. LC-MS calculated for C₂₃H₁₆FN₈: 423.1 (M+H)⁺; found: 423.3 (M+H)⁺.

Example 16. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




5 *Step A. 4,6-dichloro-3H-[1,2,3]triazolo[4,5-c]pyridine*




A solution of NaNO_2 (3.88 g, 56.2 mmol) in water (3mL) was added to a solution of 2,6-dichloropyridine-3,4-diamine (10 g, 56 mmol) in hydrochloric Acid, 37% (5 mL) at 0 °C. The solution was stirred for 30 min. Water (20 mL) was added and the white precipitate was
 10 filtered, washed with water, and dried to give the desired product. LC-MS calculated for $\text{C}_5\text{H}_3\text{Cl}_2\text{N}_4$: 189.0 (M+H)⁺; found: 189.0 (M+H)⁺.

Step B. 6-chloro-N-(2,4-dimethoxybenzyl)-3H-[1,2,3]triazolo[4,5-c]pyridin-4-amine



15 The mixture of 4,6-dichloro-3H-[1,2,3]triazolo[4,5-c]pyridine (600 mg, 3.17 mmol), (2,4-dimethoxyphenyl)methanamine (0.53mL, 3.49 mmol) and triethylamine (0.53 mL, 3.81 mmol) in 1,4-dioxane (10 mL) was stirred at 110 °C for 3 days. Direct purification on silica gel column afforded the desired product (875 mg, 86%). LC-MS calculated for $\text{C}_{14}\text{H}_{15}\text{ClN}_5\text{O}_2$: 320.1 (M+H)⁺; found: 320.3 (M+H)⁺.


Step C. 6-chloro-N-(2,4-dimethoxybenzyl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine



The mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-3H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (875 mg, 2.74 mmol), pyridin-2-ylmethanol (0.317 mL, 3.28 mmol) and triphenylphosphine (1436 mg, 5.47 mmol) in DCM (20 mL) was added diisopropyl azodicarboxylate (0.647 mL, 3.28 mmol) at 0 °C. The resulting mixture was stirred at 0 °C for 1 h. Direct purification on silica gel column afforded the desired product (375 mg, 33.4 % yield). LC-MS calculated for C₂₀H₂₀ClN₆O₂: 411.1 (M+H)⁺; found: 411.2 (M+H)⁺.

10

Step D. 3-(4-((2,4-dimethoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




To the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (375 mg, 0.913 mmol) and (3-cyanophenyl)boronic acid (268 mg, 1.825 mmol) in 1,4-dioxane (10 mL) and water (1.00 mL) was added cesium carbonate (595 mg, 1.825 mmol). The resulting mixture was purged with N₂ and then chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (71.8 mg, 0.091 mmol) was added. The reaction mixture was stirred at 120 °C under microwave irradiation for 90 min. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried

20

over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (300 mg, 68.9%). LC-MS calculated for C₂₇H₂₄N₇O₂: 478.2 (M+H)⁺; found: 478.3 (M+H)⁺.


5 *Step E. 3-(4-amino-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile*



The solution of 3-(4-((2,4-dimethoxybenzyl)amino)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (300.3 mg, 0.629 mmol) in TFA (5 mL) was stirred at 100 °C for 30 min. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO₃ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (175 mg, 85%). LC-MS calculated for C₁₈H₁₄N₇: 328.1 (M+H)⁺; found: 328.2 (M+H)⁺.

15


Step F. 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The mixture of 3-(4-amino-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (175 mg, 0.535 mmol) and 1-bromopyrrolidine-2,5-dione (100 mg, 0.561 mmol) in THF (10 mL) was stirred at 0 °C for 30 min and then quenched with saturated NaHCO₃ aqueous solution. The organic phase was separated, dried over Na₂SO₄, filtered and evaporated under reduced pressure. The resulting residue was purified on silica gel column to afford the desired product (135 mg, 62.2%). LC-MS calculated for C₁₈H₁₃BrN₇: 406.0 (M+H)⁺ and 408.0 (M+H)⁺; found: 406.1 (M+H)⁺ and 408.2 (M+H)⁺.


25

Step G. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




A mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (182 mg, 0.448 mmol), 4-(tributylstannyl)pyrimidine (496 mg, 1.344 mmol), and copper(I) chloride (53.2 mg, 0.538 mmol), lithium chloride (22.79 mg, 0.538 mmol) and tetrakis(triphenylphosphine)palladium(0) (51.8 mg, 0.045 mmol) in THF (1 ml) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction was diluted with methanol and purified with prep-LCMS (pH=2) to give the desired product. LC-MS calculated for C₂₂H₁₆N₉: 406.2 (M+H)⁺; found: 406.2 (M+H)⁺.

Example 17. 3-(4-amino-7-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Cesium carbonate (30.8 mg, 0.095 mmol) was added to the mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (20 mg, 0.047 mmol) and 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (9.88 mg, 0.047 mmol) in 1,4-dioxane (2 mL) and water (0.2 mL). The reaction was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (3.5 mg, 4.73 μmol) was added. The reaction was stirred at 120 °C under microwave irradiation for 1.5 h. The mixture was diluted with methanol. Direct purification on prep. HPLC (pH=2) afforded the desired product. LC-MS calculated for C₂₂H₁₇N₈O: 409.2 (M+H)⁺; found: 409.2 (M+H)⁺.

Example 18. 3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Cesium carbonate (292 mg, 0.896 mmol) was added to the mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (182 mg, 0.448 mmol) and (1-ethyl-1H-pyrazol-5-yl)boronic acid (62.7 mg, 0.448 mmol) in 1,4-dioxane (2 ml) and water (2 ml). The resulting reaction mixture was sparged with N₂ for 2 min and then chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (0.045 mmol) was added. The reaction mixture was stirred at 120 °C for 1.5 h. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for C₂₃H₂₀N₉: 422.2 (M+H)⁺; found: 422.3 (M+H)⁺.

Example 19. 3-(4-amino-7-(3-methylpyridin-4-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedure described in Example 18, except by using (3-methylpyridin-4-yl)boronic acid instead of using (1-ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for C₂₄H₁₉N₈: 419.2 (M+H)⁺; found: 419.3 (M+H)⁺.

Example 20. 3-(4-amino-7-(4-(hydroxymethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Cesium carbonate (80 mg, 0.246 mmol) was added to a mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (50 mg, 0.123 mmol) and 4-(((tert-butyldimethylsilyl)oxy)methyl)-2-methyloxazol-5-ylboronic acid (66.8 mg, 0.246 mmol) in 1,4-dioxane (2.0 mL) and water (0.2 mL). The reaction was sparged with N₂ for 1 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (0.012 mmol) was added. The reaction was stirred at 120 °C for 1.5 h. TFA (1.0 mL, 12.98 mmol) was added and the reaction mixture was stirred at 100 °C for 60 min. The mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for C₂₃H₁₉N₈O₂: 439.2 (M+H)⁺; found: 439.3 (M+H)⁺.

Example 21. 3-(4-amino-7-(oxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedure described in Example 18, except by using oxazol-5-ylboronic acid instead of using (1-ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for C₂₁H₁₅N₈O: 395.1 (M+H)⁺; found: 395.3 (M+H)⁺.

Example 22. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(quinolin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The above title compound was synthesized according to the procedure described in Example 18, except by using quinolin-6-ylboronic acid instead of using (1-ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for $C_{27}H_{19}N_8$: 455.2 (M+H)⁺; found: 455.3 (M+H)⁺.

Example 23. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(quinoxalin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




10

The above title compound was synthesized according to the procedure described in Example 18, except by using quinoxalin-6-ylboronic acid instead of using (1-ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for $C_{26}H_{18}N_9$: 456.2 (M+H)⁺; found: 456.3 (M+H)⁺.


15

Example 24. 3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




20

Step A 6-chloro-N-(2,4-dimethoxybenzyl)-N-(4-methoxybenzyl)-3H-[1,2,3]triazolo[4,5-c]pyridin-4-amine




The mixture of 4,6-dichloro-3H-[1,2,3]triazolo[4,5-c]pyridine (600 mg, 3.17 mmol), bis(4-methoxybenzyl)amine (899 mg, 3.49 mmol) and triethylamine (531 μ L, 3.81 mmol) in 1,4-dioxane (1.7 mL) was stirred at 110 °C for 3 days. Direct purification on silica gel column afforded the desired product (875 mg, 63%). LC-MS calculated for $C_{22}H_{23}ClN_5O_3$: 440.1 (M+H)⁺; found: 440.3 (M+H)⁺.

Step B. 6-chloro-N,N-bis(4-methoxybenzyl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine




To the mixture of 6-chloro-N,N-bis(4-methoxybenzyl)-3H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (875 mg, 2.135 mmol), (6-methylpyridin-2-yl)methanol (315 mg, 2.56 mmol) and triphenylphosphine (1120 mg, 4.27 mmol) in DCM (20 mL) was added diisopropyl azodicarboxylate (0.504 mL, 2.56 mmol) at 0 °C. The reaction mixture was stirred at 0 °C overnight. Direct purification on silica gel afforded the desired product (353 mg, 32%). LC-MS calculated for $C_{28}H_{28}ClN_6O_2$: 515.2 (M+H)⁺; found: 515.2 (M+H)⁺.

Step C. 3-(4-(bis(4-methoxybenzyl)amino)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



Cesium carbonate (447 mg, 1.371 mmol) was added to the mixture of 6-chloro-N,N-bis(4-methoxybenzyl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (353 mg, 0.685 mmol) and (3-cyanophenyl)boronic acid (201 mg, 1.371 mmol) in 1,4-dioxane (10 mL) and water (1.00 mL). The resulting reaction mixture was sparged with N₂ for 2 min. Chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (53.9 mg, 0.069 mmol) was added. The reaction was stirred at 120 °C under microwave irradiation for 1.5 h. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (280 mg, 70.3%). LC-MS calculated for C₃₅H₃₂N₇O₂: 582.3 (M+H)⁺; found: 582.3 (M+H)⁺.

Step D. 3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The solution of 3-(4-(bis(4-methoxybenzyl)amino)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (280.3 mg, 0.482 mmol) in TFA (5 mL) was stirred at 100 °C for 1h. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO₃ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure.

The residue was purified on silica gel column to afford the desired product (136 mg, 58%).


LC-MS calculated for $C_{19}H_{16}N_7$: 342.1 (M+H)⁺; found: 342.3 (M+H)⁺.

Step E. 3-(4-amino-7-bromo-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The mixture of 3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (136 mg, 0.398 mmol) and 1-bromopyrrolidine-2,5-dione (142 mg, 0.797 mmol) in THF (10 mL) was stirred at 0 °C for 1h. Direct purification on silica gel
10 afforded the desired product (133 mg, 79%). LC-MS calculated for $C_{19}H_{15}BrN_7$: 420.1 (M+H)⁺ and 422.1 (M+H)⁺; found: 420.1 (M+H)⁺ and 422.2 (M+H)⁺.

Step F. 3-(4-amino-7-bromo-2-((6-methylpyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The mixture of 3-(4-amino-7-bromo-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (182 mg, 0.433 mmol), 4-(tributylstannyl)pyrimidine (480 mg, 1.299 mmol), and copper(I) chloride (51.4 mg, 0.520 mmol), lithium chloride (22.03 mg, 0.520 mmol) and
15 tetrakis(triphenylphosphine)palladium(0) (50.0 mg, 0.043 mmol) in THF (1 ml) was first purged with N₂, and then heated and stirred at 90 °C for 2 h. The reaction was diluted with methanol and purified with prep-LCMS (pH=2) to give the desired product. LC-MS
20 calculated for $C_{23}H_{18}N_9$: 420.2 (M+H)⁺; found: 420.3 (M+H)⁺.

Example 25. 3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(pyridin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Cesium carbonate (78 mg, 0.238 mmol) was added to the mixture of 3-(4-amino-7-bromo-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (50 mg, 0.119 mmol) and pyridin-4-ylboronic acid (14.62 mg, 0.119 mmol) in 1,4-dioxane (2.0 mL) and water (0.2 mL). The resulting mixture was sparged with N₂ for 25 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (9.36 mg, 0.012 mmol) was added. The reaction was stirred at 120 °C for 1.5 h under microwave irradiation. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for C₂₄H₁₉N₈: 419.2 (M+H)⁺; found: 419.3 (M+H)⁺.

Example 26. 3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(3-methylpyridin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedure described in Example 25, except by using (3-methylpyridin-4-yl)boronic acid instead of using pyridin-4-ylboronic acid. LC-MS calculated for C₂₅H₂₁N₈: 433.2 (M+H)⁺; found: 433.3 (M+H)⁺.

Example 27. 3-(4-amino-7-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The above title compound was synthesized according to the procedure described in Example 25, except by using (1-methyl-6-oxo-1,6-dihydropyridin-3-yl)boronic acid instead of using pyridin-4-ylboronic acid. LC-MS calculated for $C_{25}H_{21}N_8O$: 449.2 (M+H)⁺; found: 449.3 (M+H)⁺.


Example 28. 3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The above title compound was synthesized according to the procedure described in Example 25, except by using (1-ethyl-1H-pyrazol-5-yl)boronic acid instead of using pyridin-4-ylboronic acid. LC-MS calculated for $C_{24}H_{22}N_9$: 436.2 (M+H)⁺; found: 436.3 (M+H)⁺.

15

Example 29. 3-(4-amino-7-(4-methyloxazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedure described in Example 25, except by using 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-


20

yl)oxazole instead of using pyridin-4-ylboronic acid. LC-MS calculated for $C_{23}H_{19}N_8O$: 423.2 (M+H)⁺; found: 423.3 (M+H)⁺.

Example 30. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Step A. 6-chloro-N-(2,4-dimethoxybenzyl)-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine




To the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-3H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (1000 mg, 3.13 mmol), (3-fluoropyridin-2-yl)methanol (477 mg, 3.75 mmol) and triphenylphosphine (1641 mg, 6.25 mmol) in DCM (1.7 mL) was added diisopropyl azodicarboxylate (739 μ L, 3.75 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 1h. Direct purification on silica gel column afforded the desired product (433 mg, 32%). LC-MS calculated for $C_{20}H_{19}ClFN_6O_2$: 429.1 (M+H)⁺; found: 429.3 (M+H)⁺.

Step B. 3-(4-((2,4-dimethoxybenzyl)amino)-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



Cesium carbonate (658 mg, 2.019 mmol) was added to the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (433 mg, 1.010 mmol) and (3-cyanophenyl)boronic acid (297 mg, 2.019 mmol) in 1,4-dioxane (10.0 mL) and water (1.0 mL). The resulting mixture was sparged with N₂ for 2 min and (SP-4-4)-[2'-Amino[1,1'-biphenyl]-2-yl]chloro[dicyclohexyl][2',4',6'-tris(1-methylethyl)[1,1'-biphenyl]-2-yl]phosphine[palladium] (79 mg, 0.101 mmol) was added. The reaction mixture was stirred at 120 °C for 1.5 h under microwave irradiation. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (357 mg, 71%). LC-MS calculated for C₂₇H₂₃FN₇O₂: 496.2 (M+H)⁺; found: 496.3 (M+H)⁺.

Step C. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The solution of 3-(4-((2,4-dimethoxybenzyl)amino)-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (357.3 mg, 0.721 mmol) in TFA (5 mL) was stirred at 100 °C for 1h. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO₃ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure.

The residue was purified on silica gel column to afford the desired product (213 mg, 61%).


LC-MS m/z calculated for $C_{18}H_{13}FN_7$: 346.1 (M+H)⁺; found: 346.3 (M+H)⁺.

Step D. 3-(4-amino-7-bromo-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The mixture of 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (213 mg, 0.617 mmol) and 1-bromopyrrolidine-2,5-dione (220 mg, 1.234 mmol) in THF (5 mL) was stirred at 0 °C for 1h. Direct purification on silica gel afforded the desired product (175 mg, 67%). LC-MS calculated for $C_{18}H_{12}BrFN_7$: 424.0 (M+H)⁺ and 426.0 (M+H)⁺; found: 424.3 (M+H)⁺ and 426.3 (M+H)⁺.

Step E. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The mixture of 3-(4-amino-7-bromo-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (220 mg, 0.519 mmol), 4-(tributylstannyl)pyrimidine (383 mg, 1.037 mmol), and copper(I) chloride (61.6 mg, 0.622 mmol), lithium chloride (26.4 mg, 0.622 mmol) and tetrakis(triphenylphosphine)palladium(0) (59.9 mg, 0.052 mmol) in THF (1 mL) was first purged with N_2 , and then heated and stirred at 90 °C for 2 h. The reaction was diluted with methanol and purified with prep-LCMS (pH=2) to give the desired product. ¹H NMR (500 MHz, DMSO- d_6) ppm 8.98 (s, 1H), 8.77 (d, J = 5.02 Hz, 1H), 8.38 (dd, J_1 = 4.60 Hz, J_2 = 1.32 Hz, 1H), 7.90-8.30 (bs, 2H), 7.76-7.89 (m, 3H), 7.66 (dd, J_1 = 5.25 Hz, J_2 = 1.25 Hz, 1H), 7.45-7.58 (m, 3H), 6.25 (s, 2H). LC-MS calculated for $C_{22}H_{15}FN_9$: 424.1 (M+H)⁺; found: 424.3 (M+H)⁺.

Example 31. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(pyridin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Cesium carbonate (46.1 mg, 0.141 mmol) was added to the mixture of 3-(4-amino-7-bromo-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (30 mg, 0.071 mmol) and pyridin-4-ylboronic acid (17.38 mg, 0.141 mmol) in 1,4-dioxane (2 mL) and water (0.2 mL). The resulting mixture was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (5.56 mg, 7.07 μmol) was added. The reaction mixture was stirred at 120 °C for 1.5 h under microwave irradiation. The reaction mixture was diluted with methanol. Direct purification on pre. HPLC afforded the desired product. LC-MS calculated for C₂₃H₁₆FN₈: 423.1 (M+H)⁺; found: 423.3 (M+H)⁺.

Example 32. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedure described in Example 31, except by using (1-methyl-6-oxo-1,6-dihydropyridin-3-yl)boronic acid instead of using pyridin-4-ylboronic acid. LC-MS calculated for C₂₄H₁₈FN₈O: 453.2 (M+H)⁺; found: 453.3 (M+H)⁺.

Example 33. 3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The above title compound was synthesized according to the procedure described in
 5 Example 31, except by using (1-ethyl-1H-pyrazol-5-yl)boronic acid instead of using pyridin-4-ylboronic acid. LC-MS calculated for $C_{23}H_{19}FN_9$: 440.2 (M+H)⁺; found: 440.3 (M+H)⁺.

Example 34. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(1-isopropyl-1H-pyrazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




10

The above title compound was synthesized according to the procedure described in
 Example 31, except by using (1-isopropyl-1H-pyrazol-5-yl)boronic acid instead of using
 pyridin-4-ylboronic acid. LC-MS calculated for $C_{24}H_{21}FN_9$: 454.2 (M+H)⁺; found: 454.3
 (M+H)⁺.


15

Example 35. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(1-methyl-1H-pyrazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The above title compound was synthesized according to the procedure described in
 20 Example 31, except by using (1-methyl-1H-pyrazol-5-yl)boronic acid instead of using
 pyridin-4-ylboronic acid. LC-MS calculated for $C_{22}H_{17}FN_9$: 426.2 (M+H)⁺; found: 426.3
 (M+H)⁺.


Example 36. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(4-methyloxazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The above title compound was synthesized according to the procedure described in


- 5 Example 31, except by using 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole instead of using pyridin-4-ylboronic acid. LC-MS calculated for $C_{22}H_{16}FN_8O$: 427.1 (M+H)⁺; found: 427.3 (M+H)⁺.

Example 37. 3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(quinoxalin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The above title compound was synthesized according to the procedure described in Example 31, except by using quinoxalin-6-ylboronic acid instead of using pyridin-4-ylboronic acid. LC-MS calculated for $C_{26}H_{17}FN_9$: 474.2 (M+H)⁺; found: 474.3 (M+H)⁺.

Example 38. 3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile




Step A. 3-(4-amino-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile



To the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (300 mg, 0.730 mmol) and (3-cyano-2-fluorophenyl)boronic acid (241 mg, 1.460 mmol) was added cesium carbonate (476 mg, 1.460 mmol). The reaction was sparged with N₂ for 2 min. The reaction was stirred at 120 °C for 1.5 h. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (260 mg, 72%).


The intermediate obtained from the previous step (260 mg, 0.525 mmol) in TFA (5.0 mL), and the reaction was stirred at 100 °C for 1h. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO₃ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (151 mg, 83%). LC-MS calculated for C₁₈H₁₃FN₇: 346.1 (M+H)⁺; found: 346.3 (M+H)⁺.

Step B. 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile




The mixture of 3-(4-amino-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile (151 mg, 0.437 mmol) and 1-bromopyrrolidine-2,5-dione (86 mg, 0.481 mmol) in THF (10 mL) was stirred at 0 °C for 1h. Direct purification on silica gel afforded the desired product (144 mg, 78%). LC-MS calculated for C₁₈H₁₂BrFN₇: 424.0 (M+H)⁺ and 426.0 (M+H)⁺; found: 424.3 (M+H)⁺ and 426.3 (M+H)⁺.

Step C. 3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile



Cesium carbonate (23.04 mg, 0.071 mmol) was added to the mixture of 3-(4-amino-
 5 7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile
 (30 mg, 0.071 mmol) and (1-ethyl-1H-pyrazol-5-yl)boronic acid (19.79 mg, 0.141 mmol) in
 1,4-dioxane (2.0 ml) and water (0.2 ml). The reaction was sparged with N₂ for 2 min and
 chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-
 biphenyl)]palladium(II) (7.07 μmol) was added. The reaction was stirred at 120 °C for 1.5 h.
 10 Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for
 C₂₃H₁₉FN₉: 440.2 (M+H)⁺; found: 440.3 (M+H)⁺.


Example 39. 3-(4-amino-7-(1-methyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile



15


The above title compound was synthesized according to the procedure described in
 Example 38, except by using (1-methyl-1H-pyrazol-5-yl)boronic acid instead of using (1-
 ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for C₂₂H₁₇FN₉: 426.2 (M+H)⁺; found:
 426.3 (M+H)⁺.

Example 40. 3-(4-amino-7-(1-isopropyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile




The above title compound was synthesized according to the procedure described in
5 Example 38, except by using (1-isopropyl-1H-pyrazol-5-yl)boronic acid instead of using (1-ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for $C_{24}H_{21}FN_9$: 454.2 (M+H)⁺; found: 454.3 (M+H)⁺.


Example 41. 3-(4-amino-7-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile



The above title compound was synthesized according to the procedure described in
Example 38, except by using (4-methyloxazol-5-yl)boronic acid instead of using (1-ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for $C_{22}H_{16}FN_8O$: 427.1 (M+H)⁺; found: 427.3
15 (M+H)⁺.


Example 42. 3-(4-amino-7-(4-(hydroxymethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile



Step A. 4-(((tert-butyl dimethylsilyl)oxy)methyl)-2-methyloxazole

In a round-bottomed flask under N₂, a solution of 2-methyloxazol-4-yl)methanol (5.40 g, 26.8 mmol) in CH₂Cl₂ (56.0 mL) was treated at rt with tert-butylchlorodimethylsilane (8.52 g, 53.7 mmol) followed by imidazole (3.69 g, 53.67 mmol) and the resulting suspension was stirred for 1 h at rt. Water was then added, and the mixture extracted with CH₂Cl₂. The organic layers were dried over Na₂SO₄, filtered, concentrated, and purified via column chromatography (heptane/ethyl acetate) to give the desired product. LC-MS calculated for C₁₁H₂₂NO₂Si: 228.1 (M+H)⁺; found: 228.3 (M+H)⁺.


10

Step B. 4-(((tert-butyl dimethylsilyl)oxy)methyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole

Under N₂ atmosphere, a flask was charged with [Ir(OMe)(1,5-cod)]₂ (13 mg, 0.019 mmol), 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (0.085 mL, 0.58 mmol), and pentane (2.0 mL). The resulting mixture was stirred at room temperature for 10 to 15 min. 4,4'-Di-tert-butyl-2,2'-dipyridyl (dtbpy) (10.5 mg, 0.039 mmol) was added, and the reaction mixture was stirred for additional 20 min. 4-(((tert-butyl dimethylsilyl)oxy)methyl)-2-methyloxazole (83 mg, 0.39 mmol) in diethyl ether (2.0 mL) was then added, and the resulting mixture was stirred at room temperature until completion. Solvent was removed under reduced pressure, and the crude material was washed with pentane to furnish the desired product. LC-MS calculated for C₁₇H₃₃BNO₄Si: 354.2 (M+H)⁺; found: 354.2 (M+H)⁺.

20


Step C. 3-(4-amino-7-(4-(hydroxymethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile



Cesium carbonate (77 mg, 0.236 mmol) was added to the mixture of 3-(4-amino-7-bromo-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile (50 mg, 0.118 mmol) and 4-(((tert-butyldimethylsilyl)oxy)methyl)-2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (83 mg, 0.236 mmol) in 1,4-dioxane (2.0 mL) and Water (0.20 mL). The reaction was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (9.20 mg, 0.012 mmol) was added. The reaction was stirred at 120 °C for 1.5 h.


To the reaction mixture was added TFA (1.0 mL, 12.98 mmol). The reaction mixture was stirred at 100 °C for 1h. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for C₂₃H₁₈FN₃O₂: 457.2 (M+H)⁺; found: 457.3 (M+H)⁺.

Example 43. 3-(4-amino-2-(pyridin-2-ylmethyl)-7-(quinoxalin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile




The title compound was synthesized according to the procedure described in Example 38, except by using quinoxalin-6-ylboronic acid instead of using (1-ethyl-1H-pyrazol-5-yl)boronic acid. LC-MS calculated for C₂₆H₁₇FN₉: 474.2 (M+H)⁺; found: 474.3 (M+H)⁺.

Example 44. 3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Step A 6-chloro-N-(2,4-dimethoxybenzyl)-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine




To the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-3H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (1000 mg, 3.13 mmol), (1-methyl-1H-pyrazol-3-yl)methanol (421 mg, 3.75 mmol) and triphenylphosphine (1641 mg, 6.25 mmol) in DCM (10 mL) was added diisopropyl azodicarboxylate (0.739 mL, 3.75 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 1h. Direct purification on silica gel column afforded the desired product (345 mg, 27%). LC-MS calculated for C₁₉H₂₁ClN₇O₂: 414.1 (M+H)⁺; found: 414.2 (M+H)⁺.

Step B. 3-(4-((2,4-dimethoxybenzyl)amino)-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Cesium carbonate (543 mg, 1.667 mmol) was added to the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (345 mg, 0.834 mmol), (3-cyanophenyl)boronic acid (245 mg, 1.667 mmol) in 1,4-dioxane (10 ml) and water (1.00 ml). The reaction was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (65.6 mg, 0.083 mmol) was added. The reaction was stirred at 120 °C for 1.5 h. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (305 mg, 76%). LC-MS calculated for C₂₆H₂₅N₈O₂: 481.2 (M+H)⁺; found: 481.3 (M+H)⁺.

Step C. 3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The solution of 3-(4-((2,4-dimethoxybenzyl)amino)-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (305.3 mg, 0.635 mmol) in TFA (5 mL) was stirred at 100 °C for 1h. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO₃ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (159 mg, 76%). LC-MS calculated for C₁₇H₁₅N₈: 331.1 (M+H)⁺; found: 331.3 (M+H)⁺.

Step D. 3-(4-amino-7-bromo-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The mixture of 3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (159 mg, 0.481 mmol) and 1-

bromopyrrolidine-2,5-dione (90 mg, 0.505 mmol) in THF (10 mL) was stirred at 0 °C for 1h. Direct purification on silica gel column afforded the desired product (149 mg, 76%). LC-MS calculated for $C_{17}H_{14}BrN_8$: 409.1 (M+H)⁺ and 411.1 (M+H)⁺; found: 409.3 (M+H)⁺ and 411.3 (M+H)⁺.

5

Step E. 3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The mixture of 3-(4-amino-7-bromo-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (220 mg, 0.538 mmol), 4-(tributylstannyl)pyrimidine (50 mg, 0.135 mmol), and copper(I) chloride (63.9 mg, 0.645 mmol), lithium chloride (27.3 mg, 0.645 mmol) and tetrakis(triphenylphosphine)palladium(0) (62.1 mg, 0.054 mmol) in THF (1 ml) was first purged with N₂, and then heated and stirred at 90 °C for 2 h.

The reaction was diluted with methanol and purified with prep-LCMS (pH 2) to give the desired compound. LC-MS calculated for $C_{21}H_{17}N_{10}$: 409.2 (M+H)⁺; found: 409.3 (M+H)⁺.

15

Example 45. 3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-7-(4-methyloxazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




20

Cesium carbonate (159 mg, 0.489 mmol) was added to the mixture of 3-(4-amino-7-bromo-2-((1-methyl-1H-pyrazol-3-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (100 mg, 0.244 mmol) and 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (102 mg, 0.489 mmol) in 1,4-dioxane (2.0 mL) and water (0.2 mL). The reaction was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (17 mg, 0.024


25

mmol) was added. The reaction was stirred at 120 °C for 1.5 h. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for $C_{21}H_{18}N_9O$: 412.2 (M+H)⁺; found: 412.3 (M+H)⁺.

5 **Example 46. 3-(4-amino-7-(4-methyloxazol-5-yl)-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile**



Step A. 6-chloro-N-(2,4-dimethoxybenzyl)-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine




10

The mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-3H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (1000 mg, 3.13 mmol), 1-(pyridin-2-yl)ethan-1-ol (462 mg, 3.75 mmol), triphenylphosphine (1641 mg, 6.25 mmol) and diisopropyl azodicarboxylate (0.739 mL, 3.75 mmol) was stirred at 0 °C for 1h. Direct purification on silica gel column afforded the desired product (359 mg, 27%). LC-MS calculated for $C_{21}H_{22}ClN_6O_2$: 425.1 (M+H)⁺; found: 425.3 (M+H)⁺.


15

Step B. 3-(4-((2,4-dimethoxybenzyl)amino)-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



Cesium carbonate (551 mg, 1.690 mmol) was added to the mixture of 6-chloro-N-(2,4-dimethoxybenzyl)-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-4-amine (359 mg, 0.845 mmol) and (3-cyanophenyl)boronic acid (248 mg, 1.690 mmol) in 1,4-dioxane (10 mL) and Water (1.00 mL). The resulting reaction mixture was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (66.5 mg, 0.084 mmol) was added. The reaction was stirred at 120 °C for 1.5 h. The reaction was quenched with 20 mL of ethyl acetate and 20 mL of water. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure. The residue was purified on silica gel column to afford the desired product (347 mg, 84%). LC-MS calculated for C₂₈H₂₆N₇O₂: 492.2 (M+H)⁺; found: 492.3 (M+H)⁺.


Step C. 3-(4-amino-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



The solution of 3-(4-((2,4-dimethoxybenzyl)amino)-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (347 mg, 0.707 mmol) in TFA (5.00 mL) was stirred at 100 °C for 1h. TFA was evaporated under reduced pressure and then 20 mL of saturated NaHCO₃ aqueous solution and 20 mL of ethyl acetate were added. The organic phase was separated and the aqueous solution was extracted with ethyl acetate twice. The combined extracts were dried over Na₂SO₄, filtered and evaporated under reduced pressure.


The residue was purified on silica gel column to afford the desired product (200 mg, 83%).
LC-MS calculated for $C_{19}H_{16}N_7$: 342.1 (M+H)⁺; found: 342.3 (M+H)⁺.

Step D. 3-(4-amino-7-bromo-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




The mixture of 3-(4-amino-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (200 mg, 0.586 mmol) and 1-bromopyrrolidine-2,5-dione (115 mg, 0.644 mmol) in THF (10 ml) was stirred at 0 °C for 1h. Direct purification on silica gel column
afforded the desired product (197 mg, 80%). LC-MS calculated for $C_{19}H_{15}BrN_7$: 420.1
(M+H)⁺ and 422.1(M+H)⁺; found: 420.3 (M+H)⁺ and 422.3 (M+H)⁺.

Step E. 3-(4-amino-7-(4-methyloxazol-5-yl)-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




Cesium carbonate (132 mg, 0.404 mmol) was added to the mixture of 3-(4-amino-7-bromo-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (85 mg, 0.202 mmol) and 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (85 mg, 0.404 mmol) in 1,4-dioxane (2.0 mL) and water (0.2 mL). The resulting mixture was sparged with N₂ for 2 min and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (0.020 mmol) was added. The reaction was stirred at 120 °C for 1.5 h. The reaction mixture was diluted with methanol. Direct purification on prep. HPLC afforded the desired product. LC-MS calculated for $C_{23}H_{19}N_8O$: 423.2 (M+H)⁺; found: 423.3 (M+H)⁺.

Example 47. 3-(4-amino-2-(1-(pyridin-2-yl)ethyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile




A mixture of 3-(4-amino-7-bromo-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile (85 mg, 0.202 mmol), 4-(tributylstannyl)pyrimidine (50 mg, 0.135 mmol), and copper(I) chloride (24.03 mg, 0.243 mmol), lithium chloride (10.29 mg, 0.243 mmol) and tetrakis(triphenylphosphine)palladium(0) (23.37 mg, 0.020 mmol) in THF (1 ml) was first purged with N₂, and then heated and stirred at 90 C for 2 h. The reaction was diluted with methanol and purified with prep-LCMS (pH=2) to give the desired product. LC-MS
 10 calculated for C₂₃H₁₈N₉: 420.2 (M+H)⁺; found: 420.3 (M+H)⁺.

Example 48. 3-(4-amino-7-(4-methyloxazol-5-yl)-2-(2-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile



15 The above title compound was synthesized according to the procedure described in Example 46, except by using 2-(pyridin-2-yl)ethan-1-ol instead of using 1-(pyridin-2-yl)ethan-1-ol in *step A*. LC-MS calculated for C₂₃H₁₉N₈O: 423.2 (M+H)⁺; found: 423.3 (M+H)⁺.

Example 49. 3-(4-amino-7-(4-methyloxazol-5-yl)-2-(2-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile



The above title compound was synthesized according to the procedure described in Example 48, except by using (2-fluoro-3-cyanophenyl)boronic acid instead of using (3-cyanophenyl)boronic acid. LC-MS calculated for $C_{23}H_{18}FN_8O$: 441.2 (M+H)⁺; found: 441.3 (M+H)⁺.

5

Example A. Adenosine A2A Receptor cyclic AMP GS Assay

Stably transfected HEK-293 cells expressing the human adenosine A2A receptor (Perkin Elmer) are maintained in MEM culture medium with 10% FBS and 400 µg/mL Geneticin (Life Technologies). 18 to 24 hours prior to assay, geneticin is removed from culture. The cisbio cAMP-GS Dynamic kit utilizing the FRET (Fluorescence Resonance Energy Transfer) technology is used to measure cAMP accumulation in the cells. Compounds of the present disclosure at an appropriate concentration are mixed with 10000 cells/well in white 96 well half area plates (Perkin Elmer) for 30 min at RT gently shaking. Agonist, CGS21680 (R&D Technologies) at 4 nM is added to each well for 60 min at room temperature gently shaking. Detection reagents, d2-labeled cAMP (acceptor) and anti-cAMP cryptate (donor) are added to each well for 60 min at room temperature gently shaking. Plates are read on Pherastar (BMG Labtech), fluorescence ratio 665/620 is calculated and EC₅₀ determination is performed by fitting the curve of percent of control versus the log of the compound concentration using GraphPad Prism.

20

Example B. Adenosine A2B Receptor cyclic AMP GS Assay

Stably transfected HEK-293 cells expressing the human adenosine A2B receptor (Perkin Elmer) were maintained in MEM culture medium with 10% FBS and 100 µg/mL Geneticin (Life Technologies). 18 to 24 hours prior to assay, geneticin was removed from culture. The cisbio cAMP-GS Dynamic kit utilizing the FRET (Fluorescence Resonance Energy Transfer) technology was used to measure cAMP accumulation in the cells. Compounds of the present disclosure at an appropriate concentration were mixed with 10000 cells/well in white 96 well half area plates (Perkin Elmer) for 30 min at room temperature gently shaking. Agonist, NECA (R&D Technologies) at 12 nM was added to each well for 60 min at room temperature gently shaking. Detection reagents, d2-labeled cAMP (acceptor) and anti-cAMP cryptate (donor) were added to each well for 60 min at RT gently shaking. Plates were read on Pherastar (BMG Labtech), fluorescence ratio 665/620 was calculated and EC₅₀ determination was performed by fitting the curve of percent of control versus the log of the compound concentration using GraphPad Prism. The EC₅₀ data for the Examples obtained via this method are shown in Table 1.

35

Table 1. The A_{2A}_Ki data and A_{2B}_cAMP_EC₅₀ data are provided below. The symbol “†” indicates A_{2A}_Ki or A_{2B}_cAMP_EC₅₀ ≤ 10 nM, “††” indicates A_{2A}_Ki or A_{2B}_cAMP_EC₅₀ > 10 nM but ≤ 100 nM. “†††” indicates A_{2A}_Ki or A_{2B}_cAMP_EC₅₀ > 100 nM but ≤ 1 μM; and “††††” indicates A_{2A}_Ki or A_{2B}_cAMP_EC₅₀ is greater than 1 μM.

Ex. No.	A _{2A} _Ki (nM)	A _{2B} _cAMP_EC ₅₀ (nM)
1	†	††
2	†	†
3	†	††
4	†	††
5	NA	NA
6	†	†
7	†	†
8	†	†
9	†	†
10	†	†
11	†	††
12	†	††
13	†	††
14	†	†
15	NA	NA
16	†	†
17	†	†
18	†	†
19	†	†
20	†	†
21	†	†
22	†	†
23	†	†††
24	†	†
25	†	†
26	†	†
27	†	††
28	†	†

29	†	†
30	†	†
31	†	†
32	†	††
33	†	†
34	†	†
35	†	†
36	†	†
37	†	††
38	†	†
39	†	†
40	†	†
41	†	†
42	†	††
43	†	†††
44	†	†
45	†	†
46	†	†
47	†	†††
48	†	††
49	†	††

Example C. A2A Tag-lite® HTRF Assay

Assays were conducted in black low volume 384-well polystyrene plates (Greiner 784076-25) in a final volume of 10 μ L. Test compounds were first serially diluted in DMSO and 100 nl added to the plate wells before the addition of other reaction components. The final concentration of DMSO was 1%. Tag-lite® Adenosine A2A labeled cells (CisBio C1TT1A2A) were diluted 1:5 into Tag-lite buffer (CisBio LABMED) and spun 1200 g for 5 mins. The pellet was resuspended at a volume 10.4 X the initial cell suspension volume in Tag-lite buffer, and Adenosine A2A Receptor Red antagonist fluorescent ligand (CisBio L0058RED) added at 12.5 nM final concentration. 10 μ L of the cell and ligand mix was added to the assay wells and incubated at room temperature for 45 minutes before reading on a PHERAstar FS plate reader (BMG Labtech) with HTRF 337/620/665 optical module. Percent binding of the fluorescent ligand was calculated; where 100 nM of A2A antagonist control

ZM 241385 (Tocris 1036) displaces the ligand 100% and 1% DMSO has 0% displacement. The % binding data versus the log of the inhibitor concentration was fitted to a one-site competitive binding model (GraphPad Prism version 7.02) where the ligand constant = 12.5 nM and the ligand K_d = 1.85 nM. The K_i data for the Examples obtained via this method are shown in **Table 1** (See Example B).

Example D. A2B Filter Binding Assay

Assays are conducted in deep well polypropylene plates (Greiner 786201) in a final volume of 550 μ L. Test compounds are first serially diluted in DMSO and 5.5 μ L is then added to the plate wells before the addition of other reaction components. The final concentration of DMSO is 3%. HEK293 cell membranes overexpressing the human adenosine receptor A2B (Perkin Elmer ES-113-M400UA) are diluted to 40 μ g/mL in 50 mM HEPES pH 7.0, 5 mM $MgCl_2$, 1 mM EDTA (Assay buffer). [3H] 8-cyclopentyl-1,3-dipropylxanthine (Perkin Elmer NET974001MC) is diluted in assay buffer + 22% DMSO to 24.2 nM, and then further diluted to 1 nM by addition to the diluted membranes. 545 μ L of the membrane and ligand mix is added to the assay wells and incubated on a shaker at room temperature for 1 hour. The membrane mix is then filtered over a UniFilter GF/C filter plate (Perkin Elmer 6005174) pre-soaked in 50 mM HEPES pH 6.5, 5 mM $MgCl_2$, 1 mM EDTA 0.5% BSA and then washed with 5 mL ice cold 50 mM HEPES pH 6.5, 5 mM $MgCl_2$, 1 mM EDTA 0.2% BSA. 50 μ L MicroScintTM cocktail (Perkin Elmer 6013621) is added and plates are read on a Topcount NXT FS (Perkin Elmer). Percent binding of the [3H] ligand is calculated, where 1000 nM of LUF 5834 (Tocris 4603) control displaces the ligand 100% and 3% DMSO has 0% displacement. The % binding data versus the log of the inhibitor concentration is fitted to a one-site competitive binding model (GraphPad Prism version 7.02) where the ligand constant = 2 nM and the ligand K_d = 13 nM.

Example E. A1 and A3 SPA Binding Assays

Both assays are conducted in white 384-well polystyrene plates (Greiner 781075) in a final volume of 50 μ L. Inhibitors are first serially diluted in DMSO and 100 nL is added to the plate wells before the addition of other reaction components. The final concentration of DMSO is 2%.

Wheatgerm agglutinin-coated yttrium silicate SPA beads (Perkin Elmer RPNQ0023) and CHO-K1 cell membranes overexpressing each human adenosine receptor are incubated in 50 mM HEPES pH 7.0, 5 mM $MgCl_2$, 1 mM EDTA (Assay buffer) on a rotary stirrer for 2 hours at 4 $^{\circ}$ C. The beads are pelleted by centrifugation at 6000 g for one minute, and then the

supernatant with unbound membrane is discarded. The beads are re-suspended to the original volume in assay buffer. Each radioligand is diluted in assay buffer + 22% DMSO at 12.2X the final concentration, and then added to the SPA bead suspension. 50 μ L of the SPA bead reaction mix is added to the assay wells and the plates shaken at 600 rpm for 1 hour at room temperature. The beads are then allowed to settle for 1 hour before reading on a Topcount NXT FS (Perkin Elmer). Percent binding of the radiolabeled ligand is calculated, where a control at >100X K_i displaces the ligand 100% and 2% DMSO has 0% displacement. The % binding data versus the log of the inhibitor concentration is fitted to a one-site competitive binding model (GraphPad Prism version 7.02). Assay conditions are provided in Table A below.


Table A.

Assay Component	A1	A3
SPA beads in Hepes buffer	3 mg/mL	1.25 mg/mL
Membrane	60 μ g/mL Perkin Elmer ES-010	20 μ g/mL Perkin Elmer ES-012
Radioligand	1 nM [^{3} H] DP-CPX (Perkin Elmer NET974) K_D = 1nM	0.1 nM [^{125} I] MECA (Perkin Elmer NEX312) K_D = 0.8nM
Control	1 μ M DPCPX (Tocris 0439)	0.1 μ M IB-MECA (Tocris 1066)

Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. Each reference, including all patent, patent applications, and publications, cited in the present application is incorporated herein by reference in its entirety.

What is claimed is:

1. A compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

X^3 is N or CR^3 ;

X^4 is N or CR^4 ;

Cy^1 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy^1 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^5 substituents;

Cy^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl of Cy^2 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^6 substituents;

R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $C(=NR^{e1})R^{b1}$, $C(=NR^{e1})NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl of R^1 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

each R^{a1} , R^{b1} , R^{c1} , and R^{d1} is independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, and C_{2-6} alkynyl of R^{a1} , R^{b1} , R^{c1} , and R^{d1} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{1A} substituents;

each R^{e1} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, and C_{1-6} haloalkoxy;

each R^{1A} is independently selected from OH, CN, halo, C_{1-3} alkyl, C_{2-3} alkenyl, C_{2-3} alkynyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, C_{1-3} haloalkoxy, amino, C_{1-3} alkylamino, and $di(C_{1-3}$ alkyl)amino;

R^2 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $C(=NR^{e2})R^{b2}$, $C(=NR^{e2})NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, and $S(O)_2NR^{c2}R^{d2}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^2 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

each R^{a2} , R^{b2} , R^{c2} , and R^{d2} is independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a2} , R^{b2} , R^{c2} , and R^{d2} , are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

or, any R^{c2} and R^{d2} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2A} substituents;

each R^{e2} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)NR^{c21}(OR^{a21})$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $NR^{c21}NR^{c21}R^{d21}$, $NR^{c21}C(O)R^{b21}$, $NR^{c21}C(O)OR^{a21}$, $NR^{c21}C(O)NR^{c21}R^{d21}$, $C(=NR^{e21})R^{b21}$, $C(=NR^{e21})NR^{c21}R^{d21}$, $NR^{c21}C(=NR^{e21})NR^{c21}R^{d21}$, $NR^{c21}C(=NR^{e21})R^{b21}$, $NR^{c21}S(O)NR^{c21}R^{d21}$, $NR^{c21}S(O)R^{b21}$, $NR^{c21}S(O)_2R^{b21}$, $NR^{c21}S(O)(=NR^{e21})R^{b21}$, $NR^{c21}S(O)_2NR^{c21}R^{d21}$, $S(O)R^{b21}$, $S(O)NR^{c21}R^{d21}$, $S(O)_2R^{b21}$, $S(O)_2NR^{c21}R^{d21}$, $OS(O)(=NR^{e21})R^{b21}$, $OS(O)_2R^{b21}$, SF_3 , $P(O)R^{f21}R^{g21}$, $OP(O)(OR^{h21})(OR^{i21})$, $P(O)(OR^{h21})(OR^{i21})$, and $BR^{j21}R^{k21}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10}

aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2A} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2B} substituents;

each R^{a21}, R^{b21}, R^{c21} and R^{d21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a21}, R^{b21}, R^{c21} and R^{d21} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2B} substituents;

or, any R^{c21} and R^{d21}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{2B} substituents;

each R^{e21} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f21} and R^{g21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h21} and Rⁱ²¹ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j21} and R^{k21} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j21} and R^{k21} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{2B} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered

heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a22}, SR^{a22}, NHOR^{a22}, C(O)R^{b22}, C(O)NR^{c22}R^{d22}, C(O)NR^{c22}(OR^{a22}), C(O)OR^{a22}, OC(O)R^{b22}, OC(O)NR^{c22}R^{d22}, NR^{c22}R^{d22}, NR^{c22}NR^{c22}R^{d22}, NR^{c22}C(O)R^{b22}, NR^{c22}C(O)OR^{a22}, NR^{c22}C(O)NR^{c22}R^{d22}, C(=NR^{e22})R^{b22}, C(=NR^{e22})NR^{c22}R^{d22}, NR^{c22}C(=NR^{e22})NR^{c22}R^{d22}, NR^{c22}C(=NR^{e22})R^{b22}, NR^{c22}S(O)NR^{c22}R^{d22}, NR^{c22}S(O)R^{b22}, NR^{c22}S(O)₂R^{b22}, NR^{c22}S(O)(=NR^{e22})R^{b22}, NR^{c22}S(O)₂NR^{c22}R^{d22}, S(O)R^{b22}, S(O)NR^{c22}R^{d22}, S(O)₂R^{b22}, S(O)₂NR^{c22}R^{d22}, OS(O)(=NR^{e22})R^{b22}, OS(O)₂R^{b22}, SF₅, P(O)R^{f22}R^{g22}, OP(O)(OR^{h22})(ORⁱ²²), P(O)(OR^{h22})(ORⁱ²²), and BR^{j22}R^{k22}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2B} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

each R^{a22}, R^{b22}, R^{c22} and R^{d22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a22}, R^{b22}, R^{c22} and R^{d22} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

or, any R^{c22} and R^{d22}, attached to the same N atom, together with the N atom to which they are attached, form a 4-7 membered heterocycloalkyl group, wherein the 4-7 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

each R^{e22} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f22} and R^{g22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h22} and Rⁱ²² is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered

heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j22} and R^{k22} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j22} and R^{k22} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{2C} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a23}, SR^{a23}, NHOR^{a23}, C(O)R^{b23}, C(O)NR^{c23}R^{d23}, C(O)NR^{c23}(OR^{a23}), C(O)OR^{a23}, OC(O)R^{b23}, OC(O)NR^{c23}R^{d23}, NR^{c23}R^{b23}, NR^{c23}NR^{c23}R^{d23}, NR^{c23}C(O)R^{b23}, NR^{c23}C(O)OR^{a23}, NR^{c23}C(O)NR^{c23}R^{d23}, C(=NR^{e23})R^{b23}, C(=NR^{e23})NR^{c23}R^{d23}, NR^{c23}C(=NR^{e23})NR^{c23}R^{d23}, NR^{c23}C(=NR^{e23})R^{b23}, NR^{c23}S(O)NR^{c23}R^{d23}, NR^{c23}S(O)R^{b23}, NR^{c23}S(O)₂R^{b23}, NR^{c23}S(O)(=NR^{e23})R^{b23}, NR^{c23}S(O)₂NR^{c23}R^{d23}, S(O)R^{b23}, S(O)NR^{c23}R^{d23}, S(O)₂R^{b23}, S(O)₂NR^{c23}R^{d23}, OS(O)(=NR^{e23})R^{b23}, OS(O)₂R^{b23}, SF₅, P(O)R^{f23}R^{g23}, OP(O)(OR^{h23})(ORⁱ²³), P(O)(OR^{h23})(ORⁱ²³), and BR^{j23}R^{k23}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2C} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2D} substituents;

each R^{a23}, R^{b23}, R^{c23} and R^{d23} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a23}, R^{b23}, R^{c23} and R^{d23} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2D} substituents;

or, any R^{c23} and R^{d23}, attached to the same N atom, together with the N atom to which they are attached, form a 4-7 membered heterocycloalkyl group, wherein the 4-7 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2D} substituents;

each R^{e23} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{f23} and R^{g23} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h23} and R^{i23} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j23} and R^{k23} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j23} and R^{k23} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl;

each R^{2D} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, (4-7 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a24} , SR^{a24} , $NHOR^{a24}$, $C(O)R^{b24}$, $C(O)NR^{c24}R^{d24}$, $C(O)NR^{c24}(OR^{a24})$, $C(O)OR^{a24}$, $OC(O)R^{b24}$, $OC(O)NR^{c24}R^{d24}$, $NR^{c24}R^{d24}$, $NR^{c24}NR^{c24}R^{d24}$, $NR^{c24}C(O)R^{b24}$, $NR^{c24}C(O)OR^{a24}$, $NR^{c24}C(O)NR^{c24}R^{d24}$, $C(=NR^{e24})R^{b24}$, $C(=NR^{e24})NR^{c24}R^{d24}$, $NR^{c24}C(=NR^{e24})NR^{c24}R^{d24}$, $NR^{c24}C(=NR^{e24})R^{b24}$, $NR^{c24}S(O)NR^{c24}R^{d24}$, $NR^{c24}S(O)R^{b24}$, $NR^{c24}S(O)_2R^{b24}$, $NR^{c24}S(O)(=NR^{e24})R^{b24}$, $NR^{c24}S(O)_2NR^{c24}R^{d24}$, $S(O)R^{b24}$, $S(O)NR^{c24}R^{d24}$, $S(O)_2R^{b24}$, $S(O)_2NR^{c24}R^{d24}$, $OS(O)(=NR^{e24})R^{b24}$, $OS(O)_2R^{b24}$, SF_5 , $P(O)R^{f24}R^{g24}$, $OP(O)(OR^{h24})(OR^{i24})$, $P(O)(OR^{h24})(OR^{i24})$, and $BR^{j24}R^{k24}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, and (4-7 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{2D} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2E} substituents;

each R^{a24} , R^{b24} , R^{c24} and R^{d24} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, phenyl, C_{3-7} cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl- C_{1-6} alkyl-, C_{3-7} cycloalkyl- C_{1-6} alkyl-, (5-6 membered heteroaryl)- C_{1-6} alkyl-, and (4-7 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6}

alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a24}, R^{b24}, R^{c24} and R^{d24} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2E} substituents;

or, any R^{e24} and R^{d24}, attached to the same N atom, together with the N atom to which they are attached, form a 4-7 membered heterocycloalkyl group, wherein the 4-7 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, or 4 independently selected R^{2E} substituents;

each R^{e24} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f24} and R^{g24} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h24} and Rⁱ²⁴ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, phenyl, C₃₋₇ cycloalkyl, 5-6 membered heteroaryl, 4-7 membered heterocycloalkyl, phenyl-C₁₋₆ alkyl-, C₃₋₇ cycloalkyl-C₁₋₆ alkyl-, (5-6 membered heteroaryl)-C₁₋₆ alkyl-, and (4-7 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j24} and R^{k24} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j24} and R^{k24} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{2E} is independently selected from OH, NO₂, CN, halo, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₃ haloalkyl, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, C₁₋₃ alkoxy, C₁₋₃ haloalkoxy, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxy carbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxy carbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

R³ is selected from H, D, halo, OH, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃

alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

R⁴ is selected from H, D, halo, OH, CN, NO₂, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, cyano-C₁₋₃ alkyl, HO-C₁₋₃ alkyl, C₁₋₃ alkoxy-C₁₋₃ alkyl, C₃₋₅ cycloalkyl, amino, C₁₋₃ alkylamino, di(C₁₋₃ alkyl)amino, thio, C₁₋₃ alkylthio, C₁₋₃ alkylsulfinyl, C₁₋₃ alkylsulfonyl, carbamyl, C₁₋₃ alkylcarbamyl, di(C₁₋₃ alkyl)carbamyl, carboxy, C₁₋₃ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, C₁₋₃ alkylcarbonylamino, C₁₋₃ alkoxycarbonylamino, C₁₋₃ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₃ alkylaminocarbonyloxy, di(C₁₋₃ alkyl)aminocarbonyloxy, C₁₋₃ alkylsulfonylamino, aminosulfonyl, C₁₋₃ alkylaminosulfonyl, di(C₁₋₃ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₃ alkylaminosulfonylamino, di(C₁₋₃ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₃ alkylaminocarbonylamino, and di(C₁₋₃ alkyl)aminocarbonylamino;

each R⁵ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a5}, SR^{a5}, NHOR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)NR^{c5}(OR^{a5}), C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(O)NR^{c5}R^{d5}, C(=NR^{e5})R^{b5}, C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})R^{b5}, NR^{c5}S(O)NR^{c5}R^{d5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)(=NR^{e5})R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, S(O)₂NR^{c5}R^{d5}, OS(O)(=NR^{e5})R^{b5}, OS(O)₂R^{b5}, SF₅, P(O)R^{f5}R^{g5}, OP(O)(OR^{h5})(ORⁱ⁵), P(O)(OR^{h5})(ORⁱ⁵), and BR^{j5}R^{k5}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁵ are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{5A} substituents;

each R^{a5}, R^{b5}, R^{c5} and R^{d5} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10

membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{a5}, R^{b5}, R^{c5} and R^{d5} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{5A} substituents;

or, any R^{c5} and R^{d5}, attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{5A} substituents;

each R^{e5} is independently selected from H, OH, CN, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, and C₂₋₆ alkynyl;

each R^{f5} and R^{g5} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{h5} and Rⁱ⁵ is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-;

each R^{j5} and R^{k5} is independently selected from OH, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy;

or any R^{j5} and R^{k5} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C₁₋₆ alkyl and C₁₋₆ haloalkyl;

each R^{5A} is independently selected from OH, NO₂, CN, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, cyano-C₁₋₆ alkyl, HO-C₁₋₆ alkyl, C₁₋₆ alkoxy-C₁₋₆ alkyl, C₃₋₆ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, amino, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, thio, C₁₋₆ alkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, carbamyl, C₁₋₆ alkylcarbamyl, di(C₁₋₆ alkyl)carbamyl, carboxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxycarbonylamino, C₁₋₆ alkylcarbonyloxy, aminocarbonyloxy, C₁₋₆ alkylaminocarbonyloxy, di(C₁₋₆ alkyl)aminocarbonyloxy, C₁₋₆ alkylsulfonylamino, aminosulfonyl, C₁₋₆ alkylaminosulfonyl, di(C₁₋₆ alkyl)aminosulfonyl, aminosulfonylamino, C₁₋₆ alkylaminosulfonylamino, di(C₁₋₆ alkyl)aminosulfonylamino, aminocarbonylamino, C₁₋₆ alkylaminocarbonylamino, and di(C₁₋₆ alkyl)aminocarbonylamino;

each R^6 is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a6} , SR^{a6} , $NHOR^{a6}$, $C(O)R^{b6}$, $C(O)NR^{c6}R^{d6}$, $C(O)NR^{c6}(OR^{a6})$, $C(O)OR^{a6}$, $OC(O)R^{b6}$, $OC(O)NR^{c6}R^{d6}$, $NR^{c6}R^{d6}$, $NR^{c6}NR^{c6}R^{d6}$, $NR^{c6}C(O)R^{b6}$, $NR^{c6}C(O)OR^{a6}$, $NR^{c6}C(O)NR^{c6}R^{d6}$, $C(=NR^{e6})R^{b6}$, $C(=NR^{e6})NR^{c6}R^{d6}$, $NR^{c6}C(=NR^{e6})NR^{c6}R^{d6}$, $NR^{c6}C(=NR^{e6})R^{b6}$, $NR^{c6}S(O)NR^{c6}R^{d6}$, $NR^{c6}S(O)R^{b6}$, $NR^{c6}S(O)_2R^{b6}$, $NR^{c6}S(O)(=NR^{e6})R^{b6}$, $NR^{c6}S(O)_2NR^{c6}R^{d6}$, $S(O)R^{b6}$, $S(O)NR^{c6}R^{d6}$, $S(O)_2R^{b6}$, $S(O)_2NR^{c6}R^{d6}$, $OS(O)(=NR^{e6})R^{b6}$, $OS(O)_2R^{b6}$, SF_6 , $P(O)R^{f6}R^{g6}$, $OP(O)(OR^{h6})(OR^{i6})$, $P(O)(OR^{h6})(OR^{i6})$, and $BR^{j6}R^{k6}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^6 are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{6A} substituents,

or, two R^6 groups together form an oxo group;

each R^{a6} , R^{b6} , R^{c6} and R^{d6} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^{a6} , R^{b6} , R^{c6} and R^{d6} are each optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{6A} substituents;

or, any R^{c6} and R^{d6} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group, wherein the 4-10 membered heterocycloalkyl group is optionally substituted with 1, 2, 3, 4, 5, or 6 independently selected R^{6A} substituents;

each R^{e6} is independently selected from H, OH, CN, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, and C_{2-6} alkynyl;

each R^{f6} and R^{g6} is independently selected from H, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{h6} and R^{i6} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

each R^{j6} and R^{k6} is independently selected from OH, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

or any R^{j6} and R^{k6} attached to the same B atom, together with the B atom to which they are attached, form a 5- or 6-membered heterocycloalkyl group optionally substituted with 1, 2, 3, or 4 substituents independently selected from C_{1-6} alkyl and C_{1-6} haloalkyl; and

each R^{6A} is independently selected from OH, NO_2 , CN, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, cyano- C_{1-6} alkyl, HO- C_{1-6} alkyl, C_{1-6} alkoxy- C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, amino, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino, thio, C_{1-6} alkylthio, C_{1-6} alkylsulfinyl, C_{1-6} alkylsulfonyl, carbamyl, C_{1-6} alkylcarbamyl, di(C_{1-6} alkyl)carbamyl, carboxy, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy carbonyl, C_{1-6} alkylcarbonylamino, C_{1-6} alkoxy carbonylamino, C_{1-6} alkylcarbonyloxy, aminocarbonyloxy, C_{1-6} alkylaminocarbonyloxy, di(C_{1-6} alkyl)aminocarbonyloxy, C_{1-6} alkylsulfonfylamino, aminosulfonyl, C_{1-6} alkylaminosulfonyl, di(C_{1-6} alkyl)aminosulfonyl, aminosulfonylamino, C_{1-6} alkylaminosulfonylamino, di(C_{1-6} alkyl)aminosulfonylamino, aminocarbonylamino, C_{1-6} alkylaminocarbonylamino, and di(C_{1-6} alkyl)aminocarbonylamino.

2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein X^3 is N and X^4 is N.

3. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein X^3 is CR^3 and X^4 is N.

4. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein X^3 is N and X^4 is CR^4 .

5. The compound of any one of claims 1 or 3, or a pharmaceutically acceptable salt thereof, wherein R^3 is selected from H and C_{1-6} alkyl.

6. The compound of any one of claims 1 or 4, or a pharmaceutically acceptable salt thereof, wherein R^4 is selected from H and C_{1-6} alkyl.

7. The compound of any one of claims 1-6, or a pharmaceutically acceptable salt thereof, wherein Cy¹ is C₆₋₁₀ aryl or 5-10 membered heteroaryl, wherein the C₆₋₁₀ aryl and 5-10 membered heteroaryl are each optionally substituted with 1, 2, 3, or 4 independently selected R⁵ substituents.

8. The compound of any one of claims 1-6, or a pharmaceutically acceptable salt thereof, wherein Cy¹ is phenyl optionally substituted with 1, 2, 3, or 4 independently selected R⁵ substituents; and

each R⁵ is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, NO₂, OR^{a5}, SR^{a5}, NHOR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)NR^{c5}(OR^{a5}), C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(O)NR^{c5}R^{d5}, C(=NR^{e5})R^{b5}, C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})NR^{c5}R^{d5}, NR^{c5}C(=NR^{e5})R^{b5}, NR^{c5}S(O)NR^{c5}R^{d5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)(=NR^{e5})R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, S(O)₂NR^{c5}R^{d5}, OS(O)(=NR^{e5})R^{b5}, and OS(O)₂R^{b5}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R⁵ are each optionally substituted with 1, 2, 3, or 4 independently selected R^{5A} substituents.

9. The compound of any one of claims 1-6, or a pharmaceutically acceptable salt thereof, wherein Cy¹ is phenyl optionally substituted with 1 or 2 independently selected R⁵ substituents; and

each R⁵ is independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and NO₂.

10. The compound of any one of claims 1-6, or a pharmaceutically acceptable salt thereof, wherein Cy¹ is cyanophenyl optionally substituted with fluoro.

11. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein Cy² is 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, or 4 independently selected R⁶ substituents.

12. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein Cy^2 is 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, 3, or 4 independently selected R^6 substituents; and

each R^6 is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a6} , SR^{a6} , $NHOR^{a6}$, $C(O)R^{b6}$, $C(O)NR^{c6}R^{d6}$, $C(O)NR^{c6}(OR^{a6})$, $C(O)OR^{a6}$, $OC(O)R^{b6}$, $OC(O)NR^{c6}R^{d6}$, $NR^{c6}R^{d6}$, $NR^{c6}NR^{c6}R^{d6}$, $NR^{c6}C(O)R^{b6}$, $NR^{c6}C(O)OR^{a6}$, $NR^{c6}C(O)NR^{c6}R^{d6}$, $C(=NR^{e6})R^{b6}$, $C(=NR^{e6})NR^{c6}R^{d6}$, $NR^{c6}C(=NR^{e6})NR^{c6}R^{d6}$, $NR^{c6}C(=NR^{e6})R^{b6}$, $NR^{c6}S(O)NR^{c6}R^{d6}$, $NR^{c6}S(O)R^{b6}$, $NR^{c6}S(O)_2R^{b6}$, $NR^{c6}S(O)(=NR^{e6})R^{b6}$, $NR^{c6}S(O)_2NR^{c6}R^{d6}$, $S(O)R^{b6}$, $S(O)NR^{c6}R^{d6}$, $S(O)_2R^{b6}$, $S(O)_2NR^{c6}R^{d6}$, $OS(O)(=NR^{e6})R^{b6}$, and $OS(O)_2R^{b6}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^6 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{6A} substituents,

or, two R^6 groups together form an oxo group.

13. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein Cy^2 is 5-10 membered heteroaryl or 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl are each optionally substituted with 1, 2, or 3 independently selected R^6 substituents; and

each R^6 is independently selected from halo, C_{1-6} alkyl, and C_{1-6} haloalkyl, wherein the C_{1-6} alkyl is optionally substituted with OH

or, two R^6 groups together form an oxo group.

14. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein Cy^2 is selected from pyrimidin-4-yl, 1-ethyl-1H-pyrazol-5-yl, pyridin-4-yl, 1-isopropyl-1H-pyrazol-5-yl, 1-methyl-1H-pyrazol-5-yl, 1-methyl-6-oxo-1,6-dihydropyridin-3-yl, 3-methylpyridin-4-yl, 4-(hydroxymethyl)-2-methyloxazol-5-yl, 4-methyloxazol-5-yl, oxazol-5-yl, quinolin-6-yl, and quinoxalin-6-yl.

15. The compound of any one of claims 1-14, or a pharmaceutically acceptable salt thereof, wherein R^1 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, and C_{1-6} haloalkyl.

16. The compound of any one of claims 1-14, or a pharmaceutically acceptable salt thereof, wherein R^1 is H.

17. The compound of any one of claims 1-16, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents.

18. The compound of any one of claims 1-16, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents; and


each R^{2A} is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a21} , SR^{a21} , $NHOR^{a21}$, $C(O)R^{b21}$, $C(O)NR^{c21}R^{d21}$, $C(O)NR^{c21}(OR^{a21})$, $C(O)OR^{a21}$, $OC(O)R^{b21}$, $OC(O)NR^{c21}R^{d21}$, $NR^{c21}R^{d21}$, $S(O)R^{b21}$, $S(O)NR^{c21}R^{d21}$, and $S(O)_2R^{b21}$.

19. The compound of any one of claims 1-16, or a pharmaceutically acceptable salt thereof, wherein R^2 is phenyl- C_{1-3} alkyl- or (5-10 membered heteroaryl)- C_{1-3} alkyl-, wherein

the phenyl- C_{1-3} alkyl- and (5-10 membered heteroaryl)- C_{1-3} alkyl- are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents; and each R^{2A} is independently selected from halo, C_{1-6} alkyl, and C_{1-6} haloalkyl.


20. The compound of any one of claims 1-16, or a pharmaceutically acceptable salt thereof, wherein R^2 is selected from pyridin-2-ylmethyl, (1-methyl-1H-pyrazol-3-yl)methyl, (3-fluoropyridin-2-yl)methyl, (6-methylpyridin-2-yl)methyl, 1-(pyridin-2-yl)ethyl, 2-(pyridin-2-yl)ethyl, and 2-fluorobenzyl.

21. The compound of claim 1, which is a compound of Formula (II):




or a pharmaceutically acceptable salt thereof.

22. The compound of claim 1, which is a compound of Formula (IIa):




or a pharmaceutically acceptable salt thereof.

23. The compound of claim 1, which is a compound of Formula (IIb):




or a pharmaceutically acceptable salt thereof.

24. The compound of claim 1, which is a compound of Formula (III):




or a pharmaceutically acceptable salt thereof.

25. The compound of claim 1, which is a compound of Formula (IIIa):




or a pharmaceutically acceptable salt thereof.

26. The compound of claim 1, which is a compound of Formula (IIIb):




or a pharmaceutically acceptable salt thereof.

27. The compound of claim 1, which is a compound of Formula (IV):




or a pharmaceutically acceptable salt thereof.

28. The compound of claim 1, which is a compound of Formula (IVa):




or a pharmaceutically acceptable salt thereof.

29. The compound of claim 1, which is a compound of Formula (IVb):



or a pharmaceutically acceptable salt thereof.

30. The compound of claim 1, which is a compound of Formula (V):



or a pharmaceutically acceptable salt thereof,

wherein n is 0, 1, or 2.

31. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein

X^3 is N or CR^3 ,

X^4 is N or CR^4 ,

wherein at least one of X^3 and X^4 is N;

Cy^1 is C_{6-10} aryl optionally substituted with 1, 2, 3, or 4 independently selected R^5 substituents;

Cy^2 is selected from 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl, wherein the 5-10 membered heteroaryl and 4-10 membered heterocycloalkyl of Cy^2 are each optionally substituted with 1, 2, 3, or 4 independently selected R^6 substituents;

R^1 is selected from H and C_{1-6} alkyl;

R^2 is selected from C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, wherein the C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10

membered heterocycloalkyl)-C₁₋₆ alkyl- of R² are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2A} substituents;

each R^{2A} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, and OR^{a21}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2A} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2B} substituents;

R^{a21} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl;

each R^{2B} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, CN, and OR^{a22}, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, and (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl- of R^{2B} are each optionally substituted with 1, 2, 3, or 4 independently selected R^{2C} substituents;

R^{a22} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl;

each R^{2C} is independently selected from D, halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C₆₋₁₀ aryl-C₁₋₆ alkyl-, C₃₋₁₀ cycloalkyl-C₁₋₆ alkyl-, (5-10 membered heteroaryl)-C₁₋₆ alkyl-, (4-10 membered heterocycloalkyl)-C₁₋₆ alkyl-, and OR^{a23}

R^{a23} is independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₆₋₁₀ aryl, C₃₋₁₀ cycloalkyl, 5-10 membered heteroaryl, and 4-10 membered heterocycloalkyl.

R³ is selected from H and C₁₋₆ alkyl;

R⁴ is selected from H and C₁₋₆ alkyl;

each R^5 is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a5} , and $NR^{c5}R^{d5}$;

each R^{a5} , R^{c5} and R^{d5} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

or, any R^{c5} and R^{d5} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group;

each R^6 is independently selected from D, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-, CN, NO_2 , OR^{a6} , and $NR^{c6}R^{d6}$, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl- of R^6 are each optionally substituted with 1, 2, 3, or 4 independently selected R^{6A} substituents,

or, two R^6 groups together form an oxo group;

each R^{a6} , R^{c6} and R^{d6} is independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{6-10} aryl, C_{3-10} cycloalkyl, 5-10 membered heteroaryl, 4-10 membered heterocycloalkyl, C_{6-10} aryl- C_{1-6} alkyl-, C_{3-10} cycloalkyl- C_{1-6} alkyl-, (5-10 membered heteroaryl)- C_{1-6} alkyl-, and (4-10 membered heterocycloalkyl)- C_{1-6} alkyl-;

or, any R^{c6} and R^{d6} , attached to the same N atom, together with the N atom to which they are attached, form a 4-10 membered heterocycloalkyl group; and

each R^{6A} is independently selected from OH, NO_2 , CN, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, cyano- C_{1-6} alkyl, HO- C_{1-6} alkyl, C_{1-6} alkoxy- C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, amino, and C_{1-6} alkylamino.

32. The compound of claim 1, selected from:

3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyridin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-(2-fluorobenzyl)-7-(pyrimidin-4-yl)-2H-pyrazolo[4,3-c]pyridin-6-yl)benzonitrile;

3-(7-amino-2-(pyridin-2-ylmethyl)-4-(pyrimidin-4-yl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(7-amino-4-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(7-amino-2-((6-methylpyridin-2-yl)methyl)-4-(pyrimidin-4-yl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(7-amino-4-(1-ethyl-1H-pyrazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(7-amino-4-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(7-amino-4-(4-methyloxazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(7-amino-4-(1-ethyl-1H-pyrazol-5-yl)-2-((3-fluoropyridin-2-yl)methyl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(7-amino-2-((3-fluoropyridin-2-yl)methyl)-4-(pyrimidin-4-yl)-2H-pyrazolo[3,4-c]pyridin-5-yl)benzonitrile;

3-(4-amino-2-(pyridin-2-ylmethyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(3-methylpyridin-4-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(4-(hydroxymethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(oxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-(pyridin-2-ylmethyl)-7-(quinolin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-(pyridin-2-ylmethyl)-7-(quinoxalin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(pyridin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((6-methylpyridin-2-yl)methyl)-7-(3-methylpyridin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(4-methyloxazol-5-yl)-2-((6-methylpyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(pyridin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(1-methyl-6-oxo-1,6-dihydropyridin-3-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-((3-fluoropyridin-2-yl)methyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(1-isopropyl-1H-pyrazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(1-methyl-1H-pyrazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;

3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(4-methyloxazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;
3-(4-amino-2-((3-fluoropyridin-2-yl)methyl)-7-(quinoxalin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;
3-(4-amino-7-(1-ethyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile;
3-(4-amino-7-(1-methyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile;
3-(4-amino-7-(1-isopropyl-1H-pyrazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile;
3-(4-amino-7-(4-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile;
3-(4-amino-7-(4-(hydroxymethyl)-2-methyloxazol-5-yl)-2-(pyridin-2-ylmethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile;
3-(4-amino-2-(pyridin-2-ylmethyl)-7-(quinoxalin-6-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile;
3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;
3-(4-amino-2-((1-methyl-1H-pyrazol-3-yl)methyl)-7-(4-methyloxazol-5-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;
3-(4-amino-7-(4-methyloxazol-5-yl)-2-(1-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;
3-(4-amino-2-(1-(pyridin-2-yl)ethyl)-7-(pyrimidin-4-yl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile;
3-(4-amino-7-(4-methyloxazol-5-yl)-2-(2-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)benzonitrile; and
3-(4-amino-7-(4-methyloxazol-5-yl)-2-(2-(pyridin-2-yl)ethyl)-2H-[1,2,3]triazolo[4,5-c]pyridin-6-yl)-2-fluorobenzonitrile,
or a pharmaceutically acceptable salt thereof.

33. A pharmaceutical composition comprising a compound of any one of claims 1 to 32, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient or carrier.

34. A method of inhibiting an activity of an adenosine receptor, comprising contacting the receptor with a compound of any one of claims 1 to 32, or a pharmaceutically acceptable salt thereof.
35. A method of treating a disease or disorder in a patient, wherein the disease or disorder is associated with abnormal expression of A2A or A2B receptors, comprising administering to said patient a therapeutically effective amount of a compound of any one of claims 1 to 32, or a pharmaceutically acceptable salt thereof.
36. The method of claim 35, wherein the disease or disorder is cancer, an inflammatory disease, a cardiovascular disease, or a neurodegenerative disease.
37. The method of claim 36, wherein the cancer is bladder cancer, lung cancer, melanoma, breast cancer, cervical cancer, ovarian cancer, colorectal cancer, pancreatic cancer, esophageal cancer, prostate cancer, kidney cancer, skin cancer, thyroid cancer, liver cancer, uterine cancer, head and neck cancer, or renal cell carcinoma.
38. The method of claim 36, wherein the cancer is non-small cell lung cancer (NSCLC), metastatic castrate-resistant prostate carcinoma (mCRPC), or colorectal carcinoma (CRC).
39. The method of claim 36, wherein the inflammatory disease is pulmonary inflammation.
40. The method of claim 39, wherein the pulmonary inflammation is bleomycin-induced pulmonary fibrosis.
41. The method of claim 36, wherein the inflammatory disease is an adenosine receptor dependent allergic reaction or adenosine receptor immune reaction.
42. The method of claim 41, wherein the adenosine receptor dependent allergic reaction is A2B receptor dependent.
43. The method of claim 36, wherein the inflammatory disease is a respiratory disorder, sepsis, reperfusion injury, or thrombosis.

44. The method of claim 36, wherein the cardiovascular disease is coronary artery disease, cerebrovascular disease, peripheral artery disease, aortic atherosclerosis, or aneurysm.
45. The method of claim 44, wherein the coronary artery disease is myocardial infarction, angina pectoris, or heart failure.
46. The method of claim 44, wherein the cerebrovascular disease is stroke or transient ischemic attack.
47. The method of claim 36, wherein the neurodegenerative disease is Parkinson's disease.
48. The method of claim 35, wherein the disease or disorder is diabetes or insulin resistance.
49. A method of treating or preventing atherosclerotic plaque formation in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a compound of any one of claims 1 to 32, or a pharmaceutically acceptable salt thereof.