




Variable Light

		50	60	0	80
2C4	GQSPKLLIY	[SASYRYT]	GVPDRFTGSGSGTDFTF	TISSVQA	
	**		* *	* *	
574	GKAPKLLIY	[SASYRYT]	GVPSRFSGSGSGTDFLT	LISSLQP	
		* *****			
hum κI	GKAPKLLIY	[AASSLES]	GVPSRFSGSGSGTDFLT	LISSLQP	

		90	100	
2C4	EDLAVYYC	[QQYYIIPYT]	FGGGTKLEIK	(SEQ ID NO:5)
	* *		* *	
574	EDFATYYC	[QQYYIIPYT]	FGQGTVKEIK	(SEQ ID NO:7)
		*** *		
hum κI	EDFATYYC	[QQYNSLPWT]	FGQGTVKEIK	(SEQ ID NO:9)

FIG. 2A

	10	20	30	40
2C4	EVQLQQSGPELVKPGTSVKISKAS	[GFTFTDYTMD]	WVKQS	
	** ** *	* * * *		**
574	EVQLVESGGGLVQPGGSLRLSCAAS	[GFTFTDYTMD]	WVRQA	
		** *	*	
hum III	EVQLVESGGGLVQPGGSLRLSCAAS	[GFTFSSYAMS]	WVRQA	

		50	a	60	70	80
2C4	HGKSLEWIG	[DVNPNSGGSIYNQRFKG]			KASLTVDRSSRIVYM	
	* * *				*** * *	
574	PGKGLEWVA	[DVNPNSGGSIYNQRFKG]			RFTLSVDRSKNTLYL	
		***** * *			* *	
hum III	PGKGLEWVA	[VISGDGGSTYYADSVKG]			RFTISRDNKNTLYL	

	abc	90	100ab	110	
2C4	ELRSLTFEDTAVYYCAR	[NLGPSFYFDY]	WGQGTTLTVSS	(SEQ ID NO:6)	
	*** **		**		
574	QMNSLRAEDTAVYYCAR	[NLGPSFYFDY]	WGQGTTLTVSS	(SEQ ID NO:8)	

hum III	QMNSLRAEDTAVYYCAR	[GRVGYSLYDY]	WGQGTTLTVSS	(SEQ ID NO:10)	

FIG. 2B

3/33

Amino Acid Sequence for Pertuzumab Light Chain

```

1      10      20      30      40      50      60
DIQMTQSPSSLSASVGDRVTITCKASQDVSIQVAWYQOKPGKAPKLLIYSASRYITGVPS
      70      80      90     100     110     120
RFSGSGSGTDFTLTISSLPEDFATYYCQYYIYPYTFGQGTKVEIKRTVAAPSVFIFPP
      130     140     150     160     170     180
SDEQLKSGTASVVCLLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT
      190     200     210
LSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 11)

```

FIG. 3A**Amino Acid Sequence for Pertuzumab Heavy Chain**

```

1      10      20      30      40      50      60
EVQLVESGGGLVQPGGSLRLSCAASGFTFTDITMDWVRQAPGKGLEWVADVNPNSSGGSIY
      70      80      90     100     110     120
NQRFKGRFTLSVDRSKNTLYLQMNSLRAEDTAVYYCARNLGPSFYFDYWGQGTLLTVSSA
      130     140     150     160     170     180
STKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSG
      190     200     210     220     230     240
LYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGP
      250     260     270     280     290     300
SVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS*
      310     320     330     340     350     360
TYRVVSVLTIVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM
      370     380     390     400     410     420
TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ
      430     440     448
QGNVFSCSVMEALHNHYTQKSLSLSPG (SEQ ID NO: 12)

```

FIG. 3B

4/33

Trastuzumab Light Chain

1 15 30 45
DIQNTQSPSSLASVGDRTITCRASQDVNTAVAWYQQKPKGKAPK
46 60 75 90
LLIYSAFLYSGVPSRPSGSRSGTDTLTITSSLPEDFATYYCQQ
91 105 120 135
HYTTPPTPGQGTKEIKRIVAAPSVFIFPPPSDEQLKSGTASVVC
136 150 165 180
LNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSITYSLSSLT
181 195 210 214
LSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 13)

FIG. 4A

Trastuzumab Heavy Chain

1 15 30 45
[EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGL
46 60 75 90
EWVARIYPTNGYTRYADSVKGRFTISADTSKNTAYLQMNSLRAED
91 105 120 135
TAVYYCSRWGCGDGFYAMDYWGQGTLVTVSSA STKGPSVFP LAPSS
136 150 165 180
KSTSGGTAA LGLV K D Y F P E P V T V S W N S G A L T S G V H T F P A V L Q S S
181 195 210 225
GLYSLSSVTVPSSSLG T Q T Y I C N V N H K P S N T K V D K K V E P K S C D K
226 240 255 270
THTCPPCPAPELLCGPSVFLFPKPKD T L M I S R T P E V T C V V D V S
271 285 300 315
HEDPEVKFNWYVDGVEVHN A K T K P R E E Q Y N S T Y R V V S V L T V L H Q D
316 330 345 360
WLN G K E Y K C K V S N K A L P A P I E K T I S K A K G Q P R E P Q V Y T L P P S R E E
361 375 390 405
M T K N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V L D S D G
406 420 435 449
S F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L H N H Y T Q K S L S L S P G

FIG. 4B

(SEQ ID NO: 14)

Pertuzumab Variant Light Chain


1 V H S D I Q M T Q S P S S L S A S V G D R V T I T C K A S Q D V S I G V A W Y Q Q K P G K 45
46 A P K L L I Y S A S Y R Y T G V P S R F S G S G S G T D F T L T I S S L Q P E D F A T Y Y 90
91 C Q Q Y Y I Y P Y T F G Q Q T K V E I K R T V A A P S V F I F P S D E Q L K S G T A S V 135
136 V C L L N N F Y P R E A K V Q W K V D N A L Q S G N S Q E S V T E Q D S K D S T Y S L S S 180
181 T L T L S K A D Y E K H K V Y A C E V T H Q G L S S P V T K S F N R G E C (SEQ ID NO: 15) 217

FIG. 5A

Pertuzumab Variant Heavy Chain

1 E V Q L V E S G G L V Q P G G S L R L S C A A S G F T F T D Y T M D W V R Q A P G K G L 45
46 E W V A D V N P N S G G S I Y N Q R F K G R F T L S V D R S K N T L Y L Q M N S L R A E D 90
91 T A V Y Y C A R N L G P S F Y F D Y W G Q G T L V T V S S A S T K G P S V F P L A P S S K 135
136 S T S G G T A A L G C L V K D Y F P E P V T V S W N S G A L T T S G V H T F P A V L Q S S G 180
181 L Y S L S S V V T V P S S S L G T Q T Y I C N V N H K P S N T K V D K K V E P K S C D K T 225
226 H T C P P C P A P E L L G G P S V F L F P P K P K D T L M I S R T P E V T C V V D V S H 270
271 E D P E V K F N W Y V D G V E V H N A K T K P R E E Q Y N S T Y R V V S V L T V L H Q D W 315
316 L N G K E Y K C K V S N K A L P A P I E K T I S K A K G Q P R E P Q V Y T L P P S R E E M 360
361 T K N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V L D S D G S 405
406 F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L H N H Y T Q K S L S L S P G K 449
(SEQ ID NO: 16)

FIG. 5B



9/33




FIG. 7

8030185 Study Overview

Cohort	Population	N	Agent	Dose (mg) Single Injection	Volume of Injection (mL)	Concentration of Halozyme (U/mL)#	Amount of Halozyme (Units)
Dose Finding							
1 (control)	HMV	6	Piv	420		NA	NA
2	HMV	6	Psc	400	3.3	2,000 U/mL	6,600 U
3	HMV	6	Psc	600	5	2,000 U/mL	10,000 U
4	HMV	6	Psc loading	1200	10	2,000 U/mL	20,000 U
5 (control)	HMV	6	Hsc	600	5	2,000 U/mL	10,000 U
6	HMV	6	Psc + Hsc (co-mix)	400 + 600	8.3	2,000 U/mL	16,600 U
7	HMV	6	Psc + Hsc (co-mix)	1200 + 600	15	2,000 U/mL	30,000 U
8 (Halozyme from Herceptin only)	HMV	6	Psc + Hsc (co-mix)	1200 + 600	15	667 U/mL	10,000 U
Dose Confirmation							
9*	EBC	20	Psc + Hsc (co-mix)	TBD + 600			
10^	EBC	20	Psc + Hsc (FDC)	TBD + 600			


Part 1

Part 2

#MabThera and Herceptin SC use 2,000 U/mL
*Enrollment planned January 2017
^Enrollment planned October 2017

FIG. 8

11/33




No PK interactions between Perjeta SC and Herceptin SC.
No differences in Herceptin PK as monotherapy, with Perjeta
SC or with different rHuPH20 concentrations.

FIG. 9

12/33

No Impact of Lower rHuPH20 Concentration/
Amount on Perjeta PK

Pertuzumab Concentrations with standard (2,000 U/m L) and low
(667 U/m L) rHuPH20



No significant differences in Perjeta PK or Herceptin PK (not shown) when administered with standard (2,000 U/mL) or low (667 U/mL) rHuPH20.

FIG. 10

13/33

Parameter Estimations – Historical IV and
BO30185 SC Data vs. BO30185 IV and BO30185 SC Data




FIG. 11

14/33




FIG. 12

Overview of Adverse Events in Part 1

Extreme Grade	Cohort 1 (P IV 420mg) N = 6	Cohort 5 (H 600mg) N = 6	Cohort 2 (P 400mg) N = 6	Cohort 3 (P 600mg) N = 6	Cohort 4 (P 1200 mg) N = 6	Cohort 6 (P 400mg+ H 600mg) N = 6	Cohort 7 (P 1200mg+ H 600mg) N = 6	Cohort 8 (P 1200mg* + H 600mg) N = 6
Total AEs n = 145 (%)	15 (10)	10 (7)	10 (7)	22 (15)	24 (17)	33 (23)	18 (13)	13 (9)
Grade 1	9 (6)	9 (6)	6 (4)	19 (13)	17 (12)	21 (15)	9 (6)	8 (6)
Grade 2	2 (1)	1 (1)	4 (3)	1 (1)	2 (1)	6 (4)	4 (3)	2 (1)
Grade >=3	1 (1)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

- One Grade 3 AE of diarrhoea , onset Day 32: unrelated, concurrent illness.
- Possible viral infection as concurrent URTI
- No SAE, AESI, AE leading to discontinuation, AE leading to death
- Some AEs ongoing so no final Extreme Grade

*No additional rHuPH20

FIG. 13

Overview of Adverse Events in Part 1
No. of subjects

No. of subjects with AE	Cohort 1 (P IV 420mg) N = 6	Cohort 5 (H 600mg) N = 6	Cohort 2 (P400mg) N = 6	Cohort 3 (P 600mg) N = 6	Cohort 4 (P 1200 mg) N = 6	Cohort 6 (P 400 mg+ H 600mg) N = 6	Cohort 7 (P 1200 mg+ H 600mg) N = 6	Cohort 8 (P 1200 mg*+ H 600mg) N = 6
Any AE, n (%)	5 (83)	5 (83)	5 (83)	6 (100)	5 (83)	6 (100)	6 (100)	6 (100)
Grade 1	4 (67)	5 (83)	4 (67)	6 (100)	5 (83)	6 (100)	6 (100)	5 (83)
Grade 2	2 (33)	1 (17)	3 (50)	1 (17)	2 (33)	5 (83)	2 (33)	2 (33)
Grade >= 3	1 (17)	0	0	0	0	0	0	0

*No additional rHuPH20

FIG. 14

17/33

Most Common Adverse Events (all grades) – incidence ≥ 5% overall in study No. of subjects

Adverse event preferred term	Cohort 1 (P IV 420mg) N = 6	Cohort 5 (H 600mg) N = 6	Cohort 2 (P 400mg) N = 6	Cohort 3 (P 600mg) N = 6	Cohort 4 (P 1200mg) N = 6	Cohort 6 (P 400 mg+ H 600mg) N = 6	Cohort 7 (P 1200 mg+ H 600mg) N = 6	Cohort 8 (P 1200 mg+ H 600mg) N = 6
UPPER RESPIRATORY TRACT INFECTION	2 (33)	0	0	2 (33)	2 (33)	3 (50)	2 (33)	2 (33)
DIARRHOEA	3 (50)	0	1 (17)	2 (33)	1 (17)	1 (17)	1 (17)	0
HEADACHE	1 (17)	1 (17)	0	1 (17)	2 (33)	0	4 (67)	0
DRUG ERUPTION	1 (17)	0	0	0	1 (17)	2 (33)	3 (50)	2 (33)
ANGULAR CHEILITIS	2 (33)	0	0	1 (17)	0	3 (50)	0	1 (17)
PAIN IN EXTREMITY	0	2 (33)	0	1 (17)	2 (33)	1 (17)	0	0
EPISTAXIS	1 (17)	0	0	0	1 (17)	3 (50)	0	0
RASH	0	1 (17)	1 (17)	2 (33)	1 (17)	0	0	0
ABDOMINAL PAIN	0	0	2 (33)	0	0	1 (17)	1 (17)	0
DRY SKIN	0	0	0	1 (17)	1 (17)	2 (33)	0	0
INJECTION RELATED REACTION	0	1 (17)	0	0	0	2 (33)	0	1 (17)
NASAL CONGESTION	0	1 (17)	1 (17)	1 (17)	0	0	0	1 (17)
OROPHARYNGEAL PAIN	1 (17)	0	0	0	2 (33)	0	0	1 (17)
ECZEMA	0	1 (17)	0	0	1 (17)	1 (17)	0	0
FOLLICULITIS	0	0	0	1 (17)	1 (17)	0	0	0

FIG. 15

EGFR Related Toxicity

AEGT basket term/ No. of subjects (%)	Cohort 1 (P IV 420mg) N = 6	Cohort 5 (H 600mg) N = 6	Cohort 2 (P 400mg) N = 6	Cohort 3 (P 600mg) N = 6	Cohort 4 (P 1200mg) N = 6	Cohort 6 (P 400 mg+ H 600mg) N = 6	Cohort 7 (P 1200 mg+ H 600mg) N = 6	Cohort 8 (P 1200 mg*+ H 600mg) N = 6
Diarrhoea	3 (50)	0	1 (17)	2 (33)	2 (33)	2 (33)	1 (17)	0
Mucositis	2 (33)	0	2 (33)	1 (17)	0	4 (67)	0	1 (17)
EGFR Rash	1 (17)	2 (33)	1 (17)	3 (50)	3 (50)	3 (50)	4 (67)	2 (33)

Diarrhoea [PT: Diarrhoea (9), Frequent Bowel Movements (2)]
Onset range: Day 2- 36 with most in first 10 days after drug

Mucositis [PT: Angular Cheilitis (7), Mouth Ulceration (2), Aphthous Ulcer (2) and Gastroenteritis (1)]
Onset range: Day 8- 38, majority occurring between 13 - 29

EGFR Associated Rash [PT: Drug Eruption (9), Rash (5), Eczema (3), Folliculitis (2), Acne (1) and Dermatitis Acneform (1)]

Onset range: Day 4 - 54, no pattern

*No additional rHuPH20

FIG. 16

19/33

Injection Related Reactions and Injection Site Reactions

No. of subjects (%)	Cohort 1 (P IV 420mg) N=6	Cohort 5 (H 600mg) N=6	Cohort 2 (P 400mg) N=6	Cohort 3 (P 600mg) N=6	Cohort 4 (P 1200mg) N=6	Cohort 6 (P 400mg+ H 600mg) N=6	Cohort 7 (P 1200mg+ H 600mg) N=6	Cohort 8 (P 1200mg*+ H 600mg) N=6
Infusion Related Reaction (Systemic)	0							
Injection Related Reaction (Systemic)	1 (17)	0	0	0	0	2 (33)	0	1 (17)
Injection Site Reaction (Local)	0	0	0	0	0	0	1 (17)	0

Injection Site Reaction (Local): any morphological or physiological change at or near the injection site

- 1 Injection Site Reaction in Cohort 7
- 1200mg P SC + 600mg H SC with rHuPH20
- Symptoms were discomfort, pain, tightness and numbness at injection site

Injection Related Reactions (Systemic)

- Symptoms included fever, chills, nausea, stiffness, lightheadedness, skin sensitivity, photophobia, temperature fluctuation, headache

*No additional rHuPH20

FIG. 17

LVEF – ECHO Assessments

No. of subjects (%)	Cohort 1 (P IV) N=6	Cohort 5 (H SC) N=6	Cohort 2, 3, 4 (P SC) N=18	Cohort 6, 7, 8 (P+H SC) N=18
LVEF measurement (%) - Baseline	6 (100)	6 (100)	18 (100)	18 (100)
Absolute value <50% and decrease from baseline ≥10%	0	0	0	0
Absolute value <50% <u>OR</u> decrease from baseline ≥10%points	0	0	1*	0

- * Cohort 3 (600mg P SC):
- o One HMV had a drop of >10% from 67% at Baseline to 56% at Day 22
 - o Follow up EF at Day 85 = 60 %
 - o Cardiologist confirmed no evidence of cardiotoxicity
 - o Drop due to variability of imaging method

FIG. 18


FIG. 19

Drug Substance to support FDCs

Trastuzumab (Herceptin SC DS)	Pertuzumab SC DS
– Trastuzumab: 120 mg/ml	– Pertuzumab: 120 mg/ml
– pH 5.5	– pH 5.5
– 20 mM His-HCl	– 20 mM His-HCl
– 210 mM trehalose	– 200 mM sucrose
– 0.04% PS20	– 0.04% PS20
– 10 mM methionine	– 10 mM methionine

rHuPH20
10 mg/mL rHuPH20 in His-HCl buffer pH 6.5, 130 mM NaCl, 0.05% (w/v) Polysorbate 80
Diluted ~ 435X in DP (negligible)


22/33



Note: Higher HMWS for higher total mAb concentration

FIG. 20


23/33

FIG. 21**Mean Serum Pertuzumab Concentration-Time Profile by Cohort (HMV)**

24/33

FIG. 22

**Geometric Mean Dose-Normalized Serum Pertuzumab
Concentration-Time Profile, With and Without Concomitant
Herceptin**



GM = geometric mean; SC = subcutaneous.


All pertuzumab concentration values normalized to a dose of 600 mg.

Vertical bars, when shown, represent 2.5 or 97.5 percentile confidence intervals on geometric mean.

25/33

FIG. 23


**Geometric Mean Serum Pertuzumab Concentration-Time Profile,
With 667 U/mL or 2,000 U/mL rHuPH20 (HMV)**



GM = geometric mean; H = Herceptin; SC = subcutaneous.

Vertical bars, when shown, represent 2.5 or 97.5 percentile confidence intervals on geometric mean.

26/33

FIG. 24**Geometric Mean Serum Trastuzumab Concentration-Time Profile,
With 667 U/mL or 2,000 U/mL rHuPH20 (HMF)**


GM = geometric mean.

Vertical bars, when shown, represent 2.5 or 97.5 percentile confidence intervals on geometric mean.

27/33

FIG. 25


**Geometric Mean Serum Pertuzumab Concentration-Time Profile
following Perjeta 600 mg SC and Perjeta 420 mg IV Doses**



EBC = early breast cancer; GM = geometric mean; HMV = healthy male volunteer; IV = intravenous SC = subcutaneous.

Vertical bars, when shown, represent 2.5 or 97.5 percentile confidence intervals on geometric mean.

28/33

FIG. 26**Geometric Mean Serum Pertuzumab Concentration-Time Profile
in HMV or EBC Patients**


EBC = early breast cancer; GM = geometric mean; HMV = healthy male volunteer;
SC = subcutaneous.

Vertical bars, when shown, represent 2.5 or 97.5 percentile confidence intervals on geometric mean.

29/33

FIG. 27

**Geometric Mean Dose-Normalized Serum Pertuzumab
Concentration-Time Profile, With 667 U/mL, 1,000 U/mL or
2,000 U/mL rHuPH20**



GM = geometric mean.


All pertuzumab concentration values normalized to a dose of 600 mg.

Vertical bars, when shown, represent 2.5 or 97.5 percentile confidence intervals on geometric mean.

30/33

FIG. 28

**Geometric Mean Serum Trastuzumab Concentration-Time Profile,
With 667 U/mL, 1,000 U/mL, or 2,000 U/mL rHuPH20**



GM = geometric mean.

Vertical bars, when shown, represent 2.5 or 97.5 percentile confidence intervals on geometric mean.

31/33

52 weeks SEC overlay of S&S test samples
stored at -20°C: trehalose (F3) vs sucrose (F4)




FIG. 29




FIG. 30

33/33




FIG. 31