
ATEREOSELECTIVE METHOD FOR THE PREPARATION OF NUCLEOSIDE ANALOGUES

The present invention relates to a novel process for producing *cis* nucleosides or nucleoside analogues and derivatives of formula (A):



(A)

wherein R1 is a pyrimidine base or a pharmaceutically acceptable derivative thereof and Q is oxygen, carbon or sulphur.


Classes of compounds of formula (A), particularly the 2-substituted 4-substituted 1,3-oxathiolanes have been found to have potent antiviral activity. In particular, these compounds have been found to act as potent inhibitors of HIV-1 replication in T-lymphocytes over a prolonged period of time with less cytotoxic side effects than compounds known in the art (see Belleau et al (1993) Bioorg. Med. Chem. Lett. Vol. 3, No. 8, pp. 1723-1728). These compounds have also been found active against 3TC-resistant HIV strains (see Taylor et al (2000) Antiviral Chem. Chemother. Vol 11, No. 4, pp. 291-301; Stoddart et al (2000) Antimicrob. Agents Chemother. Vol. 44, No. 3, pp. 783-786). These compounds are also useful in prophylaxis and treatment of hepatitis B virus infections.

Methods for the preparation of these compounds have been disclosed in PCT publications WO 92/08717 and WO 95/29176 as well as in publications by Belleau et al (1993) Bioorg. Med. Chem. Lett. Vol. 3, No. 8, pp. 1723-1728; Wang et al (1994) Tetrahedron Lett. Vol. 35, No.27, pp. 4739 -

4742; Mansour et al , (1995) J. of Med. Chem. Vol. 38, No. 1, pp. 1-4 and Caputo et al in Eur. J. Org. Chem. Vol. 6, pp. 1455-1458 (1999). These processes involve a multitude of steps that increase the cost of production and reduce the yield of the desired compounds.

Summary of the Invention

The process object of the present invention comprise the step of coupling the intermediate of formula (B):




(B)

wherein

R2 is hydrogen or a hydroxyl protecting group such as C_{7-10} arylalkyl, C_{1-16} acyl or $Si(Z^1)(Z^2)(Z^3)$, where Z^1, Z^2 and Z^3 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-6} alkoxy or C_{6-20} aryloxy; C_{7-20} aralkyl optionally substituted by halogen, C_{1-20} alkyl or C_{1-20} alkoxy; C_{6-20} aryl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-20} alkyl or C_{1-20} alkoxy; trialkylsilyl; fluoro; bromo; chloro and iodo; and


Q is carbon, oxygen or sulfur; with

R1, a pyrimidine base or a pharmaceutically acceptable derivative thereof in the presence of a catalytic amount of an element or combination of elements of group 1B or 11B, a Lewis acid and a tertiary amine. The resulting intermediate of formula (D)




(D).


is deprotected to obtain the *cis* nucleoside of formula (A):



(A)

The compound of formula (B) may be either :

(B1) 

; or (B2) 

or a mixture of the two enantiomers.

In an alternative embodiment of the present invention the deprotection of intermediate of formula (D) is achieved by dissolving said intermediate in a suitable solvent in the presence of a deprotection agent.

In an alternative embodiment of the present invention a simple two step preparation method for *cis* nucleosides of formula (A) wherein the coupling step results in a product wherein the ratio of *cis* to *trans* is greater than 2 to 1 is provided.

In a further embodiment the cis to trans ratio of the intermediate product of formula (D) is inversely proportional to the reaction temperature of the coupling step.

In an alternate embodiment of the present invention, the deprotection step results in the selective

precipitation of the cis nucleoside of formula (A) by the selection of an appropriate deprotection agent and solvent.


The processes of this invention have the advantages of allowing preparation of a nucleoside of formula (A), analogues or derivatives thereof without using expensive starting materials, cumbersome protection and deprotection steps or addition and removal of 2'- or 3'-substituents.

The process of this invention produces cis nucleosides of formula (A) in high yields, with high purity and high stereoselectivity.

The process of this invention has the further advantage of generating nucleosides whose stereochemical configuration can be easily controlled simply by the selection of the appropriate starting conditions.

Detailed Description of the Invention

The present invention discloses a stereoselective process for making predominantly cis nucleosides or nucleoside analogues and derivatives of formula (A):




(A)

wherein

R1 is a pyrimidine base or a pharmaceutically acceptable derivative thereof; and

Q is carbon, oxygen or sulfur;

consisting of coupling step of the a compound of formula (B):




(B)

wherein

R2 is hydrogen or a hydroxyl protecting group such as C_{7-10} aralalkyl, C_{1-16} acyl or $Si(Z^1)(Z^2)(Z^3)$, where Z^1, Z^2 and Z^3 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-6} alkoxy or C_{6-20} aryloxy; C_{7-20} aralkyl optionally substituted by halogen, C_{1-20} alkyl or C_{1-20} alkoxy; C_{6-20} aryl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-20} alkyl or C_{1-20} alkoxy; trialkylsilyl; fluoro; bromo; chloro and iodo;

with a base R1, wherein R1 is a pyrimidine base or a pharmaceutically acceptable derivative thereof, in a suitable coupling solvent in the presence of a catalytic amount of an element or combination of elements of group 1B or 11B ; a tertiary amine and a Lewis acid of formula (C):



(C)


wherein

R4, R5 and R6 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-6} alkoxy or C_{6-}

$_{20}$ aryloxy; C_{7-20} aralkyl optionally substituted by halogen, C_{1-20} alkyl or C_{1-20} alkoxy; C_{6-20} aryl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-20} alkyl or C_{1-20} alkoxy; trialkylsilyl; fluoro; bromo; chloro and iodo; and

R7 is selected from the group consisting of fluoro; bromo; chloro; iodo; C_{1-20} sulphonate esters, optionally substituted by fluoro, bromo, chloro or iodo; C_{1-20} alkyl esters optionally substituted by fluoro, bromo, chloro or iodo; triiodide; a silyl group of the general formula (R4)(R5)(R6)Si (wherein R4, R5, and R6 are as defined above); C_{6-20} arylselenenyl; C_{6-20} arylsulfenyl; C_{6-20} alkoxyalkyl; and trialkylsiloxy;

to yield an intermediate or formula (D):



(D).

The coupling step is followed by a deprotection step to yield the *cis* nucleosides or nucleoside analogues or derivatives of formula (A).

In an alternative embodiment of the present invention a simple two step preparation method for *cis* nucleosides of formula (A) wherein the process results in a product of formula (A) wherein the ratio of *cis* to *trans* is greater than 2 to 1. The invention includes a process wherein the ratio of *cis* to *trans* is greater than or equal to 3 to 1.

The term "alkyl", as used herein, unless otherwise specified, refers to a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon of C_{1-30}.

particularly C_{1-6}, unsubstituted or optionally mono- or di-substituted by hydroxy, N_3, CN, SH, amino, halogen (F, Cl, Br, I), C_{6-12} aryl, C_{1-6} alkyl, C_{2-12} alkoxyalkyl or nitro. It specifically includes methyl, ethyl, cyclopropyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, cyclohexylmethyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl.

The term "acyl", as used hereinafter refers to a functional group derived from an aliphatic carboxylic acid, by removal of the -OH group of 1 to 30 carbon atoms, particularly 1 to 6 carbon atoms. Like the acid to which it is related, an aliphatic acyl radical may be substituted (by a hydroxy, N_3, CN, halogen (F, Cl, Br, I), C_{6-12} aryl, C_{1-6} alkyl, C_{2-12} alkoxyalkyl or nitro) or unsubstituted, and whatever the structure of the rest of the molecule may be, the properties of the functional group remain essentially the same (e.g., acetyl, propionyl, isobutanoyl, pivaloyl, hexanoyl, butyryl, pentanoyl, 3-methylbutyryl, hydrogen succinate, mesylate, valeryl, caproic, caprylic, capric, lauric, myristic, palmitic, stearic, oleic, 3-chlorobenzoate, trifluoroacetyl, chloroacetyl, and cyclohexanoyl).

The terms "alkenyl" and "alkynyl" represent substituted (by a N_3, CN, halogen, hydroxyl or C_{6-20} aryl) or unsubstituted straight, branched or cyclic hydrocarbon chains having 2 to 30 carbon atoms and preferably from 2 to 6 carbon atoms and containing at least one unsaturated group (e.g. allyl).

The term "alkoxy" represents a substituted or unsubstituted alkyl group containing from 1 to 30 carbon atoms and preferably from 1 to 6 carbon atoms, wherein the alkyl group is covalently bonded to an oxygen atom (e.g., methoxy and ethoxy).

The term "aryl" represents a aromatic moiety which may be mono, bi, tri, tetra or penta-substituted by hydroxy, nitro, N_3, CN, halogen (F, Cl, Br, I) or combinations thereof and containing at least one benzenoid-type ring, the group may contain from 6 to 14 carbon atoms (e.g., phenyl and naphthyl), particularly 6 to 10 carbon atoms.

The term "aryloxy" represents a substituted (by a halogen, trifluoromethyl or C_{1-5} alkoxy) or unsubstituted aryl moiety, having 6 to 14 carbon atoms, covalently bonded through an oxygen atom (e.g., benzyloxy, phenoxy).

The term "aralalkyl" represents a substituent comprising an aryl moiety attached via an alkyl chain (e.g. benzyl, phenylethyl) wherein the sum total of carbon atoms for the aryl moiety and the alkyl chain is 7 to 21. The aryl or chain portion of the group is optionally mono- or di-substituted with OH, SH, amino, halogen or C_{1-6} alkyl.

The term "thiol" represents C_{1-6} alkyl, C_{6-15} aryl, C_{7-21} aralkyl, C_{2-6} alkenyl or C_{2-6} alkynyl groups covalently bonded to an adjacent sulfur atom bearing a hydrogen.

The terms "alkylthio" (e.g. thiomethy, thioethyl) and "arylthio" (e.g. thiophenyl, thiobenzyl), refers to C_{1-6} alkyl or C_{6-10} aryl groups, unsubstituted or optionally mono- or di-substituted by, hydroxy, halogen (F, Cl, Br, I),

C_{6-12} aryl, C_{1-6} alkyl, C_{2-12} alkoxyalkyl or nitro, covalently bonded to an adjacent sulfur atom.

The terms "acyloxy" and "alkoxycarbonyl" refer to 1 to 30 carbon atoms chains, particularly 1 to 6 carbon atoms, that are either saturated or unsaturated and may also be straight or branched (e.g.: acetyloxy). The chains may be unsubstituted or optionally mono- or di-substituted by hydroxy, N_3, CN, SH, amino, halogen (F, Cl, Br, I), C_{6-12} aryl, C_{1-6} alkyl, C_{2-12} alkoxyalkyl or nitro.

The term "alkoxyalkyl" represents a C_{1-6} alkoxy group attached to an adjacent C_{1-6} alkyl group (e.g., methoxymethyl, ethoxymethyl). They may be unsubstituted or optionally mono- or di-substituted by hydroxy, N_3, CN, SH, amino, halogen (F, Cl, Br, I), C_{6-12} aryl, C_{1-6} alkyl, C_{2-12} alkoxyalkyl or nitro.

The term "heterocycle" represents a saturated or unsaturated mono- or polycyclic (i.e. bicyclic) ring incorporating 1 or more (i.e. 1-4) heteroatoms selected from N, O and S. It is understood that a heterocycle is optionally mono- or di-substituted with OH, SH, amino, halogen, CF_3, oxo or C_{1-6} alkyl. Examples of suitable monocyclic heterocycles include but are not limited to pyridine, piperidine, pyrazine, piperazine, pyrimidine, imidazole, thiazole, oxazole, furan, pyran and thiophene. Examples of suitable bicyclic heterocycles include but are not limited to indole, benzimidazole, quinoline, isoquinoline, purine, and carbazole.

The term "aralkyl" represents a substituent comprising a C_{6-10} aryl moiety attached via a C_{1-6} alkyl

chain (e.g. benzyl, phenethyl). The aryl or chain portion of the group is unsubstituted or optionally mono- or di-substituted by hydroxy, N_3, CN, SH, amino, halogen (F, Cl, Br, I), C_{6-12} aryl, C_{1-6} alkyl, C_{2-12} alkoxyalkyl or nitro.

The term "amino" represents C_{1-6} alkyl, C_{6-10} aryl, C_{2-6} alkenyl, C_{2-6} alkynyl, or C_{7-12} aralkyl groups, unsubstituted or optionally mono- or di-substituted by hydroxy, N_3, CN, SH, amino, halogen (F, Cl, Br, I), C_{6-12} aryl, C_{1-6} alkyl, C_{2-12} alkoxyalkyl or nitro, wherein the carbon atoms are covalently bonded to an adjacent element through a nitrogen atom (e.g., pyrrolidine). They include primary, secondary and tertiary amines and quaternary ammonium salts.


The term "protected" as used herein and unless otherwise defined refers to a group that is added to an oxygen, nitrogen, or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis. Suitable protecting groups are described, for example, in Greene, et al., "Protective Groups in Organic Synthesis," John Wiley and Sons, Second Edition, 1991, hereby incorporated by reference.

By pyrimidine base derivative or analogue is meant a pyrimidine base found in nucleoside or an analogue thereof which mimics such bases in that their structures (the kinds of atoms and their arrangement) are similar to the normal bases but may possess additional or lack certain of the functional properties of the normal bases. Derivatives of such bases or analogues include those obtained by replacement of a CH moiety by a nitrogen atom (for example,


5-azapyrimidines such as 5-azacytosine) or may have ring substituted by halogen, hydroxyl, azido, cyano, amino, substituted amino, thiol, C_{1-6} alkyl and C_{6-10}_aryl.

By the term "pharmaceutically acceptable derivative" of a pyrimidine is meant any pyrimidine base which may give rise to a pharmaceutically acceptable salt, ester, or salt of such ester of a compound of formula (A), or any other compound which, upon administration to the recipient, is capable of providing (directly or indirectly) a compound of formula (A) or an antivirally active metabolite or residue thereof. It will be appreciated by those skilled in the art that the compounds of formula (A) may be modified to provide pharmaceutically acceptable derivatives thereof, at functional groups in the base moiety.

The compound of formula (B) may be either :



(B1); or



(B2) or a mixture of the two enantiomers. The sulfoxide may be a single enantiomer or a mixture of enantiomers including but not limited to a racemic mixture.

The coupling step of the process object of the present invention includes the addition of one or more elements of group 1B or 11B. The element or combination of elements used may be in their oxidized state. This element or combination of elements of group 1B or 11B catalyzes the coupling step. The chosen element or combination of elements of group 1B or 11B are present in amounts between about 0.25 molar % and about 100 molar %. In another

embodiment, the concentration of the element or combination of elements of group 1B or 11B may be between about 5% to about 35%.

The process object of the present invention includes a coupling step wherein the element or combination of elements of group 1B or 11B are selected from the group comprising Cu^{+}, Cu^{2+},Ag^{+}, Au^{+}, Au^{3+}, Zn^{2+} Cd^{2+} and combinations thereof.

The process object of the present invention includes a coupling step wherein the element or combination of elements of group 1B or 11B are selected from Cu^{+}, Cu^{2+} or Zn^{2+}.

The term tertiary amine includes tertiary amines with high basicity. The tertiary amine is of the form $N(Z^4)(Z^5)(Z^6)$ wherein (Z^4),(Z^5),(Z^6) are independently selected from the group consisting C_{1-6} alkyl optionally substituted with C_{1-3} alkyl, C_{6-10} aryl, halogen. Examples of the tertiary amine include triethylamine, diethylcyclohexylamine, diethylmethylamine, dimethylethylamine, dimethylisopropylamine, dimethylbutylamine, dimethylcyclohexylamine, tributylamine, diethylmethylamine, dimethylisopropylamine and diisopropylethylamine.

The amount of tertiary amine can vary between about 1 eq to about 4 eq. The amount of tertiary amine used may vary between about 1 eq and 2 eq.

The coupling step of the process object of the present invention is preformed in a suitable coupling solvent. A suitable coupling solvent includes C_{1-10} chlorinated organic solvents. Suitable coupling solvents

also include C_{1-8} chloroalkyls, C_{1-8} chloroalkenyls, C_{6-10} chloroaryls , C_{1-8} alkylonitriles and combinations thereof. The coupling solvents may be selected from chloromethanes, chloroethanes, methanonitriles or mixtures thereof. The coupling solvents of interest include dichloromethane, chloroform, acetonitrile, dichloroethane, chlorobenzene and combinations thereof.

The amount of coupling solvent used may vary between about 5 mL per g of sulfoxide to 50 mL per gram of sulfoxide. In an alternate embodiment of the invention the amount of coupling solvent is between 10 mL per g of sulfoxide to 30 mL per gram of sulfoxide.

The coupling step of the process object of the present invention is affected by the temperature of the reaction. The cis to trans ratio of the product of formula (D) is inversely proportional to the reaction temperature. The coupling step is preformed at a temperature between about 40 degrees C and about - 40 degrees C. In an alternate embodiment, the coupling step reaction temperature is between about 30 degrees C and about - 50 degrees C.


The second step in the process object of the present invention is a deprotection step. The deprotection crystallization step is preformed in a suitable solvent. Of particular interest are solvents that favor the crystallization of the product of formula (A). Suitable solvents include water, methanol, ethanol, toluene, tert-butyl methyl ether or combinations thereof.

The deprotection may also include the presence of adequate amounts a deprotection agent. Of particular interest are deprotection agents that will aid in the separation of the cis product of formula (A). Suitable


deprotection agents are selected according to the identity of the protecting group on the intermediate of formula (D) as shown in Greene, et al., "Protective Groups in Organic Synthesis," John Wiley and Sons, Second Edition, 1991. Deprotection agents may be alkaline. Deprotection agents include sodium hydroxide, sodium methoxide, ammonium hydroxide, potassium hydroxide, lithium hydroxide and methanolic ammonia.

In another embodiment of the present invention, the deprotection agent is present in catalytic amounts. In another embodiment the deprotection agent is present in concentrations between about 0.1 molar percentage and about 50 molar percentage. An alternative embodiment includes deprotection agent concentrations between about 5 to about 20 molar percentage of the deprotection agent.


Conveniently, the base R1 is selected from:




(I)




(II)




(III)




(IV)




(V)



(VI)



(VII)



(VIII)

wherein:

x is oxygen, NH or sulfur.

y is oxygen, NH or sulfur.


R8 and R9 are independently selected from hydrogen, hydroxyl, amino, C_{1-6} alkyl or C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-10} acyl, C_{6-10} aryl, C_{6-11} carbonylaryl, C_{1-7} carbonyloxyalkyl, C_{6-11} carbonyloxyaryl, C_{2-7} carbonylaminoalkyl, or amino acids.

R8 may be a saturated or unsaturated C_{3-8} carbocyclic ring optionally substituted with COOH, $C(O)NH_2$, OH, SH, NH_2, NO_2, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halogen, C(O)R14 wherein R14 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and C(O)OR15 wherein R15 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl; and R9 is chosen from H, C_{1-6} alkyl, C_{2-6} alkenyl and C_{2-6} alkynyl.

R_8R_9 can also be connected to the nitrogen atom to form a saturated or unsaturated C_{3-8} heterocyclic ring optionally substituted with C(O)OH, $C(O)NH_2$, OH, SH, NH_2, NO_2, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halogen, C(O)R14 wherein R14 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and C(O)OR15 wherein R15 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl.

R10 , R11, R12 and R13 are each independently selected from hydrogen, halogen, hydroxyl, amino, cyano, carboxyl, carbamoyl, C_{2-7} alkoxycarbonyl, hydroxymethyl, trifluoromethyl, C_{6-10} arylthio, C_{1-6} alkyl, C_{2-6} alkenyl substituted or unsubstituted with halogen or azido, C_{2-6} alkynyl, C_{1-6} acyloxy, thiocarboxy, thiocarbamoyl, carbamate, ureido, amidino, or C_{6-10} aryloxy.

In another embodiment of the present invention R1 is



(I) or (VI)

wherein R8 and R9 are independently selected from hydrogen, hydroxyl, amino, C_{1-10} alkyl or C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} acyl, C_{6-10} aryl, C_{6-16} carbonylaryl, C_{1-10} carbonyloxyalkyl, C_{6-16} carbonyloxyaryl, C_{2-12} carbonylaminoalkyl, or amino acids; R10 and R11 are each independently selected from hydrogen, halogen, hydroxyl, hydroxymethyl, trifluoromethyl, C_{1-6} alkyl, C_{2-6} alkenyl substituted or unsubstituted with halogen , azido, C_{2-6} alkynyl, or C_{6-10} aryloxy; and

X and Y are independently selected from O or S.


In an alternate embodiment R1 is a pyrimidine base selected from N^4 -alkylpyrimidines, N^4 -acylpyrimidines, 4-halopyrimidines, N^4 -acetylenic pyrimidines, 4-amino and N^4 -acyl pyrimidines, 4-hydroxyalkyl pyrimidines, 4-thioalkyl pyrimidines, thymine, cytosine, 6-azapyrimidine, including 6-azacytosine, 2- and/or 4-mercaptopyrimidine, uracil, C^5 -alkylpyrimidines, C^5 -benzylpyrimidines, C^5 halopyrimidines, C^5 -vinylpyrimidine, C^5 -acetylenic pyrimidine, C^5 -acyl pyrimidine, C^5 -amidopyrimidine, C^5 -cyanopyrimidine, C^5 -nitropyrimidine, C^5 -aminopyrimidine, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl or pyrazolopyrimidinyl.

The functional oxygen and nitrogen groups on R1 can be protected as necessary or desired. Suitable

protecting groups are well known to those skilled in the art, and include trimethylsilyl, dimethylhexylsilyl, t-butyldimethylsilyl, and t-butyldiphenylsilyl, trityl, C_{1-12} alkyl groups, C_{1-12} acyl groups such as acetyl and propionyl, benzoyl, methanesulfonyl, and p-toluenesulfonyl.

In an additional embodiment of the present invention R1 is selected from cytosine, uracil, thymine, 5-fluoropyrimidine or protected analogs of these bases.

Another embodiment of the present invention includes a stereoselective process for making predominantly cis nucleosides or nucleoside analogues and derivatives of formula (A):




(A)

wherein

R1 is a pyrimidine base selected from N^4 -alkylpyrimidines, N^4 -acylpyrimidines, 4-halopyrimidines, N^4 -acetylenic pyrimidines, 4-amino and N^4 -acyl pyrimidines, 4-hydroxyalkyl pyrimidines, 4-thioalkyl pyrimidines, thymine, cytosine, 6-azapyrimidine, including 6-azacytosine, 2- and/or 4-mercaptopyrimidine, uracil, C^5 -alkylpyrimidines, C^5 -benzylpyrimidines, C^5 halopyrimidines, C^5 -vinylpyrimidine, C^5 -acetylenic pyrimidine, C^5 -acyl pyrimidine, C^5 -amidopyrimidine, C^5 -cyanopyrimidine, C^5 -nitropyrimidine, C^5 -aminopyrimidine, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl. or a pharmaceutically acceptable derivative thereof; and

Q is oxygen ,

consisting of coupling step of the a compound of formula (B):




(B)

wherein

R2 is C_{7-10} aralalkyl, C_{1-16} acyl or $Si(Z^1)(Z^2)(Z^3)$, where Z^1, Z^2 and Z^3 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-6} alkoxy or C_{6-20} aryloxy; C_{7-20} aralkyl optionally substituted by halogen, C_{1-20} alkyl or C_{1-20} alkoxy; C_{6-20} aryl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-20} alkyl or C_{1-20} alkoxy; trialkylsilyl; fluoro; bromo; chloro and iodo;

with a base R1, as defined above, in a suitable coupling solvent in the presence of a catalytic amount of an element or combination of elements of group 1B or 11B ; a tertiary amine and a Lewis acid of formula (C):



(C)


wherein

R4, R5 and R6 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-6} alkoxy or C_{6-}

$_{20}$ aryloxy; C_{7-20} aralkyl optionally substituted by halogen, C_{1-20} alkyl or C_{1-20} alkoxy; C_{6-20} aryl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-20} alkyl or C_{1-20} alkoxy; trialkylsilyl; fluoro; bromo; chloro and iodo; and

R7 is selected from the group consisting of fluoro; bromo; chloro; iodo; C_{1-20} sulphonate esters, optionally substituted by fluoro, bromo, chloro or iodo; C_{1-20} alkyl esters optionally substituted by fluoro, bromo, chloro or iodo; triiodide; a silyl group of the general formula (R4)(R5)(R6)Si (wherein R4, R5, and R6 are as defined above); C_{6-20} arylselenenyl; C_{6-20} arylsulfenyl; C_{6-20} alkoxyalkyl; and trialkylsiloxy;


to yield an intermediate or formula (D):



(D)

wherein Q, R1 and R2 are as defined above. The coupling step is followed by a deprotection step wherein said intermediate (D) is dissolved in a suitable solvent in the presence of appropriate amounts of a deprotection agent to yield the *cis* nucleosides or nucleoside analogues or derivatives of formula (A).

The present invention includes the embodiment wherein the stereoselective process for making predominantly cis nucleosides or nucleoside analogues and derivatives of formula (A):




(A)

wherein

R1 is a pyrimidine base selected from cytosine, uracil, thymine, 5-fluoropyrimidine or a pharmaceutically acceptable derivative thereof; and

Q is oxygen ,


consisting of coupling step of the a compound of formula (B):




(B)

wherein

R2 is



wherein W is halogen, C_{1-16} alkyl, C_{2-16} alkoxyalkyl, C_{6-10} aryl, C_{1-16} alkoxy or nitro; or




wherein E is C_{6-10} aryl, C_{1-16} alkoxy, C_{2-16} alkoxyalkyl or C_{1-16} alkyl; or

$Si(Z^1)(Z^2)(Z^3)$, where Z^1, Z^2 and Z^3 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo; C_{7-10} aralalkyl optionally substituted by

fluoro, bromo, chloro or iodo; and C_{6-10} aryl optionally substituted by fluoro, bromo, chloro or iodo; with

a base R1, as defined above, in a suitable coupling solvent in the presence of a catalytic amount of Cu or Zn or mixtures thereof; a tertiary amine and a Lewis acid selected from trimethylsilyl triflate bromotrimethylsilane or iodotrimethylsilane;to yield an intermediate or formula (D):




(D)

wherein Q, R1 and R2 are as defined above. The coupling step is followed by a deprotection step wherein said intermediate (D) is dissolved in a suitable solvent in the presence of appropriate amounts of a deprotection agent to yield the *cis* nucleosides or nucleoside analogues or derivatives of formula (A).

The process object of the present invention includes the reaction scheme shown in Scheme 1:

Scheme 1



The various steps as illustrated in Scheme 1 may be briefly described as follows:

Step 1: The sulfoxide of formula (B) may be obtained using several methods including those disclosed in PCT publications WO 92/08717 and WO 95/29176, J. Med. Chem. 38(1) 1-4 (1995), Tetrahedron Lett. 35(27) 4739-4742 (1994), Bioorg. Med. Chem. Lett. 3 (8) 1723-1728 (1993) and Eur. J. Org. Chem. 6:1455-1458 (1999). The sulfoxide may be a single enantiomer or a mixture of enantiomers including but not limited to a racemic mixture.

The sulfoxide of formula (B) is coupled with base R1. The base R1 may be previously protected , for example silylated (or silylated in situ) pyrimidine base or pharmaceutically acceptable derivative may be used. The coupling reaction takes place in the presence of a tertiary amine, a Lewis acid of formula (C) and catalytic amounts of

an element of groups IB or IIB in a suitable coupling solvent to give the cis/trans pyrimidine nucleoside of formula (D). In the resulting intermediate of formula (D), the cis isomer predominates over the trans isomer in a ratio equal to greater than 2 to 1. The ratio of cis to trans isomer is inversely proportional to the temperature of the reaction temperature. The coupling reaction may be preformed at or below room temperature. The temperature of the coupling step may be between about 0 degrees C and about - 50 degrees C.

If a silylated base is used, adequate silylating agent may include t-butyldimethylsilyl triflate 1,1,1,3,3,3 hexamethyldisilazane, TMSI, N,O,bis(TMS) acetonide and trimethylsilyl triflate. Additional protective agents are disclosed in Greene, et al., "Protective Groups in Organic Synthesis," John Wiley and Sons, Second Edition, 1991.

The tertiary amine used in step 1 include triethylamine, diethylcyclohexylamine, diethylmethylamine, dimethylethylamine, dimethylisopropylamine, dimethylbutylamine, dimethylcyclohexylamine, tributylamine, diethylmethylamine, dimethylisopropylamine and diisopropylethylamine and combinations thereof.

The element or combination of elements of groups IB or IIB used in step 1 include Cu, Ag, Au, Zn, or Cd in oxidized state.

The suitable coupling solvent is an organic solvent with one to ten carbons. The suitable coupling solvents include CH_2Cl_2, CH_3CN and mixtures thereof.

Suitable Lewis acids include trimethylsilyl triflate bromotrimethylsilane iodotrimethylsilane and


combinations thereof. The amount of the Lewis acid may be between about 2 eq to about 5 eq.

Step 2: The cis/trans pyrimidine 1,3-oxathiolane nucleoside of formula (D) is dissolved in a suitable deprotection solvent in the presence of adequate amounts of a deprotection agent to yield the *cis* nucleosides or nucleoside analogues or derivatives of formula (A). The deprotection step is preformed at a temperature below the boiling point of the suitable deprotection solvent. The reaction temperature of the deprotection step may be between - 30 degrees C and 60 degrees C. The reaction may be preformed at a temperature between about 0 degrees C to about 35 degrees C.

A deprotection suitable solvent favors the crystallization of the product of formula (A). Suitable solvents include water, methanol, ethanol, toluene, tert-butyl methyl ether or combinations thereof.

Suitable deprotection agents include sodium hydroxide, sodium methoxide, ammonium hydroxide, potassium hydroxide, lithium hydroxide and methanolic ammonia. Of particular interest are deprotection agents which will aid in the separation of the product of formula (A).

Scheme 1a



The various steps as illustrated in Schemes 1a may be briefly described as follows:

Step 1: The sulfoxide of the 1,3 oxathiolane of formula (Bx) may be obtained using several methods including those disclosed in PCT publications WO 92/08717 and WO 95/29176, J. Med. Chem. 38(1) 1-4 (1995), Tetrahedron Lett. 35(27) 4739-4742 (1994), Bioorg. Med. Chem. Lett. 3 (8) 1723-1728 (1993). The asymmetric synthesis of the sulfoxide of formula (Bx) is disclosed by Caputo et al in Eur. J. Org. Chem. 6:1455-1458 (1999).

The sulfoxide of the 1,3 oxathiolane of formula (Bx) is coupled with base R1. The base R1 may be previously protected, for example silylated (or silylated in situ) pyrimidine base or pharmaceutically acceptable derivative. The coupling reaction takes place in the presence of a tertiary amine, a Lewis acid of formula (C) and catalytic

amounts of an element of groups IB or IIB in a suitable coupling solvent to give the cis/trans pyrimidine 1,3 oxathiolane nucleoside of formula (Dx). In the resulting intermediate of formula (Dx), the cis isomer predominates over the trans isomer in a ratio equal to greater than 2 to 1. The ratio of cis to trans isomer is inversely proportional to the temperature of the reaction temperature. The reaction may be preformed at or below room temperature.

The silylating agent that may be used for the protection of R1 include t-butyldimethylsilyl triflate 1,1,1,3,3,3 hexamethyldisilazane, TMSI, N,O,bisTMS acetonite and trimethylsilyl triflate.

The tertiary amine used in step 1 include triethylamine, diethylcyclohexylamine, diethylmethylamine, dimethylethylamine, dimethylisopropylamine, dimethylbutylamine, dimethylcyclohexylamine, tributylamine, diethylmethylamine, dimethylisopropylamine and diisopropylethylamine and combinations thereof.

The element or combination of elements of groups IB or IIB used in step 1 include Cu, Ag, Au, Zn, or Cd in oxidized state.

The suitable coupling solvent is an organic solvent. The suitable coupling solvents include CH_2Cl_2, CH_3CN or mixtures thereof.

The Lewis acid that may be used in this step include trimethylsilyl triflate bromotrimethylsilane iodotrimethylsilane and mixtures thereof. The amount of the Lewis acid may be between about 2 eq to about 5 eq.

Step 2: The cis/trans pyrimidine nucleoside of formula (Dx) is dissolved in a suitable deprotection solvent

in the presence of adequate amounts of a deprotection agent to yield the *cis* nucleosides or nucleoside analogues or derivatives of formula (Ax).


A suitable deprotection solvent favors the crystallization of the product of formula (A). Suitable solvents include water, methanol, ethanol, toluene, tert-butyl methyl ether or combinations thereof. Suitable combinations of solvents include methanol and water, methanol and toluene, methanol and tert-butyl methyl ether mixtures.

Suitable deprotection agents include sodium hydroxide, sodium methoxide, ammonium hydroxide, potassium hydroxide, lithium hydroxide and methanolic ammonia. Of particular interest are deprotection agents that will aid in the separation of the product of formula (A).

The deprotection step is preformed at a temperature below the boiling point of the suitable deprotection solvent

The following examples illustrate the present invention in a manner of which it can be practiced but, as such, should not be construed as limitations upon the overall scope of the processes of this invention.

1) 2-benzoyloxymethyl-1, 3-oxathiolane



(1) **(2)**

Compound (1) was dissolved in toluene and the solution was heated up to 90-100°C. Catalyst was added followed by mercaptoethanol (portionwise). 5 molar % of catalyst were used. Reactions were carried out on 15g scale at 0.3M concentration of (1). The reaction mixture was refluxed with water removal by Dean-Stark trap. Results for this step are shown in Table 1.

Table 1

Catalyst	Reaction after 20 min		Reaction after 40 min	
	Conversion %	Yield of (2) %	Conversion %	Yield of (2) %
BF_3OEt_2	100	79	100	71
pTsOH	100	82	100	80

Compound (2) was identified by ^1H- and ^{13}C-NMR.

R_f: 0.39 (hexane: ethyl acetate)

^1H-NMR: δ (ppm in $CDCl_3$)

8.03 (m, 2H, aromatic)

7.53 (m, 1H, aromatic)

7.39 (m, 2H, aromatic)

5.41 (dd, 1H, C_2-H)

4.43 (m, 2H, C_2-$CH_2OCC_6H_5$)

4.21 (m, 1H, C_5-H)


3.96 (m, 1H, C_5-H)

2.98 (m, 2H, C_4-H)

^{13}C-NMR: δ (ppm in $CDCl_3$)

166.82, 133.74, 130.35, 128.97, 83.58, 71.87, 66.62 and 32.74

2) 2-benzoyloxymethyl-1,3-oxathiolane-S-oxide



(2) (3)

Cold 30% hydrogen peroxide (46ml, 0.44 mol) was added to (2) (82g, 0.366 mol) in toluene (8 ml). 10M sulfuric acid (4.5ml, 0.044 mol, 10 mol %) was added dropwise (addition time approximately 1 min). The reaction mixture was vigorously stirred at 25-30 °C for 2 h followed by 1 h string at 30 °C. Water (100ml) was added followed by sodium bicarbonate (3.7 g 0.044 mol) followed by sodium metabisulfite (8g). Organic layer was separated and aqueous phase was extracted with dichloromethane (3x20 ml). The combined organic extracts were dried over sodium sulfite, concentrated to dryness and triturated with hexane to form a solid. 83g

(94%) of target compound (3) was obtained.

m.p.: 70-72°

^{1}H-NMR: δ(ppm in $CDCl_3$)

8.05 (m, 2H, aromatic, *cis*-isomer)
7.95 (m, 2H, aromatic, *trans*-isomer)
7.56 (m, aromatic)
7.23 (m, aromatic)
4.77 (m, 4H, C_2-H, C_5-H, and C_2-$CH_2OOCC_6H_5$)
4.43 (m, 1H, C_5-H, *trans*-isomer)

4.09 (m, 1H, C_5-H, *cis*-isomer)

3.11 (m, 2H, C_4-H, *trans*-isomer)

2.75 (m, 2H, C_4-H, *cis*-isomer)

^{13}C-NMR: δ(ppm in $CDCl_3$)


cis-isomer:

166.64, 134.02, 130.42, 129.88, 129.06, 96.16, 68.83, 59.47 and 54.30

trans-isomer:

166.36, 134.12, 130.29, 129.68, 129.15, 108.07, 70.09, 61.83 and 53.47

3) (+/-)-Cis, trans-2benzoylmethyl-4-(N-acetylcytosine-1'-yl)-1,3-oxathiolane



(3) (4)

Compound (3) was dissolved in CH_2Cl_2 (20mL/g) and cooled to -15 °C. The amine (between 1 and 2 eq) was added followed by addition of TMSI (between 2 and 5 eq) while keeping the internal temperature below -5°C. stirred at -5°C to -10°C until the compound (3) disappeared. The CuCl (20%) and the pyrimidine (1.1eq) was added. The reaction mix was warmed up and kept between 5-10°C until TLC indicated the

reaction is complete. The reaction mixture was poured into 5% NH_4OH and stirred for 10 minutes until no solid precipitate was detected. The organic layer was separated and aqueous layer was extracted with CH_2Cl_2. The organic layer was washed with water, 2% HCl, dilute $Na_2S_2O_3$. The washed organic layer was evaporated to give the product, compound (4). Results for this step are shown in Table 2.

These were characterized by ^1H and ^{13}C-NMR.

cis-isomer:

^1H-NMR: δ(ppm in $CDCl_3$)

9.61 (b, 1H, C_4'-$NHCOCH_3$)
8.29 (d, 1H, C_6'-H)
8.06 (m, 2H, aromatic)
7.65 (m, 1H, aromatic)
7.51 (m, 2H, aromatic)
7.25 (d, 1H, C_5'-H)
6.61 (d, 1H, C_4-H)
5.50 (t, 1H, C_2-H)
4.80 (m, 2H, C_2-$CH_2OOCC_6H_5$)
4.48 (d, 1H, C_5-H)
4.05 (dd, 1H, C_5-H)
2.25 (s, 3H, CH_3)

^{13}C-NMR: δ(ppm in $CDCl_3$)

170.93, 166.28, 162.80, 155.76, 146.06, 133.91, 129.90, 128.84, 97.45, 85.88, 78.25, 64.60, 63.53 and 24.71.

trans-isomer:

^{1}H-NMR: δ(ppm in DMSO d_6)

10.88 (s, 1H, $C_{4'}$-$NHCOCH_3$)

8.13 (d, 1H, $C_{6'}$-H)

7.96 (m, 2H, aromatic)

7.68 (m, 1H, aromatic)

7.52 (m, 2H, aromatic)

7.20 (d, 1H, $C_{5'}$-H)

6.35 (d, 1H, C_4-H)

5.96 (dd, 1H, C_2-H)

4.58 (dd, 1H, C_2-$CH_2OOCC_6H_5$)

4.44 (d, 1H, C_5-H)

4.29 (m, 2H, C_5-H and $CH_2OOCC_6H_5$)

2.07 (s, 3H, CH_3)

^{13}C-NMR: δ(ppm in DMSO d_6)

171.53, 165.84, 162.76, 155.21, 146.59, 134.00, 129.64, 129.23, 96.54, 83.78, 74.24, 64.58, 64.01 and 24.35

Table 2

Pyrimidine	Base	Catalyst (molar eq)	Conditions	Yield (Cis + Trans)	Cis/ Trans	Yield % Cis
N-Ac-Cy 1.1eq	DIPEA 1.2eq		CH_2Cl_2, -15 °C RT, O/N	80%	2.0 : 1	53
N-Ac-Cy 1.1eq	DIPEA 1.2eq	$CuCl_2$ (20%)	CH_2Cl_2, -15 °C, 3h RT, O/N	91%	3.9 : 1	72
N-Ac-Cy 1.1eq	TEA 1.2eq	$CuCl_2$ (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	75%	3.8 : 1	59
N-Ac-Cy 1.1eq	DMCA 1.2eq	$CuCl_2$ (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	80%	3.4 : 1	62
N-Ac-Cy 1.1eq	DECA 1.2eq	$CuCl_2$ (20%)	CH_2Cl_2, -15 °C RT, O/N	71%	3.9 : 1	57
N-Ac-Cy 1.1eq	DIPEA 1.2eq	$CuCl_2$ (2%)	CH_2Cl_2, -15 °C, 3h RT, O/N	80%	2.4 : 1	56
N-Ac-Cy 1.1eq	DIPEA 1.2eq	$CuBr_2$ (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	80%	3.7 : 1	63
N-Ac-Cy 1.1eq	DIPEA 1.2eq	$Cu(acac)_2$ (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	85%	3.7 : 1	67
N-Ac-Cy 1.1eq	DIPEA 1.2eq	CuCl (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	80%	3.6 : 1	63
N-Ac-Cy 1.1eq	DIPEA 1.2eq	CuI (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	74%	3.5 :1	58
N-Ac-Cy 1.1eq	DIPEA 1.2eq	CuSCN (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	70%	3.1:1	53
N-Ac-Cy 1.1eq	DIPEA 1.2eq	$ZnBr_2$ (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	53%	3.1: 1	40
Cy 1.1eq	DIPEA 1.2eq	$CuCl_2$ (20%)	CH_2Cl_2, -15 °C, 5h RT, O/N	72%	2.4:1	51

Cy = cytosine

DIPEA = diisopropylethylamine


TEA = Triethylamine

DECA = diethylcyclohexylamine

DMCA = dimethylcyclohexylamine

$Cu(acac)_2$ = Copper (II) acetylacetonate

4) 2-hydroxymethyl-4-(cytosin-1'-yl)-1,3-oxathiolane




A suspension of the substrate, sodium methoxide, 10 mole percentage, and the appropriate solvent was stirred at RT for 2 h before it was filtered. The filter cake was dried and weighed before checking the C/T ratio (by ^1H NMR) and yield. Results for this step are shown in table 3.

Table 3

Weight of Cis/Trans mix (g)	*Cis/ Trans* ratio	Solvent	Strong base	Conditions	% of trans (NMR)	Recovery of Cis (%)
4	3.4 : 1	MeOH,	MeONa	Mix and stir at RT	0	89
10	3.4 : 1	MeOH	MeONa	Mix and stir at RT	0	88.5
4	3.4 : 1	MeOH/tBME (7 :3)	MeONa	Mix and stir at RT	0	90
25	3.4 : 1	MeOH/tBME (7 :3)	MeONa	Mix and stir at RT	traces	91
1	3.4 : 1	MeOH/EtOH (1 :1)	MeONa	RT	0	98
4	3.4 : 1	MeOH/EtOH (1 :1)	MeONa	Mix and stir at RT	traces	92.5
25	3.4 : 1	MeOH/EtOH (1 :1), 150 mL	MeONa	Mix and stir at RT	traces	93
1	3.4 : 1	MeOH/Toluene (6 :4)	MeONa	RT	traces	92.5
4	3.4 : 1	MeOH :Toluene (6 :4)	MeONa	Mix and stir at RT	traces	90
25	3.4 : 1	MeOH/Toluene (6 :4) L	MeONa	Mix and stir at RT	traces	91
1	3.4 : 1	MeOH/H2O (95 :5)	NaOH	RT	20	90
1	3.4 : 1	MeOH/H2O (95 :5)	NaOH	At 50 °C	20	90

5) Synthiesis of 2benzoylmethyl-4-(N-acetylcytosine-1'-yl)-1,3-oxathiolane using (-) or (+) 2-benzoyloxymethyl-1,3-oxathiolane-S-oxide



(3a) (4a)

The enantiomerically pure Compound (3a) was dissolved in CH_2Cl_2 (10mL/g) and cooled to reaction temperature. The amine was added followed by addition of TMSI while keeping the internal temperature below -5°C. stirred at -5°C to -40°C until the compound (3a) disappeared. The CuCl (20%) and the pyrimidine (1.1eq) was added. The reaction mix was warmed up and kept between - 40 to 30°C until TLC indicated the reaction is complete. The reaction was cooled to 0°C-5°C. Celite (12g, 100% w/w) was added to the suspension and stirred. Concentrated ammonium hydroxyde (was added slowly and the suspension temperature was kept between 0 °C - 10 °C. Stirred at 0°C-5°C. The suspension was filtered and the cake was resuspended in dichloromethane. Stirred then filtered. The phases were separated and the aqueous was extracted with dichloromethane . The combined organic layers were washed with a 2% solution of ammonium hydroxyde, water , 2% hydrochloric solution , 2% sodium metabisulfite solution and saturated sodium bicarbonate solution. The organic phase was dried over magnesium sulfate, filtered then reduced in volume *in vacuo* to give

crude base-coupled material as a beige solid. The solids were dissolved in EtOAc and allowed to crystallize. The suspension was stirred at 0-5°C then filtered. The solids were dried *in vacuo* to give pure base-coupled product as a pale beige solid. Results for this step are shown in Table 4 and Table 5.

^{1}H NMR (300MHz)($CDCl_3$) δ: 2.27 (s, 3H, CH_3 cis), 2.29 (s, 3H, CH_3 trans), 4.06 (dd, J= 4.5 Hz & J=10.8 Hz, 1H, C5 cis), 4.30 (dd, J= 3.4 Hz & J=12.4 Hz, 1H, C5 trans), 4.49 (dd, J=3.1 Hz & J=10.8 Hz, 1H, C5 cis) 4.72 (dd, J=8.3 Hz & J=12.4 Hz, 1H, C5 trans), 4.77, (AB, J=4.5 Hz, 2H, CH_2OBz cis), 4.84 (AB, J=2.3 Hz, 2H, CH_2OBz trans), 5.50 (dd, J= 3.1 Hz & J=4.5 Hz, 1H, C4 cis), 5.92 (dd, J= 3.4 Hz & J=8.3 Hz, 1H, C4 trans), 6.61 (dd, J= 2.3 Hz trans & J=4.5 Hz cis, 1H, C2), 7.25 (d, J=7.5 Hz, 1H, C5'), 7.5-8.1 (m, 10H, aromatic cis & trans), 8.30 (d, J=7.5 Hz, 1H, C6'), 9.50 (s 1H, NH).

^{13}C NMR (300MHz)($CDCl_3$) δ: 25.9 (cis & trans), 64.3 (cis), 65.1 (trans), 65.3 (cis & trans), 76.1 (trans), 78.7 (cis), 84.7 (trans), 86.2 (cis), 97.9 (cis), 98.1 (trans), 128.6 (cis & trans), 128.7 (cis & trans), 129.2 (cis & trans), 129.4 (cis & trans), 129.8 (cis & trans), 133.5 (trans), 133.7 (cis), 145.3 (trans), 145.6 (cis), 155.2 (trans), 155.3 (cis), 162.5 (cis & trans), 162.6 (trans), 165.7 (cis), 171.0 (cis), 171.1 (trans).

Table 4











Sulfoxide	Base	Reaction Temp. (°C)	(-)Cis (%)	(+)Cis (%)	(-)Trans (%)	(+)trans (%)	Cis/ Trans (%)
	DIPEA	-15	48.9	27.4	6.7	17	3.2 :1
	TEA	-15	70.4	5.2	1.6	22.8	3.1 :1
	TEA	-15	4.9	73.6	20	1.5	3.6 :1
	DMCA	-25	72.4	4.5	1.25	21.8	3.3 :1
	TEA	- 30 (internal temp), warm up to RT after adding silylated base	1	74.4	24.6	0	3 :1

Table 5

Sulfoxide	Base	Reaction Temperature °C	Yield %	Cis/Trans (%)
	TEA +DIPEA	-20	74	3.3 : 1
	(i-pro) NMe_2	-20	72	3.7 : 1
	TEA	-20	63	4 : 1
	(i-pro) NMe_2	-20	72	3.3 : 1
	TEA	-20	70	3.3 : 1

TEA = Triethylamine


DIPEA = Diisopropylethylamine

DMCA = Dimethylcyclohexylamine

(i-pro)NMe_2 = Isoproplydimethylamine

CLAIMS:

1) A stereoselective process for making predominantly cis nucleosides or nucleoside analogues and derivatives of formula (A):




(A)

wherein

R1 is a pyrimidine base or a pharmaceutically acceptable derivative thereof; and

Q is carbon, oxygen or sulfur;

consisting of coupling step of the a compound of formula (B):




(B)

wherein

R2 is C_{7-10} aralalkyl, C_{1-16} acyl or $Si(Z^1)(Z^2)(Z^3)$, where Z^1, Z^2 and Z^3 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-6} alkoxy or C_{6-20} aryloxy; C_{7-20} aralkyl optionally substituted by halogen, C_{1-20} alkyl or C_{1-20} alkoxy; C_{6-20} aryl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-20} alkyl or C_{1-20} alkoxy; trialkylsilyl; fluoro; bromo; chloro and iodo;

with a base R1, wherein R1 is a pyrimidine base or a pharmaceutically acceptable derivative thereof;

in a C_{1-10} coupling solvent in the presence of a catalytic amount of an element or combination of elements of group 1B or 11B of the periodic table; a tertiary amine and a Lewis acid to yield an intermediate of formula (D):




(D).

wherein R1, R2 and Q are defined above; and

a second step to deprotect the intermediate for formula (D) to yield of the *cis* nucleosides or nucleoside analogues or derivatives of formula (A).

2) A process according to claim 1 wherein the Lewis acid is a compound of formula (C):



(C)


wherein

R4, R5 and R6 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-6} alkoxy or C_{6-20} aryloxy; C_{7-20} aralkyl optionally substituted by halogen, C_{1-20} alkyl or C_{1-20} alkoxy; C_{6-20} aryl optionally substituted by fluoro, bromo, chloro, iodo, C_{1-20} alkyl or C_{1-20} alkoxy; trialkylsilyl; fluoro; bromo; chloro and iodo; and


R7 is selected from the group consisting of fluoro; bromo; chloro; iodo; C_{1-20} sulphonate esters, optionally substituted by fluoro, bromo, chloro or iodo; C_{1-20}

alkyl esters optionally substituted by fluoro, bromo, chloro or iodo; triiodide; a silyl group of the general formula (R4)(R5)(R6)Si (wherein R4, R5, and R6 are as defined above); C_{6-20} arylselenenyl; C_{6-20} arylsulfenyl; C_{6-20} alkoxyalkyl; and trialkylsiloxy.


3) A process according to claim 1 wherein R1 is selected from:




(I)




(II)




(III)




(IV)




(V)



(VI)



(VII)



(VIII)

wherein:

x is oxygen, NH or sulfur;

y is oxygen, NH or sulfur;

R8 and R9 are independently selected from hydrogen, hydroxyl, amino, C_{1-6} alkyl or C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-10} acyl, C_{6-10} aryl, C_{6-11} carbonylaryl, C_{1-7} carbonyloxyalkyl, C_{6-11} carbonyloxyaryl, C_{2-7} carbonylaminoalkyl, or amino acids;

R8 may be a saturated or unsaturated C_{3-8} carbocyclic ring optionally substituted with COOH, $C(O)NH_2$,

OH, SH, NH_2, NO_2, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halogen, C(O)R14 wherein R14 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and C(O)OR15 wherein R15 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl; and R9 is chosen from H, C_{1-6} alkyl, C_{2-6} alkenyl and C_{2-6} alkynyl;

R_8R_9 can also be connected to the nitrogen atom to form a saturated or unsaturated C_{3-8} heterocyclic ring optionally substituted with C(O)OH, $C(O)NH_2$, OH, SH, NH_2, NO_2, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halogen, C(O)R14 wherein R14 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and C(O)OR15 wherein R15 is a C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl; and

R10 , R11, R12 and R13 are each independently selected from hydrogen, halogen, hydroxyl, amino, cyano, carboxyl, carbamoyl, C_{2-7} alkoxycarbonyl, hydroxymethyl, trifluoromethyl, C_{6-10} arylthio, C_{1-6} alkyl, C_{2-6} alkenyl substituted or unsubstituted with halogen or azido, C_{2-6} alkynyl, C_{1-6} acyloxy, thiocarboxy, thiocarbamoyl, carbamate, ureido, amidino, or C_{6-10} aryloxy.

4) A process according to claim 1 wherein the element or combination of elements of groups IB or IIB is Cu, Ag, Au, Zn or Cd.

5) A process according to claim 1 wherein the tertiary amine is of the form $N(Z^4)(Z^5)(Z^6)$ wherein $(Z^4),(Z^5),(Z^6)$ are independently selected from the group consisting C_{1-6} alkyl optionally substituted with C_{1-3} alkyl, C_{6-10} aryl or halogen.

6) A process according to claim 1 wherein the organic solvent is a. C_{1-8} chloroalkyl, a C_{1-8} chloroalkenyls, a C_{6-10} chloroaryl, C_{1-8} alkylonitrile or mixtures thereof.

7) A process according to claim 6 wherein the organic solvent is selected from the group consisting of chloromethanes, chloroethanes, methanonitriles and mixtures thereof.

8) A process according to claim 5 wherein the tertiary amine is triethylamine, diethylcyclohexylamine, diethylmethylamine, dimethylethylamine, dimethylisopropylamine, dimethylbutylamine, dimethylcyclohexylamine, tributylamine, diethylmethylamine, dimethylisopropylamine, diisopropylethylamine or combinations thereof.

9) A process according to claim 4 wherein the element or combination of elements of groups IB or IIB is Cu, Zn or combinations thereof.

10) A process according to claim 1 wherein Q is oxygen.

11) A process according to claim 1 wherein the coupling step is preformed at a temperature equal to or less than 30 degrees Celsius.

12) A process according to claim 1 wherein the deprotection step is preformed in the presence of a deprotecting agent and a deprotecting solvent which favors the crystallization of the *cis* nucleosides or nucleoside analogues or derivatives of formula (A).


13) A process according to claim 12 wherein the deprotection agent is alkaline.

14) A process according to claim 13 wherein the deprotection agent is sodium hydroxide, sodium methoxide, ammonium hydroxide, potassium hydroxide, lithium hydroxide, methanolic ammonia or combinations thereof.


15) A process according to claim 12 wherein the deprotecting solvent is water, methanol, ethanol, toluene, tert-butyl methyl ether or combinations thereof.

16) A process according to claim 3 wherein R1 is selected from N^4 -alkylpyrimidines, N^4 -acylpyrimidines, 4-halopyrimidines, N^4 -acetylenic pyrimidines, 4-amino and N^4 -acyl pyrimidines, 4-hydroxyalkyl pyrimidines, 4-thioalkyl pyrimidines, thymine, cytosine, 6-azapyrimidine, 2- mercaptopyrimidine, 4-mercaptopyrimidine, uracil, C^5 -alkylpyrimidines, C^5 -benzylpyrimidines, C^5 halopyrimidines, C^5-vinylpyrimidine, C^5 -acetylenic pyrimidine, C^5 -acyl pyrimidine, C^5 -amidopyrimidine, C^5 -cyanopyrimidine, C^5 -nitropyrimidine, C^5 -aminopyrimidine, 5-azacytidinyl, 5-azauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl or a pharmaceutically acceptable derivative thereof.

17) A process according to claim 3 wherein R2 is



wherein W is halogen, C_{1-16} alkyl, C_{2-16} alkoxyalkyl, C_{6-10} aryl, C_{1-16} alkoxy or nitro; or



wherein E is C_{6-10} aryl, C_{1-16} alkoxy, C_{2-16} alkoxyalkyl or C_{1-16} alkyl; or

$Si(Z^1)(Z^2)(Z^3)$, where Z^1, Z^2 and Z^3 are independently selected from the group consisting of hydrogen; C_{1-20} alkyl optionally substituted by fluoro, bromo, chloro, iodo; C_{7-10} aralalkyl optionally substituted by fluoro, bromo, chloro or iodo; and C_{6-10} aryl optionally substituted by fluoro, bromo, chloro or iodo.