

UNIT DOSAGE OF APADENOSON

5 CROSS-REFERENCE TO RELATED APPLICATIONS


The present application claims the priority benefits of U.S. Provisional Applications No. 61/078,169, filed 3 July 2008 and No. 61/155,937, filed 27 February 2009, which are expressly incorporated fully herein by reference.

10 FIELD OF THE INVENTION

The present invention relates to a unit dosage of Apadenoson, a pharmacological stress agent, and use of the same as a pharmacologic agent for myocardial perfusion imaging.

15 BACKGROUND OF THE INVENTION

Apadenoson, shown below, was first described as a pharmacologic stress agent



that can be used in clinical perfusion imaging techniques (e.g., for diagnosing
20 and assessing the extent of coronary artery disease) in US 6,322,771. This agent
has since been taken into Phase I and II clinical trials. In 2005, Dr. Hendel et al
reported to the American Heart Association on the preliminary results of 127
patient SPECT Tc^{99m} sestamibi imaging studying comparing adenosine with
Apadenoson using either 1 µg/kg or 2 µg/kg intravenous boluses of Apadenoson.
25 The report concluded that Apadenoson was safe and well-tolerated and worthy
of Phase III evaluation. In 2006, Dr. Kern et al reported to the American Heart
Association the results of a Phase II study of Apadenoson, one goal of which
was to determine an appropriate dose for Phase III clinical trials. Intravenous
bolus dosages of 0.5, 1.0, 2.0, and 2.5 µg/kg were studied. For a patients, the
30 average peak velocity for coronary blood flow was shown to increase with a
corresponding increase in dosage from 0.5 to 2 µg/kg (see Figure 1). In light of

this data, it was believed that Apadenoson would need to be administered on a weight basis, not a unit dose basis.

There are inherent limitations and opportunities for operator error when parenterally administering a pharmaceutical agent on a weight basis. This type of dosing requires calculating the amount of agent to administer based on a patient's weight, administering the calculated amount from a larger dose, and disposing of any left over agent. Thus, it is desirable and beneficial for a pharmaceutical agent to be provided in a unit dose.

10 SUMMARY OF THE INVENTION

The present invention provides a novel unit dose of Apadenoson suitable for parenteral administration.

The present invention also provides a novel method of diagnosing myocardial dysfunction using a unit dose of Apadenoson as a pharmacologic stress agent.

These and other aspects of the present invention have been accomplished in view of the discovery that no dose response curve is seen when 1 $\mu\text{g/kg}$ or 2 $\mu\text{g/kg}$ of Apadenoson is administered.

20 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the average peak flow wire velocity of one patient from a Phase II clinical trial study. In this Phase II study in 100 patients, conducted by Dr. Morton Kern, an independent investigator at UC Irvine, adenosine was administered by intracoronary injection, and coronary blood flow velocity was monitored with a flow wire. On the left, the three injections of adenosine increased blood flow velocity consistent with the well characterized pharmacology of adenosine. On the right, increasing bolus doses of Apadenoson, at doses shown to be safe, achieved peak flow equivalent to adenosine.

Figure 2 shows the average peak flow wire velocity of 33 patients from the study described in Figure 1.

DETAILED DESCRIPTION OF THE INVENTION

The previously reported increase in coronary blood flow velocity (CBFV) corresponding to an increase in dose of Apadenoson from 0.5 to 2 $\mu\text{g/kg}$ was based on the increase in CBFV in a limited number of patients (see Figure 1). However, it has now been found that if one considers the entire sample of patients, as shown in Figure 2, then at dose $\geq 1 \mu\text{g/kg}$ and in particular between 1 and 2.5 $\mu\text{g/kg}$ no dose response is seen. In light of this, Applicant has surprisingly discovered that instead of the weight-based dosing predicted by the results reported in Figure 1, Apadenoson can actually be administered via a unit dose.

Thus, in an embodiment, the present invention provides a novel unit dose of Apadenoson, comprising: (a) Apadenoson and (b) a pharmaceutically acceptable carrier, wherein the unit dose is suitable for parenteral administration.

In another embodiment, the unit dose is suitable for intravenous administration.

In another embodiment, the amount of Apadenoson present in the unit dose is selected from 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, and 175 μg . Additional examples of the weight of Apadenoson present include (a) 100, 110, 120, 130, 140, and 150 μg ; (b) 100 μg ; and (c) 150 μg .

In another embodiment, the amount of Apadenoson present in the unit dose is in the range selected from 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, to 125 μg . Additional examples of the weight of Apadenoson present include (a) 115, 120, and 125 μg and (b) 120 μg .

In another embodiment, the pharmaceutical carrier, comprises: a cyclodextrin. Examples of cyclodextrins include α -CD or derivatives thereof (e.g., α -hydroxypropyl-CD (HP- α -CD)), β -CD or derivatives thereof (e.g., β -hydroxypropyl-CD (HP- β -CD), methylated β -cyclodextrin, hydroxyethyl- β -

cyclodextrin, and sulfobutylether β -CD), and γ -CD or derivatives thereof (e.g., γ -hydroxypropyl-CD (HP- γ -CD)).

Examples of the concentration of CD (e.g., hydroxypropyl- β -cyclodextrin) include being within the range selected from (a) about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10% w/v; (b) about 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, to 4% w/v; (c) about 1% w/v; and, (d) about 2% w/v of the final formulation.

In another embodiment, the pharmaceutical carrier, comprises: buffered saline. A useful buffer is a citrate buffer (e.g., sodium citrate). Citric acid can be useful to adjust the pH of the unit dose. As an example, the pharmaceutical carrier, comprises: buffered saline, comprising: saline, sodium citrate, and citric acid. One would recognize that citric acid may not be present in the final unit dose due to ionization.

In another embodiment, the pH of the unit dose is selected from 4.6, 4.7, 4.8, 4.9, to 5.0. Another example of the pH of the unit dose is 4.8.

In another embodiment, the volume of the unit dose is selected from 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, and 5 mL. Another example of the volume is from 2, 3, 4, to 5 mL.

In another embodiment, the unit dose, comprises:

- (a) 100 μ g Apadenoson;
- (b) a pharmaceutically acceptable carrier, comprising:
 - (b_i) 2% w/v HP- β -CD;
 - (b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to pH 4.8; and,
 - (b_{iii}) saline in an amount to form a 1-5 mL unit dose.

In another embodiment, the unit dose, comprises:

- (a) 100 μ g Apadenoson;
- (b) a pharmaceutically acceptable carrier, comprising:
 - (b_i) 1% w/v HP- β -CD;
 - (b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to pH 4.8; and,
 - (b_{iii}) saline in an amount to form a 1-5 mL unit dose.

In another embodiment, the unit dose, comprises:

- (a) 150 μ g Apadenoson;
(b) a pharmaceutically acceptable carrier, comprising:
 (b_i) 2% w/v HP- β -CD;
 (b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to
5 pH 4.8; and,
 (b_{iii}) saline in an amount to form a 1-5 mL unit dose.

In another embodiment, the unit dose, comprises:

- (a) 150 μ g Apadenoson;
(b) a pharmaceutically acceptable carrier, comprising:
10 (b_i) 2% w/v HP- β -CD;
 (b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to
pH 4.8; and,
 (b_{iii}) saline in an amount to form a 1-5 mL unit dose.

The unit dose of the present invention can be filled into any useful
15 container for storage, transportation, and usage. An example of a useful
container is a syringe body.

In another embodiment, the present invention provides a novel method of
diagnosing myocardial perfusion abnormalities in a mammal, comprising:

- (a) parenterally administering to the mammal a unit dose of Apadenoson;
20 and
(b) performing a technique on the mammal to detect the presence of
coronary artery stenoses, assess the severity of coronary artery stenoses,
or a combination thereof.

In another embodiment, the patient weighs at least 40 kg.

25 In another embodiment, the administration is intravenous administration.

In another embodiment, the technique is an imaging technique selected
from: planar or single photon emission computed tomography (SPECT), gamma
camera scintigraphy, positron emission tomography (PET), nuclear magnetic
resonance (NMR) imaging, magnetic resonance imaging (MRI) imaging,
30 perfusion contrast echocardiography, digital subtraction angiography (DSA), and
ultrafast X-ray computed tomography (CINE CT).

In another embodiment, the present invention provides a prefilled
syringe, comprising: a syringe and a unit dose of Apadenoson, comprising: (a)

Apadenoson and (b) a pharmaceutically acceptable carrier, wherein the unit dose is suitable for parenteral administration. The syringe can be any known syringe useful for parenteral administration. For example, the syringe can comprise: a body and a plunger movably disposed within the body. The body can be
5 cylindrical with a first open end to receive the plunger and a second end opposite the first, with the second end modified with an opening sufficient for the unit dose to pass through. The syringe can further comprise: a needle (e.g., an injection needle). The needle can be detachably connected to or permanently fixed to the body. A needle guard can also be present

10 In another embodiment, the present invention provides a novel unit dose of Apadenoson for use in medical therapy.

In another embodiment, the present invention provides a novel use of a unit dose of Apadenoson for the manufacture of a medicament for use in diagnosing myocardial perfusion abnormalities in a mammal.

15 The Apadenoson unit dose of the present invention can be administered as a pharmacological stress agent and used in conjunction with any one of several noninvasive diagnostic procedures to measure aspects of myocardial, coronary, and/or ventricular perfusion. Aspects that can be measured include coronary artery stenoses, myocardial dysfunction (e.g., myocardial ischemia,
20 coronary artery disease, ventricular dysfunction, and differences in blood flow through disease-free coronary vessels and/or stenotic coronary vessels), myocardial contractile dysfunction, the presence of regional wall motion abnormalities, the functional significance of stenotic coronary vessels, coronary artery disease, ischemic ventricular dysfunction, and vasodilatory capacity
25 (reserve capacity) of coronary arteries in humans. Radiopharmaceuticals are typically used in diagnostic method methods. The radiopharmaceutical agent may comprise, for example, a radionuclide selected from the group consisting of thallium-201, technetium-99m, nitrogen-13, rubidium-82, iodine-123 and oxygen-15.

30 Any embodiment or feature of the present invention whether characterized as preferred or not characterized as preferred may be combined with any other aspect or feature of the invention, whether such other feature is characterized as preferred or not characterized as preferred.

Definitions

The examples provided in this application are non-inclusive unless otherwise stated. They include but are not limited to the recited groups.

Unit dose means the amount of a medication administered to a patient in a single dose. A unit dose is typically independent of the weight of the patient or may be associated with a specified weight range (e.g., ≥ 40 kg).

The indefinite articles "a" and "an" mean "at least one" or "one or more" when used in this application, including the claims, unless specifically indicated otherwise.

Mammal and patient covers warm blooded mammals that are typically under medical care (e.g., humans and domesticated animals). Examples of mammals include (a) feline, canine, equine, and bovine and (b) human.

Parenteral includes intravenous, intramuscular, and subcutaneous routes.

Dosage and formulation

Sterile injectable solutions are typically prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization (or some other form of sterilization). In the case of sterile powders for the preparation of sterile injectable solutions, the methods of preparation include vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

All patents, patent applications, books and literature cited in the specification are hereby incorporated by reference in their entirety. In the case of any inconsistencies, the present disclosure, including any definitions therein will prevail.

Numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

WHAT IS CLAIMED IS:

1. A unit dose of Apadenoson, comprising: (a) Apadenoson and (b) a pharmaceutically acceptable carrier, wherein the unit dose is suitable for parenteral administration.
- 5 2. The unit dose of claim 1, wherein the amount of Apadenoson present is selected from 76-175 μg .
3. The unit dose of claim 2, wherein the amount of Apadenoson present is
10 selected from 100-150 μg .
4. The unit dose of claim 3, wherein the amount of Apadenoson present is 100 μg .
- 15 5. The unit dose of claim 3, wherein the amount of Apadenoson present is 150 μg .
6. The unit dose of claim 1, wherein the pharmaceutical carrier, comprises: a cyclodextrin (CD).
- 20 7. The unit dose of claim 6, wherein the cyclodextrin is β -hydroxypropyl-CD.
8. The unit dose of claim 6, wherein the concentration of CD is selected
25 from 0.1-10% w/v of the final formulation.
9. The unit dose of claim 8, wherein the concentration of CD is 1% w/v.
10. The unit dose of claim 8, wherein the concentration of CD is 2% w/v.
- 30 11. The unit does of claim 1, wherein the pharmaceutical carrier, comprises: buffered saline.

12. The unit dose of claim 11, wherein the pharmaceutical carrier, comprises:
buffered saline, comprising: saline and sodium citrate.
13. The unit dose of claim 1, wherein the pH of the unit dose is selected from
4.6-5.0.
14. The unit dose of claim 13, wherein the pH is 4.8.
15. The unit dose of claim 1, wherein the volume of the unit dose is selected
from 1-5 mL.
16. The unit dose of claim 1, comprising:
(a) 100 µg Apadenoson;
(b) a pharmaceutically acceptable carrier, comprising:
(b_i) 2% w/v HP-β-CD;
(b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to
pH 4.8; and,
(b_{iii}) saline in an amount to form a 1-5 mL unit dose.
17. The unit dose of claim 1, comprising:
(a) 100 µg Apadenoson;
(b) a pharmaceutically acceptable carrier, comprising:
(b_i) 1% w/v HP-β-CD;
(b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to
pH 4.8; and,
(b_{iii}) saline in an amount to form a 1-5 mL unit dose.

18. The unit dose of claim 1, comprising:
(a) 150 µg Apadenoson;
(b) a pharmaceutically acceptable carrier, comprising:
 (b_i) 2% w/v HP-β-CD;
5 (b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to
pH 4.8; and,
 (b_{iii}) saline in an amount to form a 1-5 mL unit dose.
19. The unit dose of claim 1, comprising:
10 (a) 150 µg Apadenoson;
(b) a pharmaceutically acceptable carrier, comprising:
 (b_i) 2% w/v HP-β-CD;
 (b_{ii}) sodium citrate buffer in an amount to buffer the unit dose to
pH 4.8; and,
15 (b_{iii}) saline in an amount to form a 1-5 mL unit dose.
20. A method of diagnosing myocardial perfusion abnormalities in a
mammal, comprising:
(a) parenterally administering to the mammal a unit dose of Apadenoson;
20 and
(b) performing a technique on the mammal to detect the presence of
coronary artery stenoses, assess the severity of coronary artery stenoses,
or a combination thereof.
- 25 21. The method of claim 20, wherein the patient weighs at least 40 kg.