

ANTIBODIES THAT BIND TO SORTILIN AND INHIBIT THE BINDING OF PROGRANULIN

Field of The Invention

[0001] The present invention relates to monoclonal anti-Sortilin antibodies useful in correcting a deficient level of progranulin (PGRN). In particular these
5 antibodies can be used in the treatment of frontotemporal dementia (FTD) and amyotrophic lateral sclerosis (ALS). Furthermore, it anticipated that the monoclonal antibodies may also be useful to treat neurodegenerative disorders such as Alzheimer's Disease (AD).

Reference To Sequence Listing

10 [0002] This application includes one or more Sequence Listings pursuant to 37 C.F.R. 1.821 et seq., which are disclosed in computer-readable media (file name: 0993_ST25.txt, created on 22 June 2016, and having a size of 144 kB), which file is herein incorporated by reference in its entirety.

Background of the Invention

15 [0003] Sortilin is a receptor that has been reported to mediate pro-apoptotic effects of pro-neurotrophins and to mediate trafficking and sorting of neurotrophin receptors (Nykjær et al, 2012, Trends Neurosci. 2012;35(4):261-70; Glerup et al, Handb Exp Pharmacol, 2014;220:165-89, Carlo et al, J Mol Med (Berl). 2014 Sep;92(9):905-11). A number of sortilin ligands have been
20 identified including neurotensin for which a high affinity binding site was localized by x-ray crystallography to inside a beta propeller tunnel in the sortilin molecule (Quistgaard et al, Nat Struct Mol Biol. 2009 Jan; 16(1):96-8; Quistgaard et al, Protein Sci. 2014, Sep;23(9):1291-300). More recently, sortilin was shown to function as a high affinity receptor for the growth factor
25 progranulin (PGRN, Hu *et al.* Neuron. 2010 Nov 18;68(4):654-67.

[0004] PGRN ((proepithelin, granulin-epithelin precursor, PC-cell-derived growth factor, acrogranin)) is a secreted glycosylated protein with anti-inflammatory and neurotrophic-like actions (For a recent review, see Nguyen, Trends Endocrinol Metab. 2013 Dec;24(12):597-606). PGRN is proteolytically

cleaved to granulins, but much remains to be learned regarding the physiological role of PGRN and granulins and the identity of their receptors. PGRN has been implicated in several cellular functions including cell cycle regulation and cell motility (He, Z. & Bateman, A., J. Mol. Med. 57:600-612
5 (2003); Monami, G., *et al.*, Cancer Res. (5(5:7103-7110 (2006)), wound repair, inflammation (Zhu, J., *et al.*, Cell 777:867-878 (2002)), induction of growth factors such as vascular endothelial growth factor (VEGF) (Tangkeangsinsin, W. & Serrero, G, Carcinogenesis 25.1587-1592 (2004)), and tumorigenesis (He, Z. & Bateman, A., J. Mol. Med. 81:600- 612 (2003), Monami, G., *et al.*, Cancer
10 Res (5(5:7103-7110 (2006); Serrero, G., Biochem Biophys. Res. Commun. 505-409-413 (2003), Lu, R & Serrero, G., Proc. Natl Acad Sci U.SA 98 142-147 (2001); Liao, L M., *et al.*, Cancer Res. 60:1353-1360 (2000)). PGRN has been reported to bind the TNF receptor (Tang W *et al.*, Science 2011, 332(6028):478-84). but this observation has been challenged by others (Chen *et al.*, J
15 Neurosci. 2013, 33(21):9202-9213).

[0005] The binding of PGRN to sortilin has been mapped to the neurotensin site and reported to be mediated solely through the PGRN C-terminal domain (Zheng *et al.* PLoS One. 2011;6(6):e21023; Lee *et al.* Hum Mol Genet. 2013) in a manner similar to neurotensin and in accordance, neurotensin has been
20 shown to block the interaction of sortilin with PGRN and other ligands. Upon binding, sortilin mediates lysosomal clearance of PGRN and thereby regulates extracellular PGRN levels (Hu *et al.* 2010). Thus, knockdown or overexpression of sortilin have been shown to regulate extracellular PGRN levels in cell culture (Carrasquillo *et al.* Am J Hum Genet. 2010 Dec 10;87(6):890-7) and in mice,
25 sortilin deficiency was reported to increase PGRN levels and to restore plasma and brain PGRN-levels in PGRN +/- mice (Hu *et al.* 2010). Interestingly, a single nucleotide polymorphism (SNP) near sortilin was associated with decreased plasma PGRN and increased sortilin mRNA levels (Carrasquillo *et al.* Am J Hum Genet. 2010 Dec 10;87(6):890-7). These observations suggest that sortilin
30 is a key regulator of extracellular PGRN.

[0006] PGRN has been linked to frontotemporal dementia (FTD), a progressive dementia characterized by behavioral and semantic changes, as well as frontotemporal lobar degeneration (FTLD) and neuronal inclusions containing TAR DNA Binding Protein-43 (TDP-43) or tau inclusions (Baker et al, 5 2006, Nature. 2006 Aug 24;442(7105):916-9; Cruts et al, Nature 442: 920-924 (2006); Am J Hum Genet. 2010 Dec 10;87(6):890-7, M et al, Trends in Genetics 24: 186-194 (2008)). The majority of sporadic and familial FTD cases show TDP-43 pathology (~50%) similar to ALS and FTD-TDP43 and ALS are by some considered to constitute a disease spectrum (Ito D Neurology. 2011 Oct 10 25;77(17):1636-43; Boxer AL et al, Alzheimers Dement. 2013 Mar;9(2):176-88; Rademakers et al, Nat Rev Neurol. 2012 August; 8(8): 423–434) due to common pathologies and genetic factors and some overlap in symptomatology. No disease-modifying treatment options are available for FTD. A subset of frontotemporal dementia patients with TDP-43 pathology have loss of function mutations in the granulin gene (GRN) resulting in PGRN haplo-insufficiency. To 15 date, 69 different mutations in the granulin gene, all resulting in reduced PGRN levels and/or function, have been associated with FTD and it is believed that raising extracellular PGRN in plasma and brain would counteract the disease process.

20 [0007] PGRN mutations have also been linked with Alzheimer's disease (AD) (Sheng *et al.*, 2014, Gene. 2014 Jun 1;542(2):141-5; Brouwers et al., 2008, Neurology. 2008 Aug 26;71(9):656-64) suggesting that PGRN deficiency may play an important role in AD pathogenesis. Furthermore, neuroprotective effects of PGRN in mouse AD models have been observed (Minami et al, 2014, 25 Nat Med. 2014 Oct;20(10):1157-64) providing support for the view that enhanced PGRN may be beneficial in AD and possibly other neurodegenerative conditions.

[0008] The present application describes the generation and identification of anti-human Sortilin antibodies which can regulate PGRN in cellular models and 30 in mice. Those antibodies surprisingly bind to a region on Sortilin which is distant to the previously reported progranulin binding site, the so-called

neurotensin-site, and yet are capable of inhibiting Sortilin-PGRN interaction and of thereby increasing extracellular PGRN.

[0009] The inventors have defined six Sortilin binding regions and surprisingly identified that the most efficacious antibodies bind a region ("region D"). As PGRN has neuroprotective and anti-inflammatory effects, the inventors' findings indicate that such antibodies targeting Sortilin are likely to have a beneficial effect in the treatment of a range of neurodegenerative disorders including FTD/FTLD. A subgroup of these patients carry a mutation in the gene encoding PGRN leading to haploinsufficiency. Sortilin antibodies are therefore likely to have similar therapeutic benefits for patients suffering from other TDP-43 proteinopathies and diseases in which PGRN levels may influence TDP43-function and pathology, including ALS and AD.

Summary of invention

[0010] The inventors of the present invention have generated monoclonal antibodies which are able to inhibit the binding of PGRN to Sortilin, and which surprisingly bind to a novel Sortilin region denominated the "D-region" as defined in **SEQ ID NO:170**. Several antibodies identified by the inventors have properties similar to the D-region antibodies and experimental evidence indicates that they also bind within the D-region, and those antibodies are herein referred to as D+ antibodies. Accordingly, in one aspect, there is provided an antibody, or an antigen-binding fragment thereof, capable of binding specifically to Sortilin and inhibiting or reducing the binding of PGRN to Sortilin. In another aspect, there is provided a pharmaceutical composition comprising an antibody, or an antigen-binding fragment thereof as described herein, or the preparation as described herein, and a pharmaceutically-acceptable carrier. Accordingly, in one embodiment, the invention relates to such antibodies, to compositions and/or kits comprising such antibodies, and to methods and uses thereof.

[0011] The disclosure also relates to a method of preventing or treating a disease associated with decreased PGRN levels in the brain of a patient, comprising administering an effective dosage of an antibody or an antigen-binding fragment thereof that binds to the D-region of Sortilin. These diseases include *i.a.* FTD, ALS and TDP43 proteinopathies such as AD.

Brief Description of the Drawings

[0012] **Figure 1** provides an overview of the generation of region assignment of human antibodies based on Sortilin region binding, effect on PGRN-binding and PGRN-levels and on cross-blocking between the antibodies.

[0013] Sortilin-binding antibodies were selected and assigned to regions A-E based on binding to shuffle constructs in which the sequence corresponded to the tetraodon Sortilin sequence within selected regions of the protein (**Example 1, Figure 2**).

[0014] 20 antibodies which inhibited Sortilin-PGRN binding (as measured by HTRF analysis) were selected (see **Example 10, Figure 5** and **Figure 6**). 15 of these antibodies were D-region antibodies while 3 were D+ antibodies (**Figure 6**). Subsequent cross-blocking analysis showed that the 18 D-region and D+ antibodies (D+ antibodies have a different binding pattern to shuffle constructs than D-region antibodies. Nevertheless D+ antibodies cannot with certainty be assigned to a binding region A-E as outlined above but shared functional characteristics (cell assays etc.) similar to the D-region antibodies), all cross-blocked each other supporting that they interacted with the same Sortilin region (**Example 9, Figure 7**). When sortilin antibodies of other region classes were tested in the HTRF sortilin-PGRN binding assay, only two of 41 antibodies exhibited an inhibitory effect. One of these two antibodies cross-blocked with the D and D+, but had an atypical shuffle construct binding pattern (D-like except that it bound the hB01-05 region), while the other antibody did not cross-block with the other antibodies that inhibited PGRN-sortilin binding, thus supporting the conclusion that it binds to another sortilin region.

[0015] These observations show that antibodies binding to an area in sortilin defined by the D-region have the potential to inhibit sortilin-PGRN binding.

[0016] 19 cross-blocking antibodies, of which 18 were D-region and D+ antibodies, increased extracellular PGRN in a cellular assay (**Example 13, Figure 10** and **Figure 11**). Three of these antibodies were tested in vivo and found to increase plasma PGRN (**Figure 13, Example 15**).

[0017] The boxes in **Figure 1** illustrate steps in the selection of antibodies. A-E refer to the regions to which the respective Sortilin-binding antibodies were assigned based on shuffle constructs as described in **Example 1** and **SEQ ID NOs:171-179**. "Other" refers to an antibody which could not be assigned to one
5 region, and which may bind at the interface between the A- and B-regions. Tet refers to antibodies binding also tetraodon-Sortilin.

[0018] In addition to the human antibodies shown, a set of mouse anti-human sortilin antibodies was generated and similarly characterized. Two of these antibodies were assigned to the D Region and shown to cross block with
10 human D-region and D+ antibodies, to inhibit sortilin-PGRN binding and to increase extracellular PGRN (see **Figure 4**).

[0019] **Figure 2** shows the region assignment of antibodies based on binding to Sortilin shuffle constructs.

[0020] **Panel A** shows a linear illustration of the shuffle constructs used for
15 region assignment of antibodies as described in **Example 1**. Sortilin shuffle constructs were generated based on the human Sortilin sequence (**SEQ ID NO:169**) (sections depicted in grey) in which amino acid residues were exchanged to the corresponding amino acid from the tetraodon Sortilin sequence (depicted in black) (**SEQ ID NO:173**) (**Examples 1-3**).

20 [0021] **Panel B** shows predicted structure of the shuffle constructs illustrated linearly in A. Dark residues indicate residues changed to the corresponding tetraodon sequence in the shuffle constructs.

[0022] **Panel C** illustrates the binding pattern of antibodies assigned to the D-region and the E region classes respectively. A "+" indicates binding to a
25 given shuffle construct and a "-" indicates lack of binding. Based on the binding pattern to the different shuffle constructs, antibodies were assigned to regions. The resultant antibody region classes are indicated by A-E. For the illustrated D and E region antibodies, both bound the human sequences (all grey) as indicated by "+" and neither bound the tetraodon sequence (all black) as

indicated by “-”, whereas the E region antibody bound the hB45678 shuffle construct while the D Region antibody did not bind resulting in the localisation of binding as illustrated in Panel A. For D Region antibodies, binding to the following shuffle regions was observed: hsort, hB06-10, B12390. The antibodies
5 did not bind to hB01-05, B45678, tet. For D+ antibodies, binding to the following shuffle region was observed: hsort, B12390. The antibodies did not bind to hB01-05, hB06-10, B45678, tet. The F binding pattern was similar to the D binding pattern except that no binding to the hB06-10 was observed for D+ antibodies.

10 **[0023]** The antibodies did not bind to the fully tetraodon Sortilin protein, except two. The two antibodies capable of binding the tetraodon sequence were denoted “tet”. “Other” refers to an antibody which could not be assigned to one region.

[0024] **Figure 3** shows the binding affinities of human D-region and D+ antibodies. Binding affinities to sortilin shuffle constructs by bioLayer interferometry using Octet 384RED as described in **Example 8** (EC50, ng/ml). No shading indicates EC50 of 0.1-10 ng/ml, light grey shading indicates EC50 >10 ng/ml and grey shading indicates no binding (NB). Region assignment was based on binding patterns is illustrated in **Figure 2**. Shuffle constructs are
15 illustrated in **Figure 2** and sequences are given in **SEQ ID NOs:171-179**. mAb = monoclonal antibody.
20

[0025] **Figure 4** shows the binding affinities of mouse anti-human antibodies to Sortilin shuffle constructs as obtained by bioLayer interferometry using Octet 384RED as described in **Example 8** (EC50, ng/ml). No shading indicates
25 binding and grey shading indicates no binding (NB). Region assignment based on binding patterns is illustrated in **Figure 2**.

[0026] **Figure 5** shows the effect of Sortilin antibody on Sortilin PGRN binding. The D Region Sortilin human monoclonal (humAb) antibody 45 (filled circles) prevented PGRN binding to Sortilin, in contrast to a control Sortilin E region antibody (filled triangles) and an IgG control, IgG1-b12 (open triangles)
30

that did not interfere with the binding. The binding of antibodies was determined by measuring the displacement of PGRN binding to Sortilin using Homogenous Time Resolved Fluorescent (HTRF) (**Example 10**). Dose-response evaluation of antibodies was performed with ten concentrations covering 50pM to 1µM in a 3-fold dilution curve. The half-maximal inhibitory concentration (IC₅₀) values were calculated by non-linear regression using sigmoidal concentration response (variable slope) in XLfit 4 (IDBS, UK).

[0027] **Figure 6** Summary of effect of antibodies on Sortilin-PGRN binding determined by homogenous time resolved fluorescent (HTRF) analysis as shown in **Figure 5**. In total, 62 antibodies were tested - 15 D-region antibodies and 3 D+ antibodies were found to inhibit sortilin-PGRN binding and the IC₅₀ values were determined. For two additional antibodies (E and other regions), an inhibitory effect was observed. All remaining antibodies were negative in the test. * antibody too weak to fit a dose-response curve. 6% inhibition at 1 µM. ** control (ctrl) antibody too weak to fir a dose-response curve. 37% inhibition at 1 µM.

[0028] These observations show that sortilin antibodies characterized by their D-region or D+ assignment inhibit sortilin's binding to PGRN directly and are capable of inhibiting sortilin-PGRN binding.

[0029] **Figure 7** shows cross-blocking between antibodies. Human antibodies and the mouse antibodies were all tested in a single experiment where each antibody was bound to human wild type (WT) Sortilin (**Figure 7**). Subsequently all other antibodies were tested for binding to the preformed sortilin:antibody complex (**Example 9**). The selected 15 D-region and 3 D+ human antibodies (based on their effect in the HTRF PGRN-sortilin assay, (**Figure 5** and **Figure 6**) and two mouse D Region antibodies all inhibited binding of each other to human WT Sortilin.

[0030] The antibodies did not cross-block with antibodies designated to other region classes (as illustrated for A-region, E-region and tetraodon recognizing antibodies numbered AbA1-x, AbE1-x and Abtet in the table

respectively) except for one cross blocking A region antibody, one antibody with unknown region assignment ("other") and a partial block for a D+ antibody 548. These data support that the D-region and D+ antibodies capable of inhibiting sortilin-PGRN binding in the HTRF assay all interact with the same region in
5 sortilin.

[0031] Cross blocking between Sortilin antibodies from the same or different regions (regions based on binding to shuffle constructs as illustrated in **Figure 2**) was determined by analyzing interference with antibody-Sortilin binding. Binding of antibodies to Sortilin-ECD-His was measured by BioLayer
10 Interferometry using Octet 384RED (**Example 9**). The left column indicates primary (immobilized) antibodies and the top row indicates the secondary antibodies (antibodies being tested against the immobilized antibodies). Binding of both the primary and secondary antibodies to Sortilin-ECD-His would results in a response value higher than 0.1 and indicate that both antibodies were
15 binding to different regions of the protein. Response value less than 0.1 shows lack of binding of the secondary antibody and an effective cross blocking by the immobilized (primary) antibody, which suggests that both antibodies bind to the same region of Sortilin.

[0032] **Figure 8** shows the effect of D-region and D+ Sortilin antibodies on
20 the binding of the selective small molecule ligand AF38469 to Sortilin. The binding site for AF38469 has been shown to be similar to the binding site of neurotensin and characterized by X-ray crystallography (Schröder *et al.* Bioorg Med Chem Lett. 2014 Jan 1;24(1):177-80). PGRN has been reported to bind to the same site (Lee *et al.* Hum Mol Genet. 2013) antibodies 45 and 68, binding
25 to D-region, and D+ respectively, did not inhibit the binding of AF38469 to sortilin. This data suggests that these antibodies have a binding site for Sortilin distinct from the binding site for AF38469. Therefore, antibodies 45 and 68 inhibit PGRN-sortilin binding through a binding site distinct from the hitherto presumed PGRN binding site in sortilin.

[0033] **Figure 9** Effect of antibodies 45 and 68 on cellular binding and endocytosis of PGRN (**Example 12**). Antibodies 45 and 68 inhibited the binding and/or endocytosis of PGRN by sortilin overexpressing cells. Addition of neurotensin (NT, 10uM) similarly reduced binding or endocytosis of PGRN as reflected in reduced fluorescence as expected whereas the isotype control antibody B12 did not influence PGRN fluorescence levels.

[0034] Antibodies (100nM) to be tested were added to S18 cells 30 min before addition of recombinant PGRN for 4 hr. The cells were then fixed, stained for PGRN and analyzed by Cellomics. PGRN fluorescence was measured as mean fluorescence per cell. Data is presented as mean $\pm$ SD. Data analyzed by one-way Anova followed by Dunnett's analysis, all groups were compared to PGRN. *p<0.05; **p<0.01

[0035] **Figure 10** Extracellular PGRN levels estimated by ELISA in media from cultures of sortilin over-expressing HEK cells (S18). Sortilin D-region (45, 811) and D+ (68) antibodies increased PGRN levels and a similar effect of the sortilin ligand neurotensin was observed whereas the control antibody B12 had no effect. These observations indicate that D-region and D+ Sortilin antibodies were capable of inhibiting sortilin-mediated clearance of PGRN thereby increasing extracellular PGRN. All antibodies were tested at 100nM. Neurotensin was tested at 10uM. PGRN levels have been normalized to control. Data is presented as mean $\pm$ SD. Data was analyzed by one-way Anova followed by Dunnett's analysis, all groups were compared to CTRL *p<0.05; **p<0.01. (**Example 13**).

[0036] **Figure 11** shows the effect of antibodies on extracellular PGRN in human Sortilin over-expressing HEK cells measured by ELISA as described in **example 13**. All selected D-region antibodies and the three selected D+ antibodies increased extracellular PGRN. PGRN levels were analysed the same as above. PGRN levels are normalized to untreated controls and given in %. Two antibodies were raised in mouse against human Sortilin (1F2F4 & 5E1F6) and the rest are human antibodies. Ab= monoclonal antibody.

[0037] **Figure 12 (Upper and Lower Panels)** shows the effect of Sortilin antibody on extracellular PGRN in neuronally differentiated iPSC cells (**Example 14**). The sortilin D-region antibody 45 and the D+ antibody 68 increased PGRN levels whereas the control antibodies B12 and anti-HEL had
5 no effect.

[0038] Neuronally differentiated iPSC cells were plated into 96 wells plate. One week later, antibodies were added to the cells. Media from the cells were collected at 48 hrs or 96 hrs and analysed by human PGRN ELISA (Enzo Life sciences) and samples analysed as per the manufacturer's instructions. Sortilin
10 human antibodies 45 and 68 increased PGRN levels in the media at both time points. Control isotype antibodies B12 and Anti-Hel (negative control) did not change extracellular PGRN. Data is presented as mean $\pm$ SD. Data was analyzed by one-way Anova followed by Dunnett's analysis * $p < 0.05$; ** $p < 0.01$ (**Example 14**)

15 [0039] **Figure 13 (Panels A-C)** shows plasma PGRN levels in human Sortilin expressing knock in (KI) mice treated with Sortilin human antibody (**Example 15**). Sortilin antibody 45 increased plasma PGRN levels whereas the control antibody had no effect.

[0040] **A Time course study:** Increased plasma levels of PGRN were
20 observed after injection of antibody 45 (D Region). Mice were injected with 45 (n=5) or control (n=3) antibody sc at a dose of 10mg/kg. Each group was sacrificed at different time points. In mice treated with control antibodies (Anti-Hel) there was no change in plasma PGRN, whereas in mice treated with 45 there was a gradual increase in PGRN levels. Effect appeared to peak between
25 24 and 48hrs and gradually decreased by day 4-7.

[0041] **B Subchronic study:** Mice were treated twice a week with 10mg/kg of 45 and control antibody (Anti-Hel). Samples were collected from cheek blood every week. Plasma PGRN was elevated at week 1 and remained at approximately the same level throughout the entire study as compared to the
30 animals treated with control antibody (n=20).

[0042] **C Dose response study:** Different doses (4 doses: 0.1, 0.4, 2 and 10 mg/kg) of the Sortilin (45) and control antibodies (Anti-Hel) were injected and mice were sacrificed on day 2. Plasma PGRN was elevated in mice treated with 45 (10 and 2mg/kg). lower doses (0.4 and 0.1 mg/kg) did not have an effect on plasma PGRN. Data is presented as mean $\pm$ SD. Data was analyzed by two-way Anova followed by Bonferroni's analysis * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (Example 15).

[0043] **Figure 14** provides an illustration of sortilin regions, based on sortilin shuffle constructs, imposed onto the predicted sortilin structure. The regions constitute the sections of the sortilin protein in which the change of selected amino acid residues from the human sequence to the tetraodon sequence inhibited binding for antibodies of that region class. The arrow indicates the reported high affinity binding site of neurotensin and PGRN (Quistgaard Nat Struct Mol Biol. 2009 Jan;16(1):96-8, Lee et al, Hum Mol Genet. 2013).

[0044] **Figure 15a (Panels 1-6) and 15b (Panels 1-3)** Shows representative peptides covering the conformational epitope of antibody 45, 68 and 811. All of the shown peptides show a protection from exchange larger than 0.5D, except peptide 115-125. Peptide 115-125 is an example of a peptide which is unaffected by the presence of antibody 45, 68 or 811, and thereby is not a part of the conformational binding epitope (Example 16)

[0045] **Figure 16a (Panels 1-6) and 16b (Panels 1-3)** shows representative peptides covering the conformational epitope of antibody 30. All of the shown peptides show a protection from exchange larger than 0.5D, except peptide 563-572, peptide 646-656 and peptide 704-714. These three peptides are examples of peptides which are unaffected by the presence of antibody 30, and thereby is not a part of the conformational binding (Example 16)

[0046] **Figure 17** shows an illustration of the microdialysis for procedure.

[0047] **Figure 18a** shows a time course: effect of systemic administration of antibody 45 or PBS (50 mg/kg, 10 ml/kg, s.c.) 24h prior to the microdialysis

experiments on the levels of PRGN in the hippocampus of freely moving hSORT1 mice over time (24h). (**Example 17**)

[0048] **Figure 18b** shows the results of a pooled 24h dialysis: basal PRGN in the hippocampus of freely moving hSORT1 mice in mab#45- and PBS- treated mice, 3.3 ± 0.3 ng/ml and 1.1 ± 0.1 ng/ml, respectively, as assessed by push-pull microdialysis (**Example 17**).

[0049] **Figure 18c** presents a table showing PRGN levels (means $\pm$ SEM) in the hippocampus of freely moving hSORT1 mice measured every 2h during 24h, 1d after the animals were treated with mab#45 (n= 10) or PBS (n=8) (**Example 17**).

Detailed Description Of The Invention

[0050] As used herein, the term "Sortilin" is synonymous with the Sortilin protein (identified in for example UniProt as Q99523, 1 and 2). The amino acid numbering of Sortilin is given with respect to **SEQ ID NO:169** as shown below, Met being amino acid 1:

```

MERPWGAADG LSRWPHGLGL LLLLQLLPPS TLSQDRLDAP PPPAAPLPRW
SGPIGVSWGL RAAAAGGAFP RGGRWRRSAP GEDEECGRVR DFVAKLANNT
HQHVFDDLRLG SVSLSWVGDS TGVILVLTTF HVPLVIMTFG QSKLYRSEDY
GKNFKDITDL INNTFIRTEF GMAIGPENSG KVVLTAEVSG GSRGGGRIFRS
SDFAKNFVQT DLPFHPLTQM MYSPQNSDYL LALSTENGLW VSKNFGGKWE
EIHKAVCLAK WGSNTIIFT TYANGSCKAD LGALELWRTS DLGKSFKTIG
VKIYSFGLGG RFLFASVMAD KDTTRRIHVS TDQGDWTSMA QLPSVGQEQF
YSILAANDDM VFMHVDEPGD TGFGTIFTSD DRGIVYSKSL DRHLYTTTGG
ETDFTNVTSL RGVYITSVLS EDNSIQTMIT FDQGGRWTHL RKPENSECDA
TAKNKNECSL HIHASYSISQ KLNVPMAPLS EPNAVGIIVIA HGSVGDAISV
MVPDVYISDD GGYSWTKMLE GPHYITILDS GGIIVAIEHS SRPINVIKFS
TDEGQCWQTY TFTRDPIYFT GLASEPGARS MNISIWGFTE SFLTQWVSY
TIDFKDILER NCEEKDYTIW LAHSTDPEY EDGCILGYKE QFLRLRKSSV
CQNGRDYVVT KQPSICLCSL EDFLCDFGYR RPENDSKCVE QPELKGHDLE
FCLYGREEHL TTNGYRKIPG DKCQGGVNPV REVKDLKKKC TSNFLSPEKQ
NSKSNSVPII LAIVGLMLVT VVAGVLIVKK YVCGGRFLVH RYSVLQQHAE
ANGVDGVDAL DTASHTNKSG YHDDSDEDLLE

```

[0051] As used herein, the term “D Region” is intended to refer to the region on Sortilin (corresponding to residues 523-610 of **SEQ ID NO:169**) consisting of the amino acids in **SEQ ID NO:170** as shown below:

HYYTILDSGG IIVAIEHSSR PINVIKFSTD EGQCWQTYTF TRDPIYFTGL
5 ASEPGARSMN ISIWGFTESE LTSQWVSYTI DFKDILER

[0052] For D Region antibodies, binding to the following shuffle regions was observed: hsort, hB06-10, B12390. The antibodies did not bind to hB01-05, B45678, tet. For D+ antibodies, binding to the following shuffle regions was observed: hsort, B12390. The antibodies did not bind to hB01-05, hB06-10, B45678, tet. For the antibodies named “D+” a similar binding pattern was observed as for “D” region antibodies except that no binding to the hB06-10 was observed for D+ antibodies. Despite the different binding pattern to shuffle constructs, D+ antibodies shared functional characteristics (cell assays etc.) to D-region antibodies.

[0053] Certain D and D + binding antibodies binds particular the region as defined in **SEQ ID NO:185, 186 or 187**. Thus, the invention relates in certain embodiments to antibodies or antigen binding fragments thereof that binds to the D region sequences of **SEQ ID 170**, and within that region **SEQ ID NO:185, 186 or 187**. By binding, these antibodies or antigen-binding fragments, influence the PGRN levels and thus may be used to treat disease associated with PGRN, such as FTD, ALS and AD.

[0054] By further analysing the binding of the D and D+ antibodies of the invention, partial binding to the neighbouring region (the A region) was identified for some of the antibodies. This region corresponds to amino acids 78-254 of **SEQ ID NO:169**, as shown in **SEQ ID NO:180** and below:

SAPGEDEECG RVRDFVAKLA NNTHQHVFDL LRGSVSLSWV GDSTGVILVL
TTFHVPLVIM TFGQSKLYRS EDYGKNFKDI TDLINNTFIR TEFGMAIGPE
NSGKVVLTAE VSGGSRGGRI FRSSDFAKNF VQTDLPFHPL TQMMYSPQNS
DYLLALSTEN GLWVSKNFGG KWEEIHK

This region has been given the term “A region”.

[0055] Thus, in certain embodiments the invention relates to antibodies or antigen binding fragments thereof that binds the D region as defined above and in SEQ ID Nos 170, 185, 186 or 187, and further have affinity for the A region
5 identified in SEQ ID NO 180. Within the A region certain antibodies or antigen binding fragments thereof have affinity for the A region amino acids identified in SEQ ID NOs 181, 182, 183 or 184.

[0056] PGRN (proepithelin, granulin-epithelin precursor, PC-cell-derived growth factor, acrogranin) encodes a 68.5 kDa secreted glycoprotein that has
10 7.5 repeats of smaller granulin motifs, ranging from 6-25 kDa, which can be proteolytically cleaved from the precursor PGRN (He, Z. & Bateman, A., J. Mol. Med. 81:600-6X2 (2003)). In non- neuronal cells, PGRN has been associated with a variety of events, such as cell cycle regulation and cell motility (He, Z. & Bateman, A., J. Mol. Med. 57:600-612 (2003); Monami, G., et al., Cancer Res.
15 (5(5:7103-7110 (2006)), wound repair, inflammation (Zhu, J., et al., Cell 77:867-878 (2002)), induction of growth factors such as vascular endothelial growth factor (VEGF) (Tangkeangsinsin, W. & Serrero, G, Carcinogenesis 25:1587-1592 (2004)), and tumorigenesis (He, Z. & Bateman, A., J. Mol. Med. 81:600- 612 (2003), Monami, G., et al., Cancer Res (5(5:7103-7110 (2006);
20 Serrero, G., Biochem Biophys. Res. Commun. 505:409-413 (2003), Lu, R & Serrero, G., Proc. Natl Acad Sci U.S.A 98 142-147 (2001); Liau, L M., et al., Cancer Res. 60:1353-1360 (2000)).

[0057] PGRN mutations result in haploinsufficiency (Baker, M., et al., Nature 442:916-919 (2006); Cruts, M., et al., Nature 442:920-924 (2006)) and
25 are known to be present in nearly 50% of familial FTD cases, making PGRN mutation a major genetic contributor to FTD (Cruts, M. & Van Broeckhoven, C, Trends Genet. 24:186-194 (2008); Le Ber, I., et al., Brain 129:3051- 3065 (2006)). The loss-of-function heterozygous character of PGRN mutations implies that in healthy individuals, PGRN expression plays a dose-dependent,
30 critical role in protecting healthy individuals from the development of FTD.

[0058] The term "antibody" (Ab) in the context of the present invention refers to an immunoglobulin molecule or according to some embodiments of the invention, a fragment of an immunoglobulin molecule which has the ability to bind to an epitope of a molecule ("antigen"). Naturally occurring antibodies typically comprise a tetramer which is usually composed of at least two heavy (H) chains and at least two light (L) chains. Each heavy chain is comprised of a heavy chain variable domain (abbreviated herein as VH) and a heavy chain constant domain, usually comprised of three domains (CH1, CH2 and CH3). Heavy chains can be of any isotype, including IgG (IgG1, IgG2, IgG3 and IgG4 subtypes), IgA (IgA1 and IgA2 subtypes), IgM and IgE. Each light chain is comprised of a light chain variable domain (abbreviated herein as VL) and a light chain constant domain (CL). Light chains include kappa chains and lambda chains. The heavy and light chain variable domain is typically responsible for antigen recognition, while the heavy and light chain constant domain may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (*e.g.*, effector cells) and the first component (C1q) of the classical complement system. The VH and VL domains can be further subdivided into regions of hypervariability, termed "complementarity determining regions," that are interspersed with domains of more conserved sequence, termed "framework regions" (FR). Each VH and VL is composed of three CDR Domains and four FR Domains arranged from amino-terminus to carboxy-terminus in the following order: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4. The variable domains of the heavy and light chains contain a binding domain that interacts with an antigen. Of particular relevance are antibodies and their antigen-binding fragments that have been "isolated" so as to exist in a physical milieu distinct from that in which it may occur in nature or that have been modified so as to differ from a naturally occurring antibody in amino acid sequence.

[0059] The term "epitope" means an antigenic determinant capable of specific binding to an antibody. Epitopes usually consist of surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge

characteristics. Conformational and linear epitopes are distinguished in that the binding to the former, but not the latter, is always lost in the presence of denaturing solvents. The epitope may comprise amino acid residues directly involved in the binding and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the specifically antigen-binding peptide (in other words, the amino acid residue is within the footprint of the specifically antigen-binding peptide).

[0060] As used herein, the term “antigen-binding fragment of an antibody” means a fragment, portion, region or domain of an antibody (regardless of how it is produced (*e.g.*, via cleavage, recombinantly, synthetically, *etc.*)) that is capable of binding to an epitope, and thus the term “antigen-binding” is intended to mean the same as “epitope-binding” so that, for example, an “antigen-binding fragment of an antibody” is intended to be the same as an “epitope-binding fragment of an antibody”. An antigen-binding fragment may contain 1, 2, 3, 4, 5 or all 6 of the CDR Domains of such antibody and, although capable of binding to such epitope, may exhibit a specificity, affinity or selectivity toward such epitope that differs from that of such antibody. Preferably, however, an antigen-binding fragment will contain all 6 of the CDR Domains of such antibody. An antigen-binding fragment of an antibody may be part of, or comprise, a single polypeptide chain (*e.g.*, an scFv), or may be part of, or comprise, two or more polypeptide chains, each having an amino-terminus and a carboxyl terminus (*e.g.*, a diabody, a Fab fragment, a Fab₂ fragment, *etc.*). Fragments of antibodies that exhibit antigen-binding ability can be obtained, for example, by protease cleavage of intact antibodies. More preferably, although the two domains of the Fv fragment, VL and VH, are naturally encoded by separate genes, or polynucleotides that encode such gene sequences (*e.g.*, their encoding cDNA) can be joined, using recombinant methods, by a flexible linker that enables them to be made as a single protein chain in which the VL and VH regions associate to form monovalent antigen-binding molecules (known as single-chain Fv (scFv); see *e.g.*, Bird *et al.*, (1988) Science 242:423-426; and Huston *et al.* (1988) Proc. Natl. Acad. Sci. (U.S.A.) 85:5879-5883). Alternatively, by employing a flexible linker that is too short (*e.g.*, less than about 9 residues)

to enable the VL and VH domains of a single polypeptide chain to associate together, one can form a bispecific antibody, diabody, or similar molecule (in which two such polypeptide chains associate together to form a bivalent antigen-binding molecule) (see for instance PNAS USA 90(14), 6444-8 (1993) for a description of diabodies). Examples of antigen-binding fragments encompassed within the present invention include (i) a Fab' or Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains, or a monovalent antibody as described in WO2007059782; (ii) F(ab')₂ fragments, bivalent fragments comprising two Fab fragments linked by a disulfide bridge at the hinge domain; (iii) an Fd fragment consisting essentially of the VH and CH1 domains; (iv) a Fv fragment consisting essentially of a VL and VH domains, (v) a dAb fragment (Ward et al., Nature 341, 544-546 (1989)), which consists essentially of a VH domain and also called domain antibodies (Holt et al; Trends Biotechnol. 2003 Nov;21(11) :484-90); (vi) camelid or nanobodies (Revets et al; Expert Opin Biol Ther. 2005 Jan;5_(I): 11-24) and (vii) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they may be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH domains pair to form monovalent molecules (known as single chain antibodies or single chain Fv (scFv), see for instance Bird et al., Science 242, 423-426 (1988) and Huston et al., PNAS USA 85, 5879-5883 (1988)). These and other useful antibody fragments in the context of the present invention are discussed further herein. It also should be understood that the term antibody, unless specified otherwise, also includes antibody-like polypeptides, such as chimeric antibodies and humanized antibodies, and antibody fragments retaining the ability to bind to the antigen (antigen-binding fragments) provided by any known technique, such as enzymatic cleavage, peptide synthesis, and recombinant techniques. An antibody as generated can possess any isotype. As used herein, "isotype" refers to the immunoglobulin class (for instance IgG1, IgG2, IgG3 or IgG4) that is encoded by heavy chain constant domain genes. Such antibody fragments are obtained using conventional techniques known to those of skill in the art;

suitable fragments capable of binding to a desired epitope may be readily screened for utility in the same manner as an intact antibody.

[0061] The term "bispecific antibody" refers to an antibody containing two independent antigen-binding fragments that each target independent targets. These targets can be epitopes present on different proteins or different epitopes present on the same target. Bispecific antibody molecules can be made using compensatory amino acid changes in the constant domains of the HCs of the parent monospecific bivalent antibody molecules. The resulting heterodimeric antibody contains one Fabs contributed from two different parent monospecific antibodies. Amino acid changes in the Fc domain leads to increased stability of the heterodimeric antibody with bispecificity that is stable over time. (Ridgway et al., Protein Engineering 9, 617-621 (1996), Gunasekaran et al., JBC 285, 19637-1(2010), Moore et al., MABs 3:6 546-557 (2011), Strop et al., JMB 420, 204-219 (2012), Metz et al., Protein Engineering 25:10 571-580 (2012), Labrijn et al., PNAS 110:113, 5145 -5150 (2013), Spreter Von Kreudenstein et al., MABs 5:5 646-654 (2013)). Bispecific antibodies can also include molecules that are generated using ScFv fusions. Two monospecific scfv are then independently joined to Fc domains able to form stable heterodimers to generate a single bispecific molecule (Mabry et al., PEDS 23:3 115-127 (2010)). Bispecific molecules have dual binding capabilities.

[0062] An "anti-Sortilin antibody" or "Sortilin antibody" (used interchangeably herein, depending on the context wherein its written) is an antibody an antigen-binding fragment thereof which binds specifically to Sortilin, and especially to the Sortilin D Region, **SEQ ID NO:170**. An anti-Sortilin antibody that binds to the Sortilin D Region will usually bind to a conformational epitope or a linear epitope of 3, 4, 5, 6 or 7 consecutive amino acids within the D-Region (for example **SEQ ID NOs:185, 186 or 187**) with an affinity (IC50) at or below 22 nM, such as between 22 nM and 1 nM, between 10 nM and 1 nM or between 5 nM and 1 nM. According to some embodiments the anti-Sortilin antibodies may also bind to the A region (**SEQ ID NOs:180, 181, 182, 183 or 184**) although it's

emphasized that their main biological function is believed to be achieved by binding to the D Region.

[0063] The binding site identified is rather unique as shown with for example the binding of the selective small molecule ligand AF38469 to Sortilin. The binding site for AF38469 has been shown to be similar to the binding site of neurotensin and characterized by X-ray crystallography (Schröder *et al.* Bioorg Med Chem Lett. 2014 Jan 1;24(1):177-80). PGRN has been reported to bind to the same site (Lee *et al.* Hum Mol Genet. 2013). Antibodies 45 and 68, binding to D-region, and D+ respectively, did not inhibit the binding of AF38469 to sortilin. This data suggests that these antibodies have a binding site for Sortilin distinct from the binding site for AF38469 and neurotensin. Therefore, in certain embodiments the invention relates to an antibody, or an antigen-binding fragment thereof, capable of specifically binding to Sortilin and inhibiting the binding of PGRN to Sortilin, but which binding does not inhibit or substantially inhibit the binding of neurotensin or AF38469 to Sortilin. This can be shown using for example displacement of binding to Sortilin using a scintillation proximity assay (SPA) (Example 11). One way of explaining this finding could be that the antibodies, or antigen-binding fragments thereof, are binding to surface areas of Sortilin whereas the small molecules like neurotensine are binding inside the binding pocket.

[0064] The term "human antibody" (which may be abbreviated to "humAb" or "HuMab"), as used herein, is intended to include antibodies having variable and constant domains derived from human germline immunoglobulin sequences. The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or during gene rearrangement or by somatic mutation in vivo).

[0065] The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of single molecular composition. A conventional monoclonal antibody composition

displays a single binding specificity and affinity for a particular epitope. In certain embodiments a monoclonal antibody can be composed of more than one Fab domain thereby increasing the specificity to more than one target. The terms "monoclonal antibody" or "monoclonal antibody composition" are not intended to be limited by any particular method of production (e.g., recombinant, transgenic, hybridoma, etc.).

[0066] The antibodies of the present invention, and their sortilin antigen-binding fragments will preferably be human or, for example for the mouse antibodies (denoted 1F2, 5E1), "humanized," particularly if employed for therapeutic purposes. The term "humanized" refer to a molecule, generally prepared using recombinant techniques, having an antigen-binding site derived from an immunoglobulin from a non-human species and a remaining immunoglobulin structure based upon the structure and /or sequence of a human immunoglobulin. The antigen-binding site may comprise either complete non-human antibody variable domains fused to human constant domains, or only the complementarity determining regions (CDRs) of such variable domains grafted to appropriate human framework regions of human variable domains. The framework residues of such humanized molecules may be wild type (e.g., fully human) or they may be modified to contain one or more amino acid substitutions not found in the human antibody whose sequence has served as the basis for humanization. Humanization lessens or eliminates the likelihood that a constant domain of the molecule will act as an immunogen in human individuals, but the possibility of an immune response to the foreign variable domain remains (LoBuglio, A.F. *et al.* (1989) "Mouse/Human Chimeric Monoclonal Antibody In Man: Kinetics And Immune Response," Proc. Natl. Acad. Sci. (U.S.A.) 86:4220-4224). Another approach focuses not only on providing human-derived constant domains, but modifying the variable domains as well so as to reshape them as closely as possible to human form. It is known that the variable domains of both heavy and light chains contain three complementarity- determining regions (CDRs) which vary in response to the antigens in question and determine binding capability, flanked by four framework regions (FRs) which are relatively conserved in a given species and

which putatively provide a scaffolding for the CDRs. When nonhuman antibodies are prepared with respect to a particular antigen, the variable domains can be "reshaped" or "humanized" by grafting CDRs derived from nonhuman antibody on the FRs present in the human antibody to be modified.

5 Application of this approach to various antibodies has been reported by Sato, K. *et al.* (1993) *Cancer Res* 53:851-856. Riechmann, L. *et al.* (1988) "*Reshaping Human Antibodies for Therapy*," *Nature* 332:323-327; Verhoeyen, M. *et al.* (1988) "*Reshaping Human Antibodies: Grafting An Antilysozyme Activity*," *Science* 239:1534-1536; Kettleborough, C. A. *et al.* (1991) "*Humanization Of A*

10 *Mouse Monoclonal Antibody By CDR-Grafting: The Importance Of Framework Residues On Loop Conformation*," *Protein Engineering* 4:773-3783; Maeda, H. *et al.* (1991) "*Construction Of Reshaped Human Antibodies With HIV-Neutralizing Activity*," *Human Antibodies Hybridoma* 2:124-134; Gorman, S. D. *et al.* (1991) "*Reshaping A Therapeutic CD4 Antibody*," *Proc. Natl. Acad. Sci. (U.S.A.)* 88:4181-4185; Tempest, P.R. *et al.* (1991) "*Reshaping A Human Monoclonal Antibody To Inhibit Human Respiratory Syncytial Virus Infection in vivo*," *Bio/Technology* 9:266-271; Co, M. S. *et al.* (1991) "*Humanized Antibodies For Antiviral Therapy*," *Proc. Natl. Acad. Sci. (U.S.A.)* 88:2869-2873; Carter, P. *et al.* (1992) "*Humanization Of An Anti-p185her2 Antibody For Human Cancer*

15 *Therapy*," *Proc. Natl. Acad. Sci. (U.S.A.)* 89:4285-4289; and Co, M.S. *et al.* (1992) "*Chimeric And Humanized Antibodies With Specificity For The CD33 Antigen*," *J. Immunol.* 148:1149-1154. In some embodiments, humanized antibodies preserve all CDR sequences (for example, a humanized mouse antibody which contains all six CDRs from the mouse antibodies). In other

20 embodiments, humanized antibodies have one or more CDRs (one, two, three, four, five, six) which are altered with respect to the original antibody, which are also termed one or more CDRs "derived from" one or more CDRs from the original antibody. The ability to humanize an antigen is well known (see, *e.g.*, US Patents No. 5,225,539; 5,530,101; 5,585,089; 5,859,205; 6,407,213;

25 6,881,557).

30

[0067] The term "antibody "XX" is intended to denote an antibody or antigen-binding fragment thereof (for example antibody "5E1"), comprising or

consisting of the Light Chain, the Light Chain Variable domain, or the Light Chain Variable domain CDR1-3, as defined by its respective SEQ ID NO, and the Heavy Chain, Heavy Chain Variable Domain, or Heavy Chain Variable Domain CDR1-3 as defined by its respective SEQ ID NO. In certain
5 embodiments the antibody or antigen-binding fragment thereof are defined by their entire Heavy Chain Variable Domain comprising as defined by their SEQ ID NO and their Light Chain Variable Domain as defined by their SEQ ID NO.

[0068] The numbering of amino acid residues in this region is according to
10 IMGT®, the international ImMunoGeneTics information system® or, Kabat, E. A., Wu, T. T., Perry, H. M., Gottesmann, K. S. & Foeller, C. (1991). Sequences of Proteins of Immunological Interest, 5th edit., NIH Publication no. 91-3242 U.S. Department of Health and Human Services; Chothia, C. & Lesk, A. M. (1987). Canonical structures For The Hypervariable domains Of Immunoglobu-
15 lins. J. Mol. Biol. 196, 901-917.

[0069] As used herein, an antibody or an antigen-binding fragment thereof is said to “specifically” bind a region of another molecule (*i.e.*, an epitope) if it reacts or associates more frequently, more rapidly, with greater duration and/or with greater affinity or avidity with that epitope relative to alternative epitopes.
20 In one embodiment, the antibody, or antigen-binding fragment thereof, of the invention binds at least 10-fold more strongly to its target (Sortilin) than to another molecule; preferably at least 50-fold more strongly and more preferably at least 100-fold more strongly. Preferably, the antibody, or antigen-binding fragment thereof, binds under physiological conditions, for example, *in vivo*.
25 Thus, by “specifically binding to Sortilin”, we include the ability of the antibody, or antigen-binding fragment thereof, to bind to Sortilin with such specificity and/or under such conditions. Methods suitable for determining such binding will be known to those skilled in the art, and exemplary methods are described in the accompanying Examples. As used herein, the term “binding” in the
30 context of the binding of an antibody to a predetermined antigen typically refers to binding with an affinity corresponding to a KD of about 10^{-7} M or less, such as about 10^{-8} M or less, such as about 10^{-9} M or less when determined by for

instance surface plasmon resonance (SPR) technology in either a BIAcore® 3000 or T200 instrument using the antigen as the ligand and the antibody as the analyte, and binds to the predetermined antigen with an affinity corresponding to a KD that is at least ten-fold lower, such as at least 100 fold lower, for instance at least 1,000 fold lower, such as at least 10,000 fold lower, for instance at least 100,000 fold lower than its affinity for binding to a non-specific antigen (e.g., BSA, casein) other than the predetermined antigen or a closely-related antigen. The amount with which the affinity is lower is dependent on the KD of the antibody, so that when the KD of the antibody is very low (that is, the antibody is highly specific), then the amount with which the affinity for the antigen is lower than the affinity for a non-specific antigen may be at least 10,000 fold. In particular, the invention pertains to anti-Sortilin antibodies that exhibit a binding affinity corresponding to at or below 22 nM, such as between 22 nM and 1 nM, between 10 nM and 1 nM or between 5 nM and 1 nM, when determined by, for instance, bioLayer interferometry using an Octet 384RED (Example 8).

[0070] In certain embodiments of the invention the invention relates to an antibody or antigen-binding fragment thereof able to compete with humAb antibody 45 or humAb antibody 68 for binding to Sortilin. In another embodiment the invention relates to an antibody or antigen-binding fragment thereof that is able to compete with antibody 45 for binding to the D Region of Sortilin as defined in **SEQ ID NO:170**. Such competitive binding inhibition can be determined using assays and methods well known in the art, for example using BIAcore® chips with immobilised human Sortilin and incubating with a reference antibody (such as antibody “45” or “68”) with and without an antibody polypeptide to be tested. Alternatively, a pair-wise mapping approach can be used, in which a reference antibody (such as antibody “45” or “68”) is immobilised to the surface of the BIAcore® chip, human Sortilin antigen is bound to the immobilised antibody, and then a second antibody is tested for simultaneous binding ability to human Sortilin (see ‘BIAcore® Assay Handbook’, GE Healthcare Life Sciences, 29-0194-00 AA 05/2012; the disclosures of which are incorporated herein by reference).

[0071] The term "kd" (sec⁻¹ or 1/s), as used herein, refers to the dissociation rate constant of a particular antibody-antigen interaction. Said value is also referred to as the koff value.

[0072] The term "ka" (M⁻¹ x sec⁻¹ or 1/Msec), as used herein, refers to the association rate constant of a particular antibody-antigen interaction.

[0073] The term "KD" (M), as used herein, refers to the dissociation equilibrium constant of a particular antibody-antigen interaction and is obtained by dividing the kd by the ka.

[0074] The term "KA" (M⁻¹ or 1/M), as used herein, refers to the association equilibrium constant of a particular antibody-antigen interaction and is obtained by dividing the ka by the kd.

[0075] In one embodiment, the invention relates to an antibody, or antigen-binding fragment thereof, which exhibits one or more of the following properties:

- (i) a binding affinity (K_D) for Sortilin of between 0.5-10 nM, such as 1-5 nM or 1-2 nM;
- (ii) capability to reduce and/or inhibit PGRN binding to Sortilin;
- (iii) capability to reduce and/or inhibit clearance of PGRN by Sortilin-expressing cells;
- (iv) capability to reduce and/or inhibit the endocytosis of PGRN by Sortilin-expressing cells;
- (v) capability to increase the amount and/or concentration of PGRN in the plasma in human-Sortilin-expressing knock-in mice.

[0076] The term "capability to reduce and/or inhibit PGRN binding to Sortilin" includes an antibody that has the ability to inhibit binding to PGRN at an IC₅₀ less than 50nM but preferably between 10nM and 0.2 nM using a time resolved fluorescence assay (HTFR) disclosed in **Example 10**.

[0077] The term "capability to reduce and/or inhibit clearance of PGRN by Sortilin-expressing cells" includes the ability to increase the concentration of PGRN in the medium by at least 25%, such as between 25% and 500%,

between 25% and 400% or between 25% and 200% as measured by an ELISA assay as disclosed in **Example 13**.

[0078] The “capability to reduce and/or inhibit the endocytosis of PGRN by Sortilin-expressing cells” includes the ability reduce the intracellular
5 concentration of PGRN by at least 10% but preferably between 20 and 100 % as measured by a cellomics based assay as disclosed in **Example 12**.

[0079] The “capability to increase the amount and/or concentration of PGRN in the plasma in human-Sortilin-expressing knock-in mice” includes the ability to increase the concentration of PGRN in the plasma by at least 25% but
10 preferably between 50 and 500 percent as measured by an ELISA assay as disclosed in **Example 15**.

[0080] It's envisaged that the capability to increase PGRN in the brain may also be assayed by for example microdialysis. Thus by “capability to increase the amount and/or concentration of PGRN in the brain” includes the ability to
15 increase the concentration of PGRN in the brain by at least 25% but preferably between 50 and 500 percent as measured by microdialysis.

[0081] In some antibodies, only part of a CDR, namely the subset of CDR residues required for binding, termed the SDRs, are needed to retain binding in a humanized antibody. CDR residues not contacting the relevant epitope and
20 not in the SDRs can be identified based on previous studies (for example residues H60-H65 in CDR H2 are often not required), from regions of Kabat CDRs lying outside Chothia hypervariable loops (see, Kabat *et al.* (1992) SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST, National Institutes of Health Publication No. 91-3242; Chothia, C. *et al.* (1987) “*Canonical Structures For The Hypervariable Regions Of Immunoglobulins*,” J. Mol. Biol. 196:901-
25 917), by molecular modeling and/or empirically, or as described in Gonzales, N.R. *et al.* (2004) “*SDR Grafting Of A Murine Antibody Using Multiple Human Germline Templates To Minimize Its Immunogenicity*,” Mol. Immunol. 41:863-872. In such humanized antibodies at positions in which one or more donor
30 CDR residues is absent or in which an entire donor CDR is omitted, the amino

acid occupying the position can be an amino acid occupying the corresponding position (by Kabat numbering) in the acceptor antibody sequence. The number of such substitutions of acceptor for donor amino acids in the CDRs to include reflects a balance of competing considerations. Such substitutions are potentially advantageous in decreasing the number of mouse amino acids in a humanized antibody and consequently decreasing potential immunogenicity. However, substitutions can also cause changes of affinity, and significant reductions in affinity are preferably avoided. Positions for substitution within CDRs and amino acids to substitute can also be selected empirically.

10 **[0082]** The fact that a single amino acid alteration of a CDR residue can result in loss of functional binding (Rudikoff, S. *etc.* (1982) "*Single Amino Acid Substitution Altering Antigen-binding Specificity*," Proc. Natl. Acad. Sci. (USA) 79(6):1979-1983) provides a means for systematically identifying alternative functional CDR sequences. In one preferred method for obtaining such variant CDRs, a polynucleotide encoding the CDR is mutagenized (for example via random mutagenesis or by a site-directed method (*e.g.*, polymerase chain-mediated amplification with primers that encode the mutated locus)) to produce a CDR having a substituted amino acid residue. By comparing the identity of the relevant residue in the original (functional) CDR sequence to the identity of the substituted (non-functional) variant CDR sequence, the BLOSUM62.ijj substitution score for that substitution can be identified. The BLOSUM system provides a matrix of amino acid substitutions created by analyzing a database of sequences for trusted alignments (Eddy, S.R. (2004) "*Where Did The BLOSUM62 Alignment Score Matrix Come From?*," Nature Biotech. 22(8):1035-1036; Henikoff, J.G. (1992) "Amino acid substitution matrices from protein blocks," Proc. Natl. Acad. Sci. (USA) 89:10915-10919; Karlin, S. *et al.* (1990) "*Methods For Assessing The Statistical Significance Of Molecular Sequence Features By Using General Scoring Schemes*," Proc. Natl. Acad. Sci. (USA) 87:2264-2268; Altschul, S.F. (1991) "*Amino Acid Substitution Matrices From An Information Theoretic Perspective*," J. Mol. Biol. 219, 555-565. Currently, the most advanced BLOSUM database is the BLOSUM62 database (BLOSUM62.ijj). **Table 1** presents the BLOSUM62.ijj substitution scores (the

- higher the score the more conservative the substitution and thus the more likely the substitution will not affect function). If an antigen-binding fragment comprising the resultant CDR fails to bind to Sortilin, for example, then the BLOSUM62.ijj substitution score is deemed to be insufficiently conservative,
- 5 and a new candidate substitution is selected and produced having a higher substitution score. Thus, for example, if the original residue was glutamate (E), and the non-functional substitute residue was histidine (H), then the BLOSUM62.ijj substitution score will be 0, and more conservative changes (such as to aspartate, asparagine, glutamine, or lysine) are preferred.

Table 1																				
	A	R	N	D	C	Q	E	G	H	I	L	K	M	F	P	S	T	W	Y	V
A	+4	-1	-2	-2	0	-1	-1	0	-2	-1	-1	-1	-1	-2	-1	+1	0	-3	-2	0
R	-1	+5	0	-2	-3	+1	0	-2	0	-3	-2	+2	-1	-3	-2	-1	-1	-3	-2	-3
N	-2	0	+6	+1	-3	0	0	0	+1	-3	-3	0	-2	-3	-2	+1	0	-4	-2	-3
D	-2	-2	+1	+6	-3	0	+2	-1	-1	-3	-4	-1	-3	-3	-1	0	-1	-4	-3	-3
C	0	-3	-3	-3	+9	-3	-4	-3	-3	-1	-1	-3	-1	-2	-3	-1	-1	-2	-2	-1
Q	-1	+1	0	0	-3	+5	+2	-2	0	-3	-2	+1	0	-3	-1	0	-1	-2	-1	-2
E	-1	0	0	+2	-4	+2	+5	-2	0	-3	-3	+1	-2	-3	-1	0	-1	-3	-2	-2
G	0	-2	0	-1	-3	-2	-2	+6	-2	-4	-4	-2	-3	-3	-2	0	-2	-2	-3	-3
H	-2	0	+1	-1	-3	0	0	-2	+8	-3	-3	-1	-2	-1	-2	-1	-2	-2	+2	-3
I	-1	-3	-3	-3	-1	-3	-3	-4	-3	+4	+2	-3	+1	0	-3	-2	-1	-3	-1	+3
L	-1	-2	-3	-4	-1	-2	-3	-4	-3	+2	+4	-2	+2	0	-3	-2	-1	-2	-1	+1
K	-1	+2	0	-1	-3	+1	+1	-2	-1	-3	-2	+5	-1	-3	-1	0	-1	-3	-2	-2
M	-1	-1	-2	-3	-1	0	-2	-3	-2	+1	+2	-1	+5	0	-2	-1	-1	-1	-1	+1
F	-2	-3	-3	-3	-2	-3	-3	-3	-1	0	0	-3	0	+6	-4	-2	-2	+1	+3	-1
P	-1	-2	-2	-1	-3	-1	-1	-2	-2	-3	-3	-1	-2	-4	+7	-1	-1	-4	-3	-2
S	+1	-1	+1	0	-1	0	0	0	-1	-2	-2	0	-1	-2	-1	+4	+1	-3	-2	-2
T	0	-1	0	-1	-1	-1	-1	-2	-2	-1	-1	-1	-1	-2	-1	+1	+5	-2	-2	0
W	-3	-3	-4	-4	-2	-2	-3	-2	-2	-3	-2	-3	-1	+1	-4	-3	-2	+11	+2	-3
Y	-2	-2	-2	-3	-2	-1	-2	-3	+2	-1	-1	-2	-1	+3	-3	-2	-2	+2	+7	-1
V	0	-3	-3	-3	-1	-2	-2	-3	-3	+3	+1	-2	+1	-1	-2	-2	0	-3	-1	+4

- 10 [0083] The invention thus contemplates the use of random mutagenesis to identify improved CDRs. In the context of the present invention, conservative substitutions may be defined by substitutions within the classes of amino acids reflected in one or more of the following three tables:

Amino Acid Residue Classes For Conservative Substitutions:

Table 2	
Acidic Residues	Asp (D) and Glu (E)
Basic Residues	Lys (K), Arg (R), and His (H)

Hydrophilic Uncharged Residues	Ser (S), Thr (T), Asn (N), and Gln (Q)
Aliphatic Uncharged Residues	Cly (G), Ala (A), Val (V), Leu (L), and Ile (I)
Non-polar Uncharged Residues	Cys (C), Met (M), and Pro (P)
Aromatic Residues	Phe (F), Tyr (Y), and Trp (W)

Alternative Conservative Amino Acid Residue Substitution Classes:

Table 3			
1	A	S	T
2	D	E	
3	N	Q	
4	R	K	
5	I	L	M
6	F	Y	W

Alternative Physical and Functional Classifications of Amino Acid Residues:

Table 4	
Alcohol Group-Containing Residues	S and T
Aliphatic Residues	I, L, V and M
Cycloalkenyl-Associated Residues	F, H, W and Y
Hydrophobic Residues	A, C, F, G, H, I, L, M, R, T, V, W and Y
Negatively Charged Residues	D and E
Polar Residues	C, D, E, H, K, N, Q, R, S and T
Positively Charged Residues	H, K and R
Small Residues	A, C, D, G, N, P, S, T and V
Very Small Residues	A, G and S
Residues Involved In Turn Formation	A, C, D, E, G, H, K, N, Q, R, S, P and T
Flexible Residues	Q, T, K, S, G, P, D, E and R

[0084] More conservative substitutions groupings include: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine.

[0085] Additional groups of amino acids may also be formulated using the principles described in, e.g., Creighton (1984) Proteins: Structure and Molecular Properties (2d Ed. 1993), W. H. Freeman and Company.

10 [0086] Phage display technology can alternatively be used to increase (or decrease) CDR affinity. This technology, referred to as affinity maturation,

employs mutagenesis or "CDR walking" and re-selection uses the target antigen or an antigenic antigen-binding fragment thereof to identify antibodies having CDRs that bind with higher (or lower) affinity to the antigen when compared with the initial or parental antibody (See, e.g. Glaser *et al.* (1992) J. Immunology 5 149:3903). Mutagenizing entire codons rather than single nucleotides results in a semi-randomized repertoire of amino acid mutations. Libraries can be constructed consisting of a pool of variant clones each of which differs by a single amino acid alteration in a single CDR and which contain variants representing each possible amino acid substitution for each CDR residue. 10 Mutants with increased (or decreased) binding affinity for the antigen can be screened by contacting the immobilized mutants with labeled antigen. Any screening method known in the art can be used to identify mutant antibodies with increased or decreased affinity to the antigen (e.g., ELISA) (See Wu *et al.* 1998, Proc. Natl. Acad. Sci. (U.S.A.) 95:6037; Yelton *et al.*, 1995, J. 15 Immunology 155:1994). CDR walking which randomizes the Light Chain may be used possible (see, Schier *et al.*, 1996, J. Mol. Bio. 263:551).

[0087] Methods for accomplishing such affinity maturation are described for example in: Krause, J.C. *et al.* (2011) "An Insertion Mutation That Distorts Antibody Binding Site Architecture Enhances Function Of A Human Antibody," 20 MBio. 2(1) pii: e00345-10. doi: 10.1128/mBio.00345-10; Kuan, C.T. *et al.* (2010) "Affinity-Matured Anti-Glycoprotein NMB Recombinant Immunotoxins Targeting Malignant Gliomas And Melanomas," Int. J. Cancer 10.1002/ijc.25645; Hackel, B.J. *et al.* (2010) "Stability And CDR Composition Biases Enrich Binder Functionality Landscapes," J. Mol. Biol. 401(1):84-96; Montgomery, D.L. *et al.* 25 (2009) "Affinity Maturation And Characterization Of A Human Monoclonal Antibody Against HIV-1 gp41," MAbs 1(5):462-474; Gustchina, E. *et al.* (2009) "Affinity Maturation By Targeted Diversification Of The CDR-H2 Loop Of A Monoclonal Fab Derived From A Synthetic Naïve Human Antibody Library And Directed Against The Internal Trimeric Coiled-Coil Of Gp41 Yields A Set Of 30 Fabs With Improved HIV-1 Neutralization Potency And Breadth," Virology 393(1):112-119; Finlay, W.J. *et al.* (2009) "Affinity Maturation Of A Humanized Rat Antibody For Anti-RAGE Therapy: Comprehensive Mutagenesis Reveals A

High Level Of Mutational Plasticity Both Inside And Outside The Complementarity-Determining Regions,” J. Mol. Biol. 388(3):541-558; Bostrom, J. et al. (2009) “Improving Antibody Binding Affinity And Specificity For Therapeutic Development,” Methods Mol. Biol. 525:353-376; Steidl, S. et al.
5 *(2008) “In Vitro Affinity Maturation Of Human GM-CSF Antibodies By Targeted CDR-Diversification,” Mol. Immunol. 46(1):135-144; and Barderas, R. et al. (2008) “Affinity Maturation Of Antibodies Assisted By In Silico Modeling,” Proc. Natl. Acad. Sci. (USA) 105(26):9029-9034.*

[0088] Thus, the sequence of CDR variants of encompassed antibodies or
10 their antigen-binding fragments may differ from the sequence of the CDR of the parent antibody through substitutions; for instance substituted 4 amino acid residue, 3 amino acid residue, 2 amino acid residue or 1 of the amino acid residues. According to an embodiment of the invention it is furthermore envisaged that the amino acids in the CDR regions may be substituted with
15 conservative substitutions, as defined in the below 3 tables.

[0089] The term "transgenic non-human animal" refers to a non-human animal having a genome comprising one or more human heavy and/or light chain transgenes or trans-chromosomes (either integrated or non-integrated into the animal's natural genomic DNA) and which is capable of expressing fully
20 human antibodies. For example, a transgenic mouse can have a human light chain transgene and either a human heavy chain transgene or human heavy chain trans-chromosome, such that the mouse produces human anti-Sortilin antibody when immunized with Sortilin antigen and/or cells expressing Sortilin. The human heavy chain transgene may be integrated into the chromosomal
25 DNA of the mouse, as is the case for transgenic mice, for instance HuMAb mice, such as HCo7 or HCo12 mice, or the human heavy chain transgene may be maintained extra-chromosomally, as is the case for trans-chromosomal KM mice as described in WO02/43478. Such transgenic and trans-chromosomal mice (collectively referred to herein as "transgenic mice") are capable of
30 producing multiple isotypes of human monoclonal antibodies to a given antigen

(such as IgG, IgA, IgM, IgD and/or IgE) by undergoing V-D-J recombination and isotype switching.

[0090] Transgenic, nonhuman animal can also be used for production of antibodies against a specific antigen by introducing genes encoding such specific antibody, for example by operatively linking the genes to a gene which
5 is expressed in the milk of the animal.

[0091] The term "treatment" or "treating" as used herein means ameliorating, slowing, attenuating or reversing the progress or severity of a disease or disorder, or ameliorating, slowing, attenuating or reversing one or
10 more symptoms or side effects of such disease or disorder. For purposes of this invention, "treatment" or "treating" further means an approach for obtaining beneficial or desired clinical results, where "beneficial or desired clinical results" include, without limitation, alleviation of a symptom, diminishment of the extent of a disorder or disease, stabilized (i.e., not worsening) disease or disorder
15 state, delay or slowing of the progression a disease or disorder state, amelioration or palliation of a disease or disorder state, and remission of a disease or disorder, whether partial or total detectable or undetectable.

[0092] An "effective amount," when applied to an antibody or antigen-binding fragment thereof of the invention, refers to an amount sufficient, at dosages and
20 for periods of time necessary, to achieve an intended biological effect or a desired therapeutic result including, without limitation, clinical results. The phrase "therapeutically effective amount," when applied to an antibody or antigen-binding fragment thereof of the invention, is intended to denote an amount of the antibody, or antigen-binding fragment thereof, that is sufficient to
25 ameliorate, palliate, stabilize, reverse, slow, attenuate or delay the progression of a disorder or disease state, or of a symptom of the disorder or disease. In an embodiment, the method of the present invention provides for administration of the antibody, or antigen-binding fragment thereof, in combinations with other compounds. In such instances, the "effective amount" is the amount of the
30 combination sufficient to cause the intended biological effect.

[0093] A therapeutically effective amount of an anti-Sortilin antibody or antigen-binding fragment thereof of the invention may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the anti-Sortilin antibody or antigen-binding fragment thereof to elicit a
5 desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects.

[0094] The antibodies are preferably a human or humanized antibody.
10

[0095] The numbering of amino acid residues in this region is according to IMGT®, the international ImMunoGeneTics information system® or, Kabat, E. A., Wu, T. T., Perry, H. M., Gottesmann, K. S. & Foeller, C. (1991). Sequences of Proteins of Immunological Interest, 5th edit., NIH Publication no. 91-3242
15 U.S. Department of Health and Human Services; Chothia, C. & Lesk, A. M. (1987). Canonical structures For The Hypervariable domains Of Immunoglobulins. J. Mol. Biol. 196, 901-917.

Antibody 5E1:

[0096] Accordingly, the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
20

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:1**;
- (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:2**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:3**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:4**;
- 25 (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:5**;
and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:6**.

[0097] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:8** and the light chain variable domain of **SEQ ID NO:7**.

Antibody 1F2:

- 5 [0098] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:9**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:10**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:11**;
 - 10 (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:12**;
 - (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:13**; and
 - 15 (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:14**.

[0099] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:16** and the light chain variable domain of **SEQ ID NO:15**.

Antibody 068:

- 20 [00100] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:17**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:18**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:19**;
 - 25 (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:20**;

(e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:21**; and

(f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:22**.

- 5 **[00101]** Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:24** and the light chain variable domain of **SEQ ID NO:23**.

Antibody 1320:

[00102] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- 10 (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:25**;
(b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:26**;
(c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:27**;
(d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID**
15 **NO:28**;

(e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:29**; and

(f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:30**.

- 20 **[00103]** Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:32** and the light chain variable domain of **SEQ ID NO:31**.

Antibody 93-05:

[00104] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- 25 (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:33**;
(b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:34**;
(c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:35**;

- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:36**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:37**; and
- 5 (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:38**.

[00105] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:40** and the light chain variable domain of **SEQ ID NO:39**.

10 **Antibody 93-01:**

[00106] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:41**;
- (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:42**;
- 15 (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:43**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:44**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:45**; and
- 20 (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:46**.

[00107] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:48** and the light chain variable domain of **SEQ ID NO:47**.

25 **Antibody 924:**

[00108] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:49**;
- (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:50**;

- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:51**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:52**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:53**; and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:54**.

[00109] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:56** and the light chain variable domain of **SEQ ID NO:55**.

Antibody 1276:

- [00110] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:57**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:58**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:59**;
 - (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:60**;
 - (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:61**; and
 - (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:62**.

[00111] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:64** and the light chain variable domain of **SEQ ID NO:63**.

Antibody 849:

[00112] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:65**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:66**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:67**;
 - (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID**
5 **NO:68**;
 - (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID**
NO:69; and
 - (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID**
NO:70.
- 10 **[00113]** Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:72** and the light chain variable domain of **SEQ ID NO:71**.

Antibody 531-02:

- 15 **[00114]** According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:73**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:74**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:75**;
 - (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID**
20 **NO:76**;
 - (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID**
NO:77; and
 - (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID**
NO:78.
- 25 **[00115]** Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:80** and the light chain variable domain of **SEQ ID NO:79**.

Antibody 548-01:

[00116] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:81**;
- 5 (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:82**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:83**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:84**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID**
10 **NO:85**; and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:86**.

[00117] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:88** and the light chain variable
15 domain of **SEQ ID NO:87**.

Antibody 548-02:

[00118] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:89**;
- 20 (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:90**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:91**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:92**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID**
25 **NO:93**; and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:94**.

[00119] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:96** and the light chain variable domain of **SEQ ID NO:95**.

Antibody 1289-02:

- 5 [00120] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:97**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:98**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:99**;
 - 10 (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:100**;
 - (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:101**; and
 - (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID**
- 15 **NO:102**.

[00121] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:104** and the light chain variable domain of **SEQ ID NO:103**.

Antibody 811-02:

- 20 [00122] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:105**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:106**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:107**;
 - 25 (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:108**;
 - (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:109**; and

(f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:110**.

[00123] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:112** and the light chain variable
5 domain of **SEQ ID NO:111**.

Antibody 566-01:

[00124] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:113**;
- 10 (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:114**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:115**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:116**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID**
15 **NO:117**; and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:118**.

[00125] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:120** and the light chain variable
20 domain of **SEQ ID NO:119**.

Antibody 562:

[00126] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:121**;
- 25 (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:122**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:123**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:124**;

- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:125**; and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:126**.
- 5 **[00127]** Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:128** and the light chain variable domain of **SEQ ID NO:127**.

Antibody 193:

- [00128]** According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- 10 (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:129**;
- (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:130**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:131**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID**
- 15 **NO:132**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:133**; and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:134**.
- 20 **[00129]** Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:136** and the light chain variable domain of **SEQ ID NO:135**.

Antibody 88:

- [00130]** According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- 25 (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:137**;
- (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:138**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:139**;

- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:140**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:141**; and
- 5 (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:142**.

[00131] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:144** and the light chain variable domain of **SEQ ID NO:143**.

10 **Antibody 045:**

[00132] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:145**;
- (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:146**;
- 15 (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:147**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:148**;
- (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:149**; and
- 20 (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:150**.

[00133] Preferably, the monoclonal antibody may comprise or consist of the heavy chain variable domain of **SEQ ID NO:152** and the light chain variable domain of **SEQ ID NO:151**.

25 **Antibody 044:**

[00134] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:

- (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:153**;

- (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:154**;
- (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:155**;
- (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:156**;
- 5 (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:157**; and
- (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:158**.

[00135] Preferably, the monoclonal antibody may comprise or consist of the
10 heavy chain variable domain of **SEQ ID NO:160** and the light chain variable domain of **SEQ ID NO:159**.

Antibody 002:

- [00136] According to another embodiment the invention relates to an antibody, or an antigen-binding fragment thereof, comprising or consisting of:
- 15 (a) a Light Chain CDR1 having the amino acid sequence of **SEQ ID NO:161**;
 - (b) a Light Chain CDR2 having the amino acid sequence of **SEQ ID NO:162**;
 - (c) a Light Chain CDR3 having the amino acid sequence of **SEQ ID NO:163**;
 - (d) a Heavy Chain CDR1 having the amino acid sequence of **SEQ ID NO:164**;
 - 20 (e) a Heavy Chain CDR2 having the amino acid sequence of **SEQ ID NO:165**; and
 - (f) a Heavy Chain CDR3 having the amino acid sequence of **SEQ ID NO:166**.

[00137] Preferably, the monoclonal antibody may comprise or consist of the
25 heavy chain variable domain of **SEQ ID NO:168** and the light chain variable domain of **SEQ ID NO:167**.

[00138] The antibodies mentioned above may, according to one embodiment, further comprise a variant with no more than 4 amino acid differences, or no more than 3 amino acid differences, or no more than 2 amino

acid differences, or no more than 1 amino acid difference from said CDR1, CDR2, and/or CDR3 (VH and/or VL) sequences.

[00139] Further, the antibodies may be in a composition together with a pharmaceutically acceptable carrier. The antibodies of the invention may be used in therapy. In particular, the antibodies of the invention may be used in treating FTD or ALS or TDP43 proteinopathies such as Alzheimer's Disease (AD).

[00140] The treatment envisioned by the present invention may be chronic and the patient may be treated at least 2 weeks, such as at least for 1 month, 6, months, 1 year or more.

[00141] The antibodies of the present invention may, for example, be monoclonal antibodies produced by the hybridoma method first described by Kohler et al., Nature 256, 495 (1975), or may be monoclonal antibodies produced by recombinant DNA or other methods. Monoclonal antibodies may also be isolated from phage antibody libraries using the techniques described in, for example, Clackson et al., Nature 352, 624-628 (1991) and Marks et al., J. Mol. Biol. 222, 581-597 (1991). Monoclonal antibodies may be obtained from any suitable source. Thus, for example, monoclonal antibodies may be obtained from hybridomas prepared from murine splenic B lymphocyte cells obtained from mice immunized with an antigen of interest, for instance, in the form of cells expressing the antigen on the surface, or a nucleic acid encoding an antigen of interest. Monoclonal antibodies may also be obtained from hybridomas derived from antibody-expressing cells of immunized humans or from non-human mammals such as rats, rabbits, dogs, sheep, goats, primates, etc.

[00142] In one embodiment, the antibody of the invention is a human antibody. Human monoclonal antibodies directed against Sortilin may be generated using transgenic or transchromosomal mice carrying parts of the human immune system rather than the mouse system. Such transgenic and

trans-chromosomic mice include mice referred to herein as HuMAb mice and KM mice, respectively.

[00143] The HuMAb mouse contains a human immunoglobulin gene minilocus that encodes unrearranged human heavy variable and constant (μ and Y) and light variable and constant (κ) chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous μ and K chain loci (Lonberg, N. et al., Nature 368, 856-859 (1994)). Accordingly, the mice exhibit reduced expression of mouse IgM or K and in response to immunization, the introduced human heavy and light chain transgenes, undergo class switching and somatic mutation to generate high affinity human IgG, κ monoclonal antibodies (Lonberg, N. et al. (1994), supra; reviewed in Lonberg, N., Handbook of Experimental Pharmacology 113, 49-101 (1994), Lonberg, N. and Huszar, D., Intern. Rev. Immunol. Vol. 13 65-93 (1995) and Harding, F. and Lonberg, N., Ann. N. Y. Acad. Sci 764 536-546 (1995)). The preparation of HuMAb mice is described in detail in Taylor, L. et al., Nucleic Acids Research 20, 6287-6295 (1992), Chen, J. et al., International Immunology 5, 647-656 (1993), Tuaillon et al., J. Immunol. 152, 2912-2920 (1994), Taylor, L. et al., International Immunology 6, 579-591 (1994), Fishwild, D. et al., Nature Biotechnology 14, 845-851 (1996). See also US 5,545,806, US 5,569,825, US 5,625,126, US 5,633,425, US 5,789,650, US 5,877,397, US 5,661,016, US 5,814,318, US 5,874,299, US 5,770,429, US 5,545,807, WO 98/24884, WO 94/25585, WO 93/1227, WO 92/22645, WO 92/03918 and WO 01/09187.

[00144] The HCo7, HCo12, HCo17 and HCo20 mice have a JKD disruption in their endogenous light chain (kappa) genes (as described in Chen et al., EMBO J. 12, 811-820 (1993)), a CMD disruption in their endogenous heavy chain genes (as described in **Example 1** of WO 01/14424), and a KCo5 human kappa light chain transgene (as described in Fishwild et al., Nature Biotechnology 14, 845-851 (1996)). Additionally, the HCo7 mice have a HCo7 human heavy chain transgene (as described in US 5,770,429), the HCo12 mice have a HCo12 human heavy chain transgene (as described in **Example 2** of WO 01/14424), the HCo17 mice have a HCo17 human heavy chain transgene

(as described in **Example 2** of WO 01/09187) and the HCo20 mice have a HCo20 human heavy chain transgene. The resulting mice express human immunoglobulin heavy and kappa light chain transgenes in a background homozygous for disruption of the endogenous mouse heavy and kappa light chain loci.

[00145] In the KM mouse strain, the endogenous mouse kappa light chain gene has been homozygously disrupted as described in Chen et al., EMBO J. 12, 811-820 (1993) and the endogenous mouse heavy chain gene has been homozygously disrupted as described in **Example 1** of WO 01/09187. This mouse strain carries a human kappa light chain transgene, KCo5, as described in Fishwild et al., Nature Biotechnology 14, 845-851 (1996). This mouse strain also carries a human heavy chain transchromosome composed of chromosome 14 fragment hCF (SC20) as described in WO 02/43478. HCo12-Balb/c, HCo17-Balb/c and HCo20-Balb/c mice can be generated by crossing HCo12, HCo17 and HCo20 to KCo5[J/K](Balb) as described in WO 09/097006.

[00146] In the KM mouse strain, the endogenous mouse kappa light chain gene has been homozygously disrupted as described in Chen et al., EMBO J. 12, 811-820 (1993) and the endogenous mouse heavy chain gene has been homozygously disrupted as described in **Example 1** of WO 01/09187. This mouse strain carries a human kappa light chain transgene, KCo5, as described in Fishwild et al., Nature Biotechnology 14, 845-851 (1996). This mouse strain also carries a human heavy chain trans-chromosome composed of chromosome 14 antigen-binding fragment hCF (SC20) as described in WO 02/43478.

[00147] Splenocytes from these transgenic mice may be used to generate hybridomas that secrete human monoclonal antibodies according to well-known techniques. Human monoclonal or polyclonal antibodies of the present invention, or antibodies of the present invention originating from other species may also be generated transgenically through the generation of another non-human mammal or plant that is transgenic for the immunoglobulin heavy and

light chain sequences of interest and production of the antibody in a recoverable form therefrom. In connection with the transgenic production in mammals, antibodies may be produced in, and recovered from, the milk of goats, cows, or other mammals. See for instance US 5,827,690, US 5,756,687, US 5,750,172
5 and US 5,741,957.

[00148] The antibody of the invention may be of any isotype. The choice of isotype typically will be guided by the desired effector functions, such as ADCC induction. Exemplary isotypes are IgG1, IgG2, IgG3, and IgG4. Either of the human light chain constant domains, kappa or lambda, may be used. If desired,
10 the class of an anti-Sortilin antibody of the present invention may be switched by known methods. For example, an antibody of the present invention that was originally IgM may be class switched to an IgG antibody of the present invention. Further, class switching techniques may be used to convert one IgG subclass to another, for instance from IgG1 to IgG2. Thus, the effector function
15 of the antibodies of the present invention may be changed by isotype switching to, e.g., an IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM antibody for various therapeutic uses. In one embodiment an antibody of the present invention is an IgG1 antibody, for instance an IgG1, κ . An antibody is said to be of a particular isotype if its amino acid sequence is most homologous to that isotype, relative
20 to other isotypes.

[00149] In one embodiment, the antibody of the invention is a full-length antibody, preferably an IgG antibody, in particular an IgG1, κ antibody. In another embodiment, the antibody of the invention is an antibody antigen-binding fragment or a single-chain antibody.

25 **[00150]** Antibodies and antigen-binding fragments thereof may e.g. be obtained by antigen-binding fragmentation using conventional techniques, and antigen-binding fragments screened for utility in the same manner as described herein for whole antibodies. For example, F(ab')₂ antigen-binding fragments may be generated by treating antibody with pepsin. The resulting F(ab')₂
30 antigen-binding fragment may be treated to reduce disulfide bridges to produce

Fab' antigen-binding fragments. Fab antigen-binding fragments may be obtained by treating an IgG antibody with papain; Fab' antigen-binding fragments may be obtained with pepsin digestion of IgG antibody. An F(ab') antigen-binding fragment may also be produced by binding Fab'-described
5 below via a thioether bond or a disulfide bond. A Fab' antigen-binding fragment is an antibody antigen-binding fragment obtained by cutting a disulfide bond of the hinge domain of the F(ab')₂. A Fab'- antigen-binding fragment may be obtained by treating an F(ab')₂ antigen-binding fragment with a reducing agent, such as dithiothreitol. Antibody antigen-binding fragment may also be generated
10 by expression of nucleic acids encoding such antigen-binding fragments in recombinant cells (see for instance Evans et al., J. Immunol. Meth. 184, 123-38 (1995)). For example, a chimeric gene encoding a portion of an F(ab')₂ antigen-binding fragment could include DNA sequences encoding the CH1 domain and hinge domain of the H chain, followed by a translational stop codon to yield
15 such a truncated antibody antigen-binding fragment molecule.

[00151] In one embodiment, the anti-Sortilin antibody is a monovalent antibody, preferably a monovalent antibody as described in WO2007059782 (which is incorporated herein by reference in its entirety) having a deletion of the hinge region. Accordingly, in one embodiment, the antibody is a monovalent
20 antibody, wherein said anti-Sortilin antibody is constructed by a method comprising : i) providing a nucleic acid construct encoding the light chain of said monovalent antibody, said construct comprising a nucleotide sequence encoding the VL region of a selected antigen specific anti-Sortilin antibody and a nucleotide sequence encoding the constant CL region of an Ig, wherein said
25 nucleotide sequence encoding the VL region of a selected antigen specific antibody and said nucleotide sequence encoding the CL region of an Ig are operably linked together, and wherein, in case of an IgG1 subtype, the nucleotide sequence encoding the CL region has been modified such that the CL region does not contain any amino acids capable of forming disulfide bonds or covalent bonds with other peptides comprising an identical amino acid
30 sequence of the CL region in the presence of polyclonal human IgG or when administered to an animal or human being; ii) providing a nucleic acid construct

encoding the heavy chain of said monovalent antibody, said construct comprising a nucleotide sequence encoding the VH region of a selected antigen specific antibody and a nucleotide sequence encoding a constant CH region of a human Ig, wherein the nucleotide sequence encoding the CH region has been
5 modified such that the region corresponding to the hinge region and, as required by the Ig subtype, other regions of the CH region, such as the CH3 region, does not comprise any amino acid residues which participate in the formation of disulphide bonds or covalent or stable non-covalent inter-heavy chain bonds with other peptides comprising an identical amino acid sequence of
10 the CH region of the human Ig in the presence of polyclonal human IgG or when administered to an animal human being, wherein said nucleotide sequence encoding the VH region of a selected antigen specific antibody and said nucleotide sequence encoding the CH region of said Ig are operably linked together; iii) providing a cell expression system for producing said monovalent
15 antibody; iv) producing said monovalent antibody by co-expressing the nucleic acid constructs of (i) and (ii) in cells of the cell expression system of (iii).

[00152] Similarly, in one embodiment, the anti-Sortilin antibody is a monovalent antibody, which comprises:

- (i) a variable domain of an antibody of the invention as described herein or an
20 antigen-binding part of the said domain, and
- (ii) a CH domain of an immunoglobulin or a domain thereof comprising the CH2 and CH3 domains, wherein the CH domain or domain thereof has been modified such that the domain corresponding to the hinge domain and, if the immunoglobulin is not an IgG4 subtype, other domains of the CH
25 domain, such as the CH3 domain, do not comprise any amino acid residues, which are capable of forming disulfide bonds with an identical CH domain or other covalent or stable non-covalent inter-heavy chain bonds with an identical CH domain in the presence of polyclonal human IgG.

[00153] In a further embodiment, the heavy chain of the monovalent antibody
30 has been modified such that the entire hinge region has been deleted.

[00154] In another further embodiment, the sequence of the monovalent antibody has been modified so that it does not comprise any acceptor sites for N-linked glycosylation.

[00155] The invention also includes "Bispecific Antibodies," wherein an anti-Sortilin binding region (e.g., a Sortilin-binding region of an anti-Sortilin monoclonal antibody) is part of a bivalent or polyvalent bispecific scaffold that targets more than one epitope, (for example a second epitope could comprise an epitope of an active transport receptor, such that the Bispecific Antibody would exhibit improved transcytosis across a biological barrier, such as the Blood Brain Barrier). Thus, in another further embodiment, the monovalent Fab of an anti-Sortilin antibody may be joined to an additional Fab or scfv that targets a different protein to generate a bispecific antibody. A bispecific antibody can have a dual function, for example a therapeutic function imparted by an anti-sortilin binding domain and a transport function that can bind to a receptor molecule to enhance transfer cross a biological barrier, such as the blood brain barrier.

[00156] Antibodies and antigen-binding fragments thereof of the invention also include single chain antibodies. Single chain antibodies are peptides in which the heavy and light chain Fv domains are connected. In one embodiment, the present invention provides a single-chain Fv (scFv) wherein the heavy and light chains in the Fv of an anti-Sortilin antibody of the present invention are joined with a flexible peptide linker (typically of about 10, 12, 15 or more amino acid residues) in a single peptide chain. Methods of producing such antibodies are described in for instance US 4,946,778, Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds. Springer-Verlag, New York, pp. 269-315 (1994), Bird et al., Science 242, 423-426 (1988), Huston et al., PNAS USA 85, 5879-5883 (1988) and McCafferty et al., Nature 348, 552-554 (1990). The single chain antibody may be monovalent, if only a single VH and VL are used, bivalent, if two VH and VL are used, or polyvalent, if more than two VH and VL are used.

[00157] The antibodies and antigen-binding fragments thereof described herein may be modified by inclusion of any suitable number of modified amino acids and/or associations with such conjugated substituents. Suitability in this context is generally determined by the ability to at least substantially retain the
5 Sortilin selectivity and/or Sortilin specificity associated with the non-derivatized parent anti-Sortilin antibody. The inclusion of one or more modified amino acids may be advantageous in, for example, increasing polypeptide serum half-life, reducing polypeptide antigenicity, or increasing polypeptide storage stability. Amino acid(s) are modified, for example, co-translationally or post-
10 translationally during recombinant production (e.g., N-linked glycosylation at N-X-S/T motifs during expression in mammalian cells) or modified by synthetic means. Non-limiting examples of a modified amino acid include a glycosylated amino acid, a sulfated amino acid, a prenylated (e. g., farnesylated, geranyl-geranylated) amino acid, an acetylated amino acid, an acylated amino acid, a
15 PEGylated amino acid, a biotinylated amino acid, a carboxylated amino acid, a phosphorylated amino acid, and the like. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature. Example protocols are found in Walker (1998) Protein Protocols On CD-Rom, Humana Press, Totowa, NJ. The modified amino acid may, for instance, be
20 selected from a glycosylated amino acid, a PEGylated amino acid, a farnesylated amino acid, an acetylated amino acid, a biotinylated amino acid, an amino acid conjugated to a lipid moiety, or an amino acid conjugated to an organic derivatizing agent.

[00158] The antibodies and antigen-binding fragments thereof of the
25 invention, may also be chemically modified by covalent conjugation to a polymer to for instance increase their circulating half-life. Exemplary polymers, and methods to attach them to peptides, are illustrated in for instance US 4,766,106; US 4,179,337; US 4,495,285 and US 4,609,546. Additional illustrative polymers include polyoxyethylated polyols and polyethylene glycol
30 (PEG) (e.g., a PEG with a molecular weight of between about 1,000 and about 40,000, such as between about 2,000 and about 20,000, e.g., about 3,000-12,000 g/mol).

[00159] The antibodies and antigen-binding fragments thereof of the present invention may further be used in a diagnostic method or as a diagnostic imaging ligand.

[00160] In one embodiment, antibodies and antigen-binding fragments thereof of the invention comprising one or more radiolabeled amino acids are provided. A radiolabeled anti-Sortilin antibody may be used for both diagnostic and therapeutic purposes (conjugation to radiolabeled molecules is another possible feature). Non-limiting examples of such labels include, but are not limited to bismuth (^{213}Bi), carbon (^{11}C , ^{13}C , ^{14}C), chromium (^{51}Cr), cobalt (^{57}Co , ^{60}Co), copper (^{64}Cu), dysprosium (^{165}Dy), erbium (^{169}Er), fluorine (^{18}F), gadolinium (^{153}Gd , ^{159}Gd), gallium (^{68}Ga , ^{67}Ga), germanium (^{68}Ge), gold (^{198}Au), holmium (^{166}Ho), hydrogen (^3H), indium (^{111}In , ^{112}In , ^{113}In , ^{115}In), iodine (^{121}I , ^{123}I , ^{125}I , ^{131}I), iridium (^{192}Ir), iron (^{59}Fe), krypton ($^{81\text{m}}\text{Kr}$), lanthanum (^{140}La), lutetium (^{177}Lu), manganese (^{54}Mn), molybdenum (^{99}Mo), nitrogen (^{13}N , ^{15}N), oxygen (^{15}O), palladium (^{103}Pd), phosphorus (^{32}P), potassium (^{42}K), praseodymium (^{142}Pr), promethium (^{149}Pm), rhenium (^{186}Re , ^{188}Re), rhodium (^{105}Rh), rubidium (^{81}Rb , ^{82}Rb), ruthenium (^{82}Ru , ^{97}Ru), samarium (^{153}Sm), scandium (^{47}Sc), selenium (^{75}Se), sodium (^{24}Na), strontium (^{85}Sr , ^{89}Sr , ^{92}Sr), sulfur (^{35}S), technetium (^{99}Tc), thallium (^{201}Tl), tin (^{113}Sn , ^{117}Sn), xenon (^{133}Xe), ytterbium (^{169}Yb , ^{175}Yb , ^{177}Yb), yttrium (^{90}Y) and zinc (^{65}Zn). Methods for preparing radiolabeled amino acids and related peptide derivatives are known in the art (see for instance Junghans et al., in *Cancer Chemotherapy and Biotherapy* 655-686 (2nd edition, Chafner and Longo, eds., Lippincott Raven (1996)) and US 4,681,581, US 4,735,210, US 5,101,827, US 5,102,990 (US RE35,500), US 5,648,471 and US 5,697,902. For example, a radioisotope may be conjugated by a chloramine T method (Lindgren, S. et al. (1998) "Chloramine-T In High-Specific-Activity Radioiodination Of Antibodies Using N-Succinimidyl-3-(Trimethylstannyl)Benzoate As An Intermediate," Nucl. Med. Biol. 25(7):659-665; Kurth, M. et al. (1993) "Site-Specific Conjugation Of A Radioiodinated Phenethylamine Derivative To A Monoclonal Antibody Results In Increased Radioactivity Localization In Tumor," J. Med. Chem. 36(9):1255-1261; Rea,

D.W. *et al.* (1990) "Site-specifically radioiodinated antibody for targeting tumors," *Cancer Res.* 50(3 Suppl):857s-861s).

[00161] The invention also provides anti-Sortilin antibodies and antigen-binding fragments thereof that are detectably labeled using a fluorescent label (such as a rare earth chelate (*e.g.*, a europium chelate)), a fluorescein-type label (*e.g.*, fluorescein, fluorescein isothiocyanate, 5-carboxyfluorescein, 6-carboxy fluorescein, dichlorotriazinylamine fluorescein), a rhodamine-type label (*e.g.*, ALEXA FLUOR® 568 (Invitrogen), TAMRA® or dansyl chloride), VIVOTAG 680 XL FLUOROCHROME™ (Perkin Elmer), phycoerythrin; umbelliferone, Lissamine; a cyanine; a phycoerythrin, Texas Red, BODIPY FL-SE® (Invitrogen) or an analogue thereof, all of which are suitable for optical detection. Chemiluminescent labels may be employed (*e.g.*, luminol, luciferase, luciferin, and aequorin). Such diagnosis and detection can also be accomplished by coupling the diagnostic molecule of the present invention to detectable substances including, but not limited to, various enzymes, enzymes including, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase, or to prosthetic group complexes such as, but not limited to, streptavidin/biotin and avidin/biotin.

[00162] Chemiluminescent labels may be employed (*e.g.*, luminol, luciferase, luciferin, and aequorin). Such diagnosis and detection can also be accomplished by coupling the diagnostic molecule of the present invention to detectable substances including, but not limited to, various enzymes, enzymes including, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase, or to prosthetic group complexes such as, but not limited to, streptavidin/biotin and avidin/biotin. Paramagnetic labels can also be employed, and are preferably detected using Positron Emission Tomography (PET) or Single-Photon Emission Computed Tomography (SPECT). Such paramagnetic labels include, but are not limited to compounds containing paramagnetic ions of Aluminum (Al), Barium (Ba), Calcium (Ca), Cerium (Ce), Dysprosium (Dy), Erbium (Er), Europium (Eu), Gadolinium (Gd), Holmium (Ho), Iridium (Ir), Lithium (Li), Magnesium (Mg),

Manganese (Mn), Molybdenum (M), Neodymium (Nd), Osmium (Os), Oxygen (O), Palladium (Pd), Platinum (Pt), Rhodium (Rh), Ruthenium (Ru), Samarium (Sm), Sodium (Na), Strontium (Sr), Terbium (Tb), Thulium (Tm), Tin (Sn), Titanium (Ti), Tungsten (W), and Zirconium (Zi), and particularly, Co^{+2} , Cr^{+2} , Cr^{+3} , Cu^{+2} , Fe^{+2} , Fe^{+3} , Ga^{+3} , Mn^{+3} , Ni^{+2} , Ti^{+3} , V^{+3} , and V^{+4} , positron emitting metals using various positron emission tomographies, and non-radioactive paramagnetic metal ions.

[00163] Thus in one embodiment the anti-Sortilin antibody or Sortilin-binding fragment thereof of the invention may be labelled with a fluorescent label, a chemiluminescent label, a paramagnetic label, a radioisotopic label or an enzyme label. The labelled antibody or fragment may be used in detecting or measuring the presence or amount of said Sortilin in the brain of a subject. This method may comprise the detection or measurement of in vivo imaging of anti-Sortilin antibody or Sortilin-binding fragment bound to said Sortilin and may comprises ex vivo imaging of said anti-Sortilin antibody or Sortilin-binding fragment bound to such Sortilin.

[00164] In a further aspect, the invention relates to an expression vector encoding one or more polypeptide chains of an antibody of the invention or an antigen-binding-domain thereof. Such expression vectors may be used for recombinant production of the antibodies and antigen-binding fragments of the invention.

[00165] An expression vector in the context of the present invention may be any suitable DNA or RNA vector, including chromosomal, non-chromosomal, and synthetic nucleic acid vectors (a nucleic acid sequence comprising a suitable set of expression control elements). Examples of such vectors include derivatives of SV40, bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors derived from combinations of plasmids and phage DNA, and viral nucleic acid (RNA or DNA) vectors. In one embodiment, an anti-Sortilin antibody-encoding nucleic acid is comprised in a naked DNA or RNA vector, including, for example, a linear expression element (as described in, for

instance, Sykes and Johnston, Nat Biotech 12, 355-59 (1997)), a compacted nucleic acid vector (as described in for instance US 6,077,835 and/or WO 00/70087), a plasmid vector such as pBR322, pUC 19/18, or pUC 118/119, a "midge" minimally-sized nucleic acid vector (as described in, for instance, 5 Schakowski et al., Mol Ther 3, 793-800 (2001)), or as a precipitated nucleic acid vector construct, such as a CaPO₄-precipitated construct (as described in, for instance, WO 00/46147, Benvenisty and Reshef, PNAS USA 83, 9551-55 (1986), Wigler et al., Cell 14, 725 (1978), and Coraro and Pearson, Somatic Cell Genetics 2, 603 (1981)). Such nucleic acid vectors and the usage thereof 10 are well known in the art (see for instance US 5,589,466 and US 5,973,972).

[00166] In one embodiment, the vector is suitable for expression of anti-Sortilin antibodies or antigen-binding fragments thereof in a bacterial cell. Examples of such vectors include expression vectors such as BlueScript (Stratagene), pIN vectors (Van Heeke & Schuster, J Biol Chem 264, 5503-5509 15 (1989), pET vectors (Novagen, Madison, WI) and the like).

[00167] An expression vector may also or alternatively be a vector suitable for expression in a yeast system. Any vector suitable for expression in a yeast system may be employed. Suitable vectors include, for example, vectors comprising constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH (reviewed in: F. Ausubel *et al.*, ed. Current Protocols in 20 Molecular Biology, Greene Publishing and Wiley InterScience New York (1987), Grant *et al.*, Methods in Enzymol 153, 516-544 (1987), Mattanovich, D. *et al.* Methods Mol. Biol. 824, 329-358 (2012), Celik, E. *et al.* Biotechnol. Adv. 30(5), 1108-1118 (2012), Li, P. *et al.* Appl. Biochem. Biotechnol. 142(2), 105-124 25 (2007), Böer, E. *et al.* Appl. Microbiol. Biotechnol. 77(3), 513-523 (2007), van der Vaart, J.M. Methods Mol. Biol. 178, 359-366 (2002), and Holliger, P. Methods Mol. Biol. 178, 349-357 (2002)).

[00168] In an expression vector of the invention, anti-Sortilin antibody-encoding nucleic acids may comprise or be associated with any suitable 30 promoter, enhancer, and other expression-facilitating elements. Examples of

such elements include strong expression promoters (e. g., human CMV IE promoter/enhancer as well as RSV, SV40, SL3-3, MMTV, and HIV LTR promoters), effective poly (A) termination sequences, an origin of replication for plasmid product in E. coli, an antibiotic resistance gene as selectable marker, and/or a convenient cloning site (e.g., a polylinker). Nucleic acids may also comprise an inducible promoter as opposed to a constitutive promoter such as CMV IE (the skilled artisan will recognize that such terms are actually descriptors of a degree of gene expression under certain conditions).

[00169] In an even further aspect, the invention relates to a recombinant eukaryotic or prokaryotic host cell, such as a transfectoma, which produces an antibody or antigen-binding fragment thereof of the invention as defined herein or a bispecific molecule of the invention as defined herein. Examples of host cells include yeast, bacteria, and mammalian cells, such as CHO or HEK cells. For example, in one embodiment, the present invention provides a cell comprising a nucleic acid stably integrated into the cellular genome that comprises a sequence coding for expression of an anti-Sortilin antibody of the present invention or an antigen-binding fragment thereof. In another embodiment, the present invention provides a cell comprising a non-integrated nucleic acid, such as a plasmid, cosmid, phagemid, or linear expression element, which comprises a sequence coding for expression of an anti-Sortilin antibody or antigen-binding fragment thereof of the invention.

[00170] In a further aspect, the invention relates to a method for producing an anti-Sortilin antibody of the invention, said method comprising the steps of a) culturing a hybridoma or a host cell of the invention as described herein above, and b) purifying the antibody of the invention from the culture media.

[00171] In one embodiment, the invention relates to a preparation that, as such term is used herein, comprises an anti-Sortilin antibody as defined herein, and that is substantially free of naturally-arising antibodies that are either not capable of binding to sortilin or that do not materially alter the anti-Sortilin functionality of the preparation. Thus, such a preparation does not encompass

naturally-arising serum, or a purified derivative of such serum, that comprises a mixture of an anti-Sortilin antibody and another antibody that does not alter the functionality of the anti-Sortilin antibody of the preparation, wherein such functionality is:

- 5 (i) a binding affinity (K_D) for Sortilin;
- (ii) a capability to reduce and/or inhibit PGRN binding to Sortilin;
- (iii) a capability to reduce and/or inhibit clearance of PGRN by Sortilin-expressing cells;
- (iv) a capability to reduce and/or inhibit the endocytosis of PGRN by
10 Sortilin-expressing cells;
- (v) a capability to increase the amount and/or concentration of PGRN in the plasma in human-Sortilin-expressing knock-in mice; a capability to increase the amount and/or concentration of PGRN in the brain and/or
- 15 (vi) a capability, when administered chronically, to provide treatment of frontotemporal dementia (FTD) and/or amyotrophic lateral sclerosis (ALS).

[00172] The invention particularly relates to preparations of such an anti-Sortilin antibody having a structural change in its amino acid sequence (in any
20 of its CDRs, variable domains, framework residues and/or constant domains) relative to the structure of a naturally-occurring anti-Sortilin antibody, wherein said structural change causes the anti-Sortilin antibody monoclonal antibody to exhibit a markedly altered functionality (i.e., more than a 20% difference, more than a 40% difference, more than a 60% difference, more than an 80%
25 difference, more than a 100% difference, more than a 150% difference, more than a 2-fold difference, more than a 4-fold difference, more than a 5-fold difference, or more than a 10-fold difference in functionality) relative to the functionality exhibited by said naturally-occurring anti-Sortilin antibody; wherein such functionality is:

- 30 (i) a binding affinity (K_D) for Sortilin;
- (ii) a capability to reduce and/or inhibit PGRN binding to Sortilin;

- (iii) a capability to reduce and/or inhibit clearance of PGRN by Sortilin-expressing cells;
- (iv) a capability to reduce and/or inhibit the endocytosis of PGRN by Sortilin-expressing cells;
- 5 (vi) a capability to increase the amount and/or concentration of PGRN in the plasma in human-Sortilin-expressing knock-in mice;
- (vii) a capability to increase the amount and/or concentration of PGRN in the brain and/or
- (vi) a capability, when administered chronically, to provide treatment of
- 10 frontotemporal dementia (FTD), amyotrophic lateral sclerosis (ALS) and/or Alzheimer's Disease (AD).

especially wherein such altered functionality is a result of the structural change and thus is inseparable from it.

- 15 **[00173]** The term "substantially free" of naturally-arising antibodies refers to the complete absence of such naturally-arising antibodies in such preparations, or of the inclusion of a concentration of such naturally-arising antibodies in such preparations that does not materially affect the Sortilin-binding properties of the preparations. An antibody is said to be "isolated" if it has no naturally-arising
- 20 counterpart or has been separated or purified from components which naturally accompany it.

- [00174]** The term "naturally-arising antibodies," as it relates to such preparations, refers to antibodies (including naturally-arising autoantibodies) elicited within living humans or other animals, as a natural consequence to the
- 25 functioning of their immune systems.

- [00175]** Thus, the preparations of the present invention do not exclude, and indeed explicitly encompass, such preparations that contain an anti-Sortilin antibody and a deliberately added additional antibody capable of binding to an epitope that is not possessed by Sortilin. Such preparations particularly include
- 30 embodiments thereof wherein the preparation exhibits enhanced efficacy in

treating frontotemporal dementia (FTD) and/or amyotrophic lateral sclerosis (ALS).

[00176] The antibodies of antigen-binding fragments thereof of the present invention may be produced in different cell lines, such as a human cell line, a mammal non-human cell line, and insect cell line, for example a CHO cell line, HEK cell line, BHK-21 cell line, murine cell line (such as a myeloma cell line), fibrosarcoma cell line, PER.C6 cell line, HKB-11 cell line, CAP cell line and HuH-7 human cell line (Dumont et al, 2015, Crit Rev Biotechnol. Sep 18:1-13., the contents which is included herein by reference).

[00177] In an even further aspect, the invention relates to a pharmaceutical composition comprising:

- (i) an anti-Sortilin antibody or antigen-binding fragment thereof, both as defined herein, or a preparation, as such term is defined herein, that comprises such an anti-Sortilin antibody or antigen-binding fragment thereof; and
- (ii) a pharmaceutically-acceptable carrier.

[00178] The pharmaceutical compositions may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 22nd Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 2013.

[00179] The pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients should be suitable for the chosen compound of the present invention and the chosen mode of administration. Suitability for carriers and other components of pharmaceutical compositions is determined based on the lack of significant negative impact on the desired biological properties of the chosen compound or pharmaceutical composition of the present invention (e.g., less than a substantial impact (10% or less relative inhibition, 5% or less relative inhibition, etc.)) on epitope binding.

[00180] A pharmaceutical composition of the present invention may also include diluents, fillers, salts, buffers, detergents (e.g., a non-ionic detergent, such as Tween-20 or Tween- 80), stabilizers (e.g., sugars or protein-free amino acids), preservatives, tissue fixatives, solubilizers, and/or other materials
5 suitable for inclusion in a pharmaceutical composition. The diluent is selected to not to affect the biological activity of the combination. Examples of such diluents are distilled water, physiological phosphate-buffered saline, Ringer's solutions, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation may also include other carriers, or non-toxic,
10 nontherapeutic, non-immunogenic stabilizers and the like. The compositions may also include large, slowly metabolized macromolecules, such as proteins, polysaccharides like chitosan, polylactic acids, polyglycolic acids and copolymers (e.g., latex functionalized sepharose, agarose, cellulose, and the like), polymeric amino acids, amino acid copolymers, and lipid aggregates (e.g.,
15 oil droplets or liposomes).

[00181] The actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of
20 administration. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs,
25 compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

[00182] The pharmaceutical composition may be administered by any
30 suitable route and mode, including: parenteral, topical, oral or intranasal means for prophylactic and/or therapeutic treatment. In one embodiment, a

pharmaceutical composition of the present invention is administered parenterally. The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and include epidermal, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, intratendinous, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, intracranial, intrathoracic, epidural and intrasternal injection and infusion. Additional suitable routes of administering a compound of the present invention in vivo and in vitro are well known in the art and may be selected by those of ordinary skill in the art. In one embodiment that pharmaceutical composition is administered by intravenous or subcutaneous injection or infusion.

[00183] Pharmaceutically acceptable carriers include any and all suitable solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonicity agents, antioxidants and absorption delaying agents, and the like that are physiologically compatible with a compound of the present invention.

[00184] Examples of suitable aqueous and non-aqueous carriers which may be employed in the pharmaceutical compositions of the present invention include water, saline, phosphate buffered saline, ethanol, dextrose, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, corn oil, peanut oil, cottonseed oil, and sesame oil, carboxymethyl cellulose colloidal solutions, tragacanth gum and injectable organic esters, such as ethyl oleate, and/or various buffers. Other carriers are well known in the pharmaceutical arts.

[00185] Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active

compound, use thereof in the pharmaceutical compositions of the present invention is contemplated.

[00186] Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in
5 the case of dispersions, and by the use of surfactants.

[00187] Pharmaceutical compositions of the present invention may also comprise pharmaceutically acceptable antioxidants for instance (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble
10 antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha- tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

[00188] Pharmaceutical compositions of the present invention may also
15 comprise isotonicity agents, such as sugars, polyalcohols, such as mannitol, sorbitol, glycerol or sodium chloride in the compositions.

[00189] The pharmaceutical compositions of the present invention may also contain one or more adjuvants appropriate for the chosen route of administration such as preservatives, wetting agents, emulsifying agents,
20 dispersing agents, preservatives or buffers, which may enhance the shelf life or effectiveness of the pharmaceutical composition. The compounds of the present invention may be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Such carriers
25 may include gelatin, glyceryl monostearate, glyceryl distearate, biodegradable, biocompatible polymers such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid alone or with a wax, or other materials well known in the art. Methods for the preparation of such formulations are generally known to those skilled in the art. See, e.g.,

Sustained and Controlled Release Drug Delivery Systems, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

[00190] In one embodiment, the compounds of the present invention may be formulated to ensure proper distribution in vivo. Pharmaceutically acceptable carriers for parenteral administration include sterile aqueous solutions or
5 dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use
10 thereof in the pharmaceutical compositions of the present invention is contemplated. Supplementary active compounds may also be incorporated into the compositions.

[00191] Pharmaceutical compositions for injection must typically be sterile and stable under the conditions of manufacture and storage. The composition
15 may be formulated as a solution, micro-emulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier may be an aqueous or non-aqueous solvent or dispersion medium containing for instance water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and
20 injectable organic esters, such as ethyl oleate. The proper fluidity may be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as glycerol, mannitol, sorbitol, or
25 sodium chloride in the composition. Prolonged absorption of the injectable compositions may be brought about by including in the composition an agent that delays antibody absorption, for example, monostearate salts and gelatin. Sterile injectable solutions may be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a
30 combination of ingredients e.g. as enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating

the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients e.g. from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, examples of methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional
5 desired ingredient from a previously sterile-filtered solution thereof.

[00192] Sterile injectable solutions may be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by
10 sterilization microfiltration. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, examples of methods of preparation are vacuum drying and freeze-drying
15 (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[00193] Dosage regimens in the above methods of treatment and uses described herein are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered,
20 several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the
25 subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the present invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic
30 effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

[00194] The effective dosages and the dosage regimens for the anti-Sortilin antibodies depend on the disease or condition to be treated and may be determined by the persons skilled in the art. An exemplary, non-limiting range for a therapeutically effective amount of an antibody of the present invention is
5 about 0.1-10 mg/kg/body weight, such as about 0.1-5 mg/kg/body weight, for example about 0.1-2 mg/kg/body weight, such as about 0.1-1 mg/kg/body weight, for instance about 0.15, about 0.2, about 0.5, about 1, about 1.5 or about 2 mg/kg/body weight.

[00195] A physician or veterinarian having ordinary skill in the art may readily
10 determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the anti-Sortilin antibody employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In
15 general, a suitable daily dose of a composition of the present invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Administration may e.g. be intravenous, intramuscular, intraperitoneal, or subcutaneous, and for instance administered proximal to the
20 site of the target. If desired, the effective daily dose of a pharmaceutical composition may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the
25 compound as a pharmaceutical composition as described above.

[00196] The labelled antibodies or antigen-binding fragments thereof of the invention can be used for diagnostic purposes to detect, diagnose, or monitor diseases or disorders. The invention provides for the detection or diagnosis of a neurodegenerative or cognitive disease or disorder, including but not limited to
30 FTD, ALS or TDP43 proteinopathies such as Alzheimer's Disease (AD), comprising: (a) assaying the existence of pyroglutamated A β fragments in cells

or tissue samples of a subject using one or more antibodies that specifically bind to Sortilin; and (b) comparing the level of the antigen with a control level, e.g. levels in normal tissue samples, whereby an increase in the assayed level of antigen compared to the control level of antigen is indicative of the disease or disorder, or indicative of the severity of the disease or disorder.

[00197] The antibodies or antigen-binding fragments thereof of the invention can be used to assay Sortilin or antigen-binding fragments of Sortilin in a biological sample using immuno-histochemical methods well-known in the art. Other antibody-based methods useful for detecting protein include immunoassays such as the enzyme linked immunoassay (ELISA) and the radioimmunoassay assay (RIA) and mesoscale discovery platform based assays (MSD). Suitable antibody labels may be used in such kits and methods, and labels known in the art include enzyme labels, such as alkaline phosphatase and glucose oxidase; radioisotope labels, such as iodine (^{125}I , ^{131}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{121}In), and technetium ($^{99\text{m}}\text{Tc}$); and luminescent labels, such as luminol and luciferase; and fluorescent labels, such as fluorescein and rhodamine.

[00198] The presence of labeled anti-Sortilin antibodies or their Sortilin-binding fragments may be detected in vivo for diagnostic purposes. In one embodiment, diagnosis comprises: a) administering to a subject an effective amount of such labeled molecule; b) waiting for a time interval following administration to allow the labeled molecule to concentrate at sites (if any) of A β deposition and to allow for unbound labeled molecule to be cleared to background level; c) determining a background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level is indicative that the subject has the disease or disorder, or is indicative of the severity of the disease or disorder. In accordance with such embodiment, the molecule is labeled with an imaging moiety suitable for detection using a particular imaging system known to those skilled in the art. Background levels may be determined by various methods known in the art, including comparing the amount of labeled antibody detected to a standard

value previously determined for a particular imaging system. Methods and systems that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as positron emission tomography (PET), magnetic resonance imaging (MRI), and
5 sonography.

[00199] In a further aspect, the invention relates to an antibody, or antigen-binding fragment thereof, of the invention, for use in medicine.

[00200] In a further aspect, the invention relates to an antibody, or antigen-binding fragment thereof, of the invention, for use in treating a disease
10 associated with decreased PGRN levels in the brain of a patient,

[00201] In a further aspect, the invention relates to the use of the antibody, or antigen-binding fragment thereof, of the invention, in the manufacture of a medicament for treating a disease associated with decreased PGRN levels in the brain of a patient,

15 **[00202]** In a further aspect, the invention relates to a method of preventing or treating a disease associated with decreased PGRN levels in the brain of a patient, comprising administering an effective dosage of an antibody of the invention, or an antigen-binding fragment thereof.

[00203] It is preferred that in the uses and methods of those aspects of the invention the disease is: FTD; ALS; or TDP43 proteinopathies, such as AD.

[00204] Preferably, in the uses and methods of those aspects of the
20 invention, the treatment is chronic, and is preferably for at least 2 weeks, such as at least for 1 month, 6, months, 1 year or more.

[00205] In a further aspect, the invention provides a kit comprising the antibody, or antigen-binding fragment thereof, of the invention.

25

Table 5 - Antibody Sequences

Ab 5E1		
Seq ID No:	1	5E1 CDR1 Light Chain
Seq ID No:	2	5E1 CDR2 Light Chain
Seq ID No:	3	5E1 CDR3 Light Chain
Seq ID No:	4	5E1 CDR1 Heavy Chain
Seq ID No:	5	5E1 CDR2 Heavy Chain
Seq ID No:	6	5E1 CDR3 Heavy Chain
Seq ID No:	7	5E1 VL
Seq ID No:	8	5E1 VH
Ab 1F2		
Seq ID No:	9	1F2 CDR1 Light Chain
Seq ID No:	10	1F2 CDR2 Light Chain
Seq ID No:	11	1F2 CDR3 Light Chain
Seq ID No:	12	1F2 CDR1 Heavy Chain
Seq ID No:	13	1F2 CDR2 Heavy Chain
Seq ID No:	14	1F2 CDR3 Heavy Chain
Seq ID No:	15	1F2 VL
Seq ID No:	16	1F2 VH
Ab 068		
Seq ID No:	17	068 CDR1 Light Chain
Seq ID No:	18	068 CDR2 Light Chain
Seq ID No:	19	068 CDR3 Light Chain
Seq ID No:	20	068 CDR1 Heavy Chain
Seq ID No:	21	068 CDR2 Heavy Chain
Seq ID No:	22	068 CDR3 Heavy Chain
Seq ID No:	23	068 VL
Seq ID No:	24	068 VH
Ab 1320		
Seq ID No:	25	1320 CDR1 Light Chain
Seq ID No:	26	1320 CDR2 Light Chain
Seq ID No:	27	1320 CDR3 Light Chain
Seq ID No:	28	1320 CDR1 Heavy Chain
Seq ID No:	29	1320 CDR2 Heavy Chain
Seq ID No:	30	1320 CDR3 Heavy Chain
Seq ID No:	31	1320 VL
Seq ID No:	32	1320 VH
Ab 93-05		
Seq ID No:	33	93-05 CDR1 Light Chain
Seq ID No:	34	93-05 CDR2 Light Chain
Seq ID No:	35	93-05 CDR3 Light Chain
Seq ID No:	36	93-05 CDR1 Heavy Chain
Seq ID No:	37	93-05 CDR2 Heavy Chain

Seq ID No: 38	93-05 CDR3 Heavy Chain
Seq ID No: 39	93-05 VL
Seq ID No: 40	93-05 VH
Ab 93-01	
Seq ID No: 41	93-01 CDR1 Light Chain
Seq ID No: 42	93-01 CDR2 Light Chain
Seq ID No: 43	93-01 CDR3 Light Chain
Seq ID No: 44	93-01 CDR1 Heavy Chain
Seq ID No: 45	93-01 CDR2 Heavy Chain
Seq ID No: 46	93-01 CDR3 Heavy Chain
Seq ID No: 47	93-01 VL
Seq ID No: 48	93-01 VH
Ab 924	
Seq ID No: 49	924 CDR1 Light Chain
Seq ID No: 50	924 CDR2 Light Chain
Seq ID No: 51	924 CDR3 Light Chain
Seq ID No: 52	924 CDR1 Heavy Chain
Seq ID No: 53	924 CDR2 Heavy Chain
Seq ID No: 54	924 CDR3 Heavy Chain
Seq ID No: 55	924 VL
Seq ID No: 56	924 VH
Ab 1276	
Seq ID No: 57	1276 CDR1 Light Chain
Seq ID No: 58	1276 CDR2 Light Chain
Seq ID No: 59	1276 CDR3 Light Chain
Seq ID No: 60	1276 CDR1 Heavy Chain
Seq ID No: 61	1276 CDR2 Heavy Chain
Seq ID No: 62	1276 CDR3 Heavy Chain
Seq ID No: 63	1276 VL
Seq ID No: 64	1276 VH
Ab 849	
Seq ID No: 65	849 CDR1 Light Chain
Seq ID No: 66	849 CDR2 Light Chain
Seq ID No: 67	849 CDR3 Light Chain
Seq ID No: 68	849 CDR1 Heavy Chain
Seq ID No: 69	849 CDR2 Heavy Chain
Seq ID No: 70	849 CDR3 Heavy Chain
Seq ID No: 71	849 VL
Seq ID No: 72	849 VH
Ab 531-02	
Seq ID No: 73	531-02 CDR1 Light Chain
Seq ID No: 74	531-02 CDR2 Light Chain
Seq ID No: 75	531-02 CDR3 Light Chain

Seq ID No: 76	531-02 CDR1 Heavy Chain
Seq ID No: 77	531-02 CDR2 Heavy Chain
Seq ID No: 78	531-02 CDR3 Heavy Chain
Seq ID No: 79	531-02 VL
Seq ID No: 80	531-02 VH
Ab 548-01	
Seq ID No: 81	548-01 CDR1 Light Chain
Seq ID No: 82	548-01 CDR2 Light Chain
Seq ID No: 83	548-01 CDR3 Light Chain
Seq ID No: 84	548-01 CDR1 Heavy Chain
Seq ID No: 85	548-01 CDR2 Heavy Chain
Seq ID No: 86	548-01 CDR3 Heavy Chain
Seq ID No: 87	548-01 VL
Seq ID No: 88	548-01 VH
Ab 548-02	
Seq ID No: 89	548-02 CDR1 Light Chain
Seq ID No: 90	548-02 CDR2 Light Chain
Seq ID No: 91	548-02 CDR3 Light Chain
Seq ID No: 92	548-02 CDR1 Heavy Chain
Seq ID No: 93	548-02 CDR2 Heavy Chain
Seq ID No: 94	548-02 CDR3 Heavy Chain
Seq ID No: 95	548-02 VL
Seq ID No: 96	548-02 VH
Ab1289-02	
Seq ID No: 97	1289-02 CDR1 Light Chain
Seq ID No: 98	1289-02 CDR2 Light Chain
Seq ID No: 99	1289-02 CDR3 Light Chain
Seq ID No: 100	1289-02 CDR1 Heavy Chain
Seq ID No: 101	1289-02 CDR2 Heavy Chain
Seq ID No: 102	1289-02 CDR3 Heavy Chain
Seq ID No: 103	1289-02 VL
Seq ID No: 104	1289-02 VH
Ab 811-02	
Seq ID No: 105	811-02 CDR1 Light Chain
Seq ID No: 106	811-02 CDR2 Light Chain
Seq ID No: 107	811-02 CDR3 Light Chain
Seq ID No: 108	811-02 CDR1 Heavy Chain
Seq ID No: 109	811-02 CDR2 Heavy Chain
Seq ID No: 110	811-02 CDR3 Heavy Chain
Seq ID No: 111	811-02 VL
Seq ID No: 112	811-02 VH
Ab 566-01	
Seq ID No: 113	566-01 CDR1 Light Chain

Seq ID No: 114	566-01 CDR2 Light Chain
Seq ID No: 115	566-01 CDR3 Light Chain
Seq ID No: 116	566-01 CDR1 Heavy Chain
Seq ID No: 117	566-01 CDR2 Heavy Chain
Seq ID No: 118	566-01 CDR3 Heavy Chain
Seq ID No: 119	566-01 VL
Seq ID No: 120	566-01 VH
Ab 562	
Seq ID No: 121	562 CDR1 Light Chain
Seq ID No: 122	562 CDR2 Light Chain
Seq ID No: 123	562 CDR3 Light Chain
Seq ID No: 124	562 CDR1 Heavy Chain
Seq ID No: 125	562 CDR2 Heavy Chain
Seq ID No: 126	562 CDR3 Heavy Chain
Seq ID No: 127	562 VL
Seq ID No: 128	562 VH
Ab 193	
Seq ID No: 129	193 CDR1 Light Chain
Seq ID No: 130	193 CDR2 Light Chain
Seq ID No: 131	193 CDR3 Light Chain
Seq ID No: 132	193 CDR1 Heavy Chain
Seq ID No: 133	193 CDR2 Heavy Chain
Seq ID No: 134	193 CDR3 Heavy Chain
Seq ID No: 135	193 VL
Seq ID No: 136	193 VH
Ab 88	
Seq ID No: 137	88 CDR1 Light Chain
Seq ID No: 138	88 CDR2 Light Chain
Seq ID No: 139	88 CDR3 Light Chain
Seq ID No: 140	88 CDR1 Heavy Chain
Seq ID No: 141	88 CDR2 Heavy Chain
Seq ID No: 142	88 CDR3 Heavy Chain
Seq ID No: 143	88 VL
Seq ID No: 144	88 VH
Ab 045	
Seq ID No: 145	045 CDR1 Light Chain
Seq ID No: 146	045 CDR2 Light Chain
Seq ID No: 147	045 CDR3 Light Chain
Seq ID No: 148	045 CDR1 Heavy Chain
Seq ID No: 149	045 CDR2 Heavy Chain
Seq ID No: 150	045 CDR3 Heavy Chain
Seq ID No: 151	045 VL
Seq ID No: 152	045 VH

Ab 044	
Seq ID No: 153	044 CDR1 Light Chain
Seq ID No: 154	044 CDR2 Light Chain
Seq ID No: 155	044 CDR3 Light Chain
Seq ID No: 156	044 CDR1 Heavy Chain
Seq ID No: 157	044 CDR2 Heavy Chain
Seq ID No: 158	044 CDR3 Heavy Chain
Seq ID No: 159	044 VL
Seq ID No: 160	044 VH
Ab 002	
Seq ID No: 161	002 CDR1 Light Chain
Seq ID No: 162	002 CDR2 Light Chain
Seq ID No: 163	002 CDR3 Light Chain
Seq ID No: 164	002 CDR1 Heavy Chain
Seq ID No: 165	002 CDR2 Heavy Chain
Seq ID No: 166	002 CDR3 Heavy Chain
Seq ID No: 167	002 VL
Seq ID No: 168	002 VH
Seq ID No: 169	Full human Sortilin sequence isoform 1
Seq ID No: 170	"D Region" as identified by present invention
Seq ID No: 171	Sortilin "hSORTECDBAP"
Seq ID No: 172	Sortilin SORTECDBAP_hBACK
Seq ID No: 173	Sortilin SORTECDBAP_tetra
Seq ID No: 174	Sortilin SORTECDBAP_hB01-05
Seq ID No: 175	Sortilin SORTECDBAP_hRIM
Seq ID No: 176	Sortilin SORTECDBAP_hB06-10
Seq ID No: 177	Sortilin SORTECDBAP_hB12390
Seq ID No: 178	Sortilin SORTECDBAP_hB45678
Seq ID No: 179	Sortilin SORTECD_HIS
Seq ID No: 180	"A region" as identified by the present invention
Seq ID No: 181	A region 109-114
Seq ID No: 182	A region 126-153
Seq ID No: 183	A region 126-144
Seq ID No: 184	A region 154-159
Seq ID No: 185	D Region 570-572
Seq ID No: 186	D Region 588-597
Seq ID No: 187	D Region 593-597
Seq ID No: 188	Sequences used for HDX

The listing or discussion in this specification of an apparently prior-published document should not necessarily be taken as an acknowledgement that the document is part of the state of the art or is common general knowledge.

EMBODIMENTS

[00206] As would be apparent from the text and the Examples the invention further relates to the below embodiments:

1. An antibody, or an antigen-binding fragment thereof, capable of specifically binding to Sortilin and inhibiting binding of PGRN to Sortilin.
2. The antibody, or antigen-binding fragment thereof, according to Embodiment 1, wherein the antibody comprises or consists of an intact antibody.
3. The antibody, or antigen-binding fragment thereof, according to Embodiment 1 or 2, wherein the antigen-binding fragment comprises or consists of an antigen-binding fragment selected from the group consisting of: an Fv fragment (e.g. single chain Fv or a disulphide-bonded Fv); a Fab-like fragment (e.g. Fab fragment or F(ab')₂ fragment); and a domain antibody (e.g. a single V_H variable domain or V_L variable domain).
4. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein the antibody is selected from the group consisting of: an antibody of subtype IgG1, IgG2, IgG3 or IgG4.
5. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein said antibody or antigen-binding fragment thereof binds specifically to the D Region of Sortilin as defined in **SEQ ID NO:170**.
6. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein said antibody or fragment thereof binds specifically to at least 3 consecutive amino acids, such as 4, 5, 6 or 7 consecutive amino acids, of the D Region of Sortilin as defined in **SEQ ID NO:170**.

7. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein the antibody or antigen-binding fragment exhibits one or more of the following properties:
- (i) a binding affinity (K_D) for Sortilin of between 0.5-10 nM, such as 1-5 nM or 1-2 nM
 - (ii) capability to reduce and/or inhibit PGRN binding to Sortilin;
 - (iii) capability to reduce and/or inhibit clearance of PGRN by Sortilin-expressing cells;
 - (iv) capability to reduce and/or inhibit the endocytosis of PGRN by Sortilin-expressing cells;
 - (v) capability to increase the amount and/or concentration of PGRN in the plasma in human-Sortilin-expressing knock-in mice.
8. The antibody, or antigen-binding fragment thereof, according to Embodiment 7, wherein the capability of the antibody or fragment thereof to reduce PGRN binding to Sortilin comprises reducing PGRN binding to Sortilin by 10% or more; for example, by 20% or more; or by 30% or more.
9. The antibody, or antigen-binding fragment thereof, according to Embodiment 7 or 8, wherein the capability of said antibody or fragment thereof the antibody or fragment thereof to reduce and/or inhibit PGRN binding to Sortilin comprises reducing and/or inhibiting PGRN binding to Sortilin with an IC_{50} at or below 22 nM, such as between 22 nM and 1 nM, or between 10 nM and 1 nM, or between 5 nM and 1 nM.
10. The antibody, or antigen-binding fragment thereof, according to any previous Embodiment, wherein the antibody or antigen-binding fragment thereof is human or is humanized.
11. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, comprising a light chain variable domain comprising one or more of the CDR 1-3 Light Chain as listed for each of the antibodies defined in Table 5, or an amino acid sequence having no

more than 4 amino acid differences, or no more than 3 amino acid differences, or no more than 2 amino acid differences, or no more than 1 amino acid difference.

- 5 12. The antibody, or antigen-binding fragment thereof, according to Embodiment 11, comprising a light chain variable domain comprising the CDR 1-3 Light Chain as listed for each of the antibodies defined in Table 5.
- 10 13. The antibody, or antigen-binding fragment thereof, according to Embodiment 11 or 12, comprising a light chain variable domain comprising or consisting of the amino acid sequence VL as listed for each of the antibodies defined in Table 5.
- 15 14. The antibody, or antigen-binding fragment thereof, according to any of Embodiments 11 to 13, comprising a light chain comprising or consisting of the amino acid sequence of VL as listed for each of the antibodies defined in Table 5.
- 20 15. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, comprising a heavy chain variable domain comprising one or more CDR 1-3 Heavy Chain as listed for each of the antibodies defined in Table 5, or an amino acid sequence having no more than 4 amino acid differences, or no more than 3 amino acid differences, or no more than 2 amino acid differences, or no more than 1 amino acid difference.
- 25 16. The antibody, or antigen-binding fragment thereof, according to Embodiment 15, comprising a heavy chain variable domain comprising the CDR 1-3 Heavy Chain as listed for each of the antibodies defined in Table 5.
17. An antibody, or antigen-binding fragment thereof, according to Embodiment 15 or 16 comprising a heavy chain variable domain

comprising or consisting of the amino acid sequence of VH as listed for each of the antibodies defined in Table 5.

18. The antibody, or antigen-binding fragment thereof, according to any of Embodiments 15 to 17, comprising a heavy chain comprising or consisting of the amino acid sequence VL as listed for each of the antibodies defined in Table 5.
19. The antibody, or antigen-binding fragment thereof, according to any preceding embodiment, comprising a light chain variable domain comprising or consisting of the amino acid sequence of VL as listed for each of the antibodies defined in Table 5, and a heavy chain variable domain comprising or consisting of the amino acid sequence of VH as listed for each of the antibodies defined in Table 5.
20. The antibody, or antigen-binding fragment thereof, according to any preceding embodiment, comprising a light chain comprising or consisting of the amino acid sequence of VL as listed for each of the antibodies defined in Table 5, and a heavy chain comprising or consisting of the amino acid sequence of VH as listed for each of the antibodies defined in Table 5.
21. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein said antibody or antigen-binding fragment thereof competes with the antibody or antigen-binding fragment thereof defined in Embodiment 20 for binding to Sortilin.
22. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein the antibody or antigen-binding fragment comprises an Fc region.
23. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein the antibody or antigen-binding fragment further comprises a moiety for increasing *in vivo* half-life.

24. The antibody, or antigen-binding fragment thereof, according to Embodiment 22, wherein the moiety for increasing the *in vivo* half-life is selected from the group consisting of polyethylene glycol (PEG), human serum albumin, glycosylation groups, fatty acids and dextran.
- 5 25. The antibody, or antigen-binding fragment thereof, according to any preceding Embodiment, wherein the antibody or antigen-binding fragment further comprises a detectable moiety.
26. The antibody, or antigen-binding fragment thereof, according to Embodiment 25, wherein the detectable moiety is selected from the group consisting of: a fluorescent label; a chemiluminescent label; a
10 paramagnetic label; a radio-isotopic label; or an enzyme label.
27. The antibody, or antigen-binding fragment thereof, according to Embodiment 25 or 26, wherein the detectable moiety comprises or consists of a radioisotope.
- 15 28. The antibody, or antigen-binding fragment thereof, according to Embodiment 26 or 27, wherein the radioisotope is selected from the group consisting of ^{99m}Tc , ^{111}In , ^{67}Ga , ^{68}Ga , ^{72}As , ^{89}Zr , ^{123}I and ^{201}Tl .
29. The antibody, or antigen-binding fragment thereof, according to Embodiment 25, wherein the detectable moiety comprises or consists of
20 a paramagnetic isotope.
30. The antibody, or antigen-binding fragment thereof, according to Embodiment 29 wherein the paramagnetic isotope is selected from the group consisting of ^{157}Gd , ^{55}Mn , ^{162}Dy , ^{52}Cr and ^{56}Fe .
31. The antibody, or antigen-binding fragment thereof, according to any of
25 Embodiments 25 to 30, wherein the detectable moiety is detectable by an imaging technique such as SPECT, PET, MRI, optical or ultrasound imaging.

32. The antibody, or antigen-binding fragment thereof, according to any of Embodiments 25 to 31, wherein the detectable moiety is joined to the antibody or antigen-binding fragment thereof indirectly, via a linking moiety.
- 5 33. The antibody, or antigen-binding fragment thereof, according to Embodiment 32 wherein the linking moiety is selected from the group consisting of: derivatives of 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA); deferoxamine (DFO); derivatives of diethylenetriaminepentaacetic acid (DTPA); derivatives of S-2-(4-
10 Isothiocyanatobenzyl)-1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA); and derivatives of 1,4,8,11-tetraazacyclododecane-1,4,8,11-tetraacetic acid (TETA).
34. An isolated nucleic acid molecule encoding an antibody or antigen-binding fragment thereof as defined in any of Embodiments 1-33.
- 15 35. A nucleic acid molecule according to Embodiment 34 wherein the molecule is a cDNA molecule.
36. A vector comprising a nucleic acid molecule as defined in Embodiment 34 or 35.
37. A recombinant host cell comprising a nucleic acid molecule as defined in
20 any of Embodiments 34-36.
38. A method for producing an antibody or antigen-binding fragment as defined in any of Embodiments 1-33, the method comprising culturing a host cell as defined in Embodiment 37 under conditions which permit expression of the encoded antibody or antigen-binding fragment thereof.
- 25 39. A preparation comprising the antibody or antigen-binding fragment thereof according to any one of the previous Embodiments, wherein said preparation is substantially free of naturally-arising antibodies that are either not capable of binding to Sortilin or that do not materially alter an an-

ti-Sortilin functionality of the preparation, said functionality being selected from the group consisting of:

- (i) a binding affinity (K_D) for Sortilin;
 - (ii) a capability to reduce and/or inhibit PGRN binding to Sortilin;
 - 5 (iii) a capability to reduce and/or inhibit clearance of PGRN by Sortilin-expressing cells;
 - (iv) a capability to reduce and/or inhibit the endocytosis of PGRN by Sortilin-expressing cells;
 - (v) a capability to increase the amount and/or concentration of PGRN
10 in the plasma in human-Sortilin-expressing knock-in mice; and/or
 - (vi) a capability, when administered chronically, to provide treatment of frontotemporal dementia (FTD) and/or amyotrophic lateral sclerosis (ALS).
40. A preparation comprising the monoclonal antibody or antigen-binding
15 fragment thereof according to any one of the previous Embodiments, wherein said monoclonal antibody possesses a structural change in its amino acid sequence, relative to the structure of a naturally-occurring anti-Sortilin antibody, wherein said structural change causes said monoclonal antibody to exhibit an altered functionality relative to the functionality
20 exhibited by said naturally-occurring anti-Sortilin antibody, wherein said functionality is:
- (i) a binding affinity (K_D) for Sortilin;
 - (ii) a capability to reduce and/or inhibit PGRN binding to Sortilin;
 - (iii) a capability to reduce and/or inhibit clearance of PGRN by Sortilin-
25 expressing cells;
 - (iv) a capability to reduce and/or inhibit the endocytosis of PGRN by Sortilin-expressing cells;
 - (v) a capability to increase the amount and/or concentration of PGRN in the plasma in human-Sortilin-expressing knock-in mice; and/or
 - 30 (vi) a capability, when administered chronically, to provide treatment of frontotemporal dementia (FTD) and/or amyotrophic lateral sclerosis (ALS).

41. A pharmaceutical composition comprising an antibody, or antigen-binding fragment thereof, as defined in any of Embodiments 1-33, or the preparation of any one of embodiments 39-40, and a pharmaceutically-acceptable carrier.
- 5 42. The antibody, or antigen-binding fragment thereof, as defined in any of Embodiments 1-33, or the preparation of any one of embodiments 39-40, for use in medicine.
43. The antibody, or antigen-binding fragment thereof, as defined in any of Embodiments 1-33, or the preparation of any one of embodiments 39-40,
10 for use in preventing and/or treating a disease associated with decreased PGRN levels in the brain of a patient.
44. Use of an antibody, or antigen-binding fragment thereof, as defined in any of Embodiments 1-33, or the preparation of any one of embodiments 39-40, in the manufacture of a medicament for preventing and/or treating
15 a disease associated with decreased PGRN levels in the brain of a patient.
45. The antibody or antigen-binding fragment thereof for use according to Embodiment 43, or the use according to Embodiment 44, wherein the disease is selected from the group consisting of: FTD; ALS; TDP43
20 proteinopathies, such as AD.
46. A method of preventing or treating a disease associated with decreased PGRN levels in the brain of a patient, comprising administering an effective dosage of an antibody or a fragment thereof as defined in any of Embodiments 1-33, the preparation of any one of Embodiments 39-40, or
25 the pharmaceutical composition of Embodiment 41.
47. The antibody, or antigen-binding fragment thereof, for use according to Embodiment 43, or the use according to Embodiment 44, or the method according to Embodiment 46, wherein the disease is selected from the group consisting of: FTD; ALS; or TDP43 proteinopathies, such as AD.

48. The antibody, or antigen-binding fragment thereof, for use; or the use; or the method according to Embodiment 46 or 47, wherein the treatment is chronic.
49. The antibody, or antigen-binding fragment thereof, for use; or the use; or the method, according to Embodiment 48, wherein the chronic treatment is for at least 2 weeks, such as at least for 1 month, 6, months, 1 year or more
50. The antibody, or antigen-binding fragment thereof, as defined in any of Embodiments 1-33, the preparation of any one of Embodiments 39-40, or the pharmaceutical composition of Embodiment 41, which is capable of specifically binding to Sortilin and inhibiting the binding of PGRN to Sortilin, but which binding does not inhibit or substantially inhibit the binding of neurotensin or AF38469 to Sortilin.
51. A kit comprising the antibody, or antigen-binding fragment thereof, as defined in any of Embodiments 1-33, the preparation, as defined in any one of Embodiments 39-40, or the pharmaceutical composition as defined in Embodiment 41.

[00207] Preferred, non-limiting examples which embody certain aspects of the invention will now be described, with reference to the accompanying figures.

EXAMPLES

Examples 1-3 describe the generation of sortilin constructs

[00208] **Example 1** discloses the shuffle constructs. **Example 2** discloses the
5 expression of sortilin constructs. **Example 3** discloses the purification of sortilin
constructs.

Examples 4-7 describe the generation of sortilin antibodies

[00209] **Example 4** discloses the immunization and the hybridomas.
Example 5 discloses the sequence analysis. **Example 6** discloses the
10 purification of antibodies. **Example 7** discloses the generation of mouse
antibodies.

Examples 8-17 describe the characterization of sortilin antibodies

[00210] **Example 8** discloses the binding to sortilin. **Example 9** discloses the
cross blocking ability of Sortilin antibodies. **Example 10** discloses HTRF PGRN-
15 sortilin binding. **Example 11** discloses NTS binding. **Example 12** discloses
cellular PGRN binding and endocytosis. **Example 13** discloses extracellular
PGRN levels. **Example 14** discloses iPSC PGRN levels. **Example 15** discloses
a plasma PGRN levels. **Example 16** disclose epitope mapping by HDX.
Example 17 disclose microdialysis of PGRN in the brain.

20 **Example 1**

[00211] For use in both the hybridoma screening process and as a
diversification of the panel of antibodies, so called 'shuffle constructs" were
designed, constructed and produced, making a set of chimeric sortilin
molecules containing amino acid sequences derived from both human sortilin
25 and a distantly related species (tetraodon) with significantly reduced sequence
homology. The rationale being that the overall sortilin structure and functionality
of these chimeric constructs would be retained but that loss of binding of
antibodies to certain chimeric constructs would indicate the involvement of the

specific exchanged regions in binding. Soluble extracellular region (ECD, aa 1-755) constructs were tagged with either a BAP tag (biotin acceptor peptide), enabling the "in vitro" biotinylation of the proteins by co-expression of biotin ligase or a His tag, enabling easy purification. Expression vectors encoding the following proteins were prepared: SORT-ECDBAP, SORT-ECDBAP-hB01-05, SORT-ECDBAP-hB06-10, SORT-ECDBAP-hB12390, SORT-ECDBAP-hB45678, SORT-ECDBAP-tetra, SORT, SORT-tetra.

[00212] The Sortilin sequences can be found in **SEQ ID NOs:169-180** and **Figure 2** shows schematic presentation of the region assignment of antibodies based on binding to Sortilin shuffle constructs.

Example 2

[00213] In the case of antibody expression, the appropriate heavy chain and light chain vectors, as described in Examples 4, 5 and 6, were co-expressed in HEK-293F cells.

15 Example 3: Purification of His-tagged Sortilin

[00214] SORTECDHis was expressed in HEK-293F cells. The His-tag in the proteins enables purification with immobilized metal affinity chromatography. In this process NiNTA Superflow Cartridge (Qiagen) is equilibrated with 50mM NAH_2PO_4 , 300 mM NaCl and 10mM Imidazole pH 8.0. Column is loaded with His tagged protein with a residence time of 1 minute. Column is washed with 50mM NAH_2PO_4 , 300 mM NaCl and 20mM Imidazole pH 8.0. Protein is eluted with 50mM NAH_2PO_4 , 300 mM NaCl and 250mM Imidazole pH 8.0. Subsequently the protein is dialyzed to PBS using a Slide-A-Lyzer with a cut off of 10.000 mwco (Thermo Scientific). After dialyzing the protein is sterile filtered using a 0.2 micron SFCA filter(Thermo Scientific).

[00215] The S18-HEK cell line was generated by transfecting HEK293 cells with a human wild type (WT) sortilin expression vector. Stable transfected cells were derived after passage in the presence of a selection agent. Individual

clones were selected by dilution cloning. Clones were characterized for sortilin mRNA expression using QPCR. Highest expressing clones were then analyzed by FACS (Guava, Millipore) using an anti-sortilin polyclonal antibody (Polyclonal Goat Sortilin Biotinylated Ab, Cat.No: BAF2934_(R&D Systems)) to determine the surface expressed levels of Sortilin.

Example 4

A - Immunization procedure of transgenic mice

[00216] Antibodies HuMab Sortilin were derived from the immunizations of HuMab mouse strains HCo12, HCo17, HCo20, HCo12-BALB/c, HCo17-BALB/c and HCo20-BALB/c (human monoclonal antibody; Medarex Inc., San Jose, CA, USA). These mice are double knock out for the mouse immunoglobulin (Ig) heavy and mouse kappa light chain, which substantially inactivate the expression of antibodies that are completely murine. The various mouse strains were made transgenic by the insertion of human Ig heavy and human Ig kappa light chain loci and differ in the number of human VH (variable domain of heavy chain) and VL (variable domain of light chain) genes. HCo12-BALB/c mice were derived by crossbreeding with KCo5-BALB/c (kappa light chain transgenic) mice.

[00217] 48 mice were immunized alternating intraperitoneally (IP) with 20 µg SORTECDHis (SEQ ID NO: 179) and subcutaneously (SC, at the tail base) with the same protein, with an interval of 14 days. A maximum of eight immunizations were performed, 4 IP and 4 SC.

[00218] In one protocol, the first immunization was performed with SORTECDHis in complete Freund's adjuvant (CFA; Difco Laboratories, Detroit, MI, USA), the following immunizations in incomplete Freund's adjuvant (IFA). A second protocol used SAS as an adjuvant in all immunization steps. When serum titers were found to be sufficient (dilution of serum of 1/50 or lower found positive in antigen specific screening assay on at least two sequential, biweekly,

screening events), mice were additionally boosted twice intravenously (IV) with 10 µg SORTECDHis protein in 100 µL PBS, four and three days before fusion.

B - HuMab hybridoma-generation

[00219] HuMab mice with sufficient antigen-specific titer development as
5 defined above were sacrificed and the spleen and lymph nodes flanking the
abdominal aorta and caval vein were collected. Fusion of splenocytes and
lymph node cells with a mouse myeloma cell line was done by electrofusion
using a CEEF 50 Electrofusion System (Cyto Pulse Sciences, Glen Burnie, MD,
USA), essentially according to the manufacturer's instructions. Fused cells were
10 seeded in fusion medium containing 10% Fetal Clone I Bovine serum (Perbio),
1 mM sodium pyruvate (Cambrex), 0.5 U/mL penicillin, 0.5 U/mL streptomycin
(Cambrex), 50 µM 2-mercaptoethanol (Invitrogen), 600 ng/mL interleukin 6
(IL-6) (Strathmann), 1 x HAT (Sigma) and 0.5 mg/mL kanamycin (Invitrogen) in
HyQ mADCF-Mab (Perbio). After ten days, supernatant was harvested and
15 cells were refreshed with harvest medium, containing 10 % Fetal Clone I Bovine
serum, 0.5 U/mL penicillin, 0.5 U/mL streptomycin, 600 ng/mL IL-6 and 1 x
proHT (Cambrex) in HyQ mADCF-Mab. Supernatants of the hybridoma cultures
were screened by primary screening assays and streptavidin beads coupled to
SORTECDBAP (SEQ ID NO 171), SORTECDBAPhB06-10 (SEQ ID NO 176),
20 SORTECDBAPhB12390 (SEQ ID NO 177), to detect hybridomas producing
human (or chimeric) anti-Sortilin antibodies. Hybridoma cells from the best
primary wells were seeded in semisolid medium made from 40% CloneMedia
(Genetix, Hampshire, UK) and 60% HyQ 2x complete medium (Hyclone,
Waltham, USA). For each primary well, a well of a Genetix black 6-well plate
25 was seeded. From each well, 25 sub clones were picked, using the ClonePix
system (Genetix). The sub clones were picked in harvest medium. After seven
days, the supernatants of the sub clones were screened again for
Sortilin-specific human IgG binding and the human IgG concentration was
measured using Octet 384red (Fortebio, Menlo Park, USA). From each primary
30 well, the best sub clone was selected and expanded in expansion medium
containing only 600 ng/mL IL-6, 0.5 U/mL penicillin, 0.5 U/mL streptomycin and

1 x proHT. The sub clones were expanded from one 96-well plate well to one 24-well plate well to four 24-well plate wells to six 6-well plate wells. Clones derived by this process were designated as primary clones (PC).

[00220] The anti-sortilin HuMab antibodies of the invention were identified
5 and subjected to sequence analysis.

Example 5: Sequence analysis of the Sortilin-specific HuMab variable domains and cloning in expression vectors

[00221] Total RNA was prepared from 0.2 to 5x10⁶ hybridoma cells and 5'-RACE-Complementary DNA (cDNA) was prepared from 100 ng total RNA,
10 using the SMART RACE cDNA Amplification kit (Clontech), according to the manufacturer's instructions. VH and VL coding regions were amplified by PCR and cloned directly, in frame, in the p33G1f and p33Kappa expression vectors (containing the human IgG1./ kappa constant domain encoding sequences), by ligation independent cloning (Aslanidis, C. and P.J. de Jong, Nucleic Acids Res
15 1990;18(20): 6069-74). For each antibody, 16 VL clones and 16 VH clones were sequenced. Clones with a correct Open Reading Frame (ORF) were selected for further study and expression. Vectors of all combinations of heavy chains and light chains were transiently co-expressed in Freestyle™ 293-F cells using 293fectin.

20 [00222] The resulting sequences are shown in the Sequence Listing (**SEQ ID NOs:1-168**) herein. CDR sequences were defined according to the published guidelines.

Example 6: Purification of antibodies

[00223] Culture supernatant was filtered over 0.2 µm dead-end filters, loaded
25 on 5 mL Protein A columns (rProtein A FF, Amersham Bioscience) and eluted with 0.1 M citric acid-NaOH, pH 3. The eluate was immediately neutralized with 2M Tris-HCl, pH 9 and dialyzed to 12.6 mM NaH₂PO₄, 140 mM NaCl, pH 7.4 (B.Braun), O/N (over night). After dialysis, samples were sterile-filtered over 0.2 µm dead-end filters. Purity was determined by SDS-PAGE and concentration

was measured by nephelometry and absorbance at 280 nm. Purified antibodies were aliquoted and stored at -80°C. Once thawed, purified antibody aliquots were kept at 4°C. Mass spectrometry was performed to identify the molecular mass of the antibody heavy and light chains expressed by the hybridomas.

5 **Example 7: Generation of mouse antibodies (1F2 and 5E1)**

Immunogen

[00224] A synthetic gene coding for the chimeric immunogen hSortilin-FC, (human Sortilin AA (78-756) from **SEQ ID NO:169**) and human IgG1-FC AA (104-330) from **SEQ ID NO:169** was cloned into pcDNA3.1 and used for
10 expression using the freestyle system from Invitrogen. The antigen was purified from cell culture supernatants by protein-A affinity chromatography using standard procedures for antibody purification as described above for human antibodies.

Hybridoma generation

15 [00225] hSortilin-FC was used as immunogen and 5 BALB/c mice were immunized. A mouse with satisfactory immune response was selected for cell fusion and hybridoma generation. Hybridoma supernatants were screened by ELISA using hSortilin-ECD as coating antigen. A total of eighteen hybridoma cell lines derived from nine parental clones were generated.

20 **Expression**

[00226] Hybridomas were initially grown in complete growth medium, DMEM with 10%FBS+antibiotics, and subsequently adapted to CDhybridoma media (Invitrogen) for expression experiments.

Purification

25 [00227] Mouse monoclonal antibodies were purified from hybridoma cell culture supernatants by protein-G sepharose according to standard procedures recommended by the supplier (GE healthcare).

Example 8: Affinity of Sortilin specific HuMab and mouse antibodies to recombinant extracellular region of Sortilin

[00228] Binding kinetics of anti-Sortilin HuMab antibodies to Sortilin were determined using Octet 384RED (ForteBio, Menlo Park, USA). HuMab solutions of 2 µg/ml were made by dilution in sample diluent (ForteBio, art. No. 18-5028).
5 Prot A sensors (ForteBio, art.no. 18-0004) were prewetted with kinetics buffer (1:10 sample diluent in PBS) for at least 600 seconds. Subsequently sensors were immobilized with HuMab solution for 600 seconds. A baseline response was obtained by dipping in kinetics buffer for 120 seconds. Association of
10 SORTECD constructs was performed during a 1000 seconds incubation. This was followed by dissociation in kinetics buffer for 100 seconds. After dissociation, sensors were regenerated (10 mM Glycine pH 1.0) and neutralized (kinetics buffer) 3 times for 5 seconds. All HuMab were analysed using four concentrations of SORTECD constructs (10, 5, 2.5 and 1.25 µg/ml). A
15 molecular weight of 76.8 kDa was used for SORTECDHis. Data was fitted with ForteBio Analysis 6.4 software, using a global full fit. Results are shown in **Figure 3** and **Figure 4**.

Example 9: Antibody cross block of anti-Sortilin HuMabs

[00229] Antibody cross-block studies were performed using Octet 384RED (ForteBio, Menlo Park, USA). HuMab antibody solutions of 2 µg/ml were made by dilution in sample diluent (ForteBio, art. No. 18-5028). Amine reactive
20 sensors (ForteBio, art.no. 18-0008) were used for immobilization of HuMabs. Prior to coupling to amine reactive sensors, HuMabs were diluted in MES pH 6.0 buffer (18-5027). Coupling was performed at 30°C and 1000 rpm as follows:
25 Amine reactive sensors were prewet in PBS and subsequently activated with EDC/NHS (ForteBio. Art.no. 18-1033/18-1034) activation solution (according to manufacturer's instruction) for 300 seconds. Activated sensors were immobilized with HuMabs during 600 seconds. Immobilized sensors were quenched for remaining amine reactivity with Ethanolamine (ForteBio, cat no.
30 18-1039). After quenching sensors were placed in PBS until use. Cross block analysis starts with establishing a baseline response at 30°C and 1000 rpm.

Baseline response was obtained by dipping in sample diluent for 120 seconds. Association of SORTECDHis was performed during 300 seconds directly followed by association of HuMab for 300 seconds. After association of HuMab, sensors were regenerated (10 mM Glycine pH 1.0) and neutralized (sample
5 diluent) 3 times for 5 seconds. Data was processed using ForteBio Analysis 6.4 software.

[00230] Antibodies were grouped based on their binding profiles on the different Sortilin shuffle constructs (**Figure 2**, **Figure 3** and **Figure 4**). To confirm that all the antibodies from Region D (and region F) bind to the same
10 region on human wild type Sortilin ECD, their ability to block each other's binding to the wild type human Sortilin ECD was characterised in a cross blocking study using the Octet384 red. For example, when antibodies from the same region were tested, the primary antibody would block binding of the secondary antibody and vice versa. Whereas, when antibodies from different
15 regions were tested, there would be no cross blocking as only one region is blocked by the primary antibody and the remaining regions are available for the secondary antibody to bind. **Figure 7** shows that all D-region and D+ antibodies cross block each other which confirms classification of the antibodies to Region D and D+ based on shuffle constructs. Further, these data also confirm that the
20 chimeric Sortilin constructs retain similarity to the native human wild type Sortilin ECD.

Example 10: Characterization of sortilin - PGRN ligand binding in the presence of anti-sortilin antibodies.

[00231] IC₅₀ values for antibodies were determined by measuring the
25 displacement of PGRN binding to Sortilin using a homogenous time resolved fluorescent (HTRF, CisBio) assay, see **Figure 5** and **Figure 6**.

[00232] Experiments were performed in assay buffer (50mM Phosphate, pH 7.0, 0.1% BSA) in a total volume of 20µl in a Greiner 384 well, white, low volume microtiter plate, (784075, Greiner).

[00233] The antibodies were pre-incubated for 15min at room temperature with 50nM HIS-tagged Sortilin ECD and 4nM PGRN (SULU20110924) before 7nM anti-6HIS-d2 and 0.7nM anti-PGRN-Eu cryptate (Cisbio) diluted in conjugate buffer (50mM Phosphate, pH 7.0, 0.8mM KF, 0.1% BSA) were added. 20µM neurotensin was used as positive control and DMSO in buffer was used as negative control.

[00234] The assay plate was incubated for 60min at room temperature and overnight at 4°C before the plate was read in EnVision reader (Perkin Elmer).

[00235] Unlabelled neurotensin and DMSO blank were used as positive and negative controls for the assay set up, respectively. Dose-response evaluation of antibodies was performed with ten concentrations between 1µM and 50pM in a 3-fold dilution curve.

[00236] The half-maximal inhibitory concentration (IC₅₀) were calculated by non-linear regression using a sigmoidal concentration response (variable slope) in XLfit 4 (IDBS, UK). (Figures 5 and 6).

Example 11: Characterization of sortilin –neurotensin binding in the presence of anti-sortilin antibodies

[00237] The IC₅₀ for the Sortilin specific compound AF38469 (Schröder et al, Bioorg Med Chem Lett. 2014 Jan 1;24(1):177-80, 2014) was determined by measuring the displacement of ³H-Neurotensin binding to Sortilin using a scintillation proximity assay (SPA).

[00238] Experiments were performed in assay buffer (50mM HEPES, pH 7.4, 100mM NaCl, 2mM CaCl₂, 0.1% BSA, 0.1% Tween-20) in a total volume of 40µl in a 384-well Optiplate, white opaque, (6007299, Perkin Elmer).

[00239] 150nM HIS-tagged Sortilin was pre-incubated for 15min at room temperature with or without 1µM Sortilin specific antibody (IgG1-6003-045 or IgG1-6003-068) or with human IgG1 isotype control before the protein solutions were added to wells containing AF38469 in a concentration series 50µM to 2.5nM. The mixture was incubated for 15min at room temperature on a shaker

before 5nM ^3H -Neurotensin and Ni-chelate imaging beads (RPNQ0266, Perkin Elmer) were added. The assay was incubated additionally for 60min under same conditions.

[00240] After 6h, the plate was read on ViewLux (360s exposure time).

5 Unlabelled neurotensin and DMSO blank were used as positive and negative controls, respectively.

[00241] Dose-response evaluation for AF38469 was performed with ten concentrations between 50 μM and 2.5nM in a 3-fold dilution curve. The half-maximal inhibitory concentration (IC_{50}) were calculated by non-linear regression using sigmoidal concentration response (variable slope) in *XLfit 4* (IDBS, UK). Results can be seen in **Figure 8**.

10

Example 12: Characterization of sortilin - PGRN ligand binding on the surface of cells in the presence of anti-sortilin antibodies.

15 [00242] Both Sortilin transiently transfected cells and the stable cell line S18-HEK cells (human Sortilin over-expressing HEK cells) were used in this assay. Cells were trypsinized and plated at density of 42,000 cells per well in a 96 well plate. In the case of transiently transfected cells, the cells were plated 24 hrs after transfection in 96 well plates. Next day, media was changed completely and test compounds diluted in media were added to cells for 30 min. followed by addition of PGRN for 4hr. At the end of the study (after 4.5 hrs), the cells were fixed and stained for PGRN. All stained plates were analyzed by Cellomics Array Scan (Thermo Fischer) and average staining intensity for PGRN/cell/well was used for analysis.

20

25 [00243] PGRN used in the assay was harvested from media following transient transfection of PGRN expression plasmids in HEK 293 cells. PGRN levels were measured using PGRN ELISA kit (R&D).

[00244] PGRN added to cells was readily bound and endocytosed which led to an increased fluorescence signal in Sortilin transfected wells. Addition of

neurotensin, prevented Sortilin binding to PGRN and a PGRN fluorescence intensity is similar to control levels indicating that PGRN was not bound and endocytosed in the presence of neurotensin.

5 [00245] Both Sortilin HumAbs (45 and 68) blocked the uptake of PGRN with an efficacy similar to neurotensin. The isotype control antibody, B12, did not have any effect on PGRN endocytosis or binding. Results can be seen in **Figure 9**.

Example 13: Effect of Antibodies on extracellular PGRN levels

10 [00246] Both HEK293 cells and S18-HEK cells were found to secrete PGRN continuously into the medium without any stimulation.

[00247] Antibodies and control agents were added to S18-HEK cells to evaluate the effect on PGRN. Addition of Neurotensin, a known peptide sortilin ligand, or human antibodies, 45, 68 and 811 to S18-HEK cells led to an increase of PGRN in the cell culture medium. Two of the Sortilin human
15 antibodies (45 and 68) had an effect similar to neurotensin elevating PGRN levels to 202% and 201 % respectively. The antibody 811 increased PGRN to 146% in the medium as compared to the controls B12, an isotype control antibody was used as negative control in all our studies and did not show any effect on PGRN levels. These observations indicate that the tested sortilin
20 antibodies inhibited sortilin-mediated internalization of PGRN, thereby increasing extracellular PGRN.

[00248] On day1, S18-HEK cells were seeded in a 96 well plate. After 24hrs, medium was completely replaced with either medium alone (control) or medium supplemented with test compound. All compounds were tested at 10uM and
25 antibodies at 100nM unless otherwise specified. Medium was collected on day 3 and analyzed using PGRN ELISA (R&D). Cell viability was assessed by Cell TiterGlo (Pro Mega) to evaluate cytotoxic effect of the compounds. PGRN levels in media were analysed by ELISA and values were normalized to control wells. Results can be seen in **Figure 10**.

Example 14: ELISA assay for extracellular PGRN in iPSCs

[00249] Induced pluripotent stem cells (iPSCs) were generated by non-integrative reprogramming of human fibroblasts (Normal human dermal fibroblasts 18y male; Lonza) as described elsewhere (Rasmussen et al., Stem Cell Reports. 2014 Sep 9;3(3):404-13.). The NHDF K1_shp53 line was used for these studies. The iPSCs were initially generated in mTESR media and subsequently cultured in monolayer in Pluripro (Cell Guidance System). Neuronal differentiation was initiated day 0 by replating the cells on poly-L-ornithine/laminin coated dishes and culturing them in N3 media (50% DMEM/F12 + 50% Neurobasal media supplemented with 0.5% N2, 1% B27 with RA, 0,5 mM GlutaMAX, 0,5% NEA, 50 µM 2-mercaptoethanol and 2.5 mg/mL insulin) with 500 ng/mL noggin and 10 µM SB431542. The media was refreshed every day. After 11 days of noggin/SB431542 induction, the cells were split with dispase and re-plated on poly-L-ornithine/laminin in N3 media. From that point forward, N3 media was refreshed every 2-3 days and the cells were split approximately every 10-14 day using accutase.

[00250] Neuronal differentiated iPSC cells were plated into 96 wells plate. One week later, antibodies were added to the cells. Media from the cells were collected at 48hrs or 96 hrs and analysed by human PGRN ELISA (Enzo Life sciences) and samples analysed as per the manufacturer's instructions.

[00251] The tested Sortilin human antibodies (45 and 68) increased PGRN levels at varying levels in the media at 48hrs and 96 hrs. B12 and Anti-Hel are the control isotype antibodies (negative control). Data is presented as mean $\pm$ SD. Data was analyzed by one-way Anova followed by Dunnett's analysis * $p < 0.05$; ** $p < 0.01$. Results can be seen in **Figure 12**.

Example 15

[00252] To analyze the effect of antibodies on PGRN levels in plasma, humanized Sortilin KI mice were given a single or multiple injections (10mg/kg) of the sortilin antibodies or isotype control by subcutaneous injections. The

animals were anaesthetized and sacrificed at various time points after dosing and plasma PGRN levels determined by ELISA.

- 5 A. Time course study: Mice were treated with antibodies (a Sortilin humab or a control ab) and sacrificed at different time points. Mice treated with control antibodies (Anti-Hel), did not show change in plasma PGRN whereas in mice treated with Sortilin humab 45, there was an increase in PGRN levels which seemed to peak between 24 and 48hrs and then gradually decreased from around day 4. PGRN levels were still elevated at day 7.
- 10 B. Subchronic study: Based on the data from time course study, Sortilin KI mice were dosed twice a week with 10mg/kg, s.c, with either Sortilin human antibody 45 or isotype control antibody, in order to maintain a steady antibody level for a subchronic study (4 weeks). Blood samples were collected at the start of the study and every week to follow plasma PGRN changes. Plasma PGRN levels at the start of the study were similar in both group of animals. Higher levels of plasma PGRN were seen in mice treated with Sortilin antibody45 from week 1 and remain elevated throughout the study. Mice treated with control ab, did not show any increase in plasma PGRN and remained at baseline levels (week 0).
- 15 C. Dose response study: Different doses (4 doses: 10, 2, 0.4 and 0.1 mg/kg) of the Sortilin (45) and control antibody (Anti-Hel) were injected and mice sacrificed on day 2. Plasma PGRN was elevated with 10mg/kg and 2mg/kg in mice treated with Sortilin humab and the lower doses did not have an effect on the plasma PGRN which clearly shows a dose dependent effect of the Sortilin antibody on the plasma PGRN levels.
- 20 Mice treated with control antibody did not show any change in PGRN levels.
- 25

[00253] Mice were anaesthetized with 0,4 ml Avertin IP and heart blood was collected and transferred to a 500 ul kEDTA vial. Samples were kept on ice until centrifuged at 3600G for 15 min at 4C. The plasma was pipetted in to a micronic vial and frozen at -20C. PGRN in the samples was measured using PGRN

30

ELISA kit (Adipogen) as per the manufacturer's instructions. Results can be seen in **Figure 13**.

Example 16: Epitope mapping of antibodies targeting the progranulin-sortilin interaction by hydrogen/deuterium exchange followed by mass spectrometry

[00254] In hydrogen/deuterium exchange followed by mass spectrometry (HDX-MS) the exchange rate of backbone amide hydrogens in a protein is measured. Hereby, it is possible to probe the conformational dynamics of the entire protein backbone except at proline residues. The rate of the exchange reaction is determined by the hydrogen bonding status of the backbone amide and to a lesser extent its solvent accessibility. Subtle changes in these two parameters e.g. caused by the presence of a ligand can be observed as a change in deuterium incorporation.

[00255] To sub-localize the changes in deuterium incorporation the protein is treated with an acid stable protease (e.g. pepsin), which generates local regions of typically ten to fifteen amino acids. Regions that shows a perturbation in the presence of a ligand is either directly involved in the binding interface or allosterically affected by the binding event.

Epitope mapping of antibodies

[00256] The deuterium incorporation of the extra cellular region of Sortilin (**SEQ ID NO:188**) was measured in the absence and presence of mAb45, mAb68, mAb811 and an antibody denominated mAb30 which does not bind the D region. To secure that the measurements were conducted at steady-state conditions the complexes were equilibrated for 15min at 25°C before the exchange reaction was initiated. The exchange reaction was initiated by dilution of the protein samples 1:9 (v/v) into deuterated buffer (99% D₂O, 20mM tris, 150mM NaCl, pDread = 7.6). After various time points (15s, 1min, 10min, 1h and 8h) the exchange reaction was quenched by 1:1 (v/v) dilution with ice-cold quench buffer (2M glycine, 0.8M tris-(2-carboxyethyl)phosphine (TCEP), pH = 2.3), thereby decreasing the pH to 2.46. The quenched samples were

immediately placed inside a -80° freezer and stored until analysis. Fully deuterated control samples were prepared by diluting sortilin samples 1:9 (v/v) into a deuterated denaturation buffer (6M guanidinium chloride, 99% D₂O, 20mM tris, 150mM NaCl, pDread = 7.6) followed by incubation at 25°C for 16h
5 before they were quenched and handled as described above.

[00257] The quenched samples were thawed and injected into a cooled (0°C) reverse-phase UPLC-HDX-system (Waters Inc., USA) equipped with a home-packed pepsin column (internal volume of 60µL, pepsin beads acquired from Thermo Scientific Inc.). Here, the deuterated protein samples were subjected to
10 online pepsin digestion at 20°C, and the resulting peptic peptides were separated by reverse-phase UPLC. The peptides were ionized by electrospray ionization into a mass spectrometer (Synapt G2 mass spectrometer, Waters Inc, UK), where the peptides were further separated by ion mobility before final mass determination.

15 [00258] The Identification of peptides was performed on fully reduced and non-deuterated samples by tandem mass spectrometry using a combination of data independent (MSe) and data dependent acquisition.

Data analysis

Identification of peptides

20 [00259] The acquired mass spectra were lock mass corrected against GFP and analyzed in PLGS 3.0, which matched precursor and fragment ions to a local protein database. All peptide identifications were carefully assessed manually.

[00260] Determination of deuterium incorporation: The acquired mass
25 spectra were lock mass corrected against GFP and the software DynamX 3.0 (Waters Inc., USA) was used to determine the deuterium incorporation for all peptides of sortilin either in absence or presence of antibodies.

[00261] A peptide was considered to be a part of the binding epitope if a protection from exchange larger than 0.5D was observed in presence of an antibody.

Table 1 – Table of identified conformational epitopes by HDX-MS.

Antibody	Epitope mapping by HDX-MS relative to SEQ ID NO:169				
	109-114	126-153		570-572	588-597
45	109-114	126-144	154-159	570-572	593-597
68	109-114	126-144			593-597
811-02	109-114	126-144			593-597

5 **Example 17: Microdialysis to assess progranulin levels in the brain of awake freely moving animals**

[00262] Push-pull microdialysis method was used to assess brain ISF progranulin (PRGN) from awake and freely moving mice. Mice were single-housed in controlled temperature ($22 \pm 1.5^{\circ}\text{C}$) and humidity conditions (55-65%) and kept in a 12:12 hour light/dark cycle (lights on at 06:00h). Food and water were available ad libitum. The current study was performed in the hippocampus of human sortilin knock-in (hSORT1) mice (22 weeks old). To enable microdialysis in the hippocampus, mice were anaesthetized with isoflurane and an intracerebral guide cannula (CMA) was stereotactically implanted into the brain, positioning the microdialysis probe in the hippocampus (co-ordinates of probe tip: 3.1 mm posterior and 2.8 mm lateral from bregma, and 1.3 mm relative dura mater) according to the atlas of Paxinos and Franklin 2001. Acrylic cement was used for the fixation of the guide cannulas. After implantation of the cannula mice were allowed to recover from the surgery for 7 days before dialysis. During the first 5 days, including the surgery day, animals had pain and antibiotics treatments (Rimadyl and Noromox Prolongatum). 24h before the starting of microdialysis experiments pump was also connected to the outlet tubing in order to prevent perfusion fluid loss from the probe, by pulling the fluid through the tubing. As a perfusion buffer, 25% bovine albumin fraction V (Sigma) was diluted to 0.2 % with artificial CSF (aCSF; in mM: 147 NaCl, 2.7 KCl, 1.2 CaCl_2 , 0.85 MgCl_2) on the day of use and filtered through a 0.1- μm

- membrane. The actual flow rate of the pump was determined without having the probe connected. The sample tubes were weighed before and after sampling for a given time period and the flow rate was calculated. The pump was then set to have a constant flow of 1 $\mu\text{L}/\text{min}$. A 120-min sampling regimen was used throughout the experiment period and 12 samples (12h of collection) were collected (**Figure 17**, for procedure). At the end of experiments, blood was taken from animals, animals were perfused and brains collected. The dialysates, plasma and brains were stored at $-80\text{ }^{\circ}\text{C}$ until PRGN determination by ELISA.
- 10 **[00263]** The measurement of PRGN levels every 2h during 24h is depicted in **Figure 17**. At every time period, except 24h after starting collecting dialysates, PRGN levels are significantly increased in animals-treated with mab #45 increased when compared to the ones from animals treated with PBS (**Figure 18a**). PRGN levels are stable over time from 4h until 16h after probe insertion in hippocampus. In the first dialysate PRGN are elevated likely due to the probe insertion into hippocampus. It is speculated that PRGN levels are decreasing $\geq 18\text{h}/20\text{h}$ after probe insertion, likely due to the clogging of the probe membrane, as it occurred in both groups (and has been previously observed in other push-pull studies).
- 20 **[00264]** The average $\pm$ SEM of the 12 dialysis samples 24h after antibody or vehicle treatment, for each animal and then all animals pooled, was taken as baseline (**Figure 18b**). Differences between animals-treated with mab #45 and PBS were analyzed with unpaired t-test. The basal levels of PRGN in animals-treated with mab #45 were significantly increased when compared to the ones from animals treated with PBS ($p<0.001$, $F_{10,0}$, DF_n , 9 DF_d 7; $3.3 \pm 0.3\text{ ng/ml}$, $n=10$ versus $1.1 \pm 0.1\text{ ng/ml}$, $n=8$) (**Figure 18b**).
- 25

CLAIMS

1. An anti-Sortilin antibody, or an antigen-binding fragment thereof, selected from the group consisting of antibodies (1)-(21);

Antibody (1) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:1**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:2**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:3**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:4**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:5**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:6**.

Antibody (2) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:9**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:10**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:11**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:12**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:13**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:14**.

Antibody (3) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:17**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:18**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:19**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:20**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:21**;
and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:22**.

Antibody (4) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:25**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:26**;

- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:27**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:28**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:29**;
- and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:30**.

Antibody (5) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:33**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:34**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:35**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:36**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:37**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:38**.

Antibody (6) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:41**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:42**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:43**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:44**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:45**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:46**.

Antibody (7) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:49**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:50**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:51**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:52**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:53**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:54**.

Antibody (8) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:57**;

- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:58**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:59**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:60**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:61**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:62**.

Antibody (9) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:65**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:66**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:67**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:68**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:69**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:70**.

Antibody (10) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:73**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:74**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:75**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:76**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:77**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:78**.

Antibody (11) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:81**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:82**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:83**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:84**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:85**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:86**.

Antibody (12) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:89**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:90**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:91**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:92**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:93**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:94**.

Antibody (13) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:97**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:98**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:99**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:100**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:101**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:102**.

Antibody (14) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:105**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:106**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:107**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:108**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:109**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:110**.

Antibody (15) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:113**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:114**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:115**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:116**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:117**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:118**.

Antibody (16) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:121**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:122**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:123**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:124**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:125**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:126**.

Antibody (17) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:129**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:130**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:131**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:132**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:133**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:134**.

Antibody (18) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:137**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:138**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:139**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:140**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:141**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:142**.

Antibody (19) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:145**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:146**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:147**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:148**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:149**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:150**.

Antibody (20) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:153**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:154**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:155**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:156**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:157**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:158**.

Antibody (21) is characterized by:

- a. a light chain variable domain L-CDR1 comprising **SEQ ID NO:161**;
- b. a light chain variable domain L-CDR 2 comprising **SEQ ID NO:162**;
- c. a light chain variable domain L-CDR 3 comprising **SEQ ID NO:163**;
- d. a heavy chain variable domain H-CDR 1 comprising **SEQ ID NO:164**;
- e. a heavy chain variable domain H-CDR 2 comprising **SEQ ID NO:165**; and
- f. a heavy chain variable domain H-CDR 3 comprising **SEQ ID NO:166**.

2. An antibody according to claim 1, wherein said antibody is further characterized:

Antibody (1) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:8** and a light chain variable domain comprising **SEQ ID NO:7**.

Antibody (2) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:16** and a light chain variable domain comprising **SEQ ID NO:15**.

Antibody (3) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:24** and a light chain variable domain comprising **SEQ ID NO:23**.

Antibody (4) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:32** and a light chain variable domain comprising **SEQ ID NO:31**.

Antibody (5) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:40 and** a light chain variable domain comprising **SEQ ID NO:39**.

Antibody (6) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:48 and** a light chain variable domain comprising **SEQ ID NO:47**.

Antibody (7) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:56 and** a light chain variable domain comprising **SEQ ID NO:55**.

Antibody (8) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:64 and** a light chain variable domain comprising **SEQ ID NO:63**.

Antibody (9) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:72 and** a light chain variable domain comprising **SEQ ID NO:71**.

Antibody (10) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:80 and** a light chain variable domain comprising **SEQ ID NO:79**.

Antibody (11) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:88 and** a light chain variable domain comprising **SEQ ID NO:87**.

Antibody (12) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:96 and** a light chain variable domain comprising **SEQ ID NO:95**.

Antibody (13) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:104 and** a light chain variable domain comprising **SEQ ID NO:103**.

Antibody (14) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:112 and** a light chain variable domain comprising **SEQ ID NO:111**.

Antibody (15) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:120 and** a light chain variable domain comprising **SEQ ID NO:119**.

Antibody (16) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:128 and** a light chain variable domain comprising **SEQ ID NO:127**.

Antibody (17) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:136 and** a light chain variable domain comprising **SEQ ID NO:135**.

Antibody (18) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:144 and** a light chain variable domain comprising **SEQ ID NO:143**.

Antibody (19) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:152 and** a light chain variable domain comprising **SEQ ID NO:151**.

Antibody (20) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:160 and** a light chain variable domain comprising **SEQ ID NO:159**.

Antibody (21) is further characterized by a heavy chain variable domain comprising **SEQ ID NO:168 and** a light chain variable domain comprising **SEQ ID NO:167**.

3. A pharmaceutical composition comprising an antibody as defined in any of claims 1 or 2 and a pharmaceutically-acceptable carrier.