(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
8 May 2003 (08.05.2003)

PCT

(10) International Publication Number
WO 03/038112 A2

(51) International Patent Classification[7]: C12Q

(21) International Application Number: PCT/US02/34275

(22) International Filing Date: 25 October 2002 (25.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/357,808 26 October 2001 (26.10.2001) US

(71) Applicant: **BAYLOR COLLEGE OF MEDICINE** [US/US]; One Baylor Plaza, Houston, TX 77030 (US).

(72) Inventors: **DRAGHIA-AKLI, Ruxandra**; 5215 Starkridge Drive, Houston, TX 77035 (US). **SCHWARTZ, Robert, J.**; 4019 Marlowe, Houston, TX 77005 (US).

(74) Agents: **CHWANG, Ling, T.** et al.; Jackson Walker L.L.P., 2435 N. Central Expressway, Suite 600, Richardson, TX 75080 (US).


(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A COMPOSITION AND METHOD TO ALTER LEAN BODY MASS AND BONE PROPERTIES IN A SUBJECT



(57) Abstract: The present invention pertains to a method for decreasing the body fat proportion, increasing lean body mass ("LBM"), increasing bone density, or improving the rate of bone healing, or all, of a subject. Overall, the embodiments of the invention can be accomplished by delivering a heterologous nucleic acid sequence encoding GHRH or functional biological equivalent thereof into the cells of the subject and allowing expression of the encoded gene to occur while the modified cells are within the subject. For instance, when such a nucleic acid sequence is delivered into the specific cells of the subject tissue specific constitutive expression is achieved. Furthermore, external regulation of the GHRH or functional biological equivalent thereof gene can be accomplished by utilizing inducible promoters that are regulated by molecular switch molecules, which are given to the subject. The preferred method to deliver the constitutive or inducible nucleic acid encoding sequences of GHRH or the functional biological equivalents thereof is directly into the cells of the subject by the process of *in vivo* electroporation. A decrease the body fat proportion and an increase in lean body mass ("LBM"), or both of a subject is achieved by the delivery of GHRH or functional biological equivalent thereof as described herein by into the subject as recombinant proteins. In addition, an increase in bone density and improvement in the rate of bone healing is also achieved.

WO 03/038112 A2

A COMPOSITION AND METHOD TO ALTER LEAN BODY MASS AND BONE PROPERTIES IN A SUBJECT

RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Patent Application, Serial Number 60/357,808 entitled "increase Body mass, decrease body fat proportion, increase bone density and improve bone healing rate," filed on 10/26/2001, the entire content of which is hereby incorporated by reference.

BACKGROUND

[0002] The present invention pertains to compositions and methods for plasmid-mediated gene supplementation. The present invention relates to a method of decreasing body fat proportions and increasing lean body mass ("LBM") in an animal subject. Overall, the embodiments of the invention can be accomplished by delivering a nucleic acid expression construct that encodes a GHRH or functional biological equivalent thereof into a tissue of a subject and allowing expression of the encoded gene in the subject. For example, when such a nucleic acid sequence is delivered into the specific cells of the subject tissue specific constitutive expression is achieved. Furthermore, external regulation of the GHRH or functional biological equivalent thereof gene can be accomplished by utilizing inducible promoters that are regulated by molecular switch molecules, which are given to the subject. The preferred method to deliver the constitutive or inducible nucleic acid encoding sequences of GHRH or the functional biological equivalents thereof is directly into the cells of the subject by the process of *in vivo* electroporation. In addition, this invention also relates to a method of increasing bone density and improvising the rate of bone healing in an animal subject. More specifically, this invention pertains to both an *in vivo* and an *ex vivo* method for delivering a heterologous nucleic acid sequence encoding growth hormone releasing hormone "GHRH" or functional biological equivalent thereof into the cells of the subject and allowing expression of the encoded gene to occur while the modified cells are within the subject. Another embodiment of the present invention relates to regulating the expression of the GHRH using a molecular switch (e.g mifepistone).

[0003] Growth Hormone ("GH") and Immune Function: The central role of growth hormone ("GH") in controlling somatic growth in humans and other vertebrates, and the physiologically relevant pathways regulating GH secretion from the pituitary are well known. The GH production pathway is composed of a series of interdependent genes whose

products are required for normal growth. The GH pathway genes include: (1) ligands, such as GH and insulin-like growth factor-I ("IGF-I"); (2) transcription factors such as prophet of pit 1, or prop 1, and pit 1: (3) agonists and antagonists, such as growth hormone releasing hormone ("GHRH") and somatostatin ("SS"), respectively; and (4) receptors, such as GHRH receptor ("GHRH-R") and the GH receptor ("GH-R"). These genes are expressed in different organs and tissues, including the hypothalamus, pituitary, liver, and bone. Effective and regulated expression of the GH pathway is essential for optimal linear growth, as well as homeostasis of carbohydrate, protein, and fat metabolism GH synthesis and secretion from the anterior pituitary is stimulated by GHRH and inhibited by somatostatin, both hypothalamic hormones. GH increases production of IGF-I, primarily in the liver, and other target organs. IGF-I and GH, in turn, feedback on the hypothalamus and pituitary to inhibit GHRH and GH release. GH elicits both direct and indirect actions on peripheral tissues, the indirect effects being mediated mainly by IGF-I.

[0004] The principal feature of GH deficiencies in children is short stature. Similar phenotypes are produced by genetic defects at different points in the GH axis, as well as non-GH-deficient short stature. Non-GH-deficiencies have different etiology, such as: (1) genetic diseases, Turner syndrome, hypochondroplasia; and (2) chronic renal insufficiency. Cases where the GH axis is unaffected (*i.e*, patients have normal hormones, genes and receptors) account for more than 50% of the total cases of growth retardation. In these cases GHRH and GH therapy has been shown to be effective (Gesundheit and Alexander, 1995).

[0005] Reduced GH secretion from the anterior pituitary causes skeletal muscle mass to be lost during aging from 25 years to senescence. The GHRH-GH-IGF-I axis undergoes dramatic changes through aging and in the elderly with decreased GH production rate and GH half-life, decreased IGF-I response to GH and GHRH stimuli leads to loss of skeletal muscle mass (sarcopenia), osteoporosis, and increase in fat and decrease in lean body mass. Previous studies have shown that in elderly the level of GH secretion is significant reduced by 70-80% of teenage level. It has been demonstrated that the development of sarcopenia can be offset by exogenous GH therapy. However, this remains a controversial therapy in the elderly because of its cost and frequent side effects.

[0006] The production of recombinant proteins allows a useful tool for the treatment of these conditions. Although GH replacement therapy is widely used in patients

with growth deficiencies and provides satisfactory growth, and may have positive psychological effects on the children being treated, this therapy has several disadvantages, including an impractical requirement for frequent administration of GH and undesirable secondary effects.

[0007] GH is released in a distinctive pulsatile pattern that has profound importance for its biological activity (Argente et al., 1996). Secretion of GH is stimulated by the natural GH secretagogue, GHRH, and inhibited by somatostatin (SS), and both hypothalamic hormones (Thorner et al., 1990). GH pulses are a result of GHRH secretion that is associated with a diminution or withdrawal of somatostatin secretion. In addition, the pulse generator mechanism is timed by GH-negative feedback. The endogenous rhythm of GH secretion becomes entrained to the imposed rhythm of exogenous GH administration. Effective and regulated expression of the GH and insulin-like growth factor I ("IGF-I") pathway is essential for optimal linear growth, homeostasis of carbohydrate, protein, and fat metabolism, and for providing a positive nitrogen balance (Murray and Shalet, 2000). Numerous studies in humans, sheep or pigs showed that continuous infusion with recombinant GHRH protein restores the normal GH pattern without desensitizing GHRH receptors or depleting GH supplies as this system is capable of feed-back regulation, which is abolished in the GH therapies (Dubreuil et al., 1990a; Vance et al., 1985b; Vance, 1990; Vance et al., 1985a). Thus, GHRH recombinant protein treatment may be more physiologically relevant than GH therapy. However, due to the short half-life of GHRH *in vivo*, frequent (one to three times per day) intravenous, subcutaneous or intranasal (requiring 300-fold higher dose) administrations are necessary (Evans et al., 2001; Thorner et al., 1986). Thus, as a chronic therapy, recombinant GHRH protein administration is not practical.

[0008] Extracranially secreted GHRH, as mature peptide or truncated molecules (as seen with pancreatic islet cell tumors and variously located carcinoids) are often biologically active and can even produce acromegaly (Thorner et al., 1984). Administration of recombinant GHRH to GH-deficient children or adult humans augments IGF-I levels, increases GH secretion proportionally to the GHRH dose, yet still invokes a response to bolus doses of GHRH (Bercu et al., 1997). Thus, GHRH administration represents a more physiological alternative of increasing subnormal GH and IGF-I levels (Corpas et al., 1993).

[0009] Although GHRH protein therapy entrains and stimulates normal cyclical GH secretion with virtually no side effects, the short half-life of GHRH *in vivo* requires frequent (one to three times a day) intravenous, subcutaneous or intranasal (requiring 300-fold higher dose) administration. Thus, as a chronic treatment, GHRH administration is not practical. Extracranially secreted GHRH, as processed protein species GHRH(l-40) hydroxy or GHRH(l-44) amide or even as shorter truncated molecules, are biological active (Thorner et al., 1984). It has been reported that a low level of GHRH (100 pg/ml) in the blood supply stimulates GH secretion (Corpas et al., 1993). Direct plasmid DNA gene transfer is currently the basis of many emerging therapy strategies and thus does not require viral genes or lipid particles (Aihara and Miyazaki, 1998; Muramatsu et al., 1998). Skeletal muscle is target tissue, because muscle fiber has a long life span and can be transduced by circular DNA plasmids that express over months or years in an immunocompetent host (Davis et al., 1993; Tripathy et al., 1996). Previous reports demonstrated that human GHRH cDNA could be delivered to muscle by an injectable myogenic expression vector in mice where it transiently stimulated GH secretion to a modes extent over a period of two weeks (Draghia-Akli et al., 1997).

[0010] Wild type GHRH has a relatively short half-life in the circulatory system, both in humans (Frohman et al., 1984) and in farm animals. After 60 minutes of incubation in plasma 95% of the GHRH(1-44)NH2 is degraded, while incubation of the shorter (1-40)OH form of the hormone, under similar conditions, shows only a 77% degradation of the peptide after 60 minutes of incubation (Frohman et al., 1989). Incorporation of cDNA coding for a particular protease-resistant GHRH analog in a therapeutic nucleic acid vector results in a molecule with a longer half-life in serum, increased potency, and provides greater GH release in plasmid-injected animals (Draghia-Akli et al., 1999), herein incorporated by reference). Mutagenesis *via* amino acid replacement of protease sensitive amino acids prolongs the serum half-life of the GHRH molecule. Furthermore, the enhancement of biological activity of GHRH is achieved by using super-active analogs that may increase its binding affinity to specific receptors (Draghia-Akli et al., 1999).

[0011] Administering novel GHRH analog proteins (U.S. Pat Nos. 5,847,066; 5846,936; 5,792,747; 5,776,901; 5,696,089; 5,486,505; 5,137,872; 5,084,442, 5,036,045; 5,023,322; 4,839,344; 4,410,512, RE33,699) or synthetic or naturally occurring peptide fragments of GHRH (U.S. Pat. Nos. 4,833,166; 4,228,158; 4,228,156; 4,226,857; 4,224,316; 4,223,021; 4,223,020; 4,223, 019) for the purpose of increasing release of growth hormone

have been reported. A GHRH analog containing the following mutations has been reported (U.S. Patent No. 5,846,936): Tyr at position 1 to His; Ala at position 2 to Val, Leu, or others; Asn at position 8 to Gln, Ser, or Thr; Gly at position 15 to Ala or Leu; Met at position 27 to Nle or Leu; and Ser at position 28 to Asn. The GHRH analog is the subject of U.S. Patent Application Serial No. 09/624,268 ("the '268 patent application"), which teaches application of a GHRH analog containing mutations that improve the ability to elicit the release of growth hormone. In addition, the '268 patent application relates to the treatment of growth deficiencies; the improvement of growth performance; the stimulation of production of growth hormone in an animal at a greater level than that associated with normal growth; and the enhancement of growth utilizing the administration of growth hormone releasing hormone analog and is herein incorporated by reference. In the embodiments of the '268 patent application and specific embodiments herein, the mutated GHRH-encoding molecules lack the Gln, Ser or Thr mutations of the Asn at position 8.

[0012] U.S. Patent No. 5,061,690 is directed toward increasing both birth weight and milk production by supplying to pregnant female mammals an effective amount of human GHRH or one of it analogs for 10-20 days. Application of the analogs lasts only throughout the lactation period. However, multiple administrations are presented, and there is no disclosure regarding administration of the growth hormone releasing hormone (or factor) as a DNA molecule, such as with plasmid mediated supplementation techniques.

[0013] U.S. Patent Nos. 5,134,120 (" the '120 patent") and 5,292,721 (" the '721 patent") teach that by deliberately increasing growth hormone in swine during the last 2 weeks of pregnancy through a 3 week lactation resulted in the newborn piglets having marked enhancement of the ability to maintain plasma concentrations of glucose and free fatty acids when fasted after birth. In addition, the '120 and '721 patents teaches that treatment of the sow during lactation results in increased milk fat in the colostrum and an increased milk yield. These effects are important in enhancing survivability of newborn pigs and weight gain prior to weaning. However the '120 and '721 patents provide no teachings regarding administration of the growth hormone releasing hormone as a DNA form.

[0014] In contrast to protein therapy, nucleic acid transfer delivers polynucleotides to somatic tissue in a manner that, in some embodiments, can correct inborn or acquired deficiencies and imbalances. In other embodiments, vectors such as plasmids are

used to supplement basal levels of an expressed endogenous gene product. Gene-based drug delivery offers a number of advantages over the administration of recombinant proteins. These advantages include the conservation of native protein structure, improved biological activity, avoidance of systemic toxicities, and avoidance of infectious and toxic impurities. In addition, nucleic acid transfer, for therapeutic purposes, and plasmid-mediated supplementation of an endogenous gene product allow for prolonged exposure to the protein in the therapeutic range, because the newly secreted protein is present continuously in the blood circulation.

[0015] The primary limitation of using recombinant protein is the limited availability of protein after each administration. Plasmid-mediated gene supplementation using injectable DNA plasmid vectors overcomes this, because a single injection into the patient's skeletal muscle permits physiologic expression for extensive periods of time (WO 99/05300 and WO 01/06988). Injection of the vectors promotes the production of enzymes and hormones in animals in a manner that more closely mimics the natural process. Furthermore, among the non-viral techniques for gene transfer *in vivo*, the direct injection of plasmid DNA into muscle tissue is simple, inexpensive, and safe.

[0016] In a plasmid-mediated supplementation expression system, a non-viral nucleic acid vector, such as a plasmid, may comprise a synthetic nucleic acid delivery system in addition to a nucleic acid encoding the GHRH being supplemented. In this way, the risks associated with the use of most viral vectors can be avoided. The non-viral expression vector products generally have low toxicity due to the use of "species-specific" components for nucleic acid delivery, which minimizes the risks of immunogenicity generally associated with viral vectors. Additionally, no integration of plasmid sequences into host chromosomes has been reported *in vivo* to date, so that this type of plasmid-mediated supplementation of GHRH, should neither activate oncogenes nor inactivate tumor suppressor genes. As episomal systems residing outside the chromosomes, plasmids have defined pharmacokinetics and elimination profiles, leading to a finite duration of gene expression in target tissues. Furthermore, the plasmid DNA could be engineered so it would be delivered to the cells in a linear rather than circular form (which would further prevent any genomic integration event); the plasmid could be deleted of the antibiotic resistance gene and bacterial origin of replication, making it completely safe for *in vivo* therapy.

[0017] Efforts have been made to enhance the delivery of plasmid DNA to cells by physical means including electroporation, sonoporation, and pressure. Injection by electroporation involves the application of a pulsed electric field to create transient pores in the cellular membrane without causing permanent damage to the cell. It thereby allows for the introduction of exogenous molecules (Smith and Nordstrom, 2000). By adjusting the electrical pulse generated by an electroporetic system, nucleic acid molecules can travel through passageways or pores in the cell that are created during the procedure. U.S. Patent No. 5,704,908 describes an electroporation apparatus for delivering molecules to cells at a selected location within a cavity in the body of a patient. These pulse voltage injection devices are also described in U.S. Patent Nos. 5,439,440 and 5,702,304, and PCT WO 96/12520, 96/12006, 95/19805, and 97/07826.

[0018] Recently, significant progress has been obtained using electroporation to enhance plasmid delivery *in vivo*. Electroporation has been used very successfully to transfect tumor cells after injection of plasmid (Lucas et al., 2002; Matsubara et al., 2001) or to deliver the anti-tumor drug bleomycin to cutaneous and subcutaneous tumors in humans (Gehl et al., 1998; Heller et al., 1996). Electroporation also has been extensively used in mice (Lesbordes et al., 2002; Lucas et al., 2001; Vilquin et al., 2001), rats (Terada et al., 2001; Yasui et al., 2001), and dogs (Fewell et al., 2001) to deliver therapeutic genes that encode for a variety of hormones, cytokines or enzymes. Our previous studies using growth hormone releasing hormone ("GHRH") showed that plasmid therapy with electroporation is scalable and represents a promising approach to induce production and regulated secretion of proteins in large animals and humans (Draghia-Akli et al., 1999; Draghia-Akli et al., 2002).

[0019] The ability of electroporation to enhance plasmid uptake into the skeletal muscle has been well documented, as described above. In addition, plasmid formulated with poly-L-glutamate ("PLG") or polyvinylpyrolidone (PVP) has been observed to increase plasmid transfection and consequently expression of the desired transgene. The anionic polymer sodium PLG could enhance plasmid uptake at low plasmid concentrations, while reducing any possible tissue damage caused by the procedure. The ability of electroporation to enhance plasmid uptake into the skeletal muscle has been well documented, as previously described. PLG is a stable compound and resistant to relatively high temperatures (Dolnik et al., 1993). PLG has been previously used to increase stability in vaccine preparations (Matsuo et al., 1994) without increasing their immunogenicity. It also has been used as an anti-toxin

post-antigen inhalation or exposure to ozone (Fryer and Jacoby, 1993). In addition, plasmid formulated with PLG or polyvinylpyrrolidone (PVP) has been observed to increase gene transfection and consequently gene expression to up to 10 fold in the skeletal muscle of mice, rats and dogs (Fewell et al., 2001; Mumper et al., 1998). PLG has been used to increase stability of anti-cancer drugs (Li et al., 2000) and as "glue" to close wounds or to prevent bleeding from tissues during wound and tissue repair (Otani et al., 1996; Otani et al., 1998).

[0020] Although not wanting to be bound by theory, PLG will increase the transfection of the plasmid during the electroporation process, not only by stabilizing the plasmid DNA, and facilitating the intracellular transport through the membrane pores, but also through an active mechanism. For example, positively charged surface proteins on the cells could complex the negatively charged PLG linked to plasmid DNA through protein-protein interactions. When an electric field is applied, the surface proteins reverse direction and actively internalize the DNA molecules, process that substantially increases the transfection efficiency.

[0021] The use of directly injectable DNA plasmid vectors has been limited in the past. The inefficient DNA uptake into muscle fibers after simple direct injection has led to relatively low expression levels (Prentice et al., 1994; Wells et al., 1997) In addition, the duration of the transgene expression has been short (Wolff et al., 1990). The most successful previous clinical applications have been confined to vaccines (Danko and Wolff, 1994; Tsurumi et al., 1996).

[0022] Although there are references in the art directed to electroporation of eukaryotic cells with linear DNA (McNally et al., 1988; Neumann et al., 1982) (Toneguzzo et al., 1988) (Aratani et al., 1992; Nairn et al., 1993; Xie and Tsong, 1993; Yorifuji and Mikawa, 1990), these examples illustrate transfection into cell suspensions, cell cultures, and the like, and the transfected cells are not present in a somatic tissue.

[0023] U.S. Patent No. 4,956,288 is directed to methods for preparing recombinant host cells containing high copy number of a foreign DNA by electroporating a population of cells in the presence of the foreign DNA, culturing the cells, and killing the cells having a low copy number of the foreign DNA.

[0024] U.S. Patent No. 5,874,534 ("the '534 patent") and U.S. Patent No. 5,935,934 ("the '934 patent") describe mutated steroid receptors, methods for their use and a molecular switch for nucleic acid transfer, the entire content of each is hereby incorporated by reference. A molecular switch for regulating expression in nucleic acid transfer and methods of employing the molecular switch in humans, animals, transgenic animals and plants (e.g GeneSwitch®) are described in the '534 patent and the '934 patent. The molecular switch is described as a method for regulating expression of a heterologous nucleic acid cassette for nucleic acid transfer and is comprised of a modified steroid receptor that includes a natural steroid receptor DNA binding domain attached to a modified ligand binding domain. The modified binding domain usually binds only non-natural ligands, anti-hormones or non-native ligands. One skilled in the art readily recognizes natural ligands do not readily bind the modified ligand-binding domain and consequently have very little, if any, influence on the regulation and/or expression of the gene contained in the nucleic acid cassette.

[0025] Thus, the present invention is directed to a novel method of increasing lean body mass, decreasing body fat proportions, increasing bone density, and/or increasing the rate of bone healing in an animal by plasmid-mediated supplementation of GHRH

SUMMARY

[0026] One embodiment of the present invention pertains to a method for decreasing the body fat proportion, increasing lean body mass ("LBM"), increasing bone density, and increasing the rate of bone healing of a subject by utilizing a nucleic acid sequence containing both a constitutive promoter and an encoding sequence for growth hormone releasing hormone ("GHRH") or analog thereof. When this nucleic acid sequence is delivered into the specific cells of the subject (e.g. somatic cells, stem cells, or germ cells), tissue specific constitutive expression of GHRH is achieved. The preferred method to deliver the nucleic acid sequence with the constitutive promoter and the encoding sequence of GHRH or the analog thereof is directly into the cells of the subject by the process of *in vivo* electroporation. Electroporation may involve externally supplied electrodes, or in the case of needles, internally supplied electrodes to aid in the inclusion of desired nucleotide sequences into the cells of a subject while the cells are within a tissue of the subject.

[0027] Another embodiment of the present invention pertains to a method for decreasing the body fat proportion, increasing LBM, increasing bone density, and increasing bone healing rate of a subject by utilizing the ability to regulate the expression of GHRH or analog thereof. Regulation is achieved by delivering into the cells of the subject a first nucleic acid sequence, and a second nucleic acid sequence, followed by a molecular switch; where the first nucleic acid sequence contains an inducible-promoter with a coding region for a growth-hormone-releasing-hormone ("inducible-GHRH") or an analog thereof and the second nucleic acid sequence has a constitutive promoter with a coding region for an inactive regulator protein. By delivering a molecular switch molecule (e.g. mifepistone) into the subject, the inactive regulator protein becomes active and initiates transcription of the inducible-GHRH in the subject. The expression and ensuing release of GHRH or analog thereof by the modified-cells within the subject will decrease the body fat proportion and increase the LBM of the subject in a manner that can be regulated by external molecular switch molecules (e.g. mifepistone). The delivery of the nucleic acid sequences that allow external regulation of GHRH or the analog thereof directly into the cells of the subject can be accomplished by the process of *in vivo* electroporation.

[0028] A further embodiment of the present invention pertains to a method for increasing lean body mass, decreasing body fat proportion, increasing bone density, increasing

the rate of bone healing, or a combination thereof, of a subject by utilizing therapy that introduces specific recombinant GHRH-analog protein into the subject.

According to one aspect of the present invention, there is provided the use of a nucleic acid expression construct that encodes a growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for decreasing body fat proportion, wherein the composition is suitable for delivery into cells of a subject.

According to another aspect of the present invention, there is provided the use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for decreasing body fat proportion in a subject wherein the composition is suitable for delivery into cells of the subject, wherein the first nucleic acid expression construct is suitable for encoding growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof, wherein the second nucleic acid expression construct has an encoding region of a regulator protein and wherein upon delivery of the molecular switch molecule into the subject, the molecular switch molecule is capable of governing activation of the regulator protein.

According to another aspect of the present invention, there is provided the use of a recombinant growth-hormone-releasing-hormone ("GHRH") or a biological functional equivalent thereof in the preparation of a composition for decreasing body fat proportion in a subject, wherein the composition is suitable for delivery into the subject.

According to another aspect of the present invention, there is provided the use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for increasing lean body mass in a subject, wherein the composition is suitable for delivery into cells of the subject.

According to another aspect of the present invention, there is provided the use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for increasing lean body mass in a subject and wherein the composition is suitable for delivery into cells of

the subject wherein the first nucleic acid expression construct is suitable for encoding growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof, wherein the second nucleic acid expression construct has an encoding region of a regulator protein and wherein upon delivery of the molecular switch molecule into the subject, the molecular switch molecule is capable of activating the regulator protein.

According to another aspect of the present invention, there is provided the use of a recombinant growth-hormone-releasing-hormone ("GHRH") or a biological functional equivalent thereof in the preparation of a composition for increasing lean body mass in a subject, wherein the composition is suitable for delivery into the subject.

According to another aspect of the present invention, there is provided the use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for increasing bone density in a subject, wherein the composition is suitable for delivery into cells of the subject.

According to another aspect of the present invention, there is provided the use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for increasing bone density in a subject, and wherein the composition is suitable for delivery into the cells of the subject, wherein the first nucleic acid expression construct is suitable for encoding growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof; and wherein the second nucleic acid expression construct has an encoding region of a regulator protein, wherein upon delivery of the molecular switch molecule into the subject, the molecular switch molecule is capable of activating the regulator protein.

According to another aspect of the present invention, there is provided the use of a recombinant growth-hormone-releasing-hormone ("GHRH") or a biological functional equivalent thereof in the preparation of a composition for increasing bone density in a subject, wherein the composition is suitable for delivery into the subject.

According to another aspect of the present invention, there is provided the use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for increasing a bone healing rate in a subject, wherein the composition is suitable for delivery into cells of the subject.

According to another aspect of the present invention, there is provided the use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for increasing a bone healing rate in a subject, and wherein the composition is suitable for delivery into cells of the subject, wherein the first nucleic acid expression construct is suitable for encoding growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof, wherein the second nucleic acid expression construct has an encoding region of a regulator protein, wherein upon delivery of the molecular switch molecule into the subject, the molecular switch molecule is capable of activating the regulator protein.

According to another aspect of the present invention, there is provided the use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for altering lean body mass in a subject, wherein subsequent expression of the GHRH or biological equivalent is suitable for increasing lean body mass or decreasing body fat proportion, and wherein the composition is suitable for delivery into a tissue of the subject.

According to another aspect of the present invention, there is provided a peptide comprising: SeqID No: 2; SeqID No: 3; or SeqID No: 4.

BRIEF DESCRIPTION OF THE DRAWINGS

[0029] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[0030] FIGURE 1 shows the amino acid sequence of GHRH or functional biological equivalent thereof. All mutant sequences were obtained by site directed mutagenesis of the porcine wild type sequence.

[0031] FIGURE 2 shows the body weight of SCID mice that were injected with 7.5 micrograms of pSP-GHRH mutants.

[0032] FIGURE 3 shows the body composition of SCID mice that were injected with 7.5 micrograms of plasmid expressing either one of the GHRH mutants or a pSP-beta-galactosidase as control.

[0033] FIGURE 4 shows the bone area of SCID mice that were injected with 7.5 micrograms of plasmid expressing either one of the GHRH mutants or a pSP-beta-galactosidase as control.

[0034] FIGURE 5 shows the IGF-I levels of SCID mice that were injected with 7.5 micrograms of plasmid expressing either one of the GHRH mutants or a pSP-beta-galactosidase as control.

[0035] FIGURE 6 shows a schematic of the mifepristone-dependent GHRH/GeneSwitch® system in primary myoblasts *in vitro*. Plasmid structures and schematic for how the GeneSwitch® system works are illustrated. Plasmid p1633 encodes for the GeneSwitch® regulator protein, which is a chimera of yeast GAL4 DNA binding domain ("GAL4"), truncated human progesterone receptor ligand-binding domain ("hPR LBD"), and activation domain from the p65 subunit of human NF-κB ("p65"). The protein is synthesized as an inactive monomer. Binding of mifepristone triggers a conformational change that leads to activation and dimerization. Activated homodimers bind to GAL4 sites in the inducible promoter and stimulate transcription of the GHRH gene.

[0036] FIGURE 7 shows the function of a mifepristone-dependent GHRH/GeneSwitch® system in primary myoblasts *in vitro*. Northern blot analysis of inducible GHRH constructs. Primary chicken myoblast cultures were obtained and transfected as described previously (Bergsma et al., 1986; Draghia-Akli et al., 1997), with 4 micrograms of a 10:1 mixture of inducible GHRH ("pGR1774") and GeneSwitch® plasmids ("pGS1633"). A Muscle specific synthetic promoter (Li et al., 1999) driven construct coding for *E. coli* beta-galactosidase, □ βgal, is used as a negative control. As a positive control, cells were transfected with a constitutively active pSP-GHRH construct (Draghia-Akli et al., 1999). In the figure, Nt = non-transfected cells; β-gal = cells transfected with pSP-β-gal construct; SP-GHRH = cells transfected with a constitutively active GHRH construct; +MFP = mifepristone was added to the culture media; and - MFP = mifepristone was not added to the culture media. Ethidium bromide gels are included as loading controls.

[0037] FIGURE 8 shows that the mifepristone dosing induces serum IGF-I levels in SCID mice that received a single administration of GHRH/GeneSwitch® plasmids. Values are presented as fold activation over control levels. The area under the dark line represents normal variability of IGF-I levels in adult animals. The table contains the p values for the induced peaks. The p values C v. A indicate comparison between animals injected with the β-gal construct versus animals injected with the IS + MFP; C v. B. indicates comparison between animals injected with the IS with and without the MFP.

[0038] FIGURE 9 shows the enhanced weight gain during a chronic 149 day MFP induction. Average weight increased in injected mice upon chronic activation of the GHRH/GeneSwitch® system (* p <0.027).

[0039] FIGURE 10 shows the increase in pituitary weight with a chronic 149 day MFP induction. Pituitary weight / total body weight in +MFP injected animals (* p <0.035).

[0040] FIGURE 11 shows the improved body composition in chronically induced GHRH/GeneSwitch® mice. Body composition measurements were performed either under anesthesia, at day 149 post-injection ("PIXImus") or post-mortem (organ, carcass, body fat, direct dissection of the body). Lean non-bone mass is significantly increased (* p <0.022).

[0041] FIGURE 12 shows the improved fat body mass/ total weight in chronically induced GHRH/GeneSwitch® mice. Fat body mass/ total weight measurements were performed either under anesthesia, at day 149 post-injection ("PIXImus") or post-mortem (organ, carcass, body fat, direct dissection of the body). Fat body mass/ total weight is significantly decreased in induced animals (* $p < 0.05$).

[0042] FIGURE 13 shows the increased bone area in chronically induced GHRH/GeneSwitch® mice. Bone area measurements were performed either under anesthesia, at day 149 post-injection ("PIXImus") or post-mortem (organ, carcass, body fat, direct dissection of the body). Bone area is increased by PIXImus (*$p < 0.0006$).

[0043] FIGURE 14 shows the increased mineral content in chronically induced GHRH/GeneSwitch® mice. Bone mineral content measurements were performed either under anesthesia, at day 149 post-injection ("PIXImus") or post-mortem (organ, carcass, body fat, direct dissection of the body). Bone mineral content is increased in induced animals (* $p < 0.002$).

[0044] FIGURE 15 shows the secreted embryonic alkaline phosphatase ("SEAP") plasma concentration in pigs at 0 to 7 days post-injection. Different needle-type electrodes were compared with calipers electrodes following the plasmid injection into the muscle. In the figure, N6 = Six-needle array electrode: 21 gauge needles, 2 cm length mounted on a 1 cm - diameter array; N3 = three-needle array device: two solid needles, 1 median hypodermic needle, 21 gauge, 2 cm length; and C = caliper electrode: 2 solid square plate electrodes, 1.5 cm. Voltage and number of pulses are also indicated. At 7 days post-injection $p < 0.006$ for N3/ 200V/ 6 pulses and N6/100V/ 6 pulses groups, and $p < 0.0035$ for N6/200V/ 6 pulses group.

[0045] FIGURE 16 shows the body weights of pigs injected at 10 days of age with 3, 1 and 0.1 mg of pSP-HV-GHRH or vehicle. The greatest weight gain was achieved by pigs injected with the lowest dose, with statistically significant differences from the controls at all time points tested ($p < 0.02$). Values are means ± s.e.m.

[0046] FIGURE 17 shows the body weights of pigs injected with 2 milligrams of pSP-HV-GHRH at 0, 7, 14 and 21 days of age. Animal injected at 14 days of age showed

the greatest weight gain, statistically different from the controls at all time points tested ($p < 0.02$). Values are means ± s.e.m.

[0047] FIGURE 18 shows the plasma IGF-I concentration after direct intramuscular injection of the different quantities of pSP-HV-GHRH construct. Values are means ± s.e.m.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0048] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

DEFINITIONS

[0049] The term "a" or "an" as used herein in the specification may mean one or more. As used herein in the claim(s), when used in conjunction with the word "comprising", the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

[0050] The term "any range derivable therein" as used herein means a range selected from the numbers described in the specification, and "any integer derivable therein" means any integer between such a range.

[0051] The term "analog" as used herein includes any mutant of GHRH, or synthetic or naturally occurring peptide fragments of GHRH, such as HV-GHRH (SeqID No: 1), TI-GHRH (SeqID No: 2), TV-GHRH (SeqID No: 3), 15/27/28-GHRH (SeqID No: 4), (1-44)NH_2 (SeqID No: 5) or (1-40)OH (SeqID No: 6) forms, or any shorter form to no less than (1-29) amino acids.

[0052] The term "bone density" as used herein is defined as the density of minerals in the bone as measured by a standard means in the art, such as x-ray, MRI, dual-energy x-ray absorbitometry (DEXA), or any advanced imaging system in the art.

[0053] The term "cassette" as used herein is defined as one or more transgene expression vectors.

[0054] The term "cell-transfecting pulse" as used herein is defined as a transmission of a force which results in transfection of a vector, such as a linear DNA fragment, into a cell. In some embodiments, the force is from electricity, as in electroporation, or the force is from vascular pressure.

[0055] The term "coding region" as used herein refers to any portion of the DNA sequence that is transcribed into messenger RNA (mRNA) and then translated into a sequence of amino acids characteristic of a specific polypeptide.

[0056] The term "delivery" or "delivering" as used herein is defined as a means of introducing a material into a tissue, a subject, a cell or any recipient, by means of chemical or biological process, injection, mixing, electroporation, sonoporation, or combination thereof, either under or without pressure.

[0057] The term "donor-cells" as used herein refers to any cells that have been removed and maintained in a viable state for any period of time outside the donor-subject.

[0058] The term "donor-subject" as used herein refers to any species of the animal kingdom wherein cells have been removed and maintained in a viable state for any period of time outside the subject.

[0059] The term "DNA fragment" or "nucleic acid expression construct" as used herein refers to a substantially double stranded DNA molecule. Although the fragment may be generated by any standard molecular biology means known in the art, in some embodiments the DNA fragment or expression construct is generated by restriction digestion of a parent DNA molecule. The terms "expression vector," "expression cassette," or "expression plasmid" can also be used interchangeably. Although the parent molecule may be any standard molecular biology DNA reagent, in some embodiments the parent DNA molecule is a plasmid.

[0060] The term "electroporation" as used herein refers to a method that utilizes electrical pulses to deliver a nucleic acid sequence into cells.

[0061] The terms "electrical pulse" and "electroporation" as used herein refer to the administration of an electrical current to a tissue or cell for the purpose of taking up a nucleic acid molecule into a cell. A skilled artisan recognizes that these terms are associated with the terms "pulsed electric field" "pulsed current device" and "pulse voltage device." A skilled artisan recognizes that the amount and duration of the electrical pulse is dependent on the tissue, size, and overall health of the recipient subject, and furthermore knows how to determine such parameters empirically.

[0062] The term "encoded GHRH" as used herein is a biologically active polypeptide of GHRH.

[0063] The term "functional biological equivalent" of GHRH as used herein is a polypeptide that has a distinct amino acid sequence from a wild type GHRH polypeptide while simultaneously having similar or improved biological activity when compared to the GHRH polypeptide. The functional biological equivalent may be naturally occurring or it may be modified by an individual. A skilled artisan recognizes that the similar or improved biological activity as used herein refers to facilitating and/or releasing growth hormone or other pituitary hormones. A skilled artisan recognizes that in some embodiments the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity when compared to the GHRH polypeptide. Methods known in the art to engineer such a sequence include site-directed mutagenesis.

[0064] The term "GeneSwitch® " (which is a registered trademark of Valentis, Inc. (Burlingame, CA)) as used herein refers to the technology of mifepristone-inducible heterologous nucleic acid sequences encoding regulator proteins, GHRH, functional biological equivalent or combination thereof. Such a technology is schematically diagramed in FIGURE 1A. A skilled artisan recognizes that antiprogesterone agent alternatives to mifepristone are available, including onapristone, ZK112993, ZK98734, and 5α pregnane-3,2-dione.

[0065] The term "growth hormone" ("GH") as used herein is defined as a hormone that relates to growth and acts as a chemical messenger to exert its action on a target cell. In a specific embodiment, the growth hormone is released by the action of growth hormone releasing hormone.

[0066] The term "growth hormone releasing hormone" ("GHRH") as used herein is defined as a hormone that facilitates or stimulates release of growth hormone, and in a lesser extent other pituitary hormones, such as prolactin. It is understood that the GHRH, the recombinant GHRH, or a functional biological equivalent are biologically active.

[0067] The term "heterologous nucleic acid sequence" as used herein is defined as a DNA sequence comprising differing regulatory and expression elements.

[0068] The term "lean body mass" ("LBM") as used herein is defined as the mass of the body of an animal attributed to non-fat tissue such as muscle.

[0069] The term "modified cells" as used herein is defined as the cells from a subject that have an additional nucleic acid sequence introduced into the cell.

[0070] The term "modified-donor-cells" as used herein refers to any donor-cells that harbor a GHRH encoding nucleic acid sequence.

[0071] The term "molecular switch" as used herein refers to a molecule that is delivered into a subject that can regulate transcription of a gene. A skilled artisan recognizes that there are many such switches known in the art, such as a tetracycline switch, a zinc finger switch, a glucocorticoid switch, and so forth.

[0072] The term "nucleic acid expression construct" as used herein refers to any type of genetic construct comprising a nucleic acid coding for a RNA capable of being transcribed. The term "expression vector" can also be used interchangeably herein. In specific embodiments, the nucleic acid expression construct comprises: a promoter; a nucleotide sequence of interest; and a 3′ untranslated region; wherein the promoter, the nucleotide sequence of interest, and the 3' untranslated region are operatively linked; and *in vivo* expression of the nucleotide sequence of interest is regulated by the promoter.

[0073] The term "operatively linked" as used herein refers to elements or structures in a nucleic acid sequence that are linked by operative ability and not physical location. The elements or structures are capable of, or characterized by accomplishing a desired operation. It is recognized by one of ordinary skill in the art that it is not necessary for elements or structures in a nucleic acid sequence to be in a tandem or adjacent order to be operatively linked.

[0074] The term "poly-L-glutamate ("PLG")" as used herein refers to a biodegradable polymer of L-glutamic acid that is suitable for use as a vector or adjuvant for DNA transfer into cells with or without electroporation.

[0075] The term "post-injection" as used herein refers to a time period following the introduction of a nucleic acid cassette (that contains heterologous nucleic acid sequence encoding GHRH or functional biological equivalent thereof) into the cells of a subject and

the allowing of the expression of the encoded gene to occur while the modified cells are within the living organism.

[0076] The term "plasmid" as used herein refers generally to a construction comprised of extra-chromosomal genetic material, usually of a circular duplex of DNA that can replicate independently of chromosomal DNA. Plasmids, or fragments thereof, may be used as vectors. Plasmids are double-stranded DNA molecule that occur or are derived from bacteria and (rarely) other microorganisms. However, mitochondrial and chloroplast DNA, yeast killer and other cases are commonly excluded.

[0077] The term "plasmid mediated gene supplementation" as used herein refers a method to allow a subject to have prolonged exposure to a therapeutic range of a therapeutic protein by utilizing a nucleic acid expression construct *in vivo*.

[0078] The term "pulse voltage device," or "pulse voltage injection device" as used herein relates to an apparatus that is capable of causing or causes uptake of nucleic acid molecules into the cells of an organism by emitting a localized pulse of electricity to the cells. The cell membrane then destabilizes, forming passageways or pores. Conventional devices of this type are calibrated to allow one to select or adjust the desired voltage amplitude and the duration of the pulsed voltage. The primary importance of a pulse voltage device is the capability of the device to facilitate delivery of compositions of the invention, particularly linear DNA fragments, into the cells of the organism.

[0079] The term "plasmid backbone" as used herein refers to a sequence of DNA that typically contains a bacterial origin of replication, and a bacterial antibiotic selection gene, which are necessary for the specific growth of only the bacteria that are transformed with the proper plasmid. However, there are plasmids, called mini-circles, that lack both the antibiotic resistance gene and the origin of replication (Darquet et al., 1997; Darquet et al., 1999; Soubrier et al., 1999). The use of *in vitro* amplified expression plasmid DNA (i.e. non-viral expression systems) avoids the risks associated with viral vectors. The non-viral expression systems products generally have low toxicity due to the use of "species-specific" components for gene delivery, which minimizes the risks of immunogenicity generally associated with viral vectors. One aspect of the current invention is that the plasmid backbone does not contain viral nucleotide sequences.

[0080] The term "promoter" as used herein refers to a sequence of DNA that directs the transcription of a gene. A promoter may direct the transcription of a prokaryotic or eukaryotic gene. A promoter may be "inducible", initiating transcription in response to an inducing agent or, in contrast, a promoter may be "constitutive", whereby an inducing agent does not regulate the rate of transcription. A promoter may be regulated in a tissue-specific or tissue-preferred manner, such that it is only active in transcribing the operable linked coding region in a specific tissue type or types.

[0081] The term "rate of bone healing" as used herein is defined as the time required to repair a bone fracture.

[0082] The term "recipient-subject" as used herein refers to any species of the animal kingdom wherein modified-donor-cells can be introduced from a donor-subject.

[0083] The term "regulator protein" as used herein refers to a protein that increases or facilitates transcription of a target nucleic acid sequence.

[0084] The term "residual linear plasmid backbone" as used herein comprises any fragment of the plasmid backbone that is left at the end of the process making the nucleic acid expression plasmid linear.

[0085] The terms "subject" or "animal" as used herein refers to any species of the animal kingdom. In preferred embodiments, it refers more specifically to humans and domesticated animals used for: pets (*e.g.* cats, dogs, *etc.*); work (*e.g.* horses, *etc.*); food (cows, chicken, fish, lambs, pigs, etc); and all others known in the art.

[0086] The term "tissue" as used herein refers to a collection of similar cells and the intercellular substances surrounding them. A skilled artisan recognizes that a tissue is an aggregation of similarly specialized cells for the performance of a particular function. For the scope of the present invention, the term tissue does not refer to a cell line, a suspension of cells, or a culture of cells. In a specific embodiment, the tissue is electroporated *in vivo*. In another embodiment, the tissue is not a plant tissue. A skilled artisan recognizes that there are four basic tissues in the body: 1) epithelium; 2) connective tissues, including blood, bone, and cartilage; 3) muscle tissue; and 4) nerve tissue. In a specific embodiment, the methods

and compositions are directed to transfer of linear DNA into a muscle tissue by electroporation.

[0087] The term "therapeutic element" as used herein comprises nucleic acid sequences that will lead to an *in vivo* expression of an encoded gene product. One skilled in the art of molecular biology will recognize that the therapeutic element may include, but is not limited to a promoter sequence, a transgene, a poly A sequence, or a 3' or 5' UTR.

[0088] The term "transfects" as used herein refers to introduction of a nucleic acid into a eukaryotic cell. In some embodiments, the cell is not a plant tissue or a yeast cell.

[0089] The term "viral backbone" as used herein refers to a nucleic acid sequence that does not contain a promoter, a gene, and a 3' poly A signal or an untranslated region, but contain elements including, but not limited at site-specific genomic integration Rep and inverted terminal repeats ("ITRs") or the binding site for the tRNA primer for reverse transcription, or a nucleic acid sequence component that induces a viral immunogenicity response when inserted in vivo, allows integration, affects specificity and activity of tissue specific promoters, causes transcriptional silencing or poses safety risks to the subject.

[0090] The term "vascular pressure pulse" refers to a pulse of pressure from a large volume of liquid to facilitate uptake of a vector into a cell. A skilled artisan recognizes that the amount and duration of the vascular pressure pulse is dependent on the tissue, size, and overall health of the recipient animal, and furthermore knows how to determine such parameters empirically.

[0091] The term "vector" as used herein refers to a construction comprised of genetic material designed to direct transformation of a targeted cell by delivering a nucleic acid sequence into that cell. A vector may contain multiple genetic elements positionally and sequentially oriented with other necessary elements such that an included nucleic acid cassette can be transcribed and when necessary translated in the transfected cells. These elements are operatively linked. The term "expression vector" refers to a DNA plasmid that contains all of the information necessary to produce a recombinant protein in a heterologous cell.

[0092] The present invention concerns a method for decreasing the body fat proportion, increasing lean body mass ("LBM"), increasing bone density, increasing the rate

of bone healing, or a combination thereof, of an animal subject. In general the present invention can be accomplished by delivering a nucleic acid sequence encoding GHRH or functional biological equivalent thereof into the cells of the subject (e.g. somatic, stem, or germ cells) and allowing expression of the encoded gene to occur while the modified cells are within the living organism. The subsequent expression of the GHRH or functional biological equivalent thereof is regulated by a tissue specific promoter (e.g. muscle), and/or by a regulator protein that contains a modified ligand binding domain (e.g. molecular switch), which will only be active when the correct modified ligand (e.g. mifepistone) is administered to the subject. The expression and ensuing release of GHRH or functional biological equivalent thereof by the modified cells within the living organism will decrease the body fat proportion, increase the LBM, increase the bone density, and/or increase the bone healing rate of the subject.

[0093] One aspect of the current invention is a method for altering lean body mass in a subject by utilizing a nucleic acid expression vector regulated by a constitutive promoter. The method comprises delivering into cells of the subject the nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof. In a specific embodiment, the nucleic acid expression construct is delivered into the cells of the subject via electroporation, wherein the cells comprise somatic, stem or germ cells. In another specific embodiment the nucleic acid expression construct comprises SeqID No: 11, SeqID No: 12, SeqID No: 13, SeqID No: 14, SeqID No: 17, SeqID No: 18, SeqID No: 19, SeqID No: 20, or SeqID No: 21. Transfection of the nucleic acid expression construct can be expedited by utilizing a transfection-facilitating polypeptide (e.g. charged polypeptide or poly-L-glutamate). The encoded GHRH or functional biological equivalent thereof are expressed in tissue specific cells of the subject, which comprises muscle cells. The encoded GHRH or the encoded functional biological equivalent of GHRH are biologically active polypeptides that have been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide. In a preferred embodiment the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6). Additionally, the encoded GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion in the subject.

[0094] A second aspect of the current invention is a method for altering lean body mass in a subject by utilizing a nucleic acid expression vector regulated by a molecular switch molecule. The method comprises steps of delivering into cells of the subject a first nucleic acid expression construct (SeqID No: 26), a second nucleic acid expression construct (SeqID No: 27), and a molecular switch; wherein the first nucleic acid expression construct encodes growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof; and wherein the second nucleic acid expression construct has an encoding region of a regulator protein; and delivering a molecular switch molecule into the subject, wherein the molecular switch molecule governs activation of the regulator protein and the regulator protein governs the activation of the first nucleic acid expression construct. In some specific embodiments, the nucleic acid expression construct further comprises a transfection-facilitating polypeptide (e.g. a charged polypeptide or poly-L-glutamate). The delivering step of the first nucleic acid and the second nucleic acid expression construct into the cells of the subject is via electroporation. A specific embodiment of this method comprises that delivering the nucleic acid expression construct into the cells of the subject initiates expression of the encoded regulatory protein, but the regulatory protein is inactive. However, upon delivering a molecular switch (e.g. mifepristone) into the subject, the regulatory protein becomes active, and the active regulatory protein initiates expression of the GHRH or functional biological equivalent encoded on the first nucleic acid sequence. The encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide. The encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6). The encoded GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion in the subject. In another specific embodiment, the first nucleic acid expression vector encodes a polypeptide of sequence SeqID No: 1, SeqID No: 2, SeqID No: 3, or SeqID No: 4.

[0095] A third aspect of the current invention is a altering lean body mass in a subject comprising the steps of: delivering into a subject a recombinant growth-hormone-releasing-hormone ("GHRH") or a biological functional equivalent thereof, wherein the recombinant GHRH is a biologically active polypeptide. In specific embodiments, the recombinant functional biological equivalent of GHRH is a polypeptide that has been

engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide. In another specific embodiment, the recombinant GHRH or functional biological equivalent thereof is of formula (SEQID No: 6). The recombinant GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion in the subject.

[0096] A fourth aspect of the current invention is a method for altering bone properties in a subject by utilizing a nucleic acid expression vector regulated by a constitutive promoter. The method comprises delivering into cells of the subject the nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof. In a specific embodiment, the nucleic acid expression construct is delivered into the cells of the subject via electroporation, wherein the cells comprise somatic, stem or germ cells. In another specific embodiment the nucleic acid expression construct comprises SeqID No: 11, SeqID No: 12, SeqID No: 13, SeqID No: 14, SeqID No: 17, SeqID No: 18, SeqID No: 19, SeqID No: 20, or SeqID No: 21. Transfection of the nucleic acid expression construct can be expedited by utilizing a transfection-facilitating polypeptide (e.g. charged polypeptide or poly-L-glutamate). The encoded GHRH or functional biological equivalent thereof are expressed in tissue specific cells of the subject, which comprises muscle cells. The encoded GHRH or the encoded functional biological equivalent of GHRH are biologically active polypeptides that have been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide. In a preferred embodiment the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6). Additionally, the encoded GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion in the subject.

[0097] A fifth aspect of the current invention is a method for altering bone properties in a subject by utilizing a nucleic acid expression vector regulated by a molecular switch molecule. The method comprises steps of delivering into cells of the subject a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch; wherein the first nucleic acid expression construct encodes growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof; and wherein the second nucleic acid expression construct has an encoding region of a regulator protein; and delivering a molecular switch molecule into the subject, wherein the molecular switch

molecule governs activation of the regulator protein and the regulator protein governs the activation of the first nucleic acid expression construct. In some specific embodiments, the nucleic acid expression construct further comprises a transfection-facilitating polypeptide (e.g. a charged polypeptide or poly-L-glutamate). The delivering step of the first nucleic acid and the second nucleic acid expression construct into the cells of the subject is via electroporation. A specific embodiment of this method comprises that delivering the nucleic acid expression construct into the cells of the subject initiates expression of the encoded regulatory protein, but the regulatory protein is inactive. However, upon delivering a molecular switch (e.g. mifepristone) into the subject, the regulatory protein becomes active, and the active regulatory protein initiates expression of the GHRH or functional biological equivalent encoded on the first nucleic acid sequence. The encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide. The encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6). The encoded GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion in the subject. In another specific embodiment, the first nucleic acid expression vector encodes a polypeptide of sequence SeqID No: 1, SeqID No: 2, SeqID No: 3, or SeqID No: 4.

[0098] A sixth aspect of the current invention is a method for altering bone properties in a subject comprising the steps of: delivering into a subject a recombinant growth-hormone-releasing-hormone ("GHRH") or a biological functional equivalent thereof, wherein the recombinant GHRH is a biologically active polypeptide. In specific embodiments, the recombinant functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide. In another specific embodiment, the recombinant GHRH or functional biological equivalent thereof is of formula (SEQID No: 6). The recombinant GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion in the subject.

[0099] The plasmid-mediated supplementation of GHRH approach described herein offers advantages over the limitations of directly injecting recombinant GHRH protein. Expression of nucleic acid sequences encoding novel functional biological

equivalents of GHRH that are serum protease resistant can be directed by an expression plasmid controlled by a synthetic muscle-specific promoter. Expression of such GHRH or functional biological equivalent thereof elicited high GH and IGF-I levels in pigs following delivery by intramuscular injection and *in vivo* electroporation (Draghia-Akli et al., 1999). The process of *in vivo* electroporation may involve externally supplied electrodes, or in the case of needles, internally supplied electrodes to aid in the inclusion of desired nucleotide sequences into the cells of the subject within the living organism. Although *in vivo* electroporation is the preferred method of introducing the heterologous nucleic acid encoding system into the cells of the subject, other methods exist and are known by a person skilled in the art (*e.g.* electroporation, lipofectamine, calcium phosphate, *ex vivo* transformation, direct injection, DEAE dextran, sonication loading, receptor mediated transfection, microprojectile bombardment, *etc.*). For example, it is also possible to introduce the nucleic acid sequence that encodes the GHRH or functional biological equivalent thereof directly into the cells of the subject by first removing the cells from the body of the subject or donor, maintaining the cells in culture, then introducing the nucleic acid encoding system by a variety of methods (*e.g.* electroporation, lipofectamine, calcium phosphate, *ex vivo* transformation, direct injection, DEAE dextran, sonication loading, receptor mediated transfection, microprojectile bombardment, *etc.*), and finally reintroducing the modified cells into the original subject or other host subject (the *ex vivo* method). The GHRH sequence can be cloned into an adenovirus vector or an adeno-associated vector and delivered by simple intramuscular injection, or intravenous or intra-arterial injection. Plasmid DNA carrying the GHRH sequence can be complexed with cationic lipids or liposomes and delivered intramuscularly, intravenously or subcutaneously.

[0100] Administration as used herein refers to the route of introduction of a vector or carrier of DNA into the body. Administration can be directly to a target tissue or by targeted delivery to the target tissue after systemic administration. In particular, the present invention can be used for supplementing GHRH by administration of the vector, such as a plasmid, to the body in order to establish controlled expression of the specific nucleic acid sequence within tissues at certain useful levels.

[0101] The preferred means for administration of vector and use of formulations for delivery are described above. The preferred embodiment is by *in vivo* electroporation.

[0102] The route of administration of any selected vector construct will depend on the particular use for the expression vectors. In general, a specific formulation for each vector construct used will focus on vector uptake with regard to the particular targeted tissue, followed by demonstration of efficacy. Uptake studies will include uptake assays to evaluate cellular uptake of the vectors and expression of the tissue specific DNA of choice. Such assays will also determine the localization of the target DNA after uptake, and establishing the requirements for maintenance of steady-state concentrations of expressed protein. Efficacy and cytotoxicity can then be tested. Toxicity will not only include cell viability but also cell function.

[0103] Muscle cells have the unique ability to take up DNA from the extracellular space after simple injection of DNA particles as a solution, suspension, or colloid into the muscle. Expression of DNA by this method can be sustained for several months. DNA uptake in muscle cells is further enhanced by utilizing *in vivo* electroporation.

[0104] Delivery of formulated DNA vectors involves incorporating DNA into macromolecular complexes that undergo endocytosis by the target cell. Such complexes may include lipids, proteins, carbohydrates, synthetic organic compounds, or inorganic compounds. The characteristics of the complex formed with the vector (size, charge, surface characteristics, composition) determines the bioavailability of the vector within the body. Other elements of the formulation function as ligand which interact with specific receptors on the surface or interior of the cell. Other elements of the formulation function to enhance entry into the cell, release from the endosome, and entry into the nucleus.

[0105] Delivery can also be through use of DNA transporters. DNA transporters refers to molecules which bind to DNA vectors and are capable of being taken up by epidermal cells. DNA transporters contain a molecular complex capable of non-covalently binding to DNA and efficiently transporting the DNA through the cell

membrane. It is preferable that the transporter also transport the DNA through the nuclear membrane. See, *e.g.*, the following applications all of which (including drawings) are hereby incorporated by reference herein: (1) Woo *et al.*, U.S. Patent No. 6,150,168 entitled: "A DNA Transporter System and Method of Use;" (2) Woo *et al.*, PCT/US93/02725, entitled "A DNA Transporter System and method of Use", filed Mar. 19, 1993; (3) Woo *et al.*, U.S. Patent No. 6,177,554 "Nucleic Acid Transporter Systems and Methods of Use;" (4) Szoka *et al.*, U.S. Patent No. 5,955,365 entitled "Self-Assembling Polynucleotide Delivery System;" and (5) Szoka *et al.*, PCT/US93/03406, entitled "Self-Assembling Polynucleotide Delivery System", filed Apr. 5, 1993.

[0106] Another method of delivery involves a DNA transporter system. The DNA transporter system consists of particles containing several elements that are independently and non-covalently bound to DNA. Each element consists of a ligand which recognizes specific receptors or other functional groups such as a protein complexed with a cationic group that binds to DNA. Examples of cations which may be used are spermine, spermine derivatives, histone, cationic peptides and/or polylysine. One element is capable of binding both to the DNA vector and to a cell surface receptor on the target cell. Examples of such elements are organic compounds which interact with the asialoglycoprotein receptor, the folate receptor, the mannose-6-phosphate receptor, or the carnitine receptor. A second element is capable of binding both to the DNA vector and to a receptor on the nuclear membrane. The nuclear ligand is capable of recognizing and transporting a transporter system through a nuclear membrane. An example of such a ligand is the nuclear targeting sequence from SV40 large T antigen or histone. A third element is capable of binding to both the DNA vector and to elements which induce episomal lysis. Examples include inactivated virus particles such as adenovirus, peptides related to influenza virus hemagglutinin, or the GALA peptide described in the Skoka patent cited above.

[0107] Administration may also involve lipids. The lipids may form liposomes which are hollow spherical vesicles composed of lipids arranged in unilamellar, bilamellar, or multilamellar fashion and an internal aqueous space for entrapping water soluble compounds, such as DNA, ranging in size from 0.05 to several microns in diameter. Lipids may be useful without forming liposomes. Specific examples include the

use of cationic lipids and complexes containing DOPE which interact with DNA and with the membrane of the target cell to facilitate entry of DNA into the cell.

[0108] Gene delivery can also be performed by transplanting genetically engineered cells. For example, immature muscle cells called myoblasts may be used to carry genes into the muscle fibers. Myoblast genetically engineered to express recombinant human growth hormone can secrete the growth hormone into the animal's blood. Secretion of the incorporated gene can be sustained over periods up to 3 months.

[0109] Myoblasts eventually differentiate and fuse to existing muscle tissue. Because the cell is incorporated into an existing structure, it is not only tolerated but nurtured. Myoblasts can easily be obtained by taking muscle tissue from an individual who needs supplementation of GHRH, and the genetically engineered cells can also be easily put back with out causing damage to the patient Is muscle. Similarly, keratinocytes may be used to delivery genes to tissues. Large numbers of keratinocytes can be generated by cultivation of a small biopsy. The cultures can be prepared as stratified sheets and when grafted to humans, generate epidermis which continues to improve in histotypic quality over many years. The keratinocytes are genetically engineered while in culture by transfecting the keratinocytes with the appropriate vector. Although keratinocytes are separated from the circulation by the basement membrane dividing the epidermis from the dermis, human keratinocytes secrete into circulation the protein produced.

[0110] Delivery may also involve the use of viral vectors. For example, an adenoviral vector may be constructed by replacing the E1 region of the virus genome with the vector elements described in this invention including promoter, 5'UTR, 3'UTR and nucleic acid cassette and introducing this recombinant genome into 293 cells which will package this gene into an infectious virus particle. Virus from this cell may then be used to infect tissue ex vivo or *in vivo* to introduce the vector into tissues leading to expression of the gene in the nucleic acid cassette.

[0111] Although not wanting to be bound by theory, it is believed that in order to provide an acceptable safety margin for the use of such heterologous nucleic acid sequences in humans, a regulated gene expression system is mandated to possess low

levels of basal expression of GHRH, and still retain a high inducibility. Thus, target gene expression can be regulated by incorporating molecular switch technology as schematically diagramed in FIGURE 1A. The commercially available GeneSwitch® system for ligand-dependent induction of transgene expression is based on a C-terminally truncated progesterone receptor that fails to bind to its natural agonist, progesterone, but instead is activated by antiprogestins, such as mifepristone ("MFP") (Vegeto et al., 1992; Xu et al., 1996). Thus, the heterologous nucleic acid sequence introduced into the cells of the subject requires MFP to be transcriptionally activated. The chimeric regulator protein of the GeneSwitch® system consists of the ligand binding domain of the truncated human progesterone receptor that has been fused to the DNA binding domain of the yeast GAL4 protein (which binds a specific 17 bp recognition sequence) and a transcriptional activation domain from the p65 subunit of human NF-kB (Abruzzese et al., 1999; Abruzzese et al., 2000). The gene for the GeneSwitch® regulator protein was inserted into a myogenic expression vector, designated pGS1633, which is expressed constitutively under the control of a muscle-specific skeletal alpha-actin ("SK") promoter The GHRH plasmid, designated, p6xGal4/TATA-GHRH, or pGHRH1633 contains an inducible promoter that consists of six copies of the 17-mer Gal4 binding site fused to a minimal TATA box promoter. The GHRH coding sequence is a 228-bp fragment of super-porcine mutated GHRH cDNA, termed HV-GHRH (Draghia-Akli et al., 1999). The HV-GHRH molecule displays a high degree of stability in serum, with a half-life of 6 hours, versus the natural GHRH, that has a 6-12 min half-life. The muscle-specific GeneSwitch® and inducible GHRH plasmids both have a 5' untranslated region that contains a synthetic intron, and a 3' untranslated region/poly(A) site that is from the human GH gene.

[0112] Recombinant GH replacement therapy is widely used clinically, with beneficial effects, but generally, the doses are supraphysiological. Such elevated doses of recombinant GH are associated with deleterious side-effects, for example, up to 30% of the recombinant GH treated patients report a higher frequency of insulin resistance (Blethen, 1995; Verhelst et al., 1997) or accelerated bone epiphysis growth and closure in pediatric patients (Blethen and Rundle, 1996). In addition, molecular heterogeneity of circulating GH may have important implications in growth and homeostasis, which can lead to a less potent GH that has a reduced ability to stimulate the prolactin receptor; it has

also been described that the 20kDa form of GH has less potency to cause urine retention than the 22 kDa form (Satozawa et al., 2000; Tsunekawa et al., 1999; Wada et al., 1998). These unwanted side effects result from the fact that treatment with recombinant exogenous GH protein raises basal levels of GH and abolishes the natural episodic pulses of GH. In contradistinction, no side effects have been reported for recombinant GHRH therapies. The normal levels of GHRH in the pituitary portal circulation range from about 150-to-800 pg/ml, while systemic circulating values of the hormone are up to about 100-500 pg/ml. Some patients with acromegaly caused by extracranial tumors have level that is nearly 10 times as high (*e.g.* 50 ng/ml of immunoreactive GHRH) (Thorner et al., 1984). Long-term studies using recombinant GHRH therapies (1-5 years) in children and elderly humans have shown an absence of the classical GH side-effects, such as changes in fasting glucose concentration or, in pediatric patients, the accelerated bone epiphysal growth and closure or slipping of the capital femoral epiphysis (Chevalier et al., 2000) (Duck et al., 1992; Vittone et al., 1997). Numerous studies in humans, sheep or pigs showed that continuous infusion with recombinant GHRH protein restores the normal GH pattern without desensitizing GHRH receptors or depleting GH supplies (Dubreuil et al., 1990b). As this system is capable of a degree of feed-back which is abolished in the GH therapies, GHRH recombinant protein therapy may be more physiological than GH therapy. However, due to the short half-life of GHRH *in vivo*, frequent (one to three times per day) intravenous, subcutaneous or intranasal (requiring 300-fold higher dose) administrations are necessary (Evans et al., 1985; Thorner et al., 1986). Thus, as a chronic therapy, recombinant GHRH protein administration is not practical. A gene transfer approach, however could overcome this limitations to GHRH use. Moreover, a wide range of doses can be therapeutic. The choice of GHRH for a gene therapeutic application is favored by the fact that the gene, cDNA and native and several mutated molecules have been characterized for the pig and other species (Bohlen et al., 1983; Guillemin et al., 1982), and the measurement of therapeutic efficacy is straightforward and unequivocal.

[0113] Among the non-viral techniques for gene transfer *in vivo*, the direct injection of plasmid DNA into muscle is simple, inexpensive, and safe. The inefficient DNA uptake into muscle fibers after simple direct injection hag led to relatively low expression levels (Prentice et al., 1994; Wells et al., 1997) In addition, the duration of the

transgene expression has been short (Wolff et al., 1990). The most successful previous clinical applications have been confined to vaccines (Danko and Wolff, 1994; Tsurumi et al., 1996). Recently, significant progress to enhance plasmid delivery *in vivo* and subsequently to achieve physiological levels of a secreted protein was obtained using the electroporation technique. Recently, significant progress has been obtained using electroporation to enhance plasmid delivery *in vivo*. Electroporation has been used very successfully to transfect tumor cells after injection of plasmid (Lucas et al., 2002; Matsubara et al., 2001) or to deliver the anti-tumor drug bleomycin to cutaneous and subcutaneous tumors in humans (Gehl et al., 1998; Heller et al., 1996). Electroporation also has been extensively used in mice (Lesbordes et al., 2002; Lucas et al., 2001; Vilquin et al., 2001), rats (Terada et al., 2001; Yasui et al., 2001), and dogs (Fewell et al., 2001) to deliver therapeutic genes that encode for a variety of hormones, cytokines or enzymes. Our previous studies using growth hormone releasing hormone (GHRH) showed that plasmid therapy with electroporation is scalable and represents a promising approach to induce production and regulated secretion of proteins in large animals and humans (Draghia-Akli et al., 1999; Draghia-Akli et al., 2002). Electroporation also has been extensively used in rodents and other small animals (Bettan et al., 2000; Yin and Tang, 2001). It has been observed that the electrode configuration affects the electric field distribution, and subsequent results (Gehl et al., 1999; Miklavcic et al., 1998). Preliminary experiments indicated that for a large animal model, needle electrodes give consistently better reproducible results than external caliper electrodes.

[0114] Combining the powerful electroporation delivery method with an improved plasmid DNA vector system produced significant changes that decreased the body fat proportion, increased lean body mass ("LBM"), or both, in an animal, such as a large animal, at very low plasmid dosage.

I. Vectors

[0115] The term "vector" is used to refer to a carrier nucleic acid molecule into which a nucleic acid sequence can be inserted for introduction into a cell wherein, in some embodiments, it can be replicated. A nucleic acid sequence can be native to the animal, or it can be "exogenous," which means that it is foreign to the cell into which the vector is

being introduced or that the sequence is homologous to a sequence in the cell but in a position within the host cell nucleic acid in which the sequence is ordinarily not found. Vectors include plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), linear DNA fragments, and artificial chromosomes (*e.g.*, YACs), although in a preferred embodiment the vector contains substantially no viral sequences. One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (see, for example, (Sambrook et al., 1989).

[0116] The term "expression vector" refers to any type of genetic construct comprising a nucleic acid coding for a RNA capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. In other cases, these sequences are not translated, for example, in the production of antisense molecules or ribozymes. Expression vectors can contain a variety of "control sequences," which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operatively linked coding sequence in a particular host cell. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described *infra*.

II. Plasmid Vectors

[0117] In certain embodiments, a linear DNA fragment from a plasmid vector is contemplated for use to transfect a eukaryotic cell, particularly a mammalian cell. In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. In a non-limiting example, *E. coli* is often transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, for example, promoters which can be used by the microbial organism for expression of its own proteins. A skilled artisan recognizes that any plasmid in the art may be modified for use in the methods of the present invention. In a specific embodiment, for example, a GHRH vector

used for the therapeutic applications is derived from pBlueScript KS+ and has a kanamycin resistance gene.

[0118] In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEM™-11 may be utilized in making a recombinant phage vector which can be used to transform host cells, such as, for example, *E. coli* LE392.

[0119] Further useful plasmid vectors include pIN vectors (Inouye et al., 1985); and pGEX vectors, for use in generating glutathione S-transferase ("GST") soluble fusion proteins for later purification and separation or cleavage. Other suitable fusion proteins are those with β-galactosidase, ubiquitin, and the like.

[0120] Bacterial host cells, for example, *E. coli*, comprising the expression vector, are grown in any of a number of suitable media, for example, LB. The expression of the recombinant protein in certain vectors may be induced, as would be understood by those of skill in the art, by contacting a host cell with an agent specific for certain promoters, *e.g.*, by adding IPTG to the media or by switching incubation to a higher temperature. After culturing the bacteria for a further period, generally of between 2 and 24 h, the cells are collected by centrifugation and washed to remove residual media.

III. Promoters and Enhancers

[0121] A promoter is a control sequence that is a region of a nucleic acid sequence at which initiation and rate of transcription of a gene product are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind, such as RNA polymerase and other transcription factors, to initiate the specific transcription a nucleic acid sequence. The phrases "operatively positioned," "operatively linked," "under control," and "under transcriptional control" mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and/or expression of that sequence.

[0122] A promoter generally comprises a sequence that functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as, for example, the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation. Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. To bring a coding sequence "under the control of" a promoter, one positions the 5′ end of the transcription initiation site of the transcriptional reading frame "downstream" of (*i.e.*, 3′ of) the chosen promoter. The "upstream" promoter stimulates transcription of the DNA and promotes expression of the encoded RNA.

[0123] The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

[0124] A promoter may be one naturally associated with a nucleic acid sequence, as may be obtained by isolating the 5′ non-coding sequences located upstream of the coding segment and/or exon. Such a promoter can be referred to as "endogenous." Similarly, an enhancer may be one naturally associated with a nucleic acid sequence, located either downstream or upstream of that sequence. Alternatively, certain advantages will be gained by positioning the coding nucleic acid segment under the control of a recombinant, synthetic or heterologous promoter, which refers to a promoter that is not normally associated with a nucleic acid sequence in its natural environment. A recombinant, synthetic or heterologous enhancer refers also to an enhancer not normally associated with a nucleic acid sequence in its natural environment. Such promoters or

enhancers may include promoters or enhancers of other genes, and promoters or enhancers isolated from any other virus, or prokaryotic or eukaryotic cell, and promoters or enhancers not "naturally occurring," *i.e.*, containing different elements of different transcriptional regulatory regions, and/or mutations that alter expression. For example, promoters that are most commonly used in recombinant DNA construction include the β-lactamase (penicillinase), lactose and tryptophan (trp) promoter systems. In addition to producing nucleic acid sequences of promoters and enhancers synthetically, sequences may be produced using recombinant cloning and/or nucleic acid amplification technology, including PCR™, in connection with the compositions disclosed herein (see U.S. Patent Nos. 4,683,202 and 5,928,906, each incorporated herein by reference). Furthermore, it is contemplated the control sequences that direct transcription and/or expression of sequences within non-nuclear organelles such as mitochondria, chloroplasts, and the like, can be employed as well.

[0125] Naturally, it will be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the organelle, cell type, tissue, organ, or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression, (see, for example (Sambrook et al., 1989)). The promoters employed may be constitutive, tissue-specific, inducible, and/or useful under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins and/or peptides. The promoter may be heterologous or endogenous.

[0126] Additionally any promoter/enhancer combination (as per, for example, the Eukaryotic Promoter Data Base EPDB, http://www.epd.isb-sib.ch/) could also be used to drive expression. Use of a T3, T7 or SP6 cytoplasmic expression system is another possible embodiment. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

[0127] Tables 1 and 2 list non-limiting examples of elements/promoters that may be employed, in the context of the present invention, to regulate the expression of a

RNA. Table 2 provides non-limiting examples of inducible elements, which are regions of a nucleic acid sequence that can be activated in response to a specific stimulus.

TABLE 1
Promoter and/or Enhancer

Promoter/Enhancer	Relevant References
Immunoglobulin Heavy Chain	
Immunoglobulin Light Chain	
T-Cell Receptor	
HLA DQ a and/or DQ β	
β-Interferon	
Interleukin-2	
Interleukin-2 Receptor	
MHC Class II 5	
MHC Class II HLA-Dra	
β-Actin	(Kawamoto et al., 1988; Kawamoto et al., 1989)
Muscle Creatine Kinase (MCK)	(Horlick and Benfield, 1989; Jaynes et al., 1988)
Prealbumin (Transthyretin)	
Elastase I	
Metallothionein (MTII)	(Inouye et al., 1994; Narum et al., 2001; Skroch et al., 1993)
Collagenase	
Albumin	(Pinkert et al., 1987; Tronche et al., 1989)
α-Fetoprotein	
γ-Globin	
β-Globin	(Tronche et al., 1990; Trudel and Costantini, 1987)
c-fos	
c-HA-*ras*	
Insulin	(German et al., 1995; Ohlsson et al., 1991)
Neural Cell Adhesion Molecule (NCAM)	
α_1-Antitrypsin	
H2B (TH2B) Histone	
Mouse and/or Type I Collagen	
Glucose-Regulated Proteins (GRP94 and GRP78)	
Rat Growth Hormone	(Larsen et al., 1986)
Human Serum Amyloid A (SAA)	
Troponin I (TN I)	(Lin et al., 1991; Yutzey and Konieczny, 1992)
Platelet-Derived Growth Factor (PDGF)	(Pech et al., 1989)
Duchenne Muscular Dystrophy	(Klamut et al., 1990; Klamut et al., 1996)
SV40	
Polyoma	
Retroviruses	
Papilloma Virus	
Hepatitis B Virus	

TABLE 1 Promoter and/or Enhancer	
Promoter/Enhancer	Relevant References
Human Immunodeficiency Virus	
Cytomegalovirus (CMV)	(Boshart et al., 1985; Dorsch-Hasler et al., 1985)
Gibbon Ape Leukemia Virus	
Synthetic muscle specific promoters (c5-12, c1-28)	(Draghia-Akli et al., 1999; Draghia-Akli et al., 2002; Li et al., 1999)

TABLE 2 Element/Inducer	
Element	Inducer
MT II	Phorbol Ester (TFA) Heavy metals
MMTV (mouse mammary tumor virus)	Glucocorticoids
β-Interferon	Poly(rI)x / Poly(rc)
Adenovirus 5 E2	EIA
Collagenase	Phorbol Ester (TPA)
Stromelysin	Phorbol Ester (TPA)
SV40	Phorbol Ester (TPA)
Murine MX Gene	Interferon, Newcastle Disease Virus
GRP78 Gene	A23187
α-2-Macroglobulin	IL-6
Vimentin	Serum
MHC Class I Gene H-2κb	Interferon
HSP70	EIA, SV40 Large T Antigen
Proliferin	Phorbol Ester-TPA
Tumor Necrosis Factor α	PMA
Thyroid Stimulating Hormone α Gene	Thyroid Hormone

[0128] The identity of tissue-specific promoters or elements, as well as assays to characterize their activity, is well known to those of skill in the art. Nonlimiting examples of such regions include the human LIMK2 gene (Nomoto et al., 1999), the somatostatin receptor 2 gene (Kraus et al., 1998), murine epididymal retinoic acid-binding gene (Lareyre et al., 1999), human CD4 (Zhao-Emonet et al., 1998), mouse alpha2 (XI) collagen (Liu et al., 2000; Tsumaki et al., 1998), D1A dopamine receptor gene (Lee et al., 1997), insulin-like growth factor II (Dai et al., 2001; Wu et al., 1997), and human platelet endothelial cell adhesion molecule-1 (Almendro et al., 1996).

[0129] In a preferred embodiment, a synthetic muscle promoter is utilized, such as SPc5-12 (Li et al., 1999), which contains a proximal serum response element ("SRE") from skeletal α-actin, multiple MEF-2 sites, MEF-1 sites, and TEF-1 binding sites, and greatly exceeds the transcriptional potencies of natural myogenic promoters.

The uniqueness of such a synthetic promoter is a significant improvement over, for instance, issued patents concerning a myogenic promoter and its use (*e.g.* U.S. Pat. No. 5,374,544) or systems for myogenic expression of a nucleic acid sequence (*e.g.* U.S. Pat. No. 5,298,422). In a preferred embodiment, the promoter utilized in the invention does not get shut off or reduced in activity significantly by endogenous cellular machinery or factors. Other elements, including *trans*-acting factor binding sites and enhancers may be used in accordance with this embodiment of the invention. In an alternative embodiment, a natural myogenic promoter is utilized, and a skilled artisan is aware how to obtain such promoter sequences from databases including the National Center for Biotechnology Information ("NCBI") GenBank database or the NCBI PubMed site. A skilled artisan is aware that these databases may be utilized to obtain sequences or relevant literature related to the present invention.

IV. Initiation Signals and Internal Ribosome Binding Sites

[0130] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be "in-frame" with the reading frame of the desired coding sequence to ensure translation of the entire insert. The exogenous translational control signals and initiation codons can be either natural or synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements.

[0131] In certain embodiments of the invention, the use of internal ribosome entry sites ("IRES") elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5′ methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988). IRES elements from two members of the picornavirus family (polio and encephalomyocarditis) have been described (Pelletier and Sonenberg, 1988), as well an IRES from a mammalian message (Macejak and Sarnow, 1991). IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be

transcribed together, each separated by an IRES, creating polycistronic messages. By virtue of the IRES element, each open reading frame is accessible to ribosomes for efficient translation. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message (see U.S. Patent Nos. 5,925,565 and 5,935,819, each herein incorporated by reference).

V. Multiple Cloning Sites

[0132] Vectors can include a MCS, which is a nucleic acid region that contains multiple restriction enzyme sites, any of which can be used in conjunction with standard recombinant technology to digest the vector (see, for example, (Carbonelli et al., 1999; Cocea, 1997; Levenson et al., 1998) incorporated herein by reference.) "Restriction enzyme digestion" refers to catalytic cleavage of a nucleic acid molecule with an enzyme that functions only at specific locations in a nucleic acid molecule. Many of these restriction enzymes are commercially available. Use of such enzymes is widely understood by those of skill in the art. Frequently, a vector is linearized or fragmented using a restriction enzyme that cuts within the MCS to enable exogenous sequences to be ligated to the vector. "Ligation" refers to the process of forming phosphodiester bonds between two nucleic acid fragments, which may or may not be contiguous with each other. Techniques involving restriction enzymes and ligation reactions are well known to those of skill in the art of recombinant technology.

VI. Splicing Sites

[0133] Most transcribed eukaryotic RNA molecules will undergo RNA splicing to remove introns from the primary transcripts. Vectors containing genomic eukaryotic sequences may require donor and/or acceptor splicing sites to ensure proper processing of the transcript for protein expression (see, for example, (Chandler et al., 1997), herein incorporated by reference.)

VII. Termination Signals

[0134] The vectors or constructs of the present invention will generally comprise at least one termination signal. A "termination signal" or "terminator" is comprised of the DNA sequences involved in specific termination of an RNA transcript by

an RNA polymerase. Thus, in certain embodiments a termination signal that ends the production of an RNA transcript is contemplated. A terminator may be necessary *in vivo* to achieve desirable message levels.

[0135] In eukaryotic systems, the terminator region may also comprise specific DNA sequences that permit site-specific cleavage of the new transcript so as to expose a polyadenylation site. This signals a specialized endogenous polymerase to add a stretch of about 200 A residues ("polyA") to the 3' end of the transcript. RNA molecules modified with this polyA tail appear to more stable and are translated more efficiently. Thus, in other embodiments involving eukaryotes, it is preferred that that terminator comprises a signal for the cleavage of the RNA, and it is more preferred that the terminator signal promotes polyadenylation of the message. The terminator and/or polyadenylation site elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

[0136] Terminators contemplated for use in the invention include any known terminator of transcription described herein or known to one of ordinary skill in the art, including but not limited to, for example, the termination sequences of genes, such as for example the bovine growth hormone terminator or viral termination sequences, such as for example the SV40 terminator. In certain embodiments, the termination signal may be a lack of transcribable or translatable sequence, such as due to a sequence truncation.

VIII. Polyadenylation Signals

[0137] In expression, particularly eukaryotic expression, one will typically include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed. Preferred embodiments include the SV40 polyadenylation signal, skeletal alpha actin 3'UTR or the human or bovine growth hormone polyadenylation signal, convenient and known to function well in various target cells. Polyadenylation may increase the stability of the transcript or may facilitate cytoplasmic transport.

IX. Origins of Replication

[0138] In order to propagate a vector in a host cell, it may contain one or more origins of replication sites (often termed "ori"), which is a specific nucleic acid sequence at which replication is initiated. Alternatively an autonomously replicating sequence ("ARS") can be employed if the host cell is yeast.

X. Selectable and Screenable Markers

[0139] In certain embodiments of the invention, cells containing a nucleic acid construct of the present invention may be identified *in vitro* or *in vivo* by including a marker in the expression vector. Such markers would confer an identifiable change to the cell permitting easy identification of cells containing the expression vector. Generally, a selectable marker is one that confers a property that allows for selection. A positive selectable marker is one in which the presence of the marker allows for its selection, while a negative selectable marker is one in which its presence prevents its selection. An example of a positive selectable marker is a drug resistance marker.

[0140] Usually the inclusion of a drug selection marker aids in the cloning and identification of transformants, for example, genes that confer resistance to neomycin, puromycin, hygromycin, DHFR, GPT, zeocin and histidinol are useful selectable markers. In addition to markers conferring a phenotype that allows for the discrimination of transformants based on the implementation of conditions, other types of markers including screenable markers such as GFP, whose basis is colorimetric analysis, are also contemplated. Alternatively, screenable enzymes such as herpes simplex virus thymidine kinase ("tk") or chloramphenicol acetyltransferase ("CAT") may be utilized. One of skill in the art would also know how to employ immunologic markers, possibly in conjunction with FACS analysis. The marker used is not believed to be important, so long as it is capable of being expressed simultaneously with the nucleic acid encoding a gene product. Further examples of selectable and screenable markers are well known to one of skill in the art.

XI. Electroporation

[0141] In certain embodiments of the present invention, a nucleic acid is introduced into an organelle, a cell, a tissue or an organism *via* electroporation. Electroporation involves the exposure of a suspension of cells and DNA to a high-voltage electric discharge. In some variants of this method, certain cell wall-degrading enzymes, such as pectin-degrading enzymes, are employed to render the target recipient cells more susceptible to transformation by electroporation than untreated cells (U.S. Patent No.5,384,253, incorporated herein by reference). Alternatively, recipient cells can be made more susceptible to transformation by mechanical wounding and other methods known in the art.

[0142] Transfection of eukaryotic cells using electroporation has been quite successful. Mouse pre-B lymphocytes have been transfected with human kappa-immunoglobulin genes (Potter et al., 1984), and rat hepatocytes have been transfected with the chloramphenicol acetyltransferase gene (Tur-Kaspa et al., 1986) in this manner.

[0143] The underlying phenomenon of electroporation is believed to be the same in all cases, but the exact mechanism responsible for the observed effects has not been elucidated. Although not wanting to be bound by theory, the overt manifestation of the electroporative effect is that cell membranes become transiently permeable to large molecules, after the cells have been exposed to electric pulses. There are conduits through cell walls, which under normal circumstances, maintain a resting transmembrane potential of ca. 90 mV by allowing bi-directional ionic migration.

[0144] Although not wanting to be bound by theory, electroporation makes use of the same structures, by forcing a high ionic flux through these structures and opening or enlarging the conduits. In prior art, metallic electrodes are placed in contact with tissues and predetermined voltages, proportional to the distance between the electrodes are imposed on them. The protocols used for electroporation are defined in terms of the resulting field intensities, according to the formula $E=V/d$, where ("E") is the field, ("V") is the imposed voltage and ("d") is the distance between the electrodes.

[0145] The electric field intensity *E* has been a very important value in prior art when formulating electroporation protocols for the delivery of a drug or macromolecule into the cell of the subject. Accordingly, it is possible to calculate any electric field intensity for a variety of protocols by applying a pulse of predetermined voltage that is proportional to the distance between electrodes. However, a caveat is that an electric field can be generated in a tissue with insulated electrodes (i.e. flow of ions is not necessary to create an electric field). Although not wanting to be bound by theory, it is the current that is necessary for successful electroporation not electric field per se.

[0146] During electroporation, the heat produced is the product of the interelectrode impedance, the square of the current, and the pulse duration. Heat is produced during electroporation in tissues and can be derived as the product of the inter-electrode current, voltage and pulse duration. The protocols currently described for electroporation are defined in terms of the resulting field intensities *E*, which are dependent on short voltage pulses of unknown current. Accordingly, the resistance or heat generated in a tissue cannot be determined, which leads to varied success with different pulsed voltage electroporation protocols with predetermined voltages. The ability to limit heating of cells across electrodes can increase the effectiveness of any given electroporation voltage pulsing protocol. For example, prior art teaches the utilization of an array of six needle electrodes utilizing a predetermined voltage pulse across opposing electrode pairs. This situation sets up a centralized pattern during an electroporation event in an area where congruent and intersecting overlap points develop. Excessive heating of cells and tissue along electroporation path will kill the cells, and limit the effectiveness of the protocol. However, symmetrically arranged needle electrodes without opposing pairs can produce a decentralized pattern during an electroporation event in an area where no congruent electroporation overlap points can develop.

[0147] Controlling the current flow between electrodes allows one to determine the relative heating of cells. Thus, it is the current that determines the subsequent effectiveness of any given pulsing protocol, and not the voltage across the electrodes. Predetermined voltages do not produce predetermined currents, and prior art does not provide a means to determine the exact dosage of current, which limits the usefulness of the technique. Thus, controlling an maintaining the current in the tissue

between two electrodes under a threshold will allow one to vary the pulse conditions, reduce cell heating, create less cell death, and incorporate macromolecules into cells more efficiently when compared to predetermined voltage pulses.

[0148] One embodiment of the present invention to overcome the above problem by providing a means to effectively control the dosage of electricity delivered to the cells in the inter-electrode space by precisely controlling the ionic flux that impinges on the conduits in the cell membranes. The precise dosage of electricity to tissues can be calculated as the product of the current level, the pulse length and the number of pulses delivered. Thus, a specific embodiment of the present invention can deliver the electroporative current to a volume of tissue along a plurality of paths without, causing excessive concentration of cumulative current in any one location, thereby avoiding cell death owing to overheating of the tissue.

[0149] Although not wanting to be bound by theory, the nature of the voltage pulse to be generated is determined by the nature of tissue, the size of the selected tissue and distance between electrodes. It is desirable that the voltage pulse be as homogenous as possible and of the correct amplitude. Excessive field strength results in the lysing of cells, whereas a low field strength results in reduced efficacy of electroporation. Some electroporation devices utilize the distance between electrodes to calculate the electric field strength and predetermined voltage pulses for electroporation. This reliance on knowing the distance between electrodes is a limitation to the design of electrodes. Because the programmable current pulse controller will determine the impedance in a volume of tissue between two electrodes, the distance between electrodes is not a critical factor for determining the appropriate electrical current pulse. Therefore, an alternative embodiment of a needle electrode array design would be one that is non-symmetrical. In addition, one skilled in the art can imagine any number of suitable symmetrical and non-symmetrical needle electrode arrays that do not deviate from the spirit and scope of the invention. The depth of each individual electrode within an array and in the desired tissue could be varied with comparable results. In addition, multiple injection sites for the macromolecules could be added to the needle electrode array.

XII. Restriction Enzymes

[0150] In some embodiments of the present invention, a linear DNA fragment is generated by restriction enzyme digestion of a parent DNA molecule. Examples of restriction enzymes are provided below.

Name	Recognition Sequence
AatII	GACGTC
Acc65 I	GGTACC
Acc I	GTMKAC
Aci I	CCGC
Acl I	AACGTT
Afe I	AGCGCT
Afl II	CTTAAG
Afl III	ACRYGT
Age I	ACCGGT
Ahd I	GACNNNNNGTC
Alu I	AGCT
Alw I	GGATC
AlwN I	CAGNNNCTG
Apa I	GGGCCC
ApaL I	GTGCAC
Apo I	RAATTY
Asc I	GGCGCGCC
Ase I	ATTAAT
Ava I	CYCGRG
Ava II	GGWCC
Avr II	CCTAGG
Bae I	NACNNNNGTAPyCN
BamH I	GGATCC
Ban I	GGYRCC
Ban II	GRGCYC
Bbs I	GAAGAC
Bbv I	GCAGC
BbvC I	CCTCAGC
Bcg I	CGANNNNNNTGC
BciV I	GTATCC
Bcl I	TGATCA
Bfa I	CTAG
Bgl I	GCCNNNNNGGC
Bgl II	AGATCT
Blp I	GCTNAGC
Bmr I	ACTGGG
Bpm I	CTGGAG
BsaA I	YACGTR
BsaB I	GATNNNNATC

BsaH I	GRCGYC
Bsa I	GGTCTC
BsaJ I	CCNNGG
BsaW I	WCCGGW
BseR I	GAGGAG
Bsg I	GTGCAG
BsiE I	CGRYCG
BsiHKA I	GWGCWC
BsiW I	CGTACG
Bsl I	CCNNNNNNNGG
BsmA I	GTCTC
BsmB I	CGTCTC
BsmF I	GGGAC
Bsm I	GAATGC
BsoB I	CYCGRG
Bsp1286 I	GDGCHC
BspD I	ATCGAT
BspE I	TCCGGA
BspH I	TCATGA
BspM I	ACCTGC
BsrB I	CCGCTC
BsrD I	GCAATG
BsrF I	RCCGGY
BsrG I	TGTACA
Bsr I	ACTGG
BssH II	GCGCGC
BssK I	CCNGG
Bst4C I	ACNGT
BssS I	CACGAG
BstAP I	GCANNNNNTGC
BstB I	TTCGAA
BstE II	GGTNACC
BstF5 I	GGATGNN
BstN I	CCWGG
BstU I	CGCG
BstX I	CCANNNNNNTGG
BstY I	RGATCY
BstZ17 I	GTATAC
Bsu36 I	CCTNAGG
Btg I	CCPuPyGG
Btr I	CACGTG
Cac8 I	GCNNGC
Cla I	ATCGAT
Dde I	CTNAG
Dpn I	GATC
Dpn II	GATC
Dra I	TTTAAA
Dra III	CACNNNGTG

Drd I	GACNNNNNNGTC
Eae I	YGGCCR
Eag I	CGGCCG
Ear I	CTCTTC
Eci I	GGCGGA
EcoN I	CCTNNNNNAGG
EcoO109 I	RGGNCCY
EcoR I	GAATTC
EcoR V	GATATC
Fau I	CCCGCNNNN
Fnu4H I	GCNGC
Fok I	GGATG
Fse I	GGCCGGCC
Fsp I	TGCGCA
Hae II	RGCGCY
Hae III	GGCC
Hga I	GACGC
Hha I	GCGC
Hinc II	GTYRAC
Hind III	AAGCTT
Hinf I	GANTC
HinP1 I	GCGC
Hpa I	GTTAAC
Hpa II	CCGG
Hph I	GGTGA
Kas I	GGCGCC
Kpn I	GGTACC
Mbo I	GATC
Mbo II	GAAGA
Mfe I	CAATTG
Mlu I	ACGCGT
Mly I	GAGTCNNNNN
Mnl I	CCTC
Msc I	TGGCCA
Mse I	TTAA
Msl I	CAYNNNNRTG
MspA1 I	CMGCKG
Msp I	CCGG
Mwo I	GCNNNNNNNGC
Nae I	GCCGGC
Nar I	GGCGCC
Nci I	CCSGG
Nco I	CCATGG
Nde I	CATATG
NgoMI V	GCCGGC
Nhe I	GCTAGC
Nla III	CATG
Nla IV	GGNNCC

Not I	GCGGCCGC
Nru I	TCGCGA
Nsi I	ATGCAT
Nsp I	RCATGY
Pac I	TTAATTAA
PaeR7 I	CTCGAG
Pci I	ACATGT
PflF I	GACNNNGTC
PflM I	CCANNNNNTGG
PleI	GAGTC
Pme I	GTTTAAAC
Pml I	CACGTG
PpuM I	RGGWCCY
PshA I	GACNNNNGTC
Psi I	TTATAA
PspG I	CCWGG
PspOM I	GGGCCC
Pst I	CTGCAG
Pvu I	CGATCG
Pvu II	CAGCTG
Rsa I	GTAC
Rsr II	CGGWCCG
Sac I	GAGCTC
Sac II	CCGCGG
Sal I	GTCGAC
Sap I	GCTCTTC
Sau3A I	GATC
Sau96 I	GGNCC
Sbf I	CCTGCAGG
Sca I	AGTACT
ScrF I	CCNGG
SexA I	ACCWGGT
SfaN I	GCATC
Sfc I	CTRYAG
Sfi I	GGCCNNNNNGGCC
Sfo I	GGCGCC
SgrA I	CRCCGGYG
Sma I	CCCGGG
Sml I	CTYRAG
SnaB I	TACGTA
Spe I	ACTAGT
Sph I	GCATGC
Ssp I	AATATT
Stu I	AGGCCT
Sty I	CCWWGG
Swa I	ATTTAAAT
Taq I	TCGA
Tfi I	GAWTC

Tli I	CTCGAG
Tse I	GCWGC
Tsp45 I	GTSAC
Tsp509 I	AATT
TspR I	CAGTG
Tth111 I	GACNNNGTC
Xba I	TCTAGA
Xcm I	CCANNNNNNNNNTG G
Xho I	CTCGAG
Xma I	CCCGGG
Xmn I	GAANNNNTTC

[0151] The term "restriction enzyme digestion" of DNA as used herein refers to catalytic cleavage of the DNA with an enzyme that acts only at certain locations in the DNA. Such enzymes are called restriction endonucleases, and the sites for which each is specific is called a restriction site. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors, and other requirements as established by the enzyme suppliers are used. Restriction enzymes commonly are designated by abbreviations composed of a capital letter followed by other letters representing the microorganism from which each restriction enzyme originally was obtained and then a number designating the particular enzyme. In general, about 1 μg of plasmid or DNA fragment is used with about 1-2 units of enzyme in about 20 μl of buffer solution. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation of about 1 hour at 37°C is ordinarily used, but may vary in accordance with the supplier's instructions. After incubation, protein or polypeptide is removed by extraction with phenol and chloroform, and the digested nucleic acid is recovered from the aqueous fraction by precipitation with ethanol. Digestion with a restriction enzyme may be followed with bacterial alkaline phosphatase hydrolysis of the terminal 5′ phosphates to prevent the two restriction cleaved ends of a DNA fragment from "circularizing" or forming a closed loop that would impede insertion of another DNA fragment at the restriction site. Unless otherwise stated, digestion of plasmids is not followed by 5′ terminal dephosphorylation. Procedures and reagents for dephosphorylation are conventional as described in the art.

EXAMPLES

[0152] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLE 1

CONSTRUCTION OF DNA VECTORS AND METHODS IN ANIMAL SUBJECT

[0153] In order to increase lean body mass, decrease body fat proportions, increase bone density, and improve bone healing rate, it was first necessary to design several GHRH expression constructs. Briefly, the plasmid vectors contained the muscle specific synthetic promoter SPc5-12 (Li et al., 1999) attached to a wild type or analog porcine GHRH. The analog GHRH sequences were generated by site directed mutagenesis as described in methods section. Nucleic acid sequences encoding GHRH or analog were cloned into the BamHI/ HindIII sites of pSPc5-12 plasmid, to generate pSP-GHRH. Other elements contained in the plasmids include a 3' untranslated region ("3'UTR") (SEQID No: 8) of growth hormone and an SV40 3'UTR from pSEAP-2 Basic Vector as described in the methods section. The unique nucleic acid sequences for the constructs used are shown in Figure 1.

[0154] **DNA constructs:** Plasmid vectors containing the muscle specific synthetic promoter SPc5-12 (SeqID No: 7) were previously described (Li et al., 1999). Wild type and mutated porcine GHRH cDNAs were generated by site directed mutagenesis of GHRH cDNA (SeqID No: 9) (Altered Sites II *in vitro* Mutagenesis System, Promega, Madison, WI), and cloned into the BamHI/ Hind III sites of pSPc5-12, to generate pSP-wt-GHRH (SeqID No: 15), or pSP-HV-GHRH (SeqID No: 11), respectively. The 3' untranslated region (3'UTR) of growth hormone was cloned downstream of GHRH cDNA. The resultant plasmids contained mutated coding region

for GHRH, and the resultant amino acid sequences were not naturally present in mammals. Although not wanting to be bound by theory, the effects on increased bone density, and increased healing rate of bone in the animals are determined ultimately by the circulating levels of mutated hormones. Several different plasmids that encoded different mutated amino acid sequences of GHRH or functional biological equivalent thereof are as follows:

Plasmid	Encoded Amino Acid Sequence
wt-GHRH	YADAIFTNSYRKVLGQLSARKLLQDIMSRQQGERNQEQGA-OH (SeqID No: 10)
HV-GHRH	HVDAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH (SeqID No: 1)
TI-GHRH	YIDAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH (SeqID No: 2)
TV-GHRH	YVDAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH (SeqID No: 3)
15/27/28-GHRH	YADAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH (SeqID No: 4)

[0155] In general, the encoded GHRH or functional biological equivalent thereof is of formula:

$-X_1-X_2-$DAIFTNSYRKVL$-X_3-$QLSARKLLQDI$-X_4-X_5-$RQQGERNQEQGA-OH
(SeqID No: 6)

wherein: X_1 is a D-or L-isomer of an amino acid selected from the group consisting of tyrosine ("Y"), or histidine ("H"); X_2 is a D-or L-isomer of an amino acid selected from the group consisting of alanine ("A"), valine ("V"), or isoleucine ("I"); X_3 is a D-or L-isomer of an amino acid selected from the group consisting of alanine ("A") or glycine ("G"); X_4 is a D-or L-isomer of an amino acid selected from the group consisting of methionine ("M"), or leucine ("L"); X_5 is a D-or L-isomer of an amino acid selected from the group consisting of serine ("S") or asparagine ("N").

[0156] Another plasmid that was utilized included the pSP-SEAP construct (SeqID No: 16) that contains the SacI/ HindIII SPc5-12 fragment, SEAP gene and SV40 3'UTR from pSEAP-2 Basic Vector (Clontech Laboratories, Inc.; Palo Alto, CA).

[0157] The plasmids described above do not contain polylinker, IGF-I gene, a skeletal alpha-actin promoter or a skeletal alpha actin 3' UTR /NCR. Furthermore, these plasmids were introduced by muscle injection, followed by *in vivo* electroporation, as described below.

[0158] In terms of "functional biological equivalents", it is well understood by the skilled artisan that, inherent in the definition of a "biologically functional equivalent" protein, polypeptide, and/or polynucleotide, is the concept that there is a limit to the number of changes that may be made within a defined portion of the molecule while retaining a molecule with an acceptable level of equivalent biological activity. Functional biological equivalents are thus defined herein as those proteins (and polynucleotides) in selected amino acids (or codons) that may be substituted. A peptide comprising a functional biological equivalent of GHRH is a polypeptide that has been engineered to contain distinct amino acid sequences while simultaneously having similar or improved biologically activity when compared to GHRH. For example, one biological activity of GHRH is to facilitate growth hormone ("GH") secretion in the subject.

[0159] **Electroporation devices.** A BTX T820 generator (BTX, division of Genetronics Inc., CA) was used to deliver square wave pulses in all experiments. Voltage conditions of 100-200 V/cm, 6 pulses, 60 milliseconds per pulse were used. Caliper and needle electrodes (BTX, division of Genetronics Inc., CA) were used to deliver *in vivo* electric pulses. The plate (caliper) electrodes consisted of 1.5 cm square metallic blocks mounted on a ruler, so the distance between the plates could be easily assessed; the 6-needle device consists of a circular array (1 cm diameter) of six equally spaced filled 21-gauge needles mounted on a non-conductive material. The 3-needle device consists of two filled and one cannular needle, the last one being used both as an electrode and to deliver the plasmid. All needles were 2 cm in length. In all injections the needles were completely inserted into the muscle.

[0160] A skilled artisan recognizes that any similar electroporation device and parameters may be used in the present invention so long as the device delivers the nucleic acid sequence to the cell, tissue, or organism.

[0161] **Intramuscular injection of plasmid DNA in porcine.** Two- to three-week-old hybrid barrows (Yorkshire x Landrace x Hampshire x Duroc)(Huntsville, TX), 4-5 kg in weight, or Yorkshire x Landrace pigs were used in the secreted embryonic alkaline phosphatase studies (n = 3 /group). For the GHRH plasmid studies, time-pregnant sows (Yorkshire x Landrace) were brought three weeks before the scheduled parturition date to the Children Nutrition Research Center at Baylor College of Medicine. Piglets were born in the facility. Piglets were assigned randomly to one of the experimental (n = 2 pigs/ group/ series) or controls (n = 3) groups. All experiments were repeated three times. The animals were suckled for the first 21 days and then individually housed with ad-lib access to water. For GHRH studies, after weaning, pigs were fed a 24% protein diet (Producers Cooperative Association, Bryan, TX) at 6% of their body weight daily. The animals were weighed twice a week, at the same time of day, and the amount of feed was subsequently determined. Animals were maintained in accordance with NIH Guide, USDA and Animal Welfare Act guidelines, and approved by the Baylor College of Medicine IACUC.

[0162] Endotoxin-free plasmid (Qiagen Inc., Chatsworth, CA, USA) preparations were diluted in PBS, pH 7.4 to 1 mg/ml. Plasmid DNA was injected through the intact skin into the semitendinosus or the longissimus dorsi muscle using a 21 g needle. Two minutes later, external caliper electrodes or injectable electrodes (6-needle array or 3-needle array) were applied to the injected muscle, and 6 pulses of 200V/cm, 60 millisecond/pulse were applied. The polarity of the pulses was either constant or inverted between the needles.

[0163] Blood was collected by jugular puncture before plasmid injection, and at 3, 7, 14, 21, 35 and 45 days post-injection. At 50 days post-injection, animals were sacrificed and internal organs and the injected muscle were collected, weighed, frozen in liquid nitrogen, and stored at -80°C, or placed in 10% buffered formalin for histology.

[0164] Although *in vivo* electroporation is the preferred method for delivering the nucleic acid constructs into the cells of the subject, suitable methods for nucleic acid delivery for transformation of an organelle, a cell, a tissue or an organism for use with the current invention are believed to include virtually any method by which a nucleic acid

(*e.g.*, DNA) can be introduced into an organelle, a cell, a tissue or an organism, as described herein or as would be known to one of ordinary skill in the art. Such methods include, but are not limited to, direct delivery of DNA such as by *ex vivo* transfection (Nabel et al., 1989; Wilson et al., 1989), by injection (U.S. Patent Nos. 5,994,624, 5,981,274, 5,945,100, 5,780,448, 5,736,524, 5,702,932, 5,656,610, 5,589,466 and 5,580,859, each incorporated herein by reference), including microinjection (Harland and Weintraub, 1985) U.S. Patent No. 5,789,215, incorporated herein by reference); by electroporation (U.S. Patent No. 5,384,253, incorporated herein by reference; (Potter et al., 1984; Tur-Kaspa et al., 1986); by calcium phosphate precipitation (Chen and Okayama, 1987; Graham and van der Eb, 1973; Rippe et al., 1990); by using DEAE-dextran followed by polyethylene glycol (Gopal, 1985); by direct sonic loading (Fechheimer et al., 1987); by liposome mediated transfection (Hafez et al., 2001; Hamm et al., 2002; Madry et al., 2001; Raghavachari and Fahl, 2002; Wiethoff et al., 2001) and receptor-mediated transfection (Wu and Wu, 1988a; Wu and Wu, 1988b); by microprojectile bombardment (PCT Application Nos. WO 94/09699 and 95/06128; U.S. Patent Nos. 5,610,042; 5,322,783 5,563,055, 5,550,318, 5,538,877 and 5,538,880, and each incorporated herein by reference); by agitation with silicon carbide fibers ((Johnson et al., 1992); U.S. Patent Nos. 5,302,523 and 5,464,765, each incorporated herein by reference); by *Agrobacterium*-mediated transformation (U.S. Patent Nos. 5,591,616 and 5,563,055, each incorporated herein by reference); by PEG-mediated transformation of protoplasts (Omirulleh et al., 1993); U.S. Patent Nos. 4,684,611 and 4,952,500, each incorporated herein by reference); by desiccation/inhibition-mediated DNA uptake (Potrykus et al., 1985), and any combination of such methods. Through the application of techniques such as these, organelle(s), cell(s), tissue(s) or organism(s) may be stably or transiently transformed.

[0165] **Porcine plasma IGF-I and insulin concentrations.** Porcine IGF-I was measured by heterologous human radioimmunometric assay (Diagnostic System Lab., Webster, TX). The sensitivity of the assay was 0.8 ng/ml; intra-assay and inter-assay variation was 3.4% and 4.5%, respectively. Porcine insulin was measured using a heterologous human radioimmunoassay (Linco Research Inc., St. Charles, MO). The

sensitivity level of the assay was 2 μU/ml; intra-assay and inter-assay variation was 3.2% and 3.9% respectively.

[0166] **Body composition data.** Weights were measured on the same calibrated scales (certified to have an accuracy to ± 0.2 kg and a coefficient of variation of 0.3%) throughout the study, twice a week. Body composition measurements were performed *in vivo*, 50 days after birth. The piglets were anesthetized using a combination of xylazine (15 mg/kg) and ketamine (2 mg/kg) and the total body content of fat, percent of fat, non-bone lean tissue mass and bone mineral content was measured by dual-energy x-ray absorptiometry (Hologic QDR-2000, Waltham, Massachusetts) ("DEXA") (Toner et al., 1996). Total body potassium was measured in a potassium chamber ("K40") using a whole body detector (Cohn et al., 1984).

[0167] **Statistics** Data are analyzed using STATISTICA analysis package (StatSoft, Inc. Tulsa, OK). Values shown in the figures are the mean ± s.e.m. Specific P values were obtained by comparison using ANOVA. A $P < 0.05$ was set as the level of statistical significance.

EXAMPLE 2

CONSTITUTIVE GHRH SYSTEM IN VIVO

[0168] To test the constitutive GHRH system *in vivo*, 7.5 micrograms of pSP-GHRH or functional biological equivalents (FIGURE 1) were delivered into SCID mice. All GHRH analog sequences were obtained by site directed mutagenesis of the porcine wild type sequence. Groups of five mice were injected with 7.5 micrograms of plasmid expressing either one of the GHRH analogs, or a pSP-beta-galactosidase as control. At 45 days post-injection, animals were analyzed by PIXImus (Draghia-Akli et al., 2002)(DEXA for mice), sacrificed, blood and organs were collected and weighed. At the end of the experiment, the TI-GHRH and HV-GHRH animals were significantly bigger than controls (FIGURE 2). The body composition of the injected SCID mice was also altered. At 45 days post-injection, animals that were injected with the TI mutant had a significant increase in lean body mass of 11% versus controls, $p < 0.036$. The HV-GHRH injected animals had a significant increase of the lean body mass of almost 5% (FIGURE 3). All

GHRH injected groups had larger bone areas than the control animals, up to 10.7%, in the TI-GHRH injected group, $p < 0.027$. (FIGURE 4). At 14 and 28 days post-injection, blood was collected and IGF-I levels were measured (FIGURE 5). All GHRH injected groups had significantly increased IGF-I levels compared with control animals, up to $p < 0.005$. Some groups developed neutralizing antibodies, and in these cases the IGF-I levels dropped at the second time point. The animals injected with TI-GHRH did not develop any antibodies, and their GHRH expression continued to 45 days, correlating with significant changes in their body composition.

EXAMPLE 3

INDUCIBLE GHRH SYSTEM IN VITRO

[0169] To test the inducible GHRH system *in vitro*, primary chicken myoblasts were transfected as described previously (Bergsma et al., 1986; Draghia-Akli et al., 1997) with 4 micrograms of a mixture of the GHRH/GeneSwitch® plasmids, pGR1774 (inducible GHRH)/pGS1633 (Gene Switch®) in a 10:1 w/w ratio, which gave the best overall expression in skeletal muscle cells, and cells were allowed to differentiate into post-mitotic myotubes. At 24 and 48 hours after transfection, cells were washed in PBS, and MFP was added, where indicated, to the culture media. Media and cells were harvested 72 hours post-differentiation. 20μg of total RNA was DNase I treated, size separated in 1.5% agarose-formaldehyde gel and transferred to nylon membrane. The membranes were hybridized with a specific GHRH cDNA probe ^{32}P-labeled by random priming. Negative controls were cells transfected by the GeneSwitch® and GHRH plasmids, but not treated with MFP, or cells transfected by the inducible GHRH plasmid alone. The positive control was cells transfected by a constitutively expressed GHRH plasmid that was driven by a synthetic muscle-specific promoter ("SP-GHRH"). GHRH transcripts of the expected size of 0.35kb were only observed in cells transfected with the GeneSwitch®/inducible GHRH plasmids and treated with MFP, and in cells transfected with the positive control (FIGURE 7). No GHRH transcripts were detected in cells not treated with MFP or in cells transfected by the inducible GHRH plasmid alone.

EXAMPLE 4

GHRH/GENESWITCH® SYSTEM IN VIVO—IMPROVED BODY COMPOSITION AND FAT BODY MASS/TOTAL WEIGHT

[0170] For the *in vivo* experiments, the plasmids for the GHRH/GeneSwitch® system were delivered to the muscles of SCID mice. The left tibialis anterior muscle was injected with 10 μg of a 10:1 mixture of pGR1774/pGS1633, followed by caliper electroporation (Draghia-Akli et al., 1999). At twenty-one days post-injection, animals were injected inter perineum ("i.p.") with 250 micrograms/kg of MFP for 3 days. On the fourth day, the animals were bled and serum was used to measure IGF-I levels. Mouse IGF-I was measured by heterologous, 100% cross-reacting rat radioimmunoassay. The sensitivity of the assay was 0.8 ng/ml; intra-assay and inter-assay variation was 3.4% and 4.5% respectively. Following administration of MFP for 4 consecutive days, IGF-I levels increased from 1100.86 ± 33.67 ng/ml to 1797.28 ± 164.96 ng/ml ($p < 0.0005$). Significant changes in the IGF-I levels were seen when the MFP group was compared with the control group 1086.78 ± 65.34 ng/ml, $p < 0.0006$ (animals that received a control beta-galactosidase plasmid), 1171.79 ± 42 ng/ml, $p < 0.001$ (animals that received the GHRH/GeneSwitch® plasmids but were not dosed with MFP). Upon repeated administration of MFP to the animals using the same protocol followed by recovery to background 7 days over 149 days, serum IGF-I levels rose repeatedly 1.1-1.7 fold over the uninjected controls (FIGURE 8). Animals induced with MFP had statistically significant higher IGF-I levels.

[0171] Body weight was similar for all of the groups during the first 125 days of the study (FIGURE 9). However, from day 125 to day 149, mice were dosed with MFP every day. A 7.5% increased body weight was observed in the chronically MFP-induced GHRH/GeneSwitch® animals, averaging 31.84 ± 0.12g ($p < 0.027$), compared with β-gal controls, 29.62 ± 0.98g, and with animals that were not induced with MFP, 30.53 ± 0.59g. All values are average ± SEM. Organs (lungs, heart, liver, kidney, stomach, intestine, adrenals, gonads, brain) were collected and weighed. No organomegaly or associated pathology was observed in any of the animals. Pituitary glands were dissected within the first minutes post-mortem, and weighed. Pituitary weight/ total body weight (FIGURE 10) increased upon chronic stimulation of the GHRH/GeneSwitch® by 20% ($7.35 \pm 0.31 \times 10^{-5}$), compared with β-gal controls ($6.13 \pm 0.46 \times 10^{-5}$), and animals not dosed with MFP

($6.3 \pm 0.22 \times 10^{-5}$), $p < 0.035$. There was no significantly statistical difference between the β-gal injected animals and animals that were injected with the GHRH/GeneSwitch® system, but not given MFP. Although not to be bound by theory, the increase in pituitary weight was probably due to somatotrophs hypertrophy, as it is known that GHRH is capable of stimulating the synthesis/secretion of GH from the anterior pituitary and has a specific hypertrophic effect on somatotrophs (Morel et al., 1999; Murray et al., 2000). At the end of the experiment, body composition was analyzed *in vivo*, by dual-energy x-ray absorptiometry ("DEXA"), using a high resolution PIXImus scanner. Body composition studies by PIXImus (total body fat, non-bone lean tissue mass and bone mineral area, content and density) showed significant changes in chronically MFP induced animals injected with the GHRH/GeneSwitch® system. Lean body mass (non-bone) (FIGURE 11) increased by 2.5% in GHRH/GeneSwitch® animals + MFP (87.44 ± 0.65%, versus β-gal 84.94 ± 0.6%, and no MFP animals 84.88 ± 0.3%), $p < 0.022$. Fat mass (FIGURE 12) decreased by 2 % in GHRH/GeneSwitch® animals (12.59 ± 0.62%, versus β-gal 14.57 ± 0.75%, and no MFP animals 15.09 ± 0.3%), $p < 0.05$.

EXAMPLE 5

GHRH/GENESWITCH® IN VIVO—INCREASED BONE AREA AND MINERAL CONTENT

[0172] One aspect of the present invention involves demonstrating that the introduction of mifepristone-inducible heterologous nucleic acid sequences encoding GHRH or functional biological equivalent thereof into the cells of subjects can lead to high levels of circulating hormones (Mir et al., 1999), without the disadvantages (*e.g.* high production costs, safety concerns with the virus backbone, or *ex vivo* manipulation) associated with viral vector delivery or organoids (Barr and Leiden, 1991; Dhawan et al., 1991; Draghia-Akli et al., 1999). In addition, the invention must demonstrate that animal growth and body composition can be efficiently regulated by mifepristone following *in vivo* electroporation of the GeneSwitch® technology (*i.e.* mifepristone-inducible heterologous nucleic acid sequences encoding GHRH or functional biological equivalent thereof) into skeletal muscle of the subject, as schematically diagrammed in FIGURE 6. Enhanced biological potency, delivery and proper gene expression regulation was observed over 149 days post-injection, and effectively reduced the theoretical quantity of

GHRH needed to achieve physiological levels of GH secretion when compared to the recombinant GHRH therapies. Post-injected subjects did not experience any side effects from the GeneSwitch® technology therapy. For example, mice had normal biochemical profiles, and no associated pathology or organomegaly. From a functional standpoint, the IGF-I levels increased, growth was enhanced by 7.5%, and changes in body composition (*e.g.* with increased lean body mass by 2.5% and decreased fat by 2%) were observed following chronic induction of the GHRH/Gene Switch system. In addition, bone mineral density increased by 6%, and the stimulation of GHRH on bone metabolism were even more remarkable. Although not to be bound by theory, the observed pituitary hypertrophy was indicative that ectopic expression of myogenic GHRH plasmids operates through the natural GH axis (stimulation of GH synthesis and secretion at the pituitary level). This long-lasting regulated therapy has the potential to replace classical GH therapy regimens and may stimulate the GH axis in a more physiologically appropriate manner. It is known that GHRH stimulates bone formation (Dubreuil et al., 1996), and the described GeneSwitch® therapy may be used to promote post-fracture bone growth.

[0173] Upon chronic stimulation of the GHRH/ GeneSwitch® system, significant changes occurred in bone area (FIGURE 13), that increased by 7 %, (12.81 ± 0.14 cm^2, versus B=Beta β-gal injected controls 11.98 ± 0.3 cm^2, or no MFP animals 12.07 ± 0.26 cm^2), $p < 0.0006$, bone mineral content (FIGURE 14) increased by 14.6% (0.755 ± 0.012g, versus β-gal injected controls 0.659 ± 0.019g, or no MFP animals 0.694 ± 0.023 cm^2), $p < 0.002$, and bone mineral density increased by 6% (0.059 ± 0.0007 g/cm^2, versus β-gal injected controls 0.056 ± 0.0009 g/cm^2, or no MFP animals 0.057 ± 0.0007 g/cm^2), $p < 0.012$. Practically, there is no overall difference between the β-gal injected animals and animals that were injected with the GHRH/ GeneSwitch®, but were not given MFP, which supports the absence of GHRH expression by the GHRH/GeneSwitch® plasmids in the absence of MFP dosing.

EXAMPLE 6

LOW VOLTAGE ELECTROPORATION INCREASES PLASMID UPTAKE AND EXPRESSION IN ADULT PIGS

[0174] Direct intra-muscular plasmid DNA injection followed by electroporation is a method for the local and controlled delivery of plasmid DNA into skeletal muscle. It has the advantage that is uses low plasmid quantities (as low as 0.1 mg in pigs), rather than the high quantities typically used with passive delivery modalities. Although not wanting to be bound by theory, the mechanism of the increased plasmid uptake by electroporation probably occurs through newly created membrane pores with or without protein active transport. It has been shown that the degree of permeabilization of the muscle cells is dependent on the electric field intensity, length of pulses, shape and type of electrodes (Bureau et al., 2000; Gilbert et al., 1997), and cell size (Somiari et al., 2000). Classical electrode configuration, plates or a pair of wire electrodes placed 4 mm apart were shown to be effective in rodents, but in large mammals as pigs or humans the increased resistance of the skin, the thickness of the subcutaneous fat tissue, and the concern for tissue damage if the intensity of the electric field would be proportionally increased, make these types of electrodes unpractical. The porcine muscle fibers are quite large and consequently more suitable for electropermeabilization than rodent muscle. Data provided herein indicate that a single injection of an optimum dosage of plasmid followed by electroporation with intramuscular applicators is sufficient to produce therapeutic plasma hormone levels in a large mammal with biologically significant effects on the body fat distribution and lean body mass of the subject.

[0175] External caliper electrodes and injectable electrodes were evaluated to determine the type of electrode needed to achieve a physiologically relevant level of a secreted reporter protein in 4-5 kg hybrid pigs. Reporter vectors expressing secreted embryonic alkaline phosphatase ("SEAP") were used in these studies at a dose of 2 mg pSP-SEAP/animal. Six-needle and 3-needle array electrodes were compared with standard caliper electrodes (FIGURE 15). Conditions of 6 pulses, 200V/cm, 60 milliseconds/pulse, previously tested as being the most effective in pigs (Draghia-Akli et al., 1999) were applied in all tests. For the three-needle electrode, three pulses were applied in one direction, then the polarity was changed and the next three pulses were delivered in the opposite direction. SEAP values were measured at day 0, day 3 and day 7 post-injection. Seven days post-injection, the SEAP levels were 9.33 ± 2.26 ng/(ml·kg) in plasmid-injected and caliper electroporated animals, compared to 0.02 ± 0.005 ng/(ml·kg)

in vehicle-injected animals. Using the 3-needle and 6-needle arrays, a 12.4 and 19 fold increase in SEAP values was obtained compared to caliper delivery (116.07 ± 44.36 ng/(ml·kg), and 177.41 ± 18.44 ng/(ml·kg), respectively). When using the same number of pulses, but lower voltage (100V/cm), and the 6-needle electrodes, the average SEAP increased to 144.64 ± 11.82 ng/(ml·kg) after seven days. When longissimus dorsi and semitendinosus muscles were injected using similar conditions, expression in the semitendinosus muscle was slightly higher. Skin and muscle from the injected pigs were collected at the end of the experiment (at 50 days post-injection) and histologically analyzed. At 100-200 V/cm used in the injectable electrodes experiments, no skin or muscle damage was seen for any of the needle-type electrodes used.

EXAMPLE 7

INCREASED EFFICIENCY USING NEEDLE-TYPE ELECTROPORATION DELIVERY FOR THERAPEUTIC PROTEINS

[0176] Not wanting to be bound by theory, growth hormone releasing hormone ("GHRH") stimulates the production and release from the anterior pituitary of growth hormone ("GH"), which in turn stimulates the production of IGF-I from the liver and other target organs (Frohman et al., 1968). In previous studies (Draghia-Akli et al., 1999), young pigs weighing 4-5 kg, were injected with 10 mg myogenic vector expressing a mutated form of GHRH, stable to proteases ("pSP-HV-GHRH") and electroporated using a caliper electrode.

[0177] The present invention involves determination of the best age for treatment of young pigs. Groups of 2 piglets were injected with 2 mg pSP-HV-GHRH using the 6-needle array electrodes at different time points: birth, 7, 14 and 21 days of age (FIGURE 16). Each animal received one injection. The group injected at 14 days of age demonstrated the best weight gain, (statistically significant and different from PBS controls (n=3) at every time point (final weights: 25.8 ± 1.5 kg versus 19.7 ± 0.03 kg, $p < 0.013$)). The next best group was injected at 7 days of age, and weighed 21.9 ± 1.5 kg at age 50 days, $p < 0.02$.

[0178] In a parallel study, the reduction in the plasmid quantity needed to achieve improved growth and changes in the metabolic and hormonal profile of pigs was

explored. Groups of two piglets each (Yorkshire x Landrace) were injected at 10 days of age with pSP-HV-GHRH (3 mg, 1 mg, 0.1 mg), and electroporated using a 6-needle array electrode (FIGURE 17). The group injected with 0.1 mg of plasmid had the greater weight gain, with statistically significant differences to controls (n=3) to 50 days of age (22.4 ± 0.8 kg versus 19.7 ± 0.03 kg, $p < 0.012$). One animal in the group injected at 21 days and one animal injected with the highest plasmid dose (3 mg) developed neutralizing antibodies against the mutated HV-GHRH and showed significant reduced rates of weight gain (at 50 days post-injection 15.6 kg and 15.95 kg, respectively, versus more than 21 kg for the paired animal in the same treatment group). No other group developed neutralizing antibodies. Thus, the minimal plasmid dosage (0.1 mg) and injection at optimum age using the 6-needle electrodes resulted in the best growth performances. It is noteworthy that in previous studies the inventors used 100-fold less, *i.e.*, 10 mg pSP-HV-GHRH with the caliper electrodes to produce similar changes.

[0179] An indication of increased systemic levels of GHRH and GH is an increase in serum IGF-I concentration. The level of serum IGF-I started to rise at 3 days post-injection in pigs that received the 0.1 and 1 mg doses of pSP-HV-GHRH (FIGURE 18). By 35 days after the injection (age of animals: 45 days), serum IGF-I concentrations were approximately 10-fold higher in pigs injected with 0.1 mg and 7-fold higher in pigs injected with 1 mg plasmid compared with controls ($p < 0.007$ and $p < 0.04$ respectively).

[0180] In pSP-HV-GHRH injected pigs, under optimum conditions (Table I) serum urea decreased (8.36±1.33 to 9.67±1.27 mg/ml in pSP-HV-GHRH injected pigs versus 11.14±1.9 mg/ml in controls, respectively ($p<0.05$), indicating decreased amino acid catabolism. Serum glucose levels were similar between the plasmid pSP-HV-GHRH injected pigs and controls; insulin levels were normal and within the control range.

Table 1
The plasma metabolic profile of pSP-HV-GHRH injected and control pigs.

Group Age	Glucose (mg/ml)	Urea (mg/ml)	Creatine (mg/ml)	Total Protein (g/dl)
Day 0	125.82 ± 5.64	8.36 ± 1.33 *p < 0.01*	0.85 ± 0.05	4.81 ± 0.11
Day 7	122.43 ± 5.05	9.43 ± 1.67 *p< 0.03*	0.87 ± 0.04	5.21 ± 0.19
Day 14	129.54 ± 6.39	9.62 ± 1.72 *p < 0.02*	1.00 ± 0.04	5.22 ± 0.20
Day 21	110.25 ± 5.02	13.83 ± 1.2	0.93 ± 0.05	4.81 ± 0.21
Dose				
3 mg	111.07 ± 3.88	10.50 ± 1.87 *p < 0.05*	0.9 ± 0.09	3.89 ± 0.16
1 mg	121.63 ± 2.93	9.44 ± 1.07 *p < 0.02*	0.81 ± 0.06	4.11 ± 0.11
0.1 mg	120.73 ± 2.53	9.67 ± 1.27 *p < 0.02*	0.95 ± 0.05	4.05 ± 0.19
Control	119.77 ± 3.67	12.81 ± 2.01	0.98 ± 0.08	4.00 ± 0.11

[0181] The fact that these animals have a normal carbohydrate metabolism is very important, as most livestock and/or patients under recombinant GH therapy develop impaired glucose metabolism and insulin resistance. Creatinine concentration (a measure of kidney function) was normal in all animals. Pigs that developed antibodies to GHRH showed a tendency to increased urea levels and decreased glucose levels.

[0182] Body composition studies by dual-energy x-ray absorptiometry (total body fat, non-bone lean tissue mass and bone mineral content), K40 potassium (lean body mass) and carcass neutron activation analysis (nitrogen) showed a proportional increase of all internal organs in GHRH injected animals (heart, lung, liver, spleen, brain, adrenals, stomach, kidney, pancreas, intestine). Nevertheless, the final body composition was different: animals injected with pSP-HV-GHRH at different ages gained proportionally less fat than controls and were leaner at the end of the study (4.34 ± 0.04 g of fat gained/ kg of fat free mass gained per day for injection at birth, 4.4 ± 0.04 g for injection at 7 days,

versus controls 5.63 ± 0.34 g, $p < 0.05$). Bone mineral density was higher in animals injected at 14 days after birth, and correlates with increased efficacy of the treatment: 0.363 ± 0.005 g/cm^2 versus 0.329 ± 0.003 g/cm^2 in controls, $p < 0.004$.

[0183] Treated pigs did not experience any side effects from the therapy, had normal biochemical profiles, and had no associated pathology or organomegaly. From a functional standpoint, the increases in IGF-I levels and enhancement in growth and changes in body composition (with decreased fat deposition by 22%) were dramatic in extent. The effects of the stimulation of GHRH on bone metabolism were even more remarkable, with an increase in bone mineral density by 10%. These results indicate that ectopic expression of myogenic HV-GHRH vectors has the potential to replace classical GH therapy regimens and may stimulate the GH axis in a more physiologically appropriate manner. The HV-GHRH molecule, which displays a high degree of stability and GH secretory activity in pigs, may also be useful in human clinical medicine. However, a skilled artisan is aware that a minimal plasmid dose should be determined on a pertinent model and used in order to avoid the unwanted pathology associated with antibody development, and routine methods in the art and/or described herein teach how to achieve this goal.

[0184] The molecular techniques used to produce alterations in any conceivable encoded nucleic acid sequences are well established, and exemplified by the large number of scientific publications and patents in the field of molecular biology. Despite the accuracy of the molecular techniques used to create distinctive nucleic acid sequences, a skilled artisan recognizes that the expression of any given nucleic acid will influence the complex biochemistry of an entire organism. Thus, the highly predictable nature of constructing unique nucleic acid sequences must not be confused with unknown facts of an associated biological effect.

[0185] The invention described herein involves the utilization of several distinctive GHRH or analog nucleic acid sequences. Based upon the current understanding of protein-protein interactions, it is neither obvious nor possible to accurately speculate upon the *in vivo* parameters (*e.g.* half life, efficacy, post-translational modifications, *etc.*) of a GHRH sequence that contains a point mutation which alters a

single amino acid in the polypeptide chain. As seen in the Examples provided herein, mutation of a few base pairs gave rise to GHRH mutants with significantly longer bio-availability. The endogenous GHRH has a half-life of 6-12 minutes in different species. The HV-GHRH has a half-life of 6 hours. In further analysis, the TI-GHRH (that has only two base pair difference with the HV-GHRH) has been shown to have a much higher effect *in vivo* on lean body mass than the HV-GHRH (from simple to double). This property was not evident in extensive *in vitro* studies on pituitary cell. Correspondingly, one skilled in the art would know how to perform the plasmid-mediated supplementation of GHRH or the related recombinant protein experimentation(s), characterizing variations and permutations on a unique nucleic acid sequence in a specific tissue to accurately evaluate the *in vivo* effect within a living organism. Therefore, the utilization of the distinctive nucleic acid sequence encoding GHRH or functional biological equivalent thereof or corresponding recombinant protein as a method to decrease body fat proportion and increase lean body mass could not have been predicted based on speculation.

[0186] Although not wanting to be bound by theory, it is believed that an increase in GHRH or functional biological equivalent will increase the GH levels to decrease body fat proportion and increase lean body mass. Hormones (*e.g.* GHRH and GH) often contain a complex feedback-regulated pathway. Without direct experimentation of GHRH or analogs used in gene or recombinant protein therapy, it could not have been predicted by one skilled in the art to determine which concentrations of non-native encoded sequences will yield desired results. Ideal regulation of a nucleic acid sequence encoding GHRH or functional biological equivalent thereof is further complicated by the tissue used for polynucleotide delivery, and would not have been obvious to one skilled in the art without actual experimentation with the distinctive sequence in a particular tissue. The invention described herein contains the descriptions and results of essential experimentation that explored tissue specific and inducible regulation of distinctive nucleic acid sequences that encoded GHRH or functional biological equivalent thereof, which was not obvious based upon prior art. The present invention is a significant step forward in developing non-viral therapy for large animals, including humans. In order for gene therapies to be transferred from rodents to large mammals, and ultimately to humans, it was surprising that extremely low quantities of

plasmid were effective. It is shown herein that as little as 0.1 mg plasmid delivered under the proper electroporation conditions had an important biological impact that decreases the body fat proportion, increases lean body mass ("LBM"), or both of a subject. This plasmid quantity was 100 fold lower than the theoretical one, and could not have been predicted from the relative doses used in rodents (in average 1mg/kg). Although not wanting to be bound by theory, unlike other therapies using growth factors (GH and/ or IGF-I), GHRH is stimulating the endogenous secretion of hormones, and enhancing the own bio-potential of the animal, with no adverse effects. This experimental finding cannot be theoretically predicted, as the three hormones are members of the same growth axis.

[0187] The increase in lean body mass, decrease in body fat proportions, increase in bone density, and/or increase in bone rate of healing are a consequence of the GHRH molecules present in the subjects circulation, regardless of the means of the delivery. For example, one would obtain the same effect by delivering appropriate quantities of GHRH or analog thereof, outlined in FIGURE 1, by classical recombinant protein therapy or nucleic acid transfer. Accordingly, successful plasmid-mediated supplementation of GHRH requires accurate delivery of the encoded sequences to the cells of a subject, resulting in expression of the gene product at levels appropriate to produce a biological effect. The duration of treatment will extend through the course of the disease symptoms, and possibly continuously. Since the method to deliver nucleic acid sequences to the cells of a subject is highly dependent on specific diseases and the encoded gene, it could not have been predicted by one skilled in the art which method and conditions are appropriate without laborious and failed experimentations. Thus, the preferred method of outlined for this invention is *in vivo* electroporation.

[0188] One skilled in the art readily appreciates that the present invention is well adapted to carry out the objectives and obtain the ends and advantages mentioned as well as those inherent therein. Methods, procedures, techniques, and kits described herein are presently representative of the preferred embodiments and are intended to be exemplary and are not intended as limitations of the scope. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention or defined by the scope of the invention.

[0189] All of the methods and compositions disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the methods and compositions of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the systems and compositions without departing from the concept, spirit, and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically, structurally and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope, and concept of the invention as defined by the appended claims.

CLAIMS:

1. Use of a nucleic acid expression construct that encodes a growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for decreasing body fat proportion, wherein the composition is suitable for delivery into cells of a subject.

2. A use of claim 1, wherein the nucleic acid expression construct is suitable for delivery into cells of the subject via electroporation.

3. A use of claim 1, wherein the cells of the subject are somatic cells, stem cells, or germ cells.

4. A use of claim 1, wherein the nucleic acid expression construct comprises SeqID No: 11, SeqID No: 12, SeqID No: 13, SeqID No: 14, SeqID No: 17, SeqID No: 18, SeqID No: 19, SeqID No: 20, or SeqID No: 21.

5. A use of claim 1, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

6. A use of claim 5, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.

7. A use of claim 5, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

8. A use of claim 1, wherein upon delivery of the nucleic acid expression construct into the cells of the subject, the nucleic acid expression construct is capable of initiating expression of the encoded GHRH or functional biological equivalent thereof.

9. A use of claim 8, wherein the encoded GHRH or functional biological equivalent thereof is suitable for being expressed in tissue specific cells of the subject.

10. A use of claim 9, wherein the tissue specific cells of the subject comprises muscle cells.

11. A use of claim 8, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

12. A use of claim 8, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D-or L- isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D-or L- isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

13. A use of claim12, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

14. A use of claim 1, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

15. Use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for decreasing body fat proportion in a subject wherein the composition is suitable for delivery into cells of the subject, wherein the first nucleic acid expression construct is suitable for encoding growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof, wherein the second nucleic acid expression construct has an encoding region of a regulator protein and wherein upon delivery of the molecular switch molecule into the subject, the molecular switch molecule is capable of governing activation of the regulator protein.

16. A use of claim 15, wherein the first nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

17. A use of claim 15, wherein the second nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

18. A use of claim 15, wherein the first nucleic acid expression construct encodes a polypeptide of a sequence comprising SeqID No: 1, SeqID No: 2, SeqID No: 3, or SeqID No: 4.

19. A use of claim 15, wherein the first nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

20. A use of claim 19, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.

21. A use of claim 19, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

22. A use of claim 15, wherein upon delivery of the second nucleic acid expression construct into the cells of the subject, the second nucleic acid expression construct is capable of initiating expression of the encoded regulatory protein.

23. A use of claim 22, wherein the encoded regulatory protein is suitable for expression in the cells of the subject, but is inactive.

24. A use of claim 23, wherein upon delivery of the molecular switch into the subject the molecular switch is capable of activating the regulatory protein, and wherein the molecular switch comprises mifepristone or its functional equivalent.

25. A use of claim 24, wherein the active regulatory protein is capable of initiating expression of the GHRH or functional biological equivalent encoded on the first nucleic acid sequence.

26. A use of claim 25, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

27. A use of claim 25, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D- or L- isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D- or L- isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

28. A use of claim 27, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

29. A use of claim 15, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

30. A use of claim 15, wherein the first nucleic acid expression construct comprises SeqID No: 26.

31. A use of claim 15 wherein the second nucleic acid expression construct comprises SeqID No: 27.

32. Use of a recombinant growth- hormone- releasing-hormone ("GHRH") or a biological functional equivalent thereof in the preparation of a composition for decreasing body fat proportion in a subject, wherein the composition is suitable for delivery into the subject.

33. A use of claim 32, wherein the recombinant GHRH is a biologically active polypeptide; and the recombinant functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

34. A use of claim 32, wherein the recombinant GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D- or L- isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

35. A use of claim 32, wherein the recombinant GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

36. A use of claim 32, wherein the recombinant GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

37. Use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for increasing lean body mass in a subject, wherein the composition is suitable for delivery into cells of the subject.

38. A use of claim 37, wherein the nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

39. A use of claim 37, wherein the cells of the subject are somatic cells, stem cells, or germ cells.

40. A use of claim 37, wherein the nucleic acid expression construct comprises SeqID No: 11, SeqID No: 12, SeqID No: 13, SeqID No: 14, SeqID No: 17, SeqID No: 18, SeqID No: 19, SeqID No: 20, or SeqID No: 21.

41. A use of claim 37, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

42. A use of claim 41, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.

43. A use of claim 41, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

44. A use of claim 37, wherein upon delivery of the nucleic acid expression construct into the cells of the subject, the nucleic acid expression construct is capable of initiating expression of the encoded GHRH or functional biological equivalent thereof.

45. A use of claim 44, wherein the encoded GHRH or functional biological equivalent thereof is suitable for being expressed in tissue specific cells of the subject.

46. A use of claim 45, wherein the tissue specific cells of the subject comprises muscle cells.

47. A use of claim 44, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

48. A use of claim 44, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D- or L- isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D- or L- isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

49. A use of claim 48, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

50. A use of claim 37, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

51. Use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for increasing lean body mass in a subject and wherein the composition is suitable for delivery into cells of the subject wherein the first nucleic acid expression construct is suitable for encoding growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof, wherein the second nucleic acid expression construct has an encoding region of a regulator protein and wherein upon delivery of the molecular switch molecule into the subject, the molecular switch molecule is capable of activating the regulator protein.

52. A use of claim 51, wherein the first nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

53. A use of claim 51, wherein the second nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

54. A use of claim 51, wherein the first nucleic acid expression construct encodes a polypeptide of a sequence comprising SeqID No: 1, SeqID No: 2, SeqID No: 3, or SeqID No: 4.

55. A use of claim 51, wherein the first nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

56. A use of claim 55, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.

57. A use of claim 55, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

58. A use of claim 51, wherein upon delivery of the second nucleic acid expression construct into the cells of the subject, the second nucleic acid expression construct is capable of initiating expression of the encoded regulatory protein.59.

59. A use of claim 58, wherein the encoded regulatory protein is suitable for expression in the cells of the subject, but is inactive.

60. A use of claim 59, wherein upon delivery of the molecular switch into the subject the molecular switch is capable of activating the regulatory protein, and wherein the molecular switch comprises mifepristone or its functional equivalent.

61. A use of claim 60, wherein the active regulatory protein is suitable for initiating expression of the GHRH or functional biological equivalent encoded on the first nucleic acid sequence.

62. A use of claim 51, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

63. A use of claim 51, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D- or L- isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D- or L- isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N"); or a combination thereof.

64. A use of claim 63 wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

65. A use of claim 51, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

66. A use of claim 51 wherein the first nucleic acid expression construct comprises SeqID No: 26.

67. A use of claim 51 wherein the second nucleic acid expression construct comprises SeqID No: 27.

68. Use of a recombinant growth-hormone- releasing-hormone ("GHRH") or a biological functional equivalent thereof in the preparation of a composition for increasing lean body mass in a subject, wherein the composition is suitable for delivery into the subject.

69. A use of claim 68, wherein the recombinant GHRH is a biologically active polypeptide; and the recombinant functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

70. A use of claim 68, wherein the recombinant GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARl(LLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D- or L- isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D- or L- isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

71. A use of claim 70, wherein the recombinant GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

72. A use of claim 68, wherein the recombinant GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

73. Use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for increasing bone density in a subject, wherein the composition is suitable for delivery into cells of the subject.

74. A use of claim 73, wherein the nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

75. A use of claim 73, wherein the cells of the subject are somatic cells, stem cells, or germ cells.

76. A use of claim 73, wherein the nucleic acid expression construct comprises SeqID No: 11, SeqID No: 12, SeqID No: 13, SeqID No: 14, SeqID No: 17, SeqID No: 18, SeqID No: 19, SeqID No: 20, or SeqID No: 21.

77. A use of claim 73, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

78. A use of claim 73, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.

79. A use of claim 73, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

80. A use of claim 73, wherein upon delivery of the nucleic acid expression construct into the cells of the subject, the nucleic acid expression construct is capable of initiating expression of the encoded GHRH or functional biological equivalent thereof.

81. A use of claim 80, wherein the encoded GHRH or functional biological equivalent thereof is suitable for expression in tissue specific cells of the subject.

82. A use of claim 81, wherein the tissue specific cells of the subject comprises muscle cells.

83. A use of claim 80, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

84. A use of claim 80, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-$\mathbf{X_1}$ -$\mathbf{X_2}$- DAIFTNSYRKVL-$\mathbf{X_3}$-QLS ARKLLQDI-$\mathbf{X_4}$-$\mathbf{X_5}$-RQQGERNQEQ GA- OH

wherein the formula has the following characteristics:

$\mathbf{X_1}$ is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

$\mathbf{X_2}$ is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

$\mathbf{X_3}$ is a D- or L- isomer of the amino acid alanine ("A") or glycine ("G");

$\mathbf{X_4}$ is a D- or L- isomer of the amino acid methionine ("M"), or leucine ("L");

$\mathbf{X_5}$ is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

85. A use of claim 84, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

86. A use of claim 73, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

87. Use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for increasing bone density in a subject, and wherein the composition is suitable for delivery into the cells of the subject, wherein the first nucleic acid expression construct is suitable for encoding growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof; and wherein the second nucleic acid expression construct has an encoding region of a regulator protein, wherein upon delivery of the molecular switch molecule into the subject, the molecular switch molecule is capable of activating the regulator protein.

88. A use of claim 87, wherein the first nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

89. A use of claim 87, wherein delivering step of the second nucleic acid expression construct is suitable for delivery into cells via electroporation.

90. A use of claim 87, wherein the first nucleic acid expression construct is suitable for encoding a polypeptide of a sequence comprising SeqID No: 1, SeqID No: 2, SeqID No: 3, or SeqID No: 4.

91. A use of claim 87, wherein the first nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

92. A use of claim 87, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.

93. A use of claim 87, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

94. A use of claim 87, wherein upon delivery of the second nucleic acid expression construct into the cells of the subject, the second nucleic acid expression construct is capable of initiating expression of the encoded regulatory protein.

95. A use of claim 94, wherein the encoded regulatory protein is suitable for expression in the cells of the subject, but is inactive.

96. A use of claim 95, wherein upon delivery of the molecular switch into the subject the molecular switch is capable of activating the regulatory protein, and wherein the molecular switch comprises mifepristone or its functional equivalent.

97. A use of claim 96, wherein the active regulatory protein is suitable for initiating expression of the GHRH or functional biological equivalent encoded on the first nucleic acid sequence.

98. A use of claim 97, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

99. A use of claim 97, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D- or L- isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D- or L- isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

100. A use of claim 99 wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

101. A use of claim 87, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

102. A use of claim 87 wherein the first nucleic acid expression construct comprises SeqID No: 26.

103. A use of claim 87 wherein the second nucleic acid expression construct comprises SeqID No: 27.

104. Use of a recombinant growth- hormone- releasing-hormone ("GHRH") or a biological functional equivalent thereof in the preparation of a composition for increasing bone density in a subject, wherein the composition is suitable for delivery into the subject.

105. A use of claim 104, wherein the recombinant GHRH is a biologically active polypeptide; and the recombinant functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while

simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

106. A use of claim 104, wherein the recombinant GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2-DA TNSY VL-X_3-QLSA LLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D- or L- isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

107. A use of claim 106, wherein the recombinant GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

108. A use of claim 104, wherein the recombinant GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

109. Use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for increasing a bone healing rate in a subject, wherein the composition is suitable for delivery into cells of the subject.

110. A use of claim 109, wherein the nucleic acid expression construct is suitable for delivery into cells via electroporation.

111. A use of claim 109, wherein the cells of the subject are somatic cells, stem cells, or germ cells.

112. A use of claim 109, wherein the nucleic acid expression construct comprises SeqID No: 11, SeqlD No: 12, SeqID No: 13, SeqID No: 14, SeqID No: 17, SeqID No: 18, SeqID No: 19, SeqID No: 20, or SeqID No: 21.

113. A use of claim 109, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

114. A use of claim 113, wherein the transfection- facilitating polypeptide comprises a charged polypeptide.

115. A use of claim 113, wherein the transfection- facilitating polypeptide comprises poly-L-glutamate.

116. A use of claim 115, wherein upon delivery of the nucleic acid expression construct into the cells of the subject, the nucleic acid expression construct is capable of initiating expression of the encoded GHRH or functional biological equivalent thereof.

117. A use of claim 109, wherein the encoded GHRH or functional biological equivalent thereof is suitable for expression in tissue specific cells of the subject.

118. A use of claim 117, wherein the tissue specific cells of the subject comprises muscle cells.

119. A use of claim 109, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously

having similar or improved biologically activity when compared to the GHRH polypeptide.

120. A use of claim 109, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2-DA1 FTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D- or L- isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ('I");

or a combination thereof.

121. A use of claim 120, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

122. A use of claim 109, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

123. Use of a first nucleic acid expression construct, a second nucleic acid expression construct, and a molecular switch in the preparation of a composition for increasing a bone healing rate in a subject, and wherein the composition is suitable for delivery into cells of the subject, wherein the first nucleic acid expression construct is suitable for encoding growth hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof, wherein the second nucleic acid expression construct has an encoding region of a regulator protein, wherein upon delivery of the molecular switch

molecule into the subject, the molecular switch molecule is capable of activating the regulator protein.

124. A use of claim 123, wherein the first nucleic acid expression construct is suitable for delivery into cells of the subject via electroporation.

125. A use of claim 123, wherein the second nucleic acid expression construct is suitable for delivery into the cells of the subject via electroporation.

126. A use of claim 123, wherein the first nucleic acid expression construct encodes a polypeptide of a sequence comprising SeqID No: 1, SeqID No: 2, SeqID No: 3, or SeqID No: 4.

127. A use of claim 123, wherein the first nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.

128. A use of claim 127, wherein the transfection- facilitating polypeptide comprises a charged polypeptide.

129. A use of claim 127, wherein the transfection- facilitating polypeptide comprises poly-L-glutamate.

130. A use of claim 123, wherein upon delivery of the second nucleic acid expression construct into the cells of the subject, the second nucleic acid expression construct is capable of initiating expression of the encoded regulatory protein.

131. A use of claim 130, wherein the encoded regulatory protein is suitable for expression in the cells of the subject, but is inactive.

132. A use of claim 131, wherein upon delivery of the molecular switch into the subject the molecular switch is capable of activating the regulatory protein, and wherein the molecular switch comprises mifepristone or its functional equivalent.

133. A use of claim 132, wherein the active regulatory protein is suitable for initiating expression of the GHRH or functional biological equivalent encoded on the first nucleic acid sequence.

134. A use of claim 133, wherein the encoded GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

135. A use of claim 133, wherein the encoded GHRH or functional biological equivalent thereof is of formula (SEQID No: 6):

-X_1-X_2- DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5-RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D-or L- isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D-or L- isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ('I");

or a combination thereof.

136. A use of claim 135 wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

137. A use of claim 123, wherein the encoded GHRH or functional biological equivalent thereof is suitable for facilitating growth hormone ("GH") secretion in the subject.

138. A use of claim 123 wherein the first nucleic acid expression construct comprises SeqID No: 26.

139. A use of claim 123 wherein the second nucleic acid expression construct comprises SeqID No: 27.

140. Use of a nucleic acid expression construct that encodes a growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof in the preparation of a composition for altering lean body mass in a subject, wherein subsequent expression of the GHRH or biological equivalent is suitable for increasing lean body mass or decreasing body fat proportion, and wherein the composition is suitable for delivery into a tissue of the subject.

141. A use of claim 140, wherein applying a cell- transfecting pulse to the tissue, the cell-transfecting pulse is suitable for delivery to an area in the tissue comprising the delivered nucleic acid expression construct.

142. A use of claim 141, wherein placing a plurality of electrodes in the tissue before applying the cell-transfecting pulse to the tissue, the nucleic acid expression construct is suitable for delivery to the tissue in an area that interposes the plurality of electrodes.

143. A use of claim 141, wherein the cell-transfecting pulse is an electrical pulse or a vascular pressure pulse.

144. A use of claim 140, wherein the nucleic acid expression construct is suitable for delivery into a tissue of the subject via electroporation.

145. A use of claim 140, wherein the delivering step is by injection, gene gun, or gold particle bombardment.

146. A use of claim 140, wherein the subject is a domesticated animal; a food animal; or a work animal.

147. A use of claim 140, wherein the subject is a human.

148. A use of claim 140, wherein the nucleic acid expression construct is substantially free of a viral backbone.

149. A use of claim 140, wherein a promoter of the nucleic acid expression construct comprises a tissue-specific promoter.

150. A use of claim 149, wherein the tissue-specific promoter comprises a muscle-specific promoter.

151. A use of claim 149, wherein the promoter comprises SPc5-12 (SeqID No: 7).

152. A use of claim 140, wherein a 3' untranslated region of the nucleic expression construct is a human growth hormone 3' UTR, bovine growth hormone 3' UTR, skeletal alpha actin 3' UTR, or SV40 polyadenylation signal.

153. A use of claim 140, wherein the tissue comprises muscle.

154. A use of claim 140, wherein the nucleic acid expression construct is SeqID No: 11, SeqID No: 12, SeqID No: 13, SeqID No: 14, SeqID No: 17, SeqID No: 18, SeqID No: 19, SeqID No: 20, or SeqID No: 21.

155. A use of claim 140, wherein the encoded GHRH or functional biological equivalent comprises SeqID No: 2

156. A use of claim 140, wherein the encoded GHRH or functional biological equivalent comprises SeqID No: 3

157. A use of claim 140, wherein the encoded GHRH or functional biological equivalent comprises SeqID No: 4

158. A use of claim 140, wherein the encoded GHRH or functional biological equivalent comprises SeqID No: 10

159. A use of claim 140, wherein the encoded GHRH or functional biological equivalent thereof comprises the amino acid formula (SeqID No: 6):

-X_1-X_2-DAIFTNSYRKVL-X_3-QLSARKLLQDI-X_4-X_5- RQQGERNQEQGA-OH

wherein the formula has the following characteristics:

X_1 is a D- or L- isomer of the amino acid tyrosine ("Y"), or histidine ("H");

X_2 is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I");

X_3 is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G");

X_4 is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L");

X_5 is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N");

or a combination thereof.

160. A use of claim 140, wherein the nucleic acid expression construct further comprises a transfection-facilitating polypeptide.

161. A use of claim 160, wherein the transfection- facilitating polypeptide comprises a charged polypeptide.

162. A use of claim 160, wherein the transfection- facilitating polypeptide comprises poly-L-glutamate.

163. A use of claim140, wherein the encoded functional biological equivalent of GHRH is a polypeptide having similar or improved biological activity when compared to the GHRH polypeptide.

164. A use of claim 140, wherein the encoded GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion in the subject.

165. A peptide comprising: SeqID No: 2; SeqID No: 3; or SeqID No: 4.

REFERENCES CITED

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. PATENT DOCUMENTS

U.S. Patent No. 5,847,066 issued on December 8, 1998 with Coy *et al.* listed as inventors.

U.S. Patent No. 5,846,936 issued on December 8, 1998 with Felix *et al.* listed as inventors.

U.S. Patent No. 5,792,747 issued on August 11, 1998 with Schally *et al.* listed as inventors.

U.S. Patent No. 5,776,901 issued on July 7, 1998 with Bowers *et al.* listed as inventors.

U.S. Patent No. 5,756,264 issued on May 26, 1998 with Schwartz *et al.* listed as inventors.

U.S. Patent No. 5,696,089 issued on December 9, 1997 with Felix *et al.* listed as inventors.

U.S. Patent No. 5,486,505 issued on January 23, 1996 with Bowers *et al.* listed as inventors.

U.S. Patent No. 5,292,721 issued on March 8, 1994 with Boyd *et al.* listed as inventors.

U.S. Patent No. 5,137,872 issued on August 11, 1992 with Seely *et al.* listed as inventors.

U.S. Patent No. 5,134.120 issued on July 28, 1992 with Boyd *et al.* listed as inventors.

U.S. Patent No. 5,084,442 issued on January 28, 1992 with Felix *et al.* listed as inventors.

U.S. Patent No. 5,061,690 issued on October 29, 1991 with Kann *et al.* listed as inventors.

U.S. Patent No. 5,036,045 issued on July 30, 1991 with Thorner listed as the inventor.

U.S. Patent No. 5,023,322 issued on June 11, 1991 with Kovacs *et al.* listed as inventors.

U.S. Patent No. 4,839,344 issued on June 13, 1989 with Bowers *et al.* listed as inventors.

U.S. Patent No. 4,410,512 issued on October 18, 1983 with Bowers *et al.* listed as inventors.

U.S. Patent No. RE33,699 issued on September 24, 1991 with Drengler listed as the inventor.

U.S. Patent No. 4,833,166 issued on May 23, 1989 with Grosvenor *et al.* listed as inventors.

U.S. Patent No. 4,228,158 issued on October 14, 1980 with Momany *et al.* listed as inventors.

U.S. Patent No. 4,228,156 issued on October 14, 1980 with Momany *et al.* listed as inventors.

U.S. Patent No. 4,226,857 issued on October 7, 1980 with Momany *et al.* listed as inventors.

U.S. Patent No. 4,224,316 issued on September 23, 1980 with Momany *et al.* listed as inventors.

U.S. Patent No. 4,223,021 issued on September 16, 1980 with Momany *et al.* listed as inventors.

U.S. Patent No. 4,223,020 issued on September 16, 1980 with Momany *et al.* listed as inventors.

U.S. Patent No. 4,223,019 issued on September 16, 1980 with Momany *et al.* listed as inventors.

U.S. Patent No. 4,956,288 issued on September 11, 1990 with Barsoum listed as inventor.

U.S. Patent No. 5,704,908 issued on January 6, 1998 with Hofmann, *et al.* listed as inventors.

U.S. Patent No. 5,702,384 issued on December 30, 1997 with Umeyama, *et al.* listed as inventors.

U.S. Patent No. 5,439,440 issued on August 8, 1995 with Hofmann listed as inventor.

U.S. Patent No. 5,874,534 issued on February 23, 1999 with Vegeto, *et al* as inventors.

U.S. Patent No. 5,935,934 issued on August 10, 1999 with Vegeto, *et al* as inventors.

U.S. Patent No. 5,789,215 issued on August 4, 1998 with Berns , *et al.* as inventors.

U.S. Patent No. 5,384,253 issued on January 24, 1995 with Krzyzek , *et al.* as inventors.

U.S. Patent No. 5,994,624 issued on November 30, 1999 with Trolinder , *et al.* as inventors.

U.S. Patent No. 5,981,274 issued on November 9, 1999 with Tyrrell , *et al.* as inventors.

U.S. Patent No. 5,945,100 issued on August 31, 1999 with Fick as inventor.

U.S. Patent No. 5,780,448 issued on July 14, 1998 with Davis as inventor.

U.S. Patent No. 5,736,524 issued on April 7, 1998 with Content , *et al.* as inventors.

U.S. Patent No. 5,702,932 issued on December 30, 1997 with Hoy, *et al.* as inventors.

U.S. Patent No. 5,656,610 issued on August 12, 1997 with Shuler , *et al.* as inventors.

U.S. Patent No. 5,589,466 issued on December 31, 1996 with Felgner , *et al.* as inventors.

U.S. Patent No. 5,580,859 issued on December 3, 1996 with Felgner , *et al.* as inventors.

U.S. Patent No. 5,610,042 issued on March 11, 1997 with Chang, *et al.* as inventors.

U.S. Patent No. 5,322,783 issued on June 21, 1994 with Tomes, *et al.* as inventors.

U.S. Patent No. 5,563,055 issued on October 8, 1996 with Townsend, *et al.* as inventors.

U.S. Patent No. 5,550,318 issued on August 27, 1996 with Adams, *et al.* as inventors.

U.S. Patent No. 5,538,877 issued on July 23, 1996 with Lundquist, *et al.* as inventors.

U.S. Patent No. 5,538,880 issued on July 23, 1996 with Lundquist, *et al.* as inventors.

U.S. Patent No. 5,302,523 issued on April 12, 1994 with Coffee, *et al.* as inventors.

U.S. Patent No. 5,464,765 issued on November 7, 1995 with Coffee , *et al.* as inventors.

U.S. Patent No. 5,591,616 issued on January 7, 1997 with Hiei, *et al.* as inventors.

U.S. Patent No. 4,684,611 issued on August 4, 1987 with Schilperoort, *et al.* as inventors.

U.S. Patent No. 4,952,500 issued on August 28, 1990 with Finnerty , *et al.* as inventors.

WO 94/09699

WO95/06128;

Reference List

Abruzzese, R. V., D. Godin, M. Burcin, V. Mehta, M. French, Y. Li, B. W. O'Malley, and J. L. Nordstrom. 1999. Ligand-dependent regulation of plasmid-based transgene expression in vivo. Hum. Gene Ther. 10:1499-1507.

Abruzzese, R. V., D. Godin, V. Mehta, J. L. Perrard, M. French, W. Nelson, G. Howell, M. Coleman, B. W. O'Malley, and J. L. Nordstrom. 2000. Ligand-dependent regulation of vascular endothelial growth factor and erythropoietin expression by a plasmid-based autoinducible GeneSwitch system. Mol. Ther. 2:276-287.

Aihara, H. and J. Miyazaki. 1998. Gene transfer into muscle by electroporation in vivo. Nat. Biotechnol. 16:867-870.

Almendro, N., T. Bellon, C. Rius, P. Lastres, C. Langa, A. Corbi, and C. Bernabeu. 1996. Cloning of the human platelet endothelial cell adhesion molecule-1 promoter and its tissue-specific expression. Structural and functional characterization. J. Immunol. 157:5411-5421.

Aratani, Y., R. Okazaki, and H. Koyama. 1992. End extension repair of introduced targeting vectors mediated by homologous recombination in mammalian cells. Nucleic Acids Res. 20:4795-4801.

Argente, J., J. Pozo, and J. A. Chowen. 1996. The growth hormone axis: control and effects. Hormone Research 45 Suppl 1:9-11.

Barr, E. and J. M. Leiden. 1991. Systemic delivery of recombinant proteins by genetically modified myoblasts. Science 254:1507-1509.

Bercu, B. B., R. F. Walker 1997. Growth Hormone Secretagogues In Children With Altered Growth. Acta Paediatrica 86:102-106.

Bergsma, D. J., J. M. Grichnik, L. M. Gossett, and R. J. Schwartz. 1986. Delimitation and characterization of cis-acting DNA sequences required for the regulated expression and transcriptional control of the chicken skeletal alpha-actin gene. Molecular & Cellular Biology 6:2462-2475.

Bettan, M., F. Emmanuel, R. Darteil, J. M. Caillaud, F. Soubrier, P. Delaere, D. Branelec, A. Mahfoudi, N. Duverger, and D. Scherman. 2000. High-level protein secretion into blood circulation after electric pulse-mediated gene transfer into skeletal muscle. Mol. Ther. 2:204-210.

Blethen, S. L. 1995. Complications of growth hormone therapy in children. Curr. Opin. Pediatr. 7:466-471.

Blethen, S. L. and A. C. Rundle. 1996. Slipped capital femoral epiphysis in children treated with growth hormone. A summary of the National Cooperative Growth Study experience. Horm. Res. 46:113-116.

Bohlen, P., F. Esch, P. Brazeau, N. Ling, and R. Guillemin. 1983. Isolation and characterization of the porcine hypothalamic growth hormone releasing factor. Biochem. Biophys. Res. Commun. 116:726-734.

Boshart, M., F. Weber, G. Jahn, K. Dorsch-Hasler, B. Fleckenstein, and W. Schaffner. 1985. A very strong enhancer is located upstream of an immediate early gene of human cytomegalovirus. Cell 41:521-530.

Bureau, M. F., J. Gehl, V. Deleuze, L. M. Mir, and D. Scherman. 2000. Importance of association between permeabilization and electrophoretic forces for intramuscular DNA electrotransfer. Biochim. Biophys. Acta 1474:353-359.

Carbonelli, D. L., E. Corley, M. Seigelchifer, and J. Zorzopulos. 1999. A plasmid vector for isolation of strong promoters in Escherichia coli. FEMS Microbiol. Lett. 177:75-82.

Chandler, S. D., A. Mayeda, J. M. Yeakley, A. R. Krainer, and X. D. Fu. 1997. RNA splicing specificity determined by the coordinated action of RNA recognition motifs in SR proteins. Proc. Natl. Acad. Sci. U. S. A 94:3596-3601.

Chen, C. and H. Okayama. 1987. High-efficiency transformation of mammalian cells by plasmid DNA. Mol. Cell Biol. 7:2745-2752.

Chevalier, R. L., S. Goyal, A. Kim, A. Y. Chang, D. Landau, and D. LeRoith. 2000. Renal tubulointerstitial injury from ureteral obstruction in the neonatal rat is attenuated by IGF-1. Kidney Int. 57:882-890.

Cocea, L. 1997. Duplication of a region in the multiple cloning site of a plasmid vector to enhance cloning-mediated addition of restriction sites to a DNA fragment. Biotechniques 23:814-816.

Cohn, S. H., A. N. Vaswani, S. Yasumura, K. Yuen, and K. J. Ellis. 1984. Improved models for determination of body fat by in vivo neutron activation. Am. J Clin. Nutr. 40:255-259.

Corpas, E., S. M. Harman, M. A. Pineyro, R. Roberson, and M. R. Blackman. 1993. Continuous subcutaneous infusions of growth hormone (GH) releasing hormone 1-44 for 14 days increase GH and insulin-like growth factor-I levels in old men. Journal of Clinical Endocrinology & Metabolism 76:134-138.

Dai, B., H. Wu, E. Holthuizen, and P. Singh. 2001. Identification of a novel cis element required for cell density-dependent down-regulation of insulin-like growth factor-2 P3 promoter activity in Caco2 cells. J. Biol. Chem. 276:6937-6944.

Danko, I. and J. A. Wolff. 1994. Direct gene transfer into muscle. [Review]. Vaccine 12:1499-1502.

Darquet, A. M., B. Cameron, P. Wils, D. Scherman, and J. Crouzet. 1997. A new DNA vehicle for nonviral gene delivery: supercoiled minicircle. Gene Ther. 4:1341-1349.

Darquet, A. M., R. Rangara, P. Kreiss, B. Schwartz, S. Naimi, P. Delaere, J. Crouzet, and D. Scherman. 1999. Minicircle: an improved DNA molecule for in vitro and in vivo gene transfer. Gene Ther. 6:209-218.

Davis, H. L., R. G. Whalen, and B. A. Demeneix. 1993. Direct gene transfer into skeletal muscle in vivo: factors affecting efficiency of transfer and stability of expression. Human Gene Therapy 4:151-159.

Dhawan, J., L. C. Pan, G. K. Pavlath, M. A. Travis, A. M. Lanctot, and H. M. Blau. 1991. Systemic delivery of human growth hormone by injection of genetically engineered myoblasts. Science 254:1509-1512.

Dolnik, V., M. Novotny, and J. Chmelik. 1993. Electromigration behavior of poly-(L-glutamate) conformers in concentrated polyacrylamide gels. Biopolymers 33:1299-1306.

Dorsch-Hasler, K., G. M. Keil, F. Weber, M. Jasin, W. Schaffner, and U. H. Koszinowski. 1985. A long and complex enhancer activates transcription of the gene coding for the highly abundant immediate early mRNA in murine cytomegalovirus. Proc. Natl. Acad. Sci. U. S. A 82:8325-8329.

Draghia-Akli, R., M. L. Fiorotto, L. A. Hill, P. B. Malone, D. R. Deaver, and R. J. Schwartz. 1999. Myogenic expression of an injectable protease-resistant growth hormone-releasing hormone augments long-term growth in pigs. Nat. Biotechnol. 17:1179-1183.

Draghia-Akli, R., X. G. Li, and R. J. Schwartz. 1997. Enhanced growth by ectopic expression of growth hormone releasing hormone using an injectable myogenic vector. nature biotechnology 15:1285-1289.

Draghia-Akli, R., P. B. Malone, L. A. Hill, K. M. Ellis, R. J. Schwartz, and J. L. Nordstrom. 2002. Enhanced animal growth via ligand-regulated GHRH myogenic-injectable vectors. FASEB J. 16:426-428.

Dubreuil, P., T. Abribat, B. Broxup, and P. Brazeau. 1996. Long-term growth hormone-releasing factor administration on growth hormone, insulin-like growth factor-I concentrations, and bone healing in the Beagle. Canadian Journal of Veterinary Research 60:7-13.

Dubreuil, P., Y. Couture, G. Pelletier, D. Petitclerc, L. Delorme, H. Lapierre, P. Gaudreau, J. Morisset, and P. Brazeau. 1990a. Effect of long-term administration of porcine growth hormone-releasing factor and(or) thyrotropin-releasing factor on growth hormone, prolactin and thyroxine concentrations in growing pigs. J. Anim Sci. 68:95-107.

Dubreuil, P., D. Petitclerc, G. Pelletier, P. Gaudreau, C. Farmer, Mowles, TF, and P. Brazeau. 1990b. Effect of dose and frequency of administration of a potent analog of human growth hormone-releasing factor on hormone secretion and growth in pigs. Journal of Animal Science 68:1254-1268.

Duck, S. C., H. P. Schwarz, G. Costin, R. Rapaport, S. Arslanian, A. Hayek, M. Connors, and J. Jaramillo. 1992. Subcutaneous growth hormone-releasing hormone therapy in growth hormone-deficient children: first year of therapy. Journal of Clinical Endocrinology & Metabolism 75:1115-1120.

Evans, W. S., S. M. Anderson, L. T. Hull, P. P. Azimi, C. Y. Bowers, and J. D. Veldhuis. 2001. Continuous 24-hour intravenous infusion of recombinant human growth hormone (GH)-releasing hormone-(1-44)-amide augments pulsatile, entropic, and daily rhythmic GH secretion in postmenopausal women equally in the estrogen-withdrawn and estrogen-supplemented states. J. Clin. Endocrinol. Metab 86:700-712.

Evans, W. S., M. L. Vance, D. L. Kaiser, R. P. Sellers, J. L. Borges, T. R. Downs, L. A. Frohman, J. Rivier, W. Vale, and M. O. Thorner. 1985. Effects of intravenous, subcutaneous, and intranasal administration of growth hormone (GH)-releasing hormone-40 on serum GH concentrations in normal men. Journal of Clinical Endocrinology & Metabolism 61:846-850.

Fechheimer, M., J. F. Boylan, S. Parker, J. E. Sisken, G. L. Patel, and S. G. Zimmer. 1987. Transfection of mammalian cells with plasmid DNA by scrape loading and sonication loading. Proc. Natl. Acad. Sci. U. S. A 84:8463-8467.

Fewell, J. G., F. MacLaughlin, V. Mehta, M. Gondo, F. Nicol, E. Wilson, and L. C. Smith. 2001. Gene therapy for the treatment of hemophilia B using PINC-formulated plasmid delivered to muscle with electroporation. Mol. Ther. 3:574-583.

Frohman, L. A., T. R. Downs, E. P. Heimer, and A. M. Felix. 1989. Dipeptidylpeptidase IV and trypsin-like enzymatic degradation of human growth hormone-releasing hormone in plasma. J. Clin. Invest. 83:1533-1540.

Frohman, L. A., L. L. Nernardis, and K. J. Kant. 1968. Hypothalamic stimulation of growth hormone secretion. Science 162:580-582.

Frohman, L. A., J. L. Thominet, C. B. Webb, M. L. Vance, H. Uderman, J. Rivier, W. Vale, and M. O. Thorner. 1984. Metabolic clearance and plasma disappearance rates of human pancreatic tumor growth hormone releasing factor in man. J. Clin. Invest. 73:1304-1311.

Fryer, A. D. and D. B. Jacoby. 1993. Effect of inflammatory cell mediators on M2 muscarinic receptors in the lungs. Life Sci. 52:529-536.

Gehl, J., T. Skovsgaard, and L. M. Mir. 1998. Enhancement of cytotoxicity by electropermeabilization: an improved method for screening drugs. Anticancer Drugs 9:319-325.

Gehl, J., T. H. Sorensen, K. Nielsen, P. Raskmark, S. L. Nielsen, T. Skovsgaard, and L. M. Mir. 1999. In vivo electroporation of skeletal muscle: threshold, efficacy and relation to electric field distribution. Biochim. Biophys. Acta 1428:233-240.

German, M., S. Ashcroft, K. Docherty, H. Edlund, T. Edlund, S. Goodison, H. Imura, G. Kennedy, O. Madsen, D. Melloul, and . 1995. The insulin gene promoter. A simplified nomenclature. Diabetes 44:1002-1004.

Gesundheit, N. and J. K. Alexander. 1995. Endocrine Therapy with Recombinant Hormones and Growth Factors. In: B. D. Weintraub (Ed.) Molecular Endocrinology: Basic Concepts and Clinical Correlations. pp. 491-507. Raven Press,Ltd., New York.

Gilbert, R. A., M. J. Jaroszeski, and R. Heller. 1997. Novel electrode designs for electrochemotherapy. Biochim. Biophys. Acta 1334:9-14.

Gopal, T. V. 1985. Gene transfer method for transient gene expression, stable transformation, and cotransformation of suspension cell cultures. Mol. Cell Biol. 5:1188-1190.

Graham, F. L. and A. J. van der Eb. 1973. Transformation of rat cells by DNA of human adenovirus 5. Virology 54:536-539.

Guillemin, R., P. Brazeau, P. Bohlen, F. Esch, N. Ling, and W. B. Wehrenberg. 1982. Growth hormone-releasing factor from a human pancreatic tumor that caused acromegaly. Science 218:585-587.

Hafez, I. M., N. Maurer, and P. R. Cullis. 2001. On the mechanism whereby cationic lipids promote intracellular delivery of polynucleic acids. Gene Ther. 8:1188-1196.

Hamm, A., N. Krott, I. Breibach, R. Blindt, and A. K. Bosserhoff. 2002. Efficient transfection method for primary cells. Tissue Eng 8:235-245.

Harland, R. and H. Weintraub. 1985. Translation of mRNA injected into Xenopus oocytes is specifically inhibited by antisense RNA. J. Cell Biol. 101:1094-1099.

Heller, R., M. J. Jaroszeski, L. F. Glass, J. L. Messina, D. P. Rapaport, R. C. DeConti, N. A. Fenske, R. A. Gilbert, L. M. Mir, and D. S. Reintgen. 1996. Phase I/II trial for the treatment of cutaneous and subcutaneous tumors using electrochemotherapy. Cancer 77:964-971.

Horlick, R. A. and P. A. Benfield. 1989. The upstream muscle-specific enhancer of the rat muscle creatine kinase gene is composed of multiple elements. Mol. Cell Biol. 9:2396-2413.

Inouye, C., P. Remondelli, M. Karin, and S. Elledge. 1994. Isolation of a cDNA encoding a metal response element binding protein using a novel expression cloning procedure: the one hybrid system. DNA Cell Biol. 13:731-742.

Inouye, S., A. Nakazawa, and T. Nakazawa. 1985. Determination of the transcription initiation site and identification of the protein product of the regulatory gene xylR for xyl operons on the TOL plasmid. J. Bacteriol. 163:863-869.

Jaynes, J. B., J. E. Johnson, J. N. Buskin, C. L. Gartside, and S. D. Hauschka. 1988. The muscle creatine kinase gene is regulated by multiple upstream elements, including a muscle-specific enhancer. Mol. Cell Biol. 8:62-70.

Johnson, N. F., M. D. Hoover, D. G. Thomassen, Y. S. Cheng, A. Dalley, and A. L. Brooks. 1992. In vitro activity of silicon carbide whiskers in comparison to other industrial fibers using four cell culture systems. Am. J. Ind. Med. 21:807-823.

Kawamoto, T., K. Makino, H. Niwa, H. Sugiyama, S. Kimura, M. Amemura, A. Nakata, and T. Kakunaga. 1988. Identification of the human beta-actin enhancer and its binding factor. Mol. Cell Biol. 8:267-272.

Kawamoto, T., K. Makino, S. Orita, A. Nakata, and T. Kakunaga. 1989. DNA bending and binding factors of the human beta-actin promoter. Nucleic Acids Res. 17:523-537.

Klamut, H. J., L. O. Bosnoyan-Collins, R. G. Worton, P. N. Ray, and H. L. Davis. 1996. Identification of a transcriptional enhancer within muscle intron 1 of the human dystrophin gene. Hum. Mol. Genet. 5:1599-1606.

Klamut, H. J., S. B. Gangopadhyay, R. G. Worton, and P. N. Ray. 1990. Molecular and functional analysis of the muscle-specific promoter region of the Duchenne muscular dystrophy gene. Mol. Cell Biol. 10:193-205.

Kraus, J., M. Woltje, N. Schonwetter, and V. Hollt. 1998. Alternative promoter usage and tissue specific expression of the mouse somatostatin receptor 2 gene. FEBS Lett. 428:165-170.

Lareyre, J. J., T. Z. Thomas, W. L. Zheng, S. Kasper, D. E. Ong, M. C. Orgebin-Crist, and R. J. Matusik. 1999. A 5-kilobase pair promoter fragment of the murine epididymal retinoic acid-binding protein gene drives the tissue-specific, cell-specific, and androgen-regulated expression of a foreign gene in the epididymis of transgenic mice. J. Biol. Chem. 274:8282-8290.

Larsen, P. R., J. W. Harney, and D. D. Moore. 1986. Sequences required for cell-type specific thyroid hormone regulation of rat growth hormone promoter activity. J. Biol. Chem. 261:14373-14376.

Lee, S. H., W. Wang, S. Yajima, P. A. Jose, and M. M. Mouradian. 1997. Tissue-specific promoter usage in the D1A dopamine receptor gene in brain and kidney. DNA Cell Biol. 16:1267-1275.

Lesbordes, J. C., T. Bordet, G. Haase, L. Castelnau-Ptakhine, S. Rouhani, H. Gilgenkrantz, and A. Kahn. 2002. In vivo electrotransfer of the cardiotrophin-1 gene into skeletal muscle slows down progression of motor neuron degeneration in pmn mice. Hum. Mol. Genet. 11:1615-1625.

Levenson, V. V., E. D. Transue, and I. B. Roninson. 1998. Internal ribosomal entry site-containing retroviral vectors with green fluorescent protein and drug resistance markers. Hum. Gene Ther. 9:1233-1236.

Li, C., S. Ke, Q. P. Wu, W. Tansey, N. Hunter, L. M. Buchmiller, L. Milas, C. Charnsangavej, and S. Wallace. 2000. Tumor irradiation enhances the tumor-specific distribution of poly(L-glutamic acid)-conjugated paclitaxel and its antitumor efficacy. Clin. Cancer Res. 6:2829-2834.

Li, X., E. M. Eastman, R. J. Schwartz, and R. Draghia-Akli. 1999. Synthetic muscle promoters: activities exceeding naturally occurring regulatory sequences. nature biotechnology 17:241-245.

Lin, H., K. E. Yutzey, and S. F. Konieczny. 1991. Muscle-specific expression of the troponin I gene requires interactions between helix-loop-helix muscle regulatory factors and ubiquitous transcription factors. Mol. Cell Biol. 11:267-280.

Liu, Y., H. Li, K. Tanaka, N. Tsumaki, and Y. Yamada. 2000. Identification of an enhancer sequence within the first intron required for cartilage-specific transcription of the alpha2(XI) collagen gene. J. Biol. Chem. 275:12712-12718.

Lucas, M. L., L. Heller, D. Coppola, and R. Heller. 2002. IL-12 plasmid delivery by in vivo electroporation for the successful treatment of established subcutaneous B16.F10 melanoma. Mol. Ther. 5:668-675.

Lucas, M. L., M. J. Jaroszeski, R. Gilbert, and R. Heller. 2001. In vivo electroporation using an exponentially enhanced pulse: a new waveform. DNA Cell Biol. 20:183-188.

Macejak, D. G. and P. Sarnow. 1991. Internal initiation of translation mediated by the 5' leader of a cellular mRNA. Nature 353:90-94.

Madry, H., R. Reszka, J. Bohlender, and J. Wagner. 2001. Efficacy of cationic liposome-mediated gene transfer to mesangial cells in vitro and in vivo. J. Mol. Med. 79:184-189.

Matsubara, H., Y. Gunji, T. Maeda, K. Tasaki, Y. Koide, T. Asano, T. Ochiai, S. Sakiyama, and M. Tagawa. 2001. Electroporation-mediated transfer of cytokine genes into human esophageal tumors produces anti-tumor effects in mice. Anticancer Res. 21:2501-2503.

Matsuo, A., I. Tooyama, S. Isobe, Y. Oomura, I. Akiguchi, K. Hanai, J. Kimura, and H. Kimura. 1994. Immunohistochemical localization in the rat brain of an epitope corresponding to the fibroblast growth factor receptor-1. Neuroscience 60:49-66.

McNally, M. A., J. S. Lebkowski, T. B. Okarma, and L. B. Lerch. 1988. Optimizing electroporation parameters for a variety of human hematopoietic cell lines. Biotechniques 6:882-886.

Miklavcic, D., K. Beravs, D. Semrov, M. Cemazar, F. Demsar, and G. Sersa. 1998. The importance of electric field distribution for effective in vivo electroporation of tissues. Biophys. J 74:2152-2158.

Mir, L. M., M. F. Bureau, J. Gehl, R. Rangara, D. Rouy, J. M. Caillaud, P. Delaere, D. Branellec, B. Schwartz, and D. Scherman. 1999. High-efficiency gene transfer into skeletal muscle mediated by electric pulses. Proc. Natl. Acad. Sci. U. S. A 96:4262-4267.

Morel, G., R. Gallego, L. Boulanger, E. Pintos, T. Garcia-Caballero, and P. Gaudreau. 1999. Restricted presence of the growth hormone-releasing hormone receptor to somatotropes in rat and human pituitaries. Neuroendocrinology 70:128-136.

Mumper, R. J., J. Wang, S. L. Klakamp, H. Nitta, K. Anwer, F. Tagliaferri, and A. P. Rolland. 1998. Protective interactive noncondensing (PINC) polymers for enhanced plasmid distribution and expression in rat skeletal muscle. J. Control Release 52:191-203.

Muramatsu, T., A. Nakamura, and H. M. Park. 1998. In vivo electroporation: A powerful and convenient means of nonviral gene transfer to tissues of living animals. Int. J. Mol. Med. 1:55-62.

Murray, R. A., H. G. Maheshwari, E. J. Russell, and G. Baumann. 2000. Pituitary hypoplasia in patients with a mutation in the growth hormone-releasing hormone receptor gene. AJNR Am. J Neuroradiol. 21:685-689.

Murray, R. D. and S. M. Shalet. 2000. Growth hormone: current and future therapeutic applications. Expert. Opin. Pharmacother. 1:975-990.

Nabel, E. G., G. Plautz, F. M. Boyce, J. C. Stanley, and G. J. Nabel. 1989. Recombinant gene expression in vivo within endothelial cells of the arterial wall. Science 244:1342-1344.

Nairn, R. S., G. M. Adair, T. Porter, S. L. Pennington, D. G. Smith, J. H. Wilson, and M. M. Seidman. 1993. Targeting vector configuration and method of gene transfer influence targeted correction of the APRT gene in Chinese hamster ovary cells. Somat. Cell Mol. Genet. 19:363-375.

Narum, D. L., S. Kumar, W. O. Rogers, S. R. Fuhrmann, H. Liang, M. Oakley, A. Taye, B. K. Sim, and S. L. Hoffman. 2001. Codon optimization of gene fragments encoding Plasmodium falciparum merzoite proteins enhances DNA vaccine protein expression and immunogenicity in mice. Infect. Immun. 69:7250-7253.

Neumann, E., M. Schaefer-Ridder, Y. Wang, and P. H. Hofschneider. 1982. Gene transfer into mouse lyoma cells by electroporation in high electric fields. EMBO J. 1:841-845.

Nomoto, S., Y. Tatematsu, T. Takahashi, and H. Osada. 1999. Cloning and characterization of the alternative promoter regions of the human LIMK2 gene responsible for alternative transcripts with tissue-specific expression. Gene 236:259-271.

Ohlsson, H., S. Thor, and T. Edlund. 1991. Novel insulin promoter- and enhancer-binding proteins that discriminate between pancreatic alpha- and beta-cells. Mol. Endocrinol. 5:897-904.

Omirulleh, S., M. Abraham, M. Golovkin, I. Stefanov, M. K. Karabaev, L. Mustardy, S. Morocz, and D. Dudits. 1993. Activity of a chimeric promoter with the doubled CaMV 35S enhancer element in protoplast-derived cells and transgenic plants in maize. Plant Mol. Biol. 21:415-428.

Otani, Y., Y. Tabata, and Y. Ikada. 1996. Rapidly curable biological glue composed of gelatin and poly(L-glutamic acid). Biomaterials 17:1387-1391.

Otani, Y., Y. Tabata, and Y. Ikada. 1998. Hemostatic capability of rapidly curable glues from gelatin, poly(L-glutamic acid), and carbodiimide. Biomaterials 19:2091-2098.

Pech, M., C. D. Rao, K. C. Robbins, and S. A. Aaronson. 1989. Functional identification of regulatory elements within the promoter region of platelet-derived growth factor 2. Mol. Cell Biol. 9:396-405.

Pelletier, J. and N. Sonenberg. 1988. Internal initiation of translation of eukaryotic mRNA directed by a sequence derived from poliovirus RNA. Nature 334:320-325.

Pinkert, C. A., D. M. Ornitz, R. L. Brinster, and R. D. Palmiter. 1987. An albumin enhancer located 10 kb upstream functions along with its promoter to direct efficient, liver-specific expression in transgenic mice. Genes Dev. 1:268-276.

Potrykus, I., J. Paszkowski, M. W. Saul, J. Petruska, and R. D. Shillito. 1985. Molecular and general genetics of a hybrid foreign gene introduced into tobacco by direct gene transfer. Mol. Gen. Genet. 199:169-177.

Potter, H., L. Weir, and P. Leder. 1984. Enhancer-dependent expression of human kappa immunoglobulin genes introduced into mouse pre-B lymphocytes by electroporation. Proc. Natl. Acad. Sci. U. S. A 81:7161-7165.

Prentice, H., R. A. Kloner, T. Prigozy, T. Christensen, L. Newman, Y. Li, and L. Kedes. 1994. Tissue restricted gene expression assayed by direct DNA injection into cardiac and skeletal muscle. Journal of Molecular & Cellular Cardiology 26:1393-1401.

Raghavachari, N. and W. E. Fahl. 2002. Targeted gene delivery to skin cells in vivo: a comparative study of liposomes and polymers as delivery vehicles. J. Pharm. Sci. 91:615-622.

Rippe, R. A., D. A. Brenner, and H. L. Leffert. 1990. DNA-mediated gene transfer into adult rat hepatocytes in primary culture. Mol. Cell Biol. 10:689-695.

Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular Cloning - A Laboratory Manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

Satozawa, N., K. Takezawa, T. Miwa, S. Takahashi, M. Hayakawa, and H. Ooka. 2000. Differences in the effects of 20 K- and 22 K-hGH on water retention in rats. Growth Horm. IGF. Res. 10:187-192.

Skroch, P., C. Buchman, and M. Karin. 1993. Regulation of human and yeast metallothionein gene transcription by heavy metal ions. Prog. Clin. Biol. Res. 380:113-28.:113-128.

Smith, L. C. and J. L. Nordstrom. 2000. Advances in plasmid gene delivery and expression in skeletal muscle. Curr. Opin. Mol. Ther. 2:150-154.

Somiari, S., J. Glasspool-Malone, J. J. Drabick, R. A. Gilbert, R. Heller, M. J. Jaroszeski, and R. W. Malone. 2000. Theory and in vivo application of electroporative gene delivery. Mol. Ther. 2:178-187.

Soubrier, F., B. Cameron, B. Manse, S. Somarriba, C. Dubertret, G. Jaslin, G. Jung, C. L. Caer, D. Dang, J. M. Mouvault, D. Scherman, J. F. Mayaux, and J. Crouzet. 1999. pCOR: a new design of plasmid vectors for nonviral gene therapy. Gene Ther. 6:1482-1488.

Terada, Y., H. Tanaka, T. Okado, S. Inoshita, M. Kuwahara, T. Akiba, S. Sasaki, and F. Marumo. 2001. Efficient and ligand-dependent regulated erythropoietin production by naked dna injection and in vivo electroporation. Am. J Kidney Dis. 38:S50-S53.

Thorner, M. O., L. A. Frohman, D. A. Leong, J. Thominet, T. Downs, P. Hellmann, J. Chitwood, J. M. Vaughan, and W. Vale. 1984. Extrahypothalamic growth-hormone-releasing factor (GRF) secretion is a rare cause of acromegaly: plasma GRF levels in 177 acromegalic patients. Journal of Clinical Endocrinology & Metabolism 59:846-849.

Thorner, M. O., M. L. Vance, W. S. Evans, A. D. Rogol, J. Rivier, W. Vale, Blizzard, and RM. 1986. CLINICAL STUDIES WITH GHRH IN MAN. Hormone Research 24:91-98.

Thorner, M. O., M. L. Vance, M. L. Hartman, R. W. Holl, W. S. Evans, J. D. Veldhuis, E. Van Cauter, G. Copinschi, and C. Y. Bowers. 1990. Physiological role of somatostatin on growth hormone regulation in humans. Metabolism: Clinical & Experimental 39:40-42.

Toneguzzo, F., A. Keating, S. Glynn, and K. McDonald. 1988. Electric field-mediated gene transfer: characterization of DNA transfer and patterns of integration in lymphoid cells. Nucleic Acids Res. 16:5515-5532.

Toner, M. S., R. H. King, F. R. Dunshea, H. Dove, and C. S. Atwood. 1996. The effect of exogenous somatotropin on lactation performance of first-litter sows. J. Anim Sci. 74:167-172.

Tripathy, S. K., E. C. Svensson, H. B. Black, E. Goldwasser, M. Margalith, Hobart, PM, and J. M. Leiden. 1996. Long-term expression of erythropoietin in the systemic circulation of mice after intramuscular injection of a plasmid DNA vector. Proc. Natl. Acad. Sci. USA 93:10876-10880.

Tronche, F., A. Rollier, I. Bach, M. C. Weiss, and M. Yaniv. 1989. The rat albumin promoter: cooperation with upstream elements is required when binding of APF/HNF1 to the proximal element is partially impaired by mutation or bacterial methylation. Mol. Cell Biol. 9:4759-4766.

Tronche, F., A. Rollier, P. Herbomel, I. Bach, S. Cereghini, M. Weiss, and M. Yaniv. 1990. Anatomy of the rat albumin promoter. Mol. Biol. Med. 7:173-185.

Trudel, M. and F. Costantini. 1987. A 3' enhancer contributes to the stage-specific expression of the human beta-globin gene. Genes Dev. 1:954-961.

Tsumaki, N., T. Kimura, K. Tanaka, J. H. Kimura, T. Ochi, and Y. Yamada. 1998. Modular arrangement of cartilage- and neural tissue-specific cis-elements in the mouse alpha2(XI) collagen promoter. J. Biol. Chem. 273:22861-22864.

Tsunekawa, B., M. Wada, M. Ikeda, H. Uchida, N. Naito, and M. Honjo. 1999. The 20-kilodalton (kDa) human growth hormone (hGH) differs from the 22-kDa hGH in the effect on the human prolactin receptor. Endocrinology 140:3909-3918.

Tsurumi, Y., S. Takeshita, D. Chen, M. Kearney, S. T. Rossow, J. Passeri, J. R. Horowitz, J. F. Symes, and J. M. Isner. 1996. Direct intramuscular gene transfer of naked DNA encoding vascular endothelial growth factor augments collateral development and tissue perfusion [see comments]. Circulation 94:3281-3290.

Tur-Kaspa, R., L. Teicher, B. J. Levine, A. I. Skoultchi, and D. A. Shafritz. 1986. Use of electroporation to introduce biologically active foreign genes into primary rat hepatocytes. Mol. Cell Biol. 6:716-718.

Vance, M. L. 1990. Growth-hormone-releasing hormone. [Review] [52 refs]. Clinical Chemistry 36:415-420.

Vance, M. L., D. L. Kaiser, W. S. Evans, R. Furlanetto, W. Vale, J. Rivier, and M. O. Thorner. 1985a. Pulsatile growth hormone secretion in normal man during a continuous 24-hour infusion of human growth hormone releasing factor (1-40). Evidence for intermittent somatostatin secretion. J. Clin. Invest. 75:1584-1590.

Vance, M. L., D. L. Kaiser, W. S. Evans, M. O. Thorner, R. Furlanetto, J. Rivier, W. Vale, G. Perisutti, and L. A. Frohman. 1985b. Evidence for a limited growth hormone (GH)-releasing hormone (GHRH)-releasable quantity of GH: effects of 6-hour infusions of GHRH on GH secretion in normal man. Journal of Clinical Endocrinology & Metabolism 60:370-375.

Vegeto, E., G. F. Allan, W. T. Schrader, M. J. Tsai, D. P. McDonnell, and B. W. O'Malley. 1992. The mechanism of RU486 antagonism is dependent on the conformation of the carboxy-terminal tail of the human progesterone receptor. Cell 69:703-713.

Verhelst, J., R. Abs, M. Vandeweghe, J. Mockel, J. J. Legros, G. Copinschi, C. Mahler, B. Velkeniers, L. Vanhaelst, A. Van Aelst, D. De Rijdt, A. Stevenaert, and A. Beckers. 1997. Two years of replacement therapy in adults with growth hormone deficiency. Clin. Endocrinol. (Oxf) 47:485-494.

Vilquin, J. T., P. F. Kennel, M. Paturneau-Jouas, P. Chapdelaine, N. Boissel, P. Delaere, J. P. Tremblay, D. Scherman, M. Y. Fiszman, and K. Schwartz. 2001. Electrotransfer of naked DNA in the skeletal muscles of animal models of muscular dystrophies. Gene Ther. 8:1097-1107.

Vittone, J., M. R. Blackman, J. Busby-Whitehead, C. Tsiao, K. J. Stewart, J. Tobin, T. Stevens, M. F. Bellantoni, M. A. Rogers, G. Baumann, J. Roth, S. M. Harman, and R. G. S. Spencer. 1997. Effects of single nightly injections of growth hormone-releasing hormone (GHRH 1-29) in healthy elderly men. Metabolism: Clinical and Experimental 46:89-96.

Wada, M., H. Uchida, M. Ikeda, B. Tsunekawa, N. Naito, S. Banba, E. Tanaka, Y. Hashimoto, and M. Honjo. 1998. The 20-kilodalton (kDa) human growth hormone (hGH) differs from the 22-kDa hGH in the complex formation with cell surface hGH receptor and hGH-binding protein circulating in human plasma. Mol. Endocrinol. 12:146-156.

Wells, K. E., J. Maule, R. Kingston, K. Foster, J. McMahon, E. Damien, A. Poole, and D. J. Wells. 1997. Immune responses, not promoter inactivation, are responsible for decreased long-term expression following plasmid gene transfer into skeletal muscle. FEBS Lett. 407:164-168.

Wiethoff, C. M., J. G. Smith, G. S. Koe, and C. R. Middaugh. 2001. The potential role of proteoglycans in cationic lipid-mediated gene delivery. Studies of the interaction of cationic lipid-DNA complexes with model glycosaminoglycans. J. Biol. Chem. 276:32806-32813.

Wilson, J. M., L. K. Birinyi, R. N. Salomon, P. Libby, A. D. Callow, and R. C. Mulligan. 1989. Implantation of vascular grafts lined with genetically modified endothelial cells. Science 244:1344-1346.

Wolff, J. A., R. W. Malone, P. Williams, W. Chong, G. Acsadi, A. Jani, Felgner, and PL. 1990. Direct gene transfer into mouse muscle in vivo. Science 247:1465-1468.

Wu, G. Y. and C. H. Wu. 1988a. Evidence for targeted gene delivery to Hep G2 hepatoma cells in vitro. Biochemistry 27:887-892.

Wu, G. Y. and C. H. Wu. 1988b. Receptor-mediated gene delivery and expression in vivo. J. Biol. Chem. 263:14621-14624.

Wu, H. K., J. A. Squire, Q. Song, and R. Weksberg. 1997. Promoter-dependent tissue-specific expressive nature of imprinting gene, insulin-like growth factor II, in human tissues. Biochem. Biophys. Res. Commun. 233:221-226.

Xie, T. D. and T. Y. Tsong. 1993. Study of mechanisms of electric field-induced DNA transfection. V. Effects of DNA topology on surface binding, cell uptake, expression, and integration into host chromosomes of DNA in the mammalian cell. Biophys. J. 65:1684-1689.

Xu, J., Z. Nawaz, S. Y. Tsai, M. J. Tsai, and B. W. O'Malley. 1996. The extreme C terminus of progesterone receptor contains a transcriptional repressor domain that functions through a putative corepressor. Proc. Natl. Acad. Sci. USA 93:12195-12199.

Yasui, A., K. Oda, H. Usunomiya, K. Kakudo, T. Suzuki, T. Yoshida, H. M. Park, K. Fukazawa, and T. Muramatsu. 2001. Elevated gastrin secretion by in vivo gene electroporation in skeletal muscle. Int. J Mol. Med. 8:489-494.

Yin, D. and J. G. Tang. 2001. Gene therapy for streptozotocin-induced diabetic mice by electroporational transfer of naked human insulin precursor DNA into skeletal muscle in vivo. FEBS Lett. 495:16-20.

Yorifuji, T. and H. Mikawa. 1990. Co-transfer of restriction endonucleases and plasmid DNA into mammalian cells by electroporation: effects on stable transformation. Mutat. Res. 243:121-126.

Yutzey, K. E. and S. F. Konieczny. 1992. Different E-box regulatory sequences are functionally distinct when placed within the context of the troponin I enhancer. Nucleic Acids Res. 20:5105-5113.

Zhao-Emonet, J. C., O. Boyer, J. L. Cohen, and D. Klatzmann. 1998. Deletional and mutational analyses of the human CD4 gene promoter: characterization of a minimal tissue-specific promoter. Biochim. Biophys. Acta 1442:109-119.

SEQUENCE LISTING

<110> Baylor College of Medicine

<120> A COMPOSITION AND METHOD TO ALTER LEAN BODY MASS AND BONE PROPERTIES IN A SUBJECT.

<130> 108328.00071 - AVSI-0008

<160> 27

<170> PatentIn version 3.1

<210> 1
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is a growth hormone releasing hormone ("GHRH") with a His1 and Val2 substituting the Try1 and Ala2, Gly15 substituted with A la15, and Met27, Ser28 with Leu27 and Asn28.

<400> 1

His Val Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
20 25 30

Glu Arg Asn Gln Glu Gln Gly Ala
35 40

<210> 2
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is a modified amino acid sequence for growth hormone releasi ng hormone (GHRH). Alpha-helical conformation was increased by su bstituting Gly15 to Ala15. This is the TI-GHRH sequence so it als o has the Ile2 instead of Ala2, Gly15 substituted with Ala15, and Met27, Ser28 with Leu27 and Asn28.

<400> 2

Tyr Ile Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
20 25 30

Glu Arg Asn Gln Glu Gln Gly Ala
35 40

<210> 3
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is a growth hormone releasing hormone that has a Val2 substi tution for a Ala2, Gly15 substituted with Ala15, and Met27, Ser28 with Leu27 and Asn28.

<400> 3

Tyr Val Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
20 25 30

Glu Arg Asn Gln Glu Gln Gly Ala
35 40

<210> 4
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is a porcine growth hormone releasing hormone ("GHRH") that has the following substitutions: Gly15 substituted with Ala15, Met27, Ser28 with Leu27 and Asn28.

<400> 4

Tyr Ala Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
20 25 30

Glu Arg Asn Gln Glu Gln Gly Ala
35 40

<210> 5
<211> 44
<212> PRT
<213> artificial sequence

<220>
<223> This is the artificial sequence for the (1-44)NH2

<400> 5

Thr Ala Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Gly Gln
1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Met Ser Arg Gln Gln Gly
20 25 30

Glu Ser Asn Gln Glu Arg Gly Ala Arg Ala Arg Leu
35 40

<210> 6
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is the artificial sequence for GHRH (1-40)OH.

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Xaa at position 1 may be tyrosine, or histidine

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> Xaa at position 2 may be alanine, valine, or isoleucine.

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa at position 15 may be alanine, valine, or isoleucine.

<220>
<221> MISC_FEATURE
<222> (27)..(27)
<223> Xaa at position 27 may be methionine, or leucine.

<220>
<221> MISC_FEATURE
<222> (28)..(28)
<223> Xaa at position 28 may be serine or asparagine.

<220>
<221> MISC_FEATURE
<222> (34)..(34)
<223> ARG may also be SER

<220>
<221> MISC_FEATURE
<222> (38)..(38)
<223> Gln may also be Arg

<400> 6

Xaa Xaa Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Xaa Gln
1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Xaa Xaa Arg Gln Gln Gly
20 25 30

Glu Arg Asn Gln Glu Gln Gly Ala
35 40

<210> 7
<211> 323
<212> DNA
<213> artificial sequence

<220>
<223> This is a nucleic acid sequence of a eukaryotic promoter c5-12.

<400> 7
cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg gtgaggaatg 60

gtggggagtt atttttagag cggtgaggaa ggtgggcagg cagcaggtgt tggcgctcta 120

aaaataactc ccgggagtta tttttagagc ggaggaatgg tggacaccca aatatggcga 180

cggttcctca cccgtcgcca tatttgggtg tccgccctcg gccgggggccg cattcctggg 240

ggccgggcgg tgctcccgcc cgcctcgata aaaggctccg gggccggcgg cggcccacga 300

gctacccgga ggagcgggag gcg 323

<210> 8
<211> 190
<212> DNA
<213> artificial sequence

<220>
<223> This is a nucleic acid sequence of a human growth hormone ("hGH") poly A tail.

<400> 8
gggtggcatc cctgtgaccc ctccccagtg cctctcctgg ccctggaagt tgccactcca 60

gtgcccacca gccttgtcct aataaaatta agttgcatca ttttgtctga ctaggtgtcc 120

ttctataata ttatggggtg gaggggggtg gtatggagca aggggcaagt tgggaagaca 180

acctgtaggg 190

<210> 9
<211> 219
<212> DNA
<213> artificial sequence

<220>
<223> This is the cDNA for Porcine growth hormone releasing hormone

<400> 9
atggtgctct gggtgttctt ctttgtgatc ctcaccctca gcaacagctc ccactgctcc 60

ccacctcccc ctttgaccct caggatgcgg cggcacgtag atgccatctt caccaacagc 120

taccggaagg tgctggccca gctgtccgcc cgcaagctgc tccaggacat cctgaacagg 180

cagcagggag agaggaacca agagcaagga gcataatga 219

<210> 10
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is the amino acid sequence for porcine growth hormone releasing hormone.

<400> 10

Tyr Ala Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Gly Gln
1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Met Ser Arg Gln Gln Gly
20 25 30

Glu Arg Asn Gln Glu Gln Gly Ala
35 40

<210> 11
<211> 3534
<212> DNA
<213> artificial sequence

<220>
<223> This is the nucleic acid sequence for the operatively linked components of the HV-GHRH plasmid.

<400> 11
gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60

accgcggtgg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120

gtgaggaatg gtggggagtt atttttagag cggtgaggaa ggtgggcagg cagcaggtgt 180

tggcgctcta aaaataactc ccgggagtta tttttagagc ggaggaatgg tggacaccca 240

aatatggcga cggttcctca cccgtcgcca tatttgggtg tccgccctcg gccggggccg 300

cattcctggg ggccgggcgg tgctcccgcc cgcctcgata aaaggctccg gggccggcgg 360

cggcccacga gctacccgga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420

ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480

ctgggtgttc ttctttgtga tcctcaccct cagcaacagc tcccactgct ccccacctcc 540

ccctttgacc ctcaggatgc ggcggcacgt agatgccatc ttcaccaaca gctaccggaa 600

ggtgctggcc cagctgtccg cccgcaagct gctccaggac atcctgaaca ggcagcaggg 660

agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720

ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt gccactccag 780

tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct 840

tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt gggaagacaa 900

cctgtagggc ctgcggggtc tattgggaac caagctggag tgcagtggca caatcttggc 960

tcactgcaat ctccgcctcc tgggttcaag cgattctcct gcctcagcct cccgagttgt 1020

tgggattcca ggcatgcatg accaggctca gctaattttt gtttttttgg tagagacggg 1080

gtttcaccat attggccagg ctggtctcca actcctaatc tcaggtgatc tacccacctt 1140

ggcctcccaa attgctggga ttacaggcgt gaaccactgc tcccttccct gtccttctga 1200

ttttaaaata actataccag caggaggacg tccagacaca gcataggcta cctggccatg 1260

cccaaccggt gggacatttg agttgcttgc ttggcactgt cctctcatgc gttgggtcca 1320

ctcagtagat gcctgttgaa ttcgataccg tcgacctcga ggggggggcc ggtaccagct 1380

tttgttccct ttagtgaggg ttaatttcga gcttggcgta atcatggtca tagctgtttc 1440

ctgtgtgaaa ttgttatccg ctcacaattc cacacaacat acgagccgga agcataaagt 1500

gtaaagcctg gggtgcctaa tgagtgagct aactcacatt aattgcgttg cgctcactgc 1560

ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg 1620

ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct 1680

cggtcgttcg gctgcggcga gcggtatcag ctcactcaaa ggcggtaata cggttatcca 1740

cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga 1800

accgtaaaaa ggccgcgttg ctggcgtttt tccataggct ccgcccccct gacgagcatc 1860

acaaaaatcg acgctcaagt cagaggtggc gaaacccgac aggactataa agataccagg 1920

cgtttccccc tggaagctcc ctcgtgcgct ctcctgttcc gaccctgccg cttaccggat 1980

acctgtccgc ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt 2040

atctcagttc ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa ccccccgttc 2100

agcccgaccg ctgcgcctta tccggtaact atcgtcttga gtccaacccg gtaagacacg 2160

acttatcgcc actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg 2220

gtgctacaga gttcttgaag tggtggccta actacggcta cactagaaga acagtatttg 2280

gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg 2340

gcaaacaaac caccgctggt agcggtggtt tttttgtttg caagcagcag attacgcgca 2400

gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagaaga 2460

actcgtcaag aaggcgatag aaggcgatgc gctgcgaatc gggagcggcg ataccgtaaa 2520

gcacgaggaa gcggtcagcc cattcgccgc caagctcttc agcaatatca cgggtagcca 2580

acgctatgtc ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa 2640

agcggccatt ttccaccatg atattcggca agcaggcatc gccatgggtc acgacgagat 2700

cctcgccgtc gggcatgcgc gccttgagcc tggcgaacag ttcggctggc gcgagcccct 2760

gatgctcttc gtccagatca tcctgatcga caagaccggc ttccatccga gtacgtgctc 2820

gctcgatgcg atgtttcgct tggtggtcga atgggcaggt agccggatca agcgtatgca 2880

gccgccgcat tgcatcagcc atgatggata ctttctcggc aggagcaagg tgagatgaca 2940

ggagatcctg ccccggcact tcgcccaata gcagccagtc ccttcccgct tcagtgacaa 3000

cgtcgagcac agctgcgcaa ggaacgcccg tcgtggccag ccacgatagc cgcgctgcct 3060

cgtcctgcag ttcattcagg gcaccggaca ggtcggtctt gacaaaaaga accgggcgcc 3120

cctgcgctga cagccggaac acggcggcat cagagcagcc gattgtctgt tgtgcccagt 3180

catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttgtt 3240

caatcatgcg aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc 3300

agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag 3360

agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca 3420

actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg 3480

gatgtgctgc aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgac 3534

<210> 12
<211> 3534
<212> DNA
<213> artificial sequence

<220>
<223> This is the nucleic acid sequence for the operatively linked components of the TI-GHRH plasmid.

<400> 12

gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60

accgcggtgg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120

gtgaggaatg gtggggagtt atttttagag cggtgaggaa ggtgggcagg cagcaggtgt 180

tggcgctcta aaaataactc ccgggagtta tttttagagc ggaggaatgg tggacaccca 240

aatatggcga cggttcctca cccgtcgcca tatttgggtg tccgccctcg gccggggccg 300

cattcctggg ggccgggcgg tgctcccgcc cgcctcgata aaaggctccg gggccggcgg 360

cggcccacga gctacccgga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420

ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480

ctgggtgttc ttctttgtga tcctcaccct cagcaacagc tcccactgct ccccacctcc 540

ccctttgacc ctcaggatgc ggcggtatat cgatgccatc ttcaccaaca gctaccggaa 600

ggtgctggcc cagctgtccg cccgcaagct gctccaggac atcctgaaca ggcagcaggg 660

agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720

ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt gccactccag 780

tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct 840

tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt gggaagacaa 900

cctgtagggc ctgcggggtc tattgggaac caagctggag tgcagtggca caatcttggc 960

tcactgcaat ctccgcctcc tgggttcaag cgattctcct gcctcagcct cccgagttgt 1020

tgggattcca ggcatgcatg accaggctca gctaattttt gtttttttgg tagagacggg 1080

gtttcaccat attggccagg ctggtctcca actcctaatc tcaggtgatc tacccacctt 1140

ggcctcccaa attgctggga ttacaggcgt gaaccactgc tcccttccct gtccttctga 1200

ttttaaaata actataccag caggaggacg tccagacaca gcataggcta cctggccatg 1260

cccaaccggt gggacatttg agttgcttgc ttggcactgt cctctcatgc gttgggtcca 1320

ctcagtagat gcctgttgaa ttcgataccg tcgacctcga gggggggccc ggtaccagct 1380

tttgttccct ttagtgaggg ttaatttcga gcttggcgta atcatggtca tagctgtttc 1440

ctgtgtgaaa ttgttatccg ctcacaattc cacacaacat acgagccgga agcataaagt 1500

gtaaagcctg gggtgcctaa tgagtgagct aactcacatt aattgcgttg cgctcactgc 1560

ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg 1620

ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct 1680

cggtcgttcg gctgcggcga gcggtatcag ctcactcaaa ggcggtaata cggttatcca 1740

cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga 1800

accgtaaaaa ggccgcgttg ctggcgtttt tccataggct ccgcccccct gacgagcatc 1860

acaaaaatcg acgctcaagt cagaggtggc gaaacccgac aggactataa agataccagg 1920

cgtttccccc tggaagctcc ctcgtgcgct ctcctgttcc gaccctgccg cttaccggat 1980

acctgtccgc ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt 2040

atctcagttc ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa ccccccgttc 2100

agcccgaccg ctgcgccttа tccggtaact atcgtcttga gtccaacccg gtaagacacg 2160

acttatcgcc actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg 2220

gtgctacaga gttcttgaag tggtggccta actacggcta cactagaaga acagtatttg 2280

gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg 2340

gcaaacaaac caccgctggt agcggtggtt tttttgtttg caagcagcag attacgcgca 2400

gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagaaga 2460

actcgtcaag aaggcgatag aaggcgatgc gctgcgaatc gggagcggcg ataccgtaaa 2520

gcacgaggaa gcggtcagcc cattcgccgc caagctcttc agcaatatca cgggtagcca 2580

acgctatgtc ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa 2640

agcggccatt ttccaccatg atattcggca agcaggcatc gccatgggtc acgacgagat 2700

cctcgccgtc gggcatgcgc gccttgagcc tggcgaacag ttcggctggc gcgagcccct 2760

gatgctcttc gtccagatca tcctgatcga caagaccggc ttccatccga gtacgtgctc 2820

gctcgatgcg atgtttcgct tggtggtcga atgggcaggt agccggatca agcgtatgca 2880

gccgccgcat tgcatcagcc atgatggata ctttctcggc aggagcaagg tgagatgaca 2940

ggagatcctg ccccggcact tcgcccaata gcagccagtc ccttcccgct tcagtgacaa 3000

cgtcgagcac agctgcgcaa ggaacgcccg tcgtggccag ccacgatagc cgcgctgcct 3060

cgtcctgcag ttcattcagg gcaccggaca ggtcggtctt gacaaaaaga accgggcgcc 3120

cctgcgctga cagccggaac acggcggcat cagagcagcc gattgtctgt tgtgcccagt 3180

catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttgtt 3240

caatcatgcg aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc 3300

agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag 3360

agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca 3420

actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg 3480

gatgtgctgc aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgac 3534

<210> 13
<211> 3534
<212> DNA
<213> artificial sequence

<220>
<223> This is the nucleic acid sequence for the operatively linked components of the TV-GHRH plasmid.

<400> 13
gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60

accgcggtgg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120

gtgaggaatg gtggggagtt atttttagag cggtgaggaa ggtgggcagg cagcaggtgt 180

tggcgctcta aaaataactc ccgggagtta tttttagagc ggaggaatgg tggacaccca 240

aatatggcga cggttcctca cccgtcgcca tatttgggtg tccgccctcg gccggggccg 300

cattcctggg ggccgggcgg tgctcccgcc cgcctcgata aaaggctccg gggccggcgg 360

cggcccacga gctacccgga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420

ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480

ctgggtgttc ttctttgtga tcctcaccct cagcaacagc tcccactgct ccccacctcc 540

ccctttgacc ctcaggatgc ggcggtatgt agatgccatc ttcaccaaca gctaccggaa 600

ggtgctggcc cagctgtccg cccgcaagct gctccaggac atcctgaaca ggcagcaggg 660

agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720

ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt gccactccag 780

tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct 840

tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt gggaagacaa 900

cctgtagggc ctgcggggtc tattgggaac caagctggag tgcagtggca caatcttggc 960

tcactgcaat ctccgcctcc tgggttcaag cgattctcct gcctcagcct cccgagttgt 1020

tgggattcca ggcatgcatg accaggctca gctaattttt gtttttttgg tagagacggg 1080

gtttcaccat attggccagg ctggtctcca actcctaatc tcaggtgatc tacccacctt 1140

ggcctcccaa attgctggga ttacaggcgt gaaccactgc tcccttccct gtccttctga 1200

ttttaaaata actataccag caggaggacg tccagacaca gcataggcta cctggccatg 1260

cccaaccggt gggacatttg agttgcttgc ttggcactgt cctctcatgc gttgggtcca 1320

ctcagtagat gcctgttgaa ttcgataccg tcgacctcga gggggggccc ggtaccagct 1380

tttgttccct ttagtgaggg ttaatttcga gcttggcgta atcatggtca tagctgtttc 1440

ctgtgtgaaa ttgttatccg ctcacaattc cacacaacat acgagccgga agcataaagt 1500

gtaaagcctg gggtgcctaa tgagtgagct aactcacatt aattgcgttg cgctcactgc 1560

ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg 1620

ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct 1680

cggtcgttcg gctgcggcga gcggtatcag ctcactcaaa ggcggtaata cggttatcca 1740

cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga 1800

accgtaaaaa ggccgcgttg ctggcgtttt tccataggct ccgcccccct gacgagcatc 1860

acaaaaatcg acgctcaagt cagaggtggc gaaacccgac aggactataa agataccagg 1920

cgtttccccc tggaagctcc ctcgtgcgct ctcctgttcc gaccctgccg cttaccggat 1980

acctgtccgc ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt 2040

atctcagttc ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa cccccgttc 2100

agcccgaccg ctgcgccttа tccggtaact atcgtcttga gtccaacccg gtaagacacg 2160

acttatcgcc actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg 2220

gtgctacaga gttcttgaag tggtggccta actacggcta cactagaaga acagtatttg 2280

gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg 2340

gcaaacaaac caccgctggt agcggtggtt tttttgtttg caagcagcag attacgcgca 2400

gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagaaga 2460

actcgtcaag aaggcgatag aaggcgatgc gctgcgaatc gggagcggcg ataccgtaaa 2520

gcacgaggaa gcggtcagcc cattcgccgc caagctcttc agcaatatca cgggtagcca 2580

acgctatgtc ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa 2640

agcggccatt ttccaccatg atattcggca agcaggcatc gccatgggtc acgacgagat 2700

cctcgccgtc gggcatgcgc gccttgagcc tggcgaacag ttcggctggc gcgagcccct 2760

gatgctcttc gtccagatca tcctgatcga caagaccggc ttccatccga gtacgtgctc 2820

gctcgatgcg atgtttcgct tggtggtcga atgggcaggt agccggatca agcgtatgca 2880

gccgccgcat tgcatcagcc atgatggata ctttctcggc aggagcaagg tgagatgaca 2940

ggagatcctg ccccggcact tcgcccaata gcagccagtc ccttcccgct tcagtgacaa 3000

cgtcgagcac agctgcgcaa ggaacgcccg tcgtggccag ccacgatagc cgcgctgcct 3060

cgtcctgcag ttcattcagg gcaccggaca ggtcggtctt gacaaaaaga accgggcgcc 3120

cctgcgctga cagccggaac acggcggcat cagagcagcc gattgtctgt tgtgcccagt 3180

catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttgtt 3240

caatcatgcg aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc 3300

agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag 3360

agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca 3420

actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg 3480

gatgtgctgc aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgac 3534

<210> 14
<211> 3534
<212> DNA
<213> artificial sequence

<220>
<223> This is the nucleic acid sequence for the operatively linked comp
onents of the 15/27/28 GHRH plasmid.

<400> 14
gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60

accgcggtgg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120

gtgaggaatg gtggggagtt atttttagag cggtgaggaa ggtgggcagg cagcaggtgt 180

tggcgctcta aaaataactc ccgggagtta tttttagagc ggaggaatgg tggacaccca 240

aatatggcga cggttcctca cccgtcgcca tatttgggtg tccgccctcg gccggggccg 300

cattcctggg ggccgggcgg tgctcccgcc cgcctcgata aaaggctccg gggccggcgg 360

cggcccacga gctacccgga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420

ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480

ctgggtgttc ttctttgtga tcctcaccct cagcaacagc tcccactgct ccccacctcc 540

ccctttgacc ctcaggatgc ggcggtatat cgatgccatc ttcaccaaca gctaccggaa 600

ggtgctggcc cagctgtccg cccgcaagct gctccaggac atcctgaaca ggcagcaggg 660

agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720

ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt gccactccag 780

tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct 840

tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt gggaagacaa 900

cctgtagggc ctgcggggtc tattgggaac caagctggag tgcagtggca caatcttggc 960

tcactgcaat ctccgcctcc tgggttcaag cgattctcct gcctcagcct cccgagttgt 1020

tgggattcca ggcatgcatg accaggctca gctaattttt gtttttttgg tagagacggg 1080

gtttcaccat attggccagg ctggtctcca actcctaatc tcaggtgatc tacccacctt 1140

ggcctcccaa attgctggga ttacaggcgt gaaccactgc tcccttccct gtccttctga 1200

ttttaaaata actataccag caggaggacg tccagacaca gcataggcta cctggccatg 1260

cccaaccggt gggacatttg agttgcttgc ttggcactgt cctctcatgc gttgggtcca 1320

ctcagtagat gcctgttgaa ttcgataccg tcgacctcga ggggggggcc ggtaccagct 1380

tttgttccct ttagtgaggg ttaatttcga gcttggcgta atcatggtca tagctgtttc 1440

ctgtgtgaaa ttgttatccg ctcacaattc cacacaacat acgagccgga agcataaagt 1500

gtaaagcctg gggtgcctaa tgagtgagct aactcacatt aattgcgttg cgctcactgc 1560

ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg 1620

ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct 1680

cggtcgttcg gctgcggcga gcggtatcag ctcactcaaa ggcggtaata cggttatcca 1740

cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga 1800

accgtaaaaa ggccgcgttg ctggcgtttt tccataggct ccgcccccct gacgagcatc 1860

acaaaaatcg acgctcaagt cagaggtggc gaaacccgac aggactataa agataccagg 1920

cgtttccccc tggaagctcc ctcgtgcgct ctcctgttcc gaccctgccg cttaccggat 1980

acctgtccgc ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt 2040

atctcagttc ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa ccccccgttc 2100

agcccgaccg ctgcgcctta tccggtaact atcgtcttga gtccaacccg gtaagacacg 2160

acttatcgcc actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg 2220

gtgctacaga gttcttgaag tggtggccta actacggcta cactagaaga acagtatttg 2280

gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg 2340

gcaaacaaac caccgctggt agcggtggtt tttttgtttg caagcagcag attacgcgca 2400

gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagaaga 2460

actcgtcaag aaggcgatag aaggcgatgc gctgcgaatc gggagcggcg ataccgtaaa 2520

gcacgaggaa gcggtcagcc cattcgccgc caagctcttc agcaatatca cgggtagcca 2580

acgctatgtc ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa 2640

agcggccatt ttccaccatg atattcggca agcaggcatc gccatgggtc acgacgagat 2700

cctcgccgtc gggcatgcgc gccttgagcc tggcgaacag ttcggctggc gcgagcccct 2760

gatgctcttc gtccagatca tcctgatcga caagaccggc ttccatccga gtacgtgctc 2820

gctcgatgcg atgtttcgct tggtggtcga atgggcaggt agccggatca agcgtatgca 2880

gccgccgcat tgcatcagcc atgatggata ctttctcggc aggagcaagg tgagatgaca 2940

ggagatcctg ccccggcact tcgcccaata gcagccagtc ccttcccgct tcagtgacaa 3000

cgtcgagcac agctgcgcaa ggaacgcccg tcgtggccag ccacgatagc cgcgctgcct 3060

cgtcctgcag ttcattcagg gcaccggaca ggtcggtctt gacaaaaaga accgggcgcc 3120

cctgcgctga cagccggaac acggcggcat cagagcagcc gattgtctgt tgtgcccagt 3180

catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttgtt 3240

caatcatgcg aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc 3300

agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag 3360

agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca 3420

actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg 3480

gatgtgctgc aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgac 3534

<210> 15
<211> 3534
<212> DNA
<213> artificial sequence

<220>
<223> This is the entire plasmid sequence for wildtype GHRH.

<400> 15
gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60

accgcggtgg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120

gtgaggaatg gtggggagtt atttttagag cggtgaggaa ggtgggcagg cagcaggtgt 180

tggcgctcta aaaataactc ccgggagtta ttttagagc ggaggaatgg tggacaccca 240

aatatggcga cggttcctca cccgtcgcca tatttgggtg tccgccctcg gccggggccg 300

cattcctggg ggccgggcgg tgctcccgcc cgcctcgata aaaggctccg gggccggcgg 360

cggcccacga gctacccgga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420

ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480

ctgggtgttc ttctttgtga tcctcaccct cagcaacagc tcccactgct ccccacctcc 540

ccctttgacc ctcaggatgc ggcggtatgc agatgccatc ttcaccaaca gctaccggaa 600

ggtgctgggc cagctgtccg cccgcaagct gctccaggac atcatgagca ggcagcaggg 660

agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720

ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt gccactccag 780

tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct 840

tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt gggaagacaa 900

cctgtagggc ctgcggggtc tattgggaac caagctggag tgcagtggca caatcttggc 960

tcactgcaat ctccgcctcc tgggttcaag cgattctcct gcctcagcct cccgagttgt 1020

tgggattcca ggcatgcatg accaggctca gctaattttt gtttttttgg tagagacggg 1080

gtttcaccat attggccagg ctggtctcca actcctaatc tcaggtgatc tacccacctt 1140

ggcctcccaa attgctggga ttacaggcgt gaaccactgc tcccttccct gtccttctga 1200

ttttaaaata actataccag caggaggacg tccagacaca gcataggcta cctggccatg 1260

cccaaccggt gggacatttg agttgcttgc ttggcactgt cctctcatgc gttgggtcca 1320

ctcagtagat gcctgttgaa ttcgataccg tcgacctcga ggggggggccc ggtaccagct 1380

tttgttccct ttagtgaggg ttaatttcga gcttggcgta atcatggtca tagctgtttc 1440

ctgtgtgaaa ttgttatccg ctcacaattc cacacaacat acgagccgga agcataaagt 1500

gtaaagcctg gggtgcctaa tgagtgagct aactcacatt aattgcgttg cgctcactgc 1560

ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg 1620

ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct 1680

cggtcgttcg gctgcggcga gcggtatcag ctcactcaaa ggcggtaata cggttatcca 1740

cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga 1800

accgtaaaaa ggccgcgttg ctggcgtttt tccataggct ccgcccccct gacgagcatc 1860

acaaaaatcg acgctcaagt cagaggtggc gaaacccgac aggactataa agataccagg 1920

cgtttccccc tggaagctcc ctcgtgcgct ctcctgttcc gaccctgccg cttaccggat 1980

acctgtccgc ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt 2040

atctcagttc ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa ccccccgttc 2100

agcccgaccg ctgcgccttа tccggtaact atcgtcttga gtccaacccg gtaagacacg 2160

acttatcgcc actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg 2220

gtgctacaga gttcttgaag tggtggccta actacggcta cactagaaga acagtatttg 2280

gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg 2340

gcaaacaaac caccgctggt agcggtggtt tttttgtttg caagcagcag attacgcgca 2400

gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagaaga 2460

actcgtcaag aaggcgatag aaggcgatgc gctgcgaatc gggagcggcg ataccgtaaa 2520

gcacgaggaa gcggtcagcc cattcgccgc caagctcttc agcaatatca cgggtagcca 2580

acgctatgtc ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa 2640

agcggccatt ttccaccatg atattcggca agcaggcatc gccatgggtc acgacgagat 2700

cctcgccgtc gggcatgcgc gccttgagcc tggcgaacag ttcggctggc gcgagcccct 2760

gatgctcttc gtccagatca tcctgatcga caagaccggc ttccatccga gtacgtgctc 2820

gctcgatgcg atgtttcgct tggtggtcga atgggcaggt agccggatca agcgtatgca 2880

gccgccgcat tgcatcagcc atgatggata ctttctcggc aggagcaagg tgagatgaca 2940

ggagatcctg ccccggcact tcgcccaata gcagccagtc ccttcccgct tcagtgacaa 3000

cgtcgagcac agctgcgcaa ggaacgcccg tcgtggccag ccacgatagc cgcgctgcct 3060

cgtcctgcag ttcattcagg gcaccggaca ggtcggtctt gacaaaaaga accgggcgcc 3120

cctgcgctga cagccggaac acggcggcat cagagcagcc gattgtctgt tgtgcccagt 3180

catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttgtt 3240

caatcatgcg aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc 3300

agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag 3360

agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca 3420

actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg 3480

gatgtgctgc aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgac 3534

<210> 16
<211> 4260
<212> DNA
<213> Artificial sequence

<220>
<223> This is the sequence for the pSP-SEAP cDNA construct

<400> 16
ggccgtccgc cttcggcacc atcctcacga cacccaaata tggcgacggg tgaggaatgg 60

tggggagtta tttttagagc ggtgaggaag gtgggcaggc agcaggtgtt ggcgctctaa 120

aaataactcc cgggagttat ttttagagcg gaggaatggt ggacacccaa atatggcgac 180

ggttcctcac ccgtcgccat atttgggtgt ccgccctcgg ccggggccgc attcctgggg 240

gccgggcggt gctcccgccc gcctcgataa aaggctccgg ggccggcggc ggcccacgag 300

ctacccggag gagcgggagg cgccaagctc tagaactagt ggatcccccg ggctgcagga 360

attcgatatc aagcttcgaa tcgcgaattc gcccaccatg ctgctgctgc tgctgctgct 420

gggcctgagg ctacagctct ccctgggcat catcccagtt gaggaggaga acccggactt 480

ctggaaccgc gaggcagccg aggccctggg tgccgccaag aagctgcagc ctgcacagac 540

agccgccaag aacctcatca tcttcctggg cgatgggatg ggggtgtcta cggtgacagc 600

tgccaggatc ctaaaagggc agaagaagga caaactgggg cctgagatac ccctggccat 660

ggaccgcttc ccatatgtgg ctctgtccaa gacatacaat gtagacaaac atgtgccaga 720

cagtggagcc acagccacgg cctacctgtg cggggtcaag ggcaacttcc agaccattgg 780

cttgagtgca gccgcccgct ttaaccagtg caacacgaca cgcggcaacg aggtcatctc 840

cgtgatgaat cgggccaaga aagcagggaa gtcagtggga gtggtaacca ccacacgagt 900

gcagcacgcc tcgccagccg gcacctacgc ccacacggtg aaccgcaact ggtactcgga 960

cgccgacgtg cctgcctcgg cccgccagga ggggtgccag gacatcgcta cgcagctcat 1020

ctccaacatg gacattgacg tgatcctagg tggaggccga aagtacatgt ttcgcatggg 1080

aaccccagac cctgagtacc cagatgacta cagccaaggt gggaccaggc tggacgggaa 1140

gaatctggtg caggaatggc tggcgaagcg ccagggtgcc cggtatgtgt ggaaccgcac 1200

tgagctcatg caggcttccc tggacccgtc tgtgacccat ctcatgggtc tctttgagcc 1260

tggagacatg aaatacgaga tccaccgaga ctccacactg gacccctccc tgatggagat 1320

gacagaggct gccctgcgcc tgctgagcag gaacccccgc ggcttcttcc tcttcgtgga 1380

gggtggtcgc atcgaccatg gtcatcatga aagcagggct taccgggcac tgactgagac 1440

gatcatgttc gacgacgcca ttgagagggc gggccagctc accagcgagg aggacacgct 1500

gagcctcgtc actgccgacc actcccacgt cttctccttc ggaggctacc ccctgcgagg 1560

gagctccatc ttcgggctgg cccctggcaa ggcccgggac aggaaggcct acacggtcct 1620

cctatacgga aacggtccag gctatgtgct caaggacggc gcccggccgg atgttaccga 1680

gagcgagagc gggagccccg agtatcggca gcagtcagca gtgcccctgg acgaagagac 1740

ccacgcaggc gaggacgtgg cggtgttcgc gcgcggcccg caggcgcacc tggttcacgg 1800

cgtgcaggag cagaccttca tagcgcacgt catggccttc gccgcctgcc tggagcccta 1860

caccgcctgc gacctggcgc ccccgccgg caccaccgac gccgcgcacc cgggttactc 1920

tagagtcggg gcggccggcc gcttcgagca gacatgataa gatacattga tgagtttgga 1980

caaaccacaa ctagaatgca gtgaaaaaaa tgctttattt gtgaaatttg tgatgctatt 2040

gctttatttg taaccattat aagctgcaat aaacaagtta acaacaacaa ttgcattcat 2100

tttatgtttc aggttcaggg ggaggtgtgg gaggtttttt aaagcaagta aaacctctac 2160

aaatgtggta aaatcgataa ggatccgtcg accgatgccc ttgagagcct tcaacccagt 2220

cagctccttc cggtgggcgc ggggcatgac tatcgtcgcc gcacttatga ctgtcttctt 2280

tatcatgcaa ctcgtaggac aggtgccggc agcgctcttc cgcttcctcg ctcactgact 2340

cgctgcgctc ggtcgttcgg ctgcggcgag cggtatcagc tcactcaaag gcggtaatac 2400

ggttatccac agaatcaggg gataacgcag gaaagaacat gtgagcaaaa ggccagcaaa 2460

aggccaggaa ccgtaaaaag gccgcgttgc tggcgttttt ccataggctc cgcccccctg 2520

acgagcatca caaaaatcga cgctcaagtc agaggtggcg aaacccgaca ggactataaa 2580

gataccaggc gtttcccccc ggaagctccc tcgtgcgctc tcctgttccg accctgccgc 2640

ttaccggata cctgtccgcc tttctccctt cgggaagcgt ggcgctttct catagctcac 2700

gctgtaggta tctcagttcg gtgtaggtcg ttcgctccaa gctgggctgt gtgcacgaac 2760

ccccgttca gcccgaccgc tgcgccttat ccggtaacta tcgtcttgag tccaacccgg 2820

taagacacga cttatcgcca ctggcagcag ccactggtaa caggattagc agagcgaggt 2880

atgtaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac actagaagga 2940

cagtatttgg tatctgcgct ctgctgaagc cagttacctt cggaaaaaga gttggtagct 3000

cttgatccgg caaacaaacc accgctggta gcggtggttt ttttgtttgc aagcagcaga 3060

ttacgcgcag aaaaaaagga tctcaagaag atcctttgat cttttctacg gggtctgacg 3120

ctcagtggaa cgaaaactca cgttaaggga ttttggtcat gagattatca aaaaggatct 3180

tcacctagat cctttttaaat taaaaatgaa gttttaaatc aatctaaagt atatatgagt 3240

aaacttggtc tgacagttac caatgcttaa tcagtgaggc acctatctca gcgatctgtc 3300

tatttcgttc atccatagtt gcctgactcc ccgtcgtgta gataactacg atacgggagg 3360

gcttaccatc tggccccagt gctgcaatga taccgcgaga cccacgctca ccggctccag 3420

atttatcagc aataaaccag ccagccggaa gggccgagcg cagaagtggt cctgcaactt 3480

tatccgcctc catccagtct attaattgtt gccgggaagc tagagtaagt agttcgccag 3540

ttaatagttt gcgcaacgtt gttgccattg ctacaggcat cgtggtgtca cgctcgtcgt 3600

ttggtatggc ttcattcagc tccggttccc aacgatcaag gcgagttaca tgatccccca 3660

tgttgtgcaa aaaagcggtt agctccttcg gtcctccgat cgttgtcaga agtaagttgg 3720

ccgcagtgtt atcactcatg gttatggcag cactgcataa ttctcttact gtcatgccat 3780

ccgtaagatg cttttctgtg actggtgagt actcaaccaa gtcattctga gaatagtgta 3840

tgcggcgacc gagttgctct tgcccggcgt caatacggga taataccgcg ccacatagca 3900

gaactttaaa agtgctcatc attggaaaac gttcttcggg gcgaaaactc tcaaggatct 3960

taccgctgtt gagatccagt tcgatgtaac ccactcgtgc acccaactga tcttcagcat 4020

cttttacttt caccagcgtt tctgggtgag caaaaacagg aaggcaaaat gccgcaaaaa 4080

agggaataag ggcgacacgg aaatgttgaa tactcatact cttccttttt caatattatt 4140

gaagcattta tcagggttat tgtctcatga gcggatacat atttgaatgt atttagaaaa 4200

ataaacaaat aggggttccg cgcacatttc cccgaaaagt gccacctgac gcgccctgta 4260

<210> 17
<211> 2710
<212> DNA
<213> artificial sequence

<220>
<223> This is a plasmid vector with an analog growth hormone releasing hormone ("GHRH") sequence codon optimized for mouse.

<400> 17
tgtaatacga ctcactatag ggcgaattgg agctccaccg cggtggcggc cgtccgccct 60

cggcaccatc ctcacgacac ccaaatatgg cgacgggtga ggaatggtgg ggagttattt 120

ttagagcggt gaggaaggtg ggcaggcagc aggtgttggc gctctaaaaa taactcccgg 180

gagttatttt tagagcggag gaatggtgga cacccaaata tggcgacggt tcctcacccg 240

tcgccatatt tgggtgtccg ccctcggccg gggccgcatt cctgggggcc gggcggtgct 300

cccgcccgcc tcgataaaag gctccggggc cggcggcggc ccacgagcta cccggaggag 360

cgggaggcgc caagcggatc ccaaggccca actccccgaa ccactcaggg tcctgtggac 420

agctcaccta gctgccatgg tgctctgggt gctctttgtg atcctcatcc tcaccagcgg 480

cagccactgc agcctgcctc ccagccctcc cttcaggatg cagaggcacg tggacgccat 540

cttcaccacc aactacagga agctgctgag ccagctgtac gccaggaagg tgatccagga 600

catcatgaac aagcagggcg agaggatcca ggagcagagg gccaggctga gctgataagc 660

ttatcggggt ggcatccctg tgacccctcc ccagtgcctc tcctggccct ggaagttgcc 720

actccagtgc ccaccagcct tgtcctaata aaattaagtt gcatcatttt gtctgactag 780

gtgtccttct ataatattat ggggtggagg ggggtggtat ggagcaaggg gcaagttggg 840

aagacaacct gtagggctcg agggggggcc cggtaccagc ttttgttccc tttagtgagg 900

gttaatttcg agcttggtct tccgcttcct cgctcactga ctcgctgcgc tcggtcgttc 960

ggctgcggcg agcggtatca gctcactcaa aggcggtaat acggttatcc acagaatcag 1020

gggataacgc aggaaagaac atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa 1080

aggccgcgtt gctggcgttt ttccataggc tccgcccccc tgacgagcat cacaaaaatc 1140

gacgctcaag tcagaggtgg cgaaacccga caggactata aagataccag gcgtttcccc 1200

ctggaagctc cctcgtgcgc tctcctgttc cgaccctgcc gcttaccgga tacctgtccg 1260

cctttctccc ttcgggaagc gtggcgcttt ctcatagctc acgctgtagg tatctcagtt 1320

cggtgtaggt cgttcgctcc aagctgggct gtgtgcacga acccccccgtt cagcccgacc 1380

gctgcgcctt atccggtaac tatcgtcttg agtccaaccc ggtaagacac gacttatcgc 1440

cactggcagc agccactggt aacaggatta gcagagcgag gtatgtaggc ggtgctacag 1500

agttcttgaa gtggtggcct aactacggct acactagaag aacagtattt ggtatctgcg 1560

ctctgctgaa gccagttacc ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa 1620

ccaccgctgg tagcggtggt ttttttgttt gcaagcagca gattacgcgc agaaaaaaag 1680

gatctcaaga agatcctttg atcttttcta cggggctagc gcttagaaga actcatccag 1740

cagacggtag aatgcaatac gttgagagtc tggagctgca ataccataca gaaccaggaa 1800

acggtcagcc cattcaccac ccagttcctc tgcaatgtca cgggtagcca gtgcaatgtc 1860

ctggtaacgg tctgcaacac ccagacgacc acagtcaatg aaaccagaga aacgaccatt 1920

ctcaaccatg atgttcggca ggcatgcatc accatgagta actaccaggt cctcaccatc 1980

cggcatacga gctttcagac gtgcaaacag ttcagccggt gccagaccct gatgttcctc 2040

atccaggtca tcctggtcaa ccagacctgc ttccatacgg gtacgagcac gttcaatacg 2100

atgttttgcc tggtggtcaa acggacaggt agctgggtcc agggtgtgca gacgacgcat 2160

tgcatcagcc atgatagaaa ctttctctgc cggagccagg tgagaagaca gcaggtcctg 2220

acccggaact tcacccagca gcagccagtc acgaccagct tcagtaacta catccagaac 2280

tgcagcacac ggaacaccag tggttgccag ccaagacaga cgagctgctt catcctgcag 2340

ttcattcaga gcaccagaca ggtcagtttt aacaaacaga actggacgac cctgtgcaga 2400

cagacggaaa acagctgcat cagagcaacc aatggtctgc tgtgcccagt cataaccaaa 2460

cagacgttca acccaggctg ccggagaacc tgcatgcaga ccatcctgtt caatcatgcg 2520

aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc agatccttgg 2580

cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag agggcgcccc 2640

agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca actgttggga 2700

agggcgatcg 2710

<210> 18
<211> 2713

<212> DNA
<213> artificial sequence

<220>
<223> This is a plasmid vector with an analog growth hormone releasing hormone ("GHRH") sequence codon optimized for rat.

<400> 18
tgtaatacga ctcactatag ggcgaattgg agctccaccg cggtggcggc cgtccgccct 60

cggcaccatc ctcacgacac ccaaatatgg cgacgggtga ggaatggtgg ggagttattt 120

ttagagcggt gaggaaggtg ggcaggcagc aggtgttggc gctctaaaaa taactcccgg 180

gagttatttt tagagcggag gaatggtgga cacccaaata tggcgacggt tcctcacccg 240

tcgccatatt tgggtgtccg ccctcggccg gggccgcatt cctgggggcc gggcggtgct 300

cccgcccgcc tcgataaaag gctccggggc cggcggcggc ccacgagcta cccggaggag 360

cgggaggcgc caagcggatc ccaaggccca actccccgaa ccactcaggg tcctgtggac 420

agctcaccta gctgccatgg ccctgtgggt gttcttcgtg ctgctgaccc tgaccagcgg 480

aagccactgc agcctgcctc ccagccctcc cttcagggtg cgccggcacg ccgacgccat 540

cttcaccagc agctacagga ggatcctggg ccagctgtac gctaggaagc tcctgcacga 600

gatcatgaac aggcagcagg gcgagaggaa ccaggagcag aggagcaggt tcaactgata 660

agcttatcgg ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt 720

gccactccag tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac 780

taggtgtcct tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt 840

gggaagacaa cctgtagggc tcgaggggggg gcccggtacc agcttttgtt cccttttagtg 900

agggttaatt tcgagcttgg tcttccgctt cctcgctcac tgactcgctg cgctcggtcg 960

ttcggctgcg gcgagcggta tcagctcact caaaggcggt aatacggtta tccacagaat 1020

caggggataa cgcaggaaag aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta 1080

aaaaggccgc gttgctggcg tttttccata ggctccgccc ccctgacgag catcacaaaa 1140

atcgacgctc aagtcagagg tggcgaaacc cgacaggact ataaagatac caggcgtttc 1200

cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc ggatacctgt 1260

ccgcctttct cccttcggga agcgtggcgc tttctcatag ctcacgctgt aggtatctca 1320

gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc gttcagcccg 1380

accgctgcgc cttatccggt aactatcgtc ttgagtccaa cccggtaaga cacgacttat 1440

cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 1500

cagagttctt gaagtggtgg cctaactacg gctacactag aagaacagta tttggtatct 1560

gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac 1620

aaaccaccgc tggtagcggt ggtttttttg tttgcaagca gcagattacg cgcagaaaaa 1680

aaggatctca agaagatcct ttgatctttt ctacggggct agcgcttaga agaactcatc 1740

cagcagacgg tagaatgcaa tacgttgaga gtctggagct gcaataccat acagaaccag 1800

gaaacggtca gcccattcac cacccagttc ctctgcaatg tcacgggtag ccagtgcaat 1860

gtcctggtaa cggtctgcaa cacccagacg accacagtca atgaaaccag agaaacgacc 1920

attctcaacc atgatgttcg gcaggcatgc atcaccatga gtaactacca ggtcctcacc 1980

atccggcata cgagctttca gacgtgcaaa cagttcagcc ggtgccagac cctgatgttc 2040

ctcatccagg tcatcctggt caaccagacc tgcttccata cgggtacgag cacgttcaat 2100

acgatgtttt gcctggtggt caaacggaca ggtagctggg tccagggtgt gcagacgacg 2160

cattgcatca gccatgatag aaactttctc tgccggagcc aggtgagaag acagcaggtc 2220

ctgacccgga acttcaccca gcagcagcca gtcacgacca gcttcagtaa ctacatccag 2280

aactgcagca cacggaacac cagtggttgc cagccaagac agacgagctg cttcatcctg 2340

cagttcattc agagcaccag acaggtcagt tttaacaaac agaactggac gaccctgtgc 2400

agacagacgg aaaacagctg catcagagca accaatggtc tgctgtgccc agtcataacc 2460

aaacagacgt tcaacccagg ctgccggaga acctgcatgc agaccatcct gttcaatcat 2520

gcgaaacgat cctcatcctg tctcttgatc agatcttgat cccctgcgcc atcagatcct 2580

tggcggcaag aaagccatcc agtttacttt gcagggcttc ccaaccttac cagagggcgc 2640

cccagctggc aattccggtt cgcttgctgt ccataaaacc gcccagtcta gcaactgttg 2700

ggaagggcga tcg 2713

<210> 19
<211> 2704
<212> DNA
<213> artificial sequence

<220>
<223> This is a plasmid vector with an analog growth hormone releasing hormone ("GHRH") sequence codon optimized for bovine.

<400> 19
tgtaatacga ctcactatag ggcgaattgg agctccaccg cggtggcggc cgtccgccct 60

cggcaccatc ctcacgacac ccaaatatgg cgacgggtga ggaatggtgg ggagttattt 120

ttagagcggt gaggaaggtg ggcaggcagc aggtgttggc gctctaaaaa taactcccgg 180

gagttatttt tagagcggag gaatggtgga cacccaaata tggcgacggt tcctcacccg 240

tcgccatatt tgggtgtccg ccctcggccg gggccgcatt cctgggggcc gggcggtgct 300

cccgcccgcc tcgataaaag gctccggggc cggcggcggc ccacgagcta cccggaggag 360

cgggaggcgc caagcggatc ccaaggccca actccccgaa ccactcaggg tcctgtggac 420

agctcaccta gctgccatgg tgctgtgggt gttcttcctg gtgaccctga ccctgagcag 480

cggctcccac ggctccctgc cctcccagcc tctgcgcatc cctcgctacg ccgacgccat 540

cttcaccaac agctaccgca aggtgctcgg ccagctcagc gcccgcaagc tcctgcagga 600

catcatgaac cggcagcagg gcgagcgcaa ccaggagcag ggagcctgat aagcttatcg 660

gggtggcatc cctgtgaccc ctccccagtg cctctcctgg ccctggaagt tgccactcca 720

gtgcccacca gccttgtcct aataaaatta agttgcatca ttttgtctga ctaggtgtcc 780

ttctataata ttatggggtg gaggggggtg gtatggagca aggggcaagt tgggaagaca 840

acctgtaggg ctcgaggggg ggcccggtac cagcttttgt tcccttttagt gagggttaat 900

ttcgagcttg gtcttccgct tcctcgctca ctgactcgct gcgctcggtc gttcggctgc 960

ggcgagcggt atcagctcac tcaaaggcgg taatacggtt atccacagaa tcaggggata 1020

acgcaggaaa gaacatgtga gcaaaaggcc agcaaaaggc caggaaccgt aaaaaggccg 1080

cgttgctggc gtttttccat aggctccgcc cccctgacga gcatcacaaa aatcgacgct 1140

caagtcagag gtggcgaaac ccgacaggac tataaagata ccaggcgttt cccccctggaa 1200

gctccctcgt gcgctctcct gttccgaccc tgccgcttac cggatacctg tccgcctttc 1260

tcccttcggg aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcggtgt 1320

aggtcgttcg ctccaagctg ggctgtgtgc acgaacccce cgttcagccc gaccgctgcg 1380

ccttatccgg taactatcgt cttgagtcca acccggtaag acacgactta tcgccactgg 1440

cagcagccac tggtaacagg attagcagag cgaggtatgt aggcggtgct acagagttct 1500

tgaagtggtg gcctaactac ggctacacta gaagaacagt atttggtatc tgcgctctgc 1560

tgaagccagt taccttcgga aaaagagttg gtagctcttg atccggcaaa caaaccaccg 1620

ctggtagcgg tggttttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc 1680

aagaagatcc tttgatcttt tctacggggc tagcgcttag aagaactcat ccagcagacg 1740

gtagaatgca atacgttgag agtctggagc tgcaatacca tacagaacca ggaaacggtc 1800

agcccattca ccacccagtt cctctgcaat gtcacgggta gccagtgcaa tgtcctggta 1860

acggtctgca acacccagac gaccacagtc aatgaaacca gagaaacgac cattctcaac 1920

catgatgttc ggcaggcatg catcaccatg agtaactacc aggtcctcac catccggcat 1980

acgagctttc agacgtgcaa acagttcagc cggtgccaga ccctgatgtt cctcatccag 2040

gtcatcctgg tcaaccagac ctgcttccat acgggtacga gcacgttcaa tacgatgttt 2100

tgcctggtgg tcaaacggac aggtagctgg gtccagggtg tgcagacgac gcattgcatc 2160

agccatgata gaaactttct ctgccggagc caggtgagaa gacagcaggt cctgacccgg 2220

aacttcaccc agcagcagcc agtcacgacc agcttcagta actacatcca gaactgcagc 2280

acacggaaca ccagtggttg ccagccaaga cagacgagct gcttcatcct gcagttcatt 2340

cagagcacca gacaggtcag ttttaacaaa cagaactgga cgaccctgtg cagacagacg 2400

gaaaacagct gcatcagagc aaccaatggt ctgctgtgcc cagtcataac caaacagacg 2460

ttcaacccag gctgccggag aacctgcatg cagaccatcc tgttcaatca tgcgaaacga 2520

tcctcatcct gtctcttgat cagatcttga tcccctgcgc catcagatcc ttggcggcaa 2580

gaaagccatc cagtttactt tgcagggctt cccaacctta ccagagggcg ccccagctgg 2640

caattccggt tcgcttgctg tccataaaac cgcccagtct agcaactgtt gggaagggcg 2700

atcg 2704

<210> 20
<211> 2704
<212> DNA
<213> artificial sequence

<220>
<223> This is a plasmid vector with an analog growth hormone releasing hormone ("GHRH") sequence codon optimized for ovine.

<400> 20
tgtaatacga ctcactatag ggcgaattgg agctccaccg cggtggcggc cgtccgccct 60

cggcaccatc ctcacgacac ccaaatatgg cgacgggtga ggaatggtgg ggagttattt 120

ttagagcggt gaggaaggtg ggcaggcagc aggtgttggc gctctaaaaa taactcccgg 180

gagttatttt tagagcggag gaatggtgga cacccaaata tggcgacggt tcctcacccg 240

tcgccatatt tgggtgtccg ccctcggccg gggccgcatt cctgggggcc gggcggtgct 300

cccgcccgcc tcgataaaag gctccggggc cggcggcggc ccacgagcta cccggaggag 360

cgggaggcgc caagcggatc ccaaggccca actccccgaa ccactcaggg tcctgtggac 420

agctcaccta gctgccatgg tgctgtgggt gttcttcctg gtgaccctga ccctgagcag 480

cggaagccac ggcagcctgc ccagccagcc cctgaggatc cctaggtacg ccgacgccat 540

cttcaccaac agctacagga agatcctggg ccagctgagc gctaggaagc tcctgcagga 600

catcatgaac aggcagcagg gcgagaggaa ccaggagcag ggcgcctgat aagcttatcg 660

gggtggcatc cctgtgaccc ctccccagtg cctctcctgg ccctggaagt tgccactcca 720

gtgcccacca gccttgtcct aataaaatta agttgcatca ttttgtctga ctaggtgtcc 780

ttctataata ttatggggtg gaggggggtg gtatggagca aggggcaagt tgggaagaca 840

acctgtaggg ctcgaggggg ggcccggtac cagcttttgt tcccttttagt gagggttaat 900

ttcgagcttg gtcttccgct tcctcgctca ctgactcgct gcgctcggtc gttcggctgc 960

ggcgagcggt atcagctcac tcaaaggcgg taatacggtt atccacagaa tcaggggata 1020

acgcaggaaa gaacatgtga gcaaaaggcc agcaaaaggc caggaaccgt aaaaaggccg 1080

cgttgctggc gtttttccat aggctccgcc cccctgacga gcatcacaaa aatcgacgct 1140

caagtcagag gtggcgaaac ccgacaggac tataaagata ccaggcgttt cccccctggaa 1200

gctccctcgt gcgctctcct gttccgaccc tgccgcttac cggatacctg tccgcctttc 1260

tcccttcggg aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcggtgt 1320

aggtcgttcg ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc gaccgctgcg 1380

ccttatccgg taactatcgt cttgagtcca acccggtaag acacgactta tcgccactgg 1440

cagcagccac tggtaacagg attagcagag cgaggtatgt aggcggtgct acagagttct 1500

tgaagtggtg gcctaactac ggctacacta gaagaacagt atttggtatc tgcgctctgc 1560

tgaagccagt taccttcgga aaaagagttg gtagctcttg atccggcaaa caaaccaccg 1620

ctggtagcgg tggtttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc 1680

aagaagatcc tttgatcttt tctacggggc tagcgcttag aagaactcat ccagcagacg 1740

gtagaatgca atacgttgag agtctggagc tgcaatacca tacagaacca ggaaacggtc 1800

agcccattca ccacccagtt cctctgcaat gtcacgggta gccagtgcaa tgtcctggta 1860

acggtctgca acacccagac gaccacagtc aatgaaacca gagaaacgac cattctcaac 1920

catgatgttc ggcaggcatg catcaccatg agtaactacc aggtcctcac catccggcat 1980

acgagctttc agacgtgcaa acagttcagc cggtgccaga ccctgatgtt cctcatccag 2040

gtcatcctgg tcaaccagac ctgcttccat acgggtacga gcacgttcaa tacgatgttt 2100

tgcctggtgg tcaaacggac aggtagctgg gtccagggtg tgcagacgac gcattgcatc 2160

agccatgata gaaactttct ctgccggagc caggtgagaa gacagcaggt cctgacccgg 2220

aacttcaccc agcagcagcc agtcacgacc agcttcagta actacatcca gaactgcagc 2280

acacggaaca ccagtggttg ccagccaaga cagacgagct gcttcatcct gcagttcatt 2340

cagagcacca gacaggtcag ttttaacaaa cagaactgga cgaccctgtg cagacagacg 2400

gaaaacagct gcatcagagc aaccaatggt ctgctgtgcc cagtcataac caaacagacg 2460

ttcaacccag gctgccggag aacctgcatg cagaccatcc tgttcaatca tgcgaaacga 2520

tcctcatcct gtctcttgat cagatcttga tcccctgcgc catcagatcc ttggcggcaa 2580

gaaagccatc cagtttactt tgcagggctt cccaacctta ccagagggcg ccccagctgg 2640

caattccggt tcgcttgctg tccataaaac cgcccagtct agcaactgtt gggaagggcg 2700

atcg 2704

<210> 21
<211> 2713
<212> DNA
<213> artificial sequence

<220>
<223> This is a plasmid vector with an analog growth hormone releasing hormone ("GHRH") sequence codon optimized for chicken.

<400> 21
tgtaatacga ctcactatag ggcgaattgg agctccaccg cggtggcggc cgtccgccct 60

cggcaccatc ctcacgacac ccaaatatgg cgacgggtga ggaatggtgg ggagttattt 120

ttagagcggt gaggaaggtg ggcaggcagc aggtgttggc gctctaaaaa taactcccgg 180

gagttatttt tagagcggag gaatggtgga cacccaaata tggcgacggt tcctcacccg 240

tcgccatatt tgggtgtccg ccctcggccg gggccgcatt cctgggggcc gggcggtgct 300

cccgcccgcc tcgataaaag gctccggggc cggcggcggc ccacgagcta cccggaggag 360

cgggaggcgc caagcggatc ccaaggccca actccccgaa ccactcaggg tcctgtggac 420

agctcaccta gctgccatgg ccctgtgggt gttctttgtg ctgctgaccc tgacctccgg 480

aagccactgc agcctgccac ccagcccacc cttccgcgtc aggcgccacg ccgacggcat 540

cttcagcaag gcctaccgca agctcctggg ccagctgagc gcacgcaact acctgcacag 600

cctgatggcc aagcgcgtgg gcagcggact gggagacgag gccgagcccc tgagctgata 660

agcttatcgg ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt 720

gccactccag tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac 780

taggtgtcct tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt 840

gggaagacaa cctgtagggc tcgaggggcg gcccggtacc agcttttgtt ccctttagtg 900

agggttaatt tcgagcttgg tcttccgctt cctcgctcac tgactcgctg cgctcggtcg 960

ttcggctgcg gcgagcggta tcagctcact caaaggcggt aatacggtta tccacagaat 1020

caggggataa cgcaggaaag aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta 1080

aaaaggccgc gttgctggcg tttttccata ggctccgccc ccctgacgag catcacaaaa 1140

atcgacgctc aagtcagagg tggcgaaacc cgacaggact ataaagatac caggcgtttc 1200

cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc ggatacctgt 1260

ccgcctttct cccttcggga agcgtggcgc tttctcatag ctcacgctgt aggtatctca 1320

gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc gttcagcccg 1380

accgctgcgc cttatccggt aactatcgtc ttgagtccaa cccggtaaga cacgacttat 1440

cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 1500

cagagttctt gaagtggtgg cctaactacg gctacactag aagaacagta tttggtatct 1560

gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac 1620

aaaccaccgc tggtagcggt ggtttttttg tttgcaagca gcagattacg cgcagaaaaa 1680

aaggatctca agaagatcct ttgatctttt ctacggggct agcgcttaga agaactcatc 1740

cagcagacgg tagaatgcaa tacgttgaga gtctggagct gcaataccat acagaaccag 1800

gaaacggtca gcccattcac cacccagttc ctctgcaatg tcacgggtag ccagtgcaat 1860

gtcctggtaa cggtctgcaa cacccagacg accacagtca atgaaaccag agaaacgacc 1920

attctcaacc atgatgttcg gcaggcatgc atcaccatga gtaactacca ggtcctcacc 1980

atccggcata cgagctttca gacgtgcaaa cagttcagcc ggtgccagac cctgatgttc 2040

ctcatccagg tcatcctggt caaccagacc tgcttccata cgggtacgag cacgttcaat 2100

acgatgtttt gcctggtggt caaacggaca ggtagctggg tccagggtgt gcagacgacg 2160

cattgcatca gccatgatag aaactttctc tgccggagcc aggtgagaag acagcaggtc 2220

ctgacccgga acttcaccca gcagcagcca gtcacgacca gcttcagtaa ctacatccag 2280

aactgcagca cacggaacac cagtggttgc cagccaagac agacgagctg cttcatcctg 2340

cagttcattc agagcaccag acaggtcagt tttaacaaac agaactggac gaccctgtgc 2400

agacagacgg aaaacagctg catcagagca accaatggtc tgctgtgccc agtcataacc 2460

aaacagacgt tcaacccagg ctgccggaga acctgcatgc agaccatcct gttcaatcat 2520

gcgaaacgat cctcatcctg tctcttgatc agatcttgat cccctgcgcc atcagatcct 2580

tggcggcaag aaagccatcc agtttacttt gcagggcttc ccaaccttac cagagggcgc 2640

cccagctggc aattccggtt cgcttgctgt ccataaaacc gcccagtcta gcaactgttg 2700

ggaagggcga tcg 2713

<210> 22
<211> 55
<212> DNA

<213> artificial sequence

<220>
<223> This is a nucleic acid sequence of a human growth hormone ("hGH") 5' untranslated region ("5' UTR).

<400> 22
caaggcccaa ctccccgaac cactcagggt cctgtggaca gctcacctag ctgcc 55

<210> 23
<211> 782
<212> DNA
<213> artificial sequence

<220>
<223> This is a nucleic acid sequence of a plasmid pUC-18 origin of replication

<400> 23
tcttccgctt cctcgctcac tgactcgctg cgctcggtcg ttcggctgcg gcgagcggta 60

tcagctcact caaaggcggt aatacggtta tccacagaat caggggataa cgcaggaaag 120

aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta aaaaggccgc gttgctggcg 180

tttttccata ggctccgccc ccctgacgag catcacaaaa atcgacgctc aagtcagagg 240

tggcgaaacc cgacaggact ataaagatac caggcgtttc cccctggaag ctccctcgtg 300

cgctctcctg ttccgaccct gccgcttacc ggatacctgt ccgcctttct cccttcggga 360

agcgtggcgc tttctcatag ctcacgctgt aggtatctca gttcggtgta ggtcgttcgc 420

tccaagctgg gctgtgtgca cgaaccccc gttcagcccg accgctgcgc cttatccggt 480

aactatcgtc ttgagtccaa cccggtaaga cacgacttat cgccactggc agcagccact 540

ggtaacagga ttagcagagc gaggtatgta ggcggtgcta cagagttctt gaagtggtgg 600

cctaactacg gctacactag aaggacagta tttggtatct gcgctctgct gaagccagtt 660

accttcggaa aaagagttgg tagctcttga tccggcaaac aaaccaccgc tggtagcggt 720

ggtttttttg tttgcaagca gcagattacg cgcagaaaaa aaggatctca agaagatcct 780

tt 782

<210> 24
<211> 5
<212> DNA
<213> artificial sequence

<220>
<223> This is a NEO ribosomal binding site

<400> 24
tcctc 5

<210> 25
<211> 29
<212> DNA
<213> artificial sequence

<220>
<223> This is a nucleic acid sequence of a prokaryotic PNEO promoter.

<400> 25
accttaccag agggcgcccc agctggcaa 29

<210> 26
<211> 3558
<212> DNA
<213> artificial sequence

<220>
<223> This is the plasmid sequence for the inducible pGR1774 containing human (1-40) GHRH

<400> 26
atgcctggag acgccatcca cgctgttttg acctccatag aagacaccgg gaccgatcca 60

gcctccgcgg ccgggaacgg tgcattggaa cgcggattcc ccgtgttaat taacaggtaa 120

gtgtcttcct cctgtttcct tcccctgcta ttctgctcaa ccttcctatc agaaactgca 180

gtatctgtat ttttgctagc agtaatacta acggttcttt ttttctcttc acaggccacc 240

atgtagaact agtgatccca aggcccaact ccccgaacca ctcagggtcc tgtggacagc 300

tcacctagct gccatggtgc tctgggtgtt cttctttgtg atcctcaccc tcagcaacag 360

ctcccactgc tccccacctc ccccttgac cctcaggatg cggcggtatg cagatgccat 420

cttcaccaac agctaccgga aggtgctggg ccagctgtcc gcccgcaagc tgctccagga 480

catcatgagc aggcagcagg gagagagcaa ccaagagcga ggagcataat gactgcagga 540

attcgatatc aagcttatcg gggtggcatc cctgtgaccc ctccccagtg cctctcctgg 600

ccctggaagt tgccactcca gtgcccacca gccttgtcct aataaaatta agttgcatca 660

ttttgtctga ctaggtgtcc ttctataata ttatggggtg gaggggggtg gtatggagca 720

aggggcaagt tgggaagaca acctgtaggg cctgcggggt ctattgggaa ccaagctgga 780

gtgcagtggc acaatcttgg ctcactgcaa tctccgcctc ctgggttcaa gcgattctcc 840

tgcctcagcc tcccgagttg ttgggattcc aggcatgcat gaccaggctc agctaatttt 900

tgtttttttg gtagagacgg ggtttcacca tattggccag gctggtctcc aactcctaat 960

ctcaggtgat ctacccacct tggcctccca aattgctggg attacaggcg tgaaccactg 1020

ctcccttccc tgtccttctg attttaaaat aactatacca gcaggaggac gtccagacac 1080

agcataggct acctggccat gcccaaccgg tgggacattt gagttgcttg cttggcactg 1140

tcctctcatg cgttgggtcc actcagtaga tgcctgttga attcgatacc gtcgacctcg 1200

agggggggcc cggtaccagc ttttgttccc tttagtgagg gttaatttcg agcttggcgt 1260

aatcatggtc atagctgttt cctgtgtgaa attgttatcc gctcacaatt ccacacaaca 1320

tacgagccgg aagcataaag tgtaaagcct ggggtgccta atgagtgagc taactcacat 1380

taattgcgtt gcgctcactg cccgctttcc agtcgggaaa cctgtcgtgc cagctgcatt 1440

aatgaatcgg ccaacgcgcg gggagaggcg gtttgcgtat tgggcgctct tccgcttcct 1500

cgctcactga ctcgctgcgc tcggtcgttc ggctgcggcg agcggtatca gctcactcaa 1560

aggcggtaat acggttatcc acagaatcag gggataacgc aggaaagaac atgtgagcaa 1620

aaggccagca aaaggccagg aaccgtaaaa aggccgcgtt gctggcgttt ttccataggc 1680

tccgcccccc tgacgagcat cacaaaaatc gacgctcaag tcagaggtgg cgaaacccga 1740

caggactata aagataccag gcgtttcccc ctggaagctc cctcgtgcgc tctcctgttc 1800

cgaccctgcc gcttaccgga tacctgtccg cctttctccc ttcgggaagc gtggcgcttt 1860

ctcatagctc acgctgtagg tatctcagtt cggtgtaggt cgttcgctcc aagctgggct 1920

gtgtgcacga acccccccgtt cagcccgacc gctgcgcctt atccggtaac tatcgtcttg 1980

agtccaaccc ggtaagacac gacttatcgc cactggcagc agccactggt aacaggatta 2040

gcagagcgag gtatgtaggc ggtgctacag agttcttgaa gtggtggcct aactacggct 2100

acactagaag aacagtattt ggtatctgcg ctctgctgaa gccagttacc ttcggaaaaa 2160

gagttggtag ctcttgatcc ggcaaacaaa ccaccgctgg tagcggtggt ttttttgttt 2220

gcaagcagca gattacgcgc agaaaaaaag gatctcaaga agatcctttg atcttttcta 2280

cggggtctga cgctcagaag aactcgtcaa gaaggcgata gaaggcgatg cgctgcgaat 2340

cgggagcggc gataccgtaa agcacgagga agcggtcagc ccattcgccg ccaagctctt 2400

cagcaatatc acgggtagcc aacgctatgt cctgatagcg gtccgccaca cccagccggc 2460

cacagtcgat gaatccagaa aagcggccat tttccaccat gatattcggc aagcaggcat 2520

cgccatgggt cacgacgaga tcctcgccgt cgggcatgcg cgccttgagc ctggcgaaca 2580

gttcggctgg cgcgagcccc tgatgctctt cgtccagatc atcctgatcg acaagaccgg 2640

cttccatccg agtacgtgct cgctcgatgc gatgtttcgc ttggtggtcg aatgggcagg 2700

tagccggatc aagcgtatgc agccgccgca ttgcatcagc catgatggat actttctcgg 2760

caggagcaag gtgagatgac aggagatcct gccccggcac ttcgcccaat agcagccagt 2820

ccctttcccgc ttcagtgaca acgtcgagca cagctgcgca aggaacgccc gtcgtggcca 2880

gccacgatag ccgcgctgcc tcgtcctgca gttcattcag ggcaccggac aggtcggtct 2940

tgacaaaaag aaccgggcgc ccctgcgctg acagccggaa cacggcggca tcagagcagc 3000

cgattgtctg ttgtgcccag tcatagccga atagcctctc cacccaagcg gccggagaac 3060

ctgcgtgcaa tccatcttgt tcaatcatgc gaaacgatcc tcatcctgtc tcttgatcag 3120

atcttgatcc cctgcgccat cagatccttg gcggcaagaa agccatccag tttactttgc 3180

agggcttccc aaccttacca gagggcgccc cagctggcaa ttccggttcg cttgctgtcc 3240

ataaaaccgc ccagtctagc aactgttggg aagggcgatc ggtgcgggcc tcttcgctat 3300

tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagttgggta acgccagggt 3360

tttcccagtc acgacgttgt aaaacgacgg ccagtgaatt gtaatacgac tcactatagg 3420

gcgaattaat tcgagcttgc atgcctgcag ggtcgaagcg gagtactgtc ctccgagtgg 3480

agtactgtcc tccgagcgga gtactgtcct ccgagtcgag ggtcgaagcg gagtactgtc 3540

ctccgagtgg agtactgt 3558

<210> 27
<211> 4855
<212> DNA
<213> artificial Sequence

<220>
<223> This is the sequence for the muscle-specific GeneSwitch plasmid, pGS1633

<400> 27
aggggccgct ctagctagag tctgcctgcc ccctgcctgg cacagcccgt acctggccgc 60

acgctccctc acaggtgaag ctcgaaaact ccgtccccgt aaggagcccc gctgcccccc 120

gaggcctcct ccctcacgcc tcgctgcgct cccggctccc gcacggccct gggagaggcc 180

cccaccgctt cgtccttaac gggcccggcg gtgccggggg attatttcgg ccccggcccc 240

gggggggccc ggcagacgct ccttatacgg cccggcctcg ctcacctggg ccgcggccag 300

gagcgccttc tttgggcagc gccgggccgg ggccgcgccg ggcccgacac ccaaatatgg 360

cgacggccgg ggccgcattc ctgggggccg ggcggtgctc ccgcccgcct cgataaaagg 420

ctccggggcc ggcgggcgac tcagatcgcc tggagacgcc atccacgctg ttttgacctc 480

catagaagac accgggaccg atccagcctc cgcggccggg aacggtgcat tggaacgcgg 540

attccccgtg ttaattaaca ggtaagtgtc ttcctcctgt ttccttcccc tgctattctg 600

ctcaaccttc ctatcagaaa ctgcagtatc tgtatttttg ctagcagtaa tactaacggt 660

tctttttttc tcttcacagg ccaccaagct accggtccac catggactcc cagcagccag 720

atctgaagct actgtcttct atcgaacaag catgcgatat ttgccgactt aaaaagctca 780

agtgctccaa agaaaaaccg aagtgcgcca agtgtctgaa gaacaactgg gagtgtcgct 840

actctcccaa aaccaaaagg tctccgctga ctagggcaca tctgacagaa gtggaatcaa 900

ggctagaaag actggaacag ctatttctac tgatttttcc tcgagaccag aaaaagttca 960

ataaagtcag agttgtgaga gcactggatg ctgttgctct cccacagcca gtgggcgttc 1020

caaatgaaag ccaagcccta agccagagat tcacttttтc accaggtcaa gacatacagt 1080

tgattccacc actgatcaac ctgttaatga gcattgaacc agatgtgatc tatgcaggac 1140

atgacaacac aaaacctgac acctccagtt ctttgctgac aagtcttaat caactaggcg 1200

agaggcaact tctttcagta gtcaagtggt ctaaatcatt gccaggtttt cgaaacttac 1260

atattgatga ccagataact ctcattcagt attcttggat gagcttaatg gtgtttggtc 1320

taggatggag atcctacaaa cacgtcagtg ggcagatgct gtattttgca cctgatctaa 1380

tactaaatga acagcggatg aaagaatcat cattctattc attatgcctt accatgtggc 1440

agatcccaca ggagtttgtc aagcttcaag ttagccaaga agagttcctc tgtatgaaag 1500

tattgttact tcttaataca attcctttgg aagggctacg aagtcaaacc cagtttgagg 1560

agatgaggtc aagctacatt agagagctca tcaaggcaat tggtttgagg caaaaaggag 1620

ttgtgtcgag ctcacagcgt ttctatcaac ttacaaaact tcttgataac ttgcatgatc 1680

ttgtcaaaca acttcatctg tactgcttga atacatttat ccagtcccgg gcactgagtg 1740

ttgaatttcc agaaatgatg tctgaagtta ttgctgggtc gacgcccatg gaattccagt 1800

acctgccaga tacagacgat cgtcaccgga ttgaggagaa acgtaaaagg acatatgaga 1860

ccttcaagag catcatgaag aagagtcctt tcagcggacc caccgacccc cggcctccac 1920

ctcgacgcat tgctgtgcct tcccgcagct cagcttctgt ccccaagcca gcaccccagc 1980

cctatcccct tacgtcatcc ctgagcacca tcaactatga tgagtttccc accatggtgt 2040

ttccttctgg gcagatcagc caggcctcgg ccttggcccc ggcccctccc caagtcctgc 2100

cccaggctcc agcccctgcc cctgctccag ccatggtatc agctctggcc caggccccag 2160

cccctgtccc agtcctagcc ccaggccctc ctcaggctgt ggccccacct gccccccaagc 2220

ccacccaggc tggggaagga acgctgtcag aggccctgct gcagctgcag tttgatgatg 2280

aagacctggg ggccttgctt ggcaacagca cagacccagc tgtgttcaca gacctggcat 2340

ccgtcgacaa ctccgagttt cagcagctgc tgaaccaggg catacctgtg gcccccccaca 2400

caactgagcc catgctgatg gagtaccctg aggctataac tcgcctagtg acaggggccc 2460

agaggccccc cgacccagct cctgctccac tgggggcccc ggggctcccc aatggcctcc 2520

tttcaggaga tgaagacttc tcctccattg cggacatgga cttctcagcc ctgctgagtc 2580

agatcagctc ctaaggatcc tccggactag aaaagccgaa ttctgcagga attgggtggc 2640

atccctgtga cccctcccca gtgcctctcc tggccctgga agttgccact ccagtgccca 2700

ccagccttgt cctaataaaa ttaagttgca tcattttgtc tgactaggtg tccttctata 2760

atattatggg gtggaggggg gtggtatgga gcaaggggca agttgggaag acaacctgta 2820

gggctcgagg gggggcccgg taccagcttt tgttcccttt agtgagggtt aatttcgagc 2880

ttggcgtaat catggtcata gctgtttcct gtgtgaaatt gttatccgct cacaattcca 2940

cacaacatac gagccggaag cataaagtgt aaagcctggg gtgcctaatg agtgagctaa 3000

ctcacattaa ttgcgttgcg ctcactgccc gctttccagt cgggaaacct gtcgtgccag 3060

ctgcattaat gaatcggcca acgcgcgggg agaggcggtt tgcgtattgg gcgctcttcc 3120

gcttcctcgc tcactgactc gctgcgctcg gtcgttcggc tgcggcgagc ggtatcagct 3180

cactcaaagg cggtaatacg gttatccaca gaatcagggg ataacgcagg aaagaacatg 3240

tgagcaaaag gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc 3300

cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 3360

aacccgacag gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct 3420

cctgttccga ccctgccgct taccggatac ctgtccgcct ttctcccttc gggaagcgtg 3480

gcgctttctc atagctcacg ctgtaggtat ctcagttcgg tgtaggtcgt tcgctccaag 3540

ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct gcgccttatc cggtaactat 3600

cgtcttgagt ccaacccggt aagacacgac ttatcgccac tggcagcagc cactggtaac 3660

aggattagca gagcgaggta tgtaggcggt gctacagagt tcttgaagtg gtggcctaac 3720

tacggctaca ctagaaggac agtatttggt atctgcgctc tgctgaagcc agttaccttc 3780

ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cggtggtttt 3840

tttgtttgca agcagcagat tacgcgcaga aaaaaaggat ctcaagaaga tcctttgatc 3900

ttttctacgg ggtctgacgc tcagaagaac tcgtcaagaa ggcgatagaa ggcgatgcgc 3960

tgcgaatcgg gagcggcgat accgtaaagc acgaggaagc ggtcagccca ttcgccgcca 4020

agctcttcag caatatcacg ggtagccaac gctatgtcct gatagcggtc cgccacaccc 4080

agccggccac agtcgatgaa tccagaaaag cggccatttt ccaccatgat attcggcaag 4140

caggcatcgc catgcgtcac gacgagatcc tcgccgtcgg gcatgcgcgc cttgagcctg 4200

gcgaacagtt cggctggcgc gagcccctga tgctcttcgt ccagatcatc ctgatcgaca 4260

agaccggctt ccatccgagt acgtgctcgc tcgatgcgat gtttcgcttg gtggtcgaat 4320

gggcaggtag ccggatcaag cgtatgcagc cgccgcattg catcagccat gatggatact 4380

ttctcggcag gagcaaggtg agatgacagg agatcctgcc ccggcacttc gcccaatagc 4440

agccagtccc ttcccgcttc agtgacaacg tcgagcacag ctgcgcaagg aacgcccgtc 4500

gtggccagcc acgatagccg cgctgcctcg tcctgcagtt cattcagggc accggacagg 4560

tcggtcttga caaaaagaac cgggcgcccc tgcgctgaca gccggaacac ggcggcatca 4620

gagcagccga ttgtctgttg tgcccagtca tagccgaata gcctctccac ccaagcggcc 4680

ggagaacctg cgtgcaatcc atcttgttca atcatgcgaa acgatcctca tcctgtctct 4740

tgatcagatc ttgatccccc gcgccatcag atccttggcg gcaagaaagc catccagttt 4800

actttgcagg gcttcccaac cttaccagag ggcgaattcg agcttgcatg cctgc 4855