




[図1]




[図2]



[図3]




治療前



治療5ヶ月



[図4A]



[図4B]

ASC 0.1 μg 投与群
n=4


①初回			②6~9ヶ月後			③2年1ヶ月~3年		
3.15	1.65	3.18	4.08	2.70	4.08	3.67	2.53	3.73
3.33	1.75	3.38	4.09	2.75	4.03	3.63	2.50	3.77
1.80	1.75	1.83	2.08	2.90	2.08	1.77	2.67	1.87
↓			↓			↓		
②-①			③-②			③-①		
0.93	1.05	0.90	-0.41	-0.17	-0.34	0.52	0.88	0.56
0.75	1.00	0.65	-0.44	-0.25	-0.26	0.31	0.75	0.39
0.28	1.15	0.25	-0.31	-0.23	-0.21	-0.03	0.92	0.04
			↙					
			③-①					
			0.52	0.88	0.56			
			0.31	0.75	0.39			
			-0.03	0.92	0.04			

[図4C]


ASC 0.4 μg 投与群
n=8

①初回	→			②6～9ヶ月後	→			③2年1ヶ月～3年		
4.13	3.05	3.98		4.61	3.69	4.43		4.66	3.83	4.64
3.89	3.05	3.71	→	4.43	3.05	4.28	→	4.64	3.41	4.57
3.00	2.73	3.13		3.50	3.26	3.58		4.33	3.87	4.34
↓				↓						
②-①				③-②						
0.49	0.64	0.45		0.04	0.14	0.22				
0.54	0.00	0.56		0.22	0.36	0.30				
0.50	0.53	0.45		0.83	0.61	0.76				


[図5A]




[図5B]




[図5C]



[図6]




[図7]



[図8A]

患者C

レンズ計測例(C)



[図8B]

