

and execute relevant motor and cognitive patterns while suppressing unwanted or irrelevant patterns (Graybiel, A.M. Curr. Biol. 2000, 10, R509–R511 (2000).

Papaverine, a relatively specific PDE10A inhibitor, and PDE10A-knockout mice have been used to explore the physiology of this enzyme and the possible therapeutic utility of PDE10A inhibition. Inhibition of this enzyme pharmacologically or through gene disruption causes a reduction in activity and a reduced response to psychomotor stimulants. Inhibition also reduces the conditioned avoidance response, a behavioural response that is predictive of clinical antipsychotic activity (Siuciak, J.A.; et al., Neuropharmacology 2006, 51 (2), 386-396; Siuciak, J.A.; et al., Neuropharmacology 2006, 51 (2), 374-385).

In addition PDE10A inhibition bears the potential to improve the negative and cognitive symptoms associated to schizophrenia. Indeed papaverine have been shown to attenuate the deficits in the extra-dimensional shift learning induced in rats by sub-chronic treatment with PCP, an animal paradigm of NMDA receptor hypofunction (Rodefer, J.S., et al., Eur. J. Neuroscience 2005, 2,: 1070-1076). In addition increased social interaction in PDE10A2-deficient mice have been observed (Sano, H. J. Neurochem. 2008, 105, 546-556).

Diseases that can be treated with PDE10A inhibitors include, but are not limited to, diseases thought to be mediated in part by dysfunction of the basal ganglia, of other parts of the central nervous system and of other PDE10A expressing tissues. In particular, diseases can be treated, where inhibition of PDE10A can have therapeutic effects.

These diseases include, but are not limited to, certain psychotic disorders such as schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder or substance-induced psychotic disorder, anxiety disorders such as panic disorder, obsessive-compulsive disorder, acute stress disorder or generalized anxiety disorder, obsessive/compulsive disorders, drug addictions, movement disorders such as Parkinson's disease or restless leg syndrome, cognition deficiency disorders such as Alzheimer's disease or multi-infarct dementia, mood disorders such as depression or bipolar disorders, or neuropsychiatric conditions such as psychosis, attention-deficit/hyperactivity disorder (ADHD) or related attentional disorders.


The compounds of the present ~~invention~~ disclosure are also suitable for the treatment of diabetes and related disorders such as obesity by regulating the cAMP signaling system.

PDE10A inhibitors might also be useful in preventing neurons from undergoing apoptosis by raising cAMP and cGMP levels and, thus, might possess anti-inflammatory properties. Neurodegenerative disorders treatable with PDE10A inhibitors include, but are not

limited to, as Alzheimer's disease, Huntington's disease, Parkinson's disease, multiple sclerosis, stroke or spinal cord injury.

The growth of cancer cells is inhibited by cAMP and cGMP. Thus by raising cAMP and cGMP, PDE10A inhibitors can also be used for the treatment of different solid tumors and hematological malignancies such as renal cell carcinoma or breast cancer.

According to a first aspect, there is provided a compound of formula (I)



wherein

R^1 is aryl, heteroaryl or NR^4R^5 , wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;

R^2 is hydrogen, halogen or C_1 - C_7 alkyl;

R^3 is aryl or heteroaryl, wherein said aryl and said heteroaryl can be substituted by 1 to 3 substituents independently selected from the group consisting of C_1 - C_7 alkyl, hydroxyl, halogen, $-C(O)-NR^6R^7$ and $-C(O)-O-R^8$;

R^4 and R^5 are independently C_1 - C_7 alkyl, C_3 - C_8 cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;

R^6 and R^7 , are independently C_1 - C_7 alkyl, C_3 - C_8 cycloalkyl, together with the nitrogen atom to which they are attached, form a heterocycloalkyl which can be substituted by 1 to 3 substituents independently selected from the group consisting of hydroxyl, halogen, C_1 - C_7 alkyl, C_1 - C_7 alkoxy and C_1 - C_7 haloalkyl;


R^8 is hydrogen, C_1 - C_7 alkyl, cycloalkyl;

or a pharmaceutically acceptable salt thereof.

According to a second aspect, there is provided a pharmaceutical composition comprising a compound as defined above and a therapeutically inert carrier.


According to a third aspect, there is provided a process for the manufacture of compounds of formula I as defined above, which process comprises

- 5 a) reacting a compound of formula (2)



with

- b) a compound of formula (3)



- 10 wherein R¹, R², R³ are as defined above and, if desired, converting the compounds into pharmaceutically acceptable salts thereof.

- According to a fourth aspect, there is provided use of a compound as defined above for the preparation of a medicament for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia,
15 delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder,
20 attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

- According to a fifth aspect, there is provided a compound as defined above for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or
25 cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct

dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

DETAILED DESCRIPTION OF THE INVENTION

Unless otherwise indicated, the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

It must be noted that, as used in the specification and the claims, the singular forms "a", "an" and "the" include plural referents unless the context clearly dictates otherwise.

When indicating the number of substituents, the term "one or more" means from one substituent to the highest possible number of substitution, i.e. replacement of one hydrogen up to replacement of all hydrogens by substituents.


The term "optional" or "optionally" denotes that a subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not.

The term "halogen" refers to fluorine, chlorine, bromine and iodine, more specifically fluorine, chlorine and bromine.

The term "alkyl" denotes a monovalent linear or branched saturated hydrocarbon group of 1 to 12 carbon atoms. In particular embodiments, alkyl has 1 to 7 carbon atoms, and in more particular embodiments 1 to 4 carbon atoms. Examples of alkyl include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, or tert-butyl.


The term "aromatic" denotes the conventional idea of aromaticity as defined in the literature, in particular in IUPAC - Compendium of Chemical Terminology, 2nd, A. D. McNaught & A. Wilkinson (Eds). Blackwell Scientific Publications, Oxford (1997).

The term "cycloalkyl" denotes a monovalent saturated monocyclic or bicyclic hydrocarbon group of 3 to 10 ring carbon atoms. In particular embodiments cycloalkyl denotes a monovalent saturated monocyclic hydrocarbon group of 3 to 8 ring carbon atoms. Bicyclic means consisting of two saturated carbocycles having one or more carbon atoms in common. Particular cycloalkyl groups are monocyclic. Examples for monocyclic cycloalkyl are



with

b) a compound of formula (3)



wherein R^1 , R^2 , R^3 are as defined above and, if desired, converting the compounds into pharmaceutically acceptable salts thereof.

~~10. The use of a compound according to any one of claims 1 - 7 for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.~~

~~11.10.~~ The use of a compound according to any one of claims 1 - 7 for the preparation of a medicament for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

1211. A compound according to any one of claims 1 - 7 for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer.

~~13. — A method for the treatment or prophylaxis of psychotic disorders, schizophrenia, positive, negative and/or cognitive symptoms associated with schizophrenia, delusional disorder, substance-induced psychotic disorder, anxiety disorders, panic disorder, obsessive/compulsive disorders, acute stress disorder, generalized anxiety disorder, drug addictions, movement disorders, Parkinson's disease, restless leg syndrome, cognition deficiency disorders, Alzheimer's disease, multi-infarct dementia, mood disorders, depression, bipolar disorders, neuropsychiatric conditions, psychosis, attention-deficit/hyperactivity disorder, attentional disorders, diabetes and related disorders, type 2 diabetes mellitus, neurodegenerative disorders, Huntington's disease, multiple sclerosis, stroke, spinal cord injury, solid tumors, hematological malignancies, renal cell carcinoma or breast cancer, which method comprises administering an effective amount of a compound as defined in any one of claims 1 — 7.~~