

**CRYSTALS OF LAQUINIMOD SODIUM, AND PROCESS FOR THE
MANUFACTURE THEREOF**

5

This application claims the benefit of U.S. Provisional Application No. 60/728,657, filed October 19, 2005, the entire contents of which are hereby incorporated by reference.

10

Throughout this application various publications, published patent applications, and published patents are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

15

Background of the Invention

Laquinimod is a compound which has been shown to be effective in the acute experimental autoimmune encephalomyelitis (aEAE) model (U.S. Patent No. 6,077,851). Its chemical name is N-ethyl-N-phenyl-1,2,-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxoquinoline-3-carboxamide, and its Chemical Registry number is 248281-84-7. The processes of synthesis of laquinimod and the preparation of its sodium salt are disclosed in U.S. Patent No. 6,077,851. An additional process of synthesis of laquinimod is disclosed in U.S. Patent No. 6,875,869.

20

25

In the preparation of laquinimod sodium disclosed in U.S. Patent No. 6,077,851, laquinimod acid was suspended in ethanol, and 5M sodium hydroxide solution was added. After stirring, the resulting precipitate was filtered, washed with ethanol, and dried. The method used to make laquinimod sodium in U.S. Patent No. 6,077,851 is commonly referred to as a slurry-to-slurry salt formation.

30

35

In the slurry-to-slurry salt formation method of laquinimod sodium, the laquinimod sodium is not dissolved in solution. Any solid impurities, if present in the laquinimod sodium
5 suspension, are therefore not removed by filtration.

Applicants have found that the slurry-to-slurry formation of laquinimod sodium usually results in a product contaminated with other compounds and/or metals. Disclosed is a process
10 for the preparation of laquinimod sodium which addresses this.

Summary of the Invention

Disclosed is a process for the preparation of laquinimod sodium which removes the impurities present after the salt formation step, thus resulting in crystals of higher purity
5 as well as crystals having improved crystalline characteristics.

The subject invention provides a mixture of crystalline laquinimod sodium particles, wherein 10% or more of the
10 total amount by volume of the laquinimod sodium particles have a size of greater than 40 microns.

The subject invention also provides a mixture of crystalline laquinimod sodium particles, having a tapped density of at
15 least 0.6 g/mL.

The subject invention also provides a composition comprising laquinimod sodium and no more than 2 ppm of a heavy metal calculated based on the total amount of laquinimod sodium in
20 the composition.

The subject invention also provides a process of recrystallization of laquinimod sodium comprising:

- 25 a) dissolving laquinimod sodium in water to form an aqueous solution;
- b) concentrating the aqueous solution to form a concentrated solution;
- c) adding a water-miscible anti-solvent to the concentrated solution to form laquinimod sodium
30 crystals; and
- d) isolating the laquinimod sodium crystals.

The subject invention also provides a process for making a pharmaceutical composition comprising laquinimod sodium
35 comprising:

- a) obtaining a batch of laquinimod sodium;
 - b) determining whether insoluble matter is present in the batch of step a) by mixing a sample from the batch in deionized water at room temperature at a ratio of at least 110 mg of sample to 1.0 ml of water, and inspecting the resulting mixture for the presence of insoluble matter; and
 - c) mixing the batch of step a) with at least one pharmaceutically acceptable carrier if in step b) insoluble matter is determined to be present below a predetermined amount.
- 5
- 10

Brief Description of Figures

Figure 1: Scanning electron micrograph of laquinimod sodium
prepared according to Example 1 (batch B) from
5 Example 14, before recrystallization.

Figure 2: Scanning electron micrograph of recrystallized
crystals from Example 15.

10 **Figure 3:** Scanning electron micrograph of recrystallized
crystals from Example 16.

Detailed Description of the Invention

The subject invention provides a mixture of crystalline laquinimod sodium particles, wherein 10% or more of the
5 total amount by volume of the laquinimod sodium particles have a size of greater than 40 microns.

In an embodiment of the mixture, 50% or more of the total amount by volume of the laquinimod sodium particles have a
10 size of greater than 15 microns.

In a further embodiment, the mixture has a tapped density of at least 0.6 g/mL, at least 0.5 g/mL, or at least 0.4 g/mL.

15 In another embodiment, the mixture has a bulk density of at least 0.4 g/mL, at least 0.3 g/mL, or at least 0.2 g/mL.

In yet another embodiment, the mixture has a tapped density of less than 0.8 g/mL, or less than 0.7 g/mL.
20

In a further embodiment, the mixture comprises no more than 2 ppm of a heavy metal. The heavy metal may be iron, nickel or chromium.

25 In an embodiment, the mixture comprises no more than 2 ppm of iron, no more than 1.5 ppm of iron, or no more than 1 ppm of iron.

In a further embodiment, the mixture comprises no more than
30 0.2 ppm of nickel, no more than 0.15 ppm of nickel, or no more than 0.1 ppm of nickel.

In yet a further embodiment, the mixture comprises no more than 0.3 ppm of chromium, no more than 0.25 ppm of chromium,
35 no more than 0.2 ppm of chromium, no more than 0.15 ppm of

chromium, or no more than 0.1 ppm of chromium.

5 The subject invention also provides a pharmaceutical composition comprising any of the disclosed mixtures and a pharmaceutically acceptable carrier. The pharmaceutical composition may be in the form of a tablet or capsule.

10 The subject invention also provides a composition comprising laquinimod sodium and no more than 2 ppm of a heavy metal calculated based on the total amount of laquinimod sodium in the composition. The heavy metal may be iron, nickel or chromium.

15 In an embodiment, the iron content of the composition is no more than 2 ppm, no more than 1.5 ppm, or no more than 1 ppm.

20 In a further embodiment of the composition, the nickel content is no more than 0.2 ppm, no more than 0.15 ppm, or no more than 0.1 ppm.

25 In yet a further embodiment of the composition, the chromium content is no more than 0.3 ppm, no more than 0.25 ppm, no more than 0.2 ppm, no more than 0.15 ppm, or no more than 0.1 ppm.

30 In another embodiment, the composition is in crystalline form. A composition in the crystalline form may be in the form of any of the disclosed embodiments.

The subject invention also provides a process of recrystallization of laquinimod sodium comprising:

- a) dissolving laquinimod sodium in water to form an aqueous solution;
- 5 b) concentrating the aqueous solution to form a concentrated solution;
- c) adding a water-miscible anti-solvent to the concentrated solution to form laquinimod sodium crystals; and
- 10 d) isolating the laquinimod sodium crystals.

In an embodiment of the process, step a) is performed by heating the aqueous solution to a temperature of 40-80°C.

- 15 In a further embodiment of the process, the concentrated solution comprises 1-4 milliliters of water per gram of laquinimod sodium.

In a further embodiment of the process, the concentrated
20 solution comprises 1-2 milliliters of water per gram of laquinimod sodium.

In another embodiment of the process, the anti-solvent is one, or a mixture of more than one, of the group consisting
25 of ethanol, isopropanol, and acetone.

In another embodiment of the process, the anti-solvent is acetone.

- 30 In yet another embodiment of the process, the anti-solvent is added in an amount between 3 and 15 milliliters per gram of laquinimod sodium.

In a further embodiment of the process, step c) is followed
35 by cooling the solution to a temperature of below 10°C.

In yet a further embodiment of the process, step b) is followed by seeding the concentrated solution with laquinimod sodium.

5

The subject invention also provides laquinimod sodium prepared by any one of the disclosed processes.

A process for making a pharmaceutical composition comprising
10 laquinimod sodium comprising:

- a) obtaining a batch of laquinimod sodium;
- b) determining whether insoluble matter is present in the batch of step a) by mixing a sample from the batch in deionized water at room temperature
15 at a ratio of at least 110 mg of sample to 1.0 ml of water, and inspecting the resulting mixture for the presence of insoluble matter; and
- c) mixing the batch of step a) with at least one pharmaceutically acceptable carrier if in step b)
20 insoluble matter is determined to be present below a predetermined amount.

In one embodiment of the process, if insoluble matter in the mixture of step b) is determined not to be present below a
25 predetermined amount, the process further comprises:

- d) dissolving the batch of step a) in water to form an aqueous solution;
- e) filtering the aqueous solution of step d) to reduce the amount of insoluble matter to below
30 the predetermined amount;
- f) concentrating the aqueous solution of step e) to form a concentrated solution;
- g) adding a water-miscible anti-solvent to the concentrated solution of step f) to form
35 laquinimod sodium crystals; and

- h) isolating the laquinimod sodium crystals of step g).

As used herein, a "pharmaceutically acceptable" component is one that is suitable for use with humans and/or animals without undue adverse side effects (such as toxicity, irritation, and allergic response) commensurate with a reasonable benefit/risk ratio.

Thus, a "pharmaceutically acceptable carrier" is a pharmaceutically acceptable solvent, suspending agent or vehicle, for delivering the instant compounds to the animal or human. The carrier is selected with the planned manner of administration in mind. Liposomes are also a pharmaceutical carrier.

A dosage unit may comprise a single compound or mixtures of compounds thereof. A dosage unit can be prepared for oral dosage forms, such as tablets, capsules, pills, powders, and granules.

Drug substance can be administered in admixture with suitable pharmaceutical diluents, extenders, excipients, or carriers (collectively referred to herein as a pharmaceutically acceptable carrier) suitably selected with respect to the intended form of administration and as consistent with conventional pharmaceutical practices. The unit will be in a form suitable for oral administration. The drug substance can be administered alone but are generally mixed with a pharmaceutically acceptable carrier, and co-administered in the form of a tablet or capsule, liposome, or as an agglomerated powder. Examples of suitable solid carriers include lactose, sucrose, gelatin and agar. Capsule or tablets can be easily formulated and can be made easy to swallow or chew; other solid forms

include granules, and bulk powders. Tablets may contain suitable binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents flow-inducing agents, and melting agents.

5

Specific examples of the techniques, pharmaceutically acceptable carriers and excipients that may be used to formulate oral dosage forms of the present invention are described, e.g., in U.S. Patent Application Publication No. 10 2005/0192315. For instance, the oral dosage form of the present invention may comprise an alkaline-reacting component, said component preferably amounting from about 1 to 20% by weight of the formulation in order to keep the pH above 8.

15

General techniques and compositions for making dosage forms useful in the present invention are described in the following references: 7 Modern Pharmaceutics, Chapters 9 and 10 (Banker & Rhodes, Editors, 1979); Pharmaceutical 20 Dosage Forms: Tablets (Lieberman et al., 1981); Ansel, Introduction to Pharmaceutical Dosage Forms 2nd Edition (1976); Remington's Pharmaceutical Sciences, 17th ed. (Mack Publishing Company, Easton, Pa., 1985); Advances in Pharmaceutical Sciences (David Ganderton, Trevor Jones, 25 Eds., 1992); Advances in Pharmaceutical Sciences Vol 7. (David Ganderton, Trevor Jones, James McGinity, Eds., 1995); Aqueous Polymeric Coatings for Pharmaceutical Dosage 30 Forms (Drugs and the Pharmaceutical Sciences, Series 36 (James McGinity, Ed., 1989); Pharmaceutical Particulate Carriers: Therapeutic Applications: Drugs and the Pharmaceutical Sciences, Vol 61 (Alain Rolland, Ed., 1993); Drug Delivery to the Gastrointestinal Tract (Ellis Horwood Books in the Biological Sciences. Series in Pharmaceutical 35 Technology; J. G. Hardy, S. S. Davis, Clive G. Wilson,

Eds.); Modern Pharmaceutics Drugs and the Pharmaceutical Sciences, Vol 40 (Gilbert S. Banker, Christopher T. Rhodes, Eds.).

5 Tablets may contain suitable binders, lubricants, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. For instance, for oral administration in the dosage unit form of a tablet or capsule, the active drug component can be combined with an
10 oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, gelatin, agar, starch, sucrose, glucose, methyl cellulose, dicalcium phosphate, calcium sulfate, mannitol, sorbitol, microcrystalline cellulose and the like. Suitable binders include starch, gelatin, natural
15 sugars such as glucose or beta-lactose, corn starch, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, povidone, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium
20 stearate, sodium benzoate, sodium acetate, sodium chloride, stearic acid, sodium stearyl fumarate, talc and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, croscarmellose sodium, sodium starch glycolate and the like.

25 As used herein, an "anti-solvent" is a solvent in which laquinimod sodium is slightly soluble, very slightly soluble, practically insoluble, or insoluble at room temperature (20-25°C). The solubility terms are defined
30 below, in accordance with the United States Pharmacopoeia XXV.

<u>Term</u>	<u>Parts of solvent required for 1 part solute</u>
Slightly soluble	From 100 to 1000
Very slightly soluble	From 1000 to 10,000
Practically insoluble	10,000 and over
Insoluble	10,000 and over

As used herein, "density" is a measurement defined as the mass of a substance per unit volume.

5

As used herein, "bulk density" or "BD" refers to a density measurement of a loose, uncompacted substance, wherein the volume of the substance includes the air trapped between particles.

10

As used herein, "tapped density" or "TD" refers to a density measurement of a substance that has been tapped or vibrated, thus minimizing the volume of the substance by eliminating or minimizing the air trapped between particles.

15

The purification of impure crystalline compounds is usually attained by recrystallization from a suitable solvent or mixture of solvents. (Vogel's Textbook of Practical Organic Chemistry. 5th edition. Longman Scientific & Technical, 1989.) The recrystallization process generally comprises the following steps: a) dissolving the impure crystalline substance in a suitable solvent near the boiling point; b) filtering the hot solution from particles of insoluble material and dust; c) allowing the hot solution to cool to cause the dissolved substance to crystallize out; and d) separating the crystals from the supernatant solution. (*Id.*)

20

25

However, standard recrystallization techniques were accompanied by low or no yields when applied to laquinimod

sodium. As shown in Examples 1-10, attempts to recrystallize laquinimod sodium resulted in poor yields, if any. The present invention provides an industrially reproducible recrystallization process that results in high yields of
5 laquinimod sodium.

The process of the present invention uses an anti-solvent in which laquinimod sodium is practically insoluble. In addition, the process of the present invention concentrates
10 the laquinimod sodium aqueous solution before the addition of the anti-solvent.

The laquinimod sodium manufactured by the recrystallization processes of the present invention has increased purity over
15 the laquinimod sodium disclosed in the prior art. U.S. Patent No. 6,875,869 discloses a process of preparing the base compound laquinimod in high yield and low level of impurities. However, the process in U.S. Patent No. 6,875,869 is only for synthesis of the base compound and not
20 the salt. As such, the slurry-to-slurry salt formation process would still be needed to form the sodium salt. The slurry-to-slurry salt formation process previously disclosed is not efficient in removing any impurities present in the starting material.

25

A second advantage of the recrystallization process of the present invention is environmentally-friendliness, as water is used as the primary solvent.

30 A third advantage of the recrystallization process of the present invention is that laquinimod sodium crystals of a higher density than the laquinimod sodium crystals disclosed in the prior art are produced. Low tapped density is anathema to certain prized qualities in a drug substance
35 such as compressibility, the ability of a powder to decrease

in volume under pressure, and compactibility, the ability of a powder to be compressed into a tablet of certain strength or hardness. Crystals with low tapped density are also known to have poor flowability, which results in a lack of
5 uniformity of content in finished dosage forms, especially in tablets. (Rudnic et al. Chpt. 45, *Remington's Pharmaceutical Sciences*, 20th Edition, Lippincott Williams & Wilkins, Baltimore, MD. (2000)) Problems of uniformity of content are especially important in tablets in which the
10 amount of active pharmaceutical ingredient within the tablet is low.

A fourth advantage of the recrystallization process of the present invention is that the resulting laquinimod sodium
15 crystals have increased particle size. Larger particles of laquinimod sodium are more processable when making pharmaceutical compositions. Smaller particles are often associated with dust-like properties which may interfere with processing in manufacture of pharmaceutical
20 compositions. In addition, smaller particles are sometimes associated with flowability problems which may interfere with the manufacture of pharmaceutical compositions. Furthermore, in some instances, chemical stability has been shown to be decreased by the increase in surface area that
25 results from smaller particle size. (Felmeister, A. Chpt 88, *Remington's Pharmaceutical Sciences*, 15th Edition, Mack Publishing Company, Easton, PA (1975)).

Experimental Details

Determination of Powder Density

Bulk Density

- 5 1. Mix powder;
2. Tare a 50 ml empty cylinder on a 0.01 g sensitivity balance;
3. Transfer the powder, without compacting, to the cylinder being held at approximately a 45 degree angle
- 10 to achieve an untapped apparent volume of 40 to 50 ml.
4. Bring the cylinder containing the sample to a vertical position by a sharp move in order to level the volume for reading.
5. Read the apparent volume (V_a) to the nearest
- 15 graduated unit;
6. Weigh the cylinder with sample (the balance gives sample weight M);
7. Calculate bulk density in g/ml according to the following equation: $BD = M/V_a$;
- 20 8. Perform steps 1-7 again and report the average data of duplicates.

Tapped Density

1. Put the same cylinder used to calculate Bulk Density
- 25 in a Quantachrome Dual Autotap instrument;
2. Perform 1250 taps;
3. Read the tapped volume (V_f) to the nearest graduated unit;
4. Calculate the tapped density in g/ml according to the
- 30 following equation: $TD = M/V_f$;
5. Perform steps 1-4 again and report the average data of duplicates.

Determination of Particle Size

The particle size distributions were measured by Malvern Laser Diffraction, using the Mastersizer S model. Laser diffraction relies on the fact that diffraction angle of light is inversely proportional to particle size. Properties of particles are measured and interpreted as measurements of a sphere (a sphere being the only shape that can be described by one unique number). In addition, laser diffraction calculates a particle size distribution based around volume terms, thus eliminating particle count from the determination of particle size. The Mastersizer S model measures particles using a single technique and a single range setting.

D(0.1) is the particle size, in microns, below which 10% by volume distribution of the population is found. D(0.5) is the particle size, in microns, below which 50% by volume distribution of the population is found. D(0.9) is the particle size, in microns, below which 90% by volume distribution of the population is found.

Determination of Heavy Metals

Metal content was measured using inductively coupled plasma atomic emission spectrometry using an inductively coupled plasma atomic emission spectrometry ("ICP-AES") system manufactured by Spectro (Kleve, Germany). Sample digestion was performed in 65% nitric acid, and the internal standard used was scandium.

Note: In the following examples the volumes of solvents used are calculated relative to starting weight of laquinimod sodium. The yields are calculated in weight percent.

Determination of Purity

Laquinimod sodium and polar impurity/degradation products were determined by isocratic reversed phase high performance liquid chromatography (RP-HPLC), using an ODS-
5 3V column and a mobile phase comprised of a mixture of ammonium acetate buffer at pH 7.0 (80%) and acetonitrile (20%). The detection technique was ultraviolet absorption at 240nm.

10 Example 1 (method of preparing laquinimod sodium)

Laquinimod acid was prepared according to the method described in U.S. Patent No. 6,875,869: 5-Chloro-1,2-dihydro-4-hydroxy-1-methyl-2-oxo-quinoline-3-carboxylic
acid methyl ester (3.0 g), N-ethylaniline (2 eq 2-2.88 ml),
15 and heptane (60 ml) were heated and the volatiles, mainly heptane and formed methanol, (32 ml) distilled off during 6 hours and 35 minutes. After cooling to room temperature the crystalline suspension was filtered and the crystals were washed with heptane and dried in vacuum to yield laquinimod
20 acid (3.94 g, 98%) as white to off-white crystals.

Laquinimod acid was converted into laquinimod sodium using the method described in U.S. Patent No. 6,077,851, Example 2: A solution of 5 M sodium hydroxide was prepared by
25 dilution of a 50% by weight sodium hydroxide solution (10.0 g) with sterile water to the total volume of 25 ml. N-Ethyl-N-phenyl-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxo-quinoline-3-carboxamide (10.0 g) was suspended in ethanol (150 ml) and the previously prepared 5 M sodium
30 hydroxide solution was added to pH of 8-12 (5.6 ml). The reaction mixture was stirred for another 30 minutes at ambient temperature. The resulting precipitation was filtered off and rapidly washed twice with ethanol (2x150 ml). The precipitate was then dried in vacuum over P₂O₅ to
35 give the title compound (9.5 g), yield 90%. This process is

known as a "slurry-to-slurry process."

Example 2

5 Laquinimod sodium prepared according to Example 1 was added
to 6.1 volumes of water at 50°C. The pH was adjusted to 12.5
by the addition of NaOH and the mixture was stirred until
complete dissolution. 50.0 volumes of ethanol were added.
The solution was cooled to 2°C but no crystallization
occurred.

10

Example 3

Laquinimod sodium prepared according to Example 1 was added
to 6.1 volumes of water at 50°C. The pH was adjusted to 12.5
by the addition of NaOH and the mixture was stirred until
15 complete dissolution. 100.0 volumes of ethanol were added.
The solution was cooled to -18°C but no crystallization
occurred.

Example 4

20 Laquinimod sodium prepared according to Example 1 was added
to 6.1 volumes of water at 50°C. The pH was adjusted to 12.5
by the addition of NaOH and the mixture was stirred until
complete dissolution. 50.0 volumes of ethanol were added.
The solution was cooled to -18°C but no crystallization
25 occurred.

Example 5

Laquinimod sodium prepared according to Example 1 was added
to 6.1 volumes of water at 50°C. The pH was adjusted to 12.5
30 by the addition of NaOH and the mixture was stirred until
complete dissolution. 50.0 volumes of ethanol were added.
The solution was acidified to a pH of 5.0 by the addition of
HCl. The solution was cooled to 4°C, and crystallization
occurred. The compound which crystallized was filtered and
35 washed with 20 mL of ethanol: water solution 1:1 and was

dried at 50°C under vacuum to a constant weight and was determined to be laquinimod acid, in a yield of 56.2%.

Discussion of Examples 2-5

5 In Examples 2-5, recrystallization was attempted by dissolving the laquinimod sodium in a small amount of water and by addition of ethanol as an anti-solvent. Although the solubility of laquinimod sodium in ethanol is low (laquinimod sodium is slightly soluble in ethanol at room
10 temperature), nevertheless, no crystallization of laquinimod sodium was attained even though large quantities (as much as 100 volumes) of ethanol were added.

Example 6

15 Laquinimod sodium prepared according to Example 1 was added to 9.9 volumes of water at 76°C. The pH was adjusted to 10.5-11 by the addition of NaOH and the mixture was stirred until complete dissolution. The solution was cooled to 3°C but no crystallization occurred.

20

Example 7

Laquinimod sodium prepared according to Example 1 was added to 9.9 volumes of water at 76°C. The pH was adjusted to 10.5-11 by the addition of NaOH and the mixture was stirred
25 until complete dissolution. 30.6 volumes of isopropanol were added. The solution was cooled to 5°C but no crystallization occurred.

Discussion of Examples 2-7

30 Example 6 shows that merely cooling an aqueous solution of laquinimod sodium does not cause crystallization. Example 7 shows that addition of isopropanol anti-solvent (laquinimod sodium is very slightly soluble in isopropanol at room temperature) to an aqueous solution of laquinimod sodium
35 does not cause crystallization.

Examples 2-7 show that standard methods of recrystallization were not effective when used for recrystallization of laquinimod, as either no crystallization occurred or low yields were attained.

Example 8

The solution of Example 7 was then concentrated by evaporation under vacuum to 3.8 volumes. A small amount of solid crystallized out of solution. The mixture was cooled at 7°C overnight. The mixture was filtered, and the solid crystals were washed with 20 ml of isopropanol and were dried at 50°C under vacuum to a constant weight. The yield was determined to be 11.2%.

Example 9

The filtrate from Example 8 was collected, and 8.2 volumes (relative to starting laquinimod sodium in Example 7) isopropanol were added to the filtrate at room temperature. The filtrate with isopropanol was then cooled to 7°C, and a solid crystallized out of solution. The crystals were washed with 10 ml of isopropanol and were dried at 50°C under vacuum to a constant weight and the yield was determined to be 29.8% (of starting laquinimod sodium of Example 7.)

Example 10

Laquinimod sodium prepared according to Example 1 was added to 9.9 volumes of water at 76°C. The pH was adjusted to 10.5-11 by the addition of NaOH and the mixture was stirred until complete dissolution. 91 volumes of acetone were added. The solution was cooled to 5°C and a crystallization of a small amount of solid was noticed. The solid was washed with acetone and filtered and dried at 50°C under vacuum to a constant weight. The yield was determined to be 10.2%.

Example 11

Laquinimod sodium prepared according to Example 1 was added to 9.9 volumes of water at 76°C. The pH was adjusted to 10.5-11 by the addition of NaOH and the mixture was stirred
5 until complete dissolution. The solution was concentrated to 1.4 volumes using a rotation evaporator.

8.0 volumes of acetone were added to the solution, and crystallization occurred. The mixture was cooled to 7°C
10 overnight. The solid was filtered and dried at 50°C under vacuum to a constant weight. The solid was determined to be laquinimod sodium, with a yield of 90.3%.

Discussion of Examples 10 and 11

15 Example 10 shows that addition of even large amounts of acetone (laquinimod sodium is practically insoluble in acetone at room temperature) to a non-concentrated aqueous solution of laquinimod sodium provides low yields of crystalline laquinimod sodium.

20

On the other hand, Example 11 shows that if laquinimod sodium aqueous solution is first concentrated, and then anti-solvent is added, the yields of laquinimod sodium crystal are high. Large amounts of anti-solvent are not
25 required to attain high yields in this case.

Example 12

Laquinimod sodium prepared according to Example 1 was added to 11.1 volumes of water at 78°C. The pH was adjusted to 12
30 by the addition of NaOH and the mixture was stirred until complete dissolution. The solution was concentrated to 1.9 volumes using a rotary evaporator. The solution was transferred to a warmed reactor (jacket temperature 50°C.)

35 9.5 volumes of acetone were slowly added to the solution,

and crystallization occurred. The mixture was cooled to 3°C and mixed for 1.5 hours in the reactor. The solid was filtered and washed with fresh acetone, and dried at 50°C under vacuum to a constant weight and was determined to be
5 laquinimod sodium, with a yield of 79.5%.

Example 13 (recrystallization without seeding)

46.7g of laquinimod sodium prepared by a scaled-up process following the procedure of Example 1 (batch A), and 500ml of
10 deionized water were introduced into a laboratory glass reactor. The mixture was stirred and heated to 50°C until complete dissolution of the solids was observed. The solution was filtered through filter paper, and the filter was washed with 10ml of water and the wash was combined with
15 the filtrate.

The resulting solution was introduced into a laboratory reactor equipped with a vacuum distillation system. The solution was concentrated by evaporation under vacuum (35-38
20 mbar) to a volume of 112ml. After evaporation, the pressure was adjusted to atmospheric pressure and the jacket temperature was raised to 50°C, and 295g of acetone were added to the batch over 2 hours. Solid crystallization was observed during the acetone addition. The batch was cooled
25 to 2°C and stirred at this temperature for 12 hours. The solid product was isolated by filtration, washed twice with acetone and dried under vacuum at 35-40°C to constant weight. 35.7g of dry solid was obtained, yield 76.4%.

30 The starting material prepared according to Example 1 (batch A) and the dry recrystallized product were sampled and analyzed for particle size distribution, powder density and chemical purity. The results are presented in Table 1:

Table 1. Properties and purity of laquinimod sodium, Example 13

Quality parameters		Laquinimod Sodium prepared according to Example 1 (batch A)	Re-crystallized product
Particle size distribution by Malvern, micron	d(0.1)	1.5	13.3
	d(0.5)	7.1	51.1
	d(0.9)	23.2	105.1
Powder density, g/ml	BD	0.166	0.498
	TD	0.347	0.758
Heavy metals by ICP, ppm	Fe	7	< 2
	Ni	0.6	< 0.5
	Cr	0.7	0.3
Color		Grey	Off-white
Purity by HPLC, area %	Impurity 1, RT=5.49	0.06	Not detectable

HPLC = High Performance Liquid Chromatography

RT = Retention Time

5

The method of Example 13 was accompanied by high yields which are industrially reproducible.

Example 13 shows that the recrystallization process increased the purity of the laquinimod sodium, as the impurity peak 1 was no longer detectable after recrystallization, and the color was changed. In addition, the content of heavy metals Fe, Ni, and Cr was decreased.

15 In addition, the powder density of the laquinimod sodium was increased, and the size of the particles was also increased.

Example 14 (laquinimod sodium recrystallization with spontaneous crystallization - nucleation in water)

71.4g of laquinimod sodium prepared by a scaled-up process following the procedure of Example 1 (batch B) and 750ml
5 deionized water were introduced into a laboratory glass reactor. The mixture was stirred and heated to 60°C, and complete dissolution of the solids was observed. The solution was filtered through filter paper, the filter was washed with 36ml water and the wash was combined with the
10 filtrate.

The resulting solution was introduced into a laboratory reactor equipped with a vacuum distillation system. The batch was concentrated by evaporation under vacuum (37-
15 38mbar) to a volume of 153ml. After the evaporation completion, the reactor pressure was adjusted to atmospheric pressure and the jacket temperature was adjusted to 50°C. The batch was stirred for 25 minutes. At this stage spontaneous crystallization of solids was observed. Then
20 450.5g acetone was added to the batch over 2 hours. The batch was cooled to 2°C and stirred at this temperature for 12hrs, then the solid product was isolated by filtration, washed twice with acetone and dried under vacuum at 35-40°C to a constant weight. 64.2g of dry solid was obtained, yield
25 89.9%.

The starting material prepared according to Example 1 (batch B) and the dry, recrystallized product were sampled and analyzed for particle size distribution, powder density and
30 chemical purity. The results are presented in Table 2:

Table 2. Properties and purity of Laquinimod Sodium, Example 14

Quality parameters		Laquinimod Sodium prepared according to Example 1 (batch B)	Re-crystallized product
Particle size distribution by Malvern, micron	d(0.1)	2.1	3.5
	d(0.5)	10.8	15.7
	d(0.9)	35.3	43.2
Powder density, g/ml	BD	0.189	0.224
	TD	0.452	0.429
Heavy metals by ICP, ppm	Fe	4	< 2
	Ni	< 0.5	< 0.2
	Cr	1	0.2
Color		White	White
Purity by HPLC, area %	Impurity 1, RT=5.52	0.03	0.00
	Impurity 2, RT=8.48	0.05	0.01
	Impurity 3, RT=12.19	0.03	0.00

Example 15 (recrystallization method with seeded crystallization - controlled nucleation in water)

25.0g of laquinimod sodium prepared by a scaled-up process following the procedure of Example 1 (batch C) and 260ml of deionized water were introduced into a laboratory glass reactor. The mixture was stirred and heated to 60°C, and complete dissolution of the solids was observed. The solution was filtered through filter paper, the filter was washed with 15ml of water and the wash was combined with the filtrate. The resulting solution was concentrated by evaporation in a rotation evaporator under vacuum (20-

25mbar) to a residual weight of 60.0g. After the evaporation completion, the residue was introduced into a laboratory glass reactor which was pre-heated to 50°C (jacket temperature). The batch was seeded with 0.2 g of solid
5 laquinimod sodium and stirred for one hour, and crystallization of solids was observed. Then 157.7g of acetone were added to the batch over 2 hours. The batch was cooled to 2°C and stirred for 12 hours. The solid product was isolated by filtration, washed twice with acetone, and
10 dried under vacuum at 35-40°C to a constant weight. 22.6g of dry solid was obtained, yield 90.4%.

The starting material prepared according to Example 1 (batch C) and the dry, recrystallized product were sampled and
15 analyzed for particle size distribution, powder density and chemical purity. The results are presented in Table 3:

Table 3. Properties and purity of Laquinimod Sodium

Quality parameters		Laquinimod Sodium prepared according to Example 1 (batch C)	Re-crystallized product
Particle size distribution by Malvern, micron	d(0.1)	1.3	6.1
	d(0.5)	5.9	21.2
	d(0.9)	19.4	51.8
Powder density, g/ml	BD	0.158	No data
	TD	0.362	No data
Heavy metals by ICP, ppm	Fe	25	8
	Ni	2.9	1.1
	Cr	3.5	1.5
Color		Grey	Off-white
Purity by HPLC, area %	Impurity 1, RT=5.49	0.02	Not detectable
	Impurity 2, RT=8.38	0.03	Not detectable

Discussion of Examples 14 and 15

The methods of Examples 14 and 15 were accompanied by high yields which are industrially reproducible.

Examples 14 and 15 show that the recrystallization process increased the purity of the laquinimod sodium, as the impurity peaks were no longer detectable after recrystallization. In addition, the content of heavy metals Fe, Ni, and Cr was decreased. The crystals which resulted after recrystallization in examples 14 and 15 were larger than the crystals before the recrystallization.

Example 16 (crystallization without seeding - nucleation in the presence of acetone)

Water (532 mL) and laquinimod sodium (52.3 g) were introduced into a laboratory glass reactor (0.5 L). The suspension was heated to 70-73°C until a clear solution was obtained. The hot solution was cooled to 50°C and then filtered through a 0.2 micron filter. The filter was washed with 10ml of water and the wash was combined with the filtrate. The resulting solution was concentrated to a volume of 112 mL in a 1 liter reactor by evaporation while stirring under a vacuum of 30-50 mbar while maintaining the jacket temperature at 60°C, and the temperature of the reactor at about 35-40°C. Immediately after the evaporation completion and adjustment of pressure, acetone (417mL) was added to the evaporation residue over 2 hours while the jacket temperature was maintained at 50°C. The crystallization mixture was cooled to a temperature of 2°C over 2 hours and was kept at this temperature for 5-10 hours. The solid formed was collected by filtration and washed twice with 50 mL of acetone. The wet material was dried in a dryer at 30-40°C under vacuum to give 47.6 gram (90.6% yield) of dried material. The results are presented in Table 4:

Table 4. Properties of Laquinimod Sodium, Example 16

Quality parameters		Laquinimod Sodium according to Example 1 (batch B)	Re-crystallized product
Particle size distribution by Malvern, micron	d(0.1)	2.1	15.7
	d(0.5)	10.8	65.5
	d(0.9)	35.3	156.4
Color		White	White

The crystals produced by the recrystallization process were larger than the crystals of the starting material.

5

Example 17 (recrystallization of crude laquinimod sodium with insoluble impurities)

10 A 55 mg sample of laquinimod sodium prepared by a scaled-up process following the procedure of Example 1 (batch D) was mixed in 0.5 mL of deionized water at ambient temperature. The sample did not completely dissolve in water.

15 Purification by recrystallization of a sample of the batch was performed as follows:

20 Water (391 mL) and laquinimod sodium of Example 1 (batch D) (39.1 g) were introduced into a laboratory glass reactor (0.5 L). The suspension was heated by raising the jacket temperature to 73°C. After 20 min the solution was not clear. The suspension was warmed further by raising the jacket temperature to 75°C and a clear solution was still not obtained. The hot solution was cooled to 50°C and filtered through laboratory filter paper over a Buchner
25 funnel. 0.3 grams of solid residue remained on the filter paper. A sample of the solid residue was tested for

impurity content. The filter papers were washed with 47 ml of water and the wash was combined with the filtrate. The resulting solution was cooled by lowering the jacket temperature to 25°C and the solution was then concentrated under vacuum ($P < 45$ mmHg) while heating over the course of 30 min by raising the jacket temperature to 65°C. After completion of evaporation, the residue (82.1 ml, 93.2 g, $d=1.135$ g/ml) was cooled by lowering the jacket temperature to 50°C and agitated for 10 min. The batch was then seeded with solid laquinimod sodium and was stirred while maintaining the jacket temperature at 50°C for 1 hour. Acetone (316.7mL, 250.2 g) was then added to the crystallizing mixture over 2 hours at 50°C. The resulting suspension was cooled to 2°C over 4 hours and kept at this temperature for another 11 hours. The solid formed was collected by filtration and was washed twice with 31.3 g of acetone. The wet material was dried in a dryer at 30-40°C under vacuum to yield 31.7 gram (81.1%) of dried crystalline laquinimod sodium. The impurity content of the crude laquinimod sodium was tested by ICP before and after recrystallization.

Table 5. Impurity content in PPM of Laquinimod Sodium

Impurity	Laquinimod Sodium prepared according to Example 1 (batch D)	Re-crystallized product	Solid Residue
Al	14.0	5.6	411
Ca	165	65	860
Cr	2.6	<0.5	99
Cu	2.8	1.3	64
Fe	31.5	5.8	1544
Ni	5.5	<0.5	69
S	466	<1	193
Zn	20.5	7.5	352

Discussion of Example 17

Even though crude laquinimod sodium had high insoluble impurity levels before recrystallization, the recrystallization process had lowered the impurity levels. The high impurity content in the solid residue shows the importance of filtration of aqueous laquinimod solution in order to lower levels of impurities. Thus it is desirable to lower the amount of insoluble matter to below an amount that has been predetermined to cause deleterious effects on, e.g., stability of the laquinimod formulation.

What is claimed is:

1. A mixture of crystalline laquinimod sodium particles wherein 10% or more of the total amount by volume of the laquinimod sodium particles have a size of greater than 40 microns, having a bulk density of at least 0.2 g/mL or a tapped density of at least 0.4 g/mL.
2. The mixture of claim 1, having a tapped density of at least 0.4 g/mL and a bulk density of at least 0.2 g/mL.
3. The mixture of claim 1, having a tapped density of at least 0.5 g/mL.
4. The mixture of claim 1, having a tapped density of less than 0.7 g/mL.
5. The mixture of claim 1, having a tapped density of at least 0.5 g/mL and less than 0.7 g/mL.
6. The mixture of claim 5, having a tapped density of at least 0.6 g/mL and less than 0.7 g/mL.
7. The mixture of claim 1, having a bulk density of at least 0.3 g/mL.
8. The mixture of claim 7, having a tapped density of at least 0.4 g/mL.
9. The mixture of claim 8, having a tapped density of at least 0.5 g/mL.
10. The mixture of claim 7, having a tapped density of less than 0.7 g/mL.

11. The mixture of claim 7, having a tapped density of at least 0.5 g/mL and less than 0.7 g/mL.

12. The mixture of claim 11, having a tapped density of at least 0.6 g/mL and less than 0.7 g/mL.

13. The mixture of claim 1, comprising no more than 2 ppm of a heavy metal, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm, the nickel is no more than 0.2 ppm and the chromium content is no more than 0.3 ppm.

14. The mixture of claim 5, comprising no more than 2 ppm of a heavy metal, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm, the nickel content is no more than 0.2 ppm and the chromium content is no more than 0.3 ppm.

15. The mixture of claim 6, comprising no more than 2 ppm of a heavy metal, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm, the nickel content is no more than 0.2 ppm and the chromium content is no more than 0.3 ppm.

16. The mixture of claim 7, comprising no more than 2 ppm of a heavy metal, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm, the nickel content is no more than 0.2 ppm and the chromium content no more than 0.3 ppm.

17. The mixture of claim 11, comprising no more than 2 ppm of a heavy metal, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm,

the nickel content is no more than 0.2 ppm and the chromium content is no more than 0.3 ppm.

18. The mixture of claim 12, comprising no more than 2 ppm of a heavy metal, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm, the nickel content is no more than 0.2 ppm and the chromium content is no more than 0.3 ppm.

19. A pharmaceutical composition comprising the mixture of claim 1 and a pharmaceutically acceptable carrier.

20. The pharmaceutical composition of claim 19, wherein the mixture has a tapped density of at least 0.5 g/mL and less than 0.7 g/mL.

21. The pharmaceutical composition of claim 20, wherein the mixture has a tapped density of at least 0.6 g/mL and less than 0.7 g/mL.

22. The pharmaceutical composition of claim 19, wherein the mixture has a bulk density of at least 0.3 g/mL.

23. The pharmaceutical composition of claim 22, wherein the mixture has a tapped density of at least 0.5 g/mL and less than 0.7 g/mL.

24. The pharmaceutical composition of claim 23, wherein the mixture has a tapped density of at least 0.6 g/mL and less than 0.7 g/mL.

25. The pharmaceutical composition of claim 19 in the form of a tablet, capsule, pill, powder or granule.

26. The pharmaceutical composition of claim 25, wherein the

mixture has a tapped density of at least 0.5 g/mL and less than 0.7 g/mL.

27. The pharmaceutical composition of claim 26, wherein the mixture has a tapped density of at least 0.6 g/mL and less than 0.7 g/mL.

28. The pharmaceutical composition of claim 25, wherein the mixture has a bulk density of at least 0.3 g/mL.

29. The pharmaceutical composition of claim 28, wherein the mixture has a tapped density of at least 0.5 g/mL and less than 0.7 g/mL.

30. The pharmaceutical composition of claim 29, wherein the mixture has a tapped density of at least 0.6 g/mL and less than 0.7 g/mL.

31. A composition comprising a mixture of crystalline laquinimod sodium particles according to claim 1 and no more than 2 ppm of a heavy metal calculated based on total amount of laquinimod sodium in the composition, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm the nickel content is no more than 0.2 ppm and the chromium content is no more than 0.3 ppm.

32. The composition of claim 31 in crystalline, powder or granule form.

33. The composition of claim 31 further comprising a pharmaceutically acceptable carrier.

34. The composition of claim 31 in the form of a tablet, or capsule or pill.

35. The mixture of claim 2, having a tapped density of at least 0.5 g/mL and less than 0.7 g/mL.

36. The mixture of claim 35, comprising no more than 2 ppm of a heavy metal, wherein the heavy metal is iron, nickel, chromium or a mixture thereof, and the iron content is no more than 2 ppm, the nickel content is no more than 0.2 ppm and the chromium content is no more than 0.3 ppm.

37. The mixture of claim 35, wherein 50% or more of the total amount by volume of the laquinimod sodium particles have a size of greater than 15 microns.

38. The mixture of claim 1, wherein an amount of a polar impurity having an approximate retention time of 5.52 minutes when determined by isocratic reversed phase HPLC, using an ODS-3V column and a mobile phase comprised of a mixture of ammonium acetate buffer at pH 7.0 (80%) and acetonitrile (20%) is less than 0.01 of the total amount of laquinimod sodium as measured by ultraviolet absorption at 240 nm, or the impurity is not detectable.

39. The mixture of claim 1, wherein an amount of a polar impurity having an approximate retention time of 8.48 minutes when determined by isocratic reversed phase HPLC, using an ODS-3V column and a mobile phase comprised of a mixture of ammonium acetate buffer at pH 7.0 (80%) and acetonitrile (20%) is less than 0.01 of the total amount of laquinimod sodium as measured by ultraviolet absorption at 240 nm, or the impurity is not detectable.

40. The mixture of claim 1, wherein an amount of a polar impurity having an approximate retention time of 12.19 minutes when determined by isocratic reversed phase HPLC, using an ODS-

3V column and a mobile phase comprised of a mixture of ammonium acetate buffer at pH 7.0 (80%) and acetonitrile (20%) is less than 0.01 of the total amount of laquinimod sodium as measured by ultraviolet absorption at 240 nm, or the impurity is not detectable.

41. The pharmaceutical composition of claim 19, wherein the pharmaceutically acceptable carrier is mannitol.

42. The pharmaceutical composition of claim 33, wherein the pharmaceutically acceptable carrier is mannitol.

43. The pharmaceutical composition of claim 19, further comprising an alkaline-reacting component.

44. The pharmaceutical composition of claim 33, further comprising an alkaline-reacting component.