

POWDER SOLID COMPOSITIONS, PROCESS FOR THEIR PREPARATION, FORMULATIONS AND USE THEREOF

Technical field of the invention

The present invention relates to powder solid compositions comprising an extract of Bergamot and phospholipids.

5 The invention also relates to a process for the preparation of said compositions in form of powder solid. Furthermore, the invention concerns pharmaceutical, nutraceutical and cosmetic formulations comprising said compositions as well as the use of said compositions and formulations.

Background of the invention

10 Flavonoids are an important family of polyphenolic compounds naturally occurring in fruit and vegetables.

The positive effect of flavonoids in the prevention and amelioration of metabolic syndrome and in the treatment of associated pathologies, like cardiovascular diseases, hyperlipidemia and type 2 diabetes, has been confirmed by several studies (Galleano M. et al. "Flavonoids and metabolic
15 syndrome", Ann. NY Acad. Sci. (2012); 1259: 87-94).

Citrus fruits are an important source of flavonoids. The most important flavonoids of citrus fruits include diosmetin, diosmin, hesperidin, naringin, neohesperidin, nobiletin, quercetin, rutin and the flavone tangeritin. The potential beneficial effects of citrus fruits flavonoids in the treatment of
20 dysmetabolic syndromes, in the normalization of dyslipidemias and in the management of type 2 diabetes is well documented (Ashraful Alam M. et al. "Effect of Citrus Flavonoids, Naringin and Naringenin on Metabolic Syndrome and Their Mechanisms of Action", Adv. Nutr. (2014), vol. 5: 404-417; Assini et al. "Citrus flavonoids in lipid metabolism", Curr. Opin. Lipidol. (2013), vol.
25 24(1): 34-40; Un Ju Jung et al., "Effect of citrus flavonoids on lipid metabolism

and glucose-regulating enzyme mRNA levels in type-2 diabetic mice", Int. J. Biochem. Cell. Biol. (2006), vol. 38(7), 1134-145).

Among the citrus fruits, bergamot represents an important source of specific flavanone-7-O-glycosides, such as naringin, neohesperidin, brutelidin
5 and melitidin, that cannot be found in any other citrus fruits, which have demonstrated potential health benefits in clinical testing (Mollace V., et al. *"Hypolipemic and hypoglycaemic activity of bergamot polyphenols: From animal models to human studies"*, Fitoterapia (2011), 3: 309-316).

Recently an hydroalcoholic extract of Bergamot fruit has been developed
10 by Lombardo et al. (US 8,741,362 B2) with a specific polyphenol composition which proved to be effective in reducing cholesterol, blood glucose and treating metabolic syndrome.

As for all the other flavonoids, flavanones-7-O-glycosides of bergamot fruit extracts are characterized by poor oral bioavailability; consequently, high
15 oral doses of bergamot fruit extracts are required to obtain significant health benefits.

Therefore, there is still the need to find alternative Bergamot fruit extract derivatives having improved oral bioavailability.

Summary of the invention

20 The present invention relates to a process for the preparation of powder solid compositions comprising the following steps:

- a) a hydroalcoholic extract of Bergamot fruit is dispersed in at least one organic solvent and kept under mixing until a solution or a dispersion is obtained; heating is optionally applied;
- 25 b) at least one phospholipid is then added to the solution, or dispersion, of the extract and the mixture is kept under mixing; heating is optionally applied;
- c) the organic solvent is then removed to obtain the powder

composition.

The invention also relates to the powder solid compositions comprising a hydroalcoholic extract of Bergamot fruit (*Citrus aurantium* var. *bergamia*) and at least one phospholipid obtainable by the process of the invention and it
5 also relates to pharmaceutical, nutraceutical and cosmetic formulations comprising said compositions.

The powder compositions according to the invention and the formulations containing said compositions are useful in the prevention and/or treatment of metabolic syndromes, dyslipidemias and type 2 diabetes.

10 **Detailed description of the invention**

The present invention relates to powder solid compositions comprising a hydroalcoholic extract of Bergamot fruit (*Citrus aurantium* var. *bergamia*) and at least one phospholipid obtainable by the process of the invention.

The hydroalcoholic extract of Bergamot fruit is characterized by the
15 presence of three main flavanone-7-O-glycosides: neoeriocitrin, naringin and neohesperidin, that are the main responsible of the biological activity of the extract.

It has been surprisingly found that the compositions according to the present invention are characterized by an improved oral bioavailability of the
20 active ingredients (flavonoids) of the Bergamot fruit extract and, in particular, of the flavanone-7-O-glycosides, in particular naringin and neohesperidin, which were analyzed in the plasma samples of the pharmacokinetic study on rats.

The phospholipid may be selected from the group comprising lecithins
25 from soy, sunflower or egg, phosphatidyl choline, phosphatidyl serine, phosphatidyl ethanolamine, wherein the acyl groups being the same or different are mostly derived from palmitic, stearic, oleic, linoleic, linolenic acids; or combinations thereof. Preferably, the phospholipid is lecithin.

The at least one phospholipid to Bergamot fruit extract flavanone-7-O-glycosides (neohesperidin, naringin and neohesperidin) weight ratio is in the range from 6 to 30, preferably from 6 to 20, more preferably from 6 to 12.

Thanks to this ratio an improved oral bioavailability of the active ingredients (flavonoids) of Bergamot fruit extract is achieved.

The powder solid compositions may also comprise an additional surfactant, other than lecithin, with a HLB value equal or higher than 12. This additional surfactant enhances the solubilisation of Bergamot fruit extract in the organic solvent during the manufacturing process and maximizes the interaction of the extract with phospholipids. Furthermore, the additional surfactant enhances the wettability and fast dispersion of the powder solid dispersion in the gut fluids and promotes a faster and higher absorption, hence a higher bioavailability of the active ingredient of Bergamot fruit extract.

The additional surfactant may be selected from the group comprising sucrose esters, polysorbates, polyoxyethylene castor oil derivatives, polyoxyethylene stearates, D- α -tocopheryl polyethylene glycol succinate, or combinations thereof. Sucrose esters and D- α -tocopheryl polyethylene glycol succinate are the preferred ones.

Additional ingredients may be also added to the powder solid dispersion with the purpose to improve its physical and technological characteristics, allowing an easier incorporation in conventional dosage forms, such as tablets and capsules.

These additional ingredients may include soluble and insoluble fillers, such as cellulose powder, microcrystalline cellulose, calcium carbonate, calcium phosphate, mannitol, maltodextrins, sorbitol, xylitol, fructose, isomalt, inulin; binders, such as methylcellulose, hydroxypropylmethyl cellulose, hydroxypropyl cellulose, natural gums; glidants and lubricants, such as silicon dioxide, talc, stearic acid, magnesium stearate.

The present invention also relates to a manufacturing process for the preparation of the powder solid compositions comprising hydroalcoholic Bergamot fruit extract and at least one phospholipid.

The manufacturing process for preparing a powder solid composition according to the present invention comprises the following steps:

a) a hydroalcoholic extract of Bergamot fruit is dispersed in at least one suitable organic solvent and kept under mixing until a solution or a dispersion is obtained; heating is optionally applied, if required;

b) at least one phospholipid is then added to the solution/dispersion of the Bergamot fruit extract and the mixture is kept under mixing; heating is optionally applied;

c) the organic solvent is then removed to obtain the composition in the form of powder solid.

The suitable organic solvent is a polar organic solvent that leads to a total or at least substantial solubilisation of the extract, such as polar protic solvent or a polar aprotic solvent.

Preferably the polar protic solvent is a, straight or branched, C₁-C₈ alkyl alcohol and the polar aprotic solvent is a, straight or branched, C₁-C₈ alkyl ester or a C₁-C₈ dialkylketone.

A total or event partial solubilisation of the phospholipids in the selected organic solvent is also desirable.

The organic solvent may be selected from the group comprising ethyl alcohol, ethyl acetate, acetone, isobutyl alcohol, isopropyl alcohol, and combinations thereof. Ethyl alcohol and ethyl acetate are preferred.

Heating is optionally applied to facilitate solubilisation without causing any degradation of the active ingredients.

After step b) the additional surfactant and/or additional ingredients may be added to the obtained solution (or dispersion), which is kept under mixing

for a suitable period of time to facilitate the interaction of the different ingredients.

The solvent may be removed under vacuum. Alternative drying method may also be used to remove the organic solvent, such as spray drying and freeze
5 drying.

The obtained powder composition usually is then calibrated and eventually grinded to obtain the desired particle size distribution.

The present invention also relates to pharmaceutical, nutraceutical and cosmetic formulations comprising a composition of the invention and at least
10 one physiologically acceptable excipient and/or carrier.

Preferably the formulations are for oral administration or for topical administration.

Physiologically acceptable excipients and/or carriers may be, for example, disintegrant, lubricant, binder, coating agent, colorant, absorption
15 enhancer, solubilizing agent, stabilizer, flavor, sweetener, antiseptic, preservative, antioxidant and the like.

Examples of dosage forms of the formulations of the invention may be, for example tablets, chewable tablets, hard gelatin capsules, powder for reconstitution, extemporaneous or ready-to-use suspensions.

20 The powder solid compositions of the invention and formulations thereof may be used, alone or in combination with other botanical extracts, for the prevention and/or treatment of dysmetabolic syndromes, dyslipidemias and type 2 diabetes.

The improved bioavailability of active ingredients (flavonoids) of the
25 hydroalcoholic extract of Bergamot fruit allows a significant reduction of the daily dosage and an improvement of the pharmacological performance of Bergamot fruit extract.

The powder solid composition was tested in an *in vivo* pharmacokinetic

study on rats compared to the hydroalcoholic Bergamot fruit extract. The results of the study, reported in the examples, clearly show the increase of the bioavailability of the active ingredients of Bergamot fruit extract when administered as compositions of the invention.

- 5 A preliminary clinical study on volunteers was also performed to evaluate the cholesterol lowering effect of the composition of the invention compared to the hydroalcoholic Bergamot fruit extract.

The following examples further describe the invention.

Examples

- 10 **Example 1** - Preparation of a lecithin based powder solid composition of a hydroalcoholic extract of Bergamot fruit.

350 g of hydroalcoholic extract of Bergamot fruit, 420 g of sunflower lecithin and 70 g of maltodextrin are suspended in 8500 mL of ethyl alcohol in a reactor. The suspension is heated at 70°C for 2 hours under mixing.

- 15 Solvent is then removed under vacuum (about 200 mBar) at 70°C until a soft mass is obtained.

Drying is completed in an oven under vacuum at 50°C for about 12 hours until a dry mass is obtained.

- 20 The dry mass is calibrated through a 2 mm screen and mixed with 2% of silicon dioxide to obtain a freely flowable powder.

Example 2 - Preparation of a lecithin based powder solid composition of a hydroalcoholic extract of Bergamot fruit containing an additional surfactant.

- 25 500 g of hydroalcoholic extract of Bergamot fruit, 700 g of sunflower lecithin and 50 g of sucrose monopalmitate are solubilized in about 10 Lt of ethyl acetate in a reactor.

100 g of microcrystalline cellulose are added to the obtained solution and the suspension is heated at 70°C for 2 hours under mixing.

Solvent is then removed under vacuum (about 200 mBar) at 70°C until a

soft mass is obtained.

Drying is completed in an oven under vacuum at 50°C for about 12 hours until a dry mass is obtained.

The dry mass is calibrated through a 2 mm screen and mixed with 2% of silicon dioxide and 1% of talc to obtain a freely flowable powder.

Example 3 - Pharmacokinetic study on rats

Plasma concentration of naringin and neohesperidin were determined in rats after the oral administration of a single dose of the composition obtained in Example 1 and compared to the plasma concentration obtainable administering the hydroalcoholic extract of Bergamot fruit. Male Sprague-Dawley rats, weighting about 300 g were used for the pharmacokinetic experiments. A single dose of 500 mg/Kg of hydroalcoholic extract of Bergamot fruit and a single dose of 500 mg/Kg of the composition of obtained in example 1, corresponding to about 200 mg/Kg of hydroalcoholic extract of Bergamot fruit, were administered as water suspension by intragastric gavage. Blood sample were collected after 1 and 2 hours from the administration. Plasma was obtained from blood samples by centrifugation at 2.000 rpm for 10 minutes at 4°C in presence of EDTA. After a suitable extraction procedure, samples were analyzed by HPLC equipped with fluorescent detector for naringin and neohesperidin concentration.

The analytical results are summarized in the following table 1:

Table 1

	Naringin (ppm)		Neohesperidin (ppm)	
	Hydroalcoholic extract of Bergamot fruit	Composition of Example 1	Hydroalcoholic extract of Bergamot fruit	Composition of Example 1
Rat 1 (1 hour)	0.2029	1.4409	0.2728	1.5621
Rat 2 (1 hour)	0.2286	1.545	0.2751	1.1917
Rat 3 (1 hour)	0.2186	1.7878	0.3201	1.543
Rat 1 (2 hour)	0.1089	1.9046	0.1648	3.5433
Rat 2 (2 hour)	0.1029	2.334	0.1447	3.0571
Rat 3 (2 hour)	0.1067	2.0516	0.1417	3.2322

Example 4 - Formulations containing the composition obtained in example 1 (tablets)

5	Composition obtained in Example 1	500.0 mg
	Microcrystalline cellulose	150.0 mg
	Dicalcium phosphate anhydrous	104.0 mg
	Sodium croscarmellose	30.0 mg
	Silicon dioxide	8.0 mg
10	Magnesium stearate	8.0 mg

1.0 Kg of the composition obtained in Example 1, 0.3 Kg of microcrystalline cellulose (direct compression grade), 0.208 Kg of dicalcium phosphate anhydrous (direct compression grade) and 0.06 Kg of sodium croscarmellose were blended in a V-mixer for 10 minutes. 16 g of silicon dioxide and 16 g of magnesium stearate were added to the powder mixture and blended for 2 additional minutes. The obtained mixture was compressed in a single punch tableting machine, equipped with a round concave punch with a diameter of 11 mm.

Example 5 - Formulations containing the composition obtained in example 2 (size 0 hard gelatin capsules)

	Composition obtained in example 2	350.0 mg
	Silicon dioxide	5.0 mg
5	Magnesium stearate	5.0 mg

2.0 Kg of the composition obtained in example 2 were blended with 28.5 g of silicon dioxide and 28.5 g of magnesium stearate in a V-mixer for 2 minutes. The mixture was filled in size 0 hard gelatin capsules.

Example 6 - Clinical study

10 A pilot clinical study on volunteers was also performed to evaluate the cholesterol lowering effect of the composition of the invention.

Ten (10) patients were treated with 2 hard gelatin capsules of 500 mg, one in the morning and one in the evening (for a total of 1000 mg daily) for 30 consecutive days. The primary objective of the study was the evaluation of the
15 clinical activity on the reduction of cardiovascular risk in patients with dyslipidemia associated or not with hyperglycemia, by measuring the modulation of total cholesterol (tChol), low-density lipoproteins (LDL), triglycerides (TG), high-density lipoproteins (HDL) and blood glucose before and after 30-day treatment. Safety was also assessed.

20 Inclusion criteria were: signed informed consent; age between 30 and 90 years old; a diagnosis of type II diabetes mellitus; fasting blood glucose ≥ 110 mg/dl; isolated hypercholesterolemia (cholesterol bound to low-density lipoprotein cLDL ≥ 130 mg/dl), with or without hypertriglyceridemia (> 175 mg/dl); medium cardiovascular risk measured by the “European
25 Guidelines on Cardiovascular disease prevention on clinical practice” (2012). All concomitant treatments must have begun at least 3 months before the beginning of the study and maintained in a constant regimen for the entire duration of the study.

The exclusion criteria were: patients with overt liver disease, serious gastrointestinal disorders, chronic renal failure, hypercalcemia, myopathy, uncontrolled diabetes, myocardial ischemic, presence of heart failure NYHA class III and IV, alcohol abuse, history of psychiatric disorders and major depression, HIV or serious infections or neoplasia, statin therapy stabilized by 4 months.

The patients were visited every 7 days during the study and the compliance was monitored. Serum aspartate aminotransferase, alanine aminotransferase, creatine were measured in order to monitor possible side effects,

Treatment with 1000 mg daily for 30 consecutive days resulted in a strong reduction for total cholesterol, LDL, TG, fasting blood glucose and a significant increase in HDL in majority of subjects. The results are reported in Table 2.

The initial mean values of 277 mg/dl tChol, 184 mg/dl LDL and 255 mg/dl TG, decreased to 195, 113 and 164 mg/dl, respectively after treatment.

In addition, a reduction in fasting blood glucose was also observed.

Table 2

mg/dl	tChol		Trigl		LDL		HDL		Gly	
	before	after	before	after	before	after	before	after	before	after
Mean	277	195	255	164	184	113	41	50	111	101
SD	15	11	24	11	15	10	2	15	8	3

CLAIMS

1. A manufacturing process for preparing a powder solid composition comprising the following steps:
 - 5 a) a hydroalcoholic extract of Bergamot fruit is dispersed in at least one polar organic solvent and kept under mixing until a solution or a dispersion is obtained; heating is optionally applied; wherein the extract contains the flavonoids neoeriocitrin, naringin and neohesperidin;
 - b) at least one phospholipid is then added to the solution/dispersion of the
10 Bergamot fruit extract and the mixture is kept under mixing; wherein the weight ratio between the at least one phospholipid and the flavonoids of the extract ranges from 6 to 30; heating is optionally applied;
 - c) the organic solvent is then removed to obtain the powder solid composition.
- 15 2. A manufacturing process according to claim 1, wherein the weight ratio ranges from 6 to 20, preferably from 6 to 12.
3. A manufacturing process according to claim 1 or 2, wherein the at least one
20 phospholipid is selected from the group consisting of lecithins from soy, sunflower or egg, phosphatidyl choline, phosphatidyl serine, phosphatidyl ethanolamine, wherein the acyl groups being the same or different are mostly derived from palmitic, stearic, oleic, linoleic, linolenic acids; or combinations thereof.
- 25 4. A manufacturing process according to claim 3, wherein the at least one phospholipid is lecithin.
5. A manufacturing process according to claims 1-4, further comprising an additional surfactant, other than lecithin, with a HLB value equal or higher than
30 12.
6. A manufacturing process according to claim 5, wherein the additional

surfactant is selected from the group consisting of sucrose esters, polysorbates, polyoxyethylene castor oil derivatives, polyoxyethylene stearates, D- α -tocopheryl polyethylene glycol succinate, or combinations thereof.

- 5 7. A manufacturing process according to claim 6, wherein the additional surfactant is selected from sucrose esters and D- α -tocopheryl polyethylene glycol succinate.
8. A manufacturing process according to claim 1, wherein the polar organic
10 solvent is a polar protic solvent or a polar aprotic solvent.
9. A manufacturing process according to claim 8, wherein the polar protic solvent is a, straight or branched, C₁-C₈ alkyl alcohol.
- 15 10. A manufacturing process according to claim 8, wherein the polar aprotic solvent is selected from a, straight or branched, C₁-C₈ alkyl ester or C₁-C₈ dialkylketone.
11. A manufacturing process according to claims 1-10, wherein the polar
20 organic solvent is selected from the group consisting of ethyl alcohol, ethyl acetate, acetone, isobutyl alcohol, isopropyl alcohol, and combinations thereof.
12. A manufacturing process according to claim 11, wherein the organic solvent
is selected from ethyl alcohol and ethyl acetate.
- 25 13. A powder solid composition obtainable by the manufacturing process according to any one of claims 1-12, wherein the powder solid composition comprises at least one phospholipid and a hydroalcoholic extract of Bergamot fruit containing flavonoids, neoeriocitrin, naringin and neohesperidin.
- 30 14. Pharmaceutical, nutraceutical and cosmetic formulations comprising a powder solid composition according to claim 13 and at least one physiologically

acceptable excipient and/or carrier.

15. A powder solid composition according to claim 13 or a formulation according to claim 14 for use in the prevention and/or treatment of metabolic syndromes, dyslipidemias and type 2 diabetes.

16. Use of a powder solid composition according to claim 13 or a formulation according to claim 14 in the manufacture of a medicament for the prevention and/or treatment of metabolic syndromes, dyslipidemias and type 2 diabetes.

17. The powder solid composition according to claim 13 wherein the at least one phospholipid is selected from the group consisting of lecithins from soy, sunflower or egg, phosphatidyl choline, phosphatidyl serine, phosphatidyl ethanolamine, wherein the acyl groups being the same or different are mostly derived from palmitic, stearic, oleic, linoleic, linolenic acids; or combinations thereof.

18. The powder solid composition according to claim 13 wherein the at least one phospholipid is lecithin.

19. The powder solid composition according to claim 18 further comprising an additional surfactant, other than lecithin, with a HLB value equal or higher than 12.

20. The powder solid composition according to claim 19, wherein the additional surfactant is selected from the group consisting of sucrose esters, polysorbates, polyoxyethylene castor oil derivatives, polyoxyethylene stearates, D- α -tocopheryl polyethylene glycol succinate, or combinations thereof.

21. The powder solid composition according to claim 20, wherein the additional surfactant is selected from sucrose esters and D- α -tocopheryl polyethylene glycol succinate.

22. Pharmaceutical, nutraceutical or cosmetic formulation according to claim 14, for oral administration or for topical administration.