~~R^{10} is selected from hydrogen, alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclyl;~~

wherein the term "cycloalkyl" refers to a hydrocarbon group selected from saturated and partially unsaturated cyclic hydrocarbon groups.

6. The compound of Claim 5, wherein p' is an integer of 1; R^1 and R^2, which may be the same or different, are each selected from hydrogen, halogen, alkyl, and aryl; Z is independently selected from halogen and alkyl.

7. The compound of Claim 5, wherein p' is an integer of 3; R^1 and R^2, which may be the same or different, are each selected from hydrogen, halogen, alkyl, and aryl; Z is independently selected from halogen and alkyl.

8. The compound of Claim 1, which is selected from a compound of Formula **(III)**:



(III)

or a pharmaceutically acceptable salt thereof,

wherein:

Y, at each occurrence, is independently selected from $-CR^1R^2-$, $-R^3C{=}CR^4-$, $-NR^5-$, -O-, and -S-;

p" is an integer ranging from 1 to 11;

Z, at each occurrence, is independently selected from hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, -CN, $-NO_2$, $-OR^6$, $-NR^6R^7$, $-NR^6COR^7$, $-NR^6-NR^7COR^8$, $-NR^6SO_2R^7$, $-CONR^6R^7$, $-COOR^6$, and $-SO_2R^6$, wherein each of the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and heterocyclyl is optionally substituted with at least one substituent R^9;

n is an integer ranging from 0 to 3;

R^1 and R^2, which may be the same or different, are each selected from hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, $-NR^6R^7$, $-OR^6$, $-COR^6$, $-CO_2R^6$, $-CONR^6R^7$, $-NR^6CONR^7R^8$, $-NR^6CO_2R^7$, $-NR^6SO_2R^7$, and $-SO_2R^6$, wherein each of the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and heterocyclyl is optionally substituted with at least one substituent R^9; and when $(Y)_{p"}$ comprises $-CR^1R^2-CR^1R^2-$, optionally R^1 or R^2 substituted on each of the two carbons, together with the two carbons to which they are attached,