

TITLE

Recombinant viral-based malaria vaccines

FIELD OF THE INVENTION

5 The invention relates to the field of medicine. More
in particular, the invention relates to the use of a
recombinantly produced viral vector as a carrier of an
antigenic determinant selected from a group of malaria
pathogens for the development of a vaccine against
10 malaria infections.

BACKGROUND OF THE INVENTION

Malaria currently represents one of the most
prevalent infections in tropical and subtropical areas
15 throughout the world. Per year, malaria infections lead
to severe illnesses in hundreds of million individuals
worldwide, while it kills 1 to 3 million people in
developing and emerging countries every year. The
widespread occurrence and elevated incidence of malaria
20 are a consequence of the increasing numbers of drug-
resistant parasites and insecticide-resistant parasite
vectors. Other factors include environmental and climatic
changes, civil disturbances and increased mobility of
populations. Malaria is caused by the mosquito-borne
25 hematoprotezoan parasites belonging to the genus
Plasmodium. Four species of *Plasmodium* protozoa
(*P.falciparum*, *P.vivax*, *P.ovale* and *P.malariae*) are
responsible for the disease in man; many others cause
disease in animals, such as *P.yoelii* and *P.berghei* in
30 mice. *P.falciparum* accounts for the majority of
infections and is the most lethal type ("tropical
malaria"). Malaria parasites have a life cycle consisting
of several stages. Each one of these stages is able to
induce specific immune responses directed against the

corresponding occurring stage-specific antigens. Malaria parasites are transmitted to man by several species of female Anopheles mosquitoes. Infected mosquitoes inject the 'sporozoite' form of the malaria parasite into the mammalian bloodstream. Sporozoites remain for few minutes in the circulation before invading hepatocytes. At this stage the parasite is located in the extra-cellular environment and is exposed to antibody attack, mainly directed to the 'circumsporozoite' (CS) protein, a major component of the sporozoite surface. Once in the liver, the parasites replicate and develop into so-called 'schizonts'. These schizonts occur in a ratio of up to 20,000 per infected cell. During this intra-cellular stage of the parasite, main players of the host immune response are T-lymphocytes, especially CD8+ T-lymphocytes (Romero et al. 1998). After about one week of liver infection, thousands of so-called 'merozoites' are released into the bloodstream and enter red blood cells, becoming targets of antibody-mediated immune response and T-cell secreted cytokines. After invading erythrocytes, the merozoites undergo several stages of replication and transform into so-called 'trophozoites' and into schizonts and merozoites, which can infect new red blood cells. This stage is associated with overt clinical disease. A limited amount of trophozoites may evolve into 'gametocytes', which is the parasite's sexual stage. When susceptible mosquitoes ingest erythrocytes, gametocytes are released from the erythrocytes, resulting in several male gametocytes and one female gametocyte. The fertilization of these gametes leads to zygote formation and subsequent transformation into ookinetes, then into oocysts, and finally into salivary gland sporozoites.

Targeting antibodies against gametocyte stage-specific surface antigens can block this cycle within the

mosquito mid gut. Such antibodies will not protect the mammalian host, but will reduce malaria transmission by decreasing the number of infected mosquitoes and their parasite load.

5 Current approaches to malaria vaccine development can be classified according to the different stages in which the parasite can exist, as described above. Three types of possible vaccines can be distinguished:

- 10 - Pre-erythrocytic vaccines, which are directed against sporozoites and/or schizont-infected cells. These types of vaccines are mostly CS-based and should ideally confer sterile immunity, mediated by humoral and cellular immune response, preventing malaria infection.
- 15 - Asexual blood stage vaccines, which are designed to minimize clinical severity. These vaccines should reduce morbidity and mortality and are meant to prevent the parasite from entering and/or developing in the erythrocytes.
- 20 - Transmission-blocking vaccines, which are designed to hamper the parasite development in the mosquito host. This type of vaccine should favor the reduction of population-wide malaria infection rates.

25 Next to these vaccines, the feasibility of developing malaria vaccines that target multiple stages of the parasite life cycle is being pursued in so-called multi-component and/or multi-stage vaccines. Currently no
30 commercially available vaccine against malaria is available, although the development of vaccines against malaria has already been initiated more than 30 years ago: immunization of rodents, non-human primates and humans with radiation-attenuated sporozoites conferred

protection against a subsequent challenge with sporozoites (Nussenzweig et al. 1967; Clyde et al. 1973). However, the lack of a feasible large-scale culture system for the production of sporozoites prevents the widespread application of such vaccines.

To date the most promising vaccine candidates tested in humans have been based on a small number of sporozoite surface antigens. The CS protein is the only *P.falciparum* antigen demonstrated to consistently prevent malaria when used as the basis of active immunization in humans against mosquito-borne infection, albeit it at levels that is often insufficient. Theoretical analysis has indicated that the vaccine coverage as well as the vaccine efficiency should be above 85%, or otherwise mutants that are more virulent may escape (Gandon et al. 2001).

One way of inducing an immune response in a mammal is by administering an infectious carrier, which harbors the antigenic determinant in its genome. One such carrier is a recombinant adenovirus, which has been replication-defective by removal of regions within the genome that are normally essential for replication, such as the E1 region. Examples of recombinant adenoviruses that comprise genes encoding antigens are known in the art (WO 96/39178), for instance HIV-derived antigenic components have been demonstrated to yield an immune response if delivered by recombinant adenoviruses (WO 01/02607; WO 02/22080). Also for malaria, recombinant adenovirus-based vaccines have been developed. These vectors express the entire CS protein of *P.yoelii*, which is a mouse-specific parasite and these vectors have been shown to be capable of inducing sterile immunity in mice, in response to a single immunizing dose (Bruña-Romero et al. 2001a). Furthermore, a similar vaccine vector using CS from

P.berghei was recently shown to elicit long-lasting protection when used in a prime-boost regimen, in combination with a recombinant vaccinia virus (Gilbert et al. 2002) in mice. It has been demonstrated that CD8+ T cells primarily mediate the adenovirus-induced protection. It is unlikely the *P.yoelii*- and *P.berghei* based adenoviral vectors would work well in humans, since the most dramatic malaria-related illnesses in humans are not caused by these two parasites. Moreover, it is preferred to have a vaccine which is potent enough to generate long-lasting protection after one round of vaccination, instead of multiple vaccination rounds using either naked DNA injections and/or vaccinia based vaccines as boosting or priming agents.

Despite all efforts to generate a vaccine that induces an immune response against a malaria antigenic determinant and protects from illnesses caused by the malaria parasite, many vaccines do not fulfill all requirements as described above. Whereas some vaccines fail to give a protective efficiency of over 85% in vaccinated individuals, others perform poorly in areas such as production or delivery to the correct cells of the host immune system.

DESCRIPTION OF THE FIGURES

Figure 1 shows the newly synthesized clone 02-148 (pCR-script.Pf), which is based on a range of known *Plasmodium falciparum* genes, and which encodes the novel circumsporozoite protein (A) (SEQ ID NO:3), plus the codon-optimized nucleic acid sequence (B) (SEQ ID NO:1). SEQ ID NO:2 is the translated protein product translated from the Coding Sequence of SEQ ID NO:1 as generated by PatentIn 3.1. SEQ ID NO:2 is identical to SEQ ID NO:3 in content.

Figure 2 shows the amino acid sequence (A) (SEQ ID NO:6) and nucleic acid sequence (B) (SEQ ID NO:4) of synthetic clone named 02-659 (pf-aa-sub), which is the

5 circumsporozoite gene of the *P.falciparum* strain 3D7, lacking the C-terminal 14 amino acids. SEQ ID NO:5 is the translated protein product translated from the Coding Sequence of SEQ ID NO:4 as generated by PatentIn 3.1. SEQ ID NO:5 is identical to SEQ ID NO:6 in content.

10

Figure 3 shows the amino acid sequence (A) (SEQ ID NO:9) and nucleic acid sequence (B) (SEQ ID NO:7) of the codon-optimized circumsporozoite gene of *P.yoelii*. SEQ ID NO:8 is the translated protein product translated from the

15 Coding Sequence of SEQ ID NO:7 as generated by PatentIn 3.1. SEQ ID NO:8 is identical to SEQ ID NO:9 in content.

Figure 4 shows the (A) cellular immune response and the (B) humoral immune response in mice upon immunization

20 with Ad5- and Ad35-based vectors harboring the *P.yoelii* circumsporozoite gene, administered via two routes: intra muscular and subcutaneous in different doses.

Figure 5 shows the inhibition in mice of a *P.yoelii* sporozoite challenge following immunization with Ad5- and Ad35-based vectors harboring the *P.yoelii* circumsporozoite gene, administered in different doses, depicted in percentage of inhibition (A), and in the presence of a parasite specific RNA molecules in the

25 liver (B).

30

Figure 6 shows the cellular immune response raised by immunization with an Ad5-based vector harboring the full

length *P.falciparum* circumsporozoite gene, and two deletion mutants, administered in different doses.

DESCRIPTION OF THE INVENTION

5 The present invention relates to different kinds of replication-defective recombinant viral vectors comprising a heterologous nucleic acid encoding an antigenic determinant of several *Plasmodium* protozoa. Preferably it relates to viral vectors that comprise
10 nucleic acids encoding the circumsporozoite (CS) protein of *P.falciparum* and *P.yoelii*. More preferably, said viral vector is an adenovirus, preferably based on a serotype that is efficient in delivering the gene of interest, that encounters low numbers of neutralizing antibodies in
15 the host and that binds to the relevant immune cells in an efficient manner. In a preferred embodiment the CS protein is generated such that it will give rise to a potent immune response in mammals, preferably humans. In one aspect, the expression of the protein is elevated due
20 to codon-optimization and thus altering the codon-usage such that it fits the host of interest. The novel CS proteins of the present invention are depicted in figure 1A (SEQ ID NO:3), 2A (SEQ ID NO:6) and 3A (SEQ ID NO:9), while the codon-optimized genes encoding said proteins
25 are depicted in figure 1B (SEQ ID NO:1), 2B (SEQ ID NO:4) and 3B (SEQ ID NO:7) respectively.

 The invention also relates to vaccine compositions comprising a replication-defective recombinant viral vector according to the invention, and a pharmaceutically
30 acceptable carrier, further comprising preferably an adjuvant. Furthermore, the invention relates to the use of a vaccine composition according to the invention in the therapeutic, prophylactic or diagnostic treatment of malaria.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the use of recombinant viruses as carriers of certain specific antigenic determinants selected from a group of malaria antigens. It is the goal of the present invention to provide a solution to at least a part of the problems outlined above for existing vaccines against malaria.

The present invention relates to a replication-defective recombinant viral vector comprising a heterologous nucleic acid encoding an antigenic determinant of *Plasmodium falciparum*. In a preferred embodiment said viral vector is an adenovirus, an alphavirus or a vaccinia virus. In a more preferred embodiment said viral vector is an adenovirus, wherein said adenovirus is preferably derived from a serotype selected from the group consisting of: Ad5, Ad11, Ad26, Ad34, Ad35, Ad48, Ad49 and Ad50. In one particular aspect of the invention the replication-defective recombinant viral vector according to the invention, comprises an antigenic determinant that is the circumsporozoite (CS) protein, or an immunogenic part thereof. Preferably, said heterologous nucleic acid is codon-optimized for elevated expression in a mammal, preferably a human. Codon-optimization is based on the required amino acid content, the general optimal codon usage in the mammal of interest and a number of provisions of aspects that should be avoided to ensure proper expression. Such aspects may be splice donor or -acceptor sites, stop codons, Chi-sites, poly(A) stretches, GC- and AT-rich sequences, internal TATA boxes, etcetera.

In a preferred embodiment, the invention relates to a replication-defective recombinant viral vector according to the invention, wherein the adenine plus

thymine content in said heterologous nucleic acid, as compared to the cytosine plus guanine content, is less than 87%, preferably less than 80%, more preferably less than 59% and most preferably equal to approximately 45%.

5 The invention provides in one embodiment a replication-defective recombinant viral vector, wherein said circumsporozoite protein is the circumsporozoite protein as depicted in figure 1A and in another embodiment a codon-optimized heterologous nucleic acid as depicted in
10 figure 1B. The proteins can be in a purified form, but also expressed *in vivo* from nucleic acid delivery vehicles such as the recombinant viral vectors of the present invention. In a purified form, such proteins can be applied in other types of vaccines, wherein the
15 protein is for instance enclosed in liposomes or other carriers used in the art. The nucleic acid can be cloned into other vectors than as disclosed herein, but also be applied as naked DNA in other vaccine-settings.

In another embodiment, the invention relates to a
20 replication-defective recombinant viral vector according to the invention, wherein the circumsporozoite protein, or the immunogenic part thereof, is lacking a functional GPI anchor sequence.

Apart from the use of new genes and proteins that
25 can be applied for use in humans, the invention also discloses novel genes that may be used in humans as well as in other mammals. Therefore, the invention also relates to a replication-defective recombinant viral vector comprising a heterologous nucleic acid encoding
30 the circumsporozoite protein of *Plasmodium yoelii*, wherein said nucleic acid is codon-optimized for elevated expression in a mammal.

In a more preferred embodiment said viral vector is an adenovirus, an alphavirus or a vaccinia virus, and it

is even more preferred to use a recombinant adenovirus, which is preferably selected from the group consisting of: Ad5, Ad11, Ad26, Ad34, Ad35, Ad48, Ad49 and Ad50. As in *P.falciparum* it is also for *P.yoelii* preferred to use
5 a codon-optimized gene for proper expression in the host of interest. Therefore, in a preferred embodiment the adenine plus thymine content in said nucleic acid, as compared to the cytosine plus guanine content, is less than 87%, preferably less than 80%, more preferably less
10 than 59% and most preferably equal to approximately 45%.

The invention provides in one embodiment a replication-defective recombinant viral vector according to the invention, wherein said circumsporozoite protein is the circumsporozoite protein as depicted in figure 3A,
15 while in another embodiment, a replication-defective recombinant viral vector is provided, wherein said nucleic acid is the nucleic acid as depicted in figure 3B. In a preferred aspect, the circumsporozoite protein, or the immunogenic part thereof, is lacking a functional
20 GPI anchor sequence.

The invention further relates to an isolated nucleic acid encoding a circumsporozoite protein of *Plasmodium falciparum* as depicted in figure 1B, wherein said nucleic acid is codon-optimized, and to an isolated nucleic acid
25 encoding a circumsporozoite protein of *Plasmodium falciparum* strain 3D7, as depicted in figure 2B, wherein said nucleic acid is codon-optimized. Such isolated nucleic acids can be applied in subcloning procedures for the generation of other types of viral-based vaccines,
30 apart for the types as disclosed herein. Furthermore, such isolated nucleic acids can be used for naked DNA vaccines or in cloning procedures to generate vectors for *in vitro* production of the encoded protein, which, in itself can be further used for vaccination purposes and

the like. The production can be in all kinds of systems, such as bacteria, yeasts or mammalian cells known in the art.

In another embodiment of the present invention, an
5 isolated nucleic acid encoding a circumsporozoite protein of *Plasmodium yoelii* as depicted in figure 3B is provided, wherein said nucleic acid is codon-optimized. Furthermore, a vaccine composition comprising a replication-defective recombinant viral vector according
10 to the invention, and a pharmaceutically acceptable carrier is provided. Pharmaceutically acceptable carriers are well known in the art and used extensively in a wide range of therapeutic products. Preferably, carriers are applied that work well in vaccines. More preferred are
15 vaccines, further comprising an adjuvant. Adjuvants are known in the art to further increase the immune response to an applied antigenic determinant. The invention also relates to the use of a vaccine composition according to the invention in the therapeutic, prophylactic or
20 diagnostic treatment of malaria.

Another embodiment of the present invention relates to a method of treating a mammal for a malaria infection or preventing a malaria infection in a mammal, said method comprising (in either order, or simultaneously)
25 the steps of administering a vaccine composition according to the invention, and administering a vaccine composition comprising at least one purified malaria-derived protein or peptide. The invention also relates to a method of treating a mammal for a malaria infection or
30 preventing a malaria infection in a mammal, said method comprising (in either order, or simultaneously) the steps of administering a vaccine composition comprising a replication-defective recombinant viral vector comprising a malaria circumsporozoite antigen according to the

invention; and administering a vaccine composition comprising a replication-defective recombinant viral vector comprising another antigen, such as LSA-1 or LSA-3 according to the invention.

5

The advantages of the present invention are multi-fold. Next to the knowledge that recombinant viruses, such as recombinant adenoviruses can be produced to very high titers using cells that are considered safe, and that can grow in suspension to very high volumes, using medium that does not contain any animal- or human derived components, the present invention combines these features with a vector harboring the circumsporozoite gene of *Plasmodium falciparum*. *P.falciparum* is the parasite that causes tropical malaria. Moreover, the gene has been codon-optimized to give an expression level that is suitable for giving a proper immune response in humans. The present invention provides a vaccine against malaria infections, making use of for instance adenoviruses that do not encounter high titers of neutralizing antibodies. Examples of such adenoviruses are serotype 11 and 35 (Ad11 and Ad35, see WO 00/70071 and WO 02/40665).

The nucleic acid content between the malaria-causing pathogen, such as *P.falciparum* and the host of interest, such as *Homo sapiens* is very different. The invention now provides a solution to some of the disadvantages of vaccines known in the art, such as expression levels that are too low to elicit a significant immune response in the host of interest, preferably humans.

Recombinant viral vectors have been used in vaccine set-ups. This has been demonstrated for vaccinia-based vaccines and for adenovirus-based vaccines. Moreover, a platform based on alphaviruses is being developed for vaccines as well. In a preferred embodiment, the

invention relates to the use of recombinant adenoviruses that are replication defective through removal of at least part of the E1 region in the adenoviral genome, since the E1 region is required for replication-, transcription-, translation- and packaging processes of newly made adenoviruses. E1 deleted vectors are generally produced on cell lines that complement for the deleted E1 functions. Such cell lines and the use thereof for the production of recombinant viruses have been described extensively and are well known in the art. Preferably, PER.C6™ cells, as represented by the cells deposited under ECACC no. 96022940 at the European Collection of Animal Cell Cultures (ECACC) at the Centre for Applied Microbiology and Research (CAMR, UK), are being used to prevent the production of replication competent adenoviruses (rca). In another preferred embodiment, cells are being applied that support the growth of recombinant adenoviruses other than those derived of adenovirus serotype 5 (Ad5). Reference is made to publications WO 97/00326, WO 01/05945, WO 01/07571, WO 00/70071, WO 02/40665 and WO 99/55132, for methods and means to obtain rca-free adenoviral stocks for Ad5 as well as for other adenovirus serotypes.

Adenoviral-based vectors that have been used in the art mainly involved the use of Ad5 vectors. However, as has been described (WO 00/03029, WO 02/24730, WO 00/70071, WO 02/40665 and in other reports in the art), administration of Ad5 and efficient delivery to the target cells of interest, responsible for sufficient immunogenic responses, is hampered by the presence of high titres of neutralizing antibodies circulating in the bloodstream if a subject previously encountered an Ad5 infection. It has been investigated what serotypes are better suited for therapeutic use, and it turned out that

a limited number of serotypes encountered neutralizing antibodies in only a small percentage of individuals in the human population. These experiments have been described in WO 00/70071. Therefore, in a preferred embodiment, the invention relates to the use of adenovirus serotype 11, 26, 34, 35, 48 and 50, and more preferably to Ad11 and Ad35, since these serotypes encountered no neutralizing antibodies in the vast majority of tested samples.

10 Apart from avoiding the presence of neutralizing antibodies directed against certain serotypes, it might also be beneficial to target the replication-deficient recombinant viral vectors to a certain subset of cells involved in the immune response. Such cells are for instance dendritic cells. It was found that certain adenovirus serotypes, such as Ad16, Ad35 and Ad50, carry capsid proteins that specifically bind to certain receptors present on dendritic cells (WO 02/24730). Ad5 is a serotype that is mainly homing to the liver, which may be a disadvantage if sufficient numbers of viral particles should infect cells of the immune system. It was found that at least in *in vitro* experiments some of the serotypes, different from Ad5 could infect dendritic cells multi-fold better than Ad5, suggesting that also *in vivo* the delivery to such cells is more efficient. It still remains to be seen whether this *in vitro* to *in vivo* translation holds up, and if serotypes other than Ad5 will give rise to the required protection level. It is also part of the invention to provide the serotypes of choice, as far as neutralizing antibodies are concerned, with capsid proteins, such as the fiber or a part thereof from a serotype that is able to selectively recognize dendritic cells. It must be noted here that in the published documents WO 00/03029, WO 02/24730, WO 00/70071

and WO 02/40665, Ad50 was mistakenly named Ad51. The Ad51 serotype that was referred to in the mentioned publications is the same as serotype Ad50 in a publication by De Jong et al. (1999), wherein it was
5 denoted as a B-group adenovirus. For the sake of clarity, Ad50 as used herein, is the B-group Ad50 serotype as mentioned by De Jong et al. (1999).

It is now known that a first administration with a specific adenoviral serotype elicits the production of
10 neutralizing antibodies in the host against that specific vector. Thus, it is desirable to use in a subsequent setting (a follow-up boost or in the administration of another, non-related vaccine) a composition based on a different adenovirus serotype, which is not neutralized
15 by antibodies raised in the first administration. Therefore, the invention further relates to methods for vaccinating mammalian individuals in which a priming vaccine composition comprises a replication-defective recombinant adenovirus of a first serotype, while in a
20 boosting vaccine composition a replication-defective recombinant adenovirus of a second serotype are used. Prime/boost settings have been described in more detail in international patent applications PCT/NL02/00671 and PCT/EP03/50748 (not published); these applications relate
25 to the use of a recombinant adenovirus vector of a first serotype for the preparation of a medicament for the treatment or prevention of a disease in a human or animal treated with a recombinant adenovirus vector of a second serotype, wherein said first serotype is different from
30 said second serotype, and wherein said first serotype is selected from the group consisting of: Ad11, Ad26, Ad34, Ad35, Ad46 and Ad49, and wherein said second serotype is preferably adenovirus serotype 5. Thus, it relates to the use of different adenoviral serotypes that encounter low

pre-existing immunities in subjects that are to be treated. Preferred examples of such serotypes are the recombinant mentioned, wherein Ad5 is not excluded for individuals that have never experienced an Ad5 infection.

5 The settings described and claimed in the applications mentioned above relate to the use of adenoviral vectors carrying transgenes such as those from measles, or gag from HIV (for treatment of humans) or SIV (for treatment and studies in monkeys).

10 One non-limiting example of a prime-boost set-up towards Malaria is a setting in which, next to different adenovirus serotypes, also different antigenic determinants may be used. One non-limiting example of an antigen different from CS is the Liver Specific Antigen 1
15 (LSA-1, Kurtis et al. 2001). Such set-ups are at least for one reason useful, namely that the CS antigen is expressed mainly during the blood-stage of the parasite, while its expression goes down in the liver-stage. For LSA-1, this situation is more or less the opposite; it is
20 expressed to low levels during the blood-stage, but is highly expressed during the liver-stage. Although one could use both antigens in subsequent administrations, it may also be used at the same time to provide protection against the parasite at the blood-stage as well as at the
25 liver-stage. In a further embodiment of the present invention both antigens may be delivered by one adenovirus serotype (either cloned together in the same vector, or separately in separate vectors of the same serotype). In another embodiment both antigens are
30 delivered by different serotypes that may be delivered at the same time or separately in time, for instance in a prime-boost setting. The vaccines of the present invention may also be used in settings in which prime-boosts are being used in combination with naked DNA or

other delivery means, unrelated to the replication-defective viral vectors of the present invention, such as purified proteins or peptides. Examples of such proteins that may be used in prime-boosts (Ad/protein; protein/Ad; protein/Ad/Ad; Ad/protein/Ad; Ad/Ad/protein, etc) are CS, LSA-1, LSA-3, MSP-1, MSP-119, MSP-142 (see below), or the hepatitis B particles-containing and CS-derived vaccine composition known as RTS,S (see Gordon et al. (1995; US 6,306,625 and WO 93/10152).

Although the invention is exemplified herein with the use of adenoviruses, it is to be understood that the invention is by no means intended to be limited to adenoviruses but also relates to the use of other recombinant viruses as delivery vehicles. Examples of viruses that can also be used for administering the antigenic determinants of the present invention are poxviruses (vaccinia viruses, such as MVA) and flaviviruses such as alphaviruses. Non-limiting examples of alphaviruses that may be applied for delivering the immunogenic *Plasmodium* components of the present invention are: Ndumu virus, Buggy Creek virus, Highland J. virus, Fort Morgan virus, Babanki virus, Kyzylagach virus, Una virus, Aura virus, Whataroa virus, Bebaru virus, South African Arbovirus No. 86, Mayaro virus, Sagiyama virus, Getah virus, Ross River virus, Barmah Forest virus, Chikungunya virus, O'nyong-nyong virus, Western Equine Encephalitis virus (WEE), Middelburg virus, Everglades virus, Eastern Encephalitis virus (EEE), Mucambo virus and Pixuna virus. Preferably, when an alphavirus is the virus of choice, Semliki Forest Virus, Sindbis virus or Venezuelan Equine Encephalitis virus are applied.

A sequence is 'derived' as used herein if a nucleic acid can be obtained through direct cloning from wild-

type sequences obtained from wild-type viruses, while they can for instance also be obtained through PCR by using different pieces of DNA as a template. This means also that such sequences may be in the wild-type form as well as in altered form. Another option for reaching the same result is through combining synthetic DNA. It is to be understood that 'derived' does not exclusively mean a direct cloning of the wild type DNA. A person skilled in the art will also be aware of the possibilities of molecular biology to obtain mutant forms of a certain piece of nucleic acid. The terms 'functional part, derivative and/or analogue thereof' are to be understood as equivalents of the nucleic acid they are related to. A person skilled in the art will appreciate the fact that certain deletions, swaps, (point) mutations, additions, etcetera may still result in a nucleic acid that has a similar function as the original nucleic acid. It is therefore to be understood that such alterations that do not significantly alter the functionality of the nucleic acids are within the scope of the present invention. If a certain adenoviral vector is derived from a certain adenoviral serotype of choice, it is also to be understood that the final product may be obtained through indirect ways, such as direct cloning and synthesizing certain pieces of genomic DNA, using methodology known in the art. Certain deletions, mutations and other alterations of the genomic content that do not alter the specific aspects of the invention are still considered to be part of the invention. Examples of such alterations are for instance deletions in the viral backbone to enable the cloning of larger pieces of heterologous nucleic acids. Examples of such mutations are for instance E3 deletions or deletions and/or alterations in the regions coding for the E2 and/or E4 proteins of

adenovirus. Such changes applied to the adenoviral backbone are known in the art and often applied, since space is a limiting factor for adenovirus to be packaged; this is a major reason to delete certain parts of the adenoviral genome. Other reasons for altering the E2, E3 and/or E4 regions of the genome may be related to stability or integrity of the adenoviral vector, as for instance described in international patent applications PCT/NL02/00280, PCT/EP03/50125, PCT/NL02/00281, PCT/EP03/50126 (non published). These applications relate amongst others to the use of an E4orf6 gene from a serotype from one subgroup in the backbone of an adenovirus from another subgroup, to ensure compatibility between the E4orf6 activity and the E1B-55K activity during replication and packaging in a packaging cell line. They further relate to the use of a proper functioning pIX promoter for obtaining higher pIX expression levels and a more stable recombinant adenoviral vector.

'Replication defective' as used herein means that the viral vectors do not replicate in non-complementing cells. In complementing cells, the functions required for replication, and thus production of the viral vector, are provided by the complementing cell. The replication defective viral vectors of the present invention do not harbor all elements enabling replication in a host cell other than a complementing cell.

'Heterologous' as used herein in conjunction with nucleic acids means that the nucleic acid is not found in wild type versions of the viral vectors in which the heterologous nucleic acid is cloned. For instance in the case of adenoviruses, the heterologous nucleic acid that is cloned in the replication defective adenoviral vector, is not an adenoviral nucleic acid.

'Antigenic determinant' as used herein means any antigen derived from a pathogenic source that elicits an immune response in a host towards which the determinant is delivered (administered). Examples of antigenic determinants of *Plasmodium* that can be delivered by using the replication defective recombinant viruses of the present invention are the circumsporozoite protein, the SE36 polypeptide, the merozoite surface protein 119 kDa C-terminal polypeptide (MSP-119), MSP-1, MSP-142, Liver Stage Antigen 1 or 3 (LSA-1 or -3), or a fragment of any of the aforementioned. In a preferred aspect the invention relates to the circumsporozoite (CS) protein from *P.falciparum*.

'Codon-optimized' as used herein means that the nucleic acid content has been altered to reach sufficiently high expression levels of the protein of interest in a host of interest to which the gene encoding said protein is delivered. Sufficiently high expression levels in this context means that the protein levels should be high enough to elicit an immune response in the host in order to give protection to a malaria-inducing parasite that may enter the treated host before or after treatment. It is known in the art that some vaccines give an immune response in humans, through which approximately 60% of the vaccinated individuals is protected against illnesses induced by subsequent challenges with the pathogen (e.g., sporozoites). Therefore the expression levels are considered to be sufficient if 60% or more of the treated individuals is protected against subsequent infections. It is believed that with the combinations of adenoviral aspects that can be applied and the choice of antigen as disclosed herein, such percentages may be reached. Preferably, 85% of the individuals are protected, while it is most preferred to have protection

to a subsequent challenge in more than 90% of the vaccinated hosts. The nucleic acids disclosed in the present invention are codon-optimized for expression in humans. According to Narum et al. (2001), the content of adenine plus thymine (A+T) in DNA of *Homo sapiens* is approximately 59%, as compared to the percentage cytosine plus guanine (C+G). The adenine plus thymine content in *P.falciparum* is approximately 80%. The adenine plus thymine content in the CS gene of *P.falciparum* is approximately 87%. To obtain sufficient protection it is believed to be necessary to improve production levels in the host. One way to achieve this is to optimize codon usage by altering the nucleic acid content of the antigenic determinant in the viral-based vector, without altering the amino acid sequence thereof. For this, the replication-defective recombinant viral vectors according to the invention have an adenine plus thymine content in the heterologous nucleic acids of the present invention of less than 87%, preferably less than 80%, and more preferably less than or equal to approximately 59%. Based on codon-usage in human and the amino acid content of the CS genes of *P.falciparum* and *yoelii*, the percentages of the codon-optimized genes were even lower, reaching approximately 45% for the amino acid content as disclosed by the present invention. Therefore, as far as the CS genes are concerned it is preferred to have an adenine plus thymine content of approximately 45%. It is to be understood, that if another species than humans is to be treated, which may have a different adenine plus thymine concentration (less or more than 59%), and/or a different codon usage, that the genes encoding the CS proteins of the present invention may be adjusted to fit the required content and give rise to suitable expression levels for that particular host. Of course, it cannot be excluded

either, that slight changes in content may result in slight expression level changes in different geographical areas around the world. It is also to be understood that slight changes in the number of repeats included in the amino acid sequence of the proteins, that percentages may differ accordingly. All these adjusted contents are part of the present invention.

EXAMPLES

10

Example 1. Assembly of the *Plasmodium falciparum* circumsporozoite synthetic gene.

Comparative studies conducted with DNA vaccines based on native and codon-optimized genes encoding merozoites proteins of *P.falciparum* have indicated a direct correlation with expression levels and immunogenicity (Narum et al. 2001). A new sequence of the gene encoding the *Plasmodium falciparum* circumsporozoite (CS) protein was designed. Studies on populations of malaria parasites obtained from widely separated geographical regions have revealed the presence of CS sequence polymorphism. The new *P.falciparum* CS sequence was assembled by alignment of the different available protein sequences present in the GeneBank database (listed in Table I). First, all the different sequences were placed in order of subgroups based on global location or by lab-strain where the isolates originated. All CS complete or partial sequences were used in order to identify variation between the different geographical areas and identified lab-strains. The final amino acid consensus sequence determined was thoroughly examined. The inventors of the present invention subsequently adjusted this consensus sequence and to have a new CS gene synthesized (Fig. 1). The novel amino acid sequence

is shown in Fig. 1A. The new CS protein harbors the aspects listed below (from N-terminus to C-terminus):

- 5 - The N-terminal signal sequence, which would direct the protein to the endoplasmic reticulum, is left unchanged.
- 10 - The HLA binding peptide amino acid (31-40), as well as region 1 (predominant B-cell epitope) are conserved, therefore these sequences are left unchanged.
- 15 - A number of repeats: there are 14-41 NANP (SEQ ID NO:10) repeats in the different isolates and 4 NVDP (SEQ ID NO:11) repeats. It was chosen to incorporate 27 NANP repeats, a cluster of 3 NVDP repeats and one separate NVDP repeat.
- 20 - The ENANANNAVKN (SEQ ID NO:12) sequence directly downstream of the repeats mentioned above, was found to be reasonably conserved between strains.
- 25 - The Th2R region and the immunodominant CD8 epitope (Lockyer et al. 1989; Zevering et al. 1994): a single consensus sequence that differs in some respects from that of the known, and frequently used lab-strain 3D7 sequence was determined. This sequence is sometimes referred to as the 'universal epitope' in literature (Nardin et al. 2001).
- 30 - The region 2, overlapping with the Th2R region, remained conserved.
- The TH3R region, which is considered to be a less important CD8 epitope, is used in the form of a consensus sequence, since only point mutations were found.
- The C-terminal 28 amino acids, which constitute a GPI signal anchor sequence, that is inefficient in mammalian cells (Moran and Caras, 1994), and not

hydrophobic by itself to serve as a stable membrane anchor. The gene was constructed such that the whole sequence can be removed, but also leaving open the possibility of remaining present. This allows a comparison of the antigenicity of adenovirus vectors carrying a full length CS versus those expressing the protein deleted in the GPI signal anchor sequence. In fact it has been described that removal of the GPI signal sequence from a CS DNA vaccine enhanced induction on immune response against malaria infection in rodents (Scheiblhofer et al. 2001).

- Substitution S to A at position 373: this amino acid substitution was introduced to eliminate a potential glycosylation site recognized by mammalian cells.

Since the malaria parasite residue usage (87% A and T) is significantly different from that of the *Homo sapiens*, the gene encoding the newly designed CS protein was codon-optimized in order to improve its expression in mammalian cells, taking care of the following aspects to avoid cis-acting sequences: no premature poly(A) sites and internal TATA boxes should be present; Chi-sites, ribosomal entry sites and AT-rich sequence clusters should be avoided; no (cryptic) splice acceptor and -donor sites should be present; repetitive sequence stretches should be avoided as much as possible; and GC-rich sequences should also be avoided. The final codon-optimized gene is shown in Fig. 1B.

The newly designed CS consensus sequence was synthesized and cloned into pCR-script (Stratagene) by GeneArt (Regensburg, Germany), using methodology known to persons skilled in the art of synthetic DNA generation,

giving rise to a clone named 02-148 (pCR-script.Pf) (SEQ ID NO:1).

Next to this synthetic clone, another synthetic gene was generated, wherein a number of mutations were introduced in the 3' end, to obtain an amino acid sequence that is identical to the *P.falciparum* CS protein of the 3D7 strain, which is deleted in the last 14 amino acids (Fig. 2). This gene was also codon-optimized using the same provisions as described above and subsequently synthesized and cloned into pCR-script (Stratagene) by GeneArt. The clone was named 02-659 (pf-aa-sub) (SEQ ID NO:4).

Example 2. Codon-optimization of the circumsporozoite gene of the rodent-specific malaria parasite *Plasmodium yoelii*.

Malaria species that have been adapted to robust rodent models, such as *P.berghei* and *P.yoelii*, have been powerful tools for identification and testing of malaria candidate vaccines. Since infectivity of *P.yoelii* sporozoites resembles that of *P.falciparum*, it was decided to make use of the *P.yoelii* model for exemplification of the capability of Ad35 vectors carrying codon-optimize CS proteins to provide sterile immunity and therefore protection against malaria infection. The *P.yoelii* CS gene, encoding for residues 1-356 as previously described (Rodrigues et al. 1997), was codon-optimized using the same provisions as described above and synthesized by GeneArt (GmbH-Regensburg, Germany). The sequence of the codon-optimized *P.yoelii* CS gene (plasmid 02-149) is depicted in Fig. 3.

Example 3. Generation of recombinant adenoviral vectors based on Ad5.

RCA-free recombinant adenoviruses can be generated very efficiently using adapter plasmids, such as pAdApt, and adenovirus plasmid backbones, such as pWE/Ad.AflIII-rITRsp. Methods and tools have been described extensively elsewhere (WO 97/00326, WO 99/55132, WO 99/64582, WO 00/70071, WO 00/03029). Generally, the adapter plasmid containing the transgene of interest in the desired expression cassette is digested with suitable enzymes to free the recombinant adenovirus sequences from the plasmid vector backbone. Similarly, the adenoviral complementation plasmid pWE/Ad.AflIII-rITRsp is digested with suitable enzymes to free the adenovirus sequences from the vector plasmid DNA.

The cloning of the gene encoding the CS protein from *P.yoelii* parasite into pIPspAdapt1 was performed as follows. Plasmid 02-149 (GeneArt, see above) containing the codon optimized CS gene was digested with HindIII and BamHI restriction enzymes. The 1.1 Kb fragment corresponding to the *P.yoelii* CS gene was isolated from agarose gel and ligated to HindIII and BamHI-digested pIPspAdapt1 vector (described in WO 99/64582). The resulting plasmid was named pAdapt.CS.Pyoel and contains the CS gene under the transcriptional control of full length human immediate-early (IE) cytomegalovirus (CMV) promoter and a downstream SV40 poly(A) signal.

The cloning of the gene encoding the full length CS protein from *P.falciparum* parasite into pIPspAdapt1 was performed as follows. Plasmid 02-148 pCR-script.Pf (see above) containing the codon optimized CS gene was digested with HindIII and BamHI restriction enzymes. The 1.2 kb fragment corresponding to the CS gene was isolated from agarose gel and ligated to HindIII and BamHI-digested pIPspAdapt1 vector. The resulting plasmid was named pAdapt.CS.Pfalc and contains the CS gene under the

transcriptional control of full length human immediate-early (IE) cytomegalovirus (CMV) promoter and the downstream SV40 poly(A) signal.

The cloning of the gene encoding the CS *P.falciparum* protein minus the GPI anchor sequence, thus with the deletion of the last 28 amino acids, into pIPspAdapt1 was performed as follows. A 1.1 kb PCR fragment was amplified using plasmid 02-148 as template, with the primers Forw.Falc (5'-CCA AGC TTG CCA CCA TGA TGA GG-3') (SEQ ID NO:13) and Rev.Falc.CS-28 (5'-CCG GAT CCT CAG CAG ATC TTC TTC TCG-3') (SEQ ID NO:14). Primers were synthesized by Invitrogen. For the PCR, the enzyme Pwo DNA polymerase (Inno-train Diagnostic) was used, while the following program was applied: 1 cycle of 5 min at 94°C, 1 min at 50°C, 2 min 30 sec at 72°C; 5 cycles of 1 min at 94°C, 1 min at 50°C, 2 min 50 sec at 72°C; 20 cycles of 1 min at 94°C, 1 min at 54°C, 2 min 50 sec at 72°C; and 1 cycle of 1 min at 94°C, 1 min at 54°C, followed by 10 min at 72°C. The amplified PCR product was digested with the restriction enzymes HindIII and BamHI and then cloned into pIPspAdapt1 which was also digested with HindIII and BamHI. The resulting plasmid was designated pAdapt.CS.Pfalc(-28) and contains the CS gene under the transcriptional control of full length human immediate-early (IE) cytomegalovirus (CMV) promoter and the downstream SV40 poly(A) signal.

The cloning of the gene encoding the CS *P.falciparum* protein minus the GPI anchor, thus with the deletion of the last 14 amino acids, into pIPspAdapt1 was performed as follows. A 1.1 kb PCR fragment was amplified using plasmid 02-148 as template, with the primers Forw.Falc (SEQ ID NO:13) and Rev.Falc.CS-14 (5'-CCG GAT CCT CAG CTG

TTC ACC ACG TTG-3') (SEQ ID NO:15). Primers were synthesized by Invitrogen. For the PCR, the enzyme Pwo DNA polymerase (Inno-train Diagnostic) was used, while the following program was applied: 1 cycle of 5 min at 94°C, 1 min at 50°C, 2 min 30 sec at 72°C; 5 cycles of 1 min at 94°C, 1 min at 50°C, 2 min 50 sec at 72°C; 20 cycles of 1 min at 94°C, 1 min at 54°C, 2 min 50 sec at 72°C; and 1 cycle of 1 min at 94°C, 1 min at 54°C, followed by 10 min at 72°C. The amplified PCR product was digested with the restriction enzymes HindIII and BamHI and then cloned into pIPspAdapt1 also digested with HindIII and BamHI. The resulting plasmid was designated pAdapt.CS.Pfalc(-14) and contains the CS gene under the transcriptional control of full length human immediate-early (IE) cytomegalovirus (CMV) promoter and the downstream SV40 poly(A) signal.

The cloning of the gene encoding the CS *P.falciparum* protein minus the GPI anchor, displaying a C-terminal sequence as in the 3D7 strain, into pIPspAdapt1 is performed as follows. Plasmid 02-659 pf-aa-sub (see above) containing the codon-optimized CS gene is digested with HindIII and BamHI restriction enzymes. The 1.1 kb fragment corresponding to the CS gene is ligated to HindIII and BamHI-digested pIPspAdapt1 vector. The resulting plasmid is designated pAdapt.CS.Pfalc (pf-aa-sub) and contains the CS gene under the transcriptional control of full length human immediate-early (IE) cytomegalovirus (CMV) promoter and the downstream SV40 poly(A) signal.

The generation of the recombinant virus named Ad5ΔE3.CS.Pyoel was performed as follows. pAdapt.CS.Pyoel

was digested by PacI restriction enzyme to release the left-end portion of the Ad genome. Plasmid pWE/Ad.AflIII-rITRsp containing the remaining right-end part of the Ad genome has a deletion of 1878 bp in the E3 region (XbaI deletion). This construct was also digested with PacI. 5 pAdapt.CS.Pyoel was separately transfected with PacI digested pWE/Ad.AflIII-rITRsp into PER-E1B55K producer cells (cells have been described in WO 02/40665) using lipofectamine transfection reagent (Invitrogen) using 10 methods known in the art and as described in WO 00/70071. Homologous recombination between overlapping sequences led to generation of the recombinant virus named Ad5ΔE3.CS.Pyoel. It is to be understood that Ad5-based vectors can also be produced on PER.C6™ cells, which 15 cells are represented by the cells deposited under ECACC no. 96022940 (see above). The adenoviral vector, in crude lysates, resulting from this transfection were plaque purified using methods known to persons skilled in the art. Single plaques were analyzed for the presence of the 20 CS transgene and amplified for large-scale production in triple-layer flasks (3x175 cm²/flask). Upon amplification cells are harvested at full CPE and the virus is purified by a two-step Cesium Chloride (CsCl) purification procedure as routinely done by those skilled in the art 25 and generally as described in WO 02/40665.

The generation of the recombinant virus named Ad5ΔE3.CS.Pfalc was performed as follows. pAdapt.CS.Pfalc was digested by PacI restriction enzyme to release the left-end portion of the Ad genome. Plasmid pWE/Ad.AflIII-rITRsp containing the remaining right-end part of the Ad genome has a deletion of 1878 bp in the E3 region (XbaI deletion). This construct was also digested with PacI. 30 pAdapt.CS.Pfalc was transfected with PacI digested pWE/Ad.AflIII-rITRsp into PER-E1B55K producer cells using

lipofectamine transfection reagent. Homologous recombination between overlapping sequences led to generation of the recombinant virus named Ad5ΔE3.CS.Pfalc. The adenoviral vector, in crude lysates, resulting from this transfection was plaque purified using methods known to persons skilled in the art. Single plaques were analyzed for the presence of the CS transgene and amplified for large-scale production in triple-layer flasks (3x175 cm²/flask). Cells were harvested at full CPE and the virus was purified by a two-step CsCl purification procedure as routinely done by those skilled in the art and generally as described in WO 02/40665.

The generation of the recombinant viruses named Ad5ΔE3.CS.Pfalc(-28) and Ad5ΔE3.CS.Pfalc(-14) was performed as follows. pAdapt.CS.Pfalc(-28) and pAdapt.CS.Pfalc(-14) were separately digested by PacI restriction enzyme to release the left-end portion of the Ad genome. Plasmid pWE/Ad.AflII-rITRsp containing the remaining right-end part of the Ad genome has a deletion of 1878 bp in the E3 region (XbaI deletion). This construct was also digested with PacI. pAdapt.CS.Pfalc(-28) and pAdapt.CS.Pfalc(-14) were separately transfected with PacI digested pWE/Ad.AflII-rITRsp into PER-E1B55K producer cells using lipofectamine transfection reagent. Homologous recombination between overlapping sequences led to generation of recombinant viruses named respectively Ad5ΔE3.CS.Pfalc(-28) and Ad5ΔE3.CS.Pfalc(-14). Adenoviral vectors in crude lysates resulting from these transfections are plaque purified using methods known to persons skilled in the art. Single plaques are analyzed for the presence of the CS transgene and amplified for large-scale production in triple-layer flasks (3x175 cm²/flask). Cells are harvested at full CPE .

and the virus is purified by a two-step CsCl purification procedure as routinely done by those skilled in the art and generally as described in WO 02/40665.

The generation of the recombinant virus named

5 Ad5ΔE3.CS.Pfalc(pf-aa-sub) is performed as follows. pAdapt.CS.Pfalc(pf-aa-sub) is digested by PacI restriction enzyme to release the left-end portion of the Ad genome. Plasmid pWE/Ad.AflIII-rITRsp containing the remaining right-end part of the Ad genome is also

10 digested with PacI. pAdapt.CS.Pfalc(pf-aa-sub) is transfected with PacI digested pWE.Ad.AflIII-rITRspΔE3 into PER.C6™ or PER-E1B55K producer cells using lipofectamine transfection reagent, or by other means such as electroporation or other transfection methods

15 known to persons skilled in the art. Homologous recombination between overlapping sequences leads to generation of the recombinant virus named Ad5ΔE3.CS.Pfalc(pf-aa-sub). The adenoviral vector, in crude lysates, resulting from this transfection is plaque purified using

20 methods known to persons skilled in the art. Single plaques are analyzed for the presence of the CS transgene and amplified for large-scale production in triple-layer flasks (3x175 cm²/flask). Cells are harvested at full CPE and the virus is purified by a two-step CsCl purification

25 procedure as routinely done by those skilled in the art and generally as described in WO 02/40665.

Next to these procedures, generation of the control recombinant adenovirus named Ad5ΔE3.empty was carried out as described above, using as adapter the plasmid pAdapt,

30 lacking a transgene.

Example 4. Generation of recombinant adenoviral vaccine vectors based on Ad35.

A first 101 bp PCR fragment containing the Ad5 pIX promoter (nucleotides 1509-1610) was generated with the
5 primers SV40for (5'-CAA TGT ATC TTA TCA TGT CTA G-3')
(SEQ ID NO:16) and pIX5Rmfe (5'-CTC TCT CAA TTG CAG ATA
CAA AAC TAC ATA AGA CC-3') (SEQ ID NO:17). The reaction
was done with Pwo DNA polymerase according to
manufacturers instructions but with 3% DMSO in the final
10 mix. pAdApt was used as a template. The program was set
as follows: 2 min at 94°C; 30 cycles of: 30 sec at 94°C,
30 sec at 52°C and 30 sec at 72°C; followed by 8 min at
72°C. The resulting PCR fragment contains the 3' end of
the SV40 poly(A) signal from pAdApt and the Ad5-pIX
15 promotor region as present in Genbank Accession number
M73260 from nucleotide 3511 to nucleotide 3586 and an
MfeI site at the 3' end. A second PCR fragment was
generated as described above but with primers pIX35Fmfe
(5'-CTC TCT CAA TTG TCT GTC TTG CAG CTG TCA TG-3') (SEQ
20 ID NO:18) and 35R4 (for reference to the sequence of the
35R4 primer, see WO 00/70071). pAdApt35IP1 (described in
WO 00/70071) was used as a template, the annealing was set
at 58°C for 30 sec and the elongation of the PCR program
was set at 72°C for 90 sec. This PCR procedure amplifies
25 Ad35 sequences from nucleotide 3467 to nucleotide 4669
(sequence numbering as in WO 00/70071) and adds an MfeI
site to the 5' end. Both PCR fragments were digested with
MfeI and purified using the Qiagen PCR purification kit
(Qiagen). Approximate equimolar amounts of the two
30 fragments were used in a ligation reaction. Following an
incubation of two hours with ligase enzyme in the correct
buffers, at room temperature, the mixture was loaded on
an agarose gel and the DNA fragments of 1.4 kb length
were isolated with the Geneclean II kit (BIO101, Inc).

The purified DNA was used in a PCR amplification reaction with primers SV40for and 35R4. The PCR was done as described above with an annealing temperature of 52°C and an elongation time of 90 sec. The resulting product was
5 isolated from gel using the Qiagen gel extraction kit and digested with AgeI and BglII. The resulting 0.86 kb fragment containing the complete 100 nucleotide pIX promoter from Ad5, the MfeI site and the pIX ORF (fragment MfeI-AgeI, including the ATG start site) from
10 Ad35, but without a poly(A) sequence, was isolated from gel using the Geneclean II kit.

RCA-free recombinant adenoviruses based on Ad35 can be generated very efficiently using adapter plasmids, such as pAdApt535 (described below) and adenovirus
15 plasmid backbones, such as pWE/Ad35.pIX-rITRAE3 (described in WO 02/40665). To generate pAdApt535, pAdApt35.Luc (described in WO 00/70071) was digested with BglII and AgeI and the resulting 5.8 kb vector was isolated from gel. This fragment was ligated with the
20 isolated 0.86 kb BglII-AgeI fragment containing the Ad5-Ad35 chimeric pIX promoter described above, to result in a plasmid named pAdApt535.Luc, which was subsequently digested with BglII and ApaI. The resulting 1.2 kb insert was purified over gel. pAdApt35IP1 was digested with
25 BglII and ApaI and the 3.6-kb vector fragment was isolated over gel. Ligation of the 1.2 kb BglII-ApaI insert from pAdApt535.Luc and the 3.6 kb BglII-ApaI digested vector resulted in pAdApt535.

The cloning of the gene encoding the CS protein from
30 *P.yoelii* parasite into pAdapt535 was performed as follows. Plasmid 02-149 containing the codon optimized *P.yoelii* CS gene (see above) was digested with the restriction enzymes HindIII and BamHI. The 1.1 kb fragment corresponding to the CS gene was isolated over

agarose gel and ligated to the HindIII and BamHI digested pAdapt535 vector. The resulting plasmid was named pAdapt535-CS.Pyoel and contains the CS gene under the transcriptional control of the full length human CMV promoter and the downstream SV40 poly(A) signal.

The cloning of the gene encoding the full length CS protein from *P.falciparum* parasite into pAdapt535 was performed as follows. Plasmid 02-148 (pCR-script.Pf) containing the codon optimized CS gene of *P.falciparum* was digested with the restriction enzymes HindIII and BamHI. The 1.2 kb fragment corresponding to the CS gene was isolated over agarose gel and ligated to the HindIII and BamHI digested pAdapt535 vector. The resulting plasmid was named pAdapt535-CS.Pfalc and contains the CS gene under the transcriptional control of the full length human CMV promoter and the downstream SV40 poly(A) signal.

The cloning of the gene encoding the CS *P.falciparum* protein minus the GPI anchor sequence, thus with the deletion of the last 28 amino acids, into pAdapt535 is performed as follows. The 1.1 kb PCR fragment obtained as described above using primers Forw.Falc and Rev.Falc-CS-28, is digested with the restriction enzymes HindIII and BamHI and then cloned into pAdapt535 vector also digested with HindIII and BamHI. The resulting plasmid is designated pAdapt535-CS.Pfalc(-28) and contains the CS gene under the transcriptional control of the full length human CMV promoter and the downstream SV40 poly(A) signal.

The cloning of the gene encoding the CS *P.falciparum* protein minus the GPI anchor sequence, now with the deletion of the last 14 amino acids, into pAdapt535 is

performed as follows. The 1.1 kb PCR fragment obtained as described above using primers Forw.Falc and Rev.Falc.CS-14, is digested with the restriction enzymes HindIII and BamHI and then cloned into pAdapt535 vector also digested with HindIII and BamHI. The resulting plasmid is designated pAdapt535.CS.Pfalc(-14) and contains the CS gene under the transcriptional control of the full length human CMV promoter and the downstream SV40 poly(A) signal.

10 The cloning of the gene encoding the CS *P.falciparum* protein minus the GPI anchor sequence, displaying a C-terminus sequence as in the 3D7 strain, into pAdapt535 is performed as follows. Plasmid 02-659 pf-aa-sub (see above) containing the codon optimized CS gene is digested with HindIII and BamHI restriction enzymes. The 1.1 kb fragment corresponding to the CS gene is ligated to HindIII and BamHI digested pAdapt535 vector. The resulting plasmid is designated pAdapt535.CS.Pfalc(pf-aa-sub) and contains the CS gene under the transcriptional control of the full length human CMV promoter and the downstream SV40 poly(A) signal.

The generation of the recombinant virus named Ad35AE3.CS.Pyoel was performed as follows.

25 pAdapt535.CS.Pyoel was digested by PacI restriction enzyme to release the left-end portion of the Ad genome. Plasmid pWE.Ad35.pIX-rITRAE3, containing the remaining right-end part of the Ad genome with a deletion of 2673 bp in the E3 region is digested with NotI.

30 pAdapt535.CS.Pyoel was transfected with NotI digested pWE.Ad35.pIX-rITRAE3 into PER-E1B55K producer cells using lipofectamine transfection reagent. The generation of the cell line PER-E1B55K has been described in detail in WO 02/40665. In short, this publication describes that

PER.C6™ cells were stably transfected with ScaI linearised pIG35-55K DNA, carrying the E1B-55K gene of adenovirus serotype 35, after which a selection procedure with G418 yielded in 196 picked colonies. Further
5 culturing of a limited number of well growing colonies resulted in stable cell lines that upon numerous subcultures stably expressed the Ad35 E1B-55K gene and supported the growth of recombinant Ad35 viruses, while the original PER.C6™ cell were very inefficient in
10 supporting this.

Homologous recombination between overlapping sequences led to generation of the recombinant virus named Ad35ΔE3.CS.Pyoel. The adenoviral vector, in crude lysates, resulting from this transfection was plaque
15 purified using methods known to persons skilled in the art. Single plaques were analyzed for the presence of the CS transgene and amplified for large-scale production in triple-layer flasks (3x175 cm²/flask). Upon amplification cells were harvested at full CPE and the virus was
20 purified by a two-step CsCl purification procedure as routinely done by those skilled in the art and generally as described in WO 02/40665.

The generation of the recombinant virus named Ad35ΔE3.CS.Pfalco was performed as follows.
25 pAdapt535.CS.Pfalco was digested by PacI restriction enzyme to release the left-end portion of the Ad genome. Plasmid pWE.Ad35.pIX-rITRAE3, containing the remaining right-end part of the Ad genome with a deletion of 2673 bp in the E3 region was digested with NotI.
30 pAdapt535.CS.Pfalco was transfected with NotI digested pWE.Ad35.pIX-rITRAE3 into PER-E1B55K producer cells using lipofectamine transfection reagent. Homologous recombination between overlapping sequences led to generation of the recombinant virus named

Ad35ΔE3.CS.Pfalc. The adenoviral vector, in crude lysates, resulting from this transfection was plaque purified using methods known to persons skilled in the art. Single plaques were analyzed for the presence of the CS transgene and amplified for large-scale production in triple-layer flasks (3x175 cm²/flask). Upon amplification cells were harvested at full CPE and the virus was purified by a two-step CsCl purification procedure as routinely done by those skilled in the art and generally as described in WO 02/40665.

The generation of the recombinant viruses named Ad35ΔE3.CS.Pfalc(-28) and Ad35ΔE3.CS.Pfalc(-14) is performed as follows. pAdapt535.CS.Pfalc(-28) and pAdapt535.CS.Pfalc(-14) were separately digested by PacI restriction enzyme to release the left-end portion of the Ad genome. Plasmid pWE.Ad35.pIX-rITRAE3 containing the remaining right-end part of the Ad genome is digested with NotI. pAdapt535.CS.Pfalc(-28) and pAdapt535.CS.Pfalc(-14) are separately transfected with NotI digested pWE.Ad35.pIX-rITRAE3 into PER-E1B55K producer. Homologous recombination between overlapping sequences leads to generation of recombinant viruses named respectively Ad35ΔE3.CS.Pfalc(-28) and Ad35ΔE3.CS.Pfalc(-14). Adenoviral vectors in crude lysates resulting from these transfections are plaque purified using methods known to persons skilled in the art. Single plaques are analyzed for the presence of the CS transgene and amplified for large-scale production in triple-layer flasks (3x175 cm²/flask). Cells are harvested at full CPE and the virus is purified by a two-step CsCl purification procedure as routinely done by those skilled in the art and generally as described in WO 02/40665.

The generation of the recombinant virus named Ad35ΔE3.CS.Pfalc(pf-aa-sub) is performed as follows. pAdapt535.CS.Pfalc(pf-aa-sub) is digested by PacI restriction enzyme to release the left-end portion of the

5 Ad genome. Plasmid pWE.Ad35.pIX-rITRAE3 containing the remaining right-end part of the Ad genome is digested with NotI. pAdapt535.CS.Pfalc(pf-aa-sub) is transfected with NotI digested pWE.Ad35.pIX-rITRAE3 into PER-E1B55K producer cells using lipofectamine transfection reagent

10 (Invitrogen) using methods known in the art and as described in WO 00/70071 or by electroporation or other transfection methods known to those skilled in the art. Homologous recombination between overlapping sequences leads to generation of the recombinant virus named

15 Ad35ΔE3.CS.Pfalc(pf-aa-sub). The adenoviral vector, in crude lysates, resulting from this transfection is plaque purified using methods known to persons skilled in the art. Single plaques are analyzed for the presence of the CS transgene and amplified for large-scale production in

20 triple-layer flasks (3x175 cm²/flask). Cells are harvested at full CPE and the virus is purified by a two-step CsCl purification procedure as routinely done by those skilled in the art and generally as described in WO 02/40665.

25 **Example 5. Inducing protection against *P.yoelii* malaria infection using recombinant adenoviral-based vaccines in vivo.**

Adenovirus serotype 5 (Ad5)-based vectors genetically engineered to express the CS antigen of the

30 rodent malaria *P.yoelii* have been shown capable to induce complete protection against *P.yoelii* infection (Rodrigues et al. 1997). A side-by-side comparison between Ad5 and Ad35 vectors carrying the codon-optimized *P.yoelii* CS gene was designed to investigate the immune response that

is induced, and to investigate their ability in raising protection against *P.yoelii* parasite infection in mice. The study enrolled Balb/C mice that were immunized by intra-muscular or subcutaneous injection of 10^8 - 10^{10} viral particles (vp) of Ad5ΔE3- or Ad35ΔE3-based viral vectors (as described above) carrying either the *P.yoelii* CS gene (Ad5ΔE3-CS.Pyoel and Ad35ΔE3-CS.Pyoel) or no transgene (Ad5ΔE3-empty and Ad35ΔE3-empty). Figure 4 shows the results of the experiments wherein the administration route was compared using both vectors. The number of IFN- γ -secreting cells in a population of 10^6 splenocytes was determined (Fig. 4A) as well as the antibody titers in the serum (Fig. 4B). The experiments were performed on mice that were sacrificed two weeks after injection with the recombinant adenoviruses. Each of the bars represents the average of 5 mice. If mice were not sacrificed they were used for a challenge with live sporozoites, after which the rate of protection was determined (Fig. 5A and B). Each of these bars represents the average of 5 mice. The experiments on humoral and cellular immune responses are performed with immunological assays well known to persons skilled in the art and as described for instance by Bruña-Romero et al. (2001a). The immunization, challenge and read out are scheduled in Table II and III. Antibodies titers against sporozoites can be determined by an indirect immunofluorescence assay or with an ELISA. Figure 4B shows the results as calculated with an ELISA. Cellular immune responses were determined by ex-vivo ELISPOT assay measuring the relative number of CS-specific, IFN- γ -secreting, CD8+ and CD4+ T cells. Protection against malaria infection was monitored by determining the levels of parasite inhibition in the

livers of immunized mice through reverse transcriptase PCR quantification of *P.yoelii* ribosomal RNA copies.

The immunization with Ad5- and Ad35-based vectors was performed as follows. Aliquots of recombinant adenoviruses that were stored at -70°C were gradually thawed on ice and diluted to 100 µl in the desired concentration in PBS with 1% heat-inactivated Normal Mouse Serum. Subsequently the samples were sonicated for 5 sec. Sub-cutaneous administration was performed at both sides of the tail base with a volume of 50 µl at each side. Intra muscular administration was performed in both thighs with a volume of 50 µl at each thigh.

The Indirect Immunofluorescence Assay (IFA) is performed according to Bruña-Romero et al. (2001a). First, infected mosquitoes are generated by initially having a native mouse infected with an infected mosquito by having the mouse bitten at three different sites. Blood is removed from the mouse after 8 days when parasitemia is 4-8% and diluted to 1%. Then, other naïve mice are injected i.p. with the diluted blood sample. After 3 days the blood is taken which serves as a blood meal for starved mosquitoes. These are fed for 2 days. After 14 days *P.yoelii* sporozoites are isolated from the blood-fed mosquitoes by anaesthetizing infected mosquitoes on ice and subsequently saturating them in 70% ethanol. Then, the mosquitoes are transferred to PBS pH 7.4 and the salivary glands are dissected. These are subsequently grinded on ice and the sporozoites are separated from the debris by centrifugation. Using this method, approximately 35,000 *P.yoelii* sporozoites can be obtained from 1 mosquito. Then, glass-slides in a 12-multi-well plate are coated with approximately 10,000 *P.yoelii* sporozoites each in Dulbecco's Modified Eagle's

Medium (DMEM) plus 10% Fetal Bovine Serum by air-drying. A range of dilutions of sera of the vaccinated mice (in a volume of 10 μ l in PBS plus 5% FBS) is subsequently incubated with the air-dried sporozoites for 30 min at room temperature in a moistures environment. Then, the slides are aspirated, washed twice with PBS and 10 μ l of a 30-fold diluted FITC conjugated Goat-anti-Mouse antibody (Kirkegaard & Perry Laboratories, USA, catalogue no. 02-18-06) is added and incubated for 30 min at room temperature. Wells were again aspirated and washed twice. For counterstaining, a solution of 100 μ g/ml Ethidium Bromide is incubated for 10 min, after which the aspiration step is repeated and the wells are washed with water. Slides are mounted using permount containing phenylenediamine/anti fade. The anti-sporozoite antibody titers are determined as the highest serum dilution producing fluorescence. For the determination of antibody titers, one can also use an ELISA. For this, ELISA plates (Immulon II, Dynatech) were coated with 2 μ g/ml antigen in PBS by adding 100 μ l per well of this solution and leaving it overnight at 4°C. The antigen that was used is a 3x6 amino acid repeat of the *P.yoelii* CS protein: QGPGAPQGPGAPQGPGAP (SEQ ID NO:19). The plates were subsequently washed three times with washing buffer (1x PBS, 0.05% Tween), and 200 μ l blocking buffer (10% FCS in washing solution) was added per well. Plates were incubated for 1-2 h at room temperature. Then, plates were washed three times again with washing buffer including 5% FCS. Dilutions of the sera were made as follows: 50 μ l washing buffer plus 5% FCS was added to wells 2-12. Then 100 μ l washing buffer plus 5% FCS is added to the first well and 1:2 serial dilutions are made by transferring 50 μ l from well 1 to 2, then from 2 to 3, etc. Plates are incubated for 1 h at room temperature.

Then the plates are washed three times with washing buffer and 100 μ l of a 1:2000 diluted peroxidase-labeled Goat anti-Mouse IgG (anti Heavy and Light chain, human absorbed, Kirkegaard & Perry Laboratories, catalogue no. 074-1806) is added per well and incubated. Then, plates are washed with washing buffer three times and once with PBS and then 100 μ l ABTS substrate solution (ABTS 1-Component, Kirkegaard & Perry Laboratories, catalogue number 50-66-18) is added to each well. The reaction is terminated by the addition of 50 μ l 1% SDS, and plates are read at 405 nm in an ELISA reader.

The ELISPOT assay to determine the relative number of CS-specific IFN- γ -secreting, CD8+ and CD4+ T cells in the spleen, and the reverse transcriptase PCR and real-time PCR to quantify the amount of parasite specific RNA present in the liver of the challenged mice were all performed as described by Bruña-Romero et al. (2001a and 2001b) except for the fact that the number of cycles in the real-time PCR was 45.

While attenuation *P.yoelii* infection in Ad5 Δ E3-CS.Pyoel vaccine recipients is predicted (Rodrigues et al. 1997), vaccination with Ad35 Δ E3-CS.Pyoel is expected to be superior or at least equally effective.

Figure 4A shows that with an administration of 10^9 and 10^{10} viral particles per mouse the Ad35-based vector is at least as effective in inducing a cellular immune response as the Ad5-based vector, if not superior. It can be concluded that with this set-up that there is no dramatic difference in cellular response as indicated by the number of IFN- γ -secreting cells after intra muscular and subcutaneous delivery.

Figure 4B shows the antibody titers in the same experiment and performed on the same sera using the indirect immuno-fluorescence experiment outlined above. If compared to the results shown in Figure 4A it is clear that at a dose of 10^9 viral particles, the Ad35 based vector induces a significant cellular immune response but does not give rise to very high titers of anti-sporozoite antibodies. Again, there is not a significant difference between the two routes of administration.

Animals that received different doses of Ad5- and Ad35-based vectors expressing the codon-optimized *P.yoelii* CS antigen, were subsequently challenged i.v. with 10^5 sporozoites purified as described above. The results of these experiments are shown in Figure 5A and B. The percentage of inhibition was calculated as compared to naïve mice that were not immunized.

Mice that were immunized received s.c. 10^9 or 10^{10} viral particles (vp) and were challenged after 14 days with the sporozoites and then sacrificed after 48 h. Negative controls were empty vectors without antigen and non-immunized mice. Clearly, a high percentage of inhibition is obtained when using the Ad5-based vector as well as with the Ad35-based vector, applying the two doses, while no protection was found in the negative controls (Fig. 5A). Importantly, only a low number of parasite-specific 18S ribosomal RNA's could be determined in the liver of the immunized mice, while the mice that received no adenoviral vector or empty vectors contained large numbers of these RNA's (Fig. 5B). This strongly indicates that the Ad35-based vector, like the Ad5-based vector can give rise to significant protection against the malaria parasite, even after a single round of immunization.

Example 6. Inducing immunity against *P.falciparum* malaria infection using recombinant adenoviral-based vaccines in vivo.

A side-by-side comparison between Adenovirus
5 serotype 5 (Ad5) and Adenovirus serotype 35 (Ad35)
vectors is designed to investigate the ability to induce
humoral and cellular immune responses against the CS
antigen of the *P.falciparum* parasite in mice. In
addition, immunogenicities of Adenovirus vectors
10 containing full length and GPI minus CS are compared.
This study enrolls B10.BR mice. Animals are immunized by
intra-muscular injection of 10^8 - 10^{10} vp of Ad5ΔE3 or
Ad35ΔE3 viral vectors carrying either the full length CS
gene (Ad5ΔE3-CS.Pfalc and Ad35ΔE3-CS.Pfalc) or the GPI-
15 anchor sequence minus CS gene (Ad5ΔE3-CS.Pfalc.(-28)/(-
14) and Ad35ΔE3-CS.Pfalc.(-28)/(-14) or no transgene
(Ad5ΔE3-empty and Ad35ΔE3-empty). At two weeks and six to
eight weeks post-vaccination, cellular and humoral
responses are monitored with immunological assays well
20 known to persons skilled in the art as described above.
The immunization, challenge and read out is scheduled in
Table IV and V.

Immunogenicity of the Ad35-based vectors is expected
to be superior or at least comparable to the
25 immunogenicity triggered by Ad5-based vectors. Figure 6
shows the results that were obtained by using the Ad5-
based vector containing the full length gene encoding the
P.falciparum CS protein, the gene encoding the protein
with the 14 amino acid deletion and the gene encoding the
30 protein with the 28 amino acid deletion. The results
indicate that all three (Ad5-based) vectors are able to
induce a cellular immune response as measured by the
number of CS-specific IFN- γ -secreting cells in a

population of splenocytes, determined by the *ex-vivo* ELISPOT assay described above, and generally as in Bruña-Romero et al. (2001a).

5 Example 7. Inducing a long-lasting protection against *P.yoelii* malaria infection by prime-boost regimens with different adenovirus serotype-based vaccines.

Recombinant Adenovirus serotype 5 expressing a CS antigen of *P.yoelii* was shown to elicit protection when
10 used in prime-boost regimen in combination with a recombinant vaccinia virus carrying the same antigen (Bruña-Romero et al. 2001a). An experiment to investigate the capability of prime/boost regimens based on adenovirus vectors carrying codon-optimized CS and
15 derived from two different serotypes to induce long-lasting protection against the *P. yoelii* CS antigen was designed. This study enrolls Balb/C mice distributed in experimental groups of 12 mice each. Animals are immunized by intra-muscular injection of an optimal dose
20 of Ad5ΔE3 or Ad35ΔE3 viral vectors carrying either the *P.yoelii* CS gene (Ad5ΔE3-CS.Pyoel and Ad35ΔE3-CS.Pyoel) or no transgene (Ad5ΔE3-empty and Ad35ΔE3-empty). One group of animals is primed at week 0 with Ad5ΔE3-CS.Pyoel and boosted at week 8 with Ad35ΔE3-CS.Pyoel. Another
25 group of mice is primed at week 0 with Ad35ΔE3-CS.Pyoel and boosted at week 8 with Ad5ΔE3-CS.Pyoel. Other groups of mice are primed at week 0 with Ad35ΔE3-CS.Pyoel or Ad5ΔE3-CS.Pyoel and boosted at week 8 with the same vector. Finally, a control group of mice is primed at
30 week 0 with Ad5ΔE3-empty and boosted at week 8 with Ad35ΔE3-empty. At week 2 post-boost, 6 mice of each group are sacrificed to allow evaluation and characterization of humoral and cellular immune responses with

immunological assays well known to persons skilled in the art, the remaining 6 mice from each group are challenged with live sporozoites. The immunization, challenge and read out are scheduled in Table VI. Protection against malaria infection will be monitored and measured using assays well known to people skilled in the art as described above. Vaccine regimens based on Ad35 alone or Ad5/Ad35 combinations are expected to be superior or at least comparable in efficacy as compared to regimens based solely on Ad5.

Example 8. Inducing a long-lasting immunity against *P.falciparum* malaria infection by prime-boost regimens with different adenovirus serotype-based vaccines.

An experiment to investigate the ability of prime/boost regimens based on adenovirus vectors derived from two different serotypes to induce long-lasting immunity against the *P.falciparum* CS antigen was designed. The study enrolls B10.BR mice distributed in experimental groups of 24 mice each. Animals are immunized by intra-muscular injection of an optimal dose of adenoviral vectors carrying either the full length CS gene (Ad5ΔE3-CS.Pfalc and Ad35ΔE3-CS.Pfalc) or the GPI-anchor sequence minus CS gene (Ad5ΔE3-CS.Pfalc(-28)/(-14) and Ad35ΔE3-CS.Pfalc(-28)/(-14)) or no transgene (Ad5ΔE3-empty and Ad35ΔE3-empty). One group of animals is primed at week 0 with Ad5ΔE3-CS.Pfalc or Ad5ΔE3-CS.Pfalc(-28)/(-14) and boosted at week 8 with Ad35ΔE3-CS.Pfalc or Ad35ΔE3-CS.Pfalc(-28)/(-14). Another group of mice is primed at week 0 with Ad35ΔE3-CS.Pfalc or Ad35ΔE3-CS.Pfalc(-28)/(-14) and boosted at week 8 with Ad5ΔE3-CS.Pfalc or Ad5ΔE3-CS.Pfalc(-28)/(-14). Another group of mice is primed at week 0 with Ad35ΔE3-CS.Pfalc or

Ad35ΔE3-CS.Pfalc(-28)/(-14) and boosted at week 8 with the same vector. Finally, a control group of mice is primed at week 0 with Ad5ΔE3-empty and boosted at week 8 with Ad35ΔE3-empty. At week 2 and 6 or 10 or 16 post-boost, 6 mice are sacrificed at each time point and cellular and humoral responses are monitored with immunological assays well known to persons skilled in the art and as described above. The immunization, challenge and read out are scheduled in Table VII. Vaccine regimens based on Ad35 alone or Ad5/Ad35 combinations are expected to be superior or at least comparable in efficacy as compared to regimens based solely on Ad5.

Example 9. Inducing an immune response against the *P.falciparum* CS antigen by prime/boost regimens using different Adenovirus serotype-based vaccines in non-human primates.

An example of an experiment useful to investigate the capability of prime/boost regimens based on adenovirus vectors derived from two different serotypes to elicit immunity against the *P.falciparum* CS antigen in non-human primates is described. Moreover, the effect of two different routes of vaccine administration, intra-muscular and intra-dermal, is evaluated.

Rhesus monkeys are vaccinated with adenoviral vectors carrying either the full-length CS gene (Ad5ΔE3-CS.Pfalc or Ad35ΔE3-CS.Pfalc) or the GPI-anchor sequence minus CS gene (Ad5ΔE3-CS.Pfalc(pf-aa-sub) or Ad35ΔE3-CS.Pfalc(pf-aa-sub)). Prime/boost regimens (Ad5 followed by Ad35 or Ad35 followed by Ad5) are compared to generally applied prime/boost regimens (Ad5 followed by Ad5 or Ad35 followed by Ad35). Humoral and cellular immune responses are monitored using immunological assays

well known to persons skilled in the art. Serum of immunized monkeys is tested by ELISA assay to determine the nature and magnitude of the antibody response against the repeat region of CS. Cellular immune response is
5 measured by ELISPOT assay to determine the amount of antigen-specific IFN- γ secreting cells.

Table I. Names and Genbank database entry numbers of the *P.falciparum* circumsporozoite amino acid sequences used to generate the final consensus sequence.

Wild type isolates	Entry numbers	Lab strains	Entry numbers
China	AAG37074	3D7	CAA33421
Thailand	CAB64171		CAB38998
	CSP_PLAFT		CSP_PLAFO
	AAA29542 - AAA29552		NP_473175
	AAA29555 - AAA29576	7G8	CSP_PLAFA
Brazil	CAB64167		C60657
	CAB64190 - CAB64197		AAA29524
Senegal	CAB64180 - CAB64189	NF54	AAA29521
Myanmar	CAB64237 - CAB64243		AAA29527
India	CAB64169		S05428
Tanzania	CAB64168		CSP_PLAFL
	CAB64170	WELLCOME	A54529
	CAB64172		AAA29554
Gambia	AAF03134 - AAF03136		CSP_PLAFW
	A38869		D60657
	B60657	LE5	CSP_PLAFL
	B38869		AAA57043
	H60657		B29765
Uganda	CAA27599		
	CAB64177		
Liberia	CAB64176		
Hondouras	CAB64174		
South East Asia	AAA29516 - AAA29519		
	CAB64175		
	CAB64178		
	CAB64179		

Table II. Immunization, challenge and read-out schedule for mice vaccinations with Ad5.CS.Pyoel (Ad5-PyCS), vp = viral particles per mouse.

Immunization schedule	Viral vector	vp	# mice	ELISPOT/serum	Challenge
Prime/challenge	Ad5-PyCS	10^8	12	2 weeks (6 mice)	2 weeks (6 mice)
Prime/challenge	Ad5-PyCS	10^9	12	2 weeks (6 mice)	2 weeks (6 mice)
Prime/challenge	Ad5-PyCS	10^{10}	12	2 weeks (6 mice)	2 weeks (6 mice)
Prime/challenge	Ad5-empty	10^{10}	8	2 weeks (4 mice)	2 weeks (4 mice)

5

Table III. Immunization, challenge and read-out schedule for mice vaccinations with Ad35.CS.Pyoel (Ad35-PyCS), vp = viral particles per mouse.

Immunization schedule	Viral vector	vp	# mice	ELISPOT/serum	Challenge
Prime/challenge	Ad35-PyCS	10^8	12	2 weeks (6 mice)	2 weeks (6 mice)
Prime/challenge	Ad35-PyCS	10^9	12	2 weeks (6 mice)	2 weeks (6 mice)
Prime/challenge	Ad35-PyCS	10^{10}	12	2 weeks (6 mice)	2 weeks (6 mice)
Prime/challenge	Ad35-empty	10^{10}	8	2 weeks (4 mice)	2 weeks (4 mice)

10

Table IV. Immunization, challenge and read-out schedule for mice vaccinations with Ad5.CS.Pfalc (Ad5-PfCS, with or without anchor), vp = viral particles per mouse.

Immunization schedule	Viral vector	vp	# mice	ELISPOT/serum	ELISPOT/serum
Prime	Ad5-PfCS	10^8	12	2 weeks (6 mice)	6-8 weeks (6 mice)
Prime	Ad5-PfCS	10^9	12	2 weeks (6 mice)	6-8 weeks (6 mice)
Prime	Ad5-PfCS	10^{10}	12	2 weeks (6 mice)	6-8 weeks (6 mice)
Prime	Ad5-empty	10^{10}	8	2 weeks (4 mice)	6-8 weeks (4 mice)

15

Table V. Immunization, challenge and read-out schedules for mice vaccinations with Ad35.CS.Pfalc (Ad35-PfCS, with or without anchor), vp = viral particles per mouse.

Immunization schedule	Viral vector	vp	# mice	ELISPOT/serum	ELISPOT/serum
Prime	Ad35-PfCS	10^8	12	2 weeks (6 mice)	6-8 weeks (6 mice)
Prime	Ad35-PfCS	10^9	12	2 weeks (6 mice)	6-8 weeks (6 mice)
Prime	Ad35-PfCS	10^{10}	12	2 weeks (6 mice)	6-8 weeks (6 mice)
Prime	Ad35-empty	10^{10}	8	2 weeks (4 mice)	6-8 weeks (4 mice)

20

Table VI. Immunization, challenge and read-out schedule for mice in a prime-boost vaccination set-up using Ad5.CS.Pyoel (Ad5-PyCS) and Ad35.CS.Pyoel (Ad35-PyCS).

Immunization schedule	Viral vector-prime	Viral vector-boost (after 8 weeks)	# mice	ELISPOT/serum	challenge
Prime-boost/challenge	Ad5-PyCS	Ad5-PyCS	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)
Prime-boost/challenge	Ad35-PyCS	Ad35-PyCS	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)
Prime-boost/challenge	Ad5-PyCS	Ad35-PyCS	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)
Prime-boost/challenge	Ad35-PyCS	Ad5-PyCS	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)
Prime-boost/challenge	Ad5- empty	Ad35- empty	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)

5

Table VII. Immunization, challenge and read-out schedule for mice in a prime-boost vaccination set-up using Ad5.CS.Pfalc (Ad5-PfCS) and Ad35.CS.Pfalc (Ad35-PfCS),
with or without GPI anchor.

10

Immunization schedule	Viral vector-prime	Viral vector-boost (after 8 weeks)	# mice	ELISPOT/serum	ELISPOT/serum
Prime-boost/challenge	Ad5-PfCS	Ad5-PfCS	12	2 wks post boost (6 mice)	6/10/16 wks post boost (6 mice)
Prime-boost/challenge	Ad35-PfCS	Ad35-PfCS	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)
Prime-boost/challenge	Ad5-PfCS	Ad35-PfCS	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)
Prime-boost/challenge	Ad35-PfCS	Ad5-PfCS	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)
Prime-boost/challenge	Ad5- empty	Ad35- empty	12	2 wks post boost (6 mice)	2 wks post boost (6 mice)

15

REFERENCES

- Bruña-Romero O, Gonzalez-Aseguinolaza G, Hafalla JCR, et al. (2001a) Complete, long-lasting protection against malaria of mice primed and boosted with two distinct viral vectors expressing the same plasmodial antigen. Proc Natl Acad Sci USA 98:11491-11496
- 5 Bruña-Romero O, Hafalla JC, Gonzalez-Aseguinolaza G, sano G, Tsjuji M, Zavala F. (2001b) Detection of malaria liver-stages in mice infected through the bite of a single Anopheles mosquito using a highly sensitive real-time PCR. Int J Parasitol 31:1499-1502
- 10 Clyde DF, Most H, McCarthy VC, Vanderberg JP. (1973) Immunization of men against sporozoite-induced falciparum malaria. Am J Med Sci 266:169-177
- 15 De Jong JC, Wermenbol AG, Verweij-Uijterwaal MW, Slaterus KW, Wertheim-Van Dillen P, Van Doornum GJ, Khoo SH, Hierholzer JC. (1999) Adenoviruses from human immunodeficiency virus-infected individuals, including two strains that represent new candidate serotypes Ad50 and Ad51 of species B1 and D, respectively. J Clin Microbiol 37:3940-3945
- 20 Gandon S, Mackinnon MJ, Nee S, Read AF. (2001) Imperfect vaccines and the evolution of pathogen virulence. Nature 414:751-756
- 30 Gilbert SC, Schneider J, Hannan CM, et al. (2002) Enhanced CD8 T cell immunogenicity and protective efficacy in a mouse malaria model using a recombinant adenoviral vaccine in heterologous prime-boost immunisation regimes. Vaccine 20:1039-1045
- 35 Gordon DM, McGovern TW, Krzych U, Cohen JC, Schneider I, LaChance R, Heppner DG, Yuan G, Hollingdale M, Slaoui M et al. (1995) Safety, immunogenicity, and efficacy of a recombinantly produced Plasmodium falciparum circumsporozoite protein-hepatitis B surface antigen subunit vaccine. J Infect Dis 171:1576-1585
- 40 Kurtis JD, Hollingdale MR, Luty AJF, Lanar DE, Krzych U and Duffy PE (2001) Pre-erythrocytic immunity to Plasmodium falciparum: the case for an LSA-1 vaccine. Trends in Parasitology 17:219-223
- 45 Lockyer MJ, Marsh K, Newbold CI. (1989) Wild isolates of Plasmodium falciparum show extensive polymorphism in T
- 50

cell epitopes of the circumsporozoite protein. Mol
Biochem Parasitol 37:275-280

- 5 Moran P and Caras IW. (1994) Requirements for
glycosylphosphatidylinositol attachment are similar but
not identical in mammalian cells and parasitic protozoa.
J Cell Biol 125:333-343
- 10 Nardin EH, Calvo-Calle JM, Oliveira GA, et al. (2001) A
totally synthetic polyoxime malaria vaccine containing
Plasmodium falciparum B cell and universal T cell
epitopes elicits immune responses in volunteers of
diverse HLA types. J Immunol 166:481-489
- 15 Narum DL, Kumar S, Rogers WO, et al. (2001) Codon
optimization of gene fragments encoding *Plasmodium*
falciparum merzoite proteins enhances DNA vaccine protein
expression and immunogenicity in mice. Infect and Immun
69:7250-7253
- 20 Nussenzweig RS, Vanderberg J, Most H, Orton C. (1967)
Protective immunity produced by the injection of X-
irradiated sporozoites of *Plasmodium berghei*. Nature
216:160-162
- 25 Romero P, Marayanski JL, Corradin G, et al. (1989) Cloned
cytotoxic T cells recognize an epitope in the
circumsporozoite protein and protect against malaria.
Nature 341:323-326
- 30 Rodrigues EG, Zavala F, Eichinger D, et al. (1997) Single
immunizing dose of recombinant adenovirus efficiently
induces CD8+ T cell-mediated protective immunity against
malaria. J Immunol 158:1268-1274
- 35 Scheiblhofer S, Chen D, Weiss R, et al. (2001) Removal of
the circumsporozoite protein (CSP)
glycosylphosphatidylinositol signal sequence from a CSP
DNA vaccine enhances induction of CSP-specific Th2 type
40 immune responses and improves protection against malaria
infection. Eur J Immunol 31:692-698
- 45 Zevering Y, Khamboonruang C, Good MF. (1994) Effect of
polymorphism of sporozoite antigens on T-cell activation.
Res Immunol 145:469-476

SEQUENCE LISTING

5 <110> Crucell Holland B.V.
 Crucell Holland B.V.
 Pau, Maria Grazia
 Stegmann, Antonius J.H.
 Kaspers, Jorn
 Holterman, Lennart

10 <120> Recombinant viral -based malaria vaccines
 <130> 0084W0000RD

15 <140> PCT/EP03/xxxxx
 <141> 2003-12-17
 <150> EP 02102781.8
 <151> 2002-12-17

20 <150> PCT/EP03/50222
 <151> 2003-06-12
 <160> 19

25 <170> PatentIn version 3.1
 <210> 1
 <211> 1193
 <212> DNA
 30 <213> Artificial

<220>
 <223> Description of Artificial Sequence: Codon -optimised circumsporozoite gene of Plasmodium falciparum, clone 02 -148

35 <220>
 <221> CDS
 <222> (13)..(1170)
 <223>

40 <220>
 <221> sig_peptide
 <222> (13)..(102)
 <223>

45 <400> 1
 aagcttgcca cc atg atg agg aaa ctg gcc atc ctg agc gtg agc agc ttc 51
 1 5 10
 Met Met Arg Lys Leu Ala Ile Leu Ser Val Ser Ser Phe

50 ctg ttc gtg gag gcc ctg ttt cag gag tac cag tgc tac ggc agc agc 99
 Leu Phe Val Glu Ala Leu Phe Gln Glu Tyr Gln Cys Tyr Gly Ser Ser
 15 20 25

55 agc aac acc cgg gtg ctg aac gag ctg aac tac gac aac gcc ggc acc 147
 Ser Asn Thr Arg Val Leu Asn Glu Leu Asn Tyr Asp Asn Ala Gly Thr
 30 35 40 45

60 aac ctg tac aac gag ctg gag atg aac tac tac ggc aag cag gag aac 195
 Asn Leu Tyr Asn Glu Leu Glu Met Asn Tyr Tyr Gly Lys Gln Glu Asn
 50 55 60

65 tgg tac agc ctg aag aag aac agc cgg tct ctg ggc gag aac gac gac 243
 Trp Tyr Ser Leu Lys Lys Asn Ser Arg Ser Leu Gly Glu Asn Asp Asp
 65 70 75

	ggc aac aac aac aac ggc gac aac ggc cgg gag ggc aag gac gag gac Gly Asn Asn Asn Asn Gly Asp Asn Gly Arg Glu Gly Lys Asp Glu Asp 80 85 90	291
5	aag cgg gac ggc aac aac gag gac aac gag aag ctg cgg aag ccc aag Lys Arg Asp Gly Asn Asn Glu Asp Asn Glu Lys Leu Arg Lys Pro Lys 95 100 105	339
10	cac aag aaa ctt aag cag ccc gcc gac ggc aac ccc gac ccc aac gcc His Lys Lys Leu Lys Gln Pro Ala Asp Gly Asn Pro Asp Pro Asn Ala 110 115 120 125	387
15	aac ccc aac gtg gac ccc aac gcc aat cct aat gtc gac ccc aat gcc Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn Val As p Pro Asn Ala 130 135 140	435
20	aat cgg aac gtt gat ccc aat ggg aat cct aac gct aac ccc aat gcc Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala 145 150 155	483
25	aac cca aat gcc aat cca aat gca aat ccc aac gcc aat cca aac gca Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala 160 165 170	531
30	aac cct aat gct aat cca aac gct aat cct aat gcc aat ccc aat gct Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala 175 180 185	579
35	aac cca aac gtc gat cct aac gca aat cgg aac gct aac ccc aac gca Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala 190 195 200 205	627
40	aat ccc aac gct aac cgg aac gca aac cct aac gcc aat cgg aat gcc Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala 210 215 220	675
45	aac cca aac gcc aac cgg aac gct aat cgg aat gct aac cgg aat gct Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala 225 230 235	723
50	aat cct aac gca aac cca aat gca aac ccc aat gca aac cgg aac gcc Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala 240 245 250	771
55	aat ccc aac gcc aat cct aat gcc aac aag aac aat cag ggc aac ggc Asn Pro Asn Ala Asn Pro Asn Ala Asn Lys Asn Asn Gln Gly Asn Gly 255 260 265	819
60	cag ggc cac aac atg ccc aac gac ccc aac cgg aac gtg gac gag aac Gln Gly His Asn Met Pro Asn Asp Pro Asn Arg Asn Val Asp Glu Asn 270 275 280 285	867
65	gcc aac gcc aac aac gcc gtg aag aac aac aac aac gag gag ccc agc Ala Asn Ala Asn Asn Ala Val Lys Asn Asn Asn Asn Glu Glu Pro Ser 290 295 300	915
70	gac aag cac atc gag cag tac ctg aag aag atc cag aac agc ctg agc Asp Lys His Ile Glu Gln Tyr Leu Lys Lys Ile Gln Asn Ser Leu Ser 305 310 315	963
75	acc gag tgg agc ccc tgc agc gtg acc tgc ggc aac ggc att cag gtg Thr Glu Trp Ser Pro Cys Ser Val Thr Cys Gly Asn Gly Ile Gln Val 320 325 330	1011
80	cgg atc aag ccc ggc agc gcc aac aag ccc aag gac gag ctg gac tac Arg Ile Lys Pro Gly Ser Ala Asn Lys Pro Lys Asp Glu Leu Asp Tyr 335 340 345	1059

gag aat gac atc gag aag aag atc tgc aag atg gag aag tgc agc agc 1107
 Glu Asn Asp Ile Glu Lys Lys Ile Cys Lys Met Glu Lys Cys Ser Ser
 350 355 360 365

5 gtg ttc aac gtg gtg aac agc gcc atc ggc ctg att atg gtg ctg agc 1155
 Val Phe Asn Val Val Asn Ser Ala Ile Gly Leu Ile Met Val Leu Ser
 370 375 380

10 ttc ctg ttc ctg aac tgaagatctg ctgataagga tcc 1193
 Phe Leu Phe Leu Asn
 385

15 <210> 2
 <211> 386
 <212> PRT
 <213> Artificial

20 <220>
 <223> Description of Artificial Sequence: Codon -optimised circumsporozoite gene of Plasmodium falciparum, clone 02 -148
 <400> 2

25 Met Met Arg Lys Leu Ala Ile Leu Ser Val Ser Ser Phe Leu Phe Val
 1 5 10 15

30 Glu Ala Leu Phe Gln Glu Tyr Gln Cys Tyr Gly Ser Ser Ser Asn Thr
 20 25 30

35 Arg Val Leu Asn Glu Leu Asn Tyr Asp Asn Ala Gly Thr Asn Leu Tyr
 35 40 45

40 Asn Glu Leu Glu Met Asn Tyr Tyr Gly Lys Gln Glu Asn Trp Tyr Ser
 50 55 60

45 Leu Lys Lys Asn Ser Arg Ser Leu Gly Glu Asn Asp Asp Gly Asn Asn
 65 70 75 80

50 Asn Asn Gly Asp Asn Gly Arg Glu Gly Lys Asp Glu Asp Lys Arg Asp
 85 90 95

55 Gly Asn Asn Glu Asp Asn Glu Lys Leu Arg Lys Pro Lys His Lys Lys
 100 105 110

60 Leu Lys Gln Pro Ala Asp Gly Asn Pro Asp Pro Asn Ala Asn Pro Asn
 115 120 125

65 Val Asp Pro Asn Ala Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn
 130 135 140

70 Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 145 150 155 160

75 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 165 170 175

Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
180 185 190

5 Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
195 200 205

10 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
210 215 220

15 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
225 230 235 240

20 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
245 250 255

Ala Asn Pro Asn Ala Asn Lys Asn Asn Gln Gly Asn Gly Gln Gly His
260 265 270

25 Asn Met Pro Asn Asp Pro Asn Arg Asn Val Asp Glu Asn Ala Asn Ala
275 280 285

30 Asn Asn Ala Val Lys Asn Asn Asn Asn Glu Glu Pro Ser Asp Lys His
290 295 300

35 Ile Glu Gln Tyr Leu Lys Lys Ile Gln Asn Ser Leu Ser Thr Glu Trp
305 310 315 320

Ser Pro Cys Ser Val Thr Cys Gly Asn Gly Ile Gln Val Arg Ile Lys
325 330 335

40 Pro Gly Ser Ala Asn Lys Pro Lys Asp Glu Leu Asp Tyr Glu Asn Asp
340 345 350

45 Ile Glu Lys Lys Ile Cys Lys Met Glu Lys Cys Ser Ser Val Phe Asn
355 360 365

50 Val Val Asn Ser Ala Ile Gly Leu Ile Met Val Leu Ser Phe Leu Phe
370 375 380

55 Leu Asn
385

<210> 3
<211> 386
<212> PRT
60 <213> Plasmodium falciparum

<220>
<221> SIGNAL
<222> (1)..(30)
65 <223>

<220>
<221> REPEAT

<222> (124)..(259)
 <223> Region of 4 NVDP and in total 30 NAVP repeats

5 <220>
 <221> LIPID
 <222> (359)..(386)
 <223> GPI-ANCHOR

10 <220>
 <221> MUTAGEN
 <222> (373)..(373)
 <223> Ser to Ala substitution, elimination of glycosylation site

15 <400> 3

20 Met Met Arg Lys Leu Ala Ile Leu Ser Val Ser Ser Phe Leu Phe Val
 1 5 10 15

Glu Ala Leu Phe Gln Glu Tyr Gln Cys Tyr Gly Ser Ser Ser Asn Thr
 20 25 30

25 Arg Val Leu Asn Glu Leu Asn Tyr Asp Asn Ala Gly Thr Asn Leu Tyr
 35 40 45

30 Asn Glu Leu Glu Met Asn Tyr Tyr Gly Lys Gln Glu Asn Trp Tyr Ser
 50 55 60

35 Leu Lys Lys Asn Ser Arg Ser Leu Gly Glu Asn Asp Asp Gly Asn Asn
 65 70 75 80

40 Asn Asn Gly Asp Asn Gly Arg Glu Gly Lys Asp Glu Asp Lys Arg Asp
 85 90 95

Gly Asn Asn Glu Asp Asn Glu Lys Leu Arg Lys Pro Lys His Lys Lys
 100 105 110

45 Leu Lys Gln Pro Ala Asp Gly Asn Pro Asp Pro Asn Ala Asn Pro Asn
 115 120 125

50 Val Asp Pro Asn Ala Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn
 130 135 140

55 Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 145 150 155 160

60 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 165 170 175

Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 180 185 190

65 Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 195 200 205

5 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 210 215 220
 10 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 225 230 235 240
 15 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
 245 250 255
 20 Ala Asn Pro Asn Ala Asn Lys Asn Asn Gln Gly Asn Gly Gln Gly His
 260 265 270
 25 Asn Met Pro Asn Asp Pro Asn Arg Asn Val Asp Glu Asn Ala Asn Ala
 275 280 285
 Asn Asn Ala Val Lys Asn Asn Asn Asn Glu Glu Pro Ser Asp Lys His
 290 295 300
 30 Ile Glu Gln Tyr Leu Lys Lys Ile Gln Asn Ser Leu Ser Thr Glu Trp
 305 310 315 320
 35 Ser Pro Cys Ser Val Thr Cys Gly Asn Gly Ile Gln Val Arg Ile Lys
 325 330 335
 40 Pro Gly Ser Ala Asn Lys Pro Lys Asp Glu Leu Asp Tyr Glu Asn Asp
 340 345 350
 Ile Glu Lys Lys Ile Cys Lys Met Glu Lys Cys Ser Ser Val Phe Asn
 355 360 365
 45 Val Val Asn Ser Ala Ile Gly Leu Ile Met Val Leu Ser Phe Leu Phe
 370 375 380
 Leu Asn
 385
 50 <210> 4
 <211> 1154
 <212> DNA
 <213> Artificial
 55 <220>
 <223> Description of Artificial Sequence: Codon -optimised circumsporozo-
 ite gene of Plasmodium falciparum strain 3D7, clone 02-659
 60 <220>
 <221> CDS
 <222> (13)..(1128)
 <223>
 65 <220>
 <221> sig_peptide
 <222> (13)..(102)
 <223>

<400> 4
aagcttgcca cc atg atg agg aaa ctg gcc atc ctg agc gtg agc agc ttc 51
Met Met Arg Lys Leu Ala Ile Leu Ser Val Ser Ser Phe
5 1 5 10

ctg ttc gtg gag gcc ctg ttt cag gag tac cag tgc tac ggc agc agc 99
Leu Phe Val Glu Ala Leu Phe Gln Glu Tyr Gln Cys Tyr Gly Ser Ser
15 20 25

10 agc aac acc cgg gtg ctg aac gag ctg aac tac gac aac gcc ggc acc 147
Ser Asn Thr Arg Val Leu Asn Glu Leu Asn Tyr Asp Asn Ala Gly Thr
30 35 40 45

15 aac ctg tac aac gag ctg gag atg aac tac tac ggc aag cag gag aac 195
Asn Leu Tyr Asn Glu Leu Glu Met Asn Tyr Tyr Gly Lys Gln Glu Asn
50 55 60

20 tgg tac agc ctg aag aa g aac agc cgg tct ctg ggc gag aac gac gac 243
Trp Tyr Ser Leu Lys Lys Asn Ser Arg Ser Leu Gly Glu Asn Asp Asp
65 70 75

25 ggc aac aac aac aac ggc gac aac ggc cgg gag ggc aag gac ga g gac 291
Gly Asn Asn Asn Asn Gly Asp Asn Gly Arg Glu Gly Lys Asp Glu Asp
80 85 90

aag cgg gac ggc aac aac gag gac aac gag aag ctg cgg aag ccc aag 339
Lys Arg Asp Gly Asn Asn Glu Asp Asn Glu Lys Leu Arg Lys Pro Lys
95 100 105

30 cac aag aaa ctt aag cag ccc gcc gac ggc aac ccc gac ccc aac gcc 387
His Lys Lys Leu Lys Gln Pro Ala Asp Gly Asn Pro Asp Pro A sn Ala
110 115 120 125

35 aac ccc aac gtg gac ccc aac gcc aat cct aat gtc gac ccc aat gcc 435
Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn Val Asp Pro Asn Ala
130 135 140

40 aat cgg aac gtt gat ccc aat gcg aat cct aac gct aac ccc aat gcc 483
Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
145 150 155

45 aac cca aat gcc aat cca aat gca aat ccc aac gcc aat cca aac gca 531
Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
160 165 170

aac cct aat gct aa t cca aac gct aat cct aat gcc aat ccc aat gct 579
Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
175 180 185

50 aac cca aac gtc gat cct aac gca aat cgg aac gct aac cc c aac gca 627
Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
190 195 200 205

55 aat ccc aac gct aac cgg aac gca aac cct aac gcc aat cgg aat gcc 675
Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
210 215 220

60 aac cca aac gcc aac cgg aac gct aat cgg aat gct aac cgg aat gct 723
Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn P ro Asn Ala
225 230 235

aat cct aac gca aac cca aat gca aac ccc aat gca aac cgg aac gcc 771
Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
65 240 245 250

aat ccc aac gcc aat cct aat gcc aac aag aac aat cag gcc aac gcc 819
 Asn Pro Asn Ala Asn Pro Asn Ala Asn Lys Asn Asn Gln Gly Asn Gly
 255 260 265

5 cag gcc cac aac atg ccc aac gac ccc aac cgg aac gtg gac gag aac 867
 Gln Gly His Asn Met Pro Asn Asp Pro Asn Arg Asn Val Asp Glu Asn
 270 275 280 285

10 gcc aac gcc aac agc gcc gtg aag aac aac aac aac gag gag ccc agc 915
 Ala Asn Ala Asn Ser Ala Val Lys Asn Asn Asn Asn Glu Glu Pro Ser
 290 295 300

15 gac aag cac atc aag gag tac ctg aac aag atc cag aac agc ctg agc 963
 Asp Lys His Ile Lys Glu Tyr Leu Asn Lys Ile Gln Asn Ser Leu Ser
 305 310 315

20 acc gag tgg agc ccc tgc agc gtg acc tgc gcc aac gcc att cag gtg 1011
 Thr Glu Trp Ser Pro Cys Ser Val Thr Cys Gly Asn Gly Ile Gln Val
 320 325 330

cgg atc aag ccc gcc agc gcc aac aag ccc aag gac gag ctg gac tac 1059
 Arg Ile Lys Pro Gly Ser Ala Asn Lys Pro Lys Asp Glu Leu Asp Tyr
 335 340 345

25 gcc aat gac atc gag aag aag atc tgc aag atg gag aag tgc agc agc 1107
 Ala Asn Asp Ile Glu Lys Lys Ile Cys Lys Met Glu Lys Cys Ser Ser
 350 355 360 365

30 gtg ttc aac gtg gtg aac tcc tgataaagat ctgctgataa ggatcc 1154
 Val Phe Asn Val Val Asn Ser
 370

35 <210> 5
 <211> 372
 <212> PRT
 <213> Artificial

40 <220>
 <223> Description of Artificial Sequence: Codon -optimised circumsporozo-
 ite gene of Plasmodium falciparum strain 3D7, clone 02 -659
 <400> 5

45 Met Met Arg Lys Leu Ala Ile Leu Ser Val Ser Ser Phe Leu Phe Val
 1 5 10 15

50 Glu Ala Leu Phe Gln Glu Tyr Gln Cys Tyr Gly Ser Ser Ser Asn Thr
 20 25 30

55 Arg Val Leu Asn Glu Leu Asn Tyr Asp Asn Ala Gly Thr Asn Leu Tyr
 35 40 45

Asn Glu Leu Glu Met Asn Tyr Tyr Gly Lys Gln Glu Asn Trp Tyr Ser
 50 55 60

60 Leu Lys Lys Asn Ser Arg Ser Leu Gly Glu Asn Asp Asp Gly Asn Asn
 65 70 75 80

65 Asn Asn Gly Asp Asn Gly Arg Glu Gly Lys Asp Glu Asp Lys Arg Asp
 85 90 95

	Gly	Asn	Asn	Glu	Asp	Asn	Glu	Lys	Leu	Arg	Lys	Pro	Lys	His	Lys	Lys	
				100					105					110			
5	Leu	Lys	Gln	Pro	Ala	Asp	Gly	Asn	Pro	Asp	Pro	Asn	Ala	Asn	Pro	Asn	
			115					120					125				
10	Val	Asp	Pro	Asn	Ala	Asn	Pro	Asn	Val	Asp	Pro	Asn	Ala	Asn	Pro	Asn	
			130				135					140					
15	Val	Asp	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	
			145			150				155						160	
20	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	
				165					170					175			
25	Val	Asp	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	
			195				200					205					
30	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	
			210				215					220					
35	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	
			225			230				235					240		
40	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	
				245					250					255			
45	Asn	Met	Pro	Asn	Asp	Pro	Asn	Arg	Asn	Val	Asp	Glu	Asn	Ala	Asn	Ala	
			275				280						285				
50	Asn	Ser	Ala	Val	Lys	Asn	Asn	Asn	Asn	Glu	Glu	Pro	Ser	Asp	Lys	His	
			290				295					300					
55	Ile	Lys	Glu	Tyr	Leu	Asn	Lys	Ile	Gln	Asn	Ser	Leu	Ser	Thr	Glu	Trp	
		305				310					315					320	
60	Ser	Pro	Cys	Ser	Val	Thr	Cys	Gly	Asn	Gly	Ile	Gln	Val	Arg	Ile	Lys	
				325						330					335		
65	Pro	Gly	Ser	Ala	Asn	Lys	Pro	Lys	Asp	Glu	Leu	Asp	Tyr	Ala	Asn	Asp	
				340					345					350			
	Ile	Glu	Lys	Lys	Ile	Cys	Lys	Met	Glu	Lys	Cys	Ser	Ser	Val	Phe	Asn	
			355					360						365			

Val Val Asn Ser
370

5 <210> 6
<211> 372
<212> PRT
<213> Plasmodium falciparum

10 <220>
<221> SIGNAL
<222> (1)..(30)
<223>

15 <220>
<221> REPEAT
<222> (124)..(259)
<223> Region of 4 NVDF and in total 30 NAVP repeats

20 <220>
<221> LIPID
<222> (359)..(372)
<223> GPI-ANCHOR

25 <400> 6

30 Met Met Arg Lys Leu Ala Ile Leu Ser Val Ser Ser Phe Leu Phe Val
1 5 10 15

Glu Ala Leu Phe Gln Glu Tyr Gln Cys Tyr Gly Ser Ser Ser Asn Thr
20 25 30

35 Arg Val Leu Asn Glu Leu Asn Tyr Asp Asn Ala Gly Thr Asn Leu Tyr
35 40 45

40 Asn Glu Leu Glu Met Asn Tyr Tyr Gly Lys Gln Glu Asn Trp Tyr Ser
50 55 60

45 Leu Lys Lys Asn Ser Arg Ser Leu Gly Glu Asn Asp Asp Gly Asn Asn
65 70 75 80

50 Asn Asn Gly Asp Asn Gly Arg Glu Gly Lys Asp Glu Asp Lys Arg Asp
85 90 95

Gly Asn Asn Glu Asp Asn Glu Lys Leu Arg Lys Pro Lys His Lys Lys
100 105 110

55 Leu Lys Gln Pro Ala Asp Gly Asn Pro Asp Pro Asn Ala Asn Pro Asn
115 120 125

60 Val Asp Pro Asn Ala Asn Pro Asn Val Asp Pro Asn Ala Asn Pro Asn
130 135 140

65 Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
145 150 155 160

Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
165 170 175

5 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
180 185 190

10 Val Asp Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
195 200 205

15 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
210 215 220

Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
225 230 235 240

20 Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
245 250 255

25 Ala Asn Pro Asn Ala Asn Lys Asn Asn Gln Gly Asn Gly Gln Gly His
260 265 270

30 Asn Met Pro Asn Asp Pro Asn Arg Asn Val Asp Glu Asn Ala Asn Ala
275 280 285

35 Asn Ser Ala Val Lys Asn Asn Asn Asn Glu Glu Pro Ser Asp Lys His
290 295 300

Ile Lys Glu Tyr Leu Asn Lys Ile Gln Asn Ser Leu Ser Thr Gl u Trp
305 310 315 320

40 Ser Pro Cys Ser Val Thr Cys Gly Asn Gly Ile Gln Val Arg Ile Lys
325 330 335

45 Pro Gly Ser Ala Asn Lys Pro Lys Asp Glu Leu Asp Ty r Ala Asn Asp
340 345 350

50 Ile Glu Lys Lys Ile Cys Lys Met Glu Lys Cys Ser Ser Val Phe Asn
355 360 365

55 Val Val Asn Ser
370

60 <210> 7
<211> 1115
<212> DNA
<213> Artificial

65 <220>
<223> Description of Artificial Sequence: Codon -optimised cirrcumsporoz
oite gene of Plasmodium yoelii, clone 02 -149

<220>
<221> CDS
<222> (17) .. (1084)

cag cag ccc ccc cag caa cct ccc caa cag cct ccc cag caa cca cca 772
 Gln Gln Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro
 240 245 250

5 cag cag cca cca cag caa ccc agg ccc cag ccc gac ggc aac aac aac 820
 Gln Gln Pro Pro Gln Gln Pro Arg Pro Gln Pro Asp Gly Asn Asn Asn
 255 260 265

10 aac aat aac aac aac ggc aac aac aac gag gac agc tac gtg ccc agc 868
 Asn Asn Asn Asn Asn Gly Asn Asn Asn Glu Asp Ser Tyr Val Pro Ser
 270 275 280

15 gcc gag cag atc ctg gag ttc gtg aag cag atc agc agc caa ctg acc 916
 Ala Glu Gln Ile Leu Glu Phe Val Lys Gln Ile Ser Ser Gln Leu Thr
 285 290 295 300

20 gag gag tgg agc cag tgc agc gtg acc tgc ggc agc ggc gtg cgg gtg 964
 Glu Glu Trp Ser Gln Cys Ser Val Thr Cys Gly Ser Gly Val Arg Val
 305 310 315

cgg aag cgg aag aac gtg aac aag cag ccc gag aac ctg acc ctg gag 1012
 Arg Lys Arg Lys Asn Val Asn Lys Gln Pro Glu Asn Leu Thr Leu Glu
 320 325 330

25 gac atc gac acc gag atc tgc aag atg gac aag tgc agc agc atc ttc 1060
 Asp Ile Asp Thr Glu Ile Cys Lys Met Asp Lys Cys Ser Ser Ile Phe
 335 340 345

30 aac atc gtg agc aac agc ctg ggc tgaagatctg ctgataagtt taaacggatc 1114
 Asn Ile Val Ser Asn Ser Leu Gly
 350 355

c 1115

35 <210> 8
 <211> 356
 <212> PRT
 <213> Artificial

40 <220>
 <223> Description of Artificial Sequence: Codon -optimised cirrcumsporoz
 oite gene of Plasmodium yoelii, clone 02 -149

45 <400> 8
 Met Lys Lys Cys Thr Ile Leu Val Val Ala Ser Leu Leu Leu Val Asp
 1 5 10 15

50 Ser Leu Leu Pro Gly Tyr Gly Gln Asn Lys Ser Val Gln Ala Gln Arg
 20 25 30

55 Asn Leu Asn Glu Leu Cys Tyr Asn Glu Glu Asn Asp Asn Lys Leu Tyr
 35 40 45

60 His Val Leu Asn Ser Lys Asn Gly Lys Ile Tyr Asn Arg Asn Ile Val
 50 55 60

65 Asn Arg Leu Leu Gly Asp Ala Leu Asn Gly Lys Pro Glu Glu Lys Lys
 65 70 75 80

Asp Asp Pro Pro Lys Asp Gly Asn Lys Asp Asp Leu Pro Lys Glu Glu
 85 90 95

5 Lys Lys Asp Asp Leu Pro Lys Glu Glu Lys Lys Asp Asp Pro Pro Lys
 100 105 110
 Asp Pro Lys Lys Asp Asp Pro Pro Lys Glu Ala Gln Asn Lys Leu Asn
 115 120 125
 10 Gln Pro Val Val Ala Asp Glu Asn Val Asp Gln Gly Pro Gly Ala Pro
 130 135 140
 15 Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly
 145 150 155 160
 20 Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly
 165 170 175
 25 Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro
 180 185 190
 Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly
 195 200 205
 30 Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly
 210 215 220
 35 Pro Gly Ala Pro Gln Glu Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro
 225 230 235 240
 40 Gln Gln Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro
 245 250 255
 45 Gln Gln Pro Arg Pro Gln Pro Asp Gly Asn Asn Asn Asn Asn Asn Asn
 260 265 270
 Asn Gly Asn Asn Asn Glu Asp Ser Tyr Val Pro Ser Ala Glu Gln Ile
 275 280 285
 50 Leu Glu Phe Val Lys Gln Ile Ser Ser Gln Leu Thr Glu Glu Trp Ser
 290 295 300
 55 Gln Cys Ser Val Thr Cys Gly Ser Gly Val Arg Val Arg Lys Arg Lys
 305 310 315 320
 60 Asn Val Asn Lys Gln Pro Glu Asn Leu Thr Leu Glu Asp Ile Asp Thr
 325 330 335
 65 Glu Ile Cys Lys Met Asp Lys Cys Ser Ser Ile Phe Asn Ile Val Ser
 340 345 350
 Asn Ser Leu Gly
 355

5 <210> 9
 <211> 356
 <212> PRT
 <213> Plasmodium yoelii

 10 <220>
 <221> SIGNAL
 <222> (1) .. (19)
 <223>

 15 <220>
 <221> REPEAT
 <222> (138) .. (225)
 <223> Region of 15 QGPGAP repeats

 20 <220>
 <221> REPEAT
 <222> (229) .. (254)
 <223> Region of 7 PQQP repeats

 25 <400> 9
 Met Lys Lys Cys Thr Ile Leu Val Val Ala Ser Leu Leu Leu Val Asp
 1 5 10 15

 30 Ser Leu Leu Pro Gly Tyr Gly Gln Asn Lys Ser Val Gln Ala Gln Arg
 20 25 30

 35 Asn Leu Asn Glu Leu Cys Tyr Asn Glu Glu Asn Asp Asn Lys Leu Tyr
 35 40 45

 40 His Val Leu Asn Ser Lys Asn Gly Lys Ile Tyr Asn Arg Asn Ile Val
 50 55 60

 45 Asn Arg Leu Leu Gly Asp Ala Leu Asn Gly Lys Pro Glu Glu Lys Lys
 65 70 75 80

 50 Asp Asp Pro Pro Lys Asp Gly Asn Lys Asp Asp Leu Pro Lys Glu Glu
 85 90 95

 55 Lys Lys Asp Asp Leu Pro Lys Glu Glu Lys Lys Asp Asp Pro Pro Lys
 100 105 110

 60 Asp Pro Lys Lys Asp Asp Pro Pro Lys Glu Ala Gln Asn Lys Leu Asn
 115 120 125

 65 Gln Pro Val Val Ala Asp Glu Asn Val Asp Gln Gly Pro Gly Ala Pro
 130 135 140

 Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly
 145 150 155 160

 Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly
 165 170 175

5 Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro
 180 185 190
 Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly
 195 200 205
 10 Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly
 210 215 220
 15 Pro Gly Ala Pro Gln Glu Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro
 225 230 235 240
 20 Gln Gln Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro Gln Gln Pro Pro
 245 250 255
 25 Gln Gln Pro Arg Pro Gln Pro Asp Gly Asn Asn Asn Asn Asn Asn Asn
 260 265 270
 Asn Gly Asn Asn Asn Glu Asp Ser Tyr Val Pro Ser Ala Glu Gln Ile
 275 280 285
 30 Leu Glu Phe Val Lys Gln Ile Ser Ser Gln Leu Thr Glu Glu Trp Ser
 290 295 300
 35 Gln Cys Ser Val Thr Cys Gly Ser Gly Val Arg Val Arg Lys Arg Lys
 305 310 315 320
 40 Asn Val Asn Lys Gln Pro Glu Asn Leu Thr Leu Glu Asp Ile Asp Thr
 325 330 335
 45 Glu Ile Cys Lys Met Asp Lys Cys Ser Ser Ile Phe Asn Ile Val Ser
 340 345 350
 Asn Ser Leu Gly
 355
 50 <210> 10
 <211> 4
 <212> PRT
 <213> Plasmodium falciparum
 55 <400> 10
 Asn Ala Asn Pro
 1
 60 <210> 11
 <211> 4
 <212> PRT
 65 <213> Plasmodium falciparum
 <400> 11

Asn Val Asp Pro
1

5 <210> 12
<211> 11
<212> PRT
<213> Plasmodium falciparum

10 <400> 12
Glu Asn Ala Asn Ala Asn Asn Ala Val Lys Asn
1 5 10

15 <210> 13
<211> 23
<212> DNA
<213> Artificial

20 <220>
<223> Description of Artificial Sequence: oligonucleotide Forw.Falc

25 <400> 13
ccaagcttgc caccatga tg agg 23

30 <210> 14
<211> 27
<212> DNA
<213> Artificial

35 <220>
<223> Description of Artificial Sequence: oligonucleotide Rev.Falc.CS -2
8

40 <400> 14
ccggatcctc agcagatctt cttctcg 27

45 <210> 15
<211> 27
<212> DNA
<213> Artificial

50 <220>
<223> Description of Artificial Sequence: oligonucleotide Rev.Falc.CS -1
4

55 <400> 15
ccggatcctc agctgttcac cacgttg 27

60 <210> 16
<211> 22
<212> DNA
<213> Artificial

65 <220>
<223> Description of Artificial Sequence: oligonucleotide SV40for

<400> 16
caatgtatct tatcatgtot ag 22

65 <210> 17
<211> 35
<212> DNA

<213> Artificial

<220>
<223> Description of Artificial Sequence: oligonucleotide pIXRmfe

5 <400> 17
ctctctcaat tgcagataca aaactacata agacc 35

10 <210> 18
<211> 32
<212> DNA
<213> Artificial

15 <220>
<223> Description of Artificial Sequence: oli gonucleotide pIX35mfe

20 <400> 18
ctctctcaat tgtctgtctt gcagctgtca tg 32

25 <210> 19
<211> 18
<212> PRT
<213> Plasmodium yoelii

<400> 19
Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly Ala Pro Gln Gly Pro Gly
30 1 5 10 15

Ala Pro

35

Claims

1. A replication-defective recombinant adenovirus derived from a serotype selected from the group consisting of: Ad11, Ad26, Ad34, Ad35, Ad48, Ad49 and Ad50, wherein said recombinant adenovirus comprises a heterologous nucleic acid encoding an antigenic determinant of *Plasmodium falciparum*.
2. A replication-defective recombinant adenovirus according to claim 1, wherein said antigenic determinant is the circumsporozoite protein, or an immunogenic part thereof.
3. A replication-defective recombinant adenovirus according to claim 1 or 2, wherein said heterologous nucleic acid is codon-optimized for elevated expression in a mammal.
4. A replication-defective recombinant adenovirus according to claim 3, wherein said mammal is a human.
5. A replication-defective recombinant adenovirus according to claim 3 or 4, wherein the adenine plus thymine content in said heterologous nucleic acid, as compared to the cytosine plus guanine content, is less than 87%, preferably less than 80%, more preferably less than 59% and most preferably equal to approximately 45%.
6. A replication-defective recombinant adenovirus according to claim 2, wherein said circumsporozoite

protein is the circumsporozoite protein as depicted in figure 1A (SEQ ID NO:3).

7. A replication-defective recombinant adenovirus according to claim 3, wherein said codon-optimized heterologous nucleic acid is the nucleic acid as depicted in figure 1B (SEQ ID NO:1).

8. A replication-defective recombinant adenovirus according to any one of claims 2 to 7, wherein the circumsporozoite protein, or the immunogenic part thereof, is lacking a functional GPI anchor sequence.

9. A replication-defective recombinant adenovirus according to claim 1, wherein said antigenic determinant is selected from the group consisting of: LSA-1, LSA-3, MSP-1, MSP-119 and MSP-142, or combinations thereof.

10. A replication-defective recombinant adenovirus according to any one of claims 1 to 9, wherein said adenovirus further comprises a gene encoding a liver specific antigen for *Plasmodium falciparum* as antigenic determinant, or an immunogenic part thereof.

11. A replication-defective recombinant adenovirus according to claim 13, wherein said liver specific antigen is LSA-1.

12. A vaccine composition comprising a replication-defective recombinant adenovirus according to any one of claims 1 to 11, and a pharmaceutically acceptable carrier.
13. A vaccine composition according to claim 12, further comprising an adjuvant.
14. Use of a vaccine composition according to claim 12 and a vaccine composition comprising at least one purified malaria-derived protein or peptide in the manufacture of a medicament for therapeutic and/or prophylactic treatment of a malaria infection in a mammal.
15. Use of a vaccine composition according to claim 13 and a vaccine composition comprising at least one purified malaria-derived protein or peptide in the manufacture of a medicament for therapeutic and/or prophylactic treatment of a malaria infection in a mammal.
16. Use of a vaccine composition comprising a replication-defective recombinant adenovirus according to any one of claims 1 to 8 and a vaccine composition comprising a replication-defective recombinant adenovirus according to any one of claims 9 to 11 in the manufacture of a medicament for therapeutic and/or prophylactic treatment of a malaria infection in a mammal.