

(51) International Patent Classification:
C07D 215/20 (2006.01)(21) International Application Number:
PCT/SG2010/000188(22) International Filing Date:
17 May 2010 (17.05.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/178,722 15 May 2009 (15.05.2009) US(71) Applicant (for all designated States except US):
NANYANG TECHNOLOGICAL UNIVERSITY [SG/
SG]; 50 Nanyang Avenue, Singapore 639798 (SG).


(72) Inventors; and

(75) Inventors/Applicants (for US only): LOH, Teck Peng
[MY/SG]; c/o Nanyang Technological University, 50
Nanyang Avenue, Singapore 639798 (SG). XIAO, Jian
[CN/SG]; c/o Nanyang Technological University, 50
Nanyang Avenue, Singapore 639798 (SG).(74) Agent: GOH, Su Lin, Audrey; Viering, Jentschura &
Partner LLP, P.O. Box 1088, Rochor Post Office, Rochor
Road, Singapore 911833 (SG).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,
ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: OCTAHYDRO BIQUINOLINE COMPOUND



(57) Abstract: The present invention relates to an octahydro biquinoline compound. Provided is also a method of separating the octahydro biquinoline compound into enantiomers. The octahydro biquinoline compound is of the general formula (V): In formula (V) R^1 is one of H, a protective group and an aliphatic group, with the aliphatic group having a main chain of a length of 1 to about 10 carbon atoms, comprising 0 to about 6 heteroatoms selected from the group consisting of N, O, S, Se and Si. R^2 and R^3 are independent from one another selected from the group consisting of (i) H, (ii) one of an aliphatic, an alicyclic, an aromatic, an arylaliphatic, and an arylalicyclic group comprising 0 to about 6 heteroatoms selected from the group consisting of N, O, S, Se and Si, an ester, a carbonate group, a carbamoyl group and a phosphate ester. R^4 and R^5 are independent from one another H, an aliphatic, an alicyclic, an aromatic, an arylaliphatic or an arylalicyclic group comprising 0 to about 6 heteroatoms selected from the group consisting of N, O, S, Se and Si.