

PYRROLIDIN-2-ONES AS HDM2 LIGANDS

BACKGROUND OF THE INVENTION

5 MDM2 (also known as HDM2) plays a central role in regulating and influencing important cell-signalling pathways. HDM2 is known to interact with a range of different proteins that control cell cycle progression, cellular apoptosis, proliferation and survival.

10 Thus, amongst other proteins, HDM2 binds to the tumor suppressor protein p53 and targets this protein for ubiquitination and degradation; facilitate translocation of p53 from the nucleus to cytosole and further translocation to the proteosomes. Thereby, HDM2 prevents transactivation of p53 target genes that are implicated in the regulation of cell cycle and apoptosis. The p53 protein is a potent cell cycle inhibitor that prevents
15 propagation of permanently damaged cell clones by the induction of growth arrest or apoptosis, resulting in the protection against development of cancer by guarding cellular and genomic integrity.

Both p53 as well as HDM2 can be associated with cancer: about 50% of all human
20 tumors harbor a mutation or deletion in the p53 gene that impairs normal p53 function. In many cancers with wild-type p53, HDM2 is overexpressed, disabling the normal p53 function (Momand et al. *Nucleic Acids Res.* **1998**, 26, 3453-3459).

The HDM2 gene has a p53-responsive promoter element and elevated levels of p53 that
25 translocate to the nucleus induce expression of HDM2. Induction of HDM2 by p53 forms an autoregulatory feedback loop, ensuring low levels of both HDM2 and p53 in normally proliferating cells (Vousden and Lu *Nature Reviews Cancer* **2002**, 2, 594-604). However, in many cancers this normal ratio of HDM2 to p53 is changed and misregulated.

30

Inhibiting the interaction of HDM2 with p53 in cells with wild-type p53 should lead to an increase of p53 levels in the nucleus, facilitating cell cycle arrest and/or apoptosis

- 2 -

and restoring the tumor suppressor role of p53. The feasibility of this strategy has been shown by the use of different macromolecular tools for inhibition of HDM2-p53 interaction (e.g. antibodies, antisense oligonucleotides, peptides).

- 5 Besides p53, a number of proteins have been found to interact with HDM2, performing either effectors (regulating HDM2 functions) or effectors (regulated by HDM2). Totally about 20 interacting with HDM2 proteins have been described (Ganguli and Wasylyk, *Mol. Cancer Research*, 2003, v.1, 1027-1035), Zhu et al. *Mol. Cell*, 2009, 35, 316-326). Among them, HDM2 binds to the tumor suppressor pRB, as well as E2F-1 (Yang et al. *Clinical Cancer Research* 1999, 5, 2242-2250).
- 10

E2F-1 is a transcription factor that regulates S phase entry and has been shown to cause apoptosis in some cell types when overexpressed. HDM2 binds to E2F through a conserved binding region at p53, activating E2F-dependent transcription of cyclin A, and suggesting that HDM2 small molecule ligands or antagonists might have also anti-tumor effects in cells independent of their role of restoring p53 function.

15

HDM2 can associate in vitro and in vivo with the mammalian Numb protein. The association occurs through the N-terminal domain of HDM2, which is the region also involved in p53 binding. The Numb protein is involved in the regulation of cell fate and in a variety of developmental processes, most notably in the nervous system. Through its interaction with Numb, HDM2 may influence processes such as differentiation and survival. This could also contribute to the altered properties of tumor cells, which overexpress HDM2 (Juven-Gershon et al. *Mol. Cell. Biol.* 1998, 18, 3974-3982).

20

Similarly, small molecules that block the HDM2 interaction with p53 also block the interaction of HDM2 with hypoxia inducible factor 1 α (HIF-1 α), a protein that induces vascular endothelial growth factor (VEGF) under normoxic or hypoxic conditions. As VEGF is proangiogenic, inhibition of HDM2 by small molecules will also prevent blood vessel formation to cancer metastases and primary tumors (G.A. LaRusch et al. *Cancer Res.* 2007, 67, 450-454).

25

30

- 3 -

There is also evidence that HDM2 has a direct role in the regulation of p21, a cyclin-dependent kinase inhibitor. The inhibition of HDM2 with anti-HDM2 antisense oligonucleotide or Short Interference RNA targeting HDM2 significantly elevates p21 protein levels in p53 null PC3 cells. In contrast, overexpression of HDM2 diminishes p21 levels by shortening the p21 half-life, an effect reversed by HDM2 antisense inhibition. HDM2 facilitates p21 degradation independent of ubiquitination and the E3 ligase function of HDM2. Instead, HDM2 promotes p21 degradation by facilitating binding of p21 with the proteasomal C8 subunit. The p21 and HDM2 bind through 180–the 298 amino acids region of the HDM2 protein (Zhang et al. *J. Biol. Chem.* **2004**, 279, 16000–16006).

There is also evidence for a malfunctioning HDM2 regulation having effect on a proper p53 function and causing cancer, beyond mutated p53 or overexpression of HDM2. Thus, when E2F signals the growth of a cancer, P14ARF is dispatched to break down HDM2, freeing p53 to kill the cancer cell. In certain cancers P14ARF is lacking (Moule et al. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 14063-6). P14ARF binds to HDM2 and promotes the rapid degradation of HDM2. ARF-mediated HDM2 degradation is associated with HDM2 modification and concurrent p53 stabilization and accumulation.

Small molecule HDM2 inhibitors also induce senescence dependent on the presence of functional p53, whereas cells lacking p53 were completely insensitive (A. Efeyan, A. Ortega-Molina, S. Velasco-Miguel, D. Herranz, L.T. Vassilev, M. Serrano, *Cancer Res.* **2007**, *67*, 7350-7357).

The validity of inhibiting HDM2 as a therapeutic concept has been first demonstrated by antisense HDM2 inhibitors that exhibit significant antitumor activity in multiple human cancer models with various p53 statuses (Zhang et al. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 11636–11641).

Small molecule antagonists of the HDM2 protein interactions may therefore offer a viable approach towards cancer therapy, either as single agents or in combination with a broad variety of other anti-tumour therapies.

There is also growing evidence that HDM2 plays an important role in viral infections. First, it is known that viruses rely on changing normal p53 signalling (O'shea and Fried M. *Cell Cycle* **2005**; Machida et al. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *23*, 101, 4262-7).

Second, HDM2 directly interacts with viral proteins, for example HDM2 is a target of simian virus 40 in cellular transformation and during lytic infection (Henning et al. *J. Virol.* **1997**, *71*, 7609–7618). Furthermore, the HDM2 protein, like p53, becomes metabolically stabilized in SV40-transformed cells. This suggests the possibility that the specific targeting of HDM2 by SV40 is aimed at preventing HDM2-directed proteasomal degradation of p53 in SV40-infected and -transformed cells, thereby leading to metabolic stabilization of p53 in these cells. A trimeric LT-p53-HDM2 complex is formed with simian virus 40 large tumour antigen (LT) in SV40-transformed cells.

The human immunodeficiency virus type 1 (HIV-1) encodes a potent transactivator, Tat. HDM2 has been shown to interact with Tat and mediating its ubiquitination in vitro and in vivo. In addition, HDM2 is a positive regulator of Tat-mediated transactivation, indicating that the transcriptional properties of Tat are stimulated by ubiquitination (Bres et al. *Nat Cell Biol.* **2003**, *5*, 754-61).

Small molecule inhibitors of the HDM2 interaction have been reported and show pro-apoptotic effects in *in vitro* models and an antitumor effect in animal models of cancer. Thus, benzodiazepines have been used as a chemical scaffold to achieve HDM2 inhibitory activity (Grasberger et al. *J. Med. Chem.* **2005**, *48*, 909-912; Parks et al. *Bioorganic & Medicinal Chemistry Letters* **2005**, *15*, 765–770). Similarly, imidazolines (Vassilev et al. *Science* **2004**, *303*, 844-848), isoindolones (Hardcastle et al. *Bioorganic & Medicinal Chemistry Letters* **2005**, *15*, 1515–1520), norbornanes (Zhao et al. *Cancer Letters* **2002**, *183*, 69–77) and sulfonamides (Galatin and Abraham *J. Med. Chem.* **2004**, *47*, 4163-4165) have been reported as small molecule HDM2 inhibitors.

- 5 -

It has also been reported that HDM2 ligands have a cytoprotective effect. Thus, HDM2 inhibitors can be employed in methods of inducing cytoprotection and are useful to protect non-target cells against the harmful effects of chemotherapeutic agents. The amount of HDM2 inhibitor that provides such an effect can be about 5 to about 10 fold
5 lower than the amount needed to induce apoptosis (Koblish et al. WO03095625, METHOD FOR CYTOPROTECTION THROUGH HDM2 AND HDM2 INHIBITION, 2003-11-20).

Pyrrolidin-2-ones have already been described as therapeutically useful compounds to
10 treat viral infections (US 6509359, PYRROLIDIN-2-ONE COMPOUNDS AND THEIR USE AS NEURAMINIDASE INHIBITORS, 1999-03-25), to inhibit factor Xa for the treatment of cardiovascular disorders (US 7226929, Pyrrolidin-2-one derivatives as inhibitors of factor Xa, 2006-03-17; Watson et al., Design and Synthesis of Orally Active Pyrrolidin-2-one-Based Factor Xa Inhibitors, Bioorganic & Medicinal Chemistry
15 Letters 2006, 16, 3784-3788), as inhibitors of 11 β HSD1 for the treatment of diabetes (WO/2005/108360, PYRROLIDIN-2-ONE AND PIPERIDIN-2-ONE DERIVATIVES AS 11-BETA HYDROXYSTEROID DEHYDROGENASE INHIBITORS, 2005-04-29). Pyrrolidin-2-ones are scaffolds for established therapeutic compounds such as rolipram, an antidepressant agent and oxiracetam, piracetam or nebracetam, being
20 nootropic drugs for the Alzheimer's disease. These compounds have low toxicity, good pharmacokinetic properties and render the chemical class of pyrrolidin-2-ones an interesting scaffold for new drug candidates.

MDMX (also known as MDM4 or HDMX) is a relative of MDM2 that was identified
25 on the basis of its ability to physically interact with p53. An increasing body of evidence, including recent genetic studies, suggests that MDMX also acts as a key negative, independent regulator of p53. Aberrant expression of MDMX may contribute to tumor formation and is observed for example in gliomas, breast cancers, retinoblastomas and in a large subset of cervical and ovarian cancer cell lines. A
30 systemic analysis of 500 human tumors (Danovi et al, MCB, 2004) of HDMX expression in primary tumors of different origins revealed a broad spectrum of human cancers with HDMX overexpression such as breast cancer, colon cancer, lung cancer,

- 6 -

prostate cancer, stomach cancer, testis cancer, larynx cancer, uterus cancer, melanoma, and sarcoma.

Specific MDMX antagonists should therefore be developed as a pharmaceutical product
5 to ensure activation of 'dormant' p53 activity in tumors that retain wild-type p53.

Although MDMX is highly homologous to MDM2, it does not possess ubiquitin ligase capability and its expression level is not p53 dependent. It was shown that MDMX could inhibit p53 transcriptional activity even stronger than MDM2 and both proteins
10 cooperate in the inactivation of p53. Therefore, to achieve full activation of p53 in tumor cells, compounds that exhibit dual specificity for MDMX and MDM2 may be superior over MDM2 or MDMX specific binders alone.

The 3-dimensional structure of human MDMX protein bound to optimized p53 peptides
15 have been solved by Kallen et al., *JBC*, 2009, 284, 8812-8821. The crystal structure of humanized zebra fish MDMX to p53 peptide by Popowicz et al., *Cell Cycle* 6:19, 2386-2392, 1 October 2007 reveals that the principle features of the p53 and MDM2 interaction are preserved in the p53/MDMX complex and that "hybrid" MDM2/MDMX inhibitors could be developed. Thus, the structures of p53/MDMX and p53/MDM2
20 complexes show that both MDMX and MDM2 utilize the same p53-binding motif and many of the same residues for binding to p53. The overall shape of the binding sites is similar in terms of general shape and orientation of hydrophobic binding pockets, but the exact sizes respectively depths of these pockets are somewhat different. Thus, in MDMX, the hydrophobic cleft on which the p53 peptide binds appears slightly more
25 flexible than in MDM2.

SUMMARY OF THE INVENTION

In one embodiment of the present invention, novel small molecules are described that
30 bind to HDM2 and/or MDMX, are inhibitors of HDM2 and/or MDMX mediated biology and can be used as novel therapeutic agents, especially for the treatment of cancer and/or viral infections.

The present invention provides at least one compound selected from formula (I), (Ia), (Ic), (Id), (Ie) or (If) and pharmaceutically acceptable salts and esters thereof. Such a compound preferably is a ligand binding to HDM2 and/or MDMX protein, inducing
5 apoptosis and inhibiting proliferation, and having therapeutic utility in cancer therapy and prevention. This therapeutic effect can be achieved by using one or more compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) alone or in combination with other agents that are used to treat or prevent cancer.

10 One or more compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) can also be used to treat or prevent cancer e.g. by protecting non-cancer cells from the deleterious effects of cytotoxic cancer treating drugs or radiotherapy. In such a treatment, a combination of either an antineoplastic agent or radiotherapy and a cytoprotective amount of at least one compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If), and preferably one or more
15 pharmaceutically acceptable excipients are used. Preferably, the compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If) (also called HDM2 and/or MDMX ligand herein) is administered prior to, concurrently or after administration of the antineoplastic agent. Additionally, the HDM2 and/or MDMX inhibitor can be administered continuously or at repeated regular intervals.

20

A compound selected from compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) can e.g. be used as a therapeutic agent in methods of treating stroke, myocardial infarction, ischemia, multi-organ failure, spinal cord injury, Alzheimer's Disease, injury from ischemic events, heart valvular degenerative disease or decreasing the side effects from
25 cytotoxic agents, such as hair loss or cardio toxicity induced by doxorubicin or paclitaxel.

A compound selected from compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) of the present invention can further be used to treat viral infections, especially in a
30 pharmaceutical combination comprising a known antiviral compound.

- 8 -


Further, the present invention is directed to a pharmaceutical composition comprising a cytoprotective amount of a compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If), and one or more pharmaceutically acceptable excipients that is applied before, concomitantly and or subsequent to the treatment of a patient with a cytotoxic cancer treatment such as radiation or a cytotoxic antineoplastic agent.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides one or more compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) that are preferably small molecule ligands of the HDM2 and/or MDMX protein and prevent or reduce binding of other proteins to HDM2 and/or MDMX.

In *in vitro* cell-based assays, one or more compounds of the present invention inhibit the interaction of the HDM2 and/or MDMX protein with the p53 protein. In such cell-based assays, such compounds demonstrate mechanistic activity such as induction of apoptosis and inhibition of proliferation. Incubation of cancer cells with one or more compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) leads to an accumulation of p53 protein, induction of p53-regulated p21 gene, and cell cycle arrest in G1 and G2 phase, resulting in potent antiproliferative activity against wild-type p53 cells *in vitro*. In contrast, these activities were not observed in cancer cells with missing p53 at comparable compound concentrations. Therefore, the activity of HDM2 and/or MDMX antagonists is likely linked to its mechanism of action. These compounds are therefore potent and selective anticancer agents.

The present invention provides one or more compounds of formula (I)



- 9 -

wherein

V is C=O, C=S or CH₂ (especially C=O);

- 5 X is sulphur, oxygen or a group of formula CH₂, CR^{4b}R^{4c}, NH, NR^{4b}, SO or SO₂ or a bond;

- Y is a group of formula CONR⁶, CH₂NR⁶, CO, COO, CH₂O, SO₂NR⁶, NR⁶CO, NR⁶SO₂, NR^{5a}CONR⁶, NR⁶COO, OCONR⁶, CONR^{5a}NR⁶, CONR^{5a}OR⁶, CH₂CO
10 CH₂CONR⁶, CH₂COO, COCR^{5a}R⁶ or a bond;

n is 1, 2, 3 or 0;

- R¹ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl,
15 cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-
aralkyl radical;

- R² is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl,
cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-
20 aralkyl radical;

- R³ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl,
cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-
aralkyl radical;

- 25 R⁴ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl,
cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-
aralkyl radical;

- 30 R^{4b} is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl,
cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-
aralkyl radical;

R^{4c} is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

5

R⁵ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

10 R^{5a} is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

15 R⁶ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

the residues R⁷ are independently from each other a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

the residues R⁸ are independently from each other a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

25

or two of the radicals R¹, R², R³, R⁴, R^{4b}, R^{4c}, R⁵, R^{5a}, R⁶, R⁷ and R⁸ together are part of an optionally substituted cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heteroaryl, aralkyl or heteroarylalkyl ring system;

30 or a pharmaceutically acceptable salt, ester, solvate or hydrate or a pharmaceutically acceptable formulation thereof.

The expression alkyl refers to a saturated, straight-chain or branched hydrocarbon group that contains from 1 to 20 carbon atoms, preferably from 1 to 12 carbon atoms, especially from 1 to 6 (e.g. 1, 2, 3 or 4) carbon atoms, for example a methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, iso-pentyl, n-hexyl,
5 2,2-dimethylbutyl or n-octyl group.

The expressions alkenyl and alkynyl refer to at least partially unsaturated, straight-chain or branched hydrocarbon groups that contain from 2 to 20 carbon atoms, preferably from 2 to 12 carbon atoms, especially from 2 to 6 (e.g. 2, 3 or 4) carbon atoms, for
10 example an ethenyl (vinyl), propenyl (allyl), iso-propenyl, butenyl, ethinyl, propinyl, butinyl, acetylenyl, propargyl, isoprenyl or hex-2-enyl group. Preferably, alkenyl groups have one or two (especially preferably one) double bond(s), and alkynyl groups have one or two (especially preferably one) triple bond(s).

15 Furthermore, the terms alkyl, alkenyl and alkynyl refer to groups in which one or more hydrogen atoms have been replaced by a halogen atom (preferably F or Cl) such as, for example, a 2,2,2-trichloroethyl or a trifluoromethyl group.

The expression heteroalkyl refers to an alkyl, alkenyl or alkynyl group in which one or
20 more (preferably 1, 2 or 3) carbon atoms have been replaced by an oxygen, nitrogen, phosphorus, boron, selenium, silicon or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom). The expression heteroalkyl furthermore refers to a carboxylic acid or to a group derived from a carboxylic acid, such as, for example, acyl, acylalkyl, alkoxy-carbonyl, acyloxy, acyloxyalkyl, carboxyalkylamide or alkoxycarbonyloxy.

25 Preferably, a heteroalkyl group contains from 1 to 12 carbon atoms and from 1 to 4 hetero atoms selected from oxygen, nitrogen and sulphur (especially oxygen and nitrogen). Especially preferably, a heteroalkyl group contains from 1 to 6 (e.g. 1, 2, 3 or 4) carbon atoms and 1, 2 or 3 (especially 1 or 2) hetero atoms selected from oxygen,
30 nitrogen and sulphur (especially oxygen and nitrogen). The term C₁-C₆ heteroalkyl refers to a heteroalkyl group containing from 1 to 6 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S and/or N (especially O and/or N). The term C₁-C₄

heteroalkyl refers to a heteroalkyl group containing from 1 to 4 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S and/or N (especially O and/or N). Furthermore, the term heteroalkyl refers to groups in which one or more hydrogen atoms have been replaced by a halogen atom (preferably F or Cl).

5

Examples of heteroalkyl groups are groups of formulae: R^a-O-Y^a -, R^a-S-Y^a -,
 $R^a-N(R^b)-Y^a$ -, R^a-CO-Y^a -, $R^a-O-CO-Y^a$ -, $R^a-CO-O-Y^a$ -, $R^a-CO-N(R^b)-Y^a$ -,
 $R^a-N(R^b)-CO-Y^a$ -, $R^a-O-CO-N(R^b)-Y^a$ -, $R^a-N(R^b)-CO-O-Y^a$ -, $R^a-N(R^b)-CO-N(R^c)-Y^a$ -,
 $R^a-O-CO-O-Y^a$ -, $R^a-N(R^b)-C(=NR^d)-N(R^c)-Y^a$ -, R^a-CS-Y^a -, $R^a-O-CS-Y^a$ -,
10 $R^a-CS-O-Y^a$ -, $R^a-CS-N(R^b)-Y^a$ -, $R^a-N(R^b)-CS-Y^a$ -, $R^a-O-CS-N(R^b)-Y^a$ -,
 $R^a-N(R^b)-CS-O-Y^a$ -, $R^a-N(R^b)-CS-N(R^c)-Y^a$ -, $R^a-O-CS-O-Y^a$ -, $R^a-S-CO-Y^a$ -,
 $R^a-CO-S-Y^a$ -, $R^a-S-CO-N(R^b)-Y^a$ -, $R^a-N(R^b)-CO-S-Y^a$ -, $R^a-S-CO-O-Y^a$ -,
 $R^a-O-CO-S-Y^a$ -, $R^a-S-CO-S-Y^a$ -, $R^a-S-CS-Y^a$ -, $R^a-CS-S-Y^a$ -, $R^a-S-CS-N(R^b)-Y^a$ -,
 $R^a-N(R^b)-CS-S-Y^a$ -, $R^a-S-CS-O-Y^a$ -, $R^a-O-CS-S-Y^a$ -, wherein R^a being a hydrogen
15 atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group; R^b being a hydrogen
atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group; R^c being a hydrogen
atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group; R^d being a hydrogen
atom, a C_1-C_6 alkyl, a C_2-C_6 alkenyl or a C_2-C_6 alkynyl group and Y^a being a direct
20 bond, a C_1-C_6 alkylene, a C_2-C_6 alkenylene or a C_2-C_6 alkynylene group, wherein each
heteroalkyl group contains at least one carbon atom and one or more hydrogen atoms
may be replaced by fluorine or chlorine atoms.

Specific examples of heteroalkyl groups are methoxy, trifluoromethoxy, ethoxy, n-pro-
pyloxy, isopropoxy, butoxy, *tert*-butoxy, methoxymethyl, ethoxymethyl,
25 $-CH_2CH_2OH$, $-CH_2OH$, methoxyethyl, 1-methoxyethyl, 1-ethoxyethyl, 2-methoxyethyl
or 2-ethoxyethyl, methylamino, ethylamino, propylamino, isopropylamino, dimethyl-
amino, diethylamino, isopropylethylamino, methylamino methyl, ethylamino methyl,
diisopropylamino ethyl, methylthio, ethylthio, isopropylthio, enol ether, dimethylamino
methyl, dimethylamino ethyl, acetyl, propionyl, butyryloxy, acetyloxy,
30 methoxycarbonyl, ethoxycarbonyl, propionyloxy, acetyl amino or propionyl amino,
carboxymethyl, carboxyethyl or carboxypropyl, N-ethyl-N-methylcarbamoyl or N-

methylcarbamoyl. Further examples of heteroalkyl groups are nitrile, isonitrile, cyanate, thiocyanate, isocyanate, isothiocyanate and alkyl nitrile groups.

The expression cycloalkyl refers to a saturated or partially unsaturated (for example, a
5 cycloalkenyl group) cyclic group that contains one or more rings (preferably 1 or 2),
and contains from 3 to 14 ring carbon atoms, preferably from 3 to 10 (especially 3, 4, 5,
6 or 7) ring carbon atoms. The expression cycloalkyl refers furthermore to groups in
which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine
or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups, thus, for
10 example, cyclic ketones such as, for example, cyclohexanone, 2-cyclohexenone or
cyclopentanone. Further specific examples of cycloalkyl groups are a cyclopropyl,
cyclobutyl, cyclopentyl, spiro[4,5]decanyl, norbornyl, cyclohexyl, cyclopentenyl,
cyclohexadienyl, decalanyl, bicyclo[4.3.0]nonyl, tetraline, cyclopentylcyclohexyl,
fluorocyclohexyl or cyclohex-2-enyl group.

15 The expression heterocycloalkyl refers to a cycloalkyl group as defined above in which
one or more (preferably 1, 2 or 3) ring carbon atoms have been replaced by an oxygen,
nitrogen, silicon, selenium, phosphorus or sulfur atom (preferably by an oxygen, sulfur
or nitrogen atom). A heterocycloalkyl group has preferably 1 or 2 ring(s) containing
20 from 3 to 10 (especially 3, 4, 5, 6 or 7) ring atoms (preferably selected from C, O, N
and S). The expression heterocycloalkyl refers furthermore to groups in which one or
more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms
or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups. Examples are a piperidyl,
prolinyl, imidazolidinyl, piperazinyl, morpholinyl, urotropinyl, pyrrolidinyl, tetrahydro-
25 thiophenyl, tetrahydropyranyl, tetrahydrofuryl or 2-pyrazolinyl group and also lactames,
lactones, cyclic imides and cyclic anhydrides.

The expression alkylcycloalkyl refers to groups that contain both cycloalkyl and also
alkyl, alkenyl or alkynyl groups in accordance with the above definitions, for example
30 alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkenyl, alkenylcycloalkyl and alkynyl-
cycloalkyl groups. An alkylcycloalkyl group preferably contains a cycloalkyl group that
contains one or two ring systems having from 3 to 10 (especially 3, 4, 5, 6 or 7) ring

carbon atoms, and one or two alkyl, alkenyl or alkynyl groups having 1 or 2 to 6 carbon atoms.

- The expression heteroalkylcycloalkyl refers to alkylcycloalkyl groups as defined above
- 5 in which one or more (preferably 1, 2 or 3) carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom). A heteroalkylcycloalkyl group preferably contains 1 or 2 ring systems having from 3 to 10 (especially 3, 4, 5, 6 or 7) ring atoms, and one or two alkyl, alkenyl, alkynyl or heteroalkyl groups having from 1 or 2 to 6 carbon atoms.
- 10 Examples of such groups are alkylheterocycloalkyl, alkylheterocycloalkenyl, alkenylheterocycloalkyl, alkynylheterocycloalkyl, heteroalkylcycloalkyl, heteroalkylheterocycloalkyl and heteroalkylheterocycloalkenyl, the cyclic groups being saturated or mono-, di- or tri-unsaturated.
- 15 The expression aryl refers to an aromatic group that contains one or more rings containing from 6 to 14 ring carbon atoms, preferably from 6 to 10 (especially 6) ring carbon atoms. The expression aryl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, SH, NH₂, N₃ or NO₂ groups. Examples are the phenyl, naphthyl, biphenyl,
- 20 2-fluorophenyl, aniliny, 3-nitrophenyl or 4-hydroxyphenyl group.

- The expression heteroaryl refers to an aromatic group that contains one or more rings containing from 5 to 14 ring atoms, preferably from 5 to 10 (especially 5 or 6) ring atoms, and contains one or more (preferably 1, 2, 3 or 4) oxygen, nitrogen, phosphorus
- 25 or sulfur ring atoms (preferably O, S or N). The expression heteroaryl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, SH, N₃, NH₂ or NO₂ groups. Examples are pyridyl (e.g. 4-pyridyl), imidazolyl (e.g. 2-imidazolyl), phenylpyrrolyl (e.g. 3-phenylpyrrolyl), thiazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl,
- 30 oxadiazolyl, thiadiazolyl, indolyl, indazolyl, tetrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl, isoxazolyl, indazolyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzthiazolyl, pyridazinyl, quinolinyl,

isoquinolinyl, pyrrolyl, purinyl, carbazolyl, acridinyl, pyrimidyl, 2,3'-bifuryl, pyrazolyl (e.g. 3-pyrazolyl) and isoquinolinyl groups.

The expression aralkyl refers to groups containing both aryl and also alkyl, alkenyl, alkynyl and/or cycloalkyl groups in accordance with the above definitions, such as, for example, arylalkyl, arylalkenyl, arylalkynyl, arylcycloalkyl, arylcycloalkenyl, alkylaryl-cycloalkyl and alkylaryl-cycloalkenyl groups. Specific examples of aralkyls are toluene, xylene, mesitylene, styrene, benzyl chloride, o-fluorotoluene, 1H-indene, tetraline, dihydronaphthalene, indanone, phenylcyclopentyl, cumene, cyclohexylphenyl, fluorene and indane. An aralkyl group preferably contains one or two aromatic ring systems (1 or 2 rings) containing from 6 to 10 carbon atoms and one or two alkyl, alkenyl and/or alkynyl groups containing from 1 or 2 to 6 carbon atoms and/or a cycloalkyl group containing 5 or 6 ring carbon atoms.

The expression heteroaralkyl refers to an aralkyl group as defined above in which one or more (preferably 1, 2, 3 or 4) carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus, boron or sulfur atom (preferably oxygen, sulfur or nitrogen), that is to say to groups containing both aryl or heteroaryl, respectively, and also alkyl, alkenyl, alkynyl and/or heteroalkyl and/or cycloalkyl and/or heterocycloalkyl groups in accordance with the above definitions. A heteroaralkyl group preferably contains one or two aromatic ring systems (1 or 2 rings) containing from 5 or 6 to 10 ring carbon atoms and one or two alkyl, alkenyl and/or alkynyl groups containing 1 or 2 to 6 carbon atoms and/or a cycloalkyl group containing 5 or 6 ring carbon atoms, wherein 1, 2, 3 or 4 of these carbon atoms have been replaced by oxygen, sulfur or nitrogen atoms.

Examples are arylheteroalkyl, arylheterocycloalkyl, arylheterocycloalkenyl, arylalkyl-heterocycloalkyl, arylalkenylheterocycloalkyl, arylalkynylheterocycloalkyl, arylalkyl-heterocycloalkenyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, heteroaryl-heteroalkyl, heteroarylcycloalkyl, heteroarylcycloalkenyl, heteroarylheterocycloalkyl, heteroarylheterocycloalkenyl, heteroarylalkylcycloalkyl, heteroarylalkylheterocycloalkenyl, heteroarylheteroalkylcycloalkyl, heteroarylheteroalkylcycloalkenyl and hetero-

arylheteroalkylheterocycloalkyl groups, the cyclic groups being saturated or mono-, di- or tri-unsaturated. Specific examples are a tetrahydroisoquinoliny, benzoyl, 2- or 3-ethylindolyl, 4-methylpyridino, 2-, 3- or 4-methoxyphenyl, 4-ethoxyphenyl, 2-, 3- or 4-carboxyphenylalkyl group.

5

As already stated above, the expressions cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aryl, heteroaryl, aralkyl and heteroaralkyl also refer to groups in which one or more hydrogen atoms of such groups have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups.

10

The expression "optionally substituted" especially refers to groups in which one, two, three or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups. This expression refers furthermore to groups that are substituted by one, two, three or more unsubstituted C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ heteroalkyl, C₃-C₁₀ cycloalkyl, C₂-C₉ heterocycloalkyl, C₆-C₁₀ aryl, C₁-C₉ heteroaryl, C₇-C₁₂ aralkyl or C₂-C₁₁ heteroaralkyl groups.

15

Preferred substituents are F, Cl, Br, Me, OMe, CN or CF₃.

20

Preferably, all alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aralkyl and heteroaralkyl groups described herein may optionally be substituted.

25

Preferred are compounds of formula (I) wherein the radicals R⁵ and R⁶ together are part of an optionally substituted heterocycloalkyl, heteroalkylcycloalkyl, heteroaryl or heteroarylalkyl ring system, and/or wherein R² and R³ together are part of an optionally substituted cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl ring system.

30

- 17 -

Preferred are compounds of formula (I) wherein X is sulphur, oxygen, NH, CH₂, SO, SO₂, especially sulphur.


Further preferred are compounds of formula (I) wherein Y is a group of formula
5 CONR⁶.

Further preferred are compounds of formula (I) wherein n is 0 or 1, especially 1.

Further preferred are compounds of formula (I) wherein R⁷ is hydrogen.
10

Moreover preferred are compounds of formula (I) wherein R⁸ is hydrogen.

Especially preferred are compounds of formula (Ia)



15

(Ia)

wherein

R¹ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl,
20 cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-
aralkyl radical;

R² is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl,
cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-
25 aralkyl radical;

- 18 -

R³ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

5 R⁴ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

10 R⁵ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

15 R⁶ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

or the radicals R⁵ and R⁶ together are part of an optionally substituted heterocycloalkyl, heteroalkylcycloalkyl, heteroaryl or heteroarylalkyl ring system, and/or R² and R³ together are part of an optionally substituted cycloalkyl, alkylcycloalkyl, 20 heterocycloalkyl, heteroalkylcycloalkyl, aralkyl or heteroaralkyl ring system;

or a pharmaceutically acceptable salt, ester, solvate or hydrate or a pharmaceutically acceptable formulation thereof.

25 Further preferred are compounds of formula (I) or (Ia), wherein R¹ is a cycloalkyl, alkylcycloalkyl, heterocycloalkyl, heteroalkylcycloalkyl, aryl, heteroaryl, aralkyl or heteroaralkyl radical.

Further preferred are compounds of formula (I) or (Ia), wherein R¹ is an aryl, heteroaryl, 30 aralkyl or heteroaralkyl radical.

- 19 -

Further preferred are compounds of formula (I) or (Ia), wherein R^1 is a group of formula $-A-Ar$ or $-A-Cy$ (especially $-A-Ar$) wherein A is a bond, C_1-C_4 alkyl (especially a bond, CH_2 or $CH(CH_3)$) or C_1-C_6 heteroalkyl (e.g. $CH(CH_2N(CH_3)_2)$), or wherein A is a group of formula $-CHR^{1a}$ wherein R^{1a} is a C_1-C_6 heteroalkyl group, Cy is an optionally substituted C_3-C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing from 3 to 7 ring atoms and Ar is an optionally substituted (e.g. by 1, 2 or 3 substituents) phenyl ring or an optionally substituted (e.g. by 1, 2 or 3 substituents) heteroaryl ring containing 5 or 6 ring atoms (especially including from 1 to 3 heteroatoms selected from O, S and N), especially preferably Ar is an optionally substituted phenyl or an optionally substituted pyridyl residue (e.g. a 4-bromobenzyl residue). Preferred substituents are F, Cl, Br, CN, CH_3 , OCH_3 and CF_3 .

Further preferred are compounds of formula (I) or (Ia), wherein R^1 is a group of formula $-A-Ar$ wherein A is a bond or C_1-C_4 alkyl (especially a bond, CH_2 or $CHCH_3$) and Ar is an optionally substituted (e.g. by 1, 2 or 3 substituents) phenyl ring or an optionally substituted (e.g. by 1, 2 or 3 substituents) heteroaryl ring containing 5 or 6 ring atoms (especially containing from 1 to 3 heteroatoms selected from O, S and N), especially preferably Ar is an optionally substituted phenyl or an optionally substituted pyridyl residue (e.g. a 4-bromobenzyl residue).

Especially preferably, R^1 is a group of formula $-A$ -phenyl (especially $-CH_2$ -phenyl) which is optionally substituted, preferably by one or two halogen atoms selected from F, Cl and Br and wherein A is preferably a group of formula $-CHR^{1a}$ wherein R^{1a} is a C_1-C_6 heteroalkyl group (e.g. $COOH$, CH_2COOH)

Further preferred, R^1 is cyclopropylmethyl, picolyl, phenylbenzyl or phenoxybenzyl, all of which may optionally be substituted.


Further preferred are compounds of formula (I) or (Ia), wherein R^2 is hydrogen.

Further preferred are compounds of formula (I) or (Ia), wherein R^3 is C_1-C_6 alkyl, an aryl (especially phenyl), heteroaryl, aralkyl or heteroaralkyl residue, all of which may

- 20 -

be substituted (e.g. by 1, 2 or 3 substituents). Especially preferably, R^3 is an optionally substituted phenyl group, an optionally substituted benzyl group or an optionally substituted heteroaryl residue having 1 or 2 rings and from 5 to 10 ring atoms (especially 2 rings and a total of 9 ring atoms) including 1, 2, 3 or 4 heteroatoms selected from O, S and N (especially N). Preferred substituents are F, Cl, Br, C_1 - C_4 alkyl groups (e.g. CH_3) and C_1 - C_6 heteroalkyl groups (e.g. $CH_2SO_3^-$, $(CH_2)_5NH_2$).

Further preferred are compounds of formula (I) or (Ia), wherein R^3 has the following structure




10

wherein E is N or CH, R^{3a} is H, C_1 - C_6 alkyl or C_1 - C_6 heteroalkyl (especially H or CH_3), R^{3b} is H, F, Cl, Br, CH_3 , OCH_3 or CF_3 and R^{3c} is H, F, Cl, Br, CH_3 , OCH_3 or CF_3 (especially preferably, E is CH, R^{3a} is H, R^{3b} is Cl and R^{3c} is H).

Further preferred are compounds of formula (I) or (Ia), wherein R^3 is an aryl (especially phenyl), heteroaryl, aralkyl or heteroaralkyl residue, all of which may be substituted (e.g. by 1, 2 or 3 substituents); especially preferably, R^3 is an optionally substituted heteroaryl residue having 1 or 2 rings and from 5 to 10 ring atoms (especially 2 rings and a total of 9 ring atoms) including 1, 2, 3 or 4 heteroatoms selected from O, S and N (especially N).

20


Further preferred are compounds of formula (I) or (Ia), wherein R^3 has the following structure



- 21 -

wherein E is N or CH, R^{3a} is H or CH₃, R^{3b} is F, Cl or Br and R^{3c} is H, F, Cl or Br (especially preferably, E is CH, R^{3a} is H, R^{3b} is Cl and R^{3c} is H).

Further preferred, R² and R³ together are part of an optionally substituted
5 heterocycloalkyl or heteroaralkyl ring. Moreover preferred, R² and R³ together are part of a group having the following structure:



Further preferred are compounds of formula (I) or (Ia), wherein R⁴ is C₁-C₆ alkyl, C₂-C₆
10 alkenyl, optionally substituted C₁-C₄ alkyl-C₃-C₇ cycloalkyl, an optionally substituted phenyl ring, an optionally substituted benzyl group or an optionally substituted heteroaryl ring having 5 or 6 ring atoms including from 1 to 3 heteroatoms selected from O, S and N (e.g. pyridyl). Especially preferably, R⁴ is a phenyl ring which is substituted by 1 or 2 substituents, preferably selected from F, Br, Cl, I, NO₂, methyl or
15 cyanide (e.g. 4-methylphenyl), or an unsubstituted phenyl ring. Moreover preferably, R⁴ is phenyl or 4-methylphenyl.

Further preferred are compounds of formula (I) or (Ia), wherein R⁴ is C₁-C₆ alkyl or an
optionally substituted phenyl ring or an optionally substituted heteroaryl ring having 5
20 or 6 ring atoms and containing from 1 to 3 heteroatoms selected from O, S and N. Especially preferably, R⁴ is a phenyl ring which is substituted by 1 or 2 substituents, preferably selected from F, Br, Cl, I, methyl or cyanide (e.g. 4-methylphenyl).

Further preferred are compounds of formula (I) or (Ia), wherein R⁵ is an alkyl, hetero-
25 alkyl, heterocycloalkyl, heteroalkylcycloalkyl or heteroaralkyl group, all of which groups may be substituted.

Further preferred are compounds of formula (I) or (Ia), wherein R⁵ is selected from the
following groups: C₁-C₆ alkyl; heteroalkyl containing 1-6 carbon atoms and 1, 2 or 3
30 heteroatoms selected from O, S and N; heteroalkylcycloalkyl comprising a C₁-C₄ alkyl

- 22 -

group or a C₁-C₄ heteroalkyl group and an optionally substituted heterocycloalkyl group containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N; heteroaralkyl comprising a C₁-C₄ alkyl group or a C₁-C₄ heteroalkyl group and an optionally substituted heteroaryl group containing 5 or 6 ring atoms including 1, 2 or 3
5 heteroatoms selected from O, S and N; optionally substituted heteroaryl containing 5 or 6 ring atoms including 1, 2 or 3 heteroatoms selected from O, S and N; and optionally substituted heterocycloalkyl containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N.

10 Further preferred are dimers of compounds of formulas (I) and/or (Ia) that are linked via a heteroalkyl, heteroalkylcycloalkyl or a heteroaralkyl group, preferably via R⁵.

Further preferred are compounds of formula (I) or (Ia), wherein R⁵ is C₁-C₆ alkyl; heteroalkyl containing 1-6 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S,
15 N; heteroalkylcycloalkyl comprising a C₁-C₄ alkyl group and an optionally substituted heterocycloalkyl group containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N; heteroaralkyl comprising a C₁-C₄ alkyl group and an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N.


20

Further preferred are compounds of formula (I) or (Ia), wherein R⁶ is hydrogen or C₁-C₄ alkyl, especially hydrogen.

Further preferred are compounds of formula (I) or (Ia), wherein R⁵ and R⁶ together with
25 the nitrogen atom to which they are bound form an optionally substituted (e.g. by 1, 2 or 3 substituents) heterocycloalkyl ring containing 4, 5, 6 or 7 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N.

Moreover preferred, R⁵ and R⁶ together with the nitrogen atom to which they are bound
30 form the following group:

- 23 -



Therein, m is 0, 1 or 2; o is 0, 1 or 2; the sum of m and o is preferably from 0 to 3; Q is N-R^{6x}, CR^{6y}R^{6z}, C=O, -CO-NR^{6x}-, -NR^{6x}-CO-NR^{6y}-, -SO₂-NR^{6x}-, -SO-NR^{6x}- or -O-CO-NR^{6x}-, wherein R^{6x}, R^{6y} and R^{6z} independently from each other are a hydrogen atom, OH, NH₂, SH or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical.

Preferably, Q is N-CO-R^{6a}, NR^{6b} or CHR^{6c} wherein R^{6a} is C₁-C₆ alkyl, C₁-C₆ heteroalkyl, NH₂, optionally substituted phenyl or hydrogen; R^{6b} is optionally substituted phenyl or optionally substituted heteroaryl containing 5 or 6 ring atoms including one or two heteroatoms selected from O, S or N; R^{6c} is C₁-C₆ heteroalkyl, NH₂ or OH.

Further preferred, Q is N-CO-NHR^{6d}, N-COOR^{6e}, N-SO₂R^{6f}, N-SO₂NHR^{6g}, N-NHCOR^{6h}, CH-NH₂, CH-OH, CH-SH, CH-NH-COR⁶ⁱ, CO, CONH, NHCONH, SO₂NH, OCONH, CH-COOH, CH-COOR^{6j}, CH-COR^{6k} or CH-SO₂R^{6l}, wherein R^{6d}, R^{6e}, R^{6f}, R^{6g}, R^{6h}, R⁶ⁱ, R^{6j}, R^{6k} and R^{6l} independently from each other are a hydrogen atom, OH, NH₂, SH or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical, especially hydrogen or a C₁-C₆ alkyl group or a C₁-C₆ heteroalkyl group.

Preferably, Q is N-CO-R^{6a} wherein R^{6a} is preferably NH₂, C₁-C₆ alkyl, NH-C₁-C₆ alkyl or N(C₁-C₆ alkyl)₂.

Further preferably, group Q contains a hydrogen bond acceptor (especially an atom or group having a lone electron pair like e.g. an electronegative atom such as fluorine, oxygen, or nitrogen).

Further preferred, o is 1 and m is 1.


- 24 -

Especially preferably, m is 1, o is 1 and Q is N-CO-R^{6a}. Thereby R^{6a} is preferably C₁-C₄ alkyl or NH-C₁-C₄ alkyl (e.g. CH₃ or NHCH₂CH₃).

- 5 Especially preferred are compounds of formula (I) or (Ia), wherein R⁵ and R⁶ together are part of an optionally substituted (e.g. by 1, 2 or 3 substituents like e.g. =O) piperazine ring.

- Further preferred are compounds of formula (I) or (Ia), wherein R⁵ and R⁶ together are
 10 part of an optionally substituted (e.g. by 1, 2 or 3 substituents) heterocycloalkyl ring containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N

Further preferred are compounds of formula (Ic):



15

(Ic)

wherein W is an optionally substituted phenyl ring or an optionally substituted heteroaryl group having 5 or 6 ring atoms including 1 or 2 heteroatoms selected from O, S and N; and wherein R¹, R^{3a}, R⁴, R⁵, R⁷, R⁸, E, X, Y and n are defined as above.


20

- Especially preferred are compounds of formula (Ic) wherein W is an optionally substituted phenyl ring (preferably substituted by 1, 2 or 3 halogen atoms selected from F, Cl and Br); R¹ is an optionally substituted benzyl group (preferably substituted by 1, 2 or 3 halogen atoms selected from F, Cl and Br); X is S; Y is CONR⁶; n is 1; R^{3a} is
 25 hydrogen; R⁴ is an optionally substituted phenyl group (preferably unsubstituted or substituted by a methyl group, especially in the para position); R⁷ and R⁸ are hydrogen;

- 25 -

E is CH; and R⁵ and R⁶ are defined as above; especially preferably, R⁵ and R⁶ together are part of an optionally substituted piperazine ring (especially as defined above).

Further preferred are compounds of formula (Id):




(Id)

wherein R¹, R^{3a}, R^{3b}, R^{3c}, R⁴, R⁵, R⁶, E and X are defined as above.

- 10 Especially preferred are compounds of formula (Id) wherein R¹ is an optionally substituted benzyl group (preferably substituted by 1, 2 or 3 halogen atoms selected from F, Cl and Br); X is S; R^{3a} is hydrogen; R^{3b} is Cl; R^{3c} is hydrogen, R⁴ is an optionally substituted phenyl group (preferably unsubstituted or substituted by a methyl group, especially in the para position); E is CH; and R⁵ and R⁶ are defined as above;
- 15 especially preferably, R⁵ and R⁶ together are part of an optionally substituted piperazine ring (especially as defined above).

Further preferred are compounds of formula (Ie):




(Ie)

wherein R^1 , R^{3a} , R^{3b} , R^{3c} , R^4 , E, Q, m, o and X are defined as above.

Especially preferred are compounds of formula (Ie) wherein R^1 is an optionally substituted benzyl group (preferably substituted by 1, 2 or 3 halogen atoms selected from F, Cl and Br); X is S; R^{3a} is hydrogen; R^{3b} is Cl; R^{3c} is hydrogen, R^4 is an optionally substituted phenyl group (preferably unsubstituted or substituted by a methyl group, especially in the para position); E is CH; Q is N-CO- R^{6a} wherein R^{6a} is preferably C₁-C₆ alkyl, NH-C₁-C₆ alkyl or N(C₁-C₆ alkyl)₂; m is 0, 1 or 2; o is 0, 1 or 2; and the sum of m and o is preferably from 0 to 3. Especially preferably, m is 1, o is 1 and Q is N-CO- R^{6a} , wherein R^{6a} is preferably C₁-C₄ alkyl or NH-C₁-C₄ alkyl (e.g. CH₃ or NHCH₂CH₃).

Further preferred are compounds of formula (If)



(If)

wherein A, Ar, R^{3a} , R^{3b} , R^{3c} , E, X, R^{4a} and R^{6a} are defined as above.

Especially preferred are compounds of formula (If) wherein X is S; R^{3a} is hydrogen; R^{3b} is Cl; R^{3c} is hydrogen, R^{4a} is hydrogen or a methyl group; E is CH; A is CH₂; Ar is phenyl which is substituted by one or two halogen atoms selected from F, Cl and Br; and R^{6a} is C₁-C₄ alkyl or NH-C₁-C₄ alkyl (e.g. CH₃ or NHCH₂CH₃).

A further preferred embodiment of the present invention relates to compounds of formula (I) or (Ia), wherein

R¹ is aryl, heteroaryl, arylalkyl or heteroarylalkyl, all of which may be substituted by F, Br, Cl, I, methyl or cyanide,

5 R² is hydrogen,

R³ is aryl, heteroaryl, arylalkyl or heteroarylalkyl, all of which may be substituted by F, Br, Cl, I, methyl or cyanide,

R⁴ is aryl, heteroaryl, arylalkyl or heteroarylalkyl,

10 R⁵ and R⁶ are independently selected from hydrogen, alkyl, heteroalkyl, aryl, alkenyl, akynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, arylalkyl or heteroarylalkyl, or wherein R⁵ and R⁶ may be part of one heteroaryl, heterocycloalkyl, heteroalkylcycloalkyl or heteroarylalkyl ring system, and wherein the other residues and groups are defined as above.

15 A further preferred embodiment of the present invention relates to compounds of formula (I) or (Ia), wherein

R¹ is aryl, heteroaryl, arylalkyl or heteroarylalkyl, all of which may be substituted by F, Br, Cl, I, methyl or cyanide,

20 R² and R³ together are part of a heteroaryl, heteroalkyl, heterocycloalkyl or heteroalkylcycloalkyl ring system such as but not restricted to 1,3-dihydroindole, 1,3-dihydro-indol-2-one, 2,3-dihydro-1H-indazole, tetrahydro-quinoline, tetrahydro-quinoline-2-one, 3,4-dihydro-1H-quinolin-2-one, 3,4-dihydro-1H-quinazolin-2-one, all of which may be substituted by F, Br, Cl, I, methyl or cyanide,

R⁴ is selected from aryl, heteroaryl, arylalkyl or heteroarylalkyl,

25 R⁵ and R⁶ are independently selected from hydrogen, alkyl, heteroalkyl, aryl, alkenyl, akynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, arylalkyl or heteroarylalkyl, or wherein R⁵ and R⁶ may also be part of one heteroaryl, heterocycloalkyl, heteroalkylcycloalkyl or heteroarylalkyl ring system, and wherein the other residues and groups are defined as above.


30

- 28 -

A further preferred embodiment of the present invention relates to compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If), wherein R^3 and the sulfanyl group (i.e. the group carrying X) bearing the R^4 group are in cis position (especially when R^2 is H).


- 5 An especially preferred embodiment are enantiomerically pure compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If).

Further preferred is the following compound:




10

i.e. the following diastereomers:




- 29 -

especially the following diastereomer:




5 Further preferred is the following compound:




i.e. the following diastereomers:

- 30 -




especially the following diastereomer:




5

Further preferred is the following compound:

- 31 -




i.e. the following diastereomers.



5

especially the following diastereomer:




Preferably, the compounds described in:

1. Ng et al. Angew. Chem. Int. Ed. 2007, 46, 5352-5355 and
2. Ng et al. Organic Letters 2006, Vol. 8, No. 18, 3999-4002 (and supporting information thereof) are excluded from the scope of the present application and/or patent.


5

Further preferred, one or more of the following compounds are excluded from the present application and/or patent:




10


Further preferred, also the following compounds are excluded from the present application and/or patent:



wherein R^3 is $p\text{-C}_6\text{H}_5\text{CH}_2\text{OH}$ (or a derivative thereof which is bound to the solid phase as described in Ng et al. Angew. Chem. Int. Ed. 2007, 46, 5352-5355), R^1 is selected from the following groups:



R^4 is $\text{para-C}_6\text{H}_5\text{CH}_3$ and R^5 is selected from the following groups:




All those compounds are described in Ng et al. Angew. Chem. Int. Ed. 2007, 46, 5352-5355.

- 34 -

Further preferred, compounds of formula (I) or (Ia) are excluded wherein R³ is p-C₆H₅CH₂OH (or a derivative thereof which is bound to the solid phase as described in Ng et al. Angew. Chem. Int. Ed. 2007, 46, 5352-5355).

- 5 Further preferred, also the following compounds are excluded from the present application and/or patent (X = H, I):



- The present invention further provides pharmaceutical compositions comprising a
 10 compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If) as defined herein or a pharmaceutically acceptable ester, prodrug, hydrate, solvate or salt thereof, optionally in combination with a pharmaceutically acceptable carrier.

- A further preferred embodiment of the present invention relates to pharmaceutical
 15 compositions comprising one or more compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) as defined herein or a pharmaceutically acceptable ester, prodrug, hydrate, solvate or salt thereof, optionally in combination with a pharmaceutically acceptable carrier, further comprising one or more other anti-tumor agents, wherein the anti-tumor agent is especially selected from 16-Aza-epothilone B, Aldesleukin, Amifostine, Aranose,
 20 Bevacizumab, Bleocin, Bleomycin, BMS-184476, Bortezomib, Calcitriol, Carmustine, Canertinib, Canfosfamide, Capecitabine, Carboplatin, Carmustine, Cefixime, Ceftriaxone, Celecoxib, Celmoleukin, Cetuximab, Ciclosporin, Cisplatin, Clodronate, Cyclophosphamide, Cytarabine, Doxorubicin, Edatrexate, Efaproxirala, EKB-569, Epirubicin,
 25 Epratuzumab, Erlotinib, Etoposide, ET-18-OCH₃, Exatecan, Fludarabine, Fluorouracil, Folinic acid, Galarubicin, Gefinitib, Gemcitabine, Gemtuzumab, Gimatecan,

Glufosfamide, Granisetron, Homoharringtonine, Hyaluronic acid, Ibandronate, Ibritumomab, Ifosfamide, Imatinib, Interferon alfa, Interferon alfa-2a, Interferon alfa-2b, Irinotecan, Isoflavone, Isotretinoin, Ixabepilone, Ketoconazole, Lapatinib, Leflunomide, Lenograstim, Leucovorin, Lexidronam, Linezolid, Lometrexol, 5 Lurtotecan, MEN10755, Methotrexate, Mitomycin, Neridronate, Nimesulide, Nitroglycerin, 06-Benzylguanine, Omeprazole, Ortaxel, Oxaliplatin, Paclitaxel, Patupilone, Pegfilgrastim, PEG-filgrastim, Pelitinib, Pemetrexed, Pentostatin, Perifosine, Plevitrexed, Polyprenoic acid, Quinupristin, Raloxifene, Raltitrexed, Ramosetron, Retinoic acid, Risedronate, Rituximab, Rofecoxib, Rubitecan, S-9788, 10 Sabarubicin, Sargramostim, Satraplatin, SN-38, Sorafenib, Suberanilohydroxamic acid, Sutent, Tamoxifen, Taxotere, Tazarotene, Tegafur, Temozolamide, Tesmilifene, Tetrodotoxin, Thalidomide, Tipifarnib, Topotecan, Trabectedin, Trastuzumab, Trastuzumab, Tretinoin, Vatalanib, Vincristine, Vinorelbine, Vinscrisone, ZD-6474, Zoledronate or Zosuquidar.

15

In a preferred embodiment, the compounds of the present invention sensitize cancer cells for radio and/or chemotherapy whereas they display chemoprotective and/or radioprotective activity on healthy cells. Thereby the dosage of these therapies can be better adjusted.

20

A further preferred embodiment of the present invention relates to a pharmaceutical composition comprising one or more compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) as defined herein or one or more pharmaceutically acceptable esters, prodrugs, hydrates, solvates or salts thereof, optionally in combination with a pharmaceutically acceptable carrier. The pharmaceutical composition optionally comprises one or more 25 antiviral agents. Preferably, the antiviral agent is selected from 3TC, Abacavir, Adefovir, Didanosine, Acyclovir, Amprenavir, Amantadine, Amoxovir, AZT, Clevudine, Delavirdine, d4T, Emtricitabine, Entecavir, Famciclovir, Ganciclovir, Indinavir, Lamivudine, Nelfinavir, Nevirapine, Oseltamavir, Rimantadine, Ritonavir, Saquinavir, 30 Septrin, Telbivudine, Tenofovir, Valacyclovir, Valganciclovir, Valopicitabine or Zanamivir.

It is a further object of the present invention to provide a compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If) as defined herein or a pharmaceutical composition as defined herein for the preparation of a medicament for the treatment of cancer and/or viral infections

5

The compounds selected from formula (I), (Ia), (Ic), (Id), (Ie) or (If) of the present invention are e.g. HDM2 and/or MDMX ligands and show binding affinities from about 1 nM to about 100 μ M to HDM2 and/or MDMX, preferably from about 1 nM to about 10 μ M, especially to about 1 μ M, preventing binding of p53 and other proteins,
10 inhibition of proliferation and induction of apoptosis in cell based assays, especially in the assays described herein.

The compounds of the present invention are useful in the treatment or control of cell proliferative disorders, in particular oncological disorders. These compounds and
15 formulations containing said compounds are useful in the treatment or control of solid tumors, such as, for example, breast, colon, lung and prostate tumors, as well as osteosarcoma, acute myeloid leukaemia, sporadic endometrial cancer, melanoma, malignant melanoma, soft tissue Sarcoma, B-cell chronic lymphocytic leukaemia, gastric cancers, cervical cancer, hepatocellular carcinoma, and colorectal cancer.

20

The compounds described herein are especially useful for the treatment and/or prevention of cancers associated with overexpression of HDM2 and/or MDMX.

Accordingly the compounds of the present invention are especially useful for the
25 treatment and/or prevention of the following cancers associated with MDM2 and/or MDMX:

MDM2 is amplified in 7% of all human cancers. Gene amplification was observed in 19 tumor types, with the highest frequency observed in soft tissue tumors (20%),
30 osteosarcomas (16%) and esophageal carcinomas (13%). Tumors which showed a higher incidence of MDM2 amplification than p53 mutation were soft tissue tumors, testicular germ cell cancers and neuro-blastomas (Momand et al, NAR, 1998). Naturally

occurring polymorphism (SNP309) occurring within the MDM2 promoter leads to an increase in MDM2 transcription and translation. The overall frequency of MDM2 amplification in these human tumors was 7%. It is a common event in hematological malignancies. A list of cancers with a wild type of p53 gene that is sensitive to MDM2 inhibitors includes: B-cell CLL (chronic lymphocytic leukemia) (Coll-Muler et al, Blood, 2006), AML (Kojima et al, Blood, 2005), multiple myeloma (Shruhmer et al, Blood, 2005), neuroblastoma (Cattelani et al, CCR, 2008), Hodgkin lymphoma (Drakos et al, CCR, 2007), osteosarcoma and prostate cancer (Vassilev et al, Science, 2004), Kaposi's sarcoma (Sarek, J. Clinic. Invest., 2007), rhabdomyosarcoma (Miyachi et al, CCR, 2009), RCC (renal cell carcinoma) (Roberts et al, CR, 2009), squamous cell carcinoma and esophageal cancers (Cescon et al, CCR, 2009), cutaneous melanoma (Firoz et al, CCR, 2009), retinoblastoma (Laurie et al, Nature, 2006). There are evidences that pancreatic cancer with wild type p53 gene could be sensitive to MDM2 inhibitors as well (submitted for publication).

15

Table 1. Summary of MDM2 gene amplification frequencies from 28 human tumors (Momand et al, NAR, 1998).

Tumor type	MDM2 amplification (<i>n</i> ^a) (%)	References
Brain tumors	6.7 (239)	57-60
Astrocytomas	8.1 (37)	57,60
Glioblastomas	6.8 (191)	57,58,60
Medulloblastomas	0 (8)	59
Other	0 (3)	60
Breast carcinomas	5.9 (1774)	61-65
Cervical carcinomas	1.1 (88)	19,66
Esophageal carcinomas	13 (96)	14,67
Leukemias/lymphomas	0 (304)	68-70

- 38 -

Hepatoblastomas	0 (38)	71
Lung	5.7 (88)	72-74
Lung cancers (NSCLC)	6.0 (83)	72,74
Lung (not specified)	0 (5)	73
Nasopharyngeal carcinomas	2.1 (46)	75
Neuroblastoma	2.0 (51)	76-78
Osteosarcomas	16 (207)	3,79-82
Ovarian carcinomas	3.1 (190)	64,83
Pancreatic carcinomas	0 (25)	84
Soft tissue tumors	20 (479)	3,36,76,79,80,85-90
Ewing's sarcomas	10 (30)	85
Leiomyosarcomas	0 (46)	79,86,88
Lipomas (benign)	30 (43)	80,86
Liposarcomas	29 (87)	3,79,80,87,89
Malignant fibrous histiocytomas	21 (163)	3,79,80,86,90
Malignant Schwannomas	19 (16)	79
Sarcomas (non-specific) ^b	13 (85)	36,76
Various ^c	33 (9)	76,79,86
Testicular tumors	4.6 (64)	91,92
Thyroid carcinomas	0 (22)	93
Urothelial carcinomas	2.2 (137)	94,95
Wilms' tumors	0 (40)	76

Total number of tumor samples analyzed was 3889 and the average MDM2 gene amplification frequency was 7.2%.

^aNumber of samples analyzed.

^bSarcomas of soft tissue origin that were not specified.

^cSoft tissue tumors that did not fall into the listed classes. The number of samples was less than five in any individual class.

- 5 Human MDMX gene maps in chromosomal region 1q32, which is frequently amplified in human cancers. It has been documented in 4% of gliomas (Riemenschneider, CR 1999) and 5% breast cancers (Danovi et al, MCB, 2004). Recently, ~60% of retinoblastomas (Laurie, Nature 2006) have been found to bear HDMX overexpression. Moreover, HDMX gene was found overexpressed in a large subset of cervical and
- 10 ovarian cancer cell lines (Ramos, CR 2001). A systemic analysis of HDMX expression in primary tumors of different origins revealed broad spectrum of human cancers with HDMX overexpression

Table 1. Summary of HDMX gene amplification frequencies from 500 human tumors (Danovi et al, MCB, 2004).

Tumor type	Total	HDMX overexpressed
Breast cancer	218	41
Colon cancer	27	5
Lung cancer	88	16
Prostate cancer	25	0
Stomach cancer	14	6
Testis cancer	11	3
Larynx cancer	13	3
Uterus cancer	13	2
Melanoma	14	2
Sarcoma	10	0

A therapeutically effective amount of a compound in accordance with this invention means an amount of compound that is effective to prevent, alleviate or ameliorate

symptoms of disease or prolong the survival of the subject being treated. Determination of a therapeutically effective amount is within the skill in the art.

The therapeutically effective amount or dosage of a compound according to this

5 invention can vary within wide limits and may be determined in a manner known in the art. Such dosage may be adjusted to the individual requirements in each particular case including the specific compound being administered, the route of administration, the condition being treated, as well as the patient being treated.

10 Examples of pharmacologically acceptable salts of sufficiently basic compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) are salts of physiologically acceptable mineral acids like hydrochloric, hydrobromic, sulfuric and phosphoric acid; or salts of organic acids like methanesulfonic, p-toluenesulfonic, lactic, acetic, trifluoroacetic, citric, succinic, fumaric, maleic and salicylic acid. Further, a sufficiently acidic compound of
15 formula (I), (Ia), (Ic), (Id), (Ie) or (If) may form alkali or earth alkali metal salts, for example sodium, potassium, lithium, calcium or magnesium salts; ammonium salts; or organic base salts, for example methylamine, dimethylamine, trimethylamine, triethylamine, ethylenediamine, ethanolamine, choline hydroxide, meglumin, piperidine, morpholine, tris-(2-hydroxyethyl)amine, lysine or arginine salts; all of which
20 are also further examples of salts of formula (I), (Ia), (Ic), (Id), (Ie) or (If). Compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) may be solvated, especially hydrated. The hydratization/hydration may occur during the process of production or as a consequence of the hygroscopic nature of the initially water free compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If). The solvates and/or hydrates may e.g. be present in solid or liquid form.

25

It should be appreciated that certain compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) may have tautomeric forms from which only one might be specifically mentioned or depicted in the following description, different geometrical isomers (which are usually denoted as cis/trans isomers or more generally as (E) and (Z) isomers) or different
30 optical isomers as a result of one or more chiral carbon atoms (which are usually nomenclatured under the Cahn-Ingold-Prelog or R/S system). All these tautomeric forms, geometrical or optical isomers (as well as racemates and diastereomers) and

polymorphous forms are included in the invention. Since the compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) may contain asymmetric C-atoms, they may be present either as achiral compounds, mixtures of diastereomers, mixtures of enantiomers or as optically pure compounds. The present invention comprises both all pure enantiomers
5 and all pure diastereomers, and also the mixtures thereof in any mixing ratio.

The therapeutic use of compounds according to formula (I), (Ia), (Ic), (Id), (Ie) or (If), their pharmacologically acceptable salts, solvates and hydrates, respectively, as well as formulations and pharmaceutical compositions also lie within the scope of the present
10 invention.

The pharmaceutical compositions according to the present invention comprise at least one compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If) as an active ingredient and, optionally, carrier substances and/or adjuvants.

15

The present invention also relates to pro-drugs which are composed of a compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If) and at least one pharmacologically acceptable protective group which will be cleaved off under physiological conditions, such as an alkoxy-, arylalkyloxy-, acyl-, acyloxymethyl group (e.g. pivaloyloxymethyl), an 2-alkyl-, 2-aryl- or 2-arylalkyl-oxycarbonyl-2-alkylidene ethyl group or an acyloxy group
20 as defined herein, e.g. ethoxy, benzyloxy, acetyl or acetyloxy or, especially for a compound of formula (I), (Ia), (Ic), (Id), (Ie) or (If), carrying a hydroxy group (-OH): a sulfate, a phosphate (-OPO₃ or -OCH₂OPO₃) or an ester of an amino acid. Especially preferred are pro-drugs of the hydroxy group of a compound of formula (I), (Ia), (Ic),
25 (Id), (Ie) or (If).

As mentioned above, therapeutically useful agents that contain compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If), their solvates, salts or formulations are also comprised in the scope of the present invention. In general, compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) will be administered by using the known and acceptable modes known in the
30 art, either alone or in combination with any other therapeutic agent.


For oral administration such therapeutically useful agents can be administered by one of the following routes: oral, e.g. as tablets, dragees, coated tablets, pills, semisolids, soft or hard capsules, for example soft and hard gelatine capsules, aqueous or oily solutions, emulsions, suspensions or syrups, parenteral including intravenous, intramuscular and subcutaneous injection, e.g. as an injectable solution or suspension, rectal as suppositories, by inhalation or insufflation, e.g. as a powder formulation, as microcrystals or as a spray (e.g. liquid aerosol), transdermal, for example via an transdermal delivery system (TDS) such as a plaster containing the active ingredient or intranasal. For the production of such tablets, pills, semisolids, coated tablets, dragees and hard, e.g. gelatine, capsules the therapeutically useful product may be mixed with pharmaceutically inert, inorganic or organic excipients as are e.g. lactose, sucrose, glucose, gelatine, malt, silica gel, starch or derivatives thereof, talc, stearinic acid or their salts, dried skim milk, and the like. For the production of soft capsules one may use excipients as are e.g. vegetable, petroleum, animal or synthetic oils, wax, fat, polyols. For the production of liquid solutions, emulsions or suspensions or syrups one may use as excipients e.g. water, alcohols, aqueous saline, aqueous dextrose, polyols, glycerin, lipids, phospholipids, cyclodextrins, vegetable, petroleum, animal or synthetic oils. Especially preferred are lipids and more preferred are phospholipids (preferred of natural origin; especially preferred with a particle size between 300 to 350 nm) preferred in phosphate buffered saline (pH = 7 to 8, preferred 7.4). For suppositories one may use excipients as are e.g. vegetable, petroleum, animal or synthetic oils, wax, fat and polyols. For aerosol formulations one may use compressed gases suitable for this purpose, as are e.g. oxygen, nitrogen and carbon dioxide. The pharmaceutically useful agents may also contain additives for conservation, stabilization, e.g. UV stabilizers, emulsifiers, sweetener, aromatizers, salts to change the osmotic pressure, buffers, coating additives and antioxidants.

In general, in the case of oral or parenteral administration to adult humans weighing approximately 80 kg, a daily dosage of about 10 mg to about 10,000 mg, preferably from about 20 mg to about 1,000 mg, should be appropriate, although the upper limit may be exceeded when indicated. The daily dosage can be administered as a single

- 43 -

dose or in divided doses, or for parenteral administration, it may be given as continuous infusion or subcutaneous injection.

The compounds of the present invention can be prepared according to the following procedure:



The synthesis of the 4-sulfanyl-pyrrolidin-2-one scaffold is based on a four-component reaction (4CR) between a primary amine (II), an aldehyde or ketone (III) with maleic anhydride (IV), and a thiol (V). The reaction is preferably performed in toluene at reflux with a stoichiometric amount of the starting materials, according to J. Wei, J. T. Shaw *Org. Lett.* **2007**, *9*, 4077. The resulting 4-sulfanyl-pyrrolidin-2-one (VI) is formed in acceptable to good yields as a diastereoisomeric mixture. Generally, the two diastereoisomers are separated and isolated by preparative HPLC-chromatography. The final 4-sulfanyl-pyrrolidin-2-one amide of formula (I), (Ia), (Ic), (Id), (Ie) or (If) was obtained via aminolysis using amine (VIII) of the corresponding pentafluorophenyl esters of formula (VII) that were synthesized according to M. Bodanszky, A. Bodanszky, *The practice of Peptide Synthesis* 2nd Edition, p 102, Springer-Verlag Berlin Heidelberg New York (1994). These compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If) can be further modified such as the conversion into esters or salts from acids, salts from amines or by cleaving off protecting groups found in substituents R¹ to R⁶.

- 44 -

Further compounds of formula (I) wherein n is 0 can be prepared following the procedures described in:

- 1) M. R. Linder, J. Podlech, Organic Letters 2001, Vol. 3, No. 12, 1849-1851;
- 2) J. Podlech, M. R. Linder, J. Org. Chem. 1997, 62, 5873-5883;
- 5 3) J. Cesar, M. Sollner Dolenc, Tetrahedron Letters 42 (2001) 7099-7102.

The reaction procedures described herein may also be carried out in the presence of a chiral catalyst like e.g. proline-derived catalysts (as e.g. described in www.organic-chemistry.org/Highlights/2007/25March.shtm) in order to obtain the
10 corresponding enantiomerically pure compounds.

EXAMPLES

The present invention encompasses the following Examples:

15

Example 1

General procedure for the synthesis of 5-oxo-3-sulfanyl-pyrrolidin-3-carboxamides (I):

Maleic anhydride (IV, 1 mmol), a thiol (V, 1 mmol), aldehyde or ketone (III, 1 mmol)
20 and amine (II, 1 mmol) in toluene (8 mL) were heated to 150°C in a sealed tube for 24 hours. After cooled to room temperature, the solution was concentrated in vacuo. Purification on silica gel using an eluent (ethyl acetate: methanol = 9:1 to 1:1) yielded compounds of formula (VI) as a diastereoisomeric mixture. Afterwards, the two diastereoisomers were separated by preparative HPLC chromatography. Preparative
25 separations were usually performed with an acetonitrile-water eluent (+0.1 % formic acid) on a RP Polaris C18 column (length: 250 mm, diameter: 21 mm; particle size: 5 µm). Generally, good separations were observed (retention times of the two cis/trans diastereoisomers differed by 1 to 2 minutes) by using isocratic systems (70 % acetonitrile: 30 % water).

30

To a suspension of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride EDCI (1.5 mmol) in 8 mL ethyl acetate was added pentafluorophenol (3 mmol) at 0°C.

- 45 -

After 10 minutes, 5-oxo-pyrrolidine-3-carboxylic acid VI (1 mmol) was added at 0°C and the reaction mixture was stirred for 1 hour at room temperature. After evaporation of the solvent, the crude product was purified by chromatography on silica gel (ethyl acetate:hexane = 1:2) to yield the corresponding 5-oxo-pyrrolidine-3-carboxylic acid pentafluorophenyl ester VII as a colourless oil.

To a suspension of 5-oxo-pyrrolidine-3-carboxylic acid pentafluorophenyl ester VII (0.5 mmol) in 2 mL dry THF was added the desired amine VIII (0.5 mmol) at room temperature. The reaction mixture was stirred for 1 hour at room temperature.

Afterwards, 20 mL methylene chloride were added. The resulting organic layer was washed with 20 mL of a saturated aqueous solution of sodium hydrogencarbonate, dried over magnesium sulfate and the solvent was removed in vacuo. Finally, the crude product was purified by chromatography on silica gel with a suitable eluent to afford the desired 5-oxo-pyrrolidine-3-carboxamide I as a white solid.

15

Example 2

According to the general procedure in example 1, the following compounds were prepared:

- 2.1 cis-2-(6-chloro-1-methyl-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(pyridin-2-ylmethyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₀Cl₂N₄O₂S. Molecular Weight = 629.599. [M+H]⁺ observed = 629.1. Isolated yield 34.08 %.
- 2.2 trans-2-(6-chloro-1-methyl-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(pyridin-2-ylmethyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₀Cl₂N₄O₂S. Molecular Weight = 629.599. [M+H]⁺ observed = 629.1. Isolated yield 3.78 %.
- 2.3 trans-2-(6-chloro-1-methyl-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(thiophen-2-ylmethyl)pyrrolidine-3-carboxamide.

- 46 -

Molecular Formula = C₃₃H₂₉Cl₂N₃O₂S₂. Molecular Weight = 634.638. [M+H]⁺ observed = 656.0. Isolated yield 3.04 %.

5 **2.4** cis-2-(6-chloro-1-methyl-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(thiophen-2-ylmethyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₂₉Cl₂N₃O₂S₂. Molecular Weight = 634.638. [M+Na]⁺ observed = 656.0. Isolated yield 27.34 %.

10 **2.5** trans-2-(6-chloro-1-methyl-1H-indol-3-yl)-1-(4-chlorophenyl)-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(pyridin-2-ylmethyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₂₈Cl₂N₄O₂S. Molecular Weight = 615.572. [M+H]⁺ observed = 615.1. Isolated yield 2.78 %.

15 **2.6** cis-2-(6-chloro-1-methyl-1H-indol-3-yl)-1-(4-chlorophenyl)-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(pyridin-2-ylmethyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₂₈Cl₂N₄O₂S. Molecular Weight = 615.572. [M+H]⁺ observed = 615.1. Isolated yield 25.06 %.

20 **2.7** cis-2-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(pyridin-2-ylmethyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₂₈Cl₂N₄O₂S. Molecular Weight = 615.572. [M+H]⁺ observed = 615.2. Isolated yield 12.50 %.

25 **2.8** trans-2-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-5-oxo-N-(pyridin-2-ylmethyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₂₈Cl₂N₄O₂S. Molecular Weight = 615.572. [M+H]⁺ observed = 615.2. Isolated yield 4.56 %.

30 **2.9** trans-2-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄Cl₂N₄O₃S. Molecular Weight = 637.619. [M+H]⁺ observed = 637.2. Isolated yield 7.30 %.

2.10 cis-2-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄Cl₂N₄O₃S. Molecular Weight = 637.619. [M+H]⁺ observed = 637.2. Isolated
5 yield 7.18 %.

2.11 trans-2-(6-chloro-1H-indol-3-yl)-1-[(4-chloro-2-methylphenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆Cl₂N₄O₃S. Molecular Weight = 651.646. [M+H]⁺
10 observed = 651.2. Isolated yield 5.18 %.

2.12 cis-2-(6-chloro-1H-indol-3-yl)-1-[(4-chloro-2-methylphenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆Cl₂N₄O₃S. Molecular Weight = 651.646. [M+H]⁺
15 observed = 651.2. Isolated yield 7.74 %.

2.13 trans-2-(6-chloro-1H-indol-3-yl)-1-[(3-chlorophenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄Cl₂N₄O₃S. Molecular Weight = 637.619. [M+H]⁺ observed = 637.2. Isolated
20 yield 2.82 %.

2.14 cis-2-(6-chloro-1H-indol-3-yl)-1-[(3-chlorophenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄Cl₂N₄O₃S. Molecular Weight = 637.619. [M+H]⁺ observed = 637.2. Isolated
25 yield 3.81 %.

2.15 trans-2-(6-chloro-1H-indol-3-yl)-1-[(1R)-1-(4-chlorophenyl)ethyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆Cl₂N₄O₃S. Molecular Weight = 651.646. [M+H]⁺
30 observed = 651.2. Isolated yield 2.65 %.

2.16 cis-2-(6-chloro-1H-indol-3-yl)-1-[(1R)-1-(4-chlorophenyl)ethyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆Cl₂N₄O₃S. Molecular Weight = 651.646. [M+H]⁺ observed = 651.2. Isolated yield 1.36 %.

5

2.17 trans-2-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-chlorophenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₃Cl₃N₄O₃S. Molecular Weight = 672.064. [M+H]⁺ observed = 671.1. Isolated yield 6.65 %.

10

2.18 cis-2-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-chlorophenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₃Cl₃N₄O₃S. Molecular Weight = 672.064. [M+H]⁺ observed = 673.1. Isolated yield 19.04 %.

15

2.19 trans-1-benzyl-2-(6-chloro-1H-indol-3-yl)-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₅ClN₄O₃S. Molecular Weight = 603.174. [M+H]⁺ observed = 603.0. Isolated yield 4.31 %.

2.20 cis-1-benzyl-2-(6-chloro-1H-indol-3-yl)-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₅ClN₄O₃S. Molecular Weight = 603.174. [M+H]⁺ observed = 603.0. Isolated yield 4.59 %.

2.21 trans-2-(6-chloro-1H-indol-3-yl)-1-[(1S)-1-(4-chlorophenyl)ethyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆Cl₂N₄O₃S. Molecular Weight = 651.646. [M+H]⁺ observed = 650.9. Isolated yield 4.91 %.

2.22 cis-2-(6-chloro-1H-indol-3-yl)-1-[(1S)-1-(4-chlorophenyl)ethyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆Cl₂N₄O₃S. Molecular Weight = 651.646. [M+H]⁺ observed = 651.0. Isolated yield 4.86 %.

30

2.23 trans-2-(6-bromo-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆BrClN₄O₃S. Molecular Weight = 696.097. [M+H]⁺

5 observed = 697.1. Isolated yield 9.27%.

2.24 cis-2-(6-bromo-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₄H₃₆BrClN₄O₃S. Molecular Weight = 696.097. [M+H]⁺

10 observed = 697.0. Isolated yield 11.14%.

2.25 trans-2-(5-bromo-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄BrClN₄O₃S. Molecular Weight = 682.07. [M+H]⁺ observed = 682.9. Isolated

15 yield 3.71 %.

2.26 cis-2-(5-bromo-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄BrClN₄O₃S. Molecular Weight = 682.07. [M+H]⁺ observed = 682.8. Isolated

20 yield 4.65 %.

2.27 cis-2-(6-chloro-1H-indol-3-yl)-1-[(6-chloropyridin-3-yl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₂H₃₃Cl₂N₅O₃S. Molecular Weight = 638.607. [M+H]⁺ observed = 638.0. Isolated

25 yield 3.76 %.

2.28 trans-2-(6-chloro-1H-indol-3-yl)-1-[(6-chloropyridin-3-yl)methyl]-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide.

Molecular Formula = C₃₂H₃₃Cl₂N₅O₃S. Molecular Weight = 638.607. Isolated yield

30 1.46 %.

2.29 trans-4- {[2-(6-chloro-1H-indol-3-yl)-1-[(1S)-1-(4-chlorophenyl)ethyl]-5-oxo-3-(phenylsulfanyl)pyrrolidin-3-yl]carbonyl}piperazin-2-one. Molecular Formula = C₃₁H₂₈Cl₂N₄O₃S. Molecular Weight = 607.55. [M+H]⁺ observed = 607.2. Isolated yield 4.44 %.

5

2.30 cis-4- {[2-(6-chloro-1H-indol-3-yl)-1-[(1S)-1-(4-chlorophenyl)ethyl]-5-oxo-3-(phenylsulfanyl)pyrrolidin-3-yl]carbonyl}piperazin-2-one. Molecular Formula = C₃₁H₂₈Cl₂N₄O₃S. Molecular Weight = 607.55. [M+H]⁺ observed = 607.2. Isolated yield 4.09 %.

10

2.31 trans-4- {[2-(6-chloro-1H-indol-3-yl)-1-[(1R)-1-(4-chlorophenyl)ethyl]-5-oxo-3-(phenylsulfanyl)pyrrolidin-3-yl]carbonyl}piperazin-2-one. Molecular Formula = C₃₁H₂₈Cl₂N₄O₃S. Molecular Weight = 607.55. [M+H]⁺ observed = 608.8. Isolated yield 2.62 %.

15

2.32 cis-4- {[2-(6-chloro-1H-indol-3-yl)-1-[(1R)-1-(4-chlorophenyl)ethyl]-5-oxo-3-(phenylsulfanyl)pyrrolidin-3-yl]carbonyl}piperazin-2-one. Molecular Formula = C₃₁H₂₈Cl₂N₄O₃S. Molecular Weight = 607.55. [M+H]⁺ observed = 606.9. Isolated yield 1.89 %.

20

2.33 trans-1-[(4-chlorophenyl)methyl]-2-(6-fluoro-1H-indol-3-yl)-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄ClFN₄O₃S. Molecular Weight = 621.164. [M+H]⁺ observed = 621.0. Isolated yield 2.98 %.

25

2.34 cis-1-[(4-chlorophenyl)methyl]-2-(6-fluoro-1H-indol-3-yl)-N-[3-(morpholin-4-yl)propyl]-5-oxo-3-(phenylsulfanyl)pyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄ClFN₄O₃S. Molecular Weight = 621.164. [M+H]⁺ observed = 621.0. Isolated yield 4.72 %.

30

2.35 cis-4- {[2-(6-bromo-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-5-oxopyrrolidin-3-yl]carbonyl}piperazin-2-one. Molecular

Formula = C₃₁H₂₈BrClN₄O₃S. Molecular Weight = 652.001. [M+H]⁺ observed = 651.3. Isolated yield 8.00 %.

5 **2.36** cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₆BrClN₄O₃S. Molecular Weight = 696.097. [M+H]⁺ observed = 697.2. Isolated yield 8.39 %.

10 **2.37** trans-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₆BrClN₄O₃S. Molecular Weight = 696.097. [M+H]⁺ observed = 697.1. Isolated yield 3.45 %.

15 **2.38** cis-2-(6-bromo-1H-indol-3-yl)-1-[(4-bromophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₆Br₂N₄O₃S. Molecular Weight = 740.548. [M+H]⁺ observed = 740.5. Isolated yield 10.00 %.

20 **2.39** cis-2-(6-chloro-1H-indol-3-yl)-1-[(4-chloro-3-fluorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₅Cl₂FN₄O₃S. Molecular Weight = 669.636. [M+H]⁺ observed = 669.1. Isolated yield 14.85 %.

25 **2.40** trans-2-(6-chloro-1H-indol-3-yl)-1-[(4-chloro-3-fluorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₅Cl₂FN₄O₃S. Molecular Weight = 669.636. [M+H]⁺ observed = 669.0. Isolated yield 4.48 %.

30 **2.41** cis-2-(6-chloro-5-fluoro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₅Cl₂FN₄O₃S. Molecular Weight = 669.636. [M+H]⁺ observed = 669.1. Isolated yield 4.88 %.

2.42 trans-2-(6-chloro-5-fluoro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₅Cl₂FN₄O₃S. Molecular Weight = 669.636. [M+H]⁺

5 observed = 669.1. Isolated yield 1.08 %.

2.43 cis-5-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-4-{[4-(2-hydroxyethyl)piperazin-1-yl]carbonyl}-4-[(4-methylphenyl)sulfanyl]pyrrolidin-2-one.
Molecular Formula = C₃₃H₃₄Cl₂N₄O₃S. Molecular Weight = 637.619. [M+H]⁺

10 observed = 637.0. Isolated yield 7.67 %.

2.44 cis-2-(6-chloro-1H-indazol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₃H₃₅Cl₂N₅O₃S. Molecular Weight = 652.634. [M+H]⁺

15 observed = 652.0. Isolated yield 6.04 %.

2.45 cis-2-(6-chloro-1H-indol-3-yl)-1-[(4-chlorophenyl)methyl]-3-[(4-methylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₄H₃₆Cl₂N₄O₃S. Molecular Weight = 651.646. [M+H]⁺

20 observed = 651.1. Isolated yield 7.97 %.

2.46 cis-1-[(4-bromophenyl)methyl]-5-(6-chloro-1H-indol-3-yl)-4-{[4-(2-hydroxyethyl)piperazin-1-yl]carbonyl}-4-[(4-methylphenyl)sulfanyl]pyrrolidin-2-one.
Molecular Formula = C₃₃H₃₄BrClN₄O₃S. Molecular Weight = 682.07. [M+H]⁺

25 observed = 683.0. Isolated yield 5.35 %.

2.47 cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-N-(2,3-dihydroxypropyl)-3-[(4-methylphenyl)sulfanyl]-5-oxopyrrolidine-3-carboxamide.
Molecular Formula = C₃₀H₂₉BrClN₃O₄S. Molecular Weight = 642.991. [M+Na]⁺

30 observed = 668.1. Isolated yield 6.92 %.

2.48 cis-4- {[1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(4-methylphenyl)sulfanyl]-5-oxopyrrolidin-3-yl]carbonyl}piperazin-2-one. Molecular Formula = C₃₁H₂₈BrClN₄O₃S. Molecular Weight = 652.001. [M+H]⁺ observed = 653.1. [M+Na]⁺ observed = 675.3. Isolated yield 3.39 %.

5

2.49 cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(4-methylphenyl)sulfanyl]-N-[2-(morpholin-4-yl)ethyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄BrClN₄O₃S. Molecular Weight = 682.07. [M+H]⁺ observed = 683.1. Isolated yield 7.30 %.

10

2.50 cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(4-methylphenyl)sulfanyl]-N-[4-(morpholin-4-yl)butyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₅H₃₈BrClN₄O₃S. Molecular Weight = 710.123. [M+H]⁺ observed = 711.0. Isolated yield 7.08 %.

15

2.50 cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-N-(4-hydroxybutyl)-3-[(4-methylphenyl)sulfanyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₁H₃₁BrClN₃O₃S. Molecular Weight = 641.018. [M+Na]⁺ observed = 664.2. Isolated yield 7.32 %.

20

2.51 cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(4-methylphenyl)sulfanyl]-N-(1-methylpiperidin-4-yl)-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₄BrClN₄O₂S. Molecular Weight = 666.071. [M+H]⁺ observed = 668.2. Isolated yield 7.51 %.

25

2.52 cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(4-ethylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₅H₃₈BrClN₄O₃S. Molecular Weight = 710.123. [M+H]⁺ observed = 711.1. Isolated yield 17.16 %.

30

2.53 cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(3,4-dimethylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-

carboxamide. Molecular Formula = C₃₅H₃₈BrClN₄O₃S. Molecular Weight = 710.123. [M+H]⁺ observed = 711.0. Isolated yield 9.74 %.

5 **2.52** cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-3-[(2,4-dimethylphenyl)sulfanyl]-N-[3-(morpholin-4-yl)propyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₅H₃₈BrClN₄O₃S. Molecular Weight = 710.123. [M+H]⁺ observed = 711.0. Isolated yield 6.38 %.

10 **2.53** trans-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-N-[3-(morpholin-4-yl)propyl]-3-[(4-nitrophenyl)sulfanyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₃BrClN₅O₅S. Molecular Weight = 727.068. [M+H]⁺ observed = 726.0. Isolated yield 3.00 %.

15 **2.54** cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-N-[3-(morpholin-4-yl)propyl]-3-[(4-nitrophenyl)sulfanyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₃H₃₃BrClN₅O₅S. Molecular Weight = 727.068. [M+H]⁺ observed = 727.9. Isolated yield 0.41 %.

20 **2.55** cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-N-[(2R)-1-hydroxypropan-2-yl]-3-[(4-methylphenyl)sulfanyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₀H₂₉BrClN₃O₃S. Molecular Weight = 626.992. [M+Na]⁺ observed = 650.6. Isolated yield 8.24 %.

25 **2.56** cis-1-[(4-bromophenyl)methyl]-2-(6-chloro-1H-indol-3-yl)-N-[(2S)-1-hydroxypropan-2-yl]-3-[(4-methylphenyl)sulfanyl]-5-oxopyrrolidine-3-carboxamide. Molecular Formula = C₃₀H₂₉BrClN₃O₃S. Molecular Weight = 626.992. [M+Na]⁺ observed = 650.2. Isolated yield 8.04 %.


Example 3

30

1) Synthesis of 4-[5-oxo-pyrrolidine-3-carbonyl]-piperazine-1-carboxylic acid ethylamide compounds

Synthesis of (cis)-4-[1-(4-Bromo-benzyl)-2-(6-chloro-1H-indol-3-yl)-5-oxo-3-p-tolylsulfanyl-pyrrolidine-3-carbonyl]-piperazine-1-carboxylic acid ethylamide [PXN727-d1]

5



Multicomponent reaction (step 1):

Maleic anhydride **2** (6 mmol), thiol **4** (6 mmol), aldehyde **3** (6 mmol) and amine **1** (6 mmol) in toluene (50 mL) were heated to 150 °C under Dean-Stark conditions for 24 hours. After being cooled to room temperature, the solution was concentrated in vacuo. Purification on silica gel (ethyl acetate: methanol = 9:1 to 1:1) yielded **5** as a diastereoisomeric mixture (1.48 g, 46 %).

Literature: J. Wei, J. T. Shaw *Org. Lett.* **2007**, *9*, 4077.

Separation of the diastereoisomeric mixture:

- 56 -

The above described reaction sequence yielded two diastereoisomers **d1** and **d2** in a 50:50 ratio. They were separated by preparative HPLC chromatography using the following conditions:

- Column RP Polaris C18 (length: 250 mm, Ø: 21 mm; particle size: 5 µm).
- 5 - Isocratic elution (70 % acetonitrile: 30 % water, 0.1% HCOOH), 21mL/min, Rt=7.62min.

The separation can also be performed with methanol/water mixtures.

- 10 Concentration of the solution in vacuo yielded the desired pure diastereoisomer **6** (*cis*, **d1**) as a light yellow solid (528.9 mg, 33.6 %).

Overall yield for the preparation of **6**: 15.47 % (MCR and isolation of the *cis*-isomer **d1**)

15 **Pentafluorophenyl ester formation:**

- To a suspension of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride EDCI (267 mg, 1.4 mmol) in 8 mL ethyl acetate was added pentafluorophenol (512 mg, 2.8 mmol) at 0 °C. After 10 minutes, compound **6** (528.9 mg, 0.9 mmol) was added at 0 °C and the reaction mixture was
- 20 stirred for 1 hour at room temperature. After evaporation of the solvent, the crude product was purified by chromatography on silica gel (ethyl acetate:hexane = 1:2 → 1:1) to yield the corresponding pentafluorophenyl ester **7** as a colourless oil (632.0 mg, 92.5%).

25 **Literature:**

M. Bodanszky, A. Bodanszky, The practice of Peptide Synthesis 2nd Edition, p 102, Springer-Verlag Berlin Heidelberg New York (1994).

Amide coupling:

- 30 To a suspension of pentafluorophenyl ester **7** (1.3 g, 1.8 mmol) in 16 mL dry THF was added piperazine (7.2 mmol) at room temperature. The reaction mixture was stirred for 20 hours at room temperature. Afterwards, 20 mL

- 57 -

methylene chloride were added. The resulting organic layer was washed with 20 mL of a saturated aqueous solution of sodium hydrogenocarbonate, dried over magnesium sulfate and the solvent was removed in vacuo. Finally, the crude product was purified by chromatography on silica gel (ethyl acetate:methanol 9:1 → 1:1) to afford the desired piperazine amide **8** as a white solid (977.8 mg, 84.20 %).

Reaction with ethyl isocyanate:

To a solution of compound **8** (848.3 mg, 1.3 mmol) in 15 mL THF extra dry was added ethyl isocyanate (283.6 mg, 4 mmol) at -30°C. After 1h of stirring at -30°C, 20 mL methylene chloride were added. The resulting organic layer was washed with 20 mL of a saturated aqueous solution of sodium hydrogenocarbonate, dried over magnesium sulfate and the solvent was removed in vacuo. Finally, the crude product was purified by chromatography on silica gel with the system ethyl acetate:methanol 19:1 to yield **PXN727-d1** as a white solid (853.4 mg, 90.5 %).

mp = 263.7-267.2 °C

¹H-NMR (400 MHz, DMSO) δ 11.57 (s, 1H), 7.57 (s, 1H), 7.50 (s, 1H), 7.44 (d, 2H, J = 6.70 Hz), 7.25 (d, 1 H, J = 7.39 Hz), 7.09 (m, 5 H), 6.90 (d, 2H, J = 7.39 Hz), 6.53 (s, 1 H), 4.93 (s, 1H), 4.74 (d, 1H, J = 15.29 Hz), 3.84 (s, 2 H), 3.60-3.20 (m, 4H), 3.45 (d, 1H, J = 15.29), 3.15-2.80 (m, 4H), 3.09-3.06 (m, 2 H), 2.25 (s, 3H), 1.03 (t, 3H, J = 7.05 Hz).


IR: 3397, 3174, 2923, 1674, 1625, 1535, 1487, 1401, 1361, 1241, 1174, 1118, 1002, 794.

MS (+ESI): m/z = 709.9 [M+H], 730.2 [M+Na].


Overall yield (four preparative steps and diastereoisomer separation): **10.91 %**

Example 4**2) Replacement of the sulfur S (group X) by methylene CH₂**

Synthesis of (cis)4-[3-Benzyl-1-(4-chloro-3-fluoro-benzyl)-2-(6-chloro-1H-indol-3-yl)-5-oxo-pyrrolidine-3-carbonyl]-piperazine-1-carboxylic acid ethylamide [PXN790-d1]



Synthesis of α -benzylsuccinic anhydride 10



The commercially available α -benzylsuccinic acid **9** (1g, 4,8 mmol) was refluxed for 1h in 30 mL trifluoroacetic anhydride. Afterwards, the solvent was removed in vacuo. The crude residue was washed with cold hexane to yield α -benzylsuccinic anhydride **10** as a white solid (858.2mg, 93.95 %).

15

Multicomponent reaction

Alpha-benzylsuccinic anhydride **10** (850 mg, 4.5 mmol) , aldehyde **3** (1 mmol) and amine **11** (1 mmol) in toluene (16 mL) were heated to 150 °C in a sealed tube for 24 hours. After cooled to room temperature, the solution was concentrated in vacuo. Purification on silica gel (ethyl acetate: methanol = 9:1 to 1:1) yielded MCR-product **12** as a diastereoisomeric mixture (210.3 mg, 9.20 %).

Pentafluorophenyl ester formation

To a suspension of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride EDCI (118.2 mg, 0.617 mmol) in 5 mL ethyl acetate was added pentafluorophenol (227.1 mg, 1.23 mmol) at 0 °C. After 10 minutes, 5-oxo-pyrrolidine-3-carboxylic acid **12** (210.3 mg, 0.411 mmol) was added at 0 °C and the reaction mixture was stirred for 1 hour at room temperature. After evaporation of the solvent, the crude product was purified by chromatography on silica gel (ethyl acetate:hexane = 1:2 → 1:1) to yield the corresponding (cis)-5-oxo-pyrrolidine-3-carboxylic acid pentafluorophenyl ester **13** as a colourless oil (78.2 mg, 28.9 %).

Amide coupling

To a suspension of (cis)-5-oxo-pyrrolidine-3-carboxylic acid pentafluorophenyl ester **13** (78.2 mg, 0.1154 mmol) in 2 mL THF extra dry was added piperazine (39.8 mg, 0.4616 mmol) at room temperature. The reaction mixture was stirred for 10 hours at room temperature. Afterwards, 20 mL methylene chloride were added. The resulting organic layer was washed with 20 mL of a saturated aqueous solution of sodium hydrogenocarbonate, dried over magnesium sulfate and the solvent was removed in vacuo. Finally, the crude product was purified by chromatography on silica gel (ethyl acetate:methanol 9:1 → 1:1) to afford the desired (cis)-4-(piperazine-1-carbonyl)-pyrrolidin-2-one **14** as a white solid (38.5 mg, 57.55 %).

- 60 -

Reaction with ethyl isocyanate

To a solution of compound **14** (38.5 mg, 0,066 mmol) in 3 mL THF extra dry was added ethyl isocyanate (14.2 mg, 0.199 mmol) at -30°C. After 1h of stirring at -30°C, 20 mL methylene chloride were added. The resulting organic layer was washed with 20 mL of a saturated aqueous solution of sodium hydrogenocarbonate, dried over magnesium sulfate and the solvent was removed in vacuo. Finally, the crude product was crystallized from ethyl acetate:methanol 19:1 to yield **PXN790-d1** as a white solid (26.9 mg, 62.24 %).

¹H-NMR (400 MHz, DMSO) δ 11.57 (s, 1H), 7.58-7.43 (m, 3H), 7.28-7.11 (m, 5H), 6.95-6.83 (m, 4H), 6.55 (s, 1H), 4.97-4.91 (m, 1H), 4.71 (d, 1H, J = 15.24 Hz), 3.63-3.57 (m, 5H), 3.10-3.08 (m, 3H), 2.91-2.85 (m, 3H), 2.69-2.65 (m, 1H), 1.03 (t, 3H, J = 7.02 Hz).

IR: 3043, 3165, 3033, 2964, 2930, 1677, 1615, 1538, 1449, 1401, 1262, 1240, 1207, 1119, 796, 698.


MS (+ESI): m/z = 650.1 [M+H], 672.1 [M+Na].

Overall yield (four preparative steps): **0.93 %**

Example 5**Replacement of the sulfur S (group X) by oxygene O**


Synthesis of (cis)-4-[1-(4-Chloro-3-fluoro-benzyl)-2-(6-chloro-1H-indol-3-yl)-5-oxo-3-p-tolyloxy-pyrrolidine-3-carbonyl]-piperazine-1-carboxylic acid ethylamide [**PXN789-d1**]

- 61 -



Synthesis of the oxo-substituted anhydride

5 The oxo-substituted anhydride was obtained via a three-step synthesis:



10 1-Phenyl-2,5-dihydro-1H-pyrrole-2,5-dione **19** (5.19 g, 3 mmol), p-cresol **20** (3.24 g, 3 mmol), and triethylamine (3.03 g, 3 mmol) were added in 20 ml toluene extra dry and heated at 100°C for 6h. Afterwards, the mixture was cooled to 0°C. The precipitated solid was filtered and washed with cold

toluene and hexane to yield compound **21** as a purple solid (2.895 g, 34.30 %).

Analytic data for Compound 21:

5 ¹H NMR (DMSO, 399.83 MHz): 2.26 (s, 3H), 2.89-2.94 (m, 1H), 3.31-3.46 (m, 1H), 5.44-5.47 (m, 1H), 6.97 (d, 2H, J=8.4 Hz), 7.14 (d, 2H, J=7.6 Hz), 7.33 (d, 2H, J=7.2 Hz), 7.44-7.53 (m, 3H).

MS (+ESI): *m/z*= 282 [M+H].

10 Compound **21** (610.3 mg, 2.17 mmol) was dissolved in 30 ml of a mixture of aqueous HCl 37%:HCOOH 1:1. The mixture was heated for 3h at 110°C. Afterwards, the mixture was cooled to room temperature and the aqueous phase was washed 3 times with DCM and then evaporated. The resulting solid was washed 3 times with cold ether and the resulting ether phase
15 evaporated to yield the succinic acid **22** as a white solid. Finally, the succinic acid **22** was solved in 10ml of trifluoroacetic anhydride (TFAA) and heated for 6h at 100°C. Then TFAA was evaporated and the resulting solid was washed with cold hexane to yield the corresponding succinic anhydride **15** as a white solid (170.4 mg, 95.56 %).

20

Analytic data for compound 15 :

¹H NMR (DMSO, 399.43 MHz): 2.25 (s, 3H), 3.21–3.27 (m, 1H), 3.52–3.59 (m, 1H), 5.57–5.61 (m, 1H), 6.92 (d, 2H, J=8.26 Hz), 7.14 (d, 2H, J=8.22 Hz).

25 IR: 3001, 2920, 1865, 1781, 1608, 1508, 1396, 1213, 1178, 1086, 1021, 903, 806.

MS (+ESI): *m/z*=207[M+H].

Multicomponent reaction

30 First, aldehyde **3** (646.6 mg, 3.6 mmol) and amine **11** (478.8 mg, 3 mmol) were condensed in 3 mL trimethylorthoformate for 10 hours at room temperature. Then, the solvent was removed in vacuo and the residue was

- 63 -

solved in 25 mL o-xylene. Afterwards, succinic anhydride **15** (850 mg, 4.5 mmol) was added and the mixture was heated to 150 °C for 24 hours under Dean-Stark conditions. After cooled to room temperature, the solution was concentrated in vacuo. Purification on silica gel (ethyl acetate: methanol =
5 9:1 → 1:1) yielded MCR-product **16** as a diastereoisomeric mixture (33.9 mg, 2.11 %).

Pentafluorophenyl ester formation

To a suspension of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
10 hydrochloride EDCI (18.5 mg, 0.096 mmol) in 2 mL ethyl acetate was added pentafluorophenol (35.6 mg, 0.193 mmol) at 0 °C. After 10 minutes, 5-oxo-pyrrolidine-3-carboxylic acid **16** (33.9 mg, 0.064 mmol) was added at 0 °C and the reaction mixture was stirred for 1 hour at room temperature. After
15 evaporation of the solvent, the crude product was purified by chromatography on silica gel (ethyl acetate:hexane = 1:2) to yield the corresponding 5-oxo-pyrrolidine-3-carboxylic acid pentafluorophenyl ester **17** as a colourless oil (40.1 mg, 89.80 %).

Amide coupling

20 To a suspension of 5-oxo-pyrrolidine-3-carboxylic acid pentafluorophenyl ester **17** (40.1 mg, 0.0578 mmol) in 2 mL THF extra dry was added piperazine (19.9 mg, 0.231 mmol) at room temperature. The reaction mixture was stirred for 10 hours at room temperature. Afterwards, 20 mL methylene
25 chloride were added. The resulting organic layer was washed with 20 mL of a saturated aqueous solution of sodium hydrogenocarbonate, dried over magnesium sulfate and the solvent was removed in vacuo. Finally, the crude product was purified by chromatography on silica gel (ethyl acetate:methanol 9:1 → 1:1) to afford the desired 4-(piperazine-1-carbonyl)-pyrrolidin-2-one
30 **18** as a white solid (12.2 mg, 45.91 %).

Reaction with ethyl isocyanate

- 64 -

To a solution of compound **18** (12.2 mg, 0.0204 mmol) in 3 mL THF extra dry was added ethyl isocyanate (4.4 mg, 0.0612 mmol) at -30°C. After 1h of stirring at -30°C, 20 mL methylene chloride were added. The resulting organic layer was washed with 20 mL of a saturated aqueous solution of sodium hydrogenocarbonate, dried over magnesium sulfate and the solvent was removed in vacuo. Finally, the crude product was purified by chromatography on silica gel (methylene chloride:methanol 95:5) to yield **PXN789-d1** as a yellow solid (9.6 mg, 70.60 %).


MS (+ESI): m/z =666.1[M+H].

Overall yield (four preparative steps): 0.61 %

Further examples

Further examples which have been prepared according to one of the procedures described above: All products were obtained and tested as racemates. The cellular activity was measured on p53 wild type ovarian teratocarcinoma cells (PA-1) and measured IC₅₀ are given in micromolar. CCA is the abbreviation of Cell Cycle Arrest.


PXN610



C₂₈H₂₈Cl₂N₂O₃S; MW: 543.52; found (HPLC MS):
[M+H⁺] = 543.0; Yield: 47 %; IC₅₀=15


PXN611

- 65 -



$C_{29}H_{31}Cl_2N_3O_2S$; MW: 556.56; found (HPLC MS):
 $[M+H^+] = 556.1$; Yield: 28 %; $IC_{50}=8.3$


PXN612



5

$C_{27}H_{26}Cl_2N_2O_3S$; MW: 529.49; found (HPLC MS):
 $[M+H^+] = 529.0$; Yield: 17 %; $IC_{50}=10.3$

PXN613




10

$C_{28}H_{29}Cl_2N_3O_2S$; MW: 542.53; found (HPLC MS):
 $[M+H^+] = 542.0$; Yield: 13 %; $IC_{50}=8$


PXN617

- 66 -



$C_{30}H_{29}Cl_2N_3O_3S$; MW: 582.55; found (HPLC MS): $[M+H^+] = 582.1$; $[M+Na^+] = 604.0$; Yield: 32 %; $IC_{50}=3.1$


PXN618



5

$C_{31}H_{32}Cl_2N_4O_2S$; MW: 595.60; found (HPLC MS): $[M+H^+] = 595.1$; Yield: 35 %; $IC_{50}=4.2$

PXN619




10

$C_{31}H_{31}Cl_2N_3O_3S$; MW: 596.58; found (HPLC MS): $[M+Na^+] = 618.1$; Yield: 40 %; $IC_{50}=28.7$

PXN620


- 67 -



$C_{30}H_{29}Cl_2N_3O_3S$; MW: 582.55; found (HPLC MS):

$[M+H^+] = 581.9$; Yield: 15 %; $IC_{50}=15.3$

PXN623




5

$C_{32}H_{36}Cl_2N_4O_2S$; MW: 611.64; found (HPLC MS): $[M+H^+]$

$= 611.1$; Yield: 49 %; $IC_{50}=7.8$

PXN624




10

$C_{31}H_{27}Cl_2N_3O_2S$; MW: 576.55; found (HPLC MS): $[M+H^+]$

$= 576.0$; Yield: 41 %; $IC_{50}=22.4$


PXN625

- 68 -



$C_{30}H_{26}Cl_2N_2O_2S_2$; MW: 581.59; found (HPLC MS): $[M+H^+]$
= 581.1; Yield: 37 %; $IC_{50}>60$


PXN626



5

$C_{31}H_{34}Cl_2N_4O_2S$; MW: 597.61; found (HPLC MS):
 $[M+H^+] = 597.1$; Yield: 21 %; $IC_{50}=7.4$

PXN627




10

$C_{30}H_{25}Cl_2N_3O_2S$; MW: 562.52; found (HPLC MS): $[M+H^+]$
= 562.0; Yield: 17 %; $IC_{50}=10.2$

PXN628


- 69 -



$C_{29}H_{24}Cl_2N_2O_2S_2$; MW: 567.56; found (HPLC MS): $[M+H^+]$

= 567.0; Yield: 20 %; $IC_{50} > 60$


PXN629



5

$C_{35}H_{39}Cl_2N_5O_2S$; MW: 664.70; found (HPLC MS): $[M+H^+] = 664.2$; Yield: 40 %; $IC_{50} = 7.2$

PXN630-d1




10

$C_{34}H_{30}Cl_2N_4O_2S$; MW: 629.61; found (HPLC MS): $[M+H^+] = 629.1$; Yield: 38 %; $IC_{50} > 60$


PXN631-d1

- 70 -



$C_{33}H_{29}Cl_2N_3O_2S_2$; MW: 634.65; found (HPLC MS):
 $[M+Na^+] = 656.0$; Yield: 30 %; $IC_{50} > 60$


PXN632



5

$C_{34}H_{37}Cl_2N_5O_2S$; MW: 650.68; found (HPLC MS):
 $[M+H^+] = 650.2$; Yield: 36 %; $IC_{50} = 5.2$

PXN633-d1




10

$C_{33}H_{28}Cl_2N_4O_2S$; MW: 615.59; found (HPLC MS):
 $[M+H^+] = 615.1$; Yield: 17 %; $IC_{50} = 23.3$


PXN633-d2

- 71 -



$C_{33}H_{28}Cl_2N_4O_2S$; MW: 615.59; found (HPLC MS):
 $[M+H^+] = 615.1$; Yield: 11 %; $IC_{50}=22.2$


PXN634



5

$C_{32}H_{27}Cl_2N_3O_2S_2$; MW: 620.62; found (HPLC MS):
 $[M+H^+] = 620.0$; Yield: 27 %; $IC_{50}>60$

PXN635




$C_{23}H_{26}Cl_2N_2O_3S$; MW: 481.45

10


PXN636

- 72 -



$C_{27}H_{25}Cl_3N_2O_3S$; MW: 563.93; found (HPLC MS): $[M+H]^+ = 563.0$; Yield: 26 %; $IC_{50} > 60$


PXN637



5

$C_{30}H_{24}Cl_3N_3O_2S$; MW: 596.97; found (HPLC MS): $[M+H]^+ = 596.0$; Yield: 24 %; $IC_{50} > 60$

PXN638




10

$C_{30}H_{29}Cl_2N_3O_5S$; MW: 614.55; found (HPLC MS): $[M+H]^+ = 614.1$; Yield: 11 %; $IC_{50} = 60.4$


PXN639

- 73 -



$C_{33}H_{28}Cl_2N_4O_4S$; MW: 647.59; found (HPLC MS):
 $[M+H^+] = 647.1$; Yield: 5 %; $IC_{50}=65.7$


PXN640-d1



5

$C_{33}H_{28}Cl_2N_4O_2S$; MW: 615.59; found (HPLC MS):
 $[M+H^+] = 615.2$; Yield: 13 %; $IC_{50}=18.7$

PXN640-d2




10

$C_{33}H_{28}Cl_2N_4O_2S$; MW: 615.59; found (HPLC MS):
 $[M+H^+] = 615.2$; Yield: 5 %; $IC_{50}=24.5$

PXN641


- 74 -



$C_{34}H_{36}Cl_2N_4O_3S$; MW: 651.66; found (HPLC MS):

$[M+H^+] = 651.2$; Yield: 9 %; $IC_{50}=9.4$

PXN642




5

$C_{29}H_{27}Cl_2N_3O_2S$; MW: 552.53; found (HPLC MS): $[M+H^+] =$

552.1; $[M+Na^+] = 574.1$; Yield: 8 %; $IC_{50}=5.7$

PXN643




10

$C_{31}H_{28}Cl_2N_4O_3S$; MW: 607.56; found (HPLC MS):

$[M+H^+] = 607.2$; Yield: 4 %; $IC_{50}=3.9$


PXN644

- 75 -



$C_{28}H_{25}Cl_2N_3O_3S$; MW: 554.50; found (HPLC MS):
 $[M+Na^+] = 576.1$; Yield: 10 %; $IC_{50}=13.5$


PXN645



5

$C_{27}H_{31}Cl_2N_3O_3S$; MW: 548.54; found (HPLC MS):
 $[M+Na^+] = 570.2$; Yield: 1 %


PXN646



10


$C_{26}H_{24}Cl_2N_2O_3S$; MW: 515.46; found (HPLC MS):
 $[M+H^+] = 515.0$; Yield: 11 %; $IC_{50}=14.6$

PXN647




$C_{25}H_{30}Cl_2N_2O_3S$; MW: 509.50


PXN649

 $C_{26}H_{24}Cl_2N_2O_3S$; MW: 515.46

5 PXN650

 $C_{28}H_{25}Cl_2N_3O_3S$; MW: 554.50; found (HPLC MS): $[M+Na^+] = 576.1$; Yield: 8 %; $IC_{50}=17.1$

PXN651




10

 $C_{32}H_{32}Cl_2N_4O_3S$; MW: 623.61; found (HPLC MS): $[M+H^+] = 623.2$; Yield: 9 %; $IC_{50}=9.2$

PXN652

PXM653




$\text{C}_{31}\text{H}_{31}\text{Cl}_2\text{N}_4\text{O}_4\text{S}$; MW: 625.58; found (HPLC MS):
 $[\text{M}+\text{H}^+] = 625.1$; Yield: 2 %

O=C1NC(=O)C2(C(=O)Nc3ccc(Cl)cc3)C(=O)N(C2C(=O)c4ccc(Cl)cc4)C1Sc5ccccc5NCCCN6CCOCC6


$\text{C}_{31}\text{H}_{30}\text{Cl}_2\text{N}_4\text{O}_4\text{S}$; MW: 625.58; found (HPLC MS):
 $[\text{M}+\text{H}^+] = 625.1$; Yield: 3 %; $\text{IC}_{50}=29.9$

PXM655

- 78 -


 $C_{29}H_{34}Cl_2N_4O_4S$; MW: 605.59


PXN656


 $C_{31}H_{38}Cl_2N_4O_3S$; MW: 617.64; found (HPLC MS):

 $[M+H^+] = 617.2$; Yield: 1 %; $IC_{50}=17.8$

5

PXN657



 $C_{28}H_{26}ClN_3O_4S$; MW: 536.05; found (HPLC MS):

 $[M+H^+] = 536.1$; Yield: 11 %


10

PXN658

- 79 -



 $C_{32}H_{32}Cl_2N_4O_4S$; MW: 639.61

PXN659-d1


 $C_{33}H_{34}Cl_2N_4O_3S$; MW: 637.63; found (HPLC MS): $[M+H^+] = 637.2$; Yield: 7 %; $IC_{50}=6.8$

5


PXN659-d2


 $C_{33}H_{34}Cl_2N_4O_3S$; MW: 637.63; found (HPLC MS): $[M+H^+] = 637.2$; Yield: 7 %; $IC_{50}=12$

10


PXN660-d1

- 80 -




C₃₄H₃₆Cl₂N₄O₃S; MW: 651.66; found (HPLC MS): [M+H⁺] = 651.2; Yield: 8 %; IC₅₀=13

PXM660-d2



C₃₄H₃₆Cl₂N₄O₃S; MW: 651.66; found (HPLC MS): [M+H⁺] = 651.2; Yield: 5 %; IC₅₀=8.3


PXM661-d1



C₃₃H₃₄Cl₂N₄O₃S; MW: 637.63; found (HPLC MS): [M+H⁺] = 637.2; Yield: 4 %; IC₅₀=13.8


PXM661-d2

- 81 -



$C_{33}H_{34}Cl_2N_4O_3S$; MW: 637.63; found (HPLC MS): $[M+H^+] = 637.2$; Yield: 3 %; $IC_{50}=10.2$


PXN662-d1



5

$C_{34}H_{36}Cl_2N_4O_3S$; MW: 651.66; found (HPLC MS): $[M+H^+] = 651.2$; Yield: 1 %; $IC_{50}=17.2$

PXN662-d2




10

$C_{34}H_{36}Cl_2N_4O_3S$; MW: 651.66; found (HPLC MS): $[M+H^+] = 651.2$; Yield: 3 %; $IC_{50}=11.5$


PXN663-d1

- 82 -



$C_{33}H_{33}Cl_3N_4O_3S$; MW: 672.08; found (HPLC MS): $[M+H^+] = 673.1$; Yield: 19 %; $IC_{50}=9.2$


PXN663-d2



5

$C_{33}H_{33}Cl_3N_4O_3S$; MW: 672.08; found (HPLC MS): $[M+H^+] = 671.1$; Yield: 7 %; $IC_{50}=11.2$

PXN666-d1




10


$C_{30}H_{29}Cl_2N_3O_3S$; MW: 582.55; PXN617-enantiomer1; $IC_{50}=2.9$

PXN667-d1

- 83 -

 $C_{30}H_{29}Cl_2N_3O_3S$; MW: 582.55; PXN617-enantiomer2; $IC_{50}=39$


PXN668-d1



5

 $C_{33}H_{35}ClN_4O_3S$; MW: 603.19; found (HPLC MS): $[M+H^+] = 603.0$; Yield: 5 %; $IC_{50}=18.3$

PXN668-d2




10

 $C_{33}H_{35}ClN_4O_3S$; MW: 603.19; found (HPLC MS): $[M+H^+] = 603.0$; Yield: 4 %; $IC_{50}=26.9$


PXN669-d1

- 84 -



$C_{34}H_{36}Cl_2N_4O_3S$; MW: 651.66; found (HPLC MS): $[M+H^+] = 651.0$; Yield: 5 %; $IC_{50}=16.1$


PXN669-d2



5

$C_{34}H_{36}Cl_2N_4O_3S$; MW: 651.66; found (HPLC MS): $[M+H^+] = 650.9$; Yield: 5 %; $IC_{50}=10.6$

PXN671-d1




10

$C_{34}H_{36}BrClN_4O_3S$; MW: 696.11; found (HPLC MS): $[M+H^+] = 697.0$; Yield: 11 %; $IC_{50}=8.7$


PXN671-d2

- 85 -



$C_{34}H_{36}BrClN_4O_3S$; MW: 696.11; found (HPLC MS): $[M+H^+] = 697.1$; Yield: 9 %; $IC_{50}=8.5$


PXN672-d1



5

$C_{33}H_{34}BrClN_4O_3S$; MW: 682.08; found (HPLC MS): $[M+H^+] = 682.8$; Yield: 5 %; $IC_{50}=29.5$

PXN672-d2




10

$C_{33}H_{34}BrClN_4O_3S$; MW: 682.08; found (HPLC MS): $[M+H^+] = 682.9$; Yield: 4 %; $IC_{50}=8.8$


PXN673-d2

- 86 -



$C_{32}H_{33}Cl_2N_5O_3S$; MW: 638.62; found (HPLC MS): $[M+H^+] = 637.9$; Yield: 1 %; $IC_{50}=163.4$


PXN673-d1



5

$C_{32}H_{33}Cl_2N_5O_3S$; MW: 638.62; found (HPLC MS): $[M+H^+] = 638.0$; Yield: 4 %; $IC_{50}=9.3$

PXN674-d1




10

$C_{31}H_{28}Cl_2N_4O_3S$; MW: 607.56; found (HPLC MS): $[M+H^+] = 607.2$; Yield: 4 %; $IC_{50}=17.5$


PXN674-d2

- 87 -



$C_{31}H_{28}Cl_2N_4O_3S$; MW: 607.56; found (HPLC MS):
[M+H⁺] = 607.2; Yield: 4 %; IC₅₀=28.9


PXN675-d1



5

$C_{31}H_{28}Cl_2N_4O_3S$; MW: 607.56; found (HPLC MS):
[M+H⁺] = 606.9; Yield: 2 %; IC₅₀=25.5

PXN675-d2




10

$C_{31}H_{28}Cl_2N_4O_3S$; MW: 607.56; found (HPLC MS):
[M+H⁺] = 608.8; Yield: 3 %; IC₅₀=17.8


PXN676-d1

- 88 -



$C_{33}H_{34}ClFN_4O_3S$; MW: 621.18; found (HPLC MS): $[M+H^+] = 621.0$; Yield: 5 %; $IC_{50}=15.6$


PXN676-d2



5

$C_{33}H_{34}ClFN_4O_3S$; MW: 621.18; found (HPLC MS): $[M+H^+] = 621.0$; Yield: 3 %; $IC_{50}=18.5$

PXN677-d1




10

$C_{31}H_{28}BrClN_4O_3S$; MW: 652.01; found (HPLC MS): $[M+H^+] = 651.3$; Yield: 8 %; $IC_{50}=5.45$


PXN678-d1

- 89 -



$C_{33}H_{33}Cl_3N_4O_5S$; MW: 704.08; found (HPLC MS): $[M+H^+] = 702.9$; Yield: 5 %; $IC_{50}=49.3$


PXN679-d2



5

$C_{34}H_{36}BrClN_4O_3S$; MW: 696.11; found (HPLC MS): $[M+H^+] = 697.1$; Yield: 3 %; $IC_{50}=9.3$

PXN679-d1




10

$C_{34}H_{36}BrClN_4O_3S$; MW: 696.11; found (HPLC MS): $[M+H^+] = 697.2$; Yield: 8 %; $IC_{50}=7.7$


PXN680-d1

- 90 -



$C_{34}H_{36}Br_2N_4O_3S$; MW: 740.56; found (HPLC MS): $[M+H^+] = 740.5$; Yield: 10 %; $IC_{50}=8.2$


PXN681-d2



5

$C_{34}H_{35}Cl_2FN_4O_3S$; MW: 669.65; found (HPLC MS): $[M+H^+] = 669.0$; Yield: 4 %; $IC_{50}=11.8$

PXN681-d1




10

$C_{34}H_{35}Cl_2FN_4O_3S$; MW: 669.65; found (HPLC MS): $[M+H^+] = 669.1$; Yield: 15 %; $IC_{50}=9.5$


PXN682-d2

- 91 -



$C_{34}H_{35}Cl_2FN_4O_3S$; MW: 669.65; found (HPLC MS): $[M+H^+] = 669.1$; Yield: 1 %; $IC_{50}=16.2$


PXN682-d1



5

$C_{34}H_{35}Cl_2FN_4O_3S$; MW: 669.65; found (HPLC MS): $[M+H^+] = 669.1$; Yield: 5 %; $IC_{50}=8.6$

PXN683-d1




10

$C_{33}H_{34}Cl_2N_4O_3S$; MW: 637.63; found (HPLC MS): $[M+H^+] = 637.0$; Yield: 8 %; $IC_{50}=6.0$


PXN684

- 92 -



$C_{30}H_{29}Cl_2N_3O_4S$; MW: 598.55; found (HPLC MS):
 $[M+H^+] = 598.2$; Yield: 7 %; $IC_{50}=5.7$


PXN686-d1



5

$C_{33}H_{35}Cl_2N_5O_3S$; MW: 652.65; found (HPLC
 MS): $[M+H^+] = 652.0$; Yield: 6 %; $IC_{50}=6.8$

PXN687




10

$C_{28}H_{24}Cl_2N_2O_3S$; MW: 539.49; found (HPLC MS):
 $[M+H^+] = 539.1$


PXN688

- 93 -



$C_{34}H_{36}Cl_2N_4O_5S$; MW: 683.66; found (HPLC MS): $[M+H^+] = 683.1$; Yield: 5 %; $IC_{50} > 60$


PXN689-d1



5

$C_{34}H_{36}Cl_2N_4O_3S$; MW: 651.66; found (HPLC MS): $[M+H^+] = 651.1$; Yield: 8 %; $IC_{50} = 8.6$

PXN690




10

$C_{33}H_{33}Cl_2N_3O_7S$; MW: 686.62; found (HPLC MS): $[M+Na^+] = 710.1$; Yield: 9 %; $IC_{50} = 33.8$


PXN691

- 94 -




$C_{34}H_{37}Cl_2N_3O_7S$; MW: 702.66; found (HPLC MS): $[M+H^+] = 702.3$; $[M+Na^+] = 724.1$; Yield: 4 %; $IC_{50}=17.6$

5 PXN693-d1



$C_{34}H_{36}BrClN_4O_5S$; MW: 728.11; found (HPLC MS): $[M+H^+] = 729.1$; Yield: 4 %

PXN694-d1




10

$C_{34}H_{35}Cl_2FN_4O_5S$; MW: 701.65; found (HPLC MS): $[M+H^+] = 703.1$; Yield: 7 %


- 95 -

PXN695-d1




$C_{33}H_{34}BrClN_4O_3S$; MW: 682.08; found (HPLC MS): $[M+H^+] = 683.0$; Yield: 5 %; $IC_{50}=5.4$

5 PXN696-d1



$C_{30}H_{29}BrClN_3O_4S$; MW: 643.00; found (HPLC MS): $[M+H^+] = 668.1$; Yield: 7 %; $IC_{50}=4.1$


PXN697-d1



$C_{31}H_{28}BrClN_4O_3S$; MW: 652.01; found (HPLC MS): $[M+H^+] = 653.1$; $[M+Na^+] = 675.3$; Yield: 3 %; $IC_{50}=3.6$


- 96 -

PXN698-d1




$C_{33}H_{34}BrClN_4O_3S$; MW: 682.08; found (HPLC MS): $[M+H^+] = 683.1$; Yield: 7 %; $IC_{50}=6.3$

5 PXN699-d1



$C_{35}H_{38}BrClN_4O_3S$; MW: 710.14; found (HPLC MS): $[M+H^+] = 711.0$; Yield: 7 %; $IC_{50}=4.2$


PXN700-d1



$C_{31}H_{31}BrClN_3O_3S$; MW: 641.03; found (HPLC MS): $[M+Na^+] = 664.2$; Yield: 7 %; $IC_{50}=4.3$


- 97 -

PXN701-d1




$C_{33}H_{34}BrClN_4O_2S$; MW: 666.09; found (HPLC MS):
[M+H⁺] = 668.2; Yield: 8 %; IC₅₀=4.1

5 PXN702-d1



$C_{35}H_{38}BrClN_4O_3S$; MW: 710.14; found (HPLC MS): [M+H⁺] = 711.1; Yield: 17 %; IC₅₀=18.6

PXN703-d1




10

$C_{35}H_{38}BrClN_4O_3S$; MW: 710.14; found (HPLC MS): [M+H⁺] = 711.0; Yield: 10 %; IC₅₀=9.5


- 98 -

PXN704-d1




$C_{35}H_{38}BrClN_4O_3S$; MW: 710.14; found (HPLC MS): $[M+H^+] = 711.0$; Yield: 6 %; $IC_{50}=11.7$

5 PXN705-d1



$C_{33}H_{33}BrClN_5O_5S$; MW: 727.08; found (HPLC MS): $[M+H^+] = 727.9$; Yield: 1 %; $IC_{50}=16.4$

PXN706-d1




10

$C_{31}H_{29}BrClN_3O_3S$; MW: 639.02; found (HPLC MS): $[M+H^+] = 640.3$; $[M+Na^+] = 662.3$; Yield: 8 %; $IC_{50}=4.7$


- 99 -

PXN707-d1




$C_{31}H_{29}BrClN_3O_3S$; MW: 639.02; found (HPLC MS):
[M+H⁺] = 640.3; [M+Na⁺] = 662.3; Yield: 11 %; IC₅₀=5.4

5 PXN708-d1



$C_{30}H_{29}BrClN_3O_3S$; MW: 627.00; found (HPLC MS):
[M+Na⁺] = 650.6; Yield: 8 %; IC₅₀=6.1

PXN709-d1




10

$C_{30}H_{29}BrClN_3O_3S$; MW: 627.00; found (HPLC MS):
[M+Na⁺] = 650.2; Yield: 8 %; IC₅₀=7.4


- 100 -

PXN705-d2




$C_{33}H_{33}BrClN_5O_5S$; MW: 727.08; found (HPLC MS): $[M+H^+] = 726.0$; Yield: 3 %; $IC_{50}=9.5$

5 PXN710



$C_{35}H_{40}BrClN_4O_2S$; MW: 696.16; found (HPLC MS): $[M+H^+] = 697.3$; Yield: 6 %


PXN711-d1



$C_{33}H_{33}BrClFN_4O_3S$; MW: 700.08; found (HPLC MS): $[M+H^+] = 701.2$; Yield: 3 %; $IC_{50}=7.3$


- 101 -

PXN712-d1




$C_{33}H_{34}BrClN_4O_3S$; MW: 682.08; found (HPLC MS): $[M+H^+] = 683.1$; Yield: 6 %

5 PXN713-d1



$C_{32}H_{40}BrClN_4O_3S$; MW: 676.12; found (HPLC MS): $[M+H^+] = 677.2$; Yield: 4 %; $IC_{50}=8.1$

PXN714-d1




10

$C_{34}H_{36}BrClN_4O_3S$; MW: 696.11; found (HPLC MS): $[M+H^+] = 697.2$; Yield: 2 %; $IC_{50}=6.8$


PXN715-d1

- 102 -



$C_{33}H_{32}BrClF_2N_4O_3S$; MW: 718.07; found (HPLC MS): $[M+H^+] = 719.2$; Yield: 4 %; $IC_{50}=7.9$


PXN716-d1



5

$C_{31}H_{28}BrClN_4O_3S$; MW: 652.01; found (HPLC MS): $[M+H^+] = 653.1$; $[M+Na^+] = 675.3$; Yield: 4 %

PXN717-d1




10

$C_{31}H_{29}BrClN_3O_3S$; MW: 639.02; found (HPLC MS): $[M+H^+] = 640.2$; $[M+Na^+] = 662.1$; Yield: 10 %; $IC_{50}=8.9$


PXN718-d1

- 103 -



$C_{31}H_{30}BrClN_4O_2S$; MW: 638.03; found (HPLC MS):
[M+H⁺] = 639.2; Yield: 9 %; IC₅₀=3.5


PXN719-d1



5

$C_{33}H_{34}BrClN_4O_2S$; MW: 666.09; found (HPLC MS):
[M+H⁺] = 665.4; Yield: 8 %; IC₅₀=8.2

PXN720-d1




10

$C_{34}H_{36}BrClN_4O_3S$; MW: 696.11; found (HPLC MS):
[M+H⁺] = 695.4; Yield: 5 %; IC₅₀=6.6


PXN721-d1

- 104 -



$C_{32}H_{32}BrClN_4O_2S$; MW: 652.06; found (HPLC MS):
 $[M+H^+] = 651.4$; Yield: 5 %; $IC_{50}=7.2$


PXN722-d1



5

$C_{33}H_{36}BrClN_4O_2S$; MW: 668.10; found (HPLC
 MS): $[M+H^+] = 667.3$; Yield: 2 %; $IC_{50}=13.2$

PXN725-d1




10

$C_{32}H_{32}BrClN_4O_4S_2$; MW: 716.12; found (HPLC MS):
 $[M+Na^+] = 737.2$; Yield: 5 %; $IC_{50}>60$


PXN726-d1

- 105 -



$C_{33}H_{32}BrClN_4O_3S$; MW: 680.07; found (HPLC MS):
 $[M+H^+] = 681.1$; $[M+Na^+] = 703.2$; Yield: 9 %; $IC_{50}=2.1$


PXN727-d1



5

$C_{34}H_{35}BrClN_5O_3S$; MW: 709.11; found (HPLC MS):
 $[M+H^+] = 710.0$; $[M+Na^+] = 732.2$; Yield: 11 %; $IC_{50}=1.4$

PXN728-d1




10


$C_{31}H_{32}BrClN_4O_3S$; MW: 656.05; found (HPLC
 MS): $[M+H^+] = 657.2$; Yield: 4 %

PXN729

- 106 -


 $C_{58}H_{52}Br_2Cl_2N_6O_5S_2$; MW:
1207.94; found (HPLC MS): $[M+Na^+] = 1229.5$; Yield: 10 %

PXN730-d1




5

 $C_{32}H_{30}BrClN_4O_3S$; MW: 666.04; found (HPLC MS):

 $[M+H^+] = 666.9$; $[M+Na^+] = 689.0$; Yield: 6 %; $IC_{50}=4.8$

PXN731-d1




10

 $C_{34}H_{34}BrClN_4O_4S$; MW: 710.10; found (HPLC MS):

 $[M+H^+] = 711.5$; $[M+Na^+] = 733.5$; Yield: 4 %; $IC_{50}=118.7$


PXN732-d1

- 107 -



$C_{33}H_{32}BrClN_4O_4S$; MW: 696.07; found (HPLC MS):
 $[M+H^+] = 697.4$; $[M+Na^+] = 719.4$; Yield: 4 %; $IC_{50}=2.1$


PXN733-d1



5

$C_{36}H_{38}BrClN_4O_4S$; MW: 738.15; found (HPLC
 MS): $[M+H^+] = 739.1$; $[M+Na^+] = 759.5$; Yield: 4 %

PXN734-d1




$C_{32}H_{30}BrClN_4O_3S$; MW: 666.04

10


PXN735-d1

- 108 -



$C_{35}H_{36}BrClN_4O_3S$; MW: 708.12; found (HPLC MS):
 $[M+H^+] = 711.0$; $[M+Na^+] = 731.0$; Yield: 8 %; $IC_{50}=3.4$


PXN736-d1



5

$C_{36}H_{38}BrClN_4O_3S$; MW: 722.15; found (HPLC
 MS): $[M+H^+] = 723.4$; $[M+Na^+] = 745.4$; Yield: 7 %; $IC_{50}>60$

PXN737-d1




10


$C_{36}H_{38}BrClN_4O_3S$; MW: 722.15; found (HPLC MS):
 $[M+H^+] = 721.4$; $[M+Na^+] = 743.5$; Yield: 7 %; $IC_{50}=15.6$

PXN738-d1

- 109 -

 $C_{33}H_{33}BrClN_5O_3S$; MW: 695.08


PXN739-d1

 $C_{34}H_{35}BrClN_5O_3S$; MW: 709.11; found (HPLC MS):

5


 $[M+H^+] = 710.2$; Yield: 2 %; IC₅₀=2.0

PXN740-d1

 $C_{35}H_{37}BrClN_5O_3S$; MW: 723.14


10 PXN741-d1

- 110 -



$C_{33}H_{33}BrClN_3O_3S$; MW: 667.07; found (HPLC MS):
 $[M+H^+] = 668.5$; $[M+Na^+] = 690.4$; Yield: 5 %; $IC_{50}=7.0$


PXN742-d1



5

$C_{34}H_{35}Cl_2N_5O_3S$; MW: 664.66; found (HPLC MS):
 $[M+H^+] = 664.2$; $[M+Na^+] = 686.1$; Yield: 6 %; $IC_{50}=1.1$

PXN743-d1




10

$C_{34}H_{34}Cl_2FN_5O_3S$; MW: 682.65; found (HPLC MS):
 $[M+H^+] = 682.7$; $[M+Na^+] = 704.2$; Yield: 6 %; $IC_{50}=1.5$


PXN744-d1

- 111 -



$C_{35}H_{38}ClN_5O_3S$; MW: 644.24; found (HPLC MS):
 $[M+H^+] = 644.4$; $[M+Na^+] = 666.3$; Yield: 5 %; $IC_{50}=107.1$


PXN745



5

$C_{31}H_{32}Cl_2N_4O_3S$; MW: 611.60; found (HPLC MS):
 $[M+H^+] = 611.3$; $[M+Na^+] = 633.2$; Yield: 2 %

PXN746




10


$C_{32}H_{34}Cl_2N_4O_3S$; MW: 625.62; found (HPLC MS):
 $[M+H^+] = 625.3$; $[M+Na^+] = 649.3$; Yield: 22 %; $IC_{50}=14.7$

PXN747

- 112 -




PXN748-d1



5


$[\text{M}+\text{H}^+] = 739.3$; $[\text{M}+\text{Na}^+] = 762.1$; Yield: 5 %; $\text{IC}_{50}=16.6$

PXN749-d1




10 PXN750-d1

- 113 -


 $C_{34}H_{33}BrClN_4NaO_6S_2$; MW: 796.14


PXN751-d1


 $C_{32}H_{31}Cl_2N_5O_3S$; MW: 636.61; found (HPLC MS):

5

 $[M+H^+] = 636.3$; $[M+Na^+] = 658.2$; Yield: 2 %; $IC_{50}=5.8$

PXN752-d1



 $C_{32}H_{30}Cl_2N_4O_3S$; MW: 621.59; found (HPLC MS):

 $[M+H^+] = 621.1$; Yield: 6 %; $IC_{50}=2.0$

10


PXN753-d1

- 114 -



$C_{34}H_{36}ClN_5O_3S$; MW: 630.21; found (HPLC MS):
 $[M+H^+] = 630.3$; $[M+Na^+] = 652.3$; Yield: 5 %; $IC_{50}=170.6$


PXN754-d1



5

$C_{31}H_{36}ClN_5O_3S$; MW: 594.18; found (HPLC MS):
 $[M+H^+] = 594.3$; Yield: 12 %; $IC_{50}=13.6$

PXN755-d1




10

$C_{33}H_{33}Cl_2N_5O_3S$; MW: 650.63; found (HPLC MS):
 $[M+H^+] = 650.1$; $[M+Na^+] = 672.3$; Yield: 1 %; $IC_{50}=3.6$


PXN756-d1

- 115 -



$C_{34}H_{35}BrClN_5O_3S$; MW: 709.11; found (HPLC MS):
 $[M+H^+] = 710.0$; $[M+Na^+] = 732.2$; Yield: 8 %; $IC_{50}=2.6$


PXN757-d1



5

$C_{35}H_{35}BrClN_5O_4S$; MW: 709.11; found (HPLC MS):
 $[M+Na^+] = 730.4$; Yield: 12 %; $IC_{50}=206.1$

PXN758-d1




10

$C_{34}H_{35}BrClN_5O_3S$; MW: 709.11; found (HPLC MS):
 $[M+H^+] = 710.0$; $[M+Na^+] = 732.2$; Yield: 13 %

PXN759-d1


- 116 -



$C_{34}H_{36}ClN_5O_3S$; MW: 630.21; found (HPLC MS):

$[M+H^+] = 630.2$; $[M+Na^+] = 652.3$; Yield: 6 %; $IC_{50}=182.6$


PXN760-d1



5

$C_{34}H_{35}BrClN_5O_4S$; MW: 725.11; found (HPLC MS): $[M+H^+] = 726.2$; $[M+Na^+] = 748.0$; Yield: 4 %; $IC_{50}=3.2$

PXN761-d1




10

$C_{34}H_{36}BrCl_2N_5O_3S$; MW: 745.57; found (HPLC MS): $[M+H^+] = 708.1$; Yield: 10 %; $IC_{50}=3.1$


PXN762-d1

- 117 -



$C_{33}H_{34}Cl_3N_5O_3S$; MW: 687.09; found (HPLC MS): $[M+H^+] = 650.2$; Yield: 4 %; $IC_{50}=4.5$


PXN763-d1



5

$C_{35}H_{38}Cl_2N_6O_3S$; MW: 693.70; found (HPLC MS): $[M+H^+] = 693.2$; Yield: 2 %

PXN764-d1




10

$C_{36}H_{39}Cl_2N_5O_4S$; MW: 708.71; found (HPLC MS): $[M+H^+] = 708.2$; $[M+Na^+] = 730.2$; Yield: 5 %; $IC_{50}=3.3$


PXN765-d1

- 118 -



$\text{C}_{35}\text{H}_{37}\text{Cl}_2\text{N}_5\text{O}_4\text{S}$; MW: 694.69; found (HPLC MS): $[\text{M}+\text{H}^+] = 694.2$; $[\text{M}+\text{Na}^+] = 716.3$; Yield: 4 %; $\text{IC}_{50}=2.6$


PXN766-d1



5

$\text{C}_{33}\text{H}_{32}\text{BrClN}_4\text{O}_4\text{S}$; MW: 696.07; found (HPLC MS): $[\text{M}+\text{Na}^+] = 717.2$; Yield: 6 %; $\text{IC}_{50}=3.8$

PXN767-d1




10

$\text{C}_{36}\text{H}_{33}\text{Cl}_2\text{N}_5\text{O}_2\text{S}$; MW: 670.67; found (HPLC MS): $[\text{M}+\text{H}^+] = 670.2$; Yield: 9 %; $\text{IC}_{50}>60$


PXN768-d1

- 119 -



$C_{35}H_{32}Cl_2N_6O_2S$; MW: 671.65; found (HPLC MS):
[M+H⁺] = 671.4; Yield: 6 %; IC₅₀=118.7


PXN769-d1



5

$C_{35}H_{32}Cl_2N_6O_2S$; MW: 671.65; found (HPLC MS):
[M+H⁺] = 671.1; Yield: 8 %; IC₅₀>60

PXN770-d1




10

$C_{38}H_{35}Cl_2N_5O_3S$; MW: 712.70; found (HPLC MS):
[M+H⁺] = 712.2; Yield: 5 %; IC₅₀=191.7


PXN771-d1

- 120 -



$C_{34}H_{35}BrClN_5O_3S$; MW: 709.11; found (HPLC MS):
 $[M+H^+] = 710.0$; $[M+Na^+] = 732.2$; Yield: 10 %


PXN775-d1



5

$C_{35}H_{38}BrN_5O_3S$; MW: 688.69; found (HPLC MS):
 $[M+H^+] = 690.2$; Yield: 2 %; CCA

PXN776-d1




10

$C_{33}H_{33}BrClN_5O_3S$; MW: 695.08; found (HPLC MS):
 $[M+Na^+] = 716.1$; Yield: 11 %; $IC_{50}=2.1$


PXN777-d1

- 121 -



$C_{35}H_{37}BrClN_5O_3S$; MW: 723.14; found (HPLC MS): $[M+H^+] = 722.1$; Yield: 10 %; $IC_{50}=5.5$


PXN779-d1



5

$C_{33}H_{34}BrClN_6O_3S$; MW: 710.10; found (HPLC MS): $[M+H^+] = 711.2$; Yield: 6 %; $IC_{50}=5.0$

PXN780-d1




10


$C_{35}H_{35}ClF_3N_5O_3S$; MW: 698.21; found (HPLC MS): $[M+H^+] = 698.2$; $[M+Na^+] = 720.2$; Yield: 6 %; $IC_{50}=5.3$

PXN781-d1

- 122 -




PXN782-d1



5


PXN783-d1



10

PXN784-d1


- 123 -



$C_{39}H_{46}BrClN_6O_3S$; MW: 794.26; found (HPLC MS):

$[M+H^+] = 795.3$; Yield: 5 %

PXN785-d1




5

$C_{39}H_{47}BrClN_5O_3S$; MW: 781.26; found (HPLC MS): $[M+H^+] = 781.6$; $[M+Na^+] = 803.8$; Yield: 10 %


PXN787-d1

- 124 -



$C_{35}H_{39}BrClN_5O_2S$; MW: 709.15; found
(HPLC MS): $[M+H^+] = 709.5$; Yield: 13 %; $IC_{50}=8.3$


PXN791-d1



5

$C_{34}H_{35}BrClN_5O_3S$; MW: 709.11; found (HPLC
MS): $[M+H^+] = 710.2$; $[M+Na^+] = 732.0$; Yield: 1 %; $IC_{50}=6.6$

PXN792-d1




10

$C_{34}H_{35}BrClN_5O_3S$; MW: 709.11; found (HPLC
MS): $[M+H^+] = 708.0$; $[M+Na^+] = 730.1$; Yield: 8 %; $IC_{50}=5.3$


PXN793-d1

- 125 -



$C_{33}H_{32}BrClN_4O_3S$; MW: 680.07; found (HPLC MS): $[M+Na^+] = 703.1$; Yield: 6 %; $IC_{50}=4.7$


PXN794-d1



5

$C_{33}H_{32}BrClN_4O_3S$; MW: 680.07; found (HPLC MS): $[M+H^+] = 679.0$; $[M+Na^+] = 701.0$; Yield: 4 %; $IC_{50}=6.0$

PXN795-d1




10


$C_{34}H_{34}BrClN_4O_4S$; MW: 710.10; found (HPLC MS): $[M+H^+] = 711.1$; $[M+Na^+] = 733.1$; Yield: 3 %; $IC_{50}=80.0$

PXN796-d1

- 126 -


 $C_{34}H_{34}BrClN_4O_4S$; MW: 710.10


PXN797-d1


 $C_{34}H_{35}BrFN_5O_3S$; MW: 692.66; found (HPLC MS):

5


 $[M+H^+] = 693.5$; $[M+Na^+] = 716.2$; Yield: 11 %; CCA

PXN798-d1


 $C_{37}H_{42}BrClN_6O_3S$; MW: 766.21


10 PXN799-d1

- 127 -



$C_{32}H_{31}BrClN_3O_3S$; MW: 653.04; found (HPLC MS):
 $[M+H^+] = 651.8$; $[M+Na^+] = 675.7$; Yield: 13 %; $IC_{50}=4.1$


PXN800-d1



5

$C_{34}H_{34}BrClN_4O_4S$; MW: 710.10; found (HPLC
 MS): $[M+H^+] = 711.3$; $[M+Na^+] = 731.2$; Yield: 2 %; $IC_{50}=11.7$

PXN801-d1




10

$C_{35}H_{37}BrClN_5O_3S$; MW: 723.14; found
 (HPLC MS): $[M+H^+] = 724.3$; $[M+Na^+] = 746.3$; Yield: 5 %; $IC_{50}>60$


PXN802-d1

- 128 -



$C_{34}H_{34}BrClN_4O_3S$; MW: 694.10; found (HPLC MS): $[M+H^+] = 695.3$; Yield: 6 %; $IC_{50}=4.5$


PXN803-d1



5

$C_{35}H_{36}BrClN_4O_4S$; MW: 724.12

PXN805-d1




$C_{32}H_{32}BrClN_4O_2S$; MW: 652.06; found (HPLC MS): $[M+H^+] = 653.4$; $[M+Na^+] = 675.4$; Yield: 7 %; $IC_{50}=5.5$


10

PXN806-d1

- 129 -



 $C_{33}H_{32}BrClN_4O_4S$; MW: 696.07

PXN807-d1


 $C_{33}H_{32}BrClN_4O_4S$; MW: 696.07


5

PXN808-d1


 $C_{32}H_{29}Cl_2FN_4O_3S$; MW: 639.58; found (HPLC MS):
 $[M+H^+] = 639.2$; $[M+Na^+] = 660.9$; Yield: 16 %; $IC_{50}=1.3$


10 PXN811-d1

- 130 -



$C_{30}H_{26}Cl_2FN_3O_3S$; MW: 598.53; found (HPLC MS):
 $[M+H^+] = 598.1$; $[M+Na^+] = 620.0$; Yield: 7 %; $IC_{50}=5.6$


PXN813-d1



5

$C_{30}H_{27}BrClN_3O_3S$; MW: 624.99; found (HPLC MS):
 $[M+H^+] = 624.0$; $[M+Na^+] = 646.0$; Yield: 11 %; $IC_{50}=5.4$

PXN814-d1




$C_{28}H_{25}BrClN_3O_3S$; MW: 598.95

10


PXN815-d1

- 131 -



$C_{33}H_{32}ClF_2N_5O_3S$; MW: 652.17; found (HPLC MS):
 $[M+H^+] = 652.1$; $[M+Na^+] = 674.2$; Yield: 4 %; $IC_{50}=5.0$


PXN816-d1



5

$C_{33}H_{32}Cl_2FN_5O_3S$; MW: 668.62; found (HPLC MS):
 $[M+H^+] = 668.2$; Yield: 5 %; $IC_{50}=3.2$

PXN817-d2




10

$C_{34}H_{33}ClN_6O_3S$; MW: 641.20; found (HPLC MS):
 $[M+H^+] = 641.1$; $[M+Na^+] = 663.2$; Yield: 1 %; $IC_{50}=1.0$


PXN820-d1

- 132 -



$C_{31}H_{28}Cl_2FN_3O_3S$; MW: 612.56; found (HPLC MS):
 $[M+H^+] = 612.1$; $[M+Na^+] = 634.1$; Yield: 20 %; $IC_{50}=3.7$


PXN821-d1



5

$C_{32}H_{31}Cl_2FN_4O_3S$; MW: 641.60; found (HPLC
 MS): $[M+H^+] = 641.2$; Yield: 19 %; $IC_{50}=8.6$

PXN822-d1




10

$C_{33}H_{32}Cl_2FN_5O_3S$; MW: 668.62; found (HPLC
 MS): $[M+H^+] = 668.2$; $[M+Na^+] = 690.1$; Yield: 16 %; $IC_{50}=1.2$


PXN825-d1

- 133 -



$C_{31}H_{27}Cl_2FN_4O_3S$; MW: 625.55; found (HPLC MS):
 $[M+H^+] = 627.2$; $[M+Na^+] = 647.1$; Yield: 23 %; $IC_{50}=1.5$


PXN826-d1



5

$C_{32}H_{30}Cl_2FN_3O_3S$; MW: 626.58; found (HPLC MS):
 $[M+Na^+] = 648.1$; Yield: 19 %; $IC_{50}=3.5$

PXN833-d1




10

$C_{35}H_{37}Cl_2FN_6O_4S$; MW: 727.69;
 found (HPLC MS): $[M+H^+] = 727.2$; Yield: 5 %


PXN834-d1

- 134 -



$C_{34}H_{34}Cl_2FN_5O_4S$; MW: 698.65; found
(HPLC MS): $[M+H^+] = 698.1$; Yield: 23 %


PXN835-d1



5

$C_{34}H_{34}Cl_2FN_5O_3S$; MW: 682.65; found (HPLC
MS): $[M+H^+] = 684.3$; $[M+Na^+] = 703.9$; Yield: 5 %; $IC_{50}=3.4$

PXN836-d2




10

$C_{33}H_{30}ClN_5O_3S$; MW: 612.16; found (HPLC MS):
 $[M+H^+] = 612.5$; $[M+Na^+] = 634.5$; Yield: 3 %

PXN849-d1


- 135 -



$C_{33}H_{31}Cl_2FN_4O_3S$; MW: 653.61; found (HPLC MS):

$[M+H^+] = 653.4$; $[M+Na^+] = 677.4$; Yield: 6 %

PXN850-d1




5

$C_{34}H_{34}Cl_2FN_5O_3S$; MW: 682.65; found (HPLC

MS): $[M+H^+] = 682.4$; Yield: 11 %

PXN670-d1




10

$C_{33}H_{34}Cl_2N_4O_3$; MW: 605.57; found (HPLC MS):

$[M+H^+] = 605.0$; Yield: 6 %; $IC_{50}=12.5$


PXN670-d2

- 136 -


 $C_{33}H_{34}Cl_2N_4O_3$; MW: 605.57; found (HPLC MS):

 $[M+H^+] = 605.0$; Yield: 7 %; $IC_{50}=11.2$

PXN778-d1




5

 $C_{33}H_{33}BrClN_5O_3$; MW: 663.02; found (HPLC MS):

 $[M+H^+] = 664.1$; Yield: 11 %; $IC_{50}=8.1$

PXN788-d1




10

 $C_{33}H_{39}BrClN_5O_3$; MW: 669.07; found (HPLC MS):

 $[M+H^+] = 670.3$; $[M+Na^+] = 692.2$; Yield: 1 %; $IC_{50}=6.6$


PXN790-d1

- 137 -


 $C_{34}H_{34}Cl_2FN_5O_3$; MW: 650.59; found (HPLC MS):

 $[M+H^+] = 650.1$; $[M+Na^+] = 672.1$; Yield: 1 %; $IC_{50}=3.0$

PXN804-d1




5

 $C_{30}H_{33}BrClN_5O_3$; MW: 626.99; found (HPLC MS):

 $[M+H^+] = 628.2$; $[M+Na^+] = 650.2$; Yield: 9 %; $IC_{50}=9.0$


PXN809-d1


 $C_{27}H_{26}Cl_2FN_3O_3$; MW: 530.43


10

PXN810-d1

- 138 -

 $C_{28}H_{28}Cl_2FN_3O_3$; MW: 544.46


PXN812-d1

 $C_{31}H_{28}Cl_2FN_3O_3$; MW: 580.49; found (HPLC MS):

5


 $[M+H^+] = 580.1$; $[M+Na^+] = 602.1$; Yield: 4 %; $IC_{50}=11.9$

PXN823-d1


 $C_{32}H_{30}Cl_2FN_3O_3$; MW: 594.52

10 PXN824-d1

- 139 -


 $C_{33}H_{31}Cl_2FN_4O_3$; MW: 621.54

PXN827-d1

 $C_{32}H_{29}Cl_2FN_4O_3$; MW: 607.52


5

PXN828-d1

 $C_{32}H_{35}Cl_2FN_4O_3$; MW: 613.57


PXN829-d1

- 140 -



$C_{33}H_{38}Cl_2FN_5O_3$; MW: 642.61; found (HPLC MS):
 $[M+H^+] = 642.2$; $[M+Na^+] = 664.2$; Yield: 1 %; $IC_{50}=12.5$


PXN830-d1



5

$C_{31}H_{33}Cl_2FN_4O_3$; MW: 599.54; found (HPLC MS):
 $[M+H^+] = 599.2$; $[M+Na^+] = 621.2$; Yield: 1 %; $IC_{50}=10.2$

PXN831-d1




10

$C_{31}H_{34}Cl_2FN_5O_3$; MW: 614.55; found (HPLC
 MS): $[M+H^+] = 614.2$; $[M+Na^+] = 636.2$; Yield: 1 %; $IC_{50}=4.9$


PXN832-d1

- 141 -



$C_{31}H_{36}Cl_2FN_5O_3$; MW: 616.57; found (HPLC MS): $[M+H^+] = 616.3$; Yield: 1 %; $IC_{50}=7.9$


PXN789-d1



5

$C_{34}H_{34}Cl_2FN_5O_4$; MW: 666.59; found (HPLC MS): $[M+H^+] = 666.1$; Yield: 1 %; $IC_{50}=11.7$

PXN723-d1




10

$C_{31}H_{31}BrClN_3O_2S$; MW: 625.03; found (HPLC MS):

$[M+H^+] = 626.2$; Yield: 2 %; $IC_{50}=15.1$


PXN724-d1

- 142 -



$C_{31}H_{33}BrClN_3OS$; MW: 611.05; found (HPLC MS):
 $[M+H^+] = 612.3$; Yield: 1 %; $IC_{50}=20.6$


PXN818-d1



5

$C_{27}H_{24}BrClN_2O_2S$; MW: 555.93; found (HPLC MS): $[M+H^+]$
 $= 556.9$; $[M+Na^+] = 577.1$; Yield: 10 %; $IC_{50}=8.4$

PXN819-d1




$C_{34}H_{37}BrClN_5O_2S$; MW: 695.13

10

PXN837-d1


- 143 -



$C_{22}H_{24}BrNO_3S$; MW: 462.41; found (HPLC MS): $[M+H^+] = 464.2$;

Yield: 5 %

PXN838-d1




$C_{25}H_{22}BrNO_3S$; MW: 496.43; found (HPLC MS):

5

$[M+H^+] = 496.9$; Yield: 2 %

PXN839-d1




$C_{34}H_{30}BrClN_4O_2S$; MW: 674.06; found (HPLC MS): $[M+H^+] = 675.4$; $[M+Na^+] = 696.9$; Yield: 10 %

10


PXN840-d1

- 144 -



$C_{34}H_{29}BrClN_3O_3S$; MW: 675.05; found (HPLC MS): $[M+H^+] = 676.7$; $[M+Na^+] = 698.3$; Yield: 9 %


PXN841-d1



5

$C_{29}H_{26}BrClN_2O_3S$; MW: 597.96; found (HPLC MS): $[M+H^+] = 599.3$; $[M+Na^+] = 620.9$; Yield: 11 %

PXN842-d1




10

$C_{34}H_{35}BrClN_5O_2S_2$; MW: 725.18; found (HPLC MS): $[M+H^+] = 726.5$; $[M+Na^+] = 749.7$; Yield: 9 %


PXN843-d1

- 145 -



$C_{26}H_{22}N_4O_2S$; MW: 454.55; found (HPLC MS): $[M+H^+] = 455.3$; $[M+Na^+] = 477.2$; Yield: 22 %


PXN844-d1



5

$C_{27}H_{25}N_3O_4S$; MW: 487.58; found (HPLC MS): $[M+H^+] = 488.2$; $[M+Na^+] = 510.3$; Yield: 22 %

PXN845-d1




10

$C_{28}H_{24}BrClN_2O_3S$; MW: 583.94; found (HPLC MS): $[M+H^+] = 585.2$; $[M+Na^+] = 607.4$; Yield: 5 %


PXN846-d1

- 146 -



$\text{C}_{25}\text{H}_{21}\text{Cl}_3\text{N}_2\text{O}_3$; MW: 503.82; found (HPLC MS): $[\text{M}+\text{H}^+] = 505.2$; Yield: 4 %


PXN847-d1



5

$\text{C}_{29}\text{H}_{30}\text{BrClN}_4\text{O}_3$; MW: 597.94; found (HPLC MS): $[\text{M}+\text{H}^+] = 597.3$; Yield: 7 %

PXN848-d1




10

$\text{C}_{26}\text{H}_{21}\text{Cl}_2\text{NO}_5\text{S}$; MW: 530.43; found (HPLC MS): $[\text{M}+\text{H}^+] = 530.2$; $[\text{M}+\text{Na}^+] = 552.4$; Yield: 13 %


PXN 1000

- 147 -



$C_{28}H_{29}ClN_4O_3S$; MW: 537.1; found (HPLC MS):
 $[M+H^+] = 537.1$; Yield: 9 %


PXN 1001



5

$C_{28}H_{30}ClN_3O_3S$; MW: 524.1; found (HPLC MS):
 $[M+Na^+] = 546.2$; Yield: 19 %


PXN 1002



10

$C_{28}H_{30}N_4O_3S$; MW: 502.64; found (HPLC MS):
 $[M+H^+] = 503.2$, $[M+Na^+] = 525.2$; Yield: 11 %

PXN 1003




15

$C_{28}H_{31}N_3O_3S$; MW: 489.64; found (HPLC MS):
 $[M+H^+] = 490.5$, $[M+Na^+] = 512.2$; Yield: 12 %


- 148 -

PXN 1004


 $C_{28}H_{29}ClN_4O_3$; MW: 505.02; found (HPLC MS):


 $[M+H^+] = 505.2$; Yield: 11 %

5 PXN 1005


 $C_{28}H_{30}ClN_3O_3$; MW: 492.02; found (HPLC MS):

 $[M+H^+] = 492.2$; Yield: 13 %


PXN 1006



10

 $C_{29}H_{31}ClN_4O_4$; MW: 535.05

PXN 1007



 $C_{26}H_{28}ClN_3O_3S$; MW: 498.05; found (HPLC MS):

 $[M+H^+] = 498.3$; Yield: 5 %


15

- 149 -


PXN 1008

 $C_{26}H_{29}N_3O_3S$; MW: 463.60; found (HPLC MS): $[M+H^+] = 464.3$; Yield: 31 %

5 PXN 1009

 $C_{31}H_{29}ClN_4O_3S$; MW: 573.12; found (HPLC MS): $[M+H^+] = 574.4$, $[M+Na^+] = 595.4$; Yield: 1 %

PXN 1010




10


 $C_{31}H_{30}N_4O_3S$; MW: 538.67; found (HPLC MS): $[M+H^+] = 539.4$; Yield: 3%

PXN 1011

- 150 -

 $C_{29}H_{28}ClN_3O_3S$; MW: 534.08; found (HPLC MS): $[M+H^+] = 534.3$; Yield: 13 %


PXN 1012



5

 $C_{29}H_{29}N_3O_3S$; MW: 499.64; found (HPLC MS): $[M+H^+] = 500.3$; Yield: 31 %

PXN 1013




10


 $C_{28}H_{27}ClN_4O_3S$; MW: 535.07; found (HPLC MS): $[M+H^+] = 535.3$; Yield: 2 %

PXN 1014

- 151 -

 $C_{30}H_{34}ClN_5O_3S$; MW: 580.2; found (HPLC MS): $[M+Na^+] = 602.4$; Yield: 14 %


PXN 1015



5

 $C_{30}H_{29}N_5O_3S$; MW: 539.66; found (HPLC MS): $[M+H^+] = 540.4$; Yield: 1 %

PXN 1016




10

 $C_{28}H_{33}ClN_4O_3S$; MW: 541.12; found (HPLC MS): $[M+H^+] = 541.4$; Yield: 4 %


PXN 1017

- 152 -



$C_{37}H_{37}ClN_4O_4S$; MW: 669.2; found (HPLC MS): $[M+H^+] = 671.7$, $[M+Na^+] = 691.3$; Yield: 3 %


PXN 1018



5

$C_{30}H_{35}N_5O_3S$; MW: 545.7; found (HPLC MS):
 $[M+H^+] = 546.3$; Yield: 18 %

PXN 1019




10


$C_{31}H_{35}N_3O_4S$; MW: 545.7; found (HPLC MS):
 $[M+H^+] = 546.3$, $[M+Na^+] = 568.3$; Yield: 27 %

PXN 1020

- 153 -

C₂₉H₃₁N₃O₄S; MW: 517.7


PXN 1021



C₃₁H₃₄ClN₃O₄S; MW: 580.15; found (HPLC
MS): [M+H⁺] = 580.3; Yield: 20 %


5

PXN 1022

C₂₉H₃₀ClN₃O₄S; MW: 552.1


10 PXN 1023

- 154 -



$C_{40}H_{40}ClN_5O_4S$; MW: 722.3; found (HPLC MS): $[M+H^+] = 722.6$, $[M+Na^+] = 744.3$; Yield: 4 %

PXN 1024




5

$C_{40}H_{40}ClN_5O_3S$; MW: 706.3; found (HPLC MS): $[M+H^+] = 708.2$; Yield: 1 %

Further modifications of the pyrrolidin-2-one scaffold are possible by using the following procedures:


10

1) Redox variations:

**Scheme 1**

In the presence of NaBH_4 , it is possible to reduce the carboxylic acid function of the MCR-product to the corresponding alcohol (see PXN818-d1). The isolated alcohol can
 5 be further converted to the corresponding ether (alkylation with various halogens) or to the corresponding ester (acylation with acyl chloride) (Scheme 1).


Furthermore, the obtained alcohol can be oxidized to the corresponding aldehyde (Swern oxidation). Alternatively, this aldehyde can also be obtained by selective
 10 reduction of the carboxylic acid. The aldehyde can be converted to numerous further compounds.

**Scheme 2**

15

As can be seen from scheme 2, amines are accessible via a reductive amination process. Further, a Knoevenagel condensation is also possible for modification yielding new substituted pyrrolidin-2-ones as shown in Scheme 3.


- 156 -



Scheme 3


2) Amide variations

- 5 Different amides have been synthesized by aminolysis of the pentafluorophenyl ester using various amines. Other nucleophilic compounds are also suitable to attack the activated carbonyl of the pentafluorophenyl ester, leading to new pyrrolidin-2-one derivatives as shown in Scheme 4.



Scheme 4


3) Reduction of amide



PXN717-d1 has been treated with BMS (dimethylsulfide borane) yielding a mixture of the two compounds PXN723-d1 and PXN724-d1.


4) Homologation of carboxylic acids (Arndt-Eistert Reaction)

- 157 -




Under the conditions of the Arndt-Eistert homologation reaction, formation of the desired product has been observed through HPLC-MS analysis (see PXN845-d1). The obtained carboxylic acid can be further modified as described above.

Using a substituted succinic anhydride in the multicomponent reaction, compounds of formula (I) can be prepared wherein R^7 and/or R^8 are other than hydrogen (see also Org. Lett.2007, 9(20), 4077-4080).



5) Elimination




PXN736-d1 has been treated with 1,1 eq of sodium hydride at room temperature. Thereby, elimination product PXN847-d1 has been isolated and characterized by HPLC-MS. This Product can be used for further modifications (for example Michael-addition).

6) Synthesis of compounds wherein X is N

These compounds may be prepared according to the following scheme:

- 158 -



In addition, compounds of formulas (I), (Ia), (Ic), (Id), (Ie) and (If) may be prepared following the procedures described e.g. in: Synlett, (11), 1883-1885, 2002; Organic Letters, 9(20), 4077-4080, 2007; Organic Letters, 8(18), 3999-4002, 2006; Tetrahedron, 50(36), 10701-8, 1994; Journal of the Chemical Society, Chemical Communications, (5), 386-7, 1987; Journal of the Chemical Society, Chemical Communications, (5), 386-7, 1987; Tetrahedron Letters, 49(35), 5217-5219, 2008 and Journal of Organic Chemistry, 73(14), 5566-5569, 2008.

Synthesis of the starting material 6-chloro-1H-indole-3-carbaldehyde:

This Vilsmeier reaction was performed according to H. G. O. Becker, Organikum, pp. 364-365, Johann Ambrosius Barth Verlag, Heidelberg-Leipzig (1996). To 5 mL DMF in a three-necked flask equipped with a thermometer, 1.8 mL POCl₃ was added dropwise in a temperature range between 15°C and 20°C. Then, a solution of 1g (6.6 mMol) of 6-chloro-1H-indole in 2 mL DMF was added dropwise in a temperature range between 20°C and 30°C. The reaction mixture was stirred for 45 minutes at 37°C. Afterwards, the reaction mixture was poured in a mixture of 15g ice in 10 mL water under stirring. 3.4g NaOH in 18 mL were added between in a temperature range between 20 and 30°C. The resulting mixture was then refluxed for 5 minutes. After cooling to room temperature, the precipitate was filtered off and washed with 10 mL cold water. Crystallization from ethanol yielded 6-chloro-1H-indole-3-carbaldehyde as a white solid (1.04 g, 88%).

Proliferation assay:

5000 PA-1 or PA-1/E6 cells were plated in each well of 96-well flat bottom plates, and incubated overnight at 37°C in 5% CO₂. The growth of plated cells was measured by adding 7.5 microMol of WST-1 reagent (Roche Applied Sciences, Germany) to 3 control wells and measuring the OD₆₅₀ and OD₄₅₀ absorbances with a SpectraMax250

- 159 -

plate reader. If the OD₆₅₀-OD₄₅₀ values were above 0.5, the remainder of the plate was used for incubation with the compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If), other pharmacological agents or solvent controls for 48 hours. After this incubation, the WST-1 reagent was added to the wells of the plate and OD₆₅₀-OD₄₅₀ values were
5 calculated as before. Triplicate wells were assayed for each conditions and standard deviation was determined: all experiments were performed at least three times independently.

Apoptosis Annexin V and Tunel assay:

10 Annexin V and BrdU-incorporation levels were determined with Guava Nexin and Guava Tunel kits using a Guava Personal Cell Analysis System (PCAS, Guava Technologies, Hayward, CA) according to the manufacturer's instruction. 1×10^6 PA-1 and PA-1/E6 cells were cultured in BME media supplemented with 10%FBS and various concentrations of the compounds of formula (I) or DMSO for 24 h. Nutlin-3,
15 racemic (Calbiochem, Roche) at 10 μ M was applied as positive control. For the Guava Nexin assay, cells were trypsinized and collected by centrifuging at 1000 rpm for 5 min at 4 °C. After washing with ice-cold 1 $\times$ Nexin buffer, cells were resuspended in the same buffer, labeled with Annexin V-PE and 7-aminoactinomycin D in the dark on ice for 20 min, and then analyzed with the PCAS. According to the manufacturer protocol
20 for Guava Tunel assay cells were resuspended in 1% paraformaldehyde, incubated on ice for 60 min, washed in ice-cold PBS buffer. Than cells were fixed in ice-cold 70% ethanol for at least 16h at -20°C. After incubation, cells were labeled with BrdU DNA labeling mix for 60 min at 37°C, collected by centrifugation at 1000 rpm for 5 min. Cells were resuspended in anti-BrdU staining mix and incubated at room temperature
25 for 45 min in the dark, and then analyzed with the PCAS.

Apoptosis Assays

Temperature-sensitive H1299 clones were seeded onto 6-well plates at a density of 50,000 cells/well. Saos2 cells were plated at 1×10^6 cells/100-mm plate. Cells were
30 shifted to 32 °C and harvested at the times indicated after temperature shift. Control cells were maintained at 39 °C. TUNEL and multi-caspase assays were conducted


- 160 -

using the Guava Personal Cytometer (Guava Technologies) using the Guava TUNEL and multi-caspase detection kits, using protocols provided by the manufacturer (Guava Technologies) with the compounds of formula (I), (Ia), (Ic), (Id), (Ie) or (If).

- 161 -

Claims

1. A compound of formula (I):



(I)

wherein

V is C=O, C=S or CH₂;

X is sulphur, oxygen or a group of formula CH₂, CR^{4b}R^{4c}, NH, NR^{4b}, SO, SO₂ or a bond;

Y is a group of formula CONR⁶, CH₂NR⁶, CO, COO, CH₂O, SO₂NR⁶, NR⁶CO, NR⁶SO₂, NR^{5a}CONR⁶, NR⁶COO, OCONR⁶, CONR^{5a}NR⁶, CONR^{5a}OR⁶, CH₂CO, CH₂CONR⁶, CH₂COO, COCR^{5a}R⁶ or a bond;

n is 1, 2, 3 or 0;

R¹ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical;

R² is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical;

- 162 -

R^3 is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical;

5 R^4 is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical;

10 R^{4b} is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical;

15 R^{4c} is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical;

20 R^5 is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

R^{5a} is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

25 R^6 is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl radical;

30 the residues R^7 are independently from each other a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;


- 163 -

the residues R^8 are independently from each other a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

- 5 or two of the radicals R^1 , R^2 , R^3 , R^4 , R^{4b} , R^{4c} , R^5 , R^{5a} , R^6 , R^7 and R^8 together are part of an optionally substituted cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heteroaryl, aralkyl or heteroarylalkyl ring system;

10 or a pharmaceutically acceptable salt, ester, solvate or hydrate or a pharmaceutically acceptable formulation thereof.

2. A compound of formula (Ia)



15 wherein

R^1 is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

20

R^2 is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

25

R^3 is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

- 164 -

R⁴ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

- 5 R⁵ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

- 10 R⁶ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl radical;

- 15 or the radicals R⁵ and R⁶ together are part of an optionally substituted heterocycloalkyl, heteroalkylcycloalkyl, heteroaryl or heteroarylalkyl ring system, and/or R² and R³ together are part of an optionally substituted cycloalkyl, alkylcycloalkyl, heterocycloalkyl, heteroalkylcycloalkyl, aralkyl or heteroaralkyl ring system;

or a pharmaceutically acceptable salt, ester, solvate or hydrate or a pharmaceutically acceptable formulation thereof.

20


3. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R¹ is an aryl, heteroaryl, aralkyl or heteroaralkyl radical.

- 25 4. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R¹ is a group of formula –A-Ar or A-Cy wherein A is a bond, C₁-C₄ alkyl or C₁-C₆ heteroalkyl or wherein A is a group of formula –CHR^{1a}- wherein R^{1a} is a C₁-C₆ heteroalkyl group, Cy is an optionally substituted C₃-C₇ cycloalkyl group or an optionally substituted heterocycloalkyl group containing from 3 to 7 ring atoms and Ar is an optionally substituted phenyl ring or an optionally substituted heteroaryl ring
30 containing 5 or 6 ring atoms, especially preferably Ar is an optionally substituted phenyl or an optionally substituted pyridyl residue.

5. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R^2 is hydrogen.

6. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R^3 is an optionally substituted phenyl group, an optionally substituted benzyl group or an optionally substituted heteroaryl residue having 1 or 2 rings and from 5 to 10 ring atoms including 1, 2, 3 or 4 heteroatoms selected from O, S and N.

7. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R^3 has the following structure



wherein E is N or CH, R^{3a} is H, C_1 - C_6 alkyl or C_1 - C_6 heteroalkyl, R^{3b} is H, F, Cl, Br, CH_3 , OCH_3 or CF_3 and R^{3c} is H, F, Cl, Br, CH_3 , OCH_3 or CF_3 .

8. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R^4 is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, optionally substituted C_1 - C_4 alkyl- C_3 - C_7 cycloalkyl, an optionally substituted phenyl ring, an optionally substituted benzyl group or an optionally substituted heteroaryl ring having 5 or 6 ring atoms and including from 1 to 3 heteroatoms selected from O, S and N.

9. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R^5 is an alkyl, heteroalkyl, heterocycloalkyl, heteroalkylcycloalkyl or heteroaralkyl group, all of which groups may be substituted.

10. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R^5 is selected from the following groups: C_1 - C_6 alkyl; heteroalkyl containing 1-6 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S and N; heteroalkylcycloalkyl comprising a C_1 - C_4 alkyl group or a C_1 - C_4 heteroalkyl group and an optionally substituted heterocycloalkyl group containing 5 or 6 ring atoms and 1, 2 or

3 heteroatoms selected from O, S and N; heteroaralkyl comprising a C₁-C₄ alkyl group or a C₁-C₄ heteroalkyl group and an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N; optionally substituted heteroaryl containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected
5 from O, S and N; and optionally substituted heterocycloalkyl containing 5 or 6 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N.

11. A compound of formula (I) or (Ia), according to any one of the preceding claims wherein R⁶ is hydrogen or C₁-C₄ alkyl.

10

12. A compound of formula (I) or (Ia), according to any one of claims 1 to 8 wherein R⁵ and R⁶ together are part of an optionally substituted heterocycloalkyl ring containing 4, 5, 6 or 7 ring atoms and 1, 2 or 3 heteroatoms selected from O, S and N.

15

13. A compound of formula (I) or (Ia), according to any one of claims 1 to 8 wherein R⁵ and R⁶ together are part of an optionally substituted piperazine or piperidine ring.

20

14. A pharmaceutical composition comprising a compound according to anyone of the preceding claims or a pharmaceutically acceptable ester, prodrug, hydrate, solvate or salt thereof, optionally in combination with a pharmaceutically acceptable carrier.

25

15. Use of a compound or a pharmaceutical composition according to anyone of the preceding claims for the preparation of a medicament for the treatment of cancer and/or viral infections.

16. Compound or pharmaceutical composition according to anyone of claims 1 to 14 for use in the treatment of cancer and/or viral infections.