

2,4-PYRIMIDINEDIAMINE COMPOUNDS AND THEIR USES

1. CROSS REFERENCE TO RELATED APPLICATIONS

This application claims benefit under 35 U.S.C. § 119(e) to application Serial No. 60/353,333 filed February 1, 2002; application Serial No. 60/353,267 filed February 1, 2002; application Serial No. 60/399,673 filed July 29, 2002; and application Serial No. 60/434,277 filed December 17, 2002.

2. FIELD OF THE INVENTION

The present invention relates generally to 2,4-pyrimidinediamine compounds, pharmaceutical compositions comprising the compounds, intermediates and synthetic methods of making the compounds and methods of using the compounds and compositions in a variety of contexts.

3. BACKGROUND OF THE INVENTION


Crosslinking of Fc receptors, such as the high affinity receptor for IgE (FcεRI) and/or the high affinity receptor for IgG (FcγRI) activates a signaling cascade in mast, basophil and other immune cells that results in the release of chemical mediators responsible for numerous adverse events. For example, such crosslinking leads to the release of preformed mediators of Type I (immediate) anaphylactic hypersensitivity reactions, such as histamine, from storage sites in granules *via* degranulation. It also leads to the synthesis and release of other mediators, including leukotrienes, prostaglandins and platelet-activating factors (PAFs), that play important roles in inflammatory reactions. Additional mediators that are synthesized and released upon crosslinking Fc receptors include cytokines and nitric oxide.

The signaling cascade(s) activated by crosslinking Fc receptors such as FcεRI and/or FcγRI comprises an array of cellular proteins. Among the most important intracellular signal propagators are the tyrosine kinases. And, an important tyrosine kinase involved in the signal transduction pathways associated with crosslinking the FcεRI and/or FcγRI receptors, as well as other signal transduction cascades, is Syk kinase (*see* Valent *et al.*, 2002, *Intl. J. Hematol.* 75(4):257-362 for review).

As the mediators released as a result of FcεRI and FcγRI receptor cross-linking are responsible for, or play important roles in, the manifestation of numerous adverse events, the availability of compounds capable of inhibiting the signaling cascade(s) responsible for their release would be highly desirable. Moreover, owing to the critical role that Syk kinase plays these and other receptor signaling cascade(s), the availability of compounds capable of inhibiting Syk kinase would also be highly desirable.

4. SUMMARY OF THE INVENTION

In one aspect, the present invention provides novel 2,4-pyrimidinediamine compounds that, as will be discussed in more detail below, have myriad biological activities. The compounds generally comprise a 2,4-pyrimidinediamine “core” having the following structure and numbering convention:




The compounds of the invention are substituted at the C2 nitrogen (N2) to form a secondary amine and are optionally further substituted at one or more of the following positions: the C4 nitrogen (N4), the C5 position and/or the C6 position. When substituted at N4, the substituent forms a secondary amine. The substituent at N2, as well as the optional substituents at the other positions, may range broadly in character and physico-chemical properties. For example, the substituent(s) may be a branched, straight-chained or cyclic alkyl, a branched, straight-chained or cyclic heteroalkyl, a mono- or polycyclic aryl a mono- or polycyclic heteroaryl or combinations of these groups. These substituent groups may be further substituted, as will be described in more detail below.

The N2 and/or N4 substituents may be attached directly to their respective nitrogen atoms, or they may be spaced away from their respective nitrogen atoms *via* linkers, which may be the same or different. The nature of the linkers can vary widely, and can include virtually any combination of atoms or groups useful for spacing one molecular moiety from another. For example, the linker may be an acyclic hydrocarbon bridge (*e.g.*, a saturated or unsaturated alkylene such as methano, ethano, etheno, propano, prop[1]eno, butano, but[1]eno, but[2]eno, buta[1,3]dieno, and the like), a monocyclic or polycyclic hydrocarbon bridge (*e.g.*, [1,2]benzeno, [2,3]naphthaleno, and the like), a simple acyclic heteroatomic or

heteroalkyldiyl bridge (*e.g.*, -O-, -S-, -S-O-, -NH-, -PH-, -C(O)-, -C(O)NH-, -S(O)-, -S(O)₂-, -S(O)NH-, -S(O)₂NH-, -O-CH₂-, -CH₂-O-CH₂-, -O-CH=CH-CH₂-, and the like), a monocyclic or polycyclic heteroaryl bridge (*e.g.*, [3,4]furano, pyridino, thiopheno, piperidino, piperazino, pyrazidino, pyrrolidino, and the like) or combinations of such
 5 bridges.

The substituents at the N2, N4, C5 and/or C6 positions, as well as the optional linkers, may be further substituted with one or more of the same or different substituent groups. The nature of these substituent groups may vary broadly. Non-limiting examples of suitable substituent groups include branched, straight-chain or cyclic alkyls, mono- or
 10 polycyclic aryls, branched, straight-chain or cyclic heteroalkyls, mono- or polycyclic heteroaryls, halos, branched, straight-chain or cyclic haloalkyls, hydroxyls, oxos, thioxos, branched, straight-chain or cyclic alkoxys, branched, straight-chain or cyclic haloalkoxys, trifluoromethoxys, mono- or polycyclic aryloxys, mono- or polycyclic heteroaryloxys, ethers, alcohols, sulfides, thioethers, sulfanyls (thiols), imines, azos, azides, amines
 15 (primary, secondary and tertiary), nitriles (any isomer), cyanates (any isomer), thiocyanates (any isomer), nitrosos, nitros, diazos, sulfoxides, sulfonyls, sulfonic acids, sulfamides, sulfonamides, sulfamic esters, aldehydes, ketones, carboxylic acids, esters, amides, amidines, formadines, amino acids, acetylenes, carbamates, lactones, lactams, glucosides, gluconurides, sulfones, ketals, acetals, thioketals, oximes, oxamic acids, oxamic esters, etc.,
 20 and combinations of these groups. Substituent groups bearing reactive functionalities may be protected or unprotected, as is well-known in the art.

In one illustrative embodiment, the 2,4-pyrimidinediamine compounds of the invention are compounds according to structural formula (I):



25 including salts, hydrates, solvates and N-oxides thereof, wherein:

L¹ and L² are each, independently of one another, selected from the group consisting of a direct bond and a linker;

R² is selected from the group consisting of (C1-C6) alkyl optionally substituted with one or more of the same or different R⁸ groups, (C3-C8) cycloalkyl optionally substituted

with one or more of the same or different R^8 groups, cyclohexyl optionally substituted with one or more of the same or different R^8 groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R^8 groups, (C5-C15) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups;

R^4 is selected from the group consisting of hydrogen, (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C3-C8) cycloalkyl optionally substituted with one or more of the same or different R^8 groups, cyclohexyl optionally substituted with one or more of the same or different R^8 groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R^8 groups, (C5-C15) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups;

R^5 is selected from the group consisting of R^6 , (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C1-C4) alkanyl optionally substituted with one or more of the same or different R^8 groups, (C2-C4) alkenyl optionally substituted with one or more of the same or different R^8 groups and (C2-C4) alkynyl optionally substituted with one or more of the same or different R^8 groups;

each R^6 is independently selected from the group consisting of hydrogen, an electronegative group, $-OR^d$, $-SR^d$, (C1-C3) haloalkyloxy, (C1-C3) perhaloalkyloxy, $-NR^cR^c$, halogen, (C1-C3) haloalkyl, (C1-C3) perhaloalkyl, $-CF_3$, $-CH_2CF_3$, $-CF_2CF_3$, $-CN$, $-NC$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $-N_3$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)NR^cR^c$, $-S(O)_2NR^cR^c$, $-OS(O)R^d$, $-OS(O)_2R^d$, $-OS(O)_2OR^d$, $-OS(O)NR^cR^c$, $-OS(O)_2NR^cR^c$, $-C(O)R^d$, $-C(O)OR^d$, $-C(O)NR^cR^c$, $-C(NH)NR^cR^c$, $-OC(O)R^d$, $-SC(O)R^d$, $-OC(O)OR^d$, $-SC(O)OR^d$, $-OC(O)NR^cR^c$, $-SC(O)NR^cR^c$, $-OC(NH)NR^cR^c$, $-SC(NH)NR^cR^c$, $-[NHC(O)]_nR^d$, $-[NHC(O)]_nOR^d$, $-[NHC(O)]_nNR^cR^c$ and $-[NHC(NH)]_nNR^cR^c$, (C5-C10) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups, (C6-C16) arylalkyl optionally substituted with one or more of the same or different R^8 groups, 5-10 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups and 6-16 membered heteroarylalkyl optionally substituted with one or more of the same or different R^8 groups;

R^8 is selected from the group consisting of R^a , R^b , R^a substituted with one or more of the same or different R^a or R^b , $-OR^a$ substituted with one or more of the same or different R^a or R^b , $-B(OR^a)_2$, $-B(NR^cR^c)_2$, $-(CH_2)_m-R^b$, $-(CHR^a)_m-R^b$, $-O-(CH_2)_m-R^b$, $-S-(CH_2)_m-R^b$, $-O-CHR^aR^b$, $-O-CR^a(R^b)_2$, $-O-(CHR^a)_m-R^b$, $-O-(CH_2)_m-CH[(CH_2)_mR^b]R^b$, $-S-(CHR^a)_m-R^b$,
 5 $-C(O)NH-(CH_2)_m-R^b$, $-C(O)NH-(CHR^a)_m-R^b$, $-O-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$,
 $-S-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$, $-O-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$,
 $-S-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$, $-NH-(CH_2)_m-R^b$, $-NH-(CHR^a)_m-R^b$, $-NH[(CH_2)_mR^b]$,
 $-N[(CH_2)_mR^b]_2$, $-NH-C(O)-NH-(CH_2)_m-R^b$, $-NH-C(O)-(CH_2)_m-CHR^bR^b$ and
 $-NH-(CH_2)_m-C(O)-NH-(CH_2)_m-R^b$;

10 each R^a is independently selected from the group consisting of hydrogen, (C1-C6) alkyl, (C3-C8) cycloalkyl, cyclohexyl, (C4-C11) cycloalkylalkyl, (C5-C10) aryl, phenyl, (C6-C16) arylalkyl, benzyl, 2-6 membered heteroalkyl, 3-8 membered cycloheteroalkyl, morpholinyl, piperazinyl, homopiperazinyl, piperidinyl, 4-11 membered cycloheteroalkylalkyl, 5-10 membered heteroaryl and 6-16 membered heteroarylalkyl;

15 each R^b is a suitable group independently selected from the group consisting of $=O$, $-OR^d$, (C1-C3) haloalkyloxy, $-OCF_3$, $=S$, $-SR^d$, $=NR^d$, $=NOR^d$, $-NR^cR^c$, halogen, $-CF_3$, $-CN$, $-NC$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)NR^cR^c$, $-S(O)_2NR^cR^c$, $-OS(O)R^d$, $-OS(O)_2R^d$, $-OS(O)_2OR^d$, $-OS(O)_2NR^cR^c$, $-C(O)R^d$, $-C(O)OR^d$, $-C(O)NR^cR^c$, $-C(NH)NR^cR^c$, $-C(NR^a)NR^cR^c$, $-C(NOH)R^a$, $-C(NOH)NR^cR^c$, $-OC(O)R^d$,
 20 $-OC(O)OR^d$, $-OC(O)NR^cR^c$, $-OC(NH)NR^cR^c$, $-OC(NR^a)NR^cR^c$, $-[NHC(O)]_nR^d$,
 $-[NR^aC(O)]_nR^d$, $-[NHC(O)]_nOR^d$, $-[NR^aC(O)]_nOR^d$, $-[NHC(O)]_nNR^cR^c$, $-[NR^aC(O)]_nNR^cR^c$,
 $-[NHC(NH)]_nNR^cR^c$ and $-[NR^aC(NR^a)]_nNR^cR^c$;

each R^c is independently a protecting group or R^a , or, alternatively, each R^c is taken together with the nitrogen atom to which it is bonded to form a 5 to 8-membered
 25 cycloheteroalkyl or heteroaryl which may optionally include one or more of the same or different additional heteroatoms and which may optionally be substituted with one or more of the same or different R^a or suitable R^b groups;

each R^d is independently a protecting group or R^a ;

each m is independently an integer from 1 to 3; and

30 each n is independently an integer from 0 to 3.


In another aspect, the present invention provides prodrugs of the 2,4-pyrimidinediamine compounds. Such prodrugs may be active in their prodrug form, or may be inactive until converted under physiological or other conditions of use to an active drug

form. In the prodrugs of the invention, one or more functional groups of the 2,4-pyrimidinediamine compounds are included in promoieties that cleave from the molecule under the conditions of use, typically by way of hydrolysis, enzymatic cleavage or some other cleavage mechanism, to yield the functional groups. For example, primary or secondary amino groups may be included in an amide promoiety that cleaves under conditions of use to generate the primary or secondary amino group. Thus, the prodrugs of the invention include special types of protecting groups, termed "progroups," masking one or more functional groups of the 2,4-pyrimidinediamine compounds that cleave under the conditions of use to yield an active 2,4-pyrimidinediamine drug compound. Functional groups within the 2,4-pyrimidinediamine compounds that may be masked with progroups for inclusion in a promoiety include, but are not limited to, amines (primary and secondary), hydroxyls, sulfanyls (thiols), carboxyls, carbonyls, phenols, catechols, diols, alkynes, phosphates, etc. Myriad progroups suitable for masking such functional groups to yield promoieties that are cleavable under the desired conditions of use are known in the art. All of these progroups, alone or in combinations, may be included in the prodrugs of the invention. Specific examples of promoieties that yield primary or secondary amine groups that can be included in the prodrugs of the invention include, but are not limited to amides, carbamates, imines, ureas, phosphenyls, phosphoryls and sulfenyls. Specific examples of promoieties that yield sulfanyl groups that can be included in the prodrugs of the invention include, but are not limited to, thioethers, for example S-methyl derivatives (monothio, dithio, oxythio, aminothio acetals), silyl thioethers, thioesters, thiocarbonates, thiocarbamates, asymmetrical disulfides, etc. Specific examples of promoieties that cleave to yield hydroxyl groups that can be included in the prodrugs of the invention include, but are not limited to, sulfonates, esters and carbonates. Specific examples of promoieties that yield carboxyl groups that can be included in the prodrugs of the invention included, but are not limited to, esters (including silyl esters, oxamic acid esters and thioesters), amides and hydrazides.

In one illustrative embodiment, the prodrugs of the invention are compounds according to structural formula (I) in which the protecting group of R^c and R^d is a progroup.

Replacing the hydrogens attached to N2 and N4 in the 2,4-pyrimidinediamines of structural formula (I) with substituents adversely affects the activity of the compounds. However, as will be appreciated by skilled artisans, these nitrogens may be included in promoieties that, under conditions of use, cleave to yield 2,4-pyrimidinediamines according

to structural formula (I). Thus, in another illustrative embodiment, the prodrugs of the invention are compounds according to structural formula (II):




including salts, hydrates, solvates and N-oxides thereof, wherein:

5 R^2 , R^4 , R^5 , R^6 , L^1 and L^2 are as previously defined for structural formula (I); and R^{2b} and R^{4b} are each, independently of one another, a progroup.

In another aspect, the present invention provides compositions comprising one or more compounds and/or prodrugs of the invention and an appropriate carrier, excipient or diluent. The exact nature of the carrier, excipient or diluent will depend upon the desired use for the composition, and may range from being suitable or acceptable for veterinary
10 uses to being suitable or acceptable for human use.


In still another aspect, the present invention provides intermediates useful for synthesizing the 2,4-pyrimidinediamine compounds and prodrugs of the invention. In one embodiment, the intermediates are 4-pyrimidineamines according to structural formula (III):



15


including salts, hydrates, solvates and N-oxides thereof, wherein R^4 , R^5 , R^6 and L^2 are as previously defined for structural formula (I); LG is a leaving group such as, for example, $-S(O)_2Me$, $-SMe$ or halo (e.g., F, Cl, Br, I); and R^{4c} is hydrogen or a progroup.

In another embodiment, the intermediates are 2-pyrimidineamines according to
20 structural formula (IV):




including salts, hydrates, solvates and N-oxides thereof, wherein R^2 , R^5 , R^6 and L^1 are as previously defined for structural formula (I); LG is a leaving group, such as, for example, $-S(O)_2Me$, $-SMe$ or halo (*e.g.*, F, Cl, Br, I) and R^{2c} is hydrogen or a progroup.

In yet another embodiment, the intermediates are 4-amino- or 4-hydroxy-2-
5 pyrimidineamines according to structural formula (V):



including salts, hydrates, solvates and N-oxides thereof, wherein R^2 , R^5 , R^6 and L^1 are as previously defined for structural formula (I), R^7 is an amino or hydroxyl group and
10 R^{2c} is hydrogen or a progroup.

In another embodiment, the intermediates are N4-substituted cytosines according to structural formula (VI):



15 including salts, hydrates, solvates and N-oxides thereof, wherein R^4 , R^5 , R^6 and L^2 are as previously defined for structural formula (I) and R^{4c} is hydrogen or a progroup.

In yet another aspect, the present invention provides methods of synthesizing the 2,4-pyrimidinediamine compounds and prodrugs of the invention. In one embodiment, the method involves reacting a 4-pyrimidineamine according to structural formula (III) with an
20 amine of the formula $HR^{2c}N-L^1-R^2$, where L^1 , R^2 and R^{2c} are as previously defined for structural formula (IV) to yield a 2,4-pyrimidinediamine according to structural formula (I) or a prodrug according to structural formula (II).

In another embodiment, the method involves reacting a 2-pyrimidineamine according to structural formula (IV) with an amine of the formula $R^4-L^2-NHR^{4c}$ where L^4 ,
25 R^4 and R^{4c} are as previously defined for structural formula (III) to yield a 2,4-

pyrimidinediamine according to structural formula (I) or a prodrug according to structural formula (II).

In yet another embodiment, the method involves reacting a 4-amino-2-pyrimidineamine according to structural formula (V) (in which R^7 is an amino group) with an amine of the formula $R^4-L^2-NHR^{4c}$, where L^2 , R^4 and R^{4c} are as defined for structural formula (III), to yield a 2,4-pyrimidinediamine according to structural formula (I) or a prodrug according to structural formula (II). Alternatively, the 4-amino-2-pyrimidineamine may be reacted with a compound of the formula R^4-L^2-LG , where R^4 and L^2 are as previously defined for structural formula (I) and LG is a leaving group.

In still another embodiment, the method involves halogenating a 4-hydroxy-2-pyrimidineamine according to structural formula (V) (R^7 is a hydroxyl group) to yield a 2-pyrimidineamine according to structural formula (IV) and reacting this pyrimidineamine with an appropriate amine, as described above.

In yet another embodiment, the method involves halogenating an N4-substituted cytosine according to structural formula (VI) to yield a 4-pyrimidineamine according to structural formula (III) and reacting this pyrimidineamine with an appropriate amine, as described above.

The 2,4-pyrimidinediamine compounds of the invention are potent inhibitors of degranulation of immune cells, such as mast, basophil, neutrophil and/or eosinophil cells. Thus, in still another aspect, the present invention provides methods of regulating, and in particular inhibiting, degranulation of such cells. The method generally involves contacting a cell that degranulates with an amount of a 2,4-pyrimidinediamine compound or prodrug of the invention, or an acceptable salt, hydrate, solvate, N-oxide and/or composition thereof, effective to regulate or inhibit degranulation of the cell. The method may be practiced in *in vitro* contexts or in *in vivo* contexts as a therapeutic approach towards the treatment or prevention of diseases characterized by, caused by or associated with cellular degranulation.

While not intending to be bound by any theory of operation, biochemical data confirm that the 2,4-pyrimidinediamine compounds exert their degranulation inhibitory effect, at least in part, by blocking or inhibiting the signal transduction cascade(s) initiated by crosslinking of the high affinity Fc receptors for IgE ("Fc ϵ RI") and/or IgG ("Fc γ RI"). Indeed, the 2,4-pyrimidinediamine compounds are potent inhibitors of both Fc ϵ RI-mediated and Fc γ RI-mediated degranulation. As a consequence, the 2,4-pyrimidine compounds may be used to inhibit these Fc receptor signalling cascades in any cell type expressing such

FcεRI and/or FcγRI receptors including but not limited to macrophages, mast, basophil, neutrophil and/or eosinophil cells.

The methods also permit the regulation of, and in particular the inhibition of, downstream processes that result as a consequence of activating such Fc receptor signaling cascade(s). Such downstream processes include, but are not limited to, FcεRI-mediated and/or FcγRI-mediated degranulation, cytokine production and/or the production and/or release of lipid mediators such as leukotrienes and prostaglandins. The method generally involves contacting a cell expressing an Fc receptor, such as one of the cell types discussed above, with an amount of a 2,4-pyrimidinediamine compound or prodrug of the invention, or an acceptable salt, hydrate, solvent, N-oxide and/or composition thereof, effective to regulate or inhibit the Fc receptor signaling cascade and/or a downstream process effected by the activation of this signaling cascade. The method may be practiced in *in vitro* contexts or in *in vivo* contexts as a therapeutic approach towards the treatment or prevention of diseases characterized by, caused by or associated with the Fc receptor signaling cascade, such as diseases effected by the release of granule specific chemical mediators upon degranulation, the release and/or synthesis of cytokines and/or the release and/or synthesis of lipid mediators such as leukotrienes and prostaglandins.

In yet another aspect, the present invention provides methods of treating and/or preventing diseases characterized by, caused by or associated with the release of chemical mediators as a consequence of activating Fc receptor signaling cascades, such as FcεRI and/or FcγRI- signaling cascades. The methods may be practiced in animals in veterinary contexts or in humans. The methods generally involve administering to an animal subject or human an amount of a 2,4-pyrimidinediamine compound or prodrug of the invention, or an acceptable salt, hydrate, solvate, N-oxide and/or composition thereof, effective to treat or prevent the disease. As discussed previously, activation of the FcεRI or FcγRI receptor signaling cascade in certain immune cells leads to the release and/or synthesis of a variety of chemical substances that are pharmacological mediators of a wide variety of diseases. Any of these diseases may be treated or prevented according to the methods of the invention.

For example, in mast cells and basophil cells, activation of the FcεRI or FcγRI signaling cascade leads to the immediate (*i.e.*, within 1-3 min. of receptor activation) release of preformed mediators of atopic and/or Type I hypersensitivity reactions (*e.g.*, histamine, proteases such as tryptase, etc.) *via* the degranulation process. Such atopic or Type I hypersensitivity reactions include, but are not limited to, anaphylactic reactions to

environmental and other allergens (e.g., pollens, insect and/or animal venoms, foods, drugs, contrast dyes, etc.), anaphylactoid reactions, hay fever, allergic conjunctivitis, allergic rhinitis, allergic asthma, atopic dermatitis, eczema, urticaria, mucosal disorders, tissue disorders and certain gastrointestinal disorders.

5 The immediate release of the preformed mediators *via* degranulation is followed by the release and/or synthesis of a variety of other chemical mediators, including, among other things, platelet activating factor (PAF), prostaglandins and leukotrienes (e.g., LTC₄) and the *de novo* synthesis and release of cytokines such as TNF α , IL-4, IL-5, IL-6, IL-13, etc. The first of these two processes occurs approximately 3-30 min. following receptor activation;
10 the latter approximately 30 min. – 7 hrs. following receptor activation. These “late stage” mediators are thought to be in part responsible for the chronic symptoms of the above-listed atopic and Type I hypersensitivity reactions, and in addition are chemical mediators of inflammation and inflammatory diseases (e.g., osteoarthritis, inflammatory bowel disease, ulcerative colitis, Crohn’s disease, idiopathic inflammatory bowel disease, irritable bowel
15 syndrome, spastic colon, etc.), low grade scarring (e.g., scleroderma, increased fibrosis, keloids, post-surgical scars, pulmonary fibrosis, vascular spasms, migraine, reperfusion injury and post myocardial infarction), and sicca complex or syndrome. All of these diseases may be treated or prevented according to the methods of the invention.

 Additional diseases which can be treated or prevented according to the methods of
20 the invention include diseases associated with basophil cell and/or mast cell pathology. Examples of such diseases include, but are not limited to, diseases of the skin such as scleroderma, cardiac diseases such as post myocardial infarction, pulmonary diseases such as pulmonary muscle changes or remodeling and chronic obstructive pulmonary disease (COPD) and diseases of the gut such as inflammatory bowel syndrome (spastic colon).

25 The 2,4-pyrimidinediamine compounds of the invention are also potent inhibitors of the tyrosine kinase Syk kinase. Thus, in still another aspect, the present invention provides methods of regulating, and in particular inhibiting, Syk kinase activity. The method generally involves contacting a Syk kinase or a cell comprising a Syk kinase with an amount of a 2,4-pyrimidinediamine compound or prodrug of the invention, or an acceptable
30 salt, hydrate, solvate, N-oxide and/or composition thereof, effective to regulate or inhibit Syk kinase activity. In one embodiment, the Syk kinase is an isolated or recombinant Syk kinase. In another embodiment, the Syk kinase is an endogenous or recombinant Syk kinase expressed by a cell, for example a mast cell or a basophil cell. The method may be

practiced in *in vitro* contexts or in *in vivo* contexts as a therapeutic approach towards the treatment or prevention of diseases characterized by, caused by or associated with Syk kinase activity.

While not intending to be bound by any particular theory of operation, it is believed that the 2,4-pyrimidinediamine compounds of the invention inhibit cellular degranulation and/or the release of other chemical mediators primarily by inhibiting Syk kinase that gets activated through the gamma chain homodimer of FcεRI (*see, e.g.*, FIG. 2). This gamma chain homodimer is shared by other Fc receptors, including FcγRI, FcγRIII and FcαRI. For all of these receptors, intracellular signal transduction is mediated by the common gamma chain homodimer. Binding and aggregation of those receptors results in the recruitment and activation of tyrosine kinases such as Syk kinase. As a consequence of these common signaling activities, the 2,4-pyrimidinediamine compounds described herein may be used to regulate, and in particular inhibit, the signaling cascades of Fc receptors having this gamma chain homodimer, such as FcεRI, FcγRI, FcγRIII and FcαRI, as well as the cellular responses elicited through these receptors.

Syk kinase is known to play a critical role in other signaling cascades. For example, Syk kinase is an effector of B-cell receptor (BCR) signaling (Turner *et al.*, 2000, *Immunology Today* 21:148-154) and is an essential component of integrin beta(1), beta(2) and beta(3) signaling in neutrophils (Mocsai *et al.*, 2002, *Immunity* 16:547-558). As the 2,4-pyrimidinediamine compounds described herein are potent inhibitors of Syk kinase, they can be used to regulate, and in particular inhibit, any signaling cascade where Syk plays a role, such as, for example, the Fc receptor, BCR and integrin signaling cascades, as well as the cellular responses elicited through these signaling cascades. The particular cellular response regulated or inhibited will depend, in part, on the specific cell type and receptor signaling cascade, as is well known in the art. Non-limiting examples of cellular responses that may be regulated or inhibited with the 2,4-pyrimidinediamine compounds include a respiratory burst, cellular adhesion, cellular degranulation, cell spreading, cell migration, phagocytosis (*e.g.*, in macrophages), calcium ion flux (*e.g.*, in mast, basophil, neutrophil, eosinophil and B-cells), platelet aggregation, and cell maturation (*e.g.*, in B-cells).

Thus, in another aspect, the present invention provides methods of regulating, and in particular inhibiting, signal transduction cascades in which Syk plays a role. The method generally involves contacting a Syk-dependent receptor or a cell expressing a Syk-dependent receptor with an amount of a 2,4-pyrimidinediamine compound or prodrug of the

invention, or an acceptable salt, hydrate, solvate, N-oxide and/or composition thereof, effective to regulate or inhibit the signal transduction cascade. The methods may also be used to regulate, and in particular inhibit, downstream processes or cellular responses elicited by activation of the particular Syk-dependent signal transduction cascade. The methods may be practiced to regulate any signal transduction cascade where Syk is not known or later discovered to play a role. The methods may be practiced in *in vitro* contexts or in *in vivo* contexts as a therapeutic approach towards the treatment or prevention of diseases characterized by, caused by or associated with activation of the Syk-dependent signal transduction cascade. Non-limited examples of such diseases include those previously discussed.

5. BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 provides a cartoon illustrating allergen-induced production of IgE and consequent release of preformed and other chemical mediators from mast cells;

FIG. 2 provides a cartoon illustrating the FcεR1 signal transduction cascade leading to degranulation of mast and/or basophil cells;

FIG. 3 provides a cartoon illustrating the putative points of action of compounds that selectively inhibit upstream FcεRI-mediated degranulation and compounds that inhibit both FcεRI-mediated and ionomycin-induced degranulation;

FIG. 4 provides graphs illustrating the effects of certain 2,4-pyrimidinediamine compounds, DMSO (control) and ionomycin on Ca^{2+} flux in CHMC cells;

FIG. 5 provides graphs illustrating the immediacy of the inhibitory activity of compounds R921218 and R926495;

FIG. 6 provides a graph illustrating the effect of washout on the inhibitory activity of compounds R921218 and R921302;

FIG. 7 provides data showing that varying concentrations of compounds R921218 (A) and R921219 (B) inhibit phosphorylation of various proteins downstream of Syk kinase in the IgE receptor signal transduction cascade in activated BMHC cells;

FIG. 8 provides data showing dose responsive inhibition of Syk kinase phosphorylation of an endogenous substrate (LAT) and a peptide substrate in the presence of increasing concentrations of compounds R921218 (X), R921219 (Y) and R921304 (Z);

FIG. 9 provides data showing that the inhibition of Syk kinase by compound R921219 is ATP competitive;

FIG. 10 provides data showing that varying concentrations of compounds R921219 (A) and R218218 (B) inhibit phosphorylation of proteins downstream of Syk kinase, but not LYN kinase, in the FcεRI signal transduction cascade in activated CHMC cells; also shown is inhibition of phosphorylation of proteins downstream of LYN kinase but not Syk kinase, in the presence of a known LYN kinase inhibitor (PP2); and

FIGS. 11A-D provide data showing inhibition of phosphorylation of proteins downstream of Syk kinase in the FcεRI signal transduction cascade in BMDC cells.

6. DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

6.1 Definitions

As used herein, the following terms are intended to have the following meanings:

“Alkyl” by itself or as part of another substituent refers to a saturated or unsaturated branched, straight-chain or cyclic monovalent hydrocarbon radical having the stated number of carbon atoms (*i.e.*, C1-C6 means one to six carbon atoms) that is derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane, alkene or alkyne.

Typical alkyl groups include, but are not limited to, methyl; ethyls such as ethanyl, ethenyl, ethynyl; propyls such as propan-1-yl, propan-2-yl, cyclopropan-1-yl, prop-1-en-1-yl, prop-1-en-2-yl, prop-2-en-1-yl, cycloprop-1-en-1-yl; cycloprop-2-en-1-yl, prop-1-yn-1-yl, prop-2-yn-1-yl, etc.; butyls such as butan-1-yl, butan-2-yl, 2-methyl-propan-1-yl, 2-methyl-propan-2-yl, cyclobutan-1-yl, but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl, etc.; and the like. Where specific levels of saturation are intended, the nomenclature “alkanyl,” “alkenyl” and/or “alkynyl” is used, as defined below. In preferred embodiments, the alkyl groups are (C1-C6) alkyl.

“Alkanyl” by itself or as part of another substituent refers to a saturated branched, straight-chain or cyclic alkyl derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane. Typical alkanyl groups include, but are not limited to, methanyl; ethanyl; propanyls such as propan-1-yl, propan-2-yl (isopropyl), cyclopropan-1-yl, etc.; butanyls such as butan-1-yl, butan-2-yl (*sec*-butyl), 2-methyl-propan-1-yl (isobutyl), 2-methyl-propan-2-yl (*t*-butyl), cyclobutan-1-yl, etc.; and the like. In preferred embodiments, the alkanyl groups are (C1-C6) alkanyl.

“Alkenyl” by itself or as part of another substituent refers to an unsaturated branched, straight-chain or cyclic alkyl having at least one carbon-carbon double bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkene. The group may be in either the *cis* or *trans* conformation about the double bond(s). Typical alkenyl groups include, but are not limited to, ethenyl; propenyls such as prop-1-en-1-yl , prop-1-en-2-yl, prop-2-en-1-yl, prop-2-en-2-yl, cycloprop-1-en-1-yl; cycloprop-2-en-1-yl ; butenyls such as but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, etc.; and the like. In preferred embodiments, the alkenyl group is (C2-C6) alkenyl.

“Alkynyl” by itself or as part of another substituent refers to an unsaturated branched, straight-chain or cyclic alkyl having at least one carbon-carbon triple bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkyne. Typical alkynyl groups include, but are not limited to, ethynyl; propynyls such as prop-1-yn-1-yl , prop-2-yn-1-yl, etc.; butynyls such as but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl , etc.; and the like. In preferred embodiments, the alkynyl group is (C2-C6) alkynyl.

“Alkyldiyl” by itself or as part of another substituent refers to a saturated or unsaturated, branched, straight-chain or cyclic divalent hydrocarbon group having the stated number of carbon atoms (*i.e.*, C1-C6 means from one to six carbon atoms) derived by the removal of one hydrogen atom from each of two different carbon atoms of a parent alkane, alkene or alkyne, or by the removal of two hydrogen atoms from a single carbon atom of a parent alkane, alkene or alkyne. The two monovalent radical centers or each valency of the divalent radical center can form bonds with the same or different atoms. Typical alkyldiyl groups include, but are not limited to, methandiyl; ethyldiyls such as ethan-1,1-diyl, ethan-1,2-diyl, ethen-1,1-diyl, ethen-1,2-diyl; propyldiyls such as propan-1,1-diyl, propan-1,2-diyl, propan-2,2-diyl, propan-1,3-diyl, cyclopropan-1,1-diyl, cyclopropan-1,2-diyl, prop-1-en-1,1-diyl, prop-1-en-1,2-diyl, prop-2-en-1,2-diyl, prop-1-en-1,3-diyl, cycloprop-1-en-1,2-diyl, cycloprop-2-en-1,2-diyl, cycloprop-2-en-1,1-diyl, prop-1-yn-1,3-diyl, etc.; butyldiyls such as, butan-1,1-diyl, butan-1,2-diyl, butan-1,3-diyl, butan-1,4-diyl, butan-2,2-diyl, 2-methyl-propan-1,1-diyl, 2-methyl-propan-1,2-diyl, cyclobutan-1,1-diyl; cyclobutan-1,2-diyl, cyclobutan-1,3-diyl, but-1-en-1,1-diyl, but-1-en-1,2-diyl, but-1-en-1,3-diyl, but-1-en-1,4-diyl, 2-methyl-prop-1-en-1,1-diyl, 2-methanylidene-propan-1,1-diyl, buta-1,3-dien-1,1-diyl,

buta-1,3-dien-1,2-diyl, buta-1,3-dien-1,3-diyl, buta-1,3-dien-1,4-diyl, cyclobut-1-en-1,2-diyl, cyclobut-1-en-1,3-diyl, cyclobut-2-en-1,2-diyl, cyclobuta-1,3-dien-1,2-diyl, cyclobuta-1,3-dien-1,3-diyl, but-1-yn-1,3-diyl, but-1-yn-1,4-diyl, buta-1,3-diyn-1,4-diyl, etc.; and the like. Where specific levels of

5 saturation are intended, the nomenclature alkanyldiyl, alkenyldiyl and/or alkynyldiyl is used. Where it is specifically intended that the two valencies are on the same carbon atom, the nomenclature "alkylidene" is used. In preferred embodiments, the alkylidiyl group is (C1-C6) alkylidiyl. Also preferred are saturated acyclic alkanyldiyl groups in which the radical centers are at the terminal carbons, *e.g.*, methandiyl (methano); ethan-1,2-diyl

10 (ethano); propan-1,3-diyl (propano); butan-1,4-diyl (butano); and the like (also referred to as alkylenos, defined *infra*).

"Alkyleno" by itself or as part of another substituent refers to a straight-chain saturated or unsaturated alkylidiyl group having two terminal monovalent radical centers derived by the removal of one hydrogen atom from each of the two terminal carbon atoms

15 of straight-chain parent alkane, alkene or alkyne. The locant of a double bond or triple bond, if present, in a particular alkyleno is indicated in square brackets. Typical alkyleno groups include, but are not limited to, methano; ethylenos such as ethano, etheno, ethyno; propylenos such as propano, prop[1]eno, propa[1,2]dieno, prop[1]yno, etc.; butylenos such as butano, but[1]eno, but[2]eno, buta[1,3]dieno, but[1]yno, but[2]yno, buta[1,3]diyno, etc.;

20 and the like. Where specific levels of saturation are intended, the nomenclature alkano, alkeno and/or alkyno is used. In preferred embodiments, the alkyleno group is (C1-C6) or (C1-C3) alkyleno. Also preferred are straight-chain saturated alkano groups, *e.g.*, methano, ethano, propano, butano, and the like.

"Heteroalkyl," "Heteroalkanyl," "Heteroalkenyl," "Heteroalkynyl," "Heteroalkylidiyl" and "Heteroalkyleno" by themselves or as part of another substituent refer to alkyl, alkanyl, alkenyl, alkynyl, alkylidiyl and alkyleno groups, respectively, in which one or more of the carbon atoms are each independently replaced with the same or different heteroatoms or heteroatomic groups. Typical heteroatoms and/or heteroatomic groups which can replace the carbon atoms include, but are not limited to, -O-, -S-, -S-O-, -NR'-, -PH-, -S(O)-,

25 -S(O)₂-, -S(O)NR'-, -S(O)₂NR'-, and the like, including combinations thereof, where each R' is independently hydrogen or (C1-C6) alkyl.

"Cycloalkyl" and "Heterocycloalkyl" by themselves or as part of another substituent refer to cyclic versions of "alkyl" and "heteroalkyl" groups, respectively. For heteroalkyl groups, a heteroatom can occupy the position that is attached to the remainder of the

molecule. Typical cycloalkyl groups include, but are not limited to, cyclopropyl; cyclobutyls such as cyclobutanyl and cyclobutenyl; cyclopentyls such as cyclopentanyl and cyclopentenyl; cyclohexyls such as cyclohexanyl and cyclohexenyl; and the like. Typical heterocycloalkyl groups include, but are not limited to, tetrahydrofuranyl (*e.g.*,
5 tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, etc.), piperidiny (*e.g.*, piperidin-1-yl, piperidin-2-yl, etc.), morpholinyl (*e.g.*, morpholin-3-yl, morpholin-4-yl, etc.), piperazinyl (*e.g.*, piperazin-1-yl, piperazin-2-yl, etc.), and the like.

“Acyclic Heteroatomic Bridge” refers to a divalent bridge in which the backbone atoms are exclusively heteroatoms and/or heteroatomic groups. Typical acyclic
10 heteroatomic bridges include, but are not limited to, -O-, -S-, -S-O-, -NR’-, -PH-, -S(O)-, -S(O)₂-, -S(O) NR’-, -S(O)₂NR’-, and the like, including combinations thereof, where each R’ is independently hydrogen or (C1-C6) alkyl.

“Parent Aromatic Ring System” refers to an unsaturated cyclic or polycyclic ring system having a conjugated π electron system. Specifically included within the definition
15 of “parent aromatic ring system” are fused ring systems in which one or more of the rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example, fluorene, indane, indene, phenalene, tetrahydronaphthalene, etc. Typical parent aromatic ring systems include, but are not limited to, aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene,
20 fluorene, hexacene, hexaphene, hexalene, indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, tetrahydronaphthalene, triphenylene, trinaphthalene, and the like, as well as the various hydro isomers thereof.

25 “Aryl” by itself or as part of another substituent refers to a monovalent aromatic hydrocarbon group having the stated number of carbon atoms (*i.e.*, C5-C15 means from 5 to 15 carbon atoms) derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. Typical aryl groups include, but are not limited to, groups derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene,
30 benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, *as*-indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene,

trinaphthalene, and the like, as well as the various hydro isomers thereof. In preferred embodiments, the aryl group is (C5-C15) aryl, with (C5-C10) being even more preferred. Particularly preferred aryls are cyclopentadienyl, phenyl and naphthyl.

“Arylaryl” by itself or as part of another substituent refers to a monovalent hydrocarbon group derived by the removal of one hydrogen atom from a single carbon atom of a ring system in which two or more identical or non-identical parent aromatic ring systems are joined directly together by a single bond, where the number of such direct ring junctions is one less than the number of parent aromatic ring systems involved. Typical arylaryl groups include, but are not limited to, biphenyl, triphenyl, phenyl-naphthyl, binaphthyl, biphenyl-naphthyl, and the like. Where the number of carbon atoms in an arylaryl group are specified, the numbers refer to the carbon atoms comprising each parent aromatic ring. For example, (C5-C15) arylaryl is an arylaryl group in which each aromatic ring comprises from 5 to 15 carbons, *e.g.*, biphenyl, triphenyl, binaphthyl, phenyl-naphthyl, etc. Preferably, each parent aromatic ring system of an arylaryl group is independently a (C5-C15) aromatic, more preferably a (C5-C10) aromatic. Also preferred are arylaryl groups in which all of the parent aromatic ring systems are identical, *e.g.*, biphenyl, triphenyl, binaphthyl, trinaphthyl, etc.

“Biaryl” by itself or as part of another substituent refers to an arylaryl group having two identical parent aromatic systems joined directly together by a single bond. Typical biaryl groups include, but are not limited to, biphenyl, binaphthyl, bianthracyl, and the like. Preferably, the aromatic ring systems are (C5-C15) aromatic rings, more preferably (C5-C10) aromatic rings. A particularly preferred biaryl group is biphenyl.

“Arylalkyl” by itself or as part of another substituent refers to an acyclic alkyl group in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with an aryl group. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, 2-phenylethen-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, 2-naphthylethen-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. Where specific alkyl moieties are intended, the nomenclature arylalkanyl, arylakenyl and/or arylalkynyl is used. In preferred embodiments, the arylalkyl group is (C6-C21) arylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is (C1-C6) and the aryl moiety is (C5-C15). In particularly preferred embodiments the arylalkyl group is (C6-C13), *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is (C1-C3) and the aryl moiety is (C5-C10).

“Parent Heteroaromatic Ring System” refers to a parent aromatic ring system in which one or more carbon atoms are each independently replaced with the same or different heteroatoms or heteroatomic groups. Typical heteroatoms or heteroatomic groups to replace the carbon atoms include, but are not limited to, N, NH, P, O, S, S(O), S(O)₂, Si, etc. Specifically included within the definition of “parent heteroaromatic ring systems” are fused ring systems in which one or more of the rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example, benzodioxan, benzofuran, chromane, chromene, indole, indoline, xanthene, etc. Also included in the definition of “parent heteroaromatic ring system” are those recognized rings that include common substituents, such as, for example, benzopyrone and 1-methyl-1,2,3,4-tetrazole. Specifically excluded from the definition of “parent heteroaromatic ring system” are benzene rings fused to cyclic polyalkylene glycols such as cyclic polyethylene glycols. Typical parent heteroaromatic ring systems include, but are not limited to, acridine, benzimidazole, benzisoxazole, benzodioxan, benzodioxole, benzofuran, benzopyrone, benzothiadiazole, benzothiazole, benzotriazole, benzoxazine, benzoxazole, benzoxazoline, carbazole, β -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazole, xanthene, and the like.

“Heteroaryl” by itself or as part of another substituent refers to a monovalent heteroaromatic group having the stated number of ring atoms (e.g., “5-14 membered” means from 5 to 14 ring atoms) derived by the removal of one hydrogen atom from a single atom of a parent heteroaromatic ring system. Typical heteroaryl groups include, but are not limited to, groups derived from acridine, benzimidazole, benzisoxazole, benzodioxan, benzodioxole, benzofuran, benzopyrone, benzothiadiazole, benzothiazole, benzotriazole, benzoxazine, benzoxazole, benzoxazoline, carbazole, β -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole,

thiophene, triazole, xanthene, and the like, as well as the various hydro isomers thereof. In preferred embodiments, the heteroaryl group is a 5-14 membered heteroaryl, with 5-10 membered heteroaryl being particularly preferred.

“Heteroaryl-Heteroaryl” by itself or as part of another substituent refers to a monovalent heteroaromatic group derived by the removal of one hydrogen atom from a single atom of a ring system in which two or more identical or non-identical parent heteroaromatic ring systems are joined directly together by a single bond, where the number of such direct ring junctions is one less than the number of parent heteroaromatic ring systems involved. Typical heteroaryl-heteroaryl groups include, but are not limited to, bipyridyl, tripyridyl, pyridylpurinyl, bipurinyl, etc. Where the number of atoms are specified, the numbers refer to the number of atoms comprising each parent heteroaromatic ring systems. For example, 5-15 membered heteroaryl-heteroaryl is a heteroaryl-heteroaryl group in which each parent heteroaromatic ring system comprises from 5 to 15 atoms, *e.g.*, bipyridyl, tripyridyl, etc. Preferably, each parent heteroaromatic ring system is independently a 5-15 membered heteroaromatic, more preferably a 5-10 membered heteroaromatic. Also preferred are heteroaryl-heteroaryl groups in which all of the parent heteroaromatic ring systems are identical.

“Biheteroaryl” by itself or as part of another substituent refers to a heteroaryl-heteroaryl group having two identical parent heteroaromatic ring systems joined directly together by a single bond. Typical biheteroaryl groups include, but are not limited to, bipyridyl, bipurinyl, biquinoliny, and the like. Preferably, the heteroaromatic ring systems are 5-15 membered heteroaromatic rings, more preferably 5-10 membered heteroaromatic rings.

“Heteroarylalkyl” by itself or as part of another substituent refers to an acyclic alkyl group in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with a heteroaryl group. Where specific alkyl moieties are intended, the nomenclature heteroarylalkanyl, heteroarylakenyl and/or heteroarylalkynyl is used. In preferred embodiments, the heteroarylalkyl group is a 6-21 membered heteroarylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the heteroarylalkyl is (C1-C6) alkyl and the heteroaryl moiety is a 5-15-membered heteroaryl. In particularly preferred embodiments, the heteroarylalkyl is a 6-13 membered heteroarylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety is (C1-C3) alkyl and the heteroaryl moiety is a 5-10 membered heteroaryl.

“Halogen” or “Halo” by themselves or as part of another substituent, unless otherwise stated, refer to fluoro, chloro, bromo and iodo.

“Haloalkyl” by itself or as part of another substituent refers to an alkyl group in which one or more of the hydrogen atoms is replaced with a halogen. Thus, the term “haloalkyl” is meant to include monohaloalkyls, dihaloalkyls, trihaloalkyls, etc. up to perhaloalkyls. For example, the expression “(C1-C2) haloalkyl” includes fluoromethyl, difluoromethyl, trifluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl, 1,2-difluoroethyl, 1,1,1-trifluoroethyl, perfluoroethyl, etc.

The above-defined groups may include prefixes and/or suffixes that are commonly used in the art to create additional well-recognized substituent groups. As examples, “alkyloxy” or “alkoxy” refers to a group of the formula -OR”, “alkylamine” refers to a group of the formula -NHR” and “dialkylamine” refers to a group of the formula -NR”R”, where each R” is independently an alkyl. As another example, “haloalkoxy” or “haloalkyloxy” refers to a group of the formula -OR’”, where R’” is a haloalkyl.

“Protecting group” refers to a group of atoms that, when attached to a reactive functional group in a molecule, mask, reduce or prevent the reactivity of the functional group. Typically, a protecting group may be selectively removed as desired during the course of a synthesis. Examples of protecting groups can be found in Greene and Wuts, *Protective Groups in Organic Chemistry*, 3rd Ed., 1999, John Wiley & Sons, NY and Harrison *et al.*, *Compendium of Synthetic Organic Methods*, Vols. 1-8, 1971-1996, John Wiley & Sons, NY. Representative amino protecting groups include, but are not limited to, formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (“CBZ”), *tert*-butoxycarbonyl (“Boc”), trimethylsilyl (“TMS”), 2-trimethylsilyl-ethanesulfonyl (“TES”), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (“Fmoc”), nitro-veratryloxycarbonyl (“NVOC”) and the like. Representative hydroxyl protecting groups include, but are not limited to, those where the hydroxyl group is either acylated or alkylated such as benzyl and trityl ethers, as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers (*e.g.*, TMS or TIPPS groups) and allyl ethers.

“Prodrug” refers to a derivative of an active 2,4-pyrimidinediamine compound (drug) that requires a transformation under the conditions of use, such as within the body, to release the active 2,4-pyrimidinediamine drug. Prodrugs are frequently, but not necessarily, pharmacologically inactive until converted into the active drug. Prodrugs are typically obtained by masking a functional group in the 2,4-pyrimidinediamine drug believed to be in part required for activity with a progroup (defined below) to form a promoiety which

undergoes a transformation, such as cleavage, under the specified conditions of use to release the functional group, and hence the active 2,4-pyrimidinediamine drug. The cleavage of the promoiety may proceed spontaneously, such as by way of a hydrolysis reaction, or it may be catalyzed or induced by another agent, such as by an enzyme, by light, by acid or base, or by a change of or exposure to a physical or environmental parameter, such as a change of temperature. The agent may be endogenous to the conditions of use, such as an enzyme present in the cells to which the prodrug is administered or the acidic conditions of the stomach, or it may be supplied exogenously.

A wide variety of progroups, as well as the resultant promoieties, suitable for masking functional groups in the active 2,4-pyrimidinediamines compounds to yield prodrugs are well-known in the art. For example, a hydroxyl functional group may be masked as a sulfonate, ester or carbonate promoiety, which may be hydrolyzed *in vivo* to provide the hydroxyl group. An amino functional group may be masked as an amide, carbamate, imine, urea, phosphenyl, phosphoryl or sulfenyl promoiety, which may be hydrolyzed *in vivo* to provide the amino group. A carboxyl group may be masked as an ester (including silyl esters and thioesters), amide or hydrazide promoiety, which may be hydrolyzed *in vivo* to provide the carboxyl group. Other specific examples of suitable progroups and their respective promoieties will be apparent to those of skill in the art.

“Progroup” refers to a type of protecting group that, when used to mask a functional group within an active 2,4-pyrimidinediamine drug to form a promoiety, converts the drug into a prodrug. Progroups are typically attached to the functional group of the drug *via* bonds that are cleavable under specified conditions of use. Thus, a progroup is that portion of a promoiety that cleaves to release the functional group under the specified conditions of use. As a specific example, an amide promoiety of the formula —NH—C(O)CH_3 comprises the progroup —C(O)CH_3 .

“Fc Receptor” refers to a member of the family of cell surface molecules that binds the Fc portion (containing the specific constant region) of an immunoglobulin. Each Fc receptor binds immunoglobulins of a specific type. For example the $\text{Fc}\alpha$ receptor (“ $\text{Fc}\alpha\text{R}$ ”) binds IgA, the $\text{Fc}\epsilon$ R binds IgE and the $\text{Fc}\gamma$ R binds IgG.

The $\text{Fc}\alpha\text{R}$ family includes the polymeric Ig receptor involved in epithelial transport of IgA/IgM, the myeloid specific receptor $\text{R}\alpha\text{RI}$ (also called CD89), the $\text{Fc}\alpha/\mu\text{R}$ and at least two alternative IgA receptors (for a recent review see Monteiro & van de Winkel, 2003, Annu. Rev. Immunol, advanced e-publication. The $\text{Fc}\alpha\text{RI}$ is expressed on neutrophils, eosinophils, monocytes/macrophages, dendritic cells and kupfer cells. The

Fc α RI includes one alpha chain and the Fc γ R gamma homodimer that bears an activation motif (ITAM) in the cytoplasmic domain and phosphorylates Syk kinase.

The Fc ϵ R family includes two types, designated Fc ϵ RI and Fc ϵ RII (also known as CD23). Fc ϵ RI is a high affinity receptor (binds IgE with an affinity of about 10^{10}M^{-1}) found on mast, basophil and eosinophil cells that anchors monomeric IgE to the cell surface. The Fc ϵ RI possesses one alpha chain, one beta chain and the gamma chain homodimer discussed above. The Fc ϵ RII is a low affinity receptor expressed on mononuclear phagocytes, B lymphocytes, eosinophils and platelets. The Fc ϵ RII comprises a single polypeptide chain and does not include the gamma chain homodimer.

The Fc γ R family includes three types, designated Fc γ RI (also known as CD64), Fc γ RII (also known as CD32) and Fc γ RIII (also known as CD16). Fc γ RI is a high affinity receptor (binds IgG1 with an affinity of 10^8M^{-1}) found on mast, basophil, mononuclear, neutrophil, eosinophil, dendritic and phagocyte cells that anchors monomeric IgG to the cell surface. The Fc γ RI includes one alpha chain and the gamma chain dimer shared by Fc α RI and Fc ϵ RI.

The Fc γ RII is a low affinity receptor expressed on neutrophils, monocytes, eosinophils, platelets and B lymphocytes. The Fc γ RII includes one alpha chain, and does not include the gamma chain homodimer discussed above.

The Fc γ RIII is a low affinity (binds IgG1 with an affinity of $5 \times 10^5\text{M}^{-1}$) expressed on NK, eosinophil, macrophage, neutrophil and mast cells. It comprises one alpha chain and the gamma homodimer shared by Fc α RI, Fc ϵ RI and Fc γ RI.

Skilled artisans will recognize that the subunit structure and binding properties of these various Fc receptors, cell types expressing them, are not completely characterized. The above discussion merely reflects the current state-of-the-art regarding these receptors (*see, e.g.*, Immunobiology: The Immune System in Health & Disease, 5th Edition, Janeway et al., Eds, 2001, ISBN 0-8153-3642-x, Figure 9.30 at pp. 371), and is not intended to be limiting with respect to the myriad receptor signaling cascades that can be regulated with the compounds described herein.

“Fc Receptor-Mediated Degranulation” or “Fc Receptor-Induced Degranulation” refers to degranulation that proceeds *via* an Fc receptor signal transduction cascade initiated by crosslinking of an Fc receptor.

“IgE-Induced Degranulation” or “Fc ϵ RI-Mediated Degranulation” refers to degranulation that proceeds *via* the IgE receptor signal transduction cascade initiated by

crosslinking of FcεRI-bound IgE. The crosslinking may be induced by an IgE-specific allergen or other multivalent binding agent, such as an anti-IgE antibody. Referring to FIG. 2, in mast and/or basophil cells, the FcεRI signaling cascade leading to degranulation may be broken into two stages: upstream and downstream. The upstream stage includes all of the processes that occur prior to calcium ion mobilization (illustrated as “Ca²⁺” in FIG. 2; see also FIG. 3). The downstream stage includes calcium ion mobilization and all processes downstream thereof. Compounds that inhibit FcεRI-mediated degranulation may act at any point along the FcεRI-mediated signal transduction cascade. Compounds that selectively inhibit upstream FcεRI-mediated degranulation act to inhibit that portion of the FcεRI signaling cascade upstream of the point at which calcium ion mobilization is induced. In cell-based assays, compounds that selectively inhibit upstream FcεRI-mediated degranulation inhibit degranulation of cells such as mast or basophil cells that are activated or stimulated with an IgE-specific allergen or binding agent (such as an anti-IgE antibody) but do not appreciably inhibit degranulation of cells that are activated or stimulated with degranulating agents that bypass the FcεRI signaling pathway, such as, for example the calcium ionophores ionomycin and A23187.

“IgG-Induced Degranulation” or “FcγRI-Mediated Degranulation” refers to degranulation that proceeds *via* the FcγRI signal transduction cascade initiated by crosslinking of FcγRI-bound IgG. The crosslinking may be induced by an IgG-specific allergen or another multivalent binding agent, such as an anti-IgG or fragment antibody. Like the FcεRI signaling cascade, in mast and basophil cells the FcγRI signaling cascade also leads to degranulation which may be broken into the same two stages: upstream and downstream. Similar to FcεRI-mediated degranulation, compounds that selectively inhibit upstream FcγRI-mediated degranulation act upstream of the point at which calcium ion mobilization is induced. In cell-based assays, compounds that selectively inhibit upstream FcγRI-mediated degranulation inhibit degranulation of cells such as mast or basophil cells that are activated or stimulated with an IgG-specific allergen or binding agent (such as an anti-IgG antibody or fragment) but do not appreciably inhibit degranulation of cells that are activated or stimulated with degranulating agents that bypass the FcγRI signaling pathway, such as, for example the calcium ionophores ionomycin and A23187.

“Ionophore-Induced Degranulation” or “Ionophore-Mediated Degranulation” refers to degranulation of a cell, such as a mast or basophil cell, that occurs upon exposure to a calcium ionophore such as, for example, ionomycin or A23187.

“Syk Kinsase” refers to the well-known 72kDa non-receptor (cytoplasmic) spleen protein tyrosine kinase expressed in B-cells and other hematopoietic cells. Syk kinase includes two consensus Src-homology 2 (SH2) domains in tandem that bind to phosphorylated immunoreceptor tyrosine-based activation motifs (“ITAMs”), a “linker” domain and a catalytic domain (for a review of the structure and function of Syk kinase see Sada *et al.*, 2001, J. Biochem. (Tokyo) 130:177-186); see also Turner *et al.*, 2000, Immunology Today 21:148-154). Syk kinase has been extensively studied as an effector of B-cell receptor (BCR) signaling (Turner *et al.*, 2000, *supra*). Syk kinase is also critical for tyrosine phosphorylation of multiple proteins which regulate important pathways leading from immunoreceptors, such as Ca^{2+} mobilization and mitogen-activated protein kinase (MAPK) cascades (see, e.g., FIG. 2) and degranulation. Syk kinase also plays a critical role in integrin signaling in neutrophils (*see, e.g.*, Mocsai *et al.* 2002, Immunity 16:547-558).


As used herein, Syk kinase includes kinases from any species of animal, including but not limited to, homosapiens, simian, bovine, porcine, rodent, etc., recognized as belonging to the Syk family. Specifically included are isoforms, splice variants, allelic variants, mutants, both naturally occurring and man-made. The amino acid sequences of such Syk kinases are well known and available from GENBANK. Specific examples of mRNAs encoding different isoforms of human Syk kinase can be found at GENBANK accession no. gi|21361552|ref|NM__003177.2|, gi|496899|emb|Z29630.1|HSSYKPTK[496899] and gi|15030258|gb|BC011399.1|BC011399[15030258], which are incorporated herein by reference.

Skilled artisans will appreciate that tyrosine kinases belonging to other families may have active sites or binding pockets that are similar in three-dimensional structure to that of Syk. As a consequence of this structural similarity, such kinases, referred to herein as “Syk mimics,” are expected to catalyze phosphorylation of substrates phosphorylated by Syk. Thus, it will be appreciated that such Syk mimics, signal transduction cascades in which such Syk mimics play a role and biological responses effected by such Syk mimics and Syk mimic-dependent signaling cascades may be regulated, and in particular inhibited, with the 2,4-pyrimidinediamine compounds described herein.

“Syk-Dependent Signaling Cascade” refers to a signal transduction cascade in which Syk kinase plays a role. Non-limiting examples of such Syk-dependent signaling cascades include the $\text{Fc}\alpha\text{RI}$, $\text{Fc}\epsilon\text{RI}$, $\text{Fc}\gamma\text{RI}$, $\text{Fc}\gamma\text{RIII}$, BCR and integrin signaling cascades.

6.2 The 2,4-Pyrimidinediamine Compounds

The compounds of the invention are generally 2,4-pyrimidinediamine compounds according to structural formula (I):



5 including salts, hydrates, solvates and N-oxides thereof, wherein:

L¹ and L² are each, independently of one another, selected from the group consisting of a direct bond and a linker;

R² is selected from the group consisting of (C1-C6) alkyl optionally substituted with one or more of the same or different R⁸ groups, (C3-C8) cycloalkyl optionally substituted with one or more of the same or different R⁸ groups, cyclohexyl optionally substituted with one or more of the same or different R⁸ groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R⁸ groups, (C5-C15) aryl optionally substituted with one or more of the same or different R⁸ groups, phenyl optionally substituted with one or more of the same or different R⁸ groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R⁸ groups;

R⁴ is selected from the group consisting of hydrogen, (C1-C6) alkyl optionally substituted with one or more of the same or different R⁸ groups, (C3-C8) cycloalkyl optionally substituted with one or more of the same or different R⁸ groups, cyclohexyl optionally substituted with one or more of the same or different R⁸ groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R⁸ groups, (C5-C15) aryl optionally substituted with one or more of the same or different R⁸ groups, phenyl optionally substituted with one or more of the same or different R⁸ groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R⁸ groups;

R⁵ is selected from the group consisting of R⁶, (C1-C6) alkyl optionally substituted with one or more of the same or different R⁸ groups, (C1-C4) alkanyl optionally substituted with one or more of the same or different R⁸ groups, (C2-C4) alkenyl optionally substituted with one or more of the same or different R⁸ groups and (C2-C4) alkynyl optionally substituted with one or more of the same or different R⁸ groups;

each R^6 is independently selected from the group consisting of hydrogen, an electronegative group, $-OR^d$, $-SR^d$, (C1-C3) haloalkyloxy, (C1-C3) perhaloalkyloxy, $-NR^cR^c$, halogen, (C1-C3) haloalkyl, (C1-C3) perhaloalkyl, $-CF_3$, $-CH_2CF_3$, $-CF_2CF_3$, $-CN$, $-NC$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $-N_3$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)NR^cR^c$, $-S(O)_2NR^cR^c$, $-OS(O)R^d$, $-OS(O)_2R^d$, $-OS(O)_2OR^d$, $-OS(O)NR^cR^c$, $-OS(O)_2NR^cR^c$, $-C(O)R^d$, $-C(O)OR^d$, $-C(O)NR^cR^c$, $-C(NH)NR^cR^c$, $-OC(O)R^d$, $-SC(O)R^d$, $-OC(O)OR^d$, $-SC(O)OR^d$, $-OC(O)NR^cR^c$, $-SC(O)NR^cR^c$, $-OC(NH)NR^cR^c$, $-SC(NH)NR^cR^c$, $-[NHC(O)]_nR^d$, $-[NHC(O)]_nOR^d$, $-[NHC(O)]_nNR^cR^c$ and $-[NHC(NH)]_nNR^cR^c$, (C5-C10) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups, (C6-C16) arylalkyl optionally substituted with one or more of the same or different R^8 groups, 5-10 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups and 6-16 membered heteroarylalkyl optionally substituted with one or more of the same or different R^8 groups;

R^8 is selected from the group consisting of R^a , R^b , R^a substituted with one or more of the same or different R^a or R^b , $-OR^a$ substituted with one or more of the same or different R^a or R^b , $-B(OR^a)_2$, $-B(NR^cR^c)_2$, $-(CH_2)_m-R^b$, $-(CHR^a)_m-R^b$, $-O-(CH_2)_m-R^b$, $-S-(CH_2)_m-R^b$, $-O-CHR^aR^b$, $-O-CR^a(R^b)_2$, $-O-(CHR^a)_m-R^b$, $-O-(CH_2)_m-CH[(CH_2)_mR^b]R^b$, $-S-(CHR^a)_m-R^b$, $-C(O)NH-(CH_2)_m-R^b$, $-C(O)NH-(CHR^a)_m-R^b$, $-O-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$, $-S-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$, $-O-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$, $-S-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$, $-NH-(CH_2)_m-R^b$, $-NH-(CHR^a)_m-R^b$, $-NH[(CH_2)_mR^b]$, $-N[(CH_2)_mR^b]_2$, $-NH-C(O)-NH-(CH_2)_m-R^b$, $-NH-C(O)-(CH_2)_m-CHR^bR^b$ and $-NH-(CH_2)_m-C(O)-NH-(CH_2)_m-R^b$;

each R^a is independently selected from the group consisting of hydrogen, (C1-C6) alkyl, (C3-C8) cycloalkyl, cyclohexyl, (C4-C11) cycloalkylalkyl, (C5-C10) aryl, phenyl, (C6-C16) arylalkyl, benzyl, 2-6 membered heteroalkyl, 3-8 membered cycloheteroalkyl, morpholinyl, piperazinyl, homopiperazinyl, piperidinyl, 4-11 membered cycloheteroalkylalkyl, 5-10 membered heteroaryl and 6-16 membered heteroarylalkyl;

each R^b is a suitable group independently selected from the group consisting of $=O$, $-OR^d$, (C1-C3) haloalkyloxy, $-OCF_3$, $=S$, $-SR^d$, $=NR^d$, $=NOR^d$, $-NR^cR^c$, halogen, $-CF_3$, $-CN$, $-NC$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)NR^cR^c$, $-S(O)_2NR^cR^c$, $-OS(O)R^d$, $-OS(O)_2R^d$, $-OS(O)_2OR^d$, $-OS(O)_2NR^cR^c$, $-C(O)R^d$, $-C(O)OR^d$, $-C(O)NR^cR^c$, $-C(NH)NR^cR^c$, $-C(NR^a)NR^cR^c$, $-C(NOH)R^a$, $-C(NOH)NR^cR^c$, $-OC(O)R^d$, $-OC(O)OR^d$, $-OC(O)NR^cR^c$, $-OC(NH)NR^cR^c$, $-OC(NR^a)NR^cR^c$, $-[NHC(O)]_nR^d$,

$-\text{[NR}^a\text{C(O)]}_n\text{R}^d$, $-\text{[NHC(O)]}_n\text{OR}^d$, $-\text{[NR}^a\text{C(O)]}_n\text{OR}^d$, $-\text{[NHC(O)]}_n\text{NR}^c\text{R}^c$, $-\text{[NR}^a\text{C(O)]}_n\text{NR}^c\text{R}^c$,
 $-\text{[NHC(NH)]}_n\text{NR}^c\text{R}^c$ and $-\text{[NR}^a\text{C(NR}^a)]_n\text{NR}^c\text{R}^c$;

each R^c is independently R^a , or, alternatively, each R^c is taken together with the nitrogen atom to which it is bonded to form a 5 to 8-membered cycloheteroalkyl or

5 heteroaryl which may optionally include one or more of the same or different additional heteroatoms and which is optionally substituted with one or more of the same or different R^a or suitable R^b groups;

each R^d is independently R^a ;

each m is independently an integer from 1 to 3; and


10 each n is independently an integer from 0 to 3.

In the compounds of structural formula (I), L^1 and L^2 represent, independently of one another, a direct bond or a linker. Thus, as will be appreciated by skilled artisans, the substituents R^2 and/or R^4 may be bonded either directly to their respective nitrogen atoms or, alternatively, spaced away from their respective nitrogen atoms by way of a linker. The identity of the linker is not critical and typical suitable linkers include, but are not limited to, (C1-C6) alkylidiyls, (C1-C6) alkanos and (C1-C6) heteroalkylidiyls, each of which may be optionally substituted with one or more of the same or different R^8 groups, where R^8 is as previously defined for structural formula (I). In a specific embodiment, L^1 and L^2 are each, independently of one another, selected from the group consisting of a direct bond, (C1-C3) alkylidiyl optionally substituted with one or more of the same or different R^a , suitable R^b or R^9 groups and 1-3 membered heteroalkylidiyl optionally substituted with one or more of the same or different R^a , suitable R^b or R^9 groups, wherein R^9 is selected from the group consisting of (C1-C3) alkyl, $-\text{OR}^a$, $-\text{C(O)OR}^a$, (C5-C10) aryl optionally substituted with one or more of the same or different halogens, phenyl optionally substituted with one or more of the same or different halogens, 5-10 membered heteroaryl optionally substituted with one or more of the same or different halogens and 6 membered heteroaryl optionally substituted with one or more of the same or different halogens; and R^a and R^b are as previously defined for structural formula (I). Specific R^9 groups that may be used to substitute L^1 and L^2 include $-\text{OR}^a$, $-\text{C(O)OR}^a$, phenyl, halophenyl and 4-halophenyl, wherein R^a is as previously defined for structural formula (I).

In another specific embodiment, L^1 and L^2 are each, independently of one another, selected from the group consisting of methano, ethano and propano, each of which may be optionally monosubstituted with an R^9 group, where R^9 is as previously defined above.


In all of the above embodiments, specific R^a groups that may be included in R^9 groups are selected from the group consisting of hydrogen, (C1-C6) alkyl, phenyl and benzyl.

In still another specific embodiment, L^1 and L^2 are each a direct bond such that the 2,4-pyrimidinediamine compounds of the invention are compounds according to structural formula (Ia):



including salts, hydrates, solvates and N-oxides thereof, wherein R^2 , R^4 , R^5 and R^6 are as previously defined for structural formula (I). Additional specific embodiments of the 2,4-pyrimidinediamine compounds of the invention are described below.

In a first embodiment of the compounds of structural formulae (I) and (Ia), R^2 , R^4 , R^5 , R^6 , L^1 and L^2 are as previously defined for their respective structures (I) and (Ia), with the proviso that R^2 is not 3,4,5-trimethoxyphenyl, 3,4,5-tri (C1-C6) alkoxyphenyl or




where R^{21} , R^{22} and R^{23} are as defined for R^1 , R^2 and R^3 , respectively of U.S. Patent No. 6,235,746, the disclosure of which is incorporated by reference. In a specific embodiment of this first embodiment, R^{21} is hydrogen, halo, straight-chain or branched (C1-C6) alkyl optionally substituted with one or more of the same or different R^{25} groups, hydroxyl, (C1-C6) alkoxy optionally substituted with one or more of the same or different phenyl or R^{25} groups, thiol (-SH), (C1-C6) alkylthio optionally substituted with one or more of the same or different phenyl or R^{25} groups, amino (-NH₂), -NHR²⁶ or -NR²⁶R²⁶; R^{22} and R^{23} are each, independently of one another, a (C1-C6) straight-chain or branched alkyl optionally substituted with one or more of the same or different R^{25} groups; R^{25} is selected from the group consisting of halo, hydroxyl, (C1-C6) alkoxy, thiol, (C1-C6) alkylthio, (C1-C6) alkylamino and (C1-C6) dialkylamino; and each R^{26} is independently a (C1-C6) alkyl optionally substituted with one or more of the same or different phenyl or R^{25} groups or a

-C(O)R²⁷, where R²⁷ is a (C1-C6) alkyl optionally substituted with one or more of the same or different phenyl or R²⁵ groups.

In another specific embodiment of this first embodiment, R²¹ is methoxy optionally substituted with one or more of the same or different halo groups and/or R²² and R²³ are each, independently of one another, a methyl or ethyl optionally substituted with one or more of the same or different halo groups.

In a second embodiment of the compounds of structural formulae (I) and (Ia), R², R⁴, R⁵ and L² are as previously described for their respective structures (I) and (Ia), L¹ is a direct bond and R⁶ is hydrogen, with the proviso that R² is not 3,4,5-trimethoxyphenyl, 3,4,5-tri (C1-C6) alkoxyphenyl or




where R²¹, R²² and R²³ are as defined above, in connection with the first embodiment.

In a third embodiment, the 2,4-pyrimidinediamine compounds of structural formulae (I) and (Ia) exclude one or more of the following compounds:

- N2,N4-bis(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R070790);
- N2,N4-bis(2-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R081166);
- N2,N4-bis(4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R088814);
- N2,N4-bis(2-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R088815);
- N2,N4-bisphenyl-5-fluoro-2,4-pyrimidinediamine (R091880);
- N2,N4-bis(3-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R092788);
- N2,N4-bis(3-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R067962);
- N2,N4-bis(2,5-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067963);
- N2,N4-bis(3,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067964);
- N2,N4-bis(4-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R0707153);
- N2,N4-bis(2,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R070791);
- N2,N4-bis(3-bromophenyl)-5-fluoro-2,4-pyrimidinediamine (R008958);
- N2,N4-bis(phenyl)-5-fluoro-2,4-pyrimidinediamine;
- N2,N4-bis(morpholino)-5-fluoro-2,4-pyrimidinediamine; and
- N2,N4-bis[(3-chloro-4-methoxyphenyl)]-5-fluoro-2,4-pyrimidinediamine.

In a fourth embodiment, the compounds of structural formulae (I) and (Ia) exclude compounds according to the following structural formula (Ib):



wherein R^{24} is (C1-C6) alkyl; and R^{21} , R^{22} and R^{23} are as previously defined in connection with the first embodiment.

In a fifth embodiment, the compounds of structural formulae (I) and (Ia) exclude the compounds described in Examples 1-141 of U.S. Patent No. 6,235,746, the disclosure of which is incorporated herein by reference.

In a sixth embodiment, the compounds of structural formulae (I) and (Ia) exclude compounds defined by formula (1) or formula 1(a) of this U.S. Patent No. 6,235,746 (see, e.g., the disclosure at Col. 1, line 48 through Col. 7, line 49 and Col. 8, lines 9-36, which is incorporated by reference).

In a seventh embodiment, the compounds of structural formulae (I) and (Ia) exclude compounds in which R^5 is cyano or $-C(O)NHR$, where R is hydrogen or (C1-C6) alkyl, when R^2 is a substituted phenyl; R^4 is a substituted or unsubstituted (C1-C6) alkyl, (C₃-C₈) cycloalkyl, 3-8 membered cycloheteralkyl or 5-15 membered heteroaryl; and R^6 is hydrogen.

In an eighth embodiment, the compounds of structural formulae (I) and (Ia) exclude the compounds defined by formulae (I) and (X) of WO 02/04429 or any compound disclosed in WO 02/04429, the disclosure of which is incorporated herein by reference.

In a ninth embodiment of the compounds of structural formulae (I) and (Ia), when R^5 is cyano or $-C(O)NHR$, where R is hydrogen or (C1-C6) alkyl; and R^6 is hydrogen, then R^2 is other than a substituted phenyl group.

In a tenth embodiment, the compounds of structural formulae (I) and (Ia) exclude compounds in which R^2 and R^4 are each independently a substituted or unsubstituted pyrrole or indole ring which is attached to the remainder of the molecule *via* its ring nitrogen atom.

In an eleventh embodiment, the compounds of structural formulae (I) and (Ia) exclude compounds defined by formulae (I) and (IV) of U.S. Patent No. 4,983,608 or any

compound disclosed in U.S. Patent No. 4,983,608, the disclosure of which is incorporated herein by reference.

Those of skill in the art will appreciate that in the compounds of formulae (I) and (Ia), R^2 and R^4 may be the same or different, and may vary broadly. When R^2 and/or R^4 are optionally substituted rings, such as optionally substituted cycloalkyls, cycloheteroalkyls, aryls and heteroaryl, the ring may be attached to the remainder of the molecule through any available carbon or heteroatom. The optional substituents may be attached to any available carbon atoms and/or heteroatoms.

In a twelfth embodiment of the compounds of structural formulae (I) and (Ia), R^2 and/or R^4 is an optionally substituted phenyl or an optionally substituted (C5-C15) aryl, subject to the provisos that (1) when R^6 is hydrogen, then R^2 is not 3,4,5-trimethoxyphenyl or 3,4,5-tri (C1-C6) alkoxyphenyl; (2) when R^2 is a 3,4,5-trisubstituted phenyl, then the substituents at the 3- and 4-positions are not simultaneously methoxy or (C1-C6) alkoxy; or (3) when R^6 is hydrogen and R^4 is (C1-C6) alkyl, (C₃-C₈) cycloalkyl, 3-8 membered cycloheteroalkyl or 5-15 membered heteroaryl, then R^5 is other than cyano. Alternatively, R^2 is subject to the provisos described in connection with the first or second embodiments. The optionally substituted aryl or phenyl group may be attached to the remainder of the molecule through any available carbon atom. Specific examples of optionally substituted phenyls include phenyls that are optionally mono-, di- or tri-substituted with the same or different R^8 groups, where R^8 is as previously defined for structural formula (I) and subject to the above provisos. When the phenyl is mono-substituted, the R^8 substituent may be positioned at either the *ortho*, *meta* or *para* position. When positioned at the *ortho*, *meta* or *para* position, R^8 is preferably selected from the group consisting of (C1-C10) alkyl, (C1-C10) branched alkyl, $-OR^a$ optionally substituted with one or more of the same or different R^b groups, $-O-C(O)OR^a$, $-O-(CH_2)_m-C(O)OR^a$, $-C(O)OR^a$, $-O-(CH_2)_m-NR^cR^c$, $-O-C(O)NR^cR^c$, $-O-(CH_2)_m-C(O)NR^cR^c$, $-O-C(NH)NR^cR^c$, $-O-(CH_2)_m-C(NH)NR^cR^c$ and $-NH-(CH_2)_m-NR^cR^c$, where m , R^a and R^c are as previously defined for structural formula (I). In one embodiment of these compounds, $-NR^cR^c$ is a 5-6 membered heteroaryl which optionally includes one or more of the same or different additional heteroatoms. Specific examples of such 5-6 membered heteroaryls include, but are not limited to, oxadiazolyl, triazolyl, thiazolyl, oxazolyl, tetrazolyl and isoxazolyl.

In another embodiment of these compounds, $-NR^cR^c$ is a 5-6 membered saturated cycloheteroalkyl ring which optionally includes one or more of the same or different

heteroatoms. Specific examples of such cycloheteroalkyls include, but are not limited to, pyrrolidinyl, pyrazolidinyl, imidazolidinyl, piperidinyl, piperazinyl and morpholinyl.

In still another embodiment of these compounds, each R^a is independently a (C1-C6) alkyl and/or each $-NR^cR^c$ is $-NHR^a$, where R^a is a (C1-C6) alkyl. In one specific

5 embodiment, R^8 is $-O-CH_2-C(O)NHCH_3$. In another specific embodiment R^8 is $-OH$.

When the phenyl is di-substituted or tri-substituted, the R^8 substituents may be positioned at any combination of positions. For example, the R^8 substituents may be positioned at the 2,3-, 2,4-, 2,5-, 2,6-, 3,4-, 3,5-, 2,3,4-, 2,3,5-, 2,3,6-, 2,4,5-, 2,4,6-, 2,5,6- or 3,4,5-positions. In one embodiment of compounds including a disubstituted phenyl, the

10 substituents are positioned other than 3,4. In another embodiment they are positioned 3,4. In one embodiment of compounds including a trisubstituted phenyl, the substituents are positioned other than 3,4,5 or, alternatively, no two of the substituents are positioned 3,4. In another embodiment, the substituents are positioned 3,4,5.

Specific examples of R^8 substituents in such di- and trisubstituted phenyls include

15 the various R^8 substituents described above in connection with the *ortho*, *meta* and *para* substituted phenyls.

In another specific embodiment, R^8 substituents useful for substituting such di- and trisubstituted phenyls include (C1-C6) alkyl, (C1-C6) alkoxy, methoxy, halo, chloro, (C1-C6) perhaloalkyl, $-CF_3$, (C1-C6) perhaloalkoxy and $-OCF_3$. In a preferred embodiment,

20 such R^8 substituents are positioned 3,4 or 3,5. Specific examples of preferred di-substituted phenyl rings include 3-chloro-4-methoxy-phenyl, 3-methoxy-4-chlorophenyl, 3-chloro-4-trifluoromethoxy-phenyl, 3-trifluoromethoxy-4-chloro-phenyl, 3,4-dichloro-phenyl, 3,4-dimethoxyphenyl and 3,5-dimethoxyphenyl, with the provisos that: (1) when R^4 is one of the above-identified phenyls, and R^5 and R^6 are each hydrogen, then


25 R^2 is not 3,4,5-tri(C1-C6)alkoxyphenyl or 3,4,5-trimethoxyphenyl; (2) when R^2 is 3,4-dimethoxyphenyl and R^5 and R^6 are each hydrogen, then R^4 is not 3-(C1-C6)alkoxyphenyl, 3-methoxyphenyl, 3,4-di-(C1-C6) alkoxyphenyl or 3,4-dimethoxyphenyl; (3) when R^4 is 3-chloro-4-methoxyphenyl and R^5 is halo or fluoro, and optionally R^6 is hydrogen, then R^2 is not 3-chloro-4-(C1-C6)alkoxyphenyl or

30 3-chloro-4-methoxyphenyl; (4) when R^4 is 3,4-dichlorophenyl, R^5 is hydrogen, (C1-C6) alkyl, methyl, halo or chloro and optionally R^6 is hydrogen, then R^2 is not a phenyl mono substituted at the *para* position with a (C1-C6) alkoxy group which is optionally substituted with one or more of the same or different R^b , $-OH$ or $-NR^cR^c$ groups, where R^b and R^c are as

previously described for structural formula (I); and/or (5) R^2 and/or R^4 is not 3,4,5-tri(C1-C6)alkoxyphenyl or 3,4,5-trimethoxyphenyl, especially when R^5 and R^6 are each hydrogen..

In another embodiment of compounds including a trisubstituted phenyl, the trisubstituted phenyl has the formula:

5



wherein: R^{31} is methyl or (C1-C6) alkyl; R^{32} is hydrogen, methyl or (C1-C6) alkyl; and R^{33} is a halo group.

10

In a thirteenth embodiment of the compounds of structural formulae (I) and (Ia), R^2 and/or R^4 is an optionally substituted heteroaryl. Typical heteroaryl groups according to this thirteenth embodiment comprise from 5 to 15, and more typically from 5 to 11 ring atoms, and include one, two, three or four of the same or different heteratoms or heteroatomic groups selected from the group consisting of N, NH, O, S, S(O) and S(O)₂.

15


The optionally substituted heteroaryl may be attached to its respective C2 or C4 nitrogen atom or linker L^1 or L^2 through any available carbon atom or heteroatom, but is typically attached *via* a carbon atom. The optional substituents may be the same or different, and may be attached to any available carbon atom or heteroatom. In one embodiment of these compounds, R^5 is other than bromo, nitro, trifluoromethyl, cyano or $-C(O)NHR$, where R is


20

hydrogen or (C1-C6) alkyl. In another embodiment of these compounds, when R^2 and R^4 are each a substituted or unsubstituted pyrrole or indole, then the ring is attached to the remainder of the molecule *via* a ring carbon atom. In still another embodiment of

25

compounds including an optionally substituted heteroaryl group, the heteroaryl is unsubstituted or substituted with from one to four of the same or different R^8 groups, where R^8 is as previously defined for structural formula (I). Specific examples of such optionally substituted heteroaryls include, but are not limited to, the following heteroaryl groups:





wherein:

p is an integer from one to three;

each --- independently represents a single bond or a double bond;

5 R^{35} is hydrogen or R^8 , where R^8 is as previously defined for structural formula (I);

X is selected from the group consisting of CH, N and N-O;

each Y is independently selected from the group consisting of O, S and NH;

each Y^1 is independently selected from the group consisting of O, S, SO,

10 SO_2 , $SONR^{36}$, NH and NR^{37} ;

each Y^2 is independently selected from the group consisting of CH, CH_2 , O, S, N, NH and NR^{37} ;

R^{36} is hydrogen or alkyl;

R^{37} is selected from the group consisting of hydrogen and a progroup, preferably hydrogen or a progroup selected from the group consisting of aryl, arylalkyl, heteroaryl, R^a , R^b - CR^aR^b -O-C(O) R^8 , - CR^aR^b -O-PO(OR^8)₂, -CH₂-O-PO(OR^8)₂,
 5 -CH₂-PO(OR^8)₂, -C(O)- CR^aR^b -N(CH₃)₂, - CR^aR^b -O-C(O)- CR^aR^b -N(CH₃)₂, -C(O) R^8 ,
 -C(O)CF₃ and -C(O)-NR⁸-C(O) R^8 ;

A is selected from the group consisting of O, NH and NR³⁸;

R^{38} is selected from the group consisting of alkyl and aryl;

R^9 , R^{10} , R^{11} and R^{12} are each, independently of one another, selected from
 10 the group consisting of alkyl, alkoxy, halogen, haloalkoxy, aminoalkyl and hydroxyalkyl,
 or, alternatively, R^9 and R^{10} and/or R^{11} and R^{12} are taken together form a ketal;

each Z is selected from the group consisting of hydroxyl, alkoxy, aryloxy, ester, carbamate and sulfonyl;

Q is selected from the group consisting of -OH, OR^8 , -NR^c R^c ,
 15 -NHR³⁹-C(O) R^8 , -NHR³⁹-C(O) OR^8 , -NR³⁹-CHR⁴⁰- R^b , -NR³⁹-(CH₂)_m- R^b and
 -NR³⁹-C(O)-CHR⁴⁰-NR^c R^c ;

R^{39} and R^{40} are each, independently of one another, selected from the group consisting of hydrogen, alkyl, aryl, alkylaryl; arylalkyl and NHR⁸; and

R^a , R^b and R^c are as previously defined for structural formula (I). Preferred
 20 R^b substituents for Q are selected from -C(O) OR^8 , -O-C(O) R^8 , -O-P(O)(OR^8)₂ and
 -P(O)(OR^8)₂.

In one embodiment of the above-depicted heteroaryls, as well as other 5-15 membered heteroaryls according to this embodiment of the invention, each R^8 is independently selected from the group consisting of R^d , -NR^c R^c , -(CH₂)_m-NR^c R^c ,
 25 -C(O)NR^c R^c , -(CH₂)_m-C(O)NR^c R^c , -C(O) OR^d , -(CH₂)_m-C(O) OR^d and -(CH₂)_m- OR^d , where
 m , R^c and R^d are as previously defined for structural formula (I).

In a specific embodiment, R^d and/or R^c is selected from the group consisting of R^a and (C3-C8) cycloalkyl optionally substituted with one or more of the same or different hydroxyl, amino or carboxyl groups.

30 In another embodiment of the above-depicted heteroaryls, each R^{35} is hydrogen or (C1-C6) ethyl or methyl.


In still another embodiment of the above-depicted heteroaryls, the aromatic ring connectivity is either at the 5 or 6 position. It should be understood that either R^2 or R^4 can utilize the heteroaryl groups discussed throughout this specification.

In a fourteenth embodiment of the compounds of structural formulae (I) and (Ia), R^2 and R^4 are each, independently of one another, an optionally substituted phenyl, aryl or heteroaryl, with the provisos that: (1) when L^1 is a direct bond and R^6 and optionally R^5 is hydrogen, then R^2 is other than 3,4,5-trimethoxyphenyl or 3,4,5-tri(C1-C6) alkoxyphenyl; (2) when L^1 and L^2 are each a direct bond, R^6 is hydrogen and R^5 is halo, then R^2 and R^4 are not each simultaneously 3,4,5-trimethoxyphenyl or 3,4,5-tri(C1-C6) alkoxyphenyl; (3) when R^4 is 3-methoxyphenyl or 3-(C1-C6) alkoxyphenyl and R^2 is a 3,4,5-trisubstituted phenyl, the substituents positioned at the 3 and 4 positions are not both simultaneously methoxy or (C1-C6) alkoxy; (4) when R^2 is a substituted phenyl and R^6 is hydrogen, then R^5 is other than cyano or $-C(O)NHR$, where R is hydrogen or (C1-C6) alkyl; and/or (5) when R^2 and R^4 are each independently a substituted or unsubstituted pyrrole or indole, then the pyrrole or indole is attached to the remainder of the molecule *via* a ring carbon atom. Alternatively, R^2 is subject to the provisos described in connection with the first or second embodiment.

In this fourteenth embodiment of the invention, the R^2 and R^4 substituents may be the same or different. Specific optionally substituted phenyl, aryl and/or heteroaryls include those illustrated above in connection with the twelfth and thirteenth embodiments.

In a fifteenth embodiment of the compounds of structural formulae (I) and (Ia), including the above-described first through fourteenth embodiments thereof, R^6 is hydrogen and R^5 is an electronegative group. As will be recognized by skilled artisans, electronegative groups are atoms or groups of atoms that have a relatively great tendency to attract electrons to themselves. Specific examples of electronegative groups according to this fourteenth embodiment include, but are not limited to, $-CN$, $-NC$, $-NO_2$, halo, bromo, chloro, fluoro, (C1-C3) haloalkyl, (C1-C3) perhaloalkyl, (C1-C3) fluoroalkyl, (C1-C3) perfluoroalkyl, $-CF_3$, (C1-C3) haloalkoxy, (C1-C3) perhaloalkoxy, (C1-C3) fluoroalkoxy, (C1-C3) perfluoroalkoxy, $-OCF_3$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)CF_3$ and $-C(O)OCF_3$. In a specific embodiment, the electronegative group is a halogen-containing electronegative group, such as $-OCF_3$, $-CF_3$, bromo, chloro or fluoro. In another specific embodiment, R^5 is fluoro, subject to the proviso that the compound is not any compound according to the third embodiment.

In a sixteenth embodiment, the compounds of structural formulae (I) and (Ia) are compounds according to structural formula (Ib):




and salts, hydrates, solvates and N-oxides thereof, wherein R^{11} , R^{12} , R^{13} and R^{14} are each, independently of one another, selected from the group consisting of hydrogen,

- 5 hydroxy, (C1-C6) alkoxy and $-NR^cR^c$; and R^5 , R^6 and R^c are as previously defined for structural formula (I), with the proviso that when R^{13} , R^5 and R^6 are each hydrogen, then R^{11} and R^{12} are not simultaneously methoxy, (C1-C6) alkoxy or (C1-C6) haloalkoxy

In a seventeenth embodiment, the compounds of structural formulae (I) and (Ia) are compounds according to structural formula (Ic):

10



and salts, hydrates, solvates and N-oxides thereof, wherein:

R^4 is selected from the group consisting of 5-10 membered heteroaryl and 3-hydroxyphenyl;

- 15 R^5 is F or $-CF_3$; and

R^8 is $-O(CH_2)_m-R^b$, where m and R^b are as previously defined for structural formula (I). In a specific embodiment, R^8 is $-O-CH_2-C(O)NH-CH_3$ and/or R^4 is a heteroaryl according to the thirteenth embodiment.

- In an eighteenth embodiment, the compounds of structural formulae (I) and (Ia) include any compound selected from TABLE 1 that inhibits an Fc receptor signal transduction cascade, a Syk kinase activity, a Syk-kinase dependent receptor signal transduction cascade or cell degranulation as measured in an *in vitro* assay, optionally subject to the proviso that the compound is not a compound excluded by the above-described third embodiment and/or other embodiments. In a specific embodiment, such compounds have an IC_{50} of about 20 μM or less as measured in an *in vitro* degranulation assay, such as one of the degranulation assays described in the Examples section.
- 20
- 25

In a nineteenth embodiment, the compounds of structural formulae (I) and (Ia) include any compound selected from TABLE 1 that inhibits the $Fc\gamma R1$ or $Fc\epsilon R1$ receptor

cascade with an IC_{50} of about 20 μ M or less as measured in an *in vitro* assay, such as one of the *in vitro* assays provided in the Examples section, optionally subject to the proviso that the compound is not a compound excluded by the above-described third embodiment and/or other embodiments.

5 Also specifically described are combinations of the above first through nineteenth specific embodiments.


Those of skill in the art will appreciate that the 2,4-pyrimidinediamine compounds described herein may include functional groups that can be masked with progroups to create prodrugs. Such prodrugs are usually, but need not be, pharmacologically inactive until
10 converted into their active drug form. Indeed, many of the active 2,4-pyrimidinediamine compounds described in TABLE 1, *infra*, include promoieties that are hydrolyzable or otherwise cleavable under conditions of use. For example, ester groups commonly undergo acid-catalyzed hydrolysis to yield the parent carboxylic acid when exposed to the acidic conditions of the stomach, or base-catalyzed hydrolysis when exposed to the basic
15 conditions of the intestine or blood. Thus, when administered to a subject orally, 2,4-pyrimidinediamines that include ester moieties may be considered prodrugs of their corresponding carboxylic acid, regardless of whether the ester form is pharmacologically active. Referring to TABLE 1, numerous ester-containing 2,4-pyrimidinediamines of the invention are active in their ester, "prodrug" form.

20 In the prodrugs of the invention, any available functional moiety may be masked with a progroup to yield a prodrug. Functional groups within the 2,4-pyrimidinediamine compounds that may be masked with progroups for inclusion in a promoiety include, but are not limited to, amines (primary and secondary), hydroxyls, sulfanyls (thiols), carboxyls, etc. Myriad progroups suitable for masking such functional groups to yield promoieties that
25 are cleavable under the desired conditions of use are known in the art. All of these progroups, alone or in combinations, may be included in the prodrugs of the invention.

In one illustrative embodiment, the prodrugs of the invention are compounds according to structural formula (I) in which R^c and R^d may be, in addition to their previously-defined alternatives, a progroup.

30 Replacing the hydrogens attached to N2 and N4 in the 2,4-pyrimidinediamines of structural formula (I) with substituents adversely effects the activity of the compounds. However, as will be appreciated by skilled artisans, these nitrogens may be included in promoieties that, under conditions of use, cleave to yield 2,4-pyrimidinediamines according

to structural formula (I). Thus, in another embodiment, the prodrugs of the invention are compounds according to structural formula (II):



including salts, hydrates, solvates and N-oxides thereof, wherein:

R^2 , R^4 , R^5 , R^6 , L^1 and L^2 are as previously defined for structural formula (I); and R^{2b} and R^{4b} are each, independently of one another, a progroup. Specific examples of progroups according to this embodiment of the invention include, but are not limited to, (C1-C6) alkyl, $-C(O)CH_3$, $-C(O)NHR^{36}$ and $-S(O)_2R^{36}$, where R^{36} is (C1-C6) alkyl, (C5-C15) aryl and (C3-C8) cycloalkyl.

In the prodrugs of structural formula (II), the various substituents may be as described for the various first through twentieth embodiments previously described for the compounds of structural formulae (I) and (Ia), or combinations of such embodiments.

Those of skill in the art will appreciate that many of the compounds and prodrugs of the invention, as well as the various compound species specifically described and/or illustrated herein, may exhibit the phenomena of tautomerism, conformational isomerism, geometric isomerism and/or optical isomerism. For example, the compounds and prodrugs of the invention may include one or more chiral centers and/or double bonds and as a consequence may exist as stereoisomers, such as double-bond isomers (i.e., geometric isomers), enantiomers and diastereomers and mixtures thereof, such as racemic mixtures. As another example, the compounds and prodrugs of the invention may exist in several tautomeric forms, including the enol form, the keto form and mixtures thereof. As the various compound names, formulae and compound drawings within the specification and claims can represent only one of the possible tautomeric, conformational isomeric, optical isomeric or geometric isomeric forms, it should be understood that the invention encompasses any tautomeric, conformational isomeric, optical isomeric and/or geometric isomeric forms of the compounds or prodrugs having one or more of the utilities described herein, as well as mixtures of these various different isomeric forms. In cases of limited rotation around the 2,4-pyrimidinediamine core structure, atrop isomers are also possible and are also specifically included in the compounds of the invention.

Moreover, skilled artisans will appreciate that when lists of alternative substituents include members which, owing to valency requirements or other reasons, cannot be used to substitute a particular group, the list is intended to be read in context to include those members of the list that are suitable for substituting the particular group. For example, skilled artisans will appreciate that while all of the listed alternatives for R^b can be used to substitute an alkyl group, certain of the alternatives, such as =O, cannot be used to substitute a phenyl group. It is to be understood that only possible combinations of substituent-group pairs are intended.

The compounds and/or prodrugs of the invention may be identified by either their chemical structure or their chemical name. When the chemical structure and the chemical name conflict, the chemical structure is determinative of the identity of the specific compound.

Depending upon the nature of the various substituents, the 2,4-pyrimidinediamine compounds and prodrugs of the invention may be in the form of salts. Such salts include salts suitable for pharmaceutical uses ("pharmaceutically-acceptable salts"), salts suitable for veterinary uses, etc. Such salts may be derived from acids or bases, as is well-known in the art.

In one embodiment, the salt is a pharmaceutically acceptable salt. Generally, pharmaceutically acceptable salts are those salts that retain substantially one or more of the desired pharmacological activities of the parent compound and which are suitable for administration to humans. Pharmaceutically acceptable salts include acid addition salts formed with inorganic acids or organic acids. Inorganic acids suitable for forming pharmaceutically acceptable acid addition salts include, by way of example and not limitation, hydrohalide acids (*e.g.*, hydrochloric acid, hydrobromic acid, hydriodic, etc.), sulfuric acid, nitric acid, phosphoric acid, and the like. Organic acids suitable for forming pharmaceutically acceptable acid addition salts include, by way of example and not limitation, acetic acid, trifluoroacetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, oxalic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, palmitic acid, benzoic acid, 3-(4-hydroxybenzoyl) benzoic acid, cinnamic acid, mandelic acid, alkylsulfonic acids (*e.g.*, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, etc.), arylsulfonic acids (*e.g.*, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, etc.), 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid,

glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like.

Pharmaceutically acceptable salts also include salts formed when an acidic proton
5 present in the parent compound is either replaced by a metal ion (*e.g.*, an alkali metal ion, an alkaline earth metal ion or an aluminum ion) or coordinates with an organic base (*e.g.*, ethanolamine, diethanolamine, triethanolamine, N-methylglucamine, morpholine, piperidine, dimethylamine, diethylamine, etc.).

The 2,4-pyrimidinediamine compounds and of the invention, as well as the salts
10 thereof, may also be in the form of hydrates, solvates and N-oxides, as are well-known in the art.


6.3 Methods of Synthesis

The compounds and prodrugs of the invention may be synthesized *via* a variety of different synthetic routes using commercially available starting materials and/or starting
15 materials prepared by conventional synthetic methods. Suitable exemplary methods that may be routinely adapted to synthesize the 2,4-pyrimidinediamine compounds and prodrugs of the invention are found in U.S. Patent No. 5,958,935, the disclosure of which is incorporated herein by reference. Specific examples describing the synthesis of numerous compounds and prodrugs of the invention, as well as intermediates therefor, are provided in
20 the Examples section. All of the compounds of structural formulae (I), (Ia) and (II) may be prepared by routine adaptation of these methods.

A variety of exemplary synthetic routes that can be used to synthesize the 2,4-pyrimidinediamine compounds of the invention are described in Schemes (I)-(XI), below. In Schemes (I)-(XI), like-numbered compounds have similar structures. These methods
25 may be routinely adapted to synthesize the prodrugs according to structural formula (II).

In one exemplary embodiment, the compounds can be synthesized from substituted or unsubstituted uracils or thiouracils as illustrated in Scheme (I), below:

Scheme (I)



In Scheme (I), R^2 , R^4 , R^5 , R^6 , L^1 and L^2 are as previously defined for structural formula (I), X is a halogen (*e.g.*, F, Cl, Br or I) and Y and Y' are each, independently of one another, selected from the group consisting of O and S. Referring to Scheme (I), uracil or thioracil **2** is dihalogenated at the 2- and 4-positions using standard halogenating agent POX_3 (or other standard halogenating agent) under standard conditions to yield 2,4-bishalo pyrimidine **4**. Depending upon the R^5 substituent, in pyrimidine **4**, the halide at the C4 position is more reactive towards nucleophiles than the halide at the C2 position. This differential reactivity can be exploited to synthesize 2,4-pyrimidinediamines according structural formula (I) by first reacting 2,4-bishalopyrimidine **4** with one equivalent of amine **10**, yielding 4N-substituted-2-halo-4-pyrimidineamine **8**, followed by amine **6** to yield a 2,4-pyrimidinediamine according structural formula (I). 2N,4N-bis(substituted)-2,4-pyrimidinediamines **12** and **14** can be obtained by reacting 2,4-bishalopyrimidine **4** with excess **6** or **10**, respectively.

In most situations, the C4 halide is more reactive towards nucleophiles, as illustrated in the Scheme. However, as will be recognized by skilled artisans, the identity of the R⁵ substituent may alter this reactivity. For example, when R⁵ is trifluoromethyl, a 50:50 mixture of 4N-substituted-4-pyrimidineamine **8** and the corresponding 2N-substituted-2-pyrimidineamine is obtained. Regardless of the identity of the R⁵ substituent, the regioselectivity of the reaction can be controlled by adjusting the solvent and other synthetic conditions (such as temperature), as is well-known in the art.

The reactions depicted in Scheme (I) may proceed more quickly when the reaction mixtures are heated *via* microwave. When heating in this fashion, the following conditions may be used: heat to 175°C in ethanol for 5-20 min. in a Smith Reactor (Personal Chemistry) in a sealed tube (at 20 bar pressure).


The uracil or thiouracil **2** starting materials may be purchased from commercial sources or prepared using standard techniques of organic chemistry. Commercially available uracils and thiouracils that can be used as starting materials in Scheme (I) include, by way of example and not limitation, uracil (Aldrich #13,078-8; CAS Registry 66-22-8); 2-thio-uracil (Aldrich #11,558-4; CAS Registry 141-90-2); 2,4-dithiouracil (Aldrich #15,846-1; CAS Registry 2001-93-6); 5-acetouracil (Chem. Sources Int'l 2000; CAS Registry 6214-65-9); 5-azidouracil; 5-aminouracil (Aldrich #85,528-6; CAS Registry 932-52-5); 5-bromouracil (Aldrich #85,247-3; CAS Registry 51-20-7); 5-(trans-2-bromovinyl)-uracil (Aldrich #45,744-2; CAS Registry 69304-49-0); 5-(trans-2-chlorovinyl)-uracil (CAS Registry 81751-48-2); 5-(trans-2-carboxyvinyl)-uracil; uracil-5-carboxylic acid (2,4-dihydroxypyrimidine-5-carboxylic acid hydrate; Aldrich #27,770-3; CAS Registry 23945-44-0); 5-chlorouracil (Aldrich #22,458-8; CAS Registry 1820-81-1); 5-cyanouracil (Chem. Sources Int'l 2000; CAS Registry 4425-56-3); 5-ethyluracil (Aldrich #23,044-8; CAS Registry 4212-49-1); 5-ethenyluracil (CAS Registry 37107-81-6); 5-fluorouracil (Aldrich #85,847-1; CAS Registry 51-21-8); 5-iodouracil (Aldrich #85,785-8; CAS Registry 696-07-1); 5-methyluracil (thymine; Aldrich #13,199-7; CAS Registry 65-71-4); 5-nitouracil (Aldrich #85,276-7; CAS Registry 611-08-5); uracil-5-sulfamic acid (Chem. Sources Int'l 2000; CAS Registry 5435-16-5); 5-(trifluoromethyl)-uracil (Aldrich #22,327-1; CAS Registry 54-20-6); 5-(2,2,2-trifluoroethyl)-uracil (CAS Registry 155143-31-6); 5-(pentafluoroethyl)-uracil (CAS Registry 60007-38-3); 6-aminouracil (Aldrich #A5060-6; CAS Registry 873-83-6) uracil-6-carboxylic acid (orotic acid; Aldrich #0-840-2; CAS Registry 50887-69-9); 6-methyluracil (Aldrich #D11,520-7; CAS Registry 626-48-2); uracil-5-amino-6-carboxylic acid (5-aminoorotic acid; Aldrich #19,121-3; CAS Registry

#7164-43-4); 6-amino-5-nitrosouracil (6-amino-2,4-dihydroxy-5-nitrosopyrimidine; Aldrich #27,689-8; CAS Registry 5442-24-0); uracil-5-fluoro-6-carboxylic acid (5-fluoroorotic acid; Aldrich #42,513-3; CAS Registry 00000-00-0); and uracil-5-nitro-6-carboxylic acid (5-nitroorotic acid; Aldrich #18,528-0; CAS Registry 600779-49-9). Additional 5-, 6- and 5,6-substituted uracils and/or thiouracils are available from General Intermediates of Canada, Inc., Edmonton, CA (www.generalintermediates.com) and/or Interchim, Cedex, France (www.interchim.com), or may be prepared using standard techniques. Myriad textbook references teaching suitable synthetic methods are provided *infra*.

Amines **6** and **10** may be purchased from commercial sources or, alternatively, may be synthesized utilizing standard techniques. For example, suitable amines may be synthesized from nitro precursors using standard chemistry. Specific exemplary reactions are provided in the Examples section. See also Vogel, 1989, *Practical Organic Chemistry*, Addison Wesley Longman, Ltd. and John Wiley & Sons, Inc.

Skilled artisans will recognize that in some instances, amines **6** and **10** and/or substituents R⁵ and/or R⁶ on uracil or thiouracil **2** may include functional groups that require protection during synthesis. The exact identity of any protecting group(s) used will depend upon the identity of the functional group being protected, and will be apparent to those of skill in the art. Guidance for selecting appropriate protecting groups, as well as synthetic strategies for their attachment and removal, may be found, for example, in Greene & Wuts, *Protective Groups in Organic Synthesis*, 3d Edition, John Wiley & Sons, Inc., New York (1999) and the references cited therein (hereinafter "Greene & Wuts").

A specific embodiment of Scheme (I) utilizing 5-fluorouracil (Aldrich #32,937-1) as a starting material is illustrated in Scheme (Ia), below:




In Scheme (Ia), R², R⁴, L¹ and L² are as previously defined for Scheme (I).


- 5 According to Scheme (Ia), 5-fluorouracil 3 is halogenated with POCl₃ to yield 2,4-dichloro-5-fluoropyrimidine 5, which is then reacted with excess amine 6 or 10 to yield N²,N⁴-bis substituted 5-fluoro-2,4-pyrimidinediamine 11 or 13, respectively. Alternatively, asymmetric 2N,4N-disubstituted-5-fluoro-2,4-pyrimidinediamine 9 may be obtained by reacting 2,4-dichloro-5-fluoropyrimidine 5 with one equivalent of amine 10 (to yield 2-chloro-N⁴-substituted-5-fluoro-4-pyrimidineamine 7) followed by one or more equivalents
- 10 of amine 6.

In another exemplary embodiment, the 2,4-pyrimidinediamine compounds of the invention may be synthesized from substituted or unsubstituted cytosines as illustrated in Schemes (IIa) and (IIb), below:

Scheme (IIa)



Scheme (IIb)



5


In Schemes (IIa) and (IIb), R^2 , R^4 , R^5 , R^6 , L^1 , L^2 and X are as previously defined for Scheme (I) and PG represents a protecting group. Referring to Scheme (IIa), the C4 exocyclic amine of cytosine **20** is first protected with a suitable protecting group PG to yield N4-protected cytosine **22**. For specific guidance regarding protecting groups useful in this context, see Vorbrüggen and Ruh-Pohlenz, 2001, *Handbook of Nucleoside Synthesis*, John Wiley & Sons, NY, pp. 1-631 ("Vorbrüggen"). Protected cytosine **22** is halogenated at the C2 position using a standard halogenation reagent under standard conditions to yield 2-chloro-4N-protected-4-pyrimidineamine **24**. Reaction with amine **6** followed by deprotection of the C4 exocyclic amine and reaction with amine **10** yields a 2,4-pyrimidinediamine according to structural formula (I).

Alternatively, referring to Scheme (IIb), cytosine **20** may be reacted with amine **10** or protected amine **21** to yield N4-substituted cytosine **23** or **27**, respectively. These substituted cytosines may then be halogenated as previously described, deprotected (in the case of N4-substituted cytosine **27**) and reacted with amine **6** to yield a 2,4-pyrimidinediamine according to structural formula (I).

Commercially-available cytosines that may be used as starting materials in Schemes (IIa) and (IIb) include, but are not limited to, cytosine (Aldrich #14,201-8; CAS Registry 71-30-7); N⁴-acetylcytosine (Aldrich #37,791-0; CAS Registry 14631-20-0); 5-fluorocytosine (Aldrich #27,159-4; CAS Registry 2022-85-7); and 5-(trifluoromethyl)-cytosine. Other suitable cytosines useful as starting materials in Schemes (IIa) are available from General Intermediates of Canada, Inc., Edmonton, CA (www.generalintermediates.com) and/or Interchim, Cedex, France (www.interchim.com), or may be prepared using standard techniques. Myriad textbook references teaching suitable synthetic methods are provided *infra*.

In still another exemplary embodiment, the 2,4-pyrimidinediamine compounds of the invention may be synthesized from substituted or unsubstituted 2-amino-4-pyrimidinols as illustrated in Scheme (III), below:

Scheme (III)




In Scheme (III), R², R⁴, R⁵, R⁶, L¹, L² and X are as previously defined for Scheme (I) and Z is a leaving group as discussed in more detail in connection with Scheme IV, *infra*. Referring to Scheme (III), 2-amino-4-pyrimidinol 30 is reacted with amine 6 (or optionally protected amine 21) to yield N2-substituted-4-pyrimidinol 32, which is then halogenated as previously described to yield N2-substituted-4-halo-2-pyrimidineamine 34. Optional deprotection (for example if protected amine 21 was used in the first step) followed by reaction with amine 10 affords a 2,4-pyrimidinediamine according to structural formula (I). Alternatively, pyrimidinol 30 can be reacted with acylating agent 31.

Suitable commercially-available 2-amino-4-pyrimidinols 30 that can be used as starting materials in Scheme (III) include, but are not limited to, 2-amino-6-chloro-4-pyrimidinol hydrate (Aldrich #A4702-8; CAS Registry 00000-00-0) and 2-amino-6-hydroxy-4-pyrimidinol (Aldrich #A5040-1; CAS Registry 56-09-7). Other 2-amino-4-pyrimidinols 30 useful as starting materials in Scheme (III) are available from General Intermediates of Canada, Inc., Edmonton, CA (www.generalintermediates.com) and/or Interchim, Cedex, France (www.interchim.com), or may be prepared using standard techniques. Myriad textbook references teaching suitable synthetic methods are provided *infra*.

Alternatively, the 2,4-pyrimidinediamine compounds of the invention may be prepared from substituted or unsubstituted 4-amino-2-pyrimidinols as illustrated in Scheme (IV), below:

5

Scheme (IV)




In Scheme (IV), R², R⁴, R⁵, R⁶, L¹ and L² are as previously defined for Scheme (I) and Z represents a leaving group. Referring to Scheme (IV), the C2-hydroxyl of 4-amino-2-pyrimidinol 40 is more reactive towards nucleophiles than the C4-amino such that reaction with amine 6 yields N2-substituted-2,4-pyrimidinediamine 42. Subsequent reaction with compound 44, which includes a good leaving group Z, or amine 10 yields a 2,4-pyrimidinediamine according to structural formula (I). Compound 44 may include virtually any leaving group that can be displaced by the C4-amino of N2-substituted-2,4-pyrimidinediamine 42. Suitable leaving groups Z include, but are not limited to, halogens, methanesulfonyloxy (mesyloxy; "OMs"), trifluoromethanesulfonyloxy ("OTf") and *p*-toluenesulfonyloxy (tosyloxy; "OTs"), benzene sulfonyloxy ("besylate") and metanitro benzene sulfonyloxy ("nosylate"). Other suitable leaving groups will be apparent to those of skill in the art.

Substituted 4-amino-2-pyrimidinol starting materials may be obtained commercially or synthesized using standard techniques. Myriad textbook references teaching suitable synthetic methods are provided *infra*.

In still another exemplary embodiment, the 2,4-pyrimidinediamine compounds of the invention can be prepared from 2-chloro-4-aminopyrimidines or 2-amino-4-chloropyrimidines as illustrated in Scheme (V), below:

Scheme (V)




In Scheme (V), R^2 , R^4 , R^5 , R^6 , L^1 , L^2 and X are as defined for Scheme (I) and Z is as defined for Scheme (IV). Referring to Scheme (V), 2-amino-4-chloropyrimidine **50** is reacted with amino **10** to yield 4N-substituted-2-pyrimidineamine **52** which, following reaction with compound **31** or amine **6**, yields a 2,4-pyrimidinediamine according to structural formula (I). Alternatively, 2-chloro-4-amino-pyrimidine **54** may be reacted with compound **44** followed by amine **6** to yield a compound according to structural formula (I).

A variety of pyrimidines **50** and **54** suitable for use as starting materials in Scheme (V) are commercially available, including by way of example and not limitation, 2-amino-4,6-dichloropyrimidine (Aldrich #A4860-1; CAS Registry 56-05-3); 2-amino-4-chloro-6-methoxy-pyrimidine (Aldrich #51,864-6; CAS Registry 5734-64-5); 2-amino-4-chloro-6-methylpyrimidine (Aldrich #12,288-2; CAS Registry 5600-21-5); and 2-amino-4-chloro-6-methylthiopyrimidine (Aldrich #A4600-5; CAS Registry 1005-38-5). Additional pyrimidine starting materials are available from General Intermediates of Canada, Inc., Edmonton, CA (www.generalintermediates.com) and/or Interchim, Cedex, France (www.interchim.com), or may be prepared using standard techniques. Myriad textbook references teaching suitable synthetic methods are provided *infra*.

Alternatively, 4-chloro-2-pyrimidineamines **50** may be prepared as illustrated in Scheme (Va):

Scheme (Va)




In Scheme (Va), R^5 and R^6 are as previously defined for structural formula (I). In Scheme (Va), dicarbonyl **53** is reacted with guanidine to yield 2-pyrimidineamine **51**.

- 5 Reaction with peracids like m-chloroperbenzoic acid, trifluoroperacetic acid or urea hydrogen peroxide complex yields N-oxide **55**, which is then halogenated to give 4-chloro-2-pyrimidineamine **50**. The corresponding 4-halo-2-pyrimidineamines may be obtained by using suitable halogenation reagents.

10 In yet another exemplary embodiment, the 2,4-pyrimidinediamine compounds of the invention can be prepared from substituted or unsubstituted uridines as illustrated in Scheme (VI), below:


Scheme (VI)



In Scheme (VI), R^2 , R^4 , R^5 , R^6 , L^1 , L^2 and X are as previously defined for Scheme (I) and the superscript PG represents a protecting group, as discussed in connection with Scheme (IIb). According to Scheme (VI), uridine 60 has a C4 reactive center such that reaction with amine 10 or protected amine 21 yields N4-substituted cytidine 62 or 64, respectively. Acid-catalyzed deprotection of N4-substituted 62 or 64 (when "PG" represents an acid-labile protecting group) yields N4-substituted cytosine 28, which may be subsequently halogenated at the C2-position and reacted with amine 6 to yield a 2,4-pyrimidinediamine according to structural formula (I).

Cytidines may also be used as starting materials in an analogous manner, as illustrated in Scheme (VII), below:

Scheme (VII)



In Scheme (VII), R², R⁴, R⁵, R⁶, L¹, L² and X are as previously defined in Scheme (I) and the superscript PG represents a protecting group as discussed above. Referring to Scheme (VII), like uridine 60, cytidine 70 has a C4 reactive center such that reaction with amine 10 or protected amine 21 yields N4-substituted cytidine 62 or 64, respectively. These cytidines 62 and 64 are then treated as previously described for Scheme (VI) to yield a 2,4-pyrimidinediamine according to structural formula (I).

Although Schemes (VI) and (VII) are exemplified with ribosynucleosides, skilled artisans will appreciate that the corresponding 2'-deoxyribo and 2',3'-dideoxyribo nucleosides, as well as nucleosides including sugars or sugar analogs other than ribose, would also work.


Numerous uridines and cytidines useful as starting materials in Schemes (VI) and (VII) are known in the art, and include, by way of example and not limitation,

5-trifluoromethyl-2'-deoxycytidine (Chem. Sources #ABCR F07669; CAS Registry 66,384-66-5); 5-bromouridine (Chem. Sources Int'l 2000; CAS Registry 957-75-5); 5-iodo-2'-deoxyuridine (Aldrich #1-775-6; CAS Registry 54-42-2); 5-fluorouridine (Aldrich #32,937-1; CAS Registry 316-46-1); 5-iodouridine (Aldrich #85,259-7; CAS


Registry 1024-99-3); 5-(trifluoromethyl)uridine (Chem. Sources Int'l 2000; CAS Registry 70-00-8); 5-trifluoromethyl-2'-deoxyuridine (Chem. Sources Int'l 2000; CAS Registry 70-00-8). Additional uridines and cytidines that can be used as starting materials in Schemes (VI) and (VII) are available from General Intermediates of Canada, Inc., Edmonton, CA (www.generalintermediates.com) and/or Interchim, Cedex, France (www.interchim.com), or may be prepared using standard techniques. Myriad textbook references teaching suitable synthetic methods are provided *infra*.

The 2,4-pyrimidinediamine compounds of the invention can also be synthesized from substituted pyrimidines, such as chloro-substituted pyrimidines, as illustrated in Schemes (VIII) and (IX), below:

Scheme (VIII)



Scheme (IX)



In Schemes (VIII) and (IX), R², R⁴, L¹, L² and R^a are as previously defined for structural formula (I) and “Ar” represents an aryl group. Referring to Scheme (VIII), reaction of 2,4,6-trichloropyrimidine **80** (Aldrich #T5,620-0; CAS#3764-01-0) with amine **6** yields a mixture of three compounds: substituted pyrimidine mono-, di- and triamines **81**, **82** and **83**, which can be separated and isolated using HPLC or other conventional techniques. Mono- and diamines **81** and **82** may be further reacted with amines **6** and/or **10** to yield N2,N4,N6-trisubstituted-2,4,6-pyrimidinetriamines **84** and **85**, respectively.


N2,N4-bis-substituted-2,4-pyrimidinediamines can be prepared in a manner analogous to Scheme (VIII) by employing 2,4-dichloro-5-methylpyrimidine or 2,4-dichloropyrimidine as starting materials. In this instance, the mono-substituted pyrimidineamine corresponding to compound **81** is not obtained. Instead, the reaction proceeds to yield the N2,N4-bis-substituted-2,4-pyrimidinediamine directly.

Referring to Scheme (IX), 2,4,5,6-tetrachloropyrimidine **90** (Aldrich #24,671-9; CAS#1780-40-1) is reacted with excess amine **6** to yield a mixture of three compounds: **91**, **92**, and **93**, which can be separated and isolated using HPLC or other conventional techniques. As illustrated, N2,N4-bis-substituted-5,6-dichloro-2,4-pyrimidinediamine **92** may be further reacted at the C6 halide with, for example a nucleophilic agent **94** to yield compound **95**. Alternatively, compound **92** can be converted into N2,N4-bis-substituted-5-

chloro-6-aryl-2,4-pyrimidinediamine **97** via a Suzuki reaction. 2,4-Pyrimidinediamine **95** may be converted to 2,4-pyrimidinediamine **99** by reaction with Bn_3SnH .

As will be recognized by skilled artisans, 2,4-pyrimidinediamines according to the invention, synthesized via the exemplary methods described above or by other well-known means, may also be utilized as starting materials and/or intermediates to synthesize additional 2,4-pyrimidinediamine compounds of the invention. A specific example is illustrated in Scheme (X), below:


Scheme (X)



In Scheme (X), R^4 , R^5 , R^6 , L^2 and R^a are as previously defined for structural formula (I). Each $\text{R}^{a'}$ is independently an R^a , and may be the same or different from the illustrated R^a . Referring to Scheme (X), carboxylic acid or ester **100** may be converted to amide **104** by reaction with amine **102**. In amine **102**, $\text{R}^{a'}$ may be the same or different than R^a of acid or ester **100**. Similarly, carbonate ester **106** may be converted to carbamate **108**.

A second specific example is illustrated in Scheme (XI), below:

Scheme (XI)



In Scheme (XI), R^4 , R^5 , R^6 , L^2 and R^c are as previously defined for structural formula (I). Referring to Scheme (XI), amide 110 or 116 may be converted to amine 114 or 118, respectively, by borane reduction with borane methylsulfide complex 112. Other suitable reactions for synthesizing 2,4-pyrimidinediamine compounds from 2,4-pyrimidinediamine starting materials will be apparent to those of skill in the art.

Although many of the synthetic schemes discussed above do not illustrate the use of protecting groups, skilled artisans will recognize that in some instances substituents R^2 , R^4 , R^5 , R^6 , L^1 and/or L^2 may include functional groups requiring protection. The exact identity of the protecting group used will depend upon, among other things, the identity of the functional group being protected and the reaction conditions used in the particular synthetic scheme, and will be apparent to those of skill in the art. Guidance for selecting protecting groups and chemistries for their attachment and removal suitable for a particular application can be found, for example, in Greene & Wuts, *supra*.

Prodrugs according to structural formula (II) may be prepared by routine modification of the above-described methods. Alternatively, such prodrugs may be prepared by reacting a suitably protected 2,4-pyrimidinediamine of structural formula (I) with a suitable progroup. Conditions for carrying out such reactions and for deprotecting the product to yield a prodrug of formula (II) are well-known.

Myriad references teaching methods useful for synthesizing pyrimidines generally, as well as starting materials described in Schemes (I)-(IX), are known in the art. For

specific guidance, the reader is referred to Brown, D. J., "The Pyrimidines", in *The Chemistry of Heterocyclic Compounds, Volume 16* (Weissberger, A., Ed.), 1962, Interscience Publishers, (A Division of John Wiley & Sons), New York ("Brown I"); Brown, D. J., "The Pyrimidines", in *The Chemistry of Heterocyclic Compounds, Volume 16, Supplement I* (Weissberger, A. and Taylor, E. C., Ed.), 1970, Wiley-Interscience, (A Division of John Wiley & Sons), New York (Brown II"); Brown, D. J., "The Pyrimidines", in *The Chemistry of Heterocyclic Compounds, Volume 16, Supplement II* (Weissberger, A. and Taylor, E. C., Ed.), 1985, An Interscience Publication (John Wiley & Sons), New York ("Brown III"); Brown, D. J., "The Pyrimidines" in *The Chemistry of Heterocyclic Compounds, Volume 52* (Weissberger, A. and Taylor, E. C., Ed.), 1994, John Wiley & Sons, Inc., New York, pp. 1-1509 (Brown IV"); Kenner, G. W. and Todd, A., in *Heterocyclic Compounds, Volume 6*, (Elderfield, R. C., Ed.), 1957, John Wiley, New York, Chapter 7 (pyrimidines); Paquette, L. A., *Principles of Modern Heterocyclic Chemistry*, 1968, W. A. Benjamin, Inc., New York, pp. 1 – 401 (uracil synthesis pp. 313, 315; pyrimidine synthesis pp. 313-316; amino pyrimidine synthesis pp. 315); Joule, J. A., Mills, K. and Smith, G. F., *Heterocyclic Chemistry*, 3rd Edition, 1995, Chapman and Hall, London, UK, pp. 1 – 516; Vorbrüggen, H. and Ruh-Pohlenz, C., *Handbook of Nucleoside Synthesis*, John Wiley & Sons, New York, 2001, pp. 1-631 (protection of pyrimidines by acylation pp. 90-91; silylation of pyrimidines pp. 91-93); Joule, J. A., Mills, K. and Smith, G. F., *Heterocyclic Chemistry*, 4th Edition, 2000, Blackwell Science, Ltd, Oxford, UK, pp. 1 – 589; and *Comprehensive Organic Synthesis*, Volumes 1-9 (Trost, B. M. and Fleming, I., Ed.), 1991, Pergamon Press, Oxford, UK.

6.4 Inhibition of Fc Receptor Signal Cascades

Active 2,4-pyrimidinediamine compounds of the invention inhibit Fc receptor signalling cascades that lead to, among other things, degranulation of cells. As a specific example, the compounds inhibit the FcεRI and/or FcγRI signal cascades that lead to degranulation of immune cells such as neutrophil, eosinophil, mast and/or basophil cells. Both mast and basophil cells play a central role in allergen-induced disorders, including, for example, allergic rhinitis and asthma. Referring to FIG. 1, upon exposure allergens, which may be, among other things, pollen or parasites, allergen-specific IgE antibodies are synthesized by B-cells activated by IL-4 (or IL-13) and other messengers to switch to IgE class specific antibody synthesis. These allergen-specific IgEs bind to the high affinity FcεRI. Upon binding of antigen, the FcεR1-bound IgEs are cross-linked and the IgE

receptor signal transduction pathway is activated, which leads to degranulation of the cells and consequent release and/or synthesis of a host of chemical mediators, including histamine, proteases (e.g., tryptase and chymase), lipid mediators such as leukotrienes (e.g., LTC₄), platelet-activating factor (PAF) and prostaglandins (e.g., PGD₂) and a series of
5 cytokines, including TNF- α , IL-4, IL-13, IL-5, IL-6, IL-8, GM-CSF, VEGF and TGF- β . The release and/or synthesis of these mediators from mast and/or basophil cells accounts for the early and late stage responses induced by allergens, and is directly linked to downstream events that lead to a sustained inflammatory state.

The molecular events in the Fc ϵ RI signal transduction pathway that lead to release
10 of preformed mediators *via* degranulation and release and/or synthesis of other chemical mediators are well-known and are illustrated in FIG. 2. Referring to FIG. 2, the Fc ϵ RI is a heterotetrameric receptor composed of an IgE-binding alpha-subunit, a beta subunit, and two gamma subunits (gamma homodimer). Cross-linking of Fc ϵ RI-bound IgE by
15 multivalent binding agents (including, for example IgE-specific allergens or anti-IgE antibodies or fragments) induces the rapid association and activation of the Src-related kinase Lyn. Lyn phosphorylates immunoreceptor tyrosine-based activation motifs (ITAMS) on the intracellular beta and gamma subunits, which leads to the recruitment of additional Lyn to the beta subunit and Syk kinase to the gamma homodimer. These
20 receptor-associated kinases, which are activated by intra- and intermolecular phosphorylation, phosphorylate other components of the pathway, such as the Btk kinase, LAT, and phospholipase C-gamma (PLC-gamma). Activated PLC-gamma initiates pathways that lead to protein kinase C activation and Ca²⁺ mobilization, both of which are required for degranulation. Fc ϵ RI cross-linking also activates the three major classes of mitogen activated protein (MAP) kinases, *i.e.* ERK1/2, JNK1/2, and p38. Activation of
25 these pathways is important in the transcriptional regulation of proinflammatory mediators, such as TNF- α and IL-6, as well as the lipid mediator leukotriene CA (LTC₄).

Although not illustrated, the Fc γ RI signaling cascade is believed to share some common elements with the Fc ϵ RI signaling cascade. Importantly, like Fc ϵ RI, the Fc γ RI includes a gamma homodimer that is phosphorylated and recruits Syk, and like Fc ϵ RI,
30 activation of the Fc γ RI signaling cascade leads to, among other things, degranulation. Other Fc receptors that share the gamma homodimer, and which can be regulated by the active 2,4-pyrimidinediamine compounds include, but are not limited to, Fc α RI and Fc γ RIII.

The ability of the 2,4-pyrimidinediamine compounds of the invention to inhibit Fc receptor signaling cascades may be simply determined or confirmed in *in vitro* assays. Suitable assays for confirming inhibition of FcεRI-mediated degranulation are provided in the Examples section. In one typical assay, cells capable of undergoing FcεRI-mediated degranulation, such as mast or basophil cells, are first grown in the presence of IL-4, Stem Cell Factor (SCF), IL-6 and IgE to increase expression of the FcεRI, exposed to a 2,4-pyrimidinediamine test compound of the invention and stimulated with anti-IgE antibodies (or, alternatively, an IgE-specific allergen). Following incubation, the amount of a chemical mediator or other chemical agent released and/or synthesized as a consequence of activating the FcεRI signaling cascade may be quantified using standard techniques and compared to the amount of the mediator or agent released from control cells (*i.e.*, cells that are stimulated but that are not exposed to test compound). The concentration of test compound that yields a 50% reduction in the quantity of the mediator or agent measured as compared to control cells is the IC₅₀ of the test compound. The origin of the mast or basophil cells used in the assay will depend, in part, on the desired use for the compounds and will be apparent to those of skill in the art. For example, if the compounds will be used to treat or prevent a particular disease in humans, a convenient source of mast or basophil cells is a human or other animal which constitutes an accepted or known clinical model for the particular disease. Thus, depending upon the particular application, the mast or basophil cells may be derived from a wide variety of animal sources, ranging from, for example, lower mammals such as mice and rats, to dogs, sheep and other mammals commonly employed in clinical testing, to higher mammals such as monkeys, chimpanzees and apes, to humans. Specific examples of cells suitable for carrying out the *in vitro* assays include, but are not limited to, rodent or human basophil cells, rat basophil leukemia cell lines, primary mouse mast cells (such as bone marrow-derived mouse mast cells "BMMC") and primary human mast cells isolated from cord blood ("CHMC") or other tissues such as lung. Methods for isolating and culturing these cell types are well-known or are provided in the Examples section (*see, e.g.*, Demo *et al.*, 1999, Cytometry 36(4):340-348 and copending application Serial No. 10/053,355, filed November 8, 2001, the disclosures of which are incorporated herein by reference). Of course, other types of immune cells that degranulate upon activation of the FcεRI signaling cascade may also be used, including, for example, eosinophils.

As will be recognized by skilled artisans, the mediator or agent quantified is not critical. The only requirement is that it be a mediator or agent released and/or synthesized as a consequence of initiating or activating the Fc receptor signaling cascade. For example,

referring to FIG. 1, activation of the FcεRI signaling cascade in mast and/or basophil cells leads to numerous downstream events. For example, activation of the FcεRI signal cascade leads to the immediate release (i.e., within 1-3 min. following receptor activation) of a variety of preformed chemical mediators and agents *via* degranulation. Thus, in one embodiment, the mediator or agent quantified may be specific to granules (i.e., present in granules but not in the cell cytoplasm generally). Examples of granule-specific mediators or agents that can be quantified to determine and/or confirm the activity of a 2,4-pyrimidinediamine compound of the invention include, but are not limited to, granule-specific enzymes such as hexosaminidase and tryptase and granule-specific components such as histamine and serotonin. Assays for quantifying such factors are well-known, and in many instances are commercially available. For example, tryptase and/or hexosaminidase release may be quantified by incubating the cells with cleavable substrates that fluoresce upon cleavage and quantifying the amount of fluorescence produced using conventional techniques. Such cleavable fluorogenic substrates are commercially available. For example, the fluorogenic substrates Z-Gly-Pro-Arg-AMC (Z=benzyloxycarbonyl; AMC=7-amino-4-methylcoumarin; BIOMOL Research Laboratories, Inc., Plymouth Meeting, PA 19462, Catalog No. P-142) and Z-Ala-Lys-Arg-AMC (Enzyme Systems Products, a division of ICN Biomedicals, Inc., Livermore, CA 94550, Catalog No. AMC-246) can be used to quantify the amount of tryptase released. The fluorogenic substrate 4-methylumbelliferyl-N-acetyl-β-D-glucosaminide (Sigma, St. Louis, MO, Catalog #69585) can be used to quantify the amount of hexosaminidase released. Histamine release may be quantified using a commercially available enzyme-linked immunosorbent assay (ELISA) such as Immunotech histamine ELISA assay #IM2015 (Beckman-Coulter, Inc.). Specific methods of quantifying the release of tryptase, hexosaminidase and histamine are provided in the Examples section. Any of these assays may be used to determine or confirm the activity of the 2,4-pyrimidinediamine compounds of the invention.

Referring again to FIG. 1, degranulation is only one of several responses initiated by the FcεRI signaling cascade. In addition, activation of this signaling pathway leads to the *de novo* synthesis and release of cytokines and chemokines such as IL-4, IL-5, IL-6, TNF-α, IL-13 and MIP1-α), and release of lipid mediators such as leukotrienes (e.g., LTC4), platelet activating factor (PAF) and prostaglandins. Accordingly, the 2,4-pyrimidinediamine compounds of the invention may also be assessed for activity by

quantifying the amount of one or more of these mediators released and/or synthesized by activated cells.

Unlike the granule-specific components discussed above, these “late stage” mediators are not released immediately following activation of the FcεRI signaling cascade.

5 Accordingly, when quantifying these late stage mediators, care should be taken to insure that the activated cell culture is incubated for a time sufficient to result in the synthesis (if necessary) and release of the mediator being quantified. Generally, PAF and lipid mediators such as leukotriene C4 are released 3-30 min. following FcεRI activation. The cytokines and other late stage mediators are released approx. 4-8 hrs. following FcεRI
10 activation. Incubation times suitable for a specific mediator will be apparent to those of skill in the art. Specific guidance and assays are provided in the Examples section.

The amount of a particular late stage mediator released may be quantified using any standard technique. In one embodiment, the amount(s) may be quantified using ELISA assays. ELISA assay kits suitable for quantifying the amount of TNFα, IL-4, IL-5, IL-6
15 and/or IL-13 released are available from, for example, Biosource International, Inc., Camarillo, CA 93012 (see, e.g., Catalog Nos. KHC3011, KHC0042, KHC0052, KHC0061 and KHC0132). ELISA assay kits suitable for quantifying the amount of leukotriene C4 (LTC4) released from cells are available from Cayman Chemical Co., Ann Arbor, MI 48108 (see, e.g., Catalog No. 520211).

20 Typically, active 2,4-pyrimidinediamine compounds of the invention will exhibit IC₅₀s with respect to FcεRI-mediated degranulation and/or mediator release or synthesis of about 20 μM or lower, as measured in an *in vitro* assay, such as one of the *in vitro* assays described above or in the Examples section. Of course, skilled artisans will appreciate that compounds which exhibit lower IC₅₀s, for example on the order of 10 μM, 1 μM, 100 nM,
25 10 nM, 1 nM, or even lower, are particularly useful.

Skilled artisans will also appreciate that the various mediators discussed above may induce different adverse effects or exhibit different potencies with respect to the same adverse effect. For example, the lipid mediator LTC4 is a potent vasoconstrictor – it is approximately 1000-fold more potent at inducing vasoconstriction than histamine. As
30 another example, in addition to mediating atopic or Type I hypersensitivity reactions, cytokines can also cause tissue remodeling and cell proliferation. Thus, although compounds that inhibit release and/or synthesis of any one of the previously discussed chemical mediators are useful, skilled artisans will appreciate that compounds which inhibit

the release and/or synthesis of a plurality, or even all, of the previously described mediators find particular use, as such compounds are useful for ameliorating or avoiding altogether a plurality, or even all, of the adverse effects induced by the particular mediators. For example, compounds which inhibit the release of all three types of mediators—granule-specific, lipid and cytokine—are useful for treating or preventing immediate Type I hypersensitivity reactions as well as the chronic symptoms associated therewith.

Compounds of the invention capable of inhibiting the release of more than one type of mediator (*e.g.*, granule-specific or late stage) may be identified by determining the IC_{50} with respect to a mediator representative of each class using the various *in vitro* assays described above (or other equivalent *in vitro* assays). Compounds of the invention which are capable of inhibiting the release of more than one mediator type will typically exhibit an IC_{50} for each mediator type tested of less than about 20 μM . For example, a compound which exhibits an IC_{50} of 1 μM with respect to histamine release ($IC_{50}^{histamine}$) and an IC_{50} of 1 nM with respect to leukotriene LTC₄ synthesis and/or release ($IC_{50}^{LTC_4}$) inhibits both immediate (granule-specific) and late stage mediator release. As another specific example, a compound that exhibits an $IC_{50}^{tryptase}$ of 10 μM , an $IC_{50}^{LTC_4}$ of 1 μM and an IC_{50}^{IL-4} of 1 μM inhibits immediate (granule-specific), lipid and cytokine mediator release. Although the above specific examples utilize the IC_{50} s of one representative mediator of each class, skilled artisans will appreciate that the IC_{50} s of a plurality, or even all, mediators comprising one or more of the classes may be obtained. The quantity(ies) and identity(ies) of mediators for which IC_{50} data should be ascertained for a particular compound and application will be apparent to those of skill in the art.

Similar assays may be utilized to confirm inhibition of signal transduction cascades initiated by other Fc receptors, such as Fc α RI, Fc γ RI and/or Fc γ RIII signaling, with routine modification. For example, the ability of the compounds to inhibit Fc γ RI signal transduction may be confirmed in assays similar to those described above, with the exception that the Fc γ RI signaling cascade is activated, for example by incubating the cells with IgG and an IgG-specific allergen or antibody, instead of IgE and an IgE-specific allergen or antibody. Suitable cell types, activating agents and agents to quantify to confirm inhibition of other Fc receptors, such as Fc receptors that comprise a gamma homodimer, will be apparent to those of skill in the art.

One particularly useful class of compounds includes those 2,4-pyrimidinediamine compounds that inhibit the release of immediate granule-specific mediators and late stage

mediators with approximately equivalent IC_{50} s. By approximately equivalent is meant that the IC_{50} s for each mediator type are within about a 10-fold range of one another. Another particularly useful class of compounds includes those 2,4-pyrimidinediamine compounds that inhibit the release of immediate granule-specific mediators, lipid mediators and cytokine mediators with approximately equivalent IC_{50} s. In a specific embodiment, such compounds inhibit the release of the following mediators with approximately equivalent IC_{50} s: histamine, tryptase, hexosaminidase, IL-4, IL-5, IL-6, IL-13, $TNF\alpha$ and LTC₄. Such compounds are particularly useful for, among other things, ameliorating or avoiding altogether both the early and late stage responses associated with atopic or immediate Type I hypersensitivity reactions.

Ideally, the ability to inhibit the release of all desired types of mediators will reside in a single compound. However, mixtures of compounds can also be identified that achieve the same result. For example, a first compound which inhibits the release of granule specific mediators may be used in combination with a second compound which inhibits the release and/or synthesis of cytokine mediators.

In addition to the Fc ϵ RI or Fc γ RI degranulation pathways discussed above, degranulation of mast and/or basophil cells can be induced by other agents. For example, ionomycin, a calcium ionophore that bypasses the early Fc ϵ RI or Fc γ RI signal transduction machinery of the cell, directly induces a calcium flux that triggers degranulation. Referring again to FIG. 2, activated PLC γ initiates pathways that lead to, among other things, calcium ion mobilization and subsequent degranulation. As illustrated, this Ca²⁺ mobilization is triggered late in the Fc ϵ RI signal transduction pathway. As mentioned above, and as illustrated in FIG. 3, ionomycin directly induces Ca²⁺ mobilization and a Ca²⁺ flux that leads to degranulation. Other ionophores that induce degranulation in this manner include A23187. The ability of granulation-inducing ionophores such as ionomycin to bypass the early stages of the Fc ϵ RI and/or Fc γ RI signaling cascades may be used as a counter screen to identify active compounds of the invention that specifically exert their degranulation-inhibitory activity by blocking or inhibiting the early Fc ϵ RI or Fc γ RI signaling cascades, as discussed above. Compounds which specifically inhibit such early Fc ϵ RI or Fc γ RI-mediated degranulation inhibit not only degranulation and subsequent rapid release of histamine, tryptase and other granule contents, but also inhibit the pro-inflammatory activation pathways causing the release of $TNF\alpha$, IL-4, IL-13 and the lipid mediators such as LTC₄. Thus, compounds which specifically inhibit such early Fc ϵ RI and/or Fc γ RI-

mediated degranulation block or inhibit not only acute atopic or Type I hypersensitivity reactions, but also late responses involving multiple inflammatory mediators.

Compounds of the invention that specifically inhibit early FcεRI and/or FcγRI-mediated degranulation are those compounds that inhibit FcεRI and/or FcγRI-mediated degranulation (for example, have an IC₅₀ of less than about 20 μM with respect to the release of a granule-specific mediator or component as measured in an *in vitro* assay with cells stimulated with an IgE or IgG binding agent) but that do not appreciably inhibit ionophore-induced degranulation. In one embodiment, compounds are considered to not appreciably inhibit ionophore-induced degranulation if they exhibit an IC₅₀ of ionophore-induced degranulation of greater than about 20 μM, as measured in an *in vitro* assay. Of course, active compounds that exhibit even higher IC₅₀s of ionophore-induced degranulation, or that do not inhibit ionophore-induced degranulation at all, are particularly useful. In another embodiment, compounds are considered to not appreciably inhibit ionophore-induced degranulation if they exhibit a greater than 10-fold difference in their IC₅₀s of FcεRI and/or FcγRI-mediated degranulation and ionophore-induced degranulation, as measured in an *in vitro* assay. Assays suitable for determining the IC₅₀ of ionophore-induced degranulation include any of the previously-described degranulation assays, with the modification that the cells are stimulated or activated with a degranulation-inducing calcium ionophore such as ionomycin or A23187 (A.G. Scientific, San Diego, CA) instead of anti-IgE antibodies or an IgE-specific allergen. Specific assays for assessing the ability of a particular 2,4-pyrimidinediamine compound of the invention to inhibit ionophore-induced degranulation are provided in the Examples section.

As will be recognized by skilled artisans, compounds which exhibit a high degree of selectivity of FcεRI-mediated degranulation find particular use, as such compounds selectively target the FcεRI cascade and do not interfere with other degranulation mechanisms. Similarly, compounds which exhibit a high degree of selectivity of FcγRI-mediated degranulation find particular use, as such compounds selectively target the FcγRI cascade and do not interfere with other degranulation mechanisms. Compounds which exhibit a high degree of selectivity are generally 10-fold or more selective for FcεRI- or FcγRI-mediated degranulation over ionophore-induced degranulation, such as ionomycin-induced degranulation.

Biochemical and other data confirm that the 2,4-pyrimidinediamine compounds described herein are potent inhibitors of Syk kinase activity. For example, in experiments

with an isolated Syk kinase, of twenty four 2,4-pyrimidinediamine compounds tested, all but two inhibited the Syk kinase catalyzed phosphorylation of a peptide substrate with IC₅₀s in the submicromolar range. The remaining compounds inhibited phosphorylation in the micromolar range. In addition, of sixteen compounds tested in an *in vitro* assay with mast cells, all inhibited phosphorylation of Syk kinase substrates (*e.g.*, PLC-gamma1, LAT) and proteins downstream of Syk kinase (*e.g.*, JNK, p38, Erk1/2 and PKB, when tested), but not proteins upstream of Syk kinase in the cascade (*e.g.*, Lyn). Phosphorylation of Lyn substrates was not inhibited by the 2,4-pyrimidinediamine compounds tested. Moreover, for the following compounds, a high correlation was observed between their inhibition of Syk kinase activity in biochemical assays (IC₅₀s in the range of 3 to 1850 nM) and their inhibition of FcεR1-mediated degranulation in mast cells (IC₅₀s in the range of 30 to 1650 nM): R950373, R950368, R921302, R945371, R945370, R945369, R945365, R921304, R945144, R945140, R945071, R940358, R940353, R940352, R940351, R940350, R940347, R921303, R940338, R940323, R940290, R940277, R940276, R940275, R940269, R940255, R935393, R935372, R935366, R935310, R935309, R935307, R935304, R935302, R935293, R935237, R935198, R935196, R935194, R935193, R935191, R935190, R935138, R927050, R926968, R926956, R926931, R926891, R926839, R926834, R926816, R926813, R926791, R926782, R926780, R926757, R926753, R926745, R926715, R926508, R926505, R926502, R926501, R926500, R921218, R921147, R920410, R909268, R921219, R908712, R908702.

Accordingly, the activity of the 2,4-pyrimidinediamine compounds of the invention may also be confirmed in biochemical or cellular assays of Syk kinase activity. Referring again to FIG. 2, in the FcεRI signaling cascade in mast and/or basophil cells, Syk kinase phosphorylates LAT and PLC-gamma1, which leads to, among other things, degranulation. Any of these activities may be used to confirm the activity of the 2,4-pyrimidinediamine compounds of the invention. In one embodiment, the activity is confirmed by contacting an isolated Syk kinase, or an active fragment thereof with a 2,4-pyrimidinediamine compound in the presence of a Syk kinase substrate (*e.g.*, a synthetic peptide or a protein that is known to be phosphorylated by Syk in a signaling cascade) and assessing whether the Syk kinase phosphorylated the substrate. Alternatively, the assay may be carried out with cells that express a Syk kinase. The cells may express the Syk kinase endogenously or they may be engineered to express a recombinant Syk kinase. The cells may optionally also express the Syk kinase substrate. Cells suitable for performing such confirmation assays, as well as methods of engineering suitable cells will be apparent to those of skill in the art. Specific

examples of biochemical and cellular assays suitable for confirming the activity of the 2,4-pyrimidinediamine compounds are provided in the Examples section.

Generally, compounds that are Syk kinase inhibitors will exhibit an IC_{50} with respect to a Syk kinase activity, such as the ability of Syk kinase to phosphorylate a synthetic or endogenous substrate, in an *in vitro* or cellular assay in the range of about 20 μ M or less. Skilled artisans will appreciate that compounds that exhibit lower IC_{50} s, such as in the range of 10 μ M, 1 μ M, 100 nM, 10 nM, 1 nM, or even lower, are particularly useful.

6.5 Uses and Compositions

As previously discussed, the active compounds of the invention inhibit Fc receptor signaling cascades, especially those Fc receptors including a gamma homodimer, such as the Fc ϵ RI and/or Fc γ RI signaling cascades, that lead to, among other things, the release and/or synthesis of chemical mediators from cells, either *via* degranulation or other processes. As also discussed, the active compounds are also potent inhibitors of Syk kinase. As a consequence of these activities, the active compounds of the invention may be used in a variety of *in vitro*, *in vivo* and *ex vivo* contexts to regulate or inhibit Syk kinase, signaling cascades in which Syk kinase plays a role, Fc receptor signaling cascades, and the biological responses effected by such signaling cascades. For example, in one embodiment, the compounds may be used to inhibit Syk kinase, either *in vitro* or *in vivo*, in virtually any cell type expressing Syk kinase. They may also be used to regulate signal transduction cascades in which Syk kinase plays a role. Such Syk-dependent signal transduction cascades include, but are not limited to, the Fc ϵ RI, Fc γ RI, Fc γ RIII, BCR and integrin signal transduction cascades. The compounds may also be used *in vitro* or *in vivo* to regulate, and in particular inhibit, cellular or biological responses effected by such Syk-dependent signal transduction cascades. Such cellular or biological responses include, but are not limited to, respiratory burst, cellular adhesion, cellular degranulation, cell spreading, cell migration, cell aggregation, phagocytosis, cytokine synthesis and release, cell maturation and Ca^{2+} flux. Importantly, the compounds may be used to inhibit Syk kinase *in vivo* as a therapeutic approach towards the treatment or prevention of diseases mediated, either wholly or in part, by a Syk kinase activity. Non-limiting examples of Syk kinase mediated diseases that may be treated or prevented with the compounds are those discussed in more detail, below.

In another embodiment, the active compounds may be used to regulate or inhibit the Fc receptor signaling cascades and/or Fc ϵ RI- and/or Fc γ RI-mediated degranulation as a therapeutic approach towards the treatment or prevention of diseases characterized by,

caused by and/or associated with the release or synthesis of chemical mediators of such Fc receptor signaling cascades or degranulation. Such treatments may be administered to animals in veterinary contexts or to humans. Diseases that are characterized by, caused by or associated with such mediator release, synthesis or degranulation, and that can therefore be treated or prevented with the active compounds include, by way of example and not limitation, atopy or anaphylactic hypersensitivity or allergic reactions, allergies (e.g., allergic conjunctivitis, allergic rhinitis, atopic asthma, atopic dermatitis and food allergies), low grade scarring (e.g., of scleroderma, increased fibrosis, keloids, post-surgical scars, pulmonary fibrosis, vascular spasms, migraine, reperfusion injury and post myocardial infarction), diseases associated with tissue destruction (e.g., of COPD, cardiobronchitis and post myocardial infarction), diseases associated with tissue inflammation (e.g., irritable bowel syndrome, spastic colon and inflammatory bowel disease), inflammation and scarring.

When used to treat or prevent such diseases, the active compounds may be administered singly, as mixtures of one or more active compounds or in mixture or combination with other agents useful for treating such diseases and/or the symptoms associated with such diseases. The active compounds may also be administered in mixture or in combination with agents useful to treat other disorders or maladies, such as steroids, membrane stabilizers, 5LO inhibitors, leukotriene synthesis and receptor inhibitors, inhibitors of IgE isotype switching or IgE synthesis, IgG isotype switching or IgG synthesis, β -agonists, tryptase inhibitors, aspirin, COX inhibitors, methotrexate, anti-TNF drugs, retuxin, PD4 inhibitors, p38 inhibitors, PDE4 inhibitors, and antihistamines, to name a few. The active compounds may be administered *per se* in the form of prodrugs or as pharmaceutical compositions, comprising an active compound or prodrug.

Pharmaceutical compositions comprising the active compounds of the invention (or prodrugs thereof) may be manufactured by means of conventional mixing, dissolving, granulating, dragee-making levigating, emulsifying, encapsulating, entrapping or lyophilization processes. The compositions may be formulated in conventional manner using one or more physiologically acceptable carriers, diluents, excipients or auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically.

The active compound or prodrug may be formulated in the pharmaceutical compositions *per se*, or in the form of a hydrate, solvate, N-oxide or pharmaceutically

acceptable salt, as previously described. Typically, such salts are more soluble in aqueous solutions than the corresponding free acids and bases, but salts having lower solubility than the corresponding free acids and bases may also be formed.

5 Pharmaceutical compositions of the invention may take a form suitable for virtually any mode of administration, including, for example, topical, ocular, oral, buccal, systemic, nasal, injection, transdermal, rectal, vaginal, etc., or a form suitable for administration by inhalation or insufflation.

For topical administration, the active compound(s) or prodrug(s) may be formulated as solutions, gels, ointments, creams, suspensions, etc. as are well-known in the art.

10 Systemic formulations include those designed for administration by injection, *e.g.*, subcutaneous, intravenous, intramuscular, intrathecal or intraperitoneal injection, as well as those designed for transdermal, transmucosal oral or pulmonary administration.

Useful injectable preparations include sterile suspensions, solutions or emulsions of the active compound(s) in aqueous or oily vehicles. The compositions may also contain
15 formulating agents, such as suspending, stabilizing and/or dispersing agent. The formulations for injection may be presented in unit dosage form, *e.g.*, in ampules or in multidose containers, and may contain added preservatives.

Alternatively, the injectable formulation may be provided in powder form for reconstitution with a suitable vehicle, including but not limited to sterile pyrogen free water,
20 buffer, dextrose solution, etc., before use. To this end, the active compound(s) may be dried by any art-known technique, such as lyophilization, and reconstituted prior to use.

For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are known in the art.

For oral administration, the pharmaceutical compositions may take the form of, for
25 example, lozenges, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (*e.g.*, lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, talc or silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); or
30 wetting agents (*e.g.*, sodium lauryl sulfate). The tablets may be coated by methods well known in the art with, for example, sugars, films or enteric coatings. Compounds which are particularly suitable for oral administration include Compounds R940350, R935372, R935193, R927050 and R935391.

Liquid preparations for oral administration may take the form of, for example, elixirs, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as
5 suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol, cremophoreTM or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, preservatives, flavoring, coloring and sweetening agents as appropriate.

10 Preparations for oral administration may be suitably formulated to give controlled release of the active compound or prodrug, as is well known.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For rectal and vaginal routes of administration, the active compound(s) may be
15 formulated as solutions (for retention enemas) suppositories or ointments containing conventional suppository bases such as cocoa butter or other glycerides.

For nasal administration or administration by inhalation or insufflation, the active compound(s) or prodrug(s) can be conveniently delivered in the form of an aerosol spray from pressurized packs or a nebulizer with the use of a suitable propellant, e.g.,
20 dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, fluorocarbons, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges for use in an inhaler or insufflator (for example capsules and cartridges comprised of gelatin) may be formulated containing a powder mix of the compound and a suitable
25 powder base such as lactose or starch.

A specific example of an aqueous suspension formulation suitable for nasal administration using commercially-available nasal spray devices includes the following ingredients: active compound or prodrug (0.5-20 mg/ml); benzalkonium chloride (0.1-0.2 mg/mL); polysorbate 80 (TWEEN® 80; 0.5-5 mg/ml); carboxymethylcellulose sodium or
30 microcrystalline cellulose (1-15 mg/ml); phenylethanol (1-4 mg/ml); and dextrose (20-50 mg/ml). The pH of the final suspension can be adjusted to range from about pH5 to pH7, with a pH of about pH 5.5 being typical.

Another specific example of an aqueous suspension suitable for administration of the compounds *via* inhalation, and in particular for such administration of Compound

R921218, contains 1-20 mg/mL Compound or prodrug, 0.1-1% (v/v) Polysorbate 80 (TWEEN®80), 50 mM citrate and/or 0.9% sodium chloride.

For ocular administration, the active compound(s) or prodrug(s) may be formulated as a solution, emulsion, suspension, etc. suitable for administration to the eye. A variety of vehicles suitable for administering compounds to the eye are known in the art. Specific non-limiting examples are described in U.S. Patent No. 6,261,547; U.S. Patent No. 6,197,934; U.S. Patent No. 6,056,950; U.S. Patent No. 5,800,807; U.S. Patent No. 5,776,445; U.S. Patent No. 5,698,219; U.S. Patent No. 5,521,222; U.S. Patent No. 5,403,841; U.S. Patent No. 5,077,033; U.S. Patent No. 4,882,150; and U.S. Patent No. 4,738,851.

For prolonged delivery, the active compound(s) or prodrug(s) can be formulated as a depot preparation for administration by implantation or intramuscular injection. The active ingredient may be formulated with suitable polymeric or hydrophobic materials (e.g., as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, e.g., as a sparingly soluble salt. Alternatively, transdermal delivery systems manufactured as an adhesive disc or patch which slowly releases the active compound(s) for percutaneous absorption may be used. To this end, permeation enhancers may be used to facilitate transdermal penetration of the active compound(s). Suitable transdermal patches are described in for example, U.S. Patent No. 5,407,713.; U.S. Patent No. 5,352,456; U.S. Patent No. 5,332,213; U.S. Patent No. 5,336,168; U.S. Patent No. 5,290,561; U.S. Patent No. 5,254,346; U.S. Patent No. 5,164,189; U.S. Patent No. 5,163,899; U.S. Patent No. 5,088,977; U.S. Patent No. 5,087,240; U.S. Patent No. 5,008,110; and U.S. Patent No. 4,921,475.

Alternatively, other pharmaceutical delivery systems may be employed. Liposomes and emulsions are well-known examples of delivery vehicles that may be used to deliver active compound(s) or prodrug(s). Certain organic solvents such as dimethylsulfoxide (DMSO) may also be employed, although usually at the cost of greater toxicity.

The pharmaceutical compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active compound(s). The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

6.6 Effective Dosages

The active compound(s) or prodrug(s) of the invention, or compositions thereof, will generally be used in an amount effective to achieve the intended result, for example in an amount effective to treat or prevent the particular disease being treated. The compound(s) may be administered therapeutically to achieve therapeutic benefit or prophylactically to achieve prophylactic benefit. By therapeutic benefit is meant eradication or amelioration of the underlying disorder being treated and/or eradication or amelioration of one or more of the symptoms associated with the underlying disorder such that the patient reports an improvement in feeling or condition, notwithstanding that the patient may still be afflicted with the underlying disorder. For example, administration of a compound to a patient suffering from an allergy provides therapeutic benefit not only when the underlying allergic response is eradicated or ameliorated, but also when the patient reports a decrease in the severity or duration of the symptoms associated with the allergy following exposure to the allergen. As another example, therapeutic benefit in the context of asthma includes an improvement in respiration following the onset of an asthmatic attack, or a reduction in the frequency or severity of asthmatic episodes. Therapeutic benefit also includes halting or slowing the progression of the disease, regardless of whether improvement is realized.

For prophylactic administration, the compound may be administered to a patient at risk of developing one of the previously described diseases. For example, if it is unknown whether a patient is allergic to a particular drug, the compound may be administered prior to administration of the drug to avoid or ameliorate an allergic response to the drug. Alternatively, prophylactic administration may be applied to avoid the onset of symptoms in a patient diagnosed with the underlying disorder. For example, a compound may be administered to an allergy sufferer prior to expected exposure to the allergen. Compounds may also be administered prophylactically to healthy individuals who are repeatedly exposed to agents known to one of the above-described maladies to prevent the onset of the disorder. For example, a compound may be administered to a healthy individual who is repeatedly exposed to an allergen known to induce allergies, such as latex, in an effort to prevent the individual from developing an allergy. Alternatively, a compound may be administered to a patient suffering from asthma prior to partaking in activities which trigger asthma attacks to lessen the severity of, or avoid altogether, an asthmatic episode.

The amount of compound administered will depend upon a variety of factors, including, for example, the particular indication being treated, the mode of administration,

whether the desired benefit is prophylactic or therapeutic, the severity of the indication being treated and the age and weight of the patient, the bioavailability of the particular active compound, etc. Determination of an effective dosage is well within the capabilities of those skilled in the art.

5 Effective dosages may be estimated initially from *in vitro* assays. For example, an initial dosage for use in animals may be formulated to achieve a circulating blood or serum concentration of active compound that is at or above an IC_{50} of the particular compound as measured in as *in vitro* assay, such as the *in vitro* CHMC or BMMC and other *in vitro* assays described in the Examples section. Calculating dosages to achieve such circulating
10 blood or serum concentrations taking into account the bioavailability of the particular compound is well within the capabilities of skilled artisans. For guidance, the reader is referred to Fingl & Woodbury, "General Principles," *In: Goodman and Gilman's The Pharmaceutical Basis of Therapeutics*, Chapter 1, pp. 1-46, latest edition, Pagamonon Press, and the references cited therein.

15 Initial dosages can also be estimated from *in vivo* data, such as animal models. Animal models useful for testing the efficacy of compounds to treat or prevent the various diseases described above are well-known in the art. Suitable animal models of hypersensitivity or allergic reactions are described in Foster, 1995, *Allergy* 50(21Suppl):6-9, discussion 34-38 and Tumas *et al.*, 2001, *J. Allergy Clin. Immunol.* 107(6):1025-1033.
20 Suitable animal models of allergic rhinitis are described in Szelenyi *et al.*, 2000, *Arzneimittelforschung* 50(11):1037-42; Kawaguchi *et al.*, 1994, *Clin. Exp. Allergy* 24(3):238-244 and Sugimoto *et al.*, 2000, *Immunopharmacology* 48(1):1-7. Suitable animal models of allergic conjunctivitis are described in Carreras *et al.*, 1993, *Br. J. Ophthalmol.* 77(8):509-514; Saiga *et al.*, 1992, *Ophthalmic Res.* 24(1):45-50; and Kunert *et al.*, 2001,
25 Invest. Ophthalmol. Vis. Sci. 42(11):2483-2489. Suitable animal models of systemic mastocytosis are described in O'Keefe *et al.*, 1987, *J. Vet. Intern. Med.* 1(2):75-80 and Bean-Knudsen *et al.*, 1989, *Vet. Pathol.* 26(1):90-92. Suitable animal models of hyper IgE syndrome are described in Claman *et al.*, 1990, *Clin. Immunol. Immunopathol.* 56(1):46-53. Suitable animal models of B-cell lymphoma are described in Hough *et al.*, 1998, *Proc. Natl.*
30 Acad. Sci. USA 95:13853-13858 and Hakim *et al.*, 1996, *J. Immunol.* 157(12):5503-5511. Suitable animal models of atopic disorders such as atopic dermatitis, atopic eczema and atopic asthma are described in Chan *et al.*, 2001, *J. Invest. Dermatol.* 117(4):977-983 and Suto *et al.*, 1999, *Int. Arch. Allergy Immunol.* 120(Suppl 1):70-75. Ordinarily skilled

artisans can routinely adapt such information to determine dosages suitable for human administration. Additional suitable animal models are described in the Examples section.

Dosage amounts will typically be in the range of from about 0.0001 or 0.001 or 0.01 mg/kg/day to about 100 mg/kg/day, but may be higher or lower, depending upon, among other factors, the activity of the compound, its bioavailability, the mode of administration and various factors discussed above. Dosage amount and interval may be adjusted individually to provide plasma levels of the compound(s) which are sufficient to maintain therapeutic or prophylactic effect. For example, the compounds may be administered once per week, several times per week (e.g., every other day), once per day or multiple times per day, depending upon, among other things, the mode of administration, the specific indication being treated and the judgment of the prescribing physician. In cases of local administration or selective uptake, such as local topical administration, the effective local concentration of active compound(s) may not be related to plasma concentration. Skilled artisans will be able to optimize effective local dosages without undue experimentation.

Preferably, the compound(s) will provide therapeutic or prophylactic benefit without causing substantial toxicity. Toxicity of the compound(s) may be determined using standard pharmaceutical procedures. The dose ratio between toxic and therapeutic (or prophylactic) effect is the therapeutic index. Compounds(s) that exhibit high therapeutic indices are preferred.

The invention having been described, the following examples are offered by way of illustration and not limitation.

7. EXAMPLES

7.1 Synthesis of Starting Materials and Intermediates Useful for Synthesizing The 2,4-Pyrimidinediamine Compounds According to Schemes (I)–(V)

A variety of starting materials and N4-monosubstituted-2-pyrimidineamines and N2-monosubstituted-4-pyrimidinediamines [mono Substitution Nucleophilic Aromatic Reaction (SNAR) products] useful for synthesizing the 2,4-pyrimidinediamine compounds of the invention according to Schemes (I)–(V) were prepared as described below.

Conditions suitable for synthesizing the mono SNAR products are exemplified with 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (**R926087**).

7.1.1 2,4-Dichloro-5-fluoropyrimidine

To a dry reaction flask equipped with a stir bar and a reflux condenser was placed 5-fluorouracil (0.65g, 5mmol) followed by phosphorus oxychloride (POCl_3) (1.53g, 10mmol). The resultant mixture was heated at 110 °C for 8 hours under a nitrogen atmosphere. The reaction was cooled to room temperature, phosphorus pentachloride (PCl_5) (3.12g, 15mmol) was added and heated to 110 °C for a period of 12 hours. After cooling to room temperature, the mixture was poured into ice-water, saturated with sodium chloride and left for 1 hour at 0 °C to complete the decomposition of POCl_3 and PCl_5 . The solid of 2,4-dichloro-5-fluoropyrimidine was collected by rapid filtration, dried using blotting paper and stored at low temperature. ^1H NMR (CDCl_3): δ 8.47 (s, 1H); ^{13}C NMR (CDCl_3): δ 155.42, 151.87, 147.43 and 147.13; ^{19}F NMR (CDCl_3): $-\text{38149}$.

7.1.2 2,4-Dichloro-5-nitropyrimidine (Aldrich D6, 930-0)

A suspension of 5-nitrouracil (10g, 63 mmol) in POCl_3 (100 mL) was refluxed for 5h in the presence of N,N-dimethylaniline (10 mL), cooled to room temperature and poured on to crushed ice with vigorous stirring. The aqueous layer was extracted with ethyl acetate. The combined organic layers were dried over MgSO_4 and the solvent was evaporated under reduce pressure. The residue was purified by chromatography on silica gel (hexane/ethyl acetate; 1/1; v/v) to give the desired 2,4-dichloro-5-nitropyrimidine. LCMS: ret. time: 23.26 min.; purity: 95%; ^1H NMR (CDCl_3): δ 9.16 (1H, s).

7.1.3 2,4-Dichloro-5-cyanopyrimidine

In like manner to the preparation of 2,4-dichloro-5-nitropyrimidine, the reaction of 5-cyanouracil with POCl_3 and N,N-dimethylaniline gave 2,4-dichloro-5-cyanopyrimidine. LCMS: ret. time: 13.75 min.; purity: 95%.

7.1.4 2,4-Dichloro-5-trifluoromethylpyrimidine

In like manner to the preparation of 2,4-dichloro-5-nitropyrimidine, the reaction of 5-cyanouracil with POCl_3 and N,N-dimethylaniline gave 2,4-dichloro-5-cyanopyrimidine. ^1H NMR (CD_3OD): δ 9.07; LCMS: ret. time: 16.98 min. (fast method); purity: 70%.

7.1.5 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (R926087)

The reaction flask equipped with a magnetic stirring bar and a rubber septum (to prevent loss of 2,4-dichloro-5-fluoropyrimidine and N₂ inlet was charged with 3,4-ethylenedioxyaniline (34 g, 225 mmol), MeOH (100 mL), H₂O (300 mL) and 2,4-dichloro-5-fluoropyrimidine (25 g, 150 mmol). The reaction mixture was stirred at room temperature for 1h, diluted with H₂O (1.5 liter), acidified with 2N HCl (200 mL) and sonicated. The solid obtained was filtered, washed with H₂O and dried to obtain 33 g (78%) of the desired product, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (**R926087**).¹H NMR (CDCl₃): δ 8.02 (1H, d, J= 3Hz), 7.25 (d, 1H, J= 1.2 Hz), 6.98 (dd, 1H, J= 2.4 and 8.1 Hz), 6.85 (d, 1H, J= 5.7 Hz), 4.27 (m, 4H); ¹⁹F NMR (CDCl₃): - 44570; LCMS: ret. time: 26.70 min.; purity 100%; MS (m/e): 283 (MH⁺).

7.1.6 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-nitro-4-pyrimidineamine (R940094)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-nitropyrimidine and 3,4-ethylenedioxyaniline were reacted to prepare 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-nitro-4-pyrimidineamine. LCMS: ret. time: 28.79 min.; purity: 90%; MS (m/e): 308 (M⁺); ¹HNMR (CDCl₃): δ 10.07 (1H, s), 9.15 (1H, s), 7.02-6.88 (3H, m), 4.29 (4H, s).

7.1.7 2-Chloro-N4-(3-hydroxyphenyl)-5-nitro-4-pyrimidineamine (R940097)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-nitropyrimidine and 3-hydroxyaniline were reacted to prepare 2-chloro-N4-(3-hydroxyphenyl)-5-nitro-4-pyrimidineamine. LCMS: ret. time: 24.21 min.; purity: 93%; MS (m/e): 267 (MH⁺); ¹HNMR (CDCl₃): δ 10.20 (1H, s), 9.19 (1H, s), 7.32 (1H, t, J= 2.2 Hz), 7.28 (1H, d, J= 7.8 Hz), 7.11 (1H, dd, J= 7.8 and 1.8 Hz), 7.76 (1H, dd, J= 8.4 and 2.4 Hz), 5.20 (1H, s).

7.1.8 2-Chloro-N4-(3-hydroxyphenyl)-5-fluoro-4-pyrimidineamine (R926111)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-hydroxyaniline were reacted to prepare product 2-chloro-N4-(3-hydroxyphenyl)-5-fluoro-4-pyrimidineamine. ¹H

NMR (CD₃OD): δ 8.06 (bd, 1H), 7.26 (bd, 1H), 7.20-7.00 (m, 2H), 6.57 (d, 1H, J= 7.2 Hz);
¹⁹F NMR (CD₃OD): - 44374; LCMS: ret. time: 22.02; purity: 100%, MS (m/e): 240 (M⁺).

7.1.9 2-Chloro-N4-(3,4-dimethoxyphenyl)-5-fluoro-4-pyrimidineamine (R926073)

5 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3,4-dimethoxyaniline were reacted to prepare 2-chloro-N4-(3,4-dimethoxyphenyl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.02 (d, 1H, J= 2.7 Hz), 7.38 (d, 1H, J= 2.4 Hz), 7.05 (dd, 1H, J= 2.4 and 9.0 Hz), 6.89 (bs, 1H), 6.88 (d, 1H, J= 9 Hz), 3.91 (s, 3H), 3.89 (s, 3H); ¹⁹F NMR (CDCl₃):
10 - 44593; LCMS: ret. time: 24.95 min.; purity: 98%; MS (m/e): 285 (MH⁺).

7.1.10 2-Chloro-N4-(4-ethoxyphenyl)-5-fluoro-4-pyrimidineamine (R926066)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-ethoxyaniline were
15 reacted to prepare 2-chloro-N4-(4-ethoxyphenyl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.01 (d, 1H, J= 3Hz), 7.49 (bdd, 2H, J= 8.7 Hz), 6.92 (bdd, 2H, J= 9.6 Hz), 4.03 (q, 2H, J= 7.2 Hz), 1.42 (t, 3H, J= 7.2 Hz); ¹⁹F NMR (CDCl₃): - 44627; LCMS: ret. time: 29.50 min.; purity: 99%, MS (m/e): 268 (MH⁺).

7.1.11 2-Chloro-N4-(4-chlorophenyl)-5-fluoro-4-pyrimidineamine (R926207)

20 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-chloroaniline were reacted to prepare 2-chloro-N4-(4-chlorophenyl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.1 (bs, 1H), 8.60 (bdd, 2H), 8.36 (bdd, 2H), 6.90 (bs, 1H); ¹⁹F NMR (CDCl₃): -
25 44407; LCMS: ret. time: 31.63 min.; purity: 85%; MS (m/e): 258 (MH⁺).

7.1.12 2-Chloro-5-fluoro-N4-(3-hydroxy-4-methoxycarbonylmethyleneoxyphenyl)-4-pyrimidineamine (R926393)

30 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-hydroxy-4-methoxycarbonylmethyleneoxyaniline were reacted to prepare 2-chloro-5-fluoro-N4-(3-hydroxy-4-methoxycarbonylmethyleneoxyphenyl)-4-pyrimidineamine. ¹H NMR (CD₃OD):

δ 8.03 (d, 1H, J= 3.6 Hz), 7.35 (dd, 1H, J= 2.4 Hz), 7.12 (dd, 1H, J= 2.4 and 8.7 Hz), 6.82 (d, 1H, J= 8.1 Hz), 4.86 (s, 2H), 3.81 (s, 3H).

7.1.13 N4-(4-tert-Butoxycarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (R926573)

5 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and tert-butyl 4-aminophenoxyacetate were reacted to prepare product N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.02 (d, 1H, J= 2.7 Hz), 7.51 (d, 1H, J= 8.7 Hz), 6.93 (d, 1H, J= 8.7 Hz), 4.52 (s, 10 2H), 1.49 (s, 9H); LCMS: ret. time: 29.50 min.; purity: 97%; MS (m/e): 354 (MH⁺).

7.1.14 2-Chloro-5-fluoro-N4-(indol-5-yl)-4-pyrimidineamine (R926581)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 5-aminoindole were reacted
15 to prepare 2-chloro-5-fluoro-N4-(indol-5-yl)-4-pyrimidineamine. ¹H NMR (CDCl₃ + CD₃OD): δ 9.45 (bs, 1H), 8.00 (bs, 1H), 7.82 (bd, 1H), 7.75 (s, 1H), 7.38-7.10 (m, 3H), 6.40 (bs, 1H); LCMS: ret. time: 23.85 min.; purity: 100%; MS (m/e): 263 (MH⁺).

7.1.15 2-Chloro-5-fluoro-N4-(4-methoxymethyl-coumarin-7-yl)-4-pyrimidineamine (R926618)

20 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-methoxymethyl-7-aminocoumarin were reacted to prepare 2-chloro-5-fluoro-N4-(4-methoxymethyl-coumarin-7-yl)-4-pyrimidineamine. ¹H NMR (CD₃OD): δ 8.05 (d, 1H), 7.90 (s, 1H), 7.70 (dd, 1H, J= 2.4 and 8.7 Hz), 7.53 (d, 1H, J= 8.7 Hz), 6.42 (s, 1H), 4.61 (s, 2H), 3.49 (s, 3H); LCMS: ret.
25 time: 26.38 min.; purity: 87%; MS (m/e): 336 (MH⁺).

7.1.16 2-Chloro-N4-(2,5-dimethyl-4-hydroxyphenyl)-5-fluoro-4-pyrimidineamine (R926619)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 2,5-dimethyl-4-hydroxyaniline were reacted to prepare 2-chloro-N4-(2,5-dimethyl-4-hydroxyphenyl)-5-
30

fluoro-4-pyrimidineamine. LCMS: ret. time: 23.31 min.; purity: 96%; MS (m/e): 268 (MH⁺).

7.1.17 2-Chloro-N4-(5-chloropyrid-2-yl)-5-fluoro-4-pyrimidineamine (R926061)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 5-chloro-2-aminopyridine were reacted to prepare 2-chloro-N4-(5-chloropyrid-2-yl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.40 (d, 1H, J= 8.7 Hz), 8.28 (d, 1H, J= 1.8 Hz), 8.17 (d, 1H, J= 2.1 and 9 Hz); LCMS: ret. time: 28.58 min.; purity: 100%; MS (m/e): 259 (MH⁺).

7.1.18 2-Chloro-5-fluoro-N4-(5-methylpyrid-2-yl)-4-pyrimidineamine (R926062)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 5-methyl-2-aminopyridine were reacted to prepare 2-chloro-5-fluoro-N4-(5-methylpyrid-2-yl)-5-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 9.20 (s, 1H), 8.51 (s, 1H), 7.63 (d, 1H, J= 5.7 Hz), 7.45 (dd, 1H, J= 1.8 and 9.3 Hz), 2.43 (s, 3H); LCMS: ret. time: 21.29 min.; purity: 97%; MS (m/e): 239 (MH⁺).

7.1.19 N4-[6-(1,4-Benzoxazinyl)]-N2-chloro-5-fluoro-4-pyrimidineamine

In like manner to 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-1,4-benzoxazine were reacted (in methanol or methanol:water) to yield N4-[6-(1,4-benzoxazinyl)]-N2-chloro-5-fluoro-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.05 (m, 2H), 3.2 (m, 2H); LCMS: ret. time: 20.8 min.; purity: 95 %; MS (m/e): 295 (MH⁺).

7.1.20 N2-Chloro-N4-(2,3-dihydrobenzofuran-5-yl)-5-fluoro-4-pyrimidinediamine

In like manner to 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 5-amino-2,3-dihydrobenzofuran were reacted to yield N2-chloro-N4-(2,3-dihydrobenzofuran-5-yl)-5-fluoro-4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.09 (d, 1H), 8.00 (m, 1H), 7.42 (m, 2H), 7.05

(m, 1H), 4.53 (m, 2H), 4.25 (s, 2H), 3.15 (m, 2H); LCMS: ret. time: 20.35 min.; purity: 90 %; MS (m/e): 266 (MH⁺).

7.1.21 2-Chloro-N4-(2-carboxy-4-chlorophenyl)-5-fluoro-4-pyrimidineamine (R940050)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 2-carboxy-4-chloroaniline were reacted to prepare 2-chloro-N4-(2-carboxy-4-chlorophenyl)-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 20.83 min.; purity: 98%; ¹H NMR (CDCl₃): δ 8.64 (1H, d, J= 4.8 Hz), 8.24 (1H, d, J= 2.7 Hz), 7.76 (1H, dd, J= 8.7 and 2.7 Hz), 7.70 (1H, dd, J= 8.7 and J= 0.9 Hz).

7.1.22 N-(2-Chloro-5-fluoro-4-pyrimidinyl)-L-tyrosine Methyl Ester (R940108)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and L-tyrosine methyl ester were reacted to prepare N-(2-chloro-5-fluoro-4-pyrimidinyl)-L-tyrosine Methyl Ester. LCMS: ret. time: 23.32 min.; purity: 83%; MS (m/e): 325 (M⁺); ¹H NMR (CDCl₃): δ 7.90 (1H, d, J= 2.7 Hz), 6.95 (2H, d, J= 8.7 Hz), 6.75 (2H, d, J= 8.7 Hz), 5.95 (1H, s), 5.72 (1H, d, J= 7.5 Hz), 5.05 (1H, dt, J= 7.5 and 5.3 Hz), 3.77 (3H, s), 3.16 (2H, m).

7.1.23 2-Chloro-N4-[3-(5-cyano-2-methyl-4-thiomethyl-6-pyrimidinyl)phenyl]-5-fluoro-4-pyrimidineamine (R940141)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-(5-cyano-2-methyl-4-thiomethyl-6-pyrimidinyl)aniline were reacted to prepare 2-chloro-N4-[3-(5-cyano-2-methyl-4-thiomethyl-6-pyrimidinyl)phenyl]-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 18.23 min.; purity: 84%; MS (m/e): 386 (M⁺); ¹H NMR (CDCl₃): δ 8.19 (1H, t, J= 1.9 Hz), 8.11 (1H, d, J= 3.1 Hz), 7.98 (1H, dd, J= 8.1 and J= 2.4 Hz), 7.82 (1H, dd, J= 7.8 and 1.8 Hz), 7.57 (1H, t, J= 7.8 Hz), 7.11 (1H, s), 2.79 (3H, s), 2.69 (3H, s).

7.1.24 N4-[4-(N-Benzylpiperazino)phenyl]-2-chloro-5-fluoro-4-pyrimidineamine (R945154)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 4-(N-benzylpiperazino)aniline and 2,4-dichloro-5-

fluoropyrimidine gave N4-[4-(N-benzylpiperazino)phenyl]-2-chloro-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 2.81 (m, 4 H), 3.37 (m, 6 H), 6.85 (br, 1 H), 6.93 (d, J= 9.0 Hz, 2 H), 7.40 (m, 5 H), 7.50 (d, J= 9.3 Hz, 2 H), 8.02 (d, J= 2.7 Hz, 1 H); LCMS: ret. time: 20.56 min, purity: 97.75%; MS (m/e): 398.00 (MH⁺).

5 **7.1.25 2-Chloro-N4-(4-cyanomethyleneoxyphenyl)-5-fluoro-4-pyrimidineamine (R945069)**

In a manner analogous to the preparation of N2,N4-bis(4-cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N4-(4-aminocarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (178 mg, 0.6 mmol), trifluoroacetic anhydride (0.17 mL, 1.2 mmol) and pyridine (0.15 mL, 1.84 mmol) gave 2-chloro-N4-(4-cyanomethyleneoxyphenyl)-5-fluoro-4-pyrimidineamine (110 mg, 66%). ¹H NMR (acetone-*d*₆): δ 5.22 (s, 2 H), 7.24 (d, J= 9.3 Hz, 2 H), 7.62 (d, J= 9.0 Hz, 2 H), 8.94 (d, J= 1.8 Hz, 1 H); ¹⁹F NMR (acetone-*d*₆): -137.60; LCMS: ret. time: 26.19 min.; purity: 89.93%; MS (m/e): 279.06 (MH⁺).

15 **7.1.26 N4-(4-Acetoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (R940210)**

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-acetoxyaniline were reacted to prepare N4-(4-acetoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 25.97 min.; purity: 98%; MS (m/e): 281 (M⁺); ¹H NMR (CDCl₃): δ 8.07 (1H, d, J= 2.7 Hz), 7.64 (2H, d, J= 9 Hz), 7.12 (2H, d, J= 9 Hz), 7.00 (1H, s), 2.31 (3H, s).

7.1.27 2-Chloro-5-fluoro-N4-(4-hydroxyphenyl)-4-pyrimidineamine (R940211)

25 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-hydroxyaniline were reacted to prepare 2-chloro-5-fluoro-N4-(4-hydroxyphenyl)-4-pyrimidineamine. LCMS: ret. time: 20.10 min.; purity: 98%; MS (m/e): 240 (MH⁺); ¹H NMR (CDCl₃): δ 8.02 (1H, d, J= 2.7 Hz), 7.46 (2H, d, J= 8.7 Hz), 6.86 (2H, d, J= 9 Hz), 6.85 (1H, s), 4.94 (1H, s).

7.1.28 2-Chloro-N4-(2,3-dimethyl-4-hydroxyphenyl)-5-fluoro-4-pyrimidineamine (R940213)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 2,3-dimethyl-4-hydroxyaniline were reacted to prepare 2-chloro-N4-(2,3-dimethyl-4-hydroxyphenyl)-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 23.29 min.; purity: 93%; MS (m/e): 268 (MH⁺); ¹H NMR (CDCl₃): δ 8.00 (1H, d, J= 2.7 Hz), 7.16 (1H, d, J= 8.7 Hz), 6.68 (1H, d, J= 8.7 Hz), 6.61 (1H, s), 4.87 (1H, s), 2.21 (3H, s), 2.16 (3H, s).

7.1.29 2-Chloro-N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-4-pyrimidineamine (R940230)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-chloro-4-hydroxy-5-methylaniline were reacted to prepare 2-chloro-N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 26.26 min.; purity: 90%; ¹H NMR (DMSO-d₆): δ 9.94 (1H, s), 9.21 (1H, s), 8.37 (1H, d, 3.6 Hz), 7.68 (1H, s), 7.41 (1H, s), 2.30 (3H, s).

7.1.30 2-Chloro-5-fluoro-N4-[4-[3-(N-morpholino)propyl]oxyphenyl]-4-pyrimidineamine (R940247)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-[3-(N-morpholino)propyl]oxyaniline were reacted to prepare 2-chloro-5-fluoro-N4-[4-[3-(N-morpholino)propyl]oxyphenyl]-4-pyrimidineamine. LCMS: ret. time: 17.15 min.; purity: 99%; MS (m/e): 367 (MH⁺); ¹H NMR (CDCl₃): δ 8.02 (1H, d, J= 2.7 Hz), 7.49 (2H, d, J= 8.7 Hz), 6.92 (2H, d, J= 9 Hz), 6.85 (1H, s), 4.03 (2H, t, J= 6.3 Hz), 3.73 (4H, t, J= 4.6 Hz), 2.53 (2H, t, J= 6.7 Hz), 2.47 (4H, m), 1.98 (2H, m).

7.1.31 N4-[2-[4-(N-Benzylpiperazino)ethyl]]-2-chloro-5-fluoro-4-pyrimidineamine (R940259)

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 2-[4-(N-benzylpiperazino)ethyl]amine were reacted to prepare N4-[2-[4-(N-benzylpiperazino)ethyl]]-2-chloro-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 21.11 min.; purity: 96%; MS (m/e): 349 (M⁺); ¹H NMR (CDCl₃): δ 7.88 (1H, d, J= 2.6 Hz), 7.31-7.17 (4H, m), 7.14 (1H, d, J=

1.7 Hz), 7.10 (1H, s), 3.76 (2H, m), 3.24 (2H, m), 2.90 (2H, m), 2.59 (2H, m), 2.34 (2H, m), 1.76 (4H, m).

7.1.32 N4-(3-*tert*-Butylphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (R940268)

5 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-*tert*-butylaniline were reacted to prepare N4-(3-*tert*-butylphenyl)-2-chloro-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 33.96 min.; purity: 98 %; MS (m/e): 279 (M⁺); ¹H NMR (CDCl₃): δ 8.05 (1H, d, J= 3 Hz), 7.62 (1H, t, J= 1.3 Hz), 7.50 (1H, m), 7.34 (1H, t, J= 7.8 Hz), 7.22 (1H, m), 6.96
10 (1H, sl), 1.34 (9H, s).

7.1.33 2-Chloro-5-fluoro-N4-[3-(hydroxymethyl)phenyl]-4-pyrimidineamine (R925756)

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-aminobenzylalcohol were
15 reacted to yield 2-chloro-5-fluoro-N4-[3-(hydroxymethyl)phenyl]-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.45 (bs, 1H), 7.96 (d, 1H, J= 2.9 Hz), 7.65 (d, 1H, J= 8.2 Hz), 7.34 (s, 1H), 7.31 (t, 1H, J= 8.2 Hz), 7.07 (d, 1H, J= 8.2), 4.52 (s, 2H); ¹⁹F NMR (CDCl₃): -44394 (s, 1F); LCMS: ret. time: 20.29 min.; purity: 100 %; MS (m/e): 254 (MH⁺).

20 **7.1.34 2-Chloro-5-fluoro-N4-[4-(hydroxymethyl)phenyl]-4-pyrimidineamine (R925759)**

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-aminobenzylalcohol were
reacted to yield 2-chloro-5-fluoro-N4-[4-(hydroxymethyl)phenyl]-4-pyrimidineamine. ¹H
25 NMR (CDCl₃): δ 8.08 (d, 1H, J= 2.7 Hz), 7.62 (d, 2H, J= 9.0 Hz), 7.40 (d, 2H, J= 8.1 Hz), 6.99 (bs, 1H), 4.70 (s, 2H); ¹⁹F NMR (CDCl₃): -44570 (s, 1F); LCMS: ret. time: 19.57 min.; purity: 99%; MS (m/e): 254 (MH⁺).

7.1.35 2-Chloro-N4-(3,3-dihydroisobenzofuranylnl-1-one-6-yl)-5-fluoro-4-pyrimidineamine R940279

30 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-3,3-

dihydroisobenzofuran-1-one were reacted to give 2-chloro-N4-(3,3-dihydroisobenzofuranylnl-1-one-6-yl)-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 21.15 min.; purity: 94.7 %; MS (m/e): 280 (MH⁺).

7.1.36 2-Chloro-5-fluoro-N4-((2R)-hydroxy-(1S)-methyl-2-phenylethyl)-4-pyrimidineamine (R925762)

5

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and (1R,2S)-(-)-norephedrine were reacted to yield 2-chloro-5-fluoro-N4-(2R-hydroxy-1S-methyl-2-phenylethyl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.85 (d, 1H, J= 3.0 Hz), 7.38 (m, 5H), 5.56 (d, 1H, J= 7.5 Hz), 5.00 (d, 1H, J= 3.0 Hz), 4.54 (m, 1H), 2.87 (bs, 1H), 1.10 (d, 1H, J= 6.9 Hz); ¹⁹F NMR (CDCl₃): - 44408.

10

7.1.37 N-(2-Chloro-6-ethoxycarbonyl-5-nitro-4-pyrimidinyl)glycine Ethyl Ester (R925850)

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-6-ethoxycarbonyl-5-nitropyrimidine and glycine ethyl ester hydrochloride salt were reacted to yield N-(2-chloro-6-ethoxycarbonyl-5-nitro-4-pyrimidinyl)glycine Ethyl Ester. ¹H NMR (CDCl₃): δ 8.87 (bs, 1H), 4.48 (q, 2H, J= 7.2 Hz), 4.39 (d, 2H, J= 5.1 Hz), 1.40 (t, 3H, J= 6.9 Hz), 1.33 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 28.27 min.; purity: 97%; MS (m/e): 332 (M⁺).

15

7.1.38 2-Chloro-5-fluoro-N4-(2-hydroxy-2-phenylethyl)-4-pyrimidineamine (R925763)

20

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 2-amino-1-phenylethanol were reacted to yield 2-chloro-5-fluoro-N4-(2-hydroxy-2-phenylethyl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.88 (d, 1H, J= 3.0 Hz), 7.41-7.32 (m, 5H), 5.71 (bs, 1H), 4.97 (d, 1H, J= 8.1 Hz), 3.98 (m, 1H), 3.56 (m, 1H), 2.57 (s, 1H); ¹⁹F NMR (CDCl₃): - 45149; LCMS: ret. time: 22.27 min.; purity: 98%; MS (m/e): 263 (M⁺).

25

7.1.39 2-Chloro-5-fluoro-N4-(furfuryl)-4-pyrimidineamine (R925764)

30

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and

furfurylamine were reacted to yield 2-chloro-5-fluoro-N4-(furfuryl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.91 (d, 1H, J= 1.8 Hz), 7.39 (d, 1H, J= 1.2 Hz), 6.35 (m, 2H), 5.50 (bs, 1H), 4.69 (d, 2H, J= 5.1 Hz); ¹⁹F NMR (CDCl₃): - 45163; LCMS: ret. time: 24.52 min.; purity: 97%; MS (m/e): 228 (M⁺).

5 **7.1.40 R935010: (±)-2-Chloro-5-fluoro-N4-[1-(4-hydroxyphenyl)ethyl]-4-pyrimidineamine**

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with 1-(4-hydroxyphenyl)ethylamine to provide (±)-2-chloro-5-fluoro-N4-[1-(4-hydroxyphenyl)ethyl]-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.88 (d, 1H, J= 2.3 Hz), 7.50-7.47 (dd, 2H, J= 1.7 and 8.7 Hz), 7.26-7.23 (dd, J= 8.7 and 1.7 Hz), 5.35-5.28 (m, 2H), 1.59 (d, 3H, J= 7.0 Hz).

15 **7.1.41 R935011: (±)-N4-[1-(4-Bromophenyl)ethyl]-2-chloro-5-fluoro-4-pyrimidineamine**

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with 1-(4-bromophenyl)ethylamine to provide (±)-N4-[1-(4-bromophenyl)ethyl]-2-chloro-5-fluoro-4-pyrimidineamine: ¹H NMR (CDCl₃): δ 7.88 (d, 1H, J= 2.3 Hz), 7.49 (d, 2H, J= 8.7 Hz), 7.25 (d, 2H, J= 8.7 Hz), 4.45-5.26 (m, 2H), 1.59 (d, 3H, J=7.0 Hz).

20 **7.1.42 R935007: 2-chloro-5-fluoro-N4-[1-[(1S)-phenyl]ethyl]-4-pyrimidineamine**

In like manner to the preparation of of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 1-(1S)-phenyl ethylamine were reacted to produce 2-chloro-5-fluoro-N4-[1-[(1S)-phenyl]ethyl]-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.86 (d, 1H, J = 2.9 Hz), 7.37 (d, 4H, J = 4.7 Hz), 7.34-7.30 (m, 1H), 5.40-5.32 (m, 2H), 1.62 (d, 3H, J = 6.4 Hz); LCMS: ret. time: 29.5 min.; purity: 98%; MS (m/e): 252 (MH⁺).

25 **7.1.43 R935008: 2-Chloro-5-fluoro-N4-[1-[(1R)-phenyl]ethyl]-4-pyrimidineamine**

30 In like manner to the preparation of of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 1-(1R)-phenyl ethylamine

were reacted to produce 2-chloro-5-fluoro-N4-[1-[(1*R*)-phenyl]ethyl]-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.87 (d, 1H, J = 2.9 Hz), 7.37 (d, 4H, J = 4.1 Hz), 7.34-7.30 (m, 1H), 5.38-5.31 (m, 2H), 1.62 (d, 3H, J = 6.4 Hz).

7.1.44 R935012: 2-Chloro-N4-[[di(3,5-di(trifluoromethyl)phenyl)]methyl]-5-fluoro-4-pyrimidineamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with di[3,5-di(trifluoromethyl)phenyl]methylamine to provide 2-chloro-N4-[[di(3,5-di(trifluoromethyl)phenyl)]methyl]-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.06 (d, 1H, J = 2.3 Hz), 7.92 (s, 2H), 7.74 (s, 4H), 6.75 (d, 1H, J = 7.6 Hz), 5.80 (d, 1H, J = 7.0 Hz).

7.1.45 R935014: 2-Chloro-5-fluoro-N4-[1-[(1*R*)-4-methoxyphenyl]ethyl]-4-pyrimidineamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with (*R*)-(+)-1-(4-methoxyphenyl)ethylamine to provide 2-chloro-5-fluoro-N4-[1-[(1*R*)-4-methoxyphenyl]ethyl]-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.84 (d, 1H, J = 2.3 Hz), 7.30 (d, 2H, J = 8.8 Hz), 6.89 (d, 2H, J = 8.8 Hz), 5.39-5.26 (m, 2H), 3.80 (s, 3H), 1.59 (d, 3H, J = 6.4 Hz).

7.1.46 R935015: 2-Chloro-5-fluoro-N4-[1-[(1*S*)-4-methoxyphenyl]ethyl]-4-pyrimidineamine

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with (*S*)-(-)-1-(4-methoxyphenyl)ethylamine to provide 2-chloro-5-fluoro-N4-[1-[(1*S*)-4-methoxyphenyl]ethyl]-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.85 (d, 1H, J = 2.3 Hz), 7.31 (d, 2H, J = 8.8 Hz), 6.89 (d, 2H, J = 8.8 Hz), 5.38-5.29 (m, 2H), 3.80 (s, 3H), 1.59 (d, 3H, J = 7.7 Hz).

7.1.47 R935013: 2-Chloro-N-(fluoren-9-yl)-5-fluoro-4-pyrimidineamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 9-aminofluorene hydrochloride and 2,4-dichloro-5-fluoropyrimidine

with added diisopropylethylamine were reacted to produce 2-chloro-N-(fluoren-9-yl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.97 (d, 1H, J= 2.3 Hz), 7.73 (d, 2H, J= 7.6 Hz), 7.59(d, 2H, J= 7.6 Hz), 7.44 (t, 2H, J= 7.6 Hz), 7.32 (app t, 2H, J= 7.6 Hz), 6.50 (d, 1H, J= 8.8 Hz), 5.45 (d, 1H, J= 8.4 Hz).

5 **7.1.48 R935210: 2-Chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-4-pyrimidineamine**

In like manner to the preparation of 2-chloro-N-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, experiment, 2,4-dichloro-5-fluoropyrimidine was reacted with 4-(methoxycarbonylmethyleneoxy)aniline to produce 2-chloro-5-fluoro-N-[4-(methoxycarbonylmethyleneoxy)phenyl]-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 10.17 (s, 1H), 8.33 (d, 1H, J= 3.5 Hz), 8.05 (s, 1H), 7.91 (s, 1H), 7.74 (d, 1H, J= 8.2 Hz), 7.40 (d, 1H, J= 7.6 Hz), 5.31 (s, 2H), 3.66 (s, 3H).

7.1.49 R935200: 2-Chloro-5-fluoro-N-(1-methyl-indazoline-5-yl)-4-pyrimidineamine:

15 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 5-amino-1-methyl-indazoline were reacted to provide 2-chloro-5-fluoro-N-(1-methyl-indazoline-5-yl)-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 10.01 (s, 1H), 8.27 (d, 1H, J= 3.5 Hz), 8.04 (d, 1H, J= 1.7 Hz), 7.98 (d, 1H, J= 1.7 Hz), 7.64 (d, 1H, J= 8.8 Hz), 7.56 (dd, 1H, J= 1.7 and 8.8 Hz), 4.02 (s, 3H). LCMS: ret. time: 21.72 min.; purity: 99%; MS (m/e): 278 (MH⁺).

7.1.50 R935017: N-(5-Bromo-2-chloropyrimidinyl)-4-fluorophenylethylamine

25 In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 4-fluoro-α-methylbenzylamine and 5-bromo-2,4-dichloropyrimidine were reacted to produce N-(5-bromo-2-chloropyrimidinyl)-4-fluorophenylethylamine. ¹H NMR (CDCl₃): δ 8.12 (s, 1H), 7.35-7.25 (dd, 2H, J= 3.5 and 8.7 Hz), 7.05 (t, 1H, J= 8.7 Hz), 5.63 (d, 1H, J= 6.4 Hz), 5.36 (dq, 1H, 1H, J= 6.4 and 7.0 Hz), 1.60 (d, 3H, J= 7.0 Hz); LCMS: ret. time: 30.73 min.; purity: 94%; MS (m/e): 331 (MH⁺).

7.1.51 R935009: ($\pm$)-N-(2-Chloro-5-fluoropyrimidinyl)-1-(4-fluorophenyl)ethylamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 4-fluoro- α -methylbenzylamine and 2,4-dichloro-5-fluoropyrimidine were reacted to produce ($\pm$)-N-(2-chloro-5-fluoropyrimidinyl)-1-(4-fluorophenyl)ethylamine. ^1H NMR (CDCl_3): δ 7.87 (d, 1H, J = 2.3 Hz), 7.37-7.33 (dd, 2H, J = 5.4 and 8.4 Hz), 7.04 (t, 2H, J = 8.4 Hz), 5.35-5.31 (m, 2H), 1.60 (d, 3H, J = 6.4 Hz); LCMS: ret. time: 32.90 min.; purity: 98%; MS (m/e): 270 (MH^+).

7.1.52 R935022: 5-Bromo-2-chloro-N4-[4-(N-methyl-2-methoxycarbonyl)pyrrolyl]-4-pyrimidineamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 5-bromo-2,4-dichloropyrimidine and N-methyl-2-carbomethoxy-4-aminopyrrole hydrochloride with added diisopropylethylamine were reacted to produce the desired product 5-bromo-2-chloro-N-(N-methyl-2-carbomethoxypyrrol-4-yl)-4-pyrimidineamine. ^1H NMR (CDCl_3): δ 8.21 (s, 1H), 7.43 (d, 1H, J = 1.8 Hz), 7.13 (br s, 1H), 6.84 (d, 1H, J = 1.8 Hz), 3.95 (s, 3H), 3.82 (s, 3H); LCMS: ret. time: 26.96 min.; purity: 91%; MS (m/e): 346 (MH^+).

7.1.53 R935234: 2-Chloro-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 5-(4-aminophenoxymethyl)-3-phenyl-1,2,4-oxadiazole were reacted to produce 2-chloro-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine. ^1H NMR ($\text{DMSO}-d_6$): δ 9.92 (s, 1H), 8.26 (d, 1H, J = 3.5 Hz), 8.02-7.99 (m, 2H), 7.60-7.56 (m, 5H), 7.11 (d, 2H, J = 8.8 Hz), 5.58 (s, 2H); LCMS: ret. time: 32.09 min.; purity: 96%; MS (m/e): 398 (MH^+).

7.1.54 R935235: 2-Chloro-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 5-(4-aminophenoxymethyl)-3-methyl-1,2,4-oxadiazole were reacted to produce 2-chloro-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine. ^1H

NMR (DMSO- d_6): δ 9.91 (s, 1H), 8.26 (d, 1H, J = 3.5 Hz), 7.56 (d, 2H, J = 8.8 Hz), 7.05 (d, 2H, J = 8.8 Hz), 5.46 (s, 2H), 2.34 (s, 3H); LCMS: ret. time: 25.05 min.; purity: 98%; MS (m/e): 336 (MH^+).

7.1.55 R935236: 2-Chloro-5-fluoro-N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-4-pyrimidineamine

In like manner to the preparation of 2-chloro-N4-(3,4-ethyleneioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-[1-ethoxycarbonyl-1-methyl)ethyl]aniline were reacted to produce 2-chloro-5-fluoro-N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-4-pyrimidineamine. ^1H NMR (DMSO- d_6): δ 9.99 (s, 1H), 8.30 (d, 1H, J = 3.5 Hz), 7.60 (d, 2H, J = 8.8 Hz), 7.30 (d, 2H, J = 8.8 Hz), 4.04 (qt, 2H, J = 7.0 Hz), 1.47 (s, 6H), 1.10 (t, 3H, J = 7.0 Hz); LCMS: ret. time: 31.07 min.; purity: 97%; MS (m/e): 338 (MH^+).

7.1.56 2,4-Dichloro-5-ethoxycarbonylpyrimidine

A dry reaction flask equipped with a stirring bar and a reflux condenser was charged with 5-ethoxycarbonyluracil (1.84g, 10 mmol), $POCl_3$ (10 mL) and N,N -dimethylaniline (1 mL) and heated at 90 °C for 2h. The excess $POCl_3$ was removed under a reduced pressure and quenched with ice-water (100 g). The aqueous solution was extracted with ethyl ether (3 x 100 mL), washed with saturated aqueous $NaHCO_3$ solution and water (100 mL, each). After drying over sodium sulfate, the ethyl ether was removed and the residue was dried under a high vacuum to afford 2,4-dichloro-5-ethoxycarbonylpyrimidine. ^1H NMR ($CDCl_3$): δ 9.00 (s, 1H), 4.45 (q, 2H, J = 6.9 Hz), 1.42 (t, 3H, J = 6.9 Hz).

7.1.57 N-(2-Chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester (R926518) and N-(4-Chloro-5-ethoxycarbonyl-2-pyrimidinyl)-L-phenylalanine Ethyl Ester (R926519)

A mixture of L-phenylalanine Ethyl Ester Hydrochloride (0.137g, 0.6 mmol) 2,4-dichloro-5-ethoxycarbonylpyrimidine (0.112g, 0.5 mmol), triethylamine (0.7 mL, 0.6 mmol) in THF (4 mL) in a sealed tube was heated at 100 °C for 3h. The reaction was diluted with H_2O (20 mL), extracted with CH_2Cl_2 (3 x 50 mL), washed with 2N HCl (10 mL), water (10 mL) and solvent was evaporated. The residue obtained was purified by preparative TLC using 15% EtOAc in hexanes to obtain two products mainly, N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester (**R926518**). ^1H NMR ($CDCl_3$): δ 8.72 (d, 1H,

J= 6.92 Hz), 8.66 (s, 1H), 7.32-7.17 (m, 5H), 5.05 (dq, 1H, J= 1.2 and 5.7 Hz), 4.34 (q, 2H, J= 6.9 Hz), 4.20 (q, 2H, J= 5.1 Hz), 3.24 (dd, 1H, J= 5.4 Hz), 3.16 (dd, 1H, J= 7.5 Hz), 1.35 (t, 3H, J= 7.2 Hz), 1.24 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 37.15 min.; purity: 99%; MS (m/e): 378 (MH⁺) and N-(4-chloro-5-ethoxycarbonyl-2-pyrimidinyl)-L-phenylalanine Ethyl Ester (**R926519**). ¹H NMR (CDCl₃): δ 8.83 (s, 1H), 7.28 (m, 3H), 7.18 (m, 2H), 6.00 (bt, 1H), 4.99 (bdq, 1H), 4.36 (q, 2H, J= 7.8 Hz), 4.19 (q, 2H, J= 6.9 Hz), 3.20 (t, 2H, J= 6.9 Hz), 1.38 (t, 3H, J= 4.5 Hz), 1.24 (t, 3H, J= 6 Hz); LCMS: ret. time: 34.80 min.; purity: 88%; MS (m/e): 378 (M⁺).

7.1.58 N-(2-Chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-valine Ethyl Ester (R926520) and

N-(4-Chloro-5-ethoxycarbonyl-2-pyrimidinyl)-L-valine Ethyl Ester (R926521)

In like manner to the preparation of N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester, 2,4-dichloro-5-ethoxycarbonylpyrimidine and L-valine Ethyl Ester were reacted to prepare N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-valine Ethyl Ester (**R926520**). ¹H NMR (CDCl₃): δ 8.80 (d, 1H, J= 8.1 Hz), 8.68 (s, 1H), 4.77 (dd, 1H, J= 4.8 Hz), 4.36 (q, 2H, J= 7.2 Hz), 4.24 (q, 2H, J= 6.6 Hz), 2.38 (m, 1H), 1.39 (t, 3H, J= 6.9 Hz), 1.29 (t, 3H, J= 7.2 Hz), 1.03 (d, 3H, J= 3 Hz), 1.00 (d, 3H, J= 2.7 Hz); LCMS: ret. time: 36.54 min.; purity: 89%; MS (m/e): 330 (MH⁺) and N-(4-chloro-5-ethoxycarbonyl-2-pyrimidinyl)-L-valine Ethyl Ester (**R926521**). ¹H NMR (CDCl₃): δ 8.82 (s, 1H), 6.02 (m, 1H), 4.69 (dd, 1H, J= 4.8 and 4.5 Hz), 4.33 (q, 2H, J= 7.5 Hz), 4.23 (q, 2H, J= 7.5 Hz), 2.28 (sept, 1H), 1.34 (t, 3H, J= 6.9 Hz), 1.28 (t, 3H, J= 7 Hz), 1.00 (d, 6H, J= 7.2 Hz); LCMS: ret. time: 33.53 min.; purity: 91%; MS (m/e): 330 (M⁺).

7.1.59 N-(2-Chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-leucine Ethyl Ester (R926522)

In like manner to the preparation of N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester, 2,4-dichloro-5-ethoxycarbonylpyrimidine and L-leucine Ethyl Ester were reacted to prepare N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-leucine Ethyl Ester. ¹H NMR (CDCl₃): δ 8.69 (s, 1H), 8.64 (d, 1H, 7.8 Hz), 4.84 (s, 1H), 4.38 (q, 2H, J= 7.2 Hz), 3.75 (s, 3H), 1.73 (m, 2H), 1.39 (t, 3H, J= 6.9 Hz), 0.97 (d, 3H, J= 4.2 Hz), 0.95 (d, 3H, J= 4.8 Hz); LCMS: ret. time: 36.09 min.; purity: 92%; MS (m/e): 330 (MH⁺).

7.1.60 N-(2-Chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-alanine Ethyl Ester (R926523) and

N-(4-Chloro-5-ethoxycarbonyl-2-pyrimidinyl)-L-alanine Ethyl Ester (R926524)

5 In like manner to the preparation of N-(2-chloro-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester, 2,4-dichloro-5-ethoxycarbonylpyrimidine and L-valine Ethyl Ester were reacted to prepare N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-alanine Ethyl Ester (**R926523**). ¹H NMR (CDCl₃): δ 8.80 (bd, 1H), 8.68 (s, 1H), 4.79 (q, 1H, J = 7.2 Hz), 4.35 (q, 2H, J = 7.2 Hz), 4.24 (m, 2H), 1.53 (d, 3H, J = 7.2 Hz), 1.38 (t, 3H, J = 7.2 Hz), 1.29 (t, 3H, J = 7.2 Hz); LCMS: ret. time: 31.89 min.; purity: 94%; MS (m/e): 303 (MH⁺) and N-(4-chloro-5-ethoxycarbonyl-2-pyrimidinyl)-L-alanine Ethyl Ester (**R926524**). ¹H NMR (CDCl₃): δ 8.80 (s, 1H), 6.01 (bs, 1H), 4.65 (bq, 1H), 4.35 (q, 2H), 4.20 (q, 2H), 1.55, t, 3H), 1.40 (t, 3H), 1.25 (t, 3H); LCMS: ret. time: 28.78 min.; purity: 84%; MS (m/e): 302 (M⁺).

7.1.61 2-Chloro-N4-(4-n-butyloxyphenyl)-5-fluoro-4-pyrimidineamine

To a solution of 2,4-dichloro-5-fluoropyrimidine (0.5 g, 3.0 mmol) and 4-n-butoxyaniline (0.49 g, 3 mmol) in acetone/H₂O (1:9 mL) at room temperature was added concentrated HCl (0.1 mL). The mixture was heated at reflux for 1 h, cooled to room temperature, and made basic with 2 N NaOH (2 mL). The aqueous layer was extracted with EtOAc (2 x 50 mL) and the combined organic extracts were dried (Na₂SO₄), filtered, and concentrated in vacuo. The crude black solid was purified by chromatography (4:1 hexanes/EtOAc) to afford 2-chloro-N4-(4-n-butyloxyphenyl)-5-fluoro-4-pyrimidineamine (0.71 g, 80%) as a brown oil: ¹H NMR (300 MHz, CDCl₃) δ 8.01 (d, J = 2.7 Hz, 1H), 7.51-7.46 (m, 2H), 6.95-6.89 (m, 2H), 6.83 (bs, 1H), 3.99-3.95 (t, J = 6.5 Hz, 2H), 1.82-1.57 (m, 2H), 1.53-1.43 (m, 2H), 0.98 (t, J = 7.2 Hz, 3H).

7.1.62 2-Chloro-N4-(4-n-hexyloxyphenyl)-5-fluoro-4-pyridineamine

In like manner to the preparation of 2-chloro-N4-(4-n-butyloxyphenyl)-5-fluoro-4-pyrimidineamine, the reaction of 2,4-dichloro-5-fluoropyrimidine with 4-n-hexyloxyaniline gave 2-chloro-N4-(4-n-hexyloxyphenyl)-5-fluoro-4-pyridineamine. The crude product was purified by chromatography (4:1 CHCl₃/EtOAc) to afford (14) (0.74 g, 76%) as a red-brown oil that solidified upon standing: ¹H NMR (300 MHz, CDCl₃) δ 8.01 (d, J = 2.7 Hz, 1H),

7.50 (d, $J = 9.0$ Hz, 2H), 6.92 (d, $J = 9.0$ Hz, 2H), 6.84 (bs, 1H), 3.96 (t, $J = 6.5$ Hz, 2H), 1.83-1.74 (m, 2H), 1.48-1.41 (m, 2H), 1.36-1.34 (m, 4H), 0.93-0.89 (m, 3H).

7.1.63 N4-(3-Benzyloxyphenyl)-2-chloro-4-pyrimidineamine

A mixture of 2,6-dichloropyrimidine (2.00 g, 13.4 mmol), 3-benzyloxyaniline (2.07 g, 13.4 mmol) and triethylamine (2.72 g, 26.8 mmol) in 1-butanol (20 mL) was stirred at 50 °C for 17 h. The reaction mixture was concentrated to remove most of the 1-butanol, the crude product was preadsorbed onto silica gel using chloroform and purified by flash chromatography (95:5 chloroform/ methanol) to afford N4-(3-benzyloxyphenyl)-2-chloro-4-pyrimidineamine (1.70 g, 40%) as colorless oil: ^1H NMR (300 MHz, DMSO- d_6) δ 10.2 (s, 1H), 8.16 (d, $J = 6.0$ Hz, 1H), 7.48-7.24 (m, 7H), 7.12 (d, $J = 9.0$ Hz, 1H), 6.78 (m, 2H), 5.11 (s, 2H); ESI MS m/z 312 [$\text{C}_{17}\text{H}_{14}\text{ClN}_3\text{O} + \text{H}$] $^+$.

7.1.64 N4-[4-(tert-Butoxycarbonylmethyleneoxy)phenyl]-3-chloro-5-ethoxycarbonyl-4-pyrimidineamine (R926578)

In like manner to the preparation of N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester, 5-carboxyethoxy-2,4-dichloropyrimidine and tert-butyl 4-aminophenoxyacetate were reacted to prepare N4-[4-(tert-butoxycarbonylmethyleneoxy)phenyl]-2-chloro-5-ethoxycarbonyl-2-chloro-4-pyrimidineamine. LCMS: MS (m/e): 407 (MH^+).

7.1.65 N4-(4-Ethoxyphenyl)-5-ethoxycarbonyl-2-trifluoromethyl-4-pyrimidineamine (R926059)

In like manner to the preparation of N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester, 4-chloro-5-ethoxycarbonyl-2-trifluoromethylpyrimidine and 4-ethoxyaniline were reacted to prepare N4-(4-ethoxyphenyl)-5-ethoxycarbonyl-2-trifluoromethyl-4-pyrimidineamine. ^1H NMR (CDCl_3): δ 10.39 (s, 1H), 9.02 (s, 1H), 7.59 (dd, 2H, $J = 2.1$ and 7.2 Hz), 6.91 (dd, 2H, $J = 1.8$ and 6.6 Hz), 4.44 (q, 2H, $J = 7.5$ Hz), 4.06 (q, 2H, $J = 7.2$ Hz), 1.44 (m, 6H); LCMS: ret. time: 38.49 min.; purity: 100%; MS (m/e): 356 (MH^+).

7.1.66 N2-(4-Ethoxyphenyl)-5-methoxycarbonyl-4-trifluoromethyl-2-pyrimidineamine (R926060)

In like manner to the preparation of N-(2-chloro-5-ethoxycarbonyl-4-pyrimidinyl)-L-phenylalanine Ethyl Ester, 2-chloro-5-methoxycarbonyl-4-trifluoromethylpyrimidine and

4-ethoxyaniline were reacted to prepare N2-(2-ethoxyphenyl)-5-methoxycarbonyl-4-trifluoromethyl-2-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.98 (s, 1H), 7.47 (m, 3H), 6.91 (dd, 2H, J= 2.1 and 6.9 Hz), 4.05 (q, 2H, 6.9 Hz), 1.42 (t, 3H, J= 6.8 Hz); ¹⁹F NMR (CDCl₃): -19105; LCMS: ret. time: 33.87 min; purity: 100%; MS (m/e): 342 (MH⁺).

5 **7.1.67 2-Chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine (R926853)**

A reaction mixture containing 2,4-dichloro-5-fluoropyrimidine (1.2 equivalents) and 3-(tetrazol-5-yl)aniline (1 equivalents) in methanol:water (1:1; v/v) was heated at 60 °C for 24 h. Upon dilution with water and acidification, the solid formed was filtered, washed
10 with water, dried and analyzed to give 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine (R926853). Alternatively this reaction can be achieved by treating 2,4-dichloro-5-fluoropyrimidine (1 equivalent) with 3-(tetrazol-5-yl)aniline (3 equivalents) in methanol:water (1:1; v/v) at 60 °C for 2-3 hours or at room temperature for 24 h to give 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine. ¹H NMR (DMSO-
15 d₆): δ 10.25 (s, 1H), 8.43 (s, 1H), 8.37 (d, 1H, J= 3.6 Hz), 7.90 (dd, 1H, J= 0.9 and 9 Hz), 7.75 (d, 1H, J= 7.5 Hz), 7.61 (t, 1H, J= 7.8 Hz); LCMS: purity: 90%; MS (m/e): 292 (MH⁺).

7.1.68 2-Chloro-N4-(2,5-dimethoxy-4-chlorophenyl)-5-fluoro-4-pyrimidineamine (R926858)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 2,5-
20 dimethoxy-4-chloroaniline gave 2-chloro-N4-(2,5-dimethoxy-4-chlorophenyl)-5-fluoro-4-pyrimidineamine. LCMS: purity: 97%; MS (m/e): 316 (M-2H) and 320 (M+2H).

7.1.69 2-Chloro-5-fluoro-N4-(3-methoxycarbonyl-5-trifluoromethylphenyl)-4-pyrimidineamine (R926861)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-methoxycarbonyl-5-trifluoromethylaniline gave 2-chloro-5-fluoro-N4-(3-methoxycarbonyl-5-trifluoromethylphenyl)-4-pyrimidineamine. ¹H NMR (CD₃OD): δ 8.60 (s, 1H), 8.43 (s, 1H), 8.20 (d, 1H, J= 3 Hz), 7.99 (s, 1H), 3.96 (s, 3H); ¹⁹F NMR (CD₃OD): -18332, - 18374;
25 and - 44259; LCMS: purity: 91%; MS (m/e): 350 (MH⁺).

7.1.70 2-Chloro-5-fluoro-N4-[3-(2-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-4-pyrimidineamine (R926869)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-(2-phenyl-1,3,4-oxadiazol-5-yl)aniline gave 2-chloro-5-fluoro-N4-[3-(2-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 10.28 (s, 1H), 8.62 (s, 1H), 8.39 (d, 1H, J= 3.3 Hz), 8.11 (m, 2H), 7.98 (bd, 1H, J= 6.9 Hz), 7.88 (bd, 1H, J= 8.4 Hz), 7.65 (m, 4H); LCMS: purity: 76%; MS (m/e): 76%.

7.1.71 2-Chloro-N4-[3-(2-ethoxycarbonylmethylene-1,3,4-oxadiazol-5-yl)phenyl]-5-fluoro-4-pyrimidineamine (R926873)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-(2-ethoxycarbonylmethylene-1,3,4-oxadiazol-5-yl)aniline gave 2-chloro-N4-[3-(2-ethoxycarbonylmethylene-1,3,4-oxadiazol-5-yl)phenyl]-5-fluoro-4-pyrimidineamine. ¹H NMR (CD₃OD): δ 8.42 (t, 1H, J= 1.8 Hz), 8.19 (d, 1H, J= 3.3 Hz), 7.99 (dt, 1H, J= 1.2 and 8.1 Hz), 7.82 (dt, 1H, J= 1.2 and 8.1 Hz), 7.58 (t, 1H, J= 9 Hz), 4.24 (q, 2H, J= 3.9 Hz), 4.17 (s, 2H), 1.28 (t, 3H, J= 7.2 Hz); LCMS: purity: 85%; MS (m/e): 379 (MH⁺).

7.1.72 2-Chloro-5-fluoro-N4-(4-trifluoromethoxyphenyl)-4-pyrimidineamine (R926875)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 4-trifluoromethoxyaniline gave 2-chloro-5-fluoro-N4-(4-trifluoromethoxyphenyl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.11 (d, 1H, J= 2.1 Hz), 7.68 (dd, 2H, J= 2.4 and 7.6 Hz), 7.26 (dd, 2H, J= 3 and 8.7 Hz), 7.0 (bs, 1H); ¹⁹F NMR (CD₃OD): δ -16517 and -44523; LCMS: purity: 94%; MS (m/e): 308 (MH⁺).

7.1.73 2-Chloro-5-fluoro-N4-(4-trifluoromethylphenyl)-4-pyrimidineamine (R926876)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 4-trifluoromethylaniline gave 2-chloro-5-fluoro-N4-(4-trifluoromethylphenyl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.15 (d, 2.1 Hz), 7.80 (d, 2H, J= 7.1 Hz), 7.66 (d,

2H, J= 9 Hz), 7.10 (bs, 1H); ^{19}F NMR (CDCl_3): -17682 and - 44362; LCMS: purity: 91% and MS (m/z): 292 (MH^+).

7.1.74 2-Chloro-N4-(4-chloro-3-trifluoromethylphenyl)-5-fluoro-4-pyrimidineamine (R926877)

5 In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 4-chloro-3-trifluoromethylaniline gave 2-chloro-N4-(4-chloro-3-trifluoromethylphenyl)-5-fluoro-4-pyrimidineamine. ^1H NMR (CDCl_3): δ 8.15 (d, 1H, J= 2.1 Hz), 7.96 (d, 1H, J= 3 Hz), 7.91 (dd, 1H, J= 2.7 Hz and 8.7 Hz), 7.53 (d, 1H, J= 8.1 Hz), 7.06 (bs, 1H); ^{19}F NMR (CDCl_3): - 17892 and - 44402; LCMS: purity: 93%; MS (m/e): 326 (M^+).

7.1.75 2-Chloro-5-fluoro-N4-(6-methoxypyridin-3-yl)-4-pyrimidineamine (R926878)

15 In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-amino-6-methoxypyridine gave 2-chloro-5-fluoro-N4-(6-methoxypyridin-3-yl)-4-pyrimidineamine. ^1H NMR (CD_3OD): δ 8.39 (d, 1H, J= 3.0 Hz), 8.10 (d, 1H, J= 3.6 Hz), 7.95 (dd, 1H, J= 2.4 and 9 Hz), 8.30 (d, 1H, J= 9 Hz), 3.91 (s, 3H); ^{19}F NMR (CD_3OD): - 44737; LCMS: purity: 97%; MS (m/e): 255 (M^+).

7.1.76 2-Chloro-N4-(3,4-difluorophenyl)-5-fluoro-4-pyrimidineamine (R926882)

20 In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3,4-difluoroaniline gave 2-chloro-N4-(3,4-difluorophenyl)-5-fluoro-4-pyrimidineamine. ^1H NMR (CDCl_3): δ 8.10 (d, 1H, J= 2.1 Hz), 7.72 (m, 1H), 7.22 (m, 2H), 6.95 (bs, 1H); LCMS: purity: 93%; MS (m/e): 260 (M^+).

7.1.77 2-Chloro-N4-(3,4-Dichlorophenyl)-5-fluoro-4-pyrimidineamine (R926884)

30 In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3,4-dichloroaniline gave 2-chloro-N4-(3,4-dichlorophenyl)-5-fluoro-4-pyrimidineamine. LCMS: purity: 95%; MS (m/e): 294 (M^+ , 2H).

7.1.78 2-Chloro-5-fluoro-N4-(6-methylpyridin-2-yl)-4-pyrimidineamine (R926888)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 2-amino-6-methylpyridine gave 2-chloro-5-fluoro-N4-(6-methylpyridin-2-yl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.23 (s, 1H), 8.19 (s, 1H), 8.12 (d, 1H, J= 3 Hz), 7.55 (bs, 1H), 7.69 (t, 1H, J= 7.4 Hz), 9.35 (d, 1H, J= 7.5 Hz); ¹⁹F NMR (CDCl₃): - 44073; LCMS: purity: 96%; MS (m/e): 239 (M⁺).

7.1.79 2-Chloro-N4-(2,6-Dimethoxypyridin-3-yl)-5-fluoro-4-pyrimidineamine (R926889)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-amino-2,6-dimethoxypyridine gave 2-chloro-N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.57 (d, 1H, J= 8.7 Hz), 8.02 (d, 1H, J= 2.7 Hz), 6.40 (d, 1H, J= 8.1 Hz), 4.03 (s, 3H), 3.98 (s, 3H); ¹⁹F NMR (CDCl₃): - 44640; LCMS: purity: 90%; MS (m/e): 285 (M⁺).

7.1.80 2-Chloro-N4-(6-chloropyridin-3-yl)-5-fluoro-4-pyrimidineamine (R920400)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-amino-6-chloropyridine gave 2-chloro-N4-(6-chloropyridin-3-yl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.53 (d, 1H, J= 3 Hz), 8.25 (dd, 1H, J= 3 and 9 Hz), 8.15 (d, 1H, J= 2.4 Hz), 7.39 (d, 1H, J= 8.7 Hz), 7.00 (bs, 1H); LCMS: purity: 98%; MS (m/e): 259 (M⁺).

7.1.81 2-Chloro-5-fluoro-N4-(4-methylpyridin-2-yl)-4-pyrimidineamine (R920401)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 2-amino-4-methylpyridine gave 2-chloro-5-fluoro-N4-(4-methylpyridin-2-yl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.22 (s, 1H), 8.16 (d, 1H, J= 8.4 Hz), 8.13 (d, 1H, J= 2.4 Hz), 6.91 (d, 1H, J= 5.4 Hz), 2.42 (s, 3H); LCMS: purity: 87%; MS (m/e): 239 (M⁺).

7.1.82 2-Chloro-5-fluoro-N4-(3-trifluoromethoxyphenyl)-4-pyrimidineamine (R920402)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-trifluoromethoxyaniline gave 2-chloro-5-fluoro-N4-(3-trifluoromethoxyphenyl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.12 (d, 1H, J= 3 Hz), 7.68 (bs, 1H), 7.53 (dd, 1H, J= 1.2 and 8.4 Hz), 7.41 (t, 1H, J= 8.1 Hz), 7.04 (bdt, 2H); ¹⁹F NMR (CDCl₃): -16430 and -44463; LCMS: purity: 89%; MS (m/e): 308 (MH⁺).

7.1.83 2-Chloro-N4-(3,4-Difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (R920403)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3,4-difluoromethylenedioxyaniline gave 2-chloro-N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.09 (d, 1H, J= 3 Hz), 7.70 (d, 1H, J= 2.4 Hz), 7.10 (dd, 1H, J= 2.4 and 8.7 Hz), 7.06 (t, 1H, J= 8.1 Hz), 6.97 (bs, 1H); ¹⁹F NMR (CDCl₃): - 14175 and - 44562; LCMS: purity: 95%; MS (m/e): 304 (MH⁺).

7.1.84 2-Chloro-5-fluoro-N4-(quinolin-6-yl)-4-pyrimidineamine (R920409)

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 6-aminoquinoline gave 2-chloro-5-fluoro-N4-(quinolin-6-yl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.02 (dd, 1H, J= 2.7 Hz), 8.00 (dd, 1H, J= 2.4 Hz), 7.73 (d, 1H, J= 9 Hz), 7.68 (dd, 1H, J= 2.4 and 8.7 Hz), 7.28 (t, 1H, J= 10.5 Hz), 6.42 (d, 1H, J= 9.3 Hz); ¹⁹F NMR (CDCl₃): - 44344; LCMS: purity: 91%; MS (m/e): 292 (M⁺).

7.1.85 2-Chloro-N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-chloro-4-trifluoromethoxyaniline gave 2-chloro-N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.15 (d, 1H, J= 3.0 Hz), 7.86 (d, 1H, J= 2.1 Hz), 7.61 (dd, 1H, J= 2.1 and 8.7 Hz), 7.35 (dd, 1H, J= 1.2 and 8.7 Hz), 6.98 (bs, 1H); LCMS: purity: 97%; MS (m/e): 342 (M+2H).

7.1.86 2-Chloro-N4-(4-chloro-3-methoxyphenyl)-5-fluoro-4-pyrimidineamine

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 4-chloro-3-methoxyaniline gave 2-chloro-N4-(4-chloro-3-methoxyphenyl)-5-fluoro-4-aminopyrimidine. LCMS: purity: 88%; MS (m/e): 288 (MH⁺).

7.1.87 2-Chloro-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-4-pyrimidinediamine

In like manner to the preparation of 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine the reaction of 2,4-dichloro-5-fluoropyrimidine with 5-amino-2-(2-hydroxyethyloxy)pyridine gave 2-chloro-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.28 (d, 1H, J= 2.4 Hz), 8.08 (m, 1H), 7.99 (m, 1H), 7.00 (bs, 1H), 6.87 (bd, 1H), 4.47 (m, 2H), 3.97 (m, 2H).

7.1.88 2-Chloro-N4-[2-(2-chloro-5-fluoropyrimidin-4-yl)-1,2,3,4-tetrahydroisoquinolin-7-yl]-5-fluoro-4-pyrimidineamine (R926910)

In a like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-1,2,3,4-tetrahydroisoquinoline were reacted to provide 2-chloro-N4-[2-(2-chloro-5-fluoropyrimidin-4-yl)-1,2,3,4-tetrahydroisoquinolin-7-yl]-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.08 (d, 1H, J= 3.0 Hz), 7.95 (d, 1H, J= 6.0 Hz), 7.50-7.42 (m, 2H), 7.21 (d, 1H, J= 8.4 Hz), 6.96-6.90 (m, 1H), 4.95 (s, 2H), 4.04 (t, 2H, J= 5.7 Hz), 2.99 (t, 2H, J= 5.7 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -42555, -44573; LCMS: purity: 98%; MS (m/e): 410(MH⁺).

7.1.89 2-Chloro-5-fluoro-N4-[2-(t-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-7-yl]-4-pyrimidineamine (R926911)

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-2-(t-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinoline were reacted to provide 2-chloro-5-fluoro-N4-[2-(t-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-7-yl]-4-pyrimidineamine.

¹H NMR (CDCl₃): δ 8.03 (s, 1H), 7.50-7.26 (m, 2H), 7.19-7.11 (m, 2H), 4.57 (s, 2H), 3.64 (t, 2H, J= 5.7 Hz), 2.80 (t, 2H, J= 5.7 Hz), 1.48 (s, 9H); LCMS: purity: 89%; MS (m/e): 379(M⁺).

5

7.1.90 2-Chloro-5-fluoro-N4-(1,2,3,4-tetrahydroisoquinolin-7-yl)-4-pyrimidineamine (R926912)

A solution of 2-chloro-5-fluoro-N4-[2-(t-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-7-yl]-4-pyrimidineamine in 40% trifluoroacetic acid/dichloromethane was stirred at rt for 30 min. Removal of the solvent left an oily residue which was suspended in water, made basic with NaHCO₃, and extracted with ethyl acetate.

10

Purification by column chromatography over silica gel provided 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydroisoquinolin-7-yl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.04 (d, 1H, J= 3.0 Hz), 7.37 (dd, 1H, J= 2.4 and 8.4 Hz), 7.27 (d, 1H, J= 1.5 Hz), 7.11 (d, 1H, J= 8.4 Hz), 6.92 (s, 1H), 4.04 (s, 2H), 3.15 (t, 2H, J= 6.0 Hz), 2.79 (t, 2H, J= 6.0 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -44648; LCMS: purity: 97%; MS (m/e): 279(MH⁺).

15

7.1.91 2-Chloro-5-fluoro-N4-(4-methyl-3-trifluoromethylphenyl)-4-pyrimidineamine (R926920)

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-methyl-3-trifluoromethylaniline were reacted to provide 2-chloro-5-fluoro-N4-(4-methyl-3-

20

trifluoromethylphenyl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.10 (d, 1H, J= 3.0 Hz), 7.85-7.78 (m, 2H), 7.33 (d, 1H, J= 9.3 Hz), 6.96 (bs, 1H), 2.48 (d, 3H, J= 1.2 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -17641, -44541; LCMS: purity: 97%; MS (m/e): 306(MH⁺).

7.1.92 2-Chloro-5-fluoro-N4-(4-fluoro-3-methylphenyl)-4-pyrimidineamine (R926921)

25

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-fluoro-3-methylaniline were reacted to provide 2-chloro-5-fluoro-N4-(4-fluoro-3-methylphenyl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.06 (d, 1H, J= 2.4 Hz), 7.48-7.43 (m, 1H), 7.39 (dd, 1H, J= 2.7 and 6.3 Hz), 7.03 (t, 1H, J= 9.0 Hz), 6.84 (bs, 1H), 2.30 (d, 1H, J= 1.8 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -34285, -44676; LCMS: purity: 95%; MS (m/e): 257(MH⁺).

30

7.1.93 N4-[3-[(N-t-butoxycarbonyl)aminomethyl]-4-methylphenyl]-2-chloro-5-fluoro-4-pyrimidineamine (R926924)

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-[(N-t-butoxycarbonyl)aminomethyl]-4-methylaniline were reacted to provide N4-[3-[(N-t-butoxycarbonyl)aminomethyl]-4-methylphenyl]-2-chloro-5-fluoro-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.05 (d, 1H, J= 3.0 Hz), 7.52 (d, 1H, J= 9.3 Hz), 7.45 (s, 1H), 7.19 (d, 1H, J= 8.1 Hz), 6.96-6.89 (m, 1H), 4.80 (bs, 1H), 2.31 (s, 2H), 1.46 (s, 9H); LCMS: purity: 97%; MS (m/e): 311 (M – (t-butyl)⁺).

7.1.94 2-Chloro-N4-[3-[[4-(ethoxycarbonyl)piperidino]methyl]phenyl]-5-fluoro-4-pyrimidineamine

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and ethyl 1-(3-aminobenzyl)piperidine-4-carboxylate were reacted to provide 2-chloro-N4-[3-[[4-(ethoxycarbonyl)piperidino]methyl]phenyl]-5-fluoro-4-pyrimidineamine. LCMS: purity: 97%; MS (m/e): 394(MH⁺).

7.1.95 2-Chloro-N4-[3-[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-4-pyrimidineamine

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-[[4-(ethoxycarbonyl)piperidino]carbonyl]aniline were reacted to provide 2-chloro-N4-[3-[[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-4-pyrimidineamine. LCMS: purity: 96%; MS (m/e): 407(M⁺).

7.1.96 2-Chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-4-pyrimidineamine

In a manner similar to the preparation of N4-(3,4-ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-oxonaphthalen-7-yl)-4-pyrimidineamine was reduced with Dibal-H to yield 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-4-pyrimidineamine.

¹H NMR (CDCl₃): δ 8.05 (d, 1H, J= 3.0 Hz), 7.59 (d, 1H, J= 2.4 Hz), 7.14 (d, 1H, J= 8.1 Hz), 6.93 (bs, 1H), 4.82-4.78 (m, 1H), 2.82-2.71 (m, 2H), 2.08-1.74 (m, 5H) ; ¹⁹F NMR (282 MHz, CDCl₃): -44661; LCMS: purity: 94%; MS (m/e): 294(MH⁺).

7.1.97 2-Chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-oxonaphthalen-7-yl)-4-pyrimidineamine.

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-1-tetralone were reacted to provide 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-oxonaphthalen-7-yl)-4-pyrimidineamine. ¹H NMR (DMSO-*d*₆): δ 10.08 (s, 1H), 8.31 (d, 1H, J= 3.3 Hz), 8.15 (d, 1H, J= 2.4 Hz), 7.82 (dd, 1H, J= 2.4 and 8.1 Hz), 7.36 (d, 1H, J= 8.1 Hz), 2.91 (t, 2H, J= 6.0 Hz), 2.59 (t, 2H, J= 6.0 Hz), 2.07-1.98 (m, 2H); LCMS: purity: 93%; MS (m/e): 294(MH⁺).

7.1.98 2-Chloro-5-fluoro-N4-[3-(trifluoromethylthio)phenyl]-4-pyrimidineamine

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-(trifluoromethylthio)aniline were reacted to provide 2-chloro-5-fluoro-N4-[3-(trifluoromethylthio)phenyl]-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 8.13 (bs, 1H), 7.92 (bs, 1H), 7.89-7.84 (m, 1H), 7.48-7.45 (m, 2H), 7.04 (bs, 1H); LCMS: purity: 97%; MS (m/e): 325(MH⁺).

7.1.99 2-Chloro-5-fluoro-N4-[(3-dihydroxyboryl)phenyl]-4-pyrimidineamine

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-aminobenzenboronic acid were reacted to provide 2-chloro-5-fluoro-N4-[(3-dihydroxyboryl)phenyl]-4-pyrimidineamine.

7.1.100 2-Chloro-5-fluoro-N4-[(1H)-indol-6-yl]-4-pyrimidineamine

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-aminoindole were reacted to provide 2-chloro-5-fluoro-N4-[(1H)- indol-6-yl]-4-pyrimidineamine. LCMS: purity: 92%; MS (m/e): 263(MH⁺).

7.1.101 2-Chloro-5-fluoro-N4-(2-hydroxy-4-methylphenyl)-4-pyrimidineamine

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-hydroxy-4-methylaniline were reacted to provide 2-chloro-5-fluoro-N4-(2-hydroxy-4-methylphenyl)-4-pyrimidineamine. LCMS: purity: 97%; MS (m/e): 255(MH⁺).

7.1.102 2-Chloro-5-fluoro-N4-[2-(methoxycarbonyl)-(1H)-indol-6-yl]-4-pyrimidineamine

In a like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-2-(methoxycarbonyl)-(1H)-indole were reacted to provide 2-chloro-5-fluoro-N4-[2-(methoxycarbonyl)-(1H)-indol-6-yl]-4-pyrimidineamine which was used without further purification. LCMS: purity: 65%; MS (m/e): 322(MH⁺).

7.1.103 N4-[3-(4-(2-Chloro-5-fluoropyrimidine)-N-aminomethylene)-phenyl]-2-chloro-5-fluoro-4-pyrimidineamine (R940298)

The reaction flask equipped with a magnetic stirring bar and a rubber septum (to prevent loss of 2,4-dichloro-5-fluoropyrimidine and N₂ inlet was charged 3-aminobenzylamine (0.22 g, 1.79 mmol), MeOH (1 mL), H₂O (3 mL) and 2,4-dichloro-5-fluoropyrimidine (0.3 g, 1.79 mmol). The reaction mixture was stirred at 80°C for 30 min., cool to room temperature, diluted with H₂O (30 mL). Upon saturation with sodium chloride it was extracted with ethyl acetate (3 x 20 mL), dried over anhydrous sodium sulfate and the solvent was removed. The resulting residue was filtered through a pad of silica gel (200-400 mesh) using 1 to 3% MeOH in CH₂Cl₂ to obtain N4-[3-(4-(2-chloro-5-fluoropyrimidine)-N-methylaminomethylene)-phenyl]-2-chloro-5-fluoro-4-pyrimidineamine **R940298**. ¹H NMR (DMSO-d₆): δ 10.09 (1H, s), 8.88 (1H, t, J= 5.85 Hz), 8.40 (1H, d, J= 3.6 Hz), 8.23 (1H, d, J= 3.3 Hz), 7.74 (1H, s), 7.70 (1H, d, J= 8.1 Hz), 7.44 (1H, t, J= 7.8 Hz), 7.19 (1H, d, J= 8.1 Hz), 4.69 (2H, d, J= 5.7 Hz ; purity 92 %).

7.1.104 2-Chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine (R940302)

The reaction flask equipped with a magnetic stirring bar and a rubber septum (to prevent loss of 2,4-dichloro-5-fluoropyrimidine and N₂ inlet was charged with 3-methyloxycarbonyl-4-methoxyaniline (0.88 g, 4.86 mmol), MeOH (3 mL), H₂O (7 mL) and

2,4-dichloro-5-fluoropyrimidine (0.81 g, 4.86 mmol). The reaction mixture was stirred at 60°C for 30 min., diluted with H₂O (50 mL), acidified with 2N HCl (6 mL) and sonicated. The solid obtained was filtered, washed with H₂O and dried to produce 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine **R940302**. ¹H NMR (DMSO-d₆): δ 10.10 (1H, s), 8.39 (1H, d, *J* = 3.6 Hz), 8.04 (1H, d, *J* = 2.7 Hz), 7.98-7.93 (1H, m), 7.30 (1H, d, *J* = 9 Hz), 3.92 (3H, s), 3.89 (3H, m) ; purity 96% ; MS (m/e): 312 (MH⁺).

7.1.105 2-Chloro-5-fluoro-N4-(4-phahthlimide)-4-pyrimidineamine (R940303)

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 4-aminophthalimide were reacted to produce 2-chloro-5-fluoro-N4-(4-phahthlimide)-4-pyrimidineamine **R940303**. ¹H NMR (DMSO-d₆): δ 11.38 (1H, s), 10.60 (1H, s), 8.57 (1H, d, *J* = 3.3 Hz), 8.39 (1H, d, *J* = 1.8 Hz), 8.18 (1H, dd, *J* = 8.4 Hz, *J* = 2.1 Hz), 7.93 (1H, d, *J* = 8.1 Hz) ; purity 90% ; MS (m/e): 293 (MH⁺).

7.1.106 2-Chloro-5-fluoro-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-4-pyrimidineamine (R940305)

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-methylaminocarbonyl-4-methoxyaniline were reacted to produce 2-chloro-5-fluoro-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-4-pyrimidineamine **R940305**. ¹H NMR (DMSO-d₆): δ 9.91 (1H, s), 8.31 (1H, d, *J* = 3.6 Hz), 8.11 (1H, d, *J* = 2.7 Hz), 7.78 (1H, dd, *J* = 9 Hz, *J* = 2.7 Hz), 7.59 (1H, m), 6.87 (1H, d, *J* = 9 Hz), 3.90 (3H, s), 2.96 (3H, d, *J* = 4.5 Hz) ; purity 93%.

7.1.107 N2-Chloro-5-fluoro-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-4-pyrimidineamine (R940313)

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-(N-morpholinomethylene)-4-methoxyaniline were reacted to produce 2-chloro-5-fluoro-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-4-pyrimidineamine **R940313**. ¹H NMR (DMSO-d₆): δ 10.00 (1H, s), 8.35 (1H, d, *J* = 3.3 Hz), 7.72 (1H, d, *J* = 3 Hz), 7.58 (1H, d,

$J = 9.3$ Hz), 7.12 (1H, d, $J = 8.4$ Hz), 3.89 (3H, s), 3.8-3.5 (6H, m), 2.58 (4H, m); purity 96%; MS (m/e): 352 (M).

7.1.108 N4-[3-(N-*tert*-Butoxycarbonyl-N-methylaminomethylene)-phenyl]-2-chloro-5-fluoro-4-pyrimidineamine (R940315)

5 In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-(N-*tert*-butoxycarbonyl-N-methylaminomethylene)-aniline were reacted to produce N4-[3-(N-*tert*-butoxycarbonyl-N-methylaminomethylene)-phenyl]-2-chloro-5-fluoro-4-pyrimidineamine **R940315**. ^1H NMR (DMSO- d_6): δ 10.13 (1H, s), 8.42 (1H, d, $J = 3.6$ Hz), 7.69 (1H, m), 7.64 (1H, s), 7.45 (1H, t, $J = 7.6$ Hz), 7.09 (1H, d, $J = 7.8$ Hz), 4.48 (2H, s), 2.90 (3H, s), 1.49 (9H, m); purity 92%; MS (m/e): 367 (MH+).

7.1.109 N4-(3-(N-*tert*-Butoxycarbonyl-N-*iso*-propylaminomethylene)-4-methoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (R940320)

15 In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 3-(N-*tert*-butoxycarbonyl-N-*iso*-propylaminomethylene)-4-methoxy-aniline were reacted to produce N4-(3-(N-*tert*-butoxycarbonyl-N-*iso*-propylaminomethylene)-4-methoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine **R940320**. ^1H NMR (DMSO- d_6): δ 10.01 (1H, s), 8.34 (1H, d, $J = 3.6$ Hz), 7.52 (2H, m), 7.08 (1H, d, $J = 8.7$ Hz), 4.33 (3H, m), 3.90 (3H, s), 1.50-1.30 (9H, m), 1.18 (6H, d, $J = 6.9$ Hz); purity 95%.

7.1.110 2-Chloro-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine (R940322)

25 In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-2,2-dimethyl-4H-benzo[1,4]oxazin-3-one were reacted to produce 2-chloro-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine **R940322**. ^1H NMR (DMSO- d_6): δ 10.89 (1H, s), 10.04 (1H, s), 8.38 (1H, d, $J = 3.6$ Hz), 7.35 (2H, m), 7.04 (1H, d, $J = 8.4$ Hz), 1.50 (6H, s); purity 91.4%; MS (m/e): 322 (M).

7.1.111 2-Chloro-N4-[3-dihydro-2,2-dimethyl-4-(2-(pyridyl-1-oxide)-benzo[1,4]oxazin-6-yl]-5-fluoro-4-pyrimidineamine (R940328)

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 2-(6-amino-3-dihydro-2,2-dimethyl-benzo[1,4]oxazin-4-yl)pyridine 1-Oxide were reacted to produce 2-chloro-N4-[3-dihydro-2,2-dimethyl-4-(2-(pyridyl-1-oxide)-benzo[1,4]oxazin-6-yl]-5-fluoro-4-pyrimidineamine **R940328**. ¹H NMR (DMSO-d₆): δ 9.82 (1H, s), 8.39 (1H, dd, *J*= 6.3 Hz, *J*= 1.2 Hz), 8.30 (1H, d, *J*= 3.6 Hz), 7.63 (1H, dd, *J*= 8.4 Hz, *J*= 2.4 Hz), 7.47 (1H, td, *J*= 7.5 Hz, *J*= 1.8 Hz), 7.34 (1H, m), 7.21 (1H, dd, *J*= 8.7 Hz, *J*= 2.4 Hz), 7.07 (1H, d, *J*= 2.7 Hz), 6.91 (1H, d, *J*= 8.7 Hz), 3.64 (2H, s), 1.41 (6H, s) ; purity 95.8% ; MS (m/e): 402 (MH⁺).

7.1.112 2-Chloro-N4-[3-dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-4-pyrimidineamine (R940336)

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-3-dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazine were reacted to produce 2-chloro-N4-[3-dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-4-pyrimidineamine **R940336**. ¹H NMR (DMSO-d₆): δ 9.95 (1H, s), 8.38 (1H, dd, *J*= 4.8 Hz, *J*= 1.8 Hz), 8.33 (1H, d, *J*= 3.6 Hz), 7.84 (1H, d, *J*= 2.1 Hz), 7.79 (1H, ddd, *J*= 15.6 Hz, *J*= 7.2 Hz, *J*= 2.1 Hz), 7.57 (1H, d, *J*= 8.4 Hz), 7.19 (1H, dd, *J*= 8.4 Hz, *J*= 2.4 Hz), 7.01-6.95 (2H, m), 3.96 (2H, s), 1.32 (6H, s) ; purity 99.3% ; MS (m/e): 386 (MH⁺).

7.1.113 2-Chloro-N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine (R940342)

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-2,2-difluoro-4H-benzo[1,4]oxazin-3-one were reacted to produce 2-chloro-N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine **R940342**. ¹H NMR (DMSO-d₆): δ 12.24 (1H, s), 10.23 (1H, s), 8.45 (1H, dd, *J*= 3.3 Hz, *J*= 0.9 Hz), 7.66 (1H, dd, *J*= 4.2 Hz, *J*= 2.4 Hz), 7.55 (1H, dt, *J*= 9 Hz, *J*= 2.5 Hz), 7.43 (1H, d, *J*= 9 Hz); ¹⁹F NMR (DMSO-d₆): δ -21582, -43415 ; purity 96.2% ; MS (m/e) : 331 (MH⁺).

7.1.114 2-Chloro-N4-[(2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-7-yl]-5-fluoro-4-pyrimidineamine (R940344)

In like manner to the preparation of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one were reacted to produce 2-chloro-N4-[(2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-7-yl]-5-fluoro-4-pyrimidineamine **R940344**. ¹H NMR (DMSO-d₆): δ 11.32 (1H, s), 10.20 (1H, s), 8.45 (1H, d, *J* = 3.6 Hz), 8.33 (1H, d, *J* = 2.1 Hz), 7.84 (1H, d, *J* = 2.1 Hz), 1.54 (6H, s); purity 90.8%; MS (m/e): 324 (MH⁺).

7.1.115 N4-(4-Aminocarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (R945028)

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine (250 mg, 1.50 mmol) and 4-aminocarbonylmethyleneoxyaniline (540 mg, 3.25 mmol) were reacted to yield N4-(4-aminocarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine. LCMS: ret. time: 18.34 min.; purity: 100%; MS (m/e): 298.47 (MH⁺).

7.1.116 2-Chloro-5-fluoro-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-6-yl]-4-pyrimidineamine (R945298)

In a manner similar to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one were reacted to yield 2-chloro-5-fluoro-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-6-yl]-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 4.63 (s, 2H), 7.34 (d, *J* = 8.7 Hz, 1H), 7.44 (d, *J* = 8.4 Hz, 1H), 8.33 (d, *J* = 3.3 Hz, 1H), 10.14 (s, 1H, NH), 11.19 (s, 1H, NH); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 152.35; LCMS: ret. time: 26.74 min.; purity: 85.90%; MS (m/e): 296.13 (MH⁺).

7.1.117 N4-(1,4-Benzoxazin-6-yl)-N2-chloro-5-fluoropyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-1,4-benzoxazine were reacted to yield N4-(1,4-Benzoxazin-6-yl)-N2-chloro-5-fluoropyrimidineamine ¹H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.05 (m, 2H), 3.2 (m, 2H) purity 95 % MS (m/e): 281 (MH⁺).

7.1.118 N4-(1,4-Benzoxazin-7-yl)-N2-chloro-5-fluoropyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-1,4-benzoxazine were reacted to yield N4-(1,4-Benzoxazin-7-yl)-N2-chloro-5-fluoropyrimidineamine 1H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.05 (m, 2H), 3.2 (m, 2H) purity 94 % MS (m/e): 281 (MH⁺).

7.1.119 N4-(1,4-Benzoxazin-3-on-6-yl)-N2-chloro-5-fluoropyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-1,4-benzoxazine-3-one were reacted to yield N4-(1,4-Benzoxazin-3-on-6-yl)-N2-chloro-5-fluoropyrimidineamine 1H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.73 (s, 2H) purity 96 % MS (m/e): 295 (MH⁺).

7.1.120 N4-(1,4-Benzoxazin-3-on-7-yl)-N2-chloro-5-fluoropyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-1,4-benzoxazine-3-one were reacted to yield N4-(1,4-Benzoxazin-3-on-7-yl)-N2-chloro-5-fluoropyrimidineamine 1H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.79 (m, 1H), 6.6 (m, 1H), 4.68 (s, 2H) purity 93 % MS (m/e): 295 (MH⁺).

7.1.121 N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-6-yl)-pyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-4-N-methyl-1,4-benzoxazine were reacted to yield N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-6-yl)-pyrimidineamine 1H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.05 (m, 2H), 3.2 (m, 2H) 2.8 (s, 3H) purity 95 % MS (m/e): 295 (MH⁺).

7.1.122 N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-7-yl)-pyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-4-N-methyl-1,4-

benzoxazine were reacted to yield N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-7-yl)-pyrimidineamine 1H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.05 (m, 2H), 3.2 (m, 2H) 2.8 (s, 3H) purity 94 % MS (m/e): 295 (MH⁺).

7.1.123 N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-6-yl)-pyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-4-N-methyl-1,4-benzoxazine-3-one were reacted to yield N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-6-yl)-pyrimidineamine 1H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.73 (s, 2H) 2.8 (s, 3H) purity 96 % MS (m/e): 309 (MH⁺).

7.1.124 N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)-pyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-4-N-methyl-1,4-benzoxazine-3-one were reacted to N2-Chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)-pyrimidineamine 1H DMSO 8.2 (d, 1H), 6.8 (m, 1H), 6.75 (m, 1H), 6.60 (m, 1H), 4.68 (s, 2H) 2.8 (s, 3H) purity 93 % MS (m/e): 309 (MH⁺).

7.1.125 N2-chloro-N4-(3-ethylcarboxy-4H-imidazo[5,1-c]-1,4-benzoxazin-6-yl)-5-fluoropyrimidinediamine (R909258) :

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and ethyl 6-amino-3-carboxy-4H-imidazo[5,1-c]-1,4-benzoxazine were reacted to yield N2-chloro-N4-(3-ethylcarboxy-4H-imidazo[5,1-c]-1,4-benzoxazin-6-yl)-5-fluoropyrimidinediamine 1H (DMSO-d₆) 8.42 (s, 1H), 8.30 (m, 1H), , 8.05 (m, 1H), 7.43 (m, 1H), 5.53 (s, 2H), 4.25 (q, 2H J=6.5 Hz), 1.28 (t, 2H, J=6.5 Hz), purity 90 % MS (m/e): 390 (MH⁺).

7.1.126 N2-Chloro-N4-(3,3-dimethyl-1,4-benzoxazin-6-yl)-5-fluoropyrimidineamine

In like manner to 2-Chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine and 6-Amino-3,3-dimethyl-1,4-benzoxazine were reacted to yield N2-Chloro-N4-(3,3-dimethyl-1,4-benzoxazin-6-yl)-5-fluoro-pyrimidineamine 1H DMSO 8.18 (d, 1H), 6.8 (d, 1H), 6.67 (m, 2H), 3.76 (s, 2H), 1.05 (s, 6H) purity 99 % MS (m/e): 309 (MH⁺)

7.1.127 2-Chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-4-pyrimidineamine (R935241)

In like manner to the preparation of 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with 5-amino-1-(methoxycarbonyl)methyl-indazoline to produce 2-chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 10.04 (s, 1H), 8.28 (d, 1H, J = 3.5 Hz), 8.12 (s, 1H), 8.00 (dd, 1H, J = 1.2 and 4.1 Hz), 7.64 (d, 1H, J = 8.8 Hz), 7.58-7.54 (m, 1H), 5.39 (s, 2H), 3.66 (s, 3H).

7.1.128 2-Chloro-5-fluoro-N-[4*H*-imidazo[2,1-*c*][1,4]-benzoxazin-8-yl]-4-pyrimidineamine (R935257)

In like manner to the preparation of 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with 8-amino-4*H*-imidazo[2,1-*c*][1,4]-benzoxazine to produce 2-chloro-5-fluoro-N-[4*H*-imidazo[2,1-*c*][1,4]-benzoxazin-8-yl]-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 10.08 (s, 1H), 8.31 (s, 1H), 7.91 (d, 1H, J = 2.3 Hz), 7.74 (d, 1H, J = 1.2 Hz), 7.37 (dd, 1H, J = 2.3 and 8.8 Hz), 7.16 (d, 1H, J = 8.8 Hz), 7.14 (d, 1H, J = 1.2 Hz), 5.29 (s, 2H). LCMS: ret. time: 18.74 min.; purity: 99%; MS (*m/e*): 318 (MH⁺).

7.1.129 2-Chloro-5-fluoro-N-(indazoline-6-yl)-4-pyrimidineamine (R935260)

In like manner to the preparation of 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with 6-aminoindazole to produce 2-chloro-5-fluoro-N-(indazoline-6-yl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 13.03 (s, 1H), 10.07 (s, 1H), 8.32 (d, 1H, J = 3.5 Hz), 8.07 (s, 1H), 7.99 (s, 1H), 7.71 (d, 1H, J = 8.8 Hz), 7.34 (dd, 1H, J = 1.7 and 8.8 Hz). LCMS: ret. time: 18.52 min.; purity: 99%; MS (*m/e*): 263 (MH⁺).

7.1.130 2-Chloro-5-fluoro-N-(indazoline-5-yl)-4-pyrimidineamine (R935265)

In like manner to the preparation of 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with 5-aminoindazole. ¹H NMR (CDCl₃): δ 9.99 (s, 1H), 8.26 (d, 1H, J = 3.5 Hz), 8.07 (s, 1H), 7.99 (d, 1H, J = 1.1 Hz), 7.53 (dd, 2H, J = 1.7 and 8.8 Hz). LCMS: ret. time: 18.03 min.; purity: 97%; MS (*m/e*): 264 (MH⁺).

7.1.131 2-Chloro-5-fluoro-N-(1H-pyrrol-1-yl)-4-pyrimidineamine (R935275)

In like manner to the preparation of 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 2,4-dichloro-5-fluoropyrimidine was reacted with 1-aminopyrrole to produce 2-chloro-5-fluoro-N-(1H-pyrrol-1-yl)-4-pyrimidineamine. ¹H NMR (CDCl₃): δ 11.39 (s, 1H), 8.35 (d, 1H, J = 3.5 Hz), 6.83 (t, 2H, J = 2.3 Hz), 6.07 (t, 2H, J = 2.3 Hz). LCMS: ret. time: 18.95 min.; purity: 97%; MS (m/e): 213 (MH⁺).

7.1.132 2-Chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine (R926853)

A reaction mixture containing 2,4-dichloro-5-fluoropyrimidine (1.2 equivalents) and 3-(tetrazol-5-yl)aniline (1 equivalents) in methanol:water (1:1; v/v) was heated at 60 °C for 24 h. Upon dilution with water and acidification, the solid formed was filtered, washed with water, dried and analyzed to give 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine (R926853). Alternatively this reaction can be achieved by treating 2,4-dichloro-5-fluoropyrimidine (1 equivalent) with 3-(tetrazol-5-yl)aniline (3 equivalents) in methanol:water (1:1; v/v) at 60 °C for 2-3 hours or at room temperature for 24 h to give 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 10.25 (s, 1H), 8.43 (s, 1H), 8.37 (d, 1H, J = 3.6 Hz), 7.90 (dd, 1H, J = 0.9 and 9 Hz), 7.75 (d, 1H, J = 7.5 Hz), 7.61 (t, 1H, J = 7.8 Hz); LCMS: purity: 90%; MS (m/e): 292 (MH⁺).

7.1.133 2-Chloro-N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-2,4-pyrimidineamine (R950297)

A solution of 3,4-dihydro-4-hydroxy-6-amino-2H-1-benzopyran and 2,4-dichloro-5-fluoropyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-2,4-pyrimidineamine as a pale brown solid. LCMS: purity: 99.3%; MS (m/e): 296.1 (MH⁺).

7.1.134 2-Chloro-N4-(4-methoxycarbonyl ethyleneoxyphenyl)-5-fluoro-2,4-pyrimidineamine (R950375)

A solution of 3-(p-aminophenyl)-propionic acid and 2,4-dichloro-5-fluoropyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(4-

methoxycarbonylethyleneoxyphenyl)-5-fluoro-2,4-pyrimidineamine as a pale brown solid. LCMS: purity: 93.3%; MS (m/e): 311.98 (M⁺).

7.1.135 2-Chloro-N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidineamine (R950298)

A solution of 3-carboxy-4-hydroxyaniline and 2,4-dichloro-5-fluoro-pyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidineamine as a pale brown solid. LCMS: purity: 87.4%; MS (m/e): 284.1 (MH⁺).

7.1.136 2-Chloro-N4-(4-trifluoromethyl-3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidineamine (R950390)

A solution of 4-trifluoromethyl-3-methoxycarbonylaniline and 2,4-dichloro-5-fluoro-pyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(4-trifluoromethyl-3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidineamine as a pale brown solid. LCMS: purity: 96.4%; MS (m/e): 366.34 (MH⁺).

7.1.137 2-Chloro-N4-(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidineamine (R950369)

A solution of 3-methylcarbonylaniline and 2,4-dichloro-5-fluoro-pyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidineamine as a pale brown solid. LCMS: purity: 99.1%; MS (m/e): 266.12 (MH⁺).

7.1.138 2-Chloro-N4-(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidineamine (R950370)

A solution of 3-phenylcarbonylaniline and 2,4-dichloro-5-fluoro-pyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidineamine as a pale brown solid. LCMS: purity: 78.5%; MS (m/e): 328.16 (MH⁺).

7.1.139 2-Chloro-N4-(3-nitrophenyl)-5-fluoro-2,4-pyrimidineamine

A solution of 3-nitroaniline and 2,4-dichloro-5-fluoro-pyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(3-nitrophenyl)-5-fluoro-2,4-pyrimidineamine as a pale

brown solid. ^1H NMR (DMSO): δ 10.34 (s, 1H), 8.73 (d, 1H, $J = 2.4$ Hz), 7.66-8.29 (m, 4H).

7.1.140 2-Chloro-N4-(3-hydroxymethylen-4-methoxyphenyl)-5-fluoro-4-aminopyridine (R950384)

5 A solution of 3-hydroxymethylen-4-methoxyaniline and 2,4-dichloro-5-fluoropyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(3-hydroxymethylen-4-methoxyphenyl)-5-fluoro-4-aminopyridine as a pale brown solid. LCMS: purity: 91.8%; MS (m/e): 266.03 (MH⁺).

10 **7.1.141 2-Chloro-N4-(3-amino-4-ethoxyphenyl)-5-fluoro-4-aminopyridine (R950387)**

A solution of 3-amino-4-ethoxyaniline and 2,4-dichloro-5-fluoropyrimidine in MeOH was stirred for 2 hours at 70°C. The mixture was diluted with water and the resulting precipitate was filtered to give 2-chloro-N4-(3-amino-4-ethoxyphenyl)-5-fluoro-4-aminopyridine as a pale brown solid. LCMS: purity: 93.2%; MS (m/e): 252.06 (MH⁺).

7.2 Synthesis of Amines and Amine Precursors

7.2.1 5-Amino-2-(2-hydroxyethyleneoxy)pyridine

A methanolic solution (50 mL) of 2-(2-hydroxyethyleneoxy)-5-nitropyridine (0.5 g) was hydrogenated in the presence of Pd/C (10%; 0.05 g) using a balloon filled with hydrogen for 2h. After the filtration through a pad of celite and washing with methanol the solution was concentrated to give the 5-amino-2-(2-hydroxyethyloxy)pyridine. ^1H NMR (CDCl₃): δ 7.58 (d, 1H, $J = 3$ Hz), 7.05 (dd, 1H, $J = 2.7$ and 8.1 Hz), 6.64 (d, 1H, $J = 8.7$ Hz), 4.36 (m, 2H), 3.89 (m, 2H).

7.2.2 4-Chloro-3-methoxyaniline

25 In like manner to the preparation of 5-amino-2-(2-hydroxyethyleneoxy)pyridine, the hydrogenation of 4-chloro-3-methoxynitrobenzene gave 4-chloro-3-methoxyaniline. LCMS: purity: 98%; MS: 199 (M⁺ acetonitrile).

7.2.3 2-[5-Amino-2-oxo-1,3-benzoxazol-3(2H)-yl]acetamide

In like manner to the preparation of 5-amino-2-(2-hydroxyethyleneoxy)pyridine, the hydrogenation of 2-[1,3-benzoxazol-2-oxo-5-nitro-3(2H)-yl]acetamide gave 2-[5-amino-2-oxo-1,3-benzoxazol-3(2H)-yl]acetamide. LCMS: purity: 96%; MS: 208 (MH⁺).

5 7.2.4 7-nitro-1,2,3,4-tetrahydroisoquinoline

7-nitro-1,2,3,4-tetrahydroisoquinoline was prepared by nitration of 1,2,3,4-tetrahydroisoquinoline according to the following reference: Grunewald, Gary L.; Dahanukar, Vilas H.; Caldwell, Timothy M.; Criscione, Kevin R.; Journal of Medicinal Chemistry (1997), 40(25), 3997-4005.

10 7.2.5 2-(t-Butoxycarbonyl)-7-nitro-1,2,3,4-tetrahydroisoquinoline

A mixture of 7-nitro-1,2,3,4-tetrahydroisoquinoline (0.55g, 3.1 mmole), di-t-butylidicarbonate (0.70g, 3.2 mmole), triethylamine (1.0 mL, 7.7 mmole) in dichloromethane (8 mL) was stirred at rt for 8h. The reaction mixture was diluted with water (50 mL) and stirred for 1h. The organic phase was separated and washed with brine. Concentration of the organic phase gave 2-(t-butoxycarbonyl)-7-nitro-1,2,3,4-tetrahydroisoquinoline. ¹H NMR (CDCl₃): δ 8.03-7.95 (m, 2H), 7.28 (d, 1H, J= 8.4 Hz), 4.66 (s, 2H), 3.68 (t, 2H, J= 6.0 Hz), 2.92 (t, 2H, J= 6.0 Hz), 1.49 (s, 9H).

7.2.6 2,3-Dihydro-6-nitro-4-benzopyranon

3-(p-Nitrophenyl)-propionic acid is dissolved in concentrated sulfuric acid and treated with P₂O₅. The mixture is stirred for 1 hr at room temperature and poured onto ice. Filtration gave 2,3-dihydro-6-nitro-4-benzopyranon as a white solid. ¹H NMR (DMSO): δ 8.47 (d, J = 3.0 Hz, 1H), 8.35 (dd, J = 3.0, 9.0 Hz, 1H), 7.29 (d, J = 9.0 Hz, 1H), 4.70 (t, J = 7.2 Hz, 1H), 2.90 (t, J = 7.2 Hz, 1H).

7.2.7 3,4-Dihydro-4-hydroxy-6-amino-2H-1-benzopyran

A mixture 2,3-dihydro-6-nitro-4-benzopyranon and Pd/C (10%) in MeOH was hydrogenated at 22°C for 3 hours (40psi). The mixture was filtered and concentrated to dryness to give 3,4-dihydro-4-hydroxy-6-amino-2H-1-benzopyran as a brown oil. ¹H NMR (DMSO): δ 6.40-6.56 (m, 3H), 5.05 (bs, 1H), 4.45 (bs, 1H), 3.94-4.09 (m, 2H), 1.76-1.98 (m, 2H).

7.2.8 N4-(3,4-Ethylenedioxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R950287)

A solution of 2-Chloro-5-ethoxycarbonyl-N4-(3,4-ethylenedioxyphenyl)-2,4-pyrimidineamine in EtOH was treated with a 25% aqueous solution of NH₃. The mixture was stirred for 30 min at 100°C and purified by flash chromatography on silica gel to give N4-(3,4-ethylenedioxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine as a white solid. LCMS: purity: 92.3%; MS (m/e): 317.28 (MH⁺, 100).

7.2.9 3-(N-morpholinocarbonyl)aniline

To a 0°C solution of 3-nitrobenzoylchloride (0.50g, 2.7 mmole) and pyridine (0.27 mL, 3.2 mmole) in anhydrous dichloromethane (15 mL) was added morpholine (0.28 mL, 3.2 mmole). The reaction mixture was allowed to warm to rt and was stirred for 20h. The solvents were removed under vacuum and the residue suspended in ethyl acetate and washed with 1N HCl. The organic layer was washed with a saturated solution of NaHCO₃ and brine. Removal of the solvents under vacuum provided 1-(N-morpholinocarbonyl)-3-nitrobenzene which was used without further purification.

A mixture of 1-(N-morpholinocarbonyl)-3-nitrobenzene (0.64 g) and 10% Pd on activated carbon (60 mg) in degassed methanol (65 mL) was stirred under a balloon of H₂ for 2h. The reaction mixture was filtered through Celite® filter aid and then concentrated under reduced pressure to provide 3-(N-morpholinocarbonyl)aniline in quantitative yield. ¹H NMR (CDCl₃): δ 7.19-7.14 (m, 1H), 6.75-6.69 (m, 3H), 3.58-3.71 (m, 10H).

7.2.10 3-(N-propylcarbonyl)aniline

In like manner to the preparation of 3-(N-morpholinocarbonyl)aniline, 3-nitrobenzoylchloride and n-propylamine were reacted to prepare 1-[(N-propylamino)carbonyl]-3-nitrobenzene which underwent hydrogenation to provide 3-(N-propylcarbonyl)aniline. ¹H NMR (CDCl₃): δ 7.18 (t, 1H, J= 7.5 Hz), 7.13 (t, 1H, J= 1.8 Hz), 7.05-7.01 (m, 1H), 6.78 (ddd, 1H, J= 1.2, 2.4, and 7.5 Hz), 6.10 (bs, 1H), 3.58-3.53 (bs, 2H), 3.43-3.34 (m, 2H), 1.68-1.57 (m, 2H), 0.97 (t, 3H, J= 7.2 Hz).

7.2.11 3-[4-(Ethoxycarbonyl)piperidinocarbonyl]aniline

In like manner to the preparation of 3-(N-morpholinocarbonyl)aniline, 3-nitrobenzoylchloride and ethyl isonipecotate were reacted to prepare 1-[4-

(ethoxycarbonyl)piperidinocarbonyl]-3-nitrobenzene which underwent hydrogenation to provide 3-[4-(ethoxycarbonyl)piperidinocarbonyl]aniline.

7.2.12 3-(N-methylcarbonyl)aniline

In like manner to the preparation of 3-(N-morpholinocarbonyl)aniline, 3-nitrobenzoylchloride and methylamine hydrochloride were reacted to prepare 1-[(N-methylamino)carbonyl]-3-nitrobenzene which underwent hydrogenation to provide 3-(N-methylcarbonyl)aniline. ¹H NMR (CDCl₃): δ 7.18 (t, 1H, J= 7.5 Hz), 7.13 (t, 1H, J= 1.8 Hz), 7.04-6.99 (m, 1H), 6.81-6.75 (m, 1H), 6.05 (bs, 1H), 3.84 (bs, 2H), 2.99 (d, 3H, J= 4.8 Hz).

7.2.13 7-Amino-1-tetralone

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of 7-nitro-1-tetralone was carried out to prepare 7-amino-1-tetralone. ¹H NMR (CDCl₃): δ 7.32 (d, 1H, J= 2.4 Hz), 7.05 (d, 1H, J= 8.1 Hz), 6.82 (dd, 1H, J= 2.4 and 8.1 Hz), 2.85 (t, 2H, J= 6.6 Hz), 2.61 (t, 2H, J= 6.6 Hz), 2.14-2.04 (m, 2H).

7.2.14 7-Amino-2-(t-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinoline

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of 2-(t-butoxycarbonyl)-7-nitro-1,2,3,4-tetrahydroisoquinoline was carried out to prepare 7-amino-2-(t-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinoline. ¹H NMR (CDCl₃): δ 6.92 (d, 1H, J= 8.4 Hz), 6.52 (dd, 1H, J= 2.4 and 8.4 Hz), 6.44 (bs, 1H), 4.47 (s, 2H), 3.63-3.48 (m, 2H), 2.71 (t, 2H, J= 5.1 Hz), 1.45 (s, 9H).

7.2.15 7-Amino-1,2,3,4-tetrahydroisoquinoline

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of 7-nitro-1,2,3,4-tetrahydroisoquinoline was carried out to prepare 7-amino-1,2,3,4-tetrahydroisoquinoline. ¹H NMR (DMSO-*d*₆): δ 9.35 (bs, 1H), 6.82 (d, 1H, J= 8.1 Hz), 6.45 (dd, 1H, J= 2.4 and 8.4 Hz), 6.30 (d, 1H, J= 2.4 Hz), 5.05 (s, 2H), 4.05 (s, 2H), 3.24 (t, 2H, J= 6.6 Hz), 2.78 (t, 2H, J= 6.6 Hz).

7.2.16 2-(3-aminophenoxy)-N,2-dimethylpropanamide

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of N,2-dimethyl-2-(3-nitrophenoxy)propanamide was carried out to prepare 2-(3-

aminophenoxy)-N,2-dimethylpropanamide. ¹H NMR (CDCl₃): δ 7.03 (t, 1H, J= 7.8 Hz), 6.71 (bs, 1H), 6.39 (dd, 1H, J= 1.2 and 6.9 Hz), 6.29 (dd, 1H, J= 2.4 and 9.6 Hz), 6.25-6.22 (m, 1H), 2.86 (d, 3H, J= 4.2 Hz), 2.86 (d, 3H, J= 4.2 Hz), 1.50 (s, 6H).

7.2.17 Ethyl 2-(3-aminophenoxy)-2-methylpropanate

5 In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of ethyl 2-methyl-2-(3-nitrophenoxy)propanate was carried out to prepare ethyl 2-(3-aminophenoxy)-2-methylpropanate. ¹H NMR (CDCl₃): δ 6.99 (t, 2H, J= 8.7 Hz), 6.32 (dt, 1H, J= 1.2 and 7.2 Hz), 6.24-6.18 (m, 2H), 4.23 (q, 2H, J= 7.2 Hz), 1.58 (s, 6H), 1.24 (t, 3H, J= 6.9 Hz).

10 7.2.18 N-methyl-2-(5-amino-2-methylphenoxy)acetamide

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of N-methyl-2-(2-methyl-5-nitrophenoxy)acetamide was carried out to prepare N-methyl-2-(5-amino-2-methylphenoxy)acetamide. ¹H NMR (CD₃OD): δ 6.86 (d, 1H, J= 7.5 Hz), 6.32-6.25 (m, 2H), 4.43 (s, 2H), 2.82 (s, 3H), 2.14 (s, 3H).


15 7.2.19 6-Amino-2-(methoxycarbonyl)-(1H)-indole

6-Amino-2-(methoxycarbonyl)-(1H)-indole was prepared according to the following references:

1. Adams, Richard E.; Press, Jeffery B.; Deegan, Edward G.; Synthetic Communications (1991), 12 (5), 675-681.

20 2. Boger, Dale L.; Yun, Weiya; Han, Nianhe; Johnson, Douglas S.; Bioorganic & Medicinal Chemistry (1995), 3(6), 611-621

7.2.20 Preparation of 3-hydroxy-5-(methoxycarbonylmethyleneoxy)aniline and 3,5-bis(methoxycarbonylmethyleneoxy)aniline



Benzyl N-(3,5-dihydroxyphenyl)carbamate

To a mixture of 5-aminobenzene-1,3-diol (0.60 g, 3.7 mmole) and sodium hydrogencarbonate (1.4 g, 16 mmole) in THF/water (15 mL, 1:1 v/v) was added dropwise benzyl chloroformate 1.6 mL, 11 mmole). After 3h at rt, THF was removed under vacuum and the remaining aqueous layer was extracted with ethyl acetate. Purification by column chromatography over silica gel provided benzyl N-(3,5-dihydroxyphenyl)carbamate. ¹H NMR (CD₃OD): δ 7.42-7.25 (m, 5H), 6.46 (d, 2H, J = 2.4 Hz), 5.97-5.94 (m, 1H), 5.14 (s, 2H).

10 Benzyl N-[3-hydroxy-5-(methoxycarbonylmethylenoxy) phenyl]carbamate and Benzyl N-[3,5-bis(methoxycarbonylmethylenoxy)phenyl]carbamate

In like manner to the preparation of ethyl 4-nitrophenoxyacetate, benzyl N-(3,5-dihydroxyphenyl)carbamate and methyl bromoacetate were reacted to give a mixture of benzyl N-[3-hydroxy-5-(methoxycarbonylmethylenoxy)phenyl]carbamate ¹H NMR (DMSO-*d*₆): δ 9.62 (s, 1H), 9.44 (s, 1H), 7.42-7.31 (m, 5H), 6.63 (s, 1H), 6.50 (t, 1H, J = 2.4 Hz), 5.93 (t, 1H, J = 2.4 Hz), 5.10 (s, 2H), 4.63 (s, 2H), 3.67 (s, 3H), and benzyl N-[3,5-bis(methoxycarbonylmethylenoxy)phenyl]carbamate ¹H NMR (CDCl₃): δ 7.38-7.32 (m, 5H), 6.86 (s, 1H), 6.67 (d, 2H, J = 1.8 Hz), 6.19 (t, 1H, J = 2.4 Hz), 5.16 (s, 2H), 4.57 (s, 4H), 3.78 (s, 6H).

20 which were separated by column chromatography over silica gel.

3-Hydroxy-5-(methoxycarbonylmethyleneoxy)aniline

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of benzyl N-[3-hydroxy-5-(methoxycarbonylmethyleneoxy)phenyl]carbamate was carried out to prepare 3-hydroxy-5-(methoxycarbonylmethyleneoxy)aniline. ¹H NMR (CD₃OD): δ

5 5.87-5.80 (m, 2H), 5.78-5.72 (m, 1H), 4.56 (s, 2H), 3.76 (s, 3H).

3,5-Bis(methoxycarbonylmethyleneoxy)aniline

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of benzyl N-[3,5-bis(methoxycarbonylmethyleneoxy)phenyl]carbamate was carried out to prepare 3,5-bis(methoxycarbonylmethyleneoxy)aniline. ¹H NMR (CD₃OD): δ 5.92 (d, 2H,

10 J= 2.4 Hz), 5.83 (t, 1H, J= 2.4 Hz), 4.58 (s, 4H), 3.78 (s, 6H).

7.2.21 N4-(3,4-Ethylenedioxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R950287)

A solution of 2-Chloro-5-ethoxycarbonyl-N4-(3,4-ethylenedioxyphenyl)-2,4-pyrimidineamine in EtOH was treated with a 25% aqueous solution of NH₃. The mixture
15 was stirred for 30 min at 100°C and purified by flash chromatography on silica gel to give N4-(3,4-ethylenedioxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine as a white solid. LCMS: purity: 92.3%; MS (m/e): 317.28 (MH⁺, 100).

7.2.22 Ethyl 6-Nitro-3-carboxy-4H-imidazo[5,1-c]-1,4-benzoxazine

Was prepared according to J. of Heterocyclic Chemistry, 26, 205, (1989)

7.2.23 Ethyl 6-Amino-3-carboxy-4H-imidazo[5,1-c]-1,4-benzoxazine

Ethyl 6-Nitro-3-carboxy-4H-imidazo[5,1-c]-1,4-benzoxazine was reduced shaken in MeOH under 40 p.s.i. H₂ with 20 weight percent of 10% Pd/C (Degussa) for 1 h then filtered and the solvent evaporated. The compound was purified directly by column chromatograph (EtOAc/hexane) to yield Ethyl 6-Amino-3-carboxy-4H-imidazo[5,1-c]-1,4-
25 benzoxazine 1H (DMSO-d₆) 8.41 (s, 1H), 6.98 (m, 1H), 6.82 (m, 1H), 6.43 (m, 1H), 5.28 ((s, 2H), 4.23 (q, 2H, J=6.2 Hz), 1.27 (t, 2H, J=6.2 Hz) purity 92 % MS (m/e): 232 (MH⁺).

7.2.24 6-Amino-3,3-dimethyl-1,4-benzoxazine

A mixture of 15 g 2-Amino-4-nitrophenol and 40 g Boc₂O in 300 mL CHCl₃ was refluxed overnight filtered and the filtrate was evaporated to near dryness. The residue was

trituated with hexanes, collected by suction filtration, and dried to yield 2-N-Boc-amino-4-nitrophenol. The 2-N-Boc-amino-4-nitrophenol was refluxed in acetone with 15.6 mL of 1-Chloro-2-methylpropene and 25 g potassium carbonate overnight. The reaction mixture was poured into ice-slush, the solid was collected by suction filtration and washed with water. The solid was dissolved in EtOAc and the organic was washed with 10% NaOH solution, water, then brine and dried over MgSO₄. The organic was filtered to remove the drying agent and evaporated to yield 18 g 1-(2-N-Boc-amino-4-nitrophenoxy)-2-methyl-2-propene. 7.8 g of 1-(2-N-Boc-amino-4-nitrophenoxy)-2-methyl-2-propene was stirred overnight in methanolic HCl in a round-bottom flask with a septum wired on, and then heated with a reflux condenser attached at 80° C for 10 minutes. The reaction was cooled and the methanol was removed by rotary-evaporation. The residue was dissolved in 30 mL of 4N HCl, transferred to a new vessel to leave behind any undissolved solids and cooled to 0° C. 1.83 g of NaNO₂ in 5 mL water was added drop wise and the solution was neutralized with solid sodium bicarbonate. A solution of 1.64 g NaN₃ in 17 mL water was added slowly drop wise and the reaction was stirred 30 minutes. The precipitate was collected by suction filtration, washed well with water and dried on the funnel to yield 5.7 g 1-(2-Azido-4-nitrophenoxy)-2-methyl-2-propene. 7 g of 1-(2-Azido-4-nitrophenoxy)-2-methyl-2-propene was refluxed in 300 mL benzene overnight, cooled then evaporated. The crude product was recrystallized from EtOAc/Hexanes to yield 3-Methyl-6-nitro-azirino[2,1-*c*]-1,4-benzoxazine in two crops with a combined mass of 5.1 g of 3-Methyl-6-nitro-azirino[2,1-*c*]-1,4-benzoxazine was dissolved in 500 mL of MeOH/5% THF, 200 mg of 10% Pd/C (Degussa) was added and the resulting mixture was shaken under 30 p.s.i. H₂ atmosphere for 8 hours. The reaction mixture was filtered through a pad of celite and the solvent evaporated. The residue was dissolved in a minimum amount of DCM/THF/MeOH and loaded onto a 5 cm by 20 cm 3% MeOH/DCM SiO₂ column and the compound was eluted isocratically with a small amount of positive pressure. The appropriate fractions were combined and evaporated to yield 590 mg of 6-Amino-3,3-dimethyl-1,4-benzoxazine. ¹H (DMSO-d₆) 6.30 (d, 1H), 5.75 (d, 1H), 5.65 (dd, 1H), 3.58 (s, 2H), 1.08 (s, 6H) purity 99 % MS (m/e): 179 (MH⁺).

7.2.25 Ethyl 4-Aminophenoxyacetate

Ethyl 4-Nitrophenoxyacetate

A dry reaction flask equipped with a reflux condenser, N₂ inlet and a magnetic stirring bar was charged with 3-nitrophenol (76.45 g, 550 mmol), K₂CO₃ (76.45 g, 550 mmol) and dry acetone (500 mL) under N₂ atmosphere. To this at room temperature was added ethyl bromoacetate (55.44 mL, 500 mmol) over a period of 15 min. The reaction mixture was refluxed for 16h, cooled and poured over ice-water (4 Kg). The resulting aqueous solution was extracted with CH₂Cl₂ (3 x 500 mL), dried over anhydrous Na₂SO₄ and solvent was removed to obtain 103g (92%) of the desired ethyl 4-nitrophenoxyacetate. ¹H NMR (CDCl₃): δ 8.20 (d, 2H, J= 8.2 Hz), 6.95 (d, 2H, J= 8.1 Hz), 4.72 (s, 2H), 4.25 (q, 2H), 1.23 (t, 3H); LCMS: ret. time: 27.07 min.; purity: 100%; MS: 267 (M+ acetonitrile).

Ethyl 4-Aminophenoxyacetate

A solution of ethyl 4-nitrophenoxyacetate (15 g) in EtOH (400 mL) was hydrogenated at 40 PSI for 40 minutes in the presence of 10% Pd/C (1.5 g, 10% by weight). After the filtration through a celite the solvent was removed under a reduced pressure to obtain ethyl 4-aminophenoxyacetate. ¹H NMR (CDCl₃): δ 6.77 (d, 2H, 8.1 Hz), 6.60 (d, 2H, J= 8.0 Hz), 4.50 (s, 2H), 4.24 (q, 2H), 1.24 (t, 3H); LCMS: ret. time: 12.00 min.; purity: 100%; MS (m/e): 196 (MH⁺).

7.2.26 tert-Butyl 4-Aminophenoxyacetate

tert-Butyl 4-Nitrophenoxyacetate

In like manner to the preparation of ethyl 4-nitrophenoxyacetate, 4-nitrophenol and tert-butyl bromoacetate were reacted to prepare tert-butyl 4-nitrophenoxyacetate. ¹H NMR (CDCl₃): δ 8.2 (d, 2H, J= 8.1 Hz), 6.95 (d, 2H, J= 8.2 Hz), 4.60 (s, 2H), 1.42 (s, 9H).

tert-Butyl 4-Aminophenoxyacetate

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of tert-butyl 4-nitrophenoxyacetate was carried out to prepare tert-butyl 4-aminophenoxyacetate. ¹H NMR (CDCl₃): δ 6.74 (d, 2H, J= 9 Hz), 6.62 (d, 2H, J= 9 Hz), 4.42 (s, 2H), 1.42 (s, 9H); LCMS: ret. time: 16.35 min.; purity: 94%; MS (m/e): 224 (MH⁺).

7.2.27 Ethyl 3-Aminophenoxyacetate

Ethyl 3-Nitrophenoxyacetate

In like manner to the preparation of ethyl 4-nitrophenoxyacetate, 3-nitrophenol and ethyl bromoacetate were reacted to prepare ethyl 3-nitrophenoxyacetate. ¹H NMR (CDCl₃):
5 δ 7.88 (dt, 1H, J= 1.2 and 8.7 Hz), 7.71 (t, 1H, J= 2.4 Hz), 7.45 (t, 1H, J= 8.4 Hz), 7.27 (dt, 1H, J= 2.4 and 8.4 Hz), 4.70 (s, 2H), 4.29 (q, 2H, J= 6.9 Hz), 1.30 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 27.28 min.; purity: 96%.

Ethyl 3-Aminophenoxyacetate

10 In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of ethyl 3-nitrophenoxyacetate was carried out to prepare ethyl 3-aminophenoxyacetate. ¹H NMR (CDCl₃): δ 7.05 (t, 1H, J= 7.2 Hz), 6.30 (m, 3H), 4.56 (s, 2H), 4.25 (q, 2H, J= 7.2 Hz), 1.29 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 10.69 min.; purity: 96%; MS (m/e): 196 (MH⁺).

15 7.2.28 (+)-Ethyl 2-(4-Aminophenoxy)propionate

In like manner to the preparation of ethyl 4-aminophenoxyacetate, hydrogenation of ethyl (+)-2-(4-nitrophenoxy)propionate was carried out to prepare (+) ethyl 2-(4-aminophenoxy)propionate. ¹H NMR (CDCl₃): δ 6.70 (d, 2H), 6.58 (d, 2H), 4.60 (m, 1H), 4.20 (q, 2H), 3.2 (bs, 2H), 1.45 (d, 3H), 1.22 (t, 3H).

20 7.2.29 N-Methyl 3-Aminophenoxyacetamide

N-Methyl 3-Nitrophenoxyacetamide

A mixture of ethyl 3-nitrophenoxyacetate (9.12g, 40 mmol), methylamine hydrochloride (26.8g, 400 mmol) and diisopropylethylamine (35.5 mL, 200 mL) in MeOH (100 mL) was stirred in a pressure vial at 90 °C for 6h. The reaction was cooled to room
25 temperature, diluted with water (1 liter), the solid formed was filtered, washed with water and dried to get the desired N-methyl 3-nitrophenoxyacetamide (8g, 95%). ¹H NMR CDCl₃): δ 7.91 (dd, 1H, J= 1.8 and 8.1 Hz), 7.78 (t, 1H, J= 2.4 Hz), 7.50 (t, 1H, J= 8.7 Hz), 7.29 (dd, 1H, J= 1.8 and 8.4 Hz), 6.50 (bs, 2H), 4.57 (s, 2H), 2.95 and 2.93 (2s, 3H); LCMS: ret. time: 17.54 min.; purity: 100%; MS (m/e): 211 (MH⁺).

N-Methyl 3-Aminophenoxyacetamide

In like manner to the preparation of ethyl 4-aminophenoxyacetate, the hydrogenation of N-methyl 3-nitrophenoxyacetamide (8 g, 39 mmol) was conducted to give the desired N-methyl 3-aminophenoxyacetamide (6g, 86%). ¹H NMR (CD₃OD): δ 6.99 (t, 1H, J= 8.1 Hz), 6.37-6.25 (m, 3H), 4.41 (s, 2H), 2.80 (s, 3H); LCMS: ret. time: 19.80 min.; purity: 100%.

7.2.30 2-Methoxycarbonyl-5-aminobenzofuran (R926610)**2-Methoxycarbonyl-5-nitrobenzofuran (R926609)**

To a suspension of 5-nitro-2-benzofurancarboxylic acid (5 g, 24.15 mmol) in CH₂Cl₂ (250 mL) at 0 °C was added DMF (0.100 mL) followed by (COCl)₂ (2M in CH₂Cl₂, 36.23 mL, 72.46 mL) over a period of 10 min. The reaction was stirred at 0 °C for 1h and then at room temperature for 30 min. The reaction solvent was removed under a reduced pressure, dried under high vacuum and again suspended in CH₂Cl₂ (250 mL). The solution was cooled to 0 °C, were added pyridine (4.8 mL, 48.03 mmol) followed by MeOH (10 mL, excess) and stirred overnight. The extractive work-up with CH₂Cl₂ gave the expected 2-methoxycarbonyl-5-nitrobenzofuran (R926609). ¹H NMR (CDCl₃): δ 8.66 (d, 1H, J= 2.4 Hz), 8.36 (dd, 1H, J= 2.4 and 9.6 Hz), 7.71 (d, 1H, J= 9.3 Hz), 7.65 (s, 1H), 4.01 (s, 3H); LCMS: ret. time: 26.94 min.

2-Methoxycarbonyl-5-aminobenzofuran (R926610)

In like manner to the preparation of ethyl 4-aminophenoxyacetate, the hydrogenation of 2-methoxycarbonyl-5-nitrobenzofuran (2 g) in MeOH gave 2-methoxycarbonyl-5-aminobenzofuran. ¹H NMR (CDCl₃): δ 7.38 (bt, 2H), 6.90 (bd, 1H), 6.85 (bdd, 1H), 3.98 (s, 3H).

7.2.31 Methyl 2-(2-methyl-5-nitrophenoxy)acetate

In like manner to the preparation of ethyl 4-nitrophenoxyacetate, 2-methyl-5-nitrophenol and methyl bromoacetate were reacted to prepare methyl 2-(2-methyl-5-nitrophenoxy)acetate. ¹H NMR (CD₃OD): δ 7.80 (dd, 1H, J= 2.4 and 8.1 Hz), 7.65 (d, 1H, J= 2.4 Hz), 7.38 (d, 1H, J= 8.1 Hz), 4.90 (s, 2H), 3.80 (s, 3H), 2.36 (s, 3H).

7.2.32 Ethyl 2-methyl-2-(3-nitrophenoxy)propanate

A mixture of 3-nitrophenol (0.50g, 3.6 mmole), ethyl bromodimethylacetate (0.64g, 3.3 mmole), K₂CO₃ (1.3 g, 9.4 mmole), potassium iodide (catalytic) in absolute ethanol (8 mL) was heated at 70°C for 18h. The reaction mixture was cooled, poured into a saturated solution of NaHCO₃, and extracted with dichloromethane. The product, ethyl 2-methyl-2-(3-nitrophenoxy)propanate, was obtained after purification by column chromatography over silica gel. ¹H NMR (CDCl₃): δ 7.85 (dt, 1H, J= 1.2 and 8.1 Hz), 7.68 (t, 1H, J= 2.4 Hz), 7.40 (t, 1H, J= 8.4 Hz), 7.19-7.13 (m, 1H), 4.26 (q, 2H, J= 7.2 Hz), 1.64 (s, 6H), 1.26 (t, 3H, J= 7.21),

7.2.33 N-Methyl-2-(2-methyl-5-nitrophenoxy)acetamide

In like manner to the preparation of N-methyl 3-nitrophenoxyacetamide, methyl 2-methyl-5-nitrophenoxyacetate and methylamine hydrochloride were reacted to prepare N-methyl-2-(2-methyl-5-nitrophenoxy)acetamide. ¹H NMR (CD₃OD): δ 7.82 (dd, 1H, J= 2.4 and 8.1 Hz), 7.69 (d, 1H, J= 2.4 Hz), 7.40 (d, 1H, J= 8.1 Hz), 4.66 (s, 2H), 2.83 (s, 3H), 2.40 (s, 3H).

7.2.34 N,2-Dimethyl-2-(3-nitrophenoxy)propanamide

In like manner to the preparation of ethyl 2-methyl-2-(3-nitrophenoxy)propanate, 3-nitrophenol and N,2-dimethyl-2-bromopropanamide (prepared according to the following reference: Guziec, Frank S., Jr.; Torres, Felix F. Journal of Organic Chemistry (1993), 58(6), 1604-6) were reacted to prepare N,2-dimethyl-2-(3-nitrophenoxy)propanamide. ¹H NMR (CDCl₃): δ 7.94 (dt, 1H, J= 1.2 and 8.1 Hz), 7.78 (t, 1H, J= 2.4 Hz), 7.45 (t, 1H, J= 8.4 Hz), 7.22 (ddd, 1H, J= 1.2, 2.4, and 8.1 Hz), 6.61 (bs, 1H), 2.89 (d, 3H, J= 5.1 Hz), 1.55 (s, 6H).

7.2.35 4-Amino-[(1H,1,2,3,4-tetrazolyl)methyleneoxy]benzene**4-Nitro-[(1H,1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene**

A mixture of 2-cyanomethoxy-4-nitrophenyl (5.8 g, 32.6 mmol), sodium azide (6.3 g, 98.0 mmol) and ammonium chloride (8.5 g, 163.3 mmol) was suspended in DMF (100 mL) containing acetic acid (1 mL) and the mixture heated at 70 °C. After 17 h, the reaction was cooled to room temperature and 2 N aqueous hydrochloric acid (100 mL) was added.

The solid which precipitated out of the reaction mixture was collected by filtration, washed with water (2 x 20 mL) then hexane (30 mL), affording compound 4-nitro-[(1H,1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene (6.7 g, 99%) as an orange solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.25 (d, *J* = 9.2 Hz, 2H), 7.29 (d, *J* = 9.1 Hz, 2H), 5.68 (s, 2H); ESI MS *m/z* 220 [C₈H₇N₅O₃ - H]⁻.

4-Amino-[(1H,1,2,3,4-tetrazolyl)methyleneoxy]benzene

A mixture of 4-nitro-[(1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene (6.7 g, 30.4 mmol) and 5 wt % palladium on carbon (700 mg) suspended in ethanol/concentrated hydrochloric acid (14:1, 150 mL) was hydrogenated in a sealed vessel at 50 psi. The mixture was shaken until no further hydrogen uptake was observed, after which the reaction was filtered through diatomaceous earth with chloroform and the filtrate concentrated to afford crude product. Purification by flash chromatography (7:2.5:0.5 CHCl₃/CH₃OH/NH₄OH) afforded 4-amino-[(1H,1,2,3,4-tetrazolyl)methyleneoxy]benzene as a brown solid: ¹H NMR (300 MHz, DMSO-*d*₆) δ 6.76 (d, *J* = 8.7 Hz, 2H), 6.52 (d, *J* = 8.7 Hz, 2H), 5.07 (s, 2H); ESI MS *m/z* 190 [C₈H₉N₅O - H]⁻.

7.2.36 4-Amino-[(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene

4-Nitro-[(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene and 4-Nitro-[(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene

A mixture of 4-nitro-[(1H,1,2,3,4-tetrazolyl)methyleneoxy]benzene (10.00 g, 45.2 mmol), cesium carbonate (22.09 g, 67.8 mmol) and methyl iodide (7.70 g, 54.3 mmol) in DMF (200 mL) was stirred at room temperature for 24 h. The reaction mixture was concentrated to remove most of the DMF and the crude residue was partitioned between chloroform (100 mL) and water (50 mL). The organic phase was separated, washed with brine, dried (Na₂SO₄) and concentrated to afford crude product as a orange solid. Purification by flash chromatography (chloroform) afforded 4-nitro-[(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene: ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.26 (d, *J* = 9.2 Hz, 2H), 7.31 (d, *J* = 9.2 Hz, 2H), 5.72 (s, 2H), 4.15 (s, 3H); and 4-nitro-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene: ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.24 (d, *J* = 9.3 Hz, 2H), 7.29 (d, *J* = 9.3 Hz, 2H), 5.58 (s, 2H), 4.41 (s, 3H).

4-Amino-[(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene

A mixture of 4-nitro-[(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene (3.60 g, 15.3 mmol) and 5% Pd/C (0.40 g) in 14:1 ethanol/concentrated hydrochloric acid (75 mL) was shaken at room temperature in a atmosphere of hydrogen at 50 psi. After 4 h no further hydrogen uptake was observed. The reaction mixture was filtered through diatomaceous earth, the solids washed with a 6:3:1 chloroform/methanol/concentrated ammonium hydroxide solution and the filtrate concentrated to afford crude 4-amino-[(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene, which was purified by flash chromatography (95:5 chloroform/ methanol): ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.48 (br s, 2H), 6.79 (d, *J* = 6.9 Hz, 2H), 6.55 (d, *J* = 6.9 Hz, 2H), 5.36 (s, 2H), 4.10 (s, 3H).


7.2.37 4-Amino-[(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene

A mixture of 4-nitro-[(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene (3.60 g, 15.3 mmol) and 5% Pd/C (0.40 g) in 14:1 ethanol/concentrated hydrochloric acid (75 mL) was shaken at room temperature in a hydrogen atmosphere at 50 psi. After 3 h no further hydrogen uptake was observed. The reaction mixture was filtered through diatomaceous earth, the solids washed with a 6:3:1 chloroform/methanol/concentrated ammonium hydroxide solution and the filtrate concentrated to afford crude 4-amino-[(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene, which was purified by flash chromatography (95:5 chloroform/ methanol): ¹H NMR (300 MHz, DMSO-*d*₆) δ 6.80 (br s, 2H), 6.75 (d, *J* = 9.0 Hz, 2H), 6.50 (d, *J* = 9.0 Hz, 2H), 5.17 (s, 2H), 4.37 (s, 3H).

7.2.38 2-Ethoxycarbonyl-5-aminoindole (R926611)

In like manner to the preparation of ethyl 4-aminophenoxyacetate, the hydrogenation of 2-ethoxycarbonyl-5-nitroindole gave the 2-ethoxycarbonyl-5-aminoindol. LCMS: ret. time: 13.44 min.; purity: 93%; MS (m/e): 205 (MH⁺).

7.2.39 5-[(4-Aminophenoxy)methyl]-3-phenyl-1,2,4-oxadiazole

**Preparation of 5-[4-(Nitrophenoxy)methyl]-3-phenyl-1,2,4-oxadiazole**

4-Nitrophenol (0.36 g, 2.56 mmole), 5-(chloromethyl)-3-phenyl-1,2,4-oxadiazole (0.5 g, 2.56 mmole) and anhydrous K_2CO_3 (0.39 g, 2.82 mmole) were dissolved in anhydrous acetone (20 mL) and heated to reflux for 12 h. Reaction mixture was cooled and the solvent removed under vacuum. The crude solid formed was collected by filtration, washed with water and dried under vacuum to provide 5-[(4-nitrophenoxy)methyl]-3-phenyl-1,2,4-oxadiazole (0.70 g, 92%). ^1H NMR ($CDCl_3$): δ 8.25 (d, 2H, $J = 8.8$ Hz), 8.08 (dd, 2H, $J = 8.2$ Hz), 7.52-7.49 (m, 3H), 7.13 (d, 2H, $J = 8.8$ Hz), 5.45 (s, 2H).

Preparation of 5-[(4-Aminophenoxy)methyl]-3-phenyl-1,2,4-oxadiazole


The 5-[(4-nitrophenoxy)methyl]-3-phenyl-1,2,4-oxadiazole (0.5 g, 1.68 mmole) was dissolved in methanol:methylenechloride (1:1) (120 mL). Aqueous solution of (15 mL) sodium hydrosulfite (0.88g, 5.05 mmole) and K_2CO_3 (0.70g, 5.06 mmole) was added dropwise under nitrogen for 10 min. The contents were allowed to stir at room temperature. After consumption of starting material, reaction mixture was concentrated, diluted with water till the homogeneous layer formed. The aqueous layer was extracted with several times with ethylacetate and methylene chloride. The turbid organic layers were combined, dried with anhydrous Na_2SO_4 and concentrated. Purification of the solid concentrate by silica gel chromatography provided 5-[(4-aminophenoxy)methyl]-3-phenyl-1,2,4-oxadiazole (0.23g, 51%). ^1H NMR ($CDCl_3$): δ 8.11 (m, 2H), 7.52-7.46 (m, 3H), 6.87 (d, 2H, $J = 8.8$ Hz), 6.64 (d, 2H, $J = 8.8$ Hz), 5.26 (s, 2H), 3.49 (br s, 2H).

Preparation of 5-[(4-Nitrophenoxy)methyl]-3-methyl-1,2,4-oxadiazole

A mixture of 4-nitrophenoxy acetic acid (2.25 g, 11.4 mmole), acetamidoxime, triethylamine hydrochloride (3.85g, 27.62 mmole), EDCl.HCl (4.37g, 22.79 mmole) and diisopropylethylamine (7.42g, 57.40 mmole) in anhydrous THF (250 ml) was refluxed for 18h. The unhomogenous brown colored reaction mixture was quenched with water and extracted with EtOAc (3 x 300 mL). The combined organic layers washed successively with aqueous NaHCO₃, brine and dried over anhydrous Na₂SO₄. Removal of solvent and purified by chromatographic purification provided 5-[(4-nitrophenoxy)methyl]-3-methyl-1,2,4-oxadiazole (1.62 g, 60 %). ¹H NMR (CDCl₃): δ 8.24 (d, 2H, J= 8.8 Hz), 7.08 (d, 2H, J= 8.8 Hz), 5.36 (s, 2H), 2.44 (s, 3H).

Preparation of 5-[(4-Aminophenoxy)methyl]-3-methyl-1,2,4-oxadiazole

In like manner to the preparation of 5-[(4-aminophenoxy)methyl]-3-phenyl-1,2,4-oxadiazole, 5-(4-nitrophenoxy)methyl-3-methyl-1,2,4-oxadiazole was reacted with aqueous solution of sodium hydrosulfite and K₂CO₃ to prepare 5-[(4-aminophenoxy)methyl]-3-methyl-1,2,4-oxadiazole. ¹H NMR (CDCl₃): δ 6.82 (d, 2H, J= 8.8 Hz), 6.63 (d, 2H, J= 8.8 Hz), 5.15 (s, 2H), 3.38 (br s, 2H), 2.41 (s, 3H).

7.2.40 Ethyl 2-(4-aminophenyl)-2-methylpropionate**Ethyl 2-methyl-2-(4-nitrophenyl)propionate**


A dry reaction flask charged with ethyl 4-nitrophenylacetate (5.0 g, 23.89 mmole), iodomethane (8.48 g, 3.72 mL, 59.74 mmole), 18-crown-6 (1.57 g, 5.93 mmole) in dry THF (200 mL) was cooled to -78 °C under nitrogen atmosphere. While stirring the contents, *t*-BuOK (5.90 g, 52.57 mmole) was added portionwise. The resulting violet precipitate was stirred at -78 °C for 2h and allowed the contents to warm to room temperature. The reaction was stirred at room temperature for 6h. At this time, once again the contents were cooled to -78 °C another portion of iodomethane, *t*-BuOK, and 18-crown-6 were added successively and stirred at the same temperature for 2h. The reaction was allowed to warm

to room temperature and stirred overnight. The reaction was quenched with saturated aq. NH_4Cl (75 mL), the resulting homogenous mixture extracted with ether (4 x 200 mL), dried over anhydrous Na_2SO_4 , and concentrated. The concentrate was purified by silica gel column chromatography with 1% EtOAc/hexanes to provide ethyl 2-methyl-2-(4-nitrophenyl)propionate as a pale yellow oil (2.38, 42%). ^1H NMR (CDCl_3): δ 8.17 (d, 2H, $J = 8.8$ Hz), 7.49 (d, 2H, $J = 8.8$ Hz), 4.12 (qt, 2H, $J = 7.0$ Hz), 1.60 (s, 6H), 1.17 (t, 3H, $J = 7.0$ Hz).

Ethyl-2-(4-aminophenyl)-2-methylpropionate

In like manner to the preparation of ethyl 4-aminophenoxyacetate, the hydrogenation of ethyl 2-methyl-2-(4-nitrophenyl)propionate provided ethyl-2-(4-aminophenyl)-2-methylpropionate. ^1H NMR (CDCl_3): δ 7.16 (d, 2H, $J = 8.8$ Hz), 6.63 (d, 2H, $J = 8.8$ Hz), 4.09 (qt, 2H, $J = 7.0$ Hz), 3.62 (br s, 2H), 1.52 (s, 6H), 1.17 (t, 3H, $J = 7.0$ Hz).

7.2.41 Anilines substituted with 1,3,4-oxadiazole moieties



15 N'1-(3-Chlorobenzoyl)-3-nitrobenzene-1-carbohydrazide

To a solution of 3-chlorobenzohydrazide (1 equivalent) and pyridine (2 equivalents) in CH_2Cl_2 at 0°C was added a CH_2Cl_2 solution of 3-nitrobenzoyl chloride (1 equivalents) and stirred at 0°C for 1 h and then at room temperature for overnight. The resulting solution was concentrated and diluted with water, basified with NaHCO_3 , the solid was filtered, washed with water, dried and analyzed to obtain N'1-(3-chlorobenzoyl)-3-nitrobenzene-1-carbohydrazide. ^1H NMR ($\text{DMSO}-d_6$): δ 10.99 (s, 1H), 10.79 (s, 1H), 8.73 (bs, 1H), 8.43 (bdd, 1H, $J = 1.2$ and 8.1 Hz), 8.33 (bdd, 1H, $J = 8.4$ Hz), 7.95 (s, 1H), 7.87 (m,

2H), 7.67 (bdd, 1H, J= 1.2 and 8.1 Hz), 7.57 (t, 1H, J= 7.8 Hz); LCMS: purity: 85%; MS (m/e): 320 (MH⁺).

[2-(3-Chlorophenyl)-1,3,4-oxadiazol-5-yl]-3-nitrobenzene

A suspension of N¹-(3-chlorobenzoyl)-3-nitrobenzene-1-carbohydrazide (0.321 g) in POCl₃ (3 mL) was stirred at 90 °C for 24 h. The resulting clear solution was quenched with ice-water, solid obtained was filtered washed with water, dried and analyzed to give [2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl]-3-nitrobenzene. ¹H NMR (DMSO-d₆): δ 8.86 (t, 1H, J= 1.8 Hz), 8.59 (dt, 1H, J= 1.8 and 8.4 Hz), 8.48 (m, 1H), 8.25 (t, 1H, J= 1.8 Hz), 8.16 (dt, 1H, J= 1.2 and 7.5 Hz), 7.93 (t, 1H, J= 8.1 Hz), 7.75 (m, 1H), 7.66 (t, 1H, J= 7.5 Hz), LCMS: purity: 86%; MS (m/e): 302 (MH⁺).

Reduction of [2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl]-3-nitrobenzene


The hydrogenation of [2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl]-3-nitrobenzene (0.2 g) using 10% Pd/C (0.04 g) in MeOH (200 mL) at 15 PSI for 1 h gave a mixture of two products viz. 3-amino-[2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl]benzene and 3-amino-(2-phenyl-1,3,4-oxadiazol-5-yl)benzene which were separated by silica gel column chromatography using n-hexanes then n-hexanes: 5-10% EtOAc as a solvent system. **3-Amino-[2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl]benzene:** ¹H NMR (DMSO-d₆): δ 8.08 (m, 2H), 7.64 (m, 4H), 7.42 (m, 1H), 7.10 (m, 1H); LCMS: purity: 82%; MS (m/e): 272 (MH⁺). **3-Amino-(2-phenyl-1,3,4-oxadiazol-5-yl)benzene:** ¹H NMR (DMSO-d₆): δ 8.13 (m, 1H), 7.54 (m, 5H), 7.30 (m, 1H), 6.86 (dd, 1H, J= 1.5 and 8.1 Hz); LCMS: purity: 93%; MS (m/e): 238 (MH⁺).

N¹-(Ethoxycarbonylmethylenecabonyl)-3-nitrobenzene-1-carbohydrazide

In like manner to the preparation of N¹-(3-chlorobenzoyl)-3-nitrobenzene-1-carbohydrazide, the reaction of 3-nitrobenzoyl chloride with ethoxycarbonylmethylenecarbohydrazide gave N¹-(ethoxycarbonylmethylenecabonyl)-3-nitrobenzene-1-carbohydrazide. ¹H NMR (CD₃OD): δ 8.74 (m, 1H), 8.44 (dd, 1H, 1.8 and 8.1 Hz), 8.25 (bd, 1H, J= 8.4 Hz), 7.76 (t, 1H, J= 8.4 Hz), 4.22 (q, 2H, J= 6.9 Hz), 3.44 (bs, 2H), 1.29 (t, 3H, J= 6.8 Hz); LCMS: purity: 93%; MS (m/e): 296 (MH⁺).

[2-(Ethoxycarbonylmethylene)-1,3,4-oxadiazol-5-yl]-3-nitrobenzene

In like manner to the preparation of [2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl]-3-nitrobenzene the reaction of POCl₃ with N'-1-(ethoxycarbonylmethylenecarbonyl)-3-nitrobenzene-1-carbohydrazide gave [2-(ethoxycarbonylmethylene)-1,3,4-oxadiazol-5-yl]-3-nitrobenzene. ¹H NMR (CDCl₃): δ 8.88 (t, 1H, J= 1.8 Hz), 8.42 (m, 2H), 7.74 (t, 1H, J= 7.5 Hz), 4.27 (q, 2H, J= 7.2 Hz), 4.08 (s, 2H), 1.31 (t, 3H, J= 7.2 Hz); LCMS: purity: 95%; MS (m/e): 278 (MH⁺).

7.2.42 Synthesis of (±)-5-Amino-(2,3-dihydro-2-methoxycarbonyl)benzofuran**2-Methoxycarbonyl-5-nitrobenzofuran**

A mixture of 2-carboxy-5-nitrobenzofuran (2.0 g), MeOH (10 mL) and Concentrated H₂SO₄ (2.1 mL) was heated in a sealed tube at 60 °C for 3 h. Upon cooling to the room temperature it was quenched with ice-water and carefully basified with addition of NaHCO₃. The solid obtained was filtered, washed with water, dried and analyzed to give 2-methoxycarbonyl-5-nitrobenzofuran. ¹H NMR (CDCl₃): δ 8.66 (d, 1H, J= 2.4 Hz), 8.36 (dd, 1H, J= 2.4 and 9.6 Hz), 7.71 (d, 1H, J= 9.3 Hz), 7.65 (s, 1H), 4.01 (s, 3H); LCMS: purity: 97%; MS (m/e): 222 (MH⁺).

15

(±)-5-Amino-(2,3-dihydro-2-methoxycarbonyl)benzofuran

A suspension of 2-methoxycarbonyl-5-nitrobenzofuran (2.0 g), 10% Pd/C (2.0 g), Na₂SO₄ (2.0 g) in MeOH (500 mL) was hydrogenated at 55 PSI for 3 days. The resulting solution was filtered through a pad of celite, concentrated and chromatographed using n-hexanes then 10%, 20% EtOAc/n-hexanes to give (±)-5-amino-(2,3-dihydro-2-methoxycarbonyl)benzofuran. ¹H NMR (CDCl₃): δ 6.69 (d, 1H, J= 8.1 Hz), 6.56 (d, 1H, J= 1.2 Hz), 6.48 (dd, 1H, J= 1.8 and 7.5 Hz), 5.14 (dd, 1H, J= 6.6 and 7.2 Hz), 3.79 (s, 3H), 3.47 (dd, 1H, J= 10.5 and 10.8 Hz), 3.26 (dd, 1H, J= 7.2 and 6.6 Hz); LCMS: purity: 100%; MS (m/e): 194 (MH⁺).

20

25

7.2.43 3-[1-Bis(ethoxycarbonyl)ethoxy]aniline


Preparation of Diethyl 2-methyl-2-(3-nitrophenoxy)malonate

Diethyl 2-bromo-2-methylmalonate (1.0 g, 3.95 mmole) was added to a stirred suspension of potassium fluoride (0.57 g, 9.8 mmole) in dry DMF (5 mL). After stirring for 5 20 min at room temperature, 3-nitrophenol (0.55 g, 3.95 mmole) was added. The resulting mixture was stirred at 60 °C for 6 h, cooled to room temperature, diluted with water (30 mL) and extracted with ethyl acetate (3 X 200 mL). The organic layer was washed with aq. 1N NaOH (2 X 75 mL), dried over anhydrous Na₂SO₄, filtered and evaporated in vacuo to provide diethyl 2-methyl-2-(3-nitrophenoxy)malonate (0.89 g, 80%). ¹H NMR (CDCl₃): 10 δ 7.92 (dd, 1H, J = 2.3 and 8.2 Hz), 7.82 (t, 1H, J = 2.3 Hz), 7.41 (t, 1H, J = 8.2 Hz), 7.30 (dd, 1H, J = 2.3 and 8.2 Hz), 4.28 (qt, 4H, J = 7.0 Hz), 1.81 (s, 3H), 1.26 (t, 6H, J = 7.0 Hz).

Preparation of 3-[1-Bis(ethoxycarbonyl)ethoxy]aniline

Diethyl 2-methyl-2-(3-nitrophenoxy)malonate (0.75 g, 2.40 mmole) was dissolved in toluene: ethanol (1:1, 100 mL), transferred to par shaker bottle containing Pd/C (0.15 g) and anhydrous Na₂SO₄ (5.0 g) in the presence of nitrogen atmosphere. The resulting 15 mixture was treated with hydrogen (30 PSI) till the disappearance of diethyl 2-methyl-2-(3-nitrophenoxy)malonate (2 h). The mixture was filtered through celite covered with anhydrous Na₂SO₄ followed by washing the celite pad with EtOAc. The filtrate was concentrated and dried under vacuo to furnish 3-[1-bis(ethoxycarbonyl)ethoxy]aniline in 20 quantitative yield. ¹H NMR (CDCl₃): δ 6.98 (t, 1H, J = 8.2 Hz), 6.37-6.28 (m, 3H), 4.26 (qt, 4H, J = 7.0 Hz), 3.65 (br s, 2H), 1.72 (s, 3H), 1.24 (t, 6H, J = 7.0 Hz).

7.2.44 Preparation of 4-(4-aminophenoxy)methyl)-2-methoxycarbonyl-furan



25 Preparation of 4-(4-nitrophenoxy)methyl)-2-methoxycarbonyl-furan


3-Nitrophenol (1.0 g, 7.19 mmole), methyl 5-(chloromethyl)-2-furoate (1.38 g, 7.90 mmole) and anhydrous K₂CO₃ (1.19 g, 8.60 mmole) in acetone (30 mL) were refluxed for 8

h. The reaction mixture was cooled and diluted with water. The resultant white solid was filtered, washed with water and air dried overnight to give 1.81 g (90%) of the desired product. ^1H NMR (CDCl_3): δ 7.86 (dd, 1H, $J = 2.3$ and 8.2 Hz), 7.80 (t, 1H, $J = 2.3$ Hz), 7.45 (t, 1H, $J = 8.2$ Hz), 7.27 (dd, 1H, $J = 2.3$ and 8.2 Hz), 7.17 (d, 1H, $J = 3.5$ Hz), 6.58 (d, 1H, $J = 3.5$ Hz), 5.13 (s, 2H), 3.90 (s, 3H).

Preparation of 4-(4-aminophenoxymethyl)-2-methoxycarbonyl-furan

In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, 4-(4-nitrophenoxymethyl)-2-methoxycarbonyl-furan was reduced to provide 4-(4-aminophenoxymethyl)-2-methoxycarbonyl-furan. ^1H NMR (CDCl_3): δ 7.15 (d, 1H, $J = 3.5$ Hz), 7.05 (t, 1H, $J = 8.2$ Hz), 6.50 (d, 1H, $J = 3.5$ Hz), 6.37-6.27 (m, 3H), 5.01 (s, 2H), 3.89 (s, 3H).

7.2.45 Preparation of 6-amino-1-(methoxycarbonyl)methylindazoline



Preparation of 1-(methoxycarbonyl)methyl-6-nitroindazoline

To a solution of 6-nitroindazoline (2.0 g, 12.25 mmole) in dry DMF was added anhydrous K_2CO_3 (1.84 g, 13.31 mmole) and methyl 2-bromoacetate (2.04 g, 13.33 mmole). The resulting mixture was stirred at room temperature for 12 h. The reaction mixture was quenched with water and the resulting solid was collected by filtration, washed with excessive water, and air dried. The yellow solid collected was purified by silica gel column chromatography using gradient solvent system to furnish two products. The desired product (1.12 g, 41%) with high R_f value on the TLC in 30% EtOAc : hexanes was collected.

In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, 1-(Methoxycarbonyl)methyl-6-nitro-indazoline was reduced to provide 6-amino-1-

(methoxycarbonyl)methylindazoline. ^1H NMR (CDCl_3): δ 7.73 (d, 1H, $J = 1.1$ Hz), 7.35 (d, 1H, $J = 8.2$ Hz), 6.49 (dd, 1H, $J = 1.8$ and 8.8 Hz), 6.39 (s, 1H), 5.34 (br s, 2H), 5.10 (s, 2H), 3.64 (s, 3H).

Preparation of 1-(methoxycarbonyl)methyl-5-nitroindazoline

5 In like manner to the preparation of 1-(methoxycarbonyl)methyl-6-nitroindazoline, 1-(methoxycarbonyl)methyl-5-nitroindazoline was prepared by alkylation of 5-nitroindazoline with methyl 2-bromoacetate in presence of K_2CO_3 . The desired product (1.34 g, 46%) with high R_f value on the TLC in 30% EtOAc : hexanes was collected by silica gel column chromatographic purification. ^1H NMR (CDCl_3): δ 8.75 (d, 1H, $J = 1.8$ Hz), 8.30 (dd, 1H, $J = 2.3$ and 8.2 Hz), 8.26 (s, 1H), 7.40 (d, 1H, $J = 8.2$ Hz), 5.22 (s, 2H), 3.78 (s, 3H).

Preparation of 5-amino-1-(methoxycarbonyl)methylindazoline

In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, 1-(Methoxycarbonyl)methyl-5-nitro-indazoline was reduced to provide 5-amino-1-(methoxycarbonyl)methylindazoline. ^1H NMR (CDCl_3): δ 7.84 (d, 1H, $J = 2.3$ Hz), 7.15 (d, 1H, $J = 8.8$ Hz), 6.95 (d, 1H, $J = 2.3$ Hz), 6.88 (dd, 1H, $J = 2.3$ and 8.8 Hz), 5.09 (s, 2H), 3.73 (s, 3H).

Preparation of 1-(2-ethoxycarbonylethyl)-6-nitroindazoline

20 In like manner to the preparation of 1-(methoxycarbonyl)methyl-6-nitroindazoline, 1-(ethoxycarbonyl)ethyl-6-nitroindazoline was prepared by alkylation of 6-nitroindazoline with ethyl 3-bromopropionate in presence of K_2CO_3 . The desired product (58%) with high R_f value on the TLC in 30% EtOAc : Hexanes was collected by silica gel column chromatographic purification. ^1H NMR (CDCl_3): δ 8.49 (s, 1H), 8.12 (s, 1H), 8.01 (dd, 1H, $J = 1.7$ and 8.8 Hz), 7.82 (d, 1H, $J = 8.8$ Hz), 4.74 (t, 2H, $J = 6.4$ Hz), 4.09 (qt, 2H, $J = 7.0$ Hz), 3.03 (t, 2H, $J = 6.4$ Hz), 1.18 (t, 3H, $J = 7.0$ Hz).

Preparation of 6-amino-1-(2-ethoxycarbonylethyl)indazoline

In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, 1-(2-ethoxycarbonylethyl)-6-nitroindazoline was reduced to provide 6-amino-1-(2-ethoxycarbonylethyl)indazoline. ^1H NMR (CDCl_3): δ 7.81 (s, 1H), 7.46 (d, 1H, $J = 8.8$ Hz),

6.60 (app s, 1H), 6.55 (dd, 1H, J = 2.3 and 8.8 Hz), 4.51 (t, 2H, J = 7.0 Hz), 4.11 (qt, 2H, J = 7.0 Hz), 3.52 (br s, 2H), 2.91 (t, 2H, J = 7.0 Hz), 1.18 (t, 3H, J = 7.0 Hz).

Preparation of 1-(2-ethoxycarbonyl-ethyl)-5-nitroindazole

In like manner to the preparation of 1-(methoxycarbonyl)methyl-5-nitroindazole, 1-(ethoxycarbonyl)ethyl-5-nitroindazole was prepared by alkylation of 5-nitroindazole with ethyl 3-bromopropionate in presence of K_2CO_3 . The desired product (43%) with high Rf value on the TLC in 30% EtOAc : Hexanes was collected by silica gel column chromatographic purification. ^1H NMR ($CDCl_3$): δ 8.70 (d, 1H, J = 1.7 Hz), 8.27 (dd, 1H, J = 2.3 and 8.8 Hz), 8.20 (d, 1H, J = 1.7 Hz), 7.59 (d, 1H, J = 8.8 Hz), 4.70 (t, 2H, J = 6.4 Hz), 4.07 (qt, 2H, J = 7.0 Hz), 3.01 (t, 2H, J = 6.4 Hz), 1.16 (t, 3H, J = 7.0 Hz).


Preparation of 5-amino-1-(2-ethoxycarbonyl-ethyl)indazole

In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, 1-(2-ethoxycarbonyl-ethyl)-5-nitroindazole was reduced to provide 5-amino-1-(2-ethoxycarbonyl-ethyl)indazole. ^1H NMR ($CDCl_3$): δ 7.78 (s, 1H), 7.30 (d, 1H, J = 8.8 Hz), 6.91 (d, 1H, J = 2.3 Hz), 6.87 (dd, 1H, J = 2.3 and 8.8 Hz), 4.59 (t, 2H, J = 6.4 Hz), 4.08 (qt, 2H, J = 7.0 Hz), 3.02 (br s, 2H), 2.92 (t, 2H, J = 7.0 Hz), 1.16 (t, 3H, J = 7.0 Hz).

Preparation of 5-amino-2-methylindazole

In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, commercially available 2-methyl-5-nitroindazole was reduced to provide 5-amino-2-methylindazole. ^1H NMR ($CDCl_3$): δ 7.61 (s, 1H), 7.53 (d, 1H, J = 8.8 Hz), 6.81 (dd, 1H, J = 2.3 and 8.8 Hz), 6.75 (d, 1H, J = 2.3 Hz), 4.13 (s, 3H), 3.85 (br s, 2H).

7.2.46 Preparation of methyl 3-methoxy-4-[(6-nitroindazol-1-yl)methyl]benzoate



In like manner to the preparation of 1-(methoxycarbonyl)methyl-6-nitro-indazoline, methyl 3-methoxy-4-[(6-nitroindazol-1-yl)methyl]benzoate was prepared by alkylation of 6-nitroindazoline with methyl (4-bromomethyl)-3-methoxybenzoate in presence of K_2CO_3 .

The desired product (48%) with high R_f value on the TLC in 30% EtOAc : hexanes was

5 collected by silica gel column chromatographic purification. ^1H NMR ($CDCl_3$): δ 8.50 (d, 1H, $J = 1.7$ Hz), 8.14 (s, 1H), 8.00 (dd, 1H, $J = 1.8$ and 8.8 Hz), 7.82 (d, 1H, $J = 8.8$ Hz), 7.56 (s, 1H), 7.54 (d, 1H, $J = 1.8$ Hz), 7.07 (d, 1H, $J = 8.2$ Hz), 5.70 (s, 2H), 3.96 (s, 3H), 3.88 (s, 3H). Low R_f : **Methyl 3-methoxy-4-[(6-nitroindazol-2-yl)methyl]benzoate**: ^1H NMR ($CDCl_3$): δ 8.68 (br s, 1H), 8.07 (s, 1H), 7.86 (dd, 1H, $J = 1.8$ and 9.0 Hz), 7.72 (d, 1H, $J = 9.0$ Hz), 7.61 (d, 1H, $J = 7.7$ Hz), 7.58 (s, 1H), 7.19 (d, 1H, $J = 7.7$ Hz), 5.69 (s, 2H), 10 3.93 (s, 3H), 3.90 (s, 3H).

Preparation of Methyl 4-[(6-aminoindazol-1-yl)methyl]benzoate


In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, methyl 3-methoxy-4-[(6-nitroindazol-1-yl)methyl]benzoate was reduced to provide methyl

15 4-[(6-aminoindazol-1-yl)methyl]benzoate. ^1H NMR ($CDCl_3$): δ 7.88 (s, 1H), 7.53 (d, 1H, $J = 8.8$ Hz), 7.51 (d, 1H, $J = 8.8$ Hz), 7.50 (d, 1H, $J = 1.7$ Hz), 6.67 (d, 1H, $J = 8.8$ Hz), 6.56 (dd, 1H, $J = 1.7$ and 8.8 Hz), 6.45 (d, 1H, $J = 1.2$ Hz), 5.50 (s, 2H), 3.94 (s, 3H), 3.87 (s, 3H), 3.79 (br s, 2H).

Preparation of Methyl 4-[(6-aminoindazol-2-yl)methyl]benzoate

20 In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, methyl 3-methoxy-4-[(6-nitroindazol-2-yl)methyl]benzoate was reduced to provide methyl 4-[(6-aminoindazol-2-yl)methyl]benzoate. ^1H NMR ($CDCl_3$): δ 7.78 (s, 1H), 7.56-7.53 (m, 2H), 7.43 (d, 1H, $J = 8.8$ Hz), 6.98 (d, 1H, $J = 8.2$ Hz), 6.81 (app s, 1H), 6.58 (dd, 1H, $J = 1.8$ and 8.8 Hz), 5.53 (s, 2H), 3.91 (s, 3H), 3.89 (s, 3H).

7.2.47 Preparation of 6-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline



Preparation of 6-nitro-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline

Ester hydrolysis of methyl 3-methoxy-4-[(6-nitroindazol-1-yl)methyl]benzoate in presence of LiOH:H₂O produced the corresponding acid. The acid (1.65 g, 5.04 mmole) thus formed was converted to the acid chloride by reacting with SOCl₂ (3.68 mL, 50.45 mmole) at reflux temperature for 5 h. The reaction mixture was cooled to room temperature and concentrated under vacuo. To acid chloride concentrate dissolved in dry CH₂Cl₂ (75 mL), *o*-toluylbenzenesulfonamide (0.95 g, 5.54 mmole) and 4-(dimethylamino)-pyridine (0.67 g, 5.54 mmole) were added successively at room temperature and stirred for 12 h. The reaction mixture was concentrated, dissolved in EtOAc (700 mL) and successively treated with 2 N HCl (2 X 100 mL), water (150 mL) and brine (100 mL). Usual workup and purification by silica gel column chromatography provided the product (1.57 g, 64%). ¹H NMR (DMSO-d₆): δ 8.75 (s, 1H), 8.31 (s, 1H), 8.00 (d, 1H, J = 8.8 Hz), 7.95-7.91 (m, 2H), 7.50 (d, 1H, J = 1.2 Hz), 7.46-7.27 (m, 4H), 6.92 (d, 1H, J = 7.6 Hz), 5.76 (s, 2H), 3.81 (s, 3H), 2.54 (s, 3H).

Preparation of 6-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline

In like manner to the reduction of diethyl 2-methyl-2-(3-nitrophenoxy)malonate, 6-nitro-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline was reduced to provide 6-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline. ¹H NMR (CDCl₃): δ 7.96 (dd, 1H, J = 1.2 and 8.2 Hz), 7.76 (s, 1H), 7.51 (d, 1H, J = 1.2 Hz), 7.49-7.44 (m, 1H), 7.37 (d, 2H, J = 8.8 Hz), 7.34-7.32 (m, 1H), 7.30 (d, 1H, J = 8.8 Hz), 6.51-6.47 (m, 2H), 6.35 (s, 1H), 5.35 (s, 2H), 3.89 (s, 3H), 2.54 (s, 3H).

Preparation of methyl 3-methoxy-4-[(5-nitroindazol-1-yl)methyl]benzoate

In like manner to the preparation of methyl 3-methoxy-4-[(6-nitroindazol-1-yl)methyl]benzoate, methyl 3-methoxy-4-[(5-nitroindazol-1-yl)methyl]benzoate was prepared by alkylation of 5-nitroindazoline with methyl (4-bromomethyl)-3-methoxybenzoate in presence of K_2CO_3 . The desired product (47%) with high R_f value on the TLC in 30% EtOAc : Hexanes as eluent was collected by silica gel column chromatographic purification. ^1H NMR ($CDCl_3$): δ 8.73 (d, 1H, $J = 1.8$ Hz), 8.26-8.22 (m, 2H), 7.56 (s, 1H), 7.54 (dd, 1H, $J = 1.8$ and 8.2 Hz), 7.49 (d, 1H, $J = 9.4$ Hz), 6.98 (d, 1H, $J = 8.2$ Hz), 5.66 (s, 2H), 3.91 (s, 3H), 3.89 (s, 3H). Low R_f : Methyl 3-methoxy-4-[(5-nitroindazol-2-yl)methyl]benzoate.


Preparation of 5-nitro-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline

In like manner to the preparation of 6-nitro-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline, 5-nitro-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline was prepared from methyl 3-methoxy-4-[(5-nitroindazol-1-yl)methyl]benzoate. ^1H NMR ($DMSO-d_6$): δ 8.81 (d, 1H, $J = 2.3$ Hz), 8.39 (s, 1H), 8.21 (dd, 1H, $J = 1.8$ and 8.8 Hz), 7.87 (dd, 2H, $J = 3.6$ and 8.8 Hz), 7.48 (d, 1H, $J = 1.2$ Hz), 7.39 (dd, 1H, $J = 1.2$ and 8.2 Hz), 7.33-7.15 (m, 3H), 6.85 (d, 1H, $J = 8.2$ Hz), 5.65 (s, 2H), 3.76 (s, 3H), 2.49 (s, 3H).

Preparation of 5-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline

In like manner to the preparation of 6-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline, 5-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline was prepared by reduction of 5-nitro-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline. ^1H NMR ($DMSO-d_6$): δ 7.87 (dd, 1H, $J = 1.2$ and 7.7 Hz), 7.73 (s, 1H), 7.50 (s, 1H), 7.35-7.14 (m, 5H), 6.78 (d, 1H, $J = 1.8$ Hz), 6.75 (s, 1H), 6.53 (d, 1H, $J = 8.2$ Hz), 5.44 (s, 2H), 3.82 (s, 3H), 2.50 (s, 3H).

7.2.48 Preparation of 8-amino-4*H*-imidazo[2,1-*c*][1,4]-benzoxazine



7.3 Synthesis of 2,4-Pyrimidinediamines

A variety of 2,4-pyrimidinediamines of the invention were synthesized from the above starting materials and intermediates and other commercially available reagents. Conditions suitable for synthesizing N2,N4-bis-substituted-2,4-pyrimidinediamine compounds ("general S_NAr " reaction conditions; Substitution Nucleophilic Aromatic Reaction) are exemplified with N2,N4-bis(4-ethoxyphenyl)-2,4-pyrimidinediamine (**R926069**) and N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (**R921218**). Conditions suitable for synthesizing asymmetric N2,N4-disubstituted-2,4-pyrimidinediamines are exemplified by N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (**R926210**).

7.3.1 N2,N4-Bis(4-ethoxyphenyl)-2,4-pyrimidinediamine (**R926069**)

To a solution of 2,4-dichloropyrimidine (0.015 g, 0.1 mmol) in EtOH (1 mL) was added 4-ethoxyaniline (0.034 g, 0.025 mmol) and heated in a sealed tube at 70-80 °C for 24h. Upon cooling the reaction was diluted with H₂O (10 mL), acidified with 2N HCl, the solid obtained was filtered, washed with H₂O and dried to give N2,N4-bis(4-ethoxyphenyl)-2,4-pyrimidinediamine (**R926069**). ¹H NMR (CD₃OD): δ 7.63 (d, 1H), 7.45 (d, 2H), J= 9 Hz), 7.32 (d, 2H, J= 9.3 Hz), 6.95 (d, 2H, J= 6.9 Hz), 6.87 (d, 2H, J= 8.7 Hz), 6.23 (d, 1H, J= 7.2 Hz), 4.04 (m, 4H), 1.38 (m, 6H); LCMS: ret. time: 25.91 min.; purity: 99.5%; MS (m/e): 351 (MH⁺).

7.3.2 N2,N4-Bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (**R921218**)

A mixture of 2,4-dichloro-5-fluoropyrimidine (0.0167 g, 0.1 mmol) and 3-aminophenol (0.033 g, 0.3 mmol) in MeOH: H₂O (1.8:0.2 mL; v/v) was shaken in a sealed tube at 100 °C for 24h (or 80 °C for 3 days), cooled to room temperature, diluted with water (15 mL), acidified with 2N HCl (pH >2). Upon saturation with sodium chloride it was

extracted with ethyl acetate (3 x 20 mL), dried over anhydrous sodium sulfate and solvent was removed. The resulting residue was filtered through a pad of silica gel (200-400 mesh) using CH₂Cl₂ - >1 >10% MeOH in CH₂Cl₂ to obtain the desired N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (**R921218**). If the reaction scale is large enough, solid of the resulting product can be isolated by filtration. ¹H NMR (CDCl₃): δ 7.73 (d, 1H, J= 5.1 Hz), 7.12-6.90 (m, 6H), 6.64 (dd, 1H, J= 1.8 and 8.1 Hz), 6.53 (dd, 1H, J= 1.2 and 5.7 Hz); LCMS: ret. time: 16.12 min.; purity: 100%; MS (m/e): 313 (MH⁺).

7.3.3 N₂,N₄-Bis(4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (**R926017**)

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-methoxyaniline were reacted to yield N₂,N₄-bis(4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.67 (d, 1H, J= 4.8 Hz), 7.43 (d, 2H, J= 9.3 Hz), 7.67 (d, 2H, J= 8.7 Hz), 6.87 (d, 2H, J= 9.6 Hz), 6.83 (d, 2H, J= 8.7 Hz), 3.83 (s, 3H), 3.81 (s, 3H); LCMS: ret. time: 22.53 min.; purity: 100%; MS (m/e): 341 (MH⁺).

7.3.4 N₂,N₄-Bis(3-fluoro-4-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine (**R926018**)

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-fluoro-4-trifluoromethylaniline were reacted to yield N₂,N₄-bis(3-fluoro-4-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.01 (d, 1H, J= 3 Hz), 7.77 (m, 3H), 7.61 (dt, 1H, J= 4.2 and 3 Hz), 7.20 (t, 1H, 8.7 Hz), 7.12 (t, 1H, J= 9.3 Hz), 6.95 (s, 1H), 6.82 (s, 1H); ¹⁹F NMR (CDCl₃): δ -17505 (s, 3F), -17517 (s, 3F), -17525 (s, F), -17537 (s, F), -46835 (s, 1F); LCMS: ret. time: 32.39 min.; purity: 95%; MS (m/e): 453 (MH⁺).

7.3.5 N₂,N₄-Bis(3,4-tetrafluoroethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (**R926037**)

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3,4-tetrafluoroethylenedioxyaniline were reacted to yield N₂,N₄-bis(3,4-tetrafluoroethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.01 (d, 1H, J= 3.0 Hz), 7.71 (d, 1H, J= 2.4 Hz), 7.70 (1H, d, J= 2.4 Hz), 7.18 (dd, 2H, J= 2.4 and 6 Hz), 7.07 (d, 2H, J= 1.8 Hz), 7.00 (1H, bs), 6.81 (d, 1H, J= 2.7 Hz); ¹⁹F NMR (CDCl₃): -

26029 (sept, 8F), -46791 (s, C5-F); LCMS: ret. time: 38.20 min.; purity: 85%; MS (m/e): 541 (MH⁺).

7.3.6 N2,N4-Bis(3-trifluoromethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926038)

5 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-trifluoromethoxyaniline were reacted to yield N2,N4-bis(3-trifluoromethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.03 (bd, 1H), 7.62 (bs, 2H), 7.48 (bd, 1H), 7.39 (t, 1H, J= 8.1 Hz), 7.34 (m, 1H), 7.29 (t, 1H, J= 7.5 Hz), 7.01 (m, 2H), 6.88 (m, 2H); ¹⁹F NMR (CDCl₃): -16447 (s, 3F), -16459 (s, 3F), -46738 (s, 1F); LCMS: ret. time: 33.77 min.; purity: 93%; MS (m/e): 449 (MH⁺).

7.3.7 N2,N4-Bis(4-chloro-3-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926039)

15 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-chloro-3-trifluoromethylaniline were reacted to yield N2,N4-bis(4-chloro-3-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.05 (bs, 1H), 7.89 (bd, 1H), 7.77 (dd, 1H, J= 2.4 and 9 Hz), 7.65 (dd, 1H, J= 2.4 and 8.7 Hz), 7.49 (d, J= 8.1 Hz), 7.40 (d, 1H, J= 6.2 Hz), 7.03 (s, 1H), 6.91 (s, 1H); ¹⁹F NMR (CDCl₃): δ -17864 (s, 3F), -17894 (s, 3F), -46550 (s, 1F); LCMS: ret. time: 38.81 min.; purity: 75%; MS (m/e): 485 (MH⁺).

7.3.8 N2,N4-Bis(3-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926064)

25 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-ethoxyaniline were reacted to yield N2,N4-bis(3-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.96 (1H, d, J= 4.8 Hz), 7.22 (m, 6H), 7.07 (t, 1H, J= 1.8 Hz), 6.95 (dt, 1H, J= 1.2 and 7.2 Hz), 6.77 (m, 2H), 3.88 (q, 4H, J= 6.3 Hz), 1.33 (two t, 6H, J= 6.3 Hz); ¹⁹F NMR (CDCl₃): -46175; LCMS: ret. time: 26.86 min.; purity: 97%; MS (m/e): 369 (MH⁺).

7.3.9 N2,N4-Bis(3-hydroxy-4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926339)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-hydroxy-4-methoxyaniline were reacted to yield N2,N4-bis(3-hydroxy-4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.82 (d, 1H J= 4 Hz), 7.18 (m, 2H), 6.95 (m, 2H), 6.83 (m, 2H) 3.93 (s, 6H); LCMS: ret. time: 16.63 min.; purity: 97 %; MS (m/e): 373 (MH⁺).

7.3.10 N2,N4-Bis(4-ethoxycarbonylamino-3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926340)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-ethoxycarbonylamino-3-hydroxyaniline were reacted to yield N2,N4-bis(4-ethoxycarbonylamino-3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.86 (d, 1H J= 4 Hz), 7.67 (m, 2H), 7.20 (dd, 1H, J= 8 Hz, J= 4.1 Hz), 7.13 (d, 1H), 6.90 (m, 2H), 4.2 (m, 4H), 1.32 (m, 6H); LCMS: ret. time: 20.92 min.; purity: 98 %; MS (m/e): 487 (MH⁺).

7.3.11 N2,N4-Bis(-3-hydroxy-4-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926341)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-hydroxy-4-methylaniline were reacted to yield N2,N4-bis(-3-hydroxy-4-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.83 (d, 1H J= 4 Hz), 7.11 (m, 4H), 6.81 (m, 2H), 2.19 (m, 6H); LCMS: ret. time: 20.69 min.; purity: 98 %; MS (m/e): 341 (MH⁺).

7.3.12 N2,N4-Bis[4-(2-methoxyethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926342)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-(2-methoxyethoxy)aniline were reacted to yield N2,N4-bis[4-(2-methoxyethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.89 (d, 1H J= 4 Hz), 7.54 (dd, 2H, J= 6.8 and 2.7 Hz), 7.38 (dd, 2H, J= 6.8 and 2.7 Hz), 6.87 (dd, 2H, J= 6.8 and 2.7 Hz), 6.82 (dd, 2H, J= 6.8 and 2.7 Hz) 4.6 (m, 4H), 4.11 (m, 4H), 3.35 (m, 6H); LCMS: ret. time: 21.76 min.; purity: 97 %; MS (m/e): 429 (MH⁺).

7.3.13 N2,N4-Bis(dihydrobenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine (R909237)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 5-amino-2,3-dihydrobenzofuran were reacted to yield N2,N4-bis(dihydrobenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.99 (d, 1H J= 4 Hz), 7.22 (m, 4H), 6.81 (m, 2H), 4.55 (m, 4H), 3.22 (m, 4H); LCMS: ret. time: 23.80 min.; purity: 98 %; MS (m/e): 438 (MH⁺).

7.3.14 N2,N4-Bis(3-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926065)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-methoxyaniline were reacted to yield N2,N4-bis(3-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.96 (d, 1H, J= 5.4 Hz), 7.24 (m, 6H), 7.06 (t, 1H, J= 2.4 Hz), 7.00 (dt, 1H, J= 1.2 Hz), 6.79 (m, 1H), 3.72 (s, 3H), 3.70 (s, 3H); ¹⁹F NMR (CD₃OD): δ - 46112; LCMS: ret. time: 23.46 min.; purity: 99%; MS (m/e): 341 (MH⁺).

7.3.15 N2,N4-Bis[4-(N,N-dimethylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926086)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-N,N-dimethylaniline were reacted to yield N2,N4-bis[4-(N,N-dimethylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.84 (d, 1H, J= 3.6 Hz), 7.43 (d, 2H, J= 8.7 Hz), 7.34 (d, 2H, J= 8.7 Hz), 7.25 (s, 1H), 6.73 (m, 4H), 6.55 (s, 1H), 2.95 (s, 6H), 2.90 (s, 6H); ¹⁹F NMR (CDCl₃): - 47770; LCMS: ret. time: 12.48 min.; purity: 99%; MS (m/e): 367 (MH⁺).

7.3.16 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926109)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3,4-ethylenedioxyaniline were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.88 (d, 1H, J= 3.6 Hz), 7.23 (d, 1H, J= 2.3 Hz), 7.15 (d, 1H, J= 2.4 Hz), 7.00 (dd, 1H, J= 3 and 8.1 Hz), 6.98 (dd, 1H, J= 3 and 8 Hz), 6.83 (d, 1H, J=8.7 Hz), 6.81 (d, 1H, J= 8.7 Hz), 6.7(s, 1H), 6.58 (s, 1H), 4.23 (m, 4H), 4.24(m, 4H); ¹⁹F NMR (CDCl₃): δ - 47445; LCMS: ret. time: 21.81 min.; purity: 96%; MS (m/e): 397 (MH⁺).

7.3.17 N2,N4-Bis(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926110)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3,4-dimethoxyaniline were reacted to yield N2,N4-bis(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.90 (d, 1H, J= 1.8 Hz), 7.13 (d, 2H, J= 4.8 Hz), 7.08 (d, 1H, J= 8.7 Hz), 6.94 (d, 2H, J= 10.5 Hz), 6.81 (d, 1H, J= 8.7 Hz), 6.76 (d, 1H, J= 8.7 Hz), 6.70 (s, 1H), 3.87 (s, 3H), 3.84 (s, 3H), 3.74 (s, 3H), 3.71 (s, 3H); ¹⁹F NMR (CDCl₃): δ - 47433; LCMS: ret. time: 19.64 min.; purity: 95%; MS (m/e): 401 (MH⁺).

7.3.18 N2,N4-Bis[4-(N-morpholino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926114)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-N-morpholinylaniline were reacted to yield N2,N4-bis[4-(N-morpholino)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.80 (s, 1H), 7.78 (s, 1H, partially exchanged), 7.76 (bs, 1H, partially exchanged), 7.53 (d, 2H, J= 8.1 Hz), 7.39 (d, 2H, J= 9 Hz), 6.93 (d, 2H, J= 8.7 Hz), 6.86 (bd, 2H), 3.84 (m, 8H), 3.11 (m, 8H); ¹⁹F NMR (CD₃OD): δ - 47697; LCMS: ret. time: 18.15 min.; purity: 99.55%; MS (m/e): 451 (MH⁺).

7.3.19 N2,N4-Bis(4-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R926206)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-chloroaniline were reacted to yield N2,N4-bis(4-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃ + CD₃OD): δ 7.80 (d, 1H, J= 4.2 Hz), 7.45 (d, 2H, J= 8.7 Hz), 7.33 (d, 2H, J= 9 Hz), 7.20 (d, 2H, J= 8.7 Hz), 7.14 (d, 2H, J= 9.6 Hz); LCMS: ret. time: 28.84 min.; purity: 87%; MS (m/e): 349 (MH⁺).

7.3.20 N2,N4-Bis(3-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R926209)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-chloroaniline were reacted to yield N2,N4-bis(3-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.08 (d, 1H, J= 5.4 Hz), 7.70 (t, 1H, J= 1.8 Hz), 7.57 (t, 1H, J= 1.2 Hz), 7.54 (m, 1H), 7.35

(m, 4H), 7.28 (t, 1H, J= 1.8 Hz), 7.24 (m, 1H), 7.22 (t, 1H, J= 1.8 Hz); ^{19}F NMR (CD_3OD): - 43631; LCMS: ret. time: 28.99 min.; purity: 99%; MS (m/e): 349 (M^+).

7.3.21 N2,N4-Bis(4-tert-butylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926222)

5 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-tert-butylaniline were reacted to yield N2,N4-bis(4-tert-butylphenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.77 (d, 1H, J= 3.9 Hz), 7.47 (d, 2H, J= 9Hz), 7.38 (m, 4H), 7.30 (d, 2H, J= 8.7 Hz), 1.34 (s, 9H), 1.32 (s, 9H); LCMS: ret. time: 34.09 min.; purity: 93%; MS: 393 (MH^+).

10 **7.3.22 N2,N4-Bis(3-chloro-4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine (R926223)**

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-chloro-4-fluoroaniline were reacted to yield N2,N4-bis(3-chloro-4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$): δ 7.81 (d, 1H), 7.60 (m, 1H), 7.58 (m, 1H), 7.38 (m, 1H), 7.19 (m, 1H), 7.0 (m, 2H); LCMS: ret. time: 28.98 min.; purity: 97%; MS (m/e): 385 (M^+).

7.3.23 N2,N4-Bis(4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine (R926224)

20 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-fluoroaniline were reacted to yield N2,N4-bis(4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 8.79 (d, 2H, J= 5.4 Hz), 7.40 (m, 2H), 7.30 (m, 2H), 6.90 (m, 4H); ^{19}F NMR (CDCl_3): - 32425 (s, 1F), -32940 (s, 1F), -45525 (s, 1F); LCMS: ret. time: 23.53 min.; purity: 100%; MS (m/e): 317 (MH^+).

25 **7.3.24 N2,N4-Bis(4-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926225)**

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-methylaniline were reacted to yield N2,N4-bis(4-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 30 7.73 (d, 1H, J= 4.2 Hz), 7.43 (d, 2H, J= 8.1 Hz), 7.36 (d, 2H, J= 8.4 Hz), 7.14 (d, 2H, J= 8.4

Hz), 7.10 (d, 2H, J= 8.1 Hz), 2.39 (s, 3H), 2.35 (s, 3H); LCMS: ret. time: 25.81 min.; purity: 99.65%; MS (m/e): 309 (MH⁺).

7.3.25 N2,N4-Bis[(4-methoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926240)

5 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and ethyl 4-aminophenoxyacetate were reacted to yield N2,N4-bis[(4-methoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.8 (bs, 1H), 7.50 (d, 2H, J= 9.3 Hz), 7.32 (d, 2H, J= 8.41 Hz), 6.88 (m, 4H), 4.72 (s, 2H), 4.70 (s, 2H), 3.79 (s, 3H), 3.78 (s, 3H); ¹⁹F
10 NMR (CDCl₃): -47570; LCMS: ret. time: 21.17 min.; purity: 95%; MS (m/e): 457 (MH⁺).

7.3.26 (+)-N2,N4-Bis[4-methoxycarbonyl(α-methyl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R926254)

15 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and (±)-ethyl 2-(4-aminophenoxy)propionate were reacted to yield (±)-N2,N4-bis[4-methoxycarbonyl(α-methyl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.89 (bs, 1H), 7.48 (dd, 2H, J= 2.4 and 6.9 Hz), 7.40 (dd, 2H, J= 1.8 and 6.9 Hz), 6.85 (m, 4H), 6.76 (s, 1H), 6.63 (s, 1H), 4.75 (hex, 2H, J= 6.3 Hz), 3.77 (s, 3H), 3.76 (s, 3H), 1.62 (t, 6H, J= 7.5 Hz); LCMS: ret. time: 23.76 min.; purity: 97%; MS (m/e): 485 (MH⁺).

7.3.27 N2,N4-Bis[(3-methoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926255)

25 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and ethyl 3-aminophenoxyacetate were reacted to yield N2,N4-bis[(3-methoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.96 (d, 1H, J= 2.4 Hz), 7.71 (t, 1H, J= 2.4 Hz), 7.44 (m, 2H), 7.21 (m, 3H), 6.96 (dd, 1H, J= 1.2 and 7.8 Hz), 6.86 (d, 1H, J= 3 Hz), 6.53 (m, 1H), 4.64 (s, 2H), 4.60 (s, 2H), 3.79 (s, 6H); LCMS: ret. time: 21.72 min.; purity: 87%; MS (m/e): 457 (MH⁺).

7.3.28 N2,N4-Bis(3-acetyloxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926387)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-acetoxyaniline were reacted to yield N2,N4-bis[(3-acetoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. Alternatively, N2,N4-bis[(3-acetoxyphenyl)-5-fluoro-2,4-pyrimidinediamine can be prepared by acetylation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine with acetyl chloride in the presence of pyridine in CH₂Cl₂. ¹H NMR (CDCl₃): δ 8.00 (bs, 1H), 7.51-7.25 (m, 8H), 2.32 (s, 3H), 2.28 (s, 3H); LCMS: ret. time: 22.14 min; purity: 100%; MS (m/e): 397 (MH⁺).

7.3.29 N2,N4-Bis(3-benzyloxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926394)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-benzyloxyaniline were reacted to yield N2,N4-bis(3-benzyloxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.98 (bs, 1H), 7.42-6.99 (m, 16H), 6.75 (d, 1H, J= 2.4 Hz), 6.71 (m, 1H), 6.60 (dd, 1H, J= 2.4 and 8.4 Hz), 6.32 (m, 1H), 4.97 (s, 2H), 4.94 (s, 2H); LCMS: ret. time: 32.56 min.; purity: 98%; MS (m/e): 493 (MH⁺).

7.3.30 N2,N4-Bis(2-phenylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926398)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-phenylaniline were reacted to yield N2,N4-bis[(2-phenylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.35 (m, 1H), 8.0 (s, 1H), 7.85 (s, 1H), 7.45-7.00 (m, 18H); LCMS: ret. time: 30.29 min.; purity: 68%; MS (m/e): 433 (MH⁺).

7.3.31 (R926404) N2, N4-Bis(2-phenylphenyl)-5-methyl-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N2,N4-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-aminobiphenyl and 2,4-dichloro-5-methylpyrimidine were reacted to provide N2, N4-bis(2-phenylphenyl)-5-methyl-2,4-pyrimidinediamine. LCMS: ret. time: 30.47 min.; purity: 91%; MS (m/e): 429 (MH⁺).

7.3.32 N2,N4-Bis[(4-methoxy-3-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926399)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-methoxy-3-phenylaniline were reacted to yield N2,N4-bis[(4-methoxy-3-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.83 (d, 1H, J= 4.2 Hz), 7.57 (bd, 1H, J= 8.7 Hz), 7.48 (d, 1H, J= 2.7 Hz), 7.47-7.22 (m, 12H), 6.85 (d, 1H, J= 8.7 Hz), 6.78 (d, 1H, 9.3 Hz), 3.72 (s, 3H), 3.69 (s, 3H); LCMS: ret. time: 29.97 min.; purity: 92%; MS (m/e): 493 (MH⁺).

7.3.33 N2,N4-Bis[(2-methoxy-5-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926400)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-methoxy-5-phenylaniline were reacted to yield N2,N4-bis[(2-methoxy-5-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.03 (d, 1H, J= 6.6 Hz), 7.76 (t, 1H, J= 2.4 Hz), 7.28-7.10 (m, 13H), 7.07 (d, 1H, J= 9 Hz), 7.01 (d, 1H, J= 8.1 Hz), 3.91 (s, 3H), 3.86 (s, 3H); LCMS: ret. time: 18.58 min.; purity: 96%; MS (m/e): MH⁺.

7.3.34 N2,N4-Bis[(2-methoxy-5-methyl-4-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926401)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-methoxy-5-methyl-4-phenylaniline were reacted to yield N2,N4-bis[(2-methoxy-5-methyl-4-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.00 (d, 1H, J= 4.8 Hz), 7.73 (s, 1H), 7.66 (s, 1H), 7.43-7.24 (m, 9H), 6.91 (s, 1H), 6.82 (s, 1H), 3.86 (s, 3H), 3.85 (s, 3H), 2.14 (s, 3H), 1.99 (s, 3H); LCMS: ret. time: 19.98 min.; purity: 99%; MS (m/e): 521 (MH⁺).

7.3.35 N2,N4-Bis[(2-methyl-5-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926402)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-methyl-5-phenylaniline were reacted to yield N2,N4-bis[(2-methyl-5-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.84 (bd, 1H), 7.51-7.20 (m, 16H), 2.30 (s, 3H), 2.24 (s, 3H); LCMS: ret. time: 18.57 min.; purity: 87%; MS (m/e): 461 (MH⁺).

7.3.36 N₂,N₄-Bis[(3-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926403)

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-phenylaniline were reacted to yield N₂,N₄-bis[(3-phenyl)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.02 (d, 1H, J= 5.1 Hz), 7.82 (t, 1H, J= 1.5 Hz), 7.67 (t, 1H, J= 1.8 Hz), 7.58 (dd, 1H, J= 1.2 and 7.2 Hz), 7.42-7.24 (m, 15H); LCMS: ret. time: 32.06 min.; purity: 94%; MS (m/e): 433 (MH⁺).

7.3.37 N₂,N₄-Bis(4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926405)

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-acetoxylaniline were reacted to yield N₂,N₄-bis[(4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. After the work up it was observed that the acetoxy group was hydrolyzed to afford the N₂,N₄-bis(4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine instead of the corresponding acetate derivative. ¹H NMR (CD₃OD): δ 7.74 (d, 1H, J= 5.6 Hz), 7.43 (dd, 2H, J= 2.1 and 6.6 Hz), 7.28 (dd, 2H, J= 2.4 and 6.3 Hz), 6.74 (dd, 2H, J= 2.4 and 6.3 Hz), 6.66 (dd, 2H, J= 2.4 and 7.2 Hz); ¹⁹F NMR (CD₃OD): - 48116 (d, 1F); LCMS: ret. time: 16.15 min; purity: 100%; MS (m/e): 313 (MH⁺).

7.3.38 N₂,N₄'-Bis(4-hydroxy-3-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926469)

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-hydroxy-3-methylaniline were reacted to yield N₂,N₄-bis[(4-hydroxy-3-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.64 (d, 1H, J= 3.6 Hz), 7.11 (t, 2H, J= 9 Hz), 6.70-6.45 (m, 4H), 2.15 (s, 3H), 2.09 (s, 3H); ¹⁹F NMR (CD₃OD): - 46278; LCMS: ret. time: 15.53; purity: 84%; MS (m/e): 341 (MH⁺).

7.3.39 N₂,N₄-Bis[4-(tert-butoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926574)

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and tert-butyl 4-aminophenoxyacetate were reacted to yield N₂,N₄-bis[4-(tert-butoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-

pyrimidinediamine ^1H NMR (CDCl_3): δ 7.88 (s, 1H), 7.48 (d, 2H, $J=8.4$ Hz), 7.40 (d, 2H, $J=8.7$ Hz), 6.86 (m, 4H), 4.52 (s, 2H), 4.48 (s, 2H), 1.49 (s, 9H), 1.48 (s, 9H); LCMS: ret. time: 28.48 min.; purity: 95%; MS (m/e): 541 (MH^+).

7.3.40 N2,N4-Bis(indol-5-yl)-5-fluoro-2,4-pyrimidinediamine (R926582)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 5-aminoindole were reacted to yield N2,N4-bis(indol-5-yl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 20.26 min.; purity: 99%; MS (m/e): 359 (MH^+).

7.3.41 N2,N4-Bis(4-cyanomethylphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926319)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine, 2,4-dichloro-5-ethoxycarbonylpyrimidine and 4-cyanomethylaniline were reacted to yield N2,N4-bis(4-cyanomethylphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine. ^1H NMR (DMSO-d_6): δ 8.72 (s, 1H), 7.64 (m, 4H), 7.32 (d, 2H, $J=8.7$ Hz), 7.21 (d, 2H, $J=8.4$ Hz), 4.3 (q, 2H, $J=7.0$ Hz), 3.97 (s, 2H), 3.89 (s, 2H), 1.32 (3H, $J=7$ Hz); LCMS: ret. time: 30.83 min.; purity: 90 %; MS (m/e): 413 (MH^+).

7.3.42 N2,N4-Bis(3-indazol-6-yl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926320)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine, 2,4-dichloro-5-ethoxycarbonylpyrimidine and 6-aminoindazole were reacted to yield N2,N4-bis(6-indazolyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine. ^1H NMR (DMSO-d_6): δ 8.76 (s, 1H), 7.73 (d, 2H, $J=8.8$), 7.54 (m, 4H), 7.36 (d, 2H, $J=9.5$ Hz), 4.3 (q, 2H, $J=7.0$ Hz), 1.34 (3H, $J=7$ Hz); LCMS: ret. time 27.59 min.; purity: 95 %; MS (m/e): 415 (MH^+).

7.3.43 N2,N4-Bis(3-indazol-7-yl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926321)

In like manner to N2,N4-bis(3-hydroxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine, 2,4-dichloro-5-ethoxycarbonylpyrimidine and 7-aminoindazole were reacted to yield N2,N4-bis(7-indazolyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine. ^1H NMR

(DMSO-d₆): δ 8.70 (s, 1H), 7.54 (d, 2H J= 8.4 Hz), 7.37 (m, 6H), 4.3 (q, 2H, J= 7.0 Hz), 1.33 (3H, J= 7 Hz); LCMS: ret. time 23.61 min.; purity: 94 %; MS (m/e): 415 (MH⁺).

7.3.44 N₂,N₄-Bis[6-(1,4-benzoxazine-3-onyl)]-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926325)

5 In like manner to to N₂,N₄-bis(3-hydroxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine, 2,4-dichloro-5-ethoxycarbonylpyrimidine and 6-amino-1,4-benzoxazine-3-one were reacted to yield N₂,N₄-bis[6-(1,4-benzoxazine-3-onyl)]-5-ethoxycarbonyl-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.66 (s, 1H), 7.21 (dd, 2H J= 8.8 and J= 2.2 Hz), 6.89 (d, 2H J= 8.4 Hz), 4.54 (s, 2H) 4.49 (s, 2H) 4.3 (q, 2H, J= 7.0 Hz), 1.33 (3H, J= 7 Hz); LCMS: ret. time 23.08 min.; purity: 88 %; MS (m/e): 477 (MH⁺).

7.3.45 N₂,N₄-Bis(4-ethoxycarbonylmethyleneaminophenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926331)

In like manner to to N₂,N₄-bis(3-hydroxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine, 2,4-dichloro-5-ethoxycarbonylpyrimidine and 4-ethoxycarbonylmethyleneaminoaniline were reacted to yield N₂,N₄-bis(4-ethoxycarbonylaminophenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.72 (s, 1H), 7.70 (d, 2H J= 8.8 Hz), 7.28 (d, 2H J= 8.4 Hz), 7.05 (d, 2H, J= 8.4 Hz) 6.82 (d, 2H J= 8.4 Hz) 4.5 (m, 4H), 4.23 (m, 6H) 1.53 (m, 9H); LCMS: ret. time 18.08 min.; purity: 85%; MS (m/e): 537 (MH⁺).

7.3.46 N₂,N₄-Bis(4-ethoxyphenyl)-6-methoxycarbonyl-2,4-pyrimidinediamine (R926058)

20 In a manner analogous to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-6-methoxycarbonylpyrimidine with 4-ethoxyaniline gave N₂,N₄-bis(4-ethoxyphenyl)-6-methoxycarbonyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.42 (bs, 1H), 7.35 (bd, 4H), 6.85 (bs, 1H), 6.75 (bd, 4H), 3.97 (q, 4H, J= 4.8 Hz), 3.92 (s, 3H), 1.36 (t, 6H, J= 6.3 Hz); LCMS: ret. time: 27.47 min.; purity: 97%; MS (m/e): 409 (MH⁺).

7.3.47 N₂,N₄-Bis(4-ethoxyphenyl)-5-methyl-2,4-pyrimidinediamine (R926068)

30 In a manner analogous to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-methylpyrimidine with 4-

ethoxyaniline gave N2,N4-bis(4-ethoxyphenyl)-5-methyl-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.55 (s, 1H), 7.40 (d, 2H), 7.21 (d, 2H, J= 8.7 Hz), 6.90 (dd, 4H, J= 8.7 Hz), 4.04 (q, 4H, J= 6.6 Hz), 2.17 (m, 6H); LCMS: ret. time: 26.51 min.; purity: 95%; MS (m/e): 365 (MH⁺).

5 **7.3.48 N2,N4-Bis(4-ethoxyphenyl)-6-chloro-2,4-pyrimidinediamine (R926072)**

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4,6-trichloropyrimidine with 4-ethoxyaniline gave N2,N4-bis(4-ethoxyphenyl)-6-chloro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.42 (d, 10 2H, J= 9 Hz), 7.18 (d, 2H, J= 8.7 Hz), 6.89 (d, 2H, J= 6.3 Hz), 6.84 (d, 2H, J= 8.7 Hz), 6.58 (bs, 1H), 4.02 (m, 4H), 1.43 (m, 6H); LCMS: ret. time: 83.21 min.; purity: 87%; MS (m/e): 385 (MH⁺).

7.3.49 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-methyl-2,4-pyrimidinediamine (R926242)

15 In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-methylpyrimidine with 3,4-ethyleneoxyaniline gave N2,N4-bis(3,4-ethylenedioxyphenyl)-5-methyl-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.75 (bs, 1H), 7.06 (d, 1H, J= 2.4 Hz), 6.96 (d, 1H, J= 2.1 Hz), 6.94 (d, 1H, J= 2.1 Hz), 6.85-6.77 (m, 2H), 6.70 (d, 1H, J= 9 Hz), 4.23 (s, 20 4H), 4.19 (s, 4H), 2.09 (s, 3H); LCMS: ret. time: 22.01 min.; purity: 100%; MS (m/e): 393 (MH⁺).

7.3.50 N2,N4-Bis(3,4-ethylenedioxyphenyl)-2,4-pyrimidinediamine (R926243)

25 In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloropyrimidine with 3,4-ethyleneoxyaniline gave N2,N4-bis(3,4-ethylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.95 (s, 1H), 10.50 (s, 1H), 7.84 (bd, 2H), 7.24 (bd, 2H), 6.79 (bd, 2H), 6.40 (bd, 2H), 4.24 (s, 8H); LCMS: ret. time: 21.68 min.; purity: 100%; MS (m/e): 379 (MH⁺).

7.3.51 N2,N4-Bis(3-hydroxyphenyl)-5-methyl-2,4-pyrimidinediamine (R926248)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-methylpyrimidine with 3-hydroxyaniline gave N2,N4-bis(3-hydroxyphenyl)-5-methyl-2,4-pyrimidinediamine. LCMS: ret. time: 16.76 min.; purity: 100%, MS (m/e): 309 (MH⁺).

7.3.52 N2,N4-Bis(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926249)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloropyrimidine with 3-hydroxyaniline gave N2,N4-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 16.21 min.; purity: 100%; MS (m/e): 295 (MH⁺).

7.3.53 N2,N4-Bis[(4-methoxycarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R926256)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloropyrimidine with methyl 4-aminophenoxyacetate gave N2,N4-bis[(4-methoxycarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.7 (bs, 1H), 10.28 (bs, 1H), 7.84 (d, 1H, J= 6.9 Hz), 7.48 (bd, 2H), 7.35 (d, 2H, J= 8.7 Hz), 6.95 (d, 2H, J= 9 Hz), 6.90 (d, 2H, J= 8.7 Hz), 6.35 (d, 1H, J= 6.9 Hz), 4.81 (s, 2H), 4.79 (s, 2H), 3.69 (s, 3H), 3.68 (s, 3H); LCMS: ret. time: 21.27 min.; purity: 98%; MS (m/e): 439 (MH⁺).

7.3.54 (+)-N2,N4-Bis[4-methoxycarbonyl(alpha-methyl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R926257)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloropyrimidine with (+)-methyl 2-(4-aminophenoxy)propionate gave (+)-N2,N4-bis[4-methoxycarbonyl(alpha-methyl)methyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 24.09 min.; purity: 90%; MS (m/e): 467 (MH⁺).

7.3.55 N2,N4-Bis(4-methoxycarbonylmethyleneoxyphenyl)-5-methyl-2,4-pyrimidinediamine (R926258)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-methylpyrimidine with methyl-4-aminophenoxyacetate gave N2,N4-bis(4-methoxycarbonylmethyleneoxyphenyl)-5-methyl-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.21 (s, 1H), 9.65 (s, 1H), 7.78 (s, 1H), 7.42 (dd, 2H, J= 2.7 and 8.7 Hz), 7.28 (dd, 2H, J= 8.1 Hz), 6.94 (d, 2H, J= 8.47 Hz), 6.85 (d, 2H, J= 8.7 Hz), 4.82 (s, 2H), 4.77 (s, 2H), 3.69 (s, 3H), 3.68 (s, 3H), 2.12 (s, 3H); LCMS: ret. time: 21.76 min.; purity: 100%; MS (m/e): 453 (MH⁺).

7.3.56 (±)-N2,N4-Bis[4-ethoxycarbonyl(α-methyl)methyleneoxyphenyl]-5-methyl-2,4-pyrimidinediamine (R926259)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-methylpyrimidine with (±)-ethyl 2-(4-aminophenoxy)propionate gave (±)-N2,N4-bis[4-ethoxycarbonyl(α-methyl)methyleneoxyphenyl]-5-methyl-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.9 (bs, 1H), 9.35 (bs, 1H), 7.79 (s, 1H), 7.43 (dd, 2H, J= 3.6 and 8.7 Hz), 7.32 (d, 2H, J= 7.5 Hz), 6.86 (d, 2H, J= 9 Hz), 6.78 (d, 2H, J= 8.7 Hz), 4.95 (q, 1H, J= 7.2 Hz), 4.90 (q, 1H, J= 7.2 Hz), 4.12 (2q, 4H, J= 5.7 Hz), 2.10 (s, 3H), 1.51 (d, 3H, J= 6.3 Hz), 1.47 (d, 3H, J= 6.3 Hz), 1.16 (2t, 6H, J= 5.7 Hz); LCMS: ret. time: 27.41 min.; purity: 96%; MS (m/e): 509 (MH⁺).

7.3.57 N2,N4-Bis[2-(4-hydroxyphenyl)ethyl]-5-methyl-2,4-pyrimidinediamine (R926397)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-methylpyrimidine with 2-(4-hydroxyphenyl)ethylamine gave N2,N4-bis[2-(4-hydroxyphenyl)ethyl]-5-methyl-2,4-pyrimidinediamine. LCMS: ret. time: 19.94 min.; purity: 100%; MS (m/e): 365 (MH⁺).

7.3.58 N2,N4-Bis-(3,4-dimethoxyphenyl)-5-nitro-2,4-pyrimidinediamine (R940089)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-nitropyrimidine with 3,4-dimethoxyaniline gave N2,N4-bis-(3,4-dimethoxyphenyl)-5-nitro-2,4-pyrimidinediamine.

LCMS: ret. time: 28.30 min.; purity: 100%; MS (m/e): 428 (MH⁺); ¹H NMR (CDCl₃): δ 10.30 (1H, s), 9.14 (1H, s), 7.52 (1H, s), 7.08 (3H, m), 7.00 (1H, d, J= 8.4 Hz), 6.84 (1H, d, J= 8.4 Hz), 6.76 (1H, d, J= 8.4 Hz), 3.90 (3H, s), 3.87 (3H, s), 3.68 (3H, s), 3.60 (3H, s).

7.3.59 N2,N4-Bis-(4-ethoxyphenyl)-5-nitro-2,4-pyrimidinediamine (R940090)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-nitropyrimidine with 4-ethoxyaniline gave N2,N4-bis-(4-ethoxyphenyl)-5-nitro-2,4-pyrimidinediamine. LCMS: ret. time: 35.91 min.; purity: 100%; MS (m/e): 396 (MH⁺); ¹H NMR (CDCl₃): δ 10.25 (1H, s), 9.11 (1H, s), 7.44 (2H, d, J= 8.6 Hz), 7.37 (2H, d, J= 9Hz), 6.88 (2H, d, J= 8.6 Hz), 6.80 (2H, d, J= 8.6 Hz), 4.06 (2H, q, J= 7.2 Hz), 4.02 (2H, q, J= 7.2 Hz), 1.45 (3H, t, J= 7.2 Hz), 1.42 (3H, t, J= 7.2 Hz).

7.3.60 N2,N4-Bis-(3,4-ethylenedioxyphenyl)-5-nitro-2,4-pyrimidinediamine (R940095)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-nitropyrimidine with 3,4-ethylenedioxyaniline gave N2,N4-bis-(3,4-ethylenedioxyphenyl)-5-nitro-2,4-pyrimidinediamine. LCMS: ret. time: 30.78 min.; purity: 100%; MS (m/e): 424 (MH⁺); ¹H NMR (CDCl₃): δ 10.21 (1H, s), 9.10 (1H, s), 7.40 (1H, s), 7.11-6.71 (6H, m), 4.29 (4H, s), 4.25 (4H, s).

7.3.61 N2,N4-Bis-[(4-ethoxycarbonylmethyleneoxy)phenyl]-5-nitro-2,4-pyrimidinediamine (R940096)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-nitropyrimidine with ethyl-4-aminophenoxyacetate gave N2,N4-bis-[(4-ethoxycarbonylmethyleneoxy)phenyl]-5-nitro-2,4-pyrimidinediamine. LCMS: ret. time: 32.48 min.; purity: 94%; MS (m/e): 512 (MH⁺); ¹H NMR (CDCl₃): δ 10.22 (1H, s), 9.13 (1H, s), 7.50 (1H, s), 7.45 (2H, d, J= 8.7 Hz), 7.38 (2H, d, J= 8.7 Hz), 6.93 (2H, d, J= 8.7 Hz), 6.83 (2H, d, J= 8.7 Hz), 4.67 (2H, s), 4.63 (2H, s), 4.29 (2H, q, J= 7.2 Hz), 4.28 (2H, q, J= 7.2 Hz), 1.31 (3H, t, J= 7.2 Hz), 1.30 (3H, t, J= 7.2 Hz).

7.3.62 N2,N4-Bis-(2,2-difluoro-1,3-benzodioxol-5-yl)-5-nitro-2,4-pyrimidinediamine (R940100)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-nitropyrimidine with 2,2-difluoro-5-amino-1,3-benzodioxole gave N2,N4-bis-(2,2-difluoro-1,3-benzodioxol-5-yl)-5-nitro-2,4-pyrimidinediamine. LCMS: ret. time: 38.15 min.; purity: 96%; MS (m/e): 467 (M⁺); ¹H NMR (CDCl₃): δ 10.76 (1H, s), 10.49 (1H, s), 9.20 (1H, s), 7.74 (2H, s), 7.56 (1H, d, J=11.4 Hz), 7.33 (2H, m), 7.20 (1H, m).

7.3.63 N2,N4-Bis-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940215)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-fluoropyrimidine with 3,5-dichloro-4-hydroxyaniline gave N2,N4-bis-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 21.26 min.; purity: 88%; MS (m/e): 450 (M⁺); ¹H NMR (DMSO-d₆): δ 9.96 (1H, s), 9.59 (1H, s), 9.47 (1H, s), 9.37 (1H, s), 8.22 (1H, d, J=3.6 Hz), 7.79 (2H, s), 7.74 (2H, s).

7.3.64 N2,N4-Bis-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R940216)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-fluoropyrimidine with 3-chloro-4-hydroxy-5-methylaniline gave N2,N4-bis-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 20.55 min.; purity: 99%; MS (m/e): 410 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.23 (1H, s), 9.07 (1H, s), 8.99 (1H, s), 8.66 (1H, s), 8.13 (1H, d, J=3.6 Hz), 7.59 (2H, t, J=3.1 Hz), 7.50 (1H, d, J=2.3 Hz), 7.34 (1H, d, J=2.3 Hz), 2.27 (3H, s), 2.18 (3H, s).

7.3.65 N2,N4-Bis-(2,3-dimethyl-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940217)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-fluoropyrimidine with 2,3-dimethyl-4-hydroxyaniline gave N2,N4-bis-(2,3-dimethyl-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 19.07 min.; purity: 99%; MS (m/e): 369 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.21 (1H, s), 8.99 (1H, s), 8.63 (1H, s), 7.92 (1H, s), 7.84 (1H, d, J=

3.6 Hz), 6.94 (1H, d, J= 8.5 Hz), 6.85 (1H, d, J= 8.5 Hz), 6.70 (1H, d, J= 8.5 Hz), 6.58 (1H, d, J= 8.5 Hz), 2.12 (3H, s), 2.06 (3H, s), 2.02 (3H, s), 1.94 (3H, s).

7.3.66 N2,N4-Bis-(4-Acetamidophenyl)-5-fluoro-2,4-pyrimidinediamine (R940222)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-fluoropyrimidine with 4-acetamidoaniline gave N2,N4-bis-(4-acetamidophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 14.82 min.; purity: 95%; MS (m/e): 395 (MH⁺); ¹H NMR (DMSO-d₆): δ 10.33 (1H, s), 10.14 (1H, s), 10.07 (2H, s), 8.39 (1H, d, J= 5.1 Hz), 7.64 (8H, m), 2.15 (3H, s), 2.13 (3H, s).

7.3.67 N2,N4-Bis(3-isopropylphenyl)-5-fluoro-2,4-pyrimidinediamine R940297

In like manner to the preparation of N2,N4-bis-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-isopropylaniline were reacted to give N2,N4-bis-(3-isopropylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 29.58 min.; Purity: 98 %; MS (m/e): 365 (MH⁺); ¹H NMR (DMSO-d₆): δ 10.5 (1H, s), 10.34 (1H, s), 8.41 (1H, d, J= 5.1 Hz), 7.62 (1H, d, J= 8.1 Hz), 7.53 (1H, s), 7.43 (1H, d, J= 8.1 Hz), 7.37 (2H, m), 7.29 (1H, t, J= 8.1 Hz), 7.19 (1H, d, J= 7.8 Hz), 7.08 (1H, d, J= 7.8 Hz), 2.88 (2H, m), 1.25 (6H, d, J= 7.2 Hz), 1.201 (6H, d, J= 7.2 Hz).

7.3.68 N2,N4-Bis(3,4,5-trimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926688)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3,4,5-trimethoxyaniline were reacted to yield N2,N4-bis(3,4,5-trimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 19.55 min.; purity: 99 %; MS (m/e): 461 (MH⁺).

7.3.69 N2,N4-Bis(2-methyl-5-phenylphenyl)-5-bromo-2,4-pyrimidinediamine R925800

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and 5-phenyl-*ortho*-toluidine were reacted to yield N2,N4-bis(2-methyl-5-phenylphenyl)-5-bromo-2,4-pyrimidinediamine. LCMS: ret. time: 19.54 min.; purity: 90 %; MS (m/e): 422 (MH⁺).

7.3.70 N2,N4-Bis(2-methoxy-5-methyl-4-phenylphenyl)-5-bromo-2,4-pyrimidinediamine (R925801)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine 5-methyl-4-phenyl-*ortho*-anisidine
5 were reacted to yield N2,N4-bis(2-methoxy-5-methyl-4-phenylphenyl)-5-bromo-2,4-pyrimidinediamine. LCMS: ret. time: 20.99 min.; purity: 85 %; MS (m/e): 583 (MH⁺).

7.3.71 N2,N4-Bis(indol-6-yl)-5-fluoro-2,4-pyrimidinediamine (R926594)

10 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 6-aminoindole were reacted to yield N2,N4-bis(indol-6-yl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 22.39 min.; purity: 85%; MS (m/e): 359 (MH⁺).

7.3.72 N2,N4-Bis(2-methoxycarbonyl benzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine (R926604)

15 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-methoxycarbonyl-5-aminobenzofuran were reacted to yield N2,N4-bis(2-methoxycarbonyl benzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.3 (bs, 1H), 10.05 (bs, 1H), 8.25 (d, 1H, J= 5.4 Hz), 8.06 (s, 1H), 7.94 (s, 1H), 7.77-7.49 (m, 5H), 7.36 (bs, 1H), 3.89 (s, 3H),
20 3.87 (s, 3H).

7.3.73 N2,N4-Bis[4-(methoxycarbonylmethyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926605)

25 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and ethyl 4-aminophenyl acetate were reacted to yield N2,N4-bis[4-(methoxycarbonylmethyl)phenyl]-5-fluoro-2,4-pyrimidinediamine. The cross esterification reaction of ethyl ester to obtain the corresponding methyl ester was observed. ¹H NMR (CDCl₃): δ 10.62 (s, 1H), 8.06 (s, 1H), 7.69 (d, 1H, J= 4.5 Hz), 7.53 (d, 2H, J= 8.1 Hz), 7.43 (d, 2H, J= 8.7 Hz), 7.30 (d, 2H, J= 8.4 Hz), 7.20 (d, 2H, J= 8.4 Hz), 3.73 (s, 3H), 3.72 (s, 3H), 3.67 (s, 2H), 3.63 (s, 2H).

7.3.74 N2,N4-Bis(2-ethoxycarbonylindol-5-yl)-5-fluoro-2,4-pyrimidinediamine (R926616)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-ethoxycarbonyl-5-indoleamine were reacted to yield N2,N4-bis(2-ethoxycarbonylindol-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 11.83 (s, 1H), 11.63 (s, 1H), 9.21 (s, 1H), 8.99 (s, 1H), 8.08 (s, 1H), 8.01 (m, 2H), 7.49-7.22 (m, 4H), 6.92 (s, 1H), 6.63 (s, 1H), 4.29 (q, 4H, J= 7.2 Hz), 1.32 (m, 6H); LCMS: ret. time: 24.74 min.; purity: 99%; MS (m/e): 503 (MH⁺).

7.3.75 N2,N4-Bis(coumarin-6-yl)-5-fluoro-2,4-pyrimidinediamine (R926617)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 6-aminocoumarin were reacted to yield N2,N4-bis(coumarin-6-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.17 (d, 2H, J= 3.6 Hz), 7.97-7.74 (m, 5H), 7.40 (1H, d, J= 8.7 Hz), 7.30 (d, 1H, J= 9Hz), 6.50 (d, 1H, J= 10.2 Hz), 6.40 (d, 1H, J= 9.3 Hz); LCMS: ret. time: 19.05 min.; purity: 94%; MS (m/e): 417 (MH⁺).

7.3.76 N2,N4-Bis(4-methoxymethyl)coumarin-7-yl)-5-fluoro-2,4-pyrimidinediamine (R926620)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 7-amino-4-methoxymethylcoumarin were reacted to yield N2,N4-bis(coumarin-7-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.38 (s, 1H), 8.42 (d, 1H, J= 3 Hz), 8.28 (m, 1H), 8.05-7.93 (m, 2H), 7.77-7.50 (m, 4H), 6.31 (s, 1H), 6.29 (s, 1H), 4.66 (s, 2H), 4.65 (s, 2H), 3.43 (s, 3H), 3.41 (s, 3H); LCMS: MS (m/e): 505 (MH⁺).

7.3.77 N2,N4-Bis(3-(hydroxymethyl)phenyl)-5-fluoro-2,4-pyrimidinediamine (R925757)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-aminobenzylalcohol were reacted to yield N2,N4-bis(3-(hydroxymethyl)phenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.90 (d, 1H, J= 3.3 Hz), 7.71 (m, 1H), 7.61 (d, 1H, J= 6.9 Hz), 7.50 (d,

1H, J= 6.0), 7.47 (s, 1H), 7.31 (t, 1H, J= 8.1 Hz), 7.22 (t, 1H, J= 8.1 Hz), 7.10 (d, 1H, J= 6.9), 6.97 (d, 1H, J= 7.5 Hz), 4.63 (s, 4H); LCMS: ret. time: 15.36 min.; purity: 100%; MS (m/e): 342 (MH⁺).

7.3.78 N2,N4-Bis[(2R)-hydroxy-(1S)-methyl-2-phenylethyl]-5-fluoro-2,4-pyrimidinediamine (R925767)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and (1R,2S)-(-)-norephedrine were reacted to yield N2,N4-bis[(2R)-hydroxy-(1S)-methyl-2-phenylethyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (acetone-d₆): δ 7.67 (s, 1H), 7.49-7.42 (m, 4H), 7.38-7.19 (m, 6H), 6.09 (d, 1H, J= 9.0 Hz), 5.73 (d, 1H, J= 7.5 Hz), 5.61 (d, 1H, J= 9.3 Hz), 5.04 (d, 1H, J= 3.6 Hz), 4.97 (d, 1H, J= 2.7 Hz), 4.74 (bs, 1H), 4.48 (bs, 1H), 4.30-4.25 (m, 1H), 1.09 (d, 1H, J= 6.9 Hz), 1.03 (d, 1H, J= 6.6 Hz); LCMS: ret. time: 21.56 min.; purity: 98%; MS (m/e): 397(MH⁺).

7.3.79 N2,N4-Bis(2-hydroxy-2-phenylethyl)-5-fluoro-2,4-pyrimidinediamine (R925768)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-amino-1-phenylethanol were reacted to yield N2,N4-bis(2-hydroxy-2-phenylethyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (acetone-d₆): δ 8.15 (s, 1H), 7.46-7.22 (m, 10H), 5.01 (dd, 1H), 4.91 (dd, 1H), 4.78 (dd, 1H), 3.86-3.18 (m, 5H); LCMS: ret. time: 19.64 min.; purity: 89 %; MS (m/e): 369 (MH⁺).

7.3.80 N2,N4-Bis(furfuryl)-5-fluoro-2,4-pyrimidinediamine (R925769)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and furfurylamine were reacted to yield N2,N4-bis(furfuryl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.72 (bs, 1H), 7.38 (dd, 2H, J= 1.8 and 7.5 Hz), 6.34-6.30 (m, 2H), 6.22 (dd, 2H, J= 2.4 and 9.9 Hz), 5.163 (bs, 2H), 4.63 (d, 2H, J= 6.0), 4.54 (d, 2H, J= 6.0).; ¹⁹F NMR (CDCl₃): - 48621; LCMS: ret. time: 97.27min.; purity: 97%; MS (m/e): 289 (MH⁺).

7.3.81 N2,N4-Bis(piperonyl)-5-fluoro-2,4-pyrimidineamine (R925770)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and piperonylamine were reacted to yield N2,N4-bis(piperonyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.60 (bs, 1H), 6.78-6.69 (m, 6H), 5.93 (s, 2H), 5.91 (s, 2H), 4.51 (d, 2H, J= 5.7 Hz), 4.43 (d, 2H, J= 5.1 Hz); ¹⁹F NMR (CDCl₃): - 45257; LCMS: ret. time: 22.06 min.; purity: 96%; MS (m/e): 397 (MH⁺).

7.3.82 N2,N4-Dibenzyl-5-fluoro-2,4-pyrimidinediamine (R925772)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and benzylamine were reacted to yield N2,N4-bis(benzyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.69 (bs, 1H), 7.35-7.24 (m, 10H), 5.63 (bs, 1H), 5.27 (bs, 1H), 4.61 (d, 2H, J= 6.0 Hz), 4.55 (d, 2H, J= 6.0 Hz); ¹⁹F NMR (CDCl₃): - 48580; LCMS: ret. time: 23.73 min.; purity: 100%; MS (m/e): 309 (MH⁺).

7.3.83 N2,N4-Bis(3,4-methylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R925776)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3,4-methylenedioxyaniline were reacted to yield N2,N4-bis(3,4-methylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.86 (bs, 1H), 7.27 (m, 1H), 7.19 (m, 1H), 6.89 (dd, 2H, J= 2.1 and 8.1 Hz), 6.80 (dd, 2H, J= 1.8 and 8.1 Hz), 6.73 (t, 2H, J= 8.1 Hz), 5.97 (s, 2H), 5.92 (s, 2H); ¹⁹F NMR (CDCl₃): - 47591; LCMS: ret. time: 21.74 min.; purity: 97%; MS (m/e): 369 (MH⁺).

7.3.84 N2,N4-Bis[2-(4-hydroxyphenyl)ethyl]-5-fluoro-2,4-pyrimidinediamine (R925791)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and tyramine were reacted to yield N2,N4-bis[2-(4-hydroxyphenyl)ethyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.17 (bs, 1H), 8.22 (bs, 1H), 6.99 (d, 4H, J= 8.1 Hz), 6.65 (d, 4H, J= 8.1 Hz), 3.48-3.43 (m, 4H), 2.72 (t, 4H, J= 7.7 Hz); LCMS: ret. time: 19.19 min.; purity: 100 %; MS (m/e): 369 (MH⁺).

7.3.85 N2,N4-Bis(4-cyanophenyl)-5-fluoro-2,4-pyrimidinediamine (R945057)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-4-pyrimidineamine, 4-aminobenzonitrile and 2,4-dichloro-5-fluoropyrimidine gave N2,N4-bis(4-cyanophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 7.26 (d, J= 8.7 Hz, 2 H), 7.36 (d, J= 9.0 Hz, 2 H), 7.43 (d, J= 8.7 Hz, 2 H), 7.60 (d, J= 8.7 Hz, 2 H), 7.86 (d, J= 3.6 Hz, 1 H), 9.49 (br, 1 H, NH), 9.51 (br, 1 H, NH); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 161.48; LC: 27.15 min.; 100%; MS (m/e): 331.00 (MH⁺).

7.3.86 N2,N4-Bis(4-ethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926234)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-ethylaniline were reacted to yield N2,N4-bis(4-ethylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.83 (bs, 1H), 7.77 (d, 1H, J= 3.9 Hz), 7.48 (d, 2H, J= 8.7 Hz), 7.40 (d, 2H, J= 8.7 Hz), 7.31 (bs, 1H), 7.18 (d, 2H, J= 8.7 Hz), 7.11 (d, 2H, J= 8.7 Hz), 2.68-2.61 (m, 4H), 1.28-1.21 (m, 6H); LCMS: ret. time: 29.17 min.; purity: 100 %; MS (m/e): 337(MH⁺).

7.3.87 N2,N4-Bis(3-chloro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926675)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-chloro-4-hydroxyaniline were reacted to yield N2,N4-bis(3-chloro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.83 (d, 1H, J= 4.2 Hz), 7.59 (d, 1H, J= 2.4 Hz), 7.53 (d, 1H, J= 2.4 Hz), 7.40 (dd, 1H, J= 2.4 and 8.7 Hz), 7.20 (dd, 1H, J= 2.4 and 8.7 Hz), 6.89 (d, 1H, J= 8.7 Hz), 6.81 (d, 1H, J= 8.7 Hz); ¹⁹F NMR (CD₃OD): - 47862; LCMS: ret. time: 17.89 min.; purity: 99 %; MS (m/e): 382 (MH⁺).

7.3.88 N2,N4-Bis[3-chloro-4-(ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R926676)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-chloro-4-(ethoxycarbonylmethyleneoxy)aniline were reacted to yield N2,N4-bis[3-chloro-4-(ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.93 (bs, 1H), 7.67-7.65 (m, 2H), 7.41 (dd, 1H, J= 3.0 and 9.3 Hz), 7.26 (dd, 1H, J= 2.7

and 9.3 Hz), 6.92-6.85 (m, 3H), 6.69 (d, 1H, J= 2.4 Hz), 4.71 (s, 2H), 4.66 (s, 2H), 4.32-4.23 (m, 4H), 1.33-1.27 (m, 6H); ^{19}F NMR (CDCl_3): - 47274; LCMS: ret. time: 27.51 min.; purity: 97 %; MS (m/e): 553 (M^+).

7.3.89 N2,N4-Bis(3-fluoro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926681)

5

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-fluoro-4-hydroxyaniline were reacted to yield N2,N4-bis(3-fluoro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.83 (d, 1H), 7.53 (dd, 1H), 7.42 (dd, 1H), 7.22 (dq, 1H), 7.03 (dq, 1H), 6.89 (d, 1H), 6.83 (s, 1H), 6.80 (s, 1H), 6.78 (d, 1H); ^{19}F NMR (CDCl_3): - 390060, - 39165, - 47835; LCMS: ret. time: 15.27 min.; purity: 95 %; MS (m/e): 349 (MH^+).

10

7.3.90 N2,N4-Bis(3-acetamidophenyl)-5-fluoro-2,4-pyrimidinediamine (R926682)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-aminoacetanilide were reacted to yield N2,N4-bis(3-acetamidophenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR ($\text{DMSO}-d_6$): δ 10.24 (bs, 1H), 10.03 (s, 1H), 9.94 (s, 1H), 8.20 (d, 1H, J= 4.8 Hz), 7.91 (bs, 1H), 7.68 (bs, 1H), 7.43 (d, 1H, J= 8.1 Hz), 7.35-7.30 (m, 2H), 7.24-7.19 (m, 2H), 7.11 (t, 1H, J= 8.1 Hz), 2.03 (s, 3H), 2.01 (s, 3H); LCMS: ret. time: 15.10 min.; purity: 99 %; MS (m/e): 395 (MH^+).

15

20

7.3.91 N2,N4-Bis(2-fluoro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926683)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-fluoro-4-hydroxyaniline were reacted to yield N2,N4-bis(2-fluoro-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR ($\text{DMSO}-d_6$): δ 9.78 (s, 1H), 9.50 (s, 1H), 8.75 (s, 1H), 8.06 (s, 1H), 7.87 (d, 1H, J= 4.2 Hz), 7.25-7.18 (m, 2H), 6.61 (dd, 1H, J= 2.4 and 12.3 Hz), 6.56-6.47 (m, 2H), 6.39 (dd, 1H, J= 1.8 and 8.7 Hz); LCMS: ret. time: 15.52 min.; purity: 99 %; MS (m/e): 349 (MH^+).

25

7.3.92 N2,N4-Bis(4-isopropoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926701)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-isopropoxyaniline were reacted to yield N2,N4-bis(4-isopropoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.89 (bs, 1H), 7.47 (d, 2H, J= 8.7 Hz), 7.38 (d, 2H, J= 9.0 Hz), 6.87 (d, 2H, J= 9.0 Hz), 6.83 (d, 2H, J= 8.7 Hz); LCMS: ret. time: 27.51 min.; purity: 98 %; MS (m/e): 397 (MH⁺).

7.3.93 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine (R925771)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and 3,4-ethylenedioxyaniline were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.07 (bs, 1H), 7.16 (d, 1H, J= 3.0 Hz), 7.10 (d, 1H, J= 2.7 Hz), 6.98-6.93 (m, 2H), 6.90-6.75 (m, 3H), 4.28-4.21 (m, 8H); LCMS: ret. time: 22.61 min.; purity: 100%; MS (m/e): 458 (MH⁺).

7.3.94 N2,N4-Bis(3-hydroxyphenyl)-5-bromo-2,4-pyrimidinediamine (R925778)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and 3-aminophenol were reacted to yield N2,N4-bis(3-hydroxyphenyl)-5-bromo-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.99 (bs, 1H), 9.34 (bs, 1H), 8.30 (s, 1H), 7.15 (t, 1H, J= 8.4 Hz), 7.06-6.97 (m, 2H), 6.94-6.92 (m, 2H), 6.80 (bs, 1H), 6.62 (s, 1H, J= 8.1 Hz), 6.43 (d, 1H, J= 7.8 Hz); LCMS: ret. time: 18.48 min.; purity: 97%; MS (m/e): 374 (MH⁺).

7.3.95 N2,N4-Bis[4-(ethoxycarbonylmethyleneoxy)phenyl]-5-bromo-2,4-pyrimidinediamine (R925779)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and ethyl 4-aminophenoxyacetate were reacted to yield N2,N4-bis[4-(ethoxycarbonylmethyleneoxy)phenyl]-5-bromo-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.12 (s, 1H), 8.48 (s, 1H), 8.11 (s, 1H), 7.42 (d, 4H, J= 8.7 Hz), 6.89 (d, 2H, J= 9.0 Hz), 6.71 (d, 2H, J= 9.3 Hz), 4.78 (s, 2H), 4.66 (s,

2H), 4.20-4.10 (m, 4H), 1.23-1.16 (m, 6H); LCMS: ret. time: 25.82 min.; purity: 94%; MS (m/e): 546 (MH⁺).

7.3.96 N₂,N₄-Bis[2-(4-hydroxyphenyl)ethyl]-5-bromo-2,4-pyrimidinediamine (R925792)

5 In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and tyramine were reacted to yield N₂,N₄-bis[2-(4-hydroxyphenyl)ethyl]-5-bromo-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 7.83 (s, 1H), 6.96 (d, 4H, J = 8.1 Hz), 6.63 (d, 4H, J = 8.1 Hz), 3.54-3.42 (m, 2H), 2.74-2.66 (m, 2H), 2.74-2.66 (m, 4H); ret. time: 20.10 min.; purity: 100 %; MS (m/e): 430 (MH⁺).

7.3.97 N₂,N₄-Bis(2-phenylphenyl)-5-bromo-2,4-pyrimidinediamine (R925798)

15 In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and 2-aminobiphenyl were reacted to yield N₂,N₄-bis(2-phenylphenyl)-5-bromo-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.34 (d, 1H, J = 8.1 Hz), 8.27 (d, 1H, J = 8.1 Hz), 8.00 (s, 1H), 7.51-7.18 (m, 17H), 6.95 (s, 1H); LCMS: ret. time: 18.87 min.; purity: 97 %; MS (m/e): 495 (MH⁺).

7.3.98 N₂,N₄-Bis(2-methoxy-5-phenylphenyl)-5-bromo-2,4-pyrimidinediamine (R925799)

20 In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and 5-phenyl-*ortho*-anisidine were reacted to yield N₂,N₄-bis(2-methoxy-5-phenylphenyl)-5-bromo-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.26 (m, 2H), 8.05 (m, 2H), 7.39-7.21 (m, 12H), 7.17 (dd, 1H, J = 2.4 and 8.1 Hz), 7.11 (d, 1H, J = 8.7 Hz), 7.05 (d, 1H, J = 9.0 Hz), 3.88 (s, 3H), 3.83 (s, 3H); LCMS: ret. time: 20.51 min.; purity: 98 %; MS (m/e): 554 (MH⁺).

7.3.99 N₂,N₄-Bis(4-methoxy-3-phenylphenyl)-5-bromo-2,4-pyrimidinediamine (R925802)

30 In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, with the addition of triethylamine, 5-bromo-2,4-dichloropyrimidine and 3-phenyl-*para*-anisidine hydrochloride were reacted to yield N₂,N₄-bis(4-methoxy-3-phenylphenyl)-5-bromo-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.26 (m, 2H),

8.06 (m, 2H), 7.38-7.25 (m, 12H), 7.18 (dd, 1H, J= 2.4 and 8.1 Hz), 7.12 (d, 1H, J= 8.7 Hz), 7.05 (d, 1H, 8.7 Hz), 3.89 (s, 3H), 3.83 (s, 3H); LCMS: ret. time: 36.77 min.; purity: 98 %; MS (m/e): 554 (MH⁺).

7.3.100 N2,N4-Bis(3-phenylphenyl)-5-bromo-2,4-pyrimidinediamine (R925803)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-bromo-2,4-dichloropyrimidine and 3-aminobiphenyl were reacted to yield N2,N4-bis(3-phenylphenyl)-5-bromo-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.86 (bs, 1H), 9.20 (bs 1H), 8.33 (s, 1H), 7.79 (bs, 1H), 7.18 (bs, 1H), 7.61 (d, 1H), 7.56-7.51 (m, 2H), 7.48-7.23 (m, 11H), 7.17-7.04 (m, 2H); LCMS: ret. time: 19.52 min.; purity: 80 %; MS (m/e): 494 (MH⁺).

7.3.101 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-cyano-2,4-pyrimidinediamine (R925773)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-cyanopyrimidine and 3,4-ethylenedioxyaniline were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-cyano-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.69 (bs, 1H), 9.28 (bs, 1H), 8.40 (s, 1H), 7.16-6.89 (m, 4H), 6.79 (d, 1H, J= 9.0 Hz), 6.65 (bs, 1H), 4.22 (s, 4H), 4.16 (s, 4H); LCMS: ret. time: 24.42 min.; purity: 93 %; MS (m/e): 404 (MH⁺).

7.3.102 N2,N4-Bis(3-hydroxyphenyl)-5-cyano-2,4-pyrimidinediamine (R925774)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-cyanopyrimidine and 3-hydroxyaniline were reacted to yield N2,N4-bis(3-hydroxyphenyl)-5-cyano-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.73 (bs, 1H), 9.40 (s, 1H), 9.33 (bs, 1H), 9.24 (s, 1H), 8.47 (s, 1H), 7.20 (d, 1H, J= 7.5 Hz), 7.11 (t, 1H, J= 7.5 Hz), 7.09-7.02 (m, 2H), 6.99-6.89 (m, 3H), 6.54 (d, 1H, J= 7.2 Hz), 6.37 (dd, 1H, J= 1.8 and 8.4 Hz); LCMS: ret. time: 19.71 min.; purity: 97%; MS (m/e): 320 (MH⁺).

7.3.103 N2,N4-Bis[4-(ethoxycarbonylmethyleneoxy)phenyl]-5-cyano-2,4-pyrimidinediamine (R925775)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-cyanopyrimidine and ethyl 4-aminophenoxyacetate were reacted to yield N2,N4-bis[4-(ethoxycarbonylmethyleneoxy)phenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.80 (s, 1H), 7.40 (d, 4H, J= 8.7 Hz), 6.90 (4H, J= 9.0 Hz), 6.82-6.75 (m, 2H), 4.60 (bs, 4H), 4.29-4.25 (m, 4H), 1.32-1.26 (m, 5H), LCMS: ret. time: 28.50 min.; purity: 100 %; MS (m/e): 493 (MH⁺).

7.3.104 R935192: N2, N4-Bis(1-methyl-indazolin-5-yl)-5-fluoro-2,4-pyrimidinediamine:

In like manner to the preparation of N2, N4-bis (3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoropyrimidine and 1-methyl-5-aminoindazole were reacted to produce N2, N4-bis(1-methyl-indazolin-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.65 (s, 1H), 10.41 (s, 1H), 8.29 (d, 1H, J= 5.3 Hz), 7.98 (s, 1H), 7.79 (d, 2H, J= 9.4 Hz), 7.69-7.54 (m, 4H), 7.35 (dd, 1H, J= 1.7 and 9.4 Hz), 4.03 (s, 3H), 4.01 (s, 3H). LCMS: ret. time: 16.86 min.; purity: 99%; MS (m/e): 389 (MH⁺).

7.3.105 R935205: N2, N4-Bis[1-(methoxycarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N2, N4-bis (3-hydroxyphenyl)-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-1-(methoxycarbonyl)methyl-indazoline were reacted to produce N2, N4-bis[1-(methoxycarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.59 (s, 1H), 9.45 (s, 1H), 8.18 (d, 1H, J= 3.5 Hz), 8.11 (s, 1H), 8.04 (s, 1H), 7.95 (s, 1H), 7.93 (s, 1H), 7.69 (d, 1H, J= 8.8 Hz), 7.58 (d, 1H, J= 8.8 Hz), 7.48 (dd, 1H, J= 1.7 and 8.8 Hz), 7.32 (d, 1H, J= 8.8 Hz), 5.17 (s, 2H), 4.88 (s, 1H), 3.58 (s, 3H), 3.58 (s, 3H). LCMS: ret. time: 17.80 min.; purity: 99%; MS (m/e): 505 (MH⁺).

7.3.106 R935211: N2, N4-Bis[1-(methoxycarbonyl)methyl-indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 6-amino-1-(methoxycarbonyl)methyl-indazoline was reacted to produce N2, N4-bis[1-(methoxycarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR

(DMSO-d₆): δ 9.37 (s, 1H), 9.17 (s, 1H), 8.11-8.06 (m, 3H), 7.94 (s, 1H), 7.70 (s, 1H), 7.63 (s, 2H), 7.46 (s, 2H), 5.40 (s, 2H), 5.31 (s, 2H), 3.67 (s, 3H), 3.64 (s, 3H). LCMS: ret. time: 17.06 min.; purity: 96%; MS (*m/e*): 505 (MH⁺).

7.3.107 R935188: N₂,N₄-Bis(indazolin-6-yl)-5-fluoro-2,4-pyrimidinediamine:

5

In like manner to the preparation of 5-fluoro-N₂, N₄-bis (3-hydroxyphenyl)-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 6-aminoindazoline were reacted to produce N₂,N₄-bis(indazolin-6-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.80 (s, 1H), 9.65 (s, 1H), 8.20 (d, 1H, J= 4.1 Hz), 8.01 (s, 1H), 7.96 (s, 1H), 7.93 (s, 1H), 7.89 (s, 1H), 7.69 (d, 1H, J= 8.8 Hz), 7.57 (d, 1H, J= 8.3 Hz), 7.54 (dd, 1H, J= 1.7 and 8.8 Hz), 7.29 (dd, 1H, J= 1.7 and 8.8 Hz); LCMS: ret. time: 15.17 min.; purity: 94%; MS (*m/e*): 361 (MH⁺).

10

7.3.108 R935189: N₂, N₄-Bis(indazolin-5-yl)-5-fluoro-2,4-pyrimidinediamine:

15

In like manner to the preparation of N₂, N₄-bis (3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 5-aminoindazole were reacted to produce N₂, N₄-bis(indazolin-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.05 (s, 1H), 9.76 (s, 1H), 8.16 (d, 1H, J= 4.7 Hz), 8.05 (s, 1H), 7.92 (s, 1H), 7.82 (s, 1H), 7.68 (s, 1H), 7.52-7.52 (m, 2H), 7.44 (d, 1H, J= 8.8 Hz), 7.34 (dd, 1H, J= 1.7 and 8.8 Hz); LCMS: ret. time: 14.33 min.; purity: 100%; MS (*m/e*): 361 (MH⁺).

20

7.3.109 N₂,N₄-Bis(1-ethoxycarbonyl-2-methylpropyl)-5-cyano-2,4-pyrimidinediamine (R925814)

In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-cyanopyrimidine and valine ethyl ester were reacted to yield N₂,N₄-bis(1-ethoxycarbonyl-2-methylpropyl)-5-cyano-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.15 (s, 1H), 6.10 (d, 1H, J= 8.4 Hz), 5.67 (d, 1H, J= 8.1 Hz), 4.66-4.62 (m, 1H), 4.50-4.46 (m, 1H), 4.25-4.13 (m, 4H), 2.27-2.14 (m, 2H), 1.31-1.24 (m, 6H), 1.00-0.94 (m, 12H); LCMS: ret. time: 30.41 min.; purity: 98 %; MS (*m/e*): 392 (MH⁺).

25

7.3.110 N2,N4-Bis(1-methoxycarbonyl-3-methylbutyl)-5-cyano-2,4-pyrimidinediamine (R925815)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-cyanopyrimidine and leucine methyl ester were reacted to yield N2,N4-bis(1-methoxycarbonyl-3-methylbutyl)-5-cyano-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): mixture of rotamers δ 8.15 (s, 1H), 6.10 and 5.49 (2d, 1H, J= 8.1 Hz), 5.53 (d, 1H, J= 8.4 Hz), 4.80-4.67 (m, 1H), 4.57-4.48 (m, 1H), 3.73 (s, 3H), 3.71 (s, 3H), 1.78-1.60 (m, 6H), 0.97-0.89 (m, 12H); LCMS: ret. time: 30.33 min.; purity: 91 %; MS (m/e): 392 (MH⁺).

7.3.111 N2,N4-Bis(methoxycarbonylbenzyl)-5-cyano-2,4-pyrimidinediamine (R925819)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-cyanopyrimidine and phenyl glycine methyl ester were reacted to yield N2,N4-bis(methoxycarbonylbenzyl)-5-cyano-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): mixture of rotamers δ 8.15 (s, 1H), 7.69-7.60 (m, 1H), 7.42-7.32 (m, 10H), 6.20 and 5.73 (2d, 1H, J= 6.6 Hz), 6.14 and 5.65 (2d, 1H, J= 6.3 Hz), 5.55 (d, 1H, J= 6.3 Hz), 5.39 (t, 1H, J= 7.2 Hz), 3.79 and 3.78 (2s, 3H), 3.67 and 3.65 (2s, 3H); LCMS: ret. time: 30.22 min.; purity: 91 %; MS (m/e): 432 (MH⁺).

7.3.112 N2,N4-Bis[4-(ethoxycarbonylmethyl)phenyl]-5-cyano-2,4-pyrimidinediamine (R926662)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-cyanopyrimidine and ethyl 4-aminophenylacetate were reacted to yield N2,N4-bis[4-(ethoxycarbonylmethyl)phenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.29 (bs, 1H), 7.46 (2d, 4H, J= 7.8 Hz), 7.28 (d, 2h, J= 8.1 Hz), 7.19 (d, 2H, J= 8.1 Hz), 4.16 (2q, 4H, J= 6.3 Hz), 3.64 (s, 2H), 3.59 (s, 2H), 1.30-1.23 (m, 6H); LCMS: ret. time: 29.29 min.; purity: 93%; MS (m/e): 461 (MH⁺).

7.3.113 R935000: N2,N4-Bis(2-methoxy-5-phenylphenyl)-5-methyl-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N2,N4-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine, 5-phenyl-2-anisidine and 2,4-dichloro-5-methylpyrimidine were reacted to provide N2,N4-bis(2-methoxy-5-phenylphenyl)-5-methyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃ + CD₃OD): δ 7.76 (d, 1H, J= 2.3 Hz), 7.57 (s, 1H), 7.56 (s, 1H), 7.02-6.85

(m, 8H), 6.86-6.80 (m, 4H), 6.72 (d, 2H), 3.73 (s, 3H), 3.72 (s, 3H), 2.07 (s, 3H); LCMS: ret. time: 31.53 min.; purity: 97%; MS (*m/e*): 489 (MH⁺).

7.3.114 R935001: N₂,N₄-Bis[(2-methyl-5-phenyl)phenyl]-5-methyl-2,4-pyrimidinediamine

5 In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-phenyl-2-toluidine and 2,4-dichloro-5-methylpyrimidine were reacted to produce N₂,N₄-bis[(2-methyl-5-phenyl)phenyl]-5-methyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.59-7.55 (m, 1H), 7.45 (d, 2H, J= 3.6 Hz), 7.26-7.17 (m, 6H), 7.09-6.98 (m, 8H), 2.36 (s, 3H), 2.22 (s, 3H), 2.21 (s, 3H); LCMS: ret. time: 32.44 min.; purity: 90%;
10 MS (*m/e*): 457 (MH⁺).

7.3.115 R935002: N₂,N₄-Bis[(4-methoxy-3-phenyl)phenyl]-5-methyl-2,4-pyrimidinediamine

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 3-phenyl-4-anisidine hydrochloride and 2,4-dichloro-5-methylpyrimidine with an added diisopropylethylamine were reacted to produce N₂,N₄-bis[(4-methoxy-3-phenyl)phenyl]-5-methyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.15 (d, 1H, J= 2.3 Hz), 7.76 (t, 1H, J= 2.3 Hz), 7.71 (s, 1H), 7.59 (s, 1H), 7.16-7.03 (m, 8H), 6.98-6.81 (5H), 3.96 (s, 3H), 3.89 (s, 3H), 2.21 (s, 3H); LCMS: ret. time: 32.01 min.; purity: 90%; MS (*m/e*): 489 (MH⁺).

7.3.116 R935003: N₂,N₄-Bis[(4-phenyl-2-methoxy-5-methyl)phenyl]-5-methyl-2,4-pyrimidinediamine

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 5-methyl-4-phenyl-2-anisidine and 2,4-dichloro-5-methylpyrimidine were reacted to produce N₂,N₄-bis[(4-phenyl-2-methoxy-5-methyl)phenyl]-5-methyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 9.25 (br s, 1H), 8.17 (s, 1H), 7.77 (t, 1H, J= 6.4 Hz), 7.66 (s, 2H), 7.43-7.25 (m, 10H), 6.79 (s, 2H), 3.91 (s, 3H), 3.85 (s, 3H), 2.20 (s, 3H), 2.09 (s, 3H), 2.02 (s, 3H); LCMS: ret. time: 31.10 min.; purity: 100%; MS (*m/e*): 517 (MH⁺).

7.3.117 R935004: N2,N4-Bis[[di-(4-methoxyphenyl)]methyl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 1,1-di(4-anisyl)methylamine and 2,4-dichloro-5-fluoropyrimidine were reacted to produce N2,N4-bis[[di-(4-methoxyphenyl)]methyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃ + CD₃OD): δ 7.91 (d, 1H, J= 2.3 Hz), 7.18 (d, 8H, J= 9.0 Hz), 6.85 (d, 8H, J= 9.0 Hz), 6.40 (d, 1H, J= 8.2 Hz), 5.39 (d, 1H, J= 7.1 Hz), 3.81 (s, 6H), 3.78 (s, 6H); LCMS: ret. time: 32.76 min.; purity: 95%; MS (*m/e*): 581 (MH⁺).

7.3.118 R935005: N2,N4-Bis(diphenylmethyl)-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 1,1-diphenyl methylamine and 2,4-dichloro-5-fluoropyrimidine were reacted to produce N2, N4-bis(diphenylmethyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.91 (d, 1H, J= 2.3 Hz), 7.39-7.25 (m, 20H), 6.51 (d, 1H, J= 8.2 Hz), 5.77 (d, 1H, J= 7.0 Hz); LCMS: ret. time: 33.46 min.; purity: 92%; MS (*m/e*): 461 (MH⁺).

7.3.119 R935006: N2,N4-Bis[di-(4-chlorophenyl)methyl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, benzhydrylamine and 2,4-dichloro-5-fluoropyrimidine were reacted to yield N2,N4-bis[di-(4-chlorophenyl)methyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃ + CD₃OD): δ 7.94 (d, 1H, J= 2.3 Hz), 7.40-7.20 (m, 16H), 6.46 (d, 1H, J= 8.2 Hz), 5.69 (d, 1H, J= 7.0 Hz); LCMS: ret. time: 32.83 min.; purity: 90%; MS (*m/e*): 599 (MH⁺).

7.3.120 R935016: N2,N4-Bis[1(*R*)-4-methoxyphenylethyl]-5-bromo-2,4-pyrimidineamine

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, (R)-(+)-1-(4-methoxyphenyl)ethylamine and 5-bromo-2,4-dichloropyrimidine were reacted to produce N2,N4-bis[1(*R*)-4-methoxyphenylethyl]-5-bromo-2,4-pyrimidineamine. ¹H NMR (CDCl₃): δ 7.81 (s, 1H), 7.25 (d, 4H, J= 8.4 Hz), 6.86 (app t, 4H, J= 8.4 and 8.7 Hz), 5.27-5.20 m (2H), 5.09 (dq, 1H, J= 6.4 and 7.0 Hz), 4.89 (dq, 1H, J= 6.4 and 7.0 Hz), 3.80 (s, 3H), 3.79 (s, 3H), 1.40 (d, 6H, J= 7.0 Hz).

7.3.121 R935075: N2, N4-Bis[3-(2-hydroxyethoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-(3-aminophenoxy)ethanol were reacted to produce N2,N4-bis[3-(2-hydroxyethoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.50 (br s, 1H), 9.35 (br s, 1H), 8.13 (d, 1H, J= 4.1 Hz), 7.44 (d, 1H, J= 7.6 Hz), 7.26-7.19 (m, 4H), 7.10 (t, 1H, J= 7.6 Hz), 6.65 (dd, 1H, J= 2.3 and 8.2 Hz), 6.50 (dd, 1H, J= 2.3 and 8.2 Hz), 5.0 (br s, 2H), 3.91 (t, 2H, J= 5.2 Hz), 3.85 (t, 2H, J= 5.2 Hz), 3.68 (qt, 2H, J= 5.2 Hz), 3.66 (qt, 2H, J= 5.2 Hz); LCMS: ret. time: 15.76 min.; purity: 97%; MS (*m/e*): 401 (MH⁺).

7.3.122 R935076: N2,N4-Bis[3-(2-methoxyethyl)oxyphenyl]-5-fluoro-2,4-pyrimidinediamine:

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-(2-methoxyethoxy)aniline were reacted to produce N2,N4-bis[3-(2-methoxyethyl)oxyphenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.96 (d, 1H, J= 2.9 Hz), 7.36 (t, 1H, J= 1.7 Hz), 7.28 (t, 1H, J= 1.7 Hz), 7.25-7.06 (m, 4H), 6.98 (br s, 1H), 6.75 (d, 1H, J= 2.3 Hz), 6.70 (dd, 1H, J= 1.7 and 8.2 Hz), 6.58 (dd, 1H, J= 1.7 and 8.2 Hz), 4.08-4.03 (m, 4H), 3.74-3.69 (m, 4H), 3.44 (s, 3H), 3.43 (s, 3H); LCMS: ret. time: 21.01 min.; purity: 97%; MS (*m/e*): 429 (MH⁺).

7.3.123 R935077: N2,N4-Bis(5-hydroxy-2-isopropylphenyl)-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 3-amino-4-isopropylphenol and 2,4-dichloro-5-fluoropyrimidine were reacted to produce N2,N4-bis(5-hydroxy-2-isopropylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.93 (d, 1H, J= 3.5 Hz), 7.79 (br s, 1H), 7.64 (br s, 1H), 7.13 (d, 1H, J= 8.7 Hz), 7.06 (d, 1H, J= 2.3 Hz), 7.05 (d, 1H, J= 8.7 Hz), 6.89 (d, 1H, J= 2.3 Hz), 6.66 (d, 1H, J= 2.3 and 8.7 Hz), 6.57 (d, 1H, J= 2.3 and 8.7 Hz), 2.96 (m, 2H), 1.25 (d, 6H, J= 7.0 Hz), 1.13 (dd, 6H, J= 7.0 Hz); LCMS: ret. time: 24.27 min.; purity: 97%; MS (*m/e*): 397 (MH⁺).

7.3.124 R935114: N₂,N₄-Bis(3-methoxycarbonylmethylenephényl)-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-

5 (methoxycarbonylmethylene)aniline were reacted to produce the desired N₂,N₄-bis(3-methoxycarbonylmethylenephényl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.23 (br s, 1H), 10.05 (br s, 1H), 8.26 (d, 1H, J= 4.6 Hz), 7.64 (d, 1H, J= 8.2 Hz), 7.51 (br s, 1H), 7.46 (d, 1H, J= 8.2 Hz), 7.33 (br s, 1H), 7.29 (t, 1H, J= 7.6 Hz), 7.20 (t, 1H, J= 7.6 Hz), 7.06 (d, 1H, J= 7.6 Hz), 6.93 (d, 1H, J= 7.6 Hz), 3.63 (s, 2H), 3.58 (s, 3H), 3.57 (s, 10 3H), 3.56 (s, 2H); LCMS: ret. time: 21.74 min.; purity: 92%; MS (*m/e*): 425 (MH⁺).

7.3.125 R935162: N₂, N₄-Bis(3,4-propylenedioxyphenyl)- 5-fluoro-2,4-pyrimidinediamine:

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and (3,4-propylenedioxy)aniline were
15 reacted to give N₂,N₄-bis(3,4-propylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.18 (s, 1H), 9.07 (s, 1H), 8.03 (d, 1H, J= 3.5 Hz), 7.38 (dd, 1H, J= 2.3 and 8.2 Hz), 7.35 (d, 1H, J= 2.3 Hz), 7.33 (d, 1H, J= 2.3 Hz), 7.18 (dd, 1H, J= 2.3 and 8.8 Hz), 6.90 (d, 1H, J= 8.8 Hz), 6.80 (d, 1H, J= 8.2 Hz), 4.11-3.98 (m, 8H), 2.09-2.01 (m, 4H); LCMS: ret. time: 21.40 min.; purity: 97%; MS (*m/e*): 425 (MH⁺).

20 **7.3.126 R935163: N₂,N₄-Bis(3-chloro-4-fluorophényl)-2,4-pyrimidinediamine:**

In like manner to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-chloro-4-fluoroaniline were reacted to produce N₂, N₄-bis(3-chloro-4-fluorophényl)-2,4-pyrimidinediamine. ¹H NMR
25 (DMSO-d₆): δ 9.58 (s, 1H), 9.48 (s, 1H), 8.17 (d, 1H, J= 4.1 Hz), 7.94-7.90 (m, 2H), 7.73-7.67 (m, 1H), 7.51-7.45 (m, 1H), 7.38 (t, 1H, J= 8.8 Hz), 7.26 (t, 1H, J= 8.8 Hz); LCMS: ret. time: 27.83 min.; purity: 99%; MS (*m/e*): 386 (MH⁺).

7.3.127 N₂,N₄-Bis(3-hydroxyphenyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine (R925849)

30 In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-6-ethoxycarbonyl-5-nitropyrimidine and 3-aminophenol were reacted to yield N₂,N₄-bis(3-hydroxyphenyl)-6-ethoxycarbonyl-5-nitro-2,4-

pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.56 (bs, 1H), 10.32 (bs, 1H), 9.54 (s, 1H), 9.32 (bs, 1H), 7.22-7.15 (m, 2H), 7.02-6.96 (m, 1H), 6.93-6.82 (m, 2H), 6.81-6.74 (m, 1H), 6.67 (d, 1H, J= 9.3 Hz), 6.43 (d, 1H, J= 8.1 Hz), 4.35 (q, 2H, J= 6.9 Hz), 1.30 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 26.01 min.; purity: 96 %; MS (m/e): 412 (MH⁺).

5 **7.3.128 N2,N4-Bis(3,4-ethylenedioxyphenyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine (R925852)**

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-6-ethoxycarbonyl-5-nitropyrimidine and 3,4-ethylenedioxyaniline were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.52 (s, 1H), 10.28 (s, 1H), 7.07-7.01 (m, 2H), 6.96 (dd, 1H, J= 1.8 and 8.7 Hz), 6.90-6.84 (m, 2H), 6.61 (d, 1H, J= 8.7 Hz), 4.33 (q, 2H, J= 6.9 Hz), 4.24 (s, 4H), 4.17 (s, 4H), 1.29 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 30.40 min.; purity: 100 %; MS (m/e): 496 (MH⁺).

15 **7.3.129 N2,N4-Bis(ethoxycarbonylmethyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine (R925864)**

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, with the addition of triethylamine, 2,4-dichloro-6-ethoxycarbonyl-5-nitropyrimidine and glycine ethyl ester hydrochloride were reacted to yield N2,N4-bis(ethoxycarbonylmethyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): mixture of rotamers δ 8.99 and 8.80 (2bs, 1H), 6.22 and 6.00 (2bs, 1H), 4.45 (t, 2H, J= 7.2 Hz), 4.31-4.21 (m, 6H), 4.14 (d, 2H, J= 5.1 Hz), 1.39 (t, 3H, J= 7.2 Hz), 1.34-1.28 (m, 6H); LCMS: ret. time: 26.06 min.; purity: 99 %; MS (m/e): 400 (MH⁺).

7.3.130 N2,N4-Bis[2-(4-hydroxyphenyl)ethyl]-2,4-pyrimidinediamine (R925790)

25 In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and tyramine were reacted to yield N2,N4-bis[2-(4-hydroxyphenyl)ethyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 11.56 (bs, 1H), 9.23 (s, 1H), 8.89 (bs, 1H), 7.92 (bs, 1H), 7.60 (d, 1H, J= 6.9 Hz), 6.99 (d, 4H, J= 8.1 Hz), 6.65 (d, 4H, J= 8.1 Hz), 6.00 (d, 1H, J= 7.2 Hz), 3.59-3.42 (m, 4H), 2.76-2.67 (m, 4H); LCMS: ret. time: 17.93 min.; purity: 95 %; MS (m/e): 351 (MH⁺).

7.3.131 N2,N4-Bis(2-phenylphenyl)-2,4-pyrimidinediamine (R925804)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 2-aminobiphenyl were reacted to yield N2,N4-bis(2-phenylphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.36 (d, 1H, J= 8.1 Hz), 7.97 (d, 1H, J= 5.7 Hz), 7.80 (d, 1H, J= 7.5 Hz), 7.50-7.21 (m, 15H), 7.12-7.05 (m, 1H), 6.91 (bs, 1H), 6.38 (bs, 1H), 6.07 (d, 1H, J= 6.0 Hz); LCMS: ret. time: 29.94 min.; purity: 100 %; MS (m/e): 415 (MH⁺).

7.3.132 N2,N4-Bis(2-methoxy-5-phenylphenyl)-2,4-pyrimidinediamine (R925805)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 5-phenyl-*ortho*-anisidine were reacted to yield N2,N4-bis(2-methoxy-5-phenylphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.88-7.84 (m, 2H), 7.82 (d, 1H, J= 6.9 Hz), 7.30-7.14 (m, 14H), 7.10 (dd, 2H, J= 3.0 and 8.1 Hz), 6.48 (d, 1H, J= 6.9 Hz), 3.93 (s, 3H), 3.92 (s, 3H); LCMS: ret. time: 30.09 min.; purity: 94 %; MS (m/e): 476 (MH⁺).

7.3.133 N2,N4-Bis(3-carboxy-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945041)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, from 5-amino-2-hydroxybenzoic acid (458 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N2,N4-bis(3-carboxy-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (235 mg, 98%). ¹H NMR (DMSO-d₆): δ 6.76 (d, J= 9.0 Hz, 1 H), 6.88 (d, J= 9.6 Hz, 1 H), 7.75 (dd, J= 3.0, 9.0 Hz, 1 H), 7.90-7.94 (m, 3 H), 8.02 (d, J= 3.9 Hz, 1 H), 9.04 (s, 1 H, NH), 9.28 (s, 1 H, NH); ¹⁹F NMR (282 MHz, DMSO-d₆): δ -165.79; LC: 16.02 min, 86.82%; MS (m/z): 400.94 (MH⁺).

7.3.134 N2,N4-Bis(4-methoxy-3-phenylphenyl)-2,4-pyrimidinediamine (R925806)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, with the addition of triethylamine, 2,4-dichloropyrimidine and 3-phenyl-*para*-anisidine hydrochloride were reacted to yield N2,N4-bis(4-methoxy-3-phenylphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.93 (d, 1H, J= 2.4 Hz), 7.88 (d, 1H, J= 2.4 Hz), 7.29 (dd, 1H, J= 1.8 and 9.0 Hz), 7.26-7.18 (m, 13H), 7.10 (d, 2H, J= 8.7

Hz), 6.46 (d, 1H, J= 7.2 Hz), 3.93 (s, 3H), 3.92 (s, 3H); LCMS: ret. time: 29.99 min.; purity: 92%; MS (m/e): 476 (MH⁺).

7.3.135 N₂,N₄-Bis(2-methyl-5-phenylphenyl)-2,4-pyrimidinediamine (R925807)

5 In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 5-phenyl-*ortho*-toluidine were reacted to yield N₂,N₄-bis(2-methyl-5-phenylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.45 (bs, 1H), 10.01 (bs, 1H), 7.86 (bs, 1H), 7.69-7.22 (m, 17H), 2.28 (s, 6H); LCMS: ret. time: 18.69 min.; purity: 98 %; MS (m/e): 443 (MH⁺).

10 **7.3.136 N₂,N₄-Bis(2-methoxy-5-methyl-4-phenylphenyl)-2,4-pyrimidinediamine (R925808)**

In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 5-methyl-4-phenyl-*ortho*-anisidine were reacted to yield N₂,N₄-bis(2-methoxy-5-methyl-4-phenylphenyl)-2,4-pyrimidinediamine.
15 ¹H NMR (DMSO-d₆): δ 9.99 (bs, 1H), 9.22 (bs, 1H), 7.98 (d, 1H, J= 6.3 Hz), 7.75 (s, 1H), 7.59 (s, 1H), 7.46-7.29 (m, 10H), 6.92 (s, 1H), 6.87 (s, 1H), 6.49 (d, 1H, J= 5.4 Hz), 3.82 (s, 3H), 3.81 (s, 3H), 2.07 (s, 3H), 1.98 (s, 3H); LCMS: ret. time: 19.69 min.; purity: 93 %; MS (m/e): 503 (MH⁺).

20 **7.3.137 N₂,N₄-Bis[4-(ethoxycarbonylmethyleneoxy)phenyl]-5-trifluoromethyl-2,4-pyrimidinediamine (R925862)**

In a manner similar to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-trifluoromethylpyrimidine and ethyl 4-aminophenoxyacetate were reacted to yield N₂,N₄-bis[4-(ethoxycarbonylmethyleneoxy)phenyl]-5-trifluoromethyl-2,4-pyrimidinediamine. ¹H NMR
25 (DMSO-d₆): δ 9.64 (bs, 1H), 8.80 (bs, 1H), 8.29 (s, 1H), 7.36 (d, 2H, J= 8.1 Hz), 7.31 (d, 2H, J= 9.3 Hz), 6.93 (d, 2H, J= 8.7 Hz), 6.70 (d, 2H, J= 9.0 Hz), 4.80 (s, 2H), 4.67 (s, 2H), 4.18 (q, 2H, J= 6.9 Hz), 4.15 (q, 2H, J= 6.9 Hz), 1.20 (t, 3H, J= 6.9 Hz), 1.19 (t, 3H, J= 6.9 Hz); ¹⁹F NMR (DMSO-d₆): -16932; LCMS: ret. time: 26.33 min.; purity: 98 %; MS (m/e): 535 (MH⁺).

7.3.138 N2,N4-Bis(3-hydroxyphenyl)-5-trifluoromethyl-2,4-pyrimidinediamine (R925863)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-trifluoromethylpyrimidine and 3-aminophenol were reacted to yield N2,N4-bis(3-hydroxyphenyl)-5-trifluoromethyl-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.82 (bs, 1H), 8.88 (bs, 1H), 8.36 (s, 1H), 7.18-7.11 (m, 2H), 6.96 (m, 4H), 6.63 (dd, 1H, J= 2.4 and 8.1 Hz), 6.38 (d, 1H, J= 8.1 Hz); ¹⁹F NMR (DMSO-d₆): -16979; LCMS: ret. time: 19.04 min.; purity: 95 %; MS (m/e): 363 (MH⁺).

7.3.139 N2,N4-Bis[4-(ethoxycarbonylmethyl)phenyl]-5-trifluoromethyl-2,4-pyrimidinediamine (R926663)

In a manner similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-trifluoromethylpyrimidine and ethyl 4-aminophenylacetate were reacted to yield N2,N4-bis[4-(ethoxycarbonylmethyl)phenyl]-5-trifluoromethyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.31 (s, 1H), 7.46 (d, 2H, J= 9.0 Hz), 7.45 (d, 2H, J= 8.7 Hz), 7.30 (d, 2H, J= 9.0 Hz), 7.18 (d, 2H, J= 8.7 Hz), 7.16 (bs, 1H), 6.82 (bs, 1H), 4.16 (2q, 4H, J= 7.8 Hz), 3.64 (s, 2H), 3.57 (s, 2H), 1.27 (t, 3H, J= 7.8 Hz), 1.26 (t, 3H, J= 7.8 Hz); ¹⁹F NMR (CDCl₃): -17223; LCMS: ret. time: 28.07 min.; purity: 99 %; MS (m/e): 504 (MH⁺).

7.3.140 N2,N4-Bis(2,5-dimethyl-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926623)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2,5-dimethyl-4-hydroxyaniline were reacted to yield N2,N4-bis(2,5-dimethyl-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.63 (d, 1H, J= 4.2 Hz), 7.05 (s, 1H), 6.97 (s, 1H), 6.64 (1H), 6.54 (s, 1H), 2.12 (s, 6H), 2.06 (s, 3H), 2.03 (s, 3H), 2.03 (s, 3H); ¹⁹F NMR (CD₃OD): -48488; LCMS: ret. time: 18.28; purity: 94%; MS (m/e): 369 (MH⁺).

7.3.141 N2,N4-Bis(3-sodiumphenoxy)-5-fluoro-2,4-pyrimidinediamine (R926461)

The reaction of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine with 2 equivalents of sodium methoxide in methanol followed by removal of solvent gave the requisite compound, N2,N4-bis(3-sodiumphenoxy)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (D₂O): δ 7.65 (bd, 1H), 7.00-6.90 (m, 2H), 6.71 (m, 2H), 6.55 (dd, 1H, J= 1.2 and 6.3

Hz), 6.31 (bd, 1H, J= 8.1 Hz), 6.23 (bd, 1H, J= 8.7 Hz); ^{19}F NMR (D_2O): -47016; LCMS: ret. time: 15.68 min.; purity: 99%; MS (m/e): 313 (MH^+).

7.3.142 N2,N4-Bis(3-cyanophenyl)-5-fluoro-2,4-pyrimidinediamine (R945051)

5 In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 3-aminobenzonitrile (177 mg, 1.5 mmol) and 2,4-dichloro-5-fluoropyrimidine (50 mg, 0.3 mmol) gave N2,N4-bis(3-cyanophenyl)-5-fluoro-2,4-pyrimidinediamine (75 mg, 76%). ^1H NMR (acetone- d_6): δ 7.33 (dt, J= 1.8, 7.8 Hz, 1 H), 7.46-7.52 (m, 2 H), 7.59 (t, J= 7.8 Hz, 1 H), 7.90 (ddd, J= 0.9, 2.1 and 8.4 Hz, 1 H), 8.09
10 (ddd, J= 1.2, 2.4 and 8.4 Hz, 1 H), 8.17 (d, J= 3.3 Hz, 1 H), 8.31 (m, 1 H), 8.35 (t, J= 2.1 Hz, 1 H), 8.98 (br, 1 H, NH), 9.02 (br, 1 H, NH); ^{19}F NMR (282 MHz, acetone- d_6): δ -165.80; LCMS: 24.64 min.; purity: 98.02%; MS (m/e): 331.01 (MH^+).

7.3.143 N2,N4-Bis(benzothiophen-3-ylmethyl)-5-fluoro-2,4-pyrimidinediamine (R945145)

15 Using procedure similar to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, benzothiophen-3-ylmethylamine and 2,4-dichloro-5-fluoropyrimidine gave N2,N4-bis(benzothiophen-3-ylmethyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 4.82 (dd, J= 0.9 and 5.7 Hz, 2 H), 4.86 (dd, J= 0.9
and 5.7 Hz, 2 H), 5.14 (br, 2 H), 7.31-7.40 (m, 6 H), 7.75-7.89 (m, 5 H); ^{19}F NMR (282
20 MHz, CDCl_3): δ -172.12; LCMS: 27.79 min.; purity: 96.47%; MS (m/e): 420.92 (MH^+).

7.3.144 N2,N4-Bis[4-(N-benzylpiperazino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R945152)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 4-(N-benzylpiperazino)aniline (400 mg, 1.5 mmol)
25 and 2,4-dichloro-5-fluoropyrimidine (50 mg, 0.3 mmol) resulted N2,N4-bis[4-(N-benzylpiperazino)phenyl]-5-fluoro-2,4-pyrimidinediamine (120 mg, 64%). ^1H NMR (CDCl_3): δ 2.63 (p, J= 2.4 Hz, 8 H), 3.14 (t, J= 4.8 Hz, 4 H), 3.19 (t, J= 4.8 Hz, 4 H), 3.58 (s, 4 H), 6.58 (d, 1 H, NH), 6.67 (br, 1 H, NH), 6.87 (d, J= 9.3 Hz, 2 H), 6.90 (d, J= 9.0 Hz, 2 H), 7.33-7.39 (m, 12 H), 7.46 (d, J= 9.0 Hz, 2 H), 7.87 (d, J= 3.3 Hz, 1 H); ^{19}F NMR (282
30 MHz, CDCl_3): δ -169.06; LCMS: 16.82 min.; purity: 96.88%; MS (m/e): 629.12 (MH^+).

7.3.145 N2,N4-Bis(3-hydroxy-2-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R945038)

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 3-hydroxy-2-methylaniline (369 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N2,N4-bis(3-hydroxy-2-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (180 mg, 88%). ¹H NMR (acetone-*d*₆): δ 2.14 (s, 3 H), 2.22 (s, 3 H), 6.61 (d, J= 8.1 Hz, 1 H), 6.78 (t, J= 8.7 Hz, 1 H), 6.87 (d, J= 7.8 Hz, 1 H), 6.99 (d, J= 9.0 Hz, 1 H), 7.08 (t, J= 7.8 Hz, 1 H), 7.13 (dd, J= 3.9, 8.4 Hz, 1 H), 8.24 (d, J= 5.1 Hz, 1 H), 8.32 (br, 1 H, NH), 8.57 (br, 1 H, NH); LCMS: ret. time: 16.51 min.; purity: 90.47%; MS (m/e): 341.07 (MH⁺).

7.3.146 N2,N4-Bis(3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950160)

2,4-Dichloro-5-fluoropyrimidine (4.7 g, 28.1 mmol) was dissolved in a mixture of MeOH (150 ml) and H₂O (15 ml). 3-nitroaniline (15.5 g, 112 mmol) was added and the mixture was refluxed for 20 hours (100 °C oil-bath temperature). The mixture was cooled to 22 °C and filtered. The residue was washed carefully with 200 ml MeOH-H₂O (1:1; v/v) and dried under vacuum to give 7.89 g (76%) of N2,N4-bis(3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine as yellow crystals. ¹H NMR (DMSO-*d*₆ + D₂O): δ 8.63 (m, 2H), 8.21 (m, 1H), 8.08 (d, 1H, J= 8.41 Hz), 7.88 (d, 1H, J= 8.4 Hz), 7.79 (d, 1H, J= 8.4 Hz), 7.70 (d, 1H, J= 8.4 Hz), 7.57 (d, 1H, J= 8.4 Hz), 7.45 (t, 1H, J= 8.4 Hz); LCMS: purity: 100%; MS (m/e): 371.30 (M⁺, 100).

7.3.147 N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine (R921302)

N2,N4-Bis(3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (4.0 g, 10.8 mmol) and Pd/C 10% (1.2 g, 50% water content) were suspended in 300 ml EtOH-10% aqueous HCl (1 : 1) and hydrogenated in a Parr apparatus for 6 hours (22 °C, 50 psi). The suspension was filtered over celite and carefully washed with 20 ml DMF-H₂O (1:1; v/v) followed by 50 ml H₂O. The combined filtrates were concentrated under reduced pressure to give pale yellow oil, which was triturated with MeOH to give the product as fine white needles. The precipitate was filtered off and washed with MeOH followed by Et₂O. The remaining crystals were dried under vacuum to give 4.00 g of pure material (100%) as determined by LCMS. The free amine was obtained by adding 10 ml 1 N NaOH to a solution of 1g HCl-

salt in 5 ml H₂O. The resulting precipitate was filtered, washed with H₂O and dried under vacuum for 24 hours to give N₂,N₄-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine (770 mg) as a white solid. ¹H NMR (CD₃OD): δ 7.92 (d, 1H, J= 3.6 Hz), 7.31 (t, 1H, J= 2.1 Hz), 7.21 (t, 1H, J= 2.4 Hz), 7.08, (t, 1H, J= 8.1 Hz), 6.99 (t, 1H, J= 8.1 Hz), 6.88 (m, 1H),
5 6.77 (m, 1H), 6.47 (m, 1H), 6.34 (m, 1H); LCMS: purity: 100%; MS (m/e): 311.07 (M⁺, 100).

7.3.148 N₂,N₄-Bis(4-aminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950122)

In like manner to the preparation of N₂,N₄-bis(3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 1,4-diaminobenzene were reacted
10 to prepare N₂,N₄-bis(4-aminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 11.15 min.; purity: 100%; MS (m/e): 311.09 (MH⁺).

7.3.149 N₂,N₄-Bis[3-(dimethylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R950182)

2,4-Dichloro-5-fluoropyrimidine (50 mg, 0.30 mmol) was dissolved in a mixture of MeOH (0.3 ml) and H₂O (0.03 ml). N,N-3-dimethyldiaminoaniline (163 mg, 1.2 mmol) was added and the mixture was refluxed for 24 hours (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 2 : 1) to give N₂,N₄-bis[3-(dimethylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS purity: 99.0%; MS (m/e):
15 367.13 (M⁺, 100).
20

7.3.150 N₂,N₄-Bis(3-amino-4-methylphenyl)-2,4-pyrimidinediamine (R950130)

2,4-Dichloropyrimidine (45 mg, 0.30 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 3-amino-4-methylaniline (146 mg, 1.2 mmol) was added and the mixture was refluxed for 20 hours (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 2:1) to give N₂,N₄-bis(3-amino-4-methylphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.13 (s, 1H), 6.95 (d, 2H, J= 7.5 Hz), 6.82 (d, 2H, J= 1.8 Hz), 6.60 (dd, 2H, J= 1.8, 7.5 Hz), 6.17 (s, 1H), 2.12 (s, 6H);
25 LCMS purity: 97.3%; MS (m/e): 321.09 (M⁺, 100).
30

7.3.151 N₂,N₄-Bis(3-amino-4-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950129)

2,4-Dichloro-5-fluoropyrimidine (50 mg, 0.30 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 3-amino-4-methylaniline (146 mg, 1.2 mmol) was added and the mixture was refluxed for 20 hours (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 2:1) to give N₂,N₄-bis(3-amino-4-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.11 (d, 1H, J= 5.1 Hz), 7.98 (bs, 1H) (7.68 (dd, 1H, J= 2.4, 8.1 Hz), 7.40-7.55 (m, 4H), 2.43 (s, 3H), 2.42 (s, 3H); LCMS: purity: 95.0%; MS (m/e): 338.66 (M⁺, 70).

7.3.152 N₂,N₄-Bis[(4-methylsulfonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R950083)

2,4-Dichloro-5-fluoropyrimidine (50 mg, 0.30 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 4-methylsulfonylaminoaniline (335 mg, 1.8 mmol) was added and the mixture was refluxed for 24 hours (100 °C oil-bath temperature). The mixture was cooled to 22 °C and filtered. The residue was washed carefully with MeOH–H₂O (1:1) and dried under vacuum to give N₂,N₄-bis[(4-methylsulfonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.86 (s, 1H), 8.65 (s, 1H), 8.53 (bs, 1H), 8.39 (bs, 1H), 7.32 (d, 1H, J= 3.3 Hz), 7.12 (d, 1H, J= 8.7 Hz), 6.98 (d, 1H, J= 8.7 Hz), 6.62 (d, 1H, J= 8.7 Hz), 6.52 (d, 1H, J= 8.7 Hz), 2.32 (s, 3H), 2.27 (s, 3H); LCMS: purity: 96.8%; MS (m/e): 466.94 (M⁺, 100).

7.3.153 N₂,N₄-Bis(4-benzyloxy-3-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950090)

2,4-Dichloro-5-fluoropyrimidine (50 mg, 0.30 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 4-benzyloxy-3-trifluoromethylaniline (481 mg, 1.8 mmol) was added and the mixture was refluxed for 2 days (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 9:1) to give N₂,N₄-bis(4-benzyloxy-3-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.51 (s, 1H), 8.05 (s, 1H), 7.38-7.64 (m, 5H), 6.94-7.14 (m, 11H), 6.44-6.73 (m, 4H), 4.84 (s, 2H), 4.79 (s, 2H); LCMS purity: 94.7%; MS (m/e): 628.93 (M⁺, 100).

7.3.154 N2,N4-Bis(3-cyano-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950092)

2,4-Dichloro-5-fluoropyrimidine (50 mg, 0.30 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 3-cyano-4-hydroxyaniline (241 mg, 1.8 mmol) was added and the mixture was refluxed for 2 days (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 9:1) to give N2,N4-bis(4-hydroxy-3-cyanophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.96 (d, 1H, J= 3.5 Hz), 7.82 (d, 1H, J= 3.0 Hz), 7.79 (d, 1H, J= 3.0 Hz), 7.71 (dd, 1H, J= 3.0, 8.8 Hz), 7.54 (dd, J= 3.0, 8.8 Hz), 6.94 (d, 1H, J= 8.8 Hz), 6.84 (d, 1H, J= 8.8 Hz); LCMS: purity: 97.2%; MS (m/e): 362.98 (M⁺, 100).

7.3.155 N2,N4-Bis[3-methylsulfonylamino]phenyl]-5-fluoro-2,4-pyrimidinediamine (R950100)

2,4-Dichloro-5-fluoropyrimidine (50 mg, 0.3 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 3-methylsulfonylaminoaniline (300 mg, 1.5 mmol) was added and the mixture was refluxed for 24 hours (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 9:1) to give N2,N4-bis[3-methylsulfonylamino]phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD): δ 8.01 (d, 1H, J= 3.5 Hz), 7.46-7.68 (m, 4H), 7.49 (t, 1H, J= 8.2 Hz), 7.13 (t, 1H, J= 8.2 Hz), 6.89 (dd, 1H, J= 2.4, 8.2 Hz), 6.72 (m, 1H), 2.95 (s, 3H), 2.91 (s, 3H); LCMS: purity: 97.2%; MS (m/e): 466.89 (M⁺, 100).

7.3.156 N2,N4-Bis[3-(tert-butoxycarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R950108)

2,4-Dichloro-5-fluoropyrimidine (75 mg, 0.45 mmol) was dissolved in a mixture of MeOH (2 ml) and H₂O (0.2 ml). 3-tert-butoxycarbonylaminoaniline (374 mg, 1.8 mmol) was added and the mixture was refluxed for 40 hours (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 9:1) to give N2,N4-bis[3-(tert-butoxycarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD): δ 7.96 (d, 1H, J= 4.1 Hz), 7.83 (m, 1H), 7.60 (m, 1H), 7.34-7.42 (m, 2H), 7.15-

7.19 (m, 2H), 7.06 (t, 1H, J= 8.2 Hz), 6.93 (d, 1H, J= 8.2 Hz), 1.43 (s, 9H), 1.40 (s, 9H); LCMS: purity: 93.2%; MS (m/e): 511.06 (M^+ , 100).

7.3.157 N2,N4-Bis[4-(tert-butoxycarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R950120)

5 2,4-Dichloro-5-fluoropyrimidine (75 mg, 0.45 mmol) was dissolved in a mixture of MeOH (2 ml) and H₂O (0.2 ml). 4-tert-butoxycarbonylaminoaniline (374 mg, 1.8 mmol) was added and the mixture was refluxed for 24 hours (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃–Acetone, 9:1) to give N2,N4-bis[4-(tert-butoxycarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD): δ 7.96 (d, 1H, J= 3.5 Hz), 7.63 (d, 2H, J= 8.8 Hz), 7.49 (d, 2H, J= 8.8 Hz), 7.37 (d, 2H, J= 8.8 Hz), 7.24 (d, 2H, J= 8.8 Hz), 1.45 (s, 9H), 1.43 (s, 9H); LCMS: purity: 97.9%; MS (m/e): 511.04 (M^+ , 100).

7.3.158 N2,N4-Bis[2-[2-(methylamino)ethyleneaminocarbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine (R950170)

15 N2,N4-Bis[2-(ethoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine (10 mg, 0.02 mmol) was dissolved in EtOH. To this was added N-methyl-1,2-aminoethane (0.1 ml : 0.1 ml) and the mixture was refluxed for 3 days (70 °C oil-bath temperature). The mixture was cooled to 22 °C, diluted with water and filtered. The residue was subjected to column chromatography on silica gel (CHCl₃–Acetone, 2:1) to give N2,N4-bis[2-[2-(methylamino)ethyleneaminocarbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD): δ 8.14 (s, 1H), 8.02 (s, 1H), 7.99 (d, 1H, J= 2.4 Hz), 7.35-7.68 (m, 5H), 7.17 (s, 1H), 3.41 (m, 2H), 2.75 (m, 2H), 2.35 (s, 3H); LCMS: purity: 84.2%; MS (m/e): 561.08 (M^+ , 100).

7.3.159 N2,N4-Bis[2-(2-hydroxyethyleneamoinocarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine (R950167)

25 In like manner to the preparation of N2,N4-bis[2-[2-(methylamino)ethyleneamino carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis[2-(ethoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine and 2-aminoethanol
30 were reacted to prepare N2,N4-bis[2-(2-hydroxyethyleneamoinocarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 14.22 min.; purity: 95.7%; MS (m/e): 535.01 (MH^+).

7.3.160 N2,N4-Bis[2-(2-aminoethyleneamino)carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine (R950168)

In like manner to the preparation of N2,N4-bis[2-[2-(methylamino)ethyleneamino carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis[2-(ethoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine and 1,2-diaminoethane were reacted to prepare N2,N4-bis[2-(2-aminoethyleneamino)carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 13.15 min.; purity: 95.8%; MS (m/e): 532.99 (MH⁺).

7.3.161 N2,N4-Bis[2-(2-(N-benzylamino)ethyleneamino)carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine (R950169)

In like manner to the preparation of N2,N4-bis[2-[2-(methylamino)ethyleneamino carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis[2-(ethoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine and N-benzyl-1,2-diaminoethane were reacted to prepare N2,N4-bis[2-(2-(N-benzylamino)ethyleneamino)carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 13.15 min.; purity: 95.8%; MS (m/e): 713.10 (MH⁺).

7.3.162 N2,N4-Bis[2-(N-morpholinocarbonyl)benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine (R950172)

In like manner to the preparation of N2,N4-bis[2-[2-(methylamino)ethyleneamino carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis[2-(ethoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine and morpholine were reacted to N2,N4-bis[2-(N-morpholinocarbonyl)benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD): δ 8.13 (d, 1H, J= 2.7 Hz), 8.06 (d, 1H, J= 2.4 Hz), 8.03 (d, 1H, J= 3.6 Hz), 7.63 (dd, 1H, J= 2.4, 8.8 Hz), 7.57 (d, 1H, J= 9.3 Hz), 7.49 (dd, 1H, J= 2.4, 8.4 Hz), 7.42 (d, 1H, J= 8.8 Hz), 7.25 (s, 1H), 7.05 (s, 1H), 4.09 (m, 4H), 3.65 (m, 4H); LCMS: ret. time: 18.04 min.; purity: 83.2%; MS (m/e): 587.04 (MH⁺).

7.3.163 N2,N4-Bis[2-(2-N-morpholinoethyleneamino)carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine (R950173)

In like manner to the preparation of N2,N4-bis[2-[2-(methylamino)ethyleneamino carbonyl]-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis[2-(ethoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine and N-(2-aminoethyleneamino)morpholine were reacted to prepare N2,N4-bis[2-(2-N-

morpholinoethyleneamino carbonyl)-benzofuran-5-yl]-5-fluoro-2,4-pyrimidinediamine.
¹H NMR (DMSO-d₆ + CD₃OD): δ 8.16 (d, 1H, J = 2.4 Hz), 8.03-8.05 (m, 2H), 7.71 (dd, 1H, J = 1.8, 8.8 Hz), 7.56 (d, 1H, J = 8.8 Hz), 7.42 (d, 1H, J = 8.8 Hz), 7.36 (s, 1H), 7.19 (s, 1H), 4.19 (m, 4H), 3.38 (m, 4H), 3.16 (t, 2H, J = 6.3 Hz), 2.28 (t, 2H, J = 6.3 Hz); LCMS: 5 ret. time: 12.85 min.; purity: 93.8%; MS (m/e): 673.35 (MH⁺).

7.3.164 N₂,N₄-Bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950135)

2,4-Dichloro-5-fluoropyrimidine (50 mg, 0.3 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 3-amino-4-nitroaniline (184 mg, 1.2 mmol) was added and 10 the mixture was refluxed for 3 days (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃-Acetone, 2:1) to give N₂,N₄-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD): δ 8.21 (d, 1H, J = 2.9 Hz), 7.89 (m, 3H), 7.56 (d, 1H, J = 2.3 Hz), 7.01 (m, 1H), 6.81 (dd, 1H, J = 2.3, 15 9.4 Hz); LCMS: purity: 91.1%; MS (m/e): 401.00 (M⁺, 100).

7.3.165 N₂,N₄-Bis(3-amino-2,4-difluorophenyl)-5-fluoro-2,4-pyrimidinediamine (R950138)

In like manner to the preparation of N₂,N₄-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-amino-2,4-difluoroaniline were 20 reacted to prepare N₂,N₄-bis(3-amino-2,4-difluorophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 16.98 min.; purity: 91.7%; MS (m/e): 382.97 (MH⁺).

7.3.166 N₂,N₄-Bis(3-amino-4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950139)

In like manner to the preparation of N₂,N₄-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-amino-4-ethoxyaniline were 25 reacted to prepare N₂,N₄-bis(3-amino-4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 14.29 min.; purity: 93.4%; MS (m/e): 399.09 (MH⁺).

7.3.167 N₂,N₄-Bis(3-amino-5-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950134)

In like manner to the preparation of N₂,N₄-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-amino-5- 30

methoxycarbonylaniline were reacted to prepare N2,N4-bis(3-amino-5-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 14.72 min.; purity: 93.8%; MS (m/e): 427.02 (MH⁺).

7.3.168 N2,N4-Bis(3-amino-5-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950140)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-amino-5-trifluoromethylaniline were reacted to prepare N2,N4-bis(3-amino-5-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 23.35 min.; purity: 100%; MS (m/e): 446.92 (MH⁺).

7.3.169 N2,N4-Bis(3-amino-5-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R950141)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-amino-5-chloroaniline were reacted to prepare N2,N4-bis(3-amino-5-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine.

LCMS: ret. time: 19.25 min.; purity: 99.3%; MS (m/e): 378.91 (MH⁺).

7.3.170 N2,N4-Bis(4-hydroxy-3-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950093)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-hydroxy-3-trifluoromethylaniline were reacted to prepare N2,N4-bis(4-hydroxy-3-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 22.06 min.; purity: 99.1%; MS (m/e): 448.88 (MH⁺).

7.3.171 N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine Hydrogen Chloride salt (R950107)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine was treated with 2 equivalents of HCl in dioxane. The volatiles were removed under reduced pressure to give N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine hydrogen chloride salt. LCMS: ret. time: 9.74 min.; purity: 91.3%; MS (m/e): 311.06 (MH⁺).

7.3.172 N2,N4-Bis(4-aminophenyl)-5-fluoro-2,4-pyrimidinediamine Hydrogen Chloride Salt (R950121)

N2,N4-Bis(4-aminophenyl)-5-fluoro-2,4-pyrimidinediamine was treated with 2 equivalents of HCl in dioxane. The volatiles were removed under reduced pressure to give
5 N2,N4-bis(4-aminophenyl)-5-fluoro-2,4-pyrimidinediamine Hydrogen Chloride Salt.
LCMS: ret. time: 11.15 min.; purity: 100%; MS (m/e): 311.09 (MH⁺).

7.3.173 N2,N4-Bis(3-aminophenyl)-2,4-pyrimidinediamine (R950109)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 3-aminoaniline were reacted to prepare
10 N2,N4-bis(3-aminophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 8.90 min.; purity: 91%; MS (m/e): 293.06 (MH⁺).

7.3.174 N2,N4-Bis(3-amino-2,4-difluorophenyl)-2,4-pyrimidinediamine (R950131)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 3-amino-2,4-difluoroaniline were reacted to
15 prepare N2,N4-bis(3-amino-2,4-difluorophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 16.62 min.; purity: 96.7%; MS (m/e): 364.99 (MH⁺).

7.3.175 N2,N4-Bis(3-amino-4-ethoxyphenyl)-2,4-pyrimidinediamine (R950142)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 3-amino-4-ethoxyaniline were reacted to
20 prepare N2,N4-bis(3-amino-4-ethoxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 14.38 min.; purity: 99.7%; MS (m/e): 381.07 (MH⁺).

7.3.176 N2,N4-Bis(3-amino-5-methoxycarbonylphenyl)-2,4-pyrimidinediamine (R950132)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 3-amino-5-methoxycarbonylaniline were
25 reacted to prepare N2,N4-bis(3-amino-5-methoxycarbonylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 15.25 min.; purity: 93.6%; MS (m/e): 409.02 (MH⁺).

7.3.177 N2,N4-Bis(3-amino-5-trifluoromethylphenyl)-2,4-pyrimidinediamine (R950143)

5 In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 3-amino-5-trifluoromethylaniline were reacted to prepare N2,N4-bis(3-amino-5-trifluoromethylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 23.23 min.; purity: 99.1%; MS (m/e): 428.95 (MH⁺).

7.3.178 N2,N4-Bis(3-amino-5-chlorophenyl)-2,4-pyrimidinediamine (R950133)

10 In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 3-amino-5-chloroaniline were reacted to prepare N2,N4-bis(3-amino-5-chlorophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 19.45 min.; purity: 100%; MS (m/e): 360.93 (MH⁺).

7.3.179 N2,N4-Bis[3-amino-4-(N-phenylamino)-phenyl]-5-fluoro-2,4-pyrimidinediamine (R950125)

15 In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-amino-4-(N-phenylamino)-aniline were reacted to prepare N2,N4-bis[3-amino-4-(N-phenylamino)-phenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 23.67 min.; purity: 100%; MS (m/e): 476.36 (MH⁺).

20 7.3.180 N2,N4-Bis[3-amino-4-(N-phenylamino)-phenyl]-2,4-pyrimidinediamine (R950123)

In like manner the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 3-amino-4-(N-phenylamino)-aniline were reacted to prepare N2,N4-bis[3-amino-4-(N-phenylamino)-phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 23.77 min.; purity: 77.8%; MS (m/e): 475.04 (MH⁺).

25 7.3.181 N2,N4-Bis(5-amino-2-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950157)

In like manner to the preparation of N2,N4-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 5-amino-2-methylaniline were reacted to prepare N2,N4-bis(5-amino-2-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 10.61 min.; purity: 83.4%; MS (m/e): 339.13 (MH⁺).

7.3.182 N₂,N₄-Bis(5-amino-2-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine (R950158)

In like manner to the preparation of N₂,N₄-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 5-amino-2-fluoroaniline were reacted to prepare N₂,N₄-bis(5-amino-2-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 11.48 min.; purity: 95.6%; MS (m/e): 347.04 (MH⁺).

7.3.183 N₂,N₄-Bis(3-amino-4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine (R950159)

In like manner to the preparation of N₂,N₄-bis(3-amino-4-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-amino-4-fluoroaniline were reacted to prepare N₂,N₄-bis(3-amino-4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 18.74 min.; purity: 95.6%; MS (m/e): 347.29 (MH⁺).

7.3.184 N₂,N₄-Bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950146)

2,4-Dichloro-5-fluoropyrimidine (33 mg, 0.2 mmol) was dissolved in a mixture of MeOH (1 ml) and H₂O (0.1 ml). 2-Methyl-5-nitroaniline (122 mg, 0.8 mmol) was added and the mixture was refluxed for 2 days (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃-Acetone, 9:1) to give N₂,N₄-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD): δ 8.31 (d, 1H, J= 2.3 Hz), 8.20 (d, 1H, J= 2.3 Hz), 8.06 (d, 1H, J= 3.5 Hz), 7.91 (dd, 1H, J= 2.3, 8.2 Hz), 7.65 (dd, 1H, J= 2.9, 8.8 Hz), 7.41 (m, 1H), 7.28 (d, 1H, J= 8.2 Hz), 2.28 (s, 3H), 2.24 (s, 3H); LCMS purity: 87.4%; MS (m/e): 399.20 (M⁺, 100).

7.3.185 N₂,N₄-Bis(2-fluoro-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950147)

In like manner to the preparation of N₂,N₄-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-fluoro-5-nitroaniline were reacted to prepare N₂,N₄-bis(2-fluoro-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 31.07 min.; purity: 93.6%; MS (m/e): 407.14 (MH⁺).

7.3.186 N₂,N₄-Bis(4-fluoro-3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950148)

In like manner to the preparation of N₂,N₄-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-fluoro-3-nitroaniline were reacted to prepare N₂,N₄-bis(4-fluoro-3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 27.17 min.; purity: 94.3%; MS (m/e): 406.96 (MH⁺).

7.3.187 N₂,N₄-Bis(4-methyl-3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950144)

In like manner to the preparation of N₂,N₄-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-methyl-3-nitroaniline were reacted to prepare N₂,N₄-bis(4-methyl-3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 27.40 min.; purity: 96.6%; MS (m/e): 399.00 (MH⁺).

7.3.188 N₂,N₄-Bis(4-chloro-3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950149)

In like manner to the preparation of N₂,N₄-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-chloro-3-nitroaniline were reacted to prepare N₂,N₄-bis(4-chloro-3-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 35.63 min.; purity: 98.9%; MS (m/e): 439.09 (MH⁺).

7.3.189 N₂,N₄-Bis(2-hydroxyethyleneamino-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950150)

In like manner to the preparation of N₂,N₄-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-hydroxyethyleneamino-5-nitroaniline were reacted to prepare N₂,N₄-bis(2-hydroxyethyleneamino-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 17.90 min.; purity: 97.8%; MS (m/e): 489.19 (MH⁺).

7.3.190 N₂,N₄-Bis(2-methoxy-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine (R950151)

In like manner to the preparation of N₂,N₄-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 2-methoxy-5-nitroaniline were reacted to prepare N₂,N₄-bis(2-methoxy-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 31.46 min.; purity: 95.9%; MS (m/e): 431.22 (MH⁺).

7.3.191 N2,N4-Bis(4-fluoro-3-nitrophenyl)-2,4-pyrimidinediamine (R950152)

In like manner to the preparation of N2,N4-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 4-fluoro-3-nitroaniline were reacted to prepare N2,N4-bis(4-fluoro-3-nitrophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 30.92 min.; purity: 94.4%; MS (m/e): 389.31 (MH⁺).

7.3.192 N2,N4-Bis(4-methyl-3-nitrophenyl)-2,4-pyrimidinediamine (R950153)

In like manner to the preparation of N2,N4-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 4-methyl-3-nitroaniline were reacted to prepare N2,N4-bis(4-methyl-3-nitrophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 31.22 min.; purity: 99.6%; MS (m/e): 381.35 (MH⁺).

7.3.193 N2,N4-Bis(4-chloro-3-nitrophenyl)-2,4-pyrimidinediamine (R950154)

In like manner to the preparation of N2,N4-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 4-chloro-3-nitroaniline were reacted to prepare N2,N4-bis(4-chloro-3-nitrophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 37.24 min.; purity: 99.1%; MS (m/e): 421.30 (MH⁺).

7.3.194 N2,N4-Bis(2-hydroxy-5-nitrophenyl)-2,4-pyrimidinediamine (R950155)

In like manner to the preparation of N2,N4-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 2-hydroxy-5-nitroaniline were reacted to prepare N2,N4-bis(2-hydroxy-5-nitrophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 23.26 min.; purity: 100%; MS (m/e): 385.33 (MH⁺).

7.3.195 N2,N4-Bis(2-hydroxyethyleneamino-5-nitrophenyl)-2,4-pyrimidinediamine (R950156)

In like manner to the preparation of N2,N4-bis(2-methyl-5-nitrophenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloropyrimidine and 2-hydroxyethyleneamino-5-nitroaniline were reacted to prepare N2,N4-bis(2-hydroxyethyleneamino-5-nitrophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 17.87 min.; purity: 97.2%; MS (m/e): 470.99 (MH⁺).

7.3.196 N2,N4-Bis[3-(N-isopropyl)aminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950166)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, acetone and sodiumcyanoborohydride were reacted together to give N2,N4-bis[3-(N-isopropyl)aminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 14.07 min.; purity: 90.3%; MS (m/e): 395.14 (MH⁺).

7.3.197 N2,N4-Bis[3-N-(2-hydroxy-1-methylethyl)aminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950171)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, 1-hydroxyacetone and sodiumcyanoborohydride were reacted to give N2,N4-bis[3-N-(2-hydroxy-1-methylethyl)aminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 11.97 min.; purity: 79.01%; MS (m/e): 427.12 (MH⁺).

7.3.198 N2,N4-Bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950177)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and tert-butyl bromoacetate were reacted together to give N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 29.34 min.; purity: 97.2%; MS (m/e): 427.07 (MH⁺).

7.3.199 N4-(3-Aminophenyl)-N2-(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950178)

In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and tert-butyl bromoacetate were reacted together to give N4-(3-aminophenyl)-N2-(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 18.33 min.; purity: 94.5%; MS (m/e): 369.09 (MH⁺).

7.3.200 N2-(3-Aminophenyl)-N4-(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950179)

In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-

aminophenyl)-5-fluoro-2,4-pyrimidinediamine and tert-butyl bromoacetate were reacted together to give N2-(3-aminophenyl)-N4-(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 18.82 min.; purity: 85.8%; MS (m/e): 369.11 (MH⁺).

5 **7.3.201 N2,N4-Bis(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950184)**

In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and ethyl bromoacetate were reacted
10 together to give N2,N4-bis(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 23.41 min.; purity: 96.3%; MS (m/e): 483.08 (MH⁺).

7.3.202 N2,N4-Bis(3-ethoxycarbonylmethyleneaminophenyl)-N2-(ethoxycarbonylmethyl)-5-fluoro-2,4-pyrimidinediamine (R950183)

15 In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and ethyl bromoacetate were reacted together to give N2,N4-bis(3-ethoxycarbonylmethyleneaminophenyl)-N2-(ethoxycarbonylmethyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 25.65 min.;
20 purity: 92.5%; MS (m/e): 569.08 (MH⁺).

7.3.203 N2-(3-Aminophenyl)-N4-(3-hydroxyethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine and N4-(3-Aminophenyl)-N2-(3-hydroxyethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950180)

25 In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and 1-bromo-2-hydroxyethane were reacted together to give a unseparable mixture of N2-(3-aminophenyl)-N4-(3-hydroxyethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine and N4-(3-aminophenyl)-
30 N2-(3-hydroxyethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 9.84 min.; purity: 89.5%; MS (m/e): 355.10 (MH⁺).

7.3.204 N2,N4-Bis(3-hydroxyethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950181)

In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and 1-bromo-2-hydroxyethane were reacted together to give N2,N4-bis(3-hydroxyethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 11.46 min.; purity: 83.3%; MS (m/e): 399.12 (MH⁺).

7.3.205 N2,N4-Bis[3-(N-benzyloxyethyleneamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R950174)

In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and 1-benzyloxy-2-bromoethane were reacted together to give N2,N4-bis[3-(N-benzyloxyethyleneamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 32.92 min.; MS (m/e): 579.17 (MH⁺).

7.3.206 N2-(3-Aminophenyl)-N4-[3-(N-benzyloxyethyleneamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R950175)

In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and 1-benzyloxy-2-bromoethane were reacted together to give N2-(3-aminophenyl)-N4-[3-(N-benzyloxyethyleneamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 23.79 min.; MS (m/e): 445.11 (MH⁺).

7.3.207 N4-(3-Aminophenyl)-N2-[3-(N-benzyloxyethyleneamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R950176)

In like manner to the preparation of N2,N4-bis(3-tert-butoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and 1-benzyloxy-2-bromoethane were reacted together to give N4-(3-aminophenyl)-N2-[3-(N-benzyloxyethyleneamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 23.64 min.; MS (m/e): 445.13 (MH⁺).

7.3.208 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926210)

To a solution of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (0.028g, 0.1 mmol) in MeOH: H₂O (1.8: 0.2 mL) was added 3-hydroxyaniline (0.033g, 0.3 mmol) and heated in a sealed tube at 100 °C for 24h. The resulting reaction was diluted with H₂O (10 mL), acidified with 2N HCl (pH >2), saturated and the resulting solid was filtered to give the desired product, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (**R926210**). Purification can be done by filtration through a pad of silica gel using 1-5% MeOH in CH₂Cl₂ or by crystallization using an appropriate solvent system. ¹H NMR (CDCl₃ + CD₃OD): δ 7.76 (bs, 1H), 7.30 (d, 1H, J= 2.4 Hz), 7.10 (m, 1H), 7.03 (t, 1H, J= 8.1 Hz), 6.89 (dd, 2H, J= 2.4 and 9 Hz), 6.78 (d, 1H, J= 8.7 Hz), 6.42 (dd, 1H, J= 2.4 and 9 Hz), 4.22 (m, 4H); ¹⁹F NMR (CDCl₃ + CD₃OD): - 47196; LCMS: ret. time: 19.55 min.; purity: 95%; MS (m/e): 355 (MH⁺).

Note: When the substrate has ethyl, butyl, benzyl etc. ester functions and the reaction is carried out in methanol as a solvent, the cross esterification to produce methyl ester was observed.

7.3.209 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[3-(hydroxymethyl)phenyl]-2,4-pyrimidinediamine (R925758)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[3-(hydroxymethyl)phenyl]-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[3-(hydroxymethyl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.92 (d, 1H, J= 3.0 Hz), 7.78 (bs, 1H), 7.41-7.31 (m, 3H), 7.12 (d, 1H, J= 7.2 Hz), 6.94 (bs, 1H), 6.81-6.75 (m, 3H), 4.68 (s, 2H), 4.25 (s, 4H); ¹⁹F NMR (CDCl₃): - 47438; LCMS: ret. time: 17.73 min.; purity: 100 %; MS (m/e): 369 (MH⁺).

7.3.210 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-(hydroxymethyl)phenyl]-2,4-pyrimidinediamine (R925760)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(hydroxymethyl)phenyl]-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-(hydroxymethyl)phenyl]-2,4-

pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.92 (bs, 1H), 7.62 (d, 2H, $J = 8.7$ Hz), 7.36 (d, 2H, $J = 8.7$ Hz), 7.19 (d, 1H, $J = 2.1$), 6.87 (dd, 1H, $J = 2.7$ and 8.7 Hz), 6.79 (d, 1H, $J = 8.7$ Hz), 4.68 (s, 2H), 4.28-4.23 (m, 4H); ^{19}F NMR (CDCl_3): - 4.7466; LCMS: ret. time: 17.86 min.; purity: 93 %; MS (m/e): 369 (MH^+).

5 **7.3.211 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-hydroxy-2-phenylethyl)-2,4-pyrimidinediamine (R925765)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(2-hydroxy-2-phenylethyl)-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-hydroxy-2-phenylethyl)-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.79 (s, 1H), 7.48 (m, 5H), 6.89-6.71 (m, 3H), 5.41-5.38, 4.97 (dd, 1H, $J = 3.6$ and 7.5 Hz), 4.28-4.22 (m, 4H), 3.88 (ddd, 1H, $J = 4.2$, 7.2, and 14.1), 3.64-3.55 (m, 1H); ^{19}F NMR (CDCl_3): - 47910; LCMS: ret. time: 20.47 min.; purity: 88 %; MS (m/e): 383 (MH^+).

15 **7.3.212 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[(2R)-hydroxy-(1S)-methyl-2-phenylethyl]-2,4-pyrimidinediamine (R925766)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(2R)-hydroxy-(1S)-methyl-2-phenylethyl]-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[(2R)-hydroxy-(1S)-methyl-2-phenylethyl]-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.80 (bs, 1H), 7.71 (bs, 1H), 7.36-7.23 (m, 6H), 6.91 (dd, 1H, $J = 3.0$ and 9.0 Hz), 6.80 (d, 1H, $J = 9.0$ Hz), 5.17 (d, 1H, $J = 8.1$ Hz), 5.01 (d, 1H, $J = 3.0$ Hz), 4.56-4.50 (m, 1H), 4.24 (s, 4H), 1.10 (d, 3H, $J = 6.3$ Hz); ^{19}F NMR (CDCl_3): - 47840; LCMS: ret. time: 21.43 min.; purity: 99 %; MS (m/e): 397 (MH^+).

7.3.213 N4-Cyclohexyl-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R925794)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-cyclohexyl-5-fluoro-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N4-cyclohexyl-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.62 (d, 1H, $J = 4.2$ Hz), 7.31 (d, 1H, $J = 2.1$ Hz), 6.86 (dd, 1H, $J = 2.4$ and 8.7 Hz), 6.68 (d, 1H, $J =$

8.7 Hz), 4.23-4.16 (m, 4H), 3.99-3.89 (m, 1H), 2.03 (dd, 2H, J= 2.1 and 12.3 Hz), 1.80 (dt, 2H, J= 3.0 and 13.5 Hz), 1.72-1.65 (m, 1H), 1.49-1.20 (m, 5H); ¹⁹F NMR (CD₃OD): - 48332; LCMS: ret. time: 24.54 min.; purity: 95 %; MS (m/e): 345 (MH⁺).

7.3.214 N4-(4-Carboxycyclohexyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R925795)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(4-carboxycyclohexyl)-2-chloro-5-fluoro-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N4-(4-carboxycyclohexyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.62 (d, 1H, J= 4.2 Hz), 7.31 (d, 1H, J= 2.4 Hz), 6.84 (dd, 1H, J= 2.4 and 8.7 Hz), 6.70 (d, 1H, J= 8.7 Hz), 4.23-4.18 (m, 4H), 3.99-4.08 (m, 1H), 2.59 (t, 1H, J= 3.9 Hz), 2.16-2.09 (m, 2H), 1.91-1.84 (m, 2H), 1.78-1.57 (m, 4H); ¹⁹F NMR (CD₃OD): - 48152; LCMS: ret. time: 19.31 min.; purity: 96 %; MS (m/e): 389 (MH⁺).

7.3.215 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925796)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.30 (s, 1H), 9.12 (bs, 1H), 8.91 (bs, 1H), 8.02 (d, 1H, J= 3.3 Hz), 7.35-7.30 (m, 1H), 7.24-7.21 (m, 1H), 7.12 (t, 1H, J= 1.8 Hz), 7.09-7.04 (m, 2H), 6.67 (d, 1H, J= 9.0), 6.46 (dd, 1H, J= 1.8 and 8.4 Hz), 4.18-4.12 (m, 4H); ¹⁹F NMR (DMSO-d₆): - 46594; LCMS: ret. time: 18.43 min.; purity: 97 %; MS (m/e): 355 (MH⁺).

7.3.216 N2-Allyl-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R925823)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and allylamine were reacted to yield N2-allyl-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.71 (bs, 1H), 7.37 (d, 1H, J= 2.4 Hz), 7.07 (dd, 1H, J= 2.4 and 8.7 Hz), 6.75 (d, 1H, J= 8.7 Hz), 5.98-5.85 (m, 1H), 5.19 (dq, 1H, J= 1.8 and 16.8 Hz), 5.06 (dq, 1H, J= 1.8 and 10.5 Hz), 4.24-4.18 (m,

4H), 3.92-3.68 (m, 2H); ^{19}F NMR (CD_3OD): - 48552; LCMS: ret. time: 19.36 min.; purity: 95 %; MS (m/e): 303 (MH^+).

7.3.217 N4-(3,4-Ethylenedioxyphenyl)-N2-(4-ethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926237)

5 In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-ethylaniline were reacted to yield N4-(3,4-ethylenedioxyphenyl)-N2-(4-ethylphenyl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.87 (bs, 1H), 7.42 (d, 2H, $J = 8.7$ Hz), 7.26 (d, 1H, $J = 3.0$ Hz), 7.13-7.08 (m, 3H), 6.95 (dd, 1H, $J = 2.4$ and 8.7 Hz), 6.82 (d, 1H, $J = 9.0$ Hz), 6.60 (bs, 1H), 4.23 (s, 4H), 2.59 (q, 2H, $J = 7.5$ Hz), 1.20 (t, 3H, $J = 7.5$ Hz); ^{19}F NMR (CDCl_3): - 47549; LCMS: ret. time: 25.31 min.; purity: 99 %; MS (m/e): 367 (MH^+).

7.3.218 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine (R926690)

15 In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2-methoxycarbonyl-5-aminobenzofuran were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine. ^1H NMR ($\text{DMSO}-d_6$): δ 9.68 (bs, 1H), 8.13-8.10 (m, 2H), 7.63-7.54 (m, 3H), 7.27 (bs, 1H), 7.10 (d, 1H, $J = 8.7$ Hz), 6.80 (d, 1H, $J = 8.1$ Hz), 4.21 (s, 4H), 3.88 (s, 3H); LCMS: ret. time: 23.22 min.; purity: 95 %; MS (m/e): 437 (MH^+).

7.3.219 5-Fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R926704)

25 In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(isopropoxy)phenyl]-4-pyrimidineamine and 2-methoxycarbonyl-5-aminobenzofuran were reacted to yield 5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 8.04 (d, 1H, $J = 1.8$ Hz), 7.49-7.41 (m, 4H), 7.35 (dd, 1H, $J = 2.4$ and 8.7 Hz), 7.14 (bs, 1H), 6.90 (d, 2H, $J = 9.3$ Hz), 6.70 (bs, 1H), 4.56 (2q, 1H, $J = 5.7$ Hz), 3.98 (s, 3H), 1.37 (d, 6H, $J = 5.7$ Hz); LCMS: ret. time: 25.52 min.; purity: 98 %; MS (m/e): 437 (MH^+).

7.3.220 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(2-hydroxyethyl)oxyphenyl]-2,4-pyrimidinediamine (R926376)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 4-(2-hydroxyethyloxy)aniline were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(2-hydroxyethyl)oxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (D₂O): δ 8.40 (d, 1H J= 4 Hz), 7.57 (m, 6H), 7.12 (m, 2H), 6.90 (m, 2H), 4.40 (m, 4H), 2.2 (s, 3H); LCMS: ret. time: 13.61 min.; purity: 97 %; MS (m/e): 357 (MH⁺).

7.3.221 N2-[4-(2-N,N-Dimethylamino)ethoxyphenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R909236)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 4-(2-N,N-dimethylamino)ethoxyaniline were reacted to yield N2-[4-(2-N,N-dimethylamino)ethoxyphenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.80 (d, 1H J= 4 Hz), 7.47 (dd, 1H, J= 6.8 Hz, 2.7 Hz), 7.44 (m, 1H), 7.05 (m, 1H), 6.85 (m, 1H), 6.78 (m, 2H), 4.16 (m, 2H), 3.03 (m, 2H), 2.55 (s, 6H); LCMS: ret. time: 12.74 min.; purity: 98 %; MS (m/e): 384 (MH⁺).

7.3.222 N2-(1,4-Benzoxazin-3-on-6-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R909238)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 6-amino-1,4-benzoxazin-3-one were reacted to yield N2-(1,4-benzoxazin-3-on-6-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.18 (d, 1H J= 4 Hz), 7.17 (m, 3H), 7.09 (m, 1H), 7.06 (m, 1H), 6.58 (m, 1H), 4.52 (s, 3H); LCMS: ret. time: 17.18 min.; purity: 99 %; MS (m/e): 368 (MH⁺).

7.3.223 N2-(1,4-Benzoxazin-6-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R909241)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 6-amino-1,4-benzoxazine were reacted to yield N2-(1,4-benzoxazin-6-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.18 (d, 1H, J= 4 Hz), 7.15

(m, 3H), 6.68 (m, 2H), 6.52 (m, 2H), 6.52 (m, 1H), 4.18 (m, 2H), 3.37 (m, 2H); LCMS: ret. time 17.42 min.; purity: 95%; MS (m/e): 354 (MH⁺).

7.3.224 N4-(1,4-Benzoxazin-6-yl)-N2-[3-ethoxyocarbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R909242)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-chloro-5-fluoro-4-pyrimidineamine and 3-ethoxyocarbonylmethyleneoxyaniline were reacted to yield N4-(1,4-benzoxazin-6-yl)-N2-(3-ethoxyocarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H

NMR (CD₃OD): δ □□(d, 1H, J= 4 Hz), 7.15 (m, 4H), 6.84 (m, 2H), 6.62 (m, 1H), 4.65 (s, 2H), 4.15 (m, 4H), 3.28 (m, 2H), 1.19 (t, 3H, J= 7 Hz); LCMS: ret. time 22.6 min.; purity: 94%; MS (m/e): 439 (MH⁺).

7.3.225 N2-(1,4-Benzoxazin-6-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R909243)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-chloro-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to yield N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ □□□(d, 1H, J= 4 Hz),

7.18 (m, 3H), 6.68 (m, 2H), 6.45 (m, 2H), 6.52 (m, 1H), 4.22 (m, 2H), 3.31 (m, 2H);

LCMS: ret. time: 17.24; purity: 96%; MS (m/e): 354 (MH⁺).

7.3.226 N4-(1,4-Benzoxazin-6-yl)-N2-(3,5-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R909245)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-chloro-5-fluoro-4-pyrimidineamine and 3,5-dimethoxyaniline were reacted to yield N4-(1,4-benzoxazin-6-yl)-N2-(3,5-

dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ □□□(d, 1H, J= 4 Hz), 6.80 (m, 4H), 6.60 (m, 1H), 6.05 (m, 1H), 4.02 (m, 2H), 3.65 (s, 6H), 3.31 (m, 2H); LCMS: ret. time: 22.38 min.; purity: 99 %; MS (m/e): 398 (MH⁺).

7.3.227 N4-(1,4-Benzoxazin-6-yl)-N2-(3-*tert*-butylphenyl)-5-fluoro-2,4-pyrimidinediamine (R909246)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-chloro-5-fluoro-4-pyrimidineamine

and 3-*tert*-butylaniline were reacted to yield N4-(1,4-benzoxazin-6-yl)-N2-(3-*tert*-butylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ □□ (d, 1H, J= 4 Hz), 7.5 (m, 1H), 7.4 (m, 1H), 7.18 (m, 1H), 7.02 (m, 1H), 6.80 (m, 2H), 6.60 (m, 1H), 4.02 (m, 2H), 3.31 (m, 2H), 1.2 (s, 9H); LCMS: ret. time: 26.64 min.; purity: 99 %; MS (m/e): 508 (MH⁺).

7.3.228 N4-(1,4-Benzoxazin-6-yl)-5-fluoro-N2-[4-(2-hydroxyethyl)oxyphenyl]-2,4-pyrimidinediamine (R909248)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-chloro-5-fluoro-4-pyrimidineamine and 4-(2-hydroxyethyl)oxyaniline were reacted to yield N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[4-(2-hydroxyethyl)oxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ □□□ (d, 1H, J= 4 Hz), 7.52 (m, 1H), 7.4 (m, 3H), 6.90 (m, 2H), 6.68 (m, 1H), 4.56 (s, 2H), 4.02 (m, 2H), 3.75 (m, 2H), 3.31 (m, 4H); LCMS: ret. time: 26.67 min.; purity: 93 %; MS(m/e): 399 (MH⁺).

7.3.229 N2-(2,3-Dihydrobenzofuran-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R909250)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 5-amino-2,3-dihydrobenzofuran were reacted to yield N2-(2,3-dihydrobenzofuran-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.09 (d, 1H), 8.00 (m, 1H), 7.82 (m, 1H), 7.57 (m, 1H), 7.22 (m, 1H), 7.08 (m, 1H), 6.99 (m, 1H), 6.82 (m, 1H), 6.70 (m, 1H), 6.42 (m, 1H), 4.49 (m, 2H), 3.15 (m, 2H); LCMS: ret time: 19.39 min.; MS (m/e): 338 (MH⁺).

7.3.230 N4-(1,4-Benzoxazin-6-yl)-N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R909255)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-chloro-5-fluoro-4-pyrimidineamine and 3-chloro-4-hydroxy-5-methylaniline were reacted to yield N4-(1,4-benzoxazin-6-yl)-N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ □□□ (d, 1H, J= 4 Hz), 7.25 (m, 1H), 7.14 (m, 1H), 6.80 (m, 2H), 6.82 (m, 1H), 4.29 (s, 2H), 3.35 (m, 2H), 2.20 (s, 3H); LCMS: ret. time: 17.05 min.; purity: 99 %; MS(m/e): 402 (MH⁺).

7.3.231 5-Fluoro-N2-(2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R926706)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine and 5-amino-2,3-dihydro-2-(methoxycarbonyl)benzofuran were reacted to yield 5-fluoro-N2-(2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.87 (d, 1H, J= 3.0 Hz), 7.47-7.42 (m, 3H), 7.12 (dd, 1H, J= 2.4 and 8.4 Hz), 6.87 (d, 2H, J= 9.6 Hz), 6.80 (d, 1H, J= 8.7 Hz), 6.63 (d, 1H, J= 2.4 Hz), 5.21 (dd, 1H, J= 6.3 and 10.5 Hz), 4.53 (2q, 1H, J= 5.7 Hz), 3.80 (s, 3H), 3.52 (dd, 1H, J= 10.5 and 15.9 Hz), 3.35 (dd, 1H, J= 6.3 and 15.9 Hz), 1.34 (d, 6H, J= 5.7 Hz); ¹⁹F NMR (CDCl₃): - 47664; LCMS: ret. time: 23.78 min.; purity: 95 %; MS (m/e): 439 (MH⁺).

7.3.232 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926699)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-hydroxyphenyl)-5-fluoro-4-pyrimidineamine and 4-[2-(N-morpholino)ethyleneoxy]aniline were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.34 (s, 1H), 9.17 (bs, 1H), 8.95 (bs, 1H), 8.02 (d, 1H, J= 3.3 Hz), 7.53 (d, 2H, J= 9.0 Hz), 7.28-7.23 (m, 1H), 7.12-7.04 (m, 2H), 6.79 (d, 2H, J= 9.0 Hz), 6.47 (dd, 1H, J= 1.2 and 5.7 Hz), 4.00 (t, 2H, J= 6.0 Hz), 3.56 (t, 4H, J= 4.5 Hz), 2.64 (t, 2H, J= 6.0 Hz), 2.44 (t, 4H, J= 4.5 Hz); ¹⁹F NMR (DMSO-d₆): - 46715; LCMS: ret. time: 12.66 min.; purity: 95 %; MS (m/e): 426 (MH⁺).

7.3.233 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926709)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-[2-(N-morpholino)ethyleneoxy]aniline were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.80 (d,

1H, J= 3.6 Hz), 7.72 (bs, 1H), 7.62 (bs, 1H), 7.41 (d, 1H, J= 9.3 Hz), 7.24 (d, 1H, J= 5.4 Hz), 7.05 (dd, 1H, J= 2.4 and 8.7 Hz), 6.84 (d, 2H, J= 8.7 Hz), 6.75 (d, 1H, J= 9.0 Hz), 4.24 (bs, 4H), 4.11 (t, 2H, J= 5.4 Hz), 3.74-3.69 (m, 4H), 2.80 (t, 2H, J= 5.4 Hz), 2.62-2.58 (m, 4H); ¹⁹F NMR (CD₃OD): - 47912; LCMS: ret. time: 15.16 min.; purity: 91 %; MS (m/e): 468 (MH⁺).

7.3.234 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926710)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-4-pyrimidineamine and 3-aminophenol were reacted to yield 5-fluoro-N2-(3-hydroxyphenyl)-N4-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.84 (d, 1H, J= 4.2 Hz), 7.60 (d, 1H, J= 9.3 Hz), 7.09 (t, 1H, J= 2.4 Hz), 7.04-6.96 (m, 2H), 6.93 (d, 2H, J= 9.3 Hz), 6.40 (dt, 1H, J= 1.8 and 7.5 Hz), 4.15 (t, 2H, J= 5.4 Hz), 3.75-3.70 (m, 4H), 2.81 (t, 2H, J= 5.1 Hz), 2.63-2.59 (m, 4H); LCMS: ret. time: 14.16 min.; purity: 98 %; MS (m/e): 426 (MH⁺).

7.3.235 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926711)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.80 (d, 1H, J= 4.2 Hz), 7.56 (d, 2H, J= 8.7 Hz), 7.13 (d, 1H, J= 2.4 Hz), 6.91 (d, 2H, J= 9.6 Hz), 6.86 (dd, 1H, J= 2.4 and 9.0 Hz), 6.67 (d, 1H, J= 9.0 Hz), 4.23-4.18 (m, 4H), 4.14 (t, 3H, J= 5.4 Hz), 3.74-3.70 (m, 4H), 2.82 (t, 3H, J= 5.4 Hz), 2.64-2.59 (m, 4H); ¹⁹F NMR (CDCl₃): - 47914; LCMS: ret. time: 15.97 min.; purity: 94 %; MS (m/e): 468 (MH⁺).

7.3.236 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-(tetrahydro(1H)-pyrrol-1-ylsulfonyl)phenyl]-2,4-pyrimidinediamine (R926716)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-

fluoro-4-pyrimidineamine and 4-(tetrahydro-(1H)-pyrrol-1-ylsulfonyl)aniline were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-(tetrahydro-(1H)-pyrrol-1-ylsulfonyl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.11 (bs, 1H), 9.76 (bs, 1H), 8.19 (d, 1H, J = 3.9 Hz), 7.82 (d, 2H, J = 8.7 Hz), 7.62 (d, 2H, J = 8.7 Hz), 7.27 (d, 1H, J = 2.4 Hz), 7.08 (dd, 1H, J = 2.4 and 8.7 Hz), 6.85 (d, 1H, J = 8.7 Hz), 4.23 (s, 4H), 3.10-3.06 (m, 4H), 1.64-1.58 (m, 4H); LCMS: ret. time: 22.68 min.; purity: 93 %; MS (m/e): 472 (MH⁺).

7.3.237 N2-[3-[4-(2-Chloro-6-fluorobenzyl)piperazino]propyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926717)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-[4-(2-chloro-6-fluorobenzyl)piperazino]propylamine were reacted to yield N2-[3-[4-(2-chloro-6-fluorobenzyl)piperazino]propyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.79 (d, 1H, J = 3.0 Hz), 7.37 (d, 1H, J = 2.4 Hz), 7.19-7.15 (m, 2H), 7.00-6.93 (m, 2H), 6.81 (d, 1H, J = 8.7 Hz), 6.56 (d, 1H, J = 2.7 Hz), 5.48 (bs, 1H), 4.27-4.21 (m, 4H), 3.70 (d, 2H, J = 1.8 Hz), 3.36 (q, 2H, J = 6.3 Hz), 2.68-2.35 (m, 10H), 1.75 (q, 2H, J = 6.3 Hz); ¹⁹F NMR (CDCl₃): -31693, -48483; LCMS: ret. time: 18.20 min.; purity: 97 %; MS (m/e): 532 (MH⁺).

7.3.238 N2-(4-*tert*-Butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine (R926719)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(4-*tert*-butylphenyl)-2-chloro-5-fluoro-4-pyrimidineamine and 5-amino-2,3-dihydro-2-(methoxycarbonyl)benzofuran were reacted to yield N2-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.16 (bs, 1H), 9.84 (bs, 1H), 8.16 (d, 1H, J = 5.4 Hz), 7.56 (d, 2H, J = 8.1 Hz), 7.49 (s, 1H), 7.35 (d, 2H, J = 8.7 Hz), 7.13 (dd, 1H, J = 1.8 and 8.7 Hz), 6.78 (d, 1H, J = 8.7 Hz), 5.35 (dd, 1H, J = 6.6 and 10.5 Hz), 3.52 (dd, 1H, J = 10.5 and 16.5 Hz), 3.20 (dd, 1H, J = 6.6 and 16.5 Hz), 1.27 (s, 9H); LCMS: ret. time: 26.52 min.; purity: 96 %; MS (m/e): 437 (MH⁺).

7.3.239 N4-[(5-Chloro-1-benzothiophen-3-yl)methyl]-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926721)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(5-chloro-1-benzothiophen-3-yl)methyl]-5-fluoro-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield N4-[(5-chloro-1-benzothiophen-3-yl)methyl]-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.08 (d, 1H, J= 1.8 Hz), 8.02 (d, 1H, J= 8.7 Hz), 7.97 (d, 1H, J= 4.8 Hz), 7.63 (s, 1H), 7.42 (dd, 1H, J= 1.8 and 9.3 Hz), 7.07 (bs, 1H), 6.85 (dd, 1H, J= 2.4 and 8.7 Hz), 6.56 (d, 1H, J= 8.7 Hz), 4.77 (s, 1H), 4.75 (s, 1H), 4.14 (s, 4H); LCMS: ret. time: 25.89 min.; purity: 97 %; MS (m/e): 444 (MH⁺).

7.3.240 N4-[(5-Chloro-1-benzothiophen-3-yl)methyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926722)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(5-chloro-1-benzothiophen-3-yl)methyl]-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to yield N4-[(5-chloro-1-benzothiophen-3-yl)methyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.47 (bs, 1H), 9.33 (bs, 1H), 8.78 (bs, 1), 8.11 (d, 1H, J= 2.1 Hz), 8.02 (d, 1H, J= 8.7 Hz), 7.98 (d, 1H, J= 4.5 Hz), 7.69 (s, 1H), 7.41 (dd, 1H, J= 1.8, 8.1 Hz), 7.07 (bs, 1H), 6.92 (d, 1H, J= 8.4 Hz), 6.82 (t, 1H, J= 8.1 Hz), 6.34 (d, 1H, J= 6.9 Hz), 4.80 (s, 1H), 4.78 (s, 1H); LCMS: ret. time: 23.32 min.; purity: 93 %; MS (m/e): 402 (MH⁺).

7.3.241 N4-[2-[(2-Chloro-6-fluorobenzyl)thio]ethyl]-N2-(3,4-ethylenedioxy)-5-fluoro-2,4-pyrimidinediamine (R926723)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[2-[(2-chloro-6-fluorobenzyl)thio]ethyl]-5-fluoro-4-pyrimidineamine and 1,4-benzodioxan-6-amine were reacted to yield N4-[2-[(2-chloro-6-fluorobenzyl)thio]ethyl]-N2-(3,4-ethylenedioxy)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.09 (bs, 1H), 7.94 (bs, 1H), 7.87 (d, 1H, J= 4.2 Hz), 7.34-7.30 (m, 2H), 7.24-7.18 (m, 2H), 7.01 (dd, 1H, J= 2.4 and 8.7 Hz), 6.68 (d, 1H, J= 8.7 Hz), 4.11 (s, 4H), 3.83 (d, 2H, J= 1.2 Hz), 3.63-3.56 (m, 2H), 2.74 (t, 2H, J= 7.5 Hz); LCMS: ret. time: 25.17 min.; purity: 92 %; MS (m/e): 466 (MH⁺).

7.3.242 N2-(2,3-Dihydro-1,4-benzodioxin-6-ylmethyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945168)

In a manner analogous to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 2,3-dihydro-1,4-benzodioxin-6-ylmethylamine gave N2-(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃) δ 4.24 (s, 4 H), 4.45 (d, J= 6.0 Hz, 2 H), 6.55 (ddd, J= 0.9, 2.4 and 8.4 Hz, 1 H), 6.66 (d, 1 H), 6.84 (m, 4 H), 6.90 (m, 1 H), 7.14 (t, J= 8.1 Hz, 1 H), 7.30 (m, 1 H), 7.86 (d, J= 3.3 Hz, 1 H); ¹⁹F NMR (282 MHz, CDCl₃) δ -170.44; LCMS: ret. time: 18.33 min.; purity: 96.75%; MS (m/e): 369.03 (MH⁺).

7.3.243 N4-[2-[(2-Chloro-6-fluorobenzyl)thio]ethyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926724)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[2-[(2-chloro-6-fluorobenzyl)thio]ethyl]-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to yield N4-[2-[(2-chloro-6-fluorobenzyl)thio]ethyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (methyl sulfoxide-d₆): δ 9.76 (bs, 1H), 9.42 (bs, 1H), 8.70 (bs, 1H), 8.02 (d, 1H, J= 5.1 Hz), 7.33-7.30 (m, 2H), 7.24-7.18 (m, 1H), 7.08-6.96 (m, 2H), 6.42 (d, 1H, J= 4.6 Hz), 3.82 (d, 2H, J= 1.2 Hz), 3.68-3.61 (m, 2H), 2.77 (t, 2H, J= 7.2 Hz); LCMS: ret. time: 23.00 min.; purity: 93 %; MS (m/e): 424 (MH⁺).

7.3.244 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-phenyl-5-methylisoxazol-4-yl)-2,4-pyrimidinediamine (R926743)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 5-methyl-3-phenyl-4-isoxazamine were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-phenyl-5-methylisoxazol-4-yl)-2,4-pyrimidinediamine. LCMS: ret. time: 20.90 min.; purity: 96 %; MS (m/e): 420 (MH⁺).

7.3.245 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3,5-dimethylisoxazol-4-yl)-2,4-pyrimidinediamine (R926744)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3,5-dimethyl-4-isoxazamine were reacted to yield N4-(3,4-

ethylenedioxyphenyl)-5-fluoro-N2-(3,5-dimethylisoxazol-4-yl)-2,4-pyrimidinediamine.

LCMS: ret. time: 18.89 min.; purity: 98 %; MS (m/e): 358 (MH⁺).

7.3.246 N2-[2-(Ethoxycarbonylmethylenethio)pyridin-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926727)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 5-amino-2-(ethoxycarbonylmethylenethio)pyridine were reacted to yield N2-[2-(ethoxycarbonylmethylenethio)pyridin-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.30 (s, 1H), 9.22 (s, 1H), 8.62 (d, 1H, J= 2.4 Hz), 8.06-8.01 (m, 2H), 7.25 (d, 1H, J= 2.4 Hz), 7.18-7.14 (m, 2H), 6.80 (d, 1H, J= 6.0 Hz), 4.22 (bs, 4H), 4.07 (q, 2H, J= 6.9 Hz), 3.95 (s, 2H), 1.14 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 21.60 min.; purity: 97 %; MS (m/e): 458(MH⁺).

7.3.247 N2-[2-(Ethoxycarbonylmethyleneoxy)pyridin-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926740)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 5-amino-2-(ethoxycarbonylmethyleneoxy)pyridine were reacted to yield N2-[2-(ethoxycarbonylmethyleneoxy)pyridin-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.54 (bs, 1H), 9.14 (bs, 1H), 8.05 (s, 1H), 7.88 (d, 1H, J= 2.4 Hz), 7.54 (dd, 1H, J= 2.7 and 10.2 Hz), 7.22 (d, 1H, J= 1.8 Hz), 7.10 (dd, 1H, J= 1.8 and 8.7 Hz), 6.75 (d, 1H, J= 9.0 Hz), 6.40 (d, 1H, J= 9.9 Hz), 4.55 (s, 2H), 4.20 (bs, 4H), 4.10 (q, 2H, J= 7.2 Hz), 1.18 (t, 2H, J= 7.2 Hz).

7.3.248 5-Bromo-N2-(3,4-ethylenedioxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925797)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 5-bromo-2-chloro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield 5-bromo-N2-(3,4-ethylenedioxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 9.33 (s, 1H), 9.06 (s, 1H), 8.34 (s, 1H), 8.14 (s, 1H), 7.13-7.06-(m, 4H), 6.94 (bs, 1H), 6.61 (d, 1H, J= 8.7 Hz), 6.54-6.50 (m, 1H), 4.17-4.13 (m, 4H); LCMS: ret. time: 20.01 min.; purity: 93 %; MS (m/e): 416 (MH⁺).

7.3.249 N2-Allyl-5-bromo-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925822)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 5-bromo-2-chloro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and allylamine were reacted to yield N2-allyl-5-bromo-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.08 (s, 1H), 7.21 (t, 1H, J= 8.1 Hz), 7.02-6.97 (m, 2H), 6.71 (dd, 1H, J= 2.4 and 8.7 Hz), 5.91-5.77 (m, 1H), 5.19-5.09 (m, 2H), 3.94-3.89 (m, 2H); LCMS: ret. time: 18.33 min.; purity: 99 %; MS (m/e): 322 (MH⁺).

7.3.250 5-Cyano-N2-(3,4-ethylenedioxyphenyl)-N4-(methoxycarbonylbenzyl)-2,4-pyrimidinediamine (R925820)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-cyano-N4-(methoxycarbonylbenzyl)-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to yield 5-cyano-N2-(3,4-ethylenedioxyphenyl)-N4-(methoxycarbonylbenzyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.23 (s, 1H), 7.41-7.32 (m, 5H), 7.01 (d, 1H, J= 3.0 Hz), 6.86-6.71 (m, 3H), 6.54 (bs, 1H), 5.48 (d, 1H, J= 6.3 Hz), 4.31 (bs, 4H), 3.68 (s, 3H); LCMS: ret. time: 25.53 min.; purity: 97 %; MS (m/e): 418 (MH⁺).

7.3.251 (R935172): N4-[4-[Ethoxycarbonyl(dimethyl)methyl]phenyl]-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[4-[ethoxycarbonyl(dimethyl)methyl]phenyl]-5-fluoro-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to produce N4-[4-[ethoxycarbonyl(dimethyl)methyl]phenyl]-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.31 (s, 1H), 8.97 (s, 1H), 8.03 (d, 1H, J = 3.5 Hz), 7.70 (d, 2H, J = 8.8 Hz), 7.29 (d, 1H, J = 2.3 Hz), 7.23 (d, 2H, J = 8.8 Hz), 6.98 (dd, 1H, J = 2.1 and 8.8 Hz), 6.66 (d, 1H, J = 8.2 Hz), 4.19-4.15 (m, 4H), 4.07 (qt, 2H, J = 7.0 Hz), 1.48 (s, 6H), 1.10 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 24.51 min.; purity: 100%; MS (m/e): 453 (MH⁺).

7.3.252 (R935173): N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-pyrimidine-2,4-diamine, N4-[4-

5 [ethoxycarbonyl(dimethyl)methyl]phenyl]-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine was reduced with DIBALH to give N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.23 (s, 1H), 8.94 (s, 1H), 8.01 (d, 1H, J = 3.5 Hz), 7.63 (d, 2H, J = 8.8 Hz), 7.31-7.27 (m, 3H), 6.98 (dd, 1H, J = 2.9 and 8.8 Hz), 6.65 (d, 1H, J = 8.8 Hz), 4.65 (t, 1H, J = 5.3 Hz), 4.17-4.16 (m, 4H), 3.39 (d, 2H, J = 5.2 Hz), 1.20 (s, 6H). 8.9 Hz, LCMS: ret. time: 19.52 min.; purity: 100%; MS (*m/e*): 411 (MH⁺).

7.3.253 R935182: 5-Fluoro-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine

15 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3,4-propylenedioxyphenyl)-4-pyrimidineamine and 4-(methoxycarbonylmethyleneoxy)aniline were reacted to produce 5-fluoro-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.16 (s, 1H), 9.01 (s, 1H), 8.10 (d, 1H, J = 4.1 Hz), 7.51 (d, 2H, J = 8.8 Hz), 7.37 (d, 1H, J = 2.9 Hz), 7.32 (dd, 1H, J = 2.9 and 8.8 Hz), 6.98 (d, 1H, J = 8.3 Hz), 6.80 (d, 2H, J = 8.3 Hz), 4.70 (s, 2H), 4.12-4.05 (app qt, 4H, J = 5.3 Hz), 3.68 (s, 3H), 2.07 (q, 2H, J = 5.3 Hz); LCMS: ret. time: 20.51 min.; purity: 97%; MS (*m/e*): 441 (MH⁺).

7.3.254 R935185: 5-Fluoro-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine

25 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3,4-propylenedioxyphenyl)-4-pyrimidineamine and 3-(methoxycarbonylmethyleneoxy)aniline were reacted to produce 5-fluoro-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.22 (s, 1H), 9.18 (s, 1H), 8.07 (d, 1H, J = 3.5 Hz), 7.41-7.35 (m, 2H), 7.32-7.28 (m, 2H), 7.09 (t, 1H, J = 8.2 Hz), 6.90 (d, 1H, J = 8.2 Hz), 6.43 (dd, 1H, J = 2.3 and 8.8 Hz), 4.65 (s, 2H), 4.11-4.04 (app q, 4H, J = 5.3 Hz), 3.67

(s, 3H), 2.06 (q, 2H, J= 5.3 Hz); LCMS: ret. time: 20.57 min.; purity: 97%; MS (*m/e*): 441 (MH⁺).

7.3.255 R935187: N4-[3-(1-Bis(ethoxycarbonyl)ethoxy)phenyl]-5-fluoro-N2-[4-isopropoxyphenyl]-2,4-pyrimidinediamine

5 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine and 3-[1-bis(ethoxycarbonyl)ethoxy]aniline were reacted to provide N4-[3-(1-bis(ethoxycarbonyl)ethoxy)phenyl]-5-fluoro-N2-[4-isopropoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.08 (s, 1H), 9.98 (s, 1H), 8.19 (d, 1H, J= 4.7 Hz), 7.55 (d, 2H, J= 8.8 Hz), 7.25 (d, 1H, J= 8.8 Hz), 7.15 (d 1H, J= 8.3 Hz), 7.13 (d, 1H, J= 8.3 Hz), 6.91 (d, 2H, J= 8.8 Hz), 6.51 (dd, 1H, J= 1.7 and 8.3 Hz), 4.56 (q, 1H, J= 5.8 Hz), 4.19 (qt, 4H, J= 7.0 Hz), 1.61 (s, 3H), 1.23 (d, 6H, J= 5.8 Hz), 1.14 (t, 6H, J= 7.0 Hz); LCMS: ret. time: 15.23 min.; purity: 94%; MS (*m/e*): 527 (MH⁺).

15 **7.3.256 R935190: N4-(3,4-Ethylenedioxyphenyl)-5-fluoro -N2-(indazolin-6-yl) - 2,4-pyrimidinediamine.**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 6-aminoindazole were reacted to produce N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(indazolin-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.69 (s, 1H), 9.62 (s, 1H), 8.14 (d, 1H, J= 4.7 Hz), 7.93 (s, 1H), 7.92 (s, 1H), 7.60 (d, 1H, J= 8.8 Hz), 7.33-7.31 (m 1H), 7.24 (dd, 2H, J= 1.7 and 8.8 Hz), 6.79 (d, J= 8.8 Hz), 4.20 (s, 4H); LCMS: ret. time: 17.66 min.; purity: 99%; MS (*m/e*): 379 (MH⁺)

7.3.257 R935191: 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(indazolin-6-yl)-2,4-pyrimidinediamine

25 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine and 5-aminoindazole were reacted to give 5-fluoro N4-(3-hydroxyphenyl)-N2-(indazolin-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.74 (s, 1H), 9.66 (s, 1H), 8.18 (d, 1H, J= 4.1 Hz), 7.95 (s, 1H), 7.93 (s, 1H), 7.59 (d, 1H, J= 8.8 Hz), 7.33-7.26 (m, 2H), 7.12-7.07 (m, 2H), 6.52 (dd, 1H, J= 2.3 and 8.2 Hz); LCMS: ret. time: 15.27 min.; purity: 99%; MS (*m/e*): 337 (MH⁺)

7.3.258 R935193: N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine and 1-methyl-5-aminoindazole were reacted to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.42 (s, 2H), 8.25 (d, 1H, J = 5.2 Hz), 7.92 (s, 1H), 7.86 (app s, 1H), 7.61 (d, 1H, J = 8.8 Hz), 7.38 (dd, 1H, J = 2.3 and 9.3 Hz), 7.21 (d, 1H, J = 2.3 Hz), 7.09 (dd, 1H, J = 2.3 and 8.8 Hz), 6.79 (d, 1H, J = 8.8 Hz), 4.20 (s, 4H), 4.02 (s, 3H); LCMS: ret. time: 19.09 min.; purity: 99%; MS (*m/e*): 393 (MH⁺).

7.3.259 R935194: 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 1-methyl-5-aminoindazole to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.56 (s, 1H), 10.49 (s, 1H), 8.29 (d, 1H, J = 5.2 Hz), 7.98 (d, 1H, J = 1.7 Hz), 7.92 (s, 1H), 7.59 (d, 1H, J = 8.8 Hz), 7.36 (dd, 1H, J = 1.7 and 8.8 Hz), 7.10 (br m, 3H), 6.66 (td, 1H, J = 1.7 and 7.0 Hz), 4.01 (s, 3H). LCMS: ret. time: 16.62 min.; purity: 98%; MS (*m/e*): 351 (MH⁺).

7.3.260 R935197: 5-Fluoro-N2-(indazoline-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine:

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine was reacted with 5-aminoindazole to produce 5-fluoro-N2-(indazoline-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.96 (s, 1H), 9.76 (s, 1H), 8.12 (d, 1H, J = 4.6 Hz), 7.94 (s, 1H), 7.92 (s, 1H), 7.53 (d, 2H, J = 9.8 Hz), 7.46 (d, 1H, J = 8.8 Hz), 7.34 (dd, 1H, J = 1.7 and 9.8 Hz), 6.83 (d, 2H, J = 9.8 Hz), 4.55 (q, 1H, J = 5.8 Hz), 1.24 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 18.96 min.; purity: 100%; MS (*m/e*): 379 (MH⁺).

7.3.261 R935198: N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine and 5-aminoindazole were reacted to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.91 (s, 1H), 9.82 (s, 1H), 8.13 (d, 1H, J= 4.6 Hz), 7.94 (app s, 2H), 7.47 (d, 1H, J= 8.8 Hz), 7.36 (dd, 1H, J= 1.7 and 8.8 Hz), 7.23 (d, 1H, J= 2.3 Hz), 7.13 (dd, 1H, J= 2.3 and 8.8 Hz), 6.76 (d, 1H, J= 8.8 Hz), 4.20 (s, 4H); LCMS: ret. time: 16.17 min.; purity: 99%; MS (*m/e*): 379 (MH⁺).

7.3.262 R935199: 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine and 5-aminoindazole were reacted to give 5-fluoro-N4-(3-hydroxyphenyl)-N2-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.78 (s, 1H), 9.68 (s, 1H), 9.49 (br s, 1H), 8.13 (d, 1H, J= 4.6 Hz), 8.06 (s, 1H), 7.93 (s, 1H), 7.50 (d, 1H, J= 8.8 Hz), 7.38 (dd, 1H, J= 1.7 and 8.8 Hz), 7.17 (d, 1H, J= 8.2 Hz), 7.11-7.06 (m, 2H), 6.57 (dd, 1H, J= 1.1 and 8.2 Hz). LCMS: ret. time: 13.79 min.; purity: 96%; MS (*m/e*): 337 (MH⁺).

7.3.263 R935203: 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1-methyl-indazoline-5-yl)-4-pyrimidineamine and 4-isopropoxyaniline were reacted to produce 5-fluoro-N2-(4-isopropoxyphenyl)-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.57 (s, 1H), 10.12 (s, 1H), 8.24 (d, 1H, J= 5.3 Hz), 8.04 (s, 1H), 7.95 (s, 1H), 7.63 (d, 1H, J= 9.3 Hz), 7.55 (dd, 1H, J= 1.7 and 8.8 Hz), 7.30 (d, 2H, J= 9.4 Hz), 6.82 (d, 2H, J= 8.8 Hz), 4.53 (q, 1H, J= 6.4 Hz), 4.02 (s, 3H), 1.22 (d, 6H, J= 6.4 Hz). LCMS: ret. time: 20.56 min.; purity: 99%; MS (*m/e*): 393 (MH⁺).

7.3.264 R935204: 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1-methyl-indazoline-5-yl)-4-pyrimidineamine and 3-aminophenol were reacted to produce 5-fluoro-N2-(3-hydroxyphenyl)-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine.. LCMS: ret. time: 15.55 min.; purity: 98%; MS (*m/e*): 351 (MH⁺).

7.3.265 R935207: N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-methoxycarbonyl-fur-4-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro- N-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 4-(4-aminophenoxymethyl)-2-methoxycarbonyl-furan to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-methoxycarbonyl-fur-4-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.48 (s, 1H), 9.41 (s, 1H), 8.08 (d, 1H, J= 3.4 Hz), 7.37-7.10 (m, 6H), 6.74 (d, 2H, J= 8.2 Hz), 6.61 (d, 1H, J= 8.2 Hz), 5.00 (s, 2H), 4.19 (br s, 4H), 3.79 (s, 3H). LCMS: ret. time: 22.85 min.; purity: 97%; MS (*m/e*): 493 (MH⁺).

7.3.266 R935208: N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine was reacted with 6-amino-1-(methoxycarbonyl)methyl-indazoline to produce N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.39 (s, 1H), 9.19 (s, 1H), 8.08 (d, 1H, J= 3.5 Hz), 7.95 (s, 1H), 7.91 (s, 1H), 7.56 (d, 1H, J= 8.2 Hz), 7.32 (d, 2H, J= 8.9 Hz), 7.22 (dd, 1H, J= 2.9 and 8.2 Hz), 6.78 (d, 1H, J= 8.8 Hz), 5.06 (s, 2H), 4.21 (s, 4H), 3.61 (s, 3H). LCMS: ret. time: 19.39 min.; purity: 93%; MS (*m/e*): 451 (MH⁺).

7.3.267 R935209: 5-Fluoro-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1-methyl-indazoline-5-yl)-4-pyrimidineamine and 4-(methoxycarbonylmethyleneoxy)aniline were reacted to provide 5-fluoro-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.31 (s, 1H), 8.99 (s, 1H), 8.17 (s, 1H), 8.02 (d, 1H, J= 3.5 Hz), 7.92 (s, 1H), 7.59 (s, 2H), 7.50 (d, 2H, J= 8.8 Hz), 6.73 (d, 2H, J= 8.8 Hz), 4.69 (s, 2H), 4.03 (s, 3H), 3.68 (s, 3H). LCMS: ret. time: 17.60 min.; purity: 99%; MS (m/e): 423 (MH⁺).

7.3.268 R935214: 5-Fluoro-N2-(3,5-dimethoxyphenyl)-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1-methyl-indazoline-5-yl)-4-pyrimidineamine and 3,5-dimethoxyaniline were reacted to produce 5-fluoro-N2-(3,5-dimethoxyphenyl)-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.34 (s, 1H), 9.09 (s, 1H), 8.20 (d, 1H, J= 5.3 Hz), 8.07 (d, 1H, J= 3.5 Hz), 7.90 (s, 1H), 7.63-7.55 (m, 2H), 6.89 (d, 2H, J= 1.7 Hz), 6.02 (t, 1H, J= 2.3 Hz), 4.02 (s, 3H), 3.54 (s, 6H). LCMS: ret. time: 18.81 min.; purity: 97%; MS (m/e): 395 (MH⁺).

7.3.269 R935215: 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 6-amino-1-(methoxycarbonyl)methyl-indazoline to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 16.08 min.; purity: 90%; MS (m/e): 408 (MH⁺).

7.3.270 R935218: 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-

indazoline-6-yl]-4-pyrimidineamine was reacted with 4-isopropoxyaniline to provide 5-fluoro-N2-(4-isopropoxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.47 (s, 1H), 8.99 (s, 1H), 8.10 (s, 1H), 8.07 (d, 1H, J= 4.1 Hz), 8.02 (s, 1H), 7.68 (d, 1H, J= 8.8 Hz), 7.50-7.46 (m, 3H), 6.74 (d, 2H, 8.8 Hz), 5.26 (s, 2H), 4.47 (q, 1H, J= 5.8 Hz), 3.62 (s, 3H), 1.21 (d, 6H, J= 5.8 Hz). LCMS: ret. time: 21.76 min.; purity: 97%; MS (*m/e*): 451 (MH⁺).

7.3.271 R935219: N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-4-pyrimidineamine was reacted with 3,4-ethylenedioxyaniline to provide N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.48 (s, 1H), 9.01 (s, 1H), 8.10 (s, 1H), 8.09 (d, 1H, J= 3.5 Hz), 8.01 (s, 1H), 7.68 (d, 1H, J= 8.8 Hz), 7.48-7.43 (m, 1H), 7.29 (d, 1H, J= 2.3 Hz), 6.99 (d, 1H, J= 2.3 and 8.2 Hz), 6.67 (dd, 1H, J= 2.3 and 8.8 Hz), 5.27 (s, 2H), 4.15 (s, 4H), 3.62 (s, 3H). LCMS: ret. time: 18.99 min.; purity: 93%; MS (*m/e*): 451 (MH⁺).

7.3.272 R935220: 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-4-pyrimidineamine was reacted with 3-aminophenol to provide 5-fluoro-N2-(3-hydroxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.51 (s, 1H), 9.19 (s, 1H), 9.10 (s, 1H), 8.21 (s, 1H), 8.12 (d, 1H, J= 3.5 Hz), 8.02 (s, 1H), 7.68 (d, 1H, J= 8.8 Hz), 7.49-7.45 (m 1H), 7.16 (s, 1H), 7.09 (d, 1H, J= 7.6 Hz), 6.95 (app t, 1H, J= 7.6 and 8.2 Hz), 6.31 (dd, 1H, J= 1.7 and 7.6 Hz), 5.29 (s, 2H), 3.62 (s, 3H). LCMS: ret. time: 16.16 min.; purity: 97%; MS (*m/e*): 409 (MH⁺).

7.3.273 N4-(3,4-Ethylenedioxyphenyl)-N2-(3-furanylmethylene)-5-fluoro-2,4-pyrimidinediamine (R950203)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-aminomethylenefurane were reacted to give N4-(3,4-ethylenedioxyphenyl)-N2-(3-furanylmethylene)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 19.99 min.; purity: 88.4%; MS (m/e): 343.07 (MH⁺).

7.3.274 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[(4-methoxyphenyloxy)ethyl]-2,4-pyrimidinediamine (R950204)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2-(4-methoxyphenyloxy)ethyl amine were reacted to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[(4-methoxyphenyloxy)ethyl]-2,4-pyrimidinediamine. LCMS: ret. time: 22.74 min.; purity: 91.9%; MS (m/e): 413.05 (MH⁺).

7.3.275 N2-[2,3-Dihydrobenzo[b]furan-5-ylmethyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950205)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2,3-dihydrobenzo[b]furan-5-ylmethylamine were reacted to give N2-[2,3-dihydrobenzo[b]furan-5-ylmethyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 21.43 min.; purity: 97.5%; MS (m/e): 395.05 (MH⁺).

7.3.276 N2-(2,3-Dihydro-1,4-benzodioxin-2-ylmethyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950206)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2,3-dihydro-1,4-benzodioxin-2-ylmethylamine were reacted to give N2-(2,3-dihydro-1,4-benzodioxin-2-ylmethyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 22.49 min.; purity: 87.6%; MS (m/e): 411.01 (MH⁺).

7.3.277 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(methylthio)-1,3-benzothiaz-6-yl]-2,4-pyrimidinediamine (R950201)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-N4-(3-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2-(methylthio)-1,3-benzothiazol-6-amine were reacted to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(methylthio)-1,3-benzothiaz-6-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 22.67 min.; purity: 76.9%; MS (m/e): 441.91 (MH⁺).

7.3.278 N2-[2,3-Dihydrobenzo[b]furan-5-ylmethyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950213)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 2,3-dihydrobenzo[b]furan-5-ylmethylamine were reacted to N2-[2,3-dihydrobenzo[b]furan-5-ylmethyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 17.80 min.; purity: 99.2%; MS (m/e): 353.08 (MH⁺).

7.3.279 N2-(2,3-Dihydro-1,4-benzodioxin-2-ylmethyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950214)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 2,3-dihydro-1,4-benzodioxin-2-ylmethylamine were reacted to give N2-(2,3-dihydro-1,4-benzodioxin-2-ylmethyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 19.26 min.; purity: 96.2%; MS (m/e): 369.08 (MH⁺).

7.3.280 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(methylthio)-1,3-benzothiaz-6-yl]-2,4-pyrimidinediamine (R950212)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 2-(methylthio)-1,3-benzothiazol-6-amine were reacted to give 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(methylthio)-1,3-benzothiaz-6-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 19.83 min.; purity: 98.9%; MS (m/e): 399.98 (MH⁺).

7.3.281 N2-(3-Aminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950227)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 1,3-diaminobenzene were reacted to give N2-(3-aminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 11.89 min.; purity: 97.6%; MS (m/e): 312.05 (MH⁺).

7.3.282 N2-(1,4-Benzoxazin-6-yl)-5-fluoro-N4-(3-nitrophenyl)-2,4-pyrimidinediamine (R950253)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 6-amino-1,4-benzoxazine were reacted to give N2-(1,4-benzoxazin-6-yl)-5-fluoro-N4-(3-nitrophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 18.52 min.; purity: 99.5%; MS (m/e): 382.93 (MH⁺).

7.3.283 N2-(Ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950215)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-ethoxycarbonylmethyleneaminophenylaniline were reacted to N2-(ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 18.90 min.; purity: 83.4%; MS (m/e): 398.06 (MH⁺).

7.3.284 N2-(Ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine (R950229)

In like manner to the preparation of N4-(3-aminophenyl)-N2-[2-(methoxycarbonyl)-benzofurane-5-yl]-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-ethoxycarbonylmethyleneaminophenylaniline were reacted to N2-(ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 16.37 min.; purity: 78.3%; MS (m/e): 441.03 (MH⁺).

7.3.285 5-Cyano-N2-(3-hydroxyphenyl)-N4-(methoxycarbonylbenzyl)-2,4-pyrimidinediamine (R925821)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-cyano-N4-(methoxycarbonylbenzyl)-4-pyrimidineamine and 3-hydroxyaniline were reacted to yield 5-cyano-N2-(3-hydroxyphenyl)-N4-(methoxycarbonylbenzyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.27 (s, 1H), 7.38-7.28 (m, 5H), 7.19-7.07 (m, 2H), 6.98-6.91 (m, 2H), 6.64 (d, 1H, J= 6.6 Hz), 3.55 (s, 3H); LCMS: ret. time: 24.18 min.; purity: 98 %; MS (m/e): 376 (MH⁺).

7.3.286 5-Fluoro-N4-[2-fluoro-4-(methoxymethyleneoxy)phenyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926680)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(2-fluoro-4-methoxymethyleneoxyphenyl)-4-pyrimidineamine and 3-hydroxyaniline were reacted to yield 5-fluoro-N4-(2-fluoro-4-methoxymethyleneoxyphenyl)-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine.

7.3.287 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[(1H)-indol-5-yl]-2,4-pyrimidinediamine (R926748)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 5-aminoindole were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[(1H)-indol-5-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 20.37 min.; purity: 97 %; MS (m/e): 378 (MH⁺).

7.3.288 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[(1H)-indol-5-yl]-2,4-pyrimidinediamine (R926749)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 5-aminoindole were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[(1H)-indol-5-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 17.31 min.; purity: 94 %; MS (m/e): 366 (MH⁺).

7.3.289 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[(1H)-indol-6-yl]-2,4-pyrimidinediamine (R926750)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 6-aminoindole were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[(1H)-indol-6-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 20.80 min.; purity: 91 %; MS (m/e): 378 (MH⁺).

7.3.290 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[(1H)-indol-6-yl]-2,4-pyrimidinediamine (R926751)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 6-aminoindole were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[(1H)-indol-6-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 18.13 min.; purity: 96 %; MS (m/e): 336 (MH⁺).

7.3.291 N4-[4-(Aminocarbonylmethyleneoxy)phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945063)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 3-hydroxyaniline (110 mg, 1 mmol) and N4-[4-(aminocarbonylmethyleneoxy)phenyl]-2-chloro-5-fluoro-4-pyrimidineamine (80 mg, 0.27 mmol) gave N4-[4-(aminocarbonylmethyleneoxy)phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (75 mg, 76%). ¹H NMR (acetone-*d*₆): δ 4.51 (s, 2 H), 6.64 (dm, J= 8.4 Hz, 1 H), 7.06-7.14 (m, 5 H), 7.70 (dd, J= 2.4 and 9.0 Hz, 2 H), 8.27 (d, J= 6.0 Hz, 1 H); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ - 164.00; LCMS: ret. time: 14.66 min.; purity: 88.63%; MS (m/e): 370.00 (MH⁺).

7.3.292 N4-[4-(Cyanomethyleneoxy)phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945071)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 3-hydroxyaniline (94 mg, 0.86 mmol) and 2-chloro-N4-[4-(cyanomethyleneoxy)phenyl]-5-fluoro-4-pyrimidineamine (80 mg, 0.29 mmol) gave N4-[4-(cyanomethyleneoxy)phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (65 mg, 64%) as a off-white solid. ¹H NMR (acetone-*d*₆): δ 5.16 (s, 2 H), 6.64 (ddd, J= 1.8, 2.4 and 7.5 Hz, 1 H), 7.03 (t, J= 2.1 Hz, 1 H), 7.08-7.16 (m, 2 H),

7.19 (d, J= 9.3 Hz, 2 H), 7.77 (d, J= 9.3 Hz, 2 H), 8.30 (d, J= 5.4 Hz, 1 H), 10.04 (s, 1 H, NH), 11.33 (s, 1 H, NH); ^{19}F NMR (282 MHz, acetone- d_6): δ - 163.52; LCMS: ret. time: 17.08 min.; purity: 100%; MS (m/e): 352.13 (MH^+).

7.3.293 N4-(3-Cyanophenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945109)

5

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 3-aminobenzonitrile (142 mg, 1.2 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave 2-chloro-N4-(3-cyanophenyl)-5-fluoro-4-pyrimidineamine (128 mg, 86%) as a white solid. The reaction of
10 2-chloro-N4-(3-cyanophenyl)-5-fluoro-4-pyrimidineamine (50 mg, 0.2 mmol) and 3-aminophenol (66 mg, 0.6 mmol) gave N4-(3-cyanophenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (40 mg, 62%). ^1H NMR (acetone- d_6): δ 6.48 (ddd, J= 0.9, 2.4 and 7.8 Hz, 1 H), 7.10 (t, J= 8.1 Hz, 1 H), 7.18 (ddd, J= 1.2, 2.1 and 8.1 Hz, 1 H), 7.33 (t, J= 2.1 Hz, 1 H), 7.45 (dt, J= 1.2 and 7.8 Hz, 1 H), 7.54 (t, J= 8.1 Hz, 1 H), 8.08 (d, J= 3.3 Hz, 1 H),
15 8.14 (ddd, J= 1.5, 2.7 and 8.4 Hz, 1 H), 8.39 (t, J= 2.1 Hz, 1 H), 8.58 (s, 1 H, NH), 8.84 (s, 1 H, NH); ^{19}F NMR (282 MHz, acetone- d_6): δ - 167.41; LCMS: ret. time: 17.75 min.; purity: 92.39%; MS (m/e): 322.59 (MH^+).

7.3.294 N4-(3-Cyanophenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945110)

20

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-cyanophenyl)-5-fluoro-4-pyrimidineamine (50 mg, 0.2 mmol) and 4-(methoxycarbonylmethyleneoxy)aniline (109 mg, 0.6 mmol) gave N4-(3-cyanophenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (30 mg, 38%). ^1H NMR (acetone- d_6): δ 3.74 (s, 3 H), 4.72 (s, 2 H), 6.93 (d, J= 9.0 Hz, 2 H), 7.46 (dt, J= 1.5 and 7.5 Hz, 1 H), 7.54 (t, J= 7.8 Hz, 1 H), 7.60 (dd, J= 1.8 and 9.0 Hz, 2 H), 8.03-8.07 (m, 2 H), 8.43 (m, 1 H), 8.48 (br, 1 H, NH), 8.80 (br, 1 H, NH); ^{19}F NMR (282 MHz, acetone- d_6): δ - 168.2; LCMS: ret. time: 20.24 min.; purity: 94.79%; MS (m/e): 393.98 (MH^+).

25

7.3.295 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(indol-3-yl)ethyl]-2,4-pyrimidinediamine (R945117)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (50 mg, 0.21 mmol) and tryptamine (100 mg, 0.62 mmol) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(indol-3-yl)ethyl]-2,4-pyrimidinediamine (40 mg, 53%). ¹H NMR (CD₃OD): δ 3.01 (t, J= 7.2 Hz, 2 H), 3.61 (t, J= 7.2 Hz, 2 H), 6.51 (ddd, J= 0.9, 2.1 and 8.1 Hz, 1 H), 6.96 (td, J= 0.9 and 7.2 Hz, 1 H), 7.03-7.09 (m, 3 H), 7.22 (d, J= 7.5 Hz, 1 H), 7.28-7.32 (m, 2 H), 7.53 (d, J= 7.8 Hz, 1 H), 7.72 (d, J= 4.5 Hz, 1 H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 171.72; LCMS: ret. time: 20.17 min.; 95.66%; MS (m/e): 364.05 (MH⁺).

7.3.296 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945118)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (80 mg, 0.33 mmol) and 3-methoxycarbonylmethyleneoxyaniline (180 mg, 0.99 mmol) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (130 mg). ¹H NMR (acetone-*d*₆): δ 3.74 (s, 3 H), 4.64 (s, 2 H), 6.71 (m, 1 H), 6.80 (m, 1 H), 7.23-7.32 (m, 6 H), 8.32 (d, J= 5.1 Hz, 1 H); LCMS: ret. time: 18.37 min.; purity: 100%; MS (m/e): 384.70 (MH⁺).

7.3.297 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945124)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (80 mg, 0.28 mmol) and 3-methoxycarbonylmethyleneoxyaniline (154 mg, 0.85 mmol) gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (90 mg, 74%). ¹H NMR (CDCl₃): δ 3.80 (s, 3H), 4.27 (q, J= 0.9 Hz, 4H), 4.58 (s, 2H), 6.54 (ddd, J= 0.9, 2.7 and 8.1 Hz, 1H), 6.65 (d, J= 2.7 Hz, 1H), 6.86 (d, J= 8.7 Hz, 1H), 6.98 (dd, J= 2.4 and 8.4 Hz, 1H), 6.98 (br, 1 H), 7.09 (ddd, J= 1.2, 2.1 and

8.1 Hz, 1H), 7.18 (t, J= 8.1 Hz, 1H), 7.24 (d, J= 2.4 Hz, 1H), 7.32 (t, J= 2.1 Hz, 1H), 7.92 (d, J= 3.3 Hz, 1H); ^{19}F NMR (282 MHz, CDCl_3): δ - 167.52; LCMS: ret. time: 21.64 min.; purity: 98.07%; MS (m/e): 426.99 (MH^+).

7.3.298 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945125)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine (80 mg, 0.28 mmol) and methyl 3-aminophenoxyacetate (154 mg, 0.85 mmol) gave 5-fluoro-N4-(4-isopropoxyphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (80 mg, 66%). ^1H NMR (CDCl_3) δ 1.33 (s, 3H), 1.35 (s, 3H), 3.80 (s, 3H), 4.52 (p, J= 6.0 Hz, 1H), 4.55 (s, 2H), 6.53 (ddd, J= 0.9, 2.4 and 8.1 Hz, 1H), 6.69 (d, J= 2.4 Hz, 1H), 6.90 (d, J= 9.0 Hz, 2H), 7.04-7.08 (m, 2H), 7.16 (t, J= 8.1 Hz, 1H), 7.32 (t, J= 2.1 Hz, 1H), 7.47 (d, J= 8.7 Hz, 2H), 7.92 (d, J= 3.0 Hz, 1H); ^{19}F NMR (282 MHz, CDCl_3): δ - 167.64; LCMS: ret. time: 24.70 min.; purity: 100%; MS (m/e): 427.00 (MH^+).

7.3.299 N2-[4-(Aminocarbonylmethyleneoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945064)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-N2-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 4-(aminocarbonylmethyleneoxy)aniline (198 mg, 1.2 mmol) and 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (95 mg, 0.4 mmol) gave N2-[4-(aminocarbonylmethyleneoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (60 mg, 41%). ^1H NMR (CD_3OD): δ 4.55 (s, 2H), 6.75 (dm, J= 7.5 Hz, 1H), 7.08 (d, J= 9.3 Hz, 2H), 7.18 (m, 2H), 7.22 (d, J= 8.7 Hz, 1H), 7.46 (d, J= 9.0 Hz, 2H), 8.09 (d, 1H); LCMS: ret. time: 14.38 min.; purity: 100%; MS (m/e): 370.00 (MH^+).

7.3.300 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945132)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyaniline (490 mg, 2.4 mmol) and 2,4-dichloro-5-fluoropyrimidine (200

mg, 1.2 mmol) gave 2-chloro-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-4-pyrimidineamine. The reaction of 2-chloro-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-4-pyrimidineamine (40 mg, 0.12 mmol) and 3-aminophenol (40 mg, 0.36 mmol) gave 5-fluoro-N2-(3-hydroxyphenyl)-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (30 mg, 62%).
¹H NMR (CDCl₃): δ 2.61 (s, 3H), 5.21 (s, 2H), 6.50 (ddd, J= 0.9, 2.4 and 7.8 Hz, 1H), 6.76 (ddd, J= 0.6, 2.4 and 9.0 Hz, 1H), 6.80-6.85 (m, 3H), 7.12 (t, J= 8.1 Hz, 1H), 7.23 (t, J= 7.8 Hz, 1H), 7.50-7.52 (m, 2H), 7.94 (d, J= 3.3 Hz, 1H), 7.98 (t, J= 2.4 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 167.19; LCMS: ret. time: 18.88 min.; purity: 100%; MS (m/e): 408.97 (MH⁺).

7.3.301 N2-[4-(Aminocarbonylmethoxy)phenyl]-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945133)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-4-pyrimidineamine (30 mg, 0.09 mmol) and 4-(aminocarbonylmethyleneoxy)aniline (45 mg, 0.27 mmol) gave N2-[4-(aminocarbonylmethyleneoxy)phenyl]-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (10 mg, 24%).
¹H NMR (acetone-*d*₆): δ 2.62 (s, 3H), 4.43 (s, 2H), 5.19 (s, 2H), 6.77 (ddd, J= 1.2, 2.4 and 8.1 Hz, 1H), 6.94 (d, J= 9.0 Hz, 2H), 7.25 (t, J= 8.1 Hz, 1H), 7.34 (ddd, J= 0.9, 1.8, 9.0 Hz, 1H), 7.68 (d, J= 9.0 Hz, 2H), 7.81 (t, J= 2.1 Hz, 1H), 7.99 (d, J= 3.6 Hz, 1H), 8.45 (br, 1H, NH), 8.57 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ - 168.20; LCMS: ret. time: 16.80 min.; purity: 84.91%; MS (m/e): 466.05 (MH⁺).

7.3.302 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945128)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (40 mg, 0.14 mmol) and 3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyaniline (87 mg, 0.42 mmol) gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-

yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (30 mg, 47%). ¹H NMR (CDCl₃): δ 2.62 (s, 3H), 4.26 (q, J= 2.1 Hz, 4H), 5.09 (s, 2H), 6.63-6.67 (m, 2H), 6.85 (d, J= 8.4 Hz, 1H), 6.95-6.99 (m, 2H), 7.09 (dt, J= 0.9 and 6.9 Hz, 1H), 7.19 (t, J= 8.4 Hz, 1H), 7.23 (d, J= 2.4 Hz, 1H), 7.42 (t, J= 2.4 Hz, 1H), 7.92 (d, J= 3.0 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 167.47; LCMS: ret. time: 21.26 min.; purity: 96.72%; MS (m/e): 451.01 (MH⁺).

7.3.303 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945129)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine (40 mg, 0.14 mmol) and 3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyaniline (87 mg, 0.42 mmol) gave 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (40 mg, 63%). ¹H NMR (CDCl₃): δ 1.32 (s, 3H), 1.34 (s, 3H), 2.61 (s, 3H), 4.52 (p, J= 6.0 Hz, 1H), 5.08 (s, 2H), 6.64 (ddd, J= 1.2, 2.7 and 8.1 Hz, 1H), 6.70 (d, J= 2.4 Hz, 1H), 6.89 (d, J= 9.0 Hz, 2H), 7.07-7.11 (m, 2H), 7.16 (t, J= 8.1 Hz, 1H), 7.38 (t, J= 2.1 Hz, 1H), 7.46 (d, J= 9.0 Hz, 2H), 7.91 (d, J= 3.3 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 167.55; LCMS: ret. time: 24.49 min.; 96.15%; MS (m/e): 451.08 (MH⁺).

7.3.304 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945137)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-4-pyrimidineamine (40 mg, 0.12 mmol) and 3,4-ethylenedioxyaniline (55 mg, 0.36 mmol) reacted to give N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 2.60 (s, 3H), 4.24 (q, J= 2.7 Hz, 4H), 5.21 (s, 2H), 6.74-6.78 (m, 2H), 6.81 (d, J= 8.4 Hz, 1H), 6.90 (dd, J= 1.2, 7.8 Hz, 1H), 7.01 (dd, J= 2.4 and 8.4 Hz, 1H), 7.22 (t, J= 8.4 Hz, 1H), 7.30 (d, J= 2.4 Hz, 1H), 7.48 (br, 1H), 7.94 (d, J= 3.3 Hz, 1H), 7.98 (br, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 168.23; LCMS: ret. time: 21.20 min.; purity: 91.09%; MS (m/e): 450.99 (MH⁺).

7.3.305 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945138)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-4-pyrimidineamine (40 mg, 0.12 mmol) and 4-isopropoxyaniline (55 mg, 0.36 mmol) gave 5-fluoro-N2-(4-isopropoxyphenyl)-N4-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 1.31 (s, 3H), 1.33 (s, 3H), 2.60 (s, 3H), 4.48 (p, J= 6.0 Hz, 1H), 5.20 (s, 2H), 6.74-6.78 (m, 2H), 6.87 (d, J= 9.0 Hz, 2H), 6.92 (dd, J= 1.2 and 8.4 Hz, 1H), 7.22 (t, J= 8.4 Hz, 1H), 7.50 (m, 3H), 7.94 (d, J= 3.0 Hz, 2H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 168.46; LCMS: ret. time: 24.95 min.; purity: 73.74%; MS (m/e): 451.06 (MH⁺).

7.3.306 N4-(3,5-Dimethyl-4-hydroxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945139)

Using general hydrogenation conditions, 2,6-dimethyl-4-nitrophenol was reduced to 4-amino-2,6-dimethylphenol. In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 4-amino-2,6-dimethylphenol (823 mg, 6 mmol) and 2,4-dichloro-5-fluoropyrimidine (500 mg, 3 mmol) gave 2-chloro-N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-4-pyrimidineamine. Compound 2-chloro-N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-4-pyrimidineamine (500 mg, 1.87 mmol) and 3-(methoxycarbonylmethyleneoxy)aniline (500 mg, 2.76 mmol) reacted to give N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (500 mg, 65%). ¹H NMR (CD₃OD): δ 2.16 (s, 6H), 3.76 (s, 3H), 4.51 (s, 2H), 6.79 (ddd, J= 0.9, 2.4 and 8.1 Hz, 1H), 7.01-7.06 (m, 2H), 7.15 (s, 2H), 7.26 (t, J= 8.1 Hz, 1H), 7.93 (d, J= 5.7 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 163.31; LCMS: ret. time: 20.44 min.; purity: 84.25%; MS (m/e): 413.03 (MH⁺).

7.3.307 N4-(Benzothiophen-3-ylmethyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945146)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of benzothiophen-3-ylmethylamine (244 mg, 1.5 mmol) and 2,4-dichloro-5-fluoropyrimidine (50 mg, 0.3 mmol)

gave N4-(benzothiophen-3-ylmethyl)-2-chloro-5-fluoro-4-pyrimidineamine. The reaction of N4-(benzothiophen-3-ylmethyl)-2-chloro-5-fluoro-4-pyrimidineamine and 3-aminophenol (200 mg, 1.83 mmol) gave N4-(benzothiophen-3-ylmethyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. (40 mg, 36%). ¹H NMR (CDCl₃): δ 4.45 (br, 1H), 4.95 (dd, J= 1.2 and 5.4 Hz, 2H), 5.33 (br, 1H), 6.40 (ddd, J= 1.2, 2.4 and 8.1 Hz, 1H), 6.85 (ddd, J= 0.9, 2.1 and 8.1 Hz, 1H), 6.91 (br, 1H), 7.05 (t, J= 8.1 Hz, 1H), 7.26 (m, 1H), 7.39-7.47 (m, 3H), 7.81 (dd, J= 1.2 and 5.1 Hz, 1H), 7.84 (d, J= 3.3 Hz, 1H), 7.92 (m, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 168.89; LCMS: ret. time: 21.91 min.; purity: 99.34%; MS (m/e): 366.96 (MH⁺).

10 **7.3.308 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(3-pyridylmethyl)-2,4-pyrimidinediamine (R945147)**

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 3-pyridylmethylamine (162 mg, 1.5 mmol) and 2,4-dichloro-5-fluoropyrimidine (50 mg, 0.3 mmol) were reacted to give 2-chloro-5-fluoro-N4-(3-pyridylmethyl)-4-pyrimidineamine. Then 2-chloro-5-fluoro-N4-(3-pyridylmethyl)-4-pyrimidineamine and 3-aminophenol (200 mg, 1.83 mmol) reacted to give 5-fluoro-N2-(3-hydroxyphenyl)-N4-(3-pyridylmethyl)-2,4-pyrimidinediamine (40 mg, 43%). ¹H NMR (CD₃OD): δ 4.71 (s, 2H), 6.38 (ddd, J= 0.9, 2.4 and 8.1 Hz, 1H), 6.88 (ddd, J= 0.9, 2.1 and 8.1 Hz, 1H), 7.00 (t, J= 8.1 Hz, 1H), 7.14 (t, J= 2.4 Hz, 1H), 7.37 (dd, J= 4.8 and 7.8 Hz, 1H), 7.73 (d, J= 3.6 Hz, 1H), 7.87 (dt, J= 2.1 and 7.5 Hz, 1H), 8.39 (dd, J= 1.2 and 7.8 Hz, 1H), 8.57 (d, J= 2.1 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 170.99; LCMS: ret. time: 8.82 min.; purity: 92.90%; MS (m/e): 312.05 (MH⁺).

25 **7.3.309 N4-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945148)**

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 4-amino-2-chloro-6-methylphenol and 2,4-dichloro-5-fluoropyrimidine resulted 2-chloro-N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-4-pyrimidineamine. The reaction of 2-chloro-N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-4-pyrimidineamine and 3-methoxycarbonylmethyleneoxyaniline (1.95 g, 11 mmol) gave N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (850 mg, 55%). ¹H NMR

(CD₃OD): δ 2.22 (s, 3H), 3.76 (s, 3H), 4.52 (s, 2H), 6.50 (dt, J= 2.7 and 6.3 Hz, 1H), 7.09-7.14 (m, 2H), 7.24 (t, J= 1.8 Hz, 1H), 7.30 (t, J= 1.2 Hz, 1H), 7.49 (d, J= 2.4 Hz, 1H), 7.88 (d, J= 3.9 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 168.70; LCMS: ret. time: 20.63 min.; purity: 98.56%; MS (m/e): 432.96 (MH⁺).

5 **7.3.310 N4-[(2,5-Dimethyl-3-furyl)methyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945151)**

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of (2,5-dimethyl-3-furyl)methylamine (188 mg, 1.5 mmol) and 2,4-dichloro-5-fluoropyrimidine (50 mg, 0.3 mmol) gave 2-chloro-N4-[(2,5-dimethyl-3-furyl)methyl]-5-fluoro-4-pyrimidineamine. The reaction of 2-chloro-N4-[(2,5-dimethyl-3-furyl)methyl]-5-fluoro-4-pyrimidineamine and 3-aminophenol (200 mg, 1.83 mmol) gave N4-[(2,5-dimethyl-3-furyl)methyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (50 mg, 51%). ¹H NMR (CDCl₃): δ 2.22 (s, 3H), 2.23 (s, 3H), 4.39 (d, J= 5.1 Hz, 2H), 5.24 (br, 1H), 5.90 (s, 1H), 6.52 (d, J= 6.6 Hz, 1H), 6.99 (d, J= 8.1 Hz, 1H), 7.13 (t, J= 8.1 Hz, 1H), 7.29 (s, 1H), 7.71 (m, 2H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 167.84; LCMS: ret. time: 19.83 min.; purity: 96.32%; MS (m/e): 329.05 (MH⁺).

20 **7.3.311 N4-(3,5-Dimethyl-4-methoxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945153)**

In a manner analogous to the preparation of N2,N4-bis[3-methoxy-4-(methoxycarbonyl)phenyl]-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,6-dimethyl-4-nitrophenol (1.67 g, 10 mmol), potassium carbonate (13 g, 0.1 mol) and iodomethane (2.5 mL, 50 mmol) gave 2,6-dimethyl-1-methoxy-4-nitrobenzene. Hydrogenation of 2,6-dimethyl-1-methoxy-4-nitrobenzene gave 3,5-dimethyl-4-methoxyaniline.

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-N2-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 3,5-dimethyl-4-methoxyaniline (400 mg, 2.6 mmol) and 2,4-dichloro-5-fluoropyrimidine (200 mg, 1.2 mmol) gave 2-chloro-N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine. The reaction of 2-chloro-N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine and 3-(methoxycarbonylmethyleneoxy)aniline (650 mg, 3.6 mmol) gave N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-

pyrimidinediamine (180 mg, 35%). ^1H NMR (CD_3OD): δ 2.20 (s, 6H), 3.70 (s, 3H), 3.74 (s, 3H), 4.52 (s, 2H), 6.76 (ddd, $J = 0.9, 2.4$ and 8.4 Hz, 1H), 7.03-7.08 (m, 2H), 7.24 (m, 3H), 7.96 (d, $J = 5.4$ Hz, 1H); ^{19}F NMR (282 MHz, CD_3OD): δ - 162.92; LCMS: ret. time: 23.13 min.; purity: 100%; MS (m/e): 427.04 (MH^+).

5 **7.3.312 N4-[4-(N-Benzylpiperazino)phenyl]-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945155)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-N2-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidineamine, the reaction of N4-[4-(N-benzylpiperazino)phenyl]-2-chloro-5-fluoro-4-pyrimidineamine (50 mg, 0.12 mmol) and 3,4-ethylenedioxyaniline (0.045 mL, 0.36 mmol) gave N4-[4-(N-benzylpiperazino)phenyl]-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (40 mg, 63%). ^1H NMR (CDCl_3): δ 2.64 (t, $J = 4.8$ Hz, 4H), 3.20 (t, $J = 4.8$ Hz, 4H), 3.59 (s, 2H), 4.24 (m, 4H), 6.61 (d, 1H, NH), 6.68 (br, 1H, NH), 6.76 (d, $J = 8.7$ Hz, 1H), 6.88 (dd, $J = 2.4$ and 8.7 Hz, 1H), 6.93 (d, $J = 8.7$ Hz, 2H), 7.19 (d, $J = 2.4$ Hz, 1H), 7.28-7.36 (m, 5H), 7.47 (d, $J = 8.7$ Hz, 2H), 7.87 (d, $J = 3.3$ Hz, 1H); ^{19}F NMR (282 MHz, CDCl_3): δ -168.66; LCMS: ret. time: 18.05 min.; purity: 100%; MS (m/e): 513.10 (MH^+).

7.3.313 N2-[(2,5-Dimethyl-3-furyl)methyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945162)

20 In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (50 mg, 0.21 mmol) and (2,5-dimethyl-3-furyl)methylamine (80 mg, 0.63 mmol) gave N2-[(2,5-dimethyl-3-furyl)methyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (40 mg, 59%). ^1H NMR (acetone- d_6): δ 2.14 (s, 6H), 4.37 (d, $J = 4.2$ Hz, 2H), 5.96 (s, 1H), 6.77 (d, $J = 6.6$ Hz, 1H), 7.23-7.28 (m, 2H), 7.44 (s, 1H), 8.11 (d, $J = 4.8$ Hz, 1H), 9.05 (br, 1H), 9.75 (br, 1H); ^{19}F NMR (282 MHz, acetone- d_6): δ - 165.77; LCMS: ret. time: 19.23 min.; purity: 94.89%; MS (m/e): 329.08 (MH^+).

7.3.314 N2-[4-(N-Benzylpiperazino)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945163)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (50 mg, 0.18 mmol) and 4-(4-benzylpiperazino)aniline (142 mg, 0.53 mmol) resulted N2-[4-(N-benzylpiperazino)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (30 mg, 33%). ¹H NMR (CDCl₃): δ 2.63 (t, J= 4.8 Hz, 4H), 3.16 (t, J= 4.8 Hz, 4H), 3.58 (s, 2H), 4.27 (m, 4H), 6.56 (d, 1H, NH), 6.70 (br, 1H, NH), 6.82 (d, J= 8.7 Hz, 1H), 6.89 (d, J= 9.0 Hz, 2H), 6.96 (dd, J= 2.7 and 8.7 Hz, 1H), 7.28 (d, J= 2.7 Hz, 1H), 7.30-7.36 (m, 5H), 7.39 (d, J= 8.7 Hz, 2H), 7.88 (d, J= 3.3 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 168.94; LCMS: ret. time: 18.12 min.; purity: 98.42%; MS (m/e): 512.95 (MH⁺).

7.3.315 N2-(Benzothiophen-3-ylmethyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945164)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (50 mg, 0.21 mmol) and benzothiophen-3-ylmethylamine (100 mg, 0.61 mmol) gave N2-(benzothiophen-3-ylmethyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (40 mg, 53%). ¹H NMR (CDCl₃): δ 4.82 (d, J= 6.0 Hz, 2H), 6.45 (dd, J= 8.1 Hz, 1H), 6.70 (m, 1H), 6.80 (d, J= 8.4 Hz, 1H), 7.03 (t, J= 8.1 Hz, 1H), 7.22 (m, 1H), 7.34 (s, 1H), 7.39-7.46 (m, 2H), 7.82 (m, 1H), 7.89-7.92 (m, 2H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 170.02; LCMS: ret. time: 21.29 min.; purity: 92.97%; MS (m/e): 367.03 (MH⁺).

7.3.316 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(3-pyridylmethyl)-2,4-pyrimidinediamine (R945165)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (50 mg, 0.21 mmol) and 3-pyridylmethylamine (68 mg, 0.63 mmol) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-pyridylmethyl)-2,4-pyrimidinediamine (40 mg, 62%). ¹H NMR (CDCl₃): δ 4.40 (d, J= 6.3 Hz, 2H), 5.60 (br, 1H), 6.62-6.70 (m, 3H), 7.05 (br, 1H), 7.14 (t, J= 8.1 Hz, 1H), 7.30 (dd, J= 5.1 and 7.8 Hz,

1H), 7.73 (d, J= 7.5 Hz, 1H), 7.80 (d, J= 3.3 Hz, 1H), 8.49 (d, J= 4.5 Hz, 1H), 8.66 (s, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 169.52; LCMS: ret. time: 9.41 min.; purity: 99.25%; MS (m/e): 312.01 (MH⁺).

5 **7.3.317 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(2-pyridylmethyl)-2,4-pyrimidinediamine (R945166)**

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (50 mg, 0.21 mmol) and 2-pyridylmethylamine (68 mg, 0.63 mmol) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-pyridylmethyl)-2,4-pyrimidinediamine (40 mg, 62%). ¹H NMR (CDCl₃): δ 4.73 (d, J= 6.3 Hz, 2H), 5.85 (t, J= 6.0 Hz, 1H, NH), 6.48 (d, J= 6.9 Hz, 1H), 6.61 (dd, J= 2.7 and 8.1 Hz, 1H), 6.67 (s, 1H), 7.13 (t, J= 8.1 Hz, 1H), 7.21 (dd, J= 5.1 and 7.5 Hz, 1H), 7.49 (d, J= 7.5 Hz, 1H), 7.69 (td, J= 1.8 and 7.8 Hz, 1H), 7.85 (d, J= 3.6 Hz, 1H), 8.38 (br, 1H), 8.56 (dd, J= 1.2 and 3.9 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ -170.49; LCMS: ret. time: 10.10 min.; purity: 100%; MS (m/e): 312.08 (MH⁺).

7.3.318 N4-(3,5-Dimethoxyphenyl)-N2-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926802)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,5-dimethoxyphenyl)-5-fluoro-4-pyrimidineamine with 3-hydroxyaniline gave N4-(3,5-dimethoxyphenyl)-N2-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 18.98 min.; purity: 90%; MS (m/e): 357 (MH⁺).

7.3.319 N4-(3,5-Dimethoxyphenyl)-N2-(2-ethoxycarbonylindol-7-yl)-5-fluoro-2,4-pyrimidinediamine (R926803)

25 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,5-dimethoxyphenyl)-5-fluoro-4-pyrimidineamine with 2-ethoxycarbonyl-7-aminoindole gave N4-(3,5-dimethoxyphenyl)-N2-(2-ethoxycarbonylindol-7-yl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 24.21 min.; purity: 91%; MS (m/e): 452 (MH⁺).

7.3.320 N2-(3,4-Dimethoxyphenyl)-N4-(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926108)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(4-ethoxyphenyl)-5-fluoro-4-pyrimidineamine with 3,4-dimethoxyaniline gave N2-(3,4-dimethoxyphenyl)-N4-(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.89 (d, 1H, J= 3 Hz), 7.45 (bd, 2H, J= 9 Hz), 7.20 (d, 1H, J= 2.4 Hz), 6.96-6.77 (m, 5H), 6.63 (bs, 1H), 4.03 (q, 2H, J= 7.2 Hz), 3.86 (s, 3H), 3.72 (s, 3H), 1.42 (t, 3H, J= 7.2 Hz); ¹⁹F NMR (CDCl₃): -47473.

7.3.321 N4-(4-Ethoxyphenyl)-N2-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926146)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(4-ethoxyphenyl)-5-fluoro-4-pyrimidineamine with 3-hydroxyaniline gave N4-(4-ethoxyphenyl)-N2-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.79 (d, 1H, J= 4.2 Hz), 7.54 (dd, 2H, J= 2.4 and 7.2 Hz), 7.05-6.97 (m, 3H), 6.87 (dd, 2H, J= 2.4 and 4.2 Hz), 6.41 (m, 1H), 4.02 (q, 2H, J= 6.6 Hz), 1.38 (t, 3H, J= 6.9 Hz); ¹⁹F NMR (CD₃OD): -47444; LCMS: ret. time: 21.15 min.; purity: 98%; MS (m/e): 341 (MH⁺).

7.3.322 N4-(4-Ethoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926213)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(4-ethoxyphenyl)-5-fluoro-4-pyrimidineamine with 3,4-ethylenedioxyaniline gave N4-(4-ethoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.87 (d, 1H, J= 3Hz), 7.47 (dd, 2H, J= 2.4 and 5.1 Hz), 7.18 (d, 1H, J= 2.4 Hz), 6.91-6.85 (m, 3H), 6.79-6.73 (m, 2H), 6.64 (bs, 1H), 4.25 (bs, 4H), 4.05 (q, 2H, J= 6.9 Hz), 1.43 (t, 3H, J= 7.2 Hz); ¹⁹F NMR (CDCl₃): -47467; LCMS: ret. time: 24.32 min.; purity: 90%; MS (m/e): 383 (MH⁺).

7.3.323 N4-(3,4-Dimethoxyphenyl)-N2-(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926145)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-

dimethoxyphenyl)-5-fluoro-4-pyrimidineamine with 4-ethoxyaniline gave N4-(3,4-dimethoxyphenyl)-N2-(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.90 (bs, 1H), 7.37 (dd, 2H, J= 2.4 and 6.3 Hz), 7.21 (d, 1H, J= 2.4 Hz), 7.03 (dd, 1H, J= 2.4 and 8.1 Hz), 6.86-6.80 (m, 4H), 6.65 (bs, 1H), 4.00 (q, 2H, J= 7.2 Hz), 3.89 (s, 3H), 3.75 (s, 3H), 1.39 (t, 3H, J= 6.9 Hz); ¹⁹F NMR (CDCl₃): -47501; LCMS: ret. time: 22.69 min.; purity: 98%; MS (m/e): 385 (MH⁺).

7.3.324 N4-(3,4-Dimethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926147)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-dimethoxyphenyl)-5-fluoro-4-pyrimidineamine with 3-hydroxyaniline gave N4-(3,4-dimethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.77 (d, 1H, J= 3.3 Hz), 7.15 (d, 1H, J= 2.4 Hz), 7.05 (dd, 1H, J= 2.4 and 8.4 Hz), 7.00-6.90 (m, 4H), 6.80 (d, 1H, J= 8.1 Hz), 6.40 (m, 1H), 4.05 (q, 2H), 3.80 (s, 3H), 3.75 (s, 3H), 1.20 (t, 3H); ¹⁹F NMR (CD₃OD): - 47223; LCMS: ret. time: 17.94 min.; purity: 99%; MS (m/e): 357 (MH⁺).

7.3.325 N2-(3,4-Dimethoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926113)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3,4-dimethoxyaniline gave N2-(3,4-dimethoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.90 (d, 1H, J= 6.6 Hz), 7.59 (bs, 1H), 7.30 (s, 1H), 7.20-7.10 (m, 2H), 7.00-6.75 (m, 4H), 6.59 (bd, 1H, J= 7.8 Hz), 3.87 (s, 3H), 3.84 (s, 3H); ¹⁹F NMR (CDCl₃): - 47229; LCMS: ret. time: 17.77 min.; purity: 78%; MS (m/e): 357 (MH⁺).

7.3.326 N2-(4-Ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926395)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with ethyl-4-aminophenoxyacetate gave N2-(4-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.90 (d, 1H, J= 5.1 Hz), 7.35 (dd, 2H, J= 2.1 and

7.2 Hz), 7.13 (t, 1H, J= 7.2 Hz), 7.10 (d, 1H, J= 6.6 Hz), 6.96 (dd, 2H, J= 2.4 and 7.2 Hz), 6.67 (m, 1H), 4.72 (s, 2H), 4.25 (q, 2H, J= 7.2 Hz), 1.29 (t, 3H, J= 7.2 Hz); ¹⁹F NMR (CD₃OD): - 21885; LCMS: ret. time: 20.18 min.; purity: 92%; MS (m/e): 399 (MH⁺).

7.3.327 5-Bromo-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926396)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with ethyl 4-aminophenoxyacetate gave 5-bromo-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine.

LCMS: ret. time: 21.64 min.; purity: 92%; MS (m/e): 459 (MH⁺).

7.3.328 N2-(4-Ethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926211)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-ethoxyaniline were reacted to yield N2-(4-ethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.88 (bs, 1H), 7.40 (bd, 2H, J= 8.7 Hz), 7.27 (bd, 2H, J= 6.3 Hz), 6.95 (dd, 1H, J= 3 and 9 Hz), 6.86-6.77 (m, 3H), 6.58 (s, 1H), 4.28 (bs, 4H), 4.01 (q, 2H, J= 6.9 Hz), 1.40 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 24.46 min.; purity: 90%; MS (m/e): 383 (MH⁺).

7.3.329 N2-(3,4-Dimethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926212)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3,4-dimethoxyaniline were reacted to yield N2-(3,4-dimethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 20.98 min.; purity: 74%; MS (m/e): 399 (MH⁺).

7.3.330 N2-(3-Chloro-4-fluorophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926218)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-chloro-4-fluoroaniline were reacted to yield N2-(3-chloro-4-fluorophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR

(CD₃OD): δ 7.75 (bd, 1H), 7.70 (bd, 1H), 7.18 (m, 1H), 7.10 (m, 1H), 6.90 (m, 2H), 6.75 (m, 1H), 4.20 (bs, 4H); LCMS: ret. time: 25.04 min.; purity: 99%; MS (m/e): 392 (MH⁺).

7.3.331 N2-(4-tert-Butylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926219)

5 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-tert-butylaniline were reacted to yield N2-(4-tert-butylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.85 (d, 1H, J= 3.6 Hz), 7.44 (bdd, 2H, J= 6.3 Hz), 7.35-7.31 (m, 3H), 6.93 (dd, 1H, J= 2.7 and 8.7 Hz), 6.83 (d, 1H, J= 9 Hz), 6.80 (bs, 1H), 4.23 (s, 4H), 1.28 (s, 9H); LCMS: ret. time: 27.67 min.; purity: 98%; MS (m/e): 395 (MH⁺).

7.3.332 N4-(3,4-Ethylenedioxyphenyl)-N2-(4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine (R926220)

15 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-fluoroaniline were reacted to yield N4-(3,4-ethylenedioxyphenyl)-N2-(4-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.92 (bs, 1H), 7.80 (bs, 1H), 7.60 (bd, 2H), 6.90 (m, 2H), 6.80 (bs, 1H), 6.65 (bs, 1H), 4.25 (s, 4H); LCMS: ret. time: 22.87 min.; purity: 97%; MS (m/e): 357 (MH⁺).

20 **7.3.333 N4-(3,4-Ethylenedioxyphenyl)-N2-(3-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine (R926221)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-fluoroaniline were reacted to yield N4-(3,4-ethylenedioxyphenyl)-N2-(3-fluorophenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.76 (d, 1H, J= 5.6 Hz), 7.39 (m, 2H), 7.14 (d, 1H, J= 2.4 Hz), 6.94-6.85 (m, 3H), 6.75 (d, 1H, J= 9 Hz), 4.21 (s, 4H); LCMS: ret. time: 22.60 min.; purity: 100%; MS (m/e): 357 (MH⁺).

7.3.334 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxyethyl)-2,4-pyrimidinediamine (R926229)

30 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-

4-pyrimidineamine and 2-methoxyethylamine were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxyethyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.81 (bs, 1H), 7.33 (d, 1H, J= 2.4 Hz), 6.93 (dd, 1H, J= 2.4 Hz and 9 Hz), 6.81 (d, 1H, J= 9 Hz), 6.53 (s, 1H), 4.25 (bs, 2H), 3.54 (bs, 2H), 3.36 (s, 3H); LCMS: ret. time: 18.01 min.; purity: 100%; MS (m/e): 321 (MH⁺).

7.3.335 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(4-methoxybenzyl)-2,4-pyrimidinediamine (R926230)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-methoxybenzylamine were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(4-methoxybenzyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.81 (d, 1H, J= 2.7 Hz), 7.27 (m, 3H), 6.86 (m, 3H), 6.52 (s, 1H), 5.14 (s, 1H), 4.46 (d, 2H, J= 5.4 Hz), 4.24 (s, 4H), 3.78 (s, 3H); LCMS: ret. time: 23.06 min.; purity: 94%; MS (m/e): 383 (MH⁺).

7.3.336 N2-(2,2-Difluorobenzodioxol-5-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926386)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2,2-difluoro-5-aminobenzodioxole were reacted to yield N2-(2,2-difluorobenzodioxol-5-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 9.39 (s, 1H), 9.24 (s, 1H), 8.06 (d, 1H, J= 5.6 Hz), 7.87 (d, 1H, J= 1.8 Hz), 7.27-7.19 (m, 3H), 7.08 (dd, 1H, J= 2.4 and 8.7 Hz), 6.80 (d, 1H, J= 9 Hz), 4.21 (bs, 4H); ¹⁹F NMR (CDCl₃): -14012 and -46487; LCMS: ret. time: 25.32 min.; purity: 100%; MS (m/e): 419 (MH⁺).

7.3.337 N2-(2-Ethoxycarbonylindol-5-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926476)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2-ethoxycarbonyl-5-aminoindole were reacted to yield N2-(2-ethoxycarbonylindol-5-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine.

¹H NMR (CDCl₃): δ 7.84 (d, 1H, J= 5.4 Hz), 7.76 (d, 1H, J= 3.6 Hz), 7.50 (d, 1H, J= 9 Hz), 7.23-7.15 (m, 3H), 7.03 (bd, 1H, J= 8.7 Hz), 6.78 (d, 1H, J= 8.7 Hz), 4.38 (q, 2H, J= 7.2 Hz), 4.22 (s, 4H), 1.41 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 23.58 min; purity: 100%; MS (m/e): 451 (MH⁺).

5 **7.3.338 N2-(4-Cyanomethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926480)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-10 4-pyrimidineamine and 4-cyanomethyleneoxyaniline were reacted to yield N2-(4-cyanomethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.87 (d, 1H, J= 3.6 Hz), 7.52 (d, 1H, J= 8.7 Hz), 7.38 (bs, 1H), 7.28 (d, 1H, J= 2.4 Hz), 6.96-6.86 (m, 3H), 6.65 (bd, 1H), 4.73 (s, 2H), 4.29 (m, 4H); ¹⁹F NMR (CDCl₃): - 47416; LCMS: ret. time: 20.49 min.; purity: 100%; MS (m/e): 394 (MH⁺).

15 **7.3.339 N2-(3-Ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926482)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-20 4-pyrimidineamine and ethyl-3-aminophenoxyacetate were reacted to yield N2-(3-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 10.53 (s, 1H), 8.18 (s, 1H), 7.67 (d, 1H, J= 4.8 Hz), 7.19-7.02 (m, 5H), 6.86 (d, 1H, 9Hz), 6.71 (dd, 1H, J= 1.8 and 9 Hz), 4.51 (s, 2H), 4.25 (m, 6H), 1.29 (t, 3H, J= 7.5 Hz); ¹⁹F NMR (CDCl₃): - 45640; LCMS: ret. time: 22.71 25 min.; purity: 99%; MS (m/e): 441 (MH⁺).

7.3.340 N2-(3-Ethoxycarbonylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R925745)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-30 4-pyrimidineamine and 3-ethoxycarbonylaniline gave N2-(3-ethoxycarbonylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.04 (bs, 1H), 7.94 (bs, 1H), 7.90 (bd, 1H), 7.68 (bd, 1H, J= 7.5 Hz), 7.35 (t, 1H, J= 8.1 Hz), 7.28 (d,

1H, J= 2.4 Hz), 7.07 (s, 1H), 6.93 (dd, 1H, J= 3 and 8.7 Hz), 6.83 (d, 1H, J= 9 Hz), 6.64 (bs, 1H), 4.36 (q, 2H, J= 7.2 Hz), 4.26 (s, 4H), 1.35 (t, 3H, J= 7.5 Hz); ¹⁹F NMR (CDCl₃): -47247; LCMS: ret. time: 15.88.; purity: 100%; MS (m/e): 411 (MH⁺).

7.3.341 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-hydroxyethyl)-2,4-pyrimidinediamine (R925746)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2-hydroxyethylamine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-hydroxyethyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.7 (bs, 1H), 7.32 (d, 1H, J= 2.4 Hz), 7.05 (dd, 1H, J= 2.4 and 9 Hz), 6.75 (d, 1H, J= 8.9 Hz), 4.21 (s, 4H), 3.67 (t, 2H, J= 5.7 Hz), 3.38 (t, 2H, J= 5.4 Hz); ¹⁹F NMR (CD₃OD): -48518; LCMD: ret. time: 15.54 min.; purity: 100%; MS (m/e): 307 (MH⁺).

7.3.342 N2-(4-Ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R925747)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and ethyl-4-aminophenoxyacetate gave N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.88 (bs, 1H), 7.42 (dd, 2H, J= 2.4 and 6.9 Hz), 7.28 (d, 1H, J= 3 Hz), 6.95-6.81 (m, 4H), 6.59 (s, 1H), 4.59 (s, 4H), 4.28 (q, 2H, J= 6.2 Hz), 1.30 (t, 3H, J= 6.1 Hz); ¹⁹F NMR (CDCl₃): -47570; LCMS: ret. time: 22.74 min.; purity: 100%; MS (m/e): 441 (MH⁺).

7.3.343 N2-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940233)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-chloro-4-hydroxy-5-methylaniline gave N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: retn, time: 19.20 min.; purity: 94%; MS (m/e): 360 (M⁺); ¹H NMR (CDCl₃): δ 7.93 (1H, d, J= 3.1 Hz), 7.54 (1H, d, J= 2.6 Hz), 7.30 (1H, t, J= 2.1 Hz), 7.21 (1H, t, J= 7.9 Hz), 7.02 (3H, m), 6.78 (1H, s), 6.61 (1H, dd, J= 7.9 Hz, J= 2.1 Hz), 2.26 (3H, s).

7.3.344 N4-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940235)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-chloro-4-hydroxy-5-methylphenyl)-4-pyrimidineamine with 3-hydroxyaniline gave N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: retn, time: 18.20 min.; purity: 94%; MS (m/e): 360 (M⁺); ¹H NMR (DMSO-d₆): δ 9.26 (1H, s), 9.23 (1H, s), 9.16 (1H, s), 8.89 (1H, s), 8.14 (1H, d, J= 4.5 Hz), 7.66 (1H, d, J= 2.1 Hz), 7.60 (1H, d, J= 2.1 Hz), 7.29 (1H, d, J= 8.4 Hz), 7.11 (1H, s), 7.06 (1H, t, J= 8.4 Hz), 6.41 (1H, d, J= 8.4 Hz), 2.30 (3H, s).

7.3.345 N2-(3,4-Dimethoxyphenyl)-5-fluoro-N4-[4-[3-(N-morpholinyl)propyl]oxyphenyl]-2,4-pyrimidinediamine (R940250)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-[4-[3-(N-morpholinyl)propyl]oxyphenyl]-4-pyrimidineamine with 3,4-dimethoxyaniline gave N2-(3,4-dimethoxyphenyl)-5-fluoro-N4-[4-[3-(N-morpholinyl)propyl]oxyphenyl]-2,4-pyrimidinediamine. LCMS: retn, time: 14.72 min.; purity: 94%; MS (m/e): 484 (MH⁺); ¹H NMR (CDCl₃): δ 7.89 (1H, d, J= 3.3 Hz), 7.47 (2H, d, J= 9 Hz), 7.22 (1H, d, J= 2.2 Hz), 6.93-6.76 (5H, m), 6.64 (1H, d, J= 2.2 Hz), 4.01 (2H, t, J= 5.6 Hz), 3.86 (3H, s), 3.72 (3H, s), 3.71 (4H, m), 2.58-2.44 (6H, m), 1.97 (2H, m).

7.3.346 N2-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-[4-[3-(N-morpholinyl)propyl]oxyphenyl]-2,4-pyrimidinediamine (R940251)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-[4-[3-(N-morpholinyl)propyl]oxyphenyl]-4-pyrimidineamine with 2-chloro-4-hydroxy-5-methylaniline gave N2-(2-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-[4-[3-(N-morpholinyl)propyl]oxyphenyl]-2,4-pyrimidinediamine. LCMS: retn, time: 15.19 min.; purity: 94%; MS (m/e): 488 (MH⁺); ¹H NMR (CDCl₃): δ 7.89 (1H, d, J= 3.3 Hz), 7.52 (1H, d, J= 2.5 Hz), 7.44 (2H, d, 8.7 Hz), 6.97 (1H, d, J= 2.5 Hz), 6.91 (2H, d, 9 Hz), 6.71 (1H, s), 6.64 (1H, 2.5 Hz), 4.03 (2H, t, J= 6.03 Hz), 3.74 (4H, t, J= 4.65 Hz), 2.60-2.43 (6H, m), 2.23 (3H, s), 1.49 (2H, m).

7.3.347 N4-(3,5-Dimethyl-4-hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940253)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-4-pyrimidineamine with ethyl 3-aminophenoxyacetate gave N4-(3,5-dimethyl-4-hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: retn, time: 21.79 min.; purity: 91 %; MS (m/e): 427 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.80 (1H, s), 8.30 (1H, s), 8.23 (1H, d, J= 4.5 Hz), 7.37-7.17 (5H, m), 6.66 (1H, d, J= 9 Hz), 4.73 (2H, s), 4.25 (2H, q, J= 7.2 Hz), 2.23 (6H, s), 1.29 (3H, t, J= 7.0 Hz).

7.3.348 N2-(3-*tert*-Butylphenyl)-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-4-pyrimidinediamine (R940266)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-4-pyrimidineamine with 3-*tert*-butylaniline gave N2-(3-*tert*-butylphenyl)-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-4-pyrimidinediamine. LCMS: retn, time: 28.17 min.; purity: 96 %; MS (m/e): 439 (M⁺), 440 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.40 (1H, s), 9.19 (1H, s), 8.21 (1H, d, J= 3.6 Hz), 7.78 (1H, d, J= 8.5 Hz) 7.60 (2H, m), 7.48 (1H, t, J= 2 Hz), 7.31 (1H, t, J= 8.5 Hz), 7.25 (1H, t, J= 8.5 Hz), 7.02 (1H, d, J= 8.5 Hz), 6.70 (1H, dd, J= 8.5 and 2 Hz), 4.79 (2H, s), 4.26 (2H, q, J= 7.2 Hz), 1.33 (9H, s), 1.29 (3H, t, J= 7.2 Hz).

7.3.349 5-Fluoro-N4-(3-isopropylphenyl)-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine and 5-fluoro-N2-(2-ethoxycarbonylbenzofur-5-yl)-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine R940284

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-isopropylphenyl)-4-pyrimidineamine and ethyl 3-aminophenoxyacetate were reacted to give the mixture of 5-fluoro-N4-(3-isopropylphenyl)-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine and 5-fluoro-N2-(2-ethoxycarbonylbenzofur-5-yl)-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine. (R = CO₂Me). LCMS: retn, time: 25.41 min.; purity: 60.61 %; MS (m/e): 411 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.38 (1H, s), 9.29 (1H, s),

8.20 (1H, d, J= 3.9 Hz), 7.85 (1H, d, J= 9.3 Hz), 7.58 (1H, t, J= 1.6 Hz), 7.43-7.33 (3H, m), 7.18 (1H, t, J= 8.2 Hz), 7.05 (1H, d, J= 7.8 Hz), 6.53 (1H, dd, J= 8.4 Hz, J= 2.1 Hz), 4.72 (2H, s), 3.79 (3H, s), 2.95 (1H, quint, J= 7.2 Hz), 1.26 (6H, d, J= 7.2 Hz) (R = CO₂Et)
 LCMS: retn, time: 26.99 min.; purity: 39 %; MS (m/e): 425 (MH⁺); ¹H NMR (DMSO-d₆):
 5 δ 9.38 (1H, s), 9.29 (1H, s), 8.20 (1H, d, J= 3.9 Hz), 7.85 (1H, d, J= 9.3 Hz), 7.58 (1H, t, J= 1.6 Hz), 7.43-7.33 (3H, m), 7.18 (1H, t, J= 8.2 Hz), 7.05 (1H, d, J= 7.8 Hz), 6.53 (1H, dd, J= 8.4 and 2.1 Hz), 4.71 (2H, s), 4.25 (2H, q, J= 7.2 Hz), 2.95 (1H, quint, J= 7.2 Hz), 1.31 (3H, t, J= 7.2 Hz), 1.26 (6H, d, J= 7.2 Hz).

10 **7.3.350 N4-(3-*tert*-Butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine**
R940281

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(3-*tert*-butylphenyl)-2-chloro-5-fluoro-4-pyrimidineamine and 2-methoxycarbonyl-5-aminobenzofuran were reacted to give N4-(3-*tert*-butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine.
 15 LCMS: retn, time: 26.76 min.; purity: 97 %; MS (m/e): 435 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.41 (2H, sl), 8.27 (1H, s), 8.21 (1H, d, J 3.9 Hz), 7.98 (1H, m), 7.77-7.60 (3H, m), 7.37 (1H, t, J 8.1 Hz), 7.22 (1H, d, J 8.1 Hz), 3.98 (3H, s), 1.34 (9H, s).

20 **7.3.351 5-fluoro-N4-(3-isopropylphenyl)-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine**
R940283

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-isopropylphenyl)-4-pyrimidineamine and 2-methoxycarbonyl-5-aminobenzofuran were reacted to give 5-fluoro-N4-(3-isopropylphenyl)-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine.
 25 LCMS: retn, time: 26.05 min.; purity: 99 %; MS (m/e): 420 (M⁺), 422 (MH⁺); ¹H NMR (DMSO-d₆): δ 10.00 (1H, s), 9.95 (1H, s), 8.31 (1H, d, J= 4.8 Hz), 8.11 (1H, s), 7.74 (3H, m), 7.35 (1H, s), 7.35 (1H, t, J= 7.2 Hz), 7.12 (1H, d, J= 7.2 Hz), 3.99 (3H, s), 2.83 (1H, sept, J= 6.9 Hz), 1.20 (6H, d, J= 6.9 Hz).

7.3.352 N2-(1,1-Dihydroisobenzofuran-1-one-6-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926786)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 6-amino-1,1-dihydroisobenzofuran-1-one gave N2-(1,1-dihydroisobenzofuran-1-one-6-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.20 (s, 1H), 9.85 (s, 1H), 8.22 (d, 1H, J= 4.8 Hz), 8.10 (d, 1H, J= 1.2 Hz), 7.86 (dd, 1H, J= 2.4 and 8.7 Hz), 7.54 (d, 1H, J= 8.4 Hz), 7.22 (d, 1H, J= 2.4 Hz), 7.13 (dd, 1H, J= 2.1 and 9 Hz), 6.81 (d, 1H, J= 8.7 Hz), 5.34 (s, 2H), 4.20 (s, 4H); LCMS: ret. time: 17.40 min.; purity: 83%; MS (m/e): 395 (MH⁺).

7.3.353 N2-[3-(3-Acetamidophenoxy)propyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926787)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 3-N-acetamidophenoxy-3-propylamine gave N2-[3-(3-acetamidophenoxy)propyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.45 (bs, 1H), 10.07 (s, 1H), 8.42 (s, 1H), 8.20 (s, 1H), 7.37 (d, 1H, J= 3 Hz), 7.31 (s, 1H), 7.20-7.05 (m, 3H), 6.83 (d, 1H, J= 9Hz), 6.53 (d, 1H, J= 6.6 Hz), 4.18 (s, 4H), 3.95 (t, 2H, J= 6 Hz), 2.48 (m, 2H), 2.07 (s, 3H), 1.96 (t, 3H, J= 7.8 Hz); LCMS: ret. time: 19.58 min.; purity: 99%; MS (m/e): 454 (MH⁺).

7.3.354 N2-[4-(4,5-Dichloro-1H-imidazol-1-yl)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926788)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 4,5-dichloro-1H-imidazoleamine gave N2-[4-(4,5-dichloro-1H-imidazol-1-yl)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.10 (s, 1H), 9.85 (s, 1H), 8.20 (d, 1H, J= 4.2 Hz), 8.01 (s, 1H), 7.78 (d, 1H, J= 8.7 Hz), 7.36 (d, 1H, J= 9 Hz), 7.25 (d, 1H, J= 3 Hz), 7.14 (dd, 1H, J= 2.1 and 9 Hz), 6.85 (d, 1H, J= 8.7 Hz); LCMS: ret. time: 23.59 min.; purity: 95%; MS (m/e): 474 (MH⁺).

7.3.355 N2-(2,4-Dimethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926789)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 2,4-dimethoxyaniline gave N2-(2,4-dimethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.35 (s, 1H), 8.14 (bd, 1H), 7.38 (d, 1H, J= 9 Hz), 7.23 (s, 1H), 7.09 (d, 1H, J= 8.7 Hz), 6.79 (d, 1H, J= 8.7 Hz), 6.66 (d, 1H, J= 2.4 Hz), 6.49 (dd, 1H, J= 2.4 and 9 Hz), 4.22 (s, 4H), 3.77 (s, 6H); LCMS: ret. time: 20.93 min.; purity: 98%; MS (m/e): 399 (MH⁺).

7.3.356 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(4-isopropylphenyl)-2,4-pyrimidinediamine (R926790)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 4-isopropylaniline gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(4-isopropylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.30 (s, 1H), 10.50 (s, 1H), 8.22 (d, 1H, J= 5.4 Hz), 7.37 (d, 1H, J= 8.4 Hz), 7.26 (d, 1H, J= 3 Hz), 7.18 (s, 1H), 7.15 (s, 1H), 7.06 (dd, 1H, J= 3.3 and 8.7 Hz), 6.81 (d, 1H, J= 8.7 Hz), 4.23 (s, 4H), 2.85 (sept., 1H, J= 7.2 Hz), 1.17 (d, 6H, J= 6.9 Hz); LCMS: ret. time: 24.91 min.; purity: 95%; MS (m/e): 381 (MH⁺).

7.3.357 N2-(3,5-Dimethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926791)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 3,4-dimethoxyaniline gave N2-(3,5-dimethoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.08 (s, 1H), 9.99 (s, 1H), 8.19 (m, 1H), 7.21 (d, 1H, J= 2.4 Hz), 7.14 (dd, 1H, J= 2.1 and 8.7 Hz), 6.79 (d, 1H, J= 9 Hz), 6.72 (s, 1H), 6.20 (d, 1H, J= 1.8 Hz), 4.21 (s, 4H); LCMS: ret. time: 21.19 min.; purity: 93%; MS (m/e): 399 (MH⁺).

7.3.358 N2-(2,5-Dimethyl-4-hydroxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926792)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 2,5-dimethyl-4-hydroxyaniline gave N2-(2,5-dimethyl-4-hydroxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.69 (d, 1H, J= 3.9 Hz), 7.16 (d, 1H, J= 2.4 Hz), 7.05 (d, 1H, J= 2.4 Hz), 7.02 (d, 1H, J= 1.2 Hz), 6.66 (s, 1H), 6.63 (s, 1H), 6.62 (s, 1H), 4.19 (s, 4H), 2.12 (s, 3H), 2.10 (s, 3H); LCMS: ret. time: 19.80 min.; purity: 90%; MS (m/e): 383 (MH⁺).

7.3.359 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(5-methyl-3-phenyl-4-oxazolyl)-2,4-pyrimidinediamine (R926793)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 5-methyl-3-phenyl-4-oxazolylamine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(5-methyl-3-phenyl-4-oxazolyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.80-7.65 (m, 2H), 7.45 (bd, 1H), 7.20 (m, 1H), 7.00 (m, 1H), 6.65 (bd, 1H), 4.20 (s, 4H), 2.35 (s, 3H); LCMS: ret. time: 20.61 min.; purity: 78%; MS (m/e): 420 (MH⁺).

7.3.360 N4-(3,5-Dimethoxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926795)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,5-dimethoxyphenyl)-5-fluoro-4-pyrimidineamine with ethyl-3-aminophenoxyacetate gave N4-(3,5-dimethoxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 21.02 min.; purity: 84%; MS (m/e): 429 (MH⁺).

7.3.361 N4-(3,4-Ethylenedioxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926797)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-

dimethoxyphenyl)-5-ethoxycarbonyl-4-pyrimidineamine with ethyl-3-aminophenoxyacetate gave N4-(3,4-ethylenedioxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine. LCMS: ret. time: 27.60 min.; purity: 82%; MS (m/e): 495 (MH⁺).

5 **7.3.362 N4-(3-Hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926798)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-4-pyrimidineamine with ethyl-3-aminophenoxyacetate gave N4-(3-hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine. LCMS: ret. time: 24.78 min.; purity: 85%; MS (m/e): 453 (MH⁺).

15 **7.3.363 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine (R926614)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 2-methoxycarbonyl-5-aminobenzofuran gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 9.42 (s, 1H), 9.33 (s, 1H), 9.23 (s, 1H), 8.26 (s, 1H), 8.09 (d, 1H, J= 3.6 Hz), 7.59 (m, 3H), 7.13 (m, 3H), 6.53 (d, 1H, J= 7.5 Hz), 3.87 (s, 3H), 3.87 (s, 3H).

25 **7.3.364 N2-(2-Ethoxycarbonylindol-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926615)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 2-ethoxycarbonyl-5-aminoindole gave N2-(2-ethoxycarbonylindol-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.95 (d, 1H), 7.84 (d, 1H, J= 3.9 Hz), 7.34 (s, 1H), 7.33 (d, 1H, J= 1.8 Hz), 7.22-7.19 (m, 2H), 7.11-7.05 (m, 2H), 6.55 (m, 1H), 4.62 (s, 2H), 4.38 (q, 1H, J= 6.9 Hz), 1.40 9t, 3H, J= 7.5 Hz).

7.3.365 N2-[4-(4,5-Dichloro-1H-imidazol-1-yl)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926777)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with (4,5-dichloro-1H-imidazol-1-yl)-4-aniline gave N2-[4-(4,5-dichloro-1H-imidazol-1-yl)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 22.09 min.; purity: 98%; MS (m/e): 431 (MH⁺).

7.3.366 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(4-isopropylphenyl)-2,4-pyrimidinediamine (R926778)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 4-isopropylaniline gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-(4-isopropylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 23.08 min.; purity: 99%; MS (m/e): 439 (MH⁺).

7.3.367 5-Fluoro N4-(3-hydroxyphenyl)-N2-(5-methyl-4-oxazolyl-2-phenyl)-2,4-pyrimidinediamine (R926779)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 5-methyl-4-oxazolyl-2-phenyl-1-amine gave 5-fluoro N4-(3-hydroxyphenyl)-N2-(5-methyl-4-oxazolylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 23.08 min.; purity: 99%; MS (m/e): 439 (MH⁺). LCMS: ret. time: 19.17 min.; purity: 81%; MS (m/e): 378 (MH⁺).

7.3.368 N2-(3,5-Dimethoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926780)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3,5-dimethoxyaniline gave N2-(3,5-dimethoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 19.61 min.; purity: 97%; MS (m/e): 357 (MH⁺).

7.3.369 N4-(4-tert-Butoxycarbonylmethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R926572)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine with methyl 4-aminophenoxyacetate gave N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.49 (d, 2H, J= 8.7 Hz), 7.40 (d, 2H, J= 9.3 Hz), 6.89 (d, 2H, J= 9.3 Hz), 6.85 (d, 2H, J= 8.7 Hz), 4.62 (s, 2H), 4.52 (s, 2H), 3.81 (s, 3H), 1.49 (s, 9H); LCMS: ret. time: 24.68 min.; purity: 100%; MS (m/e): 499 (MH⁺).

7.3.370 5-Fluoro-N4-(3-isopropoxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine (R926487)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-isopropoxyphenyl)-4-pyrimidinediamineamine with 2-methoxycarbonyl-5-aminobenzofuran gave 5-fluoro-N4-(3-isopropoxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.09 (d, 1H, J= 2.4 Hz), 7.96 (d, 1H, J= 3 Hz), 7.52 (s, 1H), 7.48 (t, 1H, J= 1.8 Hz), 7.40 (dd, 1H, J= 6.3 Hz), 7.24 (m, 2H), 7.10 (m, 1H), 6.97 (bs, 1H), 6.74 (d, 1H, J= 2.4 Hz), 6.68 (dd, 1H, J= 2.1 and 6.9 Hz), 4.49 (sept., 1H, J= 5.7 Hz), 3.98 (s, 3H), 1.30 (d, 6H, J= 5.7 Hz); LCMS: ret. time: 25.86 min.; purity: 94%; MS (m/e): 437 (MH⁺).

7.3.371 N4-(4-tert-Butylphenyl)-N2-(2-ethoxycarbonylindol-5-yl)-5-fluoro-2,4-pyrimidinediamine (R926474)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of N4-(tert-butylcarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine with 2-ethoxycarbonyl-5-aminolindole gave N4-(4-tert-butylphenyl)-N2-(2-ethoxycarbonylindol-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.05 (d, 1H, J= 1.8 Hz), 7.85 (d, 1H, J= 3.9 Hz), 7.58 (d, 2H, J= 9 Hz), 7.36-7.10 (m, 4H), 7.03 (s, 1H), 6.95 (bd, 1H), 6.84 (dd, 1H, J= 7.2 Hz), 4.36 (q, 2H, J= 7.2 Hz), 1.40 (t, 3H, J= 7.5 Hz), 1.33 (s, 9H); LCMS: ret. time: 28.67 min.; purity: 100%; MS (m/e): 449 (MH⁺).

7.3.372 N4-(4-tert-Butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine (R926477)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of N4-(tert-butylcarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine with 2-methoxycarbonyl-5-aminobenzofuran gave N4-(4-tert-butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.6 (s, 1H), 8.09 (d, 1H, J= 1.8 Hz), 7.86 (d, 1H, J= 3.3 Hz), 7.54-7.36 (m, 6H), 6.90 (m, 1H), 3.97 (s, 3H), 1.36 (s, 9H), ¹⁹F NMR (CDCl₃): - 47188; LCMS: ret. time: 29.69 min.; purity: 84%; MS (m/e): 393 (M-41).

7.3.373 N2-(3,4-Ethylenedioxyphenyl)-N4-(2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine (R926485)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidineamine with 2-methoxycarbonyl-5-aminobenzofuran gave N2-(3,4-ethylenedioxyphenyl)-N4-(2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.07 (s, 1H), 7.76 (s, 1H), 7.44 (m, 3H), 7.13 (m, 1H), 6.68 (m, 2H), 4.18 (s, 4H), 3.95 (s, 3H); LCMS: ret. time: 26.63 min.; purity: 100%; MS (m/e): 437 (MH⁺).

7.3.374 N4-(3-Ethoxycarbonylmethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926774)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidineamine with 3,4-ethylenedioxyaniline gave N4-(3-ethoxycarbonylmethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.92 (d, 1H, J= 3.6 Hz), 7.67 (s, 1H), 7.40 (s, 1H), 7.28-7.21 (m, 2H), 7.01-6.96 (m, 2H), 6.80 (m, 2H), 6.68 (bd, 1H, 1H), 4.61 (s, 2H), 4.25 (m, 6H), 1.25 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 22.03 min.; purity: 84%; MS (m/e): 441 (MH⁺).

7.3.375 N4-(3-Ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926775)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidineamine with 3-hydroxyaniline gave N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 19.50 min.; purity: 84%; MS (m/e): 399 (MH⁺).

7.3.376 N4-(4-Aminocarbonylmethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945171)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(4-aminocarbonylmethyleneoxyphenyl)-5-fluoro-4-pyrimidineamine and 3,4-ethylenedioxyaniline gave N4-(4-aminocarbonylmethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (acetone-d₆): δ 4.24-4.31 (m, 4H), 4.51 (s, 2H), 6.77 (d, J= 8.7 Hz, 1H), 6.95 (dm, J= 8.7 Hz, 1H), 7.06 (d, J= 9.3 Hz, 2H), 7.28 (m, 1H), 7.71 (d, J= 9.0 Hz, 2H), 8.15 (m, 1H); LCMS: 15.23 min, 97.05%; MS (m/e): 412.01 (MH⁺).

7.3.377 (R935019): 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[di-(4-chlorophenyl)methyl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-4-pyrimidinediamine, 3-aminophenol and N-(2-chloro-5-fluoro-pyrimidinyl)-1,1-di(4-chlorophenyl)methylamine produced 5-fluoro-N2-(3-hydroxyphenyl)-N4-[di-(4-chlorophenyl)methyl]-2,4-pyrimidinediamine. LCMS: ret. time: 25.59 min.; purity: 91%; MS (m/e): 421 (MH⁺-Cl).

7.3.378 (R935020): N4-(Fluoren-9-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine:

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-4-pyrimidinediamine, 2-chloro-N-(fluoren-9-yl)-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to produce N4-(fluoren-9-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.85 (d, 1H, J= 2.9 Hz), 7.74 (d, 2H, J= 7.6 Hz), 7.64 (d, 2H, J= 7.6 Hz), 7.41-7.28 (m, 6H), 7.14-7.05 (m, 2H),

6.56 (d, 1H, J= 8.8 Hz), 5.28 (d, 1H, J= 8.8 Hz); LCMS: ret. time: 23.27 min.; purity: 89%; MS (*m/e*): 385 (MH⁺).

7.3.379 (R935021): (+)-5-Fluoro-N4-[1-(4-fluorophenyl)ethyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine

5 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-4-pyrimidinediamine, 3-aminophenol and (+)-N-(2-chloro-5-fluoropyrimidinyl)-1-(4-fluorophenyl)ethylamine were reacted to produce the desired (+)-5-fluoro- N4-[1-(4-fluorophenyl)ethyl]-N2-(3-hydroxyphenyl)-2, 4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.79 (d, 1H, J= 3.3 Hz), 7.38-7.34 (dd, 2H, J= 5.2 and 8.5 Hz), 7.14 (t, 1H, J= 4.5 Hz), 7.09 (d, 1H, J= 8.5 Hz), 7.03 (d, 1H, J= 8.5 Hz), 6.84 (br s, 1H), 6.84-6.78 (ddd, 1H, J= 0.8, 2.0, and 8.2 Hz), 6.46-6.42 (ddd, 1H, J= 0.8, 2.0 and 8.2 Hz), 5.26 (overlapped dq, 1H, J= 7.1 and 7.9 Hz), 5.18 (d, 1H, J=7.1 Hz), 1.59 (d, 3H, J= 7.1 Hz); LCMS: ret. time: 21.52 min.; purity: 92%; MS (*m/e*): 343 (MH⁺).

7.3.380 (R935023): (+)-5-Bromo-N4-[1-(4-fluorophenyl)ethyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine

15 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-4-pyrimidinediamine, 3-aminophenol and (+)-5-bromo-2-chloro-N4-[1-(4-fluorophenyl)ethyl]-4-pyrimidineamine were reacted to produce (+)-5-bromo- N4-[1-(4-fluorophenyl)ethyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.97 (s, 1H), 7.36-7.31 (m, 2H), 7.17 (s, 1H), 7.09-7.01 (m, 4H), 6.82 (dd, 1H, J= 2.2 and 8.2 Hz), 6.46 (d, 1H, J= 2.2 and 8.2 Hz), 5.50 (br d, 1H, J= 7.0), 5.27 (overlapped dq, 1H, J= 7.1 and 7.9 Hz), 1.58 (d, 3H, J= 7.0 Hz); LCMS: ret. time: 22.64 min.; purity: 94%; MS (*m/e*): 404 (MH⁺)

7.3.381 (R935025): 5-Bromo-N2-(3-hydroxyphenyl)-N4-(N-methyl-2-carbomethoxypyrrol-4-yl)-2,4-pyrimidinediamine

25 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-4-pyrimidinediamine, 3-aminophenol and 5-bromo-2-chloro-N-(N-methyl-2-carbomethoxypyrrol-4-yl)-4-pyrimidineamine were reacted to give 5-bromo-N2-(3-hydroxyphenyl)-N4-(N-methyl-5-carbomethoxypyrrol-4-yl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃ + CD₃OD): δ 7.92 (s, 1H), 7.58 (d, 1H, J= 8.0 Hz), 7.09 (d, 1H, J= 8.5 Hz), 7.04 (d, 1H, J= 8.5 Hz), 6.90 (d, 1H, J= 4.5 Hz), 6.81 (d, 1H, J= 1.8 Hz), 6.5 (m, 1H), 3.82 (s, 3H), 3.75 (s, 3H); LCMS: ret. time: 19.73 min.; purity: 90%; MS (*m/e*): 419 (MH⁺)

7.3.382 (R935029): 4-Amino-5-bromo-N2-(3-hydroxyphenyl)-2-pyrimidineamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-4-pyrimidinediamine, 4-amino-5-bromo-2-chloropyrimidine and 3-aminophenol were reacted to give 4-amino-5-bromo-N2-(3-hydroxyphenyl)-2-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 10.33 (br s, 1H), 8.27 (s, 1H), 7.14-6.06 (m, 2H), 7.01 (d, 1H, J= 1.7 Hz), 6.54 (td, 1H, J= 1.7 Hz and 7.0 Hz).

7.3.383 R935134: 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

The reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 5-(4-aminophenoxyethyl)-3-phenyl-1,2,4-oxadiazole were reacted in microwave at 180 °C for 10-20 minutes at 20 bar. Upon concentration and addition of 2N HCl provided 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.21 (br s, 1H), 9.91 (br s, 1H), 8.18 (d, 1H, J= 5.2 Hz), 8.03-7.99 (m, 2H), 7.61-7.53 (m, 3H), 7.46 (br d, 2H, J= 7.9 Hz), 7.14-7.01 (m, 5H), 6.54 (app d, 1H, J= 7.96 Hz), 5.56 (s, 2H); LCMS: ret. time: 24.61 min.; purity: 100%; MS (*m/e*): 471 (MH⁺).

7.3.384 R935135: 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine and 5-(4-aminophenoxyethyl)-3-phenyl-1,2,4-oxadiazole were reacted to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 10.21 (br s, 1H), 9.93 (br s, 1H), 8.17 (d, 1H, J= 5.2 Hz), 8.02-7.98 (m, 2H), 7.60-7.49 (m, 5H), 7.42 (app d, 2H, J= 7.0 Hz), 7.04 (d, 2H, J= 9.4 Hz), 6.89 (app d, 2H, J= 9.4 Hz), 5.56 (s, 2H), 4.58 (septet, 1H, J= 6.4 Hz), 1.23 (app d, 6H, J= 6.4 Hz); LCMS: ret. time: 26.90 min.; purity: 97%; MS (*m/e*): 513 (MH⁺).

7.3.385 R935136: N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxy)phenyl-4-pyrimidineamine and 5-(4-aminophenoxymethyl)-3-phenyl-1,2,4-oxadiazole were reacted provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 10.18 (br s, 1H), 9.12 (br s, 1H), 8.14 (d, 1H, 4.7 Hz), 8.02-7.97 (m, 2H), 7.65-7.52 (m, 3H), 7.44 (d, 2H, J= 8.8 Hz), 7.25-7.23 (m, 1H), 7.15-7.08 (m, 1H), 7.03 (d, 2H, J= 8.8 Hz), 6.81 (d, 1H, J= 8.8 Hz), 5.56 (s, 2H), 4.24-4.20 (m, 4H); LCMS: ret. time: 26.90 min.; purity: 97%; MS (*m/e*): 513 (MH⁺).

7.3.386 R935137: 5-Fluoro-N4-(2-methoxycarbonylbenzofura-5-yl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(2-methoxycarbonylbenzofura-5-yl)-4-pyrimidineamine and 5-(4-aminophenoxymethyl)-3-phenyl-1,2,4-oxadiazole were reacted to provide 5-fluoro-N4-(2-methoxycarbonylbenzofura-5-yl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.21 (br s, 1H), 9.79 (br s, 1H), 8.19 (d, 1H, J= 4.7 Hz), 8.09 (br s, 1H), 7.99 (dd, 2H, J= 2.3 and 8.2 Hz), 7.76-7.67 (m, 2H), 7.59-7.52 (m, 4H), 7.44 (d, 2H, J= 8.8 Hz), 7.02 (d, 2H, J= 8.8 Hz), 5.55 (s, 2H), 3.85 (s, 3H); LCMS: ret. time: 27.61 min.; purity: 92%; MS (*m/e*): 553 (MH⁺).

7.3.387 R935138: 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 3-aminophenol were reacted to provide 5-fluoro-N2-(3-hydroxyphenyl)-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 8.12 (d, 1H, J= 4.7 Hz), 8.03-7.99 (m, 2H), 7.69 (dd, 2H, J=

3.5 and 8.8 Hz), 7.61-7.53 (m, 3H), 7.06 (d, 2H, J= 9.9 Hz), 6.98 (m, 3H), 6.38 (br s, 1H), 5.58 (s, 2H). LCMS: ret. time: 24.83 min.; purity: 96%; MS (*m/e*): 471 (MH⁺).

7.3.388 R935139: 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 4-isopropoxyaniline were reacted to provide 5-fluoro- N2-(4-isopropoxyphenyl)-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 10.21 (br s, 1H), 9.78 (br s, 1H), 8.13 (d, 1H, J= 4.7Hz), 8.02-7.98 (m 2H), 7.65-7.53 (m, 5H), 7.34 (d, 2H, J= 7.6 Hz), 7.07 (d, 2H, J= 9.3 Hz), 6.86 (d, 2H, J= 8.8 Hz), 5.59 (s, 2H), 4.54 (sept, 1H, J= 5.8 Hz), 1.22 (d, 6H, J= 5.8 Hz); LCMS: ret. time: 29.64 min.; purity: 97%; MS (*m/e*): 513 (MH⁺).

7.3.389 R935140: N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to provide N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.31 (br s, 1H), 9.59 (br s, 1H), 8.11 (d, 1H, J= 4.7 Hz), 8.03-7.99 (m, 2H), 7.68-7.49 (m, 5H), 7.14-7.08 (m, 1H), 7.06 (d, 2H, J= 8.8 Hz), 6.90 (d, 1H, J= 8.8 Hz), 6.76 (d, 1H, J= 8.8 Hz), 5.59 (s, 2H), 4.22-4.17 (m, 4H); LCMS: ret. time: 21.35 min.; purity: 95%; MS (*m/e*): 513 (MH⁺).

7.3.390 R935141: 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine:

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 5-(4-aminophenoxymethyl)-3-methyl-1,2,4-oxadiazole were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-

methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 10.91 (br s, 1H), 9.91 (br s, 1H), 8.18 (d, 1H, J= 4.7 Hz), 7.43 (d, 2H, J= 8.8 Hz), 7.15-7.04 (m, 3H), 6.96 (d, 2H, J= 8.8 Hz), 6.58 (app d, 1H, J= 7.6 Hz), 5.43 (s, 2H), 2.34 (s, 3H); LCMS: ret. time: 18.68 min.; purity: 95%; MS (*m/e*): 409 (MH⁺).

7.3.391 R935142: 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine and 5-(4-aminophenoxymethyl)-3-methyl-1,2,4-oxadiazole were reacted to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 8.16 (d, 1H, J= 5.2 Hz), 7.52 (dd, 2H, J= 3.5 Hz and 9.3 Hz), 7.40 (d, 2H, J= 8.8 Hz), 6.98 (d, 2H, J= 8.8 Hz), 6.88 (d, 2H, J= 9.3 Hz), 5.44 (s, 2H), 4.58 (sept, 1H, J= 5.8 Hz), 2.34 (s, 3H), 1.24 (d, 6H, J= 5.8 Hz); LCMS: ret. time: 24.47 min.; purity: 93%; MS (*m/e*): 451 (MH⁺).

7.3.392 R935143: N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxy)phenyl-4-pyrimidineamine and 5-(4-aminophenoxymethyl)-3-methyl-1,2,4-oxadiazole were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 9.12 (br s, 1H), 9.04 (br s, 1H), 7.99 (d, 1H, J= 3.5 Hz), 7.55 (d, 2H, J= 1.7 and 8.8 Hz), 7.30 (d, 1H, J= 2.9 Hz), 7.17 (td, 1H, J= 2.9 and 8.8 Hz), 6.88 (d, 2H, J= 8.8 Hz), 6.77 (d, 1H, J= 8.8 Hz), 5.38 (s, 2H), 4.24-4.20 (m, 4H), 2.34 (s, 3H); LCMS: ret. time: 21.34 min.; purity: 97%; MS (*m/e*): 451 (MH⁺).

7.3.393 R935144: 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 4-isopropoxyaniline were reacted to provide 5-fluoro-N2-(4-isopropoxyphenyl)-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 10.11 (br s, 1H), 9.72 (br s, 1H), 8.12 (s, 1H, J= 5.3 Hz), 7.61 (dd, 2H, J= 8.8 Hz), 7.34 (d, 2H, J= 7.3 Hz), 7.01 (d, 2H, J= 8.8 Hz), 6.84 (d, 2H, J= 8.8 Hz), 5.47 (s, 2H), 4.54 (septet, 1H, J= 5.8 Hz), 2.34 (s, 3H), 1.23 (d, 6H, J= 6.4 Hz); LCMS: ret. time: 24.31 min.; purity: 96%; MS (m/e): 451 (MH⁺).

7.3.394 R935145: N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to provide N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.81 (br s, 1H), 9.67 (br s, 1H), 8.13 (d, 1H, J= 4.7 Hz), 7.63 (dd, 2H, J= 4.1 and 8.8 Hz), 7.07 (m, 1H), 7.00 (d, 2H, J= 8.8 Hz), 6.89 (d, 1H, J= 8.8 Hz), 6.76 (d, 1H, J= 8.8 Hz), 5.46 (s, 2H), 4.22-4.18 (m, 4H), 2.34 (s, 3H); LCMS: ret. time: 21.54 min.; purity: 97%; MS (m/e): 451 (MH⁺).

7.3.395 R935146: 5-Fluoro-N2-(2-methoxycarbonylbenzofura-5-yl)-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 2-methoxycarbonyl-5-aminobenzofuran were reacted to provide 5-fluoro-N2-(2-methoxycarbonylbenzofura-5-yl)-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.14 (d, 1H, J=

4.7 Hz), 8.02 (s, 1H), 7.63-7.56 (m, 5H), 7.02 (d, 2H, $J = 8.8$ Hz), 5.47 (s, 2H), 3.85 (s, 3H), 2.34 (s, 3H); LCMS: ret. time: 22.46 min.; purity: 97%; MS (m/e): 491 (MH^+).

7.3.396 R935147: 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 3-hydroxyaniline were reacted to provide 5-fluoro-N2-(3-hydroxyphenyl)-N4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine as fine flakes of the product. ^1H NMR (DMSO- d_6): δ 8.11 (d, 1H, $J = 4.6$ Hz), 7.66 (d, 2H, $J = 5.8$ Hz), 7.06-6.97 (m, 5H), 6.42-4.0 (m, 1H), 5.46 (s, 2H), 2.35 (s, 3H); LCMS: ret. time: 19.00 min.; purity: 95%; MS (m/e): 409 (MH^+).

7.3.397 R935148: N2-(3,4-Ethylenedioxyphenyl)-N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-Chloro--[4-[ethoxycarbonyl(dimethyl)methyl]phenyl]-5-fluoro-2, 4-pyrimidine amine and 3,4-ethylenedioxyaniline were reacted to produce N2-(3,4-ethylenedioxyphenyl)-N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 9.31 (s, 1H), 8.97 (s, 1H), 8.03 (d, 1H, $J = 3.5$ Hz), 7.70 (d, 2H, $J = 8.8$ Hz), 7.29 (d, 1H, $J = 2.3$ Hz), 7.23 (d, 2H, $J = 8.8$ Hz), 6.98 (dd, 1H, $J = 2.1$ and 8.8 Hz), 6.66 (d, 1H, 8.2 Hz); 4.19-4.15 (m, 4H), 4.07 (qt, 2H, $J = 7.0$ Hz), 1.48 (s, 6H), 1.10 (t, 3H, $J = 7.0$ Hz); LCMS: ret. time: 24.51 min.; purity: 100%; MS (m/e): 453 (MH^+).

7.3.398 R935150: N2-[4-[(1-Ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (or it can be prepared similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine), 2-chloro-5-fluoro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine and 4-[ethoxycarbonyl(dimethyl)methyl]aniline were reacted to produce N2-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-

pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.18 (br s, 1H), 9.11 (br s, 1H), 8.01 (d, 1H, J= 3.5 Hz), 7.56 (d, 2H, J= 8.8 Hz), 7.54 (d, 2H, J= 8.8 Hz), 7.09 (d, 2H, J= 8.8 Hz), 6.86 (d, 2H, J= 8.8 Hz), 4.56 (sept, 1H, J= 5.8 Hz), 4.02 (qt, 2H, J= 7.0 Hz), 1.43 (s, 6H), 1.26 (d, 6H, J= 7.0 Hz), 1.09 (t, 3H, J= 7.0 Hz); LCMS: ret. time: 28.49 min.; purity: 98%; MS (m/e): 453 (MH⁺).

7.3.399 R935179: N2-[4-(2,3-Dihydroxypropoxy)phenyl]- N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine and 4-(2,3-dihydroxypropoxy)aniline were reacted to produce N2-[4-(2,3-dihydroxypropoxy)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.09 (s, 1H), 8.95 (s, 1H), 7.98 (d, 1H, J= 3.5 Hz), 7.51 (d, 2H, J= 8.8 Hz), 7.32 (d, 1H, J= 2.3 Hz), 7.17 (dd, 1H, J= 2.3 and 8.8 Hz), 6.77 (dd, 3H, J= 8.8 Hz), 4.90 (d, 1H, J= 5.3 Hz), 4.64 (t, 1H, J= 5.8 Hz), 4.23-4.19 (m, 4H), 3.91-3.89 (m, 1H), 3.80-3.73 (m, 2H), 3.41 (t, 2H, J= 5.3 Hz); LCMS: ret. time: 15.04 min.; purity: 96%; MS (m/e): 429 (MH⁺).

7.3.400 R935180: N2-[4-(2,3-Dihydroxypropoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine and 4-(2,3-dihydroxypropoxy)aniline were reacted to produce N2-[4-(2,3-dihydroxypropoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.38 (s, 1H), 9.18 (s, 1H), 8.98 (s, 1H), 8.12 (d, 1H, J= 3.5 Hz), 7.58 (d, 2H, J= 8.8 Hz), 7.22 (d, 1H, J= 2.3 Hz), 7.12 (dd, 2H, J= 2.3 and 8.8 Hz), 6.79 (d, 2H, J= 8.8 Hz), 6.45 (d, 1H, J= 8.8 Hz), 4.91 (d, 1H, J= 5.3 Hz), 4.65 (t, 1H, J= 5.8 Hz), 3.92-3.89 (m, 1H), 3.79-3.74 (m, 2H), 3.44 (t, 2H, J= 5.3 Hz); LCMS: ret. time: 12.79 min.; purity: 89%; MS (m/e): 387 (MH⁺).

7.3.401 R935175: N2-[4-(2,3-Dihydroxypropoxy)phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine and 4-(2,3-dihydroxypropoxy)aniline were reacted to produce N2-[4-(2,3-

dihydroxypropoxy)phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine: ^1H NMR (DMSO- d_6): δ 9.12 (s, 1H), 8.91 (s, 1H), 7.97 (d, 1H, $J = 3.5$ Hz), 7.58 (d, 2H, $J = 8.8$ Hz), 7.49 (d, 2H, $J = 8.8$ Hz), 6.85 (d, 2H, $J = 8.8$ Hz), 6.76 (d, 2H, $J = 8.8$ Hz); 4.89 (d, 1H, $J = 4.7$ Hz), 4.63 (t, 1H, $J = 5.2$ Hz), 4.56 (septet, 1H, $J = 5.8$ Hz), 3.90-3.89 (m, 1H), 3.76-3.73 (m, 2H), 3.41 (t, 2H, $J = 5.3$ Hz), 1.25 (d, 6H, $J = 5.8$ Hz); LCMS: ret. time: 17.48 min.; purity: 98%; MS (m/e): 429 (MH^+).

7.3.402 R935169: N4-[4-[(1-Ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to produce N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 7.87 (d, 1H, $J = 3.5$ Hz), 7.56 (d, 2H, $J = 8.8$ Hz), 7.35 (d, 2H, $J = 8.8$ Hz), 7.25-7.23 (m, 1H), 7.08 (t, 1H, $J = 8.2$ Hz), 6.91 (d, 1H, $J = 2.3$ Hz), 6.83 (d, 1H, $J = 7.6$ Hz), 6.50 (dd, 1H, $J = 1.7$ and 8.2 Hz), 4.13 (qt, 2H, $J = 7.0$ Hz), 1.58 (s, 6H), 1.19 (t, 3H, $J = 7.0$ Hz); LCMS: ret. time: 22.97 min.; purity: 98%; MS (m/e): 411 (MH^+).

7.3.403 R935164: 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[(N-methyl-2-methoxycarbonyl)pyrrol-4-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine, 2-chloro-5-fluoro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine and N-methyl-2-methoxycarbonyl-4-aminopyrrole hydrochloride with added diisopropylethylamine were reacted to produce the desired product 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[(N-methyl-2-carbomethoxy)pyrrol-4-yl]-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.87 (br s, 1H), 7.44 (d, 2H, $J = 8.8$ Hz), 7.08 (br s, 1H), 6.99-6.85 (m, 3H), 6.70 (d, 1H, $J = 2.3$ Hz), 6.63 (d, 1H, $J = 1.7$ Hz), 4.52 (septet, 1H, $J = 5.8$ Hz), 3.80 (s, 3H), 3.79 (s, 3H), 1.34 (d, 6H, $J = 5.8$ Hz); LCMS: ret. time: 23.89 min.; purity: 99%; MS (m/e): 400 (MH^+).

7.3.404 R935165: 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-[(N-methyl-2-carbomethoxy)pyrrole-4-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-2-

carbomethoxypyrrol-4-yl)-4-pyrimidineamine and 4-isopropoxyaniline were reacted to produce 5-fluoro-N2-(4-isopropoxyphenyl)-N4-[(N-methyl-5-carbomethoxy)pyrrol-4-yl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.84 (d, 1H, J= 2.3 Hz), 7.36 (d, 2H, J= 8.8 Hz), 7.22 (d, 1H, J= 1.1 Hz), 6.87 (d, 2H, J= 8.8 Hz), 6.84 (s, 1H), 6.77 (d, 1H, J= 1.7 Hz), 6.61 (br s, 1H), 4.49 (septet, 1H, J= 5.8 Hz), 3.82 (d, 3H), 3.81 (s, 3H), 1.33 (d, 6H, J= 5.8 Hz); LCMS: ret. time: 23.36 min.; purity: 96%; MS (*m/e*): 400 (MH⁺).

7.3.405 R935166: N2-(3,4-Ethylenedioxyphenyl)- 5-fluoro-N4-[(N-methyl-2-methoxycarbonyl)pyrrol-4-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-2-methoxycarbonylpyrrol-2-yl)-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to produce 5-fluoro-N2-(3,4-ethylenedioxyphenyl)-N4-[(N-methyl-2-carbomethoxy)pyrrol-4-yl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.84 (d, 1H, J= 3.5 Hz), 7.34 (s, 1H), 7.21 (s, 1H), 6.82 (d, 2H, J= 8.8 Hz), 6.76 (d, 2H, J= 8.8 Hz), 6.58 (s, 1H), 4.27-4.18 (m, 4H), 3.90 (s, 3H), 3.81 (s, 3H); LCMS: ret. time: 20.02 min.; purity: 93%; MS (*m/e*): 400 (MH⁺).

7.3.406 R935167: N4-[4-[(1-Ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[4-[1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-4-pyrimidineamine and 4-isopropoxyaniline were reacted to produce N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.29 (s, 1H), 8.95 (s, 1H), 8.02 (d, 1H, J= 4.1 Hz), 7.68 (d, 2H, J= 8.8 Hz), 7.46 (d, 2H, J= 8.8 Hz), 7.22 (d, 2H, J= 8.8 Hz), 6.75 (d, 2H, J= 8.8 Hz), 4.48 (septet, 1H, J= 5.8 Hz), 4.04 (qt, 2H, J= 7.0 Hz), 1.47 (s, 6H), 1.22 (d, 6H, J= 5.8 Hz), 1.10 (t, 3H, J= 7.0 Hz); LCMS: ret. time: 28.11 min.; purity: 99%; MS (*m/e*): 453 (MH⁺).

7.3.407 R935159: 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[3-hydroxyphenyl]-pyrimidine-2,4-diamine, 2-chloro-5-fluoro-N4-(4-isopropoxyphenyl)-4-

pyrimidineamine and methyl 4-aminophenoxyacetate were reacted to produce 5-fluoro-N4-(4-isopropoxyphenyl)-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.88 (d, 1H, J= 3.5 Hz), 7.46 (d, 2H, J= 8.8 Hz), 7.42 (d, 2H, J= 8.8 Hz), 6.88 (d, 2H, J= 9.3 Hz), 6.85 (d, 2H, J= 9.3 Hz), 6.78 (br s, 1H), 6.63 (br d, 1H, J= 2.3 Hz), 4.61 (s, 2H), 4.53 (septet, 1H, J= 6.4 Hz), 3.81 (s, 3H), 1.35 (d, 6H, J= 6.4 Hz); LCMS: ret. time: 23.19 min.; purity: 97%; MS (*m/e*): 427 (MH⁺).

7.3.408 R935157: N4-[4-[(1-Ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[3-hydroxyphenyl]-pyrimidine-2,4-diamine, 2-chloro-N4-[4-[1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-4-pyrimidineamine was reacted with 4-(methoxycarbonylmethyleneoxy)aniline to produce N4-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.92 (s, 1H), 7.55 (d, 2H, J= 8.7 Hz), 7.43 (d, 2H, J= 9.3 Hz), 7.33 (d, 2H, J= 8.7 Hz), 6.87 (d, 2H, J= 9.3 Hz), 6.79 (s, 1H), 6.73 (d, 1H, J= 2.3 Hz), 4.62 (s, 2H), 4.13 (qt, 2H, J= 7.0 Hz), 3.81 (s, 3H), 1.59 (s, 6H), 1.20 (t, 3H, 7.0 Hz); LCMS: ret. time: 25.20 min.; purity: 97%; MS (*m/e*): 483 (MH⁺).

7.3.409 R935152: N2-[4-[(1-Ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-[4-(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 4-[1-ethoxycarbonyl-1-methyl)ethyl]aniline were reacted to give N2-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.89 (d, 1H, J= 2.9 Hz), 7.24-7.10 (m, 5H), 6.93 (d, 1H, J= 7.6 Hz), 6.68 (d, 2H, J= 8.2 Hz), 4.08 (qt, 2H, J= 7.0 Hz), 1.52 (s, 3H), 1.49 (s, 3H), 1.16 (t, 3H, J= 7.0 Hz); LCMS: ret. time: 22.15 min.; purity: 96%; MS (*m/e*): 411 (MH⁺).

7.3.410 N2-(3-*tert*-Butylphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940257)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-

hydroxyphenyl)-4-pyrimidineamine with 3-*tert*-buthylaniline gave N2-(3-*tert*-butylphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 23.82 min.; purity: 100%; MS (m/e): 353 (MH⁺); ¹H NMR (CDCl₃): δ 7.96 (1H, d, J= 3 Hz), 7.61 (1H, ddd, J= 7.5, 2.2 and 0.9 Hz), 7.49 (1H, t, J= 2.5 Hz), 7.27 (1H, m), 7.18 (1H, t, J= 8.1 Hz), 7.99 (1H, m), 6.94 (1H, s), 6.91 (1H, dd, J= 7.5 and 2.5 Hz), 6.80 (1H, d, J= 7.5 Hz), 6.72 (2H, m), 6.58 (1H, ddd, J= 7.5, 2.5 and 0.9 Hz), 6.52 (1H, ddd, J= 7.5, 2.5 and 0.9 Hz), 1.28 (9H, s).

7.3.411 N4-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and N4-(3-chloro-4-hydroxy-5-methylphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940258)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-chloro-4-hydroxy-5-methylphenyl)-4-pyrimidineamine with ethyl 3-aminophenoxyacetate gave a mixture of N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and N4-(3-chloro-4-hydroxy-5-methylphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 20.34 min. (CO₂Me); purity: 17%; MS (m/e): 432 (M⁺); LCMS: ret. time: 21.83 min; purity 78%; MS (m/e): 446 (M⁺).

7.3.412 N2-(3-*tert*-Butylphenyl)-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940260)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3,4-dimethoxyphenyl)-4-pyrimidineamine with ethyl 3-*tert*-buthylaniline gave N2-(3-*tert*-butylphenyl)-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 24.87 min.; purity: 99%; MS (m/e): 397 (MH⁺); ¹H NMR (CDCl₃): δ 7.92 (1H, d, J= 3.4 Hz), 7.50 (1H, d, J= 8 Hz), 7.28 (1H, t, J= 2.3 Hz), 7.21 (1H, d, J= 8 Hz), 7.18 (1H, m), 7.08-7.01 (2H, m), 6.99 (1H, s), 6.84 (2H, d, J= 9.2 Hz), 6.65 (1H, s), 3.89 (3H, s), 3.72 (3H, s), 1.26 (9H, s).

7.3.413 N2-[2-(N-Benzylpiperazino)ethyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940261)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 4-(N-benzylpiperazino)ethylamine gave N2-[2-(N-benzylpiperazino)ethyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 17.15 min.; purity: 90 %; MS (m/e): 422 (M⁺), 423 (MH⁺); ¹H NMR (CDCl₃): δ 8.42 (1H, s), 7.82 (1H, d, J= 3.9 Hz), 7.32-7.08 (6H, m), 6.73 (1H, s), 6.61 (1H, dd, J= 8.1 and 2.1 Hz), 6.51 (1H, d, J= 7.5 Hz), 5.18 (1H, s), 3.59 (2H, m), 3.02 (2H, m), 2.71-2.41 (3H, m), 2.10-1.16 (5H, m).

7.3.414 N2-[2-(N-Benzylpiperazino)ethyl]-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940262)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3,4-dimethoxyphenyl)-4-pyrimidineamine with 4-(N-benzylpiperazino)ethylamine gave N2-[2-(N-benzylpiperazino)ethyl]-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 17.48 min.; purity: 99 %; MS (m/e): 466 (M⁺), 467 (MH⁺); ¹H NMR (CDCl₃): δ 7.82 (1H, d, J= 3.9 Hz), 7.44 (1H, s), 7.33-7.10 (6H, m), 7.04 (1H, dd, J= 8.9 and 2.5 Hz), 6.84 (1H, d, J= 8.9 Hz), 6.58 (1H, s), 5.40 (1H, s), 3.91 (3H, s), 3.87 (3H, s), 3.41 (2H, m), 2.87 (2H, m), 2.51 (3H, m), 1.80 (2H, m), 1.60 (4H, m), 1.30 (1H, m).

7.3.415 N2-[4-(N-Benzylpiperidino)]-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940263)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3,4-dimethoxyphenyl)-4-pyrimidineamine with N-benzyl-4-aminopiperidine gave N2-[4-(N-benzylpiperidino)]-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 15.52 min.; purity: 99 %; MS (m/e): 438 (MH⁺); ¹H NMR (CDCl₃): δ 7.81 (1H, d, 3.3 Hz), 7.35-7.18 (5H, m), 7.10 (1H, dd, J= 8.7 and 2.6 Hz), 6.84 (1H, d, J= 8.7 Hz), 6.56 (1H, s), 4.73 (1H, d, J= 6.9 Hz), 3.89 (6H, s), 3.75 (1H, m), 3.51 (2H, m), 2.81 (2H, m), 2.15 (2H, m), 2.00 (2H, m), 1.66-1.44 (4H, m).

7.3.416 N2-[4-(N-Benzylpiperidino)]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940264)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with N-benzyl-4-aminopiperidine gave N2-[4-(N-benzylpiperidino)]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 14.00 min.; purity: 96 %; MS (m/e): 394 (M⁺), 395 (MH⁺); ¹H NMR (CDCl₃): δ 7.81 (1H, d, J= 3.6 Hz), 7.40-7.28 (5H, m), 7.21-7.14 (2H, m), 6.69 (1H, m), 6.62 (1H, m), 6.59 (1H, m), 5.20 (1H, s), 3.65 (2H, s), 3.50 (1H, s), 3.03 (1H, m), 2.83 (1H, m), 2.13 (1H, m), 1.95-1.70 (1H, m), 1.58 (4H, m).

7.3.417 N4-(3-*tert*-Butylphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940270)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of N4-(3-*tert*-butylphenyl)-2-chloro-5-fluoro-4-pyrimidineamine with ethyl 3-aminophenoxyacetate gave N4-(3-*tert*-butylphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 27.30 min.; purity: 98 %; MS (m/e): 439 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.50 (1H, s), 9.33 (1H, s), 8.11 (1H, dd, J=4.2 and 1.8 Hz), 7.81 (1H, d, J= 7.2 Hz), 7.49 (1H, t, 2.4 Hz), 7.30-7.28 (3H, m), 7.14-7.03 (2H, m), 6.46 (1H, d, J= 7.8 Hz), 4.57 (2H, s), 4.13 (2H, q, J= 7.2 Hz), 1.23 (9H, s), 1.18 (3H, t, J= 7.2 Hz).

7.3.418 N4-(3-*tert*-Butylphenyl)-N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R940271)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of N4-(3-*tert*-butylphenyl)-2-chloro-5-fluoro-4-pyrimidineamine with 3-chloro-4-hydroxy-5-methylaniline gave N4-(3-*tert*-butylphenyl)-N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 25.46 min.; purity: 100 %; MS (m/e): 400 (M⁺); ¹H NMR (DMSO-d₆): δ 9.63 (1H, s), 9.30 (1H, s), 8.82 (1H, s), 8.20 (1H, d, J= 3.9 Hz), 7.92 (1H, d, J= 8.8 Hz), 7.58 (2H, m), 7.40-7.20 (3H, m), 2.22 (3H, s), 1.35 (9H, s).

7.3.419 N2-(3-*tert*-Butylcarbonylaminophenyl)-N4-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R940275)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-butylcarbonylaminoaniline gave N2-(3-*tert*-butylcarbonylaminophenyl)-N4-(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine.

LCMS: ret. time: 20.19 min.; purity: 91 %; MS (m/e): 396 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.42 (1H, s), 9.28 (1H, s), 9.21 (1H, s), 9.18 (1H, s), 8.17 (1H, d, J= 3.9 Hz), 7.90 (1H, s), 7.55 (1H, dt, J= 6.9 and 2.1 Hz), 7.51 (1H, dd, J= 7.8 and 1.5 Hz), 7.26-7.13 (4H, m), 6.57 (1H, dd, J= 7.5 and 1.5 Hz), 1.30 (9H, s).

7.3.420 N4-(3,3-Dihydroisobenzofuranyl-1-one-6-yl)-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine R940294

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,3-dihydroisobenzofuranyl-1-one-6-yl)-5-fluoro-4-pyrimidineamine and 2-methoxycarbonyl-5-aminobenzofuran were reacted to give N4-(3,3-dihydroisobenzofuranyl-1-one-6-yl)-5-fluoro-N2-(2-

methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine. LCMS: ret. time: 21.34 min.; purity: 97 %; MS (m/e): 434 (M⁺); ¹H NMR (DMSO-d₆): δ 9.90 (1H, s), 9.61 (1H, s), 8.4-8.12 (4H, m), 7.35-7.67 (4H, m), 5.50 (2H, s), 3.98 (3H, s).

7.3.421 N2-[3-Ethoxycarbonylmethyleneoxyphenyl]-N4-(3,3-dihydroisobenzofuranyl-1-one-6-yl)-5-fluoro-2,4-pyrimidinediamine R940285

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,3-dihydroisobenzofuranyl-1-one-6-yl)-5-fluoro-4-pyrimidineamine and ethyl 3-aminophenoxyacetate were reacted to give N2-(3-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,3-dihydroisobenzofuranyl-1-one-6-yl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 20.55 min.; purity: 76 %; MS (m/e): 438 (M⁺), 440 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.70 (1H, s), 9.30 (1H, s), 8.23-8.06 (1H, m), 8.05 (1H, s), 7.63 (1H, d, J= 8.1 Hz), 7.30 (1H, s), 7.22 (1H, m), 7.08 (1H, t, J= 8.1 Hz), 6.43 (1H, d, J= 8.1 Hz), 5.37 (1H, s), 5.37 (2H, s), 4.60 (2H, s), 4.13 (2H, q, J= 7.2 Hz), 1.18 (3H, t, J= 7.2 Hz).

7.3.422 N2-(3,5-Dimethoxyphenyl)-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926804)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-4-pyrimidineamine with 3,5-dimethoxyaniline gave N2-(3,5-dimethoxyphenyl)-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 24.12 min.; purity: 86%; MS (m/e): 443 (MH⁺).

7.3.423 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-trifluoromethylphenyl)-2,4-pyrimidinediamine (R926805)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 3-trifluoromethylaniline gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-trifluoromethylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 25.88 min.; purity: 89%; MS (m/e): 407 (MH⁺).

7.3.424 N2-(2-Ethoxycarbonylindol-7-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyridinediamine (R926808)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 2-ethoxycarbonyl-7-aminoindole gave N2-(2-ethoxycarbonylindol-7-yl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyridinediamine. LCMS: ret. time: 24.11 min.; purity: 88%, MS (m/e): 450 (MH⁺).

7.3.425 N4-[4-(4,5-Dichloro-1H-imidazol-1-yl)phenyl]-5-fluoro-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R926809)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of N-4-[4-(4,5-dichloro-1H-imidazol-1-yl)phenyl]-2-chloro-5-fluoro-4-pyrimidineamine with ethyl-3-aminophenoxyacetate gave N4-[4-(4,5-dichloro-1H-imidazol-1-yl)phenyl]-5-fluoro-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 25.22 min, purity: 77%; MS (m/e): 519 (MH⁺).

7.3.426 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(1,3-oxazol-5-yl)phenyl]-2,4-pyrimidinediamine (R926813)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 3-(1,3-oxazol-5-yl)aniline gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(1,3-oxazol-5-yl)phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 20.25 min.; purity: 81%, MS (m/e): 406 (MH⁺).

7.3.427 N2-(2-Ethoxycarbonylindol-7-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyridinediamine (R926814)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 2-ethoxycarbonyl-7-aminoindol gave N2-(2-ethoxycarbonylindol-7-yl)-5-fluoro N4-(3-hydroxyphenyl)-2,4-pyridinediamine. LCMS: ret. time: 25.94 min.; purity: 91%.

7.3.428 N2-(3-Aminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950207)

N4-(3,4-Ethylenedioxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (50 mg, 0.18 mmol) was dissolved in dry MeOH (1 ml), to it was added 3-aminoaniline (163 mg, 1.2 mmol) and the mixture was refluxed for 4 days (70 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃-Acetone, 9:1) to give N2-(3-aminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.66 (d, 1H, J= 3.6 Hz), 7.18 (d, 1H, J= 2.1 Hz), 7.09 (t, 1H, J= 2.1 Hz), 6.80-6.90, (m, 1H), 6.69 (d, 1H, J= 8.1 Hz), 6.57 (m, 1H), 6.20 (m, 1H), 6.60 (m, 1H), 4.10 (m, 4H); LCMS purity: 90.7%; MS (m/e): 354.13 (M⁺, 100).

7.3.429 N4-(3,4-Ethylenedioxyphenyl)-N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine (R950186)

In like manner to the preparation of N2-(3-aminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine and 3-ethoxycarbonylmethyleneaminophenylaniline were reacted to prepare N4-(3,4-ethylenedioxyphenyl)-N2-(3-

ethoxycarbonylmethyleneaminophenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 23.29 min.; purity: 95.7%; MS (m/e): 440.41 (MH⁺).

7.3.430 N4-(3,5-Dichloro-4-hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950185)

In like manner to the preparation of N2-(3-aminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2-chloro-N4-(3,5-dichlorophenyl-4-hydroxy)-5-fluoro-4-pyrimidineamine and ethyl 3-aminophenoxyacetate were reacted to prepare N4-(3,5-dichloro-4-hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 22.51 min.; purity: 96.1%; MS (m/e): 466.88 (MH⁺).

7.3.431 N4-(3-Aminophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofurane-5-yl)-2,4-pyrimidinediamine (R950162)

A mixture of N4-(3-aminophenyl)-2-chloro-5-fluoro-4-pyrimidineamine (10 mg, 0.06 mmol) and 2-methoxycarbonyl-5-aminobenzofuran (36 mg, 0.18 mmol) in dry MeOH (0.5 ml) was refluxed for 2 days (100 °C oil-bath temperature). The mixture was cooled to 22 °C, concentrated to dryness under reduced pressure and subjected to column chromatography on silica gel (CHCl₃:Acetone, 9:1) to give N4-(3-aminophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofurane-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.24 (s, 1H), 7.96 (dd, 1H, J= 1.7, 3.5 Hz), 7.46-7.59 (m, 3H), 6.93-6.99 (m, 2H), 6.84 (d, 1H, J= 8.2 Hz), 6.35 (m, 1H), 3.84 (s, 3H); LCMS purity: 97.8%; MS (ES) m/e 394.02 (M⁺, 70).

7.3.432 N4-(3-Aminophenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950163)

In like manner to the preparation of N4-(3-aminophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofurane-5-yl)-5-fluoro-2,4-pyrimidinediamine, N4-(3-aminophenyl)-2-chloro-5-fluoro-4-pyrimidineamine and 3-hydroxyaniline were reacted to prepare N4-(3-aminophenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 7.94 (d, 1H, J= 4.1 Hz), 7.20 (m, 2H), 6.89-7.00 (m, 4H), 6.30 (m, 2H); LCMS: ret. time: 11.92 min.; purity: 95.0%; MS (m/e): 312.09 (MH⁺).

7.3.433 N4-(3-Aminophenyl)-5-fluoro-N2-(3-isopropoxyphenyl)-2,4-pyrimidinediamine (R950164)

In like manner to the preparation of N4-(3-aminophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofurane-5-yl)-5-fluoro-2,4-pyrimidinediamine, N4-(3-aminophenyl)-2-chloro-5-fluoro-4-pyrimidineamine and 3-isopropoxyaniline were reacted to prepare N4-(3-aminophenyl)-5-fluoro-N2-(3-isopropoxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 17.52 min.; purity: 98.9%; MS (m/e): 354.13 (MH⁺).

7.3.434 N4-(3-Aminophenyl)-5-fluoro-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R950165)

In like manner to the preparation of N4-(3-aminophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofurane-5-yl)-5-fluoro-2,4-pyrimidinediamine, N4-(3-aminophenyl)-2-chloro-5-fluoro-4-pyrimidineamine and 4-isopropoxyaniline were reacted to prepare N4-(3-aminophenyl)-5-fluoro-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-D₆-MeOD, 300 MHz): δ 7.90 (d, 1H, J= 4.1 Hz), 7.47 (m, 2H), 7.03 (t, 1H, J= 1.7 Hz), 6.60-6.95 (m, 5H), 6.29 (m, 1H), 4.43 (septett, 1H, J= 6.0 Hz), 1.18 (d, 6H, J= 6.0 Hz); LCMS: ret. time: 17.11 min.; purity: 88.4%; MS (m/e): 354.09 (MH⁺).

7.3.435 N2-(3-Furylmethylene)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950210)

In like manner to the preparation of N4-(3-aminophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofurane-5-yl)-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-furylmethylamine were reacted to prepare N2-(3-furylmethylene)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 16.03 min.; purity: 93.5%; MS (m/e): 301.10 (MH⁺).

7.3.436 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(4-methoxyphenyloxyethyleneamino)-2,4-pyrimidinediamine (R950211)

In like manner to the preparation of N4-(3-aminophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofurane-5-yl)-5-fluoro-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 2-(4-methoxyphenyl)ethylamine were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-N2-(4-methoxyphenyloxyethyleneamino)-2,4-pyrimidinediamine. LCMS: ret. time: 18.88 min.; purity: 97.6%; MS (m/e): 371.09 (MH⁺).

7.3.437 N4-(3-Aminophenyl)-N2-[[N3-[N4-(3-aminophenyl)]-5-fluoro-2,4-pyrimidinediamine]aminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950137)

2,4-Dichloro-5-fluoropyrimidine and 3-aminoaniline were reacted to prepare N4-(3-aminophenyl)-N2-[[N3-[N4-(3-aminophenyl)]-5-fluoro-2,4-pyrimidinediamine]aminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 13.10 min.; purity: 96.4%; MS (m/e): 513.01 (MH⁺).

7.3.438 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(hydroxyethylamino)phenyl]-2,4-pyrimidinediamine (R950208)

N2-(3-Aminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and 2-bromoethanol were reacted together to give N4-(3,4-ethylenedioxyphenyl)-N2-[3-(hydroxyethylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 15.44 min.; purity: 98.6%; MS (m/e): 398.05 (MH⁺).

7.3.439 N2-[3-Bis(hydroxyethyl)aminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950209)

N2-(3-Aminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and 2-bromoethanol were reacted together to give N2-[3-bis(hydroxyethyl)aminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 15.64 min.; purity: 97.8%; MS (m/e): 442.06 (MH⁺).

7.3.440 6-Ethoxycarbonyl-N4-(ethoxycarbonylmethyl)-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-5-nitro-2,4-pyrimidinediamine (R925858)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-hydroxyphenyl]-2,4-pyrimidinediamine, N-(2-chloro-6-ethoxycarbonyl-5-nitro-4-pyrimidinyl)glycine ethyl ester and ethyl 4-aminophenoxyacetate were reacted to yield 6-ethoxycarbonyl-N4-(ethoxycarbonylmethyl)-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-5-nitro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 9.00 (bs, 1H), 7.49 (bs, 1H), 7.41 (d, 2H, J= 9.0 Hz), 6.89 (d, 2H, J= 9.0 Hz), 4.62 (s, 2H), 4.46 (q, 2H, J= 7.2 Hz), 4.31-4.19 (m, 6H), 1.40 (t, 3H, J= 7.2 Hz), 1.33-1.25 (m, 6H); LCMS: ret. time: 30.00 min.; purity: 98 %; MS (m/e): 492 (MH⁺).

7.3.441 N4-Benzylloxy-5-ethoxycarbonyl-N2-(3,4-ethylenedioxyphenyl)-2,4-pyrimidinediamine (R925837)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-benzylloxy-2-chloro-5-ethoxycarbonyl-4-pyrimidineamine and 1,4-benzodioxan-6-amine were reacted to yield N4-benzylloxy-5-ethoxycarbonyl-N2-(3,4-ethylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.55 (s, 1H), 7.49-7.44 (m, 3H), 7.39-7.34 (m, 4H), 7.30-7.22 (m, 1H), 6.67 (d, 1H, J= 8.4 Hz), 4.98 (s, 2H), 4.23-4.17 (m, 6H), 1.26 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 26.14 min.; purity: 95%; MS (m/e): 423 (MH⁺).

7.3.442 N4-Benzylloxy-5-ethoxycarbonyl-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925824)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-benzylloxy-2-chloro-5-ethoxycarbonyl-4-pyrimidineamine and 3-hydroxyaniline were reacted to yield N4-benzylloxy-5-ethoxycarbonyl-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 24.28 min.; purity: 88 %; MS (m/e): 381 (MH⁺).

7.3.443 N2,N4-Bis[4-(aminocarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R945025)

A mixture of 4-nitrophenol (7.65 g, 55 mmol), 2-bromoacetamide (6.90 g, 50 mmol) and K₂CO₃ (13.8 g, 0.1 mol) in acetone (50 mL) was stirred at room temperature for 24 h. The reaction mixture was diluted with water, and acetone was removed under reduced pressure. The formed light-yellow precipitate was collected by filtration, washed with water and dried to give 1-aminocarbonylmethyleneoxy-4-nitrobenzene (8.28 g, 84%).

Hydrogenation of 1-aminocarbonylmethyleneoxy-4-nitrobenzene (3 g, 15 mmol) in methanol (50 mL) catalyzed by 10% Pd-C (500 mg) and Na₂SO₄ (500 mg) at 50 psi for 2h gave 4-(aminocarbonylmethyleneoxy)aniline (2.59 g, quant.).

4-(Aminocarbonylmethyleneoxy)aniline (500 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (200 mg, 1.2 mmol) were dissolved in methanol (10 mL) and water (1 mL) and was stirred at 70 °C for 24 h. Then methanol was removed under reduced pressure. The remaining aqueous solution was acidified with 1 N HCl (80 mL). The formed white precipitate was collected by filtration to give N2,N4-bis[4-(aminocarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (370 mg, 72%). ¹H

NMR (acetone- d_6): δ 4.46 (s, 2H), 4.50 (s, 2H), 6.81 (br, NH, 2H), 6.91 (d, J = 9.0 Hz, 2H), 6.98 (d, J = 9.0 Hz, 2H), 7.20 (br, 2H, NH), 7.63 (d, J = 9.3 Hz, 2H), 7.72 (d, J = 8.7 Hz, 2H), 7.93 (d, J =3.6 Hz, 1H), 8.27 (br, 1H, NH), 8.44 (br, 1H, NH); LCMS: ret. time: 13.91 min.; purity: 100%; MS (m/e): 427.02 (MH^+).

5 **7.3.444 N2,N4-Bis[4-(cyanomethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R945032)**

To a solution of N2,N4-bis[4-(aminocarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (200 mg, 0.47 mmol) in THF (10 mL) was added trifluoroacetic anhydride (0.33 mL, 2.35 mmol) and pyridine (0.38 mL, 4.7 mmol) at room temperature and was stirred at room temperature overnight. The mixture was diluted with ethyl acetate (80 mL) and 1 N HCl (80 mL). The organic layer was washed with 1 N HCl (2 x 60 mL), water (2 x 60 mL) and brine (60 mL). The ethyl acetate layer was dried and evaporated. The residue was recrystallized from ethyl acetate and hexanes to give N2,N4-bis[4-(cyanomethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (159 mg, 87%) as a white solid. ^1H NMR (acetone- d_6): δ 5.09 (s, 2H), 5.16 (s, 2H), 7.08 (d, J = 9.3 Hz, 2H), 7.17 (d, J = 9.0 Hz, 2H), 7.63 (d, J = 9.0 Hz, 2H), 7.77 (d, J = 9.3 Hz, 2H), 8.17 (d, J = 4.8 Hz, 1H), 9.55 (br, 1H, NH), 11.00 (br, 1H, NH); LCMS: 21.47 min.; 96.11%; MS (m/e): 391.20 (MH^+).

20 **7.3.445 N2,N4-Bis[4-(1H-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R945033)**

To a solution of N2,N4-bis[4-(cyanomethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (100 mg, 0.26 mmol) in DMF (10 mL) was added NH_4Cl (136 mg, 2.54 mmol), sodium azide (100 mg, 1.54 mmol), and one drop of acetic acid and was stirred at 70 °C overnight. Then it was titrated with ethyl acetate (80 mL) to give precipitation. The precipitate was collected by filtration, washed with 1 N HCl and water to give N2,N4-bis[4-(1H-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (66 mg, 54%) as a white solid. ^1H NMR (CD_3OD): δ 5.31 (s, 2H), 5.34 (s, 2H), 6.93 (d, J = 9.0 Hz, 2H), 7.00 (d, J = 9.3 Hz, 2H), 7.04 (d, J = 9.0 Hz, 2H), 7.57 (d, J = 9.0 Hz, 2H), 7.81 (d, J = 4.2 Hz, 1H); LCMS: 16.54 min.; purity: 88.34%; MS (m/e): 477.02 (MH^+).

7.3.446 N₂,N₄-Bis(4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R945034)

A mixture of 4-Aminobenzoic acid (410 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) in methanol (10 mL) and water (1 mL) was stirred at 100 °C for 24 h to yield N₂,N₄-bis(4-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine after methanol was removal. This residue was redissolved in DMF (10 mL) and to it was added potassium carbonate (1.65 g, 12 mmol) and iodomethane (0.37 mL, 6 mmol), stirred at room temperature overnight, and then diluted with 1 N HCl (80 mL) and ethyl acetate (80 mL). The ethyl acetate layer was washed with 1N HCl (60 mL) and water (60 mL). The organic layer was separated, dried, evaporated and the resulting residue was recrystallized from ethyl acetate/hexanes to give N₂,N₄-bis(4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (150 mg, 63%). ¹H NMR (acetone-*d*₆): δ 3.85 (s, 3H), 3.88 (s, 3H), 7.88-7.97 (m, 4H), 7.98-8.05 (m, 4H), 8.18 (d, J= 3.0 Hz, 1H), 9.00 (br, 1H, NH), 9.04 (br, 1H, NH); LCMS: ret. time: 27.07 min.; purity: 95.54%; MS (m/e): 397.04 (MH⁺).

7.3.447 N₂,N₄-Bis(3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R945035)

In a manner analogous to the preparation of N₂,N₄-bis(4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 3-aminobenzoic acid (410 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N₂,N₄-bis(3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (180 mg, 76%) as a white solid. ¹H NMR (acetone-*d*₆): δ 3.81 (s, 3H), 3.83 (s, 3H), 7.37 (t, J= 8.1 Hz, 1H), 7.47 (t, J= 8.1 Hz, 1H), 7.60 (d, J= 7.8 Hz, 1H), 7.75 (d, J= 7.5 Hz, 1H), 8.02 (d, J= 6.3 Hz, 1H), 8.10 (d, J= 3.6 Hz, 1H), 8.24 (d, J= 8.4 Hz, 1H), 8.36 (d, J= 11.4 Hz, 2H), 8.74 (br, 1H, NH), 8.82 (br, 1H, NH); LCMS: ret. time: 22.77 min.; purity: 91.04%; MS (m/e): 397.00 (MH⁺).

7.3.448 N₂,N₄-Bis(3-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945036)

A solution of N₂,N₄-bis(3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (100 mg, 0.25 mmol) and NaOH (140 mg, 3.5 mmol) in THF:H₂O (5 mL, each) was stirred at room temperature overnight. The reaction mixture was diluted with water (60 mL) and ethyl acetate (60 mL). The aqueous layer was separated, acidified with 1N HCl solution to pH 3. The formed precipitate was collected by filtration and recrystallized from methanol to give N₂,N₄-bis(3-carboxyphenyl)-5-fluoro-2,4-

pyrimidinediamine (54 mg, 58%) as a white solid. ¹H NMR (CD₃OD): δ 7.31 (t, J= 8.1 Hz, 1H), 7.42 (t, J= 8.1 Hz, 1H), 7.61 (dm, J= 7.8 Hz, 1H), 7.76 (dm, J= 8.4 Hz, 1H), 7.89 (dm, J= 7.2 Hz, 1H), 7.98 (d, J= 3.6 Hz, 1H), 8.01 (m, 1H), 8.20 (m, 1H), 8.37 (m, 1H); LCMS: ret. time: 15.77 min.; purity: 98.84%; MS (m/e): 369.03 (MH⁺).

5 **7.3.449 N2,N4-Bis(4-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945037)**

In a manner analogous to the preparation of N2,N4-bis(3-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (100 mg, 0.25 mmol) and NaOH (200 mg, 5 mmol) gave N2,N4-bis(4-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine (55 mg, 59%) as a white solid. ¹H NMR (CD₃OD): δ 7.77 (d, J= 8.7 Hz, 2H), 7.92 (d, J= 8.7 Hz, 2H), 7.94 (d, J= 8.4 Hz, 2H), 8.02 (d, J= 8.7 Hz, 2H), 8.07 (d, J= 3.6 Hz, 1H); LCMS: ret. time: 16.34 min.; purity: 100%; MS (m/e): 368.87 (MH⁺).

15 **7.3.450 N2,N4-Bis(3-isopropylaminocarbonyloxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926412)**

The reaction of 1 equivalent of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine with 3 equivalents of isopropyl isocyanate in the presence of pyridine in CH₂Cl₂ at room temperature for 24 h followed by extractive work up using CH₂Cl₂ gave the desired N2,N4-bis(3-isopropylaminocarbonyloxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃ + CD₃OD): δ 7.82 (d, 1H, J= 3.6 Hz), 7.66 (bd, 1H), 7.48 (bd, 1H), 7.15-7.02 (m, 2H), 6.76-6.76 (m, 2H), 6.56 (bd, 1H, J= 8.1 Hz), 6.45 (dd, 1H, J= 1.8 and 8.4 Hz), 4.70 (m, 2H), 1.05 (d, 12H, J= 6.3 Hz); ¹⁹F NMR (CDCl₃ + CD₃OD): - 47206; LCMS: ret. time: 15.40 min.; purity: 90%.

25 **7.3.451 N2,N4-Bis[4-(ethylaminocarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R945040)**

A mixture of 1,4-diaminobenzene (4 g, 37 mmol), ethyl isocyanate (1 mL, 12.6 mmol) and potassium carbonate (8.72 g, 63 mmol) in THF (20 mL) was stirred at room temperature overnight. The reaction mixture was partitioned in 1N HCl solution (80 mL) and ethyl acetate (80 mL). The aqueous layer was extracted with ethyl acetate (4 x 80 mL). The combined organic layers was dried, evaporated, recrystallized from MeOH/CH₂Cl₂/hexanes to give 4-(ethylaminocarbonylamino)aniline (1.4 g, 62%) as a beige solid.

In a manner analogous to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 4-(ethylaminocarbonylamino)aniline (537 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N2,N4-bis[4-(ethylaminocarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (180 mg, 66%) as a white solid. ¹H NMR (CD₃OD): δ 1.16 (t, J= 7.2 Hz, 6H), 3.24 (q, J= 7.2 Hz, 4H), 7.29 (d, J= 9.0 Hz, 2H), 7.40 (t, J= 9.0 Hz, 4H), 7.55 (d, J= 9.0 Hz, 2H), 7.87 (s, 1H, NH), 7.89 (s, 1H, NH); LCMS: ret. time: 16.93 min.; purity: 93.43%; MS (m/e): 453.03 (MH⁺).

7.3.452 N2,N4-Bis[3-(ethylaminocarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (R945045)

In a manner analogous to the preparation of N2,N4-bis[4-(ethylaminocarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine, the reaction of 1,3-diaminobenzene (2 g, 18.5 mmol), ethyl isocyanate (0.5 mL, 6.3 mmol) and potassium carbonate (4.36 g, 31.5 mmol) gave 3-(ethylaminocarbonylamino)aniline (940 mg, 83%). The reaction of 3-(ethylaminocarbonylamino)aniline (537 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N2,N4-bis[3-(ethylaminocarbonylamino)phenyl]-5-fluoro-2,4-pyrimidinediamine (180 mg, 66%) as a white solid. ¹H NMR (CD₃OD): δ 1.14 (t, J= 6.9 Hz, 3H), 1.15 (t, J= 7.5 Hz, 3H), 3.21 (q, J= 7.2 Hz, 2H), 3.22 (q, J= 7.5 Hz, 2H), 7.06 (ddd, J= 0.9, 2.1, 7.8 Hz, 1H), 7.10-7.28 (m, 5H), 7.53 (t, J= 2.1 Hz, 1H), 7.80 (m, 1H), 7.92 (d, J= 5.7 Hz, 1H); LCMS: ret. time: 17.17 min.; purity: 89.63%; MS (m/e): 453.38 (MH⁺).

7.3.453 N2,N4-Bis(4-hydroxy-3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R945043)

A solution of N2,N4-bis(3-carboxy-4-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine (70 mg, 0.17 mmol) and thionyl chloride (0.04 mL, 0.55 mmol) in MeOH (10 mL) was refluxed overnight. Methanol was removed *in vacuo*. The residue was diluted with EtOAc (60 mL) and sodium hydrogen carbonate solution (60 mL). The EtOAc layer was washed with NaHCO₃ aqueous solution (60 mL) and water (60 mL). The organic layer was dried, evaporated and crystallized from MeOH/Et₂O to give N2,N4-bis(4-hydroxy-3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (58 mg, 77%). ¹H NMR (DMSO-d₆): δ 3.69 (s, 3H), 3.71 (s, 3H), 6.81 (d, J= 9.3 Hz, 1H), 6.92 (d, J= 9.0 Hz, 1H), 7.64 (dd, J= 2.7, 9.0 Hz, 1H), 7.84 (dd, J= 2.1 and 8.4 Hz, 1H), 8.03-8.07 (m, 3H), 9.14 (s, 1H, NH), 9.34 (s, 1H, NH), 10.16 (s, 1H, OH), 10.29 (s, 1H, OH); ¹⁹F NMR (282

MHz, DMSO-d₆): δ - 165.60; LCMS: ret. time: 22.24 min.; purity: 100%; MS (m/e): 428.98 (MH⁺).

7.3.454 N₂,N₄-Bis[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine(R945046)

5-Fluoro-N₂,N₄-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl],[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945047)

N₂,N₄-Bis[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R945048)

Compound N₂,N₄-bis[4-(1H-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (30 mg, 0.063 mmol), iodomethane (0.024 mL, 0.38 mmol) and K₂CO₃ (88 mg, 0.64 mmol) in DMF (5 mL) was stirred at room temperature overnight.

Then it was diluted with ethyl acetate (50 mL) and water (50 mL). The organic layer was washed with water (50 mL) and brine (50 mL). After separation, the ethyl acetate layer was dried, evaporated and purified by flash column chromatography (EtOAc/hexanes = 2/1, 1/1, EtOAc) to give a mixture of following compounds: N₂,N₄-bis[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine **R945046** (6 mg, 19%),

¹H NMR (CDCl₃): δ 4.37 (s, 3H), 4.38 (s, 3H), 5.33 (s, 2H), 5.36 (s, 2H), 6.65 (d, J= 3.0 Hz, 1H), 6.76 (s, 1H), 6.98 (d, J= 9.0 Hz, 2H), 7.04 (d, J= 9.3 Hz, 2H), 7.42 (d, J= 9.0 Hz, 2H), 7.51 (d, J= 9.0 Hz, 2H), 7.90 (br, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 168.52;

LCMS: ret. time: 20.44 min.; purity: 94.92%; MS (m/e): 505.02 (MH⁺); 5-fluoro-N₂,N₄-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl],[4-(2-methyl-1,2,3,4-tetrazol-5-

yl)methyleneoxyphenyl]-2,4-pyrimidinediamine **R945047** (8 mg, 25%), ¹H NMR (CDCl₃):

δ 4.18 (s, 3H), 4.20 (s, 3H), 4.36 (s, 3H), 4.37 (s, 3H), 5.34 (s, 2H), 5.37 (s, 2H), 5.42 (s, 2H), 5.46 (s, 2H), 6.69 (br, 2H), 6.80 (s, 1H), 6.83 (s, 1H), 6.91 (d, J= 9.3 Hz, 2H), 6.98 (d, J= 9.0 Hz, 2H), 6.99 (d, J= 9.3 Hz, 2H), 7.04 (d, J= 9.0 Hz, 2H), 7.41 (d, J= 9.9 Hz, 2H), 7.44 (d, J= 9.3 Hz, 2H), 7.50 (d, J= 9.0 Hz, 2H), 7.54 (d, J= 9.0 Hz, 2H), 7.91 (br, 2H); ¹⁹F

NMR (282 MHz, CDCl₃): δ - 168.39, - 168.16; LCMS: ret. time: 19.42 min.; purity:

91.18%; MS (m/e): 504.99 (MH⁺), and N₂,N₄-bis[4-(1-methyl-1,2,3,4-tetrazol-5-

yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine **R945048** (6 mg, 19%), ¹H NMR

(CD₃OD + CDCl₃): δ 4.20 (s, 3H), 4.22 (s, 3H), 5.50 (s, 2H), 5.55 (s, 2H), 6.95 (d, J= 9.0 Hz, 2H), 7.02 (d, J= 9.3 Hz, 2H), 7.52 (d, J= 9.0 Hz, 2H), 7.66 (d, J= 9.3 Hz, 2H), 7.84 (d, J= 3.6 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD+CDCl₃): δ - 163.12; LCMS: ret. time: 18.32 min.; purity: 83.41%; MS (m/e): 504.99 (MH⁺).

5 **7.3.455 N4-(4-Aminocarbonylmethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945052)**

In like manner to the preparation of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 4-(aminocarbonylmethyleneoxy)aniline (398 mg, 2.4 mmol) and
 10 2,4-dichloro-5-fluoropyrimidine (200 mg, 1.2 mmol) gave N4-(4-aminocarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (270 mg, 76%).
 In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of methyl 4-aminophenoxyacetate (183 mg, 1 mmol) and
 N4-(4-aminocarbonylmethyleneoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine (100 mg,
 15 0.34 mmol) gave N4-(4-aminocarbonylmethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (120 mg, 80%). ¹H NMR (acetone-*d*₆): δ 3.25 (s, 3H), 3.98 (s, 2H), 4.33 (s, 2H), 6.45 (d, J= 8.7 Hz, 2H), 6.49 (d, J= 9.3 Hz, 2H), 6.93 (d, J= 8.7 Hz, 2H), 7.13 (d, J= 9.0 Hz, 2H), 7.71 (d, J= 5.1 Hz, 1H), 9.46 (br, 1H, NH), 9.78 (br, 1H, NH); LCMS: ret. time: 16.65 min.; purity: 100%; MS (m/e):
 20 442.01 (MH⁺).

7.3.456 N4-(4-Cyanomethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945053)

In a manner analogous to the preparation of N2,N4-bis(4-
 25 cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of N4-(4-aminocarbonylmethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (80 mg, 0.18 mmol), trifluoroacetic anhydride (0.13 mL, 0.92 mmol) and pyridine (0.15 mL, 1.84 mmol) gave N4-(4-cyanomethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (52 mg, 68%)
 30 as a white solid. ¹H NMR (DMSO-*d*₆): δ 3.24 (s, 3H), 4.26 (s, 2H), 4.71 (s, 2H), 6.36 (d, J= 9.3 Hz, 2H), 6.59 (d, J= 9.0 Hz, 2H), 7.06 (d, J= 9.0 Hz, 2H), 7.28 (d, J= 9.0 Hz, 2H), 7.58 (d, J= 3.6 Hz, 1H), 8.59 (br, 1H, NH), 8.85 (br, 1H, NH); ¹⁹F NMR (282 MHz,

DMSO-d₆): δ - 166.26; LCMS: ret. time: 21.37 min.; purity: 100%; MS (m/e): 424.01 (MH⁺).

7.3.457 N₂,N₄-Bis[3-hydroxy-4-(methoxycarbonyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (R945056)

5 A solution of 4-amino-2-hydroxybenzoic acid (1 g, 6.5 mmol) in MeOH (15 mL) and concentrated sulfuric acid (1 mL) was refluxed overnight. The reaction mixture was quenched with NaHCO₃ aqueous solution (60 mL) and EtOAc (60 mL). The organic layer was separated, dried, evaporated to give 3-hydroxy-4-methoxycarbonylaniline.

In a manner analogous to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 3-hydroxy-4-methoxycarbonylaniline (500 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N₂,N₄-bis-[3-hydroxy-4-(methoxycarbonyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (105 mg, 41%). ¹H NMR (DMSO-d₆): δ 3.90 (s, 3H), 3.93 (s, 3H), 7.31 (dd, J = 2.4, 9.0 Hz, 1H), 7.56 (dd, J = 2.1, 8.7 Hz, 1H), 7.63 (d, J = 2.1 Hz, 1H), 7.67 (d, J = 2.1 Hz, 1H), 7.67 (d, J = 9.0 Hz, 1H), 7.79 (d, J = 9.0 Hz, 1H), 8.28 (d, J = 3.6 Hz, 1H), 9.72 (s, 1H, NH), 9.82 (s, 1H, NH), 10.77 (s, 1H, OH), 10.80 (s, 1H, OH); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 161.74; LCMS: ret. time: 31.47 min.; purity: 96.03%; MS (m/e): 428.99 (MH⁺).

7.3.458 N₂-(4-Aminocarbonylmethyleneoxyphenyl)-5-fluoro-N₄-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945060)

In a manner analogous to the preparation of N₄-(4-aminocarbonylmethyleneoxyphenyl)-5-fluoro-N₂-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N₄-(4-methoxycarbonylmethyleneoxyphenyl)-4-pyrimidineamine (150 mg, 0.48 mmol) and 4-(aminocarbonylmethyleneoxy)aniline (240 mg, 1.44 mmol) gave N₂-(4-aminocarbonylmethyleneoxyphenyl)-5-fluoro-N₄-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (145 mg, 68%). ¹H NMR (DMSO-d₆): δ 3.70 (s, 3H), 4.40 (s, 2H), 4.81 (s, 2H), 6.91 (d, J = 8.4 Hz, 2H), 6.93 (d, J = 8.4 Hz, 2H), 7.36 (d, J = 8.4 Hz, 2H), 7.54 (d, J = 8.7 Hz, 2H), 8.21 (d, J = 4.8 Hz, 1H), 10.13 (br, 1H, NH), 10.39 (br, 1H, NH); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 162.26; LCMS: ret. time: 15.37 min.; purity: 78.49%; MS (m/e): 442.07 (MH⁺).

7.3.459 N2,N4-Bis(3-hydroxy-4-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945061)

In a manner analogous to the preparation of N2,N4-bis(3-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of N2,N4-bis[3-hydroxy-4-(methoxycarbonyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (70 mg, 0.16 mmol) and NaOH (100 mg, 2.5 mmol) gave N2,N4-bis(3-hydroxy-4-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine (50 mg, 77%) as a white solid. ¹H NMR (DMSO-d₆): δ 7.21 (dd, J= 1.5 and 8.7 Hz, 1H), 7.46-7.52 (m, 3H), 7.63 (d, J= 8.7 Hz, 1H), 7.72 (d, J= 8.7 Hz, 1H), 8.28 (d, J= 3.3 Hz, 1H), 9.71 (s, 1H, NH), 9.79 (s, 1H, NH), 11.34 (br, 2H); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 161.10; LCMS: ret. time: 20.76 min.; purity: 84.65%; MS (m/e): 400.95 (MH⁺).

7.3.460 N2-(4-Cyanomethyleneoxyphenyl)-5-fluoro-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945062)

In a manner analogous to the preparation of N2,N4-bis(4-cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N2-(4-aminocarbonylmethyleneoxyphenyl)-5-fluoro-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (100 mg, 0.23 mmol), trifluoroacetic anhydride (0.16 mL, 1.13 mmol) and pyridine (0.18 mL, 2.21 mmol) gave N2-(4-cyanomethyleneoxyphenyl)-5-fluoro-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (66 mg, 69%) as a white solid. ¹H NMR (acetone-d₆): δ 3.75 (s, 3H), 4.67 (s, 2H), 4.89 (s, 2H), 6.88 (d, J= 9.0 Hz, 2H), 6.90 (d, J= 9.3 Hz, 2H), 7.48 (d, J= 9.0 Hz, 2H), 7.54 (d, J= 9.0 Hz, 2H), 7.84 (d, J= 4.2 Hz, 1H), 9.17 (br, 1H, NH), 10.59 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-d₆): δ - 164.65; LCMS: ret. time: 20.69 min.; purity: 94.35%; MS (m/e): 424.02 (MH⁺).

7.3.461 N2,N4-Bis(3-methoxy-4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R945065)

In a manner analogous to the preparation of N2,N4-bis[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine, 2-methoxy-4-nitrobenzoic acid (1 g, 5 mmol), potassium carbonate (1.4 g, 10 mmol) and iodomethane (0.47 mL, 7.5 mmol) gave methyl 2-methoxy-4-nitrobenzoate (820 mg, 77%) as a white solid.

The hydrogenation of methyl 2-methoxy-4-nitrobenzoate (700 mg, 3.3 mmol) in methanol (10 mL) catalyzed by 5% Pd-C (100 mg) and Na₂SO₄ (100 mg) at 50 psi for 1h gave methyl 4-amino-2-methoxybenzoate (600 mg, quant.) as a white solid.

In a manner analogous to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, methyl 4-amino-2-methoxybenzoate (542 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N₂,N₄-bis(3-methoxy-4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (180 mg, 66%) as a white solid. ¹H NMR (acetone-*d*₆): δ 3.76 (s, 3H), 3.77 (s, 3H), 3.81 (s, 6H), 7.36 (dd, J= 1.8, 8.7Hz, 1H), 7.57 (s, 1H), 7.58 (dd, J= 2.1 and 7.2 Hz, 1H), 7.69 (d, J= 2.1 Hz, 1H), 7.73 (d, J= 8.4 Hz, 1H), 7.75 (d, J= 9.0 Hz, 1H), 8.17 (d, J= 3.3 Hz, 1H), 8.89 (s, 2H, NH); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ - 165.18; LCMS: ret. time: 23.17 min.; purity: 100%; MS (m/e): 456.96 (MH⁺).

7.3.462 N₂,N₄-Bis(4-methoxy-3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R945066)

In a manner analogous to the preparation of N₂,N₄-bis(3-methoxy-4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine, 2-hydroxy-5-nitrobenzoic acid (1 g, 5.5 mmol), potassium carbonate (3 g, 22 mmol) and iodomethane (1 mL, 16 mmol) gave methyl 2-hydroxy-5-nitrobenzoate (880 mg, 77%).

The hydrogenation of methyl 2-hydroxy-5-nitrobenzoate (700 mg, 3.3 mmol) using 10% Pd-C (100 mg) and Na₂SO₄ (100 mg) in MeOH at 50 psi gave methyl 5-amino-2-methoxybenzoate (600 mg).

In a manner analogous to the preparation of N₂,N₄-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine, methyl 5-amino-2-methoxybenzoate (542 mg, 3 mmol) and 2,4-dichloro-5-fluoropyrimidine (100 mg, 0.6 mmol) gave N₂,N₄-bis(4-methoxy-3-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (170 mg, 62%) as a pink solid. ¹H NMR (acetone-*d*₆): δ 3.76 (s, 3H), 3.77 (s, 3H), 3.88 (s, 3H), 3.93 (s, 3H), 7.08 (dd, J= 0.8, 9.0 Hz, 1H), 7.18 (d, J= 9.0 Hz, 1H), 7.66 (dd, J= 3.0 and 8.7 Hz, 1H), 7.78 (dd, J= 1.5 and 3.0 Hz, 1H), 7.86 (dt, J= 2.7 and 9.0 Hz, 1H), 7.98 (t, J= 2.7 Hz, 1H), 8.32 (d, J= 5.1 Hz, 1H); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ -163.88; LCMS: ret. time: 19.07 min.; purity: 98.17%; MS (m/e): 456.94 (MH⁺).

7.3.463 N2,N4-Bis(3-carboxy-4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945067)

In a manner analogous to the preparation of N2,N4-bis(3-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis[4-methoxy-3-(methoxycarbonyl)phenyl]-5-fluoro-2,4-pyrimidinediamine (80 mg, 0.18 mmol) and NaOH (200 mg, 5 mmol) gave N2,N4-bis(3-carboxy-4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (80 mg). ¹H NMR (DMSO-d₆): δ 3.75 (s, 3H), 3.80 (s, 3H), 6.94 (d, J = 9.6 Hz, 1H), 7.05 (d, J = 9.3 Hz, 1H), 7.78-7.80 (m, 3H), 7.94 (dd, J = 9.3 Hz, 1H), 8.04 (d, J = 3.6 Hz, 1H), 9.10 (s, 1H, NH), 9.30 (s, 1H, NH); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 165.56; LCMS: ret. time: 14.65 min.; purity: 100%; MS (m/e): 428.83 (MH⁺).

7.3.464 N2,N4-Bis(4-carboxy-3-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945068)

In a manner analogous to the preparation of N2,N4-bis(3-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N2,N4-bis(3-methoxy-4-methoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (30 mg, 0.06 mmol) and NaOH (200 mg, 5 mmol) gave N2,N4-bis(4-carboxy-3-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (18 mg, 64%) as a white solid. ¹H NMR (DMSO-d₆): δ 3.66 (s, 3H), 3.73 (s, 3H), 7.37 (d, J = 8.4 Hz, 1H), 7.47 (s, 1H), 7.49 (s, 1H), 7.61-7.71 (m, 3H), 8.25 (d, J = 3.6 Hz, 1H), 9.65 (s, 1H, NH), 9.70 (s, 1H, NH); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 162.11; LCMS: ret. time: 17.25 min.; purity: 100%; MS (m/e): 429.04 (MH⁺).

7.3.465 N2-(4-Cyanomethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945070)

In a manner analogous to the preparation of N2,N4-bis(4-cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N2-(4-(aminocarbonylmethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (60 mg, 0.16 mmol), trifluoroacetic anhydride (0.11 mL, 0.8 mmol) and pyridine (0.13 mL, 1.6 mmol) gave N2-(4-cyanomethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (30 mg, 53%). ¹H NMR (acetone-d₆): δ 5.04 (s, 2H), 6.60 (ddd, J = 0.9, 2.4 and 8.1 Hz, 1H), 7.02 (d, J = 9.3 Hz, 2H), 7.15 (t, J = 8.1 Hz, 1H), 7.31 (ddd, J = 1.2, 2.1 and 8.1 Hz, 1H), 7.38 (t, J = 2.1 Hz, 1H), 7.78 (d, J = 9.3 Hz, 2H), 7.98 (d, J = 3.6 Hz, 1H), 8.34 (s, 1H, NH), 8.42 (s, 1H, NH); ¹⁹F NMR (282 MHz, acetone-d₆): δ - 168.06; LCMS: ret. time: 18.17 min.; purity: 97.47%; MS (m/e): 352.05 (MH⁺).

7.3.466 N4-(4-Cyanomethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945172)

In a manner analogous to the preparation of N2,N4-bis(4-cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N4-(4-aminocarbonylmethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, trifluoroacetic anhydride and pyridine in THF gave N4-(4-cyanomethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 4.27 (m, 4H), 4.82 (s, 2H), 6.70 (dd, J= 2.4 and 8.4 Hz, 1H), 6.75 (d, J= 2.4 Hz, 1H), 6.86 (d, J= 8.4 Hz, 1H), 7.02 (d, J= 9.0 Hz, 2H), 7.32 (d, J= 9.0 Hz, 2H), 8.64 (d, J= 1.8 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 135.58; LCMS: ret. time: 19.92 min.; purity: 98.02%; MS (m/e): 393.98 (MH⁺).

7.3.467 N2,N4-Bis[4-[2-methoxyimino(amino)ethyleneoxy]phenyl]-5-fluoro-2,4-pyrimidinediamine (R945096)

N2,N4-Bis(4-cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (50 mg, 0.13 mmol), methoxyamine HCl salt (54 mg, 0.65 mmol) and sodium bicarbonate (54 mg, 0.65 mmol) were dissolved in methanol (5 mL). The reaction solution was stirred at 70 °C for 7 days. Then methanol was removed under reduced pressure. The residue was partitioned in EtOAc (60 ml) and water (60 mL). The ethyl acetate layer was washed with water (2 x 60 mL), dried, evaporated and purified by flash column chromatography (EtOAc/hexanes; 1:1; EtOAc) to give N2,N4-bis[4-[2-methoxyimino(amino)ethyleneoxy]phenyl]-5-fluoro-2,4-pyrimidinediamine (30 mg, 48%). ¹H NMR (acetone-*d*₆): δ 3.70 (s, 3H), 3.71 (s, 3H), 4.44 (s, 2H), 4.49 (s, 2H), 5.43 (br, 2H), 5.47 (br, 2H), 6.93 (d, J= 9.0 Hz, 2H), 7.00 (d, J= 9.0 Hz, 2H), 7.62 (d, J= 9.0 Hz, 2H), 7.71 (d, J= 9.0 Hz, 2H), 7.93 (d, J= 3.6 Hz, 1H), 8.26 (br, 1H, NH), 8.40 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ - 169.08; LCMS: ret. time: 14.41 min.; purity: 100%; MS (m/e): 484.97 (MH⁺).

7.3.468 N2-(4-Carboxymethyleneoxyphenyl)-N4-(4-cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R945097)

In a manner analogous to the preparation of N2,N4-bis(3-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine, N4-(4-cyanomethyleneoxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (10 mg, 0.024 mmol) and

LiOH (2 mg, 0.048 mmol) gave N2-(4-carboxymethyleneoxyphenyl)-N4-(4-cyanomethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (5 mg, 52%) as a white solid. ¹H NMR (CD₃OD): δ 4.60 (s, 2H), 4.99 (s, 2H), 6.88 (d, J= 9.0 Hz, 2H), 7.02 (d, J= 9.0 Hz, 2H), 7.39 (d, J= 8.7 Hz, 2H), 7.65 (d, J= 9.0 Hz, 2H), 7.84 (d, J= 3.9 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 168.81; LCMS: ret. time: 17.95 min.; purity: 86.04%; MS (m/e): 409.99 (MH⁺).

7.3.469 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945127)

A mixture of 3-nitrophenol (4 g, 29 mmol), bromoacetonitrile (2.5 mL, 36 mmol) and K₂CO₃ (8 g, 58 mmol) in acetone (20 mL) was stirred at room temperature overnight. The reaction mixture was diluted with water (80 mL) and acetone was removed under reduced pressure. The light-yellow precipitate was collected by filtration, washed with water and dried to give 1-cyanomethyleneoxy-3-nitrobenzene.

1-Cyanomethyleneoxy-3-nitrobenzene (2 g, 11 mmol) was dissolved in methanol (20 mL) and to the solution was added hydroxyamine HCl salt (1 g, 14 mmol) and triethylamine (3 mL, 22 mmol). The reaction mixture was refluxed for 2 h and the solvent was removed under reduced pressure. The residue was redissolved in THF (30 mL). To the solution was added acetyl chloride (4 mL, 56 mmol) and pyridine (9 mL, 0.11 mol). The reaction mixture was stirred at room temperature overnight, then added THF (10 mL), water (10 mL) and NaOH (3 g, 75 mmol). The reaction solution was refluxed overnight, diluted with water (80 mL). The aqueous solution was extracted with EtOAc (3 x 60 mL). After separation, the combined EtOAc layers was dried, evaporated to give 1-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxy-3-nitrobenzene.

1-(5-Methyl-1,2,4-oxadiazol-3-yl)methyleneoxy-3-nitrobenzene was dissolved in THF (10 mL) and water (10 mL) and to it were added sodium bisulfite (1 g, 5.7 mmol) and sodium bicarbonate (1 g, 12 mmol). The resulting mixture was stirred at room temperature for 30 min, then diluted with EtOAc (80 mL) and water (80 mL). The aqueous solution was extracted with EtOAc (80 mL). The organic layers were combined, dried, evaporated to give 3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyaniline (500 mg, 22% in four steps).

The reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (40 mg, 0.17 mmol) and 3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyaniline (102 mg, 0.50 mmol) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-

yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (35mg, 51%). ¹H NMR (CDCl₃): δ 2.61 (s, 3H), 5.09 (s, 2H), 6.58-6.62 (m, 2H), 6.76 (dt, J= 1.2, 8.1 Hz, 1H), 6.84 (dt, J= 1.2 and 7.8 Hz, 1H), 6.92 (d, J= 3.0 Hz, 1H), 7.139 (t, J= 8.1 Hz, 1H), 7.145 (t, J= 8.1 Hz, 1H), 7.25 (m, 1H), 7.54 (dt, J= 2.1, 8.7 Hz, 2H), 7.88 (d, J= 3.3 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): -166.52; LCMS: ret. time: 19.33 min.; purity: 84.80%; MS (m/e): 409.35 (MH⁺).

7.3.470 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[3-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945130)

1-Methoxycarbonylmethyleneoxy-3-nitrobenzene (2 g, 9.5 mmol) was dissolved in THF (10 mL) and water (10 mL). To the solution was added NaOH (1 g, 25 mmol). The reaction mixture was stirred at room temperature overnight. The solution was diluted with water (60 mL) and EtOAc (60 mL). After extraction, the aqueous layer was separated, acidified with 1N HCl to pH 3. The formed white precipitate was collected by filtration, washed with water, dried to give 1-carboxymethyleneoxy-3-nitrobenzene.

Acetonitrile (2.25 mL, 43 mmol) was dissolved in methanol (10 mL) and to the solution was added hydroxyamine HCl salt (2 g, 29 mmol) and triethylamine (8 mL, 57 mmol). The reaction mixture was refluxed for 2 days and the solvent was removed under reduced pressure to give acetamide oxime as white solid.

Acetamide oxime (0.75 g, 10 mmol), 1-carboxymethyleneoxy-3-nitrobenzene (1 g, 5 mmol), EDC HCl (1.45 g, 7.5 mmol) and diisopropylethylamine (2.65 mL, 15 mmol) were dissolved in THF (15 mL) and refluxed for 4h. The reaction mixture was diluted with EtOAc (60 mL) and water (60 mL). The EtOAc layer was washed with sodium bicarbonate aqueous solution (2 x 60 mL), 1N HCl (2 x 60 mL) and water (60 mL). After separation, the EtOAc layer was dried, evaporated to give 1-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxy-3-nitrobenzene.

Sodium bisulfite (1.5 g, 8.6 mmol), sodium bicarbonate (1.5 g, 18 mmol) and 1-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxy-3-nitrobenzene (1 g, 4 mmol) were dissolved in THF (15 mL) and water (15 mL). It was stirred at room temperature for 20 min, diluted with EtOAc (60 mL) and water (60 mL). The aqueous solution was extracted with EtOAc (2 x 60 mL). The organic layers were combined, dried, evaporated to give 3-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyaniline.

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of 3-(3-methyl-1,2,4-oxadiazol-

5-yl)methyleneoxyaniline (369 mg, 1.8 mmol) and 2,4-dichloro-5-fluoropyrimidine (150 mg, 0.9 mmol) gave 2-chloro-5-fluoro-N4-[3-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine. The reaction of 2-chloro-5-fluoro-N4-[3-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine (20 mg, 0.06 mmol) and 3-hydroxyaniline (20 mg, 0.18 mmol) gave 5-fluoro-N2-(3-hydroxyphenyl)-N4-[3-(3-methyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (10 mg, 42%). ¹H NMR (CDCl₃): δ 2.42 (s, 3H), 5.28 (s, 2H), 6.49 (ddd, J= 0.9, 2.7 and 8.4 Hz, 1H), 6.73 (ddd, J= 0.9, 2.7 and 8.4 Hz, 1H), 6.81-6.84 (m, 2H), 6.88 (ddd, J= 0.6, 2.1 and 8.1 Hz, 1H), 7.13 (t, J= 8.1 Hz, 1H), 7.26 (t, J= 8.1 Hz, 1H), 7.40 (br, 1H), 7.49 (t, J= 2.1 Hz, 1H), 7.94-7.97 (m, 2H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 167.11; LCMS: ret. time: 18.80 min.; purity: 92.01%; MS (m/e): 409.01 (MH⁺).

7.3.471 5-Fluoro-N4-(2-methoxycarbonylbenzofuran-5-yl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945131)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, the reaction of N4-(2-carboxybenzofuran-5-yl)-2-chloro-5-fluoro-4-pyrimidineamine (50 mg, 0.16 mmol) and 3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyaniline (100 mg, 0.49 mmol) gave N4-(2-carboxybenzofuran-5-yl)-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine.

In a manner analogous to the preparation of N2,N4-bis[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine, the reaction of N4-(2-carboxybenzofuran-5-yl)-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, potassium carbonate (100 mg, 0.7 mmol) and iodomethane (0.03 mL, 0.5 mmol) gave 5-fluoro-N4-(2-methoxycarbonylbenzofuran-5-yl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (40 mg, 50%). ¹H NMR (acetone-*d*₆): δ 2.63 (s, 3H), 3.94 (s, 3H), 5.04 (s, 2H), 6.65 (ddd, J= 0.9, 2.4 and 7.8 Hz, 1H), 7.16 (t, J= 7.8 Hz, 1H), 7.24 (ddd, J= 1.2, 1.8 and 8.1 Hz, 1H), 7.58 (d, J= 1.2 Hz, 1H), 7.64 (d, J= 9.3 Hz, 1H), 7.67 (t, J= 2.1 Hz, 1H), 7.88 (dd, J= 2.1 and 9.0 Hz, 1H), 8.04 (d, J= 3.6 Hz, 1H), 8.26 (d, J= 1.8 Hz, 1H), 8.47 (br, 1H, NH), 8.71 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ - 167.73; LCMS: ret. time: 22.55 min.; purity: 85.43%; MS (m/e): 490.97 (MH⁺).

7.3.472 N4-(2-Carboxybenzofuran-5-yl)-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945134)

In a manner analogous to the preparation of N2,N4-bis(3-carboxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(2-methoxycarbonylbenzofuran-5-yl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (20 mg, 0.04 mmol) and NaOH (10 mg, 0.25 mmol) gave N4-(2-carboxybenzofuran-5-yl)-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (acetone-*d*₆): δ 2.63 (s, 3H), 5.04 (s, 2H), 6.64 (d, J= 8.1 Hz, 1H), 7.17 (t, J= 8.1 Hz, 1H), 7.26 (d, J= 7.8 Hz, 1H), 7.56 (s, 1H), 7.62 (d, J= 9.3 Hz, 1H), 7.67 (t, 1H), 7.86 (dd, J= 1.8 and 9.0 Hz, 1H), 8.04 (d, J= 3.3 Hz, 1H), 8.26 (d, 1H), 8.48 (br, 1H, NH), 8.71 (br, 1H, NH); LCMS: ret. time: 18.00 min.; purity: 75.13%; MS (m/e): 476.70 (MH⁺).

7.3.473 N4-(2-Aminocarbonylbenzofuran-5-yl)-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945135)

A mixture of 5-fluoro-N4-(2-methoxycarbonylbenzofuran-5-yl)-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (20 mg, 0.04 mmol) and concentrated NH₄OH (5 mL) in methanol (5 mL) was stirred at room temperature overnight. The solvent was evaporated to give N4-[2-(aminocarbonyl)benzofuran-5-yl]-5-fluoro-N2-[3-(5-methyl-1,2,4-oxadiazol-3-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (acetone-*d*₆): δ 2.61 (s, 3H), 5.04 (s, 2H), 6.64 (ddd, J= 0.9, 2.4 and 8.1 Hz, 1H), 7.16 (t, J= 8.1 Hz, 1H), 7.27 (ddd, J= 0.9, 1.8 and 8.4 Hz, 1H), 7.44 (d, J= 0.6 Hz, 1H), 7.55 (dd, J= 0.6 and 8.1 Hz, 1H), 7.64 (t, J= 2.4 Hz, 1H), 7.79 (dd, J= 2.4 and 9.0 Hz, 1H), 8.03 (d, J= 3.6 Hz, 1H), 8.24 (d, J= 2.4 Hz, 1H), 8.48 (br, 1H, NH), 8.68 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ - 167.80; LCMS: ret. time: 17.43 min.; purity: 100%; MS (m/e): 475.62 (MH⁺).

7.3.474 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(2-methoxyimino(amino)ethyleneoxy)phenyl]-2,4-pyrimidinediamine (R945167)

In a manner analogous to the preparation of N2,N4-bis[4-(2-methoxyimino(amino)ethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine, the reaction of N2-(4-cyanomethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (50 mg, 0.14 mmol), methoxyamine HCl salt (0.71 mmol) and triethylamine (0.2 mL, 1.4

mmol) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(2-methoxyimino(amino)ethyleneoxy)phenyl]-2,4-pyrimidinediamine (40 mg, 70%). ¹H NMR (CDCl₃): δ 3.82 (s, 3H), 4.50 (s, 2H), 4.87 (br, 2H, NH), 6.60 (ddd, J= 0.9, 2.4 and 8.1 Hz, 1H), 6.79-6.84 (m, 2H), 6.86 (d, J= 8.7 Hz, 2H), 7.00 (s, 1H), 7.14 (t, J= 8.1 Hz, 1H), 7.34 (d, J= 9.0 Hz, 2H), 7.47 (t, J= 2.1 Hz, 1H), 7.87 (d, J= 3.3 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 167.67; LCMS: ret. time: 13.69 min.; purity: 92.51%; MS (m/e): 399.01 (MH⁺).

7.3.475 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[4-methoxyimino(amino)ethyleneoxyphenyl]-2,4-pyrimidinediamine (R945175)

In a manner analogous to the preparation of N2,N4-bis[4-(2-methoxyimino(amino)ethyleneoxyphenyl)]-5-fluoro-2,4-pyrimidinediamine, N4-(4-cyanomethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, methoxyamine hydrochloride salt and triethylamine gave N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-methoxyimino(amino)ethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (acetone-d₆): δ 3.70 (s, 3H), 4.21-4.28 (m, 4H), 4.48 (s, 2H), 5.46 (br, 2H), 6.71 (d, J= 8.7 Hz, 1H), 6.99 (d, J= 9.0 Hz, 2H), 7.06 (dd, J= 2.4 and 8.7 Hz, 1H), 7.42 (d, J= 2.4 Hz, 1H), 7.72 (d, J= 9.3 Hz, 2H), 7.93 (d, J= 3.3 Hz, 1H), 8.22 (br, 1H, NH), 8.40 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-d₆): δ - 169.05; LCMS: ret. time: 16.49 min.; purity: 96.47%; MS (m/e): 440.96 (MH⁺).

7.3.476 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926495)

A mixture of N2-(3-ethoxy/or methoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (19.8g, 45 mmol), methylamine hydrochloride (30.63g, 450 mmol) and diisopropylethylamine (78.07 mL, 450 mmol) in MeOH (450 mL) was stirred in a pressure bottle at 100 °C for 8h (followed by TLC). The reaction was cooled to room temperature, diluted with H₂O (6 lit), the solid obtained was filtered, washed with H₂O and dried to obtain 18 g of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxyphenyl)]-2,4-pyrimidinediamine.

Alternatively, the reaction of equimolar amount of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-aminopyridine with 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110 °C for 24h and or in EtOH using microwave at 175 °C for 10-20

min followed by aqueous work up gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.90 (s, 1H), 7.89 (bs, 1H), 7.38 (d, 1H, J= 2.4 Hz), 7.28 (d, 1H, J= 2.4 Hz), 7.17-7.09 (m, 2H), 6.79 (d, 1H, J= 9 Hz), 6.57 (m, 1H), 4.38 (s, 2H), 4.24 (s, 4H), 2.81 (s, 3H); LCMS: ret. time: 18.20 min.; purity: 98%; MS (m/e): 426 (MH⁺).

7.3.477 N4-(1,4-Benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R921219)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.8 (d, 1H), 7.4 (m, 1H), 7.05 (m, 2H), 7.0 (s, 1H), 6.8 (dd, 1H), 6.66 (d, 1H), 6.56 (dd, 1H), 4.35 (s, 2H), 4.18 (m, 2H), 3.25 (m, 2H), 2.8 (s, 3H); LCMS: ret time: 18.0 min. purity: 97 %; MS (m/e): 425 (MH⁺).

7.3.478 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-(N-2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R909239)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 2-hydroxyethylamine were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-(N-2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (D₂O): δ 8.02 (d, 1H, J= 4 Hz), 7.40 (m, 2H), 7.28 (m, 1H), 7.05 (m, 5H), 4.83 (s, 2H), 4.5 (m, 2H), 4.23 (m, 2H), 4.03 (m, 2H), 3.87 (m, 2H); LCMS: ret. time: 17.17 min.; purity: 94%; MS (m/e): 456 (MH⁺).

7.3.479 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R909240)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-

ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-(N-methylamino)carbonylmethyleneoxyaniline were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (D₂O): δ 8.02 (d, 1H, J= 4 Hz), 7.40 (m, 2H), 7.28 (m, 1H), 7.05 (m, 5H), 4.83 (s, 2H), 4.5 (m, 2H), 4.23 (m, 2H), 3.87 (s, 3H); LCMS: ret. time: 18.43 min.; purity: 94%; MS (m/e): 426 (MH⁺).

7.3.480 N4-(1,4-Benzoxazin-6-yl)-5-fluoro-N2-[3-(N-2-hydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R909251)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-(3-ethoxyocarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine and 2-hydroxypropylamine were reacted to yield N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-2-hydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.02 (d, 1H, J= 4 Hz), 7.25 (m, 2H), 7.04 (m, 1H), 6.82 (m, 2H), 6.58 (m, 1H), 6.45 (m, 1H), 4.36 (s, 2H), 4.02 (m, 2H), 3.75 (m, 1H), 3.31 (m, 2H), 3.00 (m, 2H), 1.00 (m, 3H); LCMS: ret. time: 17.33 min.; purity: 97 %; MS(m/e): 469 (MH⁺).

7.3.481 N4-(1,4-Benzoxazin-6-yl)-5-fluoro-N2-[3-(N-3-hydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R909252)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-[3-ethoxyocarbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine and 3-hydroxypropylamine were reacted to yield N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-3-hydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.02 (d, 1H, J= 4 Hz), 7.39 (m, 2H), 7.04 (m, 1H), 6.87 (m, 2H), 6.55 (m, 1H), 6.41 (m, 1H), 4.29 (s, 2H), 4.02 (m, 2H), 3.35 (m, 2H), 3.31 (m, 2H), 3.09 (m, 2H), 1.50 (m, 3H); LCMS: ret. time: 17.11 min.; purity: 94 %; MS (m/e): 469 (MH⁺).

7.3.482 N4-(1,4-Benzoxazin-6-yl)-N2-[3-(N-isopropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R909254)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-

6-yl)]-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-pyrimidinediamine and isopropylamine were reacted to yield N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-isopropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 7.25 (d, 1H, J= 4 Hz), 7.14 (m, 1H), 7.02 (m, 1H), 6.85 (m, 3H), 6.63 (m, 1H), 4.39 (s, 2H), 4.12 (m, 2H), 4.05 (m, 1H), 3.38 (m, 2H), 1.20 (m, 6H); LCMS: ret. time: 20.83 min.; purity: 96 %; MS (m/e): 453 (MH⁺).

7.3.483 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[2-(N-pyrrolidino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926703)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine and pyrrolidine were reacted to yield 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[2-(N-pyrrolidino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.83 (s, 1H), 7.79 (d, 1H, J= 5.4 Hz), 7.42 (bs, 1H), 7.39 (d, 2H, J= 8.7 Hz), 7.28-7.24 (m, 2H), 6.81 (d, 2H, J= 8.7 Hz), 4.52 (2q, 1H, J= 6.0 Hz), 3.92 (t, 2H, J= 6.9 Hz), 3.67 (t, 2H, J= 6.9 Hz), 2.05-1.90 (m, 4H), 1.32 (d, 6H, J= 6.6 Hz); ¹⁹F NMR (CDCl₃): - 24000; LCMS: ret. time: 23.49 min.; purity: 97 %; MS (m/e): 476 (MH⁺).

7.3.484 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926708)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.10 (bs, 1H), 9.88 (bs, 1H), 8.15 (t, 1H, J= 4.5 Hz), 8.05 (bs, 1H), 7.40 (d, 2H, J= 8.7 Hz), 7.23 (d, 1H, J= 2.1 Hz), 7.11 (dd, 1H, J= 2.4 and 8.7 Hz), 6.89 (d, 2H, J= 8.7 Hz), 6.81 (d, 1H, J= 8.7 Hz), 4.42 (s, 2H), 4.23 (s, 4H), 2.64 (d, 3H, J= 4.5 Hz); LCMS: ret. time: 17.60 min.; purity: 96 %; MS (m/e): 426 (MH⁺).

7.3.485 N4-(4-*tert*-Butylphenyl)-5-fluoro-N2-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926494)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine with methylamine hydrochloride gave N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.04 (d, 1H, J = 2.4 Hz), 7.88 (d, 1H, 4.2 Hz), 7.58-7.30 (m, 7H), 2.94 (s, 3H), 1.33 (s, 9H); LCMS: ret. time: 22.86 min.; purity: 94%; MS (m/e): 434 (MH⁺).

7.3.486 N4-(4-*tert*-Butylphenyl)-5-fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926712)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-[4-(*tert*-butylphenyl)-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.92 (d, 1H, J = 5.4 Hz), 7.53 (d, 2H, J = 8.7 Hz), 7.40 (d, 2H, J = 8.7 Hz), 7.34 (d, 2H, J = 8.7 Hz), 7.03 (d, 2H, J = 8.7 Hz), 4.52 (s, 2H), 2.82 (s, 3H), 1.35 (s, 9H); ¹⁹F NMR (CD₃OD): - 46174; LCMS: ret. time: 23.34 min.; purity: 94%; MS (m/e): 424 (MH⁺).

7.3.487 N4-(3-*tert*-Butylphenyl)-5-fluoro-N2-[3-(N-2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine R940295

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3-*tert*-butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 2-hydroxyethylamine were reacted to give N4-(3-*tert*-butylphenyl)-5-fluoro-N2-[3-(N-2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 21.34 min.; purity: 97 %; MS (m/e): 453 (M⁺); 454 (MH⁺); ¹H NMR (CDCl₃): δ 10.34 (1H, s), 7.76 (1H, m), 7.52 (1H, m), 7.4-7.1 (5H, m), 6.98 (1H, m), 6.7 (1H, m), 4.36 (2H, s), 3.77 (2H, t, J 5 Hz), 3.51 (2H, m), 1.27 (9H, s).

7.3.488 N2,N4-Bis[4-(N-pyrrolidino)carbonylmethyleneoxyphenyl]-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926562)

In like manner of the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of
5 N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine with pyrrolidine gave N2,N4-bis[4-(N-pyrrolidino)carbonylmethyleneoxyphenyl]-5-ethoxycarbonyl-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.17 (s, 1H), 8.73 (bs, 1H), 7.50 (bd, 2H, J= 9.0 Hz), 7.43 (d, 2H, J= 2.4 and 6.9 Hz), 6.91 (m, 4H), 4.64 (s, 2H), 4.62 (s, 2H), 4.34 (q, 2H, J= 7.2 Hz), 3.53 (m,
10 8H), 1.95 (m, 4H), 1.86 (m, 4H), 1.38 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 22.54 min.; purity: 100%; MS (m/e): 590 (MH⁺).

7.3.489 N2,N4-Bis(4-N-pyrrolidinocarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926563)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of
15 N2,N4-bis(4-methoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with pyrrolidine gave N2,N4-bis[4-(N-pyrrolidino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.90 (s, 1H), 7.50 (bd, 2H, J= 7.8 Hz), 7.41 (bd, 2H, J= 7.2 Hz), 6.93 (m, 4H), 6.73 (s, 1H), 6.64 (s, 1H), 4.65 (s, 1H), 4.65 (s, 1H), 3.54 (m,
20 8H), 1.96 (m, 4H), 1.87 (m, 4H).

7.3.490 N4-(3-*tert*-Butylphenyl)-N2-[3-(N-1,3-dihydroxypropyl-2-amino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R940296)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3-*tert*-
25 butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 2-amino-1,3-propanediol were reacted to give N4-(3-*tert*-butylphenyl)-N2-[3-(1,3-dihydroxypropyl-2-amino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 20.26 min.; purity: 97.67 %; MS (m/e): 484 (M⁺); 485 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.75 (1H, s), 9.57 (1H, s), 8.25 (1H, m), 7.92 (1H, m), 7.62 (2H, m), 7.37 (3H, m), 7.23 (1H, m), 6.66 (1H, m), 4.46 (2H, s), 3.87 (1H, m), 3.55 (4H, m), 1.36 (9H, s).

7.3.491 N2-[3-(N-2,3-Dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine R940290

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-isopropylphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 3-amino-1,2-propanediol were reacted to give N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 20.04 min.; purity: 98 %; MS (m/e): 470 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.54 (1H, s), 9.41 (1H, s), 8.22 (1H, m), 7.95 (1H, m), 7.85 (1H, d, J= 10 Hz), 7.58 (1H, s), 7.43-7.32 (3H, m), 7.25 (1H, t, J= 7.75 Hz), 7.06 (1H, d, J= 7.75 Hz), 6.64 (1H, d, J= 10 Hz), 4.47 (2H, s), 3.38 (4H, m), 3.16 (1H, m), 2.96 (1H, m), 1.28 (6H, d, J=6.9 Hz).

7.3.492 5-Fluoro-N4-(3-isopropylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine R940288

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-isopropylphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to give 5-fluoro-N4-(3-isopropylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 23.43 min.; purity: 99 %; MS (m/e): 409 (M⁺), 411 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.90 (1H, s), 9.74 (1H, s), 8.28 (1H, d, J= 4.8 Hz), 8.06 (1H, m), 7.78 (1H, d, J= 7.2 Hz), 7.58 (1H, s), 7.4-7.3 (3H, m), 7.24 (1H, t, J= 8.4 Hz), 7.00 (1H, d, J= 7.25 Hz), 6.70 (1H, d, J= 7.25 Hz), 4.44 (2H, s), 2.93 (1H, sept, J= 6.9 Hz), 2.74 (3H, d, J= 4.8 Hz), 1.27 (6H, d, J= 6.9 Hz).

7.3.493 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-dimethylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926718)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and dimethylamine were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-

dimethylamino)carbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 8.06 (d, 1H, $J = 2.1$ Hz), 7.91 (d, 1H, $J = 3.6$ Hz), 7.57 (t, 1H, $J = 2.4$ Hz), 7.37 (d, 1H, $J = 9.0$ Hz), 7.28 (s, 1H), 7.19 (t, 1H, $J = 7.8$), 7.06 (s, 1H), 6.82-6.76 (m, 2H), 6.71 (dd, 1H, $J = 2.4$ and 7.8 Hz), 3.31 (s, 3H), 3.09 (s, 3H); ^{19}F NMR (CDCl_3): - 47292; LCMS: ret. time: 17.29 min.; purity: 92 %; MS (m/e): 408 (MH^+).

7.3.494 N4-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R945149)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (700 mg, 1.6 mmol) and piperazine (4 g, 46 mmol) gave N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (520 mg, 66%). ^1H NMR (CD_3OD): δ 2.22 (s, 3H), 2.75 (t, $J = 5.4$ Hz, 4H), 3.40 (t, $J = 4.8$ Hz, 2H), 3.54 (t, $J = 5.1$ Hz, 2H), 4.62 (s, 2H), 6.57 (ddd, $J = 1.5, 2.7$ and 7.5 Hz, 1H), 7.09 (dt, $J = 1.5$ and 8.1 Hz, 1H), 7.14 (t, $J = 7.8$ Hz, 1H), 7.28 (t, $J = 2.1$ Hz, 1H), 7.31 (dd, $J = 0.9$ and 2.7 Hz, 1H), 7.50 (d, $J = 2.7$ Hz, 1H), 7.88 (d, $J = 3.9$ Hz, 1H); ^{19}F NMR (282 MHz, CD_3OD): δ - 168.63; LCMS: ret. time: 14.99 min.; 93.88%; MS (m/e): 486.96 (MH^+).

7.3.495 N4-(4-*tert*-Butylphenyl)-5-fluoro-N2-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926713)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2-(N-methylaminocarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 8.05 (d, 1H, $J = 2.4$ Hz), 7.88 (d, 1H, $J = 4.2$ Hz), 7.57 (d, 2H, $J = 8.7$ Hz), 7.51-7.41 (m, 2H), 7.34-7.31 (m, 3H), 2.94 (s, 3H), 1.33 (s, 9H); ^{19}F NMR (CD_3OD): - 47682; LCMS: ret. time: 23.02 min.; purity: 90 %; MS (m/e): 434 (MH^+).

7.3.496 N4-(3,5-Dimethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926796)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,5-dimethoxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyridinediamine with methylamine hydrochloride gave N4-(3,5-dimethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.92 (d, 1H, J= 4.2 Hz), 7.42 (t, 1H, J= 1.8 Hz), 7.12 (m, 2H), 6.91 (d, 1H, J= 2.4 Hz), 6.59 (m, 1H), 6.22 (t, 1H, J= 1.8 Hz), 4.35 (s, 2H), 3.69 (s, 6H), 2.81 (s, 3H); LCMS: ret. time: 18.35 min.; purity: 93%; MS (m/e): 428 (MH⁺).

7.3.497 5-Ethoxycarbonyl-N4-(3,4-ethylenedioxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926800)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-ethoxycarbonyl-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-2,4-pyridinediamine with methylamine hydrochloride gave 5-ethoxycarbonyl-N4-(3,4-ethylenedioxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.05 (s, 1H), 9.34 (s, 1H), 8.69 (s, 1H), 7.95 (d, 1H, J= 4.8 Hz), 7.34 (dd, 1H, J= 1.2 and 7.8 Hz), 7.25 (bs, 2H), 7.13 (t, 1H, J= 8.1 Hz), 7.00 (bd, 1H, J= 9Hz), 6.81 (d, 1H, J= 8.7 Hz), 6.59 (dd, 1H, J= 1.5 and 8.4 Hz), 4.32 (s, 2H), 4.30 (q, 2H, J= 7.2 Hz), 4.21 (s, 4H), 2.63 and 2.62 (2s, 3H), 1.31 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 24.12 min.; purity: 91%; MS (m/e): 481 (MH⁺).

7.3.498 N4-(3,5-Dimethoxyphenyl)-5-ethoxycarbonyl-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926801)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxyphenyl)]-2,4-pyrimidinediamine, the reaction of N4-(3,5-dimethoxyphenyl)-5-ethoxycarbonyl-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-2,4-pyridinediamine with methylamine hydrochloride gave N4-(3,5-dimethoxyphenyl)-5-ethoxycarbonyl-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-

pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.20 (s, 1H), 9.96 (s, 1H), 8.73 (s, 1H), 7.90 (bs, 1H), 7.36 (d, 1H, J= 8.7 Hz), 7.28 (bs, 1H), 7.12 (t, 1H, J= 7.5 Hz), 6.84 (s, 2H), 6.59 (dd, 1H, J= 1.8 and 8.1 Hz), 6.25 (t, 1H, J= 2.4 Hz), 4.31 (m, 4H), 3.67 (s, 6H), 2.63 and 2.62 (2s, 3H), 1.31 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 25.50 min.; purity: 96%; MS (m/e): 482 (MH⁺).

7.3.499 N4-(4-*tert*-Butylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926714)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.90 (d, 1H, J= 3.3 Hz), 7.61 (d, 2H, J= 8.7 Hz), 7.40-7.33 (m, 3H), 7.14-7.11 (m, 2H), 6.62-6.57 (m, 1H), 4.36 (s, 2H), 2.79 (s, 3H), 1.31 (s, 9H); ¹⁹F NMR (CD₃OD): -47514; LCMS: ret. time: 23.70 min.; purity: 93 %; MS (m/e): 424 (MH⁺).

7.3.500 N4-(3-Hydroxyphenyl)-5-trifluoromethyl-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926742)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3-hydroxyphenyl)-5-trifluoromethyl-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(3-hydroxyphenyl)-5-trifluoromethyl-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 19.11 min.; purity: 99 %; MS (m/e): 434 (MH⁺).

7.3.501 5-Fluoro-N4-[(1H)-indol-6-yl]-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926745)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(1H)-indol-6-yl]-4-pyrimidineamine and 3-(N-methylamino)carbonylmethyleneoxyaniline were reacted to yield 5-fluoro-N4-[(1H)-indol-

6-yl]-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine.

LCMS: ret. time: 17.41 min.; purity: 93 %; MS (m/e): 407(MH⁺).

7.3.502 N4-(3,5-Dimethyl-4-methoxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R945156)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and piperazine gave N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 2.23 (s, 6H), 3.24 (m, 4H), 3.71 (s, 3H), 3.72-3.81 (m, 4H), 4.75 (s, 2H), 6.81 (dt, J= 1.2 and 8.1 Hz, 1H), 7.10-7.13 (m, 2H), 7.24 (d, J= 8.7 Hz, 1 H), 7.29 (s, 2H), 7.98 (d, J= 4.8 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ -163.88; LCMS: ret. time: 15.94 min.; purity: 100%; MS (m/e): 481.12 (MH⁺).

7.3.503 N4-(3-*tert*-Butylphenyl)-5-fluoro-N2-[2-(N-methylamino)carbonylbenzofur-5-yl]-2,4-pyrimidinediamine R940291

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3-*tert*-butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to give N4-(3-*tert*-butylphenyl)-5-fluoro-N2-[2-(N-methylamino)carbonylbenzofur-5-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 23.05 min.; purity: 100 %; MS (m/e): 434 (MH⁺); ¹³C NMR (DMSO-d₆): δ 9.6 (1H, s), 9.57 (1H, s), 8.75 (1H, m), 8.25 (1H, s), 8.15 (1H, s), 7.93 (1H, d, J= 8.5 Hz), 7.47 (3H, m), 7.44 (1H, s), 7.36 (1H, t, J= 8.5 Hz), 7.25 (1H, d, J= 8.5 Hz), 2.89 (3H, d, J= 4.5 Hz), 1.33 (9H, s).

7.3.504 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926505)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxy)phenyl)-2,4-pyrimidinediamine and 2-hydroxyethylamine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H

NMR (CD₃OD): δ 7.87 (d, 1H, J= 3.6 Hz), 7.37 (t, 1H, J= 1.8 Hz), 7.24 (d, 1H, J= 2.4 Hz), 7.13 (m, 2H), 7.08 (dd, 1H, J= 2.1 and 8.1 Hz), 6.77 (m, 1H), 4.38 (s, 2H), 4.22 (s, 3H), 3.63 (t, 2H), 3.40 (t, 2H, J= 6 Hz); LCMS: ret. time: 16.72 min.; purity: 98%; MS (m/e): 456 (MH⁺).

5 **7.3.505 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926746)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 3-amino-1,2-propanediol were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.84 min.; purity: 96 %; MS (m/e): 444 (MH⁺).

15 **7.3.506 5-Fluoro-N2-[2-(2-hydroxy-1,1-dimethylethylamino)carbonylbenzofuran-5-yl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926715)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and 2-amino-2-methylpropanol were reacted to yield 5-fluoro-N2-[2-(2-hydroxy-1,1-dimethylethylamino)carbonylbenzofuran-5-yl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.41 (s, 1H), 9.28 (s, 1H), 9.22 (s, 1H), 8.18 (t, 1H, J= 2.4, Hz), 8.09 (d, 1H, J= 3.6 Hz), 7.56 (dd, 1H, J= 2.4 and 8.7 Hz), 7.47 (d, 1H, J= 8.7 Hz), 7.32 (s, 1H), 7.26-7.21 (m, 1H), 7.13- 7.07 (m, 2H), 6.53 (d, 1H, J= 8.7 Hz), 5.05 (t, 1H, J= 5.7 Hz), 3.46 (d, 2H, J= 5.7 Hz), 1.32 (s, 6H); LCMS: ret. time: 17.93 min.; purity: 97 %; MS (m/e): 452 (MH⁺).

30 **7.3.507 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926730)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield 5-fluoro-N4-(4-

isopropoxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.93 (d, 1H, J= 3.0 Hz), 7.47 (d, 2H, J= 9.3 Hz), 7.42 (t, 1H, J= 1.8 Hz), 7.17 (t, 1H, J= 8.1 Hz), 7.10 (bs, 1H), 7.00 (dd, 1H, J= 1.8 and 9.3 Hz), 6.89 (d, 2H, J= 9.3 Hz), 6.80 (d, 1H, J= 1.8 Hz), 6.58 (bs, 1H), 6.50 (dd, 1H, J= 1.5 and 8.1 Hz), 4.51 (2q, 1H, J= 5.7 Hz), 4.44 (s, 2H), 2.88 (d, 3H, J= 4.5 Hz), 1.33 (d, 6H, J= 5.7 Hz); ¹⁹F NMR (CDCl₃): - 47198; LCMS: ret. time: 19.66 min.; purity: 97 %; MS (m/e): 426 (MH⁺).

7.3.508 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R945170)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)methyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2-(4-cyanomethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 2.91 (d, J= 5.1 Hz, 3H), 4.48 (s, 2H), 6.61 (ddd, J= 0.9, 2.7 and 8.1 Hz, 1H), 6.63 (br, 1H), 6.76 (d, J= 3.0 Hz, 1H), 6.84-6.89 (m, 4H), 7.18 (t, J= 8.1 Hz, 1H), 7.44 (d, J= 8.7 Hz, 2H), 7.51 (t, J= 2.1 Hz, 1H), 7.92 (d, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ -167.70; LCMS: ret. time: 14.32 min.; purity: 100%; MS (m/e): 383.98 (MH⁺).

7.3.509 5-Fluoro-N4-(3-isopropoxyphenyl)-N2-[2-(N-morpholino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926489)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-isopropoxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidineamine with morpholine gave 5-fluoro-N4-(3-isopropoxyphenyl)-N2-[2-(N-morpholino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.01 (d, 1H, J= 1.2 Hz), 7.95 (bs, 1H), 7.43-7.38 (m, 2H), 7.29 (s, 1H), 7.25-7.11 (m, 4H), 6.97 (bs, 1H), 6.73 (m, 1H), 6.67 (bdd, 1H), 4.48 (sept, 1H, J= 5.7 Hz), 3.87 (m, 4H), 3.79 (m, 4H), 1.30 (d, 6H, J= 5.7 Hz), LCMS: ret. time: 22.12 min.; purity: 98%; MS (m/e): 492 (MH⁺).

7.3.510 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926772)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine with piperazine gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.91 (d, 1H, J= 3.6 Hz), 7.42 (t, 1H, J= 2.4 Hz), 7.20-7.07 (m, 5H), 6.55 (m, 2H), 4.63 (s, 2H), 3.54 (t, 2H, J= 6 Hz), 3.40 (t, 2H, J= 5.1 Hz), 2.76 (t, 4H, J= 5.4 Hz); LCMS: ret. time: 12.98 min.; purity: 92%; MS (m/e): 439 (MH⁺).

7.3.511 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926506)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with 2-hydroxyethylamine gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 14.95 min.; purity: 96%; MS (m/e): 414 (MH⁺).

7.3.512 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926508)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-ethoxy or methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with methylamine hydrochloride gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.64 (bs, 1H), 9.58 (bs, 1H), 8.15 (d, 1H, J= 4.2 Hz), 7.95 (bd, 1H), 7.25 (bd, 2H, J= 6.6 Hz), 7.16-7.07 (m, 4H), 6.53 (m, 2H), 4.35 (s, 2H), 2.64 and 2.62 (2s, 3H); LCMS: ret. time: 15.66 min.; purity: 98%; MS (m/e): 384 (MH⁺).

7.3.513 5-Fluoro-N4-[3,4-(1,1,2,2-tetrafluoroethylenedioxy)phenyl]-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926732)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[3,4-(1,1,2,2-tetrafluoroethylenedioxy)phenyl]-4-pyrimidineamine and methylamine hydrochloride were reacted to yield 5-fluoro-N4-[3,4-(1,1,2,2-tetrafluoroethylenedioxy)phenyl]-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.65 (s, 1H), 9.37 (s, 1H), 8.16 (d, 1H, J= 3.6 Hz), 8.14 (d, 1H, J= 2.4 Hz), 7.97 (d, 1H, J= 4.8 Hz), 7.65 (dd, 1H, J= 2.4 and 8.7 Hz), 7.41 (d, 1H, J= 9.3 Hz), 7.34 (t, 1H, J= 2.4 Hz), 7.27 (d, 1H, J= 8.1 Hz), 7.13 (t, 1H, J= 8.1 Hz), 6.51 (dd, 1H, J= 2.1 and 7.5 Hz), 4.36 (s, 2H), 2.63 (d, 3H, J= 4.8 Hz); ¹⁹F NMR (DMSO-d₆): - 25765 (pent, 2F), - 25830 (pent, 2F), - 46309; LCMS: ret. time: 24.85 min.; purity: 95 %; MS (m/e): 497 (MH⁺).

7.3.514 N4-(3,5-Dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R940254)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,5-dimethyl-4-hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with morpholine gave N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-(N-morpholinocarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 18.38 min.; purity: 92 %; MS (m/e): 468 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.20 (1H, s), 9.10 (1H, s), 8.15 (1H, s), 8.11 (1H, d, J= 3.9 Hz), 7.43 (1H, d, J= 8.1 Hz), 7.32 (3H, m), 7.14 (1H, t, J= 8.1 Hz), 6.54 (1H, dd, J= 8.1 and 2.0 Hz), 4.77 (2H, s), 3.64 (4H, m), 3.54-3.45 (4H, m), 2.24 (6H, s).

7.3.515 N4-(3-*tert*-Butylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R940276)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3-*tert*-butylphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with methylamine hydrochloride gave N4-(3-*tert*-butylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS:

ret. time: 22.90 min.; purity: 99 %; MS (m/e): 424 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.46 (1H, s), 9.34 (1H, s), 8.08 (1H, d, J= 3.9 Hz), 7.90 (1H, m), 7.30 (1H, d, J= 8.1 Hz), 7.46 (1H, m), 7.26 (1H, m), 7.20 (2H, m), 7.10-7.03 (2H, m), 6.47 (1H, d, J= 8.1 Hz), 4.26 (2H, s), 2.59 (3H, d, J= 4.5 Hz), 1.20 (9H, s).

5 **7.3.516 N4-(3-*tert*-Butylphenyl)-N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R940277)**

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3-*tert*-butylphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with 2,3-dihydroxypropylamine gave N4-(3-*tert*-butylphenyl)-N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: retn. time: 20.46 min.; purity: 100 %; MS (m/e): 484 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.38 (1H, s), 9.29 (1H, s), 8.20 (1H, d, J= 3.9 Hz), 8.00 (1H, d, J= 8.3 Hz), 7.93 (1H, t, J= 5.5 Hz), 7.60 (1H, m), 7.47 (1H, m), 7.41-7.17 (4H, m), 6.59 (1H, dd, J= 8.3 and 2 Hz), 3.43 (2H, s), 3.39 (4H, m), 3.16 (1H, m), 1.36 (9H, s).

7.3.517 N4-(3,3-Dihydroisobenzofuran-1-one-6-yl)-N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine R940293

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-[3-(ethoxycarbonylmethyleneoxy)phenyl]-N4-(3,3-dihydroisobenzofuran-1-one-6-yl)-5-fluoro-2,4-pyrimidinediamine and 3-amino-1,2-propanediol were reacted to N4-(3,3-dihydroisobenzofuran-1-one-6-yl)-N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 13.92 min.; purity: 92 %; MS (m/e): 483 (M⁺); ¹H NMR (DMSO-d₆): δ 9.80 (1H, s), 9.46 (1H, s), 8.37-8.27 (2H, m), 8.21 (1H, s), 7.96 (1H, t, J= 4.6 Hz), 7.24 (1H, d, J= 9 Hz), 7.44 (1H, s), 7.37 (1H, d, J= 9 Hz), 7.23 (1H, t, J= 8 Hz), 6.60 (1H, dd, J= 7 and 3.75 Hz), 5.49 (2H, s), 4.46 (2H, s), 3.38 (4H, m), 3.2-3.1 (1H, m).

7.3.518 N4-(3,4-Dimethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926733)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-dimethoxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(3,4-dimethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.95 (d, 1H, J= 3.6 Hz), 7.45 (t, 1H, J= 1.8 Hz), 7.21-7.17 (m, 2H), 7.05 (dd, 1H, J= 2.7 and 8.7 Hz), 6.96-6.90 (m, 2H), 6.87 (d, 1H, J= 9.0 Hz), 6.72 (d, 1H, J= 2.4 Hz), 6.67-6.58 (m, 1H), 6.52 (dd, 1H, J= 3.6 and 8.1 Hz), 4.39 (s, 2H), 3.89 (s, 3H), 3.78 (s, 3H), 2.90 (d, 3H, J= 4.8 Hz); LCMS: ret. time: 17.09 min.; purity: 98 %; MS (m/e): 428 (MH⁺).

7.3.519 N2-[3-(N-2,3-Dihydroxypropylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926734)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-dimethoxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 3-amino-1,2-propanediol were reacted to yield N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-dimethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.05 (d, 1H, J= 4.2 Hz), 7.38-7.34 (m, 2H), 7.31-7.26 (m, 2H), 7.07 (t, 1H, J= 8.4 Hz), 6.89 (d, 1H, J= 8.7 Hz), 6.46 (dd, 1H, J= 2.4 and 8.4 Hz), 4.36 (s, 2H), 3.72 (s, 3H), 3.68 (s, 3H), 3.32-3.24 (m, 3H), 3.03 (dd, 1H, J= 6.9 and 13.5 Hz); ¹⁹F NMR (DMSO-d₆): - 46574; LCMS: ret. time: 14.85 min.; purity: 94 %; MS (m/e): 488 (MH⁺).

7.3.520 5-Fluoro-N4-(3-methoxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926738)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-methoxyphenyl)-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield 5-fluoro-N4-(3-

methoxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 18.40 min.; purity: 98 %; MS (m/e): 398 (MH⁺).

7.3.521 N2-[3-(N-2,3-Dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine (R926739)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine and 3-amino-1,2-propanediol were reacted to yield N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 16.66 min.; purity: 99 %; MS (m/e): 458 (MH⁺).

7.3.522 N4-(3,5-Dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R945140)

In a manner analogous to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and piperazine gave N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 2.18 (s, 6H), 2.72 (q, J= 5.1 Hz, 4H), 3.32 (t, 2H), 3.52 (t, J= 5.1 Hz, 2H), 4.55 (s, 2H), 6.56 (ddd, J= 1.2, 2.4 and 8.1 Hz, 1 H), 7.03 (ddd, J= 1.2, 1.8 and 8.1 Hz, 1H), 7.11 (t, J= 8.1 Hz, 1H), 7.20 (s, 2H), 7.35 (t, J= 2.1 Hz, 1H), 7.84 (d, J= 3.9 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 168.78; LCMS: ret. time: 14.32 min.; purity: 88.37%; MS (m/e): 467.06 (MH⁺).

7.3.523 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-morpholino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926488)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine with morpholine gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-morpholino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.19 (t, 1H, J= 1.5 Hz), 7.90 (d, 1H, J= 3.9 Hz), 7.44 (d, 2H, J= 0.9 Hz), 7.28 (s, 1H), 7.21 (t, 1H,

J= 2.4 Hz), 7.15 (t, 1H, J= 7.5 Hz), 7.08 (m, 1H), 7.61 (bd, 1H, J= 6.9 Hz), 3.8 (m, 4H), 3.65 (m, 4H); LCMS: ret. time: 17.21 min.; purity: 83%; MS (m/e): 450 (MH⁺).

7.3.524 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926493)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine with methylamine hydrochloride gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.71 (d, 1H, J= 4.8 Hz), 8.00-7.92 (m, 2H), 7.56-7.52 (m, 1H), 7.44-7.39 (m, 2H), 7.12 (m, 2H), 6.69 (bdd, 1H), 2.96 and 2.94 (2s, 3H).

7.3.525 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-2-hydroxyethylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926497)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine with 2-hydroxyethylamine gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-2-hydroxyethylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.18 (d, 1H, J= 1.8 Hz), 7.80 (bs, 1H), 7.60 (m, 1H), 7.34-7.16 (m, 3H), 7.10 9t, 1H, 8.4 Hz), 6.85 (bdd, 1H), 6.62 (dd, 1H, J= 1.5 and 8.1 Hz), 3.70 (t, 2H, J= 4.8 Hz), 3.52 (t, 2H, J= 4.0 Hz); LCMS: ret. time: 14.49 min.; purity: 97%; MS (m/e): 424 (MH⁺).

7.3.526 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926500)

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine with piperazine gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.19 (t, 1H, J= 1.2 Hz), 7.90 (d, 1H, J= 3.9 Hz), 7.43 (d, 2H, J= 1.2 Hz), 7.25-7.06 (m, 4H), 6.59

(m, 1H), 3.80 (m, 4H), 2.95 (m, 4H); LCMS: ret. time: 12.97 min.; purity: 79%; MS (m/e): 449 (MH⁺).

7.3.527 5-Cyano-N4-(3-hydroxyphenyl)-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R925844)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-N-methylaminocarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-cyano-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine with methylamine hydrochloride gave 5-cyano-N4-(3-hydroxyphenyl)-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 18.83 min.; purity: 96 %; MS (m/e): 391 (MH⁺).

7.3.528 5-Cyano-N4-[4-(N-cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925845)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-cyano-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine reacted with cyclopropylmethylamine to give 5-cyano-N4-[4-(N-cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 22.47 min.; purity: 100 %; MS (m/e): 431 (MH⁺).

7.3.529 5-Cyano-N4-(3-hydroxyphenyl)-N2-[4-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R925846)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-cyano-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine reacted with 2,3-dihydroxypropylamine to give 5-cyano-N4-(3-hydroxyphenyl)-N2-[4-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 15.84 min.; purity: 100 %; MS (m/e): 451 (MH⁺).

7.3.530 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3-trifluoromethylphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-N4-(3-trifluoromethylphenyl)-2,4-pyrimidineamine with methylamine hydrochloride gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3-trifluoromethylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 21.98 min., purity: 86%, MS (m/e): 436 (MH⁺).

7.3.531 N4-[4-(4,5-Dichloro-1H-imidazol-1-ylphenyl)]-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926812)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N-4-[4-(4,5-dichloro-1H-imidazol-1-ylphenyl)]-5-fluoro-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with methylamine hydrochloride gave N4-[4-(4,5-dichloro-1H-imidazol-1-ylphenyl)]-5-fluoro N2-(3-[N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 21.02 min., purity: 100%, MS (m/e): 502 (MH⁺).

7.3.532 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-methylaminocarbonyllindol-7-yl)- 2,4-pyridinediamine (R926815)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-N4-(3-trifluorophenyl)-2,4-pyrimidineamine with methylamine hydrochloride gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-methylaminocarbonyllindol-7-yl)- 2,4-pyridinediamine. LCMS: ret. time: 17.97 min., purity: 97%, MS (m/e): 435 (MH⁺).

7.3.533 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926484)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2-

(3-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and morpholine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.94 (bs, 1H), 7.35 (t, 1H, J= 2.4 Hz), 7.24 (m, 1H), 7.19 (t, 1H, J= 8.1 Hz), 7.10 (bdd, 1H, J= 6.9 Hz), 6.95 (m, 2H), 6.85 (d, 1H, J= 8.1 Hz), 6.94 (s, 1H), 6.58 (dd, 1H, J= 1.8 and 2.8 Hz), 4.64 (s, 2H), 4.27 9s, 4H), 3.62 (m, 4H), 3.55 (m, 4H); LCMS: ret. time: 18.45 min.; purity: 100%; MS (m/e): 482 (MH⁺).

7.3.534 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-morpholino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926492)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine with morpholine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-morpholino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.27 (s, 1H), 9.17 (s, 1H), 8.14 (d, 1H, J= 2.4 Hz), 8.05 (d, 1H, J= 5.6 Hz), 7.58-7.46 (m, 2H), 7.27 (m, 1H), 7.15 (dd, 1H, J= 2.4 and 9 Hz), 6.80 9m, 1H), 4.24 (s, 4H), 3.80-3.45 (m, 8H); LCMS: ret. time: 19.97 min.; purity: 76%; MS (m/e): 492 (MH⁺).

7.3.535 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926496)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and methylamine hydrochloride gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.06 (s, 1H), 7.85 (d, 1H, J= 3.3 Hz), 7.42 (d, 2H, J= 1.2 Hz), 7.35 (s, 1H), 7.29 (d, 1H, J= 2.4 Hz), 6.99 (dd, 1H, J= 3.3 and 8.7 Hz), 6.78 (d, 1H, J= 8.7 Hz), 4.24 (s, 4H), 2.94 (s, 3H); LCMS: ret. time: 18.05 min.; purity: 99%; MS (m/e): 436 (MH⁺).

7.3.536 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-2-hydroxyethylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926498)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine with 2-hydroxyethylamine yielded N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-2-hydroxyethylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.07 (d, 1H, J= 1.2 Hz), 7.86 (d, 1H, J= 3.9 Hz), 7.43 (d, 2H, J= 1.5 Hz), 7.38 (s, 1H), 7.29 (d, 1H, J= 2.4 Hz), 6.98 (dd, 1H, J= 2.1 and 9 Hz), 6.78 (d, 1H, J= 8.7 Hz), 4.23 (s, 4H), 3.72 (t, 2H, J= 5.7 Hz), 3.53 (t, 2H, J= 6.0 Hz); LCMS: ret. time: 16.21 min.; purity: 97%; MS (m/e): 466 (MH⁺).

7.3.537 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926499)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and piperazine yielded N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.26 (s, 1H), 9.16 (s, 1H), 8.12 (d, 1H, J= 1.8 Hz), 8.04 (d, 1H, J= 3.6 Hz), 7.49 (d, 2H), 7.30 (d, 1H, J= 2.4 Hz), 7.20 (s, 1H), 7.15 (bdd, 1H, J= 3 Hz), 6.79 (d, 1H, J= 8.7 Hz), 4.22 (s, 4H), 2.48 (s, 3H); LCMS: ret. time: 14.61 min.; purity: 94%; MS (m/e): 491 (MH⁺).

7.3.538 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926503)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and piperazine were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 9.14 (bs, 2H), 8.04 (d, 3.6 Hz), 7.32-7.20 (m, 4H), 7.06 (t, 1H, J= 8.1

Hz), 6.79 (d, d, 1H, J= 9 Hz), 6.43 (bd, 1H, J= 9.9 Hz), 4.64 (s, 2H), 4.20 (bs, 4H), 3.29 (m, 4H), 2.59 (m, 4H); LCMS: ret. time: 14.92 min.; purity: 99%; MS (m/e): 481 (MH⁺).

7.3.539 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-2-hydroxy-1,1-dimethylethylamino)carboxymethyleneoxyphenyl]-2,4-pyrimidinediamine (R926764)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 2-amino-2-methylpropanol gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-2-hydroxy-1,1-dimethylethylamino)carboxymethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.95 (d, 1H, J= 2.7 Hz), 7.47 (t, 1H, J= 2.4 Hz), 7.20 (t, 1H, J= 8.1 Hz), 7.03 (dd, 1H, J= 1.2 and 8.1 Hz), 6.98 (dd, 1H, J= 3 and 8.2 Hz), 6.93 (s, 1H), 6.84 (d, 1H, J= 8.7 Hz), 6.66 (d, 1H, J= 3 Hz), 6.57 (bs, 1H), 6.53 (m, 1H), 4.65 (m, 1H), 4.39 (s, 2H), 4.28 (s, 4H), 3.63 (d, 2H, J= 5.7 Hz), 1.31 (s, 6H); LCMS: ret. time: 19.19 min.; purity: 89%; MS (m/e): 484 (MH⁺).

7.3.540 N2-[3-(N-Cyclohexylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926765)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and cyclohexylamine gave N2-[3-(N-cyclohexylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.94 (d, 1H, J= 3.3 Hz), 7.41 (t, 1H, J= 2.4 Hz), 7.28 (d, 1H, J= 2.4 Hz), 7.20 (t, 1H, J= 7.5 Hz), 7.04 (dd, 1H, J= 1.2 and 8.1 Hz), 6.95 (m, 2H), 6.85 (d, 1H, J= 8.7 Hz), 6.68 (d, 1H, J= 3.0 Hz), 6.53 (dd, 1H, J= 2.4 and 8.4 Hz), 6.45 (bd, 1H, J= 8.1 Hz), 4.43 (s, 2H), 4.24 (s, 4H), 3.85 (m, 1H), 1.90 (m, 2H), 1.75-1.55 (m, 2H), 1.45-1.05 (m, 6H); LCMS: ret. time: 23.70 min.; purity: 97%; MS (m/e): 494 (MH⁺).

7.3.541 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-[N-methyl-N-(2-hydroxyethyl)amino]carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926766)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-

ethylenedioxyphenyl)-5-fluoro-N2-[3-methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and N-methyl-N-2-hydroxyethylamine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[N-methyl-N-(2-hydroxyethyl)amino]carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.93 (d, 1H, J= 3 Hz), 7.92 (bs, 1H), 7.35 (t, 1H, J= 2.4 Hz), 7.18 (m, 1H), 7.06 (dd, 1H, J= 1.2 and 8.7 Hz), 6.97 (t, 1H, J= 2.4 Hz), 6.94 (m, 1H), 6.85 (d, 1H, J= 8.7 Hz), 6.70 (bd, 1H), 6.59 (dd, 1H, J= 1.8 and 8.1 Hz), 4.66 (s, 2H), 4.28 (s, 4H), 3.79 (t, 2H, J= 5.4 Hz), 3.56 (t, 3H, J= 5.4 Hz), 3.10 (s, 3H); LCMS: ret. time: 16.64 min.; purity: 97%; MS (m/e): 470 (MH⁺).

7.3.542 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-homopiperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926767)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and homopiperazine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-homopiperazinocarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.27 (s, 1H), 9.17 (d, 1H, J= 1.2 Hz), 8.14 (s, 1H), 8.05 (d, 1H, J= 3.6 Hz), 7.54-7.46 (m, 2H), 7.30 (d, 1H, J= 2.4 Hz), 7.24 (s, 1H), 7.17 (dd, 1H, J= 2.4 and 8.7 Hz), 6.80 (d, 1H, J= 8.7 Hz), 4.22 (s, 4H), 3.79 (m, 2H), 3.65 (m, 2H), 3.01 (m, 2H), 2.89 (m, 2H), 1.90 (m, 1H), 1.80 (m, 1H); ¹⁹F NMR (DMSO-d₆): - 46687; LCMS: ret. time: 14.99 min.; purity: 77%; MS (m/e): 505 (MH⁺).

7.3.543 N4-(3,4-Ethylenedioxyphenyl)-N2-[3-(N,N-dimethylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R925755)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and N,N-dimethylamine hydrochloride gave N4-(3,4-ethylenedioxyphenyl)-N2-[3-(N,N-dimethylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.16 (d, 1H, J= 1.2 Hz), 9.15 (s, 1H), 8.04 (d, 1H, J= 5.6 Hz), 7.30-7.21 (m, 4H), 7.06 (t, 1H, J= 9Hz), 6.78 (d, 1H, J= 9Hz), 6.43 (m, 1H), 4.65 (s, 2H), 4.21 (s, 4H), 2.94 (s, 3H), 2.82 (s, 3H); LCMS: ret. time: 18.70 min.; purity: 83%; MS (m/e): 440 (MH⁺).

**7.3.544 N2-[3-[N,N-Bis-(2-hydroxyethylamino)]
carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-
5-fluoro-2,4-pyrimidinediamine (R926781)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and N,N-bis(hydroxyethyl)amine gave N2-[3-[N,N-bis-(2-hydroxyethylamino)]carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.86 (d, 1H, J= 3.6 Hz), 7.25 (m, 2H), 7.17-7.03 (m, 3H), 6.78 (d, 1H, J= 9Hz), 6.58 (bd, 1H), 4.80 (s, 2H), 4.23 (s, 4H), 3.71 (t, 4H, J= 4.8 Hz), 3.53 (t, 2H, J= 6Hz), 3.49 (t, 3H, J= 5.4 Hz); LCMS: ret. time: 16.25 min.; purity: 94%; MS (m/e): 500 (MH⁺).

**7.3.545 N2-[3-(N-2,3-Dihydroxypropylamino)
carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-
5-fluoro-2,4-pyrimidinediamine (R926782)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and 2,3-dihydroxypropylamine gave N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.86 (d, 1H, J= 4.2 Hz), 7.37 (t, 1H, J= 1.8 Hz), 7.24 (d, 1H, J= 2.4 Hz), 7.14 (m, 2H), 7.09 (dd, 1H, J= 2.4 and 9 Hz), 6.78 (d, 1H, J= 8.7 Hz), 6.59 (m, 1H), 4.39 (s, 2H), 4.22 (s, 4H), 3.73 (m, 1H), 3.48 (m, 4H); ¹⁹F NMR (CD₃OD): - 47575; LCMS: ret. time: 15.97; purity: 98%; MS (m/e): 486 (MH⁺).

7.3.546 N2-[2-(N-2,3-Dihydroxypropylamino)carbonylbenzofuran-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926783)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and 2,3-dihydroxypropylamine gave N2-[2-(N-2,3-dihydroxypropylamino)carbonylbenzofuran-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.86 (d, 1H, J= 4.2 Hz), 7.35 (t, 1H, J= 1.2

Hz), 7.24 (d, 1H, J= 3 Hz), 7.15 (m, 2H), 7.07 (dd, 1H, J= 2.1 and 8.7 Hz), 6.78 (d, 1H, J= 8.7 Hz), 6.59 (m, 1H), 4.40 (s, 1H), 4.23 (s, 4H), 4.03 (t, 1H, J= 5.7 Hz), 3.67 (d, 2H, 3.6 Hz), 3.65 (d, 2H, J= 4.2 Hz); ¹⁹F NMR (CD₃OD): -47578; LCMS: ret. time: 15.72 min.; purity: 99%; MS (m/e): 486 (MH⁺).

5 **7.3.547 N2-[3-(N-1,3-Dihydroxy-2-propylamino) carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926784)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethylenedioxyphenyl)-2,4-pyrimidinediamine and 2-amino-1,3-propanediol gave N2-[3-(N-1,3-dihydroxy-2-propylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.08 (bd, 1H), 7.86 (bs, 1H), 7.44 (s, 2H), 7.39 (s, 1H), 7.29 (d, 1H, J= 2.4 Hz), 6.97 (dd, 1H, J= 2.4 and 8.7 Hz), 6.78 (d, 1H, J= 8.7 Hz), 4.24 (s, 4H), 3.84 (m, 1H), 3.56 (m, 2H), 3.44 (m, 2H); LCMS: ret. time: 16.63 min.; purity: 97%; MS (m/e): 496 (MH⁺).

7.3.548 N2-[2-(N-1,3-Dihydroxy-2-propylamino)carbonylbenzofuran-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926785)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and 2-amino-1,3-propanediol gave N2-[2-(N-1,3-dihydroxy-2-propylamino)carbonylbenzofuran-5-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.08 (t, 1H, J= 1.8 Hz), 7.86 (d, 1H, J= 3.9 Hz), 7.45 (s, 2H), 7.41 (s, 1H), 7.29 (d, 1H, J= 2.4 Hz), 6.97 (dd, 1H, J= 3 and 8.7 Hz), 6.77 (d, 1H, J= 8.7 Hz), 4.24 (s, 4H), 4.19 (t, 1H, J= 5.7 Hz), 3.75 (d, 4H, J= 5.4 Hz); ¹⁹F NMR (CD₃OD): - 47745; LCMS: ret. time: 15.09 min., purity: 97%; MS (m/e): 496 (MH⁺).

30 **7.3.549 N4-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R940265)**

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3-

chloro-4-hydroxy-5-methylphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with morpholine gave N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine.

LCMS: ret. time: 18.66 min.; purity: 92 %; MS (m/e): 487 (M^+), 489 (MH^+); ^1H NMR (DMSO-d₆): 9.28 (2H, s), 9.01 (1H, s), 8.17 (1H, d, J = 3.6 Hz), 7.65 (1H, d, J = 2.4 Hz), 7.5

(1H, d, J = 2.7 Hz), 7.42 (1H, d, J = 6.6 Hz), 7.29 (1H, s), 7.18 (1H, t, J = 8.1 Hz), 6.57 (1H, dd, J = 6.6 and 2.2 Hz), 4.79 (2H, s), 3.67 (4H, m), 3.52 (4H, m), 2.29 (3H, s).

7.3.550 N4-(3,5-Dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950187)

N4-(3,5-Dichlorophenyl-4-hydroxy)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (0.5 g, 1.1 mmol) was dissolved in EtOH:morpholine (4 ml : 4ml) and the mixture was refluxed for 1 day (100 °C oil-bath temperature). The mixture was cooled to 22 °C, diluted with water and brine, filtered, and dried under reduced pressure to give N4-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ^1H NMR (DMSO-d₆): δ 9.35 (s, 1H), 9.22 (s, 1H), 8.09 (d, 1H, J = 3.6 Hz), 7.94 (m, 1H), 7.75 (m, 1H), 7.27 (m, 1H), 7.18 (m, 1H), 7.12 (t, 1H, J = 8.4 Hz), 6.44 (m, 1H), 4.64 (s, 2H), 3.39 (m, 4H), 2.68 (m, 4H); LCMS purity: 92.6%; MS (m/e): 507.89 (M^+ , 100).

7.3.551 N4-(3,5-Dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950188)

In like manner to the preparation of N4-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,5-dichloro-4-hydroxyphenyl)-N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine and piperazine were reacted to prepare N4-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 15.26 min.; purity: 88.5%; MS (m/e): 506.89 (MH^+).

7.3.552 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926776)

In like manner to the preparation of N4-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3-

ethoxycarbonylmethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine with methylamine hydrochloride gave N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 16.94 min.; purity: 73%; MS (m/e): 426 (MH⁺).

5 **7.3.553 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-(4-methylaminocarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R945173)**

In a manner analogous to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(4-methylaminocarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine, N4-(4-cyanomethyleneoxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and methylamine hydrochloride salt gave N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-(4-methylaminocarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (acetone-d₆): δ 2.80 (d, 3H), 4.21-4.28 (m, 4H), 4.47 (s, 2H), 6.71 (d, J= 8.7 Hz, 1H), 6.96 (d, J= 9.0 Hz, 2H), 7.06 (dd, J= 2.7 and 9.0 Hz, 1H), 7.41 (d, J= 2.4 Hz, 1H), 7.74 (d, J= 9.0 Hz, 2H), 7.93 (d, J= 3.6 Hz, 1H), 8.20 (br, 1H, NH), 8.41 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-d₆): δ - 169.05; LCMS: ret. time: 17.47 min.; purity: 98.99%; MS (m/e): 425.89 (MH⁺).

7.3.554 N2-[4-(2-N,N-Dimethylaminoethyl)oxyphenyl]-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R909253)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N2-chloro-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-4-pyrimidineamine and 4-(2-N,N-dimethylaminoethyl)oxyaniline were reacted to yield N2-[4-(2-N,N-dimethylaminoethyl)oxyphenyl]-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.0 (d, 1H J= 4 Hz), 7.42 (m, 2H), 7.24 (m, 2H), 7.05 (m, 2H), 6.85 (m, 1H), 4.39 (s, 2H), 4.30 (m, 2H), 3.66 (m, 2H), 3.04 (s, 6H), 2.83 (s, 3H); LCMS: ret. time: 14.0 min.; purity: 96 %; MS (m/e): 455 (MH⁺).

7.3.555 N2-(1,4-Benzoxazin-6-yl)-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R909247)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N2-chloro-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-4-pyrimidineamine and 6-amino-1,4-benzoxazine were reacted to yield N2-(1,4-

benzoxazin-6-yl)-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H (DMSO-d₆): δ 8.0 (d, 1H), 7.6 (m, 1H), 7.42 (m, 1H), 7.20 (m, 1H), 6.95 (m, 1H), 6.76 (m, 1H), 6.56 (m, 1H), 4.43 (s, 2H), 4.05 (m, 2H), 3.25 (s, 3H), 3.13 (m, 2H); LCMS: ret time: 17.67 min.; MS (m/e): 425 (MH⁺).

5 **7.3.556 N2-(4-Dihydrobenzofuranyl)-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R909249)**

10 In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N2-chloro-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-4-pyrimidineamine and 5-amino-2,3-dihydrobenzofuran were reacted to yield N2-(4-dihydrobenzofuranyl)-5-fluoro-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.09 (d, 1H), 8.00 (m, 1H), 7.42 (m, 2H), 7.05 (m, 1H), 6.96 (m, 1H), 6.76 (m, 1H), 6.58 (m, 1H), 4.53 (m, 2H), 4.25 (s, 2H), 3.15 (m, 2H), 2.70 (m, 3H); LCMS: ret time: 19.24 min; MS (m/e): 410 (MH⁺).

15 **7.3.557 N2-(3-*tert*-Butylphenyl)-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R940267)**

20 In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2-(3-*tert*-butylphenyl)-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with methylamine hydrochloride gave N2-(3-*tert*-butylphenyl)-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 22.22 min.; purity: 97 %; MS (m/e): 424 (MH⁺); ¹□□□□ (CDCl₃): δ 7.98 (2H, m), 7.76 (2H, m), 7.56 (1H, t, J= 1.3 Hz), 7.28-7.22 (1H, m), 7.04 (1H, d, J= 7.8 Hz), 6.90 (1H, dd, J= 9 Hz, J= 1.3 Hz), 6.80 (1H, 2.6 Hz), 6.66 (1H, dd, J= 9 and 2.6 Hz), 6.46 (1H, s), 4.53 (2H, s), 2.88 (3H, d, J= 5.1 Hz), 1.31 (9H, s).

30 **7.3.558 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926491)**

In like manner to the preparation of N4-(ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2-(3,4-ethylenedioxyphenyl)-N4-(2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine with methylamine hydrochloride gave N2-(3,4-ethylenedioxyphenyl)-5-

fluoro-N4-[2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.10 (s, 1H), 7.94 (d, 1H, J= 5.1 Hz), 7.59 (s, 2H), 7.44 (s, 1H), 6.96 (d, 1H, J= 2.4 Hz), 6.82 (d, 1H, J= 8.4 Hz), 6.76 (dd, 1H, J= 3.6 and 8.1 Hz), 4.22 (s, 2H), 4.21 (s, 2H), 2.95 (s, 3H); LCMS: ret. time: 17.76 min.; purity: 97%; MS (m/e): 436 (MH⁺).

5 **7.3.559 N2-(3,5-Dimethoxyphenyl)-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine (R926810)**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2-
10 (3,5-dimethoxyphenyl)-N4-(3-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with methylamine hydrochloride gave N2-(3,5-dimethoxyphenyl)-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.93 (d, 1H, J= 3.9 Hz), 7.72 (t, 1H, J= 1.8 Hz), 7.27-7.19 9m, 2H), 6.88 (d, 2H, J= 2.4 Hz), 6.72 (m, 1H), 6.01 (t, 1H, J= 2.4 Hz), 4.44 (s, 2H), 3.67 (s, 6H), 2.80 (s,
15 3H).

7.3.560 5-Bromo-N2-(3,4-ethylenedioxyphenyl)-N4-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R925851)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-bromo-N2-(3,4-ethylenedioxyphenyl)-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and methylamine hydrochloride were reacted to yield 5-bromo-N2-(3,4-ethylenedioxyphenyl)-N4-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.01 (s, 1H), 7.48 (d, 2H, J= 8.7 Hz), 7.09 (d,
20 1H, J= 3.0 Hz), 7.08 (d, 2H, J= 8.7 Hz), 6.81 (dd, 1H, J= 8.7 Hz), 6.64 (d, 1H, J= 8.7 Hz), 4.52 (s, 2H), 4.20 (bs, 4H), 2.83 (s, 3H); LCMS: ret. time: 19.13 min.; purity: 94 %; MS (m/e): 487 (MH⁺).

30 **7.3.561 N2-(3-Hydroxyphenyl)-5-trifluoromethyl-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926741)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-(3-hydroxyphenyl)-5-trifluoromethyl-N4-(3-N-methoxycarbonylmethyleneoxyphenyl)-2,4-

pyrimidinediamine and methylamine hydrochloride were reacted to yield N2-(3-hydroxyphenyl)-5-trifluoromethyl-N4-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 18.52 min.; purity: 96%; MS (m/e): 434 (MH⁺).

7.3.562 N2,N4-Bis[4-(N-n-butylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine (R925860)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-cyano-2,4-pyrimidinediamine and n-butylamine were reacted to yield N2,N4-bis[4-(N-n-butylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.77 (bs, 1H), 9.38 (bs, 1H), 8.42 (s, 1H), 8.09 (t, 1H, J= 5.4 Hz), 8.02 (t, 1H, J= 5.7 Hz), 7.48-7.34 (m, 4H), 6.93 (d, 2H, J= 9.3 Hz), 6.82-6.72 (m, 2H), 4.47 (s, 2H), 4.38 (s, 2H), 3.14-3.06 (m, 4H), 1.42-1.33 (m, 4H), 1.28-1.18 (m, 4H), 0.83 (t, 6H, J= 6.9 Hz); LCMS: ret. time: 26.40 min.; purity: 97 %; MS (m/e): 546 (MH⁺).

7.3.563 N2,N4-Bis[4-(N-isopropylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine (R925861)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-cyano-2,4-pyrimidinediamine and isopropylamine were reacted to yield N2,N4-bis[4-(N-isopropylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.41 (s, 1H), 7.90 (d, 1H, J= 7.5 Hz), 7.81 (d, 1H, J= 7.5 Hz), 7.50-7.36 (m, 4H), 6.93 (d, 2H, J= 8.7 Hz), 6.84-6.75 (m, 2H), 4.45 (s, 2H), 4.36 (s, 2H), 3.99-3.87 (m, 2H), 1.08 (d, 6H, J= 3.0 Hz), 1.06 (d, 6H, J= 2.4 Hz); LCMS: ret. time: 23.45 min.; purity: 89 %; MS (m/e): 518 (MH⁺).

7.3.564 N2,N4-Bis[4-(N-n-propylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine (R925853)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-cyano-2,4-pyrimidinediamine and n-propylamine were reacted to yield N2,N4-bis[4-(N-n-propylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.78 (bs, 1H), 9.38 (bs, 1H), 8.41

(s, 1H), 8.07 (dt, 2H, J= 6.0 and 22.5 Hz), 7.48-7.36 (m, 4H), 6.93 (d, 2H, J= 8.7 Hz), 6.78 (d, 2H, J= 8.1 Hz), 4.48 (s, 2H), 4.39 (s, 2H), 3.07 (2q, 4H, J= 7.2 Hz), 1.47-1.38 (m, 4H), 0.90-0.77 (m, 6H); LCMS: ret. time: 23.67 min.; purity: 94 %; MS (m/e): 519 (MH⁺).

7.3.565 N2,N4-Bis[4-(N-morpholino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine (R925854)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-cyano-2,4-pyrimidinediamine and morpholine were reacted to yield N2,N4-bis[4-(N-morpholino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.78 (bs, 1H), 9.31 (bs, 1H), 8.41 (s, 1H), 7.43 (d, 4H, J= 8.1 Hz), 6.89 (d, 2H, J= 9.3 Hz), 6.75 (d, 2H, J= 8.4 Hz), 4.84 (s, 2H), 4.74 (s, 2H), 3.76 (t, 4H, J= 5.1 Hz), 3.62-3.50 (m, 4H), 3.49-3.38 (m, 4H), 3.08-3.01 (m, 4H); LCMS: ret. time: 19.25 min.; purity: 89 %; MS (m/e): 574 MH⁺.

7.3.566 N2,N4-Bis[4-(N-piperidino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine (R925855)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-cyano-2,4-pyrimidinediamine and piperidine were reacted to yield N2,N4-bis[4-(N-piperidino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (acetone-d₆): δ 8.86 (bs, 1H), 8.48 (bs, 1H), 8.34 (s, 1H), 7.61-7.50 (m, 4H), 6.98 (d, 2H, J= 8.7 Hz), 6.90 (d, 2H, J= 9.3 Hz), 4.84 (s, 2H), 4.75 (s, 2H), 3.59-3.48 (m, 8H), 1.68-1.44 (m, 12H); LCMS: ret. time: 24.76 min.; purity: 98 %; MS (m/e): 571 (MH⁺).

7.3.567 N2,N4-Bis[4-(N-cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine (R925859)

In a manner similar to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-cyano-2,4-pyrimidinediamine and cyclopropylmethylamine were reacted to yield N2,N4-bis[4-(N-cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-5-cyano-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.78 (bs, 1H), 9.36 (bs, 1H), 8.41 (s, 1H), 8.18 (t, 1H, J= 5.1 Hz),

8.10 (t, 1H, J= 5.1 Hz), 7.52-7.38 (m, 4H), 6.94 (d, 2H, J= 8.7 Hz), 6.84-6.76 (m, 2H), 4.48 (s, 2H), 4.40 (s, 2H), 3.00 (q, 4H, J= 6.3 Hz), 0.97-0.88 (m, 2H), 0.40-0.33 (m, 4H), 0.18-0.03 (m, 4H); ¹⁹F NMR (CDCl₃): LCMS: ret. time: 24.58 min.; purity: 100 %; MS (m/e): 543 (MH⁺).

5 **7.3.568 N4-(3-Aminophenyl)-N2-(1,4-benzoxazin-6-yl)-5-fluoro-2,4-pyrimidinediamine (R950254)**

N4-(3-Nitrophenyl)-N2-[(2H)1,4-benzoxazin-3(4H)-one-6-yl]-5-fluoro-2,4-pyrimidinediamine (940 mg, 2.5 mmol) and Pd/C 10% (300 mg, 50% water content) were suspended in EtOH (7 mL) and 10% aqueous HCl (5 mL) and hydrogenated in a Parr
10 apparatus for 3 hours (22 °C, 60 psi). The suspension was filtered over celite and neutralized by addition of K₂CO₃. The solvents were removed and the resulting black slurry was suspended in MeOH. Silica gel (4 g) was added and the volatiles were removed under reduced pressure. The residue was subjected to column chromatography on silica gel (CHCl₃-Acetone, 2:1) to give 186 mg of N4-(3-aminophenyl)-N2-(1,4-benzoxazin-6-yl)-5-
15 fluoro-2,4-pyrimidinediamine as brownish solid. ¹H NMR (DMSO-d₆): δ 8.92 (s, 1H), 8.64 (s, 1H), 7.95 (d, 1H, J= 3.6 Hz), 7.11 (s, 1H), 6.84-6.95 (m, 3H), 6.66 (dd, 1H, J= 2.4, 9.0 Hz), 6.46 (d, 1H, J= 8.1 Hz), 6.28 (d, 1H, J= 8.1 Hz), 5.62 (s, 1H), 4.98 (s, 2H), 4.03 (m, 2H), 3.31 (m, 2H); LCMS purity: 98.4%; MS (m/e): 352.7 (M⁺, 100).

20 **7.3.569 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-2-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950200)**

N2-(3-Ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (50 mg, 0.11 mmol) was dissolved in EtOH:4-(2-aminoethyl)morpholine (0.5 ml : 0.5 ml) and the mixture was refluxed for 3 hours (100 °C
25 oil-bath temperature). The mixture was cooled to 22 °C, diluted with water and washed with EtOAc. The organic phase was dried over MgSO₄, concentrated under reduced pressure, and the residue was subjected to column chromatography on silica gel (CHCl₃:Acetone, 2:1) to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-2-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆ + CD₃OD):
30 δ 7.92 (d, 1H, J= 4.1 Hz), 7.31 (d, 1H, J= 2.3 Hz), 7.20 (dd, 1H, J= 2.7, 8.8 Hz), 6.87-6.99 (m, 2H), 6.74 (d, 1H, J= 8.8 Hz), 6.09 (m, 1H), 4.19 (m, 4H), 3.38 (m, 4H), 3.16 (t, 2H, J= 6.3 Hz), 2.28 (t, 2H, J= 6.3 Hz); LCMS purity: 99.2%; MS (m/e): 524.01 (M⁺, 100).

7.3.570 N4-(3,4-Ethylenedioxyphenyl)-N2-[3-N-methylamino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950191)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and methylamine were reacted to prepare N4-(3,4-ethylenedioxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 17.32 min.; purity: 99.3%; MS (m/e): 425.04 (MH⁺).

7.3.571 N2-[3-(N-Amino)carbonylmethyleneaminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950192)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and aqueous ammonia were reacted to prepare N2-[3-(N-amino)carbonylmethyleneaminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 16.59 min.; purity: 98.8%; MS (m/e): 411.02 (MH⁺).

7.3.572 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950193)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and morpholine were reacted to prepare N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 18.70 min.; purity: 85.8%; MS (m/e): 481.05 (MH⁺).

7.3.573 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methyl)piperazino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950194)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine,

N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and N-methylpiperazine were reacted to prepare N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-(N-methyl)piperazino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 15.75 min.; purity: 99.1%; MS (m/e): 494.06 (MH⁺).

7.3.574 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-2-hydroxyethyleneamino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950195)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and 2-aminoethanol were reacted to prepare N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-2-hydroxyethyleneamino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 16.23 min.; purity: 97.3%; MS (m/e): 455.02 (MH⁺).

7.3.575 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)ethyleneaminocarbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950196)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and N-methyl-ethylen-1,2-diamine were reacted to prepare N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)ethyleneaminocarbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 15.34 min.; purity: 98.2%; MS (m/e): 468.06 (MH⁺).

7.3.576 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950197)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and piperazine were reacted to prepare N4-(3,4-ethylenedioxyphenyl)-5-

fluoro-N2-[3-(N-piperazino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 15.38 min.; purity: 93.2%; MS (m/e): 479.99 (MH⁺).

7.3.577 N2-[3-(N-Benzylamino)ethyleneaminocarbonylmethyleneaminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950198)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and N-benzyl-ethylen-1,2-diamine were reacted to prepare N2-[3-(N-benzylamino)ethyleneaminocarbonylmethyleneaminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 17.70 min.; purity: 92.5%; MS (m/e): 544.04 (MH⁺).

7.3.578 N2-[3-(N,N'-Bis(2-N-hydroxyethyl)amino)carbonylmethyleneaminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950199)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and N,N'-bis(2-hydroxyethylene)amine were reacted to N2-[3-(N,N'-bis(2-N-hydroxyethyl)amino)carbonylmethyleneaminophenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 15.81 min.; purity: 99.4%; MS (m/e): 499.01 (MH⁺).

7.3.579 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950217)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and methylamine were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 14.41 min.; purity: 93.0%; MS (m/e): 383.02 (MH⁺).

7.3.580 N2-(3-Aminocarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950219)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and aqueous ammonia were reacted to prepare N2-(3-aminocarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 14.23 min.; purity: 95.0%; MS (m/e): 369.03 (MH⁺).

7.3.581 N2-[3-(N,N-Dimethylamino)carbonylmethyleneaminophenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R950220)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and dimethylamine were reacted to prepare N2-[3-(N,N-dimethylamino)carbonylmethyleneaminophenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 16.59 min.; purity: 96.5%; MS (m/e): 397.06 (MH⁺).

7.3.582 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-morpholino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950221)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and morpholine were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-morpholino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 16.29 min.; purity: 91.5%; MS (m/e): 439.03 (MH⁺).

7.3.583 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-piperazino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950222)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and piperazine were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-

N2-[3-(N-piperazino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 13.04 min.; purity: 89.9%; MS (m/e): 438.06 (MH⁺).

7.3.584 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-[N-(N-methyl)piperazino]carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950223)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and N-methylpiperazine were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-[N-(N-methyl)piperazino]carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.06 min.; purity: 98.7%; MS (m/e): 452.06 (MH⁺).

7.3.585 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-2-hydroxyethylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950224)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and 2-aminoethanol were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-2-hydroxyethylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 13.28 min.; purity: 97.3%; MS (m/e): 413.04 (MH⁺).

7.3.586 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-methylamino)ethylamino]carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950225)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and N-methyl-ethylen-1,2-diamine were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-methylamino)ethylamino]carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.31 min.; purity: 94.7%; MS (m/e): 426.01 (MH⁺).

7.3.587 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-2-morpholinoethylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950226)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and N-morpholinylethylamine were reacted to prepare 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-2-morpholinoethylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.66 min.; MS (*m/e*): 482.39 (MH⁺).

7.3.588 R935184: 5-Fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine was reacted with Me₂NH.HCl and *i*-Pr₂NEt in methanol to produce 5-fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 6.91 min.; purity: 98%; MS (*m/e*): 440 (MH⁺).

7.3.589 R935196: N2-[3-(1-Bis(N-methylaminocarbonyl)ethoxy)phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidineamine:

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-[3-(1-bis(ethyloxycarbonyl)ethoxy)phenyl]-5-fluoro-N2-[4-isopropoxyphenyl]-2,4-pyrimidinediamine was reacted with Me₂NH.HCl and *i*-Pr₂NEt in presence of methanol to produce N2-[3-(1-bis(N-methylaminocarbonyl)ethoxy)phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidineamine. ¹H NMR (DMSO-d₆): δ 9.18 (s, 1H), 9.15 (s, 1H), 8.07 (app qt, 2H, J= 4.7 Hz), 8.01 (d, 1H, J= 3.5 Hz), 7.65-7.62 (m 2H), 7.36 (br s, 1H), 7.28 (dd, 1H, J= 1.1 and 8.2 Hz), 7.03 (t, 1H, J= 8.2 Hz), 6.87 (d, 2H, J= 8.8 Hz), 6.35 (dd, 1H, J= 1.1 and 8.8 Hz), 4.54 (q, 1H, J= 6.4 Hz), 2.62 (d, 6H, J= 4.7 Hz), 1.49 (s, 3H), 1.23 (d, 6H, J= 5.8 Hz). LCMS: ret. time: 19.40 min.; purity: 94%; MS (*m/e*): 497 (MH⁺).

7.3.590 R935202: 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine:

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine was reacted with Me₂NH.HCl to give 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.21 (s, 1H), 9.19 (s, 1H), 8.06 (d, 1H, J= 4.1 Hz), 7.94 (q, 1H, J= 3.5 Hz), 7.42-7.38 (m, 2H), 7.30 (d, 2H, J= 7.6 Hz), 7.12 (t, 1H, J= 7.6 Hz), 6.89 (d, 1H, J= 8.2 Hz), 6.47 (dd, 1H, J= 2.3 and 8.8 Hz), 4.33 (s, 2H), 4.11-4.03 (m, 4H), 2.63 (d, 3H, J= 4.7 Hz), 2.08-2.03 (m, 2H). LCMS: ret. time: 17.33 min.; purity: 98%; MS (*m/e*): 440 (MH⁺).

7.3.591 R935206: N2, N4-Bis[1-(N-methylaminocarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2, N4-Bis[1-(methoxycarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine and was reacted with Me₂NH.HCl and *i*-PrN₂Et in presence of methanol to produce N2, N4-bis[1-(N-methylaminocarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.56 (s, 1H), 9.40 (s, 1H), 8.17 (d, 1H, J= 3.5 Hz), 8.12 (s, 1H), 7.99 (s, 1H), 7.96 (s, 2H), 7.90 (s, 2H), 7.66 (d, 1H, J= 8.8 Hz), 7.56 (d, 1H, J= 8.8 Hz), 7.49 (dd, 1H, J= 1.7 and 8.8 Hz), 7.34 (dd, 1H, J= 1.7 and 8.8 Hz), 4.90 (s, 2H), 4.66 (s, 2H), 2.56 (d, 6H, J= 4.11 Hz). LCMS: ret. time: 13.85 min.; purity: 98%; MS (*m/e*): 503 (MH⁺).

7.3.592 R935212: N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(N-methylaminocarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine and Me₂NH.HCl was reacted to produce N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(N-methylaminocarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.35 (s, 1H), 9.17 (s, 1H), 8.07 (d, 1H, J= 4.8 Hz), 7.92 (s, 1H),

7.89 (s, 1H), 7.66 (q, 1H, J= 4.7 Hz), 7.54 (d, 1H, J= 8.8 Hz), 7.35-7.24 (m, 3H), 6.76 (d, 1H, J= 8.8 Hz), 4.77 (s, 2H), 4.20 (s, 4H), 2.57 (d, 3H, J= 4.7 Hz). LCMS: ret. time: 15.82 min.; purity: 94%; MS (*m/e*): 450 (MH⁺).

5

7.3.593 R935213: N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-(N-methylamino)carbonyl-fur-4-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-methoxycarbonyl-fur-4-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine was reacted with Me₂NH.HCl and *i*-Pr₂NEt. to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-(N-methylamino)carbonyl-fur-4-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.17 (s, 2H), 8.30 (q, 1H, J= 4.7 Hz), 8.05 (d, 1H, J= 3.5 Hz), 7.42 (s, 1H), 7.29-7.19 (m, 2H), 7.09 (t, 1H, J= 8.2 Hz), 7.02 (d, 1H, J= 2.9 Hz), 6.76 (d, 1H, J= 8.8 Hz), 6.67 (d, 1H, J= 2.9 Hz), 6.54 (dd, 1H, J= 1.7 and 8.2 Hz), 4.94 (s, 2H), 4.21-4.18 (m, 4H), 2.70 (d, 3H, J= 4.7 Hz). LCMS: ret. time: 18.85 min.; purity: 91%; MS (*m/e*): 492 (MH⁺).

10

15

7.3.594 R935216: 5-Fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N2-[4-(N-methylamino)carbonylmethyleneoxy)phenyl]-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.31 (s, 1H), 9.00 (s, 1H), 8.17 (s, 1H), 8.02 (d, 1H, J= 3.5 Hz), 7.99 (m, 1H), 7.93 (s, 1H), 7.59 (m, 2H), 7.52 (d, 2H, J= 8.8 Hz), 6.78 (d, 2H, J= 8.8 Hz), 4.36 (s, 2H), 4.03 (s, 3H), 2.63 (d, 3H, J= 4.7 Hz). LCMS: ret. time: 14.81 min.; purity: 99%; MS (*m/e*): 422 (MH⁺).

20

25

30

7.3.595 R935217: N2, N4-Bis[1-(N-methylaminocarbonyl)methyl-indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2, N4-bis[1-(methoxycarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine and

Me₂NH.HCl were reacted to produce N2, N4-bis[1-(N-methylaminocarbonyl)methyl-indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.35 (s, 1H), 9.15 (s, 1H), 8.09-8.06 (m, 2H), 7.97-7.96 (m, 2H), 7.91 (s, 1H), 7.70 (s, 1H), 7.69 (s, 1H), 7.64-7.55 (m, 2H), 7.48-7.40 (m, 2H), 5.06 (s, 2H), 4.97 (s, 2H), 2.62 (d, 3H, J= 4.7 Hz), 2.61 (d, 3H, J= 4.7 Hz). LCMS: ret. time: 12.54 min.; purity: 95%; MS (*m/e*): 503 (MH⁺).

7.3.596 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926486)

A dry reaction vial equipped with a rubber septum was charged with N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (0.019 g, 0.04 mmol) and THF (1 mL). To this was added boranemethyl sulfide complex (0.044 mL, 0.088 mmol) and stirred at room temperature for 2h. The amount of boranemethyl sulfide complex was evaporated and the reaction was quenched with MeOH (CAUTION: vigorous evolution of hydrogen gas occurs during the addition of MeOH), heated for 30 min. The solvent was removed and again the residue was suspended in MeOH, extracted with EtOAc, EtOAc was evaporated and the residue was purified by preparative TLC to obtain N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.20 (s, 1H), 8.01 (d, 1H, J= 6 Hz), 7.26-7.05 (m, 3H), 7.05-6.97 (m, 3H), 6.82 (d, 1H, J= 9.3 Hz), 6.67 (dd, 1H, J= 1.8 and 8.1 Hz), 4.44 (t, 2H), 4.27 (s, 4H), 4.14 (m, 2H), 3.76 (m, 2H), 3.22 (t, 2H, J= 5.4 Hz), 3.05 (m, 2H), 2.88 (m, 2H).

7.3.597 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-morpholinomethylene)benzofuran-5-yl]-2,4-pyrimidinediamine (R926490)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-morpholino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine with boranemethyl sulfide complex gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-morpholinomethylene)benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.65 (d, 2H, J= 2.1 Hz), 8.30 (dd, 2H, J= 2.1 and 9.6 Hz), 7.73 (d, 2H, J= 9.3 Hz), 7.49 (bs, 2H), 7.32 (m, 1H), 6.74 (m, 1H), 4.24 (s, 4H), 3.97 (s, 2H), 3.78 (m, 4H), 3.56 (m, 4H).

7.3.598 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-methylamino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926510)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and boranemethyl sulfide complex gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-methylamino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.00 (d, 1H, J= 5.2 Hz), 7.50-7.30 (m, 2H), 7.16- 6.80 (m, 5H), 4.28 (m, 1H), 4.27 (bs, 4H), 4.22 (m, 1H), 3.44 (m, 2H), 2.79 (d, 3H, J= 3Hz); LCMS: ret. time: 15.64 min.; purity: 96%; MS (m/e): 412 (MH⁺).

7.3.599 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazinomethylene)benzofuran-5-yl]-2,4-pyrimidinediamine (R926770)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine with boranemethyl sulfide complex gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazinomethylene)benzofuran-5-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.06 min.; purity: 75%; MS (m/e): 435 (MH⁺).

7.3.600 N4-(3,5-Dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine (R940255)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine with boranemethyl sulfide complex gave N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 15.94 min.; purity: 99 %; MS (m/e): 454 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.16 (1H, s), 9.07 (1H, s), 8.15 (1H, s), 8.11 (1H, d, J= 3.9 Hz), 7.40-7.30 (4H, m), 7.13 (1H, t, 8.1 Hz), 6.55 (1H, dd, J= 8.1 Hz, 3.2 Hz), 4.01 (2H, t, J= 5.7 Hz), 3.65 (4H, t, J= 4.2 Hz), 2.72 (2H, t, J= 5.7 Hz), 2.515 (4H, t, J= 4.5 Hz), 2.24 (6H, s).

7.3.601 N4-(3,5-Dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethoxy]phenyl]-2,4-pyrimidinediamine bis Hydrogen Chloride Salt (R945142)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine was treated with boranemethyl sulfide complex to give N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethoxy]phenyl]-2,4-pyrimidinediamine, which was then treated with 4N HCl in dioxane (3 mL) followed by crystallization from MeOH/EtOAc to give N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethoxy]phenyl]-2,4-pyrimidinediamine bis Hydrogen Chloride Salt. ¹H NMR (CD₃OD): δ 2.17 (s, 6H), 3.66 (m, 10H), 4.26 (t, J= 4.5 Hz, 2H), 6.93 (dd, J= 1.5, 7.2 Hz, 1H), 7.10-7.13 (m, 2H), 7.17 (s, 2H), 7.31 (t, J= 8.4 Hz, 1H), 7.98 (d, J= 6.0 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 162.93; LCMS: ret. time: 13.25 min.; purity: 96.08%; MS (m/e): 453.09 (MH⁺).

7.3.602 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine (R945144)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, the reaction of N2-(4-carboxymethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and boranemethyl sulfide complex gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (acetone-*d*₆): δ 3.86 (t, J= 4.8 Hz, 2H), 4.04 (t, J= 4.8 Hz, 2H), 4.28 (m, 4H), 6.78 (d, J= 9.0 Hz, 1H), 6.86 (d, J= 9.0 Hz, 2H), 7.18 (dd, J= 2.7, 8.7 Hz, 1H), 7.47 (d, J= 2.7 Hz, 1H), 7.63 (d, J= 9.0 Hz, 2H), 7.91 (d, J= 3.6 Hz, 1H), 8.29 (br, 1H, NH), 8.31 (br, 1H, NH); ¹⁹F NMR (282 MHz, acetone-*d*₆): δ - 169.18; LCMS: ret. time: 17.41 min.; purity: 98.36%; MS (m/e): 399.01 (MH⁺).

7.3.603 N4-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethoxy]phenyl]-2,4-pyrimidinediamine Dihydrochloride Salt (R945150)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-

pyrimidinediamine was treated with boranemethyl sulfide complex to give N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethyloxy]phenyl]-2,4-pyrimidinediamine, which was then treated with 4N HCl in dioxane (3 mL) followed crystallization from MeOH/EtOAc to give N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethoxy]phenyl]-2,4-pyrimidinediamine bis Hydrogen Chloride Salt. ¹H NMR (CD₃OD): δ 2.21 (s, 3H), 3.72 (m, 10H), 4.35 (t, J= 4.5 Hz, 2H), 6.95 (dt, J= 1.5 and 9.0 Hz, 1H), 7.11-7.14 (m, 2H), 7.26 (dd, J= 0.9 and 2.7 Hz, 1H), 7.34 (t, J= 8.4 Hz, 1H), 7.50 (d, J= 2.4 Hz, 1H), 8.03 (d, J= 5.4 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 162.74; LCMS: ret. time: 14.50 min.; purity: 94.75%; MS (m/e): 472.98 (MH⁺).

7.3.604 N4-(3,5-Dimethyl-4-methoxyphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethoxy]phenyl]-2,4-pyrimidinediamine Dihydrochloride Salt (R945157)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine was treated with boranemethyl sulfide complex to give N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethyloxy]phenyl]-2,4-pyrimidinediamine, which was then treated with 4N HCl in dioxane (3 mL) followed crystallization from MeOH/EtOAc to give N4-(3,5-dimethyl-4-methoxyphenyl)-5-fluoro-N2-[3-[2-(N-piperazino)ethoxy]phenyl]-2,4-pyrimidinediamine bis Hydrogen Chloride Salt. ¹H NMR (CD₃OD): δ 2.23 (s, 6H), 3.66 (m, 10H), 3.72 (s, 3H), 4.31 (t, J= 4.5 Hz, 2H), 6.95 (dd, J= 1.8 and 8.4 Hz, 1H), 7.09-7.15 (m, 2H), 7.27 (s, 2H), 7.32 (t, J= 8.1 Hz, 1H), 8.01 (d, J= 5.4 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 162.71; LCMS: ret. time: 16.41 min.; purity: 97.50%; MS (m/e): 467.12 (MH⁺).

7.3.605 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926501)

The reaction of equivalent amount of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine with hydrogen chloride (4M, dioxane) in methanol at 0 °C followed by dilution with dry ethyl ether or ethyl acetate gave the precipitate. The resulting precipitate was isolated by filtration (and/or using centrifuse technique) to give N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-

piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. ¹H NMR (CD₃OD): δ 7.97 (d, 1H, J= 5.4 Hz), 7.92 (d, 1H, J= 1.8 Hz), 7.62 (d, 1H, J= 8.2 Hz), 7.48 (s, 1H), 7.43 (dd, 1H, J= J= 2.4 and 8.7 Hz), 7.17 (d, 1H, J=2.4 Hz), 6.98 (dd, 1H, J= 2.4 and 8.7 Hz), 6.77 (d, 1H, J= 8.7 Hz), 4.13 (m, 4H), 4.22 (s, 4H), 3.38 (t, 4H, J= 5.7 Hz); LCMS: ret. time: 15.12 min; purity: 89%; MS (m/e): 491 (MH⁺).

7.3.606 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926504)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydrogen chloride gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. ¹H NMR (DMSO-d₆): δ 9.6 (bs, 1H), 9.04 (bs, 1H), 8.12 (d, 1H, J= 3.6 Hz), 7.25-7.00 (m, 5H), 7.81 (d, 1H, J= 8.7 Hz), 6.54 (d, 1H, J= 8.4 Hz), 4.74 (s, 2H), 4.22 (s, 4H), 3.64 (m, 4H), 3.11 (m, 4H); LCMS: ret. time: 15.34 min.; purity: 100%; MS (m/e): 481 (MH⁺).

7.3.607 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-N-methylaminoethyl)phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926509)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-methylamino)ethyloxy]phenyl]-2,4-pyrimidinediamine with hydrogen chloride (4M, dioxane) gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-methylamino)ethyloxy]phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. LCMS: ret. time: 15.88 min.; purity: 92%; MS (m/e): 412 (MH⁺).

7.3.608 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926511)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-

morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine and hydrogen chloride gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. ¹H NMR (CD₃OD): δ 7.98 (d, 1H, J= 5.4 Hz), 7.34 (t, 1H, 8.4 Hz), 7.16-6.81 (m, 6H), 4.42 (m, 1H), 4.40 (m, 2H), 4.25 (m, 5H), 4.10 (m, 2H), 3.90 (bs, 2H), 3.60 (m, 4H); LCMS: ret. time: 16.39 min.; purity: 100%; MS (m/e): 468 (MH⁺).

7.3.609 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-homopiperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926768)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-homopiperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine with hydrogen chloride treatment gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-homopiperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. ¹H NMR (DMSO-d₆): δ 9.98 (bs, 1H), 9.05 (bs, 1H), 8.18 (d, 1H, J= 4.8 Hz), 8.01 (s, 1H), 7.58 (d, 1H, J= 8.7 Hz), 7.50 (bd, 1H), 7.35 (s, 1H), 7.24 (d, 1H, J= 2.4 Hz), 7.11 (dd, 1H, J= 3 and 9 Hz), 6.80 (d, 1H, J= 8.7 Hz), 4.22 (s, 4H), 4.20-3.60 (m, 8H), 3.20 (m, 2H); LCMS: ret. time: 14.91 min.; purity: 86%; MS (m/e): 505 (MH⁺).

7.3.610 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt R926502)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine upon treatment with hydrogen chloride (4M, dioxane) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. ¹H NMR (CDCl₃OD): δ 8.00 (s, 1H), 7.89 (s, 1H), 7.98 (s, 1H), 7.60 (d, 1H, J= 8.7 Hz), 7.45 (m, 3H), 7.16 (t, 1H, J= 8.1 Hz), 7.10 (m, 1H), 7.02 (dd, 1H, J= 1.2 and 7.2 Hz), 6.70 (dd, 1H, J= 2.4 and 8.4 Hz), 4.13 (m, 4H), 3.37 (t, 4H, J= 5.4 Hz), 3.38 (t, 4H, J= 5.7 Hz); LCMS: ret. time: 13.40 min; purity: 79%; MS (m/e): 450 (MH⁺).

7.3.611 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazinomethylene)benzofuran-5-yl]-2,4-pyrimidinediamine Dihydrochloride Salt (R926769)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazinomethylene)benzofuran-5-yl]-2,4-pyrimidinediamine with hydrogen chloride (4M, dioxane) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(N-piperazinomethylene)benzofuran-5-yl]-2,4-pyrimidinediamine Dihydrochloride Salt. ¹H NMR (CD₃OD): δ 8.00 (d, 1H), 7.85 (bd, 1H), 7.75 (m, 3H), 7.60 (m, 2H), 7.40-7.15 (m, 4H), 7.05 (s, 1H), 7.00-6.800 (m, 3H), 4.65 (dd, 2H), 3.60 (m, 8H).

7.3.612 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926773)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydrogen chloride (4M, dioxane) gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-piperazino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. ¹H NMR (CD₃OD): δ 7.99 (d, 1H, J= 5.1 Hz), 7.29 (t, 1H, J= 8.1 Hz), 7.21-7.05 (m, 5H), 6.83 (dd, 1H, J= 2.4 and 8.7 Hz), 6.77 (bd, 1H), 4.79 (s, 2H), 3.83 (m, 2H), 3.78 (m, 2H), 3.25 (m, 2H); LCMS: ret. time: 12.27 min.; purity: 91%; MS (m/e): 439 (MH⁺).

7.3.613 N2-[3-[2-(N, N-Dimethylamino)ethyloxy]phenyl]-N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926771)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the treatment of N4-(3,4-ethylenedioxyphenyl)-N2-[3-[2-(N, N-dimethylamino)ethyloxy]phenyl]-5-fluoro-2,4-pyrimidinediamine with equivalent amount of hydrogen chloride (4M, dioxane) gave N4-(3,4-ethylenedioxyphenyl)-N2-[3-[2-(N, N-dimethylamino)ethyloxy]phenyl]-5-fluoro-2,4-pyrimidinediamine Hydrogen Chloride Salt. LCMS: ret. time: 15.37 min.; purity: 93%; MS (m/e): 426 (MH⁺).

7.3.614 N4-(3,5-Dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R940256)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the reaction of N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine with hydrogen chloride (4M, dioxane) gave N4-(3,5-dimethyl-4-hydroxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. LCMS: ret. time: 15.78 min.; purity: 98 %; MS (M/e): 454 (MH⁺); ¹H NMR (DMSO-d₆): δ 10.60 (1H, s), 9.58 (1H, s), 8.29 (1H, s), 8.20 (1H, s), 7.43 (1H, d, J= 9Hz), 7.38-7.30 (3H, m), 7.24 (1H, t, J= 9 Hz), 6.70 (1H, d, J= 9 Hz), 4.35 (2H, m), 4.05 (2H, m), 3.84 (4H, m), 3.65-3.50 (2H, m), 3.26 (2H, m), 2.25 (6H, s).

7.3.615 N4-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R940269)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the reaction of N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine with hydrogen chloride (4M, dioxane) gave N4-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyloxy]phenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt. LCMS: ret. time: 14.74 min.; purity: 96 %; MS (m/e): 474 (M⁺), 475 (MH⁺); ¹H NMR (DMSO-d₆): δ 10.03 (1H, s), 9.35 (2H, s), 9.06 (1H, s), 8.17 (1H, d, J= 3.9 Hz), 7.67 (1H, m), 7.52 (1H, m), 7.46 (1H, d, J= 8.7 Hz), 7.39 (1H, s), 7.24 (1H, t, J= 8.1 Hz), 6.66 (1H, d, J= 8.1 Hz), 4.33 (1H, m), 4.07 (1H, d, J= 13 Hz), 3.79 (1H, t, J= 12.5 Hz), 3.56 (4H, m), 3.49 (4H, m), 3.29 (1H, t, J= 12.5 Hz), 2.29 (3H, s).

7.3.616 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R926816)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(N-piperazino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine Hydrogen Chloride Salt, the treatment of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine with equivalent

amount of hydrogen chloride (4M, dioxane) gave the N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride salt. LCMS: ret. time: 17.04 min., purity: 96%, MS (m/e): 426 (MH⁺).

7.3.617 N4-(3,4-Ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine (R926696)

A dry reaction flask charged with N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine was reacted with diisobutylaluminum hydride (DIBALH) (5 equivalents) in CH₂Cl₂ at -78 °C (reaction was monitored by TLC) followed by treatment with Rochell's salt to yield N4-(3,4-ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.11 (s, 1H), 8.02 (d, 1H, J= 3.3 Hz), 7.96 (t, 1H, J= 1.8 Hz), 7.40-7.30 (m, 3H), 7.19 (dt, 1H, J= 3.6 and 8.1 Hz), 6.78 (d, 1H, J= 8.7 Hz), 6.59 (s, 1H), 4.52 (d, 2H, J= 5.1 Hz), 4.22 (s, 4H); ¹⁹F NMR (DMSO-d₆): - 46802; LCMS: ret. time: 19.14 min.; purity: 95 %; MS (m/e): 409 (MH⁺).

7.3.618 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(hydroxymethyl)-(1H)-indol-5-yl]-2,4-pyrimidinediamine (R926700)

In a manner similar to the preparation of N4-(3,4-ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(methoxycarbonyl)-(1H)-indol-5-yl]-2,4-pyrimidinediamine was reduced with DIBALH to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(hydroxymethyl)-(1H)-indol-5-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.81 (d, 1H, J= 4.2 Hz), 7.23 (d, 1H, J= 1.8 Hz), 7.28-7.23 (m, 2H), 7.19 (t, 1H, J= 2.4 Hz), 7.12 (dd, 1H, J= 1.8 and 9.0 Hz), 7.07 (t, 1H, J= 8.4 Hz), 6.52 (ddd, 1H, J= 1.2 and 8.1 Hz), 6.30 (s, 1H), 4.71 (s, 2H); ¹⁹F NMR (CD₃OD): - 47971; LCMS: ret. time: 15.36 min.; purity: 100 %; MS (m/e): 366 (MH⁺).

7.3.619 5-Fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-N4-[4-(isopropoxy)phenyl]-2,4-pyrimidinediamine (R926705)

In a manner similar to the preparation of N4-(3,4-ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine, 5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to yield 5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.83 (d, 1H, J= 3.3 Hz),

7.81 (s, 1H), 7.50 (d, 2H, J= 9.0 Hz), 7.29 (d, 1H, J= 9.0 Hz), 7.22 (dd, 1H, J= 2.4 and 8.7 Hz), 6.84 (d, 2H, J= 8.7 Hz), 6.56 (d, 1H, J= 1.2 Hz), 4.64 (s, 2H), 4.56 (2q, 1H, J= 5.7 Hz), 1.31 (d, 6H, J= 6.0 Hz); ^{19}F NMR (CD_3OD): - 47926; LCMS: ret. time: 21.03 min.; purity: 99 %; MS (m/e): 409 (MH^+).

5 **7.3.620 5-Fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926707)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine was reduced with
10 DIBALH to yield 5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ^1H NMR ($\text{DMSO}-d_6$): δ 9.37 (s, 1H), 9.17 (s, 1H), 9.12 (s, 1H), 8.06 (d, 1H, J= 3.9 Hz), 8.01 (d, 1H, J= 1.8 Hz), 7.41-7.35 (m, 2H), 7.26 (d, 1H, J= 8.1 Hz), 7.11-7.05 (m, 2H), 6.60 (s, 1H), 6.51 (dd, 1H, J= 2.4 and 8.4 Hz), 5.41 (t, 1H, J= 6.0 Hz), 4.51 (d, 2H, J= 5.7 Hz); LCMS: ret. time: 16.21 min.; purity: 95 %; MS (m/e): 367 (MH^+).

15 **7.3.621 N4-(4-*tert*-Butyl)phenyl)-5-fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-2,4-pyrimidinediamine (R926728)**

In a manner similar to the preparation of N4-(3,4-ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine, N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced
20 with DIBAL to yield N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.94 (d, 1H, J= 3.0 Hz), 7.54 (d, 2H, J= 9.0 Hz), 7.37 (d, 2H, J= 8.4 Hz), 7.29-7.35 (m, 1H), 7.19-7.14 (m, 2H), 7.06 (d, 1H, J= 8.1 Hz), 6.82 (d, 1H, J= 2.7 Hz), 6.57 (dd, 1H, J= 2.4 and 8.1 Hz), 4.04-4.00 (m, 2H), 3.93-
25 3.89 (m, 2H), 1.33 (s, 9H); ^{19}F NMR (CDCl_3): -47214; LCMS: ret. time: 22.39 min.; purity: 94 %; MS (m/e): 397 (MH^+).

7.3.622 5-(Hydroxymethyl)-N2-[3-(2-hydroxyethyleneoxy)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926735)

In a manner similar to the preparation of N4-(3,4-ethylenedioxy)-5-fluoro-N2-[2-(hydroxymethyl)benzofuran-5-yl]-2,4-pyrimidinediamine, N4-(3-hydroxyphenyl)-5-methoxycarbonyl-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine
30 was reduced with DIBALH to yield 5-(hydroxymethyl)-N2-[3-(2-

hydroxyethyleneoxy)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.87 (s, 1H), 7.35 (t, 1H, J= 1.5 Hz), 7.15-7.08 (m, 5H), 6.57-6.50 (m, 2H), 4.56 (s, 2H), 3.92-3.86 (m, 2H), 3.84-3.79 (m, 2H); LCMS: ret. time: 14.11 min.; purity: 89 %; MS (m/e): 369 (MH⁺).

5 **7.3.623 5-Fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine R940289**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-isopropylphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine
10 reacted with DIBALH to give 5-fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 23.03 min.; purity: 93 %; MS (m/e): 382 (M⁺), 384 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.36 (1H, s), 9.24 (1H, s), 8.20 (1H, d, J= 4.2 Hz), 7.85 (1H, d, J= 8.5 Hz), 7.57 (1H, s), 7.41 (1H, s), 7.33 (1H, t, J= 8.5 Hz), 7.17 (1H, t, J= 8.5 Hz), 7.05 (1H, d, J= 8.5 Hz), 6.56 (1H, dd, J= 8.5 Hz, J= 2 Hz) 4.94 (1H, t, J= 12 Hz), 3.94 (2H, t, J= 4.7 Hz), 3.76 (2H, m), 2.95 (1H, sept, J= 6.9 Hz), 1.28 (6H, dd, J= 6.9 Hz, J= 0.6Hz).

7.3.624 N4-(3-*tert*-Butylphenyl)-5-fluoro-N2-[(2-hydroxymethylene)benzofur-5-yl]-2,4-pyrimidinediamine R940287

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, N4-(3-*tert*-butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine reacted with
20 DIBALH to give N4-(3-*tert*-butylphenyl)-5-fluoro-N2-[2-(hydroxymethylene)benzofur-5-yl]-2,4-pyrimidinediamine. LCMS: retn, time: 23.15 min.; purity: 99 %; MS (m/e): 407 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.34 (1H, s), 9.22 (1H, s), 8.18 (1H, d, J= 3.9 Hz), 8.04 (1H, s), 8.00 (1H, d, J= 8.7 Hz), 7.60 (1H, t, J= 2.1 Hz), 7.47 (2H, m), 7.34 (1H, t, J= 7.8 Hz), 7.21 (1H, d, J= 8.7 Hz), 6.69 (1H, s), 5.54 (1H, t, J= 5.8 Hz), 4.63 (2H, d, J= 5.8 Hz), 1.35 (9H, s).

30 **7.3.625 5-Fluoro-N4-(3-isopropylphenyl)-N2-[(2-hydroxymethylene)benzofur-5-yl]-2,4-pyrimidinediamine R940286**

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-

isopropylphenyl)-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine reacted with DIBALH to give 5-fluoro-N4-(3-isopropylphenyl)-N2-[(2-hydroxymethylene)benzofur-5-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 21.93 min.; purity: 99 %; MS (m/e): 393 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.33 (1H, s), 9.23 (1H, s), 8.18 (1H, d, J= 3.9 Hz), 8.03 (1H, s), 7.86 (1H, d, J= 7.1 Hz), 7.57 (1H, s), 7.49 (2H, m), 7.33 (1H, t, J= 7.1 Hz), 7.05 (1H, d, J= 7.1 Hz), 6.69 (1H, s), 5.54 (1H, t, J= 5.7 Hz), 4.63 (2H, d, J= 5.7 Hz), 2.90 (1H, sept, J= 6.9 Hz), 1.26 (6H, d, J= 6.9 Hz).

7.3.626 N4-(3-*tert*-Butylphenyl)-5-fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-2,4-pyrimidinediamine R940282

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, N4-(3-*tert*-butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine reacted with DIBALH to give N4-(3-*tert*-butylphenyl)-5-fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-2,4-pyrimidinediamine.

LCMS: ret. time: 21.63 min.; Purity: 98 %; MS (m/e): 396 (M⁺).

7.3.627 N4-[3,4-Bis(hydroxymethyl)phenyl]-5-fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940292)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[2-(N-morpholino)ethyleneoxy]phenyl]-2,4-pyrimidinediamine, N2-(3-ethoxycarbonylmethyleneoxyphenyl)-N4-[6-(3,3-dihydroisobenzofuranyl-1-one)]-5-fluoro-2,4-pyrimidinediamine reacted with DIBALH to give N4-[3,4-bis(hydroxymethyl)phenyl]-5-fluoro-N2-[3-(2-hydroxyethyleneoxy)phenyl]-2,4-pyrimidinediamine. LCMS: retn, time: 13.06 min.; purity: 100 %; MS (m/e): 400 (M⁺).

7.3.628 (R935149): N2-(3,4-Ethylenedioxyphenyl)-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-5-fluoro-2,4-pyrimidinediamine

2-Chloro-5-fluoro-N4-[4-[ethoxycarbonyl(dimethyl)methyl]phenyl]-N2-(3,4-ethylenedioxyphenyl)-2,4-pyrimidinediamine was reduced with 10 eq. DIBALH (1.0 M in toluene) at 0 °C in dichloromethane. Reaction was quenched with methanol, diluted with ethylacetate followed by the addition of aqueous Rochelle's salt solution, stirred at room temperature for 30 minutes followed by the addition of anhydrous sodium sulfate. The solution was filtered through Celite, concentrated and purified the concentrated by silica gel

column chromatography to furnish the N2-(3,4-ethylenedioxyphenyl)-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.01 (br s, 1H), 9.6 (br s, 1H), 8.13 (d, 1H, J= 4.7 Hz), 7.58 (d, 2H, J= 8.2 Hz), 7.31 (d, 2H, J= 8.8 Hz), 7.18 (d, 1H, J= 2.3 Hz), 6.88 (dd, 1H, J= 2.3 and 8.8 Hz), 6.73 (d, 1H, J= 8.8 Hz), 4.21-4.19 (m, 4H), 3.56 (br s, 2H), 1.20 (s, 6H); LCMS: ret. time: 20.34 min.; purity: 98%; MS (*m/e*): 411 (MH⁺).

7.3.629 (R935151): 5-Fluoro-N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-[4-[(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to provide 5-fluoro-N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.89 (d, 1H, J= 2.9 Hz), 7.46 (d, 3H, J= 8.8 Hz), 7.27 (d, 2H, J= 8.2 Hz), 6.89 (d, 2H, J= 9.3 Hz), 6.68-6.65 (m, 1H), 4.53 (septet, 1H, J= 5.8 Hz), 3.57 (s, 2H), 1.36 (d, 6H, J= 5.8 Hz), 1.31 (s, 6H); LCMS: ret. time: 23.43 min.; purity: 99%; MS (*m/e*): 411 (MH⁺).

7.3.630 (R935153): 5-Fluoro-N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine:

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-[4-[ethoxycarbonyl(dimethyl)methyl]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to provide 5-fluoro-N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.89 (d, 1H, J= 2.9 Hz), 7.57 (s, 1H), 7.41 (d, 2H, J= 8.8 Hz), 7.29 (d, 2H, J= 8.2 Hz), 7.16 (d, 1H, J= 8.2 Hz), 7.10 (d, 1H, J= 8.8 Hz), 6.80-6.55 (m, 2H), 5.58 (s, 2H), 1.30 (s, 6H); LCMS: ret. time: 18.01 min.; purity: 98%; MS (*m/e*): 369 (MH⁺).

7.3.631 (R935154): N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyoxyphenyl)-2,4-

pyrimidinediamine was reduced with DIBALH to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.88 (d, 1H, J= 3.8 Hz), 7.34 (t, 1H, J= 2.3 Hz), 7.19 (dd, 1H, J= 2.3 and 8.2 Hz), 7.14 (d, 1H, J= 7.6 Hz), 7.01-6.97 (m, 2H), 6.84 (d, 1H, J= 8.8 Hz), 6.53 (dd, 1H, J= 1.7 and 7.6 Hz), 4.26 (s, 4H), 3.98 (t, 2H, J= 4.1 Hz), 3.89 (t, 2H, J= 4.1 Hz); LCMS: ret. time: 18.36 min.; purity: 99%; MS (*m/e*): 399 (MH⁺).

7.3.632 (R935155): 5-Fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine:

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine was reduced to 5-fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine with DIBALH. ¹H NMR (CDCl₃): δ 7.73 (d, 1H, J= 3.5 Hz), 7.33 (d, 2H, J= 8.8 Hz), 7.15 (br s, 1H), 7.04 (app t, 2H, J= 8.2 and 7.6 Hz), 6.78 (d, 2H, J= 8.8 Hz), 6.49 (d, 1H, J= 7.6 Hz), 3.95 (t, 2H, J= 4.7 Hz), 3.80 (t, 2H, J= 4.7 Hz); LCMS: ret. time: 14.49 min.; purity: 98%; MS (*m/e*): 357 (MH⁺).

7.3.633 (R935156): 5-Fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(4-isopropoxyphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to provide 5-fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.90 (d, 1H, J= 3.5 Hz), 7.45 (d, 2H, J= 8.8 Hz), 7.34 (t, 1H, J= 2.3 Hz), 7.13 (t, 1H, J= 8.2 Hz), 6.93 (m, 3H), 7.76 (d, 1H, J= 2.3 Hz), 6.52 (dd, 1H, J= 2.3 and 8.2 Hz), 4.52 (septet, 1H, J= 5.7 Hz), 3.95-3.85 (m, 4H), 1.34 (d, 6H, J= 5.7 Hz); LCMS: ret. time: 21.17 min.; purity: 98%; MS (*m/e*): 399 (MH⁺).

7.3.634 (R935158): 5-Fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N2-[4-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine:

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-[4-[1-ethoxycarbonyl]-

methyl)ethyl]phenyl]-5-fluoro-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to give 5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N2-[4-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.83 (d, 1H, J= 3.5 Hz), 7.49 (d, 2H, J= 8.8 Hz), 7.35 (d, 2H, J= 8.8 Hz), 7.31 (d, 2H, J= 8.8 Hz), 6.82 (d, 2H, J= 8.8 Hz), 4.03 (t, 2H, J= 4.7 Hz), 3.89 (t, 2H, J= 4.7 Hz), 3.56 (s, 2H), 1.30 (s, 6H); LCMS: ret. time: 16.86 min.; purity: 96%; MS (*m/e*): 413 (MH⁺).

7.3.635 (R935160): 5-Fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine:

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(4-isopropoxyphenyl)-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to give 5-fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.12 (s, 1H), 8.92 (s, 1H), 7.98 (d, 1H, J= 3.5 Hz), 7.59 (d, 2H, J= 8.8 Hz), 7.49 (d, 2H, J= 9.3 Hz), 6.86 (d, 2H, J= 8.8 Hz), 6.76 (d, 2H, J= 9.3 Hz), 4.82 (t, 1H, J= 4.9 Hz), 4.55 (septet, 1H, J= 6.4 Hz), 3.89 (t, 2H, J= 5.3 Hz), 3.67 (app q, 2H, J= 5.3 and 4.9 Hz), 1.24 (d, 6H, J= 6.4 Hz); LCMS: ret. time: 19.56 min.; purity: 100%; MS (*m/e*): 399 (MH⁺).

7.3.636 (R935161): 5-Fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N2-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-[4-(1-ethoxycarbonyl-1-methyl)ethyl]phenyl]-5-fluoro-N2-(3-methoxycarbonylmethylphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to give 5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N2-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.27 (s, 1H), 9.11 (s, 1H), 8.07 (d, 1H, J= 3.5 Hz), 7.67 (d, 2H, J= 8.8 Hz), 7.38-7.24 (m, 4H), 7.06 (t, 1H, J= 8.2 Hz), 6.46 (dd, 1H, J= 8.2 Hz), 4.83 (t, 1H, J= 5.3 Hz), 4.66 (t, 1H, J= 5.3 Hz), 3.88 (t, 2H, J= 5.3 Hz), 3.67 (t, 1H, J= 5.3 Hz), 3.66 (t, 1H, J= 5.3 Hz), 3.38 (d, 2H, J= 5.3 Hz), 1.20 (s, 6H); LCMS: ret. time: 17.17 min.; purity: 96%; MS (*m/e*): 413 (MH⁺).

7.3.637 (R935168): 5-Fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-[4-(1-ethoxycarbonyl-1-methyl)ethylphenyl]-5-fluoro-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to produce 5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.21 (s, 1H), 8.93 (s, 1H), 8.00 (d, 1H, J= 4.1 Hz), 7.62 (d, 2H, J= 8.8 Hz), 7.48 (d, 2H, J= 8.8 Hz), 7.27 (d, 2H, J= 8.8 Hz), 6.75 (d, 2H, J= 8.8 Hz), 4.65 (t, 1H, J= 5.3 Hz), 4.47 (septet, 1H, J= 5.8 Hz), 3.38 (d, 2H, J= 5.3 Hz), 1.22 (d, 6H, J= 5.8 Hz), 1.20 (s, 6H); LCMS: ret. time: 22.97 min.; purity: 99%; MS (*m/e*): 411 (MH⁺).

7.3.638 (R935170): 5-Fluoro-N4-[3-(2-hydroxyethoxy)phenyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine:

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-(3-hydroxyphenyl)-N4-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to produce 5-fluoro-N4-[3-(2-hydroxyethoxy)phenyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.23 (s, 1H), 9.14 (s, 1H), 9.06 (s, 1H), 8.07 (d, 1H, J= 4.1 Hz), 7.51 (dd, 1H, J= 1.7 and 7.6 Hz), 7.30 (app t, 1H, J= 2.3 and 1.7 Hz), 7.19 (t, 1H, J= 8.2 Hz), 7.13 (br s, 1H), 7.11 (m, 1H), 6.96 (t, 1H, J= 7.6 Hz), 6.61 (dd, 1H, J= 2.3 and 8.2 Hz), 6.28 (dd, 1H, J= 2.3 Hz and 8.2 Hz), 4.84 (t, 1H, J= 5.8 Hz), 3.92 (t, 2H, J= 5.2 Hz), 3.68 (app qt, 2H, J= 5.2 Hz); LCMS: ret. time: 14.71 min.; purity: 96%; MS (*m/e*): 357 (MH⁺).

7.3.639 (R935171): 5-Fluoro-N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-pyrimidine-2,4-diamine, N4-[4-(1-ethoxycarbonyl-1-methyl)ethylphenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to give 5-fluoro-N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.24 (s, 1H), 9.13 (s, 1H), 9.01 (s, 1H), 8.04 (d, 1H, J= 3.5 Hz), 7.68 (d, 2H, J= 8.8 Hz), 7.29 (d, 2H, J= 8.8 Hz),

7.16 (br s, 1H), 7.07 (m, 1H), 6.94 (t, 1H, 8.8 Hz), 6.30 (m, 1H), 4.64 (t, 1H, J= 5.8 Hz), 3.38 (d, 2H, J= 5.3 Hz), 1.20 (s, 6H); LCMS: ret. time: 17.36 min.; purity: 100%; MS (*m/e*): 369 (MH⁺).

5 **7.3.640 (R935174): 5-Fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-N4-(2-hydroxymethylbenzofur-5-yl)-2,4-pyrimidinediamine**

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-(2-carbomethoxybenzofur-5-yl)-5-fluoro-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to provide 5-fluoro-N2[4-(2-hydroxyethoxy)phenyl]-N2-(2-hydroxymethylbenzofur-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.26 (s, 1H), 8.94 (s, 1H), 8.01 (d, 1H, J= 4.1 H), 7.99 (s, 1H), 7.52-7.45 (m, 4H), 6.72 (d, 2H, J= 9.3 Hz), 6.66 (s, 1H), 5.46 (t, 1H, J= 5.3 Hz), 4.82 (t, 1H, J= 5.8 Hz), 4.55 (d, 2H, J= 5.8 Hz), 3.89 (t, 2H, J= 5.3 Hz), 3.67 (app qt, 2H, J= 5.3 Hz); LCMS: ret. time: 14.97 min.; purity: 91%; MS (*m/e*): 411 (MH⁺).

15 **7.3.641 (R935176): N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine:**

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to provide N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.22 (s, 1H), 8.98 (s, 1H), 8.05 (d, 1H, J= 3.5 Hz), 7.47 (dd, 1H, J= 1.1 and 8.2 Hz), 7.27 (t, 1H, J= 1.7 Hz), 7.23 (d, 1H, J= 2.3 Hz), 7.18 (t, 1H, J= 8.2 Hz), 7.05 (dd, 1H, J= 2.3 and 8.8 Hz), 6.68 (d, 1H, J= 8.2 Hz), 6.61 (dd, 1H, J= 1.7 and 8.8 Hz), 4.85 (t, 1H, J= 5.3 Hz), 4.18-4.14 (m, 4H), 3.91 (t, 2H, J= 5.3 Hz), 3.68 (qt, 2H, J= 5.3 Hz); LCMS: ret. time: 17.35 min.; purity: 92%; MS (*m/e*): 399 (MH⁺).

7.3.642 (R935177): 5-Fluoro-N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(2-hydroxymethylbenzofur-5-yl)-2,4-pyrimidinediamine

30 In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-(2-carbomethoxybenzofur-5-yl)-N2-[4-(1-ethoxycarbonyl-1-methyl)ethylphenyl]-5-fluoro-2,4-

pyrimidinediamine was reduced with DIBALH to produce 5-fluoro- N2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-N4-(2-hydroxymethylbenzofur-5-yl)-2,4-pyrimidinediamine. LCMS: ret. time: 18.17 min.; purity: 94%; MS (*m/e*): 423 (MH⁺).

7.3.643 (R935178): 5-Fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-N4-(2-hydroxymethylbenzofur-5-yl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-(2-carbomethoxybenzofur-5-yl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to provide 5-fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-N4-(2-hydroxymethylbenzofur-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.93 (s, 1H), 9.12 (s, 1H), 8.07 (d, 1H, J= 3.6 Hz), 8.01 (d, 1H, J= 2.3 Hz), 7.55-7.46 (m, 2H), 7.29 (br s, 1H), 7.23 (d, 1H, J= 8.2 Hz), 7.03 (t, 1H, J= 8.2 Hz), 6.68 (s, 1H), 6.44 (dd, 1H, J= 2.3 and 8.2 Hz), 5.47 (t, 1H, J= 5.8 Hz), 4.80 (t, 1H, J= 5.3 Hz), 4.55 (d, 2H, J= 5.3 Hz), 3.81 (qt, 2H, J= 5.3 Hz), 3.63 (qt, 2H, J= 5.3 Hz); LCMS: ret. time: 15.41 min.; purity: 88%; MS (*m/e*): 411 (MH⁺).

7.3.644 (R935181): N4-(3,5-Dimethoxyphenyl)-5-fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-(3,5-dimethoxyphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidineamine was reduced with DIBALH to give N4-(3,5-dimethoxyphenyl)-5-fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-2,4-pyrimidinediamine: ¹H NMR (DMSO-d₆): δ 9.24 (s, 1H), 9.18 (s, 1H), 8.11 (d, 1H, J= 3.5 Hz), 7.31-7.26 (m, 2H), 7.05 (d, 1H, J= 8.2 Hz), 6.99 (d, 1H, J= 2.3 Hz), 6.43 (dd, 1H, J= 2.3 Hz, 8.2 Hz), 6.20 (t, 1H, J= 2.3 Hz), 4.80 (t, 1H, J= 5.8 Hz), 3.83 (t, 2H, J= 5.3 Hz), 3.67 (s, 6H), 3.66-3.60 (m, 2H); LCMS: ret. time: 18.78 min.; purity: 95%; MS (*m/e*): 400 (MH⁺).

7.3.645 (R935183): 5-Fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-[4-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine was reduced with DIBAL-H to provide 5-fluoro-N2-[4-(2-

hydroxyethoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.15 (s, 1H), 8.97 (s, 1H), 8.00 (d, 1H, J= 3.5 Hz), 7.49 (d, 2H, J= 8.8 Hz), 7.40-7.31 (m 2H), 6.88 (d, 1H, J= 8.8 Hz), 6.80 (d, 2H, J= 8.8 Hz), 4.82 (t, 1H, J= 5.3 Hz), 4.12-4.04 (m 4H), 3.90 (t, 2H, J= 5.2 Hz), 3.70-3.65 (app qt, 2H, J= 5.3 Hz), 2.07 (q, 2H, J= 5.3 Hz); LCMS: ret. time: 17.05 min.; purity: 96%; MS (*m/e*): 413 (MH⁺).

7.3.646 (R935186): 5-Fluoro-N2-[4-(2-hydroxyethoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine was reduced with DIBALH to provide 5-fluoro-N2-[3-(2-hydroxyethoxy)phenyl]-N4-(3,4-propylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.21 (s, 1H), 9.14 (s, 1H), 8.07 (d, 1H, J= 3.5 Hz), 7.42-7.36 (m, 2H), 7.29-7.24 (m, 2H), 7.07 (t, 1H, J= 8.2 Hz), 6.90 (d, 1H, J= 8.8 Hz), 6.45 (dd, 1H, J= 1.7 and 8.3 Hz), 4.82 (t, 1H, J= 5.3 Hz), 4.12-4.04 (app q, 2H, J= 5.3 Hz), 3.86 (t, 2H, J= 5.3 Hz), 3.67 (app qt, 2H, J= 5.3 Hz), 2.07 (q, 2H, J= 5.3 Hz); LCMS: ret. time: 17.95 min.; purity: 96%; MS (*m/e*): 413 (MH⁺).

7.3.647 N4-(4-*tert*-Butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine (R926720)

The reaction of N2-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine and lithium hydroxide (LiOH) in THF:H₂O at room temperature gave N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.01 (bs, 1H), 9.69 (bs, 1H), 8.13 (d, 1H, J= 4.8 Hz), 7.57 (d, 2H, J= 8.7 Hz), 7.50 (s, 1H), 7.35 (d, 2H, J= 8.1 Hz), 7.13 (d, 1H, J= 8.7 Hz), 6.75 (d, 1H, J= 9.0 Hz), 5.21 (dd, 1H, J= 6.3 and 10.5 Hz), 3.49 (dd, 1H, J= 10.5 and 16.5 Hz), 3.17 (dd, 1H, J= 6.6 and 16.5 Hz), 1.27 (s, 9H); LCMS: ret. time: 22.53 min.; purity: 93 %; MS (*m/e*): 423 (MH⁺).

7.3.648 N4-(4-*tert*-Butylphenyl)-N2-(3-carboxymethyleneoxyphenyl)-5-fluor-2,4-pyrimidinediamine (R926726)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and lithium

hydroxide were reacted to yield N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(3-carboxymethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.88 (bs, 1H), 9.29 (s, 1H), 9.16 (s, 1H), 8.07 (d, 1H, J= 3.3 Hz), 7.68 (d, 2H, J= 8.7 Hz), 7.35-7.31 (m, 3H), 7.26 (d, 1H, J= 8.4 Hz), 7.06 (t, 1H, J= 8.4 Hz), 6.41 (dd, 1H, J= 2.4 and 8.4 Hz), 4.54 (s, 2H), 1.27 (s, 9H); ¹⁹F NMR (DMSO-d₆): - 46463; LCMS: ret. time: 22.94 min.; purity: 97 %; MS (m/e): 411 (MH⁺).

7.3.649 5-Fluoro-N2-[3-(carboxymethyleneoxy)phenyl]- N4-[4-(isopropoxy)phenyl]-2,4-pyrimidinediamine (R926731)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, 5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine and lithium hydroxide were reacted to yield 5-fluoro-N2-(3-carboxymethyleneoxyphenyl)-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 6.19 (bs, 1H), 9.01 (s, 1H), 8.02 (d, 1H, J= 3.9 Hz), 7.63 (d, 2H, J= 9.3 Hz), 7.19-7.14 (m, 2H), 6.96 (t, 1H, J= 8.7 Hz), 6.87 (d, 2H, J= 9.6 Hz), 6.28 (dd, 1H, J= 2.45 and 9.0 Hz), 4.56 (2q, 1H, J= 6.6 Hz), 3.94 (s, 2H), 1.24 (d, 6H, J= 6.6 Hz); LCMS: ret. time: 20.13 min.; purity: 100 %; MS (m/e): 413 (MH⁺).

7.3.650 N2,N4-Bis(4-carboxymethyleneoxy)phenyl-5-fluoro-2,4-pyrimidinediamine (R926560)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, the hydrolysis of N2,N4-bis(4-methoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine with LiOH gave N2,N4-bis(4-carboxymethyleneoxy)phenyl-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.86 (bs, 1H), 7.55 (d, 2H, J= 9.0 Hz), 7.32 (bd, 2H, J= 9.3 Hz), 6.95 (m, 4H), 4.66 (s, 2H); ¹⁹F NMR (CDCl₃): - 21852; LCMS: ret. time: 15.16 min.; purity: 77%; MS (m/e): 429 (MH⁺).

7.3.651 N2-(3-Carboxymethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926483)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of N2-(3-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-

pyrimidinediamine with LiOH gave N2-(3-carboxymethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.90 (s, 1H), 9.20 (s, 2H), 8.05 (d, 1H, J= 1.2 Hz), 7.32-7.21 (m, 3H), 7.08 (t, 1H, J= 8.1 Hz), 6.80 (d, 1H, J= 8.4 Hz), 6.40 (dd, 1H, J= 1.8 and 8.2 Hz), 4.53 (s, 2H), 4.20 (s, 4H); LCMS: ret. time: 18.26 min.; purity: 100%; MS (m/e): 413 (MH⁺).

7.3.652 N2-(3-Carboxymethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945126)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-(methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with LiOH gave N2-(3-carboxymethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 4.55 (s, 2H), 6.43 (dd, J= 2.1, 8.1 Hz, 1H), 6.48 (dd, J= 2.1 and 7.2 Hz, 1H), 7.06-7.13 (m, 3H), 7.28-7.34 (m, 3H), 8.09 (d, J= 3.6 Hz, 1H), 9.22 (br, 1H), 9.28 (br, 1H), 9.34 (br, 1H); ¹⁹F NMR (282 MHz, DMSO-d₆): δ -163.85; LCMS: ret. time: 15.88 min.; purity: 100%; MS (m/e): 370.63 (MH⁺).

7.3.653 N2-(4-Carboxymethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926238)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine with LiOH gave N2-(carboxymethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.16 (d, 1H, J= 4.8 Hz), 7.37 (bd, 2H, J= 9 Hz), 7.25 (d, 1H, J= 3 Hz), 7.08 (m, 1H), 6.83 (m, 3H), 4.64 (s, 2H), 4.23 (s, 4H); LCMS: ret. time: 19.15 min.; purity: 100%; MS (m/e): 413 (MH⁺).

7.3.654 N2-(4-Carboxymethyleneoxyphenyl)-5-Fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926564)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, 5-fluoro-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine upon treatment with LiOH gave 5-fluoro-N2-(4-carboxymethyleneoxyphenyl)-N4-(3-

hydroxyphenyl)-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.89 (d, 1H, $J=5.1$ Hz), 7.34 (dd, 2H, $J=2.1$ and 9.3 Hz), 7.19-7.08 (m, 2H), 6.98 (dd, 2H, $J=2.4$ and 8.4 Hz), 6.69 (m, 1H), 4.68 (s, 2H); ^{19}F NMR (CD_3OD): - 21860; LCMS: ret. time: 15.69 min.; purity: 99%; MS (m/e): 371 (MH^+).

5 **7.3.655 N2-(2-Carboxybenzofuran-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926478)**

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methoxycarbonylbenzofuran-5-yl)-4-pyrimidinediamine upon LiOH treatment gave N2-(2-carboxybenzofuran-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.97 (bd, 2H), 7.60-7.44 (m, 4H), 7.20-7.05 (m, 3H), 6.69 (bd, 1H); ^{19}F NMR (CD_3OD): - 21844; LCMS: ret. time: 16.77 min.; purity: 100%; MS (m/e): 381 (MH^+).

15 **7.3.656 N2-(2-Carboxyindol-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926479)**

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, N2-(2-ethoxycarbonylindol-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine upon LiOH treatment gave N2-(2-carboxyindol-5-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.83 (m, 1H), 7.73 (s, 1H), 7.50 (bd, 1H, $J=8.7$ Hz), 7.30-7.11 (m, 5H), 6.68 (bd, 1H); LCMS: ret. time: 16.50 min.; purity: 97%; MS (m/e): 380 (MH^+).

7.3.657 N4-(4-*tert*-Butylphenyl)-N2-(2-carboxybenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine (R926481)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, LiOH treatment with N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine gave N4-(4-*tert*-butylphenyl)-N2-(2-carboxybenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 9.3 (bd, 2H), 8.25 (s, 1H), 8.10 (s, 1H), 7.65-7.30 (m, 5H), 1.25 (s, 9H); ^{19}F NMR (CD_3OD): - 21844; LCMS: ret. time: 23.32 min.; purity: 100%; MS (m/e): 421 (MH^+).

7.3.658 N4-(3-*tert*-Butylphenyl)-N2-[3-carboxymethyleneoxyphenyl]-5-fluoro-2,4-pyrimidinediamine R940280

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-[2,3-dihydro-2-(carboxy)benzofuran-5-yl]-2,4-pyrimidinediamine, N4-(3-*tert*-butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine was reacted with LiOH to give N4-(3-*tert*-butylphenyl)-N2-(3-carboxymethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 23.61 min.; purity: 99 %; MS (m/e): 410 (M⁺), 412 (MH⁺); ¹H NMR (DMSO-d₆): δ 9.45 (1H, s), 9.33 (1H, s), 8.21 (1H, d, J= 3.9 Hz), 7.98 (1H, d, J= 6.6 Hz), 7.60 (1H, t, J= 2 Hz), 7.44-7.34 (3H, m), 7.24-7.15 (2H, m), 6.54 (1H, d, J= 7.8 Hz), 4.68 (2H, s), 1.36 (9H, s).

7.3.659 N2-(3-Carboxymethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950190)

The reaction of N2-(3-ethoxycarbonylmethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (0.1 g) and LiOH (10 equivalents) in MeOH:water (1:1, v/v) for 1h at room temperature followed by treatment with aqueous HCl gave the solid. The resulting solid was filtered, washed with water and dried to give N2-(3-carboxymethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 18.23 min.; purity: 87.6%; MS (m/e): 412.01 (MH⁺).

7.3.660 N2-(Carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethyloxy)phenyl]-2,4-pyrimidinediamine (R950230)

In a manner similar to the preparation of N2-(3-carboxymethyleneaminophenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the hydrolysis of N2-(ethoxycarbonylmethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethyloxy)phenyl]-2,4-pyrimidinediamine with LiOH gave N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethyloxy)phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.15 min.; purity: 78.3%; MS (m/e): 413.01 (MH⁺).

7.3.661 5-Fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950231)

A mixture of N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine (10 mg), 2-aminoethanol (10 equiv.) and PyBroP (2 equiv.) was stirred in 0.5 ml DMF for 24 hours at room temperature. The

mixture was diluted with water, extracted with EtOAc and the organic phase was dried over MgSO₄. The solvent was removed under reduced pressure and the residue was subjected to column chromatography on silica gel (CHCl₃:Acetone, 2:1) to give 5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.98 min.; purity: 92.6%; MS (m/e): 455.97 (MH⁺).

7.3.662 N2-[3-(N-2-Aminoethylamino)carbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylenoxy)phenyl]-2,4-pyrimidinediamine (R950232)

In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and 1,2-ethylenediamine were reacted to afford N2-[3-(N-2-aminoethylamino)carbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylenoxy)phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 11.31 min.; purity: 93.6%; MS (m/e): 454.94 (MH⁺).

7.3.663 5-Fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-(N-methylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950233)

In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and methylamine were reacted to give 5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-(N-methylamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 14.93 min.; purity: 92.9%; MS (m/e): 426.27 (MH⁺).

7.3.664 5-Fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-methylamino)ethylamino]carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950234)

In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and N-methylethylenediamine were reacted to give 5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-methylamino)ethylamino]

carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 11.39 min.; purity: 97.7%; MS (m/e): 468.96 (MH⁺).

5 **7.3.665 N2-[3-[N-(2-N-Benzylamino)ethylamino]carbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine (R950235)**

10 In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and N-benzylethylenediamine were reacted to give N2-[3-[N-(2-N-benzylamino)ethylamino]carbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 14.39 min.; purity: 97.3%; MS (m/e): 545.01 (MH⁺).

15 **7.3.666 5-Fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-(N-morpholino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950236)**

20 In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and morpholine were reacted to afford 5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-(N-morpholino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 15.24 min.; purity: 94.6xx%; MS (m/e): 482.40 (MH⁺).

25 **7.3.667 N2-[3-(3-N,N-Dimethylaminopropyl)aminocarbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine (R950237)**

30 In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and N,N-dimethylpropanediamine were reacted to give N2-[3-(3-N,N-Dimethylaminopropyl)aminocarbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 13.33 min.; purity: 91.4%; MS (m/e): 497.47 (MH⁺).

7.3.668 N2-[3-[N-(2,3-Dihydroxypropyl)amino]carbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine (R950238)

5 In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and 1-amino-2,3-propanediol were reacted to give N2-[3-[N-(2,3-dihydroxypropyl)amino]carbonylmethyleneaminophenyl]-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 12.86 min.; purity: 90.0%; MS (m/e): 486.40 (MH⁺).

7.3.669 5-Fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine (R950239)

15 In like manner to the preparation of N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-[N-(2-hydroxyethylamino)]carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and 4-(2-aminoethyl)morpholine were reacted to give 5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-N2-[3-(N-morpholinoethyleneamino)carbonylmethyleneaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 13.52 min.; purity: 92.4%; MS (m/e): 525.47 (MH⁺).

7.3.670 2,4-Bis[N-(L)-tyrosine methyl ester]-5-ethoxycarbonylpyrimidine (R926514)

and

25 5-Ethoxycarbonyl-2-methoxy-4-[N-(L)-tyrosine methyl ester]pyrimidine (R926513)

A mixture of tyrosine methyl ester (58 mg, 0.3 mmol), 2,4-dichloro-5-ethoxycarbonylpyrimidine (44 mg, 0.1 mmol) in MeOH (2mL) was heated in a sealed tube at 100 °C for a period of overnight, diluted with H₂O (20 mL), acidified with 2N HCl and extracted with ethyl acetate (3 x 25 mL). The solvent was evaporated and the residue was purified by preparative TLC using 30% EtOAc/Hexanes to obtain a mixture of 2,4-bis[N-(L)-tyrosine methyl ester]-5-ethoxycarbonylpyrimidine (R926514). ¹H NMR (CDCl₃): δ

8.60 (1H, J= 6.6 Hz), 8.36 (s, 1H), 7.05 (d, 2H, J= 8.7 Hz), 6.84 (d, 2H, J= 8.1 Hz), 6.74 (d, 2H, J= 9 Hz), 6.54 (d, 2H, J= 9 Hz), 4.82 (t, 2H, J= 6 Hz), 4.25 (q, 2H, J= 6.3 Hz), 3.73 (s, 3H), 3.72 (s, 3H), 3.06 (m, 4H), 1.31 (t, 3H, J= 7.2 Hz) and 5-ethoxycarbonyl-2-methoxy-4-[N-(L)-tyrosine methyl ester]pyrimidine (**R926513**): ¹H NMR (CDCl₃): δ 8.78 (s, 1H), 8.65 (d, 1H, J= 6.9 Hz), 7.02 (dd, 2H, J= 2.1 and 6.3 Hz), 6.77 (dd, 2H, J= 2.4 and 6.6 Hz), 4.93 (q, 1H, J= 1.5 and 6.9 Hz), 4.30 (q, 2H, J= 8.1 Hz), 3.90 (s, 3H), 3.70 (s, 3H), 3.17 (dd, 1H, J= 5.4 Hz), 3.06 (dd, 1H, J= 7.5 and 7.8 Hz), 1.33 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 22.58 min.; purity: 99%; MS (m/e): 376 (M⁺).

7.3.671 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926252)

In like manner to the preparation of N2,N4-bis[N-(L)-tyrosine methyl ester]-5-ethoxycarbonylpyrimidine, the reaction of 2,4-dichloro-5-ethoxycarbonylpyrimidine with 3,4-ethylenedioxyaniline gave N2,N4-bis(3,4-ethylenedioxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.01 (s, 1H), 9.65 (bs, 1H), 8.62 (s, 1H), 7.18 (bs, 2H), 7.04 (dd, 1H, J= 1.8 and 8.7 Hz), 6.93 (d, 1H, J= 7.5 Hz), 6.76 (d, 1H, J= 8.7 Hz), 6.65 (d, 1H, J= 8.7 Hz), 4.28 (q, 2H, J= 6.9 Hz), 1.31 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 27.25 min.; purity: 100%; MS (m/e): 451 (MH⁺).

7.3.672 N2,N4-Bis(4-methoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (R926253)

In like manner to the preparation of N2,N4-bis[N-(L)-tyrosine methyl ester]-5-ethoxycarbonylpyrimidine, the reaction of 2,4-dichloro-5-ethoxycarbonylpyrimidine with ethyl 4-aminophenoxyacetate gave N2,N4-bis(4-methoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-2,4-pyrimidinediamine (**R926253**). ¹H NMR (CD₃OD): δ 8.60 (bs, 1H), 7.4 (bs, 1H), 7.33 (d, 4H, J= 9Hz), 6.94 (bd, 4H), 4.76 (s, 2H), 4.75 (s, 2H), 4.44 (q, 2H, J= 6.9 Hz), 3.79 (s, 3H), 1.40 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 25.83 min.; purity: 89%; MS (m/e): 511 (MH⁺).

7.3.673 2,4-Bis[N-(L)-phenylalaninyl ethyl ester]-5-ethoxycarbonylpyrimidine (R926526)

In like manner to the preparation of N2,N4-bis[N-(L)-tyrosine methyl ester]-5-ethoxycarbonylpyrimidine, the reaction of 2,4-dichloro-5-ethoxycarbonylpyrimidine with ethyl (L)-phenylalanine ethyl ester in MeOH or EtOAc gave 2,4-bis[N-(L)-phenylalanine ethyl ester]-5-ethoxycarbonylpyrimidine. ¹H NMR (CDCl₃): δ 8.55 (d, 1H, J= 7.2 Hz), 8.51

(s, 1H), 7.35-7.10 (m, 10H), 5.88 (d, 1H, J= 6 Hz), 4.88 (ddd, 1H, J= 6.3 Hz), 4.80 (ddd, 1H, J= 6.3 Hz), 4.23 (q, 2H, J= 7.2 Hz), 4.12 (q, 4H, J= 7.2 Hz), 3.65 (t, 2H, J= 6 Hz), 3.56 (t, 2H, J= 6.0 Hz), 1.30 (t, 2H, J= 6 Hz), 1.30 (t, 3H, J= 7.2 Hz), 1.20 (m, 6H); LCMS: ret. time: 32.22 min.; purity: 89%; MS (m/e): 535 (MH⁺).

5 **7.3.674 2,4-Bis[N-(L)-valinyl ethyl ester]-5-ethoxycarbonylpyrimidine (R926527)**

In like manner to the preparation of N2,N4-bis[N-(L)-tyrosine methyl ester]-5-ethoxycarbonylpyrimidine, the reaction of 2,4-dichloro-5-ethoxycarbonylpyrimidine with ethyl (L)-valine ethyl ester in MeOH or EtOAc gave 2,4-bis[N-(L)-valinyl ethyl ester]-5-ethoxycarbonylpyrimidine. ¹H NMR (CDCl₃): δ 8.59 (d, 1H, J= 7.8 Hz), 8.56 (s, 1H), 5.69 (d, 1H, J= 8.7 Hz), 4.62 (m, 1H), 4.51 (m, 1H), 4.25 (q, 2H, J= 7.5 Hz), 4.20 (m, 4H), 2.20 (m, 2H), 1.34 (t, 3H, J= 7.8 Hz), 1.27 (t, 6H, J= 7.5 Hz), 1.00 (m, 12H); LCMS: ret. time: 29.27 min.; purity: 97%; MS (m/e): 439 (MH⁺).

15 **7.3.675 5-Ethoxycarbonyl-N2-(3-hydroxyphenyl)-4-[N-(L)-phenylalanine ethyl ester]-2-pyrimidineamine (R926528)**

The reaction of 2-chloro-N4-(3-hydroxyphenyl)-5-ethoxycarbonylpyrimidineamine with 3 equivalents of (L)-N-phenylalanine ethyl ester in methanol at 80-100 °C for 24 h followed by dilution with water and acidification with 2N HCl have the acidic solution. The resulting solution was extracted with EtOAc and the residue was purified by silics gel column chromatography to afford 4-[N-(L)-phenylalanine ethyl ester]-N2-(3-hydroxyphenyl)-5-ethoxycarbonyl-2-pyrimidineamine. ¹H NMR (CDCl₃): δ 9.4 (bs, 1H), 9.13 (d, 1H, J= 6 Hz), 8.45 (bs, 1H), 7.59 (s, 1H), 7.34-7.25 (m, 5H), 7.15 (t, 1H, J= 8.1 Hz), 6.73 (bd, 1H, J= 7.5 Hz), 6.67 (dd, 1H, J= 1.8 and 7.8 Hz), 4.86 (dt, 1H, J= 3 and 5.1 Hz), 4.32 (q, 2H, J= 6.3 Hz), 4.19 (q, 2H, J= 7.2 Hz), 3.30 (dd, 1H, J= 4.8 and 8.7 Hz), 3.18 (dd, 1H, J= 5.1 and 8.7 Hz), 1.36 (t, 3H, J= 6.9 Hz), 1.65 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 27.49 min.; purity: 91%; MS (m/e): 451 (MH⁺).

7.3.676 N2-(3,4-Ethylenedioxyphenyl)-5-ethoxycarbonyl-4-[N-(L)-phenyl glyciny ethyl ester]-2-pyrimidineamine (R926536)

In like manner to the preparation of 4-[N-(L)-phenylalanine ethyl ester]-N2-(3-hydroxyphenyl)-5-ethoxycarbonyl-2-pyrimidineamine, the reaction of 2-chloro-5-ethoxycarbonyl-4-[N-(L)-phenyl glyciny ethyl ester]pyrimidine with 3,4-ethylenedioxyaniline in MeOH or EtOAc gave N2-(3,4-ethylenedioxyphenyl)-5-

ethoxycarbonyl-4-[N-(L)-phenyl glyciny ethyl ester]-2-pyrimidineamine. ¹H NMR (CDCl₃): δ 9.15 (s, 1H), 8.9 (s, 1H), 8.61 (s, 1H), 7.48 (m, 2H), 7.38 (m, 3H), 7.16 (bs, 1H), 6.80 (m, 2H), 5.75 (d, 1H), 4.24 (m, 6H), 3.66 (s, 3H), 1.35 (t, 3H); LCMS: ret. time: 28.16 min.; purity: 85%; MS (m/e): 465 (MH⁺).

5 **7.3.677 N4-(4-tert-Butoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R926579)**

10 In like manner to the preparation of 4-[N-(L)-phenylalanine ethyl ester]-N2-(3-hydroxyphenyl)-5-ethoxycarbonyl-2-pyrimidineamine, the reaction of N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-2-chloro-5-ethoxycarbonyl-4-pyrimidineamine with methyl 4-aminophenoxyacetate gave N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 10.17 (s, 1H), 8.73 (s, 1H), 8.45 (bs, 1H), 7.49 (d, 2H, J= 8.7 Hz), 7.43 (d, 15 2H, J= 8.7 Hz), 7.33 (bs, 1H), 6.87 (d, 2H, J= 6 Hz), 6.84 (d, 2H, J= 5.7 Hz), 4.63 (s, 2H), 4.53 (s, 2H), 4.33 (q, 2H, J= 6.9 Hz), 3.81 (s, 3H), 1.49 (s, 9H), 1.39 (t, 3H, J= 7.5 Hz); LCMS: ret. time: 27.93 min.; purity: 96%; MS (m/e): 553 (MH⁺).

20 **7.3.678 N4-(4-tert-Butoxycarbonylmethyleneoxyphenyl)-N2-(4-methoxycarbonylmethyleneoxyphenyl)-5-methoxycarbonyl-2,4-pyrimidinediamine (R926580)**

25 In like manner to the preparation of 4-[N-(L)-phenylalanine ethyl ester]-N2-(3-hydroxyphenyl)-5-ethoxycarbonyl-2-pyrimidineamine, the reaction of N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-2-chloro-5-ethoxycarbonyl-4-pyrimidineamine with methyl 4-aminophenoxyacetate gave N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-5-methoxycarbonyl-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. 5-methyl ester was obtained due to the cross esterification reaction in MeOH. ¹H NMR (CDCl₃): δ 10.13 (s, 1H), 8.73 (s, 1H), 8.45 (bs, 1H), 7.49 (d, 2H, J= 8.7 Hz), 7.43 (d, 2H, J= 8.7 Hz), 7.33 (bs, 1H), 6.87 (m, 4H), 4.63 (s, 2H), 4.53 (s, 2H), 4.33 (q, 2H, J= 6.9 Hz), 3.88 (s, 3H), 3.81 (s, 3H), 1.49 (s, 9H); LCMS: ret. time: 27.43 min.; purity: 100%; MS 30 (m/e): 539 (MH⁺).

7.3.679 N4-(4-Carboxymethyleneoxyphenyl)-5-ethoxycarbonyl-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R926583)

The treatment of N4-(4-tert-butoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with trifluoroacetic acid in THF:H₂O at room temperature afforded N4-(4-carboxymethyleneoxyphenyl)-5-ethoxycarbonyl-N2-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.03 (s, 1H), 8.65 (s, 1H), 7.49 (bd, 4H, J= 8.7 Hz), 6.89 (d, 2H, J= 9.3 Hz), 6.81 (d, 2H, J= 8.1 Hz), 4.70 (s, 2H), 4.65 (s, 2H), 4.33 (q, 2H, J= 6.9 Hz), 3.81 (s, 3H), 1.49 (s, 9H), 1.39 (t, 3H, J= 7.5 Hz); LCMS: ret. time: 22.28 min.; purity: 73%; MS (m/e): 497 (MH⁺).

7.3.680 N2-(4-Carboxymethyleneoxyphenyl)-5-ethoxycarbonyl-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine (R926584)

The treatment of N2-(4-tert-butoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with trifluoroacetic acid in THF:H₂O at room temperature afforded N2-(4-carboxymethyleneoxyphenyl)-5-ethoxycarbonyl-N4-(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.01 (s, 1H), 8.64 (s, 1H), 7.45 (bd, 4H, J= 7.2 Hz), 6.90 (d, 2H, J= 8.7 Hz), 6.75 (d, 2H, J= 8.4 Hz), 4.80 (s, 2H), 4.38 (s, 2H), 4.26 (q, 2H, J= 7.2 Hz), 3.70 (s, 3H), 1.30 (t, 3H, J= 7.2 Hz); LCMS: ret. time: 22.37 min.; purity: 100%; MS (m/e): 497 (MH⁺).

7.3.681 5-Carboxy-N2-(3-hydroxyphenyl)-N4-[N-(L)-phenylglycine]-2-pyrimidineamine (R926535)

The LiOH hydrolysis of N2-(3-hydroxyphenyl)-5-ethoxycarbonyl-4-[N-(L)-phenylglycine ethyl ester]-2-pyrimidineamine afforded 5-carboxy-N2-(3-hydroxyphenyl)-N4-[N-(L)-phenylglycine]-2-pyrimidineamine. ¹H NMR (CD₃OD): δ 8.89 (s, 1H), 8.50 (s, 1H), 7.43 (m, 2H), 7.33 (m, 3H), 7.14 (m, 2H), 6.98 (m, 2H), 6.62 (m, 1H), 5.71 (s, 1H); LCMS: ret. time: 17.75 min.; purity: 73%; MS (m/e): 382 (MH⁺).

7.3.682 5-Amino-6-ethoxycarbonyl-N2,N4-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925856)

A suspension of 6-ethoxycarbonyl-N2,N4-bis(3-hydroxyphenyl)-5-nitro-2,4-pyrimidinediamine and 10% Pd/C (10% by weight) in ethanol was prepared and reacted in

a Parr bottle under hydrogen gas (20 PSI) for 1h. The reaction mixture was filtered through Celite. Purification by column chromatography gave 5-amino-6-ethoxycarbonyl-N2,N4-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.30 (bs, 1H), 7.18-7.10 (m, 3H), 7.00 (t, 2H, J= 8.1 Hz), 6.59-6.54 (m, 1H), 6.33 (dd, 1H, J= 2.1 and 11.1 Hz), 4.39 (q, 2H, J= 6.9 Hz), 1.43 (t, 3H, J= 6.9 Hz); LCMS: ret. time: 19.24 min.; purity: 100 %; MS (m/e): 382 (MH⁺).

7.3.683 5-Amino-6-ethoxycarbonyl-N2,N4-bis(3,4-ethylenedioxyphenyl)-2,4-pyrimidinediamine (R925857)

In a manner similar to the preparation of 5-amino-6-ethoxycarbonyl-N2,N4-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine, 6-ethoxycarbonyl-N2,N4-bis(3,4-ethylenedioxyphenyl)-5-nitro-2,4-pyrimidinediamine, hydrogen, and 10% Pd/C were reacted to yield 5-amino-6-ethoxycarbonyl-N2,N4-bis(3,4-ethylenedioxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.16 (d, 1H, J= 2.4 Hz), 7.07 (d, 1H, J= 2.4 Hz), 7.04 (dd, 1H, J= 2.4 and 9.0 Hz), 6.84-6.79 (m, 2H), 6.70 (d, 1H, J= 9.0), 4.43 (q, 2H, J= 7.8 Hz), 4.25 (s, 4H), 4.21 (bs, 4H), 1.43 (t, 3H, J= 7.8 Hz); LCMS: ret. time: 23.70 min.; purity: 100 %; MS (m/e): 466 (MH⁺).

7.3.684 5-Amino-N2,N4-bis(ethoxycarbonylmethyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine (R925865)

In a manner similar to the preparation of 5-amino-6-ethoxycarbonyl-N2,N4-bis(3-hydroxyphenyl)-2,4-pyrimidinediamine, 6-ethoxycarbonyl-N2,N4-bis(ethoxycarbonylmethyl)-5-nitro-2,4-pyrimidinediamine, hydrogen, and 10% Pd/C were reacted to yield 5-amino-N2,N4-bis(ethoxycarbonylmethyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 6.25 (bs, 2H), 4.38 (q, 2H, J= 6.9 Hz), 4.23-4.14 (m, 6H), 4.05 (bs, 2H), 1.39 (t, 3H, J= 6.9 Hz), 1.30-1.22 (m, 6H); LCMS: ret. time: 17.67 min.; purity: 95 %; MS (m/e): 370 (MH⁺).

7.3.685 5-Amino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine (R926567)

Hydrogenation of N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine using Pd/C in MeOH at 40 PSI gave 5-amino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.47 (d, 2H, J= 8.7 Hz), 7.41 (d, 2H, J= 8.7 Hz), 6.88 (d, 2H, J= 8.1 Hz), 6.81 (d, 2H, J= 8.7 Hz), 4.63 (s, 2H), 4.59 (s, 2H), 4.41 (q, 2H, J=

7.5 Hz), 4.29 (m, 4H), 1.44 (t, 3H), 1.31 (m, 6H); LCMS: ret. time: 26.15 min.; purity: 97%; MS (m/e): 554 (MH⁺).

7.3.686 N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(phenylaminocarbonylamino)-2,4-pyrimidinediamine (R926571)

A dry reaction flask equipped with a rubber septum and a N₂ inlet was charged with 5-amino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine, equimolar amount of pyridine and phenyl isocyanate at room temperature. The reaction was allowed to stirred at room temperature for overnight and the resulting reaction was poured over n-hexane to precipitate the desired product, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(phenylaminocarbonylamino)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.92 (s, 1H), 7.47 (s, 1H), 7.35 (bt, 5H, J= 8.4 Hz), 7.25 (bt, 2H, J= 7.5 Hz), 7.03 (m, 2H), 6.81 (d, 2H, J= 8.7 Hz), 6.76 (d, 2H, J= 8.7 Hz), 4.60 (s, 2H), 4.58 (s, 2H), 4.29 (m, 6H), 1.45 (m, 9H); LCMS: ret. time: 27.75 min.; purity: 91%; MS (m/e): 673 (MH⁺).

7.3.687 5-Allylaminocarbonylamino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine (R926585)

In like manner to the preparation of N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(phenylaminocarbonylamino)-2,4-pyrimidinediamine, the reaction of 5-amino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine with allyl isocyanate gave 5-allylaminocarbonylamino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine. LCMS: ret. time: 25.60 min.; purity: 91%; MS (m/e): 637 (MH⁺).

7.3.688 N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(ethoxycarbonylamino)-2,4-pyrimidinetriamine (R926586)

In like manner to the preparation of N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(phenylaminocarbonylamino)-2,4-pyrimidinediamine, the reaction of 5-amino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine with ethoxycarbonyl isocyanate gave N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-

(ethoxycarbonylaminoacarbonylamino)-2,4-pyrimidinediamine. LCMS: ret. time: 26.79 min.; purity: 88%; MS (m/e): 669 (MH⁺).

7.3.689 N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(ethoxycarbonylmethyleneaminocarbonylamino)-2,4-pyrimidinediamine (R926587)

In like manner to the preparation of N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(phenylaminocarbonylamino)-2,4-pyrimidinediamine, the reaction of 5-amino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine with ethylacetyl isocyanate gave N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(ethoxycarbonylmethyleneaminocarbonylamino)-2,4-pyrimidinediamine. LCMS: ret. time: 25.76 min.; purity: 96%; MS (m/e): 683 (MH⁺).

7.3.690 N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(cyclopentylaminocarbonylamino)-2,4-pyrimidinediamine (R926588)

In like manner to the preparation of N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(phenylaminocarbonylamino)-2,4-pyrimidinediamine, the reaction of 5-amino-N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine with cyclopentyl isocyanate gave N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(cyclopentylaminoacarbonylamino)-2,4-pyrimidinediamine. LCMS: ret. time: 27.36 min.; purity: 83%; MS (m/e): 665 (MH⁺).

7.3.691 N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(chloroacetylaminocarbonylamino)-2,4-pyrimidinediamine (R926589)

In like manner to the preparation of N2,N4-bis(ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(N-phenylformyl-amino)-2,4-pyrimidinediamine, the reaction of N5-amino-N2,N4-bis(ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-2,4-pyrimidinediamine with chloroacetylformyl isocyanate gave N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-6-ethoxycarbonyl-5-(chloroacetylaminocarbonylamino)-2,4-pyrimidinediamine. LCMS: ret. time: 26.60 min.; purity: 100%; MS (m/e): 580 (MH⁺).

7.3.692 (R920669): N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-trifluoro-2,4-pyridinediamine

A mixture of 2,4-dichloro-5-trifluoromethylpyrimidine (416 mg, 1.9 mmol), 3,4-ethylenedioxyaniline (0.5 mL, 4.1 mmol), and concentrated HCl (0.1 mL) in 1:9 acetone/H₂O (10 mL) was heated to reflux. After 1 h, the reaction was complete as determined by TLC. The mixture was cooled to room temperature and EtOAc (30 mL) was added. The organic layer was washed with 2 N HCl (2 x 15 mL), water (15 mL), and dried (Na₂SO₄). The organic layer was filtered through a silica gel pad, washing the filter cake with EtOAc, and concentrated. The material was purified by chromatography (silica gel, 95:5 dichloromethane/ethyl acetate) to afford N2,N4-bis(3,4-ethylenedioxyphenyl)-5-trifluoro-2,4-pyridinediamine (380 mg, 44%): *R*_f 0.27 (silica gel, 9.5:0.5 dichloromethane/ethyl acetate); mp 141-143 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.25 (s, 1H), 7.07 (m, 2H), 6.99 (bs, 1H), 6.93-6.84 (m, 3H), 6.77-6.74 (m, 1H), 6.67 (bs, 1H), 4.29-4.24 (m, 8H); ¹³C NMR (75 MHz, CDCl₃) δ 161.2, 157.9, 155.8, 143.7, 132.6, 131.1, 117.5, 117.3, 114.4, 113.2, 110.3, 64.7, 64.5; IR (ATR) 3446 cm⁻¹; ESI MS *m/z* 447 [C₂₁H₁₇F₃N₄O₄ + H]⁺; HPLC (Method C) >99% (AUC), *t*_R = 8.5 min. Anal. Calcd for C₂₁H₁₇F₃N₄O₄: C, 56.50; H, 3.84; N, 12.55. Found: C, 56.46; H, 4.41; N, 12.57.

7.3.693 (R920668): N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-pyridyl)-2,4-pyrimidinediamine

A mixture of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (280 mg, 1 mmol), 3-aminopyridine (113 mg, 1.2 mmol), sodium *t*-butoxide (134 mg, 1.4 mmol), binap (38 mg, 0.06 mmol), and palladium(II)acetate (14 mg, 0.06 mmol) in 9 mL of toluene was purged with N₂ (3 cycles of alternating N₂ and vacuum). The mixture was heated to 80 °C (oil-bath temperature). After 24 h, the mixture was cooled to room temperature and EtOAc (30 mL) and of water (10 mL) was added. After stirring 15 min, the precipitate was collected by filtration. A ¹H NMR spectrum and ESI mass spectrum of the solid (150 mg) indicated the product (TLC analysis of the organic layer of the filtrate detected only starting materials). The crude product was slurried in 2 N HCl and the mixture was filtered. The filtrate was neutralized with 10% aqueous NaOH and concentrated. The material was slurried with MeOH and the solids removed by filtration. The concentrated material was slurried in CH₃CN and TFA was added to afford a solution. *N,N*-diisopropylethylamine was added to the solution and the solid was collected by filtration, washing with CH₃CN followed by Et₂O to afford N4-(3,4-ethylenedioxyphenyl)-

5-fluoro-N2-(3-pyridyl)-2,4-pyrimidinediamine (55 mg, 14%): R_f 0.42 (silica gel, 4:1:0.1:0.1 dichloromethane/ethyl acetate/methanol/concentrated ammonium hydroxide); mp 251-253 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 9.38 (s, 1H), 9.26 (s, 1H), 8.74 (s, 1H), 8.20-8.17 (m, 1H), 8.09-8.08 (m, 2H), 7.29-7.28 (m, 1H), 7.23-7.17 (m, 2H), 6.83-6.80 (m, 1H), 4.24 (m, 4H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 155.2, 149.8, 142.9, 141.6, 140.5, 140.0, 139.8, 139.7, 137.5, 132.1, 124.8, 123.0, 116.4, 115.1, 110.9, 64.1, 64.0; IR (ATR) 3264, 3195 cm^{-1} ; APCI MS m/z 340 [$\text{C}_{17}\text{H}_{14}\text{FN}_5\text{O}_2 + \text{H}$] $^+$. Anal. Calcd for $\text{C}_{17}\text{H}_{14}\text{FN}_5\text{O}_2 \cdot 0.5\text{H}_2\text{O}$: C, 58.70; H, 4.20; N, 20.13. Found: C, 58.71; H, 4.20; N, 19.51.

7.3.694 (R920664): N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(4-n-hexyloxyphenyl)-2,4-pyrimidindiamine

To a magnetically stirred solution of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (0.25 g, 0.89 mmol) in ethylene glycol (3.0 mL) under nitrogen at room temperature was added *N,N*-diisopropylethylamine (0.12 g, 0.89 mmol) followed by 4-hexyloxyaniline (0.27 g, 1.4 mmol). The reaction mixture was heated to 170 °C for 5.5 h, cooled to room temperature and partitioned between water (20 mL) and chloroform (20 mL). The aqueous layer was extracted with chloroform (20 mL) and the combined organic layers were dried (Na_2SO_4), filtered and concentrated in vacuo. The crude brown solid was purified by chromatography (silica gel, 2:1 hexanes/ethyl acetate) to afford N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(4-n-hexyloxyphenyl)-2,4-pyrimidindiamine (0.09 g, 23%) as a white solid: R_f 0.53 (silica gel, 4:1 chloroform/ethyl acetate); mp 115-117 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.90 (d, $J = 3.2$ Hz, 1H), 7.40 (d, $J = 8.9$ Hz, 2H), 7.29 (d, $J = 2.5$ Hz, 2H), 6.98 (d, $J = 8.8$ Hz, 1H), 6.88-6.82 (m, 3H), 6.61 (s, 1H), 4.29 (d, $J = 3.1$ Hz, 4H), 3.94 (t, $J = 6.6, 6.7$ Hz, 2H), 1.77 (m, 2H), 1.47 (m, 2H), 1.35 (m, 4H), 0.92 (m, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 156.3, 155.1, 150.3, 143.6, 142.7, 140.3, 140.0, 139.4, 133.0, 131.7, 121.9, 117.3, 115.0, 114.7, 110.8, 68.6, 64.6, 31.8, 29.5, 25.9, 22.8, 14.2; IR (ATR) 3357 cm^{-1} ; ESI MS m/z 439 [$\text{C}_{24}\text{H}_{27}\text{FN}_4\text{O}_3 + \text{H}$] $^+$; HPLC (Method B) 98.5% (AUC), $t_R = 7.9$ min. Anal. Calcd for $\text{C}_{24}\text{H}_{27}\text{FN}_4\text{O}_3$: C, 65.74; H, 6.21; N, 12.78. Found: C, 65.34; H, 6.19; N, 12.96.

7.3.695 (R920666): N2-(4-n-Butyloxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine

To a magnetically stirred solution of 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine (0.25 g, 0.89 mmol) in ethylene glycol (3.0 mL) under nitrogen at

room temperature was added *N,N*-diisopropylethylamine (0.12 g, 0.89 mmol) followed by 4-butoxyaniline (0.18 g, 1.1 mmol). The reaction mixture was heated to 185 °C for 5 h, cooled to room temperature, and partitioned between water (20 mL) and ethyl acetate (20 mL). The aqueous layer was extracted with ethyl acetate (20 mL) and the combined organic layers were dried (Na₂SO₄), filtered and concentrated in vacuo. The crude brown solid was purified by chromatography (silica gel, 2:1 hexanes/ethyl acetate) to afford N2-(4-n-Butyloxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyridinediamine (0.18 g, 49%) as a tan solid: *R*_f 0.66 (silica gel, 4:1 chloroform/ethyl acetate); mp 133-135 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.89 (d, *J* = 3.2 Hz, 1H), 7.39 (d, *J* = 8.9 Hz, 2H), 7.28 (d, *J* = 2.5 Hz, 1H), 6.95 (dd, *J* = 8.7, 2.5 Hz, 1H) 6.90-6.81 (m, 4H), 6.60 (d, *J* = 2.4 Hz, 1H), 4.27 (s, 4H), 3.94 (t, *J* = 6.5 Hz, 2H), 1.80-1.71 (m, 2H), 1.55-1.42 (m, 2H), 0.97 (t, *J* = 7.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 156.3, 155.1, 150.4, 143.6, 142.7, 140.3, 140.0, 139.4, 133.0, 131.7, 121.9, 117.3, 115.0, 114.7, 110.8, 68.2, 64.7, 64.5, 31.6, 19.4, 14.0; IR (ATR) 3356 cm⁻¹; ESI MS *m/z* 411 [C₂₂H₂₃FN₄O₃ + H]⁺; HPLC (Method A) >99% (AUC), *t*_R = 17.3 min. Anal. Calcd for C₂₂H₂₃FN₄O₃: C, 64.38; H, 5.65; N, 13.65. Found: C, 62.64; H, 5.59; N, 13.15.

7.3.696 (R920670): N4-(4-ethyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine

To a solution of 2-chloro-N4-(4-ethyloxyphenyl)-5-fluoro-4-pyrimidineamine (0.25 g, 0.93 mmol) in ethylene glycol (3 mL) under nitrogen at room temperature was added *i*-Pr₂EtN, 0.93 mmol) followed by 3,4-ethylenedioxyaniline (0.17 g, 1.12 mmol). The reaction mixture was heated to 200 °C for 5 h and then cooled to room temperature. The mixture was partitioned between H₂O (20 mL) and EtOAc (20 mL) and the aqueous layer was extracted with EtOAc (20 mL). The combined organic layers were dried (Na₂SO₄), filtered, and concentrated in vacuo. The crude brown solid was purified by chromatography (2:1 CHCl₃/EtOAc) to afford N4-(4-ethyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (0.21 g, 60%) as a tan solid: *R*_f 0.42 (4:1 CHCl₃/EtOAc); mp 163.8-167.2 °C (DSC); ¹H NMR (300 MHz, CDCl₃) δ 7.89 (d, *J* = 2.8 Hz, 1H), 7.50-7.45 (m, 2H), 7.17 (d, *J* = 2.5 Hz, 1H), 6.92-6.86 (m, 3H), 6.80-6.75 (m, 2H), 6.64 (bs, 1H), 4.26-4.21 (m, 4H), 4.03 (q, *J* = 7.0, 2H), 1.42 (t, *J* = 6.9 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 156.1, 150.6, 143.6, 142.8, 140.3, 140.0, 139.5, 139.3, 134.0, 130.8, 123.2, 117.2, 115.1, 113.6, 109.4, 64.6, 64.0, 15.1; IR (ATR) 3403 cm⁻¹; ESI MS *m/z* 383 [C₂₀H₁₉FN₄O₃

+ H]⁺; HPLC (Method A) 98.1% (AUC), *t*_R = 12.0 min. Anal. Calcd for C₂₀H₁₉FN₄O₃: C, 62.82; H, 5.01; N, 14.65. Found: C, 62.06; H, 5.01; N, 14.35.

7.3.697 (R920671): N4-(4-n-Butyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyridinediamine

5 In like manner to the preparation of N4-(4-ethyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(4-n-butyloxyphenyl)-5-fluoro-4-pyrimidineamine with 3,4-ethylenedioxyaniline gave N4-(4-n-butyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyridinediamine. The crude product was purified by chromatography (2:1 CHCl₃/EtOAc) ; (0.17 g, 52%) as a tan solid:
 10 *R*_f 0.51 (4:1 CHCl₃/EtOAc); mp 149.6-151.4 °C (DSC); ¹H NMR (300 MHz, CDCl₃) δ 7.88 (d, *J* = 3.4 Hz, 1H), 7.47 (d, *J* = 8.8 Hz, 2H), 7.18 (d, *J* = 2.4 Hz, 1H), 6.91-6.86 (m, 3H), 6.78-6.75 (m, 2H), 6.62 (bs, 1H), 4.26-4.22 (m, 4H), 3.96 (t, *J* = 6.5, 2H), 1.82-1.73 (m, 2H), 1.56-1.44 (m, 2H), 0.98 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 156.1, 150.8, 143.6, 142.8, 140.2, 139.9, 139.5, 139.2, 133.9, 130.7, 123.1, 117.1, 115.0, 113.5, 109.4,
 15 68.2, 64.6, 31.6, 19.4, 14.0; IR (ATR) 3365 cm⁻¹; ESI MS *m/z* 411 [C₂₂H₂₃FN₄O₃ + H]⁺; HPLC (Method A) 99.0% (AUC), *t*_R = 13.2 min. Anal. Calcd for C₂₂H₂₃FN₄O₃: C, 64.38; H, 5.65; N, 13.65. Found: C, 63.63; H, 5.60; N, 13.38.

7.3.698 (R920672): N4-(4-n-Hexyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyridinediamine

20 In like manner to the preparation of N4-(4-ethyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2-chloro-N4-(4-n-hexyloxyphenyl)-5-fluoro-4-pyrimidineamine with 3,4-ethylenedioxyaniline gave N4-(4-n-hexyloxyphenyl)-N2-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyridinediamine. The crude product was purified by chromatography (2:1 CHCl₃/EtOAc) (0.22 g, 69%) as a tan solid: *R*_f
 25 0.54 (4:1 CHCl₃/EtOAc); mp 124.0-125.2 °C (DSC); ¹H NMR (300 MHz, CDCl₃) δ 7.88 (d, *J* = 3.2 Hz, 1H), 7.47 (d, *J* = 8.9 Hz, 2H), 7.18 (d, *J* = 2.4 Hz, 1H), 6.91-6.86 (m, 3H), 6.78-6.74 (m, 2H), 6.62 (d, *J* = 1.8 Hz, 1H), 4.26-4.22 (m, 4H), 3.96 (t, *J* = 6.5, 2H), 1.83-1.74 (m, 2H), 1.51-1.42 (m, 2H), 1.36-1.32 (m, 4H), 0.93-0.89 (t, *J* = 6.7 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 156.1, 150.5, 143.5, 143.0, 142.8, 140.2, 139.9, 139.5, 139.2, 133.9,
 30 130.7, 123.1, 117.1, 115.0, 113.5, 109.3, 68.5, 64.7, 64.5, 31.8, 29.5, 25.9, 22.8, 14.2; IR (ATR) 3378 cm⁻¹; ESI MS *m/z* 439 [C₂₄H₂₇FN₄O₃ + H]⁺; HPLC (Method A) >99% (AUC),

$t_R = 14.6$ min. Anal. Calcd for $C_{24}H_{27}FN_4O_3$: C, 65.74; H, 6.21; N, 12.78. Found: C, 65.52; H, 6.23; N, 12.66.

7.3.699 (R920818): 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

5 To a mixture of 4-amino-[(1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene (1.2 g, 6.2 mmol), 1-propanol (40 mL) and trifluoroacetic acid (1 mL) was added 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyridineamine (1.5 g, 6.2 mmol). The mixture was heated at 110 °C for 17 h and then cooled to room temperature. The purple solid that formed was collected by filtration, washing with 1-propanol (30 mL) to afford 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (1.6 g, 65%) as an off-white solid: R_f 0.55 (6:3:1 $CHCl_3/CH_3OH/NH_4OH$); mp (DSC) 191.2-193.7 °C, 257.2-260.0 °C, 344.7-345.2 °C; ^1H NMR (300 MHz, $DMSO-d_6$) δ 9.39 (s, 1H), 9.21 (s, 1H), 9.10 (s, 1H), 8.04 (d, $J = 3.8$ Hz, 1H), 7.59 (d, $J = 9.1$ Hz, 2H), 7.38 (s, 1H), 7.23 (t, $J = 8.1$ Hz, 1H), 7.17 (d, $J = 1.8$ Hz, 1H), 7.05 (t, $J = 8.1$ Hz, 1H), 6.93 (d, $J = 9.1$ Hz, 2H), 6.50 (dd, $J = 1.8, 8.1$ Hz, 1H), 5.40 (s, 2H); ^{13}C NMR (75 MHz, $DMSO-d_6$) δ 157.3, 155.3, 153.5, 151.9, 149.8, 149.7, 141.0 (d, $J_{C-F} = 150.0$ Hz), 139.7, 138.7, 135.0, 128.9, 120.2, 114.8, 110.3, 108.7, 59.6; IR (ATR) 3338, 2923, 2581, 1724, 1661, 1580, 1557 cm^{-1} ; ESI MS m/z 395 [$C_{18}H_{15}FN_8O_2 + H$] $^+$; HPLC (Method A) 96.5% (AUC), $t_R = 6.9$ min.

7.3.700 (R920819): N4-(3-Hydroxyphenyl)-N2-[4-(1H,1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

20 To a mixture of 4-amino-[(1H,1,2,3,4-tetrazolyl)methyleneoxy]benzene (0.1 g, 0.5 mmol), 1-propanol (2 mL) and trifluoroacetic acid (0.2 mL) was added 2-chloro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (0.1 g, 0.5 mmol). The mixture was heated at 110 °C for 17 h and then cooled to room temperature. The purple solid that formed was collected by filtration, washing with 1-propanol (5 mL) to afford N4-(3-hydroxyphenyl)-N2-[4-(1H,1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (59.4 mg, 30%) as an off-white solid: R_f 0.51 (6:3:1 $CHCl_3/CH_3OH/NH_4OH$); mp 292-295 °C dec; ^1H NMR (300 MHz, $DMSO-d_6$) δ 9.34 (s, 2H), 9.13 (s, 1H), 7.95 (d, $J = 5.8$ Hz, 1H), 7.64 (d, $J = 8.9$ Hz, 2H), 7.39 (s, 1H), 7.19 (t, $J = 8.1$ Hz, 1H), 7.05 (t, $J = 8.1$ Hz, 1H), 6.96 (d, $J = 8.9$ Hz, 2H), 6.43 (dd, $J = 1.4, 8.1$ Hz, 1H), 6.20 (d, $J = 5.8$ Hz, 1H), 5.40 (s, 2H); ^{13}C NMR (75 MHz, $DMSO-d_6$) δ 160.4, 158.5, 157.5, 154.0, 153.7, 152.2, 140.6, 134.4, 129.1, 120.9,

114.7, 111.0, 109.5, 107.2, 98.4, 59.6; IR (ATR) 3321, 2920, 2581, 1649, 1605, 1487 cm^{-1} ; ESI MS m/z 377 [$\text{C}_{18}\text{H}_{16}\text{N}_8\text{O}_2 + \text{H}$] $^+$; HPLC (Method A) 97.6% (AUC), $t_R = 7.6$ min.

7.3.701 (R920820): N4-(3-Hydroxyphenyl)-5-methyl-N2-[4-(1H, 1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine

To a mixture of 4-amino-[(1H,1,2,3,4-tetrazolyl)methyleneoxy]benzene (0.2 g, 0.9 mmol), 1-propanol (4 mL) and trifluoroacetic acid (0.2 mL) was added 2-chloro-N4-(3-hydroxyphenyl)-5-methyl-4-pyrimidineamine (0.2 g, 0.9 mmol). The mixture was heated at 110 °C for 17 h and then cooled to room temperature. The purple solid that formed was collected by filtration, washing with 1-propanol (10 mL) to afford N4-(3-hydroxyphenyl)-5-methyl-N2-[4-(1H,1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine (0.3 g, 89%) as an off-white solid: R_f 0.44 (6:3:1 $\text{CHCl}_3/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$); mp (DSC) 255.3-262.4 °C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 10.32 (s, 1H), 9.65 (s, 2H), 7.85 (s, 1H), 7.38 (d, $J = 10.5$ Hz, 2H), 7.17 (s, 1H), 7.12 (t, $J = 7.9$ Hz, 1H), 7.06 (s, 1H), 6.90 (d, $J = 10.5$ Hz, 2H), 6.68 (d, $J = 7.9$ Hz, 1H), 5.45 (s, 2H), 2.14 (s, 3H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 161.6, 157.9, 154.5, 153.7, 151.2, 140.4, 138.2, 130.1, 129.4, 123.3, 115.9, 115.4, 113.5, 112.4, 107.5, 59.8, 13.7; IR (ATR) 3214, 3051, 2157, 1632, 1596, 1547 cm^{-1} ; ESI MS m/z 391 [$\text{C}_{19}\text{H}_{18}\text{N}_8\text{O}_2 + \text{H}$] $^+$; HPLC (Method A) >99% (AUC), $t_R = 7.9$ min.

7.3.702 N4-(3-Benzyloxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine [NEED R NO.]

A mixture of N4-(3-benzyloxyphenyl)-2-chloro-4-pyrimidineamine (0.25 g, 0.82 mmol), 4-amino-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy-benzene (0.17 g, 0.82 mmol) and trifluoroacetic acid (0.2 mL) in 1-propanol (10 mL) was heated to 110 °C for 24 h. The reaction was concentrated to remove most of the 1-propanol, the crude product was preadsorbed onto silica gel using 95:5 methylene chloride /methanol and purified by flash chromatography (95:5 methylene chloride /methanol) to give N4-(3-benzyloxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine as a tan solid (0.20 g, 52%): ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.00 (br s, 1H), 7.86 (d, $J = 6.1$ Hz, 1H), 7.53-7.20 (m, 13H), 7.14 (d, $J = 9.0$ Hz, 2H), 6.93 (d, $J = 6.1$ Hz, 1H), 6.13 (d, $J = 6.1$ Hz, 1H), 5.27 (s, 2H), 4.04 (s, 3H); ESI MS m/z 481 [$\text{C}_{26}\text{H}_{24}\text{N}_8\text{O}_2 + \text{H}$] $^+$.

7.3.703 (R920917): N4-(3-hydroxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine

A mixture of N4-(3-benzyloxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine (0.20 g, 0.42 mmol) and 5% Pd/C (0.10 g) in 14:1 ethanol/concentrated hydrochloric acid (40 mL) was at room temperature was shaken in a hydrogen atmosphere at 50 psi. After 3 h no further hydrogen uptake was observed. The reaction mixture was filtered through diatomaceous earth, the solids washed with 95:5 methylene chloride/methanol and the filtrate concentrated to afford N4-(3-hydroxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine (0.16 g, 95%) as a tan solid: R_f 0.23 (95:5 methylene chloride/methanol); mp (DSC) 207.1-212.8, 287.4-295.7 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 10.87 (br s, 1H), 10.81 (br s, 1H), 9.62 (br s, 1H), 8.08-8.06 (m, 1H), 7.72 (d, J = 9.0 Hz, 2H), 7.24 (br s, 1H), 7.20-7.00 (m, 3H), 6.61 (m, 2H), 6.46 (d, J = 6.0 Hz, 1H), 5.38 (s, 2H), 4.40 (s, 3H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 161.3, 160.1, 157.0, 154.3, 151.6, 141.7, 137.6, 129.1, 128.6, 123.4, 114.4, 111.9, 111.5, 108.3, 98.6, 59.6, 38.0; IR (ATR) 2975, 1639, 1602, 1521 cm^{-1} ; ESI MS m/z 391 [$\text{C}_{19}\text{H}_{18}\text{N}_8\text{O}_2 + \text{H}$] $^+$; HPLC (Method A) 94.9 % (AUC), t_R = 8.19 min.

7.3.704 N4-(3-Benzyloxyphenyl)-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine [NEED R NO.]

A mixture of N4-(3-benzyloxyphenyl)-2-chloro-4-pyrimidineamine (0.52 g, 1.69 mmol), 4-amino-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]benzene (0.34 g, 1.69 mmol) and trifluoroacetic acid (0.4 mL) in 1-propanol (10 mL) was heated to 110 °C for 24 h. The reaction was concentrated to remove most of the 1-propanol. The crude product was preadsorbed onto silica gel using 95:5 methylene chloride /methanol and purified by flash chromatography (95:5 methylene chloride /methanol) affording the requisite product N4-(3-benzyloxyphenyl)-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine as a tan solid (0.41 g, 51%): ^1H NMR (300 MHz, DMSO- d_6) δ 7.85 (d, J = 6.1 Hz, 1H), 7.49-7.04 (m, 14H), 6.93 (d, J = 9.0 Hz, 2H), 6.60-6.72 (m, 1H), 6.11 (d, J = 6.1 Hz, 1H), 5.14 (s, 2H), 4.34 (s, 3H); ESI MS m/z 481 [$\text{C}_{26}\text{H}_{24}\text{N}_8\text{O}_2 + \text{H}$] $^+$.

7.3.705 (R920910): N4-(3-Hydroxyphenyl)-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine

A mixture of N4-(3-benzyloxyphenyl)-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine (0.40 g, 0.42 mmol) and 5% Pd/C (0.10 g) in 14:1 ethanol/concentrated hydrochloric acid (40 mL) at room temperature was shaken in an

atmosphere of hydrogen at 50 psi. After 3 h no further hydrogen uptake was observed. The reaction mixture was filtered through diatomaceous earth, the solids washed with 95:5 methylene chloride/methanol and the filtrate concentrated to afford N4-(3-hydroxyphenyl)-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine (0.29 mg, 89%) as a beige solid: R_f 0.43 (95:5 methylene chloride/methanol); mp 140-152 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 10.24 (br s, 1H), 9.98 (br s, 1H), 9.52 (br s, 1H), 7.94 (d, J = 6.6 Hz, 2H), 7.54 (d, J = 8.8 Hz, 2H), 7.43 (s, 1H), 7.26 (s, 1H), 7.18-7.01 (m, 3H), 6.53, (d, J = 7.5 Hz, 1H), 6.37, (d, J = 6.6 Hz, 1H), 5.52 (s, 2H), 4.13 (s, 3H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 160.2, 157.2, 154.5, 153.0, 151.2, 146.8, 139.9, 131.8, 128.7, 122.3, 114.7, 111.4, 110.5, 107.5, 99.5, 59.5, 33.3; IR (ATR) 3042, 1578, 1504, 1459 cm^{-1} ; ESI MS m/z 391 [$\text{C}_{19}\text{H}_{18}\text{N}_8\text{O}_2 + \text{H}$] $^+$; HPLC (Method A) 95.8 % (AUC), t_R = 8.82 min.

7.3.706 (R920861): 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine

A mixture of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (0.22 g, 0.93 mmol, 4-amino-[(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene (0.19 g, 0.93 mmol) and trifluoroacetic acid (0.2 mL) in 1-propanol (8 mL) was heated to 110 °C for 24 h. The reaction was concentrated to remove most of the 1-propanol. The crude product was preadsorbed onto silica gel using 95:5 methylene chloride/methanol and purified by flash chromatography (95:5 methylene chloride /methanol) affording the requisite product 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine as a purple solid (0.18 g, 49%): R_f 0.47 (95:5 methylene chloride/methanol); mp 219-224 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 9.36 (s, 1H), 9.18 (s, 1H), 9.06 (s, 1H), 8.05 (d, J = 6.0 Hz, 1H), 7.60 (d, J = 9.0 Hz, 2H), 7.27 (d, J = 9.0 Hz, 1H), 7.09 (t, J = 8.0 Hz, 2H), 6.94 (d, J = 9.0 Hz, 2H), 6.49 (dd, J = 8.0, 2.1 Hz, 1H), 5.45 (s, 2H), 4.11 (s, 3H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 157.4, 155.5, 151.7, 151.6, 149.6, 149.5, 142.0, 142.0, 139.3 (d, $J_{\text{C-F}}$ = 127.5 Hz), 135.3, 128.9, 120.1, 114.9, 112.3, 110.3, 108.5, 58.5, 33.9; IR (ATR) 3278, 1586, 1542, 1508 cm^{-1} ; ESI MS m/z 409 [$\text{C}_{19}\text{H}_{17}\text{FN}_8\text{O}_2 + \text{H}$] $^+$; HPLC (Method A) 98.2 % (AUC), t_R = 7.69 min. Anal. Calcd for $\text{C}_{19}\text{H}_{17}\text{FN}_8\text{O}_2 \cdot 0.5 \text{H}_2\text{O}$: C, 54.74; H, 4.23; N, 26.88. Found: C, 54.55; H, 4.02; N, 26.62.

7.3.707 (R920860): 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine

A mixture of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine (0.31 g, 1.28 mmol), 4-amino-[(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene (0.26 g, 1.28 mmol) and trifluoroacetic acid (0.2 mL) in 1-propanol (8 mL) was heated at 110 °C for 24 h. The reaction was concentrated to remove most of the 1-propanol. The crude product was preadsorbed onto silica gel using 95:5 methylene chloride/methanol and purified by flash chromatography (95:5 methylene chloride /methanol) to give 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine as a purple solid (0.20 g, 37 %): R_f 0.63 (95:5 methylene chloride/methanol); mp 220-224 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 9.36 (s, 1H), 9.17 (s, 1H), 9.02 (s, 1H), 8.05 (d, J = 2.8 Hz, 1H), 7.57 (d, J = 9.1 Hz, 2H), 7.27 (d, J = 8.0 Hz, 1H), 7.10 (dt, J = 2.8, 8.0 Hz, 2H), 6.91 (d, J = 9.1 Hz, 2H), 6.49 (dd, J = 8.0, 2.8 Hz, 1H), 5.29 (s, 2H), 4.39 (s, 3H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 162.2, 157.4, 155.5, 152.1, 149.6, 149.5, 140.9 (d, $J_{\text{C-F}}$ = 142.0 Hz), 140.5, 140.2, 138.7, 134.8, 128.9, 120.2, 114.5, 112.2, 110.2, 108.5, 60.5, 38.5; IR (ATR) 3274, 1587, 1507 cm^{-1} ; ESI MS m/z 409 $[\text{C}_{19}\text{H}_{17}\text{FN}_8\text{O}_2 + \text{H}]^+$; HPLC (Method A) 97.2 % (AUC), t_R = 8.04 min. Anal. Calcd for $\text{C}_{19}\text{H}_{17}\text{FN}_8\text{O}_2$: C, 55.88; H, 4.20; N, 27.44. Found: C, 55.56; H, 4.10; N, 27.17.

7.3.708 (R920894): N4-(3-Hydroxyphenyl)-5-methyl-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine

A mixture of 2-chloro-N4-(3-hydroxyphenyl)-5-methyl-4-pyrimidineamine (0.20 g, 0.85 mmol), 4-amino-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene (0.17 g, 0.85 mmol) and trifluoroacetic acid (0.2 mL) in 1-propanol (8 mL) was heated at 110 °C for 24 h. The reaction was concentrated to remove most of the 1-propanol. The crude product was preadsorbed onto silica gel using 95:5 methylene chloride/methanol and purified by flash chromatography (95:5 methylene chloride /methanol) to give N4-(3-hydroxyphenyl)-5-methyl-N2-[4-(1-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine as a purple solid (0.18 g, 52%): R_f 0.61 (95:5 methylene chloride/methanol); mp 209-211 °C; ^1H NMR (300 MHz, DMSO- d_6) δ 9.30 (s, 1H), 8.82 (s, 1H), 8.13 (s, 1H), 7.83 (s, 1H), 7.60 (d, J = 9.0 Hz, 2H), 7.18-7.05 (m, 3H), 6.89 (d, J = 9.0 Hz, 2H), 6.48 (t, J = 7.1 Hz, 1H), 5.27 (s, 2H), 4.39 (s, 3H), 2.08 (s, 3H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 161.7, 158.6, 157.5, 156.7, 154.7, 151.2, 140.2, 134.6, 134.6, 128.1, 119.3, 114.0, 112.6, 109.4, 108.9,

104.7, 59.8, 38.0, 12.9; IR (ATR) 3003, 1602, 1581, 1531, 1507 cm^{-1} ; ESI MS m/z 405 $[\text{C}_{20}\text{H}_{20}\text{N}_8\text{O}_2 + \text{H}]^+$; HPLC (Method A) 96.8 % (AUC), $t_R = 8.23$ min.

7.3.709 (R920893): N4-(3-Hydroxyphenyl)-5-methyl-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine

5

A mixture of 2-chloro-N4-(3-hydroxyphenyl)-5-methyl-4-pyrimidineamine (0.20 g, 0.85 mmol), 4-amino-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxy]-benzene (0.17 g, 0.85 mmol) and trifluoroacetic acid (0.2 mL) in 1-propanol (8 mL) was heated at 110 °C for 24 h. The reaction was concentrated to remove most of the 1-propanol. The crude product was preadsorbed onto silica gel using 95:5 methylene chloride/methanol and purified by flash chromatography (95:5 methylene chloride /methanol) to give N4-(3-Hydroxyphenyl)-5-methyl-N2-[4-(2-methyl-1,2,3,4-tetrazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidineamine as a purple solid (0.14 g, 42%): R_f 0.44 (95:5 methylene chloride/methanol); mp 219-221 °C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 9.32 (s, 1H), 8.85 (s, 1H), 8.13 (s, 1H), 7.85 (s, 1H), 7.64 (d, $J = 9.0$ Hz, 2H), 7.20-7.07 (m, 3H), 6.91 (d, $J = 9.0$ Hz, 2H), 6.50 (dd, $J = 8.0, 1.2$ Hz, 1H), 5.45 (s, 2H), 4.12 (s, 3H), 2.09 (s, 3H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 158.0, 157.0, 156.1, 154.3, 150.6, 150.0, 139.6, 134.6, 127.5, 118.6, 113.7, 112.0, 108.8, 108.2, 104.2, 57.4, 32.7, 12.3; IR (ATR) 3428, 1595, 1567, 1509 cm^{-1} ; ESI MS m/z 405 $[\text{C}_{20}\text{H}_{20}\text{N}_8\text{O}_2 + \text{H}]^+$; HPLC (Method A) 98.5 % (AUC), $t_R = 7.89$ min. Anal. Calcd for $\text{C}_{20}\text{H}_{20}\text{N}_8\text{O}_2 \cdot \text{H}_2\text{O}$: C, 57.00; H, 5.02; N, 26.59. Found: C, 56.86; H, 4.92; N, 26.50.

20

7.3.710 N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-(1,2,3,4-tetrazol-5-yl)-2,4-pyrimidinediamine (R925810)

In a manner similar to experiment #, N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-cyano-2,4-pyrimidinediamine and sodium azide were reacted to yield N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-(1,2,3,4-tetrazol-5-yl)-2,4-pyrimidinediamine. LCMS: ret. time: 25.8 min.; purity: 95%; MS: 535 (MH^+).

25

7.3.711 N2-[4-(N-Cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925838)

30

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-N2-(methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with cyclopropylmethylamine gave N2-[4-(N-

cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: MS (m/e): 478 (MH⁺).

7.3.712 5-Ethoxycarbonyl-N4-(3-hydroxyphenyl)-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R925839)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-N2-(methoxycarbonylmethyleneoxyphenyl)-2,4-pyridinediamine with methylamine hydrochloride gave 5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-N2-[4-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: MS (m/e): 438 (MH⁺).

7.3.713 N2-[4-(N-2,3-Dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925840)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-N2-(methoxycarbonylmethyleneoxyphenyl)-2,4-pyridinediamine with 3-amino-1,2-propanediol gave N2-[4-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-ethoxycarbonyl-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: MS (m/e): 498 (MH⁺).

7.3.714 N2,N4-Bis[4-[N-(3-methoxybenzylamino)carbonylmethyleneoxy]phenyl]-5-bromo-2,4-pyrimidinediamine (R925841)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N2,N4-bis[4-ethoxycarbonylmethyleneoxyphenyl]-5-bromo-2,4-pyrimidinediamine with 3-methoxybenzylamine gave N2,N4-bis[4-[N-(3-methoxybenzylamino)carbonylmethyleneoxy]phenyl]-5-bromo-2,4-pyrimidinediamine. LCMS: ret. time: 25.94 min.; purity: 95 %; MS (m/e): 727 (MH⁺).

7.3.715 5-Bromo-N4-[4-[(N-cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R925842)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-bromo-N4-(4-ethoxycarbonylmethyleneoxyphenyl)-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine with cyclopropylmethylamine gave 5-bromo-N4-[4-(N-cyclopropylmethylamino)carbonylmethyleneoxyphenyl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 20.63 min.; purity: 100 %; MS (m/e): 485 (MH⁺).

7.3.716 5-Bromo-N2-(3-hydroxyphenyl)-N4-[4-(N-3-methoxybenzylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R925843)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-bromo-N4-(4-ethoxycarbonylmethyleneoxyphenyl)-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine with 3-methoxybenzylamine gave 5-bromo-N2-(3-hydroxyphenyl)-N2-(3-hydroxyphenyl)-N4-[4-(N-3-methoxybenzylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 22.34 min.; purity: 90 %; MS (m/e): 551 (MH⁺).

7.3.717 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-carboxybenzofuran-5-yl)-2,4-pyrimidinediamine (R926698)

In a manner similar to the preparation of N4-(4-*tert*-butylphenyl)-5-fluoro-N2-(2,3-dihydro-2-carboxybenzofuran-5-yl)-2,4-pyrimidinediamine, N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine and LiOH were reacted to yield N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-carboxybenzofuran-5-yl)-2,4-pyrimidinediamine.

7.3.718 N2,N4-Bis(4-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926016)

In a manner similar to the preparation of N2-N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidineamine, the reaction of 2,4-dichloro-5-fluoropyrimidine with 4-trifluoromethylaniline gave N2,N4-bis(4-trifluoromethylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.06 (bs, 1H), 7.75 (d, 2H, J= 9 Hz), 7.67 (d, 2H, J= 9Hz), 7.63 (d, 2H, J= 9Hz), 7.54 (d, 2H, J= 9 Hz), 7.19 (bs, 1H), 6.96 (s, 1H); ¹⁹F NMR (CDCl₃): δ -17598 (s, 3F), -17676 (s, 3F), -46549 (s, 1F); HPLC: 85% pure.

7.3.719 N2-(3,4-Ethylenedioxyphenyl)-N4-(3,4-methylenedioxyphenylhydrazinyl)-5-fluoro-2-pyrimidineamine (R926406)

In a manner similar to the preparation of N2-(3-hydroxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro N4-(3,4-methylenedioxyphenylhydrazinyl)-4-pyrimidineamine with 3,4-ethylenedioxyaniline gave N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-(3,4-methylenedioxyphenylhydrazinyl)-2-pyrimidineamine. ¹H NMR (CD₃OD): δ 7.82 (d, 1H, J= 3.6 Hz), 7.52 (dd, 1H, J= 1.8 and 7.5 Hz), 7.40 (d, 1H, J= 1.2 Hz), 7.14 (d, 1H, J= 2.4 Hz), 6.92 (d, 1H, J= 8.4 Hz), 6.85 (dd, 1H, J= 2.1 and 8.7 Hz), 6.45 (d, 1H, J= 9Hz), 6.06 (s, 2H), 4.10 (s, 4H); LCMS: ret. time: 12.14 min.; purity: 88%; MS (m/e): 426 (MH⁺).

7.3.720 N2,N4-Bis(4-ethoxycarbonylmethylenedioxyphenyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine (R926566)

To a solution of 2,4-dichloro-5-nitropyrimidine (0.264 g, 1 mmol) in EtOAc (10 mL) at 0 °C was added diisopropylethyl amine (0.200 mL) followed by ethyl 4-aminophenoxy acetate (0.585 g, 3 mmol) and then shaken at room temperature for 2h. The reaction was quenched with water and extracted with EtOAc. The EtOAc extract was washed with 2N HCl and water. The solvent was evaporated and the residue was purified by crystallization using EtOAc/hexanes to afford N2,N4-bis(4-ethoxycarbonylmethylenedioxyphenyl)-6-ethoxycarbonyl-5-nitro-2,4-pyrimidinediamine (R926566). ¹H NMR (CDCl₃): 10.32 (s, 1H), 7.42 (s, 1H), 7.40 (d, 2H, J= 8.7 Hz), 7.32 (d, 2H, J= 8.7 Hz), 6.93 (d, 2H, J= 8.7 Hz), 6.82 (d, 2H, J= 8.7 Hz), 4.67 (s, 2H), 4.62 (s, 2H), 4.47 (q, 2H, J= 7.5 Hz), 4.30 (m, 4H), 1.42 (t, 3H, J= 6.9 Hz), 1.31 (m, 6H); LCMS: ret. time: 32.10 min.; purity: 100%; MS (m/e): 584 (MH⁺).

7.3.721 N2,N4-Bis[2-(methylthio)-1,3-benzothiaz-6-yl]-5-fluoro-2,4-pyrimidinediamine (R950202)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-fluoropyrimidine and 2-(methylthio)-1,3-benzothiazol-6-amine were reacted to prepare N2,N4-bis[2-(methylthio)-1,3-benzothiaz-6-yl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 24.98 min.; purity: 84.6%; MS (m/e): 486.80 (MH⁺).

7.3.722 N4-[3-(2-Hydroxyethyleamino)phenyl]-N2-[3-(N-(N-methyl)-piperazino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950240)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethylamino)phenyl]-2,4-pyrimidinediamine and N-methylpiperazine were reacted to give N4-[3-(2-hydroxyethylenoxy)phenyl]-N2-[3-(N-(N-methyl)-piperazino)carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 13.36 min.; purity: 97.6%; MS (m/e): 495.42 (MH⁺).

7.3.723 N4-[3-(2-Hydroxyethyleamino)phenyl]-N2-[3-(N-piperazino)-carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine (R950241)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-(carboxymethyleneaminophenyl)-5-fluoro-N4-[3-(2-hydroxyethyleneamino)phenyl]-2,4-pyrimidinediamine and piperazine were reacted to give N4-[3-(2-hydroxyethyleneamino)phenyl]-N2-[3-(N-piperazino)-carbonylmethyleneaminophenyl]-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 13.21 min.; purity: 100%; MS (m/e): 481.40 (MH⁺).

7.3.724 (±)-N4-(3-Aminophenyl)-5-fluoro-N2-(3-(3-carboxy-3-D,L-N-phtaloylamino)propylenecarbonylaminophenyl)-2,4-pyrimidinediamine (R950251)

N2,N4-Bis(4-aminophenyl)-5-fluoro-2,4-pyrimidinediamine and N-phtaloyl-DL-glutamic anhydride were reacted in DMF to give N4-(3-aminophenyl)-5-fluoro-N2-(3-(3-carboxy-3-D,L-N-phtaloylamino)propylenecarbonylaminophenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 19.41 min.; purity: 95.7%; MS (m/e): 569.98 (MH⁺).

7.3.725 (±)-N4-(3-Aminophenyl)-5-fluoro-N2-[3-(3-carboxy-3-amino)propylenecarbonylaminophenyl]-2,4-pyrimidinediamine (R950255)

(±)-N4-(3-Aminophenyl)-5-fluoro-N2-[3-(3-carboxy-3-D,L-N-phtaloylamino)propylenecarbonylaminophenyl]-2,4-pyrimidinediamine was reacted with hydrazine to give N4-(3-aminophenyl)-5-fluoro-N2-[3-(3-carboxy-3-

amino)propylenecarbonylaminophenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 11.98 min.; purity: 90.1%; MS (m/e): 440.3 (MH⁺).

7.3.726 5-Methoxycarbonyl-N2,N4-bis[4-(N-pyrrolidino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926559)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-ethoxycarbonyl-N2,N4-bis(4-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with pyrrolidine gave 5-methoxycarbonyl-N2,N4-bis[4-(N-pyrrolidino) methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. The ethyl ester at 5-position was exchanged to methyl ester in methanol as a solvent. MS (m/e): 575 (MH⁺).

7.3.727 N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyridinediamine (R925565)

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-fluoropyrimidine with ethyl 4-aminophenoxyacetate gave N2,N4-bis(4-ethoxycarbonylmethyleneoxyphenyl)-5-fluoro-2,4-pyridinediamine. MS (m/e): 485 (MH⁺).

7.3.728 N2-(3-Ethoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-N4-(3,4-tetrafluoroethylenedioxyphenyl)-2,4-pyrimidinediamine (R926799)

In a manner similar to the preparation of N2-(3-hydroxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidineamine, the reaction of ethyl 3-aminophenoxyacetate with 2-chloro-5-ethoxycarbonyl-N4-(3,4-tetrafluoroethylenedioxyphenyl)-4-pyrimidineamine gave N2-(3-ethoxycarbonylmethyleneoxyphenyl)-5-ethoxycarbonyl-N4-(3,4-tetrafluoroethylenedioxyphenyl)-2,4-pyrimidinediamine. MS (m/e): 567 (MH⁺).

7.3.729 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-[N-2-(D)-(+)-biotinylethylamino]carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926811)

To a solution of D-(+)-biotin and N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-2-hydroxyethylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in DMF at -20 °C was added diisopropylethylamine and the mixture was shaken for 10 minutes. To this mixture was added benzotriazole-1-yl-oxy-tris(dimethylamino)-

phosphoniumhexafluorophosphate (BOP) and shaken at room temperature for 24 h. The reaction was quenched with water and extracted with ethyl acetate. The ethyl acetate extract was washed with aqueous solution of NaHCO₃ and finally with water. The residue obtained after the removal of solvent was purified by preparative TLC to obtain the desired

5 N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-[N-2-(D)-(+)-biotinylethylamino] carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: ret. time: 19.29 min.; purity: 99%; MS (m/e): 682 (M⁺).

7.3.730 5-Fluoro-N4-(3-hydroxyphenyl)-N2[2-(N-methyl-N-2-hydroxyethyl)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R926725)

10

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of 5-fluoro-N4-(3-hydroxyphenyl)-N2[2-methoxycarbonylbenzofuran-5-yl]-2,4-pyrimidinediamine with 2-(N-methyl)ethanolamine gave 5-fluoro-N4-(3-hydroxyphenyl)-

15 N2[2-(N-methyl-N-2-hydroxyethyl)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 14.87 min.; purity: 98%; MS: 438 (MH⁺).

7.3.731 N2,N4-Bis(3-ethoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926228)

20

In like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, the reaction of 2,4-dichloro-5-fluoropyrimidine and 3-ethoxycarbonylaniline gave N2,N4-bis(3-ethoxycarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 26.55 min.; purity: 100%; MS (m/e): 425 (MH⁺).

7.3.732 N2-(3-chloro-4-methylbenzyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R908696)

25

In a manner similar to the preparation of N2-(3-hydroxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidineamine, the reaction of 2-chloro-N4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with 3-chloro-4-methylbenzylamine gave N2-(3-chloro-4-methylbenzyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 25.38 min.; purity: 99 %; MS (m/e): 401 (MH⁺).

7.3.733 (+)-N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-phenylethyl)-2,4-pyrimidinediamine (R908697)

In a manner similar to the preparation of N2-(3-hydroxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidineamine, the reaction of 2-chloro-N4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine with (+)-2-aminoethylbenzene gave (+)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-phenylethyl)-2,4-pyrimidinediamine. LCMS: ret. time: 23.48 min.; purity: 99 %; MS (m/e): 367 (MH⁺).

7.3.734 N2-(3-Ethoxycarbonylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R925745)

In like manner to preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-ethoxycarbonylaniline gave N2-(3-ethoxycarbonylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.04 (bs, 1H), 7.94 (bs, 1H), 7.90 (bd, 1H), 7.68 (bd, 1H, J= 7.5 Hz), 7.35 (t, 1H, J= 8.1 Hz), 7.28 (d, 1H, J= 2.4 Hz), 7.07 (s, 1H), 6.93 (dd, 1H, J= 3 and 8.7 Hz), 6.83 (d, 1H, J= 9 Hz), 6.64 (bs, 1H), 4.36 (q, 2H, J= 7.2 Hz), 4.26 (s, 4H), 1.35 (t, 3H, J= 7.5 Hz); ¹⁹F NMR (CDCl₃): -47247; LCMS: ret. time: 15.88.; purity: 100%; MS (m/e): 411 (MH⁺).

7.3.735 N4-(3,4-Difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920394)

A solution of N-methyl 3-aminophenoxyacetamide (1 equivalent) and 2-chloro-N4-(3,4-difluorophenyl)-5-fluoro-4-pyrimidineamine (1.2 equivalents) in MeOH was shaken in a sealed tube at 100 °C for 24 hours for 24 h. Upon cooling to the room temperature, it was diluted with ethyl acetate. The resulting solid was filtered and washed with a mixture of ethyl acetate: n-hexanes (1:1; v/v) to obtain N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.05 (bs, 1H), 9.83 (bs, 1H), 8.23 (d, 1H, J= 2.7 Hz), 7.98 (m, 2H), 7.52 (m, 1H), 7.39 (m, 1H), 7.20 (m, 3H), 6.60 (m, 1H), 4.37 (s, 2H), 2.63 (d, 3H, J= 3.3 Hz); LCMS: purity: 94%; MS (m/e): 404 (MH⁺).

7.3.736 N4-(4-Chlorophenyl)-5-fluoro-N2-[3-(N-methylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920396)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(4-chlorophenyl)-5-fluoro-4-pyrimidineamine gave N4-(4-chlorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.21 (bs, 1H), 10.00 (bs, 1H), 8.26 (d, 1H, J= 4.8 Hz), 8.00 (bd, 1H, J= 4.2 Hz), 7.77 (dd, 2H, J= 2.1 and 7.6 Hz), 7.37 (dd, 2H, J= 2.1 and 7.6 Hz), 7.17 9m, 3H), 8.63 (dd, 1H, J= 1.8 and 8.1 Hz), 4.37 (s, 2H), 2.64 (d, 3H, 4.5 Hz); LCMS: purity: 92%; MS (m/e): 402 (MH⁺).

7.3.736.1 N4-(3,4-Dichlorophenyl)-5-fluoro-N2-[3-(N-methylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920397)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(3,4-dichlorophenyl)-5-fluoro-4-pyrimidineamine gave N4-(3,4-dichlorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.02 (bs, 1H), 9.76 (bs, 1H), 8.24 (d, 1H, J= 4.2 Hz), 8.08 (m, 1H), 7.97 (bd, 1H, J= 4.8 Hz), 7.77 (m, 1H), 7.55 (d, 1H, J= 8.7 Hz), 7.18 (m, 3H), 6.58 (m, 1H), 4.36 (s, 1H), 2.63 (d, 1H, J= 2.7 Hz); LCMS: purity: 91%; MS: 434 (MH⁺).

7.3.737 5-Fluoro-N4-(5-methylpyridin-2-yl)-N2-[3-(N-methylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920398)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(5-methylpyridin-2-yl)-4-pyrimidineamine gave 5-fluoro-N4-(5-methylpyridin-2-yl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 11.35 (bs, 1H), 10.70 (bs, 1H), 8.58 (s, 1H), 8.42 (d, 1H, J= 3.0 Hz), 8.12 (bd, 1H, J= 9.3 Hz), 8.03 (bd, 1H, J= 4.2 Hz), 7.82 (d, 1H, J= 8.7 Hz), 7.56 (s, 1H), 7.30 (bdd, 1H, J= 8.1

Hz), 7.19 (t, 1H, J= 8.1 Hz), 6.55 (dd, 1H, J= 1.8 and 8.1 Hz), 4.41 (s, 2H), 2.63 (d, 3H, J= 3.6 Hz), 2.36 (s, 3H); LCMS: purity: 99%; MS (m/e): 382 (M⁺).

7.3.738 5-Fluoro-N4-(6-methylpyridin-2-yl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920399)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(6-methylpyridin-2-yl)-4-pyrimidineamine gave 5-fluoro-N4-(6-methylpyridin-2-yl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.00 (bs, 1H), 9.60 (bs, 1H), 8.25 (s, 1H), 7.95 (m, 3H), 7.30 (s, 1H), 7.10 (m, 3H), 6.55 (d, 1H, J= 7.2 Hz), 4.40 (s, 2H), 2.62 (d, 3H, J= 3.6 Hz), 2.45 (s, 3H); LCMS: purity: 92%; MS (m/e): 383 (MH⁺).

7.3.739 N4-(5-Chloropyridin-2-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920405)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(5-chloropyridin-2-yl)-5-fluoro-4-pyrimidineamine gave N4-(5-chloropyridin-2-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.04 (bs, 1H), 9.53 (bs, 1H), 8.40 (d, 1H, J= 2.4 Hz), 8.22 (m, 2H), 7.88 (bd, 1H, J= 4.5 Hz), 7.86 (dd, 1H, J= 2.4 and 8.7 Hz), 7.40 (d, 1H, J= 1.8 Hz), 7.19 (m, 2H), 6.51 (bdd, 1H, J= 1.2 and 9 Hz), 4.38 (s, 2H), 2.64 (d, 3H, J= 3.3 Hz); LCMS: purity: 95%; MS (m/e): 403 (MH⁺).

7.3.740 N4-(6-Chloropyridin-3-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920406)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(6-chloropyridin-3-yl)-5-fluoro-4-pyrimidineamine gave N4-(6-chloropyridin-3-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆):

δ 9.72 (s, 1H), 9.38 (s, 1H), 8.93 (t, 1H, J= 3.0 Hz), 8.28 (m, 1H), 8.18 (d, 1H, J= 3.6 Hz), 7.95 (m, 1H), 7.45 (d, 1H, J= 8.7 Hz), 7.39 (m, 1H), 7.21 (m, 1H), 7.14 (t, 1H, J= 4.8 Hz), 6.50 (bdd, 1H, J= 7.8 Hz), 4.4 (s, 2H), 2.63 (d, 3H); LCMS: purity: 100%; MS (m/e): 403 (MH⁺).

5 **7.3.741 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(4-methylpyridin-2-yl)-2,4-pyrimidinediamine (R927016)**

10 In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(4-methylpyridin-2-yl)-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(4-methylpyridin-2-yl)-2,4-pyrimidinediamine. LCMS: purity: 95%; MS (m/e): 383 (MH⁺).

15 **7.3.742 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R920407)**

20 In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(3-trifluoromethylphenyl)-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.835 (bs, 1H), 9.54 (bs, 1H), 8.20 (d, 1H, J= 3.6 Hz), 7.94 (m, 2H), 7.78 (bs, 1H), 7.43 (t, 1H, J= 8.4 Hz), 7.25 (m, 2H), 7.15 (t, 1H, J= 7.5 Hz), 7.03 (bd, 1H, J= 9.3 Hz), 6.55 (bd, 1H, J= 7.5 Hz), 4.36 (s, 2H), 2.63 (d, 3H, J= 4.5 Hz); LCMS: purity: 91%; MS (m/e): 452 (MH⁺).

25 **7.3.743 N4-(3,4-Difluoromethylenedioxyphenyl)-5-fluoro- N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920408)**

30 In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine gave N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro- N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.91 (bs, 1H), 9.64 (bs, 1H), 8.19 (d, 1H, J= 3.9 Hz), 8.03 (s, 1H), 7.96 (bd, 1H, J= 4.8 Hz), 7.46 (m, 1H), 7.36 (d, 1H, J= 8.7 Hz), 7.27 (bs, 1H), 7.17 (m, 2H), 6.57 (bdd, 1H,

J= 7.2 Hz), 4.36 (s, 1H), 2.62 (d, 3H, J= 4.5 Hz); LCMS: purity: 96%; MS (m/e): 448 (MH⁺).

7.3.744 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920410)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine gave N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.08 (d, 1H, J= 5.4 Hz), 7.99 (d, 1H, J= 3.6 Hz), 7.67 (dd, 1H, J= 2.4 and 9.0 Hz), 7.40 (m, 3H), 7.06 (m, 2H), 6.92 (dd, 1H, J= 2.4 and 8.4 Hz), 4.44 (s, 2H), 2.80 (s, 3H); ¹⁹F NMR (CD₃OD): - 16973 and - 45983; LCMS: purity: 96%; MS (m/e): 486 (MH⁺).

7.3.745 N4-(4-Ethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926827)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(4-ethoxyphenyl)-5-fluoro-4-pyrimidineamine gave N4-(4-ethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 96%; MS: 412 (MH⁺).

7.3.746 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[4-methoxy-3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926828)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-amino-6-methoxyphenoxyacetamide with 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[4-methoxy-3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.83 (s, 1H), 7.80 (d, 1H, J= 4.2 Hz), 7.30 (d, 1H, J= 2.4 Hz), 7.23 (d, 1H, J= 2.4 Hz), 7.06 (m, 2H), 6.90 (d, 1H, J= 5.7 Hz), 6.73 (d, 1H, J= 5.2

Hz), 4.32 (s, 2H), 4.22 (s, 4H), 3.86 (s, 3H), 2.83 (s, 3H); LCMS: purity: 97%; MS (m/e): 455 (MH⁺).

7.3.747 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-methoxy-3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926829)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-amino-4-methoxyphenoxyacetamide with 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-methoxy-3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.86 (d, 1H, J= 4.2 Hz), 7.35 (d, 1H, J= 2.4 Hz), 7.19 (m, 1H), 7.12 (m, 3H), 6.93 (d, 1H, J= 8.7 Hz), 6.52 (m, 1H), 4.37 (s, 2H), 3.85 (s, 3H), 2.82 (s, 3H); ¹⁹F NMR (CD₃OD): - 47650; LCMS: purity: 100%; MS: 414 (MH⁺).

7.3.748 N4-(3-Chlorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926832)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of 3 N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(3-chlorophenyl)-5-fluoro-4-pyrimidineamine gave N4-(3-chlorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.12 (s, 1H), 9.93 (s, 1H), 8.27 (d, 1H, J= 4.2 Hz), 7.98 (d, 1H, J= 4.9 Hz), 7.85 (s, 1H), 7.73 (d, 1H, J= 8.1 Hz), 7.35 (t, 1H, J= 8.4 Hz), 7.19 (m, 3H), 6.62 (m, 1H), 4.36 (s, 2H), 2.63 (d, 3H, J= 4.2 Hz); LCMS: purity: 95%; MS: 402 (MH⁺).

7.3.749 5-Fluoro-N4-(3-methoxy-5-trifluoromethylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926833)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(3-methoxy-5-trifluoromethylphenyl)-4-pyrimidineamine gave 5-fluoro-N4-(3-methoxy-5-trifluoromethylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 95%; MS (m/e): 466 (MH⁺).

7.3.750 5-Fluoro-N4-(3-hydroxy-4-methoxyphenyl)-N2-[3-(N-methylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926834)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(3-hydroxy-4-methoxyphenyl)-4-pyrimidineamine gave 5-fluoro-N4-(3-hydroxy-4-methoxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.70 (bs, 2H), 8.12 (d, 1H, J= 4.8 Hz), 7.96 (m, 1H), 7.12 (m, 5H), 6.85 (d, 1H, J= 8.7 Hz), 6.57 (bd, 1H, J= 8.1 Hz), 4.35 (s, 2H), 3.74 (s, 3H), 2.63 (d, 3H, J= 4.5 Hz); LCMS: purity: 99%; MS (m/e): 414 (MH⁺).

7.3.751 5-Fluoro-N4-(4-methoxy-3-trifluoromethylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926835)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(4-methoxy-3-trifluoromethylphenyl)-4-pyrimidineamine gave 5-fluoro-N4-(4-methoxy-3-trifluoromethylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.9 Bs, 1H), 9.62 (bs, 1H), 8.17 (d, 1H, J= 4.2 Hz), 8.04 (bdd, 1H, J= 7.2 Hz), 7.82 (t, 1H, 2.7 Hz), 7.18 (m, 3H), 7.11 (t, 1H, J= 8.1 Hz), 6.55 (bd, 1H, J= 6.9 Hz); 4.33 (s, 2H), 3.86 (s, 3H), 2.61 (d, 3H, J= 4.0 Hz); LCMS: purity: 93%; MS: 466 (MH⁺).

7.3.752 5-Fluoro-N4-(4-fluoro-3-trifluoromethylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926838)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(4-fluoro-3-trifluoromethylphenyl)-4-pyrimidineamine gave 5-fluoro-N4-(4-fluoro-3-trifluoromethylphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.80 (s, 1H), 9.44 (s, 1H), 8.25 (m, 1H), 8.18 (d, 1H, J= 3.9 Hz), 8.00 (m, 1H), 7.97 (m, 1H), 7.47 (t, 1H, J= 9.6 Hz), 7.26 (s, 1H), 7.21

(m, 1H), 7.11 (t, 1H, J= 8.4 Hz), 6.51 (bd, 1H, J= 9.9 Hz), 4.34 (s, 2H), 2.62 (d, 3H, J= 4.8 Hz); LCMS: purity: 88%; MS: 454 (MH⁺).

7.3.753 N4-(3-Chloro-4-methylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926839)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(3-chloro-4-methylphenyl)-4-pyrimidineamine gave N4-(3-chloro-4-methylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.69 (s, 1H), 9.52 (s, 1H), 8.16 (d, 1H, J= 4.2 Hz), 7.96 (bs, 1H), 7.81 (d, 1H, J= 2.1 Hz), 7.67 (bd, 1H, J= 8.4 Hz), 7.26 (m, 3H), 7.15 (t, 1H, J= 8.1 Hz), 6.54 (bd, 1H, J= 7.2 Hz), 4.34 (s, 2H), 2.63 (d, 3H, J= 4.2 Hz), 2.27 (s, 3H); LCMS: purity: 80%; MS (m/e): 415 (M⁺).

7.3.754 N4-(2-Chloro-5-methylphenyl)-5-fluoro-N2-[3-(N-methylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926840)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(2-chloro-5-methylphenyl)-4-pyrimidineamine gave N4-(2-chloro-5-methylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.80 (bs, 2H), 8.21 (d, 1H, J= 4.8 Hz), 7.92 (d, 1H, J= 4.8 Hz), 7.46 (m, 1H), 7.31 (m, 2H), 7.04 (m, 2H), 6.53 (bd, 1H, J= 8.1 Hz), 4.30 (s, 1H), 2.18 (s, 3H); LCMS: purity: 93%; MS (m/e): 416 (MH⁺).

7.3.755 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-isopropylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926830)

The reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(ethoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with isopropylamine (5 equivalents) in the presence of diisopropylethylamine (5 equivalents) in MeOH in a sealed tube at 80 °C for 24 hours gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-isopropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-

d6): δ 9.15 (s, 1H), 8.04 (d, 1H, J = 4.2 Hz), 7.77 (d, 1H, J = 7.5 Hz), 7.28 (m, 4H), 7.08 (t, 1H, J = 8.1 Hz), 6.78 (d, 1H, J = 8.7 Hz), 6.45 (dd, 1H, J = 1.8 and 7.8 Hz), 4.30 (s, 2H), 4.20 (s, 4H), 3.92 (m, 1H), 1.06 (d, 6H, J = 6.6 Hz); LCMS: purity: 95%; MS (m/e): 454 (MH⁺).

5 **7.3.756 N2-[3-(N-Cyclopropylamino)carbonylmethyleneoxyphenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926848)**

10 In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-isopropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(ethoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine with cyclopropylamine gave 5-fluoro-N4-(3,4-ethylenedioxyphenyl)-N2-[3-(N-cyclopropylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.17 (bs, 2H), 8.05 (m, 2H), 7.27 (m, 4H), 7.08 (t, 1H, J = 8.1 Hz), 7.67 (d, 1H, J = 8.7 Hz), 6.42 (dd, 1H, J = 2.4 and 8.1 Hz), 4.3 (s, 2H), 4.2 (bs, 4H), 2.65 (m, 1H), 0.6 (m, 2H), 0.45 (m, 2H); LCMS: purity: 91%; MS (m/e): 452 (MH⁺).

15 **7.3.757 N4-(4-Cyano-3-methylphenyl)-5-fluoro-N2-[3-(N-methylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926851)**

20 In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(4-cyano-3-methylphenyl)-4-pyrimidineamine gave N4-(4-cyano-3-methylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.7 (s, 1H), 9.40 (s, 1H), 8.2 (s, 1H), 8.00-7.50 (m, 3H), 7.40-7.00 (m, 3H), 6.50 (bm, 1H), 4.35 (s, 2H), 2.60 (s, 3H), 2.35 (s, 3H); LCMS: purity: 91%; MS (m/e): 407 (MH⁺).

25 **7.3.758 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[3-(tetrazol-5-yl)phenyl]-2,4-pyrimidinediamine (R926855)**

30 In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-[3-(1H-tetrazol-5-yl)phenyl]-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[3-(tetrazol-5-yl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.04 (bs, 1H), 9.65 (bs, 1H), 8.35 (s, 1H), 8.23 (d, 1H, J = 3.9 Hz), 8.00 (bd, 1H, J = 6.6 Hz), 7.91 (bd,

J= 3.6 Hz), 7.77 (d, 1H, J= 8.1 Hz), 7.57 (t, 1H, J= 8.1 Hz), 7.23 (m, 2H), 6.95 (t, 1H, J= 8.4 Hz), 6.46 (bdd, 1H, J= 1.8 and 8.1 Hz), 4.22 (s, 2H), 2.62 (d, 3H, 4.2 Hz); LCMS: purity: 83%; MS (m/e): 436 (MH⁺).

7.3.759 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(N-methylphthalimido-4-yl)-2,4-pyrimidinediamine (R926856)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(N-methylphthalimido-4-yl)-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(N-methylphthalimido-4-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.95 (s, 1H), 9.44 (s, 1H), 8.29 (m, 1H), 8.25 (m, 1H), 8.18 (d, 1H, J= 1.8 Hz), 7.88 (bd, 1H, J= 4.5 Hz), 7.75 (d, 1H, J= 6.6 Hz), 7.38 (bs, 1H), 7.22 (bd, 1H, J= 8.1 Hz), 7.14 (t, 1H, J= 7.8 Hz), 6.50 (dd, 1H, J= 1.8 and 9.0 Hz), 4.28 (s, 2H), 2.99 (s, 3H), 2.60 (d, 3H, J= 4.5 Hz); LCMS: purity: 92%; MS (m/e): 451 (MH⁺).

7.3.760 N4-(2,5-Dimethoxy-4-chlorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926859)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with N4-(2,5-dimethoxy-4-chlorophenyl)-2-chloro-5-fluoro-4-pyrimidineamine gave N4-(2,5-dimethoxy-4-chlorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.05 (d, 1H, J= 5.4 Hz), 7.29 (s, 1H), 7.24 (t, 1H, J= 8.1 Hz), 7.18 (s, 1H), 7.02 (t, 1H, J= 2.1 Hz), 6.92 (dd, 1H, J= 1.8 and 8.1 Hz), 6.83 (dd, 1H, J= 2.4 and 8.4 Hz), 4.29 (s, 2H), 3.81 (s, 3H), 3.59 (s, 3H), 2.81 (s, 3H); LCMS: purity: 96%; MS (m/e): 460 (MH⁻); 462 (MH⁻).

7.3.761 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3-methoxycarbonyl-5-trifluoromethylphenyl)-2,4-pyrimidinediamine (R926862)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(3-methoxycarbonyl-5-

trifluoromethylphenyl)-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(3-methoxycarbonyl-5-trifluoromethylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.95 (s, 1H), 9.41 (s, 1H), 8.57 (s, 1H), 8.33 (s, 1H), 8.23 (d, 1H, J = 3 Hz), 7.83 (s and d, 2H), 7.22 (m, 2H), 7.02 (t, 1H, J = 8.7 Hz), 6.48 (1H, J = 2.4 and 7.5 Hz), 4.27 (s, 2H), 3.80 (s, 3H), 2.60 (d, 3H, J = 4.8 Hz); ¹⁹F NMR (DMSO-d₆): -17446; LCMS: purity: 94%; MS (m/z): 494 (MH⁺).

7.3.762 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[3-(2-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-2,4-pyrimidinediamine (R926870)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-[3-(2-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[3-(2-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-2,4-pyrimidinediamine. LCMS: purity: 86%; MS (m/e): 512 (MH⁺).

7.3.763 N4-[3-(2-(3-Chlorophenyl)-1,3,4-oxadiazol-5-yl)phenyl]-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926871)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-[3-(2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl)phenyl]-4-pyrimidineamine gave N4-[3-(2-(3-chlorophenyl)-1,3,4-oxadiazol-5-yl)phenyl]-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 100%; MS (m/e): 546 (MH⁺).

7.3.764 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[4-trifluoromethoxyphenyl]-2,4-pyrimidinediamine (R926879)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(4-trifluoromethoxyphenyl)-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[4-trifluoromethoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.05 (bs, 1H), 9.74 (bd, 1H, J = 1.5 Hz), 8.22 (d, 1H, J = 4.2 Hz), 7.99 (bd, 1H, J = 4.5 Hz), 7.86 (m,

2H), 7.32 (d, 2H, J= 8.1 Hz), 7.26 (s, 1H), 7.16 (m, 2H), 6.58 (m, 1H), 4.36 (s, 2H), 2.65 (bd, 3H); LCMS: purity: 92%; MS (m/e): 452 (MH⁺).

7.3.765 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[4-trifluoromethylphenyl]-2,4-pyrimidinediamine (R926880)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(4-trifluoromethylphenyl)-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-[4-trifluoromethylphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.10 (bs, 1H), 9.72 (d, 1H, J= 1.2 Hz), 8.26 (d, 1H, J= 4.2 Hz), 8.00 (m, 3H), 7.65 (d, 2H, J= 8.1 Hz), 7.31 (bs, 1H), 7.17 (d, 2H, J= 5.4 Hz), 6.59 (m, 1H), 4.36 (s, 2H), 2.62 (d, 3H, J= 4.8 Hz); LCMS: purity: 92%; MS (m/e): 436 (MH⁺).

7.3.766 N4-(4-Chloro-3-trifluoromethylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926881)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(4-chloro-3-trifluoromethylphenyl)-5-fluoro-4-pyrimidineamine gave N4-(4-chloro-3-trifluoromethylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.20 (bs, 1H), 9.81 (bs, 1H), 8.28 (d, 1H, J= 3.9 Hz), 8.23 (bdd, 1H, J= 8.7 Hz), 8.11 (d, 1H, J= 2.4 Hz), 7.98 (bd, 1H, J= 4.5 Hz), 7.65 (d, 1H, J= 8.7 Hz), 7.17 (m, 3H), 6.59 (m, 1H), 4.35 (s, 2H), 2.63 (d, 3H, J= 4.2 Hz); LCMS: purity: 87%; MS (m/e): 470 (MH⁺).

7.3.767 5-Fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(quinolin-6-yl)-2,4-pyrimidinediamine (R926883)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of 3 N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(quinolin-6-yl)-4-pyrimidineamine gave 5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-N4-(quinolin-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.17 (bs, 1H), 9.83 (s, 1H), 8.24 (d, 1H, J= 4.8 Hz), 8.17 (m, 1H), 7.94 (m, 2H), 7.86 (m, 1H), 7.39 (d, 1H, J= 9.3

Hz), 7.25 (s, 1H), 7.16 (m, 2H), 6.60 (m, 1H), 6.50 (d, 1H, J= 9.6 Hz), 4.32 (s, 2H), 2.60 (d, 3H, J= 3.6 Hz); LCMS: purity: 98%; MS 9m/e): 436 (MH⁺).

7.3.768 5-Fluoro-N4-(2-methoxypyridin-5-yl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R926886)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro-N4-(2-methoxypyridin-5-yl)-4-pyrimidineamine gave 5-fluoro-N4-(2-methoxypyridin-5-yl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.36 (bs, 1H), 9.19 (s, 1H), 8.59 (d, 1H, J= 3 Hz), 8.05 (m, 3H), 7.38 (m, 1H), 7.24 (bd, 1H, J= 8.1 Hz), 7.08 (t, 1H, J= 8.4 Hz), 6.79 (d, 1H, J= 8.7 Hz), 6.46 (dd, 1H, J= 2.4 and 7.8 Hz), 4.34 (s, 2H), 3.82 (s, 3H), 2.63 (d, 3H, J= 4.5 Hz); LCMS: purity: 95%; MS (m/e): 399 (MH⁺).

7.3.769 5-Fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R927023)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-5-fluoro- N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-4-pyrimidineamine gave 5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl] - 2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.65 (bs, 1H), 9.45 (bs, 1H), 8.55 (s, 1H), 8.12 (d, 1H, J= 3.6 Hz), 7.99 (m, 2H), 7.28 (m, 1H), 7.19 (m, 2H), 7.11 (t, 1H, J= 8.4 Hz), 6.81 (d, 1H, J= 8.7 Hz), 6.52 (m, 2H), 4.35 (s, 2H), 4.23 (t, 2H, J= 5.1 Hz), 3.69 (t, 2H, J= 4.5 Hz), 2.63 (d, 3H, J= 2.7 Hz); LCMS: purity: 95%; MS (m/e): 429 (MH⁺).

7.3.770 N4-(2,6-Dimethoxypyridin-3-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R920404)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-4-pyrimidineamine gave N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-N2-[3-(N-

methylamino)carbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.05 (d, 1H, J= 1.8 Hz), 8.62 (s, 1H), 8.01 (d, 1H, J= 3.6 Hz), 7.91 (bd, 1H, J= 4.8 Hz), 7.77 (m, 1H), 7.18 (m, 2H), 6.96 (t, 1H, J= 8.1 Hz), 6.40 (d, 2H, J= 8.1 Hz), 4.29 (s, 2H), 3.86 (s, 3H), 3.85 (s, 3H), 2.63 (d, 3H, J= 4.5 Hz); LCMS: purity: 86%; MS (m/e): 429 (MH⁺).

7.3.771 N4-(4-Chloro-3-methoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R927042)

In like manner to preparation of N4-(3,4-difluorophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine the reaction of N-methyl 3-aminophenoxyacetamide with 2-chloro-N4-(4-chloro-3-methoxyphenyl)-5-fluoro-4-pyrimidineamine gave N4-(4-chloro-3-methoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.89 (bs, 1H), 9.66 (bs, 1H), 8.20 (d, 1H, J= 4.2 Hz), 7.95 (bd, 1H), 7.48 (m, 2H), 7.33 (d, 1H, J= 9.1 Hz), 7.26 (bs, 1H), 7.17 (m, 2H), 6.57 (bd, 1H, J= 7.8 Hz), 4.34 (s, 2H), 3.72 (s, 3H), 2.62 (d, 3H); LCMS: purity: 97%; MS (m/e): 432 (MH⁺).

7.3.772 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R920411)

A solution of 2-chloro-N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine (1.1 equivalents) and 3-hydroxyaniline (1 equivalent) in a sealed tube was heated at 100 °C for 24 hours. The resulting solution was diluted with EtOAc and the solid obtained was filtered, washed with a mixture of EtOAc:n-hexanes (1:1; v/v), dried and analyzed to give N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine. ¹H NMR (CD₃OD): δ 8.02 (d, 1H, J= 5.1 Hz), 7.98 (d, 1H, J= 3.0 Hz), 7.72 (dd, 1H, J= 3.0 and 9.3 Hz), 7.42 (dd, 1H, J= 1.2 and 9.0 Hz), 7.22 (t, 1H, J= 8.4 Hz), 6.85 (m, 2H), 6.73 (dd, 1H, J= 2.4 and 8.7 Hz); ¹⁹F NMR (CD₃OD): - 16967 and - 46027; LCMS: purity: 97%; MS (m/e): 415 (MH⁺).

7.3.773 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[3-(3-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-2,4-pyrimidinediamine (R926866)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-5-fluoro-N4-[3-(2-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-4-pyrimidineamine gave

5-fluoro-N2-(3-hydroxyphenyl)-N4-[3-(2-phenyl-1,3,4-oxadiazol-5-yl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.72 (bs, 1H), 7.96 (bd, 3H), 7.85 (m, 2H), 7.56 (m, 4H), 7.14 (d, 1H, J= 2.1 Hz), 6.91 (m, 2H), 6.28 (dd, 1H, J= 1.8 and 6.9 Hz); LCMS: purity: 80%; MS (m/e): 441 (MH⁺).

5 **7.3.774 N4-(3,4-Difluoromethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926794)**

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine gave N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. LCMS: purity: 85%; MS (m/e): 377 (MH⁺).

7.3.775 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R926885)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-5-fluoro-N4-(3-trifluoromethoxyphenyl)-4-pyrimidineamine gave 5-fluoro-N2-(3-hydroxyphenyl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.99 (bs, 1H), 9.61 (bs, 1H), 8.21 (d, 1H, J= 4.2 Hz), 7.93 (bd, 1H, J= 7.5 Hz), 7.78 (s, 1H), 7.43 (t, 1H, J= 8.4 Hz), 7.03 (m, 4H), 6.43 (m, 1H); ¹⁹F NMR (DMSO-d₆): -16097; LCMS: purity: 85%; MS (m/e): 381 (MH⁺).

7.3.776 N4-(2,6-Dimethoxypyridin-3-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926887)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-4-pyrimidineamine gave N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.98 (bs, 2H), 8.20 (d, 1H, J= 5.4 Hz), 7.72 (m, 1H), 6.90 (t, 1H, J= 7.8 Hz), 6.81 (m, 2H), 6.42 (m, 2H), 3.88 (s, 3H), 3.86 (s, 3H); LCMS: purity: 94%; MS (m/e): 358 (MH⁺).

7.3.777 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(5-methylpyridin-2-yl)-2,4-pyrimidinediamine (R927017)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-5-fluoro-N4-(5-methylpyridin-2-yl)-4-pyrimidineamine gave 5-fluoro-N2-(3-hydroxyphenyl)-N4-(5-methylpyridin-2-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 11.39 (bs, 1H), 10.59 (bs, 1H), 8.58 (s, 1H), 8.41 (d, 1H, J = 3 Hz), 8.12 (d, 1H, J = 8.7 Hz), 7.82 (d, 1H, J = 8.7 Hz), 7.29 (s, 1H), 7.16 (d, 1H, J = 9 Hz), 7.05 (t, 1H, J = 8.4 Hz), 6.38 (dd, 1H, 1.2 and 6.9 Hz); LCMS: purity: 99%; MS (m/e): 312 (MH⁺).

7.3.778 N4-(6-Chloropyridin-3-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R927018)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-N4-(6-chloropyridin-3-yl)-5-fluoro-4-pyrimidineamine gave N4-(6-chloropyridin-3-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.10 (bs, 1H), 9.64 (bs, 1H), 8.85 (m, 1H), 8.30 (m, 2H), 8.22 (d, 1H, J = 4.2 Hz), 7.43 (d, 1H, J = 8.7 Hz), 7.01 (m, 3H), 6.42 (bd, 1H, J = 8.4 Hz); LCMS: purity: 93%; MS (m/e): 332 (MH⁺).

7.3.779 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(quinolin-6-yl)-2,4-pyrimidinediamine (R927019)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-5-fluoro-N4-(quinolin-6-yl)-4-pyrimidineamine gave 5-fluoro-N2-(3-hydroxyphenyl)-N4-(quinolin-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.50 (s, 1H), 10.14 (s, 1H), 8.29 (d, 1H, J = 4.8 Hz), 8.14 (d, 1H, J = 1.8 Hz), 7.96 (d, 1H, J = 9.3 Hz), 7.83 (dd, 1H, J = 2.4 and 9.0 Hz), 7.40 (d, 1H, J = 8.7 Hz), 7.04 (t, 1H, J = 8.1 Hz), 6.93 (m, 2H), 6.52 (m, 2H); LCMS: purity: 93%; MS (m/e): 365 (MH⁺).

7.3.780 N4-(5-Chloropyridin-2-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R927020)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-N4-(5-chloropyridin-2-yl)-5-fluoro-4-pyrimidineamine gave N4-(5-chloropyridin-

2-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.80 (bs, 1H), 9.77 (bs, 1H), 8.45 (bd, 1H), 8.26 (d, 1H, J= 3.9 Hz), 8.15 (d, 1H, J= 8.7 Hz), 7.85 (dd, 1H, J= 2.4 and 8.7 Hz), 7.06 (m, 3H), 6.43 (bd, 1H, J= 7.2 Hz); LCMS: purity: 97%; MS (m/e): 332 (MH⁺).

5 **7.3.781 N4-(4-Chloro-2,5-dimethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926860)**

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 3-hydroxyaniline with 2-chloro-N4-(4-chloro-2,5-dimethoxyphenyl)-5-fluoro-4-pyrimidineamine gave N4-(4-chloro-2,5-dimethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.96 (d, 1H, J= 4.8 Hz), 7.66 (s, 1H), 7.13 (s, 1H), 7.07 (t, 1H, J= 8.7 Hz), 8.86 (m, 2H), 6.57 (dd, 1H, J= 3.2 and 8.1 Hz), 3.48 (s, 3H), 3.66 (s, 3H); ¹⁹F NMR (CD₃OD): - 46968.

15 **7.3.782 N4-(4-Chlorophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine (R927026)**

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 5-amino-2-methoxycarbonylbenzofuran with 2-chloro-N4-(4-chlorophenyl)-5-fluoro-4-pyrimidineamine gave N4-(4-chlorophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.28 (bs, 1H), 10.18 (bs, 1H), 8.25 (d, 1H, J= 4.5 Hz), 7.96 (bs, 1H), 7.84 (m, 1H), 7.67 (m, 3H), 7.57 (m, 1H), 7.37 (bd, 2H, J= 9.0 Hz), 3.88 (s, 3H); LCMS: purity: 96%; MS (m/e): 413 (MH⁺).

25 **7.3.783 N4-(3,4-Dichlorophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine (R927027)**

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 5-amino-2-methoxycarbonylbenzofuran with 2-chloro-N4-(3,4-dichlorophenyl)-5-fluoro-4-pyrimidineamine gave N4-(3,4-dichlorophenyl)-5-fluoro-N2-(2-methoxycarbonylbenzofuran-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.70

(bs, 1H), 9.50 (bs, 1H), 8.20 (d, 1H, J= 4.5 Hz), 8.09 (m, 1H), 7.80 (m, 3H), 7.62 (m, 2H), 7.53 (m, 1H), 7.38 (m, 1H), 3.88 (s, 3H); LCMS: purity: 94%; MS (m/e): 448 (MH⁺).

7.3.784 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(3-methoxycarbonyl-5-trifluoromethylphenyl)-2,4-pyrimidinediamine (R926863)

5 In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-methoxycarbonyl-5-trifluoromethylaniline gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-methoxycarbonyl-5-trifluoromethylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.98 (s, 1H), 9.52 (s, 1H), 8.53 (s, 1H),
10 8.38 (s, 1H), 8.20 (d, 1H, J= 4.2 Hz), 7.69 (s, 1H), 7.27 (d, 1H, J= 8.1 Hz), 7.14 (s, 1H), 7.05 (t, 1H, 7.8 Hz), 6.49 (dd, 1H, J= 1.8 and 8.4 Hz), 3.80 (s, 3H); LCMS: purity: 82%; MS (m/e): 423 (MH⁺).

7.3.785 N2-(4-Chloro-2,5-dimethoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926857)

15 In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 4-chloro-2,5-dimethoxyaniline gave N2-(4-chloro-2,5-dimethoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.04 (d, 1H, J= 5.4 Hz), 7.46 (s, 1H), 7.17 (m, 2H), 7.03 (m, 2H), 6.72
20 (dd, 1H, J= 1.8 and 7.8 Hz), 3.85 (s, 3H), 3.52 (s, 3H); LCMS: purity: 98%; MS (m/e): 390 (MH⁺).

7.3.786 N2-(3-Bromo-5-trifluorophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926846)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-bromo-5-trifluoromethylaniline gave N2-(3-bromo-5-trifluorophenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H
25 NMR (DMSO-d₆): δ 9.70 (s, 1H), 9.36 (s, 1H), 9.34 (s, 1H), 8.31 (s, 1H), 8.18 (d, 1H, J= 3.6 Hz), 8.02 (s, 1H), 7.35 (s, 1H), 7.28 (bd, 1H, J= 7.2 Hz), 7.11 (t, 1H, J= 8.4 Hz), 7.02
30 (m, 1H), 6.49 (dd, 1H, J= 1.8 and 7.8 Hz); LCMS: purity: 94%; MS (m/e): 442 (MH⁺).

7.3.787 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(1H-pyrazol-3-yl)phenyl]-2,4-pyrimidinediamine (R926841)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-(1H-pyrazol-3-yl)aniline gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(1H-pyrazol-3-yl)phenyl]-2,4-pyrimidinediamine. LCMS: purity: 84%; MS 363 (MH⁺)

7.3.788 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(tetrazol-5-yl)phenyl]-2,4-pyrimidinediamine (R926842)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-(tetrazol-5-yl)aniline gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(tetrazol-5-yl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.05 (bs, 1H), 9.80 (bs, 1H), 8.27 (s, 1H), 8.23 (d, 1H, J= 3.3 Hz), 7.86 (d, 1H, J= 8.1 Hz) 7.65 (d, 1H, J= 6.9 Hz), 7.44 (t, 1H, J= 7.5 Hz), 7.19 (m, 2H), 6.93 (t, 1H, J= 7.5 Hz), 6.49 (dd, 1H, J= 2.4 and 8.1 Hz); LCMS: purity: 89%; MS (m/e): 364 (MH⁺).

7.3.789 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(1,3-oxazol-5-yl)phenyl]-2,4-pyrimidinediamine (R926831)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-(1,3-oxazol-5-yl)aniline gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-(1,3-oxazol-5-yl)phenyl)-2,4-pyrimidinediamine. LCMS: purity: 76%; MS (m/e): 364 (MH⁺).

7.3.790 N2-(3-Chloro-4-trifluoromethylphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinedimine (R926844)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-chloro-4-trifluoromethoxyaniline gave N2-(3-chloro-4-trifluoromethylphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinedimine. ¹H NMR (DMSO-d₆): δ 9.70 (bs, 1H), 9.48 (bs, 1H), 8.15 (bd, 1H, J= 3.6 Hz), 8.06 (s, 1H), 7.62 (dd, 1H, J= 2.4 and 9.3 Hz), 7.37 (d, 1H, J= 9.0 Hz), 7.20 9m, 1H), 7.11 (m, 3H), 6.53 (bd, 1H, J= 8.1 Hz); LCMS: purity: 93%; MS (m/e): 414 (MH⁺).

7.3.791 5-Fluoro-N4-(3,4-ethylenedioxyphenyl)-N2-[3-(tetrazol-5-yl)phenyl]-2,4-pyrimidinediamine (R926843)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine with 3-(tetrazol-5-yl)aniline gave 5-fluoro-N4-(3,4-ethylenedioxyphenyl)-N2-[3-(tetrazol-5-yl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.91 (s, 1H), 9.74 (s, 1H), 8.29 (s, 1H), 8.18 (d, 1H, J= 4.5 Hz), 7.76 (bdd, 1H, J= 1.5 and 8.1 Hz), 7.64 (d, 1H, J= 8.1 Hz), 7.46 (t, 1H, J= 8.1 Hz), 7.29 (m, 1H), 7.13 (dd, 1H, J= 2.4 and 8.7 Hz), 6.64 (d, 1H, J= 8.7 Hz), 4.11 (m, 4H); LCMS: purity: 91%; MS (m/e): 407 (MH⁺).

7.3.792 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(4-methoxy-2-methylphenyl)-2,4-pyrimidinediamine (R926845)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine with 4-methoxy-2-methylaniline gave N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(4-methoxy-2-methylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.30 (bs, 1H), 9.10 (bs, 1H), 8.22 (d, 1H, J= 5.1 Hz), 7.55 (m, 1H), 7.15 (m, 1H), 7.07 (m, 1H), 6.92 (m, 2H), 6.82 (d, 1H, J= 8.7 Hz), 4.22 (bs, 4H), 3.80 (s, 3H), 2.15 (s, 3H); LCMS: purity: 94%; MS (m/e): 383 (MH⁺).

7.3.793 N2-[5-(N-Aminocarbonylmethylene-2-oxo-1,3-oxazol-3(2H)-yl)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926847)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine with 2-[5-amino-2-oxo-1,3-benzoxazol-3(2H)-yl]acetamide gave N2-[5-(N-aminocarbonylmethylene-2-oxo-1,3-oxazol-3(2H)-yl)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.95 (d, 1H, J= 8.4 Hz), 7.32 (dd, 1H, J= 2.4 and 8.1 Hz), 7.24 (d, 1H, J= 2.4 Hz), 7.19 (m, 2H), 6.95 (dd, 1H, J= 2.7 and 9 Hz), 6.80 (d, 1H, J= 9 Hz), 4.51 (s, 2H), 4.21 (m, 4H).

7.3.794 N2-[3-(2-Ethoxycarbonylmethylene-1,3,4-oxadiazol-5-yl)phenyl-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926874)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine with 3-(2-ethoxycarbonylmethylene-1,3,4-oxadiazol-5-yl)aniline gave N2-[3-(2-ethoxycarbonylmethylene-1,3,4-oxadiazol-5-yl)phenyl-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.52 (s, 1H), 9.31 (s, 1H), 9.28 (s, 1H), 8.30 (s, 1H), 8.12 (d, 1H, J= 3.6 Hz), 8.00 (m, 1H), 7.49 (d, 1H, J= 7.5 Hz), 7.42 (d, 1H, J= 8.4 Hz), 7.30 (m, 1H), 7.12 (bs, 1H), 7.03 (t, 1H, J= 8.1 Hz), 6.46 (m, 1H), 4.21 (s, 2H), 4.15 (q, 2H, J= 6.9 Hz), 1.19 (t, 3H, J= 7.2 Hz); LCMS: purity: 90%; MS (m/e): 451 (MH⁺).

7.3.795 N2,N4-Bis(3-boronylphenyl)-5-fluoro-2,4-pyrimidinediamine (R926836)

A mixture of 2,4-dichloro-5-fluoro-pyrimidine (1 equivalents) and 3-aminophenylboronic acid (3 equivalents) in MeOH was heated in a sealed tube at 100 °C for 24 hours. The resulting mixture was cooled to room temperature, acidified with 2N HCl and the solid obtained was filtered, washed with water, dried and analyzed to give N2,N4-bis(3-boronylphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.40 (s, 1H), 10.07 (s, 1H), 8.25 (d, 8.4 Hz), 7.85 (s, 1H), 7.73 (d, 1H, J= 7.5 Hz), 7.63 (bt, 3H), 7.48 (d, 1H, J= 6.9 Hz), 7.30 (t, 1H, J= 8.4 Hz), 7.12 (t, 1H, J= 2.5 Hz); LCMS: purity: 85%; MS (m/e): 368 (MH⁺).

7.3.796 N2-(3-Boronylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926837)

In like manner to the preparation of N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidineamine, the reaction of 2-chloro-5-fluoro-N4-(3,4-ethylenedioxyphenyl)-4-pyrimidineamine with 3-aminophenylboronic acid gave N2-(3-boronylphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: purity: 99%; MS (m/e): 383 (MH⁺).

7.3.797 (+)-N4-(3,4-Difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine (R927030)

A mixture of equivalent amount of 2-chloro-N4-(3,4-difluorophenyl)-5-fluoro-4-pyrimidineamine and (+)-5-amino-2,3-dihydro-2-methoxycarbonylbenzofuran in MeOH was shaken in a sealed tube at 80 °C for 48 h, cooled to room temperature and diluted with a mixture of n-hexanes:EtOAc (1:1; v/v). The resulting solid formed was filtered, washed with a mixture of EtOAc:n-hexanes (1:1; v/v), dried and analyzed to give (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.21 (bs, 1H), 9.80 (bs, 1H), 8.20 (d, 1H, J= 4.8 Hz), 7.94 (bs, 1H), 7.43 (m, 3H), 7.15 (bd, 1H, J= 8.4 Hz), 6.81 (d, 1H, J= 8.1 Hz), 5.35 (dd, 1H, J= 6.0 and 6.3 Hz), 3.69 (s, 3H), 3.52 (dd, 1H, J= 10.5), 3.22 (dd, 1H, J= 9.0 and 6.0 Hz); LCMS: purity: 99%; MS (m/e): 417 (MH⁺).

7.3.798 (+)-N4-(4-Chlorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine (R927024)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine, the reaction of (+)-5-amino-2,3-dihydro-2-methoxycarbonylbenzofuran with 2-chloro-N4-(4-chlorophenyl)-5-fluoro-4-pyrimidineamine gave (+)-N4-(4-chlorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.29 (bs, 1H), 9.89 (bs, 1H), 8.21 (d, 1H, J= 4.8 Hz), 7.69 (m, 2H), 7.38 (m, 3H), 7.13 (bd, 1H, J= 8.1 Hz), 6.83 (d, 1H, J= 8.4 Hz), 5.36 (dd, 1H, J= 6.3 and 5.7 Hz), 3.70 (s, 3H), 3.52 (dd, 1H, J= 10.5 Hz), 3.20 (dd, 1H, J= 5.4 and 5.7 Hz); LCMS: purity: 98%; MS (m/e): 415 (MH⁺).

7.3.799 (+)-N4-(3,4-Dichlorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine (R927031)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine, the reaction of (+)-5-amino-2,3-dihydro-2-methoxycarbonylbenzofuran with 2-chloro-N4-(3,4-dichlorophenyl)-5-fluoro-4-pyrimidineamine gave (+)-N4-(3,4-dichlorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆):

δ 10.13 (bs, 1H), 9.70 (bs, 1H), 8.21 (d, 1H, $J = 4.8$ Hz), 8.04 (d, 1H, $J = 2.4$ Hz), 7.68 (m, 1H), 7.54 (d, 1H, $J = 9.0$ Hz), 7.37 (bs, 1H), 7.19 (m, 1H), 6.80 (d, 1H, $J = 8.7$ Hz), 5.35 (dd, 1H, $J = 6.6$ Hz), 3.69 (s, 3H), 3.53 (dd, 1H, $J = 10.5$ and 11.1 Hz), 3.21 (dd, 1H, $J = 6.0$ Hz); LCMS: purity: 100%; MS (m/e): 450 (MH^+).

5 **7.3.800 (+)-N2-(2,3-Dihydro-2-methoxycarbonylbenzofuran-5-yl)-N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-2,4-pyrimidinediamine (R927032)**

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine, the reaction of (+)-5-amino-2,3-dihydro-2-methoxycarbonylbenzofuran with 2-chloro-N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-4-pyrimidineamine gave (+)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-N4-(2,6-dimethoxypyridin-3-yl)-5-fluoro-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 10.03 (bs, 2H), 8.18 (d, 1H, $J = 4.8$ Hz), 7.68 (bd, 1H, $J = 8.1$ Hz), 7.27 (bs, 1H), 6.98 (bd, 1H, $J = 8.1$ Hz), 6.69 (d, 1H, $J = 8.7$ Hz), 6.44 (d, 1H, $J = 8.1$ Hz), 5.33 (dd, 1H, $J = 5.7$ Hz), 3.88 (s, 3H), 3.86 (s, 3H), 3.69 (s, 3H), 3.42 (dd, 1H, $J = 10.8$ and 11.1 Hz), 3.10 (dd, 1H, $J = 6.3$ and 6.6 Hz); LCMS: purity: 99%; MS (m/e): 442 (MH^+).

20 **7.3.801 (+)-N2-(2,3-Dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-5-fluoro-2,4-pyrimidinediamine (R927025)**

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine, the reaction of (+)-5-amino-2,3-dihydro-2-methoxycarbonylbenzofuran with 2-chloro-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-4-pyrimidineamine gave (+)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 10.10 (bs, 1H), 9.70 (bs, 1H), 8.46 (m, 1H), 8.13 (d, 1H, $J = 4.8$ Hz), 7.92 (m, 1H), 7.41 (bs, 1H), 7.12 (bdd, 1H, $J = 8.4$ Hz), 6.79 (m, 2H), 5.35 (dd, 1H, $J = 5.7$ and 6.0 Hz), 4.24 (t, 2H, $J = 5.1$ Hz), 3.70 (s, 3H), 3.69 (t, 2H, $J = 5.1$ Hz), 3.52 (dd, 1H, $J = 11.1$ Hz), 3.24 (dd, 1H, $J = 6.6$ Hz); LCMS: purity: 92%; MS (m/e): 442 (MH^+).


7.3.802 (+)-N2-(2,3-Dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-N4-(3-trifluorophenyl)-2,4-pyrimidinediamine (R927028)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine, the reaction of (+)-5-amino-2,3-dihydro-2-methoxycarbonylbenzofuran with 2-chloro-5-fluoro-N4-(3-trifluorophenyl)-4-pyrimidineamine gave (+)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-N4-(3-trifluorophenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.32 (bs, 1H), 9.90 (bs, 1H), 8.23 (d, 1H, J= 4.8 Hz), 7.80 (bd, 1H, J= 6.9 Hz), 7.73 (bs, 1H), 7.43 (t, 1H, J= 8.1 Hz), 7.36 (bs, 1H), 7.16 (m, 2H), 6.79 (d, 1H, J= 8.1 Hz), 5.33 (dd, 1H, J= 6.0 and 6.6 Hz), 3.69 (s, 3H), 3.51 (dd, 1H, J= 10.5 Hz), 3.20 (dd, 1H, J= 6.0 Hz); LCMS: purity: 98%; MS (m/e): 465 (MH⁺).

7.3.803 (+)-N2-(2,3-Dihydro-2-methoxycarbonylbenzofuran-5-yl)-N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R927029)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine, the reaction of (+)-5-amino-2,3-dihydro-2-methoxycarbonylbenzofuran with 2-chloro-N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine gave (+)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-N4-(3,4-difluoromethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.36 (bs, 1H), 9.93 (bs, 1H), 8.22 (d, 1H, J= 4.8 Hz), 7.91 (bs, 1H), 7.38 (m, 3H), 7.15 (bd, 1H, J= 8.7 Hz), 6.79 (d, 1H, J= 6.0 Hz), 5.33 (dd, 1H, J= 6.3 and 6.6 Hz), 3.69 (s, 3H), 3.50 (dd, 1H, J= 10.5 and 10.8 Hz), 3.22 (dd, 1H, J= 6.0 Hz); LCMS: purity: 100%; MS (m/e): 461 (MH⁺).

Esters were transformed to amides allowing to the scheme illustrated below:



7.3.804 (+)-N4-(3,4-Difluorophenyl)-5-fluoro-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine (R927035)

A mixture of (+)-N4-(3,4-difluorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine, methylamine Hydrogen

Chloride (5 equivalents) and diisopropylethylamine (5 equivalents) in MeOH was shaken in a sealed tube at 80 °C for 24 h. The resulting solution was diluted with water and the precipitate obtained was filtered, washed with water, dried and analyzed to afford (+)-N4-(3,4-difluorophenyl)-5-fluoro-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.46 (s, 1H), 9.07 (s, 1H), 8.05 (m, 3H), 7.48 (m, 2H), 7.35 (m, 1H), 7.22 (m, 1H), 6.72 (d, 1H, J= 8.1 Hz), 5.07 (dd, 1H, J= 6.6 and 6.3 Hz), 3.40 (dd, 1H), 3.15 (dd, 1H), 2.60 (d, 3H, J= 4.5 Hz); LCMS: purity: 98%; MS (m/e): 416 (MH⁺).

7.3.805 (+)-N4-(4-Chlorophenyl)-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-2,4-pyrimidinediamine (R927036)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-5-fluoro-N2-[2-(N-methylamino)carbonyl-2,3-dihydrobenzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of methyl amine Hydrogen Chloride with (+)-N4-(4-chlorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine gave (+)-N4-(4-chlorophenyl)-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.40 (s, 1H), 9.02 (s, 1H), 8.05 (m, 2H), 7.84 (dd, 2H, J= 2.7 and 9.3 Hz), 7.51 (bs, 1H), 7.32 (bd, 2H, J= 8.7 Hz), 7.23 (bd, 1H, J= 8.7 Hz), 6.72 (d, 1H, J= 8.7 Hz), 5.07 (dd, 1H, J= 6.0 and 6.3 Hz), 3.39 (dd, 1H), 3.17 (dd, 1H), 2.60 (d, 3H, J= 4.8 Hz); LCMS: purity: 99%; MS (m/e): 414 (MH⁺).

7.3.806 (+)-N4-(3,4-Dichlorophenyl)-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-2,4-pyrimidinediamine (R927037)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-5-fluoro-N2-[2-(N-methylamino)carbonyl-2,3-dihydrobenzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of methyl amine Hydrogen Chloride with (+)-N4-(3,4-dichlorophenyl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine gave (+)-N4-(3,4-dichlorophenyl)-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.52 (s, 1H), 9.09 (s, 1H), 8.08 (m, 3H), 7.76 (bd, 1H, J= 9.3 Hz), 7.50 (d, 1H, J= 9.0 Hz), 7.43 (bs, 1H), 7.24 (bd, 1H, J= 8.7 Hz), 6.73 (d, 1H, J= 8.1 Hz), 5.07 (dd, 1H, J= 6.3 and 6.6 Hz), 3.39 (dd, 1H, J= 10.5 Hz), 3.15 (dd, 1H, J= 6.3 Hz), 2.60 (d, 3H, J= 4.8 Hz); LCMS: purity: 99%; MS (m/e): 450 (MH⁺).

7.3.807 (+)-N4-(2,6-Dimethoxypyridin-3-yl)-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-2,4-pyrimidinediamine (R927038)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-5-fluoro-N2-[2-(N-methylamino)carbonyl-2,3-dihydrobenzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of methyl amine Hydrogen Chloride with (+)-N4-(2,6-dimethoxypyridin-3-yl)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-2,4-pyrimidinediamine gave (+)-N4-(2,6-dimethoxypyridin-3-yl)-N2-[2,3-dihydro-2-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.98 (d, 1H, J= 8.1 Hz), 7.81 (d, 1H, J= 3.6 Hz), 7.39 (bd, 1H, J= 2.4 Hz), 7.06 (dd, 1H, J= 2.4 and 8.7 Hz), 6.72 (d, 1H, J= 8.1 Hz), 6.31 (d, 1H, J= 8.7 Hz), 5.07 (dd, 1H, J= 6.3 Hz), 3.96 (s, 3H), 3.93 (s, 3H), 3.46 (dd, 1H, J= 7.8 and 10.5 Hz), 3.19 (dd, 1H, J= 5.7 and 6.3 Hz), 2.77 (d, 3H, J= 4.8 Hz); LCMS: purity: 98%; MS (m/e): 441 (MH⁺).

7.3.808 (+)-N2-[2,3-Dihydro-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-2,4-pyrimidinediamine (R927039)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-5-fluoro-N2-[2-(N-methylamino)carbonyl-2,3-dihydrobenzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of methyl amine Hydrogen Chloride with (+)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-2,4-pyrimidinediamine gave (+)-5-fluoro-N4-[2-(2-hydroxyethyleneoxy)pyridin-5-yl]-N2-[2-(N-methylamino)carbonyl-2,3-dihydrobenzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.26 (s, 1H), 8.99 (s, 1H), 8.50 (bd, 1H, J= 3.0 Hz), 8.02 (bd, 2H, J= 3.6 Hz), 7.94 (dd, 2H, J= 2.7 and 5.1 Hz), 7.52 (bs, 1H), 7.20 (bd, 1H, J= 8.7 Hz), 6.78 (d, 1H, J= 8.7 Hz), 6.67 (d, 1H, J= 8.7 Hz), 5.05 (dd, 1H, J= 6.3 and 6.6 Hz), 4.80 (t, 1H), 4.23 (t, 2H, J= 5.1 Hz), 3.69 (q, 2H, J= 5.4 Hz), 3.40 (dd, 1H), 3.15 (dd, 1H, J= 6.3 and 9.9 Hz), 2.60 (d, 3H, J= 4.5 Hz); LCMS: purity: 86%; MS (m/e): 441 (MH⁺).

7.3.809 (+)-N2-[2,3-Dihydro-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R927040)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-5-fluoro-N2-[2-(N-methylamino)carbonyl-2,3-dihydrobenzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of methyl amine Hydrogen Chloride with (+)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine gave (+)-N2-[2,3-

dihydro-(N-methylamino)carbonylbenzofuran-5-yl]-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. LCMS: purity: 94%; MS (m/e): 464 (MH⁺).

7.3.810 (+)-N2-[2,3-Dihydro-(N-methylamino)carbonylbenzofuran-5-yl]-N4-(3,4-difluoromethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R927041)

In like manner to the preparation of (+)-N4-(3,4-difluorophenyl)-5-fluoro-N2-[2-(N-methylamino)carbonyl-2,3-dihydrobenzofuran-5-yl]-2,4-pyrimidinediamine, the reaction of methyl amine Hydrogen Chloride with (+)-N2-(2,3-dihydro-2-methoxycarbonylbenzofuran-5-yl)-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine gave (+)-N2-[2,3-dihydro-(N-methylamino)carbonylbenzofuran-5-yl]-N4-(3,4-difluoromethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.46 (s, 1H), 9.05 (s, 1H), 8.05 (m, 3H), 7.43 (m, 2H), 7.31 (d, 1H, J= 8.7 Hz), 7.23 (bd, 1H, J= 7.5 Hz), 6.70 (d, 1H, J= 9.0 Hz), 5.04 (dd, 1H, J= 6.6 Hz), 3.40 (dd, 1H), 3.14 (dd, 1H, J= 5.7 and 6.6 Hz), 2.60 (d, 3H, J= 3.9 Hz); LCMS: purity: 94%; MS (m/e): 460 (MH⁺).

7.3.811 N2-(4-Carboxymethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926238)

The reaction of N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine with LiOH in THF:H₂O at room temperature gave N2-(carboxymethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.16 (d, 1H, J= 4.8 Hz), 7.37 (bd, 2H, J= 9 Hz), 7.25 (d, 1H, J= 3Hz), 7.08 (m, 1H), 6.83 (m, 3H), 4.64 (s, 2H), 4.23 (s, 4H); LCMS: ret. time: 19.15 min.; purity: 100%; MS (m/e): 413 (MH⁺).

7.3.812 N4-(1,4-Benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride Salt (R920395)

To a solution of N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (1 equivalent) in MeOH at 0 °C was added HCl (4M, dioxane, 1.1 equivalents) dropwise and shaken for 5 minutes. The resulting solution was diluted with EtOAc and the solid obtained was filtered washed with EtOAc, dried and analyzed to give N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride

Salt. ¹H NMR (DMSO-d₆): δ 9.80 (bs, 2H), 8.12 (d, 1H, J= 4.8 Hz), 7.89 (bd, 1H, J= 4.5 Hz), 7.18 (m, 3H), 8.24 (m, 2H), 6.60 (bd, 2H, J= 8.1 Hz), 4.36 (s, 2H), 4.10 (t, 2H, J= 3.9 Hz), 3.27 (t, 2H, J= 3.9 Hz), 2.62 (d, 3H, J= 4.5 Hz); LCMS: purity: 98%, MS (m/e): 425 (MH⁺).

5 **7.3.813 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Trifluoro Acetic Acid Salt (R926826)**

In like manner to the synthesis of N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Hydrogen Chloride
 10 Salt the reaction of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine with trifluoroacetic acid gave N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine Trifluoro Acetic Acid
 15 Salt. ¹H NMR (DMSO-d₆): δ 9.40 (bs, 1H), 9.36 (bs, 1H), 8.07 (d, 1H, J= 4.2 Hz), 7.94 (bd, 1H), 7.22 (m, 4H), 7.11 (t, 1H, J= 7.5 Hz), 6.79 (d, 1H, J= 8.7 Hz), 6.51 (bd, 1H, J= 7.5 Hz), 4.33 (s, 2H), 4.21 (bs, 4H), 2.63 (d, 3H, 3.3 Hz).

20 **7.3.814 5-Fluoro-N4-[(1H)-indol-6-yl]-N2-[4-methoxy-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926752)**

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(1H)indol-6-yl]-4-pyrimidineamine and 4-methoxy-3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to produce 5-fluoro-N4-[(1H)-indol-6-yl]-N2-[4-methoxy-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ
 25 7.83 (d, 1H, J= 3.6 Hz), 7.73 (d, 1H, J= 0.9 Hz), 7.49 (d, 1H, J= 8.1 Hz), 7.39 (d, 1H, J= 3.0 Hz), 7.20 (d, 1H, J= 3.6 Hz), 7.15 (dd, 1H, J= 1.8 and 8.1 Hz), 7.05 (dd, 1H, J= 2.1 and 8.7 Hz), 6.81 (d, 1H, J= 8.7 Hz), 6.41 (d, 1H, J= 4.2 Hz), 4.09 (s, 2H), 3.81 (s, 3H), 2.76 (s, 3H); LCMS: purity: 100%; MS (m/e): 437(MH⁺).

7.3.815 5-Fluoro-N4-(3-hydroxy-4-methylphenyl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926753)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(2-hydroxy-4-methylphenyl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy] aniline were reacted to produce 5-fluoro-N4-(3-hydroxy-4-methylphenyl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.95 (bs, 1H), 9.83 (bs, 1H), 9.38 (bs, 1H), 8.17 (d, 1H, J= 4.4 Hz), 7.97 (d, 1H, J= 4.8 Hz), 7.24-7.17 (m, 2H), 7.16 (d, 1H, J= 8.4 Hz), 7.10 (dd, 1H, J=1.8 and 8.4 Hz), 7.03 (d, 1H, J= 2.4 Hz), 7.00 (d, 1H, J= 9.0 Hz), 6.61 (d, 1H, J= 8.7 Hz), 4.34 (s, 2H), 2.63 (d, 3H, J= 4.5 Hz), 2.08 (s, 3H); LCMS: purity: 96%; MS (m/e): 398(MH⁺).

7.3.816 5-Fluoro-N4-(3-dihydroxyborylphenyl)- N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926754)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-dihydroxyborylphenyl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to produce 5-fluoro-N4-(3-dihydroxyborylphenyl)- N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.38 (bs, 1H), 9.22 (bs, 1H), 8.08 (d, 1H, J= 3.6 Hz), 8.06-7.81 (m, 4H), 7.51 (d, 1H, J= 8.1 Hz), 7.33-7.28 (m, 3H), 7.06 (t, 1H, J= 8.1 Hz), 6.44 (dd, 1H, J= 2.4 and 7.5 Hz), 4.33 (s, 2H), 2.63 (d, 3H, J= 4.8 Hz); LCMS: purity: 95%; MS (m/e): 412(MH⁺).

7.3.817 5-Fluoro-N4-(3-dihydroxyborylphenyl)-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926755)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-dihydroxyborylphenyl)-4-pyrimidineamine and 3-hydroxyaniline were reacted to produce 5-Fluoro-N4-(3-dihydroxyborylphenyl)-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.68 (bs, 1H), 9.35 (bs, 1H), 9.22 (bs, 1H), 8.10 (d, 1H, J= 3.9 Hz), 7.88-7.80 (m, 2H), 7.54 (d, 1H, J= 7.2 Hz), 7.31 (t, 1H, J= 7.2 Hz), 7.08 (d, 1H, J= 8.4 Hz), 6.98-6.93 (m, 2H), 6.35 (d, 1H, J= 8.4 Hz); LCMS: purity: 96%; MS (m/e): 341(MH⁺).

7.3.818 N2-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-(3-hydroxyborylphenyl)-2,4-pyrimidinediamine (R926756)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-dihydroxyborylphenyl)-4-pyrimidineamine and 3,4-ethylenedioxyaniline were reacted to produce N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-(3-hydroxyborylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.46 (bs, 1H), 9.11 (bs, 1H), 8.05 (d, 1H, J= 4.2 Hz), 7.95 (bs, 1H), 7.88 (s, 1H), 7.78 (d, 1H, J= 7.5 Hz), 7.52 (d, 1H, J= 7.5 Hz), 7.29 (t, 1H, J= 7.5 Hz), 7.16 (s, 1H), 7.02 (d, 1H, J= 8.7 Hz), 6.65 (d, 1H, J= 8.7 Hz), 3.40 (s, 4H); LCMS: purity: 98%; MS (m/e): 383(MH⁺).

7.3.819 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926757)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.32 (s, 1H), 9.17 (s, 1H), 9.04 (s, 1H), 8.04 (d, 1H, J= 4.2 Hz), 7.76 (d, 1H, J= 4.8 Hz), 7.32 (td, 2H, J= 1.8 and 8.1 Hz), 7.13-7.04 (m, 3H), 6.95 (d, 1H, J= 8.4 Hz), 6.46 (dd, 1H, J= 1.8 and 8.4 Hz), 4.31 (s, 2H), 2.65 (d, 3H, J= 4.8 Hz), 2.14 (s, 3H); LCMS: purity: 99%; MS (m/e): 398(MH⁺).

7.3.820 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro- N2-[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926758)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to produce N4-(3,4-ethylenedioxyphenyl)-5-fluoro- N2-[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.13 (bs, 1H), 9.05 (s, 1H), 8.01 (d, 1H, J= 4.2 Hz), 7.76 (d, 1H, J= 4.8 Hz), 7.32 (d, 1H, J= 2.4 Hz), 7.27 (dd, 1H, J= 2.4 and 8.1 Hz), 7.21 (dd, 1H, J= 2.4 and 8.7 Hz), 7.13 (d,

1H, J= 1.8 Hz), 6.95 (d, 1H, J= 8.1 Hz), 6.76 (d, 1H, J= 8.7 Hz), 4.28 (s, 2H), 4.20 (s, 4H), 2.65 (d, 3H, J= 4.8 Hz), 2.15 (s, 3H); LCMS: purity: 97%; MS (m/e): 440(MH⁺).

7.3.821 5-Fluoro-N4-(3-hydroxy-4-methylphenyl)-N2-[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926759)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(2-hydroxy-4-methylphenyl)-4-pyrimidineamine and 4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to produce 5-fluoro-N4-(3-hydroxy-4-methylphenyl)-N2-[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 10.09 (bs, 1H), 9.96 (bs, 1H), 9.44 (bs, 1H), 8.16 (d, 1H, J= 4.8 Hz), 7.81 (d, 1H, J= 4.8 Hz), 7.13-6.94 (m, 6H), 4.29 (s, 2H), 2.64 (d, 3H, J= 4.5 Hz), 2.17 (s, 3H), 2.07 (s, 3H); LCMS: purity: 99%; MS (m/e): 412(MH⁺).

7.3.822 5-Fluoro-N2,N4-bis[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926760)

In a like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N2,N4-bis[4-methyl-3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.30 (s, 1H), 9.02 (s, 1H), 8.06 (d, 1H, J= 3.6 Hz), 7.94 (d, 1H, J= 4.5 Hz), 7.80 (d, 1H, J= 4.2 Hz), 7.58 (bs, 1H), 7.31-7.22 (m, 3H), 7.05 (d, 1H, J= 9.0 Hz), 6.97 (d, 1H, J= 7.5 Hz), 4.41 (s, 2H), 4.27 (s, 2H), 2.66 (d, 3H, J= 4.2 Hz), 2.63 (d, 3H, J= 4.2 Hz), 2.18 (s, 3H), 2.14 (s, 3H); LCMS: purity: 100%; MS (m/e): 483(MH⁺).

7.3.823 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine (R926761)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3,4,5-trimethoxyaniline were reacted to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.33 (s, 1H), 9.17 (s, 1H), 8.99 (s, 1H), 8.06 (d, 1H, J= 3.3 Hz), 7.27 (d, 1H, J= 7.5

Hz), 7.08-7.02 (m, 4H), 6.46 (dd, 1H, J= 1.8 and 7.8 Hz), 3.60 (s, 6H), 3.57 (s, 3H); LCMS: purity: 99%; MS (m/e): 387(MH⁺).

7.3.824 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine (R926762)

5 In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3,4,5-trimethoxyaniline were reacted to produce N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine
1H NMR (DMSO-*d*₆): δ 8.08 (d, 1H, J= 4.8 Hz), 7.29 (d, 1H, J= 2.4 Hz), 7.15 (dd, 1H, J=3.0 and 9.0 Hz), 6.91 (s, 1H), 6.76 (d, 1H, J= 8.7 Hz), 4.20 (s, 4H), 3.61 (s, 6H), 3.59 (s, 3H); LCMS: purity: 97%; MS (m/e): 429(MH⁺).

7.3.825 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3,5-dichloro-4-hydroxyphenyl)-2,4-pyrimidinediamine (R926763)

15 In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3,5-dichloro-4-hydroxyaniline were reacted to produce N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3,5-dichloro-4-hydroxyphenyl)-2,4-pyrimidinediamine. 1H NMR (DMSO-*d*₆): δ 9.50 (bs, 1H), 9.26 (bd, 2H, J= 7.5 Hz), 8.06 (d, 1H, J= 3.9 Hz), 7.65 (s, 2H), 7.18-7.13 (m, 2H), 6.80 (d, 1H, J= 9.0 Hz), 4.20 (s, 4H);
20 LCMS: purity: 100%; MS (m/e): 424(MH⁺).

7.3.826 5-Fluoro-N2-(3,5-dichloro-4-hydroxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926890)

25 In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro -N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3,5-dichloro-4-hydroxyaniline were reacted to produce 5-Fluoro-N2-(3,5-dichloro-4-hydroxyphenyl)-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. 1H NMR (DMSO-*d*₆): δ 9.47 (bs, 1H), 9.35 (bs, 1H), 9.22 (bs, 2H), 8.09 (d, 1H, J= 3.6 Hz), 7.70 (s, 2H), 7.31 (dd, 1H, J= 1.2 and 9.3 Hz), 7.10 (t, 1H, J= 7.5 Hz), 7.00 (bs, 1H), 6.48 (dd, 1H, J= 1.2 and 6.9 Hz); LCMS: purity: 93%; MS (m/e): 382(MH⁺).

7.3.827 N4-(3-Chloro-4-methoxyphenyl)-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926891)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-chloro-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to produce N4-(3-chloro-4-methoxyphenyl)-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.85 (bs, 1H), 9.70 (bs, 1H), 8.17 (d, 1H, J= 4.8 Hz), 7.98 (d, 1H, J= 3.9 Hz), 7.79 (d, 1H, J= 2.4 Hz), 7.65 (dd, 1H, J= 3.0 and 9.3 Hz), 7.24-7.09 (m, 4H), 6.57 (d, 1H, J= 5.7 Hz), 4.34 (s, 2H), 3.82 (s, 3H), 2.62 (d, 3H, J= 4.8 Hz); LCMS: purity: 95%; MS (m/e): 433(MH⁺).

7.3.828 5-Fluoro-N4-(3-fluoro-4-methoxyphenyl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926892)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-fluoro-4-methoxyphenyl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to produce 5-fluoro-N4-(3-fluoro-4-methoxyphenyl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.68 (bs, 1H), 9.53 (bs, 1H), 8.13 (d, 1H, J= 4.2 Hz), 7.97 (d, 1H, J= 4.8 Hz), 7.76 (dd, 1H, J= 2.4 and 13.5 Hz), 7.47 (d, 1H, J= 7.5 Hz), 7.27-7.08 (m, 4H), 6.54 (d, 1H, J= 8.4 Hz), 4.35 (s, 2H), 3.80 (s, 3H), 2.63 (d, 3H, J= 4.8 Hz); LCMS: purity: 94%; MS (m/e): 416(MH⁺).

7.3.829 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N4-(3-hydroxy-5-methylphenyl)-2,4-pyrimidinediamine (R926893)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine, 4-amino-*m*-cresol hydrochloride salt, and diisopropylethylamine were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-(3-hydroxy-5-methylphenyl)-2,4-pyrimidinediamine.

¹H NMR (DMSO-*d*₆): δ 9.06 (s, 1H), 8.94 (s, 1H), 8.11 (s, 1H), 7.86 (d, 1H, J= 3.9 Hz), 7.21-7.15 (m, 2H), 7.03 (d, 1H, J= 8.1 Hz), 6.59 (bd, 2H, J= 8.7 Hz), 6.52 (dd, 1H, J= 3.0 and 8.1 Hz), 4.17 (s, 4H), 2.05 (s, 3H); LCMS: purity: 99%; MS (m/e): 369(MH⁺).

7.3.830 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-fluoro-5-trifluoromethylphenyl)-2,4-pyrimidinediamine (R926894)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-amino-5-fluorobenzotrifluoride were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-fluoro-5-trifluoromethylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.75 (s, 1H), 9.32 (d, 1H, J= 1.2 Hz), 8.13 (d, 1H, J= 3.6 Hz), 7.99 (d, 1H, J= 12.3 Hz), 7.77 (s, 1H), 7.21 (d, 1H, J= 2.4 Hz), 7.13 (dd, 1H, J= 2.1 and 8.7 Hz), 7.03 (d, 1H, J= 9.0 Hz), 6.80 (d, 1H, J= 8.7 Hz), 4.21 (s, 4H); LCMS: purity: 97%; MS (m/e): 425(MH⁺).

7.3.831 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3-methyl-5-trifluoromethylphenyl)-2,4-pyrimidinediamine (R926895)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-amino-5-methylbenzotrifluoride were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-methyl-5-trifluoromethylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.57 (bs, 1H), 9.39 (bs, 1H), 8.12 (d, 1H, J= 3.6 Hz), 7.77 (s, 2H), 7.25-7.13 (m, 2H), 7.02 (s, 1H), 6.79 (d, 1H, J= 9.0 Hz), 4.20 (s, 4H), 2.27 (s, 3H); LCMS: purity: 100%; MS (m/e): 421(MH⁺).

7.3.832 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(5-methoxy-2-methylphenyl)-2,4-pyrimidinediamine (R926896)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 5-methoxy-2-methylaniline were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(5-methoxy-2-methylphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.91 (bs, 1H), 7.61 (d, 1H, J= 2.1 Hz), 7.17 (d, 1H, J= 3.0 Hz), 7.05 (d, 1H, J= 9.3 Hz), 7.03 (dd, 1H, J= 3.0 and 8.7 Hz),

6.82 (d, 1H, J= 8.1 Hz), 6.68-6.60 (m, 2H), 6.55 (dd, 1H, J= 2.1 and 8.1 Hz), 4.26 (s, 4H), 3.70 (s, 3H), 2.22 (s, 3H); ^{19}F NMR (282 MHz, CDCl_3): -47450; LCMS: purity: 99%; MS (m/e): 383(MH^+).

7.3.833 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-fluoro-5-methylphenyl)-2,4-pyrimidinediamine (R926897)

5

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 2-fluoro-5-methylaniline were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(2-fluoro-5-methylphenyl)-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 8.11 (dd, 1H, J= 1.8 and 8.1 Hz), 7.94 (d, 1H, J= 2.7 Hz), 7.08-6.84 (m, 4H), 6.74-6.67 (m, 1H), 6.64-6.59 (m, 1H), 4.27 (s, 4H), 2.28 (s, 3H); ^{19}F NMR (282 MHz, CDCl_3): -38659, -47267; LCMS: purity: 100%; MS (m/e): 371(MH^+).

10

7.3.834 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(3,5-difluorophenyl)-2,4-pyrimidinediamine (R926898)

15

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3,5-difluoroaniline were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3,5-difluorophenyl)-2,4-pyrimidinediamine. ^1H NMR (CDCl_3): δ 7.94 (d, 1H, J= 3.3 Hz), 7.20-7.11 (m, 3H), 7.02 (s, 1H), 6.92-6.90 (m, 2H), 6.65 (s, 1H), 6.39 (tt, 1H, J= 2.4 and 9.0 Hz), 4.31 (s, 4H); ^{19}F NMR (282 MHz, CDCl_3): -31142, -47002; LCMS: purity: 97%; MS (m/e): 375(MH^+).

20

7.3.835 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro-N2-(4-trifluoromethylthiophenyl)-2,4-pyrimidinediamine (R926899)

25

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 4-(trifluoromethylthio)aniline were reacted to provide N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(4-trifluoromethylthiophenyl)-2,4-pyrimidinediamine. ^1H NMR ($\text{DMSO}-d_6$): δ 9.73 (s, 1H), 9.47 (s, 1H), 8.13 (d, 1H, J= 3.6 Hz), 7.79 (d, 2H, J= 9.0 Hz), 7.51 (d, 2H, J= 9.0 Hz), 7.28

30

(d, 1H, J= 2.1 Hz), 7.12 (dd, 1H, J= 2.4 and 9.0 Hz), 6.83 (d, 1H, J= 8.7 Hz), 4.23 (s, 4H); ¹⁹F NMR (282 MHz DMSO-*d*₆): -12306; LCMS: purity: 97%; MS (m/e): 439(MH⁺).

7.3.836 N4-[3-(Benzothiazol-2-yl)-4-chlorophenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926900)

5

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, N4-[3-(Benzothiazol-2-yl)-4-chlorophenyl]-2-chloro-5-fluoro-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide N4-[3-(benzothiazol-2-yl)-4-chlorophenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.77 (s, 1H), 9.30 (s, 1H), 8.49 (d, 1H, J= 3.0 Hz), 8.25 (dd, 1H, J= 3.0 and 9.0), 8.21-8.16 (m, 2H), 8.06 (d, 1H, J= 7.8 Hz), 7.92 (d, 1H, J= 4.8 Hz), 7.63-7.48 (m, 3H), 7.30 (t, 1H, J= 1.8 Hz), 7.22 (dd, 1H, J= 1.8 and 7.5 Hz), 6..95 (t, 1H, J= 8.1 Hz), 6.32 (dd, 1H, J= 1.2 and 8.1 Hz), 4.29 (s, 2H), 2.62 (d, 1H, J= 4.8 Hz); LCMS: purity: 100%; MS (m/e): 536(MH⁺).

15

7.3.837 5-Fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(3-methoxy-4-methylphenyl)-2,4-pyrimidinediamine (R926902)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-Chloro-5-fluoro-N4-(3-methoxy-4-methylphenyl)-4-pyrimidineamine and 3-methoxy-4-methylaniline were reacted to provide 5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(3-methoxy-4-methylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.78 (bs, 1H), 9.63 (bs, 1H), 8.15 (d, 1H, J= 4.5 Hz), 7.94 (d, 1H, J= 4.5 Hz), 7.94 (d, 1H, J= 4.5 Hz), 7.30 (dd, 1H, J= 1.8 and 8.4 Hz), 7.25-7.04 (m, 5H), 6.57 (d, 1H, J= 8.1 Hz), 4.31 (s, 2H), 3.66 (s, 3H), 2.62 (d, 1H, J= 4.8 Hz), 2.09 (s, 3H); LCMS: purity: 95%; MS (m/e): 412(MH⁺).

25

7.3.838 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(methoxycarbonyl)-(1H)-indol-6-yl]-2,4-pyrimidinediamine (R926903)

30

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 6-amino-2-(methoxycarbonyl)-(1H)-

indole were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(methoxycarbonyl)-(1H)-indol-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 11.53 (s, 1H), 9.37 (s, 1H), 9.18 (d, 2H, J= 9.9 Hz), 8.08 (d, 1H, J= 3.6 Hz), 7.96 (bs, 1H), 7.46 (d, 1H, J= 9.0 Hz), 7.39-7.35 (m, 2H), 7.16 (t, 1H, J= 2.4 Hz), 7.10-7.04 (m, 2H), 6.48 (dd, 1H, J= 2.4 and 7.5 Hz), 3.82 (s, 3H); LCMS: purity: 95%; MS (m/e): 394(MH⁺).

7.3.839 5-Fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-[2-(methoxycarbonyl)-(1H)-indol-6-yl]-2,4-pyrimidinediamine (R926904)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro N4-[2-(methoxycarbonyl)-(1H)-indol-6-yl]-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-[2-(methoxycarbonyl)-(1H)-indol-6-yl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 9.05 (bs, 1H), 8.35 (s, 1H), 8.00 (bs, 1H), 7.66-7.62 (m, 2H), 7.27-7.17 (m, 3H), 7.01-6.90 (m, 3H), 6.64 (dd, 1H, J= 2.4 and 8.1 Hz), 6.40 (bs, 1H), 4.49 (s, 2H), 3.94 (s, 3H), 2.75 (d, 3H, J= 5.1 Hz); LCMS: purity: 86%; MS (m/e): 465(MH⁺).

7.3.840 N4-[3-[[4-(Ethoxycarbonyl)piperidino]methyl]phenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926905)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[3-[[4-(ethoxycarbonyl)piperidino]methyl]phenyl]-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N4-[3-[[4-(ethoxycarbonyl)piperidino]methyl]phenyl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): 9.33 (s, 1H), 9.20 (s, 1H), 8.09 (d, 1H, J= 4.2 Hz), 7.93 (d, 1H, J= 4.8 Hz), 7.82 (d, 1H, J= 8.1 Hz), 7.55 (s, 1H), 7.35 (t, 1H, J= 2.4 Hz), 7.29-7.22 (m, 2H), 7.09 (t, 1H, J= 8.1 Hz), 6.96 (d, 1H, J= 7.8 Hz), 6.47 (dd, 1H, J= 2.4 and 8.1 Hz), 4.32 (s, 2H), 4.02 (q, 2H, J= 6.9 Hz), 3.39 (s, 2H), 2.73 (bd, 2H, J=11.1 Hz), 2.63 (d, 3H, J= 4.5 Hz), 2.30-2.20 (m, 1H),

1.94 (t, 2H, J= 11.1 Hz), 1.74 (d, 2H, J= 9.9 Hz), 1.60-1.50 (m, 2H), 1.14 (t, 3H, J= 6.9 Hz); LCMS: purity: 99%; MS (m/e): 537(M - CH₂⁺).

7.3.841 N2-[3-(Ethoxycarbonyl-1,1-dimethylmethylenedioxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926906)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methylenedioxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-(ethoxycarbonyl-1,1-dimethylmethylenedioxy)aniline were reacted to provide N2-[3-(ethoxycarbonyl-1,1-dimethylmethylenedioxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.91 (d, 1H, J= 4.8 Hz), 7.20-7.03 (m, 6H), 6.67 (td, 1H, J= 2.1 and 7.5 Hz), 6.57-6.53 (m, 1H), 4.19 (q, 2H, J= 6.9 Hz), 1.53 (s, 6H), 1.20 (t, 3H, J= 6.9 Hz); ¹⁹F NMR (282 MHz, CD₃OD): -46120; LCMS: purity: 91%; MS (m/e): 427(MH⁺).

7.3.842 N2-[3-(Ethoxycarbonyl-1,1-dimethylmethylenedioxy)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926907)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methylenedioxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-(ethoxycarbonyl-1,1-dimethylmethylenedioxy)aniline were reacted to provide N2-[3-(ethoxycarbonyl-1,1-dimethylmethylenedioxy)phenyl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.92 (d, 1H, J= 3.0 Hz), 7.21-7.08 (m, 4H), 7.00 (dd, 1H, J= 2.4 and 8.4 Hz), 6.93 (bs, 1H), 6.86 (d, 1H, J= 8.7 Hz), 6.99 (d, 1H, J= 2.4 Hz), 6.45 (ddd, 1H, J= 1.2, 1.2, and 7.8 Hz), 4.27 (s, 4H), 4.23 (q, 2H, J= 6.9 Hz), 1.60 (s, 6H), 1.23 (t, 3H, J= 6.9 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -47216; LCMS: purity: 85%; MS (m/e): 469(MH⁺).

7.3.843 N2-[3-(Ethoxycarbonyl-1,1-dimethylmethylenedioxy)phenyl]-5-fluoro-N4-(3-hydroxy-4-methylphenyl)-2,4-pyrimidinediamine (R926908)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methylenedioxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(2-hydroxy-4-methylphenyl)-4-pyrimidineamine and 3-(ethoxycarbonyl-1,1-dimethylmethylenedioxy)aniline were reacted to provide N2-[3-(ethoxycarbonyl-1,1-

dimethylmethylenedioxy)phenyl]-5-fluoro-N4-(3-hydroxy-4-methylphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.86 (bs, 1H), 7.80 (bs, 1H), 7.53 (s, 1H), 7.16-6.86 (m, 4H), 6.54 (d, 2H, J = 7.5 Hz), 4.21 (q, 2H, J = 6.9 Hz), 3.48 (s, 2H), 2.20 (s, 3H), 1.60 (s, 6H), 1.22 (t, 3H, J = 6.9 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -46808; LCMS: purity: 96%; MS (m/e): 441(MH⁺).

7.3.844 N2-[3-(Ethoxycarbonyl-1,1-dimethylmethylenedioxy)phenyl]-5-fluoro-N4-[(1H)-indol-6-yl]-2,4-pyrimidinediamine (R926909)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methylenedioxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(1H)indol-6-yl]-4-pyrimidineamine and 3-(ethoxycarbonyl-1,1-dimethylmethylenedioxy)aniline were reacted to provide N2-[3-(Ethoxycarbonyl-1,1-dimethylmethylenedioxy)phenyl]-5-fluoro-N4-[(1H)-indol-6-yl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 9.43 (bs, 1H), 8.64 (s, 1H), 7.92 (d, 1H, J = 3.6 Hz), 7.66 (t, 1H, J = 2.4 Hz), 7.54 (d, 1H, J = 8.4 Hz), 7.44 (s, 1H), 7.19 (t, 1H, J = 3.0 Hz), 7.15 (d, 1H, J = 8.1 Hz), 6.96 (d, 1H, J = 3.0 Hz), 6.80 (dd, 1H, J = 1.8 and 7.5 Hz), 6.77 (dd, 1H, J = 1.8 and 8.1 Hz), 6.52 (dd, 1H, J = 1.8 and 7.5 Hz), 6.49-6.46 (m, 1H), 4.32 (q, 2H, J = 7.2 Hz), 1.57 (s, 6H), 1.31 (t, 3H, J = 7.2 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -47190; LCMS: purity: 93%; MS (m/e): 450(MH⁺).

7.3.845 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-methylamino)carbonyl-1,1-dimethylmethylenedioxy]phenyl]-2,4-pyrimidinediamine (R926913)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methylenedioxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-(N-methylamino)carbonyl-1,1-dimethylmethylenedioxyaniline were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-methylamino)carbonyl-1,1-dimethylmethylenedioxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.35 (s, 1H), 9.20 (s, 1H), 9.17 (s, 1H), 8.07 (d, 1H, J = 3.3 Hz), 7.93 (d, 1H, J = 3.9 Hz), 7.40-7.29 (m, 3H), 7.13-7.02 (m, 3H), 6.47 (d, 1H, J = 7.5 Hz), 6.33 (d, 1H, J = 7.5 Hz), 2.60 (s, 3H), 1.37 (s, 6H); LCMS: purity: 97%; MS (m/e): 412(MH⁺).

7.3.846 5-Fluoro-N4-(1,2,3,4-tetrahydroisoquin-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926914)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[2-(t-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-7-yl]-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N4-(1,2,3,4-tetrahydroisoquin-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 7.90 (d, 1H, J= 3.3 Hz), 7.47 (d, 1H, J= 2.4 Hz), 7.42-7.37 (m, 2H), 7.16 (t, 1H, J= 8.4 Hz), 7.10-7.04 (m, 2H), 6.50 (ddd, 1H, J= 1.2, 2.4, and 8.1 Hz), 4.26 (s, 2H), 3.93 (s, 2H), 3.12 (t, 2H, J= 6.3 Hz), 2.84-2.76 (m, 5H), ; ¹⁹F NMR (282 MHz, CD₃OD): -47489; LCMS: purity: 87%; MS (m/e): 423(MH⁺).

7.3.847 N4-(3,4-Ethylenedioxyphenyl)-5-fluoro- N2-[3-[(N-methylamino)carbonyl-1,1-dimethylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926915)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 3-(N-methylamino)carbonyl-1,1-dimethylmethyleneoxyaniline were reacted to provide N4-(3,4-Ethylenedioxyphenyl)-5-fluoro- N2-[3-[(N-methylamino)carbonyl-1,1-dimethylmethyleneoxy]phenyl]- 2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.26 (t, 1H, J= 7.5 Hz), 7.19 (d, 1H, J= 9.3 Hz), 7.13 (d, 1H, J= 2.4 Hz), 7.06 (dd, 1H, J= 2.4 and 8.7 Hz), 7.04-7.03 (m, 1H), 6.83 (d, 1H, J= 9.0 Hz), 6.75 (d, 1H, J= 7.2 Hz), 4.25 (s, 4H), 2.76 (s, 3H), 1.43 (s, 6H); LCMS: purity: 97%; MS (m/e): 454(MH⁺).

7.3.848 5-Fluoro-N4-[3-[(N-allylamino)carbonyloxy]phenyl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]- 2,4-pyrimidinediamine (R926917)

A mixture of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (20 mg, 0.052 mmol), allyl isocyanate (13mg, 0.16 mmol), and 2-(N,N-dimethylamino)pyridine (18 mg, 0.15 mmol) in anhydrous THF (1 mL) were heated at 60°C in a sealed vial for 2 days. The reaction was diluted with ethyl acetate and washed with 1N HCl and brine. Concentration gave an oily residue which was

purified by preparative TLC (5% methanol/dichloromethane) to give the product 5-fluoro-N4-[3-[(N-allylamino)carbonyloxy]phenyl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.93 (d, 1H, J= 3.6 Hz), 7.62-7.55 (m, 2H), 7.32 (s, 1H), 7.30 (t, 1H, J= 8.1 Hz), 7.19-7.15 (m, 2H), 6.82 (dd, 1H, J= 2.4 and 8.1 Hz), 6.61 (m, 1H), 5.96-5.82 (m, 1H), 5.24 (dd, 1H, J= 1.8 and 16.8 Hz), 5.13 (dd, 1H, J= 1.8 and 11.7 Hz), 4.41 (s, 2H), 3.79 (d, 1H, J= 5.4 Hz), 2.80 (s, 3H); ¹⁹F NMR (282 MHz, CD₃OD): -47357; LCMS: purity: 99%; MS (m/e): 468(MH⁺).

7.3.849 5-Fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-[3-[(N-isopropylamino)carbonyl]-N-isopropylamino)carbonyloxy]phenyl]-2,4-pyrimidinediamine (R926916)

In a like manner to the preparation of 5-fluoro-N4-[3-[(N-allylamino)carbonyloxy]phenyl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine and isopropyl isocyanate were reacted to provide 5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-[3-[(N-isopropylamino)carbonyl]-N-isopropylamino)carbonyloxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.40 (bs, 1H), 9.27 (bs, 1H), 8.12 (d, 1H, J= 3.6 Hz), 7.94 (d, 1H, J= 3.9 Hz), 7.78 (d, 1H, J= 8.7 Hz), 7.64 (d, 1H, J= 7.5 Hz), 7.46 (s, 1H), 7.36-7.26 (m, 3H), 7.12 (t, 1H, J= 8.1 Hz), 6.81-6.74 (m, 1H), 6.47 (dd, 1H, J= 2.4 and 8.1 Hz), 5.43 (d, 1H, J= 3.9 Hz), 4.36 (s, 2H), 3.65-3.55 (m, 2H), 3.14 (s, 2H), 2.63 (d, 3H, J= 3.9 Hz), 1.10 (d, 6H, J= 7.2 Hz), 0.97 (d, 6H, J= 6.6 Hz).

7.3.850 N4-[3-[[N-(Ethoxycarbonylmethyl)amino]carbonyloxy]phenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926918)

In a like manner to the preparation of 5-fluoro-N4-[3-[(N-allylamino)carbonyloxy]phenyl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine and ethyl isocyanatoacetate were reacted to provide N4-[3-[[N-(ethoxycarbonylmethyl)amino]carbonyloxy]phenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.94 (d, 1H, J= 3.3 Hz), 7.69 (t, 1H, J= 1.8 Hz), 7.56 (ddd, 1H, J= 1.2, 1.2, and 8.1 Hz), 7.35 (m, 1H), 7.31 (t, 1H, J= 8.1 Hz), 7.18 (d, 1H, J= 2.4 Hz), 7.17 (d, 1H, J= 1.2 Hz), 6.84 (dd, 1H, J= 2.4 and 8.1 Hz), 6.63-6.58 (m, 1H), 4.42 (s, 2H), 4.20 (q, 2H, J= 7.2 Hz), 3.93

(s, 2H), 2.80 (s, 3H), 1.27 (t, 3H, J= 7.2 Hz); ¹⁹F NMR (282 MHz, CD₃OD): -47371; LCMS: purity: 89%; MS (m/e): 513(MH⁺).

7.3.851 N4-[3-[(N-(Ethylamino)carbonyloxy]phenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926919)

In a like manner to the preparation of 5-fluoro-N4-[3-[(N-allylamino)carbonyloxy]phenyl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine and ethyl isocyanate were reacted to provide N4-[3-[(N-(ethylamino)carbonyloxy]phenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.94 (d, 1H, J= 3.3 Hz), 6.84-6.79 (m, 2H), 7.61-7.55 (m, 2H), 6.62-6.56 (m, 2H), 7.33-7.27 (m, 1H), 7.19-7.17 (m, 1H), 4.41 (s, 2H), 3.23 (q, 2H, J= 7.2 Hz), 2.80 (s, 3H), 1.17 (t, 3H, J= 7.2 Hz); ¹⁹F NMR (282 MHz, CD₃OD): -47378; LCMS: purity: 100%; MS (m/e): 455(MH⁺).

7.3.852 5-Fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(4-methyl-3-trifluoromethylphenyl)-2,4-pyrimidinediamine (R926922)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(4-methyl-3-trifluoromethylphenyl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(4-methyl-3-trifluoromethylphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.79 (bs, 1H), 9.48 (bs, 1H), 8.17 (d, 1H, J= 4.2 Hz), 8.10 (d, 1H, J= 6.3 Hz), 7.96 (d, 1H, J= 4.8 Hz), 7.89 (d, 1H, J= 2.1 Hz), 7.38 (d, 1H, J= 9.0 Hz), 7.26-7.20 (m, 2H), 7.11 (t, 1H, J= 8.4 Hz), 6.53 (d, 1H, J= 8.4 Hz), 4.33 (s, 2H), 2.62 (d, 3H, J= 4.8 Hz), 2.39 (s, 3H); LCMS: purity: 94%; MS (m/e): 450(MH⁺).

7.3.853 5-Fluoro-N4-(4-fluoro-3-methylphenyl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926923)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(4-fluoro-3-methylphenyl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-Fluoro-N4-(4-

fluoro-3-methylphenyl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.67 (bs, 1H), 9.51 (bs, 1H), 8.14 (d, 1H, J= 4.8 Hz), 7.95 (d, 1H, J= 4.2 Hz), 7.64 (dd, 1H, J= 2.7 and 6.9 Hz), 7.57-7.50 (m, 1H), 7.23-7.06 (m, 4H), 6.55 (d, 1H, J= 7.5 Hz), 4.33 (s, 2H), 2.63 (d, 3H, J= 4.8 Hz), 2.19 (s, 3H);

5 LCMS: purity: 94%; MS (m/e): 400(MH⁺).

7.3.854 5-Fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(3-trifluoromethylthiophenyl)-2,4-pyrimidinediamine (R926925)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-trifluoromethylthiophenyl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(3-trifluoromethylthiophenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.83 (bs, 1H), 9.49 (bs, 1H), 8.21-8.15 (m, 2H), 8.01 (s, 1H), 7.94 (bs, 1H), 7.49 (t, 1H, J= 7.8 Hz), 7.38 (d, 1H, J= 7.8 Hz), 7.29 (s, 1H), 7.22 (d, 1H, J= 7.5 Hz), 7.14 (t, 1H, J= 8.4 Hz), 6.54 (d, 1H, J= 9.9 Hz), 4.34 (s, 2H), 2.62 (d, 3H, J= 4.8 Hz); LCMS: purity: 98%; MS (m/e): 468(MH⁺).

7.3.855 N2-[3,5-Bis(methoxycarbonylmethyleneoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926926)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3,5-bis(methoxycarbonylmethyleneoxy)aniline were reacted to provide N2-[3,5-bis(methoxycarbonylmethyleneoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.92 (d, 1H, J= 4.2 Hz), 7.20-7.10 (m, 3H), 6.92 (d, 2H, J= 2.4 Hz), 6.52 (ddd, 1H, J= 1.8, 1.8, and 7.5 Hz), 6.12 (t, 1H, J= 2.4 Hz), 4.55 (s, 4H), 3.77 (s, 6H); ¹⁹F NMR (282 MHz, CD₃OD): -47342; LCMS: purity: 92%; MS (m/e): 473(MH⁺).

7.3.856 5-Fluoro-N2-[3-hydroxy-5-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926927)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-hydroxy-5-(methoxycarbonylmethyleneoxy)aniline were reacted to produce 5-fluoro-N2-[3-hydroxy-5-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 8.13 (d, 1H, J= 4.8 Hz), 7.37-7.33 (m, 1H), 7.11 (t, 1H, J= 8.4 Hz), 7.07-7.05 (m, 1H), 6.73-6.65 (m, 2H), 6.51 (dd, 1H, J= 2.1 and 8.1 Hz), 5.97 ((s, 1H), 4.59 (s, 2H), 3.67 (s, 3H); LCMS: purity: 93%; MS (m/e): 401(MH⁺).

7.3.857 N2-[3-[(N-Ethylamino)carbonyloxy]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926928)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-[(N-ethylamino)carbonyloxy]aniline were reacted to provide N2-[3-[(N-ethylamino)carbonyloxy]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.92 (d, 1H, J= 3.0 Hz), 7.67-7.55 (m, 2H), 7.24 (t, 1H, J= 7.5 Hz), 7.16 (t, 1H, J= 7.5 Hz), 7.07-6.98 (m, 2H), 6.84-6.79 (m, 2H), 6.67 (m, 2H), 6.60 (d, 1H, J= 7.5 Hz), 5.22-5.14 (m, 1H), 3.36-3.27 (m, 2H), 2.95 (s, 1H), 2.88 (s, 1H), 1.20 (t, 3H, J= 7.5 Hz); ¹⁹F NMR (282 MHz, CDCl₃): -47012; LCMS: purity: 99%; MS (m/e): 384(MH⁺).

7.3.858 5-Fluoro-N2-[3-hydroxy-5-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926929)

A solution of 5-fluoro-N2-[3-hydroxy-5-(methoxycarbonylmethyleneoxy)phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (56 mg, 0.13 mmol), methylamine hydrochloride (90 mg, 1.3 mmol), and diisopropylethylamine (0.12 mL, 0.70 mmol) in methanol (2 mL) was heated at 100°C for 8h. The cooled reaction mixture was poured into 1N HCl (20 mL) saturated with NaCl, and extracted with ethyl acetate. Purification by preparative TLC (5% methanol/dichloromethane) gave the product, 5-fluoro-N2-[3-hydroxy-5-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.29 (bs, 1H), 9.16 (s, 1H), 9.01 (s, 1H), 8.06

(d, 1H, J= 3.3 Hz), 7.87 (d, 1H, J= 4.8 Hz), 7.42 (dd, 1H, J= 1.5 and 8.1 Hz), 7.13-7.05 (m, 2H), 6.89-6.81 (m, 2H), 6.45 (dd, 1H, J= 2.4 and 8.4 Hz), 5.92 (t, 1H, J= 2.4 Hz), 4.28 (s, 2H), 3.30(bs, 1H), 2.63 (s, 3H); LCMS: purity: 94%; MS (m/e): 400(MH⁺).

7.3.859 N2-[3,5-Bis[(N-methylamino)carbonylmethyleneoxy]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926930)

In a like manner to the preparation of 5-fluoro-N2-[3-hydroxy-5-[(N-methylamino)carbonylmethyleneoxy]phenyl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N2-[3,5-bis(methoxycarbonylmethyleneoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine, methylamine hydrochloride, and diisopropylethylamine were reacted to give N2-[3,5-Bis[(N-methylamino)carbonylmethyleneoxy]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.91 (bs, 1H), 7.25 (t, 1H, J= 1.8 Hz), 7.14-7.11 (m, 1H), 6.98 (s, 1H), 6.97 (s, 1H), 6.55-6.50 (m, 1H), 6.26-6.23 (m, 1H), 4.39 (s, 4H), 2.81 (s, 6H); ¹⁹F NMR (282 MHz, CD₃OD): -47307; LCMS: purity: 99%; MS (m/e): 471=(MH⁺).

7.3.860 5-Fluoro-N4-[(1H)-indol-5-yl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926931)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(1H)-indol-5-yl]-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N4-[(1H)-indol-5-yl]-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 11.09 (bs, 1H), 9.93 (bs, 1H), 9.67 (bs, 1H), 8.12 (d, 1H, J= 4.81 Hz), 7.94-7.82 (m, 2H), 7.37-7.22 (m, 4H), 7.13 (bs, 1H), 7.07 (t, 1H, J= 8.1 Hz), 6.58 (d, 1H, J= 7.8 Hz), 6.37 (s, 1H), 4.32 (s, 2H), 2.61 (d, 3H, J= 4.2 Hz); LCMS: purity: 92%; MS (m/e): 407(MH⁺).

7.3.861 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[(1H)-indol-5-yl]-2,4-pyrimidinediamine (R926932)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-

fluoro-N4-[(1H)-indol-5-yl]-4-pyrimidineamine and 3-hydroxyaniline were reacted to provide 5-fluoro-N2-(3-hydroxyphenyl)-N4-[(1H)-indol-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 11.13 (s, 1H), 10.25 (bs, 1H), 9.87 (bs, 1H), 9.43 (bs, 1H), 8.16 (d, 1H, J= 5.1 Hz), 7.89 (d, 1H, J= 0.09 Hz), 7.39-7.27 (m, 3H), 7.03-6.94 (m, 2H), 6.83 (s, 1H), 6.48 (d, 1H, J= 7.5 Hz), 6.40 (t, 1H, J= 2.1 Hz); LCMS: purity: 92%; MS (m/e): 336(MH⁺).

7.3.862 5-Fluoro-N4-[(1H)-indol-6-yl]-N2-[3-[(N-methylamino)carbonyl]phenyl]-2,4-pyrimidinediamine (R926933)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(1H)indol-6-yl]-4-pyrimidineamine and 3-[(N-methylamino)carbonyl]aniline were reacted to provide 5-fluoro-N4-[(1H)indol-6-yl]-N2-[3-[(N-methylamino)carbonyl]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.99 (t, 1H, J= 1.8 Hz), 7.89 (d, 1H, J= 3.6 Hz), 7.78-7.76 (m, 1H), 7.70 (ddd, 1H, J= 1.2, 2.4, and 8.4 Hz), 7.50 (d, 1H, J= 9.0 Hz), 7.31 (td, 1H, J= 1.2 and 7.5 Hz), 7.23-7.17 (m, 3H), 6.43 (dd, 1H, J= 1.2 and 3.6 Hz), 2.73 (s, 3H); ¹⁹F NMR (282 MHz, CD₃OD): -47513; LCMS: purity: 99%; MS (m/e): 377(MH⁺).

7.3.863 5-Fluoro-N4-[(1H)-indol-6-yl]-N2-[3-(N-morpholinocarbonyl)phenyl]-2,4-pyrimidinediamine (R926934)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(1H)indol-6-yl]-4-pyrimidineamine and 3-(N-morpholinocarbonyl)aniline were reacted to provide 5-fluoro-N4-[(1H)-indol-6-yl]-N2-[3-(N-morpholinocarbonyl)phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.96 (d, 1H, J= 4.8 Hz), 7.73 (t, 1H, J= 2.4 Hz), 7.66 (d, 1H, J= 1.2 Hz), 7.52 (d, 1H, J= 8.1 Hz), 7.49 (ddd, 1H, J= 0.09, 2.1, and 8.1 Hz), 7.33-7.26 (m, 2H), 7.19 (dd, 1H, J= 1.8 and 8.7 Hz), 7.12-7.06 (m, 1H), 6.45 (dd, 1H, J= 1.3 and 3.0 Hz), 3.62-3.15 (m, 8H); ¹⁹F NMR (282 MHz, CD₃OD): -46545; LCMS: purity: 91%; MS (m/e): 433(MH⁺).

7.3.864 N2-[3-[[4-(Ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N4-[(1H)-indol-6-yl]-2,4-pyrimidinediamine (R926935)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[(1H)indol-6-yl]-4-pyrimidineamine and 3-[4-(ethoxycarbonyl)piperidino]aniline were reacted to provide N2-[3-[[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N4-[(1H)-indol-6-yl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.99 (d, 1H, J= 5.1 Hz), 7.64-7.58 (m, 2H), 7.52 (d, 1H, J= 8.7 Hz), 7.48 (ddd, 1H, J= 1.2, 2.4, and 8.1 Hz), 7.34-7.27 (m, 2H), 7.19-7.13 (m, 2H), 6.46 (dd, 1H, J= 1.2 and 4.2 Hz), 4.40-4.27 (m, 1H), 4.13 (q, 2H, J= 6.9 Hz), 3.56-3.41 (m, 1H), 2.95-2.82 (m, 2H), 2.58-2.47 (m, 1H), 1.98-1.82 (m, 1H), 1.75-7.60 (m, 1H), 1.58-1.39 (m, 2H), 1.24 (t, 3H, J= 6.9 Hz); ¹⁹F NMR (282 MHz, CD₃OD): -46101; LCMS: purity: 90%; MS (m/e): 503(MH⁺).

7.3.865 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-methylamino)carbonyl]phenyl]-2,4-pyrimidinediamine (R926936)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-[(N-methylamino)carbonyl]aniline were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-methylamino)carbonyl]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.01 (d, 1H, J= 5.4 Hz), 7.84 (t, 1H, J= 1.8 Hz), 7.68-7.61 (m, 2H), 7.45 (t, 1H, J= 8.4 Hz), 7.16-7.03 (m, 3H), 6.68 (td, 1H, J= 1.2 and 8.7 Hz), 2.90 (s, 3H); LCMS: purity: 95%; MS (m/e): 354(MH⁺).

7.3.866 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-propylamino)carbonyl]phenyl]-2,4-pyrimidinediamine (R926937)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-[(N-propylamino)carbonyl]aniline were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-[(N-propylamino)carbonyl]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.00 (d, 1H, J= 5.4 Hz), 7.84 (t, 1H, J= 1.8 Hz), 7.69-7.59 (m, 2H), 7.44 (t, 1H, J= 7.5 Hz), 7.16-7.05 (m, 3H), 6.67 (td, 1H, J= 2.4 and 7.2 Hz), 3.34-3.29 (m, 2H), 1.65-1.56 (m, 2H), 0.96 (t, 3H, J=

7.5 Hz); ^{19}F NMR (282 MHz, CD_3OD): -46049; LCMS: purity: 94%; MS (m/e): 382(MH^+).

7.3.867 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-morpholinocarbonyl)phenyl]-2,4-pyrimidinediamine (R926938)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-(N-morpholinocarbonyl)aniline were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[3-(N-

morpholinocarbonyl)phenyl]-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.93 (d, 1H, $J = 3.6$ Hz), 7.84 (t, 1H, $J = 1.8$ Hz), 7.62 (ddd, 1H, $J = 1.2, 2.4$, and 8.1 Hz), 7.32 (t, 1H, $J = 8.4$ Hz), 7.19-7.10 (m, 3H), 6.96 (dd, 1H, $J = 1.2$ and 7.8 Hz), 6.56 (ddd, 1H, $J = 1.2, 3.0$, and 6.9 Hz), 3.78-3.34 (m, 8H); ^{19}F NMR (282 MHz, CD_3OD): -47323; LCMS: purity: 100%; MS (m/e): 410(MH^+).

7.3.868 N2-[3-[[4-(Ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926939)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-[[4-

(ethoxycarbonyl)piperidino]carbonyl]aniline were reacted to provide N2-[3-[[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.92 (d, 1H, $J = 3.6$ Hz), 7.82 (s, 1H), 7.62 (td, 1H, $J = 1.2$ and 8.4 Hz), 7.30 (t, 1H, $J = 8.4$ Hz), 7.19-7.09 (m, 3H), 6.93 (d, 1H, $J = 7.5$ Hz), 6.55 (td, 1H, $J = 1.2$ and 7.5 Hz), 4.43 (bd, 1H, $J = 12.3$ Hz), 4.13 (q, 2H, $J = 6.9$ Hz), 3.7 (bd, 1H, $J = 11.7$ Hz), 3.10-2.92 (m, 2H), 2.67-2.55 (m, 1H), 2.06-1.50 (m, 4H), 1.24 (t, 3H, $J = 6.9$ Hz); ^{19}F NMR (282 MHz, CD_3OD): -47299; LCMS: purity: 99%; MS (m/e): 480(MH^+).

7.3.869 N4-[3-[[4-(Ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926940)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-[3-[[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-4-pyrimidineamine and 3-

hydroxyaniline were reacted to provide N4-[3-[[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.93 (d, 1H, J= 3.6 Hz), 7.89 (t, 1H, J= 1.8 Hz), 7.83 (td, 1H, J= 1.2 and 8.4 Hz), 7.41 (t, 1H, J= 7.8 Hz), 7.11-6.95 (m, 4H), 6.41 (td, 1H, J= 1.8 and 7.2 Hz), 4.44 (bd, 1H, J= 12.9 Hz), 4.10 (q, 2H, J= 7.2 Hz), 3.73 (bd, 1H, J= 12.3 Hz), 3.18-2.98 (m, 2H), 2.67-2.55 (m, 1H), 2.05-1.53 (m, 4H), 1.23 (t, 3H, J= 7.2 Hz); ¹⁹F NMR (282 MHz, CD₃OD): -47483; LCMS: purity: 99%; MS (m/e): 480(MH⁺).

7.3.870 N4-[3-[[4-(Ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926941)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-[3-[[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide N4-[3-[[4-(ethoxycarbonyl)piperidino]carbonyl]phenyl]-5-fluoro-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.95 (d, 1H, J= 3.3 Hz), 7.90 (t, 1H, J= 1.8 Hz), 7.80 (ddd, 1H, J= 0.09, 2.1, 8.1 Hz), 7.39 (t, 1H, J= 7.5 Hz), 7.31 (t, 1H, J= 1.2 Hz), 7.17-7.06 (m, 3H), 6.60-6.54 (m, 1H), 4.48-4.38 (m, 3H), 4.10 (q, 2H, J= 6.9 Hz), 3.78-3.65 (m, 1H), 3.17-2.95 (m, 2H), 2.79 (s, 3H), 2.65-2.53 (m, 1H), 2.01-1.52 (m, 4H), 1.22 (t, 3H, J= 6.9 Hz); ¹⁹F NMR (282 MHz, CD₃OD): -47309; LCMS: purity: 99%; MS (m/e): 551(MH⁺).

7.3.871 Reaction of 3-hydroxyaniline and 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-4-pyrimidineamine

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-4-pyrimidineamine and 3-hydroxyaniline were reacted to provide two products, R926942 and R926943.

7.3.872 N4-(1-Ethoxy-1,2,3,4-tetrahydronaphthalen-7-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926942)

¹H NMR (DMSO-*d*₆): δ 9.23 (bs, 1H), 9.14 (bs, 1H), 8.97 (bs, 1H), 8.04 (d, 1H, J= 3.6 Hz), 7.71 (dd, 1H, J= 2.4 and 7.5 Hz), 7.56 (bs, 1H), 7.14-6.98 (m, 3H), 6.93 (t, 1H, J=

8.1 Hz), 6.29 (bd, 1H, J= 7.2 Hz), 4.35 (bs, 1H), 3.59-3.36 (m, 2H), 2.69-2.60 (m, 2H), 1.89-1.78 (m, 2H), 1.72-1.56 (m, 2H), 1.08 (t, 3H, J= 6.9 Hz); LCMS: purity: 96%; MS (m/e): 395(MH⁺).

5

7.3.873 5-Fluoro-N4-(3,4-dihydronaphthalen-7-yl)-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926943)

¹H NMR (DMSO-*d*₆): δ 9.19 (bs, 2H), 9.01 (s, 1H), 8.04 (d, 1H, J= 3.6 Hz), 7.56-7.46 (m, 2H), 7.16-7.03 (m, 3H), 6.94 (t, 1H, J= 8.1 Hz), 6.46 (d, 1H, J= 9.6 Hz), 6.03 (dd, 1H, J= 1.8 and 8.1 Hz), 6.09-6.01 (m, 1H), 2.69 (t, 2H, J= 8.4 Hz), 2.28-2.20 (m, 2H); ¹⁹F NMR (282 MHz, DMSO-*d*₆): -46541; LCMS: purity: 98%; MS (m/e): 349(MH⁺).

10

7.3.874 5-Fluoro-N4-(3,4-dihydronaphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926944)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N4-(3,4-dihydronaphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 8.07 (d, 1H, J= 3.9 Hz), 7.53-7.45 (m, 2H), 7.32-7.29 (m, 2H), 7.11-7.01 (m, 2H), 6.49-6.40 (m, 2H), 6.08-6.00 (m, 1H), 4.32 (s, 2H), 2.69 (t, 2H, J= 8.4 Hz), 2.62 (s, 3H); LCMS: purity: 99%; MS (m/e): 420(MH⁺).

20

7.3.875 N4-(3-Chloro-4-methoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R926945)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-chloro-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine and 3-hydroxyaniline were reacted to produce N4-(3-chloro-4-methoxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.91 (d, 1H, J= 5.4 Hz), 7.71 (d, 1H, J= 2.4 Hz), 7.58 (dd, 1H, J= 3.0 and 9.0 Hz), 7.15 (t, 1H, J= 8.4 Hz), 7.06 (d, 1H, J= 8.7 Hz), 6.92 (td, 1H, J= 1.8 and 9.9 Hz), 6.88 (t, 1H, J= 1.8 Hz), 6.61 (ddd, 1H, J= 1.2, 2.4, and 8.1 Hz), 3.89 (s, 3H), ; ¹⁹F NMR (282 MHz, CD₃OD): -46612; LCMS: purity: 98%; MS (m/e): 362(MH⁺).

30

7.3.876 N2,N4-Bis(3-chloro-4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926946)

In a like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-chloro-4-methoxyaniline were reacted to provide N2,N4-Bis(3-chloro-4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.90 (bs, 1H), 9.68 (bs, 1H), 8.16 (d, 1H, J= 4.8 Hz), 7.72 (d, 1H, J= 2.4 Hz), 7.65 (d, 1H, J= 2.1 Hz), 7.58 (dd, 1H, J= 2.4 and 9.0 Hz), 7.38 (dd, 1H, J= 2.7 and 9.3 Hz), 7.12 (d, 1H, J= 8.7 Hz), 7.12 (d, 1H, J= 8.7 Hz), 7.05 (d, 1H, J= 8.7 Hz), 3.83 (s, 3H), 3.79 (s, 3H); LCMS: purity: 99%; MS (m/e): 410(MH⁺).

7.3.877 5-Fluoro-N4-(1,2,3,4-tetrahydro-1-oxonaphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926947)

In like manner to the preparation of 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(1,2,3,4-tetrahydro-1-oxo-naphthalen-7-yl)-4-pyrimidineamine and 3-[(N-methylamino)carbonylmethyleneoxy]aniline were reacted to provide 5-fluoro-N4-(1,2,3,4-tetrahydro-1-oxo-naphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.89 (bs, 1H), 9.55 (bs, 1H), 8.17 (d, 1H, J= 4.2 Hz), 8.04-7.93 (m, 3H), 7.32 (d, 1H, J= 8.7 Hz), 7.25-7.16 (m, 2H), 7.09 (t, 1H, J= 7.5 Hz), 6.52 (dd, 1H, J= 2.4 and 8.1 Hz), 4.28 (s, 2H), 2.90 (t, 2H, J= 6.0 Hz), 2.63 (d, 3H, J= 4.8 Hz), 2.59 (t, 2H, J= 6.6 Hz), 2.02 (t, 2H, J= 6.6 Hz); LCMS: purity: 93%; MS (m/e): 436(MH⁺).

7.3.878 5-Fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxyiminonaphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926948)

A solution of 5-fluoro-N4-(1,2,3,4-tetrahydro-1-oxo-naphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (42 mg, 0.095 mmole) and hydroxylamine hydrochloride (8.5 mg, 0.12 mmole) in DMF (1 mL) was heated at 60°C for 12h. The reaction mixture was cooled to rt and then poured into brine (20 mL). A brown solid was collected by suction filtration and further purified by reverse phase chromatography to provide 5-fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxyiminonaphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 8.13-8.05 (m, 2H), 7.99-7.92 (m, 2H), 7.77-7.72 (m, 1H), 7.33-7.21

(m, 2H), 7.14 (d, 1H, J= 8.7 Hz), 7.10-7.02 (m, 1H), 6.47 (dd, 1H, J= 2.4 and 7.5 Hz), 4.30 (s, 2H), 2.90 (t, 1H, J= 6.0 Hz), 2.70-2.40 (m, 6H), 2.07-1.98 (m, 1H), 1.74 (t, 1H, J= 6.6 Hz); LCMS: purity: 96%; MS (m/e): 451(MH⁺).

7.3.879 5-Fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (R926949)

To a 0°C suspension of 5-fluoro-N4-(1,2,3,4-tetrahydro-1-oxo-naphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine (50mg, 0.11 mmol) in anhydrous THF (2.0 mL) was added lithiumborohydride (5 mg, 0.23 mmole).

The reaction mixture was warmed to rt, stirred for 8h, and then quenched with methanol. The reaction mixture was poured into water and then extracted with ethyl acetate.

Purification by preparative TLC (5% methanol/dichloromethane) provided 5-fluoro-N4-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-N2-[3-[(N-methylamino)carbonylmethyleneoxy]phenyl]-2,4-pyrimidinediamine. LCMS: purity: 96%; MS (m/e): 438(MH⁺).

7.3.880 N4-(3-Chloro-4-methoxyphenyl)-5-fluoro-N2-[2-(methoxycarbonyl) benzofuran-5-yl]-2,4-pyrimidinediamine (R926950)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-chloro-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine and 5-amino-2-(methoxycarbonyl)benzofuran were reacted to produce N4-(3-chloro-4-methoxyphenyl)-5-fluoro-N2-[2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.34 (bs, 2H), 8.10-8.07 (m, 2H), 7.78 (t, 1H, J= 2.7 Hz), 7.66-7.53 (m, 4H), 7.12 (d, 1H, J= 9.3 Hz), 3.87 (s, 3H), 3.85 (s, 3H); LCMS: purity: 99%; MS (m/e): 443(MH⁺).

7.3.881 N4-(3-Chloro-4-methoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine (R926951)

In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-chloro-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine and 5-amino-2,3-dihydro-2-(methoxycarbonyl)benzofuran were reacted to produce N4-(3-chloro-4-methoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 10.31

(bs, 1H), 10.04 (bs, 1H), 8.21 (d, 1H, J= 4.8 Hz), 7.75 (t, 1H, J= 3.0 Hz), 7.54 (td, 1H, J= 3.0 and 9.0 Hz), 7.34 (s, 1H), 7.20-7.15 (m, 2H), 6.80 (d, 1H, J= 8.1 Hz), 5.38-5.31 (m, 1H), 3.85 (s, 3H), 3.69 (s, 3H), 3.49 (dd, 1H, J= 11.1 and 16.5 Hz); LCMS: purity: 99%; MS (m/e): 446(MH⁺).

5 **7.3.882 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine (R926953)**

10 In a like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine and 5-amino-2,3-dihydro-2-(methoxycarbonyl)benzofuran were reacted to produce N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-(methoxycarbonyl)benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.99 (bs, 1H), 9.49 (bs, 1H), 8.18 (d, 1H, J= 4.5 Hz), 8.08 (t, 1H, J= 2.4 Hz), 7.81-7.74 (m, 1H), 7.49 (d, 1H, J= 8.1 Hz), 7.42 (s, 1H), 15 7.20 (d, 1H, J= 8.1 Hz), 6.78 (d, 1H, J= 8.7 Hz), 5.36 (m, 1H), 3.80-3.47 (m, 4H), 3.20 (dd, 1H, J= 6.0 and 16.5 Hz); LCMS: purity: 100%; MS (m/e): 500(MH⁺).

7.3.883 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-[(N-methylamino)carbonyl]benzofuran-5-yl]-2,4-pyrimidinediamine (R926954)

20 In a like manner to the preparation of N4-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine, methylamine hydrogen chloride salt, and diisopropylethylamine were reacted to provide N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-[(N-methylamino)carbonyl]benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.59 (s, 1H), 9.10 (s, 2H), 8.13-8.10 (m, 1H), 8.08-7.98 (m, 1H), 7.82 (d, 1H, J= 8.1 Hz), 7.48-7.42 (m, 2H), 7.24 (d, 1H, J= 8.7 Hz), 6.72 (d, 1H, J= 8.7 Hz), 5.06 (dd, 1H, J= 5.4 and 9.3 Hz), 3.39 (dd, 1H, J= 10.5 and 15.6 Hz), 3.15 (dd, 1H, J= 6.3 and 15.9 Hz), 2.59 (d, 3H, J= 4.5 Hz); LCMS: 25 purity: 95%; MS (m/e): 499(MH⁺).

7.3.884 N4-(3-Chloro-4-methoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-[(N-methylamino)carbonyl]benzofuran-5-yl]-2,4-pyrimidinediamine (R926955)

In a like manner to the preparation of N4-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-N4-(3-chloro-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine, methylamine hydrochloride, and diisopropylethylamine were reacted to provide N4-(3-chloro-4-methoxyphenyl)-5-fluoro-N2-[2,3-dihydro-2-[(N-methylamino)carbonyl]benzofuran-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.24 (s, 1H), 8.99 (s, 2H), 8.02 (d, 1H, J= 3.0 Hz), 7.80-7.75 (m, 1H), 7.63 (d, 1H, J= 9.0 Hz), 7.47 (s, 1H), 7.23 (d, 1H, J= 8.1 Hz), 7.07 (d, 1H, J= 8.7 Hz), 6.69 (d, 1H, J= 8.1 Hz), 5.05 (dd, 1H, J= 2.1 and 9.9 Hz), 3.37 (dd, 1H, J= 10.5 and 15.9 Hz), 3.13 (dd, 1H, J= 6.0 and 15.9 Hz), 2.59 (d, 3H, J= 4.5 Hz); LCMS: purity: 95%; MS (m/e): 445(MH⁺).

7.3.885 5-Fluoro-N2-[2,3-dihydro-2-[(N-methylamino)carbonyl]benzofuran-5-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R926956)

In a like manner to the preparation of N4-(3,5-dichloro-4-hydroxyphenyl)-5-fluoro-N2-[3-(N-morpholino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(4-isopropoxyphenyl)-4-pyrimidineamine, methylamine hydrochloride, and diisopropylethylamine were reacted to provide 5-fluoro-N2-[2,3-dihydro-2-[(N-methylamino)carbonyl]benzofuran-5-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.11 (s, 1H), 8.92 (s, 1H), 8.06-7.98 (m, 1H), 7.97 (d, 1H, J= 4.2 Hz), 7.60-7.52 (m, 3H), 7.20 (d, 1H, J= 8.1 Hz), 6.85 (d, 2H, J= 8.7 Hz), 6.67 (d, 1H, J= 9.0 Hz), 5.04 (dd, 1H, J= 5.7 and 9.9 Hz), 4.56 (quintet, 1H, J= 6.6 Hz), 3.36 (dd, 1H, J= 10.5 and 16.5 Hz), 3.10 (dd, 1H, J= 5.7 and 15.3 Hz), 2.59 (d, 1H, J= 4.5 Hz), 1.24 (d, 6H, J= 6.6 Hz); LCMS: purity: 96%; MS (m/e): 438(MH⁺).

7.3.886 N2,N4-Bis(3-phenylphenyl)-2,4-pyrimidinediamine (R925809)

In a like manner to the preparation of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine, 2,4-dichloro-5-fluoropyrimidine and 3-aminobiphenyl were reacted to provide N2,N4-Bis(3-phenylphenyl)-2,4-pyrimidinediamine. LCMS: purity: 98%; MS (m/e): 415(MH⁺).

7.3.887 2-Dimethylamine-5-fluoro-N4-(thyrosinyl methyl ester) pyrimidine (R940110)

A solution of 2,4-dichloro-5-fluoropyrimidine (0.03 g, 0.18 mmol) and L-tyrosine methyl ester (0.14 g, 0.7 mmol) in DMF was heated at 100°C for 3 days. The reaction mixture was cooled to room temperature and diluted with H₂O (10 mL). Upon saturation with sodium chloride it was extracted with ethyl acetate (3 x 15 mL), dried over anhydrous sodium sulfate and the solvent was removed. The resulting residue was filtered through a pad of silica gel (200-400 mesh, hexanes/EtOAc 2/8) to obtain 2-dimethylamine-5-fluoro-N4-(thyrosinyl methyl ester) pyrimidine **R940110**. ¹H NMR (CDCl₃): δ 7.76 (1H, d, *J* = 3.2 Hz), 7.00 (2H, d, *J* = 7.5 Hz), 6.76 (2H, d, *J* = 7.5 Hz), 5.20 (1H, d, *J* = 7.5 Hz), 4.90 (1H, q, *J* = 5.0 Hz), 3.71 (3H, s), 3.14 (2H, m), 3.08 (6H, s); purity: 98%; MS (*m/e*): 335 (*M*+H).

7.3.888 5-Fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine (R940299)

To a solution of 2-chloro-5-fluoro-N4-(3-aminocarbonylphenyl)-4-pyrimidineamine (0.050g, 0.18 mmol) in (2 mL) was added 3-(methylaminocarbonylmethyleneoxy)aniline (0.1g, 0.5 mmol). The mixture was heated in a sealed tube at 100 °C for 24h. The resulting reaction was diluted with H₂O (10 mL), acidified with 2N HCl (pH >2), saturated with sodium chloride and the resulting solid was filtered to give the desired product 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine **R940299**. Purification can be done by filtration through a pad of silica gel using 1-5% MeOH in CH₂Cl₂ or by crystallization using an appropriate solvent system. Alternatively, the reaction of equimolar amount of 2-chloro-5-fluoro-N4-(3-aminocarbonylphenyl)-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline in MeOH in a pressure tube at 110 °C for 24h or, in EtOH using microwave at 175 °C for 30-60 min followed by aqueous work up, also gave the desired product. ¹H NMR (DMSO-*d*₆): δ 9.79 (1H, s), 9.49 (1H, s), 8.26 (1H, d, *J* = 3.9 Hz), 8.15 (1H, t, *J* = 1.8 Hz), 8.10-8.02 (3H, m), 7.68 (1H, d, *J* = 7.5 Hz), 7.51 (1H, t, *J* = 7.9 Hz), 7.48 (1H, s), 7.38 (2H, m), 7.20 (1H, t, *J* = 8.4 Hz), 6.60 (1H, d, *J* = 9.3 Hz), 4.45 (2H, s), 2.74 (3H, d, *J* = 4.8 Hz); purity: 95%; MS (*m/e*): 411 (*M*H⁺).

7.3.889 5-Fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine (R940300)

In like manner to the preparation of 5-fluoro-N2-[3-

5 (methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine **R940300**. ¹H NMR (DMSO-d₆): δ 9.66 (1H, s), 9.45 (1H, s), 8.21 (1H, d, *J* = 3.9 Hz), 8.06 (2H, m), 8.01 (1H, t, *J* = 2.7 Hz), 7.35 (2H, m), 10 7.23 (1H, d, *J* = 9 Hz), 7.18 (1H, t, *J* = 8.1 Hz), 6.60 (1H, d, *J* = 7.8 Hz), 4.45 (2H, s), 3.91 (3H, s), 3.84 (3H, s), 2.74 (3H, d, *J* = 3.6 Hz) ; purity: 93% ; MS (m/e): 456 (MH⁺).

7.3.890 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(3-methyloxycarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine (R940301)

15 In like manner to the preparation of 5-fluoro-N2-[3-

(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)-4-pyrimidineamine and 3-methyloxycarbonyl-4-methoxyaniline were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-(3-methyloxycarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine **R940301**. ¹H NMR (DMSO-d₆): δ 9.93 (1H, s), 9.79 (1H, s), 9.54 (1H, s), 8.26 (1H, s, *J* = 4.5 Hz), 7.92 (1H, s), 7.81 (1H, dd, *J* = 9.3 Hz, *J* = 2.7 Hz), 7.32 (1H, d, *J* = 8.1 Hz), 7.20- 20 7.13 (3H, m), 6.64 (1H, d, *J* = 8.1 Hz), 3.89 (3H, s), 3.84 (3H, s) ; purity: 97% ; MS (m/e): 385 (MH⁺).

7.3.891 5-Fluoro-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-N2-methyl-2,4-pyrimidinediamine (R940304)

25

A mixture of 2-chloro-5-fluoro-N4-(3-methyloxycarbonyl-4-methoxyphenyl)-4-pyrimidineamine (0.15 g, 0.4 mmol), methylamine hydrochloride (0.324 g, 48 mmol) and diisopropylethylamine (0.84 mL, 48 mmol) in MeOH (2 mL) was heated in a sealed tube at 100 °C for 24h (followed by TLC). The reaction was cooled to room temperature and 30 diluted with H₂O (20 mL). The solid was filtered, washed with H₂O and dried to obtain 5-fluoro-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-N2-methyl-2,4-pyrimidinediamine **R940304**. ¹H NMR (DMSO-d₆): δ 10.65 (1H, s), 8.48 (1H, s), 8.29 (2H, m), 7.93 (1H, m),

7.28 (1H, d, $J = 9$ Hz), 4.00 (3H, s), 2.94 (3H, s), 2.90 (3H, d, $J = 4.5$ Hz) ; purity: 90% ; MS (m/e): 306 (MH⁺);

7.3.892 5-Fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine (R940306)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine **R940306**. ¹H NMR (DMSO-d₆): δ 9.28 (1H, s), 9.21 (1H, s), 8.12 (1H, d, $J = 3.9$ Hz), 8.06 (1H, d, $J = 2.7$ Hz), 7.99 (1H, m), 7.89 (1H, dd, $J = 9.3$ Hz, $J = 2.7$ Hz), 7.52 (1H, q, $J = 4.9$ Hz), 7.41 (1H, t, $J = 2.1$ Hz), 7.37 (1H, d, $J = 7.5$ Hz), 7.10 (1H, t, $J = 8.1$ Hz), 6.83 (1H, d, $J = 9$ Hz), 6.53 (1H, dd, $J = 8.1$ Hz, $J = 1.8$ Hz), 4.40 (2H, s), 3.82 (3H, s), 2.96 (3H, d, $J = 5.1$ Hz), 2.73 (3H, d, $J = 4.5$ Hz) ; purity: 93% ; MS (m/e): 455 (MH⁺).

7.3.893 (R)-N2-[3-(dihydroxypropylaminocarbonylmethyleneoxy)phenyl]-5-fluoro-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine (R940307)

In like manner to the preparation of 5-fluoro-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-N2-methyl-2,4-pyrimidinediamine, 5-fluoro-N4-(3-isopropylphenyl)-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and (R)-3-amino-1,2-propanediol were reacted to give (R)-N2-[3-(N-2,3-dihydroxypropylamino)carbonylmethyleneoxyphenyl]-5-fluoro-N4-(3-isopropylphenyl)-2,4-pyrimidinediamine **R940307**. ¹H NMR (DMSO-d₆): δ 9.96 (1H, s), 9.80 (1H, s), 8.29 (1H, d, $J = 4.5$ Hz), 7.98 (1H, t, $J = 5.5$ Hz), 7.77 (1H, d, $J = 7.2$ Hz), 7.57 (1H, s), 7.37 (1H, t, $J = 7.8$ Hz), 7.30-7.22 (3H, m), 7.12 (1H, d, $J = 7.8$ Hz), 6.70 (1H, d, $J = 7.5$ Hz), 4.47 (2H, s), 3.62 (1H, m), 3.38 (3H, m), 3.15 (1H, m), 2.94 (1H, quint, $J = 6.9$ Hz), 1.27 (6H, d, 6.9 Hz) ; purity: 99%; MS (m/e): 469 (M), 470 (MH⁺).

7.3.894 N4-(3-*tert*-Butylphenyl)-5-fluoro-N2-[3-(1,1-dimethyl-2-hydroxyethylaminocarbonylmethyleneoxy)-phenyl]-2,4-pyrimidinediamine (R940308)

In like manner to the preparation of 5-fluoro-N4-(3-methylaminocarbonyl-4-methoxyphenyl)-N2-methyl-2,4-pyrimidinediamine, N4-(3-*tert*-butylphenyl)-5-fluoro-N2-(3-methoxycarbonylmethyleneoxyphenyl)-2,4-pyrimidinediamine and 2-amino-2-methyl-1-propanol were reacted to give N4-(3-*tert*-butylphenyl)-5-fluoro-N2-[3-(1,1-dimethyl-2-hydroxyethylaminocarbonylmethyleneoxy)-phenyl]-2,4-pyrimidinediamine **R940308**. ¹H NMR (DMSO-*d*₆): δ 9.38 (1H, s), 9.28 (1H, s), 8.20 (1H, d, *J* = 3.9 Hz), 7.99 (1H, d, *J* = 7.5 Hz), 7.60 (1H, t, *J* = 2.1 Hz), 7.46 (1H, s), 7.37 (2H, t, *J* = 7.9 Hz), 7.30 (1H, s), 7.19 (2H, t, *J* = 7.9 Hz), 6.56 (1H, dd, *J* = 7.5 Hz, *J* = 1.5 Hz), 5.06 (1H, t, *J* = 5.7 Hz), 4.37 (2H, s), 3.40 (2H, m), 1.36 (9H, s), 1.32 (6H, s) ; purity: 93% ; MS (*m/e*): 482 (MH⁺).

7.3.895 N4-(3-Aminomethylenephenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940309)

A mixture of N4-[3-(*N-tert*-butoxycarbonyl-*N*-aminomethylene)-phenyl]-2-chloro-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline in MeOH was heated in a sealed tube at 100 °C for 12h. The reaction was cool to room temperature and the solvent was removed under reduce pressure. The resulting residue was filtered through a pad of silica gel (200-400 mesh, EtOAc/MeOH (2M NH₃) 95:5) to obtain the desired product N4-(3-aminomethylenephenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine **R940309**. ¹H NMR (DMSO-*d*₆): δ 9.41 (1H, s), 9.23 (1H, s), 8.20 (1H, d, *J* = 3.9 Hz), 8.00 (1H, m), 7.78 (1H, s), 7.72 (1H, d, *J* = 7.2 Hz), 7.46 (1H, s), 7.42-7.33 (2H, m), 7.21 (1H, t, *J* = 7.8 Hz), 7.14 (1H, d, *J* = 7.8 Hz), 6.59 (1H, dd, *J* = 8.1 Hz, *J* = 2.4 Hz), 4.42 (2H, s), 3.79 (2H, s), 2.74 (3H, d, *J* = 4.8 Hz) ; purity: 98% ; MS (*m/e*): 397 (MH⁺).

7.3.896 N4-[3-(2-(N4-(3-aminomethylenephenyl)-5-fluoro-4-pyrimidineamine)-*N*-methylaminomethylene)-phenyl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidineamine (R940311)

A mixture of N4-[3-(*N*-methylaminomethylene)-phenyl]-2-chloro-5-fluoro-4-pyrimidineamine (0.05 g, 0.18 mmol) and 3-(methylaminocarbonylmethyleneoxy)aniline (0.04 g, 0.22 mmol) in EtOH (0.5 mL), was heated at 175 °C for 35 min using microwave. An aqueous work up gave the desired N4-[3-(2-(N4-(3-aminomethylenephenyl)-5-fluoro-4-

pyrimidineamine)-N-methylaminomethylene)-phenyl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidineamine **R940311**. ¹H NMR (DMSO-d₆): δ 9.48 (1H, s), 9.31 (1H, s), 9.26 (1H, s), 8.20 (1H, d, *J* = 3.6 Hz), 8.10-8.05 (4H, m), 7.62 (1H, s), 7.49 (2H, m), 7.41 (1H, t, *J* = 8.1 Hz), 7.36 (2H, m), 7.22 (1H, t, *J* = 8.4 Hz), 7.17 (1H, t, *J* = 8.4 Hz), 7.06 (1H, d, *J* = 7.5 Hz), 6.59 (1H, dd, *J* = 8.4 Hz, *J* = 2.4 Hz), 6.54 (1H, dd, *J* = 7.8 Hz, *J* = 2.4 Hz), 4.93 (2H, s), 4.46 (2H, s), 4.45 (2H, s), 3.28 (3H, d, *J* = 3 Hz), 2.73 (6H, m) ; purity: 98%; MS (m/e) : 684 (M), 685 (MH⁺).

7.3.897 5-Fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-*iso*-propylaminocarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine (R940312)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3- *N-iso*-propylaminomethylene-4-methoxyphenyl)-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-*iso*-propylaminocarbonyl-4-methoxyphenyl)-2,4-pyrimidinediamine **R940312**. ¹H NMR (DMSO-d₆): δ 10.09 (1H, s), 9.88 (1H, s), 8.25 (1H, d, *J* = 4.8 Hz), 8.07 (1H, d, *J* = 2.7 Hz), 8.05 (1H, m), 7.81 (1H, dd, *J* = 9 Hz, *J* = 2.7 Hz), 7.63 (1H, s), 7.25 (2H, m), 7.17 (1H, t, *J* = 8.25 Hz), 6.91 (1H, d, *J* = 9 Hz), 6.68 (1H, d, *J* = 8.1 Hz), 4.42 (2H, s), 3.85 (1H, m), 3.81 (3H, s), 2.72 (3H, d, *J* = 4.2 Hz), 1.30 (6H, d, *J* = 6 Hz) ; purity: 97% ; MS (m/e): 483 (MH⁺).

7.3.898 5-Fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-2,4-pyrimidinediamine (R940314)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, N2-chloro-5-fluoro-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-2,4-pyrimidinediamine **R940314**. ¹H NMR (DMSO-d₆): δ 9.33 (1H, s), 9.21 (1H, s), 8.15 (1H, d, *J* = 3.6 Hz), 8.04 (1H, d, *J* = 4.8 Hz), 7.82 (1H, dd, *J* = 9 Hz, *J* = 2.7 Hz), 7.57 (1H, d, *J* = 3 Hz), 7.47 (1H, t, *J* = 1.95 Hz), 7.34 (1H, m), 7.18 (1H, t, *J* = 8.1 Hz), 7.04 (1H, d, *J* = 9 Hz), 6.56 (1H, dd, *J* = 8.4 Hz, *J* = 2.1

Hz), 4.40 (2H, s), 3.86 (3H, s), 3.63 (4H, t, $J = 4.5$ Hz), 3.53 (2H, s), 2.74 (3H, d, $J = 4.5$ Hz), 2.46 (4H, m) ; purity: 97%; MS (m/e): 497 (MH⁺).

7.3.899 N2-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-2,4-pyrimidinediamine (R940316)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, N2-chloro-5-fluoro-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-4-pyrimidineamine and 4-amino-2-chloro-6-methylphenol were reacted to produce N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-[3-(N-morpholinomethylene)-4-methoxyphenyl]-2,4-pyrimidinediamine R940316. ¹H NMR (DMSO-d₆): δ 9.28 (1H, s), 9.01 (1H, s), 8.65 (1H, s), 8.11 (1H, d, $J = 3.9$ Hz), 7.76 (1H, dd, $J = 9$ Hz, $J = 3$ Hz), 7.61 (1H, d, $J = 2.4$ Hz), 7.50 (1H, d, $J = 2.7$ Hz), 7.30 (1H, d, $J = 2.1$ Hz), 7.04 (1H, d, $J = 8.7$ Hz), 3.87 (3H, s), 3.63 (4H, t, $J = 4.3$ Hz), 3.52 (2H, s), 2.45 (4H, m), 2.17 (3H, s) ; purity: 97% ; MS (m/e): 474 (MH⁺).

7.3.900 N4-(3-N-methylaminomethylenephenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940317)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, N4-[3-(N-*tert*-butoxycarbonyl-N-methylaminomethylene)-phenyl]-2-chloro-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-(3-N-methylaminomethylenephenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine R940317. ¹H NMR (DMSO-d₆): δ 9.41 (1H, s), 9.31 (1H, s), 9.29 (1H, s), 8.20 (1H, d, $J = 3$ Hz), 8.05 (1H, m), 7.80 (1H, d, $J = 7.8$ Hz), 7.74 (1H, s), 7.45-7.35 (3H, m), 7.21 (1H, t, $J = 8.1$ Hz), 7.13 (1H, d, $J = 7.5$ Hz), 6.59 (1H, d, $J = 9.6$ Hz), 4.43 (2H, s), 3.71 (2H, s), 2.75 (3H, d, $J = 4.2$ Hz), 2.35 (3H, s) ; purity: 83.9% ; MS (m/e): 411 (MH⁺).

7.3.901 N2-(3-Chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-[3-(N-piperazinomethylene)-4-methoxyphenyl]-2,4-pyrimidinediamine (R940318)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-

pyrimidinediamine, N4-[3-(N-piperazinomethylene)-4-methoxyphenyl]-2-chloro-5-fluoro-4-pyrimidineamine and 4-amino-2-chloro-6-methylphenol were reacted to produce N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoro-N4-[3-(N-piperazinomethylene)-4-methoxyphenyl]-2,4-pyrimidinediamine **R940318**. ¹H NMR (DMSO-d₆): δ 9.27 (1H, s), 9.00 (1H, s), 8.10 (1H, d, *J* = 3.6 Hz), 7.75 (1H, dd, *J* = 8.7 Hz, *J* = 2.7 Hz), 7.61 (1H, d, *J* = 2.4 Hz), 7.49 (1H, d, *J* = 2.4 Hz), 7.31 (1H, d, *J* = 2.4 Hz), 7.03 (1H, d, *J* = 9 Hz), 3.86 (3H, s), 3.49 (2H, s), 2.75 (4H, t, *J* = 4.65 Hz), 2.39 (4H, m), 2.17 (3H, s); purity: 95%; MS (m/e): 473 (MH⁺).

7.3.902 N4-(3-(N-*tert*-Butoxycarbonyl-N-*iso*-propylaminomethylene)-4-methoxyphenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940319)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, N4-(3-(N-*tert*-butoxycarbonyl-N-*iso*-propylaminomethylene)-4-methoxyphenyl)-2-chloro-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-(3-(N-*tert*-butoxycarbonyl-N-*iso*-propylaminomethylene)-4-methoxyphenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine **R940319**. ¹H NMR (DMSO-d₆): δ 9.44 (1H, s), 8.95 (1H, s), 8.15 (1H, d, *J* = 3.6 Hz), 8.06 (1H, m), 7.83 (1H, m), 7.74 (1H, m), 7.56 (1H, m), 7.37 (1H, m), 7.20 (1H, t, *J* = 7.9 Hz), 7.02 (1H, d, *J* = 9.3 Hz), 6.57 (1H, d, *J* = 7.8 Hz), 4.44 (2H, s), 4.42 (1H, m), 4.33 (2H, s), 3.89 (3H, s), 2.74 (3H, d, *J* = 4.8 Hz), 1.52-1.30 (9H, m), 1.16 (6H, d, *J* = 6.9 Hz); purity: 98%; MS (m/e): 569 (MH⁺).

7.3.903 N4-(3-N,N-Dimethylaminomethylene-4-methoxyphenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940321)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-(3-N,N-dimethylaminomethylene-4-methoxyphenyl)-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-(3-N,N-dimethylaminomethylene-4-methoxyphenyl)-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine **R940321**.

¹H NMR (DMSO-d₆): δ 9.32 (1H, s), 9.23 (1H, s), 8.14 (1H, d, *J* = 3.9 Hz), 8.05 (1H, m), 7.83 (1H, dd, *J* = 8.7 Hz, *J* = 2.4 Hz), 7.55 (1H, d, *J* = 2.4 Hz), 7.45 (1H, s), 7.36 (1H, d, *J* = 8.4 Hz), 7.18 (1H, t, *J* = 8.1 Hz), 7.03 (1H, d, *J* = 9 Hz), 6.56 (1H, dd, *J* = 7.2 Hz, *J* = 1.5 Hz), 4.41 (2H, s), 3.86 (3H, s), 2.73 (3H, d, *J* = 4.5 Hz), 2.24 (6H, s); purity: 91.8%;
 5 MS (m/e): 455 (MH⁺).

7.3.904 N4-[(2,2-Dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940323)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine **R940323**.

¹H NMR (DMSO-d₆): δ 10.70 (1H, s), 9.45 (1H, s), 9.19 (1H, s), 8.17 (1H, d, *J* = 3.9 Hz), 8.05 (1H, m), 7.43-7.34 (4H, m), 7.17 (1H, t, *J* = 8.25 Hz), 6.98 (1H, d, *J* = 8.4 Hz), 6.56 (1H, dd, *J* = 7.8 Hz, *J* = 2.1 Hz), 4.25 (2H, s), 2.74 (3H, d, *J* = 4.5 Hz), 1.5 (6H, s); purity: 98.7%; MS (m/e): 467 (MH⁺).

7.3.905 N4-[3-Dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940337)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[3-dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-[3-dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine

R940337. ¹H NMR (DMSO-d₆): δ 9.28 (1H, s), 9.20 (1H, s), 8.34 (1H, dd, *J* = 4.8 Hz, *J* = 1.2 Hz), 8.14 (1H, d, *J* = 3.8 Hz), 8.03 (1H, m), 7.64-7.60 (2H, m), 7.51-7.46 (3H, m), 7.37 (1H, d, *J* = 8.4 Hz), 7.17 (1H, t, *J* = 8.1 Hz), 6.94-6.91 (2H, m), 6.55 (1H, dd, *J* = 8.4 Hz, *J* = 3 Hz), 4.42 (2H, s), 3.93 (2H, s), 2.74 (3H, d, *J* = 4.5 Hz), 1.32 (6H, s); purity: 98.2%; MS (m/e): 530 (MH⁺);

7.3.906 N4-[(2,2-Dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(1-methylindazolin-5-yl)-2,4-pyrimidinediamine (R940338)

In like manner to the preparation of 5-fluoro-N2-[3-

5 (methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 5-amino-1-methyl-1-indazole were reacted to produce N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(1-methylindazolin-5-yl)-2,4-pyrimidinediamine **R940338**. ¹H NMR (DMSO-d₆): δ 10.73 (1H, s), 9.39 (1H, s), 9.17 (1H, s), 8.21 (1H, s), 8.16 (1H, d, *J* = 3.9 Hz), 7.87 (1H, s), 7.56 (2H, m), 7.41 (1H, m), 7.32 (1H, s), 7.00 (1H, d, *J* = 8.4 Hz), 4.07 (3H, s), 1.51 (6H, s); purity: 99.2%; MS (*m/e*): 434 (MH⁺).

7.3.907 N4-[(2,2-Difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R921303)

In like manner to the preparation of 5-fluoro-N2-[3-

15 (methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine **R921303**. ¹H NMR (DMSO-d₆): δ 12.05 (1H, s), 9.67 (1H, s), 9.27 (1H, s), 8.24 (1H, d, *J* = 3.6 Hz), 8.05 (1H, m), 7.73-7.68 (1H, m), 7.56 (1H, t, *J* = 2.7 Hz), 7.50 (1H, s), 7.36 (2H, d, *J* = 8.7 Hz), 7.19 (1H, t, *J* = 8.2 Hz), 6.58 (1H, dd, *J* = 8.4 Hz, *J* = 2.4 Hz), 4.34 (2H, s), 2.74 (3H, d, *J* = 4.5 Hz); ¹⁹F NMR (DMSO-d₆): δ -21643, -46385; purity: 100%; MS (*m/e*): 475 (MH⁺).

7.3.908 N4-[(2,2-Dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-7-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940345)

In like manner to the preparation of 5-fluoro-N2-[3-

30 (methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-7-yl]-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-[(2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-7-yl]-5-fluoro-N2-[3-

(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine **R940345**. ¹H NMR (DMSO-d₆): δ 11.23 (1H, s), 9.69 (1H, s), 9.54 (1H, s), 8.50 (1H, s), 8.25 (1H, d, *J* = 3.3 Hz), 8.06 (1H, m), 7.96 (1H, t, *J* = 2.5 Hz), 7.41-7.36 (2H, m), 7.24 (1H, t, *J* = 8.25 Hz), 6.34 (1H, d, *J* = 8.7 Hz), 4.47 (2H, s), 2.74 (3H, d, *J* = 3.3 Hz), 1.53 (6H, s); purity: 98.4%; MS (m/e): 468 (MH⁺).

7.3.909 N4-[(2,2-Dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940346)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to produce N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine **R940346**. ¹H NMR (DMSO-d₆): δ 10.75 (1H, s), 8.25 (1H, d, *J* = 4.5 Hz), 7.42-7.37 (1H, m), 7.34 (1H, s), 7.10 (3H, m), 7.00 (1H, d, *J* = 8.4 Hz), 6.53 (1H, m), 1.50 (6H, s); purity: 97.5%; MS (m/e): 396 (MH⁺).

7.3.910 N4-[(2,2-Dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R940347)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 3-(methylaminocarbonylmethyleneoxy)aniline were reacted to produce N4-[(2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine **R940347**. ¹H NMR (DMSO-d₆): δ 11.20 (1H, s), 9.46 (1H, s), 8.26 (1H, d, *J* = 3.6 Hz), 8.06 (1H, s), 7.71 (1H, m), 7.49 (1H, d, *J* = 8.4 Hz), 7.45 (1H, s), 7.38 (1H, d, *J* = 9 Hz), 7.21 (1H, t, *J* = 8.1 Hz), 6.61 (1H, d, *J* = 8.7 Hz), 4.47 (2H, s), 2.74 (3H, s), 1.52 (6H, s); purity: 100%; MS (m/e): 468 (MH⁺).

7.3.911 N4-[3-Dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940348)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-

pyrimidinediamine, 2-chloro-N4-[3-dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to produce N4-[3-dihydro-2,2-dimethyl-4-(2-pyridyl)-benzo[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine **R940348**. ¹H NMR (DMSO-d₆): δ 9.25 (1H, s), 9.23 (1H, s), 9.02 (1H, s), 8.34 (1H, d, *J* = 4.5 Hz), 8.11 (1H, d, *J* = 3.3 Hz), 7.62 (2H, m), 7.52 (2H, m), 7.22 (1H, s), 7.19 (1H, d, *J* = 7.5 Hz), 7.03 (1H, t, *J* = 7.9 Hz), 6.93 (2H, m), 6.38 (1H, d, *J* = 7.8 Hz), 3.93 (2H, s), 1.32 (6H, s) ; purity: 96.5%.

7.3.912 N4-[(2,2-Difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R940349)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 3-aminophenol were reacted to produce N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine **R940349**. ¹H NMR (DMSO-d₆): δ 12.03 (1H, s), 9.63 (1H, s), 9.26 (1H, s), 9.09 (1H, s), 8.21 (1H, d, *J* = 3.6 Hz), 7.70 (1H, dd, *J* = 9 Hz, *J* = 2.4 Hz), 7.59 (1H, d, *J* = 2.7 Hz), 7.34 (1H, d, *J* = 9.3 Hz), 7.26 (1H, s), 7.16 (1H, d, *J* = 7.8 Hz), 7.04 (1H, t, *J* = 8.2 Hz), 6.41 (1H, d, *J* = 10.2 Hz) ; ¹⁹F NMR (DMSO-d₆): δ -21646, -46516 ; purity: 95.8% ; MS (m/e): 404 (MH⁺);

7.3.913 N2,N4-[(2,2-Dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-2,4-pyrimidinediamine (R940350)

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-5-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 6-amino-2,2-dimethyl-4H-benzo[1,4]oxazin-3-one were reacted to produce N2,N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-2,4-pyrimidinediamine **R940350**. ¹H NMR (DMSO-d₆): δ 10.68 (1H, s), 10.62 (1H, s), 9.38 (1H, s), 9.04 (1H, s), 8.11 (1H, d, *J* = 3.6 Hz), 7.46 (1H, dd, *J* = 8.1 Hz, *J* = 1.8 Hz), 7.33-7.26 (3H, m), 6.95 (1H, d, *J* = 8.7 Hz), 6.84 (1H, d, *J* = 8.4 Hz), 1.49 (6H, s), 1.45 (6H, s) ; purity: 95.4% ; MS (m/e): 479 (MH⁺).

7.3.914 N2-[(2,2-Difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-2,4-pyrimidinediamine (R940351)

In like manner to the preparation of 5-fluoro-N2-[3-

- 5 (methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-5-pyrido[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 6-amino-2,2-difluoro-4H-benzo[1,4]oxazin-3-one were reacted to produce N2-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-2,4-pyrimidinediamine **R940351**. ¹H NMR (DMSO-d₆): δ 11.99 (1H, s), 10.74 (1H, s), 9.64 (1H, s), 9.50 (1H, s), 8.19 (1H, d, *J* = 3.9 Hz), 7.50 (2H, m), 7.43 (1H, dd, *J* = 8.4 Hz, *J* = 1.8 Hz), 7.32 (1H, s), 7.20 (1H, d, *J* = 9.3 Hz), 6.98 (1H, d, *J* = 8.7 Hz), 1.49 (6H, s) ; purity: 94.77% ; MS (m/e): 487 (MH⁺).
- 10

7.3.915 N2,N4-[(2,2-Difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-2,4-pyrimidinediamine (R940352)

In like manner to the preparation of 5-fluoro-N2-[3-

- 15 (methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and 6-amino-2,2-difluoro-4H-benzo[1,4]oxazin-3-one were reacted to produce N2,N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-2,4-
- 20 pyrimidinediamine **R940352**. ¹H NMR (DMSO-d₆): δ 12.08 (1H, s), 12.00 (1H, s), 9.72 (1H, s), 9.44 (1H, s), 8.23 (1H, d, *J* = 3.6 Hz), 7.73 (1H, dd, *J* = 11.1 Hz, *J* = 1.5 Hz), 7.6 (1H, s), 7.56 (1H, s), 7.51 (1H, dd, *J* = 9.6 Hz, *J* = 2.4 Hz), 7.35 (1H, d, *J* = 9 Hz), 7.24 (1H, d, *J* = 8.7 Hz); ¹⁹F NMR (DMSO-d₆): δ -21670, -21722, -4651 ; purity: 100% ; MS (m/e): 495 (MH⁺).

7.3.916 N4-[(2,2-Difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine (R940353)

In like manner to the preparation of 5-fluoro-N2-[3-

- 30 (methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and methyl 5-aminobenzofuran-2-carboxylate were reacted to produce N4-[(2,2-difluoro-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine **R940353**. ¹H NMR (DMSO-d₆):

δ 12.05 (1H, s), 9.69 (1H, s), 9.43 (1H, s), 8.28 (1H, s), 8.25 (1H, d, $J = 3.6$ Hz), 7.40-7.64 (4H, m), 7.54 (1H, s), 7.38 (1H, d, $J = 9$ Hz), 3.97 (3H, s); ^{19}F NMR (DMSO- d_6): δ -21707, -46489; purity: 97.77%; MS (m/e): 486 (MH $^+$).

5 **7.3.917 N4-[(2,2-Dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine (R940354)**

In like manner to the preparation of 5-fluoro-N2-[3-(methylaminocarbonylmethyleneoxy)phenyl]-N4-(3-aminocarbonylphenyl)-2,4-pyrimidinediamine, 2-chloro-N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-4-pyrimidineamine and methyl 5-aminobenzofuran-2-carboxylate were reacted to produce N4-[(2,2-dimethyl-4H-benzo[1,4]oxazin-3-one)-6-yl]-5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine **R940354**. ^1H NMR (DMSO- d_6): δ 10.75 (1H, s), 9.67 (1H, s), 9.53 (1H, s), 8.25 (1H, s), 8.21 (1H, d, $J = 4.2$ Hz), 7.66 (2H, s), 7.59 (1H, s), 7.31 (1H, d, $J = 8.7$ Hz), 7.26 (1H, s), 7.03 (1H, d, $J = 8.1$ Hz), 3.97 (3H, s), 1.52 (6H, s); purity: 95.58%; MS (m/e): 478 (MH $^+$).

7.3.918 N2,N4-Bis(3-N-acetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950244)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour. The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl_3 :Acetone, 2:1) to give N2,N4-bis(3-N-acetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine. ^1H NMR (MeOD, 300 MHz): δ 8.65 (d, 1H, $J = 2.4$ Hz), 7.15-7.58 (m, 8H), 2.24 (s, 3H), 2.22 (s, 3H), 2.14 (s, 3H), 2.09 (s, 3H); LCMS: ret. time: 17.03 min.; purity: 87.0%; MS (m/e): 478.89 (MH $^+$).

25 **7.3.919 N4-(3-N,N-Diacetylaminophenyl)-N2-(3-N-acetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950245)**

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour. The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl_3 :Acetone, 2:1) to give N4-(3-N,N-diacetylaminophenyl)-N2-(3-N-acetylaminophenyl)-5-fluoro-N2,N4-

pyrimidinediacetylamine. ^1H NMR (MeOD, 300 MHz): δ 8.65 (d, 1H, $J = 2.4$ Hz), 7.03-7.66 (m, 8H), 2.21 (s, 6H), 2.14 (s, 3H), 2.12 (s, 3H), 2.08 (s, 3H); LCMS: ret. time: 19.27 min.; purity: 92.6%; MS (m/e): 521.01 (MH^+).

5 **7.3.920 N4-(3-N-Acetylaminophenyl)-N2-(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950246)**

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour. The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl_3 :Acetone, 2:1) to give N4-[3-N-acetylaminophenyl]-N2-(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine. ^1H NMR (MeOD, 300 MHz): δ 8.66 (d, 1H, $J = 2.4$ Hz), 6.88-7.57 (m, 8H), 2.22 (s, 6H), 2.11 (s, 3H), 2.10 (s, 3H), 2.09 (s, 3H); LCMS: ret. time: 18.89 min.; purity: 83.0%; MS (m/e): 520.97 (MH^+).

15 **7.3.921 N2,N4-Bis(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950247)**

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour. The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl_3 :Acetone, 2:1) to give N2,N4-bis(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine. ^1H NMR (MeOD, 300 MHz): δ 8.58 (d, 1H, $J = 2.4$ Hz), 6.75-7.53 (m, 8H), 2.04 (s, 3H), 2.03 (s, 3H), 2.01 (s, 6H), 1.99 (s, 6H); LCMS: ret. time: 21.51 min.; purity: 91.8%; MS (m/e): 563.00 (MH^+).

25 **7.3.922 N4-(3-Nitrophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950261)**

A mixture of equimolar amounts of 2-chloro-N4-(3-nitrophenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-nitrophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 92.7%; MS (m/e): 412.94 (MH^+).

7.3.923 N4-(3-Aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine HCl salt (R950262)

N4-(3-Nitrophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and Pd/C 10% (50% water content) were suspended in EtOH-10% aqueous HCl (1 : 1) and hydrogenated in a Parr apparatus for 2 hours (22 °C, 50 psi). The suspension was filtered over celite and carefully washed with MeOH. The combined filtrates were concentrated under reduced pressure to give the HCl salt of N4-(3-aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 92.7%; MS (m/e): 383.07 (M-Cl⁺, 100).

7.3.924 N4-(3-Aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950263)

The HCl salt of N4-(3-aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine was neutralized with aqueous sodium carbonate solution and extracted with EtOAc. The organic phase was dried and concentrated to give N4-(3-aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a pale yellow solid. ¹H NMR (DMSO): δ 10.00 (s, 1H), 9.92 (s, 1H), 8.07 (d, 1H, J= 2.4 Hz), 8.15 (bs, 2H), 7.91-8.07 (m, 3H), 7.08-7.21 (m, 5H), 6.56 (d, 1H, J = 7.2 Hz), 4.32 (s, 2H), 2.72 (d, 3H, J = 4.8 Hz); LCMS: purity: 92.7%; MS (m/e): 383.17 (MH⁺, 100).

7.3.925 N4-(3-Bis-N-methylaminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950264)

A solution of N4-(3-aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in DME-DMF (1 : 1) was treated with 10 equivalents of MeI and sodium bicarbonate. The mixture was stirred for 1.5 hours at 70°C and purified by flash chromatography on silica gel to give N4-(3-bis-N-methylaminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 90.2%; MS (m/e): 411.04 (MH⁺, 100).

7.3.926 N4-(3-N-Hydroxyethylaminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950265)

A solution of N4-(3-aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in DME-DMF (1 : 1) was treated with 10 equivalents of 2-bromoethanol and sodium bicarbonate. The mixture was stirred for 16 hours at 70°C and purified by flash chromatography on silica gel to give N4-(3-N-hydroxyethylaminophenyl)-5-fluoro. LCMS: purity: 90.2%; MS (m/e): 427.33 (MH⁺, 100).

7.3.927 N4-(3-Bis(N-hydroxyethyl)aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950266)

A solution of N4-(3-aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in DME-DMF (1 : 1) was treated with 10 equivalents of 2-bromoethanol and sodium bicarbonate. The mixture was stirred for 16 hours at 70°C and purified by flash chromatography on silica gel to give N4-(3-bis(N-hydroxyethyl)aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 94.2%; MS (m/e): 471.46 (MH⁺, 100).

7.3.928 N4-(3-N-Methylaminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950267)

A solution of N4-(3-aminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in DME-DMF (1 : 1) was treated with 10 equivalents of MeI and sodium bicarbonate. The mixture was stirred for 1.5 hours at 70°C and purified by flash chromatography on silica gel to give N4-(3-N-methylaminophenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 92.3%; MS (m/e): 397.02 (MH⁺, 100).

7.3.929 N4-(3-Carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950290)

A mixture of equimolar amounts of 2-chloro-N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-4-aminopyridine and 3-ethoxycarbonylmethyleneoxyaniline in MeOH in a pressure

tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 97.8%; MS (m/e): 443.20 (MH⁺).

5 **7.3.930 N4-(3-Carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-carboxymethyleneoxyphenyl]-2,4-pyrimidinediamine (R950291)**

10 The reaction of N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (0.1 g) and LiOH (10 equivalents) in MeOH:water (1:1, v/v) for 1h at room temperature followed by treatment with aqueous HCl gave the solid. The resulting solid was filtered, washed with water and dried to give N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-carboxymethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 91.5%; MS (m/e): 415.16 (MH⁺).

15 **7.3.931 N4-(3-Methoxycarbonyl-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950293)**

20 A solution of N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeOH was treated with a 4 M solution of HCl in dioxane. The mixture was stirred for 1 hour at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(3-methoxycarbonyl-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 96.8%; MS (m/e): 457.25 (MH⁺).

25 **7.3.932 N4-(4-Methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950294)**

30 A mixture of equimolar amounts of 2-chloro-N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-4-aminopyridine and 3-ethoxycarbonylmethyleneoxyaniline in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 92.1%; MS (m/e): 469.26 (MH⁺).

7.3.933 N4-(4-Methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950295)

A mixture of equimolar amounts of 2-chloro-N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-4-aminopyridine and 3-ethoxycarbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h followed by aqueous work up gave N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 87.6%; MS (m/e): 455.26 (MH⁺).

7.3.934 N4-(4-Ethoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950296)

A solution of N4-(4-ethoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in EtOH was treated with the HCl salt of methylamine. The mixture was stirred for 4 hours at 100°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-ethoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 87.4%; MS (m/e): 468.29 (MH⁺).

7.3.935 N4-(4-Carboxyethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950344)

A mixture of equimolar amounts of 2-chloro-N4-(4-carboxyethyleneoxyphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(4-carboxyethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 97.8%; MS (m/e): 456.32 (MH⁺).

7.3.936 N4-(2,3-Dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950345)

A solution of N4-(4-Methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in TFOH was heated

for 2 hours at 100°C. Aqueous work up followed by flash chromatography on silica gel gave N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 92.2%; MS (m/e): 435.95 (MH⁺).

5 **7.3.937 N4-(4-Methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950346)**

10 A solution of N4-(4-carboxyethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeOH was treated with a 4 M solution of HCl in dioxane. The mixture was stirred for 1 hour at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 85.2%; MS (m/e): 468.01 (MH⁺).

15 **7.3.938 N4-(4-Hydroxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950347)**

20 The reaction of N4-(4-methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and LiOH (10 equivalents) in MeOH:water (1:1, v/v) for 1h at room temperature followed by treatment with aqueous HCl gave a pale yellow solid. The resulting solid was filtered, washed with water and dried to give N4-(4-hydroxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 94.7%; MS (m/e): 382.03 (MH⁺).

25 **7.3.939 N4-(2,3-Dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950348)**

30 A mixture N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(2,3-dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 89.5%; MS (m/e): 451.00 (MH⁺).

7.3.940 N4-(4-Hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950349)

A solution of N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeOH was treated with a sodiumcyanoborohydride. The mixture was stirred for 1 hour at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-Hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.19 (s, 1H), 9.09 (s, 1H), 8.03 (d, 1H, J = 2.4 Hz), 7.28-7.93 (m, 5H), 7.07 (t, 1H, J = 7.2 Hz), 6.71 (d, 1H, J = 7.2 Hz), 6.44 (dd, 1H, J = 2.6, 7.2 Hz), 5.31 (d, 1H, J = 5.1 Hz), 4.14-4.59 (m, 3H), 4.30 (s, 2H), 2.63 (d, 3H, J = 4.8 Hz), 1.82-2.03 (m, 2H); LCMS: purity: 93.3%; MS (m/e): 440.15 (MH⁺).

7.3.941 N4-(2,3-Dihydro-4-O-methyloxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950356)

A mixture N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and methoxyamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(2,3-dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 85.5%; MS (m/e): 465.10 (MH⁺).

7.3.942 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950368)

A mixture N4-(4-azido-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and Pd/C (10%) in MeOH was hydrogenated at 22°C for 6 hours (40psi). The mixture was filtered and concentrated to dryness to give N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.60 (s, 1H), 9.46 (s, 1H), 8.73 (bs, 3H), 8.00-8.10 (m, 3H), 7.47 (s, 1H), 7.42 (m, 1H), 7.29 (d, 1H, J = 7.2 Hz), 7.11 (t, 1H, J = 7.2 Hz), 6.82 (d, 1H, J 7.0 Hz), 6.46 (m, 1H), 4.23-4.46 (m, 3H), 4.31 (s, 3H), 2.63 (d, 3H, J = 4.8 Hz), 2.09-2.29 (m, 2H); LCMS: purity: 97.6%; MS (m/e): 438.98 (MH⁺).

7.3.943 N4-(3-Methylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950371)

A mixture of equimolar amounts of 2-chloro-N4-(3-methylcarbonylphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-methylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 10.16 (s, 1H), 9.82 (s, 1H), 8.24 (d, 1H, J = 2.4 Hz), 8.15 (s, 1H), 7.91-8.07 (m, 2H), 7.70 (d, 1H, J = 7.0 Hz), 7.49 (t, 1H, J = 7.2 Hz), 7.08-7.21 (m, 3H), 6.56 (d, 1H, J = 7.2 Hz), 4.30 (s, 3H), 2.62 (d, 3H, J = 4.8 Hz), 2.48 (s, 3H); LCMS: purity: 93.8%; MS (m/e): 410.50 (MH⁺).

7.3.944 N4-(3-Phenylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950372)

A mixture of equimolar amounts of 2-chloro-N4-(3-phenylcarbonylphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-phenylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 86.0%; MS (m/e): 472.50 (MH⁺).

7.3.945 N4-(3-Methylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950373)

A mixture N4-(3-methylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(3-methylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 11.21 (s, 1H), 10.11 (s, 1H), 9.85 (s, 1H), 6.54-8.23 (m, 9H), 4.32 (s, 2H), 2.63 (d, J = 7.0 Hz, 3H), 2.47 (s, 3H); LCMS: purity: 92.4%; MS (m/e): 425.28 (MH⁺).

7.3.946 N4-(3-Phenylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950374)

A mixture N4-(3-phenylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(3-phenylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 11.63 (s, 1H), 10.30 (s, 1H), 9.85 (s, 1H), 6.44-8.43 (m, 14H), 4.42 (s, 2H), 2.63 (d, J = 7.0 Hz, 3H); LCMS: purity: 92.4%; MS (m/e): 487.31 (MH⁺).

7.3.947 N2,N4-Bis(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950376)

A mixture of 2,4-dichloro-5-fluoropyridine and three equivalents of 3-acetophenone in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N2,N4-bis(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 93.1%; MS (m/e): 365.19 (MH⁺).

7.3.948 N2,N4-Bis(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950377)

A mixture of 2,4-dichloro-5-fluoropyridine and three equivalents of 3-benzophenone in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N2,N4-bis(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 95.7%; MS (m/e): 489.29 (MH⁺).

7.3.949 N2,N4-Bis(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-2,4-pyrimidinediamine (R950378)

A solution of N2,N4-bis(4-methoxycarbonylethyleneoxyphenyl)-5-fluoro-2,4-pyrimidinediamine in TfOH was heated for 2 hours at 100°C. Aqueous work up followed by flash chromatography on silica gel gave N2,N4-bis(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.36 (s, 1H), 9.14 (s, 1H), 8.06 (d, 1H, J = 2.4 Hz), 7.72-7.99 (m, 3H), 6.97 (d, J = 7.2 Hz, 1H), 6.87 (d, J = 7.2 Hz, 1H), 4.42-4.52 (m, 4H), 2.70-2.78 (m, 4H); LCMS: purity: 94.3%; MS (m/e): 484.50 (MH⁺).

7.3.950 N2,N4-Bis(3-methylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine (R950379)

A mixture of N2,N4-bis(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N2,N4-bis(3-methylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 11.21 (s, 1H), 10.11 (s, 1H), 9.85 (s, 1H), 6.54-8.23 (m, 9H), 4.32 (s, 2H), 2.63 (d, J = 7.0 Hz, 3H), 2.47 (s, 3H); LCMS: purity: 92.1%; MS (m/e): 393.06 (M-H⁺).

7.3.951 N2,N4-Bis(3-phenylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine (R950380)

A mixture of N2,N4-bis(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N2,N4-bis(3-phenylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 93.3%; MS (m/e): 486.05 (M-H⁺).

7.3.952 N2,N4-Bis(2,3-dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-2,4-pyrimidinediamine (R950381)

A mixture of N2,N4-bis(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N2,N4-bis(2,3-dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 98.1%; MS (m/e): 449.03 (M-H⁺).

7.3.953 N4-(4-Acetyloxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950382)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in pyridine was treated with acetic anhydride at 22°C for 16 hours. Aqueous work up gave N4-(4-acetyloxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 10.43 (bs, 1H), 9.62 (bs, 1H), 8.03 (d, 1H, J = 2.4 Hz), 7.10-7.83 (m, 7H), 6.83 (d, 1H, J = 7.4 Hz), 6.52 (d, 1H, J = 7.2 Hz), 5.01 (m, 1H), 4.75 (s, 2H), 4.03-4.32 (m, 2H), 2.62 (s, 3H), 2.23 (s, 3H), 1.93-2.13 (m, 2H); LCMS: purity: 92.1%; MS (m/e): 393.06 (M-H⁺).

7.3.954 N4-(4-Azido-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950383)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry THF was treated with 2 equivalents of DPPA and DBU. The mixture was stirred for 3 hours at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-azido-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 10.09 (bs, 1H), 9.83 (bs, 1H), 8.18 (d, 1H, J = 2.4 Hz), 7.97 (m, 1H), 7.11-7.61 (m, 6H), 6.82 (d, 1H, J = 7.8 Hz), 6.62 (d, 1H, J = 7.2 Hz), 4.78 (s, 2H), 4.03-4.33 (m, 3H), 2.62 (s, 3H), 1.93-2.13 (m, 2H); LCMS: purity: 97.9%; MS (m/e): 463.07 (MH⁺).

7.3.955 N4-(4-Benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950385)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in THF was treated with boron trifluoride etherate at 80°C for 8 hours. Aqueous work up gave N4-(4-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.18 (s, 1H), 9.14 (s, 1H), 8.03 (d, J = 2.4 Hz, 1H), 7.93 (bs, 1H), 5.86-7.48 (m, 9H), 4.73-4.74 (m, 2H), 4.33 (s, 2H), 2.62 (s, 3H); LCMS: purity: 96.5%; MS (m/e): 420.07 (M-H⁺).

7.3.956 N4-(3-Hydroxymethylen-4-methoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950386)

A mixture of equimolar amounts of 2-chloro-N4-(3-hydroxymethylen-4-methoxyphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-hydroxymethylen-4-methoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 89.2%; MS (m/e): 410.5 (MH⁺).

7.3.957 N4-(3-Amino-4-ethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950388)

A mixture of 2-chloro-N4-(3-amino-4-ethoxyphenyl)-5-fluoro-4-aminopyridine and 3 equivalents of 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-amino-4-ethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 95.1%; MS (m/e): 427.18 (MH⁺).

7.3.958 N4-(4-Ethoxy-3-hydroxysulfonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950389)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in HOAc was treated with sodium nitrate followed by addition of concentrated aqueous HCl and copper dichloride. The mixture was stirred for 2 hours at 22°C for 8 hours and purified by aqueous work up followed by column chromatography on silica gel to give N4-(4-ethoxy-3-hydroxysulfonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 82.3%; MS (m/e): 474.09 (M-H⁻).

7.3.959 N2,N4-Bis(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950391)

A mixture of 2,4-dichloro-5-fluoropyridine and three equivalents of 3-methoxycarbonyl-4-trifluoromethoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up N2,N4-bis(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.96 (s, 1H), 9.82 (s, 1H), 8.16-8.26 (m, 4H), 7.91 (dd, 1H, J = 3.0, 7.2 Hz), 7.42 (d, 1H, J = 7.2 Hz), 7.31 (d, 1H, J = 7.2 Hz), 3.77 (s, 3H), 3.75 (s, 3H); LCMS: purity: 93.0%; MS (m/e): 565.37 (MH⁺).

7.3.960 N4-(3-Methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950392)

A mixture of equimolar amounts of 2-chloro-N4-(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-4-aminopyridine and 3-(N-

methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid.

5 LCMS: purity: 95.8%; MS (m/e): 510.41 (MH⁺).

7.3.961 N4-(4-Acetylamino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950393)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeCN was treated with concentrated sulfuric acid. The mixture was stirred for 3 hours at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-acetylamino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H
10 NMR (DMSO): δ 10.46 (bs, 1H), 9.52 (bs, 1H), 7.98 (d, 1H, J = 2.4 Hz), 7.12-7.73 (m, 7H), 6.66 (d, 1H, J = 7.2 Hz), 6.49 (d, 1H, J = 7.2 Hz), 4.75 (s, 2H), 4.03-4.32 (m, 2H), 3.80 (m, 1H), 2.64 (s, 3H), 2.143 (s, 3H), 1.90-2.11 (m, 2H); LCMS: purity: 92.1%; MS (m/e): 393.06 (M-H⁺). LCMS: purity: 96.2%; MS (m/e): 479.13 (M-H⁺).

7.3.962 N4-[2,4-Dihydro-1-oxo-4H-imidazo[2,1-c][1,4]benzoxazin-8-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R945236)

N4-[2H-1,4-Benzoxazin-3(4H)-one-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (800 mg, 2.18 mmol) and phosphorus pentasulfide (800 mg, 1.80 mmol) were stirred in pyridine (5 mL) at 70 °C for 2h. The reaction solution was treated with 1N
25 HCl solution to pH 5. The precipitation was collected with filtration, washed with water, dried to give N4-[2H-1,4-benzoxazin-3(4H)-thione-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine.

N4-[2H-1,4-Benzoxazin-3(4H)-thione-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (400 mg, 1.04 mmol), glycine (500 mg) and triethylamine (0.5 mL)
30 were stirred in methanol (10 mL) at 70 °C overnight. The undissolved salt was filtered off, washed with methanol. The filtrate was evaporated and redissolved in THF (5mL) and DMF (5 mL). To the solution were added EDC (200 mg), HOAt (200 mg) and diisopropylethylamine (0.2 mL). The reaction solution was stirred at 70 °C for 0.5 h. The

mixture was diluted with ethyl acetate (60 mL) and washed with water (2 x 60 mL). The organic layer was separated, dried, evaporated and purified by flash column chromatography (EtOAc/hexanes = 1:1, EtOAc) to give N4-[2,4-dihydro-1-oxo-4H-imidazo[2,1-c][1,4]benzoxazin-8-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-

5 pyrimidinediamine as a white solid. ¹H NMR (CDCl₃): δ 4.35 (t, J= 2.1 Hz, 2H), 4.92 (t, J= 2.1 Hz, 2H), 6.44 (dd, J= 1.5 and 8.1 Hz, 1H), 6.81 (m, 2H), 6.99 (s, 1H), 7.11 (m, 2H), 7.39 (m, 2H), 7.97 (d, J= 3.0 Hz, 1H), 8.02 (s, 1H), 8.57 (d, J= 2.4 Hz, 1H); ¹⁹F NMR (282 MHz, CDCl₃): δ - 167.46; LCMS: ret. time: 13.71 min.; purity: 93.18%; MS (m/e): 407.10 (MH⁺).

10 **7.3.963 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[1-oxo-1,2,3,6-tetrahydropyrimido[2,1-c][1,4]benzoxazin-9-yl]-2,4-pyrimidinediamine (R945237)**

In a manner analogous to the preparation of N4-[2,4-dihydro-1-oxo-4H-imidazo[2,1-c][1,4]benzoxazin-8-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-[2H-1,4-benzoxazin-3(4H)-thione-6-yl]-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (400 mg, 1.04 mmol) and β-alanine (500 mg) gave 5-fluoro-N2-(3-hydroxyphenyl)-N4-[1-oxo-1,2,3,6-tetrahydropyrimido[2,1-c][1,4]benzoxazin-9-yl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (acetone-d₆): δ 2.68 (t, J= 7.2 Hz, 2H), 3.71 (t, J= 7.2 Hz, 2H), 4.62 (t, J= 1.2 Hz, 2H), 6.42 (ddd, J= 1.2 and 2.4 and 7.5 Hz, 1H), 6.98-7.08 (m, 3H), 7.38 (t, J= 2.4 Hz, 1H), 7.62 (dd, J= 2.4 and 8.7 Hz, 1H), 7.96 (d, J= 3.3 Hz, 1H), 8.12 (s, 1H), 8.16 (s, 1H), 8.52 (d, J= 2.7 Hz, 1H), 8.65 (s, 1H); ¹⁹F NMR (282 MHz, acetone-d₆): δ - 168.04.

25 **7.3.964 5-Fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-7-yl]-2,4-pyrimidinediamine (R945242)**

2H-Pyrido[3,2-b]-1,4-oxazin-3(4H)-one (500 mg) was treated with nitric acid (5 mL) and sulfuric acid (5 mL). The reaction mixture was heated to 70 °C for 30 min and then poured into ice-water. The solution was neutralized with sodium bicarbonate to pH 6. The yellow precipitation was collected by filtration, washed with water and dried to give a mixture of nitrated products (regio-isomers).

30 The mixture of nitrated compounds was reduced under hydrogenolysis conditions using 10% Pd-C in methanol at 40 psi for 30 min. The catalyst was filtered off. The filtrate was evaporated and treated with 2,4-dichloro-5-fluoropyrimidine (200 mg) in methanol (5

mL), water (5 mL). The reaction mixture was heated at 70 °C overnight, then evaporated. The residue was reacted with 3-methylaminocarbonylmethyleneoxyaniline (300 mg) in methanol (5 mL) and water (1 mL) at 100 °C overnight. The reaction mixture was diluted with 1N HCl solution (60 mL). The brown precipitation was collected by filtration, washed
5 with water and dried to give 5-fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-7-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 2.62 (d, J= 4.8 Hz, 3H), 4.33 (s, 2H), 4.63 (s, 2H), 6.48 (dd, J= 2.4 and 7.5 Hz, 1H), 7.11 (t, J= 8.1 Hz, 1H), 7.27 (d, J= 7.8 Hz, 1H), 7.36 (s, 1H), 7.86 (d, J= 2.1 Hz, 1H), 7.97 (m, 1H), 8.12 (d, J= 3.6 Hz, 1H), 8.38 (d, J= 2.1 Hz, 1H), 9.33 (s, 1H), 9.46 (s,
10 1H), 11.18 (s, 1H); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 164.49; LCMS: ret. time: 13.16 min.; purity: 79.30%; MS (m/e): 440.16 (MH⁺).

7.3.965 5-Fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-7-yl]-2,4-pyrimidinediamine (R945263)

2H-Pyrido[3,2-b]-1,4-oxazin-3(4H)-one (1g, 6.66 mmol) was refluxed with boron
hydride methyl sulfide complex (2 mL) in THF (10 mL) for 30 min to give 2H-pyrido[3,2-
b]-1,4-oxazine. In a manner analogous to the preparation of 5-fluoro-N2-(3-
methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-7-
yl]-2,4-pyrimidinediamine, 2H-pyrido[3,2-b]-1,4-oxazine was nitrated, reduced and reacted
20 with 2,4-dichloro-5-fluoropyrimidine (400 mg) and 3-
methylaminocarbonylmethyleneoxyaniline (500 mg) to give 5-fluoro-N2-(3-
methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-7-yl]-2,4-
pyrimidinediamine as a gray solid. ¹H NMR (CDCl₃): δ 2.91 (d, J= 4.8 Hz, 3H), 3.55 (t, J= 4.2 Hz, 2H), 4.24 (t, J= 4.5 Hz, 2H), 4.49 (s, 2H), 4.90 (br, 1H), 6.51 (dd, J= 2.7 and 8.1 Hz,
25 1H), 6.64 (s, 1H), 6.90 (dd, J= 2.1 and 8.1 Hz, 1H), 7.08 (s, 1H), 7.14 (br, 1H), 7.18 (t, J= 8.1 Hz, 1H), 7.28 (d, J= 2.1 Hz, 1H), 7.51 (t, J= 2.1 Hz, 1H), 7.93 (d, J= 3.0 Hz, 1H), 7.95 (d, J= 2.4 Hz, 1H); LCMS: ret. time: 11.91 min.; purity: 100%; MS (m/e): 426.12 (MH⁺).

7.3.966 5-Fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-6-yl]-2,4-pyrimidinediamine (R921304)

2H-Pyrido[3,2-b]-1,4-oxazin-3(4H)-one (2.5 g) was dissolved in acetic acid (6 mL) and acetic anhydride (30 mL). Fuming nitric acid (3 mL) was added dropwise to the reaction solution in ice-bath. The reaction solution was stirred in ice-bath overnight.

Solution was poured into crashed ice. The light yellow precipitation was collected by filtration, washed with water and dried to give a mixture of nitrated products (regio-isomers). The mixture was crystallized from dichloromethane to give 6-nitro-2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one (1 g) as a light yellow solid.

6-Nitro-2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one (1 g) was reduced under hydrogenolysis conditions using 10% Pd-C in methanol (50 mL) and 1N HCl solution (10 mL) at 50 psi for 2 h. The catalyst was filtered off and washed with methanol and 1N HCl solution. The filtrate was evaporated to give 6-amino-2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one.

In a manner analogous to the preparation of 5-fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-7-yl]-2,4-pyrimidinediamine, 6-amino-2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one was reacted with 2,4-dichloro-5-fluoropyrimidine (500 mg) and 3-methylaminocarbonylmethyleneoxyaniline (500 mg) to give 5-fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-6-yl]-2,4-pyrimidinediamine as a beige solid. ¹H NMR (DMSO-d₆): δ 2.63 (d, J= 4.5 Hz, 3H), 4.35 (s, 2H), 4.62 (s, 2H), 6.47 (dd, J= 1.8 and 8.1 Hz, 1H), 7.10 (t, J= 8.1 Hz, 1H), 7.25 (d, J= 8.1 Hz, 1H), 7.37 (m, 2H), 7.59 (d, J= 8.4 Hz, 1H), 7.96 (d, J= 5.1 Hz, 1H), 8.13 (d, J= 3.6 Hz, 1H), 9.26 (s, 1H), 9.29 (s, 1H), 11.13 (s, 1H); ¹⁹F NMR (282 MHz, DMSO-d₆): δ - 163.20; LCMS: ret. time: 25.22 min.; purity: 97.55%; MS (m/e): 440.25 (MH⁺).

7.3.967 5-Fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-6-yl]-2,4-pyrimidinediamine (R945299)

6-Nitro-2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one (500 mg) was refluxed with boron hydride methyl sulfide complex (1 mL) in THF (10 mL) for 30 min to give 6-nitro-2H-pyrido[3,2-b]-1,4-oxazine. In a manner analogous to the preparation of 5-fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-3(4H)-one-7-yl]-2,4-pyrimidinediamine, 6-nitro-2H-pyrido[3,2-b]-1,4-oxazine was reduced and reacted with 2,4-dichloro-5-fluoropyrimidine (500 mg) and 3-methylaminocarbonylmethyleneoxyaniline (500 mg) to give 5-fluoro-N2-(3-methylaminocarbonylmethyleneoxyphenyl)-N4-[2H-pyrido[3,2-b]-1,4-oxazin-6-yl]-2,4-pyrimidinediamine as a gray solid. ¹H NMR (CD₃OD): δ 2.81 (s, 3H), 3.48 (t, J= 4.5 Hz, 2H), 4.14 (t, J= 4.5 Hz, 2H), 4.44 (s, 2H), 6.60 (ddd, J= 1.5 and 2.7 and 7.5 Hz, 1H), 6.94

(d, J= 8.1 Hz, 1H), 7.14 (d, J= 3.0 Hz, 1H), 7.17 (t, J= 7.8 Hz, 1H), 7.40 (d, J= 8.9 Hz, 1H), 7.42 (t, J= 2.1 Hz, 1H), 7.92 (d, J= 3.3 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 168.20; LCMS: ret. time: 25.49 min.; purity: 97.56%; MS (m/e): 426.23 (MH⁺).

5 **7.3.968 N4-(1,4-Benzoxazin-3-on-7-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R908698):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(1,4-benzoxazin-3-on-7-yl)pyrimidineamine and 3-aminophenol were reacted to yield N4-(1,4-Benzoxazin-3-on-7-yl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. 1H (DMSO-d₆) 8.2 (d, 1H, J=4 Hz), 7.30 (m, 2H),
10 7.09 (m, 4H), 6.5 (m, 1H), 4.6 (s, 2H) purity 95 %; MS (m/e): 368 (MH⁺)

7.3.969 N2-(1,4-Benzoxazin-3-on-7-yl)- 5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R908699):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)] pyrimidineamine and 7-amino-1,4-benzoxazine-3-one were reacted to yield N2-(1,4-Benzoxazin-3-on-7-yl)- 5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. 1H (DMSO-d₆) 8.20 (d, 1H, J=4 Hz), 7.10 (m, 5H), 6.65 (m, 1H), 4.54 (s, 2H) purity 95 % MS (m/e): 368 (MH⁺)
15

7.3.970 N4-(1,4-Benzoxazine-3-on-7-yl)-5-fluoro-N2-((N-methyl acetamido-2)-3-phenoxy)-2,4-pyrimidinediamine (R908700):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-1,4-benzoxazin-3-on-7-yl)phenylpyrimidineamine and 3-(N-methylaminocarbonylmethyleneoxy)aniline were reacted to yield N4-(1,4-Benzoxazine-3-on-7-yl)-5-fluoro-N2-((N-methyl acetamido-2)-3-phenoxy)-2,4-pyrimidinediamine 1H (DMSO-d₆) 8.2 (d, 1H, J=4 Hz), 8.00 (m, 1H), 7.19 (m, 1H), 7.09 (m, 3H), 6.55 (m, 1H), 4.64 (s, 2H), 4.27 (s, 2H), 3.18 (s, 3H), 2.63 (m, 3H)
20 25 purity 95 % MS (m/e):439 (MH⁺)

7.3.971 N4-(1,4-Benzoxazine-3-on-6-yl)-5-fluoro-N2-[(N-methyl acetamido-2)-3-phenoxy)]-2,4-pyrimidinediamine (R908701):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[6-(1,4-benzoxazin-3-on-yl)]phenylpyrimidineamine and 3-(N- methylaminocarbonylmethyleneoxy)aniline were
30

reacted to yield N4-(1,4-Benzoxazine-3-on-6-yl)-5-fluoro-N2-[(N-methyl acetamido-2)-3-phenoxy]-2,4-pyrimidinediamine 1H (DMSO-d₆) 8.2 (d, 1H, J=4 Hz), 8.00 (m, 1H), 7.13 (m, 3H), 6.95 (m, 1H), 6.55 (m, 1H), 4.64 (s, 2H), 4.27 (s, 2H), 3.18 (s, 3H), 2.63 (m, 3H) purity 96 % MS (m/e): 439 (MH⁺)

5 **7.3.972 N4-(1,4-Benzoxazine-3-on-6-yl)- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R908702):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(1,4-benzoxazin-3-on-6-yl)phenylpyrimidineamine and 3-aminophenol were reacted to yield N4-(1,4-Benzoxazine-3-on-6-yl)- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine 1H (DMSO-d₆) 8.20 (d, 1H, J=4 Hz), 7.22 (m, 2H), 7.03 (m, 4H), 6.55 (m, 1H), 4.64 (s, 2H) purity 98 %MS (m/e): 368 (MH⁺)

7.3.973 5-Fluoro-N4-(3-hydroxyphenyl)- N2-(N-methyl-1,4-benzoxazine-3-on-6-yl)-2,4-pyrimidinediamine (R908703):

15 In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-6-yl)]phenylpyrimidineamine and 3-(N- methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-Fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl] -N4-(N-methyl-1,4-benzoxazin-3-on-6-yl)]pyrimidinediamine 1H (DMSO-d₆) 8.20 (d, 1H, J=4 Hz), 7.23 (m, 6H), 6.55 (m, 1H), 4.64 (s, 2H), 3.18 (s, 3H) purity 96 %; MS (m/e): 382(MH⁺)

7.3.974 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(N-methyl-1,4-benzoxazin-7-yl)-2,4-pyrimidinediamine (R908704):

25 In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)]phenylpyrimidineamine and 3-(N- methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(N-methyl-1,4-benzoxazin-7-yl)-2,4-pyrimidinediamine 1H (DMSO-d₆) 8.8.13 (d, 1H, J=4 Hz), 7.13 (m, 3H), 6.72 (m, 3H), 6.59 (m, 1H), 4.24 (m, 2H), 4.27 (s, 2H), 3.28 (m, 2H), 2.83 (m, 3H) purity 93 %; MS (m/e): 367 (MH⁺)

7.3.975 5-Fluoro-N2-[(N-methyl acetamido-2)-3-phenoxy]-N4-(N-methyl-1,4-benzoxazin-7-yl)-2,4-pyrimidinediamine (R908705):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)]phenylpyrimidineamine and 3-(N-methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-Fluoro-N2-[(N-methyl acetamido-2)-3-phenoxy]-N4-(N-methyl-1,4-benzoxazin-7-yl)-2,4-pyrimidinediamine 1H (DMSO-d6) 8.20 (d, 1H, J=4 Hz), 7.13 (m, 5H), 6.75 (m, 2H), 4.44 (s, 2H), 4.27 (m, 2H), 3.22 (m, 2H), 2.83 (s, 3H), 2.63 (m, 3H) purity 96 %; MS (m/e): 439 (MH+)

7.3.976 N2-(1,4-Benzoxazin-7-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R908706):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)]pyrimidineamine and 7-amino-1,4-benzoxazine were reacted to yield N2-(1,4-Benzoxazin-7-yl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. 1H (DMSO-d6) 7.95 (d, 1H, J=4 Hz), 7.43 (m, 1H), 7.02 (m, 4H), 6.42 (m, 2H), 4.17 (m, 2H), 3.33 (m, 2H) purity 96 %; MS (m/e): 353 (MH+)]

7.3.977 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(N-methyl-1,4-benzoxazin-7-yl)-2,4-pyrimidinediamine (R908707):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-7-yl)pyrimidineamine and 3-aminophenol were reacted to yield 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(N-methyl-1,4-benzoxazin-7-yl)-2,4-pyrimidinediamine. 1H (DMSO-d6) 8.20 (d, 1H, J=4 Hz), 7.10 (m, 5H), 6.65 (m, 1H), 4.54 (s, 2H) purity 95 % MS (m/e): 368 (MH+)

7.3.978 5-Fluoro-N4-(3-hydroxyphenyl) N2-(N-Methyl-1,4-benzoxazine-3-on-7-yl)-2,4-pyrimidinediamine (R908708):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)pyrimidineamine and 7-amino-4-N-methyl-1,4-benzoxazine-3-one were reacted to yield 5-Fluoro-N4-(3-hydroxyphenyl) N2-(N-Methyl-1,4-benzoxazine-3-on-7-yl)-2,4-pyrimidinediamine. 1H

(DMSO-d₆) 8.20 (d, 1H, J=4 Hz), 7.23 (m, 1H), 7.15 (m, 5H), 6.62 (m, 1H), 4.64 (s, 2H), 4.27 (s, 2H), 3.18 (s, 3H) purity 95 %; MS (m/e): 380 (MH⁺)

7.3.979 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(N-methyl-1,4-benzoxazin-3-on-6-yl)-2,4-pyrimidinediamine (R908709):

5 In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)] pyrimidineamine and 6-amino-1,4-benzoxazine were reacted to yield 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(N-methyl-1,4-benzoxazin-3-on-6-yl)-2,4-pyrimidinediamine. ¹H (DMSO-d₆) 8.20 (d, 1H, J=4 Hz), 7.43 (m, 2H), 7.19 (m, 4H), 6.55 (m, 1H), 4.64 (s, 2H), 3.25 (s, 3H) purity 95 %; MS (m/e): 382 (MH⁺)

7.3.980 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(N-methyl-1,4-benzoxazin-6-yl)-2,4-pyrimidinediamine (R908710):

15 In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)pyrimidineamine and 6-amino-1,4-benzoxazine were reacted to yield 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(N-methyl-1,4-benzoxazin-6-yl)-2,4-pyrimidinediamine. ¹H (MeOD-d₄) 8.20 (d, 1H, J=4 Hz), 7.43 (m, 3H), 6.90 (m, 2H), 6.75 (m, 1H), 4.25 (m, 2H), 3.25 (m, 2H), 2.85 (bs, 1 H) purity 96 %; MS (m/e): 382 (MH⁺)

7.3.981 N4-(1,4-Benzoxazin-7-yl)-5-fluoro-N2-(3-hydroxyphenyl) - pyrimidinediamine (R908711):

20 In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-[6-(1,4-benzoxazinyl)]-N2-chloro-5-fluoropyrimidineamine and 3-ethoxycarbonylmethyleneoxyaniline were reacted to yield N4-(1,4-Benzoxazin-7-yl)-5-fluoro-N2-(3-hydroxyphenyl) -pyrimidinediamine ¹H NMR (MeOD-d₄): δ 8.2 (d, 1H, J=4 Hz), 7.15 (m, 4H), 6.84 (m, 2H), 6.62 (m, 1H), 4.65 (s, 2H), 4.15 (m, 4H), 3.28 (m, 2H), 1.19 (t, 3H, J=7 Hz) purity 94 %; MS (m/e): 439 (MH⁺).

7.3.982 (+/-)-5-Fluoro-N2-[(N-methyl acetamido-2)-3-phenoxy]- N4-(2-methyl-1,4-benzoxazin-6-yl)-2,4-pyrimidinediamine (R908712):

30 In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, (+/-)-2-chloro-5-fluoro-N4-(2-methyl-1,4-benzoxazin-6-yl)]phenylpyrimidineamine and 3-(N-methylaminocarbonylmethyleneoxy)aniline were

reacted to yield (+/-)-5-Fluoro-N2-[(N-methyl acetamido-2)-3-phenoxy]- N4-(2-methyl-1,4-benzoxazin-6-yl)- 2,4-pyrimidinediamine 1H (DMSO-d6) 8.20 (d, 1H, J=4 Hz), 8.13 (m, 1H), 7.1 (m, 5H), 6.96 (m, 1H), 6.63 (m, 1H), 4.62 (m, 1H), 4.40 (s, 3H), 2.63 (m, 3H), 1.25 (m, 3H) purity 93 %; MS (m/e): 453 (MH⁺)

5 **7.3.983 N2-(N-Ethylcarbonylmethyleneoxy-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-hydroxyphenyl]phenyl]pyrimidinediamine (R908734):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-Chloro-5-fluoro-N4-(3-hydroxyphenyl)pyrimidineamine and 6-Amino-N-carbomethoxy-1,4-benzoxazine were reacted to yield N2-(N-Ethylcarbonylmethyleneoxy-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-hydroxyphenyl]phenyl]pyrimidinediamine 1H NMR (DMSO-d6): δ 8.23 (m, 1H), 7.20 (m, 1H), 7.14 (m, 4H), 6.95(m, 1H), 6.76 (m, 1H), 4.66 (s, 1H), 4.48 (s, 1H), 4.25 (q, 2H J=6.5 Hz), 1.28 (t, 2H, J=6.5 Hz), purity 95 % MS (m/e): 454(MH⁺).

15 **7.3.984 N4-(1,4-Benzoxazin-6-yl)-N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoropyrimidinediamine (R909255):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-[6-(1,4-benzoxazinyl)]-N2-chloro-5-fluoropyrimidineamine and 3-chloro-4-hydroxy-5-methylaniline were reacted to yield N4-(1,4-benzoxazin-6-yl)-N2-(3-chloro-4-hydroxy-5-methylphenyl)-5-fluoropyrimidinediamine ¹H NMR (DMSO-d6): 8.789 (d, 1H, J=4 Hz), 7.25 (m, 1H), 7.14 (m, 1H), 6.80 (m, 2H), 6.82 (m, 1H), 4.29 (s, 2H), 3.35 (m, 2H), 2.20 (s, 3H) purity 99 %; MS (m/e): 402 (MH⁺).

25 **7.3.985 5-Fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl] -N4-(N-methyl-1,4-benzoxazin-6-yl)pyrimidinediamine (R909259):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[6-(N-methyl-1,4-benzoxazinyl)]phenyl pyrimidineamine and 3-(N-methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-Fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-N4-(N-methyl-1,4-benzoxazin-6-yl)pyrimidinediamine 1H (DMSO-d6) 8.01 (d, 1H, J=4 Hz), 7.33 (m, 2H), 7.22 (m, 1H), 7.02 (m, 2H), 6.65 (m, 1H), 6.42 (m, 1H), 4.37 (s, 2H), 4.22 (m, 2H), 3.18 (m, 2H), 2.78 (s, 3H) 2.63 (m, 3H) purity 98 %; MS (m/e): 439 (MH⁺)

7.3.986 5-Fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-N4-[6-(N-methyl-1,4-benzoxazin-3-onyl)]pyrimidinediamine (R909260):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[6-(N-methyl-1,4-benzoxazin-3-onyl)]phenylpyrimidineamine and 3-(N-methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-Fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-N4-[6-(N-methyl-1,4-benzoxazin-3-onyl)]pyrimidinediamine ¹H (DMSO-d₆) 8.01 (d, 1H, J=4 Hz), 7.33 (m, 2H), 7.22 (m, 1H), 7.02 (m, 2H), 6.65 (m, 1H), 6.42 (m, 1H), 4.37 (s, 2H), 4.22 (s, 2H), 3.18 (s, 3H), 2.78 (s, 3H) 2.63 (m, 3H) purity 88%; MS (m/e): 453 (MH⁺)

7.3.987 5-Fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)-2,4-pyrimidinediamine (R909261):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)phenylpyrimidineamine and 3-(N-methylaminocarbonylmethyleneoxy)aniline were reacted to yield 5-Fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)pyrimidinediamine ¹H (DMSO-d₆) 8.08 (d, 1H, J=4 Hz), 7.43 (m, 2H), 7.19 (m, 1H), 7.09 (m, 3H), 6.55 (m, 1H), 4.64 (s, 2H), 4.27 (s, 2H), 3.18 (s, 3H), 2.63 (m, 3H) MS (m/e): 453 (MH⁺)

7.3.988 (+/-)-5-Fluoro-N4-(3-hydroxyphenyl)-N2-(2-methyl-1,4-benzothiazin-3-on-6-yl)pyrimidinediamine (R909263):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(3-hydroxyphenyl)pyrimidineamine and 6-amino-2-methyl-1,4-benzothiazin-3-one were reacted to yield (+/-)-5-Fluoro-N4-(3-hydroxyphenyl)-N2-(2-methyl-1,4-benzothiazin-3-on-6-yl)pyrimidinediamine ¹H NMR (MeOD-d₄): 8.02 (d, 1H, J=4 Hz), 7.30 (m, 3H), 7.08 (m, 3H), 6.52 (m, 1H), 3.57 (m, 1H), 1.25 (m, 3H) purity 92 %; MS (m/e): 398 (MH⁺).

7.3.989 5-Fluoro-N2-[3-hydroxyphenyl]-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)-2,4-pyrimidinediamine (R909264):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)phenylpyrimidineamine and 3-aminophenol were reacted to yield 5-Fluoro-N2-[3-

hydroxyphenyl)-N4-(N-methyl-1,4-benzoxazin-3-on-7-yl)]-2,4-pyrimidinediamine 1H (DMSO-d6) 8.08 (d, 1H, J=4 Hz), 7.53 (m, 2H), 7.09 (m, 4H), 6.42 (m, 1H), 4.64 (s, 2H), 3.27 (s, 3H) purity 95 % MS (m/e): 382 (MH⁺)

5 **7.3.990 N4-(3-Ethylcarboxy-4*H*-imidazo[5,1-*c*]-1,4-benzoxazin-6-yl)-5-fluoro -N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]pyrimidinediamine (R909265):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N2-Chloro-N4-(3-ethylcarboxy-4*H*-imidazo[5,1-*c*]-1,4-benzoxazin-6-yl)-5-fluoropyrimidineamine and 3-(N-methylaminocarbonylmethyleneoxy)aniline were reacted to yield N4-(3-Ethylcarboxy-4*H*-imidazo[5,1-*c*]-1,4-benzoxazin-6-yl)-5-fluoro -N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]pyrimidinediamine ¹H NMR (DMSO-d6): δ 8.23 (m, 2H), 8.08 (d, J=4 Hz, 1H), 7.92 (m, 1H), 7.43 (m, 1H), 7.38(m, 2H), 7.18 (m, 1H), 6.99 (t, 1H), 6.41 (m, 1H), 5.43 (s, 2H) purity 92 %; MS (m/e): 534 (MH⁺).

15 **7.3.991 N4-(1,4-Benzoxazin-7-yl)-N2-[3-ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R909266):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, N4-(1,4-Benzoxazin-7-yl)-2-chloro-5-fluoro- pyrimidineamine and 3-(ethoxycarbonylmethyleneoxy)aniline were reacted to yield N4-(1,4-Benzoxazin-7-yl)-N2-[3-ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine. 1H (DMSO-d6) 8.2 (d, 1H, J=4 Hz), 7.43 (m, 1H), 7.12 (m, 4H), 6.68 (m, 2H), 4.7 (s, 2H) 4.17 (m, 2H), 3.33 (m, 2H), 3.13 (m, 2H) 1.87 (m, 3H) purity 89 %; MS (m/e): 439 (MH⁺)

25 **7.3.992 N2-(3-Ethylcarboxy-4*H*-imidazo[5,1-*c*]-1,4-benzoxazin-6-yl)-5-fluoro-N4-(3-hydroxyphenyl)pyrimidinediamine (R909267):**

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-Chloro-5-fluoro-N4-(3-hydroxyphenyl)pyrimidineamine and 3 Ethyl 6-Amino-(3-carboxy-4*H*-imidazo[5,1-*c*]-1,4-benzoxazine were reacted to yield N2-(3-Ethylcarboxy-4*H*-imidazo[5,1-*c*]-1,4-benzoxazin-6-yl)-5-fluoro-N4-(3-hydroxyphenyl)pyrimidinediamine 1H NMR (DMSO-d6): δ 8.18 (m, 1H), 8.04 (m, 1H), 7.38 (m, 1H), 7.22 (m, 1H), 7.04 (m, 2H), 6.96 (m, 1H), 6.53 (m, 1H), 5.42 (s, 2H), 4.25 (q, 2H J=6.5 Hz), 1.28 (t, 2H, J=6.5 Hz), purity 92 % MS (m/e): 409 (MH⁺).

7.3.993 N2-(1,4-Benzoxazin-3-on-6-yl)- 5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R909268)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine N4-[6-(1,4-benzoxazinyl)]-N2-chloro-5-fluoropyrimidineamine and 6-amino-1,4-benzoxazin-3-one were reacted to yield N2-(1,4-benzoxazin-3-on-6-yl)-5-fluoro-N4-(6-(1,4-benzoxazinyl))-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.18 (d, 1H J= 4 Hz), 7.17 (m, 2H), 6.88 (m, 2H), 6.80 (m, 1H), 6.58 (m, 1H) 4.52 (s, 2H), 4.11 (m, 2H), 3.33 (m, 2H) purity: 97 %; MS (m/e): 409 (MH⁺).

7.3.994 N2-[3-(N,N-Dimethylaminocarbonylmethyleneoxy) phenyl]-N4-(1,4-benzoxazin-6-yl)-5-fluoro--2,4-pyrimidinediamine (R909290)

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(1,4-benzoxazin-6-yl)-N2-(3-ethoxyocarbonylmethyleneoxyphenyl)-5-fluoro-pyrimidinediamine and dimethylamine hydrochloride were reacted to yield N2-[3-(N,N-Dimethylaminocarbonylmethyleneoxy)phenyl] -N4-(1,4-benzoxazin-6-yl) -5-fluoro--2,4-pyrimidinediamine ¹H NMR (CD₃OD): δ 7.8 (d, 1H), 7.4 (m, 1H), 7.05 (m, 2H), 7.0 (s, 1H), 6.8 (dd, 1H), 6.66 (d, 1H), 6.56 (dd, 1H), 4.35 (s, 2H), 4.18 (m, 2H), 3.25 (m, 2H), 2.8 (s, 6H); purity: 95 %; MS (m/e): 439 (MH⁺)

7.3.995 N4-(4N-Carboxamidino-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl] -2,4-pyrimidinediamine (R909292)

To a solution in 2 mL THF at 0° Celsius containing 250 mg (0.59 mmol) of N4-(1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 1.4 eq, 115 uL TEA, and catalytic DMAP was added 0.4 eq, 70 mg of triphosgene. After 30 min 15 mL of aqueous ammonia and stirred for 30 min at RT. The THF was evaporated and the reaction was diluted with water, and the resulting precipitate collection by suction filtration. The crude precipitate was purified by preparative TLC (5% MeOH/EtOAc) to yield N4-(4N-Carboxamidino-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.83 (m, 1H), 7.42 (m, 1H), 7.12 (m, 2H), 7.08 (s, 1H), 6.84 (m, 1H), 6.66 (m, 1H), 6.48 (m, 1H), 4.30 (s, 2H), 4.15 (m, 2H), 3.22 (m, 2H), 2.82 (s, 3H); purity: 87 %; MS (m/e): 468 (MH⁺).

7.3.996 N4-(3,3-Dimethyl-1,4-benzoxazin-6-yl)-N2-[3-ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-pyrimidinediamine (R909308):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-
 5 2,4-pyrimidinediamine, N2-Chloro-N4-(3,3-dimethyl-1,4-benzoxazin-6-yl)-5-fluoro-
 pyrimidineamine and 3-(ethoxycarbonylmethyleneoxy)aniline were reacted to yield N4-
 (3,3-Dimethyl-1,4-benzoxazin-6-yl)-N2-[3-ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-
 2,4-pyrimidinediamine. ¹H (DMSO-d₆) 8.00 (m, 1H), 7.43 (m, 2H), 7.05 (m, 1H), 6.82 (m,
 2H), 6.68 (m, 1H), 6.41 (m, 1H), 4.80 (s, 2H), 4.18 (q, 2H), 3.74 (s, 2 H), 1.03 (t, 3H),
 10 1.00 (s, 6H) purity 99 %; MS (m/e): 467 (MH⁺)

7.3.997 N4-(3,3-Dimethyl-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R909309):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-
 15 methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,3-Dimethyl-1,4-
 benzoxazin-6-yl)-N2-[3-ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-
 pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(3,3-
 Dimethyl-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylaminocarbonyl
 methyleneoxy)phenyl]-2,4-pyrimidinediamine. ¹H (DMSO-d₆) 8.04 (d, 1H), 7.93 (m, 1H),
 20 7.45 (m, 2H), 7.09 (m, 1H), 6.93 (m, 2H), 6.62 (m, 1H), 6.43 (m, 1H), 4.37 (s, 2H), 3.74
 (s, 2 H), 2.62 (d, 3H), 1.07 (s, 6H) purity 99 %; MS (m/e): 453 (MH⁺)

7.3.998 N4-(3,3-Dimethyl-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R909309):

In like manner to N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-
 25 methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3,3-Dimethyl-1,4-
 benzoxazin-6-yl)-N2-[3-ethoxycarbonylmethyleneoxy)phenyl]-5-fluoro-2,4-
 pyrimidinediamine and methylamine hydrochloride were reacted to yield N4-(3,3-
 Dimethyl-1,4-benzoxazin-6-yl)-5-fluoro-N2-[3-(N-methylaminocarbonyl
 30 methyleneoxy)phenyl]-2,4-pyrimidinediamine. ¹H (DMSO-d₆) 8.04 (d, 1H), 7.93 (m, 1H),
 7.45 (m, 2H), 7.09 (m, 1H), 6.93 (m, 2H), 6.62 (m, 1H), 6.43 (m, 1H), 4.37 (s, 2H), 3.74
 (s, 2 H), 2.62 (d, 3H), 1.07 (s, 6H) purity 99 %; MS (m/e): 453 (MH⁺)

7.3.999 N4-(2,4-Diiodo-3-hydroxyphenyl)-5-fluoro-N2-(3-iodo-1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine (R935221)

To 5-fluoro-N2-(3-hydroxyphenyl)-N4-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine (34.4 mg, 0.098 mmole) in ethanol (2.0 mL) and aq. NH₄OH (2.0 mL),
5 I₂ (0.126 g, 0.99 mmole atom) was added and stirred at room temperature overnight. Reaction mixture was concentrated, dissolved in EtOAc and treated with aq. hypo solution. Organic layer was separated, dried with anhydrous Na₂SO₄ and concentrated. The crude material was purified by silica gel column chromatography to provide N4-(2,4-diiodo-3-hydroxyphenyl)-5-fluoro-N2-[3-iodo-1-methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H
10 NMR (DMSO-d₆): δ 9.86 (s, 1H), 9.51 (s, 1H), 9.12 (s, 1H), 8.28 (s, 1H), 8.07 (d, 1H, J = 3.5 Hz), 7.79 (s, 1H), 7.63 (s, 1H), 7.32 (d, 1H, J = 8.8 Hz), 7.37 (d, 1H, J = 8.8 Hz), 3.92 (s, 3H). LCMS: ret. time: 20.88 min.; purity: 91%; MS (*m/e*): 729 (MH⁺).

7.3.1000 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935222)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine was reacted with 5-amino-1-(methoxycarbonyl)methylindazoline to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.17 (s, 1H), 9.13 (s, 1H), 8.10 (s, 1H),
20 8.03 (d, 1H, J = 4.1 Hz), 7.85 (s, 1H), 7.58 (d, 2H, J = 8.8 Hz), 7.46 (s, 2H), 6.87 (s, 2H, J = 8.8 Hz), 5.31 (s, 2H), 4.57 (sep, 1H, J = 5.8 Hz), 3.65 (s, 3H), 1.25 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 21.33 min.; purity: 96%; MS (*m/e*): 451 (MH⁺).

7.3.1001 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935223)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 5-amino-1-(methoxycarbonyl)methyl-indazoline to provide N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-
30 indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.16 (s, 1H), 9.14 (s, 1H), 8.13 (s, 1H), 8.04 (d, 1H, J = 4.1 Hz), 7.89 (s, 1H), 7.48 (s, 2H), 7.30 (d, 1H, J = 2.9 Hz),

7.20 (dd, 1H, J = 2.9 and 8.8 Hz), 6.79 (d, 1H, J = 8.8 Hz), 5.32 (s, 2H), 4.22 (s, 4H), 3.65 (s, 3H). LCMS: ret. time: 21.33 min.; purity: 96%; MS (*m/e*): 451 (MH⁺).

7.3.1002 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-[1-(*N*-methylaminocarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine (R935224)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-(4-isopropoxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N2-(4-isopropoxyphenyl)-N4-[1-(*N*-methylaminocarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.46 (s, 1H), 8.98 (s, 1H), 8.07 (d, 1H, J = 4.1 Hz), 8.02 (d, 1H, J = 4.7 Hz), 7.98 (s, 2H), 7.66 (d, 1H, J = 8.8 Hz), 7.50 (d, 2H, J = 8.8 Hz), 7.46 (app s, 1H), 6.74 (d, 2H, J = 8.8 Hz), 4.96 (s, 2H), 4.46 (sept, 1H, J = 5.8 Hz), 2.58 (d, 3H, J = 4.7 Hz), 1.21 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 18.22 min.; purity: 93%; MS (*m/e*): 450 (MH⁺).

7.3.1003 N2-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N4-[1-(*N*-methylaminocarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine (R935225)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[1-(*N*-methylaminocarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.47 (s, 1H), 8.99 (s, 1H), 8.08 (d, 1H, J = 3.5 Hz), 8.06 (s, 1H), 8.01 (d, 1H, J = 4.7 Hz), 7.98 (d, 1H, J = 1.1 Hz), 7.66 (d, 1H, J = 8.8 Hz), 7.45 (dd, 1H, J = 1.1 and 8.8 Hz), 7.31 (d, 1H, J = 2.3 Hz), 7.01 (dd, 1H, J = 2.9 and 8.8 Hz), 6.66 (d, 1H, J = 8.8 Hz), 4.95 (s, 2H), 4.14 (s, 4H), 2.57 (d, 3H, J = 4.1 Hz). LCMS: ret. time: 15.55 min.; purity: 94%; MS (*m/e*): 450 (MH⁺).

7.3.1004 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935237)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-

pyrimidineamine was reacted with 5-amino-1-(methoxycarbonyl)methyl-indazoline to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.40 (s, 1H), 9.19 (s, 1H), 9.17 (s, 1H), 8.23 (s, 1H), 8.07 (d, 1H, J = 3.5 Hz), 7.90 (s, 1H), 7.47 (s, 2H), 7.25 (d, 1H, J = 7.6 Hz), 7.11 (d, 1H, J = 7.6 Hz), 7.08 (d, 1H, J = 8.2 Hz), 6.53 (d, 1H, J = 8.2 Hz), 5.31 (s, 2H), 3.64 (s, 3H). LCMS: ret. time: 15.82 min.; purity: 96%; MS (*m/e*): 409 (MH⁺).

7.3.1005 N2, N4-Bis[1-(2-hydroxyethyl)indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine (R935238)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2, N4-bis[1-(methoxycarbonyl)methyl-indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine was reacted with DIBAL-H to produce N2, N4-bis[1-(2-hydroxyethyl)indazoline-6-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.56 (s, 1H), 9.43 (s, 1H), 8.19 (d, 1H, J = 3.5 Hz), 8.06 (s, 1H), 8.03 (s, 1H), 7.98 (s, 1H), 7.86 (s, 1H), 7.66 (d, 1H, J = 8.8 Hz), 7.52 (d, 1H, J = 8.8 Hz), 7.42 (dd, 1H, J = 1.7 and 8.8 Hz), 7.23 (dd, 1H, J = 1.7 and 8.8 Hz), 4.75 (t, 1H, J = 5.3 Hz), 4.68 (t, 1H, J = 5.3 Hz), 4.09-4.02 (m, 2H), 3.81-3.74 (m, 2H), 3.63-3.60 (m, 2H), 3.56-3.52 (m, 2H). LCMS: ret. time: 13.73 min.; purity: 90%; MS (*m/e*): 449 (MH⁺).

7.3.1006 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[1-(*N*-methylaminocarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935239)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[1-(*N*-methylaminocarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.27 (s, 1H), 9.21 (s, 1H), 8.07 (s, 1H), 8.04 (d, 1H, J = 4.1 Hz), 7.90 (qt, 1H, J = 4.7 Hz), 7.83 (s, 1H), 7.58 (d, 2H, J = 8.8 Hz), 7.44 (s, 2H), 6.87 (d, 2H, J = 8.8 Hz), 4.98 (s, 2H), 4.57 (q, 1H, J = 5.8 Hz), 2.59 (d, 3H, J = 4.1 Hz), 1.21 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 17.74 min.; purity: 94%; MS (*m/e*): 450 (MH⁺).

7.3.1007 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-(*N*-methylaminocarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935240)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(*N*-methylaminocarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.36 (br s, 2H), 8.06 (d, 1H, J = 3.5 Hz), 8.05 (s, 1H), 7.99 (qt, 1H, J = 4.7Hz), 7.87 (s, 1H), 7.46 (s, 2H), 7.30-7.28 (m, 1H), 7.20-7.17 (m, 1H), 6.79 (d, 1H, J = 8.8 Hz), 4.99 (s, 2H), 4.22 (s, 4H), 2.59 (d, 3H, J = 4.7 Hz). LCMS: ret. time: 15.06 min.; purity: 91%; MS (*m/e*): 450 (MH⁺).

7.3.1008 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935242)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-4-pyrimidineamine was reacted with 4-isopropoxyaniline to provide 5-fluoro-N2-(4-isopropoxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.58 (s, 1H), 10.09 (s, 1H), 8.23 (d, 1H, J = 5.3 Hz), 8.04 (s, 1H), 8.02 (s, 1H, J = 5.8 Hz), 7.68–7.63 (m 1H), 7.58-7.55 (s, 1H), 7.30 (d, 2H, J = 8.8 Hz), 6.82 (d, 2H, J = 8.8 Hz), 5.41 (s, 2H), 4.53 (sept, 1H, J = 5.8 Hz), 3.66 (s, 3H), 1.21 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 19.30 min.; purity: 93%; MS (*m/e*): 451 (MH⁺).

7.3.1009 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-(2-hydroxyethyl)indazoline-6-yl]-2,4-pyrimidinediamine (R935248)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(methoxycarbonyl)methyl-indazoline-6-yl]-2,4-pyrimidinediamine was reacted with DIBAL-H to produce N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(2-hydroxyethyl)indazoline-6-yl]-2,4-pyrimidinediamine. LCMS: ret. time: 19.42 min.; purity: 94%; MS (*m/e*): 423 (MH⁺).

7.3.1010 N2-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935249)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-4-pyridinamine was reacted with 3, 4-ethylenedioxyaniline to provide N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.32 (s, 1H), 8.94 (s, 1H), 8.14 (d, 1H, J = 4.7 Hz), 8.03 (d, 1H, J = 4.7 Hz), 8.01 (s, 1H), 7.65-7.57 (m, 2H), 7.23 (d, 1H, J = 1.7 Hz), 7.02 (dd, 1H, J = 1.9 and 8.8 Hz), 6.63 (d, 1H, J = 8.8 Hz), 5.38 (s, 2H), 4.14 (s, 4H), 3.66 (s, 3H). LCMS: ret. time: 18.94 min.; purity: 91%; MS (*m/e*): 451 (MH⁺).

7.3.1011 5-Fluoro-N2-(3-hydroxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine (R935250)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-4-pyrimidineamine was reacted with 3-aminophenol to provide 5-fluoro-N2-(3-hydroxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.34 (s, 1H), 9.16 (s, 1H), 8.25 (d, 1H, J = 4.7 Hz), 8.05 (d, 1H, J = 4.7 Hz), 8.02 (s, 1H), 7.65-7.57 (m, 2H), 7.10 (d, 2H, J = 5.8 Hz), 6.93 (d, 1H, J = 8.8 Hz), 6.90 (d, 1H, J = 8.8 Hz), 6.28 (app d, 1H, J = 8.8 Hz), 5.37 (s, 2H), 3.66 (s, 3H). LCMS: ret. time: 17.87 min.; purity: 97%; MS (*m/e*): 409 (MH⁺).

7.3.1012 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine (R935251)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine was reacted with 1-aminopyrrole to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.93 (s, 1H), 9.21 (s, 1H), 7.97 (d, 1H, J = 4.1 Hz), 7.47 (d, 2H, J = 8.8 Hz), 6.70 (dd, 2H, J = 2.3 and 4.7 Hz), 6.67 (d, 2H, J = 8.8 Hz), 6.02 (dd, 2H, J = 2.3 and 4.7 Hz), 4.48 (sept, 1H, J = 5.8 Hz), 1.21 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 23.44 min.; purity: 90%; MS (*m/e*): 328 (MH⁺).

7.3.1013 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine (R935252)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 1-aminopyrrole to provide N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.95 (s, 1H), 9.16 (s, 1H), 7.95 (d, 1H, J = 3.5 Hz), 7.16-7.12 (m, 2H), 6.69 (dd, 2H, J = 2.3 and 4.7 Hz), 6.61 (d, 1H, J = 8.8 Hz), 5.99 (dd, 2H, J = 2.3 and 4.7 Hz), 4.12-4.15 (m, 4H). LCMS: ret. time: 19.86 min.; purity: 92%; MS (*m/e*): 328 (MH⁺).

7.3.1014 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine (R935253)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 1-aminopyrrole to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.95 (s, 1H), 9.22 (s, 1H), 9.19 (s, 1H), 7.99 (d, 1H, J = 3.5 Hz), 7.22 (d, 1H, J = 8.2 Hz), 6.94 (br s, 1H), 6.89 (t, 1H, J = 8.2 Hz), 6.70 (dd, 2H, J = 2.3 and 4.7 Hz), 6.38 (d, 1H, J = 8.2 Hz), 5.99 (t, 2H, J = 2.3 and 4.7 Hz). LCMS: ret. time: 18.23 min.; purity: 94%; MS (*m/e*): 286 (MH⁺).

7.3.1015 5-Fluoro-N2-[1-(2-hydroxyethyl)indazoline-5-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R935255)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N2-[1-(2-hydroxyethyl)indazoline-5-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.16 (s, 1H), 9.10 (s, 1H), 8.09 (s, 1H), 8.02 (s, 1H, J = 4.0 Hz), 7.79 (s, 1H), 7.59 (d, 2H, J = 8.8 Hz), 7.48 (d, 1H, J = 8.8 Hz), 7.42 (dd, 1H, J = 1.7 and 8.8 Hz), 6.87 (d, 2H, J = 8.8 Hz), 4.83 (t, 1H, J = 5.8 Hz), 4.57 (sept, 1H, J = 5.8 Hz), 4.35 (t, 2H, J = 5.8 Hz), 3.75 (app qt, 2H, J = 5.8 Hz), 1.26 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 20.90 min.; purity: 94%; MS (*m/e*): 423 (MH⁺).

7.3.1016 5-Fluoro-N2-[1-(2-hydroxyethyl)indazoline-5-yl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R935256)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N2-[1-(2-hydroxyethyl)indazoline-5-yl]-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.39 (s, 1H), 9.18 (s, 1H), 9.14 (s, 1H), 8.19 (s, 1H), 8.07 (d, 1H, J = 4.1 Hz), 7.84 (s, 1H), 7.50-7.42 (m, 2H), 7.26 (d, 1H, J = 8.2 Hz), 7.12-7.06 (m, 2H), 6.52 (d, 1H, J = 8.2 Hz), 4.83 (t, 1H, J = 5.8 Hz), 4.35 (t, 2H, J = 5.8 Hz), 3.75 (app qt, 2H, J = 5.8 Hz). LCMS: ret. time: 15.97 min.; purity: 95%; MS (*m/e*): 381 (MH⁺).

7.3.1017 N2-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N4-[1-(2-hydroxyethyl)indazoline-5-yl]-2,4-pyrimidinediamine (R935258)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine was reacted with DIBAL-H to produce N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[1-(2-hydroxyethyl)indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.20 (s, 1H), 8.93 (s, 1H), 8.12 (s, 1H), 8.02 (d, 1H, J = 3.5 Hz), 7.94 (s, 1H), 7.59 (s, 2H), 7.23 (d, 1H, J = 0.9 Hz), 7.02 (dd, 1H, J = 1.0 and 8.8 Hz), 6.64 (d, 1H, J = 8.8 Hz), 4.86 (t, 1H, J = 5.3 Hz), 4.40 (t, 2H, J = 5.8 Hz), 4.15 (s, 4H), 3.78 (app qt, 2H, J = 5.3 and 5.8 Hz). LCMS: ret. time: 18.07 min.; purity: 93%; MS (*m/e*): 423 (MH⁺).

7.3.1018 5-Fluoro-N4-[1-(2-hydroxyethyl)indazoline-5-yl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R935259)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-(3-hydroxyphenyl)-N4-[1-(methoxycarbonyl)methyl-indazoline-5-yl]-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N4-[1-(2-hydroxyethyl)indazoline-5-yl]-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.31 (s, 1H), 9.16 (s, 1H), 9.01 (s, 1H), 8.23 (s, 1H), 8.05 (d, 1H, J = 3.5 Hz), 7.96 (s, 1H), 7.60 (s, 2H), 7.10 (app s, 2H), 6.92 (t, 1H, J = 8.8 Hz), 6.31 (d, 1H, J = 8.8 Hz), 4.86 (t, 1H, J = 5.3 Hz), 4.40

(t, 2H, J = 5.8 Hz), 3.79 (app qt, 2H, J = 5.3 and 5.8 Hz). LCMS: ret. time: 16.09 min.; purity: 89%; MS (*m/e*): 381 (MH⁺).

7.3.1019 N2-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N4-(indazoline-6-yl)-2,4-pyrimidinediamine (R935261)

5 In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-6-yl)-4-pyrimidineamine was reacted 3, 4-ethylenedioxyaniline to produce N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.85 (s, 1H), 9.40 (s, 1H), 9.01 (s, 1H), 8.07 (d, 1H, J = 3.5 Hz), 7.97 (s, 1H), 7.86 (s, 1H), 7.65 (d, 1H, J = 8.8 Hz), 7.47 (dd, 1H, J = 2.3 and 8.8 Hz), 7.27 (d, 1H, J = 2.3 Hz), 7.07 (dd, 1H, J = 2.3 and 8.8 Hz), 6.64 (dd, 1H, J = 1.7 and 8.8 Hz), 4.14 (s, 4H). LCMS: ret. time: 15.90 min.; purity: 100%; MS (*m/e*): 379 (MH⁺).

7.3.1020 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(indazoline-6-yl)-2,4-pyrimidinediamine (R935262)

15 In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-6-yl)-4-pyrimidineamine was reacted with 3-aminophenol to produce 5-fluoro-N2-(3-hydroxyphenyl)-N4-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.80 (s, 1H), 10.49 (s, 1H), 8.35 (d, 1H, J = 5.3 Hz), 8.06 (s, 1H), 7.78 (d, 1H), 7.75 (d, 1H, J = 8.8 Hz), 7.42 (dd, 1H, J = 1.7 and 8.8 Hz), 6.99-6.97 (m, 2H), 6.80 (s, 1H), 6.52-6.48 (m, 1H). LCMS: ret. time: 13.78 min.; purity: 100%; MS (*m/e*): 379 (MH⁺).

7.3.1021 N2-(3-Chloro-4-hydroxy-3-methylphenyl)-5-fluoro-N4-[4H-imidazo[2,1-c][1,4]-benzoxazin-8-yl]-2,4-pyrimidinediamine (R935263)

25 In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[4H-imidazo[2,1-c][1,4]-benzoxazin-8-yl]-4-pyrimidineamine was reacted with 4-amino-2-chloro-6-methylphenol to produce N2-(3-chloro-4-hydroxy-3-methylphenyl)-5-fluoro-N4-[4H-imidazo[2,1-c][1,4]-benzoxazin-8-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.40 (s, 1H), 9.04 (s, 1H), 8.51 (s, 1H), 8.07 (d, 1H, J = 3.5 Hz), 7.98 (d, 1H, J = 2.3 Hz), 7.67 (s, 1H), 7.53 (s, 1H), 7.41-7.36 (m, 1H), 7.20 (s, 1H), 7.10 (d, 1H, J = 8.8 Hz), 7.07 (s, 1H), 5.24 (s, 2H), 1.98 (s, 3H). LCMS: ret. time: 13.36 min.; purity: 90%; MS (*m/e*): 439 (MH⁺).

7.3.1022 N2-(3-Chloro-4-hydroxy-3-methylphenyl)-5-fluoro-N4-(indazoline-6-yl)-2,4-pyrimidinediamine (R935264)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-6-yl)-4-pyrimidineamine was reacted with 4-amino-2-chloro-6-methylphenol to produce N2-(3-chloro-4-hydroxy-3-methylphenyl)-5-fluoro-N4-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.62 (s, 1H), 9.19 (s, 1H), 8.61 (s, 1H), 8.12 (s, 1H), 8.02 (s, 1H), 7.77 (s, 1H), 7.67 (d, 1H, J = 8.8 Hz), 7.50-7.45 (m, 2H), 7.26 (s, 1H), 1.98 (s, 3H). LCMS: ret. time: 13.78 min.; purity: 100%; MS (*m/e*): 385 (MH⁺).

7.3.1023 5-Fluoro-N4-(indazoline-5-yl)-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R935266)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-5-yl)-4-pyrimidineamine was reacted with 4-isopropoxyaniline to produce 5-fluoro-N4-(indazoline-5-yl)-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.30 (s, 1H), 9.80 (s, 1H), 8.16 (d, 1H, J = 5.3 Hz), 8.06 (s, 1H), 7.98 (s, 1H), 7.51 (s, 2H), 7.32 (d, 2H, J = 8.8 Hz), 6.79 (d, 2H, J = 8.8 Hz), 4.51 (sept, 1H, J = 5.8 Hz), 1.22 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 17.65 min.; purity: 98%; MS (*m/e*): 379 (MH⁺).

7.3.1024 N2-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N4-(indazoline-5-yl)-2,4-pyrimidinediamine (R935267)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-5-yl)-4-pyrimidineamine was reacted with 3, 4-ethylenedioxyphenylaniline to produce N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.20 (s, 1H), 9.61 (s, 1H), 8.13 (d, 1H, J = 5.3 Hz), 8.08 (s, 1H), 7.98 (s, 1H), 7.54-7.48 (m, 2H), 7.06 (d, 1H, J = 2.3 Hz), 6.90 (dd, 1H, J = 2.3 and 8.8 Hz), 6.72 (d, 1H, J = 8.8 Hz), 4.17 (s, 4H). LCMS: ret. time: 15.16 min.; purity: 100%; MS (*m/e*): 379 (MH⁺).

7.3.1025 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(indazoline-5-yl)-2,4-pyrimidinediamine (R935268)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-5-yl)-4-

pyrimidineamine was reacted with 3-aminophenol to produce 5-fluoro-N2-(3-hydroxyphenyl)-N4-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.64 (s, 1H), 10.33 (s, 1H), 8.29 (d, 1H, J = 5.3 Hz), 8.12 (s, 1H), 8.03 (s, 1H), 7.55 (dd, 2H, J = 1.7 and 8.8 Hz), 7.00 (d, 1H, J = 8.8 Hz), 6.90 (d, 1H, J = 8.8 Hz), 6.85 (d, 1H, J = 1.7 Hz), 6.53 (d, 1H, J = 8.8 Hz). LCMS: ret. time: 12.80 min.; purity: 98%; MS (*m/e*): 337 (MH⁺).

7.3.1026 5-Fluoro-N4-(indazoline-5-yl)-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R935269)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-5-yl)-4-pyrimidineamine was reacted with 3-(methoxycarbonylmethyleneoxy)aniline to produce 5-fluoro-N4-(indazoline-5-yl)-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.64 (s, 1H), 9.82 (s, 1H), 8.20 (d, 1H, J = 4.6 Hz), 8.10 (s, 1H), 8.08 (s, 1H), 7.99 (s, 1H), 7.57 (m, 2H), 7.13-7.6 (m, 3H), 6.56 (d, 1H, J = 8.8 Hz), 4.60 (s, 2H), 3.65 (s, 3H). LCMS: ret. time: 15.36 min.; purity: 94%; MS (*m/e*): 409 (MH⁺).

7.3.1027 5-Fluoro-N4-(indazoline-5-yl)-N2-(indazoline-6-yl)-2,4-pyrimidinediamine (R935270)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(indazoline-5-yl)-4-pyrimidineamine was reacted with 6-aminoindazoline to produce 5-fluoro-N4-(indazoline-5-yl)-N2-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.35 (s, 1H), 9.19 (s, 1H), 8.25 (d, 1H, J = 4.1 Hz), 8.12 (s, 1H), 8.01 (s, 1H), 7.81 (s, 1H), 7.73 (s, 1H), 7.64 (d, 1H, J = 8.8 Hz), 7.60 (dd, 2H, J = 1.7 and 8.9 Hz), 7.51 (d, 1H, J = 8.9 Hz), 7.21 (dd, 1H, J = 1.7 and 8.8 Hz). LCMS: ret. time: 13.45 min.; purity: 95%; MS (*m/e*): 361 (MH⁺).

7.3.1028 5-Fluoro-N4-[4H-imidazo[2,1-*c*][1,4]-benzoxazin-8-yl]-N2-[3-(*N*-methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R935271)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-[4H-imidazo[2,1-*c*][1,4]-benzoxazin-8-yl]-4-pyrimidineamine was reacted with 3-(*N*-

methylaminocarbonylmethyleneoxy)aniline to produce 5-fluoro-N4-[4H-imidazo[2,1-c][1,4]-benzoxazin-8-yl]-N2-[3-(N-methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.44 (s, 1H), 9.25 (s, 1H), 8.11 (d, 1H, J = 3.5 Hz), 8.03 (d, 1H, J = 2.3 Hz), 7.90 (qt, 1H, J = 4.6 Hz), 7.69 (d, 1H, J = 1.2 Hz), 7.47–7.42 (m, 1H), 7.33 (m, 1H), 7.26 (dd, 1H, J = 1.2 and 8.2 Hz), 7.12 (s, 1H), 7.09 (d, 1H, J = 1.7 Hz), 6.97 (t, 1H, J = 8.2 Hz), 6.40 (dd, 1H, J = 2.3 and 8.2 Hz), 5.25 (s, 2H), 4.26 (s, 2H), 2.61 (d, 3H, J = 4.6 Hz). LCMS: ret. time: 15.45 min.; purity: 97%; MS (*m/e*): 462 (MH⁺).

7.3.1029 5-Fluoro-N2-(4-isopropoxyphenyl)-N4-(1H-pyrrol-1-yl)- 2,4-pyrimidinediamine (R935276)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1H-pyrrol-1-yl)-4-pyrimidineamine was reacted with 4-isopropoxyaniline to produce 5-fluoro-N2-(4-isopropoxyphenyl)-N4-(1H-pyrrol-1-yl)- 2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.69 (s, 1H), 9.03 (s, 1H), 8.06 (d, 1H, J = 3.5 Hz), 7.30 (d, 2H, J = 9.3 Hz), 6.82 (t, 2H, J = 2.3 Hz), 6.58 (d, 2H, J = 9.3 Hz), 6.11 (t, 2H, J = 2.3 Hz), 4.41 (sept, 1H, J = 5.8 Hz), 1.18 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 21.21 min.; purity: 90%; MS (*m/e*): 328 (MH⁺).

7.3.1030 N2-(3, 4-Ethylenedioxyphenyl)-5-Fluoro-N4-(1H-pyrrol-1-yl)- 2,4-pyrimidinediamine (R935277)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1H-pyrrol-1-yl)-4-pyrimidineamine was reacted with 3, 4-ethylenedioxyaniline to produce N2-(3, 4-ethylenedioxyphenyl)-5-Fluoro-N4-(1H-pyrrol-1-yl)- 2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 11.63 (s, 1H), 9.94 (s, 1H), 8.23 (d, 1H, J = 4.7 Hz), 6.86 (m, 4H), 6.58 (d, 1H, J = 8.8 Hz), 6.12 (t, 2H, J = 2.3 Hz), 4.15 (s, 4H). LCMS: ret. time: 17.36 min.; purity: 96%; MS (*m/e*): 328 (MH⁺).

7.3.1031 5-Fluoro-N2-(3-hydroxyphenyl)-N4-(1H-pyrrol-1-yl)- 2,4-pyrimidinediamine (R935278)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1H-pyrrol-1-yl)-4-pyrimidineamine was reacted with 3-aminophenol to produce 5-fluoro-N2-(3-hydroxyphenyl)-N4-(1H-pyrrol-1-yl)- 2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.68 (s, 1H), 9.04 (s, 1H), 9.00 (s, 1H), 8.08 (d, 1H, J = 4.11 Hz), 7.01 (d, 1H, J = 8.2

Hz), 6.84-6.75 (m, 4H), 6.22 (dd, 1H, $J = 1.2$ and 8.2 Hz), 6.08 (t, 2H, $J = 2.3$ Hz). LCMS: ret. time: 16.24 min.; purity: 94%; MS (m/e): 286 (MH^+).

7.3.1032 5-Fluoro-N4-(indazoline-5-yl)-N2-[3-(*N*-methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine (R935279)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N4-(indazoline-5-yl)-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine and $Me_2NH.HCl$ were reacted to provide 5-fluoro-N4-(indazoline-5-yl)-N2-[3-(*N*-methylaminocarbonylmethyleneoxy)phenyl]-2,4-pyrimidinediamine. ^1H NMR ($DMSO-d_6$): δ 12.98 (s, 1H), 9.35 (s, 1H), 9.16 (s, 1H), 8.21 (s, 1H), 8.07 (d, 1H, $J = 3.5$ Hz), 7.97 (s, 1H), 7.90 (qt, 1H, $J = 4.7$ Hz), 7.59 (dd, 1H, $J = 8.8$ Hz), 7.49 (d, 1H, $J = 8.8$ Hz), 7.32-7.28 (m, 2H), 7.03 (t, 1H, $J = 8.2$ Hz), 6.45 (dd, 1H, $J = 1.7$ and 8.2 Hz), 4.31 (s, 2H), 2.61 (d, 3H, $J = 4.7$ Hz). LCMS: ret. time: 12.92 min.; purity: 90%; MS (m/e): 408 (MH^+).

7.3.1033 5-Fluoro-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-N4-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine (R935280)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(1H-pyrrol-1-yl)-4-pyrimidineamine was reacted with 3-(methoxycarbonylmethyleneoxy)aniline to produce 5-fluoro-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-N4-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine. ^1H NMR ($DMSO-d_6$): δ 11.45(s, 1H), 9.90 (s, 1H), 8.26 (d, 1H, $J = 4.7$ Hz), 7.07 (d, 1H, $J = 8.2$ Hz), 7.68 (d, 1H, $J = 8.2$ Hz), 6.94 (s, 1H), 6.85 (t, 2H, $J = 2.3$ Hz), 6.47 (dd, 1H, $J = 2.3$ and 8.2 Hz), 6.12 (t, 2H, $J = 2.3$ Hz), 4.64 (s, 2H), 3.68 (s, 3H). LCMS: ret. time: 16.24 min.; purity: 92%; MS (m/e): 358 (MH^+).

7.3.1034 5-Fluoro-N2-[3-(*N*-methylaminocarbonylmethyleneoxy)phenyl]-N4-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine (R935281)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, 5-fluoro-N2-[3-(methoxycarbonylmethyleneoxy)phenyl]-N4-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine and $Me_2NH.HCl$ were reacted to provide 5-fluoro-N2-[3-(*N*-methylaminocarbonylmethyleneoxy)phenyl]-N4-(1H-pyrrol-1-yl)-2,4-pyrimidinediamine. ^1H NMR ($DMSO-d_6$): δ 10.73 (s, 1H), 9.21 (s, 1H), 8.11 (d, 1H, $J = 4.1$ Hz), 7.89 (qt, 1H, J

= 4.7 Hz), 7.14 (d, 1H, J = 8.2 Hz), 7.09 (s, 1H), 6.93 (t, 1H, J = 8.2 Hz), 6.84 (t, 2H, J = 2.3 Hz), 6.40 (dd, 1H, J = 2.3 and 8.2 Hz), 6.09 (t, 2H, J = 2.3 Hz), 4.29 (s, 2H), 2.63 (s, 3H, J = 4.7 Hz). LCMS: ret. time: 16.16 min.; purity: 90%; MS (*m/e*): 357 (MH⁺).

7.3.1035 N2-[1-(2-ethoxycarbonylethyl)indazoline-6-yl]- N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R935286)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 6-amino-1-(2-ethoxycarbonylethyl)indazoline to provide N2-[1-(2-ethoxycarbonylethyl)indazoline-6-yl]- N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.37 (s, 1H), 9.20 (s, 1H), 8.10 (d, 1H, J = 3.5 Hz), 8.06 (s, 1H), 7.87 (s, 1H), 7.53 (d, 1H, J = 8.8 Hz), 7.33-7.21 (m, 3H), 6.77 (d, 1H, J = 8.8 Hz), 4.34 (t, 2H, J = 6.4 Hz), 4.19 (s, 4H), 3.93 (qt, 2H, J = 7.0 Hz), 2.82 (t, 2H, J = 6.4 Hz), 1.04 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 24.70 min.; purity: 90%; MS (*m/e*): 479 (MH⁺).

7.3.1036 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-6-yl]-2,4-pyrimidinediamine (R935287)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonylethyl)indazoline-6-yl]- N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine was reacted with DIBAL-H to produce N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.35 (s, 1H), 9.19 (s, 1H), 8.09 (d, 1H, J = 4.1 Hz), 8.01 (s, 1H), 7.85 (s, 1H), 7.53 (d, 1H, J = 8.8 Hz), 7.32-7.20 (m, 3H), 6.77 (d, 1H, J = 8.8 Hz), 4.20 (s, 4H), 3.27 (t, 2H, J = 6.4 Hz), 3.27 (t, 2H, J = 6.4 Hz), 1.84 (q, 2H, J = 6.4 Hz). LCMS: ret. time: 24.70 min.; purity: 90%; MS (*m/e*): 479 (MH⁺). LCMS: ret. time: 22.09 min.; purity: 90%; MS (*m/e*): 437 (MH⁺).

7.3.1037 N2-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N4-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-6-yl]-2,4-pyrimidinediamine (R935288)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-[1-(2-

ethoxycarbonyl)ethyl]indazoline-6-yl]-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[1-[2-(N-methylaminocarbonyl)ethyl]-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.35 (s, 1H), 9.19 (s, 1H), 8.10 (d, 1H, J = 3.5 Hz), 8.02 (s, 1H), 7.86 (s, 1H), 7.81 (qt, 1H, J = 4.7 Hz), 7.52 (d, 1H, J = 8.2 Hz), 7.34-7.22 (m, 3H), 6.77 (d, 1H, J = 8.8 Hz), 4.33 (t, 2H, J = 6.4 Hz), 4.19 (s, 4H), 2.57 (t, 2H, J = 6.4 Hz), 2.48 (d, 3H, J = 4.7 Hz). LCMS: ret. time: 23.10 min.; purity: 93%; MS (*m/e*): 464 (MH⁺).

7.3.1038 N4-[1-(2-Ethoxycarbonyl)ethyl]indazoline-6-yl]-5-fluoro-N2-(isopropoxyphenyl)-2,4-pyrimidinediamine (R935289)

In like manner to the preparation of N4-(3,4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine was reacted with 6-amino-1-(2-ethoxycarbonyl)ethylindazoline to provide N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]-5-fluoro-N4-(isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.74 (s, 1H), 10.55 (s, 1H), 8.35 (d, 1H, J = 5.8 Hz), 7.98 (s, 1H), 7.77 (s, 1H), 7.66 (d, 1H, J = 8.8 Hz), 7.51 (d, 2H, J = 8.8 Hz), 7.16 (dd, 1H, J = 1.2 and 8.8 Hz), 6.85 (d, 2H, J = 8.8 Hz), 4.55 (sept, 1H, J = 6.4 Hz), 4.31 (t, 2H, J = 6.4 Hz), 3.93 (qt, 2H, J = 7.0 Hz), 2.80 (t, 2H, J = 6.4 Hz), 1.22 (d, 6H, J = 7.0 Hz), 1.02 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 26.84 min.; purity: 96%; MS (*m/e*): 479 (MH⁺).

7.3.1039 5-Fluoro-N2-[1-(3-hydroxypropyl)indazoline-6-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R935290)

In like manner to the preparation of N2-(3,4-ethylenedioxyphenyl)-5-fluoro-N4-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]-5-fluoro-N4-(isopropoxyphenyl)-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-6-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.31 (s, 1H), 9.22 (s, 1H), 8.08 (d, 1H, J = 4.1 Hz), 7.98 (s, 1H), 7.85 (s, 1H), 7.62 (dd, 2H, J = 3.5 and 8.8 Hz), 7.52 (d, 1H, J = 8.8 Hz), 7.27 (d, 1H, J = 8.8 Hz), 6.86 (d, 2H, J = 8.8 Hz), 4.55 (sept, 1H, J = 7.0 Hz), 4.49 (t, 1H, J = 5.3 Hz), 4.14 (t, 2H, J = 6.4 Hz), 3.26 (t, 2H, J = 6.4 Hz), 1.84 (q, 2H, J = 6.4 Hz), 1.24 (d, 6H, J = 7.0 Hz). LCMS: ret. time: 24.13 min.; purity: 97%; MS (*m/e*): 437 (MH⁺).

7.3.1040 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[1-[2(*N*-methylamino)carbonyl]ethyl-indazoline-6-yl]-2,4-pyrimidinediamine (R935291)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N4-(isopropoxyphenyl)-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[1-[2(*N*-methylamino)carbonyl]ethyl-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.32(s, 1H), 9.24 (s, 1H), 8.10 (d, 1H, J = 3.5 Hz), 7.99 (s, 1H), 7.85 (s, 1H), 7.80 (qt, 1H, J = 4.7 Hz), 7.63 (d, 2H, J = 8.8 Hz), 7.52 (d, 1H, J = 8.8 Hz), 7.28 (d, 1H, J = 8.8 Hz), 6.84 (d, 2H, J = 8.8 Hz), 4.54 (sept, 1H, J = 5.8 Hz), 4.30 (t, 2H, J = 6.4 Hz), 2.55 (t, 2H, 7.4 Hz), 2.48 (d, 3H, J = 4.7 Hz), 1.24 (d, 6H, J = 6H). LCMS: ret. time: 23.68 min.; purity: 95%; MS (*m/e*): 464 (MH⁺).

7.3.1041 N4-[1-(2-Ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R935292)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 6-amino-1-(2-ethoxycarbonyl)ethylindazoline to provide N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.35 (s, 1H), 10.21 (s, 1H), 8.29 (d, 1H, J = 5.3 Hz), 7.96 (s, 1H), 7.90 (s, 1H), 7.63 (d, 1H, J = 8.8 Hz), 7.20-7.06 (m, 4H), 6.58 (d, 1H, J = 8.2 Hz), 4.33 (t, 2H, J = 6.4 Hz), 3.94 (qt, 2H, J = 7.0 Hz), 2.82 (t, 2H, J = 6.4 Hz), 1.03 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 22.73 min.; purity: 94%; MS (*m/e*): 437 (MH⁺).

7.3.1042 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-(3-hydroxypropyl)indazoline-6-yl]-2,4-pyrimidinediamine (R935293)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-(3-hydroxypropyl)indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.38 (s, 1H), 9.35 (s, 1H), 9.26 (s, 1H), 8.13 (d, 1H, J = 4.1 Hz), 8.05 (s, 1H), 7.85 (s, 1H), 7.52 (d, 1H, J = 8.2 Hz), 7.28 (d, 2H, J = 8.8 Hz), 7.12 (d, 1H, J = 1.7 Hz), 7.08 (t, 1H, J =

8.2 Hz), 6.49 (d, 1H, J = 8.2 Hz), 4.15 (t, 2H, J = 7.0 Hz), 3.26 (t, 2H, J = 6.4 Hz), 1.85 (q, 2H, J = 6.4 Hz). LCMS: ret. time: 24.70 min.; purity: 90%; MS (*m/e*): 479 (MH⁺). LCMS: ret. time: 20.37 min.; purity: 98%; MS (*m/e*): 395 (MH⁺).

7.3.1043 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2-(*N*-methylaminocarbonyl)ethyl]-indazoline-6-yl]-2,4-pyrimidinediamine (R935294)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2-(*N*-methylaminocarbonyl)ethyl]-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.36 (s, 1H), 9.33 (s, 1H), 9.25 (s, 1H), 8.14 (d, 1H, J = 3.5 Hz), 8.05 (s, 1H), 7.85 (s, 1H), 7.78 (qt, 1H, J = 4.7 Hz), 7.52 (d, 1H, J = 8.8 Hz), 7.30 (d, 2H, J = 8.8 Hz), 7.11 (d, 1H, J = 2.3 Hz), 7.07 (t, 1H, J = 8.2 Hz), 6.47 (d, 1H, J = 8.2 Hz), 4.32 (t, 2H, J = 6.4 Hz), 2.57 (t, 2H, J = 6.4 Hz), 2.48 (d, 3H, J = 4.7 Hz). LCMS: ret. time: 20.18 min.; purity: 93%; MS (*m/e*): 422 (MH⁺).

7.3.1044 N4-[1-(2-Ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine (R935295)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(2-methoxycarbonylbenzofur-5-yl)-4-pyrimidineamine was reacted with 6-amino-1-(2-ethoxycarbonyl)indazoline to provide N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine Purification of the crude gave two products.

N4-[1-(2-Ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine (R935295):

¹H NMR (DMSO-*d*₆): δ 9.54 (s, 1H), 9.41 (s, 1H), 8.21 (app d, 1H, J = 1.7 Hz), 8.17 (d, 1H, J = 3.5 Hz), 8.01 (s, 1H), 7.86 (s, 1H), 7.83-7.80 (m 2H), 7.68 (d, 1H, J = 8.8 Hz), 7.59 (s, 1H), 7.52 (d, 1H, J = 8.2 Hz), 7.25 (d, 1H, J = 8.2 Hz), 4.12 (t, 2H, J = 6.4 Hz), 3.91 (qt, 2H, J = 7.0 Hz), 3.88 (s, 3H), 2.72 (t, 2H, J = 6.4 Hz), 1.02 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 25.67 min.; purity: 91%; MS (*m/e*): 519 (MH⁺) and

N4-[1-(2-carboxyethyl)indazoline-6-yl]- 5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine (R935296)

¹H NMR (DMSO-d₆): δ 9.54 (s, 1H), 9.39 (s, 1H), 8.23 (app d, 1H, J = 1.7 Hz), 8.17 (d, 1H, J = 3.5 Hz), 8.00 (s, 1H), 7.86 (s, 1H), 7.83-7.80 (m 2H), 7.68 (d, 1H, J = 8.8 Hz), 7.58 (s, 1H), 7.52 (d, 1H, J = 8.2 Hz), 7.28 (d, 1H, J = 8.2 Hz), 4.13 (t, 2H, J = 6.4 Hz), 3.88 (s, 3H), 2.67 (t, 2H, J = 6.4 Hz). LCMS: ret. time: 23.28 min.; purity: 91%; MS (*m/e*): 491 (MH⁺).

7.3.1045 5-Fluoro-N4-[2-(*N*-methylaminocarbonyl)benzofuran-5-yl]-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-6-yl]-2,4-pyrimidinediamine (R935297)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N4-[2-(*N*-methylaminocarbonyl)benzofuran-5-yl]-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.00 (s, 1H), 9.90 (s, 1H), 8.70 (qt, 1H, J = 4.7 Hz), 8.24 (d, 1H, J = 4.1 Hz), 8.12 (d, 1H, J = 1.7 Hz), 7.911 (s, 1H), 7.86 (s, 1H), 7.81 (qt, 1H, J = 4.7 Hz), 7.71 (d, 2H, J = 1.7 and 8.8 Hz), 7.57 (dd, 1H, J = 3.5 and 8.8 Hz), 7.35 (s, 1H), 7.26 (dd, 1H, J = 3.5 and 8.8 Hz), 4.19 (t, 2H, J = 7.0 Hz), 2.53 (t, 2H, J = 7.0 Hz), 2.47 (d, 6H, J = 4.7 Hz). LCMS: ret. time: 20.18 min.; purity: 89%; MS (*m/e*): 503 (MH⁺).

7.3.1046 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-(2-methyl-indazoline-5-yl)-2,4-pyrimidinediamine (R935298)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine and 5-amino-2-methylindazoline were reacted to give 5-fluoro-N4-(4-isopropoxyphenyl)-N2-(2-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.15 (s, 1H), 9.03 (s, 1H), 8.03-8.00 (m, 3H), 7.60 (dd, 2H, J = 4.1 and 8.8 Hz), 7.42 (d, 1H, J = 9.3 Hz), 7.31 (d, 1H, J = 9.3 Hz), 6.86 (d, 2H, J = 8.8 Hz), 4.57 (sept, 1H, J = 6.4 Hz), 4.08 (s, 3H), 1.26 (d, 6H, J = 6.4 Hz), LCMS: ret. time: 23.89 min.; purity: 98%; MS (*m/e*): 393 (MH⁺).

7.3.1047 5-Fluoro-N4-(3-hydroxyphenyl)-N2-(2-methy-indazoline-5-yl)-2,4-pyrimidinediamine (R935299)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 5-amino-2-methylindazoline to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-(2-methy-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.35 (s, 1H), 10.30 (s, 1H), 9.62 (br s, 1H), 8.22 (d, 1H, J = 5.3 Hz), 8.13 (s, 1H), 7.85 (s, 1H), 7.49 (d, 1H, J = 8.8 Hz), 7.17 (dd, 1H, J = 1.7 and 9.3 Hz), 7.08 (d, 2H, J = 5.3 Hz), 7.03 (s, 1H), 6.64-6.60 (m, 1H), 4.09 (s, 3H). LCMS: ret. time: 20.01 min.; purity: 97%; MS (*m/e*): 351 (MH⁺).

7.3.1048 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-(2-methy-indazoline-5-yl)-2,4-pyrimidinediamine (R935300)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 5-amino-2-methylindazoline to produce N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(2-methy-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.64 (s, 1H), 10.62 (s, 1H), 8.22 (d, 1H, J = 5.3 Hz), 8.21 (s, 1H), 7.77 (s, 1H), 7.58 (d, 1H, J = 9.3 Hz), 7.23-7.19 (m, 2H), 7.10 (dd, 1H, J = 2.3 and 8.8 Hz), 6.78 (d, 1H, J = 8.8 Hz), 4.21 (s, 3H), 4.15 (s, 4H). LCMS: ret. time: 21.77 min.; purity: 92%; MS (*m/e*): 393 (MH⁺).

7.3.1049 N2-[1-(2-Ethoxycarbonyl)ethyl]indazoline-5-yl]- N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R935301)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 5-amino-1-(2-ethoxycarbonyl)ethylindazoline to provide N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]- N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.15 (s, 1H), 9.13 (s, 1H), 8.10, (s, 1H), 8.04 (d, 1H, J = 3.5 Hz), 7.83 (s, 1H), 7.50 (s, 2H), 7.30 (d, 1H, J = 2.3 and 8.8 Hz), 6.79 (d, 1H, J = 8.8 Hz), 4.55 (t, 2H, J = 6.4 Hz), 4.22 (s, 4H), 3.97 (qt, 2H, J = 7.0 Hz), 2.88 (t, 2H, J = 6.4 Hz), 1.06 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 25.19 min.; purity: 93%; MS (*m/e*): 479 (MH⁺).

7.3.1050 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine (R935302)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]- N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine was reacted with DIBAL-H to produce N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.14 (s, 1H), 9.13 (s, 1H), 8.09 (s, 1H), 8.03 (d, 1H, J = 4.1 Hz), 7.82 (s, 1H), 7.48 (s, 2H), 7.30 (d, 1H, J = 2.3 Hz), 7.18 (dd, 1H, J = 2.3 and 8.8 Hz), 6.78 (d, 1H, J = 8.8 Hz), 4.59 (t, 1H, J = 6.4 Hz), 4.37 (t, 2H, J = 6.4 Hz), 4.22 (s, 4H), 3.34 (t, 2H, J = 6.4 Hz), 1.84 (q, 2H, J = 6.4 Hz). LCMS: ret. time: 22.33 min.; purity: 100%; MS (m/e): 437 (MH⁺).

7.3.1051 N4-[1-(2-Ethoxycarbonylethyl)indazoline-5-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine (R935303)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 5-amino-1-(2-ethoxycarbonylethyl)indazoline to provide N4-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.50 (s, 1H), 10.46 (s, 1H), 9.62 (br s 1H), 8.28 (d, 1H, J = 5.8 Hz), 7.96 (s, 2H), 7.65 (d, 1H, J = 8.8 Hz), 7.36 (dd, 1H, J = 1.7 and 8.8 Hz), 7.15-7.08 (m, 3H), 6.67-6.64 (m, 1H), 4.59 (t, 2H, J = 6.4 Hz), 3.97 (qt, 2H, J = 7.0 Hz), 2.89 (t, 2H, J = 6.4 Hz), 1.06 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 23.68 min.; purity: 97%; MS (m/e): 437 (MH⁺).

7.3.1052 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine (R935304)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.39 (s, 1H), 9.18 (s, 1H), 9.15 (s, 1H), 8.20 (s, 1H), 8.07 (d, 1H, J = 4.1 Hz), 7.84 (s, 1H), 7.46 (s, 2H), 7.24 (d, 1H, J = 8.2 Hz), 7.11-7.06 (m, 2H), 6.53 (d, 1H, J = 8.8 Hz), 4.56 (t,

1H, J = 4.7 Hz), 4.37 (t, 2H, J = 6.4 Hz), 3.34 (t, 2H, J = 6.4 Hz), 1.92 (q, 2H, J = 6.4 Hz). LCMS: ret. time: 24.70 min.; purity: 90%; MS (*m/e*): 479 (MH⁺). LCMS: ret. time: 20.89 min.; purity: 98%; MS (*m/e*): 395 (MH⁺).

7.3.1053 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-5-yl]-2,4-pyrimidinediamine (R935305)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]- 5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.39 (s, 1H), 9.18 (s, 1H), 9.15 (s, 1H), 8.19 (s, 1H), 8.06 (d, 1H, J = 3.5 Hz), 7.84 (s, 1H), 7.82 (qt, 1H, J = 4.7 Hz), 7.46 (t, 2H, J = 8.2 Hz), 7.25 (d, 1H, J = 8.2 Hz), 7.11 (d, 1H, J = 8.2 Hz), 7.10 (d, 1H, J = 8.2 Hz), 6.53 (t, 1H, J = 8.2 Hz), 4.51 (t, 2H, J = 7.0 Hz), 2.61 (t, 2H, J = 7.0 Hz), 2.49 (d, 3H, J = 4.7 Hz). LCMS: ret. time: 20.66 min.; purity: 95%; MS (*m/e*): 422 (MH⁺).

7.3.1054 N4-[1-(2-Ethoxycarbonyl)ethyl]indazoline-5-yl]- 5-fluoro-N2-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R935306)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine was reacted with 5-amino-1-(2-ethoxycarbonyl)ethylindazoline to provide N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]- 5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 10.48 (s, 1H), 10.41 (s, 1H), 8.25 (d, 1H, J = 5.8 Hz), 7.93 (s, 1H), 7.84 (s, 1H), 7.66 (d, 1H, J = 8.8 Hz), 7.49 (d, 2H, J = 8.8 Hz), 7.36 (dd, 1H, J = 2.3 and 8.8 Hz), 6.86 (d, 2H, J = 8.8 Hz), 4.59 (t, 2H, J = 6.4 Hz), 4.57 (sept, 1H, J = 7.0 Hz), 3.96 (qt, 2H, J = 7.0 Hz), 2.89 (t, 2H, J = 6.4 Hz), 1.23 (d, 6H, J = 7.0 Hz), 1.05 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 27.39 min.; purity: 98%; MS (*m/e*): 479 (MH⁺).

7.3.1055 5-Fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine (R935307)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]- 5-fluoro-N4-(4-isopropoxyphenyl)-2,4-

pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.16 (s, 1H), 9.10 (s, 1H), 8.09 (s, 1H), 8.03 (d, 1H, J = 4.1 Hz), 7.79 (s, 1H), 7.57 (d, 2H, J = 8.8 Hz), 7.46 (t, 2H), 6.87 (d, 2H, J = 8.8 Hz), 4.60-4.52 (m, 2H), 4.37 (t, 2H, J = 6.4 Hz), 3.34 (t, 2H, J = 6.4 Hz), 1.84 (q, 2H, J = 6.4 Hz), 1.24 (d, 6H, J = 7.0 Hz). LCMS: ret. time: 23.71 min.; purity: 98%; MS (*m/e*): 437 (MH⁺).

7.3.1056 5-Fluoro-N4-(2-hydroxymethylbenzofur-5-yl)- N2-[1-(3-hydroxypropyl)indazoline-6-yl]- 2,4-pyrimidinediamine (R935308)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-6-yl]- 5-fluoro-N2-(2-methoxycarbonylbenzofur-5-yl)-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N4-(2-hydroxymethylbenzofur-5-yl)- N2-[1-(3-hydroxypropyl)indazoline-6-yl]- 2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.35 (s, 1H), 9.33 (s, 1H), 8.12 (d, 1H, J = 3.5 Hz), 7.99 (d, 1H, J = 1.7 Hz), 7.95 (s, 1H), 7.84 (s, 1H), 7.55-7.49 (m, 3H), 7.28 (d, 1H, J = 8.8 Hz), 6.62 (s, 1H), 5.46 (t, 1H, J = 5.8 Hz), 4.55 (d, 2H, J = 5.8 Hz), 4.45 (t, 1H, J = 4.7 Hz), 3.96 (t, 2H, J = 6.4 Hz), 3.20 (t, 2H, J = 6.4 Hz), 1.76 (q, 2H, J = 6.4 Hz). LCMS: ret. time: 20.86 min.; purity: 99%; MS (*m/e*): 449 (MH⁺).

7.3.1057 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-5-yl]-2,4-pyrimidinediamine (R935309)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]- N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.12 (s, 1H), 9.11 (s, 1H), 8.09 (s, 1H), 8.03 (d, 1H, J = 3.5 Hz), 7.82 (s, 2H), 7.47 (s, 2H), 7.32-7.30 (m, 1H), 7.22-7.17 (m, 1H), 6.80 (d, 1H, J = 8.8 Hz), 4.51 (t, 2H, J = 7.0 Hz), 4.22 (s, 4H), 2.62 (t, 2H, J = 7.0 Hz), 2.49 (d, 3H, J = 4.7 Hz). LCMS: ret. time: 18.67 min.; purity: 100%; MS (*m/e*): 464 (MH⁺).

7.3.1058 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]-indazoline-5-yl]-2,4-pyrimidinediamine (R935310)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(*N*-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]- 5-fluoro-N4-(4-isopropoxyphenyl)-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[1-[2(*N*-methylaminocarbonyl)ethyl]indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.18 (s, 1H), 9.09 (s, 1H), 8.08 (s, 1H), 8.02 (d, 1H, J = 4.1 Hz), 7.82 (qt, 1H, J = 4.7 Hz), 7.79 (s, 1H), 7.57 (d, 2H, J = 8.8 Hz), 7.45 (d, 2H, J = 8.8 Hz), 6.87 (d, 2H, J = 8.8 Hz), 4.57 (q, 2H, J = 5.8 Hz), 4.51 (t, 2H, J = 7.0 Hz), 2.61 (t, 2H, J = 7.0 Hz), 2.47 (d, 3H, J = 4.7 Hz), 1.26 (d, 6H, J = 5.8 Hz). LCMS: ret. time: 17.14 min.; purity: 99%; MS (*m/e*): 464 (MH⁺).

7.3.1059 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]- 2,4-pyrimidinediamine (R935320)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 6-amino-1-(2-methoxy-3-carbomethoxybenzyl)indazoline to provide N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[1-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.36 (s, 1H), 9.18 (s, 1H), 8.08 (d, 1H, J = 3.5 Hz), 8.04 (s, 1H), 7.56 (d, 1H, J = 8.2 Hz), 7.45 (d, 1H, J = 1.8 Hz), 7.43-7.38 (m, 1H), 7.36-7.34 (m, 1H), 7.30 (dd, 1H, J = 1.7 and 8.8 Hz), 7.20 (dd, 1H, J = 2.3 and 8.8 Hz), 6.75 (d, 1H, J = 8.8 Hz), 6.68 (d, 1H, J = 8.2 Hz), 6.48 (dd, 1H, J = 1.7 and 8.2 Hz), 5.39 (s, 2H), 4.16 (s, 4H), 3.83 (s, 3H), 3.79 (s, 3H). LCMS: ret. time: 29.92 min.; purity: 80%; MS (*m/e*): 557 (MH⁺).

7.3.1060 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]- 2,4-pyrimidinediamine (R935321)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 6-amino-1-(2-methoxy-3-carbomethoxybenzyl)indazoline to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR

(DMSO- d_6): δ 9.37 (s, 1H), 9.31(s, 1H), 9.23 (s, 1H), 8.11 (d, 1H, J = 3.5 Hz), 8.08 (s, 1H), 7.93 (s, 1H), 7.57 (d, 1H, J = 8.8 Hz), 7.45 (d, 1H, J = 1.7 Hz), 7.40 (dd, 1H, J = 1.7 and 8.8 Hz), 7.33-7.27 (, 2H), 7.13 (t, 1H, J = 1.7 HZ), 7.03 (t, 2H, J = 8.2 Hz), 6.67 (d, 1H, J = 8.2 Hz), 6.45 (dd, 1H, J = 1.7 and 8.2 Hz), 5.37 (s, 2H), 3.83 (s, 3H), 3.79 (s, 3H). LCMS: ret. time: 28.80 min.; purity: 92%; MS (m/e): 515 (MH^+).

7.3.1061 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-6-yl]-2,4-pyrimidinediamine (R935322)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine was reacted with 6-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-6-yl]-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 9.60 (s, 2H), 8.11 (d, 1H, J = 4.1 Hz), 8.00-7.92 (m, 3H), 7.61-7.53 (m, 4H), 7.47-7.24 (m, 5H), 6.81 (d, 2H, J = 8.8 Hz), 6.68 (d, 1H, J = 8.2 Hz), 5.34 (s, 2H), 4.48 (sept, 1H, J = 5.9 Hz), 3.82 (s, 3H), 2.55 (s, 3H), 1.21 (d, 6H, J = 5.9 Hz). LCMS: ret. time: 30.57 min.; purity: 95%; MS (m/e): 696 (MH^+).

7.3.1062 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N4-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-6-yl]-2,4-pyrimidinediamine (R935323)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 6-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline to provide N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-6-yl]-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 9.53 (s, 1H), 9.41 (s, 1H), 8.05 (d, 1H, J = 4.1 Hz), 7.96-7.90 (m, 3H), 7.55 (d, 1H, J = 8.8 Hz), 7.49 (dd, 1H, J = 7.6 Hz), 7.42-7.20 (m, 6H), 7.14-7.10 (m, 1H), 6.69 (d, 1H, J = 8.2 Hz), 6.60 (d, 1H, J = 8.8 Hz), 5.33 (s, 2H), 4.10 (s, 4H), 3.77 (s, 3H), 2.50 (s, 3H). LCMS: ret. time: 32.11 min.; purity: 93%; MS (m/e): 696 (MH^+).

7.3.1063 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-6-yl]-2,4-pyrimidinediamine (R935324)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 6-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.64 (s, 1H), 9.56 (s, 1H), 8.15 (d, 1H, J = 4.1 Hz), 8.00 (s, 1H), 7.97 (d, 2H, J = 8.8 Hz), 7.60 (d, 1H, J = 8.8 Hz), 7.53 (d, 1H, J = 1.2 and 8.8 Hz), 7.47-7.23 (m, 6H), 7.11 (t, 1H, J = 1.7 Hz), 7.03 (t, 1H, J = 8.2 Hz), 6.62 (d, 1H, J = 8.2 Hz), 6.48 (dd, 1H, J = 1.7 and 8.2 Hz), 5.36 (s, 2H), 3.82 (s, 3H), 2.55 (s, 3H). LCMS: ret. time: 29.79 min.; purity: 92%; MS (*m/e*): 654 (MH⁺).

7.3.1064 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[2-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]- 2,4-pyrimidinediamine (R935336)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 6-amino-2-(2-methoxy-3-carbomethoxybenzyl)indazoline to provide N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[2-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-*d*₆): δ 9.16 (s, 1H), 9.14 (s, 1H), 8.24 (s, 1H), 8.06 (s, 1H), 8.04 (d, 1H, J = 3.5 Hz), 7.51 (d, 2H, J = 7.7 Hz), 7.49 (s, 1H), 7.29-7.26 (m, 2H), 7.19 (d, 1H, J = 7.7 Hz), 6.92 (d, 1H, J = 8.8 Hz), 6.76 (d, 1H, J = 8.2 Hz), 5.58 (s, 2H), 4.22 (s, 4H), 3.92 (s, 3H), 3.82 (s, 3H). LCMS: ret. time: 10.91 min.; purity: 91%; MS (*m/e*): 557 (MH⁺).

7.3.1065 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[2-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]- 2,4-pyrimidinediamine (R935337)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 6-amino-2-(2-methoxy-3-carbomethoxybenzyl)indazoline to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[2-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]-2,4-pyrimidinediamine. ¹H NMR

(DMSO- d_6): δ 9.31 (s, 1H), 9.17 (s, 1H), 9.15 (s, 1H), 8.26 (s, 1H), 8.09 (d, 1H, $J = 5.8$ Hz), 8.08 (s, 1H), 7.52 (app t, 3H, $J = 7.6$ Hz), 7.42 (d, 1H, $J = 8.2$ Hz), 7.23 (d, 1H, $J = 8.2$ Hz), 7.08 (app s, 1H), 7.03 (d, 1H, $J = 8.2$ Hz), 6.93 (d, 1H, $J = 7.6$ Hz), 6.43 (d, 1H, $J = 8.2$ Hz), 5.57 (s, 2H), 3.90 (s, 3H), 3.82 (s, 3H). LCMS: ret. time: 10.51 min.; purity: 93%; MS (m/e): 515 (MH^+).

7.3.1066 5-Fluoro-N4-(4-isopropoxyphenyl)-N2-[2-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]-2,4-pyrimidinediamine (R935338)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(4-isopropoxyphenyl)-4-pyrimidineamine was reacted with 6-amino-2-(2-methoxy-4-carbomethoxybenzyl)indazoline to provide 5-fluoro-N4-(4-isopropoxyphenyl)-N2-[2-(2-methoxy-4-carbomethoxybenzyl)indazoline-6-yl]-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 9.20 (s, 1H), 9.16 (s, 1H), 8.26 (s, 1H), 8.16 (s, 1H), 8.06 (d, 1H, $J = 3.5$ Hz), 7.66 (d, 2H, $J = 8.8$ Hz), 7.52-7.48 (m, 3H), 7.15 (d, 1H, $J = 8.2$ Hz), 6.86 (d, 2H, $J = 8.8$ Hz), 6.81 (d, 1H, $J = 8.8$ Hz), 5.56 (s, 2H), 4.46 (sept, 1H, $J = 5.9$ Hz), 3.91 (s, 3H), 3.82 (s, 3H), 1.17 (d, 6H, $J = 5.9$ Hz). LCMS: ret. time: 11.94 min.; purity: 90%; MS (m/e): 557 (MH^+).

7.3.1067 N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-5-yl]-2,4-pyrimidinediamine (R935339)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-ethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine was reacted with 5-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline to provide N4-(3, 4-Ethylenedioxyphenyl)-5-fluoro-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-5-yl]-2,4-pyrimidinediamine. ^1H NMR (DMSO- d_6): δ 9.57 (br s, 2H), 8.08 (d, 1H, $J = 3.5$ Hz), 8.01 (s, 1H), 7.99 (d, 1H, $J = 1.0$ Hz), 7.95 (s, 1H), 7.59-7.32 (m, 3H), 7.45-7.32 (m, 4H), 7.27-7.24 (m, 1H), 7.17-7.12 (m, 1H), 6.74 (d, 1H, $J = 8.7$ Hz), 6.65 (d, 1H, $J = 8.7$ Hz), 5.58 (s, 2H), 4.15 (s, 4H), 3.88 (s, 3H), 2.56 (s, 3H). LCMS: ret. time: 11.33 min.; purity: 98%; MS (m/e): 696 (MH^+).

7.3.1068 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-5-yl]-2,4-pyrimidinediamine (R935340)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-hydroxyphenyl)-4-pyrimidineamine was reacted with 5-amino-1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline to provide 5-fluoro-N4-(3-hydroxyphenyl)-N2-[1-[2-methoxy-4-(*o*-toluylsulfonamidocarboxy)benzyl]indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.57 (s, 1H), 9.48 (s, 1H), 8.13 (app s, 2H), 8.00 (d, 1H, J = 8.2 Hz), 7.94 (s, 1H), 7.59-7.32 (m, 7H), 7.18 (d, 1H, J = 8.2 Hz), 7.06 (app t, 3H, J = 8.8 Hz), 6.64 (d, 1H, J = 8.2 Hz), 6.55 (d, 1H, J = 8.2 Hz), 5.57 (s, 2H), 3.88 (s, 3H), 2.56 (s, 3H). LCMS: ret. time: 10.16 min.; purity: 97%; MS (*m/e*): 654 (MH⁺).

7.3.1069 N4-(4-Chlorophenyl)-5-fluoro-N2-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine (R935351)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(4-chlorophenyl)-5-fluoro 4-pyrimidineamine and 1-methyl-5-aminoindazoline were reacted to give N4-(4-chlorophenyl)-5-fluoro-N2-(1-methyl-indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.86 (s, 1H), 9.61 (s, 1H), 8.17 (d, 1H, J = 4.1 Hz), 8.00 (s, 1H), 7.88 (s, 1H), 7.78 (d, 2H, J = 8.8 Hz), 7.57 (d, 1H, J = 8.8 Hz), 7.43 (d, 1H, J = 8.8 Hz), 7.34 (d, 2H, J = 8.8 Hz), 4.00 (s, 3H). LCMS: ret. time: 10.64 min.; purity: 94%; MS (*m/e*): 369 (MH⁺).

7.3.1070 N4-(4-Chlorophenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine (R935352)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(4-chlorophenyl)-5-fluoro 4-pyrimidineamine and 6-aminoindazoline were reacted to give N4-(4-chlorophenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.18 (s, 1H), 10.02 (s, 1H), 8.26 (d, 1H, J = 4.1 Hz), 7.98 (s, 1H), 7.84 (d, 1H, J = 8.8 Hz), 7.82 (d, 2H, J = 8.8 Hz), 7.65 (d, 1H, J = 8.8 Hz), 7.35 (d, 2H, J = 8.8 Hz), 7.19 (d, 1H, J = 8.8 Hz). LCMS: ret. time: 10.80 min.; purity: 90%; MS (*m/e*): 355 (MH⁺).

7.3.1071 N4-(4-Chlorophenyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine (R935353)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(4-chlorophenyl)-5-fluoro-4-pyrimidineamine was reacted with 5-amino-1-(2-ethoxycarbonylethyl)indazoline to provide N4-(4-chlorophenyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.37 (s, 1H), 10.17 (s, 1H), 8.26 (d, 1H, J = 5.3 Hz), 7.96 (s, 1H), 7.88 (s, 1H), 7.33-7.66 (m, 3H), 7.40 (d, 1H, J = 8.8 Hz), 7.35 (d, 2H, J = 8.8 Hz), 4.61 (t, 2H, J = 6.4 Hz), 3.97 (qt, 2H, J = 7.0 Hz), 2.91 (t, 2H, J = 6.4 Hz), 1.05 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 11.85 min.; purity: 95%; MS (m/e): 455 (MH⁺).

7.3.1072 N4-(3-Chloro-4-trifluoromethoxy-phenyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine (R935354)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3-chloro-trifluoromethoxy-phenyl)-5-fluoro-4-pyrimidineamine was reacted with 5-amino-1-(2-ethoxycarbonylethyl)indazoline to provide N4-(3-chloro-4-trifluoromethoxy-phenyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.63 (s, 1H), 9.30 (s, 1H), 8.17 (d, 1H, J = 3.5 Hz), 8.10 (t, 1H, J = 2.3 Hz), 8.01 (s, 1H), 7.87 (s, 1H), 7.86 (d, 1H, J = 8.2 Hz), 7.57 (d, 1H, J = 9.4 Hz), 7.47 (t, 2H, J = 10.0 Hz), 4.56 (t, 2H, J = 6.9 Hz), 3.97 (qt, 2H, J = 7.0 Hz), 2.88 (t, 2H, J = 6.9 Hz), 1.06 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 14.4 min.; purity: 95%; MS (m/e): 539 (MH⁺).

7.3.1073 N4-(3, 4-Dichlorophenyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine (R935355)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-dichlorophenyl)-5-fluoro 4-pyrimidineamine and 1-methyl-5-aminoindazoline were reacted to give N4-(3, 4-dichlorophenyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.63 (s, 1H), 9.35 (s, 1H), 8.17 (d, 1H, J = 3.5 Hz), 8.08 (s, 1H), 8.01 (s, 1H), 7.86 (s, 1H), 7.79 (d, 1H, J = 8.8 Hz), 7.53 (d, 2H, J = 8.2 Hz), 7.47 (d, 1H, J = 8.2 Hz), 3.99 (s, 3H). LCMS: ret. time: 12.30 min.; purity: 98%; MS (m/e): 404 (MH⁺).

7.3.1074 5-Fluoro-N2-(1-methylindazoline-5-yl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R935356)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-trifluoromethoxyphenyl)-4-pyrimidineamime and 1-methyl-5-aminoindazoline were reacted to give 5-fluoro-N2-(1-methylindazoline-5-yl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.37 (s, 1H), 10.17 (s, 1H), 8.25 (d, 1H, J = 4.1 Hz), 7.92 (s, 2H), 7.84 (d, 1H, J = 9.4 Hz), 7.75 (s, 1H), 7.59 (d, 1H, J = 8.8 Hz), 7.45 (d, 1H, J = 9.4 Hz), 7.38 (d, 1H, J = 9.4 Hz), 7.08 (d, 1H, J = 8.8 Hz), 3.99 (s, 3H). LCMS: ret. time: 12.13 min.; purity: 94%; MS (m/e): 419 (MH⁺).

7.3.1075 N4-(3, 4-Difluoromethylenedioxyphenyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine (R935357)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamime and 1-methyl-5-aminoindazoline were reacted to give N4-(3, 4-difluoromethylenedioxyphenyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.84 (s, 1H), 9.54 (s, 1H), 8.16 (d, 1H, J = 4.1 Hz), 8.00 (s, 2H), 7.87 (s, 1H), 7.55-7.32 (m, 4H), 3.99 (s, 3H). LCMS: ret. time: 11.26 min.; purity: 96%; MS (m/e): 415 (MH⁺).

7.3.1076 N4-(3, 4-Difluorophenyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine (R935358)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-difluorophenyl)-5-fluoro-4-pyrimidineamime and 1-methyl-5-aminoindazoline were reacted to give N4-(3, 4-difluorophenyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.50 (s, 1H), 9.27 (s, 1H), 8.13 (d, 1H, J = 4.1 Hz), 8.08 (app s, 2H), 7.85 (s, 1H), 7.50 (app s, 3H), 7.37 (q, 1H, J = 9.4 Hz), 3.99 (s, 3H). LCMS: ret. time: 10.42 min.; purity: 90%; MS (m/e): 371 (MH⁺).

7.3.1077 N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine (R935359)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3-chloro-4-trifluoromethoxyphenyl)-

5-fluoro-4-pyrimidineamine and 1-methyl-5-aminoindazole were reacted to give N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(1-methylindazole-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.79 (s, 1H), 9.45 (s, 1H), 8.19 (d, 1H, J = 4.1 Hz), 8.09 (t, 1H, J = 2.8 Hz), 8.00 (s, 1H), 7.85-7.81 (m, 2H), 7.51 (d, 1H, J = 8.8 Hz), 7.48-7.44 (m, 2H), 3.99 (s, 3H). LCMS: ret. time: 13.14 min.; purity: 92%; MS (*m/e*): 453 (MH⁺).

7.3.1078 N2-[1-(2-Ethoxycarbonylethyl)indazole-5-yl]-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R935360)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-trifluoromethoxyphenyl)-4-pyrimidineamine was reacted with 5-amino-1-(2-ethoxycarbonylethyl)indazole to provide N2-[1-(2-ethoxycarbonylethyl)indazole-5-yl]-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.55 (s, 1H), 9.26 (s, 1H), 8.15 (d, 1H, J = 3.5 Hz), 8.05 (s, 1H), 7.95 (d, 1H, J = 8.2 Hz), 7.88 (s, 1H), 7.78 (s, 1H), 7.58 (dd, 1H, J = 8.8 and 7.4 Hz), 7.39 (t, 1H, J = 8.2 Hz), 7.01 (d, 1H, J = 8.8 Hz), 4.56 (t, 2H, J = 7.0 Hz), 3.97 (q, 4H, J = 7.0 Hz), 2.88 (t, 2H, J = 7.0 Hz), 1.06 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 13.22 min.; purity: 95%; MS (*m/e*): 505 (MH⁺).

7.3.1079 5-Fluoro-N2-[1-[2(N-methylaminocarbonyl)ethyl]indazole-5-yl]-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R935361)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonylethyl)indazole-5-yl]-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide 5-fluoro-N2-[1-[2(N-methylaminocarbonyl)ethyl]indazole-5-yl]-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.55 (s, 1H), 9.25 (s, 1H), 8.15 (d, 1H, J = 3.5 Hz), 8.04 (s, 1H), 7.96 (d, 1H, J = 8.2 Hz), 7.87 (s, 1H), 7.83 (qt, 1H, J = 4.9 Hz), 7.70 (s, 1H), 7.49 (dd, 2H, J = 8.2 and 9.4 Hz), 7.40 (d, 1H, J = 8.8 Hz), 7.01 (d, 1H, J = 8.8 Hz), 4.52 (t, 2H, J = 7.0 Hz), 2.63 (t, 2H, J = 7.0 Hz), 2.49 (d, 3H, J = 4.7 Hz). LCMS: ret. time: 10.00 min.; purity: 100%; MS (*m/e*): 490 (MH⁺).

7.3.1080 5-Fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R935362)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine was reacted with DIBAL-H to produce 5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.55 (s, 1H), 9.24 (s, 1H), 8.15 (d, 1H, J = 2.9 Hz), 8.05 (s, 1H), 7.92 (d, 1H, J = 7.6 Hz), 7.87 (s, 1H), 7.78 (s, 1H), 7.50 (s, 2H), 7.39 (t, 1H, J = 8.2 Hz), 7.01 (d, 1H, J = 7.6 Hz), 4.56 (t, 1H, J = 5.2 Hz), 4.38 (t, 2H, J = 7.0 Hz), 3.35 (dd, 2H, J = 5.2 and 7.0 Hz), 1.84 (qt, 2H, J = 7.0 Hz). LCMS: ret. time: 10.42 min.; purity: 97%; MS (*m/e*): 463 (MH⁺).

7.3.1081 5-Fluoro-N2-(indazoline-6-yl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R935363)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-trifluoromethoxyphenyl)-4-pyrimidineamine and 6-aminoindazoline were reacted to give 5-fluoro-N2-(indazoline-6-yl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.72(s, 1H), 9.60 (s, 1H), 9.42 (s, 1H), 8.21 (d, 1H, J = 3.5 Hz), 8.06 (br s, 2H), 7.89 (s, 1H), 7.83 (s, 1H), 7.57 (d, 1H, J = 8.8 Hz), 7.42 (t, 1H, J = 8.2 Hz), 7.27 (d, 1H, J = 8.8 Hz), 7.00 (d, 1H, J = 8.2 Hz). LCMS: ret. time: 12.17 min.; purity: 97%; MS (*m/e*): 405 (MH⁺).

7.3.1082 5-Fluoro-N2-(indazoline-5-yl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine (R935364)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N-(3-trifluoromethoxyphenyl)-4-pyrimidineamine and 5-aminoindazoline were reacted to give 5-fluoro-N2-(indazoline-5-yl)-N4-(3-trifluoromethoxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.85 (s, 1H), 9.54 (s, 1H), 9.23 (s, 1H), 8.15 (d, 1H, J = 3.5 Hz), 8.05 (s, 1H), 7.93 (d, 1H, J = 8.2 Hz), 7.89 (s, 1H), 7.78 (s, 1H), 7.48-7.35 (m, 3H), 7.01 (d, 1H, J = 8.2 Hz). LCMS: ret. time: 10.44 min.; purity: 98%; MS (*m/e*): 405 (MH⁺).

7.3.1083 N4-(4-Chlorophenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine (R935365)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(4-chlorophenyl)-5-fluoro 4-pyrimidineamine and 5-aminoindazoline were reacted to give N4-(4-chlorophenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.85 (s, 1H), 9.43 (s, 1H), 9.19 (s, 1H), 8.11 (d, 1H, J = 3.5 Hz), 8.08 (s, 1H), 7.87 (s, 1H), 7.82 (dd, 2H, J = 3.0 and 8.8 Hz), 7.42 (dd, 2H, J = 3.0 and 8.8 Hz), 7.31 (d, 2H, J = 8.8 Hz). LCMS: ret. time: 9.07 min.; purity: 91%; MS (*m/e*): 355 (MH⁺).

7.3.1084 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine (R935366)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine and 5-aminoindazoline were reacted to give N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.90 (s, 1H), 9.45 (s, 1H), 9.27 (s, 1H), 8.15 (d, 1H, J = 3.5 Hz), 8.11 (t, 1H, J = 3.0 Hz), 8.02 (s, 1H), 7.87 (s, 1H), 7.84 (d, 1H, J = 8.8 Hz), 7.47 (d, 2H, J = 8.8 Hz), 7.44 (d, 1H, J = 8.8 Hz). LCMS: ret. time: 11.65 min.; purity: 98%; MS (*m/e*): 439 (MH⁺).

7.3.1085 5-Fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-N2-(3, 4,5-trimethoxyphenyl)-2,4-pyrimidinediamine (R935367)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-4-pyrimidineamine and 3, 4,5-trimethoxyaniline were reacted by microwave heating at 180 °C. Upon concentration of the ethanol and addition of 2N HCl provided 5-fluoro-N4-[4-(3-phenyl-1,2,4-oxadiazol-5-yl)methyleneoxyphenyl]-N2-(3, 4,5-trimethoxyphenyl)-2,4-pyrimidinediamine as fine flakes of the solid. ¹H NMR (DMSO-d₆): δ 9.59 (s, 1H), 9.25 (s, 1H), 8.09 (d, 1H, J = 3.5 Hz), 8.01 (dd, 2H, J = 5.3 and 1.2 Hz), 7.39 (dd, 2H, J = 3.1 and 8.8 Hz), 7.60-7.54 (m, 3H), 7.03 (d, 2H, J = 8.8 Hz), 6.94 (d, 2H, J = 3.1 Hz), 5.57 (s, 2H), 3.59 (s, 6H), 3.57 (s, 3H). LCMS: ret. time: 13.00 min.; purity: 97%; MS (*m/e*): 547 (MH⁺).

7.3.1086 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine (R935368)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-4-pyrimidineamine and 6-aminoindazoline were reacted to give N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.73 (s, 1H), 9.67 (s, 1H), 9.46 (s, 1H), 8.21 (d, 1H, J = 3.5 Hz), 8.17 (app d, 1H, J = 8.8 Hz), 8.04 (br s, 1H), 7.97 (dt, 1H, J = 2.4 and 9.3 Hz), 7.89 (s, 1H), 7.58 (d, 1H, J = 8.8 Hz), 7.47 (d, 1H, J = 9.3 Hz), 7.27 (dd, 1H, J = 1.7 and 8.8 Hz). LCMS: ret. time: 13.08 min.; purity: 96%; MS (m/e): 439 (MH⁺).

7.3.1087 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[1-[2(N-methylaminocarbonyl)ethyl]indazoline-5-yl]-2,4-pyrimidinediamine (R935369)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(3-chloro-4-trifluoromethoxyphenyl)-N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[1-[2(N-methylaminocarbonyl)ethyl]indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.62 (s, 1H), 9.29 (s, 1H), 8.17 (d, 1H, J = 3.5 Hz), 8.11 (t, 1H, J = 2.4 Hz), 8.02 (app s, 1H), 7.88-7.82 (m, 3H), 7.53 (d, 1H, J = 9.3 Hz), 7.47 (d, 2H, J = 8.8 Hz), 4.52 (t, 2H, J = 7.0 Hz), 2.48 (t, 2H, J = 7.0 Hz), 2.48 (d, 3H, J = 4.7 Hz). LCMS: ret. time: 10.51 min.; purity: 99%; MS (m/e): 524 (MH⁺).

7.3.1088 N4-(3-Chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine (R935370)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-(3-chloro-4-trifluoromethoxyphenyl)-N2-[1-(2-ethoxycarbonyl)ethyl]indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine was reacted with DIBAL-H to produce N4-(3-chloro-4-trifluoromethoxyphenyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.62 (s, 1H), 9.28 (s, 1H), 8.17 (d, 1H, J = 3.5 Hz), 8.11 (s, 1H), 8.02 (s, 1H), 7.85 (s, 2H), 7.53 (t, 2H, J = 8.8 Hz), 7.46 (t, 1H, J = 8.8

Hz), 4.56 (t, 1H, J = 5.8 Hz), 4.38 (t, 2H, J = 6.4 Hz), 3.35 (dd, 2H, J = 5.8 and 6.4 Hz), 1.93 (q, 2H, J = 6.4 Hz). LCMS: ret. time: 11.33 min.; purity: 99%; MS (*m/e*): 497 (MH⁺).

7.3.1089 N4-(3, 4-Dichlorophenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine (R935371)

5 In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-dichlorophenyl)-5-fluoro 4-pyrimidineamine and 5-aminoindazoline were reacted to give N4-(3, 4-dichlorophenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.90 (s, 1H), 9.60 (s, 1H), 8.20 (d, 1H, J = 4.2 Hz), 8.06 (t, 1H, J = 2.3 Hz), 7.92 (s, 2H), 7.73 (d, 1H, J = 8.8 Hz), 7.51-7.40 (m, 3H). LCMS: ret. time: 9.83 min.; purity: 98%; MS (*m/e*): 390 (MH⁺).

7.3.1090 N4-(3, 4-Dichlorophenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine (R935372)

15 In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-dichlorophenyl)-5-fluoro 4-pyrimidineamine and 6-aminoindazoline were reacted to give N4-(3, 4-dichlorophenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 12.82 (s, 1H), 9.63 (s, 1H), 9.48 (s, 1H), 8.22 (d, 1H, J = 4.3 Hz), 8.15 (t, 1H, J = 2.3 Hz), 8.02 (s, 1H), 7.92-7.90 (m, 2H), 7.59 (d, 1H, J = 8.8 Hz), 7.52 (d, 1H, J = 8.8 Hz), 7.26 (dd, 1H, J = 1.7 and 8.8 Hz). LCMS: ret. time: 11.73 min.; purity: 99%; MS (*m/e*): 390 (MH⁺).

7.3.1091 N4-(3, 4-Difluoromethylenedioxyphenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine (R935373)

25 In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 5-aminoindazoline were reacted to give N4-(3, 4-difluoromethylenedioxyphenyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.40 (s, 1H), 10.11 (s, 1H), 8.25 (d, 1H, J = 4.5 Hz), 7.95 (s, 1H), 7.89 (app s, 2H), 7.49 (d, 1H, J = 8.8 Hz), 7.37 (app d, 3H, J = 8.2 Hz). LCMS: ret. time: 8.56 min.; purity: 99%; MS (*m/e*): 401 (MH⁺).

7.3.1092 N4-(3, 4-Difluoromethylenedioxyphenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine (R935374)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(3, 4-difluoromethylenedioxyphenyl)-5-fluoro-4-pyrimidineamine and 6-aminoindazoline were reacted to give N4-(3, 4-difluoromethylenedioxyphenyl)-5-fluoro-N2-(indazoline-6-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.53 (s, 1H), 9.52 (s, 1H), 8.21 (d, 1H, J = 4.5 Hz), 8.10 (s, 1H), 8.01 (s, 1H), 7.92 (s, 1H), 7.59 (d, 1H, J = 8.8 Hz), 7.48 (dt, 1H, J = 2.3 and 8.8 Hz), 7.34 (d, 1H, J = 8.2 Hz), 7.21 (dd, 1H, J = 2.3 and 8.8 Hz). LCMS: ret. time: 11.29 min.; purity: 90%; MS (*m/e*): 401 (MH⁺).

7.3.1093 N4-(6-Chloro-3-pyridyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine (R935375)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(6-chloro-3-pyridyl)-5-fluoro-4-pyrimidineamine and 5-amino-1-methylindazoline were reacted to give N4-(6-chloro-3-pyridyl)-5-fluoro-N2-(1-methylindazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.96 (s, 1H), 9.58 (s, 1H), 8.86 (s, 1H), 8.25 (dt, 1H, J = 3.9 and 8.8 Hz), 8.20 (d, 1H, J = 4.1 Hz), 8.04 (s, 1H), 7.55 (d, 1H, J = 8.8 Hz), 7.44 (d, 2H, J = 8.8 Hz), 4.00 (s, 3H). LCMS: ret. time: 8.95 min.; purity: 100%; MS (*m/e*): 370 (MH⁺).

7.3.1094 N4-(6-Chloro-3-pyridyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine (R935376)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(6-chloro-3-pyridyl)-5-fluoro-4-pyrimidineamine and 5-aminoindazoline were reacted to give N4-(6-chloro-3-pyridyl)-5-fluoro-N2-(indazoline-5-yl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.78 (s, 1H), 9.41 (s, 1H), 8.88 (s, 1H), 8.24 (d, 1H, J = 8.2 Hz), 8.18 (d, 1H, J = 3.5 Hz), 8.06 (s, 1H), 7.92 (s, 1H), 7.42 (app s, 3H). LCMS: ret. time: 7.87 min.; purity: 90%; MS (*m/e*): 356 (MH⁺).

7.3.1095 N4-(6-Chloro-3-pyridyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine (R935377)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(6-chloro-3-pyridyl)-5-fluoro-4-pyrimidineamine and 5-amino-1-(2-ethoxycarbonylethyl)indazoline were reacted to give N4-(6-chloro-3-pyridyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 10.37 (s, 1H), 10.04 (s, 1H), 8.78 (s, 1H), 8.28 (d, 1H, J = 4.8 Hz), 8.20 (dt, 1H, J = 2.8 and 8.8 Hz), 7.96 (s, 1H), 7.92 (s, 1H), 7.65 (d, 1H, J = 8.8 Hz), 7.45 (d, 1H, J = 8.8 Hz), 7.41 (d, 1H, J = 8.8 Hz), 4.59 (t, 2H, J = 6.0 Hz), 3.97 (qt, 2H, J = 7.0 Hz), 2.90 (t, 2H, J = 6.4 Hz), 1.06 (t, 3H, J = 7.0 Hz). LCMS: ret. time: 10.87 min.; purity: 94%; MS (m/e): 456 (MH⁺).

7.3.1096 N4-(6-Chloro-3-pyridyl)-5-fluoro-N2-[1-[2(N-methylaminocarbonyl)ethyl]indazoline-5-yl]-2,4-pyrimidinediamine (R935378)

In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine, N4-(6-chloro-3-pyridyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine and Me₂NH.HCl were reacted to provide N4-(6-Chloro-3-pyridyl)-5-fluoro-N2-[1-[2(N-methylaminocarbonyl)ethyl]indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.67 (s, 1H), 9.31 (s, 1H), 8.88 (s, 1H), 8.27 (dt, 1H, J = 3.0 and 8.8 Hz), 8.17 (d, 1H, J = 3.5 Hz), 8.08 (s, 1H), 7.88 (s, 1H), 7.83 (q, 1H, J = 5.3 Hz), 7.53 (d, 1H, J = 8.8 Hz), 7.45 (d, 1H, J = 8.8 Hz), 4.53 (t, 2H, J = 7.0 Hz), 2.63 (t, 2H, J = 7.0 Hz), 2.49 (d, 3H, J = 5.3 Hz). LCMS: ret. time: 7.62 min.; purity: 89%; MS (m/e): 441 (MH⁺).

7.3.1097 N4-(6-Chloro-3-pyridyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine (R935379)

In like manner to the preparation of N2-(3, 4-ethylenedioxyphenyl)-5-fluoro-N4-[4(2-hydroxy-1,1-dimethylethyl)phenyl]-2,4-pyrimidinediamine, N4-(6-chloro-3-pyridyl)-N2-[1-(2-ethoxycarbonylethyl)indazoline-5-yl]-5-fluoro-2,4-pyrimidinediamine was reacted with DIBAL-H to produce N4-(6-chloro-3-pyridyl)-5-fluoro-N2-[1-(3-hydroxypropyl)indazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): LCMS: ret. time: 8.02 min.; purity: 98%; MS (m/e): 414 (MH⁺).

7.3.1098 N4-(2,6-Dimethoxy-3-pyridyl)-5-fluoro-N2-[1-methylindazoline-5-yl]-2,4-pyrimidinediamine (R935380)












In like manner to the preparation of N4-(3, 4-ethylenedioxyphenyl)-5-fluoro-N2-(3-hydroxyphenyl)-2,4-pyrimidinediamine, 2-chloro-N-(2,6-dimethoxy-3-pyridyl)-6-5-fluoro-4-pyrimidineamine and 1-methyl-5-aminoindazoline were reacted to give N4-(2,6-dimethoxy-3-pyridyl)-5-fluoro-N2-[1-methylindazoline-5-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.08 (s, 1H), 8.68 (s, 1H), 8.01 (d, 1H, J = 4.1 Hz), 7.96 (s, 1H), 7.76 (dd, 1H, J = 4.1 and 8.8 Hz), 7.65 (s, 1H), 7.37 (d, 1H, J = 8.8 Hz), 7.34 (d, 1H, J = 8.2 Hz), 6.46 (d, 1H, J = 8.2 Hz), 3.94 (s, 3H), 3.91 (s, 3H), 3.83 (s, 3H). LCMS: ret. time: 9.57 min.; purity: 92%; MS (m/e): 396 (MH⁺).

7.3.1099 Additional 2,4-Pyrimidinediamine

Compounds R008951, R008952, R008953, R008955, R008956, R008958, R070153 and R070790 (structures provided below) were purchased from Contact Services.

Additional compounds whose structures are provided below were synthesized using methods similar to those described in the previous examples.

R008951 R067962 R926209		R088814 R926017	
R008952 R067963		R088815	
R008953 R067964		R091880	
R008955 R081166		R092788	

R008956 R070791		R920846	
R008958			
R070153			
R070790 R926036		R926593	
R926736		R950189	
		R950216	
R935117		R950218	

7.3.1100 Synthesis of Intermediates, 2,4-Pyrimidinediamines and 2,4,6-Pyrimidinetriamines According to Schemes VIII and IX

A variety of intermediates and 2,4-pyrimidinediamine compounds were synthesized according to Schemes VIII and IX. Scheme VIII is exemplified by the reaction of 2,4,6-trichloropyrimidine with 3-hydroxyaniline to form a mixture of three compounds, which were separated and purified by chromatography. Scheme IX is exemplified by the reaction of 2,4,5,6-tetrachloridepyrimidine with 3,4-ethylenedioxyaniline to form a mixture of three compounds, which were separated and purified by chromatography.

7.3.1101 Reaction of 2,4,6-trichloropyrimidine with 3-hydroxyaniline**4,6-Dichloro-N2-(3-hydroxyphenyl)-2-pyrimidineamine (R926407)****N2,N4-Bis(3-hydroxyphenyl)-6-chloro-2,4-pyrimidinediamine (R926408) and****N2,N4,N6-Tris(3-hydroxyphenyl)-2,4,6-pyrimidinetriamine (R926409)**

A mixture of 2,4,6-trichloroaniline (0.183g, 1 mmol) and 3-hydroxyaniline (0.327g, 3 mmol) in 5 mL MeOH was heated at 100 °C in a sealed vial for 24h. The reaction mixture was diluted with H₂O, acidified with 2N HCl and extracted with EtOAc (3 x 50 mL). Upon removal of solvent the residue was purified by chromatography (as well as preparative TLC) to afford three products, mainly the mono-SNAr, 4,6-dichloro-N2-(3-hydroxyphenyl)-2-pyrimidineamine (**R926407**), ¹H NMR (CDCl₃): δ 7.16 (t, 1H, J= 8.1 Hz), 6.78 (m, 2H), 6.64 (dd, 1H, J= 1.2 and 8.1 Hz), 6.58 (s, 1H); LCMS: ret. time: 25.08 min.; purity: 99%; MS (m/e): 256 (M⁺); bis-SNAr product, N2,N4-bis(3-hydroxyphenyl)-6-chloro-2,4-pyrimidinediamine (**R926408**), ¹H NMR (CD₃OD): δ 7.21 (m, 1H), 7.14-7.03 (m, 5H), 6.50 (m, 1H), 6.44 (m, 1H), 6.16 (s, 1H); LCMS: ret. time: 25.14 min.; purity: 99%; MS (m/e): 329 (M⁺); and tris-SNAr product, N2,N4,N6-tris(3-hydroxyphenyl)-2,4,6-pyrimidinetriamine (**R926409**), ¹H NMR (CD₃OD): δ 7.29 (m, 1H), 7.12-7.05 (m, 5H), 7.02 (m, 2H), 6.88 (dd, 2H, J= 1.2 and 8.1 Hz), 6.46 (dd, 1H, J= 1.5 and 8.1 Hz), 6.41 (dt, 1H); LCMS: ret. time: 20.49 min.; purity: 94%; MS (m/e): 402 (MH⁺).

7.3.1102 N2,N4-Bis(4-methoxycarbonylmethyleneoxyphenyl)-6-chloro-2,4-pyrimidinediamine (R926411)

In like manner to the reaction of 2,4,6-trichloropyrimidine with 3-hydroxyaniline, the reaction of 2,4,6-trichloropyrimidine with methyl 4-aminophenoxyacetate gave N2,N4-bis(4-methoxycarbonylmethyleneoxyphenyl)-6-chloro-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.65 (bs, 1H), 7.40 (bd, 4H), 6.82 (bd, 4H), 6.00 (s, 1H), 6.62 (bs, 4H), 3.78 (bs, 6H); LCMS: ret. time: 29.87 min.; purity: 98%; MS (m/e): 473 (MH⁺).

7.3.1103 Reaction of 2,4,6-trichloropyrimidine with 3,4-ethylenedioxyaniline**4,6-Dichloro-N2-(3,4-ethylenedioxyphenyl)-2-pyrimidineamine (R926515)**

N2,N4-Bis(3,4-ethylenedioxyphenyl)-6-chloro-2,4-pyrimidinediamine (R926245)

N2,N4,N6 -Tris(3,4-ethylenedioxyphenyl)-2,4,6-pyrimidinetriamine (R926516)

A mixture of 2,4,6-trichloroaniline (1 mmol) and 3,4-ethylenedioxyaniline (3 mmol) in 5 mL MeOH was heated at 100 °C in a sealed vial for 24h. The reaction mixture was diluted with H₂O, acidified with 2N HCl and extracted with EtOAc (3 x 50 mL). Upon removal of solvent the residue was purified by chromatography (as well as preparative TLC) to afford three products, mainly the Mono-SNAr product, 4,6-dichloro-N2-(3,4-ethylenedioxyphenyl)-2-pyrimidineamine (**R926515**). ¹H NMR (CD₃OD): δ 7.05 (s, 1H), 6.83 (m, 2H), 6.45 (bs, 1H), 4.20 (bs, 4H); LCMS: ret. time: 29.75 min.; purity: 96%; MS (m/e): 298 (M⁺); Bis-SNAr product, N2,N4-bis(3,4-ethylenedioxyphenyl)-6-chloro-2,4-pyrimidinediamine (**R926245**):

¹H NMR (CDCl₃): δ 7.23 (d, 1H, J= 3 Hz), 6.90-6.70 (m, 6H), 6.02 (s, 1H), 4.26 (bs, 4H), 4.23 (m, 4H); LCMS: ret. time: 31.34 min.; purity: 95%; MS (m/e): 413 (MH⁺) and Tris-SNAr product, N2,N4,N6 -tris(3,4-ethylenedioxyphenyl)-2,4,6-pyrimidinetriamine (**R926516**)

¹H NMR (CD₃OD): δ 7.16 (d, 1H, J= 3Hz), 7.05 (bd, 1H), 6.99-6.90 (m, 3H), 6.80-6.70 (m, 4H), 6.03 (s, 1H), 4.22 (s, 4H), 4.20 (s, 8H); LCMS: ret. time: 27.72 min.; purity: 61%; MS (m/e): 528 (M⁺).

7.3.1104 Reaction of 2,4,6-trichloropyrimidine with ethyl-4-aminophenoxyacetate

4,6-Dichloro-N2-(4-ethoxycarbonylmethyl)-4,6-dichloro-2-pyrimidineamine (R926549)

2,6-Dichloro-N4-(ethoxycarbonylmethyl)-4-pyrimidineamine (R926550)

A mixture of 2,4,6-trichloroaniline (1 mmol) and ethyl 2-aminoacetate (3 mmol) in 5 mL MeOH was heated at 100 °C in a sealed vial for 24h. The reaction mixture was diluted with H₂O, acidified with 2N HCl and extracted with EtOAc (3 x 50 mL). Upon removal of solvent the residue was purified by chromatography (as well as preparative TLC) to afford three products, mainly the mono-SNAr product, 4,6-dichloro-N2-(4-ethoxycarbonylmethyl)-4,6-dichloro-2-pyrimidineamine (**R926549**). ¹H NMR (CDCl₃): δ 6.67 (s, 1H), 5.85 (bs, 1H), 4.23 (q, 2H, J= 7.2 Hz), 4.19 (s, 2H), 1.29 (t, 3H, J= 7.2 Hz);

LCMS: ret. time: 26.18 min.; purity: 100%; MS (m/e): 250 (MH⁺); and Mono-SNAr product, 2,6-dichloro-N4-(ethoxycarbonylmethyl)-4-pyrimidineamine (**R926550**): ¹H NMR (CDCl₃): d 6.37 (bs, 1H), 4.28 (q, 2H, J= 6.9 Hz), 4.19 (bs, 2H), 1.31 (t, 3H, J= 7.2 Hz)

7.3.1105 6-Chloro-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(methoxycarbonylmethyl)-2,4-pyrimidinediamine (R926555)

In like manner to the preparation of 4-[N-(L)-phenylalanine ethyl ester]-N2-(3-hydroxyphenyl)-5-ethoxycarbonyl-2-pyrimidineamine, the reaction of ethyl 4-aminophenoxyacetate with methyl 2-aminoacetate gave 6-chloro-N2-(4-ethoxycarbonylmethyleneoxyphenyl)-N4-(methoxycarbonylmethyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.40 (d, 2H, J= 8.7 Hz), 6.86 (d, 2H, J= 9.3 Hz), 5.97 (s, 1H), 4.64 (s, 2H), 4.26 (q, 2H, J= 7.2 Hz), 4.14 (q, 2H, J= 6.9 Hz), 4.05 (s, 2H), 1.25 (m, 6H); LCMS: ret. time: 26.21 min.; purity: 93%; MS (m/e): 409 (MH⁺).

7.3.1106 Reaction of 3,4-ethylenedioxyaniline with 2,4,5,6-tetrachloropyrimidine.

N4-(3,4-Ethylenedioxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (R926466)

N2,N4-Bis(3,4-ethylenedioxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (R926467) and

N4,N6-Bis(3,4-ethylenedioxyphenyl)-2,5-dichloro-4,6-pyrimidinediamine (R926468)

A mixture of 3,4-ethylenedioxyaniline (0.775 g, 5 mmol) and 2,4,5,6-tetrachloropyrimidine (0.434 g, 2 mmol) in the presence of DIPEA (1.043 mL, 6 mmol) in EtOAc (10 mL) was heated at 80 °C for 3 days. The reaction was diluted with water (50 mL), acidified (2N HCl) and extracted with EtOAc (3 x 50 mL). The residue obtained after removal of solvent was chromatographed using 5-30% EtOAc/hexanes to obtain three products viz. N4-(3,4-Ethylenedioxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (**R926466**): ¹H NMR (CDCl₃): δ 7.18 (d, 1H, J= 2.7 Hz), 6.92 (dd, 1H, J= 2.1 and 8.7 Hz), 6.87 (d, 1H, J= 9 Hz); LCMS: ret. time: 33.53 min.; purity: 100%; MS(m/e): 292 (MH⁺); N2,N4-Bis(3,4-ethylenedioxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (**R926467**): ¹H NMR (CDCl₃): δ 7.11 (d, 1H, J= 2.4 Hz), 7.06 (d, 1H, J= 2.1 Hz), 7.04 (s, 1H), 6.94 (m, 2H), 6.84 (d, 1H, J= 8.1 Hz), 6.76 (bd, 2H, J= 8.7 Hz), 4.27 (bs, 4H), 4.24 (bs, 1H); LCMS: ret. time:

26.54 min.; purity: 87%; MS(m/e): 364 (MH^+); and N4,N6-Bis(3,4-ethylenedioxyphenyl)-2,5-dichloro-4,6-pyrimidinediamine (**R926468**): ^1H NMR ($CDCl_3$): δ 7.07 (t, 1H, $J=2.4$ Hz), 6.99 (s, 2H), 6.83 (dd, 2H, $J=2.4$ and 8.7 Hz), 6.75 (dd, 2H, $J=1.8$ and 9 Hz), 4.19 (bs, 4H); LCMS: ret. time: 34.70 min.; purity: 99%; MS(m/e): 365 (MH^+).

5 **7.3.1107 Reaction of 2,4,5,6-tetrachloropyrimidine with ethyl-4-aminophenoxyacetate**

N4-(4-Ethoxycarbonylmethyleneoxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (R926568)

10 **N2,N4-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (R926569)**

N2,N5-Bis(4-ethoxycarbonylmethyleneoxyphenyl)-2,5-pyrimidinediamine (R926570)

In like manner to the reaction of 2,4,6-trichloropyrimidine with 3,4-ethylenedioxyaniline, the reaction of 2,4,5,6-tetrachloropyrimidine with ethyl 4-aminophenoxyacetate gave a mixture of mono-SNAr product, N4-(4-ethoxycarbonylmethyleneoxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (**R926568**): ^1H NMR ($CDCl_3$): δ 7.46 (dd, 2H, $J=2.4$ and 6.9 Hz), 7.3 (s, 1H), 6.95 (dd, 2H, $J=2.4$ and 6.9 Hz), 4.63 (s, 2H), 4.28 (q, 2H, $J=7.2$ Hz), 1.30 (t, 3H, $J=7.2$ Hz); LCMS: ret. time: 30.62 min.; purity: 99%; MS (m/e): 378 (MH^+); Bis-SNAr product, N2,N4-bis((4-ethoxycarbonylmethyleneoxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (**R926569**): ^1H NMR ($CDCl_3$): δ 7.42 (d, 2H, $J=9$ Hz), 7.35 (d, 2H, $J=8.7$ Hz), 6.90 (d, 2H, $J=9$ Hz), 6.83 (d, 2H, $J=8.7$ Hz), 4.67 (s, 2H), 4.60 (s, 2H), 4.28 (2q, 4H, $J=4.8$ Hz), 1.31 (2t, 6H, $J=6.3$ Hz); LCMS: ret. time: 33.09 min.; purity: 85%; MS (m/e): 537 (MH^+) and Bis-SNAr product, N2,N5-bis((4-ethoxycarbonylmethyleneoxyphenyl)-2,5-pyrimidinediamine (**R926570**): ^1H NMR ($CDCl_3$): δ 7.45 (d, 4H, $J=8.7$ Hz), 6.92 (d, 4H, $J=9$ Hz), 6.85 (s, 1H), 4.61 (s, 4H), 4.26 (q, 4H, $J=6.9$ Hz), 1.30 (t, 6H, $J=7.2$ Hz); LCMS: ret. time: 31.66 min.; purity: 97%; MS (m/e): 535 (MH^+).

7.3.1108 Reaction of 2,4,5,6-tetrachloropyrimidine with tert-Butyl-4-aminophenoxyacetate, N4-(4-tert-butoxyoxycarbonylmethyleneoxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (R926575), N2,N4-Bis(4-tert-butoxyoxycarbonylmethyleneoxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (R926576) and N4,N6-Bis(4-tert-butoxyoxycarbonylmethyleneoxyphenyl)-2,5-dichloro-4,6-pyrimidinediamine (R926577)

In like manner to the reaction of 2,4,6-trichloropyrimidine with 3,4-ethylenedioxyaniline, the reaction of 2,4,5,6-tetrachloropyrimidine with tert-butyl-4-aminophenoxyacetate gave a mixture of mono-SNAr product, N4-(4-tert-butoxyoxycarbonylmethyleneoxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (**R926575**): ¹H NMR (CDCl₃): δ 7.45 (dd, 2H, J= 2.4 and 7.2 Hz), 6.93 (dd, 2H, J= 2.4 and 7.2 Hz), 4.52 (s, 2H); LCMS: ret. time: 32.56 min.; purity: 100%; MS (m/e): 402 (MH⁺); Bis-SNAr product, N2,N4-bis(4-tert-butoxyoxycarbonylmethyleneoxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (**R926576**): ¹H NMR (CDCl₃): δ 7.42 (d, 2H, J= 9 Hz), 7.35 (d, 2H, 9.3 Hz), 7.08 (s, 1H), 6.90 (d, 2H, J= 9.3 Hz), 6.82 (d, 2H, J= 8.7 Hz), 4.53 (s, 2H), 4.49 (s, 2H), 1.50 (s, 9H), 1.49 (s, 9H); LCMS: ret. time: 36.04 min.; purity: 92%; MS (m/e): 591 (MH⁺) and Bis-SNAr product, N4,N6-bis(4-tert-butoxyoxycarbonylmethyleneoxyphenyl)-2,5-dichloro-4,6-pyrimidinediamine (**R926577**): ¹H NMR (CDCl₃): δ 7.43 (d, 4H, J= 8.7 Hz), 6.90 (dd, 4H, J= 9.3 Hz), 4.50 (s, 2H), 1.49 (s, 18H); LCMS: ret. time: 35.31 min.; purity: 100%; MS (m/e): 591 (MH⁺).

7.3.1109 Reaction of 2,4,5,6-tetrachloropyrimidine with 3-hydroxyaniline, N4-(3-Hydroxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (R926590), N2,N4-Bis(3-hydroxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (R926591) and N4,N6-Bis(3-hydroxyphenyl)-2,5-dichloro-4,6-pyrimidinediamine (R926592)

In like manner to the reaction of 2,4,6-trichloropyrimidine with 3,4-ethylenedioxyaniline, the reaction of 2,4,5,6-tetrachloropyrimidine with tert-butyl-4-aminophenoxyacetate gave a mixture of mono-SNAr product, N4-(3-hydroxyphenyl)-2,5,6-trichloro-4-pyrimidineamine (**R926590**): ¹H NMR (CDCl₃): δ 7.38 (bs, 1H), 7.32 (t, 1H, J= 2.4 Hz), 7.22 (s, 1H), 7.01 (dd, 1H, J= 1.2 and 8.1 Hz), 6.68 (dd, 1H, J= 1.8 and 8.1 Hz); LCMS: ret. time: 26.09 min.; purity: 99%; MS (m/e): 292 (MH⁺); Bis-SNAr product, N2,N4-bis(3-hydroxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (**R926591**): ¹H NMR

(CDCl₃): δ 7.45 (s, 1H), 7.30 (t, 1H, J= 2.4 Hz), 7.18 (t, 1H, J= 2.4 Hz), 7.07 (t, 1H, J= 6.6 Hz), 6.98 (t, 1H, J= 8.1 Hz), 6.75 (m, 2H), 6.54 (dd, 1H, J= 2.4 and 8.1 Hz); LCMS: ret. time: 26.54 min.; purity: 87%; MS (m/e): 364 (MH⁺); and Bis-SNAr product, N₄,N₆-bis(3-hydroxyphenyl)-2,5-dichloro-4,6-pyrimidinediamine (**R926592**): ¹H NMR (CDCl₃): δ 7.34 (t, 2H, J= 2.4 Hz), 7.21 (t, 2H, J= 7.5 Hz), 6.98 (m, 4H), 6.60 (m, 2H); LCMS: ret. time: 25.38 min.; purity: 73%; MS (m/e): 364 (MH⁺).

7.3.1110 N₂,N₄-Bis(3-hydroxyphenyl)-5-chloro-6-thiomethyl-2,4-pyrimidinediamine (R926595)

The reaction of N₂ N₄-bis(3-hydroxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine (18 mg, 0.05 mmol) with sodium thiomethoxide (10 mg, 0.15 mmol) in absolute EtOH (1 mL) was heated at 80 °C for 3 days, diluted with H₂O, extracted with EtOAc (3 x 10 mL), and solvent was evaporated to obtain the N₂ N₄-bis(3-hydroxyphenyl)-5-chloro-6-thiomethyl-2,4-pyrimidinediamine (**R926595**). ¹H NMR (CD₃OD): δ 7.40-7.2 (m, 2H), 7.20-6.80 (m, 3H), 6.67 (m, 1H), 6.45-6.30 (m, 2H), 2.4 (s, 3H); LCMS: ret. time: 27.78 min.; purity: 80%; MS (m/e): 376 (MH⁺).

7.3.1111 N₂,N₄-Bis(3,4-ethylenedioxyphenyl)-5-chloro-6-thiomethyl-2,4-pyrimidinediamine (R926475)

In like manner to the preparation of N₂ N₄-bis(3-hydroxyphenyl)-5-chloro-6-thiomethyl-2,4-pyrimidinediamine (**R926595**), the reaction of N₂,N₄-bis(3,4-ethylenedioxyphenyl)-5,6-dichloro-2,4-pyrimidinediamine gave N₂,N₄-bis(3,4-ethylenedioxyphenyl)-5-chloro-6-thiomethyl-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.10 (bd, 2H), 7.00-6.00 (m, 4H), 4.23 (s, 4H), 4.10 (s, 4H), 2.60 (s, 3H); LCMS: ret. time: 36.14 min; purity: 100%; MS (m/e): 459 (MH⁺).

7.3.1112 6-Chloro N₄-(3-hydroxyphenyl)-4-pyrimidineamine (R926530)

The reaction of 4,6-dichloropyrimidine with excess 3-hydroxyaniline in MeOH at 80 °C for 24 h followed by dilution with water and acidification gave the crude product which was purified by silica gel column chromatography to obtain 6-chloro N₄-(3-hydroxyphenyl)-4-pyrimidineamine. ¹H NMR (CD₃OD): δ 8.36 (d, 1H, J= 1.2 Hz), 7.15 (t, 1H, J= 8.4 Hz), 6.93 (dd, 1.2 and 8.1 Hz), 6.74 (d, 1H, J= 1.2 Hz), 6.55 (dd, 1.8 and 8.1 Hz); LCMS (m/e): ret. time: 19.75 min.; purity: 99%; MS (m/e): 222 (MH⁺).

7.3.1113 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine (R925784)

A mixture of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine (20 mg, 0.044 mmol) and phenylboronic acid (6.9 mg, 0.057 mmol) in DME (1 mL) was prepared in a sealed tube and purged with N₂. Tetrakis(triphenylphosphine) palladium(0) (0.002 mmol) was added, and the reaction tube sealed and heated at 80 °C overnight. After cooling, the reaction mixture was diluted with EtOAc, washed with 1N NaOH and brine, dried (MgSO₄), and concentrated. The residue was purified by preparative TLC (40% EtOAc/hexanes) to afford N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.77 (s, 1H), 7.52-7.36 (m, 5H), 7.10 (d, 1H, J= 2.4 Hz), 7.05 (d, 1H, J= 2.4 Hz), 6.93 (dd, 1H, J= 2.4 and 8.7 Hz), 6.87 (dd, 1H, J= 2.4 and 8.7 Hz), 6.73 (d, 1H, J= 8.7 Hz), 6.69 (d, 1H, J= 8.7 Hz), 4.23-4.20 (m, 8H); LCMS: ret. time: 25.38 min.; purity: 100 %; MS (m/e): 455 (MH⁺).

7.3.1114 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-(2-furanyl)-2,4-pyrimidinediamine (R925785)

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine and furan-2-boronic acid were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-(2-furanyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 8.13 (s, 1H), 7.61 (d, 1H, J=1.8 Hz), 7.12 (d, 1H, J= 2.4 Hz), 7.08 (d, 1H, J= 2.4 Hz), 6.93 (td, 2H, J= 2.4 and 8.7 Hz), 6.78 (d, 1H, J= 8.7 Hz), 6.68 (d, 1H, J= 8.7 Hz), 6.58 (d, 1H, J= 2.4 Hz), 6.54 (dd, 1H, J= 1.8 and 3.6), 4.24 (s, 4H), 4.20 (bs, 4H); LCMS: ret. time: 15.03 min.; purity: 88 %; MS (m/e): 445 (MH⁺).

7.3.1115 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-(4-chlorophenyl)-2,4-pyrimidinediamine (R925786)

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine and 4-chlorophenylboronic acid were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-(4-chlorophenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 8.99 (bs, 1H), 8.05 (bs, 1H), 7.85 (s, 1H), 7.50-7.42 (m, 4H), 7.23 (bs, 1H), 7.10 (dd, 1H, J= 2.4 and 8.7 Hz), 7.06 (t, 1H, J= 2.4 Hz), 7.00-6.94 (m, 1H), 6.73 (d, 1H, J= 8.7 Hz), 6.63 (d, 1H, J= 8.7 Hz); LCMS: ret. time: 16.12 min.; purity: 86 %; MS (m/e): 490 (MH⁺).

7.3.1116 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-(3-chlorophenyl)-2,4-pyrimidinediamine (R925787)

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine and 3-chlorophenylboronic acid were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-(3-chlorophenyl)-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 7.77 (s, 1H), 7.45-7.41 (m, 2H), 7.38-7.33 (m, 2H), 7.09 (d, 1H, J= 2.4 Hz), 7.01 (d, 1H, J= 2.4 Hz), 6.92 (dd, 1H, J= 2.4 and 9.0 Hz), 6.86 (dd, 1H, J= 2.4 and 8.7 Hz), 6.74 (d, 1H, J= 8.7 Hz), 6.67 (d, 1H, J= 8.7 Hz), 4.21 (s, 4H), 4.19 (s, 4H); LCMS: ret. time: 27.18 min.; purity: 95 %; MS (m/e): 490 (MH⁺).

7.3.1117 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-(4-methoxycarbonylphenyl)-2,4-pyrimidinediamine (R925813)

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine and (4-methoxycarbonylphenyl)boronic acid were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-(4-methoxycarbonylphenyl)-2,4-pyrimidinediamine. LCMS: ret. time: 26.35 min.; purity: 90 %; MS (m/e): 514 (MH⁺).

7.3.1118 N2,N4-Bis(3,4-ethylenedioxyphenyl)-5-(4-hydroxyphenyl)-2,4-pyrimidinediamine (R925816)

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3,4-ethylenedioxyphenyl)-5-bromo-2,4-pyrimidinediamine and 4-hydroxyphenylboronic acid were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-5-(4-hydroxyphenyl)-2,4-pyrimidinediamine. ¹H NMR (DMSO-d₆): δ 9.53 (s, 1H), 8.92 (s, 1H), 7.78 (s, 1H), 7.74 (bs, 1H), 7.24 (bs, 1H), 7.22 (d, 2H, J= 8.7 Hz), 7.12-7.09 (m, 2H), 6.97 (dt, 1H, J= 2.4 and 8.7 Hz), 6.83 (d, 2H, J= 8.4 Hz), 6.72 (d, 1H, J= 8.1 Hz), 6.62 (d, 1H, J= 9.0 Hz), 4.19 (s, 4H), 4.17 (s, 4H); LCMS: ret. time: 23.51 min.; purity: 95 %; MS (m/e): 471 (MH⁺).

7.3.1119 N2,N4-Bis(3-hydroxyphenyl)-5-phenyl-2,4-pyrimidinediamine (R925783)

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3-hydroxyphenyl)-5-bromo-2,4-pyrimidinediamine and phenylboronic acid were reacted to yield N2,N4-bis(3-

hydroxyphenyl)-5-phenyl-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.85 (bs, 1H), 7.54-7.38 (m, 5H), 7.13-7.11 (m, 2H), 7.10-7.04 (m, 3H), 6.97 (dt, 1H, J = 1.8 and 8.1 Hz), 6.54 (ddd, 1H, J = 1.9, 2.4, and 7.2 Hz), 6.44 (dt, 1H, J = 1.8 and 6.0 Hz); LCMS: ret. time: 20.66 min.; purity: 96 %; MS (m/e): 371 (MH^+).

5 **7.3.1120 N2,N4-Bis(3-hydroxyphenyl)-5-(3,4-methylenedioxyphenyl)-2,4-pyrimidinediamine (R925788)**

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3-hydroxyphenyl)-5-bromo-2,4-pyrimidinediamine and 3,4-methylenedioxyphenylboronic acid were reacted to yield
10 N2,N4-bis(3-hydroxyphenyl)-5-(3,4-methylenedioxyphenyl)-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.82 (s, 1H), 7.13-7.06 (m, 3H), 7.04-7.01 (m, 2H), 6.97 (dt, 1H, J = 1.2 and 8.7 Hz), 6.94-6.88 (m, 3H), 6.52 (ddd, 1H, J = 1.2, 2.4, and 6.9 Hz), 6.42 (dt, 1H, J = 2.1 and 7.5 Hz), 6.01 (s, 2H); LCMS: ret. time: 21.11 min.; purity: 99 %; MS (m/e): 415 (MH^+).

15 **7.3.1121 N2,N4-Bis(3,4-ethylenedioxyphenyl)-6-phenyl-2,4-pyrimidinediamine (R925811)**

In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3,4-ethylenedioxyphenyl)-6-chloro-2,4-pyrimidinediamine and phenylboronic acid were reacted to yield N2,N4-bis(3,4-ethylenedioxyphenyl)-6-phenyl-2,4-pyrimidinediamine. ^1H NMR (CD_3OD): δ 7.97-7.92
20 (m, 2H), 7.46-7.43 (m, 3H), 7.35 (d, 1H, J = 2.7 Hz), 7.19 (d, 1H, J = 2.4 Hz), 7.07-7.00 (m, 2H), 6.75 (t, 2H, J = 8.7 Hz), 6.50 (s, 1H), 4.24-4.19 (m, 8H); LCMS: ret. time: 26.68 min.; purity: 97 %; MS (m/e): 455 (MH^+).

7.3.1122 N2,N4-Bis(3-hydroxyphenyl)-6-phenyl-2,4-pyrimidinediamine (R925812)

25 In a manner similar to the preparation of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-phenyl-2,4-pyrimidinediamine, N2,N4-bis(3-hydroxyphenyl)-6-chloro-2,4-pyrimidinediamine and phenylboronic acid were reacted to yield N2,N4-bis(3-hydroxyphenyl)-6-phenyl-2,4-pyrimidinediamine. LCMS: ret. time: 22.13 min.; purity: 90 %; MS (m/e): 371 (MH^+).

7.3.1123 N2-(3-Aminocarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926747)

The hydrolysis of N2-(3-cyanomethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine gave N2-(3-aminocarbonylmethyleneoxyphenyl)-N4-(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. LCMS: ret. time: 16.76 min.; purity: 93 %; MS (m/e): 412 (MH⁺).

7.3.1124 N2,N4-Bis(3-sodiumphenoxy)-5-fluoro-2,4-pyrimidinediamine (R926461)

The reaction of N2,N4-bis(3-hydroxyphenyl)-5-fluoro-2,4-pyrimidinediamine with 2 equivalents of sodium methoxide in methanol gave the requisite compound, N2,N4-bis(3-sodiumphenoxy)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (D₂O): δ 7.65 (bd, 1H), 7.00-6.90 (m, 2H), 6.71 (m, 2H), 6.55 (dd, 1H, J= 1.2 and 6.3 Hz), 6.31 (bd, 1H, J= 8.1 Hz), 6.23 (bd, 1H, J= 8.7 Hz); ¹⁹F NMR (D₂O): - 47016; LCMS: ret. time: 15.68 min.; purity: 99%; MS (m/e): 313 (MH⁺).

7.3.1125 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-(1,4,5,6-tetrahydro-2-pyrimidyl)methyleneoxyphenyl]-2,4-pyrimidinediamine (R945169)

The reaction of N2-(4-cyanomethyleneoxyphenyl)-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and HCl in ethanol, followed by 1,3-diaminopropane in methanol at 100 °C gave 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-(1,4,5,6-tetrahydro-2-pyrimidyl)methyleneoxyphenyl]-2,4-pyrimidinediamine. ¹H NMR (CD₃OD): δ 2.05 (p, J= 5.7 Hz, 2H), 3.49 (t, J= 5.7 Hz, 4H), 4.84 (s, 2H), 6.56 (ddd, J= 2.1, 3.6 and 5.4 Hz, 1H), 6.93 (d, J= 9.0 Hz, 2H), 7.11-7.13 (m, 2H), 7.21 (m, 1H), 7.55 (d, J= 9.0 Hz, 2H), 7.87 (d, J= 3.9 Hz, 1H); ¹⁹F NMR (282 MHz, CD₃OD): δ - 168.66; LCMS: ret. time: 12.77 min.; purity: 97.61%; MS (m/e): 409.08 (MH⁺).

7.3.1126 5-Fluoro-N4-(3-hydroxyphenyl)-N2-[4-[(4,4-dimethyl-3-oxazolin-2-yl)methyleneoxy]phenyl]-2,4-pyrimidinediamine (R926702)

N2-[4-(cyanomethyleneoxy)phenyl]-5-fluoro-N4-(3-hydroxyphenyl)-2,4-pyrimidinediamine and 2-amino-2-methylpropanol were reacted to yield 5-fluoro-N4-(3-hydroxyphenyl)-N2-[4-[(4,4-dimethyl-3-oxazolin-2-yl)methyleneoxy]phenyl]-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 7.87 (d, 1H, J= 3.6 Hz), 7.37 (t, 1H, J= 2.4 Hz),

7.34 (d, 2H, J= 9.0 Hz), 7.14 (t, 1H, J= 8.1 Hz), 6.94 (bs, 1H), 6.90 (d, 2H, J= 9.0 Hz), 6.78 (dd, 1H, J= 2.4 and 8.4 Hz), 6.74 (d, 1H, J= 3.0 Hz), 6.62 (ddd, 1H, J= 1.2, 2.4, and 8.4 Hz), 4.67 (s, 2H), 4.02 (s, 2H), 1.25 (s, 6H); ^{19}F NMR (CDCl_3): -47399; LCMS: ret. time: 13.82 min.; purity: 98%; MS (m/e): 425 (M+2H).

5 **7.3.1127 N4-(3-Carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950290)**

10 A mixture of equimolar amounts of 2-chloro-N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-4-aminopyridine and 3-ethoxycarbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 97.8%; MS (m/e): 443.20 (MH^+).

15 **7.3.1128 N4-(3-Carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-carboxymethyleneoxyphenyl]-2,4-pyrimidinediamine (R950291)**

20 The reaction of N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (0.1 g) and LiOH (10 equivalents) in MeOH:water (1:1, v/v) for 1h at room temperature followed by treatment with aqueous HCl gave the solid. The resulting solid was filtered, washed with water and dried to give N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-carboxymethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 91.5%; MS (m/e): 415.16 (MH^+).

25 **7.3.1129 N4-(3-Methoxycarbonyl-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonyl methyleneoxyphenyl]-2,4-pyrimidinediamine (R950293)**

30 A solution of N4-(3-carboxy-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeOH was treated with a 4 M solution of HCl in dioxane. The mixture was stirred for 1 hour at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(3-methoxycarbonyl-4-hydroxyphenyl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ^1H NMR (DMSO): δ 10.30 (s, 1H), 10.13 (s, 1H), 8.22 (d, 1H, J= 5.3 Hz), 7.96 (d, 1H, J= 2.4 Hz), 7.71 (dd, J = 2.4, 9.0 Hz, 1H), 6.95-7.11

(m, 4H), 6.51 (m, 1H), 4.56 (s, 2H), 4.09 (q, $J = 7.2$ Hz, 2H), 3.72 (s, 3H), 1.14 (t, $J = 7.2$ Hz, 3H); LCMS: purity: 96.8%; MS (m/e): 457.25 (MH^+).

7.3.1130 N4-(4-Methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950294)

A mixture of equimolar amounts of 2-chloro-N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-4-aminopyridine and 3-ethoxycarbonylmethyleneoxyaniline in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 92.1%; MS (m/e): 469.26 (MH^+).

7.3.1131 N4-(4-Methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950295)

A mixture of equimolar amounts of 2-chloro-N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-4-aminopyridine and 3-ethoxycarbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h followed by aqueous work up gave N4-(4-methoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-methoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 87.6%; MS (m/e): 455.26 (MH^+).

7.3.1132 N4-(4-Ethoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino) carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950296)

A solution of N4-(4-ethoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-ethoxycarbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in EtOH was treated with the HCl salt of methylamine. The mixture was stirred for 4 hours at 100°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-ethoxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 87.4%; MS (m/e): 468.29 (MH^+).

7.3.1133 N4-(4-Carboxyethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950344)

A mixture of equimolar amounts of 2-chloro-N4-(4-carboxyethyleneoxyphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(4-carboxyethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 97.8%; MS (m/e): 456.32 (MH⁺).

7.3.1134 N4-(2,3-Dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950345)

A solution of N4-(4-Methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in TFOH was heated for 2 hours at 100°C. Aqueous work up followed by flash chromatography on silica gel gave N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 92.2%; MS (m/e): 435.95 (MH⁺).

7.3.1135 N4-(4-Methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950346)

A solution of N4-(4-carboxyethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeOH was treated with a 4 M solution of HCl in dioxane. The mixture was stirred for 1 hour at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 85.2%; MS (m/e): 468.01 (MH⁺).

7.3.1136 N4-(4-Hydroxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950347)

The reaction of N4-(4-methoxycarbonylethyleneoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and LiOH (10

equivalents) in MeOH:water (1:1, v/v) for 1h at room temperature followed by treatment with aqueous HCl gave a pale yellow solid. The resulting solid was filtered, washed with water and dried to give N4-(4-hydroxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 94.7%; MS (m/e): 382.03 (MH⁺).

7.3.1137 N4-(2,3-Dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950348)

A mixture N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(2,3-dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 89.5%; MS (m/e): 451.00 (MH⁺).

7.3.1138 N4-(4-Hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950349)

A solution of N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeOH was treated with a sodiumcyanoborohydride. The mixture was stirred for 1 hour at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-Hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.19 (s, 1H), 9.09 (s, 1H), 8.03 (d, 1H, J = 2.4 Hz), 7.28-7.93 (m, 5H), 7.07 (t, 1H, J = 7.2 Hz), 6.71 (d, 1H, J = 7.2 Hz), 6.44 (dd, 1H, J = 2.6, 7.2 Hz), 5.31 (d, 1H, J = 5.1 Hz), 4.14-4.59 (m, 3H), 4.30 (s, 2H), 2.63 (d, 3H, J = 4.8 Hz), 1.82-2.03 (m, 2H); LCMS: purity: 93.3%; MS (m/e): 440.15 (MH⁺).

7.3.1139 N4-(2,3-Dihydro-4-O-methyloxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950356)

A mixture N4-(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and methoxyamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(2,3-

dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid.
LCMS: purity: 85.5%; MS (m/e): 465.10 (MH⁺).

7.3.1140 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950368)

A mixture N4-(4-azido-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and Pd/C (10%) in MeOH was hydrogenated at 22°C for 6 hours (40psi). The mixture was filtered and concentrated to dryness to give N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.60 (s, 1H), 9.46 (s, 1H), 8.73 (bs, 3H), 8.00-8.10 (m, 3H), 7.47 (s, 1H), 7.42 (m, 1H), 7.29 (d, 1H, J = 7.2 Hz), 7.11 (t, 1H, J = 7.2 Hz), 6.82 (d, 1H, J 7.0 Hz), 6.46 (m, 1H), 4.23-4.46 (m, 3H), 4.31 (s, 3H), 2.63 (d, 3H, J = 4.8 Hz), 2.09-2.29 (m, 2H); LCMS: purity: 97.6%; MS (m/e): 438.98 (MH⁺).

7.3.1141 N4-(3-Methylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950371)

A mixture of equimolar amounts of 2-chloro-N4-(3-methylcarbonylphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-methylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 10.16 (s, 1H), 9.82 (s, 1H), 8.24 (d, 1H, J= 2.4 Hz), 8.15 (s, 1H), 7.91-8.07 (m, 2H), 7.70 (d, 1H, J= 7.0 Hz), 7.49 (t, 1H, J = 7.2 Hz), 7.08-7.21 (m, 3H), 6.56 (d, 1H, J = 7.2 Hz), 4.30 (s, 3H), 2.62 (d, 3H, J = 4.8 Hz), 2.48 (s, 3H); LCMS: purity: 93.8%; MS (m/e): 410.50 (MH⁺).

7.3.1142 N4-(3-Phenylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950372)

A mixture of equimolar amounts of 2-chloro-N4-(3-phenylcarbonylphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min

followed by aqueous work up gave N4-(3-phenylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 86.0%; MS (m/e): 472.50 (MH⁺).

7.3.1143 N4-(3-Methylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950373)

A mixture N4-(3-methylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(3-methylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 11.21 (s, 1H), 10.11 (s, 1H), 9.85 (s, 1H), 6.54-8.23 (m, 9H), 4.32 (s, 2H), 2.63 (d, J = 7.0 Hz, 3H), 2.47 (s, 3H); LCMS: purity: 92.4%; MS (m/e): 425.28 (MH⁺).

7.3.1144 N4-(3-Phenylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950374)

A mixture N4-(3-phenylcarbonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N4-(3-phenylcarbonyloximephenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 11.63 (s, 1H), 10.30 (s, 1H), 9.85 (s, 1H), 6.44-8.43 (m, 14H), 4.42 (s, 2H), 2.63 (d, J = 7.0 Hz, 3H); LCMS: purity: 92.4%; MS (m/e): 487.31 (MH⁺).

7.3.1145 N2,N4-Bis(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950376)

A mixture of 2,4-dichloro-5-fluoropyridine and three equivalents of 3-acetophenone in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N2,N4-bis(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 93.1%; MS (m/e): 365.19 (MH⁺).

7.3.1146 N2,N4-Bis(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine (R950377)

A mixture of 2,4-dichloro-5-fluoropyridine and three equivalents of 3-benzophenone in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-
5 20 min followed by aqueous work up gave N2,N4-bis(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 95.7%; MS (m/e): 489.29 (MH⁺).

7.3.1147 N2,N4-Bis(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-2,4-pyrimidinediamine (R950378)

A solution of N2,N4-bis(4-methoxycarbonylethyleneoxyphenyl)-5-fluoro-2,4-
10 pyrimidinediamine in TfOH was heated for 2 hours at 100°C. Aqueous work up followed by flash chromatography on silica gel gave N2,N4-bis(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.36 (s, 1H), 9.14 (s, 1H), 8.06 (d, 1H, J = 2.4 Hz), 7.72-7.99 (m, 3H), 6.97 (d, J = 7.2 Hz, 1H), 6.87 (d, J = 7.2 Hz, 1H), 4.42-4.52 (m, 4H), 2.70-2.78 (m, 4H); LCMS: purity: 94.3%; MS (m/e): 484.50
15 (MH⁺).

7.3.1148 N2,N4-Bis(3-methylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine (R950379)

A mixture of N2,N4-bis(3-methylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous
20 work up gave N2,N4-bis(3-methylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 11.21 (s, 1H), 10.11 (s, 1H), 9.85 (s, 1H), 6.54-8.23 (m, 9H), 4.32 (s, 2H), 2.63 (d, J = 7.0 Hz, 3H), 2.47 (s, 3H); LCMS: purity: 92.1%; MS (m/e): 393.06 (M-H⁺).

7.3.1149 N2,N4-Bis(3-phenylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine (R950380)

A mixture of N2,N4-bis(3-phenylcarbonylphenyl)-5-fluoro-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous
25 work up gave N2,N4-bis(3-phenylcarbonyloximephenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 93.3%; MS (m/e): 486.05 (M-H⁺).

7.3.1150 N2,N4-Bis(2,3-dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-2,4-pyrimidinediamine (R950381)

A mixture of N2,N4-bis(2,3-dihydro-4-benzopyranon-6-yl)-5-fluoro-2,4-pyrimidinediamine and hydroxylamine (20 equivalents) in pyridine at 22°C for 16 hours followed by aqueous work up gave N2,N4-bis(2,3-dihydro-4-oxime-benzopyran-6-yl)-5-fluoro-2,4-pyrimidinediamine as a white solid. LCMS: purity: 98.1%; MS (m/e): 449.03 (M-H⁺).

7.3.1151 N4-(4-Acetyloxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonyl methyleneoxyphenyl]-2,4-pyrimidinediamine (R950382)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in pyridine was treated with acetic anhydride at 22°C for 16 hours. Aqueous work up gave N4-(4-acetyloxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 10.43 (bs, 1H), 9.62 (bs, 1H), 8.03 (d, 1H, J = 2.4 Hz), 7.10-7.83 (m, 7H), 6.83 (d, 1H, J = 7.4 Hz), 6.52 (d, 1H, J = 7.2 Hz), 5.01 (m, 1H), 4.75 (s, 2H), 4.03-4.32 (m, 2H), 2.62 (s, 3H), 2.23 (s, 3H), 1.93-2.13 (m, 2H); LCMS: purity: 92.1%; MS (m/e): 393.06 (M-H⁺).

7.3.1152 N4-(4-Azido-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonyl methyleneoxyphenyl]-2,4-pyrimidinediamine (R950383)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry THF was treated with 2 equivalents of DPPA and DBU. The mixture was stirred for 3 hours at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-azido-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 10.09 (bs, 1H), 9.83 (bs, 1H), 8.18 (d, 1H, J = 2.4 Hz), 7.97 (m, 1H), 7.11-7.61 (m, 6H), 6.82 (d, 1H, J = 7.8 Hz), 6.62 (d, 1H, J = 7.2 Hz), 4.78 (s, 2H), 4.03-4.33 (m, 3H), 2.62 (s, 3H), 1.93-2.13 (m, 2H); LCMS: purity: 97.9%; MS (m/e): 463.07 (MH⁺).

7.3.1153 N4-(4-Benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950385)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in THF was treated with boron trifluoride etherate at 80°C for 8 hours. Aqueous work up gave N4-(4-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.18 (s, 1H), 9.14 (s, 1H), 8.03 (d, J = 2.4 Hz, 1H), 7.93 (bs, 1H), 5.86-7.48 (m, 9H) 4.73-4.74 (m, 2H), 4.33 (s, 2H), 2.62 (s, 3H); LCMS: purity: 96.5%; MS (m/e): 420.07 (M-H⁺).

7.3.1154 N4-(3-Hydroxymethylen-4-methoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950386)

A mixture of equimolar amounts of 2-chloro-N4-(3-hydroxymethylen-4-methoxyphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-hydroxymethylen-4-methoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 89.2%; MS (m/e): 410.5 (MH⁺).

7.3.1155 N4-(3-Amino-4-ethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950388)

A mixture of 2-chloro-N4-(3-amino-4-ethoxyphenyl)-5-fluoro-4-aminopyridine and 3 equivalents of 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-amino-4-ethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 95.1%; MS (m/e): 427.18 (MH⁺).

7.3.1156 N4-(4-Ethoxy-3-hydroxysulfonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950389)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in HOAc was

treated with sodium nitrate followed by addition of concentrated aqueous HCl and copper dichloride. The mixture was stirred for 2 hours at 22°C for 8 hours and purified by aqueous work up followed by column chromatography on silica gel to give N4-(4-ethoxy-3-hydroxysulfonylphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine. LCMS: purity: 82.3%; MS (m/e): 474.09 (M-H⁺).

7.3.1157 N2,N4-Bis(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R950391)

A mixture of 2,4-dichloro-5-fluoropyridine and three equivalents of 3-methoxycarbonyl-4-trifluoromethoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up N2,N4-bis(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine as a white solid. ¹H NMR (DMSO): δ 9.96 (s, 1H), 9.82 (s, 1H), 8.16-8.26 (m, 4H), 7.91 (dd, 1H, J = 3.0, 7.2 Hz), 7.42 (d, 1H, J = 7.2 Hz), 7.31 (d, 1H, J = 7.2 Hz), 3.77 (s, 3H), 3.75 (s, 3H); LCMS: purity: 93.0%; MS (m/e): 565.37 (MH⁺).

7.3.1158 N4-(3-Methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950392)

A mixture of equimolar amounts of 2-chloro-N4-(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-4-aminopyridine and 3-(N-methylamino)carbonylmethyleneoxyaniline in MeOH in a pressure tube at 110°C for 24h or in EtOH using microwave at 175°C for 10-20 min followed by aqueous work up gave N4-(3-methoxycarbonyl-4-trifluoromethoxyphenyl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 95.8%; MS (m/e): 510.41 (MH⁺).

7.3.1159 N4-(4-Acetylamino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine (R950393)

A solution of N4-(4-hydroxy-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in dry MeCN was treated with concentrated sulfuric acid. The mixture was stirred for 3 hours at 22°C, concentrated to dryness and purified by flash chromatography on silica gel to give N4-(4-acetylamino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. ¹H

NMR (DMSO): δ 10.46 (bs, 1H), 9.52 (bs, 1H), 7.98 (d, 1H, $J = 2.4$ Hz), 7.12-7.73 (m, 7H), 6.66 (d, 1H, $J = 7.2$ Hz), 6.49 (d, 1H, $J = 7.2$ Hz), 4.75 (s, 2H), 4.03-4.32 (m, 2H), 3.80 (m, 1H), 2.64 (s, 3H), 2.143 (s, 3H), 1.90-2.11 (m, 2H); LCMS: purity: 92.1%; MS (m/e): 393.06 (M-H⁺). LCMS: purity: 96.2%; MS (m/e): 479.13 (M-H⁺).

5 **7.3.1160 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethylene oxyphenyl]-2,4-pyrimidinediamine HCl salt (R950399)**

10 A solution of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in MeOH was treated with 1 equivalent of 1 N aqueous HCl. The clear solution was concentrated to dryness and the remaining solid was washed with dry acetone to give the HCl salt of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 98.2%; MS (m/e): 438.98 (MH⁺).

15 **7.3.1161 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethylene oxyphenyl]-2,4-pyrimidinediamine succinic acid salt (R950400)**

20 A solution of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in MeOH was treated with 1 equivalent of succinic acid. The clear solution was concentrated to dryness and the remaining solid was crystallized from dry acetone to give the succinic acid salt of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 98.1%; MS (m/e): 438.98 (MH⁺).

25 **7.3.1162 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethylene oxyphenyl]-2,4-pyrimidinediamine maleic acid salt (R950401)**

30 A solution of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in MeOH was treated with 1 equivalent of maleic acid. The clear solution was concentrated to dryness and the remaining solid was crystallized from dry acetone to give the maleic acid salt of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-

methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid.
LCMS: purity: 97.9%; MS (m/e): 438.98 (MH⁺).

7.3.1163 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine fumaric acid salt (R950402)

A solution of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in MeOH was treated with 1 equivalent of fumaric acid. The clear solution was concentrated to dryness and the remaining solid was crystallized from dry acetone to give the fumaric acid salt of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid.
LCMS: purity: 97.9%; MS (m/e): 438.98 (MH⁺).

7.3.1164 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine citric acid salt (R950403)

A solution of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in MeOH was treated with 1 equivalent of citric acid. The clear solution was concentrated to dryness and the remaining solid was crystallized from dry acetone to give the citric acid salt of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid.
LCMS: purity: 97.9%; MS (m/e): 438.98 (MH⁺).

7.3.1165 N4-(4-Amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine HNO₃ salt (R950404)

A solution of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine in MeOH was treated with 1 equivalent of 1 N aqueous HNO₃. The clear solution was concentrated to dryness and the remaining solid was washed with dry acetone to give the nitric acid salt of N4-(4-amino-3,4-dihydro-2H-1-benzopyran-6-yl)-5-fluoro-N2-[3-(N-methylamino)carbonylmethyleneoxyphenyl]-2,4-pyrimidinediamine as a white solid. LCMS: purity: 98.2%; MS (m/e): 438.98 (MH⁺).

7.4 Synthesis of Prodrugs

Exemplary prodrugs according to structural formula (II) were synthesized as described below.

7.4.1 N-2(4)-Acetyl-N2,N4-bis(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R926233)

A mixture of N2,N4-bis(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine, acetyl chloride (4 equivalents), pyridine (4 equivalents) in CH₂Cl₂ was stirred at room temperature for 48h. After an aqueous work up the residue was chromatographed on silica gel to give N-2(4)-acetyl-N2,N4-bis(3,4-ethylenedioxyphenyl)-5-fluoro-2,4-pyrimidinediamine. ¹H NMR (CDCl₃): δ 8.23 (d, 1H, J= 5.4 Hz), 7.03 (d, 1H, J= 2.4 Hz), 7.90-7.80 (m, 3H), 6.76 (m, 2H), 4.28 (bs, 4H), 2.10 (s, 3H); ¹⁹F NMR (CDCl₃): -42125; LCMS: ret. time: 27.94 min.; purity: 99%; MS (m/e): 439 (MH⁺).

7.4.2 N2,N4-Bis(3-N-acetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950244)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour. The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl₃:Acetone, 2:1) to give N2,N4-bis(3-N-acetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine. LCMS: ret. time: 17.03 min.; purity: 87.0%; MS (m/e): 478.89 (MH⁺).

7.4.3 N4-(3-N,N-Diacetylaminophenyl)-N2-(3-N-acetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950245)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine, dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour. The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl₃:Acetone, 2:1) to give N4-(3-N,N-diacetylaminophenyl)-N2-(3-N-acetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine. LCMS: ret. time: 19.27 min.; purity: 92.6%; MS (m/e): 521.01 (MH⁺).

7.4.4 N4-(3-N-Acetylaminophenyl)-N2-(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950246)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine,
 5 dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour. The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl₃:Acetone, 2:1) to give N4-[3-N-acetylaminophenyl]-N2-(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine. LCMS: ret. time: 18.89 min.; purity: 83.0%; MS (m/e): 520.97
 10 (MH⁺).

7.4.5 N2,N4-Bis(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine (R950247)

N2,N4-Bis(3-aminophenyl)-5-fluoro-2,4-pyrimidinediamine,
 dimethylaminopyridine (DMAP) and acetic anhydride were refluxed in pyridine for 1 hour.
 15 The mixture was cooled to room temperature, concentrated, and the residue was subjected to column chromatography on silica gel (CHCl₃:Acetone, 2:1) to give N2,N4-bis(3-N,N-diacetylaminophenyl)-5-fluoro-N2,N4-pyrimidinediacetylamine. LCMS: ret. time: 21.51 min.; purity: 91.8%; MS (m/e): 563.00 (MH⁺).

7.5 The 2,4-Pyrimidinediamine Compounds of the Invention Inhibit FcεRI Receptor-Mediated Degranulation

The ability of the 2,4-pyrimidinediamine compounds of the invention to inhibit IgE-induced degranulation was demonstrated in a variety of cellular assays with cultured human mast cells (CHMC) and/or mouse bone marrow derived cells (BMMC). Inhibition of
 25 degranulation was measured at both low and high cell density by quantifying the release of the granule specific factors tryptase, histamine and hexosaminidase. Inhibition of release and/or synthesis of lipid mediators was assessed by measuring the release of leukotriene LTC₄ and inhibition of release and/or synthesis of cytokines was monitored by quantifying TNF-α, IL-6 and IL-13. Tryptase and hexosaminidase were quantified using fluorogenic
 30 substrates as described in their respective examples. Histamine, TNFα, IL-6, IL-13 and LTC₄ were quantified using the following commercial ELISA kits: histamine (Immunotech #2015, Beckman Coulter), TNFα (Biosource #KHC3011), IL-6 (Biosource #KMC0061),

IL-13 (Biosource #KHC0132) and LTC4 (Cayman Chemical #520211). The protocols of the various assays are provided below.

7.5.1 Culturing of Human Mast and Basophil Cells

Human mast and basophil cells were cultured from CD34-negative progenitor cells as described below (see also the methods described in copending U.S. application Serial No. 10/053,355, filed November 8, 2001, the disclosure of which is incorporated herein by reference).

7.5.1.1 Preparation of STEMPRO-34 Complete Medium

To prepare STEMPRO-34 complete medium ("CM"), 250 mL STEMPRO-34TM serum free medium ("SFM"; GibcoBRL, Catalog No. 10640) was added to a filter flask. To this was added 13 mL STEMPRO-34 Nutrient Supplement ("NS"; GibcoBRL, Catalog No. 10641) (prepared as described in more detail, below). The NS container was rinsed with approximately 10 mL SFM and the rinse added to the filter flask. Following addition of 5 mL L-glutamine (200 mM; Mediatech, Catalog No. MT 25-005-CI and 5 mL 100X penicillin/streptomycin ("pen-strep"; HyClone, Catalog No. SV30010), the volume was brought to 500 mL with SFM and the solution was filtered.

The most variable aspect of preparing the CM is the method by which the NS is thawed and mixed prior to addition to the SFM. The NS should be thawed in a 37° C water bath and swirled, not vortexed or shaken, until it is completely in solution. While swirling, take note whether there are any lipids that are not yet in solution. If lipids are present and the NS is not uniform in appearance, return it to the water bath and repeat the swirling process until it is uniform in appearance. Sometimes this component goes into solution immediately, sometimes after a couple of swirling cycles, and sometimes not at all. If, after a couple of hours, the NS is still not in solution, discard it and thaw a fresh unit. NS that appears non-uniform after thaw should not be used.

7.5.1.2 Expansion of CD34+ Cells

A starting population of CD34-positive (CD34+) cells of relatively small number ($1-5 \times 10^6$ cells) was expanded to a relatively large number of CD34-negative progenitor cells (about $2-4 \times 10^9$ cells) using the culture media and methods described below. The CD34+ cells (from a single donor) were obtained from Allcells (Berkeley, CA). Because there is a degree of variation in the quality and number of CD34+ cells that Allcells

typically provides, the newly delivered cells were transferred to a 15 mL conical tube and brought up to 10 mL in CM prior to use.

On day 0, a cell count was performed on the viable (phase-bright) cells and the cells were spun at 1200 rpm to pellet. The cells were resuspended to a density of 275,000 cells/mL with CM containing 200 ng/mL recombinant human Stem Cell Factor ("SCF"; Peprotech, Catalog No. 300-07) and 20 ng/mL human flt-3 ligand (Peprotech, Catalog No. 300-19) ("CM/SCF/flt-3 medium"). On about day 4 or 5, the density of the culture was checked by performing a cell count and the culture was diluted to a density of 275,000 cells/mL with fresh CM/SCF/flt-3 medium. On about day 7, the culture was transferred to a sterile tube and a cell count was performed. The cells were spun at 1200 rpm and resuspended to a density of 275,000 cells/mL with fresh CM/SCF/flt-3 medium.

This cycle was repeated, starting from day 0, a total of 3-5 times over the expansion period.

When the culture is large and being maintained in multiple flasks and is to be resuspended, the contents of all of the flasks are combined into a single container prior to performing a cell count. This ensures that an accurate cell count is achieved and provides for a degree of uniformity of treatment for the entire population. Each flask is checked separately for contamination under the microscope prior to combining to prevent contamination of the entire population.

Between days 17-24, the culture can begin to go into decline (*i.e.*, approximately 5-10% of the total number of cells die) and fail to expand as rapidly as before. The cells are then monitored on a daily basis during this time, as complete failure of the culture can take place in as little as 24 hours. Once the decline has begun, the cells are counted, spun down at 850 rpm for 15 minutes, and resuspended at a density of 350,000 cells/mL in CM/SCF/flt-3 medium to induce one or two more divisions out of the culture. The cells are monitored daily to avoid failure of the culture.

When greater than 15% cell death is evident in the progenitor cell culture and some debris is present in the culture, the CD34-negative progenitor cells are ready to be differentiated.

7.5.1.3 Differentiation of CD34-Negative Progenitor Cells into Mucosal Mast Cells

A second phase is performed to convert the expanded CD34-negative progenitor cells into differentiated mucosal mast cells. These mucosal cultured human mast

cells ("CHMC") are derived from CD34+ cells isolated from umbilical cord blood and treated to form a proliferated population of CD34-negative progenitor cells, as described above. To produce the CD43-negative progenitor cells, the resuspension cycle for the culture was the same as that described above, except that the culture was seeded at a density of 425,000 cells/mL and 15% additional media was added on about day four or five without performing a cell count. Also, the cytokine composition of the medium was modified such that it contained SCF (200 ng/mL) and recombinant human IL-6 (200 ng/mL; Peprotech, Catalog No. 200-06 reconstituted to 100 ug/mL in sterile 10 mM acetic acid) ("CM/SCF/IL-6 medium").

Phases I and II together span approximately 5 weeks. Some death and debris in the culture is evident during weeks 1-3 and there is a period during weeks 2-5 during which a small percentage of the culture is no longer in suspension, but is instead attached to the surface of the culture vessel.

As during Phase I, when the culture is to be resuspended on day seven of each cycle, the contents of all flasks are combined into a single container prior to performing a cell count to ensure uniformity of the entire population. Each flask is checked separately for contamination under the microscope prior to combining to prevent contamination of the entire population.

When the flasks are combined, approximately 75% of the volume is transferred to the communal container, leaving behind about 10 mL or so in the flask. The flask containing the remaining volume was rapped sharply and laterally to dislodge the attached cells. The rapping was repeated at a right angle to the first rap to completely dislodge the cells.

The flask was leaned at a 45 degree angle for a couple of minutes before the remaining volume was transferred to the counting vessel. The cells were spun at 950 rpm for 15 min prior to seeding at 35-50 mL per flask (at a density of 425,000 cells/mL).

7.5.1.4 Differentiation of CD34-Negative Progenitor Cells into Connective Tissue-Type Mast Cells

A proliferated population of CD34-negative progenitor cells is prepared as above and treated to form a tryptase/chymase positive (connective tissue) phenotype. The methods are performed as described above for mucosal mast cells, but with the substitution of IL-4 for IL-6 in the culture medium. The cells obtained are typical of connective tissue mast cells.

7.5.1.5 Differentiation of CD34-Negative Progenitor Cells into Basophil Cells

A proliferated population of CD34-negative progenitor cells is prepared as described in Section 6.4.1.2, above, and used to form a proliferated population of basophil cells. The CD34-negative cells are treated as described for mucosal mast cells, but with the substitution of IL-3 (at 20-50 ng/mL) for IL-6 in the culture medium.

7.5.2 CHMC Low Cell Density IgE Activation: Tryptase and LTC4 Assays

To duplicate 96-well U-bottom plates (Costar 3799) add 65 ul of compound dilutions or control samples that have been prepared in MT [137 mM NaCl, 2.7 mM KCl, 1.8 mM CaCl₂, 1.0 mM MgCl₂, 5.6 mM Glucose, 20 mM Hepes (pH 7.4), 0.1% Bovine Serum Albumin, (Sigma A4503)] containing 2% MeOH and 1% DMSO. Pellet CHMC cells (980 rpm, 10 min) and resuspend in pre-warmed MT. Add 65 ul of cells to each 96-well plate. Depending on the degranulation activity for each particular CHMC donor, load 1000-1500 cells/well. Mix four times followed by a 1 hr incubation at 37°C. During the 1 hr incubation, prepare 6X anti-IgE solution [rabbit anti-human IgE (1 mg/ml, Bethyl Laboratories A80-109A) diluted 1:167 in MT buffer]. Stimulate cells by adding 25 ul of 6X anti-IgE solution to the appropriate plates. Add 25 ul MT to un-stimulated control wells. Mix twice following addition of the anti-IgE. Incubate at 37°C for 30 minutes. During the 30 minute incubation, dilute the 20 mM tryptase substrate stock solution [(Z-Ala-Lys-Arg-AMC2TFA; Enzyme Systems Products, #AMC-246)] 1:2000 in tryptase assay buffer [0.1 M Hepes (pH 7.5), 10 % w/v Glycerol, 10 uM Heparin (Sigma H-4898) 0.01% NaN₃]. Spin plates at 1000 rpm for 10 min to pellet cells. Transfer 25 ul of supernatant to a 96-well black bottom plate and add 100 ul of freshly diluted tryptase substrate solution to each well. Incubate plates at room temperature for 30 min. Read the optical density of the plates at 355nm/460nm on a spectrophotometric plate reader.

Leukotriene C4 (LTC4) is also quantified using an ELISA kit on appropriately diluted supernatant samples (determined empirically for each donor cell population so that the sample measurement falls within the standard curve) following the supplier's instructions.

7.5.3 CHMC High Cell Density IgE Activation: Degranulation (Tryptase, Histamine), Leukotriene (LTC₄), and Cytokine (TNF α , IL-13) Assays

Cultured human mast cells (CHMC) are sensitized for 5 days with IL-4 (20 ng/ml), SCF (200 ng/ml), IL-6 (200 ng/ml), and Human IgE (CP 1035K from Cortx Biochem, 100-500ng/ml depending on generation) in CM medium. After sensitizing, cells are counted, pelleted (1000 rpm, 5-10 minutes), and resuspended at $1-2 \times 10^6$ cells/ml in MT buffer. Add 100 μ l of cell suspension to each well and 100 μ l of compound dilutions. The final vehicle concentration is 0.5% DMSO. Incubate at 37°C (5% CO₂) for 1 hour. After 1 hour of compound treatment, stimulate cells with 6X anti-IgE. Mix wells with the cells and allow plates to incubate at 37°C (5% CO₂) for one hour. After 1 hour incubation, pellet cells (10 minutes, 1000 RPM) and collect 200 μ l per well of the supernatant, being careful not to disturb pellet. Place the supernatant plate on ice. During the 7-hour step (see next) perform tryptase assay on supernatant that had been diluted 1:500. Resuspend cell pellet in 240 μ l of CM media containing 0.5% DMSO and corresponding concentration of compound. Incubate CHMC cells for 7 hours at 37°C (5% CO₂). After incubation, pellet cells (1000 RPM, 10 minutes) and collect 225 μ l per well and place in -80°C until ready to perform ELISAS. ELISAS are performed on appropriately diluted samples (determined empirically for each donor cell population so that the sample measurement falls within the standard curve) following the supplier's instructions.

7.5.4 BMMC High Cell Density IgE Activation: Degranulation (Hexosaminidase, Histamine), Leukotriene (LTC₄), and Cytokine (TNF α , IL-6) Assays

7.5.4.1 Preparation of WEHI-Conditioned Medium

WEHI-conditioned medium was obtained by growing murine myelomonocytic WEHI-3B cells (American Type Culture Collection, Rockville, MD) in Iscove's Modified Eagles Media (Mediatech, Herndon, VA) supplemented with 10% heat-inactivated fetal bovine serum (FBS; JRH Biosciences, Kansas City, MO), 50 μ M 2-mercaptoethanol (Sigma, St. Louis, MO) and 100 IU/mL penicillin-streptomycin (Mediatech) in a humidified 37°C, 5% CO₂/95% air incubator. An initial cell suspension was seeded at 200,000 cells/mL and then split 1:4 every 3-4 days over a period of two weeks. Cell-free supernatants were harvested, aliquoted and stored at -80°C until needed.

7.5.4.2 Preparation of BMMC Medium

BMMC media consists of 20% WEHI-conditioned media, 10% heat-inactivated FBS (JHR Biosciences), 25 mM HEPES, pH7.4 (Sigma), 2mM L-glutamine (Mediatech), 0.1 mM non-essential amino acids (Mediatech), 1mM sodium pyruvate (Mediatech), 50 μ M 2-mercaptoethanol (Sigma) and 100 IU/mL penicillin-streptomycin (Mediatech) in RPMI 1640 media (Mediatech). To prepare the BMMC Media, all components are added to a sterile IL filter unit and filtered through a 0.2 μ m filter prior to use.

7.5.4.3 Protocol

Bone marrow derived mast cells (BMMC) are sensitized overnight with murine SCF (20 ng/ml) and monoclonal anti-DNP (10 ng/ml, Clone SPE-7, Sigma # D-8406) in BMMC media at a cell density of 666×10^3 cells/ml. After sensitizing, cells are counted, pelleted (1000 rpm, 5-10 minutes), and resuspended at $1-3 \times 10^6$ cells/ml in MT buffer. Add 100 μ l of cell suspension to each well and 100 μ l of compound dilutions. The final vehicle concentration is 0.5% DMSO. Incubate at 37°C (5% CO₂) for 1 hour. After 1 hour of compound treatment, stimulate cells with 6X stimulus (60 ng/ml DNP-BSA). Mix wells with the cells and allow plates to incubate at 37°C (5% CO₂) for one hour. After 1 hour incubation, pellet cells (10 minutes, 1000 RPM) and collect 200 μ l per well of the supernatant, being careful not to disturb pellet, and transfer to a clean tube or 96-well plate. Place the supernatant plate on ice. During the 4-5 hour step (see next) perform the hexosaminidase assay. Resuspend cell pellet in 240 μ l WEI-conditioned media containing 0.5% DMSO and corresponding concentration of compound. Incubate BMMC cells for 4-5 hours at 37°C (5% CO₂). After incubation, pellet cells (1000 RPM, 10 minutes) and collect 225 μ l per well and place in -80°C until ready to perform ELISAS. ELISAS are performed on appropriately diluted samples (determined empirically for each donor cell population so that the sample measurement falls within the standard curve) following the supplier's instructions.

Hexosaminidase assay: In a solid black 96-well assay plate, add 50 μ L hexosaminidase substrate (4-methylumbelliferyl-N-acetyl- β -D-glucosaminide; 2mM) to each well. Add 50 μ L of BMMC cell supernatant (see above) to the hexoseaminidase substrate, place at 37°C for 30 minutes and read the plate at 5, 10, 15, and 30 minutes on a spectrophotometer.

7.5.5 Basophil IgE or Dustmite Activation: Histamine Release Assay

The basophil activation assay was carried out using whole human peripheral blood from donors allergic to dust mites with the majority of the red blood cells removed by dextran sedimentation. Human peripheral blood was mixed 1:1 with 3% dextran T500 and RBCs were allowed to settle for 20-25min. The upper fraction was diluted with 3 volumes of D-PBS and cells were spun down for 10 min at 1500 rpm, RT. Supernatant was aspirated and cells were washed in an equal volume MT-buffer. Finally, cells were resuspended in MT-buffer containing 0.5% DMSO in the original blood volume. 80 uL cells were mixed with 20 uL compound in the presence of 0.5% DMSO, in triplicate, in a V-bottom 96-well tissue culture plate. A dose range of 8 compound concentrations was tested resulting in a 10-point dose response curve including maximum (stimulated) and minimum (unstimulated) response. Cells were incubated with compound for 1 hour at 37°C, 5% CO₂ after which 20 uL of 6x stimulus [1 ug/mL anti-IgE (Bethyl Laboratories) 667 au/mL house dustmite (Antigen Laboratories)] was added. The cells were stimulated for 30 minutes at 37°C, 5% CO₂. The plate was spun for 10 min at 1500 rpm at room temperature and 80 uL the supernatant was harvested for histamine content analysis using the histamine ELISA kit supplied by Immunotech. The ELISA was performed according to supplier's instructions.

7.5.6 Results

The results of low density CHMC assays (Section 6.4.3), the high density BMHC assays (Section 6.4.5) and the basophil assays (Section 6.4.6) are provided in TABLE 1. The results of the high density CHMC assays (Section 6.4.4) are provided in TABLE 2. In TABLES 1 and 2, all reported values are IC₅₀s (in μM). A value of "9999" indicates an IC₅₀ > 10 μM, with no measurable activity at a 10 μM concentration. Most compounds tested had IC₅₀s of less than 10 μM, with many exhibiting IC₅₀s in the sub-micromolar range.

7.6 The 2,4-Pyrimidinediamine Compounds Inhibit FcγRI Receptor-Mediated Degranulation

The ability of the 2,4-pyrimidinediamine compounds of the invention to inhibit FcγRI-mediated degranulation was demonstrated with Compounds R921218, R921302, R921303, R940347, R920410, R927050, R940350, R935372, R920323, R926971 and

R940352 in assays similar to those described in Section 6.4, with the exception that the cells were not primed with IgE and were activated with rabbit anti-human IgG Fab fragment (Bethyl Laboratories, Catalog No. A80-105).

All of the compounds tested exhibited IC_{50} s in the sub micromolar range.

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Histamine	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						CHMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R008951														
R008952														
R008953														
R008955														
R008956														
R008958														
R067934														
R067963														
R070153														
R070790	1.665	9999												
R070791														
R081166														
R088814														
R088815														
R091880														
R092788														
R908696	3.553													
R908697	9999	9999												
R909236	0.996	9999												
R909237	9999	9999												
R909238	0.174	9999							<0.22		<0.22	0.521	0.432	<0.22
R909239	0.264	9999												
R909240	0.262	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE LTC4	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R909241	0.181	9999							<0.22		<0.22	1.021	0.253	<0.22
R909242	0.567	9999												
R909243	0.263	>10												
R909245	0.255	6.242												
R909246	0.169	9999												
R909247	2.393	9999												
R909248	3.582	9999												
R909249	9999	9999												
R909250	8.025	9999												
R909251	0.138	9999												
R909252	0.248	9999												
R909253	7.955	9999												
R909254	0.136	9999												
R920664	9999	9999												
R920665	1.1	9999												
R920666	2.53	9999												
R920668	3.2	9999												
R920669	0.42	9999												
R920670	2.18	9999												
R920671	9999	9999												
R920672	9999	9999												
R920818	9999	9999												
R920819	10	9999												

TABLE 1

TABLE 1																
Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density						
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	BMMC anti-IgE hexos						BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6		
R920820	9999	9999														
R920846	9999	9999														
R920860	1.009	9999														
R920861	0.598	>10														
R920893	1.239	9999														
R920894	0.888	5.566														
R920910	0.751	7.922														
R920917	1.579	9.729														
R921218	0.499	9999	0.55	0.6	9999	0.24	9999	0.302	0.133	9999	0.203	0.766	0.274	0.100		
R921219	0.059	9999				0.025	9999	0.020	0.069		0.058	0.040	0.039	0.009		
R925734				9.2	>10				9999	9999						
R925747	1.021	3.1							3.1							
R925755	0.898	9999														
R925757	2.8	9999														
R925758	1.175	9999														
R925760	4.85	9999														
R925765	6.8	9999														
R925766	8.9	9999														
R925767	10															
R925768	9999															
R925769	9999															
R925770	9999															
R925771	0.5	2.8	0.22													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R925772	9999	9999												
R925773	0.673	9999												
R925774	0.435	9999												
R925775	0.225	9999	0.2											
R925776	2.1	9999												
R925778	0.225	9999	0.18											
R925779	0.265	9999	0.19											
R925783	2.9	9999												
R925784	3.2	9999												
R925785	2.5	9999												
R925786	1.85	9999												
R925787	9	9999												
R925788	2.4	9999												
R925790	9999	9999												
R925791	9999	9999												
R925792	6.25	9999												
R925794	9999	9999												
R925795	9999	9999												
R925796	2	9999												
R925797	0.85	9999	0.28											
R925798	9999	9999												
R925799	9999	9999												
R925800	9999	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R925801	9999	9999												
R925802	9999	9999												
R925803	9999	9999												
R925804	9999	9999												
R925805	9999	9999												
R925806	9999	9999												
R925807	9999	9999												
R925808	9999	9999												
R925810	9999	9999												
R925811	3.3	9999												
R925812	5.8	9999												
R925813	9999	9999												
R925814	9999	9999												
R925815	9999	9999												
R925816	6	9999												
R925819	9999	9999												
R925820	9999	9999												
R925821	9999	9999												
R925822	9999	9999												
R925823	9999	9999												
R925824	9999	9999												
R925837	9999	9999												
R925838	9999	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R925839	9999	9999												
R925840	9999	9999												
R925841	9999	9999												
R925842	7.3	9999												
R925843	9999	9999												
R925844	5.1	9999												
R925845	2.3	9999												
R925846	9999	9999												
R925849	8.2	9999												
R925851	0.925	9999												
R925852	3	9999												
R925853	9999	9999												
R925854	9999	9999												
R925855	4.2	9999												
R925856	9.85	9999												
R925857	5.95	9999												
R925858	8.05	7.3												
R925859	9999	9999												
R925860	9999	9999												
R925861	9999	9999												
R925862	0.7	9999												
R925863	0.274	9999												
R925864	9999	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R925865	9999	9999												
R926016						9999	9999		9999	9999				
R926017				1.43	9999	0.53	9999		1.4	9.6				
R926018						9999	10		8.5	9999				
R926037						9999	9999		9999	9999				
R926038						9999	9999		9999	9999				
R926039						9999	9999		9999	9999				
R926058						9999	9999		9999	9999				
R926064				6.2					5.9	7.3				
R926065				3.5					9999	9999				
R926068				>10					7.4	8.2				
R926069				9.1					4.5	4.4				
R926072				>10					9999	9999				
R926086						2.5	9999		2.8	7.3				
R926108			0.76	0.787	6.4	0.95	9999		0.9	9999				
R926109	0.538	5.5	0.73	0.55	>10	0.15	9999		0.6	3.2				
R926110	1.071	9999	1.42	1.2	>10	0.3	9999		1	4.5				
R926113	0.413		0.49	0.413	9999	0.27	9999		0.65	9999				
R926114				3.427	8.1	1.7	10		9999	9999				
R926145				4.764	>10				2.4	8.8				
R926146			1.59	0.761	6.7				1.35	5				
R926147				1.899	>10				2	7.1				
R926206						>10	>10		6.6	8.6				

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926209						>10	9999		10	9.1				
R926210	0.926	9999	0.8	700	9999	0.37	>10		0.6	>10				
R926211	1.299	9.8		2.7	9999	1.55	>10		3.9	>10				
R926212	0.654	9999	0.45			0.5	>10		0.5	5				
R926213	1.639	5.5				1.75	>10							
R926218				>10					9999	9999				
R926219				1.102	6.7				2.5	3.2				
R926220				>10					9999	9999				
R926221				8.5					9.9	9999				
R926222				>10					9999	9999				
R926223				>10					9999	9999				
R926224				>10					9999	9999				
R926225				>10					9999	9999				
R926228				>10					9999					
R926229				>10										
R926230				>10										
R926234				>10					9999					
R926237	1.207	6.2							1.9					
R926240	0.381	1.7	0.145											
R926241	7	9999												
R926242	4.2	9999												
R926243	3.1	9999												
R926245	3.1	9.4												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926248	0.9	9999	0.76											
R926249	0.5	9999	0.25											
R926252	2.8													
R926253	0.8		0.675											
R926254	1.3	4												
R926255	1.4	4.5												
R926256	0.275	5.1	0.23											
R926257	1.5	7.5												
R926258	0.9	9999	0.59											
R926259	2.5	6.2												
R926319	9999	9999												
R926320	9999	9999												
R926321	9999	9999												
R926325	9999	9999												
R926331	9999	9999												
R926339	0.66	9999												
R926340	3.23	9999												
R926341	0.875	9999												
R926342	10	9999												
R926376	9999													
R926386	9999	9999												
R926387	0.65	9999	0.7											
R926394	9999	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	CHMC Ionomycin Histamine	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4							BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926395	0.875	6.4	0.29												
R926396	0.7	2.6	0.16												
R926397	9999	9999													
R926398	9999	9999													
R926399	9999	9999													
R926400	9999	9999													
R926401	9999	9999													
R926402	9999	9999													
R926403	9999	9999													
R926404	9999	9999													
R926405	3.4	9999													
R926406	9999	9999													
R926408	9.6	9999													
R926409	3.15	9999													
R926411	0.69	2.5													
R926412	0.62	9999													
R926461	0.725	9999													
R926467	1.175	8.8													
R926469	9999														
R926474	2.5	9999													
R926475	2.15	>10													
R926476	0.6	7.7													
R926477	0.27	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926478	9999													
R926479	9999													
R926480	1.9	9999												
R926481	1.445	9999												
R926482	1.037	>10												
R926483	9999													
R926484	1.523	9999												
R926485	4.012	9999												
R926486	0.647	7.403												
R926487	0.554	8.867	1.25											
R926488	0.331	>10	0.752											
R926489	1.414	>10												
R926490	1.571	9999												
R926491	1.158	>10												
R926492	0.645	9999												
R926493	0.25	9.181	0.078											
R926494	0.313	9999	0.078											
R926495	0.121	>10	0.078			0.04	9999	0.038	0.056		0.089	0.24	0.077	0.028
R926496	0.571	>10												
R926497	0.138	9999				0.27	9999	0.205						
R926498	0.209	>10							<0.22		0.515	0.995	0.614	<0.22
R926499	0.29	>10												
R926500	0.418	>10												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926501	0.298	>10				0.609	9999	0.645						
R926502	0.483	>10				0.405	9999	0.491						
R926503	0.452	>10												
R926504	0.569	>10												
R926505	0.145	9999							<0.22		<0.22	<0.22	<0.22	<0.22
R926506	0.343	9999												
R926508	0.127	9999				0.065	9999	0.054	0.086		0.107	0.162	0.054	0.026
R926509	1.16	9999												
R926510	0.44	>10												
R926511	0.786	>10												
R926514	9999	9999												
R926516	1	9999												
R926526	9999	9999												
R926527	9999	9999												
R926528	8.75	9999												
R926535	9999	9999												
R926536	9999	9999												
R926555	9999	9999												
R926559	7.7	9999												
R926560	9999	9999												
R926562	9999	9999												
R926563	9999	9999												
R926564	3.75	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926565	0.625	3.3												
R926566	2.73	9999												
R926567	9.3	9999												
R926569	0.61	3.07												
R926571	9999	9999												
R926572	1.8	6.08												
R926574	1.96	2.63												
R926576	9999	9999												
R926579	9999	9999												
R926580	10	9999												
R926582	1.3	9999												
R926583	9999	9999												
R926584	9999	9999												
R926585	9999	9999												
R926586	2.75	9999												
R926587	9999	9999												
R926588	7.85	9999												
R926589	0.325	10												
R926591	2.62	9999												
R926593	0.68	8.3	0.495											
R926594	9999	9999												
R926595	4.85	9999												
R926604	2.85	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926605	2.45	9999												
R926614	0.228	9999												
R926615	0.445	9999												
R926616	0.625	3.25												
R926617	9.45	9999												
R926620	8.35	9999												
R926623	9999	9999												
R926662	9999	9999												
R926663	9999	9999												
R926675	0.63	9999												
R926676	0.76	9999												
R926680	1.71	9999												
R926681	0.775	9999												
R926682	8.41	9999												
R926683	10	9999												
R926688	2.25	>10												
R926690	0.146	>10												
R926696	0.309	>10												
R926698	9999													
R926699	0.76	9999												
R926700	0.157	>10												
R926701	2.2	9999												
R926702	0.886	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4							BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926703	0.525	9999													
R926704	0.564	9999													
R926705	0.263	9999	0.533												
R926706	0.07	2.406	0.078												
R926707	0.214	9999								<0.056		<0.056	0.39	0.088	<0.056
R926708	0.472	9999													
R926709	0.858	9999													
R926710	1.763	9999													
R926711	1.245	9999													
R926712	1.084	9999													
R926713	0.446	8.741													
R926714	0.428	>10													
R926715	0.588	>10													
R926716	1.06	9999													
R926717	7.874	9999													
R926718	1.826	9999													
R926719	0.1335	4.024													
R926720	1.555	9999													
R926721	4.441	9999													
R926722	5.96	9999													
R926723	2.591	9999													
R926724	2.059	9999													
R926725	0.431	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926726	9999	9999												
R926727	0.387	9999												
R926728	0.482	>10												
R926730	0.251	9999												
R926731	9999	9999												
R926732	0.444	9999												
R926733	1.496	9999												
R926734	4.493	9999												
R926735	3.712	9999												
R926736	0.288	9999												
R926737	0.059	9999							0.075		0.073	0.046	0.068	0.017
R926738	0.342	9999												
R926739	0.508	9999												
R926740	4.422	9999												
R926741	2.908	9999							0.961		1.025	9999	0.772	0.537
R926742	0.127					0.043	9999	0.055	0.041		0.055	0.105	0.053	0.022
R926743	9999													
R926744	9999													
R926745	0.083	9999												
R926746	0.989	9999												
R926747	0.213	>10												
R926748	0.345	>10												
R926749	0.472	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926750	0.361	>10												
R926751	0.598	9999												
R926764	0.252	5.64												
R926765	0.324	4.39												
R926766	0.756	9999												
R926767	0.387	>10												
R926768	0.443	>10												
R926769	1.067	9999												
R926770	0.583	9999												
R926771	2.049	9999												
R926772	0.337	7.501												
R926773	0.548	7.849												
R926774	1.934	7.935												
R926775	3.47	>10												
R926776	0.81	9999												
R926777	0.378	9999												
R926778	0.414	9999												
R926779	9999	9999												
R926780	0.152	>10							<0.22		<0.22	0.461	<0.22	<0.22
R926781	0.573	9999												
R926782	0.173	>10							<0.22		<0.22	1.461	0.276	<0.22
R926783	0.304	>10												
R926784	0.252	9999												

TABLE 1

Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	CHMC anti-IgE Hexos.						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926785	0.222	>10								0.989		0.561	1.411	1.312	0.513
R926786	0.504	9999													
R926787	5.422	9999													
R926788	0.336	6.341													
R926789	2.315	9999													
R926790	0.462	7.412													
R926791	0.233	>10								0.064		<0.056	0.896	0.205	<0.056
R926792	3.197	9999													
R926793	3.073	9999													
R926795	2.041	>10													
R926796	0.914	9999													
R926797	2.235	9999													
R926798	2.347	5.87													
R926799	9999	9999													
R926800	4.581	9999													
R926801	10	9999													
R926802	1.251	>10													
R926803	1.541	>10													
R926804	1.578	7.109													
R926805	0.764	9999													
R926806	0.374	9999													
R926807	0.291	9999													
R926808	0.368	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE LTC4	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926809	0.78	3.052												
R926810	1.221	9999												
R926811	3.662	9999												
R926812	0.185	>10												
R926813	0.152	9999												
R926814	1.101	9999												
R926815	1.181	9999												
R926816	0.084	9999												
R935000	9999	9999												
R935001	9999	9999												
R935002	9999	9999												
R935003	9999	9999												
R935004	9999	9999												
R935005	9999	9999												
R935006	10	9.8												
R935016	9999	9999												
R935019	8.8	9999												
R935020	9999	9999												
R935021	9999	9999												
R935023	9999	9999												
R935025	1.04	9999												
R935029	2.83	9999												
R935075	0.93	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R935076	4.15	9999												
R935077	9999	9999												
R935114	1.725	9999												
R935117	9999													
R935134	0.909	1.799												
R935135	10	9999												
R935136	0.952	2.129												
R935137	10	9999												
R935138	0.096	0.552							<0.22		<0.22	0.373	0.409	<0.22
R935139	0.846	9999												
R935140	0.275	0.959												
R935141	0.727	>10												
R935142	0.873	>10												
R935143	0.573	>10												
R935144	0.63	9999												
R935145	0.548	>10												
R935146	3.802	9999												
R935147	1.404	9999												
R935148	2.218	9.423												
R935149	0.708	>10												
R935150	1.926	9.738												
R935151	0.479	>10												
R935152	0.505	9.316												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R935153	0.238	>10												
R935154	0.127	>10							0.104		0.085	0.547	0.131	0.041
R935155	0.401	9999												
R935156	0.149	>10							<0.22		<0.22	0.433	0.22	<0.22
R935157	0.256	4.656												
R935158	0.551	>10												
R935159	0.232	4.135												
R935160	0.202	>10							<0.22		0.317	0.876	0.484	<0.22
R935161	0.277	9999												
R935162	0.269	>10												
R935163	9999	9999												
R935164	0.204	9999												
R935165	4.988	9999												
R935166	0.568	9999												
R935167	2.132	>10												
R935168	0.488	9.484												
R935169	0.999	8.007												
R935170	0.673	9999												
R935171	0.536	9999												
R935172	1.385	6.808												
R935173	0.454	>10												
R935174	1.384	9999												
R935175	0.885	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4							BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R935176	1.169	9999													
R935177	0.889	>10													
R935178	0.515	9999													
R935179	0.557	9999													
R935180	1.22	9999													
R935181	1.76	9999													
R935182	0.124	2.469													
R935183	0.729	9999													
R935184	0.605	9999													
R935185	0.351	6.642													
R935186	0.211	9999													
R935187	9.059	>10													
R935188	0.239	9999													
R935189	0.619	9999													
R935190	0.156	9999													
R935191	0.151	9999								0.068		0.043	0.213	0.071	0.027
R935192	0.337	9999													
R935193	0.136	9999								0.08		0.048	0.312	0.092	0.037
R935194	0.11	9999								0.125		0.054	0.493	0.118	0.034
R935196	0.117	9999													
R935197	0.174	>10													
R935198	0.126	>10													
R935199	0.45	>10													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Histamine	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4							BMHC anti-IgE hexos	BMHC Ionomycin Hexos.	BMHC anti-IgE histamine	BMHC anti-IgE LTC4	BMHC anti-IgE TNF-alpha	BMHC anti-IgE IL-6
R935202	0.181	9.765													
R935203	0.562	>10													
R935204	0.554	9999													
R935205	2.959	9999													
R935206	4.711	9999													
R935207	9999	9999													
R935208	1.274	9999													
R935209	0.526	1.035													
R935211	1.238	9999													
R935212	1.427	9999													
R935213	0.619	10													
R935214	0.453	5.499													
R935218	4.712	9999													
R935219	5.409	9999													
R935220	3.789	9999													
R940089	9999	9999													
R940090	9999	9999													
R940095	9999	9999													
R940100	9999	9999													
R940215	0.845	9999													
R940216	0.2675	7.3													
R940217	9999	9999													
R940222	9999	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R940233	0.132	>10												
R940235	0.8	>10												
R940250														
R940251														
R940253	1.006	>10												
R940254	0.986	9999												
R940255	1.033	9999												
R940256	1.104	9999												
R940257	0.667	9999												
R940258	0.473	5.72												
R940260	1.126	9999												
R940261	9999	9999												
R940262	9999	9999												
R940263	9999	9999												
R940264	10	9999												
R940265	0.239	>10							0.981		0.306	1.211	1.131	0.486
R940266	9999	9999												
R940267	3.151	9999												
R940269	1.654	9999												
R940270	2.144	8.739												
R940271	0.401	6.821												
R940275	0.862	9999												
R940276	0.211	9999							0.136		0.073	0.332	0.251	<0.056

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Histamine	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMHC anti-IgE hexos	BMHC Ionomycin Hexos.	BMHC anti-IgE histamine	BMHC anti-IgE LTC4	BMHC anti-IgE TNF-alpha	BMHC anti-IgE IL-6
R940277	0.141	9999							0.279		0.315	0.625	0.262	0.181
R940280	6.999	9999												
R940281	0.525	5.529												
R940282	0.401	3.015												
R940283	0.553	4.982												
R940284	0.465	3.744												
R940285	3.499	9999												
R940286	0.337	7.082												
R940287	0.288	7.684												
R940288	0.208	9999												
R940289	0.272	9999												
R940290	0.116	9999							0.255		0.545	0.59	0.246	0.1
R940291	0.396	9999												
R940292	0.683	9999												
R940293	9999	9999												
R940294	1.366	9999												
R940295	0.126	8.812												
R940296	0.41	>10												
R940297	3.465	10												
R945025	9999	9999												
R945032	0.37	9999												
R945033	9999	9999												
R945034	1.85	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density						
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6	
R945035	9999	9999													
R945036	9999	9999													
R945037	9999	9999													
R945038	9999	9999													
R945040	9999	9999													
R945041	9999	9999													
R945042	9999	9999													
R945043	9999	9999													
R945045	9999	9999													
R945046	0.82	>10													
R945047	0.845	9999													
R945048	0.76	9999													
R945051	0.95	>10													
R945052	0.425	2.48													
R945053	0.1185	1.48													
R945056	10	9999													
R945057	10	9999													
R945060	0.9375	>10													
R945061	10	9999													
R945062	0.625	>10													
R945063	1.55	>10													
R945064	0.53	>10													
R945065	1.425	>10													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	CHMC ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC ionomycin Tryptase	CHMC anti-IgE LTC4							BMMC anti-IgE hexos	BMMC ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R945066	5.2	nd													
R945067	9999	nd													
R945068	9999	nd													
R945070	0.45	>10													
R945071	0.205	>10													
R945096	1.75	>10													
R945097	10	9999													
R945109	1.025	>10													
R945110	0.602	9999													
R945117	4.077	9999													
R945118	0.668	9999													
R945124	0.69	7.852													
R945125	0.896	>10													
R945126	9999	9999													
R945127	0.704	8.955													
R945128	0.685	8.8													
R945129	1.003	>10													
R945130	1.874	9999													
R945131	0.77	9999													
R945132	0.571	8.77													
R945133	1.064	>10													
R945134	9999	9999													
R945135	0.986	8.245													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density						
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMHC anti-IgE hexos	BMHC Ionomycin Hexos.	BMHC anti-IgE histamine	BMHC anti-IgE LTC4	BMHC anti-IgE TNF-alpha	BMHC anti-IgE IL-6	
R945137	1.649	>10													
R945138	1.058	6.733													
R945139	1.016	>10													
R945140	0.573	>10													
R945142	1.049	>10													
R945144	0.244	9999													
R945145	9999	>10													
R945146	3.756	9999													
R945147	3.546	9999													
R945148	0.307	9999													
R945149	0.391	>10													
R945150	0.467	>10								>2	>2	9999	0.709	0.634	
R945151	4.07	9999													
R945152	6.94	9999													
R945153	0.688	6.561													
R945155	1.878	>10													
R945156	0.787	9999													
R945157	1.477	9999													
R945162	9999	9999													
R945163	0.922	4.251													
R945164	10	9999													
R945165	9999	9999													
R945166	9999	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Histamine	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4							BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R945167	0.761	9999													
R945168	10	9999													
R945169	10	9999													
R945170	0.661	>10													
R945171	1.327	9999													
R945172	1.179	9999													
R945173	1.419	9999													
R945175	1.648	9999													
R950082	9999	9999													
R950083	9999	9999													
R950090	9999	9999													
R921302	0.37	9999			0.19		9999	0.282							
R950092	9999	9999													
R950093	0.64	5.55													
R950100	0.71	>10													
R950107	0.46	>10													
R950108	2.075	>10													
R950109	7.95														
R950120	3	9999													
R950121	4.25	>10													
R950122	3.025	9999													
R950123	3.25	8.45													
R950125	1.375	6.3													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE LTC4	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R950129	0.665	>10												
R950130	4.9													
R950131	9999													
R950132	9													
R950133	2.2	>10												
R950134	1.875	9999												
R950135	0.85	>10												
R950137	2.23	9999												
R950138	9.5													
R950139	1.375	9999												
R950140	2.825	9999												
R950141	0.31	>10												
R950142	10													
R950143	8.23													
R950144	10													
R950145	9999													
R950146	9999													
R950147	9999													
R950148	2.275	9999												
R950149	10	9999												
R950150	9999	9999												
R950151	9999													
R950152	10													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Histamine	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R950153	9999													
R950154	2.075	9999												
R950155	9999													
R950156	9999													
R950157	9999													
R950158	9.98													
R950159	0.61	9999												
R950160	1	9999												
R950162	0.434	>10												
R950163	0.874	9999												
R950164	1.893	9999												
R950165	1.288	9999												
R950166	1.889	9999												
R950167	9999	9999												
R950168	6.496	8.653												
R950169	1.273	9.518												
R950170	9999	9999												
R950171	0.585	>10												
R950172	0.983	9999												
R950173	2.368	>10												
R950174	4.618	9999												
R950175	1.688	9999												
R950176	1.342	9999												

TABLE 1

TABLE 1															
Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	BMMC anti-IgE hexos						BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6	
R950177	2.361	8.434													
R950178	0.688	>10													
R950179	0.955	>10													
R950180	0.278	9999													
R950181	0.254	9999													
R950182	0.627	9999													
R950183	4.797	9999													
R950184	2.222	9999													
R950185	1.03	8.81													
R950186	0.558	>10													
R950187	0.724	>10													
R950188	2.327	9999													
R950189	10	9999													
R950190	1.573	9999													
R950191	0.178	9999								<0.22		>2	0.401	<0.22	<0.22
R950192	0.244	9999													
R950193	0.61	9999													
R950194	2.04	9999													
R950195	0.473	9999													
R950196	2.2	9999													
R950197	0.531	9999													
R950198	0.406	>10													
R950199	0.408	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R950200	0.245	9999												
R950201	0.261	9999												
R950202	3.218	9999												
R950203	9.035	9999												
R950204	6.285	9999												
R950205	8.997	9999												
R950206	3.66	>10												
R950207	0.164	9999							<0.22		<0.22	0.288	<0.22	<0.22
R950208	0.267	9999												
R950209	0.748	9999												
R950210	10	9999												
R950211	10	9999												
R950212	0.253	9999												
R950213	9999	9999												
R950214	10	9999												
R950215	0.409	9999												
R950216	0.327	9999												
R950217	0.34	9999												
R950218	0.292	9999												
R950219	0.439	9999												
R950220	0.489	9999												
R950221	0.636	9999												
R950222	0.865	9999												

TABLE 1

TABLE 1																
Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density						
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	BMMC anti-IgE hexos						BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6		
R950223	0.763	9999														
R950224	0.687	9999														
R950225	5.283	9999														
R950226	1.374	9999														
R950227	1.029	9999														
R950229	0.98	9999														
R950230	7.91	9999														
R950231	1.968	9999														
R950232	10	9999														
R950233	0.98	9999														
R950234	10	9999														
R950235	4.095	9999														
R950236	0.955	9999														
R950237	9999	9999														
R950238	10	9999														
R950239	2.063	9999														
R950240	1.766	9999														
R950241	3.275	9999														
R950251	9999	9999														
R950253	0.697	9999														
R950254	0.496	9999														
R950255	10	9999														
R908698	1.67	9999														

TABLE 1

TABLE 1																
Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density							
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6		
R908699	0.217	9999														
R908700	1.273	9999														
R908701	0.099	7.643														
R908702	0.104	7.395														
R908703	0.63	9999														
R908704	0.511	9999														
R908705	0.801	9999														
R908706	0.445	9999														
R908707	1.834	9999														
R908709	2.414															
R908710	1.838	99														
R908711	1.761															
R908712	0.075	99														
R908734	1.379															
R909255	0.244	9999														
R909259	0.43	9999														
R909260	1.041	9999														
R909261	0.93	9999														
R909263	0.289	9999														
R909264																
R909265	99															
R909266	99															
R909267	0.589	9999														

TABLE 1

TABLE 1																
Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density						
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	BMMC anti-IgE hexos						BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6		
R909268	0.071	9999														
R909290	0.226															
R909292	1.172															
R909308	0.671	9999														
R909309	0.083	9999														
R920394																
R920395	0.092	9999														
R920396																
R920397																
R920398																
R920399																
R920404																
R920405																
R920406																
R920407																
R920408																
R920410	0.125	9999														
R920411	0.564	9999														
R925745	1.766	9999														
R926238	9999															
R926752	0.338	9999														
R926753	0.108	9999														
R926754	0.388	9999														

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4							BMHC anti-IgE hexos	BMHC Ionomycin Hexos.	BMHC anti-IgE histamine	BMHC anti-IgE LTC4	BMHC anti-IgE TNF-alpha	BMHC anti-IgE IL-6
R926755	1.693	9999													
R926756	1.365	9999													
R926757	0.158	9999													
R926759	0.688	9999													
R926760	2.893	9999													
R926761	0.245	9999													
R926762	0.386	9999													
R926763	0.195	9999													
R926794	1.382	9999													
R926826	0.613	9999													
R926827	1.098	9999													
R926828	0.306	9999													
R926829	0.688	9999													
R926830	0.569	10													
R926831	0.133	10													
R926832	0.365	9999													
R926833	1.129	9999													
R926834	0.145	9999													
R926835	0.296	9999													
R926836	10	9999													
R926837	2.994	9999													
R926838	0.583	9999													
R926839	0.161	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMHC anti-IgE hexos	BMHC Ionomycin Hexos.	BMHC anti-IgE histamine	BMHC anti-IgE LTC4	BMHC anti-IgE TNF-alpha	BMHC anti-IgE IL-6
R926840	1.1	9999												
R926841	0.551	9999												
R926842	7.733	9999												
R926843	7.371	9999												
R926844	1.1	9999												
R926845	2.558	7.812												
R926846	0.86	6.264												
R926847	1.479	6.264												
R926848	0.254	10												
R926851	0.446													
R926855	9999	9999												
R926856	0.734	9999												
R926857	1.209	9999												
R926859														
R926860	1.949	99												
R926862	0.774	9999												
R926863														
R926866														
R926870	3.294													
R926871	2.146													
R926874	0.638	9999												
R926879	0.397	9999												
R926880														

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE LTC4	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926881														
R926883														
R926885														
R926886														
R926887	1.747													
R926890	0.361	9999												
R926891	0.152	9999												
R926892	0.685	9999												
R926893	10	9999												
R926894	9999	9999												
R926895	0.339	9999												
R926896	1.622	9999												
R926897	1.727	9999												
R926898	1.1	9999												
R926899	1.1	9999												
R926900	9999	9999												
R926902	1.37	4.586												
R926903	0.243	9999												
R926904	0.538													
R926905	99													
R926906	0.794													
R926907	0.764													
R926908	0.585													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMHC anti-IgE hexos	BMHC Ionomycin Hexos.	BMHC anti-IgE histamine	BMHC anti-IgE LTC4	BMHC anti-IgE TNF-alpha	BMHC anti-IgE IL-6
R926909	0.379													
R926913	0.548	9999												
R926914	1.86	9999												
R926915	1.713	9999												
R926916	1.958	9999												
R926917	1.169	9999												
R926918	2.521	9999												
R926919	1.413	9999												
R926922	0.305	9999												
R926923	0.346	9999												
R926925	0.307	99												
R926926	0.401	9999												
R926927	0.348	9999												
R926928	0.575	9999												
R926929	1.916	9999												
R926930	99	9999												
R926931														
R926932	0.31	9999												
R926933														
R926934														
R926935	4.44													
R926936														
R926937														

TABLE 1

TABLE 1																
Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density						
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	BMMC anti-IgE hexos						BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6		
R926938																
R926939	3.615															
R926940	7.754															
R926941	4.195															
R926942	4.81															
R926943																
R926944	0.225	99														
R926945	0.457	9999														
R926946																
R926947	0.354	9999														
R926948	0.246	9999														
R926949	0.089	9999														
R926950	99	9999														
R926951	0.183	9999														
R926953	0.049	9999														
R926954	0.284	9999														
R926955	0.36	9999														
R926956	0.211	9999														
R927016	1.408															
R927017	2.449															
R927018	1.446															
R927019	1.179															
R927020	1.316	9999														

TABLE 1

TABLE 1																
Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density						
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	BMMC anti-IgE hexos						BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6		
R927023	0.918	9999														
R935221	9999	9999														
R935222	0.52	9999														
R935223	0.469	9999														
R935224	4.578	9999														
R935225	6.495	9999														
R935237	0.24	9999														
R935238	1.854	9999														
R935239	0.609	9999														
R935240	0.606	9999														
R935242	2.855	9999														
R935248	1.1	9999														
R935249	1.1	9999														
R935250	1.1	9999														
R935251																
R935252																
R935253																
R935255	0.374	9999														
R935256	0.324	9999														
R935258	1.191	9999														
R935259	1.777	9999														
R935261	0.391	9999														
R935262	0.516	9999														

TABLE 1

Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC ionomycin Tryptase	CHMC anti-IgE LTC4	CHMC anti-IgE Hexos.						BMMC anti-IgE hexos	BMMC ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R935263	0.106	10													
R935264	0.135	9999													
R935266	2.97														
R935267	2.463														
R935268	1.059														
R935269	1.715														
R935271															
R935276	2.33														
R935277	22.883	8.9													
R935278	4.753	9999													
R935279	0.889	9999													
R935280	99														
R935281	1.399	9999													
R935286	1.158	9999													
R935287	0.403	9999													
R935288	1.58	9999													
R935289	1.688	9999													
R935290	0.34	9999													
R935291	1.364	9999													
R935292	0.483	9999													
R935293	0.141	9999													
R935294	0.388	9999													
R935295	1.943	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R935296	99	9999												
R935297	7.328	9999												
R935298	0.252	99												
R935299	0.21	9999												
R935300	0.243	9999												
R935301	4.05	99												
R935302	0.189	9999												
R935303	0.244	99												
R935304	0.188	9999												
R935305	0.495	9999												
R935306	0.345	99												
R935307	0.139	99												
R935308	0.275	9999												
R935309														
R935310														
R935320	2.769													
R935321	2.986													
R935322	3.416													
R935323	9999													
R935324	9999													
R935336	0.341	9999												
R935337	9999													
R935338	0.411	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R935339	9999													
R935340	3.606													
R935351	9999	9999												
R935352														
R935353	9999	9999												
R935354	99	9999												
R935355	9999	9999												
R935356	99													
R935357	99	9999												
R935358	9999	9999												
R935359	1.027	9999												
R935360	0.903	9999												
R935361	1.438	9999												
R935362	0.409	9999												
R935363	0.405	9999												
R935364	0.563	9999												
R935365	0.373	9999												
R935366	0.216	9999												
R935367	0.053	9999												
R940079	9999													
R940110	9999	9999												
R940299	2.497	9999												
R940300	10	9999												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R940301	1.975	9999												
R940304	9999	9999												
R940306	1.1	9999												
R940307	0.291	9999												
R940308	0.612	4.168												
R940309	1.132	9999												
R940311	1.95													
R940312	2.557													
R940314	4.197													
R940316	1.858													
R940317	0.913	9999												
R940318	3.792													
R940319	9999													
R940321	9999													
R940323	0.048	9999												
R940337	1.098													
R940338	0.073	9999												
R921303	0.033	99												
R940345	1.712													
R940346	0.142	99												
R940347	0.063	99												
R940348	2.189													
R940349	0.044	7.4												

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R940350	0.092	4												
R940351	0.12	2.7												
R940352	0.101	9999												
R940353	0.091	9999												
R940354	0.115	99												
R945236	0.562	9999												
R945237	0.461	9999												
R945242	0.247	9999												
R945263	1.642													
R921304	0.085	9999												
R945299														
R950244	9999													
R950245	9999													
R950246	9999													
R950247	9999													
R950261	0.611	9999												
R950262	0.285	9999												
R950263	0.284	3.299												
R950264	0.198	9999												
R950265	0.312	9999												
R950266	0.645	9999												
R950267	0.18	9999												
R950290	9999	9999												

TABLE 1

TABLE 1															
Test Compound	Low Density				CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4	BMMC anti-IgE hexos						BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6	
R950291	9999	9999													
R950293	3.689	8.155													
R950294	2.005	8.005													
R950295	2.041	8.795													
R950296	0.495	9999													
R950344	99														
R950345	1.962	99													
R950346	0.345	9999													
R950347	0.548														
R950348	0.066														
R950349	0.078	9999													
R950356															
R950368	0.038	9999													
R950371															
R950372	1.348	9999													
R950373															
R950374	0.599	9999													
R950376	2.539														
R950377	99														
R950378															
R950379	0.545	9999													
R950380	3	9999													
R950381	0.11	99													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC anti-IgE Hexos.	CHMC anti-IgE Histamine	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4							BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R950382															
R950383	0.114	9999													
R950385															
R950386	0.973														
R950388	2.518														
R950389	0.612	9999													
R950391	999	9999													
R950392	0.956	9999													
R950393	0.404	9999													
R945028															
R935241															
R940298															
R940302															
R940303															
R940305															
R935260	9999														
R909258															
R940313	9999														
R940315	9999														
R935275	9999														
R940320	9999														
R940322	9999	9999													
R926910	9999	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R926911	9999	9999												
R926912	9999	9999												
R926853	9999	9999												
R926852	9999	9999												
R926854	9999	9999												
R926920	9999	9999												
R926921	99	9999												
R926924	99	9999												
R926858														
R926861	9999	9999												
R945298	9999	9999												
R940328	9999													
R926869														
R926873	9999													
R926875	9999													
R926876	9999													
R926877	9999													
R940336	9999													
R926878	9999													
R926882	9999													
R926884	9999													
R926889	9999													
R920400	9999													

TABLE 1

Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R920401	9999													
R920402	9999													
R920403	9999													
R940342	99													
R920409	9999													
R940344	9999													
R926888	9999													
R926758														
R927024	0.326	99												
R927025	0.326													
R927026	9999	9999												
R927027	9999	9999												
R927028	0.208	9999												
R927029														
R927030	0.26	9999												
R927031	0.215	99												
R927032	0.899													
R927035	0.583	9999												
R927036														
R927037	0.233	9999												
R927038	1.05	9999												
R927039	1.23	9999												
R927040	1.05	9999												

TABLE 1

TABLE 1														
Test Compound	Low Density			CHMC anti-IgE Hexos.	CHMC Ionomycin Hexos.	Basophils anti-IgE Histamine	Basophils Ionomycin Histamine	Basophils Dust mite Histamine	High Density					
	CHMC anti-IgE Tryptase	CHMC Ionomycin Tryptase	CHMC anti-IgE LTC4						BMMC anti-IgE hexos	BMMC Ionomycin Hexos.	BMMC anti-IgE histamine	BMMC anti-IgE LTC4	BMMC anti-IgE TNF-alpha	BMMC anti-IgE IL-6
R927041	0.788	9999												
R927042														
R935270														
R935368	0.082	9999												
R935369	0.255	9999												
R935370														
R935371	0.794	9999												
R935372	0.06	9999												
R935373	0.274	9999												
R935374	0.356	9999												
R935375	10	9999												
R935376														
R935377														
R935378	0.566	9999												
R935379														
R935380	1.61	99												

TABLE 2

TABLE 2											
	High Density						Toxicity Jurkat Light Scat.	Toxicity Jurkat Cell Titer Glo	Toxicity BJAB Light Scat.	Toxicity BJAB Cell Titer Glo	
	CHMC high density hexos	CHMC high density tryptase	CHMC high density histamine	CHMC high density LTC4	CHMC high density TNF-alpha	CHMC high density IL-13					
R008951											
R008952											
R008953											
R008955											
R008956											
R008958											
R067934											
R067963											
R070153											
R070791											
R081166											
R088814											
R088815											
R091880											
R092788								9999		9999	
R909241									3.736		
R921219	0.124	0.121	0.162	0.034	0.190	0.175			>10	>10	
R925775								9999		9999	
R925778								9999		9999	
R925779								>10		9999	
R925797								>10		9999	
R926108								>10		>10	
R926109	0.783	0.906	1.827	0.808	1.504	1.664		>10		9999	

TABLE 2

TABLE 2											
	High Density						Toxicity Jurkat Light Scat.	Toxicity Jurkat Cell Titer Glo	Toxicity BJAB Light Scat.	Toxicity BJAB Cell Titer Glo	
	CHMC high density hexos	CHMC high density tryptase	CHMC high density histamine	CHMC high density LTC4	CHMC high density TNF-alpha	CHMC high density IL-13					
R926110							>10		>10		
R921218	0.464	0.647	0.463	0.695	1.752	2.0776	>10		>10		
R926113	1.448	1.649	1.848	0.468	5.678	3.569	>10		>10		
R926146							9999		9999		
R926210							>10		9999		
R926240							10		9999		
R926248							>10		9999		
R926249							>10		9999		
R926253							9999		9999		
R926256							>10		9999		
R926258							9999		9999		
R926387							>10		9999		
R926395							>10		9999		
R926396							>10		9999		
R926411							8.5		>10		
R926486	1.088	1.313	1.928	0.834	0.455						
R926488	0.521	0.623	0.792	0.201	2.443	1.012					
R926493	0.889	1.093	1.324	0.474	>2			>4.33			
R926494	0.640	>2	9999	0.326	9999						
R926495	0.100	0.235	0.066	0.241	0.362	0.449		>10		>10	
R926496	0.429	0.533	0.809	0.414	0.622						
R926497	1.106	1.234	1.333		1.876	9999					
R926501	>2	>2	9999		9999	9999		>4.33		>4.33	

TABLE 2

TABLE 2													
	High Density						Toxicity Jurkat Light Scat.	Toxicity Jurkat Cell Titer Glo	Toxicity BJAB Light Scat.	Toxicity BJAB Cell Titer Glo			
	CHMC high density hexos	CHMC high density tryptase	CHMC high density histamine	CHMC high density LTC4	CHMC high density TNF-alpha	CHMC high density IL-13							
R926502	>2	>2	>2		1.807	>2		1.513					
R926505								4.199					
R926508	0.170	0.434	0.105		0.505	0.763		>10		>10			
R926510	0.921	1.115	1.667		0.417	0.686		2.77					
R926511	1.183	1.474	1.73		1.307	>2		>4.33		>4.33			
R926614	>10	>10			>10	6.442							
R926696	<1.1	<1.1	<1.1	<1.1	<1.1	1.773		>5.0					
R926699	<1.1	<1.1	1.44	<1.1	<1.1	1.294							
R926700	<1.1	<1.1	<1.1	<1.1	<1.1	2.053							
R926703	1.512	1.947	>2	0.724	>2								
R926704	>2	9999	9999	9999	9999								
R926705	1.007	1.256	0.641	0.494	9999								
R926706	>2	9999	9999	1.491	9999								
R926742	0.104	0.217	0.080		0.385	0.667		9		>10			
R926745								>10		>10			
R926780								>5.0					
R926782								>4.33		>4.33			
R935075	0.647	1.212	0.443	<0.22	>2			>4.33		>4.33			
R935154								>4.33					
R935156								4.054					
R940216	<1.1	<1.1	1.176	<1.1	3.188	3.006							
R940233	0.577	0.642	0.586	0.118	2.247	1.781		>4.33		>4.33			
R945032	0.357	0.458	0.439	0.0929	1.082	0.291							

TABLE 2

TABLE 2											
	High Density						Toxicity Jurkat Light Scat.	Toxicity Jurkat Cell Titer Glo	Toxicity BJAB Light Scat.	Toxicity BJAB Cell Titer Glo	
	CHMC high density hexos	CHMC high density tryptase	CHMC high density histamine	CHMC high density LTC4	CHMC high density TNF-alpha	CHMC high density IL-13					
R945033	8.151	8.868			>10	5.983					
R945071	<1.1	<1.1	<1.1	<1.1	<1.1	<1.1					
R945128	1.279	1.749	0.547	0.729	>2	ND					
R945140	0.994	1.112	1.551		1.714	9999					
R945142	>2	>2	9999		>2	9999					
R945150								>4.33		>4.33	
R921302	0.682	0.795	1.588	0.514	1.173	1.672					
R950141	0.567	0.618	0.627	0.201	1.059	0.798					
R950207								>4.33			

7.7 The 2,4-Pyrimidinediamine Compounds of the Invention Selectively Inhibit the Upstream IgE Receptor Cascade

To confirm that many of the 2,4-pyrimidinediamine compounds of the invention exert their inhibitory activity by blocking or inhibiting the early IgE receptor signal transduction cascade, several of the compounds were tested in cellular assays for ionomycin-induced degranulation, as described below.

7.7.1 CHMC Low Cell Density Ionomycin Activation: Tryptase Assay

Assays for ionomycin-induced mast cell degranulation were carried out as described for the CHMC Low Density IgE Activation assays (Section 6.4.3, *supra*), with the exception that during the 1 hour incubation, 6X ionomycin solution [5mM ionomycin (Sigma I-0634) in MeOH (stock) diluted 1:416.7 in MT buffer (2 μ M final)] was prepared and cells were stimulated by adding 25 μ l of the 6X ionomycin solution to the appropriate plates.

7.7.2 Basophil Ionomycin Activation: Histamine Release Assay

Assays for ionomycin-induced basophil cell degranulation were carried out as described for the Basophil IgE or Dustmite Activation Assay (Section 6.4.6, *supra*), with the exception that following incubation with compound, cells were stimulated with 20 μ l of 2 μ M ionomycin.

7.7.3 Results

The results of the ionomycin-induced degranulation assays, reported as IC₅₀ values (in μ M) are provided in TABLE 1, *supra*. Of the active compounds tested (*i.e.*, those that inhibit IgE-induced degranulation), the vast majority do not inhibit ionomycin-induced degranulation, confirming that these active compounds selectively inhibit the early (or upstream) IgE receptor signal transduction cascade.

These results were confirmed for certain compounds by measuring anti-IgE-induced and ionomycin-induced calcium ion flux in CHMC cells. In these Ca²⁺ flux tests, 10 μ M R921218 and 10 μ M R902420 inhibited anti-IgE-induced Ca²⁺ flux, but had no effect on ionomycin-induced Ca²⁺ flux (See FIG. 4).

7.8 The Inhibitory Effect of the 2,4-Pyrimidinediamine Compounds of the Invention is Immediate

To test the immediacy of their inhibitory effect, certain 2,4-pyrimidinediamines of the invention were added simultaneously with anti-IgE antibody activator in the cellular assays described above. All compounds tested blocked IgE-induced degranulation of CHMC cells to the same extent as observed when the compounds were pre-incubated with CHMC cells for 10 or 30 min. prior to receptor cross-linking.

7.9 Kinetics of Pharmacological Activity *In vitro*

Compounds R921218, R921302, R921219, R926240, R940277, R926742, R926495, R909243 and R926782 were tested in washout experiments. In the experiments, CHMC cells were either activated immediately with anti-IgE antibody in the presence of 1.25 μ M compound (time zero), or the compound was washed out followed by activation with anti-IgE antibody at 30, 60 or 120 min. The inhibitory activity of these compounds was greatly diminished 30 min. after compound removal, indicating that constant exposure of mast cells to these compounds is required for maximal inhibition of degranulation. The other compounds tested yielded similar results.

7.10 Toxicity: T- and B-Cells

The ability of the compounds of the invention to exert their inhibitory activity without being toxic to cells of the immune system was demonstrated in cellular assays with B- and T-cells. The protocols for the assays are provided below.

7.10.1 Jurkat (T-Cell) Toxicity

Dilute Jurkat cells to 2×10^5 cells/ml in complete RPMI (10% heat-inactivated fetal bovine serum) media and incubate at 37°C, 5% CO₂ for 18 hours. Add 65 μ l cells at 7.7×10^5 cells/ml to a 96-well V-bottom plate (TC-treated, Costar) containing 65 μ l 2X compound (final vehicle concentration is 0.5% DMSO, 1.5% MeOH). Mix, incubate plates for 18-24 hr at 37°C, 5% CO₂. Toxicity was assessed by flow cytometric analysis of cellular light scatter

7.10.2 BJAB (B-Cell) Toxicity

The B-cell line BJAB was cultured in log phase in RPMI1640 + 10% heat-inactivated fetal bovine serum, 1x L-glutamine, 1x penicillin, 1x streptavidin and 1x beta-

mercaptoethanol at 37°C, 5% CO₂. First, BJABs were harvested, spun and resuspended in culture medium to a concentration of 7.7×10^5 cells/mL. 65 μ L cells were mixed with 65 μ L compound, in duplicate and in the presence of 0.1% DMSO in a V-bottomed 96-well tissue culture plate. Cells were incubated with compound at various dilutions at 37°C, 5% CO₂.

5 Toxicity was assessed by flow cytometric analysis of cellular light scatter.

7.10.3 Toxicity: Cell Titer Glo Assay

Seed 50 μ L cells (1×10^6 /mL) into each well containing 50 μ L compound. The final vehicle concentration is 0.5% DMSO, 1.5% MeOH. Shake plates for 1 minute to mix cells and compound. Incubate plates at 37°C (5% CO₂) for 18 hours. Next day, harvest 50 μ L cells from each well, add to 50 μ L Cell Titer Glo reagent (Invitrogen). Shake plates for 1 minute. Read on luminometer.

7.10.4 Results

The results of the T- and B-cell toxicity assays, reported as IC₅₀ values (in μ M), are presented in TABLE 2, *supra*. With a few exceptions (see TABLE 1), all compounds tested were non-toxic to both B- and T-cells at effective inhibitory concentrations. Assays performed with primary B-cells yielded similar results.

7.11 The 2,4-Pyrimidine Compounds Are Tolerated In Animals

The ability of the compounds of the invention to exert their inhibitory activity at doses below those exhibiting toxicity in animals was demonstrated with compounds R921218, R921219 and R921302.

7.11.1 R921218

R921218 was studied in an extensive program of non-clinical safety studies that concluded this agent to be well tolerated in both rodents and non-rodents. To summarize the outcome of toxicology/non-clinical safety testing with R921218; this agent produced no dose limiting toxicity by the intranasal route of administration in non-rodents (rabbits and primates) or by the oral route of administration in rodents (mice and rats) during 14-day repeat-dose toxicity studies at doses many fold above the anticipated dose expected to produce efficacy in man. There were no adverse findings in a core safety pharmacology battery of cardiovascular, respiratory and/or central nervous system function. There was no evidence for mutagenic or clastogenic potential in genetic toxicology testing

nor were there untoward effects after exposure to skin and eyes. A short discussion of key toxicology studies is provided.

A 14-day repeat-dose intranasal toxicity study in *Cynomolgus* monkeys was performed at doses of 2.1, 4.5 or 6.3 mg/kg/day. In life parameters included: clinical observations, body weights, food consumption, ophthalmology, blood pressure, electrocardiography, hematology, clinical chemistry, urinalysis, immunotoxicological assessment, gross necropsy, organ weights, toxicokinetic assessments and histopathology (including the nasal cavity). There were no adverse findings attributed to R921218 in any study parameter and the NOAEL (no observed adverse effect level) was considered 6.3 mg/kg/day.

A 14-day repeat-dose intranasal toxicity study in New Zealand White rabbits was performed at doses of 1.7, 3.4 or 5.0 mg/kg/day. In life parameters included: clinical observations, body weights, food consumption, ophthalmology, hematology, clinical chemistry, gross necropsy, organ weights, toxicokinetic assessments and histopathology (including the nasal cavity). There were no adverse findings attributed to R921218 in any study parameter and the NOAEL (no observed adverse effect level) was considered 5.0 mg/kg/day.

7.11.2 R921219

In pilot dose finding studies a single dose oral dose of 600 mg/kg was considered a NOEL (no observed effect level) while multiple (7-day) doses of 200 mg/kg/day and above were not tolerated.

In the *in vitro* *Salmonella-Escherichia coli*/Mammalian-Microsome Reverse Mutation Assay (Ames test), R921219 was found to test positive in tester strain TA1537, with and without metabolic activation, confirming the results of an earlier study. R921219 was not found to adversely affect any of the other 4 tester strains. R921219 was not found to possess clastogenic potential when studied in an *in vitro* chromosomal aberration assay.

7.11.3 R921302

Several non-GLP pilot toxicity studies have been conducted in rodents. In the mouse an oral dose of 1000 mg/kg was tolerated for up to 7-days. In a 14-day oral toxicity study in the mouse was conducted with doses of 100, 300 and 1000 mg/kg. A dose of 1000 mg/kg was not tolerated, while a dose of 300 mg/kg promoted evidence for

histopathological changes in the vulva. A dose of 100 mg/kg was considered the NOAEL (no observed adverse effect level) in the study. A 28-day oral toxicity study in the mouse was conducted at doses of 100 mg/kg q.d., 100 mg/kg b.i.d., 300 mg/kg q.d. and 300 mg/kg b.i.d. R921302 was not tolerated at 300 mg/kg q.d. or b.i.d. The lower doses (100 mg/kg q.d. or b.i.d.) appeared to be well tolerated (results of clinical and histopathology are not yet known). In the rat oral doses of 50, 150 and 300 mg/kg given for 32 days appeared to be well tolerated (results of clinical and histopathology are not yet known).

In the *in vitro* *Salmonella-Escherichia coli*/Mammalian-Microsome Reverse Mutation Assay (Ames test), R921302 was found to test positive in tester strain TA98 with S9 and TA1537, with and without metabolic activation. R921302 was not found to adversely affect any of the other 3 tester strains. R921302 was not clastogenic when assessed in an *in vitro* chromosomal aberration assay.

7.12 The 2,4-Pyrimidinediamine Compounds Are Orally Bioavailable

Over 50 2,4-pyrimidinediamine compounds of the invention were tested for oral bioavailability. For the study, compounds were dissolved in various vehicles (e.g. PEG 400 solution and CMC suspension) for intravenous and oral dosing in the rats. Following administration of the drug, plasma samples were obtained and extracted. The plasma concentrations of the compounds were determined by high performance liquid chromatography/tandem mass spectrometry (LC/MS/MS) methods. Pharmacokinetic analyses were performed based on the plasma concentration data. The pharmacokinetic parameters of interest include Clearance (CL), Volume of distribution at steady-state (V_{ss}), terminal half-life ($t_{1/2}$), and oral bioavailability (%F).

These pharmacokinetic studies indicate that many of the 2,4-pyrimidinediamine compounds are orally available, with %F up to approximately 50% (in the range of 0-50%). The half-lives ranged from 0.5 to 3 hr. In particular, Compounds R940350, R935372, R935193, R927050 and R935391 exhibited good oral bioavailabilities and half-lives in rats. Thus, these studies confirm that these 2,4-pyrimidinediamine compounds are suitable for oral administration.

7.13 The Compounds Are Effective for the Treatment of Allergies

The *in vivo* efficacy of compounds R926109, R921218, R921219, R921302, R926495, R926508, R926742, R926745 and R945150 towards allergies was evaluated in

the mouse model of passive cutaneous anaphylaxis (PCA). This model provides a direct measure of IgE-induced degranulation of tissue mast cells. In this model, IgE primed animals are exposed to an allergen challenge, and the change in permeability of dermal vasculature that results from histamine release from mast cells is measured by change in the amount of dye leakage into surrounding tissue. Inhibition of mediator release by compounds that modulate mast cell degranulation is easily measured by extracting the dye from the tissue.

7.13.1 Study Protocol and Results

In the PCA assay mice are passively sensitized by intradermal injection with anti-dinitrophenol (DNP) IgE antibodies (Day -1). At predetermined times animals are treated with the test agent (Day 0). The modulatory effect of the agent on cutaneous mast cell degranulation is measured following intravenous injection of DNP conjugated to human serum albumin (HSA-DNP), together with Evans blue dye. The resulting cross-linking of the IgE receptor and subsequent mast cell degranulation-induced increase in vascular permeability is determined by measuring the amount of dye extravasation into the tissue. Dye is extracted from the tissue by formamide, and the absorbance of this extract is read at 620 nm. The inhibitory effect of drug treatment is reported as the percent inhibition compared to vehicle treatment, that is, the percent reduction in A_{620} .

Two compounds have been tested as positive controls: the histamine antagonist diphenhydramine and the serotonin antagonist cyproheptadine. Both mediators (histamine and serotonin) are released upon IgE-mediated degranulation from the mouse mast cell. Both reference compounds inhibit the PCA response; cyproheptadine was used routinely in subsequent experiments. Cyproheptadine reproducibly inhibited the PCA response by 61% +/- 4% (8 mg/kg, i.p., 30 minutes pretreatment time, n=23 experiments).

7.13.1.1 Results

A dose-dependent inhibition of the FcεR--mediated vascular leakage was observed with increasing doses of R921218, R926109, R921219 and RR921302. These compounds were administered either in a solution formulation (67%PEG/33% citrate buffer) or an aqueous suspension (1.5% Avicel). These results demonstrate the strong correlation between compound plasma levels, in vivo efficacy, and *in vitro* potency. The most potent compound, R921219, was active with circulating exposure levels of

approximately 10 $\mu\text{g/ml}$ (68% inhibition at a dose level of 100 mg/kg) compared with R921302, a relatively less potent molecule, which reduced plasma extravasation by 42% at a dose level of 100 mg/kg. Further, the length of exposure to circulating compound was reflected in the duration of inhibitory activity. R921302, determined to be the most

5 metabolically stable compound in pharmacokinetics studie, inhibited the vascular permeability for 1-2 hours prior to antigen-induced receptor signaling, where after the efficacy began to decrease. These data are summarized in TABLE 3 and TABLE 4.

TABLE 3						
Efficacy of R921218, R926109, R921219 and R921302 in the PCA Assay						
Compound	Route	Vehicle	Pretreatment time (min)	Dose (mg/kg)	% Inhibition	Plasma level (µg/ml)
R921218	PO	67%PEG/33% citrate buffer	10	50	7	3
				100	11	4
				200	50	18
R926109	PO	67%PEG/33% citrate buffer	15	50	22	N.D.
				100	32	
				200	48	
R921219	PO	1.5% Avicel/water	15	30	25	0.4
				100	68	4
				300	92	11
R921302	PO	1.5% Avicel/water	60	50	35	25
				100	42	38
				150	56	64
				200	93	105

TABLE 4						
Duration of action of R921219 and R921302 in the PCA Assay						
Compound	Route	Vehicle	Dose (mg/kg)	Pretreatment time (min)	% Inhibition	Plasma level (µg/ml)
RR921302	PO	1.5% Avicel/water	200	30	89	88
				60	83	53
				120	82	61
				240	37	8

Similar in vivo activity was observed with compounds R926495, R926508, R926742,

5. R926745 and R926150, which were able to inhibit the PCA response after administration by the oral route in a PEG-based formulation (data not shown).

7.14 The Compounds Are Effective in the Treatment of Asthma

The efficacy of compounds R921218, R921302, R926495, R926508, R926742 and R921219 in the treatment of asthma was demonstrated in the sheep model of allergic asthma. Sheep develop bronchoconstriction within minutes of exposure to inhaled antigen (Ascaris suum), with maximal airflow obstruction during the early allergic response (EAR). Release of preformed mast cell mediators is likely responsible for this early phase of airflow obstruction. In addition to the EAR, the sheep model allows us to evaluate the effect of our compounds on the late asthmatic reaction (LAR) and non-specific airway hyperresponsiveness (AHR), which occur as a result of topical or local administration of allergen to the airway. In the sheep, AHR develops a few hours following antigen challenge, and can persist for up to 2 weeks. The results described below demonstrate the potential of the tested compounds to inhibit a cascade of events that may be a result of release of cytokines from the mast cell.

7.14.1 Study Protocol

In the sheep model of allergic asthma, sheep are administered aerosols of test article *via* an endotracheal tube, followed by an aerosol challenge with antigen extracted from the roundworm, *Ascaris suum*, to which the sheep are naturally allergic. Allergen challenge leads to direct bronchoconstriction (both EAR and LAR) and a persistent non-specific AHR. These three characteristics are similar to those seen in human allergic asthmatics. The activity of the test agent is determined by changes in the lung resistance (R_L), which is calculated from measurements of transpulmonary pressure, flow, and respiratory volume. The historical control data obtained from the same sheep following saline treatment compared with an allergen challenge show that a sharp increase of R_L occurs during the EAR and persists for approximately 2-3 hours following allergen challenge. The LAR is a less pronounced increase in R_L , which starts approximately 5-6 hours following allergen challenge and is resolved by 8 hours post-challenge. Twenty-four hours after the challenge, a dose response to carbachol is measured to determine the AHR, which is expressed as the dose of carbachol required to increase R_L by 400% over baseline. (This measurement is referred to as the provocative concentration of carbachol that elicits a 400% increase in R_L over baseline (PC_{400}). The data are compared to historical control data for the same individual when administered a saline control aerosol and challenged with *Ascaris suum*.

7.14.2 Result

All the compounds tested showed inhibitory effects in the LAR and the AHR, and several of these agents inhibited the EAR as well. The optimal response for each compound in a series of studies to evaluate activity at several pretreatment times and using several different solution and suspension formulations are shown in TABLE 5. The efficacy of R921218 on the EAR appeared to be dependent on the formulation, with the greatest effect seen at 30 mg/sheep administered as a solution aerosol in 10% ethanol. R926495, R926742, R926508 and R921219, administered in four different sheep at 45 mg/sheep in an aqueous suspension 60 minutes prior to allergen challenge, demonstrate that the LAR and AHR is blocked. In addition to these late parameters, the EAR was greatly reduced by treatment with R921219, R926508 or R926495. The efficacy of RR921302 was investigated using a 45%PEG400/55% citrate buffer vehicle. Under these conditions, R921302, administered at 30 mg/sheep 60 minutes prior to challenge, blocked the LAR and AHR, and EAR was unaffected.

These data clearly demonstrate that these compounds are able to block the asthmatic responses in allergic sheep. All compounds inhibited the AHR and LAR significantly when compared to the historical control. The EAR was significantly inhibited by R921219, R926508 and R926495 (54%, 21% and 33% respectively). In contrast, R921218, R921302 and R926742 failed to inhibit the EAR when administered in an aqueous suspension.

TABLE 5
Efficacy Of Exemplary Compounds In A Sheep Model Of Allergic Asthma

Compound	Dose (mg/sheep)	Pretreatment time (min)	Vehicle	EAR (%) inhibition)	LAR (%) inhibition)	AHR (%) inhibition)
R921218	30	15	10% ethanol	66	78	101
R926742	45	60	Aqueous suspension	-19	87	94
R926495	45	60		33	85	41
R926508	45	60		21	90	88
R921219	45	60		56	75	90
RR921302	30	60	45%PEG400/55% citrate buffer	-28	86	82

7.15 The Compounds Are Effective In The Treatment of Asthma

The efficacy of compounds R921304 and R921219 in the treatment of asthma was also demonstrated in a mouse model of allergic asthma.

7.15.1 Study protocol

5 Mice are sensitized to ovalbumin (chicken protein) in the presence of an adjuvant (Alum) by the intraperitoneal route on day 0 and day 7. One week later, mice are challenged intranasally with ovalbumin on Days 14, 15 and 16 (more stringent model) or on Day 14 (less stringent model). This sensitization and challenge regimen leads to airway hyperresponsiveness and inflammation in the lungs, which are two dominant characteristics of human allergic asthma. In the mouse model, the in vivo airway responses are measured using a whole body plethysmograph which determines the PENH (enhanced Pause, Buxco Electronics). The PENH is a dimensionless value comprised of the peak inspiratory flow (PIF), peak expiratory flow (PEF), time of inspiration, time of expiration and relaxation time, and is considered a validated parameter of airway responsiveness. Responses to allergen challenge (OVA) are compared with animals challenged with saline only. Twenty-four hours after challenge, mice are exposed to increasing doses of methacholine (muscarinic receptor agonist) which results in smooth muscle contraction. The ovalbumin-challenged mice demonstrate a significant airway hyperresponsiveness to methacholine when compared to the saline challenged mice. In addition, a cellular infiltrate in the airway is observed in ovalbumin challenged mice when compared with the saline challenged mice. This cellular infiltrate is mainly characterized by eosinophils, but a smaller influx of neutrophils and mononuclear cells is also present.

The use of this model for the evaluation of small molecule inhibitors of mast cell degranulation has been validated in several ways. First, using mast cell deficient mice (W/W^v) it has been shown that the ovalbumin-induced responses are dependent upon the presence of mast cells. In the mast cell deficient mice, ovalbumin sensitization and challenge did not result in airway hyperresponsiveness and eosinophil influx. Second, the mast cell stabilizer, Cromolyn, was able to block the ovalbumin-induced airway hyperresponsiveness and inflammation (data not shown). The use of this model to evaluate compounds for the treatment of asthmatic responses that may be mediated by mechanisms other than mast cell stabilization, is further supported by the inhibitory effect of the steroids, dexamethasone and budesonide, on methacholine-induced bronchoconstriction.

7.15.2 Results

The efficacy of R921304 was evaluated by intranasal administration on 10 consecutive days, from Day 7 through Day 16, at a dose level of 20 mg/kg, with the last 3 doses administered 30 minutes prior to either saline or ovalbumin challenge. R921304 was
5 able to inhibit the ovalbumin-induced airway hyperresponsiveness to methacholine when compared to the vehicle treated mice.

In a less stringent protocol, in which the mice were challenged with ovalbumin only once on Day 14, R921219 administered subcutaneously at 70 mg/kg in 67%PEG400/33% citrate buffer 30 minutes prior to saline or ovalbumin challenge, demonstrates that R921219
10 completely blocked the ovalbumin-induced airway hyperresponsiveness and cellular influx.

These results clearly demonstrate that R921219 and R921304 are efficacious in inhibiting the airway responses in a mouse model of allergic asthma.

7.16 2,4-Pyrimidinediamine Compounds Inhibit Phosphorylation of Proteins Downstream of Syk kinase in Activated Mast Cells

The inhibitory effect of the 2,4-pyrimidinediamine compounds on the phosphorylation of proteins downstream of Syk kinase was tested with compounds R921218, R218219 and R921304 in IgE receptor-activated BMMC cells.

For the assay, BMMC cells were incubated in the presence of varying concentrations
20 of test compound (0.08 μ M, 0.4 μ M, 2 μ M and 10 μ M) for 1 hr at 37°C. The cells were then stimulated with anti-IgE antibody as previously described. After 10 min, the cells were lysed and the cellular proteins separated by electrophoresis (SDS PAGE).

Following electrophoresis, the phosphorylation of the proteins indicated in FIGS. 7, 10 and 11A-D were assessed by immunoblot. Antibodies were purchased from Cell
25 Signaling Technology, Beverley, MA.

Referring to FIGS. 7, 10 and 11A-D, the indicated compounds tested inhibited phosphorylation of proteins downstream of Syk, but not upstream of Syk, in the IgE receptor signaling cascade, confirming both that the compounds inhibit upstream IgE induced degranulation, and that the compounds exert their inhibitory activity by inhibiting
30 Syk kinase.

7.17 2,4-Pyrimidinediamine Compounds Inhibit Syk Kinase in Biochemical Assays

Several 2,4-pyrimidinediamine compounds were tested for the ability to inhibit Syk kinase catalyzed phosphorylation of a peptide substrate in a biochemical fluorescenced polarization assay with isolated Syk kinase. In this experiment, Compounds were diluted to 1% DMSO in kinase buffer (20 mM HEPES, pH 7.4, 5 mM MgCl₂, 2 mM MnCl₂, 1 mM DTT, 0.1 mg/mL acetylated Bovine Gamma Globulin). Compound in 1% DMSO (0.2% DMSO final) was mixed with ATP/substrate solution at room temperature. Syk kinase (Upstate, Lake Placid NY) was added to a final reaction volume of 20 uL, and the reaction was incubated for 30 minutes at room temperature. Final enzyme reaction conditions were 20 mM HEPES, pH 7.4, 5 mM MgCl₂, 2 mM MnCl₂, 1 mM DTT, 0.1 mg/mL acetylated Bovine Gamma Globulin, 0.125 ng Syk, 4 uM ATP, 2.5 uM peptide substrate (biotin-EQEDEPEGDYEEVLE-CONH₂, SynPep Corporation). EDTA (10 mM final)/anti-phosphotyrosine antibody (1X final)/fluorescent phosphopeptide tracer (0.5X final) was added in FP Dilution Buffer to stop the reaction for a total volume of 40 uL according to manufacturer's instructions (PanVera Corporation) The plate was incubated for 30 minutes in the dark at room temperature. Plates were read on a Polarion fluorescence polarization plate reader (Tecan). Data were converted to amount of phosphopeptide present using a calibration curve generated by competition with the phosphopeptide competitor provided in the Tyrosine Kinase Assay Kit, Green (PanVera Corporation).

The results of the assay are shown in TABLE 6, below:

TABLE 6	
Compound No.	IC ₅₀ (in μ M)
R926505	0.0703
R926508	0.1315
R926594	0.7705
R926715	0.534
R926745	0.0925
R926782	0.1165
R926791	0.207
R926813	0.4047
R926816	0.0615
R935138	0.2288
R935190	0.0465

TABLE 6	
Compound No.	IC ₅₀ (in μ M)
R935191	0.045
R935193	0.075
R935194	0.1687
R935196	0.2655
R940255	0.7705
R940256	2.787
R940269	0.685
R940275	0.7335
R940276	0.1265
R940277	0.2143
R940290	0.187
R945071	0.4295
R945140	0.611
R945142	2.007
R945144	0.383
R921302	0.2678
R908702	0.0378
R908712	0.024
R909268	0.1253
R920410	0.157
R926753	0.108
R926757	0.5103
R926834	0.292
R926839	0.055
R926891	0.1695
R926931	0.2553
R935237	0.0455
R935293	0.0465
R935302	0.0265
R935304	0.042
R935307	0.057
R935309	0.098
R935310	0.2003
R940323	0.062
R940338	0.028

TABLE 6	
Compound No.	IC ₅₀ (in μ M)
R921303	0.00045
R940347	0.0345
R921304	0.01275
R950368	0.0107
R950373	0.0665


These data demonstrate that all of the compounds tested, except for R945142 and R909236 inhibit Syk kinase phosphorylation with IC₅₀s in the submicromolar range. All compounds tested inhibit Syk kinase phosphorylation with IC₅₀s in the micromolar range.

5 Although the foregoing invention has been described in some detail to facilitate understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims. Accordingly, the described embodiments are to be considered as illustrative and not restrictive, and the invention is not to be limited to the details given herein, but may be modified within the scope and equivalents of the appended
10 claims.

All literature and patent references cited throughout the application are incorporated by reference into the application for all purposes.

What is Claimed Is:

1. A compound according to structural formula (I):



and salts, hydrates, solvates, N-oxides and prodrugs thereof, wherein:

L^1 and L^2 are each, independently of one another, selected from the group consisting of a direct bond and a linker;

R^2 is selected from the group consisting of (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C3-C8) cycloalkyl optionally substituted with one or more of the same or different R^8 groups, cyclohexyl optionally substituted with one or more of the same or different R^8 groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R^8 groups, (C5-C15) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups;

R^4 is selected from the group consisting of hydrogen, (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C3-C8) cycloalkyl optionally substituted with one or more of the same or different R^8 groups, cyclohexyl optionally substituted with one or more of the same or different R^8 groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R^8 groups, (C5-C15) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups;

R^5 is selected from the group consisting of R^6 , (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C1-C4) alkanyl optionally substituted with one

or more of the same or different R^8 groups, (C2-C4) alkenyl optionally substituted with one or more of the same or different R^8 groups and (C2-C4) alkynyl optionally substituted with one or more of the same or different R^8 groups;

each R^6 is independently selected from the group consisting of hydrogen, an electronegative group, $-OR^d$, $-SR^d$, (C1-C3) haloalkyloxy, (C1-C3) perhaloalkyloxy, $-NR^cR^c$, halogen, (C1-C3) haloalkyl, (C1-C3) perhaloalkyl, $-CF_3$, $-CH_2CF_3$, $-CF_2CF_3$, $-CN$, $-NC$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $-N_3$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)NR^cR^c$, $-S(O)_2NR^cR^c$, $-OS(O)R^d$, $-OS(O)_2R^d$, $-OS(O)_2OR^d$, $-OS(O)NR^cR^c$, $-OS(O)_2NR^cR^c$, $-C(O)R^d$, $-C(O)OR^d$, $-C(O)NR^cR^c$, $-C(NH)NR^cR^c$, $-OC(O)R^d$, $-SC(O)R^d$, $-OC(O)OR^d$, $-SC(O)OR^d$, $-OC(O)NR^cR^c$, $-SC(O)NR^cR^c$, $-OC(NH)NR^cR^c$, $-SC(NH)NR^cR^c$, $-[NHC(O)]_nR^d$, $-[NHC(O)]_nOR^d$, $-[NHC(O)]_nNR^cR^c$ and $-[NHC(NH)]_nNR^cR^c$, (C5-C10) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups, (C6-C16) arylalkyl optionally substituted with one or more of the same or different R^8 groups, 5-10 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups and 6-16 membered heteroarylalkyl optionally substituted with one or more of the same or different R^8 groups;

R^8 is selected from the group consisting of R^a , R^b , R^a substituted with one or more of the same or different R^a or R^b , $-OR^a$ substituted with one or more of the same or different R^a or R^b , $-B(OR^a)_2$, $-B(NR^cR^c)_2$, $-(CH_2)_m-R^b$, $-(CHR^a)_m-R^b$, $-O-(CH_2)_m-R^b$, $-S-(CH_2)_m-R^b$, $-O-CHR^aR^b$, $-O-CR^a(R^b)_2$, $-O-(CHR^a)_m-R^b$, $-O-(CH_2)_m-CH[(CH_2)_mR^b]R^b$, $-S-(CHR^a)_m-R^b$, $-C(O)NH-(CH_2)_m-R^b$, $-C(O)NH-(CHR^a)_m-R^b$, $-O-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$, $-S-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$, $-O-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$, $-S-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$, $-NH-(CH_2)_m-R^b$, $-NH-(CHR^a)_m-R^b$, $-NH[(CH_2)_mR^b]$, $-N[(CH_2)_mR^b]_2$, $-NH-C(O)-NH-(CH_2)_m-R^b$, $-NH-C(O)-(CH_2)_m-CHR^bR^b$ and $-NH-(CH_2)_m-C(O)-NH-(CH_2)_m-R^b$;

each R^a is independently selected from the group consisting of hydrogen, (C1-C6) alkyl, (C3-C8) cycloalkyl, cyclohexyl, (C4-C11) cycloalkylalkyl, (C5-C10) aryl, phenyl, (C6-C16) arylalkyl, benzyl, 2-6 membered heteroalkyl, 3-8 membered cycloheteroalkyl, morpholinyl, piperazinyl, homopiperazinyl, piperidinyl, 4-11 membered cycloheteroalkylalkyl, 5-10 membered heteroaryl and 6-16 membered heteroarylalkyl;

each R^b is a suitable group independently selected from the group consisting of =O, -OR^d, (C1-C3) haloalkyloxy, -OCF₃, =S, -SR^d, =NR^d, =NOR^d, -NR^cR^c, halogen, -CF₃, -CN, -NC, -OCN, -SCN, -NO, -NO₂, =N₂, -N₃, -S(O)R^d, -S(O)₂R^d, -S(O)₂OR^d, -S(O)NR^cR^c, -S(O)₂NR^cR^c, -OS(O)R^d, -OS(O)₂R^d, -OS(O)₂OR^d, -OS(O)₂NR^cR^c, -C(O)R^d, -C(O)OR^d, -C(O)NR^cR^c, -C(NH)NR^cR^c, -C(NR^a)NR^cR^c, -C(NOH)R^a, -C(NOH)NR^cR^c, -OC(O)R^d, -OC(O)OR^d, -OC(O)NR^cR^c, -OC(NH)NR^cR^c, -OC(NR^a)NR^cR^c, -[NHC(O)]_nR^d, -[NR^aC(O)]_nR^d, -[NHC(O)]_nOR^d, -[NR^aC(O)]_nOR^d, -[NHC(O)]_nNR^cR^c, -[NR^aC(O)]_nNR^cR^c, -[NHC(NH)]_nNR^cR^c and -[NR^aC(NR^a)]_nNR^cR^c;

each R^c is independently a protecting group or R^a, or, alternatively, each R^c is taken together with the nitrogen atom to which it is bonded to form a 5 to 8-membered cycloheteroalkyl or heteroaryl which may optionally include one or more of the same or different additional heteroatoms and which may optionally be substituted with one or more of the same or different R^a or suitable R^b groups;

each R^d is independently a protecting group or R^a;

each m is independently an integer from 1 to 3; and


each n is independently an integer from 0 to 3,

with the provisos that:

- (1) when L¹ is a direct bond and R⁶ is hydrogen, then R² is not 3,4,5-tri (C1-C6) alkoxyphenyl;
- (2) when L¹ and L² are each a direct bond, R² is a substituted phenyl and R⁶ is hydrogen, then R⁵ is other than cyano or -C(O)NHR, where R is hydrogen or (C1-C6) alkyl;
- (3) when L¹ and L² are each a direct bond and R² and R⁴ are each independently a substituted or unsubstituted pyrrole or indole, then the R² and R⁴ are attached to the remainder of the molecule *via* a ring carbon atom; and
- (4) the compound is not
 - N2,N4-bis(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R070790);
 - N2,N4-bis(2-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R081166);
 - N2,N4-bis(4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R088814);
 - N2,N4-bis(2-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R088815);
 - N2,N4-bisphenyl-5-fluoro-2,4-pyrimidinediamine (R091880);

N2,N4-bis(3-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R092788);
 N2,N4-bis(3-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R067962);
 N2,N4-bis(2,5-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067963);
 N2,N4-bis(3,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067964);
 N2,N4-bis(4-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R070153);
 N2,N4-bis(2,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R070791);
 N2,N4-bis(3-bromophenyl)-5-fluoro-2,4-pyrimidinediamine (R008958);
 N2,N4-bis(phenyl)-5-fluoro-2,4-pyrimidinediamine;
 N2,N4-bis(morpholino)-5-fluoro-2,4-pyrimidinediamine;
 N2,N4-bis[(3-chloro-4-methoxyphenyl)]-5-fluoro-2,4-pyrimidinediamine; N2-(3,4,5-trimethoxyphenyl)-N4-(3,5-dimethoxyphenyl)-2,4-pyrimidinediamine; N2-(3,4-dimethoxyphenyl)-N4-(3-methoxyphenyl)-2,4-pyrimidinediamine; N2,N4-bis(3,4-dimethoxyphenyl)-2,4-pyrimidinediamine; N2,N4-Bis(3-chloro-4-methoxy-5-fluoro-2,4-pyrimidinediamine; N2-[4-(3-dimethylamino-2-hydroxy-propyloxy)phenyl]-N4-3,4-dichlorophenyl-5-chloro-2,4-pyrimidinediamine; N2-[4-(3-dimethylamino-2-hydroxy-propyloxy)phenyl]-N4-3,4-dichlorophenyl-2,4-pyrimidinediamine; N2-[4-(3-dimethylamino-2-hydroxy-propyloxy)phenyl]-N4-3,4-dichlorophenyl-5-methyl-2,4-pyrimidinediamine; N2-(4-benzoxoxyphenyl)-N4-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine; N2,N4-bis(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine; N2,N4-bis(3,4,5-trimethoxyphenyl)-5-bromo-2,4-pyrimidinediamine; N2-(1-benzyl-1H-indazol-5-yl)-N4-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine; N2-(1H-indol-1-yl)-N4-(3,4,5-trimethoxyphenyl)-2,4-pyrimidinediamine; N2-[(2-methoxy-5-aminosulfonyl)phenyl]-N4-(3-methyl-1H-indazol-6-yl)-2,4-pyrimidinediamine; N2-[2-methoxy-5-(5-methyl-3-isoxazoly-methylsulfonyl)phenyl]-N4-(3-methyl-1H-indazol-6-yl)-2,4-pyrimidinediamine; N2-[(2-methoxy-5-methylaminosulfonyl)phenyl]-N4-(3-methyl-1H-indazol-6-yl)-2,4-pyrimidinediamine; N2-[(2-methoxy-5-ethylsulfonyl)phenyl]-N4-(3-methyl-1H-indazol-6-yl)-2,4-pyrimidinediamine; N2-[(2-methoxy-5-isobutylsulfonyl)phenyl]-N4-(3-methyl-1H-indazol-6-yl)-2,4-pyrimidinediamine; N2-[(2-methoxy-5-propylcarbonyl)phenyl]-N4-(3-methyl-1H-indazol-6-yl)-2,4-pyrimidinediamine; N2-(1H-indazol-5-yl)-N4-propynyl-5-bromo-2,4-pyrimidinediamine; N2-(1-H-indol-5-yl)-N4-[1-(3-methyl-1-hydroxy)butyl]-N4-(1H-indol-5-yl)-5-bromo-2,4-pyrimidinediamine; N2-(1-dimethylaminosulfonyl-1H-indol-5-yl)-N4-[1-(2-

methyl-2-hydroxy)ethyl]-5-bromo-2,4-pyrimidinediamine, or a compound according to the formula:



wherein: R^e is (C1-C6) alkyl; R^f and R^g are each, independently of one another, a straight-chain or branched (C1-C6) alkyl which is optionally substituted with one or more of the same or different R⁸ groups; and R⁸ is as defined above.


2. The compound of Claim 1 in which L¹ and L² are each, independently of one another, selected from the group consisting of a direct bond, (C1-C3) alkylidyl optionally substituted with one or more of the same or different R⁹ groups and 1-3 membered heteroalkylidyl optionally substituted with one or more of the same or different R⁹ groups, wherein:

R⁹ is selected from the group consisting of (C1-C3) alkyl, -OR^a, -C(O)OR^a, (C5-C10) aryl optionally substituted with one or more of the same or different halogens, phenyl optionally substituted with one or more of the same or different halogens, 5-10 membered heteroaryl optionally substituted with one or more of the same or different halogens and 6 membered heteroaryl optionally substituted with one or more of the same or different halogens; and

R^a is as defined in Claim 1.

3. The compound of Claim 2 in which L¹ and L² are each, independently of one another, selected from the group consisting of methano, ethano and propano, each of which may be optionally monosubstituted with an R⁹ group.


4. The compound of Claim 3 in which the R^9 group is selected from the group consisting of $-OR^a$, $-C(O)OR^a$, halophenyl and 4-halophenyl, wherein R^a is as defined in Claim 1.
5. The compound of Claim 1 in which R^6 is hydrogen.
6. The compound of Claim 1 or 5 in which R^5 is selected from the group consisting of an electronegative group, halo, $-F$, $-CN$, $-NO_2$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)CF_3$, $-C(O)OCF_3$, (C1-C3) haloalkyl, (C1-C3) perhaloalkyl (C1-C3) haloalkoxy, (C1-C3) perhaloalkoxy, $-OCF_3$ and $-CF_3$.
7. The compound of Claim 1 in which at least one of L1 or L2 is a direct bond.
8. The compound of Claim 1 according to the structure (Ia):




and salts, hydrates, solvates, N-oxides and prodrugs thereof, wherein R^2 , R^4 , R^5 and R^6 are as defined in Claim 1.

9. The compound of Claim 8 in which R^2 is selected from the group consisting of phenyl, naphthyl, 5-10 membered heteroaryl, benzodioxanyl, 1,4-benzodioxan-(5 or 6)-yl, benzodioxolyl, 1,3-benzodioxol-(4 or 5)-yl, benzoxazinyl, 1,4-benzoxazin-(5,6,7 or 8)-yl, benzoxazolyl, 1,3-benzoxazol-(4,5,6 or 7)-yl, benzopyranyl, benzopyran-(5,6,7 or 8)-yl, benzotriazolyl, benzotrazol-(4,5,6 or 7)-yl, 1,4-benzoxazinyl-2-one, 1,4-benzoxazin-(5,6,7 or 8)-yl-2-one, 2H-1,4-benzoxazinyl-3(4H)-one, 2H-1,4-benzoxazin-(5,6,7 or 8)-yl-3(4H)-one, 2H-1,3-benzoxazinyl-2,4(3H)-dione, 2H-1,3-benzoxazin-(5,6,7 or 8)-yl-2,4(3H)-dione, benzoxazolyl-2-one, benzoxazol-(4,5,6 or 7)-yl-2-one, dihydrocoumarinyl, dihydrocoumarin-(5,6,7 or 8)-yl, 1,2-benzopyronyl, 1,2-benzopyron-(5,6,7 or 8)-yl, benzofuranyl, benzofuran-(4,5,6 or 7)-yl, benzo[b]furanyl, benzo[b]furan-(4,5,6 or 7)-yl, indolyl, indol-(4,5,6 or 7)-yl, pyrrolyl and pyrrol-(1 or 2)-yl, each of which may be optionally substituted with one or more of the same or different R^8 groups, where R^8 is as defined in Claim 1.

10. The compound of Claim 8 in which R^2 and/or R^4 are each, independently of one another, a heteroaryl selected from the group consisting of:





wherein:

p is an integer from one to three;

each --- independently represents a single bond or a double bond;

R^{35} is hydrogen or R^8 , where R^8 is as previously defined for structural formula

(I);

X is selected from the group consisting of CH, N and N-O;

each Y is independently selected from the group consisting of O, S and NH;

each Y^1 is independently selected from the group consisting of O, S, SO, SO₂, SONR³⁶, NH and NR³⁷;

each Y^2 is independently selected from the group consisting of CH, CH₂, O, S, N, NH and NR³⁷;

R³⁶ is hydrogen or alkyl;

R³⁷ is selected from the group consisting of hydrogen and a progroup, preferably hydrogen or a progroup selected from the group consisting of aryl, arylalkyl, heteroaryl, R^a, R^b-CR^aR^b-O-C(O)R⁸, -CR^aR^b-O-PO(OR⁸)₂, -CH₂-O-PO(OR⁸)₂, -CH₂-PO(OR⁸)₂, -C(O)-CR^aR^b-N(CH₃)₂, -CR^aR^b-O-C(O)-CR^aR^b-N(CH₃)₂, -C(O)R⁸, -C(O)CF₃ and -C(O)-NR⁸-C(O)R⁸;

R³⁸ is selected from the group consisting of alkyl and aryl;

A is selected from the group consisting of O, NH and NR³⁸;

R⁹, R¹⁰, R¹¹ and R¹² are each, independently of one another, selected from the group consisting of alkyl, alkoxy, halogen, haloalkoxy, aminoalkyl and hydroxyalkyl, or, alternatively, R⁹ and R¹⁰ and/or R¹¹ and R¹² are taken together form a ketal;

each Z is selected from the group consisting of hydroxyl, alkoxy, aryloxy, ester, carbamate and sulfonyl;

Q is selected from the group consisting of -OH, OR⁸, -NR^cR^c, -NHR³⁹-C(O)R⁸, -NHR³⁹-C(O)OR⁸, -NR³⁹-CHR⁴⁰-R^b, -NR³⁹-(CH₂)_m-R^b and -NR³⁹-C(O)-CHR⁴⁰-NR^cR^c;

R³⁹ and R⁴⁰ are each, independently of one another, selected from the group consisting of hydrogen, alkyl, aryl, alkylaryl; arylalkyl and NHR⁸; and

R^a, R^b and R^c are as previously defined for structural formula (I).

11. The compound of Claim 10 in which R² and R⁴ are the same.

12. The compound of Claim 10 or 11 in which each R³⁵ is independently selected from the group consisting of hydrogen, R^d, -NR^cR^c, -(CH₂)_m-NR^cR^c, -C(O)NR^cR^c, -(CH₂)_m-C(O)NR^cR^c, -C(O)OR^d, -(CH₂)_m-C(O)OR^d and -(CH₂)_m-OR^d, where *m*, R^c and R^d are as defined in Claim 1.

13. The compound of Claim 12 in which each *m* is one.

14. The compound of Claim 8 in which R^2 is an optionally substituted heteroaryl which is attached to the remainder of the molecule *via* a ring carbon atom.

15. The compound of Claim 8 in which R^4 is an optionally substituted heteroaryl which is attached to the remainder of the molecule *via* a ring carbon atom.

16. The compound of Claim 8 in which R^2 and/or R^4 are each, independently of one another, a phenyl optionally substituted with one, two or three R^8 groups, where R^8 is as defined in Claim 1.

17. The compound of Claim 16 in which R^2 and R^4 are each the same or different optionally substituted phenyl.

18. The compound of Claim 16 or 17 in which the optionally substituted phenyl is *mono* substituted.

19. The compound of Claim 18 in which the R^8 substituent is at the *ortho*, *meta* or *para* position.

20. The compound of Claim 19 in which R^8 is selected from the group consisting of (C1-C10) alkyl, (C1-C10) branched alkyl, $-OR^d$, $-O-(CH_2)_m-NR^cR^c$, $-O-C(O)NR^cR^c$, $-O-(CH_2)_m-C(O)NR^cR^c$, $-O-C(O)OR^a$, $-O-(CH_2)_m-C(O)OR^a$, $-O-C(NH)NR^cR^c$, $-O-(CH_2)_m-C(NH)NR^cR^c$, $-NH-(CH_2)_m-NR^cR^c$, $-NH-C(O)NR^cR^c$ and $-NH-(CH_2)_m-C(O)NR^cR^c$, where m , R^a , R^c and R^d are as defined in Claim 1.

21. The compound of Claim 16 or 17 in which the optionally substituted phenyl is a disubstituted phenyl.

22. The compound of Claim 21 in which the R^8 substituents are positioned 2,3-; 2,4-; 2,5-; 2,6-; 3,4-; or 3,5-.

23. The compound of Claim 21 in which each R^8 is independently selected from the group consisting of (C1-C10) alkyl, (C1-C10) branched alkyl, $-OR^a$ optionally substituted with


one or more of the same or different R^a or R^b groups, $-O-(CH_2)_m-NR^cR^c$, $-O-C(O)NR^cR^c$, $-O-(CH_2)_m-C(O)NR^cR^c$, $-O-C(O)OR^a$, $-O-(CH_2)_m-C(O)OR^a$, $-O-C(NH)NR^cR^c$, $-O-(CH_2)_m-C(NH)NR^cR^c$, $-NH-(CH_2)_m-NR^cR^c$, $-NH-C(O)NR^cR^c$ and $-NH-(CH_2)_m-C(O)NR^cR^c$, where m , R^a , R^b and R^c are as defined in Claim 1.

24. The compound of Claim 16 or 17 in which the optionally substituted phenyl is trisubstituted.

25. The compound of Claim 24 in which the R^8 substituents are positioned 2,3,4; 2,3,5; 2,3,6; 2,4,5; 2,4,6; 2,5,6; or 3,4,5.

26. The compound of Claim 25 which each R^8 is independently selected from the group consisting of (C1-C10) alkyl, (C1-C10) branched alkyl, $-OR^a$ optionally substituted with one or more of the same or different R^a or R^b groups, $-O-(CH_2)_m-NR^cR^c$, $-O-C(O)NR^cR^c$, $-O-(CH_2)_m-C(O)NR^cR^c$, $-O-C(O)OR^a$, $-O-C(NH)NR^cR^c$, $-O-(CH_2)_m-C(O)OR^a$, $-O-(CH_2)_m-C(NH)NR^cR^c$, $-NH-(CH_2)_m-NR^cR^c$, $-NH-C(O)NR^cR^c$ and $-NH-(CH_2)_m-C(O)NR^cR^c$, where m , R^a , R^b and R^c are as defined in Claim 1.


27. The compound of Claim 24 in which the trisubstituted phenyl has the formula:



wherein: R^{31} is methyl or (C1-C6) alkyl; R^{32} is hydrogen, methyl or (C1-C6) alkyl; and R^{33} is a halo group.

28. The compound of Claim 17 in which R^2 and R^4 are the same.

29. The compound of Claim 8 according to structural formula (Ib):




and salts, hydrates, solvates and N-oxides thereof, wherein R^{11} , R^{12} , R^{13} and R^{14} are each, independently of one another, selected from the group consisting of hydrogen, hydroxy, (C1-C6) alkoxy and $-NR^cR^c$; and R^5 , R^6 and R^c are as defined in Claim 1.

30. The compound of Claim 29 in which R^{11} , R^{12} , R^{13} and R^{14} are each hydrogen.

31. The compound of Claim 29 in which R^{12} and R^{13} are each hydrogen.

32. The compound of Claim 8 according to structural formula (Ic):



and salts, hydrates, solvates, N-oxides and prodrugs thereof, wherein:

R^4 is phenyl optionally substituted with from 1 to 3 of the same or different R^8 groups or 5-14 membered heteroaryl optionally substituted with from 1 to 4 of the same or different R^8 groups;

R^5 is an electronegative group, F or CF_3 ; and

R^{18} is $-O(CH_2)_m-R^b$, where m and R^b are as defined in Claim 1.


33. The compound of Claim 32 in which R^4 is an optionally substituted heteroaryl.

34. The compound of Claim 32 in which R¹⁸ is -O-CH₂-C(O)-NHCH₃.

35. The compound of Claim 1 which inhibits mast cell or basophil cell degranulation as measured in an *in vitro* assay.

36. The compound of Claim 35 which has an IC₅₀ of about 20 μM or less.

37. A compound according to structural formula (Id):



and salts, hydrates, solvates, N-oxides and prodrugs thereof, wherein:

R² and R⁴ are as defined in Claim 1; and

R¹⁵ is an electronegative group,

with the provisos that:

(1) when R² is 3,4,5-tri (C1-C6) alkoxyphenyl and R¹⁵ is halogen, then R⁴ is not 3,4,5-tri (C1-C6) alkoxyphenyl;

(2) when R² is a substituted phenyl group, then R¹⁵ is other than cyano or -C(O)NHR, where R is hydrogen or (C1-C6) alkyl; and

(3) the compound is not

N2,N4-bis(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R070790);

N2,N4-bis(2-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R081166);

N2,N4-bis(4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R088814);

N2,N4-bis(2-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R088815);

N2,N4-bisphenyl-5-fluoro-2,4-pyrimidinediamine (R091880);

N2,N4-bis(3-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R092788);

N2,N4-bis(3-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R067962);

N2,N4-bis(2,5-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067963);

N2,N4-bis(3,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067964);

N2,N4-bis(4-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R0707153);
N2,N4-bis(2,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R070791);
N2,N4-bis(3-bromophenyl)-5-fluoro-2,4-pyrimidinediamine (R008958);
N2,N4-bis(phenyl)-5-fluoro-2,4-pyrimidinediamine;
N2,N4-bis(morpholino)-5-fluoro-2,4-pyrimidinediamine; or
N2,N4-bis[(3-chloro-4-methoxyphenyl)]-5-fluoro-2,4-pyrimidinediamine.

38. The compound of Claim 37 in which when R¹⁵ is halogen or nitro, then R² is not 3,4,5-tri (C1-C6) alkoxyphenyl.

39. The compound of Claim 38 in which R¹⁵ is selected from the group consisting of –CN, –NC, –NO₂, halogen, –F, (C1-C3) haloalkyl, (C1-C3) perhaloalkyl, (C1-C3) fluoroalkyl, (C1-C3) perfluoroalkyl, –CF₃, (C1-C3) haloalkoxy, (C1-C3) perhaloalkoxy, (C1-C3) fluoroalkoxy, (C1-C3) perfluoroalkoxy and –OCF₃.

40. The compound of Claim 39 in which R¹⁵ is selected from the group consisting of halo, Br, F, –CF₃ and –NO₂.

41. The compound of Claim 37 which inhibits mast cell or basophil cell degranulation as measured in an *in vitro* assay.

42. The compound of Claim 41 which has an IC₅₀ of about 20 μM or less.


43. A compound selected from any compound in TABLE 1 which inhibits mast cell or basophil cell degranulation as measured in an *in vitro* assay, with the proviso that the compound is not

N2,N4-bis(4-ethoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R070790);
N2,N4-bis(2-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R081166);
N2,N4-bis(4-methoxyphenyl)-5-fluoro-2,4-pyrimidinediamine (R088814);
N2,N4-bis(2-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R088815);
N2,N4-bisphenyl-5-fluoro-2,4-pyrimidinediamine (R091880);

N2,N4-bis(3-methylphenyl)-5-fluoro-2,4-pyrimidinediamine (R092788);
 N2,N4-bis(3-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R067962);
 N2,N4-bis(2,5-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067963);
 N2,N4-bis(3,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R067964);
 N2,N4-bis(4-chlorophenyl)-5-fluoro-2,4-pyrimidinediamine (R070153);
 N2,N4-bis(2,4-dimethylphenyl)-5-fluoro-2,4-pyrimidinediamine (R070791); or
 N2,N4-bis(3-bromophenyl)-5-fluoro-2,4-pyrimidinediamine (R008958).

44. The compound of Claim 43 which has an IC_{50} of about 20 μM or less.

45. A pharmaceutical composition comprising a pyrimidinediamine compound and a pharmaceutically acceptable excipient, carrier or diluent, said pyrimidinediamine compound being a compound according to structural formula (I):



and salts, hydrates, solvates, N-oxides and prodrugs thereof, wherein:

L^1 and L^2 are each, independently of one another, selected from the group consisting of a direct bond and a linker;

R^2 is selected from the group consisting of (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C3-C8) cycloalkyl optionally substituted with one or more of the same or different R^8 groups, cyclohexyl optionally substituted with one or more of the same or different R^8 groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R^8 groups, (C5-C15) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more

of the same or different R^8 groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups;

R^4 is selected from the group consisting of hydrogen, (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C3-C8) cycloalkyl optionally substituted with one or more of the same or different R^8 groups, cyclohexyl optionally substituted with one or more of the same or different R^8 groups, 3-8 membered cycloheteroalkyl optionally substituted with one or more of the same or different R^8 groups, (C5-C15) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups and 5-15 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups;

R^5 is selected from the group consisting of R^6 , (C1-C6) alkyl optionally substituted with one or more of the same or different R^8 groups, (C1-C4) alkanyl optionally substituted with one or more of the same or different R^8 groups, (C2-C4) alkenyl optionally substituted with one or more of the same or different R^8 groups and (C2-C4) alkynyl optionally substituted with one or more of the same or different R^8 groups;

each R^6 is independently selected from the group consisting of hydrogen, an electronegative group, $-OR^d$, $-SR^d$, (C1-C3) haloalkyloxy, (C1-C3) perhaloalkyloxy, $-NR^cR^c$, halogen, (C1-C3) haloalkyl, (C1-C3) perhaloalkyl, $-CF_3$, $-CH_2CF_3$, $-CF_2CF_3$, $-CN$, $-NC$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $-N_3$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)NR^cR^c$, $-S(O)_2NR^cR^c$, $-OS(O)R^d$, $-OS(O)_2R^d$, $-OS(O)_2OR^d$, $-OS(O)NR^cR^c$, $-OS(O)_2NR^cR^c$, $-C(O)R^d$, $-C(O)OR^d$, $-C(O)NR^cR^c$, $-C(NH)NR^cR^c$, $-OC(O)R^d$, $-SC(O)R^d$, $-OC(O)OR^d$, $-SC(O)OR^d$, $-OC(O)NR^cR^c$, $-SC(O)NR^cR^c$, $-OC(NH)NR^cR^c$, $-SC(NH)NR^cR^c$, $-[NHC(O)]_nR^d$, $-[NHC(O)]_nOR^d$, $-[NHC(O)]_nNR^cR^c$ and $-[NHC(NH)]_nNR^cR^c$, (C5-C10) aryl optionally substituted with one or more of the same or different R^8 groups, phenyl optionally substituted with one or more of the same or different R^8 groups, (C6-C16) arylalkyl optionally substituted with one or more of the same or different R^8 groups, 5-10 membered heteroaryl optionally substituted with one or more of the same or different R^8 groups and 6-16 membered heteroarylalkyl optionally substituted with one or more of the same or different R^8 groups;

R^8 is selected from the group consisting of R^a , R^b , R^a substituted with one or more of the same or different R^a or R^b , $-OR^a$ substituted with one or more of the same or different R^a or

R^b , $-B(OR^a)_2$, $-B(NR^cR^c)_2$, $-(CH_2)_m-R^b$, $-(CHR^a)_m-R^b$, $-O-(CH_2)_m-R^b$, $-S-(CH_2)_m-R^b$,
 $-O-CHR^aR^b$, $-O-CR^a(R^b)_2$, $-O-(CHR^a)_m-R^b$, $-O-(CH_2)_m-CH[(CH_2)_mR^b]R^b$, $-S-(CHR^a)_m-R^b$,
 $-C(O)NH-(CH_2)_m-R^b$, $-C(O)NH-(CHR^a)_m-R^b$, $-O-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$,
 $-S-(CH_2)_m-C(O)NH-(CH_2)_m-R^b$, $-O-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$,
 $-S-(CHR^a)_m-C(O)NH-(CHR^a)_m-R^b$, $-NH-(CH_2)_m-R^b$, $-NH-(CHR^a)_m-R^b$, $-NH[(CH_2)_mR^b]$,
 $-N[(CH_2)_mR^b]_2$, $-NH-C(O)-NH-(CH_2)_m-R^b$, $-NH-C(O)-(CH_2)_m-CHR^bR^b$ and
 $-NH-(CH_2)_m-C(O)-NH-(CH_2)_m-R^b$;

each R^a is independently selected from the group consisting of hydrogen, (C1-C6) alkyl, (C3-C8) cycloalkyl, cyclohexyl, (C4-C11) cycloalkylalkyl, (C5-C10) aryl, phenyl, (C6-C16) arylalkyl, benzyl, 2-6 membered heteroalkyl, 3-8 membered cycloheteroalkyl, morpholinyl, piperazinyl, homopiperazinyl, piperidinyl, 4-11 membered cycloheteroalkylalkyl, 5-10 membered heteroaryl and 6-16 membered heteroarylalkyl;

each R^b is a suitable group independently selected from the group consisting of $=O$,
 $-OR^d$, (C1-C3) haloalkyloxy, $-OCF_3$, $=S$, $-SR^d$, $=NR^d$, $=NOR^d$, $-NR^cR^c$, halogen, $-CF_3$, $-CN$,
 $-NC$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $=N_2$, $-N_3$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)NR^cR^c$,
 $-S(O)_2NR^cR^c$, $-OS(O)R^d$, $-OS(O)_2R^d$, $-OS(O)_2OR^d$, $-OS(O)_2NR^cR^c$, $-C(O)R^d$, $-C(O)OR^d$,
 $-C(O)NR^cR^c$, $-C(NH)NR^cR^c$, $-C(NR^a)NR^cR^c$, $-C(NOH)R^a$, $-C(NOH)NR^cR^c$, $-OC(O)R^d$,
 $-OC(O)OR^d$, $-OC(O)NR^cR^c$, $-OC(NH)NR^cR^c$, $-OC(NR^a)NR^cR^c$, $-[NHC(O)]_nR^d$,
 $-[NR^aC(O)]_nR^d$, $-[NHC(O)]_nOR^d$, $-[NR^aC(O)]_nOR^d$, $-[NHC(O)]_nNR^cR^c$, $-[NR^aC(O)]_nNR^cR^c$,
 $-[NHC(NH)]_nNR^cR^c$ and $-[NR^aC(NR^a)]_nNR^cR^c$;

each R^c is independently R^a or, alternatively, each R^c is taken together with the nitrogen atom to which it is bonded to form a 5 to 8-membered cycloheteroalkyl or heteroaryl which may optionally include one or more of the same or different additional heteroatoms and which may optionally be substituted with one or more of the same or different R^a or suitable R^b groups;

each R^d is independently R^a ;

each m is independently an integer from 1 to 3; and


each n is independently an integer from 0 to 3, with the provisos that:

(1) when L^1 is a direct bond and R^6 is hydrogen, then R^2 is not 3,4,5-tri (C1-C6) alkoxyphenyl;

(2) when L^1 and L^2 are each direct bonds, R^2 is a substituted phenyl and R^6 is hydrogen, then R^5 is other than cyano or $-C(O)NHR$, where R is hydrogen or (C1-C6) alkyl;


(3) when R^2 and R^4 are each independently a substituted or unsubstituted pyrrole or indole, then R^2 and R^4 are attached to the remainder of the molecule *via* a ring carbon atom; and

(4) the compound is not:



wherein: R^e is (C1-C6) alkyl; R^f and R^g are each, independently of one another a straight-chain or branched (C1-C6) alkyl which is optionally substituted with one or more of the same or different R^8 groups; and R^8 is as defined above.

46. A pharmaceutical composition comprising a pyrimidinediamine compound and a pharmaceutically acceptable carrier, diluent or excipient, said pyrimidinediamine compound being a compound according to structural formula (Id):



and salts, hydrates, solvates, N-oxides and prodrugs thereof, wherein:

R^2 and R^4 are as defined for Claim 1; and

R^{15} is an electronegative group, with the provisos that:

(1) when R^2 is 3,4,5-tri (C1-C6) alkoxyphenyl and R^{15} is halogen, then R^4 is not 3,4,5-tri (C1-C6) alkoxyphenyl; and

(2) when R^2 is a substituted phenyl, then R^{15} is other than cyano or $-C(O)NHR$, where R is hydrogen or (C1-C6) alkyl.

47. A pharmaceutical composition comprising a compound according to Claim 1 and a pharmaceutically acceptable carrier, diluent or excipient.

48. A pharmaceutical composition comprising a compound selected from any compound in TABLE 1 which inhibits mast cell or basophil cell degranulation as measured in an *in vitro* assay and a pharmaceutically acceptable carrier, diluent or excipient.

49. The composition of any one of Claims 46-48 in which the compound is in the form of a pharmaceutically acceptable salt.

50. The composition of Claim 49 in which the salt is a hydrochloride salt, a hydrogen sulfate salt, a sulfate salt, a phosphate salt, an alkane sulfonate salt, a methane sulfonate salt, an ethane sulfonate salt or a *p*-toluene sulfonate salt.

51. A method of inhibiting cell degranulation, comprising contacting a cell with an amount of a compound according to any one of Claims 1, 37 or 43 effect to inhibit degranulation.

52. The method of Claim 51 in which the cell is a human mast, basophil cell, neutrophil or eosinophil cell.

53. A method of inhibiting cell degranulation, comprising contacting a mast or basophil cell with an amount of a composition according to any one of Claims 46-48 effective to inhibit degranulation.

54. The method of Claim 53 in which the cell is a human mast, basophil cell, neutrophil or eosinophil cell.

55. A method of treating a disease characterized by, caused by or associated with mast or basophil cell degranulation, comprising administering to an animal suffering from such a disease an effective amount of a composition according to any one of Claims 46-48.

56. The method of Claim 55 in which the animal is a human.

57. The method of Claim 55 in which the disease is selected from the group consisting of allergic diseases, low grade scarring, diseases associated with tissue destruction, diseases associated with tissue inflammation, inflammation, and scarring.

58. The method of Claim 57 in which the allergic disease is selected from the group consisting of conjunctivitis, rhinitis, asthma, atopic dermatitis and food allergies.

59. The method of Claim 57 in which the low grade scarring is selected from the group consisting of scleroderma, increased fibrosis, keloids, post-surgical scars, pulmonary fibrosis, vascular spasms, migraine, reperfusion injury and post myocardial infarction.

60. The method of Claim 57 in which the disease associated with tissue destruction is selected from the group consisting of COPD, cardiobronchitis and post myocardial infarction.

61. The method of Claim 57 in which the disease associated with tissue inflammation is selected from the group consisting of irritable bowel, spastic colon and inflammatory colon disease.

62. A method of inhibiting a Syk kinase, comprising the step of contacting the Syk kinase or an active fragment thereof with an effective amount of a 2,4-pyrimidinediamine compound according to Claim 1.

63. The method of Claim 62 which is practiced *in vitro* with an isolated or recombinant Syk kinase.

64. The method of Claim 62 in which the Syk kinase is practiced *in vitro* with a cell or cell population that expresses an endogenous or recombinant Syk kinase.

65. The method of Claim 62 which is practiced *in vivo*.

66. A method of inhibiting a Syk kinase in an animal, comprising the step of administering to the animal an amount of a 2,4-pyrimidinediamine compound according to Claim 1 effective to inhibit the Syk kinase.

67. A method of treating or preventing a disease mediated at least in part by Syk kinase activity, comprising the step of administering to an animal in need thereof an amount of a 2,4-pyrimidinediamine compound according to Claim 1 effective to inhibit the Syk kinase activity, thereby treating or preventing the disease.

68. A method of treating or preventing a disease mediated at least in part by Syk kinase activity, comprising the step of administering to an animal in need thereof an amount of a composition according to Claim 46 effective to inhibit the Syk kinase activity, thereby treating or preventing the disease.

69. The method of Claim 72 or 74 in which the animal is a human.

70. A method of inhibiting an Fc receptor signal transduction cascade, contacting a cell comprising an Fc receptor having a gamma homodimer with an amount of a 2,4-pyrimidinediamine compound according to Claim 1 effective to inhibit its signal transduction cascade.

71. The method of Claim 70 in which the Fc receptor is selected from the group consisting of Fc α RI, FC γ RI, FC γ RIII and Fc ϵ RI.