

Description

Asparagine plays an important role as a transport form of nitrogen and in many plants - including nitrogen fixers - it is the principle compound involved in the transfer of nitrogen from the roots into the transpiration stream. In plants asparagine is formed from glutamine, aspartate and ATP catalyzed by the asparagine synthetase ASN (E.C. 6.3.5.4) whereby glutamate, AMP and pyrophosphate are formed as by-products.

It now has been found that the prokaryotic, ammonium dependent asparagine synthetase ASN-A (E. C. 6.3.1.1) can be introduced into plant cells, resulting in transgenic plants which show a number of advantages: The plants show a more efficient net photosynthetic CO₂-fixation, an increased growth rate, accelerated plant development, earlier flower formation, increased green mass and plant dry weight. Thus, the growth area can be extended to regions with a less favourable climate and in regions with a warm climate e. g. three instead of two crops are possible. Furthermore, the transgenic plants tolerate the application of glutamine synthetase (GS) inhibitors, e. g. phosphinothricine (PPT) or methionine sulfoximine (MSX) and even show a stimulation of photosynthesis and growth upon application of such inhibitors.

In contrast to the ASN encoding gene (or genes) of higher plants (asn gene) the asnA gene of *E. coli* codes for a different type of ASN which uses ammonium rather than glutamine for the production of asparagine (Cedar and Schwartz (1969) *J. Biol. Chem.* 244, 4112-4121). The gene for this enzyme has been isolated and characterized by Nakamura et al. (1981) *Nucleic Acids Research* 9, 4669-4676. It was found that this prokaryotic enzyme is active in plants and it opens a new ammonium assimilation pathway in these transgenic plants. It results in an overall change of the plant nitrogen metabolism with a stimulatory effect on growth and green mass production.

Under normal conditions in the transformants both GS and ASH-A use ammonia. The advantage of the bacterial pathway will be more pronounced during darkness when the chloroplast GS activity is limited by the reduced availability of ATP, energy charge and magnesium ions (O'Neal and Joy (1974) *Plant Phys.* 54, 773-779; Joy (1988) *Can. J. Bot.* 66, 2103-2109). Moreover, the expression of the bacterial asn-A gene in plants even allows an assimilation of ammonia when the plant GS is blocked by specific inhibitors like PPT. In non-transformed plants the inhibition of GS activity disturbs the main road of ammonia utilization and the accumulation of ammonium is one of the key factors in lethality of the treated plants (Tachibana et al. (1986) *J. Pesticide Sci.* 11, 33-37). Thus the presence of the bacterial enzyme can reduce the ammonia accumulation in PPT-treated transgenic plants which so not only can survive doses of herbicides which are lethal for wild-type plants but rather the so treated transgenic plant shows growth stimulation.

It is apparent for a skilled person that these positive effects are not limited to the asnA gene from *E. coli* since other bacteria contain the same gene or a gene having the same capability of amidating asparaginic acid and its salts to yield asparagine.

Thus the invention relates to the use of a prokaryotic asnA gene in a plant cell, a gene construct comprising a gene encoding a prokaryotic asnA, operatively linked to a regulatory sequence which effects the expression of the said gene in a plant cell, a vector containing such a gene construct, a plant cell transformed with such a gene construct or vector and expressing a prokaryotic ammonia specific asparagine synthetase in a plant, especially a crop plant, and seeds or propagation material of such plants which contain transformed cells as hereinbefore defined.

Preferred embodiments comprise the use of the *E. coli* asnA gene encoding the said enzyme and synthetic genes encoding the said enzyme, especially genes comprising codons which are preferably used by plants. The invention also comprises genes which encode enzymes having a different amino acid composition than the natural enzymes but with essentially the same catalytic activity by deleting or adding codons or by replacing codons in the natural genes by such which encode a different amino acid. All such modifications are within the ordinary skill of persons involved in this art.

Example I

Expression of the *E. coli* asnA gene with the RUBISCO small subunit promoter in tobacco

1. Production of transgenic tobacco plants

Based on the complete nucleotide sequence of the asnA gene from *E. coli* (Nakamura et al., (1981) *Nucleic Acids Research* 18, 4673, Fig. 3) we recloned the PstI-HgaI fragment from the plasmid pMY114 into pUC9. Then the asnA gene (1.1 kb) was linked to the promoter of the small subunit gene for pea ribulose 1,5-bisphosphate carboxylase ("RUBISCO", Herrera-Estrella et al. (1984) *Nature* 310, 115-120) and the

whole fragment was introduced into the *Agrobacterium* vector pPCV001 (Koncz and Schell (1986) Mol. Gen. Genet. 204, 383-396). After leaf disc transformation of SR1 tobacco plants the transgenic plants were identified on the basis of their kanamycin resistance. Among several transformants we selected two plants (ASP4, ASP5) which showed tolerance against treatment with 1 kg/ha PPT. As result of this PPT treatment the SR1 control plants were completely killed and we could never find outgrowings with a capability for flowering and seed production. The ASP4 and ASP5 transformants showed symptoms only on the lower and older leaves while the meristematic region could overcome the inhibition. After continuation of growth these plants flowered and produced seeds.

Selfing the ASP4 and ASP5 transgenic tobacco plants has resulted in a segregating seedling population with resistant and sensitive sexual progenies. Under the *in vitro* conditions used the presence of 10 μ M L-PPT in the culture medium could clearly discriminate between the two phenotypes.

The presence of the *asnA* sequence in the genome of the transformants was also shown by Southern DNA hybridization. After digestion of plant DNAs with *EcoRI* a hybridizing fragment was revealed in the transformed plants. In Northern hybridization analysis, a low amount of mRNA which was homologous to the *asnA* gene was detected in the total RNA isolated from the *in vitro* grown ASP5 transformant.

2. Reduced ammonia accumulation in transgenic tobacco plants

The inhibition of GS activity by PPT treatment causes a rapid increase in ammonia concentration in leaves of control tobacco plants. The rate of ammonia accumulation measured with the microdiffusion method and subsequent nesslerization (Shelp et al. (1985) Can. J. Bot. 63, 1135-1140) depends on the concentration of the applied herbicide.

At a dose of 0.5 kg/ha the transgenic plants can overcome the effects of PPT treatment (Tab. 1).

Table 1

Accumulation of ammonia in control tobacco plants (SR1) and in transgenic plants with the <i>asnA</i> gene after spraying with 0.5 kg/ha PPT			
hours	ammonia concentration (mM)		
	SR1	ASP4	ASP5
up to 4	0.58	0.42	0.40
6	1.20	0.82	0.78
24	1.70	0.70	0.75
48	1.95	0.60	0.50

A reduced level of accumulation can also be seen in these plants in comparison to SR1 tobacco plants after spraying with 1 kg/ha (Tab. 2). The detected lower ammonia level will be responsible for the less pronounced damage of transformed plants.

Table 2

Effects of 1 kg/ha PPT on ammonia concentration in tobacco (SR1) and transgenic plants (ASP4, ASP5)			
hours	ammonia concentration (mM)		
	SR1	ASP4	ASP5
up to 6	0.8	0.78	0.88
8	2.2	0.87	0.95
24	4.9	2.0	1.40
48	8.6	4.1	3.6

3. Stimulation of plant growth and development

Detailed comparison of growth behaviour between control and ASP plants revealed considerable differences: An increased growth rate was characteristic for the transgenic plants but a more significant stimulation was achieved by treatment of ASP plants with low doses of PPT.

The basic as well as the PPT induced acceleration in growth could be demonstrated by various types of growth curves. Fig. 1 shows that while the spraying with 0.025 kg/ha PPT already inhibited the growth of SR1 plants a large stimulation was detected in both of the transformants. Spraying with 0.05 kg/ha PPT has a negative influence on all plants. Each point represents the average height of three plants.

The differences between the various lines under control and treated conditions are also detectable if we characterize the growth of plants by Baule Mitscherlich-curves (Fig. 2) under greenhouse conditions. The inhibition and stimulation of growth can be followed by the slope of plots with characteristic alpha angles shown in Fig. 2.

4. Increased dry weight in transgenic plants

In addition to the differences in plant height the stimulatory effects were also detectable by measuring dry weight. The data shown in Table 3 demonstrate the higher productivity of the *asnA* transformants:

Table 3

Final dry weight (gr) of control (SR1) and transformants (ASP4, ASP5)		
Lines	Treatment	
	Control	0.025 kg/ha PPT
SR1	3.19 100 %	2.76 100 %
ASP4	3.85 120 %	4.79 173 %
ASP5	3.74 117 %	5.05 183 %

Example II

Effect of the *asnA* gene driven by the CaMV35S promoter in transgenic plants

1. Selection of transgenic plants

As an alternative approach we have introduced plasmid molecules (pUC) carrying the *E. coli* *asnA* gene with the CaMV35S promoter into SR1 leaf protoplasts by direct DNA uptake (R. X. Fang et al. (1989) The Plant Cell 1, 141-150). The transformants were directly selected on the basis of their PPT resistance. Plants were regenerated from micro calli grown in the presence of 10 μ M L-PPT. The Southern hybridization confirmed the presence of the *asnA* gene in DNA isolated from the PPT resistant regenerants.

2. Reduced ammonia accumulation and improved PPT tolerance in transgenic plants

Selfing of regenerated transformants resulted in segregating progenies with various levels of PPT resistance (medium supplemented with up to 30 μ M L-PPT).

In agreement with the resistant phenotype the transformed plants accumulate less ammonia than the SR1 plants when sprayed with 1 kg/ha PPT (Tab. 4).

Table 4

Ammonia accumulation after spraying the plants with 1 kg/ha PPT			
hours	ammonia concentration (mM)		
	SR1	ASP70	ASP95
6	6.85	2.92	1.95
24	9.50	5.60	5.10
48	22.3	13.60	17.40
120	58.60	28.30	35.6
144	113.00	39.40	50.00

3. Efficiency of photosynthesis

Both the control SR1 and transgenic tobacco plants were characterized by various parameters of photosynthesis such as the CO₂ fixation rate (Szajko et al. 1971, Acta Agr. Acad. Hung. 20, 247-260) and fluorescence induction (Hideg et al., 1986, Photobiochem. Photobiophys. 12, 221-230). Under greenhouse conditions the plants were treated with various doses of PPT and the content of ammonium was also determined. As shown by Table 5 the transgenic plants with the ASN-A gene exhibit a considerable increase in efficiency of net CO₂ fixation in comparison to the control plants. Application of low dose PPT treatment can further stimulate CO₂ fixation, while the difference between SR1 plants with or without PPT (50 g/ha) treatment is not statistically significant. The Table 5 provides also evidence that in the case of tobacco plants the inhibitory concentration of PPT causes ammonium accumulation with serious damage in photosynthesis by inhibition of electron transport and a 50 % reduction of CO₂ fixation. Under the same conditions the transformed plants (ASP 70) can tolerate the treatment as the photosynthetic function is concerned.

Table 5

Parameters of photosynthesis

Lines	PPT treat- ment (g/ha)	Ammonia concen- tration (mM)	CO ₂ fixation ($\mu\text{mol CO}_2/\text{dm}^2\text{xh}$)				Fluorescence induction (in % of control SRI)		
			$\bar{x}$	$\pm s_{\bar{x}}$	n	P 1%	F_m	F_o	$F_i - F_o / F_m - F_o$
SR1	0	0.37	38.17	12.05	20	-	100	100	0.45
	50	2.00	44.23	17.73	20	-	101	102	0.44
	750	34.35	19.54	12.42	20	+	80	194	0.59
ASP70	0	0.47	49.32	7.86	20	+	98	114	0.38
	50	2.70	58.07	6.06	20	+	99	110	0.42
	750	15.80	31.61	14.16	20	-	92	144	0.49

The analysis was carried out 4 days after PPT treatment.

4. Growth behaviour of asnA transformant plants

Analysis of growth rate (mm/day) reproducibly showed accelerated growth of transformants during the early plant development. Data are shown in Table 6 for plants grown in the green house.

Table 6

Growth rate (mm/day) during various periods of plant development (green house)										
Lines	Periods (6 days)								Final plant height (cm)	
	I	II	III	IV	V	VI	VIII	VIII		
SR1	0.29	0.45	0.27	0.52	0.75	1.31	1.93	1.37	41.08	
Asp70/1	0.60	1.15	0.43	1.23	1.83	2.08	1.53	0.50	58.0	141 %
Asp70/2	0.61	0.93	0.50	0.75	1.18	1.51	1.83	0.25	48.5	118 %
SR1: average of 5 plants										
ASP70/1 and ASP70/2: individual plants										

The analysis of these plants under field conditions revealed similar differences as it was observed in the green house (Table 7). The growth rate of ASP plants during period I-III was considerably higher than in the case of SR1 plants. In this experiment the stimulatory effect of PPT on the transgenic plants was also confirmed especially in the last growing period. The Baule-Mitscherlich curves (Fig. 3) clearly demonstrate that the ASP plants exhibit faster growth than control SR1 plants grown in the field.

Table 7

Growth rate (mm/day) during various periods of plant development (field experiment)							
Treatment	Lines	Periods (7 days)				Final plant height (cm)	
		I	II	III	IV		
Control	SR1	0.36	0.85	1.31	3.24	42.5	100 %
	Asp70	0.45	1.07	1.50	3.14	47.6	112 %
	Asp95	0.48	1.14	1.92	3.24	51.5	121 %
25 g/ha PPT	SR1	0.29	0.56	1.15	2.74	35.0	100 %
	Asp70	0.44	1.02	1.69	3.84	53.8	154 %
	Asp95	0.31	0.88	1.27	3.78	48.7	139 %
Average of 5 plants							

5. Productivity of asnA transformants

As shown by Table 8 the total green mass as well as the dry weight was increased in ASP plants in comparison to SR1 plants. Here we can also see that the transgenic plants are significantly stimulated by PPT treatment. At the same time the control SR1 plants are already inhibited by the spraying.

Table 8

Green mass (gr) field test								
	Control				25 g/ha PPT			
Lines	Total	%	Leaf	%	Total	%	Leaf	%
SR1	86.5	100	57.4	100	78.7	100	43.4	100
ASP70	95.2	110	62.3	108	139.9	178	92.41	213
ASP95	103.8	120	68.4	119	105.0	133	71.56	165
Dry weight (gr) field test								
	Control				25 g/ha PPT			
Lines	Total	%	Leaf	%	Total	%	Leaf	%
SR1	6.66	100	4.83	100	6.42	100	5.05	100
ASP70	8.05	121	5.82	120	10.99	171	8.56	169
ASP95	8.48	127	6.34	131	8.96	140	6.78	134
All average data from 5 plants								

Claims

1. A plant cell expressing a prokaryotic ammonium-specific asparagine synthetase (ASN-A).
2. A plant, seeds or propagation material, containing cells as claimed in claim 1.
3. A gene construct comprising a gene encoding a prokaryotic ASN-A, operatively linked to a regulatory sequence which effects the expression of said gene in a plant cell.
4. A gene construct according to claim 3, wherein said ASN-A is an E. coli ASN-A.
5. A gene encoding E. coli ASN-A consisting of plant specific codons.
6. A gene construct comprising the gene according to claim 5 under the control of a regulatory sequence active in plants.
7. A vector containing a gene construct according to claims 3, 4 or 6.
8. A plant cell transformed with a gene construct according to claims 3, 4 or 6 or with a vector according to claim 7.
9. The use of a prokaryotic asnA gene in a plant cell to improve the phenotype of this plant cell expressing said gene.
10. The use of an E. coli asnA gene in a plant cell to improve the phenotype of this plant cell expressing said gene.

Patentansprüche

1. Pflanzenzelle, welche eine prokaryotische ammoniumspezifische Asparaginsynthetase (ASN-A) exprimiert.
2. Pflanze, Samen oder Fortpflanzungsmaterial, enthaltend Zellen wie in Anspruch 1 definiert.
3. Gen-Konstrukt umfassend ein Gen, das für eine prokaryotische ASN-A kodiert, welches operabel an eine Steuersequenz gebunden ist, die die Expression des genannten Gens in einer Pflanzenzelle

Table 8

Green mass (gr) field test								
	Control				25 g/ha PPT			
Lines	Total	%	Leaf	%	Total	%	Leaf	%
SR1	86.5	100	57.4	100	78.7	100	43.4	100
ASP70	95.2	110	62.3	108	139.9	178	92.41	213
ASP95	103.8	120	68.4	119	105.0	133	71.56	165
Dry weight (gr) field test								
	Control				25 g/ha PPT			
Lines	Total	%	Leaf	%	Total	%	Leaf	%
SR1	6.66	100	4.83	100	6.42	100	5.05	100
ASP70	8.05	121	5.82	120	10.99	171	8.56	169
ASP95	8.48	127	6.34	131	8.96	140	6.78	134
All average data from 5 plants								

Claims

1. A plant cell expressing a prokaryotic ammonium-specific asparagine synthetase (ASN-A).
2. A plant, seeds or propagation material, containing cells as claimed in claim 1.
3. A gene construct comprising a gene encoding a prokaryotic ASN-A, operatively linked to a regulatory sequence which effects the expression of said gene in a plant cell.
4. A gene construct according to claim 3, wherein said ASN-A is an E. coli ASN-A.
5. A gene encoding E. coli ASN-A consisting of plant specific codons.
6. A gene construct comprising the gene according to claim 5 under the control of a regulatory sequence active in plants.
7. A vector containing a gene construct according to claims 3, 4 or 6.
8. A plant cell transformed with a gene construct according to claims 3, 4 or 6 or with a vector according to claim 7.
9. The use of a prokaryotic asnA gene in a plant cell to improve the phenotype of this plant cell expressing said gene.
10. The use of an E. coli asnA gene in a plant cell to improve the phenotype of this plant cell expressing said gene.

Patentansprüche

1. Pflanzenzelle, welche eine prokaryotische ammoniumspezifische Asparaginsynthetase (ASN-A) exprimiert.
2. Pflanze, Samen oder Fortpflanzungsmaterial, enthaltend Zellen wie in Anspruch 1 definiert.
3. Gen-Konstrukt umfassend ein Gen, das für eine prokaryotische ASN-A kodiert, welches operabel an eine Steuersequenz gebunden ist, die die Expression des genannten Gens in einer Pflanzenzelle



bewirkt.

4. Gen-Konstrukt nach Anspruch 3, wobei die genannte ASN-A eine E.coli ASN-A ist.
- 5 5. Gen, das für E.coli ASN-A kodiert bestehend aus pflanzenspezifischen Kodons.
6. Gen-Konstrukt umfassend das Gen nach Anspruch 5 unter der Kontrolle einer Steuersequenz, die in Pflanzen wirksam ist.
- 10 7. Vektor enthaltend ein Gen-Konstrukt nach den Ansprüchen 3, 4 oder 6.
8. Mit einem Gen-Konstrukt nach den Ansprüchen 3, 4 oder 6 oder mit einem Vektor nach Anspruch 7 transformierte Pflanzenzelle.
- 15 9. Verwendung eines prokaryotischen ASN-A Gens in einer Pflanzenzelle, um den Phenotyp dieser Pflanzenzelle, welche das genannte Gen exprimiert, zu verbessern.
10. Verwendung eines E.coli ASN-A Gens in einer Pflanzenzelle, um den Phenotyp dieser Pflanzenzelle, welche das genannte Gen exprimiert, zu verbessern.

20

Revendications

1. Cellule végétale exprimant une asparagine synthétase procaryote spécifique de l'ammonium (ASN-A).
- 25 2. Plante, grains ou matériel de propagation, contenant les cellules revendiquées dans la revendication 1.
3. Construction de gène comprenant un gène codant pour une ASN-A procaryote, liée de façon opérationnelle à une séquence régulatrice qui agit sur l'expression dudit gène dans une cellule végétale.
- 30 4. Construction de gène selon la revendication 3, dans laquelle ladite ASN-A est une ASN-A d'E. coli.
5. Gène codant pour la ASN-A d'E. coli qui comprend des codons spécifiques de plantes.
- 35 6. Construction de gène comprenant un gène selon la revendication 5, sous le contrôle d'une séquence régulatrice active dans les plantes.
7. Vecteur contenant une construction de gène selon les revendications 3, 4 ou 6.
- 40 8. Cellule végétale transformée avec une construction de gène selon les revendications 3, 4 ou 6 ou avec un vecteur selon la revendication 7.
9. Utilisation d'un gène asnA procaryote dans une cellule végétale pour améliorer le phénotype de cette plante en exprimant ledit gène.
- 45 10. Utilisation d'un gène asnA d'E. coli dans une cellule végétale pour améliorer le phénotype de cette plante en exprimant ledit gène.

50

55

