

Description

This invention is relating to complex compounds combining flavonoids with phospholipids, to a process for their preparation and to pharmaceutical and cosmetic compositions containing them.

Flavonoids, a class of substances very widely distributed in the vegetal kingdom, comprise various subclasses such as the flavans, flavanones, flavones, anthocyanins, etc...

The term "flavonoids" as used herein comprises the flavonoids selected from the group consisting of quercetin, kaempferol, quercetin-3-ramnoglucoside, quercetin-3-ramnoside, quercetin-3-glucoside, hyperoside, vitexine, isovitexine, diosmin, kaempferol-3-glucoside, 3-ramnoside, apigenin, apigenin-7-glucoside, luteolin, luteolin-glucoside, ramnoglucoside, 5,7,3',4'-pentahydroxy-flavone-3D- α -rhamnopyranosyl-4-D-beta-O-(6"-transcoumaroyl)-glucopyranoside, 3,5,7,4'-tetrahydroxyflavone-3-D- α -rhamnopyranosyl-4-D-beta-O-(6"-trans-coumaroyl)-glucopyranoside, Ginkgonetine, Isoginkgonetine and Sciadopitasine and Bilobetine.

The same flavonoids may moreover be in a monomer, dimer or oligomeric form, of up to 8 base units, and may be in a free or glycosidic form, wherein one or more hydroxyl groups are involved in glycosidic linkages with one or more saccharide units.

Flavonoids are known to possess a number of recognized and investigated pharmacological properties among which are the anti-inflammatory, antispasmodic, antihistaminic, peripheral vasodilatory, platelet antiaggregating, vasoprotector in terms of altered capillary fragility and permeability, and antiallergic properties which are in connection with, on the one hand, their antioxidant activity as free radical scavengers, and, on the other hand, the interference with many enzyme systems (phosphodiesterase, lipooxygenase, cyclooxygenase, aldosereductase, protein kinase, histidine-decarboxylase).

Some Authors (E. Middleton, J. Tips 5, 335-338, 1984), in view of the wide range of activity of flavonoids and the capability thereof of modulating various physiological processes, have even formulated the hypothesis that flavonoids are semi-essential factors for human beings who daily take in small doses of flavonoids through their normal diet.


EP-A-209038 discloses a variety of specific flavonoid complexes. CA 98 : 105933h (Food Chem. 1983, 10(2), 11-20) discloses the synergistic effects of phospholipids in enhancing the antioxidant properties of polyhydroxyflavonoids in lard. EP-A-275224, filed on the same priority date of this patent application, discloses flavonoid complexes of *Vitis vinifera* extracts.

As a result, many therapeutic applications for flavonoids or the plant extracts containing them, have been devised and are known: however, a severe limitation exists and is imputable to the poor or very poor absorption of these active constituents when administered per os, or by topical application. The reasons for this poor absorption are partly due to a bacterial degradation of the phenol moiety of the molecule and a complex formation with other substances present in the gastrointestinal tract thus preventing them from being absorbed.

The therapeutic effect of flavonoids subsequent to parenteral administration shows, on the other hand, how the activity of these compounds is bound to their concentration in the plasma or the target-organ or -tissue. The oral or topical administration of flavonoids does not permit effective concentration to be achieved on all occasions.

On the contrary, the compounds of flavonoids with phospholipids, subject of the invention, thanks to their lipophile character, are perfectly absorbed through oral administration and increase the therapeutic effectiveness of the flavonoids both by oral and parenteral or topical administration.

The phospholipids that are useful to this invention may be either vegetable or synthetic in nature, with acyl residues being the same or different, as shown by the formula:



wherein R and R₁, that are the same or different, are mostly acyl residues of the palmitic, stearic, oleic, linoleic, linolenic acids while R₂ is the residue of choline, ethanolamine or serine. Particularly preferred phospholipids for use in cosmetics are the vegetal or naturally occurring phospholipids, such as those

obtained from soy or from bovine or swine cutis or brain, similar to the ones that are found in human dermis; for other uses, a phospholipid that is chemically homogeneous and defined in its structure units (acyl and phosphoryl amine group) is preferred.

More particularly, the phospholipids useful to this invention are phospholipids selected from the group consisting of soy lecithins or egg, phospholipids from bovine or swine brain or dermis, phosphatidyl choline, phosphatidyl serine, phosphatidyl ethanolamine, in which the acyl groups may be the same or different and are mostly derived from palmitic, stearic, oleic, linoleic, linolenic acids.

The complex compounds according to the invention are prepared by reacting a flavonoid in an aprotic solvent with the phospholipid dissolved in the same solvent. The mole ratio of phospholipid to flavonoid is in the range from 0.5 to 2; a more preferable mole ratio being about 1. After solubilization has been completed, the complex compound is isolated by removing the solvent under vacuum, by lyophilization or by precipitation with nonsolvents.

The complex compound thus obtained is lipophile in character and is soluble in nonpolar and aprotic solvents in which the individual components of the complex compound are normally insoluble.

The formation of a molecular complex is confirmed by a NMR spectroscopy study of the proton, carbon-13 and phosphorus, by comparing the spectra of the individual constituents with those of the reaction product. In the complex's ^1H -NMR spectrum, the signals of the protons characteristics of the flavonoid molecule are not detectable, or are very enlarged, whereas the protons belonging to the lipid can be observed. Clearly apparent in the same spectrum is a broadening of the band of $\text{N}-(\text{Me})_3$ group of choline, showing that this moiety is involved in the complex formation.

In the ^{13}C -NMR spectrum, the value of the relaxation times of the nuclei that are most involved in the complex's formation is reduced in a similar manner to the proton spectrum, until disappearance of all the signals of the flavonoid and the choline and glyceric residue of the phospholipid takes place. In the ^{31}P -NMR spectrum, a substantial broadening of the phosphorus band is observed with conspicuous peak shift.

On the basis of the spectroscopic results, in the formation of these complexes the polar head of the phospholipid is involved while the fatty acid moieties retain a high degree of mobility conferring marked lipophilia at the new molecule.

The complex compounds of the invention, as obtained from the pure compounds or the extracts, possess higher bio-availability, for instance the complexes by topical application, are 2 to 4 times as active as are the substances administered in a free form.

As it is apparent from Table I, in which the vasodilatory activity of the Ginkgo biloba extract is compared with that of the corresponding complex compound, absorption in a rat under genetic hypertension is more favourable for the complexed form; the same results are obtainable in the activity on the microcirculation and the platelet aggregation and in tests connected with topical application.

Table II gives data concerning the antiedematous activity exhibited by the Ginkgo biloba extract both alone and in complexed form, in croton oil test by topical application. It clearly appears from these data that the complexed form has substantially higher activity than normal one.

Moreover, of particular interest in the specific case of the extracts from Ginkgo biloba, Crataegus and Passiflora incarnata, is the increment of the blood flow at the cutaneous microcirculation level, that can be caused to occur in application areas of respective complexed forms, as compared with free forms; here, the action of the complexed form, besides being higher in intensity at the same dose of active principle, is longer lasting in time. It can be easily appreciated how an increment in blood flow at the cutaneous level is of importance also in cosmetic applications, in order to maintain the skin moisturised, elastic and in physiological conditions. A peculiar action of the flavonoids in general, and of the catechuic derivatives in particular, is their fibroblastic proliferation stimulating action, which action is equally useful in dermatology for ulcer healing and in cosmetic treatments where it intervenes to increase the synthesis of the mucopolysaccharides and of the collagen in the derma perivascular connective tissue.

Still in with the cosmetics field, glucoside derivative of luteolin and the Ginkgo biloba extract itself, when in complexed forms, have proved useful in bleaching melaninic freckles or pigmented maculae occurring in senescent skin, or as caused by exposure to the sun, the use of contraceptives, etc... Surprisingly, the complexed form remains a longer time in the skin to go deeper into it so that the effect is more apparent and is faster than with the free form. With other flavonoids, such as apigenin and its derivatives and luteolin and its glucosides, the anti-inflammatory activity, following topical application (already known in literature), measured as carragenin oedema-inhibition and in terms of UV radiation protection, was in all cases higher in the complex compounds of the invention than in the free flavonoids.

Quercetin, apigenin and the corresponding phosphatidylcholine-complexes have been tested for effectiveness in inhibiting the Croton oil oedema. From the results given in Table III it clearly appears that the complexed forms are higher in activity than the free flavonoids.

TABLE 1 - Arterial pressure in rat in state of spontaneous hypertension

Substance	Dose mg/kg	No. animals	Basal pressure	Systolic pressure, mm/Hg					
				4 h I Die	4 h II Die	4 h III Die	4 h IV Die	24 h IV Die	
Controls		7	206±5	203±6	200±6	203±4	199±5	199±6	199±6
Gingko biloba extract	5	7	209±4	199±6 (-7)	190±7 (-16)	189±2* (-14)	196±4* (-10)	189±5** (-17)	183±3*** (-23)
Gingko biloba/phosphatidyl choline complex	12,5	7	206±3	191±5** (-18)	186±8** (-23)	184±7** (-24)	181±7** (-28)	180±3*** (-29)	177±4*** (-32)

* = P lower than 0,05

** = P lower than 0,01

*** = P lower than 0,001

TABLE 2 - Inhibition of the Croton oil oedema in mouse.

Substances	Dose µg	Animals No.	Edema mg ± E.S.	Reduction %
Controls	-	8	5,6 ± 0,4	-
Gingko biloba extract	300	8	4,7 ± 0,9	-16
Gingko biloba/phosphatidyl choline complex	75	8	0,8 ± 0,3*	-86*

* $p \leq 0,001$

TABLE 3 - inhibition of Croton oil oedema in mouse

Substances	Dose ug	Animals No.	Edema mg $\pm$ E.S.	Reduction %
Controls	-	8	5,6 $\pm$ 0,4	
Quercetin	50	8	4,2 $\pm$ 0,6	-25
Apygenin	30	8	3,4 $\pm$ 0,5	-39
Quercetin/phosphatidyl choline complex	35	8	3,0 $\pm$ 0,6	-47
Quercetin/phosphatidyl choline complex	70	8	1,5 $\pm$ 0,4	-73
Apygenin/phosphatidyl choline complex	70	8	0,8 $\pm$ 0,3	-86

The epithelium-repairing activity and the fibroblastic proliferation-stimulating activity exhibited by the phospholipid complexes of catecholic dimeric and trimeric substances, such as those isolated from Tormentilla, are significantly greater than in the case with the free forms.

The peculiar activities of flavones, such as their capillary-protective and vasotropical activities, have been checked for the complex compounds of rutin, diosmin, quercetin and of flavonoids from whitethorn, passion flower and Ginkgo biloba.

Especially when tested by topical application, the complexes of the above-mentioned products have shown to provide significant protective effect. From an applicative point of view of pharmaceutical and cosmetic technology, the various complex compounds obtained as above can be employed as microdispersions in water by preparing them by homogenization using high-speed stirrers or ultrasonic procedures, or they may be incorporated as such into appropriate pharmaceutical or cosmetic preparations. The complexes are always provided in solid form thereby enabling them to be used for preparing tablets, capsules, powders, suppositories, creams and gels, or in aqueous microdispersions that are conveniently kept in dispersed state with thickening agents. Advantageously, in view of the higher activity exhibited by the complexed forms according to the invention, the dosage of the active principle may, under certain circumstances, be decreased while maintaining the specific activity unchanged.

The present invention also relates to pharmaceutical and/or cosmetic compositions containing as the active principle one of the complex compounds according to the invention, in admixture with an appropriate pharmaceutical or cosmetic carrier. Said compositions will be of use as medicaments or in cosmetology. Particularly preferred cosmetic compositions will contain, as the active principle, complex compounds of luteolin and its glucoside with phospholipids, in admixture with an appropriate carrier.

Suitable forms for pharmaceutical and/or cosmetic uses by topical application, are creams, gels or aqueous microdispersions containing 0.1 to 10% by weight of one or more complex compounds of the invention. Said forms will be administered once or several times daily, depending on the contemplated use. Suitable forms for pharmaceutical uses, by oral or parenteral administration, are tablets, capsules, syrups, granules, solutions and suspensions which contain unit doses of the complexate active principle in the range from 1 to 500 mg. Said pharmaceutical forms will be administered once or several times a day, depending on the severity of the pathology to be treated and the patient conditions. The compositions according to the invention can in particular be used for treating states of inflammation, of altered platelet aggregation states, of altered capillary fragility and permeability and, in general, in all the fields in which an activity of the flavonoids is recognized at present.

In the following examples, that are given for illustrative purposes and not to be constructed as limiting the invention, use has been made of a soy phosphatidylcholine containing, on an average, as fatty acids: 63% linoleic acid, 16% palmitic acid, 3.5% stearic acid and 11% oleic acid based on the total acids, or natural mixture of phospholipids extracted from vegetal sources.

EXAMPLE 1

Preparation of quercetin/di-stearoyl phosphatidylcholine equimolecular complex

3.02 g of quercetin were suspended in 100 ml of methylene chloride; 8 g of distearoyl phosphatidylcholine were added to the suspension and the mixture was heated to mild reflux to complete dissolution. The solution was concentrated to a volume of 20 ml, then diluted with 200 ml of n-hexane, under stirring. The precipitated product was filtered and dried under vacuum at 40°C. 10.6 g of a yellow powder were obtained, having

$E_{1\%} = 143.6$ at 258 nm (CHCl_3);
 $E_{1\%} = 116.3$ at 362 nm (CHCl_3).

EXAMPLE 2

Preparation of quercetin-3-galactoside/soy phosphatidylcholine equimolecular complex

4.64 g of quercetin-3-galactoside (hyperoside) were dissolved in 150 ml of a dioxane-ethanol 2:1 mixture, in the presence of 7.8 g of soy phosphatidylcholine (titre 95%). When the solution was clear, ethanol was removed by distillation under vacuum at 40°C, and the dioxane solution was freeze-dried.

12.1 g of a straw yellow powder, completely soluble in chloroform:ethyl ether, were obtained. The product showed no signals at $^1\text{H-NMR}$ (CDCl_3) which could be assigned to aromatic protons and to those of the saccharidic moiety.

$E_{1\%}$ in CHCl_3 of 160.2 at 258 nm.

EXAMPLE 3**Preparation of quercetin-3-rhamnoglucoside/soy phosphatidylcholine complex**

6.1 g of quercetin-3-rhamnoglucoside were dissolved together with 15.60 g (0.02 mole) of soy phosphatidylcholine (95% titre) in 500 ml of dioxane, heating the mixture to mild reflux for 10 minutes. When dissolution was complete, the mixture was cooled and concentrated to 40 ml under vacuum. The concentrate was poured into 500 ml of n-hexane and the resulting precipitate was filtered under vacuum, washing thoroughly with n-hexane. After drying, 20.2 g of the complex were obtained, having: m.p. = 107-109 °C; $E_{1\%}^{1\text{cm}}$ = 90.4 at 280 nm and 72.7 at 367 nm; $[\alpha]_D^{20} + 10.1^\circ$ (C = 1% in CHCl_3).

EXAMPLE 4**Preparation of kaempferol-3-glycoside/soy phosphatidylcholine complex**

4.48 g of kaempferol-3-glycoside and 7.8 g of soy phosphatidylcholine were dissolved in 60 ml of dioxane. The obtained solution was freeze-dried. 12 g of the complex were obtained, which was completely soluble in aprotic solvents, and had NMR spectra in agreement with those of complexed flavonoid products. The products had $E_{1\%}^{1\text{cm}}$ in CHCl_3 = 158.5 at 260 nm.

EXAMPLE 5**Preparation of vitexine-2"-rhamnoside-2"-acetate/soy phosphatidylcholine complex**

3.2 g of vitexine-2"-rhamnoside-2"-acetate and 3.9 g of phosphatidylcholine were dissolved in 30 ml of a dioxane:ethanol 5:1 mixture, then freeze-dried. 7 g of a light yellow powder were obtained, having $E_{1\%}^{1\text{cm}}$ = 116.9 at 272 nm (CHCl_3); $E_{1\%}^{1\text{cm}}$ = 130.5 at 330 nm (CHCl_3); and spectroscopic data in agreement.

EXAMPLE 6**Preparation of Ginkgo biloba extract/soy phosphatidylcholine complex**

10 g of Ginkgo biloba extract, having average titre in Ginkgo flavone-glycosides of about 25%, and 15 g of soy phosphatidylcholine (95% tit.) were dissolved in 200 ml of a mixture of 6 parts of methylene chloride and 1 part of methanol. When dissolution was complete, the solvent was distilled off under vacuum to small volume, the residue was diluted with 200 ml more of methylene chloride, filtering any turbidity.

Solvent was evaporated to small volume; the mixture was diluted with 300 ml of n-hexane and the product precipitated in form of a light beige solid, which was dried at 40 °C under vacuum. 22 g of product were obtained, which was completely soluble in apolar solvents, the extract was in form of a complex, as evidenced by the NMR spectra, which showed no signals of aromatic protons from flavonoid compounds. Said extract may be directly used for microdispersion or incorporation into pharmaceutical formulations.

EXAMPLE 7**Preparation of apigenin/soy phosphatidylethanolamine complex**

2.7 g of apigenin and 6.9 g of soy phosphatidylethanolamine (96 % tit.) were dissolved in 80 ml of methylene chloride. When dissolution was complete, methylene chloride was evaporated off to small volume, and the mixture was precipitated with 200 ml of n-hexane. The precipitated product was filtered and dried under vacuum at 40 °C.

8.2 g straw yellow product were obtained, having spectroscopic characteristics in agreement with the described complexes.

EXAMPLE 8**Preparation of apigenin/distearoylphosphatidylcholine complex**

2.7 g of apigenin were suspended in 100 ml of a dioxane:methanol 7:3 mixture, added with 8 g of distearoylphosphatidylcholine, and heated to reflux to complete dissolution.

The obtained solution was concentrated to dryness, the residue was dissolved in 50 ml of chloroform-methanol 9:1. The chloroform solution was evaporated to small volume and the residue was poured into 100 ml of n-hexane. 9 g of the complex were obtained, having the following characteristics: m.p. 150 °C, $E_{1\%} = 164.6$ at 270 nm; $E_{1\%} = 156.2$ at 324 nm.

EXAMPLE 9**Preparation of luteolin 7-glycoside/soy phosphatidylcholine complex**

2.24 g of luteolin-7-glucoside were suspended in 100 ml in a dioxane-methanol 7:3 mixture, and treated with 4 g of soy phosphatidylcholine (95% tit.); the mixture was refluxed till complete dissolution, solvent was removed under vacuum and the residue was taken up in 50 ml a chloroform-methanol 8:3 mixture, which was subsequently removed. The residue was dissolved in 100 ml of methylene chloride. After concentration to small volume, the residue was poured into 100 ml of n-hexane. 6.2 g of luteolin-7-glycoside and soy phosphatidylcholine complex were obtained, having the following characteristics: m.p. 120-130 °C; $E_{1\%}$ in CHCl_3 113.89 at 258 nm.

EXAMPLE 10**Preparation of diosmin/soy phosphatidylcholine complex**

6 g of diosmin and 15.5 g of soy phosphatidylcholine (95% tit.) were suspended in a dioxane-methanol 7:3 mixture, and refluxed to complete dissolution. The solution was concentrated under vacuum to small volume and the residue was redissolved in methylene chloride. After evaporation of the solution, 21 g of diosmin-phosphatidylcholine complex were obtained, having chemico-physical and spectroscopic characteristics in agreement with those of the above described complexes.

EXAMPLE 11**Preparation of ginkgo biloba depurated extract/total soy phospholipids complex**

10 g of Ginkgo biloba depurated extract having titre = 24% in Ginkgo flavone glycosides in 100 ml of methylene chloride were added with 15 g of soy phospholipids, having phosphatidylcholine content: 30%, phosphatidylethanolamine content: 20%; phosphatidylinositol content: 6% + minor phospholipids, in 15 ml of methanol, under stirring.

When the solution was clear, solvent was removed under vacuum and the residue was dissolved in 50 ml of chloroform, the solution was then poured into 500 ml of n-hexane, under strong stirring.

A viscous precipitate was obtained, which was decanted and dried under vacuum at 50 °C, for 12 hours.

A yellow-beige powder was obtained, having m.p. 105 °C, and chemico-physical and spectroscopic characteristics corresponding to a complex.

EXAMPLE 12**Preparation of Hawthorn/soy phosphatidylcholine complex**

10 g of Crataegus sp. extract, prepared starting from blossomed tops, containing 20% of flavonoidic substances expressed as hyperoside, were suspended together with 20 g of soy phosphatidylcholine (95% tit.) in 300 ml of a 9:1 chloroform-methanol mixture. When the solution was complete, the solvent was evaporated and the residue taken up with 300 ml of methylene chloride. The solvent was completely removed under vacuum.

A beige solid, having spectroscopical characteristics confirming the complex formation and easily soluble in water, was obtained.

EXAMPLE 13**Cream containing as the active ingredient the complex of Example 11**

5

10

15

20

Formulation for 100 g of cream	
Ginkgo biloba complex (Ex. 11)	2.5 g
Carboxyvinylpolymer (Carbomer 934®)	1.2 g
30% Sodium laurylsarcosinate	0.5 g
Imidazolidinylurea	0.3 g
Triton® 80	0.1 g
Polysorbate 80	3 g
Hydrogenated lanoline	5 g
Spermacetes	5 g
Polyisoprene	5 g
Wheat oil	2 g
Dimethylsilyconic oil	0.5 g
10% Sodium hydroxide sol.	2 g
Perfum	0.3 g
Water	q.s. to 100 g

25

EXAMPLE 14**Oleolyte containing as the active ingredient the complex of Example 11**

30

35

Formulation for 100 g of oleolyte	
Ginkgo biloba complex (Ex. 11)	2.5 g
Softigen® 727 (C ₈ -C ₁₂ ethoxylated triglycerides)	70 g
Volpo® 20 (polyoxyethylene 20 oleylether)	7 g
Isopropylmyristate	20 g
Preservants	0.2 g
Perfum	0.3 g

40

EXAMPLE 15**Lotion containing as the active ingredient the complex of Example 11**

45

50

55

Formulation for 100 g of lotion	
Ginkgo biloba complex	2.5 g
Softigen® 727 (C ₈ -C ₁₂ ethoxylated triglycerides)	25 g
Imidazolidinylurea	0.3 g
Octilnone	0.1 g
Volpo® 20 (polyoxyethylene 20 oleylether)	7 g
2,5% Perfumed composition	1 g
Depurated water	q.s. to 100 g

EXAMPLE 16**Gel containing as the active ingredient the complex of Example 11**

Formulation for 100 g of gel	
Ginkgo biloba complex	2.5 g
Softigen® 727 (C ₈ -C ₁₂ ethoxylated triglycerides)	25 g
Imidazolidinylurea	0.3 g
Octilnone	0.1g
Volpo® 20 (polyoxyethylene 20 oleylether)	7 g
Carboxypolymethylene (Carbopol® 934	1.5 g
Triethanolamine	2 g
2.5% Parfum	1 g
Depurated water	q.s. to 100 g
pH 7- 7.2.	

EXAMPLE 17**Capsules containing as the active ingredient the complex of Example 6**

Formulation for capsules containing 50 mg of complex	
Ginkgo biloba complex	50 mg
Silica powder	30 mg
Insoluble crosslinked polyvinylpyrrolidone	30 mg
Mais starch	20 mg
Sodium carboxymethylcellulose	10 mg
Polyvinylpyrrolidone 30000 PM	7 mg
Magnesium stearate	3 mg.

EXAMPLE 18**Capsules containing as the active ingredient the complex of Example 12**

Formulation for capsules containing 150 mg of complex	
Hawthorn complex	150 mg
Silica powder	60 mg
Mais starch	20 mg
Sodium carboxymethylcellulose	10 mg
Lactose	30 mg
Magnesium stearate	3 mg

EXAMPLE 19**Cream containing as the active ingredient the complex of Example 7**

Formulation for 100 g of cream	
Apigenin complex	2.0 g
Polyethyleneglycol	2 g
Polysorbate 80	3 g
Cetyl alcohol	10 g
Wheat oil	2 g
Silicon oil 350 cps	0.5 g
Antioxidants (oxinex 2004)	0.1 g
Carboxyvinylpolymer (Carbomer 934®)	0.8 g
Triethanolamine	1.2 g
Preservants (a mixture of methyl and propyl p-hydroxybenzoates)	0.2 g
Parfumed composition	0.1 g
Depurated water	q.s. to 100 g

EXAMPLE 20**Cream containing as the active ingredient the complex of Example 9**

Formulation for 100 g of cream	
Luteoline complex	1.5 g
Glycerol monostearate	3 g
C ₈ -C ₁₂ Liquid saturated triglycerides	10 g
Cetyl alcohol	4 g
Hydrogenated lanoline	10 g
Pentamethyleritrene	3 g
Polysorbate 80	10 g
Carboxyvinylpolymer (Carbomer 934®)	0.5 g
Triethanolamine	0.6 g
Parfumed composition	0.2 g
Preservants (a mixture of methyl and propyl p-hydroxybenzoates)	0.2 g
Depurated water	q.s. to 100 g

EXAMPLE 21**Gel containing as the active ingredient the complex of Rutine**

Formulation for 100 g of gel	
Rutine complex	1 g
Imidazolidinylurea	0.3 g
Octilnone	0.1 g
C ₈ -C ₁₂ Ethoxylated triglycerides (Softigen 767)	25 g
Polyoxyethylene 20 oleylether	5 g
Carboxyvinylpolymer (Carbomer 934)	1.5 g
Triethonalmine	2 g
Perfumed composition	0.1 g
Depurated water	65 g

EXAMPLE 22**Cream containing as the active ingredient the complex of Rutine**

Formulation for 100 g of cream	
Rutine complex	2 g
Isopropyl miristate	10 g
30% Sodium laurylsarcosinate	3 g
Carboxyvinylpolymer (Carbomer 934®)	1 g
10% Sodium hydroxide solution	2.2 g
Preservants (methyl and propyl p-hydroxybenzoates)	0.2 g
Lavender essences	0.2 g
Depurated water	q.s. to 100 g

EXAMPLE 23**Solution containing as the active ingredient the complex of Rutine**

Formulation for 100 g of solution	
Rutine complex	1 g
Imidazolidinylurea	0.3 g
Octilnone	0.1 g
Glycerides PEG-6-caprilic/capric	25 g
Polyoxyethylene 20 oleylether	5 g
Perfumed composition	0.1 g
Water	q.s. to 100 g

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Complex compounds of flavonoids with phospholipids, wherein the flavonoids are selected from the group consisting of quercetin, kaempferol, quercetin-3-ramnogluconide, quercetin-3-ramnoside, quercetin-3-glucoside, hyperoside, vitexine, isovitexine, diosmin, kaempferol-3-glucoside, 3-ramnoside,



EP 0 275 005 B1

EXAMPLE 21

Gel containing as the active ingredient the complex of Rutine

Formulation for 100 g of gel	
Rutine complex	1 g
Imidazolidinylurea	0.3 g
Octilnone	0.1 g
C ₈ -C ₁₂ Ethoxylated triglycerides (Softigen 767)	25 g
Polyoxyethylene 20 oleylether	5 g
Carboxyvinylpolymer (Carbomer 934)	1.5 g
Triethonalmine	2 g
Parfumed composition	0.1 g
Depurated water	65 g

EXAMPLE 22

Cream containing as the active ingredient the complex of Rutine

Formulation for 100 g of cream	
Rutine complex	2 g
Isopropyl miristate	10 g
30% Sodium laurylsarcosinate	3 g
Carboxyvinylpolymer (Carbomer 934®)	1 g
10% Sodium hydroxide solution	2.2 g
Preservants (methyl and propyl p-hydroxybenzoates)	0.2 g
Lavender essences	0.2 g
Depurated water	q.s. to 100 g

EXAMPLE 23

Solution containing as the active ingredient the complex of Rutine

Formulation for 100 g of solution	
Rutine complex	1 g
Imidazolidinylurea	0.3 g
Octilnone	0.1 g
Glycerides PEG-6-caprilic/capric	25 g
Polyoxyethlene 20 oleylether	5 g
Parfumed composition	0.1 g
Water	q.s. to 100 g

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Complex compounds of flavonoids with phospholipids, wherein the flavonoids are selected from the group consisting of quercetin, kaempferol, quercetin-3-ramnoglucoside, quercetin-3-ramnoside, quercetin-3-glucoside, hyperoside, vitexine, isovitexine, diosmin, kaempferol-3-glucoside, 3-ramnoside,



apigenin, apigenin-7-glucoside, luteolin, luteolin-glucoside ramnogluco-
 flavone-3D-alfa-rhamnopyranosyl-4-D-beta-O-(6"-transcoumaroyl-)glucopyranoside, 3,5,7,4'-
 tetrahydroxy-flavone-3-D-alfa-rhamnopyranosyl-4-D-beta-O-(6"-trans-coumaroyl-)glucopyranoside, Gink-
 gonetine, Isoginkgonetine and Sciadopitasine and Bilobetine.

5

2. The complex compounds according to claim 1 wherein the phospholipid to flavonoid ratio is in the range from 0.5 to 2M.

10

3. The complex compounds according to claim 1, wherein the ratio of phospholipid to flavonoid is about 1.

15

4. The complexes compounds according to any preceding claim, wherein the phospholipids are phospholipids selected from the group consisting of soy lecithins or egg, phospholipids from bovine or swine brain or dermis, phosphatidyl choline, phosphatidyl serine, phosphatidyl ethanolamine in which the acyl groups may be the same or different and are mostly derived from palmitic, stearic, oleic, linoleic, linolenic acids.

20

5. A process for preparing the complex compounds according to claims 1-4, wherein the flavonoids are reacted with the phospholipids in an inert aprotic solvent, and then the complex compound is isolated by evaporation, precipitation and/or lyophilization.

25

6. Pharmaceutical and/or cosmetic compositions containing as the active principle a complex compounds according to claims 1-4, in admixture with an appropriate pharmaceutical or cosmetic carrier.

7. The compositions according to claim 6, the compositions being in the form of aqueous microdispersions.

8. The complex compound according to claims 1-4, wherein the complex compound is for use as a medicament.

30

9. The complex compound according to claims 1-4, wherein the complex compound is for use in cosmetology.

35

10. Cosmetic compositions containing, as the active principle, complex compounds of luteolin and its glucosides with phospholipids, in admixture with an appropriate carrier.

Claims for the following Contracting States : ES, GR

40

1. A process for preparing complex compounds of flavonoids with phospholipids, wherein the flavonoids are selected from the group consisting of quercetin, kaempferol, quercetin-3-ramnogluco-
 side, 3-ramnoside, 3-glucoside, hyperoside, vitexine, isovitexine, diosmin, kaempferol-3-glucoside, 3-ram-
 noside, apigenin, apigenin-7-glucoside, luteolin, luteolin-glucoside ramnogluco-
 side, 5,7,3',4'-pentahydroxy-flavone-3D-alfa-rhamnopyranosyl-4-D-beta-O-(6"-transcoumaroyl-)glucopy-
 ranoside, 3,5,7,4'-tetrahydroxy-flavone-3-D-alfa-rhamnopyranosyl-4-D-beta-O-(6"-trans-coumaroyl-
)glucopyranoside, Ginkgonetine, Isoginkgonetine and Sciadopitasine and Bilobetine, and are reacted
 with the phospholipids in an inert aprotic solvent, and then the complex compound is isolated by
 evaporation, precipitation and/or lyophilization.

50

2. A process according to claim 1, wherein the phospholipid to flavonoid ratio is in the range from 0.5 to 2M.

3. A process according to claim 1, wherein the phospholipid to flavonoid ratio is about 1.

55

4. A process according to any preceding claim, wherein the phospholipids are phospholipids selected from the group consisting of soy lecithins or egg, phospholipids from bovine or swine brain or dermis, phosphatidyl choline, phosphatidyl serine, phosphatidyl ethanolamine in which the acyl groups may be the same or different and are mostly derived from palmitic, stearic, oleic, linoleic, linolenic acids.

**Patentansprüche****Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE**

- 5 1. Komplexverbindungen aus Flavonoiden mit Phospholipiden, dadurch **gekennzeichnet**, daß die Flavonoide aus der Gruppe bestehend aus Quercetin, Kaempferol, Quercetin-3-rhamnoglucosid, Quercetin-3-rhamnosid, Quercetin-3-glucosid, Hyperosid, Vitexin, Isovitexin, Diosmin, Kaempferol-3-glucosid, 3-Rhamnosid, Apigenin, Apigenin-7-glucosid, Luteolin, Luteolinglucosid, Rhamnoglucosid, 5,7,3',4'-Pentahydroxy-flavon-3D- α -rhamnopyranosyl-4-D- β -O-(6''-transcoumaroyl-)glucopyranosid, 3,5,7,4'-Tetrahydroxy-flavon-3-D- α -rhamnopyranosyl-4-D- β -O-(6''-trans-coumaroyl-)glucopyranosid, Ginkgonetin, Isoginkgonetin und Sciadopitasin und Bilobetin ausgewählt sind.
- 10 2. Komplexverbindungen nach Anspruch 1, dadurch **gekennzeichnet**, daß das Phospholipid:Flavonoidverhältnis im Bereich von 0,5 bis 2 M liegt.
- 15 3. Komplexverbindungen nach Anspruch 1, dadurch **gekennzeichnet**, daß das Phospholipid:Flavonoidverhältnis etwa 1 beträgt.
- 20 4. Komplexverbindungen nach einem der vorausgehenden Ansprüche, dadurch **gekennzeichnet**, daß die Phospholipide ausgewählt aus der Gruppe bestehend aus Sojalecithinen oder Ei-Phospholipiden von Rinder- oder Schweinehirn oder -haut, Phosphatidylcholin, Phosphatidylserin, Phosphatidylethanolamin, worin die Acylgruppen gleich oder verschieden sein können, sind und sich meistens von Palmitin-, Stearin-, Öl-, Linol-, Linolensäuren ableiten.
- 25 5. Verfahren zur Herstellung der Komplexverbindungen nach den Ansprüchen 1 bis 4, dadurch **gekennzeichnet**, daß die Flavonoide mit den Phospholipiden in einem inerten aprotischen Lösungsmittel umgesetzt werden, und dann die Komplexverbindung durch Verdampfung, Präzipitation und/oder Lyophilisation isoliert wird.
- 30 6. Pharmazeutische und/oder kosmetische Präparate, enthaltend als Wirkstoff eine Komplexverbindung nach den Ansprüchen 1 bis 4 im Gemisch mit einem geeigneten pharmazeutischen oder kosmetischen Träger.
- 35 7. Präparate nach Anspruch 6, dadurch **gekennzeichnet**, daß die Präparate in Form wäßriger Mikrodispersionen vorliegen.
8. Komplexverbindung nach den Ansprüchen 1 bis 4, dadurch **gekennzeichnet**, daß die Komplexverbindung als Medikament verwendet wird.
- 40 9. Komplexverbindung nach den Ansprüchen 1 bis 4, dadurch **gekennzeichnet**, daß die Komplexverbindung in der Kosmetologie verwendet wird.
10. Kosmetische Präparate, enthaltend als Wirkstoff Komplexverbindungen aus Luteolin und deren Glycosiden mit Phospholipiden im Gemisch mit einem geeigneten Träger.

45 Patentansprüche für folgende Vertragsstaaten : ES, GR

- 50 1. Verfahren zur Herstellung von Komplexverbindungen aus Flavonoiden mit Phospholipiden, dadurch **gekennzeichnet**, daß die Flavonoide aus der Gruppe bestehend aus Quercetin, Kaempferol, Quercetin-3-Rhamnoglucosid, Quercetin-3-rhamnosid, 3-Glucosid, Hyperosid, Vitexin, Isovitexin, Diosmin, Kaempferol-3-glucosid, 3-Rhamnosid, Apigenin, Apigenin-7-glucosid, Luteolin, Luteolinglucosid, Rhamnoglucosid, 5,7,3',4'-Pentahydroxy-flavon-3D- α -rhamnopyranosyl-4-D- β -O-(6''-transcoumaroyl-)glucopyranosid, 3,5,7,4'-Tetrahydroxy-flavon-3-D- α -rhamnopyranosyl-4-D- β -O-(6''-trans-coumaroyl-)glucopyranosid, Ginkgonetin, Isoginkgonetin und Sciadopitasin und Bilobetin ausgewählt werden und mit den Phospholipiden in einem inerten aprotischen Lösungsmittel umgesetzt werden, und dann die Komplexverbindung durch Verdampfung, Präzipitation und/oder Lyophilisation isoliert wird.
- 55 2. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß das Phospholipid:Flavonoidverhältnis im Bereich von 0,5 bis 2 M liegt.



3. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß das Phospholipid:Flavonoidverhältnis etwa 1 beträgt.

5 4. Verfahren nach irgendeinem der vorausgehenden Ansprüche, dadurch **gekennzeichnet**, daß die Phospholipide Phospholipide, ausgewählt aus der Gruppe bestehend aus Sojalecithinen oder Ei, Phospholipiden von Rinder- oder Schweinehirn oder -haut, Phosphatidylcholin, Phosphatidylserin, Phosphatidylethanolamin, worin die Acylgruppen gleich oder verschieden sein können, sind und sich meistens von Palmitin-, Stearin-, Öl-, Linol-, Linolensäuren ableiten.

10 Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

15 1. Composés complexes de flavonoïdes avec des phospholipides, dans lesquels les flavonoïdes sont choisis dans le groupe formé par la quercétine, le kaempférol, le quercétine-3-rhamnoglucoside, le quercétine-3-rhamnoside, le quercétine-3-glucoside, l'hyperoside, la vitexine, l'isovitexine, la diosmine, le kaempférol-3-glucoside, 3-rhamnoside, l'apigénine, l'apigénine-7-glucoside, la lutéoline, le lutéoline-glucoside rhamnoglucoside, le 5,7,3',4'-pentahydroxy flavone-3-D-alpha-rhamnopyrannosyl-4-D-bêta-O-(6"-trans-coumaroyl)glucopyrannoside, le 3,5,7,4'-tétrahydroxy-flavone-3-D-alpha-rhamnopyrannosyl-4-D-bêta-O-(6"-trans-coumaroyl)glucopyrannoside, la Ginkgonétine, l'isoginkgonétine et la Sciadopitasine
20 et la Bilobétine.

2. Les composés complexes selon la revendication 1, dans lesquels le rapport molaire du phospholipide au flavonoïde se situe dans l'intervalle de 0,5 à 2.

25 3. Les composés complexes selon la revendication 1, dans lesquels le rapport du phospholipide au flavonoïde est d'environ 1.

30 4. Les composés complexes selon l'une quelconque des revendications précédentes, dans lesquels les phospholipides sont des phospholipides choisis dans le groupe formé par les lecithines de soja ou d'oeuf, les phospholipides provenant du cerveau ou du derme de bovin ou de porc, la phosphatidylcholine, la phosphatidylsérine, la phosphatidylethanolamine dont les groupes acyles peuvent être identiques ou différents et sont principalement dérivés des acides palmitique, stéarique, oléique, linoléique, linolénique.

35 5. Un procédé pour préparer les composés complexes selon les revendications 1 à 4, dans lequel les flavonoïdes sont amenés à réagir avec les phospholipides dans un solvant aprotique inerte, puis le composé complexe est isolé par évaporation, précipitation et/ou lyophilisation.

40 6. Compositions pharmaceutiques et/ou cosmétiques contenant un composé complexe selon les revendications 1 à 4 comme principe actif, en mélange avec un support pharmaceutique ou cosmétique approprié.

45 7. Les compositions selon la revendication 6, les compositions étant sous la forme de microdispersions aqueuses.

8. Le composé complexe selon les revendications 1 à 4, le composé complexe étant destiné à être utilisé comme médicament.

50 9. Le composé complexe selon les revendications 1 à 4, le composé complexe étant destiné à être utilisé en cosmétologie.

10. Compositions cosmétiques contenant, comme principe actif, des composés complexes de lutéoline et de ses glucosides avec des phospholipides, en mélange avec un support approprié.

55 Revendications pour les Etats contractants suivants : ES, GR

1. Un procédé de préparation de composés complexes de flavonoïdes avec des phospholipides, dans lesquels les flavonoïdes sont choisis dans le groupe formé par la quercétine, le kaempférol, le



quercétine-3-rhamnoglucoside, le quercétine-3-rhamnoside, le quercétine-3-glucoside, l'hyperoside, la vitexine, l'isovitexine, la diosmine, le kaempférol-3-glucoside, 3-rhamnoside, l'apigénine, l'apigénine-7-glucoside, la lutéoline, le lutéoline-glucoside rhamnoglucoside, le 5,7,3',4'-pentahydroxy-flavone-3-D-alpha-rhamnopyrannosyl-4-D-bêta-O-(6"-*trans*-coumaroyl)glucopyrannoside, le 3,5,7,4'-tétrahydroxy-flavone-3-D-alpha-rhamnopyrannosyl-4-D-bêta-O-(6"-*trans*-coumaroyl)glucopyrannoside, la Gin kgonétine, l'Isoginkgonétine et la Sciadopitasine et la Bilobétine, et sont amenés à réagir avec les phospholipides dans un solvant aprotique inerte, puis le composé complexe est isolé par évaporation, précipitation et/ou lyophilisation.

- 5
- 10 2. Un procédé selon la revendication 1, dans lequel le rapport molaire du phospholipide au flavonoïde se situe dans l'intervalle de 0,5 à 2.
3. Un procédé selon la revendication 1, dans lequel le rapport du phospholipide au flavonoïde est d'environ 1.
- 15 4. Un procédé selon l'une quelconque des revendications précédentes, dans lequel les phospholipides sont des phospholipides choisis dans le groupe formé par les lécithines de soja ou d'œuf, les phospholipides provenant du cerveau ou du derme de bovin ou de porc, la phosphatidylcholine, la phosphatidylsérine, la phosphatidyléthanolamine dont les groupes acyles peuvent être identiques ou différents et sont principalement dérivés des acides palmitique, stéarique, oléique, linoléique, linoléique.
- 20

25

30

35

40

45

50

55