

DESCRIPTION

Title of Invention: METHOD FOR PRODUCING FLUOROALKYNE COMPOUND

Technical Field

5 [0001]

The present disclosure relates to a method for producing a fluoroalkyne compound.

Background Art

10 [0002]

Fluoroalkyne compounds are expected to be useable as cleaning gases, etching gases, refrigerants, heat transfer media, building blocks for organic synthesis, etc.

[0003]

15 As an example of a method for producing a fluoroalkyne compound, Non-patent Literature (NPL) 1 discloses that a fluoroalkyne compound is obtained from a fluoroalkane compound by a dehydrofluorination reaction, using a base such as lithium diisopropylamide (LDA).

20

Citation List

Non-patent Literature

[0004]

NPL 1: Tetrahedron, 1988, vol. 44, No. 10, p. 2865-2874

25

Summary of Invention

Technical Problem

[0005]

30 An object of the present disclosure is to provide a method for efficiently synthesizing a fluoroalkyne compound from a fluoroalkane compound.

Solution to Problem

[0006]

35 The present disclosure includes the following subject

matter.

[0007]

Item 1. A method for producing a fluoroalkyne compound represented by formula (1):

5 $R^1C\equiv CR^2$ (1)

wherein R^1 and R^2 are the same or different and each is a fluorine atom or a fluoroalkyl group,

the method comprising subjecting a fluoroalkane compound represented by formula (2):

10 $R^1CHX^1CFX^2R^2$ (2)

wherein R^1 and R^2 are as defined above; and X^1 is a fluorine atom and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine atom, to a dehydrofluorination reaction in the presence of a solvent containing an ether,

15 the method satisfying at least one requirement selected from the group consisting of the following:

(I) the fluoroalkyl group is a C_1 - C_4 fluoroalkyl group, and

(II) the ether is a chain ether, and the dehydrofluorination reaction is performed in the presence of the solvent containing a chain ether, and a base comprising a hydroxide and/or alkoxide of an alkali metal and/or alkaline earth metal.

[0008]

Item 2. The production method according to Item 1, wherein the water concentration of the solvent is 0.01 to 500 mass ppm based on the total amount of the solvent taken as 100 mass%.

[0009]

Item 3. The production method according to Item 1 or 2, wherein the number of ether bonds in the ether is 1 to 10.

[0010]

30 Item 4. The production method according to any one of Items 1 to 3, which satisfies requirement (I).

[0011]

Item 5. The production method according to Item 4, wherein the fluoroalkyl group is represented by formula (3):

35 $-CF_2R^3$ (3)

wherein R^3 is a fluorine atom or a C_1 - C_3 fluoroalkyl group.

[0012]

Item 6. The production method according to Item 4 or 5, wherein the dehydrofluorination reaction is performed in the presence of
5 a base.

[0013]

Item 7. The production method according to Item 6, wherein the base is a hydroxide and/or alkoxide of an alkali metal and/or alkaline earth metal.

10 [0014]

Item 8. The production method according to any one of Items 1 to 3, which satisfies requirement (II).

[0015]

Item 9. The production method according to Item 8, wherein the
15 fluoroalkyl group has 1 to 10 carbon atoms.

[0016]

Item 10. The production method according to Item 8 or 9, wherein the fluoroalkyl group is represented by formula (3):



20 wherein R^3 represents a fluorine atom or a C_1 - C_9 fluoroalkyl group.

[0017]

Item 11. The production method according to any one of Items 1 to 10, wherein the reaction temperature in the dehydrofluorination reaction is 0 to 300°C.

25 [0018]

Item 12. A composition comprising a fluoroalkyne compound represented by formula (1):



wherein R^1 and R^2 are the same or different and each is a fluorine
30 atom or a fluoroalkyl group, and a fluoroalkene compound represented by formula (4):



wherein R^1 and R^2 are as defined above,

the content of the fluoroalkyne compound being 25.00 to
35 75.00 mol%, the content of the fluoroalkene compound being 25.00

to 75.00 mol%, and the total content of the fluoroalkyne compound and the fluoroalkene compound being 90.00 to 100.00 mol%, based on the total amount of the composition taken as 100 mol%.

[0019]

- 5 Item 13. The composition according to Item 12, which is for use as a cleaning gas, an etching gas, a refrigerant, a heat transfer medium, or a building block for organic synthesis.

[0020]

- Item 14. Use of the composition according to Item 12 for a
10 cleaning gas, an etching gas, a refrigerant, a heat transfer medium, or a building block for organic synthesis.

Advantageous Effects of Invention

[0021]

- 15 The present disclosure provides a method for efficiently synthesizing a fluoroalkane compound from a fluoroalkane compound.

Description of Embodiments

20 [0022]

- In the present specification, the terms "comprise," "contain," and "include" encompass the concepts of comprising, consisting essentially of, and consisting of. In the present specification, a numerical range indicated by "A to B" means A or
25 more, and B or less.

[0023]

- In the present disclosure, "selectivity" refers to the ratio (mol%) of the total molar amount of the target compound contained in a gas flowing out from a reactor outlet to the total
30 molar amount of the compounds other than the starting compound in the gas flowing out from the reactor outlet.

[0024]

- In the present disclosure, "conversion" refers to the ratio (mol%) of the total molar amount of the compounds other
35 than the starting compound contained in a gas flowing out from a

reactor outlet to the molar amount of the starting compound supplied to the reactor.

[0025]

In the present disclosure, "yield" refers to the ratio
5 (mol%) of the total molar amount of the target compound contained in a gas flowing out from a reactor outlet to the molar amount of the starting compound supplied to the reactor.

[0026]

A known method for producing a fluoroalkyne compound is
10 a method in which a fluoroalkene compound, such as $\text{CF}_3\text{CF}=\text{CHCF}_3$, is synthesized from a fluoroalkane compound, such as $\text{CF}_3\text{CFHCFHCF}_3$, by a dehydrofluorination reaction, and then a dehydrofluorination reaction is further performed to synthesize a fluoroalkyne compound. In this method, it is necessary to perform a
15 dehydrofluorination reaction twice, which lengthens the production process.

[0027]

Another known method is a method in which a
fluoroalkyne compound is obtained from a fluoroalkane compound by
20 a single dehydrofluorination reaction using lithium diisopropylamide (LDA) as a base in a solvent such as hexane, diethyl ether, or monoethylene glycol dimethyl ether, as in NPL 1. In this method, a fluoroalkyne compound is obtained from a fluoroalkane compound by a single dehydrofluorination reaction
25 using $\text{KOC}(\text{CH}_3)_3$ as a base, even in a solvent such as tetrahydrofuran or dimethyl sulfoxide. However, this method is not efficient when lithium diisopropylamide (LDA) is used in this method, because lithium diisopropylamide (LDA) is expensive, and the yield is at most 41% (at most 24% when expensive LDA is not
30 used). Thus, this method is not an effective production method.

[0028]

According to the production method of the present disclosure, a method for efficiently synthesizing a fluoroalkane compound from a fluoroalkane compound in a high yield compared
35 with the conventional methods can be provided.

[0029]

1. Method for Producing Fluoroalkyne Compound (First Embodiment)

The method for producing a fluoroalkyne compound in the first embodiment of the present disclosure is a method for
5 producing a fluoroalkyne compound represented by formula (1):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a fluoroalkyl group,

the method comprising subjecting a fluoroalkane
10 compound represented by formula (2):



wherein R^1 and R^2 are as defined above; and X^1 is a fluorine atom and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine atom, to a dehydrofluorination reaction in the presence
15 of a solvent containing an ether,

the method satisfying the following:

(I) the fluoroalkyl group is a C_1 - C_4 fluoroalkyl group.

[0030]

That is, the method for producing a fluoroalkyne
20 compound in the first embodiment of the present disclosure is a method for producing a fluoroalkyne compound represented by formula (1):



wherein R^1 and R^2 are the same or different and each is a fluorine
25 atom or a C_1 - C_4 fluoroalkyl group,

the method comprising subjecting a fluoroalkane
compound represented by formula (2):



wherein R^1 and R^2 are as defined above; and X^1 is a fluorine atom
30 and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine atom, to a dehydrofluorination reaction in the presence of an ether solvent.

[0031]


According to the present disclosure, by performing a
35 dehydrofluorination reaction of the fluoroalkane compound

represented by formula (2), the fluoroalkyne compound represented by formula (1) in which 2 mol of hydrogen fluoride is eliminated per mol of the fluoroalkane compound represented by formula (2) can be obtained in only a single step.

5 [0032]

(1-1) Starting Compound (Fluoroalkane Compound)

The fluoroalkane compound as a substrate that can be used in the production method of the present disclosure (the first embodiment) is a fluoroalkane compound represented by
10 formula (2):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a C_1 - C_4 fluoroalkyl group; X^1 is a fluorine atom and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine
15 atom, as described above.

[0033]

That is, the fluoroalkane compound represented by formula (2) encompasses formulas (2A) and (2B):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a C_1 - C_4 fluoroalkyl group.

[0034]

In formula (2), the fluoroalkyl group represented by
25 each of R^1 and R^2 refers to a group formed by replacing one or more hydrogen atoms of an alkyl group with a fluorine atom, and also includes a perfluoroalkyl group formed by replacing all the hydrogen atoms of an alkyl group. The fluoroalkyl group may be a linear fluoroalkyl group or a branched fluoroalkyl group.

30 Examples include C_1 - C_4 fluoroalkyl groups, such as monofluoromethyl, difluoromethyl, trifluoromethyl, monofluoroethyl, difluoroethyl, trifluoroethyl, tetrafluoroethyl, pentafluoroethylfluoroethyl, monofluoropropyl, difluoropropyl, trifluoropropyl, tetrafluoropropyl, pentafluoropropyl,
35 hexafluoropropyl, heptafluoropropyl, monofluorobutyl,

difluorobutyl, trifluorobutyl, tetrafluorobutyl, heptafluorobutyl, hexafluorobutyl, heptafluorobutyl, octafluorobutyl, and nonafluorobutyl. From the viewpoint of producing the fluoroalkyne compound in particular with high conversion, yield, and
5 selectivity, the fluoroalkyl group is preferably a C₁-C₃ fluoroalkyl group. In addition, from the viewpoint of easily suppressing side reactions, the fluoroalkyl group preferably contains as few hydrogen atoms as possible, and is particularly preferably a perfluoroalkyl group.

10 [0035]

In the fluoroalkane compound as a substrate, R¹ and R² are preferably fluoroalkyl groups, more preferably perfluoroalkyl groups, and even more preferably trifluoromethyl groups, from the viewpoint of producing the fluoroalkyne compound in particular
15 with high conversion, yield, and selectivity.

[0036]

R¹ and R² may be the same or different.

[0037]

Specific examples of the fluoroalkane compound as a
20 substrate that satisfies the above conditions include CF₂HCF₂H, CF₃CFHCF₂H, CF₃CFHCFHCF₃, CF₃CF₂CFHCFHCF₃, CF₃CF₂CFHCFHCF₂CF₃, CFH₂CF₃, CF₃CH₂CF₃, CF₃CH₂CF₂CF₃, CF₃CF₂CH₂CF₂CF₃, CF₃CF₂CH₂CF₂CF₂CF₃, and the like. These fluoroalkane compounds can be used singly or in a combination of two or more. Such fluoroalkane compounds can be
25 known or commercially available products.

[0038]


(1-2) Dehydrofluorination Reaction

In the step of subjecting the fluoroalkane compound to a dehydrofluorination reaction in the present disclosure, for
30 example, in the fluoroalkane compound represented by formula (2) as a substrate, R¹ and R² are more preferably trifluoromethyl groups.

[0039]

That is, this reaction is preferably a
35 dehydrofluorination reaction in which 2 mol of hydrogen fluoride

is eliminated per mol of $\text{CF}_3\text{CFHCFHCF}_3$ or $\text{CF}_3\text{CH}_2\text{CF}_2\text{CF}_3$ according to the following reaction formulas:



5 [0040]

The step of subjecting the fluoroalkane compound to a dehydrofluorination reaction in the present disclosure is performed in the presence of a solvent containing an ether, that is, in a liquid phase using a solvent containing an ether, from
10 the viewpoint of conversion, selectivity, and yield. If the reaction is performed in a gas phase that does not use a solvent or in a liquid phase using a solvent other than a solvent containing an ether, the dehydrofluorination reaction of the fluoroalkene compound does not proceed, and the fluoroalkyne
15 compound cannot be efficiently obtained from the fluoroalkane compound in a single step.

[0041]

In the present disclosure, the yield of the target compound can be further improved, for example, by using a metal
20 container, and increasing the amount of the liquid phase by applying pressure and raising the boiling point of the starting material.

[0042]

Moreover, in the present disclosure, it is preferable
25 to first prepare a solution of the fluoroalkane compound represented by formula (2), and then allow the reaction to proceed in the presence of a base.

[0043]

(1-3) Solvent Containing Ether

30 In the present disclosure, in order to obtain the fluoroalkyne compound from the fluoroalkane compound by a dehydrofluorination reaction, preferably in a single step, the reaction is performed in the presence of a solvent containing an ether.

35 [0044]

However, the water content is preferably small so that the fluoroalkyne compound can be easily obtained from the fluoroalkane compound by a dehydrofluorination reaction. From such a viewpoint, the water concentration of the solvent
5 containing an ether is preferably 500 mass ppm or less, more preferably 400 mass ppm or less, and even more preferably 300 mass ppm or less, based on the total amount of the solvent containing an ether taken as 100 mass%. The lower limit of the water concentration of the solvent containing an ether is not
10 limited, and is preferably 0.01 mass ppm or more from the viewpoint that it can be easily achieved technically.
[0045]

To easily obtain the fluoroalkyne compound from the fluoroalkane compound by a dehydrofluorination reaction, the
15 number of ether bonds in the ether is preferably as small as possible. From such a viewpoint, the number of ether bonds in the ether is preferably 1 to 10, more preferably 1 to 5, even more preferably 1 to 3, and particularly preferably 1 to 2.
[0046]

20 The ether in the solvent containing an ether that satisfies the above conditions is preferably a chain ether, from the viewpoint of conversion, selectivity, and yield. The chain ether includes not only an ether compound having no cyclic structure, but also an ether compound containing a cyclic
25 structure that does not contain an ether bond. That is, the chain ether is a concept that does not include an ether compound in which an ether bond constitutes a cyclic structure, such as tetrahydrofuran, but includes an ether compound containing a cyclic structure that does not contain an ether bond, such as
30 diphenyl ether. When an ether compound having no cyclic structure is used, the yield of the fluoroalkyne compound as the target product can be particularly improved. When an ether compound containing a cyclic structure that does not contain an ether bond is used, the yield of impurities (hydrofluorocarbon compounds
35 other than fluoroalkane compounds, fluoroalkene compounds, and

fluoroalkyne compounds) can be particularly reduced. Specific preferred examples of such ethers include diethyl ether, diisopropyl ether, di(n-butyl) ether, diphenyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, diglyme, and the like. The ether solvents can be used singly or in a combination of two or more. In particular, from the viewpoint of conversion, selectivity, and yield, diethyl ether, diisopropyl ether, di(n-butyl) ether, diphenyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, and the like are preferred, diethyl ether, diisopropyl ether, di(n-butyl) ether, diphenyl ether, and the like are more preferred, and diisopropyl ether, di(n-butyl) ether, diphenyl ether, and the like are even more preferred.

[0047]

In the present disclosure, the ether may be used in combination with other solvents, such as carbonate ester solvents such as dimethyl carbonate, ethyl methyl carbonate, diethyl carbonate, methyl propyl carbonate, and ethyl propyl carbonate; ester solvents such as ethyl acetate, propyl acetate, butyl acetate, methyl propionate, ethyl propionate, and butyl propionate; ketone solvents such as acetone, ethyl methyl ketone, and diethyl ketone; lactone solvents such as γ -butyrolactone, γ -valerolactone, tetrahydrofuran, and tetrahydropyran; cyclic ether solvents such as tetrahydrofuran; nitrile solvents such as acetonitrile, propionitrile, and benzonitrile; amide solvents such as N,N-dimethylformamide; and sulfone solvents such as dimethyl sulfoxide and sulfolane. However, from the viewpoint of, for example, the conversion in reaction, selectivity, and yield, the amount of the ether for use is preferably 80 to 100 vol% (in particular 90 to 100 vol%), and the amount of other solvents is preferably 0 to 20 vol% (in particular 0 to 10 vol%), based on the total amount of the solvents for use in reaction taken as 100 vol%.

[0048]

(1-4) Base

From the viewpoint of conversion, selectivity, and

yield, the step of subjecting a fluoroalkane compound to a dehydrofluorination reaction to obtain a fluoroalkyne compound in the present disclosure is preferably performed in the presence of a base.

5 [0049]

From the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound, the base is preferably at least one member selected from the group consisting of hydroxides and alkoxides of alkali metals and
10 alkaline earth metals (i.e., at least one member selected from the group consisting of alkali metal hydroxides, alkaline earth metal hydroxides, alkali metal alkoxides, and alkaline earth metal alkoxides), more preferably at least one member selected from the group consisting of hydroxides of alkali metals and
15 alkaline earth metals and alkoxides of alkali metals (i.e., at least one member selected from the group consisting of alkali metal hydroxides, alkaline earth metal hydroxides, and alkali metal alkoxides), even more preferably a hydroxide and/or alkoxide of an alkali metal (i.e., an alkali metal hydroxide
20 and/or an alkali metal alkoxide), and particularly preferably an alkali metal hydroxide. Specific examples of bases include sodium hydroxide, potassium hydroxide, magnesium hydroxide, calcium hydroxide, sodium methoxide, potassium tert-butoxide, sodium ethoxide, sodium tert-butoxide, potassium methoxide, potassium
25 ethoxide, potassium tert-butoxide, magnesium methoxide, magnesium ethoxide, magnesium tert-butoxide, calcium methoxide, calcium ethoxide, calcium tert-butoxide, and the like. In particular, from the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound, sodium
30 hydroxide, potassium hydroxide, magnesium hydroxide, calcium hydroxide, sodium methoxide, sodium ethoxide, sodium tert-butoxide, potassium methoxide, potassium ethoxide, potassium tert-butoxide, and the like are preferred, sodium hydroxide, potassium hydroxide, sodium methoxide, sodium ethoxide, sodium
35 tert-butoxide, potassium methoxide, potassium ethoxide, potassium

tert-butoxide, and the like are more preferred, and sodium hydroxide, potassium hydroxide, and the like are even more preferred.

[0050]

5 The amount of the base for use is not limited, and is preferably 1.0 to 10.0 mol, more preferably 1.5 to 5.0 mol, and even more preferably 2.0 to 3.0 mol, per mol of the fluoroalkane compound represented by formula (2).

[0051]

10 (1-5) Reaction Conditions

Closed Reaction Process

 In the present disclosure, the fluoroalkyne compound represented by formula (1) as the target compound has a low boiling point and exists as a gas at room temperature.

15 Accordingly, in the step of performing a dehydrofluorination reaction in the present disclosure, when a closed reaction process is used as the reaction system, the pressure inside the closed reaction process naturally rises, and the reaction can be carried out under pressurized conditions. Therefore, the
20 fluoroalkyne compound represented by formula (1) as the target compound can be obtained with higher selectivity and higher conversion.

[0052]

 Thus, the closed reaction process is pressurized due to
25 the low boiling point of the target compound, and the concentration of the substrate (starting compound) in the ether solvent increases, which can improve the reactivity. To perform the closed reaction process, it is preferable to carry out the reaction while sealing the reaction system using a batch-type
30 pressure-resistant reaction vessel. When the reaction is carried out in a batch process, it is preferable, for example, to place a starting compound, an ether solvent, and if necessary a base etc. in a pressure vessel, such as an autoclave, raise the temperature to an appropriate reaction temperature with a heater, and carry
35 out the reaction with stirring for a certain period of time. As

the reaction atmosphere, it is preferable to carry out the reaction in an inert gas atmosphere, such as nitrogen, helium, or carbon dioxide.

[0053]

5 In the step of performing a dehydrofluorination reaction in the present disclosure, the reaction temperature in the closed reaction process is generally preferably 0 to 400°C, more preferably 25 to 300°C, and even more preferably 50 to 200°C, from the viewpoint of the conversion in reaction and the
10 selectivity and yield of the halogenated alkyne compound.

[0054]

 When no base is used or a hydroxide of an alkali metal and/or alkaline earth metal is used as a base, the reaction temperature in the closed reaction process is generally
15 preferably 25 to 400°C, more preferably 50 to 300°C, and even more preferably 100 to 200°C, from the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound.

[0055]

20 When an alkoxide of an alkali metal and/or alkaline earth metal is used as a base, the reaction temperature is preferably low. This is because if the reaction temperature is too high, the addition reaction of the alkoxide to the double bond proceeds. Thus, the reaction temperature in the closed
25 reaction process is generally preferably 0 to 200°C, more preferably 25 to 150°C, and even more preferably 50 to 100°C, from the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound.

[0056]

30 The reaction time is not limited, and may be a time during which the reaction can sufficiently proceed. Specifically, the reaction can proceed until there is no change in composition in the reaction system.

[0057]

35 Pressure-Increased Reaction Process

In the step of performing a dehydrofluorination reaction in the present disclosure, the reaction can be carried out in a pressure-increased reaction process by setting the reaction temperature to 0°C or more and the reaction pressure to more than 0 kPa. As a result, the fluoroalkyne compound represented by formula (1) as the target compound can be obtained with higher selectivity and higher conversion. When the reaction system is pressurized in this way, the concentration of the substrate (starting compound) in the ether solvent rises, which can improve the reactivity. In the pressure-increased reaction process, it is preferable to carry out the reaction while sealing the reaction system using a batch-type pressure-resistant reaction vessel. When the reaction is carried out in a batch process, it is preferable, for example, to place a starting compound, an ether solvent, and if necessary a base etc. in a pressure vessel, such as an autoclave, raise the temperature to an appropriate reaction temperature with a heater, and carry out the reaction with stirring for a certain period of time.

[0058]

In the step of performing a dehydrofluorination reaction in the present disclosure, the reaction pressure is preferably more than 0 kPa. In the step of performing a dehydrofluorination reaction in the present disclosure, the reaction pressure is preferably more than 0 kPa, more preferably 5 kPa or more, even more preferably 10 kPa or more, and particularly preferably 15 kPa or more. The upper limit of the reaction pressure is not limited and is generally about 2 MPa. In the present disclosure, the pressure is gauge pressure unless otherwise specified. Here, the reaction pressure when a pressure-increased reaction process is used is described. If a pressure-increased reaction process is not used, for example, the reaction can proceed under reduced pressure or ordinary pressure.

[0059]

For pressurization, the pressure in the reaction system can be increased by sending an inert gas, such as nitrogen,

helium, or carbon dioxide, into the reaction system.

[0060]

In the step of performing a dehydrofluorination reaction in the present disclosure, the reaction temperature in the pressure-increased reaction process is generally preferably 0 to 400°C, more preferably 25 to 300°C, and even more preferably 100 to 200°C, from the viewpoint that the elimination reaction can proceed more efficiently and the target compound can be obtained with higher selectivity, and from the viewpoint of suppressing the decrease in the conversion.

[0061]

When no base is used or a hydroxide of an alkali metal and/or alkaline earth metal is used as a base, the reaction temperature in the pressure-increased reaction process is generally preferably 25 to 400°C, more preferably 50 to 300°C, and even more preferably 100 to 200°C, from the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound.

[0062]

When an alkoxide of an alkali metal and/or alkaline earth metal is used as a base, the reaction temperature is preferably low. This is because if the reaction temperature is too high, the addition reaction of the alkoxide to the double bond proceeds. Thus, the reaction temperature in the pressure-increased reaction process is generally preferably 0 to 200°C, more preferably 25 to 150°C, and even more preferably 50 to 100°C, from the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound.

[0063]

Combination of Closed Reaction Process and Pressure-Increased Reaction Process

In the step of performing a dehydrofluorination reaction in the present disclosure, it is also possible to carry out the reaction in a continuous and pressurized reaction form, while extracting the liquid or extracting the gasified product,

for example, by connecting a back pressure valve to a continuous stirred-tank reactor (CSTR).

[0064]

After the dehydrofluorination reaction is completed,
5 purification can be optionally performed according to a conventional method, thereby obtaining a fluoroalkyne compound represented by formula (1).

[0065]

(1-6) Target Compound (Halogenated Alkyne Compound)

10 The target compound of the present disclosure obtained in the above manner is a fluoroalkyne compound represented by formula (1):




wherein R^1 and R^2 are the same or different and each is a fluorine
15 atom or a fluoroalkyl group.

[0066]

R^1 and R^2 in formula (1) correspond to R^1 and R^2 in formula (2) described above, respectively. Accordingly, specific examples of the fluoroalkyne compound represented by formula (1)
20 to be produced include $CF\equiv CF$, $CF_3C\equiv CF$, $CF_3C\equiv CCF_3$, $CF_3CF_2C\equiv CCF_3$, $CF_3CF_2C\equiv CCF_2CF_3$, and the like.

[0067]

According to the production method of the present invention, the fluoroalkyne compound represented by formula (1)
25 may not be obtained alone, and may be obtained in the form of a composition containing the fluoroalkyne compound represented by formula (1). Even in this case, most of the compounds other than the fluoroalkyne compound represented by formula (1) are fluoroalkene compounds represented by formula (4):



wherein R^1 and R^2 are as defined above, in which only 1 mol of hydrogen fluoride is eliminated from the fluoroalkane compound represented by formula (2). Thus, it is possible to further improve the selectivity and yield of the fluoroalkyne compound
35 represented by formula (1) by repeating the above step of

performing a dehydrofluorination reaction.

[0068]

For example, in Example 1 described later, $\text{CF}_3\text{CFHCFHCF}_3$ was used as a starting compound, and $\text{CF}_3\text{C}\equiv\text{CCF}_3$ was obtained in a
5 yield of 49.9 mol% (conversion: 97.9% $\times$ selectivity: 51.00%), and $\text{CF}_3\text{CF}=\text{CHCF}_3$ was obtained in a yield of 41.7 mol% (conversion: 97.9% $\times$ selectivity: 42.62%).

[0069]

In the present disclosure, when the step of performing
10 a dehydrofluorination reaction is repeated, that is, when the number of steps is increased, $\text{CF}_3\text{CF}=\text{CHCF}_3$ obtained as an impurity is used as a starting material, and a dehydrofluorination reaction is performed in the same manner. Thus, if the yield of $\text{CF}_3\text{CF}=\text{CHCF}_3 \rightarrow \text{CF}_3\text{C}\equiv\text{CCF}_3$ in the second step is also 49.9 mol%,
15 $\text{CF}_3\text{C}\equiv\text{CCF}_3$ is obtained in an amount that is 70.7 mol% of $\text{CF}_3\text{CFHCFHCF}_3$, which is the starting material, in total in the two steps.

[0070]

When the first step is $\text{CF}_3\text{CFHCFHCF}_3 \rightarrow \text{CF}_3\text{CF}=\text{CHCF}_3$ and the
20 second step is $\text{CF}_3\text{CF}=\text{CHCF}_3 \rightarrow \text{CF}_3\text{C}\equiv\text{CCF}_3$ as in the conventional methods, if the yield in the first step is 100 mol%, $\text{CF}_3\text{C}\equiv\text{CCF}_3$ is obtained in an amount that is 49.9 mol% of $\text{CF}_3\text{CFHCFHCF}_3$, which is the starting material, in the two steps since the yield in the second step is 49.9 mol% as described above.

25 [0071]

As described above, according to the present disclosure, it is possible to obtain a specific fluoroalkyne compound using a specific fluoroalkane compound as a starting compound by only a single step, and it is possible to improve the yield of the
30 fluoroalkyne compound when the dehydrofluorination reaction is repeated a plurality of times, compared with the conventional methods. In any case, it can be understood that the production method of the present disclosure is useful.

[0072]

35 The thus-obtained fluoroalkyne compound can be

effectively used for various applications, such as etching gases for forming cutting-edge microstructures in semiconductors, liquid crystals, etc., as well as cleaning gases, deposit gases, refrigerants, heat transfer media, and building blocks for organic synthesis. The deposit gas and the building block for organic synthesis are described later.

[0073]


2. Method for Producing Fluoroalkyne Compound (Second Embodiment)

The method for producing a fluoroalkyne compound in the second embodiment of the present disclosure is a method for producing a fluoroalkyne compound represented by formula (1):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a fluoroalkyl group,

the method comprising subjecting a fluoroalkane compound represented by formula (2):



wherein R^1 and R^2 are as defined above; and X^1 is a fluorine atom and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine atom, to a dehydrofluorination reaction in the presence of a solvent containing an ether,

(II) the ether being a chain ether, and the dehydrofluorination reaction being performed in the presence of the solvent containing a chain ether and a base comprising a hydroxide and/or alkoxide of an alkali metal and/or alkaline earth metal.

[0074]


That is, the method for producing a fluoroalkyne compound in the second embodiment of the present disclosure is a method for producing a fluoroalkyne compound represented by formula (1):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a fluoroalkyl group,

the method comprising subjecting a fluoroalkane

compound represented by formula (2):



wherein R^1 and R^2 are as defined above; and X^1 is a fluorine atom and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine atom, to a dehydrofluorination reaction in the presence of a solvent containing a chain ether and a base comprising a hydroxide and/or alkoxide of an alkali metal and/or alkaline earth metal.


[0075]

According to the present disclosure, by performing a dehydrofluorination reaction of the fluoroalkane compound represented by formula (2), the fluoroalkyne compound represented by formula (1) in which 2 mol of hydrogen fluoride is eliminated per mol of the fluoroalkane compound represented by formula (2) can be obtained in only a single step.

[0076]

(2-1) Starting Compound (Fluoroalkane Compound)

The fluoroalkane compound as a substrate that can be used in the production method of the present disclosure (the second embodiment) is a fluoroalkane compound represented by formula (2):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a fluoroalkyl group; and X^1 is a fluorine atom and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine atom, as described above.

[0077]

That is, the fluoroalkane compound represented by formula (2) encompasses formulas (2A) and (2B):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a fluoroalkyl group.

[0078]

In formula (2), the fluoroalkyl group represented by

each of R^1 and R^2 refers to a group formed by replacing one or more hydrogen atoms of an alkyl group with a fluorine atom, and also includes a perfluoroalkyl group formed by replacing all the hydrogen atoms of an alkyl group. The fluoroalkyl group may be a
5 linear fluoroalkyl group or a branched fluoroalkyl group. Examples include C_1 - C_{10} fluoroalkyl groups, such as monofluoromethyl, difluoromethyl, trifluoromethyl, monofluoroethyl, difluoroethyl, trifluoroethyl, tetrafluoroethyl, pentafluoroethylfluoroethyl, monofluoropropyl, difluoropropyl,
10 trifluoropropyl, tetrafluoropropyl, pentafluoropropyl, hexafluoropropyl, and heptafluoropropyl. From the viewpoint of producing the fluoroalkyne compound in particular with high conversion, yield, and selectivity, the fluoroalkyl group is preferably a C_1 - C_4 fluoroalkyl group, and more preferably a C_1 - C_3
15 fluoroalkyl group.
[0079]

In the fluoroalkane compound as a substrate, R^1 and R^2 are preferably fluoroalkyl groups, more preferably perfluoroalkyl groups, and even more preferably trifluoromethyl groups, from the
20 viewpoint of producing the fluoroalkyne compound in particular with high conversion, yield, and selectivity.
[0080]

R^1 and R^2 may be the same or different.
[0081]

Specific examples of the fluoroalkane compound as a
25 substrate that satisfies the above conditions include CF_2HCF_2H , CF_3CFHCF_2H , $CF_3CFHCFHCF_3$, $CF_3CF_2CFHCFHCF_3$, $CF_3CF_2CFHCFHCF_2CF_3$, CFH_2CF_3 , $CF_3CH_2CF_3$, $CF_3CH_2CF_2CF_3$, $CF_3CF_2CH_2CF_2CF_3$, $CF_3CF_2CH_2CF_2CF_2CF_3$, and the like. These fluoroalkane compounds can be used singly or in a
30 combination of two or more. Such fluoroalkane compounds can be known or commercially available products.
[0082]

(2-2) Dehydrofluorination Reaction

The "dehydrofluorination reaction" can be the same as
35 the "(1-2) Dehydrofluorination Reaction" described above. Various

conditions can also be the same as those described above.

[0083]

(2-3) Solvent Containing Chain Ether

In the present disclosure, the reaction is performed in
5 the presence of a solvent containing a chain ether to obtain a
fluoroalkyne compound from a fluoroalkane compound by a
dehydrofluorination reaction, preferably in a single step.

[0084]

However, the water content is preferably small so that
10 the fluoroalkyne compound can be easily obtained from the
fluoroalkane compound by a dehydrofluorination reaction. From
such a viewpoint, the water concentration of the solvent
containing a chain ether is preferably 500 mass ppm or less, more
preferably 400 mass ppm or less, and even more preferably 300
15 mass ppm or less, based on the total amount of the solvent
containing a chain ether taken as 100 mass%. The lower limit of
the water concentration of the solvent containing a chain ether
is not limited, and is preferably 0.01 mass ppm or more from the
viewpoint that it can be easily achieved technically.

20 [0085]

In addition, to easily obtain the fluoroalkyne compound
from the fluoroalkane compound by a dehydrofluorination reaction,
the number of ether bonds in the chain ether is preferably as
small as possible. From such a viewpoint, the number of ether
25 bonds in the chain ether is preferably 1 to 10, more preferably 1
to 5, even more preferably 1 to 3, and particularly preferably 1
to 2.

[0086]

The chain ether in the solvent containing a chain ether
30 that satisfies the above conditions includes not only an ether
compound having no cyclic structure, but also an ether compound
containing a cyclic structure that does not contain an ether bond.
That is, the chain ether is a concept that does not include an
ether compound in which an ether bond constitutes a cyclic
35 structure, such as tetrahydrofuran, but includes an ether

compound containing a cyclic structure that does not contain an ether bond, such as diphenyl ether. When an ether compound having no cyclic structure is used, the yield of the fluoroalkyne compound as the target product can be particularly improved. When
5 an ether compound containing a cyclic structure that does not contain an ether bond is used, the yield of impurities (hydrofluorocarbon compounds other than fluoroalkane compounds, fluoroalkene compounds, and fluoroalkyne compounds) can be particularly reduced. Specific preferred examples of such chain
10 ethers include diethyl ether, diisopropyl ether, di(n-butyl) ether, diphenyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, diglyme, and the like. The chain ether solvents can be used singly or in a combination of two or more. In particular, from the viewpoint of conversion, selectivity, and yield, diethyl
15 ether, diisopropyl ether, di(n-butyl) ether, diphenyl ether, 1,2-dimethoxyethane, 1,2-diethoxyethane, and the like are preferred, diethyl ether, diisopropyl ether, di(n-butyl) ether, diphenyl ether, and the like are more preferred, and diisopropyl ether, di(n-butyl) ether, diphenyl ether, and the like are even more
20 preferred.
[0087]

In the present disclosure, the chain ether may be used in combination with other solvents, such as carbonate ester solvents such as dimethyl carbonate, ethyl methyl carbonate,
25 diethyl carbonate, methyl propyl carbonate, and ethyl propyl carbonate; ester solvents such as ethyl acetate, propyl acetate, butyl acetate, methyl propionate, ethyl propionate, and butyl propionate; ketone solvents such as acetone, ethyl methyl ketone, and diethyl ketone; lactone solvents such as γ -butyrolactone, γ -valerolactone, tetrahydrofuran, and tetrahydropyran; cyclic ether
30 solvents such as tetrahydrofuran; nitrile solvents such as acetonitrile, propionitrile, and benzonitrile; amide solvents such as N,N-dimethylformamide; and sulfone solvents such as dimethyl sulfoxide and sulfolane. However, from the viewpoint of,
35 for example, the conversion in reaction, selectivity, and yield,

the amount of other solvents for use is preferably small. Specifically, the amount of the chain ether for use is preferably 80 to 100 vol% (in particular 90 to 100 vol%), and the amount of other solvents is preferably 0 to 20 vol% (in particular 0 to 10
5 vol%), based on the total amount of the solvents for use in reaction taken as 100 vol%.

[0088]

(2-4) Base

From the viewpoint of conversion, selectivity, and
10 yield, the step of subjecting a fluoroalkane compound to a dehydrofluorination reaction to obtain a fluoroalkyne compound in the present disclosure is performed in the presence of a base comprising a hydroxide and/or alkoxide of an alkali metal and/or alkaline earth metal.

15 [0089]

From the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound, the base is preferably at least one member selected from the group consisting of hydroxides and alkoxides of alkali metals and
20 alkaline earth metals (i.e., at least one member selected from the group consisting of alkali metal hydroxides, alkaline earth metal hydroxides, alkali metal alkoxides, and alkaline earth metal alkoxides), preferably at least one member selected from the group consisting of hydroxides of alkali metals and alkaline
25 earth metals and alkoxides of alkali metals (i.e., at least one member selected from the group consisting of alkali metal hydroxides, alkaline earth metal hydroxides, and alkali metal alkoxides), more preferably a hydroxide and/or alkoxide of an alkali metal (i.e., an alkali metal hydroxide and/or an alkali
30 metal alkoxide), and even more preferably an alkali metal hydroxide. Specific examples of bases include sodium hydroxide, potassium hydroxide, magnesium hydroxide, calcium hydroxide, sodium methoxide, potassium tert-butoxide, sodium ethoxide, sodium tert-butoxide, potassium methoxide, potassium ethoxide,
35 potassium tert-butoxide, magnesium methoxide, magnesium ethoxide,

magnesium tert-butoxide, calcium methoxide, calcium ethoxide, calcium tert-butoxide, and the like. In particular, from the viewpoint of the conversion in reaction and the selectivity and yield of the halogenated alkyne compound, sodium hydroxide, potassium hydroxide, magnesium hydroxide, calcium hydroxide, sodium methoxide, sodium ethoxide, sodium tert-butoxide, potassium methoxide, potassium ethoxide, potassium tert-butoxide, and the like are preferred, sodium hydroxide, potassium hydroxide, sodium methoxide, sodium ethoxide, sodium tert-butoxide, potassium methoxide, potassium ethoxide, potassium tert-butoxide, and the like are more preferred, and sodium hydroxide, potassium hydroxide, and the like are even more preferred.

[0090]

The amount of the base for use is not limited and is preferably about 1 mol per mol of the fluoroalkane compound represented by formula (2). Specifically, the amount of the base for use is preferably 1.0 to 10.0 mol, more preferably 1.5 to 5.0 mol, and even more preferably 2.0 to 3.0 mol, per mol of the fluoroalkane compound represented by formula (2).

[0091]

(2-5) Reaction Conditions and Target Compound

The "reaction conditions" can be the same as the "(1-5) Reaction Conditions" described above. The "target compound" can be the same as the "(1-6) Target Compound" described above. Various conditions can also be the same as those described above.

[0092]

3. Composition

As described above, a fluoroalkyne compound can be obtained from a fluoroalkane compound preferably in a single step, and may also be obtained in a form of a composition containing a fluoroalkyne compound. In particular, according to the production method of the present disclosure, for example, a composition containing a fluoroalkyne compound represented by formula (1) and a fluoroalkene compound represented by formula (4), in which only 1 mol of hydrogen fluoride is eliminated from a fluoroalkane

compound represented by formula (2):



wherein R^1 and R^2 are as defined above, can be produced.

[0093]

5 R^1 and R^2 in formula (4) correspond to R^1 and R^2 in formula (2) described above, respectively. Thus, specific examples of the fluoroalkene compound represented by formula (4) include $CF_2=CFH$, $CF_3CF=CFH$, $CF_3CH=CFH$, $CF_3CF=CHCF_3$, $CF_3CF_2CF=CHCF_3$, $CF_3CF_2CH=CFCF_3$, $CF_3CF_2CF=CHCF_2CF_3$, and the like. The composition may
10 contain one of these fluoroalkene compounds or a combination of two or more of these fluoroalkene compounds.

[0094]

 According to the production method of the present disclosure, a composition that contains a fluoroalkyne compound
15 represented by formula (1) and a fluoroalkene compound represented by formula (4) in similar amounts and contains a small amount of other components tends to be produced. Thus, in the composition containing a fluoroalkyne compound in the present disclosure, the content of the fluoroalkyne compound represented
20 by formula (1) is preferably 25.00 mol% or more, more preferably 30.00 mol% or more, and even more preferably 35.00 mol% or more, based on the total amount of the composition taken as 100 mol%. In the composition containing a fluoroalkyne compound in the present disclosure, the content of the fluoroalkyne compound
25 represented by formula (1) may also be, for example, 75.00 mol% or less, 70.00 mol% or less, or 65.00 mol% or less, based on the total amount of the composition taken as 100 mol%. In the composition containing a fluoroalkyne compound in the present disclosure, the content of the fluoroalkene compound represented
30 by formula (4) may be 25.00 mol% or more, 30.00 mol% or more, or 35.00 mol% or more, based on the total amount of the composition taken as 100 mol%. In the composition containing a fluoroalkyne compound in the present disclosure, the content of the fluoroalkene compound represented by formula (4) is also
35 preferably 75.00 mol% or less, more preferably 70.00 mol% or less,

and even more preferably 65.00 mol% or less, based on the total amount of the composition taken as 100 mol%. Furthermore, in the composition containing a fluoroalkyne compound in the present disclosure, the total content of the fluoroalkyne compound
5 represented by formula (1) and the fluoroalkene compound represented by formula (4) is preferably 90.00 mol% or more, more preferably 91.00 mol% or more, and even more preferably 92.00 mol% or more, based on the total amount of the composition taken as 100 mol%. In the composition containing a fluoroalkyne
10 compound in the present disclosure, the total content of the fluoroalkyne compound represented by formula (1) and the fluoroalkene compound represented by formula (4) may be 100.00 mol% or less, 99.99 mol% or less, or 99.98 mol% or less, based on the total amount of the composition taken as 100 mol%.
15 [0095]

As described above, according to the production method of the present disclosure, hydrofluorocarbon compounds, such as trifluoromethane (R23), difluoromethane (R32), tetrafluoromethane (R14), monofluoromethane (R41), 1,2-difluoroethylene (R1132), and
20 1,1,2-trifluoroethylene (R1123) may be contained as components other than the fluoroalkyne compound represented by formula (1) and the fluoroalkene compound represented by formula (4) in the composition obtained. In this case, in the composition containing a fluoroalkyne compound in the present disclosure, the content of
25 these other hydrofluorocarbon compounds may be 0.00 mol% or more, 0.01 mol% or more, or 0.02 mol% or more, based on the total amount of the composition taken as 100 mol%. In the composition containing a fluoroalkyne compound in the present disclosure, the content of these other hydrofluorocarbon compounds is preferably
30 10.00 mol% or less, more preferably 7.50 mol% or less, and even more preferably 5.00 mol% or less, based on the total amount of the composition taken as 100 mol%.
[0096]

According to the production method of the present
35 disclosure, a fluoroalkyne compound can be obtained in particular

with high selectivity and yield compared with conventional methods for obtaining a fluoroalkyne compound from a fluoroalkane compound; and as a result, purification for obtaining a fluoroalkyne compound can be efficiently performed.

5 [0097]

The composition containing a fluoroalkyne compound in the present disclosure can be effectively used for various applications, such as etching gases for forming cutting-edge microstructures in semiconductors, liquid crystals, etc., as well
10 as cleaning gases, deposit gases, refrigerants, heat transfer media, and building blocks for organic synthesis, as in the fluoroalkyne compound alone.

[0098]

The deposit gas is a gas that deposits an etching
15 resistant polymer layer.

[0099]

The building block for organic synthesis refers to a substance that can be a precursor of a compound having a highly reactive skeleton. For example, when the composition of the
20 present disclosure is reacted with a fluorine-containing organosilicon compound, such as $\text{CF}_3\text{Si}(\text{CH}_3)_3$, it is possible to introduce a fluoroalkyl group, such as a CF_3 group, to convert it into a substance that can be a cleaner or a fluorine-containing pharmaceutical intermediate.

25 [0100]

Although embodiments of the present disclosure are described above, various changes in form and details can be made without departing from the spirit and scope of the claims.


30 Examples

[0101]

The features of the present disclosure are clarified below while showing Examples. The present disclosure is not limited to these Examples.

35 [0102]

In the methods for producing a fluoroalkyne compound of Examples 1 to 6 and Comparative Example 1, the starting compound was a fluoroalkane compound represented by formula (2) wherein R¹ and R³ are trifluoromethyl groups, and a fluoroalkyne compound was
5 obtained by a dehydrofluorination reaction according to the following reaction formula:



[0103]

Examples 1 to 6 and Comparative Example 1

10 Sodium hydroxide (NaOH), potassium hydroxide (KOH), sodium tert-butoxide (t-BuONa), or potassium tert-butoxide (t-BuOK) as a base, and di(n-butyl) ether (Bu₂O), diphenyl ether (Ph₂O), diglyme, or water as a solvent were added to an autoclave, and a starting compound (CF₃CFHCFHCF₃) was further added. The
15 autoclave was closed with the lid to form a closed system. The concentration of the base added in the solvent was 2.5 mol/L. At this stage, the amount of the base for use was adjusted to 2 mol per mol of the starting compound. The pressure at that time was 0 kPa. Thereafter, the mixture was heated to 100°C or 150°C and
20 stirred to allow a reaction to proceed. After the dehydrofluorination reaction was started, sampling was suitably performed, and the reaction was considered to have completed when there was no change in composition in the reaction system. The pressure at the completion of the reaction was 500 kPa.

25 [0104]

After stirring was stopped, the reaction mixture was cooled to room temperature. Mass spectrometry was performed by gas chromatography (Shimadzu Corporation, trade name: GC-2014) according to gas chromatography-mass spectrometry (GC/MS), and
30 structural analysis was performed by using an NMR (JEOL Ltd., trade name: 400YH) based on an NMR spectrum. From the results of mass spectrometry and structural analysis, CF₃C≡CCF₃ was confirmed to have formed as a target compound. Table 1 shows the results. In Table 1, R23 denotes trifluoromethane, R1132 denotes 1,2-
35 difluoroethylene, PF2B denotes CF₃C≡CCF₃, which is the target

compound, and 1327myz denotes $\text{CF}_3\text{CF}=\text{CHCF}_3$.

[0105]

Table 1

Run	Solvent	Base	Temperature	Conversion	Selectivity [%]				
			[°C]	[%]	R23	R1132	PF2B	1327myz	Others
Ex. 1	Bu2O	KOH	150	97.9	2.94	1.53	51.00	42.62	1.9
Ex. 2	Bu2O	NaOH	150	97.6	2.55	1.26	51.40	43.60	1.2
Ex. 3	Diglyme	KOH	150	98.5	1.16	0.25	25.17	73.12	0.3
Ex. 4	Ph2O	KOH	150	99.3	0.88	0.45	47.17	49.82	1.7
Ex. 5	Bu2O	t-BuOK	100	99.5	0.037	0.034	32.5	67.2	0.3
Ex. 6	Bu2O	t-BuONa	100	99.5	0.22	0.09	34.03	64.64	1.0
Comp. Ex. 1	Water	KOH	150	96.3	1.24	1.55	0.30	95.40	1.5

[0106]

Example 7 and Comparative Example 2

Under the etching conditions of ICP (Inductive Coupled Plasma) discharge power: 800 W, bias power: 100 W, pressure: 3 mmTorr (0.399 Pa), electron density: 8×10^{10} to 2×10^{11} cm⁻³, and electron temperature: 5 to 7 eV, an SiO₂ film and a resist film were etched using the composition (containing 51.00 mol% of CF₃C≡CCF₃ and 42.62 mol% of CF₃CF=CHCF₃) obtained in Example 1, and the etching rate and the resist selectivity (etching rate of SiO₂ film/etching rate of resist) were measured (Example 7). The same experiment was performed, except that the only gas used was CF₃C≡CCF₃ (Comparative Example 2).

[0107]

Table 2 shows the results.

15 [0108]


Table 2

	Gas	Resist Selectivity
Example 7	Mixed gas of CF ₃ CCCF ₃ and CF ₃ CFCHCF ₃	5.0
Comparative Example 2	Only CF ₃ CCCF ₃	2.1

CLAIMS

[Claim 1]

A method for producing a fluoroalkyne compound represented by formula (1):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a fluoroalkyl group,

the method comprising subjecting a fluoroalkane compound represented by formula (2):



wherein R^1 and R^2 are as defined above; and X^1 is a fluorine atom and X^2 is a hydrogen atom, or X^1 is a hydrogen atom and X^2 is a fluorine atom, to a dehydrofluorination reaction in the presence of a solvent containing an ether,

15 the method satisfying at least one requirement selected from the group consisting of the following:

(I) the fluoroalkyl group is a C_1 - C_4 fluoroalkyl group, and

(II) the ether is a chain ether, and the dehydrofluorination reaction is performed in the presence of the solvent containing a chain ether, and a base comprising a hydroxide and/or alkoxide of
20 an alkali metal and/or alkaline earth metal.

[Claim 2]

The production method according to claim 1, wherein the
25 water concentration of the solvent containing an ether is 0.01 to 500 mass ppm based on the total amount of the solvent containing an ether taken as 100 mass%.

[Claim 3]

30 The production method according to claim 1 or 2, wherein the number of ether bonds in the ether is 1 to 10.

[Claim 4]

The production method according to any one of claims 1
35 to 3, which satisfies requirement (I).

[Claim 5]

The production method according to claim 4, wherein the fluoroalkyl group is represented by formula (3):

5 $-\text{CF}_2\text{R}^3$ (3)

wherein R^3 is a fluorine atom or a $\text{C}_1\text{-C}_3$ fluoroalkyl group.

[Claim 6]

10 The production method according to claim 4 or 5,
wherein the dehydrofluorination reaction is performed in the presence of a base.

[Claim 7]

15 The production method according to claim 6, wherein the
base is a hydroxide and/or alkoxide of an alkali metal and/or alkaline earth metal.

[Claim 8]

20 The production method according to any one of claims 1
to 3, which satisfies requirement (II).

[Claim 9]

25 The production method according to claim 8, wherein the
fluoroalkyl group has 1 to 10 carbon atoms.

[Claim 10]


30 The production method according to claim 8 or 9,
wherein the fluoroalkyl group is represented by formula (3):
 $-\text{CF}_2\text{R}^3$ (3)
wherein R^3 represents a fluorine atom or a $\text{C}_1\text{-C}_9$ fluoroalkyl group.

[Claim 11]

35 The production method according to any one of claims 1
to 10, wherein the reaction temperature in the
dehydrofluorination reaction is 0 to 300°C.

[Claim 12]

A composition comprising a fluoroalkyne compound represented by formula (1):



wherein R^1 and R^2 are the same or different and each is a fluorine atom or a C_1 - C_4 fluoroalkyl group, and a fluoroalkene compound represented by formula (4):



10 wherein R^1 and R^2 are as defined above,

the content of the fluoroalkyne compound represented by formula (1) being 25.00 to 75.00 mol%, the content of the fluoroalkene compound represented by formula (4) being 25.00 to 75.00 mol%, and the total content of the fluoroalkyne compound represented by formula (1) and the fluoroalkene compound represented by formula (4) being 90.00 to 100.00 mol%, based on the total amount of the composition taken as 100 mol%.

[Claim 13]

20 The composition according to claim 12, which is for use as a cleaning gas, an etching gas, a refrigerant, a heat transfer medium, or a building block for organic synthesis.