PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K

A2

(11) International Publication Number: WO 97/15269

(43) International Publication Date: 1 May 1997 (01.05.97)

(21) International Application Number: PCT/IB96/01485

(22) International Filing Date: 24 October 1996 (24.10.96)

(30) Priority Data:

60/007,071	26 October 1995 (26.10.95)	US
08/608,776	29 February 1996 (29.02.96)	US
08/733,142	16 October 1996 (16.10.96)	US

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published


Without international search report and to be republished upon receipt of that report.

(71) Applicant: BAKER NORTON PHARMACEUTICALS, INC. [US/US]; 4400 Biscayne Boulevard, Miami, FL 33137-3227 (US).

(72) Inventors: BRODER, Samuel; 759 Heron Road, Fort Lauderdale, FL 33326 (US). DUCHIN, Kenneth, L.; 2679 Cypress Lane, Fort Lauderdale, FL 33332 (US). SELIM, Sami; 17 Colonial, Irvine, CA 92720 (US).

(74) Agent: SCHIFFMILLER, Martin, W.; Kirschstein, Ottinger, Israel & Schiffmiller, P.C., 551 Fifth Avenue, New York, NY 10176 (US).

(54) Title: METHOD, COMPOSITIONS AND KITS FOR INCREASING THE ORAL BIOAVAILABILITY OF PHARMACEUTICAL AGENTS



(57) Abstract

A method of increasing the bioavailability upon oral administration of a pharmacologically active target agent, particularly an antitumor or antineoplastic agent which exhibits poor or inconsistent oral bioavailability (e.g., paclitaxel, docetaxel or etoposide), comprises the oral co-administration to a mammalian patient of the target agent and an oral bioavailability-enhancing agent (e.g., cyclosporin A, cyclosporin D, cyclosporin F or ketoconazole). The enhancing agent may be administered orally from 0.5-24 hrs. prior to the oral administration of one or more doses of the target agent, substantially simultaneously with the target agent or both prior to and substantially simultaneously with the target agent. A method of treating mammalian patients suffering from diseases responsive to target agents with poor oral bioavailability, as well as oral dosage forms containing such target agents, combination oral dosage forms containing bioavailability-enhancing agents and and target agents kits containing enhancing and target agent dosage forms and dosing information for the co-administration of the same are also disclosed.