


$$\begin{array}{c}
 \text{O} \\
 \parallel \\
 (\text{R}^0)_p - \text{C}_6\text{H}_3\text{N}(\text{R}^2)\text{C}_6\text{H}_3(\text{R}^1) - \text{CONH} - \text{C}_6\text{H}_3(\text{R}^6) - \text{A} - \text{B} - \text{CH}_2 - \text{N}(\text{R}^3) - (\text{CH}_2)_m - \text{C}_6\text{H}_3(\text{R}^4, \text{R}^5) \quad (\text{I})
 \end{array}$$

**R<sup>6</sup> represents a hydrogen atom or a C<sub>1-4</sub>alkyl or C<sub>1-4</sub>alkoxy group**

$R^7$  represents a hydrogen atom or  $R^3$  and  $R^7$  together form a group  $-(CH_2)_n-$  where  $n$  represents 1 or 2;

$R^8$  represents a hydrogen atom or a  $C_{1-4}$ alkoxy group;  
the group



10 is attached at the benzene ring 3 or 4 position relative to the carboxamide substituent, provided that when the group is attached at the benzene ring 3 position then  $R^6$  must be attached at the benzene ring 6 position;  
and salts and solvates thereof.

15 The novel compounds of formula (I) can sensitize multidrug-resistant cancer cells to chemotherapeutic agents and may be formulated for use in therapy, particularly to improve or increase the efficacy of an antitumour drug.

20

25

THIS SHEET HAS BEEN  
REPLACED

30

This Sheet Has Been Replaced

