
<15.2> The method according to <15> or <15.1>, wherein the (S)-7-(2-methoxy-3,5-dimethylpyridin-4-yl)-1-(tetrahydrofuran-3-yl)-1H-pyrazolo[4,3-c]quinoline-4(5H)-one or pharmaceutically acceptable salt thereof is (S)-7-(2-methoxy-3,5-dimethylpyridin-4-yl)-1-(tetrahydrofuran-3-yl)-1H-pyrazolo[4,3-c]quinoline-4(5H)-one maleate.


<15.3> The method according to any one of <15> to <15.2>, wherein the Alzheimer's disease is mild, moderate or severe Alzheimer's disease.

<16> A pharmaceutical composition comprising (S)-7-(2-methoxy-3,5-dimethylpyridin-4-yl)-1-(tetrahydrofuran-3-yl)-1H-pyrazolo[4,3-c]quinoline-4(5H)-one represented by formula (I):



(I)

or a pharmaceutically acceptable salt thereof, donepezil represented by formula (II):



(II)

or a pharmaceutically acceptable salt thereof,

and an excipient.

<16.1> The pharmaceutical composition according to <16>, wherein the donepezil or pharmaceutically acceptable salt thereof is donepezil hydrochloride.

<16.2> The pharmaceutical composition according to <16> or <16.1>, wherein the (S)-7-(2-methoxy-3,5-dimethylpyridin-4-yl)-1-(tetrahydrofuran-3-yl)-1H-pyrazolo[4,3-c]quinoline-4(5H)-one or pharmaceutically acceptable salt thereof is (S)-7-(2-methoxy-3,5-dimethylpyridin-4-yl)-1-(tetrahydrofuran-3-yl)-1H-pyrazolo[4,3-c]quinoline-4(5H)-one maleate.

<17> A kit comprising:

a pharmaceutical composition comprising (S)-7-(2-methoxy-3,5-dimethylpyridin-4-yl)-1-(tetrahydrofuran-3-yl)-1H-pyrazolo[4,3-c]qui